

Characterizations of 3-manifolds with diagonal metrics as almost η -Ricci solitons

Adara M. Blaga

Abstract

We characterize the 3-dimensional manifold \mathbb{R}^3 with a diagonal Riemannian metric as an almost η -Ricci soliton from the following points of view: under certain assumptions, we determine the potential vector field if η is given; we get obstructions on the metric when the potential vector field has a particular expression; and we compute the defining functions of the soliton when both the potential vector field and the 1-form are prescribed. Moreover, we find conditions for the manifold to be flat or Ricci-flat. Based on the theoretical results we provide many examples.

1 Introduction

The solitons' theory has been lately intensively investigated. The stationary solutions of the Ricci flow, namely, the *Ricci solitons*, still represent a very actual topic to be studied from various points of view. Problems like linear stability of compact Ricci solitons, curvature estimates and rigidity results for gradient Ricci solitons, etc. have been recently treated by Huai-Dong Cao *et. al.* in [9, 8, 7]. As a generalization of the notion of Ricci soliton given by Hamilton in [12], an *almost η -Ricci soliton* [4, 5, 10] is a complete Riemannian manifold (M, \tilde{g}) with a smooth vector field V which satisfies

$$(1) \quad \frac{1}{2} \mathcal{L}_V \tilde{g} + \text{Ric} + \lambda \tilde{g} + \mu \eta \otimes \eta = 0$$

2010 *Mathematics Subject Classification.* 35Q51, 53B25, 53B50.

Key words and phrases. Almost η -Ricci soliton; Killing vector field; Ricci-flat manifold.

for λ and $\mu \neq 0$ two smooth functions on M , where Ric is the Ricci curvature tensor field, \mathcal{L}_V is the Lie derivative in the direction of V and η is a 1-form on M . In particular, if $\eta = 0$, then $(M, \tilde{g}, V, \lambda)$ is called an *almost Ricci soliton* [13]. In this case, the soliton is said to be *shrinking*, *steady* or *expanding* according as λ is negative, zero or positive, respectively [11]. In both of the cases, if λ and μ are real constants, then we drop "almost". Moreover, if V is a Killing vector field, i.e., $\mathcal{L}_V \tilde{g} = 0$, then the soliton is called *trivial*, and in this case, (M, \tilde{g}) is a quasi-Einstein manifold.

The aim of our paper is to characterize 3-dimensional manifolds \mathbb{R}^3 with diagonal Riemannian metrics as almost η -Ricci solitons. More precisely, under certain assumptions, we determine the potential vector field V when η is given, we find the conditions that must be satisfied by the Riemannian metric when the potential vector field has a particular expression, and we compute the defining functions λ and μ when both the potential vector field and the 1-form are prescribed. Based on the theoretical results, we construct a lot of examples, among which the 3-dimensional Sol_3 and $\mathbb{H}^2 \times \mathbb{R}$ Lie groups.

2 \mathbb{R}^3 as an almost η -Ricci soliton

We consider the 3-dimensional manifold \mathbb{R}^3 endowed with a diagonal Riemannian metric as an almost η -Ricci soliton and, under certain assumptions, we concretely determine the potential vector field V when η is given or the 1-form η when V has a particular expression. An analogous study for the canonical metric has been previously done by the present author in [6]. More recently, some particular Riemannian manifolds, namely, the 3-dimensional Sol_3 and $\mathbb{H}^2 \times \mathbb{R}$ Lie groups, have been described as Ricci solitons by Belarbi, Atashpeykara and Haji-Badali in [3, 2, 1] from a similar point of view.

Let \tilde{g} be the Riemannian metric on \mathbb{R}^3 given by:

$$\tilde{g} = \frac{1}{f_1^2} dx^1 \otimes dx^1 + \frac{1}{f_2^2} dx^2 \otimes dx^2 + dx^3 \otimes dx^3,$$

where f_1 and f_2 are two smooth functions nowhere zero on \mathbb{R}^3 , and $\{x^1, x^2, x^3\}$ denotes the standard coordinates in \mathbb{R}^3 . Let

$$\left\{ E_1 := f_1 \frac{\partial}{\partial x^1}, \quad E_2 := f_2 \frac{\partial}{\partial x^2}, \quad E_3 := \frac{\partial}{\partial x^3} \right\}$$

be a local orthonormal frame field, and, for the sake of simplicity, we will make the following notations:

$$\frac{f_2}{f_1} \cdot \frac{\partial f_1}{\partial x^2} =: a, \quad \frac{1}{f_1} \cdot \frac{\partial f_1}{\partial x^3} =: b, \quad \frac{f_1}{f_2} \cdot \frac{\partial f_2}{\partial x^1} =: c, \quad \frac{1}{f_2} \cdot \frac{\partial f_2}{\partial x^3} =: d.$$

Computing the Lie brackets $[X, Y] := X \circ Y - Y \circ X$, we get:

$$[E_1, E_2] = -aE_1 + cE_2 = -[E_2, E_1],$$

$$[E_1, E_3] = -bE_1 = -[E_3, E_1], \quad [E_2, E_3] = -dE_2 = -[E_3, E_2].$$

The Levi-Civita connection ∇ of \tilde{g} deduced from the Koszul's formula

$$\begin{aligned} 2\tilde{g}(\nabla_X Y, Z) &= X(\tilde{g}(Y, Z)) + Y(\tilde{g}(Z, X)) - Z(\tilde{g}(X, Y)) - \\ &\quad - \tilde{g}(X, [Y, Z]) + \tilde{g}(Y, [Z, X]) + \tilde{g}(Z, [X, Y]) \end{aligned}$$

is given by:

$$\nabla_{E_1} E_1 = aE_2 + bE_3, \quad \nabla_{E_1} E_2 = -aE_1, \quad \nabla_{E_1} E_3 = -bE_1, \quad \nabla_{E_2} E_1 = -cE_2,$$

$$\nabla_{E_2} E_2 = cE_1 + dE_3, \quad \nabla_{E_2} E_3 = -dE_2, \quad \nabla_{E_3} E_1 = 0, \quad \nabla_{E_3} E_2 = 0, \quad \nabla_{E_3} E_3 = 0,$$

and the Riemannian and Ricci curvature tensor fields

$$R(X, Y)Z := \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z,$$

$$\text{Ric}(Y, Z) := \sum_{k=1}^3 \tilde{g}(R(E_k, Y)Z, E_k)$$

are:

$$R(E_1, E_2)E_2 = [E_1(c) + E_2(a) - a^2 - c^2 - bd]E_1 + [E_3(c) - cd]E_3,$$

$$R(E_2, E_1)E_1 = [E_1(c) + E_2(a) - a^2 - c^2 - bd]E_2 + [E_3(a) - ab]E_3,$$

$$R(E_1, E_3)E_3 = [E_3(b) - b^2]E_1,$$

$$R(E_2, E_3)E_3 = [E_3(d) - d^2]E_2,$$

$$R(E_3, E_1)E_1 = [E_3(a) - ab]E_2 + [E_3(b) - b^2]E_3,$$

$$R(E_3, E_2)E_2 = [E_3(c) - cd]E_1 + [E_3(d) - d^2]E_3,$$

$$\begin{aligned} R(E_1, E_2)E_3 &= \tilde{g}(R(E_2, E_1)E_1, E_3)E_1 - \tilde{g}(R(E_1, E_2)E_2, E_3)E_2 \\ &= [E_3(a) - ab]E_1 - [E_3(c) - cd]E_2, \end{aligned}$$

$$\begin{aligned} R(E_2, E_3)E_1 &= \tilde{g}(R(E_3, E_2)E_2, E_1)E_2 - \tilde{g}(R(E_2, E_3)E_3, E_1)E_3 \\ &= [E_3(c) - cd]E_2, \end{aligned}$$

$$\begin{aligned} R(E_3, E_1)E_2 &= -\tilde{g}(R(E_3, E_1)E_1, E_2)E_1 + \tilde{g}(R(E_1, E_3)E_3, E_2)E_3 \\ &= -[E_3(a) - ab]E_1, \end{aligned}$$

$$\begin{aligned}
 \text{Ric}(E_1, E_1) &= E_1(c) + E_2(a) + E_3(b) - a^2 - b^2 - c^2 - bd, \\
 \text{Ric}(E_2, E_2) &= E_1(c) + E_2(a) + E_3(d) - a^2 - c^2 - d^2 - bd, \\
 \text{Ric}(E_3, E_3) &= E_3(b) + E_3(d) - b^2 - d^2, \\
 \text{Ric}(E_1, E_2) &= 0, \quad \text{Ric}(E_1, E_3) = E_3(c) - cd, \quad \text{Ric}(E_2, E_3) = E_3(a) - ab.
 \end{aligned}$$

Let $V = \sum_{k=1}^3 V^k E_k$ and $\eta = \sum_{k=1}^3 \eta^k e_k$, where e_k is the \tilde{g} -dual 1-form of E_k for $k \in \{1, 2, 3\}$. From (1), we get the equations that define an almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$:

$$(2) \quad \frac{1}{2} \left\{ E_i(V^j) + E_j(V^i) + \sum_{k=1}^3 V^k [\tilde{g}(\nabla_{E_i} E_k, E_j) + \tilde{g}(E_i, \nabla_{E_j} E_k)] \right\} + \text{Ric}(E_i, E_j) + \lambda \delta^{ij} + \mu \eta^i \eta^j = 0$$

for any $(i, j) \in \{(1, 1), (2, 2), (3, 3), (1, 2), (1, 3), (2, 3)\}$, equivalent to:

$$(3) \quad \begin{cases} E_1(V^1) - aV^2 - bV^3 + \text{Ric}(E_1, E_1) + \lambda + \mu(\eta^1)^2 = 0 \\ E_2(V^2) - cV^1 - dV^3 + \text{Ric}(E_2, E_2) + \lambda + \mu(\eta^2)^2 = 0 \\ E_3(V^3) + \text{Ric}(E_3, E_3) + \lambda + \mu(\eta^3)^2 = 0 \\ \frac{1}{2}[E_1(V^2) + E_2(V^1) + aV^1 + cV^2] + \mu\eta^1\eta^2 = 0 \\ \frac{1}{2}[E_1(V^3) + E_3(V^1) + bV^1] + \text{Ric}(E_1, E_3) + \mu\eta^1\eta^3 = 0 \\ \frac{1}{2}[E_2(V^3) + E_3(V^2) + dV^2] + \text{Ric}(E_2, E_3) + \mu\eta^2\eta^3 = 0 \end{cases}$$

with:

$$\begin{aligned}
 \text{Ric}(E_1, E_1) &= \frac{f_1^2}{f_2} \cdot \frac{\partial^2 f_2}{\partial(x^1)^2} + \frac{f_2^2}{f_1} \cdot \frac{\partial^2 f_1}{\partial(x^2)^2} + \frac{1}{f_1} \cdot \frac{\partial^2 f_1}{\partial(x^3)^2} \\
 &\quad - 2\frac{f_1^2}{f_2} \cdot \left(\frac{\partial f_2}{\partial x^1}\right)^2 - 2\frac{f_2^2}{f_1} \cdot \left(\frac{\partial f_1}{\partial x^2}\right)^2 - 2\frac{1}{f_1} \cdot \left(\frac{\partial f_1}{\partial x^3}\right)^2 \\
 &\quad + \frac{f_1}{f_2} \cdot \frac{\partial f_1}{\partial x^1} \cdot \frac{\partial f_2}{\partial x^1} + \frac{f_2}{f_1} \cdot \frac{\partial f_1}{\partial x^2} \cdot \frac{\partial f_2}{\partial x^2} - \frac{1}{f_1 f_2} \cdot \frac{\partial f_1}{\partial x^3} \cdot \frac{\partial f_2}{\partial x^3}, \\
 \text{Ric}(E_2, E_2) &= \frac{f_1^2}{f_2} \cdot \frac{\partial^2 f_2}{\partial(x^1)^2} + \frac{f_2^2}{f_1} \cdot \frac{\partial^2 f_1}{\partial(x^2)^2} + \frac{1}{f_2} \cdot \frac{\partial^2 f_2}{\partial(x^3)^2} \\
 &\quad - 2\frac{f_1^2}{f_2} \cdot \left(\frac{\partial f_2}{\partial x^1}\right)^2 - 2\frac{f_2^2}{f_1} \cdot \left(\frac{\partial f_1}{\partial x^2}\right)^2 - 2\frac{1}{f_2} \cdot \left(\frac{\partial f_2}{\partial x^3}\right)^2 \\
 &\quad + \frac{f_1}{f_2} \cdot \frac{\partial f_1}{\partial x^1} \cdot \frac{\partial f_2}{\partial x^1} + \frac{f_2}{f_1} \cdot \frac{\partial f_1}{\partial x^2} \cdot \frac{\partial f_2}{\partial x^2} - \frac{1}{f_1 f_2} \cdot \frac{\partial f_1}{\partial x^3} \cdot \frac{\partial f_2}{\partial x^3}, \\
 \text{Ric}(E_3, E_3) &= \frac{1}{f_1} \cdot \frac{\partial^2 f_1}{\partial(x^3)^2} + \frac{1}{f_2} \cdot \frac{\partial^2 f_2}{\partial(x^3)^2} - 2\frac{1}{f_1^2} \cdot \left(\frac{\partial f_1}{\partial x^3}\right)^2 - 2\frac{1}{f_2^2} \cdot \left(\frac{\partial f_2}{\partial x^3}\right)^2,
 \end{aligned}$$

$$\begin{aligned}\operatorname{Ric}(E_1, E_3) &= \frac{f_1}{f_2} \cdot \frac{\partial^2 f_2}{\partial x^3 \partial x^1} + \frac{1}{f_2} \cdot \frac{\partial f_1}{\partial x^3} \cdot \frac{\partial f_2}{\partial x^1} - 2 \frac{f_1}{f_2^2} \cdot \frac{\partial f_2}{\partial x^1} \cdot \frac{\partial f_2}{\partial x^3}, \\ \operatorname{Ric}(E_2, E_3) &= \frac{f_2}{f_1} \cdot \frac{\partial^2 f_1}{\partial x^3 \partial x^2} + \frac{1}{f_1} \cdot \frac{\partial f_1}{\partial x^2} \cdot \frac{\partial f_2}{\partial x^3} - 2 \frac{f_2}{f_1^2} \cdot \frac{\partial f_1}{\partial x^2} \cdot \frac{\partial f_1}{\partial x^3}.\end{aligned}$$

We shall firstly determine some conditions which the two functions f_1 and f_2 must satisfy for the manifold to be flat or Ricci-flat.

THEOREM 2.1. *If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$, then $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold.*

PROOF. In this case, $a = b = c = d = 0$ and $R(E_i, E_j)E_k = 0$ for any $i, j, k \in \{1, 2, 3\}$. \square

THEOREM 2.2. *If $f_i : \mathbb{R}^2 \times I \rightarrow \mathbb{R}$, $f_i = f_i(x^3)$ for $i \in \{1, 2\}$ and $I \subset \mathbb{R}$ an open real interval, then the following assertions are equivalent:*

(1) f_1 and f_2 are constant or one of the functions is constant and the other one is of the form

$$\frac{c_1}{x^3 - c_2}$$

for $c_1 \in \mathbb{R} \setminus \{0\}$ and $c_2 \in \mathbb{R} \setminus I$;

(2) $(\mathbb{R}^2 \times I, \tilde{g}|_{\mathbb{R}^2 \times I})$ is a flat Riemannian manifold;

(3) $(\mathbb{R}^2 \times I, \tilde{g}|_{\mathbb{R}^2 \times I})$ is a Ricci-flat Riemannian manifold.

PROOF. In this case, $a = 0$, $b = \frac{f_1'}{f_1}$, $c = 0$, $d = \frac{f_2'}{f_2}$, and

$$R(E_1, E_2)E_2 = -bdE_1, \quad R(E_2, E_1)E_1 = -bdE_2,$$

$$R(E_1, E_3)E_3 = (b' - b^2)E_1, \quad R(E_3, E_1)E_1 = (b' - b^2)E_3,$$

$$R(E_2, E_3)E_3 = (d' - d^2)E_2, \quad R(E_3, E_2)E_2 = (d' - d^2)E_3,$$

$$\operatorname{Ric}(E_1, E_1) = b' - b^2 - bd, \quad \operatorname{Ric}(E_2, E_2) = d' - d^2 - bd, \quad \operatorname{Ric}(E_3, E_3) = b' + d' - b^2 - d^2,$$

$$\operatorname{Ric}(E_1, E_2) = \operatorname{Ric}(E_1, E_3) = \operatorname{Ric}(E_2, E_3) = 0.$$

Then, $R = 0$ if and only if

$$\begin{cases} bd = 0 \\ b' = b^2 \\ d' = d^2 \end{cases};$$

Ric = 0 if and only if

$$\begin{cases} b' - b^2 = bd = d' - d^2 \\ b' - b^2 = -(d' - d^2) \end{cases} ;$$

that is, $R = 0$ if and only if Ric = 0, i.e.,

$$\begin{cases} f_1' f_2' = 0 \\ \left(\frac{f_1'}{f_1}\right)' = \left(\frac{f_1'}{f_1}\right)^2 \\ \left(\frac{f_2'}{f_2}\right)' = \left(\frac{f_2'}{f_2}\right)^2 \end{cases} .$$

Therefore,

- $f_1' = 0$ and $f_2' = 0$, or
- $f_1' = 0$ and $-\frac{1}{\frac{f_2'(x^3)}{f_2(x^3)}} = x^3 + c_0$ (i.e., $f_2(x^3) = \frac{c_1}{x^3 - c_2}$ with $c_1 \in \mathbb{R} \setminus \{0\}$ and

$c_2 \in \mathbb{R} \setminus I$), or

- $-\frac{1}{\frac{f_1'(x^3)}{f_1(x^3)}} = x^3 + c_0$ (i.e., $f_1(x^3) = \frac{c_1}{x^3 - c_2}$ with $c_1 \in \mathbb{R} \setminus \{0\}$ and $c_2 \in \mathbb{R} \setminus I$) and

$f_2' = 0$. □

We immediately deduce that

COROLLARY 2.3. *If $f_1 = f_2 =: f(x^3)$, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^3, \tilde{g})$ is a Ricci-flat Riemannian manifold;
- (3) f is constant.

PROOF. (1) and (2) are equivalent to $(f')^2 = 0$, hence we get the conclusion. □

THEOREM 2.4. *If $f_1 : \mathbb{R}^3 \rightarrow \mathbb{R}$, $f_1 = f_1(x^1)$, $f_2 : \mathbb{R}^2 \times I \rightarrow \mathbb{R}$, $f_2 = f_2(x^3)$ for $I \subset \mathbb{R}$ an open real interval, then the following assertions are equivalent:*

- (1) f_2 is constant or it is of the form

$$\frac{c_1}{x^3 - c_2}$$

for $c_1 \in \mathbb{R} \setminus \{0\}$ and $c_2 \in \mathbb{R} \setminus I$;

- (2) $(\mathbb{R}^2 \times I, \tilde{g}|_{\mathbb{R}^2 \times I})$ is a flat Riemannian manifold;
- (3) $(\mathbb{R}^2 \times I, \tilde{g}|_{\mathbb{R}^2 \times I})$ is a Ricci-flat Riemannian manifold.

PROOF. In this case, $a = b = c = 0$, $d = \frac{f'_2}{f_2}$, and

$$R(E_1, E_2)E_2 = R(E_1, E_3)E_3 = R(E_2, E_1)E_1 = R(E_3, E_1)E_1 = 0,$$

$$R(E_2, E_3)E_3 = (d' - d^2)E_2, \quad R(E_3, E_2)E_2 = (d' - d^2)E_3, \quad \text{Ric}(E_2, E_2) = \text{Ric}(E_3, E_3) = d' - d^2,$$

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0.$$

Then, $R = 0$ if and only if

$$d' = d^2$$

that is, $\text{Ric} = 0$, i.e.,

$$\left(\frac{f'_2}{f_2}\right)' = \left(\frac{f'_2}{f_2}\right)^2.$$

Therefore,

- $f'_2 = 0$, or
- $-\frac{1}{\frac{f'_2(x^3)}{f_2(x^3)}} = x^3 + c_0$ (i.e., $f_2(x^3) = \frac{c_1}{x^3 - c_2}$ with $c_1 \in \mathbb{R} \setminus \{0\}$ and $c_2 \in \mathbb{R} \setminus I$). \square

THEOREM 2.5. *If $f_i = f_i(x^2)$ for $i \in \{1, 2\}$, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^3, \tilde{g})$ is a Ricci-flat Riemannian manifold;
- (3) f_1 and f_2 satisfy the equation

$$\frac{f'_1 f'_2}{f_1 f_2} + \frac{f''_1}{f_1} = 2\left(\frac{f'_1}{f_1}\right)^2.$$

PROOF. In this case, $a = f_2 \frac{f'_1}{f_1}$, $b = c = d = 0$, and

$$R(E_1, E_2)E_2 = [E_2(a) - a^2]E_1, \quad R(E_2, E_1)E_1 = [E_2(a) - a^2]E_2,$$

$$R(E_1, E_3)E_3 = R(E_3, E_1)E_1 = R(E_2, E_3)E_3 = R(E_3, E_2)E_2 = 0,$$

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_2(a) - a^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0.$$

Then, $R = 0$ if and only if

$$f_2 a' = a^2,$$

that is, $\text{Ric} = 0$, i.e.,

$$f_2 f_2' \frac{f_1'}{f_1} + f_2^2 \frac{f_1'' f_1 - (f_1')^2}{f_1^2} = f_2^2 \left(\frac{f_1'}{f_1} \right)^2,$$

which is equivalent to

$$\frac{f_1' f_2'}{f_1 f_2} + \frac{f_1''}{f_1} = 2 \left(\frac{f_1'}{f_1} \right)^2.$$

□

We immediately deduce that

COROLLARY 2.6. *If $f_1 = f_2 =: f(x^2)$, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^3, \tilde{g})$ is a Ricci-flat Riemannian manifold;
- (3) $f(x^2) = c_1 e^{c_2 x^2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$.

PROOF. (1) and (2) are equivalent to $f f'' - (f')^2 = 0$, that is, $\left(\frac{f'}{f} \right)' = 0$, with the solution $f(x^2) = c_1 e^{c_2 x^2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$, hence we get the conclusion. □

REMARK 2.7. We shall now closely look on the condition (3) from Theorem 2.5 satisfied by the two functions, more precisely,

$$\frac{f_1' f_2'}{f_1 f_2} + \frac{f_1''}{f_1} = 2 \left(\frac{f_1'}{f_1} \right)^2.$$

If $f_1' = 0$, then f_1 is a constant function. If $f_1' \neq 0$ (hence, if it is nowhere zero), then the previous relation is equivalent to

$$\frac{f_2'}{f_2} = \frac{2 \left(\frac{f_1'}{f_1} \right)^2 - \frac{f_1''}{f_1}}{\frac{f_1'}{f_1}}.$$

Let us notice that

$$\frac{f_2'}{f_2} = \frac{\left(\frac{f_1'}{f_1} \right)^2 - \left(\frac{f_1'}{f_1} \right)'}{\frac{f_1'}{f_1}},$$

which is equivalent to

$$\frac{f_2'}{f_2} = \frac{f_1'}{f_1} - \frac{\left(\frac{f_1'}{f_1} \right)'}{\frac{f_1'}{f_1}},$$

which, by integration, gives

$$\ln |f_2| = \ln |f_1| - \ln \left| \frac{f_1'}{f_1} \right| + c_0 = \ln \left(e^{c_0} \frac{f_1^2}{|f_1'|} \right),$$

where $c_0 \in \mathbb{R}$; therefore,

$$f_2(x^2) = k \frac{f_1^2(x^2)}{f_1'(x^2)}$$

with $k \in \mathbb{R} \setminus \{0\}$.

THEOREM 2.8. *If $f_1 = f_1(x^2)$ and $f_2 = f_2(x^1)$, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^3, \tilde{g})$ is a Ricci-flat Riemannian manifold;
- (3) f_1 and f_2 satisfy the equation

$$\frac{f_1'' f_1 - 2(f_1')^2}{f_1^4} = -\frac{f_2'' f_2 - 2(f_2')^2}{f_2^4} = \text{constant}.$$

PROOF. In this case, $a = f_2 \frac{f_1'}{f_1}$, $b = 0$, $c = f_1 \frac{f_2'}{f_2}$, $d = 0$, and

$$R(E_1, E_2)E_2 = [E_1(c) + E_2(a) - a^2 - c^2]E_1, \quad R(E_2, E_1)E_1 = [E_1(c) + E_2(a) - a^2 - c^2]E_2,$$

$$R(E_1, E_3)E_3 = R(E_3, E_1)E_1 = R(E_2, E_3)E_3 = R(E_3, E_2)E_2 = 0,$$

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_1(c) + E_2(a) - a^2 - c^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0.$$

Then, $R = 0$ if and only if

$$f_1 \frac{\partial c}{\partial x^1} + f_2 \frac{\partial a}{\partial x^2} = a^2 + c^2,$$

that is, $\text{Ric} = 0$, i.e.,

$$f_1^2 \frac{f_2'' f_2 - (f_2')^2}{f_2^2} + f_2^2 \frac{f_1'' f_1 - (f_1')^2}{f_1^2} = f_2^2 \left(\frac{f_1'}{f_1} \right)^2 + f_1^2 \left(\frac{f_2'}{f_2} \right)^2,$$

which is equivalent to

$$\frac{f_1'' f_1 - 2(f_1')^2}{f_1^4} = -\frac{f_2'' f_2 - 2(f_2')^2}{f_2^4}.$$

Since f_1 depends only on x^2 and f_2 depends only on x^1 , we deduce that the above ratio must be a constant. \square

REMARK 2.9. We shall now closely look on the condition (3) from Theorem 2.8 satisfied by the two functions, more precisely,

$$\frac{f''f - 2(f')^2}{f^4} = k \in \mathbb{R}.$$

Let us notice that

$$\frac{f''f - 2(f')^2}{f^4} = -\left(\frac{1}{f}\right)'' \frac{1}{f}.$$

Denoting by $h =: \frac{1}{f}$, we have

$$h''h = -k.$$

Let $r \in \mathbb{R}$ and let $J \subset \mathbb{R}$ be an open real interval such that $0 \notin J$ and $-2k \ln |y| + r > 0$ for any $y \in J$. Let F be an antiderivative on J of the function

$$y \mapsto \frac{1}{\sqrt{-2k \ln |y| + r}}.$$

Then $F'(y) > 0$ for any $y \in J$, therefore, F is strictly increasing on J , hence, it is invertible onto its image. Let $\varepsilon \in \{\pm 1\}$ and let $I := \varepsilon(F(J) - c_0)$, where $c_0 \in \mathbb{R}$. Then

$$h : I \rightarrow \mathbb{R}, \quad h(x) := F^{-1}(\varepsilon x + c_0)$$

satisfies $h''(x)h(x) = -k$ for any $x \in I$, and we get

$$f : I \rightarrow \mathbb{R}, \quad f(x) = \frac{1}{F^{-1}(\varepsilon x + c_0)}.$$

We shall further consider the cases when the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, \lambda, \mu)$ is $V = \frac{\partial}{\partial x^3}$ and respectively, when $\eta = dx^3$.

2.1 Almost η -Ricci solitons with $V = \frac{\partial}{\partial x^3}$

If $V^1 = V^2 = 0$ and $V^3 = 1$, then the system (3) becomes

$$(4) \quad \begin{cases} -b + \text{Ric}(E_1, E_1) + \lambda + \mu(\eta^1)^2 = 0 \\ -d + \text{Ric}(E_2, E_2) + \lambda + \mu(\eta^2)^2 = 0 \\ \text{Ric}(E_3, E_3) + \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = 0 \\ \text{Ric}(E_1, E_3) + \mu\eta^1\eta^3 = 0 \\ \text{Ric}(E_2, E_3) + \mu\eta^2\eta^3 = 0 \end{cases}.$$

LEMMA 2.10. Let $V = \frac{\partial}{\partial x^3}$ be the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, \lambda, \mu)$. If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$, then (4) becomes

$$(5) \quad \begin{cases} \lambda + \mu(\eta^1)^2 = 0 \\ \lambda + \mu(\eta^2)^2 = 0 \\ \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = 0 \\ \mu\eta^1\eta^3 = 0 \\ \mu\eta^2\eta^3 = 0 \end{cases}.$$

PROOF. We have $a = b = c = d = 0$ and $\text{Ric}(E_i, E_j) = 0$ for any $i, j \in \{1, 2, 3\}$ from Theorem 2.1, and the system (4) becomes (5). \square

PROPOSITION 2.11. Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$ and μ is nowhere zero, then $\eta = 0$ and V is a Killing vector field.

PROOF. Since μ is nowhere zero, we get $(\eta^1)^2 = (\eta^2)^2 = (\eta^3)^2$, hence, $\eta^i = 0$ for $i \in \{1, 2, 3\}$, and $\lambda = 0$ from (5), therefore, $\mathcal{L}_V \tilde{g} = 0$ by means of (1). \square

LEMMA 2.12. Let $V = \frac{\partial}{\partial x^3}$ be the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, \lambda, \mu)$. If $f_i = f_i(x^3)$ for $i \in \{1, 2\}$, then (4) becomes

$$(6) \quad \begin{cases} -b + b' - b^2 - bd + \lambda + \mu(\eta^1)^2 = 0 \\ -d + d' - d^2 - bd + \lambda + \mu(\eta^2)^2 = 0 \\ b' + d' - b^2 - d^2 + \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = 0 \\ \mu\eta^1\eta^3 = 0 \\ \mu\eta^2\eta^3 = 0 \end{cases}.$$

PROOF. We have $a = 0$, $b = \frac{f'_1}{f_1}$, $c = 0$, $d = \frac{f'_2}{f_2}$, and

$$\text{Ric}(E_1, E_1) = b' - b^2 - bd, \quad \text{Ric}(E_2, E_2) = d' - d^2 - bd, \quad \text{Ric}(E_3, E_3) = b' + d' - b^2 - d^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0$$

from Theorem 2.2, and the system (4) becomes (6). \square

PROPOSITION 2.13. Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_i = f_i(x^3)$ for $i \in \{1, 2\}$, $\eta^i = f$ for $i \in \{1, 2, 3\}$, and μ is nowhere zero, then $\eta = 0$.

Moreover,

(i) if one of the functions f_1 and f_2 is constant, then the other one is constant, too; in this case, $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold, $\lambda = 0$ and V is a Killing vector field;

(ii) if $\frac{f_1}{f_2}$ is constant, then f_i is of the form

$$c_i e^{c_0 e^{-x^3}}, i \in \{1, 2\}$$

for $c_0 \in \mathbb{R}$, $c_1, c_2 \in \mathbb{R} \setminus \{0\}$.

PROOF. In this case, since μ is nowhere zero, (6) is equivalent to

$$(7) \quad \begin{cases} b(d+1) = d^2 - d' \\ d(b+1) = b^2 - b' \\ \lambda = b^2 - b' + d^2 - d' \\ f^2 = 0 \end{cases},$$

which implies $\eta = 0$.

(i) f_1 is constant if and only if $b = 0$, and from (7) we deduce that $b = 0$ if and only if $d = 0$, i.e., if and only if f_2 is constant. In this case, $R = 0$, $\lambda = 0$ and $\mathcal{L}_V \tilde{g} = 0$.

(ii) If $\frac{f_1}{f_2}$ is constant, then $b = d$ and (7) becomes

$$(8) \quad \begin{cases} b' = -b \\ \lambda = 2(b^2 - b') \end{cases}.$$

If $b = d = 0$, then f_1 and f_2 are constant and $\lambda = 0$. If $b = d \neq 0$ (hence, if they are nowhere zero), then

$$(9) \quad \begin{cases} \frac{b'}{b} = -1 \\ \lambda = 2(b^2 - b') \end{cases},$$

from where we get

$$\frac{f'_i(x^3)}{f_i(x^3)} = b(x^3) = c_0 e^{-x^3}, \quad c_0 \in \mathbb{R} \setminus \{0\}, i \in \{1, 2\},$$

which, by integration, gives

$$f_i(x^3) = c_i e^{c_0 e^{-x^3}}, \quad c_i \in \mathbb{R} \setminus \{0\}, i \in \{1, 2\}. \quad \square$$

EXAMPLE 2.14. The warped product manifold $\mathbb{R} \times_f \mathbb{R}^2 := (\mathbb{R}^3, \tilde{g})$, where $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x^1) = c_1 e^{c_2 e^{-x^1}}$, $c_1, c_2 \in \mathbb{R} \setminus \{0\}$ and $\tilde{g} = dx^1 \otimes dx^1 + f^2(dx^2 \otimes dx^2 + dx^3 \otimes dx^3)$, is a non steady almost Ricci soliton with

$$V = \frac{\partial}{\partial x^1}, \quad \lambda(x^1) = 2c_2 e^{-x^1} (c_2 e^{-x^1} - 1).$$

COROLLARY 2.15. *Under the hypotheses of Proposition 2.13, if $f_1 = f_2 =: f(x^3)$, then $f(x^3) = c_1 e^{c_2 e^{-x^3}}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$.*

PROOF. It follows from Proposition 2.13 (ii). \square

LEMMA 2.16. *Let $V = \frac{\partial}{\partial x^3}$ be the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, \lambda, \mu)$. If $f_1 = f_1(x^1)$ and $f_2 = f_2(x^3)$, then (4) becomes*

$$(10) \quad \begin{cases} \lambda + \mu(\eta^1)^2 = 0 \\ -d + d' - d^2 + \lambda + \mu(\eta^2)^2 = 0 \\ d' - d^2 + \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = 0 \\ \mu\eta^1\eta^3 = 0 \\ \mu\eta^2\eta^3 = 0 \end{cases}.$$

PROOF. We have $a = b = c = 0$, $d = \frac{f_2'}{f_2}$, and

$$\text{Ric}(E_2, E_2) = \text{Ric}(E_3, E_3) = d' - d^2,$$

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0$$

from Theorem 2.4, and the system (4) becomes (10). \square

PROPOSITION 2.17. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_1 = f_1(x^1)$, $f_2 = f_2(x^3)$, $\eta^i =: f$ for $i \in \{1, 2, 3\}$, and μ is nowhere zero, then $\eta = 0$. Moreover, $\lambda = 0$, f_2 is constant and V is a Killing vector field.*

PROOF. In this case, since μ is nowhere zero, (10) is equivalent to

$$\begin{cases} d = d' - d^2 = 0 \\ \lambda = -\mu f^2 \\ f^2 = 0 \end{cases}.$$

It follows that $\eta = 0$, $\lambda = 0$ and $\mathcal{L}_V \tilde{g} = 0$. \square

LEMMA 2.18. Let $V = \frac{\partial}{\partial x^3}$ be the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, \lambda, \mu)$. If $f_i = f_i(x^2)$ for $i \in \{1, 2\}$, then (4) becomes

$$(11) \quad \begin{cases} f_2 f_2' \frac{f_1'}{f_1} + \frac{f_2^2}{f_1^2} [f_1'' f_1 - 2(f_1')^2] + \lambda + \mu(\eta^1)^2 = 0 \\ f_2 f_2' \frac{f_1'}{f_1} + \frac{f_2^2}{f_1^2} [f_1'' f_1 - 2(f_1')^2] + \lambda + \mu(\eta^2)^2 = 0 \\ \lambda + \mu(\eta^3)^2 = 0 \\ \mu \eta^1 \eta^2 = 0 \\ \mu \eta^1 \eta^3 = 0 \\ \mu \eta^2 \eta^3 = 0 \end{cases} .$$

PROOF. We have $a = f_2 \frac{f_1'}{f_1}$, $b = c = d = 0$, and

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_2(a) - a^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0$$

from Theorem 2.5, and the system (4) becomes (11). \square

PROPOSITION 2.19. Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_i = f_i(x^2)$ for $i \in \{1, 2\}$, $\eta^i =: f$ for $i \in \{1, 2, 3\}$, and μ is nowhere zero, then $\eta = 0$. Moreover, $\lambda = 0$ and

$$\frac{f_1' f_2'}{f_1 f_2} + \frac{f_1''}{f_1} = 2 \left(\frac{f_1'}{f_1} \right)^2 .$$

PROOF. In this case, since μ is nowhere zero, (11) is equivalent to

$$\begin{cases} f_2 f_2' \frac{f_1'}{f_1} + \frac{f_2^2}{f_1^2} [f_1'' f_1 - 2(f_1')^2] + \lambda + \mu f^2 = 0 \\ \lambda = -\mu f^2 \\ f^2 = 0 \end{cases} .$$

It follows that $\eta = 0$, $\lambda = 0$, and from the first equation of the previous system we get the relation between f_1 and f_2 . \square

COROLLARY 2.20. Under the hypotheses of Proposition 2.19, if $f_1 = f_2 =: f(x^2)$, then $f(x^2) = c_1 e^{c_2 x^2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$.

PROOF. It follows from Proposition 2.19 that $f f'' - (f')^2 = 0$, that is, $\left(\frac{f'}{f} \right)' = 0$, with the solution $f(x^2) = c_1 e^{c_2 x^2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$. \square

LEMMA 2.21. Let $V = \frac{\partial}{\partial x^3}$ be the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, \lambda, \mu)$. If $f_1 = f_1(x^2)$, $f_2 = f_2(x^1)$, then (4) becomes

$$(12) \quad \begin{cases} f_1 \frac{\partial c}{\partial x^1} + f_2 \frac{\partial a}{\partial x^2} - a^2 - c^2 + \lambda + \mu(\eta^1)^2 = 0 \\ f_1 \frac{\partial c}{\partial x^1} + f_2 \frac{\partial a}{\partial x^2} - a^2 - c^2 + \lambda + \mu(\eta^2)^2 = 0 \\ \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = 0 \\ \mu\eta^1\eta^3 = 0 \\ \mu\eta^2\eta^3 = 0 \end{cases} .$$

PROOF. We have $a = f_2 \frac{f_1'}{f_1}$, $b = 0$, $c = f_1 \frac{f_2'}{f_2}$, $d = 0$, and

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_1(c) + E_2(a) - a^2 - c^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0$$

from Theorem 2.8, and the system (4) becomes (12). \square

PROPOSITION 2.22. Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_1 = f_1(x^2)$, $f_2 = f_2(x^1)$, $\eta^i =: f$ for $i \in \{1, 2, 3\}$, and μ is nowhere zero, then $\eta = 0$. Moreover, $\lambda = 0$ and

$$\frac{f_1'' f_1 - 2(f_1')^2}{f_1^4} = -\frac{f_2'' f_2 - 2(f_2')^2}{f_2^4} = \text{constant}.$$

PROOF. In this case, since μ is nowhere zero, (12) is equivalent to

$$\begin{cases} f_1 \frac{\partial c}{\partial x^1} + f_2 \frac{\partial a}{\partial x^2} - a^2 - c^2 + \lambda + \mu f^2 = 0 \\ \lambda = -\mu f^2 \\ f^2 = 0 \end{cases} .$$

It follows that $\eta = 0$, $\lambda = 0$, and from the first equation of the previous system we get the relation between f_1 and f_2 . Since f_1 depends only on x^2 and f_2 depends only on x^1 , we deduce that the obtained ratio must be a constant. \square

2.2 Almost η -Ricci solitons with $\eta = dx^3$

If $\eta^1 = \eta^2 = 0$ and $\eta^3 = 1$, then the system (3) becomes

$$(13) \quad \begin{cases} E_1(V^1) - aV^2 - bV^3 + \text{Ric}(E_1, E_1) + \lambda = 0 \\ E_2(V^2) - cV^1 - dV^3 + \text{Ric}(E_2, E_2) + \lambda = 0 \\ E_3(V^3) + \text{Ric}(E_3, E_3) + \lambda + \mu = 0 \\ E_1(V^2) + E_2(V^1) + aV^1 + cV^2 = 0 \\ \frac{1}{2}[E_1(V^3) + E_3(V^1) + bV^1] + \text{Ric}(E_1, E_3) = 0 \\ \frac{1}{2}[E_2(V^3) + E_3(V^2) + dV^2] + \text{Ric}(E_2, E_3) = 0 \end{cases} .$$

LEMMA 2.23. *Let $\eta = dx^3$ be the 1-form of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$. If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$, then (13) becomes*

$$(14) \quad \begin{cases} \frac{\partial V^1}{\partial x^1} = -\frac{\lambda}{f_1} \\ \frac{\partial V^2}{\partial x^2} = -\frac{\lambda}{f_2} \\ \frac{\partial V^3}{\partial x^3} = -(\lambda + \mu) \\ f_1 \frac{\partial V^2}{\partial x^1} = -f_2 \frac{\partial V^1}{\partial x^2} \\ f_1 \frac{\partial V^3}{\partial x^1} = -\frac{\partial V^1}{\partial x^3} \\ f_2 \frac{\partial V^3}{\partial x^2} = -\frac{\partial V^2}{\partial x^3} \end{cases} .$$

PROOF. We have $a = b = c = d = 0$ and $\text{Ric}(E_i, E_j) = 0$ for any $i, j \in \{1, 2, 3\}$ from Theorem 2.1, and the system (13) becomes (14). \square

THEOREM 2.24. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an η -Ricci soliton with $\eta = dx^3$. If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$, then*

$$(15) \quad \begin{cases} V^1(x^1, x^2, x^3) = -\lambda F_1(x^1) + c_1 F_2(x^2) + c_2 x^3 + c_3 \\ V^2(x^1, x^2, x^3) = -c_1 F_1(x^1) - \lambda F_2(x^2) + c_4 x^3 + c_5 \\ V^3(x^1, x^2, x^3) = -(\lambda + \mu)x^3 - c_2 F_1(x^1) - c_4 F_2(x^2) + c_6 \end{cases} ,$$

where F_i is an antiderivative of $\frac{1}{f_i}$ for $i \in \{1, 2\}$ and $c_i \in \mathbb{R}$ for $i \in \{1, 2, 3, 4, 5, 6\}$.

PROOF. Since f_i depends only on x^i , from the first three equations of (14), we find that

$$\begin{aligned} V^1(x^1, x^2, x^3) &= -\lambda F_1(x^1) + h_1(x^2, x^3), \\ V^2(x^1, x^2, x^3) &= -\lambda F_2(x^2) + h_2(x^1, x^3), \\ V^3(x^1, x^2, x^3) &= -(\lambda + \mu)x^3 + h_3(x^1, x^2), \end{aligned}$$

where $F'_i = \frac{1}{f_i}$ for $i \in \{1, 2\}$. From the last three equations of (14) we infer

$$(16) \quad f_1(x^1) \frac{\partial h_2}{\partial x^1}(x^1, x^3) = -f_2(x^2) \frac{\partial h_1}{\partial x^2}(x^2, x^3),$$

$$(17) \quad f_1(x^1) \frac{\partial h_3}{\partial x^1}(x^1, x^2) = -\frac{\partial h_1}{\partial x^3}(x^2, x^3),$$

$$(18) \quad f_2(x^2) \frac{\partial h_3}{\partial x^2}(x^1, x^2) = -\frac{\partial h_2}{\partial x^3}(x^1, x^3).$$

Denoting $f_2(x^2) \frac{\partial h_1}{\partial x^2}(x^2, x^3) =: \bar{h}_1(x^3)$, we have

$$\frac{\partial h_1}{\partial x^2}(x^2, x^3) = \frac{\bar{h}_1(x^3)}{f_2(x^2)},$$

which, by integration, implies

$$h_1(x^2, x^3) = \bar{h}_1(x^3) F_2(x^2) + \hat{h}_1(x^3),$$

and by replacing it in (17), it gives

$$-\bar{h}'_1(x^3) F_2(x^2) - f_1(x^1) \frac{\partial h_3}{\partial x^1}(x^1, x^2) = \hat{h}'_1(x^3).$$

Differentiating the left term of the previous relation with respect to x^2 , it gives 0 and it implies

$$\bar{h}'_1(x^3) = -f_1(x^1) f_2(x^2) \frac{\partial^2 h_3}{\partial x^2 \partial x^1}(x^1, x^2),$$

which must be a constant, let's say, c_1 . Therefore,

$$\bar{h}_1(x^3) = c_1 x^3 + c_2$$

and

$$\frac{\partial^2 h_3}{\partial x^2 \partial x^1}(x^1, x^2) = -\frac{c_1}{f_1(x^1) f_2(x^2)},$$

which, by double integration, implies

$$h_3(x^1, x^2) = -c_1 F_1(x^1) F_2(x^2) + l_1(x^1) + l_2(x^2).$$

Also, from (16), we have $\frac{\partial h_2}{\partial x^1}(x^1, x^3) = -\frac{\bar{h}_1(x^3)}{f_1(x^1)}$, which gives

$$h_2(x^1, x^3) = -\bar{h}_1(x^3) F_1(x^1) + \hat{h}_2(x^3).$$

From (17), we find

$$\hat{h}'_1(x^3) = -f_1(x^1)l'_1(x^1),$$

which must be a constant, let's say, c_3 , which gives

$$\hat{h}_1(x^3) = c_3x^3 + c_4.$$

Since $f_1(x^1)l'_1(x^1) = -c_3$, we get

$$l_1(x^1) = -c_3F_1(x^1) + c_5.$$

From (18), we find

$$\hat{h}'_2(x^3) = 2c_1F_1(x^1) - f_2(x^2)l'_2(x^2),$$

which must be a constant, let's say, c_6 , which gives

$$\hat{h}_2(x^3) = c_6x^3 + c_7.$$

Since $2c_1F_1(x^1) = f_2(x^2)l'_2(x^2) + c_6$ and F_1 is not constant, we get

$$c_1 = 0, \quad l_2(x^2) = -c_6F_2(x^2) + c_8.$$

We have obtained

$$\begin{aligned} V^1(x^1, x^2, x^3) &= -\lambda F_1(x^1) + c_2F_2(x^2) + c_3x^3 + c_4, \\ V^2(x^1, x^2, x^3) &= -\lambda F_2(x^2) - c_2F_1(x^1) + c_6x^3 + c_7, \\ V^3(x^1, x^2, x^3) &= -(\lambda + \mu)x^3 - c_3F_1(x^1) + c_5 - c_6F_2(x^2) + c_8, \end{aligned}$$

which, by changing the indices of some constants, gives (15). \square

EXAMPLE 2.25. For $f_i = k_i \in \mathbb{R} \setminus \{0\}$, $i \in \{1, 2\}$, the vector field V with V^1 , V^2 and V^3 given by:

$$\begin{cases} V^1(x^1, x^2, x^3) &= -\frac{\lambda}{k_1}x^1 + \frac{c_1}{k_2}x^2 + c_2x^3 \\ V^2(x^1, x^2, x^3) &= -\frac{c_1}{k_1}x^1 - \frac{\lambda}{k_2}x^2 + c_3x^3 \\ V^3(x^1, x^2, x^3) &= -\frac{c_2}{k_1}x^1 - \frac{c_3}{k_2}x^2 - (\lambda + \mu)x^3 \end{cases},$$

where $c_i \in \mathbb{R}$ for $i \in \{1, 2, 3\}$, is the potential vector field of the η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, \lambda, \mu)$ for $\eta = dx^3$.

COROLLARY 2.26. *If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$, then*

$$\begin{cases} V^1(x^2, x^3) = c_1 F_2(x^2) + c_2 x^3 + c_3 \\ V^2(x^1, x^3) = -c_1 F_1(x^1) + c_4 x^3 + c_5 \\ V^3(x^1, x^2) = -c_2 F_1(x^1) - c_4 F_2(x^2) + c_6 \end{cases}$$

is a Killing vector field on $(\mathbb{R}^3, \tilde{g})$, where F_i is an antiderivative of $\frac{1}{f_i}$ for $i \in \{1, 2\}$ and $c_i \in \mathbb{R}$ for $i \in \{1, 2, 3, 4, 5, 6\}$.

PROOF. It follows from Lemma 2.23 (for $\lambda = \mu = 0$) and Theorem 2.24. \square

EXAMPLE 2.27. For $f_i = k_i \in \mathbb{R} \setminus \{0\}$, $i \in \{1, 2\}$, the vector field V with V^1 , V^2 and V^3 given by:

$$\begin{cases} V^1(x^2, x^3) = \frac{c_1}{k_2} x^2 + c_2 x^3 \\ V^2(x^1, x^3) = -\frac{c_1}{k_1} x^1 + c_3 x^3 \\ V^3(x^1, x^2) = -\frac{c_2}{k_1} x^1 - \frac{c_3}{k_2} x^2 \end{cases},$$

where $c_i \in \mathbb{R}$ for $i \in \{1, 2, 3\}$, is a Killing vector field.

LEMMA 2.28. *Let $\eta = dx^3$ be the 1-form of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$. If $f_i = f_i(x^3)$ for $i \in \{1, 2\}$, then (13) becomes*

$$(19) \quad \begin{cases} f_1 \frac{\partial V^1}{\partial x^1} - bV^3 + b' - b^2 - bd + \lambda = 0 \\ f_2 \frac{\partial V^2}{\partial x^2} - dV^3 + d' - d^2 - bd + \lambda = 0 \\ \frac{\partial V^3}{\partial x^3} + b' - b^2 + d' - d^2 + \lambda + \mu = 0 \\ f_1 \frac{\partial V^2}{\partial x^1} + f_2 \frac{\partial V^1}{\partial x^2} = 0 \\ f_1 \frac{\partial V^3}{\partial x^1} + \frac{\partial V^1}{\partial x^3} + bV^1 = 0 \\ f_2 \frac{\partial V^3}{\partial x^2} + \frac{\partial V^2}{\partial x^3} + dV^2 = 0 \end{cases}.$$

PROOF. We have $a = 0$, $b = \frac{f'_1}{f_1}$, $c = 0$, $d = \frac{f'_2}{f_2}$, and

$$\text{Ric}(E_1, E_1) = b' - b^2 - bd, \quad \text{Ric}(E_2, E_2) = d' - d^2 - bd, \quad \text{Ric}(E_3, E_3) = b' + d' - b^2 - d^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0$$

from Theorem 2.2, and the system (13) becomes (19). \square

THEOREM 2.29. Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $\eta = dx^3$. If $f_i = f_i(x^3)$ for $i \in \{1, 2\}$, and $V^i = V^i(x^3)$ for $i \in \{1, 2, 3\}$, we have the following cases.

(i) $\frac{f_1}{f_2}$ constant;

(1) if f_1 and f_2 are constant, then

$$\begin{cases} V^1 = c_1 \\ V^2 = c_2 \\ V^3 = F \end{cases},$$

$$\lambda = 0,$$

where $F' = -\mu$ and $c_1, c_2 \in \mathbb{R}$;

(2) if f_1 and f_2 are nonconstant, then, on any open real interval $I \subset \mathbb{R}$ where $f'_1 \neq 0$ and $f'_2 \neq 0$, we have:

$$\begin{cases} V^1 = \frac{c_1}{f_1} \\ V^2 = \frac{c_2}{f_2} \\ V^3 = \frac{b' - 2b^2 + \lambda}{b} \left(= \frac{d' - 2d^2 + \lambda}{d} \right) \end{cases},$$

$$\mu = -\left(\frac{b' + \lambda}{b}\right)' + 2b^2 - \lambda,$$

with $c_1, c_2 \in \mathbb{R} \setminus \{0\}$.

(ii) $\frac{f_1}{f_2}$ nonconstant;

(1) if f_1 is constant, then, on any open real interval $I \subset \mathbb{R}$ where $f'_2 \neq 0$, we have:

$$\begin{cases} V^1 = c_1 \\ V^2 = \frac{c_2}{f_2} \\ V^3 = \frac{d' - d^2}{d} \end{cases},$$

$$\lambda = 0, \quad \mu = -\frac{d''}{d} + \left(\frac{d'}{d}\right)^2 + d^2,$$

with $c_1 \in \mathbb{R}, c_2 \in \mathbb{R} \setminus \{0\}$;

(2) if f_2 is constant, then, on any open real interval $I \subset \mathbb{R}$ where $f'_1 \neq 0$, we have:

$$\begin{cases} V^1 = \frac{c_1}{f_1} \\ V^2 = c_2 \\ V^3 = \frac{b' - b^2}{b} \end{cases},$$

$$\lambda = 0, \quad \mu = -\frac{b''}{b} + \left(\frac{b'}{b}\right)^2 + b^2,$$

with $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$;

(3) if f_1 and f_2 are nonconstant, then, on any open real interval $I \subset \mathbb{R}$ where $f_1' \neq 0$, $f_2' \neq 0$ and $f_1 f_2' \neq f_1' f_2$, we have:

$$\begin{cases} V^1 = \frac{c_1}{f_1} \\ V^2 = \frac{c_2}{f_2} \\ V^3 = \frac{(b-d)'}{b-d} - (b+d) \end{cases},$$

$$\lambda = \frac{b'd - bd'}{b-d}, \quad \mu = -\frac{(b-d)'' + b'd - bd'}{b-d} + \left(\frac{(b-d)'}{b-d}\right)^2 + b^2 + d^2,$$

with $c_1, c_2 \in \mathbb{R} \setminus \{0\}$.

PROOF. (i) $\frac{f_1}{f_2}$ is constant if and only if $b = d$.

If f_1 and f_2 are constant, then $b = d = 0$ and (19) becomes

$$\begin{cases} \lambda = 0 \\ (V^1)' = 0 \\ (V^2)' = 0 \\ (V^3)' = -\mu \end{cases}.$$

If f_1 and f_2 are nonconstant, then $b = d \neq 0$ (on I) and (19) becomes

$$\begin{cases} (V^1)' = -bV^1 \\ (V^2)' = -bV^2 \\ (V^3)' = -2(b' - b^2) - (\lambda + \mu) \\ V^3 = \frac{b' - 2b^2 + \lambda}{b} \end{cases}.$$

From the first two equations we get either $V^i = 0$ or $\frac{(V^i)'}{V^i} = -\frac{f_i'}{f_i}$ for $i \in \{1, 2\}$, i.e., $V^i = \frac{c_i}{f_i}$ with $c_i \in \mathbb{R} \setminus \{0\}$, $i \in \{1, 2\}$. Differentiating V^3 from the last equation and using the third one, we get the expression of μ .

(ii) $\frac{f_1}{f_2}$ is nonconstant if and only if $b \neq d$.

If f_1 is constant, then $b = 0$ and since $d \neq 0$ (on I), (19) becomes

$$\begin{cases} \lambda = 0 \\ (V^1)' = 0 \\ (V^2)' = -dV^2 \\ (V^3)' = -(d' - d^2) - \mu \\ V^3 = \frac{d' - d^2}{d} \end{cases}.$$

Differentiating V^3 from the last equation and using the fourth one, we get the expression of μ .

Similarly we treat the case when f_2 is constant.

If f_1 and f_2 are nonconstant, then $b \neq 0$, $d \neq 0$ (on I) and (19) becomes

$$\left\{ \begin{array}{l} (V^1)' = -bV^1 \\ (V^2)' = -dV^2 \\ (V^3)' = -(b' - b^2 + d' - d^2) - (\lambda + \mu) \\ \frac{b' - b^2 - bd + \lambda}{b} = V^3 = \frac{d' - d^2 - bd + \lambda}{d} \end{array} \right. .$$

From the last relations we get the expression of λ , and differentiating V^3 and using the third equation, we obtain also μ . \square

EXAMPLE 2.30. For $f_i = k_i \in \mathbb{R} \setminus \{0\}$, $i \in \{1, 2\}$, the vector field

$$V = c_1 \frac{\partial}{\partial x^1} + c_2 \frac{\partial}{\partial x^2} + \left(c_3 - \frac{(x^3)^2}{2} \right) \frac{\partial}{\partial x^3}$$

is the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, 0, \mu(x^3) = x^3)$ for $\eta = dx^3$ and $c_1, c_2, c_3 \in \mathbb{R}$.

EXAMPLE 2.31. For $f_1 = k_1$, $f_2(x^3) = k_2 e^{kx^3}$ with $k, k_1, k_2 \in \mathbb{R} \setminus \{0\}$, the vector field

$$V = c_1 \frac{\partial}{\partial x^1} + c_2 \frac{\partial}{\partial x^2} - k \frac{\partial}{\partial x^3}$$

is the potential vector field of the η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, 0, k^2)$ for $\eta = dx^3$ and $c_1 \in \mathbb{R}$, $c_2 \in \mathbb{R} \setminus \{0\}$.

EXAMPLE 2.32. For $f_i(x^3) = k_i e^{kx^3}$, $i \in \{1, 2\}$, with $k, k_1, k_2 \in \mathbb{R} \setminus \{0\}$, the vector field

$$V = c_1 \frac{\partial}{\partial x^1} + c_2 \frac{\partial}{\partial x^2} + \left(\frac{\lambda}{k} - 2k \right) \frac{\partial}{\partial x^3}$$

is the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, \lambda, 2k^2 - \lambda - \frac{\lambda}{k})$ for $\eta = dx^3$ and $c_1, c_2 \in \mathbb{R} \setminus \{0\}$.

EXAMPLE 2.33. The Riemannian Sol_3 Lie group $(\mathbb{R}^3, \tilde{g})$, where $f_1(x^3) = e^{-x^3}$ and $f_2(x^3) = e^{x^3}$, is an η -Ricci soliton for $\eta = dx^3$,

$$V = c_1 \frac{\partial}{\partial x^1} + c_2 \frac{\partial}{\partial x^2}, \quad \lambda = 0, \quad \mu = 2$$

and $c_1, c_2 \in \mathbb{R} \setminus \{0\}$.

COROLLARY 2.34. *If $f_i = f_i(x^3)$ for $i \in \{1, 2\}$, then*

$$\begin{cases} V^1(x^3) &= \frac{c_1}{f_1(x^3)} \\ V^2(x^3) &= \frac{c_2}{f_2(x^3)} \\ V^3 &= 0 \end{cases}$$

is a Killing vector field on $(\mathbb{R}^3, \tilde{g})$, where $c_i \in \mathbb{R}$ for $i \in \{1, 2\}$.

PROOF. It follows from Lemma 2.28 (for $\lambda = \mu = 0$) and Theorem 2.29. \square

EXAMPLE 2.35. For $f_i(x^3) = k_i e^{-x^3}$, $k_i \in \mathbb{R} \setminus \{0\}$, $i \in \{1, 2\}$, the vector field V with V^1, V^2 and V^3 given by:

$$\begin{cases} V^1(x^3) = \frac{e^{x^3}}{k_1} \\ V^2(x^3) = \frac{e^{x^3}}{k_2} \\ V^3 = 0 \end{cases}$$

is a Killing vector field.

LEMMA 2.36. *Let $\eta = dx^3$ be the 1-form of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$. If $f_1 = f_1(x^1)$ and $f_2 = f_2(x^3)$, then (13) becomes*

$$(20) \quad \begin{cases} f_1 \frac{\partial V^1}{\partial x^1} + \lambda = 0 \\ f_2 \frac{\partial V^2}{\partial x^2} - dV^3 + d' - d^2 + \lambda = 0 \\ \frac{\partial V^3}{\partial x^3} + d' - d^2 + \lambda + \mu = 0 \\ f_1 \frac{\partial V^2}{\partial x^1} + f_2 \frac{\partial V^1}{\partial x^2} = 0 \\ f_1 \frac{\partial V^3}{\partial x^1} + \frac{\partial V^1}{\partial x^3} = 0 \\ f_2 \frac{\partial V^3}{\partial x^2} + \frac{\partial V^2}{\partial x^3} + dV^2 = 0 \end{cases}.$$

PROOF. We have $a = b = c = 0$, $d = \frac{f_2'}{f_2}$, and

$$\text{Ric}(E_2, E_2) = \text{Ric}(E_3, E_3) = d' - d^2,$$

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0$$

from Theorem 2.4, and the system (13) becomes (20). \square

THEOREM 2.37. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $\eta = dx^3$. If $f_1 = f_1(x^1)$, $f_2 = f_2(x^3)$, and $V^i = V^i(x^3)$ for $i \in \{1, 2, 3\}$, we have the following cases.*

(i) f_2 constant and

$$\begin{cases} V^1 = c_1 \\ V^2 = c_2 \\ V^3 = F \end{cases}, \\ \lambda = 0,$$

where $F' = -\mu$ and $c_1, c_2 \in \mathbb{R}$;

(ii) f_2 nonconstant and on any open real interval $I \subset \mathbb{R}$ where $f_2' \neq 0$, we have:

$$\begin{cases} V^1 = c_1 \\ V^2 = \frac{c_2}{f_2} \\ V^3 = \frac{d' - d^2}{d} \end{cases}, \\ \lambda = 0, \quad \mu = -\frac{d''}{d} + \left(\frac{d'}{d}\right)^2 + d^2,$$

with $c_1, c_2 \in \mathbb{R}$.

PROOF. (i) f_2 is constant if and only if $d = 0$, and (20) becomes

$$\begin{cases} \lambda = 0 \\ (V^1)' = 0 \\ (V^2)' = 0 \\ (V^3)' = -\mu \end{cases}.$$

(ii) f_2 is nonconstant if and only if $d \neq 0$ (on I), and (20) becomes

$$\begin{cases} \lambda = 0 \\ (V^1)' = 0 \\ (V^2)' = -dV^2 \\ (V^3)' = -(d' - d^2) - \mu \\ V^3 = \frac{d' - d^2}{d} \end{cases}.$$

Differentiating V^3 from the last equation and using the fourth one, we get μ . □

EXAMPLE 2.38. For $f_1(x^1) = e^{x^1}$, $f_2(x^3) = e^{x^3}$, the vector field

$$V = c_1 e^{x^1} \frac{\partial}{\partial x^1} + c_2 \frac{\partial}{\partial x^2} - \frac{\partial}{\partial x^3}$$

is the potential vector field of the η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, 0, 1)$ for $\eta = dx^3$ and $c_1 \in \mathbb{R}$, $c_2 \in \mathbb{R} \setminus \{0\}$.

LEMMA 2.39. *Let $\eta = dx^3$ be the 1-form of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$. If $f_i = f_i(x^2)$ for $i \in \{1, 2\}$, then (13) becomes*

$$(21) \quad \begin{cases} f_1 \frac{\partial V^1}{\partial x^1} - aV^2 + f_2 a' - a^2 + \lambda = 0 \\ f_2 \frac{\partial V^2}{\partial x^2} + f_2 a' - a^2 + \lambda = 0 \\ \frac{\partial V^3}{\partial x^3} + \lambda + \mu = 0 \\ f_1 \frac{\partial V^2}{\partial x^1} + f_2 \frac{\partial V^1}{\partial x^2} + aV^1 = 0 \\ f_1 \frac{\partial V^3}{\partial x^1} + \frac{\partial V^1}{\partial x^3} = 0 \\ f_2 \frac{\partial V^3}{\partial x^2} + \frac{\partial V^2}{\partial x^3} = 0 \end{cases} .$$

PROOF. We have $a = f_2 \frac{f_1'}{f_1}$, $b = c = d = 0$, and

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_2(a) - a^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0$$

from Theorem 2.5, and the system (13) becomes (21). \square

THEOREM 2.40. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $\eta = dx^3$. If $f_i = f_i(x^2)$ for $i \in \{1, 2\}$ and $V^i = V^i(x^3)$ for $i \in \{1, 2, 3\}$, we have the following cases.*

(i) f_1 constant and

$$\begin{cases} V^1 = c_1 \\ V^2 = c_2 \\ V^3 = F \\ \lambda = 0, \end{cases}$$

where $F' = -\mu$ and $c_1, c_2 \in \mathbb{R}$;

(ii) f_1 nonconstant and

$$\begin{cases} V^1 = 0 \\ V^2 = 0 \\ V^3 = F \end{cases} ,$$

$$\lambda = 2f_2^2 \left(\frac{f_1'}{f_1} \right)^2 - f_2 f_2' \frac{f_1'}{f_1} - f_2^2 \frac{f_1''}{f_1},$$

where $F' = -(\lambda + \mu)$.

PROOF. (i) f_1 is constant if and only if $a = 0$, and (21) becomes

$$\begin{cases} \lambda = 0 \\ (V^1)' = 0 \\ (V^2)' = 0 \\ (V^3)' = -\mu \end{cases}.$$

(ii) f_1 is nonconstant if and only if $a \neq 0$, and (21) becomes

$$\begin{cases} \lambda = -f_2 a' + a^2 \\ V^1 = 0 \\ V^2 = 0 \\ (V^3)' = -(\lambda + \mu) \end{cases}.$$

From the last equation we deduce that $\lambda + \mu$ depends only on x^3 , therefore,

$$0 = \frac{\partial(\lambda + \mu)}{\partial x^2} = \lambda' + \frac{\partial\mu}{\partial x^2},$$

which, by integration, gives

$$\mu(x^2, x^3) = -\lambda(x^2) - G(x^3),$$

and we will denote by F an antiderivative of G , i.e., $F' = G$. □

EXAMPLE 2.41. For $f_1(x^2) = f_2(x^2) = e^{x^2}$, the vector field

$$V = e^{x^3} \frac{\partial}{\partial x^3}$$

is the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, 0, \mu(x^3) = -e^{x^3})$ for $\eta = dx^3$.

LEMMA 2.42. *Let $\eta = dx^3$ be the 1-form of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$. If $f_1 = f_1(x^2)$, $f_2 = f_2(x^1)$, then (13) becomes*

$$(22) \quad \begin{cases} E_1(V^1) - aV^2 + E_1(c) + E_2(a) - a^2 - c^2 + \lambda = 0 \\ E_2(V^2) - cV^1 + E_1(c) + E_2(a) - a^2 - c^2 + \lambda = 0 \\ E_3(V^3) + \lambda + \mu = 0 \\ E_1(V^2) + E_2(V^1) + aV^1 + cV^2 = 0 \\ E_1(V^3) + E_3(V^1) = 0 \\ E_2(V^3) + E_3(V^2) = 0 \end{cases}.$$

PROOF. We have $a = f_2 \frac{f_1'}{f_1}$, $b = 0$, $c = f_1 \frac{f_2'}{f_2}$, $d = 0$, and

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_1(c) + E_2(a) - a^2 - c^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0$$

from Theorem (2.8), and the system (13) becomes (22). \square

THEOREM 2.43. Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $\eta = dx^3$. If $f_1 = f_1(x^2)$, $f_2 = f_2(x^1)$, and $V^i = V^i(x^3)$ for $i \in \{1, 2, 3\}$, then

$$\begin{cases} V^1 = c_1 \\ V^2 = c_2 \\ V^3 = F \end{cases},$$

$$\lambda = 0,$$

where $(c_1, c_2) \in \mathbb{R}^2 \setminus \{(0, 0)\}$ and $F' = -\mu$, or

$$\begin{cases} V^1 = 0 \\ V^2 = 0 \\ V^3 = F \end{cases},$$

$$\lambda = f_1^2 \left[\left(\frac{f_2'}{f_2} \right)' - \left(\frac{f_2'}{f_2} \right)^2 \right] + f_2^2 \left[\left(\frac{f_1'}{f_1} \right)' - \left(\frac{f_1'}{f_1} \right)^2 \right],$$

where $F' = -(\lambda + \mu)$.

PROOF. Since V^i for $i \in \{1, 2, 3\}$ depends only on x^3 , (22) becomes

$$\begin{cases} aV^2 = f_1 \frac{\partial c}{\partial x^1} + f_2 \frac{\partial a}{\partial x^2} - a^2 - c^2 + \lambda = cV^1 \\ aV^1 = -cV^2 \\ (V^1)' = 0 \\ (V^2)' = 0 \\ (V^3)' = -(\lambda + \mu) \end{cases}.$$

From the third and fourth equations we infer

$$V^1 = c_1, \quad V^2 = c_2,$$

for $c_1, c_2 \in \mathbb{R}$, and the previous system is equivalent to

$$\left\{ \begin{array}{l} V^1 = c_1 \\ V^2 = c_2 \\ \lambda = c_1 f_1 \frac{f_2'}{f_2} + f_1^2 \left[\left(\frac{f_2'}{f_2} \right)' - \left(\frac{f_2'}{f_2} \right)^2 \right] + f_2^2 \left[\left(\frac{f_1'}{f_1} \right)' - \left(\frac{f_1'}{f_1} \right)^2 \right] \\ c_2 a = c_1 c \\ c_1 a = -c_2 c \\ (V^3)' = -(\lambda + \mu) \end{array} \right. .$$

If $c_1 = 0$, then $c_2 = 0$ or $c_2 \neq 0$ and $a = c = 0$ (i.e., f_1 and f_2 are constant). In the first case,

$$\lambda = f_1^2 \left[\left(\frac{f_2'}{f_2} \right)' - \left(\frac{f_2'}{f_2} \right)^2 \right] + f_2^2 \left[\left(\frac{f_1'}{f_1} \right)' - \left(\frac{f_1'}{f_1} \right)^2 \right], \quad V^1 = 0, \quad V^2 = 0, \quad V^3 = F,$$

where $F' = -(\lambda + \mu)$, and in the second case,

$$\lambda = 0, \quad V^1 = 0, \quad V^2 = c_2, \quad V^3 = F,$$

where $c_2 \in \mathbb{R} \setminus \{0\}$ and $F' = -\mu$.

If $c_1 \neq 0$, then $(c_1^2 + c_2^2)a = 0$, hence $a = 0$, i.e., f_1 is constant. It follows that $c = 0$, i.e., f_2 is constant. In this case,

$$\lambda = 0, \quad V^1 = c_1, \quad V^2 = c_2, \quad V^3 = F,$$

where $c_1 \in \mathbb{R} \setminus \{0\}, c_2 \in \mathbb{R}$ and $F' = -\mu$.

Making the same discussion relative to c_2 , we get the conclusion. \square

EXAMPLE 2.44. For $f_1(x^2) = e^{x^2}$ and $f_2(x^1) = e^{x^1}$, the vector field

$$V = e^{x^2} \frac{\partial}{\partial x^1} + e^{x^1} \frac{\partial}{\partial x^2} + e^{x^3} \frac{\partial}{\partial x^3}$$

is the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, 0, \mu(x^3) = -e^{x^3})$ for $\eta = dx^3$.

2.3 Almost η -Ricci solitons with $\eta = dx^3$ and $V = \frac{\partial}{\partial x^3}$

If $V^1 = V^2 = \eta^1 = \eta^2 = 0$ and $V^3 = \eta^3 = 1$, then the system (3) becomes

$$(23) \quad \begin{cases} -b + \text{Ric}(E_1, E_1) + \lambda = 0 \\ -d + \text{Ric}(E_2, E_2) + \lambda = 0 \\ \text{Ric}(E_3, E_3) + \lambda + \mu = 0 \\ \text{Ric}(E_1, E_3) = 0 \\ \text{Ric}(E_2, E_3) = 0 \end{cases} .$$

THEOREM 2.45. *Let $V = \frac{\partial}{\partial x^3}$ and $\eta = dx^3$. If $f_i = f_i(x^3)$ for $i \in \{1, 2\}$, then $(\mathbb{R}^3, \tilde{g}, V)$ is an almost η -Ricci soliton with λ and μ as scalar functions if and only if they are given by*

$$\begin{aligned} \lambda &= \frac{f_1'}{f_1} \left(\frac{f_2'}{f_2} + 1 \right) - \frac{f_1''}{f_1} + 2 \left(\frac{f_1'}{f_1} \right)^2, \\ \mu &= -\frac{f_1'}{f_1} \left(\frac{f_2'}{f_2} + 1 \right) - \frac{f_2''}{f_2} + 2 \left(\frac{f_2'}{f_2} \right)^2, \end{aligned}$$

and f_1 and f_2 satisfy

$$(24) \quad \frac{f_1'' - f_1'}{f_1} - 2 \left(\frac{f_1'}{f_1} \right)^2 = \frac{f_2'' - f_2'}{f_2} - 2 \left(\frac{f_2'}{f_2} \right)^2 .$$

PROOF. In this case, we have $a = 0$, $b = \frac{f_1'}{f_1}$, $c = 0$, $d = \frac{f_2'}{f_2}$, and (23) becomes

$$(25) \quad \begin{cases} -b + \text{Ric}(E_1, E_1) + \lambda = 0 \\ -d + \text{Ric}(E_2, E_2) + \lambda = 0 \\ \text{Ric}(E_3, E_3) + \lambda + \mu = 0 \end{cases}$$

with:

$$\begin{aligned} \text{Ric}(E_1, E_1) &= \frac{f_1''}{f_1} - 2 \left(\frac{f_1'}{f_1} \right)^2 - \frac{f_1' f_2'}{f_1 f_2}, \\ \text{Ric}(E_2, E_2) &= \frac{f_2''}{f_2} - 2 \left(\frac{f_2'}{f_2} \right)^2 - \frac{f_1' f_2'}{f_1 f_2}, \\ \text{Ric}(E_3, E_3) &= \frac{f_1''}{f_1} - 2 \left(\frac{f_1'}{f_1} \right)^2 + \frac{f_2''}{f_2} - 2 \left(\frac{f_2'}{f_2} \right)^2. \end{aligned}$$

The two functions f_1 and f_2 must satisfy (24) due to the first two equations of (25), and we obtain the expressions of λ and μ from the first and the last equation of the same system. \square

EXAMPLE 2.46. $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an η -Ricci soliton with

$$V = \frac{\partial}{\partial x^3}, \quad \eta = dx^3, \quad \lambda = 2k^2 + k, \quad \mu = -k,$$

and $f_i(x^3) = k_i e^{kx^3}$, $i \in \{1, 2\}$, $k, k_1, k_2 \in \mathbb{R} \setminus \{0\}$. In this case, the scalar curvature is constant equal to $-6k^2$. Indeed,

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = \text{Ric}(E_3, E_3) = -2k^2.$$

COROLLARY 2.47. *Under the hypotheses of Theorem 2.45, if $f_1 = f_2 =: f(x^3)$, then $(\mathbb{R}^3, \tilde{g}, V)$ is an almost η -Ricci soliton with λ and μ as scalar functions if and only if they are given by*

$$\begin{aligned} \lambda &= 3\left(\frac{f'}{f}\right)^2 + \frac{f' - f''}{f}, \\ \mu &= \left(\frac{f'}{f}\right)^2 - \frac{f' + f''}{f}. \end{aligned}$$

PROOF. It follows immediately from Theorem 2.45. □

EXAMPLE 2.48. $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton with

$$V = \frac{\partial}{\partial x^3}, \quad \eta = dx^3,$$

$$\lambda(x^3) = 2\frac{(x^3)^3 + 5(x^3)^2 + kx^3 - k}{[(x^3)^2 + k]^2}, \quad \mu(x^3) = -2\frac{(x^3)^3 - (x^3)^2 + kx^3 + k}{[(x^3)^2 + k]^2},$$

and $f_1(x^3) = f_2(x^3) = (x^3)^2 + k$, $k \in \mathbb{R}_+ \setminus \{0\}$. In this case, the scalar curvature is upper bounded by $\frac{8}{k}$. Indeed,

$$\text{Ric}(E_1, E_1) = 2\frac{k - 5(x^3)^2}{[(x^3)^2 + k]^2} = \text{Ric}(E_2, E_2), \quad \text{Ric}(E_3, E_3) = 4\frac{k - 3(x^3)^2}{[(x^3)^2 + k]^2},$$

and

$$r = 8\frac{k - 4(x^3)^2}{[(x^3)^2 + k]^2} \leq \frac{8}{k}.$$

THEOREM 2.49. *Let $V = \frac{\partial}{\partial x^3}$ and $\eta = dx^3$. If $f_1 = f_1(x^1)$, $f_2 = f_2(x^3)$, then:*

(i) $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton if and only if

$$f_2 : \mathbb{R} \rightarrow \mathbb{R}, \quad f_2(x^3) = c_0 e^{-x^3}, \quad \lambda = 0, \quad \mu = 1$$

for $c_0 \in \mathbb{R} \setminus \{0\}$, or

$$f_2 : \mathbb{R} \rightarrow \mathbb{R}, \quad f_2(x^3) = \frac{c_1}{c_2 e^{x^3} + 1}, \quad \lambda = 0, \quad \mu(x^3) = \frac{c_2 e^{x^3}}{c_2 e^{x^3} + 1}$$

for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}_+ \setminus \{0\}$;

(ii) $(\mathbb{R}^2 \times I, \tilde{g}|_{\mathbb{R}^2 \times I}, V, \lambda, \mu)$ is an almost η -Ricci soliton if and only if

$$f_2 : I \rightarrow \mathbb{R}, \quad f_2(x^3) = \frac{c_1}{c_2 e^{x^3} - 1}, \quad \lambda = 0, \quad \mu(x^3) = \frac{c_2 e^{x^3}}{c_2 e^{x^3} - 1}$$

for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}_+ \setminus \{0\}$, for $I \subset \mathbb{R}$ an open real interval such that $-\ln(c_2) \notin I$.

PROOF. In this case, we have $a = b = c = 0$, $d = \frac{f'_2}{f_2}$, and (23) becomes

$$\begin{cases} \lambda = 0 \\ -d + d' - d^2 = 0 \\ d' - d^2 + \mu = 0 \end{cases} .$$

If $d = 0$, then f_2 is constant and $\lambda = \mu = 0$ (which contradicts $\mu \neq 0$). If $d = -1$, then $\ln |f_2(x^3)| = -x^3 + c_0$ with $c_0 \in \mathbb{R}$, and we get $f_2(x^3) = \pm e^{-x^3 + c_0}$ and $\mu = 1$. If $d \neq -1$ and $d \neq 0$ (hence, d is nowhere -1 neither 0), from the second equation of the above system we get

$$1 = \frac{d'}{d^2 + d} = \frac{d'}{d} - \frac{d'}{d + 1},$$

which, by integration, gives

$$\ln \left(\left| \frac{d(x^3)}{d(x^3) + 1} \right| \right) = x^3 + k_1$$

with $k_1 \in \mathbb{R}$, which infers

$$\frac{f'_2(x^3)}{f_2(x^3)} = d(x^3) = \frac{\pm e^{x^3 + k_1}}{1 \mp e^{x^3 + k_1}}$$

for all $x^3 \neq -k_1$. By integrating the previous relation, we get

$$f_2(x^3) = \frac{k_2}{e^{x^3 + k_1} \mp 1}$$

with $k_2 \in \mathbb{R} \setminus \{0\}$ and

$$\mu(x^3) = d^2(x^3) - d'(x^3) = -d(x^3) = \frac{e^{x^3 + k_1}}{e^{x^3 + k_1} \mp 1}. \quad \square$$

THEOREM 2.50. *Let $V = \frac{\partial}{\partial x^3}$ and $\eta = dx^3$. If $f_i = f_i(x^2)$ for $i \in \{1, 2\}$, then $(\mathbb{R}^3, \tilde{g}, V)$ is an almost η -Ricci soliton with λ and μ as scalar functions if and only if they are given by*

$$\begin{aligned}\lambda &= \frac{f_2}{f_1} \left[2 \frac{f_2}{f_1} (f_1')^2 - f_1' f_2' - f_1'' f_2 \right], \\ \mu &= -\frac{f_2}{f_1} \left[2 \frac{f_2}{f_1} (f_1')^2 - f_1' f_2' - f_1'' f_2 \right].\end{aligned}$$

PROOF. In this case, we have $a = f_2 \frac{f_1'}{f_1}$, $b = c = d = 0$, and (23) becomes

$$\begin{cases} \text{Ric}(E_1, E_1) + \lambda = 0 \\ \text{Ric}(E_2, E_2) + \lambda = 0 \\ \lambda + \mu = 0 \end{cases}$$

with:

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = f_2 f_2' \frac{f_1'}{f_1} + f_2^2 \frac{f_1''}{f_1} - 2f_2^2 \left(\frac{f_1'}{f_1} \right)^2. \quad \square$$

EXAMPLE 2.51. $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton with

$$V = \frac{\partial}{\partial x^3}, \quad \eta = dx^3, \quad \lambda(x^2) = 2k_2^2 e^{2x^2}, \quad \mu(x^2) = -2k_2^2 e^{2x^2},$$

and $f_1(x^2) = k_1 e^{-x^2}$, $f_2(x^2) = k_2 e^{x^2}$, $k_1, k_2 \in \mathbb{R} \setminus \{0\}$.

COROLLARY 2.52. *Under the hypotheses of Theorem 2.50, if $f_1 = f_2 =: f(x^2)$, then $(\mathbb{R}^3, \tilde{g}, V)$ is an almost η -Ricci soliton with λ and μ as scalar functions if and only if they are given by*

$$\begin{aligned}\lambda &= (f')^2 - f'' f, \\ \mu &= -(f')^2 + f'' f.\end{aligned}$$

PROOF. It follows immediately from Theorem 2.50. □

EXAMPLE 2.53. $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton with

$$V = \frac{\partial}{\partial x^3}, \quad \eta = dx^3, \quad \lambda(x^2) = 2[(x^2)^2 - k], \quad \mu(x^2) = -2[(x^2)^2 - k],$$

and $f_1(x^2) = f_2(x^2) = (x^2)^2 + k$, $k \in \mathbb{R} \setminus \{0\}$.

EXAMPLE 2.54. The manifold $\mathbb{H}^2 \times \mathbb{R} := \{(x^1, x^2, x^3) \in \mathbb{R}^3 : x^2 > 0\}$ with $\tilde{g} = \frac{1}{f^2} dx^1 \otimes dx^1 + \frac{1}{f^2} dx^2 \otimes dx^2 + dx^3 \otimes dx^3$ for $f(x^2) = x^2$ is an η -Ricci soliton with

$$V = \frac{\partial}{\partial x^3}, \quad \eta = dx^3, \quad \lambda = 1, \quad \mu = -1.$$

THEOREM 2.55. Let $V = \frac{\partial}{\partial x^3}$ and $\eta = dx^3$. If $f_1 = f_1(x^2)$ and $f_2 = f_2(x^1)$, then $(\mathbb{R}^3, \tilde{g}, V)$ is an almost η -Ricci soliton with λ and μ as scalar functions if and only if they are given by

$$\begin{aligned} \lambda &= 2 \left[\frac{f_2^2}{f_1^2} (f_1')^2 + \frac{f_1^2}{f_2^2} (f_2')^2 \right] - f_1^2 \frac{f_2''}{f_2} - f_2^2 \frac{f_1''}{f_1}, \\ \mu &= -2 \left[\frac{f_2^2}{f_1^2} (f_1')^2 + \frac{f_1^2}{f_2^2} (f_2')^2 \right] + f_1^2 \frac{f_2''}{f_2} + f_2^2 \frac{f_1''}{f_1}. \end{aligned}$$

PROOF. In this case, we have $a = f_2 \frac{f_1'}{f_1}$, $b = 0$, $c = f_1 \frac{f_2'}{f_2}$, $d = 0$, and (23) becomes

$$\begin{cases} \text{Ric}(E_1, E_1) + \lambda = 0 \\ \text{Ric}(E_2, E_2) + \lambda = 0 \\ \lambda + \mu = 0 \end{cases}$$

with:

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = f_2^2 \frac{f_1''}{f_1} + f_1^2 \frac{f_2''}{f_2} - 2 \left[f_2^2 \left(\frac{f_1'}{f_1} \right)^2 + f_1^2 \left(\frac{f_2'}{f_2} \right)^2 \right]. \quad \square$$

EXAMPLE 2.56. $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton with

$$V = \frac{\partial}{\partial x^3}, \quad \eta = dx^3, \quad \lambda(x^1, x^2) = e^{2x^1} + e^{2x^2}, \quad \mu(x^1, x^2) = -e^{2x^1} - e^{2x^2},$$

and $f_1(x^2) = e^{x^2}$, $f_2(x^1) = e^{x^1}$.

References

- [1] Atashpeykara, P., Haji-Badali, A. The algebraic Ricci solitons of Lie groups $\mathbb{H}^2 \times \mathbb{R}$ and Sol_3 . *Journal of Finsler Geometry and its Applications* **2020**, 1(2), 105–114. <https://doi.org/10.22098/jfga.2020.1244>
- [2] Belarbi, L. Ricci solitons of the $\mathbb{H}^2 \times \mathbb{R}$ Lie group. *Electronic Research Archive* **2020**, 28(1), 157–163. <https://doi.org/10.3934/era.2020010>

- [3] Belarbi, L. Ricci solitons of the Sol_3 Lie group. *Authorea*. July 26, 2020. <https://doi.org/10.22541/au.159576131.13724173>
- [4] Blaga, A.M. Almost η -Ricci solitons in $(LCS)_n$ -manifolds. *Bull. Belg. Math. Soc. Simon Stevin* **2018**, *25(5)*, 641–653. <https://doi.org/10.36045/bbms/1547780426>
- [5] Blaga, A.M. η -Ricci solitons on para-Kenmotsu manifolds. *Balkan J. Geom. Appl.* **2015**, *20(1)*, 1–13.
- [6] Blaga, A.M. Solutions of some types of soliton equations in \mathbb{R}^3 . *Filomat* **2019**, *33(4)*, 1159–1162. <https://doi.org/10.2298/FIL1904159B>
- [7] Cao, H.-D., Cui, X. Curvature estimates for four-dimensional gradient steady Ricci solitons. *J. Geom. Anal.* **2020**, *30(1)*, 511–525. <https://doi.org/10.1007/s12220-019-00152-z>
- [8] Cao, H.-D., Liu, T. Curvature estimates for 4-dimensional complete gradient expanding Ricci solitons. *Journal für die reine und angewandte Mathematik* **2022**, *790*, 115–135. <https://doi.org/10.1515/crelle-2022-0039>
- [9] Cao, H.-D., Zhu, M. Linear stability of compact shrinking Ricci solitons. *Math. Ann.* **2024**, 1–17. <https://doi.org/10.1007/s00208-024-02824-w>
- [10] Cho, J.T., Kimura, M. Ricci solitons and real hypersurfaces in a complex space form. *Tohoku Math. J.* **2019**, *61(2)*, 205–212. <https://doi.org/10.2748/tmj/1245849443>
- [11] Chow, B., Lu, P., Ni, L. *Hamilton's Ricci Flow*, Graduate Studies in Mathematics 77, AMS, Providence, RI, USA, 2006.
- [12] Hamilton, R.S. The Ricci flow on surfaces, Math. and general relativity (Santa Cruz, CA, 1986). *Contemp. Math.* **1988**, *71*, AMS, 237–262. <https://doi.org/10.1090/conm/071/954419>
- [13] Pigola, S., Rigoli, M., Rimoldi, M., Setti, A.G. Ricci almost solitons. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **2011**, *10(4)*, 757–799.

Department of Mathematics

West University of Timișoara

Faculty of Mathematics and Computer Science

Bld. V. Pârvan nr. 4, 300223, Timișoara, România

adarablaga@yahoo.com