arXiv:2406.12540v1 [math.CO] 18 Jun 2024

HERITABILITY OF KŐNIG'S PROPERTY FROM FINITE EDGE SETS

MARC KAUFMANN AND DOMINIC VAN DER ZYPEN

ABSTRACT. A hypergraph H = (V, E) is said to have K such that $|S \cap e| = 1$ for all $e \in M$, and S is a vertex cover of H. Aharoni posed the question whether K sing's Property is inheritable from finite subsets of E (Problem 6.7, [1]). We provide a negative answer and investigate similar questions for weaker properties.

1. INTRODUCTION

The study of min-max duality in graphs goes back at least to 1912 and the work of Frobenius on determinants of square matrices, a problem which Dénes Kőnig showed to be modelable by the search of perfect matchings in certain bipartite graphs (for an overview of the history cf. Plummer, [6]). In 1931, Kőnig proved the first milestone

Theorem 1.1. (in: [5]) Let G be a finite bipartite graph. Then there exists a vertex cover of its edges whose size is equal to a matching of maximal size.

The theorem extended his result for regular bipartite graphs, first communicated seventeen years prior. It was conjectured by Erdős that the same result should hold for infinite bipartite graphs as well and finally settled, in the affirmative, by Aharoni in 1984:

Theorem 1.2. (in: [2]) In any bipartite graph G, there exists a matching F and a cover C, such that C consists of the choice of precisely one vertex from each edge in F.

It is natural to ask when an analogous statement should hold for infinite *hyper* graphs. One approach requires that the result holds for all finite sub*hyper* graphs and transfer the property to the infinite hypergraph

²⁰¹⁰ Mathematics Subject Classification. 05C15, 05C83.

The first-named author gratefully acknowledges support by the Swiss National Science Foundation [grant number 200021 192079].

of interest. This suffices for graphs (cf. Proposition 3.2). Aharoni therefore asked if the same heritability holds for hypergraphs. In this article, we give a negative answer to this question.

2. Preliminaries

A hypergraph is a pair H = (V, E) where V is a set and $E \subseteq \mathcal{P}(V)$. The elements of E are called *edges*. For all hypergraphs in this paper we assume that all edges are non-empty (i.e., $\emptyset \notin E$).

A (vertex) cover of H is a set $C \subseteq V$ such that $C \cap e \neq \emptyset$ for all $e \in E$. The minimal cardinality that a cover can have is said to be the (vertex) covering number of H, and we denote it by $\nu(H)$.

A matching is a set $M \subseteq E$ of pairwise disjoint edges.

In the construction of examples, we will often use ω , the first infinite ordinal (which is equal to the set of non-negative integers).

Observation 2.1. If H = (V, E) is a hypergraph and $M \subseteq E$ is a matching, then $|M| \leq \nu(H)$.

Definition 2.2. A hypergraph H = (V, E) is said to have *Kőnig's Property* if there is a matching $M_0 \subseteq E$ and a cover $C_0 \subseteq V$ such that

$$|C_0 \cap e| = 1$$
 for all $e \in M_0$.

The following weakening of Kőnig's Property will be useful:

Definition 2.3. A hypergraph H = (V, E) has König's Weak Property if there is a matching $M \subseteq E$ with $|M| = \nu(H)$.

Observation 2.4. Kőnig's Property implies Kőnigs Weak Property.

Note that the contrapositive of 2.4 says that if for every matching $M \subseteq E$ we have $|M| < \nu(H)$ then H cannot have Kőnig's Property. We will use this elementary fact in the construction of the negative answer to Aharoni's question asked in [1].

The necessary condition of 2.4 is not sufficient, as the following example shows:

Example 2.5. Let G be K_{ω} , the complete graph on ω . Clearly, $\nu(G) = \aleph_0$, and

$$\left\{\{2n, 2n+1\}: n \in \omega\right\}$$

is a matching of cardinality \aleph_0 . But G does not have Kőnig's Property because every vertex cover of G either equals ω or $\omega \setminus \{k\}$ for some

 $k \in \omega$, and a cover of this form can never pick exactly 1 point of any given matching of $G = K_{\omega}$.

However, for finite graphs the necessary condition of Observation 2.4 is also sufficient:

Proposition 2.6. If G = (V, E) is a finite graph with a matching $M \subseteq E$ and $|M| = \nu(G)$, then G satisfies Kőnig's Property.

Proof. Take any maximal matching $M \subseteq E$, a minimal vertex cover $C \subseteq V$ and partition M with respect to C:

$$M = (\bigcup_{e \in M, |e \cap C| = 1} e) \sqcup (\bigcup_{e \in M, |e \cap C| > 1} e) \sqcup (\bigcup_{e \in M, e \cap C = \emptyset} e)$$

(By \sqcup we denote disjoint union.)

We now prove that only the first union may be non-empty:

• C intersects every edge $f \in E$, in particular all $e \in M \subset E$, so

$$\bigcup_{e \in M, e \cap C = \emptyset} e = \emptyset$$

• Consider $e \in M : |e \cap C| > 1$ and any of the k > 1 vertices $v \in e \cap C$. Either v is contained in another edge of M - contradicting the disjointness of elements of M - or $C - \{v\}$ yields a vertex cover of G. But |C| is finite by assumption, so $\nu(G) = |C| > |C - \{v\}|$, another contradiction. This yields

$$\bigcup_{e \in M, |e \cap C| > 1} e = \emptyset$$

We conclude

$$M = \bigcup_{e \in M, |e \cap C| = 1} e$$

Another concept related to Kőnig's Property is that of bipartiteness:

Definition 2.7. A hypergraph H = (V, E) is *bipartite* if there is $D \subseteq V$ such that both D and $V \setminus D$ intersect every $e \in E$ with |e| > 1.

It is easily seen that any hypergraph satisfying Kőnig's Property is bipartite.

3. Heritability from finite edge sets

In [1], Ron Aharoni asks the following question:

Problem 6.7. Suppose that every finite subhypergraph of H (i.e. H' = (V, E') where E' is a finite subset of E) satisfies Kőnig's Property. Does it necessarily follow that H satisfies Kőnig's Property?

The following gives a negative answer to this question:

Proposition 3.1. Let $E = \{A \subseteq \omega : (\omega \setminus A) \text{ is finite}\}$. Then

- (1) (ω, E_0) satisfies Kőnig's Property for all finite $E_0 \subseteq E$, but
- (2) (ω, E) does not satisfy Kőnig's Property.

Proof.

(1) Let $E_0 \subseteq E$ be finite. If $E_0 = \emptyset$, then the empty cover $C_0 = \emptyset$ and the empty matching $M_0 = \emptyset$ establish Kőnig's Property in a vacuous way. So we assume that $E_0 \neq \emptyset$. Since E_0 consists of co-finite subsets of ω , pick $x^* \in \bigcap E_0$. So $C := \{x^*\}$ is a cover of (ω, E_0) . Picking any member $e \in E_0$, we see that the singleton cover $C = \{x^*\}$ together with the singleton matching $M := \{e\}$ establish Kőnig's Property.

(2) Since every two members of E intersect, we see that we have |M| = 1 for every matching $M \subseteq E$. On the other hand, (ω, E) has no finite cover. (If $F \subseteq \omega$ is finite, then F does not intersect $\omega \setminus F$, which is a member of E by definition.) So $\nu((\omega, E)) = \aleph_0$. So, (ω, E) does not have Kőnig's Weak Property, and therefore it does not have Kőnig's Property (see Observation 2.4).

So we can say that Kőnig's Property is not "finitely inheritable".

However, the situation is different if we restrict ourselves to graphs:

Proposition 3.2. Kőnig's property is finitely inheritable for graphs.

Proof. Let G = (V, E) be a simple, undirected graph. If every finite subgraph satisfies Kőnig's Property, then G has no odd cycles and so G itself is bipartite. Finally, every bipartite graph satisfies Kőnig's Property, see [2].

In the remainder of this article we investigate the properties that are weaker than Kőnig's Property for finite inheritability.

We will consider bipartiteness first.

 $\mathbf{5}$

Example 3.3. Let $[\omega]^{\omega}$ denote the collection of all infinite subsets of ω . We consider the hypergraph $H = (\omega, [\omega]^{\omega})$.

(1) for all finite sets $E_0 \subseteq [\omega]^{\omega}$ the hypergraph (ω, E_0) is bipartite: If $E_0 = \emptyset$, then there is a trivial bipartition of (ω, \emptyset) . Otherwise, let

$$D = \{\min(e) : e \in E_0\}.$$

It is easy to see that the finite set D intersects every member of E_0 , and so does $\omega \setminus D$, as it is co-finite.

(2) $(\omega, [\omega]^{\omega})$ is not bipartite: Take any $D \subseteq \omega$. If D is finite, then D does not intersect the edge $\omega \setminus D$. If D is infinite, then D itself is an edge, which gives us the problem that $(\omega \setminus D) \cap D = \emptyset$. So, no matter whether D is finite or infinite, either D or $\omega \setminus D$ has empty intersection with some member of $[\omega]^{\omega}$. Therefore $H = (\omega, [\omega]^{\omega})$ is not bipartite.

For finite edge size, the situation changes, and with Tychonoff's compactness theorem [3] we can make the following positive statement.

Proposition 3.4. Let H = (V, E) be a hypergraph with edges of finite size only, such that for all finite subsets $E_0 \subseteq E$ the hypergraph (V, E_0) is bipartite. Then H is bipartite.

Proof. Suppose H = (V, E) is a hypergraph such that all members of E are finite, and we assume that H is not bipartite. We will construct $E_0 \subseteq E$ finite such that (V, E_0) is not bipartite.

Let $\{0,1\}^V$ be the set of all maps $f: V \to \{0,1\}$. Then H not being bipartite is equivalent to saying that

- (*) for every map $f \in \{0,1\}^V$, there is $e \in E$ with |e| > 1 and f is constant on e.
- Let $E_{>1} = \{e \in E : |e| > 1\}$. To every $e \in E_{>1}$ we associate the set $C_e = \{f \in \{0, 1\}^V : f \text{ is constant on } e\}.$

Claim. For every $e \in E_{>1}$, the set C_e is open in the product topology of $\{0,1\}^V$ (where we endow the base space $\{0,1\}$ with the discrete topology). - Fix $e \in E_{>1}$, keeping in mind that e is finite. For $v \in V$ let $\operatorname{pr}_v : \{0,1\}^V \to \{0,1\}$ be the projection map sending $f \in \{0,1\}^V$ to f(v). Then

$$C_e = \left(\bigcap_{v \in V} \operatorname{pr}^{-1}(\{0\})\right) \cup \left(\bigcap_{v \in V} \operatorname{pr}^{-1}(\{1\})\right).$$

Since the intersections involved are finite, C_e is open.

Statement (\star) above amounts to saying that

$$\mathcal{U} := \left\{ C_e : e \in E_{>1} \right\}$$

is an (open) cover of $\{0,1\}^V$. Using Tychonoff's theorem, we get a finite subcover $\mathcal{U}_0 \subseteq \mathcal{U}$.

So there is $n \in \mathbb{N}$ and edges $e_1, \ldots, e_n \in E_{>1}$ such that

$$\mathcal{U}_0 = \{C_{e_1}, C_{e_2}, \dots, C_{e_n}\}.$$

Set $E_0 := \{e_1, e_2, \dots, e_n\}.$

Recall that every member of E_0 has more than 1 element by definition. Since \mathcal{U}_0 is an open cover of $\{0,1\}^V$, we conclude that for every $f \in \{0,1\}^V$ there is $e \in E_0$ such that f is constant on e. This is exactly statement (\star) above, and we conclude that (V, E_0) is not bipartite. \Box

Next, we turn to Kőnig's Weak Property.

The example used in 3.1 shows that for Kőnig's Weak Property is not finitely inheritable either. But again, things change if we restrict ourselves to edges of *finite size*.

Proposition 3.5. Let H = (V, E) be a hypergraph with edges of finite size only, such that for all finite subsets $E_0 \subseteq E$ the hypergraph (V, E_0) satisfies Kőnig's Weak Property. Then H satisfies Kőnig's Weak Property.

Proof. First, note that a straightforward application of Zorn's Lemma implies that there is a matching M that is maximal with respect to set inclusion \subseteq . Maximality implies that $\bigcup M$ is a vertex cover. So by Observation 2.1 and definition of $\nu(H)$ we have

$$|M| \le \nu(H) \le |M|.$$

Case 1. If M is infinite, then the inequality above directly implies $|M| = \nu(H)$ because all edges are finite, so we get Kőnig's Weak Property.

Case 2. If M is finite, then $\nu(H)$ is finite, and a compactness argument similar to the one used in Proposition 3.4 shows that there is a finite set $E_0 \subseteq E$ with $\nu((V, E_0)) = \nu(H)$. Since (V, E_0) is balanced by assumption, there is a matching $M_0 \subseteq E_0$ such that $|M_0| = \nu((V, E_0)) = \nu(H)$, which finishes the proof. \Box

4. Other notions of bipartiteness

There are several ways to define a bipartite hypergraph. (All the definitions agree on graphs.)

Definition 4.1. Let H = (V, E) be a hypergraph with |e| > 1 for all $e \in E$. We say that H has the *choosability property* (CP) if there is $C \subseteq V$ such that $|C \cap e| = 1$ for all $e \in E$.

Clearly, the choosability property implies bipartiteness.

For showing that (CP) is not finitely inheritable in general, we need to introduce the concept of an *almost disjoint family*.

Definition 4.2. Let $A, B \in [\omega]^{\omega}$ (the collection of infinite subsets of ω). We say A, B are almost disjoint if $A \cap B$ is finite. An almost disjoint family is a set $\mathcal{A} \subseteq [\omega]^{\omega}$ consisting of pairwise almost disjoint infinite subsets of ω .

A standard application shows that every maximal almost disjoint family is contained in a maximal almost disjoint ("MAD") family.

Example 4.3. Erdős and Shelah constructed a non-bipartite MAD family $E \subseteq [\omega]^{\omega}$, see [4], Theorem 1.1. Let $H = (\omega, E)$.

- (1) Since H is not bipartite, it does not have the choosability property.
- (2) Let $E_0 \subseteq E$ be finite. If $E_0 = \emptyset$, then (CP) is satisfied vacuously. So suppose $E_0 \neq \emptyset$. Let F be the set of members of ω that are an element of more than one member of E_0 . Since E_0 is finite, and the intersection of any two distinct members of E_0 is finite, we get that F is finite (possibly empty). Let $x_0 = 1 + \max(F)$. From every member $e \in E_0$, the set

$$e \setminus \underbrace{\{0, \dots, x_0\}}_{=x_0+1}$$

is non-empty because e is infinite. So for every $e \in E_0$, pick one member of that set. This establishes the choosability property.

Again, by a compactness argument, the choosability property is finitely inheritable. But the compactness argument is quite different from the one used in Proposition 3.4, so we will state the proof in its entirety.

Proposition 4.4. Let H = (V, E) be a hypergraph with edges of finite size only, such that for all finite subsets $E_0 \subseteq E$ the hypergraph (V, E_0) has (CP). Then H has (CP).

Proof. Consider the Cartesian product

$$K = \prod \{e : e \in E\}$$

of all edges, endowed with the product topology (where every individual edge is given the discrete topology). The set K may be understood as the set of simultaneous choices $v_e \in e$ for all edges e. For every pair $e \neq e'$ we define

$$U(e, e') = \{ f \in K : f(e) \in e' \}.$$

It is a standard exercise to verify that for all $e \neq e' \in E$ the set U(e, e') is open in the product topology.

Case 1. $\bigcup \{U(e, e') : e \neq e' \in E\} \neq K$. It immediately follows that H = (V, E) has the choosability property.

Case 2. $\bigcup \{U(e, e') : e \neq e' \in E\} = K$. Then the collection $\{U(e, e') : e \neq e' \in E\}$ forms an open cover of K. So we apply Tychonoff's compactness theorem and get a finite subcover. In that case, we obtain a finite $E_0 \subseteq E$ such that (V, E_0) does not have (CP), contradicting the assumption of the Proposition. \Box

References

- Ron Aharoni, Infinite Matching Theory, Discrete Mathematics 95 (1991), 5– 22.
- [3] Andrey Nikolayevich Tychonoff, Über einen Funktionenraum, Mathematische Annalen 111 (1935), 762–766.
- [4] Paul Erdős, Saharon Shelah, Separability properties of almost-disjoint families of sets, Isr. J. Math. 12 (1972), 207–214.
- [5] Dénes Kőnig, Graphs and matrices, FMat. Fiz. Lapok 38 (1931), 116–119.
- [6] Michael Plummer, Matching Theory a sampler: From Dénes König to the present, Discrete Mathematics 100 (1992), 177–219.

ETH ZÜRICH, CH-8092 ZÜRICH, SWITZERLAND

Email address: marc.kaufmann@inf.ethz.ch

BERN, SWITZERLAND

Email address: dominic.zypen@gmail.com