

Generalized mixed and primal hybrid methods with applications to plate bending *

Norbert Heuer[†]

Abstract

We present an extended framework for hybrid finite element approximations of self-adjoint, positive definite operators. It covers the cases of primal, mixed, and ultraweak formulations, both at the continuous and discrete levels, and gives rise to conforming discretizations. Our framework allows for flexible continuity restrictions across elements, and includes the extreme cases of conforming and discontinuous hybrid methods. We illustrate an application of the framework to the Kirchhoff–Love plate bending model and present three primal hybrid and two mixed hybrid methods, four of them with numerical examples. In particular, we present conforming frameworks for (in classical meaning) non-conforming elements of Morley, Zienkiewicz triangular, and Hellan–Herrmann–Johnson types.

AMS Subject Classification: 65N30, 35J35, 74G15, 74S05 74K20,

Key words: primal hybrid method, mixed hybrid method, Kirchhoff–Love model, biharmonic operator, Morley element, Zienkiewicz triangle element, Hellan–Herrmann–Johnson method

1 Introduction

We present a framework for generalized mixed and primal hybrid methods with flexible continuity requirements of dual and primal variables. Our framework provides a conforming setting for discrete schemes that are non-conforming in standard spaces. We prove well-posedness of the continuous and discrete formulations, and quasi-optimality of the schemes. In this paper, we restrict ourselves to general positive-definite self-adjoint differential operators without lower-order terms, and consider homogeneous Dirichlet boundary conditions. We illustrate an application of the framework to the Kirchhoff–Love plate bending model which, in the simplest case, reduces to the biharmonic problem.

Different strategies are known to reduce the required regularity of discrete spaces for mixed finite element methods both of the standard (dual) and primal types. Such a regularity reduction is particularly attractive for spaces whose conformity not only requires continuity of different orders but especially for those aiming at pointwise symmetry. Examples are stress tensors of linear elasticity and bending moments in plate bending, but also (though without symmetry) deflections in plate bending of Kirchhoff–Love type. Strategies include discontinuous Galerkin (DG), non-conforming, and hybrid methods, see, e.g., [22, 15, 3, 55, 32]. This has been and continues to be a very active field of research, and an up-to-date literature discussion goes beyond the scope of this paper. DG and non-conforming methods are usually defined and analyzed at a purely discrete level in finite-dimensional spaces. Hybrid methods can also be interpreted as being of the discontinuous Galerkin type, but their setting is usually closer to the underlying variational formulation of the problem. Primal hybrid methods ignore the conformity of approximations of

*Supported by ANID-Chile through FONDECYT project 1230013

[†]Facultad de Matemáticas, Pontificia Universidad Católica de Chile, Avenida Vicuña Mackenna 4860, Santiago, Chile, email: nheuer@uc.cl

primal variables across element interfaces, and compensate this by the introduction of Lagrange multipliers. Mixed hybrid settings do this for the dual variable. The idea of using non-conforming hybrid discretizations goes back to the solid mechanics community, see Fraeijs de Veubeke [37] and Pian & Tong, Pian [53, 52]. Thomas and Raviart & Thomas provided analyses of the dual and primal hybrid schemes [56, 57, 54]. See also the early book by Brezzi and Fortin [15], predecessor of [10]. Brezzi provided an abstract framework for the appearing saddle point problems and discretizations in his seminal paper [13] and, also together with Marini [14, 16], analyzed such schemes for the biharmonic problem. Of course, there is related analysis for saddle point discretizations by Babuška [4], and Ladyzhenskaya [48] is usually cited for the continuous analysis. Brezzi specifically motivated his formulation and conditions for the analysis of hybrid finite element schemes.

There is no clear separation between different types of methods with approximations of reduced regularity, cf. the discussions in [2, 9], see also [10, Remark 10.3.4]. In this paper, we consider a finite element discretization to be of hybrid type if there is a variational formulation that renders the discretization conforming. We propose and analyze a framework both for primal and mixed hybrid methods with general continuity (conformity) requirements so as to have a functional setting that covers the extreme cases of primal/mixed Galerkin methods and primal/mixed hybrid methods, and includes intermediate cases. We also consider the case of ultraweak formulations. In fact, our presentation and analysis is in the spirit of the discontinuous Petrov–Galerkin (DPG) method with optimal test functions, proposed by Demkowicz and Gopalakrishnan as a scheme that aims at “automatic” discrete inf-sup stability [27, 28]. The DPG method is typically based on ultraweak formulations. Their use has been proposed by Després and Cessenat [31, 20]. Switching to an ultraweak formulation means that all appearing derivatives are thrown onto the test side via integrations by parts, in this way generating trace terms. Bottasso *et. el.* [11] did this in a DG setting and suggested to replace trace terms with independent variables, as proposed in the unified DG setting of [3]. As the DPG setting is essentially a functional analytic framework, the use of ultraweak formulations requires a definition of trace variables in continuous spaces. This has led to a specific analysis of such formulations [18, 41], see also [30, Appendix A] for a first abstract setting. It is now clear that the treatment of traces in the ultraweak DPG setting is equivalent to the framework set up for hybrid methods, usually working with quotient spaces and “harmonic” extensions. We again refer to Brezzi [13] for abstract saddle point formulations, there motivated by the analysis of hybrid schemes, and to the more extensive treatise [15] for specific problems. Here, we systematically work with trace spaces and norms as in the DPG spirit, and provide abstract formulations for hybrid settings. Of course, the well-posedness of our variational formulations and their discretizations follows from the Brezzi theory by verifying the now standard criteria.

An overview of the remainder is as follows. In Section 2 we present the abstract framework of our variational formulations and their discretizations. We do this for general self-adjoint positive definite operators but, for ease of presentation, restrict ourselves to homogeneous Dirichlet conditions and discard lower-order terms. The primal variable is assumed to be scalar, though an extension of our framework to vector cases is straightforward. In Subsections 2.1 and 2.2 we prove the well-posedness of generalized primal hybrid and mixed hybrid formulations, respectively, and the quasi-optimality of their discretizations. For its relevance for the DPG method, we also study a general ultraweak formulation in Subsection 2.3. A direct discretization would be a Petrov–Galerkin scheme, as used for the DPG method with optimal test functions. We briefly illustrate the path of a minimum residual discretization which is analogous to a) a fully discrete DPG scheme as proposed in [43] and b) a variational stabilization proposed in [25]. Proofs of all the abstract results are given at the end, in Section 4. In Section 3 we illustrate an application of the abstract results to the Kirchhoff–Love plate bending problem, and present five different formulations with discretizations. We include the primal and mixed hybrid versions (in §3.2 and

§3.5, respectively) which are new formulations, to the best of our knowledge, but are equivalent to the ones proposed in [13, 16], though with different discretizations. In the remaining Sections 3.3, 3.4, and 3.6 we depart from the standard hybrid settings and consider formulations that do not impose continuity restrictions between elements of principal variables. To our knowledge, these formulations and discretizations are new. In particular, we do not assume convexity of the domain and appearing traces of bending moments include corner forces in a well-posed manner. A primal hybrid scheme with continuity at vertices is the subject of §3.3. It can be interpreted as a scheme with Morley-type element and Lagrangian multipliers, cf. [50]. A primal hybrid scheme with continuous approximation is studied in §3.4. It can be seen as a conforming extension of C^0 -interior penalty (C0IP) schemes, see, e.g., [36, 12], or the Zienkiewicz triangular element, cf. [8, 49], see also the hybrid high-order (HHO) method in [33] with continuous approximation and the C0-hybrid formulation in [21], the latter also presenting a mixed hybrid discretization. In §3.6 we study a mixed hybrid formulation and discretization that imposes normal-normal continuity of bending moments, and thus is a conforming framework for elements of the Hellan–Herrmann–Johnson (HHJ) type, cf. [44, 45, 47]. Let us stress the fact that we do not give conforming settings for Morley, C0IP or HHJ elements. We do propose variational formulations of the underlying model problem (or the abstract model problem in Section 2), and conforming discretizations. This means that norms and variables, in particular trace variables, are induced by the energy spaces rather than representing L_2 -Lagrange multipliers on interfaces. In comparison with the non-conforming schemes, we usually need more degrees of freedom to have conformity and discrete stability. On the other hand, we measure energy errors of physical variables, and do have conformity which is advantageous, e.g., when performing a posteriori error analysis. Let us also note that our abstract framework covers the cases of classical primal and mixed formulations, not explicitly studied here. In the case of the Kirchhoff–Love model, composite Hsieh–Clough–Tocher (HCT) elements can be used for the conforming approximation of primal variables, see [24, 23, 34], whereas conforming elements for bending moments and the resulting mixed method have been recently presented in [38]. We do use traces of HCT elements to approximate traces of H^2 -variables, and also use a reduced element from [38] to approximate bending moments and their traces. All our discretizations are of low order and aim at a low number of degrees of freedom, though we do not claim optimality. For all but the primal hybrid method (which is very close to the nodal-continuous primal hybrid method) we present numerical results that illustrate the convergence properties of the schemes for a smooth model solution, see Section 3.7. In this paper, we generally prove quasi-optimal error estimates in energy norms, and leave the proof of specific convergence orders open. We also do not elaborate on superconvergence results which can be observed in some cases. We restrict our analysis to homogeneous Dirichlet boundary conditions. Though, we stress the fact that the analysis can be extended to include any combination of different types of boundary conditions that make physical sense. This is due to the fact that all the variables and spaces stem from well-posed variational formulations: no non-physical terms or Lagrangian multipliers are used. For an illustration of the inclusion of boundary conditions we refer to [19] which deals with the particular case of plane elasticity. At an abstract level, our framework includes the analysis provided there.

2 Abstract framework

For a bounded Lipschitz domain $\Omega \subset \mathbb{R}^d$ ($d \in \mathbb{N}$) and $U \in \{\mathbb{R}, \mathbb{R}^d, \mathbb{R}^{d \times d}\}$ or $U = \mathbb{S} := \{w \in \mathbb{R}^{d \times d}; w = w^\top\}$ we consider an unbounded U -valued differential operator $A : \text{dom}(A) \subset L_2(\Omega) \rightarrow L_2(\Omega; U)$ with unbounded formal adjoint $A^* : \text{dom}(A^*) \subset L_2(\Omega; U) \rightarrow L_2(\Omega)$, and the Hilbert spaces with

(squared) norms

$$\begin{aligned} H(A) &:= \{v \in L_2(\Omega); Av \in L_2(\Omega; U)\}, & \|v\|_A^2 &:= \|v\|^2 + \|Av\|^2, \\ H(A^*) &:= \{w \in L_2(\Omega; U); A^*w \in L_2(\Omega)\}, & \|w\|_{A^*}^2 &:= \|w\|^2 + \|A^*w\|^2. \end{aligned}$$

Here, $\|\cdot\|$ is the generic $L_2(\Omega)$ -norm and, below, (\cdot, \cdot) denotes the generic $L_2(\Omega)$ duality. There is a corresponding trace operator

$$\gamma_{A,\Gamma} : \begin{cases} H(A) & \rightarrow H(A^*)^*, \\ v & \mapsto \langle \gamma_{A,\Gamma}(v), w \rangle_\Gamma := (Av, w) - (v, A^*w) \end{cases}$$

and space with vanishing traces

$$H_0(A) := \{v \in H(A); \gamma_{A,\Gamma}(v) = 0\}.$$

Given a symmetric, positive definite tensor-field $\mathcal{C} \in L_\infty(\Omega; U \times U)$ and $f \in L_2(\Omega)$, our model problem with homogeneous Dirichlet boundary condition reads

$$u \in H_0(A) : \quad A^* \mathcal{C} A u = f. \quad (1)$$

There are three canonical variational formulations, the Euler–Lagrange equation

$$u \in H_0(A) : \quad (\mathcal{C} A u, A \delta u) = (f, \delta u) \quad \forall \delta u \in H_0(A), \quad (2)$$

the primal mixed formulation

$$w \in L_2(\Omega; U), u \in H_0(A) : \quad (\mathcal{C}^{-1} w, \delta w) - (A u, \delta w) = 0, \quad \forall \delta w \in L_2(\Omega; U), \quad (3a)$$

$$-(w, A \delta u) = -(f, \delta u) \quad \forall \delta u \in H_0(A), \quad (3b)$$

and the (dual) mixed formulation

$$w \in H(A^*), u \in L_2(\Omega) : \quad (\mathcal{C}^{-1} w, \delta w) - (u, A^* \delta w) = 0, \quad \forall \delta w \in H(A^*), \quad (4a)$$

$$-(A^* w, \delta u) = -(f, \delta u) \quad \forall \delta u \in L_2(\Omega). \quad (4b)$$

Under the standard assumptions

$$\exists C_{\text{PF}} > 0 : \quad \|v\| \leq C_{\text{PF}} \|Av\| \quad \forall v \in H_0(A) \quad (\text{Poincaré–Friedrichs}), \quad (5a)$$

$$\exists c_{\text{is}} > 0 : \quad \sup_{w \in H(A^*), \|w\|_{A^*} = 1} (A^* w, v) \geq c_{\text{is}} \|v\| \quad \forall v \in L_2(\Omega) \quad (\text{inf-sup}), \quad (5b)$$

formulations (2), (3), (4) are well posed and equivalent. Of course, for the prescribed boundary condition these properties are equivalent.

Our aim is to provide well-posed formulations that require less regularity than $u \in H(A)$ and $w \in H(A^*)$. To this end we consider a regular mesh $\mathcal{T} = \{T\}$ of polyhedrals T covering Ω and introduce the product spaces with (squared) norms

$$\begin{aligned} H(A, \mathcal{T}) &:= \{v \in L_2(\Omega); Av|_T \in L_2(T; U) \quad \forall T \in \mathcal{T}\}, & \|v\|_{A, \mathcal{T}}^2 &:= \|v\|^2 + \|Av\|_{\mathcal{T}}^2, \\ H(A^*, \mathcal{T}) &:= \{w \in L_2(\Omega; U); A^* w|_T \in L_2(T) \quad \forall T \in \mathcal{T}\}, & \|w\|_{A^*, \mathcal{T}}^2 &:= \|w\|^2 + \|A^* w\|_{\mathcal{T}}^2. \end{aligned}$$

Here, $\|\cdot\|_{\mathcal{T}}^2 := (\cdot, \cdot)_{\mathcal{T}}$ where the latter indicates the $L_2(\mathcal{T})$ -duality. In the following, $A_{\mathcal{T}}$ and $A_{\mathcal{T}}^*$ denote the corresponding \mathcal{T} -piecewise differential operators, e.g., $\|Av\|_{\mathcal{T}} = \|A_{\mathcal{T}} v\|$ for any $v \in H(A, \mathcal{T})$. Here and in the following, we identify elements of product spaces with corresponding piecewise defined function. For instance, $(v_T)_{T \in \mathcal{T}} \in \prod_{T \in \mathcal{T}} H(A, T)$ is identified with $v \in L_2(\Omega)$ defined by $v|_T := v_T$, $T \in \mathcal{T}$.

We consider closed spaces $\tilde{H}(A, \mathcal{T}) \subset H(A, \mathcal{T})$ and $\tilde{H}(A^*, \mathcal{T}) \subset H(A^*, \mathcal{T})$ that are intermediate:

$$H_0(A) \subset \tilde{H}(A, \mathcal{T}) \subset H(A, \mathcal{T}), \quad H(A^*) \subset \tilde{H}(A^*, \mathcal{T}) \subset H(A^*, \mathcal{T}).$$

These spaces induce two trace operators with support on the skeleton $\mathcal{S} = \cup\{\partial T; T \in \mathcal{T}\}$,

$$\begin{aligned} \gamma_{A, \mathcal{S}} : & \begin{cases} H(A) & \rightarrow & \tilde{H}(A^*, \mathcal{T})^*, \\ v & \mapsto & \langle \gamma_{A, \mathcal{S}}(v), w \rangle_{\mathcal{S}} := (Av, w) - (v, A^*w)_{\mathcal{T}} \end{cases} , \\ \gamma_{A^*, \mathcal{S}} : & \begin{cases} H(A^*) & \rightarrow & \tilde{H}(A, \mathcal{T})^*, \\ w & \mapsto & \langle \gamma_{A^*, \mathcal{S}}(w), v \rangle_{\mathcal{S}} := (A^*w, v) - (w, Av)_{\mathcal{T}} \end{cases} , \end{aligned}$$

and corresponding trace spaces

$$H(A, \mathcal{S}) := \gamma_{A, \mathcal{S}}(H(A)), \quad H_0(A, \mathcal{S}) := \gamma_{A, \mathcal{S}}(H_0(A)), \quad H(A^*, \mathcal{S}) := \gamma_{A^*, \mathcal{S}}(H(A^*))$$

with norms

$$\begin{aligned} \|\phi\|_{A, \mathcal{S}} &:= \inf\{\|v\|_A; v \in H(A), \gamma_{A, \mathcal{S}}(v) = \phi\}, \\ \|\phi\|_{(A^*, \sim, \mathcal{T})^*} &:= \sup_{w \in \tilde{H}(A^*, \mathcal{T}), \|w\|_{A^*, \mathcal{T}}=1} \langle \phi, w \rangle_{\mathcal{S}} \quad (\phi \in H(A, \mathcal{S})), \\ \|\psi\|_{A^*, \mathcal{S}} &:= \inf\{\|w\|_{A^*}; w \in H(A^*), \gamma_{A^*, \mathcal{S}}(w) = \psi\}, \\ \|\psi\|_{(A, \sim, \mathcal{T})^*} &:= \sup_{v \in \tilde{H}(A, \mathcal{T}), \|v\|_{A, \mathcal{T}}=1} \langle \psi, v \rangle_{\mathcal{S}} \quad (\psi \in H(A^*, \mathcal{S})). \end{aligned}$$

Here, the dualities are defined as

$$\begin{aligned} \langle \phi, w \rangle_{\mathcal{S}} &:= \langle \gamma_{A, \mathcal{S}}(v), w \rangle_{\mathcal{S}} \quad (w \in \tilde{H}(A^*, \mathcal{T}), v \in \gamma_{A, \mathcal{S}}^{-1}(\phi)) \\ \text{and } \langle \psi, v \rangle_{\mathcal{S}} &:= \langle \gamma_{A^*, \mathcal{S}}(w), v \rangle_{\mathcal{S}} \quad (v \in \tilde{H}(A, \mathcal{T}), w \in \gamma_{A^*, \mathcal{S}}^{-1}(\psi)). \end{aligned}$$

Of course, in the case that $\tilde{H}(A, \mathcal{T}) = H(A)$, $\gamma_{A, \mathcal{S}} = \gamma_{A, \Gamma}$.

2.1 Generalized primal hybrid formulation

To reformulate (1), we define $w := \mathcal{C}Au$. Testing $A^*w = f$ with $\delta u \in \tilde{H}(A, \mathcal{T})$ and applying trace operator $\gamma_{A^*, \mathcal{S}}$, we find that

$$(f, \delta u) = (w, A\delta u)_{\mathcal{T}} + \langle \gamma_{A^*, \mathcal{S}}(w), \delta u \rangle_{\mathcal{S}}.$$

We introduce the independent variable $\psi := \gamma_{A^*, \mathcal{S}}(w)$ and relax the regularity of u in the weak form of $\mathcal{C}^{-1}w = Au$. This leads to a generalized primal-mixed formulation of (1): *Find* $w \in L_2(\Omega; U)$, $u \in \tilde{H}(A, \mathcal{T})$, and $\psi \in H(A^*, \mathcal{S})$ such that

$$(\mathcal{C}^{-1}w, \delta w) - (Au, \delta w)_{\mathcal{T}} - \langle \delta \psi, u \rangle_{\mathcal{S}} = 0 \quad \forall \delta w \in L_2(\Omega; U), \quad \forall \delta \psi \in H(A^*, \mathcal{S}), \quad (6a)$$

$$-(w, A\delta u)_{\mathcal{T}} - \langle \psi, \delta u \rangle_{\mathcal{S}} = -(f, \delta u) \quad \forall \delta u \in \tilde{H}(A, \mathcal{T}). \quad (6b)$$

Elimination of w yields the *generalized primal hybrid formulation*: *Find* $u \in \tilde{H}(A, \mathcal{T})$ and $\psi \in H(A^*, \mathcal{S})$ such that

$$(\mathcal{C}Au, A\delta u)_{\mathcal{T}} + \langle \psi, \delta u \rangle_{\mathcal{S}} = (f, \delta u) \quad \forall \delta u \in \tilde{H}(A, \mathcal{T}), \quad (7a)$$

$$\langle \delta \psi, u \rangle_{\mathcal{S}} = 0 \quad \forall \delta \psi \in H(A^*, \mathcal{S}). \quad (7b)$$

Theorem 1. Let $f \in L_2(\Omega)$ be given and assume that (5a) holds. Problem (7) is well posed. Its solution (u, ψ) satisfies

$$\|u\|_{A, \mathcal{T}} + \|\psi\|_{A^*, \mathcal{S}} \leq C \|f\|$$

with a constant C that is independent of f and \mathcal{T} . Furthermore, $u \in H_0(A)$, $\psi = \gamma_{A^*, \mathcal{S}}(\mathcal{C}Au)$, and u solves (1).

A proof of this theorem is given in §4.1.

For a discretization of (7) we select finite-dimensional subspaces $\tilde{H}_h(A, \mathcal{T}) \subset \tilde{H}(A, \mathcal{T})$, $H_h(A^*, \mathcal{S}) \subset H(A^*, \mathcal{S})$, and assume the existence of a Fortin operator $\mathcal{F} : \tilde{H}(A, \mathcal{T}) \rightarrow \tilde{H}_h(A, \mathcal{T})$,

$$\langle \delta\psi, v - \mathcal{F}v \rangle_{\mathcal{S}} = 0 \quad \forall \delta\psi \in H_h(A^*, \mathcal{S}), \quad \forall v \in \tilde{H}(A, \mathcal{T}), \quad (8a)$$

$$\exists C_F > 0 : \quad \|\mathcal{F}v\|_{A, \mathcal{T}} \leq C_F \|v\|_{A, \mathcal{T}} \quad \forall v \in \tilde{H}(A, \mathcal{T}), \quad (8b)$$

to conclude the well-posedness and quasi-optimal convergence of the *generalized primal hybrid method*: Find $u_h \in \tilde{H}_h(A, \mathcal{T})$ and $\psi_h \in H_h(A^*, \mathcal{S})$ such that

$$(\mathcal{C}Au_h, A\delta u)_{\mathcal{T}} + \langle \psi_h, \delta u \rangle_{\mathcal{S}} = (f, \delta u) \quad \forall \delta u \in \tilde{H}_h(A, \mathcal{T}), \quad (9a)$$

$$\langle \delta\psi, u_h \rangle_{\mathcal{S}} = 0 \quad \forall \delta\psi \in H_h(A^*, \mathcal{S}). \quad (9b)$$

Theorem 2. Under the conditions of Theorem 1, and assuming (8) with a constant C_F independent of \mathcal{T} and the discrete subspace, scheme (9) has a unique solution (u_h, ψ_h) . It satisfies

$$\|u - u_h\|_{A, \mathcal{T}} + \|\psi - \psi_h\|_{A^*, \mathcal{S}} \leq C (\|u - v\|_{A, \mathcal{T}} + \|\psi - \eta\|_{A^*, \mathcal{S}}) \quad \forall v \in \tilde{H}_h(A, \mathcal{T}), \quad \forall \eta \in H_h(A^*, \mathcal{S})$$

with a constant C that is independent of f , \mathcal{T} and the discrete subspaces.

A proof of this theorem is standard, cf. , e.g., [10].

2.2 Generalized mixed hybrid formulation

As before, we introduce $w := \mathcal{C}Au$. Testing this relation with $\delta w \in \tilde{H}(A^*, \mathcal{T})$ and applying trace operator $\gamma_{A, \mathcal{S}}$, we find that

$$(\mathcal{C}^{-1}w, \delta w) = (u, A^* \delta w)_{\mathcal{T}} + \langle \gamma_{A, \mathcal{S}}(u), \delta w \rangle_{\mathcal{S}}.$$

We introduce the independent trace variable $\phi := \gamma_{A, \mathcal{S}}(u)$ and relax the regularity of w in the weak form of relation $A^*w = f$. This leads to the *generalized mixed hybrid formulation* of (1): Find $w \in \tilde{H}(A^*, \mathcal{T})$, $u \in L_2(\Omega)$, and $\phi \in H_0(A, \mathcal{S})$ such that

$$(\mathcal{C}^{-1}w, \delta w) - (u, A^* \delta w)_{\mathcal{T}} - \langle \phi, \delta w \rangle_{\mathcal{S}} = 0 \quad \forall \delta w \in \tilde{H}(A^*, \mathcal{T}), \quad (10a)$$

$$-(A^*w, \delta u)_{\mathcal{T}} - \langle \delta\phi, w \rangle_{\mathcal{S}} = -(f, \delta u) \quad \forall \delta u \in L_2(\Omega), \quad \forall \delta\phi \in H_0(A, \mathcal{S}). \quad (10b)$$

Theorem 3. Let $f \in L_2(\Omega)$ be given and assume that relations (5) hold. Problem (10) is well posed. Its solution (w, u, ϕ) satisfies

$$\|w\|_{A^*, \mathcal{T}} + \|u\| + \|\phi\|_{A, \mathcal{S}} \leq C \|f\|$$

with a constant C that is independent of f and \mathcal{T} . Furthermore, $u \in H_0(A)$, $w = \mathcal{C}Au \in H(A^*)$, $\phi = \gamma_{A, \mathcal{S}}(u)$, and u solves (1).

A proof of this theorem is given in §4.2.

For a discretization of (10) we select finite-dimensional subspaces $\tilde{H}_h(A^*, \mathcal{T}) \subset \tilde{H}(A^*, \mathcal{T})$, $H_h(\mathcal{T}) \subset L_2(\Omega)$, $H_h(A, \mathcal{S}) \subset H_0(A, \mathcal{S})$, and assume that there are operators

$$\mathcal{F}_1: \tilde{H}(A^*, \mathcal{T}) \rightarrow \tilde{H}_h(A^*, \mathcal{T}) \cap H(A^*), \quad (11a)$$

$$\mathcal{F}_2: \tilde{H}(A^*, \mathcal{T}) \rightarrow \tilde{H}_h(A^*, \mathcal{T}) \quad (11b)$$

that satisfy

$$(A^*(w - \mathcal{F}_1 w), \delta u)_{\mathcal{T}} = 0 \quad \forall \delta u \in H_h(\mathcal{T}), \quad \forall w \in \tilde{H}(A^*, \mathcal{T}), \quad (11c)$$

$$\langle \delta \phi, w - \mathcal{F}_2 w \rangle_{\mathcal{S}} = 0 \quad \forall \delta \phi \in H_h(A, \mathcal{S}), \quad \forall w \in \tilde{H}(A^*, \mathcal{T}), \quad (11d)$$

$$\exists C_1 > 0: \quad \|\mathcal{F}_1 w\|_{A^*, \mathcal{T}} \leq C_1 \|w\|_{A^*, \mathcal{T}} \quad \forall w \in \tilde{H}(A^*, \mathcal{T}), \quad (11e)$$

$$\exists C_2 > 0: \quad \|\mathcal{F}_2 w\|_{A^*, \mathcal{T}} \leq C_2 \|w\|_{A^*, \mathcal{T}} \quad \forall w \in \tilde{H}(A^*, \mathcal{T}) \quad (11f)$$

to conclude the well-posedness and quasi-optimal convergence of the *generalized mixed hybrid method*: Find $w_h \in \tilde{H}_h(A^*, \mathcal{T})$, $u_h \in H_h(\mathcal{T})$, and $\phi_h \in H_h(A, \mathcal{S})$ such that

$$(\mathcal{C}^{-1} w_h, \delta w) - (u_h, A^* \delta w)_{\mathcal{T}} - \langle \phi_h, \delta w \rangle_{\mathcal{S}} = 0 \quad \forall \delta w \in \tilde{H}_h(A^*, \mathcal{T}), \quad (12a)$$

$$-(A^* w_h, \delta u)_{\mathcal{T}} - \langle \delta \phi, w_h \rangle_{\mathcal{S}} = -(f, \delta u) \quad \forall \delta u \in H_h(\mathcal{T}), \quad \forall \delta \phi \in H_h(A, \mathcal{S}). \quad (12b)$$

Properties (11) imply that a discrete inf-sup condition holds, as we establish now. This result is a discrete analogue of [18, Theorem 3.3] which considers the continuous inf-sup condition.

Proposition 4. *If operators $\mathcal{F}_1, \mathcal{F}_2$ with properties (11) exist, then any $\delta u \in H_h(\mathcal{T})$ and $\delta \phi \in H_h(A, \mathcal{S})$ are bounded as*

$$\|\delta u\| + \|\delta \phi\|_{A, \mathcal{S}} \leq (C_2 + \frac{C_1}{c_{\text{is}}}(C_2 + 1)) \sup_{w_h \in \tilde{H}_h(A^*, \mathcal{T}) \setminus \{0\}} \frac{(A^* w_h, \delta u)_{\mathcal{T}} + \langle \delta \phi, w_h \rangle_{\mathcal{S}}}{\|w_h\|_{A^*, \mathcal{T}}}. \quad (13)$$

A proof of this proposition is given in §4.3. Let us state the well-posedness and quasi-optimal convergence of the generalized mixed hybrid method.

Theorem 5. *Under the conditions of Theorem 3, and assuming the existence of operators $\mathcal{F}_1, \mathcal{F}_2$ that satisfy (11) with constants C_1, C_2 independent of \mathcal{T} and the discrete subspaces, scheme (12) has a unique solution (w_h, u_h, ϕ_h) . It satisfies*

$$\|w - w_h\|_{A^*, \mathcal{T}} + \|u - u_h\| + \|\phi - \phi_h\|_{A, \mathcal{S}} \leq C(\|w - z\|_{A^*, \mathcal{T}} + \|u - v\| + \|\phi - \eta\|_{A, \mathcal{S}})$$

for any $z \in \tilde{H}_h(A^*, \mathcal{T})$, $v \in H_h(\mathcal{T})$, and $\eta \in H_h(A, \mathcal{S})$ with a constant C that is independent of f, \mathcal{T} and the discrete subspaces.

The comment to the proof of Theorem 2 applies in this case as well, noting that the discrete inf-sup property is satisfied due to Proposition 4.

2.3 General ultraweak formulation

For completeness we also consider a general ultraweak formulation of (1) and show its well-posedness. We introduce $w := \mathcal{C}Au$ and combine trace relations (6b) and (10a) with the two independent trace variables $\phi := \gamma_{A, \mathcal{S}}(u)$, $\psi := \gamma_{A^*, \mathcal{S}}(w)$. This gives the *general ultraweak formulation*: Find $u \in L_2(\Omega)$, $w \in L_2(\Omega; U)$, $\phi \in H_0(A, \mathcal{S})$, and $\psi \in H(A^*, \mathcal{S})$ such that

$$(\mathcal{C}^{-1} w, \delta w) - (u, A^* \delta w)_{\mathcal{T}} - \langle \phi, \delta w \rangle_{\mathcal{S}} = 0 \quad \forall \delta w \in \tilde{H}(A^*, \mathcal{T}), \quad (14a)$$

$$-(w, A \delta u)_{\mathcal{T}} - \langle \psi, \delta u \rangle_{\mathcal{S}} = -(f, \delta u) \quad \forall \delta u \in \tilde{H}(A, \mathcal{T}). \quad (14b)$$

Of course, w can be eliminated but this requires to introduce a test space of higher regularity than we are considering here.

Theorem 6. *Let $f \in L_2(\Omega)$ be given and assume that relations (5) hold. Problem (14) is well posed. Its solution (w, u, ϕ) satisfies*

$$\|u\| + \|w\| + \|\phi\|_{A,S} + \|\psi\|_{A^*,S} \leq C\|f\|$$

with a constant C that is independent of f and \mathcal{T} . Furthermore, $u \in H_0(A)$, $w = CAu \in H(A^*)$, $\phi = \gamma_{A,S}(u)$, $\psi = \gamma_{A^*,S}(w)$, and u solves (1).

A proof of this theorem is given in §4.4.

System (14) is unsymmetric and any direct discretization would be a Petrov–Galerkin scheme, the prototype being the discontinuous Petrov–Galerkin (DPG) method with optimal test functions [29]. We aim at symmetric formulations and extend (14) to a symmetric mixed system by introducing the residual as independent variable. This is in fact the stabilization idea of Dahmen et al., see [25, 26], and equivalent to the DPG approach, see [29, (2.21)].

To formulate the extended mixed system we introduce the linear functional $L((\delta u, \delta w)) := -(f, \delta u)$ and abbreviate the sum of the bilinear forms of (14) by $b(\mathbf{u}, \delta \mathbf{v})$ with

$$\begin{aligned} \mathbf{u} &= (u, w, \phi, \psi) \in \mathcal{U}(\mathcal{T}) := L_2(\Omega) \times L_2(\Omega; U) \times H_0(A, S) \times H(A^*, S) \\ \text{and } \delta \mathbf{v} &= (\delta u, \delta w) \in \mathcal{V}(\mathcal{T}) := \tilde{H}(A, \mathcal{T}) \times \tilde{H}(A^*, \mathcal{T}). \end{aligned}$$

Let $\langle\langle \cdot, \cdot \rangle\rangle_{\mathcal{V}(\mathcal{T})}$ denote the inner product of $\mathcal{V}(\mathcal{T})$ with norm $\|\cdot\|_{\mathcal{V}(\mathcal{T})}$. For $\mathbf{u} \in \mathcal{U}(\mathcal{T})$, we introduce the Riesz representation $\mathbf{v} \in \mathcal{V}(\mathcal{T})$ of the (negative) residual of (14),

$$\langle\langle \mathbf{v}, \delta \mathbf{v} \rangle\rangle_{\mathcal{V}(\mathcal{T})} = L(\delta \mathbf{v}) - b(\mathbf{u}, \delta \mathbf{v}) \quad \forall \delta \mathbf{v} \in \mathcal{V}(\mathcal{T}) \quad \text{and} \quad \|\mathbf{v}\|_{\mathcal{V}(\mathcal{T})} = \|b(\mathbf{u}, \cdot) - L\|_{\mathcal{V}(\mathcal{T})^*}.$$

This leads to the (trivial) mixed representation of (14): *Find $\mathbf{v} \in \mathcal{V}(\mathcal{T})$ and $\mathbf{u} \in \mathcal{U}(\mathcal{T})$ such that*

$$\langle\langle \mathbf{v}, \delta \mathbf{v} \rangle\rangle_{\mathcal{V}(\mathcal{T})} + b(\mathbf{u}, \delta \mathbf{v}) = L(\delta \mathbf{v}) \quad \forall \delta \mathbf{v} \in \mathcal{V}(\mathcal{T}), \quad (15a)$$

$$b(\delta \mathbf{u}, \mathbf{v}) = 0 \quad \forall \delta \mathbf{u} \in \mathcal{U}(\mathcal{T}). \quad (15b)$$

System (14) is well posed by Theorem 6 and therefore, mixed formulation (15) is well posed as well.

An abstract discretization is formulated in the canonical way. We select finite-dimensional subspaces $\mathcal{V}_h(\mathcal{T}) \subset \mathcal{V}(\mathcal{T})$, $\mathcal{U}_h(\mathcal{T}) \subset \mathcal{U}(\mathcal{T})$, and assume the existence of a Fortin operator $\mathcal{F} : \mathcal{V}(\mathcal{T}) \rightarrow \mathcal{V}_h(\mathcal{T})$,

$$b(\delta \mathbf{u}, \mathbf{v} - \mathcal{F}\mathbf{v}) = 0 \quad \forall \delta \mathbf{u} \in \mathcal{U}_h(\mathcal{T}), \quad \forall \mathbf{v} \in \mathcal{V}(\mathcal{T}), \quad (16a)$$

$$\exists C_F > 0 : \quad \|\mathcal{F}\mathbf{v}\|_{\mathcal{V}(\mathcal{T})} \leq C_F \|\mathbf{v}\|_{\mathcal{V}(\mathcal{T})} \quad \forall \mathbf{v} \in \mathcal{V}(\mathcal{T}), \quad (16b)$$

to conclude the well-posedness and quasi-optimal convergence of the *mixed general ultraweak (DPG) method*: *Find $\mathbf{v}_h \in \mathcal{V}_h(\mathcal{T})$ and $\mathbf{u}_h \in \mathcal{U}_h(\mathcal{T})$ such that*

$$\langle\langle \mathbf{v}_h, \delta \mathbf{v} \rangle\rangle_{\mathcal{V}(\mathcal{T})} + b(\mathbf{u}_h, \delta \mathbf{v}) = L(\delta \mathbf{v}) \quad \forall \delta \mathbf{v} \in \mathcal{V}_h(\mathcal{T}), \quad (17a)$$

$$b(\delta \mathbf{u}, \mathbf{v}_h) = 0 \quad \forall \delta \mathbf{u} \in \mathcal{U}_h(\mathcal{T}). \quad (17b)$$

Theorem 7. *Under the conditions of Theorem 6, and assuming (16) with a constant C_F independent of \mathcal{T} and the discrete subspaces, scheme (17) has a unique solution $(\mathbf{v}_h, \mathbf{u}_h)$ with components $\mathbf{u}_h = (u_h, w_h, \phi_h, \psi_h)$. It satisfies*

$$\|\mathbf{u} - \mathbf{u}_h\|_{\mathcal{U}(\mathcal{T})}^2 = \|u - u_h\|^2 + \|w - w_h\|^2 + \|\phi - \phi_h\|_{A,S}^2 + \|\psi - \psi_h\|_{A^*,S}^2 \leq C\|\mathbf{u} - \mathbf{w}\|_{\mathcal{U}(\mathcal{T})}^2$$

for any $\mathbf{w} \in \mathcal{U}_h(\mathcal{T})$ with a constant $C > 0$ that is independent of f , \mathcal{T} , and the discrete subspaces.

Proof. By the well-posedness of (15) and the existence of a Fortin operator, scheme (17) is well posed. Equation (17a) means that \mathbf{v}_h is the Riesz representation of the residual $L - b(\mathbf{u}_h, \cdot) \in \mathcal{V}_h(\mathcal{T})^*$, and (17b) implies that

$$\mathbf{u}_h = \arg \min_{\mathbf{w} \in \mathcal{U}_h(\mathcal{T})} \|b(\mathbf{w}, \cdot) - L\|_{\mathcal{V}_h(\mathcal{T})^*},$$

leading to the claimed error estimate. For details we refer to [17, Propositions 2.2, 2.3], see also [43, Theorem 2.1]. \square

3 Applications to the Kirchhoff–Love model

The Kirchhoff–Love model of plate bending reads

$$\operatorname{div} \mathbf{div} \mathbf{M} = f, \quad \mathbf{M} = \mathcal{C} D^2 u \quad \text{in } \Omega, \quad (18a)$$

$$u = \partial_n u = 0 \quad \text{on } \Gamma := \partial\Omega. \quad (18b)$$

Here, $u: \Omega \rightarrow \mathbb{R}$ is the deflection of a plate with mid-surface $\Omega \subset \mathbb{R}^2$, $\mathbf{M}: \Omega \rightarrow \mathbb{S}$ is the tensor of bending moments, and $f: \Omega \rightarrow \mathbb{R}$ represents the external vertical load. Tensor \mathcal{C} is symmetric, positive definite and represents the rigidity of the plate. Expressions $\mathbf{div} \mathbf{M}$, $D^2 u$, and $\partial_n u$ denote the row-wise divergence of \mathbf{M} , the Hessian of u , and the exterior normal derivative of u , respectively. In principle Ω can be a bounded, simply connected Lipschitz domain. For ease of discrete analysis we assume that it is a Lipschitz polygon.

3.1 Notation of spaces, operators and mesh data

Given a subdomain ω of \mathbb{R}^2 or a line segment, we use the canonical Lebesgue and Sobolev spaces $L_2(\omega)$ and $H^s(\omega)$ ($0 < s \leq 2$) of scalar functions, and need the corresponding spaces $L_2(\omega; U)$, $H^s(\omega; U)$ of functions with values in $U \in \{\mathbb{R}, \mathbb{R}^2, \mathbb{S}\}$. The $L_2(\omega; U)$ duality and norm will be generically denoted by $(\cdot, \cdot)_\omega$ and $\|\cdot\|_\omega$. We use canonical Sobolev norms in $H^s(\omega; U)$, indicated by $\|\cdot\|_{s,\omega}$, cf. [1], except for $s = 2$ when $\|\cdot\|_{2,\omega}^2 := \|\cdot\|_\omega^2 + \|D^2 \cdot\|_\omega^2$ with Hessian D^2 . We need the space of bending moments

$$H(\operatorname{ddiv}, \omega; \mathbb{S}) := \{\mathbf{M} \in L_2(\omega; \mathbb{S}); \operatorname{div} \mathbf{div} \mathbf{M} \in L_2(\omega; \mathbb{R})\}$$

with (squared) norm $\|\cdot\|_{\operatorname{ddiv}, \omega}^2 := \|\cdot\|_\omega^2 + \|\operatorname{div} \mathbf{div} \cdot\|_\omega^2$. Furthermore, $H_0^1(\omega; U)$ and $H_0^2(\omega; U)$ are the respective subspaces with homogeneous traces on the boundary $\partial\omega$ (of course, including the normal derivative for H^2). We generally drop index ω when $\omega = \Omega$.

We consider a regular (family of) mesh(es) \mathcal{T} consisting of shape-regular triangles T covering Ω , $\cup_{T \in \mathcal{T}} \bar{T} = \bar{\Omega}$. The set of (open) edges of \mathcal{T} is denoted by \mathcal{E} , $\mathcal{E}(\Omega) := \{E \in \mathcal{E}; E \subset \Omega\}$, and $\mathcal{E}(\Gamma) := \{E \in \mathcal{E}; E \subset \Gamma\}$. The set of edges of $T \in \mathcal{T}$ is $\mathcal{E}(T)$. There are corresponding sets of vertices \mathcal{N} , vertices $\mathcal{N}(T)$ of elements $T \in \mathcal{T}$, vertices $\mathcal{N}(\Omega)$ interior to Ω , and vertices $\mathcal{N}(\Gamma)$ on Γ . We also need the exterior unit normal and tangential vectors, defined almost everywhere on ∂T , generically denoted by \mathbf{n} and \mathbf{t} , respectively. Occasionally we use barycentric coordinates to specify basis functions. For every element $T \in \mathcal{T}$, we generically denote its coordinates by λ_j ($j = 1, 2, 3$), corresponding to vertices x_j and opposite edges E_j , numbered modulo 3. That is, e.g., $\{\lambda_j, \lambda_{j+1}, \lambda_{j+2}\} = \{\lambda_1, \lambda_2, \lambda_3\}$ for any integer j .

Mesh \mathcal{T} induces product spaces denoted as before, but replacing ω with \mathcal{T} , e.g., $H^1(\mathcal{T}) := \prod_{T \in \mathcal{T}} H^1(T)$. We will identify elements of product spaces with piecewise defined functions on Ω , e.g., $v \in H^1(\Omega)$ is identified with $v \in H^1(\mathcal{T})$, and $v \in H^1(\mathcal{T})$ corresponds to an element $v \in L_2(\Omega)$. Furthermore, adding index \mathcal{T} to a differential operator means that it is considered to be a piecewise operator, e.g., $\operatorname{div}_{\mathcal{T}}$ is the \mathcal{T} -piecewise divergence operator. The $L_2(\mathcal{T})$ duality is denoted by $(\cdot, \cdot)_{\mathcal{T}}$.

Throughout this section we select, consistent with the notation just introduced,

$$U = \mathbb{S}, \quad A = D^2, \quad A^* = \operatorname{div} \mathbf{div}, \quad \|\cdot\|_{2,\mathcal{T}} = \|\cdot\|_{A,\mathcal{T}}, \quad \|\cdot\|_{\operatorname{ddiv},\mathcal{T}} = \|\cdot\|_{A^*,\mathcal{T}}$$

and refer to the fixed canonical H^2 -trace operator as

$$\gamma_2 := \gamma_{A,\mathcal{S}} : H(A) = H^2(\Omega) \rightarrow H(\operatorname{ddiv}, \mathcal{T}; \mathbb{S})^*. \quad (19)$$

Up to a sign change, it is the trace operator $\operatorname{tr}^{\operatorname{Ggrad}}$ from [40].

For discretizations we need the space $P^s(T)$ that consists of polynomials of degree less than or equal to $s \in \mathbb{N}_0$ on $T \in \mathcal{T}$. According to our notation, $P^s(\mathcal{T})$ is the piecewise polynomial space (without continuity requirement). We also need the space $P_{\operatorname{hom}}^s(T)$ of homogeneous polynomials of degree s . The $L_2(\Omega)$ -projection operator onto $P^s(\mathcal{T})$ is denoted as $\Pi_{\mathcal{T}}^s$. The space of continuous piecewise polynomials is $P^{s,c}(\mathcal{T}) := P^s(\mathcal{T}) \cap H^1(\Omega)$.

3.2 Primal hybrid formulation

The following formulation gives the variational framework for a discretization that Brezzi and Fortin call a primal hybrid method, cf. [15, IV.1.3]. We select

$$\tilde{H}(A, \mathcal{T}) := H(A, \mathcal{T}) = H^2(\mathcal{T})$$

and denote

$$\gamma_{\operatorname{dDiv}} := \gamma_{A^*,\mathcal{S}}, \quad H^{-3/2,-1/2}(\mathcal{S}) := H(A^*, \mathcal{S}) = \gamma_{\operatorname{dDiv}}(H(\operatorname{ddiv}, \Omega; \mathbb{S})), \quad \|\cdot\|_{-3/2,-1/2,\mathcal{S}} := \|\cdot\|_{A^*,\mathcal{S}}.$$

In this case, with $\tilde{H}(A, \mathcal{T}) = H(A, \mathcal{T})$, $\gamma_{A^*,\mathcal{S}} = \gamma_{\operatorname{dDiv}}$ is the canonical trace operator introduced in [40].

The generalized primal hybrid formulation (7) becomes the *primal hybrid formulation* of the Kirchhoff–Love model: *Find $u \in H^2(\mathcal{T})$ and $\boldsymbol{\eta} \in H^{-3/2,-1/2}(\mathcal{S})$ such that*

$$(CD^2u, D^2\delta u)_{\mathcal{T}} + \langle \boldsymbol{\eta}, \delta u \rangle_{\mathcal{S}} = (f, \delta u), \quad (20a)$$

$$\langle \delta \boldsymbol{\eta}, u \rangle_{\mathcal{S}} = 0 \quad (20b)$$

holds for any $\delta u \in H^2(\mathcal{T})$ and $\delta \boldsymbol{\eta} \in H^{-3/2,-1/2}(\mathcal{S})$.

Theorem 8. *Let $f \in L_2(\Omega)$ be given. Problem (20) is well posed. Its solution $(u, \boldsymbol{\eta})$ satisfies*

$$\|u\|_{2,\mathcal{T}} + \|\boldsymbol{\eta}\|_{-3/2,-1/2,\mathcal{S}} \leq C\|f\|$$

with a constant C that is independent of f and \mathcal{T} . Furthermore, $u \in H_0^2(\Omega)$, $\boldsymbol{\eta} = \gamma_{\operatorname{dDiv}}(CD^2u)$, and $u, \mathbf{M} := CD^2u$ solve (18).

Proof. We note that $H_0(A) = H_0^2(\Omega)$, cf. [40, Proof of Proposition 3.8(i)]. The Poincaré–Friedrichs inequality holds by [6, Lemma 3.3]. An application of Theorem 1 proves the statements. \square

Discretization

In order to discretize trace space $H^{-3/2,-1/2}(\mathcal{S})$ we need to represent trace operator $\gamma_{\operatorname{dDiv}}$ explicitly. Integration by parts shows that $u \in H^2(\mathcal{T})$ and sufficiently smooth tensor functions $\mathbf{M} \in H(\operatorname{ddiv}, \mathcal{T}; \mathbb{S})$ satisfy

$$\begin{aligned} (\operatorname{div} \mathbf{div} \mathbf{M}, u)_{\mathcal{T}} - (\mathbf{M}, D^2u)_{\mathcal{T}} &= \sum_{T \in \mathcal{T}} \langle \mathbf{n} \cdot \operatorname{div} \mathbf{M}, u \rangle_{\partial T} - \langle \mathbf{M} \mathbf{n}, \nabla u \rangle_{\partial T} \\ &= \sum_{T \in \mathcal{T}} \left(\langle \mathbf{n} \cdot \operatorname{div} \mathbf{M} + \partial_t(\mathbf{t} \cdot \mathbf{M} \mathbf{n}), u \rangle_{\partial T} - \sum_{x \in \mathcal{N}(T)} [\mathbf{t} \cdot \mathbf{M} \mathbf{n}]_{\partial T}(x) u(x) - \langle \mathbf{n} \cdot \mathbf{M} \mathbf{n}, \partial_n u \rangle_{\partial T} \right). \end{aligned} \quad (21)$$

Here, $[\mathbf{t} \cdot \mathbf{M}\mathbf{n}]_{\partial T}(x)$ denotes the jump at node $x \in \mathcal{N}(T)$ of the trace $\mathbf{t} \cdot \mathbf{M}\mathbf{n}|_{\partial T}$ from within T (in a certain orientation), and $\partial_t = \mathbf{t} \cdot \nabla$ is the tangential derivative on ∂T in positive orientation. For details and a Sobolev space setting we refer to [40, Section 3].

Relation (21) gives rise to trace operator γ_{dDiv} by selecting tensors without jumps $\mathbf{M} \in H(\text{ddiv}, \Omega; \mathbb{S})$. Trace $\gamma_{\text{dDiv}}(\mathbf{M})$ has the following components:

$$\mathbf{n} \cdot \text{div } \mathbf{M} + \partial_t(\mathbf{t} \cdot \mathbf{M}\mathbf{n})|_E, \quad E \in \mathcal{E} \quad (\text{effective shear-force(s)}), \quad (22a)$$

$$\mathbf{n} \cdot \mathbf{M}\mathbf{n}|_E, \quad E \in \mathcal{E} \quad (\text{normal-normal traces}), \quad (22b)$$

$$[\mathbf{t} \cdot \mathbf{M}\mathbf{n}]_{\partial T}(x), \quad x \in \mathcal{N}(T), \quad T \in \mathcal{T} \quad (\text{corner forces}) \quad (22c)$$

$$\text{subject to} \quad \sum_{T \in \mathcal{T}: x \in \mathcal{N}(T)} [\mathbf{t} \cdot \mathbf{M}\mathbf{n}]_{\partial T}(x) = 0, \quad x \in \mathcal{N}(\Omega). \quad (22d)$$

The lowest-order discretization consists of edge-piecewise constants for both the effective shear forces and normal-normal traces, and the corner forces are point values subject to the constraints at interior vertices, cf. [40, (6.5)]. We represent this space as $P^0(\mathcal{S}) \times P^0(\mathcal{S}) \times \mathbb{R}_{\text{constr}}^{3|\mathcal{T}|}$ with obvious meaning of $\mathbb{R}_{\text{constr}}^{3|\mathcal{T}|}$ and where

$$P^0(\mathcal{S}) := \{\boldsymbol{\eta}: \mathcal{S} \rightarrow \mathbb{R}; \boldsymbol{\eta}|_E \text{ is a univariate constant } \forall E \in \mathcal{E}\}.$$

Interpreting trace space $H^{-3/2, -1/2}(\mathcal{S})$ as a space with three components (effective shear forces, normal-normal traces, corner forces), our discretization spaces are

$$\begin{aligned} \tilde{H}_h(A, \mathcal{T}) &:= P^3(\mathcal{T}) \subset H^2(\mathcal{T}), \\ H_h(A^*, \mathcal{S}) &:= P^0(\mathcal{S}) \times P^0(\mathcal{S}) \times \mathbb{R}_{\text{constr}}^{3|\mathcal{T}|} \subset H^{-3/2, -1/2}(\mathcal{S}). \end{aligned}$$

They have dimensions $10|\mathcal{T}|$ and $2|\mathcal{E}| + 3|\mathcal{T}| - |\mathcal{N}(\Omega)|$, respectively.

The resulting primal hybrid method reads as follows. Find $u_h \in P^3(\mathcal{T})$ and $\boldsymbol{\eta}_h \in P^0(\mathcal{S}) \times P^0(\mathcal{S}) \times \mathbb{R}_{\text{constr}}^{3|\mathcal{T}|}$ such that

$$(CD^2 u_h, D^2 \delta u)_{\mathcal{T}} + \langle \boldsymbol{\eta}_h, \delta u \rangle_{\mathcal{S}} = (f, \delta u), \quad (23a)$$

$$\langle \boldsymbol{\delta}\boldsymbol{\eta}, u_h \rangle_{\mathcal{S}} = 0 \quad (23b)$$

holds for any $\delta u \in P^3(\mathcal{T})$ and $\boldsymbol{\delta}\boldsymbol{\eta} \in P^0(\mathcal{S}) \times P^0(\mathcal{S}) \times \mathbb{R}_{\text{constr}}^{3|\mathcal{T}|}$. It converges quasi-optimally.

Theorem 9. Let $f \in L_2(\Omega)$ be given. System (23) is well posed. Its solution $(u_h, \boldsymbol{\eta}_h)$ satisfies

$$\|u - u_h\|_{2, \mathcal{T}} + \|\boldsymbol{\eta} - \boldsymbol{\eta}_h\|_{-3/2, -1/2, \mathcal{S}} \leq C \left(\|u - v\|_{2, \mathcal{T}} + \|\boldsymbol{\eta} - \boldsymbol{\psi}\|_{-3/2, -1/2, \mathcal{S}} \right)$$

for any $v \in P^3(\mathcal{T})$ and $\boldsymbol{\psi} \in P^0(\mathcal{S}) \times P^0(\mathcal{S}) \times \mathbb{R}_{\text{constr}}^{3|\mathcal{T}|}$. Here, $(u, \boldsymbol{\eta})$ is the solution of (20) and $C > 0$ is independent of \mathcal{T} , v and $\boldsymbol{\psi}$.

Proof. The proof is given in abstract form by Theorem 2, once the conditions made there are verified. The Poincaré–Friedrichs inequality holds for $H_0^2(\Omega)$ so that we only have to check the existence of a Fortin operator \mathcal{F} satisfying (8). Such an operator is given in [39, Lemma 11], see Π^{Grad} there. \square

3.3 Nodal-continuous primal hybrid formulation

We present a formulation that gives the variational framework for a Morley-type element, cf. [50]. We select

$$\tilde{H}(A, \mathcal{T}) := H_0^{2, \mathcal{N}}(\mathcal{T}) := \{v \in H^2(\mathcal{T}); v \text{ is continuous at } x \in \mathcal{N}(\Omega) \text{ and vanishes at } x \in \mathcal{N}(\Gamma)\}$$

and denote $\gamma_{\text{dDiv},J0} := \gamma_{A^*,\mathcal{S}}$ and

$$H_{J0}^{-3/2,-1/2}(\mathcal{S}) := H(A^*, \mathcal{S}) = \gamma_{\text{dDiv},J0}(H(\text{ddiv}, \Omega; \mathbb{S})), \quad \|\cdot\|_{-3/2,-1/2,J0,\mathcal{S}} := \|\cdot\|_{A^*,\mathcal{S}}. \quad (24)$$

Index notation “ $J0$ ” (“jump zero”) refers to the fact that tangential-normal jumps of bending moments do not appear, see (26) below.

The generalized primal hybrid formulation (7) becomes the *nodal-continuous primal hybrid formulation* of the Kirchhoff–Love model: *Find* $u \in H_0^{2,\mathcal{N}}(\mathcal{T})$ and $\boldsymbol{\eta} \in H_{J0}^{-3/2,-1/2}(\mathcal{S})$ such that

$$(CD^2u, D^2\delta u)_{\mathcal{T}} + \langle \boldsymbol{\eta}, \delta u \rangle_{\mathcal{S}} = (f, \delta u), \quad (25a)$$

$$\langle \boldsymbol{\delta\eta}, u \rangle_{\mathcal{S}} = 0 \quad (25b)$$

holds for any $\delta u \in H_0^{2,\mathcal{N}}(\mathcal{T})$, and $\boldsymbol{\delta\eta} \in H_{J0}^{-3/2,-1/2}(\mathcal{S})$.

Theorem 10. *Let* $f \in L_2(\Omega)$ *be given. Problem (25) is well posed. Its solution* $(u, \boldsymbol{\eta})$ *satisfies*

$$\|u\|_{2,\mathcal{T}} + \|\boldsymbol{\eta}\|_{-3/2,-1/2,J0,\mathcal{S}} \leq C\|f\|$$

with a constant C that is independent of f and \mathcal{T} . Furthermore, $u \in H_0^2(\Omega)$, $\boldsymbol{\eta} = \gamma_{\text{dDiv},J0}(CD^2u)$, and $u, \mathbf{M} := CD^2u$ solve (18).

Proof. The proof is identical to the proof of Theorem 8. □

Discretization

To discretize trace space $H_{J0}^{-3/2,-1/2}(\mathcal{S})$, we consider $u \in H_0^{2,\mathcal{N}}(\mathcal{T})$ and proceed as in (21) to find that a sufficiently smooth (piecewise polynomial) tensor $\mathbf{M} \in H(\text{ddiv}, \Omega; \mathbb{S})$ satisfies

$$\langle \gamma_{\text{dDiv},J0}(\mathbf{M}), u \rangle_{\mathcal{S}} = \sum_{T \in \mathcal{T}} (\mathbf{n} \cdot \text{div } \mathbf{M} - \partial_t(\mathbf{t} \cdot \mathbf{M} \mathbf{n}), u)_{\partial T} - \langle \mathbf{n} \cdot \mathbf{M} \mathbf{n}, \partial_n u \rangle_{\partial T} \quad \forall u \in H_0^{2,\mathcal{N}}(\mathcal{T}). \quad (26)$$

Here, we used that

$$\sum_{T \in \mathcal{T}: x \in \mathcal{V}(T)} [\mathbf{t} \cdot \mathbf{M} \mathbf{n}]_{\partial T}(x) = 0 \quad \forall x \in \mathcal{N}(\Omega)$$

in distributional sense (testing with C^∞ -functions with support in a neighborhood of x), see [40, Proposition 3.6]. We conclude that trace $\gamma_{\text{dDiv},J0}(\mathbf{M})$ has two scalar components living on the skeleton, the effective shear force $\mathbf{n} \cdot \text{div } \mathbf{M} + \partial_t(\mathbf{t} \cdot \mathbf{M} \mathbf{n})$ (22a) and the normal-normal trace $\mathbf{n} \cdot \mathbf{M} \mathbf{n}$ (22b). Corner forces $[\mathbf{t} \cdot \mathbf{M} \mathbf{n}]_{\partial T}(x)$ (22c) do not appear. We approximate both trace components by edge-piecewise constant functions, and use nodal-continuous piecewise cubic polynomials to discretize $H_0^{2,\mathcal{N}}(\mathcal{T})$. In our abstract notation, the approximation spaces are

$$\tilde{H}_h(A, \mathcal{T}) := \mathbb{X}^{3,\mathcal{N}}(\mathcal{T}) := P^3(\mathcal{T}) \cap H_0^{2,\mathcal{N}}(\mathcal{T}), \quad H_h(A^*, \mathcal{S}) := P^0(\mathcal{S}) \times P^0(\mathcal{S}).$$

The dimensions of $\mathbb{X}^{3,\mathcal{N}}(\mathcal{T})$ and $P^0(\mathcal{S}) \times P^0(\mathcal{S})$ are $|\mathcal{N}(\Omega)| + 7|\mathcal{T}|$ and $2|\mathcal{E}|$, respectively.

The resulting nodal-continuous primal hybrid method reads as follows. *Find* $u_h \in \mathbb{X}^{3,\mathcal{N}}(\mathcal{T})$ and $\boldsymbol{\eta}_h \in P^0(\mathcal{S}) \times P^0(\mathcal{S})$ such that

$$(CD^2u_h, D^2\delta u)_{\mathcal{T}} + \langle \boldsymbol{\eta}_h, \delta u \rangle_{\mathcal{S}} = (f, \delta u), \quad (27a)$$

$$\langle \boldsymbol{\delta\eta}, u_h \rangle_{\mathcal{S}} = 0 \quad (27b)$$

holds for any $\delta u \in \mathbb{X}^{3,\mathcal{N}}(\mathcal{T})$ and $\boldsymbol{\delta\eta} \in P^0(\mathcal{S}) \times P^0(\mathcal{S})$.

Remark 11. *We note that scheme (23) can be interpreted as a hybridization of scheme (27) or, vice versa, scheme (27) as a reduction of (23) by elimination of vertex discontinuities of u_h . In fact, the degrees stemming from the component $\mathbb{R}_{\text{const}}^{3|\mathcal{T}|}$ in (23) fix the vertex values of u_h .*

Scheme (27) converges quasi-optimally.

Theorem 12. *Let $f \in L_2(\Omega)$ be given. System (27) is well posed. Its solution $(u_h, \boldsymbol{\eta}_h)$ satisfies*

$$\|u - u_h\|_{2,\mathcal{T}} + \|\boldsymbol{\eta} - \boldsymbol{\eta}_h\|_{-3/2,-1/2,J_0,\mathcal{S}} \leq C \left(\|u - v\|_{2,\mathcal{T}} + \|\boldsymbol{\eta} - \boldsymbol{\psi}\|_{-3/2,-1/2,J_0,\mathcal{S}} \right)$$

for any $v \in \mathbb{X}^{3,\mathcal{N}}(\mathcal{T})$ and $\boldsymbol{\psi} \in P^0(\mathcal{S}) \times P^0(\mathcal{S})$. Here, $(u, \boldsymbol{\eta})$ is the solution of (25) and $C > 0$ is independent of \mathcal{T} , v and $\boldsymbol{\psi}$.

According to Remark 11, this theorem follows from Theorem 9 as a special case.

3.4 Continuous primal hybrid formulation

The next formulation gives rise to a scheme that uses continuous approximations of deflections, similarly to the Zienkiewicz triangular element [8], the non-conforming approach in [7], C^0 -interior penalty methods [36, 12], though without stabilization, and the HHO method in [33].

We select

$$\tilde{H}(A, \mathcal{T}) := H_0^{2,1}(\mathcal{T}) := H^2(\mathcal{T}) \cap H_0^1(\Omega) \not\subseteq H(A, \mathcal{T}) = H^2(\mathcal{T})$$

and denote

$$\gamma_{\text{dDiv},nn} := \gamma_{A^*,\mathcal{S}}, \quad H^{-1/2}(\mathcal{S}) := H(A^*, \mathcal{S}) = \gamma_{\text{dDiv},nn}(H(\text{ddiv}, \Omega; \mathbb{S})), \quad \|\cdot\|_{-1/2,\mathcal{S}} := \|\cdot\|_{A^*,\mathcal{S}}.$$

Notation $\gamma_{\text{dDiv},nn}$ is suggested by the fact that the duality with test functions that are continuous across \mathcal{S} gives rise to the normal-normal traces on \mathcal{S} , see (29) below. For localized traces in this context with Banach spaces see also [33, §2.4].

The generalized primal hybrid formulation (7) becomes the *continuous primal hybrid formulation* of the Kirchhoff–Love model: Find $u \in H_0^{2,1}(\mathcal{T})$ and $\eta \in H^{-1/2}(\mathcal{S})$ such that

$$(CD^2u, D^2\delta u)_{\mathcal{T}} + \langle \eta, \delta u \rangle_{\mathcal{S}} = (f, \delta u), \quad (28a)$$

$$\langle \delta \eta, u \rangle_{\mathcal{S}} = 0 \quad (28b)$$

holds for any $\delta u \in H_0^{2,1}(\mathcal{T})$, and $\delta \eta \in H^{-1/2}(\mathcal{S})$.

Theorem 13. *Let $f \in L_2(\Omega)$ be given. Problem (28) is well posed. Its solution (u, η) satisfies*

$$\|u\|_{2,\mathcal{T}} + \|\eta\|_{-1/2,\mathcal{S}} \leq C \|f\|$$

with a constant C that is independent of f and \mathcal{T} . Furthermore, $u \in H_0^2(\Omega)$, $\eta = \gamma_{\text{dDiv},nn}(CD^2u)$, and $u, \mathbf{M} := CD^2u$ solve (18).

Proof. The proof is identical to the proof of Theorem 8. □

Discretization

In order to discretize trace space $H^{-1/2}(\mathcal{S})$ we need an explicit representation of trace operator $\gamma_{\text{dDiv},nn}$ for sufficiently smooth (piecewise polynomial) tensors $\mathbf{M} \in H(\text{ddiv}, \Omega; \mathbb{S})$. Proceeding as in (21), we find that such a tensor \mathbf{M} satisfies

$$\langle \gamma_{\text{dDiv},nn}(\mathbf{M}), u \rangle_{\mathcal{S}} = - \sum_{T \in \mathcal{T}} \langle \mathbf{n} \cdot \mathbf{M} \mathbf{n}, \partial_n u \rangle_{\partial T} \quad \forall u \in H_0^{2,1}(\mathcal{T}). \quad (29)$$

Therefore, $\gamma_{\text{dDiv},nn}(\mathbf{M})|_E$ can be identified with $-\mathbf{n} \cdot \mathbf{M} \mathbf{n}|_E$ on edges $E \in \mathcal{E}$. We approximate these normal-normal traces by edge-piecewise constant functions, and use standard continuous,

piecewise cubic polynomials, enriched with element-bubble functions of degree 4, to discretize $H_0^{2,1}(\mathcal{T})$. Specifically, we define the following spaces,

$$\begin{aligned} P_b^4(T) &:= P^4(T) \cap H_0^1(T), & \mathbb{X}_b^4(T) &:= P^3(T) + P_b^4(T) \quad (T \in \mathcal{T}), \\ P_0^{3,c}(\mathcal{T}) &:= P^3(\mathcal{T}) \cap H_0^1(\Omega), & \mathbb{X}_b^{4,c}(\mathcal{T}) &:= P_0^{3,c}(\mathcal{T}) + P_b^4(\mathcal{T}) \end{aligned}$$

and, in our abstract notation, select the approximation spaces

$$\tilde{H}_h(A, \mathcal{T}) := \mathbb{X}_b^{4,c}(\mathcal{T}) \subset H_0^{2,1}(\mathcal{T}), \quad H_h(A^*, \mathcal{S}) := P^0(\mathcal{S}) \subset H^{-1/2}(\mathcal{S}).$$

Here, $P^0(\mathcal{S})$ is the space of edge-piecewise constant functions on \mathcal{S} introduced previously. The dimensions of $\mathbb{X}_b^{4,c}(\mathcal{T})$ and $P^0(\mathcal{S})$ are $|\mathcal{N}(\Omega)| + 2|\mathcal{E}(\Omega)| + 3|\mathcal{T}|$ and $|\mathcal{E}|$, respectively (see degrees of freedom (31) and note that the continuity of $v \in \mathbb{X}_b^{4,c}(\mathcal{T})$ requires unique vertex values $v(x)$ and edge moments $\langle v, \phi \rangle_E$).

The resulting continuous primal hybrid method reads as follows. Find $u_h \in \mathbb{X}_b^{4,c}(\mathcal{T})$ and $\eta_h \in P^0(\mathcal{S})$ such that

$$(CD^2 u_h, D^2 \delta u)_{\mathcal{T}} + \langle \eta_h, \delta u \rangle_{\mathcal{S}} = (f, \delta u), \quad (30a)$$

$$\langle \delta \eta, u_h \rangle_{\mathcal{S}} = 0 \quad (30b)$$

holds for any $\delta u \in \mathbb{X}_b^{4,c}(\mathcal{T})$ and $\delta \eta \in P^0(\mathcal{S})$. It converges quasi-optimally.

Theorem 14. *Let $f \in L_2(\Omega)$ be given. System (30) is well posed. Its solution (u_h, η_h) satisfies*

$$\|u - u_h\|_{2,\mathcal{T}} + \|\eta - \eta_h\|_{-1/2,\mathcal{S}} \leq C \left(\|u - v\|_{2,\mathcal{T}} + \|\eta - \psi\|_{-1/2,\mathcal{S}} \right)$$

for any $v \in \mathbb{X}_b^{4,c}(\mathcal{T})$ and $\psi \in P^0(\mathcal{S})$. Here, (u, η) is the solution of (28) and $C > 0$ is independent of \mathcal{T} , v and ψ .

A proof of this theorem is given by Theorem 2. We only have to verify the existence of a Fortin operator, denoted by \mathcal{F}_b^4 . This needs some preparation. For a definition of \mathcal{F}_b^4 and the verification of Fortin properties, see (34) and Lemma 17 below.

Lemma 15. *Given $T \in \mathcal{T}$, the local space $\mathbb{X}_b^4(T)$ has dimension 12. A function $v \in \mathbb{X}_b^4(T)$ has the degrees of freedom*

$$v(x), \quad \langle v, \phi \rangle_E, \quad \langle \mathbf{n} \cdot \nabla v, 1 \rangle_E \quad (31)$$

for $x \in \mathcal{V}(T)$, $\phi \in P^1(E)$, $E \in \mathcal{E}(T)$.

Proof. For $T \in \mathcal{T}$ we use the notation λ_j for barycentric coordinates, vertices x_j , and edges E_j . Let us denote $\psi_0 := \lambda_1 \lambda_2 \lambda_3$, the lowest order polynomial bubble function. Space $P_b^4(T)$ is spanned by $\{\phi \psi_0; \phi \in P^1(T)\}$ with basis $\{\psi_j := \lambda_j \psi_0; j = 1, 2, 3\}$ and therefore has dimension 3. Since $P^3(T) \cap P_b^4(T)$ is generated by ψ_0 , we conclude that $\mathbb{X}_b^4(T)$ has dimension $10 + 3 - 1 = 12$, equal to the number of claimed degrees of freedom. It is enough to show their injectivity.

Let $v \in \mathbb{X}_b^{4,c}(T)$ be given with vanishing degrees of freedom (31). The vanishing of the vertex values means that $v|_{E_j} \in (\lambda_{j+1} \lambda_{j+2})|_{E_j} P^1(E_j)$, and its orthogonality to $P^1(E_j)$ implies that $v|_{E_j} = 0$, $j = 1, 2, 3$, thus $v \in P_b^4(T)$. Therefore, function v has a representation $v = \sum_{j=1}^3 c_j \psi_j$. We calculate

$$\nabla v = 2\lambda_1 \lambda_2 \lambda_3 \sum_{j=1}^3 c_j \nabla \lambda_j + \sum_{j=1}^3 c_j \lambda_j^2 \left(\lambda_{j+1} \nabla \lambda_{j+2} + \lambda_{j+2} \nabla \lambda_{j+1} \right),$$

that is, on edge E_k ,

$$\nabla v = c_{k+1} \lambda_{k+1}^2 \lambda_{k+2} \nabla \lambda_k + c_{k+2} \lambda_{k+2}^2 \lambda_{k+1} \nabla \lambda_k = \left(c_{k+1} \lambda_{k+1} + c_{k+2} \lambda_{k+2} \right) \lambda_{k+1} \lambda_{k+2} \nabla \lambda_k.$$

Moments $\langle \mathbf{n} \cdot \nabla v, 1 \rangle_{E_k}$ vanish iff $c_{k+1} \lambda_{k+1} + c_{k+2} \lambda_{k+2}$ vanishes at the midpoint of E_k , that is, iff $c_k + c_{k+1} = 0$, $k = 1, 2, 3$, with only solution $c_1 = c_2 = c_3 = 0$, $v = 0$. \square

We prove the existence of a Fortin operator $\mathcal{F}_b^4 : H_0^{2,1}(\mathcal{T}) \rightarrow \mathbb{X}_b^{4,c}(\mathcal{T})$ in two steps.

Lemma 16. *There is an operator $\tilde{\mathcal{F}}_b^4 : H_0^{2,1}(\mathcal{T}) \rightarrow P_b^4(\mathcal{T})$ that satisfies*

$$\langle \delta\eta, \tilde{\mathcal{F}}_b^4 u \rangle_{\mathcal{S}} = \langle \delta\eta, u \rangle_{\mathcal{S}} \quad \forall \delta\eta \in P^0(\mathcal{S}), \quad u \in H_0^{2,1}(\mathcal{T}) \quad (32)$$

and

$$\| \tilde{\mathcal{F}}_b^4 u \| + \| h_{\mathcal{T}}^2 D^2 \tilde{\mathcal{F}}_b^4 u \|_{\mathcal{T}} \lesssim \| u \| + \| h_{\mathcal{T}}^2 D^2 u \|_{\mathcal{T}} \quad \forall u \in H_0^{2,1}(\mathcal{T}). \quad (33)$$

Proof. By Lemma 15, for every $u \in H_0^{2,1}(\Omega)$ there are unique polynomials $q_T \in P_b^4(T)$ with the respective degrees of freedom (edge moments of normal derivative) from $u|_T$, $T \in \mathcal{T}$. They give rise to a function $q \in H_0^1(\Omega)$ ($q|_T := q_T \quad \forall T \in \mathcal{T}$) whose normal derivatives on edges have jumps (values on edges $\subset \Gamma$) with mean-value zero, that is, $\tilde{\mathcal{F}}_b^4 u := q$ satisfies (32). By the boundedness of the associated degrees of freedom (in H^2 on a reference element) and scaling properties, one establishes (33). \square

We define

$$\mathcal{F}_b^4 : H_0^{2,1}(\mathcal{T}) \ni u \mapsto \tilde{\mathcal{F}}_b^4(u - \mathcal{I}_{\mathcal{T}}^1 u) + \mathcal{I}_{\mathcal{T}}^1 u \quad (34)$$

where $\mathcal{I}_{\mathcal{T}}^1 : H^2(\mathcal{T}) \rightarrow P^1(\mathcal{T})$ is the element-piecewise nodal interpolation operator. Of course, it maps $H_0^{2,1}(\mathcal{T}) \rightarrow P_0^{1,c}(\mathcal{T})$.

Lemma 17. *Operator \mathcal{F}_b^4 defined in (34) is a mapping $\mathcal{F}_b^4 : H_0^{2,1}(\mathcal{T}) \rightarrow \mathbb{X}_b^{4,c}(\mathcal{T})$ and satisfies*

$$\langle \delta\eta, \mathcal{F}_b^4 u \rangle_{\mathcal{S}} = \langle \delta\eta, u \rangle_{\mathcal{S}} \quad \forall \delta\eta \in P^0(\mathcal{S}), \quad u \in H_0^{2,1}(\mathcal{T}), \quad (35)$$

$$\| \mathcal{F}_b^4 u \| + \| D^2 \mathcal{F}_b^4 u \|_{\mathcal{T}} \lesssim \| u \| + \| D^2 u \|_{\mathcal{T}} \quad \forall u \in H_0^{2,1}(\mathcal{T}). \quad (36)$$

Proof. Let $u \in H_0^{2,1}(\mathcal{T})$ be given. By construction, $\mathcal{F}_b^4 u \in P_0^{1,c}(\mathcal{T}) \oplus P_b^4(\mathcal{T})$, and $\mathcal{I}_{\mathcal{T}}^1 u \in P_0^{1,c}(\mathcal{T})$. Therefore, relation (32) implies (35):

$$\langle \delta\eta, \mathcal{F}_b^4 u \rangle_{\mathcal{S}} = \langle \delta\eta, \tilde{\mathcal{F}}_b^4(u - \mathcal{I}_{\mathcal{T}}^1 u) \rangle_{\mathcal{S}} + \langle \delta\eta, \mathcal{I}_{\mathcal{T}}^1 u \rangle_{\mathcal{S}} = \langle \delta\eta, u \rangle_{\mathcal{S}} \quad \forall \delta\eta \in P^0(\mathcal{S}).$$

Property $D^2(\mathcal{I}_{\mathcal{T}}^1 u)|_T = 0$ ($T \in \mathcal{T}$), bound (33) and the approximation property $\|u - \mathcal{I}_{\mathcal{T}}^1 u\| \lesssim \|h_{\mathcal{T}}^2 D^2 u\|_{\mathcal{T}}$ show that $\mathcal{F}_b^4 u$ satisfies

$$\| D^2 \mathcal{F}_b^4 u \|_{\mathcal{T}} = \| D^2 \tilde{\mathcal{F}}_b^4(u - \mathcal{I}_{\mathcal{T}}^1 u) \| \lesssim \| h_{\mathcal{T}}^{-2}(u - \mathcal{I}_{\mathcal{T}}^1 u) \| + \| D^2(u - \mathcal{I}_{\mathcal{T}}^1 u) \|_{\mathcal{T}} \lesssim \| D^2 u \|_{\mathcal{T}}.$$

The L_2 -estimate in (36) follows with (33) and scaling arguments to bound $\| \mathcal{I}_{\mathcal{T}}^1 u \| \lesssim \| u \| + \| h_{\mathcal{T}}^2 D^2 u \|_{\mathcal{T}}$:

$$\| \mathcal{F}_b^4 u \| \leq \| \tilde{\mathcal{F}}_b^4(u - \mathcal{I}_{\mathcal{T}}^1 u) \| + \| \mathcal{I}_{\mathcal{T}}^1 u \| \lesssim \| u - \mathcal{I}_{\mathcal{T}}^1 u \| + \| h_{\mathcal{T}}^2 D^2 u \| + \| \mathcal{I}_{\mathcal{T}}^1 u \| \lesssim \| u \| + \| D^2 u \|_{\mathcal{T}}.$$

This finishes the proof. \square

3.5 Mixed hybrid formulation

The following formulation gives rise to the so-called assumed stresses hybrid method by Pian and Tong [53], cf. the analysis by Brezzi and Marini [16]. We select

$$\tilde{H}(A^*, \mathcal{T}) := H(A^*, \mathcal{T}) = H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$$

and denote

$$H_0^{3/2,1/2}(\mathcal{S}) := H_0(A, \mathcal{S}) = \gamma_2(H_0^2(\Omega)), \quad \|\cdot\|_{3/2,1/2,\mathcal{S}} := \|\cdot\|_{A,\mathcal{S}}.$$

In this case, with $\tilde{H}(A^*, \mathcal{T}) = H(A^*, \mathcal{T})$, $\gamma_{A,\mathcal{S}} = \gamma_2$ defined before in (19).

The generalized mixed hybrid formulation (10) becomes the *mixed hybrid formulation* of the Kirchhoff–Love model: Find $\mathbf{M} \in H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$, $u \in L_2(\Omega)$, and $\boldsymbol{\psi} \in H_0^{3/2,1/2}(\mathcal{S})$ such that

$$(\mathcal{C}^{-1} \mathbf{M}, \boldsymbol{\delta M}) - (u, \operatorname{div} \mathbf{div} \boldsymbol{\delta M})_{\mathcal{T}} - \langle \boldsymbol{\psi}, \boldsymbol{\delta M} \rangle_{\mathcal{S}} = 0, \quad (37a)$$

$$- (\operatorname{div} \mathbf{div} \mathbf{M}, \boldsymbol{\delta u})_{\mathcal{T}} - \langle \boldsymbol{\delta \psi}, \mathbf{M} \rangle_{\mathcal{S}} = -(f, \boldsymbol{\delta u}) \quad (37b)$$

holds for any $\boldsymbol{\delta M} \in H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$, $\boldsymbol{\delta u} \in L_2(\Omega)$, and $\boldsymbol{\delta \psi} \in H_0^{3/2,1/2}(\mathcal{S})$. It is an extended form of formulation [16, (1.23)] without an extension of right-hand side function f to an element of $H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$.

Theorem 18. *Let $f \in L_2(\Omega)$ be given. Problem (37) is well posed. Its solution $(\mathbf{M}, u, \boldsymbol{\psi})$ satisfies*

$$\|\mathbf{M}\|_{\mathbf{ddiv}, \mathcal{T}} + \|u\| + \|\boldsymbol{\psi}\|_{3/2,1/2,\mathcal{S}} \leq C \|f\|$$

with a constant C that is independent of f and \mathcal{T} . Furthermore, $u \in H_0^2(\Omega)$, $\mathbf{M} = \mathcal{C} D^2 u$, $\boldsymbol{\psi} = \gamma_2(u)$, and u, \mathbf{M} solve (18).

Proof. We have already seen that the Poincaré–Friedrichs inequality (5a) holds true. Inf-sup property (5b) reads

$$\sup_{\mathbf{Q} \in H(\mathbf{ddiv}, \Omega; \mathbb{S})} \frac{(\operatorname{div} \mathbf{div} \mathbf{Q}, v)}{\|\mathbf{Q}\|_{\mathbf{ddiv}}} \geq c_{\text{is}} \|v\| \quad \forall v \in L_2(\Omega)$$

and is satisfied by the surjectivity of $\operatorname{div} \mathbf{div} : H(\mathbf{ddiv}, \Omega; \mathbb{S}) \rightarrow L_2(\Omega)$. In fact, given $g \in L_2(\Omega)$, there is a (unique) solution $v_g \in H_0^2(\Omega)$ to $\operatorname{div} \mathbf{div} D^2 v_g = \Delta^2 v_g = g$ in Ω , and $\mathbf{Q} := D^2 v_g \in H(\mathbf{ddiv}, \Omega; \mathbb{S})$ satisfies $\operatorname{div} \mathbf{div} \mathbf{Q} = g$. Theorem 3 proves the statements. \square

Discretization

In order to discretize trace space $H_0^{3/2,1/2}(\mathcal{S})$ we need to represent trace operator γ_2 explicitly. This is dual to the setting of the primal hybrid formulation, cf. (21). By our definition of γ_2 we have a sign change:

$$\begin{aligned} \langle \gamma_2(u), \mathbf{M} \rangle_{\mathcal{S}} &= \sum_{T \in \mathcal{T}} \langle \gamma_{2,\partial T}(u), \mathbf{M} \rangle_{\partial T} := \sum_{T \in \mathcal{T}} (D^2 u, \mathbf{M})_T - (u, \operatorname{div} \mathbf{div} \mathbf{M})_{\mathcal{T}} \\ &= \sum_{T \in \mathcal{T}} \left(\langle \mathbf{n} \cdot \mathbf{M} \mathbf{n}, \partial_n u \rangle_{\partial T} - \langle \mathbf{n} \cdot \mathbf{div} \mathbf{M} + \partial_t(\mathbf{t} \cdot \mathbf{M} \mathbf{n}), u \rangle_{\partial T} + \sum_{x \in \mathcal{N}(T)} [\mathbf{t} \cdot \mathbf{M} \mathbf{n}]_{\partial T}(x) u(x) \right) \end{aligned} \quad (38)$$

for $u \in H_0^2(\Omega)$ and a sufficiently piecewise-smooth tensor $\mathbf{M} \in H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$. Here, we introduced the local trace operators $\gamma_{2,\partial T}$ for notational convenience. Trace $\boldsymbol{\psi}|_{\partial T} = \gamma_{2,\partial T}(u)$ consists of the components $u|_{\partial T}$ and $\partial_n u|_{\partial T}$, the canonical traces of u and the normal component $\mathbf{n} \cdot \nabla u$.

We approximate $\boldsymbol{\psi}$ by traces of the reduced Hsieh–Clough–Tocher (HCT) composite element which are edge-piecewise cubic polynomials, cf. [23, 40],

$$\begin{aligned} HCT(T) &:= \{v \in H^2(T); \Delta^2 v + v = 0, v|_E \in P^3(E), \mathbf{n} \cdot \nabla v|_E \in P^1(E) \ \forall E \in \mathcal{E}(T)\}, \quad (39) \\ HCT_0^2(\mathcal{S}) &:= \gamma_2(HCT(\mathcal{T}) \cap H_0^2(\Omega)). \end{aligned}$$

Bending moments are approximated by a reduction of the $H(\text{ddiv}; \mathbb{S})$ element from [38]. Specifically, for $T \in \mathcal{T}$ we use the Raviart–Thomas spaces

$$RT^s(T) = \mathbf{x}P_{\text{hom}}^s(T) \oplus P^s(T; \mathbb{R}^2) \quad (s \in \mathbb{N}_0)$$

where $\mathbf{x} : \mathbb{R}^2 \ni (x_1, x_2) \mapsto (x_1, x_2)^\top$, denote $\text{sym}(\mathbf{Q}) := (\mathbf{Q} + \mathbf{Q}^\top)/2$ and introduce

$$\mathbb{X}^{\text{dDiv}}(T) := \text{sym}(RT^0(T) \otimes RT^1(T)).$$

We approximate bending moments by the $\mathbb{X}^{\text{dDiv}}(T)$ -element reduced to constant normal-normal edge traces,

$$\mathbb{X}^{\text{dDiv}, \text{nnc}}(T) := \{\mathbf{M} \in \mathbb{X}^{\text{dDiv}}(T); \mathbf{n} \cdot \mathbf{M} \mathbf{n}|_E \in P^0(E), E \in \mathcal{E}(T)\}.$$

In our abstract notation, the approximation spaces are

$$\tilde{H}_h(A^*, \mathcal{T}) := \mathbb{X}^{\text{dDiv}, \text{nnc}}(\mathcal{T}), \quad H_h(\mathcal{T}) := P^1(\mathcal{T}), \quad H_h(A, \mathcal{S}) := HCT_0^2(\mathcal{S}).$$

Their respective dimensions are $12|\mathcal{T}|$, $3|\mathcal{T}|$, and $3|\mathcal{N}(\Omega)|$.

The resulting mixed hybrid scheme reads as follows. Find $\mathbf{M}_h \in \mathbb{X}^{\text{dDiv}, \text{nnc}}(\mathcal{T})$, $u_h \in P^1(\mathcal{T})$, and $\boldsymbol{\psi}_h \in HCT_0^2(\mathcal{S})$ such that

$$(\mathcal{C}^{-1} \mathbf{M}_h, \boldsymbol{\delta} \mathbf{M}) - (u_h, \text{div} \text{div} \boldsymbol{\delta} \mathbf{M})_{\mathcal{T}} - \langle \boldsymbol{\psi}_h, \boldsymbol{\delta} \mathbf{M} \rangle_{\mathcal{S}} = 0, \quad (40a)$$

$$-(\text{div} \text{div} \mathbf{M}_h, \delta u)_{\mathcal{T}} - \langle \delta \boldsymbol{\psi}, \mathbf{M}_h \rangle_{\mathcal{S}} = -(f, \delta u) \quad (40b)$$

holds for any $\boldsymbol{\delta} \mathbf{M} \in \mathbb{X}^{\text{dDiv}, \text{nnc}}(\mathcal{T})$, $\delta u \in P^1(\mathcal{T})$, and $\delta \boldsymbol{\psi} \in HCT_0^2(\mathcal{S})$.

It converges quasi-optimally.

Theorem 19. *Let $f \in L_2(\Omega)$ be given. System (40) is well posed. Its solution $(\mathbf{M}_h, u_h, \boldsymbol{\psi}_h)$ satisfies,*

$$\|\mathbf{M} - \mathbf{M}_h\|_{\text{ddiv}, \mathcal{T}} + \|u - u_h\| + \|\boldsymbol{\psi} - \boldsymbol{\psi}_h\|_{3/2, 1/2, \mathcal{S}} \leq C(\|\mathbf{M} - \mathbf{Q}\|_{\text{ddiv}, \mathcal{T}} + \|u - v\| + \|\boldsymbol{\psi} - \boldsymbol{\phi}\|_{3/2, 1/2, \mathcal{S}})$$

for any $\mathbf{Q} \in \mathbb{X}^{\text{dDiv}, \text{nnc}}(\mathcal{T})$, $v \in P^1(\mathcal{T})$, and $\boldsymbol{\phi} \in HCT_0^2(\mathcal{S})$. Here, $(\mathbf{M}, u, \boldsymbol{\psi})$ is the solution of (37) and $C > 0$ is independent of \mathcal{T} , \mathbf{Q} , v , and $\boldsymbol{\phi}$.

A proof of this theorem is given in abstract form by Theorem 5. We only have to check the existence of Fortin operator components $\mathcal{F}_1, \mathcal{F}_2$ that satisfy (11). This is done in the remainder of this section, see (46) and Lemma 22 below.

Element $\mathbb{X}^{\text{dDiv}}(T)$ has the following 15 degrees of freedom, cf. (22),

$$\langle \mathbf{n} \cdot \text{div} \mathbf{M} + \partial_t(\mathbf{t} \cdot \mathbf{M} \mathbf{n}), \phi \rangle_E, \quad \phi \in P^1(E), \quad E \in \mathcal{E}, \quad (41a)$$

$$\langle \mathbf{n} \cdot \mathbf{M} \mathbf{n}, \phi \rangle_E, \quad \phi \in P^1(E), \quad E \in \mathcal{E}, \quad (41b)$$

$$[\mathbf{t} \cdot \mathbf{M} \mathbf{n}]_{\partial T}(x), \quad x \in \mathcal{N}(T). \quad (41c)$$

These degrees, taken for every element $T \in \mathcal{T}$ with unique values for edges $E \in \mathcal{E}$ and vertices $x \in \mathcal{N}$, subject to constraints (22d) at interior vertices, define an interpolation operator

$$\Pi_{\mathcal{T}}^{\text{dDiv}} : H(\text{ddiv}, \Omega; \mathbb{S}) \cap H^r(\Omega; \mathbb{S}) \rightarrow \mathbb{X}^{\text{dDiv}}(\mathcal{T}) \cap H(\text{ddiv}, \Omega; \mathbb{S}) \quad (42)$$

for $r > 3/2$. According to [38, Proposition 10], this interpolation operator satisfies

$$\operatorname{div} \mathbf{div} \Pi_{\mathcal{T}}^{\operatorname{dDiv}} = \Pi_{\mathcal{T}}^1 \operatorname{div} \mathbf{div}, \quad (43)$$

$$\|\Pi_{\mathcal{T}}^{\operatorname{dDiv}} \mathbf{M}\| \lesssim \|\mathbf{M}\|_r \quad \forall \mathbf{M} \in H(\operatorname{ddiv}, \Omega; \mathbb{S}) \cap H^r(\Omega; \mathbb{S}) \quad (r > 3/2). \quad (44)$$

We use the reduced element $\mathbb{X}^{\operatorname{dDiv}, \operatorname{nnc}}(T)$ with only constant moments in (41b). Inspection of the details in [38] reveals that the corresponding interpolation operator $\Pi_{\mathcal{T}}^{\operatorname{dDiv}, \operatorname{nnc}}$ satisfies properties (43) and (44) as well. In particular, the commutativity property can be seen by evaluating $(\operatorname{div} \mathbf{div} \Pi_{\mathcal{T}}^{\operatorname{dDiv}, \operatorname{nnc}} \mathbf{M}, v)_T$ for $v \in P^1(T)$ and integrating by parts. The normal-normal trace of \mathbf{M} meets the normal derivative of v , an edge-wise constant, and thus gives only rise to the constant moments in (41b). The fact that $\Pi_{\mathcal{T}}^{\operatorname{dDiv}, \operatorname{nnc}}$ maps $H(\operatorname{ddiv}, \Omega; \mathbb{S}) \cap H^r(\Omega; \mathbb{S})$ (for $r > 1/2$) to $\mathbb{X}^{\operatorname{dDiv}, \operatorname{nnc}}(\mathcal{T}) \cap H(\operatorname{ddiv}, \Omega; \mathbb{S})$ holds by (42) and the selection of the degrees of freedom.

Element $HCT(T)$ has dimension 9. The canonical degrees of freedom of $v \in HCT(T)$ are

$$v(x), \nabla v(x) \quad (x \in \mathcal{N}(T)) \quad \text{for } T \in \mathcal{T}. \quad (45)$$

We continue to identify degrees that are dual to the degrees (41a), (41c) of $\mathbb{X}^{\operatorname{dDiv}, \operatorname{nnc}}(T)$.

Lemma 20. *A function $v \in HCT(T)$ with $T \in \mathcal{T}$ is uniquely defined by*

$$v(x) \quad (x \in \mathcal{N}(T)) \quad \text{and} \quad \langle \delta\psi, v \rangle_E \quad (\delta\psi \in P^1(E), E \in \mathcal{E}(T)).$$

Proof. Since the dimension of $HCT(T)$ is equal to the number of specified degrees of freedom, it is enough to show that an element $v \in HCT(T)$ with vanishing degrees is zero. For given $T \in \mathcal{T}$ with vertices x_j and opposite edges E_j , $j = 1, 2, 3$, let λ_j , $j = 1, 2, 3$, be the barycentric coordinates. By definition, $v|_E \in P^3(E)$ for any $E \in \mathcal{E}(T)$. Setting $v(x_j) = 0$, $j = 1, 2, 3$, means that $v|_{E_j}$ is a cubic polynomial that vanishes at the endpoints x_{j+1} , x_{j+2} , $j = 1, 2, 3$ (we use a numbering modulo 3). Therefore, $v|_{E_j} \in (\lambda_{j+1}\lambda_{j+2} \operatorname{span}\{1, \lambda_{j+2} - \lambda_{j+1}\})|_{E_j}$, $j = 1, 2, 3$. Orthogonalities $\langle 1, v \rangle_{E_j} = \langle \lambda_{j+2} - \lambda_{j+1}, v \rangle_{E_j}$ imply $v|_{E_j} = 0$, $j = 1, 2, 3$. We conclude that the canonical degrees of freedom (45) of v vanish, that is, $v = 0$. \square

Remark 21. *The degrees of freedom given by Lemma 20 are intrinsic to ∂T and therefore uniquely define functions of $HCT(T)$ by their traces. However, these degrees do not imply conformity for $HCT_0^{2,1}(\mathcal{S})$ as traces of $H^2(\Omega)$ without the use of basis functions that are conforming with the canonical degrees of freedom (45).*

We are in a position to define and analyze a Fortin operator. Let $\mathbf{I} \in \mathbb{S}$ denote the identity tensor. For given $\mathbf{M} \in H(\operatorname{ddiv}, \mathcal{T}; \mathbb{S})$ we define

$$\mathcal{F}_1(\mathbf{M}) := \Pi_{\mathcal{T}}^{\operatorname{dDiv}, \operatorname{nnc}}(z\mathbf{I}) \quad \text{with} \quad z \in H_0^1(\Omega) : \Delta z = \Pi_{\mathcal{T}}^1 \operatorname{div}_{\mathcal{T}} \mathbf{div}_{\mathcal{T}} \mathbf{M}, \quad (46a)$$

$$\begin{aligned} \mathcal{F}_2(\mathbf{M}) := \mathbf{Q} := (\mathbf{Q}_T)_T \quad \text{with} \quad \mathbf{Q}_T \in \mathbb{X}^{\operatorname{dDiv}}(T) : \mathbf{n} \cdot \mathbf{Q}_T \mathbf{n}|_{\partial T} = 0, \quad (46b) \\ \langle \gamma_{2, \partial T}(v), \mathbf{Q}_T \rangle_{\partial T} = \langle \gamma_{2, \partial T}(v), \mathbf{M}|_T \rangle_{\partial T} \quad \forall v \in HCT(T), T \in \mathcal{T}. \end{aligned}$$

Lemma 22. *Operators $\mathcal{F}_1, \mathcal{F}_2$ satisfy conditions (11).*

Proof. The construction of operator \mathcal{F}_1 stems from [38, Proof of Theorem 12]. It is well defined by the properties of $\Pi_{\mathcal{T}}^{\operatorname{dDiv}, \operatorname{nnc}}$ because $z \in H^r(\Omega)$ with $r > 3/2$. Relation (11c) follows by using commutativity property (43) for $\Pi_{\mathcal{T}}^{\operatorname{dDiv}, \operatorname{nnc}}$ and noting that $\operatorname{div}_{\mathcal{T}} \mathbf{div}_{\mathcal{T}}(z\mathbf{I}) = \Delta z$. Boundedness (11e) follows by the same relation, together with boundedness (44) for $\Pi_{\mathcal{T}}^{\operatorname{dDiv}, \operatorname{nnc}}$ and stability $\|z\|_r \lesssim \|\operatorname{div} \mathbf{div} \mathbf{M}\|_{\mathcal{T}}$.

Operator \mathcal{F}_2 is well defined (on every element $T \in \mathcal{T}$). In fact, the defining right-hand side is a duality pairing on $H^2(T) \times H(\operatorname{ddiv}, T; \mathbb{S})$, and the degrees of freedom (41a), (41c) of $\mathbb{X}^{\operatorname{dDiv}, \operatorname{nnc}}(T)$

(setting degrees (41b) to zero) and those of $HCT(T)$ are dual to each other, see representation (38) and Lemma 20. By definition, $\mathbf{Q} \in H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$. Given any $v \in HCT(\mathcal{T}) \cap H^2(\Omega)$, \mathbf{Q} satisfies

$$\langle \gamma_2(v), \mathbf{Q} \rangle_{\mathcal{S}} = \sum_{T \in \mathcal{T}} \langle \gamma_{2, \partial T}(v), \mathbf{Q}_T \rangle_{\partial T} = \sum_{T \in \mathcal{T}} \langle \gamma_{2, \partial T}(v), \mathbf{M}|_T \rangle_{\partial T} = \langle \gamma_2(v), \mathbf{M} \rangle_{\mathcal{S}}.$$

This proves (11d). It remains to check the boundedness of \mathcal{F}_2 . The bound $\|\mathbf{Q}\| \lesssim \|\mathbf{M}\|_{\mathbf{ddiv}, \mathcal{T}}$ holds by a finite-dimension argument on every element and scaling properties, cf. [39, Proof of Lemma 16]. The bound $\|\mathbf{div} \mathbf{div} \mathbf{Q}\|_{\mathcal{T}} \leq \|\mathbf{div} \mathbf{div} \mathbf{M}\|_{\mathcal{T}}$ is due to commutativity property (43). \square

Remark 23. *We have proved the discrete inf-sup condition (13) by constructing Fortin operators satisfying (11). Brezzi and Marini used an $H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$ -extension of right-hand side function f , thus avoiding the field variable u . In their case, only an inf-sup condition for bilinear form $\langle \delta\psi, \mathbf{M} \rangle_{\mathcal{S}}$ is required. Our selection of discrete spaces satisfies [16, Theorem 3.7] with $m = 1$, $r = 3$, $s = 1$ and thus proves the discrete inf-sup condition in this case. To this end we note that $P^1(T; \mathbb{S}) \subset \mathbb{X}^{\mathbf{dDiv}, nnc}(T)$ holds as for the full element $\mathbb{X}^{\mathbf{dDiv}}(T)$, cf. [38, Proposition 4].*

3.6 Normal-normal continuous mixed formulation

We present a formulation that provides the variational framework for a type of Hellan–Herrmann–Johnson method that controls bending moments in the product energy space $H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$ rather than in $L_2(\Omega; \mathbb{S})$ augmented with traces, cf. [44, 45, 47]. For a variational formulation in Banach spaces see [10, Cap. 10.3].

Recall trace operator γ_2 introduced in (19). We select

$$\begin{aligned} \tilde{H}(A^*, \mathcal{T}) &:= H_{nn}(\mathbf{ddiv}, \mathcal{T}; \mathbb{S}) := \{\mathbf{Q} \in H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S}); \langle \gamma_2(v), \mathbf{Q} \rangle_{\mathcal{S}} = 0 \ \forall v \in H_0^2(\Omega) \cap H_0^1(\mathcal{T})\} \\ &\subset H(A^*, \mathcal{T}) = H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S}) \end{aligned}$$

with $H_0^1(\mathcal{T}) := \Pi_{T \in \mathcal{T}} H_0^1(T)$ and denote

$$\gamma_{2,1} := \gamma_{A, \mathcal{S}}, \quad H_0^{3/2}(\mathcal{S}) := H_0(A, \mathcal{S}) = \gamma_{2,1}(H_0^2(\Omega)), \quad \|\cdot\|_{3/2, \mathcal{S}} := \|\cdot\|_{A, \mathcal{S}}.$$

Notation $H_{nn}(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$ indicates that sufficiently smooth elements have continuous normal-normal traces across \mathcal{S} whereas notation $\gamma_{2,1}$ refers to the fact that it is the canonical trace operator from $H^1(\Omega)$ onto \mathcal{S} , restricted to $H^2(\Omega)$ and with stronger norm. For a non-trivial mesh \mathcal{T} , $H_{nn}(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$ is a strict subspace of $H(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$.

The generalized mixed hybrid formulation (10) becomes the *normal-normal continuous mixed formulation* of the Kirchhoff–Love model: Find $\mathbf{M} \in H_{nn}(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$, $u \in L_2(\Omega)$, and $\psi \in H_0^{3/2}(\mathcal{S})$ such that

$$(\mathcal{C}^{-1} \mathbf{M}, \delta \mathbf{M}) - (u, \mathbf{div} \mathbf{div} \delta \mathbf{M})_{\mathcal{T}} - \langle \psi, \delta \mathbf{M} \rangle_{\mathcal{S}} = 0, \quad (47a)$$

$$-(\mathbf{div} \mathbf{div} \mathbf{M}, \delta u)_{\mathcal{T}} - \langle \delta \psi, \mathbf{M} \rangle_{\mathcal{S}} = -(f, \delta u) \quad (47b)$$

holds for any $\delta \mathbf{M} \in H_{nn}(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$, $\delta u \in L_2(\Omega)$, and $\delta \psi \in H_0^{3/2}(\mathcal{S})$.

Theorem 24. *Let $f \in L_2(\Omega)$ be given. Problem (47) is well posed. Its solution (\mathbf{M}, u, ψ) satisfies*

$$\|\mathbf{M}\|_{\mathbf{ddiv}, \mathcal{T}} + \|u\| + \|\psi\|_{3/2, \mathcal{S}} \leq C \|f\|$$

with a constant C that is independent of f and \mathcal{T} . Furthermore, $u \in H_0^2(\Omega)$, $\mathbf{M} = \mathcal{C}D^2u \in H(\mathbf{ddiv}, \Omega; \mathbb{S})$, $\psi = \gamma_{2,1}(u)$, and u, \mathbf{M} solve (18).

Proof. The proof is identical to the proof of Theorem 18. \square

Discretization

Relation (38) shows that $u \in H_0^2(\Omega)$ and a sufficiently smooth tensor $\mathbf{M} \in H_{nn}(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$ satisfy

$$\langle \gamma_{2,1}(u), \mathbf{M} \rangle_{\mathcal{S}} = \sum_{T \in \mathcal{T}} \left(\sum_{x \in \mathcal{N}(T)} [\mathbf{t} \cdot \mathbf{M} \mathbf{n}]_{\partial T}(x) u(x) - \langle \mathbf{n} \cdot \mathbf{div} \mathbf{M} + \partial_t(\mathbf{t} \cdot \mathbf{M} \mathbf{n}), u \rangle_{\partial T} \right). \quad (48)$$

As already indicated, we conclude that trace $\psi = \gamma_{2,1}(u)$ reduces to the canonical trace of u onto \mathcal{S} , measured in a stronger norm than the canonical trace operator acting on $H^1(\Omega)$. We approximate ψ by traces of the reduced HCT-element (39),

$$HCT_0^{2,1}(\mathcal{S}) := \gamma_{2,1} \left(HCT(\mathcal{T}) \cap H_0^2(\Omega) \right),$$

and use the reduced-element space $\mathbb{X}^{\mathbf{dDiv}, nnc}(\mathcal{T})$ with continuous normal-normal traces to approximate \mathbf{M} ,

$$\mathbb{X}_{nn}^{\mathbf{dDiv}}(\mathcal{T}) := \mathbb{X}^{\mathbf{dDiv}, nnc}(\mathcal{T}) \cap H_{nn}(\mathbf{ddiv}, \mathcal{T}; \mathbb{S}).$$

In our abstract notation, the approximation spaces are

$$\tilde{H}_h(A^*, \mathcal{T}) := \mathbb{X}_{nn}^{\mathbf{dDiv}}(\mathcal{T}), \quad H_h(\mathcal{T}) := P^1(\mathcal{T}), \quad H_h(A, \mathcal{S}) := HCT_0^{2,1}(\mathcal{S}).$$

They have dimensions $|\mathcal{E}| + 9|\mathcal{T}|$, $3|\mathcal{T}|$ and $3|\mathcal{V}(\Omega)|$, respectively.

The resulting normal-normal continuous mixed scheme reads as follows. Find $\mathbf{M}_h \in \mathbb{X}_{nn}^{\mathbf{dDiv}}(\mathcal{T})$, $u_h \in P^1(\mathcal{T})$, and $\psi_h \in HCT_0^{2,1}(\mathcal{S})$ such that

$$(\mathcal{C}^{-1} \mathbf{M}_h, \delta \mathbf{M}) - (u_h, \mathbf{div} \mathbf{div} \delta \mathbf{M})_{\mathcal{T}} - \langle \psi_h, \delta \mathbf{M} \rangle_{\mathcal{S}} = 0, \quad (49a)$$

$$-(\mathbf{div} \mathbf{div} \mathbf{M}_h, \delta u)_{\mathcal{T}} - \langle \delta \psi, \mathbf{M}_h \rangle_{\mathcal{S}} = -(f, \delta u) \quad (49b)$$

holds for any $\delta \mathbf{M} \in \mathbb{X}_{nn}^{\mathbf{dDiv}}(\mathcal{T})$, $\delta u \in P^1(\mathcal{T})$, and $\delta \psi \in HCT_0^{2,1}(\mathcal{S})$.

It converges quasi-optimally.

Theorem 25. *Let $f \in L_2(\Omega)$ be given. System (49) is well posed. Its solution $(\mathbf{M}_h, u_h, \psi_h)$ satisfies,*

$$\|\mathbf{M} - \mathbf{M}_h\|_{\mathbf{ddiv}, \mathcal{T}} + \|u - u_h\| + \|\psi - \psi_h\|_{3/2, \mathcal{S}} \leq C \left(\|\mathbf{M} - \mathbf{Q}\|_{\mathbf{ddiv}, \mathcal{T}} + \|u - v\| + \|\psi - \phi\|_{3/2, \mathcal{S}} \right)$$

for any $\mathbf{Q} \in \mathbb{X}_{nn}^{\mathbf{dDiv}}(\mathcal{T})$, $v \in P^1(\mathcal{T})$, and $\phi \in HCT_0^{2,1}(\mathcal{S})$. Here, (\mathbf{M}, u, ψ) is the solution of (47) and $C > 0$ is independent of \mathcal{T} , \mathbf{Q} , v , and ϕ .

The proof of Theorem 19 applies in this case as well. In fact, Fortin operator component \mathcal{F}_1 from (46a) maps to $H(\mathbf{ddiv}, \Omega; \mathbb{S}) \subset H_{nn}(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$ and component \mathcal{F}_2 from (46b) maps to $H_{nn}(\mathbf{ddiv}, \mathcal{T}; \mathbb{S})$ since it sets degrees of freedom (41b) to zero.

3.7 Numerical experiments

We consider a simple example of problem (18) with domain $\Omega = (0, 1)^2$ and polynomial solution $u(x_1, x_2) = x_1^2(1 - x_1)^2 x_2^2(1 - x_2)^2$. Using the identity tensor for \mathcal{C} , the bending moments and right-hand side function are $\mathbf{M} = D^2 u$ and $f = \mathbf{div} \mathbf{div} \mathbf{M} = \Delta^2 u$, respectively. We use the discretization spaces as specified in the respective sections, with uniform meshes of size $h := N^{-1/2}$ where $N := |\mathcal{T}|$. We present results for the nodal-continuous and continuous primal hybrid methods (27), (30), and the mixed hybrid and normal-normal continuous mixed methods (40), (49). In all the cases, the domain bilinear forms are calculated analytically on a reference element

and Piola–Kirchhoff transformation (with appropriate scalings) onto elements, cf. [40, 19]. We use numerical integration for the right-hand side entries $(f, \delta u)_T$ (7-point Gauss) and element error calculation (16-point Gauss), cf. [35]. We approximate the skeleton bilinear forms by the 5-point Gauss formula on every edge, and use central differences for the derivatives of the effective shear force (normal component of $\mathbf{div} \mathbf{M}$ and tangential derivative of $\mathbf{t} \cdot \mathbf{Mn}$), required for schemes (40) and (49).

All the discretizations aim at lowest-order approximations, and this is confirmed by our numerical results, with some superconverging components that we do not analyze here. Figure 1 shows the errors $\|u - u_h\|$ (“u”) and $\|D^2(u - u_h)\|_{\mathcal{T}}$ (“D²u”) for the nodal-continuous primal hybrid method (27) along with curves of orders $O(h) = O(N^{-1/2})$ and $O(h^2)$, indicating $\|u - u_h\|_{2,\mathcal{T}} = O(h)$ and superconvergence $\|u - u_h\| = O(h^2)$. For illustration we also plot weighted $L_2(\mathcal{S})$ -errors for the approximations of the traces of \mathbf{M} on the skeleton. Let us indicate by ψ_h^{nn} and ψ_h^{sf} the components of $\boldsymbol{\psi}_h$ that correspond to the normal-normal trace and the effective shear force, respectively. Curves “Mnn” and “shear” present the errors $\|h_S^{1/2}(\mathbf{n} \cdot \mathbf{Mn} - \psi_h^{nn})\|_{\mathcal{S}}$ and $\|h_S^{3/2}(\mathbf{n} \cdot \mathbf{div} \mathbf{M} + \partial_t(\mathbf{t} \cdot \mathbf{Mn}) - \psi_h^{sf})\|_{\mathcal{S}}$. Here, $h_S|_E := |E|$ for $E \in \mathcal{E}$. Both curves are of order $O(h)$. The weightings are chosen to have the respective scalings of the edge-wise $H^{-1/2}$ and $H^{-3/2}$ norms, cf. [46]. We conclude that the results indicate convergence $\|\gamma_{\text{dDiv}, J_0}(\mathbf{M}) - \boldsymbol{\psi}_h\|_{-3/2, -1/2, J_0, \mathcal{S}} = O(h)$. A numerical confirmation of this result would require to construct appropriate extensions of $\boldsymbol{\psi}_h$ to elements of $H(\mathbf{ddiv}, \Omega; \mathbb{S})$ and the calculation of the error in this norm, cf. (24). For the relevance in applications of the traces of \mathbf{M} , in particular of the effective shear force, we present their approximations on the left in Figures 2 (approximation of $\mathbf{n} \cdot \mathbf{Mn}$) and 3 (absolute value of approximation of effective shear force). In both cases, the right figures show (with the same scale as on the respective left side) their difference with the piecewise-constant L_2 -projections of the exact values (absolute difference in the latter case). In this example with smooth solution, we observe point-wise convergence of the $\mathbf{n} \cdot \mathbf{Mn}$ approximation, and point-wise control of the approximation of the effective shear force, essentially an $H^{-3/2}$ -functional. In the case of the continuous primal hybrid method (30), the results are shown in Figure 4 and are analogous. In this case, the trace variable provides only an approximation of $\mathbf{n} \cdot \mathbf{Mn}$, with behavior as before.

Figures 5 and 6 present the results for the mixed hybrid and the normal-normal continuous mixed methods, respectively. We show the curves for the errors $\|u - u_h\|$ (“u”), $\|\mathbf{M} - \mathbf{M}_h\|$ (“M”), $\|\mathbf{div} \mathbf{div}(\mathbf{M} - \mathbf{M}_h)\|_{\mathcal{T}}$ (“divDiv M”) and $\|D^2u - \varepsilon(G_h)\|$ (“D²u”), along with lines indicating $O(h)$ and $O(h^2)$. Here, G_h is the $P^1(\mathcal{T}; \mathbb{R}^2)$ approximation of ∇u , given explicitly by the gradient unknowns of trace approximation $\boldsymbol{\psi}_h$ (in the case of scheme (40)) or ψ_h (in the case of scheme (49)). The numerical results indicate convergence order $O(h)$ of $\|\mathbf{M} - \mathbf{M}_h\|$ and the trace approximations of u in both cases, and increased convergence order $O(h^2)$ of $\|u - u_h\|$ and $\|\mathbf{div} \mathbf{div}(\mathbf{M} - \mathbf{M}_h)\|_{\mathcal{T}}$. We have not shown the superconvergence of u_h but the convergence $\|\mathbf{div} \mathbf{div}(\mathbf{M} - \mathbf{M}_h)\|_{\mathcal{T}} = O(h^2)$ holds by construction, $\mathbf{div}_{\mathcal{T}} \mathbf{div}_{\mathcal{T}} \mathbf{M}_h = \Pi_{\mathcal{T}}^1 f$. We note that the observation $\|D^2u - \varepsilon(G_h)\| = O(h)$ means that the approximation of $\partial_n u|_{\mathcal{S}}$ in a skeleton space $H^{1/2}(\mathcal{S})$ (normal derivatives on \mathcal{S} of $H^2(\Omega)$ -functions) is of this order. A direct control of the approximation of $u|_{\mathcal{S}}$ in $H^{3/2}(\mathcal{S}) := H^2(\Omega)|_{\mathcal{S}}$ would require to provide an $H^2(\Omega)$ -extension of the corresponding component of $\boldsymbol{\psi}_h$ or ψ_h . There are no simple low-order polynomial elements to do this, which is precisely the reason to use composite HCT-elements for domain-based approximations of $u \in H^2(\Omega)$. For illustration, we have also implemented scheme (49) with the full $\mathbb{X}_h^{\text{dDiv}}$ -element from [38] rather than reduced element $\mathbb{X}_{nn}^{\text{dDiv}}$ for the bending moment \mathbf{M} . The curve labelled as “M(15)” in Figure 6 (15 refers to the dimension of the full element) refers to this case and indicates the improved convergence $\|\mathbf{M} - \mathbf{M}_h\| = O(h^2)$.




Figure 1: Errors for the nodal-continuous primal hybrid method (27). The curves are “u”: $\|u - u_h\|$, “ $D^2 u$ ”: $\|D^2(u - u_h)\|_{\mathcal{T}}$, “Mnn”: $L_2(\mathcal{S})$ -error for normal-normal traces of \mathbf{M} , weighted with $h^{1/2}$, “shear”: $L_2(\mathcal{S})$ -error for effective shear force approximation, weighted with $h^{3/2}$, and curves indicating $O(h)$, $O(h^2)$.




Figure 2: The approximation of trace component $\mathbf{n} \cdot \mathbf{M}\mathbf{n}|_{\mathcal{S}}$ from the nodal-continuous primal hybrid method (27) (on the left) and the difference of p/w constant L^2 -projection of $\mathbf{n} \cdot \mathbf{M}\mathbf{n}|_{\mathcal{S}}$ and its approximation (on the right). The mesh has 8192 elements and 12416 edges.




Figure 3: The approximation of the effective shear force from the nodal-continuous primal hybrid method (27) (absolute values, on the left) and the difference of the p/w constant L_2 -projection of the effective shear force and its approximation (absolute values of the difference, on the right). The mesh has 8192 elements and 12416 edges.




Figure 4: Errors for the continuous primal hybrid method (30). The curves are “u”: $\|u - u_h\|_{2,\mathcal{T}}$, “Mnn”: $L_2(\mathcal{S})$ -error for normal-normal traces of \mathbf{M} , weighted with $h^{1/2}$, and a curve indicating $O(h)$.




Figure 5: Errors for the mixed hybrid method (40). The curves are “u”: $\|u - u_h\|$, “M”: $\|\mathbf{M} - \mathbf{M}_h\|$, “divDiv M”: $\|f - \operatorname{div} \mathbf{div} \mathbf{M}_h\|_{\mathcal{T}}$, “D²u”: the L_2 -error of the approximation of the Hessian induced by ψ_h , along with curves indicating orders $O(h)$ and $O(h^2)$.




Figure 6: Errors for the normal-normal continuous mixed method (49). The curves are “u”: $\|u - u_h\|$, “M”: $\|\mathbf{M} - \mathbf{M}_h\|$, “M(15)”: L_2 -error of \mathbf{M} with $\mathbb{X}^{\operatorname{dDiv}}(\mathcal{T})$ -approximation, “divDiv M”: $\|f - \operatorname{div} \mathbf{div} \mathbf{M}_h\|_{\mathcal{T}}$, “D²u”: the L_2 -error of the approximation of the Hessian induced by ψ_h , along with curves indicating orders $O(h)$ and $O(h^2)$.

4 Proofs of abstract results

We start with some preliminary results before proving Theorems 1, 3, Proposition 4, and Theorem 6 at the end of this section.

The next statement is known for special cases, see, e.g., [15, (IV.1.43)], [10, (10.2.22)], [18], [40, Propositions 3.5, 3.9], [41, Lemma 4], and in particular [30, Lemma A.10] for an abstract version.

Lemma 26. *Any $\phi \in H(A, \mathcal{S})$ and $\psi \in H(A^*, \mathcal{S})$ satisfy $\|\phi\|_{A, \mathcal{S}} = \|\phi\|_{(A^*, \sim, \mathcal{T})^*}$ and $\|\psi\|_{A^*, \mathcal{S}} = \|\psi\|_{(A, \sim, \mathcal{T})^*}$. In particular, $\gamma_{A, \mathcal{S}}$ and $\gamma_{A^*, \mathcal{S}}$ are bounded below (with constant 1) and the trace spaces $H(A, \mathcal{S})$ and $H(A^*, \mathcal{S})$ are closed.*

Proof. We only show the norm relation for $\phi \in H(A, \mathcal{S})$. The proof for $\psi \in H(A^*, \mathcal{S})$ is analogous.

Relation $\|\phi\|_{(A^*, \sim, \mathcal{T})^*} \leq \|\phi\|_{A, \mathcal{S}}$ is due to the Cauchy–Schwarz inequality. In fact, considering $v \in H(A)$ with $\gamma_{A, \mathcal{S}}(v) = \phi$ and bounding

$$\langle \phi, w \rangle_{\mathcal{S}} = (Av, w) - (v, A^*w)_{\mathcal{T}} \leq \|v\|_A \|w\|_{A^*, \mathcal{T}} \quad \forall w \in H(A^*, \mathcal{T}),$$

we find that

$$\|\phi\|_{(A^*, \sim, \mathcal{T})^*} = \sup_{w \in \tilde{H}(A^*, \mathcal{T}), \|w\|_{A^*, \mathcal{T}}=1} \langle \phi, w \rangle_{\mathcal{S}} \leq \|v\|_A.$$

Taking the infimum with respect to $v \in H(A)$ subject to $\gamma_{A, \mathcal{S}}(v) = \phi$ gives the result.

Let $\phi \in H(A, \mathcal{S})$ given. It remains to show the inequality $\|\phi\|_{A, \mathcal{S}} \leq \|\phi\|_{(A^*, \sim, \mathcal{T})^*}$. To this end we first define $w \in \tilde{H}(A^*, \mathcal{T})$ by

$$(A^*w, A^*\delta w)_{\mathcal{T}} + (w, \delta w) = -\langle \phi, \delta w \rangle_{\mathcal{S}} \quad \forall \delta w \in \tilde{H}(A^*, \mathcal{T}) \quad (50)$$

and then $v \in H(A)$ by

$$(Av, A\delta v) + (v, \delta v) = -\langle \gamma_{A, \mathcal{S}}(\delta v), w \rangle_{\mathcal{S}} \quad \forall \delta v \in H(A). \quad (51)$$

We establish some relations between v , w , and ϕ .

1. Function v satisfies $v = A_{\mathcal{T}}^*w$. To show this, let $\tilde{v} := A_{\mathcal{T}}^*w$. Relation (50) means that $AA_{\mathcal{T}}^*w = -w$ in the distributional sense so that $\tilde{v} \in H(A)$. Then,

$$(A\tilde{v}, A\delta v) + (\tilde{v}, \delta v) = -(w, A\delta v) + (A^*w, \delta v)_{\mathcal{T}} = -\langle \gamma_{A, \mathcal{S}}(\delta v), w \rangle_{\mathcal{S}} \quad \forall \delta v \in H(A),$$

that is, $\tilde{v} = v$ is the solution to (51).

2. Function v has trace $\gamma_{A, \mathcal{S}}(v) = \phi$. This follows with the previously seen relations and (50), calculating

$$\langle \gamma_{A, \mathcal{S}}(v), \delta w \rangle_{\mathcal{S}} = (Av, \delta w) - (v, A^*\delta w)_{\mathcal{T}} = -(w, \delta w) - (A^*w, A^*\delta w)_{\mathcal{T}} = \langle \phi, \delta w \rangle_{\mathcal{S}}$$

for any $\delta w \in H(A^*, \mathcal{T})$.

3. Function v has norm $\|v\|_A = \|\phi\|_{A, \mathcal{S}}$. This is due to the definition of v , being the minimum energy extension of its trace.

We conclude the proof by setting $\delta w := w$ in (50) and $\delta v := v$ in (51) to find that

$$\|w\|_{A^*, \mathcal{T}}^2 = -\langle \phi, w \rangle_{\mathcal{S}} = \|v\|_A^2$$

so that

$$\|\phi\|_{A, \mathcal{S}} = \|v\|_A = \frac{\langle \phi, w \rangle_{\mathcal{S}}}{\|w\|_{A^*, \mathcal{T}}} \leq \|\phi\|_{(A^*, \sim, \mathcal{T})^*}.$$

□

For a variant of the next statement we refer to [30, Lemma A.9]. Specific versions have been considered, e.g., in [15, Propositions III.1.1, III.1.2], [40, Propositions 3.4(i), 3.8(i)], and [41, Proposition 5].

Lemma 27. *Any $v \in \tilde{H}(A, \mathcal{T})$ and $w \in \tilde{H}(A^*, \mathcal{T})$ satisfy*

$$\begin{aligned} v \in H_0(A) &\Leftrightarrow \langle \psi, v \rangle_{\mathcal{S}} = 0 \quad \forall \psi \in H(A^*, \mathcal{S}), \\ w \in H(A^*) &\Leftrightarrow \langle \phi, w \rangle_{\mathcal{S}} = 0 \quad \forall \phi \in H_0(A, \mathcal{S}). \end{aligned}$$

Proof. We show the statement for $v \in H_0(A)$. The other case is analogous. Any $\psi \in H(A^*, \mathcal{S})$ can be written as $\psi = \gamma_{A^*, \mathcal{S}}(w)$ for a $w \in H(A^*)$. By definition of $\gamma_{A^*, \mathcal{S}}$, $\gamma_{A, \Gamma}$, and $H_0(A)$ we find for any $v \in H_0(A)$ that

$$\langle \psi, v \rangle_{\mathcal{S}} = (A^*w, v) - (w, Av)_{\mathcal{T}} = (A^*w, v) - (w, Av) = -\langle \gamma_{A, \Gamma}(v), w \rangle_{\Gamma} = 0 \quad \forall w \in H(A^*).$$

This shows the direction “ \Rightarrow ”. To see the other direction let $v \in \tilde{H}(A, \mathcal{T})$ be given with $\langle \psi, v \rangle_{\mathcal{S}} = 0$ for any $\psi \in H(A^*, \mathcal{S})$. We calculate Av in the distributional sense,

$$Av(w) = (v, A^*w) = \langle \gamma_{A^*, \mathcal{S}}(w), v \rangle_{\mathcal{S}} + (Av, w)_{\mathcal{T}} = (Av, w)_{\mathcal{T}} \quad \forall w \in C_0^\infty(\Omega; U),$$

and conclude that $Av \in L_2(\Omega)$ so that $v \in H(A)$. To see that $v \in H_0(A)$ we calculate

$$\langle \gamma_{A, \Gamma}(v), w \rangle_{\Gamma} = (Av, w) - (v, A^*w) = (Av, w)_{\mathcal{T}} - (v, A^*w) = \langle \gamma_{A^*, \mathcal{S}}(w), v \rangle_{\mathcal{S}} = 0$$

for any $w \in H(A^*)$ by assumption. This finishes the proof. \square

The next result is [18, Theorem 3.3]. We just translate it to our notation and verify the required assumptions from [18]. For a similar abstract result see [42, Appendix A].

Lemma 28. *Assume that (5b) holds. The bilinear form*

$$b: \begin{cases} \tilde{H}(A^*, \mathcal{T}) \times (L_2(\Omega) \times H_0(A, \mathcal{S})) & \rightarrow \mathbb{R}, \\ (w; v, \phi) & \mapsto (A^*w, v)_{\mathcal{T}} + \langle \phi, w \rangle_{\mathcal{S}} \end{cases}$$

satisfies the inf-sup property

$$\sup_{w \in \tilde{H}(A^*, \mathcal{T}), \|w\|_{A^*, \mathcal{T}}=1} b(w; v, \phi) \geq C(\|v\| + \|\phi\|_{A, \mathcal{S}}) \quad \forall v \in L_2(\Omega), \quad \forall \phi \in H_0(A, \mathcal{S})$$

with a constant $C > 0$ that is independent of v , ϕ , and \mathcal{T} .

Proof. We set

$$\begin{aligned} Y &:= \tilde{H}(A^*, \mathcal{T}), \quad Y_0 := H(A^*), \quad X_0 := L_2(\Omega), \quad \hat{X} := H_0(A, \mathcal{S}), \\ b_0(v, w) &:= (A^*w, v)_{\mathcal{T}}, \quad \hat{b}(\phi, w) := \langle \phi, w \rangle_{\mathcal{S}} \quad \text{for } v \in X_0, \phi \in \hat{X}, w \in Y. \end{aligned}$$

Switching to our notation, Assumptions 3.1 and 3.2 in [18] read as

$$\sup_{0 \neq w \in Y_0} \frac{b_0(v, w)}{\|w\|_Y} := \sup_{0 \neq w \in H(A^*)} \frac{(A^*w, v)_{\mathcal{T}}}{\|w\|_{A^*}} \geq c_0 \|v\| =: c_0 \|v\|_{X_0} \quad \forall v \in X_0 \quad (52)$$

and

$$\begin{aligned} Y_0 &:= H(A^*) = \{w \in \tilde{H}(A^*, \mathcal{T}); \langle \phi, w \rangle_{\mathcal{S}} = 0 \quad \forall \phi \in H_0(A, \mathcal{S})\}, \\ &=: \{w \in Y; \hat{b}(\phi, w) = 0 \quad \forall \phi \in \hat{X}\}, \end{aligned} \quad (53a)$$

$$\sup_{0 \neq w \in Y} \frac{\hat{b}(\phi, w)}{\|w\|_Y} := \sup_{0 \neq w \in \tilde{H}(A^*, \mathcal{T})} \frac{\langle \phi, w \rangle_{\mathcal{S}}}{\|w\|_{A^*, \mathcal{T}}} \geq c \|\phi\|_{A, \mathcal{S}} =: c \|\phi\|_{\hat{X}} \quad \forall \phi \in \hat{X}. \quad (53b)$$

Inf-sup property (52) holds by assumption (5b) with constant $c_0 = c_{\text{is}}$ independent of v (and \mathcal{T}), Lemma 27 proves (53a), and (53b) holds by Lemma 26 with $c = 1$. The statement follows by [18, Theorem 3.3]. \square

4.1 Proof of Theorem 1

Problem (7) is a mixed system that satisfies the usual conditions. In particular, all (bi)linear forms are uniformly bounded and duality $\langle \delta\psi, u \rangle_{\mathcal{S}}$ satisfies the inf-sup condition with constant 1 by Lemma 26. Furthermore, by Lemma 27 we have the kernel representation

$$\{u \in \tilde{H}(A, \mathcal{T}); \langle \delta\psi, u \rangle_{\mathcal{S}} = 0 \ \forall \delta\psi \in H(A^*, \mathcal{S})\} = H_0(A)$$

and the $H_0(A)$ -coercivity of $(\mathcal{C}\cdot, \cdot)$ holds by assumption (5a). This proves the well-posedness of (7). Lemma 27 and relation (7b) imply that $u \in H_0(A)$. Relation (7a) with $\delta u \in C_0^\infty(\Omega)$ and an application of Lemma 27 to conclude that $\langle \psi, \delta u \rangle_{\mathcal{S}} = 0$ for such δu , show that $A^* \mathcal{C}Au = f$. Using this relation together with the definition of $\gamma_{A^*, \mathcal{S}}$, and again (7a), we find that

$$\langle \gamma_{A^*, \mathcal{S}}(\mathcal{C}Au), \delta u \rangle_{\mathcal{S}} = (f, \delta u)_{\mathcal{T}} - (\mathcal{C}Au, A\delta u) = \langle \phi, \delta u \rangle_{\mathcal{S}} \quad \forall \delta u \in \tilde{H}(A, \mathcal{T}),$$

that is, $\phi = \gamma_{A^*, \mathcal{S}}(\mathcal{C}Au)$. This finishes the proof.

4.2 Proof of Theorem 3

Again, problem (10) is a mixed system that satisfies the usual conditions. All (bi)linear forms are uniformly bounded. By Lemma 28, the bilinear form

$$b(z; \delta u, \delta\phi) := (A^*z, \delta u)_{\mathcal{T}} + \langle \delta\phi, z \rangle_{\mathcal{S}}$$

satisfies the inf-sup condition. By Lemma 27 we have the kernel representation

$$\{z \in \tilde{H}(A^*, \mathcal{T}); \langle \delta\phi, z \rangle_{\mathcal{S}} = 0 \ \forall \delta\phi \in H_0(A, \mathcal{S})\} = H(A^*)$$

so that

$$\begin{aligned} \ker(b) &:= \{z \in \tilde{H}(A^*, \mathcal{T}); b(z; \delta u, \delta\phi) = 0 \ \forall \delta u \in L_2(\Omega), \ \forall \delta\phi \in H_0(A, \mathcal{S})\} \\ &= \{z \in H(A^*); A^*z = 0\}. \end{aligned}$$

The coercivity

$$(\mathcal{C}^{-1}z, z) \geq C \|z\|_{A^*, \mathcal{T}}^2 \quad \forall z \in \ker(b)$$

with a constant $C > 0$ independent of z and \mathcal{T} follows. Therefore, (10) is well posed.

Relation $w = \mathcal{C}Au$ follows from (10a) by a distributional argument, also implying that $u \in H(A)$. Then, (10a) shows that

$$\langle \gamma_{A, \mathcal{S}}(u), \delta w \rangle_{\mathcal{S}} = (\delta w, Au) - (A^* \delta w, u)_{\mathcal{T}} = \langle \phi, \delta w \rangle_{\mathcal{S}} \quad \forall \delta w \in \tilde{H}(A^*, \mathcal{T}),$$

that is, $\phi = \gamma_{A, \mathcal{S}}(u)$. This implies that $u \in H_0(A)$. Indeed, since $\phi \in H_0(A, \mathcal{S})$ we find with Lemma 27 that

$$\langle \gamma_{A, \Gamma}(u), \delta w \rangle_{\Gamma} = (Au, \delta w) - (u, A^* \delta w)_{\mathcal{T}} = \langle \gamma_{A, \mathcal{S}}(u), \delta w \rangle_{\mathcal{S}} = \langle \phi, \delta w \rangle_{\mathcal{S}} = 0 \quad \forall \delta w \in H(A^*),$$

that is, $\gamma_{A, \Gamma}(u) = 0$.

By relation (10b) and Lemma 27 we have that $w \in H(A^*)$ and $A^*w = f$. This finishes the proof.

4.3 Proof of Proposition 4

We use inf-sup stability (5b), the properties of operator \mathcal{F}_1 , and relation $\langle \delta\phi, w \rangle_{\mathcal{S}} = 0$ for $\delta\phi \in H_0(A, \mathcal{S})$ and $w \in H(A^*)$ by Lemma 27, to deduce the bound

$$\begin{aligned} c_{\text{is}} \|\delta u\| &\leq \sup_{w \in H(A^*) \setminus \{0\}} \frac{(A^* w, \delta u)}{\|w\|_{A^*}} \leq C_1 \sup_{w_h \in H(A^*) \cap \tilde{H}_h(A^*, \mathcal{T}) \setminus \{0\}} \frac{(A^* w_h, \delta u) + \langle \delta\phi, w_h \rangle_{\mathcal{S}}}{\|w_h\|_{A^*}} \\ &\leq C_1 \sup_{w_h \in \tilde{H}_h(A^*, \mathcal{T}) \setminus \{0\}} \frac{(A^* w_h, \delta u)_{\mathcal{T}} + \langle \delta\phi, w_h \rangle_{\mathcal{S}}}{\|w_h\|_{A^*, \mathcal{T}}} \quad \forall \delta\phi \in H_h(A, \mathcal{S}), \quad \forall \delta u \in H_h(\mathcal{T}). \end{aligned} \quad (54)$$

Lemma 26, the properties of operator \mathcal{F}_2 , and estimate (54) show that any $\delta\phi \in H_h(A, \mathcal{S})$ satisfies

$$\begin{aligned} \|\delta\phi\|_{A, \mathcal{S}} &= \sup_{w \in \tilde{H}(A^*, \mathcal{T}) \setminus \{0\}} \frac{\langle \delta\phi, w \rangle_{\mathcal{S}}}{\|w\|_{A^*, \mathcal{T}}} \leq C_2 \sup_{w_h \in \tilde{H}_h(A^*, \mathcal{T}) \setminus \{0\}} \frac{\langle \delta\phi, w_h \rangle_{\mathcal{S}}}{\|w_h\|_{A^*, \mathcal{T}}} \\ &= C_2 \sup_{w_h \in \tilde{H}_h(A^*, \mathcal{T}) \setminus \{0\}} \left(\frac{(A^* w_h, \delta u)_{\mathcal{T}} + \langle \delta\phi, w_h \rangle_{\mathcal{S}}}{\|w_h\|_{A^*, \mathcal{T}}} - \frac{(A^* w_h, \delta u)_{\mathcal{T}}}{\|w_h\|_{A^*, \mathcal{T}}} \right) \\ &\leq C_2 \sup_{w_h \in \tilde{H}_h(A^*, \mathcal{T}) \setminus \{0\}} \frac{(A^* w_h, \delta u)_{\mathcal{T}} + \langle \delta\phi, w_h \rangle_{\mathcal{S}}}{\|w_h\|_{A^*, \mathcal{T}}} + C_2 \|\delta u\| \\ &\leq C_2 \left(1 + \frac{C_1}{c_{\text{is}}}\right) \sup_{w_h \in \tilde{H}_h(A^*, \mathcal{T}) \setminus \{0\}} \frac{(A^* w_h, \delta u)_{\mathcal{T}} + \langle \delta\phi, w_h \rangle_{\mathcal{S}}}{\|w_h\|_{A^*, \mathcal{T}}}. \end{aligned} \quad (55)$$

A combination of estimates (54) and (55) proves the discrete inf-sup property (13).

4.4 Proof of Theorem 6

System (14) is not of a (standard) mixed form but satisfies the standard properties of an operator equation [51, 5]. As before, all (bi)linear forms are uniformly bounded. Furthermore, the operator

$$\mathcal{B} : \mathcal{U} := L_2(\Omega) \times L_2(\Omega; U) \times H_0(A, \mathcal{S}) \times H(A^*, \mathcal{S}) \rightarrow \mathcal{V}^* := \tilde{H}(A, \mathcal{T})^* \times \tilde{H}(A^*, \mathcal{T})^*$$

defined by the system is injective and satisfies the inf-sup condition, as we briefly recall now.

Injectivity. Let $\delta v = (\delta u, \delta w) \in \mathcal{V}$ satisfy $\mathcal{B}^* \delta v = 0$. Lemma 27 shows that $\delta w \in H(A^*)$ and $\delta u \in H_0(A)$. Then $\mathcal{B}^* \delta v = 0$ implies $A^* \delta w = 0$, $C^{-1} \delta w = A \delta u$. We conclude that $\delta u \in H_0(A)$ solves $A^* C A \delta u = 0$, thus $\delta u = 0$ by (5a), and $\delta w = 0$.

Inf-sup property. In abstract form, we have to show that there is a constant $C > 0$, independent of \mathcal{T} and \mathbf{u} , that satisfies

$$\sup_{v \in \mathcal{V}, \|v\|_{\mathcal{V}}=1} \langle \mathcal{B} \mathbf{u}, v \rangle_{\mathcal{V}^* \times \mathcal{V}} \geq C \|\mathbf{u}\|_{\mathcal{U}} \quad \forall \mathbf{u} \in \mathcal{U}$$

with (squared) norms $\|\mathbf{u}\|_{\mathcal{U}}^2 := \|u\|^2 + \|w\|^2 + \|\phi\|_{A, \mathcal{S}}^2 + \|\psi\|_{A^*, \mathcal{S}}^2$ and $\|v\|_{\mathcal{V}}^2 := \|\delta u\|_{A, \mathcal{T}}^2 + \|\delta w\|_{A^*, \mathcal{T}}^2$ for $\mathbf{u} \in \mathcal{U}$ and $v = (\delta u, \delta w) \in \mathcal{V}$. As Lemma 28, this can be seen by [18, Theorem 3.3]. We set

$$\begin{aligned} Y &:= \mathcal{V}, \quad Y_0 := H_0(A) \times H(A^*), \quad X_0 := L_2(\Omega) \times L_2(\Omega; U), \quad \hat{X} := H_0(A, \mathcal{S}) \times H(A^*, \mathcal{S}), \\ b_0((u, w), (\delta u, \delta w)) &:= (A \delta u - C^{-1} \delta w, w)_{\mathcal{T}} + (A^* \delta w, u)_{\mathcal{T}}, \\ \hat{b}((\phi, \psi), (\delta u, \delta w)) &:= \langle \phi, \delta w \rangle_{\mathcal{S}} + \langle \psi, \delta u \rangle_{\mathcal{S}} \quad \text{for } (u, w) \in X_0, (\phi, \psi) \in \hat{X}, (\delta u, \delta w) \in Y \end{aligned}$$

and need to check conditions (52) and (53) in the current setting. Identity (53a) holds by Lemma 27 and Lemma 26 implies (53b) with $c = 1$. Inf-sup condition (52) follows by the stability of the adjoint problem: *Given $(g, G) \in X_0$ find $(\delta u, \delta w) \in Y$ such that*

$$\begin{aligned} A^* \delta w &= g, & A \delta u - C^{-1} \delta w &= G \\ \Leftrightarrow A^* C A \delta u &= g + A^* C G, & \delta w &= C A \delta u - C G. \end{aligned}$$

Of course, this is the initial (self-adjoint) problem (1) with general data. By Assumption (5a) it is well posed with solution $(\delta u, \delta w) \in H_0(A) \times H(A^*)$ bounded as

$$\begin{aligned}\|\delta u\|_A &\leq C(\|g\|^2 + \|G\|^2)^{1/2} = C\|(g, G)\|_{X_0}, \\ \|\delta w\|_{A^*}^2 &= \|\delta w\|^2 + \|A^* \delta w\|^2 = \|\mathcal{C}A\delta u - \mathcal{C}G\|^2 + \|g\|^2 \leq C^2\|(g, G)\|_{X_0}^2\end{aligned}$$

with a constant $C > 0$ that depends on \mathcal{C} but is independent of g and G . The inf-sup property follows by [18, Theorem 3.3].

We conclude that problem (14) is well posed with solution $(u, w, \phi, \psi) \in \mathcal{U}$ that satisfies the claimed stability estimate. Distributional arguments verify the properties $u \in H(A)$, $w = \mathcal{C}Au \in H(A^*)$, and $A^*w = f$. Then, applying (14a) and (14b), we find, respectively, that

$$\langle \gamma_{A, \mathcal{S}}(u), \delta w \rangle_{\mathcal{S}} = (Au, \delta w) - (u, A^* \delta w)_{\mathcal{T}} = (\mathcal{C}^{-1}w, \delta w) - (u, A^* \delta w)_{\mathcal{T}} = \langle \phi, \delta w \rangle_{\mathcal{S}}$$

for any $\delta w \in \tilde{H}(A^*, \mathcal{T})$ and

$$\langle \gamma_{A^*, \mathcal{S}}(w), \delta u \rangle_{\mathcal{S}} = (A^*w, \delta u) - (w, A\delta u)_{\mathcal{T}} = (A^*w, \delta u) - (f, \delta u) + \langle \psi, \delta u \rangle_{\mathcal{S}} = \langle \psi, \delta u \rangle_{\mathcal{S}}$$

for any $\delta u \in \tilde{H}(A, \mathcal{T})$. It follows that $\gamma_{A, \mathcal{S}}(u) = \phi$, also implying $u \in H_0(A)$, and $\gamma_{A^*, \mathcal{S}}(w) = \psi$. This finishes the proof.

References

- [1] R. A. ADAMS, *Sobolev spaces*, Academic Press, New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- [2] D. N. ARNOLD AND F. BREZZI, *Mixed and nonconforming finite element methods: implementation, postprocessing and error estimates*, RAIRO Modél. Math. Anal. Numér., 19 (1985), pp. 7–32.
- [3] D. N. ARNOLD, F. BREZZI, B. COCKBURN, AND L. D. MARINI, *Unified analysis of discontinuous Galerkin methods for elliptic problems*, SIAM J. Numer. Anal., 39 (2001/02), pp. 1749–1779.
- [4] I. BABUŠKA, *The finite element method with Lagrangian multipliers*, Numer. Math., 20 (1972/73), pp. 179–192.
- [5] I. BABUŠKA, *Error-bounds for finite element method*, Numer. Math., 16 (1970/71), pp. 322–333.
- [6] I. BABUŠKA AND J. PITKÄRANTA, *The plate paradox for hard and soft simple support*, SIAM J. Math. Anal., 21 (1990), pp. 551–576.
- [7] I. BABUŠKA AND M. ZLÁMAL, *Nonconforming elements in the finite element method with penalty*, SIAM J. Numer. Anal., 10 (1973), pp. 863–875.
- [8] G. P. BAZELEY, Y. K. CHEUNG, B. M. IRONS, AND O. C. ZIENKIEWICZ, *Triangular elements in plate bending – conforming and non-conforming solutions*, in Proceedings of the Conference on Matrix Methods in Structural Mechanics, J. S. Przemieniecki, R. M. Bader, W. F. Bozich, J. R. Johnson, and W. J. Mykytow, eds., Wright–Patterson A.F.B., Ohio, 1965, pp. 547–576. AFFDL Report No. TR-66-80, <https://contrails.library.iit.edu/item/160952>.

- [9] H. BLUM AND R. RANNACHER, *On mixed finite element methods in plate bending analysis. Part 1: The first Herrmann scheme*, *Comput. Mech.*, 6 (1990), pp. 221–236.
- [10] D. BOFFI, F. BREZZI, AND M. FORTIN, *Mixed finite element methods and applications*, vol. 44 of Springer Series in Computational Mathematics, Springer, Heidelberg, 2013.
- [11] C. L. BOTTASSO, S. MICHELETTI, AND R. SACCO, *The discontinuous Petrov-Galerkin method for elliptic problems*, *Comput. Methods Appl. Mech. Engrg.*, 191 (2002), pp. 3391–3409.
- [12] S. C. BRENNER AND L.-Y. SUNG, *C^0 interior penalty methods for fourth order elliptic boundary value problems on polygonal domains*, *J. Sci. Comput.*, 22/23 (2005), pp. 83–118.
- [13] F. BREZZI, *On the existence, uniqueness and approximation of saddle-point problems arising from Lagrangian multipliers*, *Rev. Francaise Automat. Informat. Recherche Opérationnelle Sér. Rouge*, 8 (1974), pp. 129–151.
- [14] ———, *Sur la méthode des éléments finis hybrides pour le problème biharmonique*, *Numer. Math.*, 24 (1975), pp. 103–131.
- [15] F. BREZZI AND M. FORTIN, *Mixed and hybrid finite element methods*, vol. 15 of Springer Series in Computational Mathematics, Springer-Verlag, New York, 1991.
- [16] F. BREZZI AND L. D. MARINI, *On the numerical solution of plate bending problems by hybrid methods*, *Rev. Francaise Automat. Informat. Recherche Opérationnelle Sér. Rouge*, 9 (1975), pp. 5–50.
- [17] D. BROERSEN AND R. STEVENSON, *A robust Petrov–Galerkin discretisation of convection-diffusion equations*, *Comput. Math. Appl.*, 68 (2014), pp. 1605–1618.
- [18] C. CARSTENSEN, L. F. DEMKOWICZ, AND J. GOPALAKRISHNAN, *Breaking spaces and forms for the DPG method and applications including Maxwell equations*, *Comput. Math. Appl.*, 72 (2016), pp. 494–522.
- [19] C. CARSTENSEN AND N. HEUER, *Normal-normal continuous symmetric stresses in mixed finite element elasticity*. Submitted for publication.
- [20] O. CESSENAT AND B. DESPRÉS, *Application of an ultra weak variational formulation of elliptic PDEs to the two-dimensional Helmholtz problem*, *SIAM J. Numer. Anal.*, 35 (1998), pp. 255–299.
- [21] L. CHEN AND X. HUANG, *A new div-div-conforming symmetric tensor finite element space with applications to the biharmonic equation*, *Math. Comp.* <https://doi.org/10.1090/mcom/3957>.
- [22] P. G. CIARLET, *The Finite Element Method for Elliptic Problems*, North-Holland, Amsterdam, 1978.
- [23] P. G. CIARLET, *Interpolation error estimates for the reduced Hsieh-Clough-Tocher triangle*, *Math. Comp.*, 32 (1978), pp. 335–344.
- [24] R. W. CLOUGH AND J. L. TOCHER, *Finite element stiffness matrices for analysis of plate bending*, in *Proceedings of the Conference on Matrix Methods in Structural Mechanics*, J. S. Przemieniecki, R. M. Bader, W. F. Bozich, J. R. Johnson, and W. J. Mykytow, eds., Wright–Patterson A.F.B., Ohio, 1965, pp. 515–545. AFFDL Report No. TR-66-80, <https://contrails.library.iit.edu/item/160951>.

- [25] A. COHEN, W. DAHMEN, AND G. WELPER, *Adaptivity and variational stabilization for convection-diffusion equations*, ESAIM Math. Model. Numer. Anal., 46 (2012), pp. 1247–1273.
- [26] W. DAHMEN, C. HUANG, C. SCHWAB, AND G. WELPER, *Adaptive Petrov-Galerkin methods for first order transport equations*, SIAM J. Numer. Anal., 50 (2012), pp. 2420–2445.
- [27] L. F. DEMKOWICZ AND J. GOPALAKRISHNAN, *A class of discontinuous Petrov-Galerkin methods. Part I: the transport equation*, Comput. Methods Appl. Mech. Engrg., 199 (2010), pp. 1558–1572.
- [28] ———, *A class of discontinuous Petrov-Galerkin methods. Part II: Optimal test functions*, Numer. Methods Partial Differential Eq., 27 (2011), pp. 70–105.
- [29] ———, *An overview of the discontinuous Petrov Galerkin method*, in Recent developments in discontinuous Galerkin finite element methods for partial differential equations, vol. 157 of IMA Vol. Math. Appl., Springer, Cham, 2014, pp. 149–180.
- [30] L. F. DEMKOWICZ, J. GOPALAKRISHNAN, S. NAGARAJ, AND P. SEPÚLVEDA, *A spacetime DPG method for the Schrödinger equation*, SIAM J. Numer. Anal., 55 (2017), pp. 1740–1759.
- [31] B. DESPRÉS, *Sur une formulation variationnelle de type ultra-faible*, C. R. Acad. Sci. Paris Sér. I Math., 318 (1994), pp. 939–944.
- [32] D. A. DI PIETRO AND A. ERN, *Mathematical aspects of discontinuous Galerkin methods*, vol. 69 of Mathématiques & Applications (Berlin), Springer, Heidelberg, 2012.
- [33] Z. DONG AND A. ERN, *C^0 -hybrid high-order methods for biharmonic problems*, IMA J. Numer. Anal., 44 (2024), pp. 24–57.
- [34] J. DOUGLAS, JR., T. DUPONT, P. PERCELL, AND R. SCOTT, *A family of C^1 finite elements with optimal approximation properties for various Galerkin methods for 2nd and 4th order problems*, RAIRO Anal. Numér., 13 (1979), pp. 227–255.
- [35] D. A. DUNAVANT, *High degree efficient symmetrical Gaussian quadrature rules for the triangle*, Internat. J. Numer. Methods Engrg., 21 (1985), pp. 1129–1148.
- [36] G. ENGEL, K. GARIKIPATI, T. J. R. HUGHES, M. G. LARSON, L. MAZZEI, AND R. L. TAYLOR, *Continuous/discontinuous finite element approximations of fourth-order elliptic problems in structural and continuum mechanics with applications to thin beams and plates, and strain gradient elasticity*, Comput. Methods Appl. Mech. Engrg., 191 (2002), pp. 3669–3750.
- [37] B. FRAEIJIS DE VEUBEKE, *Displacement and equilibrium models in the finite element method*, in Stress analysis, O. C. Zienkiewicz and G. S. Holister, eds., John Wiley & Sons, 1965, ch. 9, pp. 145–197. Reprint in *Internat. J. Numer. Methods Engrg.*, 52 (3), 2001, pp. 287–342.
- [38] T. FÜHRER AND N. HEUER, *Mixed finite elements for Kirchhoff–Love plate bending*, Math. Comp. <https://doi.org/10.1090/mcom/3995>, arxiv preprint 2305.08693.
- [39] ———, *Fully discrete DPG methods for the Kirchhoff–Love plate bending model*, Comput. Methods Appl. Mech. Engrg., 343 (2019), pp. 550–571.
- [40] T. FÜHRER, N. HEUER, AND A. H. NIEMI, *An ultraweak formulation of the Kirchhoff–Love plate bending model and DPG approximation*, Math. Comp., 88 (2019), pp. 1587–1619.

- [41] T. FÜHRER, N. HEUER, AND F.-J. SAYAS, *An ultraweak formulation of the Reissner–Mindlin plate bending model and DPG approximation*, Numer. Math., 145 (2020), pp. 313–344.
- [42] V. V. GARG, S. PRUDHOMME, K. VAN DER ZEE, AND G. F. CAREY, *Adjoint-consistent formulations of slip models for coupled electroosmotic flow systems*, Adv. Model. Simul. Eng., 2 (2014). Article 15.
- [43] J. GOPALAKRISHNAN AND W. QIU, *An analysis of the practical DPG method*, Math. Comp., 83 (2014), pp. 537–552.
- [44] K. HELLAN, *Analysis of elastic plates in flexure by a simplified finite element method*, Acta Polytech. Scand. Civ. Eng. Build. Constr. Ser. 46, 1 (1967).
- [45] L. R. HERRMANN, *Finite-element bending analysis for plates*, J. Eng. Mech. Div., 93 (1967), pp. 13–26.
- [46] N. HEUER, *On the equivalence of fractional-order Sobolev semi-norms*, J. Math. Anal. Appl., 417 (2014), pp. 505–518.
- [47] C. JOHNSON, *On the convergence of a mixed finite-element method for plate bending problems*, Numer. Math., 21 (1973), pp. 43–62.
- [48] O. A. LADYZHENSKAYA, *The mathematical theory of viscous incompressible flow*, vol. Vol. 2 of Mathematics and its Applications, Gordon and Breach Science Publishers, New York–London–Paris, English ed., 1969.
- [49] P. LASCAUX AND P. LESAINT, *Some nonconforming finite elements for the plate bending problem*, Rev. Française Automat. Informat. Recherche Operationnelle Sér. Rouge Anal. Numér., 9 (1975), pp. 9–53.
- [50] L. S. D. MORLEY, *The triangular equilibrium element in the solutions of plate bending problem*, Aero. Quart., 19 (1968), pp. 149–169.
- [51] J. NEČAS, *Les méthodes directes en théorie des équations elliptiques*, Masson et Cie, Éditeurs, Paris; Academia, Éditeurs, Prague, 1967.
- [52] T. H. H. PIAN, *Finite element formulation by variational principles with relaxed continuity requirements*, in The mathematical foundations of the finite element method with applications to partial differential equations (Proc. Sympos., Univ. Maryland, Baltimore, Md., 1972), Academic Press, New York-London, 1972, pp. 671–687.
- [53] T. H. H. PIAN AND P. TONG, *Basis of finite element methods for solid continua*, Internat. J. Numer. Methods Engrg., 1 (1969), pp. 3–28.
- [54] P.-A. RAVIART AND J.-M. THOMAS, *Primal hybrid finite element methods for 2nd order elliptic equations*, Math. Comp., 31 (1977), pp. 391–413.
- [55] B. RIVIÈRE, *Discontinuous Galerkin methods for solving elliptic and parabolic equations*, vol. 35 of Frontiers in Applied Mathematics, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2008.
- [56] J. M. THOMAS, *Méthode des éléments finis équilibre*, in Journées “Éléments Finis” (Rennes, 1975), Univ. Rennes, Rennes, 1975.

- [57] —, *Méthode des éléments finis hybrides d'aux pour les problèmes elliptiques du 2nd ordre*, Rev. Française Automat. Informat. Recherche Operationnelle Sér. Rouge Anal. Numér., 10 (1976), pp. 51–70.