

1-COCYCLES OF THE WITT ALGEBRA WITH COEFFICIENTS IN TENSOR PRODUCT OF MODULES

SHOULAN GAO, DONG LIU, AND YUFENG PEI*

ABSTRACT. In this paper, we classify 1-cocycles of the Witt algebra with coefficients in the tensor product of two arbitrary tensor density modules. In a special case, we recover a theorem originally established by Ng and Taft in [24]. Furthermore, by these 1-cocycles, we determine Lie bialgebra structures over certain infinite-dimensional Lie algebras containing the Witt algebra.

1. INTRODUCTION

The Witt algebra \mathcal{W} is an infinite-dimensional Lie algebra that emerges as the algebra of derivations acting on the Laurent polynomial ring in a single variable. The Virasoro algebra, which is the universal central extension of the Witt algebra, holds significant importance in the realms of both mathematics and theoretical physics. The cohomology of the Witt algebra and the Virasoro algebra has been an active area of research owing to its deep relations with representation theory [10, 16, 21, 20]. In particular, the algebraic cohomology of the Witt algebra, when considering coefficients related to the tensor density modules \mathcal{F}_α , has been the subject of study in various works [5, 6, 7, 11, 14, 25, 29]. In [10], Feigin and Fuchs achieved a highly non-trivial classification of length two extensions for the Witt algebra using the first algebraic cohomology, with coefficients in the module of the space of homomorphisms between the two arbitrary tensor density modules. Recently, in [4], Dykes computed all possible 1-cocycles for the Witt algebra by employing a compatible action of the commutative algebra of Laurent polynomials in one variable.

Motivated by Witten's study of left-invariant symplectic structures on the Virasoro group ([31]), Ng and Taft established that all Lie bialgebra structures on the Witt algebra and the Virasoro algebra are triangular coboundary through computation of 1-cocycles with coefficients in the tensor product of the adjoint modules in [24] (see also [17]). The Lie bialgebra structures on various infinite-dimensional Lie algebras related to the Witt algebra have been investigated by computing the first algebraic cohomology [8, 15, 19, 18, 27]. As mentioned in [19], a key and complicated step of characterizing these Lie bialgebras is dealing with 1-cocycles of the Witt algebra with coefficients in the tensor product of two tensor density modules. To the best knowledge of the authors, there is no general formula for these 1-cocycles in the literature.

2020 *Mathematics Subject Classification.* 17B05, 17B62, 17B65, 17B68.

Key words and phrases. Witt algebra, 1-cocycles, Lie bialgebra.

*: Corresponding author.

This paper aims to classify 1-cocycles of the Witt algebra with coefficients in the tensor product of two arbitrary tensor density modules. Our main theorem can be stated as follows:

Main Theorem (Theorem 5.7): For $\alpha, \beta \in \mathbb{C}$, we have

$$\dim H^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta) = \begin{cases} 5, & (\alpha, \beta) = (0, 0); \\ 2, & (\alpha, \beta) = (0, 1) \text{ or } (1, 0); \\ 1, & (\alpha, \beta) = (0, 2), (2, 0), (1, 1), \text{ or } \alpha + \beta = 1, \beta \neq 0, 1; \\ 0, & \text{otherwise.} \end{cases}$$

Establishing the proof for this theorem is challenging and involves numerous calculations. Nonetheless, the fundamental concept underlying the proof remains clear. First, suitable normalizations allow selecting well-defined representatives for each cohomology class, reducing the computation of $H^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)$ to that of normalized 1-cocycles. Through several technical lemmas, we then demonstrate a normalized 1-cocycle must assume a specialized form, enabling an upper bound on $\dim H^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)$. Finally, exhibiting linearly independent normalized 1-cocycles produces the result.

The interest of this classification for these 1-cocycles lies in the applications to the Lie bialgebras associated with some infinite dimensional Lie algebras. When $(\alpha, \beta) = (-1, -1)$, we recover a theorem first proven by Ng and Taft in [24]. Furthermore, as consequence of the Main Theorem, we establish Lie bialgebra structures over various infinite-dimensional Lie algebras that contain the Witt algebra, such as the Oveshenko-Roger algebra [25], the twisted Heisenberg-Virasoro algebra [1], the BMS algebra [2, 32], and the Schrödinger-Virasoro algebra [26].

It is worth noting that, as stated in [10], an isomorphism exists: $H^1(\mathcal{W}, \text{Hom}(\mathcal{F}_\alpha, \mathcal{F}_\beta)) \cong H_1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}'_\beta)$, where \mathcal{F}'_β denotes the graded dual modules of \mathcal{F}_β . Our main theorem may be considered as a dual counterpart to the 1-cocycles presented in [10]. However, there is currently no more efficient method for computing 1-cocycles via 1-cycles.

The paper is organized as follows. In Section 2, we introduce the notations and conventions used throughout the paper, we also present several technical lemmas on 1-cocycles of degree zero that will be useful later. In Section 3, we construct several 1-cocycles of the Witt algebra. Section 4 contains the proof of Theorem 5.7 for the case $(\alpha, \beta) \notin \mathbb{Z}^2$, while Section 5 handles the case $(\alpha, \beta) \in \mathbb{Z}^2$. Finally, in Section 6 we apply Theorem 5.7 to investigate several infinite-dimensional Lie bialgebras containing \mathcal{W} .

Throughout the paper, we use $\mathbb{C}, \mathbb{C}^*, \mathbb{Z}, \mathbb{Z}_+, \mathbb{N}$, and \mathbb{Z}^* to denote the sets of complex numbers, non-zero complex numbers, integers, positive integers, non-negative integers, and non-zero integers, respectively.

2. PRELIMINARIES AND NOTATIONS

In this section, we review some preliminaries and notations on 1-cocycles of the Witt algebra and obtain several useful lemmas.

Definition 2.1. *The Witt algebra \mathcal{W} is an infinite-dimensional Lie algebra spanned by $\{L_n \mid n \in \mathbb{Z}\}$ with the bracket*

$$[L_m, L_n] = (m - n)L_{m+n}, \quad \forall m, n \in \mathbb{Z}.$$

Definition 2.2. *For $\alpha \in \mathbb{C}$, the tensor density module $\mathcal{F}_\alpha = \bigoplus_{n \in \mathbb{Z}} \mathbb{C}v_n$ for \mathcal{W} is given by*

$$L_m \cdot v_n = -(\alpha m + n)v_{m+n}, \quad \forall m, n \in \mathbb{Z}.$$

For $\alpha, \beta \in \mathbb{C}$, $\mathcal{F}_\alpha \otimes \mathcal{F}_\beta$ is a \mathcal{W} -module with

$$L_m \cdot (v_i \otimes v_j) = -(i + \alpha m)v_{m+i} \otimes v_j - (j + \beta m)v_i \otimes v_{m+j}, \quad \forall m, i, j \in \mathbb{Z}.$$

In particular,

$$L_0 \cdot (v_i \otimes v_j) = -(i + j)v_i \otimes v_j, \quad \forall i, j \in \mathbb{Z}.$$

It follows that $\mathcal{F}_\alpha \otimes \mathcal{F}_\beta$ is equipped with a \mathbb{Z} -grading:

$$\mathcal{F}_\alpha \otimes \mathcal{F}_\beta = \bigoplus_{k \in \mathbb{Z}} (\mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_k, \quad \text{where} \quad (\mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_k = \bigoplus_{i \in \mathbb{Z}} \mathbb{C}v_i \otimes v_{k-i}.$$

Definition 2.3 ([30]). *Let L be a Lie algebra and M an L -module. Denote by $\text{Der}(L, M)$ the set of 1-cocycles (derivations) $d : L \rightarrow M$ with coefficients in M , which are linear maps satisfying the condition $d([x, y]) = x \cdot d(y) - y \cdot d(x)$ for $x, y \in L$. The set $\text{Inn}(L, M)$ consists of the 1-cocycles d_v for some $v \in M$, where d_v is called 1-coboundary (inner derivation) with coefficients in M defined by $d_v : x \mapsto x \cdot v$ for $x \in L$. A 1-cocycle with coefficients in M is called non-trivial if it is not a 1-coboundary with coefficients in M . We denote by $H^1(L, M) = \text{Der}(L, M)/\text{Inn}(L, M)$ the first algebraic cohomology of L with coefficients in M .*

Remark 2.4. It is a well-known fact [12] that there exists a one-to-one correspondence between elements in $H^1(L, M)$ and equivalence classes of extensions of L -modules by M .

Definition 2.5. *Let G be an abelian group, $L = \bigoplus_{g \in G} L_g$ a G -graded Lie algebra over \mathbb{C} . An L -module V is said to be G -graded if $V = \bigoplus_{g \in G} V_g$ and $L_g \cdot V_h \subset V_{g+h}$ for all $g, h \in G$.*

Proposition 2.6 ([9]). *Assume that L is finitely generated as a G -graded Lie algebra. Let V be a G -graded L -module. Then*

$$\text{Der}(L, V) = \bigoplus_{g \in G} \text{Der}(L, V)_g.$$

Lemma 2.7. *The derivation algebra $\text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)$ is \mathbb{Z} -graded, i.e.,*

$$\text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta) = \bigoplus_{k \in \mathbb{Z}} \text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_k, \quad D_k(L_m) \subseteq (\mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_{m+k}$$

for any $m \in \mathbb{Z}$, $D_k \in \text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_k$.

Proof. According to Proposition 2.6, coupled with the knowledge that the Witt algebra \mathcal{W} is both finitely generated and \mathbb{Z} -graded. □

Lemma 2.8. $\text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta) = \text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 + \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)$.

Proof. Assuming $k \neq 0$ and $D_k \in \text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_k$, we have $D_k(L_0) = \sum_{r=0}^s a(i_r)v_{i_r} \otimes v_{k-i_r}$, where $a(i_r) \in \mathbb{C}$ and $s \in \mathbb{N}$. Using the equations $D_k[L_n, L_0] = L_n \cdot D_k(L_0) - L_0 \cdot D_k(L_n)$ and $D_k(L_n) \subseteq (\mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_{k+n}$, we obtain $L_n \cdot D_k(L_0) = -kD_k(L_n)$ for any integers n, k . Therefore,

$$D_k(L_n) = -\frac{1}{k}L_n \cdot D_k(L_0) = \frac{1}{k} \sum_{r=1}^s a(i_r) \left((i_r + \alpha n)v_{n+i_r} \otimes v_{k-i_r} + (k - i_r + \beta n)v_{i_r} \otimes v_{n+k-i_r} \right).$$

This implies that

$$D_k(L_n) = L_n \cdot \left(\sum_{r=1}^s \frac{a(i_r)}{k} v_{i_r} \otimes v_{k-i_r} \right),$$

which shows that D_k is a 1-coboundary. □

Lemma 2.9. *For any $D \in \text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$, we have $D(L_0) = a\delta_{\alpha,0}\delta_{\beta,0}v_0 \otimes v_0$ for some $a \in \mathbb{C}$.*

Proof. Let $D \in \text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$ and suppose that for any $n \in \mathbb{Z}^*$, we have $D(L_n) = \sum_{i \in \mathbb{Z}} a_{n,i}v_i \otimes v_{n-i}$, where $a_{n,i} \in \mathbb{C}$. It should be noted that $L_n \cdot D(L_0) = 0$ for all $n \in \mathbb{Z}$. Additionally, assume that $D(L_0) = \sum_{i=p}^q a(i)v_i \otimes v_{-i}$, where $a(i) \in \mathbb{C}$ and p, q are integers. Then

$$L_n \cdot D(L_0) = - \sum_{i=p}^q a(i) \left((i + \alpha n)v_{n+i} \otimes v_{-i} + (-i + \beta n)v_i \otimes v_{n-i} \right), \quad \forall n \in \mathbb{Z}.$$

Taking $n = 1, -1$ respectively, it follows that

$$\sum_{i=p}^q a(i) \left((i + \alpha)v_{1+i} \otimes v_{-i} + (-i + \beta)v_i \otimes v_{1-i} \right) = \sum_{i=p}^q a(i) \left((i - \alpha)v_{i-1} \otimes v_{-i} + (-i - \beta)v_i \otimes v_{-i-1} \right) = 0.$$

Then

$$a(p)(-p + \beta) = 0, \quad a(p)(p - \alpha) = 0, \quad a(q)(q + \beta) = 0, \quad a(q)(q + \alpha) = 0; \quad (2.1)$$

$$a(i)(-i + \beta) + a(i-1)(i-1 + \alpha) = 0, \quad p+1 \leq i \leq q; \quad (2.2)$$

$$a(i+1)(i+1 - \alpha) - a(i)(i + \beta) = 0, \quad p \leq i \leq q-1. \quad (2.3)$$

It is clear that $a(p)(\alpha - \beta) = a(q)(\alpha - \beta) = 0$. Replacing i by $i-1$ in (2.3), we have

$$a(i)(i - \alpha) - a(i-1)(i-1 + \beta) = 0, \quad p+1 \leq i \leq q. \quad (2.4)$$

Using (2.2) and (2.4), we have $[a(i) - a(i-1)](\alpha - \beta) = 0$ for $p+1 \leq i \leq q$.

Case 1: $\alpha \neq \beta$. Clearly, $a(i) = 0$ for all $p \leq i \leq q$, which implies that $D(L_0) = 0$.

Case 2: $\alpha = \beta \notin \mathbb{Z}$. For any integers p, q , we have $a(p) = a(q) = 0$. Therefore, using (2.2) or (2.3), we conclude that $a(i) = 0$ for all $p \leq i \leq q$, and hence, $D(L_0) = 0$.

Case 3: $\alpha = \beta \in \mathbb{Z}$. Without loss of generality, let us assume that both $a(p)$ and $a(q)$ are non-zero. Then from (2.1), it follows that $\beta = \alpha = p = -q$.

If $\alpha = \beta \neq 0$, then we must have $p < 0$, and also, since the coefficients vanish outside the range of indices (p, q) , we get that $q > 0$. In this case,

$$D(L_0) = \sum_{i=-q}^q a(i)v_i \otimes v_{-i}.$$

Now applying the operator L_n to this expression yields

$$-\sum_{i=-q}^q \left(a(i)(i + \alpha n)v_{n+i} \otimes v_{-i} - a(i)(-i + \beta n)v_i \otimes v_{n-i} \right) = 0, \quad \forall n \in \mathbb{Z}.$$

Thus, for every positive integer value of n , we obtain two equations: $a(q)(n - q - \alpha n) = 0$ and $a(-q)(n + q - \beta n) = 0$. Since both $a(q)$ and $a(-q)$ are non-zero, we must have that $\alpha = \beta = 0$, which leads to a contradiction.

Therefore, we conclude that $\alpha = \beta = 0$. The proof is now complete. □

Let $D \in \text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$ and

$$D(L_n) = \sum_{i \in \mathbb{Z}} a_{n,i} v_i \otimes v_{n-i}, \quad \forall n \in \mathbb{Z}^*,$$

where $a_{n,i} \in \mathbb{C}$. We denote by

$$l = \max\{|i| \mid a_{1,i} \neq 0\}.$$

Lemma 2.10. *If $\beta \notin \mathbb{Z}$, there exist $c_i \in \mathbb{C}(i = -l, \dots, l)$ such that*

$$D(L_1) = L_1 \cdot \left(\sum_{i=-l}^l c_i v_i \otimes v_{-i} \right) + x_l(\alpha + l)v_{l+1} \otimes v_{-l}.$$

Proof. Suppose $\beta \notin \mathbb{Z}$. For any $l \in \mathbb{Z}_+$, we have $D(L_1) = \sum_{i=-l}^l a_{1,i} v_i \otimes v_{-i+1}$ and

$$\begin{aligned} L_1 \cdot \left(\sum_{i=-l}^l x_i v_i \otimes v_{-i} \right) &= - \sum_{i=-l}^l x_i \left((i + \alpha)v_{i+1} \otimes v_{-i} + (-i + \beta)v_i \otimes v_{-i+1} \right) \\ &= -x_{-l}(l + \beta)v_{-l} \otimes v_{l+1} - \sum_{i=-l+1}^l \left((i - 1 + \alpha)x_{i-1} + (\beta - i)x_i \right) v_i \otimes v_{-i+1} - x_l(l + \alpha)v_{l+1} \otimes v_{-l}. \end{aligned}$$

Since $\beta \notin \mathbb{Z}$, we have $\beta + i \neq 0$ for $-l \leq i \leq l$. So the following system of linear equations admits a unique solution:

$$x_{-l}(l + \beta) = -a_{1,-l}, \quad (i - 1 + \alpha)x_{i-1} + (\beta - i)x_i = -a_{1,i}, \quad -l + 1 \leq i \leq l.$$

Let $x_i = c_i(i = -l, \dots, l)$ be the solution. The proof is complete. □

Lemma 2.11. (1) If $\beta \in \mathbb{Z}$ and $|\beta| > l$, there exist some $x_i \in \mathbb{C}$ such that

$$D(L_1) = L_1 \cdot \left(\sum_{i=-l}^l x_i v_i \otimes v_{-i} \right) + x_l (\alpha + l) v_{l+1} \otimes v_{-l}.$$

(2) If $\alpha \in \mathbb{Z}$ and $\alpha < -l + 1$ or $\alpha > l + 1$, there exist $y_i \in \mathbb{C}$ such that

$$D(L_1) = L_1 \cdot \left(\sum_{i=-l-1}^{l-1} y_i v_i \otimes v_{-i} \right) + y_{-l-1} (\beta + l + 1) v_{-l-1} \otimes v_{l+2}.$$

Proof. Similar to the proof of Lemma 2.10. □

Lemma 2.12. (1) If $\beta \in \mathbb{Z}$, and $-l \leq \beta \leq l$, there exist some $x_i \in \mathbb{C} (i = -l, \dots, l)$ such that

$$D(L_1) = (a_{1,\beta} + (\alpha + \beta - 1)x_{\beta-1}) v_\beta \otimes v_{-\beta+1} + L_1 \cdot \left(\sum_{i=-l}^l x_i v_i \otimes v_{-i} \right) + x_l (\alpha + l) v_{l+1} \otimes v_{-l},$$

where $x_\beta = 0$.

(2) If $\alpha \in \mathbb{Z}$, and $-l + 2 \leq \alpha \leq l + 1$, there exist some $y_i \in \mathbb{C} (i = -l - 1, \dots, l - 1)$ such that

$$D(L_1) = (a_{1,-\alpha} + (\alpha + \beta - 1)y_{-\alpha+1}) v_{-\alpha+1} \otimes v_\alpha + L_1 \cdot \left(\sum_{i=-l-1}^{l-1} y_i v_i \otimes v_{-i} \right) + y_{-l-1} (\beta + l + 1) v_{-l-1} \otimes v_{l+2},$$

where $y_{-\alpha} = 0$.

Proof. Suppose $\beta \in \mathbb{Z}$, and $-l \leq \beta \leq l$. Then $D(L_1) = \sum_{i=-l}^l a_{1,i} v_i \otimes v_{-i+1}$ and

$$\begin{aligned} L_1 \cdot \left(\sum_{i=-l}^l x_i v_i \otimes v_{-i} \right) &= - \sum_{i=-l}^l x_i \left((i + \alpha) v_{i+1} \otimes v_{-i} + (-i + \beta) v_i \otimes v_{-i+1} \right) \\ &= -x_{-l} (l + \beta) v_{-l} \otimes v_{l+1} - \sum_{i=-l+1}^l \left((i - 1 + \alpha) x_{i-1} + (\beta - i) x_i \right) v_i \otimes v_{-i+1} - x_l (l + \alpha) v_{l+1} \otimes v_{-l}, \end{aligned}$$

where $x_\beta = 0$. Then the following system of linear equations admits a unique solution:

$$x_{-l} (l + \beta) = -a_{1,-l}, \quad (i - 1 + \alpha) x_{i-1} + (\beta - i) x_i = -a_{1,i}, \quad -l + 1 \leq i \leq l, i \neq \beta.$$

Furthermore,

$$D(L_1) = (a_{1,\beta} + (\alpha + \beta - 1)x_{\beta-1}) v_\beta \otimes v_{-\beta+1} + L_1 \cdot \left(\sum_{i=-l}^l x_i v_i \otimes v_{-i} \right) + x_l (\alpha + l) v_{l+1} \otimes v_{-l}.$$

The proof of (2) is similar. □

Lemma 2.13. If $D(L_0) = 0$, we have

$$(i + m - m\alpha) a_{m,i+m} - (i - m + m\beta) a_{m,i} = (i - m + m\alpha) a_{-m,i-m} - (i + m - m\beta) a_{-m,i} \quad (2.5)$$

for all $m, i \in \mathbb{Z}$.

Proof. If $D(L_0) = 0$, applying D on $[L_m, L_{-m}] = 2mL_0$, we have $L_m \cdot D(L_{-m}) = L_{-m} \cdot D(L_m)$ for $m \in \mathbb{Z}$. It follows that

$$(i + m - m\alpha)a_{m,i+m} - (i - m + m\beta)a_{m,i} = (i - m + m\alpha)a_{-m,i-m} - (i + m - m\beta)a_{-m,i}, \quad \forall m, i \in \mathbb{Z}.$$

□

Lemma 2.14. *For all $i \in \mathbb{Z}$, we have*

$$(i - 2 + 2\alpha)a_{-1,i-2} - (i + 1 - 2\beta)a_{-1,i} + 3a_{1,i} = (i + 1 - \alpha)a_{2,i+1} - (i - 2 + \beta)a_{2,i}, \quad (2.6)$$

$$(i + 2 - 2\alpha)a_{1,i+2} - (i - 1 + 2\beta)a_{1,i} - 3a_{-1,i} = (i - 1 + \alpha)a_{-2,i-1} - (i + 2 - \beta)a_{-2,i}. \quad (2.7)$$

Proof. Applying D on $[L_{-1}, L_2] = -3L_1$, we have

$$L_2 \cdot \left(\sum a_{-1,i} v_i \otimes v_{-i-1} \right) - 3 \sum a_{1,j} v_j \otimes v_{-j+1} = L_{-1} \cdot \left(\sum a_{2,k} v_k \otimes v_{-k+2} \right).$$

Therefore

$$(i - 2 + 2\alpha)a_{-1,i-2} - (i + 1 - 2\beta)a_{-1,i} + 3a_{1,i} = (i + 1 - \alpha)a_{2,i+1} - (i - 2 + \beta)a_{2,i}, \quad i \in \mathbb{Z}.$$

Applying D on $[L_1, L_{-2}] = 3L_{-1}$, we have

$$L_{-2} \cdot \left(\sum a_{1,i} v_i \otimes v_{-i+1} \right) + 3 \sum a_{-1,j} v_j \otimes v_{-j-1} = L_1 \cdot \left(\sum a_{-2,k} v_k \otimes v_{-k-2} \right).$$

It follows that

$$(i + 2 - 2\alpha)a_{1,i+2} - (i - 1 + 2\beta)a_{1,i} - 3a_{-1,i} = (i - 1 + \alpha)a_{-2,i-1} - (i + 2 - \beta)a_{-2,i}, \quad i \in \mathbb{Z}.$$

□

3. CONSTRUCTION OF 1-COCYCLES WITH COEFFICIENTS IN $\mathcal{F}_\alpha \otimes \mathcal{F}_\beta$

In this section, we shall construct several non-trivial 1-cocycles of \mathcal{W} with coefficients in $\mathcal{F}_\alpha \otimes \mathcal{F}_\beta$.

Proposition 3.1. *For any $\beta \in \mathbb{C}$, we define a linear operator $d_{1-\beta,\beta} : \mathcal{W} \rightarrow \mathcal{F}_{1-\beta} \otimes \mathcal{F}_\beta$ as follows:*

$$d_{1-\beta,\beta}(L_n) = \begin{cases} \sum_{i=0}^{n-1} (i - n\beta)v_i \otimes v_{-i+n}, & n \geq 1; \\ 0, & n = 0; \\ -\sum_{i=n}^{-1} (i - n\beta)v_i \otimes v_{-i+n}, & n \leq -1. \end{cases}$$

Then $d_{1-\beta,\beta}$ is a non-trivial 1-cocycle of \mathcal{W} with coefficients in $\mathcal{F}_{1-\beta} \otimes \mathcal{F}_\beta$.

Proof. It is straightforward to check that $d_{1-\beta,\beta} \in \text{Der}(\mathcal{W}, \mathcal{F}_{1-\beta} \otimes \mathcal{F}_\beta)$. Assume $d_{1-\beta,\beta}$ is a 1-coboundary. Then

$$d_{1-\beta,\beta}(L_n) = L_n \cdot \left(\sum_{i=p}^q a_i v_i \otimes v_{-i} \right) = \sum_{i=p}^q a_i \left((i - n\beta)v_i \otimes v_{-i+n} - (i + n - n\beta)v_{i+n} \otimes v_{-i} \right)$$

for some $a_i \in \mathbb{C}$, where $a_p, a_q \in \mathbb{C}^*$ and $p \leq q$. Since $d_{1-\beta, \beta}(L_1) = -\beta v_0 \otimes v_1$, we have

$$-\beta v_0 \otimes v_1 = (p - \beta)a_p v_p \otimes v_{-p+1} + \sum_{i=p+1}^q (i - \beta)[a_i - a_{i-1}]v_i \otimes v_{-i+1} - (q + 1 - \beta)a_q v_{q+1} \otimes v_{-q}.$$

Case 1: $\beta \notin \mathbb{Z}$. If $p \neq 0$, then $(p - \beta)a_p = 0$ implies that $a_p = 0$ or $p = \beta$. However, since we know that $a_p \neq 0$, it follows that $p = \beta$, which is impossible. Similarly, if $p = 0$ and $q > 0$, then $(q + 1 - \beta)a_q = 0$, which is also impossible. Therefore, we must have $p = q = 0$. In this case,

$$-\beta v_0 \otimes v_1 = (-\beta a_0)v_0 \otimes v_1 - (1 - \beta a_0)v_1 \otimes v_0,$$

which is again impossible. Hence, $d_{1-\beta, \beta}$ is a non-trivial 1-cocycle.

Case 2: $\beta \in \mathbb{Z}$.

Subcase 2.1: $p \neq 0$. Then $(p - \beta)a_p = 0$. Since $a_p \neq 0$, we have $p = \beta$.

If $q = p = \beta$, then we see that $-\beta v_0 \otimes v_1 = -a_\beta v_{\beta+1} \otimes v_{-\beta}$. Therefore, $\beta = -1$ and $a_\beta = \beta = -1$. It is clear that

$$L_n \cdot (-v_{-1} \otimes v_1) \neq \sum_{i=0}^{n-1} (i - n\beta)v_i \otimes v_{-i+n}, \quad n \geq 1.$$

Thus, we conclude that $d_{1-\beta, \beta} = d_{2, -1}$ is a non-trivial 1-cocycle.

If $q > p = \beta$, then we express $-\beta v_0 \otimes v_1$ as follows:

$$-\beta v_0 \otimes v_1 = \sum_{i=\beta+1}^q (i - \beta)(a_i - a_{i-1})v_i \otimes v_{-i+1} - (q + 1 - \beta)a_q v_{q+1} \otimes v_{-q}.$$

Since $(q + 1 - \beta)a_q \neq 0$, it follows that $q = -1$ and $a_i = -1$ for $i = \beta, \dots, -1$. This implies that $\beta \leq -1$. Furthermore,

$$L_n \cdot \left(-\sum_{i=\beta}^{-1} v_i \otimes v_{-i} \right) = -\sum_{i=\beta}^{-1} (i - n\beta)v_i \otimes v_{-i+n} + \sum_{i=\beta}^{-1} (i + n - n\beta)v_{i+n} \otimes v_{-i}.$$

However, while

$$L_\beta \cdot \left(-\sum_{i=\beta}^{-1} v_i \otimes v_{-i} \right) \neq -\sum_{i=\beta}^{-1} (i - \beta^2)v_i \otimes v_{-i+\beta},$$

it is impossible.

Subcase 2.2: $p = 0$.

If $p = 0$ and $q > 0$, we have $(q + 1 - \beta)a_q = 0$. Since $a_q \neq 0$, we have $q = \beta - 1$, which implies that $\beta > 1$ and $a_i = 1$ for $i = 0, \dots, \beta - 1$. Then

$$d_{1-\beta, \beta}(L_n) = L_n \cdot \left(\sum_{i=0}^{\beta-1} v_i \otimes v_{-i} \right) = \sum_{i=0}^{\beta-1} (i - n\beta)v_i \otimes v_{-i+n} - \sum_{i=0}^{\beta-1} (i + n - n\beta)v_{i+n} \otimes v_{-i}.$$

While $L_\beta \cdot \left(\sum_{i=0}^{\beta-1} v_i \otimes v_{-i} \right) \neq \sum_{i=0}^{\beta-1} (i - \beta^2)v_i \otimes v_{-i+\beta}$, it is impossible.

If $p = q = 0$, we have

$$-\beta v_0 \otimes v_1 = (-\beta)a_0 v_0 \otimes v_1 - (1 - \beta)a_0 v_1 \otimes v_0.$$

Then $\beta = a_0 = \beta a_0$. So $\beta = a_0 = 0$ or $\beta = a_0 = 1$. Since $a_0 \neq 0$, we have $\beta = a_0 = 1$. It is easy to see that $L_n \cdot (v_0 \otimes v_0) = -nv_0 \otimes v_n \neq d_{0,1}(L_n)$ for all $n \in \mathbb{Z}$. Hence $d_{1-\beta,\beta} = d_{0,1}$ is a non-trivial 1-cocycle.

The proof is complete. \square

Proposition 3.2. *The following linear operators $d_{0,0}^i : \mathcal{W} \rightarrow \mathcal{F}_0 \otimes \mathcal{F}_0 (i = 1, \dots, 5)$ are non-trivial 1-cocycles of \mathcal{W} with coefficients in $\mathcal{F}_0 \otimes \mathcal{F}_0$:*

$$\begin{aligned} d_{0,0}^1(L_m) &= v_0 \otimes v_m, & d_{0,0}^2(L_m) &= mv_0 \otimes v_m, \\ d_{0,0}^3(L_m) &= v_m \otimes v_0, & d_{0,0}^4(L_m) &= mv_m \otimes v_0, \\ d_{0,0}^5(L_n) &= \begin{cases} v_0 \otimes v_0, & n = 0; \\ 0, & n = 1; \\ -\sum_{i=1}^{n-1} v_i \otimes v_{n-i}, & n \geq 2; \\ \sum_{i=n}^0 v_i \otimes v_{n-i}, & n \leq -1 \end{cases} \end{aligned}$$

for $m, n \in \mathbb{Z}$. Furthermore, $d_{0,0}^1, d_{0,0}^2, d_{0,0}^3, d_{0,0}^4, d_{0,0}^5$ are linearly independent.

Proof. It is easy to verify that $d_{0,0}^i$ (for $i = 1, 2, 3, 4$) defined above are non-trivial 1-cocycles of \mathcal{W} in $\mathcal{F}_0 \otimes \mathcal{F}_0$. Next, we shall show that $d_{0,0}^5$ is a non-trivial 1-cocycle. It is straightforward to verify $d_{0,0} \in \text{Der}(\mathcal{W}, \mathcal{F}_0 \otimes \mathcal{F}_0)$. Assume $d_{0,0}^5$ is a 1-coboundary. It follows that

$$d_{0,0}^5(L_n) = L_n \cdot \left(\sum_{i=p}^q a_i v_i \otimes v_{-i} \right) = \sum_{i=p}^q ia_i (v_i \otimes v_{-i+n} - v_{i+n} \otimes v_{-i})$$

for some $a_i \in \mathbb{C}$, where $a_p, a_q \in \mathbb{C}^*$ and $p \leq q$. In particular,

$$d_{0,0}^5(L_1) = pa_p v_p \otimes v_{-p+1} + \sum_{i=p+1}^q (ia_i - (i-1)a_{i-1}) v_i \otimes v_{-i+1} - qa_q v_{q+1} \otimes v_{-q}.$$

Note that $d_{0,0}^5(L_1) = 0$. Then $pa_p = qa_q = 0, ia_i = (i-1)a_{i-1}$ for $i = p+1, \dots, q$. That is, $ia_i = 0$ for all $p \leq i \leq q$. So $p = q = 0$. While $d_{0,0}^5(L_n) = L_n \cdot (a_0 v_0 \otimes v_0) = 0$, we have $d_{0,0}^5(L_n) = 0$, which is a contradiction. Hence $d_{0,0}^5$ is a non-trivial 1-cocycle.

Let $x_1 d_{0,0}^1 + x_2 d_{0,0}^2 + x_3 d_{0,0}^3 + x_4 d_{0,0}^4 + x_5 d_{0,0}^5 = 0$. By the both sides acting on L_1, L_2 respectively, we deduce $x_i = 0, i = 1, 2, 3, 4, 5$. So $d_{0,0}^1, d_{0,0}^2, d_{0,0}^3, d_{0,0}^4, d_{0,0}^5$ are linearly independent. \square

Proposition 3.3. *For $(\alpha, \beta) = (0, 1), (1, 0), (0, 2), (2, 0), (1, 1)$, the following linear operators $d_{\alpha,\beta} : \mathcal{W} \rightarrow \mathcal{F}_\alpha \otimes \mathcal{F}_\beta$ are non-trivial 1-cocycles of \mathcal{W} :*

$$\begin{aligned} d_{0,1}^1(L_n) &= n^2 v_0 \otimes v_n, & d_{1,0}^1(L_n) &= n^2 v_n \otimes v_0, \\ d_{0,2}^1(L_n) &= n^3 v_0 \otimes v_n, & d_{2,0}^1(L_n) &= n^3 v_n \otimes v_0, \end{aligned}$$

$$d_{1,1}(L_n) = \begin{cases} \sum_{i=1}^n i(n-i)v_i \otimes v_{n-i}, & n \geq 1; \\ 0, & n = 0; \\ -\sum_{i=-1}^n i(n-i)v_i \otimes v_{n-i}, & n \leq -1 \end{cases}$$

for $n \in \mathbb{Z}$. Moreover, $d_{0,1}^1, d_{0,1}$ are linearly independent, and $d_{1,0}^1, d_{1,0}$ are linearly independent.

Proof. We directly verify that the linear operators $d_{1,0}, d_{0,1}, d_{2,0}, d_{0,2}$ defined above are non-trivial 1-cocycles. Next, we shall show that $d_{1,1}$ is a non-trivial 1-cocycle. Assume $d_{1,1}$ is a 1-coboundary, then

$$d_{1,1}(L_n) = L_n \cdot \left(\sum_{i=p}^q a_i v_i \otimes v_{-i} \right) = \sum_{i=p}^q a_i \left((i-n)v_i \otimes v_{-i+n} - (i+n)v_{i+n} \otimes v_{-i} \right)$$

for some $a_i \in \mathbb{C}$, where $a_p, a_q \in \mathbb{C}^*$ and $p \leq q$. In particular,

$$d_{1,1}(L_1) = (p-1)a_p v_p \otimes v_{-p+1} + \sum_{i=p+1}^q \left((i-1)a_i - ia_{i-1} \right) v_i \otimes v_{-i+1} - (q+1)a_q v_{q+1} \otimes v_{-q}.$$

Notice that $d_{1,1}(L_1) = 0$. It follows that $(p-1)a_p = (q+1)a_q = 0$. So $p = 1 > q = -1$, which is a contradiction. Then $d_{1,1}$ is a non-trivial 1-cocycle. \square

4. CLASSIFICATION OF 1-COCYCLES FOR $(\alpha, \beta) \notin \mathbb{Z}^2$

In this section, we shall compute $H^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)$ for $(\alpha, \beta) \notin \mathbb{Z}^2$.

Suppose $\beta \notin \mathbb{Z}$ without loss of generality. By Lemma 2.8-Lemma 2.9, it suffices to consider $D \in \text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$ and $D(L_0) = 0$. Suppose

$$D(L_n) = \sum_{i \in \mathbb{Z}} a_{n,i} v_i \otimes v_{n-i}, \quad \forall n \in \mathbb{Z}^*,$$

where $a_{n,i} \in \mathbb{C}$. According to Lemma 2.10, replace D by $D - D_v$, where D_v is a 1-coboundary induced by $v = \sum v_i \otimes v_{-i}$. We denote by

$$l = \max\{|i| \mid a_{1,i} \neq 0\}.$$

For $\beta \notin \mathbb{Z}$, we safely suppose that

$$D(L_1) = a_{1,l+1} v_{l+1} \otimes v_{-l}. \quad (4.1)$$

For the convenience of readers, we provide a scheme of our calculation in this section:

- If $D(L_{-1}) = 0$, we have $D = 0$.
- If $D(L_{-1}) \neq 0$. Set

$$p_{-1} = \min\{i \in \mathbb{Z} \mid a_{-1,i} \neq 0\}, \quad q_{-1} = \max\{i \in \mathbb{Z} \mid a_{-1,i} \neq 0\}.$$

By discussing the values of p_{-1} and q_{-1} , we shall show that $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}\delta_{\alpha+\beta,1}d_{\alpha,\beta}$.

Let $m = 1$ in (2.5). Then

$$(l + 1 - \alpha)a_{1,l+1} = (l - 1 + \alpha)a_{-1,l-1} - (l + 1 - \beta)a_{-1,l}, \quad (4.2)$$

$$-(l + \beta)a_{1,l+1} = (l + \alpha)a_{-1,l} - (l + 2 - \beta)a_{-1,l+1}, \quad (4.3)$$

$$(i - 1 + \alpha)a_{-1,i-1} = (i + 1 - \beta)a_{-1,i}, \quad i \neq l, l + 1. \quad (4.4)$$

Lemma 4.1. *If $\beta \notin \mathbb{Z}$ and $D(L_{-1}) = 0$, we have $D = 0$.*

Proof. If $\beta \notin \mathbb{Z}$ and $D(L_{-1}) = 0$, then we have $a_{1,l+1} = 0$ from (4.3), that is, $D(L_1) = 0$. Using (2.6), we have $(i - 2 + \beta)a_{2,i} = (i + 1 - \alpha)a_{2,i+1}$. Since $\{a_{2,i} \mid a_{2,i} \neq 0, i \in \mathbb{Z}\}$ is finite and $\beta \notin \mathbb{Z}$, we have $a_{2,i} = 0$ for all $i \in \mathbb{Z}$. Then $D(L_2) = 0$. Using (2.7), we have $(i - 1 + \alpha)a_{-2,i-1} = (i + 2 - \beta)a_{-2,i}$. Since $\{a_{-2,i} \neq 0\}$ is finite and $\beta \notin \mathbb{Z}$, we have $a_{-2,i} = 0$ for all $i \in \mathbb{Z}$. Then $D(L_{-2}) = 0$. Since the Witt algebra \mathcal{W} can be generated by $\{L_{\pm 1}, L_{\pm 2}\}$, we have $D(L_m) = 0$ for all $m \in \mathbb{Z}$. Then $D = 0$. \square

Next we assume $D(L_{-1}) \neq 0$. Set

$$p_{-1} = \min\{i \in \mathbb{Z} \mid a_{-1,i} \neq 0\}, \quad q_{-1} = \max\{i \in \mathbb{Z} \mid a_{-1,i} \neq 0\}.$$

Lemma 4.2. *If $\beta \notin \mathbb{Z}$, we have $p_{-1} = l$ or $p_{-1} = l + 1$. Moreover, if $q_{-1} \neq l - 1, l$, then $q_{-1} = -\alpha$.*

Proof. Assume $p_{-1} \neq l, l + 1$. Let $i = p_{-1}$ in (4.4). From $a_{-1,p_{-1}-1} = 0$, we have $a_{-1,p_{-1}}(\beta - p_{-1} - 1) = 0$. Since $\beta \notin \mathbb{Z}$, we have $a_{-1,p_{-1}} = 0$, a contradiction. Therefore $p_{-1} = l$ or $p_{-1} = l + 1$. Moreover, suppose $q_{-1} \neq l - 1, l$, that is, $q_{-1} + 1 \neq l, l + 1$. Let $i = q_{-1} + 1$ in (4.4). Then $a_{-1,q_{-1}}(\alpha + q_{-1}) = 0$. It follows that $q_{-1} = -\alpha$, which implies that $\alpha \in \mathbb{Z}$ and $-\alpha \neq l - 1, l$. \square

From Lemma 4.2, we have the following result.

Lemma 4.3. *If $\beta \notin \mathbb{Z}$ and $\alpha \notin \mathbb{Z}$, we have $p_{-1} = q_{-1} = l$.*

Lemma 4.4. *If $\beta \notin \mathbb{Z}$ and $p_{-1} = q_{-1} = l$, we have $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}\delta_{\alpha+\beta,1}d_{\alpha,\beta}$.*

Proof. If $\beta \notin \mathbb{Z}$ and $p_{-1} = q_{-1} = l$, that is, $D(L_{-1}) = a_{-1,l}v_l \otimes v_{-l-1}$, by (4.2)-(4.3), we have

$$a_{1,l+1}(\alpha - l - 1) = -a_{-1,l}(\beta - l - 1), \quad (4.5)$$

$$a_{1,l+1}(\beta + l) = -a_{-1,l}(\alpha + l). \quad (4.6)$$

Since $a_{-1,l} \neq 0$ and $\beta \notin \mathbb{Z}$, we have $a_{1,l+1} \neq 0$, $\alpha \neq l + 1$ and $(\alpha + l)(\alpha - l - 1) = (\beta + l)(\beta - l - 1)$. So $\alpha = \beta$ or $\alpha + \beta = 1$.

Case 1: $\alpha = \beta \notin \mathbb{Z}$. By (4.6), we have $a_{1,l+1} = -a_{-1,l}$, $D(L_1) = -a_{-1,l}v_{l+1} \otimes v_{-l}$ and $D(L_{-1}) = a_{-1,l}v_l \otimes v_{-l-1}$. Using (2.6), we have

$$\begin{aligned} -(l + 1 - 2\beta)a_{-1,l} &= (l + 1 - \beta)a_{2,l+1} - (l - 2 + \beta)a_{2,l}, \\ -3a_{-1,l} &= (l + 2 - \beta)a_{2,l+2} - (l - 1 + \beta)a_{2,l+1}, \\ (l + 2\beta)a_{-1,l} &= (l + 3 - \beta)a_{2,l+3} - (l + \beta)a_{2,l+2}, \\ (i + 1 - \beta)a_{2,i+1} &= (i - 2 + \beta)a_{2,i}, \quad i \neq l, l + 1, l + 2. \end{aligned}$$

Since $\{a_{2,i} \neq 0\}$ is finite and $\beta \notin \mathbb{Z}$, we have $a_{2,i} = 0$ for $i \leq l$ or $i \geq l+3$. Then

$$\begin{aligned} -(l+1-2\beta)a_{-1,l} &= (l+1-\beta)a_{2,l+1}, & -3a_{-1,l} &= (l+2-\beta)a_{2,l+2} - (l-1+\beta)a_{2,l+1}, \\ (l+2\beta)a_{-1,l} &= -(l+\beta)a_{2,l+2}. \end{aligned}$$

It follows that

$$\begin{aligned} a_{2,l+1} &= -\frac{l+1-2\beta}{l+1-\beta}a_{-1,l}, & a_{2,l+2} &= -\frac{l+2\beta}{l+\beta}a_{-1,l}, \\ \left(3 + (l-1+\beta)\frac{l+1-2\beta}{l+1-\beta} - (l+2-\beta)\frac{l+2\beta}{l+\beta}\right)a_{-1,l} &= 0. \end{aligned}$$

Since

$$\left(3 + (l-1+\beta)\frac{l+1-2\beta}{l+1-\beta} - (l+2-\beta)\frac{l+2\beta}{l+\beta}\right) = -\frac{2\beta(1-\beta)(1-2\beta)}{(l+\beta)(l+1-\beta)}$$

and $a_{-1,l} \neq 0$, we get $\beta = \frac{1}{2}$. Hence $a_{2,l+1} = -\frac{l}{l+\frac{1}{2}}a_{-1,l}$, $a_{2,l+2} = -\frac{l+1}{l+\frac{1}{2}}a_{-1,l}$ and

$$D(L_2) = -\frac{1}{l+\frac{1}{2}}a_{-1,l}(lv_{l+1} \otimes v_{-l+1} + (l+1)v_{l+2} \otimes v_{-l}).$$

Since $\{a_{-2,i} \neq 0\}$ is finite, using (2.7), we have $a_{-2,i} = 0$ for $i \neq l-1, l$, and

$$(l+\frac{1}{2})a_{-2,l-1} = la_{-1,l}, \quad (l+\frac{3}{2})a_{-2,l} - (l-\frac{1}{2})a_{-2,l-1} = 3a_{-1,l}, \quad (l+\frac{1}{2})a_{-2,l} = (l+1)a_{-1,l}.$$

Hence $a_{-2,l-1} = \frac{l}{l+\frac{1}{2}}a_{-1,l}$, $a_{-2,l} = \frac{l+1}{l+\frac{1}{2}}a_{-1,l}$. Set $a = -\frac{a_{-1,l}}{l+\frac{1}{2}}$. Then

$$D(L_{-1}) = -a(l+\frac{1}{2})v_l \otimes v_{-l-1}, \quad D(L_{-2}) = -a(lv_{l-1} \otimes v_{-l-1} + (l+1)v_l \otimes v_{-l-2}),$$

$$D(L_1) = a(l+\frac{1}{2})v_{l+1} \otimes v_{-l}, \quad D(L_2) = a(lv_{l+1} \otimes v_{-l+1} + (l+1)v_{l+2} \otimes v_{-l}).$$

Applying D on the identities $[L_k, L_1] = (k-1)L_{k+1}$ and $[L_{-k}, L_{-1}] = (-k+1)L_{-k-1}$ and using induction on $k \in \mathbb{Z}_+$, we obtain

$$D(L_n) = a \sum_{i=l+1}^{l+n} (i - \frac{n}{2})v_i \otimes v_{-i+n} = L_n \cdot \left(-a \sum_{i=0}^l v_i \otimes v_{-i}\right) + ad_{\frac{1}{2}, \frac{1}{2}}(L_n), \quad n \geq 1;$$

$$D(L_n) = -a \sum_{i=l+n+1}^l (i - \frac{n}{2})v_i \otimes v_{-i+n} = L_n \cdot \left(-a \sum_{i=0}^l v_i \otimes v_{-i}\right) + ad_{\frac{1}{2}, \frac{1}{2}}(L_n), \quad n \leq -1.$$

Therefore, $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{\frac{1}{2}, \frac{1}{2}}$.

Case 2: $\alpha = 1 - \beta$. Then $a_{1,l+1} = -\frac{l+1-\beta}{l+\beta}a_{-1,l}$ from (4.6) and $D(L_1) = -\frac{l+1-\beta}{l+\beta}a_{-1,l}v_{l+1} \otimes v_{-l}$. Using (2.6), we have

$$\begin{aligned} -(l+1-2\beta)a_{-1,l} &= (l+\beta)a_{2,l+1} - (l-2+\beta)a_{2,l}, \\ -3\frac{l+1-\beta}{l+\beta}a_{-1,l} &= (l+1+\beta)a_{2,l+2} - (l-1+\beta)a_{2,l+1}, \\ (l+2-2\beta)a_{-1,l} &= (l+2+\beta)a_{2,l+3} - (l+\beta)a_{2,l+2}, \\ (i+\beta)a_{2,i+1} &= (i-2+\beta)a_{2,i}, \quad i \neq l, l+1, l+2. \end{aligned}$$

Since $\{a_{2,i} \neq 0\}$ is finite, $a_{2,i} = 0$ for $i \leq l$ or $i \geq l+3$. Then

$$\begin{aligned} a_{2,l+1} &= -\frac{l+1-2\beta}{l+\beta}a_{-1,l}, & a_{2,l+2} &= -\frac{l+2-2\beta}{l+\beta}a_{-1,l}, \\ D(L_2) &= -\frac{l+1-2\beta}{l+\beta}a_{-1,l}v_{l+1} \otimes v_{-l+1} - \frac{l+2-2\beta}{l+\beta}a_{-1,l}v_{l+2} \otimes v_{-l}. \end{aligned}$$

Using (2.7), we have

$$a_{-2,l-1} = \frac{l-1+2\beta}{l+\beta}a_{-1,l}, \quad a_{-2,l} = \frac{l+2\beta}{l+\beta}a_{-1,l}.$$

Set $a = -\frac{a_{-1,l}}{l+\beta}$. Then

$$\begin{aligned} D(L_{-1}) &= -a(l+\beta)v_l \otimes v_{-l-1}, \\ D(L_{-2}) &= -a((l-1+2\beta)v_{l-1} \otimes v_{-l-1} + (l+2\beta)v_l \otimes v_{-l-2}), \\ D(L_1) &= a(l+1-\beta)v_{l+1} \otimes v_{-l}, \\ D(L_2) &= a((l+1-2\beta)v_{l+1} \otimes v_{-l+1} + (l+2-2\beta)v_{l+2} \otimes v_{-l}). \end{aligned}$$

Moreover, applying D on the identities $[L_k, L_1] = (k-1)L_{k+1}$ and $[L_{-k}, L_{-1}] = (-k+1)L_{-k-1}$ and using induction on $k \in \mathbb{Z}_+$, we have

$$\begin{aligned} D(L_n) &= a \sum_{i=l+1}^{l+n} (i-n\beta)v_i \otimes v_{-i+n} = L_n \cdot \left(-a \sum_{i=0}^l v_i \otimes v_{-i} \right) + a \sum_{i=0}^{n-1} (i-n\beta)v_i \otimes v_{-i+n}, \quad n \geq 1; \\ D(L_n) &= -a \sum_{i=l+n+1}^l (i-n\beta)v_i \otimes v_{-i+n} = L_n \cdot \left(-a \sum_{i=0}^l v_i \otimes v_{-i} \right) - a \sum_{i=n}^{-1} (i-n\beta)v_i \otimes v_{-i+n}, \quad n \leq -1. \end{aligned}$$

It follows that $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1-\beta,\beta}$. \square

Next we discuss the case of $q_{-1} \geq l+1$. Notice that Lemma 4.2 implies $\alpha = -q_{-1} \in \mathbb{Z}$.

Lemma 4.5. *If $\beta \notin \mathbb{Z}$, $p_{-1} = l$ and $q_{-1} = l+k_0$ for some integer $k_0 \geq 1$, then D is a 1-coboundary.*

Proof. If $\beta \notin \mathbb{Z}$, $p_{-1} = l$ and $q_{-1} = l+k_0 \geq l+1$, we have

$$D(L_{-1}) = a_{-1,l}v_l \otimes v_{-l-1} + \cdots + a_{-1,l+k_0}v_{l+k_0} \otimes v_{-l-k_0-1},$$

$\alpha = -q_{-1} = -l - k_0$ by Lemma 2.11 and

$$\begin{aligned} (2l+k_0+1)a_{1,l+1} &= -(l+1-\beta)a_{-1,l}, & (l+\beta)a_{1,l+1} &= k_0a_{-1,l} + (l+2-\beta)a_{-1,l+1}, \\ (l+k_0+1-i)a_{-1,i-1} &= -(i+1-\beta)a_{-1,i}, & i &\neq l, l+1 \end{aligned}$$

from (4.2)-(4.4). Since $\beta \notin \mathbb{Z}$ and $a_{-1,l} \neq 0$, it is easy to see that $2l+k_0+1 \neq 0$ and

$$a_{1,l+1} = -\frac{a_{-1,l}}{2l+k_0+1}(l+1-\beta), \quad (4.7)$$

$$a_{-1,l+1} = -\frac{a_{-1,l}}{2l+k_0+1} \frac{(l+k_0+\beta)(l+k_0+1-\beta)}{l+2-\beta}, \quad (4.8)$$

$$a_{-1,i} = -\frac{l+k_0+1-i}{i+1-\beta}a_{-1,i-1}, \quad i \geq l+2.$$

Then

$$a_{-1,l+i} = (-1)^{i-1} \left(\prod_{t=1}^{i-1} \frac{k_0-t}{l+t+2-\beta} \right) a_{-1,l+1}, \quad 2 \leq i \leq k_0. \quad (4.9)$$

Set $a = -\frac{a_{-1,l}}{2l+k_0+1}$. Then $D(L_1) = L_1 \cdot \left(\sum_{i=l+1}^{l+k_0} x_i v_i \otimes v_{-i} \right)$, where

$$x_{l+1} = a, \quad x_{l+i} = (-1)^{i-1} \left(\prod_{t=2}^i \frac{k_0-t+1}{l+t-\beta} \right) a, \quad 2 \leq i \leq k_0.$$

In fact, $x_{l+1}, \dots, x_{l+k_0}$ satisfy the following relations:

$$x_i = -\frac{l+k_0+1-i}{i-\beta} x_{i-1}, \quad i = l+2, \dots, l+k_0.$$

Moreover, we have

$$a_{1,l+1} = a(l+1-\beta), \quad a_{-1,l} = -a(2l+k_0+1), \quad a_{-1,l+i} = \frac{(l+k_0+\beta)(l+k_0+1-\beta)}{l+i+1-\beta} x_{l+i}, \quad 1 \leq i \leq k_0.$$

It follows that $D(L_{-1}) = L_{-1} \cdot \left(\sum_{i=l+1}^{l+k_0} x_i v_i \otimes v_{-i} \right)$. Denote by $D_0 = D - D_{\text{inn}}$, where $D_{\text{inn}}(L_n) = L_n \cdot \left(\sum_{i=l+1}^{l+k_0} x_i v_i \otimes v_{-i} \right)$. Then $D_0(L_{\pm 1}) = 0$. By Lemma 4.1, we have $D_0 = 0$, which implies that D is a 1-coboundary. \square

Lemma 4.6. *If $\beta \notin \mathbb{Z}$, $p_{-1} = l+1$, and $q_{-1} = l+k_0$, where $k_0 \geq 1$, then D is a 1-coboundary.*

Proof. Suppose $\beta \notin \mathbb{Z}$, $p_{-1} = l+1$, and $q_{-1} = l+k_0$, where $k_0 \geq 1$. Then $\alpha = -l-k_0$ and (4.2)-(4.4) become

$$(2l+k_0+1)a_{1,l+1} = 0, \quad (4.10)$$

$$(l+\beta)a_{1,l+1} = (l+2-\beta)a_{-1,l+1}, \quad (4.11)$$

$$a_{-1,i} = -\frac{l+k_0+1-i}{i+1-\beta}a_{-1,i-1}, \quad i \geq l+2. \quad (4.12)$$

Since $\beta \notin \mathbb{Z}$ and $a_{-1,l+1} \neq 0$, we have $a_{1,l+1} \neq 0$ via (4.11). Therefore, by (4.10), we have $2l+k_0+1=0$, that is, $k_0 = -2l-1$. Hence $l \leq -1$ and $\alpha = l+1$. From (4.11)-(4.12), we have $a_{1,l+1} = \frac{l+2-\beta}{l+\beta}a_{-1,l+1}$ and

$$D(L_1) = \frac{l+2-\beta}{l+\beta}a_{-1,l+1}v_{l+1} \otimes v_{-l}, \quad D(L_{-1}) = \sum_{i=l+1}^{-l-1} a_{-1,i}v_i \otimes v_{-i-l},$$

where $a_{-1,i} = \frac{l+i}{i+1-\beta}a_{-1,i-1}$ for $i \geq l+2$. Furthermore,

$$a_{-1,l+1+k} = \left(\prod_{t=1}^k \frac{2l+1+t}{l+2+t-\beta} \right) a_{-1,l+1}, \quad 1 \leq k \leq -2l-1. \quad (4.13)$$

In addition, $D(L_1) = L_1 \cdot \left(\sum_{i=l+1}^{-l-1} x_i v_i \otimes v_{-i} \right)$, where

$$x_{l+1} = \frac{l+2-\beta}{(l+\beta)(l+1-\beta)} a_{-1,l+1}, \quad x_i = \frac{l+i}{i-\beta} x_{i-1}, \quad i = l+2, \dots, -l-1.$$

Then

$$x_{l+k} = \frac{2l+k}{l+k-\beta} \cdot \frac{2l+k-1}{l+k-1-\beta} \cdots \frac{2l+3}{l+3-\beta} \cdot \frac{2l+2}{l+2-\beta} x_{l+1} = \frac{l+k+1-\beta}{(l+\beta)(l+1-\beta)} a_{-1,l+k}$$

for $1 \leq k \leq -2l-1$. Hence

$$\frac{(l+\beta)(l+1-\beta)}{i+1-\beta} x_i = a_{-1,i}$$

for $l+1 \leq i \leq -l-1$. It follows that

$$(l-i)x_{i+1} + (i+\beta)x_i = a_{-1,i}, \quad l+1 \leq i \leq -l-1.$$

Moreover,

$$\begin{aligned} L_{-1} \cdot \left(\sum_{i=l+1}^{-l-1} x_i v_i \otimes v_{-i} \right) &= - \sum_{i=l+1}^{-l-1} x_i \left((i-\alpha)v_{i-1} \otimes v_{-i} + (-i-\beta)v_i \otimes v_{-i-1} \right) \\ &= \sum_{i=l+1}^{-l-1} x_i \left((l+1-i)v_{i-1} \otimes v_{-i} + (i+\beta)v_i \otimes v_{-i-1} \right) \\ &= \sum_{i=l+1}^{-l-1} \left((l-i)x_{i+1} + (i+\beta)x_i \right) v_i \otimes v_{-i-1} \\ &= \sum_{i=l+1}^{-l-1} a_{-1,i} v_i \otimes v_{-i-1} = D(L_{-1}). \end{aligned}$$

Denote by $D_0 = D - D_{inn}$, where

$$D_{inn}(L_n) = L_n \cdot \left(\sum_{i=l+1}^{-l-1} x_i v_i \otimes v_{-i} \right).$$

Then $D_0(L_{\pm 1}) = 0$. By Lemma 4.1, we have $D_0 = 0$. Hence D is a 1-coboundary. \square

Next, we state our main result in this section.

Proposition 4.7. *If $(\alpha, \beta) \notin \mathbb{Z}^2$, then $H^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta) = \mathbb{C} \delta_{\alpha+\beta, 1} d_{\alpha, \beta}$, where $d_{\alpha, \beta}$ is defined in Proposition 3.1.*

Proof. If $\beta \notin \mathbb{Z}$, by Lemma 4.3-Lemma 4.6, we have

$$\text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 = \begin{cases} \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C} d_{\alpha, \beta}, & \alpha + \beta = 1; \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0, & \text{otherwise.} \end{cases}$$

It follows from Lemma 2.7 and Lemma 2.8 that $H^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta) = \mathbb{C} \delta_{\alpha+\beta, 1} d_{\alpha, \beta}$. The case $\alpha \notin \mathbb{Z}$ can be handled in a similar manner. \square

5. CLASSIFICATION OF 1-COCYCLES FOR $(\alpha, \beta) \in \mathbb{Z}^2$

In this section, we shall compute $H^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)$ for the case $(\alpha, \beta) \in \mathbb{Z}^2$.

By Lemma 2.8-Lemma 2.9, it suffices to consider $D \in \text{Der}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$ and $D(L_0) = a\delta_{\alpha,0}\delta_{\beta,0}v_0 \otimes v_0$ for some $a \in \mathbb{C}$. If $D(L_0) \neq 0$, we suppose that $D(L_0) = v_0 \otimes v_0$. By replacing D by $D - d_{0,0}^1$, we have $D(L_0) = 0$. Thus it suffices to deal with the case $D(L_0) = 0$.

By Lemma 2.11 and Lemma 2.12, we assume

$$D(L_1) = a_{1,\beta}v_\beta \otimes v_{-\beta+1} + a_{1,l+1}v_{l+1} \otimes v_{-l}$$

for some $l \in \mathbb{Z}$ and $\beta \leq l$.

For the convenience of readers, we provide a scheme of our proof in this section. It will be divided into three cases for discussion.

- If $D(L_{\pm 1}) = 0$, we shall classify all 1-cocycles.
- If $D(L_{\mp 1}) = 0$ and $D(L_{\pm 1}) \neq 0$, we shall determine all 1-cocycles.
- If $D(L_{\pm 1}) \neq 0$ and

$$p_{-1} = \min\{i \mid a_{-1,i} \neq 0\}, \quad q_{-1} = \max\{i \mid a_{-1,i} \neq 0\},$$

By discussing the values of p_{-1} and q_{-1} , we shall obtain all 1-cocycles.

It is important to note that the proof process for these three cases is not entirely analogous to others and needs individual treatment.

5.1. $D(L_{\pm 1}) = 0$.

Lemma 5.1. *If $D(L_0) = D(L_{\pm 1}) = 0$, we have*

$$D \in \begin{cases} \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,0}^1 \oplus \mathbb{C}d_{0,0}^2 \oplus \mathbb{C}d_{0,0}^3 \oplus \mathbb{C}d_{0,0}^4 \oplus \mathbb{C}d_{0,0}^5, & (\alpha, \beta) = (0, 0); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,1} \oplus \mathbb{C}d_{0,1}^1, & (\alpha, \beta) = (0, 1); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1,0} \oplus \mathbb{C}d_{1,0}^1, & (\alpha, \beta) = (1, 0); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,2}, & (\alpha, \beta) = (0, 2); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{2,0}, & (\alpha, \beta) = (2, 0); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1,1}, & (\alpha, \beta) = (1, 1); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0, & \text{otherwise.} \end{cases}$$

Proof. If $D(L_{\pm 1}) = 0$, we have

$$(i+1-\alpha)a_{2,i+1} = (i-2+\beta)a_{2,i}, \quad (i-1+\alpha)a_{-2,i-1} = (i+2-\beta)a_{-2,i} \quad (5.1)$$

from (2.6) and (2.7). Since both $\{a_{2,i} \neq 0 \mid i \in \mathbb{Z}\}$ and $\{a_{-2,i} \neq 0 \mid i \in \mathbb{Z}\}$ are finite, by (5.1), we have

$$D(L_2) = \sum_{i=\alpha}^{2-\beta} a_{2,i}v_i \otimes v_{-i+2}, \quad D(L_{-2}) = \sum_{i=\beta-2}^{-\alpha} a_{-2,i}v_i \otimes v_{-i-2},$$

where $\alpha \leq 2 - \beta$. Moreover,

$$a_{2,\alpha+k} = a_{2,\alpha} \prod_{t=1}^k \frac{\alpha + \beta - 3 + t}{t}, \quad 1 \leq k \leq 2 - \alpha - \beta, \quad (5.2)$$

$$a_{-2,-\alpha-k} = a_{-2,-\alpha} \prod_{t=1}^k \frac{\alpha + \beta - 3 + t}{t}, \quad 1 \leq k \leq 2 - \alpha - \beta. \quad (5.3)$$

On the other hand, by $L_2 \cdot D(L_{-2}) = L_{-2} \cdot D(L_2)$, we have

$$\begin{aligned} & \beta a_{-2,\beta-2} v_{\beta-2} \otimes v_{2-\beta} + (\beta - 1) a_{-2,\beta-1} v_{\beta-1} \otimes v_{1-\beta} \\ & + \sum_{i=\beta}^{-\alpha} \left((i - 2 + 2\alpha) a_{-2,i-2} + (-i - 2 + 2\beta) a_{-2,i} \right) v_i \otimes v_{-i} \\ & + (\alpha - 1) a_{-2,-\alpha-1} v_{-\alpha+1} \otimes v_{\alpha-1} + \alpha a_{-2,-\alpha} v_{-\alpha+2} \otimes v_{\alpha-2} \\ = & -\alpha a_{2,\alpha} v_{\alpha-2} \otimes v_{-\alpha+2} - (\alpha - 1) a_{2,\alpha+1} v_{\alpha-1} \otimes v_{-\alpha+1} \\ & + \sum_{i=\alpha}^{-\beta} \left((i + 2 - 2\alpha) a_{2,i+2} + (-i + 2 - 2\beta) a_{2,i} \right) v_i \otimes v_{-i} \\ & - (\beta - 1) a_{2,-\beta+1} v_{-\beta+1} \otimes v_{\beta-1} - \beta a_{2,-\beta+2} v_{-\beta+2} \otimes v_{\beta-2}. \end{aligned} \quad (5.4)$$

Case 1: $\alpha < \beta$. Then $-\alpha a_{2,\alpha} v_{\alpha-2} \otimes v_{-\alpha+2} = 0$ and $\alpha a_{-2,-\alpha} v_{-\alpha+2} \otimes v_{\alpha-2} = 0$ via (5.4). Hence $\alpha a_{2,\alpha} = \alpha a_{-2,-\alpha} = 0$.

Subcase 1.1: $\alpha \neq 0$. Then $a_{2,\alpha} = a_{-2,-\alpha} = 0$, which forces $D(L_{\pm 2}) = 0$. Hence $D = 0$.

Subcase 1.2: $\alpha = 0$. Since $\beta \leq 2$, we have $0 < \beta \leq 2$, i.e., $\beta = 1$ or 2 .

(1) $(\alpha, \beta) = (0, 1)$. By (5.1), we get

$$D(L_2) = a_{2,0} v_0 \otimes v_2 + a_{2,1} v_1 \otimes v_1, \quad D(L_{-2}) = a_{-2,-1} v_{-1} \otimes v_{-1} + a_{-2,0} v_0 \otimes v_{-2},$$

and $a_{2,0} = -a_{2,1}$, $a_{-2,0} = -a_{-2,-1}$. Let $m = 2$, $i = -1$ in (2.5). Then $a_{-2,-1} = a_{2,1}$. Hence

$$D(L_2) = a_{2,0} (v_0 \otimes v_2 - v_1 \otimes v_1), \quad D(L_{-2}) = -a_{2,0} (v_{-1} \otimes v_{-1} - v_0 \otimes v_{-2}).$$

Set $a = \frac{a_{2,0}}{2}$ and replace D by $D - 2ad_{0,1} - D_{inn}$, where $D_{inn}(L_n) = L_n \cdot (av_0 \otimes v_0)$ for all $n \in \mathbb{Z}$. Then

$$D(L_{-2}) = 0, \quad D(L_{-1}) = -av_0 \otimes v_{-1}, \quad D(L_1) = 3av_0 \otimes v_1, \quad D(L_2) = 8av_0 \otimes v_2.$$

Using induction on n by $(n-2)D(L_n) = D[L_{n-1}, L_1]$ and $(n-2)D(L_{-n}) = D[L_{-1}, L_{-n+1}]$, we have

$$D(L_n) = a(n^2 + 2n)v_0 \otimes v_n = ad_{0,1}^1(L_n) - L_n \cdot (2av_0 \otimes v_0), \quad \forall n \in \mathbb{Z}.$$

So $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,1} \oplus \mathbb{C}d_{0,1}^1$.

(2) $(\alpha, \beta) = (0, 2)$. Then $D(L_2) = a_{2,0} v_0 \otimes v_2$ and $D(L_{-2}) = a_{-2,0} v_0 \otimes v_{-2}$. By (2.5), we have $a_{-2,0} = -a_{2,0}$. Using induction on n by

$$(n-2)D(L_n) = D[L_{n-1}, L_1] = -L_1 \cdot D(L_{n-1}), \quad (n-2)D(L_{-n}) = D[L_{-1}, L_{-n+1}] = L_{-1} \cdot D(L_{-n+1}),$$

we obtain

$$D(L_n) = \frac{a_{-2,0}}{6}(n^3 - n)v_0 \otimes v_n = \frac{a_{-2,0}}{6}d_{0,2}(L_n) + L_n \cdot \left(\frac{a_{-2,0}}{12}v_0 \otimes v_0 \right), \quad \forall n \in \mathbb{Z}.$$

So $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,2}$.

Case 2: $\beta < \alpha$. Since $\alpha + \beta \leq 2$, we have $2\beta < \alpha + \beta \leq 2$. So $\beta \leq 0$. By (5.4), we have

$$(\beta - 1)a_{-2,\beta-1}v_{\beta-1} \otimes v_{1-\beta} = 0, \quad -(\beta - 1)a_{2,-\beta+1}v_{-\beta+1} \otimes v_{\beta-1} = 0.$$

Furthermore,

$$\left((i - 2 + 2\alpha)a_{-2,i-2} + (-i - 2 + 2\beta)a_{-2,i} \right) v_i \otimes v_{-i} = 0, \quad i = \beta, \dots, -\alpha,$$

$$\left((i + 2 - 2\alpha)a_{2,i+2} + (-i + 2 - 2\beta)a_{2,i} \right) v_i \otimes v_{-i} = 0, \quad i = \alpha, \dots, -\beta,$$

$$(\alpha - 1)a_{-2,-\alpha-1}v_{-\alpha-1} \otimes v_{\alpha-1} = 0, \quad -(\alpha - 1)a_{2,\alpha+1}v_{\alpha-1} \otimes v_{-\alpha+1} = 0,$$

$$\alpha a_{-2,-\alpha}v_{-\alpha+2} \otimes v_{\alpha-2} = 0, \quad -\alpha a_{2,\alpha}v_{\alpha-2} \otimes v_{-\alpha+2} = 0.$$

Subcase 2.1: $\beta < 0$. Then $a_{-2,\beta-2} = a_{-2,\beta-1} = 0$, $a_{2,-\beta+2} = a_{2,-\beta+1} = 0$ and we get $D(L_{\pm 2}) = 0$. Hence $D = 0$.

Subcase 2.2: $\beta = 0$. Then $\alpha = 1$ or 2 .

(1) $(\alpha, \beta) = (1, 0)$. Then

$$D(L_2) = a_{2,1}v_1 \otimes v_1 + a_{2,2}v_2 \otimes v_0, \quad D(L_{-2}) = a_{-2,-2}v_{-2} \otimes v_0 + a_{-2,-1}v_{-1} \otimes v_{-1}.$$

By (5.1), we have $a_{2,2} = -a_{2,1}$, $a_{-2,-2} = -a_{-2,-1}$. Let $m = 2, i = 1$ in (2.5). Then $a_{-2,-1} = a_{2,1}$. So $D(L_2) = a_{2,2}(v_2 \otimes v_0 - v_1 \otimes v_1)$ and $D(L_{-2}) = a_{2,2}(v_{-2} \otimes v_0 - v_{-1} \otimes v_{-1})$. Replacing D by $D + a_{2,2}d_{1,0} + D_{inn}$, where $D_{inn}(L_n) = L_n \cdot \left(\frac{a_{2,2}}{2}v_0 \otimes v_0 \right)$ for all $n \in \mathbb{Z}$. Then

$$D(L_{-1}) = \frac{3a_{2,2}}{2}v_{-1} \otimes v_0, \quad D(L_1) = -\frac{a_{2,2}}{2}v_1 \otimes v_0, \quad D(L_{-2}) = 4a_{2,2}v_{-2} \otimes v_0, \quad D(L_2) = 0.$$

Using induction on n by $(n - 2)D(L_n) = D[L_{n-1}, L_1]$ and $(n - 2)D(L_{-n}) = D[L_{-1}, L_{-n+1}]$, we have

$$D(L_n) = \frac{a_{2,2}}{2}(n^2 - 2n)v_n \otimes v_0 = \frac{a_{2,2}}{2}d_{0,1}^1(L_n) + L_n \cdot \left(a_{2,2}v_0 \otimes v_0 \right), \quad \forall n \in \mathbb{Z}.$$

Therefore, $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1,0} \oplus \mathbb{C}d_{1,0}^1$.

(2) $(\alpha, \beta) = (2, 0)$. Then $D(L_2) = a_{2,2}v_2 \otimes v_0$ and $D(L_{-2}) = a_{-2,-2}v_{-2} \otimes v_0$. Let $m = 2, i = 0$ in (2.5). Then $a_{-2,-2} = -a_{2,2}$. Using induction on n by

$$(n - 2)D(L_n) = D[L_{n-1}, L_1] = -L_1 \cdot D(L_{n-1}), \quad (n - 2)D(L_{-n}) = D[L_{-1}, L_{-n+1}] = L_{-1} \cdot D(L_{-n+1}),$$

we deduce that

$$D(L_n) = \frac{a_{2,2}}{6}(n^3 - n)v_n \otimes v_0 = \frac{a_{2,2}}{6}d_{2,0}(L_n) + L_n \cdot \left(\frac{a_{2,2}}{12}v_0 \otimes v_0 \right), \quad \forall n \in \mathbb{Z},$$

which implies that $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{2,0}$.

Case 3: $\alpha = \beta$. Then $\alpha = \beta = 0, 1$ or $\alpha = \beta < 0$.

Subcase 3.1: $\alpha = \beta = 0$. Then $D(L_2) = \sum_{i=0}^2 a_{2,i}v_i \otimes v_{-i+2}$ and $D(L_{-2}) = \sum_{i=-2}^0 a_{-2,i}v_i \otimes v_{-i-2}$. By (5.2)-(5.3), we have $a_{2,1} = -2a_{2,0}$, $a_{2,2} = a_{2,0}$, $a_{-2,-1} = -2a_{-2,0}$, $a_{-2,-2} = a_{-2,0}$. Using (5.4), we get $a_{-2,-1} = -a_{2,1}$. So $a_{-2,0} = -a_{2,0}$. Set $a = a_{2,0}$, then

$$D(L_2) = a(v_0 \otimes v_2 + v_2 \otimes v_0) - 2av_1 \otimes v_1, \quad D(L_{-2}) = -a(v_0 \otimes v_{-2} + v_{-2} \otimes v_0) + 2av_{-1} \otimes v_{-1}.$$

Using induction on n by $(n-2)D(L_n) = D[L_{n-1}, L_1]$ and $(n-2)D(L_{-n}) = D[L_{-1}, L_{-n+1}]$, we get

$$D(L_n) = \begin{cases} (n-1)a(v_0 \otimes v_n + v_n \otimes v_0) - 2a \sum_{i=1}^{n-1} v_i \otimes v_{n-i}, & n \geq 2; \\ D(L_n) = (n+1)a(v_n \otimes v_0 + v_0 \otimes v_n) + 2a \sum_{i=n+1}^{-1} v_i \otimes v_{n-i}, & n \leq -2. \end{cases}$$

Then $D = a(-d_{0,0}^1 + d_{0,0}^2 - d_{0,0}^3 + d_{0,0}^4 + 2d_{0,0}^5)$. It follows that

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,0}^1 \oplus \mathbb{C}d_{0,0}^2 \oplus \mathbb{C}d_{0,0}^3 \oplus \mathbb{C}d_{0,0}^4 \oplus \mathbb{C}d_{0,0}^5.$$

Subcase 3.2: $\alpha = \beta = 1$. Then

$$D(L_2) = a_{2,1}v_1 \otimes v_1, \quad D(L_{-2}) = a_{-2,-1}v_{-1} \otimes v_{-1} = -a_{2,1}v_{-1} \otimes v_{-1}.$$

Using induction on n by

$$(n-2)D(L_n) = D[L_{n-1}, L_1] = -L_1 \cdot D(L_{n-1}), \quad (n-2)D(L_{-n}) = D[L_{-1}, L_{-n+1}] = L_{-1} \cdot D(L_{-n+1}),$$

we obtain

$$D(L_n) = a_{2,1} \sum_{i=1}^{n-1} i(n-i)v_i \otimes v_{n-i} = a_{2,1} \sum_{i=0}^{n-1} i(n-i)v_i \otimes v_{n-i}, \quad n \geq 0;$$

$$D(L_n) = -a_{2,1} \sum_{i=n}^0 i(n-i)v_i \otimes v_{n-i} = -a_{2,1} \sum_{i=n}^{-1} i(n-i)v_i \otimes v_{n-i}, \quad n \leq -1.$$

Then $D = a_{2,1}d_{1,1} \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1,1}$.

Subcase 3.3: $\alpha = \beta < 0$. Then $a_{-2,\alpha+k-2} = -a_{2,\alpha+k}$ for $0 \leq k \leq -2\alpha + 2$. It follows that $L_{\pm 1} \cdot \left(\sum_{i=\alpha}^{-\alpha} x_i v_i \otimes v_{-i} \right) = 0$ and $D(L_{\pm 2}) = L_{\pm 2} \cdot \left(\sum_{i=\alpha}^{-\alpha} x_i v_i \otimes v_{-i} \right)$, where $x_\alpha = -\frac{a_{2,\alpha}}{\alpha}$, $x_{\alpha+k} = \left(\prod_{t=1}^k \frac{2\alpha+t-1}{t} \right) x_\alpha$ for $1 \leq k \leq -2\alpha$. Hence $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$. \square

5.2. $D(L_{\mp 1}) = 0$ and $D(L_{\pm 1}) \neq 0$.

Lemma 5.2. *If $D(L_0) = D(L_{-1}) = 0$ and $D(L_1) \neq 0$, we have*

$$D \in \begin{cases} \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,0}^1 \oplus \mathbb{C}d_{0,0}^2 \oplus \mathbb{C}d_{0,0}^3 \oplus \mathbb{C}d_{0,0}^4 \oplus \mathbb{C}d_{0,0}^5, & (\alpha, \beta) = (0, 0); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1,0}^1 \oplus \mathbb{C}d_{1,0}, & (\alpha, \beta) = (1, 0); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1-\beta, \beta}, & \alpha = 1 - \beta, \beta \leq -1; \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0, & \text{otherwise.} \end{cases}$$

Proof. By (2.5), we have

$$(i + 1 - \alpha)a_{1,i+1} = (i - 1 + \beta)a_{1,i}. \quad (5.5)$$

Let $i = \beta - 1, l, l + 1$ in (5.5) respectively. Then

$$(\beta - \alpha)a_{1,\beta} = 0, \quad (l + 1 - \alpha)a_{1,l+1} = (l - 1 + \beta)a_{1,l}, \quad (l + \beta)a_{1,l+1} = 0. \quad (5.6)$$

Case 1: $\alpha \neq \beta$. Then $a_{1,\beta} = 0$ by (5.6) and $D(L_1) = a_{1,l+1}v_{l+1} \otimes v_{-l}$. Since $D(L_1) \neq 0$, we have $a_{1,l+1} \neq 0$. Therefore, we have $a_{1,l} = 0$ and $l = -\beta, \alpha = 1 - \beta$ via (5.6). For $\beta \leq l, \beta \leq 0$ and

$$D(L_1) = a_{1,-\beta+1}v_{-\beta+1} \otimes v_{\beta} = a_{1,\alpha}v_{\alpha} \otimes v_{-\alpha+1}.$$

By (2.6), we have

$$a_{2,-\beta+2} + a_{2,-\beta+1} = 3a_{1,-\beta+1}, \quad (5.7)$$

$$(i + \beta)a_{2,i+1} = (i - 2 + \beta)a_{2,i}, \quad i \neq -\beta + 1.$$

Since $\{a_{2,i} \neq 0 | i \in \mathbb{Z}\}$ is finite, we have $a_{2,i} = 0, i \neq -\beta + 1, -\beta + 2$. By (2.7), we have

$$(1 - 2\beta)a_{-2,-\beta} - (3 - 2\beta)a_{-2,-\beta+1} = -\beta a_{1,-\beta+1}, \quad (5.8)$$

$$(-1 - 2\beta)a_{-2,-\beta-2} - (1 - 2\beta)a_{-2,-\beta-1} = (\beta - 1)a_{1,-\beta+1}, \quad (5.9)$$

$$(i - \beta)a_{-2,i-1} = (i + 2 - \beta)a_{-2,i}, \quad i \neq -\beta - 1, -\beta + 1. \quad (5.10)$$

It is clear that $a_{-2,\beta} = a_{-2,\beta-3} = 0$ and $\beta a_{-2,-\beta-1} = (\beta - 1)a_{-2,-\beta}$. By (2.5), we have

$$(i + 2\beta)a_{2,i+2} - (i - 2 + 2\beta)a_{2,i} = (i - 2\beta)a_{-2,i-2} - (i + 2 - 2\beta)a_{-2,i}. \quad (5.11)$$

Furthermore, we have

$$(\beta - 1)a_{2,-\beta+1} = -(3\beta + 1)a_{-2,-\beta-3} + (3\beta - 1)a_{-2,-\beta-1}, \quad (5.12)$$

$$\beta a_{2,-\beta+2} = -3\beta a_{-2,-\beta-2} + (3\beta - 2)a_{-2,-\beta}, \quad (5.13)$$

$$-(\beta - 1)a_{2,-\beta+1} = -(3\beta - 1)a_{-2,-\beta-1} + (3\beta - 3)a_{-2,-\beta+1}, \quad (5.14)$$

$$-\beta a_{2,-\beta+2} = -(3\beta - 2)a_{-2,-\beta} + (3\beta - 4)a_{-2,-\beta+2}, \quad (5.15)$$

$$(i - 2\beta)a_{-2,i-2} = (i - 2\beta + 2)a_{-2,i}, \quad i \neq -\beta - 1, -\beta, -\beta + 1, -\beta + 2. \quad (5.16)$$

Subcase 1.1: $\beta = 0$. Then $\alpha = 1$. From (5.8)-(5.10), we get

$$a_{-2,-2} + a_{-2,-1} = a_{1,1}, \quad a_{-2,i} = 0, \quad i \neq -2, -1.$$

Using (5.12), we get $a_{-2,-1} = a_{2,1}$. Set $a = a_{1,1}, b = a_{2,1}$. Then

$$D(L_{-1}) = 0, \quad D(L_1) = a_{1,1}v_1 \otimes v_0 = av_1 \otimes v_0,$$

$$D(L_2) = bv_1 \otimes v_1 + (3a - b)v_2 \otimes v_0, \quad D(L_{-2}) = (a - b)v_{-2} \otimes v_0 + bv_{-1} \otimes v_{-1},$$

where $a_{2,1} + a_{2,2} = 3a_{1,1}$. Using induction on n by $(n-2)D(L_n) = D[L_{n-1}, L_1]$ and $(n-2)D(L_{-n}) = D[L_{-1}, L_{-n+1}]$, we obtain

$$D(L_n) = \begin{cases} b \sum_{i=1}^n iv_i \otimes v_{n-i} + \frac{a-b}{2}(n^2+n)v_n \otimes v_0, & n \geq 1, \\ -b \sum_{i=n+1}^{-1} iv_i \otimes v_{-i+n} + \frac{a-b}{2}(n^2+n)v_n \otimes v_0, & n \leq -2. \end{cases}$$

Furthermore,

$$\begin{aligned} D(L_n) &= b \sum_{i=1}^{n-1} iv_i \otimes v_{n-i} + bnv_n \otimes v_0 + \frac{a-b}{2}(n^2+n)v_n \otimes v_0 \\ &= b \sum_{i=1}^{n-1} iv_i \otimes v_{n-i} + \frac{a+b}{2}n^2v_n \otimes v_0 - \frac{a+b}{2}nv_n \otimes v_0 \\ &= bd_{1,0}(L_n) + \frac{a-b}{2}d_{1,0}^1(L_n) - \frac{a+b}{2}L_n \cdot (v_0 \otimes v_0), \quad n \geq 1; \\ D(L_0) &= 0 = bd_{1,0}(L_0) + \frac{a-b}{2}d_{1,0}^1(L_0) - \frac{a+b}{2}L_0 \cdot (v_0 \otimes v_0); \\ D(L_{-1}) &= 0 = bv_{-1} \otimes v_0 + \frac{a-b}{2}v_{-1} \otimes v_0 - \frac{a+b}{2}v_{-1} \otimes v_0 \\ &= bd_{1,0}(L_{-1}) + \frac{a-b}{2}d_{1,0}^1(L_{-1}) - \frac{a+b}{2}L_{-1} \cdot (v_0 \otimes v_{-1}); \\ D(L_n) &= -b \sum_{i=n+1}^{-1} iv_i \otimes v_{-i+n} + \frac{a-b}{2}(n^2+n)v_n \otimes v_0 \\ &= b(d_{1,0}(L_n) + nv_n \otimes v_0) + \frac{a-b}{2}d_{1,0}^1(L_n) + \frac{a-b}{2}nv_n \otimes v_0 \\ &= bd_{1,0}(L_n) + \frac{a-b}{2}d_{1,0}^1(L_n) + \frac{a+b}{2}nv_n \otimes v_0 \\ &= bd_{1,0}(L_n) + \frac{a-b}{2}d_{1,0}^1(L_n) - \frac{a+b}{2}L_n \cdot (v_0 \otimes v_0), \quad n \leq -2. \end{aligned}$$

Then $D = bd_{1,0} + \frac{a-b}{2}d_{1,0}^1 - \frac{a+b}{2}D_{inn}$, where $D_{inn}(L_n) = L_n \cdot (v_0 \otimes v_0)$ for all $n \in \mathbb{Z}$. So

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1,0} \oplus \mathbb{C}d_{1,0}^1.$$

Subcase 1.2: $\beta = -1$. Then $\alpha = 2$. Using (5.8)-(5.16) and setting $a = \frac{a_{1,2}}{3}$, we obtain

$$D(L_{-1}) = 0, \quad D(L_1) = a(3v_2 \otimes v_{-1}),$$

$$D(L_{-2}) = a(2v_0 \otimes v_{-2} + v_1 \otimes v_{-3}), \quad D(L_2) = a(4v_2 \otimes v_0 + 5v_3 \otimes v_{-1}).$$

Using induction on n by $(n-2)D(L_n) = D[L_{n-1}, L_1]$ and $(n-2)D(L_{-n}) = D[L_{-1}, L_{-n+1}]$, we obtain

$$D(L_n) = a \sum_{i=2}^{n+1} (n+i)v_i \otimes v_{n-i}, \quad n \geq 1; \quad D(L_n) = -a \sum_{i=n+2}^1 (n+i)v_i \otimes v_{n-i}, \quad n \leq -2.$$

Then $D = ad_{2,-1} - aD_{inn}$, where $D_{inn}(L_n) = L_n \cdot \left(\sum_{i=0}^1 v_i \otimes v_{-i} \right)$ for all $n \in \mathbb{Z}$. So

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{2,-1}.$$

Subcase 1.3: $\beta \leq -2$. Then $\alpha = 1 - \beta \geq 3$, $\beta < -\beta - 2$ and

$$\dots, \beta - 3, \beta - 2, \beta - 1, \beta, \dots, -\beta - 2, -\beta - 1, -\beta, -\beta + 1, \dots$$

Since $\{a_{-2,i} \neq 0 | i \in \mathbb{Z}\}$ is finite, by (5.10), we deduce that

$$a_{-2,\beta} = 0, \quad a_{-2,i} = 0, \quad i \leq \beta - 3, \text{ or } i \geq -\beta + 1.$$

Furthermore, we have $a_{-2,i} = 0$ for $\beta + 1 \leq i \leq -\beta - 2$. Use (5.8)-(5.10), then we obtain

$$a_{-2,\beta-2} = -a_{-2,\beta-1}, \quad a_{-2,-\beta} = \frac{-\beta}{1-2\beta} a_{1,-\beta+1}, \quad a_{-2,-\beta-1} = \frac{1-\beta}{1-2\beta} a_{1,-\beta+1}. \quad (5.17)$$

Let $i = \beta, \beta + 1$ in (5.16) respectively. Then $a_{-2,\beta-2} = a_{-2,\beta-1} = 0$. Since $\beta \leq -2$, we have $\beta < -\beta - 3 < -\beta - 2$. So $a_{-2,-\beta-3} = 0$. By (5.12)-(5.13), we have

$$(\beta - 1)a_{2,-\beta+1} = (3\beta - 1)a_{-2,-\beta-1}, \quad \beta a_{2,-\beta+2} = (3\beta - 2)a_{-2,-\beta}.$$

Using (5.17),

$$a_{2,-\beta+1} = \frac{1-3\beta}{1-2\beta} a_{1,-\beta+1}, \quad a_{2,-\beta+2} = \frac{2-3\beta}{1-2\beta} a_{1,-\beta+1}.$$

Set $a = \frac{a_{1,-\beta+1}}{1-2\beta}$. It follows that $D(L_{-1}) = 0$ and

$$\begin{aligned} D(L_{-2}) &= a \left(-\beta v_{-\beta} \otimes v_{\beta-2} + (1-\beta)v_{-\beta-1} \otimes v_{\beta-1} \right), & D(L_1) &= a \left((1-2\beta)v_{-\beta+1} \otimes v_\beta \right), \\ D(L_2) &= a \left((1-3\beta)v_{-\beta+1} \otimes v_{\beta+1} + (2-3\beta)v_{-\beta+2} \otimes v_\beta \right). \end{aligned}$$

Using induction on n by $(n-2)D(L_n) = D[L_{n-1}, L_1]$ and $(n-2)D(L_{-n}) = D[L_{-1}, L_{-n+1}]$, we have

$$D(L_n) = a \sum_{i=-\beta+1}^{-\beta+n} (i-n\beta)v_i \otimes v_{n-i}, \quad n \geq 1; \quad D(L_n) = -a \sum_{i=-\beta+1+n}^{-\beta} (i-n\beta)v_i \otimes v_{n-i}, \quad n \leq -1.$$

Then $D(L_n) = ad_{1-\beta,\beta}(L_n) + L_n \cdot \left(-a \sum_{i=0}^{-\beta} v_i \otimes v_{-i} \right)$. So $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1-\beta,\beta}$.

Case 2: $\alpha = \beta$. By (5.6), we have

$$(l+1-\beta)a_{1,l+1} = (l-1+\beta)a_{1,l}, \quad (l+\beta)a_{1,l+1} = 0. \quad (5.18)$$

Subcase 2.1: $l = -\beta$. Then $(1-2\beta)a_{1,-\beta+1} = -a_{1,-\beta}$ via (5.18) and

$$D(L_1) = a_{1,\beta}v_\beta \otimes v_{-\beta+1} + a_{1,-\beta+1}v_{-\beta+1} \otimes v_\beta.$$

Assume $a_{1,-\beta+1} = 0$. Then $a_{1,-\beta} = 0$ and $D(L_1) = a_{1,\beta}v_\beta \otimes v_{-\beta+1}$, which suggests that $\beta < -\beta$, i.e., $\beta < 0$. Using (2.7), we have

$$-\beta a_{1,\beta} = (2\beta - 3)a_{-2,\beta-3}, \quad (5.19)$$

$$(i-1+\beta)a_{-2,i-1} = (i+2-\beta)a_{-2,i}, \quad i \neq \beta - 2, \beta. \quad (5.20)$$

Since $\{a_{-2,i} \neq 0 | i \in \mathbb{Z}\}$ is finite, by (5.20), we have

$$a_{-2,i} = 0, \quad i \leq \beta - 3, \quad \text{or} \quad i \geq -\beta + 1.$$

Then $a_{1,\beta} = 0$ via (5.19). Hence $D(L_1) = 0$, a contradiction. It follows that $a_{1,-\beta+1} \neq 0$. Hence $a_{1,-\beta} \neq 0$, which implies that $-\beta = \beta$. Then $\alpha = \beta = l = 0$. Hence $D(L_1) = a_{1,0}v_0 \otimes v_1 + a_{1,1}v_1 \otimes v_0$ and $a_{1,0} = -a_{1,1}$. By (2.5)-(2.7), we have

$$3a_{1,i} = (i+1)a_{2,i+1} - (i-2)a_{2,i}, \quad (5.21)$$

$$(i+2)a_{1,i+2} - (i-1)a_{1,i} = (i-1)a_{-2,i-1} - (i+2)a_{-2,i}, \quad (5.22)$$

$$(i+2)(a_{2,i+2} + a_{-2,i}) = (i-2)(a_{-2,i-2} + a_{2,i}). \quad (5.23)$$

Using (5.21), we have

$$3a_{1,0} = a_{2,1} + 2a_{2,0}, \quad 3a_{1,1} = 2a_{2,2} + a_{2,1}, \quad (5.24)$$

$$(i+1)a_{2,i+1} = (i-2)a_{2,i}, \quad i \neq 0, 1.$$

It is clear that $a_{2,i} = 0$ for $i \neq 0, 1, 2$. Using (5.22), we have

$$-a_{1,1} = a_{-2,-1} + 2a_{-2,-2}, \quad -a_{1,0} = a_{-2,-1} + 2a_{-2,0}, \quad (5.25)$$

$$(i-1)a_{-2,i-1} = (i+2)a_{-2,i}, \quad i \neq -1, 0.$$

It is clear that $a_{-2,i} = 0$ for $i \neq -2, -1, 0$. Let $i = -1, 0$ in (5.23) respectively. Then

$$a_{-2,-1} = -a_{2,1}, \quad a_{2,2} + a_{2,0} = -a_{-2,-2} - a_{-2,0}. \quad (5.26)$$

Set $a = \frac{a_{1,0}}{2}, b = \frac{a_{2,1}}{2}$. Then $a_{1,0} = 2a, a_{1,1} = -2a$ and

$$a_{2,0} = 3a - b, \quad a_{2,1} = 2b, \quad a_{2,2} = -3a - b, \quad a_{-2,-2} = a + b, \quad a_{-2,-1} = -2b, \quad a_{-2,0} = b - a$$

via (5.24)-(5.26). Therefore,

$$D(L_{-1}) = 0, \quad D(L_{-2}) = (a+b)v_{-2} \otimes v_0 - 2bv_{-1} \otimes v_{-1} - (a-b)v_0 \otimes v_{-2},$$

$$D(L_1) = 2av_0 \otimes v_1 - 2av_1 \otimes v_0, \quad D(L_2) = (3a-b)v_0 \otimes v_2 + 2bv_1 \otimes v_1 - (3a+b)v_2 \otimes v_0.$$

Using induction on n by $(n-2)D(L_n) = D[L_{n-1}, L_1]$ and $(n-2)D(L_{-n}) = D[L_{-1}, L_{-n+1}]$, we have

$$D(L_n) = (n+1)a(v_0 \otimes v_n - v_n \otimes v_0) - (n-1)b(v_0 \otimes v_n + v_n \otimes v_0) + 2b \sum_{i=1}^{n-1} v_i \otimes v_{n-i}, \quad n \geq 2;$$

$$D(L_n) = -(n+1)a(v_n \otimes v_0 - v_0 \otimes v_n) - (n+1)b(v_n \otimes v_0 + v_0 \otimes v_n) - 2b \sum_{i=n+1}^{-1} v_i \otimes v_{n-i}, \quad n \leq -2.$$

Then

$$D = a(d_{0,0}^1 + d_{0,0}^2 - d_{0,0}^3 - d_{0,0}^4) - b(-d_{0,0}^1 + d_{0,0}^2 - d_{0,0}^3 + d_{0,0}^4 + 2d_{0,0}^5).$$

Hence $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,0}^1 \oplus \mathbb{C}d_{0,0}^2 \oplus \mathbb{C}d_{0,0}^3 \oplus \mathbb{C}d_{0,0}^4 \oplus \mathbb{C}d_{0,0}^5$.

Subcase 2.2: $l \neq -\beta$. By (5.18), we have $a_{1,l+1} = 0$ and $(l-1+\beta)a_{1,l} = 0$. Then $a_{1,\beta} \neq 0$,

$$D(L_1) = a_{1,\beta}v_\beta \otimes v_{-\beta+1}.$$

If $\beta < 0$, we deduce $a_{1,\beta} = 0$ similar to the proof of Subcase 2.1. So $\beta \geq 0$.

If $\beta = 0$, similar to the proof of Subcase 2.1, we get

$$D(L_2) = a_{2,0}v_0 \otimes v_2 + a_{2,1}v_1 \otimes v_1 + a_{2,2}v_2 \otimes v_0,$$

$$D(L_{-2}) = a_{-2,-2}v_{-2} \otimes v_0 + a_{-2,-1}v_{-1} \otimes v_{-1} + a_{-2,0}v_0 \otimes v_{-2},$$

where

$$a_{-2,-1} = -2a_{-2,-2}, \quad a_{-2,0} = a_{-2,-2} - \frac{a_{1,0}}{2}, \quad a_{2,0} = a_{2,2} + \frac{3a_{1,0}}{2}, \quad a_{2,1} = -2a_{2,2}.$$

Let $m = 2, i = 1$ in (2.5). Then $a_{-2,-1} = -a_{2,1}$. So $a_{-2,-2} = -a_{2,2}$. Let $m = 2, i = 0$ in (2.5). Then $a_{2,2} + a_{-2,-2} = -a_{2,0} - a_{-2,0}$. Hence $a_{-2,0} + a_{2,0} = 0$, which forces $a_{1,\beta} = 0$, a contradiction.

If $\beta > 0$, by (2.6), we obtain

$$a_{2,\beta+1} + (2 - 2\beta)a_{2,\beta} = 3a_{1,\beta}, \quad (5.27)$$

$$(i + 1 - \beta)a_{2,i+1} = (i - 2 + \beta)a_{2,i}, \quad i \neq \beta.$$

Since $\{a_{2,i} \neq 0 | i \in \mathbb{Z}\}$ is finite, we deduce $a_{2,i} = 0$ for $i \leq \beta - 1$ or $i \geq 3 - \beta$. When $\beta - 1 < 3 - \beta$, we have $\beta < 2$. So $\beta = 1$ and $a_{1,\beta} = 0$, which is a contradiction. \square

Lemma 5.3. *If $D(L_0) = D(L_1) = 0$ and $D(L_{-1}) \neq 0$, we have*

$$D \in \begin{cases} \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,0}^1 \oplus \mathbb{C}d_{0,0}^2 \oplus \mathbb{C}d_{0,0}^3 \oplus \mathbb{C}d_{0,0}^4 \oplus \mathbb{C}d_{0,0}^5, & (\alpha, \beta) = (0, 0); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,1}^1 \oplus \mathbb{C}d_{0,1}, & (\alpha, \beta) = (0, 1); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1,0}^1 \oplus \mathbb{C}d_{1,0}, & (\alpha, \beta) = (1, 0); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1-\beta,\beta}, & \beta \in \mathbb{Z}, \beta \neq 0, 1; \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0, & \text{otherwise.} \end{cases}$$

Proof. Similar to the proof of Lemma 5.2. \square

5.3. $D(L_{\pm 1}) \neq 0$. Next, we always assume $D(L_{-1}) \neq 0$ and denote

$$p_{-1} = \min\{i | a_{-1,i} \neq 0\}, \quad q_{-1} = \max\{i | a_{-1,i} \neq 0\}.$$

Lemma 5.4. *If $D(L_0) = 0$ and $D(L_{\pm 1}) \neq 0$, we have*

$$D \in \begin{cases} \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,0}^1 \oplus \mathbb{C}d_{0,0}^2 \oplus \mathbb{C}d_{0,0}^3 \oplus \mathbb{C}d_{0,0}^4 \oplus \mathbb{C}d_{0,0}^5, & (\alpha, \beta) = (0, 0); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,1} \oplus \mathbb{C}d_{0,1}^1, & (\alpha, \beta) = (0, 1); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1,0} \oplus \mathbb{C}d_{1,0}^1, & (\alpha, \beta) = (1, 0); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,2}, & (\alpha, \beta) = (0, 2); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{2,0}, & (\alpha, \beta) = (2, 0); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1,1}, & (\alpha, \beta) = (1, 1); \\ \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0, & \text{otherwise.} \end{cases}$$

Proof. Let $m = 1$ in (2.5). Then

$$(i + 1 - \alpha)a_{1,i+1} - (i - 1 + \beta)a_{1,i} = (i - 1 + \alpha)a_{-1,i-1} - (i + 1 - \beta)a_{-1,i}. \quad (5.28)$$

Let $i = \beta - 1, \beta, \beta + 1, l, l + 1$. Then

$$(\beta - \alpha)a_{1,\beta} = (\alpha + \beta - 2)a_{-1,\beta-2}, \quad (5.29)$$

$$(\beta + 1 - \alpha)a_{1,\beta+1} - (2\beta - 1)a_{1,\beta} = (\alpha + \beta - 1)a_{-1,\beta-1} - a_{-1,\beta}, \quad (5.30)$$

$$(\beta + 2 - \alpha)a_{1,\beta+2} - 2\beta a_{1,\beta+1} = (\alpha + \beta)a_{-1,\beta} - 2a_{-1,\beta+1}, \quad (5.31)$$

$$(l + 1 - \alpha)a_{1,l+1} - (l - 1 + \beta)a_{1,l} = (l - 1 + \alpha)a_{-1,l-1} - (l + 1 - \beta)a_{-1,l}, \quad (5.32)$$

$$-(l + \beta)a_{1,l+1} = (l + \alpha)a_{-1,l} - (l + 2 - \beta)a_{-1,l+1}, \quad (5.33)$$

and

$$(i - 1 + \alpha)a_{-1,i-1} = (i + 1 - \beta)a_{-1,i}, \quad i \leq \beta - 2, \quad \text{or} \quad i \geq l + 2. \quad (5.34)$$

Assume $p_{-1} \leq \beta - 2$. Let $i = p_{-1}$ in (5.34). Then $p_{-1} = \beta - 1 \geq \beta - 2$, which is a contradiction. Therefore, $p_{-1} \geq \beta - 1$, which suggests

$$a_{-1,i} = 0, \quad i \leq \beta - 2.$$

Hence (5.29) becomes

$$(\beta - \alpha)a_{1,\beta} = 0. \quad (5.35)$$

Case 1: $\alpha \neq \beta$. Then $a_{1,\beta} = 0$ and $D(L_1) = a_{1,l+1}v_{l+1} \otimes v_{-l}$, where $\beta \leq l$. Moreover, by (4.2)-(4.4),

$$(l + 1 - \alpha)a_{1,l+1} = (l - 1 + \alpha)a_{-1,l-1} - (l + 1 - \beta)a_{-1,l},$$

$$-(l + \beta)a_{1,l+1} = (l + \alpha)a_{-1,l} - (l + 2 - \beta)a_{-1,l+1},$$

$$(i - 1 + \alpha)a_{-1,i-1} = (i + 1 - \beta)a_{-1,i}, \quad i \neq l, l + 1.$$

Similar to the proof of Lemma 4.2, we have the following result:

$$p_{-1} = \beta - 1, l, \text{ or } l + 1; \quad q_{-1} = -\alpha \quad \text{if } q_{-1} \neq l - 1, l.$$

Subcase 1.1: $p_{-1} = \beta - 1$ and $q_{-1} = l - 1$. Then $a_{-1,i} = 0$ for $i \geq l$. According to (4.2)-(4.3), we have

$$(l + 1 - \alpha)a_{1,l+1} = (l - 1 + \alpha)a_{-1,l-1}, \quad (l + \beta)a_{1,l+1} = 0. \quad (5.36)$$

Since $D(L_1) \neq 0$, that is, $a_{1,l+1} \neq 0$, we have $\beta = -l$. Since $\beta \leq l$, we have $\beta \leq 0$ and

$$D(L_1) = a_{1,-\beta+1}v_{-\beta+1} \otimes v_\beta, \quad D(L_{-1}) = \sum_{i=-\beta-1}^{-\beta-1} a_{-1,i}v_i \otimes v_{-i-1}.$$

(1.1.1) If $\beta = 0$ and $\alpha \neq 0, 1$, we have $a_{-1,-1} = -a_{1,1}$ and

$$D(L_1) = a_{1,1}v_1 \otimes v_0 = L_1 \cdot \left(-\frac{a_{1,1}}{\alpha} v_0 \otimes v_0 \right), \quad D(L_{-1}) = -a_{1,1}v_{-1} \otimes v_0 = L_{-1} \cdot \left(-\frac{a_{1,1}}{\alpha} v_0 \otimes v_0 \right).$$

Replace D by $D - D_{\text{inn}}$, where $D_{\text{inn}}(L_n) = L_n \cdot \left(-\frac{a_{1,1}}{\alpha} v_0 \otimes v_0\right)$ for all $n \in \mathbb{Z}$, then $D(L_0) = D(L_{\pm 1}) = 0$. According to the result of Lemma 5.1, we know $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \delta_{\alpha,2} \mathbb{C}d_{2,0}$.

(1.1.2) If $\beta = 0$ and $\alpha = 1$, we have

$$D(L_1) = L_1 \cdot (-a_{1,1} v_0 \otimes v_0), \quad D(L_{-1}) = L_{-1} \cdot (-a_{1,1} v_0 \otimes v_0) + (a_{1,1} + a_{-1,-1}) v_{-1} \otimes v_0.$$

Set $a = a_{1,1} + a_{-1,-1}$. Then it suffices to consider $D(L_0) = D(L_1) = 0$ and $D(L_{-1}) = av_{-1} \otimes v_0$. By Lemma 5.1 and Lemma 5.3, we have $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1,0} \oplus \mathbb{C}d_{1,0}^1$.

(1.1.3) If $\beta < 0$, we have

$$D(L_1) = a_{1,-\beta+1} v_{-\beta+1} \otimes v_\beta, \quad D(L_{-1}) = \sum_{i=\beta-1}^{-\beta-1} a_{-1,i} v_i \otimes v_{-i-1},$$

where

$$a_{-1,\beta+i} = \left(\prod_{t=1}^{i+1} \frac{\alpha + \beta - 2 + t}{t} \right) a_{-1,\beta-1}, \quad 0 \leq i \leq -2\beta - 1. \quad (5.37)$$

Since $a_{-1,-\beta-1} \neq 0$, we have $\alpha + \beta + i \neq 0$ for $-1 \leq i \leq -2\beta - 2$. Hence $\alpha - i \neq 0$ for $\beta + 2 \leq i \leq -\beta + 1$. Moreover, by (5.36), we have

$$a_{1,-\beta+1} = -\frac{\alpha - \beta - 1}{\alpha + \beta - 1} a_{-1,-\beta-1}. \quad (5.38)$$

Assume $\alpha = \beta + 1$. Then $a_{1,-\beta+1} = 0$ by (5.38) and $D(L_1) = 0$, which is a contradiction. Therefore $\alpha \neq \beta + 1$. Then $D(L_1) \neq 0$ and we deduce that

$$D(L_{\pm 1}) = L_{\pm 1} \cdot \left(\sum_{i=\beta}^{-\beta} c_i v_i \otimes v_{-i} \right),$$

where

$$c_\beta = \frac{a_{-1,\beta-1}}{\alpha - \beta}, \quad c_{\beta+i} = \left(\prod_{t=1}^i \frac{\alpha + \beta + t - 1}{t} \right) \frac{a_{-1,\beta-1}}{\alpha - \beta}, \quad 1 \leq i \leq -2\beta.$$

According to Lemma 5.1, we have $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$.

Subcase 1.2: $p_{-1} = \beta - 1$ and $q_{-1} = l$. Then

$$D(L_1) = a_{1,l+1} v_{l+1} \otimes v_{-l} \neq 0, \quad D(L_{-1}) = \sum_{i=\beta-1}^l a_{-1,i} v_i \otimes v_{-i-1} \neq 0.$$

According to (4.2)-(4.4), we have

$$(l+1-\alpha)a_{1,l+1} = (l-1+\alpha)a_{-1,l-1} - (l+1-\beta)a_{-1,l}, \quad (5.39)$$

$$(l+\beta)a_{1,l+1} = -(l+\alpha)a_{-1,l}, \quad (5.40)$$

$$(i-1+\alpha)a_{-1,i-1} = (i+1-\beta)a_{-1,i}, \quad i \neq l, l+1. \quad (5.41)$$

Assume $\beta = -l$, by (5.40), we have $(l+\alpha)a_{-1,l} = 0$. Since $a_{-1,l} = a_{-1,q_{-1}} \neq 0$, we have $\alpha = -l = \beta$, which is a contradiction. Therefore $\beta \neq -l$. By (5.40), we have

$$a_{1,l+1} = -\frac{l+\alpha}{l+\beta} a_{-1,l}, \quad D(L_1) = -\frac{l+\alpha}{l+\beta} a_{-1,l} v_{l+1} \otimes v_{-l}.$$

Since $a_{1,l+1} \neq 0$, we have $\alpha \neq -l$. Hence we obtain

$$\frac{(\alpha - \beta)(\alpha + \beta - 1)}{l + \beta} a_{-1,l} = (l - 1 + \alpha) a_{-1,l-1}. \quad (5.42)$$

Let $i = \beta + t$ ($-1 \leq t \leq l - \beta - 1$) in (5.41). Then

$$(t + 1) a_{-1,\beta+t} = (\alpha + \beta + t - 1) a_{-1,\beta+t-1}, \quad -1 \leq t \leq l - \beta - 1. \quad (5.43)$$

(1.2.1) $\alpha + \beta \neq 1$. Then $(l - 1 + \alpha) a_{-1,l-1} \neq 0$ via (5.42).

If $\beta = l$, we have $\beta \neq 0$ and

$$D(L_1) = a_{1,\beta+1} v_{\beta+1} \otimes v_{-\beta}, \quad D(L_{-1}) = a_{-1,\beta-1} v_{\beta-1} \otimes v_{-\beta} + a_{-1,\beta} v_{\beta} \otimes v_{-\beta-l},$$

where

$$a_{1,\beta+1} = -\frac{\alpha + \beta}{2\beta} a_{-1,\beta}, \quad a_{-1,\beta-1} = \frac{\alpha - \beta}{2\beta} a_{-1,\beta}.$$

Obviously,

$$D(L_{\pm 1}) = L_{\pm 1} \cdot \left(\frac{a_{-1,\beta}}{2\beta} v_{\beta} \otimes v_{-\beta} \right).$$

Using Lemma 5.1, we know when $\beta = 2$ we have $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_{\alpha} \otimes \mathcal{F}_{\beta})_0 \oplus \mathbb{C}d_{0,2}$. Otherwise, $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_{\alpha} \otimes \mathcal{F}_{\beta})_0$.

If $\beta < l$, we have

$$a_{-1,l} = \frac{(l + \beta)(l - 1 + \alpha)}{(\alpha - \beta)(\alpha + \beta - 1)} a_{-1,l-1}.$$

Using (5.43), we get

$$a_{-1,\beta+i} = \frac{\alpha + \beta + i - 1}{i + 1} \cdots \frac{\alpha + \beta}{2} \cdot \frac{\alpha + \beta - 1}{1} \cdot a_{-1,\beta-1}, \quad 0 \leq i \leq l - 1 - \beta.$$

We check that $D(L_{\pm 1}) = L_{\pm 1} \cdot \left(\sum_{i=\beta}^l c_i v_i \otimes v_{-i} \right)$, where

$$c_{\beta} = \frac{a_{-1,\beta-1}}{\alpha - \beta}, \quad c_{\beta+i} = \frac{\alpha + \beta + i - 1}{i} \cdots \frac{\alpha + \beta + 1}{2} \cdot \frac{\alpha + \beta}{1} \cdot \frac{a_{-1,\beta-1}}{\alpha - \beta}, \quad 0 \leq i \leq l - \beta.$$

Using the result of Lemma 5.1, we have $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_{\alpha} \otimes \mathcal{F}_{\beta})_0$.

(1.2.2) $\alpha + \beta = 1$. Since $\beta \leq l$, we have $l + 1 - \beta \neq 0$. Then

$$D(L_1) = a_{1,l+1} v_{l+1} \otimes v_{-l} = L_1 \cdot \left(-\frac{a_{1,l+1}}{l + 1 - \beta} \sum_{i=\beta}^l v_i \otimes v_{-i} \right).$$

Replacing D by $D - D_{\text{inn}}$, where

$$D_{\text{inn}}(L_n) = L_n \cdot \left(-\frac{a_{1,l+1}}{l + 1 - \beta} \sum_{i=\beta}^l v_i \otimes v_{-i} \right)$$

for all $n \in \mathbb{Z}$, we have $D(L_1) = 0$. From Lemma 5.1 and Lemma 5.3, we obtain

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_{\alpha} \otimes \mathcal{F}_{\beta})_0 \oplus \delta_{\alpha,0} \delta_{\beta,1} \mathbb{C}d_{0,1}^1 \oplus \delta_{\alpha,1} \delta_{\beta,0} \mathbb{C}d_{1,0}^1.$$

Subcase 1.3: $p_{-1} = \beta - 1$ and $q_{-1} = -\alpha$. Then $\alpha + \beta \leq 1$,

$$D(L_1) = a_{1,l+1}v_{l+1} \otimes v_{-l}, \quad D(L_{-1}) = \sum_{i=\beta-1}^{-\alpha} a_{-1,i}v_i \otimes v_{-i-1}.$$

Notice that $-\alpha \neq l - 1, l$. Then $-\alpha < l - 1$ or $-\alpha \geq l + 1$.

(1.3.1) If $-\alpha < l - 1$, we have $(l + \beta)a_{1,l+1} = 0$ and $(l + 1 - \alpha)a_{1,l+1} = 0$ by (4.2) and (4.3). Since $a_{1,l+1} \neq 0$, we have $l = -\beta, \alpha = 1 - \beta$. Since $\beta \leq l$, we have $\beta \leq 0$. So $D(L_1) = a_{1,-\beta+1}v_{-\beta+1} \otimes v_\beta$ and $D(L_{-1}) = a_{-1,\beta-1}v_{\beta-1} \otimes v_{-\beta}$. Then

$$\begin{aligned} D(L_{-1}) &= L_{-1} \cdot \left(\frac{a_{-1,\beta-1}}{1-2\beta} \sum_{i=\beta}^{-\beta} v_i \otimes v_{-i} \right), \\ D(L_1) &= L_1 \cdot \left(\frac{a_{-1,\beta-1}}{1-2\beta} \sum_{i=\beta}^{-\beta} v_i \otimes v_{-i} \right) + (a_{-1,\beta-1} + a_{1,-\beta+1})v_{-\beta+1} \otimes v_\beta. \end{aligned}$$

Replacing D by $D - D_{inn}$, where

$$D_{inn}(L_n) = L_n \cdot \left(\frac{a_{-1,\beta-1}}{1-2\beta} \sum_{i=\beta}^{-\beta} v_i \otimes v_{-i} \right)$$

for all $n \in \mathbb{Z}$, we have $D(L_{-1}) = 0$. By Lemma 5.1 and Lemma 5.2, we get

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1-\beta,\beta} \oplus \delta_{\alpha,1}\delta_{\beta,0}\mathbb{C}d_{1,0}^1.$$

(1.3.2) If $-\alpha \geq l + 1$, we have $\alpha + \beta \leq -1$ by $\beta \leq l$. Assume $\alpha + \beta = -k$, where $k \geq 1$. Then $-\alpha = \beta + k \geq l + 1$. So $l + 1 - k \leq \beta \leq l$. For $\beta = l + t - k$, where $1 \leq t \leq k$, we have $\alpha = -l - t$ and $D(L_1) = L_1 \cdot \left(\sum_{i=1}^t c_i v_{l+i} \otimes v_{-l-i} \right)$, where

$$c_1 = \frac{a_{1,l+1}}{k+1-t}, \quad c_i = \left(\prod_{j=2}^i \frac{t+1-j}{t-k-j} \right) \frac{a_{1,l+1}}{k+1-t}, \quad 2 \leq i \leq t.$$

Replace D by $D - D_{inn}$, where $D_{inn}(L_n) = L_n \cdot \left(\sum_{i=1}^t c_i v_{l+i} \otimes v_{-l-i} \right)$ for all $n \in \mathbb{Z}$. Then $D(L_1) = 0$. Using the results of Lemma 5.1 and Lemma 5.3, we get $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$.

Subcase 1.4: $p_{-1} = l$ and $q_{-1} = l$. Then $D(L_1) = a_{1,l+1}v_{l+1} \otimes v_{-l}$ and $D(L_{-1}) = a_{-1,l}v_l \otimes v_{-l-1}$. By (4.2)-(4.3), we have

$$(l+1-\alpha)a_{1,l+1} = -(l+1-\beta)a_{-1,l}, \quad (l+\beta)a_{1,l+1} = -(l+\alpha)a_{-1,l}, \quad (5.44)$$

$$(\alpha + \beta - 1)(a_{1,l+1} + a_{-1,l}) = 0.$$

Assume $\alpha + \beta \neq 1$. Then $a_{-1,l} = -a_{1,l+1} \neq 0$. By (5.44), we have $\alpha = \beta$, which is a contradiction. Hence $\alpha + \beta = 1$. So $\beta \neq -l, l + 1$ by (5.44) and $a_{1,l+1} = -\frac{l+1-\beta}{l+\beta}a_{-1,l}$. Then

$$D(L_1) = L_1 \cdot \left(\sum_{i=\beta}^l \frac{a_{-1,l}}{l+\beta} v_i \otimes v_{-i} \right), \quad D(L_{-1}) = L_{-1} \cdot \left(\sum_{i=\beta}^l \frac{a_{-1,l}}{l+\beta} v_i \otimes v_{-i} \right) - \frac{a_{-1,l}}{l+\beta} v_{\beta-1} \otimes v_{-\beta}.$$

Replacing D by $D - D_{inn}$, where

$$D_{inn}(L_n) = L_n \cdot \left(\sum_{i=\beta}^l \frac{a_{-1,i}}{l+\beta} v_i \otimes v_{-i} \right)$$

for all $n \in \mathbb{Z}$, we have $D(L_1) = 0$ and $D(L_{-1}) \neq 0$. Using Lemma 5.3, we get

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{1-\beta,\beta} \oplus \delta_{\alpha,1}\delta_{\beta,0}\mathbb{C}d_{1,0}^1.$$

Subcase 1.5: $p_{-1} = l$ and $q_{-1} = -\alpha > l$. Then $\alpha < -l$ and

$$D(L_1) = a_{1,l+1}v_{l+1} \otimes v_{-l}, \quad D(L_{-1}) = \sum_{i=l}^{-\alpha} a_{-1,i}v_i \otimes v_{-i-1}.$$

By (4.2)-(4.3), we have

$$(l+1-\alpha)a_{1,l+1} = -(l+1-\beta)a_{-1,l}, \quad (l+\beta)a_{1,l+1} = -(l+\alpha)a_{-1,l} + (l+2-\beta)a_{-1,l+1}, \quad (5.45)$$

$$(\alpha+\beta-1)(a_{1,l+1} + a_{-1,l}) = (l+2-\beta)a_{-1,l+1}. \quad (5.46)$$

Assume $l+1-\alpha = 0$. Since $a_{1,l+1} \neq 0$, $a_{-1,l} \neq 0$, we have $l+1-\beta = 0$ via (5.45). Hence $\alpha = \beta$, a contradiction. Therefore, $l+1-\alpha \neq 0$. Moreover, $l+1-\beta \neq 0$ and

$$a_{1,l+1} = -\frac{l+1-\beta}{l+1-\alpha}a_{-1,l} \neq -a_{-1,l}, \quad a_{1,l+1} + a_{-1,l} = \frac{\beta-\alpha}{l+1-\alpha}a_{-1,l} \neq 0.$$

(1.5.1) If $\alpha + \beta \neq 1$, by (5.46), we have $l+2-\beta \neq 0$ and

$$a_{-1,l+1} = \frac{\alpha+\beta-1}{l+2-\beta} \cdot \frac{\beta-\alpha}{l+1-\alpha}a_{-1,l} \neq 0.$$

When $-\alpha = l+1$, we have $a_{1,l+1} = -\frac{\alpha+\beta}{2\alpha}a_{-1,l}$, $a_{-1,l+1} = -\frac{\alpha-\beta}{2\alpha}a_{-1,l}$ and

$$D(L_1) = -\frac{a_{-1,l}}{2\alpha}((\alpha+\beta)v_{-\alpha} \otimes v_{\alpha+1}), \quad D(L_{-1}) = -\frac{a_{-1,l}}{2\alpha}(-2\alpha v_{-\alpha-1} \otimes v_\alpha + (\alpha-\beta)v_{-\alpha} \otimes v_{\alpha-1}).$$

Then we check that $D(L_{\pm 1}) = L_{\pm 1} \cdot \left(\frac{a_{-1,l}}{2\alpha}v_{-\alpha} \otimes v_\alpha \right)$. Replace D by $D - D_{inn}$, where $D_{inn}(L_n) = L_n \cdot \left(\frac{a_{-1,l}}{2\alpha}v_{-\alpha} \otimes v_\alpha \right)$ for all $n \in \mathbb{Z}$. Then $D(L_{\pm 1}) = 0$. Using the result of Lemma 5.1 and noticing $\beta \leq l$, we get $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$.

When $-\alpha = l+k$, where $k \geq 2$, we have

$$a_{1,l+1} = -\frac{\alpha+\beta+k-1}{2\alpha+k-1}a_{-1,l}, \quad a_{-1,l+1} = -\frac{\alpha+\beta-1}{\alpha+\beta+k-2} \cdot \frac{\alpha-\beta}{2\alpha+k-1}a_{-1,l}.$$

Moreover, by (4.4), we have

$$a_{-1,l+i} = -\left(\prod_{t=k+1-i}^{k-1} \frac{t}{\alpha+\beta+t-2} \right) \frac{\alpha+\beta-1}{\alpha+\beta+k-2} \cdot \frac{\alpha-\beta}{2\alpha+k-1}a_{-1,l}, \quad 2 \leq i \leq k.$$

Then

$$D(L_{\pm 1}) = L_{\pm 1} \cdot \left(\sum_{i=0}^{k-1} c_i v_{-\alpha-i} \otimes v_{\alpha+i} \right),$$

where

$$c_{k-1} = \frac{1}{2\alpha + k - 1} a_{-1,l}, \quad c_{i-1} = \frac{i}{\alpha + \beta + i - 1} c_i, \quad i = 1, \dots, k-1.$$

Replacing D by $D - D_{inn}$, where $D_{inn}(L_n) = L_n \cdot \left(\sum_{i=0}^{k-1} c_i v_{-\alpha-i} \otimes v_{\alpha+i} \right)$ for all $n \in \mathbb{Z}$, we have $D(L_{\pm 1}) = 0$. By Lemma 5.1 and $\beta \leq l$, we get $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$.

(1.5.2) If $\alpha + \beta = 1$, we have $(l+2-\beta)a_{-1,l+1} = 0$ by (5.46).

When $-\alpha = l+1$, we have $a_{-1,l+1} \neq 0$. So $l = \beta - 2$, $\beta \neq 1$ and $a_{1,l+1} = \frac{1}{2\beta-2} a_{-1,l}$. Then

$$D(L_1) = L_1 \cdot \left(-\frac{a_{-1,l}}{2\beta-2} v_{\beta-1} \otimes v_{-\beta+1} \right),$$

$$D(L_{-1}) = L_{-1} \cdot \left(-\frac{a_{-1,l}}{2\beta-2} v_{\beta-1} \otimes v_{-\beta+1} \right) + \left(a_{-1,l+1} + \frac{2\beta-1}{2\beta-2} a_{-1,l} \right) v_{\beta-1} \otimes v_{-\beta}.$$

Replace D by $D - D_{inn}$, where $D_{inn}(L_n) = L_n \cdot \left(-\frac{a_{-1,l}}{2\beta-2} v_{\beta-1} \otimes v_{-\beta+1} \right)$ for all $n \in \mathbb{Z}$. Then $D(L_1) = 0$. By Lemma 5.1 and Lemma 5.3 and note $\beta \leq l$, we get $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$.

When $-\alpha = l+k$, where $k \geq 2$, we have $\beta = 1 - \alpha = l+k+1$ and $l+2-\beta = 1-k \leq -1$. So $a_{-1,l+1} = 0$ by (5.46). Then we check that

$$D(L_1) = L_1 \cdot \left(-\sum_{i=l+1}^{l+k} \frac{a_{1,l+1}}{k} v_i \otimes v_{-i} \right).$$

Replace D by $D - D_{inn}$, where $D_{inn}(L_n) = L_n \cdot \left(-\sum_{i=l+1}^{l+k} \frac{a_{1,l+1}}{k} v_i \otimes v_{-i} \right)$ for all $n \in \mathbb{Z}$. Then $D(L_1) = 0$. From Lemma 5.1, Lemma 5.3 and $\beta \leq l$, we get $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$.

Subcase 1.6: $p_{-1} = l+1$ and $q_{-1} = -\alpha \geq l+1$. By (4.2)-(4.3), we have

$$(l+1-\alpha)a_{1,l+1} = 0, \quad (l+\beta)a_{1,l+1} = (l+2-\beta)a_{-1,l+1}. \quad (5.47)$$

Since $a_{1,l+1} \neq 0$, we have $\alpha = l+1$. Note that $-\alpha \geq l+1$. Then $l \leq -1$ and $\alpha \leq 0$. So $\beta \leq l \leq -1$ and $\alpha + \beta \leq -1$. Hence $D(L_1) = a_{1,\alpha} v_\alpha \otimes v_{-\alpha+1}$ and $D(L_{-1}) = \sum_{i=\alpha}^{-\alpha} a_{-1,i} v_i \otimes v_{-i-1}$. By (5.47), we have

$$(\alpha + \beta - 1)a_{1,\alpha} = (\alpha + 1 - \beta)a_{-1,\alpha}. \quad (5.48)$$

Since $\alpha + \beta \leq -1$ and $a_{1,\alpha} \neq 0$, we have $\beta \neq \alpha + 1$ according to (5.48).

(1.6.1): $\alpha = 0$. Since $\beta \leq -1$, we have $a_{-1,0} = -a_{1,0}$ via (5.48). So

$$D(L_1) = a_{1,0} v_0 \otimes v_1, \quad D(L_{-1}) = -a_{1,0} v_0 \otimes v_{-1}.$$

Then $D(L_{\pm 1}) = L_{\pm 1} \cdot \left(-\frac{a_{1,0}}{\beta} v_0 \otimes v_0 \right)$. Replacing D by $D - D_{inn}$, where $D_{inn}(L_n) = L_n \cdot \left(-\frac{a_{1,0}}{\beta} v_0 \otimes v_0 \right)$ for all $n \in \mathbb{Z}$, we have $D(L_{\pm 1}) = 0$. By Lemma 5.1, we get $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$.

(1.6.2): $\alpha < 0$. By (4.4), we have

$$(i + \alpha - 1)a_{-1,i-1} = (i + 1 - \beta)a_{-1,i}, \quad i = \alpha + 1, \dots, -\alpha.$$

Then

$$a_{-1,-\alpha-k} = \left(\prod_{t=1}^k \frac{\alpha + \beta + t - 2}{t} \right) a_{-1,-\alpha}, \quad 1 \leq k \leq -2\alpha.$$

Since $a_{-1,-\alpha} \neq 0$, we have $\alpha + \beta + i \neq 0$ for $-1 \leq i \leq -2\alpha - 2$. In addition, by (5.48), we have $a_{1,\alpha} = \frac{\alpha+1-\beta}{\alpha+\beta-1} a_{-1,\alpha}$. Then we check that $D(L_{\pm 1}) = L_{\pm 1} \cdot \left(\sum_{i=\alpha}^{-\alpha} c_i v_i \otimes v_{-i} \right)$, where

$$c_{-\alpha} = -\frac{a_{-1,-\alpha}}{\alpha - \beta}, \quad c_{-\alpha-k} = \left(\prod_{t=1}^k \frac{\alpha + \beta + t - 1}{t} \right) c_{-\alpha}, \quad 1 \leq k \leq -2\alpha.$$

Replacing D by $D - D_{inn}$, where $D_{inn}(L_n) = L_n \cdot \left(\sum_{i=\alpha}^{-\alpha} c_i v_i \otimes v_{-i} \right)$ for all $n \in \mathbb{Z}$, we have $D(L_{\pm 1}) = 0$. By Lemma 5.1, we get $D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0$.

Case 2: $\alpha = \beta$. By (5.28), we have

$$(i + 1 - \beta)(a_{1,i+1} + a_{-1,i}) = (i - 1 + \beta)(a_{1,i} + a_{-1,i-1}). \quad (5.49)$$

Since $\{a_{-1,i} \neq 0 \mid i \in \mathbb{Z}\}$ is finite, we deduce that $a_{-1,i} = 0$ for $i \leq \beta - 2$ via (5.34).

Subcase 2.1: $\alpha = \beta > 0$. Then $l > 0$ and

$$D(L_1) = a_{1,\beta} v_\beta \otimes v_{-\beta+1} + a_{1,l+1} v_{l+1} \otimes v_{-l} = a_{1,\beta} v_\beta \otimes v_{-\beta+1} + L_1 \cdot \left(\sum_{i=\beta}^l c_i v_i \otimes v_{-i} \right),$$

where

$$c_l = -\frac{a_{1,l+1}}{l + \beta}, \quad (i - \beta)c_i = (i - 1 + \beta)c_{i-1}, \quad i = \beta + 1, \dots, l.$$

Replacing D by $D - D_{inn}$, where $D_{inn}(L_n) = L_n \cdot \left(\sum_{i=\beta}^l c_i v_i \otimes v_{-i} \right)$ for all $n \in \mathbb{Z}$, we have $D(L_1) = a_{1,\beta} v_\beta \otimes v_{-\beta+1}$. By (5.49), we have

$$a_{-1,\beta} = (2\beta - 1)(a_{1,\beta} + a_{-1,\beta-1}), \quad (i + 1 - \beta)a_{-1,i} = (i - 1 + \beta)a_{-1,i-1}, \quad i \neq \beta - 1, \beta.$$

Since $\{a_{-1,i} \neq 0 \mid i \in \mathbb{Z}\}$ is finite, we deduce that $a_{-1,i} = 0$ for $i \leq \beta - 2$ or $i \geq \beta$. Furthermore, $(2\beta - 1)(a_{1,\beta} + a_{-1,\beta-1}) = 0$. So $a_{-1,\beta-1} = -a_{1,\beta}$ and $D(L_{-1}) = -a_{1,\beta} v_{\beta-1} \otimes v_{-\beta}$. By (2.6), we have

$$a_{2,\beta-1} = \frac{\beta}{2\beta - 3} a_{1,\beta}, \quad (5.50)$$

$$(i + 1 - \beta)a_{2,i+1} = (i - 2 + \beta)a_{2,i}, \quad i \neq \beta - 1, \beta, \beta + 1. \quad (5.51)$$

Since $\{a_{2,i} \neq 0 \mid i \in \mathbb{Z}\}$ is finite, we deduce that $a_{2,i} = 0$ for $i \leq \beta - 1$ via (5.51). Using (5.50), we have $a_{1,\beta} = 0$. Therefore, $D(L_{\pm 1}) = 0$. By Lemma 5.1, we have

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \delta_{\alpha,1} \delta_{\beta,1} \mathbb{C} d_{1,1}.$$

Subcase 2.2: $\alpha = \beta = 0$. Then $l \geq 0$ and $D(L_1) = a_{1,0} v_0 \otimes v_1 + a_{1,l+1} v_{l+1} \otimes v_{-l}$.

If $l = 0$, we have $D(L_1) = a_{1,0}v_0 \otimes v_1 + a_{1,1}v_1 \otimes v_0$. If $l > 0$, we have

$$D(L_1) = a_{1,0}v_0 \otimes v_1 + a_{1,l+1}v_{l+1} \otimes v_{-l} = a_{1,0}v_0 \otimes v_1 + a_{1,l+1}v_1 \otimes v_0 + L_1 \cdot \left(\sum_{i=1}^l -\frac{a_{1,l+1}}{i} v_i \otimes v_{-i} \right).$$

Then we assume that $D(L_1) = a_{1,0}v_0 \otimes v_1 + a_{1,1}v_1 \otimes v_0$. Using (5.28), we obtain

$$a_{1,0} + a_{1,1} = -a_{-1,-1} - a_{-1,0}, \quad (5.52)$$

$$a_{-1,-2} = a_{-1,1} = 0, \quad (i+1)a_{-1,i} = (i-1)a_{-1,i-1}, \quad i \neq 0, \pm 1.$$

It is easy to deduce that $a_{-1,i} = 0$ for $i \neq 0, -1$. Hence $D(L_{-1}) = a_{-1,-1}v_{-1} \otimes v_0 + a_{-1,0}v_0 \otimes v_{-1}$. By (2.6), we have

$$2a_{2,0} + a_{2,1} = 3a_{1,0} - a_{-1,0}, \quad a_{2,1} + 2a_{2,2} = 3a_{1,1} - a_{-1,-1}, \quad (5.53)$$

$$(i+1)a_{2,i+1} = (i-2)a_{2,i}, \quad i \neq 0, 1, 2.$$

Then $a_{2,i} = 0$ for $i \neq 0, 1, 2$. So $D(L_2) = a_{2,0}v_0 \otimes v_2 + a_{2,1}v_1 \otimes v_1 + a_{2,2}v_2 \otimes v_0$. By (2.7), we have

$$2a_{-2,0} + a_{-2,-1} = 3a_{-1,0} - a_{1,0}, \quad a_{-2,-1} + 2a_{-2,-2} = 3a_{-1,-1} - a_{1,1}, \quad (5.54)$$

$$(i-1)a_{-2,i-1} = (i+2)a_{-2,i}, \quad i \neq -2, -1, 0.$$

Then $a_{-2,i} = 0$ for $i \neq -2, -1, 0$. So

$$D(L_{-2}) = a_{-2,-2}v_{-2} \otimes v_0 + a_{-2,-1}v_{-1} \otimes v_{-1} + a_{-2,0}v_0 \otimes v_{-2}.$$

Let $m = 2$ and $i = 0, 1$ in (2.5). Then

$$a_{2,2} + a_{2,0} = -(a_{-2,-2} + a_{-2,0}), \quad a_{-2,-1} = -a_{2,1}. \quad (5.55)$$

Using (5.52)-(5.55), we have

$$D = (2a_{1,0} - a_{2,0})d_{0,0}^1 + (a_{2,0} - a_{1,0})d_{0,0}^2 + (2a_{1,1} - a_{2,2})d_{0,0}^3 + (a_{2,2} - a_{1,1})d_{0,0}^4 - a_{2,1}d_{0,0}^5$$

acts on $L_{\pm 1}, L_{\pm 2}$. Hence

$$D = (2a_{1,0} - a_{2,0})d_{0,0}^1 + (a_{2,0} - a_{1,0})d_{0,0}^2 + (2a_{1,1} - a_{2,2})d_{0,0}^3 + (a_{2,2} - a_{1,1})d_{0,0}^4 - a_{2,1}d_{0,0}^5.$$

That is,

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0 \oplus \mathbb{C}d_{0,0}^1 \oplus \mathbb{C}d_{0,0}^2 \oplus \mathbb{C}d_{0,0}^3 \oplus \mathbb{C}d_{0,0}^4 \oplus \mathbb{C}d_{0,0}^5.$$

Subcase 2.3: $\alpha = \beta < 0$.

(2.3.1): $\beta \leq l < -\beta$. If $l = \beta$, we have

$$D(L_1) = a_{1,\beta}v_\beta \otimes v_{-\beta+1} + a_{1,l+1}v_{l+1} \otimes v_{-l} = a_{1,\beta}v_\beta \otimes v_{-\beta+1} + L_1 \cdot \left(-\frac{a_{1,l+1}}{2\beta} v_\beta \otimes v_{-\beta} \right).$$

If $\beta < l < -\beta$, we have

$$D(L_1) = a_{1,\beta}v_\beta \otimes v_{-\beta+1} + a_{1,l+1}v_{l+1} \otimes v_{-l} = a_{1,\beta}v_\beta \otimes v_{-\beta+1} + L_1 \cdot \left(\sum_{i=\beta}^l c_i v_i \otimes v_{-i} \right),$$

where

$$c_l = -\frac{a_{1,l+1}}{l+\beta}, \quad (i-\beta)c_i = (i-1+\beta)c_{i-1}, \quad i = \beta+1, \dots, l.$$

Therefore, for $\beta \leq l < -\beta$, we assume that $D(L_1) = a_{1,\beta}v_\beta \otimes v_{-\beta+1}$.

By (5.28), we have

$$\begin{aligned} a_{-1,\beta-2} &= 0, & a_{-1,\beta} &= (2\beta-1)(a_{1,\beta} + a_{-1,\beta-1}), \\ (i-1+\beta)a_{-1,i-1} &= (i+1-\beta)a_{-1,i}, & i &\neq \beta-1, \beta. \end{aligned} \quad (5.56)$$

Then $a_{-1,i} = 0$ for $i \leq \beta-2$ or $i \geq -\beta+1$ and

$$a_{-1,\beta+i} = \left(\prod_{t=0}^{i-1} \frac{2\beta+t}{t+2} \right) a_{-1,\beta}, \quad 1 \leq i \leq -2\beta. \quad (5.57)$$

By (2.6), we have

$$(i-2+2\beta)a_{-1,i-2} - (i+1-2\beta)a_{-1,i} + 3a_{1,i} = (i+1-\beta)a_{2,i+1} - (i-2+\beta)a_{2,i}. \quad (5.58)$$

Then we obtain

$$\begin{aligned} \beta a_{-1,\beta-1} &= (3-2\beta)a_{2,\beta-1}, & \beta a_{-1,-\beta} &= (3-2\beta)a_{2,-\beta+3}, \\ (i+1-\beta)a_{2,i+1} &= (i-2+\beta)a_{2,i}, & i &\leq \beta-1 \quad \text{or} \quad i \geq -\beta+3. \end{aligned} \quad (5.59)$$

Since $\{a_{2,i} \neq 0 \mid i \in \mathbb{Z}\}$ is finite, we deduce that $a_{2,i} = 0$ for $i \leq \beta-1$ or $i \geq -\beta+3$. Using (5.59), we have

$$a_{-1,\beta-1} = a_{-1,-\beta} = 0.$$

By (5.57), $a_{-1,-\beta} = \left(\prod_{t=0}^{-2\beta-1} \frac{2\beta+t}{t+2} \right) a_{-1,\beta}$. Since $\beta < 0$, we have $\prod_{t=0}^{-2\beta-1} \frac{2\beta+t}{t+2} \neq 0$. Hence $a_{-1,\beta} = 0$ and $a_{-1,i} = 0$ for $\beta+1 \leq i \leq -\beta$ via (5.57). By (5.56), we have $a_{1,\beta} = 0$. Therefore, $D(L_{\pm 1}) = 0$. According to the result of Lemma 5.1, we have

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0.$$

(2.3.2): $l \geq -\beta$.

If $l = -\beta$, we have $D(L_1) = a_{1,\beta}v_\beta \otimes v_{-\beta+1} + a_{1,-\beta+1}v_{-\beta+1} \otimes v_\beta$. If $l > -\beta$, we have

$$D(L_1) = a_{1,\beta}v_\beta \otimes v_{-\beta+1} + (2\beta-1)x_{-\beta+1}v_{-\beta+1} \otimes v_\beta + L_1 \cdot \left(\sum_{i=-\beta+1}^l x_i v_i \otimes v_{-i} \right),$$

where

$$x_l = -\frac{a_{1,l+1}}{l+\beta}, \quad (i-\beta)x_i = (i-1+\beta)x_{i-1}, \quad i = -\beta+2, \dots, l.$$

Therefore, for $l \geq -\beta$, we assume that $D(L_1) = a_{1,\beta}v_\beta \otimes v_{-\beta+1} + a_{1,-\beta+1}v_{-\beta+1} \otimes v_\beta$.

By (5.28), we have

$$(1-2\beta)a_{1,\beta} = (2\beta-1)a_{-1,\beta-1} - a_{-1,\beta}, \quad (2\beta-1)a_{1,-\beta+1} = a_{-1,-\beta-1} - (2\beta-1)a_{-1,-\beta}, \quad (5.60)$$

$$a_{-1,\beta-2} = a_{-1,-\beta+1} = 0, \quad (i-1+\beta)a_{-1,i-1} = (i+1-\beta)a_{-1,i}, \quad i \leq \beta-2 \quad \text{or} \quad i \geq -\beta+2.$$

Then $a_{-1,i} = 0$ for $i \leq \beta-2$ or $i \geq -\beta+1$.

By (2.6), we have

$$\beta a_{-1,\beta-1} = (3 - 2\beta)a_{2,\beta-1}, \quad \beta a_{-1,-\beta} = (3 - 2\beta)a_{2,-\beta+3}, \quad (5.61)$$

$$(i + 1 - \beta)a_{2,i+1} = (i - 2 + \beta)a_{2,i}, \quad i \leq \beta - 2 \quad \text{or} \quad i \geq -\beta + 3.$$

Since $\{a_{2,i} \neq 0 \mid i \in \mathbb{Z}\}$ is finite, we get $a_{2,i} = 0$ for $i \leq \beta - 1$ or $i \geq -\beta + 3$. Using (5.61), we have $a_{-1,\beta-1} = a_{-1,-\beta} = 0$. Let $i = \beta - 1$ in (5.28). Then $a_{1,\beta} = 0$. Furthermore, using (5.28), we have

$$(2\beta - 1)a_{1,-\beta+1} = a_{-1,-\beta-1}, \quad (5.62)$$

$$(i - 1 + \beta)a_{-1,i-1} = (i + 1 - \beta)a_{-1,i}, \quad i \neq -\beta, -\beta + 1. \quad (5.63)$$

Let $i = \beta, \beta + 1, \dots, -\beta - 1$ in (5.63) and use $a_{-1,\beta-1} = 0$, then we deduce that $a_{-1,i} = 0$ for $\beta \leq i \leq -\beta - 1$. Hence $a_{1,-\beta+1} = 0$. Therefore, $D(L_{\pm 1}) = 0$. According to Lemma 5.1, we have

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta)_0.$$

□

5.4. By Lemma 2.9, we assume $D(L_0) = v_0 \otimes v_0$ in this case.

Lemma 5.5. *For $(\alpha, \beta) = (0, 0)$, if $D(L_0) = v_0 \otimes v_0$, we have*

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_0 \otimes \mathcal{F}_0)_0 \oplus \mathbb{C}d_{0,0}^1 \oplus \mathbb{C}d_{0,0}^2 \oplus \mathbb{C}d_{0,0}^3 \oplus \mathbb{C}d_{0,0}^4 \oplus \mathbb{C}d_{0,0}^5.$$

Proof. For $(\alpha, \beta) = (0, 0)$, if $D(L_0) = v_0 \otimes v_0$, replacing D by $D - d_{0,0}^1$, we have $D(L_0) = 0$. Using the results of Lemma 5.1-Lemma 5.4, we have

$$D \in \text{Inn}(\mathcal{W}, \mathcal{F}_0 \otimes \mathcal{F}_0)_0 \oplus \mathbb{C}d_{0,0}^1 \oplus \mathbb{C}d_{0,0}^2 \oplus \mathbb{C}d_{0,0}^3 \oplus \mathbb{C}d_{0,0}^4 \oplus \mathbb{C}d_{0,0}^5.$$

□

By Lemma 5.1-Lemma 5.5 our main result in this section can be stated as follows.

Proposition 5.6. *For $(\alpha, \beta) \in \mathbb{Z}^2$, we have*

$$H^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta) = \begin{cases} \bigoplus_{i=1}^5 \mathbb{C}d_{0,0}^i, & (\alpha, \beta) = (0, 0); \\ \mathbb{C}d_{0,1} \oplus \mathbb{C}d_{0,1}^1, & (\alpha, \beta) = (0, 1); \\ \mathbb{C}d_{1,0} \oplus \mathbb{C}d_{1,0}^1, & (\alpha, \beta) = (1, 0); \\ \mathbb{C}d_{0,2}, & (\alpha, \beta) = (0, 2); \\ \mathbb{C}d_{2,0}, & (\alpha, \beta) = (2, 0); \\ \mathbb{C}d_{1,1}, & (\alpha, \beta) = (1, 1); \\ \mathbb{C}d_{\alpha,\beta}, & \alpha + \beta = 1, \beta \neq 0, 1; \\ 0, & \text{otherwise,} \end{cases}$$

where $d_{0,0}^i (i = 1, 2, \dots, 5)$, $d_{1,0}^1$, $d_{0,1}^1$, $d_{\alpha,\beta}$ are defined in Proposition 3.1, Proposition 3.2, and Proposition 3.3.

By utilizing Proposition 4.7 and Proposition 5.6, we present the main theorem of this paper.

Theorem 5.7. *For $\alpha, \beta \in \mathbb{C}$, we have*

$$\dim H^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\beta) = \begin{cases} 5, & (\alpha, \beta) = (0, 0); \\ 2, & (\alpha, \beta) = (0, 1) \text{ or } (1, 0); \\ 1, & (\alpha, \beta) = (0, 2), (2, 0), (1, 1), \text{ or } \alpha + \beta = 1, \beta \neq 0, 1; \\ 0, & \text{otherwise.} \end{cases}$$

6. APPLICATIONS

In this section, we discuss some applications of Theorem 5.7.

In [3], Drinfeld introduced the notion of Lie bialgebras as a framework for investigating the Yang-Baxter equations. A Lie bialgebra, denoted as (L, δ) , consists of a Lie algebra L equipped with a Lie cobracket $\delta : L \rightarrow L \wedge L$ that satisfies the co-Jacobi identity, as described in [22, 23]. The mapping δ operates as a 1-cocycle on the Lie algebra L , with coefficients in the L -module $L \wedge L$. This action of L on $L \wedge L$ is carried out through the adjoint representation. When δ is a 1-coboundary with coefficients in $\wedge^2 L$, the Lie bialgebra (L, δ) is termed a coboundary Lie bialgebra.

Theorem 6.1 ([24, 28]). *Let \mathcal{W} be the Witt algebra. Then $H^1(\mathcal{W}, \mathcal{W} \wedge \mathcal{W}) = 0$. In particular, all Lie bialgebra structures on \mathcal{W} are coboundary.*

Proof. It follows from $\mathcal{W} \cong \mathcal{F}_{-1}$ as \mathcal{W} -modules and Theorem 5.7. \square

For $\alpha \in \mathbb{C}$, the **Ovsienko-Roger algebra** $\mathcal{W}(\alpha) = \mathcal{W} \ltimes \mathcal{F}_\alpha$ [25] (see also [13]) is the semi-direct product of \mathcal{W} and \mathcal{F}_α with the following brackets:

$$[L_m, L_n] = (m - n)L_{m+n}, \quad [L_m, v_n] = -(\alpha m + n)v_{m+n}, \quad [v_m, v_n] = 0, \quad \forall m, n \in \mathbb{Z}.$$

Proposition 6.2.

$$\dim H^1(\mathcal{W}(\alpha), \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha)) = \begin{cases} 7, & \alpha = 0; \\ 1, & \alpha = 1; \\ 1, & \alpha = \frac{1}{2}; \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Since \mathcal{F}_α is an abelian ideal of $\mathcal{W}(\alpha)$, using the Hochschild-Serre spectral sequence [30], we have the following exact sequence:

$$0 \rightarrow H^1(\mathcal{W}, (\mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha))^{\mathcal{F}_\alpha}) \rightarrow H^1(\mathcal{W}(\alpha), \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha)) \rightarrow H^1(\mathcal{F}_\alpha, \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha))^{\mathcal{W}(\alpha)}.$$

Note that the action of \mathcal{W} on $H^1(\mathcal{F}_\alpha, \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha))$ is induced by the action of \mathcal{W} on $\text{Der}(\mathcal{F}_\alpha, \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha))$ defined by

$$(x \cdot d)(z) = x \cdot d(z) - d[x, z], \quad \forall x \in \mathcal{W}(\alpha), z \in \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha).$$

Since

$$\begin{aligned} (\mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha))^{\mathcal{F}_\alpha} &= \{v \in \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha) \mid x \cdot v = 0, \forall x \in \mathcal{F}_\alpha\} = \mathcal{F}_\alpha \otimes \mathcal{F}_\alpha, \\ \mathbf{H}^1(\mathcal{F}_\alpha, \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha))^{\mathcal{W}} &= \text{Hom}_{\mathcal{W}}(\mathcal{F}_\alpha, \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha)), \end{aligned}$$

it suffices to compute $\mathbf{H}^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\alpha)$ and $\text{Hom}_{\mathcal{W}}(\mathcal{F}_\alpha, \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha))$. From Theorem 5.7, we obtain

$$\dim \mathbf{H}^1(\mathcal{W}, \mathcal{F}_\alpha \otimes \mathcal{F}_\alpha) = \begin{cases} 5, & \alpha = 0; \\ 1, & \alpha = 1; \\ 1, & \alpha = \frac{1}{2}; \\ 0, & \text{otherwise.} \end{cases}$$

By direct computation, we have

$$\text{Hom}_{\mathcal{W}}(\mathcal{F}_\alpha, \mathcal{W}(\alpha) \otimes \mathcal{W}(\alpha)) = \begin{cases} \mathbb{C}\tilde{d}_0^6 \oplus \mathbb{C}\tilde{d}_0^7, & \alpha = 0; \\ 0, & \text{otherwise,} \end{cases}$$

where

$$\tilde{d}_0^6(v_m) = v_0 \otimes v_m, \quad \tilde{d}_0^7(v_m) = v_m \otimes v_0, \quad \forall m \in \mathbb{Z}.$$

It is easy to see \tilde{d}_0^6 and \tilde{d}_0^7 lies in the image of the restriction map. This completes the proof. \square

Remark 6.3. The algebra $\mathcal{W}(0)$ is isomorphic to the twisted Heisenberg-Virasoro algebra when $\alpha = 0$. In [19], $\mathbf{H}^1(\mathcal{W}(-1), \mathcal{W}(-1) \otimes \mathcal{W}(-1))$ was calculated; however, it is regrettable that the 1-cocycle $d_{0,0}^5$ was omitted. When $\alpha = -1$, the algebra $\mathcal{W}(-1)$ is isomorphic to the BMS algebra. The calculation of $\mathbf{H}^1(\mathcal{W}(0), \mathcal{W}(0) \otimes \mathcal{W}(0))$ was successfully accomplished in [18].

Corollary 6.4.

$$\mathbf{H}^1(\mathcal{W}(\alpha), \mathcal{W}(\alpha) \wedge \mathcal{W}(\alpha)) = \begin{cases} \bigoplus_{i=1}^3 \mathbb{C}D_0^i, & \alpha = 0; \\ 0, & \text{otherwise,} \end{cases}$$

where

$$\begin{aligned} D_0^1(L_m) &= v_m \otimes v_0 - v_0 \otimes v_m, \quad D_0^2(L_m) = m(v_m \otimes v_0 - v_0 \otimes v_m), \quad D_0^3(v_m) = v_m \otimes v_0 - v_0 \otimes v_m, \\ D_0^1(v_m) &= D_0^2(v_m) = D_0^3(L_m) = 0 \end{aligned}$$

for $m \in \mathbb{Z}$. In particular, all Lie bialgebra structures on $\mathcal{W}(\alpha)$ are coboundary if and only if $\alpha \neq 0$.

Proof. By Proposition 6.2, we only need to consider $\alpha = 0, 1, \frac{1}{2}$. By utilizing the explicit non-trivial 1-cocycles in Proposition 3.1, Proposition 3.2, and Proposition 3.3, one can easily determine all non-trivial 1-cocycles with coefficients in $\mathcal{W}(\alpha) \wedge \mathcal{W}(\alpha)$ through the skew symmetry property. \square

The **twisted Schrödinger-Virasoro algebra** \mathcal{S} introduced in [26], is a Lie algebra with \mathbb{C} -basis $\{L_n, M_n, Y_n, | n \in \mathbb{Z}\}$ and the following Lie brackets,

$$\begin{aligned} [L_m, L_n] &= (m-n)L_{m+n}, & [L_m, Y_n] &= \left(\frac{m}{2} - n\right)Y_{m+n}, & [L_m, M_n] &= -nM_{n+m}, \\ [Y_m, Y_n] &= (m-n)M_{m+n}, & [Y_m, M_n] &= [M_m, M_n] = 0 \end{aligned}$$

for $m, n \in \mathbb{Z}$. Let $\mathcal{J} = \text{span}_{\mathbb{C}}\{M_n, Y_n | n \in \mathbb{Z}\}$. It is clear that \mathcal{J} is an ideal of \mathcal{S} and $\mathcal{J} \cong \mathcal{F}_0 \oplus \mathcal{F}_{-\frac{1}{2}}$ as \mathcal{W} -modules.

Proposition 6.5. $\dim H^1(\mathcal{S}, \mathcal{S} \otimes \mathcal{S}) = 7$.

Proof. For \mathcal{J} is a ideal of \mathcal{S} , we have the following exact sequence:

$$0 \rightarrow H^1(\mathcal{W}, (\mathcal{S} \otimes \mathcal{S})^{\mathcal{J}}) \rightarrow H^1(\mathcal{S}, \mathcal{S} \otimes \mathcal{S}) \rightarrow H^1(\mathcal{J}, \mathcal{S} \otimes \mathcal{S})^{\mathcal{W}}.$$

Since $(\mathcal{S} \otimes \mathcal{S})^{\mathcal{J}} = \{v \in \mathcal{S} \otimes \mathcal{S} | x \cdot v = 0, \forall x \in \mathcal{J}\} = \mathcal{F}_0 \otimes \mathcal{F}_0$. From Theorem 5.7 it follows that the dimension of the first space in the exact sequence is 5. By direct computation, we have $H^1(\mathcal{J}, \mathcal{S} \otimes \mathcal{S})^{\mathcal{W}} = \mathbb{C}\tilde{d}_6 \oplus \mathbb{C}\tilde{d}_7$, where

$$\tilde{d}_6(M_m) = M_0 \otimes M_m, \quad \tilde{d}_6(Y_m) = M_0 \otimes Y_m, \quad \tilde{d}_7(M_m) = M_m \otimes M_0, \quad \tilde{d}_7(Y_m) = Y_m \otimes M_0.$$

It is easy to see that \tilde{d}_6 and \tilde{d}_7 lie in the image of the restriction map. This completes the proof. \square

Corollary 6.6. $H^1(\mathcal{S}, \mathcal{S} \wedge \mathcal{S}) = \bigoplus_{i=1}^3 \mathbb{C}D_i$, where

$$\begin{aligned} D_1(L_m) &= M_0 \otimes M_m - M_m \otimes M_0, & D_2(L_m) &= mM_0 \otimes M_m - mM_m \otimes M_0, \\ D_3(M_m) &= M_0 \otimes M_m - M_m \otimes M_0, & D_3(Y_m) &= M_0 \otimes Y_m - Y_m \otimes M_0, \\ D_1(M_m) &= D_1(Y_m) = D_2(M_m) = D_2(Y_m) = D_3(L_m) = 0 \end{aligned}$$

for $m \in \mathbb{Z}$.

ACKNOWLEDGEMENTS

We would like to thank Professor Claude Roger for his valuable comments and Xia Xie for her suggestions. We also greatly appreciate the input from the anonymous reviewers. This research was supported by the National Natural Science Foundation of China (12071405).

REFERENCES

- [1] E. Arbarello, C. De Concini, V.G. Kac, C. Procesi, Moduli spaces of curves and representation theory, *Commun. Math. Phys.* **117** (1988), 1–36. [2](#)
- [2] A. Bagchi, R. Gopakumar, I. Mandal, A. Miwa, GCA in 2d, *J. High Energy Phys.* **2010** (2010). [2](#)
- [3] V.G. Drinfeld, Hamiltonian structures on Lie groups, Lie bialgebras and the geometric meaning of classical Yang–Baxter equations, *Dokl. Akad. Nauk SSSR* **268** (1983), 285–287. [35](#)
- [4] K. Dykes, Length two extensions of modules for the Witt algebra, *Commun. Algebra* **46** (2018), 2146–2170.

- [5] J. Ecker, The low-dimensional algebraic cohomology of the Witt and the Virasoro algebra with values in general tensor densities modules, in: *Lie Theory Appl. Phys.*, Springer Proc. Math. Stat., vol. 335, Springer, 2020, pp. 535–540. [1](#)
- [6] J. Ecker, M. Schlichenmaier, The low-dimensional algebraic cohomology of the Witt and the Virasoro algebra, *J. Phys. Conf. Ser.* **1194** (2019). [1](#)
- [7] J.M.A. Ecker, M. Schlichenmaier, The low-dimensional algebraic cohomology of the Witt and Virasoro algebra with values in natural modules, *Banach Cent. Publ.* **123** (2021), 141–174. [1](#)
- [8] H. Fa, Y. Li, J. Li, The Schrödinger-Virasoro type Lie bialgebra: a twisted case, *Front. Math. China* **6** (2011), 641–657. [1](#)
- [9] R. Farnsteiner, Derivations and extensions of finitely generated graded Lie algebras, *J. Algebra* **118** (1988), 34–45. [3](#)
- [10] B. Feigin, D. Fuchs, Homology of the Lie algebra of vector fields on the line, *Funct. Anal. Appl.* **14** (1980), 201–212. [1](#), [2](#)
- [11] A. Fialowski, Formal rigidity of the Witt and Virasoro algebra, *J. Math. Phys.* **53** (2012), 073501. [1](#)
- [12] D. Fuchs, *Cohomology of infinite-dimensional Lie algebras*, Contemp. Soviet Math., Consultants Bureau, New York, 1986. [3](#)
- [13] S. Gao, P. Jiang, Y. Pei, Low dimensional cohomology groups of the Lie algebras $W(a, b)$, *Commun. Algebra* **39** (2011), 397–423. [35](#)
- [14] L. Guieu, C. Roger, *L'Algèbre et le Groupe de Virasoro: Aspects Géométriques et Algébriques, Généralisations*, Publications CRM, 2007. [1](#)
- [15] J. Han, J. Li, Y. Su, Lie bialgebras of the Schrödinger-Virasoro algebra, *J. Math. Phys.* **50** (2009), 083504. [1](#)
- [16] K. Iohara, Y. Koga, *Representation theory of the Virasoro algebra*, Springer, 2011. [1](#)
- [17] B. Kupershmidt, O. Stoyanov, Classification of all Poisson-Lie structures on an infinite-dimensional jet group, *Lett. Math. Phys.* **37** (1996), 1–9. [1](#)
- [18] J. Li, Y. Su, B. Xin, Lie bialgebra structures on the W -algebra $W(2, 2)$, *Algebra Colloq.* **17** (2010), 181–190. [1](#), [36](#)
- [19] D. Liu, Y. Pei, L. Zhu, Lie bialgebra structures on the twisted Heisenberg-Virasoro algebra, *J. Algebra* **359** (2012), 35–48. [1](#), [36](#)
- [20] C. Martin, A. Piard, Classification of the indecomposable bounded admissible modules over the Virasoro Lie algebra with weight spaces of dimension not exceeding two, *Commun. Math. Phys.* **150** (1992), 465–493. [1](#)
- [21] A. Meurman, L. Santharoubane, Cohomology and Harish-Chandra modules over the Virasoro algebra, *Commun. Algebra* **16** (1988), 27–35. [1](#)
- [22] W. Michaelis, Lie coalgebras, *Adv. Math.* **38** (1980), 1–54. [35](#)
- [23] W. Michaelis, A class of infinite-dimensional Lie bialgebras containing the Virasoro algebra, *Adv. Math.* **107** (1994), 365–392. [35](#)
- [24] S. Ng, E. Taft, Classification of the Lie bialgebra structures on the Witt and Virasoro algebras, *J. Pure Appl. Algebra* **151** (2000), 67–88. [1](#), [2](#), [35](#)
- [25] V. Ovsienko, C. Roger, Generalizations of the Virasoro group and Virasoro algebra through extensions by modules of tensor densities on S^1 , *Indag. Math.* **9** (1998), 277–288. [1](#), [2](#), [35](#)
- [26] C. Roger, J. Unterberger, The Schrödinger-Virasoro Lie group and algebra: Representation theory and cohomological study, *Ann. Henri Poincaré* **7** (2006), 1477–1529. [2](#), [37](#)
- [27] G. Song, Y. Su, Lie bialgebras of generalized Witt type, *Sci. China Math.* **49** (2006), 533–544. [1](#)
- [28] E. Taft, Witt and Virasoro algebras as Lie bialgebras, *J. Pure Appl. Algebra* **87** (1993), 301–312. [35](#)
- [29] M. Schlichenmaier, An elementary proof of the vanishing of the second cohomology of the Witt and Virasoro algebra with values in the adjoint module, *Forum Math.* **26** (2014), 913–929. [1](#)

- [30] C. Weibel, *An introduction to homological algebra*, Cambridge University Press, 1994. 3, 35
- [31] E. Witten, Coadjoint orbits of the Virasoro group, *Commun. Math. Phys.* **114** (1988), 1–53. 1
- [32] W. Zhang, C. Dong, W -algebra $W(2, 2)$ and the vertex operator algebra $L(\frac{1}{2}, 0) \otimes L(\frac{1}{2}, 0)$, *Commun. Math. Phys.* **285** (2009), 991–1004 2

DEPARTMENT OF MATHEMATICS, HUZHOU UNIVERSITY, ZHEJIANG HUZHOU 313000, PR CHINA
Email address: gaoshoulan@zjhu.edu.cn

DEPARTMENT OF MATHEMATICS, HUZHOU UNIVERSITY, ZHEJIANG HUZHOU 313000, PR CHINA
Email address: liudong@zjhu.edu.cn

DEPARTMENT OF MATHEMATICS, HUZHOU UNIVERSITY, ZHEJIANG HUZHOU 313000, PR CHINA
Email address: pei@zjhu.edu.cn