

Constraining neutron star matter from dM/dR

Márcio Ferreira^{1,*} and Constança Providência^{1,†}

¹*CFisUC, Department of Physics, University of Coimbra, P-3004 - 516 Coimbra, Portugal*

(Dated: June 19, 2024)

We analyse the implications of information about local derivatives from the mass-radius diagram in neutron star matter. It is expected that the next generation of gravitational wave and electromagnetic detectors will allow the determination of the neutron star radius and mass with a small uncertainty. Observations of neutron stars clustered around a given neutron star mass allow the estimation of local derivatives in the $M(R)$ diagram, which can be used to constrain neutron star properties. From a model-independent description of the neutron star equation of state, it is shown that a $M(R)$ curve with a negative slope at $1.4M_\odot$ predicts a $2M_\odot$ neutron star radius below 12 km. Furthermore, a maximum mass below $2.3M_\odot$ is obtained if the $M(R)$ slope is negative in the whole range of masses above $1M_\odot$, and a maximum mass above $2.4M_\odot$ requires the $M(R)$ slope to be positive in some range of masses. Constraints on the mass-radius curve of neutron stars will place strong constraints on microscopic models.

I. INTRODUCTION

Many studies have been carried out to determine the high-density equation of state (EoS) from the knowledge of the mass-radius curve of the neutron star (NS) [1–6]. The determination of the mass and radius of an NS with sufficient precision will allow the high-density baryon EoS to be constrained. The Neutron Star Interior Composition Explorer (NICER) is expected to measure the mass and radius of an NS with an uncertainty of 5% [7]. Future detectors such as the Enhanced X-ray Timing and Polarimetry mission (eXTP) [8, 9], the (STROBE-X) [10] will improve this precision to $\sim 2\%$. Constraints on NS mass and radius are also expected from the detection of gravitational waves emitted by binary NS mergers, such as GW170817 [11] reported by the LIGO/Virgo collaboration. The third generation of gravitational wave detectors is expected to allow the determination of the NS radius with an uncertainty as small as 100 m [12]. The Square Kilometer Array [13] telescope will also play an important role in constraining the possible scenarios for the EoS of matter within the NS.

Several microscopic phenomenological models have been used to determine the mass-radius curve that links the astrophysical observations of the NS to the EoS of baryon matter. One of the features that characterise these curves is the slope of the $M(R)$ curve with respect to radius. For masses above $1M_\odot$, the slope is always negative for some models such as SLy4 and SLy9 [14], SFHo [15] and FSU or FSU2 [16], see [17–19], other models such as DD2 [20], DDME2 [21], NL3 $\omega\rho$ [22, 23], FSU2H [24] or BigApple [25] show a backbending and part of the curve above $1M_\odot$ has a positive slope, see [18], or models like NL3 [26] or IUFSU [27] have an almost infinite slope for

much of the $M(R)$ curve. For the models with a positive or infinite slope, the curve will pass through a negative slope for sufficiently large masses before the maximum mass star is reached. The transition to a negative slope can occur due to the onset of new degrees of freedom such as hyperons or quark deconfinement, if included in the NS composition, see Fig. 4 of [18]. The behavior of the $M(R)$ curve reflects the properties of the hadronic interaction, with the back-bending being generally associated with a stiffening of the EoS.

Recently, the authors of [28] selected six EoS that maximize the variance of four NS properties from their dataset of model-independent EoS constructed using Gaussian processes. Of the six EoS, five predict $M(R)$ curves with a positive derivative between $1.0M_\odot$ and $M \gtrsim 1.7M_\odot$, and the sixth has a negative slope for all stars with masses above $1.0M_\odot$, a behaviour similar to that of SLy4 and SFHo. Since the behaviour of the $M(R)$ curve is inextricably linked to the density dependence of the EoS, it makes sense to ask whether the constraints imposed to build the data sets require the $M(R)$ curves to have a positive slope in some range of NS masses, in which mass range, and what is the reason for this behaviour. Knowledge of this behaviour from observations would place strong constraints on the acceptable microscopic models.

In the present work, using a model-independent, agnostic description of NS matter, we aim to answer the question: is it possible to identify NS properties that distinguish the two sets of models that do or do not have a negative slope in the whole mass range above $1M_\odot$? Note that if sufficient precision is achieved in the determination of the NS radius, back-bending will be easily confirmed if a low-mass star has a smaller radius than a medium to high-mass star. Knowing the behaviour of the slope of $M(R)$ would place strong constraints on the microscopic model and would provide information on the high density behaviour of the QCD EoS.

The paper is organized as follows. The parametrization

* marcio.ferreira@uc.pt

† cp@uc.pt

we use to describe neutron star matter is presented in Sec. II. The results are shown and discussed in Sec. III and some conclusions are drawn in Sec. IV.

II. DATASET

To describe the equation of state (EoS) of neutron star matter, we use the piecewise polytropic parameterization $p(\rho) = K\rho^\Gamma$, where $\rho = mn$ is the rest mass density, n is the baryon number density, m is the mass of a baryon, K is the polytropic pressure coefficient, and Γ is the adiabatic index [29, 30]. The present parameterization uses five connected polytropic segments. The first segment is defined within the density range $[n_{\text{crust}}, 1.1n_0]$, where $n_0 = 0.16 \text{ fm}^{-3}$ denotes the nuclear saturation density and $n_{\text{crust}} \equiv n_0/2$, with the polytropic index Γ_0 randomly chosen from the range $1.0 < \Gamma_0 < 4.5$. This Γ_0 interval ensures that the polytrope lies within the band described by the chiral effective field theory [30]. We assume the SLy4 EoS [31] for densities $n < n_{\text{crust}}$. To obtain a flexible and robust representation of neutron star EoS, the remaining four polytropic segments start at random densities, such that $n_1 < n_2 < n_3 < n_4$, with randomly chosen polytropic indices $\{\Gamma_1, \Gamma_2, \Gamma_3, \Gamma_4\}$. The parameter space $\{\Gamma_0, \Gamma_1, n_1, \Gamma_2, n_2, \Gamma_3, n_3, \Gamma_4, n_4, \Gamma_5, n_5\}$ was covered by randomly sampling from uniform distributions consistent with $\{n_1, n_2, n_3, n_4\}$ being within n_0 and $8n_0$, $1.0 < \Gamma_0 < 4.5$ and $0.05 < \Gamma_i < 8$ for $i = 1, \dots, 4$.

To study NS properties, we solve the Tolmann-Oppenheimer-Volkoff (TOV) equations [32, 33], which describe spherically symmetric stars in hydrostatic equilibrium. Furthermore, the differential equations that determine the tidal deformability of the stars have also been solved [34]. A valid EoS must be consistent with the observation of a $M > 2M_\odot$ NS and have a speed of sound that remains less than the speed of light. We have generated a data set containing 40435 valid EoS.

Depending on the value of the slope dM/dR calculated along the TOV $M(R)$ sequence between $1M_\odot$ and M_{max} , we divide the dataset into two subsets: a) 3106 EoSs whose $M(R)$ sequences satisfy $dM/dR < 0$ in the whole mass range above $1M_\odot$ and b) 37329 EoSs that do not fulfil $dM/dR < 0$.

The following set of astrophysical constraints will be applied (see [35]): i) $R(2.0M_\odot) > 10.75$ km from NICER measurements of PSR J0740+6620 [36, 37]; ii) $R(1.1M_\odot) > 10.8$ km from NICER measurements of J0030+0451 [38, 39]; and iii) the effective tidal deformability of $\tilde{\Lambda} < 720$ (low spin-prior), where $\tilde{\Lambda} = (16/13)((12q+1)\Lambda_1 + (12+q)q^4\Lambda_2)/(1+q)^5$, estimated from the GW170817 event [40], with a binary mass ratio of $0.73 \leq q = M_2/M_1 \leq 1$ and a chirp mass of $1.186M_\odot$, where $M_{\text{chirp}} = (M_1M_2)^{3/5}/(M_1+M_2)^{1/5}$. For a review of theoretical, experimental and observational constraints for the equation of state of dense matter see the recent review [41]. After applying the above constraints we ended up with a total of 21614 EoS: 2444 with $dM/dR < 0$ and

19170 with $dM/dR \not< 0$.

III. RESULTS

The $M(R)$ and $M(\Lambda)$ sequences are shown in Fig. 1 for both subsets with (dark colors) and without (light colors) astrophysical constraints. Some comments are in order: i) EoS which satisfy $dM/dR < 0$, predict smaller maximum masses, and smaller upper bounds for the NS radii. Medium mass stars with a radius above 13 km or a maximum mass above $2.3M_\odot$ would exclude this set of EoS. Astrophysical constraints filter $M(R)$ curves with a small radius; ii) the set of EoS that show a portion with positive slope, allows for stars as massive as $3M_\odot$, and a radii above 14 km, if the astrophysical conditions are not imposed. These values are reduced, respectively, to maximum masses $\sim 2.6M_\odot$ and radii below ~ 13.5 km. Astrophysical constraints also cut the low radii $M(R)$ curves.




FIG. 1. Sequences $M(R)$ (left) and $M(\Lambda)$ (right) for the set of EoS satisfying $dM/dR < 0$ (top) and $dM/dR \not< 0$ (bottom) with (dark) and without (light) astrophysical constraints (see text for details). The boundaries of the different regions specify the extremes (minimum/maximum).

A. Neutron star equation of state

The constraint that a negative slope along the entire $M(R)$ sequence (no backbending) has on the pressure of neutron star matter is shown in Fig. 2, where we compare the $p(n)$ curves for both sets. The dark colours represent the 90% confidence intervals (CI), while the light colours

define the full data set, i.e. the extremes. The pressure has a much smoother behaviour for the $dM/dR < 0$ set, being constrained to much lower values (softening) at $n \approx 0.4 - 0.6 \text{ fm}^{-3}$: the 90% CI of the $dM/dR < 0$ is below the 90% CI of the red set. However, note that at high density this set is on average stiffer than the set with backbending. This is due to the condition that the maximum mass must be at least $2M_{\odot}$. The kink seen in the statistics of the red set at about $n = 0.55 \text{ fm}^{-3}$, which has been interpreted as a possible indication of a phase transition, is not present in the blue set. The imposition of the astrophysical constraints reduces the range covered, but the main features described above are still present.




FIG. 2. Pressure as a function of baryon density for the set $dM/dR < 0$ (blue) and $dM/dR \not< 0$ (red) with (bottom) and without (top) astrophysical constraints. The darker bands represent 90% credible intervals (CI) while the lighter bands show the extremes (minimum/maximum).

Figure 3 shows the speed-of-sound squared for both sets. We also show the 65% vertical bands representing the central densities of M_{max} for both sets. The blue set ($dM/dR < 0$, no backbending on the $M(R)$ diagram) has a $v_s^2(n)$ that increases almost monotonically up to $\sim 0.8 \text{ fm}^{-3}$, the median (solid lines) has a bend at this density, before decreasing up to the density n_{max} , but does not reach the conformal limit $v_s^2 = 1/3$. On the other hand, the red set has a median value close to $v_s^2 = 1/3$ at the central densities. The astrophysical constraints originate the peak around $3n_0$ as identified in other works [35, 42]. These astrophysical conditions remove from the dataset

the hardest EoS at the lowest densities. There is a clear distinction that can be made from the assessment of possible backbending in the $M(R)$ plot. The $dM/dR < 0$ set has larger central densities because, as the EoS is softer, $v_s^2(dM/dR < 0)$ remains below $v_s^2(dM/dR \not< 0)$ for $n \lesssim 3n_0$, matter is compressed more efficiently by gravity. Note also that the position of the central densities in the case of $dM/dR \not< 0$ ($dM/dR < 0$) shifts to larger (smaller) densities because some of the hardest (softest) EoS have been removed from the set.




FIG. 3. Speed-of-sound squared as a function of baryon density for the set $dM/dR < 0$ (blue) and $dM/dR \not< 0$ (red) with (bottom) and without (top) astrophysical constraints. The lighter bands represent 90% CI, the darker bands 50% CI, and the solid lines the median. The vertical bands specify the 65% CI for the central densities (n_{max}) at the M_{max} .

In [43] the authors consider the renormalized trace anomaly $\Delta = 1/3 - p/e$, where p is the pressure and e is the energy density, to study conformality restoration within NS. Since perturbative QCD predicts a small positive trace in the very high density limit, it was conjectured that the renormalized trace anomaly should be positive in the whole density range from low to high density. The anomaly of the normalized matter trace is shown in Fig. 4 for both sets. For the $dM/dR < 0$ set, the 50% CI crosses the zero axis and remains below zero. However, the other set of EoS has a positive Δ above the median throughout the density range. The conjecture proposed in [43] favours a $M(R)$ curve with backbending: above the median line, all EoS with backbending have a positive renormalised trace below 1.2 fm^{-3} .




FIG. 4. The normalized matter trace anomaly as a function of baryon density for the set $dM/dR < 0$ (blue) and $dM/dR \not< 0$ (red) with (bottom) and without (top) astrophysical constraints. The lighter bands represent 90% CI, the darker bands 50% CI, and the solid lines the median. The vertical bands specify the 65% CI for the central densities (n_{\max}) at the M_{\max} .

A clearer analysis is obtained from the probability distribution functions for both Δ and v_s^2 at the central densities of M_{\max} shown in Fig. 5. The speed-of-sound squared probability distributions of the two sets at the centre of the maximum mass star are quite different: the $dM/dR < 0$ set has a rather flat profile which is cut off by causality at $v_s^2/c^2 = 1$; the $dM/dR \not< 0$ set, on the other hand, presents a distribution with a well-defined peak around $v_s^2/c^2 = 0.1$, which decreases monotonically to $v_s^2/c^2 = 1$, and the causality condition has a very small effect. The renormalized trace anomaly probability distribution functions calculated at the centre of the maximum mass star also show different behaviour: for the $dM/dR \not< 0$ set, most of the EoS have a positive value, while the $dM/dR < 0$ set shows positive values for a very reduced number of EoS. As expected from Fig 4, the 50% CI for the $dM/dR < 0$ set is always positive, see the caption of Fig. 5.

To complete this study, we have identified all the EoS from both sets that have a positive trace anomaly up to 1.2 fm^{-3} and we have plotted them on top of the $M(R)$ plot, see Fig. 6. As discussed in [43], the maximum masses reached by these EoS are smaller and have a larger radius for a given mass, regardless of the behaviour of the




FIG. 5. The probability distribution functions for the normalized matter trace anomaly (top) and speed-of-sound squared (bottom) at the central densities of M_{\max} . We display the sets $dM/dR < 0$ (blue) and $dM/dR \not< 0$ (red) with (dark) and without (light) astrophysical constraints. The 50% CIs for $\Delta(n_{\max})$ are $-0.070_{+0.051}^{-0.051}$ ($dM/dR < 0$, without restrictions), $-0.057_{+0.050}^{-0.053}$ ($dM/dR < 0$, with restrictions), $0.032_{+0.073}^{-0.081}$ ($dM/dR \not< 0$, without restrictions), and $0.008_{+0.066}^{-0.073}$ ($dM/dR \not< 0$, with restrictions).

slope. The difference in radius can be of the order of 1 km (or even larger for the negative slope set).




FIG. 6. Same plots as in Fig. 1 but showing explicitly the $M(R)$ sequences for the EOS with positive $\Delta(n)$ for $0 < n < 1.2 \text{ fm}^{-3}$.

Tables I and II show several properties of NS with different masses. In Table I the median and 90% CI limits

are given for the analysis with and without additional astrophysical constraints. In Table II some extreme properties are also given.

B. Constraining R and Λ from dM/dR

Assuming that future observations will allow the estimation of dM/dR around specific neutron star masses, M_i , we analyse below what constraints can be extracted on the radius and tidal deformability of NS from the value of $dM/dR|_{M=M_i}$. Note that the onset of an exotic degree of freedom, such as hyperons or deconfined quark matter, can cause the transition from $dM/dR > 0$ to $dM/dR < 0$. It is often predicted that the opening of new degrees of freedom occurs above twice the saturation density, inside stars of mass $\gtrsim 1.4 M_\odot$. We will therefore compare some properties of the EoS that can be affected by the slope at $1.4 M_\odot$, in particular, the radius of $2 M_\odot$ star and the radius and mass of maximum mass stars.




FIG. 7. Scatter plot of $dM/dR|_{M=1.4M_\odot}$ vs. $R(2.0M_\odot)$ (top) and the respective $R(2.0M_\odot)$ PDF (bottom). The colors indicate the $dM/dR|_{M=1.4M_\odot}$ value: negative (blue), positive but finite (red), and ∞ (black) with (dark colors) and without (light colors) astrophysical constraints.

Figure 7 displays the relation between the value of dM/dR at $M = 1.4 M_\odot$ and $R_{2.0M_\odot}$ (top), and the PDF for $R_{2.0M_\odot}$ (bottom) with (dark colors) and without (light colors) astrophysical constraints considering the three possibilities for $dM/dR|_{M=1.4M_\odot}$: negative (blue), positive but finite (red), and ∞ (black). The black set represents the EoSs for which the R remains almost constant with increasing M . Figures 8 and 9 show the same information but for the maximum mass, M_{\max} , and the

tidal deformability of a $1.4 M_\odot$ NS, respectively. Considering no astrophysical constraints (light colors in Figs. 7 to 9), we have 32035 EOS in the red set, 7328 in blue set, and 1072 in black set, while considering astrophysical constraints (dark colors in the same figures) the numbers decrease to 14395 EOS in red set, 6239 in blue set, and 980 in black set. Note that the number of EoS with $dM/dR < 0$ at $1.4 M_\odot$ is larger than the number of EoS with $dM/dR < 0$ over the whole range of masses as expected (if no constraints are imposed we have 7328 in the first set and 3106 in the second). A change of slope close to $1.4 M_\odot$ may just indicate some softening of the nuclear force at large densities (in a RMF description this could be played by the ω^4 term) or the onset of new degrees of freedom such as hyperons or quarks.

From Fig. 7, we conclude that $dM/dR < 0$ at $1.4 M_\odot$ implies a $2 M_\odot$ radius below 13 km. A radius above 13 km requires a positive slope at $1.4 M_\odot$, or at least an infinite slope. However, a radius below 13 km does not distinguish the values of dM/dR at $M = 1.4 M_\odot$. From the PDF (bottom panel), we see that a radius of 12 km or below will indicate $dM/dR < 0$ or an infinite slope, with a very high probability. An analysis of Fig. 8, which relates the slope of the $M(R)$ curve at $1.4 M_\odot$ with the maximum mass, shows that a positive slope at $1.4 M_\odot$ is necessary for the condition $M_{\max} \gtrsim 2.4 M_\odot$ to be satisfied. Concerning the tidal deformability $\Lambda_{1.4}$, a value below 800 does not exclude any set, but if in the future it is possible to restrict this property to values below 500 there is a high probability that the $M(R)$ curve at $1.4 M_\odot$ does not show a back-bending. Note that astrophysical constraints limit the star maximum mass to values below $\sim 2.6 M_\odot$ and the radius of $2.0 M_\odot$ and M_{\max} stars to values below 13.5 km, or $\lesssim 12$ km if $dM/dR < 0$ at $M = 1.4 M_\odot$ (for the maximum mass below 11.5 km at 90% CI with 12.12 km the extreme and for $2.0 M_\odot$ below 11.9 km at 90% CI). This is very interesting information to constrain the microscopic modeling of the EoS.

Figure 10 displays how local information regarding the sign of the slope of the $M(R)$ sequence at $1.2 M_\odot$ and $1.8 M_\odot$ translates into constraints on $R_{2.0M_\odot}$ and $\Lambda_{1.4M_\odot}$. The most constraining conditions, both for the $R_{2.0M_\odot}$ and the $\Lambda_{1.4M_\odot}$, come from an early onset of a negative slope or infinite slope, favoring in the first case radii below 12.5 km and a tidal deformability below 600 and in the second values of the radii below 13 km and of the tidal deformability below 700. Considering the joint information from the slope of 1.2 and 1.8 M_\odot does not constraint much considering the sets $dM/dR > 0$. However, if in both cases $dM/dR < 0$ or at $1.2 M_\odot$ $dM/dR \rightarrow \infty$ and at $1.8 M_\odot$ $dM/dR < 0$ then we may conclude that $10.5 < R_{2.0M_\odot} < 12.5$ km in the first case and $10.5 < R_{2.0M_\odot} < 13$ km in the second. If for both masses $dM/dR \rightarrow \infty$, a much more restrictive scenario, a stronger constraint is obtained: there is a very large probability that $11 < R_{2.0M_\odot} < 13$ km. We also conclude that the astrophysical constraints impose $\Lambda_{1.4M_\odot} \gtrsim 200$, in accordance with [44].




FIG. 8. Scatter plot of $dM/dR|_{M=1.4M_\odot}$ vs. $\{M_{\max}, R_{\max}\}$ (top panels) and the respective $\{M_{\max}, R_{\max}\}$ PDFs (bottom panels). The colors indicate the $dM/dR|_{M=1.4M_\odot}$ value: negative (blue), positive but finite (red), and ∞ (black) with (dark colors) and without (light colors) astrophysical constraints.




FIG. 9. Scatter plot of $dM/dR|_{M=1.4M_\odot}$ vs. $\Lambda(1.4M_\odot)$ (top) and the respective $\Lambda(1.4M_\odot)$ PDF (bottom). The colors indicate the $dM/dR|_{M=1.4M_\odot}$ value: negative (blue), positive but finite (red), and ∞ (black) with (dark colors) and without (light colors) astrophysical constraints.

IV. CONCLUSIONS

Considering a model-independent set of EoS constructed from five segments of polytropes, we have analysed the information that can be extracted from the slope of the $M(R)$ curve. The set of EoS was constrained by the neutron matter EoS obtained from a chiral effective field at low densities, the description of a $2M_\odot$ star, and several observational constraints, in particular the effective tidal deformability obtained from the GW170817 detection and the lower radius limits estimated by NICER for pulsars PSR J0740+6620 and PSR J0030+0451.

We have concluded that an EoS characterized by a negative slope throughout the mass range is quite restrictive: the maximum mass predicted is $2.20M_\odot$ with a radius greater than 10.3 km at 90% CI. If this constraint is relaxed, the maximum mass increases to $2.43M_\odot$ at 90% CI and can reach $\sim 2.69M_\odot$ and a maximum radius < 13.4 km. Note that this maximum mass has also been obtained in [45] using a different description for the EoS. A negative slope also predicts a different behavior of the speed of sound, showing a peak at $4-5n_0$ instead of $3n_0$ as often obtained in different studies [35, 42]. The conjecture proposed in [43] concerning the renormalized trace anomaly, which has a positive value at all densities, favors the EoS with back-bending, and as discussed in that work, the maximum masses obtained are smaller and, for a given maximum mass, the maximum star has a larger radius.

A rough estimate of the slope of $M(R)$ at two points, which allows the sign to be extracted, is already informa-




FIG. 10. The PDFs of the radius of $2.0M_{\odot}$ (left) and the tidal deformability of a $1.4M_{\odot}$ (right) with (bottom) and without (top) astrophysical constraints applied for $dM/dR|_{M=1.2M_{\odot}}$ (dark colors) and $dM/dR|_{M=1.8M_{\odot}}$ (light colors) values: negative (blue), positive but finite (red), and ∞ (black).

tive about neutron star observables such as radius and tidal deformability. We have analysed the consequences of the value the slope takes at $1.4M_{\odot}$ and concluded that a negative slope indicates $R(2M_{\odot}) < 11.9$ km and $R(M_{\max}) < 11.5$ km at 90% CI. The simultaneous determination of the slope at two different masses can also provide additional constraints: for an EoS with a negative slope at both masses, a star of $2M_{\odot}$ mass is expected to have a radius in the range $10.5 < R_{2.0M_{\odot}} < 12.5$ km. We conclude by pointing out that knowing the slope of the $M(R)$ curve at a given mass, or at more than one mass, will provide important information for constraining the microscopic EoS of hadronic matter at high densities.

ACKNOWLEDGMENTS

This work was partially supported by national funds from FCT (Fundação para a Ciência e a Tecnologia, I.P., Portugal) under the projects 2022.06460.PTDC with the DOI identifier 10.54499/2022.06460.PTDC, and UIDB/04564/2020 and UIDP/04564/2020, with DOI identifiers 10.54499/UIDB/04564/2020 and 10.54499/UIDP/04564/2020, respectively.

Appendix A: Datasets statistical summary

-
- [1] C. A. Raithel, F. Özel, and D. Psaltis, *Astrophys. J.* **844**, 156 (2017), arXiv:1704.00737 [astro-ph.HE].
 - [2] Y. Fujimoto, K. Fukushima, and K. Murase, *Phys. Rev. D* **101**, 054016 (2020), arXiv:1903.03400 [nucl-th].
 - [3] Y. Lim, A. Bhattacharya, J. W. Holt, and D. Pati, *Phys. Rev. C* **104**, L032802 (2021), arXiv:2007.06526 [nucl-th].
 - [4] G. Raaijmakers, S. K. Greif, K. Hebeler, T. Hinderer, S. Nissanke, A. Schwenk, T. E. Riley, A. L. Watts, J. M. Lattimer, and W. C. G. Ho, *Astrophys. J. Lett.* **918**, L29 (2021), arXiv:2105.06981 [astro-ph.HE].
 - [5] C. Huang, G. Raaijmakers, A. L. Watts, L. Tolos, and C. Providência, *Mon. Not. Roy. Astron. Soc.* **529**, 4650 (2024), arXiv:2303.17518 [astro-ph.HE].
 - [6] V. Carvalho, M. Ferreira, and C. Providência, *arXiv e-prints*, arXiv:2401.05770 (2024), arXiv:2401.05770 [nucl-th].
 - [7] A. Watts, “Measuring Neutron Star Mass and Radius Using Pulse Profile Modelling – NICER and Beyond,” (2024).

TABLE I. Percentiles (5th, 50th, and 95th) for some NS and EoS properties

	Set $dM/dR < 0$						Set $dM/dR \not< 0$					
	Without constraints			With constraints			Without constraints			With constraints		
	5%	50%	95%	5%	50%	95%	5%	50%	95%	5%	50%	95%
$M_{\max} [M_{\odot}]$	2.01	2.06	2.19	2.01	2.08	2.20	2.02	2.22	2.60	2.02	2.17	2.43
$R_{\max} [\text{km}]$	10.08	10.69	11.44	10.32	10.81	11.49	10.81	12.33	13.92	10.70	11.75	12.78
$n_{\max} [\text{fm}^{-3}]$	0.856	1.057	1.187	0.837	1.039	1.148	0.571	0.789	1.047	0.682	0.873	1.072
$v_s^2(n_{\max}) [c^2]$	0.117	0.547	0.944	0.113	0.506	0.929	0.034	0.326	0.847	0.039	0.374	0.869
$p(n_{\max}) [\text{MeV}/\text{fm}^3]$	313.679	529.452	765.946	305.920	500.879	739.310	109.545	289.643	585.372	165.490	343.686	620.589
$\Delta(n_{\max})$	-0.178	-0.070	0.054	-0.176	-0.057	0.062	-0.145	0.032	0.173	-0.152	0.008	1.141
$\Lambda(1.36M_{\odot})$	270	397	519	315	419	527	385	711	1089	360	567	710
$R(1.4M_{\odot}) [\text{km}]$	11.23	11.90	12.40	11.48	12.00	12.43	11.76	12.86	13.66	11.64	12.46	12.86
$n_{\max}(1.4M_{\odot}) [\text{fm}^{-3}]$	0.450	0.509	0.587	0.448	0.497	0.549	0.276	0.359	0.489	0.354	0.410	0.502
$\Lambda(1.4M_{\odot})$	223	329	432	262	349	439	323	607	943	303	480	606
$v_s^2(1.4M_{\odot}) [c^2]$	0.350	0.408	0.565	0.348	0.403	0.549	0.281	0.469	0.641	0.317	0.450	0.626
$R(1.6M_{\odot}) [\text{km}]$	11.10	11.77	12.30	11.39	11.88	12.32	11.74	12.94	13.86	11.62	12.48	11.95
$n_{\max}(1.6M_{\odot}) [\text{fm}^{-3}]$	0.504	0.574	0.662	0.499	0.560	0.615	0.289	0.390	0.544	0.377	0.449	0.558
$R(2.0M_{\odot}) [\text{km}]$	10.34	11.14	11.86	10.81	11.28	11.90	11.35	12.90	14.16	11.22	12.29	13.04
$n_{\max}(2.0M_{\odot}) [\text{fm}^{-3}]$	0.659	0.819	1.043	0.651	0.788	0.933	0.328	0.491	0.789	0.438	0.577	0.817

TABLE II. Extremes (maximum/minimum) for some NS and EoS properties

	Set $dM/dR < 0$				Set $dM/dR \not< 0$			
	Without constraints		With constraints		Without constraints		With constraints	
	min	max	min	max	min	max	min	max
$M_{\max} [M_{\odot}]$	2.0	2.3	2.0	2.3	2.0	3.07	2.0	2.69
$R_{\max} [\text{km}]$	9.66	12.12	10.08	12.12	9.54	15.00	10.07	13.38
$R(1.4M_{\odot}) [\text{km}]$	10.49	12.88	11.07	12.88	10.06	13.98	10.92	13.00
$n_{\max} [\text{fm}^{-3}]$	0.645	1.26	0.645	1.212	0.313	1.254	0.424	1.239

- [8] A. L. Watts, W. Yu, J. Poutanen, S. Zhang, S. Bhattacharyya, S. Bogdanov, L. Ji, A. Patruno, T. E. Riley, P. Bakala, A. Baykal, F. Bernardini, I. Bombaci, E. Brown, Y. Cavecchi, D. Chakrabarty, J. Chenevez, N. Degenaar, M. Del Santo, T. Di Salvo, V. Doroshenko, M. Falanga, R. D. Ferdman, M. Feroci, A. F. Gambino, M. Ge, S. K. Greif, S. Guillot, C. Gungor, D. H. Hartmann, K. Hebeler, A. Heger, J. Homan, R. Iaria, J. i. Zand, O. Kargaltsev, A. Kurkela, X. Lai, A. Li, X. Li, Z. Li, M. Linares, F. Lu, S. Mahmoodifar, M. Méndez, M. Coleman Miller, S. Morsink, J. Nättilä, A. Possenti, C. Prescod-Weinstein, J. Qu, A. Riggio, T. Salmi, A. Sanna, A. Santangelo, H. Schatz, A. Schwenk, L. Song, E. Šrámková, B. Stappers, H. Stiele, T. Strohmayer, I. Tews, L. Tolos, G. Török, D. Tsang, M. Urbanec, A. Vacchi, R. Xu, Y. Xu, S. Zane, G. Zhang, S. Zhang, W. Zhang, S. Zheng, and X. Zhou, *Science China Physics, Mechanics, and Astronomy* **62**, 29503 (2019), [arXiv:1812.04021 \[astro-ph.HE\]](#).
- [9] S.-N. Zhang *et al.* (eXTP), *Sci. China Phys. Mech. Astron.* **62**, 29502 (2019), [arXiv:1812.04020 \[astro-ph.IM\]](#).
- [10] P. S. Ray *et al.* (STROBE-X Science Working Group), *arXiv e-prints* (2019), [arXiv:1903.03035 \[astro-ph.IM\]](#).
- [11] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. Lett.* **121**, 161101 (2018), [arXiv:1805.11581 \[gr-qc\]](#).
- [12] Z. Carson, K. Chatziioannou, C.-J. Haster, K. Yagi, and N. Yunes, *Phys. Rev. D* **99**, 083016 (2019), [arXiv:1903.03909 \[gr-qc\]](#).
- [13] A. Watts *et al.*, *Proceedings, Advancing Astrophysics with the Square Kilometre Array (AASKA14): Giardini Naxos, Italy, June 9-13, 2014*, *PoS AASKA14*, 043 (2015), [arXiv:1501.00042 \[astro-ph.SR\]](#).
- [14] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, and R. Schaeffer, *Nucl. Phys. A* **635**, 231 (1998), [Erratum: *Nucl. Phys. A* 643, 441–441 (1998)].
- [15] A. W. Steiner, J. M. Lattimer, and E. F. Brown, *Astrophys. J. Lett.* **765**, L5 (2013), [arXiv:1205.6871 \[nucl-th\]](#).
- [16] W.-C. Chen and J. Piekarewicz, *Phys. Rev. C* **90**, 044305 (2014), [arXiv:1408.4159 \[nucl-th\]](#).
- [17] F. Douchin and P. Haensel, *Astron. Astrophys.* **380**, 151 (2001), [arXiv:astro-ph/0111092](#).
- [18] M. Fortin, C. Providencia, A. R. Raduta, F. Gulminelli, J. L. Zdunik, P. Haensel, and M. Bejger, *Phys. Rev. C* **94**, 035804 (2016), [arXiv:1604.01944 \[astro-ph.SR\]](#).
- [19] M. Fortin, M. Oertel, and C. Providência, *Publ. Astron. Soc. Austral.* **35**, 44 (2018), [arXiv:1711.09427 \[astro-ph.HE\]](#).
- [20] S. Typel, G. Ropke, T. Klahn, D. Blaschke, and H. H. Wolter, *Phys. Rev. C* **81**, 015803 (2010), [arXiv:0908.2344 \[nucl-th\]](#).

- [21] G. A. Lalazissis, T. Niksic, D. Vretenar, and P. Ring, *Phys. Rev. C* **71**, 024312 (2005).
- [22] C. J. Horowitz and J. Piekarewicz, *Phys. Rev. Lett.* **86**, 5647 (2001), arXiv:astro-ph/0010227.
- [23] H. Pais and C. Providência, *Phys. Rev. C* **94**, 015808 (2016), arXiv:1607.05899 [nucl-th].
- [24] L. Tolos, M. Centelles, and A. Ramos, *Publ. Astron. Soc. Austral.* **34**, e065 (2017), arXiv:1708.08681 [astro-ph.HE].
- [25] F. J. Fattoyev, C. J. Horowitz, J. Piekarewicz, and B. Reed, *Phys. Rev. C* **102**, 065805 (2020), arXiv:2007.03799 [nucl-th].
- [26] G. A. Lalazissis, J. König, and P. Ring, *Phys. Rev. C* **55**, 540 (1997), arXiv:nucl-th/9607039.
- [27] F. J. Fattoyev, C. J. Horowitz, J. Piekarewicz, and G. Shen, *Phys. Rev. C* **82**, 055803 (2010), arXiv:1008.3030 [nucl-th].
- [28] C. Ecker, T. Gorda, A. Kurkela, and L. Rezzolla, (2024), arXiv:2403.03246 [astro-ph.HE].
- [29] J. S. Read, B. D. Lackey, B. J. Owen, and J. L. Friedman, *Phys. Rev. D* **79**, 124032 (2009), arXiv:0812.2163 [astro-ph].
- [30] K. Hebeler, J. M. Lattimer, C. J. Pethick, and A. Schwenk, *Astrophys. J.* **773**, 11 (2013), arXiv:1303.4662 [astro-ph.SR].
- [31] F. Douchin and P. Haensel, *Astron. Astrophys.* **380**, 151 (2001), arXiv:astro-ph/0111092.
- [32] J. R. Oppenheimer and G. M. Volkoff, *Physical Review* **55**, 374 (1939).
- [33] R. C. Tolman, *Physical Review* **55**, 364 (1939).
- [34] T. Hinderer, B. D. Lackey, R. N. Lang, and J. S. Read, *Phys. Rev. D* **81**, 123016 (2010), arXiv:0911.3535 [astro-ph.HE].
- [35] S. Altiparmak, C. Ecker, and L. Rezzolla, *Astrophys. J. Lett.* **939**, L34 (2022), arXiv:2203.14974 [astro-ph.HE].
- [36] M. C. Miller *et al.*, *Astrophys. J. Lett.* **918**, L28 (2021), arXiv:2105.06979 [astro-ph.HE].
- [37] T. E. Riley *et al.*, *Astrophys. J. Lett.* **918**, L27 (2021), arXiv:2105.06980 [astro-ph.HE].
- [38] T. E. Riley *et al.*, *Astrophys. J. Lett.* **887**, L21 (2019), arXiv:1912.05702 [astro-ph.HE].
- [39] M. C. Miller *et al.*, *Astrophys. J. Lett.* **887**, L24 (2019), arXiv:1912.05705 [astro-ph.HE].
- [40] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev.* **X9**, 011001 (2019), arXiv:1805.11579 [gr-qc].
- [41] R. Kumar *et al.* (MUSES), *Living Rev. Rel.* **27**, 3 (2024), arXiv:2303.17021 [nucl-th].
- [42] E. Annala, T. Gorda, A. Kurkela, J. Nättilä, and A. Vuorinen, *Nature Phys.* **16**, 907 (2020), arXiv:1903.09121 [astro-ph.HE].
- [43] Y. Fujimoto, K. Fukushima, L. D. McLerran, and M. Praszalowicz, *Phys. Rev. Lett.* **129**, 252702 (2022), arXiv:2207.06753 [nucl-th].
- [44] D. Radice and L. Dai, *Eur. Phys. J. A* **55**, 50 (2019), arXiv:1810.12917 [astro-ph.HE].
- [45] M. Ferreira and C. Providência, *Phys. Rev. D* **104**, 063006 (2021), arXiv:2110.00305 [nucl-th].