The Laplace asymptotic expansion in high dimensions: a nonasymptotic analysis

Anya Katsevich* akatsevi@mit.edu

June 19, 2024

Abstract

We study the classical Laplace asymptotic expansion of $\int_{\mathbb{R}^d} f(x)e^{-nv(x)}dx$ in high dimensions d. We derive an error bound to the expansion when truncated to arbitrary order. The error bound is fully explicit except for absolute constants, and it depends on d, n, and operator norms of the derivatives of v and f in a neighborhood of the minimizer of v.

1 Introduction

In one of its many forms, the classical Laplace approximation states [27, Chapter 2] that if f and v are sufficiently smooth and v has a unique strict minimizer $x_0 \in (a, b)$, then

$$\int_{a}^{b} f(x)e^{-nv(x)}dx = \sqrt{\frac{2\pi}{nv''(x_0)}}e^{-nv(x_0)}\left(f(x_0) + o(1)\right), \quad n \to \infty$$

More generally, the Laplace method gives a full asymptotic expansion of such "Laplace-type" integrals in any fixed dimension d. Assuming f and v are sufficiently smooth on a possibly unbounded domain $D \subseteq \mathbb{R}^d$, the integral $\int_D f(x)e^{-nv(x)}dx$ converges absolutely for all n large enough, and v has a unique strict global minimizer x_0 on D such that $\inf_{x\in D, ||x-x_0|| \ge \epsilon} v(x) - v(x_0) > 0$ for all $\epsilon > 0$, then [27, Chapter 9]

$$\int_{D} f(x)e^{-nv(x)}dx = \frac{(2\pi)^{d/2}}{\sqrt{\det(n\nabla^2 v(x_0))}}e^{-nv(x_0)}\left(f(x_0) + \sum_{k=1}^{M-1} a_k n^{-k} + \mathcal{O}(n^{-M})\right),\tag{1.1}$$

for some coefficients a_k . The order zero expansion (i.e. M = 1) is typically referred to as the Laplace approximation. Explicit expressions for the coefficients a_k have been obtained in [19, 26] in the one-dimensional case, and most explicitly by [14] in the general *d*-dimensional case. See also [4] for a procedure to compute the coefficients a_k in the multivariate setting.

Here, we are interested in studying the expansion (1.1) in high dimension. To be self-contained, we present our own derivation of the expansion and arrive at the same coefficients a_k as in [14], though we take a different route that avoids series inversion. For the expansion to be useful in this high-dimensional setting, careful error bounds on the remainder are required. The bounds should explicitly quantify the dependence of the error on d itself, and on all quantities which might increase with dimension: namely, f and v. We derive an error bound for the expansion (1.1) which is fully explicit up to absolute constants. It is given by a function of d, n, and operator norms of the derivatives of v and f in a neighborhood of the minimizer of v. In a very rough sense, the error bound reveals that $d^2 \ll n$ is sufficient for accuracy of the expansion. A more precise characterization for the range of applicability of the Laplace expansion is simply that it is accurate for any d, n, v, f for which our error bound is small.

To the best of our knowledge, this is the first work to obtain such an explicit error bound on the Laplace asymptotic expansion to arbitrary order. Explicit error bounds on the zeroth order expansion (i.e. the

^{*}This work was supported by NSF grant DMS-2202963

case L = 1 in (1.1)) do exist for the general form of the integral $\int f e^{-nv}$, but most such bounds blow up exponentially with d. Up to the f and v dependence, [17] obtains that

$$\int_{D} f(x)e^{-nv(x)}dx = \frac{(2\pi)^{d/2}}{\sqrt{\det(n\nabla^2 v(x_0))}}e^{-nv(x_0)}\left(f(x_0) + \epsilon\right), \qquad \epsilon = \mathcal{O}\left(\sqrt{d^3/n}\right)$$

However, incorporating the v dependence, the bound on ϵ actually involves $\det(\nabla^2 v(x_0))/(\lambda_{\min}(\nabla^2 v(x_0)))^d$, which will blow up exponentially with d unless the eigenvalues λ_i of $\nabla^2 v(x_0)$ scale as $\lambda_i/\lambda_{\min} \sim 1 + 1/d$. In the case of higher regularity, [16] proves that the $\mathcal{O}(\sqrt{d^3/n})$ error bound on ϵ be improved to $\mathcal{O}(d^3/n)$. This matches the powers of n^{-1} arising in (1.1). However, again, the bound on ϵ involves $\det(\nabla^2 v(x_0))/(\lambda_{\min}(\nabla^2 v(x_0)))^d$. The work [11] also obtains explicit bounds on the error, but the bounds also blow up exponentially with d, due to the term K_l .

Thus in the literature focusing on the theory of the Laplace expansion in a general context, there have not been truly high-dimensional error bounds (to our knowledge). On the other hand, in the context of Bayesian inference, tighter dimension-dependent bounds have been obtained. See e.g. [13, 12, 25] for an explanation of how Laplace-type integrals show up in Bayesian inference. The latter of these three paper is foundational in introducing the Laplace approximation in statistics. In the Bayesian context, n refers to sample size and d refers to parameter dimension. Two types of integrals are of interest in Bayesian inference: first, the posterior normalizing constant, which is of interest in Bayesian model selection, can be written as either $\int e^{-nv}$ or $\int f e^{-nv}$ (depending on whether $e^{-nv} = \text{likelihood} \times \text{prior or } e^{-nv} = \text{likelihood}$ and f = prior. In either case, the function v will be weakly dependent on n). Second, posterior expectations of an observable f can be written as the ratio of two Laplace-type integrals: $\int f e^{-nv} / \int e^{-nv}$, where the posterior is proportional to $e^{-nv} = \text{likelihood} \times \text{prior}$. The work [3] considers the posterior normalizing constant for a generalized linear model. For simplicity, we relate their results in the case of a flat prior, i.e. prior $\equiv 1$. The authors shows that if e^{-nv} is the likelihood in a generalized linear model with d coefficients, and x_0 is the minimizer of v, then

$$\int_{\mathbb{R}^d} e^{-nv(x)} dx = \frac{(2\pi)^{d/2}}{\sqrt{\det(n\nabla^2 v(x_0))}} e^{-nv(x_0)} \left(1+\epsilon\right), \qquad |\epsilon| \le C(c_1, c_2, c_3)\sqrt{(d\log n)^3/n},$$

where c_1, c_2, c_3 are defined via the assumption that $c_1 I_d \leq \nabla^2 v(x) \leq c_2 I_d$ uniformly in a neighborhood of x_0 and $\|\nabla^2 v(x) - \nabla^2 v(x')\| \leq c_3 \|x - x'\|$ for all $x, x' \in \mathbb{R}^d$. It is reasonable to assume c_1, c_2, c_3 do not blow up exponentially with d, so this bound does not have hidden exponential dimension dependence.

The work [20] also bounds the Laplace expansion error (after truncating to arbitrary order) for the normalizing constant $\int e^{-nv}$. Although the formulation of the result is general, it is unclear how to express the error bound explicitly in this general context. The author applies the general result to two-level and multi-level random intercept models, which are kinds of generalized linear mixed models. In these examples, the derivative tensors $\nabla^k v(0)$ are diagonal for $k \geq 3$.

Finally, [23] has also studied the Laplace expansion of $\int e^{-nv}$ in high dimensions using nonrigorous arguments. In particular, the authors consider the normalizing constant for a posterior in the context of an exchangeable binary array model and generalized linear models (GLMs). (The latter model is referred to as an "exponential model" in the paper). The authors show that in the first of these models, if $d = \mathcal{O}(n^{1/2})$, then the correction term a_1n^{-1} in (1.1) has order $\mathcal{O}(1)$. They also claim that for the case of GLMs, the correction term a_1n^{-1} is not small if d^3/n is not small, and therefore the standard Laplace expansion is not valid. However, this argument is based on counting the numbers of terms in a sum arising in the formula for a_1 . But in fact, we show in Appendix A that $|a_1|n^{-1} \leq (||\nabla^3 v(0)||^2 + ||\nabla^4 v(0)||)d^2/n$. Moreover, the work [13] shows that for logistic regression (a standard example of a GLM), we have $||\nabla^3 v(0)||, ||\nabla^4 v(0)|| = \mathcal{O}(1)$.

Most other works on the Laplace method in Bayesian inference do not focus on asymptotic expansions of integrals, but rather on approximating probability densities $\pi \propto e^{-nv}$ with the Gaussian density $\hat{\gamma}(x) \propto \exp(-(x-x_0)^{\mathsf{T}}\nabla^2(nv)(x_0)(x-x_0)/2)$. In fact, in the context of Bayesian inference, the term "Laplace approximation" generally refers to this approximation of a probability density, and not to the zeroth order Laplace asymptotic expansion. If π is close to $\hat{\gamma}$ in some sense, then $\int f d\pi \approx \int f d\hat{\gamma}$, and the latter integral can be more easily evaluated numerically. There have been a number of works studying the approximation $\pi \approx \hat{\gamma}$ in high dimensions; see for example [24, 13, 12, 8, 7, 10, 9]. **Organization.** In Section 2 we state our assumptions and main result on the Laplace asymptotic expansion, with some discussion. Section 3 outlines the proof, with some lemmas deferred to the Appendix.

1.1 Notation

The measure γ denotes the standard Gaussian measure on \mathbb{R}^d : $\gamma(dx) = (2\pi)^{-d/2} e^{-\|x\|^2/2} dx$. For a function $f : \mathbb{R}^d \to \mathbb{R}$, the norm $\|f\|_q$ always denotes $\|f\|_{L_q(\gamma)}$, i.e. for a function $f : \mathbb{R}^d \to \mathbb{R}$, we have

$$||f||_q = \left(\int |f(x)|^q \gamma(dx)\right)^{\frac{1}{q}}$$

We say that $a \leq b$ if $a \leq Cb$ for some absolute constant C. We say that $a \leq_L b$ if $a \leq C(L)b$ for some constant C(L) depending only on L. A tensor T of order k is an array $T = (T_{i_1i_2...i_k})_{i_1,...,i_k=1}^d$. For two order k tensors T and S we let $\langle T, S \rangle$ be the entrywise inner product. We say T is symmetric if $T_{i_1...i_k} = T_{j_1...j_k}$, for all permutations $j_1 \ldots j_k$ of $i_1 \ldots i_k$. We define

$$||T|| = \sup_{||u||=1} \langle T, u^{\otimes k} \rangle = \sup_{||u||=1} \sum_{i_1, \dots, i_k=1}^d T_{i_1 i_2 \dots i_k} u_{i_1} u_{i_2} \dots u_{i_k}$$
(1.2)

By [28, Theorem 2.1], for symmetric tensors, the definition (1.2) coincides with the standard definition of operator norm:

$$\sup_{u_1\parallel=\cdots=\parallel u_k\parallel=1} \langle T, u_1 \otimes \cdots \otimes u_k \rangle = \parallel T \parallel = \sup_{\parallel u \parallel=1} \langle T, u^{\otimes k} \rangle.$$

Finally, $\mathbb{1}_{\mathcal{U}} : \mathbb{R}^d \to \mathbb{R}$ is the indicator of the set \mathcal{U} .

2 Main results

In this section, we present our results. We start in Section 2.1 by stating our assumptions and introducing important quantities. In Section 2.2 we present our high-dimensional Laplace asymptotic expansion of $\int f e^{-nv}$, and the specialization to the case $\int e^{-nv}$. When discussing orders of magnitude throughout this section, we will focus on the case $\log n \leq d$ as a model example of a high-dimensional regime. However, our results do not require this assumption.

2.1 Assumptions and important quantities

We begin with a reparameterization: given $g : \mathbb{R}^d \to \mathbb{R}$ and a twice differentiable $\phi : \mathbb{R}^d \to \mathbb{R}$ minimized at zero with Hessian $H = \nabla^2 \phi(0) \succ 0$, note that

$$\int g(y)e^{-n\phi(y)}dy = \frac{1}{\sqrt{\det H}} \int f(x)e^{-nv(x)}dx, \qquad v(x) = \phi(H^{-1/2}x), \quad f(x) = g(H^{-1/2}x).$$

Furthermore, note that

$$\|\nabla^{k} v(x)\| = \|\nabla^{k} \phi(H^{-1/2}x)\|_{H} := \sup_{\|H^{1/2}u\|=1} \langle \nabla^{k} \phi(H^{-1/2}x), u^{\otimes k} \rangle_{H}$$
$$\sup_{\|x\| \le r} \|\nabla^{k} v(x)\| = \sup_{\|H^{1/2}y\| \le r} \|\nabla^{k} \phi(y)\|_{H}.$$

Derivatives of f can similarly be written in terms of derivatives of g. We therefore work with f and v throughout this work. All assumptions on f and v can be translated into assumptions on g and ϕ . In line with this discussion, we have the following first assumption.

Assumption 2.1 (Minimizer of v). The function v has a unique global minimizer at the origin. Furthermore, v is twice differentiable in a neighborhood of the origin to be specified, and $\nabla^2 v(0) = I_d$.

For a fixed radius R > 0 to be specified later, we let $\mathcal{U} = \{x : ||x|| \le R\sqrt{d}\}$ and $\frac{2}{\sqrt{n}}\mathcal{U} = \{\frac{2}{\sqrt{n}}x : x \in \mathcal{U}\}$. As discussed below in Remark 2.10, when $\log n \le d$, we can take R to be an absolute constant; otherwise, R should scale as $\log n/d$.

Assumption 2.2 (Regularity of f and v). The functions f and v are L and L + 2 times continuously differentiable in $\frac{2}{\sqrt{n}}\mathcal{U}$, respectively, for an even $L \ge 2$.

Assumption 2.3 (Growth of v and f at infinity). There exists $a_v \in (0, 1]$ such that

$$v(x) - v(0) \ge a_v \sqrt{d/n} \|x\| \qquad \forall \|x\| \ge \sqrt{d/n}.$$
(2.1)

For the same a_v , there exists some $b_f > 0$ such that

$$|f(x) - f(0)| \le b_f \exp(a_v \sqrt{dn} ||x||/2), \quad \forall x \in \mathbb{R}^d.$$

Remark 2.4. This assumption ensures that the tail integral of fe^{-nv} is exponentially small. However, it is clearly not the only assumption that can be imposed to achieve the desired decay. Note also that there is some C such that for all $\sqrt{d/n} \leq ||x|| \leq C$, we have

$$v(x) - v(0) \ge ||x||^2 / 4 \ge \frac{1}{4}\sqrt{d/n}||x||.$$

This follows by a Taylor expansion of v around zero, since $v(x) \sim ||x||^2/2$ near zero. Therefore, (2.1) is satisfied for small values of ||x|| with $a_v = 1/4$, so in reality, (2.1) is a condition on the growth of v far away from zero.

Definition 2.5 (Derivative bounds). For all $k \ge 1$, we define

$$c_k(R) = \sup_{\|x\| \le R\sqrt{d/n}} \|\nabla^k v(x)\|.$$

For all $k \ge 0$, we define

$$c_{k,f}(R) = \sup_{\|x\| \le R\sqrt{d/n}} \|\nabla^k f(x)\|$$

Definition 2.6 (Key error terms). For all $k \ge 1$, define

$$\epsilon_k(R) = c_{k+2}(R) \frac{\sqrt{d}^{k+2}}{\sqrt{n}^k},$$

$$\bar{\epsilon}_k(R) = (1 \vee R)^{k+1} c_{k+2}(R) \frac{\sqrt{d}^{k+1}}{\sqrt{n}^k}.$$
(2.2)

Note that $\epsilon_k(R)$ and $\bar{\epsilon}_k(R)$ both scale with n as $n^{-k/2}$. In the "high-dimensional" case (log $n \leq d$) in which R is an absolute constant (see Remark 2.10), we have $\bar{\epsilon}_k(R) \leq d^{-1/2} \epsilon_k(R)$. Next, we define compound error terms which build on the ϵ 's.

Definition 2.7 ("Compound" error terms). Let $2 \le k \le L$. For k even, we define $\delta_k(R)$ as

$$\delta_k(R) = \epsilon_k(R) = c_{k+2}(R) \frac{\sqrt{d}^{k+2}}{\sqrt{n}^k}$$
(2.3)

For k odd, we define $\delta_k(R)$ as

$$\delta_k(R) = \min\left(\epsilon_k(R), \ \bar{\epsilon}_k(0) + \epsilon_{k+1}(R)\right) = \min\left(c_{k+2}(R)\frac{\sqrt{d}^{k+2}}{\sqrt{n^k}}, \ c_{k+2}(0)\frac{\sqrt{d}^{k+1}}{\sqrt{n^k}} + c_{k+3}(R)\frac{\sqrt{d}^{k+3}}{\sqrt{n^{k+1}}}\right).$$
(2.4)

Which term is smaller in the minimum depends on the choice of R, and on how c_{k+2}, c_{k+3} scale with R. Let us write out the first four δ 's explicitly, which will arise in error bounds on the first order expansion; see Example 2.12 below. We have

$$\delta_1(R) = \min\left(c_3(R)\frac{d^{1.5}}{\sqrt{n}}, c_3(0)\frac{d}{\sqrt{n}} + c_4(R)\frac{d^2}{n}\right),$$

$$\delta_2(R) = c_4(R)\frac{d^2}{n},$$

$$\delta_3(R) = \min\left(c_5(R)\frac{d^{2.5}}{n^{1.5}}, c_5(0)\frac{d^2}{n^{1.5}} + c_6(R)\frac{d^3}{n^2}\right),$$

$$\delta_4(R) = c_6(R)\frac{d^3}{n^2}$$
(2.5)

We see that the ratios d^{α}/n^{β} which show up in $\delta_1, \ldots, \delta_4$ are each bounded by some power of d/\sqrt{n} . The only "close call" is the case k = 1, for which $\epsilon_1(R)$ contains the ratio $d^{1.5}/\sqrt{n} > d/\sqrt{n}$. But since $\delta_1(R)$ is defined as the minimum between $\epsilon_1(R)$ and $\bar{\epsilon}_1(0) + \epsilon_2(R)$, we have that $\delta_1(R)$ is no greater than a power of d/\sqrt{n} , up to the dependence on $c_3(R), c_4(R)$. This is a reflection of the fact that

$$\epsilon_k(R) \le c_{k+2}(R) \left(\frac{d}{\sqrt{n}}\right)^k, \quad \forall k \ge 2,$$

$$\bar{\epsilon}_k(0) \le c_{k+2}(0) \left(\frac{d}{\sqrt{n}}\right)^k, \quad \forall k \ge 1,$$

(2.6)

Thus up to the dependence on $c_k(R)$, all $\delta_k(R)$, $k \ge 1$ are bounded by a power of d/\sqrt{n} . In fact as k increases, the ratios $\sqrt{d}^{k+2}/\sqrt{n}^k$ and $\sqrt{d}^{k+1}/\sqrt{n}^k$ become closer and closer to a power of $\sqrt{d/n}$.

2.2 Laplace integral expansion

First, we define a few more quantities. Let

$$B_k(s_1, \dots, s_k) = k! \sum_{\substack{j_1+2j_2+\dots+kj_k=k\\j_1,\dots,j_k \ge 0}} \prod_{i=1}^k \frac{s_i^{j_i}}{(i!)^{j_i}(j_i)!}$$
(2.7)

be the kth complete Bell polynomial [1, Chapter 12]. For example, $B_0 = 1$, $B_1(s_1) = s_1$, and $B_2(s_1, s_2) = s_1^2 + s_2$. Next, we define a polynomial $b_k(0, x)$, with $x \in \mathbb{R}^d$, as follows:

$$b_k(0,x) = B_k\left(\frac{-\langle \nabla^3 v(0), x^{\otimes 3} \rangle}{2 \cdot 3}, \dots, \frac{-\langle \nabla^{k+2} v(0), x^{\otimes k+2} \rangle}{(k+1)(k+2)}\right).$$
(2.8)

The argument 0 indicates that we evaluate the derivatives of v at zero. For example, we have

$$b_1(0,x) = -\frac{1}{6} \langle \nabla^3 v(0), x^{\otimes 3} \rangle, \qquad b_2(0,x) = \frac{1}{36} \langle \nabla^3 v(0), x^{\otimes 3} \rangle^4 - \frac{1}{12} \langle \nabla^4 v(0), x^{\otimes 4} \rangle.$$

Theorem 2.8. Let L be even. Suppose Assumptions 2.1, 2.2, and 2.3 hold, and that $R \ge \frac{8}{a_v} \log \frac{2e}{a_v} \ge 8$. Then

$$\frac{e^{nv(0)}}{(2\pi/n)^{d/2}} \int_{\mathbb{R}^d} f(x)e^{-nv(x)}dx = f(0) + \sum_{k=1}^{L/2-1} A_{2k}n^{-k} + \operatorname{Rem}_L,$$
(2.9)

where the A_{2k} are given by the explicit formula in Theorem 3.6, and can be written as the following Gaussian expectation:

$$A_{2k} = \frac{1}{2k!} \sum_{\ell=0}^{2k} {\binom{2k}{\ell}} \mathbb{E} \left[\langle \nabla^{2k-\ell} f(0), Z^{\otimes 2k-\ell} \rangle b_{\ell}(0, Z) \right].$$
(2.10)

Here, $Z \sim \mathcal{N}(0, I_d)$, and $b_{\ell}(0, \cdot)$ is the function defined in (2.8). Finally, the remainder Rem_L is bounded by

$$|\operatorname{Rem}_L| \lesssim_L \kappa_L + \tau_L + \tau_{\mathcal{U}^c}, \tag{2.11}$$

where

$$\kappa_{L} = e^{\bar{\epsilon}_{1}(R)^{2} + \epsilon_{2}(R)} \sum_{\ell=0}^{L} c_{L-\ell}(R) \sqrt{d/n}^{L-\ell} B_{\ell}(\delta_{1}(R), \dots, \delta_{\ell}(R)),$$

$$\tau_{L} = \max_{0 \le m \le L-1} \|\nabla^{m} f(0)\| \sqrt{d/n}^{m} \max_{1 \le j \le L-1} (\epsilon_{j}(0) \lor 1)^{L} e^{-(R-1)^{2}d/4},$$

$$\tau_{\mathcal{U}^{c}} = (|f(0)| + b_{f}) de^{-Ra_{v}d/8},$$
(2.12)

and B_{ℓ} is as in (2.7).

Remark 2.9 (Scaling with n of κ_L). Treating v and f as fixed with respect to n (which need not always be true), the overall scaling with n of κ_L is $\kappa_L \sim n^{-L/2}$. This is because $B_\ell(\delta_1(R), \ldots, \delta_\ell(R)) \sim n^{-\ell/2}$, which follows from the fact that $\delta_i(R) \sim n^{-i/2}$ and the definition (2.7) of the Bell polynomials.

Remark 2.10 (Exponentially small error terms and choice of R). Suppose for simplicity that b_f , $\|\nabla^m f(0)\|$, $\|\nabla^{m+2}v(0)\|$, $m = 0, \ldots, L-1$ are all bounded by some polynomial of d. Then the remainder terms $\tau_{\mathcal{U}^c}$ and τ_L are exponentially small in d. In particular, for a large enough constant C_L , we can choose $R = \max(\frac{8}{a_v} \log \frac{2e}{a_v}, C_L(\log n)/d)$ to ensure $\tau_{\mathcal{U}^c}, \tau_L \leq n^{-L/2}$. Then these two terms are negligible compared to the first error term κ_L , which scales with n as $n^{-L/2}$. Supposing a_v can be chosen to be an absolute constant (see the discussion in Remark 2.1), we then have $R = C_L \max(1, (\log n)/d)$. Recall that $c_{L-\ell,f}(R)$ and $\delta_j(R)$ involve the supremum of derivatives of f and v over a neighborhood of size $R\sqrt{d/n} = C_L \max(\sqrt{d/n}, (\log n)/\sqrt{nd})$. Thus if $d \ll n$ and $\log n \ll \sqrt{nd}$ then the neighborhood is very small.

Remark 2.11 (Relationship between d and n). From the definition of κ_L in (2.12), we see that this main error term is small provided $\bar{\epsilon}_1(R)$ and $\epsilon_2(R)$ are bounded and $\delta_k(R)$, $k = 1, \ldots, L$ and $c_{k,f}(R)\sqrt{d/n}^k$, $k = 1, \ldots, L$ are small. Suppose that $c_{k,f}(R)$ and $c_k(R)$ are bounded by absolute constants for all k. Suppose also that $\log n \leq d$, so we can take R to be constant, as per the above remark. Then

$$\bar{\epsilon}_1(R) = c_3(R) \frac{R^2 d}{\sqrt{n}}, \qquad \epsilon_2(R) = c_4(R) \frac{d^2}{n}$$
(2.13)

are bounded if $d^2 \leq n$. Moreover, as we have discussed following (2.5), the $\delta_k(R)$ are no greater than some power of d^2/n under the above assumptions. We conclude that $\kappa_L \ll 1$ if $d^2 \ll n$. This is the interpretation of our claim in the introduction that "in a very rough sense.... $d^2 \ll n$ is sufficient for accuracy of the expansion".

Example 2.12 (Leading term and error for the case L = 4). Using that $B_0 = 1$, $B_1(s_1) = s_1$, and $B_2(s_1, s_2) = s_1^2 + s_2$, we obtain

$$A_{2} = \frac{1}{2} \mathbb{E} \left[\langle \nabla^{2} f(0), Z^{\otimes 2} \rangle \right] - \frac{1}{6} \mathbb{E} \left[\langle \nabla f(0), Z \rangle \langle \nabla^{3} v(0), Z^{\otimes 3} \rangle \right] + \frac{f(0)}{72} \mathbb{E} \left[\langle \nabla^{3} v(0), Z^{\otimes 3} \rangle^{2} \right] - \frac{f(0)}{24} \mathbb{E} \left[\langle \nabla^{4} v(0), Z^{\otimes 4} \rangle \right],$$

$$(2.14)$$

which can be computed explicitly to be

$$A_{2} = \frac{1}{2} \Delta f(0) - \frac{1}{2} \sum_{i,j=1}^{d} \partial_{i} f(0) \partial_{ijj}^{3} v(0) + \frac{f(0)}{12} \|\nabla^{3} v(0)\|_{F}^{2} + \frac{f(0)}{8} \sum_{i=1}^{d} (\partial_{i} \Delta v(0))^{2} - \frac{f(0)}{8} \sum_{i,j=1}^{d} \partial_{i}^{2} \partial_{j}^{2} v(0).$$

$$(2.15)$$

For the main error term κ_4 , we have

$$\kappa_{4} \lesssim \exp\left(\bar{\epsilon}_{1}^{2} + \epsilon_{2}\right) \left[c_{0,f} \left(\delta_{1}^{4} + \delta_{1}^{2} \delta_{2} + \delta_{1} \delta_{3} + \delta_{2}^{2} + \delta_{4} \right) + c_{1,f} \sqrt{d/n} (\delta_{1}^{3} + \delta_{1} \delta_{2} + \delta_{3}) + c_{2,f} \sqrt{d/n}^{2} (\delta_{1}^{2} + \delta_{2}) + c_{3,f} \sqrt{d/n}^{3} \delta_{1} + c_{4,f} \sqrt{d/n}^{4} \right],$$

$$(2.16)$$

where we have omitted the argument R for brevity. Here, $\bar{\epsilon}_1(R)$, $\epsilon_2(R)$ are as in (2.13) and the δ 's have been written out explicitly in (2.5).

By setting $f \equiv 1$, we obtain the following corollary.

Corollary 2.13 (Expansion of the partition function). Under the conditions on v from Theorem 2.8, we have

$$\frac{e^{nv(0)}}{(2\pi/n)^{d/2}} \int_{\mathbb{R}^d} e^{-nv(x)} dx = 1 + \sum_{k=1}^{L/2-1} A_{2k} n^{-k} + \operatorname{Rem}_L,$$
(2.17)

where

$$A_{2k} = \frac{1}{(2k)!} \mathbb{E} \left[b_{2k}(0, Z) \right], \qquad (2.18)$$

for $b_{2k}(0,\cdot)$ as in (2.8). The A_{2k} are also given by the explicit formula in Corollary 3.7. The remainder Rem_L is bounded by

$$|\operatorname{Rem}_L| \lesssim_L \kappa_L + \tau_L + \tau_{\mathcal{U}^c}, \tag{2.19}$$

where

$$\kappa_L = e^{\bar{\epsilon}_1(R)^2 + \epsilon_2(R)} B_L(\delta_1(R), \dots, \delta_L(R)),$$

$$\tau_L = \max_{1 \le j \le L-1} (\epsilon_j(0) \lor 1)^L e^{-(R-1)^2 d/4},$$

$$\tau_{\mathcal{U}^c} = de^{-Ra_v d/8},$$

(2.20)

and B_L is as in (2.7).

Example 2.14 (Leading term and error for the case L = 4). We can obtain the leading term and error bound by substituting $f \equiv 1$ in the formulas from Example 2.12. This gives

$$A_{2} = \frac{1}{12} \|\nabla^{3} v(0)\|_{F}^{2} + \frac{1}{8} \sum_{i=1}^{d} \left(\partial_{i} \Delta v(0)\right)^{2} - \frac{1}{8} \sum_{i,j=1}^{d} \partial_{i}^{2} \partial_{j}^{2} v(0).$$
(2.21)

Similarly, for the main error term κ_4 , we have

$$\kappa_4 \lesssim \exp\left(\bar{\epsilon}_1^2 + \epsilon_2\right) \left(\delta_1^4 + \delta_1^2 \delta_2 + \delta_1 \delta_3 + \delta_2^2 + \delta_4\right).$$
(2.22)

3 Proof of Theorem 2.8

In Section 3.1, we make some initial simplifications to the integral of interest. Then in Section 3.2, we employ a Taylor expansion in the small parameter $1/\sqrt{n}$. This leads to a decomposition of the integral into leading and remainder terms, which we analyze in Sections 3.3 and 3.4, respectively.

3.1 Initial simplifications

Define the functions $F, W, r : \mathbb{R} \times \mathbb{R}^d \to \mathbb{R}$ as follows:

$$F(t,x) = f(tx),$$

$$W(t,x) = \begin{cases} \frac{v(tx) - v(0)}{t^2}, & t \neq 0, \\ \|x\|^2/2, & t = 0, \end{cases}$$

$$r(t,x) = W(t,x) - \|x\|^2/2.$$
(3.1)

Recall that $\mathcal{U} = \{x \in \mathbb{R}^d : ||x|| \leq R\sqrt{d}\}$. Using basic manipulations, we obtain the following initial decomposition of the integral of interest.

Lemma 3.1. We have

$$\frac{e^{nv(0)}}{(2\pi/n)^{d/2}} \int f(x)e^{-nv(x)}dx
= \int_{\mathcal{U}} F(1/\sqrt{n}, x) e^{-r(1/\sqrt{n}, x)}\gamma(dx) + (2\pi)^{-d/2} \int_{\mathcal{U}^c} F(1/\sqrt{n}, x) e^{-W(1/\sqrt{n}, x)}dx
=: I(1/\sqrt{n}) + \tau_{\mathcal{U}^c},$$
(3.2)

where F, W, r, U are as above, and $\gamma(dx)$ is the standard Gaussian measure.

See Appendix **B** for the proof.

3.2 Taylor expansion of local integral

In this section, we analyze the local integral $I(1/\sqrt{n})$ from (3.2). We first state the main lemma, and then outline its proof below. Recall that $Z \sim \mathcal{N}(0, I_d)$.

Lemma 3.2. Let

$$b_{\ell}(t,x) = B_{\ell}\left(-\partial_t r(t,x), -\partial_t^2 r(t,x), \dots, -\partial_t^\ell r(t,x)\right).$$
(3.3)

Then there exists $0 \le t \le 1/\sqrt{n}$ such that

$$I(1/\sqrt{n}) = \text{Lead}_L + \kappa_L(t) + \tau_L,$$

where

$$\text{Lead}_{L} = f(0) + \sum_{k=1}^{L-1} \frac{n^{-k/2}}{k!} \sum_{\ell=0}^{k} \binom{k}{\ell} \mathbb{E} \left[\langle \nabla^{k-\ell} f(0), Z^{\otimes k-\ell} \rangle b_{\ell}(0, Z) \right]$$
(3.4)

and

$$\kappa_L(t) = \frac{n^{-L/2}}{L!} \sum_{\ell=0}^{L} {\binom{L}{\ell}} \int_{\mathcal{U}} \langle \nabla^{L-\ell} f(tx), x^{\otimes L-\ell} \rangle \, b_\ell(t,x) e^{-r(t,x)} \gamma(dx),$$

$$\tau_L = -f(0)\gamma(\mathcal{U}^c) - \sum_{k=1}^{L-1} \frac{n^{-k/2}}{k!} \sum_{\ell=0}^k {\binom{k}{\ell}} \int_{\mathcal{U}^c} \langle \nabla^{k-\ell} f(0), x^{\otimes k-\ell} \rangle \, b_\ell(0,x)\gamma(dx).$$
(3.5)

Using this lemma together with (3.2) we arrive at the following decomposition:

$$\frac{e^{nv(0)}}{(2\pi/n)^{d/2}} \int f(x)e^{-nv(x)}dx = I(1/\sqrt{n}) + \tau_{\mathcal{U}^c} = \text{Lead}_L + \kappa_L(t) + \tau_L + \tau_{\mathcal{U}^c},$$
(3.6)

the terms of which are analyzed in the next two sections. To prove Lemma 3.2, define the function

$$I(t) = \int_{\mathcal{U}} F(t, x) e^{-r(t, x)} \gamma(dx), \qquad (3.7)$$

so that the first of the two integrals in (3.2) is indeed given by $I(1/\sqrt{n})$ according to this definition of the function I. We first show that for each $x \in \mathcal{U}$, the functions $F(\cdot, x), e^{-r(\cdot, x)}$ are L times continuously differentiable for all $|t| < 2/\sqrt{n}$, and that

$$\max_{1 \le k \le L} \sup_{|t| \le 2/\sqrt{n}, x \in \mathcal{U}} \left| \partial_t^k e^{-r(t,x)} \right| + \max_{1 \le k \le L} \sup_{|t| \le 2/\sqrt{n}, x \in \mathcal{U}} \left| \partial_t^k F(t,x) \right| < \infty.$$
(3.8)

It then follows that I is L times continuously differentiable in $(-2/\sqrt{n}, 2/\sqrt{n})$, and that to differentiate I for $|t| < 2/\sqrt{n}$, we can bring the t-derivative inside the integral.

The statement about F is clear: F is jointly continuous in t and x over $t \in [-2/\sqrt{n}, 2/\sqrt{n}]$ and $x \in \mathcal{U}$, and the t derivatives of F, given by $\partial_t^k F(t, x) = \langle \nabla^k f(tx), x^{\otimes k} \rangle$, are also jointly continuous in t and x in this region. This follows by Assumption 2.2. Thus the second summand in (3.8) is indeed finite. Next, consider e^{-r} . We use the following formula for the derivatives of the exponential of a function.

Lemma 3.3 (Chapter 2 in [21]). Let $h \in C^k(\mathbb{R})$. Then

$$\frac{d^k}{dt^k}e^{h(t)} = e^{h(t)}B_k\left(h'(t), h''(t), \dots, h^{(k)}(t)\right),$$
(3.9)

where B_k is the kth complete Bell polynomial defined in (2.7).

This follows from Faa di Bruno's formula on the derivatives of $f \circ h$, with $f(x) = e^x$. See [21, 6] for more on this topic. Next, we derive the following formulas for the t-derivatives of r.

Lemma 3.4. Let r be as in (3.1). If $v \in C^{L+2}\left(\frac{2}{\sqrt{n}}\mathcal{U}\right)$ then for each $x \in \mathcal{U}$ the function $r(\cdot, x)$ is L times continuously differentiable for $|t| \leq 2/\sqrt{n}$. Furthermore, we have the following alternative representation for r:

$$r(t,x) = \frac{t}{2} \int_0^1 \langle \nabla^3 v(qtx), x^{\otimes 3} \rangle (1-q)^2 dq \qquad \forall |t| \le 2/\sqrt{n}, \ x \in \mathcal{U}$$
(3.10)

and for $1 \leq k \leq L$ we have

$$\partial_t^k r(t,x) = \int_0^1 \langle \nabla^{k+2} v(qtx), x^{\otimes k+2} \rangle q^k (1-q) dq \qquad \forall |t| \le 2/\sqrt{n}, \ x \in \mathcal{U}.$$
(3.11)

In particular, (3.11) implies

$$\partial_t^k r(0,x) = \frac{\langle \nabla^{k+2} v(0), x^{\otimes k+2} \rangle}{(k+1)(k+2)}, \qquad \forall x \in \mathcal{U}, \ \forall 1 \le k \le L.$$
(3.12)

See Appendix B for the proof. We conclude from the above two lemmas that $e^{-r(\cdot,x)}$ is also L times continuously differentiable in $\left[-2/\sqrt{n}, 2/\sqrt{n}\right]$ for each $x \in \mathcal{U}$, and

$$\partial_t^k e^{-r(t,x)} = e^{-r(t,x)} B_k \left(-\partial_t r(t,x), -\partial_t^2 r(t,x), \dots, -\partial_t^k r(t,x) \right), \qquad \forall x \in \mathcal{U}, \ \forall 1 \le k \le L$$
(3.13)

with $\partial_t^k r(t, x)$ given as in (3.11). Furthermore, we see from (3.10) and (3.11) that r and its t-derivatives are jointly continuous in t and x. We can therefore conclude (3.8). As discussed above, it follows that I is L times continuously differentiable in $(-2/\sqrt{n}, 2/\sqrt{n})$. Hence for some $t \in [0, n^{-1/2}]$ we have

$$I(1/\sqrt{n}) = \sum_{k=0}^{L-1} \frac{n^{-k/2}}{k!} I^{(k)}(0) + \frac{n^{-L/2}}{L!} I^{(L)}(t).$$
(3.14)

To prove Lemma 3.2, it remains to work out the derivatives of I more explicitly using the above results; see Appendix B for the rest of the proof.

3.3 Leading order terms

In this section, we explicitly compute the terms

$$A_k = \frac{1}{k!} \sum_{\ell=0}^k \binom{k}{\ell} \int_{\mathbb{R}^d} \langle \nabla^{k-\ell} f(0), x^{\otimes k-\ell} \rangle \, b_\ell(0, x) \gamma(dx).$$
(3.15)

We then have that $\text{Lead}_L = f(0) + \sum_{k=1}^{L-1} A_k n^{-k/2}$. As noted in the introduction, we will see that these A_k coincide with the coefficients obtained in the finite dimensional Laplace asymptotic expansion given in [14].

We first separate out the case $\ell = 0$ from the rest, and recognize that the integrals in (3.15) are Gaussian expectations. Therefore,

$$A_{k} = \frac{1}{k!} \mathbb{E}\left[\langle \nabla^{k} f(0), Z^{\otimes k} \rangle\right] + \frac{1}{k!} \sum_{\ell=1}^{k} \binom{k}{\ell} \mathbb{E}\left[\langle \nabla^{k-\ell} f(0), Z^{\otimes k-\ell} \rangle b_{\ell}(0, Z)\right]$$
(3.16)

Next, we state a useful alternative representation of the Bell polynomials; see Appendix \mathbf{E} for the proof.

Lemma 3.5. For all $k \ge 1$, the complete Bell polynomial B_k defined in (2.7) has the following equivalent form:

$$B_k(x_1, \dots, x_k) = k! \sum_{r=1}^k \frac{1}{r!} \sum_{\substack{m_1+m_2+\dots+m_r=k \ \ell=1}} \prod_{\ell=1}^r \frac{x_{m_\ell}}{m_\ell!}$$
(3.17)

We use this representation to obtain an explicit formula for $b_{\ell}(0, x)$ appearing in (3.15). We recall that $b_{\ell}(0, x)$ is defined in (3.3) and that the derivatives $\partial_t^j r(0, x)$, $j = 1, \ldots, \ell$ are given by (3.12). Combining these ingredients and using (3.17), we obtain for all $\ell \geq 1$ that

$$b_{\ell}(0,x) = B_{\ell} \left(-\partial_{t} r(0,x), \dots, -\partial_{t}^{\ell} r(0,x) \right)$$

= $B_{\ell} \left(\frac{-\langle \nabla^{3} v(0), x^{\otimes 3} \rangle}{2 \cdot 3}, \dots, \frac{-\langle \nabla^{\ell+2} v(0), x^{\otimes \ell+2} \rangle}{(\ell+1)(\ell+2)} \right)$
= $\ell! \sum_{r=1}^{\ell} \frac{(-1)^{r}}{r!} \sum_{\substack{m_{1}+m_{2}+\dots+m_{r}=\ell \\ m_{1},\dots,m_{r} \ge 1}} \prod_{j=1}^{r} \frac{\langle \nabla^{m_{j}+2} v(0), x^{\otimes m_{j}+2} \rangle}{(m_{j}+2)!}$ (3.18)

We are now in a position to compute the coefficients A_k . First, we introduce some multi-index notation. For a multi-index $\boldsymbol{\alpha} = (\alpha^1, \ldots, \alpha^d)$, with $\alpha^j \ge 0$ for all j, we define $|\boldsymbol{\alpha}| = \alpha^1 + \cdots + \alpha^d$ and $\boldsymbol{\alpha}! = \alpha^1! \ldots \alpha^d!$. Similarly, $\boldsymbol{\alpha}!! = \alpha_1!! \ldots \alpha_d!!$. If $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$ are two multi-indices, then $\boldsymbol{\alpha} + \boldsymbol{\beta}$ is their entrywise sum, and similarly for differences. We let $\mathbf{1} = (1, \ldots, 1)$ be the multi-index of all ones. If $x \in \mathbb{R}^d$ then $x^{\boldsymbol{\alpha}} = x_1^{\alpha^1} \ldots x_d^{\alpha^d}$. We let $\partial^{\boldsymbol{\alpha}} f(x) = \partial_{x_1}^{\alpha^1} \ldots \partial_{x_d}^{\alpha^d} f(x)$ whenever this partial derivative exists.

Theorem 3.6. For all $k \ge 1$, it holds

$$A_{k} = \sum_{|\boldsymbol{\beta}|=k} \frac{(\boldsymbol{\beta}-\mathbf{1})!!}{\boldsymbol{\beta}!} \partial^{\boldsymbol{\beta}} f(0) \operatorname{even}(\boldsymbol{\beta}) + \sum_{\ell=1}^{k} \sum_{r=1}^{\ell} \frac{(-1)^{r}}{r!} \sum_{\substack{m_{1}+m_{2}+\dots+m_{r}=\ell\\m_{1},\dots,m_{r}\geq1}} \sum_{|\boldsymbol{\beta}|=k-\ell, |\boldsymbol{\alpha}_{1}|=m_{1}+2,\dots,|\boldsymbol{\alpha}_{r}|=m_{r}+2} \operatorname{even}(\boldsymbol{\beta}+\boldsymbol{\alpha}_{1}+\dots+\boldsymbol{\alpha}_{r}) \frac{(\boldsymbol{\beta}+\boldsymbol{\alpha}_{1}+\dots+\boldsymbol{\alpha}_{r}-\mathbf{1})!!}{\boldsymbol{\beta}!\boldsymbol{\alpha}_{1}!\dots\boldsymbol{\alpha}_{r}!} \partial^{\boldsymbol{\beta}} f(0) \partial^{\boldsymbol{\alpha}_{1}} v(0)\dots\partial^{\boldsymbol{\alpha}_{r}} v(0),$$
(3.19)

where 0!! = 0! = 1 and $even(\alpha) = 1$ if α_i is even for each i = 1, ..., d, and $even(\alpha) = 0$ otherwise.

This coincides with the formula in Theorem 1.1 of the arXiv version of [14], and Theorem 1.2 of the published version of this work, and we have intentionally used the same notation. The only difference is

that we have factored out the $(2\pi)^{d/2}/\sqrt{\det H}$, and that we have separated out the $\ell = 0$ case into the first line of (3.19). In particular, note that if $|\boldsymbol{\alpha}|$ is odd, then all the indices α_i cannot be even, and hence even $(\boldsymbol{\alpha}) = 0$. But note that $|\boldsymbol{\beta} + \boldsymbol{\alpha}_1 + \cdots + \boldsymbol{\alpha}_r| = (k-\ell) + (m_1+2) + \cdots + (m_r+2) = k-\ell+\ell+2r = k+2r$ is odd if k is odd. Therefore the sum in the second and third line is zero when k is odd, and by the same reasoning, the sum in the first line is also zero when k is odd. We conclude that $A_k = 0$ for all odd k.

Corollary 3.7. When f = 1, the A_k are given as follows for all $k \ge 1$:

$$A_{k} = \sum_{r=1}^{\kappa} \frac{(-1)^{r}}{r!} \sum_{\substack{m_{1}+m_{2}+\dots+m_{r}=k \ |\boldsymbol{\alpha}_{1}|=m_{1}+2,\dots,|\boldsymbol{\alpha}_{r}|=m_{r}+2\\m_{1},\dots,m_{r}\geq 1}} \sum_{\substack{(\boldsymbol{\alpha}_{1}+\dots+\boldsymbol{\alpha}_{r}-1)!!\\\boldsymbol{\alpha}_{1}!\dots\boldsymbol{\alpha}_{r}!}} (3.20)$$

3.4 Remainder terms

Recall that the remainder is given by $\kappa_L(t) + \tau_L + \tau_{\mathcal{U}^c}$, where $\tau_{\mathcal{U}^c}$ is the second integral in (3.2), while $\kappa_L(t)$ and τ_L are as in (3.5). In this section, we bound each of these three terms. Since $\kappa_L(t)$ depends on some $t \in [0, 1/\sqrt{n}]$, we will bound it uniformly over t.

Theorem 3.8 (Bound on $\kappa_L, \tau_L, \tau_{\mathcal{U}^c}$). Under the assumptions of Theorem 2.8, it holds

$$\sup_{t \in [0, 1/\sqrt{n}]} \kappa_L(t) = \kappa_L \lesssim_L e^{\bar{\epsilon}_1(R)^2 + \epsilon_2(R)} \sum_{\ell=0}^L c_{L-\ell,f}(R) \sqrt{d/n}^{L-\ell} B_\ell(\delta_1(R), \dots, \delta_\ell(R)),$$

$$|\tau_{\mathcal{U}^c}| \le (|f(0)| + b_f) de^{-Ra_v d/8},$$

$$|\tau_L| \lesssim_L \max_{0 \le m \le L-1} \|\nabla^m f(0)\| \sqrt{d/n}^m \max_{1 \le j \le L-1} (\epsilon_j(0) \lor 1)^L e^{-(R-1)^2 d/4}.$$
(3.21)

Proof. Let $h_k(t, x) = \langle \nabla^k f(tx), x^{\otimes k} \rangle$. We first apply generalized Hölder's inequality:

$$|\kappa_L(t)| \le \sup_{t \in [0, 1/\sqrt{n}]} \|e^{-r(t, \cdot)} \mathbb{1}_{\mathcal{U}}\|_4 \sum_{\ell=0}^L n^{-\ell/2} \|b_\ell(t, \cdot) \mathbb{1}_{\mathcal{U}}\|_2 n^{-(L-\ell)/2} \|h_{L-\ell}(t, \cdot) \mathbb{1}_{\mathcal{U}}\|_4.$$
(3.22)

Note that we have passed the $n^{-L/2}$ inside the summation. We know by Lemma C.2 that

$$\sup_{t \in [0, 1/\sqrt{n}]} \|e^{-r(t, \cdot)} \mathbb{1}_{\mathcal{U}}\|_4 \le \exp(\bar{\epsilon}_1(R)^2 + \epsilon_2(R)).$$
(3.23)

Next, a straightforward Gaussian integral calculation gives

$$n^{-k/2} \sup_{t \in [0, 1/\sqrt{n}]} \|h_k(t, \cdot) \mathbb{1}_{\mathcal{U}}\|_4 \lesssim_k c_{k, f}(R) (d/n)^{k/2}.$$
(3.24)

Finally, Corollary C.5 gives

$$n^{-\ell/2} \|b_{\ell}(t,\cdot) \mathbb{1}_{\mathcal{U}}\|_{2} = n^{-\ell/2} \|b_{\ell}(-\partial_{t}r(t,\cdot),\ldots,-\partial_{t}^{\ell}r(t,\cdot)) \mathbb{1}_{\mathcal{U}}\|_{2}$$

$$\lesssim_{\ell} B_{\ell}(\delta_{1}(R),\ldots,\delta_{\ell}(R)), \quad \forall 0 \le t \le 1/\sqrt{n}.$$
(3.25)

Substituting (3.25), (3.24), and (3.23) into (3.22) concludes the proof of the bound on $\kappa_L(t)$. Next, consider $\tau_{\mathcal{U}^c}$. Note that Assumption 2.3 and the definition (3.1) of W implies that $W(1/\sqrt{n}, x) \geq a_v \sqrt{d} ||x||$ for all $||x|| \geq \sqrt{d}$ and therefore in particular, for all $x \in \mathcal{U}^c$. Moreover, Assumption 2.3 on f implies that $|F(1/\sqrt{n}, x)| \leq |f(0)| + b_f \exp(a_v \sqrt{d} ||x||/2)$ for all $x \in \mathbb{R}^d$. Thus

$$\begin{aligned} |\tau_{\mathcal{U}^c}| &\leq (2\pi)^{-d/2} \int_{\mathcal{U}^c} |F(1/\sqrt{n}, x)| e^{-W(1/\sqrt{n}, x)} dx \\ &\leq (|f(0)| + b_f) \ (2\pi)^{-d/2} \int_{\|x\| \geq R\sqrt{d}} e^{-a_v \sqrt{d} \|x\|/2} dx. \end{aligned}$$

The bound on $\tau_{\mathcal{U}^c}$ now follows from Lemma E.2. For the bound on τ_L , we have the following Gaussian tail bound

$$\gamma\left(\left\{x \in \mathbb{R}^d : \|x\| \ge R\sqrt{d}\right\}\right) \le \exp\left(-(R-1)^2 d/2\right), \quad \forall R \ge 1,$$
(3.26)

which follows e.g. from Example 2.12 of [5]. Also, recall the definition of $h_k(t, x)$ from the beginning of the proof. Using the generalized Hölder's inequality, we have

$$|\tau_L| \lesssim_L |f(0)|\gamma(\mathcal{U}^c) + \sum_{k=1}^{L-1} \sum_{\ell=0}^k n^{-\ell/2} \|b_\ell(0,\cdot)\|_4 n^{-(k-\ell)/2} \|h_{k-\ell}(0,\cdot)\|_4 \gamma(\mathcal{U}^c)^{1/2}.$$
 (3.27)

Now, a standard Gaussian calculation gives

$$n^{-k/2} \|h_k(0,\cdot)\|_q \lesssim_{k,q} \|\nabla^k f(0)\| \sqrt{d/n^k}.$$

Next, Corollary C.5 gives that

$$n^{-\ell/2} \|b_{\ell}(0,\cdot)\|_{4} = n^{-\ell/2} \|b_{\ell}(-\partial_{t}r(0,\cdot),\ldots,-\partial_{t}^{\ell}r(0,\cdot))\|_{4} \lesssim_{\ell} \left(1 \vee \max_{1 \le j \le \ell} \epsilon_{j}(0)\right)^{\ell}.$$

Substituting the above two bounds into (3.27) gives

$$\begin{aligned} |\tau_L| \lesssim_L \gamma(\mathcal{U}^c)^{1/2} \left(|f(0)| + \sum_{k=1}^{L-1} \sum_{\ell=0}^k \left(1 \lor \max_{1 \le j \le \ell} \epsilon_j(0) \right)^\ell \|\nabla^{k-\ell} f(0)\| \sqrt{d/n}^{k-\ell} \right) \\ \lesssim_L \max_{0 \le m \le L-1} \|\nabla^m f(0)\| \sqrt{d/n}^m \max_{1 \le j \le L-1} (\epsilon_j(0) \lor 1)^L e^{-(R-1)^2 d/4}, \end{aligned}$$

as desired.

A Comparison to [23]

The correction term ϵ_0 from (2) in [23] (which is the same as ϵ_0 from display (9), with g now playing the role of ng) is the same as our A_2n^{-1} in the Laplace expansion of the partition function,

$$\frac{n^{d/2}}{(2\pi)^{d/2}} \int e^{-nv(x)} dx = 1 + A_2 n^{-1} + \operatorname{Rem}_4.$$

Recall from (2.21) that A_2 is given by

$$A_2 = \frac{1}{12} \|\nabla^3 v(0)\|_F^2 + \frac{1}{8} \sum_{i=1}^d \left(\partial_i \Delta v(0)\right)^2 - \frac{1}{8} \sum_{i,j=1}^d \partial_i^2 \partial_j^2 v(0).$$
(A.1)

To see that A_2n^{-1} is the same as ϵ_0 from (2) of [23], note that the terms \hat{g}^{ij} are defined as $\hat{g}^{ij} = [(\partial^2 v(0))^{-1}]_{ij} = \delta_{ij}$ under our assumptions that $\nabla^2 v(0) = I_d$. The authors argue in Section 6 on GLMs, that $\epsilon_0 = A_2/n$ scales as d^3 . The argument for this is the fact that ϵ_0 involves a sum over d^3 terms of order n^{-1} . However, we claim that

$$|A_2| \le d^2 \left(\|\nabla^3 v(0)\|^2 + \|\nabla^4 v(0)\| \right),$$

and therefore, $|A_2|$ scales as d^2 if $\|\nabla^3 v(0)\|$, $\|\nabla^4 v(0)\| = \mathcal{O}(1)$. For example, it was shown in [13] that $\|\nabla^3 v(0)\|$, $\|\nabla^4 v(0)\| = \mathcal{O}(1)$ for a logistic regression model with Gaussian design, which is the prototypical example of a GLM.

The third term of (A.1) is clearly seen to be bounded by $d^2 \|\nabla^4 v(0)\|$. To bound the first term, let $T = \nabla^3 v(0)$ and T_i be the matrix whose jkth entry is T_{ijk} . Note that

$$||T_i|| = \sup_{||u|| = ||v|| = 1} \langle T_i, u \otimes v \rangle = \sup_{||u|| = ||v|| = 1} \langle T, u \otimes v \otimes e_i \rangle \le ||T||$$

for all i. Therefore,

$$||T||_F^2 = \sum_{i=1}^d \operatorname{Tr}(T_i^{\mathsf{T}}T_i) \le \sum_{i=1}^d d||T_i^{\mathsf{T}}T_i|| \le d^2 \max_{1 \le i \le d} ||T_i||^2 \le d^2 ||T||^2.$$
(A.2)

Finally, note that $\sum_{i=1}^{d} (\partial_i \Delta v(0))^2 = ||\langle T, I_d \rangle||^2$, where $\langle T, I_d \rangle_i = \sum_{j=1}^{d} T_{ijj}$. Now, we have

$$\begin{split} \|\langle T, I_d \rangle\| &= \sup_{\|u\|=1} \langle T, I_d \rangle^{\intercal} u = \sup_{\|u\|=1} \langle T, I_d \otimes u \rangle \\ &\leq \sup_{\|u\|=1} \sum_{j=1}^d |\langle T, e_j \otimes e_j \otimes u \rangle| \leq d \|T\| \end{split}$$

and hence

$$\sum_{i=1}^{d} (\partial_i \Delta v(0))^2 = \|\langle T, I_d \rangle\|^2 \le d^2 \|T\|^2.$$

B Proofs from Section 3

Proof of Lemma 3.1. First we change variables as $x = n^{-1/2}y$, and recognize that $f(y/\sqrt{n}) = F(1/\sqrt{n}, y)$, while $nv(y/\sqrt{n}) = nv(0) + W(1/\sqrt{n}, y)$. This gives

$$\int f(x)e^{-nv(x)}dx = e^{-nv(0)}n^{-d/2}\int F(1/\sqrt{n},y)e^{-W(1/\sqrt{n},y)}dy.$$
(B.1)

We now divide through by $e^{-nv(0)}n^{-d/2}(2\pi)^{d/2}$, and recognize that

$$(2\pi)^{-d/2}e^{-W(1/\sqrt{n},y)}dy = e^{-r(1/\sqrt{n},y)}\gamma(dy).$$

This gives

$$\frac{e^{nv(0)}n^{d/2}}{(2\pi)^{d/2}} \int f(x)e^{-nv(x)}dx = (2\pi)^{-d/2} \int F(1/\sqrt{n}, y)e^{-W(1/\sqrt{n}, y)}dy$$
$$= \int_{\mathcal{U}} F(1/\sqrt{n}, y)e^{-r(1/\sqrt{n}, y)}\gamma(dy) + (2\pi)^{-d/2} \int_{\mathcal{U}^c} F(1/\sqrt{n}, y)e^{-W(1/\sqrt{n}, y)}dy,$$

as desired.

Proof of Lemma 3.4. A Taylor expansion of v gives

$$v(tx) = v(0) + \frac{t^2 ||x||^2}{2} + \frac{t^3}{2!} \int_0^1 \langle \nabla^3 v(qtx), x^{\otimes 3} \rangle (1-q)^2 dq.$$

Hence for $t \neq 0$, we have

$$r(t,x) = \frac{v(tx) - v(0)}{t^2} - \frac{1}{2} ||x||^2 = \frac{t}{2!} \int_0^1 \langle \nabla^3 v(qtx), x^{\otimes 3} \rangle (1-q)^2 dq$$
(B.2)

Recall that $r(0, x) = W(0, x) - ||x||^2/2 = 0$ by definition, which coincides with the righthand side of (B.2) if we plug in t = 0. Hence r is given by (B.2) for all $|t| < 2/\sqrt{n}$ and $x \in \mathcal{U}$. Since v is at least C^3 on $(2/\sqrt{n})\mathcal{U}$, we see from this representation that $r(\cdot, x)$ is continuous in t in the range $|t| < 2/\sqrt{n}$ for each fixed $x \in \mathcal{U}$. We now show $r(\cdot, x) \in C^1((-2/\sqrt{n}, 2/\sqrt{n}))$ for all $x \in \mathcal{U}$. First, note that the t-derivative exists at zero, since

$$\frac{r(t,x) - r(0,x)}{t} = \frac{1}{2} \int_0^1 \langle \nabla^3 v(qtx), x^{\otimes 3} \rangle (1-q)^2 dq \to \frac{1}{3 \cdot 2} \langle \nabla^3 v(0), x^{\otimes 3} \rangle \tag{B.3}$$

as $t \to 0$, by continuity of $\nabla^3 v$. Next, suppose $t \neq 0$. Note that by the change of variables q = s/t, we can rewrite r in the equivalent form

$$r(t,x) = \frac{t^{-2}}{2} \int_0^t \langle \nabla^3 v(sx), x^{\otimes 3} \rangle (t-s)^2 ds.$$
 (B.4)

This function is infinitely differentiable in t for all $0 < |t| < 2/\sqrt{n}$ and $x \in \mathcal{U}$. We compute for nonzero t that

$$\partial_t r(t,x) = -t^{-3} \int_0^t \langle \nabla^3 v(sx), x^{\otimes 3} \rangle (t-s)^2 ds + t^{-2} \int_0^t \langle \nabla^3 v(sx), x^{\otimes 3} \rangle (t-s) ds$$

$$= -\int_0^1 \langle \nabla^3 v(tqx), x^{\otimes 3} \rangle (1-q)^2 dq + \int_0^1 \langle \nabla^3 v(tqx), x^{\otimes 3} \rangle (1-q) dq \qquad (B.5)$$

$$= \int_0^1 \langle \nabla^3 v(tqx), x^{\otimes 3} \rangle q (1-q) dq$$

This function is continuous in t for $x \in \mathcal{U}$, and the limit as $t \to 0$ coincides with (B.3). Hence $\partial^t r(t, x)$ exists everywhere and is continuous, so we conclude that $r(\cdot, x) \in C^1((-2/\sqrt{n}, 2/\sqrt{n}))$. But from the formula (B.5) for $\partial_t r$ we now see that $\partial_t r(\cdot, x)$ is L - 1 times continuously differentiable, and we obtain the desired formula (3.11) for $\partial_t^k r$.

Proof of Lemma 3.2. We first compute $I^{(k)}(t)$. To do so, we use the definition (3.7) of I(t), the product rule of differentiation, the formula (3.13) for the t derivatives of e^{-r} , and the fact that $\partial_t^k F(t,x) = \langle \nabla^k f(tx), x^{\otimes k} \rangle$. We also use the fact that the t derivatives can be brought inside the integral, by (3.8). We get the following expression:

$$I^{(k)}(t) = \int_{\mathcal{U}} \sum_{\ell=0}^{k} \binom{k}{\ell} \partial_{t}^{k-\ell} F(t,x) \partial_{t}^{k} e^{-r(t,x)} \gamma(dx)$$

$$= \sum_{\ell=0}^{k} \binom{k}{\ell} \int_{\mathcal{U}} \langle \nabla^{k-\ell} f(tx), x^{\otimes k-\ell} \rangle b_{\ell}(t,x) e^{-r(t,x)} \gamma(dx),$$
(B.6)

where b_{ℓ} is as in (3.3). Plugging this into the Taylor expansion (3.14) gives

$$I(1/\sqrt{n}) = f(0)\gamma(\mathcal{U}) + \sum_{k=1}^{L-1} \sum_{\ell=0}^{k} \binom{k}{\ell} \int_{\mathcal{U}} \langle \nabla^{k-\ell} f(0), x^{\otimes k-\ell} \rangle b_{\ell}(0, x)\gamma(dx)$$

$$+ \sum_{\ell=0}^{L} \binom{L}{\ell} \int_{\mathcal{U}} \langle \nabla^{L-\ell} f(tx), x^{\otimes L-\ell} \rangle b_{\ell}(t, x) e^{-r(t, x)} \gamma(dx).$$
(B.7)

Finally, we write $\gamma(\mathcal{U})$ as $1 - \gamma(\mathcal{U}^c)$ and similarly, we write each integral in the first line as $\int_{\mathcal{U}} = \int_{\mathbb{R}^d} - \int_{\mathcal{U}^c}$. *Proof of Theorem 3.6.* We use the formula (3.18) for $b_k(0, x)$ and recall the formula (3.16):

$$A_{k} = \frac{1}{k!} \mathbb{E}\left[\langle \nabla^{k} f(0), Z^{\otimes k} \rangle\right] + \frac{1}{k!} \sum_{\ell=1}^{k} \binom{k}{\ell} \mathbb{E}\left[\langle \nabla^{k-\ell} f(0), Z^{\otimes k-\ell} \rangle b_{\ell}(0, Z)\right]$$
(B.8)

Therefore

$$A_{k} = \frac{1}{k!} \mathbb{E} \left[\langle \nabla^{k} f(0), Z^{\otimes k} \rangle \right] + \sum_{\ell=1}^{k} \sum_{r=1}^{\ell} \frac{(-1)^{r}}{r!} \sum_{\substack{m_{1}+m_{2}+\dots+m_{r}=\ell\\m_{1},\dots,m_{r}\geq 1}} \mathbb{E} \left[\frac{\langle \nabla^{k-\ell} f(0), Z^{\otimes k-\ell} \rangle}{(k-\ell)!} \prod_{j=1}^{r} \frac{\langle \nabla^{m_{j}+2} v(0), Z^{\otimes m_{j}+2} \rangle}{(m_{j}+2)!} \right]$$
(B.9)

For the next part of the proof, recall the notation on multi-indices $\boldsymbol{\alpha}$ introduced before the theorem statement. Consider the inner product $\langle T, S \rangle$ for two symmetric tensors T, S of order m; that is, $T = (T_{i_1 i_2 \dots i_m})_{i_1,\dots,i_m=1}^d$ and similarly for S. Consider the set of multi-indices $\boldsymbol{\alpha} = (\alpha^1, \dots, \alpha^d)$ such that $\alpha^j \geq 0, j = 1, \dots, d$ and $|\boldsymbol{\alpha}| := \sum_{j=1}^d \alpha^j = m$. We define the map $\boldsymbol{i} = (i_1, \dots, i_m) \mapsto \boldsymbol{\alpha} = (\alpha^1, \dots, \alpha^d)$ according to $\alpha^j(\boldsymbol{i}) = \sum_{k=1}^m \delta_{i_k,j}$. In other words, $\alpha^j(\boldsymbol{i}), j = 1, \dots, d$ is the number of times \boldsymbol{j} appears among the indices i_1, \dots, i_m . Note that to each $\boldsymbol{\alpha}$ correspond $m!/\boldsymbol{\alpha}!$ different \boldsymbol{i} 's such that $\boldsymbol{\alpha}(\boldsymbol{i}) = \boldsymbol{\alpha}$. Finally, since T is symmetric we have $T_{\boldsymbol{i}} = T_{\boldsymbol{i}'}$ for any $\boldsymbol{i}, \boldsymbol{i}'$ such that $\boldsymbol{\alpha}(\boldsymbol{i}) = \boldsymbol{\alpha}(\boldsymbol{i}')$. We therefore define $T^{\boldsymbol{\alpha}} := T_{\boldsymbol{i}}$ for any \boldsymbol{i} such that $\boldsymbol{\alpha}(\boldsymbol{i}) = \boldsymbol{\alpha}$, and similarly for S. These considerations imply that the inner product $\langle T, S \rangle$ can be written as

$$\langle T, S \rangle = \sum_{i_1, \dots, i_m = 1}^d T_{i_1 \dots i_m} S_{i_1 \dots i_m} = \sum_{|\boldsymbol{\alpha}| = m} \frac{m!}{\boldsymbol{\alpha}!} T^{\boldsymbol{\alpha}} S^{\boldsymbol{\alpha}}.$$

Furthermore, if $T = \nabla^m v(0)$ and $S = Z^{\otimes m}$, then $T^{\boldsymbol{\alpha}} = \partial^{\boldsymbol{\alpha}} v(0)$ and $S^{\boldsymbol{\alpha}} = Z^{\boldsymbol{\alpha}}$. We therefore have

$$\frac{\langle \nabla^{m_{\ell}+2}v(0), Z^{\otimes m_{\ell}+2} \rangle}{(m_{\ell}+2)!} = \frac{1}{(m_{\ell}+2)!} \sum_{|\alpha|=m_{\ell}+2} \frac{(m_{\ell}+2)!}{\alpha!} \partial^{\alpha}v(0) Z^{\alpha} = \sum_{|\alpha|=m_{\ell}+2} \frac{1}{\alpha!} \partial^{\alpha}v(0) Z^{\alpha},$$

and similarly

$$\frac{\langle \nabla^{k-\ell} f(0), Z^{\otimes k-\ell} \rangle}{(k-\ell)!} = \sum_{|\boldsymbol{\beta}|=k-\ell} \frac{1}{\boldsymbol{\beta}!} \partial^{\boldsymbol{\beta}} f(0) Z^{\boldsymbol{\beta}},$$

so that

$$\frac{\langle \nabla^{k-\ell} f(0), Z^{\otimes k-\ell} \rangle}{(k-\ell)!} \prod_{\ell=1}^{r} \frac{\langle \nabla^{m_{\ell}+2} v(0), Z^{\otimes m_{\ell}+2} \rangle}{(m_{\ell}+2)!} = \sum_{|\boldsymbol{\beta}|=k-\ell, |\boldsymbol{\alpha}_{1}|=m_{1}+2, \dots, |\boldsymbol{\alpha}_{r}|=m_{r}+2} \frac{1}{\boldsymbol{\beta}! \boldsymbol{\alpha}_{1}! \dots \boldsymbol{\alpha}_{r}!} \partial^{\boldsymbol{\beta}} f(0) \partial^{\boldsymbol{\alpha}_{1}} v(0) \dots \partial^{\boldsymbol{\alpha}_{r}} v(0) Z^{\boldsymbol{\beta}+\boldsymbol{\alpha}_{1}+\dots+\boldsymbol{\alpha}_{r}}.$$
(B.10)

Finally, note that

$$\mathbb{E}\left[Z^{\boldsymbol{\beta}+\boldsymbol{\alpha}_1+\cdots+\boldsymbol{\alpha}_r}\right] = \prod_{i=1}^d \mathbb{E}\left[Z_i^{\beta^i+\alpha_1^i+\cdots+\alpha_r^i}\right] = \begin{cases} (\boldsymbol{\beta}+\boldsymbol{\alpha}_1+\cdots+\boldsymbol{\alpha}_r-\mathbf{1})!!, & \beta^i+\alpha_1^i+\cdots+\alpha_r^i \text{ even for all } i, \\ 0, & \text{otherwise.} \end{cases}$$

Thus recalling the definition of even(α) from the statement of the lemma, we conclude that

$$\mathbb{E}\left[Z^{\beta+\alpha_1+\cdots+\alpha_r}\right] = (\beta+\alpha_1+\cdots+\alpha_r-1)!!\operatorname{even}(\beta+\alpha_1+\cdots+\alpha_r).$$
(B.11)

Using (B.10) and (B.11), we now have

$$\mathbb{E}\left[\frac{\langle \nabla^{k-\ell}f(0), Z^{\otimes k-\ell} \rangle}{(k-\ell)!} \prod_{\ell=1}^{r} \frac{\langle \nabla^{m_{\ell}+2}v(0), Z^{\otimes m_{\ell}+2} \rangle}{(m_{\ell}+2)!}\right]$$

=
$$\sum_{|\beta|=k-\ell, |\alpha_{1}|=m_{1}+2, \dots, |\alpha_{r}|=m_{r}+2} \frac{(\beta+\alpha_{1}+\dots+\alpha_{r}-1)!!}{\beta!\alpha_{1}!\dots\alpha_{r}!} \partial^{\beta}f(0)\partial^{\alpha_{1}}v(0)\dots\partial^{\alpha_{r}}v(0) \operatorname{even}(\beta+\alpha_{1}+\dots+\alpha_{r})$$

By a similar argument,

$$\frac{1}{k!} \mathbb{E}\left[\langle \nabla^k f(0), Z^{\otimes k} \rangle \right] = \sum_{|\boldsymbol{\beta}| = k} \frac{(\boldsymbol{\beta} - \mathbf{1})!!}{\boldsymbol{\beta}!} \partial^{\boldsymbol{\beta}} f(0) \text{even}(\boldsymbol{\beta}).$$

Substituting these expressions into (B.9) concludes the proof.

C L^q norms of exp(-r), $\partial_t^k r$'s, p_k 's, and b_k 's

Lemma C.1. Let r be given as in (3.1), and let $\gamma_{\mathcal{U}}$ be the probability measure such that $\gamma_{\mathcal{U}}(dx) \propto \mathbb{1}_{\mathcal{U}}(x)\gamma(dx)$. Then

$$\sup_{t \in [0, 1/\sqrt{n}]} \left| \int r(t, x) \gamma_{\mathcal{U}}(dx) \right| \le \epsilon_2(R)/4.$$
(C.1)

Proof. A Taylor expansion of r about x = 0 gives

$$r(t,x) = \frac{t}{3!} \langle \nabla^3 v(0), x^{\otimes 3} \rangle + \frac{t^2}{4!} \langle \nabla^4 v(\xi tx), x^{\otimes 4} \rangle$$

for some $\xi \in [0, 1]$. Since the first summand is an odd function of x and U is a symmetric set about zero, we conclude that

$$\left| \int_{\mathcal{U}} r(t,x)\gamma(dx) \right| = \frac{t^2}{4!} \left| \int_{\mathcal{U}} \langle \nabla^4 v(\xi tx), x^{\otimes 4} \rangle \gamma(dx) \right| \le \frac{t^2}{4!} c_4(R) \mathbb{E} \left[\|Z\|^4 \right]$$

$$\le \frac{t^2 d^2}{8} c_4(R) \le c_4(R) \frac{d^2}{8n} = \epsilon_2(R)/8 \quad \forall t \in [0, 1/\sqrt{n}].$$
(C.2)

Finally, recall that we have assumed $R \geq \frac{8}{a_v} \log \frac{2e}{a_v} \geq 8$, since $a_v \leq 1$. Therefore a standard Gaussian tail bound gives $\gamma(\mathcal{U}) \geq 1/2$. Therefore dividing both sides of (C.2) by 1/2 concludes the proof.

Lemma C.2. Let r be given as in (3.1). Then

$$\sup_{t \in [0, 1/\sqrt{n}]} \|e^{-r(t, \cdot)} \mathbb{1}_{\mathcal{U}}\|_p \le \exp\left(p\bar{\epsilon}_1(R)^2/8 + \epsilon_2(R)/4\right).$$
(C.3)

Proof. Let $M = \sup_{t \in [0, 1/\sqrt{n}]} \sup_{x \in \mathcal{U}} \|\nabla_x r(t, x)\|$, which we will quantify below. Then (D.3) of Corollary D.3 gives

$$\|e^{-r(t,\cdot)}\mathbb{1}_{\mathcal{U}}\|_{p} \le \exp\left(\left|\int r(t,x)\gamma_{\mathcal{U}}(dx)\right| + M^{2}p/2\right)$$
(C.4)

We have already bounded $|\int r(t,x)\gamma_{\mathcal{U}}(dx)|$ in the previous lemma, so it remains to bound M. First recall that $r(t,x) = W(tx) - ||x||^2/2 = (v(tx) - v(0))/t^2 - ||x||^2/2$, so that $\nabla_x r(t,x) = \nabla v(tx)/t - x$. A Taylor expansion of ∇v about x = 0 now gives

$$\nabla_x r(t,x) = \frac{t}{2} \langle \nabla^3 v(\xi tx), x^{\otimes 2} \rangle$$

for some ξ between 0 and 1. Hence

$$\sup_{t \in [0, 1/\sqrt{n}]} \sup_{\|x\| \le R\sqrt{d}} \|\nabla_x r(t, x)\| \le \frac{R^2 d}{2\sqrt{n}} c_3(R) \le \frac{1}{2} \bar{\epsilon}_1(R) =: M.$$
(C.5)

Substituting the bound on M and on $\left|\int r(t, x)\gamma_{\mathcal{U}}(dx)\right|$ from Lemma C.1 into (C.4) gives the desired bound (C.3).

Lemma C.3. Let $p_k(x) = \langle \nabla^{k+2}v(0), x^{\otimes k+2} \rangle$. Define $\hat{p}_k = p_k - \int p_k d\gamma_{\mathcal{U}}$, where $\gamma_{\mathcal{U}}$ is the probability density on \mathbb{R}^d proportional to $\gamma \mathbb{1}_{\mathcal{U}}$. If $R \geq 2$, then for all $1 \leq k \leq L$ it holds

$$n^{-k/2} \|p_k\|_q \lesssim_{k,q} \begin{cases} \epsilon_k(0), & k \text{ even}, \\ \epsilon_k(0) \wedge \bar{\epsilon}_k(0) = \bar{\epsilon}_k(0), & k \text{ odd.} \end{cases},$$
(C.6)

$$a^{-k/2} \|\hat{p}_k\|_q \lesssim_{k,q} \bar{\epsilon}_k(0),$$
 (C.7)

where $\epsilon_k, \bar{\epsilon}_k$ are as in (2.2).

r

Lemma C.4. If $R \ge 2$ and L is even, then for all $1 \le k \le L$ and for all $0 \le t \le 1/\sqrt{n}$ it holds

$$n^{-k/2} \|\partial_t^k r(t, \cdot) \mathbb{1}_{\mathcal{U}}\|_q \tag{C.8}$$

$$\lesssim_{k,q} \delta_k(R) = \begin{cases} \epsilon_k(R), & k \text{ even,} \\ \min(\epsilon_k(R), \ \bar{\epsilon}_k(0) + \epsilon_{k+1}(R)), & k \text{ odd.} \end{cases}$$
(C.9)

Corollary C.5. We have

$$n^{-k/2} \left\| B_k \left(-\partial_t r(0, \cdot), \dots, -\partial_t^k r(0, \cdot) \right) \right\|_q \lesssim_{k,q} B_k(\epsilon_1(0), \dots, \epsilon_k(0)),$$
(C.10)

$$\sup_{0 \le t \le 1/\sqrt{n}} n^{-k/2} \left\| B_k \left(-\partial_t r(t, \cdot), \dots, -\partial_t^k r(t, \cdot) \right) \mathbb{1}_{\mathcal{U}} \right\|_q \lesssim_{k,q} B_k(\delta_1(R), \dots, \delta_k(R)).$$
(C.11)

The bound (C.11) requires L to be even. For each bound, we have the further upper bound

$$B_k(t_1,\ldots,t_k) \lesssim_k \max_{1 \le i \le k} (t_i \lor 1)^k.$$

Proof. Recall that $\partial_t^k r(0, \cdot) = p_k/(k+1)(k+2)$, where p_k is as in Lemma C.3. We can now apply Lemma C.3 and Lemma E.1 to prove the first bound. Similarly, we apply Lemma C.4 and E.1 to prove the second bound. The further upper bound also stems from Lemma E.1.

Proof of Lemma C.3. We have

$$\|p_k\|_q = \mathbb{E}\left[|\langle \nabla^{k+2}v(0), Z^{\otimes k+2}\rangle|^q\right]^{1/q} \lesssim_{k,q} d^{(k+2)/2} \|\nabla^{k+2}v(0)\| = c_{k+2}(0)d^{(k+2)/2}.$$
 (C.12)

Therefore,

$$n^{-k/2} \|p_k\|_q \lesssim_{k,q} c_{k+2}(0) d^{(k+2)/2} n^{-k/2} = \epsilon_k(0).$$

This holds for all k. But if k is odd, then $\hat{p}_k = p_k$, and therefore the bound $\|\hat{p}_k\|_q \leq \bar{\epsilon}_k(0)$ we are about to prove also applies to $\|p_k\|_q$. Thus we obtain $\|p_k\|_q \leq \epsilon_k(0) \wedge \bar{\epsilon}_k(0)$ for odd k.

Next, consider \hat{p}_k . We introduce another set $\mathcal{V} = \{x \in \mathbb{R}^d : ||x|| \leq S\sqrt{d}\}$, for some S > 1 that is sufficiently large to ensure $\gamma(\mathcal{V}) \geq 1/2$. We will optimized over all S in this range. We now have

$$\|\hat{p}_{k}\|_{q} = \left\|p_{k} - \int p_{k} d\gamma_{\mathcal{U}}\right\|_{q} \leq \left\|p_{k} - \int p_{k} d\gamma_{\mathcal{V}}\right\|_{q} + \left\|\int p_{k} d\gamma_{\mathcal{U}} - \int p_{k} d\gamma_{\mathcal{V}}\right\|$$

$$\leq \left\|\left(p_{k} - \int p_{k} d\gamma_{\mathcal{V}}\right) \mathbb{1}_{\mathcal{V}}\right\|_{q} + \left\|\left(p_{k} - \int p_{k} d\gamma_{\mathcal{V}}\right) \mathbb{1}_{\mathcal{V}^{c}}\right\|_{q} + \left\|\int p_{k} d\gamma_{\mathcal{U}} - \int p_{k} d\gamma_{\mathcal{V}}\right\|.$$
(C.13)

Next, note that

$$\left| \int p_{k} d\gamma_{\mathcal{U}} - \int p_{k} d\gamma_{\mathcal{V}} \right| = \left| \gamma(\mathcal{U})^{-1} \left(\int p_{k} d\gamma - \int_{\mathcal{U}^{c}} p_{k} d\gamma \right) - \gamma(\mathcal{V})^{-1} \left(\int p_{k} d\gamma - \int_{\mathcal{V}^{c}} p_{k} d\gamma \right) \right|$$

$$\lesssim \left| \gamma(\mathcal{U})^{-1} - \gamma(\mathcal{V})^{-1} \right| \|p_{k}\|_{1} + \int_{\mathcal{V}^{c}} |p_{k}| d\gamma + \int_{\mathcal{U}^{c}} |p_{k}| d\gamma$$

$$\leq \frac{\gamma(\mathcal{U}^{c}) + \gamma(\mathcal{V}^{c})}{\gamma(\mathcal{U})\gamma(\mathcal{V})} \|p_{k}\|_{1} + \int_{\mathcal{V}^{c}} |p_{k}| d\gamma + \int_{\mathcal{U}^{c}} |p_{k}| d\gamma$$

$$\lesssim (\gamma(\mathcal{U}^{c}) + \gamma(\mathcal{V}^{c})) \|p_{k}\|_{1} + \|p_{k}\|_{2} \sqrt{\gamma(\mathcal{V}^{c})} + \|p_{k}\|_{2} \sqrt{\gamma(\mathcal{U}^{c})}$$

$$\lesssim \|p_{k}\|_{2} \sqrt{\gamma(\mathcal{V}^{c})} + \|p_{k}\|_{2} \sqrt{\gamma(\mathcal{U}^{c})}$$
(C.14)

using that $\gamma(\mathcal{U}), \gamma(\mathcal{V}) \geq 1/2$. Furthermore,

$$\left\| \left(p_k - \int p_k d\gamma_{\mathcal{V}} \right) \mathbb{1}_{\mathcal{V}^c} \right\|_q \lesssim \| p_k \|_{2q} \gamma(\mathcal{V}^c)^{1/2q}.$$
(C.15)

Substituting (C.14) and (C.15) into (C.13) gives

$$\|\hat{p}_k\|_q \lesssim \left\| \left(p_k - \int p_k d\gamma_{\mathcal{V}} \right) \mathbb{1}_{\mathcal{V}} \right\|_q + \|p_k\|_{2q} \gamma(\mathcal{V}^c)^{1/2q} + \|p_k\|_2 \sqrt{\gamma(\mathcal{U}^c)}$$
(C.16)

For the third term, we have using (3.26) and (C.12) that

$$\sqrt{\gamma(\mathcal{U}^c)} \|p_k\|_2 \lesssim_k \exp(-(R-1)^2 d/4) c_{k+2}(0) d^{(k+2)/2}$$
(C.17)

For the second term, we have using (3.26) and (C.12) that

 $\|(p_k)\|$

$$\|p_k\|_{2q}\gamma(\mathcal{V}^c)^{1/2q} \lesssim_{k,q} \exp(-(S-1)^2 d/4q) c_{k+2}(0) d^{(k+2)/2}$$
(C.18)

Finally, for the first term we use (D.4) of Corollary D.3 with $f = p_k$. We have

$$\sup_{x \in \mathcal{V}} \|\nabla p_k(x)\| \le (k+2)c_{k+2}(0)(S\sqrt{d})^{k+1} =: M$$

Therefore,

and hence

$$\int_{\mathcal{V}} |p_k - \int p_k d\gamma_{\mathcal{V}}|^q d\gamma \lesssim_q M^q$$
$$-\int p_k d\gamma_{\mathcal{V}} \Big) \mathbb{1}_{\mathcal{V}} \Big\|_q \lesssim_q M \lesssim_{q,k} c_{k+2}(0) (S\sqrt{d})^{k+1}.$$
(C.19)

Using (C.19), (C.18), (C.17) in (C.16) gives

$$\begin{aligned} \|\hat{p}_{k}\|_{q} &\lesssim_{q,k} c_{k+2}(0)(S\sqrt{d})^{k+1} + \exp(-(S-1)^{2}d/4q)c_{k+2}(0)d^{(k+2)/2} + \exp(-(R-1)^{2}d/4)c_{k+2}(0)d^{(k+2)/2} \\ &= c_{k+2}(0)d^{(k+1)/2} \left(S^{k+1} + \sqrt{d}\exp(-(S-1)^{2}d/4q) + \sqrt{d}\exp(-(R-1)^{2}d/4q)\right). \end{aligned}$$
(C.20)

We now choose $S = 1 + \sqrt{2q \log(d+1)/d}$, so that $d(S-1)^2/4q = \log(d+1)/2 \ge \log\sqrt{d}$, and hence the second term in parentheses equals 1. Note that this choice of S also satisfies the required property that $\gamma(\mathcal{V}) \ge 1/2$. Indeed, using that $q \ge 1$ and $d \ge 1$, we have $\gamma(\mathcal{V}^c) \le \exp(-d(S-1)^2/2q) \le \exp(-\log(d+1)) \le 1/2$.

For this choice of S, the first term in parentheses is then bounded by some constant depending on k and q. Finally, since $R \ge 2$, the third term is bounded by an absolute constant uniformly over $d \in \mathbb{N}$. Thus in total we get the bound $\|\hat{p}_k\|_q \leq_{q,k} c_{k+2}(0)d^{(k+1)/2}$ and hence

$$n^{-k/2} \|\hat{p}_k\|_q \lesssim_{q,k} c_{k+2}(0) \frac{d^{(k+1)/2}}{n^{k/2}} = \bar{\epsilon}_k(0)$$

Proof of Lemma C.4. Recall from Lemma 3.4 that for all $1 \le k \le L$, it holds

$$\partial_t^k r(t,x) = \int_0^1 \langle \nabla^{k+2} v(\lambda tx), x^{\otimes k+2} \rangle \lambda^k (1-\lambda) d\lambda.$$
 (C.21)

Using that

$$\sup_{\lambda \in [0,1], t \in [0,1/\sqrt{n}]} \left| \langle \nabla^{k+2} v(\lambda tx), x^{\otimes k+2} \rangle \right| \lambda^k (1-\lambda) \le c_{k+2}(R) ||x||^{k+2}$$

for all $||x|| \leq R\sqrt{d}$, we get

$$\|\partial_t^k r(t,\cdot) \mathbb{1}_{\mathcal{U}}\|_q \le c_{k+2}(R) \mathbb{E} \left[\|Z\|^{(k+2)q} \right]^{\frac{1}{q}} \lesssim_{k,q} c_{k+2}(R) d^{(k+2)/2},$$

from which the bound $n^{-k/2} \|\partial_t^k r(t,\cdot) \mathbb{1}_{\mathcal{U}}\|_q \lesssim_{k,q} \epsilon_k(R)$ immediately follows, for both k odd and k even. To

bound $\|\partial_t^k r(t,\cdot)\|_q$ in the case of odd $k \leq L-1$, we first Taylor expand (C.21) further as follows:

$$\partial_t^k r(t,x) - \frac{\langle \nabla^{k+2} v(0), x^{\otimes k+2} \rangle}{(k+1)(k+2)} = \int_0^1 \langle \nabla^{k+2} v(\lambda tx) - \nabla^{k+2} v(0), x^{\otimes k+2} \rangle \lambda^k (1-\lambda) d\lambda$$

$$= t \int_0^1 d\lambda \int_0^\lambda ds \langle \nabla^{k+3} v(stx), x^{\otimes k+3} \rangle \lambda^k (1-\lambda)$$

$$= t \int_0^1 ds \langle \nabla^{k+3} v(stx), x^{\otimes k+3} \rangle \int_s^1 \lambda^k (1-\lambda) d\lambda$$

$$= \frac{t}{(k+1)(k+2)} \int_0^1 \langle \nabla^{k+3} v(stx), x^{\otimes k+3} \rangle \left(1 - (k+2)s^{k+1} + (k+1)s^{k+2}\right) ds$$

$$=: t u_{k+1}(t,x)$$

(C.22)

Therefore, recalling that $p_k(x) = \langle \nabla^{k+2} v(0), x^{\otimes k+2} \rangle$ and that $t \leq n^{-1/2}$ we have

$$n^{-k/2} \|\partial_t^k r(t,\cdot) \mathbb{1}_{\mathcal{U}}\|_q \le \frac{n^{-k/2} \|p_k\|_q}{(k+1)(k+2)} + n^{-(k+1)/2} \|u_{k+1}(t,\cdot) \mathbb{1}_{\mathcal{U}}\|_q \tag{C.23}$$

Using the bound (C.6) for odd k, we get

$$n^{-k/2} \|p_k\|_q \le \bar{\epsilon}_k(0)$$

For $u_{k+1}(t, \cdot)$ we proceed as above, noting that

$$\sup_{t\in[0,1/\sqrt{n}],s\in[0,1]} \left| \langle \nabla^{k+3}v(stx), x^{\otimes k+3} \rangle \right| \le c_{k+3}(R) \|x\|^{k+3} \qquad \forall x \in \mathcal{U},$$

and $|1 - (k+2)s^{k+1} + (k+1)s^{k+2}| = |1 - s^{k+1} - (k+1)s^{k+1}(1-s)| \le k+2$. Hence

$$\|u_{k+1}(t,\cdot)\mathbb{1}_{\mathcal{U}}\|_q \le \frac{1}{k+1} c_{k+3}(R) \mathbb{E} \left[\|Z\|^{(k+3)q} \right]^{1/q} \lesssim_{k,q} c_{k+3}(R) d^{(k+3)/2}.$$

Substituting these bounds into (C.23) gives

$$n^{-k/2} \|\partial_t^k r(t, \cdot) \mathbb{1}_{\mathcal{U}}\|_q \lesssim_{q,k} \bar{\epsilon}_k(0) + c_{k+3}(R) n^{-(k+1)/2} d^{(k+3)/2} = \bar{\epsilon}_k(0) + \epsilon_{k+1}(R),$$

as desired.

D Lipschitz concentration

Theorem D.1 (Theorem A1 in [22], Theorem 2.1 in [15]). Let $\mathcal{U} \subset \mathbb{R}^d$ be a convex set and μ be a probability measure on \mathbb{R}^d given by $\mu(dx) = Z_{\mu}^{-1} e^{-H(x)} \mathbb{1}_{\mathcal{U}}(x) dx$, where Z_{μ} is the normalization constant. Suppose $H \in C^2(\mathcal{U})$, and $\nabla^2 H(x) \succ \alpha I_d$ for all $x \in \mathcal{U}$. Then μ satisfies $\mathrm{LSI}(\alpha)$, where LSI is the log Sobolev inequality.

In particular, therefore, the measure $\gamma_{\mathcal{U}}$, the restriction of the standard Gaussian $\gamma(dx) \propto e^{-\|x\|^2/2} dx$ to \mathcal{U} , satisfies LSI(1) for any convex set \mathcal{U} . Note that we consider $\gamma_{\mathcal{U}}$ to be a measure on \mathbb{R}^d .

Theorem D.2 (Proposition 5.4.1 in [2]). Let $f : \mathbb{R}^d \to \mathbb{R}$ be *M*-Lipschitz, and $X \sim \mu$ for a probability measure μ on \mathbb{R}^d satisfying LSI(1). Then the random variable f(X) is M^2 sub-Gaussian, which by definition means that

$$\int e^{\lambda f} d\mu \le \exp\left(\lambda \int f d\mu + \lambda^2 M^2/2\right), \quad \forall \lambda \in \mathbb{R}.$$
(D.1)

This in turn implies that

$$\int \left| f - \int f d\mu \right|^q d\mu \le (2\sqrt{q}M)^q, \quad \forall q \in \mathbb{N}.$$
(D.2)

-			I.	
			I.	
			I.	
-	-	-		

Note that (D.2) follows from (D.1) by a standard Chernoff bound and the formula $\mathbb{E}[|X|^q] = \int_0^\infty qt^{q-1}\mathbb{P}(|X| \ge t)dt$.

Corollary D.3. Suppose $f \in C^1(\mathbb{R}^d)$ and let $\mathcal{U} \subset \mathbb{R}^d$ be convex. If $\sup_{x \in \mathcal{U}} \|\nabla f(x)\| \leq M$, then

$$\|e^{f}\mathbb{1}_{\mathcal{U}}\|_{\lambda} = \left(\int_{\mathcal{U}} e^{\lambda f(x)}\gamma(dx)\right)^{1/\lambda} \le \exp\left(\int f d\gamma_{\mathcal{U}} + M^{2}\lambda/2\right),\tag{D.3}$$

and

$$\int_{\mathcal{U}} \left| f - \int f d\gamma_{\mathcal{U}} \right|^q d\gamma \le (2\sqrt{q}M)^q.$$
(D.4)

Proof. Since \mathcal{U} is convex, the bound $\sup_{x \in \mathcal{U}} \|\nabla f(x)\| \leq M$ implies f is M-Lipschitz on \mathcal{U} . Let \tilde{f} be a Lipschitz extension of f to \mathbb{R}^d ; that is, $f|_{\mathcal{U}} = \tilde{f}|_{\mathcal{U}}$, and \tilde{f} is M-Lipschitz on \mathbb{R}^d . Such an extension exists by [18]. Note that if $X \sim \gamma_{\mathcal{U}}$, then f(X) is equal to $\tilde{f}(X)$ in distribution. By Theorem D.1, the measure $\gamma_{\mathcal{U}}$ satisfies LSI(1). Therefore by Theorem D.2, we have

$$\int_{\mathcal{U}} e^{\lambda f} d\gamma \leq \int e^{\lambda f} d\gamma_{\mathcal{U}} = \int e^{\lambda \tilde{f}} d\gamma_{\mathcal{U}} \leq \exp\left(\lambda \int \tilde{f} d\gamma_{\mathcal{U}} + M^2 \lambda^2/2\right) = \exp\left(\lambda \int f d\gamma_{\mathcal{U}} + M^2 \lambda^2/2\right).$$

Similarly by (D.2) we have

$$\int_{\mathcal{U}} \left| f - \int f d\gamma_{\mathcal{U}} \right|^q d\gamma \le \int \left| f - \int f d\gamma_{\mathcal{U}} \right|^q d\gamma_{\mathcal{U}} = \int \left| \tilde{f} - \int \tilde{f} d\gamma_{\mathcal{U}} \right|^q d\gamma_{\mathcal{U}} \le (2\sqrt{q}M)^q.$$

E Auxiliary results and postponed proofs

Proof of Lemma 3.5. In this proof, we use [m] to denote the set $\{1, 2, \ldots, m\}$. It is well known [6, 21] that for $k \ge 1$, we have

$$B_k(x_1, \dots, x_k) = \sum_{r=1}^k B_{k,r}(x_1, \dots, x_{k-r+1}),$$
(E.1)

where

$$B_{k,r}(x_1, \dots, x_{k-r+1}) = k! \sum_{\substack{j_1+2j_2+\dots+(k-r+1)j_{k-r+1}=k\\j_1+\dots+j_{k-r+1}=r\\j_i \ge 0 \ \forall i \in [k-r+1]}} \prod_{i=1}^{k-r+1} \frac{x_i^{j_i}}{(i!)^{j_i}(j_i)!}$$
(E.2)

are the partial Bell polynomials. For convenience, we will instead sum over all j_1, \ldots, j_k such that $j_1 + 2j_2 + \cdots + kj_k = k$ and $j_1 + \cdots + j_k = r$ and $j_1, \ldots, j_k \ge 0$ for all k. One can show that the only solutions to this equation have j_1, \ldots, j_{k-r+1} nonzero, but including j_i 's such that $j_i = 0$ does not affect the product, since $\frac{x_i^{j_i}}{(i!)^{j_i}(j_i)!} = 1$ if $j_i = 0$. Therefore another valid formula for $B_{k,r}$ is

$$B_{k,r}(x_1, \dots, x_k) = k! \sum_{\substack{j_1+2j_2+\dots+kj_k=k\\j_1+\dots+j_k=r\\j_i \ge 0 \ \forall i \in [k]}} \prod_{i=1}^k \frac{(x_i/i!)^{j_i}}{(j_i)!},$$
(E.3)

Now, we let $y_i = x_i/i!$ for i = 1, ..., k, and separate the numerator from the denominator in (E.3), to get

$$B_{k,r}(x_1, \dots, x_k) = \frac{k!}{r!} \sum_{\substack{j_1+2j_2+\dots+kj_k=k\\j_1+\dots+j_k=r\\j_i \ge 0 \ \forall i \in [k]}} \binom{r}{j_1, \dots, j_k} \prod_{i=1}^k y_i^{j_i}$$
(E.4)

Consider the product $\prod_{i=1}^{k} y_i^{j_i}$ in (E.4) for a fixed choice of j_1, \ldots, j_k satisfying the constraints. Note that there are j_1 copies of y_1 , j_2 copies of y_2 and so on, up to j_k copies of y_k . There are r total terms in the product, including copies. For example if k = 11, r = 6, $j_1 = 3$, $j_2 = 1$, $j_3 = 2$ and the other j_i 's are zero, then the terms arising in the product are

$$y_1, y_1, y_1, y_2, y_3, y_3$$

Note that there are $r!/j_1!...j_k!$ permutations of this list which correspond to the same set of y_i 's and therefore to the same product. To each permutation, assign m_i to be the index of the y in position i. In the example given above, in that order, we would have $m_1 = 1, m_2 = 1, m_3 = 1, m_4 = 2, m_5 = 3, m_6 = 3$. For the permutation

$$y_1, y_3, y_2, y_3, y_1, y_1,$$

we would take $m_1 = 1, m_2 = 3, m_3 = 2, m_4 = 3, m_5 = 1, m_6 = 1$. For each assignment, we have $m_1 + \cdots + m_r = k$, and

$$\prod_{i=1}^k y_i^{j_i} = \prod_{\ell=1}^r y_{m_\ell}$$

Let $\mathcal{M}(j_1, \ldots, j_k)$ be the set of distinct assignments m_1, \ldots, m_r ; we know that $|\mathcal{M}(j_1, \ldots, j_k)| = r!/j_1! \ldots j_k!$. We conclude that

$$\binom{r}{j_1,\ldots,j_k}\prod_{i=1}^k y_i^{j_i} = \sum_{(m_1,\ldots,m_r)\in\mathcal{M}(j_1,\ldots,j_k)}\prod_{\ell=1}^r y_{m_\ell}.$$

Finally, it is clear that for distinct j_1, \ldots, j_k , the sets $\mathcal{M}(j_1, \ldots, j_k)$ are disjoint, and

$$\bigcup_{\substack{j_1+2j_2+\dots+kj_k=k\\j_1+\dots+j_k=r\\j_k>0 \ \forall i \in [k]}} \mathcal{M}(j_1,\dots,j_k) = \{(m_1,\dots,m_r) : m_1+\dots+m_r = k, 1 \le m_\ell \le k \ \forall \ell \in [r]\}.$$

In fact, if $m_1 + \cdots + m_r = k$ and $m_\ell \ge 1$ for all ℓ , then it automatically follows that $m_\ell \le k$ for all ℓ , so we can omit this condition. We conclude that

$$B_{k,r}(x_1,\ldots,x_k) = \frac{k!}{r!} \sum_{\substack{j_1+2j_2+\cdots+kj_k=k\\j_1+\cdots+j_k=r\\j_i\ge 0 \,\,\forall i\in[k]}} \binom{r}{j_1,\ldots,j_k} \prod_{i=1}^k y_i^{j_i} = \frac{k!}{r!} \sum_{\substack{m_1+\cdots+m_r=k\\m_\ell\ge 1 \,\,\forall \ell\in[r]}} \prod_{\ell=1}^r y_{m_\ell}.$$

Summing over all $1 \le r \le k$ and recalling that $y_m = x_m/m!$ concludes the proof.

Lemma E.1. Let $f_1, \ldots, f_\ell : \mathbb{R}^d \to \mathbb{R}$. If there exist $\epsilon_1, \ldots, \epsilon_\ell \geq 0$ such that

$$\|n^{-i/2}f_i\|_{p\ell^2} \le C(p,\ell)\epsilon_i \quad \forall i = 1,\dots,\ell,$$

then

$$n^{-\ell/2} \|B_{\ell}(f_1, \dots, f_{\ell})\|_p \lesssim_{p,\ell} B_{\ell}(\epsilon_1, \dots, \epsilon_{\ell}) \lesssim_{p,\ell} \left(1 \vee \max_{1 \le i \le \ell} \epsilon_i\right)^{\ell}$$
(E.5)

Similarly, if there exist $\epsilon_1, \ldots, \epsilon_\ell \geq 0$ such that

$$\|n^{-i/2}f_i\,\mathbb{1}_{\mathcal{U}}\|_{p\ell^2} \le C(p,\ell)\epsilon_i \quad \forall i=1,\ldots,\ell,$$

then

$$n^{-\ell/2} \|B_{\ell}(f_1,\ldots,f_{\ell})\,\mathbb{1}_{\mathcal{U}}\|_p \lesssim_{p,\ell} B_{\ell}(\epsilon_1,\ldots,\epsilon_{\ell}) \lesssim_{p,\ell} \left(1 \vee \max_{1 \le i \le \ell} \epsilon_i\right)^{\ell}.$$
(E.6)

Proof. Using the definition (2.7) of the Bell polynomials, we have

$$\|B_{\ell}(f_1,\ldots,f_{\ell})\|_p \le \ell! \sum_{\substack{j_1+2j_2+\cdots+\ell j_{\ell}=\ell\\j_1,\ldots,j_{\ell}\ge 0}} \left\|\prod_{i=1}^{\ell} f_i^{j_i}\right\|_p \prod_{i=1}^{\ell} \frac{1}{j_i!(i!)^{j_i}}.$$
 (E.7)

Next, using the generalized Hölder's inequality, we have

$$\|\prod_{i=1}^{\ell} f_i^{j_i}\|_p = \left(\|\prod_{i=1}^{\ell} f_i^{pj_i}\|_1\right)^{1/p} \le \prod_{i=1}^{\ell} \|f_i^{pj_i}\|_{\ell}^{1/p} = \prod_{i=1}^{\ell} \|f_i\|_{p\ell j_i}^{j_i}\|_{\ell}^{1/p}$$

We conclude that

$$\|B_{\ell}(f_1,\ldots,f_{\ell})\|_p \le \ell! \sum_{\substack{j_1+2j_2+\cdots+\ell \\ j_1,\ldots,j_{\ell}\ge 0}} \prod_{i=1}^{\ell} \|f_i\|_{p\ell j_i}^{j_i} \prod_{i=1}^{\ell} \frac{1}{j_i!(i!)^{j_i}}$$

and hence

$$n^{-\ell/2} \|B_{\ell}(f_1, \dots, f_{\ell})\|_p \le \ell! \sum_{\substack{j_1+2j_2+\dots+\ell j_{\ell}=\ell\\j_1,\dots,j_{\ell}\ge 0}} \prod_{i=1}^{\ell} \|n^{-i/2} f_i\|_{p\ell j_i}^{j_i} \prod_{i=1}^{\ell} \frac{1}{j_i!(i!)^{j_i}}$$

Substituting the bound $||n^{-i/2}f_i||_{p\ell j_i} \le ||n^{-i/2}f_i||_{p\ell^2} \le C(p,\ell)\epsilon_i$ now gives

$$n^{-\ell/2} \|B_{\ell}(f_1,\ldots,f_{\ell})\|_p \lesssim_{p,\ell} \ell! \sum_{\substack{j_1+2j_2+\cdots+\ell j_{\ell}=\ell\\j_1,\ldots,j_{\ell}\ge 0}} \prod_{i=1}^{\ell} \frac{\epsilon_i^{j_i}}{j_i!(i!)^{j_i}} = B_{\ell}\left(\epsilon_1,\ldots,\epsilon_{\ell}\right) \lesssim \left(1 \lor \max_{1\le i\le \ell} \epsilon_i\right)^{\ell},$$

as desired. To prove (E.6), it suffices to note that $B_{\ell}(f_1, \ldots, f_{\ell})\mathbb{1}_{\mathcal{U}} = B_{\ell}(f_1\mathbb{1}_{\mathcal{U}}, \ldots, f_{\ell}\mathbb{1}_{\mathcal{U}})$, since $(\mathbb{1}_{\mathcal{U}})^k = \mathbb{1}_{\mathcal{U}}$ for any $k \in \mathbb{N}$.

Lemma E.2 (Lemma F.1 in [13]). Let a, b > 0. Then

$$I := \frac{1}{(2\pi)^{d/2}} \int_{\|x\| \ge a\sqrt{d}}^{\infty} e^{-b\sqrt{d}\|x\|} dx \le \frac{d}{e} \exp\left(\left[\log\left(\frac{e}{b}\right) - \frac{1}{2}ab\right]d\right)$$
(E.8)

In particular, if $a \geq \frac{4}{b} \log \frac{e}{b}$ then

$$I \le \frac{d}{e} e^{-abd/4}$$

Proof. Switching to polar coordinates and then changing variables, we have

$$I = \frac{S_{d-1}}{(2\pi)^{d/2}} \int_{a\sqrt{d}}^{\infty} u^{d-1} e^{-b\sqrt{d}u} du = \frac{S_{d-1}}{(2\pi)^{d/2} (b\sqrt{d})^d} \int_{abd}^{\infty} u^{d-1} e^{-u} du,$$
(E.9)

where S_{d-1} is the surface area of the unit sphere. Now, we have

$$\frac{S_{d-1}}{(2\pi)^{d/2}} = \frac{2\pi^{d/2}}{\Gamma(d/2)(2\pi)^{d/2}} \le 2\frac{(2e/d)^{d/2-1}}{2^{d/2}} = \left(\frac{e}{d}\right)^{\frac{d}{2}-1},$$

using that $\Gamma(d/2) \ge (d/2e)^{d/2-1}$. To bound the integral in (E.9), we use Lemma E.3 with $\lambda = abd$ and c = d. Combining the resulting bound with the above bound on $S_{d-1}/(2\pi)^{d/2}$, we get

$$I \le \left(\frac{e}{d}\right)^{\frac{d}{2}-1} (b\sqrt{d})^{-d} e^{-tabd} \left(\frac{d}{1-t}\right)^d = \frac{d}{e} \left(\frac{\sqrt{e}}{b(1-t)}\right)^d e^{-tabd}$$
(E.10)

Taking t = 1/2 and noting that $2\sqrt{e} \le e$ concludes the proof.

Lemma E.3 (Lemma F.2 in [13]). For all $\lambda, c > 0$ and $t \in (0, 1)$ it holds

$$\int_{\lambda}^{\infty} u^{c-1} e^{-u} du \le e^{-\lambda t} \left(\frac{c}{1-t}\right)^c$$

Proof. Let X be a random variable with gamma distribution $\Gamma(c, 1)$. Then the desired integral is given by $\Gamma(c)\mathbb{P}(X \ge \lambda)$. Now, the mgf of $\Gamma(c, 1)$ is $\mathbb{E}[e^{Xt}] = (1 - t)^{-c}$, defined for t < 1. Hence for all $t \in (0, 1)$ we have

$$\mathbb{P}(X \ge \lambda) \le e^{-\lambda t} (1-t)^{-c}.$$
(E.11)

Multiplying both sides by $\Gamma(c)$ and using that $\Gamma(c) \leq c^c$ gives the desired bound.

References

- George E. Andrews. The Theory of Partitions. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1984.
- [2] Dominique Bakry, Ivan Gentil, Michel Ledoux, et al. Analysis and geometry of Markov diffusion operators, volume 103. Springer, 2014.
- [3] Rina Foygel Barber, Mathias Drton, and Kean Ming Tan. Laplace approximation in high-dimensional Bayesian regression. In *Statistical Analysis for High-Dimensional Data: The Abel Symposium 2014*, pages 15–36. Springer, 2016.
- [4] N. Bleistein and R.A. Handelsman. Asymptotic Expansions of Integrals. Dover Books on Mathematics Series. Dover Publications, 1986.
- [5] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. Concentration Inequalities: A Nonasymptotic Theory of Independence. Oxford University Press, 02 2013.
- [6] Louis Comtet. Advanced Combinatorics: The art of finite and infinite expansions. Springer Science & Business Media, 2012.
- [7] Guillaume P Dehaene. A deterministic and computable Bernstein-von Mises theorem. arXiv preprint arXiv:1904.02505, 2019.
- [8] Adrian Fischer, Robert E Gaunt, Gesine Reinert, and Yvik Swan. Normal approximation for the posterior in exponential families. arXiv preprint arXiv:2209.08806, 2022.
- [9] Tapio Helin and Remo Kretschmann. Non-asymptotic error estimates for the Laplace approximation in Bayesian inverse problems. *Numerische Mathematik*, 150(2):521–549, 2022.
- [10] Jonathan H Huggins, Trevor Campbell, Mikolaj Kasprzak, and Tamara Broderick. Practical bounds on the error of Bayesian posterior approximations: A nonasymptotic approach. arXiv preprint arXiv:1809.09505, 2018.
- [11] Tadeusz Inglot and Piotr Majerski. Simple upper and lower bounds for the multivariate Laplace approximation. Journal of Approximation Theory, 186:1–11, 2014.
- [12] Mikolaj J Kasprzak, Ryan Giordano, and Tamara Broderick. How good is your Gaussian approximation of the posterior? Finite-sample computable error bounds for a variety of useful divergences. arXiv preprint arXiv:2209.14992, 2022.
- [13] Anya Katsevich. The Laplace approximation accuracy in high dimensions: a refined analysis and new skew adjustment. arXiv preprint arXiv:2306.07262, 2023.
- [14] William D Kirwin. Higher asymptotics of laplace's approximation. Asymptotic Analysis, 70(3-4):231–248, 2010.

- [15] Alexander V Kolesnikov and Emanuel Milman. Riemannian metrics on convex sets with applications to Poincaré and log-Sobolev inequalities. *Calculus of Variations and Partial Differential Equations*, 55(4):77, 2016.
- [16] Vassili N Kolokoltsov. Rates of convergence in Laplace's integrals and sums and conditional central limit theorems. *Mathematics*, 8(4):479, 2020.
- [17] Tomasz M Łapiński. Multivariate Laplace's approximation with estimated error and application to limit theorems. Journal of Approximation Theory, 248:105305, 2019.
- [18] E. J. McShane. Extension of range of functions. Bulletin of the American Mathematical Society, 40(12):837 – 842, 1934.
- [19] Gergő Nemes. An explicit formula for the coefficients in Laplace's method. Constructive Approximation, 38:471–487, 2013.
- [20] Helen Ogden. On the error in Laplace approximations of high-dimensional integrals. *Stat*, 10(1):e380, 2021.
- [21] John Riordan. An Introduction to Combinatorial Analysis. Princeton University Press, 1978.
- [22] André Schlichting. Poincaré and log-Sobolev inequalities for mixtures. Entropy, 21(1):89, 2019.
- [23] Zhenming Shun and Peter McCullagh. Laplace approximation of high dimensional integrals. Journal of the Royal Statistical Society: Series B (Methodological), 57(4):749–760, 1995.
- [24] Vladimir Spokoiny. Dimension free nonasymptotic bounds on the accuracy of high-dimensional laplace approximation. SIAM/ASA Journal on Uncertainty Quantification, 11(3):1044–1068, 2023.
- [25] Luke Tierney and Joseph B Kadane. Accurate approximations for posterior moments and marginal densities. Journal of the American Statistical Association, 81(393):82–86, 1986.
- [26] John Wojdylo. Computing the coefficients in Laplace's method. SIAM review, 48(1):76–96, 2006.
- [27] R. Wong. Asymptotic Approximations of Integrals. Society for Industrial and Applied Mathematics, 2001.
- [28] Xinzhen Zhang, Chen Ling, and Liqun Qi. The best rank-1 approximation of a symmetric tensor and related spherical optimization problems. SIAM Journal on Matrix Analysis and Applications, 33(3):806– 821, 2012.