

# Structure of Massive Gauge/Gravity Scattering Amplitudes, Equivalence Theorems, and Extended Double-Copy with Compactified Warped Space

---

Yanfeng Hang,<sup>a</sup> Wei-Wei Zhao,<sup>b</sup> Hong-Jian He,<sup>b,c</sup> Yin-Long Qiu<sup>b</sup>

<sup>a</sup>*Department of Physics and Astronomy,  
Northwestern University, Evanston, IL, USA*

<sup>b</sup>*T. D. Lee Institute & School of Physics and Astronomy,  
Key Laboratory for Particle Astrophysics and Cosmology,  
Shanghai Key Laboratory for Particle Physics and Cosmology,  
Shanghai Jiao Tong University, Shanghai, China*

<sup>c</sup>*Physics Department & Institute of Modern Physics, Tsinghua University, Beijing, China;  
Center for High Energy Physics, Peking University, Beijing, China*

*E-mail: [yfhang@northwestern.edu](mailto:yfhang@northwestern.edu), [weiwei\\_zhao@sjtu.edu.cn](mailto:weiwei_zhao@sjtu.edu.cn),  
[hjhe@sjtu.edu.cn](mailto:hjhe@sjtu.edu.cn), [silverdragon@sjtu.edu.cn](mailto:silverdragon@sjtu.edu.cn)*

ABSTRACT:

We study the structure of scattering amplitudes of massive Kaluza-Klein (KK) states in the compactified 5-dimensional warped gauge and gravity theories. We present systematic formulations of the gauge theory equivalence theorem (GAET) and the gravitational equivalence theorem (GRET) for warped KK theories in  $R_\xi$  gauge, where the GAET connects the scattering amplitudes of longitudinal KK gauge bosons to that of the corresponding KK Goldstone bosons and the GRET connects the scattering amplitudes of KK gravitons of helicity-zero (helicity-one) to that of the corresponding gravitational KK Goldstone bosons. We analyze the structure of 3-point and 4-point scattering amplitudes of massive KK gauge bosons and of massive KK gravitons as well as their corresponding Goldstone bosons. We first prove the GAET and GRET explicitly for the fundamental 3-point KK gauge/gravity scattering amplitudes. We then demonstrate that the validity of the GAET and GRET for 4-point gauge/gravity scattering amplitudes can be reduced to the validity of GAET and GRET for 3-point gauge/gravity scattering amplitudes at tree level. With these, we study the double-copy construction of KK scattering amplitudes in the warped gauge/gravity theories. We newly realize the double-copy for massive 3-point full gauge/gravity amplitudes at tree level under proper correspondences of color-kinematics and of gauge/gravity couplings, whereas we can construct the double-copy for 4-point KK gauge/gravity amplitudes to the leading order (LO) of high energy expansion. We further demonstrate that this LO double-copy construction can be extended to  $N$ -point KK scattering amplitudes with  $N \geq 4$ .

arXiv:2406.12713 [hep-th]

---

## Contents

<b>1 Introduction</b>	<b>3</b>
<b>2 Formulation of Gauge and Gravitational Equivalence Theorems with Compactified Warped Space</b>	<b>7</b>
2.1 Warped 5d Compactification	7
2.2 Formulating Gauge ET for Warped KK Gauge Theory	8
2.3 Formulating Gravitational ET for Warped KK Gravity Theory	13
2.3.1 Warped KK Gravity Theory with BRST Quantization	13
2.3.2 Formulating GRET for Warped KK Gravity Theory (I)	18
2.3.3 Formulating GRET for Warped KK Gravity Theory (II)	21
<b>3 Structure of Massive Scattering Amplitudes from Warped Gauge and Gravitational Equivalence Theorems</b>	<b>26</b>
3.1 Warped Gauge and Gravitational ETs for 3-Point KK Amplitudes	26
3.1.1 Warped GAET for 3-Point KK Gauge Boson Amplitudes	26
3.1.2 Warped GRET for 3-Point KK Graviton Amplitudes	30
3.2 Warped GAET and GRET: from 3-Point to 4-Point Amplitudes	34
3.2.1 Warped GAET: from 3-Point to 4-Point KK Amplitudes	34
3.2.2 Warped GRET: from 3-Point to 4-Point KK Amplitudes	42
<b>4 Extended Double-Copy for Warped KK Equivalence Theorems</b>	<b>44</b>
4.1 Double-Copy Construction of 3-Point Amplitudes and Warped GRET	45
4.2 Double-Copy Construction of 4-Point Warped GRET	49
4.2.1 Construction of 4-Point Elastic KK Graviton Amplitudes	49
4.2.2 Construction of 4-Point Inelastic KK Graviton Amplitudes	52
4.3 Extension of Warped LO Double-Copy with $N \geq 4$	59
4.4 Prescription for Warped Double-Copy Construction at Leading Order	64
<b>5 Conclusions</b>	<b>66</b>
<b>A Kinematics of KK Particle Scattering</b>	<b>70</b>
<b>B BRST Quantization for GRET Formulation</b>	<b>71</b>
B.1 BRST Quantization for Warped 5d Gravity	72
B.2 Gravitational ET of Type-I	72
B.3 Gravitational ET of Type-II	75
<b>C Feynman Rules for Warped KK Gauge and Gravity Theories</b>	<b>77</b>
C.1 Feynman Rules for Warped KK Gauge Theory	77
C.2 Feynman Rules for Warped KK Gravity Theory	79
<b>D Proving 3-Point and 4-Point Warped Identities</b>	<b>83</b>
<b>References</b>	<b>89</b>

---

# 1 Introduction

Scattering amplitude is an elementary means for studying fundamental forces in nature, and can bridge theories with experiments. There have been substantial efforts [1] for studying various massless scattering amplitudes over the past four decades. Studies on massive scattering amplitudes have attracted increasing interests in recent years, including to consider some models of massive gauge bosons and gravitons. Among these, the compactified higher dimensional gauge and gravity theories à la Kaluza-Klein (KK) [2] stand out as a fundamental approach for *consistent mass generation* of the excited KK states of gauge bosons and gravitons.<sup>1</sup>

Simple models of massive gauge bosons and gravitons include such as the massive Yang-Mills (YM) gauge theory [6] and massive Fierz-Pauli (FP) gravity [7] and alike [8], which add mass terms by hand and spoil the gauge symmetry or diffeomorphism invariance. The former is plagued with nonrenormalizability and the latter further suffers the van Dam-Veltman-Zakharov (vDVZ) discontinuity in the massive graviton propagator [9][10]. Moreover, such massive gauge and gravity theories generally exhibit bad high energy behaviors in their scattering amplitudes involving longitudinally polarized gauge bosons or gravitons. The reason behind such inconsistencies is because the nonconservation of physical degrees of freedom prohibits a smooth massless limit. In contrast, the KK gauge and gravity theories provide a truly consistent geometric “Higgs” mechanism for mass generations of KK gauge bosons and KK gravitons whose additional mass-induced physical degrees of freedom originate from absorbing the extra-dimensional components of the KK gauge bosons [11][12] and KK gravitons [14][15][16]. Hence, this mass-generation mechanism naturally conserves physical degrees of freedom for KK gauge bosons and KK gravitons, and thus has smooth massless limit. But this geometric mechanism for mass generations of KK gauge bosons and KK gravitons is realized through the compactification itself and does not invoke any additional Higgs boson of the conventional Higgs mechanism [17].

The geometric mechanism for mass generations of KK gauge bosons and KK gravitons can be formulated at the level of scattering  $S$ -matrix through the KK Equivalence Theorem (ET), as done previously for the flat 5d KK gauge theories [11][12][13] and the flat 5d KK gravity theories [15][16]. It is desirable to extend these KK ET formulations to *both* the KK gauge theories and KK gravity theories with warped 5d compactification [18][19][20], as we briefly discussed in Refs. [13][16] (cf. also a recent related study on the warped gravity KK ET [21]). The warped 5d KK gravity of Randall and Sundrum (RS1) [18] holds the promise to resolve the gauge hierarchy problem of the SM and to explain the vast discrepancy between the gravity force and other three gauge forces of the SM. It also predicts distinctive

---

<sup>1</sup>The KK theories also have important impacts on the experimental side because the KK compactification predicts an infinite tower of massive KK states for each known particle of the Standard Model (SM) and General Relativity (GR). The low-lying KK states in such KK theories have intrigued great phenomenological and experimental efforts over the past two decades [3]-[5], as they may provide the first signatures for the new physics beyond the SM, ranging from the KK states of the SM particles to the spin-2 KK gravitons and possible dark matter candidate.


KK mass spectrum and KK signatures which are important for the on-going experimental searches.

On the other hand, it is conjectured that the scattering amplitudes of gravitons and gauge bosons are connected by the striking relation of double-copy,  $\text{GR} = (\text{Gauge Theory})^2$ , which points to the common root of both the gravity force and gauge forces. It also provides a powerful means for efficiently computing the highly complex scattering amplitudes of spin-2 gravitons. Such double-copy relation was first found in string theories by Kawai-Lewellen-Tye (KLT) [23], which links the scattering amplitudes of massless closed strings to the products of scattering amplitudes of massless open strings at tree level, and further leads to the double-copy relation between the scattering amplitudes of massless gravitons and of massless gauge bosons in the field theory limit. With the inspiration of KLT, Bern, Carrasco and Johansson (BCJ) proposed the double-copy method in the quantum field theory framework via color-kinematics (CK) duality [24][25], which constructs the massless graviton amplitudes from the (squared) scattering amplitudes of massless gauge bosons. It is highly nontrivial to extend the massless double-copy formulation to the massive gauge and gravity theories. In a recent series of works, the extended massive KLT-type and BCJ-type double-copies were constructed for the compactified KK string theories [26] and for the KK gauge/gravity field theories under toroidal compactification of flat 5d (without or with orbifold) [15][16][27]. But, it is more challenging to extend double-copy to the KK gauge/gravity theories with warped extra dimension such as the RS1. This is because the double-copy for  $N$ -point ( $N \geq 4$ ) massive KK gauge/gravity scattering amplitudes was proved to directly work out only for the toroidal compactifications with flat extra dimensions [27]. In this work, we will demonstrate that double-copy can be realized for the 3-point full scattering amplitudes of warped KK gauge/gravity theories, whereas for the 4-point warped KK gauge/gravity scattering amplitudes the double-copy can hold at the leading order (LO) of high energy expansion (which are just what we need to formulate the ET in both the warped KK gauge and gravity theories).<sup>2</sup>

In this work, we study the structure of scattering amplitudes of massive KK states in the compactified 5-dimensional warped gauge and gravity theories. We present systematic formulations of the gauge theory equivalence theorem (GAET) and the gravitational equivalence theorem (GRET) for warped KK gauge/gravity theories up to loop level and in the  $R_\xi$  gauge. The GAET connects the scattering amplitudes of longitudinal KK gauge bosons to that of the corresponding KK Goldstone bosons, whereas the GRET connects the scattering amplitudes of KK gravitons of helicity-0 (helicity-1) to that of the corresponding gravitational KK Goldstone bosons. As a key idea of this work, *we take the GAET of KK Yang-Mills gauge theories as the truly fundamental ET formulation*, from

---

<sup>2</sup>In passing, the extensions of double-copy for the topologically massive Chern-Simons (CS) gauge/gravity theories were presented in [28][29]. There are other recent works in the literature tried to extend the massless double-copy formulation to the massive double-copies, including the 4d massive YM theory versus Fierz-Pauli-like gravity [30][31], the spontaneously broken YM-Einstein supergravity models with adjoint Higgs fields [32], the KK-inspired effective gauge theory with extra global U(1) [33], the 3d CS gauge/gravity theories with or without supersymmetry [34]-[36], and the massless QCD/gravity coupled to massive scalars/fermions/vectors [37]-[39].



**Figure 1:** Equivalence Theorem and Double-Copy Correspondences from 3-point (3pt) scattering amplitudes to 4-point (4pt) scattering amplitudes and from massive KK gauge scattering amplitudes to massive KK gravitational scattering amplitudes in the warped gauge and gravity theories.

which we can *reconstruct the GRET* of the corresponding KK gravity theories by using the double-copy at the leading order (LO) of high energy expansion. Then, we analyze the structure of 3-point and 4-point scattering amplitudes of massive KK gauge bosons and of massive KK gravitons, as well as their corresponding KK Goldstone bosons. We first prove explicitly the GAET and GRET for the fundamental 3-point KK gauge/gravity scattering amplitudes. We then demonstrate that the validity of the GAET and GRET for 4-point gauge/gravity scattering amplitudes can be *reduced to* the validity of GAET and GRET for 3-point gauge/gravity scattering amplitudes. With these, we further study the double-copy construction of the KK scattering amplitudes in the warped gauge/gravity theories. We newly realize the double-copy for the 3-point full gauge/gravity amplitudes at tree level under proper color-kinematics correspondence and gauge-gravity coupling correspondence without high energy expansion. Then, we construct double-copy for 4-point KK gauge/gravity scattering amplitudes to the leading order (LO) of high energy expansion. We can derive the GRET from GAET by using the LO double-copy construction for 3-point and 4-point KK scattering amplitudes. Finally, we demonstrate that this LO double-copy construction can be extended to the  $N$ -point KK gauge/gravity scattering amplitudes with  $N \geq 4$ .

In summary, we present the main logic of this study in Fig. 1. This figure shows the correspondence of the Equivalence Theorems from the GAET to the GRET, and the correspondence of Double-Copies from the 3-point KK scattering amplitudes to the 4-point KK scattering amplitudes and from the massive KK gauge scattering amplitudes to the massive KK gravitational scattering amplitudes. In the lower-left corner, we start from analyzing the most fundamental 3-point KK gauge boson scattering amplitudes and establish the GAET between 3-point scattering amplitudes of longitudinal KK gauge bosons and of the corresponding KK Goldstone bosons to the LO of high energy expansion. Then, along the bottom horizontal arrow we reach the lower-right corner where we compute the 4-point scattering amplitudes of longitudinal KK gauge bosons and of the KK Goldstone

bosons, as well as establishing the GAET for these 4-point amplitudes at the LO of high energy expansion. Here we use the word “Reconstruction” or “Reduction” to indicate that the validity of the GAET for 4-point KK gauge/Goldstone boson amplitudes can be reduced to (proved from) the validity of the GAET for 3-point KK gauge/Goldstone boson amplitudes. Next, we can go from the lower-left corner to the upper-left corner by the double-copy construction along the left vertical arrow and from the 3-point KK gauge boson amplitudes to the 3-point KK graviton amplitudes, as well as from the LO 3-point KK Goldstone boson amplitudes to the LO 3-point gravitational KK Goldstone amplitudes. Then, along the top horizontal arrow we go from the upper-left corner to reach the upper-right corner where the 4-point KK graviton amplitudes and gravitational KK Goldstone amplitudes are computed; and thus we establish the GRET for these 4-point gravitational KK amplitudes. Here we use the word “Reconstruction” or “Reduction” to indicate that along the top horizontal arrow the validity of the GRET for 4-point KK graviton/Goldstone boson amplitudes can be reduced to (proved from) the validity of the GRET for 3-point KK graviton/Goldstone boson amplitudes. Finally, we can go from the lower-right corner to the upper-right corner by the double-copy construction along the right vertical arrow and from the LO 4-point KK gauge (Goldstone) boson amplitudes to the LO 4-point KK graviton (Goldstone) amplitudes.

The rest of this paper is organized as follows. In Section 2, after introducing the warped 5d compactification with  $S^1/\mathbb{Z}_2$  orbifold (Section 2.1), we present the formulation of GAET in the warped KK gauge theory (Section 2.2), and then the formulations of the GRET of type-I,II in the warped KK gravity theory (Section 2.3). In Section 3, we study the structure of massive scattering amplitudes from the warped GAET and GRET. These include to first compute the 3-point scattering amplitudes of KK gauge bosons (KK Goldstone bosons) and of KK gravitons (KK gravitational Goldstone bosons), and to establish the GAET and GRET for 3-point KK amplitudes at the leading order of high energy expansion (Section 3.1). Then, we present the 4-point scattering amplitudes of KK gauge bosons (KK Goldstone bosons) and of KK gravitons (KK gravitational Goldstone bosons), and to establish the GAET and GRET for 4-point KK amplitudes at the leading order of high energy expansion (Section 3.2). Our key point is to demonstrate that the validity of the GRET for 4-point KK amplitude can be reduced to (proved from) the validity of the GAET for basic 3-point KK amplitudes. In Section 4, we study the double-copy construction of the KK scattering amplitudes in the warped gauge/gravity theories. We realize the double-copy for the 3-point full gauge/gravity amplitudes at tree level under the proper correspondence of color-kinematics and the gauge-gravity coupling correspondence without high energy expansion, whereas the double-copy can be constructed for the 4-point KK gauge/gravity amplitudes to the LO of high energy expansion. We further study the  $N$ -point ( $N \geq 4$ ) KK gauge/gravity scattering amplitudes at the leading order and their LO double-copy. Finally, we conclude in Section 5. The four Appendices present supporting materials for our analyses in the main text. Among these, Appendix A provides the kinematics of three and four KK-particle scattering; Appendix B presents the Becchi-Rouet-Stora-Tyutin (BRST) quantization for warped 5d KK gravity theory and further details for deriving the GRET

of type-I and type-II; Appendix C derives the Feynman Rules for warped KK gauge and gravity theories as needed for the analyses of Section 3; Appendix D proves the relevant 3-point and 4-point warped identities.

## 2 Formulation of Gauge and Gravitational Equivalence Theorems with Compactified Warped Space

In this section, we formulate the gauge and gravitational equivalence theorems for the compactified 5d warped gauge and gravity theories. In section 2.1, we present the warped 5d compactification for RS1. In section 2.2, we formulate the gauge theory ET (GAET) for the warped KK gauge theory. In section 2.3, we first present the BRST quantization (Sec. 2.3.1), and then formulate the gravitational ET (GRET) of type-I (Sec. 2.3.3) and the GRET of type-II (Sec. 2.3.2) for the warped KK gravity theory.

### 2.1 Warped 5d Compactification

We consider the warped 5d compactification à la Randall and Sundrum (RS1) [18], where a warped background metric generates weak scale from Planck scale through an exponential suppression. The background metric involves a five-dimensional non-factorizable warped geometry, derived from a slice of AdS<sub>5</sub> spacetime. This framework is realized with two 3-branes situated at fixed points of the  $S^1/\mathbb{Z}_2$  orbifold compactification, where one 3-brane is the UV brane (Planck brane) at the fixed point  $y=0$  and another 3-brane is IR brane (TeV brane) at the fixed point  $y=\bar{L}$ . Here the physical coordinate of the fifth dimension is denoted as  $y$ , so we have  $y\in[0,\bar{L}]$  with  $\bar{L}=\pi r_c$  denoting the size of fifth dimension and  $r_c$  denoting the compactification radius.

The 5-dimensional warped background metric in the physical coordinate takes the following nonfactorizable form:

$$ds^2 = e^{2A(y)}\eta_{\mu\nu}dx^\mu dx^\nu + dy^2, \quad (2.1)$$

where  $e^{2A(y)}$  is the warp factor. Then, we can transform the physical coordinates  $(x^\mu, y)$  into the conformal coordinate  $(x^\mu, z)$  by imposing the relation  $dy = e^{A(z)}dz$ . Thus, we have the conformally flat 5-dimensional spacetime metric:

$$ds^2 = e^{2A(z)}(\eta_{\mu\nu}dx^\mu dx^\nu + dz^2). \quad (2.2)$$

Setting  $A(z) = -ky$ , one can resolve the function  $A(z)$  as follows [5]:

$$A(z) = -\ln(1 + kz), \quad (2.3)$$

where  $k = \sqrt{-\Lambda/6}$  is the AdS curvature scale and has the order of Planck mass. (The parameter  $\Lambda$  denotes a negative bulk cosmological constant, as will be defined in Sec. 2.3.1.) Thus, we deduce  $z = (e^{ky} - 1)/k$ . Hence, in the conformal coordinates, we have  $z \in [0, L]$  with  $L \equiv [e^{k\bar{L}} - 1]/k$ , which corresponds to the physical coordinate  $y \in [0, \bar{L}]$  with  $\bar{L} = \pi r_c$ .

The Eq.(2.2) shows that the 5-dimensional warped metric tensor  $\hat{g}_{MN}$  is conformally flat and can be expressed as

$$\hat{g}_{MN} = e^{2A(z)}\hat{\eta}_{MN}, \quad (2.4)$$

where the flat 5d Minkowski metric  $\hat{\eta}_{MN}$  takes the form:

$$\hat{\eta}_{MN} = \hat{\eta}^{MN} = \text{diag}(-1, 1, 1, 1, 1). \quad (2.5)$$

For clarity of notations, we denote the 5d Lorentz indices by uppercase Latin letters (such as  $M, N = \mu, 5$ ) and the 4d Lorentz indices by the lowercase Greek letters (such as  $\mu, \nu = 0, 1, 2, 3$ ).

In the next subsections, we will consider the RS1 under  $S^1/\mathbb{Z}_2$  compactification for both warped 5d Yang-Mills gauge theory and the warped 5d GR theory. With these, we formulate the KK equivalence theorems for the warped 5d Yang-Mills gauge theory and the warped 5d GR theory.

## 2.2 Formulating Gauge ET for Warped KK Gauge Theory

The warped bulk gauge theories in RS1 were discussed in the literature [19][20][22]. We consider the Yang-Mills Lagrangian in warped 5d spacetime:

$$\hat{\mathcal{L}}_{\text{YM}} = \sqrt{-\hat{g}} \left( -\frac{1}{4} \hat{g}^{MP} \hat{g}^{NQ} \hat{F}_{MN}^a \hat{F}_{PQ}^a \right), \quad (2.6)$$

where the square root of the determinant of the warped 5d background metric tensor  $\hat{g}_{MN}$  is given by  $\sqrt{-\hat{g}} = \exp[5A(z)]$ . The 5d field strength tensor  $\hat{F}_{MN}^a$  takes the form:

$$\hat{F}_{MN}^a = \partial_M \hat{A}_N^a - \partial_N \hat{A}_M^a + g_5 f^{abc} \hat{A}_M^b \hat{A}_N^c, \quad (2.7)$$

where  $(a, b, c)$  denote the gauge-group indices,  $f^{abc}$  denotes the gauge-group structure constant, and  $g_5$  is the 5d gauge coupling constant having mass-dimension  $-1/2$ .

Expanding the Lagrangian (2.6) to the quadratic order, we derive the following terms:

$$\hat{\mathcal{L}}_{\text{YM}}^{(2)} = -\frac{e^{A(z)}}{4} \left\{ \left( \partial_\mu \hat{A}_\nu^a - \partial_\nu \hat{A}_\mu^a \right)^2 + 2 \left[ \left( \partial_z \hat{A}_\mu^a \right)^2 + \left( \partial_\mu \hat{A}_5^a \right)^2 \right] - 4 \partial_z \hat{A}_\mu^a \partial^\mu \hat{A}_5^a \right\}. \quad (2.8)$$

The mixing term between the 5d components  $\hat{A}_\mu^a$  and  $\hat{A}_5^a$  in Eq.(2.8) can be eliminated by including the following  $R_\xi$  gauge-fixing term:

$$\hat{\mathcal{L}}_{\text{GF}} = -\frac{e^{A(z)}}{2\xi} (\hat{\mathcal{F}}^a)^2, \quad \hat{\mathcal{F}}^a = \partial^\mu \hat{A}_\mu^a + \xi (A' + \partial_z) \hat{A}_5^a, \quad (2.9)$$

where  $\xi$  is the gauge-fixing parameter and  $A' = \partial_z A(z)$ . We note that the gauge-fixing term (2.9) contains an  $\hat{A}_\mu^a$ - $\hat{A}_5^a$  mixing term which precisely cancels the corresponding mixing term in Eq.(2.8) upon integration by part. The fifth component  $\hat{A}_5^a$  will be identified as the would-be Goldstone boson via a geometric ‘‘Higgs’’ mechanism of the KK compactification [11][13]. Using Eqs.(2.8)-(2.9), we derive the 5d equation of motion (EOM) for the free fields  $(\hat{A}_\mu^a, \hat{A}_5^a)$ :

$$(\partial_\nu^2 + e^{-A} \partial_z e^A \partial_z) \hat{A}_\mu^a - (1 - \xi^{-1}) \partial_\mu \partial_\nu \hat{A}^{a\nu} = 0, \quad (2.10a)$$

$$\partial_\mu^2 \hat{A}_5^a + \xi \partial_z (A' + \partial_z) \hat{A}_5^a = 0. \quad (2.10b)$$



Next, we compactify the warped 5d space under  $S^1/\mathbb{Z}_2$  orbifold, which corresponds to compactifying the fifth-dimension to a line segment  $y \in [0, \bar{L}]$  or  $z \in [0, L]$ , with  $\bar{L} = \pi r_c$  and  $L \equiv (e^{k\bar{L}} - 1)/k$ , where  $r_c$  denotes the compactification radius. We impose the gauge-covariant boundary conditions on the 5d gauge field strength,  $\hat{F}_{M5}^a = \hat{F}_{5M}^a = 0$  at  $z = 0, L$ . This leads to the Neumann and Dirichlet boundary conditions on the 5d gauge field components  $\hat{A}_\mu^a$  and  $\hat{A}_5^g$  as follows:

$$\partial_z \hat{A}_\mu^a(x, z) \Big|_{z=0, L} = 0, \quad \hat{A}_5^g(x, z) \Big|_{z=0, L} = 0. \quad (2.11)$$

Then, we can make KK expansions for the 5d gauge fields  $\hat{A}_\mu^a$  and  $\hat{A}_5^g$  in terms of the eigenfunctions of warped 5d space:

$$\hat{A}^{a\mu}(x, z) = \frac{1}{\sqrt{L}} \sum_{n=0}^{\infty} A_n^{a\mu}(x) f_n(z), \quad (2.12a)$$

$$\hat{A}^{a5}(x, z) = \frac{1}{\sqrt{L}} \sum_{n=1}^{\infty} A_n^{a5}(x) \tilde{f}_n(z). \quad (2.12b)$$

Substituting the KK expansions (2.12) into the 5d EOM (2.10) and using the 4d EOM for free KK gauge fields,

$$(\partial_\nu^2 - M_n^2) A_n^{a\mu} - (1 - \xi_n^{-1}) \partial_\mu \partial_\nu A_n^{a\nu} = 0, \quad (2.13a)$$

$$(\partial_\nu^2 - \tilde{M}_n^2) A_n^{a\mu} = 0, \quad (2.13b)$$

we deduce the following EOMs for the 5d eigenfunctions  $f_n(z)$  and  $\tilde{f}_n(z)$ :

$$(A' + \partial_z) \partial_z f_n(z) = -M_n^2 f_n(z), \quad (2.14a)$$

$$\partial_z (A' + \partial_z) \tilde{f}_n(z) = -\xi_n^{-1} \tilde{M}_n^2 \tilde{f}_n(z), \quad (2.14b)$$

where  $M_n$  and  $\tilde{M}_n$  are the mass eigenvalue of KK level- $n$ . Applying the partial derivative  $\partial_z$  on both sides of Eq.(2.14a) and comparing it with Eq.(2.14b), we find that  $\partial_z f_n \propto \tilde{f}_n$  and  $\tilde{M}_n^2 = \xi_n M_n^2$ . Thus, with proper normalization we infer the following equations from the above Eq.(2.14a)-(2.14b):

$$\partial_z f_n(z) = -M_n \tilde{f}_n(z), \quad (2.15a)$$

$$(A' + \partial_z) \tilde{f}_n(z) = M_n f_n(z), \quad (2.15b)$$

which connect 5d wavefunctions  $f_n(z)$  and  $\tilde{f}_n(z)$ . Substituting the KK expansions (2.12) into Eq.(2.11), we obtain the boundary conditions imposed on  $(f_n, \tilde{f}_n)$ :

$$\partial_z f_n(z) \Big|_{z=0, L} = 0, \quad \tilde{f}_n(z) \Big|_{z=0, L} = 0. \quad (2.16)$$

They also obey the orthonormal conditions:

$$\frac{1}{L} \int_0^L dz e^{A(z)} f_n(z) f_m(z) = \delta_{nm}, \quad \frac{1}{L} \int_0^L dz e^{A(z)} \tilde{f}_n(z) \tilde{f}_m(z) = \delta_{nm}. \quad (2.17)$$

The 5d wavefunctions  $f_n(z)$  and  $\tilde{f}_n(z)$  can be solved from the equation of motion (2.14) in terms of Bessel functions of order  $\nu = 1, 0$ :

$$f_n(z) = \frac{e^{-A(z)}}{N_n} \left[ J_1\left(\frac{M_n}{k} e^{-A(z)}\right) + b_{n0} Y_1\left(\frac{M_n}{k} e^{-A(z)}\right) \right], \quad (2.18a)$$

$$\tilde{f}_n(z) = \frac{e^{-A(z)}}{N_n} \left[ J_0\left(\frac{M_n}{k} e^{-A(z)}\right) + b_{n0} Y_0\left(\frac{M_n}{k} e^{-A(z)}\right) \right], \quad (2.18b)$$

where the solution  $f_n(z)$  agrees with Refs. [19][20]. In the above, the normalization factors  $N_n$  can be fixed by the orthonormal conditions (2.17), whereas the coefficient  $b_{n0}$  is derived as follows:

$$b_{n0} = -\frac{J_0\left(\frac{M_n}{k}\right)}{Y_0\left(\frac{M_n}{k}\right)}. \quad (2.19)$$

The KK mass  $M_n$  is determined by roots of the eigenvalue equation:

$$J_0\left(\frac{M_n}{k} e^{-A(L)}\right) + b_{n0} Y_0\left(\frac{M_n}{k} e^{-A(L)}\right) = 0. \quad (2.20)$$

For the massless zero-mode wavefunction  $f_0$ , we can solve it from the eigenvalue equation (2.14a) and obtain:

$$f_0 = \sqrt{\frac{1 - e^{-A(L)}}{A(L)}}. \quad (2.21)$$

Under the KK expansions for the 5d gauge fields (2.12), the 4d effective KK Lagrangian is derived by integrating over the fifth coordinate  $z$  on the interval  $[0, L]$ . The quadratic KK Yang-Mills Lagrangian is given by

$$\mathcal{L}_{\text{YM}}^{(2)} = \int_0^L dz \hat{\mathcal{L}}_{\text{YM}}^{(2)} = -\frac{1}{4} \sum_{n=0}^{\infty} \left( \partial_\mu A_\nu^{an} - \partial_\nu A_\mu^{an} \right)^2 - \frac{1}{2} \sum_{n=1}^{\infty} \left( \partial_\mu A_5^{an} + M_n A_\mu^{an} \right)^2. \quad (2.22)$$

Then, using Eq.(2.9) and making the KK expansions (2.12), we can derive the following  $R_\xi$  gauge-fixing term:

$$\mathcal{L}_{\text{GF}} = \int_0^L dz \hat{\mathcal{L}}_{\text{GF}} = \sum_{n=0}^{\infty} \frac{-1}{2\xi_n} (\mathcal{F}_n^a)^2, \quad (2.23a)$$

$$\mathcal{F}_n^a = \partial_\mu A_n^{a\mu} + \xi_n M_n A_n^{a5} = \mathbf{K}_n^T \mathbf{A}_n^a, \quad (2.23b)$$

in agreement with [11][13], where  $\mathbf{K}_n = (\partial_\mu, \xi_n M_n)^T$  and  $\mathbf{A}_n^a = (A_n^{a\mu}, A_n^{a5})^T$ . In the above gauge-fixing term, we have introduced a gauge-fixing parameter  $\xi_n$  for each KK level- $n$ .

Corresponding to the gauge-fixing (2.9), we have the 5d Faddeev-Popov ghost term,

$$\hat{\mathcal{L}}_{\text{FP}} = e^{A(z)} \hat{c}^a \hat{\mathbf{s}} \hat{\mathcal{F}}^a, \quad (2.24)$$

where  $\hat{\mathbf{s}}$  is the Becchi-Rouet-Stora-Tyutin (BRST) operator and the 5d BRST transformations take the form:

$$\hat{\mathbf{s}} \hat{A}^{aN} = D_b^{aN}(\hat{A}) \hat{c}^b, \quad \hat{\mathbf{s}} \hat{c}^a = \frac{1}{2} \hat{g}_5 C^{abc} \hat{c}^b \hat{c}^c, \quad \hat{\mathbf{s}} \hat{c}^a = -\xi^{-1} \hat{\mathcal{F}}^a. \quad (2.25)$$

We make KK expansions for ghost fields,

$$\hat{c}^a(x, z) = \frac{1}{\sqrt{L}} \sum_{n=0}^{\infty} c_n^a(x) f_n(z), \quad (2.26a)$$

$$\hat{\bar{c}}^a(x, z) = \frac{1}{\sqrt{L}} \sum_{n=1}^{\infty} \bar{c}_n^a(x) \tilde{f}_n(z), \quad (2.26b)$$

and derive the BRST transformations for the KK fields,

$$\begin{aligned} \widehat{\mathbf{S}} A_n^{a\mu} &= D_{nm}^{ab,\mu} c_m^b, & \widehat{\mathbf{S}} A_n^{a5} &= D_{nm}^{ab,5} c_m^b, \\ \widehat{\mathbf{S}} c_n^a &= \frac{1}{2} g C^{abc} \mathfrak{D}_n^{ab,m\ell} c_m^b c_\ell^c, & \widehat{\mathbf{S}} \bar{c}_n^a &= -\xi_n^{-1} \mathcal{F}_n^a, \end{aligned} \quad (2.27)$$

where the following notations are defined,

$$\begin{aligned} D_{nm}^{ab,\mu} &= -\delta^{ab} \delta_{nm} \partial^\mu + \left( \frac{1}{L} \int_0^L dz e^{2A} f_n f_m f_\ell \right) g C^{abc} A_\ell^{c\mu}, \\ D_{nm}^{ab,5} &= -\delta^{ab} \delta_{nm} M_m + \left( \frac{1}{L} \int_0^L dz e^{2A} \tilde{f}_n f_m \tilde{f}_\ell \right) g C^{abc} A_\ell^{c5}, \\ \mathfrak{D}_n^{ab,m\ell} &= \frac{1}{L} \int_0^L dz e^{2A} f_n f_m f_\ell. \end{aligned} \quad (2.28)$$

Using these, we can further derive a BRST identity for the propagator of the gauge fields  $\mathbf{A}_n^a = (A_n^{a\mu}, A_n^{a5})^T$  as follows:

$$\mathbf{K}_n^T \mathcal{D}_{nm}^{ab}(k) = -\mathbf{X}_{nm}^{ab}(k)^T, \quad (2.29)$$

where we introduce the notations,

$$\begin{aligned} \mathcal{D}_{nm}^{ab}(k) &= \langle 0 | T \mathbf{A}_n^a \mathbf{A}_m^b{}^T | 0 \rangle, & \mathcal{S}_{nm}(k) \delta^{ab} &= \langle 0 | T c_n^a \bar{c}_m^b | 0 \rangle, \\ \mathbf{X}_{nm}^{ab}(k) &= \underline{\mathbf{X}}_{mj}^{ab}(k) \mathcal{S}_{jn}(k), \\ \underline{\mathbf{X}}_{mj}^{ab}(k) &= \langle 0 | T \widehat{\mathbf{S}} \mathbf{A}_m^b | \bar{c}_j^a \rangle = \left( \langle 0 | T \widehat{\mathbf{S}} A_m^{b\mu} | \bar{c}_j^a \rangle \right) \equiv \begin{pmatrix} -i k^\mu \delta^{ab} [\delta_{jm} + \Delta_{jm}^a(k^2)] \\ -M_m \delta^{ab} [\delta_{jm} + \tilde{\Delta}_{jm}^a(k^2)] \end{pmatrix}, \end{aligned} \quad (2.30)$$

with  $\Delta_{jm}^a(k^2)$  and  $\tilde{\Delta}_{jm}^a(k^2)$  being loop-level quantities.

In the following, we derive the KK gauge boson equivalence theorem (KK GAET) for the compactified warped 5d Yang-Mills gauge theory. The KK GAET was discussed [13] for the compactified 5d Yang-Mills gauge theories with a general metric in connection to its deconstruction formulation, as an extension of the formulations in the flat 5d KK gauge theories [11][12]. Our current derivation differs from [13] since we focus on the warped RS1 and choose the conformal coordinate frame of  $(x^\mu, z)$  instead of the physical coordinate frame of  $(x^\mu, y)$ . Besides, we will further study explicitly the three-point and four-point KK gauge (Goldstone) boson scattering amplitudes in Section 3 and their double-copy in Section 4.

Due to the 5d gauge invariance, we can derive a Slavnov-Taylor (ST) type of identity:

$$\langle 0 | T \hat{\mathcal{F}}^{a_1}(\hat{x}_1) \hat{\mathcal{F}}^{a_2}(\hat{x}_2) \cdots \hat{\mathcal{F}}^{a_N}(\hat{x}_N) \hat{\Phi} | 0 \rangle = 0, \quad (2.31)$$

where  $\hat{\Phi}$  denotes any possible external physical states in 5d spacetime. This is analogous to the ST identity for external gauge-fixing functions in the 4d SM with spontaneous electroweak gauge symmetry breaking [40]. Then, after making the KK expansion, we obtain the following ST identities for the external KK gauge-fixing functions:

$$\langle 0|T\mathcal{F}_{n_1}^{a_1}(x_1)\mathcal{F}_{n_2}^{a_2}(x_2)\cdots\mathcal{F}_{n_N}^{a_N}(x_N)\tilde{\Phi}|0\rangle = 0. \quad (2.32)$$

Using the BRST identity (2.29), we can amputate each external  $\mathcal{F}^{an}$  function in the momentum space and obtain a corresponding identity for the  $S$ -matrix amplitude:

$$\mathcal{T}[\underline{\mathcal{F}}_{n_1}^{a_1}(k_1), \underline{\mathcal{F}}_{n_2}^{a_2}(k_2), \cdots, \underline{\mathcal{F}}_{n_N}^{a_N}(k_N), \Phi] = 0, \quad (2.33)$$

where each external momentum  $k_j$  ( $j = 1, 2, \cdots$ ) is set on-shell with  $k_j^2 = -M_{n_j}^2$ , each  $\underline{\mathcal{F}}_{n_j}^{a_j}(k_j)$  represents the amputated external gauge-fixing function, and  $\Phi$  denotes any possible physical states after the LSZ (Lehmann-Symanzik-Zimmermann) amputation. We can express the amputated gauge-fixing function as follows:

$$\underline{\mathcal{F}}_n^a = k_\mu A_n^{a\mu} - M_n \hat{C}_{nm}^a A_m^{a5} = M_n (A_S^{an} - \hat{C}_{nm}^a A_5^{am}) = M_n (A_L^{an} - \mathbb{Q}_n^a), \quad (2.34a)$$

$$\mathbb{Q}_n^a = \hat{C}_{nm}^a A_5^{am} + v_n^a, \quad A_S^{an} = \epsilon_S^\mu A_\mu^{an}, \quad A_L^{an} = \epsilon_L^\mu A_\mu^{an}, \quad v_n^a = v_n^\mu A_\mu^{an}, \quad (2.34b)$$

with notations defined as

$$\hat{C}_{nm}^a = i(M_m/M_n)[(1 + \tilde{\Delta}^a(k^2))(1 + \Delta^a(k^2))^{-1}]_{mn} = i\delta_{nm} + O(\text{loop}), \quad (2.35a)$$

$$\epsilon_S^\mu = k^\mu/M_n = \epsilon_L^\mu - v_n^\mu, \quad v_n^\mu = O(M_n/E_n), \quad (2.35b)$$

where the matrix form is used such that  $(\tilde{\Delta}^a)_{jj'} \equiv \tilde{\Delta}_{jj'}^a$  and  $(\Delta^a)_{jj'} \equiv \Delta_{jj'}^a$ . In the above,  $\epsilon_L^\mu$  and  $\epsilon_S^\mu$  denote the KK gauge boson's longitudinal polarization vector and scalar polarization vector respectively. Thus, from the identity (2.33) we deduce the KK gauge equivalence theorem (GAET) identity:

$$\mathcal{T}[A_{Ln_1}^{a_1}, \cdots, A_{Ln_N}^{a_N}, \Phi] = \mathcal{T}[\mathbb{Q}_{n_1}^{a_1}, \cdots, \mathbb{Q}_{n_N}^{a_N}, \Phi] \quad (2.36a)$$

$$= \hat{C}_{\text{mod}}^{n_j m_j} \mathcal{T}[A_5^{a_1 m_1}, \cdots, A_5^{a_N m_N}, \Phi] + \mathcal{T}_v, \quad (2.36b)$$

$$\mathcal{T}_v = \sum_{\ell=1}^N \tilde{C}_{\text{mod}, \ell}^{n_j m_j} \mathcal{T}[v^{a_1 n_1}, \cdots, v^{a_\ell n_\ell}, A_5^{a_{\ell+1} m_{\ell+1}}, \cdots, A_5^{a_N m_N}, \Phi], \quad (2.36c)$$

where the multiplicative modification factors are given by

$$\hat{C}_{\text{mod}}^{n_j m_j} = \hat{C}_{n_1 m_1}^{a_1} \cdots \hat{C}_{n_N m_N}^{a_N} = i^N \delta_{n_1 m_1} \cdots \delta_{n_N m_N} + O(\text{loop}), \quad (2.37a)$$

$$\tilde{C}_{\text{mod}, \ell}^{n_j m_j} = \hat{C}_{n_{\ell+1} m_{\ell+1}}^{a_{\ell+1}} \cdots \hat{C}_{n_N m_N}^{a_N} = i^{N-\ell} \delta_{n_{\ell+1} m_{\ell+1}} \cdots \delta_{n_N m_N} + O(\text{loop}). \quad (2.37b)$$

Such radiative modification factors also exist in the ET formulation for the 4d electroweak gauge theory of the SM [40]-[42]. Note that the residual term  $\mathcal{T}_v$  in the KK-ET identity is suppressed under high energy expansion due to  $v_n^\mu = O(M_n/E_n)$ . With these, we derive

the KK Equivalence Theorem for warped gauge theory (KK GAET) for the high energy scattering:

$$\mathcal{T}[A_L^{a_1 n_1}, \dots, A_L^{a_N n_N}, \Phi] = \hat{C}_{\text{mod}}^{n_j m_j} \mathcal{T}[A_5^{a_1 m_1}, \dots, A_5^{a_N m_N}, \Phi] + O(M_n/E_n), \quad (2.38)$$

where  $\Phi$  denotes any other external physical state(s) after LSZ amputation, and at tree-level the multiplicative factor  $\hat{C}_{\text{mod}}^{n_j m_j}$  just reduces to a simple factor  $(i^N \delta_{n_1 m_1} \cdots \delta_{n_N m_N})$ .

## 2.3 Formulating Gravitational ET for Warped KK Gravity Theory

In this subsection, we first present the BRST quantization for the warped KK gravity theory in Section 2.3.1. Then, we formulate the warped Gravitational Equivalence Theorem (GRET) for the KK gravitons with helicity  $\pm 1$  (called GRET of type-II) in Section 2.3.2 and for the longitudinal KK gravitons with helicity-0 (called GRET of type-I) in Section 2.3.3, respectively. Using the GRET and our generalized energy-power counting method [15], we further prove the different large energy cancellations for the scattering amplitudes of KK gravitons with helicity-0 and with helicity  $\pm 1$  around the end of this subsection.

### 2.3.1 Warped KK Gravity Theory with BRST Quantization

In this subsection, we consider the 5d RS1 with the fifth spatial dimension compactified on an interval  $[0, L]$ . The setup of RS1 contains warped 5d Einstein-Hilbert action and is compactified on a 5d interval  $[0, L]$  with the two 3-branes located at the boundaries. The relevant part of the warped 5d Lagrangian can be written as

$$\hat{\mathcal{L}}_{\text{RS}} = \frac{1}{2} \sqrt{-\hat{g}} \hat{M}_{\text{P}}^3 (\hat{R} - 2\Lambda), \quad (2.39)$$

where  $\hat{R}$  and  $\Lambda (< 0)$  are the 5d Ricci scalar curvature and cosmological constant respectively. The 5d Planck mass  $\hat{M}_{\text{P}}$  is given by  $\hat{M}_{\text{P}} = M_{\text{P}}^{\frac{2}{3}} \bar{L}^{-\frac{1}{3}}$ , where  $M_{\text{P}}$  denotes the 4d reduced Planck mass and is connected to the 4d Newton constant  $G$  via  $M_{\text{P}} = (8\pi G)^{-1/2}$ . The possible brane terms [18] are irrelevant for the current analysis.

To study the linearized theory, we apply the following weak field expansion for the 5d Lagrangian (2.39):

$$\hat{g}_{MN} = e^{2A(z)} (\hat{\eta}_{MN} + \hat{\kappa} \hat{h}_{MN}), \quad (2.40)$$

where the 5d gravitational coupling  $\hat{\kappa}$  has mass-dimension  $-3/2$  and is connected to the 4d gravitational coupling  $\kappa$  via  $\hat{\kappa} = \kappa \sqrt{L} = 2M_{\text{P}}^{-1} \sqrt{L}$ . (Here  $M_{\text{P}}$  is the 4d reduced Planck mass.) The 5d graviton field  $\hat{h}_{MN}$  can be further parametrized as follows [15, 16]:

$$\hat{h}_{MN} = \begin{pmatrix} \hat{h}_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \hat{\phi} & \hat{\mathcal{V}}_{\mu} \\ \hat{\mathcal{V}}_{\nu} & \hat{\phi} \end{pmatrix}, \quad (2.41)$$

where we denote the (off-diagonal) vector component as  $\hat{\mathcal{V}}_{\mu} \equiv \hat{h}_{\mu 5}$  and the (diagonal) scalar component as  $\hat{\phi} \equiv \hat{h}_{55}$ . Then, we derive the Lagrangian at quadratic order, which takes

the following form:

$$\begin{aligned}
\hat{\mathcal{L}}_{\text{RS}}^{(2)} = e^{3A(z)} & \left\{ -\frac{1}{4} \hat{h}^{\mu\nu} \left\{ (\eta_{\mu\alpha} \partial_\nu \partial_\beta + \eta_{\mu\beta} \partial_\nu \partial_\alpha + \eta_{\nu\alpha} \partial_\mu \partial_\beta + \eta_{\nu\beta} \partial_\mu \partial_\alpha) - 2(\eta_{\mu\nu} \partial_\alpha \partial_\beta \right. \right. \\
& + \eta_{\alpha\beta} \partial_\mu \partial_\nu) - (\eta_{\mu\alpha} \eta_{\nu\beta} + \eta_{\mu\beta} \eta_{\nu\alpha} - 2\eta_{\mu\nu} \eta_{\alpha\beta}) \left[ \partial^2 + (3A' + \partial_z) \partial_z \right] \hat{h}^{\alpha\beta} \\
& + \hat{\mathcal{V}}^\mu (\eta_{\mu\nu} \partial^2 - \partial_\mu \partial_\nu) \hat{\mathcal{V}}^\nu + \frac{3}{4} \hat{\phi} \left[ \partial^2 - 2(A' + \partial_z)(2A' + \partial_z) \right] \hat{\phi} \\
& - \frac{1}{2} \left[ \hat{h}^{\mu\nu} (\eta_{\mu\alpha} \partial_\nu + \eta_{\nu\alpha} \partial_\mu - 2\eta_{\mu\nu} \partial_\alpha) (3A' + \partial_z) \hat{\mathcal{V}}^\alpha + \hat{\mathcal{V}}^\mu (\eta_{\mu\alpha} \partial_\beta + \eta_{\mu\beta} \partial_\alpha \right. \\
& - 2\eta_{\alpha\beta} \partial_\mu) \partial_z \hat{h}^{\alpha\beta} \left. \right] + \frac{3}{4} \left[ \hat{h} (3A' + \partial_z) (2A' + \partial_z) \hat{\phi} + \hat{\phi} (A' + \partial_z) \partial_z \hat{h} \right] \\
& \left. - \frac{3}{2} \left[ \hat{\mathcal{V}}^\mu (2A' + \partial_z) \partial_\mu \hat{\phi} + \hat{\phi} (A' + \partial_z) \partial_\mu \hat{\mathcal{V}}^\mu \right] \right\}. \tag{2.42}
\end{aligned}$$

This agrees with the literature [44–47]. To eliminate the mixing terms appearing in Eq.(2.42), we choose the gauge-fixing terms as follows:

$$\hat{\mathcal{L}}_{\text{GF}} = -\frac{e^{3A(z)}}{\xi} (\hat{\mathcal{F}}_\mu^2 + \hat{\mathcal{F}}_5^2), \tag{2.43}$$

where  $\xi$  is a gauge-fixing parameter and the gauge-fixing functions  $\hat{\mathcal{F}}_\mu$  and  $\hat{\mathcal{F}}_5$  take the following form:

$$\hat{\mathcal{F}}_\mu = \partial^\nu \hat{h}_{\mu\nu} - \left(1 - \frac{1}{2\xi}\right) \partial_\mu \hat{h} + \xi (3A' + \partial_z) \hat{\mathcal{V}}_\mu, \tag{2.44a}$$

$$\hat{\mathcal{F}}_5 = \frac{1}{2} \left[ \partial_z \hat{h} - 3\xi (2A' + \partial_z) \hat{\phi} - 2\partial_\mu \hat{\mathcal{V}}^\mu \right]. \tag{2.44b}$$

In the above, we denote the gravitational gauge-fixing parameter by  $\xi$  for notational convenience, but it is independent of the gauge-fixing parameter  $\xi$  for gauge theories as defined in Eq.(2.9).

For the BRST quantization, the 5d gravitational ghost fields  $(\hat{c}^M, \hat{\bar{c}}^M)$  are introduced to compensate the gravitational gauge-fixing term (2.9) in the path integral formulation. The 5d Faddeev-Popov Lagrangian takes the following form:

$$\hat{\mathcal{L}}_{\text{FP}} = e^{3A(z)} \hat{\bar{c}}^M \hat{\mathbf{s}} \hat{\mathcal{F}}_M, \tag{2.45}$$

where the gauge-fixing functions,  $\hat{\mathcal{F}}_M = (\hat{\mathcal{F}}_\mu, \hat{\mathcal{F}}_5)$ , are defined as in Eq.(2.44), with the 5d Lorentz index  $M = \mu, 5$ .

Under the  $S^1/\mathbb{Z}_2$  orbifold compactification, the tensor component of graviton field  $\hat{h}_{\mu\nu}$  and the scalar component  $\hat{\phi}$  are assigned to be even under the  $\mathbb{Z}_2$  parity, whereas the vector component  $\hat{\mathcal{V}}_\mu$  is  $\mathbb{Z}_2$  odd:

$$\hat{h}_{\mu\nu}(x, z) = \hat{h}_{\mu\nu}(x, -z), \quad \hat{\mathcal{V}}_\mu(x, z) = -\hat{\mathcal{V}}_\mu(x, -z), \quad \hat{\phi}(x, z) = \hat{\phi}(x, -z). \tag{2.46}$$

We find that the ghost fields  $(\hat{c}^\mu, \hat{c}^5)$  obey the same equations of motion as that of the gravitational fields  $(\hat{h}^{\mu\nu}, \hat{\mathcal{V}}^\mu)$  respectively. Thus, we assign the following  $\mathbb{Z}_2$  parities for the ghost fields:

$$\hat{c}^\mu(x, z) = \hat{c}^\mu(x, -z), \quad \hat{c}^5(x, z) = -\hat{c}^5(x, -z), \tag{2.47}$$

and their corresponding anti-ghost fields share the same  $\mathbb{Z}_2$  parities.

With these, we make the following KK expansions for the 5d graviton fields in terms of Fourier series containing the zero-modes and KK-modes:

$$\hat{h}^{\mu\nu}(x, z) = \frac{1}{\sqrt{L}} \sum_{n=0}^{\infty} h_n^{\mu\nu}(x) \mathbf{u}_n(z), \quad (2.48a)$$

$$\hat{\mathcal{V}}^\mu(x, z) = \frac{1}{\sqrt{L}} \sum_{n=1}^{\infty} \mathcal{V}_n^\mu(x) \mathbf{v}_n(z), \quad (2.48b)$$

$$\hat{\phi}(x, z) = \frac{1}{\sqrt{L}} \sum_{n=0}^{\infty} \phi_n(x) \mathbf{w}_n(z). \quad (2.48c)$$

Using the above KK decompositions, we derive the following equations of motion (EOM) for the 5d gravitational KK wavefunctions:<sup>3</sup>

$$(3A' + \partial_z) \partial_z \mathbf{u}_n(z) = -\mathbb{M}_n^2 \mathbf{u}_n(z), \quad (2.49a)$$

$$\partial_z (3A' + \partial_z) \mathbf{v}_n(z) = -\mathbb{M}_n^2 \mathbf{v}_n(z), \quad (2.49b)$$

$$(2A' + \partial_z)(A' + \partial_z) \mathbf{v}_n(z) = -\mathbb{M}_n^2 \mathbf{v}_n(z), \quad (2.49c)$$

$$(A' + \partial_z)(2A' + \partial_z) \mathbf{w}_n(z) = -\mathbb{M}_n^2 \mathbf{w}_n(z). \quad (2.49d)$$

In the above and hereafter, we use the notation  $\mathbb{M}_n$  to denote the mass eigenvalue of KK gravitons at level- $n$ , which differs from the KK gauge boson mass eigenvalue  $M_n$  of level- $n$  and is in contrast to the case of compactified flat 5d extra dimension. In addition, Eqs.(2.49b) and (2.49c) are actually equivalent and we give both formulas for convenience of the later analysis. The 5d wavefunctions  $\{\mathbf{u}_n(z), \mathbf{v}_n(z), \mathbf{w}_n(z)\}$  obey the following boundary conditions,

$$\partial_z \mathbf{u}_n(z)|_{z=0,L} = 0, \quad \mathbf{v}_n(z)|_{z=0,L} = 0, \quad (2A' + \partial_z) \mathbf{w}_n(z)|_{z=0,L} = 0, \quad (2.50)$$

and the orthonormal conditions,

$$\frac{1}{L} \int_0^L dz e^{3A(z)} \chi_n(z) \chi_m(z) = \delta_{nm}, \quad (2.51)$$

where we denote  $\chi_n = \{\mathbf{u}_n, \mathbf{v}_n, \mathbf{w}_n\}$ . The 5d wavefunctions  $\mathbf{u}_n(z)$  and  $\mathbf{v}_n(z)$  are connected to each other by the equations:

$$\partial_z \mathbf{u}_n(z) = -\mathbb{M}_n \mathbf{v}_n(z), \quad (2.52a)$$

$$(3A' + \partial_z) \mathbf{v}_n(z) = \mathbb{M}_n \mathbf{u}_n(z), \quad (2.52b)$$

whereas the 5d wavefunctions  $\mathbf{w}_n(z)$  and  $\mathbf{v}_n(z)$  are connected by the equations,

$$(2A' + \partial_z) \mathbf{w}_n(z) = -\mathbb{M}_n \mathbf{v}_n(z), \quad (2.53a)$$

$$(A' + \partial_z) \mathbf{v}_n(z) = \mathbb{M}_n \mathbf{w}_n(z). \quad (2.53b)$$

---

<sup>3</sup>Note that our analyses below and elsewhere in this work are self-contained and well understood. Thus it does not invoke any additional symmetries such as SUSY and alike.

Moreover, by using Eqs.(2.52b) and (2.53b), we obtain the following identity connecting the three types of wavefunctions:

$$\mathbf{u}_n(z) = \mathbf{w}_n(z) + \frac{2A'}{\mathbb{M}_n} \mathbf{v}_n(z). \quad (2.54)$$

In passing, the relation of Eq.(2.54) to the quantum-mechanical SUSY algebra was discussed in the literature [48].

By solving the equation of motion (2.49), the 5d wavefunctions  $\mathbf{u}_n(z)$ ,  $\mathbf{v}_n(z)$  and  $\mathbf{w}_n(z)$  can be expressed in terms of the Bessel functions of order  $\nu = 2, 1, 0$  respectively:

$$\mathbf{u}_n(z) = \frac{e^{-2A(z)}}{N'_n} \left[ J_2 \left( \frac{\mathbb{M}_n}{k} e^{-A(z)} \right) + b_{n1} Y_2 \left( \frac{\mathbb{M}_n}{k} e^{-A(z)} \right) \right], \quad (2.55a)$$

$$\mathbf{v}_n(z) = \frac{e^{-2A(z)}}{N'_n} \left[ J_1 \left( \frac{\mathbb{M}_n}{k} e^{-A(z)} \right) + b_{n1} Y_1 \left( \frac{\mathbb{M}_n}{k} e^{-A(z)} \right) \right], \quad (2.55b)$$

$$\mathbf{w}_n(z) = \frac{e^{-2A(z)}}{N'_n} \left[ J_0 \left( \frac{\mathbb{M}_n}{k} e^{-A(z)} \right) + b_{n1} Y_0 \left( \frac{\mathbb{M}_n}{k} e^{-A(z)} \right) \right], \quad (2.55c)$$

where  $N'_n$  is the normalization factor and  $b_{n1}$  is given by

$$b_{n1} = -\frac{J_1 \left( \frac{\mathbb{M}_n}{k} \right)}{Y_1 \left( \frac{\mathbb{M}_n}{k} \right)}. \quad (2.56)$$

The KK mass eigenvalue  $\mathbb{M}_n$  is determined by solving the following equation:

$$J_1 \left( \frac{\mathbb{M}_n}{k} e^{-A(L)} \right) + b_{n1} Y_1 \left( \frac{\mathbb{M}_n}{k} e^{-A(L)} \right) = 0. \quad (2.57)$$

We note that the KK graviton mass  $\mathbb{M}_n$  solved from the above eigenvalue equation (2.57) differs from the KK gauge boson mass  $M_n$  solved from Eq.(2.20). Then, the zero-mode wavefunctions  $\mathbf{u}_0(z)$  and  $\mathbf{w}_0(z)$  can be solved as follows:

$$\mathbf{u}_0 = \sqrt{2} [e^{A(L)} + e^{2A(L)}]^{-\frac{1}{2}}, \quad (2.58a)$$

$$\mathbf{w}_0(z) = \sqrt{2} e^{-2A(z)} [1 + e^{-A(L)}]^{-\frac{1}{2}}, \quad (2.58b)$$

where  $\mathbf{u}_0$  has no dependence on  $z$ .

As shown in Eqs.(2.46) and (2.47), the 5d ghost fields ( $\hat{c}^\mu, \hat{c}^5$ ) share the same  $\mathbb{Z}_2$  parities as that of the gravitational fields ( $\hat{h}^{\mu\nu}, \hat{\nu}^\mu$ ) respectively. So we make the following KK expansions for the gravitational ghost and anti-ghost fields,

$$\hat{c}^\mu(x, z) = \frac{1}{\sqrt{L}} \sum_{n=0}^{\infty} c_n^\mu(x) \mathbf{u}_n(z), \quad \hat{\bar{c}}^\mu(x, z) = \frac{1}{\sqrt{L}} \sum_{n=0}^{\infty} \bar{c}_n^\mu(x) \mathbf{u}_n(z), \quad (2.59a)$$

$$\hat{c}^5(x, z) = \frac{1}{\sqrt{L}} \sum_{n=1}^{\infty} c_n^5(x) \mathbf{v}_n(z), \quad \hat{\bar{c}}^5(x, z) = \frac{1}{\sqrt{L}} \sum_{n=1}^{\infty} \bar{c}_n^5(x) \mathbf{v}_n(z). \quad (2.59b)$$



We can further derive the BRST transformations for the KK fields, which are given in Appendix B.

Making KK expansions for the 5d Lagrangian (2.42) and integrating  $z$  over the interval  $[0, L]$ , we derive the 4d KK Lagrangian at quadratic order:

$$\begin{aligned} \mathcal{L}_{\text{RS}}^{(2)} = \sum_{n=0}^{\infty} \left[ \frac{1}{2} (\partial_{\mu} h_n)^2 + \frac{1}{2} \mathbb{M}_n^2 (h_n)^2 - \frac{1}{2} (\partial^{\rho} h_n^{\mu\nu})^2 - \frac{1}{2} \mathbb{M}_n^2 (h_n^{\mu\nu})^2 - (\partial^{\mu} \mathcal{V}_n^{\nu})^2 \right. \\ \left. + (\partial_{\mu} \mathcal{V}_n^{\mu})^2 - \frac{3}{4} (\partial_{\mu} \phi_n)^2 + \frac{3}{2} \mathbb{M}_n^2 \phi_n^2 - \partial_{\mu} h_n^{\mu\nu} \partial_{\nu} h_n + \partial_{\mu} h_n^{\mu\rho} \partial^{\nu} h_{\nu\rho,n} \right. \\ \left. + 2\mathbb{M}_n h_n \partial_{\mu} \mathcal{V}_n^{\mu} - 2\mathbb{M}_n h_n^{\mu\nu} \partial_{\mu} \mathcal{V}_{\nu,n} - \frac{3}{2} \mathbb{M}_n^2 h_n \phi_n - 3\mathbb{M}_n \partial_{\mu} \mathcal{V}_n^{\mu} \phi_n \right]. \end{aligned} \quad (2.60)$$

We further derive the 4d  $R_{\xi}$ -type gauge-fixing terms under the KK expansions:

$$\mathcal{L}_{\text{GF}} = \int_0^L dz \hat{\mathcal{L}}_{\text{GF}} = - \sum_{n=0}^{\infty} \frac{1}{\xi_n} [(\mathcal{F}_n^{\mu})^2 + (\mathcal{F}_n^5)^2], \quad (2.61)$$

where  $\xi_n$  is the gauge-fixing parameter for KK level- $n$ , and the gauge-fixing functions  $\mathcal{F}_n^{\mu}$  and  $\mathcal{F}_n^5$  take the following forms:

$$\mathcal{F}_n^{\mu} = \partial_{\nu} h_n^{\mu\nu} - \left(1 - \frac{1}{2\xi_n}\right) \partial^{\mu} h_n + \xi_n \mathbb{M}_n \mathcal{V}_n^{\mu}, \quad (2.62a)$$

$$\mathcal{F}_n^5 = \frac{1}{2} (\mathbb{M}_n h_n - 3\xi_n \mathbb{M}_n \phi_n + 2\partial_{\mu} \mathcal{V}_n^{\mu}). \quad (2.62b)$$

They take the same forms as what we obtained for the flat 5d compactification [15][16]. The above gauge-fixing terms are designed to cancel the mixing terms among the KK graviton fields  $h_n^{\mu\nu}$  and the KK gravitational Goldstone fields  $\mathcal{V}_n^{\mu}$  and  $\phi_n$ . To ensure proper normalization of the kinetic terms of  $\mathcal{V}_n^{\mu}$  and  $\phi_n$ , we make the following rescaling,

$$\mathcal{V}_n^{\mu} \rightarrow \frac{1}{\sqrt{2}} \mathcal{V}_n^{\mu}, \quad \phi_n \rightarrow \sqrt{\frac{2}{3}} \phi_n, \quad (2.63)$$

under which the gauge-fixing functions (2.62a)-(2.62b) become:

$$\mathcal{F}_n^{\mu} = \partial_{\nu} h_n^{\mu\nu} - \left(1 - \frac{1}{2\xi_n}\right) \partial^{\mu} h_n + \frac{1}{\sqrt{2}} \xi_n \mathbb{M}_n \mathcal{V}_n^{\mu}, \quad (2.64a)$$

$$\mathcal{F}_n^5 = \frac{1}{2} \mathbb{M}_n h_n - \sqrt{\frac{3}{2}} \xi_n \mathbb{M}_n \phi_n + \frac{1}{\sqrt{2}} \partial_{\mu} \mathcal{V}_n^{\mu}. \quad (2.64b)$$

Thus, we derive the propagators of KK fields ( $h_n^{\mu\nu}$ ,  $\mathcal{V}_n^{\mu}$ ,  $\phi_n$ ) at tree level (as in [15]) which take the simple diagonal forms in the Feynman-'t Hooft gauge ( $\xi_n = 1$ ):

$$\mathcal{D}_{h,nm}^{\mu\nu\alpha\beta}(k) = -\frac{i\delta_{nm}}{2} \frac{\eta^{\mu\alpha}\eta^{\nu\beta} + \eta^{\mu\beta}\eta^{\nu\alpha} - \eta^{\mu\nu}\eta^{\alpha\beta}}{k^2 + \mathbb{M}_n^2}, \quad (2.65a)$$

$$\mathcal{D}_{\mathcal{V},nm}^{\mu\nu}(k) = -\frac{i\eta^{\mu\nu}\delta_{nm}}{k^2 + \mathbb{M}_n^2}, \quad (2.65b)$$

$$\mathcal{D}_{\phi,nm}(k) = -\frac{i\delta_{nm}}{k^2 + \mathbb{M}_n^2}. \quad (2.65c)$$

We will present the BRST transformations in Appendix B.

Then, we derive a general (Slavnov-Taylor-type) identity in the momentum space for the gravitational gauge-fixing functions (2.64a)-(2.64b) as external lines of the Green function [15][16]:

$$\langle 0 | \mathcal{F}_{n_1}^{\mu_1}(k_1) \mathcal{F}_{n_2}^{\mu_2}(k_2) \cdots \mathcal{F}_{m_1}^5(p_1) \mathcal{F}_{m_2}^5(p_2) \cdots \Phi | 0 \rangle = 0, \quad (2.66)$$

where each external momentum ( $k_j$  or  $p_j$ ) is chosen as incoming and  $\Phi$  denotes any possible external physical states after the LSZ reduction.

### 2.3.2 Formulating GRET for Warped KK Gravity Theory (I)

In this subsection, we first formulate a KK Gravitational Equivalence Theorem (GRET of type-II)<sup>4</sup> to quantitatively connect the scattering amplitude of KK gravitons  $h_n^{\mu\nu}$  with helicity  $\pm 1$  to that of the corresponding gravitational KK vector Goldstone bosons  $\mathcal{V}_n^\mu$ . We derived the general identity (2.66) for the gravitational gauge-fixing functions  $\mathcal{F}_n^\mu$  and  $\mathcal{F}_n^5$  of Eq.(2.64).

We reexpress the gauge-fixing function (2.64a) in matrix form:

$$\mathcal{F}_n^\mu = -\frac{i\mathbb{M}_n}{\sqrt{2}} \mathbb{F}_n^\mu, \quad \mathbb{F}_n^\mu = \mathbf{K}_n^T \mathbf{H}_n, \quad (2.67a)$$

$$\mathbf{K}_n = \left( \frac{1}{\sqrt{2}\mathbb{M}_n} [2k_\beta \eta_\alpha{}^\mu - k^\mu (2 - \xi_n^{-1}) \eta_{\alpha\beta}], i\xi_n \right)^T, \quad \mathbf{H}_n = (h_n^{\alpha\beta}, \mathcal{V}_n^\mu)^T. \quad (2.67b)$$

Then, we derive the following BRST identity including the gauge-fixing function  $\mathbb{F}_n^\mu$ ,

$$\langle 0 | T \mathbb{F}_n^\mu \mathbf{H}_m^T | 0 \rangle = -\frac{i\xi_n}{\sqrt{2}\mathbb{M}_n} \langle 0 | T \widehat{\mathbf{S}} \mathbf{H}_m^T \bar{c}_n^\mu | 0 \rangle. \quad (2.68)$$

Using Eq.(2.67), we can rewrite Eq.(2.68) as follows:

$$\mathbf{K}_n^T \mathcal{D}_{nm}(k) = -\mathbf{X}_{nm}^T(k), \quad (2.69)$$

where we have defined the notations,

$$\begin{aligned} \mathcal{D}_{nm}(k) &= \langle 0 | T \mathbf{H}_n \mathbf{H}_m^T | 0 \rangle(k), \\ \mathbf{X}_{nm}(k) &= \underline{\mathbf{X}}_{mj}(k) \mathcal{S}_{jn}(k), \quad \mathcal{S}_{jn}(k) \eta^{\lambda\mu} = \langle 0 | T c_j^\lambda \bar{c}_n^\mu | 0 \rangle(k), \\ \underline{\mathbf{X}}_{mj}(k) &= \frac{i\eta^{\lambda\mu}}{\sqrt{2}\mathbb{M}_j} \left( \langle 0 | T \widehat{\mathbf{S}} h_m^{\sigma\rho} | \bar{c}_{j,\lambda} \rangle(k) \right) = \frac{1}{\sqrt{2}\mathbb{M}_j} \begin{pmatrix} (k^\sigma \eta^{\rho\mu} + k^\rho \eta^{\sigma\mu}) [\delta_{mj} + \Delta_{mj}^{(1)}(k^2)] \\ i\sqrt{2}\mathbb{M}_m \eta^{\mu\nu} [\delta_{mj} + \tilde{\Delta}_{mj}^{(2)}(k^2)] \end{pmatrix}, \end{aligned} \quad (2.70)$$

where  $\Delta_{mj}^{(1)}(k^2)$  and  $\tilde{\Delta}_{mj}^{(2)}(k^2)$  are the loop-level quantities, and are further defined in Eq.(B.25) of Appendix B. In the above, we have chosen the Feynman-'t Hooft gauge  $\xi_n = 1$

<sup>4</sup>For convenience, we call this GRET (for KK gravitons with helicity  $\pm 1$ ) the GRET of type-II, and call the other GRET (for KK gravitons with helicity-0) the GRET of type-I as will be given in Section 2.3.3.

for simplicity of the  $\underline{\mathbf{X}}_{mj}$  formula. The case of general  $R_\xi$  gauge will be discussed in Appendix B.

Then, we analyze the identity (2.66) and construct an external state as a combination of gauge-fixing functions ( $\mathcal{F}_n^\mu$  or  $\mathbb{F}_n^\mu$ ) as given in Eq.(2.67). Thus, we amputate this external state by using the BRST identity (2.69) and obtain:

$$0 = \langle 0 | \mathbb{F}_n^\mu(k) \cdots \Phi | 0 \rangle = -\mathbf{X}_{nm}^T(k) \mathcal{M}[\underline{\mathbf{H}}_m(k), \cdots, \Phi]. \quad (2.71)$$

From the above, we can derive the following identity:

$$\mathcal{M}[\underline{\mathbb{F}}_n^\mu(k), \cdots, \Phi] = 0, \quad (2.72)$$

where  $\mathcal{M}[\cdots]$  denotes the amputated scattering amplitude. In Eq.(2.72), the amputated external state  $\underline{\mathbb{F}}_n^\mu$  takes the following form (after removing an irrelevant overall coefficient):

$$\underline{\mathbb{F}}_n^\mu(k) = \frac{\sqrt{2}}{\mathbb{M}_n} k_\nu h_n^{\mu\nu} - \hat{C}^{nm} \eta^{\mu\nu} \mathcal{V}_{m,\nu}, \quad (2.73)$$

where the loop-induced modification factor is derived as follows:

$$\hat{C}_{nm}(k^2) = -\frac{i\mathbb{M}_m}{\mathbb{M}_n} \left[ \frac{\mathbf{1} + \tilde{\Delta}^{(2)}(k^2)}{\mathbf{1} + \Delta^{(1)}(k^2)} \right]_{mn} = -i\delta_{nm} + O(\text{loop}). \quad (2.74)$$

In the above, the matrix form is used such that the loop-level quantities  $(\Delta^{(1)})_{jj'} = \Delta_{jj'}^{(1)}$  and  $(\tilde{\Delta}^{(2)})_{jj'} = \tilde{\Delta}_{jj'}^{(2)}$ . The derivations of Eqs.(2.71)-(2.74) are given in more detail around Eqs.(B.26)-(B.27) of Appendix B.

Then, we utilize a transverse polarization vector  $\epsilon_\pm^\mu$  to contract with  $\underline{\mathbb{F}}_n^\mu(k)$  in Eq.(2.73) and obtain the formula:

$$\underline{\mathbb{F}}_n(k) \equiv \epsilon_\mu^\pm \underline{\mathbb{F}}_n^\mu(k) = h_n^{S,\pm 1} - \hat{C}_{nm} \mathcal{V}_m^{\pm 1}, \quad (2.75)$$

where we have used the fact  $\epsilon_\pm^\mu k_\mu = 0$ , and defined the external states  $h_n^{S,\pm 1}$  and  $\mathcal{V}_m^{\pm 1}$  as follows:

$$h_n^{S,\pm 1} = \varepsilon_{\mu\nu}^{S,\pm 1} h_n^{\mu\nu}, \quad \mathcal{V}_m^{\pm 1} = \epsilon_\mu^\pm \mathcal{V}_m^\mu, \quad \varepsilon_{\mu\nu}^{S,\pm 1} = \frac{1}{\sqrt{2}} (\epsilon_\mu^S \epsilon_\nu^\pm + \epsilon_\nu^S \epsilon_\mu^\pm), \quad (2.76)$$

with  $\epsilon_\mu^S = k_\mu / \mathbb{M}_n$  being the scalar-polarization vector. A massive spin-2 KK graviton has 5 physical helicity states ( $\lambda = \pm 2, \pm 1, 0$ ). The polarization tensors of KK graviton with helicities  $\pm 1$  are given by

$$\varepsilon_{\pm 1}^{\mu\nu} = \frac{1}{\sqrt{2}} (\epsilon_\pm^\mu \epsilon_L^\nu + \epsilon_L^\mu \epsilon_\pm^\nu), \quad (2.77)$$

where  $(\epsilon_\pm^\mu, \epsilon_L^\mu)$  denote the 3 polarization vectors of a massive spin-1 KK gauge boson. Under high energy expansion, we have  $\epsilon_L^\mu = \epsilon_S^\mu + v^\mu$  with  $\epsilon_S^\mu = k^\mu / \mathbb{M}_n$  and  $v^\mu = O(\mathbb{M}_n / E_n)$ . Thus, we can decompose the polarization tensor  $\varepsilon_{\pm 1}^{\mu\nu}$  as follows:

$$\varepsilon_{\pm 1}^{\mu\nu} = \varepsilon_{S,\pm 1}^{\mu\nu} + v_{\pm 1}^{\mu\nu}, \quad \varepsilon_{S,\pm 1}^{\mu\nu} = \frac{1}{\sqrt{2}} (\epsilon_S^\mu \epsilon_\pm^\nu + \epsilon_S^\nu \epsilon_\pm^\mu), \quad (2.78a)$$

$$v_{\pm 1}^{\mu\nu} = \frac{1}{\sqrt{2}} (v^\mu \epsilon_\pm^\nu + v^\nu \epsilon_\pm^\mu) = O(\mathbb{M}_n / E). \quad (2.78b)$$

With these, we can reexpress the function  $\mathbb{F}_n(k)$  of Eq.(2.75) in the form:

$$\mathbb{F}_n(k) = h_n^{\pm 1} - \Theta_n, \quad \Theta_n = \hat{C}^{nm} \mathcal{V}_m^{\pm 1} + v_n^{\pm 1}, \quad (2.79)$$

where we have used the notations,

$$h_n^{\pm 1} = \varepsilon_{\mu\nu}^{\pm 1} h_n^{\mu\nu}, \quad \mathcal{V}_m^{\pm 1} = \epsilon_\mu^{\pm 1} \mathcal{V}_m^\mu, \quad v_n^{\pm 1} = v_{\mu\nu}^{\pm 1} h_n^{\mu\nu}, \quad (2.80)$$

with the tensor  $v_{\pm 1}^{\mu\nu} = O(\mathbb{M}_n/E_n)$  under high energy expansion and thus  $v_n^{\pm 1}$  is suppressed by an energy factor of  $O(\mathbb{M}_n/E_n)$ . For later analysis, we can also multiply  $v^\mu = \epsilon_L^\mu - \epsilon_S^\mu$  on both sides of Eq.(2.72) and use the expression (2.73) to obtain the following identity:

$$\epsilon_\mu^S v_\nu \mathcal{M}[h_n^{\mu\nu}(k), \dots, \Phi] = \hat{C}_{nm} v_\mu \mathcal{M}[\mathcal{V}_m^\mu(k), \dots, \Phi], \quad (2.81)$$

where we see that the right-hand side is suppressed by the external factor  $v_\mu = O(\mathbb{M}_n/E_n)$ . Hence the above identity (2.81) ensures that its left-hand side must receive a suppression such that its overall energy-power dependence is the same as the right-hand side [despite that the external factor of the left-hand side is  $\epsilon_\mu^S v_\nu = O(E_n^0)$ ].

Then, we reformulate the identity (2.72) and extend it to the general case of  $N$  external amputated gauge-fixing functions:

$$\mathcal{M}[\mathbb{F}_{n_1}(k_1), \dots, \mathbb{F}_{n_N}(k_N), \Phi] = 0. \quad (2.82)$$

Using this identity, we can further derive the gravitational equivalence theorem (GRET) identity which connects the  $h_n^{\pm 1}$  amplitude to the amplitude of  $\Theta_n$ :

$$\mathcal{M}[h_{n_1}^{\pm 1}(k_1), \dots, h_{n_N}^{\pm 1}(k_N), \Phi] = \mathcal{M}[\Theta_{n_1}(k_1), \dots, \Theta_{n_N}(k_N), \Phi]. \quad (2.83)$$

The above identity can be proven by computing the amplitude on its right-hand side and using the relation  $\Theta_n = h_n^{\pm 1} - \mathbb{F}_n$  in Eq.(2.79). Expanding  $\Theta_n (= \hat{C}_{nm} \mathcal{V}_m^{\pm 1} + v_n^{\pm 1})$  for each external state on the right-hand side of the identity (2.83), we derive the following gravitational equivalence theorem (GRET) identity:

$$\mathcal{M}[h_{n_1}^{\pm 1}, \dots, h_{n_N}^{\pm 1}, \Phi] = \hat{C}_{\text{mod}}^{n_j m_j} \mathcal{M}[\mathcal{V}_{m_1}^{\pm 1}, \dots, \mathcal{V}_{m_N}^{\pm 1}, \Phi] + \mathcal{M}_v, \quad (2.84a)$$

$$\mathcal{M}_v = \sum_{\ell=1}^N \tilde{C}_{\text{mod},\ell}^{n_j m_j} \mathcal{M}[v_{n_1}^{\pm 1}, \dots, v_{n_\ell}^{\pm 1}, \mathcal{V}_{m_{\ell+1}}^{\pm 1}, \dots, \mathcal{V}_{m_N}^{\pm 1}, \Phi], \quad (2.84b)$$

where the multiplicative modification factors are given by

$$\hat{C}_{\text{mod}}^{n_j m_j} = \hat{C}_{n_1 m_1} \cdots \hat{C}_{n_N m_N} = (-i)^N \delta_{n_1 m_1} \cdots \delta_{n_N m_N} + O(\text{loop}), \quad (2.85a)$$

$$\tilde{C}_{\text{mod},\ell}^{n_j m_j} = \hat{C}_{n_{\ell+1} m_{\ell+1}} \cdots \hat{C}_{n_N m_N} = (-i)^{N-\ell} \delta_{n_{\ell+1} m_{\ell+1}} \cdots \delta_{n_N m_N} + O(\text{loop}). \quad (2.85b)$$

Under the high energy expansion, the residual term is suppressed by the energy factor of  $O(\mathbb{M}_n/E)$ . Hence, from the identity (2.84a) we derive the following warped GRET of type-II:

$$\mathcal{M}[h_{n_1}^{\pm 1}, \dots, h_{n_N}^{\pm 1}, \Phi] = \hat{C}_{\text{mod}}^{n_j m_j} \mathcal{M}[\mathcal{V}_{m_1}^{\pm 1}, \dots, \mathcal{V}_{m_N}^{\pm 1}, \Phi] + O(\mathbb{M}_n/E_n\text{-suppressed}). \quad (2.86)$$

This connects the scattering amplitude of the KK gravitons ( $h_n^{\pm 1}$ ) with helicity  $\pm 1$  to the corresponding gravitational KK vector-Goldstone bosons ( $\mathcal{V}_m^{\pm 1}$ ).

### 2.3.3 Formulating GRET for Warped KK Gravity Theory (II)

In this subsection, we further derive the KK GRET to connect the scattering amplitude of helicity-0 longitudinal KK gravitons ( $h_L^n$ ) to the corresponding gravitational KK scalar Goldstone bosons ( $\phi_n$ ) under the high energy expansion. This GRET will be called GRET of type-I.

We use the combination of gauge-fixing functions  $\partial_\mu \mathcal{F}_n^\mu - \xi_n \mathbb{M}_n \mathcal{F}_n^5$  to eliminate the KK vector-Goldstone field  $\mathcal{V}_n^\mu$  and derive an expression in momentum space:

$$-ik_\mu \mathcal{F}_n^\mu - \xi_n \mathbb{M}_n \mathcal{F}_n^5 = -k_\mu k_\nu h_n^{\mu\nu} + \frac{1}{2} \left[ (2 - \xi_n^{-1}) k^2 - \xi_n \mathbb{M}_n^2 \right] h_n + \sqrt{\frac{3}{2}} \xi_n^2 \mathbb{M}_n^2 \phi_n, \quad (2.87)$$

where we have made the rescaling of  $\phi_n$  as in Eq.(2.63). Imposing the on-shell condition  $k^2 = -\mathbb{M}_n^2$ , we can rewrite the above formula as follows:

$$ik_\mu \mathcal{F}_n^\mu + \xi_n \mathbb{M}_n \mathcal{F}_n^5 = \sqrt{\frac{3}{2}} \mathbb{M}_n^2 \tilde{\mathbb{F}}_n, \quad (2.88a)$$

$$\tilde{\mathbb{F}}_n = \sqrt{\frac{2}{3}} h_n^S + \frac{1}{\sqrt{6}} (2 + \xi_n - \xi_n^{-1}) h_n - \xi_n^2 \phi_n = \mathbf{K}_n^T \mathbf{H}_n \quad (2.88b)$$

$$= \sqrt{\frac{2}{3}} (h_n^S + h_n) - \phi_n, \quad (\text{for } \xi_n = 1), \quad (2.88c)$$

$$\mathbf{K}_n = \left( \sqrt{\frac{2}{3}} \varepsilon_{\mu\nu}^S + \frac{1}{\sqrt{6}} (2 + \xi_n - \xi_n^{-1}) \eta_{\mu\nu}, -\xi_n^2 \right)^T, \quad \mathbf{H}_n = (h_n^{\mu\nu}, \phi_n)^T, \quad (2.88d)$$

where we have defined the scalar KK graviton  $h_n^S = \varepsilon_{\mu\nu}^S h_n^{\mu\nu}$  with its scalar-polarization tensor  $\varepsilon_{\mu\nu}^S = k_\mu k_\nu / \mathbb{M}_n^2$ . Then, we inspect the KK graviton's longitudinal polarization tensor  $\varepsilon_L^{\mu\nu}$  and make high energy expansion of it as follows:

$$\varepsilon_L^{\mu\nu} = \frac{1}{\sqrt{6}} (\epsilon_+^\mu \epsilon_-^\nu + \epsilon_+^\nu \epsilon_-^\mu + 2\epsilon_L^\mu \epsilon_L^\nu) = \sqrt{\frac{2}{3}} \varepsilon_S^{\mu\nu} + \tilde{v}^{\mu\nu}, \quad (2.89)$$

where the longitudinal polarization vector  $\epsilon_L^\mu = (k^0 / \mathbb{M}_n) (|\vec{k}| / k^0, \vec{k} / |\vec{k}|) = \epsilon_S^\mu + v^\mu$  with  $\epsilon_S^\mu = k^\mu / \mathbb{M}_n$  and  $v^\mu = \mathcal{O}(\mathbb{M}_n / E_n)$ . In the above, the residual term

$$\tilde{v}^{\mu\nu} \equiv \varepsilon_L^{\mu\nu} - \sqrt{\frac{2}{3}} \varepsilon_S^{\mu\nu} = \frac{1}{\sqrt{6}} [(\epsilon_+^\mu \epsilon_-^\nu + \epsilon_+^\nu \epsilon_-^\mu) + 2(\epsilon_S^\mu v^\nu + \epsilon_S^\nu v^\mu + v^\mu v^\nu)], \quad (2.90)$$

where  $\epsilon_S^\mu = k^\mu / \mathbb{M}_n$  and  $\tilde{v}^{\mu\nu} = \mathcal{O}(E_n^0)$  under high energy expansion.

According to Appendix B, we derive a BRST identity on the full propagator of the matrix field  $\mathbf{H}_n = (h_n^{\mu\nu}, \phi_n)^T$ ,

$$\mathbf{K}_n^T \mathcal{D}_{nm}(k) = -\mathbf{X}_{nm}^T(k), \quad (2.91)$$

where we have defined the following notations:

$$\begin{aligned} \mathcal{D}_{nm}(k) &= \langle 0 | T \mathbf{H}_n \mathbf{H}_m^T | 0 \rangle(k), \quad \mathbf{X}_{nm}(k) = \underline{\mathbf{X}}_{mj}(k) \mathcal{S}_{jn}(k), \\ \underline{\mathbf{X}}_{mj}(k) &= \frac{1}{2\sqrt{6} \mathbb{M}_j^2} \left( \langle 0 | T \hat{\mathbf{s}} h_m^{\mu\nu} | \bar{c}_j \rangle(k) \right) = \frac{1}{2\sqrt{6}} \left( \begin{array}{c} (2k^\mu k^\nu / \mathbb{M}_j^2 - \eta^{\mu\nu}) [\delta_{mj} + \Delta_{mj}^{(3)}(k^2)] \\ -\sqrt{6} [\delta_{mj} + \tilde{\Delta}_{mj}^{(4)}(k^2)] \end{array} \right), \\ \mathcal{S}_{jn}(k) &= \langle 0 | T c_j \bar{c}_n | 0 \rangle(k), \end{aligned} \quad (2.92)$$

with  $c_n \equiv i\epsilon_\mu^S c_n^\mu + c_n^5$ ,  $\bar{c}_n \equiv i\epsilon_\mu^S \bar{c}_n^\mu + \bar{c}_n^5$ , and  $\epsilon_\mu^S = k_\mu/M_n$ . We also note that the content of Eq.(2.91) actually differs from that of Eq.(2.69) although they are expressed in the same notations. In the above, we choose Feynman-'t Hooft gauge  $\xi_n=1$  for simplicity, and the quantities  $\Delta_{mj}^{(3)}(k^2)$  and  $\tilde{\Delta}_{mj}^{(4)}(k^2)$  belong to loop level and are generated by the non-linear terms of the BRST transformations (B.4a) and (B.4c) of  $h_n^{\mu\nu}$  and  $h_n^{55}(\equiv \phi_n)$ .

Using the identity (2.66) and Eq.(2.88) together with Eq.(2.91), we can obtain a new identity with external state  $\tilde{\mathbb{F}}_n(k)$ ,

$$0 = \langle 0 | \tilde{\mathbb{F}}_n(k) \cdots \Phi | 0 \rangle = -\mathbf{X}_{nm}^T(k) \mathcal{M}[\underline{\mathbf{H}}_m(k), \cdots, \Phi], \quad (2.93)$$

from which we derive the following identity:

$$\mathcal{M}[\tilde{\mathbb{F}}_n(k), \cdots, \Phi] = 0, \quad (2.94)$$

where the amputated external state  $\tilde{\mathbb{F}}_n(k)$  is given as follows:

$$\tilde{\mathbb{F}}_n = \sqrt{\frac{2}{3}} (h_n^S - \frac{1}{2} h_n) - C_{nm} \phi_m = h_n^L - \Omega_n, \quad (2.95a)$$

$$\Omega_n = C_{nm} \phi_m + \tilde{\Delta}_n, \quad \tilde{\Delta}_n = \frac{1}{\sqrt{6}} h_n + \tilde{v}_n, \quad \tilde{v}_n = v_{\mu\nu} h_n^{\mu\nu}. \quad (2.95b)$$

In the above,  $C_{nm}$  is a multiplicative modification factor induced at loop level:

$$C_{nm}(k^2) = \left[ \frac{\mathbf{1} + \tilde{\Delta}^{(4)}(k^2)}{\mathbf{1} + \Delta^{(3)}(k^2)} \right]_{mn} \equiv \delta_{nm} + O(\text{loop}), \quad (2.96)$$

where the matrix form is used such that  $(\Delta^{(3)})_{jj'} = \Delta_{jj'}^{(3)}$  and  $(\tilde{\Delta}^{(4)})_{jj'} = \tilde{\Delta}_{jj'}^{(4)}$  with the matrix elements  $(\Delta_{jj'}^{(3)}, \tilde{\Delta}_{jj'}^{(4)})$  from Eq.(2.92). More details of the derivations are given in Appendix B.2.

Then, using Eq.(2.94) we can derive the following general identity for the gravitational equivalence theorem (GRET):

$$\mathcal{M}[\tilde{\mathbb{F}}_{n_1}(k_1), \cdots, \tilde{\mathbb{F}}_{n_N}(k_N), \Phi] = 0. \quad (2.97)$$

From this, we further derive the following GRET identity:

$$\mathcal{M}[h_{n_1}^L(k_1), \cdots, h_{n_N}^L(k_N), \Phi] = \mathcal{M}[\Omega_{n_1}(k_1), \cdots, \Omega_{n_N}(k_N), \Phi], \quad (2.98)$$

which can be proven by computing the amplitude on its right-hand side and using the relation  $\Omega_n = h_n^L - \tilde{\mathbb{F}}_n$  in Eq.(2.95a). Expanding the right-hand side of the GRET identity (2.98), we connect the scattering amplitude of helicity-zero longitudinal KK gravitons ( $h_n^L$ ) to that of the corresponding KK Goldstone bosons ( $\phi_m$ ):

$$\mathcal{M}[h_{n_1}^L, \cdots, h_{n_N}^L, \Phi] = C_{\text{mod}}^{n_j m_j} \mathcal{M}[\phi_{m_1}, \cdots, \phi_{m_N}, \Phi] + \mathcal{M}_\Delta, \quad (2.99a)$$

$$\mathcal{M}_\Delta = \sum_{\ell=1}^N \tilde{C}_{\text{mod},\ell}^{n_j m_j} \mathcal{M}[\tilde{\Delta}_{n_1}, \cdots, \tilde{\Delta}_{n_\ell}, \phi_{m_{\ell+1}}, \cdots, \phi_{m_N}, \Phi], \quad (2.99b)$$

where  $\tilde{\Delta}_n = \frac{1}{\sqrt{6}}h_n + \tilde{v}_n$  with  $\tilde{v}_n = v_{\mu\nu}h_n^{\mu\nu}$ , and  $\Phi$  denotes any other external on-shell physical state(s) after the LSZ reduction. In the above, the multiplicative modification factors are defined as follows:

$$C_{\text{mod}}^{n_j m_j} = C_{n_1 m_1} \cdots C_{n_N m_N} = \delta_{n_1 m_1} \cdots \delta_{n_N m_N} + O(\text{loop}), \quad (2.100\text{a})$$

$$\tilde{C}_{\text{mod},\ell}^{n_j m_j} = C_{n_{\ell+1} m_{\ell+1}} \cdots C_{n_N m_N} = \delta_{n_{\ell+1} m_{\ell+1}} \cdots \delta_{n_N m_N} + O(\text{loop}), \quad (2.100\text{b})$$

which are renormalization-scheme-dependent and can be normalized to one under proper setup of the renormalization scheme. They reduce to one at tree level. We can further justify that in the residual term (2.99b) each external state  $\tilde{\Delta}_n (= \frac{1}{\sqrt{6}}h_n + \tilde{v}_n)$  is suppressed by an energy factor  $\mathbb{M}_n/E_n$  under high energy expansion. For this, using the expression (2.90) and the normalization condition  $\sum_{\sigma} \epsilon_{\sigma}^{\mu}(\epsilon_{\sigma}^{\nu})^* = \eta^{\mu\nu} + k^{\mu}k^{\nu}/\mathbb{M}_n^2$  (with helicity  $\sigma = \pm, L$ ), we derive the following:

$$\tilde{\Delta}_n = \frac{1}{\sqrt{6}}h_n + \tilde{v}_n = \sqrt{\frac{3}{2}}(2\epsilon_{\mu}^S v_{\nu} + v_{\mu}v_{\nu})h_n^{\mu\nu}, \quad (2.101)$$

where  $v_{\mu}v_{\nu} = O(\mathbb{M}_n^2/E_n^2)$  and  $\epsilon_{\mu}^S v_{\nu} = O(E_n^0)$ . The external state of  $(v_{\mu}v_{\nu}h_n^{\mu\nu})$  is clearly suppressed by the extra energy factor  $v_{\mu}v_{\nu} \sim \mathbb{M}_n^2/E_n^2$  in the high energy limit. As proven by Eq.(2.81), despite the external factor  $\epsilon_{\mu}^S v_{\nu} = O(E_n^0)$ , the amplitude with external state  $\epsilon_{\mu}^S v_{\nu}h_n^{\mu\nu}$  is actually suppressed by an overall energy factor  $v^{\mu} = O(\mathbb{M}_n/E_n)$ . Thus, as the external state, we can reexpress  $\tilde{\Delta}_n$  as follows:

$$\tilde{\Delta}_n = \sqrt{\frac{3}{2}}(\hat{C}_{nm}2v_{\mu}\mathcal{V}_m^{\mu} + v_{\mu}v_{\nu}h_n^{\mu\nu}), \quad (2.102)$$

where the two terms on right-hand side is suppressed by external factors of  $v_{\mu} = O(E^{-1})$  and  $v_{\mu}v_{\nu} = O(E^{-2})$ , respectively. Hence, the residual term  $\mathcal{M}_{\Delta}$  in Eq.(2.99b) must be suppressed (relative to the leading KK Goldstone amplitude) at least by an overall energy factor  $v^{\mu} = O(\mathbb{M}_n/E_n)$  under the high energy expansion. Thus, we can further formulate the following warped GRET of type-I:

$$\mathcal{M}[h_{n_1}^L, \cdots, h_{n_N}^L, \Phi] = C_{\text{mod}}^{n_j m_j} \mathcal{M}[\phi_{m_1}, \cdots, \phi_{m_N}, \Phi] + O(\mathbb{M}_n/E_n\text{-suppressed}). \quad (2.103)$$

This ensures that the GRET identity (2.99a) can establish the equivalence between the helicity-zero longitudinal KK graviton amplitude  $\mathcal{M}[h_{n_1}^L, \cdots]$  and the corresponding gravitational KK Goldstone amplitude  $C_{\text{mod}}^{n_j m_j} \mathcal{M}[\phi_{m_1}, \cdots]$  in the high energy limit. Thus, the identity (2.99) gives in Eq.(2.103) the KK gravitational equivalence theorem (GRET) for the warped 5d GR theory under high energy expansion, which extends our previous formulation of the KK GRET for the compactified flat 5d GR [15, 16].

We note that the generalized power-counting method for the compactified KK gauge theories and KK GR theories as established in Ref. [15] does hold for both flat and warped extra dimensions. Thus, we can deduce the leading energy-power dependence of  $N$ -point

helicity-zero longitudinal KK graviton amplitudes and of the corresponding  $N$ -point KK Goldstone amplitudes:

$$D_E(Nh_n^L) = 2(N+L+1), \quad D_E(N\phi_n) = 2(L+1), \quad (2.104a)$$

$$\implies D_E(Nh_n^L) - D_E(N\phi_n) = 2N, \quad (2.104b)$$

where  $L \geq 0$  denotes the number of loops involved. For the scattering amplitudes of  $N$  longitudinal KK gravitons (with helicity-zero) at tree level ( $L=0$ ), we use the right-hand side of the GRET identity (2.99a) and prove the large energy cancellations of  $E^{2N+2} \rightarrow E^2$  by an energy factor  $E^{2N}$  in the  $N$  longitudinal KK graviton amplitude.

The case of 3-point KK graviton amplitudes needs a further discussion. According to the above power counting rule (2.104a), the tree-level 3-point longitudinal KK graviton amplitude ( $3h_n^L$ ) have a possible leading energy dependence of  $E^{2(N+1)} = E^8$  for  $N=3$ . The tree-level amplitude of 3 gravitational KK Goldstone bosons ( $3\phi_n$ ) arises from trilinear KK Goldstone vertex whose coupling factor contains products of two external momenta  $p_i \cdot p_j$ . So the naive power counting shows that this KK Goldstone amplitude would have leading energy dependence of  $E^2$  in agreement with Eq.(2.104a). Thus, according to the GRET (2.99), the 3-point tree-level longitudinal KK graviton amplitude should have a total energy-power deduction of  $E^{8-2} = E^6$ , which agrees to Eq.(2.104b). But, as we will further explain at the end of the following Section 3.1.2, the  $N=3$  is a special case where the momentum conservation and on-shell condition of each external state plays additional roles to reduce the leading energy dependence. Namely, after imposing the momentum conservation at the trilinear vertex and the on-shell condition for each momentum, the individual leading-energy dependence of 3-point longitudinal KK graviton amplitude reduces to  $E^4$  and that of the corresponding KK Goldstone amplitude becomes  $E^0$ . Hence the GRET (2.99) further ensures a nontrivial energy cancellation of  $E^4 \rightarrow E^0$ .

Then, we can further consider the scattering amplitudes of  $N$  KK gravitons of  $h_n^{\pm 1}$  (with helicity  $\pm 1$ ) and of the corresponding gravitational KK vector-Goldstone bosons ( $\mathcal{V}_n^{\pm 1}$ ). Using the generalized power-counting method [15], we derive their leading energy-power dependence as follows:

$$D_E(Nh_n^{\pm 1}) = N+2(L+1), \quad D_E(N\mathcal{V}_n^{\pm 1}) = 2(L+1), \quad (2.105a)$$

$$\implies D_E(Nh_n^{\pm 1}) - D_E(N\mathcal{V}_n^{\pm 1}) = N. \quad (2.105b)$$

For the scattering amplitudes of  $N$  KK gravitons ( $h_n^{\pm 1}$ ) at tree level, we use the right-hand side of the GRET identity (2.84) [or (2.86)] and prove the large energy cancellations of  $E^{N+2} \rightarrow E^2$  by an energy factor  $E^N$  in the scattering amplitude of  $N$  KK gravitons (with helicity  $\pm 1$ ).

For the warped KK gauge theories as studied in Section 2.2, using the generalized power-counting method [15] and the GAET identity (2.36), we derive their leading energy-power dependence for  $N$  longitudinal as follows:

$$D_E(NA_L^{an}) = 4, \quad D_E(NA_5^{an}) = 4 - N - N_v - \bar{V}_3^{\min}, \quad (2.106a)$$



$$\implies D_E(NA_L^{an}) - D_E(NA_5^{an}) = N + N_v + \bar{V}_3^{\min}, \quad (2.106b)$$

where  $\bar{V}_3^{\min}$  denotes the involved minimal number of non-derivative cubic vertices in the KK Goldstone boson amplitude, with  $\bar{V}_3^{\min} = 0$  (1) for  $N = \text{even}$  (odd). Hence, this proves that the GAET identity (2.36) guarantees nontrivial energy cancellations by an energy power of  $E^N$  (or,  $E^{N+1}$ ) for  $N = \text{even}$  (or,  $N = \text{odd}$ ) in the scattering amplitude of  $N$  longitudinal KK gauge bosons. In the above,  $N_v$  denotes the number of possible external  $v_\mu A_n^{a\mu}$  lines in the KK Goldstone boson amplitude, where  $v_\mu = O(M_n/E)$  in the high energy limit. We find that the above large energy cancellations [as shown in Eqs.(2.104b), (2.105b) and (2.106b)] hold for both the flat and warped 5d KK gravity (gauge) theories. Before ending this discussion, we further comment on the  $N=3$  case for KK gauge boson (KK Goldstone boson) scattering amplitudes. The 3-point longitudinal KK gauge boson amplitude ( $3A_L^{an}$ ) has a leading energy dependence of  $E^4$  by the power counting rule (2.106a), which arises from the trilinear KK gauge boson vertex (containing one external momentum) and the three external longitudinal polarization vectors (each of which scales like  $\epsilon_L^\mu \sim p_j^\mu/M_{n_j}$  in the high energy limit). On the other hand, the corresponding 3-point KK Goldstone amplitude vanishes at tree level due to the absence of cubic vertex  $3A_5^{an}$ . According to the GAET identity (2.36), the leading nonzero contribution on its right-hand side arises from the residual terms with the trilinear vertex having two KK Goldstone bosons and one KK gauge boson (contracting with the external vector  $v^\mu$ ). So we have  $(N, N_v) = (3, 1)$  and  $\bar{V}_3^{\min} = 0$ , and thus this residual term has energy power  $D_E = 4 - 3 - 1 - 0 = 0$ , showing that the right-hand side of the GAET identity (2.36) has leading energy dependence of  $E^0$ . Hence, the total energy reduction of the 3-point longitudinal KK gauge boson amplitude is  $E^4 \rightarrow E^0$ .

We also note that for the Goldberger-Wise (GW) model [50], it further introduces a bulk scalar field  $\hat{S}$  to stabilize the size of the 5d extra dimension of the RS1. Its zero-mode  $\mathcal{S}_0$  can mix with the radion field  $h_0^{55}$ , so that the physical radion after stabilization becomes a mixture of  $\mathcal{S}_0$  and  $h_0^{55}$ , whereas the scalar KK mode  $\mathcal{S}_n$  mixes with the Goldstone KK mode  $h_n^{55}$  (denoted as  $\phi_n$  in our notation) at each KK level, leading to a mass-eigenstate  $\mathcal{S}'_n$  and the true Goldstone KK state  $h_n^{55'}$ . Thus, the gauge-fixing term (2.43)-(2.44) should be extended to contain the true KK Goldstone boson field ( $h_n^{55'}$ ). Consequently, the GRET (2.103) can be extended to hold the equivalence between the scattering amplitudes of the longitudinal KK graviton states ( $h_n^L$ ) and of the true KK Goldstone boson states ( $h_n^{55'}$ ) in the high energy limit.<sup>5</sup>

In passing, during the finalization of this paper, we received a recent paper [21] which studied the tree-level formulation of the gravitational ET for warped 5d RS1 model in the 't Hooft-Feynman gauge and the  $O(s)$  high-energy behavior of the 4-point amplitudes therein.<sup>6</sup> Our general formulations in this section significantly differ from [21] in several aspects. We first derived the new GAET for the warped KK gauge theory in Sec. 2.2, in addition to deriving the general GRET for warped KK gravity theory in Sec. 2.3. The

<sup>5</sup>In passing, Ref. [51] studied spin-2 KK scattering amplitudes with massive radion in the GW model.

<sup>6</sup>We thank the colleagues Sekhar Chivukula and Elizabeth Simmons for discussions on their paper [21] and related issues.

GAET of the warped KK gauge theory is the focus of our present work and serves as the most fundamental KK ET formulation, because we will prove in Sec. 4 that the GRET of the warped KK gravity can be derived (reconstructed) from the warped GAET by our leading-order double-copy method. This is important for us to derive the warped GRET from warped GAET *via double-copy construction* in Sec. 4.2. We derived in Sec. 2.3 two new BRST identities (2.69) and (2.91) to generally amputate the *mixed* external KK graviton states ( $h_n^{\mu\nu}$ ) and KK Goldstone states ( $\phi_n$  or  $\mathcal{V}_n^\mu$ ), which are needed because such mixings generally reappear at loop levels. We presented GRET type-II in general  $R_\xi$  gauge up to loop level (Sec. 2.3.3) and GRET type-I up to loop level in 't Hooft-Feynman gauge for simplicity (Sec. 2.3.2), whereas in Appendix B.1 we also formulated the GRET type-I (including the LSZ reduction for external states) in the general  $R_\xi$  gauge at tree level. Using the generalized power-counting method [15] for the compactified KK GR theories [with the GRET (2.99) and (2.84)] and the KK gauge theories [with the GAET (2.36)], we proved the different energy cancellations for the scattering amplitudes of longitudinal KK gravitons (with helicity-0) in Eq.(2.104), of the KK gravitons (with helicity  $\pm 1$ ) in Eq.(2.105), and of the KK gauge bosons in Eq.(2.106).

### 3 Structure of Massive Scattering Amplitudes from Warped Gauge and Gravitational Equivalence Theorems

In this section, we systematically compute the three-point and four-point scattering amplitudes of KK gauge bosons (KK gravitons) and of their corresponding KK Goldstone bosons. Then, we explicitly demonstrate the validity of KK GAET (2.38) and GRET (2.86)(2.103) by comparing the scattering amplitudes of KK gauge bosons (KK gravitons) with their corresponding KK Goldstone boson amplitudes at the leading order of high energy expansion.

#### 3.1 Warped Gauge and Gravitational ETs for 3-Point KK Amplitudes

In this subsection, we systematically compute the tree-level 3-point scattering amplitudes of massive KK gauge bosons in Section 3.1.1, and the 3-point scattering amplitudes of KK gravitons in Section 3.1.2.

##### 3.1.1 Warped GAET for 3-Point KK Gauge Boson Amplitudes

To compute the on-shell 3-point KK gauge and Goldstone scattering amplitudes, we derive the Feynman rules for cubic KK gauge/Goldstone vertices in Appendix C.1. Then, we derive the following general 3-point on-shell scattering amplitude of KK gauge bosons:

$$\mathcal{T}[A_{\lambda_1}^{an_1} A_{\lambda_2}^{bn_2} A_{\lambda_3}^{cn_3}] = -i2g a_{n_1 n_2 n_3} f^{abc} [(\epsilon_1 \cdot \epsilon_2)(\epsilon_3 \cdot p_1) + (\epsilon_2 \cdot \epsilon_3)(\epsilon_1 \cdot p_2) + (\epsilon_3 \cdot \epsilon_1)(\epsilon_2 \cdot p_3)], \quad (3.1)$$

where the subscript  $\lambda_j = \pm, 0$  ( $j = 1, 2, 3$ ) denotes the helicity of each external state and  $\epsilon_j^\mu (= \epsilon_{\lambda_j}^\mu)$  is the shorthand notation for the polarization vector of each external KK gauge boson. All the three external momenta  $\{p_j\}$  are outgoing and obey the momentum conservation  $p_1 + p_2 + p_3 = 0$ . In the above tree-level scattering amplitude,  $g$  is the Yang-Mills

gauge coupling and  $a_{n_1 n_2 n_3}$  denotes coupling coefficient induced by integrating over the compactified 5th-dimension coordinate as defined in Eq.(C.3).

With these, we compute the 3-point scattering amplitude of KK gauge bosons of *LLT* type at tree level, which contains two longitudinal KK gauge bosons [chosen as particles 1 and 2 and denoted by  $A_L^{an_1}(p_1)$  and  $A_L^{bn_2}(p_2)$ ], and one transverse KK gauge boson [chosen as the particle 3 and denoted by  $A_{\pm}^{cn_3}(p_3)$ ]. Thus, we derive the following 3-point KK gauge boson amplitude:

$$\begin{aligned}
\mathcal{T}[A_L^{an_1}A_L^{bn_2}A_{\pm}^{cn_3}] &= \mp i g f^{abc} \frac{[E_1^2 + E_2^2 + E_3^2 - 2E_1 E_2(1 + c_{2\theta}) - 2k^2] k s_{\theta}}{\sqrt{2} M_{n_1} M_{n_2}} a_{n_1 n_2 n_3}, \\
&= \mp i g f^{abc} \frac{(M_{n_3}^2 - M_{n_1}^2 - M_{n_2}^2) k s_{\theta}}{\sqrt{2} M_{n_1} M_{n_2}} a_{n_1 n_2 n_3} + O(E^0), \\
&= -i g f^{abc} (\epsilon_3 \cdot p_1) \frac{(M_{n_3}^2 - M_{n_1}^2 - M_{n_2}^2)}{M_{n_1} M_{n_2}} a_{n_1 n_2 n_3} + O(E^0), \tag{3.2}
\end{aligned}$$

where each external momentum is on-shell ( $p_{j\mu} p_j^{\mu} = -M_{n_j}^2$ ) and  $s_{\theta} = \sin \theta$  with  $\theta$  being the angle between the moving directions of particles 1 and 3. The cubic coupling coefficient of KK gauge bosons  $a_{n_1 n_2 n_3}$  is given by Eq.(C.3). We further compute the corresponding 3-point scattering amplitude with particles 1 and 2 replaced by the KK Goldstone bosons,  $A_5^{an_1}(p_1)A_5^{bn_2}(p_2)$ . Thus, we derive the following 3-point scattering amplitudes of KK Goldstone and gauge bosons:<sup>7</sup>

$$\tilde{\mathcal{T}}[A_5^{an_1}A_5^{bn_2}A_{\pm}^{cn_3}] = -i 2 g f^{abc} (\epsilon_3 \cdot p_1) \tilde{a}_{n_1 n_2 n_3} = \mp i g f^{abc} \sqrt{2} k s_{\theta} \tilde{a}_{n_1 n_2 n_3}, \tag{3.3}$$

where the cubic KK Goldstone-gauge coupling coefficient  $\tilde{a}_{n_1 n_2 n_3}$  is given by Eq.(C.3).

According to the KK GAET (2.38), the equivalence between the two KK scattering amplitudes (3.2) and (3.3) in the high energy limit,

$$\mathcal{T}_0[A_L^{an_1}A_L^{bn_2}A_{\pm}^{cn_3}] = -\tilde{\mathcal{T}}_0[A_5^{an_1}A_5^{bn_2}A_{\pm}^{cn_3}], \tag{3.4}$$

where  $\mathcal{T}_0$  and  $\tilde{\mathcal{T}}_0$  denote the leading-order KK amplitudes under high energy expansion. The 3-point GAET (3.4) is equivalent to the following sum rule condition:

$$(M_{n_1}^2 + M_{n_2}^2 - M_{n_3}^2) a_{n_1 n_2 n_3} = 2 M_{n_1} M_{n_2} \tilde{a}_{n_1 n_2 n_3}, \tag{3.5}$$

which connects the trilinear KK gauge boson coupling  $a_{n_1 n_2 n_3}$  to the corresponding trilinear KK Goldstone-gauge boson coupling  $\tilde{a}_{n_1 n_2 n_3}$ . The condition (3.5) does hold for the warped 5d KK Yang-Mills gauge theory as we directly prove it in Eq.(D.6) of Appendix D. For the special case of  $(n_1, n_2, n_3) = (n, n, j)$ , the condition (3.5) takes the following simple form:

$$(1 - \frac{1}{2} r_j^2) a_{nnj} = \tilde{a}_{nnj}, \tag{3.6}$$

---

<sup>7</sup>From this section and hereafter, we denote scattering amplitudes containing external Goldstone states with a tilde symbol, such as  $\tilde{\mathcal{T}}$  and  $\tilde{\mathcal{M}}$ , to distinguish them from the amplitudes with all external states being physical particles.

where we denote  $r_j = M_j/M_n$ .

Next, we compute the 3-point KK gauge boson scattering amplitude including one longitudinal and two transverse external states and the corresponding KK Goldstone-gauge boson scattering amplitude:

$$\begin{aligned} \mathcal{T}_0[A_{\pm}^{an_1}A_{\pm}^{bn_2}A_L^{cn_3}] &= -igf^{abc}a_{n_1n_2n_3}\left\{(\epsilon_1\cdot\epsilon_2)\left[(M_{n_1}^2-M_{n_2}^2)M_{n_3}^{-1}+(p_1-p_2)\cdot v_3\right]\right. \\ &\quad \left.+2(\epsilon_1\cdot p_2)(\epsilon_2\cdot v_3)-2(\epsilon_1\cdot v_3)(\epsilon_2\cdot p_1)\right\} \\ &= 0 + O(E^0), \end{aligned} \quad (3.7a)$$

$$\begin{aligned} \tilde{\mathcal{T}}_0[A_{\pm}^{an_1}A_{\pm}^{bn_2}A_5^{cn_3}] &= gf^{abc}(\epsilon_1\cdot\epsilon_2)(\tilde{a}_{n_1n_3n_2}M_{n_1}-\tilde{a}_{n_2n_3n_1}M_{n_2}) \\ &= gf^{abc}a_{n_1n_2n_3}(\epsilon_1\cdot\epsilon_2)(M_{n_1}^2-M_{n_2}^2)M_{n_3}^{-1} = 0 + O(E^0), \end{aligned} \quad (3.7b)$$

where  $v_3^\mu = \epsilon_{3,L}^\mu - p_3^\mu/M_{n_3} = O(M_{n_3}/E_3)$ , and we have used the sum rules (D.5) to simplify the second KK amplitude (3.7b). From the above, we see that the leading contributions of  $O(E^1)$  vanishes, and the remaining contributions are of the same  $O(E^0)$  as the residual term  $\mathcal{T}_v = \mathcal{T}[A_{\pm}^{an_1}A_{\pm}^{bn_2}A_\mu^{cn_3}]v^\mu$ .

We have also analyzed the 3-point scattering amplitudes of the pure longitudinal KK gauge bosons and of their corresponding KK Goldstone bosons. The 3-point longitudinal KK gauge boson amplitude ( $3A_L^{an}$ ) has the leading energy dependence of  $E^4$  by the power counting rule (2.106a), which arises from the trilinear KK gauge boson vertex (containing one external momentum) and the three external longitudinal polarization vectors, each of which has the high energy behavior

$$\epsilon_L^\mu = \epsilon_S^\mu + v^\mu = O(E^1) + O(E^{-1}), \quad (3.8)$$

according to Eq.(2.35b), where  $\epsilon_S^\mu = p_j^\mu/M_{n_j}$  for the  $j$ -th external state ( $j=1, 2, 3$ ). The four external momenta (from either the trilinear vertex or the  $\epsilon_S^\mu$  part of the external longitudinal polarization vector  $\epsilon_L^\mu$ ) will form two bilinear contractions like  $p_i\cdot p_j$  or  $p_j^2$ , which can be converted into squared KK-masses due to the momentum conservation  $p_1+p_2+p_3=0$  and the on-shell condition of each external momentum  $p_j^2 = -M_{n_j}^2$ . So all the leading energy dependence of  $E^4$  will be reduced to  $E^0$ . The remaining contributions arise from replacing the external longitudinal polarization vectors by one, two, and three vectors of  $v^\mu = O(E^{-1})$ , which can contribute to the 3-point KK gauge boson amplitude with leading energy dependence of  $E^0$ ,  $E^0$ , and  $E^{-2}$  respectively. Hence, the 3-point longitudinal KK gauge boson amplitude should actually have a deduction on its leading energy dependence,  $E^4 \rightarrow E^0$ . We can show that this is consistent with the requirement of the GAET identity (2.36). We note that the corresponding 3-point KK Goldstone amplitude vanishes at tree level due to the absence of cubic vertex  $A_5^{an}-A_5^{am}-A_5^{al}$ . Thus, according to the GAET identity (2.36), the leading nonzero contribution on its right-hand side arises from the residual terms with the trilinear vertex having two KK Goldstone bosons and one KK gauge boson (contracting with the external vector  $v^\mu$ ) which is apparently of  $O(E^0)$ . Note that the second power counting rule in Eq.(2.106a) gives:  $D_E = 4-3-1=0$ , where we have inputs  $(N, N_v) = (3, 1)$  and  $\bar{V}_3^{\min} = 0$ . Hence, the GAET identity (2.36) also enforces the

energy deduction of  $E^4 \rightarrow E^0$  for the 3-point longitudinal KK gauge boson amplitude on its left-hand side.

Then, we explicitly compute the 3-point scattering amplitude of pure longitudinal KK gauge bosons, in which each external longitudinal polarization vector can be split into two parts,  $\epsilon_L^\mu = \epsilon_S^\mu + v^\mu$  as shown in Eq.(3.8). Thus, we derive the expanded scattering amplitude of 3 longitudinal KK gauge bosons:

$$\begin{aligned} \mathcal{T}[A_L^{an_1} A_L^{bn_2} A_L^{cn_3}] &= -igf^{abc}(M_{n_1} M_{n_2} M_{n_3})^{-1} a_{n_1 n_2 n_3} \times \\ &\left[ (p_1 \cdot p_2)(p_3 \cdot v_2) - (p_1 \cdot p_2)(p_3 \cdot v_1) - (p_2 \cdot p_3)(p_1 \cdot v_2) + (p_1 \cdot p_3)(p_2 \cdot v_1) \right. \\ &+ (p_2 \cdot p_3)(p_1 \cdot v_3) - (p_1 \cdot p_3)(p_2 \cdot v_3) + (p_1 \cdot v_3)(p_2 \cdot v_1) - (p_1 \cdot p_2)(v_1 \cdot v_3) - (p_1 \cdot v_2)(p_2 \cdot v_3) \\ &+ (p_1 \cdot p_2)(v_2 \cdot v_3) - (p_1 \cdot v_2)(p_3 \cdot v_1) + (p_1 \cdot p_3)(v_1 \cdot v_2) + (p_1 \cdot v_3)(p_3 \cdot v_2) - (p_1 \cdot p_3)(v_2 \cdot v_3) \\ &+ (p_1 \cdot v_3)(v_1 \cdot v_2) - (p_1 \cdot v_2)(v_1 \cdot v_3) - (p_2 \cdot p_3)(v_1 \cdot v_2) + (p_2 \cdot v_1)(p_3 \cdot v_2) + (p_2 \cdot p_3)(v_1 \cdot v_3) \\ &\left. - (p_2 \cdot v_3)(p_3 \cdot v_1) - (p_2 \cdot v_3)(v_1 \cdot v_2) + (p_2 \cdot v_1)(v_2 \cdot v_3) + (p_3 \cdot v_2)(v_1 \cdot v_3) - (p_3 \cdot v_1)(v_2 \cdot v_3) \right]. \quad (3.9) \end{aligned}$$

In the above amplitude, we see that all the terms having leading energy dependence of  $E^4$  disappear as expected, and all the remaining terms can be classified into the following 4 types according to the numbers of  $v_j$ 's contained therein:

$$(p_i \cdot p_j)(p_k \cdot v_\ell), \quad (p_i \cdot v_j)(p_k \cdot v_\ell), \quad (p_i \cdot p_j)(v_k \cdot v_\ell), \quad (p_i \cdot v_j)(v_k \cdot v_\ell). \quad (3.10)$$

Due to the momentum conservation  $p_1 + p_2 + p_3 = 0$  and the on-shell condition of each external momentum, the bilinear momentum products can all be converted into squared KK-masses,  $p_i \cdot p_j = \frac{1}{2}(M_{n_i}^2 + M_{n_j}^2 - M_{n_k}^2)$  (with  $i \neq j \neq k$ ) and  $p_j^2 = -M_{n_j}^2$ . Since  $v_j = O(M_{n_j}/E_j)$ , the 3 longitudinal KK gauge boson amplitude (3.9) could only have leading energy dependence of  $E^0$  and are given by the terms of the first and second types in Eq.(3.10). Thus, the GAET identity (2.36) for the 3-point tree-level scattering amplitude of longitudinal KK gauge bosons take the following form:

$$\mathcal{T}[A_L^{an_1} A_L^{bn_2} A_L^{cn_3}] = i^3 \tilde{\mathcal{T}}[A_5^{an_1} A_5^{bn_2} A_5^{cn_3}] + \mathcal{T}_v = 0 + O(E^0), \quad (3.11)$$

where on the right-hand side the 3-point KK Goldstone amplitude vanishes and the residual term  $\mathcal{T}_v$  gives the leading contributions to the corresponding longitudinal KK gauge boson amplitude on the left-hand side. Thus, we further deduce the  $O(E^0)$  contributions to the 3-point scattering amplitude of pure longitudinal KK gauge bosons as follows:

$$\begin{aligned} \mathcal{T}_0[A_L^{an_1} A_L^{bn_2} A_L^{cn_3}] &= -igf^{abc}(M_{n_1} M_{n_2} M_{n_3})^{-1} a_{n_1 n_2 n_3} \times \\ &\left[ M_{13}(p_2 \cdot v_1) - M_{12}(p_3 \cdot v_1) - M_{23}(p_1 \cdot v_2) + M_{23}(p_1 \cdot v_3) + M_{12}(p_3 \cdot v_2) - M_{13}(p_2 \cdot v_3) \right. \\ &+ (p_1 \cdot v_3)(p_2 \cdot v_1) - (p_1 \cdot v_2)(p_2 \cdot v_3) - (p_1 \cdot v_2)(p_3 \cdot v_1) + (p_1 \cdot v_3)(p_3 \cdot v_2) \\ &\left. + (p_2 \cdot v_1)(p_3 \cdot v_2) - (p_2 \cdot v_3)(p_3 \cdot v_1) \right], \quad (3.12) \end{aligned}$$

where we have defined  $M_{ij} \equiv p_i \cdot p_j = \frac{1}{2}(M_{n_i}^2 + M_{n_j}^2 - M_{n_k}^2)$  with  $i \neq j \neq k$ .

### 3.1.2 Warped GRET for 3-Point KK Graviton Amplitudes

In this subsection, we compute the 3-point massive scattering amplitudes of KK gravitons and of the corresponding gravitational KK Goldstone bosons. With these we explicitly demonstrate that the KK GRETs (2.103) and (2.86) hold at tree level.

Using the Feynman rules of Appendix C.2, we explicitly compute the most general 3-point on-shell scattering amplitudes of KK gravitons at tree level as given in Eq.(C.15). Using this, we derive the 3-point KK graviton scattering amplitude with the external states (1, 2, 3) having helicities ( $\pm 1$ ,  $\pm 1$ ,  $\pm 2$ ). We also compute the corresponding on-shell 3-point scattering amplitudes with external states (1, 2) replaced by the gravitational KK vector Goldstone bosons having helicities  $\pm 1$ . Thus, we derive the following 3-point gravitational KK scattering amplitudes:

$$\begin{aligned}
\mathcal{M}[h_{n_1}^{\pm 1} h_{n_2}^{\pm 1} h_{n_3}^{\pm 2}] &= -\kappa \left[ (\epsilon_1 \cdot p_2)(\epsilon_1^L \cdot \epsilon_2^L)(\epsilon_2 \cdot \epsilon_3)(\epsilon_3 \cdot p_1) + (\epsilon_1 \cdot \epsilon_2^L)(\epsilon_1^L \cdot p_2)(\epsilon_2 \cdot \epsilon_3)(\epsilon_3 \cdot p_1) \right. \\
&\quad + (\epsilon_1 \cdot p_2)(\epsilon_1^L \cdot \epsilon_2)(\epsilon_2^L \cdot \epsilon_3)(\epsilon_3 \cdot p_1) + (\epsilon_1 \cdot \epsilon_2)(\epsilon_1^L \cdot p_2)(\epsilon_2^L \cdot \epsilon_3)(\epsilon_3 \cdot p_1) \\
&\quad + (\epsilon_3 \cdot \epsilon_1)(\epsilon_1^L \cdot \epsilon_2^L)(\epsilon_2 \cdot p_3)(\epsilon_3 \cdot p_1) + (\epsilon_3 \cdot \epsilon_1)(\epsilon_1^L \cdot \epsilon_2)(\epsilon_2^L \cdot p_3)(\epsilon_3 \cdot p_1) \\
&\quad + (\epsilon_1 \cdot \epsilon_2^L)(\epsilon_3 \cdot \epsilon_1^L)(\epsilon_2 \cdot p_3)(\epsilon_3 \cdot p_1) + (\epsilon_1 \cdot \epsilon_2)(\epsilon_3 \cdot \epsilon_1^L)(\epsilon_2^L \cdot p_3)(\epsilon_3 \cdot p_1) \\
&\quad + (\epsilon_1 \cdot \epsilon_2^L)(\epsilon_1^L \cdot \epsilon_2)(\epsilon_3 \cdot p_1)^2 + (\epsilon_1 \cdot \epsilon_2)(\epsilon_1^L \cdot \epsilon_2^L)(\epsilon_3 \cdot p_1)^2 \\
&\quad + (\epsilon_3 \cdot \epsilon_1)(\epsilon_1^L \cdot p_2)(\epsilon_2 \cdot p_3)(\epsilon_2^L \cdot \epsilon_3) + (\epsilon_3 \cdot \epsilon_1)(\epsilon_1^L \cdot p_2)(\epsilon_2 \cdot \epsilon_3)(\epsilon_2^L \cdot p_3) \\
&\quad + (\epsilon_1 \cdot p_2)(\epsilon_3 \cdot \epsilon_1^L)(\epsilon_2 \cdot p_3)(\epsilon_2^L \cdot \epsilon_3) + (\epsilon_1 \cdot p_2)(\epsilon_3 \cdot \epsilon_1^L)(\epsilon_2 \cdot \epsilon_3)(\epsilon_2^L \cdot p_3) \\
&\quad \left. + 2(\epsilon_1 \cdot p_2)(\epsilon_1^L \cdot p_2)(\epsilon_2 \cdot \epsilon_3)(\epsilon_2^L \cdot \epsilon_3) + 2(\epsilon_3 \cdot \epsilon_1)(\epsilon_3 \cdot \epsilon_1^L)(\epsilon_2 \cdot p_3)(\epsilon_2^L \cdot p_3) \right], \tag{3.13a}
\end{aligned}$$

$$\begin{aligned}
\widetilde{\mathcal{M}}[\mathcal{V}_{n_1}^{\pm 1} \mathcal{V}_{n_2}^{\pm 1} h_{n_3}^{\pm 2}] &= \kappa \left[ (\epsilon_1 \cdot \epsilon_2)(\epsilon_3 \cdot p_1)^2 + (\epsilon_2 \cdot \epsilon_3)(\epsilon_1 \cdot p_2)(\epsilon_3 \cdot p_1) + (\epsilon_3 \cdot \epsilon_1)(\epsilon_2 \cdot p_3)(\epsilon_3 \cdot p_1) \right. \\
&\quad \left. - (\mathbb{M}_{n_1}^2 + \mathbb{M}_{n_2} \mathbb{M}_{n_3} + \mathbb{M}_{n_3}^2)(\epsilon_3 \cdot \epsilon_1)(\epsilon_3 \cdot \epsilon_2) \right] \widetilde{\alpha}_{n_1 n_2 n_3}, \tag{3.13b}
\end{aligned}$$

where  $\epsilon_j = \epsilon_j^\pm$  denotes transverse polarization vector associated with the external KK graviton  $j$  ( $= 1, 2, 3$ ), and  $\epsilon_{j'}^L$  ( $j' = 1, 2$ ) denotes longitudinal polarization vector associated with the external KK graviton  $j'$ . In the above, the three-point gravitational KK coupling coefficients ( $\alpha_{n_1 n_2 n_3}$ ,  $\widetilde{\alpha}_{n_1 n_2 n_3}$ ) are defined in Eq.(C.3) of Appendix C.1. Then, we make high energy expansion and derive the 3-point KK scattering amplitudes at the leading order:

$$\begin{aligned}
\mathcal{M}_0[h_{n_1}^{\pm 1} h_{n_2}^{\pm 1} h_{n_3}^{\pm 2}] &= \frac{\kappa(\mathbb{M}_{n_3}^2 - \mathbb{M}_{n_1}^2 - \mathbb{M}_{n_2}^2)}{2\mathbb{M}_{n_1} \mathbb{M}_{n_2}} \left[ (\epsilon_1 \cdot \epsilon_2)(\epsilon_3 \cdot p_1)^2 + (\epsilon_2 \cdot \epsilon_3)(\epsilon_1 \cdot p_2)(\epsilon_3 \cdot p_1) \right. \\
&\quad \left. + (\epsilon_3 \cdot \epsilon_1)(\epsilon_2 \cdot p_3)(\epsilon_3 \cdot p_1) \right] \alpha_{n_1 n_2 n_3}, \tag{3.14a}
\end{aligned}$$

$$\begin{aligned}
\widetilde{\mathcal{M}}_0[\mathcal{V}_{n_1}^{\pm 1} \mathcal{V}_{n_2}^{\pm 1} h_{n_3}^{\pm 2}] &= \kappa \left[ (\epsilon_1 \cdot \epsilon_2)(\epsilon_3 \cdot p_1)^2 + (\epsilon_2 \cdot \epsilon_3)(\epsilon_1 \cdot p_2)(\epsilon_3 \cdot p_1) \right. \\
&\quad \left. + (\epsilon_3 \cdot \epsilon_1)(\epsilon_2 \cdot p_3)(\epsilon_3 \cdot p_1) \right] \widetilde{\alpha}_{n_1 n_2 n_3}. \tag{3.14b}
\end{aligned}$$

The equivalence between the two leading-order KK scattering amplitudes  $\mathcal{M}_0$  and  $\widetilde{\mathcal{M}}_0$  in Eqs.(3.14a)-(3.14b) is realized by the identity,

$$\mathcal{M}_0[h_{n_1}^{\pm 1} h_{n_2}^{\pm 1} h_{n_3}^{\pm 2}] = -\widetilde{\mathcal{M}}_0[\mathcal{V}_{n_1}^{\pm 1} \mathcal{V}_{n_2}^{\pm 1} h_{n_3}^{\pm 2}], \tag{3.15}$$

where the minus sign on the right-hand side comes from the modification factor  $\hat{C}_{\text{mod}} = (-i)^2$ , and Eq.(3.15) enforces the following condition:

$$\left(\mathbb{M}_{n_1}^2 + \mathbb{M}_{n_2}^2 - \mathbb{M}_{n_3}^2\right)\alpha_{n_1 n_2 n_3} = 2\mathbb{M}_{n_1}\mathbb{M}_{n_2}\tilde{\alpha}_{n_1 n_2 n_3}. \quad (3.16)$$

We have directly proved that the above condition (3.16) does hold for the warped 5d KK gravity theory as shown in Eq.(D.7) of Appdenix D. We compare the above Eq.(3.16) with Eq.(3.5) which holds the GAET (3.5) for 3-point KK gauge boson (KK Goldstone boson) amplitude. We observe that the condition (3.16) has *identical structure* to the condition (3.5) and can be inferred from Eq.(3.5) under the following mass and coupling correspondences (replacements):

$$M_{n_j} \rightarrow \mathbb{M}_{n_j}, \quad a_{n_1 n_2 n_3} \rightarrow \alpha_{n_1 n_2 n_3}, \quad \tilde{a}_{n_1 n_2 n_3} \rightarrow \tilde{\alpha}_{n_1 n_2 n_3}. \quad (3.17)$$

As we will see in Section 4.1, the above correspondences (replacements) are consistent with the double-copy construction of the GRET from the GAET for the 3-point KK scattering amplitudes.

Besides, for the special case of  $(n_1, n_2, n_3) = (n, n, j)$ , the condition (3.16) can be simplified as follows:

$$\left(1 - \frac{1}{2}\hat{r}_j^2\right)\alpha_{nnj} = \tilde{\alpha}_{nnj}, \quad (3.18)$$

with the defined mass ratio  $\hat{r}_j = \mathbb{M}_j/\mathbb{M}_n$ .

Next, we derive the 3-point KK graviton scattering amplitude with external states (1, 2) being longitudinal KK gravitons and the external state 3 having helicities  $\pm 2$ . We also compute the 3-point KK Goldstone scattering amplitude with the external states (1, 2) replaced by the corresponding gravitational KK Goldstone bosons. Thus, using the general 3-point KK amplitudes (C.15) of Appendix C.2, we infer the following 3-point gravitational KK scattering amplitudes at the leading order of high energy expansion:

$$\mathcal{M}_0[h_{n_1}^L h_{n_2}^L h_{n_3}^{\pm 2}] = \frac{\kappa(\epsilon_3 \cdot p_1)^2}{6\mathbb{M}_{n_1}^2 \mathbb{M}_{n_2}^2} \left[ 2\mathbb{M}_{n_1}^2 \mathbb{M}_{n_2}^2 + \left(\mathbb{M}_{n_1}^2 + \mathbb{M}_{n_2}^2 - \mathbb{M}_{n_3}^2\right)^2 \right] \alpha_{n_1 n_2 n_3}, \quad (3.19a)$$

$$\tilde{\mathcal{M}}_0[\phi_{n_1} \phi_{n_2} h_{n_3}^{\pm 2}] = \kappa(\epsilon_3 \cdot p_1)^2 \tilde{\beta}_{n_1 n_2 n_3}, \quad (3.19b)$$

where the trilinear KK coupling coefficient  $\tilde{\beta}_{n_1 n_2 n_3}$  is defined in Eq.(C.9c) of Appendix C.1 and the KK Goldstone boson amplitude (3.19b) actually equals its full amplitude without expansion, i.e.,  $\tilde{\mathcal{M}}[\phi_{n_1} \phi_{n_2} h_{n_3}^{\pm 2}] = \kappa(\epsilon_3 \cdot p_1)^2 \tilde{\beta}_{n_1 n_2 n_3}$ . Then, to hold the KK GRET (2.99) for the 3-point KK scattering amplitudes,

$$\mathcal{M}_0[h_{n_1}^L h_{n_2}^L h_{n_3}^{\pm 2}] = \tilde{\mathcal{M}}_0[\phi_{n_1} \phi_{n_2} h_{n_3}^{\pm 2}], \quad (3.20)$$

we derive the following condition to ensure the equivalence between the two gravitational KK scattering amplitudes (3.19a) and (3.19b):

$$\left[\left(\mathbb{M}_{n_1}^2 + \mathbb{M}_{n_2}^2 - \mathbb{M}_{n_3}^2\right)^2 + 2\mathbb{M}_{n_1}^2 \mathbb{M}_{n_2}^2\right] \alpha_{n_1 n_2 n_3} = 6\mathbb{M}_{n_1}^2 \mathbb{M}_{n_2}^2 \tilde{\beta}_{n_1 n_2 n_3}. \quad (3.21)$$

The proof of the condition (3.21) is highly nontrivial and we present a direct proof of it around Eq.(D.20) of Appdenix D. For the case of  $(n_1, n_2, n_3) = (n, n, j)$ , the condition (3.21) can be simplified as follows:

$$(\hat{r}_j^4 - 4\hat{r}_j^2 + 6)\alpha_{nnj} = 6\tilde{\beta}_{nnj}, \quad (3.22)$$

with the mass ratio  $\hat{r}_j = \mathbb{M}_j/\mathbb{M}_n$ .

Then, we consider a mixed 3-point scattering amplitudes including two external KK gravitons with helicities  $\pm 1$  and one external KK graviton with helicity zero. Thus, we derive the following 3-point scattering amplitudes of KK gravitons and of the corresponding KK Goldstone bosons at the leading order of high energy expansion:

$$\mathcal{M}_0[h_{n_1}^{\pm 1}h_{n_2}^{\pm 1}h_{n_3}^L] = \frac{\kappa[2 - (\hat{r}_{13}^2 + \hat{r}_{23}^2) - (\hat{r}_{13}^2 - \hat{r}_{23}^2)^2]}{2\sqrt{6}\hat{r}_{13}\hat{r}_{23}}(\epsilon_1 \cdot p_2)(\epsilon_2 \cdot p_1)\alpha_{n_1 n_2 n_3}, \quad (3.23a)$$

$$\begin{aligned} \tilde{\mathcal{M}}[\mathcal{V}_{n_1}^{\pm 1}\mathcal{V}_{n_2}^{\pm 1}\phi_{n_3}] &= -\kappa\left[\sqrt{\frac{3}{2}}(\epsilon_1 \cdot p_2)(\epsilon_2 \cdot p_1) - \frac{\mathbb{M}_{n_3}^2}{\sqrt{6}}(\epsilon_1 \cdot \epsilon_2)(\hat{r}_{13}\hat{r}_{23} + 2\hat{r}_{13} + 2\hat{r}_{23})\right]\tilde{\rho}_{n_1 n_2 n_3} \\ &= \mathcal{M}_0[\mathcal{V}_{n_1}^{\pm 1}\mathcal{V}_{n_2}^{\pm 1}\phi_{n_3}] + O(E^0), \end{aligned} \quad (3.23b)$$

$$\tilde{\mathcal{M}}_0[\mathcal{V}_{n_1}^{\pm 1}\mathcal{V}_{n_2}^{\pm 1}\phi_{n_3}] = -\kappa\sqrt{\frac{3}{2}}(\epsilon_1 \cdot p_2)(\epsilon_2 \cdot p_1)\tilde{\rho}_{n_1 n_2 n_3}, \quad (3.23c)$$

where we have defined the KK mass ratios  $\hat{r}_{13} = \mathbb{M}_{n_1}/\mathbb{M}_{n_3}$  and  $\hat{r}_{23} = \mathbb{M}_{n_2}/\mathbb{M}_{n_3}$ , and the  $\mathcal{V}_{n_1}^{\pm 1}\mathcal{V}_{n_2}^{\pm 1}\phi_{n_3}$  trilinear coupling coefficient  $\tilde{\rho}_{n_1 n_2 n_3}$  is defined in Eq.(C.9) of Appendix C. There is another helicity combination for such mixed 3-point KK scattering amplitudes, but they vanish at the leading order of high energy expansion:

$$\mathcal{M}_0[h_{n_1}^{\pm 1}h_{n_2}^{\mp 1}h_{n_3}^L] = 0, \quad \tilde{\mathcal{M}}_0[\mathcal{V}_{n_1}^{\pm 1}\mathcal{V}_{n_2}^{\mp 1}\phi_{n_3}] = 0. \quad (3.24)$$

Using the 3-point gravitational KK scattering amplitudes (3.23a)-(3.23c), we find that holding the GRET,

$$\mathcal{M}_0[h_{n_1}^{\pm 1}h_{n_2}^{\pm 1}h_{n_3}^L] = -\tilde{\mathcal{M}}_0[\mathcal{V}_{n_1}^{\pm 1}\mathcal{V}_{n_2}^{\pm 1}\phi_{n_3}], \quad (3.25)$$

will impose the following condition:

$$[2 - (\hat{r}_{13}^2 + \hat{r}_{23}^2) - (\hat{r}_{13}^2 - \hat{r}_{23}^2)^2]\alpha_{n_1 n_2 n_3} = 6\hat{r}_{13}\hat{r}_{23}\tilde{\rho}_{n_1 n_2 n_3}, \quad (3.26)$$

with the KK mass ratios  $(\hat{r}_{13}, \hat{r}_{23})$  and the trilinear KK coupling coefficient  $\tilde{\rho}_{n_1 n_2 n_3}$  are defined below Eq.(3.23). The above identity (3.26) can be directly proven as in (D.22) of Appendix D. Hence, we have explicitly established the GRET in its mixed form (3.25) for the 3-point gravitational KK scattering amplitudes.

Besides, we can consider a specific case with  $(n_1, n_2, n_3) = (n, n, j)$ . So we have  $\hat{r}_{13} = \hat{r}_{23}$  and thus the above GRET condition (3.26) is simplified as follows:

$$(\hat{r}_j^2 - 1)\alpha_{nnj} = 3\tilde{\rho}_{nnj}, \quad (3.27)$$

where the mass ratio  $\hat{r}_j = \mathbb{M}_j/\mathbb{M}_n$ . For the case of compactified flat 5d under  $S^1/\mathbb{Z}_2$ , the KK index  $j$  could only take values of  $j = 0, 2n$  in Eq.(3.27).

Finally, we discuss the 3-point scattering amplitudes of pure longitudinal KK gravitons ( $3h_n^L$ ) and of the corresponding gravitational KK Goldstone bosons ( $3\phi_n$ ). We gave



a brief comment on this in a separate paragraph below Eq.(2.104). The tree-level 3-point longitudinal KK graviton amplitude ( $3h_n^L$ ) have a possible leading energy dependence of  $E^8$  by using the power counting rule (2.104a) for  $N=3$ . But, we note that the 3-point KK graviton amplitude arises from the trilinear KK graviton vertex (whose coupling factor contains products of two external momenta) and is expected to have individual leading energy dependence of  $E^8$  after including the 3 external longitudinal polarization tensors of KK gravitons, each of which has the high energy behavior,

$$\varepsilon_L^{\mu\nu} = \sqrt{\frac{2}{3}} \varepsilon_S^{\mu\nu} + \tilde{v}^{\mu\nu} = O(E^2) + O(E^0), \quad (3.28)$$

based on Eqs.(2.89)-(2.90), where for the  $j$ -th external KK graviton the scalar polarization tensor  $\varepsilon_S^{\mu\nu}(p_j) = p_j^\mu p_j^\nu / M_{n_j}^2$  (with  $j = 1, 2, 3$ ). Thus, each leading energy term of  $O(E^8)$  is given by picking up the leading energy term  $\varepsilon_S^{\mu\nu}$  from all 3 external longitudinal polarization tensors ( $\varepsilon_L^{\mu\nu}$ ) which have contractions with the two-momentum-factors in of the trilinear KK graviton vertex. These momenta will form up 4 pairs of bilinear contraction ( $p_i \cdot p_j$ ) which can be all converted into certain combinations of squared KK-masses by using the momentum conservation  $p_1 + p_2 + p_3 = 0$  and the on-shell condition of each external momentum  $p_j^2 = -M_{n_j}^2$ . Hence, all the leading energy terms of  $O(E^8)$  are reduced to  $O(E^0)$  in this way. The remaining contributions include to replace one, two, and three external longitudinal polarization tensors ( $\varepsilon_L^{\mu\nu}$ ) by the tensor  $\tilde{v}^{\mu\nu} = O(E^0)$  respectively. For those external momentum factors contracting among themselves, they will be converted into squared KK-masses due to the momentum conservation and the on-shell conditions on all the external momenta. Hence, a given external momentum  $p_j^\mu$  could contribute a nontrivial energy factor  $E^1$  to the 3-point KK graviton amplitude only if it contracts with a  $\tilde{v}^{\mu\nu} = O(E^0)$  tensor from Eq.(3.28). The terms having maximal number of such contractions arise from picking up two  $\tilde{v}^{\mu\nu}$  tensors from two of the external KK graviton states via Eq.(3.28); so the third external longitudinal polarization tensor contributes a leading momentum term of  $\varepsilon_S^{\mu\nu}$  via Eq.(3.28) and the trilinear KK graviton vertex contributes the leading coupling factors with two external momenta; thus all the 4 external momenta may contract with the two external  $\tilde{v}^{\mu\nu}$  tensors [via Eq.(3.28)] and thus makes the leading energy contributions of  $O(E^4)$ . On the other hand, the tree-level amplitude of 3 gravitational KK Goldstone bosons ( $3\phi_n$ ) arises from trilinear KK Goldstone vertex whose coupling factor contains products of two external momenta  $p_i \cdot p_j$ . So the naive power counting shows that this KK Goldstone amplitude would have leading energy dependence of  $E^2$  in agreement with Eq.(2.104a). But the momentum conservation and the on-shell condition of each external state convert all the bilinear momentum products ( $p_i \cdot p_j$ ) in the trilinear Goldstone vertex coupling into squared KK-masses, so the 3-point KK Goldstone amplitude actually has energy dependence  $E^0$ . Then, we can apply the GRET (2.99) and note that on its right-hand side the corresponding 3-point KK Goldstone amplitude has the actual leading energy dependence  $E^0$ . Hence, the GRET ensures a large energy cancellation of  $E^4 \rightarrow E^0$  in the 3-point longitudinal KK graviton amplitude.

Then, we explicitly compute the 3-point scattering amplitudes of pure longitudinal KK gravitons ( $3h_n^L$ ) and of the corresponding gravitational KK Goldstone bosons ( $3\phi_n$ ). Under

high energy expansion, we derive these two KK amplitudes down to  $O(E^0)$ :

$$\begin{aligned} \mathcal{M}[h_{n_1}^L h_{n_2}^L h_{n_3}^L] &= \frac{-\kappa \alpha_{n_1 n_2 n_3} \hat{r}_{12}^2 \hat{r}_{13}^2 \mathbb{M}_{n_1}^2}{24\sqrt{6}} \left( -6698 + 7210 \hat{r}_{31}^2 - 20400 \hat{r}_{21}^2 - 18940 \hat{r}_{21}^4 + 14830 \hat{r}_{21}^2 \hat{r}_{31}^2 \right. \\ &\quad \left. - 26471 \hat{r}_{31}^4 - 5520 \hat{r}_{21}^6 + 7310 \hat{r}_{21}^4 \hat{r}_{31}^2 - 2950 \hat{r}_{21}^2 \hat{r}_{31}^4 + 362 \hat{r}_{31}^6 - 186 \hat{r}_{21}^8 \right. \\ &\quad \left. + 458 \hat{r}_{21}^6 \hat{r}_{31}^2 - 375 \hat{r}_{21}^4 \hat{r}_{31}^4 + 114 \hat{r}_{21}^2 \hat{r}_{31}^6 - 11 \hat{r}_{31}^8 \right), \end{aligned} \quad (3.29a)$$

$$\widetilde{\mathcal{M}}[\phi_{n_1} \phi_{n_2} \phi_{n_3}] = -\sqrt{\frac{2}{3}} \kappa \tilde{\omega}_{n_1 n_2 n_3} \mathbb{M}_{n_1}^2 (1 + \hat{r}_{21}^2 + \hat{r}_{31}^2 - \hat{r}_{21} - \hat{r}_{31} - \hat{r}_{21} \hat{r}_{31}), \quad (3.29b)$$

where we have defined the mass ratio  $\hat{r}_{ij} = \mathbb{M}_{n_i} / \mathbb{M}_{n_j}$ , and the trilinear KK graviton coupling  $\alpha_{n_1 n_2 n_3}$  and KK Goldstone coupling  $\tilde{\omega}_{n_1 n_2 n_3}$  are defined in Eq.(C.9). We see that both the above KK scattering amplitudes receive nonzero contributions at  $O(E^0)$ . Even though the power counting rule (2.104a) suggests that the 3-point gravitational KK Goldstone amplitude (3.29b) would scale like  $E^2$ , we find that its actual leading energy dependence is only of  $O(E^0)$  due to the momentum conservation and the on-shell condition for each external momentum, which agrees with our discussion in the previous paragraph.<sup>8</sup> According to the GRET (2.99) together with Eq.(2.102), the residual term contains an amplitude having two (helicity-0) KK Goldstone states ( $\phi_n$  and  $\phi_m$ ) and one external KK state  $v_\mu \mathcal{V}_\ell^\mu$  (with helicity  $\pm 1$ ) also contributes to  $O(E^0)$ . This explains why the two KK amplitudes (3.29a) and (3.29b) above do not simply equal despite that they are both of  $O(E^0)$ .

### 3.2 Warped GAET and GRET: from 3-Point to 4-Point Amplitudes

In Section 3.1 we have explicitly established the GAET for the 3-point KK gauge boson scattering amplitudes and the GRET for the 3-point KK graviton scattering amplitudes in the high energy limit. Using the GAET and GRET for the basic 3-point KK scattering amplitudes, we further prove explicitly in this subsection that the GAET and GRET hold for the 4-point KK scattering amplitudes in the high energy limit. This is realized by proving that the mass-coupling identities (sum rules) ensuring the basic 3-point-level GAET and GRET can guarantee the validity of the GAET and GRET for the 4-point KK scattering amplitudes. With these we also conjecture that this proof can be extended through iterations to the general  $N$ -point ( $N \geq 5$ ) KK scattering amplitudes in the high energy limit.

#### 3.2.1 Warped GAET: from 3-Point to 4-Point KK Amplitudes

In this section, we first compute the 4-point elastic scattering amplitude of longitudinal KK gauge bosons  $\mathcal{T}[A_L^{an} A_L^{bn} \rightarrow A_L^{cn} A_L^{dn}]$  and the corresponding KK Goldstone boson scattering amplitude  $\widetilde{\mathcal{T}}[A_5^{an} A_5^{bn} \rightarrow A_5^{cn} A_5^{dn}]$  at tree level in the 't Hooft-Feynman gauge ( $\xi_n = 1$ ). Under high energy expansion, we derive the leading-order KK amplitudes as follows:

$$\mathcal{T}_0[A_L^{an} A_L^{bn} \rightarrow A_L^{cn} A_L^{dn}] = g^2 (\mathcal{C}_s \mathcal{K}_s^0 + \mathcal{C}_t \mathcal{K}_t^0 + \mathcal{C}_u \mathcal{K}_u^0), \quad (3.30a)$$

<sup>8</sup>From the viewpoint of double-copy (cf. Sec. 4.1), this is also related to the fact that the trilinear KK Goldstone vertex  $A_5^{an} A_5^{bm} A_5^{cl}$  vanishes identically in the compactified KK Yang-Mills gauge theories as shown in Eq.(3.11).

$$\tilde{\mathcal{T}}_0[A_5^{an} A_5^{bn} \rightarrow A_5^{cn} A_5^{dn}] = g^2 (\mathcal{C}_s \tilde{\mathcal{K}}_s^0 + \mathcal{C}_t \tilde{\mathcal{K}}_t^0 + \mathcal{C}_u \tilde{\mathcal{K}}_u^0), \quad (3.30b)$$

where the non-Abelian group factors  $(\mathcal{C}_s, \mathcal{C}_t, \mathcal{C}_u)$  are defined as

$$(\mathcal{C}_s, \mathcal{C}_t, \mathcal{C}_u) = (C^{abe} C^{cde}, C^{ade} C^{bce}, C^{ace} C^{dbe}), \quad (3.31)$$

and they satisfy the color Jacobi identity:

$$\mathcal{C}_s + \mathcal{C}_t + \mathcal{C}_u = 0. \quad (3.32)$$

In Eqs.(3.30a)-(3.30b), both the leading-order KK scattering amplitudes are of  $O(E^0)$  under the high energy expansion, as required by the KK GAET (2.38) which enforces the energy cancellation of  $E^4 \rightarrow E^0$  for the 4-point longitudinal KK scattering amplitude  $\mathcal{T}[A_L^{an} A_L^{bn} \rightarrow A_L^{cn} A_L^{dn}]$  as shown in Eq.(2.106). The exact cancellations of  $O(E^4)$  and  $O(E^2)$  terms in longitudinal KK amplitude  $\mathcal{T}[A_L^{an} A_L^{bn} \rightarrow A_L^{cn} A_L^{dn}]$  impose the following sum-rule-type conditions respectively:

$$\sum_{j=0}^{\infty} a_{nnj}^2 = a_{nnnn}, \quad (3.33a)$$

$$\sum_{j=0}^{\infty} M_j^2 a_{nnj}^2 = \frac{4}{3} M_n^2 a_{nnnn}. \quad (3.33b)$$

The the above conditions (3.33a) and (3.33b) are proved respectively in Eq.(D.3a) and Eq.(D.12) of Appendix D. Combining the above two conditions, we can eliminate the quartic coupling  $a_{nnnn}$  and deduce the following sum rule on the trilinear coupling coefficient  $a_{nnj}$  alone:

$$\sum_{j=0}^{\infty} \left( \frac{4}{3} - r_j^2 \right) a_{nnj}^2 = 0. \quad (3.34)$$

In Eq.(3.30), the leading-order longitudinal KK gauge boson amplitudes for each channel are given by

$$\mathcal{K}_s^0 = -\frac{c_\theta}{4} \left[ 4a_{nn0}^2 + \sum_{j=1}^{\infty} (r_j^2 + 2)^2 a_{nnj}^2 \right], \quad (3.35a)$$

$$\mathcal{K}_t^0 = \frac{1}{(1+c_\theta)} \left\{ (6-19c_\theta-10c_{2\theta}-c_{3\theta}) a_{nn0}^2 + \sum_{j=1}^{\infty} \left[ (3r_j^4-10r_j^2+6) - (r_j^4-8r_j^2+19)c_\theta \right. \right. \\ \left. \left. + 2(r_j^2-5)c_{2\theta}-c_{3\theta} \right] a_{nnj}^2 - (2+c_\theta-2c_{2\theta}-c_{3\theta}) a_{nnnn} \right\}, \quad (3.35b)$$

$$\mathcal{K}_u^0 = \frac{-1}{(1-c_\theta)} \left\{ (6+19c_\theta-10c_{2\theta}+c_{3\theta}) a_{nn0}^2 + \sum_{j=1}^{\infty} \left[ (3r_j^4-10r_j^2+6) + (r_j^4-8r_j^2+19)c_\theta \right. \right. \\ \left. \left. + 2(r_j^2-5)c_{2\theta}+c_{3\theta} \right] a_{nnj}^2 - (2-c_\theta-2c_{2\theta}+c_{3\theta}) a_{nnnn} \right\}, \quad (3.35c)$$

and the leading-order KK Goldstone boson amplitudes for each channel are given by

$$\tilde{\mathcal{K}}_s^0 = -c_\theta \sum_{j=0}^{\infty} \tilde{a}_{nnj}^2, \quad \tilde{\mathcal{K}}_t^0 = \frac{3-c_\theta}{1+c_\theta} \sum_{j=0}^{\infty} \tilde{a}_{nnj}^2, \quad \tilde{\mathcal{K}}_u^0 = -\frac{3+c_\theta}{1-c_\theta} \sum_{j=0}^{\infty} \tilde{a}_{nnj}^2, \quad (3.36)$$

where we define the abbreviations  $(s_{n\theta}, c_{n\theta}) = (\sin n\theta, \cos n\theta)$  with  $\theta$  being the scattering angle. In the above the KK quartic coupling coefficient  $a_{nnnn}$  is defined in Eq.(C.3) of Appendix C.

The KK GAET (2.38) requires the equivalence between the two leading-order KK scattering amplitudes (3.30a) and (3.30b). To prove this explicitly, we compute the difference between the leading-order sub-amplitudes (3.35) and (3.36) for each scattering channel:

$$\mathcal{K}_s^0 - \tilde{\mathcal{K}}_s^0 = -\frac{c_\theta}{4} \left[ \sum_{j=1}^{\infty} r_j^2 (7+r_j^2) a_{nnj}^2 - \sum_{j=0}^{\infty} 4\tilde{a}_{nnj}^2 \right], \quad (3.37a)$$

$$\mathcal{K}_t^0 - \tilde{\mathcal{K}}_t^0 = \frac{-1}{4(1+c_\theta)} \left\{ \sum_{j=1}^{\infty} r_j^2 [7-3r_j^2+(7+r_j^2)c_\theta+4c_{2\theta}] a_{nnj}^2 + 4(3-c_\theta) \sum_{j=0}^{\infty} \tilde{a}_{nnj}^2 \right\}, \quad (3.37b)$$

$$\mathcal{K}_u^0 - \tilde{\mathcal{K}}_u^0 = \frac{1}{4(1-c_\theta)} \left\{ \sum_{j=1}^{\infty} r_j^2 [7-3r_j^2-(7+r_j^2)c_\theta+4c_{2\theta}] a_{nnj}^2 + 4(3+c_\theta) \sum_{j=0}^{\infty} \tilde{a}_{nnj}^2 \right\}, \quad (3.37c)$$

where we have utilized Eq.(3.33) to eliminate the quartic coupling coefficient  $a_{nnnn}$  and Eq.(3.34) to eliminate the constant coefficient of  $a_{nnj}^2$ . Inspecting the three difference expressions (3.37a)-(3.37c) and the sum rule (3.34), we find that they can be equal to each other provided the following sum rule condition is satisfied:

$$\sum_{j=0}^{\infty} \left(1 - \frac{1}{2}r_j^2\right)^2 a_{nnj}^2 = \sum_{j=0}^{\infty} \tilde{a}_{nnj}^2. \quad (3.38)$$

This condition (3.38) can be inferred directly from the identity (3.6) by squaring its two sides and then summing over KK index  $j$ . Note that Eq.(3.6) was proven to hold the GAET (3.4) [in the case of  $(n_1, n_2, n_3) = (n, n, j)$ ] for the most fundamental 3-point KK scattering amplitudes at the leading order. Hence, the validity of the GAET at 4-point-level of KK scattering amplitudes is guaranteed by the fundamental 3-point-level GAET (3.4).

With these we can further simplify the amplitude-difference in each channel of Eq.(3.37) and derive the following:

$$\mathcal{K}_i^0 - \tilde{\mathcal{K}}_i^0 = -2c_\theta \sum_{j=1}^{\infty} r_j^2 a_{nnj}^2, \quad (3.39)$$

where the index  $i \in \{s, t, u\}$  denotes the kinematic channel under consideration. Hence, we sum up the contributions of three kinematic channels and apply the color Jacobi identity (3.32) to prove the equivalence between the two leading-order KK scattering amplitudes (3.30a) and (3.30b):

$$\mathcal{T}_0[4A_L^{an}] - \tilde{\mathcal{T}}_0[4A_5^{an}] = -2c_\theta \sum_{j=1}^{\infty} r_j^2 a_{nnj}^2 \times (\mathcal{C}_s + \mathcal{C}_t + \mathcal{C}_u) = 0. \quad (3.40)$$

In summary, based upon the GAET (3.4) for the most fundamental 3-point KK scattering amplitudes, we have derived the GAET for leading-order 4-point KK gauge scattering amplitudes (3.30a)-(3.30b) as the outcome:

$$\mathcal{T}_0[A_L^{an} A_L^{bn} \rightarrow A_L^{cn} A_L^{dn}] = \tilde{\mathcal{T}}_0[A_5^{an} A_5^{bn} \rightarrow A_5^{cn} A_5^{dn}]. \quad (3.41)$$

The above analysis demonstrates that the GAET for 3-point KK scattering amplitudes is the most fundamental formulation of the GAET, with which we can further deduce the validity of the GAET for 4-point KK scattering amplitudes. Hence, we conclude that *the validity of GAET (3.41) for 4-point KK scattering amplitudes can be reduced to the validity of GAET (3.4) for 3-point KK scattering amplitudes* (which we have already proven in Sec. 3.1.1). Extending this result, we would further conjecture that the GAET for  $N$ -point KK scattering amplitudes ( $N \geq 5$ ) can be also reduced to the validity of the GAET for the 3-point KK scattering amplitudes. This is worth of more investigations in the future work.

As a consistency check, we examine the flat 5d limit by taking the warped parameter  $k \rightarrow 0$ . Substituting the flat-space couplings of Table 1 (Appendix C) into the above amplitudes (3.35)-(3.36), we obtain the following leading-order amplitudes:

$$\mathcal{K}_s^0 = -\frac{11}{2}c_\theta, \quad \tilde{\mathcal{K}}_s^0 = -\frac{3}{2}c_\theta, \quad (3.42a)$$

$$\mathcal{K}_t^0 = \frac{5-11c_\theta-4c_{2\theta}}{2(1+c_\theta)}, \quad \tilde{\mathcal{K}}_t^0 = \frac{3(3-c_\theta)}{2(1+c_\theta)}, \quad (3.42b)$$

$$\mathcal{K}_u^0 = -\frac{5+11c_\theta-4c_{2\theta}}{2(1-c_\theta)}, \quad \tilde{\mathcal{K}}_u^0 = -\frac{3(3+c_\theta)}{2(1-c_\theta)}, \quad (3.42c)$$

which agree with the previous results for the compactified flat 5d Yang-Mills gauge theories (under  $S^1/\mathbb{Z}_2$ ) as given in Refs. [15][16] and also Refs. [11][43]. Moreover, in this flat 5d limit, the elastic scattering amplitude of the longitudinal KK gauge bosons and of the corresponding KK Goldstone bosons differ by the same amount in each channel:

$$\mathcal{K}_i^0 - \tilde{\mathcal{K}}_i^0 = -4c_\theta, \quad (3.43)$$

with  $i \in \{s, t, u\}$ . They have vanishing contribution to the full amplitude due to the color Jacobi identity (3.32).

Next, we consider two inelastic scattering processes,  $(n, n) \rightarrow (m, m)$  with  $n \neq m > 0$ , and  $(n, m) \rightarrow (\ell, q)$  with  $n \neq m \neq \ell \neq q > 0$ . For the inelastic scattering process  $(n, n) \rightarrow (m, m)$ , we compute the 4-point leading-order scattering amplitudes of the longitudinal KK gauge bosons and of the corresponding KK Goldstone bosons:

$$\mathcal{T}_0[A_L^{an} A_L^{bn} \rightarrow A_L^{cm} A_L^{dm}] = g^2(\mathcal{C}_s \mathcal{K}_s^{\text{in}0} + \mathcal{C}_t \mathcal{K}_t^{\text{in}0} + \mathcal{C}_u \mathcal{K}_u^{\text{in}0}), \quad (3.44a)$$

$$\tilde{\mathcal{T}}_0[A_5^{an} A_5^{bn} \rightarrow A_5^{cm} A_5^{dm}] = g^2(\mathcal{C}_s \tilde{\mathcal{K}}_s^{\text{in}0} + \mathcal{C}_t \tilde{\mathcal{K}}_t^{\text{in}0} + \mathcal{C}_u \tilde{\mathcal{K}}_u^{\text{in}0}). \quad (3.44b)$$

In Eqs.(3.44a)-(3.44b), both the leading-order inelastic KK scattering amplitudes are of  $O(E^0)$  under the high energy expansion, as required by the KK GAET (2.38) which enforces the energy cancellations of  $E^4 \rightarrow E^0$  for the 4-point longitudinal KK scattering amplitude  $\mathcal{T}[A_L^{an} A_L^{bn} \rightarrow A_L^{cm} A_L^{dm}]$  as shown in Eq.(2.106). The exact cancellations of  $O(E^4)$  and  $O(E^2)$  terms in the inelastic longitudinal KK amplitude  $\mathcal{T}[A_L^{an} A_L^{bn} \rightarrow A_L^{cm} A_L^{dm}]$  impose the following sum-rule-type conditions:

$$\sum_{j=0}^{\infty} a_{nnj} a_{mmj} = \sum_{j=0}^{\infty} a_{nmj}^2 = a_{nnmm}, \quad (3.45a)$$

$$\sum_{j=1}^{\infty} r_j^2 a_{nmj}^2 = \sum_{j=0}^{\infty} \left[ (1+r^2) - \frac{1}{2} r_j^2 \right] a_{nnj} a_{mmj}, \quad (3.45b)$$

$$\sum_{j=1}^{\infty} \frac{(1-r^2)^2}{r_j^2} a_{nmj}^2 = \sum_{j=0}^{\infty} \left[ (1+r^2) - \frac{3}{2} r_j^2 \right] a_{nnj} a_{mmj}, \quad (3.45c)$$

where  $(r, r_j)$  denote mass ratios and are defined as  $r = M_m/M_n$  and  $r_j = M_j/M_n$ . In the above, the inelastic condition (3.45a) ensures to the energy cancellations at  $O(E^4)$  and is mass-independent, whereas the inelastic conditions (3.45b)-(3.45c) guarantees the energy cancellations at  $O(E^2)$ . We prove the three inelastic conditions respectively in Eqs.(D.3a), (D.23) and Eqs.(D.24)-(D.27) of Appendix D.

In Eq.(3.44a), for each kinematic channel, the 4-point leading-order scattering amplitudes of longitudinal KK gauge bosons are given by

$$\begin{aligned} \mathcal{K}_s^{\text{in}0} = & -\frac{c_\theta}{4r^2} \left\{ 2(5-3r^4) a_{nn0} a_{mm0} + \sum_{j=1}^{\infty} \left[ r_j^4 + 4r_j^2 + 2(5-3r^4) \right] a_{nnj} a_{mmj} \right. \\ & \left. - 2(3-2r^2-r^4) a_{nnmm} \right\}, \end{aligned} \quad (3.46a)$$

$$\begin{aligned} \mathcal{K}_t^{\text{in}0} = & \frac{1}{8r^2(1+c_\theta)} \sum_{j=1}^{\infty} \left\{ 2(3-c_\theta) r_j^4 + \left[ 4(1+3r^2) c_\theta - (3+r^2)(5-c_{2\theta}) \right] r_j^2 \right. \\ & + \left[ (15-6r^2+3r^4) - (2+32r^2+4r^4) c_\theta - (11+10r^2-r^4) c_{2\theta} - 2c_{3\theta} \right] \\ & + (1-r^2)^2 \left[ (5-r^2) + 8c_\theta + (3+r^2) c_{2\theta} \right] r_j^{-2} \left. \right\} a_{nmj}^2 - \frac{1}{8r^2} \left[ 2(3-2r^2-r^4) c_\theta \right. \\ & \left. + 4(3-2r^2-c_{2\theta}) \right] a_{nnmm}, \end{aligned} \quad (3.46b)$$

$$\begin{aligned} \mathcal{K}_u^{\text{in}0} = & -\frac{1}{8r^2(1-c_\theta)} \sum_{j=1}^{\infty} \left\{ 2(3+c_\theta) r_j^4 - \left[ 4(1+3r^2) c_\theta + (3+r^2)(5-c_{2\theta}) \right] r_j^2 \right. \\ & + \left[ (15-6r^2+3r^4) + (2+32r^2+4r^4) c_\theta - (11+10r^2-r^4) c_{2\theta} + 2c_{3\theta} \right] \\ & + (1-r^2)^2 \left[ (5-r^2) - 8c_\theta + (3+r^2) c_{2\theta} \right] r_j^{-2} \left. \right\} a_{nmj}^2 + \frac{1}{8r^2} \left[ 2(3-2r^2-r^4) c_\theta \right. \\ & \left. - 4(3-2r^2-c_{2\theta}) \right] a_{nnmm}, \end{aligned} \quad (3.46c)$$

with the mass ratios  $r = M_m/M_n$  and  $r_j = M_j/M_n$ .

For the corresponding 4-point leading-order inelastic scattering amplitude (3.44b) of KK Goldstone bosons, the sub-amplitudes in each kinematic channel are given by

$$\tilde{\mathcal{K}}_s^{\text{in}0} = -c_\theta \sum_{j=0}^{\infty} \tilde{a}_{nmj} \tilde{a}_{mmj}, \quad \tilde{\mathcal{K}}_t^{\text{in}0} = \frac{3-c_\theta}{1+c_\theta} \sum_{j=0}^{\infty} \tilde{a}_{nmj}^2, \quad \tilde{\mathcal{K}}_u^{\text{in}0} = -\frac{3+c_\theta}{1-c_\theta} \sum_{j=0}^{\infty} \tilde{a}_{nmj}^2. \quad (3.47)$$

Then, we find that in each channel, the 4-point inelastic scattering amplitudes of the longitudinal KK gauge bosons and of the corresponding KK Goldstone bosons differ by the same amount:

$$\mathcal{K}_i^{\text{in}0} - \tilde{\mathcal{K}}_i^{\text{in}0} = -\frac{c_\theta}{2r^2} \sum_{j=0}^{\infty} \left[ (3+r^2) r_j^2 + 2(1-r^4) \right] a_{nnj} a_{mmj}, \quad (3.48)$$

provided that the following sum rule conditions hold:

$$\sum_{j=0}^{\infty} \tilde{a}_{nmj}^2 = \sum_{j=0}^{\infty} \left(1 - \frac{1}{2} r_j^2\right)^2 a_{nmj}^2, \quad (3.49a)$$

$$\sum_{j=0}^{\infty} \tilde{a}_{nmj}^2 = \sum_{j=0}^{\infty} \tilde{a}_{nnj} \tilde{a}_{mmj} = \tilde{a}_{nnmm}, \quad (3.49b)$$

where the mass ratios  $r_j = M_j/M_n$  and the quartic KK Goldstone coupling coefficient  $\tilde{a}_{nnmm} = \llbracket \tilde{f}_n \tilde{f}_n \tilde{f}_m \tilde{f}_m \rrbracket$ . We note that the above condition (3.49a) is the same as the condition (3.38) (which we derived earlier for the elastic scattering channel) and thus the inelastic condition (3.49a) can be directly inferred from squaring Eq.(3.6) followed by summing over the KK index  $j$ . [We recall that Eq.(3.6) is the condition to hold the GAET (3.4) at the level of 3-point KK amplitudes.] The condition (3.49b) holds because it is just the special case of the basic relations (D.2a) which rely on the completeness condition (D.1a).

Eq.(3.48) demonstrates that the inelastic scattering amplitudes (3.44a) and (3.44b) are equal due to the color Jacobi identity (3.32). This explicitly establishes the GAET for 4-point inelastic KK scattering amplitudes:

$$\mathcal{T}_0[A_L^{an} A_L^{bn} \rightarrow A_L^{cm} A_L^{dm}] = \tilde{\mathcal{T}}_0[A_5^{an} A_5^{bn} \rightarrow A_5^{cm} A_5^{dm}]. \quad (3.50)$$

According to our discussion below Eq.(3.49), we stress that *the validity of the inelastic GAET (3.50) is guaranteed by the validity of the most fundamental GAET (3.4) at the level of 3-point KK amplitudes.*

As a consistency check, we take the flat 5d limit  $k \rightarrow 0$  and derive the following inelastic KK scattering amplitudes for each kinematic channel:

$$\mathcal{K}_s^{\text{in}0} = -c_\theta + (r^2 - r^{-2})c_\theta, \quad \tilde{\mathcal{K}}_s^{\text{in}0} = -c_\theta, \quad (3.51a)$$

$$\mathcal{K}_t^{\text{in}0} = \frac{3 - c_\theta}{1 + c_\theta} + (r^2 - r^{-2})c_\theta, \quad \tilde{\mathcal{K}}_t^{\text{in}0} = \frac{3 - c_\theta}{1 + c_\theta}, \quad (3.51b)$$

$$\mathcal{K}_u^{\text{in}0} = -\frac{3 + c_\theta}{1 - c_\theta} + (r^2 - r^{-2})c_\theta, \quad \tilde{\mathcal{K}}_u^{\text{in}0} = -\frac{3 + c_\theta}{1 - c_\theta}. \quad (3.51c)$$

Thus, the inelastic scattering amplitudes of the longitudinal KK gauge bosons and of the KK Goldstone bosons differ by the same amount in each kinematic channel:

$$\mathcal{K}_i^{\text{in}0} - \tilde{\mathcal{K}}_i^{\text{in}0} = (r^2 - r^{-2})c_\theta, \quad (3.52)$$

where each individual channel is marked by the kinematic index  $i \in \{s, t, u\}$ . Hence, the difference (3.52) gives zero contribution to the inelastic KK scattering amplitudes (3.44a)-(3.44b) due to the color Jacobi identity (3.32).

Then, we study the general inelastic scattering process  $(n, m) \rightarrow (\ell, q)$  with all the external KK indices  $(n, m, \ell, q)$  unequal. Under the high energy expansion, we compute its 4-point leading-order inelastic scattering amplitudes of the longitudinal KK gauge bosons and of the corresponding KK Goldstone bosons, which take the following form:

$$\mathcal{T}_0[A_L^{an} A_L^{bm} \rightarrow A_L^{c\ell} A_L^{dq}] = g^2 (\mathcal{C}_s \mathcal{K}_s^{\text{in}0} + \mathcal{C}_t \mathcal{K}_t^{\text{in}0} + \mathcal{C}_u \mathcal{K}_u^{\text{in}0}), \quad (3.53a)$$

$$\tilde{\mathcal{T}}_0[A_5^{an} A_5^{bm} \rightarrow A_5^{cl} A_5^{dq}] = g^2 (\mathcal{C}_s \tilde{\mathcal{K}}_s^{\text{in}0} + \mathcal{C}_t \tilde{\mathcal{K}}_t^{\text{in}0} + \mathcal{C}_u \tilde{\mathcal{K}}_u^{\text{in}0}). \quad (3.53b)$$

In each kinematic channel, we derive the leading-order scattering amplitudes of longitudinal KK gauge bosons as follows:

$$\begin{aligned} \mathcal{K}_s^{\text{in}0} = & \frac{1}{4M_n M_m M_\ell M_q} \left\{ \sum_{j=1}^{\infty} \left[ M_j^{-2} (M_n^2 - M_m^2) (M_\ell^2 - M_q^2) (M_n^2 + M_m^2 + M_\ell^2 + M_q^2) \right. \right. \\ & - c_\theta M_j^4 \left. \right] a_{nmj} a_{\ell qj} + c_\theta \left[ M_\ell^4 + M_q^4 + (M_n^2 + M_m^2) (M_n^2 + M_m^2 + M_q^2) \right. \\ & \left. \left. + M_\ell^2 (M_m^2 + M_n^2 + 2M_q^2) \right] a_{nmlq} \right\}, \end{aligned} \quad (3.54a)$$

$$\begin{aligned} \mathcal{K}_t^{\text{in}0} = & \frac{1}{16M_n M_m M_\ell M_q} \left\{ \sum_{j=1}^{\infty} \left[ -4M_j^4 + 2(5+c_\theta) M_j^2 (M_n^2 + M_m^2 + M_\ell^2 + M_q^2) \right. \right. \\ & \left. \left. - 2(1-c_\theta) M_j^{-2} (M_n^2 - M_q^2) (M_m^2 - M_\ell^2) (M_n^2 + M_m^2 + M_\ell^2 + M_q^2) \right] a_{nqj} a_{m\ell j} \right. \\ & + \frac{16}{1+c_\theta} \sum_{j=0}^{\infty} (M_n^2 + M_q^2 - M_j^2) (M_m^2 + M_\ell^2 - M_j^2) a_{nqj} a_{m\ell j} \\ & \left. - 2 \left[ (M_n^2 + M_q^2) (M_n^2 + 4M_m^2 + 4M_\ell^2 + M_q^2) - c_\theta (M_n^2 + M_m^2 + M_\ell^2 + M_q^2)^2 \right. \right. \\ & \left. \left. + (M_m^2 + M_\ell^2)^2 \right] a_{nmlq} \right\}, \end{aligned} \quad (3.54b)$$

$$\begin{aligned} \mathcal{K}_u^{\text{in}0} = & \frac{1}{16M_n M_m M_\ell M_q} \left\{ \sum_{j=1}^{\infty} \left[ 4M_j^4 - 2(5+c_\theta) M_j^2 (M_n^2 + M_m^2 + M_\ell^2 + M_q^2) \right. \right. \\ & \left. \left. + 2(1+c_\theta) M_j^{-2} (M_n^2 - M_\ell^2) (M_m^2 - M_q^2) (M_n^2 + M_m^2 + M_\ell^2 + M_q^2) \right] a_{n\ell j} a_{mqj} \right. \\ & - \frac{16}{1-c_\theta} \sum_{j=0}^{\infty} (M_n^2 + M_\ell^2 - M_j^2) (M_m^2 + M_q^2 - M_j^2) a_{n\ell j} a_{mqj} \\ & \left. + 2 \left[ (M_n^2 + M_\ell^2) (M_n^2 + 4M_m^2 + M_\ell^2 + 4M_q^2) + c_\theta (M_n^2 + M_m^2 + M_\ell^2 + M_q^2)^2 \right. \right. \\ & \left. \left. + (M_m^2 + M_q^2)^2 \right] a_{nmlq} \right\}. \end{aligned} \quad (3.54c)$$

We further compute the corresponding 4-point leading-order scattering amplitude of KK Goldstone bosons in each channel,

$$\tilde{\mathcal{K}}_s^{\text{in}0} = -c_\theta \sum_{j=0}^{\infty} \tilde{a}_{nmj} \tilde{a}_{\ell qj}, \quad (3.55a)$$

$$\tilde{\mathcal{K}}_t^{\text{in}0} = \frac{3-c_\theta}{1+c_\theta} \sum_{j=0}^{\infty} \tilde{a}_{nqj} \tilde{a}_{m\ell j}, \quad (3.55b)$$

$$\tilde{\mathcal{K}}_u^{\text{in}0} = -\frac{3+c_\theta}{1-c_\theta} \sum_{j=0}^{\infty} \tilde{a}_{n\ell j} \tilde{a}_{mqj}. \quad (3.55c)$$

Then, we demonstrate that in each kinematic channel the above inelastic scattering amplitudes of the longitudinal KK gauge bosons and of the KK Goldstone bosons differ by the same amount:

$$\mathcal{K}_i^{\text{in}0} - \tilde{\mathcal{K}}_i^{\text{in}0} = \frac{M_n^2 + M_m^2 + M_\ell^2 + M_q^2}{4M_n M_m M_\ell M_q} \sum_{j=1}^{\infty} M_j^2 \left[ (1+c_\theta) a_{nqj} a_{m\ell j} - (1-c_\theta) a_{n\ell j} a_{mqj} \right], \quad (3.56)$$



where we have imposed the following sum rule conditions,

$$a_{nmlq} = \sum_{j=0}^{\infty} a_{nmj} a_{\ell qj} = \sum_{j=0}^{\infty} a_{nlj} a_{mqj} = \sum_{j=0}^{\infty} a_{nqj} a_{\ell mj}, \quad (3.57a)$$

$$a_{nmlq} = \frac{1}{M_n^2 + M_m^2 + M_\ell^2 + M_q^2} \sum_{j=1}^{\infty} M_j^2 (a_{nmj} a_{\ell qj} + a_{nlj} a_{mqj} + a_{nqj} a_{\ell mj}), \quad (3.57b)$$

$$(M_n^2 - M_m^2)(M_\ell^2 - M_q^2) \sum_{j=1}^{\infty} \frac{a_{nmj} a_{\ell qj}}{M_j^2} = \sum_{j=0}^{\infty} M_j^2 (a_{nqj} a_{\ell mj} - a_{nlj} a_{mqj}), \quad (3.57c)$$

$$(M_n^2 - M_\ell^2)(M_m^2 - M_q^2) \sum_{j=1}^{\infty} \frac{a_{nlj} a_{mqj}}{M_j^2} = \sum_{j=0}^{\infty} M_j^2 (a_{nqj} a_{\ell mj} - a_{nmj} a_{\ell qj}), \quad (3.57d)$$

$$(M_n^2 - M_q^2)(M_\ell^2 - M_m^2) \sum_{j=1}^{\infty} \frac{a_{nqj} a_{\ell mj}}{M_j^2} = \sum_{j=0}^{\infty} M_j^2 (a_{nmj} a_{\ell qj} - a_{nlj} a_{mqj}). \quad (3.57e)$$

The relations (3.57a) can be readily proven as in Eq.(D.2a) which relies on the completeness condition (D.1a). The above formula (3.57b) is derived as in Eq.(D.11) of Appendix D and the last three conditions of (3.57) are derived around Eqs.(D.28)-(D.29), where both derivations crucially rely on the condition (3.5) [which holds the GAET (3.4) for the fundamental 3-point KK amplitudes]. With the above, we explicitly prove the GAET for the general 4-point inelastic KK scattering amplitudes with all external KK indices being different:

$$\mathcal{T}_0[A_L^{an} A_L^{bm} \rightarrow A_L^{c\ell} A_L^{dq}] = \tilde{\mathcal{T}}_0[A_5^{an} A_5^{bm} \rightarrow A_5^{c\ell} A_5^{dq}]. \quad (3.58)$$

Here we find again that the validity of the GAET (3.58) for the 4-point inelastic KK scattering amplitudes is guaranteed by the validity of the fundamental 3-point-level GAET (3.4).

Taking the flat 5d space limit  $k \rightarrow 0$ , we examine the inelastic KK scattering process  $(n, 2n) \rightarrow (3n, 4n)$  with  $n > 0$  for instance. The relevant scattering amplitudes of longitudinal KK gauge bosons and its corresponding KK Goldstone bosons can be derived from Eqs.(3.54)-(3.55) as follows:

$$\mathcal{K}_s^{\text{in}0} = \frac{3}{32}(35 + 43c_\theta), \quad \tilde{\mathcal{K}}_s^{\text{in}0} = -\frac{1}{2}c_\theta, \quad (3.59a)$$

$$\mathcal{K}_t^{\text{in}0} = \frac{451 + 468c_\theta + 145c_{2\theta}}{64(1 + c_\theta)}, \quad \tilde{\mathcal{K}}_t^{\text{in}0} = \frac{3 - c_\theta}{2(1 + c_\theta)}, \quad (3.59b)$$

$$\mathcal{K}_u^{\text{in}0} = -\frac{31 - 48c_\theta + 145c_{2\theta}}{64(1 - c_\theta)}, \quad \tilde{\mathcal{K}}_u^{\text{in}0} = -\frac{3 + c_\theta}{2(1 - c_\theta)}. \quad (3.59c)$$

From the above, we explicitly compute the difference between  $\mathcal{K}_i^{\text{in}0}$  and  $\tilde{\mathcal{K}}_i^{\text{in}0}$  which gives the same amount for all three channels:

$$\mathcal{K}_i^{\text{in}0} - \tilde{\mathcal{K}}_i^{\text{in}0} = \frac{5}{32}(21 + 29c_\theta), \quad (3.60)$$

where the index  $i \in \{s, t, u\}$  denotes the three kinematic channels. We have verified that the above difference (3.60) fully agrees with the flat 5d limit ( $k \rightarrow 0$ ) of Eq.(3.56) for the choice of external KK indices  $(n, m, \ell, q) = (n, 2n, 3n, 4n)$ .

Finally, we analyze the inelastic KK scattering channel  $(0, 0) \rightarrow (n, n)$ . Under high energy expansion, we compute the leading-order scattering amplitudes of the KK gauge bosons and of the KK Goldstone bosons as follows:

$$\mathcal{T}_0[A_{\pm}^{a0} A_{\mp}^{b0} \rightarrow A_L^{cn} A_L^{dn}] = g^2 (\mathcal{C}_s \mathcal{K}_s^{\text{in}0} + \mathcal{C}_t \mathcal{K}_t^{\text{in}0} + \mathcal{C}_u \mathcal{K}_u^{\text{in}0}), \quad (3.61a)$$

$$\tilde{\mathcal{T}}_0[A_{\pm}^{a0} A_{\mp}^{b0} \rightarrow A_5^{cn} A_5^{dn}] = g^2 (\mathcal{C}_s \tilde{\mathcal{K}}_s^{\text{in}0} + \mathcal{C}_t \tilde{\mathcal{K}}_t^{\text{in}0} + \mathcal{C}_u \tilde{\mathcal{K}}_u^{\text{in}0}). \quad (3.61b)$$

In the above, the sub-amplitudes  $\{\mathcal{K}_i^{\text{in}0}\}$  and  $\{\tilde{\mathcal{K}}_i^{\text{in}0}\}$  for each kinematic channel are derived as follows:

$$\mathcal{K}_s^{\text{in}0} = -\tilde{\mathcal{K}}_s^{\text{in}0} = 0, \quad (3.62a)$$

$$\mathcal{K}_t^{\text{in}0} = -\tilde{\mathcal{K}}_t^{\text{in}0} = -(1 - c_\theta) f_0^2, \quad (3.62b)$$

$$\mathcal{K}_u^{\text{in}0} = -\tilde{\mathcal{K}}_u^{\text{in}0} = (1 + c_\theta) f_0^2, \quad (3.62c)$$

which establish the equality between the two leading-order inelastic KK scattering amplitudes (3.61a) and (3.61b). Hence, we have GAET for the 4-point inelastic KK scattering amplitudes of  $(0, 0) \rightarrow (n, n)$ :

$$\mathcal{T}_0[A_{\pm}^{a0} A_{\mp}^{b0} \rightarrow A_L^{cn} A_L^{dn}] = -\tilde{\mathcal{T}}_0[A_{\pm}^{a0} A_{\mp}^{b0} \rightarrow A_5^{cn} A_5^{dn}]. \quad (3.63)$$

As a consistency check, we can take the flat 5d limit  $k \rightarrow 0$  and obtain:

$$\mathcal{K}_s^{\text{in}0} = -\tilde{\mathcal{K}}_s^{\text{in}0} = 0, \quad (3.64a)$$

$$\mathcal{K}_t^{\text{in}0} = -\tilde{\mathcal{K}}_t^{\text{in}0} = -(1 - c_\theta), \quad (3.64b)$$

$$\mathcal{K}_u^{\text{in}0} = -\tilde{\mathcal{K}}_u^{\text{in}0} = (1 + c_\theta). \quad (3.64c)$$

These agree with the previous inelastic KK scattering amplitudes derived for the compactified flat 5d space with orbifold  $S^1/\mathbb{Z}_2$  [27].

### 3.2.2 Warped GRET: from 3-Point to 4-Point KK Amplitudes

In this subsection, we first derive the 4-point elastic scattering amplitude of the spin-0 gravitational KK Goldstone bosons  $\phi_n \phi_n \rightarrow \phi_n \phi_n$  for the warped 5d GR under the  $S^1/\mathbb{Z}_2$  compactification. We use the Feynman rules given in Appendix C.2 and compute explicitly the KK Goldstone boson amplitude to the leading order of high energy expansion:

$$\tilde{\mathcal{M}}_0[4\phi_n] = \frac{\kappa^2 s (7 + c_{2\theta})^2 \csc^2 \theta}{64} \sum_{j=0}^{\infty} \tilde{\beta}_{nnj}^2, \quad (3.65)$$

where the leading-order scattering amplitude  $\tilde{\mathcal{M}}_0[4\phi_n] \equiv \tilde{\mathcal{M}}_0[\phi_n \phi_n \rightarrow \phi_n \phi_n]$  is of  $O(E^2)$ . To explicitly demonstrate the GRET for 4-point KK scattering amplitudes, we compare the

leading-order scattering amplitude (3.65) of the gravitational KK Goldstone bosons with the corresponding longitudinal KK graviton amplitude  $\mathcal{M}_0[4h_L^n] \equiv \mathcal{M}_0[h_L^n h_L^n \rightarrow h_L^n h_L^n]$  [49],

$$\mathcal{M}_0[4h_L^n] = \frac{\kappa^2 s (7 + c_{2\theta})^2 \csc^2 \theta}{2304} \sum_{j=1}^{\infty} \hat{r}_j^2 (\hat{r}_j^6 - 12\hat{r}_j^2 + 11) \alpha_{nnj}^2, \quad (3.66)$$

where the KK mass ratio  $\hat{r}_j = \mathbb{M}_j / \mathbb{M}_n$ . To hold the KK GRET (2.103) requires the equality between the two leading-order scattering amplitudes (3.66) and (3.65),

$$\mathcal{M}_0[4h_L^n] = \widetilde{\mathcal{M}}_0[4\phi_n], \quad (3.67)$$

which gives the following nontrivial sum-rule condition:

$$\sum_{j=1}^{\infty} \hat{r}_j^2 (\hat{r}_j^6 - 12\hat{r}_j^2 + 11) \alpha_{nnj}^2 = 36 \sum_{j=0}^{\infty} \tilde{\beta}_{nnj}^2. \quad (3.68)$$

To prove the above condition (3.68) for the 4-point GRET (3.67), we make use of the important fundamental 3-point GRET condition (3.22) to replace the coupling  $\tilde{\beta}_{nnj}$  by  $\alpha_{nnj}$  and convert Eq.(3.68) to the following form:

$$\sum_{j=1}^{\infty} (8\hat{r}_j^6 - 40\hat{r}_j^4 + 59\hat{r}_j^2 - 36) \alpha_{nnj}^2 = 0. \quad (3.69)$$

Then, the rest of the proof is to justify the condition (3.69) which can be achieved by using the properties of the 5d wavefunctions and the resultant relations. To prove the condition (3.69), we first derive the following sum rule identities:

$$\sum_{j=0}^{\infty} \alpha_{nnj}^2 = \alpha_{nnnn}, \quad (3.70a)$$

$$\sum_{j=0}^{\infty} \hat{r}_j^2 \alpha_{nnj}^2 = \frac{4}{3} \alpha_{nnnn}, \quad (3.70b)$$

where the quartic KK coupling coefficient  $\alpha_{nnnn}$  is defined in Eq.(C.11a) of Appendix D. The identities (3.70a) and (3.70b) are proved in Eqs.(D.3b) and (D.14) respectively. Combining the identities (3.70a) and (3.70b), we eliminate the quartic KK coupling coefficient  $\alpha_{nnnn}$  and obtain:

$$\sum_{j=0}^{\infty} \left(1 - \frac{3}{4} \hat{r}_j^2\right) \alpha_{nnj}^2 = 0. \quad (3.71)$$

Using Eq.(3.71), we can further simplify the identity (3.69) as follows:

$$\sum_{j=0}^{\infty} \left(\hat{r}_j^6 - 5\hat{r}_j^4 + \frac{16}{3}\right) \alpha_{nnj}^2 = 0, \quad (3.72)$$

which is proved in Eq.(D.35) of Appendix D by using the equation of motions (2.52)-(2.53). This completes the proof of the identity (3.69) and thus establishes the GRET (3.67) for the 4-point gravitational KK scattering amplitudes.

Next, we consider a mixed inelastic KK graviton scattering process  $h_0^{+2} h_0^{-2} \rightarrow h_n^{+1} h_n^{-1}$  and its corresponding KK Goldstone boson scattering channel  $h_0^{+2} h_0^{-2} \rightarrow \mathcal{V}_n^{+1} \mathcal{V}_n^{-1}$ , where

the initial state includes two massless zero-mode gravitons with helicities  $\pm 2$ , and both the final state KK gravitons and KK vector Goldstones have helicities  $\pm 1$ . By explicit calculation, we derive the following KK scattering amplitudes:

$$\mathcal{M}[h_0^{+2}h_0^{-2} \rightarrow h_n^{+1}h_n^{-1}] = -\frac{\kappa^2(1+c_\theta)(1-c_\theta)^3(12M_n^2+s_0)s_0}{16(4c_\theta^2M_n^2+s_\theta^2s_0)}\alpha_{000}\alpha_{nn0}, \quad (3.73a)$$

$$\widetilde{\mathcal{M}}[h_0^{+2}h_0^{-2} \rightarrow \mathcal{V}_n^{+1}\mathcal{V}_n^{-1}] = \frac{\kappa^2(1+c_\theta)(1-c_\theta)^3(3s_0+2M_n^2)(s_0-4M_n^2)}{48(4c_\theta^2M_n^2+s_\theta^2s_0)}\alpha_{000}\tilde{\alpha}_{nn0}. \quad (3.73b)$$

Making high energy expansion, we derive the leading-order KK scattering amplitudes:

$$\mathcal{M}_0[h_0^{+2}h_0^{-2} \rightarrow h_n^{+1}h_n^{-1}] = -\frac{\kappa^2s_0}{16}(1-c_\theta)^2\alpha_{000}\alpha_{nn0}, \quad (3.74a)$$

$$\widetilde{\mathcal{M}}_0[h_0^{+2}h_0^{-2} \rightarrow \mathcal{V}_n^{+1}\mathcal{V}_n^{-1}] = +\frac{\kappa^2s_0}{16}(1-c_\theta)^2\alpha_{000}\tilde{\alpha}_{nn0}, \quad (3.74b)$$

where  $\alpha_{000} = \alpha_{nn0} = \tilde{\alpha}_{nn0} = \mathbf{u}_0$  and  $\mathbf{u}_0$  is given by Eq.(2.58a). We see that the above KK graviton scattering amplitude equals the corresponding KK Goldstone scattering amplitude at the leading order after including an overall factor  $(-i)^2 = -1$  from the (tree-level) modification factor (2.85a) on its right-hand side,

$$\mathcal{M}_0[h_0^{+2}h_0^{-2} \rightarrow h_n^{+1}h_n^{-1}] = -\widetilde{\mathcal{M}}_0[h_0^{+2}h_0^{-2} \rightarrow \mathcal{V}_n^{+1}\mathcal{V}_n^{-1}]. \quad (3.75)$$

This agrees with the GRET (2.86).

Finally, we stress that in the proof of the condition (3.68) for the 4-point GRET (3.67), the basic 3-point GRET condition (3.22) plays a key role. Hence, we conclude that our proof of the most basic 3-point GRET identity (3.20) or its equivalent sum rule condition (3.22) does automatically guarantee the validity of the 4-point GRET identity (3.67) or its equivalent sum rule condition (3.68). We stress that a major goal of the present work is to demonstrate that the validities of the GAET and GRET for the 4-point KK scattering amplitudes can be proven from the corresponding GAET and GRET identities (sum rule conditions) at the level of the 3-point KK scattering amplitudes. We further conjecture that based upon the most fundamental GAET and GRET for 3-point KK scattering amplitudes, the GAET and GRET for general  $N$ -point ( $N \geq 5$ ) GAET and GRET should also hold. We leave this generalization to the future works.

## 4 Extended Double-Copy for Warped KK Equivalence Theorems

In this section, we study the double-copy construction of the massive gravitational KK scattering amplitudes at tree level from the corresponding massive KK gauge scattering amplitudes for the warped 5d gauge and gravity theories with  $S^1/\mathbb{Z}_2$  compactification. This is nontrivial and challenging since it was proved [27] that the double-copy construction for  $N$ -point ( $N \geq 4$ ) massive gauge/gravity KK scattering amplitudes can directly work out only for toroidal compactifications with flat extra dimensions. In this section, we demonstrate that the double-copy can be realized for 3-point ( $N = 3$ ) KK gauge/gravity scattering

amplitudes without high energy expansion. Then, we further construct the double-copy for 4-point KK gauge/gravity scattering amplitudes at the leading order of high energy expansion.

#### 4.1 Double-Copy Construction of 3-Point Amplitudes and Warped GRET

We have computed the most general 3-point on-shell scattering amplitudes of KK gauge bosons in Eq.(3.1), which takes the following form:

$$\mathcal{T}[\{\epsilon_j\}] = g f^{abc} \mathcal{N}[\{\epsilon_j\}], \quad (4.1a)$$

$$\mathcal{N}[\{\epsilon_j\}] = -i2a_{n_1 n_2 n_3} [(\epsilon_1 \cdot \epsilon_2)(\epsilon_3 \cdot p_1) + (\epsilon_2 \cdot \epsilon_3)(\epsilon_1 \cdot p_2) + (\epsilon_3 \cdot \epsilon_1)(\epsilon_2 \cdot p_3)], \quad (4.1b)$$

where  $\{\epsilon_j\} = \{\epsilon_{\lambda_j}^\mu\}$  denotes the polarization vectors for the three external KK gauge bosons  $\{A_{\lambda_j}^{an_j}\}$  with  $j = 1, 2, 3$ . In parallel, we can reexpress the most relevant 3-point scattering amplitudes (3.3) of KK Goldstone bosons with gauge bosons:

$$\tilde{\mathcal{T}}[A_5^{an_1} A_5^{bn_2} A_\pm^{cn_3}] = g f^{abc} \tilde{\mathcal{N}}[\epsilon_3], \quad \tilde{\mathcal{N}}[\epsilon_3] = -i2(\epsilon_3 \cdot p_1) \tilde{a}_{n_1 n_2 n_3}, \quad (4.2)$$

where the cubic KK Goldstone-gauge coupling coefficient  $\tilde{a}_{n_1 n_2 n_3}$  is given by Eq.(C.3).

The conventional BCJ double-copy method [24][25] applies to  $N$ -point scattering amplitudes of massless gauge bosons and gravitons with  $N \geq 4$ . For the case of  $N = 3$ , the massless double-copy can be directly realized after imposing on-shell conditions on the 3-point gauge/gravity amplitudes. For the present study, we first extend it to the 3-point ( $N = 3$ ) massive KK scattering amplitudes of the warped gauge/gravity theories. This is more nontrivial as the mass spectra of the KK gauge bosons and KK gravitons also differ from each other. Since each on-shell 3-point gauge amplitude (4.1) or (4.2) only has a single color factor and numerator without internal propagator, we may formulate the 3-point color-kinematics (CK) correspondence and the gauge/gravity coupling correspondence as follows:

$$f^{abc} \longrightarrow \mathcal{N}[\{\epsilon_j\}], \quad f^{abc} \longrightarrow \tilde{\mathcal{N}}[\epsilon_3], \quad g \longrightarrow -\frac{\kappa}{4}. \quad (4.3)$$

Moreover, at each KK level- $n$ , we should set up the KK-mass correspondence (replacement),  $M_n \rightarrow \mathbb{M}_n$ , which means to replace each mass-eigenvalue  $M_n$  of KK gauge bosons [determined by Eq.(2.20)] by mass-eigenvalue  $\mathbb{M}_n$  of KK gravitons [determined by Eq.(2.57)].

With the above, we make the double-copy construction of the on-shell 3-point KK graviton scattering amplitude from the product of the corresponding on-shell 3-point KK gauge boson scattering amplitudes (4.1):

$$\mathcal{M}^{\text{dc}}[\{e_j, \epsilon_j\}] = -\frac{\kappa \alpha_{n_1 n_2 n_3}}{4a_{n_1 n_2 n_3}^2} \frac{1}{8} \left( \mathcal{N}[\{e_j\}] \times \mathcal{N}[\{\epsilon_j\}] + \{e_j \leftrightarrow \epsilon_j\} \right), \quad (4.4)$$

where  $\{e_j\} = \{e_{\lambda_j}^\mu\}$  denotes another set of polarization vectors in parallel to the polarization vectors  $\{\epsilon_j\} = \{\epsilon_{\lambda_j}^\mu\}$ , and the overall factor  $1/8 = 1/2^3$  arises from symmetrizing the two polarization vectors associated with each external KK graviton polarization. The two polarization vectors  $\{e_j, \epsilon_j\}$  should compose the polarization tensor  $\varepsilon_j^{\mu\nu} = \varepsilon_{\lambda_j}^{\mu\nu}$  of the external

KK graviton- $j$  according to Eq.(A.9). Thus, because the graviton polarization tensor is traceless  $\varepsilon_{\hat{\lambda}_j}^{\mu\nu}\eta_{\mu\nu}=0$ , we have the corresponding products vanish:

$$e_j^\pm \cdot \epsilon_j^\pm = 0, \quad e_j^\pm \cdot \epsilon_j^L = 0, \quad e_j^\pm \cdot \epsilon_j^\mp + e_j^L \cdot \epsilon_j^L = 0. \quad (4.5)$$

This helps to simplify our explicit results of double-copy. Using the polarization tensors Eq.(A.9) for the three external KK graviton states, we can express the double-copy formula of the 3-point KK graviton scattering amplitude as follows:

$$\begin{aligned} \mathcal{M}^{\text{dc}}[h_{n_1}^{\hat{\lambda}_1} h_{n_2}^{\hat{\lambda}_2} h_{n_3}^{\hat{\lambda}_3}] &= \sum_{\lambda_j, \lambda'_j} \left( \prod_j C_{\lambda_j, \lambda'_j}^{\hat{\lambda}_j} \right) \mathcal{M}[\{e_i\}^{\lambda_j}, \{\epsilon_i\}^{\lambda'_j}] \\ &= -\frac{\kappa \alpha_{n_1 n_2 n_3}}{4 a_{n_1 n_2 n_3}^2} \sum_{\lambda_j, \lambda'_j} \left( \prod_j C_{\lambda_j, \lambda'_j}^{\hat{\lambda}_j} \right) \mathcal{N}[\{e_{\lambda_j}\}] \times \mathcal{N}[\{\epsilon_{\lambda'_j}\}] \\ &\equiv \frac{\kappa \alpha_{n_1 n_2 n_3}}{4} \sum_{\lambda_j, \lambda'_j} \left( \prod_j C_{\lambda_j, \lambda'_j}^{\hat{\lambda}_j} \right) \bar{\mathcal{M}}^{\text{dc}}, \end{aligned} \quad (4.6)$$

where the helicity index  $\hat{\lambda}_j = \{\pm 2, \pm 1, 0\} = \{\pm 2, \pm 1, L\}$  labels the 5 helicity states of each external massive KK graviton. The polarization tensors of the external KK graviton states are defined as follows:

$$\varepsilon_{\hat{\lambda}_j}^{\mu\nu} = \sum_{\lambda_j, \lambda'_j} C_{\lambda_j, \lambda'_j}^{\hat{\lambda}_j} \epsilon_{\lambda_j}^\mu \epsilon_{\lambda'_j}^\nu, \quad (4.7a)$$

$$C_{\pm 1, \pm 1}^{\pm 2} = 1, \quad C_{\pm 1, 0}^{\pm 1} = C_{0, \pm 1}^{\pm 1} = \frac{1}{\sqrt{2}}, \quad C_{\pm 1, \mp 1}^0 = \frac{1}{\sqrt{6}}, \quad C_{0, 0}^0 = \sqrt{\frac{2}{3}}. \quad (4.7b)$$

Then, we use the double-copy method to compute the on-shell 3-point scattering amplitudes of KK gravitons and derive the following formula:

$$\begin{aligned} \bar{\mathcal{M}}^{\text{dc}} &= (e_1 \epsilon_3)(e_2 p_3)(e_3 p_2)(\epsilon_1 \epsilon_2) + (e_1 e_3)(e_2 p_3)(p_2 \epsilon_3)(\epsilon_1 \epsilon_2) + (e_1 p_3)(e_2 p_3)(e_3 \epsilon_2)(\epsilon_1 \epsilon_3) \\ &+ (e_1 \epsilon_2)(e_2 p_3)(e_3 p_2)(\epsilon_1 \epsilon_3) + (e_1 p_3)(e_2 p_3)(e_3 \epsilon_1)(\epsilon_2 \epsilon_3) + (e_1 e_3)(e_2 p_3)(p_3 \epsilon_1)(\epsilon_2 \epsilon_3) \\ &+ (e_1 \epsilon_2)(e_2 p_3)(e_3 \epsilon_1)(p_2 \epsilon_3) - 2(e_1 \epsilon_2)(e_2 \epsilon_1)(e_3 p_2)(p_2 \epsilon_3) - 2(e_1 e_2)(e_3 p_2)(p_2 \epsilon_3)(\epsilon_1 \epsilon_2) \\ &+ (e_1 e_3)(e_2 \epsilon_1)(p_3 \epsilon_2)(p_2 \epsilon_3) + (e_1 e_2)(e_3 \epsilon_1)(p_3 \epsilon_2)(p_2 \epsilon_3) - (e_1 p_3)(e_2 \epsilon_3)(e_3 p_2)(\epsilon_1 \epsilon_2) \\ &- (e_1 p_3)(e_2 \epsilon_1)(e_3 p_2)(\epsilon_2 \epsilon_3) - (e_1 p_3)(e_2 \epsilon_1)(e_3 \epsilon_2)(p_2 \epsilon_3) - (e_1 p_3)(e_2 e_3)(p_2 \epsilon_3)(\epsilon_1 \epsilon_2) \\ &+ (e_1 \epsilon_3)(e_2 p_3)(e_3 \epsilon_2)(p_3 \epsilon_1) - (e_1 \epsilon_2)(e_2 \epsilon_3)(e_3 p_2)(p_3 \epsilon_1) - (e_1 e_2)(e_3 p_2)(p_3 \epsilon_1)(\epsilon_2 \epsilon_3) \\ &- (e_1 \epsilon_2)(e_2 e_3)(p_3 \epsilon_1)(p_2 \epsilon_3) - (e_1 e_2)(e_3 \epsilon_2)(p_3 \epsilon_1)(p_2 \epsilon_3) - 2(e_1 p_3)(e_2 \epsilon_3)(e_3 \epsilon_2)(p_3 \epsilon_1) \\ &- 2(e_1 p_3)(e_2 e_3)(p_3 \epsilon_1)(\epsilon_2 \epsilon_3) + (e_1 p_3)(e_2 \epsilon_3)(e_3 \epsilon_1)(p_3 \epsilon_2) + (e_1 p_3)(e_2 e_3)(p_3 \epsilon_2)(\epsilon_1 \epsilon_3) \\ &+ (e_1 e_2)(e_3 p_2)(p_3 \epsilon_2)(\epsilon_1 \epsilon_3) + (e_1 \epsilon_3)(e_2 \epsilon_1)(e_3 p_2)(p_3 \epsilon_2) - 2(e_1 \epsilon_3)(e_2 p_3)(e_3 \epsilon_1)(p_3 \epsilon_2) \\ &- 2(e_1 e_3)(e_2 p_3)(p_3 \epsilon_2)(\epsilon_1 \epsilon_3) + (e_1 \epsilon_3)(e_2 e_3)(p_3 \epsilon_1)(p_3 \epsilon_2) + (e_1 e_3)(e_2 \epsilon_3)(p_3 \epsilon_1)(p_3 \epsilon_2), \end{aligned} \quad (4.8)$$

where we have used the shorthand notation  $(ab) \equiv (a \cdot b)$ , and we also make the replacement  $M_n \rightarrow \mathbb{M}_n$  for each KK mass.

We compare the above double-copy formula (4.8) of the general 3-point KK graviton scattering amplitude with the explicit calculation result of the same KK graviton scattering

amplitude (C.15) in Appendix C.2. We find that they fully agree with each other after imposing the correspondence (4.3).

With the above, we then analyze the 3-point on-shell scattering amplitude of KK gravitons at the leading order of high energy expansion and establish the double-copy construction of the KK gravitational equivalence theorem (GRET) from the KK gauge-theory equivalence theorem (GAET) at the level of the most fundamental 3-point KK gauge/gravity scattering amplitudes. We consider a 3-point KK graviton scattering amplitude containing two external KK gravitons with helicities  $\pm 1$  and one external KK graviton with helicities  $\pm 2$ , or, including two external longitudinal polarization states (with helicity-0) plus one transverse polarization state (with helicities  $\pm 2$ ). Using the KK gauge boson scattering amplitudes (4.1) and (4.6), we can construct two types of 3-point KK graviton amplitudes at the leading order of high energy expansion. One type of the KK graviton scattering amplitudes have helicities  $(\pm 1, \pm 1, \pm 2)$ , and we derive the following:

$$\begin{aligned}\mathcal{M}^{\text{dc}}[h_{n_1}^{\pm 1} h_{n_2}^{\pm 1} h_{n_3}^{\pm 2}] &= -\frac{c_3 \kappa}{4} \left( \mathcal{N}[A_{n_1}^{\pm} A_{n_2}^{\pm} A_{n_3}^{\pm}] \mathcal{N}[A_{n_1}^L A_{n_2}^L A_{n_3}^{\pm}] + \mathcal{N}[A_{n_1}^{\pm} A_{n_2}^L A_{n_3}^{\pm}] \mathcal{N}[A_{n_1}^L A_{n_2}^{\pm} A_{n_3}^{\pm}] \right) \\ &= \mathcal{M}_0[h_{n_1}^{\pm 1} h_{n_2}^{\pm 1} h_{n_3}^{\pm 2}] + O(E^0),\end{aligned}\quad (4.9a)$$

$$\begin{aligned}\mathcal{M}_0^{\text{dc}}[h_{n_1}^{\pm 1} h_{n_2}^{\pm 1} h_{n_3}^{\pm 2}] &= -\frac{c_3 \kappa}{4} \mathcal{N}_0[A_{n_1}^{\pm} A_{n_2}^{\pm} A_{n_3}^{\pm}] \mathcal{N}_0[A_{n_1}^L A_{n_2}^L A_{n_3}^{\pm}] \\ &= c_3 a_{n_1 n_2 n_3}^2 \frac{\kappa(1 - \hat{r}_{13}^2 - \hat{r}_{23}^2)}{2\hat{r}_{13}\hat{r}_{23}} \left[ (\epsilon_1 \cdot \epsilon_2)(\epsilon_3 \cdot p_1)^2 + (\epsilon_2 \cdot \epsilon_3)(\epsilon_1 \cdot p_2)(\epsilon_3 \cdot p_1) + (\epsilon_3 \cdot \epsilon_1)(\epsilon_2 \cdot p_3)(\epsilon_3 \cdot p_1) \right],\end{aligned}\quad (4.9b)$$

where  $c_3$  is a normalization factor of couplings and will be determined shortly. As before we denote the mass ratio  $\hat{r}_{ij} = \mathbb{M}_{n_i}/\mathbb{M}_{n_j}$ . We note that on the right-hand side of Eq.(4.9a) the second amplitude product has only subleading contributions of  $O(E^0)$ , instead of the leading-order contributions of  $O(E^2)$ .

Another type of KK graviton scattering amplitudes have helicities  $(L, L, \pm 2)$ , where  $L$  stands for longitudinal polarization with helicity-0. We construct its exact tree-level scattering amplitude  $\mathcal{M}[h_{n_1}^L h_{n_2}^L h_{n_3}^{\pm 2}]$  and derive its leading-order amplitude  $\mathcal{M}_0[h_{n_1}^L h_{n_2}^L h_{n_3}^{\pm 2}]$  as follows:

$$\begin{aligned}\mathcal{M}^{\text{dc}}[h_{n_1}^L h_{n_2}^L h_{n_3}^{\pm 2}] &= -\frac{c_3 \kappa}{4} \frac{1}{3} \left\{ 2(\mathcal{N}_0[A_{n_1}^L A_{n_2}^L A_{n_3}^{\pm}])^2 + \mathcal{N}_0[A_{n_1}^+ A_{n_2}^- A_{n_3}^{\pm}] \mathcal{N}_0[A_{n_1}^- A_{n_2}^+ A_{n_3}^{\pm}] \right. \\ &\quad \left. + \mathcal{N}_0[A_{n_1}^+ A_{n_2}^+ A_{n_3}^{\pm}] \mathcal{N}_0[A_{n_1}^- A_{n_2}^- A_{n_3}^{\pm}] + 2\mathcal{N}_0[A_{n_1}^+ A_{n_2}^L A_{n_3}^{\pm}] \mathcal{N}_0[A_{n_1}^- A_{n_2}^L A_{n_3}^{\pm}] \right\},\end{aligned}\quad (4.10a)$$

$$\begin{aligned}\mathcal{M}_0^{\text{dc}}[h_{n_1}^L h_{n_2}^L h_{n_3}^{\pm 2}] &= -\frac{c_3 \kappa}{4} \left\{ 2(\mathcal{N}_0[A_{n_1}^L A_{n_2}^L A_{n_3}^{\pm}])^2 + \mathcal{N}_0[A_{n_1}^+ A_{n_2}^- A_{n_3}^{\pm}] \mathcal{N}_0[A_{n_1}^- A_{n_2}^+ A_{n_3}^{\pm}] \right\} \\ &= c_3 a_{n_1 n_2 n_3}^2 \frac{\kappa(\epsilon_3 \cdot p_1)^2}{6} \left[ 2 + \left( \frac{1 - \hat{r}_{13}^2 - \hat{r}_{23}^2}{\hat{r}_{13}\hat{r}_{23}} \right)^2 \right],\end{aligned}\quad (4.10b)$$

where the KK mass ratios  $\hat{r}_{13} = \mathbb{M}_{n_1}/\mathbb{M}_{n_3}$  and  $\hat{r}_{23} = \mathbb{M}_{n_2}/\mathbb{M}_{n_3}$ . We note that under the high energy expansion and on the right-hand side of Eq.(4.10a), the first two terms in the brackets give leading-order contributions and the last two terms are subleading.

Then, for the three-point gravitational amplitudes (3.14b) and (3.19b), we can construct them by double-copy of the two 3-point KK gauge amplitudes:

$$\begin{aligned}\widetilde{\mathcal{M}}_0^{\text{dc}}[\mathcal{V}_{n_1}^{\pm 1} \mathcal{V}_{n_2}^{\pm 1} h_{n_3}^{\pm 2}] &= -\frac{c_3 \kappa}{4} \mathcal{N}_0[A_{n_1}^{\pm} A_{n_2}^{\pm} A_{n_3}^{\pm}] \widetilde{\mathcal{N}}_0[A_{n_1}^5 A_{n_2}^5 A_{n_3}^{\pm}] \\ &= \kappa c_3 a_{n_1 n_2 n_3} \tilde{a}_{n_1 n_2 n_3} \left[ (\epsilon_1 \cdot \epsilon_2)(\epsilon_3 \cdot p_1)^2 + (\epsilon_2 \cdot \epsilon_3)(\epsilon_1 \cdot p_2)(\epsilon_3 \cdot p_1) \right]\end{aligned}$$

$$+ (\epsilon_3 \cdot \epsilon_1)(\epsilon_2 \cdot p_3)(\epsilon_3 \cdot p_1)], \quad (4.11a)$$

$$\widetilde{\mathcal{M}}_0^{\text{dc}}[\phi_{n_1} \phi_{n_2} h_{n_3}^{\pm 2}] = -\frac{c_3 \kappa}{4} (\widetilde{\mathcal{N}}_0[A_{n_1}^5 A_{n_2}^5 A_{n_3}^{\pm}])^2 = \kappa c_3 \tilde{a}_{n_1 n_2 n_3}^2 (\epsilon_3 \cdot p_1)^2. \quad (4.11b)$$

By comparing the KK scattering amplitudes (4.9)-(4.11) obtained via double-copy construction with the same type of amplitudes (3.14a)-(3.14b) and (3.19a)-(3.19b) obtained via explicit Feynman-diagram approach, we find that the two sets of KK scattering amplitudes share exactly *the same kinematic structure* and can equal to each other so long as we require that the KK gauge boson mass be replaced by the corresponding KK graviton mass at each KK level and the KK gauge coupling factor are replaced by the corresponding KK gravitational coupling factor [cf. Eq.(4.3)] together with the following replacements of the trilinear coupling factors:

$$c_3 a_{n_1 n_2 n_3}^2 \rightarrow \alpha_{n_1 n_2 n_3}, \quad c_3 a_{n_1 n_2 n_3} \tilde{a}_{n_1 n_2 n_3} \rightarrow \tilde{\alpha}_{n_1 n_2 n_3}, \quad c_3 \tilde{a}_{n_1 n_2 n_3}^2 \rightarrow \tilde{\beta}_{n_1 n_2 n_3}. \quad (4.12)$$

In general such replacements are nontrivial because the coupling ratios (such as the ratio  $\alpha_{n_1 n_2 n_3}/a_{n_1 n_2 n_3}^2$ ) do not equal one. This means that for the double-copy of KK gauge/gravity theories with warped 5d space, the coupling replacements (4.12) are necessary and nontrivial. We note that in the flat 5d limit of  $k \rightarrow 0$ , these ratios take simple values (cf. Table 1). For instance, choosing  $(n_1, n_2, n_3) = (n, n, j)$  with  $n > 0$ , we have

$$c_3 = \frac{\alpha_{nnj}}{a_{nnj}^2} = \frac{\tilde{\alpha}_{nnj}}{a_{nnj} \tilde{a}_{nnj}} = \frac{\tilde{\beta}_{nnj}}{\tilde{a}_{nnj}^2} = \begin{cases} 1, & j=0, \\ \frac{1}{\sqrt{2}}, & j=2n, \end{cases} \quad (4.13)$$

whereas for the case of  $(n_1, n_2, n_3) = (n, m, |n \pm m|)$  with  $n \neq m$  in the flat 5d limit, we have

$$c_3 = \frac{a_{nm|n \pm m}^2}{\alpha_{nm|n \pm m}} = \frac{a_{nm|n \pm m} \tilde{a}_{nm|n \pm m}}{\tilde{\alpha}_{nm|n \pm m}} = \frac{\tilde{a}_{nm|n \pm m}^2}{\tilde{\beta}_{nm|n \pm m}} = \frac{1}{\sqrt{2}}. \quad (4.14)$$

In fact, as we have checked, for the flat 5d space under the  $S^1$  toroidal compactification without orbifold [27], all the above coupling ratios will become  $c_3 = 1$ .

Finally, using the coupling replacements (4.12), we compare the leading-order KK graviton amplitudes (4.9b) and (4.10b) with the corresponding gravitational KK Goldstone boson amplitudes (4.11a) and (4.11b) respectively. We find that they share exactly the same kinematic structures and obey the GRET,

$$\mathcal{M}_0^{\text{dc}}[h_{n_1}^{\pm 1} h_{n_2}^{\pm 1} h_{n_3}^{\pm 2}] = -\widetilde{\mathcal{M}}_0^{\text{dc}}[\mathcal{V}_{n_1}^{\pm 1} \mathcal{V}_{n_2}^{\pm 1} h_{n_3}^{\pm 2}], \quad (4.15a)$$

$$\mathcal{M}_0^{\text{dc}}[h_{n_1}^L h_{n_2}^L h_{n_3}^{\pm 2}] = \widetilde{\mathcal{M}}_0^{\text{dc}}[\phi_{n_1} \phi_{n_2} h_{n_3}^{\pm 2}], \quad (4.15b)$$

provided the following conditions hold respectively,

$$(\hat{r}_{13}^2 + \hat{r}_{23}^2 - 1)\alpha_{n_1 n_2 n_3} = 2\hat{r}_{13}\hat{r}_{23}\tilde{\alpha}_{n_1 n_2 n_3}, \quad (4.16a)$$

$$[(1 - \hat{r}_{13}^2 - \hat{r}_{23}^2)^2 + 2\hat{r}_{13}^2 \hat{r}_{23}^2]\alpha_{n_1 n_2 n_3} = 6\hat{r}_{13}^2 \hat{r}_{23}^2 \tilde{\beta}_{n_1 n_2 n_3}, \quad (4.16b)$$

where the mass ratios  $\hat{r}_{13} = \mathbb{M}_{n_1}/\mathbb{M}_{n_3}$  and  $\hat{r}_{23} = \mathbb{M}_{n_2}/\mathbb{M}_{n_3}$ . Comparing the above conditions (4.16a)-(4.16b) with the conditions (3.16) and (3.21) (which we obtained by explicit Feynman diagram approach), we find that the identities (4.16a) and (3.16) are identical, and



the same is true for the two identities (4.16b) versus (3.21). This establishes our successful double-copy constructions of the GRET (4.15a) and (4.15b) (in warped KK gravity theory) from the GAET (3.4) (in the warped KK gauge theory).

In summary, we have established the double-copy construction from the GAET to GRET for the 3-point leading-order (LO) KK scattering amplitudes:

$$\begin{array}{ccc}
\mathcal{M}_0^{(3)}[h_L^n] & \xlongequal{\text{GRET}} & \widetilde{\mathcal{M}}_0^{(3)}[\phi_n] \\
\uparrow \text{DC}_3 & & \text{DC}_3 \uparrow \\
\mathcal{T}_0^{(3)}[A_L^n] & \xlongequal{\text{GAET}} & \widetilde{\mathcal{T}}_0^{(3)}[A_5^n]
\end{array} \tag{4.17}$$

In Eq.(4.17), along the horizontal directions, the lower (upper) equality presents the GAET (GRET) of 3-point scattering amplitudes of KK gauge bosons (KK gravitons). Along the direction of a given vertical arrow, each correspondence is established by the double-copy (DC<sub>3</sub>) under the extended massive color-kinematics dualities (4.3) and (4.12). The left vertical arrow represents the double-copy (DC<sub>3</sub>) construction of the 3-point LO KK graviton amplitude  $\mathcal{M}_0^{(3)}[h_L^n]$  from the corresponding LO KK gauge boson amplitude  $\mathcal{T}_0^{(3)}[A_L^n]$ . The right vertical arrow represents the double-copy (DC<sub>3</sub>) construction of the 4-point LO gravitational KK Goldstone boson amplitude  $\widetilde{\mathcal{M}}_0^{(3)}[\phi_n]$  from the corresponding (gauge theory) LO KK Goldstone boson amplitude  $\widetilde{\mathcal{T}}_0^{(3)}[A_5^n]$ . Finally, this picture also holds if we replace the two amplitudes  $\mathcal{M}_0^{(3)}[h_L^n]$  and  $\widetilde{\mathcal{M}}_0^{(3)}[\phi_n]$  in the top row by  $\mathcal{M}_0^{(3)}[h_{\pm 1}^n]$  and  $\widetilde{\mathcal{M}}_0^{(3)}[\mathcal{V}_{\pm 1}^n]$  respectively.

## 4.2 Double-Copy Construction of 4-Point Warped GRET

In this subsection, we study the construction of double-copy for 4-point KK gauge/gravity scattering amplitudes and for their corresponding 4-point KK Goldstone boson scattering amplitudes at the leading order of high energy expansion. As we will demonstrate, under high energy expansion, the leading-order KK amplitudes no longer contain summation of infinite number of KK mass-poles. By construction they can exhibit an effective single massless-pole in each channel and the associated numerators obey the kinematic Jacobi identity, which enable the realization of double-copy at the leading order. In the subsections 4.2.1 and 4.2.2, we study the double-copy constructions for the elastic and inelastic scattering amplitudes of KK gravitons respectively.

### 4.2.1 Construction of 4-Point Elastic KK Graviton Amplitudes

Under the high energy expansion, we can reexpress the 4-point KK scattering amplitudes (3.30a)-(3.30b) of the KK longitudinal gauge bosons and of the KK Goldstone bosons at the leading order:

$$\mathcal{T}_0[4A_L^n] = g^2 \left( \frac{C_s \mathcal{N}_s^0}{s_0} + \frac{C_t \mathcal{N}_t^0}{t_0} + \frac{C_u \mathcal{N}_u^0}{u_0} \right), \tag{4.18a}$$

$$\tilde{\mathcal{T}}_0[4A_5^n] = g^2 \left( \frac{\mathcal{C}_s \tilde{\mathcal{N}}_s^0}{s_0} + \frac{\mathcal{C}_t \tilde{\mathcal{N}}_t^0}{t_0} + \frac{\mathcal{C}_u \tilde{\mathcal{N}}_u^0}{u_0} \right), \quad (4.18b)$$

where the Mandelstam variables  $(s_0, t_0, u_0)$  are defined in Eq.(A.8) of Appendix A and obey the condition  $s_0 + t_0 + u_0 = 0$ . In the above, the kinematic numerators  $(\mathcal{N}_s^0, \mathcal{N}_t^0, \mathcal{N}_u^0)$  and  $(\tilde{\mathcal{N}}_s^0, \tilde{\mathcal{N}}_t^0, \tilde{\mathcal{N}}_u^0)$  are connected to the subamplitudes of Eqs.(3.30a)-(3.30b) and Eqs.(3.35)-(3.36) via the following relations:

$$(\mathcal{N}_s^0, \mathcal{N}_t^0, \mathcal{N}_u^0) = (s_0 \mathcal{K}_s^0, t_0 \mathcal{K}_t^0, u_0 \mathcal{K}_u^0), \quad (4.19a)$$

$$(\tilde{\mathcal{N}}_s^0, \tilde{\mathcal{N}}_t^0, \tilde{\mathcal{N}}_u^0) = (s_0 \tilde{\mathcal{K}}_s^0, t_0 \tilde{\mathcal{K}}_t^0, u_0 \tilde{\mathcal{K}}_u^0), \quad (4.19b)$$

Using the definitions (4.19) and the difference relation (3.39), we derive the generalized gauge transformations between the numerators of elastic scattering amplitudes of KK gauge bosons and of the KK Goldstone bosons:

$$\mathcal{N}_j^0 = \tilde{\mathcal{N}}_j^0 + s_{0j} \Theta, \quad \Theta = -2c_\theta \sum_{i=0}^{\infty} r_i^2 a_{nmi}^2, \quad (4.20)$$

where the massless Mandelstam variable  $s_{0j} \in \{s_0, t_0, u_0\}$ . Inspecting the leading-order numerators (4.19a) and (4.19b) for elastic scattering amplitudes of KK gauge bosons and of KK Goldstone bosons, we find that they satisfy the kinematic Jacobi identities:

$$\sum_j \mathcal{N}_j^0 = \sum_j \tilde{\mathcal{N}}_j^0 + \Theta \sum_j s_{0j} = 0, \quad (4.21a)$$

$$\sum_j \tilde{\mathcal{N}}_j^0 = 0, \quad (4.21b)$$

where the index  $j \in \{s, t, u\}$ . We can readily prove the above second identity (4.21b) by using the formulas (4.19b) and (3.36), whereas the first identity (4.21a) holds because of  $s_0 + t_0 + u_0 = 0$ .

We extend the conventional color-kinematics duality for massless gauge/gravity scattering amplitudes [24][25] to the massive scattering amplitudes of the longitudinal KK gauge bosons and of the corresponding KK Goldstone bosons in Eq.(4.18). We formulate the massive color-kinematics duality as follows:

$$\mathcal{C}_j \longrightarrow \mathcal{N}_j^0, \quad \tilde{\mathcal{C}}_j \longrightarrow \tilde{\mathcal{N}}_j^0, \quad g^2 \longrightarrow -(\kappa/4)^2. \quad (4.22)$$

Thus, we construct the corresponding scattering amplitudes of the longitudinal KK gravitons and of the gravitational KK Goldstone bosons, to the leading-order contributions of  $\mathcal{O}(E^2)$  under the high energy expansion. At each KK level- $n$ , we also set up the KK-mass correspondence (replacement) from KK gauge bosons to KK gravitons,  $M_n \rightarrow \mathbb{M}_n$ .

Next, we make double-copy construction of the 4-point scattering amplitudes of (helicity-zero) longitudinal KK gravitons and of the KK gravitational Goldstone bosons from the corresponding KK gauge/Goldstone boson scattering amplitudes. Applying the CK duality (4.22) to the 4-point scattering amplitudes (4.18a)-(4.18b) of the longitudinal KK gauge bosons and of the corresponding KK Goldstone bosons at the leading order of high energy

expansion, we derive the following 4-point scattering amplitudes of the longitudinal KK gravitons and of the KK gravitational Goldstone bosons:

$$\mathcal{M}_0^{\text{dc}}[4h_L^n] = -\frac{c_4\kappa^2}{16} \left[ \frac{(\mathcal{N}_s^0)^2}{s_0} + \frac{(\mathcal{N}_t^0)^2}{t_0} + \frac{(\mathcal{N}_u^0)^2}{u_0} \right], \quad (4.23a)$$

$$\widetilde{\mathcal{M}}_0^{\text{dc}}[4\phi_n] = -\frac{c_4\kappa^2}{16} \left[ \frac{(\widetilde{\mathcal{N}}_s^0)^2}{s_0} + \frac{(\widetilde{\mathcal{N}}_t^0)^2}{t_0} + \frac{(\widetilde{\mathcal{N}}_u^0)^2}{u_0} \right], \quad (4.23b)$$

where the normalization factor  $c_4$  will be determined by taking the flat 5d limit ( $k \rightarrow 0$ ). We already established [27] the exact double-copy formula for the flat 5d space under orbifold compactification of  $S^1/\mathbb{Z}_2$ , which leads to  $c_4 = 2/3$  [27].

Applying the generalized gauge transformation (4.20) to the numerators of the longitudinal KK graviton scattering amplitude (4.23a) and making use of the Jacobi identity (4.21b), we can directly prove the equivalence between the leading-order scattering amplitudes of the longitudinal KK gravitons and of the gravitational KK Goldstone bosons:

$$\mathcal{M}_0^{\text{dc}}[4h_L^n] = \widetilde{\mathcal{M}}_0^{\text{dc}}[4\phi_n]. \quad (4.24)$$

This is a proof of the KK gravitational equivalence theorem (GRET) from the KK gauge-theory equivalence theorem (GAET) by using the double-copy approach at the level of 4-point scattering amplitudes.

Based on the GRET relation (4.24), we can compute its either side to obtain the same 4-point leading-order gravitational KK scattering amplitude. Since the KK Goldstone numerators  $\{\widetilde{\mathcal{N}}_j^0\}$  [as in Eqs.(4.19b)(3.36)] are much simpler than the KK graviton numerators  $\{\mathcal{N}_j^0\}$  [as in Eqs.(4.19a)(3.35)], we can use the KK Goldstone numerators  $\{\widetilde{\mathcal{N}}_j^0\}$  to explicitly compute the leading-order amplitudes (4.23) as follows:

$$\begin{aligned} \mathcal{M}_0^{\text{dc}}[4h_L^n] &= \widetilde{\mathcal{M}}_0^{\text{dc}}[4\phi_n] = -\frac{c_4\kappa^2}{16} \left[ \frac{(\widetilde{\mathcal{N}}_s^0)^2}{s_0} + \frac{(\widetilde{\mathcal{N}}_t^0)^2}{t_0} + \frac{(\widetilde{\mathcal{N}}_u^0)^2}{u_0} \right] \\ &= \frac{c_4\kappa^2 s_0 (7+c_{2\theta})^2 \csc^2\theta}{64} \left( \sum_{j=0}^{\infty} \tilde{a}_{nnj}^2 \right)^2 \\ &= \frac{c_4\kappa^2 s_0 (7+c_{2\theta})^2 \csc^2\theta}{64} \tilde{a}_{nnnn}^2, \end{aligned} \quad (4.25)$$

where in the last step we have made use of the sum rule (D.3c) for converting the trilinear Goldstone coupling ( $\tilde{a}_{nnj}^2$ ) to the quartic Goldstone coupling ( $\tilde{a}_{nnnn}$ ). Then, using the explicit Feynman diagram calculations as shown in Eqs.(3.65)-(3.66), we have the following 4-point elastic scattering amplitudes of longitudinal KK gravitons and of KK gravitational Goldstone bosons at the leading order of high energy expansion, which are equal to each other due to the GRET (3.67):

$$\mathcal{M}_0[4h_L^n] = \widetilde{\mathcal{M}}_0[4\phi_n] = \frac{\kappa^2 s_0 (7+c_{2\theta})^2 \csc^2\theta}{2304} \sum_{j=1}^{\infty} r_j^2 (r_j^6 - 12r_j^2 + 11) \alpha_{nnj}^2$$

$$\begin{aligned}
&= \frac{\kappa^2 s_0 (7 + c_{2\theta})^2 \csc^2 \theta}{64} \sum_{j=0}^{\infty} \tilde{\beta}_{nnj}^2 \\
&= \frac{\kappa^2 s_0 (7 + c_{2\theta})^2 \csc^2 \theta}{64} \tilde{\beta}_{nnnn},
\end{aligned} \tag{4.26}$$

where the mass ratio  $r_j = \mathbb{M}_j / \mathbb{M}_n$ , and in the last step above we have used the sum rule (D.3d) with the trilinear and quartic gravitational Goldstone couplings  $\tilde{\beta}_{nnj}$  and  $\tilde{\beta}_{nnnn}$  defined respectively in Eqs.(C.9c) and (C.11b).

With the above, we compare the double-copied scattering amplitudes of KK gravitons (KK Goldstone bosons) in Eq.(4.25) with the corresponding KK graviton (KK Goldstone boson) amplitudes by explicit Feynman diagram calculations in Eq.(4.26). They have exactly the same kinematic structure except the difference between the two types of quartic coupling coefficients:

$$\frac{\mathcal{M}_0^{\text{dc}}[4h_L^n]}{\mathcal{M}_0[4h_L^n]} = \frac{\widetilde{\mathcal{M}}_0^{\text{dc}}[4\phi_n]}{\widetilde{\mathcal{M}}_0[4\phi_n]} = \frac{c_4 \tilde{a}_{nnnn}^2}{\tilde{\beta}_{nnnn}}. \tag{4.27}$$

As a consistency check, we take the flat 5d space limit  $k \rightarrow 0$  and find that the above ratio of KK amplitudes approaches one as expected:

$$\lim_{k \rightarrow 0} \frac{\mathcal{M}_0^{\text{dc}}[4h_L^n]}{\mathcal{M}_0[4h_L^n]} = \lim_{k \rightarrow 0} \frac{\widetilde{\mathcal{M}}_0^{\text{dc}}[4\phi_n]}{\widetilde{\mathcal{M}}_0[4\phi_n]} = \lim_{k \rightarrow 0} \frac{c_4 \tilde{a}_{nnnn}^2}{\tilde{\beta}_{nnnn}} = \frac{2/3 \times (3/2)^2}{3/2} = 1, \tag{4.28}$$

where the flat-space coupling coefficients  $\tilde{a}_{nnnn} = \tilde{\beta}_{nnnn} = 3/2$ , as summarized in Table 1.

Hence, according to Eqs.(4.27)-(4.28), we complete our double-copy construction of the 4-point leading-order KK gauge/gravity scattering amplitudes by further imposing the following replacements,

$$c_4 \tilde{a}_{nnnn}^2 \longrightarrow \tilde{\beta}_{nnnn}, \tag{4.29}$$

in addition to the operations in Eq.(4.22). With these, we can reexpress the leading-order double-copied gravitational KK amplitudes (4.25) as follows:

$$\begin{aligned}
\mathcal{M}_0^{\text{dc}}[4h_L^n] &= \widetilde{\mathcal{M}}_0^{\text{dc}}[4\phi_n] = -\frac{\kappa^2 \tilde{\beta}_{nnnn}}{16 \tilde{\alpha}_{nnnn}^2} \left[ \frac{(\widetilde{\mathcal{N}}_s^0)^2}{s_0} + \frac{(\widetilde{\mathcal{N}}_t^0)^2}{t_0} + \frac{(\widetilde{\mathcal{N}}_u^0)^2}{u_0} \right] \\
&= \frac{\kappa^2 s_0 (7 + c_{2\theta})^2 \csc^2 \theta}{64} \tilde{\beta}_{nnnn},
\end{aligned} \tag{4.30}$$

which agrees with Eq.(4.26).

#### 4.2.2 Construction of 4-Point Inelastic KK Graviton Amplitudes

In this subsection, we study the double-copy construction of the leading-order gravitational KK scattering amplitudes for inelastic processes. This includes the inelastic channels  $(n, n) \rightarrow (m, m)$  (with  $n \neq m$ ),  $(0, 0) \rightarrow (n, n)$ ,  $(n, m) \rightarrow (\ell, q)$  (with unequal KK indices of  $n, m, \ell, q$ ).

◆ **Inelastic Scattering Process  $(n, n) \rightarrow (m, m)$ :**

We first consider the inelastic scattering process  $(n, n) \rightarrow (m, m)$  with  $n \neq m$  and  $n, m \in \mathbb{Z}^+$ . Making high energy expansion, we reexpress the leading-order inelastic KK scattering amplitudes (3.44a)-(3.44b) as follows:

$$\mathcal{T}_0[2A_L^n 2A_L^m] = g^2 \left( \frac{\mathcal{C}_s \mathcal{N}_s^{\text{in}0}}{s_0} + \frac{\mathcal{C}_t \mathcal{N}_t^{\text{in}0}}{t_0} + \frac{\mathcal{C}_u \mathcal{N}_u^{\text{in}0}}{u_0} \right), \quad (4.31a)$$

$$\tilde{\mathcal{T}}_0[2A_5^n 2A_5^m] = g^2 \left( \frac{\mathcal{C}_s \tilde{\mathcal{N}}_s^{\text{in}0}}{s_0} + \frac{\mathcal{C}_t \tilde{\mathcal{N}}_t^{\text{in}0}}{t_0} + \frac{\mathcal{C}_u \tilde{\mathcal{N}}_u^{\text{in}0}}{u_0} \right), \quad (4.31b)$$

where the leading-order kinematic numerators  $(\mathcal{N}_s^{\text{in}0}, \mathcal{N}_t^{\text{in}0}, \mathcal{N}_u^{\text{in}0})$  and  $(\tilde{\mathcal{N}}_s^{\text{in}0}, \tilde{\mathcal{N}}_t^{\text{in}0}, \tilde{\mathcal{N}}_u^{\text{in}0})$  are given by

$$(\mathcal{N}_s^{\text{in}0}, \mathcal{N}_t^{\text{in}0}, \mathcal{N}_u^{\text{in}0}) = (s_0 \mathcal{K}_s^{\text{in}0}, t_0 \mathcal{K}_t^{\text{in}0}, u_0 \mathcal{K}_u^{\text{in}0}), \quad (4.32a)$$

$$(\tilde{\mathcal{N}}_s^{\text{in}0}, \tilde{\mathcal{N}}_t^{\text{in}0}, \tilde{\mathcal{N}}_u^{\text{in}0}) = (s_0 \tilde{\mathcal{K}}_s^{\text{in}0}, t_0 \tilde{\mathcal{K}}_t^{\text{in}0}, u_0 \tilde{\mathcal{K}}_u^{\text{in}0}). \quad (4.32b)$$

Using Eq.(4.32) and the difference relation (3.48), we derive the generalized gauge transformations between the kinematic numerators of inelastic scattering amplitudes, analogous to Eq.(3.39):

$$\mathcal{N}_j^{\text{in}0} = \tilde{\mathcal{N}}_j^{\text{in}0} + s_{0j} \Theta_{\text{in}}, \quad (4.33a)$$

$$\Theta_{\text{in}} = -\frac{c_\theta}{2r^2} \sum_{i=0}^{\infty} \left[ (3+r^2)r_i^2 + 2(1-r^4) \right] a_{nni} a_{mmi}, \quad (4.33b)$$

where the mass ratios  $r = M_m/M_n$  and  $r_i = M_i/M_n$ . Then, we sum up the leading-order kinematic numerators (4.32a) and (4.32b) respectively and deduce the following:

$$\sum_j \mathcal{N}_j^{\text{in}0} = \sum_j \tilde{\mathcal{N}}_j^{\text{in}0} + (s_0 + t_0 + u_0) \Theta_{\text{in}} = 0, \quad (4.34a)$$

$$\sum_j \tilde{\mathcal{N}}_j^{\text{in}0} = s_0 c_\theta \sum_{i=0}^{\infty} (\tilde{a}_{nmi}^2 - \tilde{a}_{nni} \tilde{a}_{mmi}) = 0, \quad (4.34b)$$

where the index  $j \in \{s, t, u\}$ , and we have used the kinematic identity  $s_0 + t_0 + u_0 = 0$  and the sum rule (D.3c). Eqs.(4.34a) and (4.34b) demonstrates that the two sets of numerators  $\{\mathcal{N}_j\}$  and  $\{\tilde{\mathcal{N}}_j\}$  satisfy the kinematic Jacobi identities respectively.

Thus, we can formulate the massive color-kinematics (CK) duality for the inelastic scattering amplitudes of the longitudinal KK gauge bosons and of the KK Goldstone bosons in Eq.(4.31) as follows:

$$\mathcal{C}_j \longrightarrow \mathcal{N}_j^{\text{in}0}, \quad \mathcal{C}_j \longrightarrow \tilde{\mathcal{N}}_j^{\text{in}0}, \quad g^2 \longrightarrow -(\kappa/4)^2. \quad (4.35)$$

Applying the color-kinematics duality (4.35) to the inelastic scattering amplitudes (4.31a)-(4.31b) of the longitudinal KK gauge bosons and of the corresponding KK Goldstone bosons at the leading order of high energy expansion, we derive the following:

$$\mathcal{M}_0^{\text{dc}}[2h_L^n 2h_L^m] = -\frac{c_4^{\text{in}} \kappa^2}{16} \left[ \frac{(\mathcal{N}_s^{\text{in}0})^2}{s_0} + \frac{(\mathcal{N}_t^{\text{in}0})^2}{t_0} + \frac{(\mathcal{N}_u^{\text{in}0})^2}{u_0} \right], \quad (4.36a)$$

$$\widetilde{\mathcal{M}}_0^{\text{dc}}[2\phi_n 2\phi_m] = -\frac{c_4^{\text{in}} \kappa^2}{16} \left[ \frac{(\widetilde{\mathcal{N}}_s^{\text{in}0})^2}{s_0} + \frac{(\widetilde{\mathcal{N}}_t^{\text{in}0})^2}{t_0} + \frac{(\widetilde{\mathcal{N}}_u^{\text{in}0})^2}{u_0} \right], \quad (4.36b)$$

where the normalization factor  $c_4^{\text{in}}$  can be determined by taking the flat 5d limit  $k \rightarrow 0$ . For the flat 5d compactification with orbifold  $S^1/\mathbb{Z}_2$ , this factor is given by  $c_4^{\text{in}} = 1$  [26][27].

Since the kinematic numerators of KK Goldstone bosons  $\{\widetilde{\mathcal{N}}_j^{\text{in}0}\}$  are much simpler than that of the longitudinal KK gauge bosons  $\{\mathcal{N}_j^{\text{in}0}\}$ , with the GRET relation (4.24) we can use the kinematic numerators of KK Goldstone bosons to explicitly compute the leading-order KK scattering amplitudes:

$$\begin{aligned} \mathcal{M}_0^{\text{dc}}[2h_L^n 2h_L^m] &= \widetilde{\mathcal{M}}_0^{\text{dc}}[2\phi_n 2\phi_m] = -\frac{c_4^{\text{in}} \kappa^2}{16} \left[ \frac{(\widetilde{\mathcal{N}}_s^{\text{in}0})^2}{s_0} + \frac{(\widetilde{\mathcal{N}}_t^{\text{in}0})^2}{t_0} + \frac{(\widetilde{\mathcal{N}}_u^{\text{in}0})^2}{u_0} \right] \\ &= -\frac{c_4^{\text{in}} \kappa^2 s_0}{32} \left[ 2c_\theta^2 - \frac{(3-c_\theta)^2}{1+c_\theta} - \frac{(3+c_\theta)^2}{1-c_\theta} \right] \tilde{a}_{nnmm}^2 \\ &= \frac{c_4^{\text{in}} \kappa^2 (7+c_{2\theta})^2 s_0}{64 \sin^2 \theta} \tilde{a}_{nnmm}^2, \end{aligned} \quad (4.37)$$

where in the second line we have utilized the sum rule (D.3c) to convert the trilinear coupling into the quartic coupling. In the above, the normalization factor  $c_4^{\text{in}}$  may be determined by taking the flat 5d space limit ( $k \rightarrow 0$ ) which gives  $c_4^{\text{in}} = 1$  [26][27]. Moreover, using the explicit Feynman diagram approach, we derive the 4-point inelastic scattering amplitudes of longitudinal KK gravitons and of KK gravitational Goldstone bosons at the leading order of high energy expansion. They are equal to each other due to the GRET (3.67) and we present them as follows:

$$\begin{aligned} \mathcal{M}_0[2h_L^n 2h_L^m] &= \widetilde{\mathcal{M}}_0[2\phi_n 2\phi_m] \\ &= \frac{\kappa^2 s_0}{128 \sin^2 \theta} \sum_{j=0}^{\infty} \left[ 128 \tilde{\beta}_{nmj}^2 - (29 - 28c_{2\theta} - c_{4\theta}) \tilde{\beta}_{nnj} \tilde{\beta}_{mmj} \right] \\ &= \frac{\kappa^2 s_0 (7+c_{2\theta})^2}{64 \sin^2 \theta} \tilde{\beta}_{nnmm}, \end{aligned} \quad (4.38)$$

where the sum rule (D.3d) is used in the last step.

With these, we compare the double-copied amplitudes of KK gravitons (KK Goldstone bosons) in Eq.(4.37) with the corresponding KK graviton (KK Goldstone bosons) amplitudes by explicit Feynman diagram calculations in Eq.(4.38). We find that they have exactly the same kinematic structure except the difference between the two types of quartic coupling coefficients:

$$\frac{\mathcal{M}_0^{\text{dc}}[2h_L^n 2h_L^m]}{\mathcal{M}_0[2h_L^n 2h_L^m]} = \frac{\widetilde{\mathcal{M}}_0^{\text{dc}}[2\phi_n 2\phi_m]}{\widetilde{\mathcal{M}}_0[2\phi_n 2\phi_m]} = \frac{c_4^{\text{in}} \tilde{a}_{nnmm}^2}{\tilde{\beta}_{nnmm}}. \quad (4.39)$$

Taking the flat 5d space limit ( $k \rightarrow 0$ ), the above ratio of KK amplitudes approaches one as expected:

$$\lim_{k \rightarrow 0} \frac{\mathcal{M}_0^{\text{dc}}[2h_L^n 2h_L^m]}{\mathcal{M}_0[2h_L^n 2h_L^m]} = \lim_{k \rightarrow 0} \frac{\widetilde{\mathcal{M}}_0^{\text{dc}}[2\phi_n 2\phi_m]}{\widetilde{\mathcal{M}}_0[2\phi_n 2\phi_m]} = \lim_{k \rightarrow 0} \frac{c_4^{\text{in}} \tilde{a}_{nnmm}^2}{\tilde{\beta}_{nnmm}} = \frac{1 \times 1^2}{1} = 1, \quad (4.40)$$

where values of the flat 5d coupling coefficients  $\tilde{a}_{nnmm}$  and  $\tilde{\beta}_{nnmm}$  are given in Table 1 of Appendix C.

Accordingly, based on Eqs.(4.39)-(4.40), we impose the following correspondence for the double-copy construction of leading-order inelastic KK gauge/gravity amplitudes,

$$c_4^{\text{in}} \tilde{a}_{nnmm}^2 \longrightarrow \tilde{\beta}_{nnmm}, \quad (4.41)$$

alongside the operations in Eq.(4.35). With these, we can reexpress the leading-order double-copy gravitational KK amplitudes (4.37) as follows:

$$\begin{aligned} \mathcal{M}_0^{\text{dc}}[2h_L^n 2h_L^m] &= \tilde{\mathcal{M}}_0^{\text{dc}}[2\phi_n 2\phi_m] = -\frac{\kappa^2 \tilde{\beta}_{nnmm}}{16 \tilde{a}_{nnmm}^2} \left[ \frac{(\tilde{\mathcal{N}}_s^{\text{in}0})^2}{s_0} + \frac{(\tilde{\mathcal{N}}_t^{\text{in}0})^2}{t_0} + \frac{(\tilde{\mathcal{N}}_u^{\text{in}0})^2}{u_0} \right] \\ &= \frac{\kappa^2 s_0 (7 + c_{2\theta})^2}{64 \sin^2 \theta} \tilde{\beta}_{nnmm}, \end{aligned} \quad (4.42)$$

which agrees with Eq.(4.38).

#### ◆ Inelastic Scattering Process $(n, m) \rightarrow (\ell, q)$ :

Consider the general inelastic scattering of the longitudinal KK gauge bosons  $A_L^n A_L^m \rightarrow A_L^\ell A_L^q$  and of the KK Goldstone bosons  $A_5^n A_5^m \rightarrow A_5^\ell A_5^q$ . Making high energy expansion, we reexpress the leading-order inelastic KK scattering amplitudes (3.53a)-(3.53b) as follows:

$$\mathcal{T}_0[A_L^n A_L^m A_L^\ell A_L^q] = g^2 \left( \frac{\mathcal{C}_s \mathcal{N}_s^{\text{in}0}}{s_0} + \frac{\mathcal{C}_t \mathcal{N}_t^{\text{in}0}}{t_0} + \frac{\mathcal{C}_u \mathcal{N}_u^{\text{in}0}}{u_0} \right), \quad (4.43a)$$

$$\tilde{\mathcal{T}}_0[A_L^n A_L^m A_L^\ell A_L^q] = g^2 \left( \frac{\mathcal{C}_s \tilde{\mathcal{N}}_s^{\text{in}0}}{s_0} + \frac{\mathcal{C}_t \tilde{\mathcal{N}}_t^{\text{in}0}}{t_0} + \frac{\mathcal{C}_u \tilde{\mathcal{N}}_u^{\text{in}0}}{u_0} \right), \quad (4.43b)$$

where the kinematic numerators  $(\mathcal{N}_s^{\text{in}0}, \mathcal{N}_t^{\text{in}0}, \mathcal{N}_u^{\text{in}0})$  and  $(\tilde{\mathcal{N}}_s^{\text{in}0}, \tilde{\mathcal{N}}_t^{\text{in}0}, \tilde{\mathcal{N}}_u^{\text{in}0})$  are given by

$$(\mathcal{N}_s^{\text{in}0}, \mathcal{N}_t^{\text{in}0}, \mathcal{N}_u^{\text{in}0}) = (s_0 \mathcal{K}_s^{\text{in}0}, t_0 \mathcal{K}_t^{\text{in}0}, u_0 \mathcal{K}_u^{\text{in}0}), \quad (4.44a)$$

$$(\tilde{\mathcal{N}}_s^{\text{in}0}, \tilde{\mathcal{N}}_t^{\text{in}0}, \tilde{\mathcal{N}}_u^{\text{in}0}) = (s_0 \tilde{\mathcal{K}}_s^{\text{in}0}, t_0 \tilde{\mathcal{K}}_t^{\text{in}0}, u_0 \tilde{\mathcal{K}}_u^{\text{in}0}). \quad (4.44b)$$

Then, we derive the generalized gauge transformations which connect the numerators of KK gauge bosons to that of the KK Goldstone bosons:

$$\mathcal{N}_j^{\text{in}0} = \tilde{\mathcal{N}}_j^{\text{in}0} + s_{0j} \Theta_{\text{in}}, \quad (4.45a)$$

$$\Theta_{\text{in}} = \frac{M_n^2 + M_m^2 + M_\ell^2 + M_q^2}{4M_n M_m M_\ell M_q} \sum_{i=1}^{\infty} M_i^2 \left[ (1 + c_\theta) a_{nqi} a_{mli} - (1 - c_\theta) a_{nli} a_{mqi} \right]. \quad (4.45b)$$

With these, we compute the sum of each set of the leading-order kinematic numerators  $\{\mathcal{N}_j^{\text{in}0}\}$  and  $\{\tilde{\mathcal{N}}_j^{\text{in}0}\}$  as follows:

$$\sum_j \tilde{\mathcal{N}}_j^{\text{in}0} = -\frac{s_0}{2} \sum_{j=0}^{\infty} \left[ 2c_\theta \tilde{a}_{nmj} \tilde{a}_{\ell qj} + (3 - c_\theta) \tilde{a}_{nqj} \tilde{a}_{m\ell j} - (3 + c_\theta) \tilde{a}_{n\ell j} \tilde{a}_{mqj} \right]$$

$$= -\frac{s_0}{2} [2c_\theta + (3 - c_\theta) - (3 + c_\theta)] \tilde{a}_{nmlq} = 0, \quad (4.46a)$$

$$\sum_j \mathcal{N}_j^{\text{in}0} = \sum_j \tilde{\mathcal{N}}_j^{\text{in}0} + (s_0 + t_0 + u_0) \Theta_{\text{in}} = 0. \quad (4.46b)$$

This proves the kinematic Jacobi identities for the leading-order inelastic KK scattering amplitudes:

$$\mathcal{N}_s^{\text{in}0} + \mathcal{N}_t^{\text{in}0} + \mathcal{N}_u^{\text{in}0} = 0, \quad \tilde{\mathcal{N}}_s^{\text{in}0} + \tilde{\mathcal{N}}_t^{\text{in}0} + \tilde{\mathcal{N}}_u^{\text{in}0} = 0. \quad (4.47)$$

Next, using the extended massive color-kinematics duality we construct the inelastic scattering amplitudes of KK gravitons and of gravitational KK Goldstone bosons at the leading order of high energy expansion, which also obey the GRET:

$$\begin{aligned} \mathcal{M}_0^{\text{dc}}[h_L^n h_L^m h_L^\ell h_L^q] &= \tilde{\mathcal{M}}_0^{\text{dc}}[\phi_n \phi_m \phi_\ell \phi_q] = -\frac{c_4^{\text{in}} \kappa^2}{16} \left[ \frac{(\tilde{\mathcal{N}}_s^{\text{in}0})^2}{s_0} + \frac{(\tilde{\mathcal{N}}_t^{\text{in}0})^2}{t_0} + \frac{(\tilde{\mathcal{N}}_u^{\text{in}0})^2}{u_0} \right] \\ &= -\frac{c_4^{\text{in}} \kappa^2 s_0}{32} \left[ 2c_\theta^2 - \frac{(3-c_\theta)^2}{1+c_\theta} - \frac{(3+c_\theta)^2}{1-c_\theta} \right] \tilde{a}_{nmlq}^2 \\ &= \frac{c_4^{\text{in}} \kappa^2 (7+c_{2\theta})^2 s_0}{64 \sin^2 \theta} \tilde{a}_{nmlq}^2 = \frac{\kappa^2 (7+c_{2\theta})^2 s_0}{64 \sin^2 \theta} \tilde{\beta}_{nmlq}, \end{aligned} \quad (4.48)$$

where in the second equality the completeness relation (D.2b) is applied and in the last equality we have used the conversion relation (4.51). In the above, the normalization factor  $c_4^{\text{in}}$  may be determined by taking the flat 5d limit ( $k \rightarrow 0$ ) which gives  $c_4^{\text{in}} = 2$  [26][27]. On the other hand, using the explicit Feynman diagram approach, we compute the inelastic scattering amplitudes of gravitational KK Goldstone bosons (KK gravitons) at the leading order of high energy expansion:

$$\begin{aligned} \mathcal{M}_0[h_L^n h_L^m h_L^\ell h_L^q] &= \tilde{\mathcal{M}}_0[\phi_n \phi_m \phi_\ell \phi_q] \\ &= -\frac{\kappa^2 s_\theta}{32} \sum_{j=0}^{\infty} \left[ (15 + c_{2\theta}) \tilde{\beta}_{nmj} \tilde{\beta}_{\ell qj} - \frac{4(5 + 2c_\theta + c_{2\theta})}{1 + c_\theta} \tilde{\beta}_{nqj} \tilde{\beta}_{m\ell j} - \frac{4(8 + 7c_\theta + c_{3\theta})}{1 - c_{2\theta}} \tilde{\beta}_{n\ell j} \tilde{\beta}_{mqj} \right] \\ &= \frac{\kappa^2 (7 + c_{2\theta})^2 s_0}{64 \sin^2 \theta} \tilde{\beta}_{nmlq}. \end{aligned} \quad (4.49)$$

As a consistency check, we take the limit of flat 5d space ( $k \rightarrow 0$ ) and find that the ratio of KK scattering amplitudes approaches one:

$$\lim_{k \rightarrow 0} \frac{c_4^{\text{in}} \tilde{a}_{nmlq}^2}{\tilde{\beta}_{nmlq}} = \frac{2 \times (1/2)^2}{1/2} = 1, \quad (4.50)$$

where the values of couplings  $\tilde{a}_{nmlq}$  and  $\tilde{\beta}_{nmlq}$  in the flat space limit are summarized in Table 1. Hence, we set up the following double-copy correspondence for the above leading-order inelastic KK gauge/gravity scattering amplitudes,

$$c_4^{\text{in}} \tilde{a}_{nmlq}^2 \longrightarrow \tilde{\beta}_{nmlq}, \quad (4.51)$$

together with Eq.(4.35).



◆ **Inelastic Scattering Process  $(0, 0) \rightarrow (n, n)$ :**

Finally, we analyze the mixed 4-point inelastic scattering amplitudes of  $(0, 0) \rightarrow (n, n)$ . According to Eq.(3.62), we verify that the kinematic Jacobi identities for the numerators  $\{\mathcal{N}_i^{\text{in}0}\}$  and  $\{\widetilde{\mathcal{N}}_i^{\text{in}0}\}$  hold. Applying the massive color-kinematics duality (4.35), we construct the following double-copied gravitational KK scattering amplitudes:

$$\mathcal{M}_0^{\text{dc}}[2h_{\pm 2}^0 2h_L^n] = \widetilde{\mathcal{M}}_0^{\text{dc}}[2h_{\pm 2}^0 2\phi_n] = \frac{c_4^{\text{in}} \kappa^2}{32} s_0 (1 - c_{2\theta}) a_{0nn}^4, \quad (4.52)$$

where  $a_{0nn} = f_0$  is given by Eq.(2.17). On the other hand, using the explicit Feynman diagram approach and the GRET identity (3.67), we compute the 4-point inelastic gravitational KK scattering amplitudes at the leading order of high energy expansion:

$$\mathcal{M}_0[2h_{\pm 2}^0 2h_L^n] = \widetilde{\mathcal{M}}_0[2h_{\pm 2}^0 2\phi_n] = \frac{\kappa^2}{32} s_0 (1 - c_{2\theta}) \alpha_{0nn}^2, \quad (4.53)$$

where  $\alpha_{0nn} = u_0$  is given by Eq.(2.51). We see that the two types of KK scattering amplitudes (4.52) and (4.53) exhibit the same kinematic structure, and thus the mixed double-copy construction (4.52) is successful. As a consistency check, we take the flat 5d limit ( $k \rightarrow 0$ ) and find that the ratio of the two KK amplitudes approaches one as expected:

$$\lim_{k \rightarrow 0} \frac{\mathcal{M}_0^{\text{dc}}[2h_{\pm 2}^0 2h_L^n]}{\mathcal{M}_0[2h_{\pm 2}^0 2h_L^n]} = \lim_{k \rightarrow 0} \frac{\widetilde{\mathcal{M}}_0^{\text{dc}}[2h_{\pm 2}^0 2\phi_n]}{\widetilde{\mathcal{M}}_0[2h_{\pm 2}^0 2\phi_n]} = \lim_{k \rightarrow 0} \frac{c_4^{\text{in}} a_{0nn}^4}{\alpha_{0nn}^2} = \frac{1 \times 1^4}{1^2} = 1, \quad (4.54)$$

where the values of the coupling coefficients  $a_{0nn}$  and  $\alpha_{0nn}$  in the flat 5d limit are given by Eq.(C.12). Hence, for the mixed inelastic KK scattering process of  $(0, 0) \rightarrow (n, n)$ , we set up the following double-copy correspondence for the leading-order inelastic KK gauge/gravity scattering amplitudes,

$$c_4^{\text{in}} a_{0nn}^4 \longrightarrow \alpha_{0nn}^2, \quad (4.55)$$

together with Eq.(4.35).

For a further demonstration, we study the double-copy constructions for the gravitational KK scattering amplitudes of  $h_0^{+2} h_0^{-2} \rightarrow h_n^{+1} h_n^{-1}$  and  $h_0^{+2} h_0^{-2} \rightarrow \mathcal{V}_n^{+1} \mathcal{V}_n^{-1}$ . For this purpose, we compute the following 4-point mixed scattering amplitudes involving two zero-mode gauge bosons and two longitudinal KK gauge bosons (two KK Goldstone bosons) at the leading order of high energy expansion:

$$\begin{aligned} \mathcal{T}_0[A_+^{a0} A_-^{b0} \rightarrow A_+^{cn} A_-^{dn}] &= g^2 \left( \frac{\mathcal{C}_s \mathcal{N}_s^{\text{in}0}}{s_0} + \frac{\mathcal{C}_t \mathcal{N}_t^{\text{in}0}}{t_0} + \frac{\mathcal{C}_u \mathcal{N}_u^{\text{in}0}}{u_0} \right) \\ &= -g^2 s_0 \left[ \frac{(1 - c_\theta)^2 \mathcal{C}_t}{2t_0} - \frac{(1 - c_\theta)^2 \mathcal{C}_u}{2u_0} \right] a_{000} a_{nn0}, \end{aligned} \quad (4.56a)$$

$$\begin{aligned} \mathcal{T}_0[A_+^{a0} A_-^{b0} \rightarrow A_L^{cn} A_L^{dn}] &= g^2 \left( \frac{\mathcal{C}_s \mathcal{N}_s^{\text{in}0}}{s_0} + \frac{\mathcal{C}_t \mathcal{N}_t^{\text{in}0}}{t_0} + \frac{\mathcal{C}_u \mathcal{N}_u^{\text{in}0}}{u_0} \right) \\ &= g^2 s_0 \left[ -\frac{(1 - c_\theta^2) \mathcal{C}_t}{2t_0} + \frac{(1 - c_\theta^2) \mathcal{C}_u}{2u_0} \right] a_{000} a_{nn0}, \end{aligned} \quad (4.56b)$$

$$\begin{aligned}
\widetilde{\mathcal{T}}_0[A_+^{a0} A_-^{b0} \rightarrow A_5^{cn} A_5^{dn}] &= g^2 \left( \frac{\mathcal{C}_s \widetilde{\mathcal{N}}_s^{\text{in}0}}{s_0} + \frac{\mathcal{C}_t \widetilde{\mathcal{N}}_t^{\text{in}0}}{t_0} + \frac{\mathcal{C}_u \widetilde{\mathcal{N}}_u^{\text{in}0}}{u_0} \right) \\
&= g^2 s_0 \left[ \frac{(1-c_\theta^2) \mathcal{C}_t}{2t_0} - \frac{(1-c_\theta^2) \mathcal{C}_u}{2u_0} \right] a_{000} \widetilde{a}_{nn0}, \tag{4.56c}
\end{aligned}$$

where the coupling coefficients  $a_{000} = a_{nn0} = \widetilde{a}_{nn0} = f_0$  according to Eq.(C.10). We see that the above leading-order amplitudes (4.56b) and (4.56c) obey the GAET (2.38) as expected.

With the leading-order (KK) gauge boson amplitudes (4.56), we construct the double-copied (KK) graviton scattering amplitude,

$$\begin{aligned}
\mathcal{M}_0^{\text{dc}}[h_0^{+2} h_0^{-2} \rightarrow h_n^{+1} h_n^{-1}] &= -\frac{c_4^{\text{in}} \kappa^2}{16} \left( \frac{\mathcal{N}_s^{\text{in}0} \mathcal{N}'_s{}^{\text{in}0}}{s_0} + \frac{\mathcal{N}_t^{\text{in}0} \mathcal{N}'_t{}^{\text{in}0}}{t_0} + \frac{\mathcal{N}_u^{\text{in}0} \mathcal{N}'_u{}^{\text{in}0}}{u_0} \right) \\
&= -\frac{c_4^{\text{in}} \kappa^2 s_0}{16} (1-c_\theta)^2 a_{000}^2 a_{nn0}^2, \tag{4.57}
\end{aligned}$$

and the corresponding scattering amplitude with gravitational KK Goldstone bosons,

$$\begin{aligned}
\widetilde{\mathcal{M}}_0^{\text{dc}}[h_0^{+2} h_0^{-2} \rightarrow \mathcal{V}_n^{+1} \mathcal{V}_n^{-1}] &= -\frac{c_4^{\text{in}} \kappa^2}{16} \left( \frac{\widetilde{\mathcal{N}}_s^{\text{in}0} \mathcal{N}'_s{}^{\text{in}0}}{s_0} + \frac{\widetilde{\mathcal{N}}_t^{\text{in}0} \mathcal{N}'_t{}^{\text{in}0}}{t_0} + \frac{\widetilde{\mathcal{N}}_u^{\text{in}0} \mathcal{N}'_u{}^{\text{in}0}}{u_0} \right) \\
&= \frac{c_4^{\text{in}} \kappa^2 s_0}{16} (1-c_\theta)^2 a_{000}^2 a_{nn0} \widetilde{a}_{nn0}, \tag{4.58}
\end{aligned}$$

where the coupling coefficients  $a_{000} = a_{nn0} = \widetilde{a}_{nn0} = f_0$  according to Eq.(C.10). Then, we explicitly compute the above scattering amplitudes by Feynman diagram approach:

$$\mathcal{M}_0[h_0^{+2} h_0^{-2} \rightarrow h_n^{+1} h_n^{-1}] = -\frac{\kappa^2 s_0}{16} (1-c_\theta)^2 \alpha_{000} \alpha_{nn0}, \tag{4.59}$$

$$\widetilde{\mathcal{M}}_0[h_0^{+2} h_0^{-2} \rightarrow \mathcal{V}_n^{+1} \mathcal{V}_n^{-1}] = \frac{\kappa^2 s_0}{16} (1-c_\theta)^2 \alpha_{000} \widetilde{\alpha}_{nn0}, \tag{4.60}$$

where the coupling coefficients  $\alpha_{000} = \alpha_{nn0} = \widetilde{\alpha}_{nn0} = u_0$  according to Eq.(C.10). Comparing the leading-order scattering amplitudes (4.58) and (4.60), we see that they share exactly the same kinematic structure and can be equal under the following correspondence:

$$c_4^{\text{in}} a_{000}^2 a_{0nn} \widetilde{a}_{nn0} \longrightarrow \alpha_{000} \widetilde{\alpha}_{nn0}. \tag{4.61}$$

As a consistency check, we take the flat 5d limit ( $k \rightarrow 0$ ) and find that the ratio of the two gravitational amplitudes (4.58) and (4.60) approaches one as expected:

$$\lim_{k \rightarrow 0} \frac{\mathcal{M}_0^{\text{dc}}[2h_{\pm 2}^0 2h_L^n]}{\mathcal{M}_0[2h_{\pm 2}^0 2h_L^n]} = \lim_{k \rightarrow 0} \frac{\widetilde{\mathcal{M}}_0^{\text{dc}}[2h_{\pm 2}^0 2\phi_n]}{\widetilde{\mathcal{M}}_0[2h_{\pm 2}^0 2\phi_n]} = \lim_{k \rightarrow 0} \frac{c_4^{\text{in}} a_{0nn}^4}{\alpha_{0nn}^2} = \frac{1 \times 1^4}{1^2} = 1, \tag{4.62}$$

where the values of the coupling coefficients  $a_{0nn}$  and  $\alpha_{0nn}$  in the flat 5d limit are given by Eq.(C.12).

In summary, we have established the double-copy construction from the GAET to the GRET for the 4-point leading-order (LO) KK scattering amplitudes:

$$\begin{aligned}
\mathcal{M}_0^{(4)}[h_L^n] &\stackrel{\text{GRET}}{=} \widetilde{\mathcal{M}}_0^{(4)}[\phi_n] \\
\uparrow \text{DC}_4 &\qquad \qquad \uparrow \text{DC}_4 \\
\mathcal{T}_0^{(4)}[A_L^n] &\stackrel{\text{GAET}}{=} \widetilde{\mathcal{T}}_0^{(4)}[A_5^n]
\end{aligned} \tag{4.63}$$

In parallel to Eq.(4.17), we discuss the meaning of the above Eq.(4.63). Along the horizontal directions, the lower (upper) equality presents the GAET (GRET) at the level of 4-point KK scattering amplitudes. On the other hand, along the direction of each vertical arrow, the correspondence is established by the double-copy (DC) under the extended massive color-kinematics dualities and the KK gauge/gravity coupling correspondences for both the elastic scattering [as shown in Eqs.(4.22)(4.29)] and the inelastic scattering [as shown in Eqs.(4.35)(4.41)(4.51)(4.55)(4.61)]. The left vertical arrow represents the double-copy (DC<sub>4</sub>) construction of the 4-point LO KK graviton amplitude  $\mathcal{M}_0^{(4)}[h_L^n]$  from the corresponding LO KK gauge boson amplitude  $\mathcal{T}_0^{(4)}[A_L^n]$ . The right vertical arrow represents the double-copy (DC<sub>4</sub>) construction of the 4-point LO gravitational KK Goldstone boson amplitude  $\widetilde{\mathcal{M}}_0^{(4)}[\phi_n]$  from the corresponding (gauge theory) LO KK Goldstone boson amplitude  $\widetilde{\mathcal{T}}_0^{(4)}[A_5^n]$ . Finally, this picture also holds if we replace the two amplitudes  $\mathcal{M}_0^{(4)}[h_L^n]$  and  $\widetilde{\mathcal{M}}_0^{(4)}[\phi_n]$  in the top row by  $\mathcal{M}_0^{(4)}[h_{\pm 1}^n]$  and  $\widetilde{\mathcal{M}}_0^{(4)}[\mathcal{V}_{\pm 1}^n]$  respectively.

### 4.3 Extension of Warped LO Double-Copy with $N \geq 4$

So far, we have explicitly demonstrated that the warped double-copy can be realized for 3-point and 4-point KK scattering amplitudes at the leading order of high energy expansion. In this subsection, we discuss the extension of the warped LO double-copy to the  $N$ -point KK scattering amplitudes with  $N \geq 4$ . Using the GAET [cf. Eq.(2.36)] and GRETs [cf. Eqs.(2.84) and (2.99)], we find that it is more convenient to analyze the LO KK Goldstone boson amplitudes.

For a given  $S$ -matrix element  $\mathbb{S}$  with  $N$  external states in (3+1)d spacetime, its mass-dimension is determined by the spacetime dimension  $d = 4$  and the number of external states  $N$  [52][15]:

$$D_{\mathbb{S}} = 4 - N. \quad (4.64)$$

Then, we first consider the pure KK Goldstone amplitude having even number of external states, with  $N \geq 4$  and  $N \in 2\mathbb{Z}$ . For the KK gauge theory, the leading energy-power dependence of the  $N$ -point KK Goldstone boson ( $A_5^{gn}$ ) amplitude  $\widetilde{\mathcal{T}}$  is given in the second equation of Eq.(2.106a) with  $\bar{V}_3^{\min} = N_v = 0$ . Thus, the mass-dimension of the amplitude  $\widetilde{\mathcal{T}}$  and its leading energy-power dependence are equal:

$$D_{\widetilde{\mathcal{T}}}(NA_5^{gn}) = D_E(NA_5^{gn}) = 4 - N. \quad (4.65)$$

Hence, Eq.(4.65) indicates that we can express the LO scattering amplitudes of KK Goldstone bosons as follows:

$$\widetilde{\mathcal{T}}_0[A_5^{a_1 n_1}, \dots, A_5^{a_N n_N}] = \tilde{a}_{n_1 \dots n_N} g^{N-2} \sum_{i=1}^{(2N-5)!!} \frac{\mathcal{C}_i \widetilde{\mathcal{N}}_i^0}{\mathcal{D}_i}, \quad (4.66)$$

where  $\mathcal{D}_i$  denotes the products of denominators of propagators and the effective  $N$ -point coupling coefficients  $\tilde{a}_{n_1 \dots n_N}$  is dimensionless because the 4d gauge coupling is dimensionless,  $[g] = 0$ . Moreover, by taking the Del Duca-Dixon-Maltoni (DDM) color decomposition [53], we can express Eq.(4.66) in terms of Kleiss-Kuijf basis [54]. Thus, we can readily

write down the  $N$ -point KK coupling coefficient  $\tilde{a}_{n_1 \dots n_N}$  as follows:

$$\begin{aligned} \tilde{a}_{n_1 \dots n_N} &= \sum_{j_1, \dots, j_{N-3}} \tilde{a}_{n_1 n_{\sigma(2)j_1}} \tilde{a}_{j_2 n_{\sigma(3)j_1}} \cdots \tilde{a}_{j_{N-4} n_{\sigma(N-2)j_{N-3}}} \tilde{a}_{n_{\sigma(N-1)n_N} j_{N-3}} \\ &= \llbracket \tilde{f}_{n_1} \cdots \tilde{f}_{n_N} \rrbracket, \end{aligned} \quad (4.67)$$

where the completeness relation (D.1) has been imposed, the external states 1 and  $N$  are fixed and  $\sigma$  represents a permutation of the remaining  $(N-2)$  external legs. Every color-ordered amplitude shares the same KK coupling coefficient, because  $\tilde{a}_{n_1 \dots n_N}$  is just the integration of the 5d wavefunctions  $\{\tilde{f}_{n_i}\}$  associated with each external state  $\{A_5^{a_i n_i}\}$  which means the KK indices of coupling coefficient  $\tilde{a}_{n_1 \dots n_N}$  are symmetric and independent of the color ordering. We also note that the LO KK Goldstone numerators  $\{\tilde{\mathcal{N}}_i^0\}$  in Eq.(4.66) should still obey the kinematic Jacobi identities for  $N$ -point amplitude since they are fully determined by kinematics (same as the flat 5d case) and the effect of the warped 5d compactification only affects the KK coupling coefficients  $\tilde{a}_{n_1 \dots n_N}$ .

As for the KK gravity theory, we first consider the scattering amplitudes of KK scalar Goldstone bosons ( $\phi_n$ ). By using the CK duality and Eq.(4.66), the  $N$ -point LO amplitudes of KK scalar Goldstone bosons can be constructed via double-copy (DC) as follows:

$$\tilde{\mathcal{M}}_0^{\text{dc}}[\phi_{n_1} \cdots \phi_{n_N}] = -c_N \tilde{a}_{n_1 \dots n_N}^2 \left(\frac{\kappa}{4}\right)^{N-2} \sum_{i=1}^{(2N-5)!!} \frac{(\tilde{\mathcal{N}}_i^0)^2}{\mathcal{D}_i}. \quad (4.68)$$

Then, we can derive the leading energy-power dependence of the KK gravitational Goldstone amplitude (4.68):

$$D_E(NA_5^{an}) \xrightarrow{\text{DC}} D_E(N\phi_n) = 2(4-N) + 2(N-3) = 2. \quad (4.69)$$

This confirms our power counting rule Eq.(2.104a) at tree level ( $L=0$ ). Moreover, based on the double-copy correspondence, we can apply the following replacement:

$$c_N \tilde{a}_{n_1 \dots n_N}^2 \longrightarrow \tilde{\beta}_{n_1 \dots n_N}. \quad (4.70)$$

Thus, the  $N$ -point Goldstone amplitude can be expressed as follows:

$$\tilde{\mathcal{M}}_0[\phi_{n_1} \cdots \phi_{n_N}] = -\tilde{\beta}_{n_1 \dots n_N} \left(\frac{\kappa}{4}\right)^{N-2} \sum_{\ell=1}^{(2N-5)!!} \frac{(\tilde{\mathcal{N}}_\ell^0)^2}{\mathcal{D}_\ell}, \quad (4.71)$$

where the KK Goldstone coupling coefficient  $\tilde{\beta}_{n_1 \dots n_N}$  is also dimensionless. To understand the dimensionlessness of  $\tilde{\beta}_{n_1 \dots n_N}$ , we compute the difference between the mass-dimension of the KK Goldstone amplitude  $\tilde{\mathcal{M}}(N\phi_n)$  and its leading energy-dependence via Eqs.(4.64) and (4.69), which yields:

$$\Delta D(\tilde{\mathcal{M}}) = D_{\tilde{\mathcal{M}}}(N\phi_n) - D_E(N\phi_n) = 2 - N = [\kappa^{N-2}]. \quad (4.72)$$

Eq.(4.72) shows that the residual mass-dimension of the LO amplitude  $\tilde{\mathcal{M}}_0$  fully comes from the gravitational coupling constant  $\kappa$  which has mass-dimension  $[\kappa] = -1$ . The LO

KK Goldstone amplitude (4.71) is mass-independent (resembling a “massless” amplitude) and is contributed by the trilinear vertices of  $A_5^{an_1}$ - $A_5^{an_2}$ - $A_\mu^{an_3}$  type (containing one partial derivative). Hence, we deduce that the KK coupling coefficient  $\tilde{\beta}_{n_1 \dots n_N}$  is dimensionless and takes the following form:

$$\tilde{\beta}_{n_1 \dots n_N} = \sum_{j_1, \dots, j_{N-3}=0}^{\infty} \tilde{\beta}_{n_1 n_2 j_1} \cdots \tilde{\beta}_{n_{N-1} n_N j_{N-3}} = \llbracket \mathbf{w}_{n_1} \cdots \mathbf{w}_{n_N} \rrbracket. \quad (4.73)$$

This shows that the coupling coefficient  $\tilde{\beta}_{n_1 \dots n_N}$  only depends on the 5d wavefunction  $\{\mathbf{w}_{n_i}\}$  associated with the external states  $\{\phi_{n_i}\}$ .

Then, we consider the case for odd number of external states with  $N \geq 4$  and  $N \in 2\mathbb{Z}+1$ , the mass-dimension of  $N$ -point KK Goldstone amplitude (in the KK gauge theory) and its leading energy-power dependence are given by

$$D_{\tilde{\mathcal{T}}}(NA_5^n) = 4 - N, \quad D_E(NA_5^n) = 3 - N. \quad (4.74)$$

Thus, the difference  $D_{\tilde{\mathcal{T}}} - D_E = 1$  for the odd number  $N$ . Actually, this difference is due to the fact that the  $N$ -point KK Goldstone amplitude (with odd  $N$ ) must contain a non-derivative cubic vertex (C.2c). Hence, its KK coupling coefficient is  $\tilde{a}_{n_1 \dots n_N} \propto M_j$  and should have mass-dimension one to make up the full mass-dimension of the KK Goldstone amplitude,  $D_{\tilde{\mathcal{T}}} = 4 - N$ . This means that for the  $N$ -point KK scattering amplitude (with odd  $N$ ), the leading KK Goldstone amplitude contains one power of KK mass and has its leading energy dependence  $D_E = 3 - N$ , which is lower than the leading energy dependence of the mass-independent KK Goldstone amplitude with even number  $N$ . For such massive amplitudes with odd number  $N$ , there is no systematic way to realize the BCJ-type double-copy for the warped KK gauge/gravity theories.

Next, we consider the double-copy construction for the  $N$ -point scattering amplitudes of KK vector Goldstone bosons ( $\mathcal{V}_n^\mu$ ). The dimensional analysis including the counting of leading energy dependence is similar to the aforementioned conclusion on the  $N$ -point amplitudes of the KK scalar Goldstone bosons ( $\phi_n$ ). For the  $N$ -point KK scattering amplitudes (with even number  $N$ ), we obtain the following scattering amplitude:

$$\begin{aligned} \tilde{\mathcal{M}}_0[\mathcal{V}_{n_1}^{\pm 1}, \dots, \mathcal{V}_{n_N}^{\pm 1}] &= -\tilde{\alpha}_{n_1 \dots n_N} \left(\frac{\kappa}{4}\right)^{N-2} \frac{(2N-5)!!}{\mathcal{D}_\ell} \frac{1}{\mathcal{D}_\ell} \sum_{\ell=1}^N \sum_{i=0}^N \sum_{j=0}^N \tilde{\mathcal{N}}_{\ell,i}^0[(N-j)A_5^{an}; jA_\pm^{an}] \\ &\quad \times \tilde{\mathcal{N}}_{\ell,i}^0[(N-j)A_\pm^{an}; jA_5^{an}], \end{aligned} \quad (4.75)$$

where for every possible product of two numerators in the sums we have replaced the relevant product of KK gauge coupling coefficients by the corresponding common gravitational KK coupling coefficient:

$$c_N \llbracket \cdots \mathbf{f}_{n_k} \cdots \tilde{\mathbf{f}}_{n_{k'}} \cdots \rrbracket \llbracket \cdots \tilde{\mathbf{f}}_{n_k} \cdots \mathbf{f}_{n_{k'}} \cdots \rrbracket \longrightarrow \tilde{\alpha}_{n_1 \dots n_N}, \quad (4.76)$$

where the  $N$ -point KK gauge coupling coefficient  $\llbracket \cdots \mathbf{f}_{n_k} \cdots \tilde{\mathbf{f}}_{n_{k'}} \cdots \rrbracket$  contains  $(N-j)$  wavefunctions associated with  $A_5^{an}$  and  $j$  wavefunctions associated with  $A_\pm^{an}$ , while the  $N$ -point

KK gauge coupling coefficient  $[[\cdots \tilde{f}_{n_k} \cdots f_{n_{k'}} \cdots]]$  includes  $j$  wavefunctions associated with  $A_5^{gn}$  and  $(N-j)$  wavefunctions associated with  $A_{\pm}^{gn}$ . The bracket notation  $[[\cdots]]$  for defining the KK coupling coefficient as an integral of 5d wavefunctions over the 5th coordinate is given in Eq.(D.10) of Appendix D. In Eq.(4.76),  $c_N$  is a normalization factor and can be determined by taking the flat 5d limit ( $k \rightarrow 0$ ). The effective  $N$ -point coupling coefficient  $\tilde{\alpha}_{n_1 \cdots n_N}$  is dimensionless and can be expressed as follows:

$$\tilde{\alpha}_{n_1 \cdots n_N} = \sum_{j_1, \dots, j_{N-3}=0}^{\infty} \tilde{\rho}_{n_1 n_2 j_1} \tilde{\rho}_{j_2 n_3 j_1} \tilde{\alpha}_{j_2 n_4 j_3} \cdots \tilde{\alpha}_{n_{N-1} n_N j_{N-3}} = [[v_{n_1} \cdots v_{n_N}]]. \quad (4.77)$$

In the above, we have chosen the case with 2  $\tilde{\rho}_{nml}$ 's and  $(N-4)$   $\tilde{\alpha}_{nml}$ 's as an example, the results computed for other combinations of  $\tilde{\rho}_{nml}$  and  $\tilde{\alpha}_{nml}$  will give the same coupling coefficient  $[[v_{n_1} \cdots v_{n_N}]]$  as in Eq.(4.77). This is because, regardless of the numbers and positions of  $\tilde{\rho}_{nml}$  and  $\tilde{\alpha}_{nml}$ , the internal lines (indices) can be summed over by using the completeness relation of Eq.(D.1). The final result only depends on the 5d eigenfunctions  $\{v_{n_i}\}$  associated with the external states  $\{\mathcal{V}_{n_i}^{\mu}\}$ . Similarly, for the scattering amplitudes containing  $N$  external states (with odd number  $N$ ), the KK scattering amplitudes will be proportional to KK masses and thus have lower energy-power dependence; so there is no clear evidence that the double-copy construction would hold for these massive amplitudes.

We can further prove the  $N$ -point LO double-copy relation by utilizing the Cachazo-He-Yuan (CHY) formalism [55]. We start by considering the  $(4+\delta)$ -dimensional spacetime with  $\delta$  extra spatial dimensions under the toroidal compactification. (Since we will study the KK scattering amplitudes at the leading order (LO) only, the current discussion under toroidal compactification will give the same results for the case of orbifold compactification of  $S^1/\mathbb{Z}_2$  at the LO of high energy expansion.) For a momentum  $\hat{p}_a$  living in the 6d spacetime, we can decompose it in the following form:

$$\hat{p}_a^M = (p_a^{\mu}; M_{n_a}, 0), \quad a \in \{1, \dots, N\}, \quad (4.78)$$

where  $p_a^{\mu}$  denotes the momentum in the  $(3+1)$ d spacetime and  $M_{n_a}$  is the momentum-component in the 5th dimension (realized as discretized KK mass-eigenvalues). For each polarization vector, we have two choices, depending on whether it is restricted to the extra dimension or not. Thus, we can define the polarizations for each KK gauge boson and its corresponding KK Goldstone boson as follows:

$$\epsilon_{a,g}^M = (\epsilon_a^{\mu}; 0, 0), \quad \epsilon_{a,s}^M = (\vec{0}; 0, 1), \quad (4.79)$$

where the subscripts ‘‘g’’ and ‘‘s’’ stand for the KK gauge boson and KK Goldstone boson respectively. The LO scattering amplitude  $\mathcal{A}_N^0$  in either gauge theory or gravity theory can be re-written as the expression of a contour integral:

$$\mathcal{A}_N = \oint_{\Gamma} z_{ij}^2 z_{jk}^2 z_{ki}^2 \prod_{\substack{a=1 \\ (a \neq i,j,k)}}^N \frac{dz_a}{\mathbb{E}_a} I_N, \quad \mathbb{E}_a = \sum_{\substack{b=1 \\ (b \neq a)}}^N \frac{p_a \cdot p_b + M_{n_a} M_{n_b}}{z_a - z_b}, \quad (4.80)$$

where the contour  $\Gamma$  encloses all the solutions for the scattering equations of  $\mathbb{E}_a$  and  $I_N$  is the theory-dependent integrand. Under high energy expansion, the above formulations can be further derived at the energy leading-order

$$\mathcal{A}_N^0 = \oint_{\Gamma} z_{ij}^2 z_{jk}^2 z_{ki}^2 \prod_{\substack{a=1 \\ (a \neq i, j, k)}}^N \frac{dz_a}{\mathbb{E}_a^0} I_N^0, \quad \mathbb{E}_a^0 = \sum_{\substack{b=1 \\ (b \neq a)}}^N \frac{p_a \cdot p_b}{z_a - z_b}. \quad (4.81)$$

We note that in high energy limit and for the KK Goldstone boson sector, the 5d KK Yang-Mills gauge theory is reduced to a 4d KK Yang-Mills-Scalar (YMS) theory, whereas the 5d KK Einstein gravity theory is reduced to a 4d KK Einstein-Maxwell-Scalar (EMS) theory. According to the dimensional compactification of the CHY formalism [56], we can express the scalar integrands for the YMS (single flavor) amplitude and the EMS amplitude as follows:

$$I_N^{0, \text{YMS:s}} = \tilde{a}_{n_1 \dots n_N} \mathcal{C}[\alpha] \text{PT}[\alpha] (\text{Pf} \mathbb{X}_N) (\text{Pf}' \mathbb{A}_N), \quad (4.82a)$$

$$I_N^{0, \text{EMS:s}} = c_N \tilde{a}_{n_1 \dots n_N}^2 (\text{Pf} \mathbb{X}_N)^2 (\text{Pf}' \mathbb{A}_N)^2, \quad (4.82b)$$

where the symbols Pf and Pf' denote the Pfaffian and reduced Pfaffian respectively. The matrix quantities  $\mathbb{X}_N$  and  $\mathbb{A}_N$  are given by

$$(\mathbb{X}_N)_{ab} = \begin{cases} \frac{1}{z_a - z_b}, & (\text{for } a \neq b), \\ 0, & (\text{for } a = b), \end{cases} \quad (\mathbb{A}_N)_{ab} = \begin{cases} \frac{p_a \cdot p_b}{z_a - z_b}, & (\text{for } a \neq b), \\ 0, & (\text{for } a = b). \end{cases} \quad (4.83)$$

In Eq.(4.82a), the color factor  $\mathcal{C}[\alpha]$  is decomposed into the DDM basis and contains  $(N-2)!$  elements, as defined below:

$$\mathcal{C}[\alpha] = \sum_{e_1, \dots, e_{N-3}} f^{a_1 a_{\alpha(2)} e_1} f^{e_1 a_{\alpha(3)} e_2} \dots f^{e_{N-3} a_{\alpha(N-1)} a_N}, \quad (4.84)$$

where the 1st and  $N$ -th labels are fixed in the color ordering  $\alpha$ , i.e.,  $\alpha = [1, \alpha(2), \dots, \alpha(N-1), N] \in S_{N-2}$ . The symbol  $\text{PT}[\alpha]$  stands for the Parke-Taylor factor and is given by

$$\text{PT}[\alpha] = \frac{1}{(z_1 - z_{\alpha(2)}) \cdots (z_{\alpha(N-1)} - z_N)(z_N - z_1)}. \quad (4.85)$$

The above CHY formulation works for even number  $N$ , whereas for the odd number  $N$  the Pfaffian is trivially zero. This is consistent with the conclusions we reached earlier in this subsection. Namely, in the case of  $N$ -point scattering (with even number  $N$ ), the leading-order double-copy holds. But, for the  $N$ -point scattering (with odd number  $N$ ), the KK GAET and GRET (which are connected by double-copy) will take the trivial form of  $0 = 0$  at the leading order of high energy expansion. Note that the scattering equations are the same for both the KK YMS theory and the KK EMS theory. So the above LO double-copy relation (4.82b) holds under the replacement  $c_N \tilde{a}_{n_1 \dots n_N}^2 \rightarrow \tilde{\beta}_{n_1 \dots n_N}$ . Using Eqs.(4.81)-(4.82)

and the fundamental BCJ relations, we derive the extended KLT-type double-copy formula for the  $N$ -point LO scattering amplitude of gravitational KK Goldstone bosons:

$$\widetilde{\mathcal{M}}_0^{\text{dc}}[\phi_{n_1}, \dots, \phi_{n_N}] = -c_N \tilde{a}_{n_1 \dots n_N}^2 \left(\frac{\kappa}{4}\right)^{N-2} \sum_{\{\alpha, \beta\} \in S_{N-3}} \tilde{\mathcal{T}}_0[A_5^{a_{\alpha(i)} n_{\alpha(i)}}] \mathcal{K}[\alpha|\beta] \tilde{\mathcal{T}}_0[A_5^{a_{\beta(i)} n_{\beta(i)}}], \quad (4.86)$$

where  $\mathcal{K}[\alpha|\beta]$  is the KLT kernel. Finally, we also note that the CHY double-copy construction of the  $N$ -point LO scattering amplitude of the vector-type gravitational KK Goldstone bosons  $\widetilde{\mathcal{M}}_0^{\text{dc}}[\mathcal{V}_{n_1}^{\pm 1}, \dots, \mathcal{V}_{n_N}^{\pm 1}]$  can be similarly done.

#### 4.4 Prescription for Warped Double-Copy Construction at Leading Order

Our previous work [27] proved that for the compactified KK gauge/gravity theories, only the toroidal compactification of flat extra dimensions can satisfy the mass spectrum condition and directly realize extended double-copy of massive KK gauge/gravity scattering amplitudes. It was also proven [26] that under the toroidal compactification of flat extra dimensions such massive KK double-copies can be derived as the field theory limit of the massive KLT relations of KK open/closed string amplitudes. But we note that the compactified warped 5d gauge/gravity theories with orbifold  $S^1/\mathbb{Z}_2$  (such as the RS1 model [18]) clearly do not meet this criterion. This is because such warped KK gauge/gravity theories have highly nonlinear mass spectra and the mass spectrum of KK gauge bosons differ from that of the KK gravitons due to the 5d warped metric, whereas coupling coefficients of the KK gauge bosons and of the KK gravitons also differ significantly. Moreover, the  $S^1/\mathbb{Z}_2$  orbifold compactification spoils the KK number conservation and the KK scattering amplitudes do not exhibit single-pole structure in each channel. Nevertheless, as demonstrated in Sections 4.1-4.2, for tree-level KK gauge/gravity scattering amplitudes, we find proper ways (or restriction) to evade these problems. These include: **(i)** the 3-point double-copied KK graviton scattering amplitudes (from that of the KK gauge bosons) exhibit exactly the same kinematic structure as the original 3-point KK graviton amplitudes even without high energy expansion except that we need to set up a proper prescription for the correspondence between the double-copied trilinear KK gauge couplings and the original trilinear KK graviton couplings, whereas under such prescription the double-copy works for the corresponding 3-point gravitational KK Goldstone amplitudes only at the leading-order of high energy expansion;<sup>9</sup> **(ii)** at the leading-order of high energy expansion, the numerators of 4-point scattering amplitudes of KK gauge bosons and of KK Goldstone bosons obey the kinematic Jacobi identities; **(iii)** the double-copied leading-order KK graviton scattering amplitudes (constructed from that of the KK gauge bosons) exhibit exactly the same kinematic structure as the original leading-order amplitudes of KK gravitons except that

---

<sup>9</sup>We found previously [27] that for the gauge/gravity KK Goldstone scattering amplitudes with toroidal compactification of flat extra dimensions, the double-copy construction also works only at the leading order of high energy expansion.



we need to set up a proper prescription for the correspondence between double-copied KK gauge couplings and the original KK graviton couplings.

Based upon the above observation and the double-copy analysis shown in Sections 4.1-4.2, we summarize the prescriptions for successful double-copy constructions of the 3-point and 4-point KK graviton scattering amplitudes in the warped 5d gauge/gravity theories under orbifold compactification of  $S^1/\mathbb{Z}_2$ :

- 1). For the present double-copy construction of  $N$ -point warped KK graviton scattering amplitudes from the corresponding  $N$ -point warped KK gauge boson scattering amplitudes, we first replace the 4d gauge coupling  $g$  by the 4d gravitational coupling  $\kappa$  as follows:

$$g^{N-2} \longrightarrow -\left(\frac{\kappa}{4}\right)^{N-2}. \quad (4.87)$$

For instance, this gives  $g \rightarrow -\kappa/4$  for  $N=3$  and  $g^2 \rightarrow -\kappa^2/16$  for  $N=4$ . This double-copy replacement (4.87) is similar to what we established for the compactified flat 5d gauge/gravity theories [26][27]. The difference is that in the present analysis, due to an overall factor of  $i$  of the  $N$ -point (for  $N=\text{odd}$ ) gauge amplitude, the corresponding double-copied gravitational amplitude has an overall factor of  $-1$ . Thus, the factor  $(-)^{N+1}$  in Refs.[26][27] will be replaced by  $-1$  in this study.

- 2). Then, for the scattering amplitudes of KK gauge bosons (KK Goldstone bosons), we apply the extended massive color-kinematics (CK) duality and make the following group factor replacements as in Eq.(4.3) for the 3-point KK amplitudes, or as in Eqs.(4.22)(4.35) for the 4-point KK amplitudes:

$$\begin{aligned} f^{abc} &\longrightarrow \mathcal{N}[\{\epsilon_j\}], & f^{abc} &\longrightarrow \widetilde{\mathcal{N}}[\epsilon_3], & (3\text{-point amplitudes}), \\ \mathcal{C}_j &\longrightarrow \mathcal{N}_j^0, & \mathcal{C}_j &\longrightarrow \widetilde{\mathcal{N}}_j^0, & (4\text{-point elastic amplitudes}), \\ \mathcal{C}_j &\longrightarrow \mathcal{N}_j^{\text{in}0}, & \mathcal{C}_j &\longrightarrow \widetilde{\mathcal{N}}_j^{\text{in}0}, & (4\text{-point inelastic amplitudes}). \end{aligned} \quad (4.88)$$

- 3). At each KK level- $n$ , we replace the mass-eigenvalue  $M_n$  of KK gauge bosons [determined by Eq.(2.20)] by mass-eigenvalue  $\mathbb{M}_n$  of KK gravitons [determined by Eq.(2.57)], i.e.,  $M_n \rightarrow \mathbb{M}_n$ .
- 4). At each KK level- $n$ , we further replace the involved KK gauge couplings by the corresponding KK gravitational couplings as in Eq.(4.12) for the 3-point KK amplitudes and as in Eqs.(4.29)(4.41)(4.55) for the 4-point KK amplitudes:

$$\begin{aligned} c_3 a_{n_1 n_2 n_3}^2 &\rightarrow \alpha_{n_1 n_2 n_3}, & c_3 a_{n_1 n_2 n_3} \tilde{a}_{n_1 n_2 n_3} &\rightarrow \tilde{\alpha}_{n_1 n_2 n_3}, & c_3 \tilde{a}_{n_1 n_2 n_3}^2 &\rightarrow \tilde{\beta}_{n_1 n_2 n_3}, \\ c_4 \tilde{a}_{nnnn}^2 &\rightarrow \tilde{\beta}_{nnnn}, & c_4^{\text{in}} \tilde{a}_{nnmm}^2 &\rightarrow \tilde{\beta}_{nnmm}, & c_4^{\text{in}} a_{0nn}^4 &\rightarrow \alpha_{0nn}^2, \\ c_4^{\text{in}} a_{000}^2 a_{nn0}^2 &\rightarrow \alpha_{000} \alpha_{nn0}, & c_4^{\text{in}} a_{000}^2 a_{nn0} \tilde{a}_{nn0} &\rightarrow \alpha_{000} \tilde{\alpha}_{nn0}, \end{aligned} \quad (4.89)$$

where the overall coefficient  $c_3$  or  $c_4$  is the relevant normalization factor and its determination is given in Sections 4.1-4.2. Finally, in Section 4.3, we studied the double-copy construction for the  $N$ -point ( $N \geq 4$ ) LO KK Goldstone boson amplitudes. Thus

for the  $N$ -point LO double-copy, we impose the following correspondence (replacement) between the KK gauge and gravity coupling coefficients:

$$c_N \llbracket \cdots f_{n_k} \cdots \tilde{f}_{n_{k'}} \cdots \rrbracket \llbracket \cdots \tilde{f}_{n_k} \cdots f_{n_{k'}} \cdots \rrbracket \longrightarrow \tilde{\alpha}_{n_1 \cdots n_N}, \quad (4.90a)$$

$$c_N \tilde{\alpha}_{n_1 \cdots n_N}^2 \longrightarrow \tilde{\beta}_{n_1 \cdots n_N}. \quad (4.90b)$$

## 5 Conclusions

In this work, we conducted comprehensive analyses on the structure of scattering amplitudes of massive Kaluza-Klein (KK) states in the compactified 5-dimensional warped gauge and gravity theories. We presented systematic formulations of the gauge theory equivalence theorem (GAET) and the gravitational equivalence theorem (GRET) within the  $R_\xi$  gauge and up to loop level. Under the high energy expansion, the GAET quantitatively connects the scattering amplitudes of longitudinal KK gauge bosons to that of the corresponding KK Goldstone bosons, whereas the GRET connects the scattering amplitudes of massive KK gravitons of helicity-0 (or helicity-1) to that of the corresponding gravitational KK Goldstone bosons. A key point of our work is *to take the GAET of KK Yang-Mills gauge theories as the truly fundamental ET formulation*, from which *the GRET of the corresponding KK gravity theories can be reconstructed* by using the double-copy at the leading order (LO) of high energy expansion. We systematically studied the double-copy construction of 3-point and 4-point KK gauge/gravity scattering amplitudes at tree level. We proved that under proper color-kinematics correspondence and gauge-gravity coupling correspondence, the double-copy of 3-point physical KK gauge-boson/graviton scattering amplitudes can be realized without high energy expansion, and for the corresponding 3-point gauge/gravity KK Goldstone scattering amplitudes the double-copy can be achieved at the leading order (LO) of high energy expansion. Moreover, we demonstrated that the double-copy can be realized for 4-point massive scattering amplitudes of KK gauge bosons and of KK gravitons (as well as their corresponding KK Goldstone bosons) to the leading order of high energy expansion. In addition, we can reduce the validity of 4-point-level GAET and GRET to the validity of 3-point-level GAET and GRET. Hence, the GAET and GRET formulations at the level of 3-point KK scattering amplitudes are the most fundamental formulations, from which the GAET and GRET formulations at 4-point-level of KK scattering amplitudes can be inferred. We further derived the GRET from GAET by LO double-copy constructions for 3-point and 4-point KK scattering amplitudes. A more elaborated summary of our results and conclusions in each sections are presented as below, which is followed by a schematic summary as shown in the last paragraph of this section together with Fig. 2.

In Section 2, we proved the formulations of the GAET and GRET within the  $R_\xi$  gauge and up to loop level. The GAET formulation was presented in Eqs.(2.36)-(2.37) and Eq.(2.38), whereas the GRET formulations were given in Eqs.(2.99)(2.103) for KK gravitons with helicity-0 (called type-I) and in Eqs.(2.84)(2.86) for KK gravitons with helicity-1 (called type-II). In essence, the GAET and GRET reflect the geometric ‘‘Higgs’’ mechanism through the KK compactifications, which hold not only for the compactified flat extra dimensions [11, 12][15, 16], but also for the compactified warped extra dimensions as

proved in Sections 2.2-2.3. They determine the high energy behaviors of massive KK graviton scattering amplitudes and ensure the interlancing cancellations among contributions of different KK levels as happened in the scattering amplitudes of KK gauge bosons and of KK gravitons.

In Section 3, we analyzed the structure of 3-point and 4-point massive scattering amplitudes of KK gauge bosons and of KK gravitons (as well as the corresponding scattering amplitudes of KK Goldstone bosons), including the interconnections between the 3-point and 4-point KK amplitudes. In Section 3.1, we explicitly proved the warped GAET and GRET for the fundamental 3-point KK gauge/gravity scattering amplitudes. We found that the nontrivial realization of GAET is given by the 3-point KK gauge boson amplitude with two longitudinal external states and one transverse external state (of *LLT* type), as shown in Eq.(3.4). We proved that to hold this 3-point-level GAET (3.4) requires the condition (3.5) which is directly proved in Eq.(D.6) of Appendix D. Then, we computed the 3-point KK graviton amplitude<sup>10</sup> of helicities  $(\pm 1, \pm 1, \pm 2)$  and its corresponding KK Goldstone amplitude in Eqs.(3.13a)-(3.13b) and Eqs.(3.14a)-(3.14b). To hold the 3-point-level GRET (3.15) imposes the condition (3.16) which we proved in Eq.(D.7) of Appdenix D. We further computed the 3-point KK graviton amplitude of helicities  $(0, 0, \pm 2)$  and its corresponding KK Goldstone amplitude in Eqs.(3.19a)-(3.19b). To hold the 3-point-level GRET (3.20) leads to the nontrivial condition (3.21) which we proved in Eq.(D.20) of Appdenix D. We also computed the mixed 3-point KK graviton amplitude of helicities  $(\pm 1, \pm 1, 0)$  and its corresponding KK Goldstone amplitude in Eq.(3.23). To hold the GRET (3.25) imposes the condition (3.26) which was proved in (D.22) of Appendix D.

In Section 3.2, we further demonstrated that the validity of the warped GAET and GRET for 4-point KK scattering amplitudes can be effectively reduced to the validity of these theorems for the 3-point KK scattering amplitudes. For the 4-point elastic and inelastic scattering amplitudes of KK gauge bosons and KK Goldstone bosons, we explicitly established the warped GAET as in Eqs.(3.41), (3.50), and (3.58) respectively. We proved that the validity of the 4-point-level GAET in these cases is ensured by the validity of the fundamental 3-point-level GAET (3.4). Then, we analyzed the 4-point scattering amplitudes of KK gravitons and gravitational KK Goldstone bosons. We explicitly established the warped GRET as in Eq.(3.67) for the 4-body elastic scattering channel. The validity of the 4-point-level GRET (3.67) relies on the sum rule condition (3.68), while the proof of Eq.(3.68) requires the sum rule condition (3.22) to play the key role, where the condition (3.22) just ensures the validity of the 3-point-level GRET (3.20) in the case of  $(n_1, n_2, n_3) = (n, n, j)$ . This shows that validity of the 4-point-level GRET (3.67) is reduced to the validity of the 3-point-level GRET (3.20). We further computed the 4-point inelastic scattering amplitudes of KK gravitons and of KK Goldstone bosons as in Eqs.(3.73a)-(3.73b). Under high energy expansion, we found that the leading-order amplitudes (3.74a)-(3.74b) are simple enough in this case. Thus, we explicitly established the 4-point-level GRET (3.75) without a need of additional sum rule condition.


In Section 4, we studied the double-copy construction of the massive gravitational KK

---

<sup>10</sup>We derived the most general 3-point scattering amplitude of KK gravitons in Eq.(C.15) of Appendix C.

scattering amplitudes from the corresponding massive KK gauge scattering amplitudes for the warped 5d gauge and gravity theories with the orbifold compactification of  $S^1/\mathbb{Z}_2$ . This is nontrivial and challenging since it was proved [27] that the direct construction of double-copy for  $N$ -point massive gauge/gravity KK scattering amplitudes (with  $N \geq 4$ ) can directly work out only for toroidal compactifications with flat extra dimensions. Nevertheless, we could realize the double-copy construction for warped gauge/gravity theories with proper restrictions and prescriptions so as to evade the previous conditions [27]. In Section 4.1, we newly proved that the double-copy can be constructed for the 3-point full scattering amplitudes of KK gravitons at tree level for warped gauge/gravity theories. We set up the 3-point color-kinematics (CK) correspondence and the gauge/gravity coupling correspondence as in Eq.(4.3), and the KK mass replacement  $M_n \rightarrow \mathbb{M}_n$  at each KK level. With these, we first presented the general 3-point double-copy formulas (with any polarization tensors of physical KK graviton states) as in Eqs.(4.6)-(4.8). Then, we explicitly constructed the 3-point KK graviton amplitudes with helicities  $(\pm 1, \pm 1, \pm 2)$  and  $(0, 0, \pm 2)$  in Eq.(4.9) and Eq.(4.10) respectively. We further derived their corresponding gravitational KK Goldstone boson amplitudes in Eqs.(4.11a)-(4.11b). The required conversions of gauge-gravity coupling constants are given in Eq.(4.12). With these we established successful double-copy constructions of the GRET (4.15a) and (4.15b) (in warped KK gravity theory) from the GAET (3.4) (in the warped KK gauge theory) at the level of 3-point KK scattering amplitudes.

In Section 4.2, we demonstrated that the double-copy construction can be achieved for the 4-point KK gauge/gravity scattering amplitudes at the leading-order (LO) of high energy expansion. For the 4-point elastic scattering  $(n, n) \rightarrow (n, n)$ , the LO amplitudes of KK gauge bosons and of KK Goldstone bosons have their effective numerators connected by the generalized gauge transformations (4.20) and obey the kinematic Jacobi identities (4.21). The double-copy of these two LO amplitudes gives the LO amplitudes of KK gravitons and of the corresponding gravitational KK Goldstone bosons as in Eqs.(4.23) and (4.25). Using the coupling conversion (4.29), we derived the double-copied LO gravitational KK amplitudes (4.30) which agree with the same amplitudes (4.26) (by explicit Feynman diagram approach). In parallel, we studied the double-copy constructions for the LO inelastic KK gauge/gravity scattering amplitudes, including the processes  $(n, n) \rightarrow (m, m)$  (with  $n \neq m$ ),  $(n, m) \rightarrow (\ell, q)$  (with  $n \neq m \neq \ell \neq q$ ), and  $(0, 0) \rightarrow (n, n)$  (with  $n > 0$ ). We found that their effective numerators satisfy the kinematic Jacobi identity respectively, as shown in Eqs.(4.34), (4.46)-(4.47), and so on. For these inelastic processes, we presented the double-copied inelastic amplitudes of KK gravitons and of gravitational KK Goldstone bosons as in Eqs.(4.42), (4.48), and (4.52)(4.55). In Section 4.3, we further established that this LO double-copy construction can be extended to the general  $N$ -point KK scattering amplitudes with  $N \geq 4$ . In Section 4.4, we summarized a set of well-defined prescriptions for the successful tree-level double-copy constructions in the warped massive KK gauge/gravity theories, which include the double-copy constructions for 3-point full KK gauge/gravity amplitudes and the double-copy constructions for the 4-point leading-order KK gauge/gravity amplitudes.



**Figure 2:** Schematic Summary of the present analyses: Equivalence Theorem and Double-Copy Correspondences from 3-point scattering amplitudes to 4-point scattering amplitudes and from massive KK gauge scattering amplitudes to massive KK gravitational scattering amplitudes at the leading order of high energy expansion.

Finally, we present a schematic summary of the present analyses as in Fig. 2. In this figure, we start from the horizontal bottom row in which all entries describe the basic 3-point KK scattering amplitudes. From the left to right, the long equality sign linking the first two entries gives the “GAET(3)” which connects the 3-point LO longitudinal KK gauge boson amplitude  $\mathcal{T}_0^{(3)}[A_L^n]$  to the corresponding LO KK Goldstone boson amplitude  $\tilde{\mathcal{T}}_0^{(3)}[A_5^n]$ ; then the long equality sign linking the last two entries gives the “GRET(3)” which connects the 3-point LO longitudinal KK graviton amplitude  $\mathcal{M}_0^{(3)}[h_L^n]$  to the corresponding LO gravitational KK Goldstone boson amplitude  $\tilde{\mathcal{M}}_0^{(3)}[\phi_n]$ ; in the middle, the horizontal arrow indicates the double-copy “DC<sub>3</sub>” which constructs the 3-point LO longitudinal KK graviton (KK Goldstone) amplitude from the 3-point LO longitudinal KK gauge boson (KK Goldstone) amplitude. After this, we see that all the entries and equality signs (or arrow) in the top row are doing the same jobs as those in the bottom row except that all entries in the top row deal with the 4-point KK gauge/gravity scattering amplitudes. Finally, the vertical arrows from bottom to top indicate the 4-point-level “GAET(4)” and “GRET(4)” can be reduced to (reconstructed from) the fundamental “GAET(3)” and “GRET(3)” for the basic 3-point KK gauge (gravity) amplitudes; and furthermore, we can construct the GRET(3) from GAET(3) through double-copy.

### Acknowledgments

This research was supported in part by the National NSF of China under grants 12175136 and 11835005. YH is supported in part by the Northwestern University Amplitudes and Insight group, the Department of Physics and Astronomy, and Weinberg College.

## Appendix

### A Kinematics of KK Particle Scattering

In this Appendix we present the kinematics of 3 and 4 KK particle scattering processes in the (3+1)d spacetime. We choose the 4d Minkowski metric tensor  $\eta^{\mu\nu} = \eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$ .

For the 3 KK particle scattering, we define the 4-momenta of the external particles as follows:

$$\begin{aligned} p_1^\mu &= (E_1, ks_\theta, 0, kc_\theta), & E_1 &= \sqrt{M_1^2 + k^2}, \\ p_2^\mu &= (E_2, -ks_\theta, 0, kc_\theta), & E_2 &= \sqrt{M_2^2 + k^2}, \\ p_3^\mu &= -(E_3, 0, 0, 2kc_\theta), & E_3 &= \sqrt{M_3^2 + 4k^2c_\theta^2}, \end{aligned} \quad (\text{A.1})$$

where  $k = |\vec{p}|$ ,  $(s_\theta, c_\theta) = (\sin\theta, \cos\theta)$ ,  $p_j^2 = -M_j^2$  (with  $j = 1, 2, 3$ ), and all momenta are outgoing by convention.

Using the energy conservation condition  $E_1 + E_2 = E_3$ , we can solve the magnitude of 3-momentum  $k = |\vec{p}|$  as a function of the scattering angle  $\theta$ :

$$\begin{aligned} k &= \left\{ \frac{1}{2} \sqrt{M_3^4 + 4 \cos^2\theta [M_1^4 + M_2^4 - (M_1^2 + M_2^2)M_3^2 + 2M_1^2M_2^2 \cos 2\theta]} \right. \\ &\quad \left. + \frac{\csc 2\theta}{2} [M_3^2 \cot 2\theta - (M_1^2 + M_2^2) \cot \theta] \right\}^{\frac{1}{2}}. \end{aligned} \quad (\text{A.2})$$

Alternatively, we can express  $\theta$  as a function of  $k$ :

$$\cos\theta = \frac{1}{2k} \left[ 2k^2 + M_1^2 + M_2^2 + 2\sqrt{(k^2 + M_1^2)(k^2 + M_2^2)} - M_3^2 \right]^{\frac{1}{2}}. \quad (\text{A.3})$$

In high energy limit, we set  $k \rightarrow \infty$  and  $\theta \rightarrow 0$ . Thus, we further expand  $k$  and  $\theta$  as follows:

$$k = \frac{M_3^2 - 2(M_1^2 + M_2^2)}{4 \sin^2\theta} + \frac{(M_1^2 - M_2^2)^2}{4[M_3^2 - 2(M_1^2 + M_2^2)]} + O(\theta^2), \quad (\text{A.4a})$$

$$\cos\theta = 1 - \frac{1}{8k^2} [M_3^2 - 2(M_1^2 + 2M_2^2)] + O(k^{-4}), \quad (\text{A.4b})$$

$$\sin\theta = \frac{1}{2k} \sqrt{M_3^2 - 2(M_1^2 + M_2^2)} + O(k^{-4}). \quad (\text{A.4c})$$

From the above formulas, we see that to have real solutions of  $k$  and  $\theta$  requires  $M_3 \geq \sqrt{2(M_1^2 + M_2^2)}$ .

For the four-body scattering of KK states  $X_1X_2 \rightarrow X_3X_4$ , the 4-momenta in the center-of-mass frame are defined as follows:

$$\begin{aligned} p_1^\mu &= -(E_1, 0, 0, k), & p_2^\mu &= -(E_2, 0, 0, -k), \\ p_3^\mu &= (E_3, k's_\theta, 0, k'c_\theta), & p_4^\mu &= (E_4, -k's_\theta, 0, -k'c_\theta), \end{aligned} \quad (\text{A.5})$$

where we define the following Mandelstam variables:

$$s = -(p_1 + p_2)^2, \quad t = -(p_1 + p_4)^2, \quad u = -(p_1 + p_3)^2, \quad (\text{A.6})$$

from which we have  $s + t + u = \sum_{j=1}^4 M_j^2$ . In addition, the momenta  $k$  and  $k'$  in Eq.(A.5) are defined as follows:

$$\begin{aligned} k &= \frac{1}{2\sqrt{s}} \left( [s - (M_1 + M_2)^2] [s - (M_1 - M_2)^2] \right)^{\frac{1}{2}}, \\ k' &= \frac{1}{2\sqrt{s}} \left( [s - (M_3 + M_4)^2] [s - (M_3 - M_4)^2] \right)^{\frac{1}{2}}. \end{aligned} \quad (\text{A.7})$$

We further define the massless Mandelstam variables as  $(s_0, t_0, u_0) = (s, t, u)|_{M_j=0}$ , and thus we have:

$$s_0 = 4k^2, \quad t_0 = -\frac{s_0}{2}(1 + c_\theta), \quad u_0 = -\frac{s_0}{2}(1 - c_\theta), \quad (\text{A.8})$$

where  $j=1, 2, 3, 4$  and the sum of these Mandelstam variables is given by  $s_0 + t_0 + u_0 = 0$ .

A massive KK graviton  $h_n^{\mu\nu}$  in 4d has 5 physical helicity states and their polarization tensors are represented by

$$\epsilon_{\pm 2}^{\mu\nu} = \epsilon_{\pm}^{\mu} \epsilon_{\pm}^{\nu}, \quad \epsilon_{\pm 1}^{\mu\nu} = \frac{1}{\sqrt{2}} (\epsilon_{\pm}^{\mu} \epsilon_L^{\nu} + \epsilon_L^{\mu} \epsilon_{\pm}^{\nu}), \quad \epsilon_L^{\mu\nu} = \frac{1}{\sqrt{6}} (\epsilon_+^{\mu} \epsilon_-^{\nu} + \epsilon_-^{\mu} \epsilon_+^{\nu} + 2\epsilon_L^{\mu} \epsilon_L^{\nu}), \quad (\text{A.9})$$

where  $\epsilon_{\pm}^{\mu}$  and  $\epsilon_L^{\mu}$  are the spin-1 polarization vectors:

$$\epsilon_{\pm}^{\mu} = \pm \frac{e^{\mp i\phi}}{\sqrt{2}} (0, c_\theta c_\phi \pm i s_\phi, c_\theta s_\phi \mp i c_\phi, -s_\theta), \quad \epsilon_L^{\mu} = \frac{1}{\mathbb{M}_n} (k, E_n s_\theta c_\phi, E_n s_\theta s_\phi, E_n c_\theta). \quad (\text{A.10})$$

In the above, the KK graviton  $h_n^{\mu\nu}$  moves in an arbitrary direction with polar and azimuthal angles  $(\theta, \phi)$ . For instance, consider the 4-body elastic scattering at the KK level- $n$ , we have  $E_{n_j} = E$  and  $\mathbb{M}_{n_j} = M$ . Then, we can define the following polarization vectors in the center-of-mass frame:

$$\begin{aligned} \epsilon_{1,\pm}^{\mu} &= \frac{1}{\sqrt{2}} (0, \mp 1, i, 0), & \epsilon_{1,L}^{\mu} &= -\frac{E}{M} (\beta, 0, 0, 1), \\ \epsilon_{2,\pm}^{\mu} &= \frac{1}{\sqrt{2}} (0, \pm 1, i, 0), & \epsilon_{2,L}^{\mu} &= -\frac{E}{M} (\beta, 0, 0, -1), \\ \epsilon_{3,\pm}^{\mu} &= \frac{1}{\sqrt{2}} (0, \pm c_\theta, -i, \mp s_\theta), & \epsilon_{3,L}^{\mu} &= \frac{E}{M} (\beta, s_\theta, 0, c_\theta), \\ \epsilon_{4,\pm}^{\mu} &= \frac{1}{\sqrt{2}} (0, \mp i c_\theta, -i, \pm s_\theta), & \epsilon_{4,L}^{\mu} &= \frac{E}{M} (\beta, -s_\theta, 0, -c_\theta), \end{aligned} \quad (\text{A.11})$$

where  $\beta = (1 - M^2/E^2)^{1/2}$ . The polarizations for the inelastic scattering processes can be derived in a similar way.

## B BRST Quantization for GRET Formulation

In this Appendix, we provide more detailed derivations to support the formulation of warped GRET in the main text of Section 2.3. This includes the BRST quantization used in Section 2.3.1, the formulation of the warped GRET type-I as presented in Section 2.3.3, and the formulation of the warped GRET type-II as given in Section 2.3.2.

## B.1 BRST Quantization for Warped 5d Gravity

For the BRST quantization approach, the ghost fields are introduced in the path integral formulation. The 5d Faddeev-Popov ghost Lagrangian for the 5d GR with warped metric takes the following form:

$$\hat{\mathcal{L}}_{\text{FP}} = e^{3A(z)} \hat{c}^M \hat{\mathbf{s}} \hat{\mathcal{F}}_M, \quad (\text{B.1})$$

where the 5d gauge-fixing functions  $\hat{\mathcal{F}}_M = (\hat{\mathcal{F}}_\mu, \hat{\mathcal{F}}_5)$  are given by Eq.(2.44). The BRST transformations for 5d graviton, ghost and anti-ghost fields take the following form:

$$\begin{aligned} \hat{\mathbf{s}} \hat{h}_{MN} &= -\partial_M \hat{c}_N - \partial_N \hat{c}_M + \hat{\kappa} (\hat{h}_{MN} \partial_P - \hat{h}_{MP} \partial_N - \hat{h}_{NP} \partial_M) \hat{c}^P, \\ \hat{\mathbf{s}} \hat{c}_M &= \hat{\kappa} \hat{c}_N \partial^N \hat{c}_M, \quad \hat{\mathbf{s}} \hat{\bar{c}}_M = -2\xi^{-1} \hat{\mathcal{F}}_M. \end{aligned} \quad (\text{B.2})$$

The BRST transformations exhibit nilpotency and obey  $\hat{\mathbf{s}}^2 = 0$ .

Next, we make KK expansions for ghost and anti-ghost fields as follows:

$$\hat{c}^\mu(x, z) = \frac{1}{\sqrt{L}} \sum_{n=0}^{\infty} c_n^\mu(x) \mathbf{u}_n(z), \quad \hat{\bar{c}}^\mu(x, z) = \frac{1}{\sqrt{L}} \sum_{n=0}^{\infty} \bar{c}_n^\mu(x) \mathbf{u}_n(z), \quad (\text{B.3a})$$

$$\hat{c}^5(x, z) = \frac{1}{\sqrt{L}} \sum_{n=1}^{\infty} c_n^5(x) \mathbf{v}_n(z), \quad \hat{\bar{c}}^5(x, z) = \frac{1}{\sqrt{L}} \sum_{n=1}^{\infty} \bar{c}_n^5(x) \mathbf{v}_n(z). \quad (\text{B.3b})$$

Using the KK expansions (2.48), we derive the BRST transformations for the KK fields in 4d spacetime:

$$\begin{aligned} \hat{\mathbf{s}} h_n^{\mu\nu} &= -\partial^\mu c_n^\nu - \frac{1}{2} \eta^{\mu\nu} \mathbb{M}_n c_n^5 + \frac{\kappa}{2} [\mathbf{u}_n \mathbf{u}_m \mathbf{u}_\ell] (h_m^{\mu\nu} \partial^\alpha - 2h_m^{\mu\alpha} \partial^\nu) c_\alpha^\ell + \frac{\kappa}{2} [\mathbf{u}_n \mathbf{u}_m \mathbf{v}_\ell] \mathbb{M}_\ell h_m^{\mu\nu} c_\ell^5 \\ &\quad + \frac{\kappa}{2\sqrt{2}} [\mathbf{u}_n \mathbf{v}_m \mathbf{u}_\ell] \eta^{\mu\nu} \mathbb{M}_\ell \mathcal{V}_m^\alpha c_\alpha^\ell - \frac{\kappa}{\sqrt{2}} [\mathbf{u}_n \mathbf{v}_m \mathbf{v}_\ell] \mathcal{V}_m^\mu \partial^\nu c_\ell^5 + \frac{\kappa}{\sqrt{6}} ([\mathbf{u}_n \mathbf{w}_m \mathbf{u}_\ell] \phi_m \partial^\mu c_\ell^\nu \\ &\quad - [\mathbf{u}_n \mathbf{w}_m \mathbf{v}_\ell] \eta^{\mu\nu} \mathbb{M}_\ell \phi_m c_\ell^5) + (\mu \leftrightarrow \nu), \end{aligned} \quad (\text{B.4a})$$

$$\begin{aligned} \hat{\mathbf{s}} \mathcal{V}_n^\mu &= -\sqrt{2} \partial^\mu c_n^5 + \sqrt{2} \mathbb{M}_n c_n^\mu + \sqrt{2} \kappa [\mathbf{v}_n \mathbf{u}_m \mathbf{u}_\ell] \mathbb{M}_\ell h_m^{\mu\nu} c_\nu^\ell + \kappa [\mathbf{v}_n \mathbf{v}_m \mathbf{u}_\ell] (\mathcal{V}_m^\mu \partial^\nu - \mathcal{V}_m^\nu \partial^\mu) c_\nu^\ell \\ &\quad - \frac{\kappa}{\sqrt{3}} ([\mathbf{v}_n \mathbf{w}_m \mathbf{u}_\ell] \mathbb{M}_\ell \phi_m c_\ell^\mu + 2[\mathbf{v}_n \mathbf{w}_m \mathbf{v}_\ell] \phi_m \partial^\mu c_\ell^5), \end{aligned} \quad (\text{B.4b})$$

$$\begin{aligned} \hat{\mathbf{s}} \phi_n &= -\sqrt{6} \mathbb{M}_n c_n^5 + \sqrt{3} \kappa [\mathbf{w}_n \mathbf{v}_m \mathbf{u}_\ell] \mathbb{M}_\ell \mathcal{V}_m^\mu c_\mu^\ell + \kappa [\mathbf{w}_n \mathbf{w}_m \mathbf{u}_\ell] \phi_m \partial_\mu c_\ell^\mu \\ &\quad - \kappa [\mathbf{w}_n \mathbf{w}_m \mathbf{v}_\ell] \mathbb{M}_\ell \phi_m c_\ell^5, \end{aligned} \quad (\text{B.4c})$$

$$\hat{\mathbf{s}} c_n^\mu = \kappa [\mathbf{u}_n \mathbf{u}_m \mathbf{u}_\ell] c_m^\nu \partial_\nu c_\ell^\mu - \kappa [\mathbf{u}_n \mathbf{v}_m \mathbf{u}_\ell] \mathbb{M}_\ell c_m^5 c_\ell^\mu, \quad (\text{B.4d})$$

$$\hat{\mathbf{s}} c_n^5 = \kappa [\mathbf{v}_n \mathbf{u}_m \mathbf{v}_\ell] c_m^\mu \partial_\mu c_\ell^5 + \kappa [\mathbf{v}_n \mathbf{v}_m \mathbf{v}_\ell] \mathbb{M}_\ell c_m^5 c_\ell^5, \quad (\text{B.4e})$$

$$\hat{\mathbf{s}} \bar{c}_n^\mu = -\frac{2}{\xi_n} \mathcal{F}_n^\mu, \quad \hat{\mathbf{s}} \bar{c}_n^5 = -\frac{2}{\xi_n} \mathcal{F}_n^5, \quad (\text{B.4f})$$

where  $\kappa = \hat{\kappa}/\sqrt{L}$  and the brackets  $[\dots]$  are defined in Appendix D.

## B.2 Gravitational ET of Type-I

In the Sections 2.3.2-2.3.3 of the main text, we have formulated the GRET which quantitatively connects the scattering amplitude of KK gravitons  $h_n^{\mu\nu}$  with helicities  $(0, \pm 1)$  to that



of the corresponding gravitational KK Goldstone bosons  $(\phi_n, \mathcal{V}_n^{\pm 1})$  respectively. In this and next sub-Appendices, we will provide more detailed derivations to support the results in the main text.

We study the combination of gauge-fixing functions  $\partial_\mu \mathcal{F}_n^\mu - \xi_n \mathbb{M}_n \mathcal{F}_n^5$  which eliminates the KK vector-Goldstone field  $\mathcal{V}_n^\mu$ . With this we derive the following formula in the momentum space:

$$-ik_\mu \mathcal{F}_n^\mu - \xi_n \mathbb{M}_n \mathcal{F}_n^5 = -k_\mu k_\nu h_n^{\mu\nu} + \frac{1}{2} \left[ (2 - \xi_n^{-1}) k^2 - \xi_n \mathbb{M}_n^2 \right] h_n + \sqrt{\frac{3}{2}} \xi_n^2 \mathbb{M}_n^2 \phi_n, \quad (\text{B.5})$$

where the rescaling  $\phi_n \rightarrow \sqrt{\frac{2}{3}} \phi_n$  is made according to Eq.(2.63). Applying the on-shell condition  $k^2 = -\mathbb{M}_n^2$ , we further derive Eq.(B.5) in the following form:

$$ik_\mu \mathcal{F}_n^\mu + \xi_n \mathbb{M}_n \mathcal{F}_n^5 = \sqrt{\frac{3}{2}} \mathbb{M}_n^2 \tilde{\mathbb{F}}_n, \quad (\text{B.6a})$$

$$\tilde{\mathbb{F}}_n = \sqrt{\frac{2}{3}} h_n^S + \frac{1}{\sqrt{6}} (2 + \xi_n - \xi_n^{-1}) h_n - \xi_n^2 \phi_n = \mathbf{K}_n^T \mathbf{H}_n \quad (\text{B.6b})$$

$$= \sqrt{\frac{2}{3}} (h_n^S + h_n) - \phi_n, \quad (\text{for } \xi_n = 1), \quad (\text{B.6c})$$

$$\mathbf{K}_n = \left( \sqrt{\frac{2}{3}} \varepsilon_{\mu\nu}^S + \frac{1}{\sqrt{6}} (2 + \xi_n - \xi_n^{-1}) \eta_{\mu\nu}, -\xi_n^2 \right)^T, \quad \mathbf{H}_n = (h_n^{\mu\nu}, \phi_n)^T, \quad (\text{B.6d})$$

$$= \left( \sqrt{\frac{2}{3}} (\varepsilon_{\mu\nu}^S + \eta_{\mu\nu}), -1 \right)^T, \quad (\text{for } \xi_n = 1). \quad (\text{B.6e})$$

We then derive a BRST identity involving the gauge-fixing function  $\tilde{\mathbb{F}}_n$  in Eq.(B.6b):

$$\langle 0 | T \tilde{\mathbb{F}}_n \mathbf{H}_m^T | 0 \rangle(k) = -\frac{\xi_n}{\sqrt{6} \mathbb{M}_n^2} \langle 0 | T \widehat{\mathbf{S}} \mathbf{H}_m^T (ik_\mu \bar{c}_n^\mu + \xi_n \mathbb{M}_n \bar{c}_n^5) | 0 \rangle(k), \quad (\text{B.7})$$

and it can be further rewritten in the following form by utilizing Eq.(B.6):

$$\mathbf{K}_n^T \mathcal{D}_{nm}(k) = -\mathbf{X}_{nm}^T(k). \quad (\text{B.8})$$

In the above, we adopt for simplicity the 't Hooft-Feynman gauge condition ( $\xi_n = 1$ ). Thus, in Eq.(B.8), we define the following notations:

$$\begin{aligned} \mathcal{D}_{nm}(k) &= \langle 0 | T \mathbf{H}_n \mathbf{H}_m^T | 0 \rangle(k), \quad \mathbf{X}_{nm}(k) = \underline{\mathbf{X}}_{mj}(k) \mathcal{S}_{jn}(k), \\ \underline{\mathbf{X}}_{mj}(k) &= \frac{1}{2\sqrt{6} M_j^2} \left( \frac{\langle 0 | T \widehat{\mathbf{S}} h_m^{\mu\nu} | \bar{c}_j \rangle(k)}{\langle 0 | T \widehat{\mathbf{S}} \phi_m | \bar{c}_j \rangle(k)} \right) = \frac{1}{2\sqrt{6}} \left( \frac{(2k^\mu k^\nu / M_j^2 - \eta^{\mu\nu}) [\delta_{mj} + \Delta_{mj}^{(3)}(k^2)]}{-\sqrt{6} [\delta_{mj} + \tilde{\Delta}_{mj}^{(4)}(k^2)]} \right), \\ \mathcal{S}_{jn}(k) &= \langle 0 | T c_j \bar{c}_n | 0 \rangle(k), \end{aligned} \quad (\text{B.9})$$

with  $c_n \equiv i\epsilon_\mu^S c_n^\mu + c_n^5$ ,  $\bar{c}_n \equiv i\epsilon_\mu^S \bar{c}_n^\mu + \bar{c}_n^5$ , and  $\epsilon_\mu^S = k_\mu / \mathbb{M}_n$ . In the above the external momentum is chosen as incoming. The formulas (B.8)-(B.9) just give Eqs.(2.91)-(2.92) in the text. In addition, we see that the quantities  $\Delta_{mj}^{(3)}(k^2)$  and  $\tilde{\Delta}_{mj}^{(4)}(k^2)$  are of loop order and are generated by the non-linear terms of the BRST transformations (B.4a) and (B.4c) of  $h_n^{\mu\nu}$  and  $\phi_n$ .

Next, we use the identity (2.66) and Eq.(B.6) to deduce an identity containing the external state  $\tilde{\mathbb{F}}_n(k)$ :

$$0 = \langle 0 | \tilde{\mathbb{F}}_n(k) \cdots \Phi | 0 \rangle = \mathbf{K}_n^T \mathcal{D}_{nm}(k) \langle 0 | \mathbf{H}_m(k) \cdots \Phi | 0 \rangle$$

$$= -\mathbf{X}_{nm}^T(k) \mathcal{M}[\underline{\mathbf{H}}_m(k), \dots, \Phi], \quad (\text{B.10})$$

which leads to the following identity,

$$\mathcal{M}[\tilde{\mathbb{F}}_n(k), \dots, \Phi] = 0, \quad (\text{B.11})$$

with the amputated external state  $\tilde{\mathbb{F}}_n(k)$  given by<sup>11</sup>

$$\tilde{\mathbb{F}}_n = \sqrt{\frac{2}{3}} (h_n^S - \frac{1}{2} h_n) - C_{nm} \phi_m = h_n^L - \Omega_n, \quad (\text{B.12a})$$

$$\Omega_n = C_{nm} \phi_m + \tilde{\Delta}_n, \quad \tilde{\Delta}_n = \frac{1}{\sqrt{6}} h_n + \tilde{v}_n, \quad \tilde{v}_n = v_{\mu\nu} h_n^{\mu\nu}. \quad (\text{B.12b})$$

In the above,  $C_{nm}$  is a multiplicative modification factor induced at loop level:

$$C_{nm}(k^2) = \left[ \frac{\mathbf{1} + \tilde{\Delta}^{(4)}(k^2)}{\mathbf{1} + \Delta^{(3)}(k^2)} \right]_{mn} = \delta_{nm} + O(\text{loop}), \quad (\text{B.13})$$

where the matrix form is used such that  $(\Delta^{(3)})_{jj'} = \Delta_{jj'}^{(3)}$  and  $(\tilde{\Delta}^{(4)})_{jj'} = \tilde{\Delta}_{jj'}^{(4)}$  with the matrix elements  $(\Delta_{jj'}^{(3)}, \tilde{\Delta}_{jj'}^{(4)})$  from Eq.(B.9). The above Eqs.(B.10)-(B.13) just reproduce the formulas (2.93)-(2.96) in the main text.

Thus, from Eq.(B.11) we deduce another general identity for gravitational equivalence theorem (GRET):

$$\mathcal{M}[\tilde{\mathbb{F}}_{n_1}(k_1), \dots, \tilde{\mathbb{F}}_{n_N}(k_N), \Phi] = 0. \quad (\text{B.14})$$

From this, we further derive the following GRET identity:

$$\mathcal{M}[h_{n_1}^L(k_1), \dots, h_{n_N}^L(k_N), \Phi] = \mathcal{M}[\Omega_{n_1}(k_1), \dots, \Omega_{n_N}(k_N), \Phi]. \quad (\text{B.15})$$

We can directly prove this identity by expanding the right-hand side of Eq.(B.15) with each external state replaced by  $\Omega_n = h_n^L - \tilde{\mathbb{F}}_n$  and further using Eq.(B.14). This just reproduces the GRET identity (2.98) in the main text.

We note that at tree level the LSZ reduction may be implemented directly. Adopting the 't Hooft-Feynman gauge ( $\xi_n = 1$ ) and using the identity (2.66), we can derive a new identity involving the external line  $\tilde{\mathbb{F}}_n(k)$  as follows:

$$\begin{aligned} 0 &= \langle 0 | \tilde{\mathbb{F}}_n(k) \dots \bar{\Phi} | 0 \rangle = \sqrt{\frac{2}{3}} (\varepsilon_{\mu\nu}^S + \eta_{\mu\nu}) \langle 0 | h_n^{\mu\nu}(k) \dots \Phi | 0 \rangle - \langle 0 | \phi_n(k) \dots \Phi | 0 \rangle \\ &= \sqrt{\frac{2}{3}} (\varepsilon_{\mu\nu}^S + \eta_{\mu\nu}) \mathcal{D}_{h, nm}^{\mu\nu\alpha\beta}(k) \mathcal{M}[h_m^{\alpha\beta}(k), \dots, \Phi] - \mathcal{D}_{\phi, nm}(k) \mathcal{M}[\phi_m(k), \dots, \Phi] \\ &= \mathcal{D}_{\phi, nm}(k) \mathcal{M}[\sqrt{\frac{2}{3}} (h_m^S - \frac{1}{2} h_m) - \phi_m, \dots, \Phi] \\ &\equiv \bar{\mathcal{D}}_\phi(k) \mathcal{M}[\tilde{\mathbb{F}}_n(k), \dots, \Phi], \end{aligned} \quad (\text{B.16})$$

where  $\mathcal{D}_{\phi, nm}(k) = \delta_{nm} \bar{\mathcal{D}}_\phi(k)$  and the LSZ-amputated external state  $\tilde{\mathbb{F}}_n(k)$  is given by Eq.(B.12) with  $C_{nm} = 1$ . We have also used the tree-level propagators (2.65a) and (2.65c)

<sup>11</sup>We note that after the LSZ amputation in general  $R_\xi$  gauge at tree-level, the coefficient of  $h_n$  in  $\tilde{\mathbb{F}}_n$  should be  $-1/2$  and corrects the coefficient of  $h_n$  as in our previous works [15, 16], but this does not affect all the conclusions therein [15, 16].

for  $\xi_n = 1$ . In addition, we can extend the above derivation to the general  $R_\xi$  gauge. Using the  $h_n^{\mu\nu}$  and  $\phi_n$  propagators in the  $R_\xi$  gauge as given by Eqs.(C.4a)-(C.4b), we derive the following identity of propagators at tree level:

$$[\varepsilon_{\mu\nu}^S + \frac{1}{2}(\xi_n - \xi_n^{-1})\eta_{\mu\nu}] \mathcal{D}_{h, nm}^{\mu\nu\alpha\beta}(k) = \xi_n^2 \mathcal{D}_{nm}(k) (\varepsilon_S^{\alpha\beta} - \frac{1}{2}\eta^{\alpha\beta}), \quad (\text{B.17})$$

we derive the following tree-level identity with amputated external state  $\tilde{\mathbb{F}}_n(k)$ :

$$\begin{aligned} 0 &= \langle 0 | \tilde{\mathbb{F}}_n(k) \cdots \bar{\Phi} | 0 \rangle \\ &= \sqrt{\frac{2}{3}} [\varepsilon_{\mu\nu}^S + \frac{1}{2}(\xi_n - \xi_n^{-1})\eta_{\mu\nu}] \mathcal{D}_{h, nm}^{\mu\nu\alpha\beta}(k) \mathcal{M}[h_m^{\alpha\beta}(k), \cdots, \Phi] - \xi_n^2 \mathcal{D}_{\phi, nm}(k) \mathcal{M}[\phi_m(k), \cdots, \Phi] \\ &= \xi_n^2 \mathcal{D}_{nm}(k) \mathcal{M}\left[\sqrt{\frac{2}{3}}(h_m^S - \frac{1}{2}h_m) - \phi_m, \cdots, \Phi\right]. \end{aligned} \quad (\text{B.18})$$

Thus, we have amputated external state  $\tilde{\mathbb{F}}_n(k)$  obey the identity:

$$\mathcal{M}[\tilde{\mathbb{F}}_n(k), \cdots, \Phi] = 0, \quad (\text{B.19})$$

where we have defined the following quantities at tree level,

$$\tilde{\mathbb{F}}_n = \sqrt{\frac{2}{3}}(h_n^S - \frac{1}{2}h_n) - \phi_n = h_n^L - \Omega_n, \quad (\text{B.20a})$$

$$\Omega_n = \phi_n + \tilde{\Delta}_n, \quad \tilde{\Delta}_n = \frac{1}{\sqrt{6}}h_n + \tilde{v}_n, \quad \tilde{v}_n = v_{\mu\nu}h_n^{\mu\nu}. \quad (\text{B.20b})$$

We can readily extend the identity (B.19) to the case with  $N$  external states  $\tilde{\mathbb{F}}_n$ , which then reproduces the form of the GRET (B.14) for  $R_\xi$  gauge and at tree level.

### B.3 Gravitational ET of Type-II

In Section 2.3.2 of the main text, we have formulated the KK GRET of type-II which connects the scattering amplitude of KK gravitons  $h_n^{\mu\nu}$  with helicity  $\pm 1$  to that of the corresponding gravitational KK vector Goldstone bosons  $\mathcal{V}_n^\mu$  with the same helicity. In this sub-Appendix, we will provide detailed derivations to support the main text.

We start with the general Slavnov-Taylor (ST) type identity (2.66) for the gravitational gauge-fixing functions given in Eq.(2.64a)-(2.64b). Then, we reexpress the gauge-fixing function (2.64a) in the following matrix notation:

$$\mathcal{F}_n^\mu = -\frac{i\mathbb{M}_n}{\sqrt{2}} \mathbb{F}_n^\mu, \quad \mathbb{F}_n^\mu = \mathbf{K}_n^T \mathbf{H}_n, \quad (\text{B.21a})$$

$$\mathbf{K}_n = \left( \frac{1}{\sqrt{2\mathbb{M}_n}} [2k_\beta \eta_\alpha^\mu - k^\mu (2 - \xi_n^{-1}) \eta_{\alpha\beta}], i\xi_n \right)^T, \quad \mathbf{H}_n = (h_n^{\alpha\beta}, \mathcal{V}_n^\mu)^T, \quad (\text{B.21b})$$

where we have also assumed the external momentum  $k^\mu$  to be incoming. Therefore, the ST-type identity associated with the gauge function  $\mathbb{F}_n^\mu$  is given by

$$\langle 0 | T \mathbb{F}_n^\mu \mathbf{H}_m^T | 0 \rangle(k) = -\frac{i\xi_n}{\sqrt{2\mathbb{M}_n}} \langle 0 | T \widehat{\mathbf{s}} \mathbf{H}_m^T \bar{c}_n^\mu | 0 \rangle(k). \quad (\text{B.22})$$

By taking Eq.(B.21), we can further rewrite Eq.(B.22) as follows:

$$\mathbf{K}_n^T \mathcal{D}_{nm}(k) = -\mathbf{X}_{nm}^T(k), \quad (\text{B.23})$$

where we have defined the following notations:

$$\begin{aligned} \mathcal{D}_{nm}(k) &= \langle 0 | T \mathbf{H}_n \mathbf{H}_m^T | 0 \rangle(k), \quad \mathbf{X}_{nm}(k) = \underline{\mathbf{X}}_{mj}(k) \mathcal{S}_{jn}(k), \\ \underline{\mathbf{X}}_{mj}(k) &= \frac{i\bar{\eta}^{\lambda\mu}}{\sqrt{2}\mathbb{M}_j} \left( \frac{\langle 0 | T \widehat{\mathbf{h}}_m^{\sigma\rho} | \bar{c}_{j,\lambda} \rangle(k)}{\langle 0 | T \widehat{\mathbf{v}}_m^\nu | \bar{c}_{j,\lambda} \rangle(k)} \right) = \frac{1}{\sqrt{2}\mathbb{M}_j} \begin{pmatrix} (k^\sigma \bar{\eta}^{\rho\mu} + k^\rho \bar{\eta}^{\sigma\mu}) [\delta_{mj} + \Delta_{mj}^{(1)}(k^2)] \\ i\sqrt{2}M_m \bar{\eta}^{\mu\nu} [\delta_{mj} + \widetilde{\Delta}_{mj}^{(2)}(k^2)] \end{pmatrix}, \\ \mathcal{S}_{jn}(k) \bar{\eta}^{\lambda\mu} &= \langle 0 | T c_j^\lambda \bar{c}_n^\mu | 0 \rangle(k), \quad \bar{\eta}^{\lambda\mu} = \eta^{\lambda\mu} - \frac{k^\lambda k^\mu (1 - \xi_n)}{k^2 + \xi_n^2 \mathbb{M}_n^2}, \end{aligned} \quad (\text{B.24})$$

which  $\Delta_{mj}^{(1)}(k^2)$  and  $\widetilde{\Delta}_{mj}^{(2)}(k^2)$  are the loop-level quantities. The above Eqs.(B.23)-(B.24) just give Eqs.(2.69)-(2.70) in the text. Moreover, according to the BRST transformations (B.4a)-(B.4b), each loop-level quantity incorporates distinct contributions from the non-linear terms of the BRST transformations (B.4a)-(B.4b):

$$\Delta_{mj}^{(1)}(k^2) = [\Delta_{h,h}^{(1)}(k^2) + \Delta_{h,\nu}^{(1)}(k^2) + \Delta_{h,\phi}^{(1)}(k^2)]_{mj}, \quad (\text{B.25a})$$

$$\widetilde{\Delta}_{mj}^{(2)}(k^2) = [\widetilde{\Delta}_{\nu,h}^{(2)}(k^2) + \widetilde{\Delta}_{\nu,\nu}^{(2)}(k^2) + \widetilde{\Delta}_{\nu,\phi}^{(2)}(k^2)]_{mj}. \quad (\text{B.25b})$$

Next, we consider the identity (2.66) with an external state given by the combination of gauge-fixing functions as in Eq.(B.21). With this, we can directly amputate the external state as follows:

$$\begin{aligned} 0 &= \langle 0 | \mathbb{F}_n^\mu(k) \cdots \Phi | 0 \rangle = \mathbf{K}_n^T \mathcal{D}_{nm}(k) \langle 0 | \underline{\mathbf{H}}_m(k) \cdots \Phi | 0 \rangle \\ &= -\mathbf{X}_{nm}^T(k) \mathcal{M}[\underline{\mathbf{H}}_m(k), \cdots, \Phi], \end{aligned} \quad (\text{B.26})$$

which leads to the following identity:

$$\mathcal{M}[\mathbb{F}_n^\mu(k), \cdots, \Phi] = 0, \quad (\text{B.27})$$

where we have used the BRST identity (B.23) and  $\mathcal{M}[\cdots]$  denotes the amputated scattering amplitude. In Eq.(B.27), we derive the following amputated external state  $\mathbb{F}_n^\mu$ :

$$\mathbb{F}_n^\mu(k) = \frac{\sqrt{2}}{\mathbb{M}_n} k_\nu h_n^{\mu\nu} - \hat{C}_{\text{mod}}^{nm} \eta^{\mu\nu} \mathcal{V}_{m,\nu}, \quad (\text{B.28})$$

with the loop-induced modification factor given by

$$\hat{C}_{\text{mod}}^{nm}(k^2) = -\frac{i\mathbb{M}_m}{\mathbb{M}_n} \left[ \frac{\mathbf{1} + \widetilde{\Delta}^{(2)}(k^2)}{\mathbf{1} + \Delta^{(1)}(k^2)} \right]_{mn} = -i\delta_{nm} + O(\text{loop}). \quad (\text{B.29})$$

The matrix form presented above represents loop-level quantities  $(\Delta^{(1)})_{jj'} = \Delta_{jj'}^{(1)}$  and  $(\widetilde{\Delta}^{(2)})_{jj'} = \widetilde{\Delta}_{jj'}^{(2)}$ . We provide the above loop-level formulation for completeness, since our

present focus in the main text is to analyze the KK scattering amplitudes at tree level. The above Eqs.(B.26)-(B.29) just give the Eqs.(2.71)-(2.74) in the main text.

By utilizing a transverse polarization vector  $\epsilon_{\pm}^{\mu}$ , we contract it with  $\mathbb{F}_n^{\mu}(k)$  in Eq.(B.28), yielding the formula:

$$\mathbb{F}_n(k) = h_n^{\pm 1} - \Theta_n, \quad \Theta_n = \hat{C}_{\text{mod}}^{nm} \mathcal{V}_m^{\pm 1} + v_n^{\pm 1}, \quad (\text{B.30})$$

where  $v_{\pm 1}^{\mu\nu} = O(\mathbb{M}_n/E_n)$ . Then, we can generalize Eq.(B.27) to incorporate  $N$  externally-amputated gauge-fixing functions:

$$\mathcal{M}[\mathbb{F}_{n_1}(k_1), \dots, \mathbb{F}_{n_N}(k_N), \Phi] = 0. \quad (\text{B.31})$$

Using this identity, we can further derive the gravitational equivalence theorem (GRET) identity which connects the  $h_n^{\pm 1}$  amplitude to the amplitude of  $\Theta_n$ :

$$\mathcal{M}[h_{n_1}^{\pm 1}(k_1), \dots, h_{n_N}^{\pm 1}(k_N), \Phi] = \mathcal{M}[\Theta_{n_1}(k_1), \dots, \Theta_{n_N}(k_N), \Phi]. \quad (\text{B.32})$$

We can prove the above identity by computing the amplitude on its right-hand side. For this, we expand  $\Theta_n$  in terms of  $\Theta_n = h_n^{\pm 1} - \mathbb{F}_n$  for each external state of Eq.(B.32) and thus deduce the left-hand side of Eq.(B.32) after applying the identity Eq.(B.31) to eliminate each external state of  $\mathbb{F}_n$ . The above Eq.(B.32) just reproduce the GRET identity (2.83) in the main text.

## C Feynman Rules for Warped KK Gauge and Gravity Theories

In this Appendix, we present the relevant Feynman rules for the warped KK gauge and gravity theories under the 5d compactification of  $S^1/\mathbb{Z}_2$ , which are needed for explicitly computing the 3-point and 4-point KK scattering amplitudes in the main text.


### C.1 Feynman Rules for Warped KK Gauge Theory

In this sub-Appendix, we present the relevant Feynman rules for the warped KK gauge theory. The propagators for KK gauge boson and the KK Goldstone boson take the following forms in the  $R_{\xi}$  gauge:

$$\mathcal{D}_{nm}^{\mu\nu}(p) = -\frac{i\delta_{nm}}{p^2 + \mathbb{M}_n^2} \left[ \eta^{\mu\nu} + (\xi_n - 1) \frac{p^{\mu} p^{\nu}}{p^2 + \xi_n \mathbb{M}_n^2} \right], \quad (\text{C.1a})$$

$$\mathcal{D}_{nm}(p) = -\frac{i\delta_{nm}}{p^2 + \xi_n \mathbb{M}_n^2}. \quad (\text{C.1b})$$

The trilinear vertices and quartic vertices for KK gauge bosons (KK Goldstone bosons) are derived as follows:



$$= -g a_{nml} f^{abc} \left[ \eta^{\mu\nu} (p_1 - p_2)^{\alpha} + \eta^{\nu\alpha} (p_2 - p_3)^{\mu} + \eta^{\alpha\mu} (p_3 - p_1)^{\nu} \right], \quad (\text{C.2a})$$

$$= -g\tilde{a}_{nm\ell}f^{abc}(p_1-p_2)^\mu, \quad (\text{C.2b})$$

$$= ig a_{nm\ell} f^{abc} \eta^{\mu\nu} (M_n^2 - M_m^2) M_\ell^{-1}, \quad (\text{C.2c})$$

$$= ig^2 a_{nm\ell q} [f^{abe} f^{cde} (\eta^{\mu\alpha} \eta^{\nu\beta} - \eta^{\mu\beta} \eta^{\nu\alpha}) + f^{ace} f^{dbe} (\eta^{\mu\beta} \eta^{\nu\alpha} - \eta^{\mu\nu} \eta^{\alpha\beta}) + f^{ade} f^{bce} (\eta^{\mu\nu} \eta^{\alpha\beta} - \eta^{\mu\alpha} \eta^{\nu\beta})], \quad (\text{C.2d})$$

where the effective cubic and quartic KK coupling coefficients ( $a_{nm\ell}$ ,  $\tilde{a}_{nm\ell}$ ,  $a_{nm\ell q}$ ,  $\tilde{a}_{nm\ell q}$ ) are defined as follows:

$$a_{nm\ell} = \frac{1}{L} \int_0^L dz e^{A(z)} f_n(z) f_m(z) f_\ell(z), \quad (\text{C.3a})$$

$$\tilde{a}_{nm\ell} = \frac{1}{L} \int_0^L dz e^{A(z)} \tilde{f}_n(z) \tilde{f}_m(z) f_\ell(z), \quad (\text{C.3b})$$

$$a_{nm\ell q} = \frac{1}{L} \int_0^L dz e^{A(z)} f_n(z) f_m(z) f_\ell(z) f_q(z), \quad (\text{C.3c})$$

$$\tilde{a}_{nm\ell q} = \frac{1}{L} \int_0^L dz e^{A(z)} \tilde{f}_n(z) \tilde{f}_m(z) \tilde{f}_\ell(z) \tilde{f}_q(z). \quad (\text{C.3d})$$

These coupling coefficients will be used for the double-copy construction in Sections 4.1-4.2 and for deriving the gravitational sum rules in Appendix D. In the above Feynman rules (C.2), the coupling  $g$  is the 4d gauge coupling and is connected to the 5d gauge coupling  $\hat{g}_5$  via  $g = \hat{g}_5/\sqrt{\bar{L}}$ . For the 5d physical coordinate  $y$  (instead of the 5d conformal coordinate  $z$ ), the 5d length is  $\bar{L} = \pi r_c$  and the corresponding 4d gauge coupling is  $\bar{g} = \hat{g}_5/\sqrt{\bar{L}}$ , which is related to the gauge coupling  $g$  via  $\bar{g} = g\sqrt{\bar{L}/L}$ . Thus, the cubic KK gauge coupling coefficient  $\bar{a}_{nm\ell}$  (defined in 5d physical coordinate) is connected to the above cubic KK gauge coupling coefficient  $a_{nm\ell}$  (defined in 5d conformal coordinate) by  $\bar{a}_{nm\ell} = a_{nm\ell} \sqrt{\bar{L}/L}$ . Similarly, for the quartic KK gauge coupling coefficient, we have the relation,  $\bar{a}_{nm\ell q} = (\bar{L}/L) a_{nm\ell q}$ .

## C.2 Feynman Rules for Warped KK Gravity Theory


The propagators for KK gravitons and gravitational KK Goldstone bosons in a general  $R_\xi$  gauge (2.43)-(2.44) are given by the following [15]:

$$\begin{aligned} \mathcal{D}_{nm}^{\mu\nu\alpha\beta}(p) = & -\frac{i\delta_{nm}}{2} \left\{ \frac{\eta^{\mu\alpha}\eta^{\nu\beta} + \eta^{\mu\beta}\eta^{\nu\alpha} - \eta^{\mu\nu}\eta^{\alpha\beta}}{p^2 + \mathbb{M}_n^2} + \frac{1}{3} \left[ \frac{1}{p^2 + \mathbb{M}_n^2} - \frac{1}{p^2 + (3\xi_n - 2)\mathbb{M}_n^2} \right] \right. \\ & \times \left( \eta^{\mu\nu} - \frac{2p^\mu p^\nu}{\mathbb{M}_n^2} \right) \left( \eta^{\alpha\beta} - \frac{2p^\alpha p^\beta}{\mathbb{M}_n^2} \right) + \frac{1}{M_n^2} \left[ \frac{1}{p^2 + \mathbb{M}_n^2} - \frac{1}{p^2 + \xi_n \mathbb{M}_n^2} \right] \left( \eta^{\mu\alpha} p^\nu p^\beta + \eta^{\mu\beta} p^\nu p^\alpha \right. \\ & \left. \left. + \eta^{\nu\alpha} p^\mu p^\beta + \eta^{\nu\beta} p^\mu p^\alpha \right) + \frac{4p^\mu p^\nu p^\alpha p^\beta}{\xi_n M_n^4} \left( \frac{1}{p^2 + \xi_n^2 \mathbb{M}_n^2} - \frac{1}{p^2 + \xi_n \mathbb{M}_n^2} \right) \right\}, \end{aligned} \quad (\text{C.4a})$$


$$\mathcal{D}_{nm}^{\mu\nu}(p) = -\frac{i\delta_{nm}}{p^2 + \xi_n \mathbb{M}_n^2} \left[ \eta^{\mu\nu} - \frac{p^\mu p^\nu (1 - \xi_n)}{p^2 + \xi_n^2 \mathbb{M}_n^2} \right], \quad (\text{C.4b})$$

$$\mathcal{D}_{nm}(p) = -\frac{i\delta_{nm}}{p^2 + (3\xi_n - 2)\mathbb{M}_n^2}. \quad (\text{C.4c})$$


The relevant trilinear gravitational KK vertices are given by




$$= i\kappa\alpha_{nm\ell}\Gamma_{nm\ell}^{\mu\nu\rho\sigma\alpha\beta}(p_1, p_2, p_3), \quad (\text{C.5a})$$



$$= \frac{i\kappa}{2}\tilde{\alpha}_{nm\ell}\Gamma_{nm\ell}^{\alpha\beta\mu\nu}(p_1, p_2), \quad (\text{C.5b})$$



$$= \frac{i\kappa}{2}\tilde{\beta}_{nm\ell}\left\{2(p_1^\mu p_1^\nu + p_2^\mu p_2^\nu) + p_1^\mu p_2^\nu + p_1^\nu p_2^\mu - [2(p_1^2 + p_2^2) + 3p_1 \cdot p_2 + \mathbb{M}_n \mathbb{M}_\ell + \mathbb{M}_m \mathbb{M}_\ell]\eta^{\mu\nu}\right\}, \quad (\text{C.5c})$$



$$= \frac{i\kappa}{\sqrt{6}}\tilde{\rho}_{nm\ell}\left\{-2(p_1^\mu p_1^\nu + p_2^\mu p_2^\nu) - 3p_2^\mu p_1^\nu + [2(p_1^2 + p_2^2) + 3p_1 \cdot p_2]\eta^{\mu\nu}\right\}, \quad (\text{C.5d})$$

$$\begin{aligned}
&= i2\sqrt{\frac{2}{3}} \kappa \tilde{\omega}_{nml} (p_1 \cdot p_2 + p_1 \cdot p_3 + p_2 \cdot p_3 - \mathbb{M}_n \mathbb{M}_m \\
&\quad - \mathbb{M}_n \mathbb{M}_\ell - \mathbb{M}_m \mathbb{M}_\ell). \tag{C.5e}
\end{aligned}$$

In the above Feynman rules (C.5), the coupling  $\kappa$  is the 4d gravitational coupling and is connected to the 5d gravitational coupling  $\hat{\kappa}_5$  via  $\kappa = \hat{\kappa}_5 / \sqrt{L}$ . For the 5d physical coordinate  $y$  (instead of the 5d conformal coordinate  $z$ ), the 5d length is  $\bar{L} = \pi r_c$  and the corresponding 4d gravitational coupling is  $\bar{\kappa} = \hat{\kappa}_5 / \sqrt{\bar{L}}$ , which is related to the gravitational coupling  $\kappa$  via  $\bar{\kappa} = \kappa \sqrt{\bar{L}/L}$ . Thus, the cubic KK gravitational coupling coefficient  $\bar{\alpha}_{nml}$  (defined in 5d physical coordinate) is connected to the above cubic KK gravitational coupling coefficient  $\alpha_{nml}$  (defined in 5d conformal coordinate) by  $\bar{\alpha}_{nml} = \alpha_{nml} \sqrt{\bar{L}/L}$ .

In the above Eq.(C.5a), the trilinear gravitational vertex function  $\Gamma_{nml}^{\mu\nu\rho\sigma\alpha\beta}$  is defined as follows:

$$\Gamma_{nml}^{\mu\nu\rho\sigma\alpha\beta}(p_1, p_2, p_3) = \frac{1}{8} [(\mathbb{M}_n^2 F_1 - p_1^2 F_2 + F_3) + (1 \leftrightarrow 2) + (1 \leftrightarrow 3)], \tag{C.6}$$

with the functions ( $F_1, F_2, F_3$ ) given by

$$\begin{aligned}
F_1 &= -\eta^{\alpha\beta} \eta^{\mu\sigma} \eta^{\nu\rho} - \eta^{\alpha\beta} \eta^{\mu\rho} \eta^{\nu\sigma} + 2\eta^{\alpha\beta} \eta^{\mu\nu} \eta^{\rho\sigma} + \eta^{\alpha\mu} \eta^{\beta\sigma} \eta^{\nu\rho} + \eta^{\alpha\mu} \eta^{\beta\rho} \eta^{\nu\sigma} - \eta^{\alpha\mu} \eta^{\beta\nu} \eta^{\rho\sigma} \\
&\quad + \eta^{\alpha\sigma} \eta^{\beta\mu} \eta^{\nu\rho} + \eta^{\alpha\rho} \eta^{\beta\mu} \eta^{\nu\sigma} + \eta^{\alpha\nu} \eta^{\beta\sigma} \eta^{\mu\rho} + \eta^{\alpha\nu} \eta^{\beta\rho} \eta^{\mu\sigma} - \eta^{\alpha\nu} \eta^{\beta\mu} \eta^{\rho\sigma} + \eta^{\alpha\sigma} \eta^{\beta\nu} \eta^{\mu\rho} \\
&\quad + \eta^{\alpha\rho} \eta^{\beta\nu} \eta^{\mu\sigma} - 3\eta^{\alpha\rho} \eta^{\beta\sigma} \eta^{\mu\nu} - 3\eta^{\alpha\sigma} \eta^{\beta\rho} \eta^{\mu\nu}, \\
F_2 &= 3\eta^{\alpha\beta} \eta^{\mu\sigma} \eta^{\nu\rho} + 3\eta^{\alpha\beta} \eta^{\mu\rho} \eta^{\nu\sigma} - 4\eta^{\alpha\beta} \eta^{\mu\nu} \eta^{\rho\sigma} - 2\eta^{\alpha\mu} \eta^{\beta\sigma} \eta^{\nu\rho} - 2\eta^{\alpha\mu} \eta^{\beta\rho} \eta^{\nu\sigma} + 3\eta^{\alpha\mu} \eta^{\beta\nu} \eta^{\rho\sigma} \\
&\quad + 3\eta^{\alpha\nu} \eta^{\beta\mu} \eta^{\rho\sigma} - 2\eta^{\alpha\nu} \eta^{\beta\sigma} \eta^{\mu\rho} - 2\eta^{\alpha\nu} \eta^{\beta\rho} \eta^{\mu\sigma} + 4\eta^{\alpha\sigma} \eta^{\beta\rho} \eta^{\mu\nu} + 4\eta^{\alpha\rho} \eta^{\beta\sigma} \eta^{\mu\nu} - 2\eta^{\alpha\rho} \eta^{\beta\nu} \eta^{\mu\sigma} \\
&\quad - 2\eta^{\alpha\rho} \eta^{\beta\mu} \eta^{\nu\sigma} - 2\eta^{\alpha\sigma} \eta^{\beta\nu} \eta^{\mu\rho} - 2\eta^{\alpha\sigma} \eta^{\beta\mu} \eta^{\nu\rho}, \\
F_3 &= 2\eta^{\mu\nu} \eta^{\rho\sigma} p_1^\alpha p_2^\beta + \eta^{\mu\nu} \eta^{\rho\sigma} p_1^\alpha p_3^\beta + \eta^{\nu\rho} \eta^{\beta\sigma} p_1^\alpha p_2^\mu + \eta^{\beta\rho} \eta^{\nu\sigma} p_1^\alpha p_2^\mu + \eta^{\nu\rho} \eta^{\beta\sigma} p_1^\alpha p_3^\mu + \eta^{\beta\rho} \eta^{\nu\sigma} p_1^\alpha p_3^\mu \\
&\quad + \eta^{\mu\rho} \eta^{\beta\sigma} p_1^\alpha p_2^\nu + \eta^{\beta\rho} \eta^{\mu\sigma} p_1^\alpha p_2^\nu + \eta^{\mu\rho} \eta^{\beta\sigma} p_1^\alpha p_3^\nu + \eta^{\beta\rho} \eta^{\mu\sigma} p_1^\alpha p_3^\nu - \eta^{\beta\sigma} \eta^{\mu\nu} p_1^\alpha p_2^\rho - \eta^{\beta\sigma} \eta^{\mu\nu} p_1^\alpha p_3^\rho \\
&\quad + \eta^{\beta\nu} \eta^{\mu\sigma} p_1^\alpha p_2^\rho + \eta^{\beta\mu} \eta^{\nu\sigma} p_1^\alpha p_2^\rho - \eta^{\mu\sigma} \eta^{\nu\rho} p_1^\alpha p_2^\beta - \eta^{\mu\sigma} \eta^{\nu\rho} p_1^\alpha p_3^\beta - \eta^{\beta\rho} \eta^{\mu\nu} p_1^\alpha p_2^\sigma - \eta^{\beta\rho} \eta^{\mu\nu} p_1^\alpha p_3^\sigma \\
&\quad + \eta^{\beta\nu} \eta^{\mu\rho} p_1^\alpha p_2^\sigma + \eta^{\beta\mu} \eta^{\nu\rho} p_1^\alpha p_2^\sigma - \eta^{\mu\rho} \eta^{\nu\sigma} p_1^\alpha p_2^\beta - \eta^{\mu\rho} \eta^{\nu\sigma} p_1^\alpha p_3^\beta - \eta^{\beta\nu} \eta^{\rho\sigma} p_1^\alpha p_2^\mu - \eta^{\beta\nu} \eta^{\rho\sigma} p_1^\alpha p_3^\mu \\
&\quad - \eta^{\beta\mu} \eta^{\rho\sigma} p_1^\alpha p_2^\nu - \eta^{\beta\mu} \eta^{\rho\sigma} p_1^\alpha p_3^\nu + 2\eta^{\mu\nu} \eta^{\rho\sigma} p_1^\beta p_2^\alpha + \eta^{\mu\nu} \eta^{\rho\sigma} p_1^\beta p_3^\alpha + \eta^{\nu\rho} \eta^{\alpha\sigma} p_1^\beta p_2^\mu + \eta^{\alpha\rho} \eta^{\nu\sigma} p_1^\beta p_2^\mu \\
&\quad + \eta^{\nu\rho} \eta^{\alpha\sigma} p_1^\beta p_3^\mu + \eta^{\alpha\rho} \eta^{\nu\sigma} p_1^\beta p_3^\mu + \eta^{\mu\rho} \eta^{\alpha\sigma} p_1^\beta p_2^\nu + \eta^{\alpha\rho} \eta^{\mu\sigma} p_1^\beta p_2^\nu + \eta^{\mu\rho} \eta^{\alpha\sigma} p_1^\beta p_3^\nu + \eta^{\alpha\rho} \eta^{\mu\sigma} p_1^\beta p_3^\nu \\
&\quad - \eta^{\alpha\sigma} \eta^{\mu\nu} p_1^\beta p_2^\rho - \eta^{\alpha\sigma} \eta^{\mu\nu} p_1^\beta p_3^\rho + \eta^{\alpha\nu} \eta^{\mu\sigma} p_1^\beta p_2^\rho + \eta^{\alpha\mu} \eta^{\nu\sigma} p_1^\beta p_2^\rho - \eta^{\mu\sigma} \eta^{\nu\rho} p_1^\beta p_2^\alpha - \eta^{\alpha\rho} \eta^{\mu\nu} p_1^\beta p_2^\sigma \\
&\quad - \eta^{\alpha\rho} \eta^{\mu\nu} p_1^\beta p_3^\sigma + \eta^{\alpha\nu} \eta^{\mu\rho} p_1^\beta p_2^\sigma + \eta^{\alpha\mu} \eta^{\nu\rho} p_1^\beta p_2^\sigma - \eta^{\mu\rho} \eta^{\nu\sigma} p_1^\beta p_2^\alpha - \eta^{\mu\rho} \eta^{\nu\sigma} p_1^\beta p_3^\alpha - \eta^{\alpha\nu} \eta^{\rho\sigma} p_1^\beta p_2^\mu \\
&\quad - \eta^{\alpha\nu} \eta^{\rho\sigma} p_1^\beta p_3^\mu - \eta^{\alpha\mu} \eta^{\rho\sigma} p_1^\beta p_2^\nu - \eta^{\alpha\mu} \eta^{\rho\sigma} p_1^\beta p_3^\nu - \eta^{\nu\rho} \eta^{\mu\sigma} p_1^\beta p_2^\alpha - \eta^{\alpha\sigma} \eta^{\beta\rho} p_1^\beta p_2^\nu - \eta^{\alpha\rho} \eta^{\beta\sigma} p_1^\beta p_2^\nu \\
&\quad + \eta^{\alpha\beta} \eta^{\rho\sigma} p_1^\mu p_2^\nu - \eta^{\alpha\sigma} \eta^{\beta\rho} p_1^\mu p_2^\nu - \eta^{\alpha\rho} \eta^{\beta\sigma} p_1^\mu p_2^\nu + \eta^{\alpha\beta} \eta^{\rho\sigma} p_1^\mu p_3^\nu - \eta^{\alpha\beta} \eta^{\nu\sigma} p_1^\mu p_2^\rho - \eta^{\alpha\beta} \eta^{\nu\rho} p_1^\mu p_2^\sigma \\
&\quad - \eta^{\alpha\nu} \eta^{\rho\sigma} p_1^\mu p_2^\beta - \eta^{\beta\nu} \eta^{\rho\sigma} p_1^\mu p_2^\alpha - \eta^{\alpha\sigma} \eta^{\beta\rho} p_1^\nu p_2^\mu - \eta^{\alpha\rho} \eta^{\beta\sigma} p_1^\nu p_2^\mu + \eta^{\alpha\beta} \eta^{\rho\sigma} p_1^\nu p_2^\mu - \eta^{\alpha\sigma} \eta^{\beta\rho} p_1^\nu p_2^\mu \\
&\quad - \eta^{\alpha\rho} \eta^{\beta\sigma} p_1^\nu p_2^\mu + \eta^{\alpha\beta} \eta^{\rho\sigma} p_1^\nu p_2^\mu - \eta^{\alpha\beta} \eta^{\mu\sigma} p_1^\nu p_2^\rho - \eta^{\alpha\beta} \eta^{\mu\rho} p_1^\nu p_2^\sigma - \eta^{\alpha\mu} \eta^{\rho\sigma} p_1^\nu p_2^\beta - \eta^{\beta\mu} \eta^{\rho\sigma} p_1^\nu p_2^\alpha \\
&\quad + \eta^{\beta\nu} \eta^{\mu\sigma} p_1^\rho p_2^\alpha + \eta^{\beta\mu} \eta^{\nu\sigma} p_1^\rho p_2^\alpha + \eta^{\alpha\nu} \eta^{\mu\sigma} p_1^\rho p_2^\beta + \eta^{\alpha\mu} \eta^{\nu\sigma} p_1^\rho p_2^\beta + \eta^{\beta\nu} \eta^{\alpha\sigma} p_1^\rho p_2^\mu + \eta^{\alpha\nu} \eta^{\beta\sigma} p_1^\rho p_2^\mu
\end{aligned}$$



$$\begin{aligned}
& + \eta^{\beta\nu} \eta^{\alpha\sigma} p_1^\rho p_3^\mu + \eta^{\alpha\nu} \eta^{\beta\sigma} p_1^\rho p_3^\mu + \eta^{\beta\mu} \eta^{\alpha\sigma} p_1^\rho p_2^\nu + \eta^{\alpha\mu} \eta^{\beta\sigma} p_1^\rho p_2^\nu - \eta^{\alpha\beta} \eta^{\mu\sigma} p_1^\rho p_2^\nu + \eta^{\beta\mu} \eta^{\alpha\sigma} p_1^\rho p_3^\nu \\
& + \eta^{\alpha\mu} \eta^{\beta\sigma} p_1^\rho p_3^\nu - \eta^{\alpha\beta} \eta^{\mu\sigma} p_1^\rho p_3^\nu - \eta^{\alpha\sigma} \eta^{\mu\nu} p_1^\rho p_2^\beta - \eta^{\alpha\sigma} \eta^{\mu\nu} p_1^\rho p_3^\beta - \eta^{\beta\sigma} \eta^{\mu\nu} p_1^\rho p_2^\alpha - \eta^{\beta\sigma} \eta^{\mu\nu} p_1^\rho p_3^\alpha \\
& - \eta^{\alpha\nu} \eta^{\beta\mu} p_1^\rho p_2^\sigma - \eta^{\alpha\mu} \eta^{\beta\nu} p_1^\rho p_2^\sigma + \eta^{\alpha\beta} \eta^{\mu\nu} p_1^\rho p_2^\sigma - \eta^{\alpha\nu} \eta^{\beta\mu} p_1^\rho p_3^\sigma - \eta^{\alpha\mu} \eta^{\beta\nu} p_1^\rho p_3^\sigma + 2\eta^{\alpha\beta} \eta^{\mu\nu} p_1^\rho p_3^\sigma \\
& - \eta^{\alpha\beta} \eta^{\nu\sigma} p_1^\rho p_2^\mu - \eta^{\alpha\beta} \eta^{\nu\sigma} p_1^\rho p_3^\mu + \eta^{\beta\nu} \eta^{\mu\rho} p_1^\sigma p_2^\alpha + \eta^{\beta\mu} \eta^{\nu\rho} p_1^\sigma p_2^\alpha + \eta^{\alpha\rho} \eta^{\mu\rho} p_1^\sigma p_2^\beta + \eta^{\alpha\mu} \eta^{\nu\rho} p_1^\sigma p_2^\beta \\
& + \eta^{\beta\nu} \eta^{\alpha\rho} p_1^\sigma p_2^\mu + \eta^{\alpha\nu} \eta^{\beta\rho} p_1^\sigma p_2^\mu + \eta^{\beta\nu} \eta^{\alpha\rho} p_1^\sigma p_3^\mu + \eta^{\alpha\nu} \eta^{\beta\rho} p_1^\sigma p_3^\mu + \eta^{\beta\mu} \eta^{\alpha\rho} p_1^\sigma p_2^\nu + \eta^{\alpha\mu} \eta^{\beta\rho} p_1^\sigma p_2^\nu \\
& - \eta^{\alpha\beta} \eta^{\mu\rho} p_1^\sigma p_2^\nu + \eta^{\beta\mu} \eta^{\alpha\rho} p_1^\sigma p_2^\nu + \eta^{\alpha\mu} \eta^{\beta\rho} p_1^\sigma p_3^\nu - \eta^{\alpha\beta} \eta^{\mu\rho} p_1^\sigma p_3^\nu - \eta^{\alpha\rho} \eta^{\mu\nu} p_1^\sigma p_2^\beta - \eta^{\alpha\rho} \eta^{\mu\nu} p_1^\sigma p_3^\beta \\
& - \eta^{\beta\rho} \eta^{\mu\nu} p_1^\sigma p_2^\alpha - \eta^{\beta\rho} \eta^{\mu\nu} p_1^\sigma p_3^\alpha - \eta^{\alpha\nu} \eta^{\beta\mu} p_1^\sigma p_2^\rho - \eta^{\alpha\mu} \eta^{\beta\nu} p_1^\sigma p_2^\rho + \eta^{\alpha\beta} \eta^{\mu\nu} p_1^\sigma p_2^\rho - \eta^{\alpha\nu} \eta^{\beta\mu} p_1^\sigma p_3^\rho \\
& - \eta^{\alpha\mu} \eta^{\beta\nu} p_1^\sigma p_3^\rho + 2\eta^{\alpha\beta} \eta^{\mu\nu} p_1^\sigma p_3^\rho - \eta^{\alpha\beta} \eta^{\nu\rho} p_1^\sigma p_2^\mu - \eta^{\alpha\beta} \eta^{\nu\rho} p_1^\sigma p_3^\mu. \tag{C.7}
\end{aligned}$$

In Eq.(C.5b), the trilinear gravitational vertex function  $\Gamma_{nml}^{\alpha\beta\mu\nu}$  is defined as follows:

$$\begin{aligned}
\Gamma_{nml}^{\alpha\beta\mu\nu}(p_1, p_2) = & -2\eta^{\alpha\beta} p_1^\mu p_1^\nu + 2\eta^{\alpha\beta} \eta^{\mu\nu} p_1^2 + \eta^{\alpha\mu} p_1^\beta p_1^\nu + \eta^{\alpha\nu} p_1^\beta p_1^\mu + \eta^{\beta\mu} p_1^\alpha p_1^\nu - \eta^{\alpha\nu} \eta^{\beta\mu} p_1^2 \\
& + \eta^{\beta\nu} p_1^\alpha p_1^\mu - \eta^{\alpha\mu} \eta^{\beta\nu} p_1^2 - 2\eta^{\mu\nu} p_1^\alpha p_1^\beta - \eta^{\alpha\beta} p_1^\nu p_1^\mu - \eta^{\alpha\beta} p_1^\mu p_1^\nu + 3\eta^{\alpha\beta} \eta^{\mu\nu} (p_1 \cdot p_2) \\
& + \eta^{\alpha\mu} p_1^\nu p_2^\beta + \eta^{\alpha\nu} p_1^\mu p_2^\beta + \eta^{\beta\mu} p_1^\alpha p_2^\nu - \eta^{\alpha\nu} \eta^{\beta\mu} (p_1 \cdot p_2) + \eta^{\beta\nu} p_1^\alpha p_2^\mu - \eta^{\alpha\mu} \eta^{\beta\nu} (p_1 \cdot p_2) \\
& - \eta^{\mu\nu} p_1^\beta p_2^\alpha - 2\eta^{\mu\nu} p_1^\alpha p_2^\beta - 2\eta^{\alpha\beta} p_2^\mu p_2^\nu + 2\eta^{\alpha\beta} \eta^{\mu\nu} p_2^2 + \eta^{\alpha\mu} p_2^\beta p_2^\nu + \eta^{\alpha\nu} p_2^\beta p_2^\mu \\
& + \eta^{\beta\mu} p_2^\alpha p_2^\nu - \eta^{\alpha\nu} \eta^{\beta\mu} p_2^2 + \eta^{\beta\nu} p_2^\alpha p_2^\mu - \eta^{\alpha\mu} \eta^{\beta\nu} p_2^2 - 2\eta^{\mu\nu} p_2^\alpha p_2^\beta. \tag{C.8}
\end{aligned}$$

Moreover, the trilinear coupling coefficients ( $\alpha_{nml}$ ,  $\tilde{\alpha}_{nml}$ ,  $\tilde{\beta}_{nml}$ ,  $\tilde{\rho}_{nml}$ ) in Eq.(C.5) are defined as bellow:

$$\alpha_{nml} = \frac{1}{L} \int_0^L dz e^{3A(z)} \mathbf{u}_n(z) \mathbf{u}_m(z) \mathbf{u}_\ell(z), \tag{C.9a}$$

$$\tilde{\alpha}_{nml} = \frac{1}{L} \int_0^L dz e^{3A(z)} \mathbf{v}_n(z) \mathbf{v}_m(z) \mathbf{u}_\ell(z), \tag{C.9b}$$

$$\tilde{\beta}_{nml} = \frac{1}{L} \int_0^L dz e^{3A(z)} \mathbf{w}_n(z) \mathbf{w}_m(z) \mathbf{u}_\ell(z), \tag{C.9c}$$

$$\tilde{\rho}_{nml} = \frac{1}{L} \int_0^L dz e^{3A(z)} \mathbf{v}_n(z) \mathbf{v}_m(z) \mathbf{w}_\ell(z), \tag{C.9d}$$

$$\tilde{\omega}_{nml} = \frac{1}{L} \int_0^L dz e^{3A(z)} \mathbf{w}_n(z) \mathbf{w}_m(z) \mathbf{w}_\ell(z). \tag{C.9e}$$

In particular, we can derive the trilinear coupling coefficients containing zero-modes,  $\alpha_{000} = \alpha_{nn0} = \tilde{\alpha}_{nn0} = \mathbf{u}_0$  and  $a_{000} = a_{nn0} = \tilde{a}_{nn0} = \mathbf{f}_0$ , based on the normalization condition Eq.(2.51). We summarize these coupling coefficients as follows:

$$\alpha_{000} = \frac{1}{L} \int_0^L dz e^{3A(z)} \mathbf{u}_0^2 \mathbf{u}_0 = \mathbf{u}_0, \tag{C.10a}$$

$$\alpha_{nn0} = \frac{1}{L} \int_0^L dz e^{3A(z)} \mathbf{u}_n^2 \mathbf{u}_0 = \mathbf{u}_0, \tag{C.10b}$$

$$\tilde{\alpha}_{nn0} = \frac{1}{L} \int_0^L dz e^{3A(z)} \mathbf{v}_n^2 \mathbf{u}_0 = \mathbf{u}_0, \tag{C.10c}$$

$$a_{000} = \frac{1}{L} \int_0^L dz e^{A(z)} \mathbf{f}_0^2 \mathbf{f}_0 = \mathbf{f}_0, \tag{C.10d}$$

Cubic and Quartic Couplings (flat 5d)	Values
$\tilde{a}_{nn2n}, \tilde{a}_{nm n\pm m }, \tilde{\alpha}_{nn2n}, \tilde{\rho}_{nn2n}, \tilde{\alpha}_{nm n\pm m }, \tilde{\rho}_{nm n\pm m }$	$-\frac{1}{\sqrt{2}}$
$a_{nm0}, \tilde{a}_{nm0}, \alpha_{nm0}, \tilde{\alpha}_{nm0}, \tilde{\beta}_{nm0}, \tilde{\rho}_{nm0}, \tilde{\omega}_{nm0}$	0
$a_{nmlq}, \tilde{a}_{nmlq}, \alpha_{nmlq}, \tilde{\beta}_{nmlq}$	$\frac{1}{2}$
$a_{nn2n}, a_{nm n\pm m }, \alpha_{nn2n}, \tilde{\beta}_{nn2n}, \tilde{\omega}_{nn2n}, \alpha_{nm n\pm m }, \tilde{\beta}_{nm n\pm m }, \tilde{\omega}_{nm n\pm m }$	$\frac{1}{\sqrt{2}}$
$a_{nn0}, \tilde{a}_{nn0}, a_{nnmm}, \alpha_{nn0}, \tilde{\alpha}_{nn0}, \tilde{\beta}_{nn0}, \tilde{\rho}_{nn0}, \tilde{\omega}_{nn0}, \alpha_{nnmm}, \tilde{\alpha}_{nnmm}, \tilde{\beta}_{nnmm}$	1
$a_{nnnn}, \alpha_{nnnn}, \tilde{\alpha}_{nnnn}, \tilde{\beta}_{nnnn}$	$\frac{3}{2}$

**Table 1:** List of relevant cubic and quartic KK coupling coefficients of the flat 5d gauge and gravity theories under  $S^1/\mathbb{Z}_2$  compactification. The subscripts are the relevant KK indices with  $n \neq m \neq \ell \neq q$  and  $(n, m, \ell, q) \in \mathbb{Z}^+$ .

$$a_{nm0} = \frac{1}{L} \int_0^L dz e^{A(z)} \mathbf{f}_n^2 \mathbf{f}_0 = \mathbf{f}_0, \quad (\text{C.10e})$$

$$\tilde{a}_{nm0} = \frac{1}{L} \int_0^L dz e^{A(z)} \tilde{\mathbf{f}}_n^2 \mathbf{f}_0 = \mathbf{f}_0. \quad (\text{C.10f})$$

In addition, we further define the following quartic coupling coefficients:

$$\alpha_{nmlq} = \frac{1}{L} \int_0^L dz e^{3A(z)} \mathbf{u}_n(z) \mathbf{u}_m(z) \mathbf{u}_\ell(z) \mathbf{u}_q(z), \quad (\text{C.11a})$$

$$\tilde{\beta}_{nmlq} = \frac{1}{L} \int_0^L dz e^{3A(z)} \mathbf{w}_n(z) \mathbf{w}_m(z) \mathbf{w}_\ell(z) \mathbf{w}_q(z), \quad (\text{C.11b})$$

which will be used for the double-copy construction of Section 4.2 and for deriving the gravitational sum rules in Appendix D.

Taking the flat space limit  $k \rightarrow 0$ , we see that all the coupling coefficients will be reduced to the simple trigonometric functions:

$$\begin{aligned} \mathbf{f}_0 &= \mathbf{u}_0 = \mathbf{w}_0 = 1, \\ \mathbf{f}_n(z) &= \mathbf{u}_n(z) = \mathbf{w}_n(z) = \sqrt{2} \cos \frac{n\pi z}{L}, \\ \tilde{\mathbf{f}}_n(z) &= \mathbf{v}_n(z) = \sqrt{2} \sin \frac{n\pi z}{L}, \end{aligned} \quad (\text{C.12})$$

where  $n \in \mathbb{Z}^+$ . Thus, together with the definitions (C.10)-(C.11), we deduce the values of these KK coupling coefficients in the flat 5d limit, which are summarized in Table 1.

Finally, we derive the on-shell 3-point KK gluon scattering amplitudes as follows:

$$\mathcal{T}[\{\epsilon_i\}] = g f^{abc} \mathcal{N}[\{\epsilon_j\}], \quad (\text{C.13a})$$

$$\mathcal{N}[\{\epsilon_j\}] = -i2g a_{nml} [(\epsilon_1 \epsilon_2)(\epsilon_3 p_1) + (\epsilon_2 \epsilon_3)(\epsilon_1 p_2) + (\epsilon_3 \epsilon_1)(\epsilon_2 p_3)], \quad (\text{C.13b})$$

which reproduces the formula (4.1) in the main text.

Then, we can express the 3-point KK graviton amplitude into following form:

$$\begin{aligned}\mathcal{M}[h_n^{\sigma_1} h_m^{\sigma_2} h_\ell^{\sigma_3}] &= \sum_{\lambda_j, \lambda'_j} \left( \prod_j C_{\lambda_j, \lambda'_j}^{\sigma_j} \right) \mathcal{M}[\{e_i\}^{\lambda_j}, \{\epsilon_i\}^{\lambda'_j}] \\ &\equiv \frac{\kappa \alpha_{nm\ell}}{4} \sum_{\lambda_j, \lambda'_j} \left( \prod_j C_{\lambda_j, \lambda'_j}^{\sigma_j} \right) \bar{\mathcal{M}}[\{e_i\}^{\lambda_j}, \{\epsilon_i\}^{\lambda'_j}],\end{aligned}\quad (\text{C.14})$$

where the helicity index  $\sigma_j = \{\pm 2, \pm 1, 0\} \equiv \{\pm 2, \pm 1, L\}$  labels the 5 helicity states of each external massive KK graviton. The polarization tensors (including the coefficients  $C_{\lambda_j, \lambda'_j}^{\sigma_j}$ ) of the external KK graviton states are defined as in Eq.(4.7) of the main text. Then, we explicitly compute the on-shell 3-point KK graviton scattering amplitudes as follows:

$$\begin{aligned}\bar{\mathcal{M}}[\{e_i\}, \{\epsilon_i\}] &= \kappa \alpha_{nm\ell} \left[ e_{1,\mu} \epsilon_{1,\nu} e_{2,\rho} \epsilon_{2,\sigma} e_{3,\alpha} \epsilon_{3,\beta} \Gamma_{nm\ell}^{\mu\nu\rho\sigma\alpha\beta}(p_1, p_2, p_3) \right] \\ &= \frac{\kappa \alpha_{nm\ell}}{4} \left[ (e_1 \epsilon_3)(e_2 p_3)(e_3 p_2)(\epsilon_1 \epsilon_2) + (e_1 \epsilon_3)(e_2 p_3)(p_2 \epsilon_3)(\epsilon_1 \epsilon_2) + (e_1 p_3)(e_2 p_3)(e_3 \epsilon_2)(\epsilon_1 \epsilon_3) \right. \\ &\quad + (e_1 \epsilon_2)(e_2 p_3)(e_3 p_2)(\epsilon_1 \epsilon_3) + (e_1 p_3)(e_2 p_3)(e_3 \epsilon_1)(\epsilon_2 \epsilon_3) + (e_1 \epsilon_3)(e_2 p_3)(p_3 \epsilon_1)(\epsilon_2 \epsilon_3) \\ &\quad + (e_1 \epsilon_2)(e_2 p_3)(e_3 \epsilon_1)(p_2 \epsilon_3) - 2(e_1 \epsilon_2)(e_2 \epsilon_1)(e_3 p_2)(p_2 \epsilon_3) - 2(e_1 \epsilon_2)(e_3 p_2)(p_2 \epsilon_3)(\epsilon_1 \epsilon_2) \\ &\quad + (e_1 \epsilon_3)(e_2 \epsilon_1)(p_3 \epsilon_2)(p_2 \epsilon_3) + (e_1 \epsilon_2)(e_3 \epsilon_1)(p_3 \epsilon_2)(p_2 \epsilon_3) - (e_1 p_3)(e_2 \epsilon_3)(e_3 p_2)(\epsilon_1 \epsilon_2) \\ &\quad - (e_1 p_3)(e_2 \epsilon_1)(e_3 p_2)(\epsilon_2 \epsilon_3) - (e_1 p_3)(e_2 \epsilon_1)(e_3 \epsilon_2)(p_2 \epsilon_3) - (e_1 p_3)(e_2 \epsilon_3)(p_2 \epsilon_3)(\epsilon_1 \epsilon_2) \\ &\quad + (e_1 \epsilon_3)(e_2 p_3)(e_3 \epsilon_2)(p_3 \epsilon_1) - (e_1 \epsilon_2)(e_2 \epsilon_3)(e_3 p_2)(p_3 \epsilon_1) - (e_1 \epsilon_2)(e_3 p_2)(p_3 \epsilon_1)(\epsilon_2 \epsilon_3) \\ &\quad - (e_1 \epsilon_2)(e_2 \epsilon_3)(p_3 \epsilon_1)(p_2 \epsilon_3) - (e_1 \epsilon_2)(e_3 \epsilon_2)(p_3 \epsilon_1)(p_2 \epsilon_3) - 2(e_1 p_3)(e_2 \epsilon_3)(e_3 \epsilon_2)(p_3 \epsilon_1) \\ &\quad - 2(e_1 p_3)(e_2 \epsilon_3)(p_3 \epsilon_1)(\epsilon_2 \epsilon_3) + (e_1 p_3)(e_2 \epsilon_3)(e_3 \epsilon_1)(p_3 \epsilon_2) + (e_1 p_3)(e_2 \epsilon_3)(p_3 \epsilon_2)(\epsilon_1 \epsilon_3) \\ &\quad + (e_1 \epsilon_2)(e_3 p_2)(p_3 \epsilon_2)(\epsilon_1 \epsilon_3) + (e_1 \epsilon_3)(e_2 \epsilon_1)(e_3 p_2)(p_3 \epsilon_2) - 2(e_1 \epsilon_3)(e_2 p_3)(e_3 \epsilon_1)(p_3 \epsilon_2) \\ &\quad \left. - 2(e_1 \epsilon_3)(e_2 p_3)(p_3 \epsilon_2)(\epsilon_1 \epsilon_3) + (e_1 \epsilon_3)(e_2 \epsilon_3)(p_3 \epsilon_1)(p_3 \epsilon_2) + (e_1 \epsilon_3)(e_2 \epsilon_3)(p_3 \epsilon_1)(p_3 \epsilon_2) \right],\end{aligned}\quad (\text{C.15})$$

where for simplicity we have introduced a shorthand notation,  $(ab) \equiv (a \cdot b)$ .

## D Proving 3-Point and 4-Point Warped Identities

In this Appendix, we will prove relevant identities and sum rules for the cubic and quartic coupling coefficients in the warped KK gauge theory and KK gravity theory.

By imposing the following completeness conditions,

$$\delta(z - z') = \sum_{j=0}^{\infty} e^{A(z)} \mathbf{f}_j(z) \mathbf{f}_j(z') = \sum_{j=0}^{\infty} e^{A(z)} \tilde{\mathbf{f}}_j(z) \tilde{\mathbf{f}}_j(z'), \quad (\text{D.1a})$$

$$\delta(z - z') = \sum_{j=0}^{\infty} e^{3A(z)} \mathbf{u}_j(z) \mathbf{u}_j(z') = \sum_{j=0}^{\infty} e^{3A(z)} \mathbf{v}_j(z) \mathbf{v}_j(z') = \sum_{j=0}^{\infty} e^{3A(z)} \mathbf{w}_j(z) \mathbf{w}_j(z'), \quad (\text{D.1b})$$

we can derive the sum rules between the cubic and quartic KK gauge/gravity coupling coefficients respectively:

$$\sum_{j=0}^{\infty} a_{nmj} a_{\ell qj} = a_{nm\ell q}, \quad \sum_{j=0}^{\infty} \alpha_{nmj} \alpha_{\ell qj} = \alpha_{nm\ell q}, \quad (\text{D.2a})$$

$$\sum_{j=0}^{\infty} \tilde{a}_{nmj} \tilde{a}_{\ell qj} = \tilde{a}_{nmlq}, \quad \sum_{j=0}^{\infty} \tilde{\beta}_{nmj} \tilde{\beta}_{\ell qj} = \tilde{\beta}_{nmlq}. \quad (\text{D.2b})$$

For the special cases with some or all of the KK indices being equal, we can recast above sum rules in the following simpler forms:

$$\sum_{j=0}^{\infty} a_{nnj}^2 = a_{nnnn}, \quad \sum_{j=0}^{\infty} a_{nnj} a_{mmj} = \sum_{j=0}^{\infty} a_{nmj}^2 = a_{nnmm}, \quad (\text{D.3a})$$

$$\sum_{j=0}^{\infty} \alpha_{nnj}^2 = \alpha_{nnnn}, \quad \sum_{j=0}^{\infty} \alpha_{nnj} \alpha_{mmj} = \sum_{j=0}^{\infty} \alpha_{nmj}^2 = \alpha_{nnmm}, \quad (\text{D.3b})$$

$$\sum_{j=0}^{\infty} \tilde{a}_{nnj}^2 = \tilde{a}_{nnnn}, \quad \sum_{j=0}^{\infty} \tilde{a}_{nnj} \tilde{a}_{mmj} = \sum_{j=0}^{\infty} \tilde{a}_{nmj}^2 = \tilde{a}_{nnmm}, \quad (\text{D.3c})$$

$$\sum_{j=0}^{\infty} \tilde{\beta}_{nnj}^2 = \tilde{\beta}_{nnnn}, \quad \sum_{j=0}^{\infty} \tilde{\beta}_{nnj} \tilde{\beta}_{mmj} = \sum_{j=0}^{\infty} \tilde{\beta}_{nmj}^2 = \tilde{\beta}_{nnmm}. \quad (\text{D.3d})$$

Many of these relations will be used in Sections 3 and 4 of the main text.

Next, by utilizing the equation of motion (2.14a) and integration by parts, we derive the following relation:

$$\begin{aligned} \int_0^L dz e^{A(z)} M_n^2 f_n f_m f_\ell &= - \int_0^L dz \partial_z (e^{A(z)} \partial_z f_n) f_m f_\ell \\ &= \int_0^L dz e^{A(z)} (f'_n f'_m f_\ell + f'_n f_m f'_\ell). \end{aligned} \quad (\text{D.4})$$

Thus, using the above relation and the definitions given in Eqs.(C.3a)-(C.3b), we obtain the following identities which are connected to each other by cycling the three KK indices  $(n, m, \ell)$ :

$$\begin{aligned} M_n^2 a_{nml} &= M_n M_m \tilde{a}_{nml} + M_n M_\ell \tilde{a}_{nlm}, \\ M_m^2 a_{nml} &= M_n M_m \tilde{a}_{nml} + M_m M_\ell \tilde{a}_{mln}, \\ M_\ell^2 a_{nml} &= M_n M_\ell \tilde{a}_{nlm} + M_m M_\ell \tilde{a}_{mln}. \end{aligned} \quad (\text{D.5})$$

From these identities, we can further derive a relation connecting the coupling coefficient  $a_{nml}$  to  $\tilde{a}_{nml}$ :

$$(M_n^2 + M_m^2 - M_\ell^2) a_{nml} = 2 M_n M_m \tilde{a}_{nml}, \quad (\text{D.6})$$

which just gives Eq.(3.5) in the main text. Similarly, we can derive the following relation between the wavefunction couplings  $\alpha_{nml}$  and  $\tilde{\alpha}_{nml}$ :

$$(M_n^2 + M_m^2 - M_\ell^2) \alpha_{nml} = 2 M_n M_m \tilde{\alpha}_{nml}, \quad (\text{D.7})$$

which further gives Eq.(3.16) in the main text.

Moreover, by utilizing Eq.(2.14a) and integration by parts, we derive the following relation containing four wavefunctions of the warped KK gauge theory:

$$\int_0^L dz e^{A(z)} M_n^2 f_n f_m f_\ell f_q = \int_0^L dz e^{A(z)} (f'_n f'_m f_\ell f_q + f'_n f_m f'_\ell f_q + f'_n f_\ell f_m f'_q). \quad (\text{D.8})$$

With this relation, we further derive the following identities by cycling the four KK indices  $(n, m, \ell, q)$ :

$$\begin{aligned}
M_n^2 a_{nm\ell q} &= M_n M_m \llbracket \tilde{f}_n \tilde{f}_m f_\ell f_q \rrbracket + M_n M_\ell \llbracket \tilde{f}_n \tilde{f}_\ell f_m f_q \rrbracket + M_n M_q \llbracket \tilde{f}_n \tilde{f}_q f_m f_\ell \rrbracket, \\
M_m^2 a_{nm\ell q} &= M_n M_m \llbracket \tilde{f}_n \tilde{f}_m f_\ell f_q \rrbracket + M_m M_\ell \llbracket \tilde{f}_m \tilde{f}_\ell f_n f_q \rrbracket + M_m M_q \llbracket \tilde{f}_m \tilde{f}_q f_n f_\ell \rrbracket, \\
M_\ell^2 a_{nm\ell q} &= M_n M_\ell \llbracket \tilde{f}_n \tilde{f}_\ell f_m f_q \rrbracket + M_m M_\ell \llbracket \tilde{f}_m \tilde{f}_\ell f_n f_q \rrbracket + M_\ell M_q \llbracket \tilde{f}_\ell \tilde{f}_q f_n f_m \rrbracket, \\
M_q^2 a_{nm\ell q} &= M_n M_q \llbracket \tilde{f}_n \tilde{f}_q f_m f_\ell \rrbracket + M_m M_q \llbracket \tilde{f}_m \tilde{f}_q f_n f_\ell \rrbracket + M_\ell M_q \llbracket \tilde{f}_\ell \tilde{f}_q f_n f_m \rrbracket.
\end{aligned} \tag{D.9}$$

In the above Eq.(D.9), we have used brackets  $\llbracket \cdots \rrbracket$  to denote the integration over product of the 5d wavefunctions:

$$\llbracket \mathbf{X}_{n_1} \cdots \mathbf{X}_{n_N} \rrbracket = \frac{1}{L} \int_0^L dz e^{rA(z)} \mathbf{X}_{n_1}(z) \cdots \mathbf{X}_{n_N}(z), \tag{D.10}$$

where  $\mathbf{X}_n$  represents the 5d wavefunctions and the parameter  $r$  is chosen as  $r=1$  ( $r=3$ ) for the warped KK gauge (gravity) theory. Then, summing up the four identities of Eq.(D.9) and making use of Eq.(D.6), we derive a new sum rule:

$$\sum_{j=0}^{\infty} M_j^2 (a_{nmj} a_{\ell qj} + a_{n\ell j} a_{mqj} + a_{nqj} a_{m\ell j}) = (M_n^2 + M_m^2 + M_\ell^2 + M_q^2) a_{nm\ell q}. \tag{D.11}$$

For the special case of  $n=m=\ell=q$ , the above identity reduces to:

$$\sum_{j=0}^{\infty} M_j^2 a_{nnj}^2 = \frac{4}{3} M_n^2 a_{nnnn}, \tag{D.12}$$

which just reproduces the identity (3.33b) in the main text. As for the warped KK gravity theory, we can derive a new sum rule in similar form:

$$\sum_{j=0}^{\infty} \mathbb{M}_j^2 (\alpha_{nmj} \alpha_{\ell qj} + \alpha_{n\ell j} \alpha_{mqj} + \alpha_{nqj} \alpha_{m\ell j}) = (\mathbb{M}_n^2 + \mathbb{M}_m^2 + \mathbb{M}_\ell^2 + \mathbb{M}_q^2) \alpha_{nm\ell q}. \tag{D.13}$$

For the special case of  $n=m=\ell=q$ , we simplify the above identity as follows:

$$\sum_{j=0}^{\infty} \mathbb{M}_j^2 \alpha_{nnj}^2 = \frac{4}{3} \mathbb{M}_n^2 \alpha_{nnnn}, \tag{D.14}$$

which reproduces the identity (3.70b) in the main text.

Next, we prove the identity (3.21) in the main text. For this purpose, we first derive the following relation,

$$\begin{aligned}
L \mathbb{M}_n \mathbb{M}_m \tilde{\beta}_{nm\ell} &= \int_0^L dz e^{3A(z)} (\mathbb{M}_n \mathbf{w}_n) (\mathbb{M}_m \mathbf{w}_m) \mathbf{u}_\ell \\
&= \int_0^L dz \left[ \partial_z (e^{A(z)} \mathbf{v}_n) \right] \left[ e^{2A(z)} (A' + \partial_z) \mathbf{v}_m \right] \mathbf{u}_\ell \\
&= - \int_0^L dz (e^{A(z)} \mathbf{v}_n) \partial_z \left[ e^{2A(z)} (A' + \partial_z) \mathbf{v}_m \right] \mathbf{u}_\ell - \int_0^L dz (e^{A(z)} \mathbf{v}_n) \left[ e^{2A} (A' + \partial_z) \mathbf{v}_m \right] \mathbf{u}'_\ell
\end{aligned}$$

$$= LM_m^2 \tilde{\alpha}_{nml} - LM_m M_\ell \tilde{\rho}_{nlm}, \quad (\text{D.15})$$

which leads to the identity:

$$M_n \tilde{\beta}_{nml} = M_m \tilde{\alpha}_{nml} - M_\ell \tilde{\rho}_{nlm}. \quad (\text{D.16})$$

For the right-hand side of Eq.(D.16), the second term contains the trilinear coupling coefficient  $\tilde{\rho}_{nml}$  for which we can derive the following relation:

$$\begin{aligned} LM_n \tilde{\rho}_{mln} &= \int_0^L dz \left[ \partial_z (e^{A(z)} \mathbf{v}_n) \right] (e^{A(z)} \mathbf{v}_m) (e^{A(z)} \mathbf{v}_\ell), \\ &= - \int_0^L dz (e^{A(z)} \mathbf{v}_n) \left[ \partial_z (e^{A(z)} \mathbf{v}_m) \right] (e^{A(z)} \mathbf{v}_\ell) - \int_0^L dz (e^{A(z)} \mathbf{v}_n) (e^{A(z)} \mathbf{v}_m) \left[ \partial_z (e^{A(z)} \mathbf{v}_\ell) \right] \\ &= -LM_m \tilde{\rho}_{nlm} - LM_\ell \tilde{\rho}_{nml}, \end{aligned} \quad (\text{D.17})$$

where we have used Eqs.(2.52) and (2.53). We can reexpress Eq.(D.17) as follows:

$$M_n \tilde{\rho}_{mln} + M_m \tilde{\rho}_{nlm} + M_\ell \tilde{\rho}_{nml} = 0. \quad (\text{D.18})$$

Using Eqs.(D.16) and (D.7) as well as cycling their three KK indices, we further derive the following relations:

$$2M_n^2 M_m^2 \tilde{\beta}_{nml} = M_m^2 (M_m^2 + M_n^2 - M_\ell^2) \alpha_{nml} - 2M_n M_m^2 M_\ell \tilde{\rho}_{nlm}, \quad (\text{D.19a})$$

$$2M_n^2 M_m^2 \tilde{\beta}_{nml} = M_n^2 (M_m^2 + M_n^2 - M_\ell^2) \alpha_{nml} - 2M_n^2 M_m M_\ell \tilde{\rho}_{mln}, \quad (\text{D.19b})$$

$$2M_m^2 M_\ell^2 \tilde{\beta}_{mln} = M_\ell^2 (M_\ell^2 + M_m^2 - M_n^2) \alpha_{nml} - 2M_n M_m M_\ell^2 \tilde{\rho}_{nml}, \quad (\text{D.19c})$$

$$2M_m^2 M_\ell^2 \tilde{\beta}_{mln} = M_m^2 (M_\ell^2 + M_m^2 - M_n^2) \alpha_{nml} - 2M_n M_n^2 M_\ell \tilde{\rho}_{nlm}, \quad (\text{D.19d})$$

$$2M_\ell^2 M_n^2 \tilde{\beta}_{lnm} = M_\ell^2 (M_\ell^2 + M_n^2 - M_m^2) \alpha_{nml} - 2M_n M_m M_\ell^2 \tilde{\rho}_{nml}, \quad (\text{D.19e})$$

$$2M_\ell^2 M_n^2 \tilde{\beta}_{lnm} = M_n^2 (M_\ell^2 + M_n^2 - M_m^2) \alpha_{nml} - 2M_n^2 M_m M_\ell \tilde{\rho}_{mln}. \quad (\text{D.19f})$$

With the six relations above, we compute the sum  $\frac{3}{2} [(a)+(b)] + \frac{1}{2} [(c)-(d)+(e)-(f)]$  and impose Eq.(D.17), with which we derive the final identity:

$$\left[ (M_n^2 + M_m^2 - M_\ell^2)^2 + 2M_n^2 M_m^2 \right] \alpha_{nml} = 6M_n^2 M_m^2 \tilde{\beta}_{nml}. \quad (\text{D.20})$$

This just reproduces the identity (3.21) in the main text.

For the derivation of Eq.(3.26) in the main text, we can sum up the third and fifth identities in Eq.(D.19) and obtain the following:

$$4M_n M_m M_\ell^2 \tilde{\rho}_{nml} = 2M_\ell^4 \alpha_{nml} - 2M_m^2 M_\ell^2 \tilde{\beta}_{mln} - 2M_n^2 M_\ell^2 \tilde{\beta}_{nlm}. \quad (\text{D.21})$$

With this and further using the identity (D.20), we arrive at

$$\left[ 2M_\ell^4 - (M_n^2 - M_m^2)^2 - M_\ell^2 (M_n^2 + M_m^2) \right] \alpha_{nml} = 6M_n M_m M_\ell^2 \tilde{\rho}_{nml}. \quad (\text{D.22})$$

This just reproduces Eq.(3.26) in the main text.

Then, we derive two sum rules Eqs.(3.45b) and (3.45c) which are used for computing the inelastic scattering amplitudes of longitudinal KK gauge bosons. For the Eq.(3.45b), we can set  $(\ell, q) = (n, m)$  in Eq.(D.11), and derive the sum rule:

$$\sum_{j=0}^{\infty} M_j^2 a_{nmj}^2 = \sum_{j=0}^{\infty} \left( M_n^2 + M_m^2 - \frac{1}{2} M_j^2 \right) a_{nnj} a_{mmj}. \quad (\text{D.23})$$

As for the derivation of Eq.(3.45c), by taking the difference between the first two identities in Eq.(D.5), we arrive at

$$(M_n^2 - M_m^2) a_{nmj} = M_j (M_n \tilde{a}_{njm} - M_m \tilde{a}_{mjn}). \quad (\text{D.24})$$

Then, squaring the above formula and sum over  $j$ , we derive the following:

$$\begin{aligned} \sum_{j=1}^{\infty} (M_n^2 - M_m^2)^2 M_j^{-2} a_{nmj}^2 &= \sum_{j=1}^{\infty} (M_n \tilde{a}_{njm} - M_m \tilde{a}_{mjn})^2 \\ &= \sum_{j=1}^{\infty} (M_n^2 \tilde{a}_{njm}^2 - 2M_n M_m \tilde{a}_{njm} \tilde{a}_{mjn} + M_m^2 \tilde{a}_{mjn}^2) \\ &= \sum_{j=1}^{\infty} (M_n^2 \tilde{a}_{nnj} a_{mmj} - 2M_n M_m \tilde{a}_{nmj} a_{nmj} + M_m^2 \tilde{a}_{mmj} a_{nnj}) \\ &= \sum_{j=1}^{\infty} \left[ \frac{1}{2} (2M_n^2 - M_j^2) a_{nnj} a_{mmj} - (M_n^2 + M_m^2 - M_j^2) a_{nmj}^2 + \frac{1}{2} (2M_m^2 - M_j^2) a_{mmj} a_{nnj} \right] \\ &= \sum_{j=0}^{\infty} M_j^2 (a_{nmj}^2 - a_{nnj} a_{mmj}), \end{aligned} \quad (\text{D.25})$$

where for the third equality sign, we have imposed the following completeness relation:

$$\sum_{j=1}^{\infty} \tilde{a}_{njm} \tilde{a}_{\ell jq} = \llbracket \tilde{\mathbf{f}}_n \tilde{\mathbf{f}}_m \tilde{\mathbf{f}}_{\ell} \tilde{\mathbf{f}}_q \rrbracket = \sum_{j=1}^{\infty} \llbracket \tilde{\mathbf{f}}_n \tilde{\mathbf{f}}_{\ell} \tilde{\mathbf{f}}_j \rrbracket \llbracket \mathbf{f}_m \mathbf{f}_q \mathbf{f}_j \rrbracket = \sum_{j=1}^{\infty} \tilde{a}_{n\ell j} a_{mqj}. \quad (\text{D.26})$$

For the fourth equality sign of Eq.(D.25), we have used Eq.(D.6). Moreover, substituting Eq.(D.23) into Eq.(D.25), we derive the sum rule identity:

$$\sum_{j=1}^{\infty} (M_n^2 - M_m^2)^2 M_j^{-2} a_{nmj}^2 = \sum_{j=0}^{\infty} (M_n^2 + M_m^2 - \frac{3}{2} M_j^2) a_{nnj} a_{mmj}, \quad (\text{D.27})$$

which is Eq.(3.45c) in the main text.

Next, we derive the sum rule identities in Eq.(3.57) of Sec. 3.2. We prove the  $s$ -channel sum rule (3.57c) as an example and other sum rules in Eq.(3.57) be readily obtained from Eq.(3.57c) by permuting the KK indices. From Eq.(D.24), we derive the following:

$$\begin{aligned} (M_n^2 - M_m^2) (M_{\ell}^2 - M_q^2) \sum_{j=1}^{\infty} M_j^{-2} a_{nmj} a_{\ell qj} \\ = \sum_{j=1}^{\infty} \left[ (M_n^2 - M_m^2) M_j^{-1} a_{nmj} \right] \left[ (M_{\ell}^2 - M_q^2) M_j^{-1} a_{\ell qj} \right] \end{aligned}$$

$$\begin{aligned}
&= \sum_{j=1}^{\infty} (M_n \tilde{a}_{njm} - M_m \tilde{a}_{mjn}) (M_\ell \tilde{a}_{\ell jq} - M_q \tilde{a}_{qj\ell}) \\
&= \sum_{j=1}^{\infty} (M_n M_\ell \tilde{a}_{njm} \tilde{a}_{\ell jq} - M_n M_q \tilde{a}_{njm} \tilde{a}_{qj\ell} - M_m M_\ell \tilde{a}_{mjn} \tilde{a}_{\ell jq} + M_m M_q \tilde{a}_{mjn} \tilde{a}_{qj\ell}) \\
&= \sum_{j=1}^{\infty} (M_n M_\ell \tilde{a}_{n\ell j} a_{mqj} - M_n M_q \tilde{a}_{nqj} a_{m\ell j} - M_m M_\ell \tilde{a}_{m\ell j} a_{nqj} + M_m M_q \tilde{a}_{mqj} a_{n\ell j}), \quad (\text{D.28})
\end{aligned}$$

where we have applied Eqs.(D.5) and (D.26) for the second and fourth equality signs, respectively. Then, substituting Eq.(D.6) into Eq.(D.28), we finally arrive at

$$\begin{aligned}
&(M_n^2 - M_m^2)(M_\ell^2 - M_q^2) \sum_{j=1}^{\infty} M_j^{-2} a_{nmj} a_{\ell qj} \\
&= \sum_{j=1}^{\infty} \left[ \frac{1}{2} (M_n^2 + M_\ell^2 - M_j^2) a_{n\ell j} a_{mqj} - \frac{1}{2} (M_n^2 + M_q^2 - M_j^2) a_{nqj} a_{m\ell j} \right. \\
&\quad \left. - \frac{1}{2} (M_m^2 + M_\ell^2 - M_j^2) a_{m\ell j} a_{nqj} + \frac{1}{2} (M_m^2 + M_q^2 - M_j^2) a_{mqj} a_{n\ell j} \right] \\
&= \sum_{j=0}^{\infty} M_j^2 (a_{nqj} a_{m\ell j} - a_{n\ell j} a_{mqj}). \quad (\text{D.29})
\end{aligned}$$

This just reproduces the sum rule identity (3.57c) given in the main text.

Finally, we derive the identity (3.72) in the main text. With Eq.(3.18) and Eq.(D.14), we first derive the following identity:

$$\sum_{j=0}^{\infty} \alpha_{nnj} \tilde{\alpha}_{nnj} = \frac{1}{3} \alpha_{nnnn}. \quad (\text{D.30})$$

By using Eq.(3.18) and Eq.(D.14), we compute:

$$\sum_{j=0}^{\infty} \hat{r}_j^6 \alpha_{nnj}^2 = \sum_{j=0}^{\infty} 4 \hat{r}_j^2 (\alpha_{nnj} - \tilde{\alpha}_{nnj})^2 = \sum_{j=0}^{\infty} 4(4 + \hat{r}_j^2) \tilde{\alpha}_{nnj}^2, \quad (\text{D.31})$$

where we have used the notation  $\hat{r}_j = \mathbb{M}_j / \mathbb{M}_n$ . Then, we compute the following:

$$\begin{aligned}
\mathbb{M}_j^2 \tilde{\alpha}_{nnj} &= \mathbb{M}_j^2 \int_0^L dz e^{3A(z)} v_n^2 u_j \\
&= \int_0^L dz e^{3A(z)} u_j \left[ -12A'' v_n^2 + 12A' u_n v_n + 2v_n^2 - 2\mathbb{M}_n^2 u_n^2 \right], \quad (\text{D.32a})
\end{aligned}$$

$$\sum_{j=0}^{\infty} \mathbb{M}_j^2 \tilde{\alpha}_{nnj}^2 = -2\mathbb{M}_n^2 \alpha_{nnj} \tilde{\alpha}_{nnj} + 2\mathbb{M}_n^2 \tilde{\alpha}_{nnj}^2 + I_1 + I_2, \quad (\text{D.32b})$$

where the  $I_1$  and  $I_2$  represent two integrals:

$$I_1 = -12\mathbb{M}_n \int_0^L dz e^{3A(z)} A' u_n v_n^3 = -2\mathbb{M}_n^2 \sum_{j=0}^{\infty} (\tilde{\alpha}_{nnj}^2 - 3\alpha_{nnj} \tilde{\alpha}_{nnj}), \quad (\text{D.33a})$$



$$I_2 = -12 \int_0^L dz e^{3A(z)} A'' v_n^4 = -\frac{1}{2} I_1. \quad (\text{D.33b})$$

In the above we have denoted  $A' = \partial_z A(z)$  and  $A'' = \partial_z^2 A(z)$ . Combining Eqs.(D.32) and (D.33), we further compute the summation of Eq.(D.31):

$$\sum_{j=0}^{\infty} \hat{r}_j^6 \alpha_{nnj}^2 = \sum_{j=0}^{\infty} 4(5\tilde{\alpha}_{nnj}^2 + \alpha_{nnj} \tilde{\alpha}_{nnj}) = \sum_{j=0}^{\infty} \left(5\hat{r}_j^4 - \frac{16}{3}\right) \alpha_{nnj}^2, \quad (\text{D.34})$$

where we have made use of Eq.(3.18) and Eq.(D.14). We can re-express Eq.(D.34) as follows:

$$\sum_{j=0}^{\infty} \left(\hat{r}_j^6 - 5\hat{r}_j^4 + \frac{16}{3}\right) \alpha_{nnj}^2 = 0, \quad (\text{D.35})$$

which just reproduces the identity (3.72) in the main text.

## References

- [1] For a recent review book, G. Travaglini *et al.*, “The SAGEX Review on Scattering Amplitudes”, J. Phys. A 55 (2022) 44, 443001 [arXiv:2203.13011 [hep-th]].
- [2] T. Kaluza, “On the Unification Problem in Physics”, Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.) 1921 (1921) 966 [Int. J. Mod. Phys. D 27 (2018) 1870001, [arXiv:1803.08616 [physics.hist-ph]]; O. Klein, “Quantum Theory and Five-Dimensional Theory of Relativity”, Z. Phys. 37 (1926) 895 [Surveys High Energy Phys. 5 (1986) 241].
- [3] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Lett. B 429 (1998) 263 [arXiv:hep-ph/9803315]; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Lett. B 436 (1998) 257 [arXiv:hep-ph/9804398].
- [4] L. Randall and R. Sundrum, Phys. Rev. Lett. 83 (1999) 3370 [arXiv:hep-ph/9905221]; Phys. Rev. Lett. 83 (1999) 4690 [arXiv:hep-th/9906064].
- [5] For reviews, e.g., R. Sundrum, arXiv:hep-th/0508134; C. Csaki, arXiv:hep-ph/0404096; R. Rattazzi, arXiv:hep-ph/0607055.
- [6] E.g., M. J. G. Veltman, Nucl. Phys. B 7 (1968) 637-650.
- [7] M. Fierz and W. Pauli, Proc. Roy. Soc. Lond. A 173 (1939) 211.
- [8] For a review, K. Hinterbichler, Rev. Mod. Phys. 84 (2012) 671 [arXiv:1105.3735 [hep-th]].
- [9] H. van Dam and M. J. G. Veltman, Nucl. Phys. B 22 (1970) 397.
- [10] V. I. Zakharov JETP Letters (Sov. Phys.) 12 (1970) 312.
- [11] R. S. Chivukula, D. A. Dicus, H.-J. He, Phys. Lett. B 525 (2002) 175 [hep-ph/0111016].
- [12] R. S. Chivukula and H.-J. He, Phys. Lett. B 532 (2002) 121 [hep-ph/0201164].
- [13] H.-J. He, Int. J. Mod. Phys. A 20 (2005) 3362 [hep-ph/0412113], presentation in the proceedings of DPF-2004: Annual Meeting of the Division of Particles and Fields, American Physical Society, Riverside, California, USA, August 26-31, 2004.
- [14] L. Dolan and M. Duff, Phys. Rev. Lett. 52 (1984) 14.

- [15] Y.-F. Hang and H.-J. He, Phys. Rev. D 105 (2022) 084005 [hep-th/2106.04568].
- [16] Y.-F. Hang and H.-J. He, Research 2022 (2022) 9860945 [hep-th/2207.11214].
- [17] F. Englert and R. Brout, Phys. Rev. Lett. 13 (1964) 321; P. W. Higgs, Phys. Rev. Lett. 13 (1964) 508; Phys. Lett. 12 (1964) 132; G. S. Guralnik, C. R. Hagen and T. Kibble, Phys. Rev. Lett. 13 (1965) 585; T. Kibble, Phys. Rev. 155 (1967) 1554.
- [18] L. Randall and R. Sundrum, Phys. Rev. Lett. 83 (1999) 3370 [hep-ph/9905221].
- [19] H. Davoudiasl, J. L. Hewett, T. G. Rizzo Phys. Lett. B 473 (2000) 43 [arXiv:hep-ph/9911262].
- [20] A. Pomarol, Phys. Lett. B 486 (2000) 153 [hep-ph/9911294].
- [21] R. S. Chivukula, J. A. Gill, K. A. Mohan, D. Sengupta, E. H. Simmons and X. Wang, Phys. Rev. D 109 (2024) 075016 [arXiv:2312.08576 [hep-ph]].
- [22] L. Randall and M. D. Schwartz, JHEP 11 (2001) 003 [hep-th/0108114].
- [23] H. Kawai, D. C. Lewellen, and S. H. H. Tye, Nucl. Phys. B 269 (1986) 1-23.
- [24] Z. Bern, J. J. M. Carrasco and H. Johansson, Phys. Rev. D 78 (2008) 085011 [hep-ph/0805.3993]; Phys. Rev. Lett. 105 (2010) 061602 [hep-th/1004.0476].
- [25] For a comprehensive review, Z. Bern, J. J. M. Carrasco, H. Johansson, and R. Roiban, (2019) [hep-th/1909.01358].
- [26] Y. Li, Y.-F. Hang, H.-J. He, S. He, JHEP 02 (2022) 120 [hep-th/2111.12042].
- [27] Y. Li, Y.-F. Hang, H.-J. He, JHEP 03 (2023) 254 [hep-th/2209.11191].
- [28] Y.-F. Hang, H.-J. He, C. Shen, JHEP 01 (2022) 153 [arXiv:2110.05399 [hep-th]]; Y.-F. Hang, H.-J. He, C. Shen, Research 6 (2023) 0072 [arXiv:2406.13671 [hep-th]].
- [29] M. C. González, A. Momeni, J. Rumbutis, JHEP 08 (2021) 116 [arXiv:2107.00611 [hep-th]].
- [30] C. de Rham and G. Gabadadze, Phys. Rev. D 82 (2010) 044020 [arXiv:1007.0443 [hep-th]]; C. de Rham, G. Gabadadze, and A. J. Tolley, Phys. Rev. Lett. 106 (2011) 231101 [arXiv:1011.1232 [hep-th]]; A. Momeni, J. Rumbutis, and A. J. Tolley, JHEP 12 (2020) 030 [2004.07853 [hep-th]].
- [31] L. A. Johnson, C. R. T. Jones, and S. Paranjape, JHEP 02 (2021) 148 [arXiv:2004.12948 [hep-th]].
- [32] M. Chiodaroli, M. Gunaydin, H. Johansson, and R. Roiban, JHEP 06 (2017) 064 [arXiv:1511.01740 [hep-th]].
- [33] A. Momeni, J. Rumbutis, and A. J. Tolley, JHEP 08 (2021) 081 [arXiv:2012.09711 [hep-th]].
- [34] T. Bargheer, S. He, and T. McLoughlin, Phys. Rev. Lett. 108 (2012) 231601 [arXiv:1203.0562 [hep-th]].
- [35] Y. t. Huang and H. Johansson, Phys. Rev. Lett. 110 (2013) 171601 [arXiv:1210.2255 [hep-th]].
- [36] N. Moynihan, JHEP 12 (2020) 163 [arXiv:2006.15957 [hep-th]]; D. J. Burger, W. T. Emond and N. Moynihan, JHEP 01 (2022) 017 [arXiv:2103.10416 [hep-th]]; N. Moynihan, JHEP 05 (2024) 310 [arXiv:2110.02209 [hep-th]].
- [37] H. Johansson and A. Ochirov, JHEP 09 (2019) 40 [arxiv:1906.12292 [hep-th]].
- [38] Y. F. Bautista and A. Guevara, JHEP 11 (2021) 184 [arxiv:1908.11349 [hep-th]].
- [39] J. Plefka, C. Shi, and T. Wang, Phys. Rev. D 101 (2020) 066004, [arxiv:1911.06785 [hep-th]].

- [40] H. J. He, Y. P. Kuang and X. Li, Phys. Rev. D 49 (1994) 4842-4872; Phys. Rev. Lett. 69 (1992) 2619.
- [41] H.-J. He and W. B. Kilgore, Phys. Rev. D 55 (1997) 1515 [hep-ph/9609326].
- [42] For a review, H. J. He, Y. P. Kuang and C. P. Yuan, hep-ph/9704276, DESY-97-056, in the proceedings of CCAST workshop on “Physics at the TeV Energy Scale”, vol.72, p.119 (1996).
- [43] R. S. Chivukula, D. A. Dicus, H.-J. He, and S. Nandi, Phys. Lett. B 562 (2003) 109 [hep-ph/0302263].
- [44] P. Callin and F. Ravndal, Phys. Rev. D 72 (2005) 064026 [hep-ph/0412109].
- [45] T. Gherghetta, M. Peloso and E. Poppitz, Phys. Rev. D 72 (2005) 104003 [hep-th/0507245].
- [46] Z. Chacko, M. Graesser, C. Grojean and L. Pilo, Phys. Rev. D 70 (2004) 084028 [hep-th/0312117].
- [47] C. S. Lim, T. Nagasawa, S. Ohay, K. Sakamoto and M. Sakamoto, Phys. Rev. D 77 (2008) 065009 [arXiv:0801.0845 [hep-th]].
- [48] R. S. Chivukula, E. H. Simmons, X. Wang, Phys. Rev. D 106 (2022) 035026 [arXiv:2207.02887 [hep-ph]].
- [49] R. S. Chivukula, D. Foren, K. A. Mohan, D. Sengupta and E. H. Simmons, Phys. Rev. D 101 (2020) 075013 [arXiv:2002.12458 [hep-ph]].
- [50] W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. 83 (1999) 4922 [hep-ph/9907447]; Phys. Lett. B 475 (2000) 275 [hep-ph/9911457].
- [51] R. S. Chivukula, D. Foren, K. A. Mohan, D. Sengupta and E. H. Simmons, Phys. Rev. D 103 (2021) 095024 [arXiv:2104.08169 [hep-ph]]; *ibid*, D 107 (2023) 035015 [arXiv:2206.10628 [hep-ph]]; R. S. Chivukula, E. H. Simmons, X. Wang, *ibid*, D 106 (2022) 035026 [arXiv:2207.02887 [hep-ph]].
- [52] S. Weinberg, Physica A 96 (1979) 327, no.1-2.
- [53] V. Del Duca, L. J. Dixon, and F. Maltoni, Nucl. Phys. B 571 (2000) 51 [hep-ph/9910563].
- [54] R. Kleiss and H. Kuijf, Nucl. Phys. B 312 (1989) 616.
- [55] F. Cachazo, S. He and E. Y. Yuan, Phys. Rev. D 90 (2014) 065001 [arXiv:1306.6575 [hep-th]]; Phys. Rev. Lett. 113 (2014) 171601 [arXiv:1307.2199 [hep-th]]; JHEP 1407 (2014) 033 [arXiv:1309.0885 [hep-th]].
- [56] F. Cachazo, S. He and E. Y. Yuan, JHEP 07, 149 (2015) [arXiv:1412.3479 [hep-th]].