

# Toward mean-field bound for critical temperature on Nishimori line

Manaka Okuyama<sup>1</sup> and Masayuki Ohzeki<sup>1,2,3,4</sup>

<sup>1</sup>*Graduate School of Information Sciences,  
Tohoku University, Sendai 980-8579, Japan*

<sup>2</sup>*International Research Frontier Initiative,  
Tokyo Institute of Technology, Tokyo 105-0023, Japan*

<sup>3</sup>*Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan and*

<sup>4</sup>*Sigma-i Co., Ltd., Tokyo 108-0075, Japan*

(Dated: June 25, 2024)

## Abstract

The critical inverse temperature of the mean-field approximation establishes a lower bound of the true critical inverse temperature in a broad class of ferromagnetic spin models. This is referred to as the mean-field bound for the critical temperature. In this study, we explored the possibility of a corresponding mean-field bound for the critical temperature in Ising spin glass models with Gaussian randomness on the Nishimori line. On this line, the critical inverse temperature of the mean-field approximation is given by  $\beta_{MF}^{NL} = \sqrt{1/z}$ , where  $z$  is the coordination number. Using the Griffiths inequalities on the Nishimori line, we proved that there is zero spontaneous magnetization in the high-temperature region  $\beta < \beta_{MF}^{NL}/2$ . In other words, the true critical inverse temperature  $\beta_c^{NL}$  on the Nishimori line is always bounded by  $\beta_c^{NL} \geq \beta_{MF}^{NL}/2$ . Unfortunately, we have not succeeded in obtaining the corresponding mean-field bound  $\beta_c^{NL} \geq \beta_{MF}^{NL}$  on the Nishimori line.

## I. INTRODUCTION

The free energy of the mean-field approximation is always an upper bound of the true free energy. This universal constraint on the free energy results from the Gibbs–Bogoliubov inequality [1, 2], which is based on the convexity of the pressure function  $\log Z$  for the inverse temperature [3].

For ferromagnetic models, additional constraints on the critical inverse temperature can be established. It has been rigorously proved that the critical inverse temperature of the mean-field approximation sets a lower bound for the true critical inverse temperature in a broad class of ferromagnetic spin models. This is referred to as the mean-field bound for the critical temperature [4–8]. It is also known that the mean-field bound for magnetization holds [9–13]. Unfortunately, the proofs of these bounds strongly rely on ferromagnetic interactions, making it seemingly impossible to extend them to spin glass models.

Concerning Ising spin glass models with Gaussian randomness on the Nishimori line [14, 15], it was recently proved that the counterpart of the Gibbs–Bogoliubov inequality holds on the Nishimori line by using the convexity of the quenched pressure function  $\mathbb{E}[\log Z]$  with respect to the parameters along the Nishimori line [16]. The Gibbs–Bogoliubov inequality on the Nishimori line shows that the quenched free energy of the mean-field approximation on the Nishimori line establishes an upper bound for the true quenched free energy on the Nishimori line [16]. This constraint on the quenched free energy of the mean-field approximation on the Nishimori line is notably similar to that of the conventional mean-field approximation resulting from the conventional Gibbs–Bogoliubov inequality. Furthermore, Ising spin glass models on the Nishimori line behave similarly to ferromagnetic models, and various exact and rigorous results have been obtained [17–41]. Then, from the mean-field bound for the critical temperature in ferromagnetic models, we have a natural question: how does the critical inverse temperature  $\beta_{MF}^{NL}$  of the mean-field approximation on the Nishimori line relate to the true critical inverse temperature  $\beta_c^{NL}$  on the Nishimori line? In particular, does the corresponding mean-field bound  $\beta_c^{NL} \geq \beta_{MF}^{NL}$  hold on the Nishimori line?

In this study, we partially answered this question. Using the Griffiths inequalities on the Nishimori line [25], we proved that there is always zero spontaneous magnetization in the high-temperature region  $\beta < \beta_{MF}^{NL}/2$ . This immediately implies that the true critical inverse temperature on the Nishimori line is always bounded by  $\beta_c^{NL} \geq \beta_{MF}^{NL}/2$ . Unfortunately, it remains an open

problem whether the corresponding mean-field bound on the Nishimori line,  $\beta_c^{NL} \geq \beta_{MF}^{NL}$ , holds.

The remainder of this paper is organized as follows. In Sec. II, we define the model and present the main result (Theorem 1). Section III presents the proof of Theorem 1. Finally, a discussion is presented in Sec. IV.

## II. DEFINITIONS AND RESULTS

Let  $\Lambda$  be any lattice invariant under translations with coordination number  $z$  (for example,  $\Lambda = Z^d$  with  $z = 2d$ ). Free or fixed boundary conditions on the Nishimori line are imposed [25]. To each site  $i \in \Lambda$ , we associate an Ising spin  $\sigma_i = \pm 1$ . The Hamiltonian of the Ising spin glass model with Gaussian randomness on the Nishimori line is defined as

$$H = - \sum_{\langle ij \rangle} J_{ij} \sigma_i \sigma_j - \sum_i h_i \sigma_i, \quad (1)$$

where  $\langle ij \rangle$  runs over interacting spin pairs on  $\Lambda$ , and  $J_{ij}$  and  $h_i$  are independent Gaussian random variables distributed as follows:

$$J_{ij} \stackrel{\text{iid}}{\sim} \mathcal{N}(\beta, 1), \quad h_i \stackrel{\text{iid}}{\sim} \mathcal{N}(\beta h, h), \quad (2)$$

where  $h \geq 0$ , and  $\beta$  is the inverse temperature. The choice of the Gaussian distribution (2) satisfies the condition of the Nishimori line [15].

The partition function is given by

$$Z = \text{Tr}(e^{-\beta H}), \quad (3)$$

where  $\text{Tr}(\dots)$  denotes the summation over all spin variables. The quenched free energy on the Nishimori line in the thermodynamic limit is defined as

$$-\beta f_{NL} = \lim_{|\Lambda| \rightarrow \infty} \frac{1}{|\Lambda|} \mathbb{E}[\log Z], \quad (4)$$

where  $|\Lambda|$  is the number of the sites, and  $\mathbb{E}[\dots]$  denotes the expectation with respect to all random variables. The thermal average  $\langle \dots \rangle$  is given by

$$\langle \dots \rangle = \frac{\text{Tr}(\dots e^{-\beta H})}{Z}. \quad (5)$$

The magnetization on the Nishimori line (equivalently, the spin-glass order parameter) in the thermodynamic limit is defined as

$$m(\beta, h) = \lim_{|\Lambda| \rightarrow \infty} \mathbb{E}[\langle \sigma_i \rangle] = \lim_{|\Lambda| \rightarrow \infty} \mathbb{E}[\langle \sigma_i \rangle^2] \geq 0, \quad (6)$$

and the spontaneous magnetization is defined as

$$m_s(\beta) = \lim_{h \rightarrow +0} m(\beta, h). \quad (7)$$

The critical inverse temperature  $\beta_c^{NL}$  is defined as

$$\beta_c^{NL} = \inf \{ \beta \geq 0 \mid m_s(\beta) > 0 \}. \quad (8)$$

The quenched free energy of the mean-field approximation on the Nishimori line is given by

$$-\beta f_{MF}^{NL} = \int Dy \log 2 \cosh(\beta \sqrt{zq + hy} + \beta^2 zq + \beta^2 h) + \frac{z\beta^2}{4}(1-q)^2 - \frac{z\beta^2 q^2}{2}, \quad (9)$$

$$q = \int Dy \tanh(\beta \sqrt{zq + hy} + \beta^2 zq + \beta^2 h). \quad (10)$$

The Gibbs–Bogoliubov inequality on the Nishimori line [16] shows that the quenched free energy of the mean-field approximation on the Nishimori line is an upper bound of the true quenched free energy

$$f^{NL} \leq f_{MF}^{NL}. \quad (11)$$

For  $h = 0$ , Eq. (10) implies that the critical inverse temperature of the mean-field approximation on the Nishimori line is given by

$$\beta_{MF}^{NL} = \sqrt{\frac{1}{z}}. \quad (12)$$

Then, our result can be expressed as follows:

**Theorem 1.** *For any positive (or vanishing)  $\beta$  and  $h$ , the magnetization on the Nishimori line satisfies the following inequality:*

$$m(\beta, h) \leq \mathbb{E}[\tanh(\beta h_i)] + 4z\beta^2 m(\beta, h). \quad (13)$$

*Thus, there is zero spontaneous magnetization, that is,*

$$m_s(\beta) = 0, \quad (14)$$

*in the high-temperature region  $\beta < \sqrt{1/(4z)} = \beta_{MF}^{NL}/2$ , and the critical inverse temperature on the Nishimori line is bounded by*

$$\beta_c^{NL} \geq \frac{\beta_{MF}^{NL}}{2}. \quad (15)$$

### III. PROOF OF THEOREM 1

Our proof is closely related to the proof of the mean-field bound for the critical temperature in the ferromagnetic Ising model using the Griffiths inequalities [6].

Let us fix the site  $i$  and introduce the interpolating parameter  $0 \leq t \leq 1$  into the Gaussian distribution of the interactions connected to the site  $i$  as

$$J_e \rightarrow J_{e,t} \stackrel{\text{iid}}{\sim} \mathcal{N}(\beta t, t) \quad (i \in e). \quad (16)$$

This interpolation does not break the condition of the Nishimori line. Notably, the Gaussian distribution of all other interactions remains unchanged. We denote the  $t$ -dependence of the Gaussian distribution of the interactions as  $\mathbb{E}[\langle \sigma_i \rangle_t]$ . The correlation function at  $t = 1$  coincides with the original correlation function,

$$\mathbb{E}[\langle \sigma_i \rangle_1] = \mathbb{E}[\langle \sigma_i \rangle], \quad (17)$$

and the spin  $\sigma_i$  is decoupled from the other spins at  $t = 0$ ,

$$\mathbb{E}[\langle \sigma_i \rangle_0] = \mathbb{E}[\tanh(\beta h_i)]. \quad (18)$$

The derivative of the correlation function with respect to  $t$  is expressed as follows [25, 34]:

$$\begin{aligned} \frac{d}{dt} \mathbb{E}[\langle \sigma_i \rangle_t] &= \sum_{j \sim i} \beta^2 \mathbb{E}[(\langle \sigma_j \rangle_t - \langle \sigma_i \rangle_t \langle \sigma_i \sigma_j \rangle_t)^2] \\ &= \sum_{j \sim i} \beta^2 \mathbb{E}[\langle \sigma_j \rangle_t^2 + \langle \sigma_i \rangle_t^2 \langle \sigma_i \sigma_j \rangle_t^2 - 2 \langle \sigma_i \rangle_t \langle \sigma_j \rangle_t], \end{aligned} \quad (19)$$

where  $\sum_{j \sim i}$  runs over the site  $j$  connected to the site  $i$ , and we used the correlation identity on the Nishimori line,  $\mathbb{E}[\langle \sigma_i \rangle_t \langle \sigma_j \rangle_t \langle \sigma_i \sigma_j \rangle_t] = \mathbb{E}[\langle \sigma_i \rangle_t \langle \sigma_j \rangle_t]$  [25]. Using a trivial inequality,  $-2\mathbb{E}[\langle \sigma_i \rangle_t \langle \sigma_j \rangle_t] \leq \mathbb{E}[\langle \sigma_i \rangle_t^2 + \langle \sigma_j \rangle_t^2]$ , we obtain

$$\begin{aligned} \frac{d}{dt} \mathbb{E}[\langle \sigma_i \rangle_t] &\leq \sum_{j \sim i} \beta^2 \mathbb{E}[\langle \sigma_j \rangle_t^2 + \langle \sigma_i \rangle_t^2 \langle \sigma_i \sigma_j \rangle_t^2 + \langle \sigma_i \rangle_t^2 + \langle \sigma_j \rangle_t^2] \\ &\leq \sum_{j \sim i} 2\beta^2 \mathbb{E}[\langle \sigma_i \rangle_t^2 + \langle \sigma_j \rangle_t^2] \\ &= \sum_{j \sim i} 2\beta^2 \mathbb{E}[\langle \sigma_i \rangle_t + \langle \sigma_j \rangle_t]. \end{aligned} \quad (20)$$

Then, the Griffiths inequalities on the Nishimori line imply [25]

$$\mathbb{E}[\langle \sigma_i \rangle_t] \leq \mathbb{E}[\langle \sigma_i \rangle_1], \quad (21)$$

which leads to

$$\frac{d}{dt}\mathbb{E}[\langle\sigma_i\rangle_t] \leq \sum_{j\sim i} 2\beta^2\mathbb{E}[\langle\sigma_i\rangle_1 + \langle\sigma_j\rangle_1]. \quad (22)$$

Thus, by integrating  $t$  from 0 to 1 and using  $\mathbb{E}[\langle\sigma_i\rangle_1] = \mathbb{E}[\langle\sigma_i\rangle]$ , we have

$$\begin{aligned} \mathbb{E}[\langle\sigma_i\rangle] &\leq \mathbb{E}[\langle\sigma_i\rangle_0] + \sum_{j\sim i} 2\beta^2\mathbb{E}[\langle\sigma_i\rangle + \langle\sigma_j\rangle] \\ &= \mathbb{E}[\tanh(\beta h_i)] + \sum_{j\sim i} 2\beta^2\mathbb{E}[\langle\sigma_i\rangle + \langle\sigma_j\rangle]. \end{aligned} \quad (23)$$

Finally, we take the thermodynamic limit in Eq. (23). The Griffiths inequalities on the Nishimori line guarantee the existence of the thermodynamic limit for the correlation function,  $\lim_{|\Lambda|\rightarrow\infty}\mathbb{E}[\langle\sigma_i\rangle]$ , under both free and fixed boundary conditions [25]. Furthermore, for these boundary conditions, the Griffiths inequalities on the Nishimori line enable us to prove that the correlation function exhibits translational invariance in the thermodynamic limit,  $\lim_{|\Lambda|\rightarrow\infty}\mathbb{E}[\langle\sigma_i\rangle] = \lim_{|\Lambda|\rightarrow\infty}\mathbb{E}[\langle\sigma_j\rangle]$ , similar to ferromagnetic systems [42]. Thus, by  $\sum_{j\sim i} = z$ , we obtain

$$\begin{aligned} m(\beta, h) &\leq \mathbb{E}[\tanh(\beta h_i)] + \sum_{j\sim i} 4\beta^2 m(\beta, h) \\ &= \mathbb{E}[\tanh(\beta h_i)] + 4z\beta^2 m(\beta, h), \end{aligned} \quad (24)$$

in the thermodynamic limit. This completes the proof of Theorem 1.

#### IV. DISCUSSIONS

We proved that the true critical inverse temperature on the Nishimori line is bounded by  $\beta_c^{NL} \geq \beta_{MF}^{NL}/2$ . Our proof is based on the proof of the mean-field bound for the critical temperature in the ferromagnetic Ising model [6], where the Griffiths inequalities [43–45] play an important role. Correspondingly, the key component of our proof is the Griffiths inequalities on the Nishimori line [25], which enables us to address the randomness in spin glass models.

Our bound is a weak correspondence of the mean-field bound for the critical temperature in ferromagnetic spin models. It remains an important future problem to investigate whether the corresponding mean-field bound on the Nishimori line,  $\beta_c^{NL} \geq \beta_{MF}^{NL}$ , holds.

In this study, we only considered Ising spin glass models with Gaussian randomness on the Nishimori line. It would be interesting to investigate whether a corresponding mean-field bound

for the critical temperature holds for other types of randomness, such as the  $\pm J$  model, on the Nishimori line.

In ferromagnetic spin models, there exists a mean-field bound for the magnetization [9–13], which is stronger than the mean-field bound for the critical temperature. An interesting problem to be addressed in the future is to obtain a corresponding mean-field bound for the magnetization on the Nishimori line.

The authors are grateful to Hal Tasaki for useful comments. This study was supported by JSPS KAKENHI Grant Nos. 24K16973 and 23H01432. Our study received financial support from the Public\Private R&D Investment Strategic Expansion Program (PRISM) and programs for bridging the gap between R&D and Ideal society (Society 5.0) and Generating Economic and social value (BRIDGE) from the Cabinet Office.

- 
- [1] I. A. Kvasnikov, Application of variational principle to the Ising model of ferromagnetism, Dokl. Akad. Nauk SSSR **110**, 755 (1956).
  - [2] N. N. Bogoliubov, On a variational principle in the many-body problem, Dokl. Akad. Nauk SSSR **119**, 244 (1958).
  - [3] R. B. Griffiths, A Proof that the Free Energy of a Spin System is Extensive, J. M. Phys. **5**, 1215 (1964).
  - [4] R. B. Griffiths, Correlations in Ising ferromagnets. III, Commun. Math. Phys. **6**, 121 (1967).
  - [5] M. E. Fisher, Critical Temperatures of Anisotropic Ising Lattices. II. General Upper Bounds, Phys. Rev. **162**, 480 (1967).
  - [6] W. Driessler, L. Landau, and J. F. Perez, Estimates of Critical Lengths and Critical Temperatures for Classical and Quantum Lattice Systems, J. Stat. Phys. **20**, 123 (1979).
  - [7] B. Simon, Mean Field Upper Bound on the Transition Temperature in Multicomponent Ferromagnets, J. Stat. Phys. **22**, 491 (1980).
  - [8] A. D. Sokal, Mean-Field Bounds and Correlation Inequalities, J. Stat. Phys. **28**, 431 (1982).
  - [9] C. J. Thompson, Upper bounds for Ising model correlation functions, Commun. Math. Phys. **24**, 61 (1971).
  - [10] P. A. Pearce, Mean-field bounds on the magnetization for ferromagnetic spin models, J. Stat. Phys. **25**, 309 (1981).
  - [11] C. M. Newman, Shock Waves and Mean Field Bounds (announcement of results, University of Ari-

- zona, 1981).
- [12] J. Slawny, On the mean field theory bound on the magnetization, *J. Stat. Phys.* **32**, 375 (1983).
  - [13] H. Tasaki and T. Hara, Mean field bound and GHS inequality, *J. Stat. Phys.* **35**, 99 (1984).
  - [14] H. Nishimori, Internal Energy, Specific Heat and Correlation Function of the Bond-Random Ising Model, *Prog. Theor. Phys.* **66**, 1169 (1981).
  - [15] H. Nishimori, *Statistical Physics of Spin Glasses and Information Processing: An Introduction*, (Oxford University Press, Oxford, 2001).
  - [16] M. Okuyama and M. Ohzeki, Gibbs–Bogoliubov inequality on the Nishimori line, *J. Phys. Soc. Jpn.* **92**, 084002 (2023).
  - [17] H. Nishimori, Exact results and critical properties of the Ising model with competing interactions, *J. Phys. C: Solid State Phys.* **13**, 4071 (1980).
  - [18] A. Georges, D. Hansel, P. Le Doussal, and J.-P. Bouchaud Exact properties of spin glasses: II. Nishimori’s line: new results and physical implications, *J. Phys. France* **46**, 1827 (1985).
  - [19] Y. Ozeki and H. Nishimori, Phase diagram of gauge glasses, *J. Phys. A: Math. Gen.* **26**, 3399 (1993).
  - [20] Y. Ozeki, Gauge transformation for dynamical systems of Ising spin glasses, *J. Phys. A: Math. Gen.* **28**, 3645 (1995).
  - [21] Y. Ozeki, The aging relation for Ising spin glasses, *J. Phys.: Condens. Matter* **9**, 11171 (1997).
  - [22] Y. Iba, The Nishimori line and Bayesian statistics, *J. Phys. A* **32**, 3875 (1998).
  - [23] I. A. Gruzberg, N. Read, and A. W. W. Ludwig, Random-bond Ising model in two dimensions: the Nishimori line and supersymmetry, *Phys. Rev. B* **63**, 104422 (2001).
  - [24] H. Nishimori and D. Sherrington, Absence of replica symmetry breaking in a region of the phase diagram of the Ising spin glass, In: *AIP Conference Proceedings*, vol. 553, p. 67 (2001).
  - [25] S. Morita, H. Nishimori, and P. Contucci, Griffiths inequalities for the Gaussian spin glass, *J. Phys. A* **37**, L203 (2004).
  - [26] S. Morita, Y. Ozeki, and H. Nishimori, Gauge theory for quantum spin glasses, *J. Phys. Soc. Jpn.* **75**, 014001 (2006).
  - [27] H. Kitatani, Griffiths Inequalities for Ising Spin Glasses on the Nishimori Line, *J. Phys. Soc. Jpn.* **78**, 044714 (2009).
  - [28] M. Ohzeki, Locations of multicritical points for spin glasses on regular lattices, *Phys. Rev. E* **79**, 021129 (2009).
  - [29] M. Ohzeki and H. Nishimori, Nonequilibrium Relations for Spin Glasses with Gauge Symmetry, *J.*



- Phys. Soc. Jpn. **79**, 084003 (2010).
- [30] M. Ohzeki, Fluctuation theorems on the Nishimori line, Phys. Rev. E **86**, 061110 (2012).
- [31] S. B. Korada and N. Macris, Exact Solution of the Gauge Symmetric p-Spin Glass Model on a Complete Graph, J. Stat. Phys. **136**, 205 (2009).
- [32] J. Barbier and N. Macris, The adaptive interpolation method for proving replica formulas. Applications to the Curie-Weiss and Wigner spike models, J. Phys. A **52**, 294002 (2019).
- [33] J. Barbier and N. Macris, The adaptive interpolation method: A simple scheme to prove replica formulas in Bayesian inference, Probab. Theory Relat. Fields **174**, 1133 (2019).
- [34] D. Alberici, F. Camilli, P. Contucci, and E. Mingione, The Solution of the Deep Boltzmann Machine on the Nishimori Line, Comm. Math. Phys. **387**, 1191 (2021).
- [35] D. Alberici, F. Camilli, P. Contucci, and E. Mingione, The Multi-species Mean-Field Spin-Glass on the Nishimori Line, J. Stat. Phys. **182**, 2 (2021).
- [36] C. Garban and T. Spencer, Continuous symmetry breaking along the Nishimori line, J. Math. Phys. **63**, 093302 (2022).
- [37] H. Nishimori, Analyticity of the energy in an Ising spin glass with correlated disorder, J. Phys. A: Math. Theor. **55**, 045001 (2022).
- [38] M. Okuyama and M. Ohzeki, Mean-field theory is exact for Ising spin glass models with Kac potential in non-additive limit on Nishimori line, J. Phys. A: Math. Theor. **56**, 325003 (2023).
- [39] C. Itoi and Y. Sakamoto, Gauge theory for mixed p-spin glasses, J. Phys. A: Math. Theor. **56**, 255002 (2023).
- [40] C. Itoi and Y. Sakamoto, Boundedness of Susceptibility in Spin Glass Transition of Transverse Field Mixed p-spin Glass Models, J. Phys. Soc. Jpn. **92**, 064004 (2023).
- [41] F. Camilli, P. Contucci, and E. Mingione, Central limit theorem for the overlaps on the Nishimori line, arXiv:2305.19943.
- [42] S. Friedli and Y. Velenik, *Statistical Mechanics of Lattice Systems: A Concrete Mathematical Introduction* (Cambridge University Press, Cambridge, United Kingdom, 2017).
- [43] R. B. Griffiths, Correlations in Ising Ferromagnets. I, J. Math. Phys. **8**, 478 (1967).
- [44] D. G. Kelly and S. Sherman, General Griffiths' inequalities on correlations in Ising ferromagnets, J. Math. Phys. **9**, 466 (1968).
- [45] J. Ginibre, General formulation of Griffiths' inequalities, Comm. Math. Phys. **16**, 310 (1970).