

Using the Haken-Strobl-Reineker Model to Determine the Temperature Dependence of the Diffusion Coefficient

William Barford*

*Department of Chemistry, Physical and Theoretical Chemistry Laboratory,
University of Oxford, Oxford, OX1 3QZ, United Kingdom*

E-mail: william.barford@chem.ox.ac.uk

Abstract

Stochastic quantum Liouville equations (SQLE) are widely used to model energy and charge dynamics in molecular systems. The Haken-Strobl-Reineker (HSR) SQLE is a particular paradigm in which the dynamical noise that destroys quantum coherences arises from a white noise (i.e., constant-frequency) spectrum. A system subject to the HSR SQLE thus evolves to its ‘high-temperature’ limit, whereby all the eigenstates are equally populated. This result would seem to imply that the predictions of the HSR model, e.g., the temperature dependence of the diffusion coefficient, have no validity for temperatures lower than the particle bandwidth. The purpose of this paper is to show that this assumption is incorrect for translationally invariant systems. In particular, provided that the diffusion coefficient is determined via the mean-squared-displacement, considerations about detailed-balance are irrelevant. Consequently, the high-temperature prediction for the temperature dependence of the diffusion coefficient may be extrapolated to lower temperatures, provided that the bath remains classical. Thus, for diagonal dynamical disorder the long-time diffusion coefficient, $D_\infty(T) = c_1/T$, while for both diagonal and off-diagonal disorder, $D_\infty(T) = c_1/T + c_2T$, where $c_2 \ll c_1$. An

appendix discusses an alternative interpretation from the HSR model of the ‘quantum to classical’ dynamics transition, whereby the dynamics is described as stochastically punctuated coherent motion.

1. Introduction

Coherent exciton dynamics in static, ordered molecular systems was described by Merrifield¹ in 1958. Assuming an exciton created at time $t = 0$ on a monomer, say $n = 0$, he showed that the subsequent wavefunction is $\Psi_n(t) = J_n(2\beta t)$, where J_n is the n th order Bessel function of the first-kind and β is the intermonomer exciton transfer integral. The wavefunction (illustrated in Fig. 2) spreads ballistically with a constant speed and a mean-squared-displacement (MSD) increasing quadratically with time.

As Merrifield observed,¹ however, dynamics on a static, ordered system is an idealization. Various physical processes, e.g., exciton-phonon coupling, and static and dynamic disorder destroy the coherent motion, eventually causing incoherent (or diffusive) motion where the MSD increases linearly with time. This topic now has a long and rich history, with many reviews describing the state of the field.²⁻⁶

The purpose of this paper is to expand on one particular rich area of investigation, namely the role of thermally-induced noise in destroying quantum coherences. A notable paradigm in this subject is the so-called HSR stochastic quantum Liouville equation (SQLE), developed and investigated by Haken, Strobl and Reineker.^{2,7} This equation was developed from the underlying time-dependent Schrödinger equation (TDSE) assuming that the dynamical fluctuations obey a white-noise spectrum, i.e., a constant power spectrum (or an Ohmic spectral function). Many important results have been derived from this model.^{2,8,9} In particular, it describes the ‘quantum to classical’ transition, in which exciton dynamics exhibits a crossover from ballistic to diffusive as a result of the noise destroying the coherent motion.

As stated, the HSR model assumes a white-noise spectrum, which implies that quantum tran-

sitions can occur between any pair of system energy eigenstates. This in turn implies that a system subject to the HSR SQLE evolves to its ‘high-temperature’ limit, whereby all the eigenstates are equally populated. This result would seem to suggest that the predictions of the HSR model, e.g., the temperature dependence of the diffusion coefficient, have no validity for temperatures lower than the particle bandwidth. The purpose of this paper is to show that this assumption is incorrect for translationally invariant systems. In particular, provided that the diffusion coefficient is determined via the mean-squared-displacement, considerations about detailed-balance are irrelevant. Consequently, the high-temperature prediction for the temperature dependence of the diffusion coefficient may be extrapolated to lower temperatures, provided that the bath remains classical.

This key result will be proved in Section 2. Since the HSR QSLE predictions for the diffusion coefficient in translationally invariant systems are valid for temperatures lower than the particle bandwidth, the predictions of the underlying stochastic TDSE are also equally valid. We use this realization to reinterpret the ‘quantum to classical’ transition as stochastically punctuated coherent motion. This is described in Appendix A.

Unlike the use of the MSD, Appendix B shows that the velocity autocorrelation function cannot be used to extrapolate the HSR predictions to temperatures lower than the bandwidth. Finally, Appendix C derives an expression for the temperature-dependence of the dephasing rate and Appendix D contains some details of the computational techniques.

2. Theory

2.1 Model of Exciton Dynamics in Linear Molecular Systems

We formulate the problem in terms of Frenkel exciton dynamics in one-dimensional molecular systems, e.g., J-aggregates or conjugated polymers. However, the analysis applies equally to triplet excitons and charges.

The total Hamiltonian is

$$\hat{H} = \hat{H}_S + \hat{H}_{SB} + \hat{H}_B, \quad (1)$$

where \hat{H}_S , \hat{H}_{SB} and \hat{H}_B are the system, system-bath and bath Hamiltonians, respectively. \hat{H}_B is defined in Appendix C (eqn (47)).

The system Hamiltonian is defined as

$$\hat{H}_S = \alpha \sum_{m=1}^N |m\rangle\langle m| + \beta \sum_{m=1}^N (|m+1\rangle\langle m| + |m\rangle\langle m+1|). \quad (2)$$

The ket $|m\rangle$ represents an exciton on monomer m , denoting a ‘site’, where N is the number of sites. α and β are the onsite potential and nearest-neighbor exciton transfer integral, respectively.

For a system with translational invariance, the eigenstates of \hat{H}_S are the Bloch states

$$|a\rangle = \frac{1}{\sqrt{N}} \sum_{m=1}^N \exp(ik_a m), \quad (3)$$

with eigenvalues $E_a = \alpha + 2\beta \cos k_a$, where $k_a = 2\pi a/N$ is the wavevector. The quantum numbers that label the eigenstates satisfy $1 \leq a \leq N$. The particle bandwidth in one-dimension is $4|\beta|$.

\hat{H}_{SB} is the system-bath Hamiltonian

$$\hat{H}_{SB} = \sum_m \delta\alpha_m(t) |m\rangle\langle m| + \sum_m \delta\beta_m(t) (|m+1\rangle\langle m| + |m\rangle\langle m+1|), \quad (4)$$

where $\delta\alpha_m(t)$ and $\delta\beta_m(t)$ represent dynamical fluctuations in α and β . These fluctuations are assumed to be uncorrelated in space, i.e.,

$$\langle \delta\alpha_m(t) \delta\alpha_n(0) \rangle = C_\alpha(t) = \sigma_\alpha^2 \exp(-t/\tau) \delta_{mn} \quad (5)$$

and

$$\langle \delta\beta_m(t) \delta\beta_n(0) \rangle = C_\beta(t) = \sigma_\beta^2 \exp(-t/\tau) \delta_{mn}. \quad (6)$$

In addition, in the white noise limit, defined by $\sigma_\times \tau \ll 1$, the bath correlation functions $C_\times(t) \rightarrow 2\gamma_\times \hbar^2 \delta(t)$, where $\gamma_\times = \sigma_\times^2 \tau / \hbar^2$ and \times indicates α or β . White noise means that the power spectrum, $I(\omega)$, is constant (or the spectral function, $J(\omega) \sim \omega I(\omega) \sim \omega$, i.e., Ohmic), which implies

that the time-dependent part of the Hamiltonian induces transitions between all pairs of eigenstates.

For a classical, harmonic bath, $\gamma_{\times} \propto \sigma_{\times}^2 \propto k_B T$. More specifically, as shown in Appendix C, for linear system-bath coupling

$$\gamma_{\times} = \pi k_B T E_{\times}^r / \hbar^2 \omega_c, \quad (7)$$

where E_{\times}^r is the reorganization energy arising from the system-bath coupling and ω_c is a high-frequency cut-off for the spectral function.

2.2 Determining the Temperature-Dependent Diffusion Coefficient

The one-dimensional thermal diffusion coefficient as a function of time is defined as

$$\langle D(t, T) \rangle = \frac{1}{2} \frac{d}{dt} \text{Tr} \{ \hat{\rho}(t, T) \hat{x}^2(t) \}, \quad (8)$$

where $\langle \dots \rangle$ indicates a thermal average, \hat{x} is the operator for the particle position, and $\hat{\rho}(t, T)$ is the system's reduced density operator. In the long-time limit the asymptotic diffusion coefficient is

$$\langle D_{\infty}(T) \rangle = \frac{1}{2} \frac{d}{dt} \text{Tr} \{ \hat{\rho}_0(T) \hat{x}^2(t) \} |_{\text{limit } t \rightarrow \infty}, \quad (9)$$

where

$$\hat{\rho}_0(T) = \frac{\exp(-\hat{H}_s/k_B T)}{\text{Tr} \{ \exp(-\hat{H}_s/k_B T) \}} \quad (10)$$

is the equilibrium density operator.

Evaluating the trace over the eigenstate basis of \hat{H}_s , Eqn (9) becomes

$$\langle D_{\infty}(T) \rangle = \sum_a p_a(T) D_a, \quad (11)$$

where

$$D_a = \frac{1}{2} \frac{d}{dt} \langle a | \hat{x}^2(t) | a \rangle |_{\text{limit } t \rightarrow \infty} \quad (12)$$

and $p_a(T)$ is the Boltzmann factor.

The mean-squared-displacement of a particle prepared in an arbitrary state $|\psi\rangle$ at $t = 0$ is defined as

$$\langle \psi | x^2(t) | \psi \rangle = \sum_m \ell^2 m^2 \rho_{mm}(t), \quad (13)$$

where ℓ is the intermonomer separation and $\rho_{mm}(t)$ are the diagonal elements of the system density matrix in the site basis, $\{|m\rangle\}$. As described in Section 2.3, the density matrix, $\rho_{mn}(t)$, evolves according to an appropriate quantum Liouville equation with the initial condition $\rho_{mn}(0) = \langle m | \psi(0) \rangle \langle \psi(0) | n \rangle$.

Transforming to the eigenstate basis via a transformation matrix, \mathbf{S} , Eqn (13) becomes

$$\langle \psi | x^2(t) | \psi \rangle = \sum_m \ell^2 m^2 \sum_{a,b} S_{ma} \tilde{\rho}_{ab}(t) S_{bm}^{-1}, \quad (14)$$

where $\tilde{\rho}_{ab}$ is the density matrix in the eigenstate basis and $S_{ma} = \langle m | a \rangle$. For a system with translational invariance, the transformation matrix elements are the Bloch factors

$$S_{ma} = (S_{am}^{-1})^* = \frac{1}{\sqrt{N}} \exp(-ik_a m). \quad (15)$$

Splitting the double sum in Eqn (14) over a and b into the separate sums of $a = b$ and $a \neq b$, and using Eqn (15) we obtain

$$\langle x^2 \rangle = \frac{1}{N} \sum_m \ell^2 m^2 \left(1 + \sum_{a \neq b} \tilde{\rho}_{ab} \exp(i(k_b - k_a)m) \right), \quad (16)$$

where we have also used $\sum_a \tilde{\rho}_{aa} = 1$. The significance of this result is that it shows that for a translationally invariant system the diffusion coefficient of a particle prepared in the state $|\psi\rangle$ depends on the evolution of eigenstate coherences and not directly on the evolution of eigenstate populations. Moreover, as will be shown in Section 2.3, for a translationally invariant system, eigenstate

coherences are decoupled from eigenstate populations, and thus for such systems the diffusion coefficient is completely independent of eigenstate populations. This means that when evaluating Eqn (11) to determine $\langle D_\infty(T) \rangle$ enforcing detailed balance – or ensuring that eigenstate populations satisfy their thermal values – is unnecessary. Thus, $\langle D_\infty(T) \rangle$ only depends on temperature *parametrically* via the temperature-dependence of the dephasing rate, Eqn (7).

Finally, as shown in Section 2.3, the asymptotic diffusion coefficient is independent of the initial state. Denoting this asymptotic value by $D_\infty(T)$, setting $D_a = D_\infty(T), \forall a$, and using $\sum_a p_a = 1$, Eqn (11) becomes $\langle D_\infty(T) \rangle = D_\infty(T)$. Thus, our task now is to determine $D_\infty(T)$ for a given dephasing rate. This is achieved via the Haken-Strobl-Reineker model as described in the next section.

2.3 The Haken-Strobl-Reineker Model

Haken and Strobl showed that ensemble averages of observables determined via the stochastic TDSE may be evaluated by a SQLE.^{2,7} For the case of diagonal noise within the site basis, the SQLE reads

$$\frac{\partial \rho_{mn}}{\partial t} = -i(\beta/\hbar)(\rho_{m\pm 1, n} - \rho_{m, n\pm 1})\delta_{mn} - 2\gamma_\alpha \rho_{mn}(1 - \delta_{mn}). \quad (17)$$

Rotating to the eigenstate basis of \hat{H}_S via Eqn (15), the SQLE becomes for the populations,

$$\frac{\partial \tilde{\rho}_{aa}}{\partial t} = - \sum_{b \neq a} (k_{ab} \tilde{\rho}_{aa} - k_{ba} \tilde{\rho}_{bb}) \quad (18)$$

and for the coherences,

$$\frac{\partial \tilde{\rho}_{ab}}{\partial t} = - (i\omega_{ab} + 2\gamma_\alpha) \tilde{\rho}_{ab} + \frac{2\gamma_\alpha}{N} \sum_{c=1}^N \tilde{\rho}_{c, c+(b-a)}, \quad (19)$$

where $\omega_{ab} = (E_a - E_b)/\hbar$.

From Eqn (18) and Eqn (19) we note the following:

1. In the HSR model $k_{ab} = k_{ba} = 2\gamma_\alpha/N$, thus guaranteeing equal eigenstate populations in the long-time limit. In principle, one could impose detailed balance on the rates, thus ensuring that the populations equilibrate to their thermal values. However, as now shown, this is unnecessary.
2. The equations of motion for the populations and coherences are decoupled. This proves, by virtue of Eqn (16), that $\langle D_\infty(T) \rangle$ is independent of temperature-dependent populations and only depends on T parametrically via $\gamma_\alpha(T)$.
3. Eqn (18) is a single N -coupled equation, whereas Eqn (19) are $(N - 1) \times N$ -coupled equations. Thus, there are $N \times N$ -coupled equations in total to solve. A numerical solution is described in Appendix D.

3. Results

Haken, Reineker and co-workers derived an expression for $D(t)$ for an initial δ -function source.^{2,10} For diagonal noise this is

$$D(t) = \frac{\beta^2 \ell^2}{\hbar^2 \gamma_\alpha} (1 - \exp(-2\gamma_\alpha t)). \quad (20)$$

Reineker² also derived the asymptotic value, D_∞ , for arbitrary initial conditions. For diagonal noise

$$D_\infty = \beta^2 \ell^2 / \hbar^2 \gamma_\alpha. \quad (21)$$

These analytical predictions were confirmed by solving both the HSR SQLE and the stochastic TDSE numerically (as described in Appendix D). Fig. 1 shows the numerically evaluated $D(t)$ for different initial conditions, namely a δ -function, Gaussian and particle-in-a-box sources. The δ -function result satisfies Eqn (20), while for all other sources, having a larger initial mean-squared-size and thus a smaller initial mean-squared-speed, the diffusion coefficient increases more slowly with time. However, Eqn (21) is satisfied for all initial conditions.

For translationally invariant systems, the evolution of the populations and coherences are also




Figure 1: Computed $D(t)$ using Eqn (19) for various initial conditions and diagonal white noise, $\gamma_\alpha = 0.1\beta/\hbar$. For the δ -function source, $D(t)$ satisfies Eqn (20). The asymptotic values satisfy Eqn (21). The Gaussian wavefunctions are defined by Eqn (26). The particle-in-a-box wavefunctions are $\psi_n = (2/(L+1))^{1/2} \sin(\pi n/(L+1))$. The diffusion coefficient at short times for the Gaussian wavepackets is superlinear, as also reported in ref.¹¹

decoupled in the presence of off-diagonal noise. Thus, for both diagonal and off-diagonal noise, the ‘high-temperature’ limit determined from the HSR SQLE can be extrapolated to temperatures lower than the bandwidth. Reineker² showed that for arbitrary initial conditions¹²

$$D_\infty = \frac{\beta^2 \ell^2}{\hbar^2 (\gamma_\alpha + 3\gamma_\beta)} + 2\gamma_\beta \ell^2, \quad (22)$$

where γ_α and γ_β are the diagonal and off-diagonal dephasing rates, respectively.

Substituting the temperature-dependence of the dephasing rates given by Eqn (7), we thus have the following prediction for the temperature-dependence of the diffusion coefficient in the presence

of white noise:

$$D_\infty(T) = \frac{c_1}{T} + c_2 T, \quad (23)$$

where

$$c_1 = \frac{\beta^2 \hbar^2 \omega_c \ell^2}{\pi k_B (E_\alpha^r + 3E_\beta^r)} \quad (24)$$

and

$$c_2 = \frac{2\pi k_B E_\beta^r \ell^2}{\hbar^2 \omega_c}. \quad (25)$$

Equation (23) is valid for translationally invariant systems subject to white noise for all temperatures – including temperatures lower than the particle bandwidth, $4|\beta|$ – provided that the bath remains classical. It remains valid if uniform long-range couplings are included, although the coefficients c_1 and c_2 are altered.

4. Conclusions

This paper has shown that for translationally invariant systems the predictions of the HSR model for the thermal diffusion coefficient can be extrapolated to temperatures lower than the particle bandwidth, provided that the bath remains classical. The proof relies on the observation that for such systems the mean-squared-displacement is independent of eigenstate populations. Consequently, considerations about detailed balance are irrelevant, and thus the diffusion coefficient depends on temperature only parametrically via the dephasing rates. When diagonal disorder dominates, $D(T) \sim T^{-1}$. This ‘high-temperature’ limit for $D(T)$ is a common prediction in many theories of charge and energy transport.¹³

Translationally invariant systems subject to white noise are an idealization of more realistic systems, where static disorder, correlated (non-Markovian) noise and electron-phonon interactions causing polaron formation are all important processes that will modify the predictions of this paper. For such systems, numerical solutions of the TDSE or QLE should explicitly ensure that detailed balance and stationarity are maintained during the system’s evolution (e.g., the Redfield quantum

Liouville equation of motion,^{4,14–16} the time-dependent wavepacket diffusion method,^{17,18} hierarchical equations for open systems,¹⁹ and stochastic Liouville equation methods²⁰). The recently proposed MASH surface-hopping schemes^{21,22} are also promising techniques for such simulations.

We conclude by noting that in the presence of static, diagonal disorder the scaling of $D(T)$ with temperature deviates from the HSR prediction.^{8,9,23} In particular, for $\gamma \sim T < \beta$, $D(T) \propto T$. This increase of the diffusion coefficient with temperature at low temperatures, sometimes known as environment-assisted quantum transport,⁸ is a consequence of the thermal fluctuations destroying the Anderson localization arising from coherent superposition of the particle wavefunction.

Appendix A. The Quantum to Classical Transition

In free space the one-dimensional TDSE reads

$$\frac{\partial \Psi(x,t)}{\partial t} = \frac{i\hbar}{2m} \frac{\partial^2 \Psi(x,t)}{\partial x^2}, \quad (26)$$

where m is the particle mass. As is well known,^{15,24} a particle prepared in a Gaussian wavepacket at $t = 0$, i.e.,

$$\Psi(x,0) = \left(\frac{1}{2\pi\sigma_x^2(0)} \right)^{1/4} \exp(-x^2/4\sigma_x^2(0)), \quad (27)$$

undergoes a ballistic (or coherent) spread. Thus, its MSD, $\sigma_x^2(t)$, increases as

$$\sigma_x^2(t) = \sigma_x^2(0) + \frac{\hbar^2 t^2}{4m\sigma_x^2(0)}. \quad (28)$$

In contrast, the diffusion equation describes the evolution of the density of an ensemble of particles, $\rho(x,t)$. In free space the one-dimensional diffusion equation reads

$$\frac{\partial \rho(x,t)}{\partial t} = D \frac{\partial^2 \rho(x,t)}{\partial x^2}. \quad (29)$$

Although this equation has precisely the same mathematical form as Eqn (26), the absence of imaginary i means that its physical predictions are quite different. In particular, an initially localized distribution of particles spreads diffusively, i.e.,

$$\rho(x,t) = \left(\frac{1}{2\pi\sigma_x^2(t)} \right)^{1/2} \exp(-x^2/2\sigma_x^2(t)), \quad (30)$$

where the MSD increases as

$$\sigma_x^2(t) = \sigma_x^2(0) + 2Dt. \quad (31)$$

The HSR equation describes how white noise causes a quantum particle's coherent motion to become incoherent. As shown by Reineker,² assuming a δ -function source on a lattice, i.e.,

$$|\Psi_n(0)|^2 = \delta_{n0}, \text{ for } t \ll 2\gamma,$$

$$|\Psi_n(t)|^2 = J_n^2(2\beta t), \quad (32)$$

(as predicted by the TDSE on a lattice¹) where J_n is the n th-order Bessel function of the first kind.

In contrast, for $t \gg 2\gamma$,

$$|\Psi_n(t)|^2 = \left(\frac{1}{4\pi Dt}\right)^{1/2} \exp(-n^2/4Dt), \quad (33)$$

(as predicted by the diffusion equation) where $D = \sigma_x^2(t)/2t = \beta^2/\gamma_\alpha$ for diagonal noise. In general for a δ -function source (where henceforth in this Appendix, we set $\hbar = \ell = 1$)^{2,25}

$$\sigma_x^2(t) = \frac{2\beta^2 t}{\gamma_\alpha} + \frac{\beta^2 t}{\gamma_\alpha^2} (\exp(-2\gamma_\alpha t) - 1). \quad (34)$$

In this interpretation, noise causes a crossover from coherent to incoherent transport. A different interpretation of the ‘quantum to classical’ transition, however, is afforded by the Lindblad formulation of the quantum Liouville equation. As we now show, in this interpretation the particle’s dynamics can be viewed as stochastically punctuated coherent motion.

1. The Lindblad dissipator is¹⁶

$$\hat{L}[\hat{\rho}(t)] = \Gamma \sum_m \left(\hat{A}_m \hat{\rho}(t) \hat{A}_m^\dagger - \frac{1}{2} \left(\hat{A}_m^\dagger \hat{A}_m \hat{\rho}(t) + \hat{\rho}(t) \hat{A}_m^\dagger \hat{A}_m \right) \right), \quad (35)$$

where \hat{A}_m is the Lindblad jump operator. For diagonal white-noise,⁸ $\hat{A}_m = |m\rangle\langle m|$ and $\Gamma = 2\gamma_\alpha$.

2. An ensemble of quantum trajectories reproduces the observables obtained via a QLE if each trajectory undergoes non-unitary evolution via an effective non-hermitian Hamiltonian,²⁶ defined by

$$\begin{aligned} \hat{H}_S \rightarrow \hat{H}_{\text{eff}} &= \hat{H}_S - \frac{i\Gamma}{2} \sum_m \hat{A}_m^\dagger \hat{A}_m \\ &= \hat{H}_S - \frac{i\Gamma}{2} \sum_m |m\rangle\langle m|. \end{aligned} \quad (36)$$

3. The simulation of a quantum trajectory then proceeds as follows:

(a) Given $|\Psi(t)\rangle$ at a time t , compute

$$|\Psi_{\text{trial}}\rangle = \exp(-i\hat{H}_{\text{eff}}\delta t)|\Psi(t)\rangle, \quad (37)$$

where δt is the time step. (See Appendix D for details.)

(b) Determine δp , defined via $\langle\Psi_{\text{trial}}|\Psi_{\text{trial}}\rangle = 1 - \delta p$.

(c) Then,

(i) with a probability $(1 - \delta p)$ define

$$|\Psi(t + \delta t)\rangle = |\Psi_{\text{trial}}\rangle / (1 - \delta p)^{1/2}. \quad (38)$$

In this case the wavefunction evolves according to \hat{H}_{eff} .

Or,

(ii) with a probability δp define

$$|\Psi(t + \delta t)\rangle = \frac{\hat{A}_m|\Psi(t)\rangle}{\langle\Psi(t)|\hat{A}_m^\dagger\hat{A}_m|\Psi(t)\rangle^{1/2}}. \quad (39)$$

Using the definition that $\hat{A}_m = |m\rangle\langle m|$, Eqn (39) implies that $|\Psi(t + \delta t)\rangle = |m\rangle$. Thus, in this case, the wavefunction collapses onto site m .

(d) If a quantum jump occurs in 3(c)(ii), the site m is chosen with a probability $P_m = |\Psi_m(t)|^2$.

Thus, the effect of the Lindblad jump operator in step is 3(c)(ii) to collapse the coherently evolving wavefunction onto the site m , after which it then again expands coherently until the next collapse. This process is illustrated in Fig. 2: At time $t = 0$ the particle is created on site $n_0 = 0$. Its wavefunction then evolves coherently until the first quantum jump at $t = t_1$. For $t \leq t_1$, $\Psi_n(t) = J_n(2\beta t)$, where J_n is the n th order Bessel function.¹ At $t = t_1$ the wavefunction collapses onto site

$n = n_1$, whereupon it again evolves coherently until a time t_2 when it again collapses onto site $n = n_2$.




Figure 2: An illustration of the evolution of a particle wavefunction subject to the TDSE with stochastic wavefunction collapses caused by white noise. The particle probability, $P_n(t) = J_n^2(2\beta t)$. The cumulative effect of the stochastic wavefunction collapses is to cause particle diffusion, where $D = \beta^2/\gamma_\alpha$, in exact agreement with the HSR model. $\gamma_\alpha = 0.1\beta/\hbar$. Time is in units of \hbar/β .

As these collapses are stochastic in space and time, their cumulative effective is to cause particle diffusion. We can determine the diffusion coefficient as follows. For a diffusive process the MSD is defined as

$$\text{MSD}(t) = \sigma_x^2(t) = N(t)\langle \ell^2 \rangle, \quad (40)$$

where $N(t) = t/\langle \tau \rangle$ is the number of jumps in a time t and $\langle \tau \rangle = 1/\Gamma = 1/2\gamma_\alpha$ is the average time interval between jumps. $\langle \ell^2 \rangle$ is the mean-squared jump size, determined by $\langle \ell^2 \rangle = \langle v^2 \tau^2 \rangle = v^2 \langle \tau^2 \rangle$, where v is the particle's coherent speed. For a δ -function source on a lattice, $v = \sqrt{2}\beta$. As determined by simulating the TDSE with Lindblad jump operators, for the dynamics described here, $\langle \tau^2 \rangle = 2\langle \tau \rangle^2 = 2/(2\gamma_\alpha)^2$. Hence, $\langle \ell^2 \rangle = \beta^2/\gamma_\alpha^2$ and therefore $D = N(t)\langle \ell^2 \rangle/2t = \beta^2/\gamma_\alpha$, thus –

on average – rederiving the prediction of the HSR quantum Liouville equation. Ensemble averaging over many particle trajectories will give precisely the same observables (e.g., average particle density) as the stochastic quantum Liouville approach, but the interpretation of the dynamics is different, namely stochastically punctuated coherent dynamics.

Appendix B. Using the Velocity Autocorrelation Function




Figure 3: The velocity autocorrelation function, $C(t) = \langle \psi | \hat{C}_v(t) | \psi \rangle$, where $\hat{C}_v(t) = \hat{v}(t)\hat{v}(0)$, computed via the stochastic TDSE with diagonal white noise ($\gamma_\alpha = 0.1\beta/\hbar$). $C(t)$ satisfies Eqn (43) for any initial condition. $C(0) = \langle \hat{v} \rangle_0$ is $2\beta^2\ell^2/\hbar^2$ for a δ -function, while it is $\beta^2/\hbar^2\sigma_x^2$ for a Gaussian wavepacket (defined in Eqn (27)), with σ_x in units of ℓ and \hat{v} is defined in Eqn (60).

An alternative definition to the thermal diffusion coefficient from Eqn (8) is

$$\langle D(t, T) \rangle = \int_0^t \text{Tr}\{\hat{\rho}(t', T)\hat{C}_v(t')\} dt', \quad (41)$$

where $\hat{C}_v(t) = \hat{v}(t)\hat{v}(0)$ is the velocity autocorrelation operator. The derivation of Eqn (41) from the identity

$$x^2(t) = \int_0^t \int_0^t v(t')v(t'')dt'dt'' \quad (42)$$

assumes *stationarity*, i.e., that the time-average of $v(t+t')v(t')$ is independent of t' .

However, if $\langle \psi | \hat{C}_v(t) | \psi \rangle$ is evaluated for an arbitrary state $|\psi\rangle$ via the HSR SQLE the stationarity condition is not met. This is because, unlike the case for the mean-squared-displacement, $\langle \psi | \hat{C}_v(t) | \psi \rangle$ depends on eigenstate populations, which become equally populated under evolution via the HSR SQLE.

Indeed, as shown in Fig. 3,

$$C(t) = \langle \psi | \hat{C}_v(t) | \psi \rangle = \langle \psi | \hat{C}_v(0) | \psi \rangle \exp(-2\gamma_\alpha t), \quad (43)$$

in agreement with the definition^{9,14}

$$\hat{v}(t) = \exp(i\hat{H}t/\hbar)\hat{v}\exp(-i\hat{H}t/\hbar)\exp(-2\gamma_\alpha t) \quad (44)$$

for the HSR model. Integrating Eqn (43) over time gives

$$D_\infty = \frac{\langle \psi | \hat{C}_v(0) | \psi \rangle}{2\gamma_\alpha}, \quad (45)$$

which erroneously predicts that the asymptotic diffusion coefficient depends on the initial condition.

Equation (45) is only correct for a δ -function source, because this maximally-localized in real-space wavefunction is maximally-delocalized in momentum-space, i.e., all the energy (momentum) eigenstates of a translationally invariant system are already equally populated at $t = 0$ so that the stationarity condition is automatically satisfied. Otherwise, for arbitrary initial conditions, eqn (41) is only valid provided detailed balance is enforced. Such a scenario is described in ref.¹⁴

Appendix C. The Dephasing Rate

For linear system-bath coupling, the system-bath Hamiltonian is¹⁵

$$\hat{H}_{SB} = \sum_m |m\rangle\langle m| \sum_j c_{jm}^\alpha u_j + \sum_m (|m\rangle\langle m+1| + |m+1\rangle\langle m|) \sum_j c_{jm}^\beta u_j, \quad (46)$$

while the bath Hamiltonian of harmonic oscillators is¹⁵

$$\hat{H}_B = \sum_j (\dot{u}_j^2 + \omega_j^2 u_j^2). \quad (47)$$

u_j are the mass-weighted oscillator displacements and c_j^\times are the weightings of each normal mode with associated spectral functions,

$$J_\times(\omega) = \frac{\pi}{2} \sum_j \frac{(c_j^\times)^2}{\omega_j} \delta(\omega - \omega_j). \quad (48)$$

The bath autocorrelation function associated with diagonal noise is $C_\alpha(t) = \langle \delta\alpha(t) \delta\alpha(0) \rangle$, where $\delta\alpha(t) = \sum_j c_j^\alpha u_j(t)$ is the dynamical fluctuation of the on-site potential energy. For a classical, harmonic bath and linear system-bath coupling¹⁵

$$C_\alpha(t) = \frac{2k_B T}{\pi} \int_0^\infty d\omega \frac{J_\alpha(\omega)}{\omega} \cos \omega t, \quad (49)$$

The spectral function, $J_\alpha(\omega)$, for an Ohmic bath with a high-frequency cut-off, ω_c is defined as

$$J_\alpha(\omega) = \left(\frac{\pi E_\alpha^r \omega}{\omega_c} \right) \exp(-\omega/\omega_c). \quad (50)$$

E_α^r is the reorganization energy of the bath modes, satisfying

$$E_\alpha^r = \frac{1}{\pi} \int_0^\infty d\omega \frac{J_\alpha(\omega)}{\omega}. \quad (51)$$

Thus, using

$$\begin{aligned} \int_0^\infty d\omega \exp(-\omega/\omega_c) \cos \omega t &= \frac{\omega_c}{1 + (\omega_c t)^2} \\ &\rightarrow \pi \delta(t), \end{aligned} \quad (52)$$

as $\omega_c \rightarrow \infty$, we obtain

$$C_\alpha(t) = \left(\frac{2\pi k_B T E_\alpha^r}{\omega_c} \right) \delta(t). \quad (53)$$

Equating Eqn (53) with the definition of γ_α given in Section 2.1, i.e., $C_\alpha(t) = 2\gamma_\alpha \hbar^2 \delta(t)$, yields an expression for the dephasing rate which is linearly proportional to temperature:

$$\gamma_\alpha = \frac{\pi k_B T E_\alpha^r}{\hbar^2 \omega_c}. \quad (54)$$

Appendix D. Computational Methods

D.1 Numerical Solution of the HSR Stochastic Quantum Liouville Equation

Since the equation of motion for the coherences in the eigenstate basis, Eqn (19), are block diagonal with respect to the quantum number c , there are $(N-1) \times N$ -coupled equations. These can be cast into the general form

$$\frac{dP_i}{dt} = \sum_{j=1}^N L_{ij} P_j, \quad (55)$$

for $1 \leq i \leq (N-1)$, where \mathbf{L} is the coupling matrix. The solution from linear algebra is

$$P_i(t) = \sum_{jk} S_{ij} \exp(\lambda_j t) S_{jk}^{-1} P_k(0), \quad (56)$$

where \mathbf{S} is the matrix whose columns are the eigenvectors of \mathbf{L} , $\{\lambda\}$ are the corresponding eigenvalues and $P_k(0)$ is an initial condition. To produce the results illustrated in Fig. 1, the complex

matrix \mathbf{L} was diagonalized via the LAPACK routine ZGEEV, and the complex matrix \mathbf{S} was inverted via the LAPACK routines ZGETRF and ZGETRI.

D.2 Numerical Solution of the Time-Dependent Schrodinger Equation

In general, given $\Psi(t)$ we require

$$\Psi(t + \delta t) = \exp(-i(H_S + H_{SB}(t))\delta t/\hbar)\Psi(t). \quad (57)$$

This is easily accomplished via the *Short Iterative Lanczos Propagator* method,²⁴ whereby a Krylov space of N vectors is generated via the Lanczos method. Diagonalizing the tridiagonal Hamiltonian, $\mathbf{H}_{\text{Lanczos}}$, within the Krylov space yields

$$\Psi(t + \delta t) = \mathbf{S} \cdot \exp(-i\mathbf{D}\delta t/\hbar) \cdot \mathbf{S}^\dagger \Psi(t), \quad (58)$$

where \mathbf{S} is the matrix whose columns are the eigenvectors of $\mathbf{H}_{\text{Lanczos}}$, the diagonal elements of \mathbf{D} are its eigenvalues, and $\Psi(t)$ is the first vector in the Krylov space.

The expectation value of the velocity autocorrelation function is

$$\begin{aligned} \langle \Psi | \hat{C}_v(t) | \Psi \rangle &= \langle \Psi | \hat{v}(t) \hat{v}(0) | \Psi \rangle \\ &= \langle \Psi(t) | \hat{v} | \Phi(t) \rangle, \end{aligned} \quad (59)$$

where $|\Phi\rangle = \hat{v}|\Psi\rangle$ and the velocity operator on a lattice is

$$\hat{v} = i(\ell\beta/\hbar) \sum_m (|m\rangle\langle m+1| - |m+1\rangle\langle m|). \quad (60)$$

References

- (1) Merrifield, R. E. Propagation of Electronic Excitation in Insulating Crystals. *Journal of Chemical Physics* **1958**, *28*, 647–650.
- (2) Kenkre, V. M.; Reineker, P. *Exciton dynamics in molecular crystals and aggregates*; Springer-Verlag: Berlin; New York, 1982.
- (3) Silinsh, E.; Capek, V. *Exciton dynamics in molecular crystals and aggregates*; American Institute of Physics: New York, 1994.
- (4) May, V.; Kühn, O. *Charge and energy transfer dynamics in molecular systems*; Wiley-VCH: Weinheim, 2011.
- (5) Dimitriev, O. P. Dynamics of Excitons in Conjugated Molecules and Organic Semiconductor Systems. *Chemical Reviews* **2022**, *122*, 8487–8593.
- (6) Barford, W. Exciton dynamics in conjugated polymer systems. *Frontiers in Physics* **2022**, *10*:1004042.
- (7) Haken, H.; Strobl, G. Exactly Solvable Model for Coherent and Incoherent Exciton Motion. *Zeitschrift Fur Physik* **1973**, *262*.
- (8) Rebentrost, P.; Mohseni, M.; Kassal, I.; Lloyd, S.; Aspuru-Guzik, A. Environment-assisted quantum transport. *New Journal of Physics* **2009**, *11*, 033003.
- (9) Kunsel, T.; Jansen, T. L. C.; Knoester, J. Scaling relations of exciton diffusion in linear aggregates with static and dynamic disorder. *Journal of Chemical Physics* **2021**, *155*, 134305.
- (10) Schwarzer, E.; Haken, H. Moments of Coupled Coherent and Incoherent Motion of Excitons. *Physics Letters A* **1972**, *A 42*, 317–318.
- (11) Tutunnikov, I.; Chuang, C. R.; Cao, J. S. Coherent Spatial Control of Wave Packet Dynamics on Quantum Lattices. *Journal of Physical Chemistry Letters* **2023**, *14*, 11632–11639.

- (12) Equation (22) was confirmed by numerical solutions of the stochastic TDSE.
- (13) Fratini, S.; Mayou, D.; Ciuchi, S. The Transient Localization Scenario for Charge Transport in Crystalline Organic Materials. *Advanced Functional Materials* **2016**, *26*, 2292–2315.
- (14) Chuang, C.; Lee, C. K.; Moix, J. M.; Knoester, J.; Cao, J. S. Quantum Diffusion on Molecular Tubes: Universal Scaling of the 1D to 2D Transition. *Physical Review Letters* **2016**, *116*, 196803.
- (15) Nitzan, A. *Chemical dynamics in condensed phases: relaxation, transfer and reactions in condensed molecular systems*; Oxford University Press: Oxford; New York, 2006.
- (16) Breuer, H.-P.; Petruccione, F. *The theory of open quantum systems*; Oxford University Press: Oxford; New York, 2002.
- (17) Zhong, X. X.; Zhao, Y. Non-Markovian stochastic Schrodinger equation at finite temperatures for charge carrier dynamics in organic crystals. *Journal of Chemical Physics* **2013**, *138*, 014111.
- (18) Han, L.; Ke, Y. L.; Zhong, X. X.; Zhao, Y. Time-Dependent Wavepacket Diffusion Method and its Applications in Organic Semiconductors. *International Journal of Quantum Chemistry* **2015**, *115*, 578–588.
- (19) Suess, D.; Strunz, W. T.; Eisfeld, A. Hierarchical Equations for Open System Dynamics in Fermionic and Bosonic Environments. *Journal of Statistical Physics* **2015**, *159*, 1408–1423.
- (20) Hsieh, C. Y.; Cao, J. S. A unified stochastic formulation of dissipative quantum dynamics. I. Generalized hierarchical equations. *Journal of Chemical Physics* **2018**, *148*, 014103.
- (21) Mannouch, J. R.; Richardson, J. O. A mapping approach to surface hopping. *Journal of Chemical Physics* **2023**, *158*, 104111.
- (22) Runeson, J. E.; Manolopoulos, D. E. A multi-state mapping approach to surface hopping. *Journal of Chemical Physics* **2023**, *159*, 094115.

- (23) Moix, J. M.; Khasin, M.; Cao, J. S. Coherent quantum transport in disordered systems: I. The influence of dephasing on the transport properties and absorption spectra on one-dimensional systems. *New Journal of Physics* **2013**, *15*, 085010.
- (24) Tannor, D. J. *Introduction to quantum mechanics: a time-dependent perspective*; University Science Books: Sausalito, Calif., 2007.
- (25) Likewise, the quantum to classical transition for an initial Gaussian wavefunction may be inferred by time-integrating the diffusion coefficient for the Gaussian initial state, as shown in Fig. 1, to obtain MSD(t).
- (26) Daley, A. J. Quantum trajectories and open many-body quantum systems. *Advances in Physics* **2014**, *63*, 77–149.