

COHOMOLOGICAL HALL ALGEBRAS FOR 3-CALABI-YAU CATEGORIES

TASUKI KINJO, HYEONJUN PARK, AND PAVEL SAFRONOV

ABSTRACT. The aim of this paper is to construct the cohomological Hall algebras for 3-Calabi–Yau categories admitting a strong orientation data. This can be regarded as a mathematical definition of the algebra of BPS states, whose existence was first mathematically conjectured by Kontsevich and Soibelman. Along the way, we prove Joyce’s conjecture on the functorial behaviour of the Donaldson–Thomas perverse sheaves for the attractor Lagrangian correspondence of (-1) -shifted symplectic stacks. This result allows us to construct a parabolic induction map for cohomological Donaldson–Thomas invariants of G -local systems on 3-manifolds for a reductive group G , which can be regarded as a 3-manifold analogue of the Eisenstein series functor in the geometric Langlands program.

CONTENTS

1. Introduction	1
2. Determinant line bundles	6
3. Shifted symplectic and d-critical structures	13
4. Donaldson–Thomas perverse sheaves	29
5. Stack of graded/filtered objects	35
6. Index line bundles and localization of orientations	48
7. Proof of the main theorem	72
8. Applications to the moduli of objects in 3CY categories	86
9. Applications to cohomological DT invariants of 3-manifolds	103
Appendix A. Braden’s theorem for stacks	116
References	117

1. INTRODUCTION

Cohomological Hall algebras. As a model of the algebra of BPS states in supersymmetric QFTs [HM98] Kontsevich and Soibelman [KS11] defined a cohomological Hall algebra (CoHA) associated to quivers with potentials which we briefly recall now. Let Q be a quiver and W a potential, which is a formal sum of cyclic paths in the quiver. Let \mathfrak{M}_Q be the moduli space of representations of Q which admits a decomposition $\mathfrak{M}_Q = \coprod_{\alpha \in \mathbb{N}^{Q_0}} \mathfrak{M}_{Q,\alpha}$ by the dimension vectors. Each $\mathfrak{M}_{Q,\alpha}$ is a quotient stack of a vector space by a product of general linear groups. The potential W defines a function $f_W: \mathfrak{M}_Q \rightarrow \mathbb{C}$ given by the trace of W in a given representation, and so we can consider the hypercohomology

$$\mathcal{H}_{Q,W,\alpha} = H^\bullet(\mathfrak{M}_{Q,\alpha}, \varphi_{f_W}(\mathbb{Q}_{\mathfrak{M}_{Q,\alpha}}[\dim \mathfrak{M}_{Q,\alpha}]))$$

of the sheaf of vanishing cycles with respect to f_W .

Let $\mathfrak{M}_Q^{2\text{-filt}}$ be the moduli space parametrizing short exact sequences $(0 \rightarrow E_1 \rightarrow E_2 \rightarrow E_3 \rightarrow 0)$ of representations of Q . We obtain a correspondence

$$\begin{array}{ccc}
 & \mathfrak{M}_Q^{2\text{-filt}} & \\
 \swarrow \text{ev}_1 \times \text{ev}_3 & & \searrow \text{ev}_2 \\
 \mathfrak{M}_Q \times \mathfrak{M}_Q & & \mathfrak{M}_Q
 \end{array}$$

of moduli stacks, where $ev_i: \mathfrak{M}_Q^{2\text{-filt}} \rightarrow \mathfrak{M}_Q$ is given by extracting E_i . If we denote by $f_W \boxplus f_W: \mathfrak{M}_Q \times \mathfrak{M}_Q \rightarrow \mathbb{C}$ the sum of the functions on the two factors, then we have $(ev_1 \times ev_3)^*(f_W \boxplus f_W) = ev_2^* f_W$. Moreover, $ev_2: \mathfrak{M}_Q^{2\text{-filt}} \rightarrow \mathfrak{M}_Q$ is proper on each connected component, so this correspondence defines a multiplication

$$(1.1) \quad \mathcal{H}_{Q,W,\alpha} \otimes \mathcal{H}_{Q,W,\beta} \longrightarrow \mathcal{H}_{Q,W,\alpha+\beta}[\chi_Q(\alpha, \beta) - \chi_Q(\beta, \alpha)],$$

where $\chi_Q(-, -)$ is the Euler form on the category of finite-dimensional representations of Q .

Given a quiver with potential (Q, W) , one can construct the corresponding Ginzburg dg algebra $\Gamma_3(Q, W)$ [Gin06] so that its stable dg category $\mathcal{C} = \text{Mod}_{\Gamma_3(Q,W)}^{\text{Perf}}$ of perfect modules carries a 3-Calabi–Yau structure. It was conjectured in [KS11] that one can generalize the construction of the CoHA to other 3-Calabi–Yau dg categories \mathcal{C} . For this one has to solve the following two problems: 1) define the space $\mathcal{H}_{\mathcal{C}}$, the *cohomological Donaldson–Thomas invariant* of \mathcal{C} , and 2) define the multiplication $\mathcal{H}_{\mathcal{C}} \otimes \mathcal{H}_{\mathcal{C}} \rightarrow \mathcal{H}_{\mathcal{C}}$.

The first problem was solved in [Bra+15; Ben+15; KL16]. Namely, if \mathcal{C} is a 3-Calabi–Yau category of finite type, its derived moduli stack of representations $\mathfrak{M}_{\mathcal{C}}$ in the sense of [TV07] carries a (-1) -shifted symplectic structure [BD21]. Consider an open substack $\mathfrak{M} \subset \mathfrak{M}_{\mathcal{C}}$ which is 1-Artin. Given an orientation of \mathfrak{M} , i.e. a square root of the canonical bundle $K_{\mathfrak{M}}|_{\mathfrak{M}}$, one can define a perverse sheaf $\varphi_{\mathfrak{M}} \in \text{Perv}(\mathfrak{M})$ locally modelled on the sheaf of vanishing cycles. Set $\Gamma = \pi_0(\mathfrak{M})$ and define the cohomological DT invariant

$$\mathcal{H} = \bigoplus_{\alpha \in \Gamma} H^{\bullet}(\mathfrak{M}_{\alpha}, \varphi_{\mathfrak{M}_{\alpha}}).$$

The monoid Γ carries an antisymmetric Euler pairing $\chi: \Gamma^2 \rightarrow \mathbb{Z}$.

The goal of this paper is to construct a multiplication on the above space. The following statement combines Corollary 8.8, Proposition 8.29 and Proposition 8.37.

Theorem A. *Consider the following data:*

- \mathcal{C} is a stable dg category of finite type equipped with a 3-Calabi–Yau structure.
- $\mathfrak{M} \subset \mathfrak{M}_{\mathcal{C}}$ is an open substack closed under direct sums and containing the zero module. Moreover, assume that the underlying classical stack \mathfrak{M}^{cl} is a quasi-separated Artin stack with affine stabilizers which is Θ -reductive (see [Hal14, Definition 5.1.1]), such that the inclusion of the zero module $\text{pt} \rightarrow \mathfrak{M}^{\text{cl}}$ is a closed immersion.
- Strong orientation data for \mathfrak{M} , i.e., an orientation on the (-1) -shifted symplectic stack \mathfrak{M} compatible with direct sums and satisfying associativity (see Definition 8.4).

Then there is an associative multiplication

$$\mathcal{H}_{\mathcal{C},\alpha} \otimes \mathcal{H}_{\mathcal{C},\beta} \longrightarrow \mathcal{H}_{\mathcal{C},\alpha+\beta}[\chi(\alpha, \beta)].$$

In the case $\mathcal{C} = \text{Mod}_{\Gamma_3(Q,W)}^{\text{Perf}}$ is the 3-Calabi–Yau category associated to a quiver with potential (Q, W) , this multiplication reduces to (1.1).

Before we explain the construction of the multiplication, let us list some examples of 3CY categories equipped with strong orientation data. In particular, one can apply our construction of the multiplication to define a CoHA for such categories.

- Given a smooth dg category \mathcal{D} , one may consider its deformed 3-Calabi–Yau completion \mathcal{C} . For instance, \mathcal{C} can be given by the dg derived category of coherent sheaves on local curves or surfaces [KM24] or $\mathcal{C} = \text{Mod}_{\Gamma_3(Q,W)}^{\text{Perf}}$ for a quiver with potential. Then $\mathfrak{M} = \mathfrak{M}_{\mathcal{C}}$ carries a strong orientation data by Proposition 8.34.
- Given a closed spin 3-manifold M , consider $\mathcal{C} = \text{LocSys}(M)^{\omega}$ the stable dg category of complexes of local systems on M . Let $\mathfrak{M} \subset \mathfrak{M}_{\mathcal{C}}$ be the substack of local systems of vector spaces. Then \mathfrak{M} carries a strong orientation data by Proposition 9.14.

For instance, let S be a smooth quasi-projective surface. Let \mathcal{C} be the dg category of perfect complexes on the Calabi–Yau threefold $\text{Tot}(K_S)$ and $\mathfrak{M} \subset \mathfrak{M}_{\mathcal{C}}$ the substack of compactly supported coherent sheaves. Combining the results of [BCS24; Kin22] we get $\mathcal{H}_{\mathcal{C}} \cong H_{\text{vd}(\text{Coh}(S))-\bullet}^{\text{BM}}(\text{Coh}(S))$, where $\text{Coh}(S)$ is the stack of compactly supported coherent sheaves on S . We expect that in this case, the cohomological Hall algebra structure we construct in the paper reduces to the one constructed in [KV23].

Note that for any compact Calabi–Yau threefold X the paper [JU21] constructs an orientation on the moduli stack of coherent sheaves on X compatible with direct sums. However, it is not known at present if

the compatibility isomorphisms can be chosen to be associative. Thus, using our results we obtain a CoHA multiplication for any compact Calabi–Yau threefold, but we know the associativity only up to some sign.

Hyperbolic localization. The construction of the CoHA multiplication goes through an intermediate step of independent interest. Given a smooth algebraic space X with a \mathbb{G}_m -action μ and a \mathbb{G}_m -equivariant function $f: X \rightarrow \mathbb{C}$ consider the *attractor correspondence*

$$\begin{array}{ccc} & X^{\mu,+} & \\ \text{gr}_\mu \swarrow & & \searrow \text{ev}_\mu \\ X^\mu & & X, \end{array}$$

where $X^\mu \subset X$ is the subscheme of fixed points and $X^{\mu,+}$ is the subscheme of points $x \in X$ such that the limit $\lim_{t \rightarrow 0} \mu(t)x$ exists. The map ev_μ is the obvious inclusion and $\text{gr}_\mu: X^{\mu,+} \rightarrow X^\mu$ is given by taking the limit. Using a theorem of Braden [Bra03; DG14] one can show (see [Nak17, Proposition 5.4.1] and [Des22, Proposition 3.3] as well as Proposition 7.8) that the functor of vanishing cycles φ_f commutes with the hyperbolic localization functor $(\text{gr}_\mu)_* \text{ev}_\mu^!$. In particular, if we denote by $I_X^\mu: X^\mu \rightarrow \mathbb{Z}$ the locally-constant function given by assigning to a fixed point $x \in X$ its index, i.e. the difference $\text{rank}(T_{X,x}^+) - \text{rank}(T_{X,x}^-)$ of the ranks of positive and negative weight subspaces of the tangent space, then

$$\varphi_{f|_{X^\mu}}(\mathbb{Q}_{X^\mu}[\dim X^\mu])[I_X^\mu] \cong (\text{gr}_\mu)_* \text{ev}_\mu^! \varphi_f(\mathbb{Q}_X[\dim X]).$$

The construction of the sheaf of vanishing cycles is globalized in [Bra+15; Ben+15] to general (-1) -shifted symplectic stacks: given a (-1) -shifted symplectic stack (X, ω_X) equipped with an orientation o , there is a perverse sheaf $\varphi_{X, \omega_X, o} \in \text{Perv}(X)$. Correspondingly, the hyperbolic localization statement was extended to (-1) -shifted symplectic algebraic spaces (and more generally to d-critical algebraic spaces) in [Des22, Theorem 4.2] (see also Theorem 7.9) as follows. Given a (-1) -shifted symplectic algebraic space (X, ω_X) equipped with an orientation o and a \mathbb{G}_m -action preserving the (-1) -shifted symplectic structure, let $u: X^\mu \rightarrow X$ be the inclusion of the fixed points. Then one can pull back the (-1) -shifted symplectic structure and the orientation to X^μ (see Lemma 3.19), so that there is an isomorphism

$$\varphi_{X^\mu, u^* \omega_X, u^* o}[I_{X^\mu}^{\text{cl}}] \cong (\text{gr}_\mu)_* \text{ev}_\mu^! \varphi_{X, \omega_X, o}.$$

Our second main result is a construction of the hyperbolic localization isomorphism for (-1) -shifted symplectic Artin stacks equipped with an orientation. The attractor correspondence for a \mathbb{G}_m -action on a stack \mathfrak{X} can be extracted from the attractor correspondence for the *trivial* \mathbb{G}_m -action on the quotient stack $[\mathfrak{X}/\mathbb{G}_m]$. So, from now on we restrict to the case of trivial \mathbb{G}_m -actions, so that the attractor correspondence becomes

$$(1.2) \quad \begin{array}{ccc} & \text{Filt}(\mathfrak{X}) & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \text{Grad}(\mathfrak{X}) & & \mathfrak{X}, \end{array}$$

where $\text{Grad}(\mathfrak{X}) = \text{Map}(B\mathbb{G}_m, \mathfrak{X})$ and $\text{Filt}(\mathfrak{X}) = \text{Map}([\mathbb{A}^1/\mathbb{G}_m], \mathfrak{X})$.

First of all, we show in Corollary 5.18 that if \mathfrak{X} carries a k -shifted symplectic structure, then the attractor correspondence (1.2) carries a natural k -shifted Lagrangian structure. For instance, when \mathfrak{X} is the moduli space of perfect complexes on an elliptic curve, the induced 0-shifted Poisson structure on $\text{Filt}(\mathfrak{X})$ is related to the Feigin–Odesskii Poisson structure [HP18].

Next, there is a locally-constant function $I_{\mathfrak{X}}: \text{Grad}(\mathfrak{X}) \rightarrow \mathbb{Z}$ given by the index of the corresponding point. Let $u: \text{Grad}(\mathfrak{X}) \rightarrow \mathfrak{X}$ be the map obtained by evaluation at the basepoint of $B\mathbb{G}_m$. The following is Corollary 7.19:

Theorem B. *Let \mathfrak{X} be a quasi-separated derived Artin stack with affine stabilizers equipped with a (-1) -shifted symplectic structure $\omega_{\mathfrak{X}}$ and an orientation o . Then there is a natural isomorphism*

$$\varphi_{\text{Grad}(\mathfrak{X}), u^* \omega_{\mathfrak{X}}, u^* o}[I_{\mathfrak{X}}^{\text{cl}}] \cong \text{gr}_* \text{ev}^! \varphi_{\mathfrak{X}, \omega_{\mathfrak{X}}, o}.$$

We call this isomorphism the *integral isomorphism* since it can be regarded as a categorification of the Kontsevich–Soibelman’s motivic integral identity conjecture [KS08, §4.4] proved later by Lê [Thu15]. The motivic integral identity was very recently generalized by Bu [Bu24] to the attractor correspondence of (-1) -shifted symplectic stacks with good moduli spaces.

Remark. While we were preparing this manuscript, Pierre Descombes let us know that he is preparing a revision of his paper [Des22] to include a statement similar to Theorem B.

The attractor correspondence reduces to several other important correspondences in particular cases. For instance, let \mathcal{C} be a dg category of finite type equipped with a 3-Calabi–Yau structure. Then by Proposition 8.26 the correspondence (1.2) for $\mathfrak{X} = \mathfrak{M}_{\mathcal{C}}$, the derived moduli stack of objects, is the union of the correspondences

$$\begin{array}{ccc} & \mathfrak{M}_{\mathcal{C}}^{n-\text{filt}} & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \mathfrak{M}_{\mathcal{C}}^n & & \mathfrak{M}_{\mathcal{C}}, \end{array}$$

where $\mathfrak{M}_{\mathcal{C}}^{n-\text{filt}}$ parametrizes sequences $x_{n-1} \rightarrow \cdots \rightarrow x_0$ and $\text{gr}: \mathfrak{M}_{\mathcal{C}}^{n-\text{filt}} \rightarrow \mathfrak{M}_{\mathcal{C}}^n$ is given by taking the cofibres of the maps $x_k \rightarrow x_{k-1}$. Moreover, by Proposition 8.28 the corresponding Lagrangian structure on the attractor correspondence reduces to the one constructed in [BD21] using relative Calabi–Yau structures. The hyperbolic localization isomorphism from Theorem B by adjunction gives rise to the multiplication on the cohomological Hall algebra in Theorem A.

Cohomological DT invariants of 3-manifolds. Let G be a connected reductive group and M a closed oriented 3-manifold. Then by Proposition 9.9 the correspondence (1.2) for $\mathfrak{X} = \text{Loc}_G(M)$, the derived character stack of M , is the union of the correspondences

$$\begin{array}{ccc} & \text{Loc}_P(M) & \\ \swarrow & & \searrow \\ \text{Loc}_L(M) & & \text{Loc}_G(M), \end{array}$$

where $P \subset G$ are parabolic subgroups and L their Levi factors. Moreover, the corresponding Lagrangian structure on the attractor correspondence reduces to the one constructed in [Saf17]. Given a spin structure on M we construct a natural orientation of this Lagrangian correspondence in Proposition 9.6 following [NS23], so the hyperbolic localization isomorphism from Theorem B gives rise to a parabolic induction map

$$\text{ind}_P: H^\bullet(\text{Loc}_L(M), \varphi_{\text{Loc}_L(M)}) \longrightarrow H^\bullet(\text{Loc}_G(M), \varphi_{\text{Loc}_G(M)}).$$

The hypercohomology groups $H^\bullet(\text{Loc}_G(M), \varphi_{\text{Loc}_G(M)})$ appear as invariants associated to 3-manifolds in the geometric Langlands program for generic quantum parameters. In particular, the parabolic induction map is a 3-dimensional analogue of the Eisenstein series functor for surfaces. There is a close relationship between these hypercohomology groups and skein modules of 3-manifolds, see [GS23, Conjecture D] and related hypercohomology groups for $G = \text{SL}_2$ were studied by Abouzaid and Manolescu [AM20].

In Theorem 9.15 we use the parabolic induction map to construct a cohomological Hall algebra structure on

$$\mathcal{H}^{\text{GL}} = \bigoplus_{n \geq 0} H^\bullet(\text{Loc}_{\text{GL}_n}(M), \varphi_{\text{Loc}_{\text{GL}_n}(M)})$$

and cohomological Hall \mathcal{H}^{GL} -module structures on

$$\mathcal{H}^{\text{SO}} = \bigoplus_{n \geq 0} H^\bullet(\text{Loc}_{\text{SO}_n}(M), \varphi_{\text{Loc}_{\text{SO}_n}(M)}), \quad \mathcal{H}^{\text{Sp}} = \bigoplus_{n \geq 0} H^\bullet(\text{Loc}_{\text{Sp}_{2n}}(M), \varphi_{\text{Loc}_{\text{Sp}_{2n}}(M)}).$$

We expect that these will be useful in establishing structural results for the cohomological DT invariants, e.g. their Langlands duality.

A conjecture of Joyce. Given an oriented Lagrangian correspondence between oriented (-1) -shifted symplectic stacks $\mathfrak{X}_1 \xleftarrow{f_1} \mathfrak{L} \xrightarrow{f_2} \mathfrak{X}_2$, Joyce has conjectured (see [JS19, Conjecture 1.1] and [AB17, Conjecture 5.22]) the existence of a natural map

$$f_1^* \varphi_{\mathfrak{X}_1}[\mathrm{vd} \mathfrak{L}] \longrightarrow f_2^! \varphi_{\mathfrak{X}_2}.$$

satisfying certain natural compatibilities. For instance, if $\mathfrak{X}_1 = \mathfrak{X}_2 = \mathrm{pt}$, it defines a virtual Lagrangian cycle $[\mathfrak{L}]_{\mathrm{vir}}^{\mathrm{Lag}} \in \mathrm{H}_{\mathrm{vd}(\mathfrak{L})}^{\mathrm{BM}}(\mathfrak{L})$ which should coincide with the one constructed in [BJ17; OT23; Pri18]. See [Kin24, §5] for a survey on this conjecture. One of the motivations for the conjecture of Joyce is to construct a cohomological Hall algebra structure.

If \mathfrak{X} is an oriented (-1) -shifted symplectic stack which is quasi-separated and has affine stabilizers, the hyperbolic localization isomorphism from Theorem B by adjunction provides a map

$$\mathrm{gr}^* \varphi_{\mathrm{Grad}(\mathfrak{X})}[\mathrm{vd} \mathrm{Filt}(\mathfrak{X})] \longrightarrow \mathrm{ev}^! \varphi_{\mathfrak{X}}$$

thus resolving the conjecture of Joyce in this case. Note, however, that Theorem B is stronger than the version of the conjecture for a general Lagrangian correspondence as it asserts that the adjoint map is an isomorphism.

Future works. The construction of the cohomological Hall algebras has striking applications in cohomological Donaldson–Thomas theory. For example, this can be used to formulate the cohomological integrality theorem for oriented (-1) -shifted symplectic stacks with symmetric weights generalizing [DM20, Theorem C] and further to prove it for the moduli space of objects in 3-Calabi–Yau categories with a choice of an orientation. This will be investigated by the first author and his collaborators in their future paper.

In joint work with Adeel Khan, the first author has constructed the cohomological Hall algebra for the 3-Calabi–Yau completion of a dg-category using a microlocal method [KK; KK23]. One advantage of the microlocal construction is that one can prove the compatibility with the Kapranov–Vasserot’s 2-dimensional cohomological Hall algebra [KV23] under the dimensional reduction isomorphism of the Donaldson–Thomas perverse sheaves [Kin22]. The first author plans to prove that the cohomological Hall algebra multiplication for 3-Calabi–Yau categories constructed in this paper coincides with the one in [KK] for 3-Calabi–Yau completion of dg-categories in his future paper.

Acknowledgement. We thank Chenjing Bu, Andrés Ibáñez Núñez, Ben Davison, Dominic Joyce and Yukinobu Toda for helpful discussions related to this paper. T.K. was supported by JSPS KAKENHI Grant Number 23K19007. H.P. was supported by Korea Institute for Advanced Study (SG089201).

Notation and Convention.

- We let \mathcal{S} denote the ∞ -category of spaces (see [Lur09, Definition 1.2.16.1]).
- By $\mathrm{cdga}^{\leq 0}$ we denote the ∞ -category of non-positively graded commutative differential graded \mathbb{C} -algebras (cdgas).
- All classical Artin stacks are assumed to be locally of finite type over the complex number field.
- All derived Artin stacks are assumed to be 1-Artin and locally of finite presentation over the complex number field.
- A derived stack that is n -geometric for some integer n and is locally of finite presentation is called a derived higher Artin stack.
- For an Artin \mathfrak{X} , we let $D_c^b(\mathfrak{X})$ denote the bounded derived category of sheaves with respect to the analytic topology in \mathbb{Q} -vector spaces on \mathfrak{X} with constructible cohomology. We let $\mathrm{Perv}(\mathfrak{X})$ denote the abelian category of perverse sheaves on \mathfrak{X} .
- Given a perfect complex E on a derived stack \mathfrak{X} , we denote by $\widehat{\det}(E) = \det(E|_{X^{\mathrm{red}}})$ the restriction of the determinant line bundle to the reduced stack.
- For a derived Artin stack \mathfrak{X} with a trivial \mathbb{G}_m -action and a \mathbb{G}_m -equivariant quasi-coherent complex $\mathcal{F} \in \mathrm{QCoh}(\mathfrak{X})$, we let \mathcal{F}^+ (resp. \mathcal{F}^-) denote the direct summand with positive (resp. negative) weights.
- For a smooth morphism $f: X_1 \rightarrow X_2$, we set $K_{X_1/X_2} := \widehat{\det}(\Omega_{X_1/X_2})$.
- If there is no confusion, we use expressions such as f_* , $f_!$, and $\mathcal{H}om$ for the derived functors Rf_* , $Rf_!$, and $R\mathcal{H}om$.

2. DETERMINANT LINE BUNDLES

The aim of this section is to discuss the construction and some properties of the determinant functor of perfect complexes following [KM76; STV15].

2.1. Algebraic K-theory of stable ∞ -categories. Here, we briefly recall some basic facts on algebraic K-theory of stable ∞ -categories: See [BGT13] for a detailed discussion.

Let \mathcal{C} be a small stable ∞ -category. The Grothendieck group $K_0(\mathcal{C})$ is defined as a group generated by elements $[E]$ labelled by objects $E \in \mathcal{C}$ and relations $[E_2] = [E_1] + [E_3]$ for fibre sequences $E_1 \rightarrow E_2 \rightarrow E_3$. One can define a spectral refinement of $K_0(\mathcal{C})$, i.e., a spectrum $K(\mathcal{C}) \in \mathrm{Sp}$ with a natural isomorphism $\pi_0(K(\mathcal{C})) \cong K_0(\mathcal{C})$, which is called the algebraic K-theory spectrum: See [BGT13, §9] for the detail.

Let $K^{\mathrm{cn}}(\mathcal{C})$ be the connective part of the algebraic K-theory spectrum. Then by the construction of $K^{\mathrm{cn}}(\mathcal{C})$ (see [BGT13, §7.1]), we have the following:

- For an object $E \in \mathcal{C}$, there exists a corresponding map

$$[E] \in \mathrm{Map}(\Delta^1, \Omega^{\infty-1}K^{\mathrm{cn}}(\mathcal{C}))$$

with homotopy $[E] \circ \delta^0 \simeq *$, $[E] \circ \delta^1 \simeq *$ where δ^i is the i -th face map and $*$ is the base point.

- For a fibre sequence $\Delta: E_1 \rightarrow E_2 \rightarrow E_3$ in \mathcal{C} , there exists a corresponding map

$$[\Delta] \in \mathrm{Map}(\Delta^2, \Omega^{\infty-1}K^{\mathrm{cn}}(\mathcal{C}))$$

with homotopy $[\Delta] \circ \delta^0 \simeq [E_3]$, $[\Delta] \circ \delta_1 \simeq [E_2]$, $[\Delta] \circ \delta_2 \simeq [E_1]$.

- For a fibre double sequence in \mathcal{C} , i.e., a fibre sequence in the ∞ -category of fibre sequences in \mathcal{C} depicted as

$$(I) \quad \begin{array}{ccccc} & & \Delta_{23} & & \Delta' \\ \Delta_{12}: E_1 & \longrightarrow & \dot{E}_2 & \longrightarrow & \dot{E}_{12} \\ & \parallel & \downarrow & & \downarrow \\ \Delta_{13}: E_1 & \longrightarrow & E_3 & \longrightarrow & E_{13} \\ & \downarrow & \downarrow & & \downarrow \\ & 0 & \longrightarrow & E_{23} & \longlongequal{\quad} E_{23}, \end{array}$$

there exists a corresponding map

$$[\Gamma] \in \mathrm{Map}(\Delta^3, \Omega^{\infty-1}K^{\mathrm{cn}}(\mathcal{C}))$$

with homotopy $[\Gamma] \circ \delta_0 \simeq [\Delta']$, $[\Gamma] \circ \delta_1 \simeq [\Delta_{23}]$, $[\Gamma] \circ \delta_2 \simeq [\Delta_{13}]$, $[\Gamma] \circ \delta_3 \simeq [\Delta_{12}]$.

In particular, for each $E \in \mathcal{C}$, there exists a corresponding point $[E] \in \Omega^{\infty}K^{\mathrm{cn}}(\mathcal{C})$ such that for a distinguished triangle $\Delta: E_1 \rightarrow E_2 \rightarrow E_3$, there exists a natural homotopy

$$(2.1) \quad I(\Delta): [E_1] \cdot [E_3] \sim [E_2]$$

and for a fibre double sequence (I), the following homotopy commutative diagram exists

$$(2.2) \quad \begin{array}{ccc} [E_1] \cdot [E_{12}] \cdot [E_{23}] & \xrightarrow[\sim]{I(\Delta_{12})} & [E_2] \cdot [E_{23}] \\ \sim \downarrow I(\Delta') & & \sim \downarrow I(\Delta_{23}) \\ [E_1] \cdot [E_{13}] & \xrightarrow[\sim]{I(\Delta_{13})} & [E_3]. \end{array}$$

2.2. Determinant functor. Here we briefly recall the notion of determinant line bundle of perfect complexes following [KM76; STV15]. Consider the following ∞ -categories for R a connective commutative dg \mathbb{C} -algebra:

- $\mathrm{Perf}(R)$ is the stable ∞ -category of perfect complexes. It has a symmetric monoidal structure given by the direct sum.
- $\mathrm{Vect}(R) \subset \mathrm{Perf}(R)$ is the subcategory of vector bundles. It inherits the symmetric monoidal structure.

- $\text{Pic}^{\text{gr}}(R)$ is the ∞ -groupoid of $\mathbb{Z}/2\mathbb{Z}$ -graded line bundles. It carries a symmetric monoidal structure given by

$$(L_1, \alpha_1) \otimes (L_2, \alpha_2) = (L_1 \otimes L_2, \alpha_1 + \alpha_2)$$

with the braiding defined using the Koszul sign rule, which we call the *standard symmetric monoidal structure*. It also carries another symmetric monoidal structure with the same monoidal product with the braiding of line bundle with no extra sign, which we call the *naive symmetric monoidal structure*. Otherwise stated, we use the standard symmetric monoidal structure.

- We define the *standard $\mathbb{Z}/2\mathbb{Z}$ -action* on $\text{Pic}^{\text{gr}}(R)$ by the monoidal dual with respect to the standard symmetric monoidal structure: $(L, \alpha) \mapsto (L, \alpha)^\vee$. We define the *naive $\mathbb{Z}/2\mathbb{Z}$ -action* on $\text{Pic}^{\text{gr}}(R)$ by the monoidal dual with respect to the naive monoidal structure: $(L, \alpha) \mapsto (L^\vee, \alpha)$. Otherwise stated, we use the standard $\mathbb{Z}/2\mathbb{Z}$ -action.
- We will fix an isomorphism $(L, \alpha)^\vee \cong (L^\vee, \alpha)$ so that the canonical trivialization $(L, \alpha) \otimes (L, \alpha)^\vee \cong (\mathcal{O}_R, 0)$ and $(L, \alpha) \otimes (L^\vee, \alpha) \cong (\mathcal{O}_R, 0)$ are identified. Under this identification, the natural isomorphism $(L_1^\vee, \alpha_1) \otimes (L_2^\vee, \alpha_2) \cong ((L_1 \otimes L_2)^\vee, \alpha_1 + \alpha_2)$ is given by the braiding of line bundles with the extra sign $(-1)^{\alpha_1 + \alpha_2}$ and the trivialization $(L^{\vee\vee}, \alpha) \cong (L, \alpha)$ is given with the intervention of the sign $(-1)^\alpha$.

The following is shown in [KM76].

Proposition 2.1. *Let R be a (discrete) commutative \mathbb{C} -algebra. Then there is a symmetric monoidal functor*

$$\det: \text{Vect}(R)^\simeq \longrightarrow \text{Pic}^{\text{gr}}(R),$$

natural in R and compatible with the $\mathbb{Z}/2\mathbb{Z}$ -actions on the left side given by passing to the dual vector bundle and on the right side the naive $\mathbb{Z}/2\mathbb{Z}$ -action.

As shown in [Hel20, Section 2] and [Elm+20, Appendix A] the derived stacks $\text{Vect}(-)$ and $\text{Pic}^{\text{gr}}(-)$ are left Kan extended from discrete commutative algebras, so we get a symmetric monoidal functor of ∞ -categories $\det: \text{Vect}(R)^\simeq \rightarrow \text{Pic}^{\text{gr}}(R)$ defined for any connective cdga R . Moreover, as $\text{Pic}^{\text{gr}}(R)$ is grouplike, it factors through algebraic K -theory:

$$\text{Vect}(R)^\simeq \longrightarrow K(\text{Vect}(R)) \longrightarrow \text{Pic}^{\text{gr}}(R).$$

Consider the commutative diagram

$$\begin{array}{ccc} \text{Perf}(R)^\simeq & \longrightarrow & K(\text{Perf}(R)) \\ \uparrow & & \uparrow \\ \text{Vect}(R)^\simeq & \longrightarrow & K(\text{Vect}(R)) \xrightarrow{\det} \text{Pic}^{\text{gr}}(R). \end{array}$$

The vertical map on the right is an equivalence by the results of [HS21, Corollary 8.1.3], [Hel20, Corollary 1.40] and [Fon18], so in total we obtain the **determinant map**

$$\det: \text{Perf}(R)^\simeq \longrightarrow \text{Pic}^{\text{gr}}(R).$$

For a derived stack \mathfrak{X} both $\text{Perf}(\mathfrak{X})$ and $\text{Pic}^{\text{gr}}(\mathfrak{X})$ are defined by a right Kan extension from their values on derived affine schemes, so we obtain the determinant map

$$\det: \text{Perf}(\mathfrak{X})^\simeq \longrightarrow \text{Pic}^{\text{gr}}(\mathfrak{X}).$$

We will often be interested in restrictions of determinant line bundles to reduced stacks, so for a derived stack \mathfrak{X} and a perfect complex $E \in \text{Perf}(\mathfrak{X})$ we introduce the notation

$$\widehat{\det}(E) := \det(E|_{\mathfrak{X}^{\text{red}}}).$$

We now prove some properties of the determinant functor and the natural transform (2.3) for perfect complexes on the derived Artin stack \mathfrak{X} . Firstly, for a fibre sequence $\Delta: E_1 \rightarrow E_2 \rightarrow E_3$, the homotopy $I(\Delta)$ in (2.1) induces a natural isomorphism

$$(2.3) \quad i(\Delta): \det(E_1) \otimes \det(E_3) \cong \det(E_2).$$

Assume that we are given a fibre double sequence as in (Γ) . Then the commutativity of the diagram (2.2) implies the commutativity of the following diagram

$$(2.4) \quad \begin{array}{ccc} \det(E_1) \otimes \det(E_{12}) \otimes \det(E_{23}) & \xrightarrow[\cong]{i(\Delta_{12})} & \det(E_2) \otimes \det(E_{23}) \\ \cong \downarrow i(\Delta') & & \cong \downarrow i(\Delta_{23}) \\ \det(E_1) \otimes \det(E_{13}) & \xrightarrow[\cong]{i(\Delta_{13})} & \det(E_3). \end{array}$$

For an object $E \in \text{Perf}(\mathfrak{X})$, consider the fibre sequence $\Delta_E: E \rightarrow 0 \rightarrow E[1]$ given by the rotation of $E \xrightarrow{\text{id}} E \rightarrow 0$. It induces an isomorphism

$$i(\Delta_E): \det(E) \otimes \det(E[1]) \cong \mathcal{O}_{\mathfrak{X}}.$$

We define a natural map

$$(2.5) \quad \theta_E: \det(E[1]) \cong \det(E)^\vee$$

so that the following diagram commutes:

$$\begin{array}{ccc} \det(E) \otimes \det(E[1]) & \xrightarrow[\cong]{i(\Delta_E)} & \mathcal{O}_{\mathfrak{X}} \\ \cong \downarrow (\text{id}, \theta_E) & \nearrow \cong & \\ \det(E) \otimes \det(E)^\vee & & \end{array}$$

For $n \in \mathbb{Z}_{>0}$, we inductively construct an isomorphism

$$\theta_{E,n}: \det(E[n]) \cong \det(E)^{\vee(n)}$$

where $(-)^{\vee(n)}$ denotes the n -th iterated dual, by the following composition

$$\det(E[n]) = \det(E[1][n-1]) \xrightarrow[\cong]{\theta_{E[1],n-1}} \det(E[1])^{\vee(n-1)} \xrightarrow[\cong]{\theta_E} \det(E)^{\vee(n)}.$$

Since the n -th iterated dual only depends on the parity of n up to the canonical isomorphism, we can define $(-)^{\vee(n)}$ for all integer $n \in \mathbb{Z}$. We define $\theta_{E,-n}: \det(E[-n]) \cong \det(E)^{\vee(-n)}$ by $(\theta_{E[-n],n})^{-1}$. By construction, for $n, m \in \mathbb{Z}$, the following diagram commutes:

$$\begin{array}{ccc} \det(E[n+m]) & \xrightarrow[\cong]{\theta_{E[n],m}} & \det(E[n])^{\vee(m)} \\ \parallel & & \cong \downarrow \theta_{E,n} \\ \det(E[n+m]) & \xrightarrow[\cong]{\theta_{E,n+m}} & \det(E)^{\vee(n+m)}. \end{array}$$

Also, it is clear from the construction that the following diagram commutes:

$$(2.6) \quad \begin{array}{ccc} \det(E[-1]) \otimes \det(E) & \xrightarrow[\cong]{\text{id} \otimes \theta_{E[-1]}} & \det(E[-1]) \otimes \det(E[-1])^\vee \\ \cong \downarrow \theta_{E,-1} \otimes \text{id} & & \cong \downarrow \\ \det(E)^\vee \otimes \det(E) & \xrightarrow[\cong]{} & \mathcal{O}_{\mathfrak{X}}. \end{array}$$

Lemma 2.2. *Let $\Delta: E_1 \xrightarrow{f} E_2 \xrightarrow{g} E_3$ be a fibre sequence in $\text{Perf}(\mathfrak{X})$. Let $\rho(\Delta): E_2 \xrightarrow{g} E_3 \xrightarrow{h} E_1[1]$ and $\rho^{-1}(\Delta): E_3[-1] \xrightarrow{h[-1]} E_1 \xrightarrow{f} E_2$ be rotated fibre sequences. Then the following diagrams commute:*

$$(2.7) \quad \begin{array}{ccc} \det(E_1) \otimes \det(E_3) \otimes \det(E_1[1]) & \xrightarrow[\cong]{i(\Delta) \otimes \text{id}} & \det(E_2) \otimes \det(E_1[1]) \\ \cong \downarrow \text{id} \otimes \theta_{E_1} & & \cong \downarrow i(\rho(\Delta)) \\ \det(E_1) \otimes \det(E_3) \otimes \det(E_1)^\vee & \xrightarrow[\cong]{} & \det(E_3), \end{array}$$

$$(2.8) \quad \begin{array}{ccc} \det(E_1) \otimes \det(E_2) \otimes \det(E_1[1]) & \xrightarrow[\cong]{\text{id} \otimes i(\rho(\Delta))} & \det(E_1) \otimes \det(E_3) \\ \cong \downarrow \text{id} \otimes \theta_{E_1} & & \cong \downarrow (-1)^{\text{rank } E_1} \cdot i(\Delta) \\ \det(E_1) \otimes \det(E_2) \otimes \det(E_1)^\vee & \xrightarrow{\cong} & \det(E_2), \end{array}$$

$$(2.9) \quad \begin{array}{ccc} \det(E_3[-1]) \otimes \det(E_1) \otimes \det(E_3) & \xrightarrow[\cong]{\text{id} \otimes i(\Delta)} & \det(E_3[-1]) \otimes \det(E_2) \\ \cong \downarrow \theta_{E_3, -1} \otimes \text{id} & & \cong \downarrow (-1)^{\text{rank } E_3} \cdot i(\rho^{-1}(\Delta)) \\ \det(E_3)^\vee \otimes \det(E_1) \otimes \det(E_3) & \xrightarrow{\cong} & \det(E_1). \end{array}$$

Proof. Consider the following fibre double sequences:

$$\begin{array}{ccccc} E_1 & \longrightarrow & 0 & \longrightarrow & E_1[1] \\ \parallel & & \downarrow & & \downarrow \begin{pmatrix} 0 \\ \text{id} \end{pmatrix} \\ E_1 & \xrightarrow{0} & E_3 & \xrightarrow{\begin{pmatrix} \text{id} \\ 0 \end{pmatrix}} & E_3 \oplus E_1[1] \\ \downarrow & & \parallel & & \downarrow (\text{id}, 0) \\ 0 & \longrightarrow & E_3 & \xlongequal{\quad} & E_3, \end{array} \quad \begin{array}{ccccc} E_1 & \xrightarrow{f} & E_2 & \xrightarrow{g} & E_3 \\ \parallel & & \downarrow g & & \downarrow \begin{pmatrix} \text{id} \\ -h \end{pmatrix} \\ E_1 & \xrightarrow{0} & E_3 & \xrightarrow{\begin{pmatrix} \text{id} \\ 0 \end{pmatrix}} & E_3 \oplus E_1[1] \\ \downarrow & & \downarrow h & & \downarrow (h, \text{id}) \\ 0 & \longrightarrow & E_1[1] & \xlongequal{\quad} & E_1[1]. \end{array}$$

Note that the horizontal middle fibre sequences are equivalent. By considering the commutative diagrams (2.4) associated with these diagrams and using the fact that the automorphism of the complex $E_3 \oplus E_1[1]$ given by $\begin{pmatrix} \text{id} & 0 \\ h & \text{id} \end{pmatrix}$ induces the identity map on the determinant line bundle, we conclude the commutativity of the diagram (2.7).

To prove the commutativity of the diagram (2.8), consider the following diagram:

$$\begin{array}{ccc} & & \det(E_1) \otimes \det(E_1) \otimes \det(E_1[1]) \otimes \det(E_3) \\ & \nearrow \text{id} \otimes \text{sw} & \cong \downarrow \text{id} \otimes \theta_{E_1} \otimes \text{id} \\ \det(E_1) \otimes \det(E_1) \otimes \det(E_3) \otimes \det(E_1[1]) & & \det(E_1) \otimes \det(E_1) \otimes \det(E_1)^\vee \otimes \det(E_3) \\ \cong \downarrow \text{id} \otimes i(\Delta) \otimes \text{id} & & \cong \downarrow \\ \det(E_1) \otimes \det(E_2) \otimes \det(E_1[1]) & \xrightarrow[\cong]{\text{id} \otimes i(\rho(\Delta))} & \det(E_1) \otimes \det(E_3) \\ \cong \downarrow \text{id} \otimes \theta_{E_1} & & \cong \downarrow (-1)^{\text{rank } E_1} \cdot i(\Delta) \\ \det(E_1) \otimes \det(E_2) \otimes \det(E_1)^\vee & \xrightarrow[\cong]{} & \det(E_2), \end{array}$$

where sw denotes the symmetrizer. The commutativity of the upper square follows from the commutativity of the diagram (2.7). The commutativity of the outer square follows from the fact that the symmetrizer for the object $\det(E_1) \otimes \det(E_1)$ is given by multiplication of $(-1)^{\text{rank } E_1}$. In particular, we conclude the commutativity of the diagram (2.8).

The commutativity of the diagram (2.9) immediately follows from the commutativity of (2.6) and (2.8). \square

Corollary 2.3. *Consider the following fibre double sequence:*

$$(2.10) \quad \begin{array}{ccccc} & & \Delta'_{23} & & \Delta'_{23} \\ & & \dot{E}'_{23} & \xlongequal{\quad} & \dot{E}'_{23} \\ & 0 & \longrightarrow & & \\ \Delta_{12} : & E_1 & \longrightarrow & E_2 & \longrightarrow & E_{12} \\ & \parallel & & \downarrow & & \downarrow \\ \Delta_{13} : & E_1 & \longrightarrow & E_3 & \longrightarrow & E_{13}. \end{array}$$

Then the following diagram commutes:

$$(2.11) \quad \begin{array}{ccc} \det(E'_{23}) \otimes \det(E_1) \otimes \det(E_{13}) & \xrightarrow[\cong]{i(\Delta')} & \det(E_1) \otimes \det(E_{12}) \\ i(\Delta_{13}) \downarrow \cong & & \cong \downarrow i(\Delta_{12}) \\ \det(E'_{23}) \otimes \det(E_3) & \xrightarrow[\cong]{i(\Delta_{23})} & \det(E_2). \end{array}$$

Proof. Consider the following diagram:

$$\begin{array}{ccc} \det(E'_{23}) \otimes \det(E_1) \otimes \det(E_{13}) & \xrightarrow[\cong]{i(\Delta')} & \det(E_1) \otimes \det(E_{12}) \\ \downarrow i(\Delta_{13}) \cong & \swarrow i(\rho(\Delta')) \cong & \uparrow \cong (-1)^{\text{rank } E'_{23} \cdot \theta_{E'_{23}}} \\ \det(E'_{23}) \otimes \det(E_1) \otimes \det(E_{12}) \otimes \det(E'_{23}[1]) & & \downarrow \cong i(\Delta_{12}) \\ \downarrow i(\Delta_{13}) \cong & \swarrow i(\rho(\Delta_{23})) \cong & \downarrow \cong (-1)^{\text{rank } E'_{23} \cdot \theta_{E'_{23}}} \\ \det(E'_{23}) \otimes \det(E_3) & \xrightarrow[\cong]{i(\Delta_{23})} & \det(E_2). \end{array}$$

The upper and lower triangles commute by the commutativity of the diagram (2.8). The middle square commutes by (2.4). Hence, we conclude. \square

The following statement can be thought of as an ∞ -categorical refinement of [KM76, Theorem 1]:

Lemma 2.4. *Let \mathfrak{X} be a derived stack. Consider the fibre double sequence in $\text{Perf}(\mathfrak{X})$:*

$$\begin{array}{ccccc} & \Delta'_1 & & \Delta'_2 & & \Delta'_3 \\ \Delta_1 : & \dot{E}'_{11} & \longrightarrow & \dot{E}'_{12} & \longrightarrow & \dot{E}'_{13} \\ & \downarrow & & \downarrow & & \downarrow \\ \Delta_2 : & E_{21} & \longrightarrow & E_{22} & \longrightarrow & E_{23} \\ & \downarrow & & \downarrow & & \downarrow \\ \Delta_3 : & E_{31} & \longrightarrow & E_{32} & \longrightarrow & E_{33}. \end{array}$$

Then the following diagram commutes:

$$(2.12) \quad \begin{array}{ccc} (\det(E_{11}) \otimes \det(E_{13})) \otimes (\det(E_{31}) \otimes \det(E_{33})) & \xrightarrow[\simeq]{i(\Delta_1) \otimes i(\Delta_3)} & \det(E_{12}) \otimes \det(E_{32}) \\ \downarrow \simeq & & \simeq \downarrow i(\Delta'_2) \\ & & \det(E_{22}) \\ & & \simeq \uparrow i(\Delta_2) \\ (\det(E_{11}) \otimes \det(E_{31})) \otimes (\det(E_{13}) \otimes \det(E_{33})) & \xrightarrow[\simeq]{i(\Delta'_1) \otimes i(\Delta'_3)} & \det(E_{21}) \otimes \det(E_{23}). \end{array}$$

Proof. Let F be the cofibre of the map $E_{11} \rightarrow E_{22}$. Consider the following fibre double sequences:

$$(2.13) \quad \begin{array}{ccccc} E_{11} & \longrightarrow & E_{12} & \longrightarrow & E_{13} \\ \parallel & & \downarrow & & \downarrow \\ E_{11} & \longrightarrow & E_{22} & \longrightarrow & F \\ \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & E_{32} & \xlongequal{\quad} & E_{32} \end{array} \quad \begin{array}{ccccc} E_{11} & \longrightarrow & E_{21} & \longrightarrow & E_{31} \\ \parallel & & \downarrow & & \downarrow \\ E_{11} & \longrightarrow & E_{22} & \longrightarrow & F \\ \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & E_{23} & \xlongequal{\quad} & E_{23} \end{array} \quad \begin{array}{ccccc} 0 & \longrightarrow & E_{13} & \xlongequal{\quad} & E_{13} \\ \downarrow & & \downarrow & & \downarrow \\ E_{31} & \longrightarrow & F & \longrightarrow & E_{23} \\ \parallel & & \downarrow & & \downarrow \\ E_{31} & \longrightarrow & E_{32} & \longrightarrow & E_{33}. \end{array}$$

Consider the following diagram:

$$\begin{array}{ccccc} & & (\det(E_{11}) \otimes \det(E_{13})) & & \\ & & \otimes (\det(E_{31}) \otimes \det(E_{33})) & & \\ & \swarrow \cong & & \searrow \cong & \\ \det(E_{11}) \otimes (\det(E_{13}) \otimes \det(E_{32})) & \xrightarrow{\cong} & \det(E_{11}) \otimes \det(F) & \xleftarrow{\cong} & \det(E_{11}) \otimes (\det(E_{31}) \otimes \det(E_{23})) \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ \det(E_{12}) \otimes \det(E_{32}) & \xrightarrow{\cong} & \det(E_{22}) & \xleftarrow{\cong} & \det(E_{21}) \otimes \det(E_{23}). \end{array}$$

The left bottom (resp. right bottom) squares commutes by the commutativity of the diagram (2.4) applied to the first (resp. second) fibre double sequence in (2.13). Also, the upper triangle commutes by applying the commutativity of (2.11) to the third fibre double sequence in (2.13). Hence, we conclude. \square

Corollary 2.5. *Let $\Delta: E_1 \rightarrow E_2 \rightarrow E_3$ be a fibre sequence. Then the following diagram commutes:*

$$\begin{array}{ccc} \det(E_1[1]) \otimes \det(E_3[1]) & \xrightarrow[\cong]{i(\Delta)} & \det(E_2[1]) \\ \downarrow \cong \theta_{E_1} \otimes \theta_{E_3} & & \downarrow \cong \theta_{E_2} \\ \det(E_1)^\vee \otimes \det(E_3)^\vee & \xrightarrow[\cong]{i(\Delta[1])} & \det(E_2)^\vee. \end{array}$$

Proof. This follows by applying Lemma 2.4 to the following fibre double sequence:

$$\begin{array}{ccccc} E_1 & \longrightarrow & E_2 & \longrightarrow & E_3 \\ \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & 0 & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ E_1[1] & \longrightarrow & E_2[1] & \longrightarrow & E_3[1]. \end{array}$$

\square

Recall by construction that the natural map

$$K(\mathrm{Perf}(\mathfrak{X})) \rightarrow \mathrm{Pic}^{\mathrm{gr}}(\mathfrak{X})$$

is $\mathbb{Z}/2\mathbb{Z}$ -equivariant, where $\mathbb{Z}/2\mathbb{Z}$ -equivariant structure on $K(\mathrm{Perf}(\mathfrak{X}))$ is induced by the duality map

$$\mathrm{Perf}(\mathfrak{X}) \rightarrow \mathrm{Perf}(\mathfrak{X})^{\mathrm{op}}, \quad E \mapsto E^\vee$$

with the canonical identification $E \simeq E^{\vee\vee}$ and similarly for $\mathrm{Pic}^{\mathrm{gr}}(\mathfrak{X})$ (with respect to the naive monoidal structure). The $\mathbb{Z}/2\mathbb{Z}$ -equivariant structure gives a natural isomorphism

$$(2.14) \quad \iota_E: \det(E^\vee) \cong \det(E)^\vee.$$

By definition, the following diagram commutes:

$$(2.15) \quad \begin{array}{ccccc} \det(E) & \xrightarrow[\cong]{(-1)^{\mathrm{rank} E} \cdot \mathrm{id}} & \det(E) & & \\ \cong \downarrow & & & & \uparrow \cong \\ \det(E^{\vee\vee}) & \xrightarrow[\cong]{\iota_{E^\vee}} & \det(E^\vee)^\vee & \xrightarrow[\cong]{(\iota_{E^\vee})^\vee} & \det(E)^{\vee\vee}. \end{array}$$

The appearance of the sign intervention is due to the fact that the trivialization of the double dual with respect to the standard and naive monoidal structure differs by a sign.

Let $\Delta: E_1 \rightarrow E_2 \rightarrow E_3$ be a fibre sequence in $\mathrm{Perf}(\mathfrak{X})$ and $\Delta^\vee: E_3^\vee \rightarrow E_2^\vee \rightarrow E_1^\vee$ be the dual fibre sequence. By the construction of algebraic K-theory, we clearly have the following equivalence of 2-cells

$$\sigma([\Delta]) \simeq [\Delta^\vee]$$

where σ denotes the non-trivial element in $\mathbb{Z}/2\mathbb{Z}$. In particular, the following diagram commutes:

$$(2.16) \quad \begin{array}{ccc} \det(E_3^\vee) \otimes \det(E_1^\vee) & \xrightarrow[\cong]{i(\Delta^\vee)} & \det(E_2^\vee) \\ \cong \downarrow \iota_{E_3} \otimes \iota_{E_1} & & \cong \downarrow \iota_{E_2} \\ \det(E_3)^\vee \otimes \det(E_1)^\vee & \xrightarrow[\cong]{i(\Delta)^\vee} & \det(E_2)^\vee. \end{array}$$

Applying this to the fibre sequence $\Delta_E: E \rightarrow 0 \rightarrow E[1]$, we see that the following diagram commutes:

$$(2.17) \quad \begin{array}{ccc} \det(E^\vee[-1]) \otimes \det(E^\vee) & \xrightarrow[\cong]{i(\Delta_E^\vee)} & \mathcal{O}_{\mathfrak{X}} \\ \cong \downarrow & & \parallel \\ \det(E[1]^\vee) \otimes \det(E^\vee) & & \\ \cong \downarrow \iota_{E[1]} \otimes \iota_E & & \\ \det(E[1]^\vee) \otimes \det(E)^\vee & \xrightarrow[\cong]{i(\Delta_E)^\vee} & \mathcal{O}_{\mathfrak{X}}. \end{array}$$

Combining this with the commutativity of the diagram (2.6), we see that the following diagram commutes:

$$(2.18) \quad \begin{array}{ccccc} \det(E^\vee[-1]) & \xrightarrow[\cong]{} & \det(E[1]^\vee) & \xrightarrow[\cong]{\iota_{E[1]}} & \det(E[1])^\vee \\ \cong \downarrow \theta_{E^\vee, -1} & & & & \cong \downarrow \theta_E^\vee \\ \det(E^\vee)^\vee & \xrightarrow[\cong]{\iota_{E^\vee}^\vee} & \det(E)^\vee & \xrightarrow[\cong]{} & \det(E)^{\vee\vee}. \end{array}$$

By repeatedly using the commutativity of the above diagram, for a positive integer $n \in \mathbb{Z}_{>0}$, we see that the following diagram commutes:

$$(2.19) \quad \begin{array}{ccccc} \det(E^\vee[-n]) & \xrightarrow{\cong} & \det(E[n]^\vee) & \xrightarrow[\cong]{\iota_{E[n]}} & \det(E[n])^\vee \\ \cong \downarrow \theta_{E^\vee, -n} & & & & \cong \downarrow \theta_{E, n}^\vee \\ \det(E^\vee)^{\vee(n)} & \xrightarrow[\cong]{\iota_{E^\vee}^\vee} & & & \det(E)^\vee{}^{(n+1)}. \end{array}$$

By substituting $E = F[-n]$, we see that the same diagram commutes for negative integers n as well.

3. SHIFTED SYMPLECTIC AND D-CRITICAL STRUCTURES

We will recall the theory of shifted symplectic structures following [Pan+13] and d-critical structures following [Joy15] which can be regarded a classical shadow of (-1) -shifted symplectic structures. These structures are a key ingredient in the construction of the Donaldson–Thomas perverse sheaves, which we will recall in the subsequent section.

3.1. Shifted symplectic structures. Here we briefly recall the theory of shifted symplectic structures. See [Pan+13] for the original source, [PY24] for a quick survey, and [Cal15; AB17; CHS21] for the material on Lagrangian correspondences.

Let A be a connective commutative dg algebra. We define the graded mixed commutative dg algebra $\mathbf{DR}(A)$ of differential forms as in [Pan+13]. Its underlying graded commutative algebra is $\mathrm{Sym}(\mathbb{L}_A[-1])$. We consider the following related objects:

- The complex of closed p -forms

$$\Omega^{\geq p}(A) = \left(\prod_{n \geq p} \mathbf{DR}(A)(n), d + d_{\mathrm{dR}} \right) [p].$$

There are natural maps $\Omega^{\geq(p+1)}(A) \rightarrow \Omega^{\geq p}(A)[1]$.

- The space of n -shifted closed p -forms $\mathcal{A}^{p, \mathrm{cl}}(A, n) = \mathrm{Map}(k[-n], \Omega^{\geq p}(A))$. There are natural forgetful maps $\mathcal{A}^{p+1, \mathrm{cl}}(A, n-1) \rightarrow \mathcal{A}^{p, \mathrm{cl}}(A, n)$.
- The space of n -shifted exact p -forms $\mathcal{A}^{p, \mathrm{ex}}(A, n)$ which fits into a fibre sequence of pointed spaces

$$\mathcal{A}^{p, \mathrm{ex}}(A, n) \xrightarrow{d_{\mathrm{dR}}} \mathcal{A}^{p, \mathrm{cl}}(A, n) \longrightarrow \mathcal{A}^{0, \mathrm{cl}}(A, p+n).$$

There are natural forgetful maps $\mathcal{A}^{p+1, \mathrm{ex}}(A, n-1) \rightarrow \mathcal{A}^{p, \mathrm{ex}}(A, n)$.

- The space of n -shifted p -forms $\mathcal{A}^p(A, n) \cong \mathrm{Map}(k[-n], \wedge^p \mathbb{L}_A)$. There are natural forgetful maps

$$(3.1) \quad \mathcal{A}^{p, \mathrm{cl}}(A, n) \rightarrow \mathcal{A}^p(A, n)$$

Proposition 3.1. *The functor \mathbf{DR} is convergent, i.e. for any connective commutative dg algebra A the natural morphism*

$$\mathbf{DR}(A) \longrightarrow \lim_n \mathbf{DR}(\tau^{\geq -n}(A))$$

is an isomorphism.

Proof. Let $A_n = \tau^{\geq -n}(A)$ and $i_n: A \rightarrow A_n$ the projection. We have to show that the natural morphism

$$\mathrm{Sym}_A^p(\mathbb{L}_A[-1]) \longrightarrow \lim_n \mathrm{Sym}_{A_n}^p(\mathbb{L}_{A_n}[-1])$$

is an isomorphism for every $p \geq 0$. This morphism factors as the composite

$$\mathrm{Sym}_A^p(\mathbb{L}_A[-1]) \longrightarrow \lim_n \mathrm{Sym}_{A_n}^p(i_n^* \mathbb{L}_A[-1]) \longrightarrow \lim_n \mathrm{Sym}_{A_n}^p(\mathbb{L}_{A_n}[-1]).$$

Since \mathbb{L}_A is connective, $\mathrm{Sym}_A^p(\mathbb{L}_A[-1])$ is bounded on the right. Therefore, the first map is an isomorphism by [Lur18, Proposition 19.2.1.5]. It remains to show that

$$\lim_n (\mathrm{fib}(\mathrm{Sym}_{A_n}^p(i_n^* \mathbb{L}_A[-1]) \longrightarrow \mathrm{Sym}_{A_n}^p(\mathbb{L}_{A_n}[-1]))) = 0.$$

Consider an integer N . By construction $\text{cofib}(i_N)$ is $(N + 2)$ -connective. Therefore, by [Lur17, Corollary 7.4.3.2] the relative cotangent complex $\mathbb{L}_{A_N/A}$ is $(N + 2)$ -connective. Using the fiber sequence

$$\mathbb{L}_{A_N/A}[-1] \longrightarrow i_N^* \mathbb{L}_A \longrightarrow \mathbb{L}_{A_N}$$

we obtain a filtration on $\text{Sym}_{A_N}^p(i_N^* \mathbb{L}_A[-1])$ with associated graded pieces

$$\text{Sym}_{A_N}^k(\mathbb{L}_{A_N/A}[-2]) \otimes \text{Sym}_{A_N}^{p-k}(\mathbb{L}_{A_N}[-1]).$$

Therefore, the fibre

$$\text{fib}(\text{Sym}_{A_N}^p(i_N^* \mathbb{L}_A[-1]) \longrightarrow \text{Sym}_{A_N}^p(\mathbb{L}_{A_N}[-1]))$$

is $(N + 1 - p)$ -connective. For an inverse system $\{x_n\}$, the inverse limit may be computed as the fibre

$$\lim_n x_n = \text{fib} \left(\prod_{n \geq N} x_n \rightarrow \prod_{n \geq N} x_n \right)$$

for any N . We obtain that the left-hand side is $(N - p)$ -connective for any N and is, therefore, 0. \square

For any derived prestack \mathfrak{X} we define $\mathbf{DR}(\mathfrak{X})$ by a right Kan extension, i.e.

$$\mathbf{DR}(\mathfrak{X}) = \lim_{\text{Spec } A \rightarrow \mathfrak{X}} \mathbf{DR}(A)$$

and similarly for the other spaces. By [Pan+13, Proposition 1.14] for any derived Artin stack \mathfrak{X} we have

$$\mathcal{A}^p(\mathfrak{X}, n) \cong \text{Map}_{\text{QCoh}(\mathfrak{X})}(\mathcal{O}_{\mathfrak{X}}, \wedge^p \mathbb{L}_{\mathfrak{X}}[n]).$$

For a morphism between derived stacks $f: \mathfrak{X}_1 \rightarrow \mathfrak{X}_2$, we can define pullback morphisms of (closed, exact) n -shifted p -forms

$$f^*: \mathcal{A}^p(\mathfrak{X}_2, n) \rightarrow \mathcal{A}^p(\mathfrak{X}_1, n), \quad f^*: \mathcal{A}^{p,\text{cl}}(\mathfrak{X}_2, n) \rightarrow \mathcal{A}^{p,\text{cl}}(\mathfrak{X}_1, n), \quad f^*: \mathcal{A}^{p,\text{ex}}(\mathfrak{X}_2, n) \rightarrow \mathcal{A}^{p,\text{ex}}(\mathfrak{X}_1, n).$$

We will particularly be interested in the case of (-1) -shifted 2-forms, in which case the following statement is useful. Recall from [GR17a, Chapter 2, Section 1.7.2] that a derived prestack \mathfrak{X} is **locally almost of finite type** if it is convergent and its n -truncation $\tau^{\leq n}(\mathfrak{X})$ is locally of finite type.

Proposition 3.2. *Let \mathfrak{X} be a derived prestack locally almost of finite type. Then there is a canonical isomorphism $\mathcal{A}^{2,\text{ex}}(\mathfrak{X}, -1) \cong \mathcal{A}^{2,\text{cl}}(\mathfrak{X}, -1) \times \Gamma(\mathfrak{X}, \mathbb{C}_X)$, so that the projection on the first factor is given by the de Rham differential, where $\Gamma(\mathfrak{X}, \mathbb{C}_X)$ is the set of locally-constant functions.*

Proof. By Proposition 3.1 the functor \mathbf{DR} is convergent, so we have

$$\mathbf{DR}(\mathfrak{X}) = \lim_n \mathbf{DR}(\tau^{\leq n}(\mathfrak{X})).$$

Since \mathfrak{X} is locally almost of finite type, $\tau^{\leq n}(\mathfrak{X})$ is locally of finite type and can, therefore, be written as a colimit (in the ∞ -category of n -truncated derived prestacks) of a diagram of n -truncated derived affine schemes of finite type. Thus, it is enough to prove the claim for $\mathfrak{X} = \text{Spec } A$ a derived affine scheme of finite type. We have a fiber sequence

$$\mathcal{A}^{0,\text{cl}}(\mathfrak{X}, 0) \longrightarrow \mathcal{A}^{2,\text{ex}}(\mathfrak{X}, -1) \xrightarrow{d_{\text{dR}}} \mathcal{A}^{2,\text{cl}}(\mathfrak{X}, -1)$$

of pointed spaces. Moreover, there is a natural forgetful map $\mathcal{A}^{2,\text{ex}}(\mathfrak{X}, -1) \rightarrow \mathcal{A}^{1,\text{ex}}(\mathfrak{X}, 0) \cong \mathcal{A}^0(\mathfrak{X}, 0)$ so that the composite

$$\mathcal{A}^{0,\text{cl}}(\mathfrak{X}, 0) \longrightarrow \mathcal{A}^{2,\text{ex}}(\mathfrak{X}, -1) \longrightarrow \mathcal{A}^0(\mathfrak{X}, 0)$$

identifies with the forgetful map (3.1). By [BBJ19, Proposition 5.6] we have the following:

- The restriction $\mathcal{A}^{0,\text{cl}}(\mathfrak{X}, 0) \rightarrow \mathcal{A}^0(\mathfrak{X}, 0) \rightarrow \mathcal{A}^0(\mathfrak{X}^{\text{red}}, 0)$ identifies $\mathcal{A}^{0,\text{cl}}(\mathfrak{X}, 0)$ with the set of locally-constant \mathbb{C} -valued functions on $\mathfrak{X}^{\text{red}}$, i.e. there is a natural isomorphism $\mathcal{A}^{0,\text{cl}}(\mathfrak{X}, 0) \cong \Gamma(\mathfrak{X}, \mathbb{C})$.
- The de Rham differential $d_{\text{dR}}: \mathcal{A}^{2,\text{ex}}(\mathfrak{X}, -1) \rightarrow \mathcal{A}^{2,\text{cl}}(\mathfrak{X}, -1)$ induces an isomorphism on π_k for $k \geq 1$.
- On the level of π_0 we have a short exact sequence of abelian groups

$$0 \longrightarrow \pi_0(\mathcal{A}^{0,\text{cl}}(\mathfrak{X}, 0)) \longrightarrow \pi_0(\mathcal{A}^{2,\text{ex}}(\mathfrak{X}, -1)) \longrightarrow \pi_0(\mathcal{A}^{2,\text{cl}}(\mathfrak{X}, -1)) \longrightarrow 0.$$

Let $\mathcal{A}_0^{2,\text{ex}}(\mathfrak{X}, -1)$ be the homotopy fibre of the composite

$$\mathcal{A}^{2,\text{ex}}(\mathfrak{X}, -1) \longrightarrow \mathcal{A}^0(\mathfrak{X}, 0) \longrightarrow \mathcal{A}^0(\mathfrak{X}^{\text{red}}, 0).$$

By the above facts we get that the composite

$$\mathcal{A}_0^{2,\text{ex}}(\mathfrak{X}, -1) \longrightarrow \mathcal{A}^{2,\text{ex}}(\mathfrak{X}, -1) \longrightarrow \mathcal{A}^{2,\text{cl}}(\mathfrak{X}, -1)$$

as well as the morphism

$$\mathcal{A}_0^{2,\text{ex}}(\mathfrak{X}, -1) \times \mathcal{A}^{0,\text{cl}}(\mathfrak{X}, 0) \longrightarrow \mathcal{A}^{2,\text{ex}}(\mathfrak{X}, -1)$$

are isomorphisms. \square

Remark 3.3. Let \mathfrak{X} be a classical prestack locally of finite type and let $\mathfrak{X}^{\text{der}}$ be the corresponding derived prestack. Let $\mathfrak{X}^{\text{der,conv}}$ be the convergent completion of $\mathfrak{X}^{\text{der}}$ (see [GR17a, Chapter 2, Section 1.4.8]). Then $\mathfrak{X}^{\text{der,conv}}$ is locally almost of finite type (see [GR17a, Chapter 2, Corollary 1.7.8]). Since \mathbf{DR} is convergent by Proposition 3.1, $\mathbf{DR}(\mathfrak{X}^{\text{der,conv}}) \cong \mathbf{DR}(\mathfrak{X}^{\text{der}})$ and so Proposition 3.2 applies to $\mathfrak{X}^{\text{der}}$ as well.

Definition 3.4. Let \mathfrak{X} be a derived higher Artin stack. An *n -shifted symplectic structure on \mathfrak{X}* is a closed n -shifted 2-form $\omega_{\mathfrak{X}} \in \mathcal{A}^{2,\text{cl}}(\mathfrak{X}, n)$ whose underlying n -shifted 2-form induces an equivalence

$$\mathbb{T}_{\mathfrak{X}} = \mathbb{L}_{\mathfrak{X}}^{\vee} \xrightarrow[\simeq]{\omega_{\mathfrak{X}}} \mathbb{L}_{\mathfrak{X}}[n].$$

The notion of shifted symplectic structures extends to locally geometric derived stacks, that is, the unions of open substacks that are derived higher Artin stacks.

A derived higher Artin stack equipped with an n -shifted symplectic structure is called an *n -shifted symplectic stack*. In this paper, we are mainly interested in (-1) -shifted symplectic stacks. Given a pair of n -shifted symplectic stacks $(\mathfrak{X}, \omega_{\mathfrak{X}}), (\mathfrak{Y}, \omega_{\mathfrak{Y}})$, their product $\mathfrak{X} \times \mathfrak{Y}$ carries an n -shifted symplectic structure

$$\omega_{\mathfrak{X}} \boxplus \omega_{\mathfrak{Y}} := \text{pr}_1^* \omega_{\mathfrak{X}} + \text{pr}_2^* \omega_{\mathfrak{Y}}.$$

Definition 3.5. Let (X_1, ω_{X_1}) and (X_2, ω_{X_2}) be a pair of n -shifted symplectic stacks. A *Lagrangian correspondence* $\mathfrak{X}_1 \xleftarrow{\tau_1} \mathfrak{L} \xrightarrow{\tau_2} \mathfrak{X}_2$ is the choice of a homotopy $\eta: \tau_1^* \omega_{X_1} \sim \tau_2^* \omega_{X_2}$ in $\mathcal{A}^{2,\text{cl}}(\mathfrak{L}, n)$ such that the naturally induced commutative square

$$\begin{array}{ccc} \mathbb{T}_{\mathfrak{L}} & \longrightarrow & \tau_1^* \mathbb{T}_{\mathfrak{X}_1} \\ \downarrow & & \downarrow \\ \tau_2^* \mathbb{T}_{\mathfrak{X}_2} & \longrightarrow & \mathbb{L}_{\mathfrak{L}}[n] \end{array}$$

is Cartesian. A *Lagrangian morphism* $\mathfrak{L} \rightarrow \mathfrak{X}$ is a Lagrangian correspondence $\text{pt} \leftarrow \mathfrak{L} \rightarrow \mathfrak{X}$.

Suppose $\mathfrak{X}_1, \mathfrak{X}_2, \mathfrak{X}_3$ are three n -shifted symplectic stacks and $\mathfrak{X}_1 \xleftarrow{\tau_1} \mathfrak{L}_1 \xrightarrow{\tau_2} \mathfrak{X}_2$ and $\mathfrak{X}_2 \xleftarrow{\sigma_1} \mathfrak{L}_2 \xrightarrow{\sigma_2} \mathfrak{X}_3$ are two Lagrangian correspondences. Consider their composition

$$\begin{array}{ccccc} & & \mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2 & & \\ & \swarrow & & \searrow & \\ & \mathfrak{L}_1 & & \mathfrak{L}_2 & \\ \swarrow \tau_1 & & & & \searrow \sigma_2 \\ \mathfrak{X}_1 & & \mathfrak{X}_2 & & \mathfrak{X}_3 \\ \nwarrow & \nearrow \tau_2 & \nwarrow \sigma_1 & & \end{array}$$

Then $\mathfrak{X}_1 \leftarrow \mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2 \rightarrow \mathfrak{X}_3$ is a Lagrangian correspondence [Cal15, Theorem 4.4].

Definition 3.6. A *2-fold Lagrangian correspondence* is a diagram

$$\begin{array}{ccccc}
 & & \mathfrak{L}_1 & & \\
 & \swarrow & \uparrow & \searrow & \\
 \mathfrak{X}_1 & & \mathfrak{M} & & \mathfrak{X}_2 \\
 & \swarrow & \downarrow & \searrow & \\
 & & \mathfrak{L}_2 & &
 \end{array}$$

τ_1 (left arrow), τ_2 (right arrow), π_1 (up arrow), π_2 (down arrow), σ_1 (left arrow from \mathfrak{L}_2), σ_2 (right arrow to \mathfrak{L}_2)

where $(\mathfrak{X}_1, \omega_{\mathfrak{X}_1})$ and $(\mathfrak{X}_2, \omega_{\mathfrak{X}_2})$ are n -shifted symplectic stacks, $\mathfrak{X}_1 \xleftarrow{\tau_1} \mathfrak{L}_1 \xrightarrow{\tau_2} \mathfrak{X}_2$ and $\mathfrak{X}_1 \xleftarrow{\sigma_1} \mathfrak{L}_2 \xrightarrow{\sigma_2} \mathfrak{X}_2$ are Lagrangian correspondences specified by the homotopies η_1 in $\mathcal{A}^{2,\text{cl}}(\mathfrak{L}_1, n)$ and η_2 in $\mathcal{A}^{2,\text{cl}}(\mathfrak{L}_2, n)$ as well as a homotopy ξ from $\pi_1^* \eta_1$ to $\pi_2^* \eta_2$ in $\mathcal{A}^{2,\text{cl}}(\mathfrak{M}, n)$ such that the naturally induced diagram

$$\begin{array}{ccccc}
 & & \mathbb{T}_{\mathfrak{M}} & & \\
 & \swarrow & & \searrow & \\
 \pi_1^* \mathbb{T}_{\mathfrak{L}_1} & & & & \pi_2^* \mathbb{T}_{\mathfrak{L}_2} \\
 \downarrow & \swarrow & & \searrow & \downarrow \\
 \pi_1^* \tau_1^* \mathbb{T}_{\mathfrak{X}_1} \simeq \pi_2^* \sigma_1^* \mathbb{T}_{\mathfrak{X}_1} & & & & \pi_1^* \tau_2^* \mathbb{T}_{\mathfrak{X}_2} \simeq \pi_2^* \sigma_2^* \mathbb{T}_{\mathfrak{X}_2} \\
 & \swarrow & & \searrow & \\
 & & \mathbb{L}_{\mathfrak{M}}[n] & &
 \end{array}$$

is a limit diagram.

We now explain some examples of n -shifted symplectic structures and Lagrangians.

Example 3.7. Let Perf denote the derived stack of perfect complexes which is locally geometric [TV07]. Let \mathcal{E} denote the universal perfect complex on Perf . Then the cotangent complex of Perf is given by

$$\mathbb{L}_{\text{Perf}} \simeq \mathcal{E}^\vee \otimes \mathcal{E}[-1].$$

It is shown in [Pan+13, Theorem 2.12] that Perf is equipped with a 2-shifted symplectic structure. The underlying 2-form corresponds to the equivalence

$$\mathbb{L}_{\text{Perf}}^\vee \simeq (\mathcal{E}^\vee \otimes \mathcal{E}[-1])^\vee \simeq \mathcal{E} \otimes \mathcal{E}^\vee[1] \simeq (\mathcal{E}^\vee \otimes \mathcal{E}[-1])[2] \simeq \mathbb{L}_{\text{Perf}}[2].$$

Example 3.8. Let G be a reductive algebraic group. The cotangent complex of BG is given by $\mathfrak{g}[-1]$, where \mathfrak{g} denote the adjoint representation of G . It is shown in [Pan+13, §1.2] that a 2-shifted symplectic structure for BG corresponds to a G -equivariant non-degenerate bilinear form for \mathfrak{g} . In particular, there exists a 2-shifted symplectic structure for BG .

Example 3.9. Let Y be a smooth projective Calabi–Yau d -fold or the Betti geometric realization of an oriented d -dimensional closed manifold. If we are given an n -shifted symplectic stack \mathfrak{X} , it is shown in [Pan+13, Theorem 2.5] that the mapping stack $\text{Map}(Y, \mathfrak{X})$ is naturally equipped with an $(n - d)$ -shifted symplectic structure. Further, assume that we are given a Lagrangian morphism $\mathfrak{L} \rightarrow \mathfrak{X}$. Then it is shown in [Cal15, Theorem 2.10] that the map $\text{Map}(Y, \mathfrak{L}) \rightarrow \text{Map}(Y, \mathfrak{X})$ is naturally equipped with a Lagrangian structure.

Example 3.10. Let \mathfrak{X} be a derived higher Artin stack. For an integer $n \in \mathbb{Z}$, the *n -shifted cotangent stack* is defined to be the total space

$$T^*[n]\mathfrak{X} := \text{Tot}_{\mathfrak{X}}(\mathbb{L}_{\mathfrak{X}}[n]).$$

Let $\lambda \in \mathcal{A}^1(T^*[n]\mathfrak{X}, n)$ denote the tautological n -form. Then it is shown in [Cal19, Theorem 2.2] that $\omega_{\mathfrak{X}} := d_{\text{dR}} \lambda$ is n -shifted symplectic.

Let $f: \mathfrak{X} \rightarrow \mathbb{A}^1[n] := \text{Tot}_{\text{pt}}(\mathbb{C}[n])$ be an n -shifted function. The de Rham differential of f defines a section $df: \mathfrak{X} \rightarrow T^*[n]\mathfrak{X}$. It is shown in [Cal19, Theorem 2.15] that the map df is canonically equipped with a Lagrangian structure. We define the derived critical locus $\text{Crit}(f)$ by the fibre product

$$\begin{array}{ccc} \text{Crit}(f) & \longrightarrow & \mathfrak{X} \\ \downarrow & \lrcorner & \downarrow df \\ \mathfrak{X} & \xrightarrow{0} & T^*[n]\mathfrak{X}. \end{array}$$

Since $\text{Crit}(f)$ is defined as the fibre product of two Lagrangian morphisms, it is equipped with an $(n-1)$ -shifted symplectic structure $\omega_{\text{Crit}(f)}$.

The following Darboux theorem plays a very important role in the study of shifted symplectic stacks:

Theorem 3.11 ([Ben+15, Corollary 2.11]).

- (i) Let (X, ω_X) be a (-1) -shifted symplectic derived algebraic space. Then for each point $x \in X$ there is an **étale derived critical chart** $\mathfrak{R} = (R, \eta, U, f, i)$, i.e. a smooth scheme U , regular function $f: U \rightarrow \mathbb{A}^1$ and an étale morphism $\eta: \text{Crit}(f) \rightarrow X$ whose image contains x such that $\eta^*\omega_X \sim \omega_{\text{Crit}(f)}$ holds.
- (ii) Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a (-1) -shifted symplectic derived Artin stack. Then for each point $x \in X$, we can construct the following diagram:

$$(3.2) \quad \begin{array}{ccc} \widehat{V} & \xrightarrow{\tau} & V = \text{Crit}(f) \\ \downarrow q & & \\ \mathfrak{X} & & \end{array}$$

and a homotopy $q^*\omega_{\mathfrak{X}} \sim \tau^*\omega_{\text{Crit}(f)}$, where q is a smooth morphism whose image contains x , f is a regular function on a smooth scheme and τ is a morphism which induces an equivalence on the classical truncations.

Remark 3.12. The homotopy $q^*\omega_{\mathfrak{X}} \sim \tau^*\omega_{\text{Crit}(f)}$ induces a morphism of fibre sequences

$$\begin{array}{ccccc} \mathbb{T}_{\widehat{V}/V}[1] & \longrightarrow & \mathbb{T}_{\widehat{V}}[1] & \longrightarrow & \tau^*\mathbb{T}_V[1] \\ \downarrow \eta & & \downarrow & & \downarrow \\ \mathbb{L}_{\widehat{V}/\mathfrak{X}}[-1] & \longrightarrow & q^*\mathbb{L}_{\mathfrak{X}} & \longrightarrow & \mathbb{L}_{\widehat{V}}. \end{array}$$

Since q is smooth and τ is an isomorphism on classical truncations, we see that η is invertible by the degree reason. In particular, the correspondence $\mathfrak{X} \leftarrow \widehat{V} \rightarrow V$ is a Lagrangian correspondence.

We also have the following equivariant version of Darboux theorem:

Proposition 3.13. *Let \mathfrak{X} be a quasi-separated derived Artin stack with affine stabilizers and $\omega_{\mathfrak{X}}$ be a (-1) -shifted symplectic structure. If $x \in \mathfrak{X}(\mathbb{C})$ is a point with linearly reductive stabilizer G , then there exists a smooth affine scheme U with a G -action, a fixed point $u \in U(\mathbb{C})$, a G -invariant function $f: U \rightarrow \mathbb{A}^1$, and a pointed étale symplectomorphism*

$$(\text{Crit}([f/G], u) \rightarrow (\mathfrak{X}, \omega_{\mathfrak{X}}, x),$$

that preserves the stabilizer at u , where $[f/G]: [U/G] \rightarrow \mathbb{A}^1$ is the induced function.

Proof. Proposition 3.2 shows that all (-1) -shifted symplectic forms admit canonical exact structures. The claim then follows from [Par24, Theorem B]. \square

Remark 3.14. If \mathfrak{X} is a (-1) -symplectic derived Artin stack with affine diagonal such that the classical truncation \mathfrak{X}^{cl} has a good moduli space Y , then we can cover \mathfrak{X} with the equivariant Darboux charts in Proposition 3.13. Indeed, for any point $y \in Y$, we can find a point $x \in \mathfrak{X}$ that maps to y and has a linearly reductive stabilizer [Alp13, Proposition 12.14]. Then we can find an étale morphism of the form $\text{Crit}([f/G]) \rightarrow \mathfrak{X}$ by Proposition 3.13, which induces an étale morphism on their good moduli spaces by [AHR20, Proposition 4.13], after shrinking U .

3.2. Orientation for (-1) -shifted symplectic stacks. We recall the notion of orientation for (-1) -shifted symplectic stacks and their Lagrangians.

Definition 3.15. Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a (-1) -shifted symplectic derived Artin stack. A (graded) **orientation of \mathfrak{X}** is the pair of a graded line bundle $\mathcal{L} \in \text{Pic}^{\text{gr}}(\mathfrak{X}^{\text{red}})$ and an isomorphism $o: \mathcal{L}^{\otimes 2} \cong \widehat{\det}(\mathbb{L}_{\mathfrak{X}})$. By abuse of notation, we simply say that o is an orientation of \mathfrak{X} .

Orientations naturally form a groupoid.

Definition 3.16. An **isomorphism of orientations** $(\mathcal{L}_1, o_1) \cong (\mathcal{L}_2, o_2)$ is an isomorphism of graded line bundles $f: \mathcal{L}_1 \cong \mathcal{L}_2$ on $\mathfrak{X}^{\text{red}}$ such that the triangle

$$\begin{array}{ccc} \mathcal{L}_1^{\otimes 2} & \xrightarrow{f^{\otimes 2}} & \mathcal{L}_2^{\otimes 2} \\ & \searrow o_1 & \swarrow o_2 \\ & \widehat{\det}(\mathbb{L}_{\mathfrak{X}}) & \end{array}$$

commutes.

Orientations of \mathfrak{X} localize to define the graded **orientation μ_2 -gerbe** $\text{or}_{\mathfrak{X}}$ on $\mathfrak{X}^{\text{red}}$, so that an orientation of \mathfrak{X} is a trivialization of $\text{or}_{\mathfrak{X}}$.

For a pair of (-1) -shifted symplectic stacks $\mathfrak{X}_1, \mathfrak{X}_2$, the natural isomorphism

$$\Psi_{\mathfrak{X}_1, \mathfrak{X}_2}: \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1}) \boxtimes \widehat{\det}(\mathbb{L}_{\mathfrak{X}_2}) \cong \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1 \times \mathfrak{X}_2})$$

of graded line bundles on $\mathfrak{X}_1^{\text{red}} \times \mathfrak{X}_2^{\text{red}}$ induces an isomorphism

$$\text{or}_{\mathfrak{X}_1} \boxtimes \text{or}_{\mathfrak{X}_2} \cong \text{or}_{\mathfrak{X}_1 \times \mathfrak{X}_2}$$

of graded orientation gerbes. In particular, if o_1 and o_2 are orientations of \mathfrak{X}_1 and \mathfrak{X}_2 , using the above isomorphism we obtain an orientation $o_1 \boxtimes o_2$ of $\mathfrak{X}_1 \times \mathfrak{X}_2$.

Assume that $\mathfrak{X}_1 \leftarrow \mathfrak{L} \rightarrow \mathfrak{X}_2$ is a Lagrangian correspondence of (-1) -shifted symplectic stacks. Then we have an equivalence

$$(3.3) \quad \mathbb{T}_{\mathfrak{L}/\mathfrak{X}_1}[1] \simeq \mathbb{L}_{\mathfrak{L}/\mathfrak{X}_2}[-1].$$

which induces an isomorphism of determinant line bundles

$$(3.4) \quad \widehat{\det}(\mathbb{L}_{\mathfrak{L}/\mathfrak{X}_1}) \cong \widehat{\det}(\mathbb{L}_{\mathfrak{L}/\mathfrak{X}_2})^{\vee}.$$

by (2.5) and (2.14). Using isomorphisms (2.3) for fibre sequences $\mathbb{L}_{\mathfrak{X}_1}|_{\mathfrak{L}} \rightarrow \mathbb{L}_{\mathfrak{L}} \rightarrow \mathbb{L}_{\mathfrak{L}/\mathfrak{X}_1}$ and $\mathbb{L}_{\mathfrak{X}_2}|_{\mathfrak{L}} \rightarrow \mathbb{L}_{\mathfrak{L}} \rightarrow \mathbb{L}_{\mathfrak{L}/\mathfrak{X}_2}$, we obtain isomorphisms

$$\widehat{\det}(\mathbb{L}_{\mathfrak{X}_1}|_{\mathfrak{L}}) \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}/\mathfrak{X}_1}) \cong \widehat{\det}(\mathbb{L}_{\mathfrak{L}}), \quad \widehat{\det}(\mathbb{L}_{\mathfrak{L}}) \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}/\mathfrak{X}_2})^{\vee} \cong \widehat{\det}(\mathbb{L}_{\mathfrak{X}_2}|_{\mathfrak{L}}).$$

By combining them and using (3.4), we obtain an isomorphism

$$(3.5) \quad \xi: \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1}|_{\mathfrak{L}}) \otimes (\widehat{\det}(\mathbb{L}_{\mathfrak{L}/\mathfrak{X}_2})^{\vee})^{\otimes 2} \cong \widehat{\det}(\mathbb{L}_{\mathfrak{X}_2}|_{\mathfrak{L}})$$

of graded line bundles on $\mathfrak{L}^{\text{red}}$. Therefore, there is an isomorphism


$$(3.6) \quad \beta_{\mathfrak{L}}: \text{or}_{\mathfrak{X}_1}|_{\mathfrak{L}} \cong \text{or}_{\mathfrak{X}_2}|_{\mathfrak{L}}$$

of orientation gerbes given by sending a pair (\mathcal{L}, o) of a graded line bundle \mathcal{L} on $\mathfrak{L}^{\text{red}}$ together with an isomorphism $o: \mathcal{L}^{\otimes 2} \simeq \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1}|_{\mathfrak{L}})$ to the graded line bundle $\mathcal{L} \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}/\mathfrak{X}_2})^{\vee}$ equipped with the isomorphism

$$(\mathcal{L} \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}/\mathfrak{X}_2})^{\vee})^{\otimes 2} \xrightarrow{\alpha} \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1}|_{\mathfrak{L}}) \otimes (\widehat{\det}(\mathbb{L}_{\mathfrak{L}/\mathfrak{X}_2})^{\vee})^{\otimes 2} \xrightarrow{\xi} \widehat{\det}(\mathbb{L}_{\mathfrak{X}_2}|_{\mathfrak{L}}).$$

Definition 3.17. An **orientation of the Lagrangian correspondence** $\mathfrak{X}_1 \xleftarrow{\tau_1} \mathfrak{L} \xrightarrow{\tau_2} \mathfrak{X}_2$ is the choice of orientations o_1 of \mathfrak{X}_1 and o_2 of \mathfrak{X}_2 together with an isomorphism ρ of the corresponding orientations related by $\beta_{\mathfrak{L}}$.

Consider a composition of Lagrangian correspondences



There is a natural fibre sequence

$$\mathbb{L}_{\mathfrak{L}_2/\mathfrak{X}_3|\mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2} \longrightarrow \mathbb{L}_{\mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2/\mathfrak{X}_3} \longrightarrow \mathbb{L}_{\mathfrak{L}_1/\mathfrak{X}_2|\mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2}.$$

Using the isomorphism (2.3) we obtain an isomorphism of determinant line bundles

$$\pi: \widehat{\det}(\mathbb{L}_{\mathfrak{L}_2/\mathfrak{X}_3|\mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2}) \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}_1/\mathfrak{X}_2|\mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2}) \cong \widehat{\det}(\mathbb{L}_{\mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2/\mathfrak{X}_3}).$$

We now show that the map π is compatible with ξ :

Lemma 3.18. *Set $\widehat{\mathfrak{L}} := \mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2$. Then the following diagram commutes:*

$$\begin{array}{ccc}
 \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1|\widehat{\mathfrak{L}}}) \otimes (\widehat{\det}(\mathbb{L}_{\mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2/\mathfrak{X}_3})^\vee)^{\otimes 2} & \xrightarrow[\cong]{\xi} & \widehat{\det}(\mathbb{L}_{\mathfrak{X}_3|\widehat{\mathfrak{L}}}) \\
 \text{id} \otimes (\pi^\vee)^{\otimes 2} \uparrow \cong & & \xi \uparrow \cong \\
 \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1|\widehat{\mathfrak{L}}}) \otimes (\widehat{\det}(\mathbb{L}_{\mathfrak{L}_1/\mathfrak{X}_2|\widehat{\mathfrak{L}}})^\vee)^{\otimes 2} \otimes (\widehat{\det}(\mathbb{L}_{\mathfrak{L}_2/\mathfrak{X}_3|\widehat{\mathfrak{L}}})^\vee)^{\otimes 2} & \xrightarrow[\cong]{\xi \otimes \text{id}} & \widehat{\det}(\mathbb{L}_{\mathfrak{X}_2|\widehat{\mathfrak{L}}}) \otimes (\widehat{\det}(\mathbb{L}_{\mathfrak{L}_2/\mathfrak{X}_3|\widehat{\mathfrak{L}}})^\vee)^{\otimes 2}.
 \end{array}$$

Proof. By the definition of the composite Lagrangian structure, we have the following equivalence of fibre sequences:

$$\begin{array}{ccccc}
 \mathbb{T}_{\mathfrak{L}_1/\mathfrak{X}_1}[1]|\widehat{\mathfrak{L}} & \longrightarrow & \mathbb{T}_{\widehat{\mathfrak{L}}/\mathfrak{X}_1}[1] & \longrightarrow & \mathbb{T}_{\mathfrak{L}_2/\mathfrak{X}_3}[1]|\widehat{\mathfrak{L}} \\
 (3.3) \downarrow \simeq & & (3.3) \downarrow \simeq & & (3.3) \downarrow \simeq \\
 \mathbb{L}_{\mathfrak{L}_1/\mathfrak{X}_2}[-1]|\widehat{\mathfrak{L}} & \longrightarrow & \mathbb{L}_{\widehat{\mathfrak{L}}/\mathfrak{X}_3}[-1] & \longrightarrow & \mathbb{L}_{\mathfrak{L}_2/\mathfrak{X}_3}[-1]|\widehat{\mathfrak{L}}.
 \end{array}$$

This equivalence of fibre sequences combined with the Lemma 2.4 implies the commutativity of the following diagram:

$$\begin{array}{ccc}
 \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1|\widehat{\mathfrak{L}}}) \otimes \widehat{\det}(\mathbb{L}_{\widehat{\mathfrak{L}}/\mathfrak{X}_3})^\vee & \xrightarrow[\cong]{(3.4)} & \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1|\widehat{\mathfrak{L}}}) \otimes \widehat{\det}(\mathbb{L}_{\widehat{\mathfrak{L}}/\mathfrak{X}_1}) \\
 \downarrow \cong & & \downarrow \cong \\
 \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1|\widehat{\mathfrak{L}}}) \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}_1/\mathfrak{X}_2|\widehat{\mathfrak{L}}})^\vee \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}_2/\mathfrak{X}_3|\widehat{\mathfrak{L}}})^\vee & & \\
 \cong \downarrow (3.4) & & \\
 \widehat{\det}(\mathbb{L}_{\mathfrak{X}_1|\widehat{\mathfrak{L}}}) \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}_1/\mathfrak{X}_1|\widehat{\mathfrak{L}}}) \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}_2/\mathfrak{X}_2|\widehat{\mathfrak{L}}}) & \xrightarrow[\cong]{} & \widehat{\det}(\mathbb{L}_{\widehat{\mathfrak{L}}}).
 \end{array} \tag{3.7}$$

Using Lemma 2.4 again, we obtain the commutativity of the following diagram:

$$\begin{array}{ccc}
 \widehat{\det}(\mathbb{L}_{\widehat{\mathfrak{L}}}) \otimes \widehat{\det}(\mathbb{L}_{\widehat{\mathfrak{L}}/\mathfrak{X}_3|\widehat{\mathfrak{L}}})^\vee & \xrightarrow[\cong]{} & \widehat{\det}(\mathbb{L}_{\mathfrak{X}_3|\widehat{\mathfrak{L}}}) \\
 \downarrow \cong & & \uparrow \cong \\
 \widehat{\det}(\mathbb{L}_{\widehat{\mathfrak{L}}}) \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}_1/\mathfrak{X}_2|\widehat{\mathfrak{L}}})^\vee \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}_2/\mathfrak{X}_3|\widehat{\mathfrak{L}}})^\vee & \xrightarrow[\cong]{} & \widehat{\det}(\mathbb{L}_{\mathfrak{L}_2|\widehat{\mathfrak{L}}}) \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{L}_2/\mathfrak{X}_3|\widehat{\mathfrak{L}}})^\vee.
 \end{array} \tag{3.8}$$

The claim of the lemma follows from the commutativity of diagrams (3.7) and (3.8). \square

By Lemma 3.18, we see that the isomorphism π induces a natural 2-isomorphism

$$(3.9) \quad \beta_{\mathfrak{L}_2} \circ \beta_{\mathfrak{L}_1} \cong \beta_{\mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2}$$

of isomorphisms $\text{or}_{\mathfrak{X}_1}|_{\mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2} \simeq \text{or}_{\mathfrak{X}_3}|_{\mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2}$ of orientation gerbes. In particular, if $\mathfrak{X}_1 \leftarrow \mathfrak{L}_1 \rightarrow \mathfrak{X}_2$ and $\mathfrak{X}_2 \leftarrow \mathfrak{L}_2 \rightarrow \mathfrak{X}_3$ are oriented, their composition $\mathfrak{X}_1 \leftarrow \mathfrak{L}_1 \times_{\mathfrak{X}_2} \mathfrak{L}_2 \rightarrow \mathfrak{X}_3$ is oriented as well.

We will often consider the following setting, which allows us to pullback orientations along Lagrangian correspondences with sections.

Lemma 3.19. *Let $\mathfrak{X}_1 \xleftarrow{\tau_1} \mathfrak{L} \xrightarrow{\tau_2} \mathfrak{X}_2$ be a (-1) -shifted Lagrangian correspondence together with a morphism $i: \mathfrak{X}_1 \rightarrow \mathfrak{L}$ such that $\tau_1 \circ i \cong \text{id}_{\mathfrak{X}_1}$. Then there is a natural isomorphism*

$$i^* \tau_2^* \text{or}_{\mathfrak{X}_2} \cong \text{or}_{\mathfrak{X}_1}.$$

Moreover, suppose $\mathfrak{L} \rightarrow \mathfrak{X}_1$ is an \mathbb{A}^1 -deformation retract, i.e. there is a map $F: \mathbb{A}^1 \times \mathfrak{L} \rightarrow \mathfrak{X}_1$ satisfying the following conditions:

- (1) $F|_{\{1\} \times X} \simeq \text{id}_X$.
- (2) $F|_{\{0\} \times X} \simeq i \circ \tau_1$.

Then the forgetful functor from the groupoid of orientations

$$\left\{ (o_1, o_2, \rho) \left| \begin{array}{l} o_1: \text{orientation for } \mathfrak{X}_1 \\ o_2: \text{orientation for } \mathfrak{X}_2 \\ \rho: \text{orientation for } \mathfrak{X}_1 \xleftarrow{\tau_1} \mathfrak{L} \xrightarrow{\tau_2} \mathfrak{X}_2 \end{array} \right. \right\}$$

of the Lagrangian correspondence $\mathfrak{X}_1 \leftarrow \mathfrak{L} \rightarrow \mathfrak{X}_2$ to the groupoid of orientations $\{o_2\}$ of \mathfrak{X}_2 is an equivalence.

Proof. The isomorphism (3.6) provides an isomorphism

$$\beta_1: \text{or}_{\mathfrak{X}_1}|_{\mathfrak{L}} \cong \text{or}_{\mathfrak{X}_2}|_{\mathfrak{L}}.$$

Restricting it along i , we obtain the isomorphism

$$\text{or}_{\mathfrak{X}_2}|_{\mathfrak{X}_1} \cong \text{or}_{\mathfrak{X}_1}.$$

Let $\mathcal{C}_{\mathfrak{X}_1}$ be the 2-groupoid of graded μ_2 -gerbes on \mathfrak{X}_1 and similarly for \mathfrak{L} . Since $\tau_1 \circ i \simeq \text{id}_{\mathfrak{X}_1}$, $i^* \circ \tau_1^* \simeq \text{id}_{\mathcal{C}_{\mathfrak{X}_1}}$. So, it is enough to prove that $\tau_1^*: \mathcal{C}_{\mathfrak{X}_1} \rightarrow \mathcal{C}_{\mathfrak{L}}$ is an equivalence. The 2-groupoid $\mathcal{C}_{\mathfrak{X}_1}$ has homotopy groups given by

$$\pi_{2-n}(\mathcal{C}_{\mathfrak{X}_1}) \simeq H_{et}^n(\mathfrak{X}_1; \mu_2) \times H_{et}^{n-1}(\mathfrak{X}_1; \mathbb{Z}/2\mathbb{Z})$$

for $n = 0, 1, 2$, so it is enough to show that $\tau_1^*: H_{et}^n(\mathfrak{X}_1; \mu_2) \rightarrow H_{et}^n(\mathfrak{X}_1; \mu_2)$ is an isomorphism for $n = 0, 1, 2$ and similarly for cohomology with $\mathbb{Z}/2\mathbb{Z}$ coefficients. The projection $\pi_2: \mathbb{A}^1 \times \mathfrak{L} \rightarrow \mathfrak{L}$ is acyclic [Mil80, Chapter VI, Corollary 4.20], so $\pi_2^* = F^*$ as maps $H_{et}^n(\mathfrak{L}; \mu_2) \rightarrow H_{et}^n(\mathbb{A}^1 \times \mathfrak{L}; \mu_2)$. Restricting to $0, 1 \in \mathbb{A}^1$, we conclude that $\tau_1^* \circ i^* = \text{id}$ and hence $\tau_1^*: H_{et}^n(\mathfrak{X}_1; \mu_2) \rightarrow H_{et}^n(\mathfrak{L}; \mu_2)$ is an isomorphism. \square

Finally, we remark that derived critical loci admit canonical orientations:

Example 3.20. Let \mathfrak{Y} be a derived higher Artin stack and $f: \mathfrak{Y} \rightarrow \mathbb{A}^1$ be a regular function. As we have seen in Example 3.10, the derived critical locus $\mathfrak{X} := \text{Crit}(f)$ admits a (-1) -shifted symplectic structure $\omega_{\mathfrak{X}}$. By construction, we have a natural fibre sequence

$$\mathbb{L}_{\mathfrak{Y}}|_{\mathfrak{X}} \rightarrow \mathbb{L}_{\mathfrak{X}} \rightarrow \mathbb{T}_{\mathfrak{X}/\mathfrak{Y}}[1].$$

where the latter map is given by the composition $\mathbb{L}_{\mathfrak{X}} \xrightarrow{(\cdot \omega_{\mathfrak{X}})^{-1}} \mathbb{T}_{\mathfrak{X}}[1] \rightarrow \mathbb{T}_{\mathfrak{X}/\mathfrak{Y}}[1]$. Applying (2.3), (2.5) and (2.14), we obtain an isomorphism

$$\bar{o}_{\mathfrak{X}}^{\text{can}}: \widehat{\det}(\mathbb{L}_{\mathfrak{Y}}|_{\mathfrak{X}})^{\otimes 2} \cong \widehat{\det}(\mathbb{L}_{\mathfrak{X}}).$$

We set $o_{\mathfrak{X}}^{\text{can}} := (-1)^{\text{vd } \mathfrak{Y}} \cdot \bar{o}_{\mathfrak{X}}^{\text{can}}$ and call it the **canonical orientation** for the derived critical locus \mathfrak{X}^1 .

¹The insertion of the sign is to make it consistent with the canonical orientation for d-critical stacks: see (3.33) and Remark 3.36

3.3. D-critical structures. Here we recall Joyce's theory of d-critical structures, which can be regarded as classical truncations of (-1) -shifted symplectic structures. The original reference is [Joy15].

For an algebraic space X , Joyce [Joy15, Theorem 2.1] defined étale sheaves

$$\mathcal{S}_X, \mathcal{S}_X^0 \in \text{Mod}_{\mathbb{C}_X}$$

of \mathbb{C} -vector spaces with the following properties:

- (1) There is a decomposition $\mathcal{S}_X \cong \mathcal{S}_X^0 \oplus \mathbb{C}_X$.
- (2) Consider an étale morphism $\eta: R \rightarrow X$ from a scheme R . Since X is locally of finite type, R is locally of finite type as well. If we denote by R^{der} the corresponding derived scheme, then there exist isomorphisms

$$(3.10) \quad \Gamma(R, \mathcal{S}_X|_R) \cong \mathcal{A}^{2, \text{ex}}(R^{\text{der}}, -1), \quad \Gamma(R, \mathcal{S}_X^0|_R) \cong \mathcal{A}^{2, \text{cl}}(R^{\text{der}}, -1).$$

Under these isomorphisms the decomposition $\mathcal{S}_X \cong \mathcal{S}_X^0 \oplus \mathbb{C}_X$ corresponds to the isomorphism $\mathcal{A}^{2, \text{ex}}(R, -1) \cong \mathcal{A}^{2, \text{cl}}(R, -1) \times \Gamma(R, \mathbb{C}_R)$ from Proposition 3.2 (see Remark 3.3 why it applies).

- (3) For any étale morphism $\eta: R \rightarrow X$ and any closed embedding $i: R \hookrightarrow U$ where U is a smooth scheme, there exists the following short exact sequence:

$$0 \rightarrow \mathcal{S}_X|_R \rightarrow i^{-1}\mathcal{O}_U/I_{R,U}^2 \xrightarrow{d_{\text{dR}}} i^{-1}\Omega_U/(I_{R,U} \cdot i^{-1}\Omega_U)$$

where $I_{R,U}$ is the ideal sheaf of $i^{-1}\mathcal{O}_U$ defining R . The composite

$$\mathcal{S}_X|_R \rightarrow i^{-1}\mathcal{O}_U/I_{R,U}^2 \rightarrow i^{-1}\mathcal{O}_U/I_{R,U} \simeq \mathcal{O}_R \rightarrow \mathcal{O}_{R^{\text{red}}}$$

factors through locally constant functions $\mathbb{C}_{R^{\text{red}}} \rightarrow \mathcal{O}_{R^{\text{red}}}$ which defines the projection $\mathcal{S}_X \rightarrow \mathbb{C}_X$.

Assume that we are given a regular function f on a smooth scheme U such that $f|_{\text{Crit}(f)^{\text{red}}} = 0$. Set $X = \text{Crit}(f)^{\text{cl}}$. Then f defines a section of \mathcal{S}_X^0 .

Definition 3.21 ([Joy15, Definition 2.5]). Let X be an algebraic space and $s \in \Gamma(X, \mathcal{S}_X^0)$ is a section.

- An *étale d-critical chart* for (X, s) is an étale morphism $\eta: R \rightarrow X$, a smooth scheme U , a closed embedding $i: R \hookrightarrow U$ and a regular function f on U with $f|_{i(R)^{\text{red}}} = 0$, $i(R) = \text{Crit}(f)$ and $f + I_{R,U}^2 = s|_R$.
- The section s is a *d-critical structure* if for each point $x \in X$ there exists an étale d-critical chart (R, η, U, f, i) such that the image of η contains x .

Let $f: X \rightarrow Y$ be a morphism of algebraic spaces. Then there exists a natural morphism of sheaves

$$f^*: f^{-1}\mathcal{S}_Y^0 \rightarrow \mathcal{S}_X^0.$$

which corresponds under the isomorphism (3.10) to the pullback of differential forms. It is shown in [Joy15, Proposition 2.8] that if f is smooth and s is a d-critical structure of Y , then f^*s is a d-critical structure on X .

For an Artin stack \mathfrak{X} , the assignment

$$\text{Sch}_{/\mathfrak{X}}^{\text{sm}} \ni T \rightarrow \Gamma(T, \mathcal{S}_T^0)$$

together with the natural transition map defines a sheaf in the lisse-étale topos of \mathfrak{X} , which will be denoted by $\mathcal{S}_{\mathfrak{X}}^0$.

Definition 3.22. Let \mathfrak{X} be an Artin stack. A *d-critical structure on \mathfrak{X}* is a choice of a section $s \in \Gamma(\mathfrak{X}, \mathcal{S}_{\mathfrak{X}}^0)$ such that the restriction $s|_T \in \Gamma(T, \mathcal{S}_T^0)$ is a d-critical structure for each $T \in \text{Sch}_{/\mathfrak{X}}^{\text{sm}}$.

By [Joy15, Proposition 2.8], $s \in \Gamma(\mathfrak{X}, \mathcal{S}_{\mathfrak{X}}^0)$ is a d-critical structure if and only if there exists a smooth surjection from an algebraic space $T \rightarrow \mathfrak{X}$ such that $s|_T$ is a d-critical structure.

The following statement summarizes the relation between (-1) -shifted symplectic structures and d-critical structures. See [Ben+15, Theorem 3.18 (a)] and also [Kin22, Theorem 4.6] for the details.

Theorem 3.23. *Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be (-1) -shifted symplectic stack and $j: \mathfrak{X}^{\text{cl}} \rightarrow \mathfrak{X}$ the inclusion of the classical truncation. Let $s \in \Gamma(X, \mathcal{S}_X^0)$ be the section corresponding to $j^*\omega_{\mathfrak{X}}$ under the isomorphism (3.10). Then s is a d-critical structure.*

- (i) If \mathfrak{X} is an algebraic space and $\mathfrak{R} = (R, \eta, U, f, i)$ is an étale derived critical chart for \mathfrak{X} , then its classical truncation $\mathfrak{R}^{\text{cl}} = (R^{\text{cl}}, \eta^{\text{cl}}, U, f, i^{\text{cl}})$ defines an étale d-critical chart for \mathfrak{X}^{cl} .
- (ii) If we are given a diagram (3.2) such that $f|_{\text{Crit}(f)^{\text{red}}} = 0$, the following equality holds:

$$q^{\text{cl},*} s = \tau^{\text{cl},*} (f + I_{\text{Crit}(f)^{\text{cl}}, U}^2).$$

For later use, we explain how to compare two étale d-critical charts following [Joy15, §2.2]. Let (X, s) be a d-critical algebraic space. An étale morphism between étale d-critical charts $(R_1, \eta_1, U_1, f_1, i_1)$ and $(R_2, \eta_2, U_2, f_2, i_2)$ is given by étale morphisms $h: U_1 \rightarrow U_2$ and $h_0: R_1 \rightarrow R_2$ such that $f_1 = f_2 \circ h$, $\eta_1 = \eta_2 \circ h_0$ and $i_2 \circ h_0 = h \circ i_1$ hold. The following technical statement will be useful to find an étale cover of a given d-critical chart:

Lemma 3.24. *Let $h_0: R_1 \rightarrow R_2$ be an étale morphism of schemes and $i_2: R_2 \hookrightarrow U_2$ be a closed embedding of R_2 to a smooth scheme. Then for each $x \in R_1$, by possibly replacing R_1 with an open neighbourhood of x , there exist an étale morphism $h: U_1 \rightarrow U_2$ and a closed embedding $R_1 \hookrightarrow U_1$ such that the natural map $R_1 \rightarrow R_2 \times_{U_2} U_1$ is an isomorphism.*

Proof. By possibly shrinking R_1 around x , we may assume that R_1, U_1 and U_2 are affine. We take a closed embedding $\iota: R_1 \hookrightarrow \mathbb{A}^n$. Since the map $(h_0, \iota): R_1 \rightarrow R_2 \times \mathbb{A}^n$ is a regular closed embedding of relative dimension n , there exists a regular functions $g_1, \dots, g_n: R_2 \times \mathbb{A}^n \rightarrow \mathbb{A}^1$ such that the natural map $R_1 \rightarrow Z(g_1, \dots, g_n)$ is an isomorphism around x . By shrinking R_1 around x again, we may assume that the natural map $R_1 \rightarrow Z(g_1, \dots, g_n)$ is an open immersion. We choose a lift $\tilde{g}_i: U_2 \times \mathbb{A}^n \rightarrow \mathbb{A}^1$ of g_i such that the morphism $\tilde{h}: Z(\tilde{g}_1, \dots, \tilde{g}_n) \rightarrow U_2$ is étale at $(h_0(x), \iota(x))$. We may take an open subset $U_1 \subset Z(\tilde{g}_1, \dots, \tilde{g}_n)$ containing $(h_0(x), \iota(x))$ such that $\tilde{h}|_{U_1}$ is étale and the natural map $R_1 \rightarrow R_2 \times_{U_2} U_1$ is an isomorphism, hence we conclude. \square

We now introduce the notion of embedding of étale d-critical charts. Let (X, s) be a d-critical algebraic space. An embedding of étale d-critical charts $(R_1, \eta_1, U_1, f_1, i_1) \hookrightarrow (R_2, \eta_2, U_2, f_2, i_2)$ is given by an open embedding $\Phi_0: R_1 \hookrightarrow R_2$ and a locally closed embedding $\Phi: U_1 \hookrightarrow U_2$ such that $\eta_1 = \eta_2 \circ \Phi_0$, $f_1 = f_2 \circ \Phi$ and $\Phi \circ i_1 = i_2 \circ \Phi_0$ hold. The following statement is useful when comparing two étale d-critical charts:

Theorem 3.25. *Let (X, s) be a d-critical algebraic space.*

- (i) *Let $\mathfrak{R}_1 = (R_1, \eta_1, U_1, f_1, i_1)$ and $\mathfrak{R}_2 = (R_2, \eta_2, U_2, f_2, i_2)$ be two étale d-critical charts and $x_1 \in R_1$ and $x_2 \in R_2$ be points such that $\eta_1(x_1) = \eta_2(x_2)$ holds. Then by possibly replacing \mathfrak{R}_1 and \mathfrak{R}_2 by étale neighbourhoods of x_1 and x_2 , we may find a third d-critical chart $\mathfrak{R}_3 = (R_3, \eta_3, U_3, f_3, i_3)$ and $x_3 \in R_3$ such that there exists embeddings $\mathfrak{R}_1 \hookrightarrow \mathfrak{R}_3$ and $\mathfrak{R}_2 \hookrightarrow \mathfrak{R}_3$ which map x_1 to x_3 and x_2 to x_3 respectively.*
- (ii) *Let $(\Phi_0, \Phi): \mathfrak{R}_1 = (R_1, \eta_1, U_1, f_1, i_1) \hookrightarrow \mathfrak{R}_2 = (R_2, \eta_2, U_2, f_2, i_2)$ be an embedding of étale d-critical charts. Then for each $x_1 \in R_1$, there exist morphisms $h_1: U'_1 \rightarrow U_1$ which contains x_1 in its image, $h_2: U'_2 \rightarrow U_2$, $\Phi': U'_1 \rightarrow U'_2$, $\alpha: U'_2 \rightarrow U_1$ and $\beta: U'_2 \rightarrow \mathbb{A}^n$ where $n = \dim U_1 - \dim U_2$, h_1, h_2 and $(\alpha, \beta): U'_2 \rightarrow U_1 \times \mathbb{A}^n$ are étale, $h_2 \circ \Phi' = \Phi \circ h_1$, $\alpha \circ \Phi' = h_1$, $\beta \circ \Phi' = 0$ and $f_2 \circ h_2 = f_1 \circ \alpha + (z_1^2 + \dots + z_n^2) \circ \beta$ hold.*

In particular, there exists an étale cover of d-critical charts $\mathfrak{R}'_2 \rightarrow \mathfrak{R}_2$ containing $\Phi_0(x_1)$ in its image and an étale morphism of d-critical charts $\mathfrak{R}'_2 \rightarrow (R_1, \eta_1, U_1 \times \mathbb{A}^n, f_1 \boxplus (z_1^2 + \dots + z_n^2), (i_1, 0))$.

Proof. The second statement is nothing but [Joy15, Theorem 2.23]. To prove the first statement, by replacing R_1 and R_2 by $R_1 \times_X R_2$ and using Lemma 3.24, we may assume that $R_1 = R_2$ holds. In this case, the statement follows from [Joy15, Theorem 2.20]. \square

The following statement is useful to understand the behaviour of d-critical structures under smooth morphisms, which can be found in the proof of [Joy15, Proposition 2.8]:

Proposition 3.26. *Let $h: X_1 \rightarrow X_2$ be a smooth morphism of algebraic spaces. Let s_2 be a d-critical structure and set $s_1 := h^* s_2$. Then for each point x_1 with $h(x_1) = x_2$, we can find an étale d-critical chart $(R_1, \eta_1, U_1, f_1, i_1)$ of x_1 , $(R_2, \eta_2, U_2, f_2, i_2)$ of x_2 , smooth morphisms $H_0: R_1 \rightarrow R_2$ and $H: U_1 \rightarrow U_2$ such that $\eta_2 \circ H_0 = h \circ \eta_1$, $f_2 \circ H = f_1$ and $i_2 \circ H_0 = H \circ i_1$.*

We define a *smooth morphism of étale d-critical charts* to be the pair of morphisms (H_0, H) as in Proposition 3.26.

Finally, we introduce the external sum of d-critical structures. Let \mathfrak{X} and \mathfrak{Y} be Artin stacks and pr_i denotes the i -th projection from $\mathfrak{X} \times \mathfrak{Y}$. For a section $s \in \Gamma(\mathfrak{X}, \mathcal{S}_{\mathfrak{X}}^0)$ and $t \in \Gamma(\mathfrak{Y}, \mathcal{S}_{\mathfrak{Y}}^0)$, we set

$$s \boxplus t := \text{pr}_1^* s + \text{pr}_2^* t.$$

Lemma 3.27. *If s and t are d-critical structures, $s \boxplus t$ is a d-critical structure for $\mathfrak{X} \times \mathfrak{Y}$.*

Proof. Since being a d-critical structure is a local condition, we may assume that \mathfrak{X} (resp. \mathfrak{Y}) is a d-critical algebraic space covered by a single étale d-critical chart $\mathfrak{R} = (\mathfrak{X}, \text{id}, U, f, i)$ (resp. $(\mathfrak{S} = (\mathfrak{Y}, \text{id}, V, g, j))$). Then the claim follows from the identity

$$s \boxplus t = (f \boxplus g) + I_{R \times S, U \times V}^2 \in \Gamma(\mathfrak{X} \times \mathfrak{Y}, \mathcal{S}_{\mathfrak{X} \times \mathfrak{Y}}^0).$$

□

3.4. Torus-equivariant d-critical spaces. Here we will discuss torus equivariant d-critical algebraic spaces.

Definition 3.28. Let T be a torus and X be an algebraic space with a T -action μ .

- A *T -invariant d-critical structure on X* is a d-critical structure on X pulled back from $[X/T]$.
- A *T -equivariant d-critical algebraic space* is a triple (X, μ, s) of algebraic spaces X with T -action μ together with a T -invariant d-critical structure s .
- A *T -equivariant étale d-critical chart* is a tuple (R, η, U, f, i) where R is a quasi-separated scheme with a T -action, $\eta: R \rightarrow X$ is a T -equivariant étale morphism, U is a smooth scheme with a T -action, f is a T -invariant function on U such that $f|_{\text{Crit}(f)^{\text{red}}} = 0$ and $i: R \hookrightarrow U$ is a T -equivariant closed embedding such that $i(R) = \text{Crit}(f)$.

Proposition 3.29. *Let (X, μ, s) be a quasi-separated T -equivariant d-critical algebraic space. Then for each $x \in X$, there exists a T -equivariant étale d-critical chart (R, η, U, f, i) of X containing x such that $\dim T_{X,x} = \dim U$ holds.*

Proof. By the étale version of Sumihiro's theorem proved in [AHR20, Theorem 4.1], there exists a T -equivariant affine scheme R and T -equivariant étale morphism $R \rightarrow X$ whose image contains x . Then the claim follows from [Joy15, Proposition 2.43]. □

For an algebraic space X with a T -action μ , we denote by $X^\mu \subset X$ the fixed point subspace.

Corollary 3.30. *Let X be a quasi-separated algebraic space with a T -action μ and s be a T -invariant d-critical structure on X . Then $s^\mu := (X^\mu \hookrightarrow X)^* s$ is a d-critical structure.*

Proof. We may assume that X is covered by a single T -equivariant étale d-critical chart (R, η, U, f, i) . Then the tuple $(R^T, \eta^T, U^T, f|_{U^T}, i^T)$ defines an étale d-critical chart for (X^μ, s^μ) . □

The notion of T -equivariant embedding of T -equivariant d-critical structures is defined in the same manner as the non-equivariant case. We have the following T -equivariant version of Theorem 3.25:

Theorem 3.31. *Let X be a quasi-separated algebraic space with a T -action μ and s be a T -invariant d-critical structure.*

- (i) *Let $\mathfrak{R}_1 = (R_1, \eta_1, U_1, f_1, i_1)$ and $\mathfrak{R}_2 = (R_2, \eta_2, U_2, f_2, i_2)$ be two étale T -equivariant d-critical charts and $x_1 \in R_1^T$ and $x_2 \in R_2^T$ be points such that $\eta_1(x_1) = \eta_2(x_2)$ holds. Then by possibly replacing \mathfrak{R}_1 and \mathfrak{R}_2 by T -equivariant étale neighbourhoods of x_1 and x_2 , we may find a third T -equivariant d-critical chart $\mathfrak{R}_3 = (R_3, \eta_3, U_3, f_3, i_3)$ and $x_3 \in R_3$ such that there exists T -equivariant embeddings $\mathfrak{R}_1 \hookrightarrow \mathfrak{R}_3$ and $\mathfrak{R}_2 \hookrightarrow \mathfrak{R}_3$ which map x_1 to x_3 and x_2 to x_3 respectively.*
- (ii) *Let $(\Phi_0, \Phi): (R_1, \eta_1, U_1, f_1, i_1) \hookrightarrow (R_2, \eta_2, U_2, f_2, i_2)$ be a T -equivariant embedding of T -equivariant étale d-critical charts. Then for each $x_1 \in R_1$, there exist T -equivariant morphisms $\eta_1: U'_1 \rightarrow U_1$ containing x_1 in its image, $\eta_2: U'_2 \rightarrow U_2$, $\Phi': U'_1 \rightarrow U'_2$, $\alpha: U'_2 \rightarrow U_1$, $\beta: U'_2 \rightarrow \mathbb{A}^n$ and a T -invariant non-degenerate bilinear form q on \mathbb{A}^n where $n = \dim U_1 - \dim U_2$, \mathbb{A}^n is equipped with some linear*

T -action, η_1, η_2 and $(\alpha, \beta): U'_2 \rightarrow U_1 \times \mathbb{A}^n$ are étale, $\eta_2 \circ \Phi' = \Phi \circ \eta_1$, $\alpha \circ \Phi' = \eta_1$, $\beta \circ \Phi' = 0$ and $f_2 \circ \eta_2 = f_1 \circ \alpha + q \circ \beta$ hold.

In particular, there exists a T -equivariant étale cover of d -critical charts $\mathfrak{X}'_2 \rightarrow \mathfrak{X}_2$ containing $\Phi_0(x_1)$ in its image and an étale morphism of d -critical charts $\mathfrak{X}'_2 \rightarrow (R_1, \eta_1, U_1 \times \mathbb{A}^n, f_1 \boxplus q, (i_1, 0))$ where q is a T -invariant non-degenerate quadratic function.

Proof. Note that we can easily make each choice in Lemma 3.24 T -equivariant. Therefore, the first claim follows by arguing as the proof of Theorem 3.25 using [Joy15, Proposition 2.44]. The latter claim follows by arguing as the proof of [Joy15, Proposition 2.23]. See [Des22, Lemma 5.1] for the proof of the corresponding statement in the analytic situation. \square

Assume that we are given algebraic spaces X_1 and X_2 with T -actions μ_1 and μ_2 , T -equivariant smooth morphism $h: X_1 \rightarrow X_2$ which fit into a commutative diagram of fixed point spaces

$$\begin{array}{ccc} X_1^{\mu_1} & \longrightarrow & X_1 \\ \downarrow h^T & & \downarrow h \\ X_2^{\mu_2} & \longrightarrow & X_2. \end{array}$$

Consider a d -critical structure s_2 of X_2 and set $s_1 := h^*s_2$. Then by construction, we have the identity

$$(3.11) \quad h^{T,*}(s_2^{\mu_2}) = s_1^{\mu_1}.$$

The following T -equivariant version of Proposition 3.26 will be useful to compare T -invariant d -critical structures related by a T -equivariant smooth morphism. See [Des22, Lemma 5.2] for the proof:

Proposition 3.32. *Let $h: X_1 \rightarrow X_2$ be a smooth T -equivariant morphism of quasi-separated algebraic spaces with T -actions. Let s_2 be a T -invariant d -critical structure and set $s_1 := h^*s_2$. Then for each point x_1 with $h(x_1) = x_2$, there exists a T -equivariant étale d -critical chart $(R_1, \eta_1, U_1, f_1, i_1)$ of x_1 , $(R_2, \eta_2, U_2, f_2, i_2)$ of x_2 , smooth T -equivariant morphisms $H_0: R_1 \rightarrow R_2$ and $H: U_1 \rightarrow U_2$ such that $\eta_2 \circ H_0 = h \circ \eta_1$, $f_2 \circ H = f_1$ and $i_2 \circ H_0 = H \circ i_1$.*

3.5. Virtual canonical bundles and orientations. We now explain the definition of the virtual canonical bundle for d -critical algebraic spaces and stacks, following [Joy15, §2.4]. Firstly, we will introduce the notion of the virtual cotangent complexes for étale d -critical charts:

Definition 3.33. For an étale d -critical chart $\mathfrak{X} = (R, \eta, U, f, i)$ of a d -critical algebraic space (X, s) , we define the *virtual cotangent complex* $\mathbb{L}_{\mathfrak{X}} \in \text{Perf}(R)$ by

$$\mathbb{L}_{\mathfrak{X}} := [T_U|_R \xrightarrow{\text{Hess}_f} \Omega_U|_R]$$

concentrated in degree $[-1, 0]$.

Let (\mathfrak{X}, s) be a d -critical Artin stack. Joyce [Joy15, Theorem 2.56] constructed a line bundle

$$K_{\mathfrak{X}, s} \in \text{Pic}(\mathfrak{X}^{\text{red}})$$

called the *virtual canonical bundle* by gluing $\widehat{\det}(\mathbb{L}_{\mathfrak{X}})$ for each étale d -critical charts $\mathfrak{X} = (R, \eta, U, f, i)$ of smooth covers of \mathfrak{X} . Throughout the paper, we equip $K_{\mathfrak{X}, s}$ with the trivial grading. The following proposition summarizes the properties of the virtual canonical bundle that we will use later:

Proposition 3.34. *The virtual canonical bundle $K_{\mathfrak{X}, s}$ for a d -critical stack (\mathfrak{X}, s) has the following properties:*

(i) *If \mathfrak{X} is an algebraic space and \mathfrak{X} is an étale d -critical chart of (\mathfrak{X}, s) , there exists a natural isomorphism*

$$(3.12) \quad \iota_{\mathfrak{X}}: K_{\mathfrak{X}, s}|_{R^{\text{red}}} \simeq \widehat{\det}(\mathbb{L}_{\mathfrak{X}}).$$

(ii) *For each point $x \in \mathfrak{X}$, there exists a natural isomorphism*

$$(3.13) \quad \kappa_x: K_{\mathfrak{X}, s}|_x \simeq \det(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, x}))^{\otimes 2}.$$

(iii) For a smooth morphism $q: X \rightarrow \mathfrak{X}$ from an algebraic space X , there exists a natural isomorphism

$$(3.14) \quad \Upsilon_q: q^{\text{red},*} K_{\mathfrak{X},s} \otimes K_{X/\mathfrak{X}}^{\otimes 2} \cong K_{X,q^*s}.$$

(iv) Assume that we are given another d -critical stack (\mathfrak{Y}, t) . Then there exists a natural isomorphism

$$(3.15) \quad \Psi_{\mathfrak{X},\mathfrak{Y}}: K_{\mathfrak{X},s} \boxtimes K_{\mathfrak{Y},t} \cong K_{\mathfrak{X} \times \mathfrak{Y}, s \boxplus t}.$$

The properties (i) – (iii) are proved in [Joy15, Theorem 2.56]. Concerning the construction of $\Psi_{\mathfrak{X},\mathfrak{Y}}$, see the discussion after (3.20).

For later use, we discuss compatibility between isomorphisms in Proposition 3.34. Assume first that \mathfrak{X} is an algebraic space and take an étale d -critical chart $\mathfrak{R} = (R, \eta, U, f, i)$. Then for each $x \in \mathfrak{X}$ with a lift $\hat{x} \in R$, the map κ_x in (3.13) is compatible with $\iota_{\mathfrak{R}}$, i.e., the following diagram commutes:

$$(3.16) \quad \begin{array}{ccc} K_{\mathfrak{X},s}|_x & \xrightarrow{\kappa_x} & \det(\Omega_{\mathfrak{X},x})^{\otimes 2} \\ & \searrow \iota_{\mathfrak{R}}|_x & \downarrow \cong \\ & & \det(\mathbb{L}_{\mathfrak{R}}|_{\hat{x}}). \end{array}$$

Here the right vertical isomorphism is constructed using the fibre sequence

$$(3.17) \quad T_{\mathfrak{X},x}[1] \rightarrow \mathbb{L}_{\mathfrak{R}}|_{\hat{x}} \rightarrow \Omega_{\mathfrak{X},x}$$

together with (2.5) and (2.14).

Let (\mathfrak{X}, s) be a d -critical stack and $q: X \rightarrow \mathfrak{X}$ be a smooth morphism from an algebraic space. Then the map Υ_q in (3.14) is the unique morphism such that the following diagram commutes for each $\hat{x} \in X$ with $x = q(\hat{x})$:

$$(3.18) \quad \begin{array}{ccc} q^{\text{red},*} K_{\mathfrak{X},s}|_x \otimes K_{X/\mathfrak{X}}|_{\hat{x}}^{\otimes 2} & \xrightarrow[\cong]{\Upsilon_q} & K_{X,q^*s}|_{\hat{x}} \\ \downarrow \kappa_x \otimes \text{id} & & \downarrow \cong \kappa_{\hat{x}} \\ \det(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},\hat{x}}))^{\otimes 2} \otimes K_{X/\mathfrak{X}}|_{\hat{x}}^{\otimes 2} & \xrightarrow[\cong]{} & \det(\Omega_{X,\hat{x}})^{\otimes 2} \end{array}$$

where the bottom horizontal arrow is induced by the fibre sequence $\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},x}) \rightarrow \Omega_{X,\hat{x}} \rightarrow \Omega_{X/\mathfrak{X}}|_{\hat{x}}$. The map Υ_h is associative in the following sense: if we are given a smooth morphism $h: X_1 \rightarrow X_2$ between algebraic spaces and a smooth morphism $q_2: X_2 \rightarrow \mathfrak{X}$, the following diagram commutes:

$$(3.19) \quad \begin{array}{ccc} h^{\text{red},*} q_2^{\text{red},*} K_{\mathfrak{X},s} \otimes h^{\text{red},*} K_{X_2/\mathfrak{X}}^{\otimes 2} \otimes K_{X_1/X_2}^{\otimes 2} & \xrightarrow[\cong]{h^{\text{red},*} \Upsilon_{q_2}} & h^{\text{red},*} K_{X_2,s_2} \otimes K_{X_1/X_2}^{\otimes 2} \\ \cong \downarrow & & \cong \downarrow \Upsilon_h \\ q_1^{\text{red},*} K_{\mathfrak{X},s} \otimes K_{X_1/\mathfrak{X}}^{\otimes 2} & \xrightarrow[\cong]{\Upsilon_{q_1}} & K_{X_1,s_1} \end{array}$$

where we write $q_1 = q_2 \circ h$.

Let (\mathfrak{X}, s) and (\mathfrak{Y}, t) be d -critical algebraic stacks. The map $\Psi_{\mathfrak{X},\mathfrak{Y}}$ in (3.15) is uniquely characterized by the property that the following diagram commutes for each $x \in \mathfrak{X}$ and $y \in \mathfrak{Y}$:

$$(3.20) \quad \begin{array}{ccc} K_{\mathfrak{X},s}|_x \otimes K_{\mathfrak{Y},t}|_y & \xrightarrow[\cong]{(\Psi_{\mathfrak{X} \times \mathfrak{Y}})|_{(x,y)}} & (K_{\mathfrak{X},s} \otimes K_{\mathfrak{Y},t})|_{(x,y)} \\ \cong \downarrow \kappa_x \otimes \kappa_y & & \cong \downarrow \kappa_{(x,y)} \\ \widehat{\det}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},x}))^{\otimes 2} \otimes \widehat{\det}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{Y},y}))^{\otimes 2} & \xrightarrow[\cong]{} & \widehat{\det}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},x} \oplus \mathbb{L}_{\mathfrak{Y},y}))^{\otimes 2}. \end{array}$$

To prove the existence of $\Psi_{\mathfrak{X},\mathfrak{Y}}$ with the above commutativity property, using the commutativity of (3.18), we may assume that \mathfrak{X} and \mathfrak{Y} are algebraic spaces covered by single critical charts $\mathfrak{R} = (R, \text{id}, U, f, i)$ and $\mathfrak{S} = (S, \text{id}, V, g, j)$. In this case, the isomorphism $\mathbb{L}_{\mathfrak{R}} \boxplus \mathbb{L}_{\mathfrak{S}} \simeq \mathbb{L}_{\mathfrak{R} \times \mathfrak{S}}$ induces an isomorphism $\widehat{\det}(\mathbb{L}_{\mathfrak{R}}) \boxtimes \widehat{\det}(\mathbb{L}_{\mathfrak{S}}) \cong \widehat{\det}(\mathbb{L}_{\mathfrak{R} \times \mathfrak{S}})$. Using (3.16), we see that it induces the isomorphism $K_{\mathfrak{X},s} \boxtimes K_{\mathfrak{Y},t} \cong K_{\mathfrak{X} \times \mathfrak{Y}, s \boxplus t}$ with the desired property.

Assume that we are given smooth morphisms $q: X \rightarrow \mathfrak{X}$ and $r: Y \rightarrow \mathfrak{Y}$ from algebraic spaces and given d-critical structures s of \mathfrak{X} and t of \mathfrak{Y} . Then the commutativity of the diagrams (3.18) and (3.20) implies the commutativity of the following diagram:

$$(3.21) \quad \begin{array}{ccc} (q^{\text{red},*} K_{\mathfrak{X},s} \otimes K_{X/\mathfrak{X}}) \boxtimes (r^{\text{red},*} K_{\mathfrak{Y},t} \otimes K_{Y/\mathfrak{Y}}) & \xrightarrow[\cong]{\Upsilon_q \boxtimes \Upsilon_r} & K_{X,q^*s} \boxtimes K_{Y,r^*t} \\ \cong \downarrow \Psi_{\mathfrak{X},\mathfrak{Y}} & & \cong \downarrow \Psi_{X,Y} \\ (q \times r)^* K_{\mathfrak{X} \times \mathfrak{Y}, s \boxplus t} \otimes K_{X \times Y/\mathfrak{X} \times \mathfrak{Y}} & \xrightarrow[\cong]{\Upsilon_{q \times r}} & K_{X \times Y, q^*s \boxplus r^*t} \end{array}$$

We now describe the virtual canonical bundle for underlying d-critical stacks of (-1) -shifted symplectic stacks. Let (X, ω_X) be a (-1) -shifted symplectic algebraic space and (X^{cl}, s) be its underlying d-critical algebraic space. For an étale derived critical chart $\mathfrak{R} = (R, \eta, U, f, i)$, we have a natural equivalence

$$(3.22) \quad \mathbb{L}_{\mathfrak{X}}|_{R^{\text{red}}} \simeq \mathbb{L}_{\mathfrak{R}^{\text{cl}}}$$

where \mathfrak{R}^{cl} denote the underlying étale d-critical chart. Under this identification, the natural equivalence $\mathbb{T}_X|_{R^{\text{red}}} \simeq \mathbb{L}_X|_{R^{\text{red}}}[-1]$ induced from the (-1) -shifted symplectic structure corresponds to the map of complexes²

$$(3.23) \quad \begin{array}{ccc} \mathbb{T}_X|_{R^{\text{red}}} & \simeq & [T_U|_{R^{\text{red}}} \xrightarrow{-\text{Hess}(f)} \Omega_U|_{R^{\text{red}}}] \\ \downarrow \cdot \omega_X|_{R^{\text{red}}} & & \downarrow \text{id} \qquad \qquad \downarrow \text{id} \\ \mathbb{L}_X[-1]|_{R^{\text{red}}} & \simeq & [T_U|_{R^{\text{red}}} \xrightarrow{-\text{Hess}(f)} \Omega_U|_{R^{\text{red}}}] \end{array}$$

The equivalence (3.22) implies an isomorphism

$$\Lambda_{X,\mathfrak{R}}: \widehat{\det}(\mathbb{L}_X)|_{R^{\text{red}}} \cong K_{X^{\text{cl}},s}|_{R^{\text{red}}}.$$

We will show that this isomorphism glues to define a global isomorphism. To see this, for a point $x \in X$ we construct an isomorphism

$$\kappa_x^{\text{der}}: \det(\mathbb{L}_{X,x}) \cong \det(\mathbb{L}_{X^{\text{cl}},x})^{\otimes 2}$$

by the following composition

$$\begin{aligned} \det(\mathbb{L}_{X,x}) &\cong \det(\tau^{\leq -1}(\mathbb{L}_{X,x})) \otimes \det(\tau^{\geq 0}(\mathbb{L}_{X,x})) \cong \det(\tau^{\geq 0}(\mathbb{L}_{X,x})^\vee[1]) \otimes \det(\tau^{\geq 0}(\mathbb{L}_{X,x})) \\ &\cong \det(\tau^{\geq 0}(\mathbb{L}_{X,x}))^{\otimes 2}. \end{aligned}$$

Here the second isomorphism is constructed using the isomorphism $(-\cdot \omega_X)^{-1}|_x: \tau^{\leq -1}(\mathbb{L}_{X,x}) \simeq \tau^{\geq 0}(\mathbb{L}_{X,x})^\vee[1]$ and the third isomorphisms from (2.5) and (2.14). By the construction and the description of the symplectic structure in (3.23), we see that the following diagram commutes:

$$(3.24) \quad \begin{array}{ccc} \det(\mathbb{L}_{X,x}) & \xrightarrow[\cong]{\Lambda_{X,\mathfrak{R}}|_x} & K_{X^{\text{cl}},s} \\ \kappa_x^{\text{der}} \downarrow \cong & & \cong \downarrow \kappa_x \\ \det(\tau^{\geq 0}(\mathbb{L}_{X,x}))^{\otimes 2} & \xlongequal{\quad\quad\quad} & \det(\tau^{\geq 0}(\mathbb{L}_{X,x}))^{\otimes 2} \end{array}$$

In particular, the map $\Lambda_{X,\mathfrak{R}}$ does not depend on the choice of \mathfrak{R} at each point, hence it glues to define an isomorphism

$$\Lambda_X: \widehat{\det}(\mathbb{L}_X) \cong K_{X^{\text{cl}},s}.$$

Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a (-1) -shifted symplectic stack and $(\mathfrak{X}^{\text{cl}}, s)$ be the underlying d-critical stack. For a point $x \in \mathfrak{X}$, we define an isomorphism

$$\kappa_x^{\text{der}}: \det(\mathbb{L}_{\mathfrak{X},x}) \cong \det(\mathbb{L}_{\mathfrak{X}^{\text{cl}},x})^{\otimes 2}$$

²See [Con00, §1.3] for the sign convention on derived categories. In particular, the appearance of the sign in the top horizontal map follows from the sign convention of the dual complex, and the fact that the right vertical map is the identity follows from (1.3.6) in loc.cit. and the explicit description of the (-1) -shifted symplectic structure on the critical locus in [BBJ19, Example 5.15].

exactly in the same manner as for algebraic spaces. We have an isomorphism of one-dimensional vector spaces

$$\Lambda_{\mathfrak{X},x}^{\text{pre}} : \widehat{\det}(\mathbb{L}_{\mathfrak{X},x}) \cong K_{\mathfrak{X}^{\text{cl}},s}|_x$$

given by $\kappa_x^{-1} \circ \kappa_x^{\text{der}}$. The following statement is proved in [Ben+15, Theorem 3.18] (see also [Kin22, Theorem 4.9] for a proof with careful choices of sign and Proposition 6.9 for a closely related statement):

Lemma 3.35. [Ben+15, Theorem 3.18] *The map $x \mapsto \Lambda_{\mathfrak{X},x}^{\text{pre}}$ is algebraic and hence defines an isomorphism of line bundles*

$$\Lambda_{\mathfrak{X}} : \widehat{\det}(\mathbb{L}_{\mathfrak{X}}) \cong K_{\mathfrak{X}^{\text{cl}},s}.$$

By the construction of the map $\Lambda_{\mathfrak{X}} : \widehat{\det}(\mathbb{L}_{\mathfrak{X}}) \cong K_{\mathfrak{X}^{\text{cl}},s}$, the following diagram commutes:

$$(3.25) \quad \begin{array}{ccc} \det(\mathbb{L}_{\mathfrak{X},x}) & \xrightarrow[\cong]{\Lambda_{\mathfrak{X}}|_x} & K_{\mathfrak{X}^{\text{cl}},s}|_x \\ \cong \downarrow \kappa_x^{\text{der}} & & \cong \downarrow \kappa_x \\ \det(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},x}))^{\otimes 2} & \xlongequal{\quad\quad\quad} & \det(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},x}))^{\otimes 2}. \end{array}$$

The map $\Lambda_{\mathfrak{X}}$ is compatible with the isomorphism $\Psi_{\mathfrak{X},\mathfrak{Y}}$: Namely, assume that we are given (-1) -shifted symplectic stacks $(\mathfrak{X}, \omega_{\mathfrak{X}})$ and $(\mathfrak{Y}, \omega_{\mathfrak{Y}})$ with underlying d-critical stacks $K_{\mathfrak{X}^{\text{cl}},s}$ and $K_{\mathfrak{Y}^{\text{cl}},t}$. Then it follows from the commutativity of diagrams (3.20) and (3.25) that the following diagram commutes:

$$(3.26) \quad \begin{array}{ccc} \widehat{\det}(\mathbb{L}_{\mathfrak{X}}) \boxtimes \widehat{\det}(\mathbb{L}_{\mathfrak{Y}}) & \xrightarrow[\cong]{} & \widehat{\det}(\mathbb{L}_{\mathfrak{X} \times \mathfrak{Y}}) \\ \cong \downarrow \Lambda_{\mathfrak{X}} \boxtimes \Lambda_{\mathfrak{Y}} & & \cong \downarrow \Lambda_{\mathfrak{X} \times \mathfrak{Y}} \\ K_{\mathfrak{X}^{\text{cl}},s} \boxtimes K_{\mathfrak{Y}^{\text{cl}},t} & \xrightarrow[\cong]{\Upsilon_{\mathfrak{X}^{\text{cl}},\mathfrak{Y}^{\text{cl}}}} & K_{\mathfrak{X}^{\text{cl}} \times \mathfrak{Y}^{\text{cl}},s \boxplus t}. \end{array}$$

3.6. Orientation for d-critical stacks. Let (\mathfrak{X}, s) be a d-critical stack. An *orientation of* (\mathfrak{X}, s) is a choice of a graded line bundle $\mathcal{L} \in \text{Pic}^{\text{gr}}(\mathfrak{X}^{\text{red}})$ and an isomorphism $\mathcal{L}^{\otimes 2} \cong K_{\mathfrak{X},s}$. An isomorphism of orientations is defined in the same manner as Definition 3.16. By abuse of notation, we simply say that o is an orientation for (\mathfrak{X}, s) . Let $q: X \rightarrow \mathfrak{X}$ be a smooth morphism from an algebraic space X . For an orientation $o: \mathcal{L}^{\otimes 2} \cong K_{\mathfrak{X},s}$, we define an orientation q^*o by the composition

$$q^*o: (\mathcal{L} \otimes \widehat{\det}(\Omega_{X/\mathfrak{X}}))^{\otimes 2} \xrightarrow{\cong} K_{\mathfrak{X},s} \otimes \widehat{\det}(\Omega_{X/\mathfrak{X}})^{\otimes 2} \xrightarrow[\cong]{\Upsilon_h} K_{X,q^*s}.$$

Assume further that we are given a smooth morphism $h: Y \rightarrow X$. Then the diagram (3.19) implies a natural equivalence

$$(3.27) \quad h^*q^*o \cong (q \circ h)^*o.$$

Let (\mathfrak{X}, s) and (\mathfrak{Y}, t) be d-critical stacks equipped with orientations $o_{\mathfrak{X}}: \mathcal{L}_{\mathfrak{X}}^{\otimes 2} \cong K_{\mathfrak{X},s}$ and $o_{\mathfrak{Y}}: \mathcal{L}_{\mathfrak{Y}}^{\otimes 2} \cong K_{\mathfrak{Y},t}$. Then we define an orientation $o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}}$ by

$$(3.28) \quad o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}}: (\mathcal{L}_{\mathfrak{X}} \boxtimes \mathcal{L}_{\mathfrak{Y}})^{\otimes 2} \xrightarrow[\cong]{o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}}} K_{\mathfrak{X},s} \boxtimes K_{\mathfrak{Y},t} \xrightarrow[\cong]{\Psi_{\mathfrak{X},\mathfrak{Y}}} K_{\mathfrak{X} \times \mathfrak{Y},s \boxplus t}.$$

Assume further that we are given another oriented d-critical stack $(\mathfrak{Z}, u, o_{\mathfrak{Z}})$. Then it easily follows from (3.20) that the isomorphism $\Psi_{\mathfrak{X},\mathfrak{Y}}$ is associative hence there exists a natural isomorphism

$$(3.29) \quad (o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}}) \boxtimes o_{\mathfrak{Z}} \cong o_{\mathfrak{X}} \boxtimes (o_{\mathfrak{Y}} \boxtimes o_{\mathfrak{Z}}).$$

Assume that we are given smooth morphisms $q: X \rightarrow \mathfrak{X}$ and $r: Y \rightarrow \mathfrak{Y}$. Then it follows from (3.21) that there exists a natural isomorphism

$$(3.30) \quad q^*o_{\mathfrak{X}} \boxtimes r^*o_{\mathfrak{Y}} \cong (q \times r)^*o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}}.$$

Let $(\mathfrak{X}, \omega_{\mathfrak{X}}, o)$ be an oriented (-1) -shifted symplectic stack with underlying d-critical structure s . By using Lemma 3.35, o induces an orientation for $(\mathfrak{X}^{\text{cl}}, s)$ which is denoted by o^{cl} . The formation of classical

truncation of orientations commutes with the product: Namely, given oriented (-1) -shifted symplectic stacks $(\mathfrak{X}, \omega_{\mathfrak{X}}, o_{\mathfrak{X}})$ and $(\mathfrak{Y}, \omega_{\mathfrak{Y}}, o_{\mathfrak{Y}})$, there exists a natural isomorphism

$$(3.31) \quad (o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}})^{\text{cl}} \cong o_{\mathfrak{X}}^{\text{cl}} \boxtimes o_{\mathfrak{Y}}^{\text{cl}}.$$

This is a consequence of the commutativity of the diagram (3.26). Further, it is clear from the construction that this isomorphism is associative under the identification (3.29).

We now introduce a d-critical version of the canonical orientation discussed in Example 3.20. Let U be a smooth scheme, $f: U \rightarrow \mathbb{A}^1$ be a regular function with $f|_{\text{Crit}(f)^{\text{red}}} = 0$ and $X = \text{Crit}(f)^{\text{cl}}$. Set $s := f + I_{X/U}^2$ and $\mathfrak{X}^{\text{taut}} = (X, \text{id}, U, f, i)$ be the tautological étale d-critical chart. Then the fibre sequence

$$(3.32) \quad \Omega_U|_{X^{\text{red}}} \rightarrow \mathbb{L}_{\mathfrak{X}^{\text{taut}}} \rightarrow T_U|_{X^{\text{red}}}[1]$$

together with (2.5) and (2.14) induce an isomorphism $\bar{o}_X^{\text{can}}: K_U|_{X^{\text{red}}}^{\otimes 2} \cong K_{X,s}$. We set

$$(3.33) \quad o_X^{\text{can}} := (-1)^{\dim U} \cdot \bar{o}_X^{\text{can}}$$

and call it the **canonical orientation** for (X, s) with respect to the critical locus description $X = \text{Crit}(f)^{\text{cl}}$. This is compatible with Example 3.20: Namely, there exists a natural isomorphism

$$(3.34) \quad (o_{\text{Crit}(f)}^{\text{can}})^{\text{cl}} \cong o_{\text{Crit}(f)^{\text{cl}}}^{\text{can}}.$$

Remark 3.36. The sign insertion seems to be unnatural, but this choice is more convenient for us in that it reduces the number of signs appearing in the paper. This is because, we did and will control the behaviour of the virtual canonical bundles using the isomorphism κ_x , and it is defined using the fibre sequence (3.17). On the other hand, the isomorphism \bar{o}_X^{can} is constructed using the fibre sequence (3.32). Therefore, it is natural to swap K_U when defining the canonical orientation, which corresponds to multiplying $(-1)^{\dim U}$ since we twist the sign of the symmetrizer: see §2.2. As an example, if we do not insert this sign, the isomorphism of underlying line bundles in (3.35) needs to be modified by $\sqrt{-1}^{\pm \dim \Omega_{U_1/U_2}}$ to define an isomorphism of orientations.

Assume that we are given a smooth morphism $H: U_1 \rightarrow U_2$ between smooth schemes and a regular function f_2 on U_2 with $f_2|_{\text{Crit}(f_2)^{\text{red}}} = 0$. Write $f_1 := f_2 \circ H$. Set $X_1 := \text{Crit}(f_1)$ and $X_2 := \text{Crit}(f_2)$ equipped with the natural d-critical structures and $h: X_1 \rightarrow X_2$ be the restriction of H . Then we see that the natural morphism $H^*K_{U_2} \otimes K_{U_1/U_2} \cong K_{U_1}$ induces an isomorphism

$$(3.35) \quad o_{X_1}^{\text{can}} \cong h^* o_{X_2}^{\text{can}}.$$

See the proof of the commutativity of the diagram (6.16) for a related discussion. This diagram is associative in the following sense: Assume further that we are given a smooth morphism $G: U_0 \rightarrow U_1$, set $X_0 = \text{Crit}(f_1 \circ G)$ and $g: X_0 \rightarrow X_1$ be the restriction of G . Then the following diagram commutes:

$$(3.36) \quad \begin{array}{ccc} o_{X_0}^{\text{can}} & \xrightarrow[\cong]{(3.35)} & g^* o_{X_1}^{\text{can}} \\ \cong \downarrow (3.35) & & \cong \downarrow (3.35) \\ (h \circ g)^* o_{X_2}^{\text{can}} & \xrightarrow[\cong]{(3.27)} & g^* h^* o_{X_2}^{\text{can}} \end{array}$$

Assume now that we are given smooth schemes U and V and regular functions f on U and g on V with $f|_{U^{\text{red}}} = 0$ and $g|_{V^{\text{red}}} = 0$. Set $X = \text{Crit}(f)$ and $Y = \text{Crit}(g)$ equipped with the natural d-critical structures. Then there exists a natural isomorphism

$$(3.37) \quad o_X^{\text{can}} \boxtimes o_Y^{\text{can}} \cong o_{X \times Y}^{\text{can}}.$$

Assume further that we are given a d-critical scheme $Z = \text{Crit}(h)$. Then by construction, the following diagram commutes:

$$(3.38) \quad \begin{array}{ccc} (o_X^{\text{can}} \boxtimes o_Y^{\text{can}}) \boxtimes o_Z^{\text{can}} & \xrightarrow[\cong]{(3.29)} & o_X^{\text{can}} \boxtimes (o_Y^{\text{can}} \boxtimes o_Z^{\text{can}}) \\ \cong \downarrow (3.37) & & \cong \downarrow (3.37) \\ o_{X \times Y}^{\text{can}} \boxtimes o_Z^{\text{can}} & \xrightarrow[\cong]{(3.37)} o_{X \times Y \times Z}^{\text{can}} \xleftarrow[\cong]{(3.37)} & o_X^{\text{can}} \boxtimes o_{Y \times Z}^{\text{can}} \end{array}$$

Assume that we are given smooth morphisms $H_U: U' \rightarrow U$ and $H_V: V' \rightarrow V$. Set $X' = \text{Crit}(f \circ H_U)$ and $Y' = \text{Crit}(g \circ H_V)$ and $h_X: X' \rightarrow X$ and $h_Y: Y' \rightarrow Y$ be the restrictions of H_U and H_V . Then the following diagram commutes:

$$(3.39) \quad \begin{array}{ccc} h_X^* o_X^{\text{can}} \boxtimes h_Y^* o_Y^{\text{can}} & \xrightarrow[\cong]{(3.35)} & o_{X'}^{\text{can}} \boxtimes o_{Y'}^{\text{can}} \\ \cong \downarrow (3.37) & & \cong \downarrow (3.37) \\ (h_X \times h_Y)^* o_{X \times Y}^{\text{can}} & \xrightarrow[\cong]{(3.35)} & o_{X' \times Y'}^{\text{can}} \end{array}$$

4. DONALDSON–THOMAS PERVERSE SHEAVES

Here we briefly recall the definition and some properties of the vanishing cycle functor (§4.1) and the construction of the Donaldson–Thomas perverse sheaves (§4.2) associated with oriented d-critical stacks following [Bra+15] which is a global version of the vanishing cycle complexes. We also prove in §4.3 the Thom–Sebastiani theorem for the Donaldson–Thomas perverse sheaves, whose detailed proof seems to be not available in the literature.

4.1. Vanishing cycle functors. In this section, we recall the theory of the vanishing cycle complexes. Let U be a smooth separated algebraic space and $f: U \rightarrow \mathbb{A}^1$ be a regular function. We set

$$U_{\leq 0} := f^{-1}(\mathbb{A}_{\leq 0}^1), \quad U_0 := f^{-1}(0)$$

where $\mathbb{A}_{\leq 0}^1 := \{z \in \mathbb{A}^1 \mid \text{Re}(z) \leq 0\}$. Define the vanishing cycle functor $\varphi_f: D(U) \rightarrow D(U_0)$ by

$$\varphi_f := (U_0 \hookrightarrow U_{\leq 0})^*(U_{\leq 0} \hookrightarrow U)^!$$

It is proved in [Ach21, Corollary 4.2.6, Theorem 4.2.8] that the functor φ_f preserves constructible complexes and perverse sheaves. For simplicity, we write

$$(4.1) \quad \varphi_f := \varphi_f(\mathbb{Q}_U[\dim U]).$$

One can easily show that the complex φ_f is supported on the critical locus.

The definition of the vanishing cycle functor can be extended to functions on an Artin stack (hence to non-separated spaces). Let \mathfrak{U} be a smooth Artin stack. We set $\mathfrak{U}_{>0} := f^{-1}(\mathbb{A}_{>0}^1)$, which is regarded as a complex analytic substack of \mathfrak{U} . Then we define the vanishing cycle functor by

$$\varphi_f := \text{fib}((\mathfrak{U}_0 \hookrightarrow \mathfrak{U})^* \rightarrow (\mathfrak{U}_0 \hookrightarrow \mathfrak{U})^*(\mathfrak{U}_{>0} \hookrightarrow \mathfrak{U})_*(\mathfrak{U}_{>0} \hookrightarrow \mathfrak{U})^*).$$

It is clear from the definition that this definition matches with the previous definition of the vanishing cycle functor for functions on a smooth separated scheme.

Assume that we are given a morphism $h: U \rightarrow V$ between smooth algebraic spaces and a regular function f on V . Set $f_0: U_0 \rightarrow V_0$ be the induced morphism on the zero locus. Then by construction, we have natural transforms

$$(4.2) \quad \begin{array}{ccc} h_{0,!} \circ \varphi_{f \circ h} \rightarrow \varphi_f \circ h_! & , & h_0^* \circ \varphi_f \rightarrow \varphi_{f \circ h} \circ h^* \\ \varphi_f \circ h_* \rightarrow h_{0,*} \circ \varphi_{f \circ h} & , & \varphi_{f \circ h} \circ h^! \rightarrow h_0^! \circ \varphi_f. \end{array}$$

The proper base change theorem implies that left maps are invertible if h is proper. The smooth base change theorem implies that right maps are invertible if h is smooth. The same base change properties hold more generally for morphisms of stacks.

Let $f: U \rightarrow \mathbb{A}^1$ and $g: V \rightarrow \mathbb{A}^1$ be regular functions on smooth separated algebraic spaces and $\mathcal{F} \in D_c^b(U)$ and $\mathcal{G} \in D_c^b(V)$ be constructible complexes. We define a function $f \boxplus g$ on $U \times V$ by $(f \boxplus g)(x, y) = f(x) + g(y)$. Then there exist natural morphisms

$$\begin{aligned} \varphi_f(\mathcal{F}) \boxtimes \varphi_g(\mathcal{G}) &= (U_0 \hookrightarrow U_{\leq 0})^*(U_{\leq 0} \hookrightarrow U)^! \mathcal{F} \boxtimes (V_0 \hookrightarrow V_{\leq 0})^*(V_{\leq 0} \hookrightarrow V)^! \mathcal{G} \\ &\rightarrow (U_0 \times V_0 \hookrightarrow U_{\leq 0} \times V_{\leq 0})^*(U_{\leq 0} \times V_{\leq 0} \hookrightarrow U \times V)^! (\mathcal{F} \boxtimes \mathcal{G}) \\ \varphi_{f \boxplus g}(\mathcal{F} \boxtimes \mathcal{G})|_{U_0 \times V_0} &\cong (U_0 \times V_0 \hookrightarrow (U \times V)_{\leq 0})^*((U \times V)_{\leq 0} \hookrightarrow U \times V)^! \\ &\leftarrow (U_0 \times V_0 \hookrightarrow U_{\leq 0} \times V_{\leq 0})^*(U_{\leq 0} \times V_{\leq 0} \hookrightarrow U \times V)^! (\mathcal{F} \boxtimes \mathcal{G}) \end{aligned}$$

which are shown to be isomorphisms in [Mas01, Proposition 1.4] and [Mas01, Lemma 1.2] respectively. In particular, we obtain an isomorphism

$$(4.3) \quad \text{TS: } \varphi_f(\mathcal{F}) \boxtimes \varphi_g(\mathcal{G}) \cong \varphi_{f \boxplus g}(\mathcal{F} \boxtimes \mathcal{G})|_{U_0 \times V_0}.$$

Assume further that we are given a regular function $h: W \rightarrow \mathbb{A}^1$ on a smooth separated algebraic space and a constructible complex $\mathcal{H} \in D_c^b(W)$. Then it is clear from the construction that the following diagram commutes:

$$(4.4) \quad \begin{array}{ccc} \varphi_f(\mathcal{F}) \boxtimes \varphi_g(\mathcal{G}) \boxtimes \varphi_h(\mathcal{H}) & \xrightarrow[\cong]{(\text{TS}, \text{id})} & \varphi_{f \boxplus g}(\mathcal{F} \boxtimes \mathcal{G})|_{U_0 \times V_0} \boxtimes \varphi_h(\mathcal{H}) \\ \cong \downarrow (\text{id}, \text{TS}) & & \cong \downarrow \text{TS} \\ \varphi_f(\mathcal{F}) \boxtimes \varphi_{g \boxplus h}(\mathcal{G} \boxtimes \mathcal{H})|_{V_0 \times W_0} & \xrightarrow[\cong]{\text{TS}} & \varphi_{f \boxplus g \boxplus h}(\mathcal{F} \boxtimes \mathcal{G} \boxtimes \mathcal{H})|_{U_0 \times V_0 \times W_0}. \end{array}$$

Assume that we are given a smooth morphism $h: U \rightarrow V$ and $h': U' \rightarrow V'$ and a regular function f on V and f' on V' . Take a constructible complex $\mathcal{F} \in D_c^b(V)$ and $\mathcal{F}' \in D_c^b(V')$. Then by the construction of the Thom–Sebastiani isomorphism, we have the following commutative diagram:

$$(4.5) \quad \begin{array}{ccc} h_0^* \varphi_f(\mathcal{F}) \boxtimes h_0'^* \varphi_{f'}(\mathcal{F}') & \xrightarrow[\cong]{\text{TS}} & (h_0 \times h_0')^* \varphi_{f \boxplus f'}(\mathcal{F} \boxtimes \mathcal{F}') \\ \cong \downarrow (4.2) & & \cong \downarrow (4.2) \\ \varphi_{f \circ h}(\mathcal{F}) \boxtimes \varphi_{f' \circ h'}(\mathcal{F}') & \xrightarrow[\cong]{\text{TS}} & \varphi_{f \circ h \boxplus f' \circ h'}(\mathcal{F} \boxtimes \mathcal{F}'). \end{array}$$

It can be easily shown that the Thom–Sebastiani theorem (4.3) holds more generally for functions on smooth Artin stacks by pulling back to a smooth chart and using (4.2).

For later use, we compute the vanishing cycle complex with respect to quadratic functions. Let $q = x_1 y_1 + \cdots + x_n y_n: \mathbb{A}^{2n} \rightarrow \mathbb{A}^1$ be a quadratic function. Set

$$\begin{aligned} V_x &:= \{(x_1, \dots, y_n) \in \mathbb{A}^{2n} \mid y_1 = \cdots = y_n = 0\}, \\ V_y &:= \{(x_1, \dots, y_n) \in \mathbb{A}^{2n} \mid x_1 = \cdots = x_n = 0\}. \end{aligned}$$

Then the inclusions $i_x: V_x \hookrightarrow \mathbb{A}^{2n}$ and $i_y: V_y \hookrightarrow \mathbb{A}^{2n}$ and the exchange map (4.2) implies natural morphisms of complexes

$$\tilde{c}_x: i_x^* \varphi_q \rightarrow \varphi_0(\mathbb{Q}_{V_x}[2n]) = \mathbb{Q}_{V_x}[2n], \quad \tilde{c}_y: i_y^* \varphi_q \rightarrow \varphi_0(\mathbb{Q}_{V_y}[2n]) = \mathbb{Q}_{V_y}[2n].$$

Note that φ_q has its support contained in the origin. Therefore, by taking the compactly supported cohomology, we obtain natural morphisms

$$(4.6) \quad c_x: \varphi_q|_0 \rightarrow \mathbb{Q}_0, \quad c_y: \varphi_q|_0 \rightarrow \mathbb{Q}_0.$$

It follows from [Dav17, Theorem A.1] that c_x and c_y are isomorphisms. We note that c_x and c_y differ by the sign $(-1)^n$ ³. This discrepancy corresponds to the ambiguity of the trivialization of the vanishing cycle complexes with respect to the quadratic function explained in [Bra+15, Example 2.14]. Assume further that

³This can be shown e.g. by constructing a proper homotopy between V_x and V_y inside $q^{-1}(0)$ which reverses the orientation induced from the complex structure. See the proof of [KK23, Proposition 2.24] for a related discussion.

we are given a quadratic function $q' = z_1 w_1 + \cdots + z_m w_m: \mathbb{A}^{2m} \rightarrow \mathbb{A}^1$. We let $V_w \subset \mathbb{A}^{2m}$ and $V_{y,w} \subset \mathbb{A}^{2n+2m}$ in the same manner as V_y . Then it is clear from the construction that the following diagram commutes:

$$(4.7) \quad \begin{array}{ccc} \varphi_q|_0 \boxtimes \varphi_{q'}|_0 & \xrightarrow[\cong]{\text{TS}} & \varphi_{q \boxplus q'}|_0 \\ \cong \downarrow (c_y, c_w) & & \cong \downarrow c_{y,w} \\ \mathbb{Q}_0 \boxtimes \mathbb{Q}_0 & \xrightarrow[\cong]{} & \mathbb{Q}_0. \end{array}$$

4.2. Donaldson–Thomas perverse sheaf. We now briefly explain the construction of the Donaldson–Thomas perverse sheaf associated with oriented d-critical algebraic stacks following [Bra+15, §6] and [Ben+15, §4]. Recall that for a d-critical scheme (X, s) and an étale d-critical chart $\mathfrak{R} = (R, \eta, U, f, i)$, we have constructed in (3.33) the canonical orientation

$$o_{\mathfrak{R}}^{\text{can}}: K_U^{\otimes 2}|_{R^{\text{red}}} \cong K_{R, \eta^* s}$$

for $(R, \eta^* s)$. For a given orientation (\mathcal{L}, o) for (X, s) such that the parity of $\dim U$ and the grading of \mathcal{L} agrees, we define a $\mathbb{Z}/2\mathbb{Z}$ -local system $Q_{\mathfrak{R}}^o$ on R parametrizing local isomorphisms of orientations $o_{\mathfrak{R}}^{\text{can}} \cong o|_R$. Assume that we are given an embedding

$$(4.8) \quad \mathfrak{R} = (R, \eta, U, f, i) \hookrightarrow \mathfrak{R}' = (R, \eta, U \times \mathbb{A}^{2n}, f \boxplus (x_1 y_1 + \cdots + x_n y_n), (i, 0)).$$

Then we construct an isomorphism $o_{\mathfrak{R}}^{\text{can}} \cong o_{\mathfrak{R}'}^{\text{can}}$ by

$$(4.9) \quad K_U \cong K_{U \times \mathbb{A}^{2n}}, \quad v \mapsto v \wedge (dx_1 \wedge \cdots \wedge dx_n \wedge dy_n \wedge \cdots \wedge dy_1).$$

This induces an isomorphism of local systems

$$b_y: Q_{\mathfrak{R}'}^o \cong Q_{\mathfrak{R}}^o.$$

Note that we can define b_x by swapping x -coordinates and y -coordinates. We have an identity $b_x = (-1)^n b_y$. Consider the following d-critical chart

$$\mathfrak{R}'' = (R, \eta, U \times \mathbb{A}^{2n+2m}, f \boxplus (x_1 y_1 + \cdots + x_n y_n) \boxplus (z_1 w_1 + \cdots + z_m w_m), (i, 0)).$$

Then we have an identity

$$(4.10) \quad b_{y,w} = b_y \circ b_w.$$

We now explain the construction of the DT perverse sheaf. It is shown [Bra+15, Theorem 6.9] that there exists a perverse sheaf called the **Donaldson–Thomas (DT) perverse sheaf**

$$\varphi_{X,s,o} \in \text{Perv}(X)$$

with a choice of an isomorphism

$$\omega_{\mathfrak{R}}: \varphi_{X,s,o} \cong \varphi_f|_R \otimes_{\mathbb{Z}/2\mathbb{Z}} Q_{\mathfrak{R}}^o$$

which satisfies the following properties:

- (1) If we are given an étale morphism $(h_0, h): \mathfrak{R} = (R, U, f, \eta, i) \rightarrow \mathfrak{R}' = (R', U', f', \eta', i')$, the following diagram commutes:

$$\begin{array}{ccc} \varphi_{X,s,o}|_R & \xrightarrow[\cong]{\omega_{\mathfrak{R}'|_R}} & \varphi_{f'}|_R \otimes_{\mathbb{Z}/2\mathbb{Z}} Q_{\mathfrak{R}'}^o \\ \parallel & & \downarrow \cong \\ \varphi_{X,s,o}|_R & \xrightarrow[\cong]{\omega_{\mathfrak{R}}} & \varphi_f|_R \otimes_{\mathbb{Z}/2\mathbb{Z}} Q_{\mathfrak{R}}^o \end{array}$$

where the right vertical map is defined in the canonical manner.

(2) If we are given an embedding (4.8), then the following diagram commutes⁴:

$$\begin{array}{ccc} \varphi_{X,s,o}|_R & \xrightarrow[\cong]{\omega_{\mathfrak{R}}} & \varphi_f|_R \otimes_{\mathbb{Z}/2\mathbb{Z}} Q_{\mathfrak{R}}^o \\ \cong \downarrow \omega_{\mathfrak{R}'} & & \cong \downarrow (c_y^{-1}, b_y^{-1}) \\ \varphi_{f \boxplus (x_1 y_1 + \dots + x_n y_n)}|_R \otimes Q_{\mathfrak{R}'}^o & \xrightarrow[\cong]{\text{TS}^{-1}} & \varphi_f|_R \boxtimes \varphi_{x_1 y_1 + \dots + x_n y_n} \otimes Q_{\mathfrak{R}'}^o. \end{array}$$

Note that the DT perverse sheaf $\varphi_{X,s,o}$ is uniquely characterized up to unique isomorphism by the above property. If there is no confusion, we simply write $\varphi_X = \varphi_{X,s,o}$.

Assume that we are given a smooth morphism $h: X_1 \rightarrow X_2$, s_2 be a d-critical structure on X_2 and o_2 be an orientation for (X_2, s_2) . We set $s_1 := h^* s_2$ and $o_1 := h^* o_2$. Then we can construct a natural isomorphism

$$\Theta_h = \Theta_{h,s_2,o_2}: h^* \varphi_{X_2,s_2,o_2} \cong \varphi_{X_1,s_1,o_1}[-\dim h]$$

characterized by the following property: If we are given étale d-critical charts $\mathfrak{R}_j = (R_j, \eta_j, U_j, f_j, i_j)$ for (X_j, s_j) and a morphism $(H_0, H): \mathfrak{R}_1 \rightarrow \mathfrak{R}_2$ as in Proposition 3.26, the following diagram commutes:

$$\begin{array}{ccc} \varphi_{X_1,s_1,o_1} & \xrightarrow[\cong]{\omega_{\mathfrak{R}_1}} & \varphi_{f_1}|_{R_1} \otimes_{\mathbb{Z}/2\mathbb{Z}} Q_{\mathfrak{R}_1}^{o_1}|_{R_1} \\ \Theta_h^{-1}|_{R_1} \cong \downarrow & & (\Theta_H, b_H) \cong \downarrow \\ h^* \varphi_{X_2,s_2,o_2}[\dim h] & \xrightarrow[\cong]{\omega_{\mathfrak{R}_2}} & (H_0^* \varphi_{f_2})|_{R_1} \otimes_{\mathbb{Z}/2\mathbb{Z}} H_0^* Q_{\mathfrak{R}_2}^{o_2}[\dim h] \end{array}$$

where Θ_H is the morphism constructed in (4.2) and the map b_H is given using (3.35). By using (3.36), one sees that the isomorphism Θ_h is associative with respect to the composition of smooth morphisms: Namely, if we are given a smooth morphism $g: X_0 \rightarrow X_1$ and if we set $s_0 := g^* s_1$ and $o_0 := g^* o_1$, the following diagram commutes:

$$(4.11) \quad \begin{array}{ccc} g^* h^* \varphi_{X_2,s_2,o_2} & \xrightarrow[\cong]{} & (h \circ g)^* \varphi_{X_2,s_2,o_2} \\ \cong \downarrow \Theta_h & & \cong \downarrow \Theta_{h \circ g} \\ g^* \varphi_{X_1,s_1,o_1}[-\dim h] & \xrightarrow[\cong]{\Theta_g} & \varphi_{X_0,s_0,o_0}[-\dim h \circ g] \end{array}$$

Now take an oriented d-critical stack (\mathfrak{X}, s, o) . Then the commutativity of the diagram (4.11) implies that there exists a perverse sheaf

$$\varphi_{\mathfrak{X}} = \varphi_{\mathfrak{X},s,o} \in \text{Perv}(\mathfrak{X})$$

with a choice of an isomorphism $\Theta_q: q^* \varphi_{\mathfrak{X},s,o} \cong \varphi_{X,q^* s, q^* o}[-\dim q]$ for a smooth morphism $q: X \rightarrow \mathfrak{X}$ from an algebraic space, uniquely characterized up to unique isomorphism by the following property: If we are given a smooth morphism $h: X_1 \rightarrow X_2$ between algebraic spaces and a smooth morphism $q_2: X_2 \rightarrow \mathfrak{X}$, the following diagram commutes:

$$(4.12) \quad \begin{array}{ccc} h^* q_2^* \varphi_{\mathfrak{X},s,o} & \xrightarrow[\cong]{} & q_1^* \varphi_{\mathfrak{X},s,o} \\ \cong \downarrow \Theta_{q_2} & & \cong \downarrow \Theta_{q_1} \\ h^* \varphi_{X_2,s_2,o_2}[-\dim q_2] & \xrightarrow[\cong]{\Theta_h} & \varphi_{X_1,s_1,o_1}[-\dim q_1] \end{array}$$

where we set $q_1 := q \circ h$, $s_1 := q_1^* s$, $s_2 := q_2^* s$, $o_1 := q_1^* o$ and $o_2 := q_2^* o$. It is shown in [Ben+15, Theorem 4.8] that the DT perverse sheaf is Verdier self-dual; namely, there exists a natural isomorphism

$$(4.13) \quad \mathbb{D}_{\mathfrak{X}}(\varphi_{\mathfrak{X},s,o}) \cong \varphi_{\mathfrak{X},s,o}$$

⁴Note that our choice for the morphism c_y and b_y are different from the one in [Ben+15] up to some sign. We do not compute the explicit difference here. However, since our choice of b_y and c_y are associative with respect to the composition of embeddings, the associativity of the Thom–Sebastiani isomorphism (4.4) implies that the difference of these signs are associative. In particular, one can glue the vanishing cycle complexes using our choice of sign.

where $\mathbb{D}_{\mathfrak{X}}$ denotes the Verdier dualizing functor. For an oriented (-1) -shifted symplectic stack $(\mathfrak{X}, \omega_{\mathfrak{X}}, o)$ whose underlying d-critical structure is denoted by s , we set

$$\varphi_{\mathfrak{X}, \omega_{\mathfrak{X}}, o} := \varphi_{\mathfrak{X}^{cl}, s, o^{cl}}.$$

4.3. Thom–Sebastiani theorem for DT perverse sheaves. The aim of this section is to prove the Thom–Sebastiani theorem for Donaldson–Thomas perverse sheaves on d-critical stacks and (-1) -shifted symplectic stacks. We first prove it for algebraic spaces:

Proposition 4.1. *Let (X_1, s_1, o_1) and (X_2, s_2, o_2) be oriented d-critical algebraic spaces. Then there exists a natural isomorphism*

$$\text{TS}: \varphi_{X_1, s_1, o_1} \boxtimes \varphi_{X_2, s_2, o_2} \cong \varphi_{X_1 \times X_2, s_1 \boxplus s_2, o_1 \boxtimes o_2}.$$

Further, if we are given another oriented d-critical algebraic space (X_3, s_3, o_3) , the following diagram commutes:

$$(4.14) \quad \begin{array}{ccc} \varphi_{X_1, s_1, o_1} \boxtimes \varphi_{X_2, s_2, o_2} \boxtimes \varphi_{X_3, s_3, o_3} & \xrightarrow[\cong]{(\text{TS}, \text{id})} & \varphi_{X_1 \times X_2, s_1 \boxplus s_2, o_1 \boxtimes o_2} \boxtimes \varphi_{X_3, s_3, o_3} \\ \cong \downarrow (\text{id}, \text{TS}) & & \cong \downarrow \text{TS} \\ \varphi_{X_1, s_1, o_1} \boxtimes \varphi_{X_2 \times X_3, s_2 \boxplus s_3, o_2 \boxtimes o_3} & \xrightarrow[\cong]{\text{TS}} & \varphi_{X_1 \times X_2 \times X_3, s_1 \boxplus s_2 \boxplus s_3, o_1 \boxtimes o_2 \boxtimes o_3}. \end{array}$$

Proof. Take étale d-critical charts $\mathfrak{R}_1 = (R_1, \eta_1, U_1, f_1, i_1)$ for X_1 and $\mathfrak{R}_2 = (R_2, \eta_2, U_2, f_2, i_2)$. As we have seen in (3.37), there exists a natural isomorphism

$$(4.15) \quad o_{\mathfrak{R}_1}^{\text{can}} \boxtimes o_{\mathfrak{R}_2}^{\text{can}} \cong o_{\mathfrak{R}_1 \times \mathfrak{R}_2}^{\text{can}}.$$

In particular, we have a natural isomorphism

$$(4.16) \quad \alpha_{\mathfrak{R}_1, \mathfrak{R}_2}: Q_{\mathfrak{R}_1}^{o_1} \boxtimes Q_{\mathfrak{R}_2}^{o_2} \cong Q_{\mathfrak{R}_1 \times \mathfrak{R}_2}^{o_1 \boxtimes o_2}.$$

Define an isomorphism $\text{TS}_{\mathfrak{R}_1, \mathfrak{R}_2}: \varphi_{X_1, s_1, o_1}|_{R_1} \boxtimes \varphi_{X_2, s_2, o_2}|_{R_2} \cong \varphi_{X_1 \times X_2, s_1 \boxplus s_2, o_1 \boxtimes o_2}|_{R_1 \times R_2}$ by the following composition:

$$(4.17) \quad \begin{aligned} \varphi_{X_1, s_1, o_1}|_{R_1} \boxtimes \varphi_{X_2, s_2, o_2}|_{R_2} &\xrightarrow[\cong]{\omega_{\mathfrak{R}_1 \boxtimes \mathfrak{R}_2}} (\varphi_{f_1}|_{R_1} \otimes_{\mathbb{Z}/2\mathbb{Z}} Q_{\mathfrak{R}_1}^{o_1}) \boxtimes (\varphi_{f_2}|_{R_2} \otimes_{\mathbb{Z}/2\mathbb{Z}} Q_{\mathfrak{R}_2}^{o_2}) \\ &\xrightarrow[\cong]{(\text{TS}, \alpha_{\mathfrak{R}_1, \mathfrak{R}_2})} \varphi_{f_1 \boxplus f_2}|_{R_1 \times R_2} \otimes_{\mathbb{Z}/2\mathbb{Z}} Q_{\mathfrak{R}_1 \times \mathfrak{R}_2}^{o_1 \boxtimes o_2} \\ &\xrightarrow[\cong]{\omega_{\mathfrak{R}_1 \times \mathfrak{R}_2}^{-1}} \varphi_{X_1 \times X_2, s_1 \boxplus s_2, o_1 \boxtimes o_2}|_{R_1 \times R_2}. \end{aligned}$$

We will show that these isomorphisms glue to define the desired Thom–Sebastiani isomorphism. Firstly, it is clear that the map $\text{TS}_{\mathfrak{R}_1, \mathfrak{R}_2}$ is compatible with étale cover of étale d-critical charts. Therefore, by using Theorem 3.25, it is enough to prove an identity

$$(4.18) \quad \text{TS}_{\mathfrak{R}_1, \mathfrak{R}_2} = \text{TS}_{\mathfrak{R}'_1, \mathfrak{R}'_2}$$

where we set $\mathfrak{R}'_1 = (R_1, \eta_1, U \times \mathbb{A}^{2n}, f_1 \boxplus (x_1 y_1 + \cdots + x_n y_n), (i_1, 0))$ and $\mathfrak{R}'_2 = (R_2, \eta_2, U \times \mathbb{A}^{2m}, f_2 \boxplus (z_1 w_1 + \cdots + z_m w_m), (i_2, 0))$. We set $q_1 = x_1 y_1 + \cdots + x_n y_n$ and $q_2 = z_1 w_1 + \cdots + z_m w_m$. By construction of the Thom–Sebastiani isomorphism of the vanishing cycle complexes and the commutativity of the diagram (4.7), we see that the following diagram commutes:

$$\begin{array}{ccc} \varphi_{f_1} \boxtimes \varphi_{q_1} \boxtimes \varphi_{f_2} \boxtimes \varphi_{q_2} & \xrightarrow[\cong]{\text{TS}} & \varphi_{f_1 \boxplus q_1 \boxplus f_2 \boxplus q_2} \\ \cong \downarrow (\text{id}, c_y, \text{id}, c_w) & & \cong \downarrow \text{TS} \\ \varphi_{f_1} \boxtimes \varphi_{f_2} & \xrightarrow[\cong]{(\text{TS}, c_{y, w}^{-1})} & \varphi_{f_1 \boxplus f_2} \boxtimes \varphi_{q_1 \boxplus q_2}. \end{array}$$

On the other hand, it follows from the construction of the isomorphism b_y and the equality (4.10) that the following diagram commutes:

$$\begin{array}{ccc} Q_{\mathfrak{R}'_1}^{o_1} \boxtimes Q_{\mathfrak{R}'_2}^{o_2} & \xrightarrow[\cong]{\alpha_{\mathfrak{R}'_1, \mathfrak{R}'_2}} & Q_{\mathfrak{R}'_1 \times \mathfrak{R}'_2}^{o_1 \boxtimes o_2} \\ \cong \downarrow (b_y, b_w) & & \cong \downarrow b_{y, w} \\ Q_{\mathfrak{R}_1}^{o_1} \boxtimes Q_{\mathfrak{R}_2}^{o_2} & \xrightarrow[\cong]{\alpha_{\mathfrak{R}_1, \mathfrak{R}_2}} & Q_{\mathfrak{R}_1 \times \mathfrak{R}_2}^{o_1 \boxtimes o_2} \end{array}$$

This commutativity implies the identity (4.18) as desired.

To prove the commutativity of the diagram (4.14), by using (4.4), it is enough to prove the commutativity of the following diagram for étale d -critical charts $\mathfrak{R}_j = (R_j, \eta_j, U_j, f_j, i_j)$ for X_j ,

$$\begin{array}{ccc} Q_{\mathfrak{R}_1}^{o_1} \boxtimes Q_{\mathfrak{R}_2}^{o_2} \boxtimes Q_{\mathfrak{R}_3}^{o_3} & \xrightarrow[\cong]{(\text{id}, \alpha_{\mathfrak{R}_2, \mathfrak{R}_3})} & Q_{\mathfrak{R}_1}^{o_1} \boxtimes Q_{\mathfrak{R}_2 \boxtimes \mathfrak{R}_3}^{o_2 \boxtimes o_3} \\ \cong \downarrow (\alpha_{\mathfrak{R}_1, \mathfrak{R}_2}, \text{id}) & & \cong \downarrow \alpha_{\mathfrak{R}_1, \mathfrak{R}_2 \times \mathfrak{R}_3} \\ Q_{\mathfrak{R}_1 \times \mathfrak{R}_2}^{o_1 \boxtimes o_2} \boxtimes Q_{\mathfrak{R}_3}^{o_3} & \xrightarrow[\cong]{\alpha_{\mathfrak{R}_1 \times \mathfrak{R}_2, \mathfrak{R}_3}} & Q_{\mathfrak{R}_1 \times \mathfrak{R}_2 \times \mathfrak{R}_3}^{o_1 \boxtimes o_2 \boxtimes o_3} \end{array}$$

which is obvious from the commutativity (3.38). \square

Next, we will prove Thom–Sebastiani theorem for d -critical Artin stacks. To do this, we first show that the Thom–Sebastiani isomorphism for algebraic spaces is compatible with smooth pullback.

Lemma 4.2. *Let $h_1: X_1 \rightarrow Y_1$ and $h_2: X_2 \rightarrow Y_2$ be smooth morphisms of algebraic spaces, t_1 and t_2 be the d -critical structures of Y_1 and Y_2 , respectively. Set $s_1 := h_1^* t_1$ and $s_2 := h_2^* t_2$. Then the following diagram commutes:*

$$\begin{array}{ccc} h_1^* \varphi_{Y_1, t_1, o_1} \boxtimes h_2^* \varphi_{Y_2, t_2, o_2} & \xrightarrow[\cong]{\text{TS}} & (h_1 \times h_2)^* (\varphi_{Y_1 \times Y_2, t_1 \boxplus t_2, o_1 \boxtimes o_2}) \\ \cong \downarrow (\Theta_{h_1}, \Theta_{h_2}) & & \cong \downarrow \Theta_{h_1 \times h_2} \\ \varphi_{X_1, s_1, h_1^* o_1}[-d_1] \boxtimes \varphi_{X_2, s_2, h_2^* o_2}[-d_2] & \xrightarrow[\cong]{\text{TS}} & \varphi_{X_1 \times X_2, s_1 \boxplus s_2, o_1 \boxtimes o_2}[-d_1 - d_2]. \end{array}$$

Here we set $d_1 = \dim h_1$ and $d_2 = \dim h_2$.

Proof. Take a point $x_1 \in X_1$ and $x_2 \in X_2$. By using Proposition 3.26, we take an étale d -critical chart $\mathfrak{R}_k = (R_k, \eta_k, U_k, f_k, i_k)$ of x_k and étale d -critical chart $\mathfrak{S}_k = (S_k, \gamma_k, V_k, g_k, j_k)$ and a smooth morphism $(H_{k,0}, H_k): \mathfrak{R}_k \rightarrow \mathfrak{S}_k$ for $k = 1, 2$. Then by the commutativity of the diagram (4.5), it is enough to prove the commutativity of the following diagram:

$$\begin{array}{ccc} H_{1,0}^* Q_{\mathfrak{S}_1}^{o_1} \boxtimes H_{2,0}^* Q_{\mathfrak{S}_2}^{o_2} & \xrightarrow[\cong]{\alpha_{\mathfrak{S}_1, \mathfrak{S}_2}} & (H_{1,0} \times H_{2,0})^* (Q_{\mathfrak{S}_1 \times \mathfrak{S}_2}^{o_1 \boxtimes o_2}) \\ b_{H_1} \boxtimes b_{H_2} \downarrow \cong & & b_{H_1 \times H_2} \downarrow \cong \\ Q_{\mathfrak{R}_1}^{h_1^* o_1} \boxtimes Q_{\mathfrak{R}_2}^{h_2^* o_2} & \xrightarrow[\cong]{\alpha_{\mathfrak{R}_1, \mathfrak{R}_2}} & Q_{\mathfrak{R}_1 \times \mathfrak{R}_2}^{h_1^* o_1 \boxtimes h_2^* o_2} \end{array}$$

This is obvious from the commutativity of (3.39). \square

By combining Lemma 4.2, the commutativity of the diagram (4.12) and (4.14), we obtain the following statement by smooth descent:

Proposition 4.3. *Let $(\mathfrak{X}_1, s_1, o_1)$ and $(\mathfrak{X}_2, s_2, o_2)$ be oriented d -critical stacks. Then there exists a natural isomorphism*

$$\text{TS}: \varphi_{\mathfrak{X}_1, s_1, o_1} \boxtimes \varphi_{\mathfrak{X}_2, s_2, o_2} \cong \varphi_{\mathfrak{X}_1 \times \mathfrak{X}_2, s_1 \boxplus s_2, o_1 \boxtimes o_2}.$$

Further, if we are given another oriented d -critical algebraic space $(\mathfrak{X}_3, s_3, o_3)$, the following diagram commutes:

$$(4.19) \quad \begin{array}{ccc} \varphi_{\mathfrak{X}_1, s_1, o_1} \boxtimes \varphi_{\mathfrak{X}_2, s_2, o_2} \boxtimes \varphi_{\mathfrak{X}_3, s_3, o_3} & \xrightarrow[\cong]{(\text{TS}, \text{id})} & \varphi_{\mathfrak{X}_1 \times \mathfrak{X}_2, s_1 \boxplus s_2, o_1 \boxtimes o_2} \boxtimes \varphi_{\mathfrak{X}_3, s_3, o_3} \\ \cong \downarrow (\text{id}, \text{TS}) & & \cong \downarrow \text{TS} \\ \varphi_{\mathfrak{X}_1, s_1, o_1} \boxtimes \varphi_{\mathfrak{X}_2 \times \mathfrak{X}_3, s_2 \boxplus s_3, o_2 \boxtimes o_3} & \xrightarrow[\cong]{\text{TS}} & \varphi_{\mathfrak{X}_1 \times \mathfrak{X}_2 \times \mathfrak{X}_3, s_1 \boxplus s_2 \boxplus s_3, o_1 \boxtimes o_2 \boxtimes o_3} \end{array}$$

Let $(\mathfrak{X}_1, \omega_{\mathfrak{X}_1}, o_1)$ and $(\mathfrak{X}_2, \omega_{\mathfrak{X}_2}, o_2)$ be oriented (-1) -shifted symplectic stacks with underlying d -critical structures s_1 and s_2 , respectively. Then, using the isomorphism of orientations (3.31), we obtain the following corollary:

Corollary 4.4. *There exists a natural isomorphism*

$$\text{TS}: \varphi_{\mathfrak{X}_1, \omega_{\mathfrak{X}_1}, o_1} \boxtimes \varphi_{\mathfrak{X}_2, \omega_{\mathfrak{X}_2}, o_2} \cong \varphi_{\mathfrak{X}_1 \times \mathfrak{X}_2, \omega_{\mathfrak{X}_1} \boxplus \omega_{\mathfrak{X}_2}, o_1 \boxtimes o_2}$$

which satisfies the associativity property as in (4.19).

5. STACK OF GRADED/FILTERED OBJECTS

In this section, we recall the notion of the stack of graded objects and filtered objects following [Hal14].

5.1. Graded and filtered objects: definition. We first recall the definition of the graded and filtered objects.

Definition 5.1. Let \mathfrak{X} be a derived higher Artin stack. For a positive integer n , we set $\text{Grad}^n(\mathfrak{X}) := \text{Map}(B\mathbb{G}_m^n, \mathfrak{X})$ (resp. $\text{Filt}^n(\mathfrak{X}) := \text{Map}([\mathbb{A}^1/\mathbb{G}_m]^n, \mathfrak{X})$) and call it the stack of n -**graded objects** (resp. n -**filtered objects**). We denote $\text{Grad}(\mathfrak{X}) = \text{Grad}^1(\mathfrak{X})$ and $\text{Filt}(\mathfrak{X}) = \text{Filt}^1(\mathfrak{X})$.

Example 5.2. Let X be a quasi-separated algebraic space with \mathbb{G}_m^n -action μ . We define the fixed point space X^μ (resp. attracting space $X^{\mu, +, \dots, +}$) by

$$\begin{aligned} X^\mu &:= \text{Map}^{\mathbb{G}_m^n}(\text{pt}, (X, \mu)) \\ X^{\mu, +, \dots, +} &:= \text{Map}^{\mathbb{G}_m^n}(\mathbb{A}^n, (X, \hat{\mu})). \end{aligned}$$

Then by [Hal14, Proposition 1.4.1], these functors are represented by quasi-separated algebraic spaces. Then the quotient stack $[X^\mu/\mathbb{G}_m^n]$ is an open and closed substack of $\text{Grad}^n([X/\mathbb{G}_m^n])$ consisting of graded points x such that the composition $B\mathbb{G}_m^n \xrightarrow{x} [X^\mu/\mathbb{G}_m^n] \rightarrow B\mathbb{G}_m^n$ is the identity map. Similarly, $[X^{\mu, +, \dots, +}/\mathbb{G}_m^n]$ is an open and closed substack of $\text{Filt}^n([X/\mathbb{G}_m^n])$.

We have the following algebraicity statement of $\text{Grad}(\mathfrak{X})$ and $\text{Filt}(\mathfrak{X})$:

Proposition 5.3 ([Hal14, Proposition 1.1.2]). *Let \mathfrak{X} be a quasi-separated (classical) Artin stack with affine stabilizers. Then $\text{Grad}^n(\mathfrak{X})$ and $\text{Filt}^n(\mathfrak{X})$ are quasi-separated Artin stacks.*

We will be interested in the following maps:

- The inclusion $i_1: \text{pt} \rightarrow [\mathbb{A}^1/\mathbb{G}_m]^n$ of 1 which induces a map $\text{ev}: \text{Filt}^n(\mathfrak{X}) \rightarrow \mathfrak{X}$.
- The inclusion $i_0: B\mathbb{G}_m^n \rightarrow [\mathbb{A}^1/\mathbb{G}_m]^n$ of the origin, which induces a map $\text{gr}: \text{Filt}^n(\mathfrak{X}) \rightarrow \text{Grad}^n(\mathfrak{X})$.
- The projection $[\mathbb{A}^1/\mathbb{G}_m]^n \rightarrow B\mathbb{G}_m^n$ which induces a map $\sigma: \text{Grad}^n(\mathfrak{X}) \rightarrow \text{Filt}^n(\mathfrak{X})$ which is a section of gr .
- The inclusion of the basepoint $\text{pt} \rightarrow B\mathbb{G}_m$ which induces a map $u: \text{Grad}(\mathfrak{X}) \rightarrow \mathfrak{X}$ which coincides with $\text{ev} \circ \sigma$.

For a derived higher Artin stack \mathfrak{X} we consider the **attractor correspondence**

$$(5.1) \quad \begin{array}{ccc} & \text{Filt}(\mathfrak{X}) & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \text{Grad}(\mathfrak{X}) & & \mathfrak{X} \end{array}$$

The commutativity of the attractor correspondence is expressed by the diagram

$$(5.2) \quad \begin{array}{ccccc} & & \text{Grad}(\text{Filt}(\mathfrak{X})) & & \text{Filt}(\mathfrak{X}) \\ & & \swarrow \text{Grad}(\text{ev}_{\mathfrak{X}}) & \searrow \text{gr}_{\mathfrak{X}} & \swarrow \text{ev}_{\mathfrak{X}} \\ & & \text{Grad}(\mathfrak{X}) & & \text{Filt}(\mathfrak{X}) \\ & \swarrow \text{Grad}(\text{gr}_{\mathfrak{X}}) & & \swarrow \text{Filt}(\text{ev}_{\mathfrak{X}}) & \\ \text{Grad}^2(\mathfrak{X}) & & \text{Filt}^2(\mathfrak{X}) & & \mathfrak{X} \\ & \swarrow \text{gr}_{\text{Filt}(\mathfrak{X})} & & \swarrow \text{ev}_{\mathfrak{X}} & \\ & & \text{Grad}(\mathfrak{X}) & & \text{Filt}(\mathfrak{X}) \\ & \swarrow \text{Filt}(\text{ev}_{\mathfrak{X}}) & \swarrow \text{ev}_{\text{Filt}(\mathfrak{X})} & \swarrow \text{gr}_{\mathfrak{X}} & \\ & \text{Filt}(\text{Grad}(\mathfrak{X})) & & \text{Filt}(\mathfrak{X}) & \\ & \swarrow \text{gr}_{\text{Grad}(\mathfrak{X})} & \swarrow \text{ev}_{\text{Grad}(\mathfrak{X})} & \swarrow \text{gr}_{\mathfrak{X}} & \\ & & \text{Grad}(\mathfrak{X}) & & \text{Filt}(\mathfrak{X}) \end{array}$$

Remark 5.4. If we consider the attractor correspondence as a morphism $\text{attr}_{\mathfrak{X}}: \mathfrak{X} \rightarrow \text{Grad}(\mathfrak{X})$ in the $(\infty, 2)$ -category of correspondences of derived stacks, the iterated correspondence (5.2) provides a two-morphism witnessing the commutativity of the diagram

$$\begin{array}{ccc} \mathfrak{X} & \xrightarrow{\text{attr}_{\mathfrak{X}}} & \text{Grad}(\mathfrak{X}) \\ \downarrow \text{attr}_{\mathfrak{X}} & & \downarrow \text{attr}_{\text{Grad}(\mathfrak{X})} \\ \text{Grad}(\mathfrak{X}) & \xrightarrow{\text{Grad}(\text{attr}_{\mathfrak{X}})} & \text{Grad}^2(\mathfrak{X}) \end{array}$$

Now let G be an affine algebraic group and

$$X_{\bullet}(G) = \text{Hom}(\mathbb{G}_m, G), \quad X_{\bullet}^2(G) = \text{Hom}(\mathbb{G}_m^2, G)$$

the sets of cocharacters of G . For a cocharacter $\lambda: \mathbb{G}_m \rightarrow G$ we denote by

$$G_{\lambda} = \{g \in G \mid \lambda(t)g\lambda(t)^{-1} = g\}$$

the fixed point subgroup and $G_{\lambda}^+ \subset G$ the subgroup of elements $g \in G$ such that the morphism $\mathbb{G}_m \rightarrow G$ given by $t \mapsto \lambda(t)g\lambda(t)^{-1}$ admits an extension to a morphism $\mathbb{A}^1 \rightarrow G$. Informally,

$$G_{\lambda}^+ = \{g \in G \mid \exists \lim_{t \rightarrow 0} \lambda(t)g\lambda(t)^{-1}\}.$$

Remark 5.5. Since G is affine and $R[t] \rightarrow R[t, t^{-1}]$ is injective for any commutative ring R , $\text{Map}(\mathbb{A}^1, G) \rightarrow \text{Map}(\mathbb{G}_m, G)$ is a monomorphism, i.e. the \mathbb{A}^1 -extension is unique if it exists.

We have a projection $G_{\lambda}^+ \rightarrow G_{\lambda}$ given by $g \mapsto \lim_{t \rightarrow 0} \lambda(t)g\lambda(t)^{-1}$ and its section $G_{\lambda} \rightarrow G_{\lambda}^+$ given by the identity map on G .

Similarly, for a pair of commuting cocharacters $\hat{\lambda} = (\lambda_1, \lambda_2): \mathbb{G}_m^2 \rightarrow G$ we denote by

$$G_{\hat{\lambda}} = \{g \in G \mid \hat{\lambda}(t_1, t_2)g\hat{\lambda}(t_1, t_2)^{-1} = g\}$$

and

$$G_{\hat{\lambda}}^{++} = \{g \in G \mid \exists \lim_{t_i \rightarrow 0} \hat{\lambda}(t_1, t_2)g\hat{\lambda}(t_1, t_2)^{-1}\}$$

the corresponding fixed point and attractor subgroups.

If X is an algebraic space with a G -action, for any $\lambda: \mathbb{G}_m \rightarrow G$ and $g \in G$ we have an isomorphism of stacks

$$[X^{\lambda}/G_{\lambda}] \longrightarrow [X^{g\lambda g^{-1}}/G_{g\lambda g^{-1}}],$$

where $X^{\lambda} \rightarrow X^{g\lambda g^{-1}}$ is given by the action of $g \in G$ and $G_{\lambda} \rightarrow G_{g\lambda g^{-1}}$ is given by the conjugation by $g \in G$. Thus, it makes sense to consider the stack $[X^{\lambda}/G_{\lambda}]$ for a conjugacy class $\lambda \in X_{\bullet}(G)/G$ of cocharacters.

Theorem 5.6. *Let G be an affine algebraic group acting on an algebraic space X and set $\mathfrak{X} := [X/G]$. Then the attractor correspondence (5.1) is equivalent to*

$$\begin{array}{ccc} & \coprod_{\lambda \in X_{\bullet}(G)/G} [X^{\lambda,+}/G_{\lambda}^{+}] & \\ & \swarrow \quad \searrow & \\ \coprod_{\lambda \in X_{\bullet}(G)/G} [X^{\lambda}/G_{\lambda}] & & [X/G] \end{array}$$

and the diagram (5.2) is equivalent to

$$\begin{array}{ccccc} & \coprod_{\hat{\lambda} \in X_{\bullet}^2(G)/G} [(X^{\lambda_2,+})^{\lambda_1}/(G_{\lambda_2}^{+})_{\lambda_1}] & & \coprod_{\lambda_1 \in X_{\bullet}(G)/G} [X^{\lambda_1,+}/G_{\lambda_1}^{+}] & \\ & \swarrow \quad \searrow & \xrightarrow{\quad} & \swarrow \quad \searrow & \\ \coprod_{\hat{\lambda} \in X_{\bullet}^2(G)/G} [X^{\hat{\lambda}}/G_{\hat{\lambda}}] & & \coprod_{\lambda_1 \in X_{\bullet}(G)/G} [X^{\lambda_1}/G_{\lambda_1}] & & [X/G] \\ & \swarrow \quad \searrow & \swarrow \quad \searrow & \swarrow \quad \searrow & \\ \coprod_{\hat{\lambda} \in X_{\bullet}^2(G)/G} [X^{\hat{\lambda},+,+}/G_{\hat{\lambda}}^{+,+}] & & \coprod_{\lambda_2 \in X_{\bullet}(G)/G} [X^{\lambda_2}/G_{\lambda_2}] & & \\ & \swarrow \quad \searrow & \swarrow \quad \searrow & \swarrow \quad \searrow & \\ \coprod_{\hat{\lambda} \in X_{\bullet}^2(G)/G} [(X^{\lambda_1,+})^{\lambda_2}/(G_{\lambda_1}^{+})_{\lambda_2}] & & \coprod_{\lambda_2 \in X_{\bullet}(G)/G} [X^{\lambda_2,+}/G_{\lambda_2}^{+}] & & \end{array}$$

where we denote by $\lambda_1, \lambda_2 \in X_{\bullet}(G)/G$ the images of $\hat{\lambda}$ under the two projections $X_{\bullet}^2(G)/G \rightarrow X_{\bullet}(G)/G$.

Proof. This follows from the description of $\text{Grad}(\mathfrak{X})$ and $\text{Filt}(\mathfrak{X})$ given in [Hal14, Theorem 1.4.8]. The claim there is stated only for schemes (not algebraic spaces), but by reducing to the case of $G = \text{GL}_n$ we can use [Hal14, Theorem 1.4.7]. \square

5.2. Perfect complexes. We will use a description of perfect complexes on $B\mathbb{G}_m$ and $[\mathbb{A}^1/\mathbb{G}_m]$. Let \mathbb{Z} be the set of integers considered as a discrete category and \mathbb{Z}_{\geq} the category where \mathbb{Z} is considered as a poset with respect to \geq . We will consider finite connected subsets $P \subset \mathbb{Z}_{\geq}$. Consider the diagram

$$\begin{array}{ccc} & \text{Fun}(P, \text{Perf}_R) & \\ & \swarrow \text{gr} \quad \searrow \text{ev} & \\ \text{Fun}(P^{\delta}, \text{Perf}_R) & & \text{Perf}_R \end{array}$$

where $\text{ev}: \text{Fun}(P, \text{Perf}_R) \rightarrow \text{Perf}_R$ is given by taking the colimit or, equivalently, by the evaluation at the minimum of P . Given an inclusion $P_1 \subset P_2$ of posets there are morphisms of correspondences

$$\left(\begin{array}{ccc} & \text{Fun}(P_1, \text{Perf}_R) & \\ & \swarrow \text{gr} \quad \searrow \text{ev} & \\ \text{Fun}(P_1^{\delta}, \text{Perf}_R) & & \text{Perf}_R \end{array} \right) \longrightarrow \left(\begin{array}{ccc} & \text{Fun}(P_2, \text{Perf}_R) & \\ & \swarrow \text{gr} \quad \searrow \text{ev} & \\ \text{Fun}(P_2^{\delta}, \text{Perf}_R) & & \text{Perf}_R \end{array} \right)$$

given by left Kan extensions.

The following statement is well-known; see e.g. [GR17b, Chapter 9, Proposition 1.3.3], [Lur15, Sections 3.1 and 3.2] and [Rak20, Section 3].

Theorem 5.7. *The correspondence*

$$\begin{array}{ccc} & \text{QCoh}([\mathbb{A}^1/\mathbb{G}_m] \times \text{Spec } R) & \\ i_0^* \swarrow & & \searrow i_1^* \\ \text{QCoh}(B\mathbb{G}_m \times \text{Spec } R) & & \text{Mod}_R \end{array}$$

is equivalent to

$$\begin{array}{ccc} & \text{Fun}(\mathbb{Z}_{\geq}, \text{Mod}_R) & \\ \text{gr} \swarrow & & \searrow \text{colim} \\ \text{Fun}(\mathbb{Z}, \text{Mod}_R) & & \text{Mod}_R \end{array}$$

such that the natural symmetric monoidal structure on the left-hand side is sent to the Day convolution symmetric monoidal structure with respect to the abelian group structure on \mathbb{Z} .

Proposition 5.8. *Let R be a commutative dg algebra. Then there is an equivalence*

$$\begin{array}{ccc} \text{Perf}([\mathbb{A}^1/\mathbb{G}_m] \times \text{Spec } R) & & \left(\begin{array}{ccc} & \text{Fun}(P, \text{Perf}_R) & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \text{Fun}(P^\delta, \text{Perf}_R) & & \text{Perf}_R \end{array} \right) \\ i_0^* \swarrow & \cong & \text{colim}_{P \subset \mathbb{Z}_{\geq}, \text{ finite, connected}} \\ \text{Perf}(B\mathbb{G}_m \times \text{Spec } R) & & \text{Perf}_R \\ i_1^* \searrow & & \end{array}$$

Under this equivalence, the natural symmetric monoidal structure on the left-hand side corresponds to the Day convolution symmetric monoidal structure.

Proof. The stacks $[\mathbb{A}^1/\mathbb{G}_m] \times \text{Spec } R$ and $B\mathbb{G}_m \times \text{Spec } R$ are perfect in the sense of [BFN10], so perfect complexes on them coincide with compact objects. The computation of compact objects is analogous for both cases, so we will stick with the case of $[\mathbb{A}^1/\mathbb{G}_m]$.

Writing \mathbb{Z}_{\geq} as a colimit over its finite connected subsets $P \subset \mathbb{Z}_{\geq}$, by Theorem 5.7 we have

$$\text{QCoh}([\mathbb{A}^1/\mathbb{G}_m] \times \text{Spec } R) \simeq \text{colim}_{P \subset \mathbb{Z}_{\geq}, \text{ finite, connected}} \text{Fun}(P, \text{Mod}_R)$$

with respect to left Kan extensions, where the colimit is taken in the ∞ -category of presentable \mathbb{C} -linear ∞ -categories. For an inclusion $P_1 \subset P_2 \subset \mathbb{Z}_{\geq}$, the left Kan extension along $P_1 \subset P_2$ preserves compact objects as it has a colimit-preserving right adjoint given by restriction. Passing to compact objects, we get

$$\text{Perf}([\mathbb{A}^1/\mathbb{G}_m] \times \text{Spec } R) \simeq \text{colim}_{P \subset \mathbb{Z}_{\geq}, \text{ finite, connected}} \text{Fun}(P, \text{Mod}_R)^\omega,$$

where $\text{Fun}(P, \text{Mod}_R)^\omega \subset \text{Fun}(P, \text{Mod}_R)$ is the full subcategory of compact objects. By [Aok23, Proposition 2.8] we may further identify

$$\text{Fun}(P, \text{Mod}_R)^\omega \simeq \text{Fun}(P, \text{Perf}_R).$$

□

Remark 5.9. Under the equivalence given by Proposition 5.8 an object $V \in \text{Perf}([\mathbb{A}^1/\mathbb{G}_m] \times \text{Spec } R)$ corresponds to a \mathbb{Z} -indexed diagram

$$\cdots \rightarrow V_1 \rightarrow V_0 \rightarrow V_{-1} \rightarrow \cdots$$

of perfect complexes of R -modules such that $V_m = 0$ for $m \gg 0$ and $V_m \rightarrow V_{m-1}$ is an isomorphism for $m \ll 0$. The Day convolution of two such objects is given by

$$(V \otimes W)_m = \text{colim}_{i+j \geq m} V_i \otimes W_j.$$

If \mathcal{F} is a quasi-coherent complex on \mathfrak{X} , its pullback along $\text{Grad}(\mathfrak{X}) \rightarrow \mathfrak{X}$ acquires a natural grading as follows. This map factors as $\text{Grad}(\mathfrak{X}) \rightarrow \text{Grad}(\mathfrak{X}) \times B\mathbb{G}_m \xrightarrow{\text{ev}} \mathfrak{X}$, where ev is the natural evaluation morphism. By Theorem 5.7 the quasi-coherent complex $\text{ev}^* \mathcal{F}$ corresponds to a graded quasi-coherent complex on $\text{Grad}(\mathfrak{X})$ and its pullback under the inclusion of the basepoint $\text{pt} \rightarrow B\mathbb{G}_m$ is given by the total complex. For a quasi-coherent complex \mathcal{F} on \mathfrak{X} , we denote by $\mathcal{F}|_{\text{Grad}(\mathfrak{X})}^0$ its weight 0 part and by $\mathcal{F}|_{\text{Grad}(\mathfrak{X})}^+$ (resp. $\mathcal{F}|_{\text{Grad}(\mathfrak{X})}^-$) its positive (resp. negative) weight part. The following is [Hal14, Lemma 1.2.2].

Proposition 5.10. *Let \mathfrak{X} be a derived stack which admits an almost perfect cotangent complex. Then $\mathrm{Grad}(\mathfrak{X})$ and $\mathrm{Filt}(\mathfrak{X})$ admit almost perfect cotangent complexes, so that there are natural isomorphisms*

$$\mathbb{L}_{\mathrm{Grad}(\mathfrak{X})} \cong \mathbb{T}_{\mathfrak{X}}^0|_{\mathrm{Grad}(\mathfrak{X})}, \quad \mathbb{L}_{\mathrm{Filt}(\mathfrak{X})/\mathrm{Grad}(\mathfrak{X})}|_{\mathrm{Grad}(\mathfrak{X})} \cong \mathbb{L}_{\mathfrak{X}}|_{\mathrm{Grad}(\mathfrak{X})}^{\vee}.$$

For later use, we record the following technical statement here:

Lemma 5.11. *Let \mathfrak{X} be a quasi-separated Artin stack with affine stabilizers. Let $\iota: \mathfrak{X} \rightarrow \mathrm{Grad}(\mathfrak{X})$ be a section of $u: \mathrm{Grad}(\mathfrak{X}) \rightarrow \mathfrak{X}$. Then ι is an open immersion and the map $\mathfrak{X} \times_{\mathrm{Grad}(\mathfrak{X})} \mathrm{Filt}(\mathfrak{X}) \rightarrow \mathfrak{X}$ defined by the base change of gr along ι is an isomorphism.*

Proof. Let $\mathfrak{X}^{\mathrm{der}}$ be \mathfrak{X} regarded as a derived stack. The composition

$$(5.3) \quad \mathbb{L}_{\mathfrak{X}^{\mathrm{der}}} \cong \iota^* u^* \mathbb{L}_{\mathfrak{X}^{\mathrm{der}}} \rightarrow \iota^* \mathbb{L}_{\mathrm{Grad}(\mathfrak{X}^{\mathrm{der}})} \rightarrow \mathbb{L}_{\mathfrak{X}^{\mathrm{der}}}$$

is the identity map. The quasi-coherent complex $u^* \mathbb{L}_{\mathfrak{X}^{\mathrm{der}}}$ carries a natural grading, which induces a grading on $\mathbb{L}_{\mathfrak{X}^{\mathrm{der}}} \cong \iota^* u^* \mathbb{L}_{\mathfrak{X}^{\mathrm{der}}}$. By Proposition 5.10 the first morphism is given by the projection to its weight 0 part, so all maps appearing in (5.3) are isomorphisms. In particular, ι is étale. By [Hal14, Lemma 1.1.5], the map u is representable. Therefore, its section ι is a monomorphism, hence an open immersion by [Sta24, Tag 025G].

By a similar argument, we see that $\sigma \circ \iota$ is étale. By using [Hal14, Lemma 1.3.5] for the open substack $\sigma \circ \iota(\mathfrak{X}) \subset \mathrm{Filt}(\mathfrak{X})$, we obtain the identity $\sigma \circ \iota(\mathfrak{X}) = \mathrm{gr}^{-1}(\iota(\mathfrak{X}))$ as desired. \square

5.3. Symplectic and Lagrangian structures. Our goal in this section is to endow the attractor correspondence with a Lagrangian structure. For this we use the AKSZ formalism developed in [Pan+13; Cal15; CHS21].

Lemma 5.12. *Let Y be a finite product of stacks $B\mathbb{G}_m$ and $[\mathbb{A}^1/\mathbb{G}_m]$. Then:*

(1) *Pullback to the origin $\mathrm{pt} \rightarrow Y$ defines an isomorphism*

$$\mathbb{R}\Gamma(Y, \mathcal{O}) \simeq \mathbb{C}.$$

(2) *The stack Y is \mathcal{O} -compact.*

Proof. Let $p: Y \rightarrow \mathrm{pt}$ be the projection. By [BFN10] the stack Y is perfect and hence p_* is colimit-preserving.

To prove the first statement we have to show that $\mathrm{id} \rightarrow p_* p^*$ is an isomorphism. Since $\mathrm{QCoh}(Y_1 \times Y_2) \simeq \mathrm{QCoh}(Y_1) \otimes \mathrm{QCoh}(Y_2)$ by [BFN10, Theorem 4.7], it is enough to show this claim for $Y = B\mathbb{G}_m$ and $Y = [\mathbb{A}^1/\mathbb{G}_m]$, where it is obvious by an explicit computation.

By [HP23a, Theorem 4.2.1, Theorem 2.4.3 and Proposition 5.1.6] the pullback $p^*: \mathrm{QCoh}(\mathrm{pt}) \rightarrow \mathrm{QCoh}(Y)$ admits a colimit-preserving left adjoint. Therefore, Y is \mathcal{O} -compact by [NS23, Proposition 1.4]. \square

In particular, there is a natural isomorphism

$$[B\mathbb{G}_m]: \mathbb{R}\Gamma(B\mathbb{G}_m, \mathcal{O}) \longrightarrow \mathbb{C}.$$

Proposition 5.13. *The morphism $[B\mathbb{G}_m]: \mathbb{R}\Gamma(B\mathbb{G}_m, \mathcal{O}) \rightarrow \mathbb{C}$ defines a 0-orientation on $B\mathbb{G}_m$.*

Proof. We have to show that, for every connective commutative dg algebra R and a perfect complex $E \in \mathrm{Perf}(B\mathbb{G}_m \times \mathrm{Spec} R)$, the induced morphism

$$\mathbb{R}\Gamma(B\mathbb{G}_m, E) \longrightarrow \mathbb{R}\Gamma(B\mathbb{G}_m, E^{\vee})^{\vee}$$

is an isomorphism.

Using Proposition 5.8 we identify $\mathrm{Perf}(B\mathbb{G}_m \times \mathrm{Spec} R)$ with collections of perfect complexes $\{E_n\}_{n \in \mathbb{Z}}$ of R -modules, where only finitely many terms are nonzero. The dual of $E \in \mathrm{Perf}(B\mathbb{G}_m \times \mathrm{Spec} R)$ corresponding to $n \mapsto E_n$ is given by $E^{\vee} \in \mathrm{Perf}(B\mathbb{G}_m \times \mathrm{Spec} R)$ which corresponds to $n \mapsto (E_{-n})^{\vee}$ with the obvious evaluation map. We have $\mathbb{R}\Gamma(B\mathbb{G}_m, E) \simeq E_0$ and the induced morphism above is the canonical isomorphism

$$E_0 \longrightarrow ((E_0)^{\vee})^{\vee}.$$

\square

Corollary 5.14. *Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a k -shifted symplectic stack. Then $u^*\omega_{\mathfrak{X}}$ defines a k -shifted symplectic structure on the stack $\text{Grad}(\mathfrak{X})$ of graded objects. Let $\tau: \mathfrak{L} \rightarrow \mathfrak{X}$ be a k -shifted Lagrangian morphism specified by a homotopy $\tau^*\omega_{\mathfrak{X}} \sim 0$. Then its restriction under u^* defines a k -shifted Lagrangian structure on the morphism $\text{Grad}(\tau): \text{Grad}(\mathfrak{L}) \rightarrow \text{Grad}(\mathfrak{X})$.*

Proof. Since $B\mathbb{G}_m$ is 0-oriented by Proposition 5.13, applying the AKSZ construction [Pan+13, Theorem 2.5] to the stack $\text{Grad}(\mathfrak{X}) = \text{Map}(B\mathbb{G}_m, \mathfrak{X})$ we obtain a k -shifted symplectic structure obtained by the composite

$$\begin{aligned} \mathbb{C}[2-k](2) &\xrightarrow{[\omega_{\mathfrak{X}}]} \mathbf{DR}(\mathfrak{X}) \\ &\xrightarrow{\text{ev}^*} \mathbf{DR}(B\mathbb{G}_m \times \text{Grad}(\mathfrak{X})) \\ &\xrightarrow{\kappa_{\mathfrak{X}, B\mathbb{G}_m}} \mathbb{R}\Gamma(B\mathbb{G}_m, \mathcal{O}) \otimes \mathbf{DR}(\text{Grad}(\mathfrak{X})) \\ &\xrightarrow{[B\mathbb{G}_m] \otimes \text{id}} \mathbf{DR}(\text{Grad}(\mathfrak{X})) \end{aligned}$$

where $\text{ev}: B\mathbb{G}_m \times \text{Grad}(\mathfrak{X}) \rightarrow \mathfrak{X}$ is the evaluation map. By construction the orientation $[B\mathbb{G}_m]: \mathbb{R}\Gamma(B\mathbb{G}_m, \mathcal{O}) \rightarrow \mathbb{C}$ is given by the pullback along $i: \text{pt} \rightarrow B\mathbb{G}_m$. Unpacking the construction of the map $\kappa_{\mathfrak{X}, B\mathbb{G}_m}$ given in [Pan+13], we see that the composite

$$\begin{aligned} \mathbf{DR}(B\mathbb{G}_m \times \text{Grad}(\mathfrak{X})) &\xrightarrow{\kappa_{\mathfrak{X}, B\mathbb{G}_m}} \mathbb{R}\Gamma(B\mathbb{G}_m, \mathcal{O}) \otimes \mathbf{DR}(\text{Grad}(\mathfrak{X})) \\ &\xrightarrow{i^* \otimes \text{id}} \mathbf{DR}(\text{Grad}(\mathfrak{X})) \end{aligned}$$

is equivalent to

$$\mathbf{DR}(B\mathbb{G}_m \times \text{Grad}(\mathfrak{X})) \xrightarrow{(i \times \text{id})^*} \mathbf{DR}(\text{Grad}(\mathfrak{X})).$$

But $\text{ev} \circ (i \times \text{id}): \mathfrak{X} \rightarrow \text{Grad}(\mathfrak{X})$ is precisely u . The claim for a Lagrangian $\mathfrak{L} \rightarrow \mathfrak{X}$ is proven analogously. \square

We will now show that the shifted symplectic structure on $\text{Grad}(\mathfrak{X})$ defined in Corollary 5.14 is compatible with taking derived critical loci. Let \mathfrak{Y} be a derived higher Artin stack locally of finite presentation and $f: \mathfrak{Y} \rightarrow \mathbb{A}^1[n]$ be an n -shifted function. Set $\mathfrak{X} := \text{Crit}(f)$ and let $\omega_{\mathfrak{X}}$ be the $(n-1)$ -shifted symplectic structure in Example 3.10.

Lemma 5.15. *There exists a natural equivalence of $(n-1)$ -shifted symplectic stacks*

$$(\text{Grad}(\mathfrak{X}), u^*\omega_{\mathfrak{X}}) \simeq (\text{Crit}(f|_{\text{Grad}(\mathfrak{Y})}), \omega_{\text{Crit}(f|_{\text{Grad}(\mathfrak{Y})})})$$

where $\omega_{\text{Crit}(f|_{\text{Grad}(\mathfrak{Y})})}$ is the $(n-1)$ -shifted symplectic structure constructed in Example 3.10.

Proof. Recall that the (-1) -shifted symplectic structure $\omega_{\mathfrak{X}}$ is constructed using the following Lagrangian intersection diagram:

$$\begin{array}{ccc} \mathfrak{X} & \longrightarrow & \mathfrak{Y} \\ \downarrow & & \downarrow df \\ \mathfrak{Y} & \xrightarrow{0} & T^*\mathfrak{Y}. \end{array}$$

Passing to the mapping stack from $B\mathbb{G}_m^n$ and using Corollary 5.14, we obtain the following Lagrangian intersection diagram:

$$(5.4) \quad \begin{array}{ccc} \text{Grad}(\mathfrak{X}) & \longrightarrow & \text{Grad}(\mathfrak{Y}) \\ \downarrow & & \downarrow \text{Grad}(df) \\ \text{Grad}(\mathfrak{Y}) & \xrightarrow{0} & \text{Grad}(T^*[n]\mathfrak{Y}) \end{array}$$

where $(df)_\alpha^0$ denote the degree zero part of $df|_{\mathfrak{U}_\alpha}$. On the other hand, we have the following Lagrangian intersection diagram:

$$(5.5) \quad \begin{array}{ccc} \text{Crit}(f|_{\text{Grad}(\mathfrak{Y})}) & \longrightarrow & \text{Grad}(\mathfrak{Y}) \\ \downarrow & & \downarrow d\text{Grad}(f) \\ \text{Grad}(\mathfrak{Y}) & \xrightarrow{0} & T^*[n]\text{Grad}(\mathfrak{Y}). \end{array}$$

It is enough to show that the diagrams (5.4) and (5.5) are identified as Lagrangian intersection diagrams. Let $\gamma_{\mathfrak{Y}}$ (resp. $\gamma_{\text{Grad}(\mathfrak{Y})}$) be the tautological one-form on $T^*[n]\mathfrak{Y}$ (resp. $T^*[n]\text{Grad}(\mathfrak{Y})$). Then we have a natural identification $u^*\gamma_{\mathfrak{Y}} \simeq \gamma_{\text{Grad}(\mathfrak{Y})}$ under the equivalence $T^*[n]\text{Grad}(\mathfrak{Y}) \simeq \text{Grad}(T^*[n]\mathfrak{Y})$, which implies that it preserves the canonical n -shifted symplectic structures. It is obvious to show that the Lagrangian structures are identified as well. \square

It is obvious that the point pt has a canonical 0-orientation

$$[\text{pt}]: \mathbb{R}\Gamma(\text{pt}, \mathcal{O}) \longrightarrow \mathbb{C}.$$

Now consider the cospan

$$\begin{array}{ccc} & [\mathbb{A}^1/\mathbb{G}_m] & \\ i_0 \nearrow & & \nwarrow i_1 \\ B\mathbb{G}_m & & \text{pt} \end{array}$$

of derived stacks.

Proposition 5.16. *There is a unique structure of a 0-preoriented cospan on*

$$\begin{array}{ccc} & [\mathbb{A}^1/\mathbb{G}_m] & \\ i_0 \nearrow & & \nwarrow i_1 \\ B\mathbb{G}_m & & \text{pt} \end{array}$$

which restricts to the orientations $[B\mathbb{G}_m]$ and $[\text{pt}]$ on $B\mathbb{G}_m$ and pt . Moreover, this cospan is oriented.

Proof. By Lemma 5.12 the functor $\mathbb{R}\Gamma(-, \mathcal{O})$ applied to this cospan gives the correspondence

$$\begin{array}{ccc} & \mathbb{C} & \\ \swarrow & & \searrow \\ \mathbb{C} & & \mathbb{C} \end{array}$$

In particular, the existence and uniqueness of a preorientation is obvious.

Let R be a connective commutative dg algebra and $E \in \text{Perf}([\mathbb{A}^1/\mathbb{G}_m] \times \text{Spec } R)$, i.e. a filtered object $\{E_n\}$ by Proposition 5.8. By [CHS21, Definition 2.5.4] we have to show that the diagram

$$\begin{array}{ccc} \mathbb{R}\Gamma([\mathbb{A}^1/\mathbb{G}_m], E) & \longrightarrow & \mathbb{R}\Gamma(\text{pt}, i_1^*E) \\ \downarrow & & \downarrow \\ \mathbb{R}\Gamma(B\mathbb{G}_m, i_0^*E) & \longrightarrow & \mathbb{R}\Gamma([\mathbb{A}^1/\mathbb{G}_m], E^\vee)^\vee \end{array}$$

is Cartesian. Any filtered object E fits into a fibre sequence

$$\tau_{\geq 0}E \rightarrow E \rightarrow \tau_{\leq -1}E,$$

where the associated graded of $\tau_{\geq 0}E$ is concentrated in weights ≥ 0 and the associated graded of $\tau_{\leq -1}E$ is concentrated in weights ≤ -1 . By stability, it is enough to prove the claim for filtered objects E with associated graded concentrated in weights ≥ 0 and in weights ≤ -1 . Replacing E by E^\vee and using the symmetry of the diagram, it is enough to prove the claim for filtered objects whose associated graded is concentrated in weights ≥ 0 . Further, by stability, it is enough to prove the claim for filtered objects whose associated graded is $V \in \text{Perf}_R$ concentrated in a single weight $n \geq 0$.

If $n > 0$, the diagram is

$$\begin{array}{ccc} V & \xrightarrow{\text{id}} & V \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & 0 \end{array}$$

while for $n = 0$ the diagram is

$$\begin{array}{ccc} V & \xrightarrow{\text{id}} & V \\ \downarrow \text{id} & & \downarrow \\ V & \longrightarrow & (V^\vee)^\vee \end{array}$$

Both of these are obviously Cartesian. □

Remark 5.17. The proof of the above proposition is sketched in [HP23b, Lemma 7.4]. While we were preparing this paper, the same statement was proved in [Bu24, Theorem 3.1.5].

Corollary 5.18. *Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a k -shifted symplectic stack. Then the attractor correspondence*

$$\begin{array}{ccc} & \text{Filt}(\mathfrak{X}) & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \text{Grad}(\mathfrak{X}) & & \mathfrak{X} \end{array}$$

carries a natural structure of a k -shifted Lagrangian correspondence.

Proof. By definition, the attractor correspondence (5.1) is obtained by applying $\text{Map}(-, \mathfrak{X})$ to the cospan from Proposition 5.16. The AKSZ construction [CHS21, Proposition 3.4.2] implies that the attractor correspondence is Lagrangian. □

Next, consider the diagram


$$\begin{array}{ccccc} & & B\mathbb{G}_m \times [\mathbb{A}^1/\mathbb{G}_m] & & [\mathbb{A}^1/\mathbb{G}_m] \\ & & \swarrow \text{id} \times i_1 & & \swarrow i_0 \\ & & B\mathbb{G}_m & & \\ & \swarrow \text{id} \times i_0 & & \searrow \text{id} \times i_1 & \\ B\mathbb{G}_m^2 & & & & [\mathbb{A}^1/\mathbb{G}_m]^2 \\ & \searrow i_0 \times \text{id} & & \swarrow i_1 \times \text{id} & \\ & & [\mathbb{A}^1/\mathbb{G}_m] \times B\mathbb{G}_m & & B\mathbb{G}_m \\ & & \swarrow i_1 \times \text{id} & & \swarrow i_0 \\ & & [\mathbb{A}^1/\mathbb{G}_m] & & \text{pt} \\ & & \swarrow i_1 & & \swarrow i_1 \end{array}$$

Proposition 5.19. *There is a unique structure of a 0-preoriented 2-fold cospan on*

$$\begin{array}{ccc} (B\mathbb{G}_m \times [\mathbb{A}^1/\mathbb{G}_m]) \amalg_{B\mathbb{G}_m} [\mathbb{A}^1/\mathbb{G}_m] & & \\ \swarrow & \downarrow & \swarrow \\ B\mathbb{G}_m^2 & [\mathbb{A}^1/\mathbb{G}_m]^2 & \text{pt} \\ \searrow & \uparrow & \searrow \\ ([\mathbb{A}^1/\mathbb{G}_m] \times B\mathbb{G}_m) \amalg_{B\mathbb{G}_m} [\mathbb{A}^1/\mathbb{G}_m] & & \end{array}$$


which restricts to the orientations $[B\mathbb{G}_m]$ and $[\text{pt}]$ on $B\mathbb{G}_m$ and pt . Moreover, this 2-fold cospan is oriented.

Proof. By Lemma 5.12 the functor $\mathbb{R}\Gamma(-, \mathcal{O})$ applied to these cospans gives the 2-fold correspondence




with all maps given by the identities. In particular, the existence and uniqueness of a preorientation is obvious.

Let R be a connective commutative dg algebra and $E \in \text{Perf}([\mathbb{A}^1/\mathbb{G}_m]^2 \times \text{Spec } R)$, i.e. a doubly filtered object $\{E_{n,m}\}$ by Proposition 5.8. By [CHS21, Definition 2.7.5] we have to show that the diagram




is a limit diagram, where we omit obvious pullbacks of E to the corresponding stacks. As in the proof of Proposition 5.16, it is enough to assume that the doubly filtered object E has associated graded given by the object $V \in \text{Perf}_R$ concentrated in a single weight (n, m) with $n, m \geq 0$.

If $n, m > 0$, we have to show that the diagram



is a limit diagram, where the nontrivial maps are all the identities. This is reduced to proving that the square



is Cartesian, which is obvious. The analysis of the other cases (when either n or m is zero) is done analogously. \square

Corollary 5.20. *Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a k -shifted symplectic stack. Then*

$$\begin{array}{ccccc}
 & & \text{Filt}(\text{Grad}(\mathfrak{X})) \times_{\text{Grad}(\mathfrak{X})} \text{Filt}(\mathfrak{X}) & & \\
 & \swarrow & \uparrow & \searrow & \\
 \text{Grad}^2(\mathfrak{X}) & & \text{Filt}^2(\mathfrak{X}) & & \mathfrak{X} \\
 & \swarrow & \downarrow & \searrow & \\
 & & \text{Grad}(\text{Filt}(\mathfrak{X})) \times_{\text{Grad}(\mathfrak{X})} \text{Filt}(\mathfrak{X}) & &
 \end{array}$$

has a natural structure of a 2-fold Lagrangian correspondence.

Proof. The 2-fold correspondence in the statement is obtained by applying $\text{Map}(-, \mathfrak{X})$ to the 2-fold iterated cospan from Proposition 5.19. The claim then follows from the AKSZ construction [CHS21, Proposition 3.4.2]. \square

5.4. Matching locus. Let $u_1, u_2: \text{Grad}^2(\mathfrak{X}) \rightarrow \text{Grad}(\mathfrak{X})$ be morphisms induced by $\mathbb{G}_m \times \{0\} \rightarrow \mathbb{G}_m^2$ and $\{0\} \times \mathbb{G}_m \rightarrow \mathbb{G}_m^2$, respectively. We will introduce the notion of matching points:

Definition 5.21. For a classical Artin stack (resp. derived Artin stack locally of finite presentation), we say that a \mathbb{C} -valued point $x \in \text{Grad}^2(\mathfrak{X})$ is **matching** if for each $(m, l) \in \mathbb{Z}^2$ with $m \cdot l < 0$ and $i = -1, 0, 1$ (resp. for any $i \in \mathbb{Z}$), the weight (m, l) -part with respect to the \mathbb{G}_m^2 -action

$$(5.6) \quad H^i(\mathbb{L}_{\mathfrak{X}, u(x)})_{(m, l)}$$

vanishes.

Roughly speaking, the matching point is a point where $u_1(x)$ is sufficiently close to $u_2(x)$. Clearly, the collection of matching points $\text{Grad}^2(\mathfrak{X})_{\text{mat}} \subset \text{Grad}^2(\mathfrak{X})$ form an open substack, which we call the matching locus. The following proposition motivates the definition of the matching points⁵:

Proposition 5.22. *Let \mathfrak{X} be a quasi-separated classical Artin stack with affine stabilizers. Consider the following commutative diagram.*

$$\begin{array}{ccccc}
 \text{Filt}(\text{Grad}(\mathfrak{X})) \times_{\text{Grad}(\mathfrak{X})} \text{Filt}(\mathfrak{X}) & \xleftarrow{\alpha} & \text{Filt}^2(\mathfrak{X}) & \xrightarrow{\beta} & \text{Grad}(\text{Filt}(\mathfrak{X})) \times_{\text{Grad}(\mathfrak{X})} \text{Filt}(\mathfrak{X}) \\
 & & \searrow & & \swarrow \\
 & & \text{Grad}^2(\mathfrak{X}) & &
 \end{array}$$

Then the maps α and β are invertible over $\text{Grad}^2(\mathfrak{X})_{\text{mat}}$. The same statement holds true for derived Artin stacks.

Proof. We first prove the statement for classical Artin stacks. Let α_{mat} denote the base change of α along the open inclusion $\text{Grad}^2(\mathfrak{X})_{\text{mat}} \rightarrow \text{Grad}^2(\mathfrak{X})$. It is enough to prove the invertibility of α_{mat} . We first show that α_{mat} is étale. To see this, it is enough to show the vanishing of the cohomology of cotangent complex $H^i(\mathbb{L}_{\alpha_{\text{mat}}}|_x)$ for each \mathbb{C} -valued points $x \in \text{Filt}^2(\mathfrak{X})$ over the matching locus and $i = -1, 0, 1$. Since the set of \mathbb{C} -valued points $x \in \text{Filt}^2(\mathfrak{X})$ over the matching locus with vanishing $H^i(\mathbb{L}_{\alpha_{\text{mat}}}|_x)$ is open, by using [Hal14, Lemma 1.3.5], it is enough to prove the vanishing for points of the form $y = \sigma(x)$ for some $y \in \text{Grad}^2(\mathfrak{X})$. In this case, the claim follows from the identification

$$H^i(\mathbb{L}_{\alpha_{\text{mat}}}|_{\sigma(y)}) \simeq \bigoplus_{m < 0, l > 0} H^i(\mathbb{L}_{\mathfrak{X}, u(y)})_{(m, l)}$$

and the vanishing (5.6).

⁵We thank Andrés Ibáñez Núñez for sharing with us the proof of a related statement.

Next, we show that α_{mat} is bijective on \mathbb{C} -valued points. Consider the following diagram of stacks

$$\begin{array}{ccc} [0/\mathbb{G}_m] \times \{1\} & \longrightarrow & [\mathbb{A}^1/\mathbb{G}_m] \times \{1\} \\ \downarrow & & \downarrow \\ [0/\mathbb{G}_m] \times [\mathbb{A}^1/\mathbb{G}_m] & \longrightarrow & [\mathbb{A}^2/\mathbb{G}_m^2]. \end{array}$$

where the horizontal arrows are canonical closed inclusions and the vertical arrows are canonical open immersions. By considering the mapping spaces from these stacks to \mathfrak{X} , we obtain the following commutative diagram:

$$\begin{array}{ccc} \text{Filt}^2(\mathfrak{X})(\mathbb{C}) & \longrightarrow & \text{Filt}(\text{Grad}(\mathfrak{X}))(\mathbb{C}) \\ \downarrow & & \downarrow \\ \text{Filt}(\mathfrak{X})(\mathbb{C}) & \longrightarrow & \text{Grad}(\mathfrak{X})(\mathbb{C}). \end{array}$$

It follows from [Hal14, Lemma 3.4.4] that this diagram is Cartesian over the matching locus as desired.

Since the diagonal map $\Delta_{\alpha_{\text{mat}}}$ is representable, étale and bijective on \mathbb{C} -valued points, it is an isomorphism by [Sta24, Tag 05W5]⁶. In particular, α_{mat} is an étale representable morphism. Using [Sta24, Tag 05W5] again, we conclude that α_{mat} is an isomorphism.

The statement for derived Artin stacks follows since α_{mat} is a classical isomorphism with a vanishing cotangent complex. \square

By Corollary 5.20, we obtain the following corollary:

Corollary 5.23. *Assume that \mathfrak{X} is a (-1) -shifted symplectic derived Artin stack. Then we have a natural equivalence between the following composite Lagrangian correspondences:*

$$\begin{array}{ccccc} & & (\text{Filt}(\text{Grad}(\mathfrak{X})) \times_{\text{Grad}(\mathfrak{X})} \text{Filt}(\mathfrak{X}))_{\text{mat}} & & \\ & \swarrow & & \searrow & \\ \text{Filt}(\text{Grad}(\mathfrak{X}))_{\text{mat}} & & & & \text{Filt}(\mathfrak{X}) \\ \swarrow & & \searrow & & \swarrow \searrow \\ \text{Grad}^2(\mathfrak{X})_{\text{mat}} & & \text{Grad}(\mathfrak{X}) & & \mathfrak{X} \\ & & & & \\ & & (\text{Grad}(\text{Filt}(\mathfrak{X})) \times_{\text{Grad}(\mathfrak{X})} \text{Filt}(\mathfrak{X}))_{\text{mat}} & & \\ & \swarrow & & \searrow & \\ \text{Grad}(\text{Filt}(\mathfrak{X}))_{\text{mat}} & & & & \text{Filt}(\mathfrak{X}) \\ \swarrow & & \searrow & & \swarrow \searrow \\ \text{Grad}^2(\mathfrak{X})_{\text{mat}} & & \text{Grad}(\mathfrak{X}) & & \mathfrak{X}. \end{array}$$

Here $(-)\text{mat}$ denotes the base change along the open inclusion $\text{Grad}^2(\mathfrak{X})_{\text{mat}} \subset \text{Grad}^2(\mathfrak{X})$. Further, as a correspondence of derived Artin stacks, they are naturally equivalent to the following correspondence:

$$\begin{array}{ccc} & \text{Filt}^2(\mathfrak{X})_{\text{mat}} & \\ & \swarrow \searrow & \\ \text{Grad}^2(\mathfrak{X})_{\text{mat}} & & \mathfrak{X}. \end{array}$$

⁶Note that a morphism between algebraic spaces locally of finite type over algebraically closed field k is universally injective if and only if injective on k -valued points: see e.g. [Ang].

5.5. Equivariant chart of stacks. In this section, we will introduce the notion of a \mathbb{G}_m^n -equivariant chart of \mathfrak{X} , which will be used to define charts for $\text{Grad}(\mathfrak{X})$.

Definition 5.24.

- (1) Let \mathfrak{X} be an Artin stack. A \mathbb{G}_m^n -**equivariant chart** is a tuple $\Omega = (X, \mu, \bar{q})$ where X is a quasi-separated algebraic space, μ is a \mathbb{G}_m^n -action on X and $\bar{q}: [X/\mathbb{G}_m^n] \rightarrow \mathfrak{X}$ is a smooth morphism.
- (2) A **morphism of \mathbb{G}_m^n -equivariant charts** from $\Omega_1 = (X_1, \mu_1, \bar{q}_1)$ to $\Omega_2 = (X_2, \mu_2, \bar{q}_2)$ is given by a smooth \mathbb{G}_m^n -equivariant morphism $f: X_1 \rightarrow X_2$ together with a homotopy $\bar{q}_1 \sim \bar{q}_2 \circ [f/\mathbb{G}_m^n]$. We let $\text{Cha}_{\mathfrak{X}}^{\mathbb{G}_m^n}$ denote the category of \mathbb{G}_m^n -equivariant charts on \mathfrak{X} .

For a \mathbb{G}_m^n -equivariant chart of \mathfrak{X} we introduce the following additional notation:

- We let $q: X \rightarrow \mathfrak{X}$ be the composite $X \rightarrow [X/\mathbb{G}_m^n] \xrightarrow{\bar{q}} \mathfrak{X}$.
- We let $q_\mu: X^\mu \rightarrow \text{Grad}(\mathfrak{X})$ be the composite $X^\mu \rightarrow \text{Grad}([X/\mathbb{G}_m^n]) \xrightarrow{\text{Grad}(\bar{q})} \text{Grad}(\mathfrak{X})$, where the first morphism is an inclusion of a component as in Theorem 5.6. It follows from [Hal14, Proposition 1.3.1 (4)] that q_μ is a smooth morphism.

The following statement will be useful to glue sections of sheaves on $\text{Grad}(\mathfrak{X})$.

Proposition 5.25 ([Hal14, Lemma 4.4.6]). *Let \mathfrak{X} be a quasi-separated Artin stack with affine stabilizers. Then the morphism*

$$\left(\coprod_{\Omega=(X,\mu,\bar{q}) \in \text{Cha}_{\mathfrak{X}}^{\mathbb{G}_m^n}} X^\mu \right) \xrightarrow{\coprod q_\mu} \text{Grad}^n(\mathfrak{X}).$$

is a smooth atlas of $\text{Grad}^n(\mathfrak{X})$.

Proof. Since q_μ is smooth, the morphism in the statement is smooth, so we just need to show that it is surjective. To see this, take a point $x \in \mathfrak{X}$ and a map $\iota: \mathbb{G}_m^n \rightarrow G_x$ where G_x denote the stabilizer group. Set $T := \iota(\mathbb{G}_m^n)$. Then by using [AHR20, Theorem 1.1], we can find an affine scheme $\text{Spec } A$ with a T -action μ together with a point $\hat{x} \in \text{Spec } A^T$ and a smooth morphism $[\text{Spec } A/T] \rightarrow \mathfrak{X}$ which restricts to the natural morphism $BT \rightarrow BG_x$. Let $\tilde{\mu}$ be a \mathbb{G}_m^n -action on $\text{Spec } A$ induced by μ and $\bar{q}: [\text{Spec } A/\mathbb{G}_m^n] \rightarrow \mathfrak{X}$ be the naturally induced morphism. Then the tuple $(\text{Spec } A, \tilde{\mu}, \bar{q})$ is a \mathbb{G}_m^n -equivariant chart of \mathfrak{X} and the morphism $q_{\tilde{\mu}}: \text{Spec } A^{\mathbb{G}_m^n} \rightarrow \text{Grad}(\mathfrak{X})$ maps \hat{x} to x . \square

Corollary 5.26. *Let \mathfrak{X} be a quasi-separated Artin stack with affine stabilizers. Assume that we are given a lisse-étale sheaf \mathcal{F} on $\text{Grad}^n(\mathfrak{X})$. Assume further that we are given the following data:*

- (1) *For each $\Omega = (X, \mu, \bar{q}) \in \text{Cha}_{\mathfrak{X}}^{\mathbb{G}_m^n}$, a choice of a section $s_\Omega \in \Gamma(X^\mu, q_\mu^* \mathcal{F})$ which satisfies the following condition: For a morphism $f: \Omega_1 = (X_1, \mu_1, \bar{q}_1) \rightarrow \Omega_2 = (X_2, \mu_2, \bar{q}_2)$ in $\text{Cha}_{\mathfrak{X}}^{\mathbb{G}_m^n}$, the following identity in $\Gamma(X_1^{\mu_1}, q_{1,\mu_1}^* \mathcal{F})$ holds:*

$$(5.7) \quad f^{\mathbb{G}_m, * } s_{\Omega_2} = s_{\Omega_1}.$$

Here $f^{\mathbb{G}_m}: X_1^{\mu_1} \rightarrow X_2^{\mu_2}$ is the induced morphism.

Then there exists a section $s \in \Gamma(\text{Grad}(\mathfrak{X}), \mathcal{F})$ such that for each $\Omega = (X, \mu, \bar{q}) \in \text{Cha}_{\mathfrak{X}}^{\mathbb{G}_m^n}$, the identity $q_\mu^* s = s_\Omega$ holds.

Proof. Consider the following diagram:

$$\left(\coprod_{\substack{\Omega_1, \Omega_2 \in \text{Cha}_{\mathfrak{X}}^{\mathbb{G}_m^n}, \\ \Omega_1 = (X_1, \mu_1, \bar{q}_1), \\ \Omega_2 = (X_2, \mu_2, \bar{q}_2)}} X_1^{\mu_1} \times_{\text{Grad}(\mathfrak{X})} X_2^{\mu_2} \right) \begin{array}{c} \xrightarrow{\text{pr}_1} \\ \xrightarrow{\text{pr}_2} \end{array} \left(\coprod_{\substack{\Omega \in \text{Cha}_{\mathfrak{X}}^{\mathbb{G}_m^n}, \\ \Omega = (X, \mu, \bar{q})}} X^\mu \right) \xrightarrow{\coprod q_\mu} \text{Grad}^n(\mathfrak{X}).$$

By using the descent of a section of a lisse-étale sheaf, it is enough to prove the identity $\text{pr}_1^* \{s_\Omega\}_\Omega = \text{pr}_2^* \{s_\Omega\}_\Omega$. We take objects $\Omega_i = (X_i, \mu_i, \bar{q}_i) \in \text{Cha}_{\mathfrak{X}}^{\mathbb{G}_m^n}$ for $i = 1, 2$. We set $Y := X_1 \times_{\mathfrak{X}} X_2$. Then Y admits

a diagonal action ν of \mathbb{G}_m^n . Let $\bar{r}: [Y/\mathbb{G}_m^n] \rightarrow \mathfrak{X}$ be the natural morphism and define a \mathbb{G}_m^n -equivariant chart $\mathfrak{R} = (Y, \nu, \bar{r})$. Consider the following Cartesian diagram:

$$\begin{array}{ccc} [Y/\mathbb{G}_m^n \times \mathbb{G}_m^n] & \longrightarrow & [X_1/\mathbb{G}_m^n] \\ \downarrow & & \downarrow \\ [X_2/\mathbb{G}_m^n] & \longrightarrow & \mathfrak{X}. \end{array}$$

By taking the mapping stack from $B\mathbb{G}_m^n$, we obtain the following Cartesian diagram:

$$\begin{array}{ccc} [Y^\nu/\mathbb{G}_m^n \times \mathbb{G}_m^n] & \longrightarrow & [X_1^{\mu_1}/\mathbb{G}_m^n] \\ \downarrow & & \downarrow \\ [X_2^{\mu_2}/\mathbb{G}_m^n] & \longrightarrow & \text{Grad}(\mathfrak{X}). \end{array}$$

In particular, there exists a natural isomorphism $Y^\nu \cong X_1^{\mu_1} \times_{\text{Grad}(\mathfrak{X})} X_2^{\mu_2}$ over $\text{Grad}(\mathfrak{X})$. Under this identification, we have an identity

$$\text{pr}_1^* s_{\Omega_1} = s_{\mathfrak{R}} = \text{pr}_2^* s_{\Omega_2}$$

as desired. □

By a similar argument, we can prove the following statement:

Corollary 5.27. *Let \mathfrak{X} be a quasi-separated Artin stack with affine stabilizers. Assume that we are given a lisse-étale stack in groupoids \mathfrak{F} on $\text{Grad}^n(\mathfrak{X})$. Assume further that we are given the following data:*

- (1) *For each $\Omega = (X, \mu, \bar{q}) \in \text{Cha}_{\mathfrak{X}^m}^{\mathbb{G}_m^n}$, there exists a section $o_\Omega \in \Gamma(X^\mu, q_\mu^* \mathfrak{F})$.*
- (2) *For a morphism $f: (\Omega_1 = (X_1, \mu_1, \bar{q}_1)) \rightarrow (\Omega_2 = (X_2, \mu_2, \bar{q}_2))$, there exists an isomorphism*

$$\Theta_f: o_{\Omega_1} \simeq f^{\mathbb{G}_m, * } o_{\Omega_2}$$

such that if we are further given a morphism $g: \Omega_2 \rightarrow \Omega_3$, the following diagram commutes:

$$(5.8) \quad \begin{array}{ccc} o_{\Omega_1} & \xrightarrow[\simeq]{\Theta_f} & f^{\mathbb{G}_m, * } o_{\Omega_2} \\ \simeq \downarrow \Theta_{g \circ f} & & \simeq \downarrow \Theta_g \\ (g^{\mathbb{G}_m} \circ f^{\mathbb{G}_m})^* o_{\Omega_3} & \xrightarrow{\simeq} & f^{\mathbb{G}_m, * } g^{\mathbb{G}_m, * } o_{\Omega_3}. \end{array}$$

Then there exists a section $o \in \Gamma(\mathfrak{X}, \mathfrak{F})$ such that for each $\Omega \in \text{Cha}_{\mathfrak{X}^m}^{\mathbb{G}_m^n}$, there exists an isomorphism $\Theta_q: o_\Omega \simeq q_\mu^ o$ such that the associativity property as in (5.8) holds.*

The following technical statements will be used for the associativity of the cohomological Hall algebras later:

Lemma 5.28. *Let \mathfrak{X} be a quasi-separated Artin stack with affine stabilizer groups. Then for any closed point $x \in \text{Grad}^2(\mathfrak{X})_{\text{mat}}$, one can find a \mathbb{G}_m^2 -equivariant chart $\Omega = (X, \hat{\mu} = (\mu_1, \mu_2), \bar{q})$ and a point $\hat{x} \in X_{\text{mat}}^{\hat{\mu}}$ such that $q_{\hat{\mu}}(\hat{x}) = x$ holds.*

Proof. By using [AHR20, Theorem 1.1], we may take a \mathbb{G}_m^2 -equivariant chart $(\text{Spec } A, \hat{\mu} = (\mu_1, \mu_2), \bar{q})$ and a point $\hat{x} \in \text{Spec } A^{\hat{\mu}}$ with $q_{\hat{\mu}}(\hat{x}) = x$. By possibly replacing $\text{Spec } A$ with its \mathbb{G}_m^2 -equivariant subspace containing x , we may assume that $T_{\text{Spec } A, \hat{x}} \cong T_{\mathfrak{X}, u(x)}$ holds. In particular, \hat{x} is contained in the matching locus. □

As an application of Proposition 5.25, we will explain that a d-critical structure is inherited to the stack of graded points:

Proposition 5.29. *Let (\mathfrak{X}, s) be a d-critical stack with \mathfrak{X} quasi-separated with affine stabilizers. Then $(\text{Grad}^n(\mathfrak{X}), u^* s)$ is a d-critical stack.*

Proof. By Proposition 5.25 the collection of \mathbb{G}_m^n -equivariant charts $\mathfrak{Q} = (X, \mu, \bar{q}) \in \text{Cha}_{\mathfrak{X}}^{\mathbb{G}_m^n}$ defines a smooth atlas $\{q_\mu: X^\mu \rightarrow \text{Grad}^n(\mathfrak{X})\}$. Therefore, by [Joy15, Proposition 2.54] it is enough to show that $q_\mu^* u^* s$ is a d-critical structure on X^μ for each equivariant chart. Using the commutative diagram

$$\begin{array}{ccc} X^\mu & \longrightarrow & X \\ \downarrow q_\mu & & \downarrow q \\ \text{Grad}^n(\mathfrak{X}) & \xrightarrow{u} & \mathfrak{X} \end{array}$$

we have $q_\mu^* u^* s = (q^* s)^\mu$. Since (\mathfrak{X}, s) is a d-critical stack, $q^* s$ is a d-critical structure and hence $(q^* s)^\mu$ is a d-critical structure by Corollary 3.30. \square

We now compare the d-critical structure constructed in Proposition 5.29 and the (-1) -shifted symplectic structure constructed in Corollary 5.14.

Proposition 5.30. *Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a (-1) -shifted symplectic Artin stack with \mathfrak{X}^{cl} quasi-separated with affine stabilizers. Let $(\mathfrak{X}^{\text{cl}}, s)$ be the underlying d-critical locus. Then the d-critical structure underlying $(\text{Grad}^n(\mathfrak{X}), u^* \omega_{\mathfrak{X}})$ is $(\text{Grad}^n(\mathfrak{X}^{\text{cl}}), u^* s)$.*

Proof. Consider the commutative diagram

$$\begin{array}{ccc} \text{Grad}^n(\mathfrak{X}^{\text{cl}}) & \xrightarrow{\text{Grad}^n(j)} & \text{Grad}^n(\mathfrak{X}) \\ \downarrow u^{\text{cl}} & & \downarrow u \\ \mathfrak{X}^{\text{cl}} & \xrightarrow{j} & \mathfrak{X}. \end{array}$$

Identifying sections of \mathcal{S}^0 with closed 2-forms of degree -1 using (3.10), the d-critical structure s becomes $j^* \omega_{\mathfrak{X}}$ by Theorem 3.23. Similarly, the d-critical structure underlying $u^* \omega_{\mathfrak{X}}$ becomes $\text{Grad}^n(j)^* u^* \omega_{\mathfrak{X}} = u^{\text{cl},*} j^* \omega_{\mathfrak{X}}$, hence we conclude. \square

6. INDEX LINE BUNDLES AND LOCALIZATION OF ORIENTATIONS

Let (\mathfrak{X}, s, o) be an oriented d-critical stack. The aim of this section is to define an orientation $u^* o$ on $\text{Grad}(\mathfrak{X})$ in a way that is compatible with a similar construction for orientations of (-1) -shifted symplectic stacks we have already seen in Lemma 3.19. To do this, we will construct the index line bundle $\mathcal{L}_{\mathfrak{X}, s}$ on $\text{Grad}(\mathfrak{X})^{\text{red}}$ together with an isomorphism $K_{\text{Grad}(\mathfrak{X}), u^* s} \otimes \mathcal{L}_{\mathfrak{X}, s}^{\otimes 2} \cong K_{\mathfrak{X}, s}|_{\text{Grad}(\mathfrak{X})^{\text{red}}}$.

In §6.2, we will introduce the notion of indices for d-critical stacks, which is a locally constant function on graded points. In §6.3 and §6.4, we will introduce index line bundles for \mathbb{G}_m -equivariant d-critical spaces and d-critical stacks, respectively. We recommend readers skip these two sections for the first read since it is technical. In §6.5, we will explain the construction of localized orientation on the stack of graded points and prove its properties.

6.1. Notation on rank and determinant for graded complexes. Throughout the section, we will use the following notations: If we are given a graded perfect complex $E^\bullet = \{E^n\}_{n \in \mathbb{Z}}$ on a derived Artin stack \mathfrak{X} , we set E^+ (resp. E^-) denote the positive (resp. negative) part of E and $E^\pm := E^+ \oplus E^-$. We define

$$\begin{aligned} \text{rank}_+(E^\bullet) &:= \text{rank}(E^+), & \text{rank}_-(E^\bullet) &:= \text{rank}(E^-), & \text{rank}_0(E^\bullet) &:= \text{rank}(E^0), \\ \widehat{\det}_+(E^\bullet) &:= \widehat{\det}(E^+), & \widehat{\det}_-(E^\bullet) &:= \widehat{\det}(E^-), & \widehat{\det}_0(E^\bullet) &:= \widehat{\det}(E^0), & \widehat{\det}_\pm(E) &:= \widehat{\det}(E^\pm). \end{aligned}$$

For a \mathbb{G}_m -equivariant algebraic space X and a \mathbb{G}_m -equivariant perfect complex E on X , we regard $E|_{X^{\mathbb{G}_m}}$ as a graded vector space using the \mathbb{G}_m -action and define E^+ to be the positive part of $E|_{X^{\mathbb{G}_m}}$. We define $\text{rank}_+(E) := \text{rank}(E^+)$ and similarly for other operations.

For an Artin stack \mathfrak{X} and a perfect complex E on \mathfrak{X} , we regard $E|_{\text{Grad}(\mathfrak{X})}$ as a graded complex by using the \mathbb{G}_m -action and define E^+ to be the positive part of $E|_{\text{Grad}(\mathfrak{X})}$. We define $\text{rank}_+(E) := \text{rank}(E^+)$ and similarly for other operations.

For a doubly graded perfect complex $E^{\bullet, \bullet}$ and a subset $\Sigma \subset \mathbb{Z}^2$, we let $E^\Sigma \subset E$ denote the subspace consisting of complexes whose grading is concentrated in Σ . We set $\text{rank}_\Sigma(E) := \text{rank } E^\Sigma$ and $\widehat{\det}_\Sigma(E) := \widehat{\det}(E^\Sigma)$. We define rank_Σ and $\widehat{\det}_\Sigma$ for \mathbb{G}_m^2 -equivariant complexes on algebraic spaces and Artin stacks similarly to rank_+ .

6.2. Indices for d-critical stacks. Let (\mathfrak{X}, s) be a d-critical Artin stack with \mathfrak{X} quasi-separated with affine stabilizers. The aim of this section is to introduce a locally constant function on $\text{Grad}(\mathfrak{X})$ called the index for \mathfrak{X} . We first deal with the case of algebraic space with a \mathbb{G}_m -action.

Definition 6.1. Let X be an algebraic space with \mathbb{G}_m -action μ . The *index function* $I_X^\mu: X^\mu \rightarrow \mathbb{Z}$ is given by

$$I_X^\mu(x) := \text{rank}_+(T_{X,x}) - \text{rank}_-(T_{X,x})$$

for a point $x \in X^\mu$.

Assume that X admits a \mathbb{G}_m -invariant étale d-critical structure $s \in \Gamma(X, \mathcal{S}_X^0)^{\mathbb{G}_m}$. Take a \mathbb{G}_m -equivariant étale d-critical chart $\mathfrak{R} = (R, \eta, U, f, i)$ of x such that $\dim T_{X,x} = \dim U$ holds. Then we have an equality $I_X^\mu(x) = \text{rank}(T_{U,x}^+) - \text{rank}(T_{U,x}^-)$. Using Theorem 3.31, we see that the same equality holds for arbitrary \mathbb{G}_m -equivariant étale d-critical chart of x . In particular, we have the following:

Proposition 6.2. *Let X be an algebraic space with \mathbb{G}_m -action μ , which admits a \mathbb{G}_m -invariant d-critical structure. Then the function*

$$I_X^\mu: X^\mu \longrightarrow \mathbb{Z}$$

is locally constant.

Remark 6.3. This proposition is false if we do not assume the existence of a \mathbb{G}_m -invariant d-critical structure. For example, consider the scheme $X = \{(x, y) \in \mathbb{A}^2 \mid xy = 0\}$ with a \mathbb{G}_m -action μ of weight $(1, 0)$. Then the function I_X^μ is not locally constant near the origin.

Assume that we are given algebraic spaces X_1 and X_2 with \mathbb{G}_m -actions μ_1 and μ_2 , \mathbb{G}_m -equivariant smooth morphism $h: X_1 \rightarrow X_2$. We define the locally constant function $I_h^{\mu_1}$ on $X_1^{\mu_1}$ by $I_h^{\mu_1} := \text{rank}_+(T_{X_1/X_2}) - \text{rank}_-(T_{X_1/X_2})$. Then we have the following identity of locally constant functions on $X_1^{\mu_1}$

$$(6.1) \quad I_{X_2}^{\mu_2} + I_h^{\mu_1} = I_{X_1}^{\mu_1}.$$

Let X be a quasi-separated algebraic space with a \mathbb{G}_m^2 -action $\hat{\mu} = (\mu_1, \mu_2)$. We say that a point $x \in X^{\hat{\mu}}$ is *matching* if its image in $[X^{\hat{\mu}}/\mathbb{G}_m^2]$ is a matching point (see §5.4 for the definition). We let $X_{\text{mat}}^{\hat{\mu}} \subset X^{\hat{\mu}}$ denote the open subspace of matching points. For an algebraic space Y over $X^{\hat{\mu}}$, we let Y_{mat} denote the base change to $X_{\text{mat}}^{\hat{\mu}}$. The following proposition is a consequence of Proposition 5.22:

Proposition 6.4. *Consider the following commutative diagram:*


$$\begin{array}{ccc} (X^{\mu_1,+})^{\mu_2} \times_{X^{\mu_2}} X^{\mu_2,+} & \xleftarrow{\alpha} & X^{\hat{\mu},+,+} & \xrightarrow{\beta} & (X^{\mu_2,+})^{\mu_1} \times_{X^{\mu_1}} X^{\mu_1,+} \\ & & \searrow & & \swarrow \\ & & X^{\hat{\mu}} & & \end{array}$$

Then the maps α and β are isomorphisms over the matching locus $X_{\text{mat}}^{\hat{\mu}}$.


The above proposition implies the following:

Corollary 6.5. *The composites of the following correspondences coincide:*

$$\begin{array}{ccccc} & & ((X^{\mu_1,+})^{\mu_2} \times_{X^{\mu_2}} X^{\mu_2,+})_{\text{mat}} & & \\ & \swarrow & & \searrow & \\ (X^{\mu_1,+})_{\text{mat}}^{\mu_2} & & & & X^{\mu_2,+} \\ \swarrow & & \searrow & \swarrow & \searrow \\ X_{\text{mat}}^{\hat{\mu}} & & X^{\mu_2} & & X, \end{array}$$



Further, these correspondences are naturally isomorphic to the following correspondence:



From the definition of matching points, we get the following identity of locally constant functions on $X^{\hat{\mu}}_{\text{mat}}$:

$$(6.2) \quad I_X^{\mu_1}|_{X^{\hat{\mu}}_{\text{mat}}} + (I_{X^{\mu_1}}^{\mu_2})|_{X^{\hat{\mu}}_{\text{mat}}} = I_X^{\mu_2}|_{X^{\hat{\mu}}_{\text{mat}}} + (I_{X^{\mu_2}}^{\mu_1})|_{X^{\hat{\mu}}_{\text{mat}}}.$$

We now introduce the index function for Artin stacks.

Definition 6.6. Let \mathfrak{X} be an Artin stack. The *index function* is the function $I_{\mathfrak{X}}: \text{Grad}(\mathfrak{X}) \rightarrow \mathbb{Z}$ given by

$$I_{\mathfrak{X}}(x) := (\text{rank}_+ H^0(\mathbb{T}_{\mathfrak{X},x}) - \text{rank}_- H^0(\mathbb{T}_{\mathfrak{X},x})) - (\text{rank}_+ H^{-1}(\mathbb{T}_{\mathfrak{X},x}) - \text{rank}_- H^{-1}(\mathbb{T}_{\mathfrak{X},x}))$$

for a point $x \in \text{Grad}(\mathfrak{X})$.

Assume that we are given a \mathbb{G}_m -equivariant chart (X, μ, \bar{q}) of \mathfrak{X} . Set

$$I_q^\mu := \text{rank}_+ T_{X/\bar{x}} - \text{rank}_- T_{X/\bar{x}}.$$

Then for a point $\hat{x} \in X^\mu$ with $q_\mu(\hat{x}) = x$, we have the following identity

$$(6.3) \quad I_{\mathfrak{X}}(x) = I_X^\mu(\hat{x}) - I_q^\mu(\hat{x}).$$

Assume that \mathfrak{X} admits a d-critical structure. Then the equality (6.3) together with Proposition 6.2 implies that the function $I_{\mathfrak{X}}$ is locally constant. Assume further that \mathfrak{X} is the underlying stack of a (-1) -shifted symplectic stack $\widehat{\mathfrak{X}}$. Since for each point $x \in \text{Grad}(\mathfrak{X})$ we have an equality

$$\text{rank}_+ H^2(\mathbb{T}_{\widehat{\mathfrak{X}},u(x)}) = \text{rank}_- H^{-1}(\mathbb{T}_{\widehat{\mathfrak{X}}}), \quad \text{rank}_+ H^1(\mathbb{T}_{\widehat{\mathfrak{X}},u(x)}) = \text{rank}_- H^0(\mathbb{T}_{\widehat{\mathfrak{X}},u(x)}),$$

we obtain an identity $I_{\mathfrak{X}} = \text{rank}_+ \mathbb{T}_{\widehat{\mathfrak{X}}}$. Since we have $(\mathbb{T}_{\text{Filt}(\widehat{\mathfrak{X}})/\text{Grad}(\widehat{\mathfrak{X}})})|_{\text{Grad}(\mathfrak{X})} \simeq \mathbb{T}_{\widehat{\mathfrak{X}}}^+$ and $\text{vd}(\text{Grad}(\widehat{\mathfrak{X}})) = 0$, we obtain

$$(6.4) \quad I_{\mathfrak{X}} = \text{vd Filt}(\widehat{\mathfrak{X}})|_{\text{Grad}(\mathfrak{X})}.$$

We set $I_{\mathfrak{X}}^1 := I_{\text{Grad}(\mathfrak{X})}: \text{Grad}^2(\mathfrak{X}) \rightarrow \mathbb{Z}$. We let $\sigma: \text{Grad}^2(\mathfrak{X}) \cong \text{Grad}^2(\mathfrak{X})$ denote the isomorphism induced by the swapping isomorphism $\mathbb{G}_m^2 \cong \mathbb{G}_m^2$ and set $I_{\mathfrak{X}}^2 := I_{\text{Grad}^2(\mathfrak{X})}^1 \circ \sigma: \text{Grad}^2(\mathfrak{X}) \rightarrow \mathbb{Z}$. By the definition of the matching locus, we have the following identity:

$$(6.5) \quad I_{\mathfrak{X}}^1|_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}} = I_{\mathfrak{X}}^2|_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}}.$$

6.3. Index line bundles for \mathbb{G}_m -equivariant d-critical spaces. Now we will introduce the index line bundle for a \mathbb{G}_m -equivariant d-critical algebraic spaces. Let X be a quasi-separated \mathbb{G}_m -equivariant d-critical algebraic space. For a \mathbb{G}_m -equivariant étale d-critical chart $\mathfrak{R} = (R, \eta, U, f, i)$ of (X, s) , define a graded line bundle on $R^{\mathbb{G}_m, \text{red}}$ by

$$\mathcal{L}_{\mu,s,\mathfrak{R}}^{\text{Ind}} := \widehat{\det}_+(\mathbb{L}_{\mathfrak{R}})$$

where $\mathbb{L}_{\mathfrak{R}}$ is the virtual cotangent complex defined in Definition 3.33. For each point $\hat{x} \in R^{\mathbb{G}_m}$ with $\eta(\hat{x}) = x$, we construct an isomorphism

$$\kappa_{\hat{x},\mathfrak{R}}^{\text{Ind}}: \mathcal{L}_{\mu,s,\mathfrak{R}}^{\text{Ind}}|_{\hat{x}} \cong \det_{\pm}(\Omega_{X,x})$$

by using the following fibre sequence

$$T_{X,x}^+[1] \rightarrow \mathbb{L}_{\mathfrak{R}}^+|_{\hat{x}} \rightarrow \Omega_{X,x}^+$$

and (2.5), (2.14). If we are given two \mathbb{G}_m -equivariant étale d-critical charts $\mathfrak{R}_1 = (R_1, \eta_1, U_1, f_1, i_1)$ and $\mathfrak{R}_2 = (R_2, \eta_2, U_2, f_2, i_2)$ and points $\hat{x}_1 \in R_1^{\mathbb{G}_m}$ and $\hat{x}_2 \in R_2^{\mathbb{G}_m}$ with $x = \eta_1(\hat{x}_1) = \eta_2(\hat{x}_2)$, we can construct the following isomorphism

$$\mathcal{L}_{\mu,s,\mathfrak{R}_1}^{\text{Ind}}|_{\hat{x}_1} \xrightarrow[\cong]{\kappa_{\hat{x}_1,\mathfrak{R}_1}^{\text{Ind}}} \widehat{\det}_{\pm}(\Omega_{X,x}) \xrightarrow[\cong]{\kappa_{\hat{x}_2,\mathfrak{R}_2}^{\text{Ind},-1}} \mathcal{L}_{\mu,s,\mathfrak{R}_2}^{\text{Ind}}|_{\hat{x}_2}.$$

We claim the composite algebraizes: namely, it induces an isomorphism of line bundles over $(R_1^{\mathbb{G}_m} \times_{X^\mu} R_2^{\mathbb{G}_m})^{\text{red}}$. To prove this, using Theorem 3.31, we may assume $R_1 = R_2$, $\eta_1 = \eta_2$, $U_2 = U_1 \times \mathbb{A}^n$ and $f_2 = f_1 \boxplus q$ where q is a \mathbb{G}_m -invariant quadratic function. Set $\mathbb{L}_{\mathfrak{R}_2/\mathfrak{R}_1}^+ := [T_{\mathbb{A}^n,0}^+ \xrightarrow{\text{Hess}_q^+} \Omega_{\mathbb{A}^n,0}^+]$. The null-homotopy $\mathbb{L}_{\mathfrak{R}_2/\mathfrak{R}_1}^+ \simeq 0$ induces a quasi-isomorphism

$$\mathbb{L}_{\mathfrak{R}_1}^+ \simeq \mathbb{L}_{\mathfrak{R}_2}^+$$

hence an isomorphism

$$(6.6) \quad \mathcal{L}_{\mu,s,\mathfrak{R}_1}^{\text{Ind}} \cong \mathcal{L}_{\mu,s,\mathfrak{R}_2}^{\text{Ind}}$$

which induces the desired isomorphism on each stalk.

By construction, the isomorphism $\mathcal{L}_{\mu,s,\mathfrak{R}_1}^{\text{Ind}}|_{(R_1^{\mathbb{G}_m} \times_{X^\mu} R_2^{\mathbb{G}_m})^{\text{red}}} \cong \mathcal{L}_{\mu,s,\mathfrak{R}_2}^{\text{Ind}}|_{(R_1^{\mathbb{G}_m} \times_{X^\mu} R_2^{\mathbb{G}_m})^{\text{red}}}$ satisfies the cocycle condition, since it does on each stalk. In particular, $\mathcal{L}_{\mu,s,\mathfrak{R}}^{\text{Ind}}$ glues to define a line bundle $\mathcal{L}_{\mu,s}^{\text{Ind}}$ on $X^{\mu,\text{red}}$ with a choice of an isomorphism

$$\iota_{\mathfrak{R}}^{\text{Ind}} : \mathcal{L}_{\mu,s}^{\text{Ind}}|_{R^{\mathbb{G}_m,\text{red}}} \cong \det_+(\mathbb{L}_{\mathfrak{R}})$$

for each \mathbb{G}_m -equivariant d-critical chart $\mathfrak{R} = (R, \eta, U, f, i)$. We call $\mathcal{L}_{\mu,s}^{\text{Ind}}$ the *index line bundle*.

We will study the behaviour of the index line bundle. The following proposition summarizes a list of properties of index line bundles:

Proposition 6.7. (i) *Let (X, μ, s) be a quasi-separated \mathbb{G}_m -equivariant d-critical algebraic space. Then for each point $x \in X^\mu$, there exists a natural isomorphism*

$$(6.7) \quad \kappa_x^{\text{Ind}} : \mathcal{L}_{\mu,s}^{\text{Ind}}|_x \cong \widehat{\det}_{\pm}(\Omega_{X,x}).$$

(ii) *Let (X, μ, s) be as above. Then there exists a natural isomorphism*

$$(6.8) \quad \xi_{\mu,s} : K_{X^\mu, s^\mu} \otimes \mathcal{L}_{\mu,s}^{\text{Ind}, \otimes 2} \cong K_{X,s}|_{X^{\mu,\text{red}}}.$$

(iii) *Let (X_1, μ_1, s_1) and (X_2, μ_2, s_2) be quasi-separated \mathbb{G}_m -equivariant d-critical algebraic spaces and $h : X_1 \rightarrow X_2$ be a smooth morphism compatible with \mathbb{G}_m -actions such that $h^*s_2 = s_1$ holds. Then there exists a natural isomorphism*

$$\Upsilon_h^{\text{Ind}} : h^{\mathbb{G}_m,\text{red},*} \mathfrak{L}_{\mu_2,s_2}^{\text{Ind}} \otimes K_{X_1/X_2}^{\pm} \cong \mathfrak{L}_{\mu_1,s_1}^{\text{Ind}}$$

where we set $K_{X_1/X_2}^{\pm} := \widehat{\det}_{\pm}(\Omega_{X_1/X_2})$.

(iv) *Assume that we are given a quasi-separated \mathbb{G}_m^2 -equivariant algebraic space $(X, \hat{\mu}(\mu_1, \mu_2), s)$. Then there exists an isomorphism*

$$\theta_{\hat{\mu},s} : \mathcal{L}_{\mu_1,s}^{\text{Ind}}|_{X^{\hat{\mu},\text{red}}} \otimes (\mathcal{L}_{\mu_2|_{X^{\mu_1},s^{\mu_1}}}^{\text{Ind}}) \cong \mathcal{L}_{\mu_2,s}^{\text{Ind}}|_{X^{\hat{\mu},\text{red}}} \otimes (\mathcal{L}_{\mu_1|_{X^{\mu_2},s^{\mu_2}}}^{\text{Ind}}).$$

(v) *Let (X, μ, s) and (Y, ν, t) be quasi-separated \mathbb{G}_m -equivariant d-critical algebraic spaces. Then there exists a natural isomorphism*

$$\Psi_{\mu,\nu}^{\text{Ind}} : \mathcal{L}_{\mu,s}^{\text{Ind}} \boxtimes \mathcal{L}_{\nu,t}^{\text{Ind}} \cong \mathcal{L}_{\mu \times \nu, s \boxplus t}^{\text{Ind}}.$$

(vi) *Let X be a quasi-separated derived algebraic space with a \mathbb{G}_m -action μ and a \mathbb{G}_m -invariant (-1) -shifted symplectic structure ω_X . Let (X^{cl}, s) be the underlying d-critical structure. Then there exists a natural isomorphism*

$$\Lambda_{\mu}^{\text{Ind}} : \widehat{\det}_+(\mathbb{L}_X) \cong \mathcal{L}_{\mu^{\text{cl}},s}^{\text{Ind}}.$$

Proof. For (i), the existence of natural isomorphisms κ_x^{Ind} is obvious from the construction. Note that for each \mathbb{G}_m -equivariant étale d-critical chart $\mathfrak{R} = (R, \eta, U, f, i)$ and a point $\hat{x} \in R^{\mathbb{G}_m}$ lifting x , the following diagram commutes:

$$(6.9) \quad \begin{array}{ccc} \mathcal{L}_{\mu,s}^{\text{Ind}}|_x & \xrightarrow[\cong]{\kappa_x^{\text{Ind}}} & \det_{\pm}(\Omega_{X,x}) \\ \cong \downarrow \iota_{\mathfrak{R}}|_{\hat{x}} & & \cong \downarrow (2.5), (2.14) \\ \det_+(\mathbb{L}_{\mathfrak{R}}|_{\hat{x}}) & \xrightarrow[\cong]{} & \det_+(T_{R,\hat{x}}[1]) \otimes \det_+(\Omega_{R,\hat{x}}). \end{array}$$

We now prove (ii). For each \mathbb{G}_m -equivariant étale d-critical chart $\mathfrak{R} = (R, \eta, U, f, i)$, there exists a natural equivalence of two-term complexes on $R^{\mathbb{G}_m}$

$$\mathbb{T}_{\mathfrak{R}}^+[1] \simeq \mathbb{L}_{\mathfrak{R}}^+$$

which is the identity map on each term. This equivalence together with the isomorphisms (2.5) and (2.14) induce an isomorphism

$$(6.10) \quad \widehat{\det}_-(\mathbb{L}_{\mathfrak{R}}) \cong \widehat{\det}_+(\mathbb{L}_{\mathfrak{R}}).$$

For each $\hat{x} \in R^{\mathbb{G}_m}$ with $\eta(\hat{x}) = x$, the following diagram commutes by (2.15):

$$(6.11) \quad \begin{array}{ccc} \det_-(\mathbb{L}_{\mathfrak{R}}|_{\hat{x}}) & \xrightarrow[\cong]{(6.10)} & \det_+(\mathbb{L}_{\mathfrak{R}}|_{\hat{x}}) \\ \cong \downarrow & & \cong \downarrow \\ \det_{\pm}(\Omega_{X,x}) & \xrightarrow[\cong]{(-1)^{\text{rank}_+ \Omega_{X,x}} \cdot \text{id}} & \det_{\pm}(\Omega_{X,x}). \end{array}$$

We define a map

$$\xi_{\mu,s,\mathfrak{R}}: K_{X^\mu, s^\mu}|_{R^{\mathbb{G}_m, \text{red}}} \otimes \mathcal{L}_{\mu,s}^{\text{Ind}, \otimes 2}|_{R^{\mathbb{G}_m, \text{red}}} \cong K_{X,s}|_{R^{\mathbb{G}_m, \text{red}}}$$

by the following composition

$$\begin{aligned} K_{X^\mu, s^\mu}|_{R^{\mathbb{G}_m, \text{red}}} \otimes \mathcal{L}_{\mu,s}^{\text{Ind}, \otimes 2}|_{R^{\mathbb{G}_m, \text{red}}} &\xrightarrow[\cong]{(\iota_{\mathfrak{R}^{\mathbb{G}_m}}, \iota_{\mathfrak{R}}^{\text{Ind}, \otimes 2})} \widehat{\det}_0(\mathbb{L}_{\mathfrak{R}}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{R}})^{\otimes 2} \\ &\xrightarrow[\cong]{\text{id} \otimes \text{id} \otimes (6.10)} \widehat{\det}_0(\mathbb{L}_{\mathfrak{R}}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{R}}) \otimes \widehat{\det}_-(\mathbb{L}_{\mathfrak{R}}) \\ &\cong \widehat{\det}(\mathbb{L}_{\mathfrak{R}}|_{R^{\mathbb{G}_m, \text{red}}}) \xrightarrow[\cong]{\iota_{\mathfrak{R}}^{-1}} K_{X,s}|_{R^{\mathbb{G}_m, \text{red}}}. \end{aligned}$$

Note by construction and commutativity of the diagram (6.11) that the following diagram commutes:

$$\begin{array}{ccc} K_{X^\mu, s^\mu}|_x \otimes \mathcal{L}_{\mu,s}^{\text{Ind}, \otimes 2}|_x & \xrightarrow[\cong]{(-1)^{\text{rank}_+ \Omega_{X,x}} \cdot \xi_{\mu,s,\mathfrak{R}}|_x} & K_{X,s}|_x \\ \cong \downarrow (\kappa_x, \kappa_x^{\text{Ind}}) & & \downarrow \kappa_x \cong \\ \det(\Omega_{X^\mu, x})^{\otimes 2} \otimes \det_{\pm}(\Omega_{X,x})^{\otimes 2} & & \det(\Omega_{X,x})^{\otimes 2} \\ \cong \downarrow & & \downarrow \cong \\ \det_{\leq 0}(\Omega_{X,x}) \otimes \det_{\geq 0}(\Omega_{X,x}) \otimes \det_{\pm}(\Omega_{X,x}) & \xrightarrow[\cong]{} & \det(\Omega_{X,x})^{\otimes 2}. \end{array}$$

Since the symmetrizer of the complex of $\det_+(\Omega_{X,x})^{\otimes 2}$ is given by the multiplication of $(-1)^{\text{rank}_+ \Omega_{X,x}}$, by combining the above diagram with the diagram (6.11), we conclude that the following diagram commutes:

$$(6.12) \quad \begin{array}{ccc} K_{X^\mu, s^\mu}|_x \otimes \mathcal{L}_{\mu,s}^{\text{Ind}, \otimes 2}|_x & \xrightarrow[\cong]{\xi_{\mu,s,\mathfrak{R}}|_x} & K_{X,s}|_x \\ \cong \downarrow (\kappa_x, \kappa_x^{\text{Ind}}) & & \downarrow \kappa_x \cong \\ \det_0(\Omega_{X,x})^{\otimes 2} \otimes \det_{\pm}(\Omega_{X,x})^{\otimes 2} & \xrightarrow[\cong]{} & \det(\Omega_{X,x})^{\otimes 2} \end{array}$$

where the bottom map is the square of the canonical isomorphism $\det_0(\Omega_{X,x}) \otimes \det_{\pm}(\Omega_{X,x}) \cong \det(\Omega_{X,x})$. In particular, $\xi_{\mu,s,\mathfrak{R}}|_x$ does not depend on the choice of \mathfrak{R} and hence $\xi_{\mu,s,\mathfrak{R}}$ glues to define $\xi_{\mu,s}$ as desired.

We now prove (iii). Assume that we are given \mathbb{G}_m -equivariant étale d-critical charts $\mathfrak{R}_j = (R_j, \eta_j, U_j, f_j, i_j)$ for (X_j, s_j) for $j = 1, 2$ and a morphism $(H_0, H): \mathfrak{R}_1 \rightarrow \mathfrak{R}_2$ as in Proposition 3.32. We define a complex on $R_1^{\mathbb{G}_m}$ concentrated in degree $[-1, 0]$ by

$$\mathbb{L}_{\mathfrak{R}_1, H}^+ := [H^*T_{U_2}|_{R_1}^+ \rightarrow \Omega_{U_1}|_{R_1}^+]$$

where the map is given by the composition

$$H^*T_{U_2}|_{R_1}^+ \xrightarrow{\text{Hess } f_2} H^*\Omega_{U_2}|_{R_1}^+ \rightarrow \Omega_{U_1}|_{R_1}^+.$$

Then we have the following fibre sequences

$$(6.13) \quad H_0^{\mathbb{G}_m, *} \mathbb{L}_{\mathfrak{R}_2}^+ \rightarrow \mathbb{L}_{\mathfrak{R}_1, H}^+ \rightarrow \mathbb{L}_H|_{R_1}^+$$

$$(6.14) \quad \mathbb{T}_H[1]|_{R_1}^+ \rightarrow \mathbb{L}_{\mathfrak{R}_1}^+ \rightarrow \mathbb{L}_{\mathfrak{R}_1, H}^+.$$

We construct an isomorphism

$$\Upsilon_H^{\text{Ind}}: h^{\mathbb{G}_m, \text{red}, *} \mathfrak{L}_{\mu_2, s_2}^{\text{Ind}}|_{R_1^{\mathbb{G}_m, \text{red}}} \otimes K_{X_1/X_2}^{\pm}|_{R_1^{\mathbb{G}_m, \text{red}}} \cong \mathfrak{L}_{\mu_1, s_1}^{\text{Ind}}|_{R_1^{\mathbb{G}_m, \text{red}}}$$

so that the following diagram commutes:

$$(6.15) \quad \begin{array}{ccc} h^{\mathbb{G}_m, \text{red}, *} \mathfrak{L}_{\mu_2, s_2}^{\text{Ind}}|_{R_1^{\mathbb{G}_m, \text{red}}} \otimes K_{X_1/X_2}^{\pm}|_{R_1^{\mathbb{G}_m, \text{red}}} & \xrightarrow[\cong]{\Upsilon_H^{\text{Ind}}} & \mathfrak{L}_{\mu_1, s_1}^{\text{Ind}}|_{R_1^{\mathbb{G}_m, \text{red}}} \\ \cong \downarrow \iota_{\mathfrak{R}_2}^{\text{Ind}} & & \cong \downarrow \iota_{\mathfrak{R}_1}^{\text{Ind}} \\ H_0^{\mathbb{G}_m, \text{red}, *} \widehat{\det}_+(\mathbb{L}_{\mathfrak{R}_2}) \otimes K_{U_1/U_2}^{\pm}|_{R_1^{\mathbb{G}_m}} & \xrightarrow[\cong]{(2.5), (2.14)} & H_0^{\mathbb{G}_m, \text{red}, *} \widehat{\det}_+(\mathbb{L}_{\mathfrak{R}_2}) \otimes \widehat{\det}_+(\mathbb{L}_H|_{R_1}) \otimes \widehat{\det}_+(\mathbb{T}_H[1]|_{R_1}) \end{array}$$

where the lower right vertical arrow is constructed using (2.3) for fibre sequences (6.13) and (6.14). Take a point $\hat{x}_1 \in R_1^{\mathbb{G}_m}$ and set $\eta_1(\hat{x}_1) = \hat{x}_2$, $H_0(\hat{x}_1) = x_1$ and $\eta_2(\hat{x}_2) = x_2$. We claim that the isomorphism Υ_H^{Ind} is compatible with $\kappa_{x_1}^{\text{Ind}}$ and $\kappa_{x_2}^{\text{Ind}}$: i.e., the following diagram commutes:

$$(6.16) \quad \begin{array}{ccc} \mathfrak{L}_{\mu_2, s_2}^{\text{Ind}}|_{x_2} \otimes K_{X_1/X_2}^{\pm}|_{x_1} & \xrightarrow[\cong]{\Upsilon_H^{\text{Ind}}|_{x_1}} & \mathfrak{L}_{\mu_1, s_1}^{\text{Ind}}|_{x_1} \\ \cong \downarrow \kappa_{x_2}^{\text{Ind}} \otimes \text{id} & & \cong \downarrow \kappa_{x_1}^{\text{Ind}} \\ \det_{\pm}(\Omega_{X_2, x_2}) \otimes K_{X_1/X_2}^{\pm}|_{x_1} & \xrightarrow[\cong]{} & \det_{\pm}(\Omega_{X_1, x_1}). \end{array}$$

To see this, consider the following fibre double sequences:

$$\begin{array}{ccccc} T_{X_2, x_2}^+[1] \longleftarrow T_{X_2, x_2}^+[1] & \longrightarrow & 0 & & \mathbb{T}_h[1]|_{x_1} \longrightarrow T_{X_1, x_1}^+[1] \longrightarrow T_{X_2, x_2}^+[1] \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{L}_{\mathfrak{R}_2}|_{\hat{x}_2}^+ \longrightarrow \mathbb{L}_{\mathfrak{R}_1, H}|_{\hat{x}_1}^+ & \longrightarrow & \mathbb{L}_{H_0}|_{\hat{x}_1}^+ & & \mathbb{T}_H[1]|_{\hat{x}_1}^+ \longrightarrow \mathbb{L}_{\mathfrak{R}_1}^+|_{\hat{x}_1} \longrightarrow \mathbb{L}_{\mathfrak{R}_1, H}^+|_{\hat{x}_1} \\ \downarrow & & \downarrow & & \downarrow \\ \Omega_{X_2, x_2}^+ \longrightarrow \Omega_{X_1, x_1}^+ & \longrightarrow & \mathbb{L}_h|_{x_1}^+ & & 0 \longrightarrow \Omega_{X_1, x_1}^+ \longleftarrow \Omega_{X_1, x_1}^+ \end{array}$$

Using Lemma 2.4, we see that the following diagrams commute:

$$\begin{array}{ccc}
\det_+(\mathbb{L}_{\mathfrak{R}_1, H_1} |_{\hat{x}_1}) & \xrightarrow{\cong} & \det_+(\mathbb{L}_{\mathfrak{R}_2} |_{\hat{x}_2}) \otimes \det_+(\mathbb{L}_{H_0} |_{\hat{x}_1}) \\
\downarrow \cong & & \downarrow \cong \\
\det_+(T_{X_2, x_2}[1]) \otimes \det_+(\Omega_{X_1, x_1}) & \xrightarrow{\cong} & \det_+(T_{X_2, x_2}[1]) \otimes \det_+(\Omega_{X_2, x_2}) \otimes \det_+(\mathbb{L}_h |_{x_1}),
\end{array}$$

$$\begin{array}{ccc}
\det_+(\mathbb{L}_{\mathfrak{R}_1} |_{\hat{x}_1}) & \xrightarrow{\cong} & \det_+(\mathbb{T}_{H_0}[1] |_{\hat{x}_1}) \otimes \det_+(\mathbb{L}_{\mathfrak{R}_1, H} |_{\hat{x}_1}) \\
\downarrow \cong & & \downarrow \cong \\
\det_+(T_{X_1, x_1}[1]) \otimes \det_+(\Omega_{X_1, x_1}) & \xrightarrow{\cong} & \widehat{\det}_+(\mathbb{T}_h[1] |_{x_1}) \otimes \det_+(T_{X_2, x_2}[1]) \otimes \widehat{\det}_+(\Omega_{X_1, x_1}).
\end{array}$$

By combining the commutativity of these diagrams and the commutativity of (6.9), we conclude the commutativity of the diagram (6.16). In particular, the isomorphism $\Upsilon_H^{\text{Ind}} |_{\hat{x}_1}$ only depends on the point $x_1 \in X_1$. Therefore, Υ_H^{Ind} glues to define the isomorphism Υ_h^{Ind} .

We now prove (iv). Take a \mathbb{G}_m^2 -equivariant étale d-critical chart $\mathfrak{R} = (R, \eta, U, f, i)$. We will construct an isomorphism $\theta_{\hat{\mu}, s, \mathfrak{R}}$ over $R^{\hat{\mu}, \text{red}}$ and show that it does not depend on the choice of \mathfrak{R} , which implies the statement (iv). For a subset $\Sigma \subset \mathbb{Z}^2$, the natural equivalence $\mathbb{T}_{\mathfrak{R}}^{\Sigma}[1] \simeq \mathbb{L}_{\mathfrak{R}}^{\Sigma}$ with the isomorphisms (2.5) and (2.14) implies an isomorphism

$$(6.17) \quad \widehat{\det}_{-\Sigma}(\mathbb{L}_{\mathfrak{R}}) \simeq \widehat{\det}_{\Sigma}(\mathbb{L}_{\mathfrak{R}}).$$

Note that we have natural equivalences

$$\begin{aligned}
\mathcal{L}_{\mu_1, s}^{\text{Ind}} |_{R^{\hat{\mu}, \text{red}}} &\cong \widehat{\det}_{\mathbb{Z}_{>0} \times \mathbb{Z}}(\mathbb{L}_{\mathfrak{R}}), & \mathcal{L}_{\mu_2, s}^{\text{Ind}} |_{R^{\hat{\mu}, \text{red}}} &\cong \widehat{\det}_{\mathbb{Z} \times \mathbb{Z}_{>0}}(\mathbb{L}_{\mathfrak{R}}), \\
(\mathcal{L}_{\mu_1 |_{X^{\mu_2}}, s^{\mu_2}}^{\text{Ind}}) |_{R^{\hat{\mu}, \text{red}}} &\cong \widehat{\det}_{\mathbb{Z}_{>0} \times \{0\}}(\mathbb{L}_{\mathfrak{R}}), & (\mathcal{L}_{\mu_2 |_{X^{\mu_1}}, s^{\mu_1}}^{\text{Ind}}) |_{R^{\hat{\mu}, \text{red}}} &\cong \widehat{\det}_{\{0\} \times \mathbb{Z}_{>0}}(\mathbb{L}_{\mathfrak{R}}).
\end{aligned}$$

Set $S_1 := \{(m, l) \mid (m, l) \neq (0, 0), m \geq 0, l \geq 0\}$ and $S_2 := \mathbb{Z}_{>0} \times \mathbb{Z}_{<0}$. Then we have the following natural decompositions

$$\begin{aligned}
(\mathbb{Z}_{>0} \times \mathbb{Z}) \coprod (\{0\} \times \mathbb{Z}_{>0}) &= S_1 \coprod S_2 \\
(\mathbb{Z} \times \mathbb{Z}_{>0}) \coprod (\mathbb{Z}_{>0} \times \{0\}) &= S_2 \coprod (-S_2).
\end{aligned}$$

Then by using (6.17), we obtain the following isomorphism

$$\theta_{\hat{\mu}, s, \mathfrak{R}} : \mathcal{L}_{\mu_1, s}^{\text{Ind}} |_{R^{\hat{\mu}, \text{red}}} \otimes (\mathcal{L}_{\mu_2 |_{X^{\mu_1}}, s^{\mu_1}}^{\text{Ind}}) |_{R^{\hat{\mu}, \text{red}}} \cong \mathcal{L}_{\mu_2, s}^{\text{Ind}} |_{R^{\hat{\mu}, \text{red}}} \otimes (\mathcal{L}_{\mu_1 |_{X^{\mu_2}}, s^{\mu_2}}^{\text{Ind}}) |_{R^{\hat{\mu}, \text{red}}}.$$

Take a point $\hat{x} \in R^{\hat{\mu}}$ and set $x := \eta(\hat{x})$. We claim that $\theta_{\hat{\mu}, s, \mathfrak{R}}$ is compatible with κ_x^{Ind} , namely, the following diagram commutes:

$$(6.18) \quad \begin{array}{ccc}
\mathcal{L}_{\mu_1, s}^{\text{Ind}} |_{\hat{x}} \otimes \left(\mathcal{L}_{\mu_2 |_{X^{\mu_1}}, s^{\mu_1}}^{\text{Ind}} \right) |_{\hat{x}} & \xrightarrow[\cong]{(-1)^{\text{rank } S_2} \Omega_{X, x} \cdot \theta_{\hat{\mu}, s, \mathfrak{R}} |_{\hat{x}}} & \mathcal{L}_{\mu_2, s}^{\text{Ind}} |_{\hat{x}} \otimes \left(\mathcal{L}_{\mu_1 |_{X^{\mu_2}}, s^{\mu_2}}^{\text{Ind}} \right) |_{\hat{x}} \\
\downarrow \cong (\kappa_{x, \mu_1}^{\text{Ind}}, \kappa_{x, \mu_2}^{\text{Ind}} |_{X^{\mu_1}}) & & \downarrow \cong (\kappa_{x, \mu_2}^{\text{Ind}}, \kappa_{x, \mu_1}^{\text{Ind}} |_{X^{\mu_2}}) \\
\det_{\mu_1 \pm}(\Omega_{X, x}) \otimes \det_{\mu_2 \pm}(\Omega_{X^{\mu_1}, x}) & & \det_{\mu_2 \pm}(\Omega_{X, x}) \otimes \det_{\mu_1 \pm}(\Omega_{X^{\mu_2}, x}) \\
& \searrow \cong & \swarrow \cong \\
& \det_{\mathbb{Z}^2 \setminus \{(0,0)\}}(\Omega_{X, x}) &
\end{array}$$

Here we use the notation $\kappa_{x,\mu_1}^{\text{Ind}}$ and $\kappa_{x,\mu_2}^{\text{Ind}}$ to clarify which action we consider and similarly for $\det_{\mu_1\pm}$ and $\det_{\mu_2\pm}$. Firstly, note that we have the following natural equivalence of fibre sequences

$$(6.19) \quad \begin{array}{ccccc} (\Omega_{X,x}^{-S_2})^\vee[1] & \longrightarrow & (\mathbb{L}_{\mathfrak{R}}|_{\hat{x}}^{-S_2})^\vee[1] & \longrightarrow & (T_{X,x}^{-S_2}[1])^\vee[1] \\ \cong \downarrow & & \cong \downarrow & & \cong \downarrow \\ T_{X,x}^{S_2}[1] & \longrightarrow & \mathbb{L}_{\mathfrak{R}}|_{\hat{x}}^{S_2} & \longrightarrow & \Omega_{X,x}^{S_2}. \end{array}$$

This combined with Corollary 2.5, (2.15) and (2.16) shows that the following diagram commutes up to the sign $(-1)^{\text{rank}_{S_2} \Omega_{X,x}}$:

$$\begin{array}{ccc} \det_{-S_2}(\mathbb{L}_{\mathfrak{R}}|_{\hat{x}}) & \xrightarrow[\cong]{(6.17)} & \det_{S_2}(\mathbb{L}_{\mathfrak{R}}|_{\hat{x}}) \\ \cong \downarrow & & \cong \downarrow \\ \det_{-S_2}(T_{X,x_2}[1]) \otimes \det_{-S_2}(\Omega_{X,x}) & & \det_{S_2}(T_{X,x_2}[1]) \otimes \det_{S_2}(\Omega_{X,x}) \\ \cong \downarrow & & \cong \downarrow \\ \det_{\pm S_2}(\Omega_{X,x}) & \xlongequal{\quad\quad\quad} & \det_{\pm S_2}(\Omega_{X,x}). \end{array}$$

This implies the commutativity of the diagram (6.18)

We now prove (v). Let $\mathfrak{R} = (R, \eta, U, f, i)$ and $\mathfrak{S} = (S, \gamma, V, g, j)$ be \mathbb{G}_m -equivariant étale d-critical charts of X and Y , respectively. We construct an isomorphism

$$\Psi_{\mu,\nu,\mathfrak{R},\mathfrak{S}} : \mathcal{L}_{\mu,s}^{\text{Ind}}|_{R^{\mathbb{G}_m,\text{red}}} \boxtimes \mathcal{L}_{\nu,t}^{\text{Ind}}|_{S^{\mathbb{G}_m,\text{red}}} \cong \mathcal{L}_{\mu \times \nu, s \boxplus t}^{\text{Ind}}|_{R^{\mathbb{G}_m,\text{red}} \times S^{\mathbb{G}_m,\text{red}}}$$

by using the natural equivalence $\mathbb{L}_{\mathfrak{R}}|_{R^{\mathbb{G}_m}}^+ \boxplus \mathbb{L}_{\mathfrak{S}}|_{S^{\mathbb{G}_m}}^+ \simeq \mathbb{L}_{\mathfrak{R} \times \mathfrak{S}}|_{R^{\mathbb{G}_m} \times S^{\mathbb{G}_m}}^+$. Take a point $\hat{x} \in R$ and $\hat{y} \in S$ with $x = \eta(\hat{x})$ and $y = \gamma(\hat{y})$. Then it follows from the construction that the following diagram commutes:

$$(6.20) \quad \begin{array}{ccc} \mathcal{L}_{\mu,s}^{\text{Ind}}|_{\hat{x}} \otimes \mathcal{L}_{\nu,t}^{\text{Ind}}|_{\hat{y}} & \xrightarrow[\cong]{\Psi_{\mu,\nu,\mathfrak{R},\mathfrak{S}}|_{(\hat{x},\hat{y})}} & \mathcal{L}_{\mu \times \nu, s \boxplus t}^{\text{Ind}}|_{(\hat{x},\hat{y})} \\ \cong \downarrow \kappa_x^{\text{Ind}} \otimes \kappa_y^{\text{Ind}} & & \cong \downarrow \kappa_{(x,y)}^{\text{Ind}} \\ \det_{\pm}(\Omega_{X,x}) \otimes \det_{\pm}(\Omega_{Y,y}) & \xrightarrow[\cong]{} & \det_{\pm}(\Omega_{X \times Y, (x,y)}). \end{array}$$

In particular, $\Psi_{\mu,\nu,\mathfrak{R},\mathfrak{S}}|_{(\hat{x},\hat{y})}$ does not depend on the choice of étale d-critical charts hence glues to define an isomorphism $\Psi_{\mu,\nu} : \mathcal{L}_{\mu,s}^{\text{Ind}} \boxtimes \mathcal{L}_{\nu,t}^{\text{Ind}} \cong \mathcal{L}_{\mu \times \nu, s \boxplus t}^{\text{Ind}}$.

We now prove (vi). Take a point $x \in X^{\mathbb{G}_m}$. By using Proposition 3.13, we can find a tuple $\mathfrak{R} = (R, \eta, U, f, i)$ where R is a derived scheme with a \mathbb{G}_m -action, η is a \mathbb{G}_m -equivariant étale morphism, U is a smooth scheme with a \mathbb{G}_m -action whose image contains x , f is a \mathbb{G}_m -invariant function on U with the property $f|_{\text{Crit}(f)^{\text{red}}} = 0$ and a \mathbb{G}_m -equivariant locally closed embedding $i: R \hookrightarrow U$ with $i(R) = \text{Crit}(f)$. Then $\mathfrak{R}^{\text{cl}} = (R^{\text{cl}}, \eta^{\text{cl}}, U, f, i^{\text{cl}})$ defines a \mathbb{G}_m -equivariant étale d-critical chart for (X, s) and we have a natural equivalence

$$\mathbb{L}_R^+ \simeq \mathbb{L}_{\mathfrak{R}^{\text{cl}}}^+.$$

This equivalence induces an isomorphism

$$\Lambda_{\mu,\mathfrak{R}}^{\text{Ind}} : \widehat{\det}_+(\mathbb{L}_X|_R) \cong \widehat{\det}_+(\mathbb{L}_R) \cong \widehat{\det}_+(\mathbb{L}_{\mathfrak{R}^{\text{cl}}}) \xrightarrow[\cong]{\iota_{\mathfrak{R}^{\text{cl}}}^{\text{Ind},-1}} \mathcal{L}_{\mu^{\text{cl}},s}^{\text{Ind}}|_{R^{\mathbb{G}_m,\text{red}}}.$$

By construction, this isomorphism is compatible with κ_x^{Ind} , i.e., the following diagram commutes for each lift $\hat{x} \in R^{\mathbb{G}_m}$ of x :

$$(6.21) \quad \begin{array}{ccc} \det_+(\mathbb{L}_{X,x}) & \xrightarrow[\cong]{\Lambda_{\mu,\mathfrak{R}}^{\text{Ind}}} & \mathcal{L}_{\mu^{\text{cl}},s}^{\text{Ind}}|_x \\ \cong \downarrow & & \cong \downarrow \kappa_x^{\text{Ind}} \\ \det_+(\tau^{\leq -1}(\mathbb{L}_{X,x})) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}_{X,x})) & \xrightarrow[\cong]{} & \det_{\pm}(\Omega_{X,x}) \end{array}$$

where the bottom arrow is induced by the natural equivalence $\omega_X|_x : T_{X,x}^+[1] \simeq \tau^{\leq -1}(\mathbb{L}_X|_x^+)$, (2.5) and (2.14). In particular, the morphism $\Lambda_{\mu, \mathfrak{R}}^{\text{Ind}}|_x$ does not depend on the choice of \mathfrak{R} and the lift \hat{x} of x . Therefore, the isomorphism $\Lambda_{\mu, \mathfrak{R}}^{\text{Ind}}$ glues to define an isomorphism Λ_μ^{Ind} . \square

By construction, the maps $\xi_{\mu,s}$, Υ_h^{Ind} , θ_μ and Λ_μ^{Ind} are compatible with κ_x^{Ind} , so that the following diagrams commute (we adopt the notation from the statement and the proof of Proposition 6.7):

$$(6.22) \quad \begin{array}{ccc} K_{X^\mu, s^\mu}|_x \otimes \mathcal{L}_{\mu, s}^{\text{Ind}, \otimes 2}|_x & \xrightarrow[\cong]{\xi_{\mu, s}|_x} & K_{X, s}|_x \\ \cong \downarrow (\kappa_x, \kappa_x^{\text{Ind}}) & & \kappa_x \downarrow \cong \\ \det(\Omega_{X^\mu, x})^{\otimes 2} \otimes \det_{\pm}(\Omega_{X, x})^{\otimes 2} & \xrightarrow[\cong]{} & \det(\Omega_{X, x})^{\otimes 2}, \end{array}$$

$$(6.23) \quad \begin{array}{ccc} \mathcal{L}_{\mu_2, s_2}^{\text{Ind}}|_{x_2} \otimes K_{X_1/X_2}^{\pm}|_{x_1} & \xrightarrow[\cong]{\Upsilon_h^{\text{Ind}}|_{x_1}} & \mathcal{L}_{\mu_1, s_1}^{\text{Ind}}|_{x_1} \\ \cong \downarrow \kappa_{x_2}^{\text{Ind}} \otimes \text{id} & & \cong \downarrow \kappa_{x_1}^{\text{Ind}} \\ \det_{\pm}(\Omega_{X_2, x_2}) \otimes K_{X_1/X_2}^{\pm}|_{x_1} & \xrightarrow[\cong]{} & \det_{\pm}(\Omega_{X_1, x_1}), \end{array}$$

(6.24)

$$\begin{array}{ccc} \mathcal{L}_{\mu_1, s}^{\text{Ind}}|_x \otimes \left(\mathcal{L}_{\mu_2|_{X^{\mu_1}}, s^{\mu_1}}^{\text{Ind}} \right)|_x & \xrightarrow[\cong]{(-1)^{\text{rank } \Omega_{X^{\mu_2}, x}^{S_2}} \cdot \theta_\mu|_x} & \mathcal{L}_{\mu_2, s}^{\text{Ind}}|_x \otimes \left(\mathcal{L}_{\mu_1|_{X^{\mu_2}}, s^{\mu_2}}^{\text{Ind}} \right)|_x \\ \cong \downarrow (\kappa_{x, \mu_1}^{\text{Ind}}, \kappa_{x, \mu_2|_{X^{\mu_1}}}^{\text{Ind}}) & & \cong \downarrow (\kappa_{x, \mu_2}^{\text{Ind}}, \kappa_{x, \mu_1|_{X^{\mu_2}}}^{\text{Ind}}) \\ \det_{\mu_1 \pm}(\Omega_{X, x}) \otimes \det_{\mu_2 \pm}(\Omega_{X^{\mu_1}, x}) & & \det_{\mu_2 \pm}(\Omega_{X, x}) \otimes \det_{\mu_1 \pm}(\Omega_{X^{\mu_2}, x}) \\ & \searrow \cong \quad \swarrow \cong & \\ & \det_{\mathbb{Z}^2 \setminus \{(0,0)\}}(\Omega_{X, x}). & \end{array}$$

$$(6.25) \quad \begin{array}{ccc} \mathcal{L}_{\mu, s}^{\text{Ind}}|_x \otimes \mathcal{L}_{\nu, t}^{\text{Ind}}|_y & \xrightarrow[\cong]{\Psi_{\mu, \nu}|_{(x, y)}} & \mathcal{L}_{\mu \times \nu, s \boxplus t}^{\text{Ind}}|_{(x, y)} \\ \cong \downarrow \kappa_x^{\text{Ind}} \otimes \kappa_y^{\text{Ind}} & & \cong \downarrow \kappa_{(x, y)}^{\text{Ind}} \\ \det_{\pm}(\Omega_{X, x}) \otimes \det_{\pm}(\Omega_{Y, y}) & \xrightarrow[\cong]{} & \det_{\pm}(\Omega_{X \times Y, (x, y)}). \end{array}$$

$$(6.26) \quad \begin{array}{ccc} \det_+(\mathbb{L}_{X, x}) & \xrightarrow[\cong]{\Lambda_\mu^{\text{Ind}}} & \mathcal{L}_{\mu^{\text{cl}}, s}^{\text{Ind}}|_x \\ \cong \downarrow & & \cong \downarrow \kappa_x^{\text{Ind}} \\ \det_+(\tau^{\leq -1}(\mathbb{L}_{X, x})) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}_{X, x})) & \xrightarrow[\cong]{} & \det_{\pm}(\Omega_{X, x}). \end{array}$$

The commutativity of the first diagram follows from the commutativity of (6.12), the second from (6.16), the third from (6.18), the fourth from (6.25) and the fifth from (6.21). Note that the diagram (6.24) commutes over the matching locus. We will prove that the isomorphisms constructed in Proposition 6.7 are compatible with each other (we only state and prove those compatibilities that we use later). Firstly, it is clear from the commutativity of the above diagrams that $\xi_{\mu,s}$, Υ_h^{Ind} , $\theta_{\hat{\mu},s}$ and Λ_μ are closed under products under the identification $\Psi_{\mu, \nu}^{\text{Ind}}$. We now claim the compatibility of $\xi_{\mu,s}$ and Υ_h . We adopt the notation from (iii) of

Proposition 6.7. Then the following diagram commutes:

$$(6.27) \quad \begin{array}{ccc} h^{\mathbb{G}_m, \text{red}, *} K_{X_2^{\mu_2, s_2}} \otimes h^{\mathbb{G}_m, \text{red}, *} \mathfrak{L}_{\mu_2, s_2}^{\text{Ind}, \otimes 2} & \xrightarrow[\cong]{\Upsilon_h^{\text{Ind}}} & h^{\mathbb{G}_m, \text{red}, *} K_{X_2^{\mu_2, s_2}} \otimes \mathfrak{L}_{\mu_1, s_1}^{\text{Ind}, \otimes 2} \\ \otimes K_{X_1/X_2}^{\otimes 2} |_{X_1^{\mu_1, \text{red}}} & & \otimes K_{X_1^{\mu_1}/X_2^{\mu_2}} |_{X_1^{\mu_1, \text{red}}} \\ \downarrow \cong \xi_{\mu_2, s_2} & & \downarrow \cong \Upsilon_{h^{\mathbb{G}_m}} \\ h^{\mathbb{G}_m, \text{red}, *} K_{X_2, s_2} |_{X_2^{\mu_2, \text{red}}} \otimes K_{X_1/X_2}^{\otimes 2} |_{X_1^{\mu_1, \text{red}}} & \xrightarrow[\cong]{\Upsilon_h} & K_{X_1, s_1} |_{X_1^{\mu_1, \text{red}}}. \end{array}$$

This is a direct consequence of the commutativity of the diagram (6.22) and (6.23). Next, we claim the compatibility of $\xi_{\mu, s}$ and $\theta_{\hat{\mu}, s}$. We adopt the notation from Proposition 6.7 (iv):

$$(6.28) \quad \begin{array}{ccc} K_{X^{\hat{\mu}, s^{\hat{\mu}}}} |_{X^{\hat{\mu}, \text{red}}} \otimes \mathcal{L}_{\mu_1, s}^{\text{Ind}, \otimes 2} |_{X^{\hat{\mu}, \text{red}}} & \xrightarrow[\cong]{\text{id} \otimes \theta_{\hat{\mu}, s}^{\otimes 2}} & K_{X^{\hat{\mu}, s^{\hat{\mu}}}} |_{X^{\hat{\mu}, \text{red}}} \otimes \mathcal{L}_{\mu_2, s}^{\text{Ind}, \otimes 2} |_{X^{\hat{\mu}, \text{red}}} \\ \otimes \mathcal{L}_{\mu_2 |_{X^{\mu_1}, s^{\mu_1}}}^{\text{Ind}, \otimes 2} & & \otimes \mathcal{L}_{\mu_1 |_{X^{\mu_2}, s^{\mu_2}}}^{\text{Ind}, \otimes 2} \\ \downarrow \cong \xi_{\mu_2 |_{X^{\mu_1}, s^{\mu_1}}} & & \downarrow \cong \xi_{\mu_1 |_{X^{\mu_2}, s^{\mu_2}}} \\ K_{X^{\mu_1, s^{\mu_1}}} |_{X^{\hat{\mu}, \text{red}}} \otimes \mathcal{L}_{\mu_1, s}^{\text{Ind}, \otimes 2} |_{X^{\hat{\mu}, \text{red}}} & & K_{X^{\mu_2, s^{\mu_2}}} |_{X^{\hat{\mu}, \text{red}}} \otimes \mathcal{L}_{\mu_2, s}^{\text{Ind}, \otimes 2} |_{X^{\hat{\mu}, \text{red}}} \\ \searrow \cong \xi_{\mu_1, s} & & \swarrow \cong \xi_{\mu_2, s} \\ & K_{X, s} |_{X^{\hat{\mu}, \text{red}}}. & \end{array}$$

This is a direct consequence of the commutativity of the diagram (6.22) and (6.24). We now prove the associativity of the isomorphism Υ_h^{Ind} . Let X_0, X_1 and X_2 be algebraic spaces with \mathbb{G}_m -action μ_0, μ_1 and μ_2 respectively. Assume that we are given \mathbb{G}_m -equivariant smooth morphisms $g: X_0 \rightarrow X_1$ and $h: X_1 \rightarrow X_2$ and take a d-critical structure s_2 of X_2 . We set $s_1 := h^*s_2$ and $s_0 := g^*s_1$. Then the following diagram commutes:

$$(6.29) \quad \begin{array}{ccc} (g^{\mathbb{G}_m, \text{red}, *} h^{\mathbb{G}_m, \text{red}, *} \mathfrak{L}_{\mu_2, s_2}^{\text{Ind}}) \otimes (g^{\mathbb{G}_m, \text{red}, *} K_{X_1/X_2}^{\pm}) \otimes K_{X_0/X_1}^{\pm} & \xrightarrow[\cong]{\Upsilon_h^{\text{Ind}}} & g^{\mathbb{G}_m, \text{red}, *} \mathfrak{L}_{\mu_1, s_1}^{\text{Ind}} \otimes K_{X_0/X_1}^{\pm} \\ \downarrow \cong & & \downarrow \cong \Upsilon_g^{\text{Ind}} \\ (h^{\mathbb{G}_m, \text{red}} \circ g^{\mathbb{G}_m, \text{red}})^* \mathcal{L}_{\mu_2, s_2}^{\text{Ind}} \otimes K_{X_0/X_2}^{\pm} & \xrightarrow[\cong]{\Upsilon_{h \circ g}^{\text{Ind}}} & \mathcal{L}_{\mu_0, s_0}^{\text{Ind}}. \end{array}$$

This is a direct consequence of the commutativity of the diagram (6.23). Finally, we prove the compatibility between Υ_h^{Ind} and $\theta_{\hat{\mu}, s}$. Assume that we are given algebraic spaces X and Y with \mathbb{G}_m^2 -actions $\hat{\mu} = (\mu_1, \mu_2)$ and $\hat{\nu} = (\nu_1, \nu_2)$. Assume further that we are given a \mathbb{G}_m^2 -equivariant smooth morphism $h: X \rightarrow Y$. Take a d-critical structure t of Y and set $s := h^*t$. Then the following diagram commutes:

$$(6.30) \quad \begin{array}{ccc} h^{\mathbb{G}_m^2, \text{red}, *} \mathcal{L}_{\nu_1, t}^{\text{Ind}} |_{Y^{\hat{\nu}, \text{red}}} \otimes h^{\mathbb{G}_m^2, \text{red}, *} \mathcal{L}_{\nu_2 |_{Y^{\nu_1}, t^{\nu_1}}}^{\text{Ind}} & \xrightarrow[\cong]{\theta_{\hat{\nu}, t}} & h^{\mathbb{G}_m^2, \text{red}, *} \mathcal{L}_{\nu_2, t}^{\text{Ind}} |_{Y^{\hat{\nu}, \text{red}}} \otimes h^{\mathbb{G}_m^2, \text{red}, *} \mathcal{L}_{\nu_1 |_{Y^{\nu_2}, t^{\nu_1}}}^{\text{Ind}} \\ \otimes K_{X/Y}^{\mu_1 \pm} |_{X^{\hat{\mu}, \text{red}}} \otimes K_{X^{\mu_2 \pm} / Y^{\nu_1}} & & \otimes K_{X/Y}^{\mu_2 \pm} |_{X^{\hat{\mu}, \text{red}}} \otimes K_{X^{\mu_1 \pm} / Y^{\nu_2}} \\ \downarrow \cong (\Upsilon_{h, \mu_1}^{\text{Ind}}, \Upsilon_{h |_{X^{\mu_1}}}^{\text{Ind}}) & & \downarrow \cong (\Upsilon_{h, \mu_2}^{\text{Ind}}, \Upsilon_{h |_{X^{\mu_2}}}^{\text{Ind}}) \\ \mathcal{L}_{\mu_1, s}^{\text{Ind}} |_{X^{\hat{\mu}, \text{red}}} \otimes \mathcal{L}_{\mu_2 |_{X^{\mu_1}, s^{\mu_1}}}^{\text{Ind}} & \xrightarrow[\cong]{(-1)^{\text{rank } S_2(\Omega_{X/Y})} \cdot \theta_{\hat{\mu}, s}} & \mathcal{L}_{\mu_2, s}^{\text{Ind}} |_{X^{\hat{\mu}, \text{red}}} \otimes \mathcal{L}_{\mu_1 |_{X^{\mu_2}, s^{\mu_2}}}^{\text{Ind}}. \end{array}$$

Here we use the notation $\Upsilon_{h, \mu_1}^{\text{Ind}}$ and $\Upsilon_{h, \mu_2}^{\text{Ind}}$ to clarify the \mathbb{G}_m -action we consider. This statement is a direct consequence of the commutativity of the diagram (6.23) and (6.24).

6.4. Index line bundles for d-critical stacks. We will introduce the index line bundle for d-critical stacks by gluing index line bundles for \mathbb{G}_m -equivariant d-critical algebraic spaces defined in §6.3:

Proposition 6.8. *Let (\mathfrak{X}, s) be a d -critical stack where \mathfrak{X} is quasi-separated with affine stabilizer groups. Then there exists a line bundle $\mathcal{L}_s^{\text{Ind}}$ on $\text{Grad}(\mathfrak{X})^{\text{red}}$ with the following properties:*

(i) *For each point $x \in \text{Grad}(\mathfrak{X})$, there exists a natural isomorphism*

$$(6.31) \quad \kappa_x^{\text{Ind}}: \mathcal{L}_s^{\text{Ind}}|_x \cong \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})).$$

(ii) *There exists a natural isomorphism*

$$(6.32) \quad \xi_s: K_{\text{Grad}(\mathfrak{X}), u^*s} \otimes \mathcal{L}_s^{\text{Ind}, \otimes 2} \cong u^{\text{red}, *}\mathcal{K}_{\mathfrak{X}, s}.$$

(iii) *For each \mathbb{G}_m -equivariant chart (X, μ, \bar{q}) , there exists a natural isomorphism*

$$(6.33) \quad \Upsilon_{q, \mu}^{\text{Ind}}: q_{\mu}^{\text{red}, *} \mathcal{L}_s^{\text{Ind}} \otimes K_{X/\mathfrak{X}}^{\pm} \cong \mathcal{L}_{\mu, q^*s}^{\text{Ind}}.$$

(iv) *Let $u_1, u_2: \text{Grad}^2(\mathfrak{X}) \rightarrow \text{Grad}(\mathfrak{X})$ be the morphisms induced from the inclusions $i_1, i_2: \mathbb{G}_m \hookrightarrow \mathbb{G}_m^2$ to the first and the second components respectively. Let $\sigma: \text{Grad}^2(\mathfrak{X}) \cong \text{Grad}^2(\mathfrak{X})$ be the isomorphism induced from the swapping isomorphism of \mathbb{G}_m^2 . Then there exists a natural isomorphism*

$$(6.34) \quad \theta_s: u_1^{\text{red}, *} \mathcal{L}_s^{\text{Ind}} \otimes \sigma^* \mathcal{L}_{u^*s}^{\text{Ind}} \cong u_2^{\text{red}, *} \mathcal{L}_s^{\text{Ind}} \otimes \mathcal{L}_{u^*s}^{\text{Ind}}.$$

Here we used the natural identification $\text{Grad}^2(\mathfrak{X}) \cong \text{Grad}(\text{Grad}(\mathfrak{X}))$ given by $\text{Map}(B\mathbb{G}_{m,1} \times B\mathbb{G}_{m,2}, \mathfrak{X}) \cong \text{Map}(B\mathbb{G}_{m,1}, \text{Map}(B\mathbb{G}_{m,2}, \mathfrak{X}))$.

(v) *Let (\mathfrak{X}, s) and (\mathfrak{Y}, t) be d -critical Artin stacks. Then there exists a natural isomorphism*

$$\Psi_{\mathfrak{X}, \mathfrak{Y}}^{\text{Ind}}: \mathcal{L}_s^{\text{Ind}} \boxtimes \mathcal{L}_t^{\text{Ind}} \cong \mathcal{L}_{s \boxplus t}^{\text{Ind}}.$$

Proof. The existence of the line bundle $\mathcal{L}_s^{\text{Ind}}$ with the property (iii) follows from Corollary 5.27 and the commutativity of the diagram (6.29). The existence of the isomorphisms κ_x^{Ind} , ξ_s and θ_s follows from Corollary 5.26 and the commutativity of the diagrams (6.23), (6.27) and (6.30) respectively. The isomorphism $\Psi_{\mathfrak{X} \times \mathfrak{Y}}^{\text{Ind}}$ exists since the map Υ_h^{Ind} in Proposition 6.7 is closed under products under the identification $\Psi_{\mu, \nu}^{\text{Ind}}$. \square

We now prove the compatibility between morphisms that we have constructed in Proposition 6.8. Firstly, the isomorphisms ξ_s , $\Upsilon_{q, \mu}^{\text{Ind}}$ are closed under products under the identification $\Psi_{\mathfrak{X} \times \mathfrak{Y}}^{\text{Ind}}$, since a similar statement is true for \mathbb{G}_m -equivariant d -critical algebraic spaces.

We now prove that κ_x^{Ind} is compatible with ξ_s . We adopt the notation from Proposition 6.8 (ii) and take $x \in \text{Grad}(\mathfrak{X})$. Then the following diagram commutes:

$$(6.35) \quad \begin{array}{ccc} K_{\text{Grad}(\mathfrak{X}), u^*s}|_x \otimes \mathcal{L}_s^{\text{Ind}, \otimes 2}|_x & \xrightarrow[\cong]{\xi_s|_x} & K_{\mathfrak{X}, s}|_{u(x)} \\ \cong \downarrow (\kappa_x, \kappa_x^{\text{Ind}}) & & \downarrow \kappa_{u(x)} \\ \det(\tau^{\geq 0}(\mathbb{L}_{\text{Grad}(\mathfrak{X}), x}))^{\otimes 2} \otimes \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, x}))^{\otimes 2} & \xrightarrow[\cong]{} & \det(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)}))^{\otimes 2}. \end{array}$$

This is a direct consequence of the commutativity of the diagram (6.22). Next, we prove that κ_x^{Ind} is compatible with $\Upsilon_{q, \mu}^{\text{Ind}}$. We adopt the notation from Proposition 6.8 (iii) and take $x \in \text{Grad}(\mathfrak{X})$ and its lift $\hat{x} \in X^{\mu}$. Then the following diagram commutes:

$$(6.36) \quad \begin{array}{ccc} \mathcal{L}_s^{\text{Ind}}|_x \otimes K_{X/\mathfrak{X}}^{\pm}|_{\hat{x}} & \xrightarrow[\cong]{\Upsilon_{q, \mu}^{\text{Ind}}|_{\hat{x}}} & \mathcal{L}_{\mu, q^*s}^{\text{Ind}}|_{\hat{x}} \\ \cong \downarrow \kappa_x^{\text{Ind}} \otimes \text{id} & & \downarrow \kappa_{\hat{x}}^{\text{Ind}} \\ \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) \otimes K_{X/\mathfrak{X}}^{\pm}|_{\hat{x}} & \xrightarrow[\cong]{} & \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{X, \hat{x}})). \end{array}$$

This is a direct consequence of the commutativity of the diagram (6.23). Next, we prove that κ_x^{Ind} is compatible with θ_s . We adopt the notation from Proposition 6.8 (iii) and take $x \in \text{Grad}^2(\mathfrak{X})$. Then the following diagram commutes:

(6.37)

$$\begin{array}{ccc}
\mathcal{L}_s^{\text{Ind}}|_{u_1(x)} \otimes \mathcal{L}_{u^*s}^{\text{Ind}}|_{\sigma(x)} & \xrightarrow[\cong]{ (-1)^{\text{rank}_{S_2}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)}) \cdot \theta_s|_x} } & \mathcal{L}_s^{\text{Ind}}|_{u_2(x)} \otimes \mathcal{L}_{u^*s}^{\text{Ind}}|_x \\
\cong \downarrow (\kappa_{u_1(x)}^{\text{Ind}}, \kappa_{\sigma(x)}^{\text{Ind}}) & & \cong \downarrow (\kappa_{u_2(x)}^{\text{Ind}}, \kappa_x^{\text{Ind}}) \\
\det_{\mathbb{Z} \neq 0 \times \mathbb{Z}}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) \otimes \det_{\{0\} \times \mathbb{Z} \neq 0}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) & & \det_{\mathbb{Z} \times \mathbb{Z} \neq 0}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) \otimes \det_{\mathbb{Z} \neq 0 \times \{0\}}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) \\
& \searrow \cong & \swarrow \cong \\
& \widehat{\det}_{\mathbb{Z} \neq (0,0)}^2(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) &
\end{array}$$

Here we set $S_2 := \mathbb{Z}_{>0} \times \mathbb{Z}_{<0}$. The commutativity of this diagram is a consequence of the commutativity of the diagram (6.24).

We now prove that the compatibility between ξ_s and $\Upsilon_{q,\mu}^{\text{Ind}}$. We adopt the notation from Proposition 6.8. Then the following diagram commutes:

$$\begin{array}{ccc}
q_\mu^{\text{red},*} K_{\text{Grad}(\mathfrak{X}), u^*s} \otimes q_\mu^{\text{red},*} \mathcal{L}_s^{\text{Ind}, \otimes 2} & \xrightarrow[\cong]{ \Upsilon_{q,\mu}^{\text{Ind}} } & q_\mu^{\text{red},*} K_{\text{Grad}(\mathfrak{X}), u^*s} \otimes \mathcal{L}_{\mu, q^*s}^{\text{Ind}, \otimes 2} \\
\otimes K_{X/\mathfrak{X}}^{\otimes 2}|_{X^{\mu, \text{red}}} & & \otimes K_{X^\mu/\text{Grad}(\mathfrak{X})}^{\otimes 2} \\
\cong \downarrow \xi_s & & \cong \downarrow \Upsilon_{q_\mu} \\
q_\mu^{\text{red},*} u^{\text{red},*} K_{\mathfrak{X}, s} \otimes K_{X/\mathfrak{X}}^{\otimes 2}|_{X^{\mu, \text{red}}} & \xrightarrow[\cong]{ \Upsilon_q } & K_{X, q^*s}|_{X^{\mu, \text{red}}} \\
& & \cong \downarrow \xi_{\mu, q^*s} \\
& & K_{X^\mu, (q^*s)^\mu} \otimes \mathcal{L}_{\mu, q^*s}^{\text{Ind}, \otimes 2}
\end{array}$$

This is a direct consequence of the construction of ξ_s . Next, we prove the compatibility between ξ_s and θ_s . We adopt the notation from Proposition 6.8. The following diagram commutes:

$$\begin{array}{ccc}
K_{\text{Grad}^2(\mathfrak{X}), u^{(2), *s}} & \xrightarrow[\cong]{ \text{id} \otimes \theta_s^{\otimes 2} } & K_{\text{Grad}^2(\mathfrak{X}), u^{(2), *s}} \\
\otimes u_1^{\text{red},*} \mathcal{L}_s^{\text{Ind}, \otimes 2} \otimes \sigma^* \mathcal{L}_{u^*s}^{\text{Ind}, \otimes 2} & & \otimes u_2^{\text{red},*} \mathcal{L}_s^{\text{Ind}, \otimes 2} \otimes \mathcal{L}_{u^*s}^{\text{Ind}, \otimes 2} \\
\cong \downarrow \sigma^* \xi_{u^*s} & & \cong \downarrow \xi_{u^*s} \\
u_1^{\text{red},*} K_{\text{Grad}(\mathfrak{X}), u^*s} \otimes u_1^{\text{red},*} \mathcal{L}_s^{\text{Ind}, \otimes 2} & & u_2^{\text{red},*} K_{\text{Grad}(\mathfrak{X}), u^*s} \otimes u_2^{\text{red},*} \mathcal{L}_s^{\text{Ind}, \otimes 2} \\
& \searrow \cong \downarrow u_1^{\text{red},*} \xi_s & \swarrow \cong \downarrow u_2^{\text{red},*} \xi_s \\
& u^{(2), \text{red},*} K_{\mathfrak{X}, s} &
\end{array}$$

Here we let $u^{(2)}: \text{Grad}^2(\mathfrak{X}) \rightarrow \mathfrak{X}$ be the canonical evaluation map and we used the identification

$$\sigma^* K_{\text{Grad}^2(\mathfrak{X}), u^{(2), *s}} \cong K_{\text{Grad}^2(\mathfrak{X}), u^{(2), *s}}$$

for the left vertical map. This is a direct consequence of the commutativity of the diagram (6.28).

We now discuss the associativity of $\Upsilon_{q,\mu}^{\text{Ind}}$. We adopt the notation from Proposition 6.8. Assume further that we are given a morphism $f: (X_1, \mu_1, \bar{q}_1) \rightarrow (X_2, \mu_2, \bar{q}_2)$ of \mathbb{G}_m -equivariant charts. Then the following diagram commutes:

$$\begin{array}{ccc}
(f^{\mathbb{G}_m, \text{red},*} q_{2, \mu_2}^{\text{red},*} \mathcal{L}_s^{\text{Ind}}) \otimes (f^{\mathbb{G}_m, \text{red},*} K_{X_2^{\mu_2}/\text{Grad}(\mathfrak{X})}^\pm) \otimes K_{X_1^{\mu_1}/X_2^{\mu_2}}^\pm & \xrightarrow[\cong]{ \Upsilon_{q_2, \mu_2}^{\text{Ind}} } & f^{\mathbb{G}_m, \text{red},*} \mathcal{L}_{\mu_2, q_2^*s}^{\text{Ind}} \otimes K_{X_1^{\mu_1}/X_2^{\mu_2}}^\pm \\
\cong \downarrow & & \cong \downarrow \Upsilon_f^{\text{Ind}} \\
(q_{2, \mu_2}^{\text{red}} \circ f^{\mathbb{G}_m, \text{red}})^* \mathcal{L}_s^{\text{Ind}} \otimes K_{X_1^{\mu_1}/\text{Grad}(\mathfrak{X})}^\pm & \xrightarrow[\cong]{ \Upsilon_{q_1, \mu_1}^{\text{Ind}} } & \mathcal{L}_{\mu_1, q_1^*s}^{\text{Ind}}
\end{array}$$

This is a direct consequence of the construction of $\Upsilon_{q,\mu}^{\text{Ind}}$. We now prove the compatibility between $\Upsilon_{q,\mu}^{\text{Ind}}$ and θ_s . We adopt the notation from Proposition 6.8. We take a \mathbb{G}_m^2 -equivariant chart $(X, \hat{\mu}, \bar{q})$. Then the following diagram commutes:

$$(6.41) \quad \begin{array}{ccc} q_{\hat{\mu}}^{\text{red},*} u_1^{\text{red},*} \mathcal{L}_s^{\text{Ind}} \otimes q_{\hat{\mu}}^{\text{red},*} \sigma^* \mathcal{L}_{u^*s}^{\text{Ind}} & \xrightarrow[q_{\hat{\mu}}^{\text{red},*} \theta_s \otimes \text{id}]{\cong} & q_{\hat{\mu}}^{\text{red},*} u_2^{\text{red},*} \mathcal{L}_s^{\text{Ind}} \otimes q_{\hat{\mu}}^{\text{red},*} \mathcal{L}_{u^*s}^{\text{Ind}} \\ \otimes K_{X/\mathfrak{X}}^{\mathbb{Z}_{\neq 0}^2} & & \otimes K_{X/\mathfrak{X}}^{\mathbb{Z}_{\neq 0}^2} \\ \cong \downarrow (\Upsilon_{q,\mu_1}^{\text{Ind}}, \Upsilon_{q\mu_1,\mu_2}^{\text{Ind}}|_{X^{\mu_1}}) & & \cong \downarrow (\Upsilon_{q,\mu_2}^{\text{Ind}}, \Upsilon_{q\mu_2,\mu_1}^{\text{Ind}}|_{X^{\mu_2}}) \\ \mathcal{L}_{\mu_1,q^*s}^{\text{Ind}}|_{X^{\hat{\mu},\text{red}}} \otimes \mathcal{L}_{\mu_2|_{X^{\mu_1}},q_{\mu_1}^*}^{\text{Ind}} u^*s & \xrightarrow[\cong]{(-1)^{\text{rank}_{S_2}(\Omega_{X/\mathfrak{X}})} \cdot \theta_{\hat{\mu},q^*s}} & \mathcal{L}_{\mu_2,q^*s}^{\text{Ind}}|_{X^{\hat{\mu},\text{red}}} \otimes \mathcal{L}_{\mu_1|_{X^{\mu_2}},q_{\mu_2}^*}^{\text{Ind}} u^*s. \end{array}$$

This is a direct consequence of the construction of $\theta_{\bar{s}}$.

We now describe the index line bundle for d-critical stacks underlying (-1) -shifted symplectic stacks. Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a (-1) -shifted symplectic derived Artin stack which is quasi-separated with affine stabilizers. Let $(\mathfrak{X}^{\text{cl}}, s)$ denote the underlying d-critical stack. For $x \in \text{Grad}(\mathfrak{X})$, we define an isomorphism

$$(6.42) \quad \kappa_x^{\text{Ind,der}} : \det_+(\mathbb{L}_{\mathfrak{X},u(x)}) \cong \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)}))$$

so that the following diagram commutes:

$$(6.43) \quad \begin{array}{ccc} \det(\mathbb{L}_{\mathfrak{X},u(x)}^+) & \xrightarrow[\cong]{\kappa_x^{\text{Ind,der}}} & \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) \\ \cong \downarrow & & \cong \uparrow (2.5), (2.14) \\ \det_+(\tau^{\leq -1}(\mathbb{L}_{\mathfrak{X},u(x)})) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) & \xrightarrow[\cong]{(-\cdot\omega_{\mathfrak{X}})^{-1}|_x \otimes \text{id}} & \det_+(\tau^{\leq 0}(\mathbb{T}_{\mathfrak{X},u(x)}[1]) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})). \end{array}$$

We will construct an isomorphism of one-dimensional vector spaces

$$\Lambda_{\mathfrak{X},x}^{\text{Ind,pre}} : \det_+(\mathbb{L}_{\mathfrak{X},u(x)}) \cong \mathcal{L}_s^{\text{Ind}}|_x$$

by $\kappa_x^{\text{Ind,-1}} \circ \kappa_x^{\text{Ind,der}}$.

Proposition 6.9. *The map $x \mapsto \Lambda_{\mathfrak{X},x}^{\text{Ind,pre}}$ is algebraic, hence it defines an isomorphism*

$$\Lambda_{\mathfrak{X}}^{\text{Ind}} : \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}}) \cong \mathcal{L}_s^{\text{Ind}}.$$

Proof. Take morphisms $q: \widehat{V} \rightarrow \mathfrak{X}$, $\tau: \widehat{V} \rightarrow V$ as in Lemma 6.10 below. We let $q_{\mu}: \widehat{V}^{\mu} \rightarrow \text{Grad}(\mathfrak{X})$ and $\tau_{\mu}: \widehat{V}^{\mu} \rightarrow V^{\mu}$ the naturally induced morphisms. It is enough to show that the map $x \mapsto \Lambda_{\mathfrak{X},x}^{\text{Ind,pre}}$ is algebraic after pulling back to \widehat{V} . Consider the following morphism of fibre sequences

$$(6.44) \quad \begin{array}{ccccc} \mathbb{T}_{\widehat{V}/\mathfrak{X}}[1] & \longrightarrow & \mathbb{T}_{\widehat{V}}[1] & \longrightarrow & q^* \mathbb{T}_{\mathfrak{X}}[1] \\ \downarrow \eta|_{\simeq} & & \downarrow & & \downarrow \\ \mathbb{L}_{\widehat{V}/V}[-1] & \longrightarrow & \tau^* \mathbb{L}_V & \longrightarrow & \mathbb{L}_{\widehat{V}} \end{array}$$

where the middle and right vertical maps are induced from (-1) -shifted 2-form $\tau^* \omega_V \sim q^* \omega_{\mathfrak{X}}$, the dotted arrow is induced by the homotopy commutativity of the right square, which is invertible by Remark 3.12. The map $\eta^{\vee}[-1]: \mathbb{L}_{\widehat{V}/\mathfrak{X}} \simeq \mathbb{T}_{\widehat{V}/V}[-2]$ induces a natural isomorphism

$$(6.45) \quad \alpha_+ : \widehat{\det}_-(\mathbb{L}_{\widehat{V}/\mathfrak{X}}) \cong \widehat{\det}_-(\mathbb{T}_{\widehat{V}/V}[-2]) \cong \widehat{\det}_+(\mathbb{L}_{\widehat{V}/V})^{\vee}$$

where the latter isomorphism is induced from (2.5) and (2.14). We construct an isomorphism

$$\Upsilon_q^{\text{Ind,der}} : q_{\mu}^{\text{red},*} \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_+(\mathbb{L}_{\widehat{V}/\mathfrak{X}}) \cong \widehat{\det}_+(\mathbb{L}_{\widehat{V}}), \quad \Upsilon_{\tau}^{\text{Ind,der}} : \widehat{\det}_+(\mathbb{L}_{\widehat{V}}) \otimes \widehat{\det}_-(\mathbb{L}_{\widehat{V}/\mathfrak{X}}) \cong \tau_{\mu}^{\text{red},*} \widehat{\det}_+(\mathbb{L}_V)$$

using fibre sequences

$$q^* \mathbb{L}_{\mathfrak{X}} \rightarrow \mathbb{L}_{\widehat{V}} \rightarrow \mathbb{L}_{\widehat{V}/\mathfrak{X}}, \quad \tau^* \mathbb{L}_V \rightarrow \mathbb{L}_{\widehat{V}} \rightarrow \mathbb{L}_{\widehat{V}/V},$$

the isomorphism (2.3) and the isomorphism (6.45). We construct an isomorphism

$$\Upsilon_{(q,\tau)}^{\text{Ind,der}}: q_\mu^{\text{red},*} \widehat{\det}_+(\mathbb{L}\mathfrak{x}) \otimes \widehat{\det}_\pm(\mathbb{L}\widehat{\mathcal{V}}/\mathfrak{x}) \cong \tau_\mu^{\text{red},*} \widehat{\det}_+(\mathbb{L}V)$$

by the composite of $\Upsilon_\tau^{\text{Ind,der}}$ and $\Upsilon_q^{\text{Ind,der}} \otimes \text{id}_{\widehat{\det}_-(\mathbb{L}\widehat{\mathcal{V}}/\mathfrak{x})}$. Finally, we construct an isomorphism

$$\Lambda_{\mathfrak{x},(q,\tau)}^{\text{Ind,pre}}: q_\mu^{\text{red},*} \widehat{\det}_+(\mathbb{L}\mathfrak{x}) \cong q_\mu^{\text{red},*} \mathcal{L}_s^{\text{Ind}}$$

so that the following diagram commutes:

$$\begin{array}{ccc} q_\mu^{\text{red},*} \widehat{\det}_+(\mathbb{L}\mathfrak{x}) \otimes \widehat{\det}_\pm(\mathbb{L}\widehat{\mathcal{V}}/\mathfrak{x}) & \xrightarrow[\cong]{\Lambda_{\mathfrak{x},(q,\tau)}^{\text{Ind,pre}} \otimes \text{id}} & q_\mu^{\text{red},*} \mathcal{L}_s^{\text{Ind}} \otimes \widehat{\det}_\pm(\mathbb{L}\widehat{\mathcal{V}}/\mathfrak{x}) \\ \cong \downarrow \Upsilon_{(q,\tau)}^{\text{Ind,der}} & & \cong \downarrow \Upsilon_{q^{\text{cl}},\mu}^{\text{Ind}} \\ \tau_\mu^{\text{red},*} \widehat{\det}_+(\mathbb{L}V) & \xrightarrow[\cong]{\Lambda_V^{\text{Ind}}} & \mathcal{L}_{\mu,q^{\text{cl}},*s}^{\text{Ind}} \end{array}$$

Take a point $\hat{v} \in \widehat{V}^{\text{Gm}}$ and set $x = q_\mu(\hat{v})$ and $v = \tau_\mu(\hat{v})$. We claim that the following diagram commutes:

$$(6.46) \quad \begin{array}{ccc} \det_\pm(\tau^{\geq 0}(\mathbb{L}\mathfrak{x}, u(x))) & \xlongequal{\quad} & \det_\pm(\tau^{\geq 0}(\mathbb{L}\mathfrak{x}, u(x))) \\ \kappa_x^{\text{Ind,der}} \uparrow \cong & & \kappa_x^{\text{Ind}} \uparrow \cong \\ \det_+(\mathbb{L}\mathfrak{x}, u(x)) & \xrightarrow[\cong]{\Lambda_{\mathfrak{x},(q,\tau)}^{\text{Ind,pre}}|_{\hat{v}}} & \mathcal{L}_s^{\text{Ind}}|_x \end{array}$$

which in particular implies that the map $x \mapsto \Lambda_{\mathfrak{x},x}^{\text{Ind,pre}}$ is algebraic as desired.

To prove this, we first show that the map $\Upsilon_{(q,\tau)}^{\text{Ind,der}}$ is compatible with maps $\kappa_x^{\text{Ind,der}}$ and $\kappa_v^{\text{Ind,der}}$. Namely, we first show that the following diagram commutes:

$$(6.47) \quad \begin{array}{ccc} \det_\pm(\tau^{\geq 0}(\mathbb{L}\mathfrak{x}, u(x))) \otimes \det_\pm(\mathbb{L}\widehat{\mathcal{V}}/\mathfrak{x}, \hat{v}) & \xleftarrow[\cong]{\kappa_x^{\text{Ind,der}} \otimes \text{id}} & \det_+(\mathbb{L}\mathfrak{x}, u(x)) \otimes \det_\pm(\mathbb{L}\widehat{\mathcal{V}}/\mathfrak{x}, \hat{v}) \\ \cong \downarrow & & \cong \downarrow \Upsilon_{(q,\tau)}^{\text{Ind,der}}|_{\hat{v}} \\ \det_\pm(\tau^{\geq 0}(\mathbb{L}V, v)) & \xleftarrow[\cong]{\kappa_v^{\text{Ind,der}}} & \det_+(\mathbb{L}V, v) \end{array}$$

Here the left vertical map is constructed using the fibre sequence $\tau^{\geq 0}(\mathbb{L}\mathfrak{x}, u(x)) \rightarrow \tau^{\geq 0}(\mathbb{L}V, v) \rightarrow \mathbb{L}\widehat{\mathcal{V}}/\mathfrak{x}, \hat{v}$. We define a map

$$\kappa_{\hat{v}}^{\text{Ind,der}}: \det_+(\mathbb{L}\widehat{\mathcal{V}}, \hat{v}) \cong \det_+(\tau^{\geq 0}(\mathbb{L}V, v)) \otimes \det_-(\tau^{\geq 0}(\mathbb{L}\mathfrak{x}, u(x)))$$

by the composition

$$\begin{aligned} \det_+(\mathbb{L}\widehat{\mathcal{V}}, \hat{v}) &\cong \det_+(\tau^{\leq -1}(\mathbb{L}\widehat{\mathcal{V}}, \hat{v})) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}\widehat{\mathcal{V}}, \hat{v})) \cong \det_+(\tau^{\leq -1}(\mathbb{L}\mathfrak{x}, u(x))) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}V, v)) \\ &\cong \det_-(\tau^{\geq 0}(\mathbb{L}\mathfrak{x}, u(x))) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}V, v)) \end{aligned}$$

where the last isomorphism is induced from the isomorphism $\tau^{\leq -1}(\mathbb{L}_{\mathcal{X}, u(x)}) \simeq \tau^{\geq 0}(\mathbb{L}_{\mathcal{X}, u(x)})^\vee[1]$ and the isomorphisms (2.5) and (2.14). The diagram (6.47) can be decomposed into the following diagram:

$$(6.48) \quad \begin{array}{ccc} \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathcal{X}, u(x)})) \otimes \det_{\pm}(\mathbb{L}_{\widehat{V}/\mathcal{X}, \widehat{v}}) & \xleftarrow[\cong]{\kappa_x^{\text{Ind, der}} \otimes \text{id}} & \det_+(\mathbb{L}_{\mathcal{X}, u(x)}) \otimes \det_{\pm}(\mathbb{L}_{\widehat{V}/\mathcal{X}, \widehat{v}}) \\ \mathbb{R} \downarrow & & \cong \downarrow \Upsilon_q^{\text{Ind, der}} \otimes \text{id} \\ \det_-(\tau^{\geq 0}(\mathbb{L}_{\mathcal{X}, u(x)})) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}_{V, v})) \otimes \det_-(\mathbb{L}_{\widehat{V}/\mathcal{X}, \widehat{v}}) & \xleftarrow[\cong]{\kappa_{\widehat{v}}^{\text{Ind, der}} \otimes \text{id}} & \det_+(\mathbb{L}_{\widehat{V}, \widehat{v}}) \otimes \det_-(\mathbb{L}_{\widehat{V}/\mathcal{X}, \widehat{v}}) \\ \cong \downarrow \text{id} \otimes \text{sw} & & \downarrow \cong \Upsilon_{\tau}^{\text{Ind, der}} \\ \det_-(\tau^{\geq 0}(\mathbb{L}_{\mathcal{X}, u(x)})) \otimes \det_-(\mathbb{L}_{\widehat{V}/\mathcal{X}, \widehat{v}}) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}_{V, v})) & & \det_+(\mathbb{L}_{V, v}) \\ \mathbb{R} \downarrow & & \downarrow \cong \Upsilon_{\tau}^{\text{Ind, der}} \\ \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{V, v})) & \xleftarrow[\cong]{\kappa_v^{\text{Ind, der}} \otimes \text{id}} & \det_+(\mathbb{L}_{V, v}). \end{array}$$

The commutativity of the upper square follows by Lemma 2.4 applied to the positive part of the following fibre double sequence

$$\begin{array}{ccccc} \tau^{\leq -1}(\mathbb{L}_{\mathcal{X}, u(x)}) & \xlongequal{\quad} & \tau^{\leq -1}(\mathbb{L}_{\mathcal{X}, u(x)}) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{L}_{\mathcal{X}, u(x)} & \longrightarrow & \mathbb{L}_{\widehat{V}, \widehat{v}} & \longrightarrow & \mathbb{L}_{\widehat{V}/\mathcal{X}, \widehat{v}} \\ \downarrow & & \downarrow & & \parallel \\ \tau^{\geq 0}(\mathbb{L}_{\mathcal{X}, u(x)}) & \longrightarrow & \tau^{\geq 0}(\mathbb{L}_{\widehat{V}, \widehat{v}}) & \longrightarrow & \mathbb{L}_{\widehat{V}/\mathcal{X}, \widehat{v}}. \end{array}$$

Now we prove the commutativity of the lower square of (6.48). Consider the following diagram:

$$\begin{array}{ccc} \det_+(\mathbb{L}_{\widehat{V}, \widehat{v}}) \otimes \det_-(\mathbb{L}_{\widehat{V}/\mathcal{X}, \widehat{v}}) & \xrightarrow[\cong]{} & \det_+(\tau^{\leq -1}(\mathbb{L}_{\widehat{V}, \widehat{v}})) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}_{\widehat{V}, \widehat{v}})) \otimes \det_-(\mathbb{L}_{\widehat{V}/\mathcal{X}, \widehat{v}}) \\ \cong \downarrow \text{id} \otimes \alpha_+ & & \cong \downarrow \\ \det_+(\mathbb{L}_{\widehat{V}, \widehat{v}}) \otimes \det_+(\mathbb{L}_{\widehat{V}/V, v})^\vee & \xrightarrow{\quad} & \det_+(\mathbb{L}_{\widehat{V}/V, \widehat{v}}) \otimes \det_+(\tau^{\leq -1}(\mathbb{L}_{V, v})) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}_{\widehat{V}, \widehat{v}})) \otimes \det_-(\mathbb{L}_{\widehat{V}/\mathcal{X}, \widehat{v}}) \\ \cong \downarrow \Upsilon_{\tau}^{\text{Ind, der}} & & \cong \downarrow \text{id} \otimes \alpha_+ \\ \det_+(\mathbb{L}_{V, v}) & \xrightarrow[\cong]{} & \det_+(\mathbb{L}_{\widehat{V}/V, \widehat{v}}) \otimes \det_+(\tau^{\leq -1}(\mathbb{L}_{V, v})) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}_{\widehat{V}, \widehat{v}})) \otimes \det_+(\mathbb{L}_{\widehat{V}/V, \widehat{v}})^\vee \\ \cong \downarrow & & \cong \downarrow \\ \det_+(\mathbb{L}_{V, v}) & \xrightarrow[\cong]{} & \det_+(\tau^{\leq -1}(\mathbb{L}_{V, v})) \otimes \det_+(\tau^{\geq 0}(\mathbb{L}_{V, v})). \end{array}$$

Here the right top vertical arrow is induced from the fibre sequence $\tau^{\leq -1}(\mathbb{L}_{V, v}) \rightarrow \tau^{\leq -1}(\mathbb{L}_{\widehat{V}, \widehat{v}}) \rightarrow \mathbb{L}_{\widehat{V}/V, \widehat{v}}$. The commutativity of this diagram follows from Lemma 2.4 applied to the positive part of the following fibre double sequence

$$\begin{array}{ccccc} \tau^{\leq -1}(\mathbb{L}_{V, v}) & \longrightarrow & \tau^{\leq -1}(\mathbb{L}_{\widehat{V}, \widehat{v}}) & \longrightarrow & \mathbb{L}_{\widehat{V}/V} \\ \downarrow & & \downarrow & & \parallel \\ \mathbb{L}_V & \longrightarrow & \mathbb{L}_{\widehat{V}} & \longrightarrow & \mathbb{L}_{\widehat{V}/V} \\ \downarrow & & \downarrow & & \downarrow \\ \tau^{\geq 0}(\mathbb{L}_{V, v}) & \xrightarrow[\cong]{} & \tau^{\geq 0}(\mathbb{L}_{\widehat{V}, \widehat{v}}) & \longrightarrow & 0 \end{array}$$

Therefore the commutativity of the lower square of the diagram (6.48) is reduced to proving that the outer square of the following diagram commutes:

(6.49)

$$\begin{array}{ccccc}
\det_-(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) & \xrightarrow{\cong} & \det_+(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})^\vee[1]) & \xrightarrow{\cong} & \det_+(\tau^{\leq -1}(\mathbb{L}_{\widehat{V}, \hat{v}})) \otimes \det_-(\mathbb{L}_{\widehat{V}/\mathfrak{X}, \hat{v}}) \\
\otimes \det_-(\mathbb{L}_{\widehat{V}/\mathfrak{X}, \hat{v}}) & & \otimes \det_+(\mathbb{T}_{\widehat{V}/\mathfrak{X}, \hat{v}}[1]) & & \downarrow \cong \\
\downarrow \cong & & \downarrow \cong & & \det_+(\mathbb{L}_{\widehat{V}/V, \hat{v}}) \otimes \det_+(\tau^{\leq -1}(\mathbb{L}_{V, v})) \otimes \det_-(\mathbb{L}_{\widehat{V}/\mathfrak{X}, \hat{v}}) \\
& & & & \downarrow \cong \text{id} \otimes \alpha_+ \\
& & & & \det_+(\mathbb{L}_{\widehat{V}/V, \hat{v}}) \otimes \det_+(\tau^{\leq -1}(\mathbb{L}_{V, v})) \otimes \det_+(\mathbb{L}_{\widehat{V}/V, \hat{v}})^\vee \\
& & & & \downarrow \cong \\
\det_-(\tau^{\geq 0}(\mathbb{L}_{\widehat{V}, \hat{v}})) & \xrightarrow{\cong} & \det_+(\tau^{\geq 0}(\mathbb{L}_{\widehat{V}, \hat{v}})^\vee[1]) & \xrightarrow{\cong} & \det_+(\tau^{\leq -1}(\mathbb{L}_{V, v})).
\end{array}$$

Here the top and bottom left horizontal maps are the isomorphisms (2.5) and (2.14) and the top (resp. bottom) right horizontal map is induced from the isomorphism $\tau^{\leq -1}(\mathbb{L}_{\widehat{V}, \hat{v}}) \simeq \tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})^\vee[1]$ (resp. $\tau^{\leq -1}(\mathbb{L}_{V, v}) \simeq \tau^{\geq 0}(\mathbb{L}_{\widehat{V}, \hat{v}})^\vee[1]$). It follows from Corollary 2.5 and (2.16) that the left square commutes. Now we will show that the right square commutes. To see this, consider the following equivalence of fibre sequences obtained by truncating the map of fibre sequences (6.44):

$$(6.50) \quad \begin{array}{ccccc}
\mathbb{T}_{\widehat{V}/\mathfrak{X}, \hat{v}}[1] & \longrightarrow & \tau^{\geq 0}(\mathbb{L}_{\widehat{V}, \hat{v}})^\vee[1] & \longrightarrow & \tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})^\vee[1] \\
\eta \downarrow \cong & & \cong \downarrow & & \cong \downarrow \\
\mathbb{L}_{\widehat{V}/V, \hat{v}}[-1] & \longrightarrow & \tau^{\leq -1}(\mathbb{L}_{V, v}) & \longrightarrow & \tau^{\leq -1}(\mathbb{L}_{\widehat{V}, \hat{v}}).
\end{array}$$

This shows that the following diagram commutes:

$$(6.51) \quad \begin{array}{ccc}
\det_+(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})^\vee[1]) \otimes \det_+(\mathbb{T}_{\widehat{V}/\mathfrak{X}, \hat{v}}[1]) & \xrightarrow{\cong} & \det_+(\tau^{\leq -1}(\mathbb{L}_{\widehat{V}, \hat{v}})) \otimes \det_+(\mathbb{L}_{\widehat{V}/V, \hat{v}}[-1]) \\
\cong \downarrow & & \cong \downarrow \\
\det_+(\tau^{\geq 0}(\mathbb{L}_{\widehat{V}, \hat{v}})^\vee[1]) & \xrightarrow{\cong} & \det_+(\tau^{\leq -1}(\mathbb{L}_{V, v})).
\end{array}$$

Next, consider the following diagram:

$$(6.52) \quad \begin{array}{ccccc}
& & \alpha_+ & & \\
& & \text{---} & & \text{---} \\
\det_-(\mathbb{L}_{\widehat{V}/\mathfrak{X}, \hat{v}}) & \xrightarrow[\cong]{\det(\eta_+[-1]^\vee|_{\hat{v}})} & \det_-(\mathbb{T}_{\widehat{V}/V, \hat{v}}[2]) & \xrightarrow[\cong]{} & \det_+(\mathbb{L}_{\widehat{V}/V, \hat{v}})^\vee \\
\cong \downarrow & & \downarrow & & \parallel \\
\det_+(\mathbb{T}_{\widehat{V}/\mathfrak{X}, \hat{v}}[1]) & \xrightarrow[\cong]{\det(\eta_+|_{\hat{v}})} & \det_+(\mathbb{L}_{\widehat{V}/V, \hat{v}}[-1]) & \xrightarrow[\cong]{} & \det_+(\mathbb{L}_{\widehat{V}/V, \hat{v}})^\vee.
\end{array}$$

The commutativity of the diagram (2.15) and (2.19) implies that this diagram commutes up to the sign $(-1)^{\text{rank } \mathbb{L}_{\widehat{V}/\mathfrak{X}, \hat{v}}^-}$.

By the commutativity of the diagram (2.9) applied to the fibre sequence $\tau^{\leq -1}(\mathbb{L}_{V,v}) \rightarrow \tau^{\leq -1}(\mathbb{L}_{\widehat{V},\widehat{v}}) \rightarrow \mathbb{L}_{\widehat{V}/V,\widehat{v}}$, we see that the following diagram commutes up to the sign $(-1)^{\text{rank } \mathbb{L}_{\widehat{V}/V,\widehat{v}}^+}$:

$$(6.53) \quad \begin{array}{ccc} \det_+(\mathbb{L}_{\widehat{V}/V,\widehat{v}}[-1]) \otimes \det_+(\tau^{\leq -1}(\mathbb{L}_{V,v})) \otimes \det_+(\mathbb{L}_{\widehat{V}/V,\widehat{v}}) & \xrightarrow{\cong} & \otimes \det_+(\tau^{\leq -1}(\mathbb{L}_{\widehat{V},\widehat{v}})) \otimes \det_+(\mathbb{L}_{\widehat{V}/V,\widehat{v}}) \\ \cong \downarrow & & \cong \downarrow \\ \det_+(\mathbb{L}_{\widehat{V}/V,\widehat{v}})^\vee \otimes \det_+(\tau^{\leq -1}(\mathbb{L}_{V,v})) \otimes \det_+(\mathbb{L}_{\widehat{V}/V,\widehat{v}}) & \xrightarrow{\cong} & \det_+(\tau^{\leq -1}(\mathbb{L}_{V,v})). \end{array}$$

By combining the commutativity properties of diagrams (6.51), (6.52) and (6.53), we see that the right square of the diagram (6.49) commutes. In particular, the diagram (6.47) commutes.

Now consider the following diagram:

$$\begin{array}{ccccc} \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) & & & & \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) \\ \otimes \det_{\pm}(\mathbb{L}_{\widehat{V}/\mathfrak{X},\widehat{v}}) & \xrightarrow{\cong} & & \xrightarrow{\cong} & \otimes \det_{\pm}(\mathbb{L}_{\widehat{V}/\mathfrak{X},\widehat{v}}) \\ \downarrow \cong & \swarrow \kappa_x^{\text{Ind,der}} \otimes \text{id} & & \searrow \cong & \downarrow \cong \\ & \det_+(\mathbb{L}_{\mathfrak{X},u(x)}) \otimes \det_{\pm}(\mathbb{L}_{\widehat{V}/\mathfrak{X},\widehat{v}}) & \xrightarrow{\Lambda_{\mathfrak{X},(q,\tau)}^{\text{Ind,pre}}|_{\widehat{v}} \otimes \text{id}} & \mathcal{L}_s^{\text{Ind}}|_x \otimes \det_{\pm}(\mathbb{L}_{\widehat{V}/\mathfrak{X},\widehat{v}}) & \\ & \downarrow \cong & & \downarrow \cong & \\ & \det_+(\mathbb{L}_{V,v}) & \xrightarrow{\Lambda_V^{\text{Ind}}|_v} & \mathcal{L}_{\mu,q^{\text{cl}},*s}^{\text{Ind}}|_v & \\ \downarrow \cong & \swarrow \kappa_v^{\text{Ind,der}} & & \searrow \cong & \downarrow \cong \\ \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{V,v})) & & & & \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{V,v})). \end{array}$$

The middle square commutes by definition, the left square is nothing but the diagram (6.47), the bottom diagram is (6.26) and the right diagram is (6.36). In particular, we see that the top diagram, and hence the diagram (6.46), commutes. \square

Lemma 6.10. *Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a (-1) -shifted symplectic stack such that \mathfrak{X} is quasi-separated with affine stabilizers and $x \in \text{Grad}^n(\mathfrak{X})$ be a point. Then we can find the following data:*

- A quasi-separated derived algebraic space \widehat{X} with a \mathbb{G}_m^n -action.
- A smooth morphism $[\widehat{X}/\mathbb{G}_m^n] \rightarrow \mathfrak{X}$ and a point $\widehat{x} \in \widehat{X}^{\mathbb{G}_m^n}$ such that the image of \widehat{x} under the natural map $\widehat{X}^{\mathbb{G}_m^n} \rightarrow \text{Grad}^n(\mathfrak{X})$ is x .
- A \mathbb{G}_m^n -equivariant quasi-separated derived algebraic space X with a morphism $i: \widehat{X} \rightarrow X$ which is an isomorphism on the classical truncations.
- A \mathbb{G}_m^n -invariant (-1) -shifted symplectic structure ω_X on X such that the correspondence $\mathfrak{X} \leftarrow \widehat{X} \rightarrow X$ carries a Lagrangian structure.

Proof. This is a combination of the Darboux theorem (Theorem 3.11) in [Ben+15] and the equivariant Darboux theorem (Proposition 3.13) in [Par24].

We first claim that there exists a pointed smooth map

$$([R/\mathbb{G}_m^n], r) \rightarrow (\mathfrak{X}, x)$$

for a derived affine scheme R with an action of \mathbb{G}_m^n and a fixed point $r \in R^{\mathbb{G}_m^n}$. Let $\text{Aut}_{\mathfrak{X}}(x)$ be the stabilizer of \mathfrak{X}^{cl} at x and let T be the image of the group homomorphism $\mathbb{G}_m^n \rightarrow \text{Aut}_{\mathfrak{X}}(x)$, which is also a torus. By [AHR20, Theorem 1.1], we can find a pointed smooth morphism $([P/T], p) \rightarrow (\mathfrak{X}^{\text{cl}}, x)$ for an affine scheme P with a T -action and a T -fixed point $p \in P$. By [Alp+24, Theorem 1.13] (and [Alp+24, Proposition 6.1]), the smooth morphism $[P/T] \rightarrow \mathfrak{X}^{\text{cl}}$ lifts to a smooth morphism $[R/T] \rightarrow \mathfrak{X}$ for a derived affine scheme R with a T -action. Consider the composition $[R/\mathbb{G}_m^n] \rightarrow [R/T] \rightarrow \mathfrak{X}$.

We then find a \mathbb{G}_m^n -invariant locally closed subscheme $\widehat{X} \subseteq R$ that contains r such that the composition

$$\widehat{X} \rightarrow R \rightarrow [R/\mathbb{G}_m^n] \rightarrow \mathfrak{X}$$

is smooth of *minimal dimension* at $\hat{x} \in \widehat{X}$, that is, $\dim(H^i(\mathbb{T}_{\widehat{X}, \hat{x}})) = \dim(H^i(\mathbb{T}_{\mathfrak{X}, x}))$ for all $i \geq 0$. This follows from (the third paragraph of) Step 1 of [Par24, Lemma 4.1.8].

By [Par24, Lemma 4.1.8], there exists a \mathbb{G}_m^n -equivariant closed embedding

$$i : \widehat{X} \hookrightarrow X,$$

into a quasi-smooth derived affine scheme X with a \mathbb{G}_m^n -action whose classical truncation $i^{\text{cl}} : \widehat{X}^{\text{cl}} \rightarrow X^{\text{cl}}$ is an isomorphism. Moreover, we may assume that X is of minimal dimension in the sense that: $\dim(H^i(\mathbb{T}_{\widehat{X}, \hat{x}})) = \dim(H^i(\mathbb{T}_{X, \hat{x}}))$ for $i = 0, 1$.

By Proposition 3.2, the symplectic form $\omega_{\mathfrak{X}}$ has a canonical exact structure. By [Par24, Lemma 4.1.9], the pullback of the exact symplectic form along $[\widehat{X}/\mathbb{G}_m^n] \rightarrow \mathfrak{X}$ lifts to a (-1) -shifted exact 2-form on $[X/\mathbb{G}_m^n]$. In particular, we have an (exact) isotropic correspondence

$$\begin{array}{ccc} & [\widehat{X}/\mathbb{G}_m^n] & \\ \swarrow & & \searrow \\ \mathfrak{X} & & [X/\mathbb{G}_m^n]. \end{array}$$

The desired \mathbb{G}_m^n -invariant closed 2-form ω_X on X can be given by the image of the (-1) -shifted closed 2-form on $[X/\mathbb{G}_m^n]$ along $\mathcal{A}^{2, \text{cl}}([X/\mathbb{G}_m^n], -1) \rightarrow \mathcal{A}^{2, \text{cl}}([X/\mathbb{G}_m^n]/B\mathbb{G}_m^n, -1)$.

By [Par24, Lemma 4.2.5], the closed 2-form ω_X is symplectic and the isotropic correspondence

$$\begin{array}{ccc} & \widehat{X} & \\ \swarrow & & \searrow \\ \mathfrak{X} & & X. \end{array}$$

is Lagrangian, after shrinking \widehat{X} . □

We now construct (-1) -shifted symplectic analogue of isomorphisms in Proposition 6.8. Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be (possibly higher) (-1) -shifted symplectic stack. Firstly, note that the natural equivalence

$$(6.54) \quad \cdot \omega_{\mathfrak{X}} : \mathbb{T}_{\mathfrak{X}}^+[1] \simeq \mathbb{L}_{\mathfrak{X}}^+$$

combined with the natural isomorphisms (2.14) and (2.5) implies an isomorphism

$$(6.55) \quad \phi : \widehat{\det}_-(\mathbb{L}_{\mathfrak{X}}) \cong \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}}).$$

We define an isomorphism

$$\xi_{\mathfrak{X}}^{\text{der}} : \widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}})^{\otimes 2} \cong \widehat{\det}(\mathbb{L}_{\mathfrak{X}}|_{\text{Grad}(\mathfrak{X})})$$

by the following composition

$$\begin{aligned} & \widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}})^{\otimes 2} \cong \widehat{\det}_0(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}})^{\otimes 2} \\ & \xrightarrow[\cong]{\text{id} \otimes \text{id} \otimes \phi^{-1}} \widehat{\det}_0(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_-(\mathbb{L}_{\mathfrak{X}}) \cong \widehat{\det}(\mathbb{L}_{\mathfrak{X}}|_{\text{Grad}(\mathfrak{X})}). \end{aligned}$$

Let $x \in \text{Grad}(\mathfrak{X})$ be a point. We now show that the map $\xi_{\mathfrak{X}}^{\text{der}}$ is compatible with the map $\kappa_x^{\text{Ind, der}}$, namely, the following diagram commutes:

$$(6.56) \quad \begin{array}{ccc} \det(\mathbb{L}_{\text{Grad}(\mathfrak{X}), x}) \otimes \det_+(\mathbb{L}_{\mathfrak{X}, u(x)})^{\otimes 2} & \xrightarrow[\cong]{\xi_{\mathfrak{X}}^{\text{der}}} & \det(\mathbb{L}_{\mathfrak{X}, u(x)}) \\ \cong \downarrow (\kappa_x^{\text{der}}, \kappa_x^{\text{Ind, der}, \otimes 2}) & & \cong \downarrow \kappa_x \\ \det_0(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)}))^{\otimes 2} \otimes \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)}))^{\otimes 2} & \xrightarrow[\cong]{} & \det(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)}))^{\otimes 2}. \end{array}$$

To see this, we construct an isomorphism

$$\kappa_x^{\text{Ind,der,-}} : \det_-(\mathbb{L}_{\mathfrak{X},u(x)}) \cong \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)}))$$

in the same manner as the construction of $\kappa_x^{\text{Ind,der}}$ in (6.42). Then the following diagram commutes:

$$(6.57) \quad \begin{array}{ccc} \det_-(\mathbb{L}_{\mathfrak{X},u(x)}) & \xrightarrow[\cong]{(-1)^{\text{rank}_+ \tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x))} \cdot \phi|_x}} & \det_+(\mathbb{L}_{\mathfrak{X},u(x)}) \\ \cong \downarrow \kappa_x^{\text{Ind,der,-}} & & \cong \downarrow \kappa_x^{\text{Ind,der}} \\ \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) & \xlongequal{\hspace{2cm}} & \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})). \end{array}$$

This follows from the commutativity of the diagrams (2.15) and (2.18). We also see that the following diagram commutes by construction:

$$(6.58) \quad \begin{array}{ccc} \det_0(\mathbb{L}_{\mathfrak{X},u(x)}) \otimes \widehat{\det}_-(\mathbb{L}_{\mathfrak{X},u(x)}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X},u(x)}) & \xrightarrow[\cong]{\xi_{\mathfrak{X}}^{\text{der}}} & \det(\mathbb{L}_{\mathfrak{X},u(x)}) \\ \cong \downarrow (\kappa_x^{\text{der}}, \kappa_x^{\text{Ind,der,-}}, \kappa_x^{\text{Ind,der}}) & & \downarrow \cong \kappa_{u(x)} \\ \det_0(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)}))^{\otimes 2} \otimes \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) \otimes \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) & & \downarrow \cong \\ \downarrow \cong & & \downarrow \cong \\ \det_{\leq 0}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) \otimes \det_{\geq 0}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) \otimes \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) & \xrightarrow[\cong]{} & \det(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)}))^{\otimes 2}. \end{array}$$

By combining the commutativity of the diagram (6.57) and (6.58) and the fact that the symmetrizer of the complex $\det_+(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)}))^{\otimes 2}$ is given by multiplying $(-1)^{\text{rank}_+ \tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})}$, we conclude that the diagram (6.56) commutes.

Now we construct a (-1) -shifted symplectic analogue of θ_s in Proposition 6.8. For a subset $\Sigma \subset \mathbb{Z}^2$, using the natural equivalence $\mathbb{T}_{\mathfrak{X}}^{\Sigma}[1] \simeq \mathbb{L}_{\mathfrak{X}}^{\Sigma}$ and (2.5) and (2.14), we see that there exists a natural isomorphism

$$\phi_{\Sigma} : \widehat{\det}_{-\Sigma}(\mathbb{L}_{\mathfrak{X}}) \cong \widehat{\det}_{\Sigma}(\mathbb{L}_{\mathfrak{X}}).$$

Further, for each point $x \in \text{Grad}^2(\mathfrak{X})$, the commutativity of the diagrams (2.15) and (2.18) implies that the following diagram commutes:

$$(6.59) \quad \begin{array}{ccc} \widehat{\det}_{-\Sigma}(\mathbb{L}_{\mathfrak{X}})|_x & \xrightarrow[\cong]{(-1)^{\text{rank}_{\Sigma} \tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x))} \cdot \phi_{\Sigma}|_x}} & \widehat{\det}_{\Sigma}(\mathbb{L}_{\mathfrak{X}})|_x \\ \cong \downarrow \kappa_x^{\text{Ind,der,-}\Sigma} & & \cong \downarrow \kappa_x^{\text{Ind,der},\Sigma} \\ \widehat{\det}_{\pm\Sigma}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) & \xlongequal{\hspace{2cm}} & \widehat{\det}_{\pm\Sigma}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X},u(x)})) \end{array}$$

where the vertical maps are constructed in the same manner as $\kappa_x^{\text{Ind,der}}$. Set

$$S_1 := \{(m, l) \mid (m, l) \neq (0, 0), m \geq 0, l \geq 0\}, \quad S_2 := \mathbb{Z}_{>0} \times \mathbb{Z}_{<0}$$

We construct an isomorphism

$$\theta_{\mathfrak{X}}^{\text{der}} : \widehat{\det}_{\mathbb{Z}_{>0} \times \mathbb{Z}}(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_{\{0\} \times \mathbb{Z}_{>0}}(\mathbb{L}_{\mathfrak{X}}) \cong \widehat{\det}_{\mathbb{Z} \times \mathbb{Z}_{>0}}(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_{\mathbb{Z}_{>0} \times \{0\}}(\mathbb{L}_{\mathfrak{X}}).$$

using the decompositions

$$(\mathbb{Z}_{>0} \times \mathbb{Z}) \coprod (\{0\} \times \mathbb{Z}_{>0}) = S_1 \coprod S_2, \quad (\mathbb{Z} \times \mathbb{Z}_{>0}) \coprod (\mathbb{Z}_{>0} \times \{0\}) = S_2 \coprod (-S_2)$$

and the map $\phi_{S_2}^{-1}$. By the commutativity of the diagram (6.59), we see that $\theta_{\mathfrak{X}}^{\text{der}}$ is compatible with $\kappa_x^{\text{Ind,der}}$, namely, the following diagram commutes:

$$(6.60) \quad \begin{array}{ccc} \widehat{\det}_{\mathbb{Z}_{>0} \times \mathbb{Z}}(\mathbb{L}_{\mathfrak{X}, u(x)}) \otimes \widehat{\det}_{\{0\} \times \mathbb{Z}_{>0}}(\mathbb{L}_{\mathfrak{X}, u(x)}) & \xrightarrow[\cong]{ (-1)^{\text{rank}_{S_2}} \tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)}) \cdot \theta_{\mathfrak{X}}^{\text{der}} } & \widehat{\det}_{\mathbb{Z} \times \mathbb{Z}_{>0}}(\mathbb{L}_{\mathfrak{X}, u(x)}) \otimes \widehat{\det}_{\mathbb{Z}_{>0} \times \{0\}}(\mathbb{L}_{\mathfrak{X}, u(x)}) \\ \cong \downarrow (\kappa_x^{\text{Ind,der}, \mathbb{Z}_{>0} \times \mathbb{Z}}, \kappa_x^{\text{Ind,der}, \{0\} \times \mathbb{Z}_{>0}}) & & \cong \downarrow (\kappa_x^{\text{Ind,der}, \mathbb{Z} \times \mathbb{Z}_{>0}}, \kappa_x^{\text{Ind,der}, \mathbb{Z}_{>0} \times \{0\}}) \\ \widehat{\det}_{\mathbb{Z}_{\neq 0} \times \mathbb{Z}}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) \otimes \widehat{\det}_{\{0\} \times \mathbb{Z}_{\neq 0}}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) & & \widehat{\det}_{\mathbb{Z} \times \mathbb{Z}_{\neq 0}}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) \otimes \widehat{\det}_{\mathbb{Z}_{\neq 0} \times \{0\}}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) \\ & \searrow \cong & \swarrow \cong \\ & \widehat{\det}_{\mathbb{Z}_{\neq (0,0)}^2}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) & \end{array}$$

We now construct a (-1) -shifted symplectic version of $\Psi_{\mathfrak{X}, \mathfrak{Y}}$ in Proposition 6.8. Assume that we are given (-1) -shifted symplectic derived Artin stacks $(\mathfrak{X}, \omega_{\mathfrak{X}})$ and $(\mathfrak{Y}, \omega_{\mathfrak{Y}})$. Then we construct isomorphisms

$$\begin{aligned} \Psi_{\mathfrak{X}, \mathfrak{Y}}^{\text{der}} &: \widehat{\det}(\mathbb{L}_{\mathfrak{X}}) \boxtimes \widehat{\det}(\mathbb{L}_{\mathfrak{Y}}) \cong \widehat{\det}(\mathbb{L}_{\mathfrak{X} \times \mathfrak{Y}}) \\ \Psi_{\mathfrak{X}, \mathfrak{Y}}^{\text{Ind,der}} &: \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}}) \boxtimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{Y}}) \cong \widehat{\det}_+(\mathbb{L}_{\mathfrak{X} \times \mathfrak{Y}}) \end{aligned}$$

using the natural equivalences $\mathbb{L}_{\mathfrak{X}} \boxplus \mathbb{L}_{\mathfrak{Y}} \simeq \mathbb{L}_{\mathfrak{X} \times \mathfrak{Y}}$ and $\mathbb{L}_{\mathfrak{X}}|_{\text{Grad}(\mathfrak{X})}^+ \boxplus \mathbb{L}_{\mathfrak{Y}}|_{\text{Grad}(\mathfrak{Y})}^+ \simeq \mathbb{L}_{\mathfrak{X} \times \mathfrak{Y}}|_{\text{Grad}(\mathfrak{X}) \times \text{Grad}(\mathfrak{Y})}^+$ respectively.

We now prove the compatibility between the morphism $\Lambda_{\mathfrak{X}}^{\text{Ind}}$ and the morphisms constructed in Proposition 6.8. Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a quasi-separated (-1) -shifted symplectic stack with affine stabilizers and $(\mathfrak{X}^{\text{cl}}, s)$ be its underlying d-critical stack. By construction, $\Lambda_{\mathfrak{X}}^{\text{Ind}}$ commutes with κ_x^{Ind} : Namely, the following diagram commutes:

$$(6.61) \quad \begin{array}{ccc} \det_+(\mathbb{L}_{\mathfrak{X}, u(x)}) & \xrightarrow[\cong]{ \Lambda_{\mathfrak{X}}^{\text{Ind}}|_x } & \mathcal{L}_s^{\text{Ind}}|_x \\ \cong \downarrow \kappa_x^{\text{Ind,der}} & & \cong \downarrow \kappa_x^{\text{Ind}} \\ \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})) & \xlongequal{\quad} & \det_{\pm}(\tau^{\geq 0}(\mathbb{L}_{\mathfrak{X}, u(x)})). \end{array}$$

Next, we will show that $\xi_{\mathfrak{X}}$ is compatible with $\Lambda_{\mathfrak{X}}^{\text{Ind}}$. Namely, we will see that the following diagram commutes:

$$(6.62) \quad \begin{array}{ccc} \widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}})^{\otimes 2} & \xrightarrow[\cong]{ \xi_{\mathfrak{X}}^{\text{der}} } & \widehat{\det}(\mathbb{L}_{\mathfrak{X}}) \\ \cong \downarrow (\Lambda_{\text{Grad}(\mathfrak{X})}, \Lambda_{\mathfrak{X}}^{\text{Ind}, \otimes 2}) & & \cong \downarrow \Lambda_{\mathfrak{X}} \\ K_{\text{Grad}(\mathfrak{X}), u^* s} \otimes \mathcal{L}_s^{\text{Ind}, \otimes 2} & \xrightarrow[\cong]{ \xi_s } & K_{\mathfrak{X}, s}|_{\text{Grad}(\mathfrak{X})}. \end{array}$$

This is a direct consequence of the commutativity of diagrams (3.25), (6.35), (6.56) and (6.61). We now show that $\theta_{\mathfrak{X}}^{\text{der}}$ is compatible with $\Lambda_{\mathfrak{X}}^{\text{Ind}}$. Namely, the following diagram commutes:

$$(6.63) \quad \begin{array}{ccc} \widehat{\det}_{\mathbb{Z}_{>0} \times \mathbb{Z}}(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_{\{0\} \times \mathbb{Z}_{>0}}(\mathbb{L}_{\mathfrak{X}}) & \xrightarrow[\cong]{ \theta_{\mathfrak{X}}^{\text{der}} } & \widehat{\det}_{\mathbb{Z} \times \mathbb{Z}_{>0}}(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_{\mathbb{Z}_{>0} \times \{0\}}(\mathbb{L}_{\mathfrak{X}}) \\ \cong \downarrow (\Lambda_{\mathfrak{X}}^{\text{Ind}}, \Lambda_{\text{Grad}(\mathfrak{X})}^{\text{Ind}}) & & \cong \downarrow (\Lambda_{\mathfrak{X}}^{\text{Ind}}, \Lambda_{\text{Grad}(\mathfrak{X})}^{\text{Ind}}) \\ u_1^{\text{red},*} \mathcal{L}_s^{\text{Ind}} \otimes \sigma^* \mathcal{L}_{u^* s}^{\text{Ind}} & \xrightarrow[\cong]{ \theta_s } & u_2^{\text{red},*} \mathcal{L}_s^{\text{Ind}} \otimes \mathcal{L}_{u^* s}^{\text{Ind}}. \end{array}$$

This is a direct consequence of the commutativity of diagrams (6.37) and (6.60). Finally, if we are given a quasi-separated (-1) -shifted symplectic $(\mathfrak{Y}, \omega_{\mathfrak{Y}})$ with affine stabilizers with classical d-critical stack $(\mathfrak{Y}^{\text{cl}}, t)$,

it is clear that the following diagram commutes:

$$(6.64) \quad \begin{array}{ccc} \widehat{\det}_+(\mathbb{L}\mathfrak{X}) \boxtimes \widehat{\det}_+(\mathbb{L}\mathfrak{Y}) & \xrightarrow[\cong]{\Psi_{\mathfrak{X},\mathfrak{Y}}^{\text{Ind,der}}} & \widehat{\det}_+(\mathbb{L}\mathfrak{X} \times \mathfrak{Y}) \\ \Lambda_{\mathfrak{X}}^{\text{Ind}} \boxtimes \Lambda_{\mathfrak{Y}}^{\text{Ind}} \downarrow \cong & & \Lambda_{\mathfrak{X} \times \mathfrak{Y}}^{\text{Ind}} \downarrow \cong \\ \mathcal{L}_s^{\text{Ind}} \boxtimes \mathcal{L}_t^{\text{Ind,der}} & \xrightarrow[\cong]{\Psi_{\mathfrak{X}^{\text{cl}},\mathfrak{Y}^{\text{cl}}}^{\text{Ind}}} & \mathcal{L}_{s \boxplus t}^{\text{Ind}} \end{array}$$

We now show that $\theta_{\mathfrak{X}}^{\text{der}}$ is compatible with $\xi_{\mathfrak{X}}^{\text{der}}$. Namely, the following diagram commutes:

$$(6.65) \quad \begin{array}{ccc} \widehat{\det}(\mathbb{L}_{\text{Grad}^2(\mathfrak{X})}) & \xrightarrow[\cong]{(\theta_{\mathfrak{X}}^{\text{der}})^{\otimes 2}} & \widehat{\det}(\mathbb{L}_{\text{Grad}^2(\mathfrak{X})}) \\ \otimes \widehat{\det}_{\mathbb{Z}_{>0} \times \mathbb{Z}}(\mathbb{L}\mathfrak{X})^{\otimes 2} \otimes \widehat{\det}_{\{0\} \times \mathbb{Z}_{>0}}(\mathbb{L}\mathfrak{X})^{\otimes 2} & & \otimes \widehat{\det}_{\mathbb{Z} \times \mathbb{Z}_{>0}}(\mathbb{L}\mathfrak{X})^{\otimes 2} \otimes \widehat{\det}_{\mathbb{Z}_{>0} \times \{0\}}(\mathbb{L}\mathfrak{X})^{\otimes 2} \\ \cong \downarrow \xi_{\text{Grad}(\mathfrak{X})}^{\text{der}} & & \cong \downarrow \xi_{\text{Grad}(\mathfrak{X})}^{\text{der}} \\ u_1^* \widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes \widehat{\det}_{\mathbb{Z}_{>0} \times \mathbb{Z}}(\mathbb{L}\mathfrak{X})^{\otimes 2} & & u_2^* \widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes \widehat{\det}_{\mathbb{Z} \times \mathbb{Z}_{>0}}(\mathbb{L}\mathfrak{X})^{\otimes 2} \\ & \xrightarrow[\cong]{u_1^* \xi_{\mathfrak{X}}^{\text{der}}} & \xrightarrow[\cong]{u_2^* \xi_{\mathfrak{X}}^{\text{der}}} \\ & \searrow & \swarrow \\ & \widehat{\det}(\mathbb{L}\mathfrak{X}) & \end{array}$$

This is an immediate consequence of the commutativity of diagrams (6.60) and (6.56). Next, we note that the map $\xi_{\mathfrak{X}}^{\text{der}}$ is compatible with $\Psi_{\mathfrak{X},\mathfrak{Y}}^{\text{Ind,der}}$:

$$(6.66) \quad \begin{array}{ccc} (\widehat{\det}_0(\mathbb{L}\mathfrak{X}) \otimes \widehat{\det}_+(\mathbb{L}\mathfrak{X})^{\otimes 2}) \boxtimes (\widehat{\det}_0(\mathbb{L}\mathfrak{Y}) \otimes \widehat{\det}_+(\mathbb{L}\mathfrak{Y})^{\otimes 2}) & \xrightarrow[\cong]{\Psi_{\text{Grad}(\mathfrak{X}),\text{Grad}(\mathfrak{Y})}^{\text{der}}, \Psi_{\mathfrak{X},\mathfrak{Y}}^{\text{Ind,der}}} & \widehat{\det}_0(\mathbb{L}\mathfrak{X} \times \mathfrak{Y}) \otimes \widehat{\det}_+(\mathbb{L}\mathfrak{X} \times \mathfrak{Y})^{\otimes 2} \\ \xi_{\mathfrak{X}}^{\text{der}} \boxtimes \xi_{\mathfrak{Y}}^{\text{der}} \downarrow \cong & & \xi_{\mathfrak{X} \times \mathfrak{Y}}^{\text{der}} \downarrow \cong \\ \widehat{\det}(\mathbb{L}\mathfrak{X} |_{\text{Grad}(\mathfrak{X})}) \boxtimes \widehat{\det}(\mathbb{L}\mathfrak{Y} |_{\text{Grad}(\mathfrak{Y})}) & \xrightarrow[\cong]{\Psi_{\mathfrak{X},\mathfrak{Y}}^{\text{der}}} & \widehat{\det}(\mathbb{L}\mathfrak{X} \times \mathfrak{Y} |_{\text{Grad}(\mathfrak{X} \times \mathfrak{Y})}). \end{array}$$

This is obvious from the construction.

Next, we will compare $\xi_{\mathfrak{X}}^{\text{der}}$ with the map ξ constructed in (3.5). Let $(\mathfrak{X}, \omega_{\mathfrak{X}})$ be a higher (-1) -shifted symplectic stack. Recall first that by Proposition 5.10 we have a natural equivalence

$$\mathbb{T}_{\text{Grad}(\mathfrak{X})} \simeq \mathbb{T}_{\mathfrak{X}}^0, \quad \mathbb{T}_{\text{Filt}(\mathfrak{X})/\text{Grad}(\mathfrak{X})|_{\text{Grad}(\mathfrak{X})}} \simeq \mathbb{T}_{\mathfrak{X}}^+.$$

The fibre sequence $\mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}[-1] \rightarrow \mathbb{L}\mathfrak{X} \rightarrow \mathbb{L}_{\text{Filt}(\mathfrak{X})}$ implies a natural equivalence

$$\mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}[-1]|_{\text{Grad}(\mathfrak{X})} \simeq \mathbb{L}\mathfrak{X}|_{\text{Grad}(\mathfrak{X})}^+.$$

In particular, we have an isomorphism

$$(6.67) \quad \widehat{\det}(\mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}|_{\text{Grad}(\mathfrak{X})})^{\vee} \cong \widehat{\det}_+(\mathbb{L}\mathfrak{X}).$$

Consider the following map of fibre sequences

$$\begin{array}{ccccc} \mathbb{T}_{\text{Filt}(\mathfrak{X})/\text{Grad}(\mathfrak{X})}[1]|_{\text{Grad}(\mathfrak{X})} & \longrightarrow & \mathbb{T}_{\text{Filt}(\mathfrak{X})}[1]|_{\text{Grad}(\mathfrak{X})} & \longrightarrow & \mathbb{T}_{\text{Grad}(\mathfrak{X})}[1]|_{\text{Grad}(\mathfrak{X})} \\ \downarrow \cong & & \downarrow & & \downarrow \\ \mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}[-1] & \longrightarrow & \mathbb{L}\mathfrak{X}|_{\text{Grad}(\mathfrak{X})} & \longrightarrow & \mathbb{L}_{\text{Filt}(\mathfrak{X})}|_{\text{Grad}(\mathfrak{X})} \end{array}$$

where the homotopy commutativity of the right diagram is induced from the Lagrangian structure of the attractor correspondence. Then the following diagram commutes by construction:

$$(6.68) \quad \begin{array}{ccc} \mathbb{T}_{\text{Filt}(\mathfrak{X})/\text{Grad}(\mathfrak{X})}[1]|_{\text{Grad}(\mathfrak{X})} & \xrightarrow{\cong} & \mathbb{T}_{\mathfrak{X}}^+[1] \\ (3.3) \downarrow \cong & & (6.54) \downarrow \cong \\ \mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}[-1]|_{\text{Grad}(\mathfrak{X})} & \xrightarrow{\cong} & \mathbb{L}_{\mathfrak{X}}^+. \end{array}$$

We prove the following statement:

Lemma 6.11. *The following diagram commutes:*

$$\begin{array}{ccc}
\widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes (\widehat{\det}(\mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}|\text{Grad}(\mathfrak{X}))^\vee)^{\otimes 2} & \xrightarrow[\cong]{(3.5)} & \widehat{\det}(\mathbb{L}_{\mathfrak{X}}) \\
\downarrow \cong & & \parallel \\
\widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}})^{\otimes 2} & \xrightarrow[\cong]{\xi_{\mathfrak{X}}^{\text{der}}} & \widehat{\det}(\mathbb{L}_{\mathfrak{X}}).
\end{array}$$

Proof. By the construction of $\xi_{\mathfrak{X}}^{\text{der}}$, the statement is equivalent to the commutativity of the following diagram:

$$\begin{array}{ccc}
\widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes (\widehat{\det}(\mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}|\text{Grad}(\mathfrak{X}))^\vee)^{\otimes 2} & \xrightarrow[\cong]{(3.5)} & \widehat{\det}(\mathbb{L}_{\mathfrak{X}}) \\
\downarrow \cong & & \uparrow \cong \\
\widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}})^{\otimes 2} & \xrightarrow[\cong]{\text{id} \otimes \text{id} \otimes \phi^{-1}} & \widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_-(\mathbb{L}_{\mathfrak{X}}).
\end{array}$$

Firstly, consider the following diagram:

$$\begin{array}{ccc}
\widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes \widehat{\det}(\mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}|\text{Grad}(\mathfrak{X}))^\vee & \xrightarrow[\cong]{(6.67)} & \widehat{\det}_0(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}}) \\
\cong \downarrow \text{id} \otimes (3.4) & & \cong \downarrow \text{id} \otimes \phi^{-1} \\
(6.69) \quad \widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}) \otimes \widehat{\det}(\mathbb{L}_{\text{Filt}(\mathfrak{X})/\text{Grad}(\mathfrak{X})}|\text{Grad}(\mathfrak{X})) & & \widehat{\det}_0(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_-(\mathbb{L}_{\mathfrak{X}}) \\
\cong \downarrow & & \downarrow \cong \\
\widehat{\det}(\mathbb{L}_{\text{Filt}(\mathfrak{X})}|\text{Grad}(\mathfrak{X})) & \xrightarrow[\cong]{} & \widehat{\det}_{\leq 0}(\mathbb{L}_{\mathfrak{X}}).
\end{array}$$

This immediately follows from the commutativity of the diagram (6.68). Next, we will show that the following diagram commutes up to sign $(-1)^{\text{rank } \mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}|\text{Grad}(\mathfrak{X})}$:

$$\begin{array}{ccc}
\widehat{\det}(\mathbb{L}_{\text{Filt}(\mathfrak{X})}|\text{Grad}(\mathfrak{X})) \otimes \widehat{\det}(\mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}|\text{Grad}(\mathfrak{X}))^\vee & \xrightarrow[\cong]{(6.67)} & \widehat{\det}_{\leq 0}(\mathbb{L}_{\mathfrak{X}}) \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}}) \\
\cong \downarrow & & \cong \downarrow \\
(6.70) \quad \widehat{\det}(\mathbb{L}_{\mathfrak{X}}|\text{Grad}(\mathfrak{X})) & \xlongequal{\quad\quad\quad} & \widehat{\det}(\mathbb{L}_{\mathfrak{X}}|\text{Grad}(\mathfrak{X})).
\end{array}$$

This follows from the commutativity of (2.8) and the following equivalence of fibre sequences:

$$\begin{array}{ccccc}
\mathbb{L}_{\mathfrak{X}}|\text{Grad}(\mathfrak{X}) & \longrightarrow & \mathbb{L}_{\text{Filt}(\mathfrak{X})}|\text{Grad}(\mathfrak{X}) & \longrightarrow & \mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}|\text{Grad}(\mathfrak{X}) \\
\downarrow \cong & & \downarrow \cong & & \downarrow \cong \\
\mathbb{L}_{\mathfrak{X}}|\text{Grad}(\mathfrak{X}) & \longrightarrow & \mathbb{L}_{\mathfrak{X}}|\text{Grad}(\mathfrak{X})^{\leq 0} & \longrightarrow & \mathbb{L}_{\mathfrak{X}}[1]|\text{Grad}(\mathfrak{X})^+.
\end{array}$$

The statement of the lemma immediately follows from the commutativity of diagrams (6.69) and (6.70) and the fact that the symmetrizer of $\widehat{\det}_+(\mathbb{L}_{\mathfrak{X}})^{\otimes 2}$ is the multiplication by $(-1)^{\text{rank } \mathbb{L}_{\text{Filt}(\mathfrak{X})/\mathfrak{X}}|\text{Grad}(\mathfrak{X})}$. \square

6.5. Localizing orientations. Here we discuss the localization of an orientation for \mathbb{G}_m -equivariant algebraic spaces and d-critical stacks.

Assume that we are given a quasi-separated \mathbb{G}_m -equivariant d-critical algebraic space (X, μ, s) . We define an isomorphism

$$\bar{\xi}_{\mu, s} : K_{X, s}|_{X^{\mu, \text{red}}} \otimes (\mathcal{L}_{\mu, s}^{\text{Ind}, \vee})^{\otimes 2} \cong K_{X^{\mu, s^{\mu}}}$$

by the following composition:

$$\begin{aligned} K_{X,s}|_{X^{\mu,\text{red}}} \otimes (\mathcal{L}_{\mu,s}^{\text{Ind},\vee})^{\otimes 2} &\xrightarrow[\cong]{\xi_{\mu,s} \otimes \text{id}} K_{X^{\mu},s^{\mu}} \otimes (\mathcal{L}_{\mu,s}^{\text{Ind}})^{\otimes 2} \otimes (\mathcal{L}_{\mu,s}^{\text{Ind},\vee})^{\otimes 2} \\ &\cong K_{X^{\mu},s^{\mu}} \otimes (\mathcal{L}_{\mu,s}^{\text{Ind}})^{\otimes 2} \otimes (\mathcal{L}_{\mu,s}^{\text{Ind},\otimes 2})^{\vee} \cong K_{X^{\mu},s^{\mu}}. \end{aligned}$$

Note that the isomorphism $(\mathcal{L}_{\mu,s}^{\text{Ind},\vee})^{\otimes 2} \cong (\mathcal{L}_{\mu,s}^{\text{Ind},\otimes 2})^{\vee}$ differs from the natural isomorphism on underlying ungraded line bundles by the sign $(-1)^{I_X^{\mu}}$, which is the symmetrizer of the graded line bundle $\mathcal{L}_{\mu,s}^{\text{Ind},\otimes 2}$ of our symmetric monoidal structure on graded line bundles. Assume now that we are given an orientation (\mathcal{L}, o) for (X, s) . We define an orientation o^{μ} for (X^{μ}, s^{μ}) by

$$(6.71) \quad o^{\mu}: (\mathcal{L}|_{X^{\mu,\text{red}}} \otimes \mathcal{L}_{\mu,s}^{\text{Ind},\vee})^{\otimes 2} \xrightarrow[\cong]{o \otimes \text{id}} K_{X,s}|_{X^{\mu,\text{red}}} \otimes (\mathcal{L}_{\mu,s}^{\text{Ind},\vee})^{\otimes 2} \xrightarrow[\cong]{\bar{\xi}_{\mu,s}} K_{X^{\mu},s^{\mu}}.$$

Let (X_1, μ_1, s_1) and (X_2, μ_2, s_2) be quasi-separated \mathbb{G}_m -equivariant d-critical algebraic spaces. Let $h: X_1 \rightarrow X_2$ be a smooth morphism such that $h^*s_2 = s_1$ holds. Assume that we are given an orientation o_2 for (X_2, s_2) . Then the commutativity of the diagram (6.27) implies that there exists a canonical isomorphism of orientations

$$(6.72) \quad (h^*o_2)^{\mu_1} \cong h^{\mathbb{G}_m,*} o_2^{\mu_2}.$$

Assume that we are given a quasi-separated \mathbb{G}_m^2 -equivariant d-critical algebraic space $(X, (\mu_1, \mu_2), s)$ and an orientation o . Then the commutativity of the diagram (6.28) implies that there exists a natural isomorphism of orientations

$$(6.73) \quad (o^{\mu_1})^{\mu_2} \cong (o^{\mu_2})^{\mu_1}.$$

Let (X, μ, s) and (Y, ν, t) be quasi-separated \mathbb{G}_m -equivariant d-critical algebraic spaces with orientations o_X and o_Y . Then using that $\xi_{\mu,s} \boxtimes \xi_{\nu,t}$ and $\xi_{\mu \times \nu, s \boxplus t}$ are equivalent after the identification $\Psi_{X,Y}$ and $\Psi_{\mu,\nu}^{\text{Ind}}$, we obtain a natural isomorphism

$$(6.74) \quad o_X^{\mu} \boxtimes o_Y^{\nu} \cong o_{X \times Y}^{\mu \times \nu}.$$

Now we discuss localization of canonical orientation (3.33). Let U be a smooth quasi-separated algebraic space with a \mathbb{G}_m -action μ_U and f be a \mathbb{G}_m -invariant regular function on U with $f|_{U^{\text{red}}} = 0$. Set $X = \text{Crit}(f)^{\text{cl}}$ and equip it with the \mathbb{G}_m -action μ induced by μ_U and the natural d-critical structure s . Then (X, μ, s) is a \mathbb{G}_m -equivariant d-critical algebraic space. Recall that we have defined the canonical orientation o_X^{can} in (3.33). Since we have

$$X^{\mu} \cong \text{Crit}(f|_{U^{\mu}})^{\text{cl}},$$

d-critical structure s^{μ} coincides with the one induced by the critical locus description, hence one can define the canonical orientation $o_{X^{\mu}}^{\text{can}}$ for (X^{μ}, s^{μ}) . Then the natural isomorphism $K_U \otimes K_U^{\pm, \vee} \cong K_{U^{\mu}}$ induces an isomorphism of orientations

$$(6.75) \quad (o_X^{\text{can}})^{\mu} \cong o_{X^{\mu}}^{\text{can}}.$$

This can be proved in the same manner as the commutativity of the diagram (6.12): see also Remark 3.36.

Assume that we are given \mathbb{G}_m -equivariant smooth morphism of smooth quasi-separated \mathbb{G}_m -equivariant algebraic spaces $H: U_1 \rightarrow U_2$ and a \mathbb{G}_m -invariant regular function $f_2: U_2 \rightarrow \mathbb{A}^1$ with $f_2|_{U_2^{\text{red}}} = 0$. Set $f_1 := f_2 \circ H$ and $h: X_1 \rightarrow X_2$ be the natural morphism induced on the critical locus. Let μ_1 and μ_2 be the \mathbb{G}_m -actions on X_1 and X_2 . Then the following diagram commutes by the construction:

$$(6.76) \quad \begin{array}{ccc} (h^*o_{X_2}^{\text{can}})^{\mu_1} & \xrightarrow[\cong]{(3.35)} & (o_{X_1}^{\text{can}})^{\mu_1} \\ \cong \downarrow (6.72) & & \downarrow \cong (6.75) \\ h^{\mathbb{G}_m,*}(o_{X_2}^{\text{can}})^{\mu_2} & & o_{X_1}^{\text{can}} \\ \cong \downarrow (6.75) & & \\ h^{\mathbb{G}_m,*}o_{X_2}^{\text{can}} & \xrightarrow[\cong]{(3.35)} & o_{X_1}^{\text{can}}. \end{array}$$

Now assume that U is a smooth quasi-separated algebraic space with a \mathbb{G}_m^2 -action $(\mu_{U,1}, \mu_{U,2})$ and f be a \mathbb{G}_m^2 -invariant regular function on U with $f|_{U^{\text{red}}} = 0$. Let $(X, (\mu_1, \mu_2), s)$ be the associated \mathbb{G}_m^2 -equivariant d-critical algebraic space. Then by repeating the proof of the commutativity of (6.18), we see that the following diagram commutes:

$$(6.77) \quad \begin{array}{ccccc} ((o_X^{\text{can}})^{\mu_1})^{\mu_2} |_{X_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{(6.75)} & (o_{X^{\mu_1}}^{\text{can}})^{\mu_2} |_{X_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{(6.75)} & o_{X^{\hat{\mu}}}^{\text{can}} |_{X_{\text{mat}}^{\hat{\mu}}} \\ \downarrow \cong & & & & \parallel \\ ((o_X^{\text{can}})^{\mu_2})^{\mu_1} |_{X_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{(6.75)} & (o_{X^{\mu_2}}^{\text{can}})^{\mu_1} |_{X_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{(6.75)} & o_{X^{\hat{\mu}}}^{\text{can}} |_{X_{\text{mat}}^{\hat{\mu}}} \end{array}$$

Now assume that we are given \mathbb{G}_m -equivariant smooth quasi-separated algebraic spaces (U, μ_U) and (V, ν_V) with \mathbb{G}_m -invariant regular functions f and g such that $f|_{U^{\text{red}}} = 0$ and $g|_{V^{\text{red}}} = 0$. Let (X, μ, s) and (Y, ν, t) be the associated \mathbb{G}_m -equivariant d-critical algebraic spaces. Then by repeating the proof of the commutativity of (6.20), we see that the following diagram commutes:

$$(6.78) \quad \begin{array}{ccc} (o_X^{\text{can}} \boxtimes o_Y^{\text{can}})^{\mu \times \nu} & \xrightarrow[\cong]{(6.74)} & (o_X^{\text{can}})^{\mu} \boxtimes (o_Y^{\text{can}})^{\nu} & \xrightarrow[\cong]{(6.75)} & o_{X^{\mu}}^{\text{can}} \boxtimes o_{Y^{\nu}}^{\text{can}} \\ \cong \downarrow (3.37) & & & & \cong \downarrow (3.37) \\ (o_{X \times Y}^{\text{can}})^{\mu \times \nu} & \xrightarrow[\cong]{(6.75)} & & & o_{X^{\mu} \times Y^{\nu}}^{\text{can}} \end{array}$$

We now explain the localization of orientations for d-critical stacks. Let (\mathfrak{X}, s) be a d-critical stack and $o: \mathcal{L}^{\otimes 2} \cong K_{\mathfrak{X}, s}$ be an orientation. We define the localized orientation u^*o on $(\text{Grad}(\mathfrak{X}), u^*s)$ by the following composition

$$u^*o: (u^{\text{red},*} \mathcal{L} \otimes \mathcal{L}_s^{\text{Ind}, \vee})^{\otimes 2} \xrightarrow{\cong} K_{\mathfrak{X}, s} \otimes (\mathcal{L}_s^{\text{Ind}, \vee})^{\otimes 2} \xrightarrow[\cong]{\xi_s} K_{\text{Grad}(\mathfrak{X}), u^*s}.$$

Assume that we are given a \mathbb{G}_m -equivariant chart (X, μ, \bar{q}) . Then the commutativity of the diagram (6.38) implies a canonical isomorphism of orientations:

$$(6.79) \quad (q^*o)^{\mu} \cong q_{\mu}^* u^*o.$$

The commutativity of the diagram (6.39) implies a canonical isomorphism of orientations:

$$(6.80) \quad u_1^* u^*o \cong u_2^* u^*o.$$

Here, u_2^*o is defined using the natural identification $\text{Grad}^2(\mathfrak{X}) \cong \text{Grad}(\text{Grad}(\mathfrak{X}))$ and we define $u_1^* := \sigma^* u_2^*$. Assume further that we are given a \mathbb{G}_m^2 -equivariant chart $(X, \hat{\mu} = (\mu_1, \mu_2), \bar{q})$. Set $\hat{\mu}' := (\mu_2, \mu_1)$. Then the commutativity of the diagram (6.41) implies that the following diagram commutes:

$$(6.81) \quad \begin{array}{ccc} ((q^*o)^{\mu_1})^{\mu_2} & \xrightarrow[\cong]{(6.73)} & ((q^*o)^{\mu_2})^{\mu_1} \\ \cong \downarrow (6.79) & & \cong \downarrow (6.79) \\ q_{\hat{\mu}}^*(u_1^* u^*o) & \xrightarrow[\cong]{(6.80)} & q_{\hat{\mu}'}^*(u_2^* u^*o) \end{array}$$

Assume now that we are given d-critical Artin stacks $(\mathfrak{X}, s, o_{\mathfrak{X}})$ and $(\mathfrak{Y}, t, o_{\mathfrak{Y}})$ such that \mathfrak{X} and \mathfrak{Y} are quasi-separated with affine stabilizers. Then the isomorphism $\Psi_{\mathfrak{X}, \mathfrak{Y}}$ and its compatibility between ξ_s, ξ_t and $\xi_{s \boxplus t}$ implies an isomorphism of orientations

$$(6.82) \quad u_{\mathfrak{X}}^* o_{\mathfrak{X}} \boxtimes u_{\mathfrak{Y}}^* o_{\mathfrak{Y}} \cong (u_{\mathfrak{X}} \times u_{\mathfrak{Y}})^* o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}}.$$

Assume further that that we are given \mathbb{G}_m -equivariant d-critical charts (X, μ, \bar{q}) of \mathfrak{X} and (Y, ν, \bar{r}) of \mathfrak{Y} . Then the compatibility between $\Upsilon_{q, \mu}^{\text{Ind}}$ and the product implies the commutativity of the following diagram:

$$(6.83) \quad \begin{array}{ccc} (q^* o_{\mathfrak{X}})^{\mu} \boxtimes (r^* o_{\mathfrak{Y}})^{\nu} & \xrightarrow[\cong]{(6.74)} & (q^* o_{\mathfrak{X}} \boxtimes r^* o_{\mathfrak{Y}})^{\mu \times \nu} \\ \cong \downarrow (6.79) & & \cong \downarrow (6.79) \\ (q_{\mu} \times r_{\nu})^* u_{\mathfrak{X}}^* o_{\mathfrak{X}} \boxtimes u_{\mathfrak{Y}}^* o_{\mathfrak{Y}} & \xrightarrow[\cong]{(6.82)} & (q_{\mu} \times r_{\nu})^* (u_{\mathfrak{X}} \times u_{\mathfrak{Y}})^* o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}} \end{array}$$

Assume that we are given an orientation $o: \mathcal{L}^{\otimes 2} \cong \widehat{\det}(\mathbb{L}_{\mathfrak{X}})$ for a (-1) -shifted symplectic derived Artin stack \mathfrak{X} whose classical truncation is quasi-separated with affine stabilizers. We define the localized orientation u^*o on $(\text{Grad}(\mathfrak{X}), u^*\omega_{\mathfrak{X}})$ by the following composition

$$u^*o: (u^{\text{red},*}\mathcal{L} \otimes \widehat{\det}_+(\mathbb{L}_{\mathfrak{X}})^\vee)^{\otimes 2} \xrightarrow{\cong} u^*\widehat{\det}(\mathbb{L}_{\mathfrak{X}}) \otimes (\det_+(\mathbb{L}_{\mathfrak{X}})^\vee)^{\otimes 2} \xrightarrow[\cong]{\xi_{\mathfrak{X}}^{\text{der}}} \widehat{\det}(\mathbb{L}_{\text{Grad}(\mathfrak{X})}).$$

Using (6.65), we see that there exists a natural isomorphism of orientations on $\text{Grad}^2(\mathfrak{X})$

$$(6.84) \quad u_1^*u^*o \cong u_2^*u^*o.$$

Remark 6.12. Using Lemma 6.11, we see that the orientation u^*o is equivalent to the orientation constructed using Lemma 3.19. Also, it follows that the natural isomorphism (6.84) restricted to the matching locus is equivalent to the one defined using (3.9) twice.

Using Proposition 6.9 and the commutativity of the diagram (6.62), we see that there exists a natural isomorphism of orientations

$$(6.85) \quad (u^*o)^{\text{cl}} \cong u^*o^{\text{cl}}.$$

By the commutativity of the diagram (6.63), we see that the following diagram commutes:

$$(6.86) \quad \begin{array}{ccc} (u_1^*u^*o)^{\text{cl}} & \xrightarrow{\cong} & (u_2^*u^*o)^{\text{cl}} \\ \downarrow \cong & & \downarrow \cong \\ u_1^*u^*o^{\text{cl}} & \xrightarrow{\cong} & u_2^*u^*o^{\text{cl}}. \end{array}$$

Assume that we are given oriented (-1) -shifted symplectic stacks $(\mathfrak{X}, \omega_{\mathfrak{X}}, o_{\mathfrak{X}})$ and $(\mathfrak{Y}, \omega_{\mathfrak{Y}}, o_{\mathfrak{Y}})$. Then the commutativity of the diagram (6.66) implies a natural isomorphism of orientations

$$(6.87) \quad u_{\mathfrak{X}}^*o_{\mathfrak{X}} \boxtimes u_{\mathfrak{Y}}^*o_{\mathfrak{Y}} \cong u_{\mathfrak{X} \times \mathfrak{Y}}^*(o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}}).$$

The commutativity of the diagram (6.64) implies that the following diagram of orientations commutes:

$$(6.88) \quad \begin{array}{ccc} (u_{\mathfrak{X}}^*o_{\mathfrak{X}} \boxtimes u_{\mathfrak{Y}}^*o_{\mathfrak{Y}})^{\text{cl}} & \xrightarrow{\cong} & (u_{\mathfrak{X} \times \mathfrak{Y}}^*(o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}}))^{\text{cl}} \\ \downarrow \cong & & \downarrow \cong \\ u_{\mathfrak{X}}^*o_{\mathfrak{X}}^{\text{cl}} \boxtimes u_{\mathfrak{Y}}^*o_{\mathfrak{Y}}^{\text{cl}} & \xrightarrow{\cong} & u_{\mathfrak{X} \times \mathfrak{Y}}^*(o_{\mathfrak{X}}^{\text{cl}} \boxtimes o_{\mathfrak{Y}}^{\text{cl}}). \end{array}$$

7. PROOF OF THE MAIN THEOREM

The aim of this section is to prove the integral isomorphism (= Theorem B).

In §7.1, we will recall some results in sheaf theory that will be used for the proof of Theorem B. In §7.2, we will recall the definition of basic properties of the hyperbolic localization functor. In §7.3, we will reprove Desombes's theorem [Des22, Theorem 4.2] stating that hyperbolic localization of the DT perverse sheaf is again a DT perverse sheaf, with a careful choice of orientations. In §7.4, we will introduce the hyperbolic localization functor for stacks using the attractor correspondence. In §7.5, we will prove Theorem B. In §7.6, we will describe the integral isomorphism for the critical locus inside a smooth stack.

7.1. Some results from sheaf theory. We will discuss some results from sheaf theory that will be used in the proof of Theorem B.

7.1.1. Contraction lemma for monodromic complexes. We first recall the definition of monodromic complexes:

Definition 7.1. Let \mathfrak{Y} be an Artin stack with a \mathbb{G}_m -action. We say that a complex $\mathcal{F} \in D(\mathfrak{Y})$ is **monodromic** if the restriction of \mathcal{F} to any \mathbb{G}_m -orbit is locally constant. We let $D_{\text{mon}}(\mathfrak{X}) \subset D(\mathfrak{X})$ denote the full subcategory consisting of monodromic complexes.

Assume that we are given an Artin stack \mathfrak{X} with an action of multiplicative monoid \mathbb{A}^1 . We let \mathfrak{X}^0 be the fixed point substack of the submonoid $\{0, 1\}$ (see [DG15, §C.5] for detail). We have a contracting morphism $\pi: \mathfrak{X} \rightarrow \mathfrak{X}_0$ and its section $s: \mathfrak{X}_0 \rightarrow \mathfrak{X}$. The following statement is called the contraction lemma:

Proposition 7.2. *The following maps are invertible on monodromic objects:*

$$\begin{aligned} \pi_* &\xrightarrow{\text{unit}} \pi_* s_* s^* \cong s^*, \\ s^! &\cong \pi_! s_! s^! \xrightarrow{\text{counit}} \pi_!. \end{aligned}$$

The statement is proved in the context of D-module in [DG15, Theorem C.5.3]. We can prove the statement by the same argument using the fact that the monodromic complex is nothing but $\tilde{\mathbb{G}}_m$ -equivariant complex where $\tilde{\mathbb{G}}_m$ is the universal cover of \mathbb{G}_m ; see [KS94, Proposition 3.7.2] for a related statement. A more down-to-earth proof will be written in [KK].

Corollary 7.3. *Let \mathfrak{X} be a quasi-separated stack with affine stabilizers. Let $\text{gr}: \text{Filt}(\mathfrak{X}) \rightarrow \text{Grad}(\mathfrak{X})$ and $\sigma: \text{Grad}(\mathfrak{X}) \rightarrow \text{Filt}(\mathfrak{X})$ be the natural morphisms defined in §5.1. Then we have the following natural equivalence of functors*

$$\text{gr}_* \cong \sigma^*, \quad \text{gr}_! \cong \sigma^!.$$

Proof. As shown in [Hal14, Lemma 1.3.8], the maps gr and σ define an \mathbb{A}^1 -deformation retract from $\text{Filt}(\mathfrak{X})$ to $\text{Grad}(\mathfrak{X})$. Therefore we can apply Proposition 7.2. \square

7.1.2. Virtual fundamental classes. We briefly recall the theory of virtual fundamental classes following [Kha19]. It will be used only for the associativity of the integral isomorphism in Theorem B, and in fact, we can avoid the use of the virtual fundamental classes for the proof. However, we will give a proof based on it since it gives a significant simplification.

Let $f: \mathfrak{X} \rightarrow \mathfrak{Y}$ be a quasi-smooth morphism of derived Artin stacks. Then Khan [Kha19, Remark 3.8] constructed a natural transform of functors between constructible derived categories

$$\text{pur}_f: f^*[2 \text{vd } f] \rightarrow f^!$$

called the **purity transform**, which is a relative version of the virtual fundamental class. When f is smooth, this isomorphism is identified by the natural isomorphism $f^*[2 \text{vd } f] \cong f^!$ induced by the trivialization $f^! \mathbb{Q}_{\mathfrak{Y}} \cong \mathbb{Q}_{\mathfrak{X}}[2 \text{vd } f]$. Assume that we are further given a quasi-smooth morphism $g: \mathfrak{Y} \rightarrow \mathfrak{Z}$. It follows from [DJK21, Proposition 2.5.4] that the following diagram commutes:

$$(7.1) \quad \begin{array}{ccc} (g \circ f)^*[2 \text{vd}(g \circ f)] & \xrightarrow{\text{pur}_{g \circ f}} & (g \circ f)^! \\ \downarrow \text{pur}_f & & \downarrow \cong \\ f^! g^*[2 \text{vd } g] & \xrightarrow{\text{pur}_g} & f^! g^!. \end{array}$$

7.1.3. Generalized base change transforms. We will introduce a generalized version of the base change transform following [FYZ23, §3]. Consider the following diagram of derived Artin stacks:

$$(7.2) \quad \begin{array}{ccc} \mathfrak{X}_{11} & \xrightarrow{g'} & \mathfrak{X}_{12} \\ \downarrow f' & & \downarrow f \\ \mathfrak{X}_{21} & \xrightarrow{g} & \mathfrak{X}_{22}. \end{array}$$

We say that this diagram is **pullable** if the natural map $a: \mathfrak{X}_{11} \rightarrow \mathfrak{X}_{12} \times_{\mathfrak{X}_{22}} \mathfrak{X}_{21}$ is quasi-smooth. We let $\tilde{f}: \mathfrak{X}_{12} \times_{\mathfrak{X}_{22}} \mathfrak{X}_{21} \rightarrow \mathfrak{X}_{21}$ and $\tilde{g}: \mathfrak{X}_{12} \times_{\mathfrak{X}_{22}} \mathfrak{X}_{21} \rightarrow \mathfrak{X}_{12}$ the base change of f and g respectively. We set $d := \text{vd } a$ and define the generalized base change transform

$$(7.3) \quad \text{Ex}_*^! : f^! g_* \rightarrow g'_* f'^![-2d]$$

by the following composition

$$f^! g_* \cong \tilde{g}_* \tilde{f}^! \xrightarrow{\text{unit}} \tilde{g}_* a_* a^* \tilde{f}^! \xrightarrow{\text{pur}_a} \tilde{g}_* a_* a^! \tilde{f}^![-2d] \cong g'_* f'^![-2d].$$

Similarly, we define the map

$$(7.4) \quad \text{Ex}_* : g'^* f^! \rightarrow f'^! g^*[-2d]$$

by the composition

$$g'^* f^! \cong a^* \tilde{g}^* f^! \rightarrow a^* \tilde{f}^! g^* \xrightarrow{\text{pur}_a} a^! \tilde{f}^! g^*[-2d] \cong f'^! g^*[-2d].$$

The following statement is an immediate consequence of (7.1):

Lemma 7.4. *Consider the diagram (7.2) where f and f' are smooth. Set $d := \dim f$ and $d' := \dim f'$. Then it is a pullable square and the following diagram commutes:*

$$\begin{array}{ccc} f^* g_* & \xrightarrow{\quad} & g'_* f'^* \\ \downarrow \cong & & \downarrow \cong \\ f^! g_*[-2d] & \xrightarrow{\text{Ex}_*} & g'_* f'^![-2d'] \end{array}$$

where the upper horizontal map is the Beck–Chevalley transform.

The following statement is a direct consequence of [FYZ23, Proposition 3.5.5]:

Lemma 7.5. *Assume that we are given the following composition of pullable squares:*

$$\begin{array}{ccccc} \mathfrak{X}_{11} & \xrightarrow{g'} & \mathfrak{X}_{12} & \xrightarrow{h'} & \mathfrak{X}_{13} \\ \downarrow f'' & & \downarrow f' & & \downarrow f \\ \mathfrak{X}_{21} & \xrightarrow{g} & \mathfrak{X}_{22} & \xrightarrow{h} & \mathfrak{X}_{23}. \end{array}$$

Let $a: \mathfrak{X}_{12} \rightarrow \mathfrak{X}_{13} \times_{\mathfrak{X}_{23}} \mathfrak{X}_{22}$ and $a': \mathfrak{X}_{11} \rightarrow \mathfrak{X}_{12} \times_{\mathfrak{X}_{22}} \mathfrak{X}_{21}$ be the natural morphisms and set $d := \text{vd } a$ and $d' := \text{vd } a'$. Then the following diagram commutes:

$$\begin{array}{ccc} f^!(h \circ g)_* & \xrightarrow{\text{Ex}_*} & (h' \circ g')_* f'^![-2d - 2d'] \\ \downarrow \cong & & \downarrow \cong \\ f^! h_* g_* & \xrightarrow{\text{Ex}_*} h'_* f'^! g_*[-2d] \xrightarrow{\text{Ex}_*} h'_* g'_* f'^![-2d - 2d'] \end{array}$$

7.2. Hyperbolic localization. We will recall the definition of the hyperbolic localization functor of Braden [Bra03] and some of its properties, such as the commutativity between the vanishing cycle functor.

Let X be a quasi-separated algebraic space with a \mathbb{G}_m^n -action μ . Consider the following attractor correspondence

$$\begin{array}{ccc} & X^{\mu, (+, +, \dots, +)} & \\ \text{gr}_\mu \swarrow & & \searrow \text{ev}_\mu \\ X^\mu & & X. \end{array}$$

We define a functor $\text{Loc}_\mu^n: D_c^b(X) \rightarrow D_c^b(X^\mu)$ by the following composition:

$$\text{Loc}_\mu^n = \text{gr}_{\mu,*} \text{ev}_\mu^!$$

When $n = 1$, we write $\text{Loc}_\mu = \text{Loc}_\mu^1$ and call it the **hyperbolic localization functor**.

Example 7.6. Assume that U is a smooth quasi-separated algebraic space with a \mathbb{G}_m -action μ . We let $\sigma_\mu: U^\mu \hookrightarrow U^{\mu,+}$ denote the natural section of gr_μ . Then Proposition 7.2 implies an isomorphism

$$(7.5) \quad \text{Loc}_\mu(\mathbb{Q}_U) \cong \sigma_\mu^* \mathbb{Q}_{U^{\mu,+}}[-2d^-] \cong \mathbb{Q}_{U^\mu}[-2d^-]$$

where we set $d^- := \text{rank } T_U|_{U^\mu}$.

Consider also the following repelling correspondence:

$$\begin{array}{ccc} & X^{\mu,-} & \\ \text{gr}_{\mu}^{-} \swarrow & & \searrow \text{ev}_{\mu}^{-} \\ X^{\mu} & & X \end{array}$$

The following result originally proved by Braden plays an important role in our paper:

Theorem 7.7 ([Bra03], [DG14]). *There exists an equivalence of functors*

$$\text{Loc}_{\mu} = \text{gr}_{\mu,*} \text{ev}^! \cong \text{gr}_{\mu,!} \text{ev}_{\mu}^{-,*}$$

on the subcategory of $D_c^b(X)$ consisting of \mathbb{G}_m -monodromic objects.

We now discuss the compatibility between the hyperbolic localization with the smooth pullback. Assume that we are given smooth algebraic spaces X and Y with \mathbb{G}_m -action μ and ν and a \mathbb{G}_m -equivariant smooth morphism $h: X \rightarrow Y$. Consider the following commutative diagram:

$$(7.6) \quad \begin{array}{ccccc} X^{\mu} & \xleftarrow{\text{gr}_{\mu}^{\nu}} & X^{\mu,+} & \xrightarrow{\text{ev}_{\mu}^{\nu}} & X \\ \downarrow h^{\mathbb{G}_m} & \swarrow \text{gr}_{\nu}^{\mu} & \downarrow \eta & \searrow & \downarrow h \\ X^{\mu} \times_{Y^{\nu}} X^{\mu,+} & & X^{\mu,+} & & \\ \downarrow h^{\mathbb{G}_m} & & \downarrow (h^{\mathbb{G}_m})' & & \downarrow h \\ Y^{\nu} & \xleftarrow{\text{gr}_{\nu}} & Y^{\nu,+} & \xrightarrow{\text{ev}_{\nu}} & Y \end{array}$$

It follows from [Hal14, Proposition 1.3.1 (4)] that $h^{\mathbb{G}_m}$ and h^+ are smooth. We claim the natural map $h^{\mathbb{G}_m,*} \text{gr}_{\nu,*} \rightarrow \text{gr}_{\mu,*} h^{+,*}$ is invertible on monodromic objects. To prove this, by the smooth base change theorem, it is enough to show that the natural map

$$\text{gr}_{\nu,*}' \rightarrow \text{gr}_{\nu,*}' \eta_* \eta^*$$

is invertible. This is a direct consequence of Proposition 7.2, as gr_{ν}' and gr_{μ} determines an \mathbb{A}^1 -deformation retract. For a monodromic object $\mathcal{F} \in D_c^b(Y)$, we define an isomorphism

$$(7.7) \quad \rho_h: h^{\mathbb{G}_m,*} \text{Loc}_{\nu}(\mathcal{F}) \cong \text{Loc}_{\mu} h^* \mathcal{F}[2d^-]$$

by the following composition

$$\begin{aligned} \rho_h: h^{\mathbb{G}_m,*} \text{Loc}_{\nu}(\mathcal{F}) &= h^{\mathbb{G}_m,*} \text{gr}_{\nu,*} \text{ev}_{\nu}^! \mathcal{F} \\ &\cong \text{gr}_{\mu,*} h^{+,*} \text{ev}_{\nu}^! \mathcal{F} \\ &\cong \text{gr}_{\mu,*} h^{+,*} \text{ev}_{\nu}^! \mathcal{F}[-2d_0 - 2d^+] \\ &\cong \text{gr}_{\mu,*} \text{ev}_{\mu}^! h^! \mathcal{F}[-2d_0 - 2d^+] \\ &\cong \text{gr}_{\mu,*} \text{ev}_{\mu}^! h^* \mathcal{F}[2d^-] = \text{Loc}_{\mu} h^* \mathcal{F}[2d^-]. \end{aligned}$$

Here we set $d^+ := \text{rank}_+ T_{X/Y}$, $d^- := \text{rank}_- T_{X/Y}$ and $d^0 := \text{rank}_0 T_{X/Y}$.

We now discuss the associativity of the hyperbolic localization functor. Assume that we are given an algebraic space X with an action $\hat{\mu} = (\mu_1, \mu_2)$ of \mathbb{G}_m^2 and a complex $\mathcal{F} \in D_c^b(X)$. Consider the following diagram:

$$\begin{array}{ccccc} & & X_{\text{mat}}^{\hat{\mu},+,+} & & \\ & & \swarrow & & \searrow \\ & (X^{\mu_2})_{\text{mat}}^{\mu_1,+} & & & X^{\mu_2,+} \\ \text{gr}_{\mu_1|X^{\mu_2}} \swarrow & & \downarrow \text{ev}_{\mu_1|X^{\mu_2}} & \searrow \text{gr}_{\mu_2} & \downarrow \text{ev}_{\mu_2} \\ X_{\text{mat}}^{\hat{\mu}} & & X^{\mu_2} & & X \end{array}$$

It follows from Proposition (6.4) that the middle square is Cartesian. By using the proper base change theorem, we obtain an isomorphism

$$\mathrm{Loc}_{\hat{\mu}}^2(\mathcal{F})|_{X_{\mathrm{mat}}^{\hat{\mu}}} \cong (\mathrm{Loc}_{\mu_1|_{X^{\mu_2}}} \circ \mathrm{Loc}_{\mu_2}(\mathcal{F}))|_{X_{\mathrm{mat}}^{\hat{\mu}}}.$$

By using this and a similar isomorphism for the action (μ_2, μ_1) , we obtain an isomorphism

$$(7.8) \quad \chi_{\hat{\mu}} : (\mathrm{Loc}_{\mu_1|_{X^{\mu_2}}} \circ \mathrm{Loc}_{\mu_2}(\mathcal{F}))|_{X_{\mathrm{mat}}^{\hat{\mu}}} \cong (\mathrm{Loc}_{\mu_2|_{X^{\mu_1}}} \circ \mathrm{Loc}_{\mu_1}(\mathcal{F}))|_{X_{\mathrm{mat}}^{\hat{\mu}}}.$$

We now discuss the compatibility between the hyperbolic localization functor and the vanishing cycle functor. Assume now that we are given a smooth quasi-separated algebraic space U with a \mathbb{G}_m -action μ and a \mathbb{G}_m -invariant regular function f on U . We let μ_0 denote the induced \mathbb{G}_m -action on the zero locus U_0 . The following statement is an easy consequence of Theorem 7.7 and the definition of the vanishing cycle functor (see e.g. [Nak17, Proposition 5.4.1] or [Des22, Proposition 3.3]):

Proposition 7.8. *The following natural morphism constructed using (4.2) is invertible on monodromic objects:*

$$\zeta_{\mu, f} : \varphi_f|_{U^\mu} \circ \mathrm{Loc}_{\mu_0} \rightarrow \mathrm{Loc}_{\mu_0} \circ \varphi_f.$$

We now prove some properties of the map $\zeta_{\mu, f}$ that we will use later. Firstly, we now discuss the map $\zeta_{\mu, f}$ for the trivial representation. Assume that we are given a trivial \mathbb{G}_m -action μ_{triv} on an algebraic space U and a regular function f on U . Note that we have an identification of the functor $\mathrm{id} \cong \mathrm{Loc}_{\mu_{\mathrm{triv}}}$. Then it is clear that the following diagram commutes:

$$(7.9) \quad \begin{array}{ccc} \varphi_f \circ \mathrm{Loc}_{\mu_{\mathrm{triv}}} & \xrightarrow[\cong]{\zeta_{\mu_{\mathrm{triv}}, f}} & \mathrm{Loc}_{\mu_{\mathrm{triv}}} \circ \varphi_f \\ \downarrow \cong & & \downarrow \cong \\ \varphi_f & \xlongequal{\quad\quad\quad} & \varphi_f. \end{array}$$

Assume now that we are given smooth algebraic spaces U and V with \mathbb{G}_m -action μ and ν and a \mathbb{G}_m -equivariant smooth morphism $h: U \rightarrow V$. Assume further that we are given a function $g: V \rightarrow \mathbb{A}^1$ and set $f = g \circ h$. Take a monodromic object $\mathcal{F} \in D_c^b(V)$. Since $\zeta_{\mu, f}$ is constructed out of the standard exchange morphisms between six-operations, one sees that the following diagram commutes:

$$(7.10) \quad \begin{array}{ccc} h_0^{\mathbb{G}_m, *} \varphi_{g|_V} \mathrm{Loc}_\nu \mathcal{F} & \xrightarrow[\cong]{h^{\mathbb{G}_m, *} \zeta_{\nu, g}} & h_0^{\mathbb{G}_m, *} \mathrm{Loc}_{\nu_0} \varphi_g \mathcal{F} \\ \cong \downarrow \rho_h & & \rho_{h_0} \downarrow \cong \\ \varphi_f|_{X^\mu} \mathrm{Loc}_\mu h^* \mathcal{F}[2d^-] & \xrightarrow[\cong]{\zeta_{\mu, f}} & \mathrm{Loc}_{\mu_0} \varphi_f h^* \mathcal{F}[2d^-]. \end{array}$$

We now discuss the associativity of the isomorphism $\zeta_{\mu, f}$, which is related to the associativity of the cohomological Hall algebras. Let U be a quasi-separated algebraic space with a \mathbb{G}_m^2 -action $\hat{\mu} = (\mu_1, \mu_2)$ and $f: U \rightarrow \mathbb{A}^1$ be a \mathbb{G}_m^2 -invariant function. Then by the construction of the map $\zeta_{\mu, f}$ and $\chi_{\hat{\mu}}$, it is obvious that the following diagram commutes:

$$(7.11) \quad \begin{array}{ccc} \varphi_f|_{U^{\hat{\mu}}} \circ \mathrm{Loc}_{\mu_1|_{U^{\mu_2}}} \circ \mathrm{Loc}_{\mu_2}(-)|_{U_{0, \mathrm{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\zeta_{\mu_2, f} \circ \zeta_{\mu_1|_{U^{\mu_2}}, f|_{U^{\mu_2}}}} & \mathrm{Loc}_{\mu_{1,0}|_{U_0^{\mu_2,0}}} \circ \mathrm{Loc}_{\mu_{2,0}} \circ \varphi_f(-)|_{U_{0, \mathrm{mat}}^{\hat{\mu}}} \\ \cong \downarrow \chi_{\hat{\mu}} & & \downarrow \chi_{\hat{\mu}_0} \\ \varphi_f|_{U^{\hat{\mu}}} \circ \mathrm{Loc}_{\mu_2|_{U^{\mu_1}}} \circ \mathrm{Loc}_{\mu_1}(-)|_{U_{0, \mathrm{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\zeta_{\mu_1, f} \circ \zeta_{\mu_2|_{U^{\mu_1}}, f|_{U^{\mu_1}}}} & \mathrm{Loc}_{\mu_{2,0}|_{U_0^{\mu_1,0}}} \circ \mathrm{Loc}_{\mu_{1,0}} \circ \varphi_f(-)|_{U_{0, \mathrm{mat}}^{\hat{\mu}}}. \end{array}$$

We now prove that $\zeta_{\mu, f}$ is compatible with the Thom–Sebastiani isomorphism. Let U and V be smooth quasi-separated algebraic spaces with \mathbb{G}_m -action μ and ν and f and g be \mathbb{G}_m -invariant functions on U and V . Take monodromic objects $\mathcal{F} \in D_c^b(U)$ and $\mathcal{G} \in D_c^b(V)$. Since $\zeta_{\mu, f}$ is constructed out of the standard

exchange morphisms between six-operations, the commutativity of the following diagram is obvious:

$$(7.12) \quad \begin{array}{ccc} \varphi_{f|_{U^\mu}}(\mathrm{Loc}_\mu(\mathcal{F})) \boxtimes \varphi_{g|_{V^\nu}}(\mathrm{Loc}_\nu(\mathcal{G})) & \xrightarrow[\cong]{\zeta_{\mu,f} \boxtimes \zeta_{\nu,g}} & \mathrm{Loc}_{\mu_0} \varphi_f(\mathcal{F}) \boxtimes \mathrm{Loc}_{\nu_0} \varphi_g(\mathcal{G}) \\ \cong \downarrow \mathrm{TS} & & \cong \downarrow \mathrm{TS} \\ \varphi_{f|_{U^\mu} \boxplus g_{V^\nu}}(\mathrm{Loc}_{\mu \times \nu}(\mathcal{F} \boxtimes \mathcal{G}))|_{U_0^{\mu_0} \times V_0^{\nu_0}} & \xrightarrow[\cong]{\zeta_{\mu \times \nu, f \boxplus g}} & \mathrm{Loc}_{\mu_0 \times \nu_0}(\varphi_{f \boxplus g}(\mathcal{F} \boxtimes \mathcal{G}))|_{U_0^{\mu_0} \times V_0^{\nu_0}}. \end{array}$$

7.3. Hyperbolic localization of DT perverse sheaves. We now discuss Descombes's Theorem [Des22, Theorem 4.2] stating that the hyperbolic localization of the DT perverse sheaf is again a DT perverse sheaf. Since we will use the construction of the isomorphism later and the paper [Des22] seems to ignore possible automorphisms of orientations, we will give a detailed proof here.

Theorem 7.9 ([Des22, Theorem 4.2]). *Let (X, μ, s) be a \mathbb{G}_m -equivariant d -critical algebraic space with an orientation o . Assume that X is quasi-separated. Then there exists a natural isomorphism*

$$(7.13) \quad \zeta_{\mu,s,o} : \varphi_{X^\mu, s^\mu, o^\mu}[I_X^\mu] \cong \mathrm{Loc}_\mu(\varphi_{X,s,o}).$$

Proof. We first construct $\zeta_{\mu,s,o}$ over each étale d -critical chart. Take a \mathbb{G}_m -equivariant étale d -critical chart $\mathfrak{R} = (R, \eta, U, f, i)$. By abuse of notation, we write the \mathbb{G}_m -action on R and U by μ . We first construct an isomorphism

$$\beta_{\mathfrak{R}} : Q_{\mathfrak{R}}^o|_{R^\mu} \cong Q_{\mathfrak{R}^\mu}^{o^\mu}.$$

Recall from §4.2 that $Q_{\mathfrak{R}}^o$ parametrizes local isomorphisms between o and the canonical orientation. Therefore, it is enough to construct an isomorphism

$$(7.14) \quad (o_{\mathfrak{R}}^{\mathrm{can}})^\mu \cong o_{\mathfrak{R}^\mu}^{\mathrm{can}}.$$

which we have already seen in (6.75).

We now construct

$$\zeta_{\mu,s,o,\mathfrak{R}} : \varphi_{X^\mu, s^\mu, o^\mu}[I_X^\mu]|_{R^\mu} \cong \mathrm{Loc}_\mu(\varphi_{X,s,o})|_{R^\mu}$$

by the following composition

$$\begin{aligned} \varphi_{X^\mu, s^\mu, o^\mu}[I_X^\mu] &\xrightarrow[\cong]{\omega_{\mathfrak{R}^\mu}} \varphi_{f|_{U^\mu}} \otimes Q_{\mathfrak{R}^\mu}^{o^\mu}[d^+ - d^-]|_{R^\mu} \\ &\xrightarrow[\cong]{\zeta_{\mu,f} \otimes \beta_{\mathfrak{R}}^{-1}} \mathrm{Loc}_\mu(\varphi_f) \otimes Q_{\mathfrak{R}}^o|_{R^\mu} \xrightarrow[\cong]{\omega_{\mathfrak{R}}^{-1}, (7.7)} \mathrm{Loc}_\mu(\varphi_{X,s,o})|_{R^\mu}. \end{aligned}$$

Here, we set $d^+ := \mathrm{rank}_+ T_U$ and $d^- := \mathrm{rank}_- T_U$. The shift in the third complex disappears by (4.1) and (7.5). We now prove that $\zeta_{\mu,s,o,\mathfrak{R}}$ does not depend on the choice of \mathfrak{R} . Since it is clear that $\zeta_{\mu,s,o,\mathfrak{R}}$ is compatible with \mathbb{G}_m -equivariant étale pullback of étale d -critical charts, by using Theorem 3.31, it is enough to prove the identity

$$\zeta_{\mu,s,o,\mathfrak{R}} = \zeta_{\mu',s,o,\mathfrak{R}'}$$

for \mathfrak{R}' the étale d -critical chart defined by

$$\mathfrak{R}' = (R, \eta, U \times \mathbb{A}^{2N}, f \boxplus q, (i, 0))$$

for some \mathbb{G}_m -equivariant non-degenerate quadratic function q on \mathbb{A}^{2N} . We write

$$q = (x_1 y_1 + \cdots + x_n y_n) + (z_1 w_1 + \cdots + z_m w_m)$$

where x_i (resp. y_i) has positive (resp. negative) \mathbb{G}_m -weights and z_j and w_j are invariant under the \mathbb{G}_m -action. We write $q_{x,y} = x_1 y_1 + \cdots + x_n y_n$ and $q_{z,w} = z_1 w_1 + \cdots + z_m w_m$. We let $\mu' = (\mu, \nu)$ denote the \mathbb{G}_m -action on $U \times \mathbb{A}^{2N}$. Consider the following diagram:

$$\begin{array}{ccc} Q_{\mathfrak{R}'}^o|_{R^\mu} & \xrightarrow[\cong]{\beta_{\mathfrak{R}'}} & Q_{\mathfrak{R}'^\mu}^{o^\mu} \\ \cong \downarrow b_{y,w} & & \cong \downarrow b_w \\ Q_{\mathfrak{R}}^o|_{R^\mu} & \xrightarrow[\cong]{\beta_{\mathfrak{R}}} & Q_{\mathfrak{R}^\mu}^{o^\mu}. \end{array}$$

To see this commutes, consider the étale d-critical chart

$$\mathfrak{X}'_{x,y} = (R, \eta, U \times \mathbb{A}^{2n}, f \boxplus q_{x,y}, (i, 0)).$$

By using the identity (4.10), it is enough to prove that the following diagram commutes:

$$\begin{array}{ccc} Q_{\mathfrak{X}'_{x,y}}^o |_{R^\mu} & \xrightarrow[\cong]{\beta_{\mathfrak{X}'_{x,y}}} & Q_{\mathfrak{X}'_{x,y}}^{o\mu} \\ \cong \downarrow b_y & & \parallel \\ Q_{\mathfrak{X}}^o |_{R^\mu} & \xrightarrow[\cong]{\beta_{\mathfrak{X}}} & Q_{\mathfrak{X}}^{o\mu}. \end{array}$$

It is equivalent to the commutativity of the following diagram:

$$(7.15) \quad \begin{array}{ccc} K_{U \times \mathbb{A}^{2n}} |_{R^{\mu, \text{red}}} \otimes K_{U \times \mathbb{A}^{2n}} |_{R^{\mu, \text{red}}}^{\pm, \vee} & \xrightarrow{\cong} & K_{U^\mu} |_{R^{\mu, \text{red}}} \\ \cong \downarrow & & \parallel \\ K_U |_{R^{\mu, \text{red}}} \otimes K_U |_{R^{\mu, \text{red}}}^{\pm, \vee} & \xrightarrow{\cong} & K_{U^\mu} |_{R^{\mu, \text{red}}} \end{array}$$

where the left vertical map is constructed using (4.9) and (6.6). Consider the following acyclic complex

$$\mathbb{L}_{\mathfrak{X}'_{x,y}/\mathfrak{X}}^+ := [T_{\mathbb{A}^{2n}}^+ \xrightarrow{\text{Hess}_q^+} \Omega_{\mathbb{A}^{2n}}^+], \quad \partial x_i \mapsto dy_i.$$

The fibre sequence

$$\Omega_{\mathbb{A}^{2n}}^+ \rightarrow \mathbb{L}_{\mathfrak{X}'/\mathfrak{X}}^+ \rightarrow T_{\mathbb{A}^{2n}}^+[1]$$

together with (2.5) and (2.14) induces a trivialization

$$(7.16) \quad K_{\mathbb{A}^{2n}} \cong K_{\mathbb{A}^{2n}}^\pm \cong \mathcal{O}_{\mathbb{A}^{2n}}.$$

By the construction of (6.6), it is enough to compare this isomorphism with (4.9). Note that we have an equivalence of fibre sequences

$$\begin{array}{ccccc} \Omega_{\mathbb{A}^{2n}}^+ & \longrightarrow & \mathbb{L}_{\mathfrak{X}'/\mathfrak{X}}^+ & \longrightarrow & T_{\mathbb{A}^{2n}}^+[1] \\ \parallel & & \simeq \downarrow & & \simeq \downarrow \text{Hess}_{q_{x,y}} \\ \Omega_{\mathbb{A}^{2n}}^+ & \longrightarrow & 0 & \longrightarrow & \Omega_{\mathbb{A}^{2n}}^+[1]. \end{array}$$

Therefore by using the construction of (2.5), (2.14) and that the double dual of a graded line bundle is trivialized with the insertion of a sign of the grading, we see that the trivialization (7.16) is given by

$$dx_1 \wedge \cdots \wedge dx_n \wedge dy_n \wedge \cdots \wedge dy_1 \mapsto 1,$$

hence we conclude the commutativity of the diagram (7.15).

On the other hand, consider the following diagram

$$\begin{array}{ccc} \varphi_f |_{U^\mu \boxplus q_{z,w}} [I_X^\mu] & \xrightarrow[\cong]{\zeta_{\mu', f \boxplus q}} & \text{Loc}_\mu(\varphi_{f \boxplus q}) \\ \cong \downarrow \text{TS}^{-1} & & \cong \downarrow \text{TS}^{-1} \\ \varphi_f |_{U^\mu} \boxtimes \varphi_{q_{z,w}} [I_X^\mu] & \xrightarrow[\cong]{\zeta_{\mu, f} \boxtimes \zeta_{\nu, q}} & \text{Loc}_\mu(\varphi_f \boxtimes \varphi_q) \\ \cong \downarrow c_w & & \cong \downarrow c_{y,w} \\ \varphi_f |_{U^\mu} [I_X^\mu] & \xrightarrow[\cong]{\zeta_{\mu, f}} & \text{Loc}_\mu(\varphi_f). \end{array}$$

The commutativity of the upper square follows from the commutativity of the diagram (7.12). To prove the commutativity of the lower diagram, using the commutativity of the diagram (4.7), it is enough to prove the

commutativity of the following diagram

$$\begin{array}{ccc}
\mathrm{Loc}_{\nu_{x,y}}(\varphi_{q_{x,y}})|_0 & \xrightarrow[\cong]{\zeta_{\nu_{x,y},q_{x,y}}} & \varphi_0 \\
\cong \downarrow c_y & & \cong \downarrow \\
\mathbb{Q}_0 & \xlongequal{\quad\quad\quad} & \mathbb{Q}_0
\end{array}$$

where $\nu_{x,y}$ denotes the \mathbb{G}_m -action on \mathbb{A}^{2n} . Note that we have an identification $(\mathbb{A}^{2n})^{\nu_{x,y},-} = V_y$ where V_y denote the subspace of \mathbb{A}^{2n} spanned by y -coordinates. Then the claim follows from the construction of c_y and $\zeta_{\nu_{x,y},q_{x,y}}$ since both of them are constructed using the fundamental class of V_y . \square

We now prove some properties of $\zeta_{\mu,s,o}$ that we will use later. Firstly, let μ^{triv} be the trivial \mathbb{G}_m -action on an oriented d-critical algebraic space (X, s, o) . Then by the commutativity of (7.9), we see that the following diagram commutes:

$$(7.17) \quad \begin{array}{ccccc}
\varphi_{X,s,o} & \xrightarrow[\cong]{\zeta_{\mu^{\mathrm{triv}},s,o}} & \mathrm{Loc}_{\mu^{\mathrm{triv}}}(\varphi_{X,s,o}) & \xrightarrow[\cong]{} & \varphi_{X,s,o} \\
& \searrow & & \swarrow & \\
& & \mathrm{id} & &
\end{array}$$

We now prove that the map $\zeta_{\mu,s,o}$ is compatible with smooth pullback. Though this is already proved in [Des22, Proposition 4.5], we will include the proof for reader's convenience:

Proposition 7.10 ([Des22, Proposition 4.5]). *Let (X, μ, s) and (Y, ν, t) be quasi-separated \mathbb{G}_m -equivariant algebraic spaces and $h: X \rightarrow Y$ be a \mathbb{G}_m -equivariant smooth morphism with $s = h^*t$. Let o be an orientation for (Y, ν, t) . Then the following diagram commutes:*

$$\begin{array}{ccc}
h^{\mathbb{G}_m,*} \varphi_{Y^{\nu,t},o^{\nu}}[I_Y^{\nu}] & \xrightarrow[\cong]{\Theta_{h^{\mathbb{G}_m},t^{\nu},o^{\nu}}} & \varphi_{X^{\mu,s^{\mu}},(h^*o)^{\mu}}[I_Y^{\nu} - d_0] \\
\cong \downarrow \zeta_{\nu,t,o} & & \cong \downarrow \zeta_{\mu,s,h^*o} \\
h^{\mathbb{G}_m,*} \mathrm{Loc}_{\nu}(\varphi_{Y,t,o}) & \xrightarrow[\cong]{\rho_h} \mathrm{Loc}_{\mu}(h^* \varphi_{Y,t,o})[2d^-] & \xrightarrow[\cong]{\Theta_{h,t,o}} \mathrm{Loc}_{\mu}(\varphi_{X,s,h^*o})[2d^- - d]
\end{array}$$

where we set $d := \dim h$, $d_0 := \dim h^{\mathbb{G}_m}$, $d^+ := \mathrm{rank}_+ T_{X/Y}$ and $d^- := \mathrm{rank}_- T_{X/Y}$ and we used the identity $I_X^{\mu} - I_Y^{\nu} = d^+ - d^-$ from (6.1).

Proof. It is enough to prove the commutativity for each \mathbb{G}_m -equivariant étale d-critical chart around each point of X^{μ} . Using Proposition 3.32, we may take \mathbb{G}_m -equivariant étale d-critical chart $\mathfrak{X} = (R, \eta, U, f, i)$ of X and $\mathfrak{S} = (S, \gamma, V, g, i)$ and a \mathbb{G}_m -equivariant smooth morphism $(H_0, H): \mathfrak{X} \rightarrow \mathfrak{S}$. By the commutativity of the diagram (7.10), it is enough to prove the commutativity of the following diagram:

$$\begin{array}{ccc}
H_0^* Q_{\mathfrak{S}}^o|_{R^{\mu}} & \xrightarrow[\cong]{b_H} & Q_{\mathfrak{X}}^{h^*o}|_{R^{\mu}} \\
\cong \downarrow \beta_{\mathfrak{S}} & & \cong \downarrow \beta_{\mathfrak{X}} \\
H_0^{\mathbb{G}_m,*} Q_{\mathfrak{S}^{\nu}}^{o^{\nu}} & \xrightarrow[\cong]{b_{H^{\mathbb{G}_m}}} & Q_{\mathfrak{X}^{\mu}}^{(h^*o)^{\mu}}
\end{array}$$

where the \mathbb{G}_m -action on R and S are denoted by μ and ν by abuse of notation, and we use the identification (6.72). This follows from the commutativity of the diagram (6.76). \square

We now prove the associativity of the morphism $\zeta_{\mu,s,o}$:

Proposition 7.11. *Let $(X, (\mu_1, \mu_2), s)$ be a quasi-separated \mathbb{G}_m^2 -equivariant d -critical algebraic space. Then the following diagram commutes:*

$$(7.18) \quad \begin{array}{ccc} \varphi_{X^{\hat{\mu}}, s^{\hat{\mu}}, (o^{\mu_2})^{\mu_1}} [I_X^{\mu_2} + I_{X^{\mu_2}}^{\mu_1}]|_{X_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\zeta_{\mu_2, s, o} \circ \zeta_{\mu_1 | X^{\mu_2}, s^{\mu_2}, o^{\mu_2}}} & \text{Loc}_{\mu_1}(\text{Loc}_{\mu_2}(\varphi_{X, s, o}))|_{X_{\text{mat}}^{\hat{\mu}}} \\ \cong \downarrow & & \cong \downarrow \chi_{\hat{\mu}} \\ \varphi_{X^{\hat{\mu}}, s^{\hat{\mu}}, (o^{\mu_1})^{\mu_2}} [I_X^{\mu_1} + I_{X^{\mu_1}}^{\mu_2}]|_{X_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\zeta_{\mu_1, s, o} \circ \zeta_{\mu_2 | X^{\mu_1}, s^{\mu_1}, o^{\mu_1}}} & \text{Loc}_{\mu_2}(\text{Loc}_{\mu_1}(\varphi_{X, s, o}))|_{X_{\text{mat}}^{\hat{\mu}}}. \end{array}$$

Here, the left vertical map is constructed using the isomorphism (6.73).

Proof. Take a point $x \in X_{\text{mat}}^{\hat{\mu}}$. It is enough to prove the commutativity for an \mathbb{G}_m^2 -equivariant étale d -critical charts around x . Take a \mathbb{G}_m^2 -equivariant étale d -critical chart $\mathfrak{X} = (R, \eta, U, f, i)$ of x . By abuse of notion, we will denote by $\hat{\mu} = (\mu_1, \mu_2)$ the \mathbb{G}_m^2 -actions on R and U . Using Proposition 3.29, we may assume that there exists a point $\hat{x} \in R^{\hat{\mu}}$ lifting x which is contained in $U^{\hat{\mu}}$. By the commutativity of the diagram (7.11), it is enough to prove the commutativity of the diagram

$$\begin{array}{ccccc} Q_{\mathfrak{X}^{\hat{\mu}}}^{(o^{\mu_2})^{\mu_1}}|_{R_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\beta_{\mathfrak{X}^{\mu_2}}} & Q_{\mathfrak{X}^{\mu_2}}^{o^{\mu_2}}|_{R_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\beta_{\mathfrak{X}, \mu_2}} & Q_{\mathfrak{X}}^o|_{R_{\text{mat}}^{\hat{\mu}}} \\ \cong \downarrow & & & & \parallel \\ Q_{\mathfrak{X}^{\hat{\mu}}}^{(o^{\mu_1})^{\mu_2}}|_{R_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\beta_{\mathfrak{X}^{\mu_1}}} & Q_{\mathfrak{X}^{\mu_1}}^{o^{\mu_1}}|_{R_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\beta_{\mathfrak{X}, \mu_1}} & Q_{\mathfrak{X}}^o|_{R_{\text{mat}}^{\hat{\mu}}}. \end{array}$$

Here the subscript of $\beta_{\mathfrak{X}, \mu_1}$ and $\beta_{\mathfrak{X}, \mu_2}$ denotes \mathbb{G}_m -action that we consider. Recall that $Q_{\mathfrak{X}}^o$ parametrizes local isomorphisms between $o_{\mathfrak{X}}^{\text{can}}$ and o . The commutativity of the diagram follows from (6.77). \square

We now show that the map $\zeta_{\mu, s, o}$ is compatible with the Thom–Sebastiani isomorphism.

Proposition 7.12. *Let (X, μ, s) and (Y, ν, t) be quasi-separated \mathbb{G}_m -equivariant d -critical algebraic spaces with orientations o_X and o_Y . Then the following diagram commutes:*

$$\begin{array}{ccc} \varphi_{X^{\mu}, s^{\mu}, o_X^{\mu}} \boxtimes \varphi_{Y^{\nu}, t^{\nu}, o_Y^{\nu}} [I_X^{\mu} + I_Y^{\nu}] & \xrightarrow[\cong]{\zeta_{\mu, s, o_X} \boxtimes \zeta_{\nu, t, o_Y}} & \text{Loc}_{\mu}(\varphi_{X, s, o_X}) \boxtimes \text{Loc}_{\nu}(\varphi_{Y, t, o_Y}) \\ \cong \downarrow \text{TS} & & \cong \downarrow \text{TS} \\ \varphi_{X^{\mu} \times Y^{\nu}, s^{\mu} \boxplus t^{\nu}, o_X^{\mu} \boxtimes o_Y^{\nu}} [I_{X \times Y}^{\mu \times \nu}] & \xrightarrow[\cong]{\zeta_{X \times Y, s \boxplus t, o_X \boxtimes o_Y}} & \text{Loc}_{\mu \times \nu}(\varphi_{X \times Y, s \boxplus t, o_X \boxtimes o_Y}). \end{array}$$

Proof. It is enough to prove the commutativity for each \mathbb{G}_m -equivariant étale d -critical charts around each point of $(x, y) \in X^{\mu} \times Y^{\nu}$. Take an étale d -critical chart $\mathfrak{X} = (R, \eta, U, f, i)$ of x and $\mathfrak{S} = (S, \gamma, V, g, j)$ of y . By the commutativity of the diagram (7.12) and the construction of Thom–Sebastiani isomorphism of d -critical algebraic spaces (4.17), it is enough to prove the commutativity of the following diagram:

$$\begin{array}{ccc} Q_{\mathfrak{X}}^{o_X} \boxtimes Q_{\mathfrak{S}}^{o_Y} |_{X^{\mu} \times Y^{\nu}} & \xrightarrow[\cong]{\beta_{\mathfrak{X}} \boxtimes \beta_{\mathfrak{S}}} & Q_{\mathfrak{X}^{\mu}}^{o_X^{\mu}} \boxtimes Q_{\mathfrak{S}^{\nu}}^{o_Y^{\nu}} \\ \cong \downarrow \alpha_{\mathfrak{X}, \mathfrak{S}} & & \cong \downarrow \alpha_{\mathfrak{X}^{\mu}, \mathfrak{S}^{\nu}} \\ Q_{\mathfrak{X} \times \mathfrak{S}}^{o_X \boxtimes o_Y} |_{X^{\mu} \times Y^{\nu}} & \xrightarrow[\cong]{\beta_{\mathfrak{X} \times \mathfrak{S}}} & Q_{\mathfrak{X}^{\mu} \times \mathfrak{S}^{\nu}}^{o_X^{\mu} \boxtimes o_Y^{\nu}}. \end{array}$$

This follows from (6.78). \square

7.4. Hyperbolic localization for stacks. Here we will introduce the hyperbolic localization functor for Artin stacks. Let \mathfrak{X} be an Artin stack with quasi-separated diagonal and affine stabilizer groups. Consider the following correspondence:

$$(7.19) \quad \begin{array}{ccc} & \text{Filt}^n(\mathfrak{X}) & \\ \swarrow \text{gr} & & \searrow \text{ev} \\ \text{Grad}^n(\mathfrak{X}) & & \mathfrak{X} \end{array}$$

We define a functor $\text{Loc}_{\mathfrak{X}}^n := D_c^b(\mathfrak{X}) \rightarrow D(\text{Grad}^n(\mathfrak{X}))$ by

$$\text{Loc}_{\mathfrak{X}}^n := \text{gr}_* \text{ev}^!$$

When $n = 1$, we write $\text{Loc}_{\mathfrak{X}} = \text{Loc}_{\mathfrak{X}}^1$ and we will call it the **hyperbolic localization functor**.

Remark 7.13. More generally, for a quasi-separated Artin stack \mathfrak{X} with affine stabilizers with a \mathbb{G}_m^n -action μ , once can define the functor $\text{Loc}_{\mu}^n: D(\mathfrak{X}) \rightarrow D(\mathfrak{X}^\mu)$ using the correspondence

$$(7.20) \quad \mathfrak{X}^\mu \leftarrow \mathfrak{X}^{\mu,+} \rightarrow \mathfrak{X}.$$

The correspondence (7.19) corresponds to the case of the trivial action. Since the correspondence (7.20) can be embedded to the correspondence (7.19) for $[\mathfrak{X}/\mathbb{G}_m^n]$, we do not lose generality by restricting to the case of the trivial action.

Let \mathfrak{X} be a quasi-separated Artin stack with affine stabilizers. By using Proposition 5.25, take a collection of \mathbb{G}_m -equivariant charts (X, μ, \bar{q}) such that $q_\mu: X^\mu \rightarrow \text{Grad}(\mathfrak{X})$ is jointly surjective. Consider the following diagram:

$$\begin{array}{ccccc} X^\mu & \xleftarrow{\text{gr}_\mu} & X^{\mu,+} & \xrightarrow{\text{ev}_\mu} & X \\ & \swarrow \text{gr}' & \searrow \eta & & \downarrow q \\ & & X^\mu \times_{\text{Grad}(\mathfrak{X})} \text{Filt}(\mathfrak{X}) & & \downarrow q \\ & & \searrow q'_\mu & & \downarrow q \\ \text{Grad}(\mathfrak{X}) & \xleftarrow{\text{gr}} & \text{Filt}(\mathfrak{X}) & \xrightarrow{\text{ev}} & \mathfrak{X} \end{array}$$

It follows from [Hal14, Proposition 1.3.1 (4)] that q_μ and q'_μ are smooth. Further, by [Hal14, Lemma 1.3.8], the map gr is \mathbb{A}^1 -deformation retract. Therefore, by repeating the construction of the isomorphism (7.7) using Corollary 7.3, we obtain an isomorphism of functors

$$(7.21) \quad \rho_q: q_\mu^* \circ \text{Loc}_{\mathfrak{X}} \cong \text{Loc}_{\mu} \circ q^*[2d^-]$$

where we set $d^- := \text{rank } T_{X/\mathfrak{X}}^-$. In particular, the functor $\text{Loc}_{\mathfrak{X}}$ preserves constructible objects.

Lemma 7.14. *Let \mathfrak{X} be a quasi-separated Artin stack with affine stabilizers and $h: (X_1, \mu_1, \bar{q}_1) \rightarrow (X_2, \mu_2, \bar{q}_2)$ be a smooth morphism of \mathbb{G}_m -equivariant charts. Then the following diagram commutes:*

$$\begin{array}{ccc} q_{1,\mu_1}^* \text{Loc}_{\mathfrak{X}} & \xrightarrow[\cong]{\rho_{q_1}} & \text{Loc}_{\mu_1} q_1^*[2d_1^-] \\ \downarrow \cong & & \uparrow \cong \\ h^{\mathbb{G}_m,*} q_{2,\mu_2}^* \text{Loc}_{\mathfrak{X}} & \xrightarrow[\cong]{\rho_{q_2}} & h^{\mathbb{G}_m,*} \text{Loc}_{\mu_2} q_2^*[2d_2^-] \end{array}$$

where we set $d_1^- := \text{rank}_- T_{X_1/\mathfrak{X}}$ and $d_2^- := \text{rank}_- T_{X_2/\mathfrak{X}}$.

Proof. The statement can be decomposed into the commutativity of the following diagrams:

$$\begin{array}{ccc} q_{1,\mu_1}^{+,*} \text{ev}^! & \xrightarrow{\cong} & \text{ev}_{\mu_1}^! q_1^*[2d_1^-] \\ \downarrow \cong & & \downarrow \cong \\ h^{+,*} q_{2,\mu_2}^{+,*} \text{ev}^! & \xrightarrow{\cong} & \text{ev}_{\mu_1}^! h^* q_2^*[2d_1^-], \end{array} \quad \begin{array}{ccc} q_{1,\mu_1}^* \text{gr}_* & \xrightarrow{\cong} & \text{gr}_{\mu_1,*} q_{1,\mu_1}^{+,*} \\ \downarrow \cong & & \downarrow \cong \\ h^{\mathbb{G}_m,*} q_{2,\mu_2}^* \text{gr}_* & \xrightarrow{\cong} & \text{gr}_{\mu_1,*} h^{+,*} q_{2,\mu_2}^{+,*}. \end{array}$$

The commutativity of the left diagram follows from the associativity with respect to the composition of the purity transform $q^*[2 \dim q] \cong q^!$ for smooth morphism q . The commutativity of the right diagram follows from the associativity of the Beck–Chevalley transform. \square

We now discuss the associativity of the hyperbolic localization functor. By repeating the construction of $\chi_{\hat{\mu}}$ in (7.8) using Proposition 5.22, we obtain the following natural isomorphism

$$\chi_{\mathfrak{X}}: \text{Loc}_{\text{Grad}(\mathfrak{X})} \circ \text{Loc}_{\mathfrak{X}}(-)|_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}} \cong \sigma^* \text{Loc}_{\text{Grad}(\mathfrak{X})} \circ \text{Loc}_{\mathfrak{X}}(-)|_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}}$$

where $\sigma: \text{Grad}^2(\mathfrak{X}) \cong \text{Grad}^2(\mathfrak{X})$ denotes the swapping isomorphism.

Lemma 7.15. *Let \mathfrak{X} be a quasi-separated Artin stack with affine stabilizers and $(X, (\mu_1, \mu_2), \bar{q})$ be a \mathbb{G}_m^2 -equivariant chart such that $q_{\hat{\mu}}: X^{\hat{\mu}} \rightarrow \text{Grad}^2(\mathfrak{X})$ restricts to $X_{\text{mat}}^{\hat{\mu}} \rightarrow \text{Grad}^2(\mathfrak{X})_{\text{mat}}$. Then the following diagram commutes:*

$$\begin{array}{ccc} q_{\hat{\mu}}^* \circ \text{Loc}_{\text{Grad}(\mathfrak{X})} \circ \text{Loc}_{\mathfrak{X}}(-)|_{X_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\chi_{\mathfrak{X}}} & q_{\hat{\mu}}^* \circ \sigma^* \text{Loc}_{\text{Grad}(\mathfrak{X})} \circ \text{Loc}_{\mathfrak{X}}(-)|_{X_{\text{mat}}^{\hat{\mu}}} \\ \cong \downarrow \rho_q \circ \rho_{q\mu_2} & & \cong \downarrow \rho_q \circ \rho_{q\mu_1} \\ \text{Loc}_{\mu_1|_{X^{\mu_2}}} \circ \text{Loc}_{\mu_2} \circ q^*(-)[2e]|_{X_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\chi_{\hat{\mu}}} & \text{Loc}_{\mu_2|_{X^{\mu_1}}} \circ \text{Loc}_{\mu_1} \circ q^*(-)[2e]|_{X_{\text{mat}}^{\hat{\mu}}}. \end{array}$$

where we set $e := \text{rank}_{\mu_1 - (T_X/\mathfrak{X})} + \text{rank}_{\mu_2 - (T_{X^{\mu_1}}/\text{Grad}(\mathfrak{X}))}$.

Proof. Note that the map $\chi_{\mathfrak{X}}$ can be written as the composition

$$\text{Loc}_{\text{Grad}(\mathfrak{X})} \circ \text{Loc}_{\mathfrak{X}}(-)|_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}} \xrightarrow[\cong]{\chi'_{\mathfrak{X}}} \text{Loc}_{\mathfrak{X}}^2(-)|_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}} \xrightarrow[\cong]{\chi''_{\mathfrak{X}}} \sigma^* \text{Loc}_{\text{Grad}(\mathfrak{X})} \circ \text{Loc}_{\mathfrak{X}}(-)|_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}}$$

where $\chi'_{\mathfrak{X}}$ and $\chi''_{\mathfrak{X}}$ are constructed using the base change transform (7.3) for the Cartesian diagram

$$\begin{array}{ccc} \text{Filt}^2(\mathfrak{X})_{\text{mat}} & \xrightarrow{\text{Filt}(\text{gr}_{\mathfrak{X}})} & \text{Filt}(\text{Grad}(\mathfrak{X}))_{\text{mat}} \\ \downarrow \text{ev}_{\text{Filt}(\mathfrak{X})} & & \downarrow \text{ev}_{\text{Grad}(\mathfrak{X})} \\ \text{Filt}(\mathfrak{X}) & \xrightarrow{\text{gr}_{\mathfrak{X}}} & \text{Grad}(\mathfrak{X}). \end{array}$$

Therefore it is enough to prove the commutativity of the following diagram:

$$\begin{array}{ccc} q_{\hat{\mu}}^* \circ \text{Loc}_{\text{Grad}(\mathfrak{X})} \circ \text{Loc}_{\mathfrak{X}}(-)|_{X_{\text{mat}}^{\hat{\mu}}} & \xrightarrow[\cong]{\chi'_{\mathfrak{X}}} & q_{\hat{\mu}}^* \circ \text{Loc}_{\mathfrak{X}}^2(-)|_{X_{\text{mat}}^{\hat{\mu}}} \\ \cong \downarrow \rho_q \circ \rho_{q\mu_2} & & \cong \downarrow \\ \text{Loc}_{\mu_1|_{X^{\mu_2}}} \circ \text{Loc}_{\mu_2} \circ q^*(-)|_{X_{\text{mat}}^{\hat{\mu}}} [2e] & \xrightarrow[\cong]{\chi'_{\hat{\mu}}} & \text{Loc}_{\mu_2}^2 \circ q^*(-)|_{X_{\text{mat}}^{\hat{\mu}}} [2e], \end{array}$$

where the right vertical map is defined in the same manner as ρ_q in (7.21). By the construction of ρ_q and $\chi'_{\mathfrak{X}}$, this can be further reduced to the commutativity of the following diagram:

$$\begin{array}{ccc} q_{\hat{\mu}}^{(+,0),*} \text{ev}_{\text{Grad}(\mathfrak{X})}^! \text{gr}_{\mathfrak{X},*}(-)|_{(X^{\mu_2})_{\text{mat}}^{\mu_1,+}} & \xrightarrow[\cong]{} & q_{\hat{\mu}}^{(+,0),*} \text{Filt}(\text{gr}_{\mathfrak{X}})_* \text{ev}_{\text{Filt}(\mathfrak{X})}^!(-)|_{(X^{\mu_2})_{\text{mat}}^{\mu_1,+}} \\ \downarrow \cong & & \downarrow \cong \\ \text{ev}_{\mu_1|_{X^{\mu_2}}}^! \text{gr}_{\mu_2,*} q_{\mu_2}^{+,*}(-)[2e']|_{(X^{\mu_2})_{\text{mat}}^{\mu_1,+}} & \xrightarrow[\cong]{} & \text{gr}_{\mu_2|_{X^{\mu_1,+}}} \text{ev}_{\mu_1|_{X^{\mu_2,+}}}^! q_{\mu_2}^{+,*}(-)[2e']|_{(X^{\mu_2})_{\text{mat}}^{\mu_1,+}} \end{array}$$

where $q_{\hat{\mu}}^{(+,0),*}: (X^{\mu_2})_{\text{mat}}^{\mu_1,+} \rightarrow \text{Filt}(\text{Grad}(\mathfrak{X}))$ is the natural morphism and we set $e' := \text{rank}_{\mu_1 - T_{X^{\mu_2}}/\text{Grad}(\mathfrak{X})}$. The horizontal map is the base change map (7.3) and the vertical map is also the base change map (7.3) after the identification $q_{\hat{\mu}}^{(+,0),*} \cong q_{\hat{\mu}}^{(+,0),!}[-2 \dim q_{\hat{\mu}}^{(+,0)}]$ and $q_{\mu_2}^{+,*} \cong q_{\mu_2}^{+,!}[-2 \dim q_{\mu_2}^+]$ by Lemma 7.4. Then the commutativity of the above diagram follows by Lemma 7.5 applied to the following composition of pullable squares:

$$\begin{array}{ccc} X_{\text{mat}}^{\hat{\mu},+,+} & \longrightarrow & (X^{\mu_2})_{\text{mat}}^{\mu_1,+} & & X_{\text{mat}}^{\hat{\mu},+,+} & \longrightarrow & (X^{\mu_2})_{\text{mat}}^{\mu_1,+} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \text{Filt}^2(\mathfrak{X})_{\text{mat}} & \longrightarrow & \text{Filt}(\text{Grad}(\mathfrak{X}))_{\text{mat}} & & X^{\mu_2,+} & \longrightarrow & X^{\mu_2} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \text{Filt}(\mathfrak{X}) & \longrightarrow & \text{Grad}(\mathfrak{X}), & & \text{Filt}(\mathfrak{X}) & \longrightarrow & \text{Grad}(\mathfrak{X}). \end{array}$$

□

7.5. Proof of the main theorem. In this section, we will prove Theorem B.

Theorem 7.16. *Let (\mathfrak{X}, s, o) be an oriented d -critical stack where \mathfrak{X} is quasi-separated and has affine stabilizers. Then there exists a natural isomorphism*

$$\zeta_{\mathfrak{X}, s, o}: \varphi_{\mathrm{Grad}(\mathfrak{X}), u^*s, u^*o}[I_{\mathfrak{X}}] \cong \mathrm{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}, s, o}).$$

Proof. Take a \mathbb{G}_m -equivariant chart $\Omega = (X, \mu, q)$. Set $d := \mathrm{rank} T_{X/\mathfrak{X}}$, $d^+ := \mathrm{rank}_+ T_{X/\mathfrak{X}}$, $d^- := \mathrm{rank}_- T_{X/\mathfrak{X}}$ and $d^0 := \mathrm{rank}_0 T_{X/\mathfrak{X}}$. We construct an isomorphism

$$\zeta_{\mathfrak{X}, s, o, \Omega}: q_{\mu}^* \varphi_{\mathrm{Grad}(\mathfrak{X}), u^*s, u^*o}[I_{\mathfrak{X}}] \cong q_{\mu}^* \mathrm{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}, s, o})$$

so that the following diagram commutes:

$$\begin{array}{ccc} q_{\mu}^* \varphi_{\mathrm{Grad}(\mathfrak{X}), u^*s, u^*o}[I_{\mathfrak{X}}] & \xrightarrow[\cong]{\zeta_{\mathfrak{X}, s, o, \Omega}} & q_{\mu}^* \mathrm{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}, s, o}) \\ \cong \downarrow \Theta_{q_{\mu}, u^*s, u^*o} & & \cong \downarrow \rho_q \\ \varphi_{X^{\mu}, (q^*s)^{\mu}, (q^*o)^{\mu}}[I_{\mathfrak{X}} - d_0] & \xrightarrow[\cong]{\zeta_{X, q^*s, q^*o}} \mathrm{Loc}_{\mu}(\varphi_{X, q^*s, q^*o})[d^- - d^+ - d_0] & \xrightarrow[\cong]{\Theta_{q, s, o}^{-1}} \mathrm{Loc}_{\mu}(q^* \varphi_{\mathfrak{X}, s, o})[2d^-]. \end{array}$$

As shown in Proposition 5.25, the family $\{q_{\mu}: X^{\mu} \rightarrow \mathrm{Grad}(\mathfrak{X})\}_{\Omega \in \mathrm{Cha}_{\mathfrak{X}}^{\mathbb{G}_m}}$ defines a covering of $\mathrm{Grad}(\mathfrak{X})$. In particular, the complex $\mathrm{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}, s, o})[-I_{\mathfrak{X}}]$ is perverse. Therefore the assignment

$$\mathrm{Sch}_{/\mathrm{Grad}(\mathfrak{X})}^{\mathrm{sm}} \ni (t: T \rightarrow \mathrm{Grad}(\mathfrak{X})) \mapsto \mathrm{Hom}(t^* \varphi_{\mathrm{Grad}(\mathfrak{X}), u^*s, u^*o}[I_{\mathfrak{X}}], t^* \mathrm{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}, s, o}))$$

defines a lisse-étale sheaf. Therefore by using Corollary 5.26, we are reduced to prove the following statement: For a morphism $h: \Omega_1 = (X_1, \mu_1, \bar{q}_1) \rightarrow \Omega_2 = (X_2, \mu_2, \bar{q}_2)$ of \mathbb{G}_m -equivariant charts, the following diagram commutes:

$$\begin{array}{ccc} q_{\mu_1}^* \varphi_{\mathrm{Grad}(\mathfrak{X}), u^*s, u^*o}[I_{\mathfrak{X}}] & \xrightarrow[\cong]{\zeta_{\mathfrak{X}, s, o, \Omega_1}} & q_{\mu_1}^* \mathrm{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}, s, o}) \\ \cong \downarrow & & \cong \downarrow \\ h^{\mathbb{G}_m, *} q_{\mu_2}^* \varphi_{\mathrm{Grad}(\mathfrak{X}), u^*s, u^*o}[I_{\mathfrak{X}}] & \xrightarrow[\cong]{\zeta_{\mathfrak{X}, s, o, \Omega_2}} & h^{\mathbb{G}_m, *} q_{\mu_2}^* \mathrm{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}, s, o}). \end{array}$$

This follows by the construction of $\zeta_{\mathfrak{X}, s, o, \Omega}$, Proposition 7.10, the commutativity of the diagram (4.12) and Lemma 7.14. \square

We now state the unitality property of $\zeta_{\mathfrak{X}, s, o}$. Consider a morphism $\iota: \mathfrak{X} \rightarrow \mathrm{Grad}(\mathfrak{X})$, which is a section of $u: \mathrm{Grad}(\mathfrak{X}) \rightarrow \mathfrak{X}$. Then by the commutativity of the diagram (7.17) as well as Lemma 5.11 we see that the following diagram commutes:

$$(7.22) \quad \begin{array}{ccc} \varphi_{\mathfrak{X}, s, o} & \xrightarrow[\cong]{\zeta_{\mathfrak{X}, s, o}} \iota^* \varphi_{\mathrm{Grad}(\mathfrak{X}), u^*s, u^*o} & \xrightarrow[\cong]{} \varphi_{\mathfrak{X}, s, o} \\ & \searrow \mathrm{id} & \nearrow \end{array}$$

where the latter isomorphism is constructed using $\iota^* u^* s = s$ and the natural isomorphism $\iota^* u^* o \cong o$.

We now prove the associativity of the morphism $\zeta_{\mathfrak{X}, s, o}$.

Proposition 7.17. *The following diagram commutes:*

$$\begin{array}{ccc} \varphi_{\mathrm{Grad}^2(\mathfrak{X})_{\mathrm{mat}}, u^{(2), *}, u_2^* u^* o}[I'_{\mathfrak{X}}] & \xrightarrow[\cong]{\zeta_{\mathfrak{X}, s, o} \circ \zeta_{\mathrm{Grad}(\mathfrak{X}), u^*s, u^*o}} & \mathrm{Loc}_{\mathrm{Grad}(\mathfrak{X})} \circ \mathrm{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}})|_{\mathrm{Grad}^2(\mathfrak{X})_{\mathrm{mat}}} \\ \cong \downarrow & & \cong \downarrow \chi_{\mathfrak{X}} \\ \varphi_{\mathrm{Grad}^2(\mathfrak{X})_{\mathrm{mat}}, u^{(2), *}, u_1^* u^* o}[I'_{\mathfrak{X}}] & \xrightarrow[\cong]{\sigma^*(\zeta_{\mathfrak{X}, s, o} \circ \zeta_{\mathrm{Grad}(\mathfrak{X}), u^*s, u^*o})} & \sigma^* \mathrm{Loc}_{\mathrm{Grad}(\mathfrak{X})} \circ \mathrm{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}})|_{\mathrm{Grad}^2(\mathfrak{X})_{\mathrm{mat}}} \end{array}$$

where we set $I'_{\mathfrak{X}} := u_1^* I_{\mathfrak{X}} + I_{\mathrm{Grad}(\mathfrak{X})}$ and the left vertical isomorphism is induced by the isomorphism of orientations (6.80).

Proof. Take a \mathbb{G}_m^2 -equivariant chart $(X, \hat{\mu}, \bar{q})$ of \mathfrak{X} . Using Lemma 5.28, it is enough to prove the statement after pulling back to $X_{\text{mat}}^{\hat{\mu}}$. Therefore the claim follows from the commutativity of the diagram (6.81), Proposition 7.11 and Lemma 7.15. \square

We now show that the isomorphism $\zeta_{\mathfrak{X}, s, o}$ is compatible with the Thom–Sebastiani isomorphism:

Proposition 7.18. *Let $(\mathfrak{X}, s, o_{\mathfrak{X}})$ and $(\mathfrak{Y}, t, o_{\mathfrak{Y}})$ be oriented d -critical stacks. Assume that \mathfrak{X} and \mathfrak{Y} are quasi-separated with affine stabilizers. Then the following diagram commutes:*

$$\begin{array}{ccc} \varphi_{\text{Grad}(\mathfrak{X}), u_{\mathfrak{X}}^* s, u_{\mathfrak{X}}^* o_{\mathfrak{X}}} \boxtimes \varphi_{\text{Grad}(\mathfrak{Y}), u_{\mathfrak{Y}}^* t, u_{\mathfrak{Y}}^* o_{\mathfrak{Y}}} [I_{\mathfrak{X}} + I_{\mathfrak{Y}}] & \xrightarrow[\cong]{\zeta_{\mathfrak{X}, s, o_{\mathfrak{X}}} \boxtimes \zeta_{\mathfrak{Y}, t, o_{\mathfrak{Y}}}} & \text{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}, s, o_{\mathfrak{X}}}) \boxtimes \text{Loc}_{\mathfrak{Y}}(\varphi_{\mathfrak{Y}, t, o_{\mathfrak{Y}}}) \\ \cong \downarrow \text{TS} & & \cong \downarrow \text{TS} \\ \varphi_{\text{Grad}(\mathfrak{X} \times \mathfrak{Y}), u_{\mathfrak{X} \times \mathfrak{Y}}^* (s \boxplus t), u_{\mathfrak{X} \times \mathfrak{Y}}^* (o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}})} [I_{\mathfrak{X} \times \mathfrak{Y}}] & \xrightarrow[\cong]{\zeta_{\mathfrak{X} \times \mathfrak{Y}, s \boxplus t, o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}}}} & \text{Loc}_{\mathfrak{X} \times \mathfrak{Y}}(\varphi_{\mathfrak{X} \times \mathfrak{Y}, s \boxplus t, o_{\mathfrak{X}} \boxtimes o_{\mathfrak{Y}}}). \end{array}$$

Proof. Take \mathbb{G}_m -equivariant charts (X, μ, \bar{q}) of \mathfrak{X} and (Y, ν, \bar{r}) of \mathfrak{Y} . By the commutativity of the diagram (6.83), Lemma 4.2 (which holds more generally for stacks by construction) and Proposition 7.12, it is enough to prove the commutativity of the following diagram of functors:

$$\begin{array}{ccc} q_{\mu}^* \text{Loc}_{\mathfrak{X}}(-) \boxtimes r_{\nu}^* \text{Loc}_{\mathfrak{Y}}(-) & \xrightarrow{\rho_q \boxtimes \rho_r} & \text{Loc}_{\mu} q^*(-)[2d^-] \boxtimes \text{Loc}_{\nu} r^*(-)[2e^-] \\ \cong \downarrow & & \cong \downarrow \\ (q_{\mu} \times r_{\nu})^* \text{Loc}_{\mathfrak{X} \times \mathfrak{Y}}(- \boxtimes -) & \xrightarrow[\cong]{\rho_{q \times r}} & \text{Loc}_{\mu \times \nu}(q \times r)^*(- \boxtimes -)[2d^- + 2e^-] \end{array}$$

where we set $d^- := \text{rank}_- T_{X/\mathfrak{X}}$ and $e^- = \text{rank}_- T_{Y/\mathfrak{Y}}$. This is obvious from the construction of the map ρ_q . \square

Now let $(\mathfrak{X}, \omega_{\mathfrak{X}}, o)$ be an oriented (-1) -shifted symplectic stack and assume that \mathfrak{X}^{cl} is quasi-separated with affine stabilizers. Then by Proposition 5.30, an isomorphism of orientations (6.85) and equality (6.4), we obtain the following corollary, which summarizes Theorem 7.16, Proposition 7.17 and Proposition 7.18 for the underlying d -critical stacks for (-1) -shifted symplectic stacks:

Corollary 7.19 (Integral isomorphism). *Let $(\mathfrak{X}, \omega_{\mathfrak{X}}, o)$ be a (-1) -shifted symplectic stack such that \mathfrak{X}^{cl} is quasi-separated with affine stabilizers. We equip $\text{Grad}(\mathfrak{X})$ with the (-1) -shifted symplectic structure $u^* \omega_{\mathfrak{X}}$ and the orientation $u^* o$. Set $d := \text{vd Filt}(\mathfrak{X})$. Then there exists a natural isomorphism*

$$\zeta_{\mathfrak{X}} = \zeta_{\mathfrak{X}, \omega_{\mathfrak{X}}, o} : \varphi_{\text{Grad}(\mathfrak{X})}[d] \cong \text{gr}_* \text{ev}^! \varphi_{\mathfrak{X}}$$

which we call the **integral isomorphism** with the following properties:

- (Unitality) : Let $\iota : \mathfrak{X} \hookrightarrow \text{Grad}(\mathfrak{X})$ be a section of $u : \text{Grad}(\mathfrak{X}) \rightarrow \mathfrak{X}$. Then

$$\iota^* \zeta_{\mathfrak{X}} = \text{id}_{\varphi_{\mathfrak{X}}}.$$

- (Associativity) : Set $d' := \text{vd Filt}(\text{Grad}(\mathfrak{X}))$. The following diagram commutes:

$$\begin{array}{ccc} \varphi_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}}[d + d'] & \xrightarrow[\cong]{\zeta_{\text{Grad}(\mathfrak{X})}} & \text{gr}_* \text{ev}^! \varphi_{\text{Grad}(\mathfrak{X})} |_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}}[d] \\ \cong \downarrow & & \cong \downarrow \zeta_{\mathfrak{X}} \\ \sigma^* \varphi_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}}[d + d'] & \xrightarrow[\cong]{\sigma^* \zeta_{\text{Grad}(\mathfrak{X})}} & \text{gr}_* \text{ev}^! \varphi_{\text{Grad}(\mathfrak{X})} |_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}}[d] \xrightarrow[\cong]{\zeta_{\mathfrak{X}}} \text{gr}_*^{(2)} \text{ev}^{(2), !} \varphi_{\mathfrak{X}} |_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}} \end{array}$$

where the left vertical map is constructed using the isomorphism of orientations (6.84).

- (Multiplicativity) : Let $(\mathfrak{Y}, \omega_{\mathfrak{Y}}, o_{\mathfrak{Y}})$ be oriented (-1) -shifted symplectic stack such that \mathfrak{Y}^{cl} is quasi-separated with affine stabilizers. Set $e := \text{vd Filt}(\mathfrak{Y})$. Then the following diagram commutes:

$$\begin{array}{ccc}
\varphi_{\text{Grad}(\mathfrak{X})} \boxtimes \varphi_{\text{Grad}(\mathfrak{Y})}[d+e] & \xrightarrow[\cong]{\zeta_{\mathfrak{X}} \boxtimes \zeta_{\mathfrak{Y}}} & \text{gr}_* \text{ev}^! \varphi_{\mathfrak{X}} \boxtimes \text{gr}_* \text{ev}^! \varphi_{\mathfrak{Y}} \\
\cong \downarrow \text{TS} & & \cong \downarrow \text{TS} \\
\varphi_{\text{Grad}(\mathfrak{X} \times \mathfrak{Y})}[d+e] & \xrightarrow[\cong]{\zeta_{\mathfrak{X} \times \mathfrak{Y}}} & \text{gr}_* \text{ev}^! (\varphi_{\mathfrak{X} \times \mathfrak{Y}}).
\end{array}$$

Proof. The existence of the map $\zeta_{\mathfrak{X}}$ follows from Theorem 7.16 combined with the isomorphism of orientations (6.85). The unitality follows from the commutativity of the diagram (7.22). The associativity follows from Proposition 7.17 together with the commutativity of the diagram (6.86). The multiplicativity follows from Proposition 7.18 together with the commutativity of the diagram (6.88). \square

Remark 7.20 (Other sheaf-theoretic contexts). Though we have proved the statement only for perverse sheaves on the analytic topology, the same argument works for ℓ -adic perverse sheaves over (-1) -shifted symplectic stacks defined over algebraically closed fields of characteristic zero. Also, using the six-functor formalism for mixed Hodge modules recently constructed by Tubach [Tub24], one can upgrade Corollary 7.19 to monodromic mixed Hodge modules with the same proof by replacing shifts with half Lefschetz twists.

7.6. Integral isomorphism for critical loci. We will describe the isomorphism $\zeta_{\mathfrak{X}}$ explicitly in the case of global critical locus. Let \mathfrak{U} be a quasi-separated smooth Artin stack with affine stabilizers. Let $f: \mathfrak{U} \rightarrow \mathbb{A}^1$ be a regular function and define $\mathfrak{X} := \text{Crit}(f)$ and equip it with the (-1) -shifted symplectic structure defined in Example 3.10 and equip it with the canonical orientation $o_{\mathfrak{X}^{\text{cl}}}^{\text{can}}$ in Example 3.20.

Lemma 7.21. *There exists a natural isomorphism*

$$(7.23) \quad \varphi_{\mathfrak{X}, \omega_{\mathfrak{X}}, o_{\mathfrak{X}^{\text{cl}}}^{\text{can}}} \cong \varphi_f(\mathbb{Q}_{\mathfrak{U}}[n]).$$

where we set $n := \dim \mathfrak{U}$.

Proof. We let $(\mathfrak{X}^{\text{cl}}, s)$ denote the underlying d-critical structure of $(\mathfrak{X}, \omega_{\mathfrak{X}})$ and $o_{\mathfrak{X}^{\text{cl}}}^{\text{can}}$ denotes the underlying orientation. Take a smooth cover $\tilde{q}: U \rightarrow \mathfrak{U}$ and let $\tilde{f}: U \rightarrow \mathbb{A}^1$ be the composition $U \rightarrow \mathfrak{U} \rightarrow \mathbb{A}^1$ and set $X := \text{Crit}(\tilde{f})$. We let $q: X^{\text{cl}} \rightarrow \mathfrak{X}^{\text{cl}}$ be the canonical morphism. Then, arguing as the proof of (3.35) and using (3.34), we see that the natural isomorphism $\tilde{q}^* K_{U/\mathfrak{U}} \otimes K_{U/\mathfrak{U}} \cong K_U$ defines an isomorphism of orientations

$$(7.24) \quad o_{X^{\text{cl}}}^{\text{can}} \cong q^* o_{\mathfrak{X}^{\text{cl}}}^{\text{can}}.$$

We let $\mathfrak{R} = (\text{Crit}(f)^{\text{cl}}, \text{id}, U, f, i)$ be the tautological étale d-critical chart for $(X^{\text{cl}}, q^* s)$. Then the $\mathbb{Z}/2\mathbb{Z}$ -local system $Q_{\mathfrak{R}}^o$ is canonically trivial. In particular, we have a natural isomorphism

$$\varphi_{X^{\text{cl}}, q^* s, o_{X^{\text{cl}}}^{\text{can}}} \cong \varphi_{\tilde{f}}.$$

Then we obtain a natural isomorphism

$$q^* \varphi_{\mathfrak{X}^{\text{cl}}, s, o_{\mathfrak{X}^{\text{cl}}}^{\text{can}}} \xrightarrow[\cong]{\Theta_{q, s, o_{\mathfrak{X}^{\text{cl}}}^{\text{can}}}} \varphi_{X^{\text{cl}}, q^* s, o_{X^{\text{cl}}}^{\text{can}}}[-d] \cong \varphi_{\tilde{f}}[-d] \cong q^* \varphi_f(\mathbb{Q}_{\mathfrak{U}}[n])$$

where we set $d := \dim q$ and the final isomorphism is constructed using (4.2). It is clear from the commutativity of the diagram (4.12) that it descends to the desired isomorphism. \square

We keep the notation. By Lemma 5.15, we have a natural equivalence of (-1) -shifted symplectic stacks

$$\text{Grad}(\mathfrak{X}) \cong \text{Crit}(f|_{\text{Grad}(\mathfrak{U})}).$$

By arguing as the construction of the map (6.75), we can construct a natural isomorphism

$$(7.25) \quad o_{\text{Grad}(\mathfrak{X})}^{\text{can}} \cong u^* o_{\mathfrak{X}}^{\text{can}}.$$

By arguing as Example 7.6, we obtain a natural isomorphism

$$(7.26) \quad \text{Loc}_{\mathfrak{U}}(\mathbb{Q}_{\mathfrak{U}}) \cong \mathbb{Q}_{\text{Grad}(\mathfrak{U})}[-2n^-]$$

where we set $n^- := \text{rank } T_{\mathfrak{U}}|_{\text{Grad}(\mathfrak{U})}^-$. With these preparations, we can explicitly describe $\zeta_{\mathfrak{X}, \omega_{\mathfrak{X}}, o_{\mathfrak{X}^{\text{cl}}}^{\text{can}}}$:

Proposition 7.22. *Set $f_0 := f|_{\text{Grad}(\mathfrak{U})}$, $n := \dim \mathfrak{U}$ and $n_0 := \dim \text{Grad}(\mathfrak{U})$. Then the following diagram commutes:*

$$(7.27) \quad \begin{array}{ccc} \varphi_{\text{Grad}(\mathfrak{X}), u^* \omega_{\mathfrak{X}}, u^* o_{\mathfrak{X}}^{\text{can}}}[\text{vd Filt}(\mathfrak{X})] & \xrightarrow[\cong]{\zeta_{\mathfrak{X}, \omega_{\mathfrak{X}}, o_{\mathfrak{X}}^{\text{can}}}} & \text{Loc}_{\mathfrak{X}}(\varphi_{\mathfrak{X}, \omega_{\mathfrak{X}}, o_{\mathfrak{X}}^{\text{can}}}) \\ \cong \downarrow (7.23) & & \cong \downarrow (7.23) \\ \varphi_{f_0}(\mathbb{Q}_{\text{Grad}(\mathfrak{U})})[\text{vd Filt}(\mathfrak{X}) + n_0] & \xrightarrow[\cong]{(7.26)} \varphi_{f_0}(\text{Loc}_{\mathfrak{U}}(\mathbb{Q}_{\mathfrak{U}}))[n] & \xrightarrow[\cong]{(4.2)} \text{Loc}_{\mathfrak{X}}(\varphi_f(\mathbb{Q}_{\mathfrak{U}}))[n] \end{array}$$

where the left vertical map is constructed using the identification (7.25).

Proof. Take a \mathbb{G}_m -equivariant chart (U, μ, \tilde{q}) of \mathfrak{U} . It is enough to prove the claim after pulling back the diagram to X^μ . Set $\tilde{f} := f \circ q$, $\tilde{f}_0 := f|_{U^\mu}$ and $X := \text{Crit}(f)$. Let $q: X^{\text{cl}} \rightarrow \mathfrak{X}^{\text{cl}}$ be the natural morphism. Arguing as the proof of the commutativity of the diagram (6.76), we see that the following diagram commutes:

$$\begin{array}{ccccc} (q^* o_{\mathfrak{X}^{\text{cl}}}^{\text{can}})^\mu & \xrightarrow[\cong]{(7.24)} & (o_{X^{\text{cl}}}^{\text{can}})^\mu & \xrightarrow[\cong]{(7.14)} & o_{X^{\mu, \text{cl}}}^{\text{can}} \\ \cong \downarrow (6.79) & & & & \parallel \\ q_\mu^* u^* o_{\mathfrak{X}^{\text{cl}}}^{\text{can}} & \xrightarrow[\cong]{(7.25)} & q_\mu^* o_{\text{Grad}(\mathfrak{X})^{\text{cl}}}^{\text{can}} & \xrightarrow[\cong]{(7.24)} & o_{X^{\mu, \text{cl}}}^{\text{can}} \end{array}$$

Using the commutativity of this diagram and the fact that each map appearing in the diagram (7.27) commutes with smooth pullback (which follows immediately from the construction), it is enough to prove the commutativity of the following diagram:

$$\begin{array}{ccc} \varphi_{X^{\mu, \text{cl}}, \tilde{s}^\mu, (o_{X^{\text{cl}}}^{\text{can}})^\mu}[I_{X^{\text{cl}}}^\mu] & \xrightarrow[\cong]{\zeta_{\mu, \tilde{s}, o_{X^{\text{cl}}}^{\text{can}}}} & \text{Loc}_\mu(\varphi_{X^{\text{cl}}, \tilde{s}, o_{X^{\text{cl}}}^{\text{can}}}) \\ \cong \downarrow & & \cong \downarrow \\ \varphi_{\tilde{f}_0}(\mathbb{Q}_{U^\mu})[I_{X^{\text{cl}}}^\mu + m_0] & \xrightarrow[\cong]{(7.5)} \varphi_{\tilde{f}_0}(\text{Loc}_\mu(\mathbb{Q}_U))[m] & \xrightarrow[\cong]{(4.2)} \text{Loc}_\mu(\varphi_{\tilde{f}}(\mathbb{Q}_U))[m] \end{array}$$

where we set $m := \dim U$, $m_0 := \dim U^\mu$, \tilde{s} is the canonical d-critical structure on X^{cl} defined using the critical locus description and the left vertical map is constructed using the natural isomorphism $(o_{X^{\text{cl}}}^{\text{can}})^\mu \cong o_{X^{\mu, \text{cl}}}^{\text{can}}$. This is immediate from the construction of $\zeta_{\mu, \tilde{s}, o_{X^{\text{cl}}}^{\text{can}}}$ in Theorem 7.9. \square

8. APPLICATIONS TO THE MODULI OF OBJECTS IN 3CY CATEGORIES

In the previous sections, we have established a hyperbolic localization statement for d-critical stacks. In this section, we explain how to use this to define a cohomological Hall algebra structure for Calabi–Yau categories.

In §8.2, we will construct the cohomological Hall algebras for a moduli-like stack introduced in §8.1. In §8.3 — §8.5, we will show that the setup in §8.2 can be used to construct the cohomological Hall algebras for 3-Calabi–Yau categories with strong orientation data. In §8.6, we will give examples of 3-Calabi–Yau categories with strong orientation data using deformed Calabi–Yau completion. In §8.7, we will show that our construction recovers Kontsevich–Soibelman’s cohomological Hall algebras for quivers with potentials.

8.1. Moduli-like stacks. In this section we introduce the geometric setting necessary to define the cohomological Hall algebra. It will be given by a (-1) -shifted symplectic stack with a $B\mathbb{G}_m$ -action and an addition map satisfying certain conditions. For a $B\mathbb{G}_m$ -action on a stack \mathfrak{X} , let $\Psi: B\mathbb{G}_m \times \mathfrak{X} \rightarrow \mathfrak{X}$ be the corresponding action map. For $n \in \mathbb{Z}$ we obtain maps

$$\iota_n: \mathfrak{X} \longrightarrow \text{Grad}(\mathfrak{X})$$

adjoint to

$$B\mathbb{G}_m \times \mathfrak{X} \xrightarrow{(-)^n \times \text{id}} B\mathbb{G}_m \times \mathfrak{X} \xrightarrow{\Psi} \mathfrak{X},$$

where the first map is induced by the n -th power map on \mathbb{G}_m . For instance, $\iota_0 = \iota: \mathfrak{X} \rightarrow \text{Grad}(\mathfrak{X})$ is induced by the projection $B\mathbb{G}_m \rightarrow \text{pt}$. Similarly, for $(n, m) \in \mathbb{Z}$ we obtain maps

$$\iota_{n,m}: \mathfrak{X} \rightarrow \text{Grad}^2(\mathfrak{X})$$

adjoint to

$$B\mathbb{G}_m \times B\mathbb{G}_m \times \mathfrak{X} \xrightarrow{(-)^n \times (-)^m \times \text{id}} B\mathbb{G}_m \times B\mathbb{G}_m \times \mathfrak{X} \xrightarrow{\text{id} \times \Psi} B\mathbb{G}_m \times \mathfrak{X} \xrightarrow{\Psi} \mathfrak{X}.$$

Definition 8.1. A *moduli-like stack* is a derived stack \mathfrak{M} which is locally geometric and locally of finite presentation, together with the following data:

- (1) A $B\mathbb{G}_m$ -action on \mathfrak{M} .
- (2) An \mathbb{E}_∞ -structure on \mathfrak{M} commuting with the $B\mathbb{G}_m$ -action. We denote by $\Phi_n: \mathfrak{M}^n \rightarrow \mathfrak{M}$ the corresponding addition maps and by $\eta: \text{pt} \rightarrow \mathfrak{M}$ the unit map.
- (3) A (-1) -shifted symplectic structure ω on \mathfrak{M} . We require it to be additive, in the sense that there is a homotopy $\Phi_2^* \omega \sim \omega \boxplus \omega$ of (-1) -shifted closed two-forms on $\mathfrak{M} \times \mathfrak{M}$ and a homotopy of homotopies of (-1) -shifted closed two-forms witnessing commutativity of the pullbacks along

$$\begin{array}{ccc} \mathfrak{M} \times \mathfrak{M} \times \mathfrak{M} & \xrightarrow{\text{id} \times \Phi_2} & \mathfrak{M} \times \mathfrak{M} \\ \downarrow \Phi_2 \times \text{id} & & \downarrow \Phi_2 \\ \mathfrak{M} \times \mathfrak{M} & \xrightarrow{\Phi_2} & \mathfrak{M} \end{array}$$

These data are required to satisfy the following conditions:

- (1) The composites

$$\tilde{\Phi}_2: \mathfrak{M}^2 \xrightarrow{\iota_0 \times \iota_1} \text{Grad}(\mathfrak{M}^2) \xrightarrow{\text{Grad}(\Phi_2)} \text{Grad}(\mathfrak{M})$$

and

$$\tilde{\Phi}_3: \mathfrak{M}^3 \xrightarrow{\iota_{0,0} \times \iota_{1,0} \times \iota_{1,1}} \text{Grad}^2(\mathfrak{M}^3) \xrightarrow{\text{Grad}^2(\Phi_3)} \text{Grad}^2(\mathfrak{M})$$

are open immersions.

- (2) The map $\eta: \text{pt} \rightarrow \mathfrak{M}$ is an open immersion.
- (3) The image of $\tilde{\Phi}_3$ lands in the matching locus of $\text{Grad}^2(\mathfrak{M})$.
- (4) The base change of $\text{Filt}(\mathfrak{M}^{\text{cl}}) \rightarrow \text{Grad}(\mathfrak{M}^{\text{cl}})$ along $\iota_n: \mathfrak{M}^{\text{cl}} \rightarrow \text{Grad}(\mathfrak{M}^{\text{cl}})$ is an isomorphism.

Remark 8.2. As we will explain in Proposition 8.29, a natural example of a moduli-like stack is the derived moduli stack of objects in a 3CY category.

Remark 8.3. A structure similar to that of a moduli-like stack was introduced in [Joy18, Assumption 3.1].

Now let \mathfrak{M} be a moduli-like stack. By Corollary 5.18 we have a (-1) -shifted Lagrangian structure on the attractor correspondence $\text{Grad}(\mathfrak{M}) \xleftarrow{\text{gr}} \text{Filt}(\mathfrak{M}) \xrightarrow{\text{ev}} \mathfrak{M}$. Its pullback along $\tilde{\Phi}_2: \mathfrak{M}^2 \rightarrow \text{Grad}(\mathfrak{M})$ gives a (-1) -shifted Lagrangian correspondence

$$(8.1) \quad \begin{array}{ccc} & \mathfrak{M}^{2\text{-filt}} & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \mathfrak{M}^2 & & \mathfrak{M} \end{array}$$

together with a splitting $\mathfrak{M}^2 \rightarrow \mathfrak{M}^{2\text{-filt}}$ of the map $\text{gr}: \mathfrak{M}^{2\text{-filt}} \rightarrow \mathfrak{M}^2$. Let

$$\chi: \mathfrak{M} \times \mathfrak{M} \rightarrow \mathbb{Z}$$

be the locally-constant function obtained by restricting the index function $I_{\mathfrak{M}}: \text{Grad}(\mathfrak{M}) \rightarrow \mathbb{Z}$ along $\tilde{\Phi}_2$.

Using Lemma 3.19 the above Lagrangian correspondence gives an isomorphism

$$(8.2) \quad \Phi_{2\text{or}\mathfrak{M}}^* \cong \text{or}_{\mathfrak{M}} \boxtimes \text{or}_{\mathfrak{M}}$$

of graded μ_2 -gerbes on \mathfrak{M}^2 . In particular, if o is an orientation of \mathfrak{M} , there is a canonical orientation $\Phi_2^* o$ of \mathfrak{M}^2 .

Moreover, by Corollary 5.23 we obtain that there is an equivalence between the composite Lagrangian correspondences

$$\begin{array}{ccccc}
 & \mathfrak{M} \times \mathfrak{M}^{2\text{-filt}} & & \mathfrak{M}^{2\text{-filt}} & \\
 \text{id} \times \text{gr} \swarrow & & \text{id} \times \text{ev} \searrow & \text{gr} \swarrow & \text{ev} \searrow \\
 \mathfrak{M}^3 & & \mathfrak{M}^2 & & \mathfrak{M}
 \end{array}$$

and

$$\begin{array}{ccccc}
 & \mathfrak{M}^{2\text{-filt}} \times \mathfrak{M} & & \mathfrak{M}^{2\text{-filt}} & \\
 \text{gr} \times \text{id} \swarrow & & \text{ev} \times \text{id} \searrow & \text{gr} \swarrow & \text{ev} \searrow \\
 \mathfrak{M}^3 & & \mathfrak{M}^2 & & \mathfrak{M}
 \end{array}$$

In particular, we get that χ satisfies a cocycle relation:

$$(8.3) \quad (\Phi_2^* \boxtimes \text{id})\chi + \text{pr}_{12}^* \chi = (\text{id} \boxtimes \Phi_2^*)\chi + \text{pr}_{23}^* \chi.$$

We denote the composite Lagrangian correspondence by

$$(8.4) \quad \begin{array}{ccc} & \mathfrak{M}^{3\text{-filt}} & \\ \text{gr}^{(2)} \swarrow & & \text{ev}^{(2)} \searrow \\ \mathfrak{M}^3 & & \mathfrak{M} \end{array}$$

Using (3.9) we obtain 2-isomorphisms witnessing commutativity of the diagram

$$(8.5) \quad \begin{array}{ccc} (\Phi_2^* \boxtimes \text{id})\Phi_2^* \text{or}_{\mathfrak{M}} & \xrightarrow{\sim} & (\text{id} \boxtimes \Phi_2^*)\Phi_2^* \text{or}_{\mathfrak{M}} \\ \downarrow (8.2) & & \downarrow (8.2) \\ (\Phi_2^* \text{or}_{\mathfrak{M}}) \boxtimes \text{or}_{\mathfrak{M}} & & \text{or}_{\mathfrak{M}} \boxtimes (\Phi_2^* \text{or}_{\mathfrak{M}}) \\ & \searrow (8.2) \quad \swarrow (8.2) & \\ & \text{or}_{\mathfrak{M}} \boxtimes \text{or}_{\mathfrak{M}} \boxtimes \text{or}_{\mathfrak{M}} & \end{array}$$

In particular, if o is an orientation of \mathfrak{M} , there is an isomorphism of orientations

$$(8.6) \quad (\Phi_2^* \boxtimes \text{id})\Phi_2^* o \cong (\text{id} \boxtimes \Phi_2^*)\Phi_2^* o$$

of \mathfrak{M}^3 .

Definition 8.4. Let \mathfrak{M} be a moduli-like stack. An **orientation data on \mathfrak{M}** is given by the following data⁷:

- (1) An orientation o of \mathfrak{M} .
- (2) An isomorphism of orientations $\lambda: \Phi_2^* o \cong o \boxtimes o$.
- (3) An isomorphism of orientations $\tau: \eta^* o \cong o_{\text{triv}}$, where o_{triv} is the trivial orientation of pt .

An orientation data is called a **strong orientation data** if the diagrams

$$\begin{array}{ccc}
 (\Phi_2^* \boxtimes \text{id})\Phi_2^* o & \xrightarrow[\sim]{(8.6)} & (\text{id} \boxtimes \Phi_2^*)\Phi_2^* o \\
 \downarrow (\Phi_2^* \boxtimes \text{id})\lambda & & \downarrow (\text{id} \boxtimes \Phi_2^*)\lambda \\
 (\Phi_2^* o) \boxtimes o & & o \boxtimes (\Phi_2^* o) \\
 \searrow \lambda \boxtimes \text{id} & & \swarrow \text{id} \boxtimes \lambda \\
 & o \boxtimes o \boxtimes o &
 \end{array}$$

⁷Our definition of the orientation data slightly differs from the one in [JU21, Definition 4.2], where they defined the orientation data to be an isomorphism class of o such that the isomorphisms λ and τ exist.

and

$$\begin{array}{ccc}
(\eta^* \boxtimes \text{id})\Phi_2^* o \xrightarrow{(\eta^* \boxtimes \text{id})^\lambda} \eta^* o \boxtimes o & & (\text{id} \boxtimes \eta^*)\Phi_2^* o \xrightarrow{(\text{id} \boxtimes \eta^*)^\lambda} o \boxtimes \eta^* o \\
\sim \downarrow & & \sim \downarrow \\
o \xrightarrow{\sim} o_{\text{triv}} \boxtimes o & & o \xrightarrow{\sim} o \boxtimes o_{\text{triv}}
\end{array}$$

commute.

Remark 8.5. Let X be a smooth projective Calabi–Yau threefold and $\mathfrak{M}_{\text{Perf}(X)}$ the moduli stack of perfect complexes on X . In [JU21], Joyce and Upmeyer constructed an orientation data for $\mathfrak{M}_{\text{Perf}(X)}$. However, it is not known whether their orientation data upgrades to a strong orientation data.

8.2. Cohomological Hall algebra. As an application of Corollary 7.19, we will construct the cohomological Hall algebra for moduli-like stacks \mathfrak{M} with a strong orientation data.

Set $\Gamma := \pi_0(\mathfrak{M})$ and write the connected component decomposition by

$$\mathfrak{M} = \coprod_{\alpha \in \Gamma} \mathfrak{M}_\alpha.$$

Since $\text{gr}: \text{Filt}(\mathfrak{M}) \rightarrow \text{Grad}(\mathfrak{M})$ is an \mathbb{A}^1 -deformation retract by [Hal14, Lemma 1.3.8], so is $\mathfrak{M}^{n\text{-filt}} \rightarrow \mathfrak{M}^n$ for $n = 2, 3$. Therefore, $\pi_0(\mathfrak{M}^{n\text{-filt}}) \cong \pi_0(\mathfrak{M}^n) \cong \pi_0(\mathfrak{M})^n$, where the first isomorphism follows from [Hal14, Lemma 1.3.7] and the second isomorphism follows from the assumption that \mathfrak{M} is locally of finite presentation. Let

$$\mathfrak{M}^{n\text{-filt}} = \coprod_{\alpha_1, \dots, \alpha_n \in \Gamma} \mathfrak{M}_{\alpha_1, \dots, \alpha_n}^{n\text{-filt}}$$

be the connected component decomposition.

The \mathbb{E}_∞ -structure on \mathfrak{M} induces an abelian monoid structure on Γ . The Lagrangian correspondences (8.1) and (8.4) restrict to the following correspondences:

$$\begin{array}{ccc}
& \mathfrak{M}_{\alpha_1, \alpha_2}^{2\text{-filt}} & \\
\text{gr} \swarrow & & \searrow \text{ev} \\
\mathfrak{M}_{\alpha_1} \times \mathfrak{M}_{\alpha_2} & & \mathfrak{M}_{\alpha_1 + \alpha_2}
\end{array}
\quad
\begin{array}{ccc}
& \mathfrak{M}_{\alpha_1, \alpha_2, \alpha_3}^{3\text{-filt}} & \\
\text{gr}^{(2)} \swarrow & & \searrow \text{ev}^{(2)} \\
\mathfrak{M}_{\alpha_1} \times \mathfrak{M}_{\alpha_2} \times \mathfrak{M}_{\alpha_3} & & \mathfrak{M}_{\alpha_1 + \alpha_2 + \alpha_3}
\end{array}$$

By (8.3) the index function $I_{\mathfrak{M}}: \text{Grad}(\mathfrak{M}) \rightarrow \mathbb{Z}$ restricts under the inclusion $\tilde{\Phi}_2: \mathfrak{M}^2 \rightarrow \text{Grad}(\mathfrak{M})$ to a 2-cocycle $\chi: \Gamma \times \Gamma \rightarrow \mathbb{Z}$.

Theorem 8.6. *Let \mathfrak{M} be a moduli-like stack with an orientation data such that \mathfrak{M}^{cl} is quasi-separated and has affine stabilizers. Then there exists a natural isomorphism*

$$\zeta_{\alpha_1, \alpha_2}: \varphi_{\mathfrak{M}_{\alpha_1}} \boxtimes \varphi_{\mathfrak{M}_{\alpha_2}} [\chi(\alpha_1, \alpha_2)] \cong \text{gr}_* \text{ev}^! \varphi_{\mathfrak{M}_{\alpha_1 + \alpha_2}}$$

with the following properties:

- (Unitality) *Let $0 \in \Gamma$ be the unit element. Then the maps*

$$\begin{aligned}
\varphi_{\mathfrak{M}_\alpha} &\cong \varphi_{\mathfrak{M}_\alpha} \boxtimes \mathbb{Q} \cong \varphi_{\mathfrak{M}_\alpha} \boxtimes \eta^* \varphi_{\mathfrak{M}_0} \xrightarrow{\cong} \varphi_{\mathfrak{M}_\alpha}, \\
\varphi_{\mathfrak{M}_\alpha} &\cong \mathbb{Q} \boxtimes \varphi_{\mathfrak{M}_\alpha} \cong \eta^* \varphi_{\mathfrak{M}_0} \boxtimes \varphi_{\mathfrak{M}_\alpha} \xrightarrow{\cong} \varphi_{\mathfrak{M}_\alpha}
\end{aligned}$$

are the identities.

- (Associativity) *Assume further that our orientation data is a strong orientation data. Then the following diagram commutes:*

$$\begin{array}{ccc}
\varphi_{\mathfrak{M}_{\alpha_1}} \boxtimes \varphi_{\mathfrak{M}_{\alpha_2}} \boxtimes \varphi_{\mathfrak{M}_{\alpha_3}} [\chi(\alpha_1, \alpha_2, \alpha_3)] & \xrightarrow[\cong]{\zeta_{\alpha_1, \alpha_2} \boxtimes \text{id}} & (\text{gr}_* \text{ev}^! \varphi_{\mathfrak{M}_{\alpha_1 + \alpha_2}}) \boxtimes \varphi_{\mathfrak{M}_{\alpha_3}} [\chi(\alpha_1 + \alpha_2, \alpha_3)] \\
\cong \downarrow \text{id} \boxtimes \zeta_{\alpha_2, \alpha_3} & & \cong \downarrow \zeta_{\alpha_1 + \alpha_2, \alpha_3} \\
\varphi_{\mathfrak{M}_{\alpha_1}} \boxtimes (\text{gr}_* \text{ev}^! \varphi_{\mathfrak{M}_{\alpha_2 + \alpha_3}}) [\chi(\alpha_1, \alpha_2 + \alpha_3)] & \xrightarrow[\cong]{\zeta_{\alpha_1, \alpha_2 + \alpha_3}} & \text{gr}_*^{(2)} \text{ev}^{(2)!} \varphi_{\mathfrak{M}_{\alpha_1 + \alpha_2 + \alpha_3}},
\end{array}$$

where we set $\chi(\alpha_1, \alpha_2, \alpha_3) := \chi(\alpha_1, \alpha_2) + \chi(\alpha_1 + \alpha_2, \alpha_3) = \chi(\alpha_2, \alpha_3) + \chi(\alpha_1, \alpha_2 + \alpha_3)$.

Proof. Consider the inclusion $\tilde{\Phi}_2: \mathfrak{M}_{\alpha_1} \times \mathfrak{M}_{\alpha_2} \rightarrow \text{Grad}(\mathfrak{M})$. By assumption, there exists a natural equivalence of symplectic structures $\omega_{\mathfrak{M}_{\alpha_1}} \boxplus \omega_{\mathfrak{M}_{\alpha_2}} \sim \tilde{\Phi}_2^* u^* \omega_{\mathfrak{M}}$. Also, by the definition of the orientation data and Remark 6.12, we have a natural isomorphism of orientations $o_{\alpha_1} \boxtimes o_{\alpha_2} \cong \tilde{\Phi}_2^* u^* o$, where o denotes the orientation for \mathfrak{M} and o_α is its restriction to \mathfrak{M}_α . Therefore, by using Corollary 4.4 and Corollary 7.19, we obtain an isomorphism $\zeta_{\alpha_1, \alpha_2}$ by the composition

$$\varphi_{\mathfrak{M}_{\alpha_1}, o_{\alpha_1}} \boxtimes \varphi_{\mathfrak{M}_{\alpha_2}, o_{\alpha_2}} [\chi(\alpha_1, \alpha_2)] \xrightarrow[\cong]{\text{TS}} \varphi_{\mathfrak{M}_{\alpha_1} \times \mathfrak{M}_{\alpha_2}, \tilde{\Phi}_2^* u^* o} [\chi(\alpha_1, \alpha_2)] \xrightarrow[\cong]{\zeta_{\mathfrak{M}_{\alpha_1 + \alpha_2}}} \text{gr}_! \text{ev}^* \varphi_{\mathfrak{M}_{\alpha_1 + \alpha_2}, o}.$$

The unitality of $\zeta_{\alpha_1, \alpha_2}$ is an immediate consequence of the unitality property of $\zeta_{\mathfrak{X}, \omega_{\mathfrak{X}}, o}$ proved in Corollary 7.19.

For the associativity of $\zeta_{\alpha_1, \alpha_2}$, consider the inclusion $\tilde{\Phi}_3: \mathfrak{M}_{\alpha_1} \times \mathfrak{M}_{\alpha_2} \times \mathfrak{M}_{\alpha_3} \rightarrow \text{Grad}^2(\mathfrak{M})$ and let $\tilde{\Phi}'_3: \mathfrak{M}_{\alpha_1} \times \mathfrak{M}_{\alpha_2} \times \mathfrak{M}_{\alpha_3} \rightarrow \text{Grad}^2(\mathfrak{M})$ its post-composition with the swapping isomorphism $\sigma: \text{Grad}^2(\mathfrak{M}) \rightarrow \text{Grad}^2(\mathfrak{M})$. By the associativity property of $\zeta_{\mathfrak{X}, \omega_{\mathfrak{X}}, o}$ proved in Corollary 7.19, we see that the following diagram commutes:

(8.7)

$$\begin{array}{ccc} \varphi_{\mathfrak{M}_{\alpha_1} \times \mathfrak{M}_{\alpha_2} \times \mathfrak{M}_{\alpha_3}, \tilde{\Phi}_3^* u_1^* u^* o} [n_{123}] & \xrightarrow[\cong]{\zeta_{\mathfrak{M}_{\alpha_1 + \alpha_2} \times \mathfrak{M}_{\alpha_3}}} & (\text{gr} \times \text{id})_* (\text{ev} \times \text{id})^! \varphi_{\mathfrak{M}_{\alpha_1 + \alpha_2}, \tilde{\Phi}_2^* u^* o} \boxtimes \varphi_{\mathfrak{M}_{\alpha_3}, o_{\alpha_3}} [n_{12,3}] \\ \cong \downarrow & & \downarrow \cong \\ \varphi_{\mathfrak{M}_{\alpha_1} \times \mathfrak{M}_{\alpha_2} \times \mathfrak{M}_{\alpha_3}, \tilde{\Phi}'_3^* u_1^* u^* o} [n_{123}] & & \zeta_{\mathfrak{M}_{\alpha_1 + \alpha_2 + \alpha_3}} \\ \cong \downarrow \zeta_{\mathfrak{M}_{\alpha_1} \times \mathfrak{M}_{\alpha_2 + \alpha_3}} & & \downarrow \\ (\text{id} \times \text{gr})_* (\text{id} \times \text{ev})^! \varphi_{\mathfrak{M}_{\alpha_1}, o_{\alpha_1}} \boxtimes \varphi_{\mathfrak{M}_{\alpha_2 + \alpha_3}, \tilde{\Phi}_2^* u^* o} [n_{1,23}] & \xrightarrow[\cong]{\zeta_{\mathfrak{M}_{\alpha_1 + \alpha_2 + \alpha_3}}} & \text{gr}^{(2)}_* \text{ev}^{(2), !} \varphi_{\mathfrak{M}_{\alpha_1 + \alpha_2 + \alpha_3}, o}. \end{array}$$

Here we set $n_{123} := \chi(\alpha_1, \alpha_2, \alpha_3)$, $n_{12,3} := \chi(\alpha_1 + \alpha_2, \alpha_3)$ and $n_{1,23} := \chi(\alpha_1, \alpha_2 + \alpha_3)$ and the upper left vertical map is induced from the isomorphism of orientations (6.84). By the definition of the strong orientation data and the associativity statement in Remark 6.12, the following diagram of orientations commutes:

$$\begin{array}{ccccc} (o_{\alpha_1} \boxtimes o_{\alpha_2}) \boxtimes o_{\alpha_3} & \xrightarrow{\cong} & (\tilde{\Phi}_2 \times \text{id})^* (u^* \boxtimes \text{id}^*) (o_{\alpha_1 + \alpha_2} \boxtimes o_{\alpha_3}) & \xrightarrow{\cong} & \tilde{\Phi}_3^* u_1^* u^* o \\ \cong \downarrow & & & & \downarrow \cong \\ o_{\alpha_1} \boxtimes (o_{\alpha_2} \boxtimes o_{\alpha_3}) & \xrightarrow{\cong} & (\text{id} \times \tilde{\Phi}_2)^* (\text{id}^* \boxtimes u^*) (o_{\alpha_1} \boxtimes o_{\alpha_2 + \alpha_3}) & \xrightarrow{\cong} & \tilde{\Phi}'_3^* u_1^* u^* o. \end{array}$$

Therefore the statement holds by the commutativity of the diagram (8.7), the associativity of the Thom–Sebastiani isomorphism in Corollary 4.4 together with the unitality and the multiplicativity of the integral isomorphism in Corollary 7.19. \square

Remark 8.7. If the orientation data is not strong, it follows from the proof of Theorem 8.6 that the associativity property of the map $\zeta_{\alpha_1, \alpha_2}$ holds true up to some choice of sign.

For Γ -graded dg vector spaces V and W , we define the twisted tensor product by

$$V \otimes^{\text{tw}} W := \bigoplus_{\alpha, \beta \in \Gamma} V_\alpha \otimes W_\beta [-\chi(\alpha, \beta)].$$

It defines a monoidal structure on the ∞ -category of Γ -graded complexes. The following corollary is an immediate consequence of Theorem 8.6:

Corollary 8.8. *Let \mathfrak{M} be a moduli-like stack with a strong orientation data satisfying the following conditions:*

- \mathfrak{M}^{cl} is a quasi-separated Artin stack with affine stabilizers.
- \mathfrak{M}^{cl} is Θ -reductive, i.e. the map $\text{ev}: \text{Filt}(\mathfrak{M}^{\text{cl}}) \rightarrow \mathfrak{M}^{\text{cl}}$ is proper over each connected component of the source.

- $\eta: \text{pt} \rightarrow \mathfrak{M}^{\text{cl}}$ is a closed immersion.

For each $\alpha \in \Gamma$, set $\mathcal{H}_\alpha := H^\bullet(\mathfrak{M}_\alpha, \varphi_{\mathfrak{M}_\alpha})$ and $\mathcal{H} := \bigoplus_{\alpha \in \Gamma} \mathcal{H}_\alpha$. Then there are natural morphisms

$$*^{\text{Hall}}: \mathcal{H} \otimes^{\text{tw}} \mathcal{H} \rightarrow \mathcal{H}, \quad \mathbb{Q} \rightarrow \mathcal{H}_0$$

which define an associative algebra, the **cohomological Hall algebra** for \mathfrak{M} .

Example 8.9. Let X be a smooth Calabi–Yau threefold and set $\mathcal{C} = \text{Perf}(X)$. In this case, points in $\mathfrak{M}_{\mathcal{C}}$ correspond to compactly supported perfect complexes on X . Let $\mathfrak{M} \subset \mathfrak{M}_{\mathcal{C}}$ be the locus consisting of compactly supported coherent sheaves (i.e., complexes concentrated in degree zero). As we will see in Proposition 8.29, \mathfrak{M} admits a moduli-like structure in a natural way. Assume that \mathfrak{M} admits an orientation data: this is satisfied for projective X by [JU21] and for local curves and surfaces by Corollary 8.36. For each $\gamma \in H_*(X)$, set

$$\mathfrak{M}_\gamma := \{[E] \in \mathfrak{M} \mid \text{ch}(E) = \gamma\}, \quad \mathcal{H}_\gamma := H^\bullet(\mathfrak{M}_\gamma, \varphi_{\mathfrak{M}_\gamma}).$$

Set $\mathcal{H}_X := \bigoplus_\gamma \mathcal{H}_\gamma$. Then Corollary 8.8 implies that there exists a $H_*(X)$ -graded multiplication

$$*^{\text{Hall}}: \mathcal{H}_X \otimes^{\text{tw}} \mathcal{H}_X \rightarrow \mathcal{H}_X$$

which is associative up to some choice of sign by Remark 8.7. If the orientation data is strong (which is satisfied for local curves and local surfaces as we will see in Corollary 8.36), the multiplication $*^{\text{Hall}}$ defines an associative algebra structure on \mathcal{H}_X .

8.3. Moduli of objects in dg categories. Let $\mathbf{dgc}at$ be the ∞ -category of \mathbb{C} -linear idempotent complete small stable ∞ -categories and exact functors; equivalently it is the ∞ -category of \mathbb{C} -linear dg categories localized along Morita equivalences [Coh, Corollary 5.7]. We will simply refer to objects of $\mathbf{dgc}at$ as dg categories. For $\mathcal{C} \in \mathbf{dgc}at$ we denote by $\text{Fun}^{\text{ex}}(\mathcal{C}, -)$ the stable ∞ -category of \mathbb{C} -linear exact functors. There is a natural symmetric monoidal structure on $\mathbf{dgc}at$ with $\text{Perf}_{\mathbb{C}} \subset \text{Mod}_{\mathbb{C}}$, the ∞ -category of perfect complexes of \mathbb{C} -vector spaces, the unit. Binary coproducts and binary products in $\mathbf{dgc}at$ coincide, and we denote them by $\mathcal{C} \oplus \mathcal{D}$.

Lemma 8.10. *Let P be a finite poset. Then its stable envelope is $\text{Perf}(P^{\text{op}}) = \text{Fun}(P^{\text{op}}, \text{Perf}_{\mathbb{C}})$. In other words, restriction along the Yoneda embedding $P \rightarrow \text{Perf}(P^{\text{op}})$ given by $p \mapsto \mathbb{C}[\text{Hom}_P(-, p)]$ induces an equivalence*

$$\text{Fun}^{\text{ex}}(\text{Perf}(P^{\text{op}}), \mathcal{C}) \longrightarrow \text{Fun}(P, \mathcal{C})$$

for any $\mathcal{C} \in \mathbf{dgc}at$.

Proof. Let $\mathcal{C} \in \mathbf{dgc}at$. Its ind-completion $\text{Ind}(\mathcal{C}) \in \text{Mod}_{\text{Mod}_{\mathbb{C}}}(\text{Pr}^L)$ is a stable presentable \mathbb{C} -linear ∞ -category. Thus, by [Lur09, Theorem 5.1.5.6] restriction along the Yoneda embedding induces an equivalence

$$\text{Fun}^L(\text{Fun}(P^{\text{op}}, \mathcal{S}), \text{Ind}(\mathcal{C})) \longrightarrow \text{Fun}(P, \text{Ind}(\mathcal{C})),$$

where on the left we consider the ∞ -category of functors preserving small colimits. By [Aok23, Corollary 2.2] the natural functor $\text{Fun}(P^{\text{op}}, \mathcal{S}) \otimes \text{Mod}_{\mathbb{C}} \rightarrow \text{Fun}(P^{\text{op}}, \text{Mod}_{\mathbb{C}})$ is an equivalence, and so we get that the restriction functor

$$\text{Fun}^{L, \mathbb{C}}(\text{Fun}(P^{\text{op}}, \mathcal{S}) \otimes \text{Mod}_{\mathbb{C}}, \text{Ind}(\mathcal{C})) \longrightarrow \text{Fun}^L(\text{Fun}(P^{\text{op}}, \mathcal{S}), \text{Ind}(\mathcal{C})),$$

where on the left we consider colimit-preserving \mathbb{C} -linear functors, is an equivalence. Since $\text{Fun}(P^{\text{op}}, \text{Mod}_{\mathbb{C}})$ is compactly generated, using [Lur09, Proposition 5.3.5.10] we obtain that the restriction

$$\text{Fun}^{\text{ex}}(\text{Fun}(P^{\text{op}}, \text{Mod}_{\mathbb{C}})^\omega, \text{Ind}(\mathcal{C})) \longrightarrow \text{Fun}(P, \text{Ind}(\mathcal{C}))$$

is an equivalence. Since P is a finite poset, by [Aok23, Proposition 2.8] we conclude that

$$\text{Fun}^{\text{ex}}(\text{Perf}(P^{\text{op}}), \mathcal{C}) \longrightarrow \text{Fun}(P, \mathcal{C})$$

is an equivalence. □

For $\mathcal{C} \in \mathbf{dgc}at$ we denote its ∞ -category of (right) modules by

$$\text{Mod}_{\mathcal{C}} = \text{Fun}^{\text{ex}}(\mathcal{C}^{\text{op}}, \text{Mod}_{\mathbb{C}}).$$

We refer to objects of $\text{Mod}_{\mathcal{C}^{\text{op}} \otimes \mathcal{D}}$ as $(\mathcal{C}, \mathcal{D})$ -bimodules. There are relative tensor product functors and internal Hom functors; we refer to [BD19, Section 2] for more details on the Morita theory of dg categories.

For $\mathcal{C} \in \mathbf{dgc}at$ we denote $\mathcal{C}^e = \mathcal{C}^{op} \otimes \mathcal{C}$. We also denote by $\mathcal{C} \in \text{Mod}_{\mathcal{C}^e}$ the diagonal bimodule $x, y \mapsto \text{Hom}_{\mathcal{C}}(x, y)$. The **Hochschild complex** is $\text{HH}(\mathcal{C}) = \mathcal{C} \otimes_{\mathcal{C}^e} \mathcal{C}$. It carries a natural S^1 -action and the **negative cyclic complex** is $\text{HC}^-(\mathcal{C}) = \text{HH}(\mathcal{C})^{S^1}$.

Definition 8.11.

- A dg category \mathcal{C} is of **finite type** if $\mathcal{C} \in \mathbf{dgc}at$ is compact.
- A dg category \mathcal{C} is **smooth** if the diagonal bimodule $\mathcal{C} \in \text{Mod}_{\mathcal{C}^e}$ is compact. In this case, we have the dual bimodule $\mathcal{C}^! = \text{Hom}_{\mathcal{C}^e}(\mathcal{C}, \mathcal{C}^e) \in \text{Mod}_{\mathcal{C}^e}$.

We have the following basic facts:

- A finite type dg category is smooth [TV07, Proposition 2.14].
- If \mathcal{C} is smooth, there is an isomorphism $\text{HH}(\mathcal{C}) \cong \text{Hom}_{\mathcal{C}^e}(\mathcal{C}^!, \mathcal{C})$.

Next, let us recall the construction of the moduli stack of objects. Let \mathcal{C} be a dg category.

Definition 8.12. The **moduli stack $\mathfrak{M}_{\mathcal{C}}$ of objects in \mathcal{C}** is the functor

$$\mathbf{cdga}^{\leq 0} \ni R \mapsto \text{Hom}_{\mathbf{dgc}at}(\mathcal{C}, \text{Perf}_R) \in \mathcal{S}.$$

We refer to functors $\mathcal{C} \rightarrow \text{Perf}_R$ as **pseudo-perfect \mathcal{C} -modules**.

We also have a similar description of mapping stacks.

Lemma 8.13. *Let X be a derived prestack and \mathcal{C} a dg category. Then there is a natural equivalence*

$$\text{Map}(X, \mathfrak{M}_{\mathcal{C}})(R) \cong \text{Hom}_{\mathbf{dgc}at}(\mathcal{C}, \text{Perf}(X \times \text{Spec } R))$$

for $R \in \mathbf{cdga}^{\leq 0}$.

Proof. It is enough to prove that $\text{Hom}(X, \mathfrak{M}_{\mathcal{C}}) \cong \text{Hom}_{\mathbf{dgc}at}(\mathcal{C}, \text{Perf}(X))$ for any derived prestack X . If $X = \text{Spec } A$ is affine, this holds by the definition of the moduli stack of objects. As any derived prestack is obtained as a colimit of derived affine schemes and both $\text{Hom}(X, \mathfrak{M}_{\mathcal{C}})$ and $\text{Hom}_{\mathbf{dgc}at}(\mathcal{C}, \text{Perf}(X))$ send colimits in X to limits, the claim follows. \square

If \mathcal{C} is of finite type, it is shown in [TV07, Theorem 3.6] that $\mathfrak{M}_{\mathcal{C}}$ is a locally geometric derived stack.

Definition 8.14. Let \mathcal{C} be a smooth dg category. A **d -dimensional Calabi–Yau structure on \mathcal{C}** is a negative cyclic class $\tilde{c} \in |\text{HC}^-(\mathcal{C})[-d]|$ whose underlying Hochschild class $c \in |\text{HH}(\mathcal{C})[-d]| \cong |\text{Hom}_{\mathcal{C}^e}(\mathcal{C}^!, \mathcal{C})[-d]|$ defines an equivalence $\mathcal{C}^! \rightarrow \mathcal{C}[-d]$ of \mathcal{C}^e -modules.

Remark 8.15. In this paper we only consider left Calabi–Yau structures in the terminology of [BD19], so the adjective “left” will often be dropped.

Example 8.16. The dg category $\text{Perf}_{\mathbb{C}}^{\oplus n}$ has $\text{HC}^-(\text{Perf}_{\mathbb{C}}^{\oplus n}) \cong \mathbb{C}[[u]]^{\oplus n}$. The class $\tilde{c}_n = (1, \dots, 1)$ defines a 0-Calabi–Yau structure on $\text{Perf}_{\mathbb{C}}^{\oplus n}$.

Given an exact functor $F: \mathcal{C} \rightarrow \text{Perf}_R$ we get a map

$$\text{HC}^-(\mathcal{C}) \xrightarrow{\text{HC}^-(F)} \text{HC}^-(R) \xrightarrow{\tilde{\kappa}_2} \Omega^{\geq 2}(R)[2],$$

where the map $\tilde{\kappa}_2$ is defined using the HKR isomorphism as in [BD21, Proposition 5.2]. By naturality, we get an induced map

$$(8.8) \quad \tilde{\kappa}_2: \text{HC}^-(\mathcal{C}) \longrightarrow \Omega^{\geq 2}(\mathfrak{M}_{\mathcal{C}})[2].$$

If \mathcal{C} is of finite type, it is shown in [BD21, Theorem 5.5] that it sends d -Calabi–Yau structures on \mathcal{C} to $(2-d)$ -shifted symplectic structures on $\mathfrak{M}_{\mathcal{C}}$.

Let $P_n = \{0 < 1 < \dots < n-1\}$ be the poset of integers between 0 and $n-1$.

Definition 8.17. The **moduli stack of n -filtered objects in \mathcal{C}** is

$$\mathfrak{M}_{\mathcal{C}}^{n\text{-filt}} = \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_n)}.$$

By Lemma 8.10 the R -points of $\mathfrak{M}_{\mathbb{C}}^{n\text{-filt}}$ are given by

$$\mathfrak{M}_{\mathbb{C}}^{n\text{-filt}}(R) = \text{Map}(P_n^{\text{op}}, \text{Fun}^{\text{ex}}(\mathbb{C}, \text{Perf}_R)),$$

so $\mathfrak{M}_{\mathbb{C}}^{n\text{-filt}}$ parametrizes sequences $M_{n-1} \rightarrow \cdots \rightarrow M_0$ of pseudo-perfect modules. We will be interested in the following functors:

- $\iota_1: \text{Perf}_{\mathbb{C}} \rightarrow \text{Perf}(P_n)$ given by the inclusion of the constant sheaf $\mathcal{O}_{P_n} \in \text{Perf}(P_n)$ sending $i \in P_n \mapsto \mathbb{C}$ and any $i \rightarrow j$ to the identity. By restriction, it induces a map $\text{ev}: \mathfrak{M}_{\mathbb{C}}^{n\text{-filt}} \rightarrow \mathfrak{M}_{\mathbb{C}}$ sending a filtered pseudo-perfect module $M_{n-1} \rightarrow \cdots \rightarrow M_0$ to M_0 .
- $P_n^{\delta} \rightarrow \text{Perf}(P_n)$, where P_n^{δ} is the same underlying set as P_n considered as a discrete category, given by sending $i \in P_n^{\delta}$ to the skyscraper sheaf i . By the universal property of $\text{Perf}(P_n^{\delta})$ it induces an exact functor $\iota_{\text{gr}}: \text{Perf}(P_n^{\delta}) \rightarrow \text{Perf}(P_n)$. Since $\mathbb{C} \otimes \text{Perf}(P_n^{\delta}) \cong \mathbb{C}^{\oplus n}$, it induces a map $\text{gr}: \mathfrak{M}_{\mathbb{C}}^{n\text{-filt}} \rightarrow \mathfrak{M}_{\mathbb{C}}^n$ sending a filtered pseudo-perfect module to its associated graded.
- The functor $\pi: \text{Perf}(P_n) \rightarrow \text{Perf}(P_n^{\delta})$ left inverse to ι_{gr} given by restriction along $P_n^{\delta} \rightarrow P_n$. By restriction, it induces a map $\sigma: \mathfrak{M}_{\mathbb{C}}^n \rightarrow \mathfrak{M}_{\mathbb{C}}^{n\text{-filt}}$ which is a section of gr . It sends M_0, \dots, M_{n-1} to $M_{n-1} \rightarrow M_{n-1} \oplus M_{n-2} \rightarrow \cdots$.
- The diagonal functor $\text{Perf}_{\mathbb{C}} \rightarrow \text{Perf}(P_n^{\delta}) = \text{Perf}_{\mathbb{C}}^{\oplus n}$ given by the inclusion of the constant sheaf $\mathcal{O}_{P_n^{\delta}} \in \text{Perf}(P_n^{\delta})$. By restriction, it induces a map $\Phi_n: \mathfrak{M}_{\mathbb{C}}^n \rightarrow \mathfrak{M}_{\mathbb{C}}$ given by sending M_0, \dots, M_{n-1} to $M_0 \oplus \cdots \oplus M_{n-1}$. We have $\Phi_n = \text{ev} \circ \sigma$.

We obtain a correspondence

$$(8.9) \quad \begin{array}{ccc} & \mathfrak{M}_{\mathbb{C}}^{n\text{-filt}} & \\ \text{gr} \nearrow & & \searrow \text{ev} \\ \mathfrak{M}_{\mathbb{C}}^n & & \mathfrak{M}_{\mathbb{C}} \\ & \sigma \searrow & \\ & & \end{array}$$

Let us now endow this correspondence with a Lagrangian structure when \mathbb{C} carries a left d -Calabi–Yau structure. The following is essentially shown in [BD19, Theorem 5.14].

Proposition 8.18. *The cospan of dg categories $\text{Perf}(P_n^{\delta}) \rightarrow \text{Perf}(P_n) \leftarrow \text{Perf}_{\mathbb{C}}$ carries a unique 0-Calabi–Yau structure which restricts to the 0-Calabi–Yau structure \tilde{c}_n on $\text{Perf}(P_n^{\delta}) \cong \text{Perf}_{\mathbb{C}}^{\oplus n}$ and the 0-Calabi–Yau structure \tilde{c}_1 on $\text{Perf}_{\mathbb{C}}$.*

Proof. Since the functor HC^- is additive, the functors

$$\pi: \text{Perf}(P_n) \longrightarrow \text{Perf}(P_n^{\delta}), \quad \iota_{\text{gr}}: \text{Perf}(P_n^{\delta}) \longrightarrow \text{Perf}(P_n)$$

induce inverse maps on HC^- [BD19, Proposition 5.15]. Therefore, a 0-Calabi–Yau structure on the cospan is given by a homotopy commutative diagram

$$\begin{array}{ccc} \mathbb{C} & \xrightarrow{c_1} & \mathbb{C}[[u]] \\ \downarrow c_n & & \downarrow \Delta \\ \mathbb{C}[[u]]^{\oplus n} & \xrightarrow{\text{id}} & \mathbb{C}[[u]]^{\oplus n} \end{array}$$

where the vertical map on the right is the diagonal. Since all complexes are concentrated in non-negative cohomological degrees, there is a unique homotopy which makes this diagram commutative. \square

Corollary 8.19 ([BD21, Corollary 6.5]). *Let \mathbb{C} be a finite type dg category equipped with a d -Calabi–Yau structure. Then there is a canonical $(2-d)$ -shifted Lagrangian structure on the correspondence*

$$(8.10) \quad \begin{array}{ccc} & \mathfrak{M}_{\mathbb{C}}^{n\text{-filt}} & \\ \text{gr} \nearrow & & \searrow \text{ev} \\ \mathfrak{M}_{\mathbb{C}}^n & & \mathfrak{M}_{\mathbb{C}} \end{array}$$

In particular, restricting this Lagrangian structure along the section $\sigma: \mathfrak{M}_{\mathbb{C}}^n \rightarrow \mathfrak{M}_{\mathbb{C}}^{n\text{-filt}}$ we get that $\Phi_n: \mathfrak{M}_{\mathbb{C}}^n \rightarrow \mathfrak{M}_{\mathbb{C}}$ is compatible with $(2-d)$ -shifted symplectic structures.

Let us now describe how to compose these Lagrangian correspondences.

Proposition 8.20. *Let \mathcal{C} be a finite type dg category. The composition of the correspondences*

$$\begin{array}{ccccc}
 & \mathfrak{M}_{\mathcal{C}}^{n\text{-filt}} \times \mathfrak{M}_{\mathcal{C}}^{m-1} & & \mathfrak{M}_{\mathcal{C}}^{m\text{-filt}} & \\
 & \swarrow \text{gr} \times \text{id} & & \swarrow \text{gr} & \searrow \text{ev} \\
 \mathfrak{M}_{\mathcal{C}}^{n+m-1} & & \mathfrak{M}_{\mathcal{C}}^n & & \mathfrak{M}_{\mathcal{C}} \\
 & \searrow \text{ev} \times \text{id} & & & \\
 & & & &
 \end{array}$$

and

$$\begin{array}{ccccc}
 & \mathfrak{M}_{\mathcal{C}}^{n-1} \times \mathfrak{M}_{\mathcal{C}}^{m\text{-filt}} & & \mathfrak{M}_{\mathcal{C}}^{n\text{-filt}} & \\
 & \swarrow \text{id} \times \text{gr} & & \swarrow \text{gr} & \searrow \text{ev} \\
 \mathfrak{M}_{\mathcal{C}}^{n+m-1} & & \mathfrak{M}_{\mathcal{C}}^n & & \mathfrak{M}_{\mathcal{C}} \\
 & \searrow \text{id} \times \text{ev} & & & \\
 & & & &
 \end{array}$$

are both equivalent to the same correspondence

$$\begin{array}{ccc}
 & \mathfrak{M}_{\mathcal{C}}^{(n+m-1)\text{-filt}} & \\
 & \swarrow \text{gr} & \searrow \text{ev} \\
 \mathfrak{M}_{\mathcal{C}}^{n+m-1} & & \mathfrak{M}_{\mathcal{C}}.
 \end{array}$$

Moreover, if \mathcal{C} is equipped with a d -Calabi–Yau structure, this equivalence is compatible with Lagrangian structures.

Proof. It is shown in [BCS24, Section 6.1.2] that the functor of moduli of objects sends compositions of Calabi–Yau cospans to compositions of Lagrangian correspondences. Thus, it is enough to prove the claim for Calabi–Yau cospans. The two assertions are proven analogously, so we will only deal with the first claim. Consider a diagram

$$\begin{array}{ccccc}
 & & \text{Perf}(P_{n+m-1}) & & \\
 & & \swarrow & & \searrow \\
 \text{Perf}_{\mathcal{C}}^{\oplus(n-1)} \oplus \text{Perf}(P_m) & & & & \text{Perf}(P_n) \\
 & \swarrow & & \swarrow & \swarrow \\
 \text{Perf}_{\mathcal{C}}^{\oplus(n+m-1)} & & \text{Perf}_{\mathcal{C}}^{\oplus n} & & \text{Perf}_{\mathcal{C}}
 \end{array}$$

of dg categories. To show that the top square is coCartesian in $\mathbf{dgc}at$, using Lemma 8.10 we have that for any dg category \mathcal{D} we have $\text{Fun}^{\text{ex}}(\text{Perf}(P_n), \mathcal{D}) \cong \text{Fun}((\Delta^{n-1})^{\text{op}}, \mathcal{D})$, so we have to show that

$$\begin{array}{ccc}
 \Delta^0 & \longrightarrow & \Delta^{m-1} \\
 \downarrow & & \downarrow \\
 \Delta^{n-1} & \longrightarrow & \Delta^{n+m-2}
 \end{array}$$

is a homotopy coCartesian square of simplicial sets with respect to the Joyal model structure. By induction, it is enough to prove the claim for $n = m = 1$ in which case we have that $\Lambda_1^2 = \Delta^1 \coprod_{\Delta^0} \Delta^1 \rightarrow \Delta^2$ is a categorical equivalence since it is inner anodyne [Lur09, Lemma 2.2.5.2]. It remains to show that the above diagram of dg categories represents a composition of Calabi–Yau cospans; but it is obvious, as there is a unique structure of a 0-Calabi–Yau cospan on $\text{Perf}_{\mathcal{C}}^{\oplus(n+m-1)} \rightarrow \text{Perf}(P_{n+m-1}) \leftarrow \text{Perf}_{\mathcal{C}}$ by Proposition 8.18. \square

8.4. From moduli of objects to mapping stacks. In this section we explain how to relate moduli of objects to mapping stacks.

Definition 8.21. Let X be a derived stack and \mathcal{D} a dg category. The ∞ -category of *perfect complexes on X with coefficients in \mathcal{D}* is

$$\mathrm{Perf}(X; \mathcal{D}) = \lim_{\mathrm{Spec} A \rightarrow X} (\mathrm{Perf}_A \otimes \mathcal{D}).$$

Example 8.22. Let R be a connective commutative dg algebra. Then

$$\mathrm{Perf}(X; \mathrm{Perf}_R) \cong \mathrm{Perf}(X \times \mathrm{Spec} R).$$

Example 8.23. If \mathcal{D} is smooth and proper, it is dualizable in **dgcat** and hence $(-) \otimes \mathcal{D}$ preserves limits. Therefore, the natural functor $\mathcal{D} \otimes \mathrm{Perf}(X) \rightarrow \mathrm{Perf}(X; \mathcal{D})$ is an equivalence.

An object $E \in \mathrm{Perf}(X; \mathcal{D})$ induces a functor

$$T_{E,R}: \mathrm{Fun}^{\mathrm{ex}}(\mathcal{D}, \mathrm{Perf}_R) \longrightarrow \mathrm{Perf}(X; \mathrm{Perf}_R) \cong \mathrm{Perf}(X \times \mathrm{Spec} R)$$

which is natural in R .

Proposition 8.24. *Let X be a derived stack and \mathcal{C}, \mathcal{D} a pair of dg categories. An object $E \in \mathrm{Perf}(X; \mathcal{D})$ gives rise to a morphism*

$$F_E: \mathfrak{M}_{\mathcal{C} \otimes \mathcal{D}} \longrightarrow \mathrm{Map}(X, \mathfrak{M}_{\mathcal{C}})$$

of derived stacks.

Proof. Using the description of the mapping stacks to the moduli of objects given in Lemma 8.13 we have to construct, for every connective cdga R , a morphism

$$\mathrm{Hom}_{\mathbf{dgcat}}(\mathcal{C} \otimes \mathcal{D}, \mathrm{Perf}_R) \longrightarrow \mathrm{Hom}_{\mathbf{dgcat}}(\mathcal{C}, \mathrm{Perf}(X \times \mathrm{Spec} R))$$

natural in R . By adjunction the left-hand side is isomorphic to

$$\mathrm{Hom}_{\mathbf{dgcat}}(\mathcal{C} \otimes \mathcal{D}, \mathrm{Perf}_R) \cong \mathrm{Hom}_{\mathbf{dgcat}}(\mathcal{C}, \mathrm{Fun}^{\mathrm{ex}}(\mathcal{D}, \mathrm{Perf}_R))$$

and the map F_E is simply given by post-composition with $T_{E,R}$. □

If X is a derived stack and \mathcal{D} a dg category, consider the morphisms

$$\begin{aligned} \mathrm{HC}^-(\mathrm{Perf}(X; \mathcal{D})) &\longrightarrow \lim_{\mathrm{Spec} A \rightarrow X} \mathrm{HC}^-(\mathrm{Perf}_A \otimes \mathcal{D}) \\ &\cong \lim_{\mathrm{Spec} A \rightarrow X} (\mathrm{HH}(A) \otimes \mathrm{HH}(\mathcal{D}))^{S^1} \\ &\longrightarrow \lim_{\mathrm{Spec} A \rightarrow X} (A \otimes \mathrm{HC}^-(\mathcal{D})) \\ &\longleftarrow \mathbb{R}\Gamma(X, \mathcal{O}) \otimes \mathrm{HC}^-(\mathcal{D}) \end{aligned}$$

where we have used the natural projection $\mathrm{HH}(A) \rightarrow A$ which is S^1 -equivariant. If X is \mathcal{O} -compact, the last backward map is an isomorphism, so in total we obtain a morphism

$$\kappa_{\mathcal{D}, X}: \mathrm{HC}^-(\mathrm{Perf}(X; \mathcal{D})) \longrightarrow \mathbb{R}\Gamma(X, \mathcal{O}) \otimes \mathrm{HC}^-(\mathcal{D}).$$

Theorem 8.25. *Consider the following data:*

- (1) X is an \mathcal{O} -compact derived stack equipped with a d_1 -preorientation $[X]: \mathbb{R}\Gamma(X, \mathcal{O}) \rightarrow \mathbb{C}[-d_1]$.
- (2) \mathcal{C} is a dg category equipped with a class $[\mathcal{C}] \in |\mathrm{HC}^-(\mathcal{C})[-d_2]|$.
- (3) $E \in \mathrm{Perf}(X; \mathcal{D})$.
- (4) \mathcal{D} is a dg category.

Consider the class $[\mathcal{D}] \in |\mathrm{HC}^-(\mathcal{D})[-d_1]|$ given by the composite

$$\begin{aligned} \mathbb{C} &\xrightarrow{[E]} \mathrm{HC}^-(\mathrm{Perf}(X; \mathcal{D})) \\ &\xrightarrow{\kappa_{\mathcal{D}, X}} \mathbb{R}\Gamma(X, \mathcal{O}) \otimes \mathrm{HC}^-(\mathcal{D}) \\ &\xrightarrow{[X] \otimes \mathrm{id}} \mathrm{HC}^-(\mathcal{D})[-d_1] \end{aligned}$$

Then the morphism

$$F_E: \mathfrak{M}_{\mathcal{C} \otimes \mathcal{D}} \longrightarrow \text{Map}(X, \mathfrak{M}_{\mathcal{C}})$$

is compatible with $(2 - d_1 - d_2)$ -shifted presymplectic structures, where the $(2 - d_1 - d_2)$ -shifted presymplectic structure on the left is defined by the image of $[\mathcal{C}] \otimes [\mathcal{D}]$ under $\text{HC}^-(\mathcal{C}) \otimes \text{HC}^-(\mathcal{D}) \rightarrow \text{HC}^-(\mathcal{C} \otimes \mathcal{D})$ and on the right induced by the AKSZ procedure from the $(2 - d_2)$ -shifted presymplectic structure on $\mathfrak{M}_{\mathcal{C}}$ defined by $[\mathcal{C}]$.

Proof. Consider an R -point of $\mathfrak{M}_{\mathcal{C} \otimes \mathcal{D}}$ specified by an exact functor $F: \mathcal{C} \otimes \mathcal{D} \rightarrow \text{Perf}_R$ and the corresponding R -point of $\text{Map}(X, \mathfrak{M}_{\mathcal{C}})$ specified by an exact functor $F': \mathcal{C} \rightarrow \text{Perf}(X \times \text{Spec } R)$. The pullback of the $(2 - d_1 - d_2)$ -shifted presymplectic structure on $\mathfrak{M}_{\mathcal{C} \otimes \mathcal{D}}$ to $\text{Spec } R$ is given by the composite

$$\begin{aligned} \mathbb{C} &\xrightarrow{[\mathcal{C}] \otimes [\mathcal{D}]} \text{HC}^-(\mathcal{C}) \otimes \text{HC}^-(\mathcal{D})[-d_1 - d_2] \\ &\rightarrow \text{HC}^-(\mathcal{C} \otimes \mathcal{D})[-d_1 - d_2] \\ &\xrightarrow{\text{HC}^-(F)} \text{HC}^-(R)[-d_1 - d_2] \\ &\xrightarrow{\tilde{\kappa}_2} \Omega^{\geq 2}(R)[2 - d_1 - d_2]. \end{aligned}$$

Similarly, the pullback of the $(2 - d_1 - d_2)$ -shifted presymplectic structure on $\text{Map}(X, \mathfrak{M}_{\mathcal{C}})$ to $\text{Spec } R$ is given by the composite

$$\begin{aligned} \mathbb{C} &\xrightarrow{[\mathcal{C}]} \text{HC}^-(\mathcal{C})[-d_2] \\ &\xrightarrow{\text{HC}^-(F')} \text{HC}^-(\text{Perf}(X \times \text{Spec } R))[-d_2] \\ &\xrightarrow{\kappa_{\text{Spec } R, X}} \mathbb{R}\Gamma(X, \mathcal{O}) \otimes \text{HC}^-(R)[-d_2] \\ &\xrightarrow{[X]} \text{HC}^-(R)[-d_1 - d_2] \\ &\xrightarrow{\tilde{\kappa}_2} \Omega^{\geq 2}(R)[2 - d_1 - d_2] \end{aligned}$$

The claim then follows from the fact that F' is given by the composite

$$\begin{aligned} \mathcal{C} &\xrightarrow{\text{id} \otimes E} \mathcal{C} \otimes \text{Perf}(X; \mathcal{D}) \\ &\rightarrow \text{Perf}(X; \mathcal{C} \otimes \mathcal{D}) \\ &\xrightarrow{F} \text{Perf}(X; \text{Perf}_R) \\ &\cong \text{Perf}(X \times \text{Spec } R). \end{aligned}$$

□

The constructions described in Proposition 8.24 and Theorem 8.25 are functorial in the following way. Suppose I is a small ∞ -category, X_{\bullet} is a diagram $I \rightarrow \mathbf{dSt}$ of derived stacks and \mathcal{D}_{\bullet} is a diagram $I \rightarrow \mathbf{dgc}$ of dg categories. Then:

- Using the description of natural transformations as an end [GHN17, Proposition 5.1], by Proposition 8.24 an object $E \in \int_{i \in I} \text{Perf}(X_i; \mathcal{D}_i)$ produces a natural transformation

$$F_E: \mathfrak{M}_{\mathcal{C} \otimes \mathcal{D}_{\bullet}} \longrightarrow \text{Map}(X_{\bullet}, \mathfrak{M}_{\mathcal{C}})$$

of functors $I^{\text{op}} \rightarrow \mathbf{dSt}$.

- If, in addition, we have a class $[\mathcal{C}] \in |\text{HC}^-(\mathcal{C})[-d_2]|$, for every $i \in I$ the derived stack X_i is \mathcal{O} -compact and we are given a d_1 -preorientation

$$[X_{\bullet}]: \text{colim}_{i \in I} \mathbb{R}\Gamma(X_i, \mathcal{O}) \rightarrow \mathbb{C}[-d_1],$$

then by Theorem 8.25 the natural transformation F_E is compatible with $(2 - d_1 - d_2)$ -shifted presymplectic structures.

8.5. Attractor correspondence for the moduli of objects. Let us now work out a description of the attractor correspondence for the moduli stack of objects. Let \mathcal{C} be a dg category of finite type. For a finite connected subset $P \subset \mathbb{Z}_{\geq}$ (which is isomorphic to P_n^{op} for some n) consider the correspondence

$$\begin{array}{ccc} & \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^{\text{op}})} & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^\delta)} & & \mathfrak{M}_{\mathcal{C}} \end{array}$$

An inclusion $P_1 \subset P_2$ produces restriction functors $\text{Perf}(P_2^{\text{op}}) \rightarrow \text{Perf}(P_1^{\text{op}})$ and hence a morphism of correspondences

$$\left(\begin{array}{ccc} & \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_1^{\text{op}})} & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_1^\delta)} & & \mathfrak{M}_{\mathcal{C}} \end{array} \right) \longrightarrow \left(\begin{array}{ccc} & \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_2^{\text{op}})} & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_2^\delta)} & & \mathfrak{M}_{\mathcal{C}} \end{array} \right)$$

Proposition 8.26. *Let \mathcal{C} be a dg category of finite type. Then*

$$\begin{array}{ccc} & \text{Filt}(\mathfrak{M}_{\mathcal{C}}) & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \text{Grad}(\mathfrak{M}_{\mathcal{C}}) & & \mathfrak{M}_{\mathcal{C}} \end{array} \cong_{P \subset \mathbb{Z}_{\geq}, \text{ finite, connected}} \text{colim} \left(\begin{array}{ccc} & \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^{\text{op}})} & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^\delta)} & & \mathfrak{M}_{\mathcal{C}} \end{array} \right)$$

Moreover, the transition maps $\mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_1^{\text{op}})} \rightarrow \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_2^{\text{op}})}$ and $\mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_1^\delta)} \rightarrow \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_2^\delta)}$ are open immersions.

Proof. Let us prove that

$$\text{Filt}(\mathfrak{M}_{\mathcal{C}}) \cong_{P \subset \mathbb{Z}_{\geq}, \text{ finite, connected}} \text{colim} \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^{\text{op}})}.$$

By Lemma 8.13 we have

$$\text{Filt}(\mathfrak{M}_{\mathcal{C}})(R) \simeq \text{Hom}_{\mathbf{dgc}at}(\mathcal{C}, \text{Perf}([\mathbb{A}^1/\mathbb{G}_m] \times \text{Spec } R))$$

for any $R \in \mathbf{cdga}^{\leq 0}$. By Proposition 5.8 we further identify

$$\text{Hom}_{\mathbf{dgc}at}(\mathcal{C}, \text{Perf}([\mathbb{A}^1/\mathbb{G}_m] \times \text{Spec } R)) \simeq \text{Hom}_{\mathbf{dgc}at}(\mathcal{C}, \text{colim}_{P \subset \mathbb{Z}_{\geq}, \text{ finite, connected}} \text{Fun}(P, \text{Perf}_R)).$$

Since \mathcal{C} is of finite type, the natural morphism

$$\text{colim}_{P \subset \mathbb{Z}_{\geq}, \text{ finite, connected}} \text{Hom}_{\mathbf{dgc}at}(\mathcal{C}, \text{Fun}(P, \text{Perf}_R)) \longrightarrow \text{Hom}_{\mathbf{dgc}at}(\mathcal{C}, \text{colim}_{P \subset \mathbb{Z}_{\geq}, \text{ finite, connected}} \text{Fun}(P, \text{Perf}_R))$$

is an equivalence. We have $\text{Fun}(P, \text{Perf}_R) \cong \text{Fun}^{\text{ex}}(\text{Perf}(P^{\text{op}}), \text{Perf}_R)$. Therefore,

$$\text{Hom}_{\mathbf{dgc}at}(\mathcal{C}, \text{Fun}(P, \text{Perf}_R)) \simeq \text{Hom}_{\mathbf{dgc}at}(\mathcal{C} \otimes \text{Perf}(P^{\text{op}}), \text{Perf}_R).$$

Finally, the right-hand side is, by definition, $\mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^{\text{op}})}(R) = \text{Fun}(P, \text{Fun}^{\text{ex}}(\mathcal{C}, \text{Perf}_R))$. Thus,

$$\text{Filt}(\mathfrak{M}_{\mathcal{C}})(R) \cong_{P \subset \mathbb{Z}_{\geq}, \text{ finite, connected}} \text{colim} \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^{\text{op}})}(R)$$

which shows that

$$\text{Filt}(\mathfrak{M}_{\mathcal{C}}) \cong_{P \subset \mathbb{Z}_{\geq}, \text{ finite, connected}} \text{colim} \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^{\text{op}})},$$

where we consider the colimit in the ∞ -category of derived prestacks. As $\text{Filt}(\mathfrak{M}_{\mathcal{C}})$ is a stack, this shows that the right-hand side coincides with the colimit computed in the ∞ -category of derived stacks.

For a subset $P_1 \subset P_2$, we will now show that $\mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_1^{\text{op}})} \rightarrow \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P_2^{\text{op}})}$ is an open immersion. On R -points this map is

$$\text{Map}(P_1, \text{Fun}^{\text{ex}}(\mathcal{C}, \text{Perf}_R)) \longrightarrow \text{Map}(P_2, \text{Fun}^{\text{ex}}(\mathcal{C}, \text{Perf}_R))$$

given by the left Kan extension. The right-hand side is given by the space of P_2 -labelled diagrams $\{V_m\}_{m \in P_2}$ of exact functors $\mathcal{C} \rightarrow \text{Perf}_R$ and the left-hand side is given by the union of connected components of those

P_2 -labelled diagrams such that $V_m = 0$ if $m > i$ for any $i \in P_1$ and $V_{m+1} \rightarrow V_m$ is an isomorphism if $m < i$ for any $i \in P_1$.

As \mathcal{C} is of finite type, it admits a generator $E \in \mathcal{C}$, so that the functor $\text{Fun}^{\text{ex}}(\mathcal{C}, \text{Perf}_R) \rightarrow \text{Perf}_R$ given by evaluation at E is conservative. Therefore, the claim is reduced to $\mathcal{C} = \text{Perf}_{\mathcal{C}}$. Therefore, we have to show the following two claims:

- (1) For any perfect complex V of R -modules the locus where it is zero is open in $\text{Spec } R$.
- (2) For a morphism of perfect complexes $V_1 \rightarrow V_2$ of R -modules, the locus where it is an isomorphism is open in $\text{Spec } R$.

By taking the cone, the second claim is reduced to the first claim. Moreover, by induction on the Tor-amplitude, we are reduced to showing the first claim for V a vector bundle on $\text{Spec } R$. Since open immersions are local on the target and a vector bundle is locally trivial, we are reduced to showing the claim for V the trivial vector bundle, where the claim is obvious.

The claims for $\text{Grad}(\mathfrak{M}_{\mathcal{C}})$ as well as for the morphisms $\text{Filt}(\mathfrak{M}_{\mathcal{C}}) \rightarrow \text{Grad}(\mathfrak{M}_{\mathcal{C}})$ and $\text{Filt}(\mathfrak{M}_{\mathcal{C}}) \rightarrow \mathfrak{M}_{\mathcal{C}}$ are proven similarly using the functoriality described in Proposition 5.8. \square

Example 8.27. Setting $\mathcal{C} = \text{Perf}_{\mathcal{C}}$ in Proposition 8.26 we get a description of the moduli stack of graded perfect complexes

$$\text{Grad}(\text{Perf}) \cong \underset{P \subset \mathbb{Z}, \text{ finite, connected}}{\text{colim}} \text{Perf}^P.$$

If we set $\text{Vect} \subset \text{Perf}$ the full substack of vector bundles, then we similarly get a description of the moduli stack of graded vector spaces

$$\text{Grad}(\text{Vect}) \cong \underset{P \subset \mathbb{Z}, \text{ finite, connected}}{\text{colim}} \text{Vect}^P.$$

Let us now show that the isomorphism exhibited in Proposition 8.26 is compatible with Lagrangian structures. Let \mathcal{C} be a finite type dg category equipped with a d -Calabi–Yau structure. For a finite connected subset $P \subset \mathbb{Z}_{\geq}$ Corollary 8.19 constructs a $(2-d)$ -shifted symplectic structure on $\mathfrak{M}_{\mathcal{C}}$ as well as a Lagrangian structure on the correspondence

$$\begin{array}{ccc} & \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^{\text{op}})} & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^{\delta})} & & \mathfrak{M}_{\mathcal{C}} \end{array}$$

Similarly, Corollary 5.18 constructs a Lagrangian structure on the correspondence

$$\begin{array}{ccc} & \text{Filt}(\mathfrak{M}_{\mathcal{C}}) & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \text{Grad}(\mathfrak{M}_{\mathcal{C}}) & & \mathfrak{M}_{\mathcal{C}} \end{array}$$

Proposition 8.28. *Let \mathcal{C} be a finite type dg category equipped with a d -Calabi–Yau structure. Let $P \subset \mathbb{Z}_{\geq}$ be a finite connected subset. Then the morphism*

$$\left(\begin{array}{ccc} & \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^{\text{op}})} & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \mathfrak{M}_{\mathcal{C} \otimes \text{Perf}(P^{\delta})} & & \mathfrak{M}_{\mathcal{C}} \end{array} \right) \longrightarrow \begin{array}{ccc} & \text{Filt}(\mathfrak{M}_{\mathcal{C}}) & \\ \text{gr} \swarrow & & \searrow \text{ev} \\ \text{Grad}(\mathfrak{M}_{\mathcal{C}}) & & \mathfrak{M}_{\mathcal{C}} \end{array}$$

is compatible with Lagrangian structures.

Proof. Let spn be the category $0 \rightarrow 1 \leftarrow 2$ with three objects. Let $X_{\bullet} : \text{spn} \rightarrow \mathbf{dSt}$ be the diagram of \mathcal{O} -compact derived stacks $B\mathbb{G}_m \xrightarrow{i_0} [\mathbb{A}^1/\mathbb{G}_m] \xleftarrow{i_1} \text{pt}$ equipped with a 0-orientation defined in Proposition 5.16. Let $\mathcal{D}_{\bullet} : \text{spn} \rightarrow \mathbf{dgc}at$ be the diagram of dg categories $\text{Perf}(P^{\delta}) \rightarrow \text{Perf}(P^{\text{op}}) \leftarrow \text{Perf}_{\mathcal{C}}$.

Let us first prove that for every $i \in \text{spn}$ the natural functor

$$\text{Fun}^{\text{ex}}(\mathcal{D}_i, \text{Perf}_{\mathcal{C}}) \otimes \text{Perf}_R \longrightarrow \text{Fun}^{\text{ex}}(\mathcal{D}_i, \text{Perf}_R)$$

is an equivalence. This is obvious for $i = 0, 2$, where \mathcal{D}_i is a finite direct sum of $\text{Perf}_{\mathbb{C}}$, so let us just show the claim for $i = 1$. Using Lemma 8.10 we are reduced to showing that

$$\text{Fun}(P, \text{Perf}_{\mathbb{C}}) \otimes \text{Perf}_R \longrightarrow \text{Fun}(P, \text{Perf}_R)$$

is an equivalence. By [Aok23, Proposition 2.8] we have $\text{Fun}(P, \text{Mod}_R)^\omega = \text{Fun}(P, \text{Perf}_R)$ (as well as the same claim with R replaced by \mathbb{C}). If we denote by $\widehat{\otimes}$ the symmetric monoidal structure in Pr^L , we are reduced to showing that

$$\text{Fun}(P, \mathcal{S}) \widehat{\otimes} \text{Mod}_R \longrightarrow \text{Fun}(P, \text{Mod}_R)$$

is an equivalence for any commutative dg algebra R , which is [Aok23, Corollary 2.2].

The natural functor

$$\text{Perf}(X_\bullet) \otimes \text{Perf}(R) \longrightarrow \text{Perf}(X_\bullet \times \text{Spec } R)$$

is also an equivalence, since X_\bullet is perfect [BFN10, Proposition 4.6].

Therefore, the equivalence established in Proposition 5.8, natural in R , is uniquely determined by its value on $R = \mathbb{C}$, so that the functor $\text{Fun}^{\text{ex}}(\mathcal{D}_\bullet, \text{Perf}_R) \rightarrow \text{Perf}(X_\bullet \times \text{Spec } R)$ defined in Proposition 5.8 is given by

$$\text{Fun}^{\text{ex}}(\mathcal{D}_\bullet, \text{Perf}_R) \xrightarrow{T_{E,R}} \text{Perf}(X_\bullet \times \text{Spec } R)$$

for a universal object

$$E \in \int_{i \in \text{spn}} \text{Perf}(X_i; \mathcal{D}_i),$$

where we note that by Example 8.23 we have $\text{Perf}(X_i; \mathcal{D}_j) \cong \text{Perf}(X_i) \otimes \mathcal{D}_j$ since \mathcal{D}_j is smooth and proper for every $j \in \text{spn}$.

Using Theorem 8.25 we deduce that the morphism of correspondences F_E is compatible with Lagrangian structures, where \mathcal{D}_\bullet is equipped with a certain structure of a 0-Calabi–Yau cospan $[\mathcal{D}_\bullet]$ described in the statement. It remains to show that this 0-Calabi–Yau structure on \mathcal{D}_\bullet is equivalent to the one defined in Proposition 8.18. By uniqueness, it is enough to identify the 0-Calabi–Yau structures on $\text{Perf}(P^\delta)$ and $\text{Perf}_{\mathbb{C}}$. The latter one is given by the composite of the identity maps, so it coincides with \tilde{c}_1 . Let us now compute $[\text{Perf}(P^\delta)]$. By definition, it is given by the composite

$$\begin{aligned} \mathbb{C} &\xrightarrow{[E_0]} \text{HC}^-(\text{Perf}(B\mathbb{G}_m; \text{Perf}(P^\delta))) \\ &\xrightarrow{\kappa_{\text{Perf}(P^\delta), B\mathbb{G}_m}} \mathbb{R}\Gamma(B\mathbb{G}_m, \mathcal{O}) \otimes \text{HC}^-(\text{Perf}(P^\delta)) \\ &\xrightarrow{[B\mathbb{G}_m] \otimes \text{id}} \text{HC}^-(\text{Perf}(P^\delta)). \end{aligned}$$

Recall that $[B\mathbb{G}_m]$ is defined by the pullback

$$\mathbb{R}\Gamma(B\mathbb{G}_m, \mathcal{O}) \xrightarrow{i^*} \mathbb{R}\Gamma(\text{pt}, \mathcal{O}) \cong \mathbb{C}$$

along the inclusion of the basepoint $i: \text{pt} \rightarrow B\mathbb{G}_m$. Using naturality of $\kappa_{\text{Perf}(P^\delta), -}$ for the morphism $i: \text{pt} \rightarrow B\mathbb{G}_m$ we get that the class $[\text{Perf}(P^\delta)]$ is given by the composite

$$\mathbb{C} \xrightarrow{[E_0]} \text{HC}^-(\text{Perf}(B\mathbb{G}_m; \text{Perf}(P^\delta))) \xrightarrow{i^*} \text{HC}^-(\text{Perf}(P^\delta)).$$

The object $E_0 \in \text{Perf}(B\mathbb{G}_m; \text{Perf}(P^\delta)) \cong \text{Fun}(P^\delta, \text{Perf}(B\mathbb{G}_m))$ is given by sending $p \in P^\delta$ to the 1-dimensional representation of \mathbb{G}_m with weight p . Thus, $i^*E_0 = \mathcal{O}_{P^\delta}$ is the constant sheaf $p \mapsto \mathbb{C}$ whose class $[i^*E_0] \in \text{HC}^-(\text{Perf}(P^\delta))$ coincides with \tilde{c}_n . \square

Proposition 8.29. *Let \mathcal{C} be a finite type dg category equipped with a 3-Calabi–Yau structure and $\mathfrak{M} \subset \mathfrak{M}_{\mathcal{C}}$ an open substack satisfying the following conditions:*

- (1) \mathfrak{M}^{cl} is a quasi-separated Artin stack with affine stabilizers.
- (2) \mathfrak{M} is closed under direct sums and it contains the zero module.

Then \mathfrak{M} is a moduli-like stack in the sense of Definition 8.1.

Proof. By [TV07, Theorem 3.6] the stack \mathfrak{M}_e is locally geometric and locally of finite presentation, so \mathfrak{M} is locally geometric and locally of finite presentation. The stack \mathfrak{M}_e carries a natural $B\mathbb{G}_m$ -action, which is given on R -points by tensoring a pseudo-perfect module $\mathcal{C} \rightarrow \text{Perf}_R$ with a line bundle on $\text{Spec } R$ considered as a constant pseudo-perfect module. Since \mathfrak{M} is open, this action restricts to \mathfrak{M} . Since \mathfrak{M} is closed under direct sums, the \mathbb{E}_∞ -structure on \mathfrak{M}_e given by direct sum restricts to \mathfrak{M} . Finally, by Corollary 8.19 we obtain a (-1) -shifted symplectic structure on \mathfrak{M} which is additive.

Let us now check the conditions:

- (1) Consider the map $\mathfrak{M}_e^2 \rightarrow \text{Grad}(\mathfrak{M}_e)$ given by the inclusion of graded objects in weights 0 and 1. By Proposition 8.26 it is an open immersion. Since \mathfrak{M}_e is locally geometric and locally of finite presentation, by Proposition 8.26 $\text{Grad}(\mathfrak{M}_e)$ is locally geometric and locally of finite presentation. Since $\mathfrak{M} \rightarrow \mathfrak{M}_e$ is a monomorphism, so is $\text{Grad}(\mathfrak{M}) \rightarrow \text{Grad}(\mathfrak{M}_e)$ (see [Hal14, Proposition 1.3.1]). Since $\mathfrak{M} \rightarrow \mathfrak{M}_e$ is formally étale, $\text{Grad}(\mathfrak{M}) \rightarrow \text{Grad}(\mathfrak{M}_e)$ is formally étale by Proposition 5.10. Therefore, by [TV08, Corollary 2.2.5.6] the morphism $\text{Grad}(\mathfrak{M}) \rightarrow \text{Grad}(\mathfrak{M}_e)$ is étale and hence flat. Therefore, $\text{Grad}(\mathfrak{M}) \rightarrow \text{Grad}(\mathfrak{M}_e)$ is an open immersion. In the diagram

$$\begin{array}{ccc} \mathfrak{M}^2 & \xrightarrow{\tilde{\Phi}_2} & \text{Grad}(\mathfrak{M}) \\ \downarrow & & \downarrow \\ \mathfrak{M}_e^2 & \longrightarrow & \text{Grad}(\mathfrak{M}_e) \end{array}$$

we have proven that all morphisms except for $\tilde{\Phi}_2$ are open immersions. Therefore, $\tilde{\Phi}_2$ is an open immersion. The statement about $\tilde{\Phi}_3$ is proven analogously.

- (2) This claim follows from Proposition 8.26.
- (3) As $\mathfrak{M} \rightarrow \mathfrak{M}_e$ is formally étale, it is enough to show that $\mathfrak{M}_e^3 \rightarrow \text{Grad}^2(\mathfrak{M}_e)$ given by the inclusion of graded objects in weights $(0, 0)$, $(1, 0)$ and $(1, 1)$ lands in the matching locus. Using the description of the tangent complex of \mathfrak{M}_e from [TV07, Corollary 3.17], we see that $\mathbb{T}_{\mathfrak{M}_e} |_{\mathfrak{M}_e^3}$ is concentrated in weights $(0, 0)$, $(\pm 1, 0)$, $(\pm 1, \pm 1)$ and $(0, \pm 1)$. In particular, the morphism $\mathfrak{M}_e^3 \rightarrow \text{Grad}^2(\mathfrak{M}_e)$ lands in the matching locus.
- (4) This claim follows from Lemma 5.11.

□

Remark 8.30. Proposition 8.28 shows that for an open substack $\mathfrak{M} \subset \mathfrak{M}_e$ as in Proposition 8.29, the (-1) -shifted Lagrangian correspondence $\mathfrak{M}^2 \leftarrow \mathfrak{M}^{2\text{-filt}} \rightarrow \mathfrak{M}$ is a restriction of the (-1) -shifted Lagrangian correspondence $\mathfrak{M}_e^2 \leftarrow \mathfrak{M}_e^{2\text{-filt}} \rightarrow \mathfrak{M}_e$ from [BD21, Corollary 6.5].

8.6. Orientation data on deformed 3-Calabi–Yau completions. We will now prove the existence of a canonical orientation data for deformed 3-Calabi–Yau completions of a dg category [Kel11]. Let \mathcal{C} be a finite type dg category and $\tilde{c} \in \text{HC}_1^-(\mathcal{C})$ be a negative cyclic class whose underlying Hochschild class is denoted by c . Following [Kel11, §5.1], we defined the deformed Calabi–Yau completion $\Pi_3(\mathcal{C}, \tilde{c})$ of \mathcal{C} associated with \tilde{c} by the following pushout diagram:

$$\begin{array}{ccc} T_e(\mathcal{C}^! [2]) & \xrightarrow{[\text{id}, 0]} & \mathcal{C} \\ \downarrow [\text{id}, c] & & \downarrow \Gamma \\ \mathcal{C} & \longrightarrow & \Pi_3(\mathcal{C}, \tilde{c}) \end{array}$$

where $\mathcal{C}^!$ denotes the inverse dualizing bimodule and $T_e(\mathcal{C}^! [2])$ denotes the tensor algebra dg category. It is shown in [BCS24, Theorem 5.9] that $\Pi_3(\mathcal{C}, \tilde{c})$ is finite type and [Kel18] that the class \tilde{c} induces a left 3-Calabi–Yau structure on $\Pi_3(\mathcal{C}, \tilde{c})$.

Example 8.31. Let Q be a (finite) quiver. We let $\text{Rep}_Q^{\text{Perf}}$ denote the dg category of perfect Q -representations. Take a potential $W \in \text{HH}_0(\text{Rep}_Q^{\text{Perf}})$. Then Ginzburg [Gin06, §5] constructed a non-positively graded dg algebra $\Gamma_3(Q, W)$ whose zeroth cohomology is the Jacobi algebra $\text{Jac}(Q, W)$. Then it is shown in [Kel11,

Theorem 6.2] that there exists a Morita equivalence

$$\Pi_3(\mathrm{Rep}_Q^{\mathrm{Perf}}, \delta W) \simeq \mathrm{Mod}_{\Gamma_3(Q, W)}^{\mathrm{Perf}}.$$

where δ denotes the mixed differential.

Example 8.32. Let S be a smooth algebraic surface and set $X := \mathrm{Tot}_S(\omega_S)$. A Calabi–Yau threefold of this form is called a **local surface**. Then it is shown in [KM24, Corollary 4.2] that there exists a natural Morita equivalence of left 3-Calabi–Yau categories

$$\Pi_3(\mathrm{Perf}(S), 0) \simeq \mathrm{Perf}(X).$$

Example 8.33. Let C be a smooth projective curve and N be a rank two vector bundle with $\det(N) \simeq \omega_C$ and set $X := \mathrm{Tot}_C(N)$. A Calabi–Yau threefold of this form is called a **local curve**. Then by [KK21, Lemma 3.3] we may find a line bundle L with a surjective morphism $N \rightarrow L$. Set $Y := \mathrm{Tot}_C(L)$. Then it is shown in [KM24, Theorem 4.3] there exists a Hochschild homology class $c \in \mathrm{HH}_0(\mathrm{Perf}(Y))$ with a Morita equivalence of left 3-Calabi–Yau categories

$$\Pi_3(\mathrm{Perf}(Y), \delta c) \simeq \mathrm{Perf}(X).$$

Let \mathcal{C} be a finite type dg category and $c \in \mathrm{HH}_0(\mathcal{C})$ be a Hochschild homology class. We have a natural map

$$(8.11) \quad |\mathrm{HH}(\mathcal{C})| \rightarrow \Gamma(\mathfrak{M}_c, \mathcal{O}_{\mathfrak{M}_c})$$

as constructed in [BD21, §5.2] and let $f_c: \mathfrak{M}_c \rightarrow \mathbb{A}^1$ be the corresponding function. Then it is shown in [BCS24, Corollary 6.19] that there exists a natural equivalence of (-1) -shifted symplectic stacks

$$(8.12) \quad \mathfrak{M}_{\Pi_3(c, \delta c)} \simeq \mathrm{Crit}(f_c)$$

where the (-1) -shifted symplectic structure for the left-hand side is constructed using (8.8) and for the right-hand side is constructed using Example 3.10. Using this equivalence, we will prove the following result:

Proposition 8.34. *The deformed 3-Calabi–Yau completion $\Pi_3(\mathcal{C}, \delta c)$ admits a strong orientation data.*

Proof. We first discuss the generality of the canonical orientation for the derived critical locus defined in Example 3.20. If we are given regular functions f_1 and f_2 on derived Artin stacks \mathfrak{Y}_1 and \mathfrak{Y}_2 with $\mathrm{Crit}(f_i) = \mathfrak{X}_i$, there exists a natural isomorphism

$$(8.13) \quad o_{\mathfrak{X}_1}^{\mathrm{can}} \boxtimes o_{\mathfrak{X}_2}^{\mathrm{can}} \simeq o_{\mathfrak{X}_1 \times \mathfrak{X}_2}^{\mathrm{can}}$$

where the canonical orientation $o_{\mathfrak{X}_1 \times \mathfrak{X}_2}^{\mathrm{can}}$ is constructed using the critical locus description $\mathfrak{X}_1 \times \mathfrak{X}_2 \simeq \mathrm{Crit}(f_1 \boxplus f_2)$. Further, if we are given yet another derived Artin stack \mathfrak{Y}_3 with a regular function f_3 on it, if we write $\mathfrak{X}_3 = \mathrm{Crit}(f_3)$, the following diagram commutes:

$$(8.14) \quad \begin{array}{ccc} o_{\mathfrak{X}_1}^{\mathrm{can}} \boxtimes o_{\mathfrak{X}_2}^{\mathrm{can}} \boxtimes o_{\mathfrak{X}_3}^{\mathrm{can}} & \xrightarrow{\simeq} & o_{\mathfrak{X}_1 \times \mathfrak{X}_2}^{\mathrm{can}} \boxtimes o_{\mathfrak{X}_3}^{\mathrm{can}} \\ \downarrow \simeq & & \downarrow \simeq \\ o_{\mathfrak{X}_1}^{\mathrm{can}} \boxtimes o_{\mathfrak{X}_2 \times \mathfrak{X}_3}^{\mathrm{can}} & \xrightarrow{\simeq} & o_{\mathfrak{X}_1 \times \mathfrak{X}_2 \times \mathfrak{X}_3}^{\mathrm{can}}. \end{array}$$

We now claim that the localization of the canonical orientation for $\mathfrak{X} = \mathrm{Crit}(f)$ to $\mathrm{Grad}(\mathfrak{X})$ is again a canonical orientation. By Lemma 5.15, there exists an equivalence of (-1) -shifted symplectic stacks $\mathrm{Grad}(\mathfrak{X}) \simeq \mathrm{Crit}(f|_{\mathrm{Grad}(\mathfrak{X})})$. The direct sum decomposition $\mathbb{L}_{\mathfrak{Y}}|_{\mathrm{Grad}(\mathfrak{Y})} = \mathbb{L}_{\mathrm{Grad}(\mathfrak{Y})} \oplus \mathbb{L}_{\mathfrak{Y}}|_{\mathrm{Grad}(\mathfrak{Y})}^{\pm}$ induces a natural isomorphism

$$\widehat{\det}(\mathbb{L}_{\mathfrak{Y}}) \otimes \widehat{\det}(\mathbb{L}_{\mathfrak{Y}}|_{\mathrm{Grad}(\mathfrak{Y})}^{\pm})^{\vee} \simeq \widehat{\det}(\mathbb{L}_{\mathrm{Grad}(\mathfrak{Y})})$$

which clearly induces an isomorphism of orientations

$$(8.15) \quad u^* o_{\mathfrak{X}}^{\mathrm{can}} \simeq o_{\mathrm{Grad}(\mathfrak{X})}^{\mathrm{can}}.$$

The following diagram commutes:

$$(8.16) \quad \begin{array}{ccccc} u_1^* u^* o_{\mathfrak{X}}^{\text{can}} |_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}} & \xrightarrow{\cong} & u_1^* o_{\text{Grad}(\mathfrak{X})}^{\text{can}} |_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}} & \xrightarrow{\cong} & o_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}}^{\text{can}} \\ \downarrow \cong & & & & \parallel \\ u_2^* u^* o_{\mathfrak{X}}^{\text{can}} |_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}} & \xrightarrow{\cong} & u_2^* o_{\text{Grad}(\mathfrak{X})}^{\text{can}} |_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}} & \xrightarrow{\cong} & o_{\text{Grad}^2(\mathfrak{X})_{\text{mat}}}^{\text{can}} \end{array}$$

If we are given derived Artin stacks $\mathfrak{X} = \text{Crit}(f: \mathfrak{Y} \rightarrow \mathbb{A}^1)$, $\mathfrak{X}' = \text{Crit}(f': \mathfrak{Y}' \rightarrow \mathbb{A}^1)$, the following diagram commutes by construction:

$$(8.17) \quad \begin{array}{ccc} (u \times u')^*(o_{\mathfrak{X}}^{\text{can}} \boxtimes o_{\mathfrak{X}'}^{\text{can}}) & \xrightarrow{\cong} & (u \times u')^*(o_{\mathfrak{X} \times \mathfrak{X}'}^{\text{can}}) \\ \downarrow \cong & & \downarrow \cong \\ u^* o_{\mathfrak{X}}^{\text{can}} \boxtimes u'^* o_{\mathfrak{X}'}^{\text{can}} & \xrightarrow{\cong} & o_{\text{Grad}(\mathfrak{X}) \times \text{Grad}(\mathfrak{X}')}^{\text{can}} \end{array}$$

We now return to the statement of the proposition. We equip $\mathfrak{M}_{\Pi_3(\mathcal{C}, \delta_c)} \simeq \text{Crit}(f_c)$ with the canonical orientation described above. Consider the diagonal map $\Delta: \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$. Under the identification of $\text{HH}(\mathcal{C} \times \mathcal{C}) \simeq \text{HH}(\mathcal{C}) \oplus \text{HH}(\mathcal{C})$, we can identify $\Delta_* c$ and (c, c) . In particular, the composition

$$\mathfrak{M}_{\mathcal{C}} \times \mathfrak{M}_{\mathcal{C}} \xrightarrow{\Phi_2} \mathfrak{M}_{\mathcal{C}} \xrightarrow{f_c} \mathbb{A}^1$$

is given by $f_c \boxplus f_c$. Then by using (8.13) and (8.15), we obtain a natural isomorphism

$$\Phi_2^*(o_{\mathfrak{M}_{\Pi_3(\mathcal{C}, \delta_c)}}^{\text{can}}) \simeq o_{\mathfrak{M}_{\Pi_3(\mathcal{C}, \delta_c)}^{\text{can}, \boxtimes 2}}.$$

Then the existence of the strong orientation data follows from the commutativity of the diagrams (8.14), (8.16) and (8.17). □

Remark 8.35. By the same proof, we can show that the deformed 3-Calabi–Yau completion $\Pi_3(\mathcal{C}, \tilde{c})$ admits a strong orientation data for arbitrary negative cyclic class $\tilde{c}: k \rightarrow |\text{HC}^-(\mathcal{C})|$.

The following statement is a consequence of Proposition 8.34 and Example 8.31, 8.32 and 8.33:

Corollary 8.36. *The following left 3-Calabi–Yau categories admit a strong orientation data:*

- (1) *The dg category of perfect modules $\text{Mod}_{\Gamma_3(Q, W)}^{\text{Perf}}$ over the Ginzburg dg algebra $\Gamma_3(Q, W)$ associated with a quiver with potential (Q, W) .*
- (2) *The dg category of perfect complexes $\text{Perf}(X)$ for a local surface $X = \text{Tot}_C(\omega_C)$.*
- (3) *The dg category of perfect complexes $\text{Perf}(X)$ for a local curve $X = \text{Tot}_C(N)$.*

8.7. Comparison with the Kontsevich–Soibelman’s CoHA for quivers with potentials. In this section, we will prove that the cohomological Hall algebra structure for the cohomological Donaldson–Thomas invariants for quivers with potentials constructed in Corollary 8.8 coincide with the one constructed in [KS11, §7.6].

Let Q be a finite quiver with Q_0 the set of vertices and Q_1 the set of edges. For a dimension vector $\mathbf{d} \in \mathbb{Z}_{\geq 0}^{Q_0}$, we let $\mathfrak{M}_{\mathbf{d}}$ denote the moduli stack of representations of Q with dimension vector \mathbf{d} . We define a pairing $(-, -)_Q$ of dimension vectors by

$$(\mathbf{d}, \mathbf{d}')_Q := \sum_{(e: i \rightarrow j) \in Q_1} \mathbf{d}_i \mathbf{d}'_j - \mathbf{d}_j \mathbf{d}'_i.$$

Consider the following correspondence:

$$\begin{array}{ccc} & \mathfrak{M}_{\mathbf{d}, \mathbf{d}'}^{2\text{-filt}} & \\ \swarrow \text{gr} & & \searrow \text{ev} \\ \mathfrak{M}_{\mathbf{d}} \times \mathfrak{M}_{\mathbf{d}'} & & \mathfrak{M}_{\mathbf{d} + \mathbf{d}'} \end{array}$$

Since all spaces appearing in this diagram are smooth, we obtain a natural isomorphism

$$(8.18) \quad \mathbb{Q}_{\mathfrak{M}_{\mathbf{d}}}^{\text{Perv}} \boxtimes \mathbb{Q}_{\mathfrak{M}_{\mathbf{d}'}}^{\text{Perv}}[(\mathbf{d}, \mathbf{d}')_Q] \cong \text{gr}_* \text{ev}^! \mathbb{Q}_{\mathfrak{M}_{\mathbf{d}+\mathbf{d}'}}^{\text{Perv}}$$

where \mathbb{Q}^{Perv} denotes the constant sheaf shifted to be perverse. Assume now that we are given a potential $W: \mathbb{C} \rightarrow |\text{HH}(\text{Rep}_Q^{\text{Perf}})|$ and $f_{W,\mathbf{d}}: \mathfrak{M}_{Q,\mathbf{d}} \rightarrow \mathbb{A}^1$ be the associated function. Then it is clear that we have an identity $f_{W,\mathbf{d}+\mathbf{d}'} \circ \text{ev} = (f_{W,\mathbf{d}} \boxplus f_{W,\mathbf{d}'}) \circ \text{gr}$. Therefore, by applying the vanishing cycle functor $\varphi_{f_{W,\mathbf{d}} \boxplus f_{W,\mathbf{d}'}}$ to the isomorphism (8.18) and using the Thom–Sebastiani isomorphism (4.3) and (4.2), we obtain the following map⁸ of shifted perverse sheaves

$$(8.19) \quad \varphi_{f_{W,\mathbf{d}}} \boxtimes \varphi_{f_{W,\mathbf{d}'}}[(\mathbf{d}, \mathbf{d}')_Q] \rightarrow \text{gr}_* \text{ev}^! \varphi_{f_{W,\mathbf{d}+\mathbf{d}'}}.$$

Write $\mathcal{H}_{Q,W,\mathbf{d}} := H^*(\mathfrak{M}_{\mathbf{d}}, \varphi_{f_{W,\mathbf{d}}})$. By passing the map (8.19) to the left adjoint and using the properness of ev , we obtain a morphism

$$*_{\text{KS}}^{\text{Hall}}: \mathcal{H}_{Q,W,\mathbf{d}} \otimes^{\text{tw}} \mathcal{H}_{Q,W,\mathbf{d}'} \rightarrow \mathcal{H}_{Q,W,\mathbf{d}+\mathbf{d}'},$$

which defines an algebra structure on $\mathcal{H}_{Q,W} := \bigoplus_{\mathbf{d}} \mathcal{H}_{Q,W,\mathbf{d}}$ and is called Kontsevich–Soibelman’s cohomological Hall algebras for (Q, W) .

Now take $\mathcal{C} = \text{Mod}_{\Gamma_3(Q,W)}^{\text{Perf}}$ where $\Gamma_3(Q, W)$ denotes the Gizburg dg algebra. Let $\mathfrak{M} \subset \mathfrak{M}_{\mathcal{C}}$ denote the locus consisting of modules concentrated in degree zero. Then there exists a natural identification $\pi_0(\mathfrak{M}) \cong \mathbb{Z}_{\geq 0}^{Q_0}$ and we have natural isomorphism of (-1) -shifted symplectic stacks

$$\mathfrak{M} \cong \coprod_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^{Q_0}} \text{Crit}(f_{W,\mathbf{d}})$$

as we have seen in (8.12). We let $\mathfrak{M}_{\mathbf{d}} \subset \mathfrak{M}$ denotes the open and closed substack corresponding to $\text{Crit}(f_{W,\mathbf{d}})$. We equip \mathfrak{M} with the strong orientation data constructed in Proposition 8.34 and write

$$\mathcal{H}_{Q,W}^{3-\text{CY}} := \bigoplus_{\mathbf{d} \in \mathbb{Z}^{Q_0}} H^{\bullet}(\mathfrak{M}_{\mathbf{d}}, \varphi_{\mathfrak{M}_{\mathbf{d}}}).$$

We denote the multiplication constructed in Corollary 8.8 by $*_{3-\text{CY}}^{\text{Hall}}$. The following statement is an immediate consequence of Lemma 7.21 and Proposition 7.22:

Proposition 8.37. *There exists a natural isomorphism of associative algebras*

$$(\mathcal{H}_{Q,W}, *_{\text{KS}}^{\text{Hall}}) \cong (\mathcal{H}_{Q,W}^{3-\text{CY}}, *_{3-\text{CY}}^{\text{Hall}}).$$

9. APPLICATIONS TO COHOMOLOGICAL DT INVARIANTS OF 3-MANIFOLDS

In this section we explain how to construct a parabolic induction map for cohomological DT invariants of 3-manifolds as in [NS23, Theorem 3.47] as well as define a cohomological Hall algebra structure.

9.1. Local systems. Let $M \in \mathcal{S}$ be a space and $\text{LocSys}(M) = \text{Fun}(M, \text{Mod}_{\mathbb{C}})$ the presentable stable ∞ -category of chain complexes of local systems of \mathbb{C} -vector spaces on M . Let $\mathcal{C} = \text{LocSys}(M)^{\omega} \subset \text{LocSys}(M)$ be the subcategory of compact objects. For a morphism $f: M \rightarrow N$ of spaces we have the restriction functor $f^*: \text{LocSys}(N)^{\omega} \rightarrow \text{LocSys}(M)^{\omega}$ and its left adjoint $f_{\#}: \text{LocSys}(M)^{\omega} \rightarrow \text{LocSys}(N)^{\omega}$. Recall that M is *finitely dominated* if it is a compact object in \mathcal{S} . In this case, we have the following objects:

- The constant local system $\mathbb{C}_M \in \text{LocSys}(M)$ with fibre \mathbb{C} is compact, i.e. it defines an object $\mathbb{C}_M \in \text{LocSys}(M)^{\omega}$.
- There is an orientation local system $\zeta_M \in \text{LocSys}(M)^{\omega}$ which represents the functor of homology $\mathbb{C}_{\bullet}(M; -): \text{LocSys}(M)^{\omega} \rightarrow \text{Perf}_{\mathbb{C}}$.

Lemma 9.1. *Let M be a connected space. Then the functor $\text{Map}(M_{\mathbb{B}}, -): \mathbf{dSt} \rightarrow \mathbf{dSt}$ commutes with coproducts.*

⁸One can show that this map is an isomorphism by repeating the proof of Theorem 7.16 or by Proposition A.1.

Proof. Let I be a set. The colimit functor $\text{Fun}(I, \mathbf{dSt}) \rightarrow \mathbf{dSt}$ lifts to an equivalence $\text{Fun}(I, \mathbf{dSt}) \cong \mathbf{dSt}/_{I_B}$. As $\text{Map}(M_B, -)$ is an M -indexed limit, we have to show that the forgetful functor $U: \mathbf{dSt}/_{I_B} \rightarrow \mathbf{dSt}$ preserves M -indexed limits. For a diagram $F: M \rightarrow \mathbf{dSt}/_{I_B}$ we have

$$U \left(\lim_{x \in M} F(x) \right) \cong \left(\lim_{x \in M} UF(x) \right) \times_{\lim_{x \in M} I_B} I_B.$$

The limit $\lim_{x \in M} I_B$ is equivalent to $\text{Hom}_{\mathcal{S}}(M, I)_B$ and since I is discrete, the constant map $I \rightarrow \text{Hom}_{\mathcal{S}}(M, I)$ is an equivalence. \square

Definition 9.2. An *oriented d -dimensional Poincaré space* is a finitely dominated space $M \in \mathcal{S}$ equipped with a fundamental class $[M] \in C_d(M; \mathbb{C})$ such that the induced morphism $\zeta_M \rightarrow \mathbb{C}_M[-d]$ is an isomorphism.

It is shown in [BD19, Proposition 5.1] that if M is finitely dominated, $\text{LocSys}(M)^\omega$ is a finite type dg category. Moreover, if M is an oriented d -dimensional Poincaré space, the image of the fundamental class under the inclusion of constant loops

$$i: C_\bullet(M; \mathbb{C}) \longrightarrow (C_\bullet(LM; \mathbb{C}))^{S^1} \simeq \text{HC}^-(\text{LocSys}(M)^\omega)$$

defines a left d -Calabi–Yau structure on $\text{LocSys}(M)^\omega$ by [BD19, Theorem 5.4].

Theorem 9.3. *Let $M \in \mathcal{S}$ be a finitely dominated space. Then there is an equivalence of derived stacks*

$$\text{Map}(M_B, \text{Perf}) \simeq \mathfrak{M}_{\text{LocSys}(M)^\omega}.$$

Moreover, if M is an oriented d -dimensional Poincaré space, this is an equivalence of $(2-d)$ -shifted symplectic stacks, where on the left the $(2-d)$ -shifted symplectic structure is induced by the AKSZ procedure from the 2-shifted symplectic structure on Perf and the d -orientation of M_B while on the right the $(2-d)$ -shifted symplectic structure is defined by the d -Calabi–Yau structure on $\text{LocSys}(M)^\omega$.

Proof. Consider the functor, natural in $M \in \mathcal{S}$,

$$M \longrightarrow \text{LocSys}(M)^\omega$$

given by $\text{pt} \mapsto \mathbb{C} \in \text{Perf}_{\mathbb{C}}$ for $M = \text{pt}$. Such a functor is unique as $M = \text{colim}_{x \in M} \text{pt}$. This functor defines a universal object

$$E_M \in \text{Perf}(M_B; \text{LocSys}(M)^\omega) \cong \lim_{x \in M} (\text{LocSys}(M)^\omega).$$

Applying Proposition 8.24 we obtain a morphism

$$F_{E_M}: \mathfrak{M}_{\text{LocSys}(M)^\omega} \longrightarrow \text{Map}(M_B, \text{Perf}).$$

The object E_M is functorial in M in the sense that it is the value on M of a universal object

$$E \in \int_{M \in \mathcal{S}} \text{Perf}(M_B; \text{LocSys}(M)^\omega).$$

The construction $M \mapsto \text{LocSys}(M)^\omega$ defines a functor $\mathcal{S} \rightarrow \mathbf{dgcst}$ which preserves colimits. Therefore, both $\text{Map}(M_B, \text{Perf})$ and $\mathfrak{M}_{\text{LocSys}(M)^\omega}$ viewed as functors $\mathcal{S}^{\text{op}} \rightarrow \mathbf{dSt}$ send colimits of spaces to limits of derived stacks. Moreover, F_{E_M} is obviously an isomorphism for $M = \text{pt}$ and hence it is an isomorphism for any M . This proves the first statement.

Next, suppose M is an oriented d -dimensional Poincaré space with fundamental class $[M] \in C_d(M; \mathbb{C})$. Using Theorem 8.25 we get that the morphism

$$\mathfrak{M}_{\text{LocSys}(M)^\omega} \longrightarrow \text{Map}(M_B, \text{Perf})$$

is compatible with $(2-d)$ -shifted symplectic structures, with a certain d -Calabi–Yau structure $[\text{LocSys}(M)^\omega]$ on $\text{LocSys}(M)^\omega$ described in the statement. It remains to show that

$$[\text{LocSys}(M)^\omega] \sim i([M]) \in |\text{HC}^-(\text{LocSys}(M)^\omega)[-d]|.$$

By definition the class $[\text{LocSys}(M)^\omega]$ is given by the composite

$$\begin{aligned} \mathbb{C} &\xrightarrow{[E_M]} \text{HC}^-(\text{Perf}(M_B; \text{LocSys}(M)^\omega)) \\ &\xrightarrow{\kappa_{\text{LocSys}(M)^\omega, M_B}} \mathbb{C}^\bullet(M) \otimes \text{HC}^-(\text{LocSys}(M)^\omega) \\ &\xrightarrow{[M] \otimes \text{id}} \text{HC}^-(\text{LocSys}(M)^\omega)[-d] \end{aligned}$$

By definition of E_M this composite is equivalent to the image of $[M]$ under a map

$$\mathbb{C}_\bullet(M; \mathbb{C}[[u]]) \longrightarrow \mathbb{C}_\bullet(LM; \mathbb{C})^{S^1} \cong \text{HC}^-(\text{LocSys}(M)^\omega)$$

which is natural in M and which is the identity for $M = \text{pt}$. The inclusion of constant loops

$$i: \mathbb{C}_\bullet(M; \mathbb{C}[[u]]) \longrightarrow \mathbb{C}_\bullet(LM; \mathbb{C})^{S^1}$$

is also natural in M and is the identity for $M = \text{pt}$. But since $\mathbb{C}_\bullet(M; \mathbb{C}[[u]])$ preserves colimits in M , such a natural transformation is unique, which implies the claim. \square

Example 9.4. Let $M = S^1$ be the circle which has a fundamental class $[S^1] \in C_1(S^1; \mathbb{C})$. Then we can identify $\text{LocSys}(S^1)^\omega \cong \text{Perf}_{\mathbb{C}[x^{\pm 1}]}$ and Theorem 9.3 in this case reduces to a 1-shifted symplectic equivalence

$$\text{Map}(S^1_B, \text{Perf}) \cong \mathfrak{M}_{\text{Perf}_{\mathbb{C}[x^{\pm 1}]}}$$

from [BCS23, Proposition 5.1].

9.2. Parabolic restriction for character stacks and its orientation. Let us now consider a more general setting. For an algebraic group G we denote by

$$\text{Loc}_G(M) = \text{Map}(M_B, BG)$$

the *character stack*. Fixing a point $x \in M$, we have the *framed character stack*

$$\text{Loc}_G^{\text{fr}}(M) = \text{Loc}_G(M) \times_{BG} \text{pt},$$

where the morphism $\text{Loc}_G(M) \rightarrow BG$ is given by evaluation at x . If M is connected, the framed character stack is represented by an affine derived scheme and

$$\text{Loc}_G(M) \cong [\text{Loc}_G^{\text{fr}}(M)/G].$$

Recall the following facts about shifted symplectic structures on classifying stacks:

- If G is equipped with a non-degenerate G -invariant symmetric bilinear pairing $c \in \text{Sym}^2(\mathfrak{g}^*)^G$ on its Lie algebra \mathfrak{g} , there is a canonical 2-shifted symplectic structure on BG .
- If G is connected reductive with $P \subset G$ a parabolic subgroup with Levi factor L , then there is a unique 2-shifted Lagrangian structure on the correspondence

$$(9.1) \quad \begin{array}{ccc} & BP & \\ & \swarrow \quad \searrow & \\ BL & & BG \end{array}$$

which restricts to the chosen 2-shifted symplectic structure on BG , see [Saf17, Lemma 3.4].

Proposition 9.5. *Let G be a connected reductive group equipped with a non-degenerate pairing c and $P_1, P_2 \subset G$ parabolic subgroups with the same Levi factor $L \subset G$. Then the diagram*

$$(9.2) \quad \begin{array}{ccccc} & & BP_1 & & \\ & \swarrow & \uparrow & \searrow & \\ BL & & B(P_1 \cap P_2) & & BG \\ & \swarrow & \downarrow & \searrow & \\ & & BP_2 & & \end{array}$$

has a unique structure of a 2-fold 2-shifted Lagrangian correspondence, which restricts to the chosen 2-shifted symplectic structure on BG .

Proof. If H is an affine algebraic group, the space $\mathcal{A}^{2,\text{cl}}(BH, 2)$ of closed two-forms on BH of degree 2 is equivalent to the set $\text{Sym}^2(\mathfrak{h}^*)^H$ of H -invariant symmetric bilinear pairings on the Lie algebra of H .

We first prove that there is a unique structure of a 2-fold 2-shifted isotropic correspondence which restricts to the chosen 2-shifted presymplectic structure on BG . Let $\mathfrak{p}_1, \mathfrak{p}_2$ and \mathfrak{p}_{12} be the Lie algebras of P_1, P_2 and $P_1 \cap P_2$. Consider the correspondence

$$\begin{array}{ccccc}
 & & \text{Sym}^2(\mathfrak{p}_1^*)^{P_1} & & \\
 & \nearrow & \downarrow & \nwarrow & \\
 \text{Sym}^2(\mathfrak{t}^*)^L & & \text{Sym}^2(\mathfrak{p}_{12}^*)^{P_1 \cap P_2} & & \text{Sym}^2(\mathfrak{g}^*)^G \\
 & \searrow & \uparrow & \swarrow & \\
 & & \text{Sym}^2(\mathfrak{p}_2^*)^{P_2} & &
 \end{array}$$

given by the pullback of closed two-forms along the diagram (9.2).

Using [Spr09, Theorem 6.2.7] we may choose Borel subgroups $B_i \subset G$ contained in P_i . By [Spr09, Corollary 8.3.10] the intersection $B_1 \cap B_2$ contains a maximal torus $T \subset G$; let \mathfrak{t} be its Lie algebra. Let

$$\mathfrak{p}_1 = \mathfrak{l} \oplus \mathfrak{u}_1, \quad \mathfrak{p}_2 = \mathfrak{l} \oplus \mathfrak{u}_2$$

be L -invariant splittings.

Let $(-, -)$ be an L -invariant pairing on \mathfrak{p}_i . For any $x, y \in \mathfrak{p}_i$ of T -weights α, β we have $(\alpha + \beta)(x, y) = 0$. In particular, this implies that $(\mathfrak{u}_i, \mathfrak{p}_i) = 0$. Thus, the map $\text{Sym}^2(\mathfrak{t}^*)^L \rightarrow \text{Sym}^2(\mathfrak{p}_i^*)^L$ is an isomorphism. We have $[\mathfrak{u}_i, \mathfrak{p}_i] \subset \mathfrak{u}_i$, so the map $\text{Sym}^2(\mathfrak{p}_i^*)^{P_i} \rightarrow \text{Sym}^2(\mathfrak{p}_i^*)^L$ is an isomorphism. An analogous argument shows that $\text{Sym}^2(\mathfrak{t}^*)^L \rightarrow \text{Sym}^2(\mathfrak{p}_{12}^*)^{P_1 \cap P_2}$ is an isomorphism. The two maps $\text{Sym}^2(\mathfrak{g}^*)^G \rightarrow \text{Sym}^2(\mathfrak{p}_i^*)^{P_i} \xleftarrow{\sim} \text{Sym}^2(\mathfrak{t}^*)^L$ are equal to the restriction along $L \subset G$. Since the restriction $\text{Sym}^2(\mathfrak{g}^*)^G \rightarrow \text{Sym}^2(\mathfrak{t}^*)$ is injective, so is the restriction $\text{Sym}^2(\mathfrak{g}^*)^G \rightarrow \text{Sym}^2(\mathfrak{p}_i^*)^{P_i}$. This finishes the construction of the unique structure of a 2-fold 2-shifted isotropic correspondence.

Let us now show that if the chosen pairing on \mathfrak{g} is non-degenerate, then this 2-fold 2-shifted isotropic correspondence is Lagrangian. The fact that $BL \leftarrow BP_i \rightarrow BG$ is a 2-shifted Lagrangian correspondence is shown in [Saf17, Lemma 3.4]. So, we have to check that $B(P_1 \cap P_2) \rightarrow BP_1 \times_{BG \times BL} BP_2$ is a 1-shifted Lagrangian morphism. This boils down to showing that the complex

$$\mathfrak{p}_{12} \rightarrow \mathfrak{p}_1 \oplus \mathfrak{p}_2 \rightarrow \mathfrak{g} \oplus \mathfrak{l} \rightarrow \mathfrak{p}_{12}^*$$

concentrated in degrees $[0, 3]$ is acyclic. Since $\mathfrak{p}_{12} \rightarrow \mathfrak{p}_1 \oplus \mathfrak{p}_2$ is injective, H^0 vanishes. Its H^1 vanishes since $(\mathfrak{p}_1 \oplus \mathfrak{p}_2)/\mathfrak{p}_{12} \rightarrow \mathfrak{g}$ is injective. Its H^3 vanishes since the composite $\mathfrak{g} \cong \mathfrak{g}^* \rightarrow \mathfrak{p}_{12}^*$ is surjective. The Euler characteristic of this complex is $\dim \mathfrak{g} + \dim \mathfrak{l} - \dim \mathfrak{p}_1 - \dim \mathfrak{p}_2$. Since $\dim \mathfrak{p}_i = (\dim \mathfrak{g} + \dim \mathfrak{l})/2$, the Euler characteristic of this complex vanishes and hence $H^2 = 0$, i.e. this complex is acyclic. \square

In particular, if M is a closed oriented 3-manifold and G is equipped with a non-degenerate pairing on its Lie algebra, there is a (-1) -shifted symplectic structure on $\text{Loc}_G(M)$ defined by the AKSZ procedure. Let us recall the orientation on character stacks following [NS23].

Proposition 9.6. *Let M be a closed spin 3-manifold.*

- (1) *For a connected reductive group G equipped with a non-degenerate pairing $c \in \text{Sym}^2(\mathfrak{g}^*)^G$ there is an orientation o_G on $\text{Loc}_G(M)$.*
- (2) *For a pair of connected reductive groups G_1, G_2 equipped with non-degenerate pairings $c_i \in \text{Sym}^2(\mathfrak{g}_i^*)^{G_i}$ there is an isomorphism $o_{G_1 \times G_2} \simeq o_{G_1} \boxtimes o_{G_2}$ of orientation on $\text{Loc}_{G_1 \times G_2}(M) \simeq \text{Loc}_{G_1}(M) \times \text{Loc}_{G_2}(M)$ which satisfies an associativity relation.*

- (3) For a parabolic subgroup $P \subset G$ whose Levi factor is L there is an orientation ρ_P of the Lagrangian correspondence

$$\begin{array}{ccc} & \text{Loc}_P(M) & \\ & \swarrow \quad \searrow & \\ \text{Loc}_L(M) & & \text{Loc}_G(M) \end{array}$$

which restricts to o_L and o_G on $\text{Loc}_L(M)$ and $\text{Loc}_G(M)$. For $P = G$ the orientation ρ_P is the identity orientation of the diagonal Lagrangian correspondence. The orientation ρ_P for conjugate parabolic subgroups are equal.

- (4) Given a pair of parabolic subgroups $P_1 \subset G_1$ and $P_2 \subset G_2$ with Levi factors L_1 and L_2 , under the isomorphisms λ_{L_1, L_2} and λ_{G_1, G_2} we have an equality $\rho_{P_1} \boxtimes \rho_{P_2} = \rho_{P_1 \times P_2}$ of orientations of the product correspondence

$$\begin{array}{ccc} & \text{Loc}_{P_1}(M) \times \text{Loc}_{P_2}(M) & \\ & \swarrow \quad \searrow & \\ \text{Loc}_{L_1}(M) \times \text{Loc}_{L_2}(M) & & \text{Loc}_{G_1}(M) \times \text{Loc}_{G_2}(M) \end{array}$$

Proof. The claim is essentially shown in [NS23, Theorem 3.45] to which we refer for details. Let us sketch how to construct the relevant compatibility isomorphisms. First, as explained in [NS23, Remark 3.15], there is a canonical homology orientation of M , i.e. a trivialization of the (graded) line $\det H_\bullet(M; \mathbb{C})$ (note that it lies in degree zero as $\chi(M) = 0$). The spin structure on M allows us to choose a canonical Euler structure in the sense of [FT99, Section 3.2], i.e. a trivialization of the homological Euler class $\xi: e(M) \sim 0$ whose characteristic class is $c(\xi) = 0 \in H_1(M; \mathbb{Z})$. Equivalently, it is the choice of a Spin^c structure with the trivial c_1 . Let G be an algebraic group and consider the derived stack $\mathfrak{X} = \text{Loc}_G(M)$. By [NS23, Proposition 3.23] we get that $\text{vd}(\text{Loc}_G(M)) = 0$ and there is a trivialization of the determinant of the cotangent complex of $\text{Loc}_G(M)$ given by the torsion volume form vol_G . Let $\overline{\text{vol}}_G$ be the corresponding section of $\widehat{\det}(\mathbb{L}_{\mathfrak{X}})$. For a pair of algebraic groups G_1, G_2 we have an equality $\overline{\text{vol}}_{G_1 \times G_2} = \overline{\text{vol}}_{G_1} \wedge \overline{\text{vol}}_{G_2}$ as sections of

$$\widehat{\det}(\mathbb{L}_{\text{Loc}_{G_1 \times G_2}(M)}) \simeq \widehat{\det}(\mathbb{L}_{\text{Loc}_{G_1}(M)}) \boxtimes \widehat{\det}(\mathbb{L}_{\text{Loc}_{G_2}(M)}).$$

Note that for sections of line bundles on classical stacks there is no room for nontrivial homotopies).

- (1) We define the orientation o_G to be given by the structure sheaf $\mathcal{O}_{\text{Loc}_G(M)^{\text{red}}}$ equipped with an isomorphism $\mathcal{O}_{\text{Loc}_G(M)^{\text{red}}}^{\otimes 2} \simeq \widehat{\det}(\mathbb{L}_{\text{Loc}_G(M)})$ given by $1 \mapsto \overline{\text{vol}}_G$.
- (2) We have an isomorphism

$$\mathcal{O}_{\text{Loc}_{G_1 \times G_2}(M)^{\text{red}}} \simeq \mathcal{O}_{\text{Loc}_{G_1}(M)^{\text{red}}} \boxtimes \mathcal{O}_{\text{Loc}_{G_2}(M)^{\text{red}}}$$

given by $1 \mapsto 1 \boxtimes 1$. The equality $\overline{\text{vol}}_{G_1 \times G_2} = \overline{\text{vol}}_{G_1} \wedge \overline{\text{vol}}_{G_2}$ then implies that it defines an isomorphism of orientations $o_{G_1 \times G_2} \simeq o_{G_1} \boxtimes o_{G_2}$. The associativity of this isomorphism follows from the associativity of the above isomorphism of structure sheaves.

- (3) We define the orientation ρ_P using the isomorphism $\mathcal{O}_{\text{Loc}_P(M)^{\text{red}}} \simeq \widehat{\det}(\mathbb{L}_{\text{Loc}_G(M)})$ given by $1 \mapsto \overline{\text{vol}}_P$. It is shown in [NS23, Theorem 3.45] that it is compatible with the orientation o_G and o_L of $\text{Loc}_G(M)$ and $\text{Loc}_L(M)$. Since it depends on the stack BP and not on the subgroup P , the orientation ρ_P is invariant under conjugation.
- (4) The multiplicativity of ρ_P follows from the relation $\overline{\text{vol}}_{P_1 \times P_2} = \overline{\text{vol}}_{P_1} \wedge \overline{\text{vol}}_{P_2}$.

□

So, if M is a closed spin 3-manifold and G a connected reductive group equipped with a non-degenerate pairing $c \in \text{Sym}^2(\mathfrak{g}^*)^G$, we obtain a perverse sheaf

$$\varphi_{\text{Loc}_G(M)} \in \text{Perv}(\text{Loc}_G(M))$$

given by the DT perverse sheaf associated to the (-1) -shifted symplectic derived Artin stack $\text{Loc}_G(M)$ equipped with the orientation o_G from Proposition 9.6. For a pair G_1, G_2 of such groups, the Thom–Sebastiani isomorphism from Proposition 4.3 as well as the isomorphism of orientations $o_{G_1 \times G_2} \simeq o_{G_1} \boxtimes o_{G_2}$

from Proposition 9.6 gives an isomorphism

$$\varphi_{\text{Loc}_{G_1 \times G_2}(M)} \cong \varphi_{\text{Loc}_{G_1}(M)} \boxtimes \varphi_{\text{Loc}_{G_2}(M)}$$

of perverse sheaves on $\text{Loc}_{G_1 \times G_2}(M) \simeq \text{Loc}_{G_1}(M) \times \text{Loc}_{G_2}(M)$ which satisfies an associativity relation.

One can compose parabolic subgroups in the following way.

Lemma 9.7. *Let G_3 be a connected reductive group with a parabolic subgroup $P_{23} \subset G_3$ with Levi factor G_2 and $P_{12} \subset G_2$ is a parabolic subgroup with Levi factor G_1 . Consider the fibre product $P_{13} = P_{12} \times_{G_2} P_{23}$. Then $P_{13} \subset G_3$ is a parabolic subgroup with Levi factor G_1 .*

Proof. Consider the diagram

$$\begin{array}{ccccc} & & BP_{13} & & \\ & \swarrow & & \searrow & \\ & BP_{12} & & BP_{23} & \\ & \swarrow \quad \searrow & & \swarrow \quad \searrow & \\ BG_1 & & BG_2 & & BG_3 \end{array}$$

where the top square is Cartesian. Since $BP_{12} \rightarrow BG_2$ is proper, the morphism $BP_{13} \rightarrow BP_{23}$ is proper. Since $BP_{23} \rightarrow BG_3$ is also proper, this implies that $BP_{13} \rightarrow BG_3$ is proper, i.e. $P_{13} \subset G_3$ is parabolic. Now, consider a Levi decomposition $P_{23} = G_2 \times U_{23}$, where U_{23} is the unipotent radical of G_2 . Therefore, $P_{13} = P_{12} \times U_{23}$. Its unipotent radical is $U_{12} \times U_{23}$, where $U_{12} \subset P_{12}$ is the unipotent radical of P_{12} . Therefore, the Levi factor of P_{13} is $P_{13}/(U_{12} \times U_{23}) \simeq G_1$. \square

Proposition 9.8. *Consider the setting of Lemma 9.7 and let M be a closed spin 3-manifold. Consider the diagram*

$$\begin{array}{ccccc} & & \text{Loc}_{P_{13}}(M) & & \\ & \swarrow & & \searrow & \\ & \text{Loc}_{P_{12}}(M) & & \text{Loc}_{P_{23}}(M) & \\ & \swarrow \quad \searrow & & \swarrow \quad \searrow & \\ \text{Loc}_{G_1}(M) & & \text{Loc}_{G_2}(M) & & \text{Loc}_{G_3}(M) \end{array}$$

where the top square is Cartesian. Then the composite of the orientation $\rho_{P_{23}}$ and $\rho_{P_{12}}$ is equal to $\rho_{P_{13}}$.

Proof. The orientation are defined using the torsion volume forms $\overline{\text{vol}}_{P_{i_j}}$ described in Proposition 9.6. The composite of the orientation $\rho_{P_{23}}$ and $\rho_{P_{12}}$ is defined using the volume form $\overline{\text{vol}}_{P_{12}}(M) \wedge \overline{\text{vol}}_{P_{23}}(M) \wedge \overline{\text{vol}}_{G_2}(M)^{-1}$ on $\text{Loc}_{P_{12}}(M) \times_{\text{Loc}_{G_2}(M)} \text{Loc}_{P_{23}}(M)$. This is precisely the glued volume form described in [NS23, Section 2.2]. But by construction the torsion volume forms are compatible with gluing, i.e. under the isomorphism

$$\text{Loc}_{P_{13}}(M) \simeq \text{Loc}_{P_{12}}(M) \times_{\text{Loc}_{G_2}(M)} \text{Loc}_{P_{23}}(M)$$

we have $\overline{\text{vol}}_{P_{13}} \mapsto \overline{\text{vol}}_{P_{12}}(M) \wedge \overline{\text{vol}}_{P_{23}}(M) \wedge \overline{\text{vol}}_{G_2}(M)^{-1}$. \square

9.3. Cohomological Hall algebra for 3-manifolds. In this section M denotes a connected closed spin 3-manifold and G a connected reductive group. Recall that for a cocharacter $\lambda: \mathbb{G}_m \rightarrow G$ we have defined the stabilizer and attractor subgroups $G_\lambda, G_\lambda^+ \subset G$ and for a pair of commuting cocharacters $\hat{\lambda} = (\lambda_1, \lambda_2): \mathbb{G}_m^2 \rightarrow G$ the subgroups $G_{\hat{\lambda}}, G_{\hat{\lambda}}^{+,+} \subset G$.

Proposition 9.9. *Let $T \subset G$ be a maximal torus and W the Weyl group. Choose a non-degenerate G -invariant symmetric bilinear pairing $c \in \text{Sym}^2(\mathfrak{g}^*)^G$ on the Lie algebra of G . Let M be a closed oriented 3-manifold.*

- (1) *For any cocharacter $\lambda: \mathbb{G}_m \rightarrow G$ the subgroup $G_\lambda^+ \subset G$ is parabolic and $G_\lambda \subset G_\lambda^+$ is the Levi factor.*
- (2) *Any parabolic subgroup $P \subset G$ arises as $G_\lambda^+ \subset G$ for some cocharacter $\lambda: \mathbb{G}_m \rightarrow G$.*

- (3) The natural morphisms $X_\bullet(T)/W \rightarrow X_\bullet(G)/G$ and $X_\bullet^2(T)/W \rightarrow X_\bullet^2(G)/G$ are isomorphisms.
(4) For a pair of commuting cocharacters $\hat{\lambda} = (\lambda_1, \lambda_2): \mathbb{G}_m^2 \rightarrow G$ the subgroup $(G_{\lambda_1}^+)_{\lambda_2} \times_{G_{\lambda_2}} G_{\lambda_1}^+ \subset G$ is parabolic with Levi factor $G_{\hat{\lambda}}$.
(5) We have

$$((G_{\lambda_1}^+)_{\lambda_2} \times_{G_{\lambda_2}} G_{\lambda_1}^+) \cap ((G_{\lambda_2}^+)_{\lambda_1} \times_{G_{\lambda_1}} G_{\lambda_2}^+) = G_{\hat{\lambda}}^{+,+}.$$

- (6) The attractor correspondence (5.1) for $\text{Loc}_G(M)$ is equivalent to

$$\begin{array}{ccc} & \coprod_{\lambda \in X_\bullet(T)/W} \text{Loc}_{G_\lambda^+}(M) & \\ \swarrow & & \searrow \\ \coprod_{\lambda \in X_\bullet(T)/W} \text{Loc}_{G_\lambda}(M) & & \text{Loc}_G(M) \end{array}$$

as a (-1) -shifted Lagrangian correspondence, where the latter structure is obtained by the AKSZ procedure from (9.1) with $P = G_\lambda^+$ and $L = G_\lambda$.

- (7) The diagram (5.2) for $\text{Loc}_G(M)$ is equivalent to

$$\begin{array}{ccccc} & & \coprod_{\hat{\lambda} \in X_\bullet^2(T)/W} \text{Loc}_{((G_{\lambda_1}^+)_{\lambda_2} \times_{G_{\lambda_2}} G_{\lambda_1}^+)}(M) & & \\ & \swarrow & \uparrow & \searrow & \\ \coprod_{\hat{\lambda} \in X_\bullet^2(T)/W} \text{Loc}_{G_{\hat{\lambda}}}(M) & & \coprod_{\hat{\lambda} \in X_\bullet^2(T)/W} \text{Loc}_{G_{\hat{\lambda}}^{+,+}}(M) & & \text{Loc}_G(M) \\ & \swarrow & \downarrow & \searrow & \\ & & \coprod_{\hat{\lambda} \in X_\bullet^2(T)/W} \text{Loc}_{((G_{\lambda_2}^+)_{\lambda_1} \times_{G_{\lambda_1}} G_{\lambda_2}^+)}(M) & & \end{array}$$

as a 2-fold (-1) -shifted Lagrangian correspondence, where the latter structure is obtained by the AKSZ procedure from Proposition 9.5 with $L = G_{\hat{\lambda}}$, $P_1 = (G_{\lambda_1}^+)_{\lambda_2} \times_{G_{\lambda_2}} G_{\lambda_1}^+$, $P_2 = (G_{\lambda_2}^+)_{\lambda_1} \times_{G_{\lambda_1}} G_{\lambda_2}^+$ and $P_1 \cap P_2 = G_{\hat{\lambda}}^{+,+}$.

Proof. The first two statements are shown in [Spr09, Proposition 8.4.5]. The third statement is shown in [CGP15, Lemma C.3.5].

The fourth statement follows from Lemma 9.7: $(G_{\lambda_1}^+)_{\lambda_2} \subset G_{\lambda_2}$ is parabolic with Levi factor $G_{\hat{\lambda}}$ and $G_{\lambda_2}^+ \subset G$ is parabolic with Levi factor G_{λ_2} .

Let us now prove the fifth statement. Since G is affine, it boils down to proving that the intersection of the two subspaces $\mathbb{C}[t_1, t_2^{\pm 1}] \times_{\mathbb{C}[t_2^{\pm 1}]} \mathbb{C}[t_2]$ and $\mathbb{C}[t_1^{\pm 1}, t_2] \times_{\mathbb{C}[t_1^{\pm 1}]} \mathbb{C}[t_1]$ of $\mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}]$ is equal to $\mathbb{C}[t_1, t_2] \subset \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}]$ which is obvious.

Now we prove the sixth statement. Specializing Theorem 5.6 to $X = \text{pt}$ we get that the attractor correspondence for BG is

$$\begin{array}{ccc} & \coprod_{\lambda \in X_\bullet(T)/W} BG_\lambda^+ & \\ \swarrow & & \searrow \\ \coprod_{\lambda \in X_\bullet(T)/W} BG_\lambda & & BG \end{array}$$

Using the 2-shifted symplectic structure on BG determined by the non-degenerate pairing c , there is an induced structure of a 2-shifted Lagrangian correspondence using Corollary 5.18. By uniqueness of a Lagrangian structure on $BG_\lambda \leftarrow BG_\lambda^+ \rightarrow BG$, it coincides with the one defined in [Saf17].

Since $\text{Loc}_G(-)$, $\text{Filt}(-)$ and $\text{Grad}(-)$ are all mapping stacks, we get that the attractor correspondence for $\text{Loc}_G(M)$ is obtained by applying $\text{Map}(M_B, -)$ to this 2-shifted Lagrangian correspondence. We can then commute the coproduct over $X_\bullet(T)/W$ with $\text{Map}(M_B, -)$ since M is connected and its homotopy type is finitely dominated.

The proof of the seventh statement is identical by using uniqueness of the 2-fold 2-shifted Lagrangian correspondence shown in Proposition 9.5. \square

Let $B \subset G$ be a Borel subgroup, $T \subset B$ a maximal torus, Φ^+ the corresponding set of positive roots of G , $\Delta \subset \Phi^+$ the subset of simple roots. Recall that a **standard parabolic subgroup** $P \subset G$ is a parabolic subgroup containing B . By [Spr09, Theorem 8.4.3] we have a bijection of sets

$$(9.3) \quad \{\text{standard parabolic subgroups } P \subset G\} \leftrightarrow \{\text{subsets } I \subset \Delta\}$$

given by assigning to a standard parabolic subgroup P with Levi factor L the subset of simple roots appearing in the root decomposition of the Lie algebra of L .

Lemma 9.10.

(1) Consider a cocharacter $\lambda: \mathbb{G}_m \rightarrow T$. Then the parabolic subgroup $G_\lambda^+ \subset G$ is standard if, and only if, $\lambda(\alpha) \geq 0$ for every $\alpha \in \Delta$. Under the bijection (9.3) the corresponding subset of simple roots is

$$I = \{\alpha \in \Delta \mid \lambda(\alpha) = 0\}.$$

(2) Consider a pair of cocharacters $\hat{\lambda} = (\lambda_1, \lambda_2): \mathbb{G}_m^2 \rightarrow T$ such that $\lambda_1(\alpha)\lambda_2(\alpha) \geq 0$ for every $\alpha \in \Phi^+$. Then the image of the inclusion $\text{Loc}_{G_{\hat{\lambda}}}(M) \subset \text{Grad}^2(\text{Loc}_G(M))$ described in Proposition 9.9(7) lies in the matching locus.

Proof. Let \mathfrak{g} be the Lie algebra of G and let

$$\mathfrak{g} = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi} \mathfrak{g}_\alpha$$

be the root decomposition.

(1) The Lie algebra of G_λ^+ has the root decomposition

$$(\mathfrak{g})_\lambda^+ = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi, \lambda(\alpha) \geq 0} \mathfrak{g}_\alpha.$$

So, G_λ^+ is standard if, and only if, $\lambda(\alpha) \geq 0$ for every $\alpha \in \Phi^+$. Since every positive root is a sum of simple roots, this condition is equivalent to $\lambda(\alpha) \geq 0$ for every $\alpha \in \Delta$. The Lie algebra of G_λ has the root decomposition

$$(\mathfrak{g})_\lambda = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi, \lambda(\alpha) = 0} \mathfrak{g}_\alpha.$$

The subset $I \subset \Delta$ of simple roots corresponding to a standard parabolic subgroup $P \subset G$ is precisely the set of simple roots appearing in the root decomposition of the Lie algebra of the Levi subgroup.

(2) Consider the bigrading of \mathfrak{g} given by $\hat{\lambda}$. By the condition in the statement, we have that the weight (m, l) -part $\mathfrak{g}_{(m, l)}$ of \mathfrak{g} is nonzero only if $ml \geq 0$. Therefore, using the description of the cotangent complex of the mapping stack (see e.g. [NS23, Proposition 1.5]) we obtain that the image of $\text{Loc}_{G_{\hat{\lambda}}}(M) \rightarrow \text{Grad}^2(\text{Loc}_G(M))$ lands in the matching locus. \square

We will now show that the hyperbolic localization for the perverse sheaf $\varphi_{\text{Loc}_G(M)}$ gives rise to a parabolic restriction isomorphism.

Proposition 9.11. *Let $P \subset G$ be a parabolic subgroup with Levi factor L and consider the correspondence*

$$\begin{array}{ccc} & \text{Loc}_P(M) & \\ \swarrow \pi_L & & \searrow \pi_G \\ \text{Loc}_L(M) & & \text{Loc}_G(M) \end{array}$$

There is an isomorphism

$$\zeta_P: \varphi_{\text{Loc}_L(M)} \cong (\pi_L)_* \pi_G^! \varphi_{\text{Loc}_G(M)}$$

invariant under conjugation of P with the following properties:

(1) *For the parabolic subgroup $G \subset G$ we have $\zeta_G = \text{id}$.*

(2) For a pair of parabolic subgroups $P_1 \subset G_1$ and $P_2 \subset G_2$ with Levi factors L_1 and L_2 the diagram

$$\begin{array}{ccc} \varphi_{\text{Loc}_{L_1}(M)} \boxtimes \varphi_{\text{Loc}_{L_2}(M)} & \xrightarrow[\cong]{\zeta_{P_1} \boxtimes \zeta_{P_2}} & (\pi_{L_1})_* \pi'_{G_1}(\varphi_{\text{Loc}_{G_1}(M)}) \boxtimes (\pi_{L_2})_* \pi'_{G_2}(\varphi_{\text{Loc}_{G_2}(M)}) \\ \cong \downarrow \text{TS} & & \cong \downarrow \text{TS} \\ \varphi_{\text{Loc}_{L_1 \times L_2}(M)} & \xrightarrow[\cong]{\zeta_{P_1 \times P_2}} & (\pi_{L_1 \times L_2})_* \pi'_{G_1 \times G_2}(\varphi_{\text{Loc}_{G_1 \times G_2}(M)}) \end{array}$$

(3) For parabolic subgroups P_{12}, P_{23}, P_{13} as in Lemma 9.7 consider the diagram

$$\begin{array}{ccccc} & & \text{Loc}_{P_{13}}(M) & & \\ & \swarrow \pi_{P_{12}} & & \searrow \pi_{P_{23}} & \\ & \text{Loc}_{P_{12}}(M) & & \text{Loc}_{P_{23}}(M) & \\ \swarrow \pi_{G_1} & & \searrow \pi'_{G_2} & \swarrow \pi''_{G_2} & \searrow \pi_{G_3} \\ \text{Loc}_{G_1}(M) & & \text{Loc}_{G_2}(M) & & \text{Loc}_{G_3}(M). \end{array}$$

Then the diagram

$$\begin{array}{ccc} \varphi_{\text{Loc}_{G_1}(M)} & \xrightarrow[\cong]{\zeta_{P_{13}}} & (\pi_{G_1})_* (\pi_{P_{12}})_* \pi'_{P_{23}} \pi'_{G_3} \varphi_{\text{Loc}_{G_3}(M)} \\ \cong \downarrow \zeta_{P_{12}} & & \cong \downarrow \\ (\pi_{G_1})_* (\pi'_{G_2})' \varphi_{\text{Loc}_{G_2}(M)} & \xrightarrow[\cong]{\zeta_{P_{23}}} & (\pi_{G_1})_* (\pi'_{G_2})' (\pi''_{G_2})_* \pi'_{G_3} \varphi_{\text{Loc}_{G_3}(M)} \end{array}$$

commutes.

Proof. By Proposition 9.9(2) we may find a cocharacter $\lambda: \mathbb{G}_m \rightarrow G$ such that $P = G_\lambda^+$ and $L = G_\lambda$. By [NS23, Proposition 3.23] we have $\text{vd}(\text{Loc}_P(M)) = 0$. In particular, we have $I_{\text{Loc}_{G_\lambda}(M)} = 0$ by (6.4).

Let $u: \text{Loc}_L(M) \rightarrow \text{Loc}_G(M)$ be the map induced by the inclusion $G_\lambda \subset G$. Let $\omega_{\text{Loc}_G(M)}$ be the natural (-1) -shifted symplectic structure on $\text{Loc}_G(M)$. The natural structure of a Lagrangian correspondence on $\text{Loc}_L(M) \leftarrow \text{Loc}_P(M) \rightarrow \text{Loc}_G(M)$ induces an equivalence between the (-1) -shifted symplectic structure $u^* \omega_{\text{Loc}_G(M)}$ and $\omega_{\text{Loc}_L(M)}$. Using the orientation $\rho_{G_\lambda^+}$ of the Lagrangian correspondence $\text{Loc}_{G_\lambda}(M) \leftarrow \text{Loc}_{G_\lambda^+}(M) \rightarrow \text{Loc}_G(M)$ from Proposition 9.6 we obtain an isomorphism between the localized orientation $u^* o_G$ and o_L . Therefore, using the isomorphism $\zeta_{\text{Loc}_G(M)}$ from Corollary 7.19 we obtain an isomorphism

$$\zeta_{G_\lambda^+}: \varphi_{\text{Loc}_{G_\lambda}(M)} \cong (\pi_{G_\lambda})_* \pi'_{G_\lambda} \varphi_{\text{Loc}_G(M)}.$$

Let us now show that it satisfies the relevant properties for *some* realizations of the parabolic subgroups as $P = G_\lambda^+$:

- (1) As $\text{Loc}_G(M_1 \amalg M_2) = \text{Loc}_G(M_1) \times \text{Loc}_G(M_2)$, it is enough to prove the claim for M connected with a chosen basepoint. In this case, we have a chart $q: X = \text{Loc}_G^{\text{fr}}(M) \rightarrow \text{Loc}_G(M)$ given by the framed character stack. Suppose the cocharacter $\lambda: \mathbb{G}_m \rightarrow G$ is such that $G_\lambda = G$. Then λ factors through the centre of G . The pair (X, λ) defines a \mathbb{G}_m -equivariant chart of $\text{Loc}_G(M)$ and, since η is surjective, it is enough to prove the statement after the pullback by q . The isomorphism $q^*(\zeta)$ constructed in Theorem 7.16 coincides with the hyperbolic localization isomorphism from Theorem 7.9 for the attractor correspondence $X^\lambda \leftarrow X^{\lambda,+} \rightarrow X$. But since the \mathbb{G}_m -action given by λ is trivial, this isomorphism is the identity by the commutativity of the diagram (7.9).
- (2) The multiplicativity of ζ follows from the multiplicativity property of the isomorphism ζ from Corollary 7.19 as well as the equality $\rho_{P_1} \boxtimes \rho_{P_2} = \rho_{P_1 \times P_2}$ of orientations of the parabolic restriction correspondence shown in Proposition 9.6(4).
- (3) Fix a minimal parabolic subgroup $B \subset P_{13}$ and a maximal torus $T \subset B$ with the corresponding set Δ of simple roots. Then $P_{23}, P_{13} \subset G$ are standard parabolic subgroups. Moreover, $B \cap G_2 \subset G_2$ is a Borel subgroup and $P_{12} \supset B \cap G_2$. Let $I_1 \subset I_2 \subset \Delta$ be the sets of simple roots of $G_1 \subset G_2 \subset G_3$.

Then $I_2 \subset \Delta$ is the subset of simple roots corresponding to the parabolic P_{23} under (9.3), $I_1 \subset \Delta$ is the subset corresponding to P_{13} and $I_1 \subset I_2$ is the subset corresponding to P_{12} .

Consider cocharacters

$$\lambda_{23}: \mathbb{G}_m \rightarrow T, \quad \lambda_{12}: \mathbb{G}_m \rightarrow T$$

satisfying the following conditions:

$$\begin{aligned} \lambda_2(\alpha) = 0 \quad \forall \alpha \in I_2, & \quad \lambda_2(\alpha) > 0 \quad \forall \alpha \in \Delta \setminus I_2 \\ \lambda_1(\alpha) = 0 \quad \forall \alpha \in I_1, & \quad \lambda_1(\alpha) > 0 \quad \forall \alpha \in \Delta \setminus I_1. \end{aligned}$$

Then using Lemma 9.10(1) as well as the fact that (9.3) is a bijection, we get

$$P_{23} = (G_3)_{\lambda_2}^+, \quad P_{12} = (G_2)_{\lambda_1}^+, \quad P_{13} = (G_3)_{\lambda_1}^+.$$

Let $\hat{\lambda} = (\lambda_1, \lambda_2): \mathbb{G}_m^2 \rightarrow T$. Then by Lemma 9.10(2) the image of $\text{Loc}_{G_{\hat{\lambda}}}(M) \rightarrow \text{Grad}^2(\text{Loc}_G(M))$ lands in the matching locus. Using the associativity property of the isomorphism ζ from Corollary 7.19 as well as the fact that the orientation of the parabolic restriction diagram is compatible with compositions shown in Proposition 9.8 we get a commutative diagram

$$\begin{array}{ccc} & \varphi_{\text{Loc}_{G_1}(M)} & \\ \zeta_{(G_2)_{\lambda_1}^+} \swarrow & & \searrow \zeta_{(G_1)_{\lambda_2}^+} \\ (\pi_{G_1})_* (\pi'_{G_2})^! \varphi_{\text{Loc}_{G_2}(M)} & & \varphi_{\text{Loc}_{G_1}(M)} \\ \zeta_{(G_3)_{\lambda_2}^+} \downarrow & & \downarrow \zeta_{(G_3)_{\lambda_1}^+} \\ (\pi_{G_1})_* (\pi'_{G_2})^! (\pi''_{G_2})^* \pi_{G_3}^! \varphi_{\text{Loc}_{G_3}(M)} & \xrightarrow{\cong} & (\pi_{G_1})_* (\pi_{P_{12}})_* \pi_{P_{23}}^! \pi_{G_3}^! \varphi_{\text{Loc}_{G_3}(M)} \end{array}$$

But by part (1) the map $\zeta_{(G_1)_{\lambda_2}^+}$ is the identity, so we get the claim.

Let us finally show that the isomorphism $\zeta_{G_{\hat{\lambda}}^+}$ is independent of the choice of λ and merely depends on the underlying subgroup $P \subset G$. Namely, given two cocharacters $\lambda_1, \lambda_2: \mathbb{G}_m \rightarrow G$ such that $P = G_{\lambda_1}^+ = G_{\lambda_2}^+$, we want to show that the isomorphisms $\zeta_{G_{\lambda_1}^+}$ and $\zeta_{G_{\lambda_2}^+}$ are equal. Since $G_{\lambda_1} = G_{\lambda_2}$ and $\lambda_i(t) \in G_{\lambda_i}$, the cocharacters λ_1 and λ_2 commute, so they define a homomorphism $\hat{\lambda} = (\lambda_1, \lambda_2): \mathbb{G}_m^2 \rightarrow G$. Consider the bigrading of \mathfrak{g} given by $\hat{\lambda}$. Since $G_{\lambda_1}^+ = G_{\lambda_2}^+$, we have that the weight (m, l) -part $\mathfrak{g}_{(m, l)}$ of \mathfrak{g} is nonzero only if $ml \geq 0$. As in the proof of Lemma 9.10(2) we obtain that the image of $\text{Loc}_{G_{\hat{\lambda}}}(M) \rightarrow \text{Grad}^2(\text{Loc}_G(M))$ lands in the matching locus. Now consider the 2-fold correspondence

$$\begin{array}{ccccc} & & \text{Loc}_{(G_{\lambda_1}^+)_{\lambda_2} \times_{G_{\lambda_2}} G_{\lambda_2}^+}(M) & & \\ & \swarrow & \uparrow & \searrow & \\ \text{Loc}_{G_{\hat{\lambda}}}(M) & & \text{Loc}_{G_{\hat{\lambda}}^{+,+}}(M) & & \text{Loc}_G(M) \\ & \swarrow & \downarrow & \searrow & \\ & & \text{Loc}_{(G_{\lambda_2}^+)_{\lambda_1} \times_{G_{\lambda_1}} G_{\lambda_1}^+}(M) & & \end{array}$$

where the maps $G_{\hat{\lambda}} \leftarrow (G_{\lambda_1}^+)_{\lambda_2} \rightarrow G_{\lambda_2}$, $G_{\hat{\lambda}} \leftarrow (G_{\lambda_2}^+)_{\lambda_1} \rightarrow G_{\lambda_1}$ and $((G_{\lambda_1}^+)_{\lambda_2} \times_{G_{\lambda_2}} G_{\lambda_2}^+) \leftarrow G_{\hat{\lambda}}^{+,+} \rightarrow ((G_{\lambda_2}^+)_{\lambda_1} \times_{G_{\lambda_1}} G_{\lambda_1}^+)$ are all equalities as subgroups of G .

Using properties (1) and (3) we get that

$$\zeta_{(G_{\lambda_1}^+)_{\lambda_2} \times_{G_{\lambda_2}} G_{\lambda_2}^+} = \zeta_{G_{\lambda_2}^+}$$

and

$$\zeta_{(G_{\lambda_2}^+)_{\lambda_1} \times_{G_{\lambda_1}} G_{\lambda_1}^+} = \zeta_{G_{\lambda_1}^+}.$$

Using the associativity property of the isomorphism ζ from Corollary 7.19 we have

$$\zeta_{(G_{\lambda_1}^+)_{\lambda_2} \times G_{\lambda_2}} G_{\lambda_2}^+ = \zeta_{(G_{\lambda_2}^+)_{\lambda_1} \times G_{\lambda_1}} G_{\lambda_1}^+.$$

□

Consider the *cohomological DT invariants*

$$H_{G,M} = H^\bullet(\mathrm{Loc}_G(M), \varphi_{\mathrm{Loc}_G(M)}).$$

Using multiplicativity of the orientation o_G we get a map

$$H_{G_1,M} \otimes H_{G_2,M} \longrightarrow H_{G_1 \times G_2, M}.$$

Using the parabolic restriction isomorphism constructed in Proposition 9.11 we obtain the following construction.

Corollary 9.12. *Let $P \subset G$ be a parabolic subgroup with Levi factor L . There is a **parabolic induction map***

$$\mathrm{ind}_P: H_{L,M} \longrightarrow H_{G,M}$$

with the following properties:

(1) For the parabolic subgroup $G \subset G$ the parabolic induction

$$\mathrm{ind}_G: H_{G,M} \longrightarrow H_{G,M}$$

is the identity.

(2) For a pair of parabolic subgroups $P_1 \subset G_1$ and $P_2 \subset G_2$ with Levi factors L_1 and L_2 we have

$$\mathrm{ind}_{P_1} \otimes \mathrm{ind}_{P_2} = \mathrm{ind}_{P_1 \times P_2}: H_{L_1, M} \otimes H_{L_2, M} \longrightarrow H_{G_1 \times G_2, M}.$$

(3) For parabolic subgroups P_{12}, P_{23}, P_{13} as in Lemma 9.7 we have

$$\mathrm{ind}_{P_{23}} \circ \mathrm{ind}_{P_{12}} = \mathrm{ind}_{P_{13}}: H_{G_1, M} \longrightarrow H_{G_3, M}.$$

Proof. Consider the correspondence

$$\begin{array}{ccc} & \mathrm{Loc}_P(M) & \\ \pi_L \swarrow & & \searrow \pi_G \\ \mathrm{Loc}_L(M) & & \mathrm{Loc}_G(M) \end{array}$$

It is shown in the proof of [NS23, Theorem 3.47] that the morphism $\pi_G: \mathrm{Loc}_P(M) \rightarrow \mathrm{Loc}_G(M)$ is representable and proper. In particular, there is a natural transformation $(\pi_G)_* \pi_G^! \rightarrow \mathrm{id}$. We define the parabolic induction map ind_P as the composite

$$\begin{aligned} H^\bullet(\mathrm{Loc}_L(M), \varphi_{\mathrm{Loc}_L(M)}) &\xrightarrow{\zeta_P} H^\bullet(\mathrm{Loc}_L(M), (\pi_L)_* \pi_G^! \varphi_{\mathrm{Loc}_G(M)}) \\ &\cong H^\bullet(\mathrm{Loc}_G(M), (\pi_G)_* \pi_G^! \varphi_{\mathrm{Loc}_G(M)}) \\ &\rightarrow H^\bullet(\mathrm{Loc}_G(M), \varphi_{\mathrm{Loc}_G(M)}). \end{aligned}$$

The three properties of ind_P follow from the corresponding properties of the isomorphism ζ_P . □

Remark 9.13. The parabolic induction map ind_P depends on the choice of a spin structure on M as follows. The volume form $\overline{\mathrm{vol}}_G$ that goes into the definition of the orientation depends on the canonical Euler structure (and hence the spin structure) as shown in [NS23, Proposition 3.23]. In particular, the orientation o_G for two different spin structures are canonically isomorphic since G is unimodular; however, the orientation ρ_P does depend on the spin structure. The difference of two spin structures s_1, s_2 defines an element of $H^1(M; \mathbb{Z}/2\mathbb{Z})$. Denote its image under the composite

$$H^1(M; \mathbb{Z}/2\mathbb{Z}) \longrightarrow H^2(M; \mathbb{Z}) \cong H_1(M; \mathbb{Z})$$

of the Bockstein homomorphism and Poincaré duality by h . It is a 2-torsion element of $H_1(M; \mathbb{Z})$ which represents the difference of the corresponding canonical Euler structures. The modular character $P \rightarrow \mathbb{G}_m$ of P factors through $\Delta_P: L \rightarrow \mathbb{G}_m$, and hence it defines a rank 1 local system on $\mathrm{Loc}_L(M)$. Denote by

$\langle \Delta_P, h \rangle$ the monodromy of this local system along h which defines a μ_2 -valued function on $\text{Loc}_L(M)$. Then the two parabolic induction maps differ by a sign:

$$\text{ind}_P^{s_2} = \text{ind}_P^{s_1} \cdot \langle \Delta_P, h \rangle.$$

We now collect all the results we have proven to define a cohomological Hall algebra for GL_n and cohomological Hall modules for SO_n and Sp_{2n} . For this, let us first describe a class of parabolic subgroups of classical groups (we denote by $\{e_i\}$ the standard basis of \mathbb{C}^n):

- Let $P_{n,m}^{\text{GL}} \subset \text{GL}_{n+m}$ for $n, m \geq 0$ be the parabolic subgroup preserving the flag $\text{span}\{e_1, \dots, e_m\} \subset \text{span}\{e_1, \dots, e_{n+m}\}$. Its Levi factor is $\text{GL}_n \times \text{GL}_m$.
- Let $P_{n_1, n_2, n_3}^{\text{GL}} \subset \text{GL}_{n_1+n_2+n_3}$ for $n_1, n_2, n_3 \geq 0$ be the parabolic subgroup preserving the flag $\text{span}\{e_1, \dots, e_{n_1}\} \subset \text{span}\{e_1, \dots, e_{n_1+n_2}\} \subset \text{span}\{e_1, \dots, e_{n_1+n_2+n_3}\}$.

Its Levi factor is $\text{GL}_{n_1} \times \text{GL}_{n_2} \times \text{GL}_{n_3}$.

- Let SO_n be the group of isometries of the symmetric bilinear form on \mathbb{C}^n with $(e_i, e_j) = \delta_{i, n-j}$ of determinant 1. Let $P_{n,m}^{\text{SO}} \subset \text{SO}_{n+2m}$ for $n, m \geq 0$ be the parabolic subgroup preserving the isotropic flag $\mathbb{C}^m = \text{span}\{e_1, \dots, e_m\} \subset \text{span}\{e_1, \dots, e_{n+2m}\}$. Its Levi factor is $\text{SO}_n \times \text{GL}_m$.
- Let $P_{n_1, n_2, n_3}^{\text{SO}} \subset \text{SO}_{n_1+2n_2+2n_3}$ for $n_1, n_2, n_3 \geq 0$ be the parabolic subgroup preserving the isotropic flag $\text{span}\{e_1, \dots, e_{n_1}\} \subset \text{span}\{e_1, \dots, e_{n_1+n_2}\} \subset \text{span}\{e_1, \dots, e_{n_1+2n_2+2n_3}\}$.

Its Levi factor is $\text{SO}_{n_1} \times \text{GL}_{n_2} \times \text{GL}_{n_3}$.

- Let Sp_{2n} be the group of invertible transformations preserving the symplectic form

$$(e_i, e_j) = \delta_{i, 2n-j}, \quad i \leq n, \quad (e_i, e_j) = -\delta_{i, 2n-j}, \quad i > n.$$

Let $P_{2n, m}^{\text{Sp}} \subset \text{Sp}_{2n+2m}$ be the parabolic subgroup preserving the isotropic flag $\text{span}\{e_1, \dots, e_m\} \subset \text{span}\{e_1, \dots, e_{2n+2m}\}$. Its Levi factor is $\text{Sp}_{2n} \times \text{GL}_m$.

- Let $P_{2n_1, 2n_2, 2n_3}^{\text{Sp}} \subset \text{Sp}_{2n_1+2n_2+2n_3}$ for $n_1, n_2, n_3 \geq 0$ be the parabolic subgroup preserving the isotropic flag $\text{span}\{e_1, \dots, e_{n_1}\} \subset \text{span}\{e_1, \dots, e_{n_1+n_2}\} \subset \text{span}\{e_1, \dots, e_{2n_1+2n_2+2n_3}\}$.

Its Levi factor is $\text{Sp}_{2n_1} \times \text{GL}_{n_2} \times \text{GL}_{n_3}$.

We have

$$(9.4) \quad P_{n_1, n_2}^{\text{GL}} \times_{\text{GL}_{n_1+n_2}} P_{n_1+n_2, n_3}^{\text{GL}} = P_{n_1, n_2, n_3}^{\text{GL}} = P_{n_2, n_3}^{\text{GL}} \times_{\text{GL}_{n_2+n_3}} P_{n_1, n_2+n_3}^{\text{GL}},$$

$$(9.5) \quad P_{n_1, n_2}^{\text{SO}} \times_{\text{SO}_{n_1+2n_2}} P_{n_1+2n_2, n_3}^{\text{SO}} = P_{n_1, n_2, n_3}^{\text{SO}} = P_{n_2, n_3}^{\text{GL}} \times_{\text{GL}_{n_2+n_3}} P_{n_1, n_2+n_3}^{\text{SO}}$$

and

$$(9.6) \quad P_{2n_1, 2n_2}^{\text{Sp}} \times_{\text{Sp}_{2n_1+2n_2}} P_{2n_1+2n_2, 2n_3}^{\text{Sp}} = P_{2n_1, 2n_2, 2n_3}^{\text{Sp}} = P_{n_2, n_3}^{\text{GL}} \times_{\text{GL}_{n_2+n_3}} P_{2n_1, 2n_2+n_3}^{\text{Sp}}.$$

Using the orientation ρ_G and ρ_P we obtain an orientation data for the category of local systems on M . Recall that by Theorem 9.3 we have

$$\mathfrak{M}_{\text{LocSys}(M)^\omega} \cong \text{Map}(M_B, \text{Perf}).$$

Let

$$\mathfrak{M} = \text{Map}(M_B, \text{Vect})$$

be the substack of local systems of vector spaces.

Proposition 9.14. *Let M be a connected closed spin 3-manifold. Let $\mathcal{C} = \text{LocSys}(M)^\omega$ and $\mathfrak{M} \subset \mathfrak{M}_{\mathcal{C}}$ the substack of local systems of vector spaces.*

- (1) *There is an isomorphism of (-1) -shifted symplectic stacks $\mathfrak{M} \cong \coprod_n \text{Loc}_{\text{GL}_n}(M)$.*
- (2) *\mathfrak{M} has affine diagonal and is Θ -reductive.*
- (3) *\mathfrak{M} is a moduli-like stack.*
- (4) *There is a strong orientation data on \mathfrak{M} .*

Proof.

(1) Write

$$\text{Vect} \simeq \text{colim}_n B \text{GL}_n.$$

Since M is connected and finitely dominated, by Lemma 9.1 we have

$$\text{Map}(M_{\mathbb{B}}, \text{Vect}) \cong \prod_n \text{Map}(M_{\mathbb{B}}, B \text{GL}_n).$$

(2) Choosing a basepoint, we may identify

$$\text{Loc}_{\text{GL}_n}(M) \cong [\text{Loc}_{\text{GL}_n}^{\text{fr}}(M) / \text{GL}_n].$$

This is a quotient of an affine scheme by an affine algebraic group, so the diagonal is affine. Moreover, by [Hal14, Example 5.1.8] it is Θ -reductive.

(3) The claim follows from Proposition 8.29.

(4) By Proposition 8.26 the moduli stack $\mathfrak{M}^{2\text{-flt}}$ parametrizes fibre sequences $\mathcal{L}_1 \rightarrow \mathcal{L}_2 \rightarrow \mathcal{L}_3$ of local systems of perfect complexes on M , such that \mathcal{L}_1 and \mathcal{L}_3 are local systems of vector spaces. Therefore, \mathcal{L}_2 is a local system of vector spaces and this is an exact sequence. Thus, the correspondence $\mathfrak{M}^2 \leftarrow \mathfrak{M}^{2\text{-flt}} \rightarrow \mathfrak{M}$ is equivalent to

$$\begin{array}{ccc} & \prod_{a,b} \text{Loc}_{P_{a,b}}(M) & \\ & \swarrow \quad \searrow & \\ \prod_{a,b} \text{Loc}_{\text{GL}_a}(M) \times \text{Loc}_{\text{GL}_b}(M) & & \prod_n \text{Loc}_{\text{GL}_n}(M) \end{array}$$

Moreover, by Proposition 9.9(6) this equivalence is compatible with (-1) -shifted Lagrangian structures. Therefore, the orientation on \mathfrak{M} follows from the orientation o_{GL_n} of $\text{Loc}_{\text{GL}_n}(M)$ and orientation $\rho_{P_{a,b}}$ of $\text{Loc}_{\text{GL}_a}(M) \times \text{Loc}_{\text{GL}_b}(M) \leftarrow \text{Loc}_{P_{a,b}}(M) \rightarrow \text{Loc}_{\text{GL}_{a+b}}(M)$ constructed in Proposition 9.6.

Using (9.4) we have

$$(P_{n_1, n_2}^{\text{GL}} \times \text{GL}_{n_3}) \times_{\text{GL}_{n_1+n_2} \times \text{GL}_{n_3}} P_{n_1+n_2, n_3}^{\text{GL}} = P_{n_1, n_2, n_3}^{\text{GL}} = (\text{GL}_{n_1} \times P_{n_2, n_3}^{\text{GL}}) \times_{\text{GL}_{n_1} \times \text{GL}_{n_2+n_3}} P_{n_1, n_2+n_3}^{\text{GL}}.$$

Thus, the associativity of the orientation data follows from Proposition 9.8. The unitality of the orientation data follows from the fact that $\rho_{P_{n,0}}$ and $\rho_{P_{0,n}}$ are the identity orientations for the correspondence $\text{Loc}_{\text{GL}_n}(M) \leftarrow \text{Loc}_{\text{GL}_n}(M) \rightarrow \text{Loc}_{\text{GL}_n}(M)$. □

Theorem 9.15. *Let M be a closed spin 3-manifold. Consider the graded vector spaces*

$$\begin{aligned} \mathcal{H}^{\text{GL}} &= \bigoplus_{n \geq 0} H_{\text{GL}_n, M} \\ \mathcal{H}^{\text{SO}} &= \bigoplus_{n \geq 0} H_{\text{SO}_n, M} \\ \mathcal{H}^{\text{Sp}} &= \bigoplus_{n \geq 0} H_{\text{Sp}_{2n}, M} \end{aligned}$$

Then the maps

$$\begin{aligned} H_{\text{GL}_n, M} \otimes H_{\text{GL}_m, M} &\rightarrow H_{\text{GL}_n \times \text{GL}_m, M} \xrightarrow{\text{ind}_{P_{n,m}^{\text{GL}}}} H_{\text{GL}_{n+m}, M} \\ H_{\text{SO}_n, M} \otimes H_{\text{GL}_m, M} &\rightarrow H_{\text{SO}_n \times \text{GL}_m, M} \xrightarrow{\text{ind}_{P_{n,m}^{\text{SO}}}} H_{\text{SO}_{n+2m}, M} \\ H_{\text{Sp}_{2n}, M} \otimes H_{\text{GL}_m, M} &\rightarrow H_{\text{Sp}_{2n} \times \text{GL}_m, M} \xrightarrow{\text{ind}_{P_{2n,m}^{\text{Sp}}}} H_{\text{Sp}_{2n+2m}, M} \end{aligned}$$

as well as the obvious isomorphism $e: \mathbb{C} \cong H_{\text{GL}_0, M}$ define a graded algebra structure on \mathcal{H}^{GL} and graded \mathcal{H}^{GL} -module structures on \mathcal{H}^{SO} and \mathcal{H}^{Sp} .

Proof. The fact that \mathcal{H}^{GL} is a unital associative algebra is shown in Corollary 8.8. The unitality and associativity of the \mathcal{H}^{GL} -actions on \mathcal{H}^{SO} and \mathcal{H}^{Sp} are checked analogously, so we will only prove the claim for \mathcal{H}^{SO} .

We begin by establishing unitality. The multiplication map

$$H_{\text{SO}_n, M} \otimes H_{\text{GL}_0, M} \longrightarrow H_{\text{SO}_n, M}$$

consists of the following two composites:

$$H_{\text{SO}_n, M} \xrightarrow{\text{id} \otimes e} H_{\text{SO}_n, M} \otimes H_{\text{GL}_0, M} \rightarrow H_{\text{SO}_n \times \text{GL}_0, M} \cong H_{\text{SO}_n, M}$$

and

$$H_{\text{SO}_n, M} \xrightarrow{\text{ind}_{P_{n,0}^{\text{SO}}}^{\text{SO}_n}} H_{\text{SO}_n, M}.$$

The first composite is clearly the identity. The second composite involves parabolic induction along the parabolic subgroup $P_{n,0}^{\text{SO}} = \text{SO}_n \subset \text{SO}_n$ and is hence also the identity.

Next, let us establish associativity. Namely, we have to show that the two composites

$$H_{\text{SO}_{n_1}, M} \otimes H_{\text{GL}_{n_2}, M} \otimes H_{\text{GL}_{n_3}, M} \xrightarrow{\text{ind}_{P_{n_1, n_2}^{\text{SO}}}^{\text{SO}_{n_1+2n_2}} \otimes \text{id}} H_{\text{SO}_{n_1+2n_2}, M} \otimes H_{\text{GL}_{n_3}, M} \xrightarrow{\text{ind}_{P_{n_1+2n_2, n_3}^{\text{SO}}}^{\text{SO}_{n_1+2n_2+2n_3}}} H_{\text{SO}_{n_1+2n_2+2n_3}, M}$$

and

$$H_{\text{SO}_{n_1}, M} \otimes H_{\text{GL}_{n_2}, M} \otimes H_{\text{GL}_{n_3}, M} \xrightarrow{\text{id} \otimes \text{ind}_{P_{n_2, n_3}^{\text{GL}}}^{\text{GL}_{n_2+n_3}}} H_{\text{SO}_{n_1}, M} \otimes H_{\text{GL}_{n_2+n_3}, M} \xrightarrow{\text{ind}_{P_{n_1, n_2+n_3}^{\text{SO}}}^{\text{SO}_{n_1+2n_2+2n_3}}} H_{\text{SO}_{n_1+2n_2+2n_3}, M}$$

are equal. Using the multiplicativity and unitality properties of the parabolic induction maps (Corollary 9.12) we are reduced to showing that the two composites

$$(9.7) \quad H_{\text{SO}_{n_1} \times \text{GL}_{n_2} \times \text{GL}_{n_3}} \xrightarrow{\text{ind}_{P_{n_1, n_2}^{\text{SO}} \times \text{GL}_{n_3}}^{\text{SO}_{n_1+2n_2} \times \text{GL}_{n_3}}} H_{\text{SO}_{n_1+2n_2} \times \text{GL}_{n_3}} \xrightarrow{\text{ind}_{P_{n_1+2n_2, n_3}^{\text{SO}}}^{\text{SO}_{n_1+2n_2+2n_3}}} H_{\text{SO}_{n_1+2n_2+2n_3}, M}$$

and

$$(9.8) \quad H_{\text{SO}_{n_1} \times \text{GL}_{n_2} \times \text{GL}_{n_3}} \xrightarrow{\text{ind}_{P_{\text{SO}_{n_1} \times \text{GL}_{n_2}, n_3}^{\text{GL}}}^{\text{SO}_{n_1} \times \text{GL}_{n_2+n_3}}} H_{\text{SO}_{n_1} \times \text{GL}_{n_2+n_3}} \xrightarrow{\text{ind}_{P_{n_1, n_2+n_3}^{\text{SO}}}^{\text{SO}_{n_1+2n_2+2n_3}}} H_{\text{SO}_{n_1+2n_2+2n_3}, M}$$

are equal.

Using (9.5) we have

$$(P_{n_1, n_2}^{\text{SO}} \times \text{GL}_{n_3}) \times_{\text{SO}_{n_1+2n_2} \times \text{GL}_{n_3}} P_{n_1+2n_2, n_3}^{\text{SO}} = P_{n_1, n_2, n_3}^{\text{SO}} = (\text{SO}_{n_1} \times P_{n_2, n_3}^{\text{GL}}) \times_{\text{SO}_{n_1} \times \text{GL}_{n_2+n_3}} P_{n_1, n_2+n_3}^{\text{SO}}.$$

Therefore, by the associativity property of parabolic induction shown in Corollary 9.12(3) the two composites (9.7) and (9.8) are both equal to $\text{ind}_{P_{n_1, n_2, n_3}^{\text{SO}}}^{\text{SO}_{n_1+2n_2+2n_3}}$. \square

APPENDIX A. BRADEN'S THEOREM FOR STACKS

The aim of this appendix is to prove a generalization of Braden's theorem on hyperbolic localization (= Theorem 7.7) to the attractor correspondences of Artin stacks. Though we have not used this result in our paper, we record it here since this can be proved analogously to Theorem 7.16 and is interesting in its own right. A similar result in the context of D-modules can be found in [DG16, Theorem 4.3.4].

Let \mathfrak{X} be a quasi-separated Artin stack with affine stabilizers. We set $\text{Filt}^-(\mathfrak{X}) := [\mathbb{A}^{1,-}/\mathbb{G}_m]$ where $\mathbb{A}^{1,-}$ denotes the affine line equipped with \mathbb{G}_m -action of weight -1 . We let $\text{gr}^- : \text{Filt}^-(\mathfrak{X}) \rightarrow \text{Grad}(\mathfrak{X})$ and $\text{ev}^- : \text{Filt}^-(\mathfrak{X}) \rightarrow \mathfrak{X}$ denote the natural maps. The following is the main result in this appendix, which generalizes Braden's theorem (= Theorem 7.7):

Proposition A.1. *There exists a natural isomorphism of functors between constructible derived categories:*

$$\text{gr}_* \text{ev}^! \cong \text{gr}_!^- \text{ev}^{-,*}.$$

We need the following lemma, which is immediate from the description of the cotangent complexes on both sides⁹:

Lemma A.2. *The natural map $\text{Grad}(\mathfrak{X}) \rightarrow \text{Filt}(\mathfrak{X}) \times_{\mathfrak{X}} \text{Filt}^-(\mathfrak{X})$ is an étale morphism.*

⁹One can more generally show that the map is an open immersion.

Proof of Proposition A.1. Let $\sigma: \text{Grad}(\mathfrak{X}) \rightarrow \text{Filt}(\mathfrak{X})$ and $\sigma^-: \text{Grad}(\mathfrak{X}) \rightarrow \text{Filt}^-(\mathfrak{X})$ be the canonical sections of gr and gr^- , respectively. Then by using Proposition 7.2, there exists a natural isomorphism

$$\text{gr}_* \cong \sigma^*, \quad \text{gr}_!^- \cong \sigma^!.$$

Therefore it is enough to construct an equivalence between $\sigma^* \text{ev}^!$ and $\sigma^{-,!} \text{ev}^{-,*}$. Consider the following diagram:

$$\begin{array}{ccc} \text{Grad}(\mathfrak{X}) & \xrightarrow{\sigma} & \text{Filt}(\mathfrak{X}) \\ \downarrow \sigma^- & & \downarrow \text{ev} \\ \text{Filt}^-(\mathfrak{X}) & \xrightarrow{\text{ev}^-} & \mathfrak{X}. \end{array}$$

By Lemma A.2, this diagram is pullable (see §7.1.3). Therefore, we can define the base change transform (7.4)

$$\eta: \sigma^* \text{ev}^! \rightarrow \sigma^{-,!} \text{ev}^{-,*}.$$

We will show that this map is invertible. To see this, take a \mathbb{G}_m -equivariant chart (X, μ, \bar{q}) of \mathfrak{X} such that $q_\mu: X^\mu \rightarrow \text{Grad}(\mathfrak{X})$ is surjective using Proposition 5.25. It is enough to show that $q_\mu^* \eta$ is invertible. Consider the following diagram:

$$\begin{array}{ccccc} X^\mu & \xrightarrow{q_\mu} & \text{Grad}(\mathfrak{X}) & \xrightarrow{\sigma} & \text{Filt}(\mathfrak{X}) \\ \downarrow \sigma_\mu^- & & \downarrow \sigma^- & & \downarrow \text{ev} \\ X^{\mu,-} & \xrightarrow{q_\mu^-} & \text{Filt}^-(\mathfrak{X}) & \xrightarrow{\text{ev}^-} & \mathfrak{X}. \end{array}$$

Since q_μ and q_{μ^-} are smooth, the left square is pullable and the base change map $q_\mu^* \sigma^{-,!} [2 \dim q_\mu] \rightarrow \sigma_\mu^{-,!} q_{\mu^-}^{-,*} [2 \dim q_\mu^-]$ is invertible. Using the associativity of the base change map (7.4) for pullable squares [FYZ23, Proposition 3.5.4], it is enough to prove the invertibility of the base change map (7.4) for the outer squares. Now consider the following diagram:

$$\begin{array}{ccccc} X^\mu & \xrightarrow{\sigma_\mu} & X^{\mu,+} & \xrightarrow{q_\mu^+} & \text{Filt}(\mathfrak{X}) \\ \downarrow \sigma_\mu^- & & \downarrow \text{ev}_\mu & & \downarrow \text{ev} \\ X^{\mu,-} & \xrightarrow{\text{ev}_\mu^-} & X & \xrightarrow{q} & \mathfrak{X}. \end{array}$$

These diagrams are obviously pullable. The base change map (7.4) for the left square is invertible by [DG14, Theorem 3.1.6] and the base change map (7.4) for the right square is invertible by the smoothness of q and q_μ . Therefore, we conclude the invertibility of the base change map (7.4) for the outer square. \square

Remark A.3. Though we have proved it for analytic sheaves on complex Artin stacks, the same method works for any sheaf theory context for which Braden’s theorem on hyperbolic localization for \mathbb{G}_m -action on algebraic spaces (= Theorem 7.7) holds. For example, one can prove the same result for ℓ -adic sheaves on Artin stacks over any field and also for mixed Hodge modules using the six-functor formalism [Tub24].

REFERENCES

- [AB17] L. Amorim and O. Ben-Bassat. “Perversely categorified Lagrangian correspondences”. *Adv. Theor. Math. Phys.* 21.2 (2017), pp. 289–381. arXiv: 1601.01536 [math.SG].
- [Ach21] P. N. Achar. *Perverse sheaves and applications to representation theory*. Vol. 258. American Mathematical Soc., 2021.
- [AHR20] J. Alper, J. Hall, and D. Rydh. “A Luna étale slice theorem for algebraic stacks”. *Annals of Mathematics* 191.3 (2020), pp. 675–738. arXiv: 1504.06467 [math.AG].
- [Alp+24] J. Alper, J. Hall, D. Halpern-Leistner, and D. Rydh. “Artin algebraization for pairs with applications to the local structure of stacks and Ferrand pushouts”. *Forum of Mathematics, Sigma*. Vol. 12. Cambridge University Press, 2024, e20. arXiv: 2205.08623 [math.AG].
- [Alp13] J. Alper. “Good moduli spaces for Artin stacks”. *Annales de l’Institut Fourier*. Vol. 63. 6. 2013, pp. 2349–2402. arXiv: 0804.2242 [math.AG].

- [AM20] M. Abouzaid and C. Manolescu. “A sheaf-theoretic model for $SL(2, \mathbb{C})$ Floer homology”. *J. Eur. Math. Soc. (JEMS)* 22.11 (2020), pp. 3641–3695. arXiv: [1708.00289 \[math.GT\]](#).
- [Ang] Angelo. *Examples of injective morphisms which are not universally injective*. MathOverflow. URL: <https://mathoverflow.net/q/114409>.
- [Aok23] K. Aoki. “Tensor triangular geometry of filtered objects and sheaves”. *Math. Z.* 303.3 (2023), Paper No. 62, 27. arXiv: [2001.00319 \[math.AT\]](#).
- [BBJ19] C. Brav, V. Bussi, and D. Joyce. “A Darboux theorem for derived schemes with shifted symplectic structure”. *J. Amer. Math. Soc.* 32.2 (2019), pp. 399–443. arXiv: [1305.6302 \[math.AG\]](#).
- [BCS23] T. Bozec, D. Calaque, and S. Scherotzke. “Calabi-Yau structures for multiplicative preprojective algebras”. *J. Noncommut. Geom.* 17.3 (2023), pp. 783–810. arXiv: [2102.12336 \[math.RT\]](#).
- [BCS24] T. Bozec, D. Calaque, and S. Scherotzke. “Relative critical loci and quiver moduli”. *Ann. Sci. École Norm. Sup. (4)* 57.2 (2024), pp. 553–614. arXiv: [2006.01069 \[math.RT\]](#).
- [BD19] C. Brav and T. Dyckerhoff. “Relative Calabi-Yau structures”. *Compos. Math.* 155.2 (2019), pp. 372–412. arXiv: [1606.00619 \[math.AG\]](#).
- [BD21] C. Brav and T. Dyckerhoff. “Relative Calabi-Yau structures II: shifted Lagrangians in the moduli of objects”. *Selecta Math. (N.S.)* 27.4 (2021), Paper No. 63, 45. arXiv: [1812.11913 \[math.AG\]](#).
- [Ben+15] O. Ben-Bassat, C. Brav, V. Bussi, and D. Joyce. “A ‘Darboux theorem’ for shifted symplectic structures on derived Artin stacks, with applications”. *Geom. Topol.* 19.3 (2015), pp. 1287–1359. arXiv: [1312.0090 \[math.AG\]](#).
- [BFN10] D. Ben-Zvi, J. Francis, and D. Nadler. “Integral transforms and Drinfeld centers in derived algebraic geometry”. *J. Amer. Math. Soc.* 23.4 (2010), pp. 909–966. arXiv: [0805.0157 \[math.AG\]](#).
- [BGT13] A. J. Blumberg, D. Gepner, and G. Tabuada. “A universal characterization of higher algebraic K-theory”. *Geometry & Topology* 17.2 (2013), pp. 733–838. arXiv: [1001.2282 \[math.KT\]](#).
- [BJ17] D. Borisov and D. Joyce. “Virtual fundamental classes for moduli spaces of sheaves on Calabi-Yau four-folds”. *Geometry & Topology* 21.6 (2017), pp. 3231–3311. arXiv: [1504.00690 \[math.AG\]](#).
- [Bra+15] C. Brav, V. Bussi, D. Dupont, D. Joyce, and B. Szendrői. “Symmetries and stabilization for sheaves of vanishing cycles”. *J. Singul.* 11 (2015). With an appendix by Jörg Schürmann, pp. 85–151. arXiv: [1211.3259 \[math.AG\]](#).
- [Bra03] T. Braden. “Hyperbolic localization of intersection cohomology”. *Transformation groups* 8 (2003), pp. 209–216. arXiv: [math/0202251](#).
- [Bu24] C. Bu. “A motivic integral identity for (-1) -shifted symplectic stacks”. preprint. 2024. arXiv: [2405.10092 \[math.AG\]](#).
- [Cal15] D. Calaque. “Lagrangian structures on mapping stacks and semi-classical TFTs”. *Stacks and categories in geometry, topology, and algebra*. Vol. 643. Contemp. Math. Amer. Math. Soc., Providence, RI, 2015, pp. 1–23. arXiv: [1306.3235 \[math.AG\]](#).
- [Cal19] D. Calaque. “Shifted cotangent stacks are shifted symplectic”. *Ann. Fac. Sci. Toulouse Math. (6)* 28.1 (2019), pp. 67–90. arXiv: [1612.08101 \[math.AG\]](#).
- [CGP15] B. Conrad, O. Gabber, and G. Prasad. *Pseudo-reductive groups*. Second. Vol. 26. New Mathematical Monographs. Cambridge University Press, Cambridge, 2015, pp. xxiv+665.
- [CHS21] D. Calaque, R. Haugseng, and C. Scheimbauer. “The AKSZ construction in derived algebraic geometry as an extended topological field theory”. preprint. 2021. arXiv: [2108.02473 \[math.CT\]](#).
- [Coh] L. Cohn. “Differential graded categories are k -linear stable infinity categories”. preprint. arXiv: [1308.2587 \[math.AT\]](#).
- [Con00] B. Conrad. *Grothendieck duality and base change*. Vol. 1750. Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2000, pp. vi+296.
- [Dav17] B. Davison. “The critical CoHA of a quiver with potential”. *Q. J. Math.* 68.2 (2017), pp. 635–703. arXiv: [1311.7172 \[math.AG\]](#).
- [Des22] P. Descombes. “Hyperbolic localization of the Donaldson-Thomas sheaf”. preprint. 2022. arXiv: [2201.12215 \[math.AG\]](#).
- [DG14] V. Drinfeld and D. Gaitsgory. “On a theorem of Braden”. *Transformation groups* 19.2 (2014), pp. 313–358. arXiv: [1308.3786 \[math.AG\]](#).
- [DG15] V. Drinfeld and D. Gaitsgory. “Compact generation of the category of D-modules on the stack of G -bundles on a curve”. *Camb. J. Math.* 3.1-2 (2015), pp. 19–125. arXiv: [1112.2402 \[math.AG\]](#).
- [DG16] V. Drinfeld and D. Gaitsgory. “Geometric constant term functor(s)”. *Selecta Mathematica* 22 (2016), pp. 1881–1951. arXiv: [1311.2071 \[math.AG\]](#).
- [DJK21] F. Déglise, F. Jin, and A. A. Khan. “Fundamental classes in motivic homotopy theory”. *J. Eur. Math. Soc. (JEMS)* 23.12 (2021), pp. 3935–3993. arXiv: [1805.05920 \[math.AG\]](#).
- [DM20] B. Davison and S. Meinhardt. “Cohomological Donaldson-Thomas theory of a quiver with potential and quantum enveloping algebras”. *Invent. Math.* 221.3 (2020), pp. 777–871. arXiv: [1601.02479 \[math.RT\]](#).
- [Elm+20] E. Elmanto, M. Hoyois, A. A. Khan, V. Sosnilo, and M. Yakerson. “Modules over algebraic cobordism”. *Forum Math. Pi* 8 (2020), e14, 44. arXiv: [1908.02162 \[math.AG\]](#).
- [Fon18] E. Fontes. “Weight structures and the algebraic K -theory of stable ∞ -categories” (2018). arXiv: [1812.09751 \[math.KT\]](#).
- [FT99] M. Farber and V. Turaev. “Absolute torsion”. *Tel Aviv Topology Conference: Rothenberg Festschrift (1998)*. Vol. 231. Contemp. Math. Amer. Math. Soc., Providence, RI, 1999, pp. 73–85. arXiv: [math/9810029](#).
- [FYZ23] T. Feng, Z. Yun, and W. Zhang. “Modularity of higher theta series I: cohomology of the generic fiber”. preprint. 2023. arXiv: [2308.10979 \[math.NT\]](#).

- [GHN17] D. Gepner, R. Haugseng, and T. Nikolaus. “Lax colimits and free fibrations in ∞ -categories”. *Doc. Math.* 22 (2017), pp. 1225–1266. arXiv: [1501.02161 \[math.CT\]](#).
- [Gin06] V. Ginzburg. “Calabi-Yau algebras”. preprint. 2006. arXiv: [math/0612139](#).
- [GR17a] D. Gaitsgory and N. Rozenblyum. *A study in derived algebraic geometry. Vol. I. Correspondences and duality*. Vol. 221. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2017, xl+533pp.
- [GR17b] D. Gaitsgory and N. Rozenblyum. *A study in derived algebraic geometry. Vol. II. Deformations, Lie theory and formal geometry*. Vol. 221. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2017, pp. xxxv+436.
- [GS23] S. Gunningham and P. Safronov. “Deformation quantization and perverse sheaves”. preprint. 2023. arXiv: [2312.07595 \[math.AG\]](#).
- [Hal14] D. Halpern-Leistner. “On the structure of instability in moduli theory”. preprint. 2014. arXiv: [1411.0627 \[math.AG\]](#).
- [Hel20] A. Heleodoro. “Determinant map for the prestack of Tate objects”. *Selecta Math. (N.S.)* 26.5 (2020), Paper No. 76, 57. arXiv: [1907.00384 \[math.AG\]](#).
- [HM98] J. A. Harvey and G. Moore. “On the algebras of BPS states”. *Comm. Math. Phys.* 197.3 (1998), pp. 489–519. arXiv: [hep-th/9609017](#).
- [HP18] Z. Hua and A. Polishchuk. “Shifted Poisson structures and moduli spaces of complexes”. *Adv. Math.* 338 (2018), pp. 991–1037. arXiv: [1706.09965 \[math.AG\]](#).
- [HP23a] D. Halpern-Leistner and A. Preygel. “Mapping stacks and categorical notions of properness”. *Compos. Math.* 159.3 (2023), pp. 530–589. arXiv: [1402.3204 \[math.AG\]](#).
- [HP23b] Z. Hua and A. Polishchuk. “Bosonization of Feigin-Odesskii Poisson varieties”. preprint. 2023. arXiv: [2306.14719 \[math.AG\]](#).
- [HS21] F. Hebestreit and W. Steimle. “Stable moduli spaces of hermitian forms”. preprint. 2021. arXiv: [2103.13911 \[math.KT\]](#).
- [Joy15] D. Joyce. “A classical model for derived critical loci”. *Journal of Differential Geometry* 101.2 (2015), pp. 289–367. arXiv: [1304.4508 \[math.AG\]](#).
- [Joy18] D. Joyce. “Ringel-Hall style vertex algebra and Lie algebra structures on the homology of moduli spaces”. preprint. 2018. URL: <https://people.maths.ox.ac.uk/joyce/hall1.pdf>.
- [JS19] D. Joyce and P. Safronov. “A Lagrangian Neighbourhood Theorem for shifted symplectic derived schemes”. *Ann. Fac. Sci. Toulouse Math. (6)* 28.5 (2019), pp. 831–908. arXiv: [1506.04024 \[math.AG\]](#).
- [JU21] D. Joyce and M. Upmeyer. “Orientation data for moduli spaces of coherent sheaves over Calabi-Yau 3-folds”. *Adv. Math.* 381 (2021), p. 107627. arXiv: [2001.00113 \[math.AG\]](#).
- [Kel11] B. Keller. “Deformed Calabi-Yau completions”. *J. Reine Angew. Math.* 654 (2011). With an appendix by Michel Van den Bergh, pp. 125–180. arXiv: [0908.3499 \[math.RT\]](#).
- [Kel18] B. Keller. “Erratum to “Deformed Calabi-Yau completions””. preprint. 2018. arXiv: [1809.01126 \[math.RT\]](#).
- [Kha19] A. A. Khan. “Virtual fundamental classes of derived stacks P”. preprint. 2019. arXiv: [1909.01332 \[math.AG\]](#).
- [Kin22] T. Kinjo. “Dimensional reduction in cohomological Donaldson-Thomas theory”. *Compos. Math.* 158.1 (2022), pp. 123–167. arXiv: [2102.01568 \[math.AG\]](#).
- [Kin24] T. Kinjo. “An introduction to cohomological Donaldson-Thomas theory”. preprint. 2024. URL: <https://drive.google.com/file/d/1fsbwvee0IXWsumqaF6NmDCInizH3rEhs/view>.
- [KK] A. Khan and T. Kinjo. “Derived microlocal geometry and cohomological Donaldson-Thomas theory”. in preparation.
- [KK21] T. Kinjo and N. Koseki. “Cohomological χ -independence for Higgs bundles and Gopakumar-Vafa invariants”. preprint. 2021. arXiv: [2112.10053 \[math.AG\]](#).
- [KK23] A. Khan and T. Kinjo. “3d cohomological Hall algebras for local surfaces”. preprint. 2023.
- [KL16] Y.-H. Kiem and J. Li. “Categorification of Donaldson-Thomas invariants via perverse sheaves”. preprint. 2016. arXiv: [1212.6444 \[math.AG\]](#).
- [KM24] T. Kinjo and N. Masuda. “Global Critical Chart for Local Calabi-Yau Threefolds”. *Int. Math. Res. Not. IMRN* 5 (2024), pp. 4062–4093. arXiv: [2112.10052 \[math.AG\]](#).
- [KM76] F. F. Knudsen and D. Mumford. “The projectivity of the moduli space of stable curves. I. Preliminaries on “det” and “Div””. *Math. Scand.* 39.1 (1976), pp. 19–55.
- [KS08] M. Kontsevich and Y. Soibelman. “Stability structures, motivic Donaldson-Thomas invariants and cluster transformations”. preprint. 2008. arXiv: [0811.2435 \[math.AG\]](#).
- [KS11] M. Kontsevich and Y. Soibelman. “Cohomological Hall algebra, exponential Hodge structures and motivic Donaldson-Thomas invariants”. *Commun. Number Theory Phys.* 5.2 (2011), pp. 231–352. arXiv: [1006.2706 \[math.AG\]](#).
- [KS94] M. Kashiwara and P. Schapira. *Sheaves on manifolds*. Vol. 292. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. With a chapter in French by Christian Houzel, Corrected reprint of the 1990 original. Springer-Verlag, Berlin, 1994, pp. x+512.
- [KV23] M. Kapranov and E. Vasserot. “The cohomological Hall algebra of a surface and factorization cohomology”. *J. Eur. Math. Soc. (JEMS)* 25.11 (2023), pp. 4221–4289. arXiv: [1901.07641 \[math.AG\]](#).
- [Lur09] J. Lurie. *Higher topos theory*. Vol. 170. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2009, pp. xviii+925.
- [Lur15] J. Lurie. “Rotation invariance in algebraic K-theory”. preprint. 2015. URL: <https://www.math.ias.edu/~lurie/papers/Waldhaus.pdf>.
- [Lur17] J. Lurie. *Higher Algebra*. 2017. URL: <https://www.math.ias.edu/~lurie/papers/HA.pdf>.

- [Lur18] J. Lurie. *Spectral Algebraic Geometry*. 2018. URL: <https://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf>.
- [Mas01] D. B. Massey. “The Sebastiani–Thom isomorphism in the derived category”. *Compositio Mathematica* 125.3 (2001), pp. 353–362. arXiv: [math/9908101](https://arxiv.org/abs/math/9908101).
- [Mil80] J. S. Milne. *Étale cohomology*. Vol. No. 33. Princeton Mathematical Series. Princeton University Press, Princeton, NJ, 1980, pp. xiii+323.
- [Nak17] H. Nakajima. “Lectures on perverse sheaves on instanton moduli spaces”. *Geometry of moduli spaces and representation theory*. Vol. 24. IAS/Park City Math. Ser. Amer. Math. Soc., Providence, RI, 2017, pp. 381–436. arXiv: [1604.06316](https://arxiv.org/abs/1604.06316) [[math.RT](https://arxiv.org/abs/math.RT)].
- [NS23] F. Naef and P. Safronov. “Torsion volume forms”. preprint. 2023. arXiv: [2308.08369](https://arxiv.org/abs/2308.08369) [[math.AG](https://arxiv.org/abs/math.AG)].
- [OT23] J. Oh and R. P. Thomas. “Counting sheaves on Calabi–Yau 4-folds, I”. *Duke Mathematical Journal* 172.7 (2023), pp. 1333–1409. arXiv: [2009.05542](https://arxiv.org/abs/2009.05542) [[math.AG](https://arxiv.org/abs/math.AG)].
- [Pan+13] T. Pantev, B. Toën, M. Vaquié, and G. Vezzosi. “Shifted symplectic structures”. *Publ. Math. Inst. Hautes Études Sci.* 117 (2013), pp. 271–328. arXiv: [1111.3209](https://arxiv.org/abs/1111.3209) [[math.AG](https://arxiv.org/abs/math.AG)].
- [Par24] H. Park. “Shifted symplectic pushforwards”. preprint. 2024.
- [Pri18] J. P. Pridham. “Deformation quantisation for (-2) -shifted symplectic structures”. preprint. 2018. arXiv: [1809.11028](https://arxiv.org/abs/1809.11028) [[math.AG](https://arxiv.org/abs/math.AG)].
- [PY24] H. Park and J. You. “An introduction to shifted symplectic structures”. preprint. 2024.
- [Rak20] A. Raksit. “Hochschild homology and the derived de Rham complex revisited”. preprint. 2020. arXiv: [2007.02576](https://arxiv.org/abs/2007.02576) [[math.AG](https://arxiv.org/abs/math.AG)].
- [Saf17] P. Safronov. “Symplectic implosion and the Grothendieck–Springer resolution”. *Transform. Groups* 22.3 (2017), pp. 767–792. arXiv: [1411.2962](https://arxiv.org/abs/1411.2962) [[math.AG](https://arxiv.org/abs/math.AG)].
- [Spr09] T. A. Springer. *Linear algebraic groups*. second. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, 2009, pp. xvi+334.
- [Sta24] Stacks project authors. *The Stacks project*. <https://stacks.math.columbia.edu>. 2024.
- [STV15] T. Schürg, B. Toën, and G. Vezzosi. “Derived algebraic geometry, determinants of perfect complexes, and applications to obstruction theories for maps and complexes”. *J. Reine Angew. Math.* 702 (2015), pp. 1–40. arXiv: [1102.1150](https://arxiv.org/abs/1102.1150) [[math.AG](https://arxiv.org/abs/math.AG)].
- [Thu15] L. Q. Thuong. “Proofs of the integral identity conjecture over algebraically closed fields”. *Duke Mathematical Journal* 164.1 (2015), pp. 157–194. arXiv: [1206.5334](https://arxiv.org/abs/1206.5334) [[math.AG](https://arxiv.org/abs/math.AG)].
- [Tub24] S. Tubach. “Mixed Hodge modules on stacks”. preprint. 2024.
- [TV07] B. Toën and M. Vaquié. “Moduli of objects in dg-categories”. *Ann. Sci. École Norm. Sup. (4)* 40.3 (2007), pp. 387–444. arXiv: [math/0503269](https://arxiv.org/abs/math/0503269).
- [TV08] B. Toën and G. Vezzosi. “Homotopical algebraic geometry. II. Geometric stacks and applications”. *Mem. Amer. Math. Soc.* 193.902 (2008), pp. x+224. arXiv: [math/0404373](https://arxiv.org/abs/math/0404373).

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN
Email address: tkinjo@kurims.kyoto-u.ac.jp

JUNE E HUH CENTER FOR MATHEMATICAL CHALLENGES, KOREA INSTITUTE FOR ADVANCED STUDY, 85 HOEGIRO, DONGDAEMUN-GU, SEOUL 02455, REPUBLIC OF KOREA
Email address: hyeonjunpark@kias.re.kr

SCHOOL OF MATHEMATICS, UNIVERSITY OF EDINBURGH, EDINBURGH, UK
Email address: p.safronov@ed.ac.uk