

# PERMANENCE AND UNIFORM ASYMPTOTIC STABILITY OF POSITIVE SOLUTIONS OF SAIQH MODELS ON TIME SCALES

NEDJOUA ZINE, BENAOUMEUR BAYOUR, AND DELFIM F. M. TORRES

**ABSTRACT.** A Susceptible–Asymptomatic–Infectious–Quarantined–Hospitalized (SAIQH) compartmental model on time scales is introduced and a suitable Lyapunov function is defined. Main results include: the proof that the system is permanent; proof of existence of solution; and sufficient conditions implying the dynamic system to have a unique almost periodic solution that is uniformly asymptotically stable. An example is presented supporting the obtained results.

## 1. INTRODUCTION

In 2019, the COVID-19 pandemic has appeared for the first time in Wuhan, China, attracting many researchers to investigate outbreaks and the spread of viruses [1, 2]. Some of the studies provided new mathematical compartmental models, illustrating well the important contributions of Mathematics to fight communicable diseases. Such models have been used to analyze the corresponding dynamics and to supply useful techniques in disease epidemiology [3, 11, 13].

Kim et al. studied a SAIQH (Susceptible–Asymptomatic–Infectious–Quarantined–Hospitalized) mathematical model to analyze the transmission dynamics of MERS and to estimate transmission rates [10]. Lemos-Paião et al. developed a SAIQH type model for COVID-19, which was important to describe and understand the pandemic in Portugal [11]. Similar to [11], here we also consider the  $H_{IC}$  class of hospitalized individuals in intensive care units.

Let the total living population under study, denoted by  $N(t)$ ,  $t \geq 0$ , be divided into six classes: (i) the susceptible individuals  $S(t)$ ; (ii) the infected individuals without (or with mild) symptoms  $A(t)$  (the Asymptomatic); (iii) infected individuals  $I(t)$  with visible symptoms; (iv) quarantined individuals  $Q(t)$  in isolation at home; (v) hospitalized individuals  $H(t)$ ; (vi) and hospitalized individuals  $H_{IC}(t)$  in intensive care units. The mathematical model introduced and studied in [11] reads:

$$\begin{cases} \dot{S}(t) = \Lambda + \omega n Q(t) - [\lambda(t)(1-p) + \phi p + \gamma] S(t), \\ \dot{A}(t) = \lambda(t)(1-p) S(t) - [q\nu + \gamma] A(t), \\ \dot{I}(t) = q\nu A(t) - [\delta_1 + \gamma] I(t), \\ \dot{Q}(t) = \phi p S(t) + \delta_1 f_1 I(t) + \delta_2(1-f_2-f_3) H(t) - [\omega m + \gamma] Q(t), \\ \dot{H}(t) = \delta_1(1-f_1) I(t) + \eta(1-k) H_{IC}(t) - [\delta_2(1-f_2-f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma] H(t), \\ \dot{H}_{IC}(t) = \delta_2 f_2 H(t) - [\eta(1-k) + \alpha_2 k + \gamma] H_{IC}(t), \end{cases} \quad (1.1)$$

where, for all time  $t \geq 0$ ,

$$\lambda(t) = \frac{\beta(l_A A(t) + I(t) + l_H H(t))}{N(t)}$$

is a bounded positive function with

$$N(t) = S(t) + A(t) + I(t) + Q(t) + H(t) + H_{IC}(t),$$

$\beta, l_A, l_H > 0$ , and all the other parameters in the model are nonnegative. In addition,

$$p, 1-p, k, 1-k, q, f_1, 1-f_1, f_2, f_3, 1-f_2-f_3 \in [0, 1].$$

---

*Date:* Submitted January 31, 2024.

*2020 Mathematics Subject Classification.* 34A12; 34N05.

*Key words and phrases.* SAIQH models; time scales; permanence; almost periodic solutions; uniform asymptotic stability.

For more details on the mathematical model (1.1) we refer the reader to [11].

Nowadays, the study of dynamic equations on time scales is a leading topic of research [4, 5, 6, 7, 9, 14, 15, 16, 17]. Motivated by the above mentioned references and the 2019 paper of Khuddush and Prasad on a  $n$ -species Lotka–Volterra system [9], here we extend (1.1) to a time scale  $\mathbb{T}$ , studying the permanence and uniform asymptotic stability of the unique positive almost periodic solution of the following SAIQH type model on time scales:

$$\begin{cases} S^\Delta(t) = \Lambda + \omega n Q(t) - [\lambda(t)(1-p) + \phi p + \gamma] S^\sigma(t), \\ A^\Delta(t) = \lambda(t)(1-p) S(t) - [q\nu + \gamma] A^\sigma(t), \\ I^\Delta(t) = q\nu A(t) - [\delta_1 + \gamma] I^\sigma(t), \\ Q^\Delta(t) = \phi p S(t) + \delta_1 f_1 I(t) + \delta_2(1-f_2-f_3) H(t) - [\omega m + \gamma] Q^\sigma(t), \\ H^\Delta(t) = \delta_1(1-f_1) I(t) + \eta(1-k) H_{IC}(t) - [\delta_2(1-f_2-f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma] H^\sigma(t), \\ H_{IC}^\Delta(t) = \delta_2 f_2 H(t) - [\eta(1-k) + \alpha_2 k + \gamma] H_{IC}^\sigma(t), \end{cases} \quad (1.2)$$

where all parameters are real nonnegative numbers and  $\lambda(t)$  is bounded positive, as mentioned before. In the particular case  $\mathbb{T} = \mathbb{R}^+$ , system (1.2) reduces to (1.1).

The text is organized as follows. In Section 2, we recall the main necessary notions and results needed in the sequel. The original results are then formulated and proved in Section 3: we show that system (1.2) is permanent (Theorem 3); we give a sufficient condition for existence of a solution to system (1.2) (Theorem 5); and we obtain conditions implying the dynamic system (1.2) to have a unique almost periodic solution that is uniformly asymptotically stable (Theorem 6).

For convenience, in the sequel we put  $x_1 = S$ ,  $x_2 = A$ ,  $x_3 = I$ ,  $x_4 = Q$ ,  $x_5 = H$ , and  $x_6 = H_{IC}$ .

## 2. PRELIMINARIES

For a function  $f(t)$  defined on  $t \in \mathbb{T}^+$ , we set

$$f^L := \inf \{f(t) : t \in \mathbb{T}^+\}, \quad f^U := \sup \{f(t) : t \in \mathbb{T}^+\}.$$

We begin by recalling some basic but fundamental concepts of the calculus on time scales [4]. A time scale  $\mathbb{T}$  is a nonempty closed subset of the real numbers  $\mathbb{R}$ ; on  $\mathbb{T}$  one considers the topology inherited from the real numbers with the standard topology. The jump operators  $\sigma, \rho : \mathbb{T} \rightarrow \mathbb{T}$  and the graininess function  $\mu : \mathbb{T} \rightarrow \mathbb{R}^+$  are defined, respectively, by

$$\sigma(t) = \inf \{\tau \in \mathbb{T} : \tau > t\}, \quad \rho(t) = \sup \{\tau \in \mathbb{T} : \tau < t\}, \quad \text{and} \quad \mu(t) = \sigma(t) - t,$$

supplemented by  $\inf \emptyset = \sup \mathbb{T}$  and  $\sup \emptyset = \inf \mathbb{T}$ . A point  $t \in \mathbb{T}$  is left-dense, left-scattered, right-dense, or right-scattered when  $\rho(t) = t$ ,  $\rho(t) < t$ ,  $\sigma(t) = t$ , and  $\sigma(t) > t$ , respectively. If  $\mathbb{T}$  has a left-scattered maximum  $m$ , then  $\mathbb{T}^\kappa := \mathbb{T} \setminus \{m\}$ ; otherwise,  $\mathbb{T}^\kappa := \mathbb{T}$ . If  $f : \mathbb{T} \rightarrow \mathbb{R}$ , then  $f^\sigma : \mathbb{T} \rightarrow \mathbb{R}$  is given by  $f^\sigma(t) = f(\sigma(t))$  for all  $t \in \mathbb{T}$ .

**Definition 1.** A function  $f : \mathbb{T} \rightarrow \mathbb{R}$  is called regressive provided  $1 + \mu(t)f(t) \neq 0$  for all  $t \in \mathbb{T}^\kappa$ ; a function  $g : \mathbb{T} \rightarrow \mathbb{R}$  is called rd-continuous provided it is continuous at right-dense points in  $\mathbb{T}$  and its left-sided limits exist (finite) at left-dense points in  $\mathbb{T}$ . The set of all regressive and rd-continuous functions  $f : \mathbb{T} \rightarrow \mathbb{R}$  will be denoted by  $\mathcal{R} = \mathcal{R}(\mathbb{T}, \mathbb{R})$ . We define the set  $\mathcal{R}^+ = \mathcal{R}^+(\mathbb{T}, \mathbb{R}) = \{f \in \mathcal{R} : 1 + \mu(t)f(t) > 0, \text{ for all } t \in \mathbb{T}\}$ .

**Lemma 1** (See [8]). *Assume that  $\alpha > 0$ ,  $b > 0$ , and  $-\alpha \in \mathcal{R}^+$ . If*

$$y^\Delta(t) \geq (\leq) b - \alpha y^\sigma(t), \quad y(t) > 0, \quad t \in [t_0, \infty)_{\mathbb{T}},$$

then

$$y(t) \geq (\leq) \frac{b}{\alpha} \left[ 1 + \left( \frac{\alpha y(t_0)}{b} - 1 \right) e_{(-\alpha)}(t, t_0) \right], \quad t \in [t_0, \infty)_{\mathbb{T}},$$

where  $e_{(\cdot)}(\cdot, \cdot)$  is the standard exponential function of the time-scale calculus [4].

**Definition 2** (See [12]). A time scale  $\mathbb{T}$  is called an almost periodic time scale if

$$\Pi = \{\tau \in \mathbb{R} : t + \tau \in \mathbb{T}, \text{ for all } t \in \mathbb{T}\} \neq \{0\}.$$

**Definition 3** (See [12]). Let  $\mathbb{T}$  be an almost periodic time scale. A function  $x \in C(\mathbb{T}, \mathbb{R}^n)$  is called an almost periodic function if the  $\varepsilon$ -translation set of  $x$ , that is,

$$E\{\varepsilon, x\} = \{\tau \in \Pi : |x(t + \tau) - x(t)| < \varepsilon \text{ for all } t \in \mathbb{T}\},$$

is a relatively dense set in  $\mathbb{T}$  for all  $\varepsilon > 0$  and there exists a constant  $l(\varepsilon) > 0$  such that each interval of length  $l(\varepsilon)$  contains a  $\tau \in E\{\varepsilon, x\}$  for which  $|x(t + \tau) - x(t)| < \varepsilon$  for all  $t \in \mathbb{T}$ . The value  $\tau$  is known as the  $\varepsilon$ -translation number of  $x$  and  $l(\varepsilon)$  is called the inclusion length of  $E\{\varepsilon, x\}$ .

**Definition 4** (See [12]). Let  $\mathbb{D}$  be an open set in  $\mathbb{R}^n$  and let  $\mathbb{T}$  be a positive almost periodic time scale. A function  $f \in C(\mathbb{T} \times \mathbb{D}, \mathbb{R}^n)$  is called an almost periodic function in  $t \in \mathbb{T}$  uniformly for  $x \in \mathbb{D}$  if the  $\varepsilon$ -translation set of  $f$ ,

$$E\{\varepsilon, f, \mathbb{S}\} = \{\tau \in \Pi : |f(t + \tau) - f(t)| < \varepsilon, \text{ for all } (t, x) \in \mathbb{T} \times \mathbb{S}\},$$

is a relatively dense set in  $\mathbb{T}$  for all  $\varepsilon > 0$  and, for each compact subset  $\mathbb{S}$  of  $\mathbb{D}$ , that is, for any given  $\varepsilon > 0$  and each compact subset  $\mathbb{S}$  of  $\mathbb{D}$ , there exists a constant  $l(\varepsilon, \mathbb{S}) > 0$  such that each interval of length  $l(\varepsilon, \mathbb{S})$  contains  $\tau(\varepsilon, \mathbb{S}) \in E\{\varepsilon, f, \mathbb{S}\}$  for which

$$|f(t + \tau, x) - f(t, x)| < \varepsilon, \text{ for all } (t, x) \in \mathbb{T} \times \mathbb{S}.$$

Consider a system

$$x^\Delta(t) = h(t, x), \quad (2.1)$$

where  $h : \mathbb{T}^+ \times \mathbb{S}_B \rightarrow \mathbb{R}^n$ ,  $\mathbb{S}_B = \{x \in \mathbb{R}^n : \|x\| < B\}$  and  $h(t, x)$  is almost periodic in  $t$  uniformly for  $x \in \mathbb{S}_B$  and is continuous in  $x$ . In [17] the question of existence of a unique almost periodic solution  $f(t) \in \mathbb{S}$  of (2.1), which is uniformly asymptotically stable, is investigated. For our model, we obtain from [17] the following result.

**Lemma 2** (cf. [17]). *Suppose that there exists a Lyapunov function  $V(t, x, z)$  defined on  $\mathbb{T}^+ \times \mathbb{S}_B \times \mathbb{S}_B$  satisfying the following conditions:*

- (i)  $a(\|x - z\|) \leq V(t, x, z) \leq b(\|x - z\|)$ , where  $a, b \in \mathbb{K}$   
with  $\mathbb{K} = \{\alpha \in C(\mathbb{R}^+, \mathbb{R}^+) : \alpha(0) = 0 \text{ and } \alpha \text{ is increasing}\}$ ;
- (ii)  $|V(t, x, z) - V(t, x_1, z_1)| \leq L(\|x - x_1\| + \|z - z_1\|)$ , where  $L > 0$  is a constant;
- (iii)  $D^+V^\Delta(t, x, z) \leq -cV(t, x, z)$ , where  $c > 0$ ,  $-c \in \mathcal{R}^+$ ,

and  $D^+V^\Delta$  is the Dini derivative of  $V$ . Furthermore, if there exists a solution  $x(t) \in \mathbb{S}$  of system

$$\begin{cases} x_1^\Delta(t) = \Lambda + \omega n x_4(t) - [\lambda(t)(1 - p) + \phi p + \gamma] x_1^\sigma(t), \\ x_2^\Delta(t) = \lambda(t)(1 - p) x_1(t) - [q\nu + \gamma] x_2^\sigma(t), \\ x_3^\Delta(t) = q\nu x_2(t) - [\delta_1 + \gamma] x_3^\sigma(t), \\ x_4^\Delta(t) = \phi p x_1(t) + \delta_1 f_1 x_3(t) + \delta_2(1 - f_2 - f_3) x_5(t) - [\omega n + \gamma] x_4^\sigma(t), \\ x_5^\Delta(t) = \delta_1(1 - f_1) x_3(t) + \eta(1 - k) x_6(t) - [\delta_2(1 - f_2 - f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma] x_5^\sigma(t), \\ x_6^\Delta(t) = \delta_2 f_2 x_5(t) - [\eta(1 - k) + \alpha_2 k + \gamma] x_6^\sigma(t), \end{cases} \quad (2.2)$$

for  $t \in \mathbb{T}^+$ , where  $\mathbb{S} \cup \mathbb{S}_B$  is a compact set, then there exists a unique almost periodic solution  $f(t) \in \mathbb{S}$  of system (2.2), which is uniformly asymptotically stable.

### 3. MAIN RESULTS

Let  $t_0 \in \mathbb{T}$  be a fixed positive initial time. Our main results are: the proof that system (2.2) is permanent (Section 3.1); a sufficient condition for existence of a solution to system (2.2) (Section 3.2); and conditions that imply the dynamic system (2.2) to have a unique almost periodic solution that is uniformly asymptotically stable (Section 3.3).

**3.1. Permanence of solutions.** We begin by introducing the notion of permanence of solutions.

**Definition 5.** System (2.2) is said to be permanent if there exist positive constants  $m$  and  $M$  such that  $m \leq \liminf_{t \rightarrow \infty} x_i(t) \leq \limsup_{t \rightarrow \infty} x_i(t) \leq M$ ,  $i = 1, 2, \dots, 6$ , for any solution  $(x_1(t), \dots, x_6(t))$  of (2.2).

**Theorem 3.** *System (2.2) is permanent.*

*Proof.* Let  $Z(t) = (x_1(t), \dots, x_6(t))$  be a positive solution of system (2.2). Then,

$$\begin{cases} x_1^\Delta(t) \geq \Lambda - [\lambda^U(1-p) + \phi p + \gamma]x_1^\sigma(t), \\ x_2^\Delta(t) \geq \lambda^L(1-p)m_1 - (q\nu + \gamma)x_2^\sigma(t), \\ x_3^\Delta(t) \geq q\nu m_2 - (\delta_1 + \gamma)x_3^\sigma(t), \\ x_4^\Delta(t) \geq \phi p m_1 + \delta_1 f_1 m_3 - (\omega n + \gamma)x_4^\sigma(t), \\ x_5^\Delta(t) \geq \delta_1(1-f_1)m_3 - [\delta_2(1-f_2-f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma]x_5^\sigma(t), \\ x_6^\Delta(t) \geq \delta_2 f_2 m_5 - [\eta(1-k) + \alpha_2 k + \gamma]x_6^\sigma(t). \end{cases}$$

From Lemma 1, it follows that

$$x_1(t) \geq \frac{\Lambda}{\lambda^u(1-p) + \phi p + \gamma} \left[ 1 + \left( \frac{\lambda^U(1-p) + \phi p + \gamma}{\Lambda} x_1(t_0) - 1 \right) e_{-(\lambda^U(1-p) + \phi p + \gamma)}(t, t_0) \right]$$

and we have  $e_{-(\lambda^U(1-p) + \phi p + \gamma)}(t, t_0) \rightarrow 0$ , as  $t \rightarrow \infty$ . Thus,

$$\begin{cases} x_1(t) \geq m_1 := \frac{\Lambda}{\lambda^U(1-p) + \phi p + \gamma}, & \text{for } t \geq T_1, \\ x_2(t) \geq m_2 := \frac{\lambda^L(1-p)m_1}{q\nu + \gamma}, & \text{for } t \geq T_2, \\ x_3(t) \geq m_3 := \frac{q\nu m_2}{\delta_1 + \gamma}, & \text{for } t \geq T_3, \\ x_4(t) \geq m_4 := \frac{\phi p m_1 + \delta_1 f_1 m_3}{\omega n + \gamma}, & \text{for } t \geq T_4, \\ x_5(t) \geq m_5 := \frac{\delta_1(1-f_1)m_3}{\delta_2(1-f_2-f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma}, & \text{for } t \geq T_5, \\ x_6(t) \geq m_6 := \frac{\delta_2 f_2 m_5}{\eta(1-k) + \alpha_2 k + \gamma}, & \text{for } t \geq T_6. \end{cases}$$

Let  $m = \min_{1 \leq i \leq 6} m_i$  and  $T = \max_{1 \leq i \leq 6} T_i$ . We can then write that  $x_i(t) \geq m$  for all  $t > T$ .  $\square$

**3.2. Existence of solution.** For system (2.2), we introduce the following assumption:

(H<sub>1</sub>)  $\lambda(t)$  is a bounded almost periodic function and satisfy  $0 < \lambda^L \leq \lambda(t) \leq \lambda^U$ .

To prove existence of solution, we first begin with a technical lemma.

**Lemma 4.** *If (H<sub>1</sub>) holds, then, for any positive solution  $Z(t) = (x_1(t), \dots, x_6(t))$  of system (2.2), there exist positive constants  $M$  and  $T$  such that  $x_i(t) < M$ ,  $i = 1, \dots, 6$ , for all  $t > T$ .*

*Proof.* Let  $Z(t) = (x_1(t), \dots, x_6(t))$  be a positive solution of system (2.2). Then,

$$\begin{cases} x_1^\Delta(t) \leq \Lambda + \omega n \frac{\Lambda}{\gamma} - [\lambda^L(1-p) + \phi p + \gamma]x_1^\sigma(t), \\ x_2^\Delta(t) \leq \lambda^U(1-p)M_1 - (q\nu + \gamma)x_2^\sigma(t), \\ x_3^\Delta(t) \leq q\nu M_2 - (\delta_1 + \gamma)x_3^\sigma(t), \\ x_4^\Delta(t) \leq \phi p M_1 + \delta_1 f_1 M_3 + \delta_2(1-f_2-f_3)\frac{\Lambda}{\gamma} - (\omega n + \gamma)x_4^\sigma(t), \\ x_5^\Delta(t) \leq \delta_1(1-f_1)M_3 + \eta(1-k)\frac{\Lambda}{\gamma} - [\delta_2(1-f_2-f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma]x_5^\sigma(t), \\ x_6^\Delta(t) \leq \delta_2 f_2 M_5 - [\eta(1-k) + \alpha_2 k + \gamma]x_6^\sigma(t). \end{cases}$$

From Lemma 1,

$$x_1(t) \leq \frac{\Lambda + \omega n \frac{\Lambda}{\gamma}}{\lambda^L(1-p) + \phi p + \gamma} \left[ 1 + \left( \frac{\lambda^L(1-p) + \phi p + \gamma}{\Lambda + \omega n \frac{\Lambda}{\gamma}} x_1(t_0) - 1 \right) e_{-(\lambda^L(1-p) + \phi p + \gamma)}(t, t_0) \right]$$

and we have  $e_{-(\lambda^L(1-p)+\phi p+\gamma)}(t, t_0) \rightarrow 0$ , as  $t \rightarrow \infty$ . Then,

$$x_1(t) \leq M_1 := \frac{\Lambda + \omega n \frac{\Delta}{\gamma}}{\lambda^L(1-p) + \phi p + \gamma},$$

$$\left\{ \begin{array}{ll} x_1(t) \leq M_1 := \frac{\Lambda + \omega n \frac{\Delta}{\gamma}}{\lambda^L(1-p) + \phi p + \gamma}, & \text{for } t \geq T_1, \\ x_2(t) \leq M_2 := \frac{\lambda^U(1-p)M_1}{q\nu + \gamma}, & \text{for } t \geq T_2, \\ x_3(t) \leq M_3 := \frac{q\nu M_2}{\delta_1 + \gamma}, & \text{for } t \geq T_3, \\ x_4(t) \leq M_4 := \frac{\phi p M_1 + \delta_1 f_1 M_3 + \delta_2(1-f_2-f_3)\frac{\Delta}{\gamma}}{\omega n + \gamma}, & \text{for } t \geq T_4, \\ x_5(t) \leq M_5 := \frac{\delta_1(1-f_1)M_3 + \eta(1-k)\frac{\Delta}{\gamma}}{\delta_2(1-f_2-f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma}, & \text{for } t \geq T_5, \\ x_6(t) \leq M_6 := \frac{\delta_2 f_2 M_5}{\eta(1-k) + \alpha_2 k + \gamma}, & \text{for } t \geq T_6. \end{array} \right.$$

Let  $M = \max_{1 \leq i \leq 6} M_i$  and  $T = \max_{1 \leq i \leq 6} T_i$ . Then  $x_i(t) \leq M$  for all  $t > T$ .  $\square$

Define

$$\Omega = \{(x_1(t), \dots, x_6(t)) : (x_1(t), \dots, x_6(t)) \text{ is a solution of (2.2) and } 0 < m \leq x_i \leq M, i = 1, \dots, 6\}.$$

**Theorem 5.** *If (H1) holds, then the set  $\Omega$  is nonempty.*

*Proof.* The almost periodicity of  $\lambda(t)$  implies that there is a sequence  $\{\xi_l\} \subseteq \mathbb{T}^+$  with  $l \rightarrow \infty$  such that

$$\lambda(t + \xi_l) \rightarrow \lambda(t) \text{ as } l \rightarrow \infty.$$

From Theorem 3 and Lemma 4, for each sufficiently small  $\varepsilon > 0$ , there exists a  $t_1 \in \mathbb{T}^+$  such that

$$m - \varepsilon \leq x_i(t) \leq M + \varepsilon, \text{ for all } t \geq t_1, \quad i = 1, \dots, 6.$$

Set  $x_{il}(t) = x_i(t + \xi_l)$  for  $t \geq t_1 - \xi_l$ ,  $l = 1, 2, \dots$ . For any positive integer  $k$ , we obtain that there exists a sequence  $\{x_{il}(t) : l \geq k\}$  such that the sequence  $\{x_{il}(t)\}$  has a subsequence, denoted by  $\{x_{il}^*(t)\}$  ( $x_{il}^*(t) = x_{il}(t + \xi_l^*)$ ), converging on any finite interval of  $\mathbb{T}^+$  as  $l \rightarrow \infty$ . So we have a sequence  $\{y_i(t)\}$  such that, for  $t \in \mathbb{T}^+$ ,

$$x_{il}^*(t) \rightarrow y_i(t), \text{ as } l \rightarrow \infty, \quad i = 1, \dots, 6. \quad (3.1)$$

It is easy to see that the above sequence  $\{\xi_l^*\} \subseteq \mathbb{T}^+$  with  $\xi_l^* \rightarrow \tau$  for  $l \rightarrow \infty$  is such that

$$\lambda(t + \xi_l^*) \rightarrow \lambda(t), \text{ as } l \rightarrow \infty,$$

which, together with (3.1) and

$$\left\{ \begin{array}{l} x_{1l}^{\Delta}(t) = \Lambda + \omega n x_{4l}^*(t) - [\lambda(t + \xi_l^*)(1-p) + \phi p + \gamma] x_{1l}^{*\sigma}(t), \\ x_{2l}^{\Delta}(t) = \lambda(t + \xi_l^*)(1-p) x_{1l}^*(t) - [q\nu + \gamma] x_{2l}^{*\sigma}(t), \\ x_{3l}^{\Delta}(t) = q\nu x_{2l}^*(t) - [\delta_1 + \gamma] x_{3l}^{*\sigma}(t), \\ x_{4l}^{\Delta}(t) = \phi p x_{1l}^*(t) + \delta_1 f_1 x_{3l}^*(t) + \delta_2(1-f_2-f_3) x_{5l}^*(t) - [\omega n + \gamma] x_{4l}^{*\sigma}(t), \\ x_{5l}^{\Delta}(t) = \delta_1(1-f_1) x_{3l}^*(t) + \eta(1-k) x_{6l}^*(t) - [\delta_2(1-f_2-f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma] x_{5l}^{*\sigma}(t), \\ x_{6l}^{\Delta}(t) = \delta_2 f_2 x_{5l}^*(t) - [\eta(1-k) + \alpha_2 k + \gamma] x_{6l}^{*\sigma}(t), \end{array} \right.$$

yields

$$\left\{ \begin{array}{l} y_1^{\Delta}(t) = \Lambda + \omega n y_4(t) - [\lambda(t)(1-p) + \phi p + \gamma] y_1^{\sigma}(t), \\ y_2^{\Delta}(t) = \lambda(t)(1-p) y_1(t) - [q\nu + \gamma] y_2^{\sigma}(t), \\ y_3^{\Delta}(t) = q\nu y_2(t) - [\delta_1 + \gamma] y_3^{\sigma}(t), \\ y_4^{\Delta}(t) = \phi p y_1(t) + \delta_1 f_1 y_3(t) + \delta_2(1-f_2-f_3) y_5(t) - [\omega n + \gamma] y_4^{\sigma}(t), \\ y_5^{\Delta}(t) = \delta_1(1-f_1) y_3(t) + \eta(1-k) y_6(t) - [\delta_2(1-f_2-f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma] y_5^{\sigma}(t), \\ y_6^{\Delta}(t) = \delta_2 f_2 y_5(t) - [\eta(1-k) + \alpha_2 k + \gamma] y_6^{\sigma}(t). \end{array} \right.$$

It is clear that  $(y_1(t), \dots, y_6(t))$  is a solution of system (2.2) and

$$m - \varepsilon \leq y_i(t) \leq M + \varepsilon, \quad \text{for all } t \in \mathbb{T}^+, \quad i = 1, \dots, 6.$$

Since  $\varepsilon$  is arbitrary, it follows that

$$m \leq y_i(t) \leq M, \quad \text{for } t \in \mathbb{T}^+, \quad i = 1, \dots, 6.$$

The proof is complete.  $\square$

**3.3. Uniform asymptotic stability.** Now, we establish sufficient conditions for the existence of a unique positive almost periodic solution to system (2.2) that is uniform asymptotically stable. We introduce some more notations. Let

$$\begin{aligned} A_1 &:= \lambda^L(1-p) + \phi p + \gamma; & A_2 &:= q\nu + \gamma; \\ A_3 &:= \delta_1 + \gamma; & A_4 &:= \omega n + \gamma; \\ A_5 &:= \delta_2(1-f_3) + \alpha_1 f_3 + \gamma; & A_6 &:= \eta(1-k) + \alpha_2 k + \gamma; \\ B_1 &:= \lambda^U(1-p) + \phi p; & B_2 &:= q\nu + 2\frac{\gamma\beta l_A(1-p)M}{\Lambda}; \\ B_3 &:= \delta_1 + 2\frac{\gamma\beta(1-p)M}{\Lambda}; & B_4 &:= \omega n; \\ B_5 &:= \delta_2(1-f_3) + 2\frac{\gamma\beta l_H(1-p)M}{\Lambda}; & B_6 &:= \eta(1-k). \end{aligned}$$

Moreover, let  $A := \min_{1 \leq i \leq 6} A_i$  and  $B := \max_{1 \leq i \leq 6} B_i$ .

In our next result (Theorem 6), we assume the following additional hypothesis:

(H2)  $B < A$ .

**Theorem 6.** *If (H1) and (H2) hold, then the dynamic system (2.2) has a unique almost periodic solution  $Z(t) = (x_1(t), \dots, x_6(t)) \in \Omega$  that is uniformly asymptotically stable.*

*Proof.* According to Theorem 3, every solution  $Z(t) = (x_1(t), \dots, x_6(t))$  of system (2.2) satisfies  $x_i^L \leq x_i \leq x_i^U$ . Hence,  $|x_i(t)| \leq K_i$ ,  $i = 1, \dots, 6$ . Suppose that  $Z(t) = (x_1(t), \dots, x_6(t))$  and  $\widehat{Z}(t) = (\widehat{x}_1(t), \dots, \widehat{x}_6(t))$  are two positive solutions of system (2.2). We have

$$\begin{cases} x_1^\Delta(t) = \Lambda + \omega n x_4(t) - [\lambda(t)(1-p) + \phi p + \gamma]x_1^\sigma(t), \\ x_2^\Delta(t) = \lambda(t)(1-p)x_1(t) - [q\nu + \gamma]x_2^\sigma(t), \\ x_3^\Delta(t) = q\nu x_2(t) - [\delta_1 + \gamma]x_3^\sigma(t), \\ x_4^\Delta(t) = \phi p x_1(t) + \delta_1 f_1 x_3(t) + \delta_2(1-f_2-f_3)x_5(t) - [\omega n + \gamma]x_4^\sigma(t), \\ x_5^\Delta(t) = \delta_1(1-f_1)x_3(t) + \eta(1-k)x_6(t) - [\delta_2(1-f_2-f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma]x_5^\sigma(t), \\ x_6^\Delta(t) = \delta_2 f_2 x_5(t) - [\eta(1-k) + \alpha_2 k + \gamma]x_6^\sigma(t), \end{cases}$$

and

$$\begin{cases} \widehat{x}_1^\Delta(t) = \Lambda + \omega n \widehat{x}_4(t) - [\widehat{\lambda}(t)(1-p) + \phi p + \gamma]\widehat{x}_1^\sigma(t), \\ \widehat{x}_2^\Delta(t) = \widehat{\lambda}(t)(1-p)\widehat{x}_1(t) - [q\nu + \gamma]\widehat{x}_2^\sigma(t), \\ \widehat{x}_3^\Delta(t) = q\nu \widehat{x}_2(t) - [\delta_1 + \gamma]\widehat{x}_3^\sigma(t), \\ \widehat{x}_4^\Delta(t) = \phi p \widehat{x}_1(t) + \delta_1 f_1 \widehat{x}_3(t) + \delta_2(1-f_2-f_3)\widehat{x}_5(t) - [\omega n + \gamma]\widehat{x}_4^\sigma(t), \\ \widehat{x}_5^\Delta(t) = \delta_1(1-f_1)\widehat{x}_3(t) + \eta(1-k)\widehat{x}_6(t) - [\delta_2(1-f_2-f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma]\widehat{x}_5^\sigma(t), \\ \widehat{x}_6^\Delta(t) = \delta_2 f_2 \widehat{x}_5(t) - [\eta(1-k) + \alpha_2 k + \gamma]\widehat{x}_6^\sigma(t). \end{cases}$$

Denote

$$\|Z\| = \|(x_1(t), \dots, x_6(t))\| = \sup_{t \in \mathbb{T}^+} \sum_{i=1}^6 |x_i(t)|.$$

Then  $\|Z\| \leq K$  and  $\|\widehat{Z}\| \leq K$  where  $K = \sum_{i=1}^6 K_i$ . Define the Lyapunov function  $V(t, Z, \widehat{Z})$  on  $\mathbb{T}^+ \times \Omega \times \Omega$  as

$$V(t, Z, \widehat{Z}) = \sum_{i=1}^6 |x_i(t) - \widehat{x}_i(t)| = \sum_{i=1}^6 V_i(t), \quad (3.2)$$

where  $V_i(t) = |x_i(t) - \widehat{x}_i(t)|$ . Then the two norms

$$\|Z - \widehat{Z}\| = \sum_{i=1}^6 |x_i(t) - \widehat{x}_i(t)|$$

and

$$\|Z - \widehat{Z}\|_* = \sup_{t \in \mathbb{R}_+} \left[ \sum_{i=1}^6 (x_i(t) - \widehat{x}_i(t))^2 \right]^{\frac{1}{2}}$$

are equivalent, that is, there exist two constants  $\eta_1, \eta_2 > 0$  such that

$$\eta_1 \|Z - \widehat{Z}\|_* \leq \|Z - \widehat{Z}\| \leq \eta_2 \|Z - \widehat{Z}\|_*.$$

Hence,

$$\eta_1 \|Z - \widehat{Z}\|_* \leq V(t, Z, \widehat{Z}) \leq \eta_2 \|Z - \widehat{Z}\|_*.$$

Let  $a, b \in C(\mathbb{R}^+, \mathbb{R}^+)$ ,  $a(x) = \eta_1 x$ , and  $b(x) = \eta_2 x$ . Then the assumption (i) of Lemma 2 is satisfied. On the other hand, we have

$$\begin{aligned} |V(t, Z(t), \widehat{Z}(t)) - V(t, Z^*(t), \widehat{Z}^*(t))| &= \left| \sum_{i=1}^6 |x_i(t) - \widehat{x}_i(t)| - \sum_{i=1}^6 |x_i^*(t) - \widehat{x}_i^*(t)| \right| \\ &\leq \sum_{i=1}^6 ||x_i(t) - \widehat{x}_i(t)| - |x_i^*(t) - \widehat{x}_i^*(t)|| \\ &\leq \sum_{i=1}^6 |x_i(t) - x_i^*(t)| + \sum_{i=1}^6 |\widehat{x}_i(t) - \widehat{x}_i^*(t)| \\ &\leq \|Z - Z^*\| + \|\widehat{Z} - \widehat{Z}^*\|, \end{aligned}$$

where  $L = 1$ , so condition (ii) of Lemma 2 is also satisfied.

Now, using Lemma 4.2 of [18], it follows that

$$D^+ V_i^\Delta(t) \leq \text{sign}(x_i^\sigma(t) - \widehat{x}_i^\sigma(t))(x_i^\Delta(t) - \widehat{x}_i^\Delta(t)), \quad i = 1, \dots, 6,$$

where  $D^+ V_i^\Delta$  is the Dini derivative of  $V_i$ . For  $i = 1$ ,

$$\begin{aligned} D^+ V_1^\Delta(t) &\leq \text{sign}(x_1^\sigma(t) - \widehat{x}_1^\sigma(t))(x_1^\Delta(t) - \widehat{x}_1^\Delta(t)) \\ &= \text{sign}(x_1^\sigma(t) - \widehat{x}_1^\sigma(t))[\omega n(x_4(t) - \widehat{x}_4(t)) - \lambda(t)(1-p)x_1^\sigma(t) \\ &\quad - [\phi p + \gamma](x_1^\sigma(t) - \widehat{x}_1^\sigma(t)) + \widehat{\lambda}(t)(1-p)\widehat{x}_1^\sigma(t)] \\ &= \text{sign}(x_1^\sigma(t) - \widehat{x}_1^\sigma(t))[\omega n(x_4(t) - \widehat{x}_4(t)) - \lambda(t)(1-p)x_1^\sigma(t) \\ &\quad - [\phi p + \gamma](x_1^\sigma(t) - \widehat{x}_1^\sigma(t)) + \widehat{\lambda}(t)(1-p)\widehat{x}_1^\sigma(t) + \lambda(t)(1-p)\widehat{x}_1^\sigma(t) - \lambda(t)(1-p)\widehat{x}_1^\sigma(t)] \\ &= \text{sign}(x_1^\sigma(t) - \widehat{x}_1^\sigma(t))[\omega n(x_4(t) - \widehat{x}_4(t)) \\ &\quad - [\lambda(t)(1-p) + \phi p + \gamma](x_1^\sigma(t) - \widehat{x}_1^\sigma(t)) \\ &\quad + (\widehat{\lambda}(t) - \lambda(t))(1-p)\widehat{x}_1^\sigma(t)] \\ &= \text{sign}(x_1^\sigma(t) - \widehat{x}_1^\sigma(t))[\omega n(x_4(t) - \widehat{x}_4(t)) - [\lambda(t)(1-p) + \phi p + \gamma](x_1^\sigma(t) - \widehat{x}_1^\sigma(t)) \\ &\quad + (\widehat{\lambda}(t) - \lambda(t))(1-p)\widehat{x}_1^\sigma(t)] \\ &\leq \omega n |x_4(t) - \widehat{x}_4(t)| - [\lambda^L(1-p) + \phi p + \gamma] |x_1^\sigma(t) - \widehat{x}_1^\sigma(t)| \\ &\quad + \frac{\gamma \beta l_A (1-p) M}{\Lambda} |x_2(t) - \widehat{x}_2(t)| + \frac{\gamma \beta (1-p) M}{\Lambda} |x_3(t) - \widehat{x}_3(t)| \\ &\quad + \frac{\gamma \beta l_H (1-p) M}{\Lambda} |x_5(t) - \widehat{x}_5(t)|; \end{aligned}$$

for  $i = 2$ ,

$$\begin{aligned} D^+ V_2^\Delta(t) &\leq \text{sign}(x_2^\sigma(t) - \widehat{x}_2^\sigma(t))(x_2^\Delta(t) - \widehat{x}_2^\Delta(t)) \\ &= \text{sign}(x_2^\sigma(t) - \widehat{x}_2^\sigma(t))[\lambda(t)(1-p)x_1(t) - [q\nu + \gamma](x_2^\sigma(t) - \widehat{x}_2^\sigma(t)) - \widehat{\lambda}(t)(1-p)\widehat{x}_1(t)] \end{aligned}$$

$$\begin{aligned}
&= \text{sign}(x_2^\sigma(t) - \widehat{x}_2^\sigma(t))[\lambda(t)(1-p)x_1(t) - [q\nu + \gamma](x_2^\sigma(t) - \widehat{x}_2^\sigma(t)) \\
&\quad - \widehat{\lambda}(t)(1-p)\widehat{x}_1(t) + \lambda(t)(1-p)\widehat{x}_1(t) - \lambda(t)(1-p)\widehat{x}_1(t)] \\
&= \text{sign}(x_2^\sigma(t) - \widehat{x}_2^\sigma(t))[-[q\nu + \gamma](x_2^\sigma(t) - \widehat{x}_2^\sigma(t)) \\
&\quad + \lambda(t)(1-p)(x_1(t) - \widehat{x}_1(t)) - (\widehat{\lambda}(t) - \lambda(t))(1-p)\widehat{x}_1(t)] \\
&\leq \lambda^U(1-p)|x_1(t) - \widehat{x}_1(t)| - [q\nu + \gamma]|x_2^\sigma(t) - \widehat{x}_2^\sigma(t)| \\
&\quad + (1-p)M|\lambda(t) - \widehat{\lambda}(t)| \\
&\leq \lambda^U(1-p)|x_1(t) - \widehat{x}_1(t)| - [q\nu + \gamma]|x_2^\sigma(t) - \widehat{x}_2^\sigma(t)| \\
&\quad + \frac{\gamma\beta l_A(1-p)M}{\Lambda}|x_2(t) - \widehat{x}_2(t)| \\
&\quad + \frac{\gamma\beta(1-p)M}{\Lambda}|x_3(t) - \widehat{x}_3(t)| + \frac{\gamma\beta l_H(1-p)M}{\Lambda}|x_5(t) - \widehat{x}_5(t)|;
\end{aligned}$$

for  $i = 3$ ,

$$\begin{aligned}
D^+V_3^\Delta(t) &\leq \text{sign}(x_3^\sigma(t) - \widehat{x}_3^\sigma(t))(x_3^\Delta(t) - \widehat{x}_3^\Delta(t)) \\
&= \text{sign}(x_3^\sigma(t) - \widehat{x}_3^\sigma(t))[q\nu(x_2(t) - \widehat{x}_2(t)) \\
&\quad - [\delta_1 + \gamma](x_3^\sigma(t) - \widehat{x}_3^\sigma(t))] \\
&\leq q\nu|x_2(t) - \widehat{x}_2(t)| - [\delta_1 + \gamma]|x_3^\sigma(t) - \widehat{x}_3^\sigma(t)|;
\end{aligned}$$

for  $i = 4$ ,

$$\begin{aligned}
D^+V_4^\Delta(t) &\leq \text{sign}(x_4^\sigma(t) - \widehat{x}_4^\sigma(t))(x_4^\Delta(t) - \widehat{x}_4^\Delta(t)) \\
&= \text{sign}(x_4^\sigma(t) - \widehat{x}_4^\sigma(t))[\phi p(x_1(t) - \widehat{x}_1(t)) + \delta_1 f_1(x_3(t) - \widehat{x}_3(t)) \\
&\quad + \delta_2(1 - f_2 - f_3)(x_5(t) - \widehat{x}_5(t)) - [\omega n + \gamma](x_4^\sigma(t) - \widehat{x}_4^\sigma(t))] \\
&\leq \phi p|x_1(t) - \widehat{x}_1(t)| + \delta_1 f_1|x_3(t) - \widehat{x}_3(t)| \\
&\quad + \delta_2(1 - f_2 - f_3)|x_5(t) - \widehat{x}_5(t)| - [\omega n + \gamma]|x_4^\sigma(t) - \widehat{x}_4^\sigma(t)|;
\end{aligned}$$

for  $i = 5$ ,

$$\begin{aligned}
D^+V_5^\Delta(t) &\leq \text{sign}(x_5^\sigma(t) - \widehat{x}_5^\sigma(t))(x_5^\Delta(t) - \widehat{x}_5^\Delta(t)) \\
&= \text{sign}(x_5^\sigma(t) - \widehat{x}_5^\sigma(t))[\delta_1(1 - f_1)(x_3(t) - \widehat{x}_3(t)) + \eta(1 - k)(x_6(t) - \widehat{x}_6(t)) \\
&\quad - [\delta_2(1 - f_2 - f_3) + \delta_1 f_1 + \alpha_1 f_3 + \gamma](x_5^\sigma(t) - \widehat{x}_5^\sigma(t))] \\
&\leq \delta_1(1 - f_1)|x_3(t) - \widehat{x}_3(t)| + \eta(1 - k)|x_6(t) - \widehat{x}_6(t)| \\
&\quad - [\delta_2(1 - f_2 - f_3) + \delta_1 f_1 + \alpha_1 f_3 + \gamma]|x_5^\sigma(t) - \widehat{x}_5^\sigma(t)|;
\end{aligned}$$

and, finally, for  $i = 6$ ,

$$\begin{aligned}
D^+V_6^\Delta(t) &\leq \text{sign}(x_6^\sigma(t) - \widehat{x}_6^\sigma(t))(x_6^\Delta(t) - \widehat{x}_6^\Delta(t)) \\
&= \text{sign}(x_6^\sigma(t) - \widehat{x}_6^\sigma(t))[\delta_2 f_2(x_5(t) - \widehat{x}_5(t)) \\
&\quad - [\eta(1 - k) + \alpha_2 k + \gamma](x_6^\sigma(t) - \widehat{x}_6^\sigma(t))] \\
&\leq \delta_2 f_2|x_5(t) - \widehat{x}_5(t)| - [\eta(1 - k) + \alpha_2 k + \gamma]|x_6^\sigma(t) - \widehat{x}_6^\sigma(t)|.
\end{aligned}$$

It follows that

$$\begin{aligned}
D^+V^\Delta(t) &\leq \omega n|x_4(t) - \widehat{x}_4(t)| - [\lambda^L(1-p) + \phi p + \gamma]|x_1^\sigma(t) - \widehat{x}_1^\sigma(t)| \\
&\quad + \frac{\gamma\beta l_A(1-p)M}{\Lambda}|x_2(t) - \widehat{x}_2(t)| + \frac{\gamma\beta(1-p)M}{\Lambda}|x_3(t) - \widehat{x}_3(t)| \\
&\quad + \frac{\gamma\beta l_H(1-p)M}{\Lambda}|x_5(t) - \widehat{x}_5(t)| + \lambda^U(1-p)|x_1(t) - \widehat{x}_1(t)| \\
&\quad - [q\nu + \gamma]|x_2^\sigma(t) - \widehat{x}_2^\sigma(t)| + \frac{\gamma\beta l_A(1-p)M}{\Lambda}|x_2(t) - \widehat{x}_2(t)| \\
&\quad + \frac{\gamma\beta(1-p)M}{\Lambda}|x_3(t) - \widehat{x}_3(t)| + \frac{\gamma\beta l_H(1-p)M}{\Lambda}|x_5(t) - \widehat{x}_5(t)| \\
&\quad + q\nu|x_2(t) - \widehat{x}_2(t)| - [\delta_1 + \gamma]|x_3^\sigma(t) - \widehat{x}_3^\sigma(t)|
\end{aligned}$$



$$\begin{aligned}
& +\phi p |x_1(t) - \widehat{x}_1(t)| + \delta_1 f_1 |x_3(t) - \widehat{x}_3(t)| \\
& +\delta_2(1 - f_2 - f_3) |x_5(t) - \widehat{x}_5(t)| - [\omega n + \gamma] |x_4^\sigma(t) - \widehat{x}_4^\sigma(t)| \\
& +\delta_1(1 - f_1) |x_3(t) - \widehat{x}_3(t)| + \eta(1 - k) |x_6(t) - \widehat{x}_6(t)| \\
& -[\delta_2(1 - f_2 - f_3) + \delta_1 f_1 + \alpha_1 f_3 + \gamma] |x_5^\sigma(t) - \widehat{x}_5^\sigma(t)| \\
& +\delta_2 f_2 |x_5(t) - \widehat{x}_5(t)| - [\eta(1 - k) + \alpha_2 k + \gamma] |x_6^\sigma(t) - \widehat{x}_6^\sigma(t)| \\
= & -[\lambda(t)(1 - p) + \phi p + \gamma] |x_1^\sigma(t) - \widehat{x}_1^\sigma(t)| - [q\nu + \gamma] |x_2^\sigma(t) - \widehat{x}_2^\sigma(t)| \\
& -[\delta_1 + \gamma] |x_3^\sigma(t) - \widehat{x}_3^\sigma(t)| - [\omega n + \gamma] |x_4^\sigma(t) - \widehat{x}_4^\sigma(t)| \\
& -[\delta_2(1 - f_2 - f_3) + \delta_1 f_1 + \alpha_1 f_3 + \gamma] |x_5^\sigma(t) - \widehat{x}_5^\sigma(t)| \\
& -[\eta(1 - k) + \alpha_2 k + \gamma] |x_6^\sigma(t) - \widehat{x}_6^\sigma(t)| \\
& + \{\lambda^U(1 - p) + \phi p\} |x_1(t) - \widehat{x}_1(t)| \\
& + \left\{ q\nu + \frac{\gamma\beta l_A(1-p)M}{\Lambda} + \frac{\gamma\beta l_A(1-p)M}{\Lambda} \right\} |x_2(t) - \widehat{x}_2(t)| \\
& + \left\{ \delta_1 f_1 + \delta_1(1 - f_1) + \frac{\gamma\beta(1-p)M}{\Lambda} + \frac{\gamma\beta(1-p)M}{\Lambda} \right\} |x_3(t) - \widehat{x}_3(t)| \\
& + \omega n |x_4(t) - \widehat{x}_4(t)| + \left\{ \delta_2(1 - f_2 - f_3) + \delta_2 f_2 + \frac{\gamma\beta l_H(1-p)M}{\Lambda} \right. \\
& \left. + \frac{\gamma\beta l_H(1-p)M}{\Lambda} \right\} |x_5(t) - \widehat{x}_5(t)| + \eta(1 - k) |x_6(t) - \widehat{x}_6(t)| \\
= & -A_1 |x_1^\sigma(t) - \widehat{x}_1^\sigma(t)| - A_2 |x_2^\sigma(t) - \widehat{x}_2^\sigma(t)| \\
& -A_3 |x_3^\sigma(t) - \widehat{x}_3^\sigma(t)| - A_4 |x_4^\sigma(t) - \widehat{x}_4^\sigma(t)| \\
& -A_5 |x_5^\sigma(t) - \widehat{x}_5^\sigma(t)| - A_6 |x_6^\sigma(t) - \widehat{x}_6^\sigma(t)| \\
& +B_1 |x_1(t) - \widehat{x}_1(t)| + B_2 |x_2(t) - \widehat{x}_2(t)| \\
& +B_3 |x_3(t) - \widehat{x}_3(t)| + B_4 |x_4(t) - \widehat{x}_4(t)| \\
& +B_5 |x_5(t) - \widehat{x}_5(t)| + B_6 |x_6(t) - \widehat{x}_6(t)| \\
= & -AV(\sigma(t)) + BV(t) \\
= & (B - A)V(t) - A\mu(t)D^+V^\Delta(t)
\end{aligned}$$

and  $D^+V^\Delta(t) \leq \frac{B-A}{1+A\mu(t)}V(t) \leq -\psi(t)V(t)$  with  $\psi = \frac{A-B}{1+A\mu^\sigma}$ . By (H2), we have  $\psi(t) = \frac{A-B}{1+A\mu^\sigma} > 0$  and  $1 - \psi\mu(t) = 1 + A(\mu^U - \mu(t)) + \mu(t)B > 0$ . Hence,  $-\psi \in \mathcal{R}^+$ . Thus, the assumption (iii) of Lemma 2 is satisfied and it follows from Lemma 2 that there exists a unique almost periodic solution  $Z(t) = (x_1(t), \dots, x_6(t))$  of the dynamic system (2.2) that is uniformly asymptotically stable with  $Z(t) \in \Omega$ .  $\square$

We illustrate our results with an example.

**Example 7.** Based on [11], let us consider the following system on the time scale  $\mathbb{T} = \mathbb{Z}_0^+$ :

$$\begin{cases}
x_1^\Delta(t) = \Lambda + \omega n x_4(t) - [\lambda(t)(1 - p) + \phi p + \gamma] x_1^\sigma(t), \\
x_2^\Delta(t) = \lambda(t)(1 - p)x_1(t) - [q\nu + \gamma] x_2^\sigma(t), \\
x_3^\Delta(t) = q\nu x_2(t) - [\delta_1 + \gamma] x_3^\sigma(t), \\
x_4^\Delta(t) = \phi p x_1(t) + \delta_1 f_1 x_3(t) + \delta_2(1 - f_2 - f_3)x_5(t) - [\omega n + \gamma] x_4^\sigma(t), \\
x_5^\Delta(t) = \delta_1(1 - f_1)x_3(t) + \eta(1 - k)x_6(t) - [\delta_2(1 - f_2 - f_3) + \delta_2 f_2 + \alpha_1 f_3 + \gamma] x_5^\sigma(t), \\
x_6^\Delta(t) = \delta_2 f_2 x_5(t) - [\eta(1 - k) + \alpha_2 k + \gamma] x_6^\sigma(t),
\end{cases} \quad (3.3)$$

subject to

$$x_1(0) = 10283785, \quad x_2(0) = 13, \quad x_3(0) = 2, \quad x_4(0) = 0, \quad x_5(0) = 0, \quad x_6(0) = 0,$$

where  $\Lambda = \frac{22614}{53}$ ,  $\omega = 1/31$ ,  $n = 0.075$ ,

$$\lambda(t) = \frac{\beta(l_A x_2(t) + x_3(t) + l_H x_5(t))}{N(t)}$$




FIGURE 1. Example 7: solution of (3.3) during 7 days.

with  $\beta = 1.93$ ,  $l_A = 1$ ,  $l_H = 0.1$ , and  $N(t) = \sum_{i=1}^6 x_i(t)$ ,  $p = 0.68$ ,  $\phi = 1/12$ ,  $\gamma = \frac{47833615}{N_0}$  with  $N_0 = N(0)$ ,  $q = 0.15$ ,  $\nu = 1/15$ ,  $\delta_1 = 1/3$ ,  $\delta_2 = 1/3$ ,  $f_1 = 0.96$ ,  $f_2 = 0.21$ ,  $f_3 = 0.03$ ,  $\eta = 1/7$ ,  $k = 0.03$ ,  $\alpha_1 = 1/7$ , and  $\alpha_2 = 1/15$ .

System (3.3) is permanent with  $\lambda^L = 1.876738171 \times 10^{-7}$ ,  $\lambda^U = 1.93$ ,  $M_1 = 65.83271997$ ,  $M_2 = 8.722416333$ ,  $M_3 = 0.017498412509$ ,  $M_4 = 4.428264471$ ,  $M_6 = 0.02788991356$ ,  $M = \max_{i=1, \dots, 6} (M_i) = M_1$ ,  $m_1 = 58.16800031$ ,  $m_2 = 7.7068897$ ,  $m_3 = 0.0154611$ ,  $m_4 = 0.7093454$ ,  $m_5 = 0.0000414$ ,  $m_6 = 6.0482208 \times 10^{-7}$ , and  $m = \min_{i=1, \dots, 6} (m_i) = m_6$ . In addition, the conditions of Theorem 6 are verified and we have

$$4.653775371 = A > B = 4.148857053, \quad \psi = 0.0505839, \quad 1 - \psi\mu(t) = 0.94941603 > 0.$$

We conclude that system (3.3) has a unique positive almost periodic solution, which is uniformly asymptotic stable. This is illustrated in Figure 1.

## ACKNOWLEDGMENTS

This work is part of first authors' PhD project. Torres is funded by the Fundação para a Ciência e a Tecnologia, I.P. (FCT, Funder ID = 50110000187) under grants UIDB/04106/2020 and UIDP/04106/2020 and the project 2022.03091.PTDC "Mathematical Modelling of Multi-scale Control Systems: applications to human diseases" (CoSysM3), financially supported by national funds (OE) through FCT/MCTES.

## REFERENCES

- [1] P. Agarwal, J. J. Nieto, M. Ruzhansky and D. F. M. Torres, Analysis of Infectious Disease Problems (Covid-19) and Their Global Impact, Infosys Science Foundation Series in Mathematical Sciences, Springer, Singapore, 2021.
- [2] P. Agarwal, J. J. Nieto and D. F. M. Torres, Mathematical Analysis of Infectious Diseases, Academic Press, London, UK, 2022.
- [3] D. I. Baleanu, H. Mohammadi and S. Rezapour, A fractional differential equation model for the COVID-19 transmission by using the Caputo-Fabrizio derivative, Adv. Difference Equ. **2020**, Paper No. 299, 27 pp.
- [4] M. Bohner and A. Peterson, *Dynamic equations on time scales*, Birkhäuser Boston, Boston, MA, 2001.
- [5] M. Bohner and A. Peterson, *Advances in dynamic equations on time scales*, Birkhäuser Boston, Inc., Boston, MA, 2003.
- [6] M. J. Bohner, S. H. Streipert and D. F. M. Torres, Exact solution to a dynamic SIR model, Nonlinear Anal. Hybrid Syst. **32** (2019), 228–238.
- [7] M. J. Bohner, J. Heim and A. L. Liu, Solow models on time scales, Cubo **15** (2013), no. 1, 13–31.
- [8] M. Hu and L. Wang, Dynamic inequalities on time scales with applications in permanence of predator-prey system, Discrete Dyn. Nat. Soc. **2012**, Art. ID 281052, 15 pp.
- [9] M. Khuddush and K. R. Prasad, Permanence and stability of multi-species nonautonomous Lotka-Volterra competitive systems with delays and feedback controls on time scales, Khayyam J. Math. **7** (2021), no. 2, 241–256.
- [10] Y. Kim, S. Lee, C. Chu, S. Choe, S. Hong and Y. Shin, The characteristics of middle eastern respiratory syndrome coronavirus transmission dynamics in South Korea, Osong Public Health Res. Perspect. **7** (2016), no. 1, 49–55.
- [11] A. P. Lemos-Paião, C. J. Silva and D. F. M. Torres, A new compartmental epidemiological model for COVID-19 with a case study of Portugal, Ecological Complexity **44** (2020), Art. 100885, 8 pp.
- [12] W.-T. Li and L. L. Wang, Existence and global attractivity of positive periodic solutions of functional differential equations with feedback control, J. Comput. Appl. Math. **180** (2005), no. 2, 293–309.
- [13] K. Shah, M. Arfan, W. Deebani, M. Shutaywi and D. Baleanu, Investigation of COVID-19 mathematical model under fractional order derivative, Math. Model. Nat. Phenom. **16** (2021), Paper No. 50, 14 pp.
- [14] K. R. Prasad and M. Khuddush, Stability of positive almost periodic solutions for a fishing model with multiple time varying variable delays on time scales, Bull. Int. Math. Virtual Inst. **9** (2019), no. 3, 521–533.
- [15] K. R. Prasad and M. Khuddush, Existence and uniform asymptotic stability of positive almost periodic solutions for three-species Lotka-Volterra competitive system on time scales, Asian-Eur. J. Math. **13** (2020), no. 3, 2050058, 24 pp.
- [16] C. C. Tisdell and A. H. Zaidi, Basic qualitative and quantitative results for solutions to nonlinear, dynamic equations on time scales with an application to economic modelling, Nonlinear Anal. **68** (2008), no. 11, 3504–3524.
- [17] H. T. Zhang and Y. Li, Almost periodic solutions to dynamic equations on time scales, J. Egyptian Math. Soc. **21** (2013), no. 1, 3–10.
- [18] Y. Li and C. Wang, Almost periodic solutions to dynamic equations on time scales and applications, J. Appl. Math. **2012**, Art. ID 463913, 19 pp.

(N. Zine) UNIVERSITY MUSTAPHA STAMBOULI OF MASCARA, ALGERIA  
*Email address:* [nadjoua.zine@univ-mascara.dz](mailto:nadjoua.zine@univ-mascara.dz)

(B. Bayour) UNIVERSITY MUSTAPHA STAMBOULI OF MASCARA, ALGERIA  
*Email address:* [b.bayour@univ-mascara.dz](mailto:b.bayour@univ-mascara.dz)

(D. F. M. Torres) CENTER FOR RESEARCH AND DEVELOPMENT IN MATHEMATICS AND APPLICATIONS (CIDMA), DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AVEIRO, 3810-193 AVEIRO, PORTUGAL  
*Email address:* [delfim@ua.pt](mailto:delfim@ua.pt)

(D. F. M. Torres) RESEARCH CENTER IN EXACT SCIENCES (CICE), FACULTY OF SCIENCES AND TECHNOLOGY (FCT), UNIVERSITY OF CAPE VERDE (UNI-CV), 7943-010 PRAIA, CAPE VERDE  
*Email address:* [delfim@unicv.cv](mailto:delfim@unicv.cv)