# ISOMORPHISMS BETWEEN LATTICES OF HYPERINVARIANT SUBSPACES

David Mingueza Nestlé Spain Esplugas de Llobregat david.mingueza@outlook.es M.Eulàlia Montoro Departament de Matemàtiques i Informàtica Universitat de Barcelona Barcelona eula.montoro@ub.edu

Alicia Roca Departamento de Matemática Aplicada, IMM Universitat Politècnica de València València aroca@mat.upv.es

# ABSTRACT

Given two nilpotent endomorphisms, we determine when their lattices of hyperinvariant subspaces are isomorphic. The study of the lattice of hyperinvariant subspaces can be reduced to the nilpotent case when the endomorphism has a Jordan-Chevalley decomposition; for example, it occurs if the underlying field is the field of complex numbers.

Keywords Hyperinvariant subspaces · Isomorphism of lattices.

# 1 Introduction

Three main lattices of subspaces are naturally associated to endomorphisms, the lattices of invariant subspaces, of characteristic subspaces and of hyperinvariant subspaces. A vector subspace  $V \subseteq \mathbb{F}^n$  is invariant with respect to the endomorphism f of  $\mathbb{F}^n$  if  $f(V) \subset V$ , the subspace is characteristic if it is invariant for every automorphism commuting with f, and it is hyperinvariant if it is invariant for every endomorphism commuting with f.

It is well-known when two lattices of invariant subspaces are isomorphic ([1]), and isomorphisms of lattices of characteristic subspaces is still an open subject of study. In this paper we characterize when two lattices of hyperinvariant subspaces are isomorphic. This problem was analyzed in P. Y. Wu, "Which Linear Transformations Have Isomorphic Hyperinvariant Subspace Lattices?" ([5]). When studying the paper we realized that, although the statement of the main theorem is correct, some of the previous lemmas, theorems, or its proofs, were not accurate, therefore they need some reformulation and, as a consequence, the inclusion of the analysis of some missed cases. Hence, this work can be regarded as a review of [5].

The paper is organized as follows. In Section 2 we state some basic definitions and properties of lattices, in particular of lattices of invariant subspaces, and of the algebra of the centralizer. In Section 3 we describe the structure and some specific properties of the lattice of hyperinvariant subspaces, wich will be used later on. Also, some necessary conditions for two lattices to be isomorphic are recalled. In Section 4, some special chains of hyperinvariant subspaces are presented, which will be a key tool to study the problem. In Section 5 the special chains are widely used, among other properties, to study all possible isomorphisms of hyperinvariant lattices, in Subsection 5.1 when the lattices have different Segre characteristic length, and in Subsection 5.2 when they have the same Segre characteristic length. Finally, in Subsection 5.3 we summarize the results in a theorem.

# 2 Preliminaries

In this section we recall some results related to endomorphisms on complex vector spaces. We assimilate an endomorphism with its matrix representation.

A partially ordered set L such that every pair of elements  $x, y \in L$  has a join  $(x \vee y)$  and a meet  $(x \wedge y)$  is called a lattice. A mapping between two lattices  $L_1$  and  $L_2$ ,  $f : L_1 \to L_2$ , is a lattice homomorphism if it preserves joints and meets. If f is an isomorphism we say that  $L_1$  and  $L_2$  are isomorphic and it is denoted by  $L_1 \simeq L_2$ . If a lattice L is finite, we denote by card(L) the number of elements of L.

Given  $a, b \in L$  with  $a \ge b$ , the set formed by  $\{x \in L : b \le x \le a\}$  is a sublattice of L and is called the quotient sublattice a/b.

An equivalent relation ( $\sim$ ) on a lattice L is a *congruence* relation if given  $a, b \in L$ , then

$$a \sim b \Rightarrow \begin{cases} (a \lor c) \sim (b \lor c), \\ (a \land c) \sim (b \land c), \end{cases} \text{ for every } c \in L.$$

$$\tag{1}$$

We denote by  $[a] = \{x \in L : x \sim a\}$  the class of elements related to a, and by  $L/\sim = \{[a] : a \in L\}$  the quotient lattice formed by all the classes in L. Moreover,

$$a \sim b \Rightarrow \forall c \in (a \lor b)/(a \land b), c \sim a.$$
 (2)

Let  $f: L \longrightarrow L_1$  be a lattice homomorphism. The congruence relation defined by  $x \sim y$  if f(x) = f(y) is called the *kernel* of f (ker f). The following isomorphism theorem is satisfied.

**Theorem 2.1.** ([3, Theorem 1.5]) Let L and  $L_1$  be lattices. Let  $f : L \to L_1$  be an homomorphism onto. Then,  $L/\ker f$  is isomorphic to  $L_1$ .

Let  $\mathbb{C}^{n \times n}$  be the algebra of  $n \times n$  matrices with entries in the complex field  $\mathbb{C}$ . Given  $A \in \mathbb{C}^{n \times n}$ , the algebra of the matrices that commute with A, i.e., the centralizer of A, denoted Z(A), is

$$Z(A) = \{ B \in \mathbb{C}^{n \times n} : AB = BA \},\$$

the lattice of invariant subspaces of A, denoted Inv(A), is

$$Inv(A) = \{ F \subset \mathbb{C}^{n \times 1} : AF \subseteq F \},\$$

and the lattice of hyperinvariant subspaces of A, denoted Hinv(A), is

$$\operatorname{Hinv}(A) = \{ F \subset \mathbb{C}^{n \times 1} : BF \subseteq F, \forall B \in Z(A) \}.$$

As mentioned before, isomorphisms of lattices of invariant subspaces are characterized. The result is stated in the next theorem.

**Theorem 2.2.** ([1]) Let  $A, B \in \mathbb{C}^{n \times n}$ . Then, the following statements are equivalent:

- 1. Inv(A) and Inv(B) are isomorphic.
- 2. A and B have the same Jordan structure.
- *3.* Z(A) and Z(B) are isomorphic.

The following theorem gives a sufficient condition for two lattices of hyperinvariant subspaces to be isomorphic. In the next section we will see that it is not necessary.

**Theorem 2.3.** ([4]) Let  $A, B \in \mathbb{C}^{n \times n}$ . If  $Inv(A) \simeq Inv(B)$ , then  $Hinv(A) \simeq Hinv(B)$ .

## **3** The lattice of hyperinvariant subspaces

The study of the lattice of hyperinvariant subspaces can be reduced to the nilpotent case when the endomorphism has a Jordan-Chevalley decomposition (see [2]). As the reduction applies on the complex field, throughout the paper we will assume that  $A \in \mathbb{C}^{n \times n}$  is nilpotent.

Let  $A \in \mathbb{C}^{n \times n}$  be a nilpotent matrix and  $\alpha = (\alpha_1, \ldots, \alpha_r) \in \mathbb{N}^r$  its Segre characteristic, where  $\alpha_1 \ge \ldots \ge \alpha_r > 0$ .

Let  $u = (u_1, \ldots, u_r) \in \mathbb{N}^r$  such that

$$u_1 \ge \ldots \ge u_r \ge 0,\tag{3}$$

$$\alpha_1 - u_1 \ge \ldots \ge \alpha_r - u_r \ge 0, \tag{4}$$

and define

$$V(\alpha_1,\ldots,\alpha_r) = \{ u \in \mathbb{N}^r : \text{ satisfying } (3) \text{ and } (4) \}.$$

Given  $u, v \in V(\alpha_1, \ldots, \alpha_r)$ , we define the following partial order:

 $u \subset v$  if  $u_i \leq v_i, i = 1, \ldots, r$ .

**Proposition 3.1.** ([2])  $V(\alpha_1, \ldots, \alpha_r)$  with the partial order relation " $\subset$ " is a lattice. The meet and the join are defined, respectively, as

$$u \wedge v = (\min(u_1, v_1), \dots, \min(u_r, v_r)),$$
$$u \vee v = (\max(u_1, v_1), \dots, \max(u_r, v_r)).$$

The elements of  $V(\alpha_1, \ldots, \alpha_r)$  are called *hypertuples*, and we refer to the lattice  $V(\alpha_1, \ldots, \alpha_r)$  as a *hyperlattice*. **Proposition 3.2.** ([2]) Given  $A \in \mathbb{C}^{n \times n}$  a nilpotent matrix with Segre Characteristic  $(\alpha_1, \ldots, \alpha_r)$ , there exists a lattice isomorphism

$$f : \operatorname{Hinv}(A) \longrightarrow V(\alpha_1, \ldots, \alpha_r).$$

As a consequence of this proposition, we identify the hyperinvariant subspaces with their corresponding hypertuples. If there is no possible confusion, given a Segre characteristic  $\alpha = (\alpha_1, \ldots, \alpha_r)$  we will also denote the corresponding hyperlattice by  $V(\alpha)$ .

Unlike the lattice of invariant subspaces Inv(A), for finite dimensional matrices the lattice Hinv(A) is always finite, and its cardinality is known; we recall it in the next theorem.

**Theorem 3.3.** ([2]) The number of hyperinvariant subspaces of a nilpotent matrix  $A \in \mathbb{C}^{n \times n}$  with Segre characteristic  $\alpha = (\alpha_1, ..., \alpha_r)$  is

$$(\alpha_1 - \alpha_2 + 1) \dots (\alpha_{r-1} - \alpha_r + 1)(\alpha_r + 1)$$

From now on, we will only consider the reduced case of Segre characteristics, i.e., we will assume that  $\alpha_1 > ... > \alpha_r$ , because there exists a lattice isomorphism between the hyperlattices of the non reduced and the reduced cases (see [5]).

**Remark 3.4.** The condition of Theorem 2.3 is not necessary. For example, let  $A, B \in \mathbb{C}^{n \times n}$  be nilpotent matrices with Segre characteristics  $\alpha = (l, l-1)$  and  $\beta = (2l-1), l > 0$ , respectively. As we can see in the following diagram  $V(\alpha) \simeq V(\beta)$ ,

(l, l - 1)	(2l - 1)
(l-1, l-1)	(2l - 2)
(l-1, l-2)	(2l - 3)
:	:
(1,0)	(1)
(0,0)	(0)

but the corresponding lattices of invariant subspaces are not isomorphic (there exist at last two different invariant subspaces of A of dimension 1, whilst there exists only one of B).

Our aim is to characterize when two lattices of hyperinvariant subspaces are isomorphic. The general scheme of this paper follows that of [5], but as announced in the Introduction, some results there were not accurate, either incomplete or defective. We fix here those flaws either completing the results or providing new proofs. Some concepts must be introduced.

We start by defining the notion of "son" of an element of a hyperlattice.

**Definition 3.5.** Given  $u = (u_1, \ldots, u_r) \in V(\alpha_1, \ldots, \alpha_r)$ , we say that  $v = (v_1, \ldots, v_r)$  is a *son* of u if there exists  $j \in \{1, \ldots, r\}$  such that  $v_i = u_i$  for  $i \in \{1, \ldots, r\} \setminus \{j\}, v_j = u_j - 1$ , and  $v \in V(\alpha_1, \ldots, \alpha_r)$ .

We denote by Son(u) the set of sons of u. Notice that if v is a *son* of u then  $v \subset u$  and there is no any  $w \in V(\alpha_1, \ldots, \alpha_r)$  such that  $v \subsetneq w \subsetneq u$ . Whenever  $v \in Son(u)$  we will say that u is a *father* of v, and if  $v, w \in Son(u)$  we will say that v and w are *brothers*.

**Definition 3.6.** Given  $V(\alpha_1, \ldots, \alpha_r)$ , a *chain* C of length t from  $w_1$  to  $w_t$  is a sequence of hypertuples  $w_1, \ldots, w_t \in V(\alpha_1, \ldots, \alpha_r)$  such that  $w_{i+1} \in \text{Son}(w_i)$ ,  $i = 1, \ldots, t-1$ . We will write the chain C as  $w_1 - \cdots - w_t$ .

In some occasions we will reorder the chain in the reverse way; i.e., a chain  $w_1 - \cdots - w_t$  from  $w_1$  to  $w_t$  will satisfy that  $w_i \in Son(w_{i+1}), i = 1, \dots, t-1$ .

The following properties are satisfied.

**Proposition 3.7.** Let  $f: V(\alpha) \longrightarrow V(\beta)$  be a hyperlattice isomorphism, and  $u, v \in V(\alpha)$ . Then,

- 1. If  $v \subset u$ , then  $f(v) \subset f(u)$ .
- 2. If  $v \in Son(u)$ , then  $f(v) \in Son(f(u))$ . As a consequence, if C is a chain in  $V(\alpha)$  of length t, its image f(C) is a chain in  $V(\beta)$  of length t.
- 3. If  $Son(u) = \{v\}$ , then  $Son(f(u)) = \{f(v)\}$ .
- *Proof.* 1. Observe that  $v \lor u = u$ . As f preserve joins,  $f(v) \lor f(u) = f(u)$ , hence  $f(v) \subset f(u)$ .
  - If v ∈ Son(u), then f(v) ⊂ f(u). Assume that f(v) is not a son of f(u). It means that there exists w ∈ V(β) such that f(v) ⊊ w ⊊ f(u). Then, v ⊊ f<sup>-1</sup>(w) ⊊ u, which implies that v ∉ Son(u).
  - 3. We know that  $f(v) \in \text{Son}(f(u))$ . If there exists  $w \in \text{Son}(f(u))$ ,  $w \neq f(v)$ , then  $f^{-1}(w) \in \text{Son}(u)$  and  $f^{-1}(w) \neq v$ , which is a contradiction.

In the following lemma we characterize the sons of a given hypertuple.

**Lemma 3.8.** Let  $V(\alpha_1, \ldots, \alpha_r)$  be a hyperlattice and  $u = (u_1, \ldots, u_r) \in V(\alpha)$ . Then:

- 1.  $(u_1 1, u_2, ..., u_r) \in Son(u)$  if and only if  $u_1 > u_2$ .
- 2. Given  $i \in \{2, ..., r-1\}$ , then  $(u_1, ..., u_i 1, ..., u_r) \in Son(u)$  if and only if  $u_i > u_{i+1}$  and  $\alpha_{i-1} u_{i-1} > \alpha_i u_i$ .
- 3.  $(u_1, u_2, ..., u_r 1) \in Son(u)$  if and only if  $u_r \neq 0$  and  $\alpha_{r-1} u_{r-1} > \alpha_r u_r$ .

*Proof.* Let  $u \in V(\alpha)$ . Notice that subtracting one from a single component of u, if the resulting tuple belongs to  $V(\alpha)$ , we obtain a son. Therefore,

- 1.  $(u_1 1, u_2, \dots, u_r) \in V(\alpha)$  if and only if  $u_1 1 \ge u_2$  and  $\alpha_1 (u_1 1) \ge \alpha_2 u_2$ . As  $u \in V(\alpha)$ , the two conditions are equivalent to  $u_1 > u_2$ .
- 2. Given  $i \in \{2, \ldots, r-1\}$ , then  $(u_1, \ldots, u_i 1, \ldots, u_r) \in V(\alpha)$  if and only if  $u_{i-1} \ge u_i 1 \ge u_i$  and  $\alpha_{i-1} u_{i-1} \ge \alpha_i (u_i 1) \ge \alpha_i u_i$ . As  $u \in V(\alpha)$ , these conditions are equivalent to  $u_i > u_{i+1}$  and  $\alpha_{i-1} u_{i-1} > \alpha_i u_i$ .
- 3.  $(u_1, u_2, \dots, u_r 1) \in Son(u)$  if and only if  $u_r > 0$ ,  $u_{r-1} \ge u_r 1$  and  $\alpha_{r-1} u_{r-1} \ge \alpha_r (u_r 1)$ . As  $u \in V(\alpha)$ , these conditions are equivalent to that  $u_r \ne 0$  and  $\alpha_{r-1} - u_{r-1} > \alpha_r - u_r$ .

**Example 3.9.** Let  $u = (3, 2, 1), v = (4, 2, 1) \in V(5, 3, 1)$ . Then,

$$Son(u) = \{(2, 2, 1), (3, 1, 1), (3, 2, 0)\}, Son(v) = \{(3, 2, 1), (4, 2, 0)\}$$

**Corollary 3.10.** Let  $u = (u_1, \ldots, u_r) \in V(\alpha)$ . Then,

 $Son(u) = \{\emptyset\} \iff if and only if u = (0, \dots, 0).$ 

As a consequence of Corollary 3.10 and Proposition 3.7, item 2, we obtain the following result.

**Corollary 3.11.** *1. Given*  $u = (u_1, \ldots, u_r) \in V(\alpha_1, \ldots, \alpha_r)$ , there exists a chain C from u to 0  $(C : u - \cdots - 0)$  of length  $t = u_1 + \cdots + u_r + 1$ .

2. Let  $f: V(\alpha_1, \ldots, \alpha_r) \longrightarrow V(\beta_1, \ldots, \beta_s)$  be a hyperlattice isomorphism, and  $v = f(u), u \in V(\alpha)$ . Then,  $u_1 + \cdots + u_r = v_1 + \cdots + v_s$ .

**Remark 3.12.** If the hyperlattices  $V(\alpha_1, \ldots, \alpha_r)$  and  $V(\beta_1, \ldots, \beta_s)$  are isomorphic, then the following two conditions must be satisfied:

- 1. Dimension condition:  $\alpha_1 + \cdots + \alpha_r = \beta_1 + \cdots + \beta_s$ .
- 2. Cardinality condition:  $\operatorname{card}(V(\alpha_1,\ldots,\alpha_r)) = \operatorname{card}(V(\beta_1,\ldots,\beta_s)).$

Along the paper we will frequently refer to these two conditions.

## 4 Special Chains and Riding Special Chains

Useful tools to study when two hyperlattices are isomorphic, are the notions of "special chains" and "riding special chains". To define them we need to characterize when a hypertuple has a unique son.

**Theorem 4.1.** Let  $u \in V(\alpha_1, \ldots, \alpha_r)$ . Let  $k = \max\{i : u_1 = \cdots = u_i\}$  and  $q = \max\{i : u_i > 0\}$ . Then,  $\operatorname{cord}(\operatorname{Sop}(u)) = 1$  if and only if  $\alpha_i = u_i$  is a set of i = 1.

$$\operatorname{card}(\operatorname{Son}(u)) = 1$$
 if and only if  $\alpha_k - u_k = \alpha_{k+1} - u_{k+1} = \ldots = \alpha_q - u_q$ 

In such a case,

$$Son(u) = \{(u_k, \dots, u_k - 1, u_{k+1}, \dots, u_q, 0, \dots, 0)\}.$$

*Proof.* If k = r the result is trivial. If k < r, we have  $u_k > u_{k+1}$ , and

$$\bar{u} = (u_k, \dots, u_k - 1, u_{k+1}, \dots, u_q, 0 \dots, 0) \in Son(u).$$

Let  $\operatorname{card}(\operatorname{Son}(u)) = 1$ . If k = q, the result is trivial. If k < q, assume that there exists  $k \leq j < q$  such that  $\alpha_j - u_j > \alpha_{j+1} - u_{j+1} = \ldots = \alpha_q - u_q$ . Then, by Lemma 3.8

$$\bar{\bar{u}} = (u_k, \dots, u_k, \dots, u_{j+1} - 1, \dots, u_q, 0, \dots, 0) \in \operatorname{Son}(u)$$

As  $\overline{\overline{u}} \neq \overline{u}$ , u has more than one son, which is a contradiction.

Conversely. Assume that u has a second son  $v = (v_1, \ldots, v_r)$ . It means that there exists only one index  $1 \le j \le r$  such that  $v_j = u_j - 1$ . Then,  $k \le j \le q$   $(j \ge k$ , otherwise  $v_j = u_j - 1 < u_{j+1} = v_{j+1}$ . Also,  $j \le q$ , otherwise  $u_j = 0$ ). If  $k < j \le q$ , then,

$$\alpha_{j-1} - u_{j-1} = \alpha_{j-1} - v_{j-1} \ge \alpha_j - v_j = \alpha_j - (u_j - 1) > \alpha_j - u_j,$$

which by hypothesis cannot occur. Therefore, j = k and  $v = \bar{u}$ .

**Example 4.2.** Let k and q be defined as in the previous theorem.

- 1. Let  $\alpha = (7, 3, 1)$ . Then,  $Son(2, 2, 0) = \{(2, 1, 0)\}$ . Here k = q = 2.
- 2. Let  $\alpha = (7, 5, 3, 1)$ . Then, Son $(3, 3, 1, 0) = \{(3, 2, 1, 0)\}$ . Here k = 2 and q = 3.
- 3. Let  $\alpha = (7, 5, 4, 3)$ . Then, Son $(3, 3, 2, 1) = \{(3, 2, 2, 1)\}$ . Here k = 2 and q = 4.
- 4. Notice that  $Son(\alpha_1, \ldots, \alpha_r) = \{(\alpha_1 1, \ldots, \alpha_r)\}$ . Here k = 1 and q = r.

Isomorphisms of hyperinvariant lattices

#### 4.1 Special Chains

The fact that a hyperlattice isomorphism preserves the parenthood, i.e., the image of a son of a given hypertuple is the son of the image of the hypertuple, leads naturally to introduce the following definition.

**Definition 4.3.** A chain  $C : w_1 - w_2 - \cdots - w_t$  in  $V(\alpha)$  is called *special* if it is of maximal length satisfying the property that  $Son(w_i) = \{w_{i+1}\}, i = 1, \dots, t-1$ .

We focus on special chains whose last element is zero. This subsection is devoted to determine the different types of special chains a hyperlattice may have. The results obtained are essentially consequence of Theorem 4.1.

Observe that  $V(\alpha_1)$  is a special chain ending at zero. We analyze next  $V(\alpha_1, \alpha_2)$ .

**Lemma 4.4.** Let  $V(\alpha_1, \alpha_2)$  a hyperlattice.

• If  $\alpha_1 - \alpha_2 = 1$ , then  $V(\alpha_1, \alpha_2)$  is the special chain ending at zero:

$$(\alpha_1, \alpha_2) - (\alpha_1 - 1, \alpha_2) - \dots - (1, 0) - (0, 0).$$

• If  $\alpha_1 - \alpha_2 > 1$ , then there exists only two special chains ending at zero:

$$(1,1) - (1,0) - (0,0),$$
 (5)

and

$$(\alpha_1 - \alpha_2, 0) - \dots - (0, 0).$$
 (6)

*Proof.* If  $\alpha_1 - \alpha_2 = 1$ , the conclusion is immediate. Assume that  $\alpha_1 - \alpha_2 > 1$ . Let  $u = (u_1, u_2) \in V(\alpha_1, \alpha_2)$ . If  $u_1 > u_2 > 0$ , then, by Theorem 4.1,  $Son(u) = \{(u_1 - 1, u_2)\}$  if and only if  $\alpha_1 - u_1 = \alpha_2 - u_2$ , and since  $\alpha_1 - \alpha_2 > 1$ ,

$$Son(Son(u)) = \{(u_1 - 2, u_2), (u_1 - 1, u_2 - 1)\},\$$

therefore u cannot be in a special chain ending at zero.

By Theorem 4.1, and following a similar argument, if  $u_1 = u_2 > 0$ , then  $(u_1, u_1)$  is in a special chain ending at zero if and only if  $u_1 = 1$ , and the special chain is (5).

Finally, if  $u_2 = 0$ , then  $(u_1, 0)$  is always in the special chain ending at zero (6).

In the following results k and q are defined as in Theorem 4.1.

**Lemma 4.5.** Let  $V(\alpha_1, \ldots, \alpha_r)$  be a hyperlattice with  $r \ge 3$ . A hypertuple of the form  $u = (u_k, \ldots, u_k, u_{k+1}, \ldots, u_q, 0, \ldots, 0), 1 < k < q$ , cannot be in any special chain ending at zero.

*Proof.* By Theorem 4.1, 
$$\operatorname{card}(\operatorname{Son}(u)) = 1$$
 if and only if  $\alpha_k - u_k = \alpha_{k+1} - u_{k+1} = \cdots = \alpha_q - u_q$ , and in that case  
 $\operatorname{Son}(u) = \{(u_1, \dots, u_k - 1, u_{k+1}, \dots, u_q, 0, \dots, 0)\}.$ 

Observe that  $\operatorname{card}(\operatorname{Son}(\operatorname{Son}(u)) = 1$  if and only if  $\alpha_{k-1} - u_{k-1} = \alpha_k - (u_k - 1) = \alpha_{k+1} - u_{k+1} = \cdots = \alpha_q - u_q$ , which is not true. Therefore, the unique-son chain stops at  $\operatorname{Son}(u)$  and does not end at zero.

**Lemma 4.6.** Let  $V(\alpha_1, \ldots, \alpha_r)$ ,  $r \geq 2$ . Then,

$$(1, 1, \dots, 1) - (1, \dots, 1, 0) - \dots - (0, \dots, 0).$$
 (7)

is a special chain ending at zero.

*Proof.* It is immediate that the chain (7) is a unique-son chain. It cannot be enlarged because the hypertuple (1, 1, ..., 1) has a brother

$$Son(2, 1, \dots, 1) = \{(1, 1, \dots, 1), (2, 1, \dots, 1, 0)\}\$$

**Lemma 4.7.** Let  $V(\alpha_1, \ldots, \alpha_r)$ ,  $r \ge 3$ , be a hyperlattice. A hypertuple  $u = (u_1, \ldots, u_q, 0, \ldots, 0) \in V(\alpha_1, \ldots, \alpha_r)$ ,  $q \le 2$ , belongs to a special chain ending at zero only in the following cases:

• If q = 1, then u belongs to the special chain ending at zero

$$(\alpha_1 - \alpha_2, 0..., 0) - (\alpha_1 - \alpha_2 - 1, 0..., 0), \dots - (1, 0..., 0) - (0, 0..., 0).$$
(8)

- If q = 2 and  $\alpha_1 \alpha_2 > 1$ , then  $u = (1, 1, 0, \dots, 0)$  and the special chain ending at zero is (7).
- If q = 2 and  $\alpha_1 \alpha_2 = 1$ , then u belongs to the special chain ending at zero

$$(\alpha_2 - \alpha_3 + 1, \alpha_2 - \alpha_3, 0, \dots, 0) - (\alpha_2 - \alpha_3, \alpha_2 - \alpha_3, 0, \dots, 0) - \dots \\ \dots - (1, 0, 0, \dots, 0) - (0, 0, 0, \dots, 0),$$
(9)

and if u = (1, 1, 0, ..., 0), then it also belongs to (7).

*Proof.* We analyze the possible special chains ending at zero containing  $u = (u_1, \ldots, u_q, 0, \ldots, 0)$  depending on the value of q.

If q = 1 it is easy to see that

$$(\alpha_1 - \alpha_2, 0..., 0) - (\alpha_1 - \alpha_2 - 1, 0..., 0), \dots - (1, 0..., 0) - (0, 0..., 0)$$

is a special chain ending at zero, and u belongs to it.

Let q = 2. Clearly, u = (1, 1, 0, ..., 0) belongs to (7).

If  $u_1 = u_2 > 1$ , then  $Son(u) = \{\bar{u} = (u_2, u_2 - 1, 0..., 0)\}$ . Moreover,  $card(Son(\bar{u})) = 1$  if and only if  $\alpha_1 - u_1 = \alpha_2 - (u_2 - 1)$  if and only if  $\alpha_1 = \alpha_2 + 1$ , and in such a case  $Son(\bar{u}) = \{\bar{u} = (u_2 - 1, u_2 - 1, 0..., 0)\}$ . Repeating the argument, we get

$$(u_2, u_2, 0..., 0) - (u_2, u_2 - 1, 0..., 0) - (u_2 - 1, u_2 - 1, 0..., 0) - \cdots$$
  
 $\cdots - (1, 0, 0..., 0) - (0, 0, 0..., 0).$ 

As  $\alpha_2 - u_2 \ge \alpha_3$ , the longest chain is obtained taking  $u_2 = \alpha_2 - \alpha_3$ , and as  $\alpha_1 = \alpha_2 + 1$ , we obtain the chain (9).

If  $u_1 > u_2$ , then  $\operatorname{Son}(u) = \{\bar{u} = (u_1 - 1, u_2, 0 \dots, 0)\}$  if and only if  $\alpha_1 - u_1 = \alpha_2 - u_2$ . In turn,  $\operatorname{card}(\operatorname{Son}(\bar{u})) = 1$  if and only if  $\alpha_1 - (u_1 - 1) = \alpha_2 - u_2$ , which is not true. Hence,  $u_1 - 1 = u_2$  and  $\operatorname{Son}(\bar{u}) = \{(u_1 - 1, u_2 - 1, 0 \dots, 0)\}$ . Notice that it implies that  $\alpha_1 = \alpha_2 + 1$ , and this case is the same as the previous one.

The analysis performed on special chains is summarized in the next theorem.

**Theorem 4.8.** In the hyperlattice  $V(\alpha_1, ..., \alpha_r)$ ,  $r \ge 2$ , there exist at most two of the three possible types of special chains ending at zero,  $C_1, C_2$  and  $C_3$ , of lengths  $r + 1, \alpha_1 - \alpha_2 + 1$  and  $2(\alpha_1 - \alpha_3)$ , respectively, defined as follows:

*Proof.* Observe that, as already seen,  $V(\alpha_1)$  is trivially a special chain ending at zero. For  $r \ge 2$ , as a consequence of Lemmas 4.4 to 4.7, if  $\alpha_1 - \alpha_2 > 1$  the only special chains in  $V(\alpha)$  are of type  $C_1$  and  $C_2$ , and if  $\alpha_1 - \alpha_2 = 1$ , they are of type  $C_1$  and  $C_3$  (we take  $\alpha_3 = 0$  when r = 2).

**Remark 4.9.** As mentioned, in a lattice there exist at most two special chains ending at 0. More precisely, for r = 1 there exists a single special chain  $C_2$  (observe that in this case the chain of type  $C_1$  is reduced to 1 - 0 and is included in  $C_2$ , therefore, 1 - 0 is not a special chain). If r = 2 and  $\alpha_1 - \alpha_2 = 1$ , then there exists only the special chain  $C_3$  (now  $C_3$  includes the chains of type  $C_2$  and  $C_1$ ). In any other case, there exist two special chains:  $C_1$  and  $C_2$  if  $r \ge 2$  and  $\alpha_1 - \alpha_2 = 1$  (in the last case  $C_1$  and  $C_3$  contain the chain of type  $C_2$ ).

#### Example 4.10.

- Example of a chain of type  $C_2$ : Let V(7, 3, 2, 1). Then, C : (4, 0, 0, 0) (3, 0, 0, 0) (2, 0, 0, 0) (1, 0, 0, 0) (0, 0, 0, 0) is a special chain of type  $C_2$ , of length $(C) = \alpha_1 \alpha_2 + 1 = 5$ .
- Example of a chain of type  $C_3$ : Let V(6, 5, 3, 2). Then, C: (3, 2, 0, 0) (2, 2, 0, 0) (2, 1, 0, 0) (1, 1, 0, 0) (1, 0, 0, 0) (0, 0, 0, 0), is a special chain of type  $C_3$ , of length(C) =  $2(\alpha_1 \alpha_3) = 6$ .
- Example of a chain of type  $C_1$ : Let V(5, 4, 3, 2). Then, C : (1, 1, 1, 1) (1, 1, 1, 0) (1, 1, 0, 0) (1, 0, 0, 0) (0, 0, 0, 0) is a special chain of type  $C_1$ , of length(C) = r + 1 = 5.

#### 4.2 Riding Special Chains

Special chains are key tools to study isomorphisms of hyperlattices, but we also need another type of chains associated to special chains.

**Definition 4.11.** Given  $p \in \{1, 2, 3\}$ , a chain  $RC_p : w_1 - w_2 - \cdots - w_l$  rides on a special chain  $C_p$  if it is of maximal length satisfying that  $\exists j \in \{1, \ldots, l-1\}$  such that  $Son(w_i) = \{w_{i+1}\}$  for  $1 \le i \le j$ , for  $j+1 \le i \le l-1$ ,  $Son(w_i) = \{w_{i+1}, w'_{i+1}\}$  with  $w'_{i+1} \in C_p$ , and  $Son(w_l) = \{w'_{l+1}\}$  with  $w'_{l+1} \in C_p$ . We will refer to  $RC_p$  as a riding special chain or as a riding chain on  $C_p$ .

To describe riding special chains in  $V(\alpha_1, \ldots, \alpha_r)$  we analyze separately the cases when r = 2, r = 3 and r > 3. **Proposition 4.12.** Let  $V(\alpha_1, \alpha_2)$  be a hyperlattice.

- *1.* Let  $\alpha_1 \alpha_2 > 1$ . Then, the riding chains are:
  - (a)  $RC_1$ :
    - If  $\alpha_2 > 1$ , then  $RC_1 : (2,2) (2,1) (2,0)$ .
    - If  $\alpha_2 = 1$  and  $\alpha_1 \neq 3$ , then  $RC_1 : (2,1) (2,0)$ .
    - If  $\alpha_2 = 1$  and  $\alpha_1 = 3$ , then  $RC_1 : (3,1) (2,1) (2,0)$ .
  - (b)  $RC_2: (\alpha_1 \alpha_2 + 1, 1) (\alpha_1 \alpha_2, 1) \ldots (1, 1).$
- 2. If  $\alpha_1 \alpha_2 = 1$ , then there exists a  $C_3$  chain, but it does not exists any  $RC_3$ .

*Proof.* 1. Let  $\alpha_1 - \alpha_2 > 1$ .

- (a) Existence of  $RC_1$  chains:
  - If  $\alpha_2 > 1$ , then  $Son(2,2) = \{(2,1)\}$ ,  $Son(2,1) = \{(2,0), (1,1)\}$  and  $Son((2,0)) = \{(1,0)\}$ . Moreover, observe that (2,2) has a unique father (3,2) and  $Son(3,2) = \{(3,1), (2,2)\}$ , therefore there exist a riding chain on  $C_1$  of length 3:

• The proof of the case  $\alpha_2 = 1$  and  $\alpha_1 > 3$  is analogous. Notice that  $Son(3,1) = \{(2,1), (3,0)\}$ :

• If  $\alpha_2 = 1$  and  $\alpha_1 = 3$ , the chain indicated cannot be enlarged, for  $V(3, 1) = C_1 \cup RC_1$ :

(b) Existence of a  $RC_2$  chain:  $Son(\alpha_1 - \alpha_2 + 1, 1) = \{(\alpha_1 - \alpha_2, 1)\}$ , for  $k = 0, ..., \alpha_1 - \alpha_2 - 2$  we have  $Son(\alpha_1 - \alpha_2 - k, 1) = \{(\alpha_1 - \alpha_2 - (k + 1), 1), (\alpha_1 - \alpha_2 - k, 0)\}$  with  $(\alpha_1 - \alpha_2 - k, 0) \in C_2$ , and for  $k = \alpha_1 - \alpha_2 - 1$ ,  $Son(1, 1) = \{(1, 0)\}$ . Moreover,

• if 
$$\alpha_2 > 1$$
, then  $(\alpha_1 - \alpha_2 + 1, 2) \in V(\alpha)$ , but  $Son(\alpha_1 - \alpha_2 + 1, 2) = \{(\alpha_1 - \alpha_2 + 1, 1), (\alpha_1 - \alpha_2, 2)\}$ :

$$\begin{array}{cccc} (\alpha_1 - \alpha_2 + 1, 2) \rightarrow & (\alpha_1 - \alpha_2 + 1, 1) \rightarrow & (\alpha_1 - \alpha_2, 1) \rightarrow & \dots & \rightarrow (1, 1) \\ \downarrow & & \downarrow & \dots & \downarrow \\ (\alpha_1 - \alpha_2, 2) & & (\alpha_1 - \alpha_2, 0) \rightarrow & \dots & \rightarrow (1, 0) & \rightarrow (0, 0) \end{array}$$

- if  $\alpha_2 = 1$ , then  $V(\alpha_1, 1) = C_2 \cup RC_2$ .
- 2. Let  $\alpha_1 \alpha_2 = 1$ . The consequence is immediate, for  $V(\alpha_1, \alpha_2) = C_3$ .

### **Proposition 4.13.** Let $V(\alpha_1, \alpha_2, \alpha_3)$ be a hyperlattice.

- *1.* Let  $\alpha_1 \alpha_2 > 1$ . Then, the riding chains are:
  - $RC_1$ : - If  $\alpha_2 - \alpha_3 > 1$ , then  $RC_1 : (2, 1, 1) - (2, 1, 0) - (2, 0, 0)$ . - If  $\alpha_2 - \alpha_3 = 1$  and  $\alpha_3 \ge 2$ , then  $RC_1$ : (2,2,2) - (2,2,1) - (2,1,1) - (2,1,0) - (2,0,0). - If  $\alpha_2 - \alpha_3 = 1$  and  $\alpha_3 = 1$ , then  $RC_1$ : (2,2,1) - (2,1,1) - (2,0,0).
  - $RC_2: (\alpha_1 \alpha_2 + 1, 1, 0) (\alpha_1 \alpha_2, 1, 0) \dots (1, 1, 0)$  is a  $RC_2$  of length  $(RC_2) = \alpha_1 \alpha_2 + 1$ , and if  $\alpha_1 \alpha_2 = 2$  and  $\alpha_2 \alpha_3 > 1$ , then  $(\alpha_1 \alpha_2, 2, 0) (\alpha_1 \alpha_2, 1, 0) \dots (1, 1, 0)$  is also a  $RC_2$  of the same length.
- 2. If  $\alpha_1 \alpha_2 = 1$ , the riding chains are:

• 
$$RC_1$$
:  
-  $If \alpha_2 - \alpha_3 > 1$ , then  $RC_1 : (2, 1, 1) - (2, 1, 0)$ .  
-  $If \alpha_2 - \alpha_3 = 1$  and  $\alpha_3 \ge 2$ , then  $RC_1$ :  
(2,2,2) - (2,2,1) - (2,1,1) - (2,1,0).  
-  $If \alpha_2 - \alpha_3 = 1$  and  $\alpha_3 = 1$ , then  $RC_1$ :  
(3,2,1) - (2,2,1) - (2,1,1) - (2,1,0).  
•  $RC_3 : (\alpha_1 - \alpha_3 + 1, \alpha_2 - \alpha_3 + 1, 1) - (\alpha_1 - \alpha_3, \alpha_2 - \alpha_3 + 1, 1) - (\alpha_1 - \alpha_3, \alpha_2 - \alpha_3, 1) - \dots - (1, 1, 1)$   
and length( $RC_3$ ) =  $2(\alpha_1 - \alpha_3)$ .

*Proof.* It is analogous to the proof of Proposition 4.12.

- 1. If  $\alpha_1 \alpha_2 > 1$ ,
  - Existence of  $RC_1$  chains:
    - If  $\alpha_2 \alpha_3 > 1$ , then (2,1,1) (2,1,0) (2,0,0) is a riding chain on  $C_1$ . Observe that it cannot be enlarged because  $(2,1,1) \in \text{Son}\{(2,2,1),(3,1,1)\}$ , but in both cases there are other sons  $(\text{Son}(2,2,1) = \{(2,1,1),(2,2,0)\}, \text{Son}(3,1,1) = \{(2,1,1),(3,1,0)\})$ , hence  $\text{length}(RC_1) = 3$ :

- If  $\alpha_2 \alpha_3 = 1$  and  $\alpha_3 \ge 2$ , then it is easy to check that the chain (2, 2, 2) (2, 2, 1) (2, 1, 1) (2, 1, 0) (2, 0, 0) is a riding chain on  $C_1$ .
- If  $\alpha_2 \alpha_3 = 1$  and  $\alpha_3 = 1$ , the chain (2, 2, 1) (2, 1, 1) (2, 1, 0) (2, 0, 0) is a riding chain on  $C_1$ .
- Existence of  $RC_2$  chains: the chain  $(\alpha_1 \alpha_2 + 1, 1, 0) (\alpha_1 \alpha_2, 1, 0) \ldots (1, 1, 0)$  rides on  $C_2$ . Notice that it cannot be enlarged:

If  $\alpha_2 - \alpha_3 > 1$ , the tuple  $(\alpha_1 - \alpha_2 + 1, 1, 0)$  has two fathers, each one of them with more than one son. Notice also that if, additionally,  $\alpha_1 - \alpha_2 > 2$ , then Son  $(\alpha_1 - \alpha_2, 2, 0) = \{(\alpha_1 - \alpha_2, 1, 0), (\alpha_1 - \alpha_2 - 1, 2, 0)\}$ . The behavior is sketched in the next scheme:

If  $\alpha_1 - \alpha_2 > 2$  and  $\alpha_2 - \alpha_3 > 1$ ,

But, if  $\alpha_1 - \alpha_2 = 2$  (and  $\alpha_2 - \alpha_3 > 1$ ), then we have Son  $(\alpha_1 - \alpha_2, 2, 0) = \{(\alpha_1 - \alpha_2 - 1, 2, 0)\}$ , therefore,  $(\alpha_1 - \alpha_2, 2, 0) - (\alpha_1 - \alpha_2, 1, 0) - \dots - (1, 1, 0)$  is also a riding chain on  $C_2$  of the same length.

If  $\alpha_1 - \alpha_2 = 2$ , then  $(\alpha_1 - \alpha_2 + 1, 1, 0)$  has only one father  $(\alpha_1 - \alpha_2 + 1, 1, 1)$ , but it has two sons, therefore the riding chain remains the same.

- 2. If  $\alpha_1 \alpha_2 = 1$ ,
  - Existence of  $RC_1$  chains: the proof is immediate.
  - Existence of a  $RC_3$  chain: albeit the proof is easy, we include a sketch of it when  $\alpha_3 > 1$ ,

$$(\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3} + 1, 2) \checkmark (\alpha_{1} - \alpha_{3} + 1, \alpha_{2} - \alpha_{3} + 1, 1) \downarrow (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3} + 1, 1) \downarrow (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3} + 1, 1) \downarrow (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 1) \rightarrow (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 0) \downarrow (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 1) \rightarrow (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 0) \downarrow (\alpha_{1} - \alpha_{3}, \alpha_{3} - \alpha_{3}, 0) \downarrow (\alpha_{1} -$$

If  $\alpha_3 = 1$ , then  $V(\alpha_1, \alpha_2, \alpha_3) = C_3 \cup RC_3$ .

	-	_	

The riding chains when r > 3 are described in the following proposition. **Proposition 4.14.** Let  $V(\alpha_1, ..., \alpha_r)$  be a hyperlattice with r > 3.

1. If  $\alpha_1 - \alpha_2 > 1$ , there exists a  $RC_1$  of length $(RC_1) = r$ :  $(2, 1, \dots, 1, 1) - (2, 1, \dots, 1, 0) - \dots - (2, 0, \dots, 0),$  (10) and a  $RC_2$  of length $(RC_2) = \alpha_1 - \alpha_2 + 1$ :

$$(\alpha_1 - \alpha_2 + 1, 1, 0, \dots, 0) - (\alpha_1 - \alpha_2, 1, 0, \dots, 0) - \dots - (1, 1, 0, \dots, 0).$$
(11)

2. If  $\alpha_1 - \alpha_2 = 1$ , there exists an  $RC_1$  of  $length(RC_1) = r - 1$ :

$$(2, 1, \dots, 1, 1) - (2, 1, \dots, 1, 0) - \dots - (2, 1, 0, \dots, 0),$$
 (12)

and a  $RC_3$  of length $(RC_3) = 2(\alpha_1 - \alpha_3)$ :

$$(\alpha_1 - \alpha_3 + 1, \alpha_2 - \alpha_3 + 1, 1, 0, 0, \dots, 0) - (\alpha_1 - \alpha_3, \alpha_2 - \alpha_3 + 1, 1, 0, \dots, 0) - \dots - (1, 1, 1, 0, \dots, 0).$$

$$(13)$$

Proof.

1. Let 
$$\alpha_1 - \alpha_2 > 1$$
. Observe that for  $i = 0, \ldots, r - 2$ ,

$$Son(2, 1, \stackrel{(r-i-1)}{\dots}, 1, 0, \stackrel{(i)}{\dots}, 0) = \{(2, 1, \stackrel{(r-i-2)}{\dots}, 1, 0, \stackrel{(i+1)}{\dots}, 0), (1, 1, \stackrel{(r-i-1)}{\dots}, 1, 0, \stackrel{(i)}{\dots}, 0)\},$$
(14)

and

$$Son(2, 0, \dots, 0) = \{(1, 0, \dots, 0)\}.$$

Moreover,

$$(2, 1, \dots, 1) \in \operatorname{Son}(2, 2, 1, \dots, 1) \cap \operatorname{Son}(3, 1, \dots, 1),$$

but in both cases it is not the unique son. Therefore, the chain (10) is a  $RC_1$  in  $V(\alpha_1, \ldots, \alpha_r)$  of length r. A sketch of the riding chain appears in the next scheme:

Concerning the chain (11), observe that:

Son
$$(\alpha_1 - \alpha_2 + 1, 1, 0, \dots, 0) = \{(\alpha_1 - \alpha_2, 1, 0, \dots, 0)\},\$$

and for  $k = 0, ..., \alpha_1 - \alpha_2 - 1$ ,

$$Son(\alpha_1 - \alpha_2 - k, 1, 0, \dots, 0) = \{(\alpha_1 - \alpha_2 - k - 1, 1, 0, \dots, 0), (\alpha_1 - \alpha_2 - k, 0, 0, \dots, 0)\}.$$

Moreover, if  $\alpha_2 - \alpha_3 > 1$ , then

$$(\alpha_1 - \alpha_2 + 1, 1, 0, \dots, 0) \in Son(\alpha_1 - \alpha_2 + 1, 2, 0, \dots, 0) \cap Son(\alpha_1 - \alpha_2 + 1, 1, 1, 0, \dots, 0).$$

Then, the chain (11) is a  $RC_2$  in  $V(\alpha_1, \ldots, \alpha_r)$  of length  $\alpha_1 - \alpha_2 + 1$ .

If  $\alpha_2 - \alpha_3 = 1$ , then  $(\alpha_1 - \alpha_2 + 1, 1, 0, \dots, 0)$  has a single father  $(\alpha_1 - \alpha_2 + 1, 1, 1, 0, \dots, 0)$ , but

$$Son(\alpha_1 - \alpha_2 + 1, 1, 1, 0, \dots, 0) = \{(\alpha_1 - \alpha_2 + 1, 1, 1, 0, \dots, 0), (\alpha_1 - \alpha_2 + 1, 1, 0, \dots, 0)\},\$$

therefore,  $(\alpha_1 - \alpha_2 + 1, 1, 1, 0, ..., 0)$  does not belong to the riding chain. Hence, the chain (11) is a  $RC_2$  in  $V(\alpha_1, ..., \alpha_r)$  of length  $\alpha_1 - \alpha_2 + 1$ . A scheme of the behaviour appears next:

$$(\alpha_{1} - \alpha_{2}, 2, 0, \dots, 0) \qquad (\alpha_{1} - \alpha_{2}, 1, 1, 0, \dots, 0) \qquad \uparrow \qquad (\alpha_{1} - \alpha_{2} + 1, 2, 0, \dots, 0) \qquad \land \qquad (\alpha_{1} - \alpha_{2} + 1, 1, 1, 0, \dots, 0) \qquad \land \qquad (\alpha_{1} - \alpha_{2} + 1, 1, 1, 0, \dots, 0) \qquad \land \qquad (\alpha_{1} - \alpha_{2}, 0, \dots, 0) \qquad \qquad (\alpha_{1} - \alpha_{$$

2. Assume that  $\alpha_1 - \alpha_2 = 1$ . Observe that for i = 0, ..., r - 3, (14) is satisfied, and Son $(2, 1, 0, ..., 0) = \{(1, 1, 0, ..., 0)\}.$ 

Since

$$(2, 1, \dots, 1), (2, 2, 1, \dots, 1, 0) \in \text{Son}(2, 2, 1, \dots, 1),$$

the chain (12) is a  $RC_1$  in  $V(\alpha_1, \ldots, \alpha_r)$  of length r - 1.

On the other hand,

$$Son(\alpha_1 - \alpha_3 + 1, \alpha_2 - \alpha_3 + 1, 1, 0, \dots, 0) = \{(\alpha_1 - \alpha_3, \alpha_2 - \alpha_3 + 1, 1, 0, \dots, 0)\},$$
  

$$Son(\alpha_1 - \alpha_3, \alpha_2 - \alpha_3 + 1, 1, 0, \dots, 0) = \{(\alpha_1 - \alpha_3, \alpha_2 - \alpha_3, 1, 0, \dots, 0)\},$$
  
for  $k = 1, \dots, \alpha_1 - \alpha_3 - 1,$   

$$Son(k + 1, k + 1, 1, 0, \dots, 0) = \{(k + 1, k, 1, 0, \dots, 0), (k + 1, k + 1, 0, \dots, 0)\},$$
  

$$Son(k + 1, k, 1, 0, \dots, 0) = \{(k, k, 1, 0, \dots, 0), (k + 1, k, 0, \dots, 0)\},$$
  

$$Son(1, 1, 1, 0, \dots, 0) = \{(1, 1, 0, \dots, 0), (k + 1, k, 0, \dots, 0)\},$$
  
and

and

$$\{ (\alpha_1 - \alpha_3 + 1, \alpha_2 - \alpha_3 + 1, 1, 0, \dots, 0), (\alpha_1 - \alpha_3, \alpha_2 - \alpha_3 + 1, 1, 1, 1, \dots, 0) \} = \\Son((\alpha_1 - \alpha_3 + 1, \alpha_2 - \alpha_3 + 1, 1, 1, 0, \dots, 0)),$$
therefore, (13) is a  $RC_3$  in  $V(\alpha_1, \dots, \alpha_r)$  of length  $2(\alpha_1 - \alpha_3)$ .

$$RC_{3} \qquad C_{3}$$

$$(\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 1, 0, \dots, 0) \qquad (\alpha_{1} - \alpha_{3}, \alpha_{1} - \alpha_{3}, 0, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 2, 0, \dots, 0) \qquad (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 1, 0, \dots, 0) \qquad (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 2, 0, \dots, 0) \qquad (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 2, 0, \dots, 0) \qquad (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 2, 0, \dots, 0) \qquad (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 2, 0, \dots, 0) \qquad (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 2, 0, \dots, 0) \qquad (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 2, 0, \dots, 0) \qquad (\alpha_{1} - \alpha_{3}, \alpha_{2} - \alpha_{3}, 2, 0, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3}, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{2} - \alpha_{3}, \alpha_{2} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - \alpha_{3} - 1, \dots, 0) \qquad (\alpha_{3} - \alpha_{3} - 1, \dots, 0)$$

**Remark 4.15.** This section could have been developed studying the hypertuple's relation of parenthood, due to the property of selfduality of hyperinvariant lattices ([2]). In that case, the components of a special chain will have only one parent, and they will end at  $u = (\alpha_1, \ldots, \alpha_r)$ . It is left to the reader to state the results with the parenthood relation.

# 5 Isomorphisms of hyperinvariant lattices

Now we have the tools to characterize isomorphisms of hyperlattices. As the lengths of the special chains and the riding chains must be preserved by isomorphisms, we will use them to disregard cases that do not meet this property.

We study first isomorphisms  $f: V(\alpha_1, \ldots, \alpha_r) \longrightarrow V(\beta_1, \ldots, \beta_s)$  with  $r \neq s$ , and afterwards when r = s. When analyzing the special chains or the riding special chains, we will denote by  $C_i, RC_i$  those in  $V(\alpha_1, \ldots, \alpha_r)$  and by  $C'_i, RC'_i$  those in  $V(\beta_1, \ldots, \beta_s)$ , for i = 1, 2, 3.

#### **5.1** Isomorphic hyperinvariant lattices when $r \neq s$ .

This section is essentially devoted to study isomorphisms of hyperlattices when  $r \neq s$ . We start with the following auxiliary lemma.

**Lemma 5.1.** There are no isomorphisms  $f: V(\alpha_1, \ldots, \alpha_r) \longrightarrow V(\beta_1, \ldots, \beta_s)$  with  $r \neq s$  such that  $f(C_1) = C'_1$ .

*Proof.*  $f(C_1) = C'_1$  implies r = s, which is a contradiction.

When r = 1 we obtain the next result.

**Proposition 5.2.**  $V(\alpha_1) \simeq V(\beta_1, ..., \beta_s)$  if and only if s = 1 and  $\alpha_1 = \beta_1$  or s = 2,  $\alpha_1 = 2\beta_1 - 1$  and  $\beta_2 = \beta_1 - 1$ .

*Proof.* If s = 1 the result is obvious. If s = 2, as  $V(\beta_1, \beta_2)$  must have only one special chain,  $\beta_1 - \beta_2 = 1$ . Since  $\alpha_1 = \beta_1 + \beta_2$ , from Remark 3.12 the conclusion is immediate. If s > 2, then  $V(\beta_1, \ldots, \beta_s)$  has always two special chains, therefore  $V(\alpha_1)$  and  $V(\beta_1, \ldots, \beta_s)$  cannot be isomorphic.

The sufficiency is immediate (see Remark 3.4).

The following proposition can be proved without using riding chains, but with them the proof becomes easier.

**Proposition 5.3.**  $V(\alpha_1, \alpha_2) \simeq V(\beta_1, \ldots, \beta_s)$ ,  $s \ge 3$ , if and only if  $\alpha = (5, 2)$  and  $\beta = (4, 2, 1)$ .

*Proof.* Let  $f: V(\alpha_1, \alpha_2) \to V(\beta_1, \dots, \beta_s)$  be an isomorphism.

If  $\alpha_1 - \alpha_2 = 1$  then, as a consequence of Proposition 5.2,  $V(\alpha_1, \alpha_2) \simeq V(2\alpha_1 - 1)$ , therefore we can assume that  $\alpha_1 - \alpha_2 > 1$ .

If  $\beta_1 - \beta_2 = 1$  then, by Lemma 5.1,  $f(C_1) = C'_3$ . It implies that  $3 = 2(\beta_1 - \beta_3)$ , which cannot occur. Hence,  $\beta_1 - \beta_2 > 1$ .

• Assume first that s = 3.

Taking into account Lemma 5.1,  $f(C_2) = C'_1$  and  $f(C_1) = C'_2$ . It means that  $\alpha_1 - \alpha_2 = 3$  and  $\beta_1 - \beta_2 = 2$ . As  $f(RC_2) = RC'_1$ , by Propositions 4.12 and 4.13 necessarily  $4 = \alpha_1 - \alpha_2 + 1 = \text{length}(RC_2) = \text{length}(RC'_1)$ , from where we conclude that  $\beta_2 - \beta_3 = \beta_3 = 1$ . Therefore,  $\beta = (4, 2, 1)$ . The dimension condition (see Remark 3.12) leads to  $2\alpha_2 + 3 = 7$ , i.e.,  $\alpha = (5, 2)$ .

• Assume now that s > 3. We prove that f cannot exist. Indeed, by Lemma 5.1,  $f(C_2) = C'_1$ , hence  $\alpha_1 - \alpha_2 = s > 3$ . Moreover, as  $f(RC_2) = RC'_1$ , by Propositions 4.12 and 4.14,  $\operatorname{length}(RC_2) = \alpha_1 - \alpha_2 + 1 = \operatorname{length}(RC'_1) = s$ , which is a contradiction.

The sufficiency is immediate.

**Proposition 5.4.** There are no isomorphisms  $f: V(\alpha_1, \ldots, \alpha_r) \to V(\beta_1, \ldots, \beta_s)$  with  $r \ge 3$ , s > 3 and  $r \ne s$ .

*Proof.* By Lemma 5.1,  $f(C_1) \neq C'_1$ .

- If  $\alpha_1 \alpha_2 > 1$ , then necessarily  $f(C_2) = C'_1$ . It means that  $\text{length}(C_2) = \text{length}(C'_1)$ , i.e.,  $\alpha_1 \alpha_2 + 1 = s + 1$ . As  $\text{length}(RC_2) = \text{length}(RC'_1)$ , by Propositions 4.13 and 4.14 we get  $\alpha_1 - \alpha_2 + 1 = s$  if  $\beta_1 - \beta_2 > 1$  or  $\alpha_1 - \alpha_2 + 1 = s - 1$  if  $\beta_1 - \beta_2 = 1$ . In both cases we have a contradiction.
- If α<sub>1</sub> − α<sub>2</sub> = 1, then necessarily f(C<sub>3</sub>) = C'<sub>1</sub>. The conclusion follows as in the previous case substituting length(C<sub>2</sub>) by length(C<sub>3</sub>) = 2(α<sub>1</sub> − α<sub>3</sub>) and length(RC<sub>2</sub>) by length(RC<sub>3</sub>) = 2(α<sub>1</sub> − α<sub>3</sub>).

Isomorphisms of hyperinvariant lattices

## **5.2** Isomorphic hyperinvariant lattices when r = s.

The first case we analyze is r = s = 2. **Proposition 5.5.**  $V(\alpha_1, \alpha_2) \simeq V(\beta_1, \beta_2)$  if and only if  $\alpha_i = \beta_i$ 

*Proof.* The sufficiency is immediate.

If  $\alpha_1 - \alpha_2 = 1$ , then  $V(\alpha_1, \alpha_2) \simeq V(2\alpha_2 + 1)$ . Hence,  $V(\beta_1, \beta_2)$  shall be isomorphic to  $V(2\beta_2 + 1)$ , and it implies that  $\beta_1 - \beta_2 = 1$ ,  $\alpha_2 = \beta_2$  and  $\alpha_1 = \beta_1$ .

It remains to analyze then, the case  $\alpha_1 - \alpha_2 > 1$  and  $\beta_1 - \beta_2 > 1$ :

- If  $f(C_2) = C'_2$ , then  $\alpha_1 \alpha_2 = \beta_1 \beta_2$ . As  $\alpha_1 + \alpha_2 = \beta_1 + \beta_2$ , the result follows.
- If  $f(C_1) = C'_2$  and  $f(C_2) = C'_1$ , then  $\beta_1 \beta_2 = 2$  and  $\alpha_1 \alpha_2 = 2$ . By the cardinality condition (Remark 3.12) we have  $(\alpha_1 \alpha_2 + 1)(\alpha_2 + 1) = (\beta_1 \beta_2 + 1)(\beta_2 + 1)$ , from where we conclude that  $\alpha_1 = \beta_1$  and  $\alpha_2 = \beta_2$ .

**Proposition 5.6.** There are no isomorphisms  $f: V(\alpha_1, \alpha_2, \alpha_3) \to V(\beta_1, \beta_2, \beta_3)$ ,  $\alpha \neq \beta$ , such that  $f(C_i) = C'_j$  for any i, j = 1, 2, 3 and  $i \neq j$ .

*Proof.* Assume that there exists an isomorphism  $f: V(\alpha_1, \alpha_2, \alpha_3) \to V(\beta_1, \beta_2, \beta_3)$  where  $\alpha \neq \beta$  and  $f(C_i) \neq C'_i$  for some *i*. We must analyze the following cases:

- If  $\alpha_1 \alpha_2 > 1$  and  $\beta_1 \beta_2 > 1$  then  $f(C_1) = C'_2$  and  $f(C_2) = C'_1$ , which implies that  $\alpha_1 \alpha_2 = \beta_1 \beta_2 = 3$ . Moreover, as  $f(RC_1) = RC'_2$ , by Proposition 4.13 we conclude that  $\text{length}(RC_1) = \text{length}(RC'_2) = \beta_1 \beta_2 + 1 = 4$  and  $\alpha_2 \alpha_3 = \alpha_3 = 1$ . Analogously,  $\beta_2 \beta_3 = \beta_3 = 1$ . Hence  $(\alpha_1, \alpha_2, \alpha_3) = (\beta_1, \beta_2, \beta_3) = (5, 2, 1)$ , which contradicts the fact that  $\alpha \neq \beta$ .
- If  $\alpha_1 \alpha_2 > 1$  and  $\beta_1 \beta_2 = 1$ , we have two possibilities:  $f(C_1) = C'_3$  and  $f(C_2) = C'_1$  or  $f(C_1) = C'_1$  and  $f(C_2) = C'_3$ .

In the first case  $2(\beta_1 - \beta_3) = 4$  and  $\alpha_1 - \alpha_2 = 3$ . As length $(RC_1) = \text{length}(RC'_3)$ , from Proposition 4.13 we conclude that  $\alpha_2 - \alpha_3 = \alpha_3 = 1$ . Hence,  $\alpha = (5, 2, 1)$  and  $\beta = (\beta_3 + 2, \beta_3 + 1, \beta_3)$ , and from the dimension condition (Remark 3.12) we get  $5 = 3\beta_3$ , which is not possible.

In the second option, from  $f(C_2) = C'_3$  we obtain that  $\alpha_1 - \alpha_2 + 1 = 2(\beta_1 - \beta_3)$ . As length $(RC_1) =$ length $(RC'_1)$ , from Proposition 4.13 we conclude that  $\alpha_2 - \alpha_3 = \alpha_3 = 1$  and  $\beta_2 - \beta_3 = 1$ . Hence,  $\alpha = (5, 2, 1)$  and  $\beta = (\beta_3 + 2, \beta_3 + 1, \beta_3)$ , and the conclusion follows as before.

• If  $\alpha_1 - \alpha_2 = 1$  and  $\beta_1 - \beta_2 = 1$ , then  $f(C_1) = C'_3$ , which implies that  $4 = 2(\beta_1 - \beta_3)$ . As length $(RC_1) =$ length $(RC'_3)$ , necessarily  $\alpha_2 - \alpha_3 = 1$ . Hence  $\alpha = (\alpha_3 + 2, \alpha_3 + 1, \alpha_3)$  and  $\beta = (\beta_3 + 2, \beta_3 + 1, \beta_3)$ . By Remark 3.12 we conclude that  $\alpha_i = \beta_i, i = 1, 2, 3$ .

**Remark 5.7.** From the proof of the above proposition we can see that there is a single hyperlattice, V(5, 2, 1), such that there exist two isomorphisms

 $f: V(5,2,1) \to V(5,2,1),$ 

which are the identity, and another one satisfying  $f(C_1) = C_2$  (see Figure 1).

Analogously, for the family of hyperlattices  $V(\alpha_3 + 2, \alpha_3 + 1, \alpha_3)$  for  $\alpha_3 = 1, 2, ...$ , there exist two possible isomorphisms

$$f: V(\alpha_3 + 2, \alpha_3 + 1, \alpha_3) \to V(\alpha_3 + 2, \alpha_3 + 1, \alpha_3),$$

the identity isomorphism, and another one such that  $f(C_1) = C_3$ .

#### Proposition 5.8. Let

$$f: V(\alpha_1, \ldots, \alpha_r) \longrightarrow V(\beta_1, \ldots, \beta_r)$$

be an isomorphism with r > 3, then  $f(C_1) = C'_1$ .



Figure 1: Hyperlattice V(5, 2, 1)

*Proof.* Assume that  $f(C_1) \neq C'_1$ . The prove is analogous to that of Proposition 5.4.

It only remains to study isomorphisms  $f : V(\alpha_1, \ldots, \alpha_r) \longrightarrow V(\beta_1, \ldots, \beta_r)$  such that  $f(C_1) = C'_1$ . To do it we need some isomorphism theorems of factor lattices, which are built as quotients by congruent relations. **Definition 5.9.** Given a hyperlattice  $V(\alpha_1, \ldots, \alpha_r)$ , we define the following congruence relations:

• If  $\alpha_1 - \alpha_2 > 1$  and  $r \ge 2$ , we denote by  $\sim_2$  the congruence relation defined as

$$(u_1,\ldots,u_r)\sim_2 (\bar{u}_1,\ldots,\bar{u}_r) \Leftrightarrow u_2=\bar{u}_2,\ldots,u_r=\bar{u}_r.$$

• If  $\alpha_1 - \alpha_2 = 1$  and  $r \ge 3$ , we denote by  $\sim_3$  the congruence relation defined as

$$(u_1,\ldots,u_r)\sim_3(\bar{u}_1,\ldots,\bar{u}_r)\Leftrightarrow u_3=\bar{u}_3,\ldots,u_r=\bar{u}_r.$$

The relation  $\sim_2$  allows us to build the factor lattice  $V(\alpha_1, \ldots, \alpha_r) / \sim_2$ . Notice that if  $(u_1, u_2, u_3, \ldots, u_r) \in V(\alpha_1, \ldots, \alpha_r)$ , all of the elements of  $V(\alpha_1, \ldots, \alpha_r)$  congruent to it are

$$\{(u_2, u_2, u_3, \ldots, u_r), \ldots, (u_2 + \alpha_1 - \alpha_2, u_2, u_3, \ldots, u_r)\},\$$

where  $u_2 \leq u_1 \leq u_2 + \alpha_1 - \alpha_2$ . Therefore, each class contains  $\alpha_1 - \alpha_2 + 1$  elements and

$$\operatorname{card}(V(\alpha_1,\ldots,\alpha_r)/\sim_2) = (\alpha_2 - \alpha_3 + 1)\ldots(\alpha_r + 1)$$

We take as representative of a class in  $V(\alpha_1, \ldots, \alpha_r) / \sim_2$  the first one of the previous elements; i.e.,

$$[(u_2, u_2, u_3, \dots, u_r)] = \{(u_2, u_2, u_3, \dots, u_r), \dots, (u_2 + \alpha_1 - \alpha_2, u_2, u_3, \dots, u_r)\}$$

The join and meet of two classes are defined, respectively, as follows

 $[(u_2, u_2, u_3, \dots, u_r)] \lor [(\bar{u}_2, \bar{u}_2, \bar{u}_3, \dots, \bar{u}_r)] = [(\max(u_2, \bar{u}_2), \max(u_2, \bar{u}_2), \dots, \max(u_r, \bar{u}_r))],$  $[(u_2, u_2, u_3, \dots, u_r)] \land [(\bar{u}_2, \bar{u}_2, \bar{u}_3, \dots, \bar{u}_r)] = [(\min(u_2, \bar{u}_2), \min(u_2, \bar{u}_2), \dots, \min(u_r, \bar{u}_r))].$  Analogously,  $\sim_3$  allows us to build the factor lattice  $V(\alpha_1, \ldots, \alpha_r) / \sim_3$ . Given  $(u_1, u_2, u_3, \ldots, u_r) \in V(\alpha_1, \ldots, \alpha_r)$ , all of the elements of  $V(\alpha_1, \ldots, \alpha_r)$  congruent to it are

$$\{(u_3, u_3, u_3, u_4, \dots, u_r), \dots, (u_3 + \alpha_1 - \alpha_3, u_3 + \alpha_2 - \alpha_3, u_3, u_4, \dots, u_r)\}$$

where  $u_3 \le u_1 \le u_3 + \alpha_1 - \alpha_3$  and  $u_3 \le u_2 \le u_3 + \alpha_2 - \alpha_3$ . Notice that each class contains  $2(\alpha_1 - \alpha_3)$  elements and

$$\operatorname{card}(V(\alpha_1,\ldots,\alpha_r)/\sim_3) = (\alpha_3 - \alpha_4 + 1)\ldots(\alpha_r + 1).$$

The join and meet in this quotient lattice are defined as in the previous case.

The following lemma is satisfied.

Lemma 5.10. We have the following isomorphisms:

- 1. If  $r \geq 2$  and  $\alpha_1 \alpha_2 > 1$ , then  $V(\alpha_1, \ldots, \alpha_r) / \sim_2 \simeq V(\alpha_2, \ldots, \alpha_r)$ .
- 2. If  $r \geq 3$  and  $\alpha_1 \alpha_2 = 1$ , then  $V(\alpha_1, \ldots, \alpha_r) / \sim_3 \simeq V(\alpha_3, \ldots, \alpha_r)$ .

*Proof.* We only prove item 1 because the proof of item 2 is analogous. To check that both lattices are isomorphic we prove that there is a bijective map between the two lattices preserving meets and joins.

The following map is well defined and is clearly bijective

$$\phi: \quad V(\alpha_1, \dots, \alpha_r) / \sim_2 \quad \longrightarrow \quad V(\alpha_2, \dots, \alpha_r) \\ [(u_2, u_2, u_3, \dots, u_r)] \quad \mapsto \quad (u_2, u_3, \dots, u_r)$$

In order to prove join preservation it is sufficient to check whether the join of representatives of two classes maps to the join of its images, which is true. The preservation of the meet can be checked analogously.  $\Box$ 

Let  $V(\alpha_1, \ldots, \alpha_r)$  and  $V(\beta_1, \ldots, \beta_r)$  be hyperlattices with  $r \ge 2$ . If  $\alpha_1 - \alpha_2 > 1$  and  $\beta_1 - \beta_2 > 1$ , we are going to prove that if there exists an isomorphism between the hyperlattices transforming  $C_2$  into  $C'_2$ , then  $V(\alpha_1, \ldots, \alpha_r)/\sim_2$  and  $V(\beta_1, \ldots, \beta_r)/\sim_2$  are also isomorphic.

Given an isomorphism

 $f: V(\alpha_1, \ldots, \alpha_r) \longrightarrow V(\beta_1, \ldots, \beta_r),$ 

satisfying  $f(C_2) = C'_2$ , and the epimorphism

$$\pi: V(\beta_1, \ldots, \beta_r) \longrightarrow V(\beta_1, \ldots, \beta_r) / \sim_2,$$

let  $\bar{f} = \pi \circ f$ . As a consequence of Theorem 2.1 we have

$$V(\alpha_1, \dots, \alpha_r) / \ker(\bar{f}) \simeq V(\beta_1, \dots, \beta_r) / \sim_2 .$$
 (15)

If  $\alpha_1 - \alpha_2 = \beta_1 - \beta_2 = 1, r \ge 3$ , let

$$g: V(\alpha_1, \ldots, \alpha_r) \longrightarrow V(\beta_1, \ldots, \beta_r),$$

be an isomorphism such that  $g(C_3) = C'_3$ , it can be proved in an analogous way that

$$V(\alpha_1, \dots, \alpha_r) / \ker(\bar{g}) \simeq V(\beta_1, \dots, \beta_r) / \sim_3, \tag{16}$$

where  $\bar{g} = \pi \circ g$ .

We only need to prove the following result.

Theorem 5.11. With the above notation

1. If 
$$r \ge 2$$
,  $\alpha_1 - \alpha_2 > 1$  and  $\beta_1 - \beta_2 > 1$ , then  

$$V(\alpha_1, \dots, \alpha_r) / \ker(\bar{f}) = V(\alpha_1, \dots, \alpha_r) / \sim_2 .$$

2. If 
$$r \ge 3$$
 and  $\alpha_1 - \alpha_2 = \beta_1 - \beta_2 = 1$ , then  
 $V(\alpha_1, \dots, \alpha_r) / \ker(\bar{g}) = V(\alpha_1, \dots, \alpha_r) / \sim_3 .$ 

*Proof.* We only prove item 1. The proof of item 2 can be done in an analogous way. We show that the classes of the two lattices  $V(\alpha_1, \ldots, \alpha_r) / \ker(\bar{f})$  and  $V(\alpha_1, \ldots, \alpha_r) / \sim_2$  coincide.

As the number of elements in a class in  $V(\beta_1, \ldots, \beta_r) / \sim_2$  is  $\beta_1 - \beta_2 + 1$ , by (15) we know that  $\operatorname{card}(V(\alpha_1, \ldots, \alpha_r) / \ker(\bar{f})) = \operatorname{card}(V(\beta_1, \ldots, \beta_r) / \sim_2)$ , and it is  $(\beta_2 - \beta_3 + 1) \ldots (\beta_{r-1} - \beta_r + 1)(\beta_r + 1)$ .

By hypothesis  $f(C_2) = C'_2$ , then  $\alpha_1 - \alpha_2 = \beta_1 - \beta_2$ . As a consequence,  $\operatorname{card}(V(\alpha_1, \ldots, \alpha_r)/\sim_2) = \operatorname{card}(V(\alpha_1, \ldots, \alpha_r)/\ker(\bar{f}))$ . Moreover, let C be a class in  $V(\beta_1, \ldots, \beta_r)/\sim_2$ . Notice that the inverse image of the class  $\bar{f}^{-1}(C)$  contains  $\alpha_1 - \alpha_2 + 1$  elements (as does  $\pi^{-1}(C)$ ), and if  $u \in \bar{f}^{-1}(C)$ , then  $\bar{f}(u) = C$ , i.e., the set  $\bar{f}^{-1}(C)$  is included in the same class of  $V(\alpha_1, \ldots, \alpha_r)/\ker(\bar{f})$ . Therefore, each class in  $V(\alpha_1, \ldots, \alpha_r)/\ker(\bar{f})$  has  $\alpha_1 - \alpha_2 + 1$  elements. Furthermore,  $C_2 = \bar{f}^{-1}(0)$ , therefore  $C_2 \in V(\alpha_1, \ldots, \alpha_r)/\ker(\bar{f})$ .

Let us see that  $V(\alpha_1, \ldots, \alpha_r) / \sim_2$  and  $V(\alpha_1, \ldots, \alpha_r) / \ker(\bar{f})$  contain the same classes.

Obviously,  $C_2 \in V(\alpha_1, \ldots, \alpha_r) / \sim_2$ . From the chain  $C_2$  we construct classes of  $V(\alpha_1, \ldots, \alpha_r) / \ker(\bar{f})$ , and we see that this quotient lattice contains all of the classes of  $V(\alpha_1, \ldots, \alpha_r) / \sim_2$ .

Since  $f(RC_2) = RC'_2$ , the set of elements of  $RC_2$ 

$$\{(1, 1, 0, \dots, 0), (2, 1, 0, \dots, 0), \dots, (\alpha_1 - \alpha_2, 1, 0, \dots, 0), (\alpha_1 - \alpha_2 + 1, 1, 0, \dots, 0)\}$$

is a class in  $V(\alpha_1, \ldots, \alpha_r) / \ker(\bar{f})$ .

Let  $[(u_2, u_2, u_3, ..., u_r)] \in V(\alpha_1, ..., \alpha_r) / \sim_2, u_2 \neq \alpha_2$ . A father of this class is of the form  $[(u_2, u_2, ..., u_i + 1, ..., u_r)]$  for some  $i \in \{3, ..., r\}$  or  $[(u_2 + 1, u_2 + 1, u_3, ..., u_r)]$ .

Assume that the father is  $[(u_2, u_2, \dots, u_i+1, \dots, u_r)]$ . Since the element  $(u_2, u_2, \dots, u_i+1, \dots, u_r) \in V(\alpha_1, \dots, \alpha_r)$ , a class parent can be constructed as

$$\begin{array}{cccc} (u_2, u_2, u_3, \dots, u_i, \dots, u_r) & \lor & (u_2, u_2, u_3, \dots, u_i + 1, \dots, u_r) \\ & \vdots & & \vdots \\ (u_2 + \alpha_1 - \alpha_2, u_2, u_3, \dots, u_i, \dots, u_r) & \lor & (u_2, u_2, u_3, \dots, u_i + 1, \dots, u_r) \end{array}$$

It means that  $[(u_2, u_2, ..., u_i + 1, ..., u_r)] \in V(\alpha_1, ..., \alpha_r) / \ker(\bar{f}).$ 

Assume now that the father is  $[(u_2 + 1, u_2 + 1, u_3, \dots, u_r)]$ . As  $(u_2 + 1, u_2 + 1, u_3, \dots, u_r) \in V(\alpha_1, \dots, \alpha_r)$ , an analogous construction proves that

$$\{(u_2+1, u_2+1, u_3, \dots, u_r), \dots, (u_2+\alpha_1-\alpha_2, u_2+1, u_3, \dots, u_r)\}$$

belong to the same class in  $V(\alpha_1, \ldots, \alpha_r) / \ker(\bar{f})$ .

As the cardinality of each class is  $\alpha_1 - \alpha_2 + 1$ , one element is missing in this class.

By (2), a class must be a chain. Therefore, the last element  $(u_2 + \alpha_1 - \alpha_2, u_2 + 1, \dots, u_r)$  must have a father in the class. The possible fathers are among the following sequence of hypertuples:  $(u_2 + \alpha_1 - \alpha_2 + 1, u_2 + 1, u_3, \dots, u_r)$ ,  $(u_2 + \alpha_1 - \alpha_2, u_2 + 2, u_3, \dots, u_r)$ ,...,  $(u_2 + \alpha_1 - \alpha_2 + 1, u_2 + 1, u_3 + 1, \dots, u_r)$ ,...:

• If 
$$(u_2 + \alpha_1 - \alpha_2, u_2 + 2, u_3, \dots, u_r) \in [(u_2 + 1, u_2 + 1, u_3, \dots, u_r)]$$
, then  
 $(u_2 + 1, u_2 + 1, u_3, \dots, u_r) \lor (u_2 + 2, u_2 + 2, u_3, \dots, u_i, \dots, u_r) \sim$   
 $\sim (u_2 + \alpha_1 - \alpha_2, u_2 + 2, u_3, \dots, u_r) \lor (u_2 + 2, u_2 + 2, u_3, \dots, u_i, \dots, u_r),$ 

that is

 $(u_2+2, u_2+2, u_3, \dots, u_r) \sim (u_2+\alpha_1-\alpha_2, u_2+2, u_3, \dots, u_r),$ 

but it means that  $(u_2 + 2, u_2 + 2, u_3, \dots, u_r) \in [(u_2 + 1, u_2 + 1, u_3, \dots, u_r)]$ , and this is not possible because then the class will have more elements than its cardinality.

• If 
$$(u_2 + \alpha_1 - \alpha_2, u_2 + 1, u_3, \dots, u_i + 1, \dots, u_r) \in [(u_2 + 1, u_2 + 1, u_3, \dots, u_r)]$$
, then

$$(u_2+1, u_2+1, u_3, \dots, u_r) \lor (u_2+1, u_2+1, u_3, \dots, u_i+1, \dots, u_r) \sim$$

$$\sim (u_2 + \alpha_1 - \alpha_2, u_2 + 1, u_3, \dots, u_i + 1, \dots, u_r) \lor (u_2 + 1, u_2 + 1, u_3, \dots, u_i + 1, \dots, u_r),$$

that is

$$(u_2+1, u_2+1, u_3, \dots, u_i+1, \dots, u_r) \sim (u_2+\alpha_1-\alpha_2, u_2+1, u_3, \dots, u_i+1, \dots, u_r),$$

and this is not possible because the class will have more elements than its cardinality.

Therefore, the new element belonging to the class is  $(u_2 + \alpha_1 - \alpha_2 + 1, u_2 + 1, u_3, \dots, u_r)$ .

Taking into account that  $C_2, RC_2 \in V(\alpha_1, \dots, \alpha_r) / \ker(\bar{f})$ , by recurrence the result follows.

These equivalence relations allow us to prove the following lemma.

**Proposition 5.12.** Let  $V(\alpha_1, \ldots, \alpha_r)$  and  $V(\beta_1, \ldots, \beta_r)$  be hyperlattices, then

$$V(\alpha_1, \ldots, \alpha_r) \simeq V(\beta_1, \ldots, \beta_r)$$
 if and only if  $\alpha_i = \beta_i, i = 1, \ldots, r$ .

*Proof.* We apply induction. For r = 1 and r = 2 the result is proved in Propositions 5.2 and 5.5. For  $r \ge 3$ , let  $f: V(\alpha_1, \ldots, \alpha_r) \longrightarrow V(\beta_1, \ldots, \beta_r)$  be an isomorphism, then by Proposition 5.8,  $f(C_1) = C'_1$ . Hence, we analyze the following cases:

• If  $\alpha_1 - \alpha_2 > 1$  and  $\beta_1 - \beta_2 > 1$ , then  $f(C_2) = C'_2$ , and by (15) and Theorem 5.11 we have

$$V(\alpha_1,\ldots,\alpha_r)/\sim_2 = V(\alpha_1,\ldots,\alpha_r)/\ker(f) \simeq V(\beta_1,\ldots,\beta_r)/\sim_2$$
.

Therefore, by Lemma 5.10 we obtain

$$V(\alpha_2,\ldots,\alpha_r)\simeq V(\beta_2,\ldots,\beta_r).$$

The result follows by induction hypothesis.

• If  $\alpha_1 - \alpha_2 = 1$  and  $\beta_1 - \beta_2 = 1$ , then  $f(C_3) = C'_3$ , and by an analogous argument to that of the previous case we obtain

$$V(\alpha_3,\ldots,\alpha_r)\simeq V(\beta_3,\ldots,\beta_r).$$

The result follows by induction hypothesis.

• If  $\alpha_1 - \alpha_2 > 1$  and  $\beta_1 - \beta_2 = 1$ , then  $f(C_2) = C'_3$ , which implies  $\alpha_1 - \alpha_2 = 2(\beta_1 - \beta_3)$  and  $f(RC_2) = RC'_3$ . It means that  $\alpha_1 - \alpha_2 + 1 = 2(\beta_1 - \beta_2)$ , and we have a contradiction.

#### 5.3 Main theorem

For the general case the result is summarized in the following main theorem, which is proved in the previous results.

**Theorem 5.13.** Let  $A, B \in \mathbb{C}^{n \times n}$  be nilpotent matrices. Let  $\alpha = (\alpha_1, \ldots, \alpha_r)$  and  $\beta = (\beta_1, \ldots, \beta_s)$  be the reduced Segre characteristic of A and B, respectively. Then,  $\operatorname{Hinv}(A) \simeq \operatorname{Hinv}(B)$  if and only if one of the following conditions are satisfied:

- 1.  $\alpha = (5, 2)$  and  $\beta = (4, 2, 1)$  or vice-versa.
- 2.  $\alpha = (l, l-1)$  and  $\beta = (2l-1)$  for some  $l \ge 2$  or vice-versa. 3.  $\alpha = \beta$ .

## Acknowledgments

The second author is partially supported by grant PID2019-104047GB-I00. The second and third authors are partially supported by grant PID2021-124827NB-I00 funded by MCIN/AEI/ 10.13039/501100011033 and by "ERDF A way of making Europe" by the "European Union".

# References

- [1] L. Brickman, P.A. Fillmore. The invariant subspace lattice of a linear transformation. Can. J. Math., 19, 35, (1967), 810-822.
- [2] P.A. Fillmore, D.A. Herrero and W.E. Longstaff. The hyperinvariant subspace lattice of a linear transformation. Linear Algebra Appl. 17 (1977), 125-132.
- [3] G. Grätzel. The Congruences of a Finite Lattice. Birkhäuser, 2023.
- [4] W. E. Longstaff. A lattice-theoretic description of the lattice of hyperinvariant subspaces of a linear transformation, Canad. /. Math. 28 (1976) 1062-1066.
- [5] P.Y. Wu. Which Linear Transformations Have Isomorphic Hyperinvariant Subspace Lattices? Linear Algebra Appl. 169 (1992) 163-178.