

skew-supersymmetric solution of the super Malcev Yang-Baxter equation and Pre-Malcev superalgebras

Fattoum Harrathi*

University of Sfax, Faculty of Sciences Sfax, BP 1171, 3038 Sfax, Tunisia

Abstract. The purpose of this paper is to introduce the notion of pre-Malcev superalgebras as the algebraic structure behind the super \mathcal{O} -operators on Malcev superalgebras. Moreover, the relations among Malcev superalgebras, pre-Malcev superalgebras and pre-alternative superalgebras are established. Then, we study the operator forms of the classical Yang-Baxter equation (CYBE) in Malcev superalgebras and give their relationship with super \mathcal{O} -operators. There are close relationships between the CYBE in Malcev superalgebras and pre-Malcev superalgebras which can be interpreted through the super \mathcal{O} -operators.

Key words: Malcev superalgebra, pre-Malcev superalgebra, super \mathcal{O} -operator, representation, super Malcev Yang-Baxter equation.

Mathematics Subject Classification (2020): 17A70, 17B38.

Contents

1	Introduction	1
2	Some basic properties of Malcev superalgebras	3
3	Super \mathcal{O}-operators of Malcev superalgebras and pre-Malcev superalgebras	7
4	An analogue of the super CYBE on Malcev superalgebras	12
4.1	Super \mathcal{O} -operators and operator forms of the super MYBE	12
4.2	Constructions of solutions of the super MYBE in Malcev superalgebras	15

1 Introduction

As a promotion of Lie algebra, the notion of Malcev algebras were introduced in 1955 by A. I. Malcev [22], who called these objects Moufang-Lie algebras because of their connection with analytic Moufang loops. Motivated by the study of physics and the geometry of smooth loops, Malcev algebras have been widely used. Similarly to the tangent algebra of a Lie group is a Lie algebra, the tangent algebra of a locally analytic Moufang loop is a Malcev algebra, see[16]. A Malcev algebra is a non-associative algebra with a skew-symmetric multiplication that satisfies the Malcev

*E-mail: harrathifattoum285@gmail.com

identity. That is to say, a vector space A with an anti-symmetric multiplication $[\cdot, \cdot]$ that satisfies, for all $x, y, z \in A$, the Malcev identity

$$J(x, y, [x, z]) = [J(x, y, z), x], \quad (1.1)$$

where $J(x, y, z) = [[x, y], z] + [[z, x], y] + [[y, z], x]$ is the Jacobian [23]. Other influential work includes [1, 2, 18, 21].

Pre-Malcev algebras have been studied extensively since [20] which are the generalization of pre-Lie algebras, in the sense that any pre-Lie algebra is a pre-Malcev algebra but the converse is not true. Studying pre-Malcev algebras independently is significant not only to its own further development, but also to develop the areas closely connected with such algebras. A pre-Malcev algebra is a vector space A endowed with a bilinear product \cdot satisfying the following identity for $x, y, z, t \in A$,

$$[y, z] \cdot (x \cdot t) + [[x, y], z] \cdot t + y \cdot ([x, z] \cdot t) - x \cdot (y \cdot (z \cdot t)) + z \cdot (x \cdot (y \cdot t)) = 0, \quad (1.2)$$

where $[x, y] = x \cdot y - y \cdot x$.

The existence of sub-adjacent Malcev algebras and compatible pre-Malcev algebras was given in [20, Proposition 5]. For a given pre-Malcev algebra (A, \cdot) , there is a Malcev algebra A^C defined by the commutator $[x, y] = x \cdot y - y \cdot x$, and the left multiplication operator in A induces a representation of Malcev algebra A^C .

In paper [7], as Lie algebras of some generalized groups, Lie superalgebras are introduced. In 1977, Kac deal as applications, Li prove that certain lowest weight modules for some well-known infinite-dimensional Lie algebras or Lie superalgebras have natural vertex operator superalgebra structures in [19]. Then as a natural generalization of Malcev algebras and Lie superalgebras, the theory of Malcev superalgebras has quite developed. In particular, Lie superalgebras are examples of Malcev superalgebras. And the definitions and basic facts of the theory of Malcev superalgebras can be found in [3, 25].

Classical Yang-Baxter equation (CYBE) first arose in the study of inverse scattering theory (see [9, 10]). It has a profound connection with many branches of mathematical physics and pure mathematics ([6], etc). In particular, CYBE can be regarded as a “classical limit” of quantum Yang-Baxter equation, which plays an important role in the study of classical integrable system ([24]). The generalization of the ordinary CYBE in the super case, namely, the CYBE in Lie superalgebras, or the graded CYBE ([12, 29]), or the super CYBE ([4]), has been studied widely since 1990s. One of the motivations to study such a structure is its relationship with the solutions of quantum Yang-Baxter equation associated to quantum supergroups or quantum superalgebras or quantized universal enveloping superalgebras ([8, 11, 27, 28]). In fact, many important results on this topic have already obtained like the close relationships between the CYBE in Lie superalgebras and Lie bi-superalgebras ([12]), Poisson-Lie supergroups ([4]) and so on.

In this paper, super Malcev Yang-Baxter equation (Super MYBE) in Malcev superalgebras and super \mathcal{O} -operators are introduced. We may concentrate on the case that $|r| = 0$ (i.e., r is even). It is hard to study the case when r is odd. Moreover, we exploit pre-Malcev superalgebras as super-versions of pre-Malcev algebras and study their relations with Malcev superalgebras and pre-alternative superalgebras. The paper is organized as follows. In Section 2, we recall some basic facts on super vector spaces and Malcev superalgebras, then present some fundamental results on representations of Malcev superalgebras, especially focusing on adjoint and co-adjoint representations. In Section 3, we introduce the super \mathcal{O} -operators and pre-Malcev superalgebras and give their relations with super \mathcal{O} -operators, Malcev superalgebras and pre-alternative superalgebras. In

Section 4, we construct a direct relation between super \mathcal{O} -operator and super Malcev Yang-Baxter equation and show that pre-Malcev algebras are the natural underlying structures.

In this paper, we consider all vector space are finite dimensional over an algebraically closed commutative field \mathbb{K} of characteristic zero.

2 Some basic properties of Malcev superalgebras

In this section we presents fundamental concepts and develop some helpful results that we will use later. Let V be a vector space over a field \mathbb{K} . The space V is called a super (i.e., \mathbb{Z}_2 -graded) vector space if $V = V_{\bar{0}} \oplus V_{\bar{1}}$. The elements in $V_{\bar{0}} \cup V_{\bar{1}}$ are called homogeneous. Denoted by $\mathcal{H}(V)$ the set of homogenous elements of V . We use the expression $|x|$ to denote the parity index of the homogeneous element x , where

$$|x| = \begin{cases} 0, & x \in V_{\bar{0}}; \\ 1, & x \in V_{\bar{1}}. \end{cases}$$

It is assumed in this paper that all elements are the homogeneous of their corresponding super vector spaces.

Definition 2.1. A left alternative superalgebra is a \mathbb{Z}_2 -graded vector space $A = A_{\bar{0}} \oplus A_{\bar{1}}$ equipped with a bilinear product $\star : A \times A \rightarrow A$ obeying $A_\alpha \star A_\beta \subset A_{\alpha+\beta}$, $\forall \alpha, \beta \in \mathbb{Z}_2$ and the left alternative super identity,

$$as(x, y, z) + (-1)^{|x||y|} as(y, x, z) = 0, \quad \forall x, y, z \in A. \quad (2.3)$$

where $as(x, y, z) = (x \star y) \star z - x \star (y \star z)$ is the associator.

Definition 2.2. A right alternative superalgebra is a \mathbb{Z}_2 -graded vector space $A = A_{\bar{0}} \oplus A_{\bar{1}}$ equipped with a bilinear product $\star : A \times A \rightarrow A$ obeying $A_\alpha \star A_\beta \subset A_{\alpha+\beta}$, $\forall \alpha, \beta \in \mathbb{Z}_2$ and the right alternative super identity

$$as(x, y, z) + (-1)^{|y||z|} as(x, z, y) = 0, \quad \forall x, y, z \in A. \quad (2.4)$$

Definition 2.3. An alternative superalgebra is one which is both left and right alternative superalgebra.

Definition 2.4. A Malcev superalgebra is a \mathbb{Z}_2 -graded vector space $A = A_{\bar{0}} \oplus A_{\bar{1}}$ over a field \mathbb{K} equipped with a \mathbb{K} -bilinear map $[\cdot, \cdot] : A \times A \longrightarrow A$ satisfying $[A_\alpha, A_\beta] \subset A_{\alpha+\beta}$, $\forall \alpha, \beta \in \mathbb{Z}_2$ and the following conditions

- (i) $[x, y] = -(-1)^{|x||y|}[y, x]$,
- (ii) $(-1)^{|y||z|}[[x, z], [y, t]] = [[[x, y], z], t] + (-1)^{|x|(|y|+|z|+|t|)}[[[y, z], t], x]$
 $+ (-1)^{(|x|+|y|)(|z|+|t|)}[[[z, t], x], y] + (-1)^{|t|(|x|+|y|+|z|)}[[[t, x], y], z]$,

for all $x, y, z, t \in \mathcal{H}(A)$. An element of $A_{\bar{0}}$ is called even and an element of $A_{\bar{1}}$ is called odd.

Theorem 2.1. Let (A, \star) be an alternative superalgebra. Then $(A, [\cdot, \cdot])$ is a Malcev admissible superalgebra, where

$$[x, y] = x \star y - (-1)^{|x||y|} y \star x, \quad \forall x, y \in \mathcal{H}(A).$$

Let A be any superalgebra and V be a \mathbb{Z}_2 -graded vector space. The space $gl(V)$ consisting of all the linear transformations on V has a natural \mathbb{Z}_2 -gradation as

$$gl(V)_\alpha = \{f \in gl(V) | f(V_\alpha) \subseteq V_{\alpha+\beta}, \alpha, \beta \in \mathbb{Z}_2\}. \quad (2.5)$$

Definition 2.5. A bimodule over an alternative superalgebra (A, \star) consists of a \mathbb{Z}_2 -graded linear space V and two even bilinear maps $\mathbf{l}, \mathbf{r} : A \rightarrow gl(V)$ such that, for any homogeneous elements $x, y \in A$,

$$\mathbf{l}(x \star y) + (-1)^{|x||y|}\mathbf{l}(y \star x) - \mathbf{l}(x)\mathbf{l}(y) - (-1)^{|x||y|}\mathbf{l}(y)\mathbf{l}(x) = 0, \quad (2.6)$$

$$\mathbf{r}(y)\mathbf{r}(x) + (-1)^{|x||y|}\mathbf{r}(x)\mathbf{r}(y) - \mathbf{r}(x \star y) - (-1)^{|x||y|}\mathbf{r}(y \star x) = 0, \quad (2.7)$$

$$\mathbf{r}(y)\mathbf{r}(x) + (-1)^{|x||y|}\mathbf{r}(y)\mathbf{l}(x) - (-1)^{|x||y|}\mathbf{l}(x)\mathbf{r}(y) - \mathbf{r}(x \star y) = 0, \quad (2.8)$$

$$\mathbf{r}(y)\mathbf{l}(x) + (-1)^{|x||y|}\mathbf{l}(x \star y) - (-1)^{|x||y|}\mathbf{l}(x)\mathbf{l}(y) - \mathbf{l}(x)\mathbf{r}(y) = 0. \quad (2.9)$$

Proposition 2.1. $(V, \mathbf{l}, \mathbf{r})$ is a bimodule of an alternative superalgebra (A, \star) if and only if the direct sum $A \oplus V$ of vector spaces is turned into an alternative superalgebra (the semidirect sum) by defining multiplication in $A \oplus V$ by

$$(x + a) \circ (y + b) = x \star y + \mathbf{l}(x)b + \mathbf{r}(y)a, \quad \forall x, y \in \mathcal{H}(A), a, b \in \mathcal{H}(V). \quad (2.10)$$

We denote it by $A \bowtie_{\mathbf{l}, \mathbf{r}} V$ or simply $A \bowtie V$.

Definition 2.6. Let A be a Malcev superalgebra.

(i) Let V be a \mathbb{Z}_2 -graded vector space. An even linear map $\rho : A \rightarrow gl(V)$ (i.e., $|\rho| = 0$) is a Malcev representation of A on V such that for any $x, y, z \in \mathcal{H}(A)$,

$$\begin{aligned} \rho([[x, y], z]) &= \rho(x)\rho(y)\rho(z) - (-1)^{|z|(|x|+|y|)}\rho(z)\rho(x)\rho(y) \\ &+ (-1)^{|x|(|y|+|z|)}\rho(y)\rho([z, x]) \\ &- (-1)^{|x|(|y|+|z|)}\rho([y, z])\rho(x). \end{aligned} \quad (2.11)$$

(ii) Consider two Malcev representations $\rho : A \rightarrow gl(V)$ and $\rho' : A \rightarrow gl(V')$ of A , where V and V' are \mathbb{Z}_2 -graded vector spaces. We say that ρ and ρ' are equivalent if there exists a bijective even linear map $\phi : V \rightarrow V'$ such that $\phi \circ \rho(x) = \rho'(x) \circ \phi, \forall x \in \mathcal{H}(A)$.

Proposition 2.2. (V, ρ) is a representation of a Malcev superalgebra A if and only if the direct sum $A \oplus V$ of vector spaces is a Malcev superalgebra by defining a map on $A \oplus V$ by

$$[x + a, y + b]_{A \oplus V} := [x, y] + \rho(x)b - (-1)^{|x||y|}\rho(y)a, \quad (2.12)$$

for all $x, y \in \mathcal{H}(A), a, b \in \mathcal{H}(V)$. In this case, $A \oplus V$ is called the semi-direct product of A and V , denoted by $A \bowtie_\rho V$ or simply $A \bowtie V$.

Note that $(A \oplus V)_{\overline{0}} = A_{\overline{0}} \oplus V_{\overline{0}}$, implying that if $x + a \in \mathcal{H}(A \oplus V)$, then $|x + a| = |x| = |a|$.

Proof. For all $x, y, z, t \in \mathcal{H}(A)$ and $a, b, c, d \in \mathcal{H}(V)$, we have

$$\begin{aligned} &[[x + a, z + c]_{A \oplus V}, [y + b, t + d]_{A \oplus V}]_{A \oplus V} \\ &= [[x, z], [y, t]] + \rho([x, z])\rho(y)d - (-1)^{|y||t|}\rho([x, z])\rho(t)b \end{aligned}$$

$$\begin{aligned}
& -(-1)^{(|x|+|z|)(|y|+|t|)} \rho([y, t]) \rho(x) c + (-1)^{(|x|+|z|)(|y|+|t|)+|x||z|} \rho([y, t]) \rho(z) a, \\
& [[x+a, y+b]_{A \oplus V}, z+c]_{A \oplus V}, t+d]_{A \oplus V} \\
& = [[[x, y], z], t] + \rho([[x, y], z]) d - (-1)^{|t|(|x|+|y|+|z|)} \rho(t) \rho([x, y]) c \\
& \quad + (-1)^{|t|(|x|+|y|+|z|)+|z|(|x|+|y|)} \rho(t) \rho(z) \rho(x) b - (-1)^{|t|(|x|+|y|+|z|)+|z|(|x|+|y|)+|x||y|} \rho(t) \rho(z) \rho(y) a, \\
& [[y+b, z+c]_{A \oplus V}, t+d]_{A \oplus V}, x+a]_{A \oplus V} \\
& = [[[y, z], t], x] + \rho([[y, z], t]) a - (-1)^{|x|(|y|+|z|+|t|)} \rho(x) \rho([y, z]) d \\
& \quad + (-1)^{|x|(|y|+|z|+|t|)+|t|(|y|+|z|)} \rho(x) \rho(t) \rho(y) c - (-1)^{|x|(|y|+|z|+|t|)+|t|(|y|+|z|)+|y||z|} \rho(x) \rho(t) \rho(z) b, \\
& [[z+c, t+d]_{A \oplus V}, x+a]_{A \oplus V}, y+b]_{A \oplus V} \\
& = [[[z, t], x], y] + \rho([[z, t], x]) b - (-1)^{|y|(|x|+|z|+|t|)} \rho(y) \rho([z, t]) a \\
& \quad + (-1)^{|y|(|x|+|z|+|t|)+|x|(|z|+|t|)} \rho(y) \rho(x) \rho(z) d - (-1)^{|y|(|x|+|z|+|t|)+|x|(|z|+|t|)+|z||t|} \rho(y) \rho(x) \rho(t) c, \\
& [[t+d, x+a]_{A \oplus V}, y+b]_{A \oplus V}, z+c]_{A \oplus V} \\
& = [[[t, x], y], z] + \rho([[t, x], y]) c - (-1)^{|z|(|x|+|y|+|t|)} \rho(z) \rho([t, x]) b \\
& \quad + (-1)^{|z|(|x|+|y|+|t|)+|y|(|x|+|t|)} \rho(z) \rho(y) \rho(t) a - (-1)^{|z|(|x|+|y|+|t|)+|y|(|x|+|t|)+|x||t|} \rho(z) \rho(y) \rho(x) d.
\end{aligned}$$

Then, $(A \oplus V, [\cdot, \cdot]_{A \oplus V})$ is a Malcev superalgebra if and only if

$$\begin{aligned}
\rho([[x, y], z]) &= \rho(x) \rho(y) \rho(z) - (-1)^{|z|(|x|+|y|)} \rho(z) \rho(x) \rho(y) \\
&\quad + (-1)^{|x|(|y|+|z|)} \rho(y) \rho([z, x]) \\
&\quad - (-1)^{|x|(|y|+|z|)} \rho([y, z]) \rho(x). \quad \square
\end{aligned}$$

Proposition 2.3. Let $(V, \mathbf{l}, \mathbf{r})$ be a bimodule of an alternative superalgebra (A, \star) . Then, $(V, \mathbf{l} - (-1)^{|x||y|} \mathbf{r})$ is a representation of the Malcev admissible superalgebra $(A, [\cdot, \cdot])$ defined in Theorem 2.1.

Proof. By Proposition 2.1, $A \ltimes_{\mathbf{l}, \mathbf{r}} V$ is an alternative superalgebra. Consider its associated Malcev superalgebra $(A \oplus V, \widehat{[\cdot, \cdot]})$,

$$\begin{aligned}
\widehat{[x+a, y+b]} &= (x+a) \circ (y+b) - (-1)^{|x||y|} (y+b) \circ (x+a) \\
&= x \star y + \mathbf{l}(x)b + \mathbf{r}(y)a - (-1)^{|x||y|} y \star x - (-1)^{|a||y|} \mathbf{l}(y)a - (-1)^{|x||b|} \mathbf{r}(x)b \\
&= [x, y] + (\mathbf{l} - (-1)^{|x||b|} \mathbf{r})(x)b - (-1)^{|a||y|} (\mathbf{l} - (-1)^{|a||y|} \mathbf{r})(y)a.
\end{aligned}$$

According to Proposition 2.2, $(V, \mathbf{l} - (-1)^{|x||y|} \mathbf{r})$ is a representation of $(A, [\cdot, \cdot])$. \square

Example 2.1. Let $A = A_{\bar{0}} \oplus A_{\bar{1}}$ be a Malcev superalgebra. It is easy to see that the map $ad : A \rightarrow gl(A)$ defined by $ad(x)(y) = [x, y], \forall x, y \in \mathcal{H}(A)$, is a Malcev representation of A in itself. It is called the adjoint representation of A . Besides, this representation induces a Malcev representation $ad : A_{\bar{0}} \rightarrow gl(A_{\bar{1}})$ of the Malcev algebra $A_{\bar{0}}$ in the vector space $A_{\bar{1}}$ defined by $ad(x)(y) = [x, y], \forall x \in A_{\bar{0}}, y \in A_{\bar{1}}$. It is called the adjoint representation of $A_{\bar{0}}$ in $A_{\bar{1}}$.

To articulate the adjoint representation and co-adjoint representation of a Malcev superalgebra, we first present some basic facts about \mathbb{Z}_2 -graded vector spaces. Let $V = V_{\bar{0}} \oplus V_{\bar{1}}$ be a \mathbb{Z}_2 -graded

vector space over a field \mathbb{K} . Then $V^* = \text{Hom}(V, \mathbb{K})$ is the dual vector space of V , whose \mathbb{Z}_2 -graduation is given by

$$V_\alpha^* = \{a^* \in V^* \mid a^*(V_{\alpha+1}) = \{0\}\}, \quad \forall \alpha \in \mathbb{Z}_2. \quad (2.13)$$

Let $\langle \cdot, \cdot \rangle : V^* \times V \rightarrow \mathbb{K}$ be the canonical pairing, which allows us to identify V with V^* by

$$\langle a^*, b \rangle = (-1)^{|a^*||b|} \langle b, a^* \rangle, \quad \forall a^* \in V^*, b \in V. \quad (2.14)$$

We extend $\langle \cdot, \cdot \rangle$ to $(V \otimes V)^* \times (V \otimes V)$ ($(V \otimes V)^* = V^* \otimes V^*$ when V is finite-dimensional) by setting

$$\langle a_1^* \otimes a_2^*, b_1 \otimes b_2 \rangle = (-1)^{|a_2^*||b_1|} \langle a_1^*, b_1 \rangle \langle a_2^*, b_2 \rangle, \quad \forall a_1^*, a_2^* \in V^*, b_1, b_2 \in V.$$

Proposition 2.4. *Let (V, ρ) be a representation of a Malcev superalgebra A and let V^* denote the dual space of V . Define a linear map $\rho^* : A \rightarrow \text{gl}(V^*)$ by*

$$\langle \rho^*(x)a^*, b \rangle = -(-1)^{|x||a|} \langle a^*, \rho(x)b \rangle, \quad \forall x \in A, a^* \in V^*, b \in V. \quad (2.15)$$

Then (V^*, ρ^*) is also a Malcev representation of A in V^* , which is called the dual representation of (V, ρ) .

Proof. For any homogenous $x, y, z \in A$, $a^* \in V^*$, $b \in V$, we have

$$\begin{aligned} \langle \rho^*([x, y], z)a^*, b \rangle &= -(-1)^{(|x|+|y|+|z|)|a|} \langle a^*, \rho([x, y], z)b \rangle, \\ \langle \rho^*(x)\rho^*(y)\rho^*(z)a^*, b \rangle &= -(-1)^{(|x|+|y|+|z|)|a|+|x||y|+|x||z|+|y||z|} \langle a^*, \rho(z)\rho(y)\rho(x)b \rangle, \\ \langle \rho^*(z)\rho^*(x)\rho^*(y)a^*, b \rangle &= -(-1)^{(|x|+|y|+|z|)|a|+|x||y|+|x||z|+|y||z|} \langle a^*, \rho(y)\rho(x)\rho(z)b \rangle, \\ \langle \rho^*(y)\rho^*([z, x])a^*, b \rangle &= (-1)^{(|x|+|y|+|z|)|a|+|x||y|+|y||z|} \langle a^*, \rho([z, x])\rho(y)b \rangle, \\ \langle \rho^*([y, z])\rho^*(x)a^*, b \rangle &= (-1)^{(|x|+|y|+|z|)|a|+|x||y|+|x||z|} \langle a^*, \rho(x)\rho([y, z])b \rangle. \end{aligned}$$

Then we have for the map ρ^* ,

$$\begin{aligned} \langle \rho^*([x, y], z)a^*, b \rangle &= \langle (\rho^*(x)\rho^*(y)\rho^*(z) - (-1)^{|z|(|x|+|y|)}\rho^*(z)\rho^*(x)\rho^*(y) \\ &\quad + (-1)^{|x|(|y|+|z|)}\rho^*(y)\rho^*([z, x]) - (-1)^{|x|(|y|+|z|)}\rho^*([y, z])\rho^*(x))a^*, b \rangle, \end{aligned}$$

what is equivalent to say that ρ^* is a Malcev representation of A . \square

Example 2.2. *Let $A = A_{\bar{0}} \oplus A_{\bar{1}}$ be a Malcev superalgebra and A^* its dual vector space. We easily show that the map $\text{ad}^* : A \rightarrow \text{gl}(A^*)$ defined by*

$$\langle \text{ad}^*(x)\alpha, y \rangle = -(-1)^{|x||\alpha|} \langle \alpha, \text{ad}(x)y \rangle, \quad \forall x, y \in \mathcal{H}(A), \alpha \in A^*,$$

is a Malcev representation of A in A^* . It is called the co-adjoint representation of A . Moreover, it is also clear that $\text{ad}^* : A_{\bar{0}} \rightarrow \text{gl}(A_{\bar{1}}^*)$ of the Malcev algebra $A_{\bar{0}}$ in the vector space $A_{\bar{1}}^*$ defined by $\langle \text{ad}^*(x)\alpha, y \rangle = -\langle \alpha, \text{ad}(x)y \rangle, \forall x \in A_{\bar{0}}, y \in A_{\bar{1}}, \alpha \in A_{\bar{1}}^*.$ It is called the co-adjoint representation of $A_{\bar{0}}$ in $A_{\bar{1}}^*$.

3 Super \mathcal{O} -operators of Malcev superalgebras and pre-Malcev superalgebras

In this section, we introduce the notion of super \mathcal{O} -operator and pre-Malcev superalgebras. Then we study the relations among Malcev superalgebras, pre-Malcev superalgebras and pre-alternative superalgebras.

Definition 3.1. A super \mathcal{O} -operator of alternative superalgebra (A, \star) with respect to the bimodule $(V, \mathbf{l}, \mathbf{r})$ is a linear map $T : V \rightarrow A$ such that, for all $a, b \in \mathcal{H}(V)$,

$$T(a) \star T(b) = T(\mathbf{l}(T(a))b + \mathbf{r}(T(b))a). \quad (3.16)$$

A super Rota-Baxter operator of weight 0 on an alternative superalgebra (A, \star) is just a super \mathcal{O} -operator associated to the bimodule (A, L, R) , where L and R are the left and right multiplication operators corresponding to the multiplication \star .

Definition 3.2. Let A be a Malcev superalgebra and (V, ρ) a representation of A . A linear map $T : V \rightarrow A$ with $|T| = 0$ is called a super \mathcal{O} -operator of A associated to (V, ρ) if T satisfies

$$[T(a), T(b)] = T(\rho(T(a))b - (-1)^{|a||b|}\rho(T(b))a), \forall a, b \in \mathcal{H}(V). \quad (3.17)$$

In particular, if \mathcal{R} is a super \mathcal{O} -operator of A associated to the adjoint representation (A, ad) , then \mathcal{R} is called a super Rota-Baxter operator (of weight 0) on A , that is, \mathcal{R} satisfies

$$[\mathcal{R}(x), \mathcal{R}(y)] = \mathcal{R}([\mathcal{R}(x), y] + (-1)^{|x||y|}[x, \mathcal{R}(y)]),$$

for all $x, y \in \mathcal{H}(A)$.

Proposition 3.1. Let T be a super \mathcal{O} -operator of an alternative superalgebra (A, \star) associated to a bimodule $(V, \mathbf{l}, \mathbf{r})$. Then, T is a super \mathcal{O} -operator of $(A, [\cdot, \cdot])$ with respect to $(V, \mathbf{l} - (-1)^{|x||y|}\mathbf{r})$.

Proof. T is a super \mathcal{O} -operator of $(A, [\cdot, \cdot])$ with respect to $(V, \mathbf{l} - (-1)^{|x||y|}\mathbf{r})$ since

$$\begin{aligned} [T(a), T(b)] &= T(a) \star T(b) - (-1)^{|a||b|}T(b) \star T(a) \\ &= T(\mathbf{l}(T(a))b + \mathbf{r}(T(b))a) - (-1)^{|a||b|}T(\mathbf{l}(T(b))a + \mathbf{r}(T(a))b) \\ &= T((\mathbf{l} - (-1)^{|a||b|}\mathbf{r})(T(a))b - (-1)^{|a||b|}(\mathbf{l} - (-1)^{|a||b|}\mathbf{r})(T(b))a). \end{aligned} \quad \square$$

Definition 3.3. A pre-Malcev superalgebra A is a super vector space $A = A_{\bar{0}} \oplus A_{\bar{1}}$ equipped with a bilinear product $(x, y) \rightarrow x \cdot y$ satisfying

$$A_\alpha \cdot A_\beta \subseteq A_{\alpha+\beta}, \quad \alpha, \beta \in \mathbb{Z}_2,$$

and the following equations ($\forall x, y, z, t \in \mathcal{H}(A)$):

$$\begin{aligned} &(-1)^{|x|(|y|+|z|)}[y, z] \cdot (x \cdot t) + [[x, y], z] \cdot t + (-1)^{|x||y|}y \cdot ([x, z] \cdot t) \\ &\quad - x \cdot (y \cdot (z \cdot t)) + (-1)^{|z|(|x|+|y|)}z \cdot (x \cdot (y \cdot t)) = 0, \end{aligned} \quad (3.18)$$

where $[x, y] = x \cdot y - (-1)^{|x||y|}y \cdot x$. The identity (3.18) is equivalent to $PM(x, y, z, t) = 0$, where

$$\begin{aligned} PM(x, y, z, t) &= (-1)^{|x|(|y|+|z|)}(y \cdot z) \cdot (x \cdot t) - (-1)^{|x|(|y|+|z|)+|y||z|}(z \cdot y) \cdot (x \cdot t) \\ &\quad + ((x \cdot y) \cdot z) \cdot t - (-1)^{|x||y|}((y \cdot x) \cdot z) \cdot t \\ &\quad - (-1)^{(|x|+|y|)|z|}(z \cdot (x \cdot y)) \cdot t + (-1)^{|x||y|+(|x|+|y|)|z|}(z \cdot (y \cdot x)) \cdot t \\ &\quad + (-1)^{|x||y|}y \cdot ((x \cdot z) \cdot t) - (-1)^{|x|(|y|+|z|)}y \cdot ((z \cdot x) \cdot t) \\ &\quad - x \cdot (y \cdot (z \cdot t)) + (-1)^{|z|(|x|+|y|)}z \cdot (x \cdot (y \cdot t)), \end{aligned} \quad (3.19)$$

for all $x, y, z, t \in \mathcal{H}(A)$.

Theorem 3.1. Let (A, \cdot) be a pre-Malcev superalgebra.

(i) Let us define, for any homogeneous elements $x, y \in A$ the operation

$$[x, y] = x \cdot y - (-1)^{|x||y|} y \cdot x. \quad (3.20)$$

Then $(A, [\cdot, \cdot])$ is a Malcev superalgebra, which is called the sub-adjacent Malcev superalgebra of (A, \cdot) and denoted by A^C . We call (A, \cdot) a compatible pre-Malcev superalgebra structure on the Malcev superalgebra.

(ii) Let L_x (for any $x \in \mathcal{H}(A)$) denote the left multiplication operator, i.e., $L_x(y) = x \cdot y, \forall y \in \mathcal{H}(A)$. Then $L: A \rightarrow A$ with $x \rightarrow L_x$ gives a representation of the Malcev superalgebra A , that is,

$$L_{[[x, y], z]} = L_x L_y L_z - (-1)^{|z|(|x|+|y|)} L_z L_x L_y + (-1)^{|x|(|y|+|z|)} L_y L_{[z, x]} - (-1)^{|x|(|y|+|z|)} L_{[y, z]} L_x,$$

for all $x, y, z \in \mathcal{H}(A)$.

Proof. For part (i) we have, for any $x, y, z, t \in \mathcal{H}(A)$,

$$\begin{aligned} & (-1)^{|y||z|} [[x, z], [y, t]] - [[[x, y], z], t] - (-1)^{|x|(|y|+|z|+|t|)} [[[y, z], t], x] \\ & \quad - (-1)^{(|x|+|y|)(|z|+|t|)} [[[z, t], x], y] - (-1)^{|t|(|x|+|y|+|z|)} [[[t, x], y], z] \\ & = (-1)^{|y||z|} [x, z] \cdot (y \cdot t) - (-1)^{|y|(|z|+|t|)} [x, z] \cdot (t \cdot y) - (-1)^{|y||x|+|t|(|x|+|z|)} [y, t] \cdot (x \cdot z) \\ & \quad + (-1)^{|x|(|y|+|z|+|t|)+|t||z|} [y, t] \cdot (z \cdot x) - [[x, y], z] \cdot t + (-1)^{|t|(|x|+|y|+|z|)} t \cdot ([x, y] \cdot z) \\ & \quad - (-1)^{|t|(|x|+|y|+|z|)+(|x|+|y|)|z|} t \cdot (z \cdot (x \cdot y)) + (-1)^{|t|(|x|+|y|+|z|)+x||y|+(|x|+|y|)|z|} t \cdot (z \cdot (y \cdot x)) \\ & \quad - (-1)^{|x|(|y|+|z|+|t|)} [[y, z], t] \cdot x + x \cdot ([y, z] \cdot t) - (-1)^{|t|(|y|+|z|)} x \cdot (t \cdot (y \cdot z)) \\ & \quad + (-1)^{|t|(|y|+|z|)+|y||z|} x \cdot (t \cdot (z \cdot y)) - (-1)^{(|x|+|y|)(|z|+|t|)} [[z, t], x] \cdot y \\ & \quad + (-1)^{|x|(|y|+|z|+|t|)} y \cdot ([z, t] \cdot x) - (-1)^{|x||y|} y \cdot (x \cdot (z \cdot t)) \\ & \quad + (-1)^{|x|(|y|+|z|+|t|)+|z||t|} y \cdot (x \cdot (t \cdot z)) \\ & \quad - (-1)^{|t|(|x|+|y|+|z|)} [[t, x], y] \cdot z + (-1)^{(|x|+|y|)(|z|+|t|)} z \cdot ([t, x] \cdot y) \\ & \quad - (-1)^{|x|(|y|+|z|+|t|)+|y||z|} z \cdot (y \cdot (t \cdot x)) + (-1)^{|x||y|+(|x|+|y|)|z|} z \cdot (y \cdot (x \cdot t)) \\ & = HPM(x, t, y, z) + HPM(y, x, z, t) + HPM(z, y, t, x) + HPM(t, z, x, y) = 0. \end{aligned}$$

For (ii) we compute

$$\begin{aligned} & \left(L_{[[x, y], z]} - L_x L_y L_z + (-1)^{|z|(|x|+|y|)} L_z L_x L_y - (-1)^{|x|(|y|+|z|)} L_y L_{[z, x]} \right. \\ & \quad \left. + (-1)^{|x|(|y|+|z|)} L_{[y, z]} L_x \right) (t) \\ & = [[x, y], z] \cdot t - x \cdot (y \cdot (z \cdot t)) + (-1)^{|z|(|x|+|y|)} z \cdot (x \cdot (y \cdot t)) \\ & \quad - (-1)^{|x|(|y|+|z|)} y \cdot ([z, x] \cdot t) + (-1)^{|x|(|y|+|z|)} [y, z] \cdot (x \cdot t) \\ & = PM(x, y, z, t) = 0. \end{aligned} \quad \square$$

Proposition 3.2. A superalgebra (A, \cdot) is a pre-Malcev superalgebra if and only if $(A, [\cdot, \cdot])$ with the commutator $[x, y] = x \cdot y - (-1)^{|x||y|} y \cdot x$ is a Malcev superalgebra and L is a representation of $(A, [\cdot, \cdot])$.

Proof. It follows from Theorem 3.1 above and the definitions of Malcev superalgebra and representation of a Malcev superalgebra. \square

Furthermore, we can construct pre-Malcev superalgebras from super \mathcal{O} -operators of Malcev superalgebras.

Theorem 3.2. *Let $(A, [\cdot, \cdot])$ be a Malcev superalgebra and (V, ρ) be its representation. Let $T : V \rightarrow A$ be a super \mathcal{O} -operator associated to (V, ρ) . Then there exists a pre-Malcev superalgebra structure on V given by*

$$a \cdot b = \rho(T(a))b, \quad \forall a, b \in A. \quad (3.21)$$

Proof. Let $a, b, c, d \in V$. Set $x = T(a), y = T(b); z = T(c)$ and $[a, b] = a \cdot b - (-1)^{|a||b|}b \cdot a$. Hence, we have

$$\begin{aligned} & (-1)^{|a|(|b|+|c|)}[b, c] \cdot (a \cdot d) + [[a, b], c] \cdot d + (-1)^{|a||b|}b \cdot ([a, c] \cdot d) \\ & \quad - a \cdot (b \cdot (c \cdot d)) + (-1)^{|c|(|a|+|b|)}c \cdot (a \cdot (b \cdot d)) \\ & = (-1)^{|a|(|b|+|c|)}(\rho(T(b))c - (-1)^{|b||c|}\rho(T(c))b) \cdot (\rho(T(a))d) + [\rho(T(a))b - (-1)^{|a||b|}\rho(T(b))a, c] \cdot d \\ & \quad + (-1)^{|a||b|}b \cdot ((\rho(T(a))c - (-1)^{|a||c|}\rho(T(c))a) \cdot d) - a \cdot (b \cdot \rho(T(c))d) \\ & \quad + (-1)^{|c|(|a|+|b|)}c \cdot (a \cdot \rho(T(b))d) \\ & = (-1)^{|a|(|b|+|c|)}\rho(T(\rho(T(b))c - (-1)^{|b||c|}\rho(T(c)))b)\rho(T(a))d + (\rho(T(\rho(T(a))b \\ & \quad - (-1)^{|a||b|}\rho(T(b))a))c - (-1)^{|c|(|a|+|b|)}\rho(T(c))(\rho(T(a))b \\ & \quad - (-1)^{|a||b|}\rho(T(b))a)) \cdot d + (-1)^{|a||b|}b \cdot (\rho(T(\rho(T(a))c - (-1)^{|a||c|}\rho(T(c))a))d) \\ & \quad - a \cdot (\rho(T(b))\rho(T(c))d) + (-1)^{|c|(|a|+|b|)}c \cdot (\rho(T(a))\rho(T(b))d) \\ & = (-1)^{|a|(|b|+|c|)}\rho([T(b), T(c)])\rho(T(a))d + \rho\left(T\left(\rho([T(a), T(b)])c - (-1)^{|c|(|a|+|b|)}\rho(T(c))(\rho(T(a))b\right.\right. \\ & \quad \left.\left. - (-1)^{|a||b|}\rho(T(b))a\right)\right)d + (-1)^{|a||b|}\rho(T(b))\rho([T(a), T(c)])d - \rho(T(a))\rho(T(b))\rho(T(c))d \\ & \quad + (-1)^{|c|(|a|+|b|)}\rho(T(c))\rho(T(a))\rho(T(b))d \\ & = (-1)^{|a|(|b|+|c|)}\rho([T(b), T(c)])\rho(T(a))d + \rho([[T(a), T(b)], T(c)])d + (-1)^{|a||b|}\rho(T(b))\rho([T(a), T(c)])d \\ & \quad - \rho(T(a))\rho(T(b))\rho(T(c))d + (-1)^{|c|(|a|+|b|)}\rho(T(c))\rho(T(a))\rho(T(b))d \\ & = (-1)^{|x|(|y|+|z|)}\rho([y, z])\rho(x)d + \rho([[x, y], z])d + (-1)^{|x||y|}\rho(y)\rho([x, z])d \\ & \quad - \rho(x)\rho(y)\rho(z)d + (-1)^{|z|(|x|+|y|)}\rho(z)\rho(x)\rho(y)d = 0. \end{aligned}$$

Thus, (V, \cdot) is a pre-Malcev superalgebra. \square

Therefore, there exists a Malcev superalgebra structure on V given by (3.20) and T is a homomorphism of Malcev superalgebras. Furthermore, there is an induced pre-Malcev superalgebra structure on $T(V)$ given by

$$T(a) \cdot T(b) = T(a \cdot b), \quad \forall a, b \in V. \quad (3.22)$$

Moreover, the corresponding associated Malcev superalgebra structure on $T(V)$ given by (3.20) is just a Malcev supersubalgebra structure of A and T becomes a homomorphism of pre-Malcev superalgebra.

Proposition 3.3. *Let $(A, [\cdot, \cdot])$ be a Malcev superalgebra. Then there exists a compatible pre-Malcev superalgebra structure on A if and only if there exists an invertible super \mathcal{O} -operator on A associated to a representation (V, ρ) .*

Proof. Let (A, \cdot) be a pre-Malcev superalgebra and $(A, [\cdot, \cdot])$ be the associated Malcev superalgebra. Then the identity map $id : A \rightarrow A$ is an invertible super \mathcal{O} -operator on $(A, [\cdot, \cdot])$ associated to (A, ad) .

Conversely, suppose that there exists an invertible super \mathcal{O} -operator T of $(A, [\cdot, \cdot])$ associated to a representation (V, ρ) . Then, using Theorem 3.2, there is a pre-Malcev superalgebra structure on $T(V) = A$ given by

$$T(a) \cdot T(b) = T(\rho(T(a))b), \quad \forall a, b \in \mathcal{H}(V).$$

If we set $x = T(a)$ and $y = T(b)$, then we get

$$x \cdot y = T(\rho(x)T^{-1}(y)), \quad \forall x, y \in \mathcal{H}(A).$$

□

An obvious consequence of Theorem 3.2 is the following construction of a pre-Malcev superalgebra in terms of super Rota-Baxter operator on a Malcev superalgebra.

Corollary 3.1. *Let $(A, [\cdot, \cdot])$ be a Malcev superalgebra and the linear map $\mathcal{R} : A \rightarrow A$ is a super Rota-Baxter operator. Then there exists a pre-Malcev superalgebra structure on A given by*

$$x \cdot y = [\mathcal{R}(x), y], \quad \forall x, y \in \mathcal{H}(A).$$

If in addition, \mathcal{R} is invertible, then there is a compatible pre-Malcev superalgebra structure on A given by

$$x \cdot y = \mathcal{R}([x, \mathcal{R}^{-1}(y)]), \quad \forall x, y \in \mathcal{H}(A).$$

Let us recall some basic notions about bilinear forms of a Malcev superalgebra $(A, [\cdot, \cdot])$. A bilinear form $\omega : A \otimes A \rightarrow \mathbb{K}$ is said to be

1. supersymmetric if $\omega(x, y) = (-1)^{|x||y|}\omega(y, x), \forall x, y \in \mathcal{H}(A)$,
2. skew supersymmetric if $\omega(x, y) = -(-1)^{|x||y|}\omega(y, x), \forall x, y \in \mathcal{H}(A)$,
3. nondegenerate if $x \in A$ satisfies $\omega(x, y) = 0, \forall y \in \mathcal{H}(A)$, then $x = 0$,
4. invariant if

$$\omega([x, y], z) = \omega(x, [y, z]), \quad \forall x, y, z \in \mathcal{H}(A). \tag{3.23}$$

Definition 3.4. *Let $(A, [\cdot, \cdot])$ be a Malcev superalgebra. A skew supersymmetric, nondegenerate bilinear form $\omega : A \otimes A \rightarrow \mathbb{K}$ is called a super two-cocycle on A if ω satisfies*

$$(-1)^{|x||z|}\omega(x, [y, z]) + (-1)^{|y||x|}\omega(y, [z, x]) + (-1)^{|z||y|}\omega(z, [x, y]), \tag{3.24}$$

$\forall x, y, z \in \mathcal{H}(A)$. Then ω is called symplectic. A Malcev superalgebra $(A, [\cdot, \cdot])$ with a symplectic form is called a symplectic Malcev superalgebra and denoted $(A, [\cdot, \cdot], \omega)$.

Let $(A, [\cdot, \cdot], \omega)$ be a symplectic Malcev superalgebra. Define the multiplication "·" on A by

$$\omega(x \cdot y, z) = (-1)^{|x|(|y|+|z|)}\omega(y, [z, x]). \tag{3.25}$$

Theorem 3.3. Under the above notations, there exists a compatible pre-Malcev superalgebra structure “ \cdot ” on A given by Eq. (3.25).

Proof. Define the linear map $T : A \rightarrow A^*$ by $\langle T(x), y \rangle = \omega(x, y)$. Since $(A, [\cdot, \cdot])$ is a Malcev superalgebra, (A^*, ad^*) is a representation of A . By the fact that ω is skew supersymmetric and using (3.24), we obtain that T is an invertible super \mathcal{O} -operator associated to the representation (A^*, ad^*) . By Proposition 3.3, there exists a compatible pre-Malcev superalgebra structure given by

$$x \cdot y = T^{-1}(ad^*(x)T(y)).$$

Hence,

$$\begin{aligned} \omega(x \cdot y, z) &= \langle T(x \cdot y), z \rangle = \langle ad^*(x)T(y), z \rangle = -(-1)^{|x||y|} \langle T(y), [x, z] \rangle \\ &= -(-1)^{|x||y|} \omega(y, [x, z]) = (-1)^{|x|(|y|+|z|)} \omega(y, [z, x]). \end{aligned}$$

□

Analogous to the connection between alternative superalgebras and Malcev superalgebras, pre-alternative superalgebras are closely related to pre-Malcev superalgebras.

Definition 3.5. A pre-alternative superalgebra A is a super vector space $A = A_{\bar{0}} \oplus A_{\bar{1}}$ equipped with two bilinear maps $\prec, \succ : A \otimes A \rightarrow A$, satisfying for all $x, y, z \in \mathcal{H}(A)$ and $x \star y = x \prec y + x \succ y$,

$$(x \star y) \succ z - x \succ (y \succ z) + (-1)^{|x||y|} (y \star x) \succ z - (-1)^{|x||y|} y \succ (x \succ z) = 0, \quad (3.26)$$

$$(x \prec y) \prec z - x \prec (y \prec z) + (-1)^{|y||z|} (x \prec z) \prec y - (-1)^{|y||z|} x \prec (z \star y) = 0, \quad (3.27)$$

$$(x \succ y) \prec z - x \succ (y \prec z) + (-1)^{|x||y|} (y \prec x) \prec z - (-1)^{|x||y|} y \prec (x \star z) = 0, \quad (3.28)$$

$$(x \succ y) \prec z - x \succ (y \prec z) + (-1)^{|y||z|} (x \star z) \succ y - (-1)^{|y||z|} x \succ (z \succ y) = 0. \quad (3.29)$$

Proposition 3.4. Let (A, \prec, \succ) be a pre-alternative superalgebra. Then the product $x \star y = x \prec y + x \succ y$ defines an alternative superalgebra A .

Theorem 3.4. Let $T : V \rightarrow A$ be a super \mathcal{O} -operator of alternative superalgebra (A, \star) with respect to the bimodule $(V, \mathbf{l}, \mathbf{r})$. Then (V, \prec, \succ) is a pre-alternative superalgebra, where for all $a, b \in \mathcal{H}(V)$,

$$a \succ b = \mathbf{l}(T(a))b \quad \text{and} \quad a \prec b = \mathbf{r}(T(b))a. \quad (3.30)$$

Moreover, if (V, \cdot) is the pre-Malcev superalgebra associated to the Malcev admissible superalgebra $(A, [\cdot, \cdot])$ on the representation $(V, \mathbf{l} - (-1)^{|x||y|} \mathbf{r})$, then $a \cdot b = a \succ b - (-1)^{|a||b|} b \prec a$.

Proof. For any $a, b, c \in \mathcal{H}(V)$, using (2.8) and (3.16) yields

$$\begin{aligned} (a \succ b) \prec c - a \succ (b \prec c) + (b \prec a) \prec c - b \prec (a \star c) \\ &= \mathbf{r}(T(c))\mathbf{l}(T(a))b - \mathbf{l}(T(a))\mathbf{r}(T(c))b \\ &\quad + (-1)^{|a||b|} \mathbf{r}(T(c))\mathbf{r}(T(a))b - (-1)^{|a||b|} \mathbf{r}(T(a \succ c + a \prec c))b \\ &= \mathbf{r}(T(c))\mathbf{l}(T(a))b - \mathbf{l}(T(a))\mathbf{r}(T(c))b \\ &\quad + (-1)^{|a||b|} \mathbf{r}(T(c))\mathbf{r}(T(a))b - (-1)^{|a||b|} \mathbf{r}(T(\mathbf{l}(T(a))c + \mathbf{r}(T(c))a))b \\ &= \mathbf{r}(T(c))\mathbf{l}(T(a))b - \mathbf{l}(T(a))\mathbf{r}(T(c))b \\ &\quad + (-1)^{|a||b|} \mathbf{r}(T(c))\mathbf{r}(T(a))b - (-1)^{|a||b|} \mathbf{r}(T(a) \star T(c)) = 0. \end{aligned}$$

The other identities for (V, \prec, \succ) being a pre-alternative superalgebras can be verified similarly. Moreover, using Eqs. (3.21) and (3.30),

$$a \cdot b = (\mathbf{l} - (-1)^{|a||b|} \mathbf{r})(T(a))b = \mathbf{l}(T(a))b - (-1)^{|a||b|} \mathbf{r}(T(a))b = a \succ b - (-1)^{|a||b|} b \prec a. \quad \square$$

Corollary 3.2. Let (A, \star) be an alternative superalgebra and $\mathcal{R} : A \rightarrow A$ be a super Rota-Baxter operator of weight 0. If multiplications \prec and \succ on A are defined for all $x, y \in \mathcal{H}(A)$ by $x \prec y = x \star \mathcal{R}(y)$ and $x \succ y = \mathcal{R}(x) \star y$, then (A, \prec, \succ) is a pre-alternative superalgebra.

Moreover, if (A, \cdot) is the pre-Malcev superalgebra associated to the Malcev admissible superalgebra $(A, [\cdot, \cdot])$, then $x \cdot y = x \succ y - (-1)^{|x||y|} y \prec x$.

Summarizing the above study, we have the following commutative diagram of categories:

$$\begin{array}{ccc}
\text{Pre-alternative} & \xleftarrow{\substack{\text{super } R-B \\ x \prec y + x \succ y}} & \text{Alternative} \\
\text{superalg.} & & \text{superalg.} \\
\downarrow & & \downarrow \\
x \succ y - (-1)^{|x||y|} y \prec x & & x \star y - (-1)^{|x||y|} y \star x \\
\downarrow & & \downarrow \\
\text{Pre-Malcev} & \xleftarrow{\substack{\text{super } R-B \\ x \cdot y - (-1)^{|x||y|} y \cdot x}} & \text{Malcev} \\
\text{superalg.} & & \text{superalg.}
\end{array} \tag{3.31}$$

4 An analogue of the super CYBE on Malcev superalgebras

We derive a close relation between super \mathcal{O} -operators and solutions to the super Malcev Yang-Baxter equations on Malcev superalgebras. We prove that under the skew-supersymmetric condition, a solution to the super MYBE in a Malcev superalgebra is equivalent to a super \mathcal{O} -operator associated to the co-adjoint representation. Further, we give an explicit constructing way to exploit solutions to the super MYBE in semidirect product Malcev superalgebras from a common given super \mathcal{O} -operator.

4.1 Super \mathcal{O} -operators and operator forms of the super MYBE

Let $(A, [\cdot, \cdot])$ be a Malcev superalgebra and $r \in A \otimes A$. The standard form of the super Malcev Yang-Baxter equation in A is given in the tensor expression as follows:

$$[r_{12}, r_{13}] + [r_{12}, r_{23}] + [r_{13}, r_{23}] = 0, \tag{4.32}$$

where r is called a solution of the super MYBE and for $r = \sum_i x_i \otimes y_i \in A^{\otimes 2}$,

$$r_{12} = \sum_i x_i \otimes y_i \otimes 1, \quad r_{13} = \sum_i x_i \otimes 1 \otimes y_i, \quad r_{23} = \sum_i 1 \otimes x_i \otimes y_i. \tag{4.33}$$

Note that Eq. (4.32) has the same form as the ordinary case, but the commutation relation is in the graded spaces, that is,

$$[r_{12}, r_{13}] = \sum_{i,j} (-1)^{|x_j||y_i|} [x_i, x_j] \otimes y_i \otimes y_j, \tag{4.34}$$

$$[r_{13}, r_{23}] = \sum_{i,j} x_i \otimes x_j \otimes [y_i, y_j], \tag{4.35}$$

$$[r_{12}, r_{23}] = \sum_{i,j} (-1)^{|x_j||y_i|} x_i \otimes [y_i, x_j] \otimes y_j. \tag{4.36}$$

Let V be a super vector space. Then $r = \sum_i x_i \otimes y_i \in A^{\otimes 2}$ is said to be **skew-supersymmetric** (resp. **supersymmetric**) if $r = -\sigma(r)$ (resp. $r = \sigma(r)$). Furthermore, r can be regarded as a map from V^* to V in the following way

$$\begin{aligned} \langle x^* \otimes y^*, r \rangle &= \langle x^* \otimes y^*, \sum_i x_i \otimes y_i \rangle = \sum_i (-1)^{|y^*||x_i|} \langle x^*, x_i \rangle \langle y^*, y_i \rangle \\ &= (-1)^{|r||x^*|} \langle y^*, r(x^*) \rangle, \quad \forall x^*, y^* \in A^*. \end{aligned} \quad (4.37)$$

Eq. (4.32) gives the tensor form of super MYBE. What we will do next is to replace the tensor form by a linear operator satisfying some conditions.

Theorem 4.1. *Let $(A, [\cdot, \cdot])$ be a Malcev superalgebra and $r \in A \otimes A$ be skew-supersymmetric with $|r| = 0$. Then r is a solution of the super MYBE in A if and only if r satisfies*

$$[r(x^*), r(y^*)] = r(ad^*r(x^*)(y^*) - (-1)^{|x^*||y^*|}ad^*r(y^*)(x^*)), \quad \forall x^*, y^* \in A^*. \quad (4.38)$$

So there is a pre-Malcev superalgebra structure on A^* given by

$$x \cdot y = ad^*r(x^*)(y^*), \quad \forall x^*, y^* \in A^*. \quad (4.39)$$

Proof. Let $\{e_1, \dots, e_m, f_1, \dots, f_n\}$ be a basis of A where $\{e_1, \dots, e_m\}$ is a basis of $A_{\bar{0}}$ and $\{f_1, \dots, f_n\}$ is a basis of $A_{\bar{1}}$ and let $\{e_1^*, \dots, e_m^*, f_1^*, \dots, f_n^*\}$ be its dual basis with $e_i^* \in A_{\bar{0}}^*$ and $f_j^* \in A_{\bar{1}}^*$. Since r is skew-supersymmetric and $|r| = 0$, we can set

$$r = \sum_{i,j}^m a_{ij} e_i \otimes e_j + \sum_{k,l}^n b_{kl} f_k \otimes f_l,$$

where $a_{ij} = -a_{ji}$ and $b_{kl} = b_{lk}$. Now, we have

$$[r_{12}, r_{13}] = \left[\sum_{i,j}^m a_{ij} e_i \otimes e_j \otimes 1 + \sum_{k,l}^n b_{kl} f_k \otimes f_l \otimes 1, \sum_{p,q}^m a_{pq} e_p \otimes 1 \otimes e_q + \sum_{s,t}^n b_{st} f_s \otimes 1 \otimes f_t \right]$$

$$\begin{aligned} &= \sum_{i,j,k,l,p}^m C_{ip}^a a_{ij} a_{pq} e_a \otimes e_j \otimes e_q + \sum_{i,j}^m \sum_{s,t,b}^n C_{is}^b a_{ij} b_{st} f_b \otimes e_j \otimes f_t \\ &+ \sum_{p,q}^m \sum_{k,l,c}^n C_{kp}^c b_{kl} a_{pq} f_c \otimes f_l \otimes e_q - \sum_d^m \sum_{k,l,s,t}^n C_{ks}^d b_{kl} b_{st} e_d \otimes f_l \otimes f_t, \end{aligned}$$

$$[r_{12}, r_{23}] = \left[\sum_{i,j}^m a_{ij} e_i \otimes e_j \otimes 1 + \sum_{k,l}^n b_{kl} f_k \otimes f_l \otimes 1, \sum_{p,q}^m 1 \otimes a_{pq} e_p \otimes e_q + \sum_{s,t}^n 1 \otimes b_{st} f_s \otimes f_t \right]$$

$$\begin{aligned} &= \sum_{i,j,k,l,p}^m C_{jp}^a a_{ij} a_{pq} e_i \otimes e_a \otimes e_q + \sum_{i,j}^m \sum_{s,t,b}^n C_{js}^b a_{ij} b_{st} e_i \otimes f_b \otimes f_t \\ &+ \sum_{p,q}^m \sum_{k,l,c}^n C_{lp}^c b_{kl} a_{pq} f_k \otimes f_c \otimes e_q + \sum_d^m \sum_{k,l,s,t}^n C_{ps}^d b_{kl} b_{st} f_k \otimes e_d \otimes f_t, \end{aligned}$$

$$[r_{13}, r_{23}] = \left[\sum_{i,j}^m a_{ij} e_i \otimes 1 \otimes e_j + \sum_{k,l}^n b_{kl} f_k \otimes 1 \otimes f_l, \sum_{p,q}^m 1 \otimes a_{pq} e_p \otimes e_q + \sum_{s,t}^n 1 \otimes b_{st} f_s \otimes f_t \right]$$

$$\begin{aligned}
&= \sum_{i,j,k,l,p}^m C_{jq}^a a_{ij} a_{pq} e_i \otimes e_p \otimes e_a + \sum_{i,j}^m \sum_{s,t,b}^n C_{jt}^b a_{ij} b_{st} e_i \otimes f_s \otimes f_b \\
&+ \sum_{p,q}^m \sum_{k,l,c}^n C_{lq}^c b_{kl} a_{pq} f_k \otimes e_p \otimes f_c - \sum_d^m \sum_{k,l,s,t}^n C_{lt}^d b_{kl} b_{st} f_k \otimes f_s \otimes e_d,
\end{aligned}$$

where C_{ij}^k 's are the structure coefficients of Malcev superalgebra A on the basis $\{e_1, \dots, e_m, f_1, \dots, f_n\}$. Then r is a solution of the super MYBE in $(A, [\cdot, \cdot])$ if and only if (for any $1 \leq a, d \leq m$ and $1 \leq b, c \leq n$)

$$\begin{aligned}
&\sum_{i,p} \left(C_{ip}^a a_{ij} a_{pq} + C_{pi}^q a_{ip} a_{ji} + C_{ip}^a a_{ai} a_{pq} \right) e_a \otimes e_j \otimes e_q \\
&+ \left(\sum_i^m \sum_s^n C_{is}^b a_{ij} b_{st} + \sum_{l,s}^n C_{ls}^j b_{bl} b_{st} + \sum_i^m \sum_s^n C_{is}^t a_{ij} b_{bs} \right) f_b \otimes e_j \otimes f_t \\
&+ \left(\sum_p^m \sum_k^n C_{kp}^c a_{pq} b_{kl} + \sum_p^m \sum_k^n C_{kp}^l b_{ck} a_{pq} - \sum_{k,t}^n C_{kt}^q b_{ck} b_{lt} \right) f_c \otimes f_l \otimes e_q \\
&+ \left(- \sum_d^m \sum_{k,s}^n C_{ks}^d b_{kl} b_{st} + \sum_j^m \sum_s^n C_{js}^l a_{dj} b_{st} + \sum_j^m \sum_s^n C_{js}^t a_{dj} b_{ls} \right) e_d \otimes f_l \otimes f_t = 0.
\end{aligned}$$

On the other hand, by (4.37), we get

$$r(e_j^*) = - \sum_{i=1}^m a_{ji} e_i, \quad r(f_k^*) = - \sum_{s=1}^n b_{ks} f_s, \quad 1 \leq j \leq m, 1 \leq k \leq n.$$

We prove the conclusion in the following four cases:

Case (1) $x^* = e_j^*$ and $y^* = e_q^*$. Then by Eq.(4.38), we have:

$$\sum_{i,p} \left(C_{ip}^a a_{ij} a_{pq} + C_{pi}^q a_{ip} a_{ji} + C_{ip}^a a_{ai} a_{pq} \right) e_a = 0.$$

Case (2) $x^* = e_q^*$ and $y^* = f_c^*$. Then by Eq.(4.38), we have:

$$\left(\sum_p^m \sum_k^n C_{kp}^c a_{pq} b_{kl} + \sum_p^m \sum_k^n C_{kp}^l b_{ck} a_{pq} - \sum_{k,t}^n C_{kt}^q b_{ck} b_{lt} \right) f_l = 0.$$

Case (3) $x^* = f_b^*$ and $y^* = e_j^*$. Then by Eq.(4.38), we have:

$$\left(\sum_i^m \sum_s^n C_{is}^b a_{ij} b_{st} + \sum_{l,s}^n C_{ls}^j b_{bl} b_{st} + \sum_i^m \sum_s^n C_{is}^t a_{ij} b_{bs} \right) f_t = 0.$$

Case (4) $x^* = f_l^*$ and $y^* = f_t^*$. Then by Eq.(4.38), we have:

$$\left(- \sum_d^m \sum_{k,s}^n C_{ks}^d b_{kl} b_{st} + \sum_j^m \sum_s^n C_{js}^l a_{dj} b_{st} + \sum_j^m \sum_s^n C_{js}^t a_{dj} b_{ls} \right) e_d = 0.$$

Therefore, it is easy to see that r is a solution of the super MYBE in A if and only if r satisfies (4.38).

Thus the first half part of the conclusion holds. It is easy to get the other results. \square

Corollary 4.1. Let $(A, [\cdot, \cdot])$ be a Malcev superalgebra and $r \in A \otimes A$. Assume r is skew-supersymmetric with $|r| = 0$ and there exists a nondegenerate symmetric invariant bilinear form B on A . Define a linear map $\varphi : A \rightarrow A^*$ by $\langle \varphi(x), y \rangle = B(x, y)$ for any $x, y \in A$. Then r is a solution of the super MYBE in A if and only if $\tilde{r} = r\varphi : A \rightarrow A$ is a Rota-Baxter operator associated to the module (A, ad) , that is, \tilde{r} satisfies the following equation

$$[\tilde{r}(x), \tilde{r}(y)] = \tilde{r}([\tilde{r}(x), y] + [x, \tilde{r}(y)]), \forall x, y \in A.$$

Hence there is a pre-Malcev superalgebra structure on A given by

$$x \cdot y = [\tilde{r}(x), y], \quad \forall x, y \in A. \quad (4.40)$$

Proof. For any $x, y \in A$, we have

$$\begin{aligned} \langle \varphi(ad(x)y), z \rangle &= B([x, y], z) = (-1)^{|z|(|x|+|y|)} B(z, [x, y]) = -(-1)^{|x||y|} B(y, [x, z]) \\ &= -(-1)^{|x||y|} \langle \varphi(y), ad(x)z \rangle = \langle ad^*(x)\varphi(y), z \rangle. \end{aligned}$$

Hence $\varphi(ad(x)y) = ad^*(x)\varphi(y)$ for any $x, y \in A$. Let $x^* = \varphi(x)$, $y^* = \varphi(y)$, then by Theorem 4.1, r is a solution of the super MYBE in A if and only if

$$[r\varphi(x), r\varphi(y)] = [r(x^*), r(y^*)] = r(ad^*r(x^*)(y^*) - (-1)^{|x||y|}ad^*r(y^*)(x^*)) = r\varphi([r\varphi(x), y] + [x, r\varphi(y)]).$$

Therefore the conclusion holds. \square

4.2 Constructions of solutions of the super MYBE in Malcev superalgebras

The following result establishes a close relation between super \mathcal{O} -operators and the super classical Yang-Baxter equation in Malcev superalgebras.

Theorem 4.2. Let $(A, [\cdot, \cdot])$ be a Malcev superalgebra. Let $\rho^* : A \rightarrow gl(V^*)$ be the dual representation of the representation $\rho : A \rightarrow gl(V)$ of the Malcev superalgebra A . Let $T : V \rightarrow A$ be a linear map which can be identified as an element in $(A \ltimes_{\rho^*} V^*) \otimes (A \ltimes_{\rho^*} V^*)$. Then T is a super \mathcal{O} -operator of A associated to (V, ρ) , with $|T| = 0$, if and only if $r = T - \sigma(T)$ is a skew-supersymmetric solution of the super MYBE in $A \ltimes_{\rho^*} V^*$.

Proof. Let $\{e_1, \dots, e_m, f_1, \dots, f_n\}$ be a basis of A as in the proof of Theorem 4.1. Let $\{u_1, \dots, u_s, v_1, \dots, v_t\}$ be a basis of V and $\{u_1^*, \dots, u_s^*, v_1^*, \dots, v_t^*\}$ be the dual basis in a similar way. Since $|T| = 0$, we set

$$T(u_i) = \sum_{a=1}^m a_{ia} e_a, \quad i = 1, \dots, s, \quad T(v_k) = \sum_{b=1}^n b_{kb} e_b, \quad k = 1, \dots, t.$$

Moreover, since $\text{Hom}(V, A) \cong A \otimes V^*$, we have

$$\begin{aligned} T &= \sum_{i=1}^s T(u_i) \otimes u_i^* + \sum_{k=1}^t T(v_k) \otimes v_k^* = \sum_{i=1}^s \sum_{a=1}^m a_{ia} e_a \otimes u_i^* + \sum_{k=1}^t \sum_{b=1}^n b_{kb} e_b \otimes v_k^* \\ &\in A \otimes V^* \subset (A \ltimes_{\rho^*} V^*) \otimes (A \ltimes_{\rho^*} V^*). \end{aligned}$$

Therefore,

$$r = T - \sigma(T) = \sum_{i=1}^s T(u_i) \otimes u_i^* + \sum_{k=1}^t T(v_k) \otimes v_k^* - \sum_{i=1}^s u_i^* \otimes T(u^i) - \sum_{k=1}^t v_k^* \otimes T(v_k).$$

Thus

$$\begin{aligned}
[r_{12}, r_{13}] &= \sum_{i,p=1}^s \{ [T(u_i), T(u_p)] \otimes u_i^* \otimes u_p * -\rho^*(T(u_i))u_p^* \otimes u_i^* \otimes T(u_p) \\
&\quad + \rho^*(T(u_p))u_i^* \otimes T(u_i) \otimes u_p^* \} \\
&\quad + \sum_{i=1}^s \sum_{q=1}^t \{ [T(u_i), T(v_q)] \otimes u_i^* \otimes v_q * -\rho^*(T(u_i))v_q^* \otimes u_i^* \otimes T(v_q) \\
&\quad + \rho^*(T(v_q))u_i^* \otimes T(u_i) \otimes v_q^* \} \\
&\quad + \sum_{k,q=1}^t \{ [T(v_k), T(v_q)] \otimes v_k^* \otimes v_q * -\rho^*(T(v_k))v_q^* \otimes v_k^* \otimes T(v_q) \\
&\quad + \rho^*(T(v_q))v_k^* \otimes T(v_k) \otimes v_q^* \} \\
&\quad + \sum_{p=1}^s \sum_{k=1}^t \{ [T(v_k), T(u_p)] \otimes v_k^* \otimes u_p * -\rho^*(T(v_k))u_p^* \otimes v_k^* \otimes T(u_p) \\
&\quad + \rho^*(T(u_p))v_k^* \otimes T(v_k) \otimes u_p^* \}, \\
\\
[r_{12}, r_{23}] &= \sum_{i,p=1}^s \{ -T(u_i) \otimes \rho^*(T(u_p))u_i^* \otimes u_p^* - u_i^* \otimes [T(u_i), T(u_p)] \otimes u_p^* \\
&\quad + u_i^* \otimes \rho^*(T(u_i))u_p^* \otimes T(u_p) \} \\
&\quad + \sum_{i=1}^s \sum_{q=1}^t \{ -T(u_i) \otimes \rho^*(T(v_q))u_i^* \otimes v_q^* - u_i^* \otimes [T(u_i), T(v_q)] \otimes v_q^* \\
&\quad + u_i^* \otimes \rho^*(T(u_i))v_q^* \otimes T(v_q) \} \\
&\quad + \sum_{k,q=1}^t \{ -T(v_k) \otimes \rho^*(T(v_q))v_k^* \otimes v_q^* - v_k^* \otimes [T(v_k), T(v_q)] \otimes v_q^* \\
&\quad + v_k^* \otimes \rho^*(T(v_k))v_q^* \otimes T(v_q) \} \\
&\quad + \sum_{p=1}^s \sum_{k=1}^t \{ -T(v_k) \otimes \rho^*(T(u_p))v_k^* \otimes u_p^* - v_k^* \otimes [T(v_k), T(u_p)] \otimes u_p^* \\
&\quad + v_k^* \otimes \rho^*(T(v_k))u_p^* \otimes T(u_p) \}, \\
\\
[r_{13}, r_{23}] &= \sum_{i,p=1}^s \{ T(u_i) \otimes u_p^* \otimes \rho^*(T(u_p))u_i^* - u_i^* \otimes T(u_p) \otimes \rho^*(T(u_i))u_p^* \\
&\quad + u_i^* \otimes u_p^* \otimes [T(u_i), T(u_p)] \} \\
&\quad + \sum_{i=1}^s \sum_{q=1}^t \{ T(u_i) \otimes v_q^* \otimes \rho^*(T(v_q))u_i^* - u_i^* \otimes T(v_q) \otimes \rho^*(T(u_i))v_q^* \\
&\quad + u_i^* \otimes v_q^* \otimes [T(u_i), T(v_q)] \} \\
&\quad + \sum_{k,q=1}^t \{ T(v_k) \otimes v_q^* \otimes \rho^*(T(v_q))v_k^* - v_k^* \otimes T(v_q) \otimes \rho^*(T(v_k))v_q^* \\
&\quad + v_k^* \otimes v_q^* \otimes [T(v_k), T(v_q)] \} \\
&\quad + \sum_{p=1}^s \sum_{k=1}^t \{ T(v_k) \otimes u_p^* \otimes \rho^*(T(u_p))v_k^* - v_k^* \otimes T(u_p) \otimes \rho^*(T(v_k))u_p^*
\end{aligned}$$

$$+v_k^* \otimes u_p^* \otimes [T(v_k), T(u_p)]\}.$$

On the other hand, from the definition of dual representation, we know

$$\rho^*(T(u_p))u_i^* = -\sum_{j=1}^s u_i^*(\rho(T(u_p))u_j)u_j^*.$$

Thus

$$\begin{aligned} -\sum_{i,p=1}^s T(u_i) \otimes \rho^*(T(u_p))u_i^* \otimes u_p^* &= \sum_{i,p=1}^s T(u_i) \otimes [\sum_{j=1}^s u_i^*(\rho(T(u_p))u_j)u_j^*] \otimes u_p^* \\ &= \sum_{i,p=1}^s \sum_{j=1}^s u_j^*(\rho(T(u_p))u_i)T(u_j) \otimes u_i^* \otimes u_p^* \\ &= \sum_{i,p=1}^s T(\sum_{j=1}^s (u_j^*(\rho(T(u_p))u_i)u_j)) \otimes u_i^* \otimes u_p^* \\ &= \sum_{i,p=1}^s T(\rho(T(u_p))u_i) \otimes u_i^* \otimes u_p^*. \end{aligned}$$

Therefore

$$\begin{aligned} &[r_{12}, r_{13}] + [r_{12}, r_{23}] + [r_{13}, r_{23}] \\ &= \sum_{i,p=1}^s \{([T(u_i), T(u_p)] + T(\rho(T(u_p))u_i) - T(\rho(T(u_i))u_p)) \otimes u_i^* \otimes u_p^* \\ &\quad - u_i^* \otimes ([T(u_i), T(u_p)] + T(\rho(T(u_p))u_i) - T(\rho(T(u_i))u_p)) \otimes u_p^* \\ &\quad + u_i^* \otimes u_p^* \otimes ([T(u_i), T(u_p)] + T(\rho(T(u_p))u_i) - T(\rho(T(u_i))u_p))\} \\ &\quad + \sum_{i=1}^s \sum_{q=1}^t \{([T(u_i), T(v_q)] + T(\rho(T(v_q))u_i) - T(\rho(T(u_i))v_q)) \otimes u_i^* \otimes v_q^* \\ &\quad + u_i^* \otimes (-[T(u_i), T(v_q)] - T(\rho(T(v_q))u_i) + T(\rho(T(u_i))v_q)) \otimes v_q^* \\ &\quad - u_i^* \otimes v_q^* \otimes (-[T(u_i), T(v_q)] - T(\rho(T(v_q))u_i) + T(\rho(T(u_i))v_q))\} \\ &\quad + \sum_{k,q=1}^t \{([T(v_k), T(v_q)] + T(\rho(T(v_q))v_k) - T(\rho(T(v_k))v_q)) \otimes v_k^* \otimes v_q^* \\ &\quad + v_k^* \otimes (-[T(v_k), T(v_q)] - T(\rho(T(v_q))v_k) + T(\rho(T(v_k))v_q)) \otimes v_q^* \\ &\quad - v_k^* \otimes v_q^* \otimes (-[T(v_k), T(v_q)] + T(\rho(T(v_k))v_q) - T(\rho(T(v_q))v_k))\} \\ &\quad + \sum_{p=1}^s \sum_{k=1}^t \{([T(v_k), T(u_p)] + T(\rho(T(u_p))v_k) - T(\rho(T(v_k))u_p)) \otimes v_k^* \otimes u_p^* \\ &\quad - v_k^* \otimes ([T(v_k), T(u_p)] + T(\rho(T(u_p))v_k) - T(\rho(T(v_k))u_p)) \otimes u_p^* \\ &\quad + v_k^* \otimes u_p^* \otimes ([T(v_k), T(u_p)] + T(\rho(T(u_p))v_k) - T(\rho(T(v_k))u_p))\}. \end{aligned}$$

Obviously, r is a solution of super MYBE in $A \bowtie_{\rho^*} V^*$ if and only if T is a super \mathcal{O} -operator associated to (V, ρ) . \square

Proposition 4.1. *Let $(A, [\cdot, \cdot])$ be a Malcev superalgebra and $r \in A \otimes A$ be skew-supersymmetric and nondegenerate with $|r| = 0$. Then r is a solution of the super MYBE in A if and only if the*

bilinear form ω on A given by

$$\omega(x, y) = \langle r^{-1}(x), y \rangle, \forall x, y \in A, \quad (4.41)$$

is a symplectic form.

Proof. Since r is nondegenerate, for $x, y, z \in A$, there exist $a^*, b^*, c^* \in A^*$ such that $x = r(a^*), y = r(b^*), z = r(c^*)$.

In the one hand, using the fact that r is skew-supersymmetric, then

$$\begin{aligned} \omega(x, y) &= \langle r^{-1}(x), y \rangle = \langle a^*, r(b^*) \rangle = -(-1)^{|a^*||b^*|} \langle b^*, r(a^*) \rangle \\ &= -(-1)^{|x||y|} \langle r^{-1}(y), x \rangle = -(-1)^{|x||y|} \omega(y, x). \end{aligned}$$

On the other hand, by Eq.(3.24) and the definition of the dual representation ad^* of the adjoint representation, we obtain

$$\begin{aligned} & (-1)^{|x||z|} \omega(x, [y, z]) + (-1)^{|y||x|} \omega(y, [z, x]) + (-1)^{|z||y|} \omega(z, [x, y]) \\ &= (-1)^{|x||z|} \langle r^{-1}(x), [y, z] \rangle + (-1)^{|y||x|} \langle r^{-1}(y), [z, x] \rangle + (-1)^{|z||y|} \langle r^{-1}(z), [x, y] \rangle \\ &= (-1)^{|a^*||c^*|} \langle a^*, [r(b^*), r(c^*)] \rangle + (-1)^{|b^*||a^*|} \langle b^*, [r(c^*), r(a^*)] \rangle + (-1)^{|c^*||b^*|} \langle c^*, [r(a^*), r(b^*)] \rangle \\ &= -(-1)^{|a^*|(|c^*|+|b^*|)} \langle ad^*(r(b^*))a^*, r(c^*) \rangle + (-1)^{|c^*||a^*|} \langle ad^*(r(a^*))b^*, r(c^*) \rangle + (-1)^{|c^*||b^*|} \langle c^*, [r(a^*), r(b^*)] \rangle \\ &= (-1)^{(|a^*|+|c^*|)|b^*|} \langle c^*, r(ad^*(r(b^*))a^*) \rangle - (-1)^{|c^*||b^*|} \langle c^*, r(ad^*(r(a^*))b^*) \rangle + (-1)^{|c^*||b^*|} \langle c^*, [r(a^*), r(b^*)] \rangle \\ &= (-1)^{|c^*||b^*|} \langle c^*, (-1)^{|a^*||b^*|} r(ad^*(r(b^*))a^*) - r(ad^*(r(a^*))b^*) + [r(a^*), r(b^*)] \rangle. \end{aligned}$$

Hence ω is a symplectic form if and only if r is a solution of the super MYBE in A . \square

Corollary 4.2. Let (A, \cdot) be a pre-Malcev superalgebra. Let $\{e_1, \dots, e_m, f_1, \dots, f_n\}$ be a basis of A where $e_i \in A_{\bar{0}}$ and $f_j \in A_{\bar{1}}$ and let $\{e_1^*, \dots, e_m^*, f_1^*, \dots, f_n^*\}$ be its dual basis with $e_i^* \in A_{\bar{0}}^*$ and $f_k^* \in A_{\bar{1}}^*$. Then

$$r = \sum_{i=1}^m (e_i \otimes e_i^* - e_i^* \otimes e_i) + \sum_{j=1}^n (f_j \otimes f_j^* + f_j^* \otimes f_j), \quad (4.42)$$

is a solution of the super MYBE in $A^C \ltimes_{ad^*} (A^C)^*$.

Proof. It follows from the fact that $T = id$ is a super \mathcal{O} -operator of A^C associated to the representation (A, ad) . \square

References

- [1] Abd el Malek, A. : On the Frattini subalgebra of a Malcev algebra. Arch. Math. (Basel), **37** (1981), 306-315.
- [2] Albuquerque, H., Elduque, A.: Classification of Mal'tsev superalgebras of small dimensions. Algebra i Logika, 35 (1996), 629-654.
- [3] Albuquerque, H.: Malcev superalgebras, in: S. Gonzalez (Ed.), Non-Associative Algebras and Its Applications, Kluwer Academic Publishers, Dordrecht, (1994).
- [4] Andruskiewitsch, N.: Lie superalgebras and Poisson-Lie supergroups, Abh. Math. Sem. Univ. Hamburg 63 (1993) 147–163.

- [5] Bai, C. M.: A unified algebraic approach to classical Yang-Baxter equation, *J. Phys. A: Math. Theor.* **40** (2007), 11073-11082.
- [6] Belavin, A. A. and Drinfeld, V. G. : Solutions of the classical Yang-Baxter equation for simple Lie algebras, *Funct. Anal. Appl.*, **16** (1982), 159-180.
- [7] Berezin, F. A. , Kac, G. I.: *Math. Sbornik* 82 (Russian), (1970), 343-351.
- [8] Bracken, A. J. , Gould, M. D. and Zhang, R. B.: Quantum supergroups and solutions of the Yang-Baxter equation, *Modern Phys. Lett. A* 5 (1990) 831–840.
- [9] Faddeev, L. D., Takhtajan, L.: The quantum inverse scattering method of the inverse problem and the Heisenberg XYZ model, *Russ. Math. Surv.* 34 (1979) 11–68.
- [10] Faddeev L. D., Takhtajan, L.: Hamiltonian methods in the theory of solitons (Springer, Berlin, 1987).
- [11] Floreanini, R. , Leites, D. A. and Vinet, L.: On the defining relations of quantum superalgebras, *Lett. Math. Phys.* 23 (1991) 127–131.
- [12] Gould, M. D. , Zhang, R. B. and Bracken, A. J.: Lie bi-super algebras and the graded classical Yang-Baxter equation, *Rev. Math. Phys.* 3 (1991) 223–240.
- [13] Harrathi, F., S. Mabrouk, O. Ncib, and S. Silvestrov. Malcev Yang-Baxter equation, weighted \mathcal{O} -operators on Malcev algebras and post-Malcev algebras. *Hacettepe J. Maths. Stats* (2022): 1-21.
- [14] Harrathi, F., Mabrouk,S., Ncib, O. and Silvestrov, S.: Kupershmidt operators on Hom-Malcev algebra. *Int. J. Geom. Methods Mod. Phys.* 20, no. 03 (2023): 2350046.
- [15] Harrathi, F., Mabrouk, S., Ncib, O. and Silvestrov, S.,: Dendrification of Hom-Malcev algebras. arXiv preprint arXiv:2205.11002(2022)(appear in Advances in Applied Clifford Algebras)(2023).
- [16] Kerdman, F. S.: Analytic Moufang loops in the large, *Alg. Logic* **18** (1980), 325-347.
- [17] Kuzmin, E. N.: The connection between Malcev algebras and analytic Moufang loops, *Alg. Logic*, **10** (1971), 1-14.
- [18] Kuzmin, E. N.: Malcev algebras and their representations, *Alg. Logic*, **7** (1968), 233-244.
- [19] Li, H.:Local Systems of Vertex Operators, Vertex Superalgebras and Modules, *J. Pure Appl. Algebra*, 109 (1996), 143-195.
- [20] Madariaga, S.: Splitting of operations for alternative and Malcev structures, *Comm. Algebra*, **45**(1)(2014), 183-197.
- [21] Mahaligesshwara, N.: A note on Malcev algebras and quasi-Lie algebras. *Yokohama Math. J.* 22 (1974), 25-29.
- [22] Malcev, A. I.: Analytic loops, *Mat. Sb.* **36**(3)(1955), 569-576.
- [23] Sagle, A. A.: Malcev algebras, *Trans. Amer. Math. Soc.* **101** (1961), 426-458.

- [24] Semenov-Tian-Shansky, M. A. : Quantum and classical Integrable systems, Lect. Not. Phys., Vol. 495 (Springer-Verlag, Berlin, 1997), pp. 314–377.
- [25] Shestakov, I. P.: Simple Mal'tsev superalgebras. (Russian) Mat. Sb., 182 (1991), 1357- 1366; translation in Math. USSR-Sb. 74 (1993), 101-110.
- [26] Wang, Y., Hou, D. and Bai, C., 2010. Operator forms of the classical Yang–Baxter equation in Lie superalgebras. International Journal of Geometric Methods in Modern Physics, 7(04), pp.583-597.
- [27] Yamane, H.: Universal R-matrices for quantum groups associated to simple Lie superalgebras, Proc. Japan Acad. A 67 (1991) 108–112.
- [28] Yamane, H.: Quantized enveloping algebra associated with simple Lie superalgebras and their universal R-matrices, Publ. RIMS. Kyoto Univ. 30 (1994) 15–87.
- [29] Zhang, R. B. , Gould , M. D. and Bracken, A. J.: Solutions of the graded classical Yang–Baxter equation and integrable models, J. Phys. A: Math. Gen. 24 (1991) 1185–1197.