CLASSICAL IDEALS THEORY OF MAXIMAL SUBRINGS IN NON-COMMUTATIVE RINGS

ALBORZ AZARANG

Department of Mathematics, Faculty of Mathematical Sciences and Computer, Shahid Chamran University of Ahvaz, Ahvaz-Iran a_azarang@scu.ac.ir

ORCID ID: orcid.org/0000-0001-9598-2411

ABSTRACT. Let R be a maximal subring of a ring T. In this paper we study relation between some important ideals in the ring extension $R \subseteq T$. In fact, we would like to find some relation between $Nil_*(R)$ and $Nil_*(T)$, $Nil^*(R)$ and $Nil^*(T)$, J(R) and J(T), $Soc(_RR)$ and $Soc(_RT)$, and finally $Z(_RR)$ and $Z(_RT)$; especially, in certain cases, for example when T is a reduced ring, R (or T) is a left Artinian ring, or R is a certain maximal subring of T. We show that either $Soc(_RR) = Soc(_RT)$ or $(R : T)_r$ (the greatest right ideal of T which is contained in R) is a left primitive ideal of R. We prove that if T is a reduced ring, then either $Z(_RT) = 0$ or $Z(_RT)$ is a minimal ideal of T, $T = R \oplus Z(_RT)$, and $(R : T) = (R : T)_l = (R : T)_r = ann_R(Z(_RT))$. If $T = R \oplus I$, where I is an ideal of T, then we completely determine relation between Jacobson radicals, lower nilradicals, upper nilradicals, socle and singular ideals of R and T. Finally, we study the relation between previous ideals of R and T when either R or T is a left Artinian ring.

1. INTRODUCTION

1.1. Motivation. Let $R \subseteq T$ be an arbitrary ring extension. Then there are many important results between certain ideals of R and T, such as prime ideals, maximal ideals, primitive ideals, Jacobson radical and lower/upper nilradicals, especially in certain cases such as commutative ring extension, integral extension of commutative rings, polynomial extension of a ring, matrix extension, finitely generated extension, central or normal extension, see [16, 20, 25, 23]. In this paper we would like to study these ideals when R is a maximal subrings of a ring T, especially in certain cases that we mentioned in the abstract of the paper. If T is a commutative ring and R is a maximal subring of T, then it is clear that $N(R) = N(T) \cap R$, where N(T) is the set of all nilpotent elements of a ring T. Also note that since in commutative case, either R is integrally closed in T or T is integral over R (i.e., T is a finitely generated R-module), then we immediately infer that if R is a maximal subring of T and T is integral over R, then $J(R) = J(T) \cap R$. In [9, Section. 1 and Section. 2 of Chapter IV], the author proved some properties between certain ideals of R and T in a minimal ring extension $R \subseteq T$ of commutative rings (these extensions are called adjacent extension in [9]). Also see [?], for some other algebraic properties which are shared between a commutative ring and its maximal subrings. In [12] the authors studied minimal ring extensions of non-commutative rings. In fact, the authors generalized the results of [10] for non-commutative rings and characterized exactly (central) minimal ring extension of prime rings and therefore simple rings, see [12, Theorem 5.1, Corollary 5.3 and Theorem 6.1]. Also, they proved that a field K has a minimal ring extension of the form $\mathbb{M}_n(D)$ where D is a centrally finite division ring and n > 1, if and only if K has a proper subfield of finite index, see [12, Lemma 6.6]. Finally, in [2] the author studied the existence of maximal subrings of non-commutative rings, and in [3], the conductor ideals of maximal subrings in non-commutative rings. We will mention some of needed results from [2] and [3] in the preliminaries subsection.

1.2. Review of the results. In the next section, we first consider the case that R is a maximal subring of a ring T such that $T = R \oplus I$, for an ideal I of T. We prove that in this case $(R : T)_l = r.ann_R(I)$ and $(R : T)_r = l.ann_R(I)$. In particular, if $I^2 \neq 0$ then $(R : T) = (R : T)_l = (R : T)_r$. We also determine when T is a prime, semiprime and left primitive ring. A structure of maximal ideals of T is given too. Next, we prove that if T is a reduced ring and R is a subring of T, then $Soc(_RT) = Soc(T_R)$ is an ideal of T.

²⁰¹⁰ Mathematics Subject Classification. 16S70; 16N20; 16N40; 16N60; 16P20; 16P40.

Key words and phrases. Maximal subrings, socle, singular ideal, Jacobson radical, upper nilradical, lower nilradical, Artinian rings.

In particular, if R is a maximal subring of T, then either Soc(R) = Soc(RT) or there exists a semisimple left (resp. right) R-submodule A (resp. B) of T such that $T = R \oplus A = R \oplus B$, moreover $(R:T)_l$ and $(R:T)_r$ are right primitive and left primitive ideals of R, respectively. Consequently, we prove that if there exists a simple left R-module S of T, such that $S^2 = S \not\subseteq R$, then there exists an idempotent e of T such that $T = R \oplus Re$. We show that if R is a maximal subring of a reduced ring T, then $Z(R) = Z(T_R)$ (which is an ideal of T, say Z) and either Z = 0 or $T = R \oplus Z$, $(R:T) = (R:T)_l = (R:T)_l = ann_R(Z)$ and $Q = ann_T(Z)$ is a minimal prime ideal of T which contract to (R:T) in R. We also prove that if R is a maximal subring of a prime ring T, then either Z(RT) = 0 or Soc(RR) = 0. If R is a prime ring which is a maximal subring of a ring T, then we determine Min(T) and the conductor ideals of $R \subseteq T$. Moreover, if R is a prime ring which is a maximal subring of a ring T and A be the set of all nonzero ideals I of T such that $I \cap R = 0$, then we prove that $|\mathcal{A}| \leq 2$ and $|\mathcal{A}| = 2$ if and only if $T \cong \Delta(R) + A \times A$ for some minimal ideal A of R and in this case Min(T) = A and $(R:T) = (R:T)_l = (R:T)_r = 0$. Conversely, we prove that if R is a ring which is either prime or reduced, A is a minimal ideal of R and $T = \Delta(R) + A \times A$, then for an ideal I of T with $I \cap \Delta(R) = 0$ and $I^2 = I$, we have $I = A \times 0$ or $I = 0 \times A$. If R is a maximal subring of a ring T and $T = R \oplus I$, where I is an ideal of T, then we prove that either $J(R) = J(T) \cap R$, or $I^2 = I, J(T) \subseteq (R:T) = (R:T)_l = (R:T)_r = l.ann_T(I)$, which is a primitive and minimal prime ideal of T, $dim(T) \ge 2$ and I is unique. A similar result holds if we replace Nil* instead of Jacobson radical in the previous result. But fortunately, we always have $Nil_*(R) = Nil_*(T) \cap R$. We also prove that, either $Soc(_RR) = Soc(_RT) \subseteq (R:T)_r$ or I is a semisimple left R-module which all of simple R-submodule of I are isomorphic as R-module, $J(R) = J(T) \cap R$ and $Nil^*(R) = Nil^*(T) \cap R$. In particular, if either R or T is a one-sided Artinian ring, then $J(R) = J(T) \cap R$ and $Nil^*(R) = Nil^*(T) \cap R$. Moreover, if R is a left Artinian ring, then $Soc(_RI) = I$. We prove that if R is a maximal subring of a left Noetherian ring T, then either $Nil^*(T) \subseteq Nil_*(R)$ or $Nil_*(R) = Nil^*(R) = Nil^*(T) \cap R$ is a nilpotent ideal of R. If in addition, T is a left Artinian ring, then either (R:T) is a maximal ideal of T, or R is a zero-dimensional ring with only finitely many prime ideals. Finally we consider the case R is a maximal subring of a ring T and Ris a left Artinian ring. First, we prove that DCC holds on Spec(T) and if R does not contain any prime ideal of T, then T is a zero-dimensional ring with only finitely many maximal ideals. Next, we observe that the algebraic properties of T depend to the properties of $(R:T)_l$ in T. (1) If $(R:T)_l$ is not an ideal of T and $(R:T)_l T = T$, then we prove that ${}_R T$ and $((R:T)_l)_R$ are finitely generated. In particular, _RT is a left Artinian R-module and hence a left Artinian ring. Moreover, either $J(R) = J(T) \cap R$ or $(R:T) \in Max(T), T(R:T)_r = T, Max(R) = \{(R:T)_l, (R:T)_r\} \cup \{P \cap R \mid P \in Max(T), P \neq (R:T)\}$ and |Max(R)| = |Max(T)| + 1. (2) If $(R:T)_l$ is not an ideal of T and $(R:T)_l T$ is a proper ideal of T, then we show that $(R:T)_l T \in Max(T), |Max(T)| \leq |Max(R)|, R$ does not contain any maximal ideal of T, either $(R:T)_l T \in Min(T)$ and $J(R) = J(T) \cap R$ or $(R:T)_l T = r.ann_T(I)$ for some ideal I of T with $I^2 = 0$, and finally either $J(R) = J(T) \cap R$ or $(R:T)_r$ is not a finitely generated right ideal of R. (3) $(R:T)_l$ is an ideal of T, which implies that $(R:T)_l = (R:T)_r = (R:T)$. In particular, we prove that in this case if T is a zero-dimensional ring, then $J(R) = J(T) \cap R$. We show that $T/(R:T) \cong \mathbb{M}_n(S)$, where S is a ring with at most two nonzero proper ideal (that are maximal ideals) and S has a maximal subring which is a division ring. Moreover, either T is a left Artinian ring or S has a unique nonzero proper ideal.

1.3. Notations and Definitions. All rings in this paper are unital and all subrings, modules and homomorphisms are also unital. If $R \subsetneq T$ is a ring extension and there exists no other subring between R and T, then R is called a maximal subring of T, or the extension $R \subseteq T$ is called a minimal ring extension. If T is a ring and R and S be subrings of T, then we can consider (R, S)-subbimodule of T. In particular, if $t \in T$, then the (R, S)-subbimodule of T which is generated by t is denoted by $RtS = \{\sum_{i=1}^{n} r_i ts_i \mid r_i \in R, s_i \in S, n \ge 0\}$. It is clear that if $R \subseteq S$ and $t \in S$, then R + RtS and R + StRare also subrings of T which contains R. In particular, for each $t \in T$, the subrings R + RtT and R + TtRof T contain R. More generally, if I is left (resp. right) ideal of T, then R + IR (resp. R + RI) is a subring of T which contains R. If T is a ring, I is an ideal of T and M is a left (resp. right) T-module, then $Min_T(I)$, $Max_r(T)$, $Max_l(T)$, Max(T), Spec(T), $l.ann_T(M)$ (resp. $r.ann_T(M)$) denote the set of all minimal prime ideals of I in T, the set of all maximal right ideals of T, the set of all maximal left ideals of T, the set of all maximal ideals of T, the set of all prime ideals of T, the left annihilator of M in T (resp. the right annihilator of M in T), respectively. We use Min(T) for $Min_T(0)$. The set of all left (resp. right) primitive ideals of T is denoted by $Prm_l(T)$ (resp. $Prm_r(T)$). The characteristic of a ring T is denoted by Char(T). If T is a ring, then dim(T) denotes the classical Krull dimension of T (i.e., the supremum of the lengths of all chains of prime ideals of T). If T is a ring and M is a left T-module, then $Soc(_TM)$, $Z(_TM)$ and $J(_TM)$ denote the socle of M, the singular submodule of M and the Jacobson radical of M, respectively (we have similar notations for the right T-modules). For a ring T, $Nil_*(T)$ and $Nil^*(T)$ are the lower nilradical and upper nilradical of T, respectively. A prime ideal P of a ring T is called strongly prime if T/P has no nonzero nil ideal. If A is a right ideal of a ring T, then $\mathbb{I}(A)$ denotes the idealizer of A in T, that is the largest subring of T which A is a two-sided ideal in it, i.e., $\mathbb{I}(A) = \{t \in T \mid tA \subseteq A\}$, the similar notation is used when A is a left ideal of T. For the properties of the previous notations and other used definitions and notations in this paper we refer the reader to [20] and [25].

1.4. Preliminaries. In this subsection we review some results from [2] and [3]. First we recall that if $R \subsetneq T$ is a ring extension which is finite as ring (module) over R, then one can easily see that T has a maximal subring which contains R, by a natural use of Zorn's Lemma. Also, a proper subring S of a ring T is maximal if and only if for each $x \in T \setminus S$, we have S[x] = T. Moreover, if R is a proper subring of T and there exists $\alpha \in T$ such that $R[\alpha] = T$, then T has a maximal subring S with $R \subseteq S$ and $\alpha \notin S$. Now let R be a ring and $\Delta(R) := \{(r, r) \mid r \in R\}$, be the diagonal subring of $T := R \times R$. Then it is not hard too see that $\Delta(R)[(1,0)] = T$. Therefore T has a maximal subring which contains $\Delta(R)$. In fact, subrings of T which contains $\Delta(R)$ has the form $\Delta_I(R) := \Delta(R) + I \times I$, where I is an ideal of R. In particular $\Delta_I(R)$ is a maximal subring of T if and only if I is a maximal ideal of R, see [2, Theorem 2.7]. Therefore a ring R is simple if and only if $\Delta(R)$ is a maximal subring of $R \times R$. More generally we have the following useful result.

Theorem 1.1. [2, Theorem 2.9, Corollary 2.10]. Let R_1 and R_2 be rings and $T = R_1 \times R_2$. Then T has a maximal subring if and only if at least one of the following conditions holds:

- (1) R_1 or R_2 has a maximal subring. (2) There exist (maximal) ideals $I_j \subsetneq R_j$, for j = 1, 2, such that $\frac{R_1}{I_1} \cong \frac{R_2}{I_2}$ as rings.

Therefore if R_1, \ldots, R_n are rings $n \ge 2$, then by the above theorem one can easily see that $T = R_1 \times I$ $\cdots \times R_n$ has a maximal subring if and only if either there exists i such that R_i has a maximal subring or there exist $i \neq j$ such and (maximal) ideals $I_k \subsetneq R_k$ for k = i, j such that $\frac{R_i}{I_i} \cong \frac{R_i}{I_i}$ as rings. Moreover if each of R_i is simple and S is a maximal subring of T, then there exists j such that either $R \cong \prod_{i \neq j} R_i$ or $R \cong S_j \times \prod_{i \neq j} R_i$, where S_j is a maximal subring of R_j .

Theorem 1.2. [2, Theorem 2.13]. Let S be a simple ring which is a maximal subring of a ring R, then every nonzero ideal of R is maximal and $|Max(R)| \leq 2$. Moreover, exactly one of the following holds:

- (1) R is simple ring which is a minimal ring extension of S.
- (2) R has distinct maximal ideals M and N such that $S \cong \frac{R}{M} \cong \frac{R}{N}$ as rings, $M \cap N = 0$ and $R \cong S \times S$ as rings.
- (3) R has exactly one nonzero proper ideal $M, S \cong \frac{R}{M}$ as rings and $M^2 = M$. In particular, either R is a two sided primitive ring or J(R) = M.
- (4) R has exactly one nonzero proper ideal M, $S \cong \frac{R}{M}$ as rings and $M^2 = 0$. In particular, J(R) = M.

Let $R \subseteq T$ be a ring extension, then we have three type of conductor ideals: $(R:T) := \{x \in T \mid TxT \subseteq x \in T \mid TxT \subseteq x \in T \mid TxT \in T\}$ R, $(R:T)_l := \{x \in T \mid Tx \subseteq R\}$ and $(R:T)_r := \{x \in T \mid xT \subseteq R\}$. In other words, (R:T) is the largest common ideal between R and T, $(R:T)_l$ (resp. $(R:T)_r$) is the largest common left (resp. right) ideal between R and T. It is clear that $(R:T)_l = r.ann_R((\frac{T}{R})_R)$ and therefore $(R:T)_l$ is a two sided ideal of R. Similarly, $(R:T)_r = l.ann_R(R(T/R))$ is an ideal of R. It is clear that $(R:T)_l(R:T)_r \subseteq (R:T)_l(R:T)_r \subseteq R$ $T \subseteq (R:T)_l \cap (R:T)_r$. In particular, if $R \neq T$, then $(R:T)_l + (R:T)_r \neq T$. Finally it is not hard to see that $(R:T) = r.ann_T(T/(R:T)_r) = l.ann_T(T/(R:T)_l) = l.ann_R(T/(R:T)_l) = r.ann_R(T/(R:T)_r),$ see [3, (6) of Corollary 2.2]. Moreover, we have the following result.

Theorem 1.3. [3, Theorem 2.1]. Let R be a maximal subring of a ring T. Then $(R:T)_l$ and $(R:T)_r$ are prime ideals of R.

Proof. Let $a, b \in R$ and $aRb \subseteq (R : T)_l$. Thus $TaRb \subseteq R$. Now assume that $a \notin (R : T)_l$, i.e., $Ta \not\subseteq R$. Thus $TaR \not\subseteq R$. Since R is a maximal subring of T we conclude that R + TaR = T and thus $Tb = Rb + TaRb \subseteq R$, i.e., $b \in (R:T)_l$. Hence $(R:T)_l$ is a prime ideal of R. Similarly, $(R:T)_r$ is a prime ideal of R. \square

We refer the reader to [3], for more results about the conductor ideals of maximal subrings (of a noncommutative rings).

2. Classical ideals

If R is a (maximal) subring of a ring T, then it is clear that for each (nonzero proper) ideal I of T either $I \cap R = 0$ or $I \cap R \neq 0$. In fact, if R is a maximal subring of a ring T, then one of the following holds:

- (1) there exists a nonzero proper ideal I of R such that $R \cap I = 0$. Therefore $R \oplus I = T$, for R is maximal.
- (2) for each proper nonzero ideal I of R, we have $R \cap I \neq 0$.

We remind the reader that the first part of item (4) of the following theorem proved in [12], we present it again for the sake of completeness and also for the presentation of the statement of it by the notation of conductor ideals.

Theorem 2.1. Let R be a maximal subring of a ring T. Assume that T has a nonzero ideal I such that $R \cap I = 0$. Then the following hold:

- (1) I is a minimal ideal of T. Moreover, I is maximal respect to the property that $R \cap I = 0$.
- (2) $(R:T)_l = r.ann_R(I)$ and $(R:T)_l = l.ann_R(I)$.
- (3) If $I^2 \neq 0$ (in particular if T is semiprime), then $(R:T) = (R:T)_l = (R:T)_r$.
- (4) T is a semiprime ring if and only if R is semiprime and $I^2 \neq 0$. In particular, in this case $Q := ann_T(I)$ is a minimal prime ideal of T and $Q \cap R = (R : T) = (R : T)_l = (R : T)_r$.
- (5) If $I^2 = 0$, then the contraction of prime (resp. semiprime, maximal, left primitive, right primitive, strongly prime) ideals of T to R remain prime (resp. semiprime, maximal, left primitive, right primitive, strongly prime). In particular, $J(R) \subseteq J(T) \cap R$, $Nil_*(R) \subseteq Nil_*(T) \cap R$ and $Nil^*(R) \subseteq Nil^*(T) \cap R$. Moreover U(T) = U(R) + I.
- (6) For each left (resp. right) ideal A of R, $TA \cap R = A$ (resp. $AT \cap R = A$).
- (7) If $I^2 = I$, then for each ideal A of R either $A \subseteq (R:T)$ or IA = AI = I and AT = TA.

Proof. Since R is a maximal subring of T, for each nonzero ideal $J \subseteq I$ of T, we infer that $T = R \oplus J$. This immediately shows that I is a minimal ideal of T. Similarly, I is maximal respect to property that $R \cap I = 0$ and hence (1) holds. As $T = R \oplus I$, we conclude that $T/R \cong I$ as left and right R-modules and therefore $(R:T)_l = r.ann_R(T/R) = r.ann_R(I)$ and $(R:T)_r = l.ann_R(T/R) = l.ann_R(I)$, thus (2) holds. Now assume that $I^2 \neq 0$. First note that $(R:T)_l I$ is an ideal of T which is contained in I. Hence by minimality of I we deduce that either $(R:T)_{l}I = I$ or $(R:T)_{l}I = 0$. If $(R:T)_{l}I = I$, then by (2) (i.e., $I(R:T)_l = 0$), we have $I^2 = ((R:T)_l I)^2 = 0$ which is impossible. Thus $(R:T)_l I = 0$ and hence $(R:T)_l \subseteq (R:T)_r$, by (2). Similarly, $(R:T)_r \subseteq (R:T)_l$ and therefore (3) is true. For (4), first assume that T is semiprime (hence $I^2 \neq 0$). Let N be an ideal of R such that $N^2 = 0$. Thus by (3) and (2), $N \subseteq (R:T) = ann_R(I)$, for $(R:T)_l$ is a prime ideal of R. Thus NI = IN = 0. Since $T = R \oplus I$, we immediately conclude that N is an ideal of T, thus N = 0, for T is semiprime. Hence R is semiprime. Conversely, assume that R is semiprime and $I^2 \neq 0$. We prove that T is semiprime too. Suppose A be a nonzero ideal of T such that $A^2 = 0$. Thus $(A \cap R)^2 = 0$ and therefore $A \cap R = 0$, for R is semiprime. Hence $T = R \oplus A$, by maximality of R. Thus $R \oplus I = T = R \oplus A$. This immediately implies that $I \cong A$ as left and right R-modules. Hence $P := l.ann_R(A) = l.ann_R(I) = r.ann_R(I) = r.ann_R(A)$, by (3) and (2). Since $A^2 = 0$ and $T = R \oplus A$, we can easily see that $l.ann_T(A) = P \oplus A = r.ann_T(A)$. Now note that since I and A are minimal ideals of T, we conclude that either $I \cap A = I$ or $I \cap A = 0$. If $I \cap A = I$, then $I \subseteq A$ and thus $I^2 = 0$ which is absurd. Therefore $I \cap A = 0$, hence AI = IA = 0. Again by $T = R \oplus A$ and AI = IA = 0, we deduce that $l.ann_T(I) = P \oplus A = r.ann_T(I)$. Hence $I \subseteq l.ann_T(A) = l.ann_I(I)$ and therefore $I^2 = 0$ which is a contradiction. Hence T is a semiprime ring. Now assume that T is a semiprime ring. If I is contained in all minimal prime ideals of T, then since T is a semiprime ring we infer that I = 0. Hence there exists a minimal prime ideal Q of T such that $I \not\subseteq Q$. Since I is a minimal ideal of T, we conclude that $I \cap Q = 0$. Hence IQ = QI = 0. From this one can easily see that $Q = ann_T(I)$. Therefore $Q \cap R = ann_R(I)$, which is equal to $(R:T) = (R:T)_l = (R:T)_r$, by (3) and (2). Thus (4) holds. For (5), assume that $I^2 = 0$, let Q be a prime ideal of T, then clearly $I \subseteq Q$. Therefore $Q \nsubseteq R$ and hence R + Q = T, by maximality of R. Therefore $R/(R \cap Q) \cong T/Q$, as rings. This immediately implies the first part of (5). Thus by the first part of (5), we conclude that for each prime (resp. maximal, left primitive, right primitive, strongly prime) ideal Q of T, $R \cap Q$ remains in R with a same property as in T. Thus the inclusions that mentioned in (5) are valid. Finally, note that since $I \subseteq J(T)$, and $T = R \oplus I$ (which implies that $R \cong T/I$ as ring), we deduce that an element t = r + i, where $r \in R$ and $i \in I$, is a unit in T if and only if r is a unit in R. Thus U(T) = U(R) + I and hence we are done for (5). For (6), it is clear that $A \subseteq TA \cap R$. Hence assume that $x \in TA \cap R$. Thus there exist $t_i \in T$ and $a_i \in A$, $1 \leq i \leq n$, such that $x = \sum_{i=1}^{n} t_i a_i$. Since $T = R \oplus I$, we conclude that each t_i has the form $r_i + c_i$, where $r_i \in R$ and $c_i \in I$. Therefore $x - \sum_{i=1}^n r_i a_i = \sum_{i=1}^n c_i a_i \in R \cap I = 0$. Thus $x = \sum_{i=1}^n r_i a_i \in A$ and therefore $A = TA \cap R$. For (7), first note that by (3), $(R : T) = (R : T)_l = (R : T)_r$. Hence $A \nsubseteq (R : T)$ if and only if $TA \nsubseteq R$ if and only if $TA \nsubseteq R$. Now assume that $A \nsubseteq (R : T)$. If AI = 0, then from $T = R \oplus I$, we conclude that $AT = A \subseteq R$, which is absurd. Thus $AI \neq 0$ and similarly $IA \neq 0$. Since AI and IA are contained in $I, I \cap R = 0$ and R is a maximal subring of T, we deduce that $T = R \oplus AI = R \oplus IA$. Again, since $IA, AI \subseteq I$ and $T = R \oplus I$, we deduce that IA = I = AI and therefore $TA = (R \oplus I)A = A \oplus IA = A \oplus AI = A(R \oplus I) = AT$.

note that in (5) of the previous theorem, since I is contained in the ideals J(T), $Nil_*(T)$ and $Nil^*(T)$, then each of them are not contained in R. Therefore by maximality of R we have R + J(T) = T, $R + Nil_*(T) = T$ and $R + Nil^*(T) = T$. Hence $R/(R \cap J(T)) \cong T/J(T)$, $R/(R \cap Nil_*(T)) \cong T/Nil_*(T)$ and $R/(R \cap Nil^*(T)) \cong T/Nil^*(T)$, as rings.

Proposition 2.2. Let R be a maximal subring of a ring T. Assume that T has a nonzero ideal I such that $R \cap I = 0$. Then the following are equivalent:

- (1) R is a prime ring and $l.ann_T(I) = 0$.
- (2) R is a prime ring and $r.ann_T(I) = 0$.
- (3) T is a prime ring and $I \in Spec(T)$.

Moreover, if any of these equivalent conditions holds, then $(R:T) = (R:T)_l = (R:T)_r = 0$.

Proof. We prove $(1) \iff (3)$. The proof of $(2) \iff (3)$ is similar. Since R is a maximal subring of T and I is a nonzero ideal of T which is not contained in R we have $R \oplus I = T$ and therefore $R \cong T/I$ as ring, for $R \cap I = 0$. This immediately implies that $(3) \Longrightarrow (1)$. Hence assume that (1) holds. It is obvious that I is a prime ideal of T for $T/I \cong R$ (as ring) and R is a prime ring. Now assume that A and B are ideals of T such that AB = 0. Since I is a prime ideal of T, the we conclude that either $A \subseteq I$ or $B \subseteq I$. If $A \subseteq I$ and $A \neq 0$, then by minimality of I (by (1) of the previous theorem), we deduce that A = I and therefore IB = 0. From $l.ann_T(I) = 0$, we conclude that $I^2 \neq 0$ and therefore T is a semiprime ring by (4) of the previous theorem. Thus $(BI)^2 = 0$ implies that BI = 0. Therefore B = 0. Similarly if $B \subseteq I$ and $B \neq 0$, we conclude that A = 0. Hence in any cases we deduce that A = 0 or B = 0, i.e., T is prime. The equalities $(R:T) = (R:T)_l = (R:T)_r = 0$ follows from (2) and (3) of the previous theorem.

Lemma 2.3. Let T be a ring and I be a left ideal of T. Then either I has a maximal left T-submodule or $I \subseteq J(T)$. In particular, if J(T) = 0, then each nonzero left ideal of T has a maximal left T-submodule.

Proof. Assume that I is not contained in J(T). Thus there exists a left maximal ideal M of T such that $I \notin M$. Therefore by maximality of M we conclude that M + I = T. Hence $I/(I \cap M) \cong T/M$ as left T-modules. Thus $I \cap M$ is a maximal submodule of I. The final part is evident.

Lemma 2.4. Let T be a ring and I be a nonzero minimal proper ideal of T. Then the following hold:

- (1) If I has a maximal left T-submodule, then $l.ann_T(I)$ is a left primitive ideal of T.
- (2) If T is a left Artinian ring (note that in this case T has a nonzero minimal ideal), then either T is a prime Artinian ring (i.e., $T \cong \mathbb{M}_n(D)$ for some division ring D), or $l.ann_T(I) \neq 0$ is a left primitive ideal.
- (3) If T is a left Noetherian ring, then either T is a left primitive ring, or $l.ann_T(I) \neq 0$ is a left primitive ideal.

Proof. (1) Let N be a maximal left T-submodule of I. Then clearly $P = l.ann_T(I/N)$ is a left primitive ideal of T. If $P = l.ann_T(I)$, then we are done. Hence assume that $PI \neq 0$. By minimality of I, we infer that PI = I. But clearly $PI \subseteq N$. Hence $I \subseteq N$ which is absurd. Thus $P = l.ann_T(I)$ is a left primitive ideal of T. (2) and (3) are immediate consequences of (1), for each cases I has a maximal T-submodule, and therefore $P = l.ann_T(I)$ is a left primitive ideal of T. Thus if P = 0, T is a left primitive ring and hence the first case of (2) or (3) holds, and otherwise $P = l.ann_T(I) \neq 0$.

Proposition 2.5. Let R be a maximal subring of a ring T. Assume that T has a nonzero ideal I such that $R \cap I = 0$. Then the following are equivalent:

- (1) R is a left primitive ring, $l.ann_T(I) = 0$ and I has a maximal left T-submodule.
- (2) T is a left primitive ring and I is a left primitive ideal of T.

Moreover, if any of these equivalent conditions holds, then $(R:T) = (R:T)_l = (R:T)_r = 0$.

Proof. First note that since R is a maximal subring of T and $T = R \oplus I$, then we conclude that $T/I \cong R$ as rings. (1) \implies (2) Since R is a left primitive we immediately conclude that I is a left primitive ideal of T. Also note that by (1) of the previous lemma, T is a left primitive ring. Conversely, assume that (2) holds. Clearly R is a left primitive ring and $l.ann_T(I) = 0$ for T is prime. If I has no maximal T-submodules, then we deduce that $I \subseteq J(T) = 0$, by Lemma 2.3, which is absurd. The equalities $(R:T) = (R:T)_l = (R:T)_r = 0$ follows from (2) and (3) of Theorem 2.1.

Let S be a ring and \mathcal{P} be an algebraic property which preserved by ring isomorphisms. Then we define the set $\Omega_{\mathcal{P}}(S)$ be the set of all (proper) ideals of S such that the ring S/I has property \mathcal{P} . For example if \mathcal{P} is the property that a ring is prime, then $\Omega_{\mathcal{P}}(S) = Spec(S)$. Now the following is in order.

Remark 2.6. Let R be a maximal subring of a ring T and I be a nonzero proper ideal of T such that $R \cap I = 0$. Therefore $T = R \oplus I$. If \mathcal{P} is an algebraic property which preserved by ring isomorphisms, then $\{K \oplus I \mid K \in \Omega_{\mathcal{P}}(R)\} \subseteq \Omega_{\mathcal{P}}(T)$. To see this, note that one can easily see that $K \oplus I$ is an ideal of T and $T/(K \oplus I) = (R \oplus I)/(K \oplus I) \cong R/K$ as rings, and therefore since the property \mathcal{P} is preserved by ring isomorphisms, we immediately conclude the inclusion. In particular, the following hold:

- (1) $\{P \oplus I \mid P \in Spec(R)\} \subseteq Spec(T).$
- (2) $\{M \oplus I \mid P \in Max(R)\} \subseteq Max(T).$
- (3) $\{P \oplus I \mid P \in Prm_l(R)\} \subseteq Prm_l(T).$

The similar inclusions hold for right primitive ideals, primitive ideals and strongly prime ideals. Moreover, $\{M \oplus I \mid M \in Max_l(R)\} \subseteq Max_l(T)$ (the same inclusion holds for Max_r instead of Max_l). To see this note that one can easily see that $M \oplus I$ is a proper left ideal of T and $T/(K \oplus I) = (R \oplus I)/(K \oplus I) \cong R/K$ holds as left R-modules. This immediately shows that $K \oplus I$ is a maximal left ideal of T.

Proposition 2.7. Let R be a maximal subring of a ring T and I be a nonzero proper ideal of T such that $R \cap I = 0$. If M is a maximal ideal of T, then either $M = (M \cap R) \oplus I$ or $I^2 = I$ is not contained in M and one of the following holds:

- (1) $(R:T) = (R:T)_l = (R:T)_r = M = ann_T(I)$ is a minimal prime ideal of T.
- (2) $(R:T) = R \cap M$ is a maximal ideal of R and $(R:T) \oplus I$ is a maximal ideal of T.

Proof. If M contains I, then M is not contained in R and therefore by maximality of R we deduce that R + M = T. Therefore $R/(R \cap M) \cong T/M$ as rings. Thus $R \cap M$ is a maximal ideal of R, therefore by the previous remark we deduce that $(R \cap M) \oplus I$ is a maximal ideal of T. Since $(R \cap M) \oplus I \subseteq M$, we infer that the equality holds. Hence assume that M does not contain I. Since I is a minimal ideal of T, we deduce that either $I^2 = 0$ or $I^2 = I$. The case $I^2 = 0$, implies that $I \subseteq M$, which is impossible. Thus $I^2 = I$ and therefore by (3) of Theorem 2.1, we conclude that $(R : T) = (R : T)_l = (R : T)_r$. Again by minimality of I and the fact that M does not contains I we deduce that $M \cap I = 0$ and therefore IM = MI = 0. Hence $M = l.ann_T(I) = r.ann_T(I) = ann_T(I)$. If Q is a prime ideal of T, which properly is contained in M, then $MI = 0 \subseteq Q$, and therefore $I \subseteq Q \subseteq M$, which is absurd. Thus M is a minimal prime ideal of T. Now we have two cases:

- (1) $M \subseteq R$ and therefore M = (R:T).
- (2) M is not contained in R. Therefore by maximality of R we deduce that R + M = T and hence $R/(R \cap M) \cong T/M$ as rings. Thus $R \cap M$ is a maximal ideal of R. Now since $R \oplus I = T$ and MI = 0, we infer that $T(R \cap M) = R \cap M$, and hence $R \cap M \subseteq (R : T)$. Thus $R \cap M = (R : T)$ is a maximal ideal of R. Clearly $(R : T) \oplus I$ is a maximal ideal of T, by the previous remark.

It is well known that if T is a semiprime ring, in particular if T is a reduced ring, then $Soc(_TT) = Soc(T_T)$. In the following result we have a more fact.

Lemma 2.8. Let R be a subring of a reduced ring T. Then the following hold:

- (1) For each $x \in T$, $r.ann_R(x) = l.ann_R(x)$. In particular, Rx is a simple left R-module if and only if xR is a simple right R-module.
- (2) For each $x \in T$ and $n \ge 1$, $l.ann_R(x) = l.ann_R(x^n)$. In particular, Rx is simple if and only if Rx^n is simple.
- (3) $Soc(_RT) = Soc(T_R)$ is an ideal of T.
- (4) If I is a simple left R-submodule of T, then $I^2 = I$ if and only if there exists a central idempotent e of T such that I = Re.

7

Proof. First note that since T is a reduced ring, we conclude that whenever $a, b \in T$, then ab = 0 if and only if ba = 0. For assume that ab = 0, then $(ba)^2 = 0$ and therefore ba = 0. This immediately implies that $A := r.ann_R(x) = l.ann_R(x)$ and therefore A is an ideal of R. Hence Rx is a simple left R-module if and only if xR is a simple right R-module. Hence (1) holds. To see (2), it suffices to show that $l.ann_R(x^n) \subseteq l.ann_R(x)$. We prove by induction on $n \ge 2$. For n = 2, if $rx^2 = 0$, then (rx)x = 0. Thus x(rx) = 0 and hence $(rx)^2 = rxrx = 0$. Therefore rx = 0, for T is a reduced ring. Hence assume that it is true for n-1, we prove it for n. Suppose that $rx^n = 0$. Therefore $(rx)x^{n-1} = 0$. Hence $rx \in l.ann_B(x^{n-1}) = l.ann_B(x)$. Therefore $rx^2 = 0$ and thus rx = 0. Hence (2) holds. For (3) it suffices to show that $Soc(_RT)$ is a right ideal of T. To see this, let S be a simple left R-submodule of T and $t \in T$. Then St (which is a homomorphic image of S) is either 0 or is isomorphic to S as left R-module. Thus $St \subseteq Soc(_RT)$. Hence $Soc(_RT)$ is a right ideal of T. Similarly $Soc(T_R)$ is a left ideal of T. No by (1), $Soc(_{R}T) = Soc(T_{R})$ and therefore (3) holds. Finally, for (4) it is clear that if e is a central idempotent of T such that I = Re, then $I^2 = I$. Conversely, assume that I is a simple left R-submodule of T, such that $I^2 = I$. Assume that $x \in I$ and $x \neq 0$. Then clearly, $0 \neq x^2 \in Ix$ and therefore $0 \neq Ix \subseteq I^2 \subseteq I$. Hence Ix = I, for I is simple. Therefore there exists $a \in I$ such that x = ax. Thus $ax = a^2x$. Now note that $a - a^2 \in I$, for $I^2 = I$ and $a - a^2 \in l.ann_T(x)$. Thus $a - a^2 \in I \cap l.ann_T(x)$. Since I is simple and Ix = I, we immediately deduce that $I \cap l.ann_T(x) = 0$ and therefore $a^2 = a$. Clearly I = Ra and a is central in T (for T is reduced).

Proposition 2.9. Let R be a maximal subring of a ring T. Then either $Soc(_RR) = Soc(_RT)$ (resp. $Soc(_RR) = Soc(_RT)$) or there exists a semisimple left (resp. right) R-submodule A (resp. B) of T such that $T = R \oplus A$ (resp. $T = R \oplus B$), in particular, $(R : T)_r$ (resp. $(R : T)_l$) is a left (resp. right) primitive ideal of R.

Proof. Assume that $Soc(_RR) \neq Soc(_RT)$. Hence there exists a semisimple left R-submodule A of T such that $Soc(_RT) = Soc(_RR) \oplus A$. Since R is a maximal subring of T and $Soc(_RT)$ (by a similar proof of the previous theorem) is a right T-submodule of T (i.e., a right ideal of T) which also a left R-submodule of T and is not contained in R, we deduce that $R + Soc(_RT) = T$. Hence $R + (Soc(_RR) \oplus A) = T$. Also, note that $R \cap A = 0$, for $R \cap A$ is a semisimple left R-submodule which is contained in $Soc(_RR) \cap A = 0$. Therefore $R \oplus A = T$. Finally, note that $R + Soc(_RT) = T$, implies that $T/R \cong Soc(_RT)/Soc(_RR)$ as left and right R-module. Hence by [3, Corollary 2.5], we immediately deduce that $(R : T)_r$ is a left primitive ideal of R, for $Soc(_RT)/Soc(_RR)$ is a nonzero semisimple left R-module.

note that by (3) of Lemma 2.8, if R is a subring of a reduced ring T, then $Soc(_RT) = Soc(T_R)$ is an ideal of T, we denote it by $Soc_R(T)$ (in particular, if R = T, then we denote it by Soc(R)).

Corollary 2.10. Let R be a maximal subring of a reduced ring T. If $Soc(R) \neq Soc_R(T)$. Then there exists $x \in T \setminus R$, such that Rx and xR are simple left and right R-submodules of T, respectively and T = R + RxT = R + TxR. In particular, RxT and TxR are semisimple left and right R-submodules of T, respectively. Moreover, if $I^2 = I \nsubseteq R$ is a simple left (right) R-submodule of T, then $T = R \oplus Re$ and $Soc_R(T) = Soc(R) \oplus Re$.

Proof. Let Rx be a simple left R-submodule of T which is not contained in R. Clearly, $Rx \cap R = 0$ and by (1) of Lemma 2.8, xR is a simple right R-module of T and therefore $xR \cap R = 0$. Hence RxTand TxR are contained in $Soc_R(T)$ but not in R (see the proof of (3) of Lemma 2.8). Hence RxT and TxR are semisimple left and right R-submodules of T, respectively. Thus by the maximality of R, we have R + RxT = T = R + TxR. Now let I be a simple left R-module of T, $I \nsubseteq R$ and $I^2 = I$. By (4) of Lemma 2.8, there exists a central idempotent e of T, such that I = Re (hence I = eR is a simple right R-submodule of T). Clearly $R \cap I = 0$, for I is a simple which is not contained in R. Thus $R + I = R \oplus Re$ is a subring of T (note that $I^2 = I$) which properly contains R. Thus we conclude that $T = R \oplus Re$. Finally note that $Soc_R(T) = (Soc(R) \oplus Re) \oplus A$ for some right R-submodule A of Tand therefore $R \oplus Re = T = R + Soc_R(T) = R \oplus (Re \oplus A)$. Hence if $x \in A$, then x = r + ae for some $r, a \in R$ and therefore $x - ae = r \in R \cap (Re \oplus A) = 0$. Hence $x = ae \in Re$ and therefore A = 0, i.e., $Soc_R(T) = Soc(R) \oplus Re$ (is a projective?) □

Lemma 2.11. Let R be a subring of a ring T. Then $Z(_RT)$ is a proper (R,T)-subbimodule of T which contains $Z(_RR)$. In particular, if R is a maximal subring of T, then either $Z(_RT) = Z(_RR)$ or $R + Z(_RT) = T$ and therefore $R/Z(_RR) \cong T/Z(_RT)$ as left and right R-modules; moreover, in this case $TZ(_RR)$ and $Z(_RR)T$ are contained in $Z(_RT)$. In particular, if T is a prime ring, then either $Soc(_RR) = 0$ or $Z(_RT) = 0$.

Proof. It is clear that $Z := Z({}_{R}T)$ is a left R-submodule of T. Also note that for each $a \in Z$ and $t \in T$, $l.ann_{R}(a) \subseteq l.ann_{R}(at)$ and therefore $at \in Z$. Thus Z is a (R,T)-subbimodule of T. Clearly $1 \notin Z$ and therefore Z is proper. It is obvious that $Z({}_{R}R) \subseteq Z$ and $Z \cap R = Z({}_{R}R)$. Now assume that R is a maximal subring of T and $Z({}_{R}R) \neq Z$. Thus by maximality of R we conclude that R + Z = T. Since in the latter equality all of the parts are left and right R-submodules of T, we obtain that $R/Z({}_{R}R) \cong T/Z$ as left and right R-modules. Now since $Z({}_{R}R)$ is an ideal of R, we deduce that $l.ann_{R}(R/Z({}_{R}R)) =$ $r.ann_{R}(R/Z({}_{R}R)) = Z({}_{R}R)$. Hence $l.ann_{R}(T/Z) = r.ann_{R}(T/Z) = Z({}_{R}R)$. This immediately shows that $TZ({}_{R}R)$ and $Z({}_{R}R)T$ are contained in Z. Since $Soc({}_{R}R)Z = 0$, we infer that $Soc({}_{R}R)TZ({}_{R}R) = 0$. The final part is evident now.

Similarly $Z(T_R)$ is a proper (T, R)-subbimodule of T, which contains $Z(R_R)$. We remind the reader that a ring T is called symmetric if for each natural number n and $a_1, \ldots, a_n \in T$, whenever $a_1 a_2 \cdots a_n = 0$, then for each permutation $\sigma \in S_n$, we have $a_{\sigma(1)}a_{\sigma(2)}\cdots a_{\sigma(n)} = 0$. Note that by [1, Theorem I.3], each reduced ring T is a symmetric ring.

Theorem 2.12. Let R be a maximal subring of a reduced ring T. Then $Z := Z(_RT) = Z(T_R)$ is a proper ideal of T. Moreover, either Z = 0 or the following hold:

- (1) $R \oplus Z = T$. In particular, Z is a minimal ideal of T. In fact for each $0 \neq x \in Z$, we have RxT = Z = TxR.
- (2) $(R:T) = (R:T)_l = (R:T)_r = ann_R(Z)$ which contains Soc(R). Moreover, $Q = ann_T(Z)$ is a minimal prime ideal of T and $Q \cap R = (R:T)$.
- (3) For each $P \in Min_T(Z)$, $T/P \cong R/(R \cap P)$ is a domain.
- (4) T/R is a left and right singular R-module.

Proof. Let $x \in Z(RT)$, then we prove that $x \in Z(T_R)$. To see this we must show that $ann_R(x)$ is essential as right ideal (note that $l.ann_R(x) = r.ann_R(x)$ for T is a symmetric ring). Hence assume that $0 \neq a \in R$. Since $ann_R(x)$ is essential as left ideal in R, we deduce that there exists $r \in R$, such that $0 \neq ra \in ann_R(x)$. Therefore rax = 0 and $ra \neq 0$. Since T is symmetric we conclude that $ar \neq 0$ and xar = 0. This immediately shows that $ann_R(x)$ is essential as right ideal. Hence $x \in Z(T_R)$. Similarly, $Z(T_R) \subseteq Z(RT)$ and hence $Z(T_R) = Z(RT)$ is a proper ideal of T, by the previous lemma. Now assume that $Z \neq 0$. Therefore $R \cap Z = Z(R) = 0$ (for R is reduced) and thus $R \oplus Z = T$, by the maximality of R. Thus Z is a minimal ideal of T by (1) Theorem 2.1. Since for each $0 \neq x \in Z$, RxT and TxR are nonzero (R, T)-subbimodule and (T, R)-subbimodule of Z, respectively. Thus by the maximality of R, we conclude that R + RxT = T = R + TxR. From these equalities we immediately deduce that Z = RxT = TxR. Therefore (1) holds. Now since T is reduced we infer that $Z^2 \neq o$. Thus by (2) and (3) of Theorem 2.1, we conclude that $(R:T) = (R:T)_l = (R:T)_r = ann_R(Z)$. Clearly, Soc(R)Z = ZSoc(R) = 0, hence $Soc(R) \subseteq (R:T)$. By (4) of Theorem 2.1, we deduce that $Q = ann_T(Z)$ is a minimal prime ideal of T which contraction to R equals to (R:T). Therefore (2) is proved. For (3), note that $T/Z \cong R$ as ring, and therefore T/Z is a reduced ring. Thus by [20, Lemma 12.6], we conclude that each minimal prime ideal of T/Z, is a completely prime ideal, i.e., each minimal prime ideal of Z is a completely prime ideal. Also, note that if P is a minimal prime ideal of Z, then $P \nsubseteq R$, for $Z \nsubseteq R$. Thus R + P = T, by the maximality of R. This conclude the final part of (3). Finally for (4), from $R \oplus Z = T$, we deduce that $T/R \cong Z$ as left and right R-modules (note that all of the part of the equality are left and right R-submodules of T). Since Z is left and right singular R-module, we deduce that T/R is left and right singular R-module. \square

In the following example we give an example of a maximal subring R of a reduced ring T with $Z(_RT) \neq 0$.

Example 2.13. Let $T = \mathbb{Z} \times \mathbb{Z}_2$ and $R = \{(n, \bar{n}) \mid n \in \mathbb{Z}\} \cong \mathbb{Z}$ be the prime subring of T. It is clear that R is a maximal subring of T and T is a reduced ring. One can easily see that $ann_R((0, \bar{1})) = 2R$ which is essential in R. Hence $Z_R(T) \neq 0$.

Theorem 2.14. Let R be a prime ring which is a maximal subring of a ring T. Then either for each nonzero ideal I of T, $R \cap I \neq 0$, or there exists a nonzero (prime) ideal I of T such that $R \cap I = 0$ and one of the following holds:

- (1) T is a prime ring. In particular, $(R:T) = (R:T)_l = (R:T)_r = 0$.
- (2) $I^2 = 0$ and $Min(T) = \{I\}.$
- (3) $I^2 = I$, T is semiprime, $Min(T) = \{I, ann_T(I)\}$. Moreover, $(R : T) = (R : T)_l = (R : T)_r = ann_T(I) \cap R$.

Proof. Assume that there exists a nonzero ideal I of T such that $R \cap I = 0$. Clearly, I must a prime ideal of T, for $R \cong T/I$ as rings. If T is prime, then $ann_T(I) = 0$ and therefore by (2) of Theorem 2.1, (1) holds. Hence assume that T is not prime. By (1) of Theorem 2.1, I is a minimal ideal of T and hence either $I^2 = 0$ or $I^2 = I$. If $I^2 = 0$, then clearly $Min(T) = \{I\}$ and therefore (2) holds. Thus assume that $I^2 = I$. By (3) of Theorem 2.1, T is a semiprime ring and $Q = ann_T(I)$ is a minimal prime ideal of T. Clearly $Q \neq 0$, for T is not prime. Now note that since $I^2 = I \neq 0$, we deduce that $I \nsubseteq Q$, therefore by minimality of I we conclude that I and Q are incomparable. Again by minimality of I we conclude that I and Q are incomparable. Again by minimality of I we conclude that I and Q are incomparable. Again by minimality of I we conclude that I and Q are incomparable. Again by minimality of I we conclude that I and Q are incomparable. Again by minimality of I we conclude that I and Q are incomparable. Again by minimality of I we conclude that I and Q are incomparable. Again by minimality of I we conclude that I and Q are incomparable. $Amin(T) = \{I, Q\}$. The final part of (3) is evident by (4) of Theorem 2.1.

Proposition 2.15. Let R be a prime ring which is a maximal subring of a ring T. Let A be the set of nonzero ideal I of T, such that $R \cap I = 0$. Then $|A| \leq 2$. In fact, |A| = 2 if and only if $T \cong \Delta(R) + A \times A$, for some minimal ideal A of R. Moreover, in this case Min(T) = A and $(R : T) = (R : T)_l = (R : T)_r = 0$.

Proof. Since for each $I \in \mathcal{A}$, $T/I \cong R$ as rings, and R is a prime ring, we infer that each element of \mathcal{A} is a prime ideal of T which is a minimal ideal of T, by (1) of Theorem 2.1. Hence, if $I \neq J$ are in \mathcal{A} , then $I \cap J = 0$ and therefore IJ = 0. This immediately shows that $\mathcal{A} = \{I, J\}$ and also $I = ann_T(J)$ and $J = ann_T(I)$. Therefore by the previous theorem we conclude that $Min(T) = \mathcal{A}$ and $(R:T) = (R:T)_l = (R:T)_r = ann_T(I) \cap R = J \cap R = 0$. Finally note that since $I \cap J = 0$, we conclude that there exists a natural ring embedding, say ϕ , such that $T \hookrightarrow T/I \times T/J = R \times R$ and since R is a maximal subring of T, by the comments in Subsection 1.4 of Preliminaries in the introduction of this paper, we conclude that $\phi(T) = \Delta(R) + A \times A$, where A is a minimal ideal of R.

It is clear that in the previous result, if $I \in \mathcal{A}$ and $I^2 = 0$, then $\mathcal{A} = \{I\}$. In other words, I is unique ideal of T respect to the property that $R \cap I = 0$. Also note that if R is note prime then either $|\mathcal{A}| \leq 1$ or $T \cong \Delta(R) + A \times A$, for some minimal ideal A of R.

Proposition 2.16. Let R be a ring and $T = \Delta(R) + A \times A$, where A is a minimal ideal of R. Assume that I be an ideal of T such that $\Delta(R) \cap I = 0$ and $I^2 = I$. If at least one of the following conditions holds, then either $I = A \times 0$ or $I = 0 \times A$ (and therefore in any cases, $A^2 = A$).

- (1) R is a prime ring.
- (2) R is a reduced ring or I is not nil.
- (3) Char(R) = 0 and $(\Delta(R) : T) = 0$ (or only, $2T \nsubseteq \Delta(R)$).
- (4) Char(R) = n > 0 is odd.
- (5) $2 \notin ann(I)$. In particular in this case, I is non-commutative (i.e., there exist $a, b \in I$ such that $ab \neq ba$).

Proof. First note that since A is a minimal ideal of R, then by the comments in Subsection 1.4 of Preliminaries in the introduction of this paper, $\Delta(R)$ is a maximal subring of $T = \Delta(R) + A \times A$. Now note that $\Delta(R) \oplus I = T$, and since $\Delta(R) \oplus (A \times 0) = T$ and $\Delta(R) \oplus (0 \times A) = T$, by (1) of Theorem 2.1, we conclude that I, $A \times 0$, and $0 \times A$ are minimal ideals of T. If $I \cap (A \times 0) \neq 0$, then by minimality we conclude that $I = A \times 0$. Thus we may assume that $I \cap (A \times 0) = 0$ and similarly $I \cap (0 \times A) = 0$. Hence by the previous result R is not a prime ring. If $(x,0) \in I$, then $(x,0) \in T = \Delta(R) \oplus (A \times 0)$. Thus there exist $r \in R$ and $a \in A$ such that (x, 0) = (r, r) + (a, 0). Therefore r = 0 and $x = a \in A$. Hence $(x,0) \in I \cap (A \times 0) = 0$, i.e., x = 0. Similarly if $(0,y) \in I$, then y = 0. Now let $(x,y) \in I$, since $(x,x), (y,y) \in \Delta(R)$ we conclude that $(x^2, xy), (x^2, yx), (xy, y^2), (yx, y^2) \in I$. Clearly $(x^2, y^2) \in I$ and therefore $(0, xy - y^2) \in I$, hence $xy = y^2$ and therefore $(xy, y^2) \in \Delta(R) \cap I = 0$. Thus $x^2 = xy = 0$. Similarly $y^2 = xy = 0$, $x^2 = yx = 0$ and $y^2 = yx = 0$. Hence $x^2 = y^2 = xy = yx = 0$ and $(x, y)^2 = 0$, i.e., I is nil. Therefore if I is not nil or if R is a reduced ring (note that if R is a reduced ring, then $R \times R$ and hence T is a reduced ring) we are done. Since for each $a \in I$, we have $a^2 = 0$, we immediately conclude that for each $a, b \in I$ we have ab = -ba (note, $a + b \in I$ and therefore $(a + b)^2 = 0$). Thus for each $a, b, c \in I$, we have abc = -bac = bca = -abc and therefore 2abc = 0. Hence $2I^3 = 0$. Now since $I = I^2$, we conclude that $I = I^3$ and therefore 2I = 0. Thus if $2 \notin ann(I)$, we have a contradiction and hence we are done. Also note that if I is commutative, then for each $a, b \in I$, we have ba = ab = -baand therefore 2ba = 0, i.e., 2I = 0 which is absurd. Hence 2I = 0 Finally note that since $T = \Delta(R) \oplus I$ and 2I = 0, we conclude that $2T = 2\Delta(R) \subseteq \Delta(R)$. Thus if Char(R) = 0, then $0 \neq 2T \subseteq (\Delta(R) : T)$, which contradiction to (3). If Char(T) = n > 0 is odd, then $2T = T \subseteq \Delta(R)$, which again is impossible. Thus if each of the cases (1) - (5), holds, we deduce that either $I = A \times 0$ or $I = 0 \times A$. \square

In what following we want to study Jacobson radicals, upper nilradicals and lower nilradicals of the ring extension $R \subseteq T$ where R is a maximal subring of T. Specially when $T = R \oplus I$, where I is a nonzero ideal of T. First we need the following fact.

Proposition 2.17. Let R be a maximal subring of a ring T. Then at least one of the following holds:

- (1) (R:T) is the unique maximal ideal of T.
- (2) $J(T_R)$ and $J(_RT)$ are proper submodules of T and hence for each left (resp. right) R-submodule A of T, either $A \subseteq J(_RT)$ (resp. $J(T_R)$) or A has a maximal left (resp. right) R-submodule.

Proof. Assume that M is a maximal ideal of T which is not contained in R, then by maximality of R we obtain that R + M = T. Therefore $R/(R \cap M) \cong T/M$ as left and right R-modules. This immediately implies that T/M has a maximal left (resp.right) R-submodule and thus T has a maximal left (resp. right) R-submodule. Thus J(RT) and $J(T_R)$ are proper submodules of T. Therefore if J(RT) = T (or $J(T_R) = T$), we conclude that each maximal ideal of T is contained in R and therefore is equal to (R:T). Hence (R:T) is the unique maximal ideal of T. For the final part of (2), if A is a left R-submodule of T which is not contained in J(RT), then there exists a maximal left R-submodule N of T such that A is not contained in N. Therefore A + N = T and thus $A/(A \cap N) \cong T/N$ as left R-modules. Hence $A \cap N$ is a maximal submodule of A and we are done.

Theorem 2.18. Let R be a maximal subring of a ring T and $T = R \oplus I$, where I is an ideal of T. Then either $J(T) \cap R = J(R)$ or the following hold:

- (1) $I^2 = I$.
- (2) $J(T) \subseteq (R:T) = (R:T)_l = (R:T)_r$.
- (3) $TJ(R) = J(R)T \notin R$; in particular, (R:T) is not a left/right primitive ideal of R and hence $dim(R) \ge 1$. IJ(R) = J(R)I = I. In particular, _RI, I_R has no maximal submodules.
- (4) $J(_RT)$ and $J(T_R)$ are proper and contains I.
- (5) $(R:T) = l.ann_T(I)$ is a primitive and minimal prime ideal of T.
- (6) (R:T) is not a maximal ideal of T. Moreover, $(R:T) \oplus I$ is a prime ideal of T which is not maximal. In particular, $\dim(T) \ge 2$.
- (7) I is contained in each maximal ideal of T.
- (8) I is unique, i.e., if I_1 is a nonzero proper ideal of T such that $R \cap I_1 = 0$, then $I_1 = I$.

Proof. First we prove that $U(R) = U(T) \cap R$ and $J(T) \cap R \subseteq J(R)$. Assume that $x \in U(T) \cap R$, thus Tx = T = xT and therefore by (6) of Theorem 2.1, we deduce that $Rx = Tx \cap R = T \cap R = R$ and similarly xR = R, which immediately implies that $x \in U(R)$. Thus $U(R) = U(T) \cap R$. Now suppose that $x \in J(T) \cap R$ and $y \in R$. Thus $1 - xy \in U(T) \cap R = U(R)$ and therefore $x \in J(R)$. Now, if for each left primitive ideal Q of T, we have $Q \nsubseteq R$, then by maximality of R, we deduce that R + Q = Tand therefore $R/(R \cap Q) \cong T/Q$ as rings. Hence $R \cap Q$ is a left primitive ideal of R and therefore $J(R) \subseteq R \cap Q$. Hence $J(R) \subseteq J(T) \cap R$, and thus $J(T) \cap R = J(R)$. Thus we may assume that there exist a left primitive ideal Q_1 and (similarly) a right primitive ideal Q_2 of T such that Q_1 and Q_2 are contained in R and therefore $Q_1, Q_2 \subseteq (R:T)$. In particular $J(T) \subseteq R$. Now since Q_i is a prime ideal of T, we deduce that $I^2 \neq 0$, for otherwise $I \subseteq Q_i \subseteq R$ which is absurd. By (1) of Theorem 2.1, I is minimal and therefore we deduce that $I^2 = I$. Thus by (3) of Theorem 2.1, we conclude that $P := (R:T) = (R:T)_l = (R:T)_r$. Hence $J(T) \subseteq Q_i \subseteq P$. Now if $J(R) \subseteq P$ (and therefore J(R)T and TJ(R) are contained in R), then since $T = R \oplus I$, we deduce that $T/R \cong I$ as left and right R-module and therefore J(R)I = 0 = J(R)I. Hence if $x \in J(R)$ and $t = r + i \in T = R \oplus I$, where $r \in R$ and $i \in I$, then $1 - tx = 1 - (r + i)x = 1 - rx \in U(R) \subseteq U(T)$, for $ix \in IJ(R) = 0$. Therefore $x \in J(T)$ and thus J(R) = J(T) and hence we are done. Thus we may assume that $J(R) \not\subseteq P$ (and therefore TJ(R)) and J(R)T are note contained in R; also note that this immediately implies that P is not a one-sided primitive ideal of R and hence is not a maximal ideal of R too, thus $dim(R) \ge 1$, hence J(R)I and IJ(R) are nonzero. Since R is a maximal subring of T we conclude that R + J(R)I = T = R + IJ(R)and since $IJ(R), J(R)I \subseteq I, I \cap R = 0$ and $R \oplus I = T$, we deduce that I = IJ(R) = J(R)I. Thus by (7) of Theorem 2.1, we have TJ(R) = J(R)T. The final part of (3) is an immediate consequence of Nakayama's Lemma. Clearly, since $T/I \cong R$ as left and right R-modules, we conclude that ${}_{R}T$ and T_R have maximal R-submodules and therefore the first part of (4) holds. Now if K is a maximal left *R*-submodule of T which does not contain I, then K + I = T. Therefore $I/(I \cap K) \cong T/K$, which means $I \cap K$ is a maximal left R-submodule of I which is a contradiction by (3). Thus $I \subseteq J(R)$ and similarly $J(T_R)$ contains I. For (5), let $A = l.ann_T(I)$, then by minimality of I, if $A \cap I = I$, then $I^2 = 0$ which is absurd. Thus $A \cap I = 0$ and therefore IA = 0. Hence $A \subseteq r.ann_T(I)$. Similarly, if $B = r.ann_T(I)$,

we have BI = 0 and therefore $B \subseteq A$. Hence A = B, i.e., $l.ann_T(I) = r.ann_T(I)$. Now note that $AI = 0 \subseteq Q_i$ and since Q_i does not contain I, we deduce that $A \subseteq Q_i \subseteq (R:T)$. Hence $A = Q_i = P$ (note that $A = l.ann_R(I) = (R : T)_l = (R : T) = (R : T)_r = r.ann_R(I)$). Hence $A = Q_1 = Q_2 = P$ is a primitive ideal of T which is also a prime ideal of R. If Q is a minimal ideal of T, which properly contained in A, then $AI = 0 \subseteq Q$ implies that $I \subseteq Q \subseteq A$ which is impossible. Thus A is a minimal prime ideal of T. For (6), if P = (R:T) = A is a maximal ideal of T, then I + P = T. If $I \cap P = 0$, then let $\alpha: T \longrightarrow T/I \times T/P$ be the ring isomorphism defined by $\alpha(t) = (t+I, t+P)$. Since $T/I \cong R$ as ring, let $\beta: T/I \times T/P \longrightarrow R \times T/P$ be the ring isomorphism defined by $\beta(t+I, s+P) = (r, s+P)$, where $t = r + i, r \in R$ and $i \in I$. Then Clearly $f = \beta \alpha$ is a ring isomorphism and $f(R) = \{(r, r + P) \mid r \in R\}$ is a maximal subring of $Im(f) = R \times T/P$, for R is a maximal subring of T. But clearly, $R \times R/P$ is a proper subring of T (note R/P is a maximal subring of T/P) which properly contains f(R) (note $(0, 1+P) \in R \times R/P$ but not in f(R). Thus P is not a maximal ideal of T and therefore $dim(T) \ge 1$. The final part of (6) is evident by second part of (3) and Remark 2.6. Now for (7), by (6) note that for each maximal ideal M of T we infer that R does not contain M and therefore by maximality of R, R + M = T. Thus $R/(R \cap M) \cong T/M$ as left (and right) R-modules. In particular, T/M has a maximal left R-submodule. Now if I is not contained in M, then $I \cap M = 0$ by minimality of I and I + M = Tby maximality of M. Therefore, $T/M \cong I$ as left (and right) R-modules. Hence $I \cong R/(R \cap M)$ as left (and right) *R*-modules. Thus *I* has a maximal left (and right) *R*-submodules which is impossible by (3). Hence I is contained in each maximal ideal M of T and therefore (7) is true. Finally, assume that $I_1 \neq I$ be a proper nonzero ideal of T such that $R \cap I_1 = 0$. Therefore $R \oplus I_1 = T$ and hence I_1 is a minimal ideal of T, by (1) of Theorem 2.1. Since $I \neq I_1$ are minimal ideals of T, we conclude that $I \cap I_1 = 0$. Therefore $II_1 = I_1I = 0$. Hence $I_1 \subseteq A \subseteq R$, which is absurd. Thus (7) holds. \square

Remark 2.19. Let S be any ring, then lower nilradical of S, $Nil_*(S)$ is the intersection of all prime ideals or only the minimal prime ideals of S. $Nil^*(S)$ is the unique maximal nil ideal of S which is the sum of all nil ideals of S and is called the upper nilradical of S. A ring S is called strongly prime if S is a prime ring with no nonzero nil ideals. An ideal P of a ring S is called a strongly prime ideal if S/P is a strongly prime ring. It is not hard to see that $Nil^*(S)$ is the intersection of all strongly prime ideals of S, see [25, Proposition 2.6.7].

Theorem 2.20. Let R be a maximal subring of a ring T. Then one of the following condition holds:

(i) There exists a strongly prime ideal Q of T such that $Q \subseteq R$. In particular, $Nil^*(T) \subseteq Q \subseteq (R : T)$.

 $(ii) \ Nil^*(T) \cap R = Nil^*(R). \ Moreover, \ either \ Nil^*(T) = Nil^*(R) \ or \ R/Nil^*(R) \cong T/Nil^*(T).$

Moreover, if there exists a nonzero ideal I of R such that $R \cap I = 0$, then either $Nil^*(T) \cap R = Nil^*(R)$, or the following hold:

- (1) $I^2 = I$.
- (2) $(R:T) = (R:T)_l = (R:T)_r = ann_T(I)$ is a strongly prime ideal of T which is a minimal prime ideal of T.
- (3) $Nil^*(R) \not\subseteq (R:T)$. In particular, IN = NI and TN = NT, where $N = Nil^*(R)$.
- (4) $(R:T) \oplus I$ is a prime ideal of T which is not strongly prime ideal of T. In particular, $(R:T) \oplus I$ is not maximal ideal of T and $\dim(T) \ge 2$.
- (5) I is unique, i.e., if I_1 is a nonzero ideal of T such that $R \cap I_1 = 0$, then $I_1 = I$.

Proof. If there exists a strongly prime ideal Q of T such that $Q \subseteq R$, then (i) holds. Hence assume that for each strongly prime ideal Q of T we have $Q \nsubseteq R$ and therefore by maximality of R, we conclude that R + Q = T. Hence $R/(R \cap Q) \cong T/Q$ as rings, which implies that $R \cap Q$ is a strongly prime ideal of R. Therefore if $\{Q_{\alpha}\}_{\alpha \in \Gamma}$ is the set of all strongly prime ideals of T, we conclude that $Nil^*(R) \subseteq$ $\bigcap_{\alpha \in \Gamma} (R \cap Q_{\alpha}) = Nil^*(T) \cap R$. Now note that if $x \in Nil^*(T) \cap R$, then TxT is a nil ideal of T and since $RxR \subseteq TxT$, we immediately deduce that RxR is a nil ideal of R and hence $Nil^*(T) \cap R \subseteq Nil^*(R)$. Therefore $Nil^*(T) \cap R = Nil^*(R)$. Hence the first part of (ii) holds. Now if $Nil^*(T) \subseteq R$, then $Nil^*(T) = Nil^*(R)$ and if $Nil^*(T) \nsubseteq R$, then by maximality of R we conclude that $R + Nil^*(T) = T$. Therefore $R/Nil^*(R) = R/(R \cap Nil^*(T)) \cong T/Nil^*(T)$. The proofs of the case $T = R \oplus I$, is very similar to the proof of the previous theorem and hence remain to the reader.

Theorem 2.21. Let R be a maximal subring of a ring T. Then at least one of the following holds:

- (1) There exists a (minimal) prime ideal Q of T such that $Q \subseteq (R:T)$. In particular, $Nil_*(T) \subseteq Q \subseteq (R:T)$.
- (2) $Nil_*(R) \subseteq Nil_*(T) \cap R \subseteq Nil^*(R).$

In particular, if there exists a nonzero ideal I of T such that $R \cap I = 0$, then $Nil_*(R) = Nil_*(T) \cap R$.

Proof. It is clear that if R contains a prime ideal of T, then (1) holds. Thus assume that R does not contain any prime ideal Q of T. Hence R + Q = T by maximality of R and therefore $R/(R \cap Q) \cong T/Q$ as ring. Thus $R \cap Q$ is a prime ideal of R. Thus $Nil_*(T) \cap R = \bigcap_{Q \in Spec(T)} (R \cap Q)$ is a semiprime ideal of R and therefore contains $Nil_*(R)$. Since $Nil_*(T)$ is a nil ideal of T, we infer that $Nil_*(T) \cap R$ is nil ideal of R and therefore is contained in $Nil^*(R)$. Now assume that I be a nonzero ideal of T such that $R \cap I = 0$ and therefore $T = R \oplus I$. By (1) of Theorem 2.1, I is minimal and hence we have two cases: (i) $I^2 = 0$ (or if $I \subseteq Nil_*(T)$). In this case first we claim that $Nil_*(T) = Nil_*(R) \oplus I$. To see this first note that since $I^2 = 0$, we deduce that for each prime ideal Q of T, we have $I \subseteq Q$ and therefore $Q \not\subseteq R$. Hence by (2), $Nil_*(R) \subseteq Nil_*(T)$. Therefore $Nil_*(R) \oplus I \subseteq Nil_*(T)$. Now since $T/(Nil_*(R) \oplus I) \cong R/Nil_*(R)$ as rings, we deduce that $Nil_*(R) \oplus I$ is a semiprime ideal of T and therefore contains $Nil_*(T)$. Thus $Nil_*(R) \oplus I = Nil_*(T)$. Finally in this case we must show that $Nil_*(T) \cap R \subseteq Nil_*(R)$. Let $x \in Nil_*(T) \cap R$, thus x = r + i, where $r \in Nil_*(R)$ and $i \in I$, for $Nil_*(T) = Nil_*(R) \oplus I$. Thus $x - r = i \in I \cap R = 0$ and therefore $x = r \in Nil_*(R)$. Thus the equality holds in this case. (ii) Hence assume that $I^2 = I$ and as we mentioned in the previous case we may also assume that $I \not\subseteq Nil_*(T)$, which immediately implies that $I \cap Nil_*(T) = 0$, for I is a minimal ideal of T. Now similar to the previous case if for each prime ideal Q of T, R does not contain Q, then by (2), we deduce that $Nil_*(R) \subseteq Nil_*(T) \cap R$. Also note that as we see in case (i), $Nil_*(R) \oplus I$ is a semiprime ideal of T and hence contains $Nil_*(T)$. Now we prove that $Nil_*(T) = Nil_*(R)$. Assume that $x \in Nil_*(T)$, thus x = r + i, where $r \in Nil_*(R)$ and $i \in I$. Therefore $x - r = i \in Nil_*(T) \cap I = 0$ and therefore $x = r \in Nil_*(T)$. Thus $Nil_*(T) = Nil_*(R)$. Hence assume that there exists a prime ideal Q of T such that $Q \subseteq R$. Thus $Q \subseteq (R:T)$. Since $I^2 = I$, we deduce that $(R:T) = (R:T)_l = (R:T)_r$, by (2) of Theorem 2.1. Therefore (R:T) is a prime ideal of R and hence $Nil_*(R) \subseteq (R:T)$. Since Q is a prime ideal of T, we also deduce that $Nil_*(T) \subseteq Q \subseteq (R:T)$. Since Q is a prime ideal of T which is contained in R and $I \cap R = 0$, we immediately conclude that $I \cap Q = 0$, by minimality of I. Hence QI = IQ = 0, and therefore $Q = l.ann_T(I) = r.ann_T(I) = ann_T(I)$. This immediate implies that Q is a minimal prime ideal of T. Now since $Q = ann_T(I)$ is contained in R, we deduce that Q = (R:T), for $(R:T) = (R:T)_l = r.ann_R(I) = r.ann_T(I) = Q$, and therefore $Nil_*(R) \subseteq Q$. Now note that similar to the previous, $Nil_*(T) \subseteq Nil_*(R) \oplus I$ and $Nil_*(T) \subseteq R$, imply that $Nil_*(T) \subseteq Nil_*(R)$. On the other hand for each minimal prime ideal $Q' \neq Q$ of T, we deduce that $Q' \nsubseteq R$, for Q = (R:T) is a minimal prime ideal of T. Therefore R + Q' = T by maximality of R and thus $R/(R \cap Q') \cong T/Q'$. Hence $R \cap Q'$ is a prime ideal of R. Therefore $Nil_*(R) \subseteq Q'$. Hence $Nil_*(R)$ is contained in each minimal prime ideal of T and therefore is contained in $Nil_*(T)$. Thus $Nil_*(T) = Nil_*(R)$. П

Remark 2.22. Similar to the proof of the previous theorem one can easily see that if R is a maximal subring of a ring T and $T = R \oplus I$, where I is an ideal of T, then if $I \subseteq J(T)$ (resp. $I \subseteq Nil^*(T)$), then $J(T) = J(R) \oplus I$ (resp. $Nil^*(T) = Nil^*(R) \oplus I$).

Proposition 2.23. Let R be a maximal subring of a ring T and I be a nonzero ideal of T such that $R \cap I = 0$. Then either $Soc(_RR) = Soc(_RT) \subseteq (R:T)_r$ or the following hold:

- (1) $Soc(_{R}T) = Soc(_{R}R) \oplus Soc(_{R}I).$
- (2) $Soc(_RI) = I$, *i.e.*, $_RI$ is a semisimple left *R*-module.
- (3) All simple R-submodules of _RI are isomorphic as a left R-module. In fact _RI $\cong \bigoplus_{\alpha \in \Gamma} S$, as left *R*-module, for a simple left R-submodule of I, and a set Γ .
- (4) $(R:T)_r$ is a left primitive ideal of R. In particular, $J(R) \subseteq (R:T)_r$.
- (5) $J(R) = J(T) \cap R$ and $Nil^*(R) = Nil^*(T) \cap R$.
- (6) if in addition, $I^2 = 0$, then I = RaR, for each $0 \neq a \in I$.

Proof. Since $T = R \oplus I$, we conclude that $Soc(_RT) = Soc(_RR) \oplus Soc(_RI)$. Hence if $Soc(_RI) = 0$, then $Soc(_RR) = Soc(_RT)$. Now note that by the proof of (3) in Lemma 2.8, $Soc(_RT)$ is a right ideal of T, which is contained in R, therefore $Soc(_RR) = Soc(_RT) \subseteq (R:T)_r$. Hence assume that $Soc(_RI) \neq 0$. Let S be a simple left R-submodule of I. Therefore ST is a (R,T)-subbimodule of I, which is not contained in R. Thus $R \oplus ST = T$, by maximality of R. Since $ST \subseteq I$ and $R \oplus I = T$, we immediately conclude that I = ST. Hence $I = ST = \sum_{t \in T} St$ and for each $t \in T$, note that either St = 0 or $St \cong S$ as a left R-module. Hence I = ST is a semisimple left R-module, all of simple left R-submodule of I are isomorphic to S, i.e., $Soc(_RI) = I$ and there exists $T' \subseteq T$ such that $I \cong \bigoplus_{t \in T'} S$, as left R-modules. Hence $T/R \cong I \cong \bigoplus_{t \in T'} S$ and therefore $(R:T)_r = l.ann_R(S)$ is a left primitive ideal of R. (5) is an immediate consequences of (3) of Theorem 2.18, (3) of Theorem 2.20, and the facts that $_RI$ has a

maximal (left) *R*-submodule in this case and also J(R)I = 0. Finally for (6), if $I^2 = 0$, then for each nonzero $a \in I$, one can easily check that $R \oplus RaR$ is a subring of *T* which properly contains *R* and therefore $R \oplus RaR = T$, by maximality of *R*. Now since $RaR \subseteq I$ and $R \oplus RaR = T = R \oplus I$, we conclude that I = RaR.

Proposition 2.24. Let R be a maximal subring of a ring T and I be a nonzero ideal of T such that $R \cap I = 0$. Then either $Z(_RT) = Z(_RR) \subseteq (R:T)_r$ or the following hold:

- (1) $Z(_{R}T) = Z(_{R}R) \oplus Z(_{R}I).$
- (2) $Z(_RI) = I$, i.e., $_RI$ is a singular left R-submodule of T (i.e., $_R(T/R)$ is a singular module).
- (3) $Soc(_RR) \subseteq (R:T)_r$.

Proof. First note that $T = R \oplus I$ implies that $Z(RT) = Z(RR) \oplus Z(RI)$. Hence if Z(RI) = 0, then Z(RR) = Z(RT) and since Z(RT) is a right ideal of T, by Lemma 2.11, we deduce that $Z(RT) = Z(RR) \subseteq (R : T)_r$. Thus assume that $Z(RI) \neq 0$. Since I is an ideal of T we deduce that Z(RI) is a right ideal of T which is contained in I, and clearly is a left nonzero R-submodule of T, therefore $R \oplus Z(RI) = T$. Hence by $R \oplus Z(RI) = T = R \oplus I$ and $Z(RI) \subseteq I$ we deduce that Z(RI) = I, i.e., I is a singular left R-submodule of T. Finally note that Soc(RR)Z(RI) = 0 implies that Soc(RR)I = 0 and since $T/R \cong I$ as left R-modules, we conclude that $Soc(RR)T \subseteq R$. Therefore $Soc(RR) \subseteq (R:T)_r$. \Box

Proposition 2.25. Let R be a maximal subring of a ring T and I be a nonzero ideal of T with $R \cap I = 0$. Then the following hold:

- (1) If either R or T is a one-sided Artinian ring, then $J(R) = J(T) \cap R$ and $Nil^*(R) = Nil^*(T) \cap R$.
- (2) If R is left Artinian ring, then $Soc(_RI) = I$. In particular, $_RT$ is Artinian if and only if $_RI$ is finitely generated.

Proof. (1) First assume that T is a one-sided Artinian ring, dim(T) = 0. Hence by (6) of Theorem 2.18, we deduce that $J(T) \cap R = J(R)$; and by (4) of Theorem 2.20, we conclude that $Nil^*(T) \cap R = Nil^*(R)$. Now assume that R is a one-sided Artinian ring, thus dim(R) = 0. Hence by (3) of Theorem 2.18, we have $J(T) \cap R = J(R)$. Finally note that by Theorem 1.1, $(R:T)_l$ is a prime ideal of R, thus by (2) and (3) of Theorem 2.20, we conclude that $Nil^*(T) \cap R = Nil^*(R)$. For (2), Since R is a left Artinian ring, we deduce that $Soc(_RI) \neq 0$. Therefore by (2) of Proposition 2.23, we conclude that $Soc(_RI) = I$. It is clear that $_RT$ is Artinian if and only if $_RI$ is finitely generated and if any of these equivalent conditions holds then T is a left Artinian ring.

If S is a one-sided Noetherian ring, it is well known that the set of minimal prime ideals of S is finite and $Nil^*(S) = Nil_*(S)$ is a nilpotent ideal of S and every one-sided nil ideal of S is nilpotent, see [20] and [25]. Moreover, if R is a subring of S, then ACC holds on the set of one-sided ann of R, see [21], and thus $Nil^*(R) = Nil_*(R)$, see [20].

Corollary 2.26. Let T be a left Noetherian ring and R be a maximal subring of T. Then either $Nil^*(T) \subseteq Nil_*(R)$ or $Nil^*(T) \cap R = Nil^*(R) = Nil_*(R)$ is a nilpotent ideal.

Proof. Since T is a left Noetherian ring, we infer that $Nil^*(T)$, and therefore $Nil^*(T) \cap R$, are nilpotent. Thus $Nil^*(T) \cap R \subseteq Nil_*(R)$. Hence, if $Nil^*(T) \subseteq R$, then $Nil^*(T) \subseteq Nil_*(R)$. Thus assume that $Nil^*(T) \notin R$. Therefore $R+Nil^*(T) = T$, by maximality of R and thus $R/(R \cap Nil^*(T)) \cong T/Nil^*(T)$ as rings. Hence $R/(R \cap Nil^*(T))$ has no nil ideal and therefore $Nil^*(R) \subseteq Nil^*(T) \cap R$. Since $Nil^*(T) \cap R \subseteq Nil_*(R)$, we conclude that $Nil^*(T) \cap R = Nil^*(R) = Nil_*(R)$ is nilpotent. \Box

Remark 2.27. If T is a left Artinian ring, then each prime ideal of T is maximal and thus minimal prime ideals, prime ideals, left/right primitive ideals and maximal ideals of T are coincided, and T has only finitely many maximal ideals, say P_1, \ldots, P_n . Hence $J(T) = Nil^*(T) = Nil_*(T) = \bigcap_{i=1}^n P_i$ is a nilpotent ideal of T, see [25].

Theorem 2.28. Let T be a left Artinian ring and R be a maximal subring of T. Then exactly one of the following holds:

- (1) There exists a (unique) maximal ideal M of T such that $M \subseteq R$. In particular, R/M is a maximal subring of Artinian simple ring T/M (which is isomorphic to $\mathbb{M}_n(D)$, for some natural number n and a division ring D). Moreover, (R : T) = M and each prime ideal of R either contains (R : T) = M or is a maximal ideal of R.
- (2) R is a zero-dimensional ring, Spec(R) is finite $(|Spec(R)| \le |Max(T)|)$ and $J(R) = Nil^*(R) = Nil_*(R) = J(T) \cap R$ is a nilpotent ideal of R and one of the following holds:

- (a) There exist maximal ideals $M \neq N$ of T such that $T/M \cong T/N$ as rings.
- (b) $R/J(R) \cong T/J(T)$ as rings. In particular, R is a semiprimary ring.

Proof. First assume that there exists a maximal ideal M of T such that $M \subseteq R$. It is clear that M must be unique for R is a proper subring of T and also (R:T) = M. Now let P be a prime ideal of R, then $R \setminus P$ is a *m*-system in T and therefore there exists a prime (maximal) ideal N of T such that $(R \setminus P) \cap N = 0$. Thus $R \cap N \subseteq P$. Now we have two cases: (i) $N \subseteq R$ and therefore $N = M = (R : T) \subseteq P$. (ii) $N \not\subseteq R$, therefore R + N = T, by the maximality of R. Hence $R/(R \cap N) \cong T/N$, as rings. Thus $R \cap N$ is a maximal ideal of R, for T/N is a simple ring. Therefore $P = R \cap N$ is a maximal ideal of R. Now assume that R does not contain any maximal ideal of T and let $Max(T) = \{M_1, \ldots, M_n\}$. Similar to the latter proof note that for each i, $R \cap M_i$ is maximal ideal of R and each prime ideal of R is of the form $R \cap M_i$ for some *i*. Therefore R is a zero-dimensional ring and clearly $|Spec(R)| \leq |Max(T)|$ in any cases. Thus $J(R) = Nil^*(R) = Nil_*(R) \subseteq J(T) \cap R$. Now note that $J(T) \cap R$ is a nilpotent ideal of R and therefore $J(T) \cap R \subseteq Nil_*(R)$. Hence $J(R) = Nil^*(R) = Nil_*(R) = J(T) \cap R$ is nilpotent. Now we have two cases: (a) $J(T) \subseteq R$, thus R/J(T) is a maximal subring of $T/J(T) = T/M_1 \times \cdots \times T/M_n$. Therefore by Theorem 1.1 and the fact that R does not contain any maximal ideals of T, we deduce that there exist distinct maximal ideals M_i and M_j of $T, i \neq j$ such that $T/M_i \cong T/M_j$ as rings, and $R/J(T) \cong \prod_{i \neq k=1}^{n} T/M_k$. (b) $J(T) \not\subseteq R$, therefore R + J(T) = T, by the maximality of R. Hence $R/J(R) = R/(J(T) \cap R) \cong T/J(T)$, as rings, and thus R/J(R) is an Artinian ring and therefore R is a semiprimary ring. \square

Remark 2.29. Let R be a left Artinian subring of a ring T, then one can easily see that $U(R) = U(T) \cap R$ and therefore $J(T) \cap R \subseteq J(R)$.

Lemma 2.30. Let R be a maximal subring of a ring T and R be a left Artinian ring. Then the following hold:

- (1) For each prime ideal P of T, either $P \subseteq R$ (i.e., $P \subseteq (R:T)$) or P is a maximal ideal of T.
- (2) Either $J(T) \subseteq J(R)$ or $J(R) = J(T) \cap R$.
- (3) If each left (resp. right) primitive ideal of T is not contained in R, then $J(R) = J(T) \cap R$ and each left (resp. right) primitive ideal of T is maximal.
- (4) Either $Nil^*(T) \subseteq J(R)$ or $Nil^*(T) \cap R = J(R) = J(T) \cap R$.
- (5) If each strongly prime ideal of T is not contained in R, then $Nil^*(T) \cap R = J(R) = J(T) \cap R$ and each strongly prime ideal of T is maximal.
- (6) Either $Nil_*(T) \subseteq J(R)$ or $Nil_*(T) \cap R = J(R) = Nil^*(T) \cap R = J(T) \cap R$.
- (7) If each prime ideal of T is not contained in R, then $Nil_*(T) \cap R = J(R) = Nil^*(T) \cap R = J(T) \cap R$ and dim(T) = 0.
- (8) DCC holds on Spec(T).
- (9) Each nonzero left R-submodule of T contains a simple left R-submodule. In fact, $Soc(_{R}T)$ is an essential left R-submodule of T. Moreover, if there exists a nonzero ideal I of T such that $I \cap R = 0$, then (1) - (6) of Proposition 2.23 hold.

Proof. (1) Let P be a prime ideal of T which is not contained in R, then by maximality of R we conclude that R+P=T and therefore $R/(R\cap P)\cong T/P$, as rings. Hence $P\cap R$ is a prime ideal of R and since R is a left Artinian ring we conclude that $P \cap R$ is a maximal ideal of R and therefore P is a maximal ideal of T. (2) If $J(T) \subseteq R$, then it is clear that $J(T) \subseteq J(R)$, by the previous remark. Hence assume that $J(T) \nsubseteq R$ and therefore by maximality of R we conclude that R + J(T) = T. Hence $R/(J(T) \cap R) \cong T/J(T)$, as rings, and therefore $R/(J(T) \cap R)$ is a J-semisimple ring. This means $J(T) \cap R$ is an intersection of a family of maximal left ideals of R and therefore $J(R) \subseteq J(T) \cap R$. Thus $J(R) = J(T) \cap R$, by the previous remark. (3) Assume that for each left primitive ideal Q of T, Q is not contained in R. Therefore by (1), we deduce that Q is a maximal ideal of T and $Q \cap R$ is a maximal ideal of R. Hence $J(R) \subseteq Q \cap R$, for each left primitive ideal Q of T and therefore $J(R) \subseteq J(T) \cap R$. Hence we are done by the previous remark. The proofs of (4) - (7) are similar to (2) and (3). (8) is an immediate consequence of (1) and the fact that R is a left Artinian ring. (9) Assume that K be a nonzero left R-submodule of T and $0 \neq x \in K$. Since R is a left Artinian ring, we infer that Rx is an Artinian submodule of K. It is clear that Rx, and therefore K, contains a simple (left) R-submodule. Thus $Soc(_{R}T)$ is an essential left R-submodule of T. The final part is evident, for $Soc(_RI) \neq 0$, and use Proposition 2.23. \Box

Lemma 2.31. Let R be a maximal subring of a ring R and M, N are distinct maximal ideals of T which are not contained in R. If $P := M \cap R = N \cap R$, then the following hold:

14

- (1) $P = (R:T) = M \cap N$ is a maximal ideal of R. Moreover, P is not a prime ideal of T.
- (2) $T/M \cong R/P \cong T/N$ as rings and left/right R-modules.
- (3) $|Max(T)| \le 1 + |Max(R)|$
- (4) If R is a left Noetherian (resp. Artinian) ring, then T is a left Noetherian (resp. Artinian) R-module; in particular T is a left Noetherian (resp. Artinian) ring.

Proof. (1) and (2), since M is not contained in R, we deduce that R + M = T and therefore $R/(M \cap R) \cong$ T/M as rings and also left/right R-modules. Thus $P = M \cap R = N \cap R$ is a maximal ideal of R. Clearly, $P = (M \cap N) \cap R$. If $M \cap N$ is not contained in R, then by maximality of R we conclude that $R + (M \cap N) = T$ and therefore $R/P \cong T/(M \cap N)$, as rings, which is impossible, for R/P is a simple ring but $T/(M \cap N)$ is not. Hence $M \cap N \subseteq R$ and therefore $P = M \cap N \subseteq (R:T)$ is a maximal ideal of R. Thus $P = M \cap N = (R:T)$. For the final part of (1), if (R:T) is a prime ideal of T, then $MN \subseteq (R:T)$, implies that $M \subseteq (R:T) \subseteq R$ or $N \subseteq (R:T) \subseteq R$, which are impossible. Hence (R:T)is not a prime ideal of T. (3) Let Q be a prime ideal of T such that $Q \cap R = P$. Hence $M \cap N = Q$ and therefore $M \subseteq Q$ or $N \subseteq Q$, for Q is prime. Thus Q = M or Q = N. Therefore for each maximal ideal Q of T, if $Q \neq M, N$, then we conclude that $Q \cap R \neq P$. Now if Q, Q' are maximal ideal of T which are not equal to M or N, then $Q \cap R \neq Q' \cap R$, for otherwise by (1) $M \cap N = Q \cap Q' \subseteq Q$, i.e., Q = M or Q = N, which is absurd. Also note that by (1), Q is not contained in R, and therefore similar to the proof of (1), we deduce that $Q \cap R$ is a maximal ideal of R. Hence M and N have a same contraction to R and other maximal ideals of T have different contractions (which also are not equal to P), therefore $|Max(T)| \leq 1 + |Max(R)|$. Finally for (4), note that $T/(M \cap N)$ embeds in $T/M \times T/N$ as left T-modules and therefore as left R-modules too. Hence by (2), $T/(M \cap N)$ embeds in $R/P \times R/P$ as left R-modules. Thus, if R is a left Noetherian (resp. Artinian) ring, then $T/(M \cap N)$ is a left Noetherian (resp. Artinian) R-module and since $M \cap N \subseteq R$, we immediately conclude that T is a left Noetherian (resp. Artinian) *R*-module. The final part is evident.

Remark 2.32. Let R be a maximal subring of a ring T and R be a left Artinian ring. Then one of the following holds:

- (1) $(R:T)_l$ is not an ideal of T. In this case $(R:T)_r$ is not an ideal of T and $(R:T)_l \neq (R:T)_r$. In particular, $R = \mathbb{I}((R:T)_l) = \mathbb{I}((R:T)_r)$.
- (2) $(R:T) = (R:T)_l = (R:T)_r$.

To see this, first note that $(R:T) \subseteq (R:T)_l, (R:T)_r$ and $(R:T)_l, (R:T)_r$ are prime ideals of R and therefore are maximal, for R is a left Artinian ring. Also note that $(R:T)_l$ is an ideal of T if and only if $(R:T)_l = (R:T)$. Since $(R:T) \subseteq (R:T)_r$ and $(R:T)_l$ is a maximal ideal of R, we infer that $(R:T)_l$ is an ideal of T if and only if $(R:T)_l = (R:T)_r$, i.e., $(R:T)_l = (R:T) = (R:T)_r$. Similarly, $(R:T)_r$ is an ideal of T if and only if $(R:T)_l = (R:T) = (R:T)_r$. Thus the first part of (1) and (2) are clear now. Finally, assume that $(R:T)_l$ and therefore $(R:T)_r$ are not ideals of T. Since these are ideals of Rand R is a maximal subring of T, we immediately conclude that $R = \mathbb{I}((R:T)_l) = \mathbb{I}((R:T)_r)$.

Also note that in Case (1) of the previous remark, either $(R:T)_l T = T$ or $(R:T)_l T$ is a proper ideal of T. In the following, we study these cases separately, to find some relations between Jacobson radicals of the ring extension $R \subseteq T$, where R is a left Artinian ring and is a maximal subring of T.

Theorem 2.33. Let R be a maximal subring of a ring T and R be a left Artinian ring. Assume that $(R:T)_l T = T$, then $_R T$ and $((R:T)_l)_R$ are finitely generated. In particular, T is a left Artinian ring. Moreover, either $J(R) = J(T) \cap R$ or the following hold:

- (1) $Q := (R:T) \in Max(T)$. In particular, $J(T) \subseteq J(R) \cap (R:T)$.
- (2) $T(R:T)_r = T$. In particular, T_R is finitely generated.
- (3) If M and N are distinct maximal ideals of T and $M, N \neq Q$, then $M \cap R \neq N \cap R$. In particular, $Max(R) = \{(R:T)_l, (R:T)_r\} \cup \{P \cap R \mid P \in Max(T), P \neq Q\}$ and |Max(R)| = |Max(T)| + 1.

Proof. Since $(R : T)_l T = T$, there exist $a_1, \ldots, a_n \in (R : T)_l$ and $t_1, \ldots, t_n \in T$ such that $a_1t_1 + \cdots + a_nt_n = 1$. Therefore for each $x \in T$ and $p \in (R : T)_l$, we have $x = xa_1t_1 + \cdots + xa_nt_n$ and $a_1t_1p + \cdots + a_nt_np = p$. Since $p, a_i \in (R : T)_l$, we infer that $xa_i, t_ip \in R$. Therefore $T = Rt_1 + \cdots + Rt_n$ and $(R : T)_l = a_1R + \cdots + a_nR$. Hence $_RT$ and $((R : T)_l)_R$ are finitely generated. It is clear that T is a left Artinian R-module and therefore is a left Artinian ring. Now by (3) of Lemma 2.30, we have two cases: (i) if R does not contain any maximal ideal of T (note R is a left Artinian ring), then $J(R) = J(T) \cap R$. (ii) hence assume that R contains a maximal ideal Q of T, which clearly is unique. Therefore Q = (R : T) and thus $J(T) \subseteq (R : T)$ is contained in R. Hence $J(T) \subseteq J(R)$, by Remark 2.29. Thus (1) holds. For

(2), since $(R:T)_l$ is not an ideal of T, we deduce that $(R:T)_r$ is not an ideal of T, by Remark 2.32. Hence $Q \subsetneq (R:T)_r \subsetneq T(R:T)_r$ and therefore by maximality of Q we conclude that $T(R:T)_r = T$. Similar to the proof of $_{R}T$ is finitely generated, we deduce that T_{R} is finitely generated. Finally for (3), assume that M and N are distinct maximal ideals of T and $M, N \neq Q$. If $M \cap R = N \cap R$, then by the first part of (1) of Lemma 2.31, we deduce that $Q = (R:T) = M \cap N$, which is absurd, by the second part of (1) of Lemma 2.31. Thus $M \cap R \neq N \cap R$. Also note that, since $M \neq Q$, we have $M \not\subseteq R$ and therefore R+M=T, for R is a maximal subring of T. Hence $R/(R\cap M)\cong T/M$ as rings and therefore $R\cap M$ is a maximal ideal of R. Also note that since $(R:T)_l T = T$ and $T(R:T)_r = T$, we immediately conclude that no maximal ideals of T contains $(R:T)_l$ or $(R:T)_r$. Hence, if $M_1 = Q, M_2, \ldots, M_k$ denote all maximal ideals of T (note, by (3) of Lemma 2.31, Max(T) is finite), then $(R:T)_l, (R:T)_r, M_2 \cap R, \ldots, M_k \cap R$ are maximal ideals of R. Conversely, if P is a maximal ideal of R, then there exists a maximal ideal Q'of T such that $Q' \cap R \subseteq R$ (note $R \setminus P$ is a m-system in R and therefore in T and T is a left Artinian ring). Now we have two cases: (a) Q' = Q and therefore $Q \subseteq P$. Since $(R:T)_l(R:T)_r \subseteq Q \subseteq P$, we deduce that $P = (R : T)_l$ or $P = (R : T)_r$, for P is a prime ideal and R is a left Artinian ring. (b) If $Q' \neq Q$, then as we see earlier $Q' \cap R$ is a maximal ideal of R and therefore $P = Q' \cap R$. Hence $Max(R) = \{(R:T)_l, (R:T)_r\} \cup \{M_2 \cap R, \dots, M_k \cap R\}$ and therefore |Max(R)| = |Max(T)| + 1. \square

Theorem 2.34. Let R be a maximal subring of a ring T and R be a left Artinian ring. Assume that $(R:T)_l$ is not an ideal of T and $(R:T)_l$ T is a proper ideal in T. Then the following hold:

- (1) $(R:T)_l T \in Max(T).$
- (2) R contains no maximal ideals of T; the contraction of distinct maximal ideals of T are distinct maximal ideals of R, in particular $|Max(T)| \leq |Max(R)|$.
- (3) If $I := l.ann_T((R:T)_l)$, then either $(R:T)_lT$ is a minimal prime ideal of T and $J(R) = J(T) \cap R$ or $I^2 = 0$, $I \subseteq Nil_*(T)$ and $(R:T)_lT = r.ann_T(I)$ (whenever $I \neq 0$).
- (4) Either $J(R) = J(T) \cap R$ or $(R:T)_l$ is not a finitely generated right ideal of R.
- (5) Either $T(R:T)_r \in Max(T)$ or there exists a natural number n such that T embeds in $((R:T)_r)^n$ as a right T-module and therefore in \mathbb{R}^n as a right R-module.

Proof. First note that by the assumptions $(R:T) \subsetneq (R:T)_l \subsetneq (R:T)_l T \subsetneq T$. Hence $R + (R:T)_l T = T$, by the maximality of R. Clearly $(R:T)_l T \cap R = (R:T)_l$ which is a maximal ideal of R. Hence $R/(R:T)_l = R/((R:T)_l T \cap R) \cong T/(R:T)_l T$, as rings and thus $(R:T)_l T$ is a maximal ideal of T. Thus (1) holds. For (2), if K is an ideal of T which is contained in R, then $K \subseteq (R:T) \subsetneq (R:T)_l T$ and therefore K is not a maximal ideal of T. Hence if M is a maximal ideal of T, then R + M = T by maximality of R, and therefore $R/(R \cap M) \cong T/M$ as rings, thus $R \cap M$ is a maximal ideal of R. Now assume that M and N are maximal ideals of T with $R \cap N = R \cap M$. Thus by (1) of Lemma 2.31, we deduce that $M \cap N = (R:T) \subseteq (R:T)_l T$. Hence by (1), we conclude that $M = (R:T)_l T$ or $N = (R:T)_l T$. In any cases, since $M \cap R = N \cap R$, we deduce that $(R:T)_l \subseteq M, N$ and thus $M = N = (R:T)_l T$. Therefore (2) holds. For (3), first suppose that $(R:T)_l T$ is a minimal prime ideal of T, then by (3) of Lemma 2.30, if Q is a left primitive ideal of T and $Q \subseteq R$, then $Q \subseteq (R:T) \subsetneq (R:T)_l \subsetneq (R:T)_l T$, which is absurd. Thus R does not contain any left primitive ideal of T and therefore $J(R) = J(T) \cap R$ by (3) of Lemma 2.30. Now assume that $(R:T)_l T$ is not a minimal prime ideal of T. For our claim we may suppose that $I \neq 0$. Since $(R:T)_{l}T$ is not a minimal prime ideal of R, we conclude that there exists a prime ideal Q of T which is properly contained in $(R:T)_l T$. By $I(R:T)_l = 0 \subseteq Q$, we deduce that $I \subseteq Q \subsetneq (R:T)_l T = P$. In particular, $I^2 = 0$ and therefore $I \subseteq Nil_*(T)$. Now note that since $I = l.ann_T((R:T)_lT)$ and $(R:T)_lT$ is a maximal ideal of T, we obtain that $(R:T)_{l}T = r.ann_{T}(I)$. Hence (3) holds. For (4), assume that $J(R) \neq J(T) \cap R$, then by (3) of Lemma 2.30, let Q be a left primitive ideal of T which is contained in R. Hence $Q \subseteq (R:T) \subsetneq (R:T)_l \subsetneq (R:T)_l T$. This immediately implies that $l.ann_T((R:T)_l) \subseteq Q$ and therefore $l.ann_T((R:T)_l)$ is a left Artinian R-module. Now suppose that $(R:T)_l$ is a finitely generated as right ideal of R. Hence $(R:T)_l = a_1R + \cdots + a_nR$, for some $a_1, \ldots, a_n \in (R:T)_l$. Therefore $l.ann_T((R:T)_l) = l.ann_T(a_1) \cap \cdots \cap l.ann_T(a_n)$, which immediately implies that $T/l.ann_T((R:T)_l)$ can be embedded in $T/l.ann_T(a_1) \times \cdots \times T/l.ann_T(a_n)$, as left T-modules. Since $T/l.ann_T(a_i) \cong Ta_i$ as left T-modules and $a_i \in (R:T)_l$, we conclude that $T/l.ann_T((R:T)_l)$ embeds in $((R:T)_l)^n$ as left T-module and therefore as left R-module too. This immediately implies that $T/l.ann_T((R:T)_l)$ is a left Artinian R-module and therefore T is a left Artinian ring. Hence Q is a maximal ideal of T, which is absurd. Thus $(R:T)_l$ is not a finitely generated right ideal of T. For (5), note that either $T(R:T)_r$ is a proper ideal of T or $T(R:T)_r = T$. First assume that $T(R:T)_r$ is a proper ideal of T. Then clearly, $T(R:T)_r \cap R = (R:T)_r$, for $(R:T)_r$ is a maximal ideal of R. By Remark 2.32, we deduce that $T(R:T)_r$ is not contained in R, for $(R:T)_l$ and $(R:T)_r$ are not ideals of T. Hence $R + T(R:T)_r = T$, by maximality of R and therefore $R/(R:T)_r \cong T/T(R:T)_r$ as rings, which immediately implies that $T(R:T)_r$ is a maximal ideal of T. Now assume that $T(R:T)_r = T$, in this case similar to the proof of (4) and the proof of Theorem 2.33, one can complete the proof.

By (4) of the previous theorem we have the following immediate corollary.

Corollary 2.35. Let R be a maximal subring of a ring T and R be an Artinian ring. Assume that $(R:T)_l$ is not an ideal of T and $(R:T)_lT$ is a proper ideal in T. Then $J(R) = J(T) \cap R$

Proposition 2.36. Let R be a maximal subring of a ring T. Assume that $(R:T)_l = (R:T) = (R:T)_r$, R is a left Artinian ring and T is a zero-dimensional ring. Then $J(R) = J(T) \cap R$.

Proof. First note by Remark 2.29, $J(T) \cap R \subseteq J(R)$. Now by (3) of Lemma 2.30, we have two cases: (*i*) If each left primitive ideal of T is not contained in R, then $J(R) = J(T) \cap R$ and hence we are done. (*ii*) There exists a left primitive ideal Q of T such that $Q \subseteq R$. Hence by our assumption Q is a maximal ideal of T and therefore Q = (R : T). Therefore in this case we infer that $J(T) \subseteq J(R) \subseteq Q$. Now let P be a left primitive ideal of T and $P \neq Q$. Therefore $P \nsubseteq R$ and thus R + P = T, by maximality of R. Thus $R/(R \cap P) \cong T/P$ as rings, and therefore $P \cap R$ is a maximal ideal of R (note that T is zero-dimensional and hence P is maximal). Thus $J(R) \subseteq P \cap R$. Hence for each left primitive ideal P of R we have $J(R) \subseteq P$ and therefore $J(R) \subseteq J(T)$. Thus J(R) = J(T).

Theorem 2.37. Let R be a maximal subring of a ring T and R be a left Artinian ring. Assume that $(R:T)_l = (R:T) = (R:T)_r$. Then $T/(R:T) \cong \mathbb{M}_n(S)$, for some ring S and a natural number n, and S has a maximal subring which is a division ring. Moreover, either T is a left Artinian ring (in particular, $J(R) = J(T) \cap R$) or S (T/(R:T)) has at most one nonzero proper ideal, and one of the following holds:

- (1) S is a simple ring, i.e., (R:T) is a maximal ideal of T.
- (2) S has a unique nonzero proper ideal M such that $M^2 = 0$. In particular, J(S) = M.
- (3) S has a unique nonzero proper ideal M such that $M^2 = M$. In particular, either S is a primitive ring or J(S) = M.

Moreover, if Q is the unique maximal ideal of T which contains (R : T) in items (1) - (3), then the following hold:

- (a) $Max(T) = \{Q\} \cup \{N \in Spec(T) \mid N \nsubseteq R\}.$
- (b) Each prime ideal of T is either maximal or is contained in Q. In particular, dim(T) = ht(Q).
- (c) If N_1 and N_2 are maximal ideals of T and $N_i \neq Q$, then $N_1 \cap R \neq N_2 \cap R$.
- (d) $|Max(T)| \le |Max(R)|$.

Proof. Let P := (R:T), then by our assumption we deduce that P is a maximal ideal of R and therefore R/P is of the form $\mathbb{M}_n(D)$ for some natural number n and a division ring D. Now since R/P is a maximal subring of T/P, we immediately conclude that T/P is of the form $\mathbb{M}_n(S)$ for a ring S where D is a maximal subring of S. Since R/P is a simple ring, by Theorem 1.2 we have two cases: (i) T/Phas exactly two nonzero proper (maximal) ideals M/P and N/P such that $(M/P) \cap (N/P) = 0$, i.e., $M \cap N = P$ and therefore by (3) of Lemma 2.31, we deduce that T is a left Artinian ring (and hence $J(R) = J(T) \cap R$, by the previous proposition). (ii) one of the items (1), (3), (4) of Theorem 1.2 holds. Now one can easily see that in case (1) of Theorem 1.2, we obtain (1), in case (4) of Theorem 1.2, we deduce (2), and in case (3) of Theorem 1.2, we have (3). Now assume that Q is the unique maximal ideal of T which contains (R:T), i.e., in case (1), Q = (R:T) and otherwise $Q = M_n(M)$ in cases (2) and (3). (a) if P is a prime ideal of T which is not contained in R, then by maximality of R we conclude that R + P = T and therefore $R/(R \cap P) \cong T/P$ as rings. Thus $R \cap P$ is a prime ideal of R, and hence is a maximal ideal of R, for R is a left Artinian ring. Hence P is a maximal ideal of T. Conversely, if N is a maximal ideal of T and $N \neq Q$, then clearly, $N \not\subseteq R$, for otherwise we conclude that $N \subseteq (R:T) \subseteq Q$ which is absurd. Thus (a) holds. For (b), assume that P is a prime ideal of T, if P is not contained in R, as we see in (a), P is a maximal ideal of T, otherwise $P \subseteq R$ and therefore $P \subseteq (R:T) \subseteq Q$. (3) is evident by our assumption in these cases and by (1) of Lemma 2.31. (d) is a consequence of (a), (c), our assumption in these cases and by Lemma 2.31, in fact the function $N \mapsto N \cap R$ is a one-one function from Max(T) into Max(R).

Acknowledgement

The author is grateful to the Research Council of Shahid Chamran University of Ahvaz (Ahvaz-Iran) for financial support (Grant Number: SCU.MM1402.721)

References

- D.D. Anderson and V. Camillo, Semigroups and rings whose zero products commute, Comm. Algebra. 27 (6) (1999) 2847-2852.
- [2] A. Azarang, Non-quasi duo rings have maximal subrings, Int. Math. Forum, 5 (20) (2010) 979-994.
- [3] Alborz Azarang, The conductors of Maximal subrings in Non-commutative Rings, preprint (2024). ArXiv:
- [4] Alborz Azarang, Conch maximal subrings, Comm. Algebra, 50 (3) (2022) 1267-1282.
- [5] A. Azarang, O.A.S. Karamzadeh, On the existence of maximal subrings in commutative artinian rings, J. Algebra Appl. 9 (5) (2010) 771-778.
- [6] A. Azarang, O.A.S. Karamzadeh, On Maximal Subrings of Commutative Rings, Algebra Colloq. 19 (Spec 1) (2012) 1125-1138.
- [7] A. Azarang, O.A.S. Karamzadeh, Most Commutative Rings Have Maximal Subrings, Algebra Colloq. 19 (Spec 1) (2012) 1139-1154.
- [8] A. Azarang, O.A.S. Karamzadeh, A. Namazi, Hereditary properties between a ring and its maximal subrings, Ukrainian. Math. J. 65 (7) (2013) 981-994.
- [9] L.I. Dechene, Adjacent Extensions of Rings, Ph.D. Dissertation, University of California, Riverside, (1978).
- [10] D.E. Dobbs and J. Shapiro, A classification of the minimal ring extensions of an integral domain, J. Algebra 305 (2006) 185-193.
- [11] D.E. Dobbs and J. Shapiro, A classification of the minimal ring extensions of certain commutative rings, J. Algebra 308 (2007) 800-821.
- [12] T. J. Dorsey, Z. Mesyan, On Minimal Extension of Rings, Comm. Algebra, 37 (10) (2009) 3463-3486.
- [13] D. Ferrand, J.-P. Olivier, Homomorphismes minimaux d'anneaux, J. Algebra 16 (1970) 461-471.
- [14] R. Gilmer, Some finiteness conditions on the set of overring an integral domain. Proc. Amer. Math. Soc. 131 (8) (2003) 2337-2346.
- [15] R. Gilmer and W.J. Heinzer, Intersection of quotient rings of an integral domain. J. Math Kyoto Univ. 7 (2) (1967) 133-150.
- [16] K. R. Goodearl, R. B. Warfield, JR, "An Introduction to Noncommutative Noetherian Rings", Camberidge University Press, Second Edition, 2004.
- [17] O.A.S. Karamzadeh, N. Nazari, On maximal commutative subrings of non-commutative rings. Comm. Algebra, 46 (12) (2018) 5083-5115.
- [18] A.A. Klein, The finiteness of a ring with a finite maximal subrings. Comm. Algebra, 21 (4) (1993) 1389-1392.
- [19] T.J. Laffey, A finiteness theorem for rings, Proc. R. Ir. Acad. 92 (2) (1992) 285-288.
- [20] T.Y. Lam, A First Course in Noncommutative Rings, Sec. Ed, Springer-Verlag, 2001.
- [21] T.Y. Lam, Lectures on Modules and Rings, Springer-Verlag, 1999.
- [22] T. Y. Lam and A. S. Dugas, Quasi-Duo Rings and Stable Range Descent, J. Pure Appl. Algebra 195 (2005) 243-259.
- [23] J. C. MacConnell and J. C. Robson, Noncommutative Noetherian Rings, Wilely-Interscience, New york, 1987.
- [24] M.L. Modica, Maximal Subrings, Ph.D. Dissertation, University of Chicago, 1975.
- [25] L. Rowen, Ring theory. Vol. I. Pure Appl. Math., 127 Academic Press, Inc., Boston, MA, 1988. xxiv+538 pp.
- [26] G. Picavet and M. Picavet-L'Hermitte. About minimal morphisms, in: Multiplicative ideal theory in commutative algebra, Springer-Verlag, New York, (2006) pp. 369-386.