

Cross-Dimensional Mathematics: A Foundation for STP/STA

Daizhan Cheng

Abstract—A new mathematical structure, called the cross-dimensional mathematics (CDM), is proposed. The CDM considered in this paper consists of three parts: hyper algebra, hyper geometry, and hyper Lie group/Lie algebra. Hyper algebra proposes some new algebraic structures such as hyper group, hyper ring, and hyper module over matrices and vectors with mixed dimensions (MVMDs). They have sets of classical groups, rings, and modules as their components and cross-dimensional connections among their components. Their basic properties are investigated. Hyper geometry starts from mixed dimensional Euclidian space, and hyper vector space. Then the hyper topological vector space, hyper inner product space, and hyper manifold are constructed. They have a joined cross-dimensional geometric structure. Finally, hyper metric space, topological hyper group and hyper Lie algebra are built gradually, and finally, the corresponding hyper Lie group is introduced. All these concepts are built over MVMDs, and to reach our purpose in addition to existing semi-tensor products (STPs) and semi-tensor additions (STAs), a couple of most general STP and STA are introduced. Some existing structures/results about STPs/STAs have also been resumed and integrated into this CDM.

Index Terms—Semi-tensor product/addition (STP/STA), MVMD, hyper group, hyper ring, hyper manifold, hyper linear algebra and hyper linear group.

I. INTRODUCTION

The semi-tensor product (STP) of matrices was initiated from 23 years ago[5]. It is a generalization of classical matrix product by removing the dimension restriction of the classical matrix product. Later on, the semi-tensor addition (STA) has also been proposed to weaken the dimension restriction of matrix addition [11]. The past two decades have witnessed the development of STP/STA[28], [41]. They have received various applications, including Boolean networks [38], finite games [14], dimension-varying systems [10], engineering problems [34], finite automata [51], coding [53], etc. In addition to thousands of papers, there are also many STP/STA monographs [9], [6], [7], [8], [10], [15], [17], [18], [20], [21],

This work is supported partly by the National Natural Science Foundation of China (NSFC) under Grants 62073315 and 62350037.

Key Laboratory of Systems and Control, Academy of Mathematics and Systems Sciences, Chinese Academy of Sciences, Beijing 100190, (e-mail: dcheng@iss.ac.cn).

[35], [37], [40], [52], and books with STP chapter or appendix [2], [46].

Roughly speaking, the STPs and STAs are cross-dimensional operators, which means they are operators acting on objects with different dimensions, i.e. MVMDs, which can formally be described as follows:

- (i) Vectors with mixed dimensions:

$$\mathbb{R}^\infty := \bigcup_{n=1}^{\infty} \mathbb{R}^n,$$

where \mathbb{R} may be replaced by \mathbb{F} , which is a field with characteristic number 0. To make the discussion and notation ease, throughout this paper \mathbb{F} is assumed to be \mathbb{R} unless elsewhere stated. It is easy to see that in the following discussions replacing \mathbb{R} by \mathbb{C} is mostly straightforward.

- (ii) Matrices with mixed dimensions:

$$\mathcal{M} = \bigcup_{m=1}^{\infty} \bigcup_{n=1}^{\infty} \mathcal{M}_{m \times n}.$$

In the long history of development of matrix theory, the algebraic structure of matrices, including group, ring, module, general linear algebra, etc., and their geometric structure, including vector space, inner product, metric, general linear group, etc., play fundamental roles in both theoretical investigations and practical applications. Since they were proposed two decades ago, the researches on STP/STA are mainly concentrated on its applications. When investigating the set of MVMDs, which are posed by cross-dimensional operators: STPs and STAs, the existing mathematical tools seem not very efficient. Hence, now seems to be the time to explore the mathematical foundation of STP/STA.

When we try to pose the existing algebraic and geometric structures on STPs/STAs with MVMDs, we found the existing mathematical framework is not suitable for them. The most critical contradiction lies on the fact that the existing algebraic or geometric structure, such as group or manifold, is for a set of objects with fixed dimensions or fixed sizes, while the STPs/STAs are cross-dimensional operators for MVMDs.

Observing carefully these objects with mixed dimensions, we found that the main difficult point for posing an algebraic

or geometric structure on them lies on “multiply unitary” trouble. For instance, if we consider the group structure, then in \mathbb{R}^∞ there are infinite zeros, precisely speaking, there are $0_n \in \mathbb{R}^n$, $n = 1, 2, \dots$. Similarly, \mathcal{M} also have infinite $0_{m \times n}$, $m, n = 1, 2, \dots$. This fact makes it almost impossible to pose a group structure to these objects (w.r.t. addition) If we consider the set of square matrices, denoted by \mathcal{M}_1 , then it has also infinite identities, I_n , $n = 1, 2, \dots$, which makes the group structure of matrices (w.r.t. product) also very difficult to define.

When we consider the geometric structure, say, vector space structure, we are also facing this “multiple zero” puzzle.

An inspiration for solving this problem comes from mechanics, where a pseudo-metric space is used.

Definition 1.1: [1] Consider $\mathcal{X} = (X, d)$, where X is a set, $d : X \times X \rightarrow \mathbb{R}$ is a mapping.

(1) \mathcal{X} is called a metric space if d satisfies

- (i) $d(x, y) \geq 0$, $x, y \in X$. Moreover, $d(x, y) = 0$, if and only if, $x = y$;
- (ii) Symmetry:

$$d(x, y) = d(y, x), \quad x, y \in X; \quad (1)$$

(iii) Triangular Inequality:

$$d(x, y) + d(y, z) \geq d(x, z), \quad x, y, z \in X. \quad (2)$$

(2) \mathcal{X} is a pseudo-metric space, if d satisfies (ii), (iii), and

- (i') $d(x, y) \geq 0$, $x, y \in X$. Moreover, there exists a zero set, $\mathbf{0}$ such that

$$d(x, x) = 0, \quad x \in \mathbf{0}. \quad (3)$$

The concept of “pseudo-metric space” seems can provide a way to address the distance of points with different dimensions. This stimulates us to define a new “group” which contains multiple identities. Instead of “pseudo-group” such a multi-identity group is called a “hyper group” later in this paper. The reason is that a hyper group contains a set of many (might be countable) groups as its components.

Motivated by this fact, we may modify the existing algebraic and geometric structures to suit the MVMDs with respect to (briefly, w.r.t.) STPs/STAs. The purpose of this paper is to propose a new mathematical framework, called the CDM, for objects with mixed dimensions. We hope that it can provide a solid mathematical foundation for STPs/STAs. Meanwhile, it may cause a development of classical fixed dimensional algebra and geometry into a new cross-dimensional algebra and geometry.

The CDM proposed in this paper consists of three parts: (1) hyper algebra; (2) hyper geometry; (3) hyper Lie algebra and




Fig. 1. Cross-Dimensional Mathematics vs Existing Math.

hyper Lie group. The hyper algebra contains hyper group, hyper ring, and hyper module; the hyper geometry includes hyper vector space, hyper inner product space, hyper manifold; and the hyper Lie algebra and hyper Lie group start from mixed dimensional Euclidian space, through constructing topological hyper group, and finally lead to hyper linear group and hyper linear algebra.

The relationship of CDM with existing mathematical structures is depicted in Figure 1.

Some key issues completed in this paper are emphasized as follows:

- (i) The most general matrix addition ($\hat{+}$) and matrix product ($\hat{\times}$) are proposed, which make the set of all matrices a hyper-ring $((\mathcal{M}, \hat{+}, \hat{\times}))$. Particularly, the hyper group structure of cross-dimensional permutations $\mathbf{S} = \bigcup_{n=1}^{\infty} \mathbf{S}_n$ is obtained.
- (ii) The most general Euclidian hyper vector space (\mathbb{R}^∞) is constructed. The vector space structure and the differentiable structure are proposed, which provide a coordinate frame and differentiable structure for hyper manifold. (This function is similar to \mathbb{R}^n for n -dimensional manifold.)
- (iii) The Lie group ($GL(m \times n, \mathbb{R})$) and Lie algebra ($gl(m \times n, \mathbb{R})$) structure for the set of $m \times n$ dimensional matrices is constructed. Their relationship with general linear group ($GL(n, \mathbb{R})$) and general linear algebra ($gl(n, \mathbb{R})$) is revealed. They provide a foundation for overall hyper matrix Lie group ($GL(\mathbb{R})$) and hyper Lie algebra ($gl(\mathbb{R})$).

The rest part of this paper is outlined as follows:

Section 2 provides some preliminaries including (i) algebraic structures of matrices/vectors such as matrix group, ring, module, and matrix Lie algebra; (ii) some commonly used STPs and STAs; (iii) dimension-keeping (DK-) STP and pseudo-STP, where pseudo-STP is newly defined. Sections 3-7 are concentrated on the hyper algebraic structure of MVMDs. Starting from lattice, Section 3 introduces hyper group, particularly, matrix/vector hyper groups. Their sub hyper groups. The group decomposition of hyper group into component groups and equivalence group is also presented. Section 4 considers the hyper group structure of permutations, including left and right permutation hyper groups. In Section 5 the hyper group homomorphisms are investigated. Three isomorphism theorems of groups have been extended to hyper groups. Symmetric and skew symmetric and squaring endomorphisms of matrix groups are defined and discussed, which extend the (skew) symmetry and eigenvalue/eigenvector of square matrices to non-square matrices. In Section 6 the hyper ring of matrices is proposed. Its sub hyper rings, homomorphisms, and some properties are investigated. Section 7 considers the hyper module of matrices. The general linear (control) systems are reconsidered under matrix hyper module framework. Sections 8-11 consider the hyper geometric structure of MVMDs. In Section 8 the hyper vector space is introduced. The inner product of vectors with different dimensions is introduced. Then the norm, the distance are also obtained. In Section 9 a metric space structure is posed on dimension-free Euclidian space \mathbb{R}^∞ , which turns this cross dimensional Euclidian space a path-wise connected topological space. Furthermore, in Section 10 a differentiable structure is also proposed to \mathbb{R}^∞ . The smooth functions, (co-) vector fields, distributions, and tensor fields are all introduced for \mathbb{R}^∞ . Section 11 considers the hyper manifold, where the topological and differentiable structure of \mathbb{R}^∞ is transferred to a topological manifold through homeomorphism, similarly to n -dimensional manifold w.r.t. \mathbb{R}^n . Sections 12-15 consider the hyper Lie group and hyper Lie algebra, which are emphasized on their expression on MVMDs as hyper linear group and hyper linear algebra. Section 12 constructs the general linear algebra ($gl(m \times n, \mathbb{R})$) and general linear group ($GL(m \times n, \mathbb{R})$) for non-square matrices. Their homomorphisms to classical general linear group ($GL(t, \mathbb{R})$) and general linear algebra ($gl(t, \mathbb{R})$) are also revealed. Section 13 considers how to construct a hyper Lie group from a topological hyper group. Section 14 investigates the structure of hyper general linear algebras based on hyper vector space and hyper rings. In Section 15 the hyper linear group is proposed. Its relationship with hyper linear algebra is revealed. Finally, Section 16 is a brief summary with some

remarks.

Before ending this section we give a list of notations.

- \mathbb{R} : The field of real numbers.
- \mathbb{C} : The field of complex numbers.
- \mathbb{Q} : The field of rational numbers.
- \mathbb{F} : A field of characteristic number 0, (in most cases $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$).
- \mathbb{Z} (\mathbb{Z}^+): Set of (positive) integers.
- $t|r$: $t, r \in \mathbb{Z}^+$ and $r/t \in \mathbb{Z}^+$.
- \mathbb{F}^n : n dimensional Euclidian space over field \mathbb{F} .
- $\mathcal{M}_{m \times n}^{\mathbb{F}}$: set of $m \times n$ matrices over \mathbb{F} (the superscript \mathbb{F} is mostly omitted).
- $\mathcal{M}_\mu := \{A_{m \times n} \mid m/n = \mu\}$ where $0 < \mu \in \mathbb{Q}$.
- $[a, b]$: the set of integers $\{a, a+1, \dots, b\}$, where $a \leq b$.
- $\text{lcm}(n, p)$ (or $n \vee p$) : the least common multiple of n and p .
- $\mathcal{D} = \{1, 2, \dots, n\}$.
- $\text{gcd}(n, p)$ (or $n \wedge p$) : the greatest common division of n and p .
- δ_n^i : the i -th column of the identity matrix I_n .
- $\Delta_n := \{\delta_n^i \mid i = 1, \dots, n\}$.
- $\mathbf{1}_k := \underbrace{(1, \dots, 1)}_k^T$.
- $\mathcal{E}_n := \frac{1}{\sqrt{n}} \mathbf{1}_n^k$.
- $\mathcal{E}_{m \times n} := E_m E_n^T$.
- $\mathcal{L}_{m \times n}$: the set of logical matrices, i.e. $\mathcal{L}_{m \times n} = \{A \in \mathcal{M}_{m \times n} \mid \text{Col}(A) \subset \Delta_m\}$.
- $\text{Col}(A)$: the set of columns of matrix A , and $\text{Col}_i(A)$ is the i -th column of A .
- $\text{Row}(A)$: the set of rows of matrix A , and $\text{Row}_i(A)$ is the i -th row of A .
- $\text{Span}(\cdot)$: subspace or dual subspace generated by \cdot .
- \mathbf{S}_n (\mathbf{S}): order n permutation group (hyper permutation group).
- \mathbf{A}_n (\mathbf{A}): order n alternative group (hyper alternative group).
- \perp : perpendicular relation between $x \in \mathbb{R}^n$ and $h \in \mathbb{R}_*^n$.
- $\bar{\dagger}$: STA w.r.t. $I_{n \times n}$.
- $\hat{\dagger}$: STA w.r.t. $\mathbf{1}_n$ or \mathcal{E}_n ($\mathcal{E}_{m \times n}$).
- \bowtie : (left) matrix-matrix semi-tensor product, which is the default one.
- \bowtie : right semi-tensor product of matrices.
- \circ : second (left) matrix-matrix semi-tensor product.
- $\vec{\bowtie}$: matrix-vector semi-tensor product.
- $\vec{\circ}$: second matrix-vector semi-tensor product.
- \times : dimension-keeping semi-tensor product of matrices.
- $\hat{\times}$: pseudo-STP.
- $\text{sup}(a, b)$ (or $a \vee b$): least common upper bound of a and

- b.
- $\inf(a, b)$ (or $a \wedge b$): greatest common lower bound of a and b .
- $\square(A)$: squaring matrix A .
- $GL(n, \mathbb{R})$ ($GL(\mathbb{R})$): (hyper) general linear group.
- $gl(n, \mathbb{R})$ ($gl(\mathbb{R})$): (hyper) general linear algebra.
- $H < G$ ($H \triangleleft G$): H is a (normal) subgroup of G .
- $H <_h G$ ($H \triangleleft_h G$): H is a (normal) hyper-subgroup of G .
- Id : identity mapping.
- Im : image set.
- Ker : kernel set.

II. PRELIMINARIES

A. Algebraic Structure on Matrices/Vectors with proper dimensions

We first review some algebraic structures over matrices/vectors.

Definition 2.1: Consider a couple $\mathcal{G} = (G, *)$, where G is a set, and $*$: $G \times G \rightarrow G$ is a binary operator.

- (i) \mathcal{G} is a semi-group, if

$$(a * b) * c = a * (b * c), \quad a, b, c \in G. \quad (4)$$

- (ii) A semi-group \mathcal{G} is a monoid (i.e., a semi-group with an identity), if there exists an identity $e \in G$, such that

$$a * e = e * a = a, \quad \forall a \in G \quad (5)$$

- (iii) A monoid \mathcal{G} is a group, if for each $a \in G$, there exists an a^{-1} , such that

$$a * a^{-1} = a^{-1} * a = e, \quad a \in G. \quad (6)$$

In addition, if $a * b = b * a$, $\forall a, b \in G$, then it is called an abelian group.

Example 2.2:

- (i) $(\mathbb{R}^n, +)$ is an abelian group with the identity $0_n \in \mathbb{R}^n$.
- (ii) $(\mathcal{M}_{m \times n}, +)$ is an abelian group with the identity $0_{m \times n} \in \mathcal{M}_{m \times n}$.
- (iii) $(\mathcal{M}_{n \times n}, +)$ is an abelian group with the identity 0_n .

– Set

$$\mathcal{S}_n := \{A \in \mathcal{M}_{n \times n} \mid A^T = A\},$$

$(\mathcal{S}_n, +) < (\mathcal{M}_{n \times n}, +)$ is the symmetric subgroup.

– Set

$$\mathcal{K}_n := \{A \in \mathcal{M}_{n \times n} \mid A^T = -A\},$$

$(\mathcal{K}_n, +) < (\mathcal{M}_{n \times n}, +)$ is the skew-symmetric subgroup.

– Set

$$\mathcal{Q}_n := \{A \in \mathcal{M}_{n \times n} \mid Tr(A) = 0\},$$

$(\mathcal{Q}_n, +) < (\mathcal{M}_{n \times n}, +)$ is the zero-trace subgroup. (There are also some other subgroups such as diagonal, upper-triangular, lower-triangular, etc.)

- (iv) $(\mathcal{M}_{n \times n}, \times)$ is a monoid with the identity I_n .

- (v) $\mathcal{T}_n := \{A \in \mathcal{M}_{n \times n} \mid A \text{ is invertible}\}$. Then (\mathcal{T}_n, \times) is a group with identity I_n .

– Set

$$\mathcal{O}_n := \{A \in \mathcal{T}_n \mid A^T = A^{-1}\},$$

$(\mathcal{O}_n, \times) < (\mathcal{T}_n, \times)$ is the orthogonal subgroup.

– Set

$$\mathcal{H}_n := \{A \in \mathcal{T}_n \mid \det(A) = 1\},$$

$(\mathcal{H}_n, \times) < (\mathcal{T}_n, \times)$ is the special linear subgroup.

– Set

$$\mathcal{P}_n := \mathcal{T}_n \cap \mathcal{L}_{n \times n},$$

where $\mathcal{L}_{n \times n}$ is the set of logical matrices (A is a logical matrix if $\text{Col}(A) \subset \Delta_n$), then $(\mathcal{P}_n, \times) < (\mathcal{T}_n, \times)$ is the permutation sub-group.

(there are also some other sub-groups such as non-singular diagonal, non-singular upper-triangular, non-singular lower-triangular, etc.)

Definition 2.3: Triple $\mathcal{R} = (R, +, *)$ is a ring, if

- (i) $(R, +)$ is an abelian group;
- (ii) $(R, *)$ is a semi-group;
- (iii) (Distributivity)

$$\begin{aligned} (a + b) * c &= a * c + b * c, & a, b, c \in R; \\ a * (b + c) &= a * b + a * c, & a, b, c \in R. \end{aligned} \quad (7)$$

In addition, if $(R, *)$ is a monoid, \mathcal{R} is a unitary ring; if $a * b = b * a$, $\forall a, b \in R$, \mathcal{R} is a commutative ring.

Definition 2.4: Assume $\mathcal{R} = (R, +, *)$ is a ring and $H \subset R$. If $(H, +, *)$ is also a ring, H is called a sub-ring of R .

Example 2.5: $(\mathcal{M}_{n \times n}, +, \times)$ is a unitary ring. It has some sub-rings as follows.

(i)

$$\mathcal{D}_n := \{A \in \mathcal{M}_{n \times n} \mid A \text{ is diagonal}\}. \quad (8)$$

Then $(\mathcal{D}_n, +, \times)$ is a sub-ring of $(\mathcal{M}_{n \times n}, +, \times)$.

(ii)

$$\mathcal{U}_n := \{A \in \mathcal{M}_{n \times n} \mid A \text{ is upper triangular}\}. \quad (9)$$

Then $(\mathcal{U}_n, +, \times)$ is a sub-ring of $(\mathcal{M}_{n \times n}, +, \times)$.

(iii)

$$\mathcal{B}_n := \{A \in \mathcal{M}_{n \times n} \mid A \text{ is lower triangular}\}. \quad (10)$$

Then $(\mathcal{B}_n, +, \times)$ is a sub-ring of $(\mathcal{M}_{n \times n}, +, \times)$.

Definition 2.6: Let $\mathcal{R} = (R, +, *)$ be a ring, and $\mathcal{G} = (G, +)$ be an abelian group. \mathcal{R}, \mathcal{G} with a mapping $\pi : \mathcal{R} \times \mathcal{G} \rightarrow \mathcal{G}$ is a (left) \mathcal{R} -module, if

(i)

$$r(a + b) = ra + rb, \quad r \in \mathcal{R}, \quad a, b \in \mathcal{G};$$

(ii)

$$(r + s)a = ra + sa, \quad r, s \in \mathcal{R}, \quad a, b \in \mathcal{G};$$

(iii)

$$r(sa) = (rs)a.$$

In addition, if there exists $e \in \mathcal{R}$ such that for all $a \in G$

$$ea = a, \quad \forall a \in G,$$

then \mathcal{G} is called a unitary \mathcal{R} -module.

Example 2.7:

(i) $(\mathcal{M}_{n \times n}, +, \times)$ is a ring with identity.(ii) $(\mathbb{R}^n, +)$ is an abelian group.(iii) Let $\pi : \mathcal{M}_{n \times n} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the classical matrix-vector product.

Then

$$((\mathcal{M}_{n \times n}, +, \times), (\mathbb{R}^n, +), \pi)$$

is a module.

Definition 2.8: [3] Let V be a vector space over \mathbb{R} and there is a mapping, called the Lie bracket and denoted by $[\cdot, \cdot] : V \times V \rightarrow V$. $(V, [\cdot, \cdot])$ is called a Lie algebra, if the followings are satisfied.

(i) Skew-Symmetry:

$$[x, y] = -[y, x], \quad x, y \in V. \quad (11)$$

(ii) Linearity:

$$[ax + by, z] = a[x, z] + b[y, z], \quad a, b \in \mathbb{R}; \quad x, y, z \in V. \quad (12)$$

(iii) Jacobi Identity:

$$[[x, y], z] + [[y, z], x] + [[z, x], y] = 0. \quad (13)$$

Example 2.9: Consider $\mathcal{M}_{n \times n}$. Define

$$[A, B] = AB - BA, \quad A, B \in \mathcal{M}_{n \times n}. \quad (14)$$

Then $(\mathcal{M}_{n \times n}, [\cdot, \cdot])$ becomes a Lie algebra, called the general linear algebra and denoted by $gl(n, \mathbb{R})$.

Note that all the above algebraic structures over matrices/vectors are restricted to a particular set of matrices with certain fixed dimensions. We refer to [31] or any other text book of abstract algebra for the related concepts. These concepts will be extended to MVMDs in this paper.

B. STP and STA of Matrices/Vectors

The main objects of linear algebra are vectors and matrices. In linear algebra both the addition and the product of two matrices, or matrix with vector, or two vectors, require the factors to meet certain dimension matching condition, here the product of two vectors means the inner product.

STP/STA are proposed to deal with MVMDs. The sets of vectors with mixed dimensions and matrices with mixed dimensions are described by \mathbb{R}^∞ and \mathcal{M} respectively. In addition we also need

$$\mathcal{M}_\mu = \{A \in \mathcal{M}_{m \times n} \mid m/n = \mu\}, \quad \mu \in \mathbb{Q}_+. \quad (15)$$

Remark 2.10: Consider \mathcal{M}_μ , where $\mu = \frac{\mu_y}{\mu_x} \in \mathbb{Q}_+$. Moreover, to be uniqueness, we assume $\gcd(\mu_y, \mu_x) = 1$. Then

(i) \mathcal{M}_1 is the set of square matrices.

(ii)

$$\mathcal{M} = \bigcup_{\mu \in \mathbb{Q}_+} \mathcal{M}_\mu$$

is a partition.

The STP and STA have been proposed to extend the matrix (vector) product and addition to overall (or larger than the original) sets. The followings are some commonly used ones.

Definition 2.11: [6], [10] Let $A, B \in \mathcal{M}$, specified by $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, and $t = \text{lcm}(n, p)$.

(i) The first type of matrix-matrix (MM-) STP of A, B , denoted by $A \times B$, is defined by

$$A \times B := (A \otimes I_{t/n}) (B \otimes I_{t/p}) \in \mathcal{M}_{mt/n \times qt/p}. \quad (16)$$

(ii) The second type of MM-STP of A, B , denoted by $A \circ B$, is defined by

$$A \circ B := (A \otimes J_{t/n}) (B \otimes J_{t/p}) \in \mathcal{M}_{mt/n \times qt/p}, \quad (17)$$

where

$$J_n := \frac{1}{n} \mathbf{1}_{n \times n}.$$

Definition 2.12: [7] Let $A \in \mathcal{M}$ and $x \in \mathbb{R}^\infty$, specified by $A \in \mathcal{M}_{m \times n}$, $x \in \mathbb{R}^p$, and $t = \text{lcm}(n, p)$.

(i) The first type of matrix-vector (MV-) STP of A and x , denoted by $A \vec{\times} x$, is defined by

$$A \vec{\times} x := (A \otimes I_{t/n}) (x \otimes \mathbf{1}_{t/p}) \in \mathbb{R}^{mt/n}. \quad (18)$$

(ii) The second type of MV-STP of A and x , denoted by $A \vec{\circ} x$, is defined by

$$A \vec{\circ} x := (A \otimes J_{t/n \times t/n}) (x \otimes \mathbf{1}_{t/p}) \in \mathbb{R}^{mt/n}. \quad (19)$$

Definition 2.13: [7] Let $x, y \in \mathbb{R}^\infty$, specified by $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, and $t = \text{lcm}(m, n)$. The vector-vector (VV-) STP of x and y , denoted by $x \bar{\cdot} y$, is defined by

$$x \bar{\cdot} y := \langle x \otimes \mathbf{1}_{t/m}, y \otimes \mathbf{1}_{t/n} \rangle \in \mathbb{R}, \quad (20)$$

where $\langle \cdot, \cdot \rangle$ is the classical inner product on Euclidian space.

Remark 2.14:

- (i) All the MM-STPs, MV-STPs, and VV-STP are the generalization of classical matrix product. That is, when the dimension-matching conditions for classical matrix-matrix (or matrix-vector or vector-vector) products are satisfied, they degenerate to classical products.
- (ii) All the MM-STPs and MV-STPs defined in(16), (18), and (20) are called the left STPs, because the main objects (A, B , etc.) are lying on left. They have their corresponding right STPs. Say,

$$(A \times B) := (I_{t/n} \otimes A) (I_{t/p} \otimes B), \quad (21)$$

etc. Such right extensions are also available for all other STPs defined in the sequel. And a parallel discussion shows similar properties of right STPs corresponding to the default left STPs. (We leave such discussions for the reader and will not mention this again.)

- (iii) All the MM-STPs, MV-STPs maintain most properties of the classical matrix product available. Particularly, the associativity and the distributivity hold. Say,

$$(A \times B) \times C = A \times (B \times C), \quad A, B, C \in \mathcal{M}; \quad (22)$$

and

$$\begin{aligned} (A + B) \times C &= A \times C + B \times C; \\ C \times (A + B) &= C \times A + C \times B, \quad A, B, C \in \mathcal{M}, \end{aligned} \quad (23)$$

etc.

Definition 2.15: Some STAs of matrices/vectors are defined as follows:

- (i) \mathbb{R}^∞ : Assume $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, and $t = \text{lcm}(m, n)$, then

$$x \pm y := (x \otimes \mathbf{1}_{t/m}) \pm (y \otimes \mathbf{1}_{t/n}), \quad (24)$$

where

$$\mathbf{1}_n := \underbrace{(1, 1, \dots, 1)}_n^T, \quad n \in \mathbb{Z}_+.$$

- (ii) \mathcal{M}_1 : Assume $A \in \mathcal{M}_{m \times m}$, $B \in \mathcal{M}_{n \times n}$, and $t = \text{lcm}(m, n)$, then

$$A \pm B := (A \otimes I_{t/m}) \pm (B \otimes I_{t/n}). \quad (25)$$

An alternative STA is defined as follows.

$$A \hat{\pm} B := (A \otimes J_{t/m}) \pm (B \otimes J_{t/n}). \quad (26)$$

- (iii) \mathcal{M}_μ : $A \in \mathcal{M}_{m\mu_y \times m\mu_x}$, $B \in \mathcal{M}_{n\mu_y \times n\mu_x}$, and $t = \text{lcm}(m, n)$: Its additivity can be define by either \pm , as in (25), or $\hat{\pm}$, as in (26).

To define ‘‘addition’’ for more general case, we need a new set of matrices as ‘‘weights’’. Set

$$\begin{aligned} \mathcal{E}_n &:= \frac{1}{\sqrt{n}} \mathbf{1}_n; \\ \mathcal{E}_{m \times n} &:= \mathcal{E}_m \mathcal{E}_n^T = \frac{1}{\sqrt{mn}} \mathbf{1}_{m \times n}, \quad m, n \in \mathbb{Z}_+. \end{aligned} \quad (27)$$

Then the following properties of this set of matrices are easily verifiable, but they are fundamental in further discussion.

Proposition 2.16:

$$\begin{cases} \mathcal{E}_n^T \mathcal{E}_n = 1; \\ \mathcal{E}_n \otimes \mathcal{E}_m = \mathcal{E}_m \otimes \mathcal{E}_n = \mathcal{E}_{mn}; \\ \mathcal{E}_{m \times n} \otimes \mathcal{E}_{p \times q} = \mathcal{E}_{p \times q} \otimes \mathcal{E}_{m \times n} = \mathcal{E}_{mp \times nq}; \\ \mathcal{E}_{m \times n}^T = \mathcal{E}_{n \times m}; \\ J_n = \mathcal{E}_{n \times n}. \end{cases} \quad (28)$$

Using this set of weights, some other STAs can be defined.

Definition 2.17:

- (i) \mathbb{R}^∞ : Assume $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, and $t = \text{lcm}(m, n)$, then

$$x \hat{\pm} y := (x \otimes \mathcal{E}_{t/m}) \pm (y \otimes \mathcal{E}_{t/n}), \quad (29)$$

- (ii) Consider \mathcal{M} , and assume $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $s = \text{lcm}(m, p)$, and $t = \text{lcm}(n, q)$. Then we define the STA over \mathcal{M} as follows:

$$A \hat{\pm} B := (A \otimes \mathcal{E}_{s/m \times t/n}) \pm (B \otimes \mathcal{E}_{s/p \times t/q}). \quad (30)$$

Note that all the STPs and STAs, defined up to this time are the generalizations of the classical matrix product/addition. That is, when the dimension matching conditions required by the classical products are satisfied, they degenerated to the classical matrix product/addition.

Before ending this subsection we give a formula for calculating equivalence transformation, which will be used in the sequel.

Proposition 2.18: Let $A \in \mathcal{M}_{m \times n}$, and $B \in \mathcal{M}_{p \times q}$. Then

$$\begin{aligned} (A \otimes \mathcal{E}_{r \times kp}) (B \otimes \mathcal{E}_{kn \times s}) &= (A \otimes \mathcal{E}_{r \times p}) (B \otimes \mathcal{E}_{n \times s}), \\ (A \otimes \mathcal{E}_{kr \times p}) (B \otimes \mathcal{E}_{n \times ks}) &= (A \otimes \mathcal{E}_{r \times p}) (B \otimes \mathcal{E}_{n \times s}). \end{aligned} \quad (31)$$

Proof. We prove the first equality only. The proof of the second one is similar.

$$\begin{aligned} \text{LHS of (31)} &= (A \otimes \mathcal{E}_{r \times p} \otimes \mathcal{E}_k^T) (B \otimes \mathcal{E}_{n \times s} \otimes \mathcal{E}_k) \\ &= [(A \otimes \mathcal{E}_{r \times p}) (B \otimes \mathcal{E}_{n \times s})] \otimes (\mathcal{E}_k^T \mathcal{E}_k) \\ &= (A \otimes \mathcal{E}_{r \times p}) (B \otimes \mathcal{E}_{n \times s}) = \text{RHS of (31)}. \end{aligned}$$

□

C. DK-STP and Pseudo-STP

The dimension-keeping (DK-) STP is defined as follows.

Definition 2.19: Let $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, and $t = \text{lcm}(n, p)$. The DK-STP of A and B , denoted as $A \times B$, is defined by

$$A \times B := \left(A \otimes \mathcal{E}_{t/n}^T \right) \left(B \otimes \mathcal{E}_{t/p} \right). \quad (32)$$

Remark 2.20: DK-STP was newly proposed in [22] as follows:

$$A \times B := \left(A \otimes \mathbf{1}_{t/n}^T \right) \left(B \otimes \mathbf{1}_{t/p} \right). \quad (33)$$

The original one defined by (33) has very clear physical meaning. Multiplying the columns of the first matrix and multiplying the rows of the second matrix to least times, such that the enlarged factor matrices meet the dimension matching condition of the classical matrix product.

In fact, the modified Definition 2.19 here uses certain weights for the multiplication, which keeps the ‘‘norms’’ of the enlarged matrices unchanged. In the sequel you will see that this modification is indeed meaningful. In [23] (32) is called a weighted DK-STP.

An obvious property of DK-STP is: if $A, B \in \mathcal{M}_{m \times n}$, the $A \times B \in \mathcal{M}_{m \times n}$. That is why it is named by ‘‘dimension-keeping’’.

The following properties about DK-STP are mimic to the one in [22].

Proposition 2.21: Define a bridge matrix as

$$\Psi_{n \times p} := \left(I_n \otimes \mathcal{E}_{t/n}^T \right) \left(I_p \otimes \mathcal{E}_{t/p} \right), \quad n, p \in \mathbb{Z}_+, \quad (34)$$

where $t = \text{lcm}(n, p)$, then

$$A \times B = A \Psi_{n \times p} B, \quad A \in \mathcal{M}_{m \times n}, B \in \mathcal{M}_{p \times q}. \quad (35)$$

Remark 2.22: Corresponding to (33), the bridge matrix becomes

$$\Psi'_{n \times p} := \left(I_n \otimes \mathbf{1}_{t/n}^T \right) \left(I_p \otimes \mathbf{1}_{t/p} \right), \quad n, p \in \mathbb{Z}_+, \quad (36)$$

which is only different from (34) by a coefficient. That is,

$$\Psi_{n \times p} := \frac{\sqrt{np}}{t} \Psi'_{n \times p}. \quad (37)$$

Using Proposition 2.21, the following results are obvious.

Proposition 2.23:

- (i) \times is a generalization of classical matrix product. That is, if $n = p$ it turns out that $A \times B = AB$.
- (ii) (Distributivity)

$$\begin{aligned} (A + B) \times C &= A \times C + B \times C, \\ C \times (A + B) &= C \times A + C \times B, \end{aligned} \quad (38)$$

(where A and B are of the same dimension).

- (iii) (Associativity)

$$(A \times B) \times C = A(B \times C). \quad (39)$$

- (iv)

$$(A \times B)^T = B^T \times A^T. \quad (40)$$

DK-STP brings a new algebraic structure to non-square matrices. The following result is interesting and useful.

Proposition 2.24: [22] $(\mathcal{M}_{m \times n}, +, \times)$ is a ring.

Using DK-STP, we can define a pseudo-STP as follows:

Definition 2.25: Consider \mathcal{M} , and assume $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $s = \text{lcm}(m, p)$, and $t = \text{lcm}(n, q)$. Then the pseudo-STP of A and B is defined as follows:

$$A \hat{\times} B := \left(A \otimes \mathcal{E}_{s/m \times t/n} \right) \times \left(B \otimes \mathcal{E}_{s/p \times t/q} \right). \quad (41)$$

Remark 2.26:

- (i) As a special case, we consider \mathcal{M}_μ , and assume $A \in \mathcal{M}_{m \mu_y \times m \mu_x}$, $B \in \mathcal{M}_{n \mu_y \times n \mu_x}$, and $t = \text{lcm}(m, n)$. Then the pseudo-STP for \mathcal{M}_μ becomes:

$$A \hat{\times} B := \left(A \otimes J_{t/m} \right) \times \left(B \otimes J_{t/n} \right). \quad (42)$$

- (ii) Comparing (41) with (30), one sees easily that in certain sense this pseudo-STP is ‘‘consistent’’ with the STA defined by (30). It will be seen in the sequel that this consistence is very meaningful.
- (iii) The reason to call (41) or (42) a pseudo-STP is: they do not always consistent with classical matrix product even if the dimension matching condition is satisfied. So they do not meet the fundamental requirement of STP.

Next, we explore some basic properties of pseudo-STP.

Proposition 2.27: Let $A, B \in \mathcal{M}$. Then

- (i)

$$(A \hat{\times} B)^T = B^T \hat{\times} A^T. \quad (43)$$

- (ii)

$$(A \hat{+} B)^T = A^T \hat{+} B^T. \quad (44)$$

Proof.

We prove (43) only. The proof of (44) is similar.

Assume $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $\xi = \text{lcm}(m, p)$, $\eta = \text{lcm}(n, q)$, and $\delta = \text{lcm}(\xi, \eta)$. Then

$$\begin{aligned} A \hat{\times} B &= \left(A \otimes \mathcal{E}_{\xi/m \times \eta/n} \right) \times \left(B \otimes \mathcal{E}_{\xi/p \times \eta/q} \right) \\ &= \left(A \otimes \mathcal{E}_{\xi/m \times \eta/n} \otimes \mathcal{E}_{\delta/\eta}^T \right) \left(B \otimes \mathcal{E}_{\xi/p \times \eta/q} \otimes \mathcal{E}_{\delta/\xi} \right) \\ &= \left(A \otimes \mathcal{E}_{\xi/m \times \delta/n} \right) \left(B \otimes \mathcal{E}_{\delta/p \times \eta/q} \right). \end{aligned}$$

It follows that

$$(A \hat{\times} B)^T = \left(B^T \otimes \mathcal{E}_{\eta/q \times \delta/p} \right) \cdot \left(A^T \otimes \mathcal{E}_{\delta/n \times \xi/m} \right)$$

On the other hand,

$$\begin{aligned} B^T \hat{\times} A^T &= (B^T \otimes \mathcal{E}_{\eta/q \times \xi/p}) \times (A^T \otimes \mathcal{E}_{\eta/n \times \xi/m}) \\ &= (B^T \otimes \mathcal{E}_{\eta/q \times \delta/p}) (A^T \otimes \mathcal{E}_{\delta/n \times \xi/m}). \end{aligned}$$

The conclusion follows. \square

Proposition 2.28: $(\mathcal{M}, \hat{\times})$ is a semi-group. That is, assume $A, B, C \in \mathcal{M}$, then

$$(A \hat{\times} B) \hat{\times} C = A \hat{\times} (B \hat{\times} C). \quad (45)$$

Proof. Assume

$$A \in \mathcal{M}_{m \times n}, B \in \mathcal{M}_{p \times q}, C \in \mathcal{M}_{r \times s},$$

and

$$\begin{aligned} m \vee p &= \xi, & n \vee q &= \eta, & \xi \vee \eta &= \delta, & \xi \vee r &= \alpha, \\ \eta \vee s &= \beta, & \alpha \vee \beta &= \theta, & p \vee r &= a, & q \vee s &= b, \\ a \vee b &= e, & m \vee a &= m \vee p \vee r &= \alpha, \\ n \vee b &= n \vee q \vee s = \beta, & n \vee p &= \mu, \\ q \vee r &= \lambda, \end{aligned}$$

Then

$$\begin{aligned} A \hat{\times} B &= (A \otimes \mathcal{E}_{\xi/m \times \eta/n}) \times (B \otimes \mathcal{E}_{\xi/p \times \eta/q}) \\ &= (A \otimes \mathcal{E}_{\xi/m \times \eta/n} \otimes \mathcal{E}_{\delta/\eta}^T) (B \otimes \mathcal{E}_{\xi/p \times \eta/q} \otimes \mathcal{E}_{\delta/\xi}) \\ &= (A \otimes \mathcal{E}_{\xi/m \times \delta/n}) (B \otimes \mathcal{E}_{\delta/p \times \eta/q}) \in \mathcal{M}_{\xi \times \eta}, \end{aligned}$$

it follows that

$$\begin{aligned} (A \hat{\times} B) \hat{\times} C &= \{[(A \otimes \mathcal{E}_{\xi/m \times \delta/n}) (B \otimes \mathcal{E}_{\delta/p \times \eta/q})] \\ &\quad \otimes \mathcal{E}_{\alpha/\xi \times \beta/\eta}\} \times (C \otimes \mathcal{E}_{\alpha/r \times \beta/s}) \\ &= \{[(A \otimes \mathcal{E}_{\xi/m \times \delta/n}) (B \otimes \mathcal{E}_{\delta/p \times \eta/q})] \\ &\quad \otimes (\mathcal{E}_{\alpha/\xi} \mathcal{E}_{\beta/\eta}^T)\} \times (C \otimes \mathcal{E}_{\alpha/r \times \beta/s}) \\ &= \{[(A \otimes \mathcal{E}_{\alpha/m \times \delta/n}) (B \otimes \mathcal{E}_{\delta/p \times \beta/q})] \\ &\quad \times (C \otimes \mathcal{E}_{\alpha/r \times \beta/s})\} \\ &= \left(\{[(A \otimes \mathcal{E}_{\alpha/m \times \delta/n}) (B \otimes \mathcal{E}_{\delta/p \times \beta/q})] \} \otimes \mathcal{E}_{\theta/\alpha}^T \right) \\ &\quad ((C \otimes \mathcal{E}_{\alpha/r \times \beta/s}) \otimes \mathcal{E}_{\theta/\alpha}) \\ &= (A \otimes \mathcal{E}_{\alpha/m \times \delta/n}) (B \otimes \mathcal{E}_{\delta/p \times \theta/q}) (C \otimes \mathcal{E}_{\theta/r \times \beta/s}) \\ &= (A \otimes \mathcal{E}_{\alpha/m \times \mu/n}) (B \otimes \mathcal{E}_{\mu/p \times \lambda/q}) (C \otimes \mathcal{E}_{\lambda/r \times \beta/s}). \end{aligned}$$

On the other hand,

$$\begin{aligned} B \hat{\times} C &= (B \otimes \mathcal{E}_{a/p \times b/q}) \times (C \otimes \mathcal{E}_{a/r \times b/s}) \\ &= (B \otimes \mathcal{E}_{a/p \times e/q}) (C \otimes \mathcal{E}_{e/r \times b/s}) \in \mathcal{M}_{a \times b}, \end{aligned}$$

and then

$$\begin{aligned} A \hat{\times} (B \hat{\times} C) &= (A \otimes \mathcal{E}_{\alpha/m \times \beta/n}) \times \\ &\quad \{[(B \otimes \mathcal{E}_{a/p \times e/q}) (C \otimes \mathcal{E}_{e/r \times b/s})] \otimes \mathcal{E}_{\alpha/a \times \beta/b}\} \\ &= (A \otimes \mathcal{E}_{\alpha/m \times \beta/n}) \times [(B \otimes \mathcal{E}_{a/p \times e/q}) (C \otimes \mathcal{E}_{e/r \times \beta/s})] \\ &= [(A \otimes \mathcal{E}_{\alpha/m \times \beta/n}) \otimes \mathcal{E}_{\theta/\beta}^T] \\ &\quad \{[(B \otimes \mathcal{E}_{a/p \times e/q}) (C \otimes \mathcal{E}_{e/r \times \beta/s})] \otimes \mathcal{E}_{\theta/\alpha}\} \\ &= (A \otimes \mathcal{E}_{\alpha/m \times \theta/n}) (B \otimes \mathcal{E}_{\theta/p \times e/q}) (C \otimes \mathcal{E}_{e/r \times \beta/s}) \\ &= (A \otimes \mathcal{E}_{\alpha/m \times \mu/n}) (B \otimes \mathcal{E}_{\mu/p \times \lambda/q}) (C \otimes \mathcal{E}_{\lambda/r \times \beta/s}). \end{aligned}$$

The conclusion follows. \square




Fig. 2. Hasse Diagram

Before end of this section, we would like to summarize the new algebraic structures over MVMDs by introducing STPs and STAs.

Remark 2.29: There are some new algebraic structures, which are caused by STPs/STAs.

(i) New Algebraic Structures:

- (\mathcal{M}, \times) is a monoid with identity 1.
- (\mathcal{M}, \circ) is a semi-group.
- $(\mathcal{M}_{m \times n}, +, \times)$ is a ring.
- $(\mathcal{M}, \hat{\times})$ is a semi-group.
- $(\mathcal{M}_\mu, \hat{\times}) < (\mathcal{M}, \hat{\times})$ is a sub-semi-group.

(ii) Observing the aforementioned STAs and STPs over MVMDs, in many cases it seems difficult to introduce some algebraic structures on them, because the MVMDs always have “multiple identities”. Recall the pseudometric space introduced in the Introduction, we may introduce a new algebraic structure, which allows multiple identities. The “hyper group”, which will be proposed in next section is one of such objects.

III. HYPER GROUP

A. Lattice

As a preparation, we briefly review the basic concepts of lattice.

We refer to [4] or [27] for some basic concepts and properties of lattice.

Definition 3.1: A partial order set Λ is a lattice, if for any two elements $\lambda, \mu \in \Lambda$ there exist a least common upper bound, denoted by $\sup(\lambda, \mu)$ (or $\lambda \vee \mu$), and a greatest common lower bound, denoted by $\inf(\lambda, \mu)$ (or $\lambda \wedge \mu$).

A lattice can be described by a graph called the Hasse diagram. A Hasse diagram is a graph with no horizontal sides. The order of nodes is decided by oblique sides. For instance, in Figure 2 the left is a lattice, say, $a \vee b = c$ and $a \wedge b = d$. The right is not a lattice, say neither the $a \vee b$ no the $a \wedge b$ exists.

In the following example, we give two lattices which will be used in the sequel.

Example 3.2:

- (i) Consider \mathbb{Z}_+ . Set $a \prec b$ if $a \neq b$ and $a|b$, i.e. a is a proper factor of b . It follows that

$$a \vee b = \text{lcm}(a, b); \quad a \wedge b = \text{gcd}(a, b).$$

Then (\mathbb{Z}_+, \prec) is a lattice. We call it the multiply-division lattice of dimension 1, briefly, MD-1 lattice.

- (ii) Consider $\mathbb{Z}_+ \times \mathbb{Z}_+$. Set $(a, c) \prec (b, d)$ if both $a \prec b$ and $c \prec d$ (defined as in (i)). It follows that

$$\begin{aligned} (a, c) \vee (b, d) &= (\text{lcm}(a, b), \text{lcm}(c, d)); \\ (a, c) \wedge (b, d) &= (\text{gcd}(a, b), \text{gcd}(c, d)). \end{aligned}$$

Then $(\mathbb{Z}_+ \times \mathbb{Z}_+, \prec)$ is a lattice, called the MD-2 lattice. Similarly, we can defined the MD-k lattice for $k > 2$.

Definition 3.3: Let L and H be two lattices.

- (i) A mapping $\pi : L \rightarrow H$ is said to be a lattice homomorphism if

$$\pi(\lambda \vee \mu) = \pi(\lambda) \vee \pi(\mu), \quad \lambda, \mu \in L; \quad (46)$$

and

$$\pi(\lambda \wedge \mu) = \pi(\lambda) \wedge \pi(\mu). \quad (47)$$

- (ii) A homomorphism $\pi : L \rightarrow H$ is said to be a lattice isomorphism if it is bijective and its inverse is also a homomorphism.
- (iii) A mapping $\pi : L \rightarrow H$ is said to be an order-preserving mapping, if

$$a \prec_L b \Rightarrow \pi(a) \prec_H \pi(b), \quad a, b \in L. \quad (48)$$

The following result comes from definition immediately.

Proposition 3.4: Let L and H be two lattices, and $\pi : L \rightarrow H$ be a bijective mapping. If both π and π^{-1} are order-preserving mappings, then π is a lattice isomorphism.

Definition 3.5: Let $(L, \text{sup}, \text{inf})$ be a lattice.

- (i) $H \subset L$, H is called a sublattice of L , if $(H, \text{sup}, \text{inf})$ is also a lattice.
- (ii) H is a sublattice of L . If for any $a \in L$, there exists an $h \in H$ such that $a \prec h$ implies $a \in H$, then H is called an ideal of L .
- (iii) H is a sublattice of L . If for any $a \in L$, there exists an $h \in H$ such that $a \succ h$ implies $a \in H$, then H is called a filter of L .

B. Hyper Group – A “Group” with Multiple Identities

This subsection proposes a “group” with a set of identities, called the identity set. To begin with, we define an identity set.

Definition 3.6: Consider a semi-group $\mathcal{G} = (G, *)$. $e \subset G$ is called an identity set of \mathcal{G} , if

- (i)

$$g * e = e * g, \quad \forall g \in G, \forall e \in e. \quad (49)$$

- (ii) There is a lattice Λ as the index set of e , that is,

$$e = \{e_\lambda \mid \lambda \in \Lambda\}.$$

Moreover,

$$e_\lambda * e_\mu = e_{\lambda \vee \mu}, \quad \lambda, \mu \in \Lambda. \quad (50)$$

- (iii) For each $g \in G$ there exists a unique $e_g = e_{\lambda_g} \in e$ such that

$$e_\lambda * g = g, \quad (51)$$

if and only if, $\lambda \prec \lambda_g$, (including $\lambda = \lambda_g$).

Definition 3.7: Consider a semi-group $\mathcal{G} = (G, *)$.

- (i) $\mathcal{G} = (G, *)$ is a hyper-monoid, if there exists an identity set $e \subset G$.
- (ii) A hyper-Monoid $\mathcal{G} = (G, *)$ is a hyper group, if for each $g \in G$, there exists a $g^{-1} \in G$ such that

$$g * g^{-1} = g^{-1} * g = e_g. \quad (52)$$

In addition, if $a * b = b * a$, $\forall a, b \in G$, then it is called an abelian hyper-monoid/hyper group.

C. Hyper Groups Over MVMDs

Recall the additions over MVMDs, defined in Definitions 2.15 and 2.17, we have the following result.

Proposition 3.8:

- (i) $(\mathbb{R}^\infty, \bar{+})$ with $\bar{+}$ defined by (24) and $(\mathbb{R}^\infty, \hat{+})$ with $\hat{+}$ defined by (29) are hyper groups, where the identity set is

$$e = \{0_n \in \mathbb{R}^n \mid n \in \mathbb{Z}_+\}, \quad (53)$$

w.r.t. MD-1 lattice.

- (ii) $(\mathcal{M}_1, \bar{+})$ is a hyper group with $\bar{+}$ defined by (25), where the identity set is

$$e = \{0_{n \times n} \in \mathcal{M}_{n \times n} \mid n \in \mathbb{Z}_+\}, \quad (54)$$

w.r.t. MD-1 lattice.

Also, $(\mathcal{M}_1, \hat{+})$ is a hyper group with $\hat{+}$ defined by (26), where the identity set is the same as (54).

(iii) $(\mathcal{M}_\mu, \bar{\dagger})$ (or $(\mathcal{M}_\mu, \hat{\dagger})$) is a hyper group with $\bar{\dagger}$ defined by (25) (or $\hat{\dagger}$ defined by (26)), where the identity set is

$$\mathbf{e} = \{0_{n\mu_y \times n\mu_x} \in \mathcal{M}_{n\mu_y \times n\mu_x} \mid n \in \mathbb{Z}_+\}, \quad (55)$$

w.r.t. MD-1 lattice.

(iv) $(\mathcal{M}, \hat{\dagger})$ is a hyper group with $\hat{\dagger}$ defined by (29), where the identity set is

$$\mathbf{e} = \{0_{m \times n} \in \mathcal{M}_{m \times n} \mid (m, n) \in \mathbb{Z}_+ \times \mathbb{Z}_+\}, \quad (56)$$

w.r.t. MD-2 lattice.

Proof. We prove (iv) only. The proofs for others are similar.

First, we show that $(\mathcal{M}, \hat{\dagger})$ is a semi-group. Assume $A \in \mathcal{M}_{m \times n}$, $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $C \in \mathcal{M}_{r \times s}$, and

$$\begin{aligned} m \vee p = a, \quad p \vee r = c, \quad m \vee p \vee r = \xi, \\ n \vee q = b, \quad q \vee s = d, \quad n \vee q \vee s = \eta. \end{aligned}$$

Then

$$\begin{aligned} (A \hat{\dagger} B) \hat{\dagger} C &= [(A \otimes \mathcal{E}_{a/m \times b/n}) + (B \otimes \mathcal{E}_{a/p \times b/q})] \hat{\dagger} C \\ &= [(A \otimes \mathcal{E}_{a/m \times b/n}) + (B \otimes \mathcal{E}_{a/p \times b/q})] \otimes \mathcal{E}_{\xi/a \times \eta/b} \\ &\quad + C \otimes \mathcal{E}_{\xi/r \times \eta/s} \\ &= (A \otimes \mathcal{E}_{\xi/m \times \eta/n}) + (B \otimes \mathcal{E}_{\xi/p \times \eta/q}) + (C \otimes \mathcal{E}_{\xi/r \times \eta/s}) \end{aligned}$$

A similar calculation shows that

$$\begin{aligned} A \hat{\dagger} (B \hat{\dagger} C) &= (A \otimes \mathcal{E}_{\xi/m \times \eta/n}) \\ &\quad + (B \otimes \mathcal{E}_{\xi/p \times \eta/q}) + (C \otimes \mathcal{E}_{\xi/r \times \eta/s}) \end{aligned}$$

Hence $(\mathcal{M}, \hat{\dagger})$ is a semi-group, which is obviously abelian.

Then we consider the identity set \mathbf{e} .

Let $A \in \mathcal{M}_{m \times n}$, $0_{p \times q} \in \mathbf{e}$, $m \vee p = a$, and $n \vee q = b$.

Then a straightforward computation shows that

$$A \hat{\dagger} 0_{p \times q} = 0_{p \times q} \hat{\dagger} A = A \otimes \mathcal{E}_{a/m \times b/n}.$$

(49) holds.

Next, it is ready to verify that the lattice structure of (56) satisfies (50).

Finally, assume A as before, then it is obvious that $e_A = 0_{m \times n} \in \mathbf{e}$, which leads to

$$A + (-A) = e_A.$$

(51) is satisfied. \square

Remark 3.9: All the additive hyper groups are abelian.

Next, we consider the product hyper groups.

Set

$$\mathcal{T} := \bigcup_{i=1}^{\infty} \mathcal{T}_n.$$

Then we have the product hyper group of matrices as follows.

Proposition 3.10: (\mathcal{T}, \times) is a hyper group with \times defined by (16), where the identity set is

$$\mathbf{e} = \{I_n \in \mathcal{M}_{n \times n} \mid n \in \mathbb{Z}_+\}, \quad (57)$$

w.r.t. MD-1 lattice.

Proof.

It is well known that (\mathcal{M}, \times) is a semi-group. Let $A, B \in \mathcal{T}$. It is easy to verify that if $A \times B \in \mathcal{T}$. Hence, (\mathcal{T}, \times) is a sub-semi-group.

Let $A \in \mathcal{T}_m$, $I_n \in \mathbf{e}$, and $t = m \vee n$. Then

$$A \times I_n = I_n \times A = A \otimes I_{t/m}.$$

(49) holds.

It is obvious that $e_A = I_m$ and

$$A \times A^{-1} = AA^{-1} = I_m.$$

(50) holds.

Finally, the lattice structure of the identity set is obvious. Moreover,

$$I_m \times I_n = I_t.$$

(51) is also true. \square

To emphasize the identity set of a hyper group, we may also denote a hyper group by a triple $(G, *, \mathbf{e})$.

D. Component Groups and Equivalence Group

This subsection considers the component groups and equivalence group of a hyper group.

Proposition 3.11: Assume $\mathcal{G} = (G, *, \mathbf{e})$ is a hyper group, where \mathbf{e} is determined by a lattice Λ . Assign for each $e_\lambda \in \mathbf{e}$, $\lambda \in \Lambda$ a set $G_\lambda \subset G$ as

$$G_\lambda := \{x \in G \mid e_x = e_\lambda\}. \quad (58)$$

Then

- (i) For each $\lambda \in \Lambda$, $\mathcal{G}_\lambda = (G_\lambda, *)$ is a group, called the component group of \mathcal{G} , and its identity is e_λ .
- (ii)

$$G = \bigcup_{\lambda \in \Lambda} G_\lambda \quad (59)$$

is a partition.

- (iii) If $x \in G_\lambda$ and $y \in G_\mu$, then $x * y \in G_{\lambda \vee \mu}$.

Proof.

- (i) To see G_λ is a group, we have to show that if $x, y \in G_\lambda$, then $x * y \in G_\lambda$. It obvious that

$$x * y * e_\lambda = e_\lambda * x * y.$$

We also have

$$x * y * y^{-1} x^{-1} = x * e_\lambda x^{-1} = e_\lambda,$$

and similarly, we have

$$y^{-1} x^{-1} * x * y = e_\lambda.$$

Hence, we have that both $(x * y)^{-1} = y^{-1} * x^{-1}$ and $x * y$ are $\in G_\lambda$.

It is clear from definition that if $x \in G_\lambda$ then $x^{-1} \in G_\lambda$.

Then the associativity and the uniqueness of the inverse are inherited from the properties of hyper group.

We conclude that G_λ is a group.

(ii) The partition (59) comes from definition immediately.

(iii) Since

$$\begin{aligned} x * y * y^{-1} * x^{-1} &= x * e_\mu * x^{-1} = e_\mu * x * x^{-1} \\ &= e_{m\mu} * e_\lambda = e_{\lambda \vee \mu}. \end{aligned}$$

By definition $x * y \in G_{\lambda \vee \mu}$.

□

Definition 3.12: Consider a hyper group $\mathcal{G} = (G, *, e)$. Let $x, y \in G$. x and y are said to be equivalent, denoted by $x \sim y$, if there exist e_α and e_β such that

$$x * e_\alpha = y * e_\beta. \quad (60)$$

The equivalence class of x , denoted by \bar{x} , is

$$\bar{x} := \{y \in G \mid y \sim x\}.$$

Note that we have to show that (60) is an equivalence relation. A straightforward verification ensures this.

Definition 3.13: Consider a hyper group $\mathcal{G} = (G, *, e)$. The equivalence \sim is said to be consistent with the group operator $*$, if the followings are satisfied.

(i) If $x_1 \sim x_2$ and $y_1 \sim y_2$, then

$$x_1 * y_1 \sim x_2 * y_2. \quad (61)$$

(ii) If there exists $e \in \mathbf{e}$ such that $x \sim e$, then $x \in \mathbf{e}$.

Then we have the identity set as one element.

Proposition 3.14: Consider a hyper group $\mathcal{G} = (G, *, e)$. If the equivalence is consistent w.r.t. the operator, then

$$\bar{e} = \mathbf{e}, \quad e \in \mathbf{e}. \quad (62)$$

Proof. By definition we have

$$e_\lambda \sim e_\mu, \quad \lambda, \mu \in \Lambda,$$

According to (ii) of Definition 3.13, we also know that if $a \notin \mathbf{e}$ then for any $e \in \mathbf{e}$, $a \not\sim e$. (62) follows. □

Using Proposition 3.14, we have the following result immediately.

Proposition 3.15: Consider a hyper group $\mathcal{G} = (G, *, e)$. Assume the equivalence \sim defined by (60) is consistent with $*$. Then

$$\bar{x} * \bar{y} := \overline{x * y}, \quad (63)$$




Fig. 3. Group Decomposition of a Hyper-Group \mathcal{G}

is properly defined. Moreover, $\bar{\mathcal{G}} = (\bar{G}, *)$ is a group, called the equivalence group of \mathcal{G} , denoted by

$$\bar{\mathcal{G}} := \mathcal{G} / \sim = (\bar{G}, *),$$

where $\bar{G} = \{\bar{x} \mid x \in G\}$.

Observing Propositions 3.11 and 3.15, a geometric picture for the structure of a hyper group, called the group decomposition of hyper group, is depicted by Figure 3.

Since the existence of the equivalence hyper group depends on the consistence of the equivalence w.r.t. the group operator, the consistence is the key issue for the structure analysis of hyper groups. It is worth showing the consistence of the hyper groups over MVMDs. To this end we need a preparation.

Lemma 3.16:

(i) Consider $x, y \in \mathbb{R}^\infty$, which are specified by $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, $m \vee n = t$, and $m \wedge n = s$. Assume there are $\mathbf{1}_p$ and $\mathbf{1}_q$ (or \mathcal{E}_p and \mathcal{E}_q) such that

$$x \otimes \mathbf{1}_p = y \otimes \mathbf{1}_q, \quad (\text{or } x \otimes \mathcal{E}_p = y \otimes \mathcal{E}_q), \quad (64)$$

then there exists a $z \in \mathbb{R}^s$ such that

$$x = z \otimes \mathbf{1}_{m/s}; \quad y = z \otimes \mathbf{1}_{n/s}. \quad (65)$$

(ii) Consider $A, B \in \mathcal{M}_\mu$, which are specified by $A \in \mathcal{M}_{m\mu_y \times m\mu_x}$, $B \in \mathcal{M}_{n\mu_y \times n\mu_x}$, $m \vee n = a$, and $m \wedge n = b$. Assume there are $H_r = I_r$ and $H_s = I_s$ (or $H_r = J_r$ and $H_s = J_s$) such that

$$A \otimes H_r = B \otimes H_s, \quad (66)$$

then there exists a $C \in \mathcal{M}_{b\mu_y \times b\mu_x}$ such that

$$A = C \otimes H_{m/b}; \quad B = C \otimes H_{n/b}. \quad (67)$$

- (iii) Consider $A, B \in \mathcal{M}$, which are specified by $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $m \wedge p = r$, and $n \wedge q = s$. Assume there are $\mathcal{E}_{a \times b}$ and $\mathcal{E}_{c \times d}$ such that

$$A \otimes \mathcal{E}_{a \times b} = B \otimes \mathcal{E}_{c \times d}, \quad (68)$$

then there exists a $C \in \mathcal{M}_{r \times s}$ such that

$$A = C \otimes \mathcal{E}_{m/r \times n/s}; \quad B = C \otimes \mathcal{E}_{p/r \times q/s}. \quad (69)$$

Proof. In fact, (i) and (ii) have been proved in [11]. The proof of (iii) is similar. We sketch it. First, note that $ma = pc$ and $nb = qd$. By deleting common factors we can assume, without loss of generality, that

$$a = \frac{m \vee p}{m}, \quad b = \frac{n \vee q}{n}, \quad c = \frac{m \vee p}{p}, \quad d = \frac{n \vee q}{q}.$$

Using the above a, b, c, d and comparing both sides of (68) carefully, the conclusion follows. The detailed arguments are similar to those for (i) and (ii) in [11]. \square

Example 3.17: Consider the hyper group $\mathcal{G} = (\mathcal{T}, \times, e)$, where the identity set is defined by (57). Then we have

- (i) The component groups are

$$\mathcal{T}_n = GL(n, \mathbb{R}), \quad n \in \mathbb{Z}_+, \quad (70)$$

where $GL(n, \mathbb{R})$ are the standard general linear groups, and

$$\mathcal{T} := \bigcup_{i=1}^{\infty} GL(i, \mathbb{R}).$$

- (ii) Using Lemma 3.16, it is easy to verify that the equivalence is consistent with the STP, hence the equivalence quotient group exists.

$$\bar{\mathcal{T}} := \bigcup_{n=1}^{\infty} \{\bar{A}_n \mid A_n \in \mathcal{M}_{n \times n} \text{ is invertible}\}.$$

If we define $\pi : \mathcal{T} \rightarrow \bar{\mathcal{T}}$ by

$$\pi(A) = \bar{A}, \quad A \in \mathcal{T},$$

then it is easy to verify that π is a hyper group homomorphism.

In the following we consider the additive hyper groups. Recall the additive hyper groups discussed in Proposition 3.8.

Example 3.18: In the following we assume the consistence of the equivalence w.r.t. addition in each case holds.

- (i) Consider $(\mathbb{R}^\infty, \bar{+})$ with $\bar{+}$ defined by (24). It is easy to verify that the component groups are $(\mathbb{R}^n, +)$, $n \in \mathbb{Z}_+$. Next, consider the equivalence hyper group. Let $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, and $t = \text{lcm}(m, n)$. Assume x and y are equivalent, denoted by $x \leftrightarrow y$. By definition, there exist 0_p and 0_q , such that

$$x \bar{+} 0_p = y \bar{+} 0_q. \quad (71)$$

Comparing the dimensions and deleting common factor $\mathbf{1}_s$, where $s = p \wedge q$, we have

$$x \otimes \mathbf{1}_{t/m} = y \otimes \mathbf{1}_{t/n}. \quad (72)$$

In fact, (72) is the definition of equivalence of two vectors in [10]. Note that the consistence of the equivalence w.r.t. $\bar{+}$ is assumed, (In fact, it has already proved in [10]), then the equivalence group exists, which is denoted by

$$\Omega = (\mathbb{R}^\infty / \leftrightarrow) = (\{\bar{x} \mid x \in \mathbb{R}^\infty\}, \bar{+}). \quad (73)$$

Similarly, we also have $x \rightleftharpoons y$, if

$$x \hat{+} 0_p = y \hat{+} 0_q, \quad (74)$$

which leads to

$$x \otimes \mathcal{E}_{t/m} = y \otimes \mathcal{E}_{t/n}. \quad (75)$$

And finally we have the equivalence group as

$$\Omega' = (\mathbb{R}^\infty / \rightleftharpoons) = (\{\bar{x} \mid x \in \mathbb{R}^\infty\}, \hat{+}). \quad (76)$$

- (ii) Consider $(\mathcal{M}_1, \bar{+})$. Let $A \in \mathcal{M}_{m \times m}$, $B \in \mathcal{M}_{n \times n}$, and $t = \text{lcm}(m, n)$. Then $A \sim B$ if there exist $0_{p \times p}$ and $0_{q \times q}$ such that

$$A \bar{+} 0_{p \times p} = B \bar{+} 0_{q \times q}, \quad (77)$$

which leads to

$$A \otimes I_{t/m} = B \otimes I_{t/n}. \quad (78)$$

And finally we have the equivalence group as

$$\Sigma = (\mathcal{M}_1 / \sim) = (\{\bar{A} \mid A \in \mathcal{M}_1\}, \bar{+}). \quad (79)$$

Similarly, consider $(\mathcal{M}_1, \hat{+})$. We have $A \approx B$ if

$$A \hat{+} 0_{p \times p} = B \hat{+} 0_{q \times q}, \quad (80)$$

which leads to

$$A \otimes J_{t/m} = B \otimes J_{t/n}. \quad (81)$$

And finally we have the equivalence group as

$$\Sigma' = (\mathcal{M}_1 / \approx) = (\{\bar{A} \mid A \in \mathcal{M}_1\}, \hat{+}). \quad (82)$$

- (iii) Consider $(\mathcal{M}_\mu, \hat{+})$. Let $A \in \mathcal{M}_{m\mu_y \times m\mu_x}$, $B \in \mathcal{M}_{n\mu_y \times n\mu_x}$, and $t = \text{lcm}(m, n)$. $A \approx B$ if

$$A \hat{+} 0_{p\mu_y \times p\mu_x} = B \hat{+} 0_{q\mu_y \times q\mu_x}, \quad (83)$$

which leads to

$$A \otimes J_{(t/m)\mu_y \times (t/m)\mu_x} = B \otimes J_{(t/n)\mu_y \times (t/n)\mu_x}. \quad (84)$$

Finally we have

$$\Sigma_\mu = (\mathcal{M}_\mu / \approx) = (\{\bar{A} \mid A \in \mathcal{M}_\mu\}, \hat{+}). \quad (85)$$

(iv) Consider $(\mathcal{M}, \hat{+})$. Let $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $A \sim B$ if there exist $0_{r \times s}$ and $0_{a \times b}$ such that

$$A \hat{+} 0_{r \times s} = B \hat{+} 0_{a \times b}. \quad (86)$$

Assume $\alpha = m \vee p$ and $\beta = n \vee q$, then (80) leads to

$$A \otimes \mathcal{E}_{(\alpha/m) \times (\beta/n)} = B \otimes \mathcal{E}_{(\alpha/p) \times (\beta/q)n}. \quad (87)$$

Finally we have

$$\Sigma = (\mathcal{M}/\approx) = (\{\bar{A} \mid A \in \mathcal{M}\}, \hat{+}). \quad (88)$$

In the above example we assume the consistence always exist. Using Lemma 3.16, this can be proved one by one:

Proposition 3.19:

- (i) Consider $(\mathbb{R}^\infty, \bar{+})$ (or $(\mathbb{R}^\infty, \hat{+})$), its equivalence is consistent with its operator $\bar{+}$ (or $\hat{+}$).
- (ii) Consider $(\mathcal{M}_1, \bar{+})$ (or $(\mathcal{M}_1, \hat{+})$), its equivalence is consistent with its operator $\bar{+}$ (or $\hat{+}$).
- (iii) Consider $(\mathcal{M}_\mu, \hat{+})$, its equivalence is consistent with its operator $\hat{+}$.
- (iv) Consider $(\mathcal{M}, \hat{+})$, its equivalence is consistent with its operator $\hat{+}$.

Proof. We prove (i) and (iv). The proofs for (ii) and (iii) are similar.

- (i) First, to use Lemma 3.16, we show that if $x, y \in \mathbb{R}^\infty$ and $x \leftrightarrow y$, then there exist $\mathbf{1}_p$ and $\mathbf{1}_q$ such that

$$x \otimes \mathbf{1}_p = y \otimes \mathbf{1}_q.$$

Assume $x \in \mathbb{R}^m$ and $y \in \mathbb{R}^n$. By definition, there exist 0_p and 0_q , such that

$$\begin{aligned} x \bar{+} 0_p &= (x \otimes \mathbf{1}_{(m \vee p)/m}) + (0_p \otimes \mathbf{1}_{(m \vee p)/p}) \\ &= (x \otimes \mathbf{1}_{(m \vee p)/m}) \\ &= y \bar{+} 0_q = (y \otimes \mathbf{1}_{(n \vee q)/n}). \end{aligned}$$

Set $t = m \vee n$. Then the above equality can be simplified to

$$x \otimes \mathbf{1}_{t/m} = y \otimes \mathbf{1}_{t/n}.$$

Next, assume $x_1 \sim x_2$ and $y_1 \sim y_2$, we have to show that

$$x_1 \bar{+} y_1 \sim x_2 \bar{+} y_2. \quad (89)$$

Suppose $x_1 \in \mathbb{R}^m$, $x_2 \in \mathbb{R}^n$, $y_1 \in \mathbb{R}^p$, $y_2 \in \mathbb{R}^q$, and

$$\begin{aligned} m \vee p &= \alpha, & n \vee q &= \beta, & m \wedge n &= r, \\ p \wedge q &= s, & r \vee s &= \xi. & \alpha &= \xi a, \\ \beta &= \xi b, \end{aligned}$$

According to Lemma 3.16 there exist $x_0 \in \mathbb{R}^r$ and $y_0 \in \mathbb{R}^s$ such that

$$\begin{aligned} x_1 &= x_0 \otimes \mathbf{1}_{m/r}, & x_2 &= x_0 \otimes \mathbf{1}_{n/r}, \\ y_1 &= y_0 \otimes \mathbf{1}_{p/s}, & y_2 &= y_0 \otimes \mathbf{1}_{q/s}. \end{aligned}$$

Then

$$\begin{aligned} x_1 \bar{+} y_1 &= x_0 \otimes \mathbf{1}_{m/r} \otimes \mathbf{1}_{\alpha/m} + y_0 \otimes \mathbf{1}_{p/s} \otimes \mathbf{1}_{\alpha/p} \\ &= x_0 \otimes \mathbf{1}_{\alpha/r} + y_0 \otimes \mathbf{1}_{\alpha/s} \\ &= (x_0 \otimes \mathbf{1}_{\xi/r} + y_0 \otimes \mathbf{1}_{\xi/s}) \otimes \mathbf{1}_a. \end{aligned}$$

Similarly,

$$\begin{aligned} x_2 \bar{+} y_2 &= x_0 \otimes \mathbf{1}_{\beta/r} + y_0 \otimes \mathbf{1}_{\beta/s} \\ &= (x_0 \otimes \mathbf{1}_{\xi/r} + y_0 \otimes \mathbf{1}_{\xi/s}) \otimes \mathbf{1}_b. \end{aligned}$$

We conclude that

$$x_1 \bar{+} y_1 \sim (x_0 \otimes \mathbf{1}_{\xi/r} + y_0 \otimes \mathbf{1}_{\xi/s}) \sim x_2 \bar{+} y_2.$$

Next, we check condition (ii) of Definition 3.13. Assume $x \in \mathbb{R}^m$ and $x \sim 0_a \in \mathbf{e}$, then there exist $\mathbf{1}_s$ and $\mathbf{1}_t$, such that

$$x \otimes \mathbf{1}_s = 0_a \otimes \mathbf{1}_t = 0_{at}.$$

It is easy to see that $x = 0_m \in \mathbf{e}$.

- (iv) To prove the consistence of the addition and the product with the equivalence we need to show that for $A_1 \sim A_2$ and $B_1 \sim B_2$ we have

$$A_1 \hat{+} B_1 \sim A_2 \hat{+} B_2. \quad (90)$$

Assume $A_1 \in \mathcal{M}_{m \times n}$, $A_2 \in \mathcal{M}_{p \times q}$, $B_1 \in \mathcal{M}_{\alpha \times \beta}$, and $B_2 \in \mathcal{M}_{\gamma \times \delta}$, and

$$\begin{aligned} m \vee \alpha &= a, & n \vee \beta &= b, & p \vee \gamma &= c, \\ q \vee \delta &= d, & m \wedge p &= s, & n \wedge q &= t, \\ \alpha \wedge \gamma &= \xi, & \beta \wedge \delta &= \eta, \\ s \vee \xi &= u, & t \vee \eta &= v. \end{aligned}$$

Using Lemma 3.16, we know that there exist $A_0 \in \mathcal{M}_{s \times t}$ and $B_0 \in \mathcal{M}_{\xi \times \eta}$ such that

$$\begin{aligned} A_1 &= A_0 \otimes \mathcal{E}_{m/s \times n/t}, & A_2 &= A_0 \otimes \mathcal{E}_{p/s \times q/t}; \\ B_1 &= B_0 \otimes \mathcal{E}_{\alpha/\xi \times \beta/\eta}, & B_2 &= B_0 \otimes \mathcal{E}_{\gamma/\xi \times \delta/\eta}. \end{aligned}$$

Now

$$\begin{aligned} A_1 \hat{+} B_1 &= A_1 \otimes \mathcal{E}_{a/m \times b/n} + B_1 \otimes \mathcal{E}_{a/\alpha \times b/\beta} \\ &= A_0 \otimes \mathcal{E}_{a/s \times b/t} + B_0 \otimes \mathcal{E}_{a/\xi \times b/\eta} \\ &= (A_0 \otimes \mathcal{E}_{u/s \times v/t} + B_0 \otimes \mathcal{E}_{u/\xi \times v/\eta}) \otimes \mathcal{E}_{a/u \times b/v}. \end{aligned}$$

$$\begin{aligned} A_2 \hat{+} B_2 &= A_2 \otimes \mathcal{E}_{c/p \times d/q} + B_2 \otimes \mathcal{E}_{c/\gamma \times d/\delta} \\ &= A_0 \otimes \mathcal{E}_{c/s \times d/t} + B_0 \otimes \mathcal{E}_{c/\xi \times d/\eta} \\ &= (A_0 \otimes \mathcal{E}_{u/s \times v/t} + B_0 \otimes \mathcal{E}_{u/\xi \times v/\eta}) \otimes \mathcal{E}_{c/u \times d/v}. \end{aligned}$$

It follows immediately that

$$A_1 \hat{+} B_1 \sim A_2 \hat{+} B_2.$$

The condition (ii) of Definition 3.13 is straightforward verifiable. \square

We conclude that for these additive hyper-vector spaces the equivalence vector spaces always exist.




Fig. 4. Sub Hyper Groups

E. Sub Hyper Group

Definition 3.20: Let $\mathcal{G} = (G, *, \mathbf{e})$ be a hyper group. $\mathcal{H} = (H, *, \mathbf{e}_0)$ is called a sub hyper group, if $H \subset G$ is a subset, $\mathbf{e}_0 \subset \mathbf{e}$ is a sub-lattice, and $(H, *, \mathbf{e}_0)$ is a hyper group. Denoted the sub hyper group by $\mathcal{H} <_h \mathcal{G}$. If $\mathcal{H} <_h \mathcal{G}$ and $gH_{e_g} = H_{e_g}g, \forall g \in G$ and $e_g \in \mathbf{e}_0$, then $(H, *, \mathbf{e}_0)$ is a normal sub hyper group, and denoted by $\mathcal{H} \triangleleft_h \mathcal{G}$.

The following result comes from the definition immediately.

Proposition 3.21: Let $\mathcal{G} = (G, *, \mathbf{e})$ be a hyper group.

(i) Assume $\mathbf{e}_0 \prec \mathbf{e}$ is a sub-lattice. Set

$$G_0 := \bigcup_{e \in \mathbf{e}_0} G_e, \quad (91)$$

then

$$\mathcal{G}_0 := (G_0, *, \mathbf{e}_0) <_h (G, *, \mathbf{e}). \quad (92)$$

(ii) Consider \mathcal{G}_0 as in (i). Assume $H_e < G_e, \forall e \in \mathbf{e}$, and

$$H^0 := \bigcup_{e \in \mathbf{e}_0} H_e. \quad (93)$$

Then

$$\mathcal{H}_0 := (H^0, *, \mathbf{e}_0) <_h \mathcal{G}_0. \quad (94)$$

(iii) Conversely, if $\mathbf{e}_0 \subset \mathbf{e}$ is a sublattice, $\emptyset \neq H_e \subset G_e, e \in \mathbf{e}_0$. H^0 is defined by (93), and $\mathcal{H}_0 := (H^0, *, \mathbf{e}_0) <_h \mathcal{G}$, then

$$\mathcal{H}_e := (H_e, *) < \mathcal{G}_e, \quad \forall e \in \mathbf{e}_0. \quad (95)$$

(iv) If $\mathcal{H}_0 := (H^0, *, \mathbf{e}_0) \triangleleft_h \mathcal{G}$, then

$$\mathcal{H}_e := (H_e, *) \triangleleft \mathcal{G}_e, \quad \forall e \in \mathbf{e}_0. \quad (96)$$

Fig. 4 depicts these relations.

Example 3.22:

(i) A lattice \mathcal{H} is constructed by

$$H = \{1, 2, 3, 4, 6, 9, 12, 18, 36\} \subset \mathbb{Z}_+$$

with sublattice structure of $(\mathbb{Z}_+, \sup = \vee, \inf = \wedge)$. Then its Hasse diagram is depicted by the left graph of Figure 2, where

$$\begin{aligned} e &= 36, & c &= 12, & f &= 18, \\ a &= 4, & h &= 6, & g &= 3, \\ i &= 2, & b &= 3, & d &= 1. \end{aligned}$$

Set

$$\mathcal{T}_H := \{\mathcal{T}_n \mid n \in H\},$$

and

$$\mathbf{e}_h = (\{I_n \mid n \in H\}, \vee, \wedge).$$

Then $\mathcal{H} := (\mathcal{T}_H, \times, \mathbf{e}_h) <_h \mathcal{G}$ is a sub-hyper group, w.r.t. MD-1 lattice. $\mathcal{G} = (\mathcal{T}, \times, \mathbf{e})$ is defined in Example 3.17.

(ii) Consider $M_1 = \bigcup_{n=1}^{\infty} \mathcal{M}_{n \times n}$, $\hat{\dagger}$ is defined by (26), and

$$\mathbf{e} = \{0_{n \times n} \in \mathcal{M}_{n \times n} \mid n \in \mathbb{Z}_+\},$$

with lattice structure (\mathbb{Z}_+, \prec) defined in (i) of Example 3.2. Then $\mathcal{M}_1 = (M_1, \hat{\dagger}, \mathbf{e})$ is a hyper group.

– Set

$$S_n := \{A \in \mathcal{M}_{n \times n} \mid A^T = A\},$$

and

$$S = \bigcup_{n=1}^{\infty} S_n.$$

Then $\mathcal{S} := (S, \hat{\dagger}, \mathbf{e})$ is a hyper group, and

$$\mathcal{S} \triangleleft_h \mathcal{M}_1.$$

– Set

$$K_n := \{A \in \mathcal{M}_{n \times n} \mid A^T = -A\},$$

and

$$K = \bigcup_{n=1}^{\infty} K_n.$$

Then $\mathcal{K} := (K, \hat{\dagger}, \mathbf{e})$ is a hyper group, and

$$\mathcal{K} \triangleleft_h \mathcal{M}_1.$$

– Set

$$D_n := \{A \in \mathcal{M}_{n \times n} \mid A \text{ is diagonal}\},$$

and

$$D = \bigcup_{n=1}^{\infty} D_n.$$

Then $\mathcal{D} := (D, \hat{\dagger}, \mathbf{e})$ is a hyper group, and

$$\mathcal{D} \triangleleft_h \mathcal{M}_1.$$

(iii) Consider $T_n, n \in \mathbb{Z}_+$ as the set of $n \times n$ non-singular matrices, and set

$$T := \bigcup_{i=1}^{\infty} T_n,$$

with \times defined by (16), and \mathbf{e} as in (57). Then $\mathcal{T} := (T, \times, \mathbf{e})$ is a hyper group.

– Set

$$O_n := \{A \in \mathcal{T}_n \mid A^T = A^{-1}\},$$

$$O = \bigcup_{n=1}^{\infty} O_n.$$

Then $\mathcal{O} := (O_n, \times, \mathbf{e})$ is a hyper group, and

$$\mathcal{O} <_h \mathcal{T}.$$

– Set

$$E_n := \{A \in \mathcal{M}_{n \times n} \mid \det(A) = 1\},$$

and

$$E = \bigcup_{n=1}^{\infty} E_n.$$

Then $\mathbf{E} = (E, \times, \mathbf{e})$ is a hyper group, and

$$\mathbf{E} <_h \mathcal{T}.$$

Proposition 3.23: Let $\mathcal{G} = (G, *, \mathbf{e})$ be a hyper group, and $H \subset G$. $\mathcal{H} = (H, *, \mathbf{e}_H)$ is a sub-hyper group, if and only if,

(i)

$$\mathbf{e}_H = \{e_x \mid x \in H\}, \quad (97)$$

and which is a sub-lattice;

(ii)

$$x * y^{-1} \in H, \quad \forall x, y \in H. \quad (98)$$

Proof. The necessity is obvious. To see the sufficiency, assume $x, y \in H_e$, then $x * y^{-1} \in H_e$. That is, $H_e < G_e$ is a sub-group. Hence, H is a sub-hyper group of G . \square

IV. PERMUTATION HYPER GROUP

A. Left Permutation Hyper Group

Permutation group is one of the most important finite groups, because any finite group can be isomorphic to a subgroup of certain permutation group [26]. In this section we construct a hyper group structure for permutations with different orders.

Let \mathbf{S}_n be the permutation group of order n with the group operator \circ as the compound of two permutations. Set

$$\mathbf{S} := \bigcup_{n=1}^{\infty} \mathbf{S}_n.$$

To build a cross-order product of permutations with different orders, we need to construct a cross-dimensional mapping.

Definition 4.1: Assume $m|n$, and $km = n$, then we define a mapping, called the left embedding, as $\varphi_n^m : \mathbf{S}_m \rightarrow \mathbf{S}_n$ as follows: Assume $\sigma \in \mathbf{S}_m$, then

$$\begin{aligned} \varphi_n^m(\sigma)((i-1)k + s) &:= (\sigma(i) - 1)k + s, \\ \sigma \in \mathbf{S}_m, s &\in [1, k], i \in [1, m]. \end{aligned} \quad (99)$$

Note that when $m = n$ φ_n^m is the identity mapping. When $m < n$, to understand this mapping, we recall permutation matrices. Recall the definition of permutation matrices in Example 2.2. A permutation matrix $P \in \mathcal{M}_{n \times n}$ is a non-singular logical matrix. A logical matrix $A \in \mathcal{L}_{n \times n}$ satisfies $\text{Col}(A) \subset$

D_n , hence it has the form that $A = [\delta_n^{i_1}, \delta_n^{i_2}, \dots, \delta_n^{i_n}]$ and can be briefly denoted by

$$A = \delta_n[i_1, i_2, \dots, i_n].$$

Recall that the set of $n \times n$ permutation matrices form a subgroup

$$\mathcal{P}_n < \mathcal{T}_n.$$

For each $\sigma \in \mathbf{S}_n$, its permutation matrix is expressed as

$$P_\sigma := \delta_n[\sigma(1), \sigma(2), \dots, \sigma(n)]. \quad (100)$$

According to the construction of a permutation matrix, we have the following result.

Proposition 4.2:

(i) Assume a permutation $\sigma \in \mathbf{S}_n$ and its permutation matrix is P_σ . Then

$$P_\sigma \delta_n^i = \delta_n^{\sigma(i)}, \quad i \in [1, n]. \quad (101)$$

(ii) Let $\sigma, \mu \in \mathbf{S}_n$. Then

$$M_{\sigma \circ \mu} = M_\sigma M_\mu. \quad (102)$$

(iii) Assume $m|n$, $km = n$, and $\sigma \in \mathbf{S}_m$. Then

$$P_{\varphi(\sigma)} = P_\sigma \otimes I_k. \quad (103)$$

Remark 4.3:

(i) Define $\pi : \mathbf{S}_n \rightarrow \mathcal{P}_n$ by $\pi : \sigma \mapsto P_\sigma$ as shown in (100). Then Proposition 4.2 shows that π is a group isomorphism.

(ii) Assume $m|n$, the φ_n^m is an one-to-one homomorphism from \mathbf{S}_m to \mathbf{S}_n . In other words, φ_n^m is an isomorphism from \mathbf{S}_m to its image $\varphi(\mathbf{S}_m) < \mathbf{S}_n$. It can also be considered as an “embedding”.

Now we are ready to define a cross-order product of permutations as follows.

Definition 4.4: Let $\sigma, \mu \in \mathbf{S}$, say, $\sigma \in \mathbf{S}_m, \mu \in \mathbf{S}_n$, and $t = \text{lcm}(m, n)$. Then the product, called the left product, of σ and μ , denoted by $\sigma \odot \mu$, is defined by

$$\sigma \odot \mu = \varphi_t^m(\sigma) \circ \varphi_t^n(\mu) \in \mathbf{S}_t. \quad (104)$$

The following result can be verified by the construction easily.

Proposition 4.5: Let $\sigma, \mu \in \mathbf{S}$ with their permutation matrices M_σ and M_μ respectively. Then

$$M_{\sigma \odot \mu} = M_\sigma \times M_\mu. \quad (105)$$

φ_n^m is called the left embedding because it is determined by this left STP of their permutation matrices.

Define the set of cross-order permutation matrices as

$$\mathcal{P} := \bigcup_{n=1}^{\infty} \mathcal{P}_n.$$

Then it is easy to verify the following.

Proposition 4.6:

- (i) $(\mathcal{P}, \times, \mathbf{e})$ is a hyper group, where \mathbf{e} is defined by (57).
(ii)

$$(\mathcal{P}, \times, \mathbf{e}) \triangleleft_h \mathcal{T}, \quad (106)$$

where $(\mathcal{T} = (T, \times, \mathbf{e}))$ is defined in Proposition 3.10.

Using Propositions 4.5 and 4.6, the following result is obvious.

Theorem 4.7: $\mathbf{S}^L := (\mathbf{S}, \odot, \mathbf{e})$ is a hyper group, where the identity set is

$$\mathbf{e} = \{e_n = Id_n \in \mathbf{S}_n \mid n \in \mathbb{Z}_+\}, \quad (107)$$

w.r.t. MD-1 lattice (\mathbb{Z}_+, \prec) . w.

Next, we consider some basic properties of hyper group \mathbf{S}^L .

Proposition 4.8: Set the alternative set of permutations by

$$\mathbf{A} := \bigcup_{n=1}^{\infty} \mathbf{A}_n. \quad (108)$$

- (i) \mathbf{A}^L is a sub-hyper group of \mathbf{S}^L . Precisely speaking,

$$(\mathbf{A}, \odot, \mathbf{e}) \triangleleft_h (\mathbf{S}, \odot, \mathbf{e}). \quad (109)$$

- (ii) \mathbf{A}^L is a normal sub-hyper group of \mathbf{S}^L . That is,

$$(\mathbf{A}, \odot, \mathbf{e}) \triangleleft_h (\mathbf{S}, \odot, \mathbf{e}). \quad (110)$$

Remark 4.9: The definition of normal hyper subgroup can not be defined in the same way as for normal subgroup. It is defined in each component groups. For instance, note that $\mathbf{A} \triangleleft_h \mathbf{S}$ does not mean

$$\sigma \circ \mathbf{A} = \mathbf{A} \circ \sigma, \forall \sigma \in \mathbf{S}.$$

It means

$$\sigma \circ \mathbf{A}_n = \mathbf{A}_n \circ \sigma, \forall \sigma \in \mathbf{S}_n, \forall n \in \mathbb{Z}_+.$$

In fact, \mathbf{A} and $g \circ \mathbf{A}$, where $g \in \mathbf{S} \setminus \mathbf{A}$, is not a partition of \mathbf{S} .

Finally, we consider the consistence of the equivalence w.r.t. \odot .

- (i) Assume $\sigma_1 \sim \sigma_2$ and $\mu_1 \sim \mu_2$, we verify if

$$\sigma_1 \odot \mu_1 \sim \sigma_2 \odot \mu_2. \quad (111)$$

Using the isomorphism between permutations and their structure matrices, (111) is equivalent to

$$M_{\sigma_1} M_{\mu_1} \sim M_{\sigma_2} M_{\mu_2}. \quad (112)$$

Assume $\sigma_1 \in \mathbf{S}_m, \sigma_2 \in \mathbf{S}_n, \mu_1 \in \mathbf{S}_p, \mu_2 \in \mathbf{S}_q, m \wedge n = \alpha, p \wedge q = \beta, m \vee p = \xi, n \vee q = \eta, \text{ and } \alpha \vee \beta = \lambda.$

According to Lemma 3.16, there exist $M_\alpha \in \mathcal{P}_\alpha$ and $M_\beta \in \mathcal{P}_\beta$ such that

$$\begin{aligned} M_{\sigma_1} &= M_\alpha \otimes I_{m/\alpha}, & M_{\sigma_2} &= M_\alpha \otimes I_{n/\alpha}, \\ M_{\mu_1} &= M_\beta \otimes I_{p/\beta}, & M_{\mu_2} &= M_\beta \otimes I_{q/\beta}. \end{aligned}$$

Then

$$\begin{aligned} M_{\sigma_1} M_{\mu_1} &= (M_\alpha \otimes I_{m/\alpha} \otimes I_{\xi/m}) (M_\beta \otimes I_{p/\beta} \otimes I_{\xi/p}) \\ &= (M_\alpha \otimes I_{xi/\alpha}) (M_\beta \otimes I_{\xi/\beta}) \\ &= [(M_\alpha \otimes I_{\lambda/\alpha}) (M_\beta \otimes I_{\lambda/\beta})] \otimes I_{\xi/\lambda}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} M_{\sigma_1} M_{\mu_1} &= [(M_\alpha \otimes I_{\lambda/\alpha}) (M_\beta \otimes I_{\lambda/\beta})] \otimes I_{\eta/\lambda}. \end{aligned}$$

(112) follows.

- (ii) Verify the (ii) of Definition 3.13. Assume $x \sim I_r$. Then there exist I_α and I_β such that

$$e \otimes I_\alpha = I_r \otimes I_\beta.$$

Then it is clear that $e = I_s \in \mathbf{e}$

□

Hence we have the following result.

Proposition 4.10: There exists the equivalence hyper group

$$\overline{\mathbf{S}}^L := \mathbf{S}^L / \sim. \quad (113)$$

Remark 4.11: We are surely interested in what are the elements in $\overline{\mathbf{S}}^L$. Let $\sigma \in \mathbf{S}$. σ is said to be reducible, if $\sigma \in \mathbf{S}_n$ and there are $m < n, m|n$ and a $\sigma_0 \in \mathbf{S}_m$ such that

$$\sigma = \varphi_n^m(\sigma_0).$$

Let $\sigma \in \mathbf{S}_m$ be irreducible and define

$$\overline{\sigma} = \{\varphi_{km}^m(\sigma) \mid k \in \mathbb{Z}_+\}.$$

Then it is easy to verify that

$$\overline{\mathbf{S}}^L := \{\overline{\sigma} \mid \sigma \in \mathbf{S} \text{ is irreducible}\}. \quad (114)$$

B. Right Permutation Hyper Group

Alternatively, we can also define a right permutation hyper group. We need to construct another cross-dimensional mapping.

Definition 4.12: Assume $m|n$, and $km = n$, then we define a mapping, called the right embedding, as $\psi_n^m : \mathbf{S}_m \rightarrow \mathbf{S}_n$ as follows: Assume $\sigma \in \mathbf{S}_m$, then

$$\begin{aligned} \psi_n^m(\sigma)((s-1)m+i) &:= (k-s)m + \sigma(i), \\ \sigma \in \mathbf{S}_m, s &\in [1, k], i \in [1, m]. \end{aligned} \quad (115)$$

Similarly to left embedding case, we have the following result.

Proposition 4.13: Assume $m|n$, $km = n$, and $\sigma \in \mathbf{S}_m$. Then

$$P_{\psi(\sigma)} = I_k \otimes P_\sigma. \quad (116)$$

Remark 4.14: Assume $m|n$, similarly to φ_n^m , the ψ_n^m is an one-to-one homomorphism from \mathbf{S}_m to \mathbf{S}_n . In other words, ψ_n^m is an isomorphism from \mathbf{S}_m to its image $\varphi(\mathbf{S}_m) < \mathbf{S}_n$. It can also be considered as an ‘‘embedding’’.

Next, we define another cross-order product of permutations as follows.

Definition 4.15: Let $\sigma, \mu \in \mathbf{S}$, say, $\sigma \in \mathbf{S}_m, \mu \in \mathbf{S}_n$, and $t = \text{lcm}(m, n)$. Then the right product of σ and μ , denoted by $\sigma \odot_r \mu$, is defined by

$$\sigma \odot_r \mu = \psi_t^m(\sigma) \circ \psi_t^n(\mu) \in \mathbf{S}_t. \quad (117)$$

The following result is obvious.

Proposition 4.16: Let $\sigma, \mu \in \mathbf{S}$ with their permutation matrices M_σ and M_μ respectively. Then

$$M_{\sigma \odot_r \mu} = M_\sigma \times M_\mu. \quad (118)$$

ψ_n^m is called the right embedding because it is determined by the right STP of their permutation matrices.

Using Propositions 4.16 and 4.6, the following result is obvious.

Theorem 4.17: $\mathbf{S}^R = (\mathbf{S}, \odot_r, \mathbf{e})$ is a hyper group, where the identity set is

$$\mathbf{e} = \{e_n = Id_n \in \mathbf{S}_n \mid n \in \mathbb{Z}_+\}, \quad (119)$$

w.r.t. MD-1 lattice (\mathbb{Z}_+, \prec) .

Correspondingly, the right hyper group \mathbf{S}^R has the following properties.

Proposition 4.18:

(i) \mathbf{A}^R is a sub-hyper group of \mathbf{S}^R . Precisely speaking,

$$(\mathbf{A}, \odot_r, \mathbf{e}) <_h (\mathbf{S}, \odot_r, \mathbf{e}). \quad (120)$$

(ii) \mathbf{A}^R is a normal sub-hyper group of \mathbf{S}^R . That is,

$$(\mathbf{A}, \odot_r, \mathbf{e}) \triangleleft_h (\mathbf{S}, \odot_r, \mathbf{e}). \quad (121)$$

σ and μ are said to be right-equivalent, denoted by $\sigma \sim_r \mu$, if there exist I_r and I_s such that

$$I_r \otimes M_\sigma = I_s \otimes M_\mu. \quad (122)$$

Then we can also prove that the equivalence is consistent with respect to the \odot_r , hence we have

Proposition 4.19: There exists the equivalence hyper group

$$\overline{\mathbf{S}}^R := \mathbf{S}^R / \sim_r. \quad (123)$$

Denote by

$$\tilde{\sigma} = \{\mu \mid \mu \sim_r \sigma\},$$

then we have

Remark 4.20:

$$\overline{\mathbf{S}}^R := \{\tilde{\sigma} \mid \sigma \in \mathbf{S} \text{ is irreducible}\}. \quad (124)$$

V. HOMOMORPHISM

A. Hyper Group Homomorphism/Isomorphism

Definition 5.1: Let $\mathcal{G}_i = (G^i, \times^i, \mathbf{e}^i)$, $i = 1, 2$ be two hyper groups. $\pi : G^1 \rightarrow G^2$ is called a hyper group homomorphism, and \mathcal{G}_1 and \mathcal{G}_2 are said to be homomorphic, denoted by $\mathcal{G}_1 \simeq_h \mathcal{G}_2$, if the followings are satisfied:

(i)

$$\pi(x \times^1 y) = \pi(x) \times^2 \pi(y), \quad x, y \in G_1. \quad (125)$$

(ii)

$$\pi(\mathbf{e}^1) \in \mathbf{e}^2, \quad \mathbf{e}^1 \in \mathbf{e}^1, \quad (126)$$

and $\pi : \mathbf{e}^1 \rightarrow \mathbf{e}^2$ is a lattice order-preserving mapping.

(iii) If a hyper group homomorphism $\pi : G_1 \rightarrow G_2$ is bijective, and $\pi^{-1} : G_2 \rightarrow G_1$ is also a homomorphism, then π is called a hyper group isomorphism, and \mathcal{G}_1 and \mathcal{G}_2 are said to be isomorphic, denoted by $\mathcal{G}_1 \cong_h \mathcal{G}_2$.

Example 5.2: Recall \mathcal{T}_H defined in Example 3.22.

- (i) Let $Id : \mathcal{T}_H \hookrightarrow \mathcal{T}$ be the including mapping. Then Id is a hyper group homomorphism, and $\mathcal{T}_H \simeq_h \mathcal{T}$.
- (ii) Let $\pi_k : \mathcal{T}_H \rightarrow \mathcal{T}$ be defined by $A \mapsto A \otimes I_k$. Then $(\pi_k(\mathcal{T}_H), \times) \simeq_h \mathcal{T}$.
- (iii) Since Id is one-to-one. Then it is easy to verify that

$$\mathcal{T}_H \cong_h (\pi_k(\mathcal{T}_H), \times).$$

Example 5.3: Consider $(\mathcal{M}_\mu, \bar{\dagger})$ (or $(\mathcal{M}_\mu, \hat{\dagger})$). Let $\pi : A \mapsto A^T$. Then π is a hyper group isomorphism. That is,

$$\begin{aligned} (\mathcal{M}_\mu, \bar{\dagger}) &\cong_h (\mathcal{M}_{1/\mu}, \bar{\dagger}); \\ \text{(or } (\mathcal{M}_\mu, \hat{\dagger}) &\cong_h (\mathcal{M}_{1/\mu}, \hat{\dagger}). \end{aligned} \quad (127)$$

Example 5.4: Consider (\mathbf{S}^L, \odot) and (\mathcal{P}, \times) (or (\mathbf{S}^R, \odot_r) and (\mathcal{P}, \times)). Then $\pi : \sigma \mapsto M_\sigma$ is a hyper group isomorphism. That is,

$$\begin{aligned} (\mathbf{S}^L, \odot) &\cong_h (\mathcal{P}, \times); \\ \text{(or } (\mathbf{S}^R, \odot_r) &\cong_h (\mathcal{P}, \times). \end{aligned} \quad (128)$$

B. Quotient Hyper Group Homomorphism/Isomorphism

Assume $\varphi : H \rightarrow G$ is a hyper group homomorphism. Set

$$\begin{aligned} Im(\varphi) &:= \{g \in G \mid \exists h \in H \text{ such that } g = \varphi(h)\}, \\ Ker(\varphi) &:= \{h \in H \mid \varphi(h) \in \mathbf{e}_G\}. \end{aligned} \quad (129)$$

Then we have the following result.

Proposition 5.5:

- (i) $Im(\varphi) <_h G$.

(ii) $\text{Ker}(\varphi) \triangleleft_h H$.

Proof.

- (i) Since $\varphi : \mathbf{e}_H \rightarrow \mathbf{e}_G$ is a lattice homomorphism, it is easy to see that $\varphi(\mathbf{e}_H)$ is a sub-lattice of \mathbf{e}_G . Let $x, y \in \text{Im}(\varphi)$. Then there exist $a, b \in H$ such that $\varphi(a) = x$, $\varphi(b) = y$. Since H is a hyper group, $y^{-1} \in H$. Then $\varphi(ab^{-1}) = xy^{-1} \in \text{Im}(\varphi)$. We conclude that $\text{Im}(\varphi) <_h G$.
- (ii) Since $\mathbf{e}_H \subset K := \ker(\varphi)$, $\mathbf{e}_K = \mathbf{e}_H$. It is obvious that $K <_h H$. To see K is a normal hyper-sub-group, we need to show that for any $a \in H$ $aKa^{-1} = K$. This is true because

$$\varphi(aka^{-1}) = \varphi(a)e(\varphi(a))^{-1} = e\varphi(a)(\varphi(a))^{-1} \in \mathbf{e}_G.$$

□

Proposition 5.6: Let $\varphi : H \rightarrow G$ be a hyper group homomorphism, and for each $e \in \mathbf{e}_H$, $\varphi_e : H_e \rightarrow G$ be the restriction of φ on H_e . Then $\varphi_e : H_e \rightarrow G_{\varphi_e}$ is a group homomorphism. Moreover, for each $e \in \mathbf{e}_H$

- (i) $\text{Im}(\varphi_e) < G_e$;
(ii) $\text{Ker}(\varphi_e) \triangleleft H$.

Proof. We claim that $\varphi_e(H_e) \subset G_{\varphi_e}$. If this claim is true, all the remaining statements are obvious. Let $h \in H_e$, then

$$\varphi(hh^{-1}) = \varphi(h)\varphi(h^{-1}) = \varphi(e) \in \mathbf{e}_G.$$

Hence $\varphi(h) \in G_{\varphi_e}$, which proves the claim. □

The three Isomorphism Theorems for groups can be modified to suit the hyper groups.

Theorem 5.7: (First Isomorphism Theorem)

- (i) Assume $\varphi : H \rightarrow G$ is a hyper group homomorphism. For each $e \in \mathbf{e}_H$, set $K_e = \text{Ker}(\varphi_e)$ and $\text{Im}_e = \text{Im}(\varphi_e)$. Define a mapping $\theta_e : H_e/K_e \rightarrow \text{Im}_e$ as $h_e K_e \mapsto \varphi_e(h_e)$, $h_e \in H_e$. Then θ_e is an isomorphism. That is.

$$H_e/K_e \cong \text{Im}_e, \quad e \in \mathbf{e}_H. \quad (130)$$

- (ii) Let $H \triangleleft_h G$ be a normal hyper-sub-group. For each $e \in \mathbf{e}_H$ define $\pi_e : G_{\varphi(e)} \rightarrow G_{\varphi(e)}/H_e$ by $g_e \mapsto g_e H_e$. Then π_e is a surjective homomorphism with $\text{Ker}(\pi_e) = H_e$.

Theorem 5.8: (Second Isomorphism Theorem) Assume H, N, G are hyper groups.

- (i) If $H <_h G$ and $N \triangleleft_h G$, then for each $e \in \mathbf{e}_H \cap \mathbf{e}_N$

$$N_e \cap H_e \triangleleft_h H_e. \quad (131)$$

- (ii) Set $NH = \{nh \mid n \in N, h \in H\}$, where N, H are as in (i). Then for each $e \in \mathbf{e}_H \cap \mathbf{e}_N$

$$N_e H_e <_h G_e. \quad (132)$$

- (iii) Define $\pi_e : H_e/(N_e \cap H_e) \rightarrow N_e H_e/N_e$ by

$$h_e(N_e \cap H_e) \mapsto h_e N_e, \quad h_e \in H_e,$$

where $e = e_h e_h$. Then π_e is an isomorphism. That is

$$H_e/(H_e \cap N_e) \cong N_e H_e/N_e. \quad (133)$$

Theorem 5.9: (Third Isomorphism Theorem) Let M, N, G be hyper groups, $M \triangleleft_h G$, $N \triangleleft_h G$, and $N <_h M$. Then for each $e \in \mathbf{e}_N$

- (i)

$$M_e/N_e \triangleleft_h G_e/N_e. \quad (134)$$

- (ii)

$$(G_e/N_e)/(M_e/N_e) \cong G_e/M_e. \quad (135)$$

Example 5.10: Consider $\Lambda = \{(2, 2), (2, 4), (4, 2), (4, 4)\}$ as a sub-lattice of $\mathbb{Z}_+ \times \mathbb{Z}_+$, $H = \bigcup_{\lambda \in \Lambda} \mathcal{M}_\lambda$, and

$$\mathcal{H} := (H, \hat{\dagger}) <_h \mathcal{G} = (\mathcal{M}, \hat{\dagger}).$$

Define a hyper group homomorphism: $\varphi : \mathcal{H} \rightarrow \mathcal{G}$ as

$$\varphi(A) = A \hat{\dagger} A^T.$$

A straightforward computation shows that

$$K := \ker(\varphi) = K_{4 \times 4} \cup K_{4 \times 2} \cup K_{2 \times 4} \cup K_{2 \times 2},$$

where

$$\begin{aligned} K_{4 \times 4} &= \{A \in \mathcal{M}_{4 \times 4} \mid A^T = A\}; \\ K_{4 \times 2} &= \{A \otimes \mathbf{1}_2 \mid A \in \mathcal{M}_{2 \times 2}, \text{ and } A^T = A\}; \\ K_{2 \times 4} &= \{A \otimes \mathbf{1}_2^T \mid A \in \mathcal{M}_{2 \times 2}, \text{ and } A^T = A\}; \\ K_{2 \times 2} &= \{A \in \mathcal{M}_{2 \times 2} \mid A^T = A\}. \end{aligned}$$

$$\text{Im} := \text{Im}(\varphi) = \text{Im}_{4 \times 4} \cup \text{Im}(2 \times 2),$$

where

$$\begin{aligned} \text{Im}_{4 \times 4} &= \{A \in \mathcal{M}_{4 \times 4} \mid A^T = -A\}; \\ \text{Im}_{2 \times 2} &= \{A \in \mathcal{M}_{2 \times 2} \mid A^T = -A\}. \end{aligned}$$

Since

$$K_{\alpha \times \beta} \hat{\dagger} K_{\gamma \times \delta} = \begin{cases} K_{4 \times 4}, & \max(\alpha, \beta, \gamma, \delta) = 4, \\ K_{2 \times 2}, & \max(\alpha, \beta, \gamma, \delta) = 2, \end{cases}$$

$K \triangleleft_h \mathcal{H}$ is a hyper-sub-group. It is normal because \mathcal{H} is abelian. Since

$$\text{Im}_{\alpha \times \alpha} \hat{\dagger} \text{Im}_{\beta \times \beta} = \begin{cases} \text{Im}_{4 \times 4}, & \max(\alpha, \beta) = 4, \\ \text{Im}_{2 \times 2}, & \max(\alpha, \beta) = 2, \end{cases}$$

$\text{Im} \triangleleft_h \mathcal{G}$ is a hyper-sub-group. Note that the lattice order-preserving mapping is:

$$\pi(e_\lambda) = \begin{cases} (4, 4), & \lambda \in \{(2, 4), (4, 2), (4, 4)\}, \\ (2, 2), & \lambda = (2, 2). \end{cases}$$

Next, we verify First Isomorphism Theorem. Consider each case when $\lambda \in \mathfrak{e}_H$.

(i) $\lambda = (4, 4)$.

$$K_\lambda = \{A \in \mathcal{M}_{4 \times 4} \mid A^T = A\}.$$

$$Im_\lambda = \{A \in \mathcal{M}_{4 \times 4} \mid A^T = -A\}.$$

Let $A \in H_\lambda = \mathcal{M}_{4 \times 4}$. Since

$$A = \frac{1}{2}(A + A^T) + \frac{1}{2}(A - A^T),$$

$$AK_\lambda = \frac{1}{2}(A + A^T) + \frac{1}{2}(A - A^T) + K_\lambda = \frac{1}{2}(A - A^T) + K_\lambda.$$

Define $\pi : H_\lambda/K_\lambda \rightarrow Im_\lambda$ as

$$\pi(AK_\lambda) := \frac{1}{2}(A - A^T).$$

Then π is an isomorphism. That is,

$$H_\lambda/K_\lambda \cong Im_\lambda.$$

(ii) $\lambda = (4, 2)$. A straightforward calculation shows that

$$K_\lambda = \left\{ \begin{bmatrix} a & a & b & b \\ b & b & c & c \end{bmatrix} \mid a, b, c \in \mathbb{R} \right\}.$$

Set

$$\begin{aligned} A &= \begin{bmatrix} a_{1,1} & a_{1,2} & a_{1,3} & a_{1,4} \\ a_{2,1} & a_{2,2} & a_{2,3} & a_{2,4} \end{bmatrix} \\ &= \begin{bmatrix} a_{1,1} & a_{1,1} & a_{1,3} & a_{1,3} \\ a_{1,3} & a_{1,3} & a_{2,3} & a_{2,3} \end{bmatrix} \\ &+ \begin{bmatrix} 0 & a_{1,2} - a_{1,1} & 0 & a_{1,4} - a_{1,3} \\ a_{2,1} - a_{1,3} & a_{2,2} - a_{1,3} & 0 & a_{2,4} - a_{1,3} \end{bmatrix} \\ &:= \begin{bmatrix} a_{1,1} & a_{1,1} & a_{1,3} & a_{1,3} \\ a_{1,3} & a_{1,3} & a_{2,3} & a_{2,3} \end{bmatrix} + \begin{bmatrix} 0 & d & 0 & e \\ f & g & 0 & h \end{bmatrix} \end{aligned}$$

Hence

$$AK_\lambda = \begin{bmatrix} 0 & d & 0 & e \\ f & g & 0 & h \end{bmatrix} + K_\lambda.$$

The Im_λ is calculated as

$$Im_\lambda = \left\{ \begin{bmatrix} 0 & d & -f & e-f \\ -d & 0 & -g & e-g \\ f & g & 0 & h \\ f-e & g-e & -h & 0 \end{bmatrix} \mid d, e, f, g, h \in \mathbb{R} \right\} \triangleleft \mathcal{K}_4.$$

That is, Im_λ is a (normal) sub-group of \mathcal{K}_4 .

Define $\pi : H_\lambda/K_\lambda \rightarrow Im_\lambda$ as

$$\begin{bmatrix} 0 & d & 0 & e \\ f & g & 0 & h \end{bmatrix} + K_\lambda \mapsto \begin{bmatrix} 0 & d & -f & e-f \\ -d & 0 & -g & e-g \\ f & g & 0 & h \\ f-e & g-e & -h & 0 \end{bmatrix}$$

Then it is easy to verify that π is an isomorphism. That is,

$$H_\lambda/K_\lambda \cong Im_\lambda.$$

(iii) $\lambda = (2, 4)$. Similarly to (ii), we have

$$K_\lambda = \left\{ \begin{bmatrix} a & b \\ a & b \\ b & c \\ b & c \end{bmatrix} \mid a, b, c \in \mathbb{R} \right\}.$$

$$AK_\lambda = \begin{bmatrix} 0 & f \\ d & g \\ 0 & 0 \\ e & h \end{bmatrix} + K_\lambda.$$

$$Im_\lambda = \left\{ \begin{bmatrix} 0 & -d & f & f-e \\ d & 0 & g & g-e \\ -f & -g & 0 & -h \\ e-f & e-g & h & 0 \end{bmatrix} \mid d, e, f, g, h \in \mathbb{R} \right\} \triangleleft \mathcal{K}_4.$$

That is, Im_λ is a sub-group of \mathcal{K}_4 .

Define $\pi : H_\lambda/K_\lambda \rightarrow Im_\lambda$ as

$$\begin{bmatrix} 0 & f \\ d & g \\ 0 & 0 \\ e & h \end{bmatrix} + K_\lambda \mapsto \begin{bmatrix} 0 & -d & f & f-e \\ d & 0 & g & g-e \\ -f & -g & 0 & -h \\ e-f & e-g & h & 0 \end{bmatrix}$$

Then π is an isomorphism. That is,

$$H_\lambda/K_\lambda \cong Im_\lambda.$$

(iv) $\lambda = (2, 2)$.

$$K_\lambda = \{A \in \mathcal{M}_{2 \times 2} \mid A^T = A\}.$$

$$Im_\lambda = \{A \in \mathcal{M}_{2 \times 2} \mid A^T = -A\}.$$

Let $A \in H_\lambda = \mathcal{M}_{2 \times 2}$. Since

$$A = \frac{1}{2}(A + A^T) + \frac{1}{2}(A - A^T),$$

$$AK_\lambda = \frac{1}{2}(A + A^T) + \frac{1}{2}(A - A^T) + K_\lambda = \frac{1}{2}(A - A^T) + K_\lambda.$$

Define $\pi : H_\lambda/K_\lambda \rightarrow Im_\lambda$ as

$$\pi(AK_\lambda) := \frac{1}{2}(A - A^T).$$

Then π is an isomorphism. That is,

$$H_\lambda/K_\lambda \cong Im_\lambda.$$

C. Symmetric and Skew symmetric Matrices

Definition 5.11: Consider the hyper group $\mathcal{G}_M := (\mathcal{M}, \hat{\cdot})$.

- (i) A mapping $\pi_s : \mathcal{G}_M \rightarrow \mathcal{G}_M$, called the symmetrizing mapping, is defined by

$$\pi_s : A \mapsto \frac{1}{2} (A \hat{+} A^T), \quad A \in \mathcal{M}. \quad (136)$$

- (ii) A mapping $\pi_a : \mathcal{G}_M \rightarrow \mathcal{G}_M$, called the alternating mapping, is defined by

$$\pi_a : A \mapsto \frac{1}{2} (A \hat{-} A^T), \quad A \in \mathcal{M}. \quad (137)$$

The following result makes the symmetry and skew-symmetry of matrices meaningful for non-square matrices.

Theorem 5.12: Consider the hyper group $\mathcal{G}_M := (\mathcal{M}, \hat{\cdot})$.

- (i) Both π_s and π_a are hyper group homomorphisms (precisely speaking, endomorphisms). According to Proposition 5.5, the kernels of π_s and π_a are hyper groups.
- (ii) The kernel of π_s , called the set of skew-symmetric matrices, denoted by \mathcal{K} , is π_a invariant in the sense

$$\pi_a(A) = A \otimes \mathcal{E}_{\alpha/m \times \alpha/n}, \quad A \in \mathcal{K}. \quad (138)$$

- (iii) $\mathcal{K}_{n \times n} = \mathcal{K}_n$. That is, when $m = n$ the component group $\mathcal{K}_{m \times n}$ coincides the classical set of skew-symmetric matrices.
- (iv) The kernel of π_a , called the set of symmetric matrices, denoted by \mathcal{S} , is π_s invariant in the sense

$$\pi_s(A) = A \otimes \mathcal{E}_{\alpha/m \times \alpha/n}, \quad A \in \mathcal{S}. \quad (139)$$

- (v) $\mathcal{S}_{n \times n} = \mathcal{S}_n$. That is, when $m = n$ the component group $\mathcal{S}_{m \times n}$ coincides the classical set of symmetric matrices.

Proof.

- (i) We show that π_s is a hyper group homomorphism. Let $A \in \mathcal{M}_{m \times n}$, and $t = \text{lcm}(m, n)$. Note that

$$A \hat{+} A^T = A \otimes \mathcal{E}_{t/m \times t/n} + A^T \otimes \mathcal{E}_{t/n \times t/m} \in \mathcal{M}_{t \times t}.$$

It is clear that $\pi_s : 0_{m \times n} \rightarrow 0_{m \vee n}$, which is order-keeping.

Next, we assume $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $s = m \vee p$, $t = n \vee q$, $\xi = s \vee t$, $\alpha = m \vee n$, and $\beta = p \vee q$. Then

$$\begin{aligned} \pi_s(A \hat{+} B) &= \pi_s [(A \otimes \mathcal{E}_{s/m \times t/n}) + (B \otimes \mathcal{E}_{s/p \times t/q})] \\ &= \frac{1}{2} [(A \otimes \mathcal{E}_{s/m \times t/n}) + (B \otimes \mathcal{E}_{s/p \times t/q})] \\ &\hat{+} \frac{1}{2} [(A^T \otimes \mathcal{E}_{t/n \times s/m}) + (B^T \otimes \mathcal{E}_{t/q \times s/p})] \\ &= \frac{1}{2} [(A \otimes \mathcal{E}_{s/m \times t/n}) + (B \otimes \mathcal{E}_{s/p \times t/q})] \otimes \mathcal{E}_{\xi/s \times \xi/t} \\ &\quad + \frac{1}{2} [(A^T \otimes \mathcal{E}_{t/n \times s/m}) + (B^T \otimes \mathcal{E}_{t/q \times s/p})] \otimes \mathcal{E}_{\xi/t \times \xi/s} \\ &= \frac{1}{2} [(A \otimes \mathcal{E}_{\xi/m \times \xi/n}) + (A^T \otimes \mathcal{E}_{\xi/n \times \xi/m})] \\ &\quad + \frac{1}{2} [(B \otimes \mathcal{E}_{\xi/p \times \xi/q}) + (B^T \otimes \mathcal{E}_{\xi/q \times \xi/p})] \\ &= \frac{1}{2} [(A \otimes \mathcal{E}_{\alpha/m \times \alpha/n}) + (A^T \otimes \mathcal{E}_{\alpha/n \times \alpha/m})] \\ &\quad \otimes \mathcal{E}_{\xi/\alpha \times \xi/\alpha} + \frac{1}{2} [(B \otimes \mathcal{E}_{\beta/p \times \beta/q}) \\ &\quad + (B^T \otimes \mathcal{E}_{\beta/q \times \beta/p})] \otimes \mathcal{E}_{\xi/\beta \times \xi/\beta} \\ &= \frac{1}{2} [(A \otimes \mathcal{E}_{\alpha/m \times \alpha/n}) + (A^T \otimes \mathcal{E}_{\alpha/n \times \alpha/m})] \\ &\quad \hat{+} \frac{1}{2} [(B \otimes \mathcal{E}_{\beta/p \times \beta/q}) + (B^T \otimes \mathcal{E}_{\beta/q \times \beta/p})] \\ &= \pi_s(A) \hat{+} \pi_s(B). \end{aligned}$$

We conclude that π_s is a hyper group homomorphism. Similarly, we can prove π_a is also a hyper group homomorphism.

- (ii) Assume $A \in \mathcal{K} = \text{Ker}(\pi_s)$. Then

$$A \otimes \mathcal{E}_{\alpha/m \times \alpha/n} = -A^T \otimes \mathcal{E}_{\alpha/n \times \alpha/m}.$$

(138) follows immediately.

All the rest parts can be proved similarly. \square

Corollary 5.13: Let $A \in \mathcal{M}_{m \times n}$ and $t = \text{lcm}(m, n)$. Then there exist unique symmetric matrix $A_s \in \mathcal{S}_t$ and a skew-symmetric matrix $A_a \in \mathcal{K}_t$ such that

$$A \otimes E_{t/m \times t/n} = A_s + A_a, \quad (140)$$

where

$$A_s = \frac{1}{2} (A \hat{+} A^T), \quad A_a = \frac{1}{2} (A \hat{-} A^T). \quad (141)$$

D. Squaring Matrix

Definition 5.14: Let $A \in \mathcal{M}_{m \times n} \subset \mathcal{M}$ and $t = \text{lcm}(m, n)$. A mapping $\square : \mathcal{M} \rightarrow \mathcal{M}_1$, called a squaring mapping, is defined by

$$\square(A) := A \hat{\times} I_t \in \mathcal{M}_{t \times t}. \quad (142)$$

Remark 5.15:

- (i) It is obvious that if A is a square matrix, then

$$\square(A) = A, \quad A \in \mathcal{M}_1. \quad (143)$$

- (ii) Let $A \in \mathcal{M}_{m \times n}$ and $t = \text{lcm}(m, n)$. Then

$$\square(A) = A \otimes \mathcal{E}_{t/m \times t/n}. \quad (144)$$

Using squaring mapping, (138)-(140) can be rewritten as (145)-(147) respectively as

$$\pi_a(A) = \square(A), \quad A \in \mathcal{K}. \quad (145)$$

$$\pi_s(A) = \square(A), \quad A \in \mathcal{S}. \quad (146)$$

$$\square(A) = A_s + A_a. \quad (147)$$

Hence, we have the following result.

Proposition 5.16: A non-square matrix A is symmetric (skew-symmetric), if and only if, its squaring matrix $\square(A)$ is symmetric (skew-symmetric).

Proposition 5.17: $\square : (\mathcal{M}, \hat{+}) \rightarrow (\mathcal{M}_1, \hat{+})$ is a hyper group homomorphism. Hence,

$$\square(A\hat{+}B) = \square(A)\hat{+}\square(B), \quad A, B \in \mathcal{M}. \quad (148)$$

The proof is similar to the proof of Theorem 5.12.

Definition 5.18: Let $A \in \mathcal{M}_{m \times n}$ and $t = \text{lcm}(m, n)$.

(i)

$$p(\lambda) := \det(\lambda I_t \hat{-} A) \quad (149)$$

is called the s-characteristic function of A .

(ii) $\sigma \in \mathbb{C}$ is called a s-eigenvalue of A , if $p(\sigma) = 0$.

(iii) $X \in \mathbb{C}^t$ is called an s-eigenvector of A w.r.t. σ , if

$$\square(A)X = \sigma X. \quad (150)$$

We have the s-Cayley-Hamilton theorem as follows.

Theorem 5.19: Let $p(\lambda)$ be the s-characteristic function of A , then

$$p(\square(A)) = 0. \quad (151)$$

Remark 5.20: The squaring mapping poses a non-square matrix all the properties of square matrix. For instance, we can define

$$\text{Tr}^\square(A) = \text{Tr}(\square(A)), \quad A \in \mathcal{M}. \quad (152)$$

Then it is clear that if $\text{Tr}^\square(A) = 0$, then $\det(\exp(\square(A))) = 1$.

E. Homomorphisms Among Component Groups

Proposition 5.21: Let $(G, *)$ be a hyper group with identity set $\mathbf{e} = \{e_\lambda \mid \lambda \in \Lambda\}$, where Λ is the lattice of identity set. Assume $H < G_\lambda$ (or $H \triangleleft G_\lambda$) and $e_\mu \in \mathbf{e}$, then

$$H * e_\mu < G_{\lambda \vee \mu}, (H * e_\mu \triangleleft G_{\lambda \vee \mu}). \quad (153)$$

Example 5.22:

(i) Since $\mathbf{A}_n \triangleleft \mathbf{S}_n$,

$$\mathbf{A} \odot Id_s \triangleleft \mathbf{S}_{n \vee s}.$$

(ii) Since $\mathcal{P}_n < \mathcal{T}_n$.

$$\mathcal{P}_n \times I_s < \mathcal{T}_{n \vee s};$$

$$\mathcal{P}_n \times I_t < \mathcal{T}_{n \vee t}.$$

Proposition 5.23: Let $(G, *)$ be a hyper group with identity set $\mathbf{e} = \{e_\lambda \mid \lambda \in \Lambda\}$, where Λ is the lattice of identity set. Assume $H < G_\lambda$, $N \triangleleft G_\mu$, then

$$HN < G_{\lambda \vee \mu}. \quad (154)$$

VI. RING AND HYPER RING OF MATRICES

A. Matrix Ring

It has been demonstrated in Example 2.5 that $(\mathcal{M}_{n \times n}, +, \times)$ is a unitary ring, and it has some sub-rings as (i) diagonal sub-ring \mathcal{D}_n ; (ii) upper triangular sub-ring \mathcal{U}_n ; (iii) lower triangular sub-ring \mathcal{B}_n ; etc.

In addition to these rings with square matrices, are there any rings with non-square matrices?

Recall the DK-STP defined by Definition 2.19. It leads to a perfect ring structure.

Theorem 6.1: [22] Consider $\mathcal{R} := (\mathcal{M}_{m \times n}, +, \times)$, where $+$ is the classical matrix addition/subtraction. Then \mathcal{R} is a ring.

B. Hyper Ring

Definition 6.2: A triple $\mathcal{R} := (R, +, \times)$ is called a hyper-ring, if the following conditions are satisfied.

(i) $(R, +)$ is an Abelian hyper group.

(ii) (R, \times) is a semi-group.

(iii)

$$\begin{aligned} (a+b) \times c &= a \times c + b \times c; \\ c \times (a+b) &= c \times a + c \times b, \quad a, b, c \in R. \end{aligned} \quad (155)$$

If (R, \times) is a hyper-monoid, \mathcal{R} is called a unitary hyper-ring. If (R, \times) is commutative, \mathcal{R} is called a commutative hyper-ring.

In the following we construct the most general matrix ring, which covers all matrices.

Theorem 6.3: The triple $(\mathcal{M}, \hat{+}, \hat{\times})$ is a hyper-ring, where the addition $\hat{+}$ is defined by (29) and the product $\hat{\times}$ is defined by (41).

Proof. It was shown in Proposition 2.28 that $(\mathcal{M}, \hat{\times})$ is a semi-group, and in Proposition 3.8 (iv) that $(\mathcal{M}, \hat{+})$ is an abelian hyper group. Hence what remaining to be proved is the distributivity, that is,

$$\begin{aligned} (A\hat{+}B) \hat{\times} C &= A \hat{\times} C \hat{+} B \hat{\times} C; \\ C \hat{\times} (A\hat{+}B) &= C \hat{\times} A \hat{+} C \hat{\times} B. \end{aligned} \quad (156)$$

We prove the first equality only. Using the first equality and Proposition 2.27, the second equality is obvious.

Assume $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $C \in \mathcal{M}_{r \times s}$, and

$$\begin{aligned} \xi &= m \vee p, & \eta &= n \vee q, \\ \alpha &= \xi \vee r = m \vee p \vee r, & \beta &= \eta \vee s = n \vee q \vee s, \\ \theta &= \alpha \vee \beta = m \vee p \vee r \vee n \vee q \vee s, \\ u &= m \vee r, & v &= n \vee s, \\ w &= u \vee v = m \vee n \vee r \vee s, \\ a &= p \vee r, & b &= q \vee s, \\ e &= a \vee b = p \vee r \vee q \vee s, \\ u \vee a &= m \vee p \vee r = \alpha, \\ v \vee b &= n \vee q \vee s = \beta, \\ \mu &= r \vee q, & \lambda &= r \vee n. \end{aligned}$$

First we have

$$A \hat{+} B = (A \otimes \mathcal{E}_{\xi/m \times \eta/n}) + (B \otimes \mathcal{E}_{\xi/p \times \eta/q}).$$

Using formula (31), we can calculate that

$$\begin{aligned} (A \hat{+} B) \hat{\times} C &= \{[(A \otimes \mathcal{E}_{\xi/m \times \eta/n}) + (B \otimes \mathcal{E}_{\xi/p \times \eta/q})] \\ &\quad \otimes \mathcal{E}_{\alpha/\xi \times \beta/\eta}\} \times (C \otimes \mathcal{E}_{\alpha/r \times \beta/s}) \\ &= [(A \otimes \mathcal{E}_{\alpha/m \times \beta/n}) + (B \otimes \mathcal{E}_{\alpha/p \times \beta/q})] \\ &\quad \times (C \otimes \mathcal{E}_{\alpha/r \times \beta/s}) \\ &= \left\{[(A \otimes \mathcal{E}_{\alpha/m \times \beta/n}) + (B \otimes \mathcal{E}_{\alpha/p \times \beta/q})] \otimes E_{\theta/\beta}^T\right\} \\ &\quad [(C \otimes \mathcal{E}_{\alpha/r \times \beta/s}) \otimes \mathcal{E}_{\theta/\alpha}] \\ &= [(A \otimes \mathcal{E}_{\alpha/m \times \theta/n}) + (B \otimes \mathcal{E}_{\alpha/p \times \theta/q})] (C \otimes \mathcal{E}_{\theta/r \times \beta/s}) \\ &= (A \otimes \mathcal{E}_{\alpha/m \times \theta/n}) (C \otimes \mathcal{E}_{\theta/r \times \beta/s}) \\ &\quad + (B \otimes \mathcal{E}_{\alpha/p \times \theta/q}) (C \otimes \mathcal{E}_{\theta/r \times \beta/s}) \\ &= (A \otimes \mathcal{E}_{\alpha/m \times \lambda/n}) (C \otimes \mathcal{E}_{\lambda/r \times \beta/s}) \\ &\quad + (B \otimes \mathcal{E}_{\alpha/p \times \mu/q}) (C \otimes \mathcal{E}_{\mu/r \times \beta/s}). \end{aligned}$$

On the other hand,

$$\begin{aligned} A \hat{\times} C &= (A \otimes \mathcal{E}_{u/m \times v/n}) \times (C \otimes \mathcal{E}_{u/r \times v/s}). \\ &= [(A \otimes \mathcal{E}_{u/m \times v/n}) \otimes \mathcal{E}_{w/v}^T] [(C \otimes \mathcal{E}_{u/r \times v/s}) \otimes \mathcal{E}_{w/u}] \\ &= (A \otimes \mathcal{E}_{u/m \times w/n}) (C \otimes \mathcal{E}_{w/r \times v/s}) \\ &= (A \otimes \mathcal{E}_{u/m \times \lambda/n}) (C \otimes \mathcal{E}_{\lambda/r \times v/s}). \end{aligned}$$

$$\begin{aligned} B \hat{\times} C &= (B \otimes \mathcal{E}_{a/p \times b/q}) \times (C \otimes \mathcal{E}_{a/r \times b/s}). \\ &= [(B \otimes \mathcal{E}_{a/p \times b/q}) \otimes \mathcal{E}_{e/b}^T] [(C \otimes \mathcal{E}_{a/r \times b/s}) \otimes \mathcal{E}_{e/a}] \\ &= (B \otimes \mathcal{E}_{a/p \times e/q}) (C \otimes \mathcal{E}_{e/r \times b/s}) \\ &= (B \otimes \mathcal{E}_{a/p \times \mu/q}) (C \otimes \mathcal{E}_{\mu/r \times b/s}). \end{aligned}$$

Then

$$\begin{aligned} (A \hat{\times} C) \hat{+} (B \hat{\times} C) &= [(A \otimes \mathcal{E}_{u/m \times \lambda/n}) (C \otimes \mathcal{E}_{\lambda/r \times v/s})] \otimes E_{\alpha/u \times \beta/v} + \\ &\quad [(B \otimes \mathcal{E}_{a/p \times \mu/q}) (C \otimes \mathcal{E}_{\mu/r \times b/s})] \otimes \mathcal{E}_{\alpha/a \times \beta/b} \\ &= (A \otimes \mathcal{E}_{\alpha/m \times \lambda/n}) (C \otimes \mathcal{E}_{\lambda/r \times \beta/s}) \\ &\quad + (B \otimes \mathcal{E}_{\alpha/p \times \mu/q}) (C \otimes \mathcal{E}_{\mu/r \times \beta/s}). \end{aligned}$$

The first equality of (186) follows. \square

Definition 6.4: Consider a hyper-ring $(R, +, \times)$. Assume the identity set of its addition hyper group is $\mathbf{e} = \{e_\lambda \mid \lambda \in \Lambda\}$.

(i) If for each $\lambda \in \Lambda$ the corresponding component group $(R_\lambda, +)$ satisfies that

$$x \times y \in R_\lambda, \quad x, y \in R_\lambda, \quad (157)$$

it is said that the addition $+$ and the product \times of R are consistent.

(ii) $x, y \in R$ are said to be equivalent, if there exist $e_\alpha, e_\beta \in \mathbf{e}$ such that

$$x + e_\alpha = y + e_\beta. \quad (158)$$

The equivalence is said to be consistent with operators, if $x_1 \sim x_2$ and $y_1 \sim y_2$ imply

$$x_1 + y_1 \sim x_2 + y_2; \quad x_1 \times y_1 \sim x_2 \times y_2. \quad (159)$$

Proposition 6.5: Consider $(\mathcal{M}, \hat{+}, \hat{\times})$. (i) $\hat{+}$ and $\hat{\times}$ are consistent; and (ii) the equivalence is consistent with operators.

Proof.

(i) Assume $A_1 \sim A_2$ and $B_1 \sim B_2$. Using all the notations/parameters in the proof of Proposition 3.19 (iv), we have

$$A_1 \hat{\times} B_1 = (A_0 \otimes \mathcal{E}_{a/s \times b/t}) \times (B_0 \otimes \mathcal{E}_{a/\xi \times b/\eta})$$

Assume

$$x := a \vee b = m \vee \alpha \vee n \vee \beta := b\mu.$$

Then

$$\begin{aligned} A_1 \hat{\times} B_1 &= (A_0 \otimes \mathcal{E}_{a/s \times b/t} \otimes \mathcal{E}_{x/b}^T) \\ &\quad (B_0 \otimes \mathcal{E}_{a/\xi \times b/\eta} \otimes \mathcal{E}_{x/a}) \\ &= (A_0 \otimes \mathcal{E}_{a/s \times x/t}) (B_0 \otimes \mathcal{E}_{x/\xi \times b/\eta}) \\ &= (A_0 \otimes \mathcal{E}_{a/s \times b/t} \otimes \mathcal{E}_\mu^T) \\ &\quad (B_0 \otimes \mathcal{E}_{b/\xi \times b/\eta} \otimes \mathcal{E}_\mu) \\ &= (A_0 \otimes \mathcal{E}_{u/s \times v/t} \otimes \mathcal{E}_{a/u \times b/v}) \\ &\quad (B_0 \otimes \mathcal{E}_{u/\xi \times v/\eta} \otimes \mathcal{E}_{a/u \times b/v}) \\ &= [(A_0 \otimes \mathcal{E}_{u/s \times v/t}) (B_0 \otimes \mathcal{E}_{u/\xi \times v/\eta})] \otimes \mathcal{E}_{a/u \times b/v}. \end{aligned}$$

Similarly,

$$\begin{aligned} A_2 \hat{\times} B_2 &= [(A_0 \otimes \mathcal{E}_{u/s \times v/t}) (B_0 \otimes \mathcal{E}_{u/\xi \times v/\eta})] \\ &\quad \otimes \mathcal{E}_{c/u \times d/v}. \end{aligned}$$

Hence

$$A_1 \hat{\times} B_1 \sim A_2 \hat{\times} B_2.$$

(ii) The fact that the equivalence is consistent w.r.t. $\hat{+}$ was proved in Proposition 3.19, and in (i) it has just been proved that the equivalence is consistent w.r.t. $\hat{\times}$. \square

Then similar to the decomposition result of hyper group, we have the following result.

Proposition 6.6: Consider a hyper-ring $(R, +, \times)$ with the identity set of its addition hyper group as $\mathbf{e} = \{e_\lambda \mid \lambda \in \Lambda\}$.




Fig. 5. Ring Decomposition of a Hyper-Ring \mathcal{R}

- (i) If the addition $+$ and the product \times of R are consistent, then each $(R_\lambda, +, \times)$ is a ring, called the component ring of the hyper-ring $(R, +, \times)$.
- (ii) In addition to (i), if the equivalence is consistent with the operators, then the operators over quotient space can be defined properly by

$$\begin{aligned}\bar{x} + \bar{y} &:= \overline{x + y}, & x, y \in R, \\ \bar{x} \times \bar{y} &:= \overline{x \times y}.\end{aligned}\quad (160)$$

Therefore, the quotient space becomes a ring, called the equivalence ring.

Similarly to hyper group, a geometric picture for the structure of a hyper-ring is shown by Figure 5.

Example 6.7: Consider $\mathcal{R} := (\mathcal{M}, \hat{+}, \hat{\times})$. To see the consistency of two operators, let $A, B \in \mathcal{M}_e$, where $e = 0_{m \times n}$. Then

$$A \hat{\times} B = A \times B \in \mathcal{M}_{m \times n} + \mathcal{M}_e.$$

The consistency of two operators is proved. Secondly, it was proved in Proposition 6.5 that the equivalence is consistent with operators. Hence, its decomposition exists. Then we have the following decomposed rings:

- (i) Component Rings:

$$\mathbf{e} = \{e_{m \times n} = 0_{m \times n} \mid (m, n) \in \mathbb{Z}_+ \times \mathbb{Z}_+\}.$$

Each component ring is:

$$\mathcal{R}_{e_{m \times n}} = (\mathcal{M}_{m \times n}, +, \times), \quad m, n \in \mathbb{Z}_+. \quad (161)$$

This set of special rings have been described in the beginning of this section.

- (ii) Equivalence Ring.

By definition, two matrices A and B are equivalent, if there are two identity elements $0_{m \times n}$ and $0_{p \times q}$, such that

$$A \hat{+} 0_{m \times n} = B \hat{+} 0_{p \times q},$$

which leads to the following equivalent statement: Let $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $s = \text{lcm}(m, p)$ and $t = \text{lcm}(n, q)$. Then $A \sim B$, if and only if,

$$A \otimes \mathcal{E}_{s/m \times t/n} = B \otimes \mathcal{E}_{s/p \times t/q}. \quad (162)$$

$A \in \mathcal{M}_{m \times n}$ is reducible, if there exist $(a, b) \prec (m, n)$ and $(a, b) \neq (m, n)$ and $A_0 \in \mathcal{M}_{a \times b}$, such that

$$A = A_0 \otimes \mathcal{E}_{m/a \times n/b}.$$

The equivalence class of A is denoted by

$$\bar{A} = \{B \mid B \sim A\}.$$

Denote the equivalence ring by

$$\bar{\mathcal{R}} := (\bar{\mathcal{M}}, \hat{+}, \hat{\times}), \quad (163)$$

then

$$\bar{\mathcal{M}} = \{\bar{A} \mid A \text{ is irreducible}\}.$$

C. Square Hyper Ring

It is well known that $(\mathcal{M}_{n \times n}, +, \times)$ is a ring, where $+$ and \times are classical matrix addition and product.

This subsection try to extend this ring structure to a larger set of matrices, which are of mixed dimensions.

Proposition 6.8: Assume $\mathcal{R} = (\mathcal{M}_1, \bar{+}, \bar{\times})$, where \mathcal{M}_1 is the set of square matrices; $\bar{+}$ is defined by (25). Then \mathcal{R} is a hyper-ring.

Proof.

It was shown in Proposition 3.8 (ii) that $(\mathcal{M}_1, \bar{+})$ is a commutative hyper group. It is obvious that $(\mathcal{M}_1, \bar{\times})$ is a semi-group.

To see that \mathcal{R} is a hyper-ring, we have only to show the distributivity.

Let $A \in \mathcal{M}_{m \times m}$, $B \in \mathcal{M}_{n \times n}$, $C \in \mathcal{M}_{r \times r}$, $\text{lcm}(m, n) = t$, $\text{lcm}(t, r) = \text{lcm}(m, n, r) = s$, $\text{lcm}(m, r) = \alpha$, $\text{lcm}(n, r) = \beta$, and $\text{lcm}(\alpha, \beta) = \text{lcm}(m, n, r) = s$. Then we have

$$\begin{aligned}(A \bar{+} B) \bar{\times} C &= (A \otimes I_{t/m} + B \otimes I_{t/n}) \bar{\times} C \\ &= [(A \otimes I_{t/m} + B \otimes I_{t/n}) \otimes I_{s/t}] (C \otimes I_{s/r}) \\ &= (A \otimes I_{s/m}) (C \otimes I_{s/r}) + (B \otimes I_{s/n}) (C \otimes I_{s/r}).\end{aligned}$$

$$\begin{aligned}(A \bar{\times} C) \bar{+} (B \bar{\times} C) &= (A \otimes I_{\alpha/m}) (C \otimes I_{\alpha/r}) \\ &\quad \bar{+} (B \otimes I_{\beta/n}) (C \otimes I_{\beta/r}) \\ &= [(A \otimes I_{\alpha/m}) (C \otimes I_{\alpha/r})] \otimes (I_{s/\alpha}) \\ &\quad + [(B \otimes I_{\beta/n}) (C \otimes I_{\beta/r})] \otimes I_{s/\beta} \\ &= (A \otimes I_{s/m}) (C \otimes I_{s/r}) + (B \otimes I_{s/n}) (C \otimes I_{s/r}).\end{aligned}$$

That is,

$$(A\bar{+}B) \times C = (A \times C)\bar{+}(B \times C).$$

Similarly, we can show

$$C \times (A\bar{+}B) = (C \times A)\bar{+}(C \times B).$$

We conclude that $\mathcal{R} = (\mathcal{M}_1, \bar{+}, \times)$ is a hyper-ring.

Finally, we consider the decomposition of this hyper-ring. The identity set is

$$\mathbf{e} = \mathbf{0} = \{0_{n \times n} \mid n = 1, 2, \dots\}.$$

Then

- (i) For each $e_n = 0_{n \times n} \in \mathbf{0}$, $\mathcal{R}_{e_n} = (\mathcal{M}_{n \times n}, \bar{+}, \times) = (\mathcal{M}_{n \times n}, +, \times)$ is a component ring.
- (ii) It is easy to see that $A \sim B$, if and only if, there exist I_α and I_β such that

$$A \otimes I_\alpha = B \otimes I_\beta.$$

This is the one defined in [10]. And the consistence of the addition with the product and the their consistence w.r.t. the equivalence have already been proved in [10]. We conclude that $\bar{\mathcal{R}} = (\bar{\mathcal{M}}_1, \bar{+}, \times)$ is the equivalence ring.

□

Proposition 6.9:

Assume $\mathcal{R} = (\mathcal{M}_1, \hat{+}, \circ)$, where the product \circ is defined by (17) and the addition is defined as follows: Assume $A \in \mathcal{M}_{m \times m}$, $B \in \mathcal{M}_{n \times n}$, and $t = \text{lcm}(m, n)$, then

$$A\hat{+}B := (A \otimes J_{t/m}) + (B \otimes J_{t/n}). \quad (164)$$

Then \mathcal{R} is a hyper-ring.

Proof.

A straightforward verification shows that

- (i) $(\mathcal{M}_1, \hat{+})$ is a hyper group, where the identity set is

$$\mathbf{e} = \{0_{n \times n} \in \mathcal{M}_{n \times n} \mid n = 1, 2, \dots\}.$$

- (ii) (\mathcal{M}_1, \circ) is a semi-group.

To see that \mathcal{R} is a hyper-ring, it is enough to show the distributivity.

Assume $C \in \mathcal{M}_{p \times p}$, and $\alpha = \text{lcm}(m, p)$, $\beta = \text{lcm}(n, p)$, and $s = \text{lcm}(m, n, p)$. Then

$$\begin{aligned} (A\hat{+}B) \circ C &= (A \otimes J_{t/m} + B \otimes J_{t/n}) \circ C \\ &= [(A \otimes J_{t/m} + B \otimes J_{t/n}) \otimes J_{s/t}](C \otimes J_{s/p}) \\ &= [(A \otimes J_{s/m} + B \otimes J_{s/n})(C \otimes J_{s/p})] \\ &= (A \otimes J_{s/m})(C + B \otimes J_{t/n}) \otimes J_{s/t}(C \otimes J_{s/p}) \\ &= (A \otimes J_{s/m})(C \otimes J_{s/p}) + (B \otimes J_{s/n})(C \otimes J_{s/p}). \end{aligned}$$

Meanwhile,

$$\begin{aligned} (A \circ C)\hat{+}(B \circ C) &= (A \otimes J_{\alpha/m})(C \otimes J_{\alpha/p}) \\ &\hat{+}(B \otimes J_{\beta/n})(C \otimes J_{\beta/p}) \\ &= (A \otimes J_{\alpha/m})(C \otimes J_{\alpha/p}) \otimes J_{s/\alpha} \\ &\quad + (B \otimes J_{\beta/n})(C \otimes J_{\beta/p}) \otimes J_{s/\beta} \\ &= (A \otimes J_{\alpha/m})(C \otimes J_{\alpha/p}) \otimes J_{s/\alpha}^2 \\ &\quad + (B \otimes J_{\beta/n})(C \otimes J_{\beta/p}) \otimes J_{s/\beta}^2 \\ &= (A \otimes J_{s/m})(C \otimes J_{s/p}) + (B \otimes J_{s/n})(C \otimes J_{s/p}). \end{aligned}$$

We conclude that

$$(A\hat{+}B) \circ C = (A \circ C)\hat{+}(B \circ C).$$

Similarly, we can also show that

$$C \circ (A\hat{+}B) = (C \circ A)\hat{+}(C \circ B).$$

We conclude that $\mathcal{R} = (\mathcal{M}_1, \hat{+}, \circ)$ is a hyper-ring.

Finally, we consider the decomposition of this hyper-ring. The identity set is

$$\mathbf{e} = \mathbf{0} = \{0_{n \times n} \mid n = 1, 2, \dots\}.$$

Then

- (i) For each $e_n = 0_{n \times n} \in \mathbf{0}$, $\mathcal{R}_{e_n} = (\mathcal{M}_{n \times n}, \hat{+}, \circ) = (\mathcal{M}_{n \times n}, +, \times)$ is a component ring.
- (ii) Since $A \sim B$ means there exist J_α and J_β such that

$$A \otimes J_\alpha = B \otimes J_\beta.$$

To prove the consistence of the addition and the product with the equivalence we need to show that for $A_1 \sim A_2$ and $B_1 \sim B_2$ we have

$$A_1\hat{+}B_1 \sim A_2\hat{+}B_2; \quad (165)$$

and

$$A_1 \circ B_1 \sim A_2 \circ B_2. \quad (166)$$

We prove (166) only. The proof of (165) is similar.

Assume $A_1 \in \mathcal{M}_{m \times m}$, $A_2 \in \mathcal{M}_{n \times n}$, $B_1 \in \mathcal{M}_{r \times r}$, and $B_2 \in \mathcal{M}_{s \times s}$ with

$$A_1 \sim A_2; \quad B_1 \sim B_2.$$

Denote $\text{gcd}(m, n) = \alpha$, and $\text{gcd}(r, s) = \beta$. Using Lemma 3.16, we know that there exist $A_0 \in \mathcal{M}_{\alpha \times \alpha}$ and $B_0 \in \mathcal{M}_{\beta \times \beta}$ such that

$$\begin{aligned} A_1 &= A_0 \otimes J_{m/\alpha}, & A_2 &= A_0 \otimes J_{n/\alpha}; \\ B_1 &= B_0 \otimes J_{r/\beta}, & B_2 &= B_0 \otimes J_{s/\beta}. \end{aligned}$$

Set $\text{lcm}(m, r) = p$, $\text{lcm}(n, s) = q$, $\text{lcm}(p, q) = \mu$, then we have

$$\begin{aligned} A_1 \circ B_1 &= (A_0 \otimes J_{m/\alpha} \otimes J_{p/m})(B_0 \otimes J_{r/\beta} \otimes J_{p/r}) \\ &= (A_0 \otimes J_{p/\alpha})(B_0 \otimes J_{p/\beta}). \end{aligned}$$

Similarly, we have

$$\begin{aligned} A_2 \circ B_2 &= (A_0 \otimes J_{n/\alpha} \otimes J_{q/n})(B_0 \otimes J_{s/\beta} \otimes J_{q/s}) \\ &= (A_0 \otimes J_{q/\alpha})(B_0 \otimes J_{q/\beta}). \end{aligned}$$

Note that

$$\begin{aligned} (A_1 \circ B_1) \otimes J_{\mu/p} &= (A_0 \otimes J_{p/\alpha})(B_0 \otimes J_{p/\beta}) \otimes J_{\mu/p}^2 \\ &= (A_0 \otimes J_{\mu/\alpha})(B_0 \otimes J_{\mu/\beta}); \end{aligned}$$

and

$$\begin{aligned} (A_2 \circ B_2) \otimes J_{\mu/q} &= (A_0 \otimes J_{q/\alpha})(B_0 \otimes J_{q/\beta}) \otimes J_{\mu/q}^2 \\ &= (A_0 \otimes J_{\mu/\alpha})(B_0 \otimes J_{\mu/\beta}). \end{aligned}$$

(166) follows.

We conclude that $\overline{\mathcal{R}} = (\overline{\mathcal{M}}_1, \hat{+}, \circ)$ is the equivalence ring.

□

D. Sub Hyper Ring

Definition 6.10: Let $(R, +, \times)$ be a hyper-Ring and $H \subset R$. $(H, +, \times)$ is said to be a sub-hyper-ring of $(R, +, \times)$, if (i) $(H, +) <_h (R, +)$ is a sub-hyper group, and (H, \times) is a sub-semi-group of (R, \times) .

Proposition 6.11:

- (i) $\mathcal{R}_\mu = (\mathcal{M}_\mu, \hat{+}, \hat{\times})$ is a hyper-ring, where $\hat{+}$ is defined by (26) and $\hat{\times}$ is defined by (42).
- (ii) \mathcal{R}_μ is a sub-hyper-ring of $\mathcal{R} = (\mathcal{M}, \hat{+}, \hat{\times})$.

Proof.

- (i) It has been proved in Proposition 3.8 that $(\mathcal{M}_\mu, \hat{+})$ is a hyper group, and in Remark 2.29 that $(\mathcal{M}_\mu, \hat{\times})$ is a hyper-semigroup. The proof of distributivity is exactly the same as the one for $(\mathcal{M}, \hat{+}, \hat{\times})$.
- (ii) A careful observation shows that the $\hat{+}$ defined by (26) and $\hat{\times}$ defined by (42) respectively are exactly the same as the restrictions of the corresponding $\hat{+}$ and $\hat{\times}$ for \mathcal{M} on \mathcal{M}_μ . Then the conclusion follows immediately.

□

Example 6.12: Define

$$\begin{aligned} U^1 &= \{A \in \mathcal{M}_1 \mid A \text{ is upper triangular.}\}; \\ B^1 &= \{A \in \mathcal{M}_1 \mid A \text{ is lower triangular.}\}; \\ D^1 &= \{A \in \mathcal{M}_1 \mid A \text{ is diagonal.}\}; \end{aligned}$$

Then (i) $(U^1, \bar{+}, \times)$ is a sub-hyper ring of $(\mathcal{M}_1, \bar{+}, \times)$. (ii) $(B^1, \bar{+}, \times)$ is a sub-hyper ring of $(\mathcal{M}_1, \bar{+}, \times)$. (iii) $(D^1, \bar{+}, \times)$ is a sub-hyper ring of $(\mathcal{M}_1, \bar{+}, \times)$.

E. Hyper Ring Homomorphism

Definition 6.13: Let $\mathcal{N} = (N, +, \times)$, and $\mathcal{R} = (R, +, \times)$ be two hyper rings. A mapping $\varphi : N \rightarrow R$ is called a hyper-ring homomorphism, if $\varphi : (N, +) \rightarrow (R, +)$ is a hyper group homomorphism, and

$$\varphi(x \times y) = \varphi(x) \times \varphi(y), \quad x, y \in N. \quad (167)$$

Proposition 6.14: Consider $\mathcal{R} = (\mathcal{M}, \hat{+}, \hat{\times})$ and $\mathcal{N} := (\mathcal{M}_1, \hat{+}, \circ)$. Let $\square : \mathcal{R} \rightarrow \mathcal{N}$. Then \square is a hyper-ring homomorphism.

Proof. It was proved in Proposition 5.17 that $\square : (\mathcal{M}, \hat{+}) \rightarrow (\mathcal{M}_1, \hat{+})$ is a hyper group homomorphism. So we have only to prove (167).

Assume $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $s = m \vee p$, $t = n \vee q$, $\alpha = m \vee n$, $\beta = p \vee q$, and $\xi = s \vee t = \alpha \vee \beta$. Then

$$\begin{aligned} \square(A \hat{\times} B) &= \square[(A \otimes \mathcal{E}_{s/m \times t/n}) \times (B \otimes \mathcal{E}_{s/p \times t/q})] \\ &= \square[(A \otimes \mathcal{E}_{s/m \times \xi/n}) (B \otimes \mathcal{E}_{\xi/p \times t/q})] \\ &= (A \otimes \mathcal{E}_{\xi/m \times \xi/n}) (B \otimes \mathcal{E}_{\xi/p \times \xi/q}). \end{aligned}$$

$$\square(A) = A \otimes \mathcal{E}_{\alpha/m \times \alpha/n};$$

$$\square(B) = B \otimes \mathcal{E}_{\beta/p \times \beta/q}.$$

$$\begin{aligned} \square(A) \circ \square(B) &= [(A \otimes \mathcal{E}_{\alpha/m \times \alpha/n}) \otimes J_{\xi/\alpha}] \\ &\quad [(B \otimes \mathcal{E}_{\beta/p \times \beta/q}) \otimes J_{\xi/\beta}] \\ &= [(A \otimes \mathcal{E}_{\alpha/m \times \alpha/n}) \otimes \mathcal{E}_{\xi/\alpha \times \xi/\alpha}] \\ &\quad [(B \otimes \mathcal{E}_{\beta/p \times \beta/q}) \otimes \mathcal{E}_{\xi/\beta \times \xi/\beta}] \\ &= (A \otimes \mathcal{E}_{\xi/m \times \xi/n}) (B \otimes \mathcal{E}_{\xi/p \times \xi/q}). \end{aligned}$$

Hence,

$$\square(A \hat{\times} B) = \square(A) \hat{\times} \square(B).$$

□

VII. HYPER MODULE AND GENERAL LINEAR (CONTROL) SYSTEM

A. Hyper Module

Similarly to the definition of module [31], a hyper module can be defined as follows.

Definition 7.1: Let R be a hyper ring. A (left) R -module is an additive (abelian) hyper group A with a function $\pi : R \times A \rightarrow A$ such that

(i)

$$r(a + b) = ra + rb, \quad r \in R, a, b \in A; \quad (168)$$

(ii)

$$(r + s)a = ra + sa, \quad r, s \in R, a, b \in A; \quad (169)$$

(iii)

$$r(sa) = (rs)a. \quad (170)$$

If R has an identity set \mathbf{e}_R for the product, such that

$$e_{Rx} \sim x \quad e_R \in \mathbf{e}_R, \forall x \in A, \quad (171)$$

and there exists at least one $e_x \in \mathbf{e}_R$, such that

$$e_x x = x, \quad (172)$$

then A is said to be a unitary hyper R -module.

Remark 7.2:

- (i) A module can be considered as a particular hyper module, where the R is degenerated to a ring.
- (ii) The equality (171) can equivalently be written as

$$e_{Rx} - x \in \mathbf{e}, \quad e_R \in \mathbf{e}_R. \quad (173)$$

- (iii) We are particularly interested in unitary (hyper-)R-module.

Consider the (unitary) module of (hyper-)matrix ring with (hyper-)vector group.

1) (Hyper) Vector Group:

- (i)

$$\mathcal{A}_1^V := (\mathbb{R}^n, +), \quad n \in \mathbb{Z}_+, \quad (174) \quad (\text{iv})$$

where $+$ is the classical vector addition. \mathcal{A}_1^V is a vector group with unique identity 0_n .

- (ii)

$$\mathcal{A}_2^V := (\mathbb{R}^\infty, \hat{+}), \quad (175)$$

where $\hat{+}$ is defined as follows: Let $x \in \mathbb{R}^m, y \in \mathbb{R}^n$, and $t = \text{lcm}(m, n)$. Then

$$x \hat{+} y := (x \otimes \mathbf{1}_{t/m}) + (y \otimes \mathbf{1}_{t/n}) \in \mathbb{R}^t. \quad (176)$$

Then \mathcal{A}_2^V is a hyper-vector group with identity set

$$\mathbf{e} = \{e_n = 0_n \mid n \in \mathbb{Z}_+\}.$$

- (iii)

$$\mathcal{A}_3^V := (\mathbb{R}^\infty, \bar{+}), \quad (177)$$

where $\bar{+}$ is the defined as follows: Let $x \in \mathbb{R}^m, y \in \mathbb{R}^n$, and $t = \text{lcm}(m, n)$. Then

$$x \hat{+} y := (x \otimes \mathcal{E}_{t/m}) + (y \otimes \mathcal{E}_{t/n}) \in \mathbb{R}^t. \quad (178)$$

Then \mathcal{A}_3^V is a hyper-vector group with identity set

$$\mathbf{e} = \{e_n = 0_n \mid n \in \mathbb{Z}_+\}.$$

2) (Hyper) Matrix Ring:

- (i)

$$\mathcal{R}_1^M := (\mathcal{M}^{n \times n}, +, \times), \quad n \in \mathbb{Z}_+, \quad (179)$$

where $+$ and \times are the classical matrix addition and production respectively. \mathcal{R}_1^M is a matrix ring with unique identity I_n .

- (ii)

$$\mathcal{R}_2^M := (\overline{\mathcal{M}}^{m \times n}, +, \times), \quad m, n \in \mathbb{Z}_+, \quad (180)$$

where $+$ is the classical matrix addition and the \times is the DK-STP, defined by (32). \mathcal{R}_2^M is a matrix ring with the unique identity $I_{m \times n}$.

- (iii)

$$\mathcal{R}_3^M := (\mathcal{M}_1, \bar{+}, \times), \quad (181)$$

where $\bar{+}$ is defined by (25) and the \times is defined by (16). \mathcal{R}_3^M is a matrix unitary hype-ring with the identity set

$$\mathbf{e} = \{0_{n \times n} \mid n \in \mathbb{Z}_+\};$$

and

$$\mathbf{e}_R = \{I_n \mid n \in \mathbb{Z}_+\}.$$

$$\mathcal{R}_4^M := (\mathcal{M}_1, \hat{+}, \circ), \quad (182)$$

where $\hat{+}$ is defined by (29) and the \circ is defined by (17). \mathcal{R}_4^M is a matrix unitary hype-ring with the identity set

$$\mathbf{e} = \{0_{n \times n} \mid n \in \mathbb{Z}_+\};$$

and

$$\mathbf{e}_R = \{I_n \mid n \in \mathbb{Z}_+\}.$$

- (v)

$$\mathcal{R}_5^M := (\overline{\mathcal{M}}_\mu, \hat{+}, \hat{\times}), \quad (183)$$

where the $\hat{\times}$ is defined by (41). \mathcal{R}_5^M is a matrix unitary hype-ring with the identity set

$$\mathbf{e} = \{0_{n\mu_y \times n\mu_x} \mid n \in \mathbb{Z}_+\};$$

and

$$\mathbf{e}_R = \{I_n \mid n \in \mathbb{Z}_+\}.$$

- (vi)

$$\mathcal{R}_6^M := (\overline{\mathcal{M}}, \hat{+}, \hat{\times}). \quad (184)$$

\mathcal{R}_6^M is a matrix unitary hype-ring with the identity set

$$\mathbf{e} = \{0_{m \times n} \mid m, n \in \mathbb{Z}_+\};$$

and

$$\mathbf{e}_R = \{I_{m \times n} \mid m, n \in \mathbb{Z}_+\}.$$

3) (Hyper) Matrix Module: We construct the following combinations of matrix (hyper) rings with vector (hyper) groups.

- (i) $\aleph_1 := (\mathcal{R}_1^M, \mathcal{A}_1^V, \times)$, where \times is the classical matrix-vector product.
- (ii) $\aleph_2 := (\mathcal{R}_2^M, \mathcal{A}_2^V, \times)$ (or $(\mathcal{R}_2^M, \mathcal{A}_3^V, \times)$), where \times is the DK-STP, defined by (32).
- (iii) $\aleph_3 := (\mathcal{R}_3^M, \mathcal{A}_2^V, \times)$ (or $(\mathcal{R}_3^M, \mathcal{A}_3^V, \times)$).
- (iv) $\aleph_4 := (\mathcal{R}_4^M, \mathcal{A}_2^V, \circ)$ (or $(\mathcal{R}_4^M, \mathcal{A}_3^V, \circ)$).
- (v) $\aleph_5 := (\mathcal{R}_5^M, \mathcal{A}_2^V, \times)$ (or $(\mathcal{R}_5^M, \mathcal{A}_3^V, \times)$).
- (vi) $\aleph_6 := (\mathcal{R}_6^M, \mathcal{A}_2^V, \times)$ (or $(\mathcal{R}_6^M, \mathcal{A}_3^V, \times)$).

Proposition 7.3:

- (i) The \aleph_1 and \aleph_2 are unitary modules.
- (ii) The $\aleph_3, \aleph_4, \aleph_5$ and \aleph_6 are unitary hyper-modules.

Proof. We prove \aleph_2 , the proof of others is similar. What do we need to verify is (168)-(170). In fact, (169) is trivial, and (170) is well know [19]. Hence, we prove (168) only.

Let $A \in \mathcal{M}_{m \times n}$, $x \in \mathbb{R}^p$, $y \in \mathbb{R}^q$, $p \vee n = \alpha$, $q \vee n = \beta$, $p \vee q = t$, and $n \vee p \vee q = s$. Then

$$\begin{aligned} A \times x &= \left(A \otimes \mathbf{1}_{\alpha/n}^T \right) (x \otimes \mathbf{1}_{\alpha/p}) \in \mathbb{R}^m; \\ A \times y &= \left(A \otimes \mathbf{1}_{\beta/n}^T \right) (y \otimes \mathbf{1}_{\beta/q}) \in \mathbb{R}^m. \end{aligned}$$

$$\begin{aligned} A \times (x \hat{+} y) &= A \times (x \otimes \mathbf{1}_{t/p} + y \otimes \mathbf{1}_{t/q}) \\ &= \left(A \otimes \mathbf{1}_{s/n}^T \right) (x \otimes \mathbf{1}_{s/p} + y \otimes \mathbf{1}_{s/q}) \\ &= \left(A \otimes \mathbf{1}_{s/n}^T \right) (x \otimes \mathbf{1}_{s/p}) + \left(A \otimes \mathbf{1}_{s/n}^T \right) (y \otimes \mathbf{1}_{s/q}) \\ &= \left(A \otimes \mathbf{1}_{\alpha/n}^T \right) (x \otimes \mathbf{1}_{\alpha/p}) + \left(A \otimes \mathbf{1}_{\beta/n}^T \right) (y \otimes \mathbf{1}_{\beta/q}) \end{aligned}$$

(168) follows.

B. Linear (Control) System

Definition 7.4: Let $\aleph = (\mathcal{R}_0, \mathcal{A}_0, \pi)$ be a (hyper-) matrix-vector (MV-) module.

- (i) A discrete-time linear system is defined as

$$\begin{aligned} x(t+1) &= M(t)x(t), \quad x(0) = x_0 \in \mathcal{V}_0, \\ M(t) &\in \mathcal{R}_0, \quad x(t) \in \mathcal{A}_0, \end{aligned} \quad (185)$$

where $M(t)x(t) = \pi(M(t), x(t))$.

- (ii) A continuous-time linear system is defined as

$$\begin{aligned} \dot{x} &= M(t)x(t), \quad x(0) = x_0 \in \mathcal{V}_0, \\ M(t) &\in \mathcal{R}_0, \quad x(t) \in \mathcal{A}_0. \end{aligned} \quad (186)$$

- (iii) A discrete-time linear control system is defined as

$$\begin{aligned} x(t+1) &= M(t)x(t) + B(t)u(t), \quad x(0) = x_0 \in \mathcal{V}_0, \\ M(t), B(t) &\in \mathcal{R}_0, \quad x(t), u(t) \in \mathcal{A}_0. \end{aligned} \quad (187)$$

- (iv) A continuous-time linear control system is defined as

$$\begin{aligned} \dot{x} &= M(t)x(t) + B(t)u(t), \quad x(0) = x_0 \in \mathcal{V}_0, \\ M(t), B(t) &\in \mathcal{R}_0, \quad x(t) \in \mathcal{A}_0. \end{aligned} \quad (188)$$

Remark 7.5:

- (i) When $M(t) = M$ (and $B(t) = B$ for control systems) the systems are called the time-invariant linear (control) systems.
- (ii) We are only interested in the case when $\aleph \in \{\aleph^i \mid i \in [1, 6]\}$, which were defined in the previous subsection.
- (iii) When $\aleph = \aleph^1$, the systems are the classical linear (control) systems, which have been investigated over 8 decades [48].
- (iv) When $\aleph = \aleph^3$ or $\aleph = \aleph^4$, the systems have been investigated by [10]. Particularly, the trajectories of the time-invariant systems have been constructed.

Example 7.6: Consider a linear control system

$$\dot{x} = Ax(t) + Bu(t) + C\eta(t), \quad (189)$$

where $x(t) \in \mathbb{R}^n$ is the state, $u(t) \in \mathbb{R}^m$ is the control, $\eta(t) \in \mathbb{R}^r$ is the disturbance, $A \in \mathcal{M}_{n \times n}$, $B \in \mathcal{M}_{n \times m}$, $C \in \mathcal{M}_{n \times r}$. Assume $\eta(t)$ is a dimension-varying disturbance, then we may replace (189) by

$$\dot{x} = Ax(t) + Bu(t) + C \times \eta(t).$$

Then the system can handle disturbance with various dimensions. If there are some state variable dimension-varying vibrations, say,

$$x(t_\tau) = x(t_{\tau-}), \quad \tau \in \{t_1, t_2, \dots\},$$

- Then (189) can be replaced by

$$\dot{x} = A \times x(t) + Bu(t) + C \times \eta(t).$$

VIII. HYPER VECTOR SPACE

Up to now the algebraic structures of MVMDs have been investigated. From this section on, we will consider the geometric structure of MVMDs.

A. From Hyper Group to Hyper Vector Space

To be precise we first review the concept of vector space in linear algebra.

Definition 8.1: [29] A vector space \mathcal{V} over \mathbb{R} is a triple $\mathcal{V} = (X, +, \cdot)$, where X is a set with elements $x \in X$ are vectors; addition $+$: $X \times X \rightarrow X$, and scalar product \cdot : $\mathbb{R} \times X \rightarrow X$, satisfying

- (1) $(X, +)$ is an abelian group, that is,

- (i) Associativity:

$$(x + y) + z = x + (y + z), \quad x, y, z \in X. \quad (190)$$

(ii) Commutativity:

$$x + y = y + x, \quad x, y \in X. \quad (191)$$

(iii) Zero: There exists a unique $\vec{0} \in X$, such that

$$x + \vec{0} = \vec{0} + x = x, \quad \forall x \in X. \quad (192)$$

(iv) Inverse: For each $x \in X$ there exists a $-x \in X$, such that

$$x + (-x) = 0, \quad x \in X. \quad (193)$$

(2) Scalar product satisfies

(i) Associativity:

$$(r_1 r_2) \cdot x = r_1 \cdot (r_2 \cdot x), \quad r_1, r_2 \in \mathbb{F}, \quad x \in X. \quad (194)$$

(ii) Distributive Law:

$$\begin{aligned} (r_1 + r_2) \cdot x &= r_1 \cdot x + r_2 \cdot x, \quad r, r_1, r_2 \in \mathbb{R}, \\ r \cdot (x + y) &= r \cdot x + r \cdot y, \quad x, y \in X. \end{aligned} \quad (195)$$

(iii) Unit:

$$1 \cdot x = x, \quad 1 \in \mathbb{R}, \quad x \in X. \quad (196)$$

When the MVMDs are considered, it is natural to consider “pseudo-” vector space, which is defined by modifying the definition of vector space a little bit.

Definition 8.2: A hyper-vector space \mathcal{V} over \mathbb{R} is a triple $\mathcal{V} = (X, +, \cdot)$ as in Definition 8.1, satisfying

- (1') $(X, +)$ is an abelian hyper group.
- (2) Same as the (2) in Definition 8.1.

Recall that each hyper group has its component groups and as the operator (it is addition now) is consistent w.r.t. equivalence then the equivalence group exists. Correspondingly, we have the following result.

Proposition 8.3: Consider a hyper-vector space $\mathcal{V} = (X, +, \cdot)$. Assume $\mathbf{e} = \{e_\lambda \mid \lambda \in \Lambda\}$. Then

- (i) For each $\lambda \in \Lambda$, X_λ is a vector space.
- (ii) Assume the addition is consistent w.r.t. equivalence, then the equivalence group is a vector space.

Proof.

(i) Let $x \in V_\lambda$. Then,

$$xe_\lambda = e_\lambda x = x, \quad x^{-1}e_\lambda = e_\lambda x^{-1} = x^{-1}.$$

It follows that for any $\alpha \in \mathbb{R}$

$$\begin{aligned} (\alpha x)e_\lambda &= e_\lambda(\alpha x) = \alpha x, \\ (\alpha x)^{-1}e_\lambda &= e_\lambda(\alpha x)^{-1} = (\alpha x)^{-1}. \end{aligned}$$

Hence $\alpha x \in X_\lambda$, which ensures that V_λ is a vector space.

(ii) Since

$$\overline{\alpha x} = \alpha \overline{x}, \quad \alpha \in \mathbb{R},$$

$(\vec{V}, +)$ is also a vector space. □

B. Hyper Vector Space over Matrices

It was mentioned in Remark 3.9 that all the addition hyper groups on MVMDs are abelian. In addition, for MVMDs the scalar product is naturally defined. Moreover, the properties of scalar product required for vector spaces are automatically satisfied. Then based on Proposition 3.8 the following results are obvious.

Proposition 8.4:

- (i) $(\mathbb{R}^\infty, \hat{+})$ is a hyper-vector space.
- (ii) $(\mathcal{M}_1, \hat{+})$ is a hyper-vector space. $(\mathcal{M}_1, \hat{+})$ is also a hyper-vector space.
- (iii) $(\mathcal{M}_\mu, \hat{+})$ is a hyper-vector space with $\hat{+}$ defined by (26), where the identity set is the same as (53).
- (iv) $(\mathcal{M}, \hat{+})$ is a hyper-vector space.

The decomposition for hyper groups can be immediately extended to hyper-vector space. Then we have

Proposition 8.5: Let $(X, +, \mathbf{e})$ be a hyper-vector space.

- (i) For each $e \in \mathbf{e}$, $(X_e, +)$ is a vector space, called the component subspace of the hyper-vector space.
- (ii) Assume the equivalence is consistent with operator $+$ and scalar product, then the quotient space is also a vector space called the equivalence vector space.

IX. METRIC SPACE STRUCTURE ON \mathbb{R}^∞

A. Hyper Inner Product Space

Replacing vector space by hyper-vector space, the hyper inner product space is defined as follows [44].

Definition 9.1: A hyper-vector space X over \mathbb{F} ($\mathbb{F} = \mathbb{C}$ or $\mathbb{F} = \mathbb{R}$) is called an hyper-inner product space if to each pair of elements $x, y \in X$ there is associated a number $\langle x, y \rangle \in \mathbb{F}$, called the hyper-inner product of x and y , with the following properties:

(i)

$$\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle, \quad x, y, z \in X. \quad (197)$$

(ii)

$$\langle x, y \rangle = \overline{\langle y, x \rangle}. \quad (198)$$

(iii)

$$\langle \alpha x, y \rangle = \alpha \langle x, y \rangle. \quad (199)$$

(iv)

$$\langle x, x \rangle \geq 0, \text{ and } \langle x, x \rangle \neq 0, \text{ if } x \notin \mathbf{0}. \quad (200)$$

Consider \mathbb{R}^∞ , an inner product has been defined as follows.

Definition 9.2: [10] Consider $x, y \in \mathbb{R}^\infty$, say, $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, and $\text{lcm}(m, n) = t$. The inner product of x and y is defined by

$$\begin{aligned} \langle x, y \rangle_{\mathcal{V}} &:= \frac{1}{t} x^T y \\ &= \frac{1}{t} (x \otimes \mathbf{1}_{t/m})^T (y \otimes \mathbf{1}_{t/n}). \end{aligned} \quad (201)$$

Recall that $(\mathbb{R}^\infty, \hat{+}, \cdot)$ is a hyper-vector space, it is readily verifying that the inner product defined by (201) is a hyper-inner product.

Note that by Definition 9.2 if $x \in \mathbb{R}^n$, then

$$\langle x, x \rangle_{\mathcal{V}} = \frac{1}{n} x^T x = \frac{1}{n} \langle x, x \rangle, \quad (202)$$

where $\langle x, x \rangle$ is the Euclidean inner product. Hence the inner product defined by (201) has the “normalized” Euclidean inner product. This is very important for comparing the vectors in Euclidean spaces of various dimensions.

Alternatively, we can also define

$$\langle x, y \rangle_{\mathcal{V}} := (x \otimes \mathcal{E}_{t/m})^T (y \otimes \mathcal{E}_{t/n}). \quad (203)$$

For a hyper-inner product space, the inner-product is said to be consistent with the operator (addition) if $x_1 \sim x_2$ and $y_1 \sim y_2$ imply

$$\langle x_1, y_1 \rangle = \langle x_2, y_2 \rangle.$$

The following result is an immediate consequence of the definition.

Proposition 9.3: Let $\mathcal{H} := (\mathcal{V}, \langle \cdot, \cdot \rangle)$ be a hyper-inner product space, where \mathcal{V} is a hyper-vector space and $\langle \cdot, \cdot \rangle$ is the inner product. If the inner-product is consistent with the addition, then the equivalence space, denoted by $\overline{\mathcal{H}} := (\overline{\mathcal{V}}, \langle \cdot, \cdot \rangle)$, is an inner product space, with the inner product as

$$\langle \bar{x}, \bar{y} \rangle := \langle x, y \rangle. \quad (204)$$

Example 9.4: Consider the hyper-inner product space $\mathcal{H} := (\mathbb{R}^\infty, \hat{+}, \langle \cdot, \cdot \rangle)$, where the inner product is defined by (201). It has been proved in [10] that this inner product is consistent with the addition, then it is clear that the equivalence space $\overline{\mathcal{H}} := (\overline{\mathbb{R}^\infty}, \hat{+}, \langle \cdot, \cdot \rangle)$ is an inner-product space. It has also been proved in [10] that this inner-product space is not complete, hence it is not a Hilbert space.

Definition 9.5:

- (1) A norm on a hyper-vector space X is a real function, whose value at $x \in X$, denoted by $\|x\|$, with the properties:

(i)

$$\|x + y\| \leq \|x\| + \|y\|, \quad x, y \in X. \quad (205)$$

(ii)

$$\|ax\| = |a|\|x\|. \quad (206)$$

(iii)

$$\|x\| \geq 0, \text{ and } \|x\| \neq 0, \text{ if } x \notin \mathbf{0}. \quad (207)$$

- (2) Assume \mathcal{V} is a hyper-inner product space, i.e., it is a hyper-vector space with an inner product $\langle \cdot, \cdot \rangle$, then a norm can be defined by

$$\|x\| := \sqrt{\langle x, x \rangle}. \quad (208)$$

- (3) The norm is said to be consistent with the equivalence, if $x \sim y$ implies $\|x\| = \|y\|$.

A hyper-vector space with a norm is called a normed hyper-vector space.

The following is an immediate consequence of definitions.

Proposition 9.6: Consider a normed hyper-vector space. If the norm is consistent with the equivalence, its equivalence space is a normed vector space.

Example 9.7: Consider the hyper-inner product space $\mathcal{H} := (\mathbb{R}^\infty, \hat{+}, \langle \cdot, \cdot \rangle)$.

- (i) Its norm has been defined in [10] as

$$\|x\|_{\mathcal{V}} := \sqrt{\langle x, x \rangle_{\mathcal{V}}}. \quad (209)$$

- (ii) It has been proved in [10] that the norm defined by (208) is consistent with the equivalence. Hence the equivalence space $(\overline{\mathbb{R}^\infty}, \hat{+}, \langle \cdot, \cdot \rangle)$ is a normed vector space.

Let X be a normed hyper-vector space, and $x, y \in X$. A natural way to define the distance between x and y is

$$d(x, y) := \|x - y\|. \quad (210)$$

As an example, consider \mathbb{R}^∞ . Let $x, y \in \mathbb{R}^\infty$. The distance of x and y is defined by [10]

$$d_{\mathcal{V}}(x, y) := \|x - y\|_{\mathcal{V}}. \quad (211)$$

It is readily verifying that the distance defined in (210), or particularly (211) for \mathbb{R}^∞ , it is a hyper-metric, which turns the hyper-vector space into a hyper-metric space.

Proposition 9.8:

- (i) The metric defined by (211) is a hyper-metric, which makes (\mathbb{R}^∞, d) a hyper-metric space.
(ii) The hyper-metric space is a special pseudo-metric space.
(iii) The metric defined by (211) is consistent with equivalence. That is, assume $x_1 \sim x_2$ and $y_1 \sim y_2$, then $d(x_1, y_1) = d(x_2, y_2)$.

(iv) The equivalence space $\bar{\mathbb{R}}^\infty$ is a metric space.

The topology deduced by the hyper-metric $d_{\mathcal{V}}$, defined by (211) and denoted by \mathcal{T}_d , makes \mathbb{R}^∞ a topological space. This topological space has the following properties [10]:

Proposition 9.9: Consider the topological space $(\mathbb{R}^\infty, \mathcal{T}_d)$.

- (i) It is separable. That is, there exists a countable subset, which is dense in \mathbb{R}^∞ .
- (ii) It is not a Hausdorff space. (In fact, it is not even T_0).

Finally, we briefly review its equivalence space. It is easy to verify the following equivalent statements.

Proposition 9.10: Let $x, y \in \mathbb{R}^\infty$. x and y are equivalent, if one of the following equivalent statements holds.

- (i) x, y are equivalent, denoted by $x \leftrightarrow y$.
- (ii) There exist 0_p and 0_q such that $x \hat{+} 0_p = y \hat{+} 0_q$.
- (iii) There exist $\mathbf{1}_p$ and $\mathbf{1}_q$ such that $x \otimes \mathbf{1}_p = y \otimes \mathbf{1}_q$.
- (iv) $d_{\mathcal{V}}(x, y) = 0$.

Denote the quotient space by

$$\Omega := \mathbb{R}^\infty / \leftrightarrow. \quad (212)$$

According to the Proposition 9.10, it is clear that Ω is both the equivalence space and the topological quotient space. Moreover, this space is a Hausdorff space.

We refer to [24] for related topics about topological spaces; and to [44], [25] for some related knowledge for inner product space.

B. Geometric Structure on \mathbb{R}^∞

First, we consider the projection from \mathbb{R}^m to \mathbb{R}^n .

Definition 9.11: [10] Let $\xi \in \mathbb{R}^m$. The projection of ξ on \mathbb{R}^n , denoted by $\pi_n^m(\xi)$, is defined as

$$\pi_n^m(\xi) := \operatorname{argmin}_{x \in \mathbb{R}^n} \|\xi \hat{-} x\|_{\mathcal{V}}. \quad (213)$$

The projection can be expressed by matrix product form as follows.

Proposition 9.12: [10]

$$x_0 := \pi_n^m(\xi) = \Pi_n^m \xi \in \mathbb{R}^n, \quad \xi \in \mathbb{R}^m, \quad (214)$$

where $(t = \operatorname{lcm}(m, n))$

$$\Pi_n^m = \frac{1}{t/n} \left(I_n \otimes \mathbf{1}_{t/n}^T \right) \left(I_m \otimes \mathbf{1}_{t/m} \right), \quad (215)$$

Moreover, $x_0 \perp (\xi \hat{-} x_0)$ (see Figure 6).

We give an example:




Fig. 6. Cross-dimensional Projection

Example 9.13: Assume $m = 4, n = 6$. Let $\xi = (\xi_1, \xi_2, \xi_3, \xi_4)^T \in \mathbb{R}^m$. We consider its projection on \mathbb{R}^6 . First, $t = \operatorname{lcm}(m, n) = 12$, then

$$\begin{aligned} \Pi_6^4 &= \frac{1}{2} \left(I_6 \otimes \mathbf{1}_2^T \right) \left(I_4 \otimes \mathbf{1}_3 \right) \\ &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.5 & 0.5 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0.5 & 0.5 \\ 0 & 0 & 0 & 1 \end{bmatrix}. \end{aligned}$$

Hence,

$$x_0 = (\xi_1, 0.5\xi_1 + 0.5\xi_2, \xi_2, \xi_3, 0.5\xi_3 + 0.5\xi_4, \xi_4)^T.$$

$$\begin{aligned} x_0 \hat{-} \xi &= (0, 0, 0.5\xi_2 - 0.5\xi_1, 0.5\xi_1 - 0.5\xi_2, 0, 0 \\ &\quad 0, 0, 0.5\xi_4 - 0.5\xi_3, 0.5\xi_3 - 0.5\xi_4, 0, 0)^T. \end{aligned}$$

It is ready to check that

$$\langle x_0 \hat{-} \xi, x_0 \rangle_{\mathcal{V}} = 0.$$

Theorem 9.14: (Generalized Pythagoras' Theorem)

- (i) Let $\xi \in \mathbb{R}^m, x_0 = \pi_n^m(\xi) \in \mathbb{R}^n, x_0 \hat{-} \xi \in \mathbb{R}^t$, where $t = \operatorname{lcm}(m, n)$. Then

$$\|\xi\|_{\mathcal{V}}^2 = \|x_0\|_{\mathcal{V}}^2 + \|x_0 \hat{-} \xi\|_{\mathcal{V}}^2. \quad (216)$$

Figure 6 depicts the relationship of above vectors.

- (ii) Let $\xi \in \mathbb{R}^m$, and the projection of ξ on \mathbb{R}^n be $x_0 = \pi_n^m(\xi)$. Then $\xi - x_0$ is perpendicular to \mathbb{R}^n . That is, for any $z \in \mathbb{R}^n$ we have

$$\langle (\xi \hat{-} x_0), (z - x_0) \rangle_{\mathcal{V}} = 0. \quad (217)$$

Moreover,


$$\|\xi \hat{-} z\|_{\mathcal{V}}^2 = \|\xi \hat{-} x_0\|_{\mathcal{V}}^2 + \|z \hat{-} x_0\|_{\mathcal{V}}^2. \quad (218)$$

Figure 7 depicts the relationship of the vectors.

Proof. We prove (ii) only. The proof of (i) is similar.

Embed \mathbb{R}^n into \mathbb{R}^t by $x \mapsto x \otimes \mathbf{1}_{t/n}$. Using Proposition 9.3, we have

$$(\xi \hat{-} x_0) \perp (\mathbb{R}^n \otimes \mathbf{1}_{t/n}).$$

Fig. 7. Projection is orthogonal to \mathbb{R}^n

Then we have

$$(\xi \hat{-} x_0) \perp (z - x_0 \otimes \mathbf{1}_{t/n}).$$

Using Pythagoras' Theorem on \mathbb{R}^t , we have

$$\|\xi \hat{-} z\|_{\mathcal{V}}^2 = \|\xi \hat{-} x_0\|_{\mathcal{V}}^2 + \|(z - x_0) \otimes \mathbf{1}_{t/n}\|_{\mathcal{V}}^2.$$

Then Proposition 9.3 yields (218) immediately. \square

Example 9.15: Recall Example 9.13, a straightforward computation shows that

$$\begin{aligned} \|\xi\|_{\mathcal{V}}^2 &= \frac{1}{4} (\xi_1^2 + \xi_2^2 + \xi_3^2 + \xi_4^2); \\ \|x_0\|_{\mathcal{V}}^2 &= \frac{1}{6} (1.25\xi_1^2 + 1.25\xi_2^2 + 1.25\xi_3^2 + 1.25\xi_4^2 \\ &\quad + 0.5\xi_1\xi_2 + 0.5\xi_3\xi_4). \end{aligned}$$

$$\begin{aligned} \|x_0 \hat{-} \xi\|_{\mathcal{V}}^2 &= \frac{1}{12} (0.5\xi_1^2 + 0.5\xi_2^2 + 0.5\xi_3^2 \\ &\quad + 0.5\xi_4^2 + \xi_1\xi_2 + \xi_3\xi_4). \end{aligned}$$

Using these equalities, (216) follows immediately.

Using projection and Pythagoras' Theorem, more geometric structure over \mathbb{R}^∞ can be explored.

Example 9.16:

- (i) The relationship between \mathbb{R}^m and \mathbb{R}^n ($m \neq n$)?

Assume $s = \gcd(m, n)$ is the great common divisor. Then it is easy to see that under the topology \mathcal{T}_d , some points of \mathbb{R}^m and \mathbb{R}^n are glued together, when they are of zero distance. Moreover,

$$d(x, y) = 0, \quad x \in \mathbb{R}^m, y \in \mathbb{R}^n,$$

if and only if, there exists $z \in \mathbb{R}^s$ such that $x = z \otimes \mathbf{1}_{m/s}$ and $y = z \otimes \mathbf{1}_{n/s}$. Roughly speaking, \mathbb{R}^m intersects \mathbb{R}^n at a subspace $Z \sim \mathbb{R}^s$. Figure 8 shows this.



- (ii) The sphere in \mathbb{R}^∞ :

Let $c_0 \in \mathbb{R}^\infty$ and $r > 0$. The sphere in \mathbb{R}^∞ , centering at c_0 with radius r , is defined by

$$S_r(c_0) := \{x \in \mathbb{R}^\infty \mid d_{\mathcal{V}}(x, c_0) = r\}. \quad (219)$$

Assume $c_0 \in \mathbb{R}^m$. Now consider \mathbb{R}^n . We project c_0 to \mathbb{R}^n , say, $\pi_n^m(c_0) = c_n$. Denote $h_n := c_0 - c_n$. Assume

$$\|h_n\|_{\mathcal{V}} \leq r. \quad (220)$$

Fig. 8. Intersection of \mathbb{R}^m and \mathbb{R}^n Fig. 9. A Sphere on \mathbb{R}^∞

Then a sphere can be constructed as

$$S_n := \left\{ x \in \mathbb{R}^n \mid d(x, c_n) = \sqrt{r^2 - \|h_n\|_{\mathcal{V}}^2} \right\}. \quad (221)$$

We conclude that the sphere $S_r(c_0)$ in \mathbb{R}^∞ consists of all spheres $S_n \in \mathbb{R}^n$, where the distance of \mathbb{R}^n with c_0 satisfies (220) and S_n is constructed by (221). Figure 9 shows this.

X. DIFFERENTIABLE STRUCTURE ON \mathbb{R}^∞

Since each component space $\mathbb{R}^n \subset \mathbb{R}^\infty$ has its differentiable structure, which provides a differentiable structure for \mathbb{R}^∞ .

Definition 10.1:

- (i) A hyper vector space is said to be differentiable, if all its component subspaces are differentiable.
(ii) Let $\Theta \subset \mathbb{Z}_+$, with (Θ, \prec) be a sublattice of MD-1. Then

$$\mathbb{R}^\Theta := \bigcup_{\theta \in \Theta} \mathbb{R}^\theta \subset \mathbb{R}^\infty,$$

as a sub hyper vector space with the inherited topological structure from \mathbb{R}^∞ and the differentiable structure from \mathbb{R}^θ , where $\theta \in \Theta$, is called an Euclidean hyper manifold.

Remark 10.2:

- (i) It follows from the definition that the differentiability is component-wise defined. Hence \mathbb{R}^∞ is obviously differentiable. Then so is \mathbb{R}^Θ .

- (ii) Unlike the differentiability, the topological structure of an Euclidean hyper manifold is globally defined.

In the following, the major objects of differential geometry will be defined over Euclidean hyper manifold.

A. Smooth Functions

Definition 10.3: A continuous function $h : \mathbb{R}^\Theta \rightarrow \mathbb{R}$ is said to be C^r (or C^∞ , or C^ω), Denoted the set of C^r (or C^∞ , or C^ω , which means the set of analytic functions.) functions by $C^r(\mathbb{R}^\Theta)$ (or $C^\infty(\mathbb{R}^\Theta)$, or $C^\omega(\mathbb{R}^\Theta)$), if for each $\theta \in \Theta$ the restriction $f|_{\mathbb{R}^\theta}$ is C^r (or C^∞ , or C^ω).

Example 10.4:

- (i) Define $f : \mathbb{R}^\infty \rightarrow \mathbb{R}$ by

$$f|_{\mathbb{R}^n}(x) = \sum_{i=1}^n |x_i|, \quad x \in \mathbb{R}^n, n \in \mathbb{Z}_+.$$

Then it is obvious that $f \in C^0(\mathbb{R}^\infty)$.

- (ii) Define $f : \mathbb{R}^\infty \rightarrow \mathbb{R}$ by

$$f(x) = \|x\|_V^2.$$

Then

$$f|_{\mathbb{R}^n}(x) = \frac{1}{n} \sum_{i=1}^n x_i^2, \quad x \in \mathbb{R}^n, n \in \mathbb{Z}_+.$$

It is clear that $f \in C^\omega(\mathbb{R}^\infty)$.

Remark 10.5: A necessary condition for h to be continuous is

$$h(x) = h(y), \quad x \leftrightarrow y. \quad (222)$$

An alternative definition for smooth functions is as follows.

Definition 10.6: A function $h : \mathbb{R}^\Theta \rightarrow \mathbb{R}$ is said to be C^r (or C^∞ , or C^ω), if

- (i) (222) holds.
(ii) For each $\theta \in \Theta$ the restriction $f|_{\mathbb{R}^\theta}$ is C^r (or C^∞ , or C^ω).

Since (222) ensures the continuity among different component spaces, the Definition 10.6 is obviously equivalent to Definition 10.3. Hereafter, for statement ease, we allow $r = \infty$ or $r = \omega$.

In general, such smooth functions are hardly constructed. Particularly, in further discussions (about vector fields etc.) and from application point of view, we may weaken the requirement.

Definition 10.7: Assume Θ has a lattice structure and $\Xi \subset \Theta$ is a filter. A function $h : \mathbb{R}^\Theta \rightarrow \mathbb{R}$ is said to be C^r w.r.t. Ξ , denoted by $h \in C_\Xi^r(\Theta)$, if

- (i) (222) holds for $x, y \in \mathbb{R}^\Xi = \bigcup_{\xi \in \Xi} \mathbb{R}^\xi$.
(ii) For each $\theta \in \Theta$ the restriction $f|_{\mathbb{R}^\theta}$ is C^r .

We need a lemma for further construction.

Lemma 10.8: Let $u, v, w \in \mathbb{Z}_+$, where $u|v$ and $v|w$. Then

$$\Pi_u^w = \Pi_v^w \Pi_u^v. \quad (223)$$

Proof. Set $s = v/u$ and $t = w/v$. Then

$$\Pi_v^w = \frac{1}{t} (I_v \otimes \mathbf{1}_t^T) (I_w \otimes \mathbf{1}) = \frac{1}{t} (I_v \otimes \mathbf{1}_t^T).$$

Similarly,

$$\Pi_u^v = \frac{1}{s} (I_u \otimes \mathbf{1}_s^T).$$

Then

$$\begin{aligned} \Pi_v^w \Pi_u^v &= \frac{1}{st} \{[(I_u \otimes \mathbf{1}_s^T) I_r] \otimes \mathbf{1}_t^T\} \\ &= \frac{1}{st} (I_u \otimes \mathbf{1}_{st}^T) = \Pi_u^w. \end{aligned}$$

□

The following example shows how to construct a C_Ξ^r function from a component space to whole hyper-manifold.

Example 10.9: Let $h(x) \in C^r(\mathbb{R}^\theta)$. Define Define

$$\Xi = \{n\theta \mid n \in \mathbb{Z}_+\}.$$

Then it is clear that (Ξ, \prec) is a sublattice of (\mathbb{Z}_+, \prec) , and it is a filter. Define

$$H(z) := \begin{cases} h(\Pi_\theta^m(z)), & z \in \mathbb{R}^m, m \in \Xi, \\ 0, & m \notin \Xi. \end{cases} \quad (224)$$

We claim that $H(z) \in C_\Xi^r(\mathbb{R}^\Theta)$. Condition (ii) of Definition 10.7 is obviously true. We have only to prove (i). Let $x \in \mathbb{R}^{r\theta}$, $y \in \mathbb{R}^{s\theta}$ and $x \leftrightarrow y$. Denote $r \wedge s = t$, then there exists $z \in \mathbb{R}^t$ such that $x = z \otimes \mathbf{1}_{r/t}$ and $y = z \otimes \mathbf{1}_{s/t}$.

Set $\Pi_n^{tn}(z) := z_0$. Using Lemma 10.8, we have

$$\begin{aligned} H(x) &= h(\Pi_n^{rn}x) = h(\Pi_n^{tn} \Pi_{tn}^r(x)) \\ &= h(\Pi_n^{tn}(z)) = h(z_0). \end{aligned}$$

Similarly,

$$H(y) = h(z_0).$$

That is, (i) follows.

Note that instead of (224) we can also define

$$H(z) := h(\Pi_\theta^m(z)). \quad (225)$$

But we can still only prove (i) for $x, y \in \mathbb{R}^\Xi$, where $\Xi = \{kn \mid k \in \mathbb{Z}_+\}$.

B. Vector Fields and Covector Fields

Definition 10.10: Consider \mathbb{R}^Θ .

- (i) The tangent space of \mathbb{R}^Θ is defined by

$$T(\mathbb{R}^\Theta) = \bigcup_{\theta \in \Theta} T(\mathbb{R}^\theta). \quad (226)$$

- (ii) A vector field $f \in V^r(\mathbb{R}^\Theta)$ if it satisfied

$$f(x) \leftrightarrow f(y), \quad x \leftrightarrow y. \quad (227)$$

$$f|_{\mathbb{R}^\theta} \in V^r(\mathbb{R}^\theta), \quad \theta \in \Theta. \quad (228)$$

Example 10.11: Let $f(x) \in V^r(\mathbb{R}^n)$ and $n \in \Theta$. Then the smallest filter containing n is

$$\Xi := \{kn \mid k \in \mathbb{Z}_+\}.$$

It is easy to extend $f(x)$ to a vector field on $V_{\Xi}^r(\mathbb{R}^\Theta)$. To this end, we define

$$F(z) := \begin{cases} \Pi_{rn}^n f(\Pi_n^m(z)), & m = kn, \\ 0, & m \neq kn, z \in \mathbb{R}^m. \end{cases} \quad (229)$$

It is ready to verify that the condition (ii) of Definition 10.7 is satisfied. We prove condition (i) is also satisfied for $x, y \in \mathbb{R}^\Xi$. Let $x \in \mathbb{R}^{r\theta}$, $y \in \mathbb{R}^{s\theta}$ and $x \leftrightarrow y$. Denote $r \wedge s = t$, then there exists $z \in \mathbb{R}^t$ such that $x = z \otimes \mathbf{1}_{r/t}$ and $y = z \otimes \mathbf{1}_{s/t}$. Then

$$\begin{aligned} F(x) &= \Pi_{rn}^n f(\Pi_n^{tn} \Pi_{tn}^r(x)) \\ &= \Pi_{rn}^n f(\Pi_n^{tn}(z)) \\ &= \Pi_{rn}^n f(z_0) \\ &= (I_n \otimes \mathbf{1}_r) f(z_0) \\ &= f(z_0) \otimes I_r. \end{aligned}$$

Similarly, we have

$$F(y) = f(z_0) \otimes I_s.$$

Hence

$$F(x) \leftrightarrow F(y).$$

Definition 10.12: Consider \mathbb{R}^Θ .

(i) The cotangent space of \mathbb{R}^Θ is defined by

$$T^*(\mathbb{R}^\Theta) = \bigcup_{\theta \in \Theta} T^*(\mathbb{R}^\theta). \quad (230)$$

(ii) A co-vector field $\omega \in V^{*r}(\mathbb{R}^\Theta)$ if it satisfied

$$\omega(x) \leftrightarrow \omega(y), \quad x \leftrightarrow y. \quad (231)$$

$$\omega|_{\mathbb{R}^\theta} \in V^{*r}(\mathbb{R}^\theta), \quad \theta \in \Theta. \quad (232)$$

Similarly to Example 10.11, we are also able to extend a co-vector field $\omega(x) \in V^{*r}(\mathbb{R}^n)$ to $V_{\Xi}^{*r}(\mathbb{R}^\Theta)$.

C. Distribution and Co-Distribution

Definition 10.13: Consider \mathbb{R}^Θ . A distribution (co-distribution) is a rule, which assign for each $x \in \mathbb{R}^\Theta$ a subspace $D_x \subset T_x(\mathbb{R}^\Theta)$ ($D_x^* \subset T_x^*(\mathbb{R}^\Theta)$), satisfying

- (i) For each $n \in \Theta$, $D|_{\mathbb{R}^n}$ is a distribution over $T(\mathbb{R}^n)$ ($D^*|_{\mathbb{R}^n}$ is a co-distribution over $T^*(\mathbb{R}^n)$);
- (ii) Assume $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, $m, n \in \Theta$, and $x \leftrightarrow y$, then $t = m \wedge n \in \Theta$ and there exists $z \in \mathbb{R}^t$ such that $x = z \otimes \mathbf{1}_{m/t}$ and $y = z \otimes \mathbf{1}_{n/t}$, and then

$$\begin{aligned} D_x &= D_z \otimes \mathbf{1}_{m/t}, & D_y &= D_z \otimes \mathbf{1}_{n/t}. \\ (D_x^* &= D_z^* \otimes \mathbf{1}_{m/t}^T, & D_y^* &= D_z^* \otimes \mathbf{1}_{n/t}^T.) \end{aligned} \quad (233)$$

- (iii) A distribution D (co-distribution D^*) is said to be C^r if it is generated by a set of C^r vector fields as

$$\begin{aligned} D(x) &= \sum_{i=1}^s c_i(x) f_i(x), \\ (D^*(x) &= \sum_{i=1}^s c_i(x) \omega_i(x),) \end{aligned} \quad (234)$$

where $f_i(x) \in V^r(\mathbb{R}^\Theta)$ ($\omega_i(x) \in V^{*r}(\mathbb{R}^\Theta)$) are C^r vector fields (co-vector fields) and the coefficients $c_i(x) \in C^r(\mathbb{R}^\Theta)$ are C^r functions.

- (iv) Let Ξ be a filter of Θ , if $D(x)$ ($D^*(x)$) is only C^r over $x \in \mathbb{R}^\Xi$, $D(x)$ ($D^*(x)$) is said to be a C^r distribution (co-distribution) w.r.t. Ξ , briefly, C_{Ξ}^r (C_{Ξ}^{*r}).

D. Tensor Fields

Definition 10.14: Consider \mathbb{R}^Θ . A tensor field with covariant order r and contra-variant order s , denoted by $\mathbf{t} \in \mathbf{T}_s^r(\mathbb{R}^\Theta)$, is a rule, which assign for each $x \in \mathbb{R}^\Theta$ a set of multi-linear mappings, satisfying

- (i) For each $n \in \Theta$, for $\mathbf{t}|_{\mathbb{R}^n}$ is a tensor field

$$\mathbf{t}|_{\mathbb{R}^n} \in \mathbf{T}_s^r(\mathbb{R}^n).$$

- (ii) Assume $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, $m, n \in \Theta$. Assume $x \leftrightarrow y$, then $t = m \wedge n \in \Theta$ and there exists $z \in \mathbb{R}^t$ such that $x = z \otimes \mathbf{1}_{m/t}$ and $y = z \otimes \mathbf{1}_{n/t}$, and then

$$\begin{aligned} M^{\mathbf{t}}(x) &= \underbrace{\mathcal{E}_{m/t}^T \otimes \cdots \otimes \mathcal{E}_{m/t}^T}_s \otimes M_z^{\mathbf{t}} \otimes \underbrace{\mathcal{E}_{n/t} \otimes \cdots \otimes \mathcal{E}_{n/t}}_r, \\ M^{\mathbf{t}}(y) &= \underbrace{\mathcal{E}_{n/t}^T \otimes \cdots \otimes \mathcal{E}_{n/t}^T}_s \otimes M_z^{\mathbf{t}} \otimes \underbrace{\mathcal{E}_{m/t} \otimes \cdots \otimes \mathcal{E}_{m/t}}_r, \end{aligned} \quad (235)$$

where $M^{\mathbf{t}}(p)$, $p = x, z, y$ are the structure matrix of \mathbf{t} at x, y, z respectively.

- (iii) A tensor field \mathbf{t} is said to be C^r if the structure matrix of \mathbf{t} is overall continuous, and within each $n \in \Theta$, $\mathbf{t}|_{\mathbb{R}^n}$ is C^r .
- (iv) Let Ξ be a filter of Θ , if the structure matrix of \mathbf{t} is continuous within Ξ (under the subspace topology),




Fig. 10. Hyper Manifold

and within each $n \in \Theta$, $\mathfrak{t}_{\mathbb{R}^n}$ is C^r , denoted by $\mathfrak{t} \in \mathbf{T}_{s \in \mathbb{R}^\Theta}^r$.

(We refer to [19] for detailed discussion on the differential structure over hyper-manifolds within an essentially the same topological space.)

XI. HYPER MANIFOLD

Definition 11.1: Let \mathbf{M} be a second countable topological space with a partition

$$\mathbf{M} = \bigcup_{\theta \in \Theta} \mathbf{M}_\theta,$$

where each \mathbf{M}_θ is a Hausdorff space (w.r.t. the inherited topology).

Assume

$$\mathcal{U} := \{\mathcal{U}_\lambda, \mid \lambda \in \Lambda\}$$

is an open cover of \mathbf{M} , such that for each $U \in \mathcal{U}$ there exists a mapping $\pi : U \rightarrow U^{\mathbb{R}}$, where $U^{\mathbb{R}} \subset \mathbb{R}^\Theta$ is an open set, and π is a homeomorphism.

Denote by $U_\theta = U \cap \mathbf{M}_\theta$, $\theta \in \Theta$. Let $\Psi : U_\theta \rightarrow U_\theta^{\mathbb{R}}$ with $\Psi : U_\theta \rightarrow U_\theta^{\mathbb{R}} \subset \mathbb{R}^\Theta$, and $\Phi : V_\theta \rightarrow V_\theta^{\mathbb{R}}$ with $\Phi : V_\theta \rightarrow V_\theta^{\mathbb{R}} \subset \mathbb{R}^\Theta$, and $U_\theta \cap V_\theta \neq \emptyset$. Then

$$\Phi \circ \Psi^{-1} : \Psi(U_\theta \cap V_\theta) \rightarrow \Phi(U_\theta \cap V_\theta) \quad (236)$$

is C^r .

Then \mathbf{M} is called a C^r hyper-manifold.

Each $\mathcal{U}_\lambda \in \mathcal{U}$ is called a coordinate chart.

Figure 10 depicts a hyper-manifold.

All the differentiable objects of \mathbf{M} can be defined over each coordinate chart U (or V) in such a way, that the image of the objects over $\Psi(U)$ are properly defined. Then it is also properly defined on each U , and hence on \mathbf{M} .

Example 11.2:

- (i) Consider \mathcal{M} . Let $\Theta = \{m \times n \mid m, n \in \mathbb{Z}_+\} \subset Z_+$. Define a mapping $\Psi : \mathcal{M} \rightarrow \mathbb{R}^\Theta$ as

$$\Psi : A \mapsto V_c(A), \quad (237)$$

then \mathbb{R}^Θ can be considered as a global coordinate chart of \mathcal{M} . Hence \mathcal{M} is a hyper-manifold.

- (ii) Consider $\overline{\mathcal{M}}$. Let $\Theta = \{m \times n + 1 \mid m, n \in \mathbb{Z}_+\} \subset Z_+$. Define a mapping $\Psi : \overline{\mathcal{M}} \rightarrow \mathbb{R}^\Theta$ as

$$\Psi : A = aI_{m \times n} + A_0 \mapsto (a, V_c(A_0)), \quad (238)$$

then \mathbb{R}^Θ can be considered as a global coordinate chart of $\overline{\mathcal{M}}$. Hence $\overline{\mathcal{M}}$ is a hyper-manifold.

XII. LIE ALGEBRA AND LIE GROUP OF NON-SQUARE MATRICES

A. Lie Algebra of Non-square Matrices

Definition 12.1: [22] Consider $\mathcal{M}_{m \times n}$. The Lie bracket is defined by

$$[A, B]_{\times} := A \times B - B \times A. \quad (239)$$

Proposition 12.2: [22] $gl(m \times n, \mathbb{R}) := (\mathcal{M}_{m \times n}, [\cdot, \cdot]_{\times})$ is a Lie algebra.

Note that when $m = n$, $gl(m \times n, \mathbb{R}) = gl(n, \mathbb{R})$. Hence $gl(m \times n, \mathbb{R})$ is a natural generalization of the classical general linear algebra from square case to non-square case.

To construct a Lie group $GL(m \times n, \mathbb{R})$ with $gl(m \times n, \mathbb{R})$ as its Lie algebra, we need some preparations.

Definition 12.3: [45]

- (i) Let $\mathfrak{g} = (G, [\cdot, \cdot])$ be a Lie algebra, and $H \subset G$ be a subspace. If

$$[h_1, h_2] \in H, \quad h_1, h_2 \in H, \quad (240)$$

then $\mathfrak{h} = (H, [\cdot, \cdot])$ is called a Lie sub-algebra of \mathfrak{g} .

- (ii) Let \mathfrak{h} be a Lie sub-algebra of \mathfrak{g} . If

$$[h, g] \in H, \quad h \in H, g \in G, \quad (241)$$

then \mathfrak{h} is called an ideal of \mathfrak{g} .

It follows from the definition immediately that

Proposition 12.4: Consider $gl(m \times n, \mathbb{R})$.

- (i) Set

$$Q_{m \times n} := \{A \in \mathcal{M}_{m \times n} \mid Tr^\Pi(A) = 0\}, \quad (242)$$

then $Q_{m \times n}$ is an ideal of $gl(m \times n, \mathbb{R})$.

- (ii) Set

$$Z_{m \times n} := \{A \in \mathcal{M}_{m \times n} \mid \Pi A = rI_m, r \in \mathbb{R}\}, \quad (243)$$

then $Z_{m \times n}$ is an ideal of $gl(m \times n, \mathbb{R})$.

Example 12.5: Consider $gl(2 \times 3, \mathbb{R})$. Solving

$$A\Psi_{3 \times 2} = \lambda I_2$$

yields

$$Z_{m \times n} = \left\{ \sqrt{6} \begin{bmatrix} 1.5 & -2 & 1 \\ 1 & -2 & 1.5 \end{bmatrix} \lambda + (1, -2, 1) \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \mid \lambda, \alpha, \beta \in \mathbb{R} \right\}. \quad (244)$$

In general, when $m < n$, $Z_{m \times n}$ is an $(n - m)m + 1$ dimensional subspace.

Definition 12.6: [45]

- (i) Let $\mathfrak{h} = (H, [\cdot, \cdot])$ and $\mathfrak{g} = (G, [\cdot, \cdot])$ be two Lie algebras. If there exists a linear mapping $\pi : H \rightarrow G$ such that

$$\pi[h_1, h_2] = [\pi(h_1), \pi(h_2)], \quad h_1, h_2 \in H, \quad (245)$$

then \mathfrak{h} and \mathfrak{g} are homomorphic and π is a homomorphism.

- (ii) $\pi : H \rightarrow G$ is an one-to-one and onto mapping, and both π and π^{-1} are homomorphisms, then \mathfrak{h} and \mathfrak{g} are isomorphic and π is an isomorphism.

Proposition 12.7: Consider $gl(m \times n, \mathbb{R})$ and $gl(m_0 \times n_0, \mathbb{R})$

Assume $m_0 | m$, $n_0 | n$, and denote $m = m_0 a$ and $n = n_0 b$. Define $\pi : gl(m_0 \times n_0, \mathbb{R}) \rightarrow gl(m \times n, \mathbb{R})$ by

$$\pi(A) := A \otimes \mathcal{E}_{a \times b}, \quad A \in gl(m_0 \times n_0). \quad (246)$$

Then π is a homomorphism.

Proof. We have to show that

$$[\pi(A_0), \pi(B_0)] = \pi([A_0, B_0]), \quad A_0, B_0 \in gl(m_0 \times n_0, \mathbb{R}).$$

It suffices to show that setting $A = A_0 \otimes \mathcal{E}_{a \times b}$ and $B = B_0 \otimes \mathcal{E}_{a \times b}$, then

$$A \times B = \pi(A_0 \times B_0), \quad A_0, B_0 \in gl(m_0 \times n_0, \mathbb{R}). \quad (247)$$

Let $t_0 = m_0 \vee n_0$ and $\xi = m \vee n$. Then

$$\begin{aligned} \pi(A_0 \times B_0) &= \pi \left[(A_0 \otimes \mathcal{E}_{t_0/n_0}^T)(B_0 \otimes \mathcal{E}_{t_0/m_0}) \right] \\ &= \left[(A_0 \otimes \mathcal{E}_{t_0/n_0}^T)(B_0 \otimes \mathcal{E}_{t_0/m_0}) \right] \otimes \mathcal{E}_{a \times b} \\ &= \left[(A_0 \otimes \mathcal{E}_{t_0/n_0}^T)(B_0 \otimes \mathcal{E}_{t_0/m_0}) \right] \otimes (\mathcal{E}_a E_b^T) \\ &= (A_0 \otimes \mathcal{E}_{a \times t_0/n_0})(B_0 \otimes \mathcal{E}_{t_0/m_0 \times b}). \end{aligned}$$

And

$$\begin{aligned} A \times B &= (A_0 \otimes \mathcal{E}_{a \times b}) \times (B_0 \otimes \mathcal{E}_{a \times b}) \\ &= (A_0 \otimes \mathcal{E}_{a \times b} \otimes \mathcal{E}_{\xi/n}^T)(B_0 \otimes \mathcal{E}_{a \times b} \otimes \mathcal{E}_{\xi/m}) \\ &= (A_0 \otimes \mathcal{E}_{a \times b \xi/n})(B_0 \otimes \mathcal{E}_{a \xi/m \times b}) \\ &= (A_0 \otimes \mathcal{E}_{a \times \xi/n_0})(B_0 \otimes \mathcal{E}_{\xi/m_0 \times b}) \\ &= (A_0 \otimes \mathcal{E}_{a \times t_0/n_0})(B_0 \otimes \mathcal{E}_{t_0/m_0 \times b}). \end{aligned}$$

The conclusion follows. \square

Corollary 12.8: Assume $gl(m \times n, \mathbb{R})$, $gl(m_0 \times n_0, \mathbb{R})$ and the mapping π are as in Proposition 12.7. Then $Im(\pi)$ is a Lie sub-algebra of $gl(m \times n, \mathbb{R})$.

Proof. Note that π is a homomorphism. Moreover, it is obvious that π is one to one, then the conclusion is obvious. \square

B. Lie Group of Non-square Matrices

Define $I_{m \times n}$ as a formal identity for the algebra $(\mathcal{M}_{m \times n} \times)$, which satisfies

$$\begin{aligned} I_{m \times n} \times I_{m \times n} &= I_{m \times n} \\ I_{m \times n} \times A &= A \times I_{m \times n} = A, \quad \forall A \in \mathcal{M}_{m \times n}. \end{aligned} \quad (248)$$

Then we have an extended matrix-ring as

$$\overline{\mathcal{M}}_{m \times n} := \{aI_{m \times n} + A_0 \mid a \in \mathbb{R}, A_0 \in \mathcal{M}_{m \times n}\}, \quad (249)$$

with

$$\begin{aligned} (aI_{m \times n} + A_0) + (bI_{m \times n} + B_0) &:= (a + b)I_{m \times n} \\ &\quad + (A_0 + B_0), \\ (aI_{m \times n} + A_0) \times (bI_{m \times n} + B_0) &:= (ab)I_{m \times n} + aB_0 \\ &\quad + bA_0 + A_0 \times B_0, \quad a, b \in \mathbb{R}, A, B \in \mathcal{M}_{m \times n}. \end{aligned} \quad (250)$$

Note that where $m \neq n$ the objects $\{aI_{m \times n} + A_0\}$ of an extended matrix-ring are not matrices unless $a = 0$.

Let $A \in \mathcal{M}_{m \times n}$. Then $\Pi_A = A\Psi_{n \times m}$, where $\Psi_{n \times m}$ defined by (34), is called the restricted (square) form of A , because

$$A \times x = \Pi_A x, \quad \forall x \in \mathbb{R}^m.$$

Define

$$A^{<k>} = \begin{cases} I_{m \times n}, & k = 0, \\ \underbrace{A \times \cdots \times A}_k, & k > 0. \end{cases} \quad (251)$$

In addition, let $p(x) = x^n + c_{n-1}x^{n-1} + \cdots + c_1x + c_0$, we define

$$\begin{aligned} p \langle A \rangle &:= A^{<n>} + c_{n-1}A^{<n-1>} + \cdots + c_1A \\ &\quad + c_0I_{m \times n} \in \overline{\mathcal{M}}_{m \times n}. \end{aligned} \quad (252)$$

Then we have the following result.

Theorem 12.9: (Generalized Cayley-Hamilton theorem)[22]

Let $p(x) = x^m + c_{m-1}x^{m-1} + \cdots + c_1x + c_0$ be the characteristic function of Π_A , which is also called the characteristic function of A . Then

$$A^{<m+1>} + c_{m-1}A^{<m>} + \cdots + c_1A^{<2>} + c_0A = 0. \quad (253)$$

Remark 12.10:

- (i) When $n < m$ we can take the characteristic function of Π_{A^T} as the characteristic function of A . This alternative choice can reduce the degree of the characteristic function.

(ii) Formally, we may rewritten (253) into the form as

$$A^{<m>} + c_{m-1}A^{<m-1>} + \cdots + c_1A + c_0I_{m \times n}^A = 0. \quad (254)$$

Then if $c_0 \neq 0$ we can define

$$I_{m \times n}^A := -\frac{1}{c_0} [A^{<m>} + c_{m-1}A^{<m-1>} + \cdots + c_1A]. \quad (255)$$

According to Generalized Cayley-Hamilton theorem, it is clear that

$$A \times I_{m \times n}^A = I_{m \times n}^A \times A = A. \quad (256)$$

Moreover, for any polynomial $p < A >$, we also have

$$(p < A >) \times I_{m \times n}^A = I_{m \times n}^A \times (p < A >) = p < A >. \quad (257)$$

Then A is said to be relatively invertible and define its relative inverse by

$$A^{<-1>} = -\frac{1}{c_0} [A^{<m-1>} + c_{m-1}A^{<m-2>} + \cdots + c_1I_{m \times n}^A]. \quad (258)$$

It follows that

$$A \times A^{<-1>} = A^{<-1>} \times A = I_{m \times n}^A. \quad (259)$$

It seems that if A is relatively invertible, then $I_{m \times n}^A|_{p < A >}$ is $I_{m \times n}^A$. Unfortunately, $I_{m \times n}^A$ depends on A . Moreover, it is a matrix while $I_{m \times n}$ is not.

Definition 12.11: $A \in \overline{\mathcal{M}}_{m \times n}$ is invertible, if there exists $B \in \overline{\mathcal{M}}_{m \times n}$, such that

$$A \times B = B \times A = I_{m \times n}.$$

Remark 12.12:

- (i) Assume $A = aI_{m \times n} + A_0$, where $A_0 \in \mathcal{M}_{m \times n}$. It is obvious that a necessary condition for A being invertible is $a \neq 0$.
- (ii) If $A = aI_{m \times n} + A_0$ is invertible and $A^{<-1>} = B$. Then $A' = \frac{1}{a}A$ is also invertible and $(A')^{<-1>} = aB$. Hence, to find the inverse of A , we can, without loss of generality, assume $a = 1$.
- (iii) Assume $A = I_{m \times n} + A_0$, A is invertible, if and only if, there exists $B = I_{m \times n} + B_0$, where $B_0 \in \mathcal{M}_{m \times n}$ such that

$$\begin{aligned} A \times B &= (I_{m \times n} + A_0) \times (I_{m \times n} + B_0) \\ &= I_{m \times n} + A_0 + B_0 + A_0 \times B_0. \end{aligned} \quad (260)$$

- (iv) Assume $m = n$. Then $\overline{\mathcal{M}}_{n \times n} = \mathcal{M}_{n \times n}$ because now $I_{n \times n} = I_n \in \mathcal{M}_{n \times n}$. Then it is easily verifiable that $A \in \mathcal{M}_{n \times n}$ is invertible (under the classical definition), if and only if, (260) holds. Hence Definition 12.11 can

be considered as an extension of the classical invertibility to non-square case.

(v) The $GL(m \times n, \mathbb{R})$ has been defined by [22] as

$$GL(m \times n, \mathbb{R}) := \{A = I_{m \times n} + A_0 \mid A_0 \in \mathcal{M}_{m \times n} \text{ and } A \text{ is invertible}\}. \quad (261)$$

Then it was proved in [22] that $GL(m \times n, \mathbb{R})$ is a Lie group, which has $gl(m \times n, \mathbb{R})$ as its Lie algebra.

Next, we consider when $A = I_{m \times n} + A_0 \in \overline{\mathcal{M}}$ is invertible. Assume the characteristic function of A_0 is

$$p(x) = x^{k-1} + c_{k-1}x^{k-2} + \cdots + c_2x + c_1. \quad (262)$$

Using generalized Cayley-Hamilton theorem, we have

$$A_0^{<k>} + c_{k-1}A_0^{<k-1>} + \cdots + c_2A_0^{<2>} + c_1A_0 = 0.$$

We assume

$$B_0 = x_{k-1}A_0^{<k-1>} + x_{k-2}A_0^{<k-2>} + \cdots + x_2A_0^{<2>} + x_1A_0. \quad (263)$$

Then

$$\begin{aligned} A_0 \times B_0 &= x_{k-1}A_0^{<k>} + x_{k-2}A_0^{<k-1>} + \cdots \\ &+ x_2A_0^{<3>} + x_1A_0^{<2>} \\ &= -x_{k-1}(c_{k-1}A_0^{<k-1>} + \cdots + c_2A_0^{<2>} + c_1A_0) \\ &+ x_{k-2}A_0^{<k-1>} + \cdots + x_2A_0^{<3>} + x_1A_0^{<2>} \end{aligned}$$

Setting

$$A_0 + B_0 + A_0 \times B_0 = 0 \quad (264)$$

and comparing the coefficients yield the following linear system

$$\begin{cases} -c_{k-1}x_{k-1} + x_{k-2} + x_{k-1} = 0 \\ -c_{k-2}x_{k-1} + x_{k-3} + x_{k-2} = 0 \\ \cdots \\ -c_2x_{k-1} + x_1 + x_2 = 0 \\ -c_1x_{k-1} + x_1 + 1 = 0. \end{cases} \quad (265)$$

Solving (265) yields

$$\begin{cases} x_1 = c_1x_{k-1} - 1, \\ x_2 = (c_2 - c_1)x_{k-1} + 1, \\ x_3 = (c_3 - c_2 + c_1)x_{k-1} - 1, \\ \cdots, \\ x_{k-2} = (c_{k-2} - c_{k-3} \pm \cdots + (-1)^{k-1}c_1)x_{k-1} + (-1)^{k-2}, \\ (c_{k-1} - c_{k-2} + c_{k-3} \pm \cdots + (-1)^k c_1 - 1)x_{k-1} \\ + (-1)^{k-1} = 0. \end{cases} \quad (266)$$

Then we have the following result.

Theorem 12.13: Consider $A = I_{m \times n} + A_0$ and assume the characteristic function of A_0 is (262). Then A is invertible, if and only if,

$$c_{k-1} - c_{k-2} + c_{k-3} \pm \cdots + (-1)^{k-2} c_1 - 1 \neq 0. \quad (267)$$

Proof. The previous constructing process is a constructive proof for the sufficiency. As for the necessity, it is reasonable to assume the inverse of A is an analytic function of A_0 . Taking the generalized Cayley-Hamilton theorem into consideration, it is obvious that the form of B_0 , expressed by (263) is a general expression of an analytic function of A_0 . \square

Remark 12.14:

- (i) If (267) holds, (266) with (263) provide the inverse of A immediately.
- (ii) It is easy to see that

$$I_{m \times n} = \exp(0_{m \times n}). \quad (268)$$

Then by continuity, one sees easily that there exists an open neighborhood \mathcal{O} of $0_{m \times n}$, say

$$0_{m \times n} \in \mathcal{O} \subset \mathbb{R}^{mn},$$

such that for each $B \in \mathcal{O}$ the characteristic function

$$p_B(x) = x^{k-1} + c_{k-1}x^{k-2} + \cdots + c_2x + c_1,$$

satisfies

$$|c_{k-1} - c_{k-2} + c_{k-3} \pm \cdots + (-1)^{k-2} c_1| < 1.$$

Hence

$$I_{m \times n} + B \in GL(m \times n, \mathbb{R}), \quad B \in \mathcal{O}.$$

- (iii) By possibly shrinking \mathcal{O} , we can assume

$$\ln \langle I_{m \times n} + B \rangle = \sum_{i=1}^{\infty} (-1)^{i-1} B^{<i>}, \quad B \in \mathcal{O} \quad (269)$$

converges. Then we define

$$GL_0(m \times n, \mathbb{R}) := \{I_{m \times n} + B \mid B \in \mathcal{O}\}. \quad (270)$$

It follows that

$$GL_0(m \times n, \mathbb{R}) \subset \exp(\mathcal{O}), \quad (271)$$

where $\exp(\mathcal{O})$ is an open sub-manifold of $GL(m \times n, \mathbb{R})$, which shows that $GL_0(m \times n, \mathbb{R})$ itself is a sub-Lie group of $GL(m \times n, \mathbb{R})$.

Example 12.15: Consider $A = I_{2 \times 3} + A_0$, where $A_0 = (a_{i,j}) \in \mathcal{M}_{2 \times 3}$. For statement ease, we let $\Psi_{3 \times 2}$ be as in (36). Then we have

(i)

$$\begin{aligned} \Pi_A &= A_0 \Psi_{3 \times 2} = A_0 \begin{bmatrix} 2 & 0 \\ 1 & 1 \\ 0 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 2a_{1,1} + a_{1,2} & a_{1,2} + 2a_{1,3} \\ 2a_{2,1} + a_{2,2} & a_{2,2} + 2a_{2,3} \end{bmatrix}. \end{aligned}$$

The characteristic function of A_0 is

$$p(x) = x^2 + c_2x + c_1,$$

where

$$\begin{aligned} c_2 &= -(2a_{1,1} + a_{1,2} + a_{2,2} + 2a_{2,3}), \\ c_1 &= (2a_{1,1} + a_{1,2})(a_{2,2} + 2a_{2,3}) \\ &\quad - (2a_{2,1} + a_{2,2})(a_{1,2} + 2a_{1,3}). \end{aligned}$$

We conclude that as long as $c_2 - c_1 - 1 \neq 0$, A is invertible. The inverse is

$$B_0 = x_2 A_0^{<2>} + x_1 A_0,$$

where

$$x_1 = \frac{c_2 - 1}{c_1 - c_2 + 1}, \quad x_2 = \frac{1}{c_1 - c_2 + 1}.$$

- (ii) To verify the above result, we calculate the inverse by definition. Assume $B_0 = (b_{i,j}) \in \mathcal{M}_{2 \times 3}$ such that

$$A_0 + B_0 + A_0 \times B_0 = 0. \quad (272)$$

A direct calculation shows that

$$A_0 \times B_0 := C_0 = (c_{i,j}) \in \mathcal{M}_{2 \times 3},$$

where

$$\begin{aligned} c_{1,1} &= 2a_{1,1}b_{1,1} + a_{1,2}b_{1,1} + a_{1,2}b_{2,1} + 2a_{1,3}b_{2,1}, \\ c_{1,2} &= 2a_{1,1}b_{1,2} + a_{1,2}b_{1,2} + a_{1,2}b_{2,2} + 2a_{1,3}b_{2,2}, \\ c_{1,3} &= 2a_{1,1}b_{1,3} + a_{1,2}b_{1,3} + a_{1,2}b_{2,3} + 2a_{1,3}b_{2,3}, \\ c_{2,1} &= 2a_{2,1}b_{1,1} + a_{2,2}b_{1,1} + a_{2,2}b_{2,1} + 2a_{2,3}b_{2,1}, \\ c_{2,2} &= 2a_{2,1}b_{1,2} + a_{2,2}b_{1,2} + a_{2,2}b_{2,2} + 2a_{2,3}b_{2,2}, \\ c_{2,3} &= 2a_{2,1}b_{1,3} + a_{2,2}b_{1,3} + a_{2,2}b_{2,3} + 2a_{2,3}b_{2,3}. \end{aligned}$$

(272) leads to a linear system as

$$\begin{bmatrix} \alpha & 0 & 0 & \beta & 0 & 0 \\ 0 & \alpha & 0 & 0 & \beta & 0 \\ 0 & 0 & \alpha & 0 & 0 & \beta \\ \gamma & 0 & 0 & \delta & 0 & 0 \\ 0 & \gamma & 0 & 0 & \delta & 0 \\ 0 & 0 & \gamma & 0 & 0 & \delta \end{bmatrix} \begin{bmatrix} b_{1,1} \\ b_{1,2} \\ b_{1,3} \\ b_{2,1} \\ b_{2,2} \\ b_{2,3} \end{bmatrix} = \begin{bmatrix} -a_{1,1} \\ -a_{1,2} \\ -a_{1,3} \\ -a_{2,1} \\ -a_{2,2} \\ -a_{2,3} \end{bmatrix}, \quad (273)$$

where

$$\begin{aligned} \alpha &= 2a_{1,1} + a_{1,2} + 1, \\ \beta &= a_{1,2} + 2a_{1,3}, \\ \gamma &= 2a_{2,1} + a_{2,2}, \\ \delta &= 2a_{2,3} + a_{2,2} + 1. \end{aligned}$$

Then it is clear that (273) has unique solution, if and only if,

$$\alpha\delta - \beta\gamma \neq 0.$$

A careful calculation shows that

$$\alpha\delta - \beta\gamma = -(c_2 - c_1 - 1).$$

That is, the condition obtained in (i) is necessary and sufficient.

In general, the following is obvious.

Proposition 12.16: Let

$$\mathcal{T}_{m \times n} := (\{A \in \overline{\mathcal{M}}_{m \times n} \mid A \text{ is invertible}\}, \times). \quad (274)$$

Then $\mathcal{T}_{m \times n}$ is a group.

Definition 12.17: Let $B \in \mathcal{M}_{m \times n}$.

(i)

$$GL(m \times n, \mathbb{R}) := \{A = I_{m \times n} + A_0 \mid A_0 \in \mathcal{M}_{m \times n} \text{ and } A \in \mathcal{T}_{m \times n}\}. \quad (275)$$

called the $m \times n$ general linear group.

(ii) Let $B \in \mathcal{M}_{m \times n}$. Define

$$\exp \langle B \rangle := I_{m \times n} + \sum_{n=1}^{\infty} \frac{1}{n!} B^{\langle n \rangle}, \quad (276)$$

Then

$$\exp \langle B \rangle \in GL(m \times n, \mathbb{R}).$$

Theorem 12.18: [22] $GL(m \times n, \mathbb{R})$ is a Lie group, which has $gl(m \times n)$ as its Lie algebra.

Remark 12.19:

(i)

$$GL(m \times n, \mathbb{R}) \subset \mathcal{T}_{m \times n}.$$

(ii) Define

$$\mathcal{B} := \{ \times_{i=1}^k \exp(B_i) \mid B_i \in \mathcal{M}_{m \times n}, k < \infty \}. \quad (277)$$

Then \mathcal{B} is a group and

$$\mathcal{B} \subset GL(m \times n, \mathbb{R})$$

is a sub-Lie group, which is a path-connected component of $GL(m \times n, \mathbb{R})$.

(iii) A_0 can be considered as the coordinate of $A = I_{m \times n} + A_0$. It is then clear that

$$\dim(GL(m \times n, \mathbb{R})) = mn. \quad (278)$$

(iv) $\overline{\mathcal{M}}_{m \times n}$ is a ring, which is also an $mn + 1$ dimensional vector space. It can also be naturally embedded into \mathbb{R}^{mn+1} by

$$A = aI_{m \times n} + A_0 \mapsto (a, A_0^{1,1}, A_0^{1,2}, \dots, A_0^{m,n})^T \in \mathbb{R}^{mn+1}.$$




Fig. 11. $gl(m \times n, \mathbb{R})$ and $GL(m \times n, \mathbb{R})$

Hence $\overline{\mathcal{M}}_{m \times n}$ can be naturally considered as a topological vector space, which has the classical topology of Euclidian space \mathbb{R}^{mn+1} . Now $gl(m \times n, \mathbb{R})$ can be considered as its mn dimensional subspace, and $GL(m \times n, \mathbb{R})$ is also its $m \times n$ dimensional sub-manifold. When $m \neq n$ $gl(m \times n) \cap GL(m \times n) = \emptyset$. This is depicted in Fig.11.

(v) It is clear that

$$I_{m \times n} = \exp(0_{m \times n}),$$

which is the identity of $GL(m \times n, \mathbb{R})$. So it is not an element in $gl(m \times n, \mathbb{R})$. When we extend it to be the identity of the ring $\overline{\mathcal{M}}$, we have the relations shown in (248).

C. Relationship with General Linear Group/Algebra

As an immediate consequence of Adi's theorem, we have the following theorem [45].

Theorem 12.20: Let \mathfrak{g} be a Lie algebra over \mathbb{R} . Then there exists an integer $n \geq 1$ and an analytic subgroup G of $GL(n, \mathbb{R})$ such that the Lie algebra of G is isomorphic to \mathfrak{g} .

According to Theorem 12.20, there is a Lie sub-algebra of $\mathfrak{g} \subset gl(k, \mathbb{R})$, such that $gl(m \times n, \mathbb{R})$ is isomorphic to $\mathfrak{g} \subset gl(k, \mathbb{R})$. This motivates the following result.

Proposition 12.21: Consider $\square : \mathcal{M}_{m \times n} \rightarrow \mathcal{M}_{t \times t}$, where $t = m \vee n$. Then $\square : gl(m \times n, \mathbb{R}) \rightarrow gl(t, \mathbb{R})$ is a Lie algebra homomorphism. Moreover, $Im(\square)$ is a sub-algebra of $gl(t, \mathbb{R})$.

Proof. Let $A, B \in \mathcal{M}_{m \times n}$. It suffices to show that

$$\square(A \times B) = \square(A)\square(B), \quad A, B \in \mathcal{M}_{m \times n}. \quad (279)$$

$$\begin{aligned}
\Box(A \times B) &= \Box\left((A \otimes \mathcal{E}_{t/n}^T)(B \otimes \mathcal{E}_{t/m})\right) \\
&= \left((A \otimes \mathcal{E}_{t/n}^T)(B \otimes \mathcal{E}_{t/m})\right) \otimes \mathcal{E}_{t/m \times t/n} \\
&= \left((A \otimes \mathcal{E}_{t/n}^T)(B \otimes \mathcal{E}_{t/m})\right) \otimes \left(\mathcal{E}_{t/m} \mathcal{E}_{t/n}^T\right) \\
&= \left((A \otimes \mathcal{E}_{t/n}^T \otimes \mathcal{E}_{t/m})\right) \left(B \otimes \mathcal{E}_{t/m} \otimes \mathcal{E}_{t/n}^T\right) \\
&= \left((A \otimes \mathcal{E}_{t/m \times t/n})\right) \left(B \otimes \mathcal{E}_{t/m \times t/n}\right) \\
&= \Box(A) \Box(B).
\end{aligned}$$

Hence $\Box : gl(m \times n, \mathbb{R}) \rightarrow gl(t, \mathbb{R})$ is a Lie algebra homomorphism. Now it is obvious that \Box is one-to-one, then

$$Im(\Box) = \Box(\mathcal{M}_{m \times n})$$

is a Lie sub-algebra of $gl(t, \mathbb{R})$.

Define

$$GL(Im(\mathcal{M}_{m \times n}), \mathbb{R}) := \left\{ \prod_{i=1}^k \exp(B_i) \mid B_i \in \Box(\mathcal{M}_{m \times n}), k < \infty \right\}. \quad (280)$$

Then

$$GL(Im(\mathcal{M}_{m \times n}), \mathbb{R}) < GL(n, \mathbb{R}) \quad (281)$$

is a Lie sub-group.

Moreover, if we choose

$$GL(m \times n, \mathbb{R}) := \left\{ \times_{i=1}^k \exp(\langle B_i \rangle) \mid B_i \in gl_{m \times n}(\mathbb{R}), k < \infty \right\}, \quad (282)$$

then

$$GL(m \times n, \mathbb{R}) \cong GL(Im(\mathcal{M}_{m \times n}), \mathbb{R}) \quad (283)$$

is a Lie group isomorphism.

XIII. FROM TOPOLOGICAL HYPER GROUP TO HYPER LIE GROUP

A. Topological Hyper Group

Definition 13.1: Assume $\mathcal{G} = (G, *, \mathbf{e})$ is both a hyper group and a topological space. If both the product and the inverse are continuous, then \mathcal{G} is called a topological hyper group.

Example 13.2:

- (i) Consider $\mathcal{T} = \{A \in \mathcal{M}_1 \mid A \text{ is invertible}\}$. Then it is easy to verify that $(\mathcal{T}, \times, \mathbf{e})$ (or $(\mathcal{T}, \times, \mathbf{e})$) is a topological hyper group, where $\mathbf{e} = \{I_n \mid n \in \mathbb{Z}_+\}$.
- (ii) Consider the permutation hyper group \mathbf{S} . By identifying $\sigma \in \mathbf{S}_n$ with its permutation matrix $M_\sigma \in \mathcal{P}_n$ as $\pi_n : \mathbf{S}_n \rightarrow \mathcal{P}_n$. Then we can define $\pi : \mathbf{S} \rightarrow \mathcal{P}$ by

$$\pi|_{\mathbf{S}_n} = \pi_n, \quad n \in \mathbb{Z}_+. \quad (284)$$

It is easy to see that π is a bijective mapping. Then we define $\circ_\ell : \mathbf{S} \times \mathbf{S} \rightarrow \mathbf{S}$ and $\circ_r : \mathbf{S} \times \mathbf{S} \rightarrow \mathbf{S}$ as

$$\begin{aligned}
\sigma \circ_\ell \mu &= \pi^{-1} [M_\sigma \times M_\mu], \\
\sigma \circ_r \mu &= \pi^{-1} [M_\sigma \times M_\mu].
\end{aligned} \quad (285)$$

(iii) Consider $(\mathcal{T}, \times, \mathbf{e})$ (or $(\mathcal{T}, \times, \mathbf{e})$), which is a topological hyper group. As the hyper-subgroup of $(\mathcal{T}, \times, \mathbf{e})$ (or $(\mathcal{T}, \times, \mathbf{e})$), the $(\mathcal{P}, \times, \mathbf{e})$ (or $(\mathcal{P}, \times, \mathbf{e})$) with its inherited topology is a topological hyper group.

(iv) Using the topology of $(\mathcal{P}, \times, \mathbf{e})$ (or $(\mathcal{P}, \times, \mathbf{e})$), $(\mathbf{S}, \circ_\ell, \mathbf{Id})$ (or $(\mathbf{S}, \circ_r, \mathbf{Id})$) is a topological hyper group.

B. Hyper Lie Group

Definition 13.3: Assume $\mathcal{G} = (G, *, \mathbf{e})$ is a topological hyper group and at the same time it is also an analytic hyper-manifold with both the product and the inverse as analytic mappings. Then \mathcal{G} is called a hyper Lie group.

Example 13.4: Consider $\mathcal{G} := (\mathbb{R}^\infty, \hat{+}, \mathbf{e})$, where $\mathbf{e} = \{0_{m \times n} \mid m, n \in \mathbb{Z}_+\}$. Equipped by its Euclidean hyper-manifold structure, \mathcal{G} is a hyper Lie group. \square

XIV. HYPER LINEAR ALGEBRA

A. Matrix Hyper Lie Algebra

Definition 14.1: Consider $\mathcal{M}_0 \subset \mathcal{M}$. Assume there are two binary operators: $+$: $\mathcal{M}_0 \times \mathcal{M}_0 \rightarrow \mathcal{M}_0$, \times : $\mathcal{M}_0 \times \mathcal{M}_0 \rightarrow \mathcal{M}_0$, such that $(\mathcal{M}_0, +)$ is a hyper-vector space; $(\mathcal{M}_0, +, \times)$ is a hyper-ring, then $(\mathcal{M}_0, +, \times)$ is called a matrix hyper-algebra.

Proposition 14.2: Assume $(\mathcal{M}_0, +, \times)$ a matrix hyper-algebra. A Lie bracket is defined by

$$[A, B] := A \times B - B \times A. \quad (286)$$

Then it satisfies the following conditions for Lie algebra.

(i) Skew Symmetry:

$$[A, B] = -[B, A], \quad A, B \in \mathcal{M}_0. \quad (287)$$

(ii) Bi-Linearity:

$$[aA + bB, C] = a[A, C] + b[B, C], \quad A, B, C \in \mathcal{M}_0. \quad (288)$$

(iii) Jacobi Identity:

$$[A, [B, C]] + [B, [C, A]] + [C, [A, B]] = 0. \quad (289)$$

That is, a matrix hyper-ring with a Lie bracket defined by (286) is called a hyper-Lie algebra.

Proof. (287) comes from definition (286) immediately. The distributivity of hyper-ring leads to (288). (289) follows from a straightforward computation, using the linearity. \square

A matrix hyper-algebra over matrices with Lie bracket defined by (286) is called a hyper linear algebra.

Moreover, the following consequence is obvious.

Proposition 14.3: Assume $(\mathcal{M}_0, +, \times)$ is a hyper linear algebra, and the identity set of $(\mathcal{M}_0, +)$ is \mathbf{e} . Then

- (i) each vector space \mathcal{M}_0^e , $e \in \mathbf{e}$, is a general linear algebra;
(ii) if $+$ and \times are consistent w.r.t. the equivalence, the equivalence classes $(\mathcal{M}_0 / \sim, +, \times)$ is a Lie algebra, called the equivalence linear algebra.

It is well known that $(\mathcal{M}_{n \times n}, +, \times)$ with the Lie bracket defined by (286) is the general linear algebra $gl(n, \mathbb{R})$. So the hyper linear algebra is a natural generalization of the classical general linear algebra.

The above argument shows that as long as a matrix hyper-ring exists, it leads to a hyper linear algebra automatically. Hence, we have the following hyper matrix Lie algebras based on the matrix hyper-rings[19]:

Example 14.4:

- (i) Consider $(\mathcal{M}_1, \bar{+}, \times)$, defined in Remark 7.2 equation (181). Define

$$[A, B] := A \times B^- B \times A. \quad (290)$$

Then $(\mathcal{M}_1, \bar{+}, \times)$ with the Lie bracket defined by (290) is a hyper-Lie algebra. Denote this hyper-Lie algebra by $gl_s^1(\mathbb{R})$.

- (ii) Consider $(\mathcal{M}_1, \hat{+}, \circ)$, defined in Remark 7.2 equation (182). Define

$$[A, B] := A \circ B^- B \circ A. \quad (291)$$

Then $(\mathcal{M}_1, \hat{+}, \circ)$ with the Lie bracket defined by (291) is a hyper-Lie algebra. Denote this hyper-Lie algebra by $gl_s^2(\mathbb{R})$.

In the following we construct the most general hyper linear algebra over the set of all matrices.

B. Hyper Linear Algebra $gl(\mathbb{R})$

Definition 14.5: Consider

$$gl(\mathbb{R}) := (\mathcal{M}, \hat{+}, \hat{\times}). \quad (292)$$

It is already known that $gl(\mathbb{R})$ is a hyper-ring. Define

$$[A, B]_{\hat{\times}} := A \hat{\times} B^- B \hat{\times} A. \quad (293)$$

Then $gl(\mathbb{R})$ becomes a hyper linear algebra.

Recall that the identity set is

$$\mathbf{e} = \{0_{m \times n} \mid m, n \in \mathbb{Z}_+\}.$$

First, for each $e = 0_{m \times n} \in \mathbf{e}$ $\mathcal{M}_e = \mathcal{M}_{m \times n}$. Hence we have

Proposition 14.6:

- (i)

$$gl(m \times n, \mathbb{R}) := ((\mathcal{M}_{m \times n}, +, \times), [\cdot, \cdot]) \quad (294)$$

is a Lie algebra.

- (ii) Particularly,

$$gl(n, \mathbb{R}) := ((\mathcal{M}_{n \times n}, +, \times), [\cdot, \cdot]) \quad (295)$$

is a Lie algebra, which is the well known n -th order general linear algebra.

- (iii)

$$gl(\mathbb{R}) = \bigcup_{m=1}^{\infty} \bigcup_{n=1}^{\infty} gl(m \times n, \mathbb{R}) \quad (296)$$

is a partition of the hyper-general linear algebra. And each Lie-algebra $gl(m \times n, \mathbb{R})$ is called its component Lie-algebra.

Then $A \sim B$ means there exist $0_{a \times b}$ and $0_{c \times d}$ such that

$$A \hat{+} 0_{a \times b} = B \hat{+} 0_{c \times d}. \quad (297)$$

Assume $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, $m \vee p = r$, and $n \wedge q = s$. By deleting common facts the (294) can be rewritten as

$$A \hat{+} 0_{r/m \times s/n} = B \hat{+} 0_{r/p \times s/q}. \quad (298)$$

It leads to

$$A \otimes \mathcal{E}_{r/m \times s/n} = B \otimes \mathcal{E}_{r/p \times s/q}. \quad (299)$$

Lemma 14.7: Consider $gl(\mathbb{R}) := (\mathcal{M}, \hat{+}, \hat{\times})$. The equivalence is consistent with $\hat{+}$ and $\hat{\times}$.

Proof.

The consistence of the equivalence w.r.t. $\hat{+}$ has already been proved in Proposition 3.14. We prove the consistence of equivalence w.r.t. $\hat{\times}$.

Assume $A_1 \in \mathcal{M}_{m \times n}$, $A_2 \in \mathcal{M}_{p \times q}$, $B_1 \in \mathcal{M}_{\alpha \times \beta}$, $B_2 \in \mathcal{M}_{\gamma \times \delta}$, and

$$\begin{aligned} m \vee \alpha &= a, & n \vee \beta &= b, & p \vee \gamma &= c, \\ q \vee \delta &= d, & m \wedge p &= s, & n \wedge q &= t, \\ \alpha \wedge \gamma &= \xi, & \beta \wedge \delta &= \eta, & \xi \vee t &= u, \\ a \vee b &= \theta = ux, & c \vee d &= \lambda = uy, \\ a &= a_0 s, & b &= b_0 \eta, & c &= c_0 s, & d &= d_0 \eta. \end{aligned}$$

Assume $A_1 \sim A_2$ and $B_1 \sim B_2$, we have to show that

$$A_1 \hat{\times} B_1 \sim A_2 \hat{\times} B_2. \quad (300)$$

We use equivalent condition (299) for matrix equivalence. Using Lemma 3.16, there exist $A_0 \in \mathcal{M}_{s \times t}$ and $B_0 \in \mathcal{M}_{\xi \times \eta}$, such that

$$\begin{aligned} A_1 &= A_0 \otimes \mathcal{E}_{m/s \times n/t}, & A_2 &= A_0 \otimes \mathcal{E}_{p/s \times q/t}, \\ B_1 &= B_0 \otimes \mathcal{E}_{\alpha/\xi \times \beta/\eta}, & B_2 &= B_0 \otimes \mathcal{E}_{\gamma/\xi \times \delta/\eta}. \end{aligned}$$

Then we have

$$\begin{aligned}
A_1 \hat{\times} B_1 &= (A_1 \otimes \mathcal{E}_{a/m \times b/n}) \times (B_1 \otimes \mathcal{E}_{a/\alpha \times b/\beta}) \\
&= (A_0 \otimes \mathcal{E}_{m/s \times n/t} \otimes \mathcal{E}_{a/m \times b/n}) \\
&\times (B_0 \otimes \mathcal{E}_{\alpha/\xi \times \beta/\eta} \otimes \mathcal{E}_{a/\alpha \times b/\beta}) \\
&= (A_0 \otimes \mathcal{E}_{a/s \times b/t}) \times (B_0 \otimes \mathcal{E}_{\alpha/\xi \times b/\eta}) \\
&= (A_0 \otimes \mathcal{E}_{a/s \times b/t} \otimes \mathcal{E}_{\theta/b}^T) (B_0 \otimes \mathcal{E}_{\alpha/\xi \times b/\eta} \otimes \mathcal{E}_{\theta/a}) \\
&= (A_0 \otimes \mathcal{E}_{a/s \times \theta/t}) (B_0 \otimes \mathcal{E}_{\theta/\xi \times b/\eta}) \\
&= (A_0 \otimes \mathcal{E}_{a_0 \times u/t}) (B_0 \otimes \mathcal{E}_{u/\xi \times b_0}) \\
&= [(A_0 \otimes \mathcal{E}_{1 \times u/t}) (B_0 \otimes \mathcal{E}_{u/\xi \times 1})] \otimes (\mathcal{E}_{a_0} \mathcal{E}_{b_0}^T). \\
&= [(A_0 \otimes \mathcal{E}_{1 \times u/t}) (B_0 \otimes \mathcal{E}_{u/\xi \times 1})] \otimes (\mathcal{E}_{a_0 \times b_0}).
\end{aligned}$$

Similarly, we also have

$$A_2 \hat{\times} B_2 = [(A_0 \otimes \mathcal{E}_{1 \times u/t}) (B_0 \otimes \mathcal{E}_{u/\xi \times 1})] \otimes (\mathcal{E}_{c_0 \times d_0}).$$

We conclude that

$$A_1 \hat{\times} B_1 \sim A_2 \hat{\times} B_2.$$

□

Denote the quotient space by

$$\Sigma = \mathcal{M} / \sim,$$

and using Lemma 14.7, the following result is obvious.

Proposition 14.8: The equivalent space $gl_{\Sigma}(\mathbb{R}) := (\Sigma, \hat{+}, \hat{\times})$ is a Lie algebra.

Definition 14.9: Assume V is a hyper vector space and there is a Lie bracket $[\cdot, \cdot] : V \times V \rightarrow V$, such that $(V, [\cdot, \cdot])$ is a Lie algebra.

- (i) If $N \subset V$ is a hyper vector subspace and $(N, [\cdot, \cdot])$ is also a Lie bracket, then $(N, [\cdot, \cdot])$ is called a hyper Lie sub-algebra of $(V, [\cdot, \cdot])$.
- (ii) If $N \subset V$ is a hyper Lie sub-algebra, and

$$[A, N] \subset N, \quad \forall A \in V, \quad (301)$$

then N is called an ideal of V .

As an immediate consequence, the following result is straightforward verifiable.

Proposition 14.10: $(\mathcal{M}_{\mu}, \hat{+}, \hat{\times})$ is a hyper Lie sub-algebra of $gl(\mathbb{R})$, denoted by $gl_{\mu}(\mathbb{R})$.

Remark 14.11: Summarizing the matrix hyper Lie algebras we obtained so far, we have the followings:

- (i) $(\mathcal{M}_1, \bar{+}, \times)$, its corresponding Lie hyper algebra is $gl_s^1(\mathbb{R})$ described in Example 14.4.
- (ii) $(\mathcal{M}_1, \hat{+}, \circ)$, its corresponding Lie hyper algebra is $gl_s^2(\mathbb{R})$ described in Example 14.4.
- (iii) $(\mathcal{M}, \hat{+}, \hat{\times})$, its corresponding Lie hyper algebra is $gl(\mathbb{R})$ described in Definition 14.5.
- (iv) $(\mathcal{M}_{\mu}, \hat{+}, \hat{\times})$, its corresponding Lie hyper algebra is $gl_{\mu}(\mathbb{R})$, which is a Lie sub-hyper algebra of $gl(\mathbb{R})$.

The following example presents an ideal of $gl(\mathbb{R})$.

Example 14.12: Define

$$T_{\square} := \{A \in \mathcal{M} \mid Tr(\square(A)) = 0\}. \quad (302)$$

Then T_{\square} is an ideal of $gl(\mathbb{R})$.

To see this, assume $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, and $m \vee n \vee p \vee q = r$. Then a straightforward computation shows that

$$\begin{aligned}
[A, B]_{\hat{\times}} &= (A \otimes \mathcal{E}_{r/m \times r/n}) (B \otimes \mathcal{E}_{r/p \times r/q}) \\
&\quad - (B \otimes \mathcal{E}_{r/p \times r/q}) (A \otimes \mathcal{E}_{r/m \times r/n}).
\end{aligned}$$

Hence,

$$Tr([A, B]_{\hat{\times}}) = 0.$$

The conclusion is obvious.

C. M -Depending Sub Hyper Algebra

This subsection considers a kind of sub-hyper algebras of $gl(\mathbb{R})$, which depend on any $M \in \mathcal{M}$. Given $M \in \mathcal{M}$, we define

$$G_M := \{A \in \mathcal{M} \mid A \hat{\times} M \hat{+} M \hat{\times} A^T = 0\}. \quad (303)$$

Theorem 14.13: Set

$$gl(M, \mathbb{R}) := (G_M, [\cdot, \cdot]_{\hat{\times}}). \quad (304)$$

Then $gl(M, \mathbb{R})$ is a hyper Lie subalgebra of $gl(\mathbb{R})$.

To prove this theorem, we need some preparations.

Lemma 14.14:

$$A \hat{\times} M \hat{+} M \hat{\times} A^T = 0, \quad (305)$$

if and only if,

$$\square(A \hat{\times} M) = -\square(M \hat{\times} A^T). \quad (306)$$

Proof. Assume $M \in \mathcal{M}_{m \times n}$, $A \in \mathcal{M}_{p \times q}$, and

$$m \vee p = \alpha, \quad n \vee q = \beta,$$

$$m \vee q = a, \quad n \vee p = b,$$

$$\alpha \vee a = m \vee p \vee q = c, \quad \beta \vee b = n \vee p \vee q = d,$$

$$\alpha \vee \beta = a \vee b = m \vee n \vee p \vee q = \xi.$$

Then

$$\begin{aligned}
A \hat{\times} M &= (A \otimes \mathcal{E}_{\alpha/p \times \beta/q}) \times (M \otimes \mathcal{E}_{\alpha/m \times \beta/n}) \\
&= (A \otimes \mathcal{E}_{\alpha/p \times \xi/q}) (M \otimes \mathcal{E}_{\xi/m \times \beta/n}) \in \mathcal{M}_{\alpha \times \beta}.
\end{aligned}$$

$$\begin{aligned}
M \hat{\times} A^T &= (M \otimes \mathcal{E}_{a/m \times b/n}) \times (A^T \otimes \mathcal{E}_{a/q \times b/p}) \\
&= (M \otimes \mathcal{E}_{a/m \times \xi/n}) (A^T \otimes \mathcal{E}_{\xi/q \times b/p}) \in \mathcal{M}_{a \times b}.
\end{aligned}$$

$$\begin{aligned}
A \hat{\times} M \hat{+} M \hat{\times} A^T &= (A \otimes \mathcal{E}_{\alpha/p \times \xi/q}) (M \otimes \mathcal{E}_{\xi/m \times \beta/n}) \hat{+} \\
&\quad (M \otimes \mathcal{E}_{a/m \times \xi/n}) (A^T \otimes \mathcal{E}_{\xi/q \times b/p}) \\
&= [(A \otimes \mathcal{E}_{\alpha/p \times \xi/q}) (M \otimes \mathcal{E}_{\xi/m \times \beta/n})] \otimes \mathcal{E}_{c/\alpha \times d/\beta} \hat{+} \\
&\quad [(M \otimes \mathcal{E}_{a/m \times \xi/n}) (A^T \otimes \mathcal{E}_{\xi/q \times b/p})] \otimes \mathcal{E}_{c/a \times d/b} \\
&= (A \otimes \mathcal{E}_{c/p \times \xi/q}) (M \otimes \mathcal{E}_{\xi/m \times d/n}) \hat{+} \\
&\quad (M \otimes \mathcal{E}_{c/m \times \xi/n}) (A^T \otimes \mathcal{E}_{\xi/q \times d/p})
\end{aligned}$$

Plugging into (305) yields

$$\begin{aligned} & (A \otimes \mathcal{E}_{c/p \times \xi/q}) (M \otimes \mathcal{E}_{\xi/m \times d/n}) + \\ & (M \otimes \mathcal{E}_{c/m \times \xi/n}) (A^T \otimes \mathcal{E}_{\xi/q \times d/p}) = 0_{c \times d}. \end{aligned} \quad (307)$$

It is clear that (307) is equivalent to (305).

Right \otimes -multiplying both sides of (307) by $\mathcal{E}_{\xi/c \times \xi/d}$, we have

$$\begin{aligned} & [(A \otimes \mathcal{E}_{c/p \times \xi/q}) (M \otimes \mathcal{E}_{\xi/m \times d/n})] \otimes \mathcal{E}_{\xi/c \times \xi/d} + \\ & [(M \otimes \mathcal{E}_{c/m \times \xi/n}) (A^T \otimes \mathcal{E}_{\xi/q \times d/p})] \otimes \mathcal{E}_{\xi/c \times \xi/d} \\ & = (A \otimes \mathcal{E}_{\xi/p \times \xi/q}) (M \otimes \mathcal{E}_{\xi/m \times \xi/n}) + \\ & (M \otimes \mathcal{E}_{\xi/m \times \xi/n}) (A^T \otimes \mathcal{E}_{\xi/q \times \xi/p}) \\ & = \square (A \hat{\times} M) + \square (M \otimes A^T) = 0_{\xi \times \xi}. \end{aligned} \quad (308)$$

Conversely, by deleting the factor $\mathcal{E}_{\xi/c \times \xi/d}$ from both sides of (308), (which is obviously "legal" as long as $\mathcal{E}_{\xi/c \times \xi/d} \neq 0$), we have (307). \square

Lemma 14.15:

$$\square(A \hat{\times} B) = \square A \circ \square(B), \quad (309)$$

where \circ is defined by (17).

Proof. Assume $A \in \mathcal{M}_{m \times n}$, $B \in \mathcal{M}_{p \times q}$, and

$$\begin{aligned} m \vee p &= \alpha, \quad n \vee q = \beta, \\ m \vee n &= a, \quad p \vee q = b, \\ \alpha \vee \beta &= a \vee b = m \vee n \vee p \vee q = \xi. \end{aligned}$$

Then

$$\begin{aligned} \square(A \hat{\times} B) &= \square [(A \otimes \mathcal{E}_{\alpha/m \times \beta/n}) \times (B \otimes \mathcal{E}_{\alpha/p \times \beta/q})] \\ &= \square [(A \otimes \mathcal{E}_{\alpha/m \times \xi/n}) \times (B \otimes \mathcal{E}_{\xi/p \times \beta/q})] \\ &= [(A \otimes \mathcal{E}_{\alpha/m \times \xi/n}) \times (B \otimes \mathcal{E}_{\xi/p \times \beta/q})] \otimes \mathcal{E}_{\xi/\alpha \times \xi/\beta} \\ &= (A \otimes \mathcal{E}_{\xi/m \times \xi/n}) (B \otimes \mathcal{E}_{\xi/p \times \xi/q}) \\ &= [(A \otimes \mathcal{E}_{a/m \times a/n}) \otimes J_{\xi/a}] [(B \otimes \mathcal{E}_{b/p \times b/q}) \otimes J_{\xi/b}] \\ &= \square(A) \circ \square(B). \end{aligned} \quad \square$$

As an immediate consequence, we have

$$\square \left(\hat{\times}_{i=1}^n A_i \right) = \circ_{i=1}^n \square(A_i), \quad A_i \in \mathcal{M}. \quad (310)$$

Proof of Theorem 14.13 Assume $A, B \in G_M$, using Lemma 14.14, (305) is equivalent to

$$\square \left[\left([A, B]_{\hat{\times}} \hat{\times} M \right) \right] + \square \left[M \hat{\times} [A, B]_{\hat{\times}}^T \right] = 0. \quad (311)$$

Using Lemma 14.15, we have

$$\begin{aligned} & \square \left[\left([A, B]_{\hat{\times}} \hat{\times} M \right) \right] + \square \left[M \hat{\times} [A, B]_{\hat{\times}}^T \right] \\ &= \square(A) \circ \square(B) \circ \square(M) - \square(B) \circ \square(A) \circ \square(M) \\ &+ \square(M) \circ \square(B^T) \circ \square(A^T) - \square(M) \circ \square(A^T) \circ \square(B^T) \\ &= -\square(A) \circ \square(M) \circ \square(B^T) + \square(B) \circ \square(M) \circ \square(A^T) \\ &+ \square(M) \circ \square(B^T) \circ \square(A^T) - \square(M) \circ \square(A^T) \circ \square(B^T) \\ &= \square(M) \circ \square(A^T) \circ \square(B^T) - \square(M) \circ \square(B^T) \circ \square(A^T) \\ &+ \square(M) \circ \square(B^T) \circ \square(A^T) - \square(M) \circ \square(A^T) \circ \square(B^T) \\ &= 0. \end{aligned} \quad \square$$

XV. HYPER LINEAR GROUP AS HYPER LIE GROUP

A. From Matrix Hyper Rings to Matrix Hyper Group

Observing that we have two well defined Lie group of matrices. One is well known general linear group $GL(n, \mathbb{R})$ and the other one is $GL(m \times n, \mathbb{R})$, which has been discussed in Section XII. It seems that the way to construct them can be described as follows: Starting from matrix ring, we first construct the Lie algebra by defining Lie bracket in a natural way, then build their corresponding group called the Lie group. Precisely, we have

$$\begin{aligned} (\mathcal{M}_{n \times n}, +, \times) &\Rightarrow gl(n, \mathbb{R}) \Rightarrow GL(n, \mathbb{R}); \\ (\mathcal{M}_{m \times n}, +, \times) &\Rightarrow gl(m \times n, \mathbb{R}) \Rightarrow GL(m \times n, \mathbb{R}). \end{aligned}$$

Recall Remark 14.11, we have 4 kinds of matrix hyper-Lie algebras. This section is aimed to build their corresponding hyper-Lie groups.

Following Remark 14.11, we consider all 4 different matrix hyper-Lie algebras.

(1) Consider case (i), which has been discussed in [19].

But we reconsider them from a new point of view: Consider them as hyper groups and then pose an \mathbb{R}^∞ differentiable structure on it.

Let $\mathcal{T} = \bigcup_{n=1}^\infty \mathcal{T}_n$, where \mathcal{T}_n is the set of $n \times n$ nonsingular matrices. Then

$$GL_s(\mathbb{R}) := (\mathcal{T}, \times, \mathbf{e}), \quad (312)$$

where $\mathbf{e} = \{I_n \mid n \in \mathbb{Z}_+\}$. Then the following facts are obvious;

- $GL_s(\mathbb{R})$ is a hyper group.
- Embedding $M_{n \times n}$ into \mathbb{R}^{n^2} by $A \mapsto V_c(A)$, and using the topology \mathcal{T}_d , then $GL_s(\mathbb{R})$ becomes a topological space and a topological hyper group, with the inherited topology from \mathbb{R}^∞ .
- Its differentiable structure comes from \mathbb{R}^Ξ with

$$\begin{aligned} \Xi &= \{n = m^2 \in \mathbb{Z}_+, \text{ and} \\ &n = s^2 \prec m = t^2 \text{ if and only if } s \prec t\}. \end{aligned}$$

(2)

The topological structure in (i) can hardly be used for this case, because $A \otimes \mathbf{1}_{k \times k}$ is no longer invertible, even if A is. We are not interested in constructing a hyper-Lie group with its hyper-Lie algebra $gl_s^2(\mathbb{R})$.

(3)

Denote by

$$\begin{aligned} G_{m \times n} &:= (\{I_{m \times n} + A_0 \mid A_0 \in \mathcal{M}_{m \times n}, \text{ and} \\ &I_{m \times n} + A_0 \text{ is invertible.}\}, \hat{\times}). \end{aligned}$$

We define

$$G_M := \bigcup_{m=1}^{\infty} \bigcup_{n=1}^{\infty} G_{m \times n} \subset \overline{\mathcal{M}}. \quad (313)$$

First we need to show that $(G_M, \hat{\times})$ is a hyper group. Set

$$\mathbf{e} := \{I_{m \times n} \mid (m, n) \in \mathbb{Z}_+ \times \mathbb{Z}_+\}, \quad (314)$$

with the lattice structure as MD-2 lattice. We need the following definition.

Definition 15.1:

(i) The product within \mathbf{e} :

$$I_{m \times n} \hat{\times} I_{p \times q} := I_{(m \vee p) \times (n \vee q)}. \quad (315)$$

(ii) The product of \mathbf{e} with \mathcal{M} : Let $A_0 \in \mathcal{M}_{p \times q}$, $m \vee p = s$, and $n \vee q = t$. Then

$$I_{m \times n} \hat{\times} A_0 = A_0 \hat{\times} I_{m \times n} := A_0 \otimes \mathcal{E}_{s/m \times t/n}. \quad (316)$$

Definition 15.2: Let $A, B \in G_M$, say, $A = I_{m \times n} + A_0$, $B = I_{p \times q} + B_0$, where $A_0 \in \mathcal{M}_{m \times n}$, $B_0 \in \mathcal{M}_{p \times q}$, $m \vee p = s$, and $n \vee q = t$. Then

$$A \hat{\times} B = I_{s \times t} + A_0 \otimes \mathcal{E}_{s/m \times t/n} + B_0 \otimes \mathcal{E}_{s/p \times t/q} + A_0 \hat{\times} B_0 \in \mathcal{M}_{s \times t}. \quad (317)$$

Proposition 15.3: $(G_M, \hat{\times}, \mathbf{e})$ is a hyper group, where $\hat{\times}$ is defined by (313) and \mathbf{e} is defined by (314).

Proof.

First, we prove the associativity. By the definition of $\hat{\times}$, It is enough to show that

$$(A \hat{\times} B) \hat{\times} C = A \hat{\times} (B \hat{\times} C), \quad A, B, C \in \mathcal{M} \bigcup \mathbf{e}. \quad (318)$$

Consider the following several cases.

(i) All $A, B, C \in \mathcal{M}$:

Then (318) has been proved in Proposition 2.28.

(ii) There are one or two factors from \mathbf{e} .

We prove one case, where $A, C \in \mathcal{M}$ and $B \in \mathbf{e}$. The proofs for all other cases are similar.

Assume $A \in \mathcal{M}_{m \times n}$, $B = I_{p \times q}$, $C \in \mathcal{M}_{r \times s}$, and

$$\begin{aligned} m \vee p &= a, \quad n \vee q = b, \quad a \vee r = c, \\ b \vee s &= d, \quad c \vee d = e, \quad n \vee r = \xi, \\ p \vee r &= u, \quad q \vee s = v, \quad m \vee u = m \vee p \vee r = c, \\ n \vee v &= n \vee q \vee s = d. \end{aligned}$$

Then

$$\begin{aligned} (A \hat{\times} I_{p \times q}) \hat{\times} C &= (A \otimes \mathcal{E}_{a/m \times b/n}) \hat{\times} C \\ &= (A \otimes \mathcal{E}_{a/m \times b/n} \otimes \mathcal{E}_{c/a \times d/b}) \times (C \otimes \mathcal{E}_{c/r \times d/s}) \\ &= (A \otimes \mathcal{E}_{c/m \times d/n}) \times (C \otimes \mathcal{E}_{c/r \times d/s}) \\ &= (A \otimes \mathcal{E}_{c/m \times d/n} \otimes \mathcal{E}_{e/d}^T) (C \otimes \mathcal{E}_{c/r \times d/s} \otimes \mathcal{E}_{e/c}) \\ &= (A \otimes \mathcal{E}_{c/m \times e/n}) (C \otimes \mathcal{E}_{e/r \times d/s}) \\ &= (A \otimes \mathcal{E}_{c/m \times \xi/n}) (C \otimes \mathcal{E}_{\xi/r \times d/s}). \end{aligned}$$

Similarly, we have

$$\begin{aligned} A \hat{\times} (I_{p \times q} \hat{\times} C) &= A \hat{\times} (C \otimes \mathcal{E}_{u/r \times v/s}) \\ &= (A \otimes \mathcal{E}_{c/m \times d/n}) \times (C \otimes \mathcal{E}_{u/r \times v/s} \otimes \mathcal{E}_{c/u \times d/v}) \\ &= (A \otimes \mathcal{E}_{c/m \times d/n} \otimes \mathcal{E}_{e/d}^T) (C \otimes \mathcal{E}_{c/r \times d/s} \otimes \mathcal{E}_{e/c}) \\ &= (A \otimes \mathcal{E}_{c/m \times e/n}) (C \otimes \mathcal{E}_{e/r \times d/s}) \\ &= (A \otimes \mathcal{E}_{c/m \times \xi/n}) (C \otimes \mathcal{E}_{\xi/r \times d/s}). \end{aligned}$$

(iii) All $A, B, C \in \mathbf{e}$: Let $A = I_{m \times n}$, $B = I_{p \times q}$, and $C = I_{r \times s}$. Then a straightforward computation shows that

$$(A \hat{\times} B) \hat{\times} C = A \hat{\times} (B \hat{\times} C) = I_{(m \vee p \vee r) \times (n \vee q \vee s)}.$$

The associativity is proved.

Next, we have to show that if $A \in \mathcal{M}_{m \times n}$ and $B \in \mathcal{M}_{p \times q}$ are invertible, then so is $A \hat{\times} B$. Using associativity, we have

$$\begin{aligned} (A \hat{\times} B) \hat{\times} (B^{-1} \hat{\times} A^{-1}) &= A \hat{\times} (B \hat{\times} B^{-1}) \hat{\times} A^{-1} \\ &= A \hat{\times} (I_{p \times q}) \hat{\times} A^{-1} \\ &= I_{p \times q} \hat{\times} (A \hat{\times} A^{-1}) \\ &= I_{p \times q} \hat{\times} I_{m \times n} = I_{(m \vee p) \times (n \vee q)}. \end{aligned}$$

Similarly, we have

$$(B^{-1} \hat{\times} A^{-1}) \hat{\times} (A \hat{\times} B) = I_{(m \vee p) \times (n \vee q)}.$$

Hence, $A \hat{\times} B$ is also invertible.

Finally, for each $e = I_{m \times n}$

$$(G_M)_{m \times n} = G_{m \times n},$$

which is a group. We conclude that $(G_M, \hat{\times})$ is a hyper group. \square

Remark 15.4:

(i) The invertibility of a matrix is the most important concept in group structure of G_M . We emphasize that the Definition 12.11 is consistent w.r.t. classical one for square matrix. Let $A \in \mathcal{M}_{n \times n}$. Then we can express

$$A = I_{n \times n} + A_0,$$

where $A_0 = A - I_n$. Then it is easy to see that A is invertible in classical sense, i.e., there exists $B = A^{-1}$ such that $AB = BA = I_n$, if and only if,

$$B = I_{n \times n} + B_0,$$

where $B_0 = B - I_n$ such that

$$A \hat{\times} B = I_{n \times n} + A_0 + B_0 + A_0 B_0 = I_{n \times n}.$$

This shows that $I_{n \times n} = I_n$ and the two invertibilities are equivalent.

(ii) Assume $A = I_{m \times n} + A_0 \in \overline{\mathcal{M}}_{m \times n}$ is invertible, where $m \neq n$. Then $A^T = I_{n \times m} + A_0^T$ is also invertible. $A \hat{\times} A^T$




Fig. 12. Matrix Hyper-Lie Group

is also invertible. Intuitively, this seems unusual. This fact shows that the product “ $\hat{\times}$ ” is different from the classical matrix product, and hence when $m \neq n$ the invertibility is different from the classical one.

B. From Matrix Hyper Group to Hyper Linear Group

To put topological and differentiable structure on G_M we have to give it coordinates. To this end, we first construct an Euclidian hyper manifold, \mathbb{R}^Ξ , where

$$\Xi = \{m \times n + 1, \mid (m, n) \in \mathbb{Z}_+ \times \mathbb{Z}_+\},$$

and $\{(m, n) \in \mathbb{Z}_+ \times \mathbb{Z}_+\}$ is the MD-2 lattice.

Then we define a mapping $\Psi : \overline{\mathcal{M}} \rightarrow \mathbb{R}^\Xi$ by

$$aI_{m \times n} + A_0 \mapsto (a, V_c(A_0)) \in \mathbb{R}^{mn+1}, \quad (m, n) \in \mathbb{Z}_+ \times \mathbb{Z}_+.$$

This mapping is described in Figure 12. Then for each $m \times n + 1 \in \Xi$ we have

$$\Psi|_{\overline{\mathcal{M}}_{m \times n}} : \overline{\mathcal{M}}_{m \times n} \rightarrow \mathbb{R}^{mn+1}.$$

And $V_c(A_0)$ is taken as the coordinate frame of $GL(m \times n, \mathbb{R})$. This assignment makes each $GL(m \times n, \mathbb{R})$ a manifold of dimension mn . As a consequence, $GL(\mathbb{R})$ becomes a matrix hyper-Lie group, called the hyper linear group.

Then each slice $GL(m \times n, \mathbb{R})$ has $gl(m \times n, \mathbb{R})$ as its Lie algebra.

XVI. CONCLUDING REMARKS

A. A Mathematics over MDMVs

To investigate linear (control) systems the classical matrix theory plays a fundamental rule. While the modern mathematics provides a solid foundation for the matrix theory. Particularly, the concepts from abstract algebra such as group, ring, and module are used to describe the matrices/vectors under certain operators; the geometric structures such as vector space, metric, inner product, topology, manifold, etc. are essential for describing the state space of the systems. Finally

the Lie group and Lie algebra structure over matrices leads to the general linear group and general linear algebra, which reveal the inside structure of the dynamics of linear (control) systems.

Recently, the STPs/STAs have been developed to overcome the dimension barrier of classical matrix theory. As the cross-dimensional operators, STPs/STAs demonstrate their dimension-free characteristic, which makes the classical algebraic and geometric structures unsuitable for MVMDs. They, therefore, are stimulating a new matrix theory to merge. This new matrix theory demands a new mathematical foundation. The purpose of this paper is to explore a new foundation, which is called the CDM. The CDM consists of three parts: hyper algebra, hyper geometry, and hyper linear algebra and hyper linear group.

The hyper algebra consists of three new concepts: the hyper group, hyper ring, and hyper module. The hyper group extends the classical group to a group with multiple identities, each identity represents a component group. The cross-dimensional operator joints all component groups together. The equivalence group shows the unified properties of component groups. Based on hyper group, the hyper ring and hyper module are also proposed in a natural way. They have similar structure: component objects and equivalence rings/modules.

The hyper geometry uses the hyper group structure to construct a hyper-vector space, which has multiple zeros. Then the hyper inner product space and hyper metric space are also obtained. Furthermore, the metric-based topological structure and a differentiable structure have been posed on \mathbb{R}^∞ , and using such topological and differentiable structure, the hyper manifold is constructed. These objects provide a geometric structure for MVMDs.

Finally, the Lie algebra and Lie group structure for hyper matrix group w.r.t. $\hat{+}$ and $\hat{\times}$ respectively, is proposed. The topological hyper group is also constructed. They eventually lead to the hyper-Lie algebra and the hyper-Lie group for MVMDs, called the hyper general linear algebra and hyper general linear group, are established.

B. Potential Applications

Dimension-varying systems appear in nature and from engineering systems. For instance, In a genetic regulatory network, cells may die or birth at any time; in the internet or some other service-based networks, some users may join in or withdraw out from time to time; some mechanical systems, such as docking, undocking, departure, and joining of spacecrafts [50], [32], modeling of biological systems [49], [30], uncertain model identification [47], etc.

There are also many dimension uncertain systems. For instance, a single electric power generator may be described by 2-dimensional model, or 3-dimensional, or even 5-, 6-, or 7-dimensional models [39]. From theoretical physics, the superstring theory proposes the space-time dimension could be 4 (Instain, Relativity), 5(Kalabi-Klein theory), 10 (Type 1 string), 11 (M-theory), or even 26 (Bosonic), etc. [33].

To deal with dimension varying and/or dimension uncertain systems, current fixed dimension mathematics seems not very efficient. In recent years, we try to use STP/STA to treat these kind of systems [15], [19], [20], [22]. Our previous approaches are based on the dynamics over quotient space, which has nice vector space structure. We then found that this approach is not suitable for dealing with networked systems, which have higher dimensions and the dimension-varying is relatively small. This fact stimulates us to investigate the structure of real spaces with mixed dimensions itself. That is the real motivation of the current paper.

By using the real space as the state-space of time-varying systems, it may provide a practically useful frame for modeling and analyzing the time-varying or uncertain systems and for designing controls.

Since the topic of this paper is wide and essential, what we did so far is very limited and very elementary. Hence, this paper is only providing a skeleton for what CDM should be. Further exploring and investigation are necessary.

REFERENCES

- [1] R. Abraham, J.E. Marden, *Foundations of Mechanics*, 2nd Ed., Benjamin/Cummings Pub., London, 1978.
- [2] T. Akutsu, *Algorithms for Analysis, Inference, and Control of Boolean networks*, World Scientific, New Jersey, 2018.
- [3] W.M. Boothby, *An Introduction to Differentiable Manifolds and Riemannian Geometry*, 2nd Ed., Academic Press, Inc., Orlando, 1986.
- [4] S. Burris, H.P. Sankappanavar, *A Course in Universal Algebra*, Springer, New York, 1981.
- [5] D. Cheng, Semi-tensor product of matrices and its application to Morgen's problem, *Sci. China Ser. F, Inf. Sci.*, Vol. 44, No. 3, 2001.
- [6] D. Cheng, H. Qi, Z. Li, *Analysis and Control of Boolean Networks, A Semi-Tensor Product Approach*, Springer, New York, 2011.
- [7] D. Cheng, H. Qi, Y. Zhao, *An Introduction to Semi-Tensor Product of Matrices and Its Applications*, World Scientific, Singapore, 2012.
- [8] D. Cheng, H. Qi, F. He, *Mappings and Dynamic Processes over Finite Sets - Semi-tensor Product Approach*, Science Press, Beijing, 2016 (in Chinese).
- [9] D. Cheng, H. Qi, *Semi-tensor Product of Matrices - Theory and Application*, Science Press, Beijing, 2007, 2-nd Ed., 2011, (in Chinese).
- [10] D. Cheng, *From Dimension-Free Matrix Theory to Cross-Dimensional Dynamic Systems*, Elsevier, London, 1019.
- [11] D. Cheng, On equivalence of Matrices, *Asian J. Math.*, Vol. 23, No. 2, 257-348, 2019.
- [12] D. Cheng, Z. Xu, T. Shen, Equivalence-based model of dimension-varying linear systems, *IEEE Trans. Aut. Contr.*, Vol. 65, No. 12, 5444-5449, 2020.
- [13] D. Cheng, H. Qi, *Lecture Notes in Semi-tensor Product of Matrices*, Vol. 1, Basic Theory and Multilinear Operation, Science Press, Beijing, 2020, (in Chinese).
- [14] D. Cheng, Y. Wu, G. Zhao, S. Fu, A comprehensive survey on STP approach to finite games, *J. Sys. Sci. Compl.*, Vol. 34, No. 5, 1666-1680, 2021.
- [15] D. Cheng, Z. Xu, T. Shen, Equivalence-based model of dimension-varying linear systems, *IEEE Trans. Aut. Contr.*, Vol. 65, No. 12, 5444-5449, 2020.
- [16] D. Cheng, H. Qi, *Lecture Notes in Semi-tensor Product of Matrices*, Vol. 1, Basic Theory and Multilinear Operation, Science Press, Beijing, 2020, (in Chinese).
- [17] D. Cheng, H. Qi, *Lecture Notes in Semi-tensor Product of Matrices*, Vol. 2, Analysis and Control of Logical Dynamic Systems, Science Press, Beijing, 2022, (in Chinese).
- [18] D. Cheng, C. Li, Y. Hao, X. Zhang, *Lecture Notes in Semi-tensor Product of Matrices*, Vol. 3, Semi-tensor Product Approach to Finite Games, Science Press, Beijing, 2022, (in Chinese).
- [19] D. Cheng, Z. Ji, From dimension-free manifolds to dimension-varying control systems, *Commun. Inform. Sys.*, Vol. 23, No. 1, 85-150, 2023.
- [20] D. Cheng, Z. Ji, *Lecture Notes in Semi-tensor Product of Matrices*, Vol. 4, *Finite and Dimension-Free Dynamic Systems*, Science Press, Beijing, 2023, (in Chinese).
- [21] D. Cheng, J. Feng, J. Zhong, Y. Wu, K. Zhang, *Lecture Notes in Semi-tensor Product of Matrices*, Vol. 5, *Application to Engineering and other Systems*, Science Press, Beijing, 2024, (in Chinese).
- [22] D. Cheng, From semi-tensor product of metrics to non-square general linear algebra and general linear group, (preprint: <http://arxiv.org/abs/2305.19794v2>, 2023), *Comm. Inform. Sys.*, to appear.
- [23] D. Cheng, X. Zhang, Z. Ji, C. Li, Observer-based realization of control systems, (preprint: <http://arxiv.org/abs/2404.15688>, 2024).
- [24] G. Choquet, *Topology*, Academic Press, New York, 1966.
- [25] J.B. Conway, *A course in Functional Analysis*, Springer-Verlag, New York, 1985.
- [26] J.D. Dixon, B. Mortimer, *Permutation Group*, Springer-Verlag, New York, 1996,.
- [27] J. Fang. *In Introduction to Lattice Theory*, High. Edu. Press, Beijing, 2014, (in Chinese).
- [28] E. Fornasini, M.E. Valcher, Recent developments in Boolean networks control, *J. Contr. Dec.*, Vol. 3, No. 1, 1-18, 2016.
- [29] W. Greub, *Linear Algebra*, 4 ed., Springer-Verlag, New York, 1981.
- [30] R. Huang, Z. Ye, An improved dimension-changeabel matrix model of simulationg the insect population dynamics, *Entomological Knowledge*, Vol. 32, No. 3, 162-164, 1995.
- [31] T.W. Hungerford, *Algebra*, Springer-verlag, New York, 1974.
- [32] P. Jiao, Y. Hao, J. Bin, Modeling and control of spacecraft formation based on impulsive switching with variable dimensions, *Computer Simulation*, Vol. 31, No. 6, 2014, 124-128.
- [33] M. Kaku, *Introduction to Superstring and M-Theory*, 2nd Ed., Springer-Verlag, New York, 1999.
- [34] H. Li, G. Zhao, M. Meng, J. Feng, A survey on applications of semi-tensor product method in engineering, *Science China*, Vol. 61, 010202:1-010202:17, 2018 .
- [35] H. Li, G. Zhao, P. Guo, Z. Liu, *Analysis and Control of Finite-Valued Systems*, CRC Press, 2018.
- [36] Z. Liu, H. Qiao, *S-System Theory of Semigroup*, 2nd Ed., Science Press, Beijing, 2008 (in Chinese).
- [37] Y. Liu, J. Liu, L. Sun, *Sampled-date Control of Logical Networks*, High Education Press, Springer, Singapore, 2023.
- [38] J. Lu, H. Li, Y. Liu, F. Li, Survey on semi-tensor product method with its applications in logical networks and other finite-valued systems, *IET Contr. Thm& Appl.*, Vol. 11, No. 13, 2040-2047, 2017.

- [39] J. Machowski, J.W. Bialek, J.R. Bumby, *Power System Dynamics and Stability*, John Wiley and Sons, Inc., Chichester, 1997.
- [40] S. Mei, F. Liu, A. Xie, *Semi-tensor Product Approach for Transient Process Analysis of Power Systems*, Tsinghua Univ. Press, Beijing, 2010, (in Chinese).
- [41] A. Muhammad, A. Rushdi, F.A. M. Ghaleb, A tutorial exposition of semi-tensor products of matrices with a stress on their representation of Boolean function, *JKAU Comp. Sci.*, Vol. 5, 3-30, 2016.
- [42] J.L. Kelley, *General Topology*, Springer-Verlag, New York, 1955.
- [43] H.H. Schaefer, *Topological Vector Spaces*, Springer-Verlag, New York, 1970.
- [44] A.E. Taylor, D.C. Lay, *Introduction to Functional Analysis*, John Wiley & Sons Inc., New York, 1980.
- [45] V.S. Varadarajan, *Lie Groups, Lie Algebras, and Their Representations*, Springer-Verlag, New York, 1984.
- [46] S.E. *Boolean Systems - Topics in Asynchronicity*, Elsevier, London, 2023.
- [47] H. Wang, J. Wang, X. Bian, G. Fu, SLAM of AUV based on the combined EKF, *Robot*, Vol. 34, No. 1, 56-64, 2012.
- [48] W.M. Wonham, *Linear Multivariable Control, A Geometric Approach*, Springer-Verlag, Berlin, 1974.
- [49] R. Xu, L. Liu, Q. Zhu, J. Shen, Application of a dimension-changeable matrix model on the simulation of the population dynamics of greenhouse whiteflies, *ACTA Ecological Sinica*, Vol. 1, No. 2, 147-158, 1981.
- [50] H. Yang, B. Jiang, V. Cocquempot, *Stabilization of Switched Nonlinear Systems with Unstable Modes*, Chapter 4. Switched Nonlinear Systems with Varying states, Springer, Switzerland, 2014.
- [51] Y. Yan, D. Cheng, J. Feng, H. Li, J. Yue, Survey on applications of algebraic state space theory of logical systems to finite state machines, *Sci. China, Inform. Sci.*, //doi.org/10.1007/s11432-022-3538-4.
- [52] K. Zhang, L. Zhang, L. Xie, *Discrete-Space Dynamic Systems*, Springer, New York, 2020.
- [53] J. Zhong, D. Lin, On minimum period of nonlinear feedback shift registers in Grain-like structure, *IEEE Trans. Inform. Theory*, Vol. 64, No. 9, 6429-6442, 2018.