

Quantum geometry of bosonic Bogoliubov quasiparticles

Isaac Tesfaye^{1,*} and André Eckardt^{1,†}

¹*Institut für Theoretische Physik, Technische Universität Berlin, Hardenbergstr. 36, D-10623 Berlin, Germany*
(Dated: July 17, 2024)

Topological and geometrical features arising in bosonic Bogoliubov-de Gennes (BdG) systems have mainly been studied by utilizing a generalized symplectic version of the Berry curvature and related Chern numbers. Here, we propose a symplectic quantum geometric tensor (SQGT), whose imaginary part leads to the previously studied symplectic Berry curvature, while the real part gives rise to a symplectic quantum metric, providing a natural distance measure in the space of bosonic Bogoliubov modes. We propose how to measure all components of the SQGT by extracting excitation rates in response to periodic modulations of the systems' parameters. Moreover, we connect the symplectic Berry curvature to a generalized symplectic anomalous velocity term for Bogoliubov Bloch wave packets. We test our results for a bosonic Bogoliubov-Haldane model.

Introduction.—Bosonic Bogoliubov-de Gennes (BBdG) Hamiltonians naturally arise in systems of weakly interacting bosons [1–4], or in photonic systems subjected to parametric driving, generating squeezing of light [5, 6]. They also describe magnonic [7–9], phononic [10], and mechanical systems [11]. Different from their fermionic (BCS) counterparts, BBdG systems are diagonalized in a symplectic manner via a paraunitary transformation [2, 12]. Their topological features [13, 14] have first been studied in the context of magnonic crystals [7, 8] by utilizing a generalized symplectic Chern number based on a symplectic Berry curvature for Bogoliubov bands. A generalization of the conventional bulk-boundary correspondence [15, 16] to the symplectic case can even lead to novel chiral edge states, whose properties are fundamentally distinct from their particle-number-conserving counterparts [17–21]. Recent works, building on the ten-fold Altland-Zirnbauer classification of topological insulators [22–26], corroborate the notions of the symplectic Berry curvature and Chern number [27, 28] and provide a topological classification of quadratic BBdG systems [29–34].

Although topological features of BBdG systems have recently received some attention [20, 21, 33, 35–54], the geometry of the underlying Bogoliubov state space is still lacking a complete characterization. In particle-number-conserving systems, the Berry curvature is related to the imaginary part of the so-called quantum geometric tensor (QGT) [55, 56] and the real part of the QGT, known as quantum metric, provides a natural distance measure in state space [56, 57]. Here, we generalize these concepts to the non-number-conserving BBdG case and propose a symplectic quantum geometric tensor (SQGT) [58]. We show that the imaginary part of the SQGT is related to the previously investigated symplectic Berry curvature and that its real part defines a natural distance measure in the space of bosonic Bogoliubov modes, which we dub symplectic quantum metric. Previous proposals




FIG. 1. (a) Illustration of the symplectic quantum metric providing a distance measure ds between infinitesimally close Bogoliubov modes $w(\lambda_0)$ and $w(\lambda_0 + d\lambda)$ [cf. Eq. (11)]. The integrated excitation rate Γ_{int}^n of a Bogoliubov system in response to (weak) periodic perturbations is dictated by the symplectic quantum geometry [cf. Eqs. (14) and (16)]. (b) Illustration of the transverse velocity a Bogoliubov Bloch wave packet experiences in response to an external force, which is shown to be proportional to the symplectic Berry curvature [cf. Eq. (18)].

for measurable signatures of the topology of BBdG systems have solely focused on probing bosonic chiral edge modes, predicted for non-zero symplectic Chern numbers [17, 21]. Here, inspired by Refs. [59, 60], we propose to measure the full SQGT from excitation rates to other Bogoliubov modes in response to weak periodic driving [Fig. 1(a)]. As a further probe, we also show that, analogous to the number-conserving case [61–63], the symplectic Berry curvature naturally arises as a transverse velocity term, when a Bogoliubov Bloch wave packet is subjected to an external force [Fig. 1(b)].

Framework.— Consider a BBdG Hamiltonian

$$\hat{H} = \sum_{i,j=1}^N K_{ij} \hat{a}_i^\dagger \hat{a}_j + \frac{1}{2} \left(G_{ij} \hat{a}_i^\dagger \hat{a}_j^\dagger + G_{ji}^* \hat{a}_i \hat{a}_j \right) = \frac{1}{2} \hat{\Psi}^\dagger \mathcal{M} \hat{\Psi} - \frac{1}{2} \text{Tr} \{K\}, \quad \mathcal{M} = \begin{pmatrix} K & G \\ G^* & K^* \end{pmatrix}, \quad (1)$$

with $K_{ij} = K_{ji}^* \in \mathbb{C}$ (hermitian), $G_{ij} = G_{ji} \in \mathbb{C}$ (symmetric) forming the hermitian $2N \times 2N$ matrix \mathcal{M} [64] and Nambu spinor $\hat{\Psi} = (\hat{a}_1 \cdots \hat{a}_N \hat{a}_1^\dagger \cdots \hat{a}_N^\dagger)^T$ of bosonic

* i.tesfaye@tu-berlin.de

† eckardt@tu-berlin.de

annihilation (creation) operators $\hat{a}_i^{(\dagger)}$ [2, 12, 65]. The Hamiltonian can be diagonalized [2, 12] (see also supplemental material (SM) [66]),

$$\hat{H} = \frac{1}{2}\hat{\Phi}^\dagger W^\dagger \mathcal{M} W \hat{\Phi} - \frac{1}{2}\text{Tr}\{K\} = \sum_{n=1}^N \omega_n \hat{b}_n^\dagger \hat{b}_n + C, \quad (2)$$

by a Bogoliubov transformation $\hat{\Psi} = W\hat{\Phi}$, where $\hat{\Phi} = (\hat{b}_1 \cdots \hat{b}_N \hat{b}_1^\dagger \cdots \hat{b}_N^\dagger)^T$ and

$$W = \begin{pmatrix} U & V^* \\ V & U^* \end{pmatrix}, \quad \text{with} \quad \begin{cases} W^\dagger \tau_z W = W \tau_z W^\dagger = \tau_z, \\ W^\dagger \mathcal{M} W = \Omega. \end{cases} \quad (3)$$

Here $C \equiv \sum_n \omega_n / 2 - \text{Tr}\{K\} / 2$; U and V are complex $N \times N$ matrices, so that $\hat{a}_i = \sum_n [U_{in} \hat{b}_n + (V^*)_{in} \hat{b}_n^\dagger]$; and $\Omega = \text{diag}(\omega_1, \dots, \omega_N, \omega_1, \dots, \omega_N)$ with real strictly positive excitation energies, $\omega_n > 0$, assumed in the following [64]. The transformation W is paraunitary, as indicated by the relation involving $\tau_z = \sigma_z \otimes \mathbb{1}_N$, with Pauli matrix σ_z and $N \times N$ identity matrix $\mathbb{1}_N$. This ensures that the new operators $\hat{b}_n^{(\dagger)}$ fulfill bosonic commutation relations. For a different ordering of the bosonic operators in the Nambu spinors $\hat{\Psi}$ and $\hat{\Phi}$, W becomes an element of the real symplectic group $\text{Sp}(2N, \mathbb{R})$ [67, 68]. Using $\tau_z^2 = \mathbb{1}_{2N}$, we identify $W^{-1} = \tau_z W^\dagger \tau_z$, such that the inverse transformation $\hat{\Phi} = W^{-1} \hat{\Psi}$ reads $\hat{b}_n = \sum_m [(U^\dagger)_{ni} \hat{a}_i - (V^\dagger)_{ni} \hat{a}_i^\dagger]$.

Symplectic quantum geometric tensor.— The Bogoliubov ground state $|\psi_0\rangle$ is the quasiparticle vacuum, $\hat{b}_n |\psi_0\rangle = 0 \forall n$. It is generally a multi-mode squeezed state with a fluctuating non-zero number of \hat{a}_i bosons [17, 45, 68]. Although $|\psi_0\rangle$ may display geometrical features itself, here we will focus on the geometrical properties of the quasiparticle excitations on top of it [17].

The Bogoliubov modes are associated with the vectors \mathbf{w}_n given by the columns of $W = (\mathbf{w}^1, \dots, \mathbf{w}^{2N})$. For $n \leq N$, they describe quasiparticles, $\hat{b}_n^\dagger = \hat{\Psi}^\dagger \tau_z \mathbf{w}^n$, and for $N > n$ to quasiholes, $\hat{b}_n = -\hat{\Psi}^\dagger \tau_z \mathbf{w}^n$ [18, 45, 66]. They play a role akin to single-particle wavefunctions [17] and solve the eigenvalue equation [66],

$$D \mathbf{w}^n = \tilde{\Omega}_n \mathbf{w}^n, \quad \text{with} \quad D \equiv \tau_z \mathcal{M} \quad \text{and} \quad \tilde{\Omega}_n = (\tau_z \Omega)_{nn}, \quad (4)$$

where D is called dynamical matrix. The excitation energies read $\tilde{\Omega}_n = (\tau_z)_{nn} \omega_n \bmod N = s_n \omega_n$, which are positive ($s_n = +1$) for quasiparticles ($n \leq N$) and negative ($s_n = -1$) for quasiholes ($n > N$). They also obey the generalized (Krein) τ_z inner product structure $(\mathbf{w}^m)^\dagger \tau_z \mathbf{w}^n = (\tau_z)_{mn} = \delta_{mn} s_n$ [69] (see SM [66]). The projector P^n onto the n th Bogoliubov mode and its orthogonal complement Q^n are naturally defined as [7]

$$P^n = W \Gamma^n W^{-1} = W \Gamma^n \tau_z W^\dagger \tau_z, \quad Q^n = \mathbb{1}_{2N} - P^n, \quad (5)$$

where $(\Gamma^n)_{lm} = \delta_{nl} \delta_{lm}$. Then $P^n P^m = \delta_{nm} P^n$ and $\sum_{n=1}^{2N} P^n = \mathbb{1}_{2N}$ [7]. We stress that P^n and Q^n

are not hermitian, but $\tau_z X^n \tau_z = (X^n)^\dagger$ for $X^n \in \{P^n, Q^n\}$, a property which is called Krein- or pseudo-hermiticity [28, 69]. Below, we assume $\tilde{\Omega}_n \neq \tilde{\Omega}_m$ for $n \neq m$, but the generalization to the degenerate case is straightforward.

Let the Hamiltonian (1) now depend on a set of dimensionless parameters, $\boldsymbol{\lambda} = (\lambda^1, \dots, \lambda^r)$, i.e. $\hat{H} \equiv \hat{H}(\boldsymbol{\lambda})$. Then, motivated by the definition of the conventional quantum geometric tensor (QGT) [55, 57], we define the symplectic quantum geometric tensor (SQGT)

$$\eta_{\mu\nu}^n = \text{Tr}\{\partial_\mu P^n Q^n \partial_\nu P^n\}, \quad (6)$$

where $\partial_\mu \equiv \partial / \partial \lambda^\mu$. The SQGT is $U(1)$ gauge-invariant, since the projectors P_n and Q_n do not depend on the phase of the modes $\mathbf{w}^n(\boldsymbol{\lambda})$. The Krein-hermiticity of P^n and Q^n (5), implies that the SQGT is hermitian, $(\eta_{\mu\nu}^n)^* = \eta_{\nu\mu}^n$. By employing completeness, $\sum_n P^n = \mathbb{1}_{2N}$, and orthogonality, $(\mathbf{w}^m)^\dagger \tau_z \mathbf{w}^n = \delta_{mn} s_n$, the SQGT can be expressed as [66]

$$\eta_{\mu\nu}^n = s_n (\partial_\mu \mathbf{w}^n)^\dagger \tau_z Q^n \partial_\nu \mathbf{w}^n. \quad (7)$$

Moreover, can be obtained [66]

$$\eta_{\mu\nu}^n = s_n \sum_{m \neq n} s_m \frac{(\mathbf{w}^n)^\dagger (\partial_\mu \mathcal{M}) \mathbf{w}^m (\mathbf{w}^m)^\dagger (\partial_\nu \mathcal{M}) \mathbf{w}^n}{(\tilde{\Omega}_n - \tilde{\Omega}_m)^2}. \quad (8)$$

A first indication that the so-defined SQGT (6) is meaningful, is that its imaginary part is proportional to the previously defined symplectic Berry-curvature $B_{\mu\nu}^n = i s_n [(\partial_\mu \mathbf{w}^n)^\dagger \tau_z (\partial_\nu \mathbf{w}^n) - (\partial_\nu \mathbf{w}^n)^\dagger \tau_z (\partial_\mu \mathbf{w}^n)]$ [7, 8, 21],

$$B_{\mu\nu}^n = -2 \text{Im}[\eta_{\mu\nu}^n] = -2 \text{Im}[\text{Tr}\{\partial_\mu P^n Q^n \partial_\nu P^n\}], \quad (9)$$

which is real-valued and antisymmetric, $B_{\mu\nu}^n = -B_{\nu\mu}^n$. From this quantity a symplectic (first) Chern number $C^n \equiv \frac{1}{2\pi} \int_S B^n$ was defined for closed two-dimensional parameter spaces S [7, 8], which describes a generalized bulk-boundary correspondence [17–21]. Like in the conventional case, we show in Ref. [66] that the symplectic Berry curvature admits a local conservation law, i.e., $\sum_n B_{\mu\nu}^n = 0$. We, moreover, show below that the symplectic Berry curvature describes the anomalous (transverse) velocity induced by an external force in Bogoliubov Bloch bands.

We now show that also the real part of the SQGT,

$$g_{\mu\nu}^n = \text{Re}[\eta_{\mu\nu}^n] = \text{Re}[\text{Tr}\{\partial_\mu P^n Q^n \partial_\nu P^n\}], \quad (10)$$

which is real-valued and symmetric, $g_{\mu\nu}^n = g_{\nu\mu}^n$, carries physical meaning. We dub it symplectic quantum metric. Unlike the symplectic Berry curvature, which is only non-zero for systems with broken time-reversal or inversion symmetry [61], the symplectic quantum metric can already be non-zero without these constraints. It generalizes the conventional quantum metric [55, 57, 70, 71], also known as the Fubini-Study metric [56, 57, 72, 73], which is given by the real part of the conventional QGT

and defines a natural distance measure in the (projective) Hilbert space. The latter is employed to describe quantum phase transitions [74–80] as well as excitations rates and quantum fluctuations [56, 60, 81] (properties which have been exploited to measure the quantum metric in various experimental setups [82–89]).

The symplectic quantum metric $g_{\mu\nu}$ naturally describes the distance ds^2 between two infinitesimally close Bogoliubov modes $\mathbf{w}(\boldsymbol{\lambda})$ and $\mathbf{w}(\boldsymbol{\lambda} + d\boldsymbol{\lambda})$,

$$ds^2 = 1 - |\mathbf{w}^\dagger(\boldsymbol{\lambda})\tau_z\mathbf{w}(\boldsymbol{\lambda} + d\boldsymbol{\lambda})|^2 = g_{\mu\nu}d\lambda^\mu d\lambda^\nu, \quad (11)$$

with summation over repeated indices. Here we first defined ds^2 via the fidelity $|\mathbf{w}^\dagger(\boldsymbol{\lambda})\tau_z\mathbf{w}(\boldsymbol{\lambda} + d\boldsymbol{\lambda})|^2$ between both modes corresponding to the inner-product structure described above. The second identity follows from expanding $\mathbf{w}(\boldsymbol{\lambda} + d\boldsymbol{\lambda})$ to second order in $d\boldsymbol{\lambda}$, giving $ds^2 = \{(\mathbf{w}^\dagger\tau_z\mathbf{w})[(\partial_\mu\mathbf{w})^\dagger\tau_z(\partial_\nu\mathbf{w})] - (\partial_\mu\mathbf{w})^\dagger\tau_z\mathbf{w}\mathbf{w}^\dagger\tau_z(\partial_\nu\mathbf{w})\}d\lambda^\mu d\lambda^\nu = \eta_{\mu\nu}d\lambda^\mu d\lambda^\nu = \frac{1}{2}(\eta_{\mu\nu} + \eta_{\nu\mu})d\lambda^\nu d\lambda^\mu = g_{\mu\nu}d\lambda^\mu d\lambda^\nu$ [66].

Relation to linear response and probing.— We will now show that the SQGT determines the transition rates of Bogoliubov excitations between different modes in linear response to weak external forcing. This provides another physical interpretation of this quantity and opens the door for measuring its components similarly to the protocols that have been used to extract the quantum geometry of free fermions [59, 60, 81, 90].

Assuming a quasiparticle in a superposition of (unperturbed) Bogoliubov modes, $\boldsymbol{\psi}(t) = \sum_{n=1}^N \psi_n(t)\mathbf{w}^n(\boldsymbol{\lambda}_0) \in \mathbb{C}^{2N}$, its dynamics is described by the effective time-dependent Schrödinger equation $-i\partial_t\boldsymbol{\psi}(t) = D(t)\boldsymbol{\psi}(t)$ [45] (see SM [66]), with $D(t)$ being the dynamical matrix (4), which becomes time-dependent in response to external forcing (external forces generally also induce terms in the Hamiltonian that do not conserve the number of Bogoliubov quasiparticles, which are not captured by $D(t)$, which will be discussed below).

We assume driving in the form of a sinusoidal modulation $\lambda^\nu = \lambda_0^\nu + 2(A_\nu/\omega)\cos(\omega t - \varphi_\nu)$ of the system parameters around a mean value $\boldsymbol{\lambda}_0$. For weak driving ($A_\nu/\omega \ll 1$), the resulting time dependence of the dynamical matrix, $D(t) = D[\boldsymbol{\lambda}(t)]$, is captured by the linear approximation $D[\boldsymbol{\lambda}(t)] \simeq D(\boldsymbol{\lambda}_0) + \sum_\nu \partial_\nu D(\boldsymbol{\lambda}_0)2(A_\nu/\omega)\cos(\omega t - \varphi_\nu)$. For the sake of simplicity and the design of a probing scheme inspired by Ref. [60], we focus on a two-parameter drive with relative phase φ , for which

$$D[\boldsymbol{\lambda}(t)] = D(\boldsymbol{\lambda}_0) + \partial_1 D(\boldsymbol{\lambda}_0)2(A_1/\omega)\cos(\omega t) \pm \partial_2 D(\boldsymbol{\lambda}_0)2(A_2/\omega)\cos(\omega t - \varphi). \quad (12)$$

The driving amplitudes A_ν shall be switched on suddenly at time $t = 0$, and we assume, for simplicity, that the system is prepared in mode n , $\boldsymbol{\psi}(0) = \mathbf{w}^n(\boldsymbol{\lambda}_0)$. Applying time-dependent perturbation theory in first order (linear

response), we obtain the probabilities

$$p_{mn}^\pm(\omega, t) = |(\mathbf{w}^m)^\dagger(A_1\partial_1\mathcal{M} \pm A_2e^{i\varphi}\partial_2\mathcal{M})\mathbf{w}^n|^2 \times (2\pi t/\omega^2)\delta(\tilde{\Omega}_{mn} - \omega). \quad (13)$$

for finding the system in state \mathbf{w}^m at time t , where $\tilde{\Omega}_{mn} \equiv \tilde{\Omega}_m - \tilde{\Omega}_n$. Note that all quantities are evaluated for $\boldsymbol{\lambda} = \boldsymbol{\lambda}_0$, whenever the argument ($\boldsymbol{\lambda}$) is dropped, i.e., $\mathbf{w}^m \equiv \mathbf{w}^m(\boldsymbol{\lambda}_0)$ and $\tilde{\Omega}_m \equiv \tilde{\Omega}_m(\boldsymbol{\lambda}_0)$. The total excitation rate $\Gamma(\omega)$ into all other Bogoliubov modes $m \neq n$ is then given by $\Gamma^{n,\pm}(\omega) = t^{-1} \sum_{m \neq n} s_m p_{mn}^\pm(\omega, t)$. For an initial quasiparticle state, $n \leq N$, the sum can be restricted to $m \leq N$, since there is no quasiparticle-to-quasihole transition for a single-quasiparticle initial state [66].

To probe the SQGT, we consider the total excitation rate integrated over all (relevant) probing frequencies ω [59, 60], $\Gamma_{\text{int}}^n = \int d\omega \Gamma^{n,\pm}(\omega)$. For the modulation of a single parameter ($A_1 \equiv A$, $A_2 = 0$), we find

$$\Gamma_{\text{int}}^n = 2\pi A^2 \sum_{m \neq n} \frac{|(\mathbf{w}^m)^\dagger(\partial_1\mathcal{M})\mathbf{w}^n|^2}{s_m(\tilde{\Omega}_m - \tilde{\Omega}_n)^2} = 2\pi A^2 g_{11}^n. \quad (14)$$

For the second identity, we have used the spectral representation (10) of the SQGT, whose diagonal elements correspond to the symmetric part given by the symplectic quantum metric, i.e. $\eta_{\nu\nu}^n = g_{\nu\nu}^n$. Thus, we can probe these diagonal elements by measuring the frequency integrated response to a modulation of λ^ν .

Guided once more by Refs. [59, 60] for conventional systems, we now consider the driving of both parameters λ^1 and λ^2 with two different phase lags φ and equal amplitudes $A_1 = A_2 = A$, to obtain the frequency integrated transition rates

$$\Gamma_{\text{int}}^{n,\pm} = 2\pi A^2 \begin{cases} g_{11}^n \pm 2g_{12}^n + g_{22}^n, & \varphi = 0 \\ g_{11}^n \mp B_{12}^n + g_{22}^n, & \varphi = \pi/2, \end{cases} \quad (15)$$

the differences of which,

$$\Delta\Gamma_{\text{int}}^n \equiv \Gamma_{\text{int}}^{n,+} - \Gamma_{\text{int}}^{n,-} = \begin{cases} 8\pi A^2 g_{12}^n, & \varphi = 0 \\ -4\pi A^2 B_{12}^n, & \varphi = \pi/2, \end{cases} \quad (16)$$

are directly proportional to the real and imaginary part of the off diagonal element η_{12}^n of the SQGT, given by g_{12}^n and $-B_{12}^n/2$. In this way, all off-diagonal components $\eta_{\mu\nu}^n = g_{\mu\nu}^n - iB_{\mu\nu}^n/2$ can be extracted from the linear response of the system, allowing (in principle) for a complete reconstruction of the SQGT from quasiparticle scattering rates.

We can generalize the above results to initial states of multiple quasiparticles. A generic perturbation, $V(t) = \sum_{ij} \{A_{ij}(t)\hat{a}_i^\dagger\hat{a}_j + \frac{1}{2}[B_{ij}(t)\hat{a}_i^\dagger\hat{a}_j^\dagger + \text{h.c.}]\} + \sum_i \{C_i(t)\hat{a}_i^\dagger + \text{h.c.}\}$, takes the form $V(t) = \sum_{nm} \{A'_{nm}(t)\hat{b}_n^\dagger\hat{b}_m + \frac{1}{2}[B'_{nm}(t)\hat{b}_n^\dagger\hat{b}_m^\dagger + \text{h.c.}]\} + \sum_n \{C'_n(t)\hat{b}_n^\dagger + \text{h.c.}\} + D'(t)$ with respect to the unperturbed Bogoliubov operators $\hat{b}_n^{(\dagger)}$. Here terms $\propto A'$ correspond to scattering between quasiparticle (or hole) states, whereas terms $\propto B'$ are associated




FIG. 2. (a) Sketch of the measurement protocol: Selecting transitions by the frequency of the drive, first the Bogoliubov mode n is populated (red), before, in a second step, a transition to another mode m is induced (green). (b) Illustration of the Bogoliubov-Haldane model. (c) Bogoliubov energy spectrum along high-symmetry path in the first Brillouin zone (inset) for $J_2 = 0.1J_1$, $\theta = \pi/2$, $\Delta = 3\sqrt{3}J_2/2$ and different interaction strengths U_n/J_1 .

with the scattering of a quasiparticle into a quasihole or vice versa. The terms $\propto C'$, which correspond to the creation of a single quasiparticle (or hole), can result from the scattering of an \hat{a}_i boson out of (or into) a Bose condensate, which is described by a macroscopic wavefunction that is not in equilibrium with respect to the perturbation. All these processes are associated with corresponding bosonic enhancement factors. To reconstruct the full SQGT, it is sufficient to consider the scattering of single quasiparticles. In case of discrete energies ω_n , this can be achieved by exploiting the fact that different processes can be addressed selectively by appropriate driving frequencies [91]. Starting from the ground state, one can first exploit the B' or C' terms to populate the state n and then modulate the system resonantly with to transitions $n \rightarrow m$ of A' type to extract information about the SQGT [Fig. 2(a)].

Application to Bogoliubov-Haldane model.— As a concrete example, we study weakly interacting bosons in a two-dimensional hexagonal tight-binding lattice with two sublattice states, $s = A, B$, per unit cell. The single-particle terms correspond to the Haldane model [92], as it was recently realized with ultracold atoms in optical lattices [90, 93, 94]. They are characterized by real (complex) tunneling matrix elements $-J_1$ ($-J_2 e^{i\theta}$) between nearest (next-nearest) neighbors, and on-site potentials Δ ($-\Delta$) on A (B) sites [Fig. 2(b)]. Due to the lattice symmetry, quasimomentum $\mathbf{q} = (q_x, q_y)$ is conserved by these processes, which is treated as a parameter in the following. By treating a weak repulsive on-site interaction U at a filling of n bosons per site in a self-consistent Bogoliubov approximation with quantum fluctuations on top of a Gross-Pitaevskii mean-field condensate, we obtain a Bogoliubov-Haldane model, $\hat{H}^{(B)} = \sum_{\mathbf{q} \neq 0} \sum_{s, s'} \hat{a}_{\mathbf{q}s}^\dagger [\mathcal{H}(\mathbf{q}) + \mathcal{H}_1]_{ss'} \hat{a}_{\mathbf{q}s'} + (\hat{a}_{\mathbf{q}s}^\dagger [\mathcal{H}_2]_{ss'} \hat{a}_{-\mathbf{q}s'}^\dagger +$




FIG. 3. Integrated and ideal symplectic quantum metric component $g_{xx}(\mathbf{q})$ and Berry curvature $B_{xy}(\mathbf{q})$ for the upper particle band of the Bogoliubov-Haldane model [66]. (a) Integrated (left) and ideal (right) symplectic quantum metric component $g_{xx}(\mathbf{q})$ for $U_n/J_1 = 3$ shown in 2D quasi-momentum space. (b) Same as in (a) but for the symplectic Berry curvature $B_{xy}(\mathbf{q})$. (c) Ideal (solid line) and integrated (crosses) symplectic quantum metric component $g_{xx}(\mathbf{q})$ along a high-symmetry path (cf. inset in Fig. 2(c)) for two interaction strengths $U_n/J_1 = 0$ (non-interacting Haldane model) and $U_n/J_1 = 3$. (d) Same as in (c) but for the symplectic Berry curvature $B_{xy}(\mathbf{q})$. Simulation parameters: $A/J_1 = 0.01$, $J_1 t/\hbar = 10$ (final integration time), $\hbar\omega/J_1 \in [0.1(0.2), 5.5]$ for $U_n/J_1 = 3(0)$ with a frequency spacing of $\delta\omega = 0.05$. Other parameters are given in Fig. 2.

h.c.), where $\mathcal{H}(\mathbf{q}) + \mathcal{H}_1$ contains the single-particle tight-binding Hamiltonian including a mean-field interaction contribution, and \mathcal{H}_2 stems exclusively from the interactions [66]. The Bogoliubov spectrum is shown in Fig. 2(c).

We test our proposal for extracting the symplectic quantum geometric tensor via excitation rates to other Bogoliubov modes by simulating the time evolution for initially populated Bogoliubov modes corresponding to the upper particle band. The parameter modulation (12) is induced by periodically shaking the Bogoliubov-Haldane model along the lattice directions x and y . In the reference frame co-moving with the lattice, the corresponding inertial forces directly induce modulations of the parameters q_x and q_y , respectively [66]. In Figs. 3(a)-(b) we present the ideal symplectic quantum metric $g_{xx}^+(\mathbf{q})$ and Berry curvature $B_{xy}^+(\mathbf{q})$ of the upper quasiparticle for $U_n/J_1 = 3$ together with the integrated rates (14),(16) obtained from the time evolution. In Figs. 3(c)-(d) we compare the ideal and integrated values for the same quantities along a high-symmetry path in quasimomentum space for $U_n/J_1 = 0$ (corresponding to a number-conserving model) and $U_n/J_1 = 3$. We find very good agreement between the ideal and integrated symplectic quantum geometric quantities, validating our scheme for extracting the components of the

SQGT. Results for $g_{xx}^+(\mathbf{q})$ and $g_{xy}^+(\mathbf{q})$ are presented in the SM [66]. Experimentally, the Bogoliubov excitation fraction can be measured by relating the presence of Bogoliubov quasiparticles to the quasimomentum distribution of the original particles, which can be observed using standard time-of-flight imaging techniques [66, 95, 96].

Symplectic anomalous velocity.— As a final result, we now show that, like in the number-conserving case [61–63, 97], a Bogoliubov lattice system acquires an anomalous velocity transverse to an applied homogeneous \mathbf{F} proportional to the symplectic Berry curvature. Consider a quasiparticle prepared in the Bogoliubov-Bloch mode $\mathbf{w}^n(\mathbf{q}_0)$. In response to the force, the quasimomentum is translated according to $\mathbf{q}(t) = \mathbf{q}_0 + \mathbf{F}t$, inducing an effective modulation of the dynamical matrix $D[\mathbf{q}(t)]$. In first-order adiabatic perturbation theory with respect to $\mathbf{q} = \mathbf{F}$, the time evolved state is given by [66]

$$\mathbf{w}^n(\mathbf{q}) + i \sum_{m \neq n} \mathbf{w}^m(\mathbf{q}) \frac{(\mathbf{w}^m(\mathbf{q}))^\dagger (\nabla_{\mathbf{q}} \mathcal{M}) \mathbf{w}^n(\mathbf{q}) \cdot \mathbf{F}}{s_m [\tilde{\Omega}_n(\mathbf{q}) - \tilde{\Omega}_m(\mathbf{q})]^2}. \quad (17)$$

In first order, the expectation value of the Bogoliubov velocity operator $\mathbf{v}(\mathbf{q}) = \nabla_{\mathbf{q}} D(\mathbf{q})$ then reads

$$\langle \mathbf{v}(\mathbf{q}) \rangle = \nabla_{\mathbf{q}} \tilde{\Omega}_n(\mathbf{q}) + \mathbf{B}_n(\mathbf{q}) \times \mathbf{F}, \quad (18)$$

with $\mathbf{B}_n(\mathbf{q}) \equiv \epsilon_{\mu\nu\kappa} B_{\mu\nu}^n(\mathbf{q}) \mathbf{e}_\kappa$. The first term corresponds to the slope of the occupied Bogoliubov energy band and the second term provides a transverse velocity component proportional to the symplectic Berry curvature, which is the symplectic generalization of the anomalous velocity of number-conserving systems [98]. The symplectic

anomalous velocity can be measured in the semiclassical dynamics of Bogoliubov-Bloch wave packets.

Conclusion.— We have introduced the manifestly gauge-invariant symplectic quantum geometric tensor (SQGT) for BBdG systems, whose imaginary part is proportional to the previously studied symplectic Berry curvature. The real part of the SQGT defines a symplectic quantum metric, which is shown to provide a natural distance measure in the space of Bogoliubov modes. All components of the symplectic quantum geometric tensor can be measured via tracking Bogoliubov excitation rates and the symplectic Berry curvature describes an anomalous velocity in Bogoliubov Bloch systems. State-of-the-art experiments with bosonic ultracold atoms [99–102] and nonlinear photonic crystal arrays [17, 40, 103–108] are well-suited to test our proposals. For future work, it will be interesting to connect recent topological notions for mixed quantum states [109–111] to thermal BBdG systems and to see whether invariants as the Euler characteristic [112–114] may be generalized to the BBdG context.

ACKNOWLEDGMENTS

We thank Christof Weitenberg, Luca Asteria, Nathan Harshman and Alexander Schnell for fruitful and insightful discussions. Part of the work has been conducted using the QuSpin project [115, 116]. This research was funded by the Deutsche Forschungsgemeinschaft via the Research Unit FOR 2414 under Project No. 277974659. I. T. acknowledges support from the Studienstiftung des deutschen Volkes.

-
- [1] N. N. Bogolyubov, *J. Phys. (USSR)* **11**, 23 (1947).
 - [2] J. Colpa, *Physica A: Statistical Mechanics and its Applications* **93**, 327 (1978).
 - [3] Y. Kawaguchi and M. Ueda, *Physics Reports* **520**, 253 (2012).
 - [4] R. Ozeri, N. Katz, J. Steinhauer, and N. Davidson, *Rev. Mod. Phys.* **77**, 187 (2005).
 - [5] A. A. Clerk, M. H. Devoret, S. M. Girvin, F. Marquardt, and R. J. Schoelkopf, *Rev. Mod. Phys.* **82**, 1155 (2010).
 - [6] T. Ozawa, H. M. Price, A. Amo, N. Goldman, M. Hafezi, L. Lu, M. C. Rechtsman, D. Schuster, J. Simon, O. Zilberberg, and I. Carusotto, *Rev. Mod. Phys.* **91**, 015006 (2019).
 - [7] R. Shindou, R. Matsumoto, S. Murakami, and J.-i. Ohe, *Phys. Rev. B* **87**, 174427 (2013).
 - [8] R. Shindou, J.-i. Ohe, R. Matsumoto, S. Murakami, and E. Saitoh, *Phys. Rev. B* **87**, 174402 (2013).
 - [9] P. A. McClarty, *Annu. Rev. Condens. Matter Phys.* **13**, 171 (2022).
 - [10] L. Zhang, J. Ren, J.-S. Wang, and B. Li, *Phys. Rev. Lett.* **105**, 225901 (2010).
 - [11] E. E. Wollman, C. U. Lei, A. J. Weinstein, J. Suh, A. Kronwald, F. Marquardt, A. A. Clerk, and K. C. Schwab, *Science* **349**, 952 (2015).
 - [12] M.-w. Xiao, Theory of transformation for the diagonalization of quadratic Hamiltonians (2009), [arXiv:0908.0787](https://arxiv.org/abs/0908.0787).
 - [13] M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045 (2010).
 - [14] X.-L. Qi and S.-C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011).
 - [15] Y. Hatsugai, *Phys. Rev. Lett.* **71**, 3697 (1993).
 - [16] Y. Hatsugai, *Phys. Rev. B* **48**, 11851 (1993).
 - [17] V. Peano, M. Houde, C. Brendel, F. Marquardt, and A. A. Clerk, *Nat Commun* **7**, 10779 (2016).
 - [18] R. Barnett, *Phys. Rev. A* **88**, 063631 (2013).
 - [19] B. Galilo, D. K. K. Lee, and R. Barnett, *Phys. Rev. Lett.* **115**, 245302 (2015).
 - [20] G. Engelhardt and T. Brandes, *Phys. Rev. A* **91**, 053621 (2015).
 - [21] S. Furukawa and M. Ueda, *New J. Phys.* **17**, 115014 (2015).
 - [22] A. Altland and M. R. Zirnbauer, *Phys. Rev. B* **55**, 1142 (1997).
 - [23] S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. W. Ludwig, *New J. Phys.* **12**, 065010 (2010).

- [24] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, *Phys. Rev. B* **78**, 195125 (2008).
- [25] A. Kitaev, *AIP Conference Proceedings* **1134**, 22 (2009).
- [26] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, *Rev. Mod. Phys.* **88**, 035005 (2016).
- [27] V. Peano and H. Schulz-Baldes, *Journal of Mathematical Physics* **59**, 031901 (2018).
- [28] M. Lein and K. Sato, *Phys. Rev. B* **100**, 075414 (2019).
- [29] Z. Zhou, L.-L. Wan, and Z.-F. Xu, *J. Phys. A: Math. Theor.* **53**, 425203 (2020).
- [30] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, *Phys. Rev. X* **8**, 031079 (2018).
- [31] K. Kawabata, K. Shiozaki, M. Ueda, and M. Sato, *Phys. Rev. X* **9**, 041015 (2019).
- [32] H. Zhou and J. Y. Lee, *Phys. Rev. B* **99**, 235112 (2019).
- [33] S. Lieu, *Phys. Rev. B* **98**, 115135 (2018).
- [34] D. Bernard and A. LeClair, in *Statistical Field Theories*, edited by A. Cappelli and G. Mussardo (Springer Netherlands, Dordrecht, 2002) pp. 207–214.
- [35] X.-Q. Wang, G.-Q. Luo, J.-Y. Liu, W. V. Liu, A. Hemmerich, and Z.-F. Xu, *Nature* **596**, 227 (2021).
- [36] Z. Jalali-mola, T. Grass, V. Kasper, M. Lewenstein, and U. Bhattacharya, *Phys. Rev. Lett.* **131**, 226601 (2023).
- [37] H. Kondo, Y. Akagi, and H. Katsura, *Phys. Rev. B* **99**, 041110 (2019).
- [38] T. Ohashi, S. Kobayashi, and Y. Kawaguchi, *Phys. Rev. A* **101**, 013625 (2020).
- [39] C.-E. Bardyn, T. Karzig, G. Refael, and T. C. H. Liew, *Phys. Rev. B* **93**, 020502 (2016).
- [40] V. Peano, M. Houde, F. Marquardt, and A. A. Clerk, *Phys. Rev. X* **6**, 041026 (2016).
- [41] A. McDonald, T. Pereg-Barnea, and A. A. Clerk, *Phys. Rev. X* **8**, 041031 (2018).
- [42] N. Okuma and M. Sato, *Annu. Rev. Condens. Matter Phys.* **14**, 83 (2023).
- [43] L.-L. Wan, Z. Zhou, and Z.-F. Xu, *Phys. Rev. A* **103**, 013308 (2021).
- [44] G. Chaudhary, M. Levin, and A. A. Clerk, *Phys. Rev. B* **103**, 214306 (2021).
- [45] Q.-R. Xu, V. P. Flynn, A. Alase, E. Cobanera, L. Viola, and G. Ortiz, *Phys. Rev. B* **102**, 125127 (2020).
- [46] V. P. Flynn, E. Cobanera, and L. Viola, *EPL* **131**, 40006 (2020).
- [47] G.-H. Huang, G.-Q. Luo, Z. Wu, and Z.-F. Xu, *Phys. Rev. A* **103**, 043328 (2021).
- [48] G.-H. Huang, Z.-F. Xu, and Z. Wu, *Phys. Rev. Lett.* **129**, 185301 (2022).
- [49] C. Chen, G.-H. Huang, and Z. Wu, *Phys. Rev. Res.* **5**, 023070 (2023).
- [50] S. Ravets, N. Pernet, N. Mostaan, N. Goldman, and J. Bloch, Thouless pumping in a driven-dissipative Kerr resonator array (2024), [arXiv:2407.02627](https://arxiv.org/abs/2407.02627).
- [51] A. Julku, G. M. Bruun, and P. Törmä, *Phys. Rev. B* **104**, 144507 (2021).
- [52] A. Julku, G. M. Bruun, and P. Törmä, *Phys. Rev. Lett.* **127**, 170404 (2021).
- [53] A. Julku, G. Salerno, and P. Törmä, *Low Temperature Physics* **49**, 701 (2023).
- [54] G. Salerno, T. Ozawa, and P. Törmä, *Phys. Rev. B* **108**, L140503 (2023).
- [55] R. Resta, *Eur. Phys. J. B* **79**, 121 (2011).
- [56] M. Kolodrubetz, D. Sels, P. Mehta, and A. Polkovnikov, *Physics Reports* **697**, 1 (2017).
- [57] J. P. Provost and G. Vallee, *Commun. Math. Phys.* **76**, 289 (1980).
- [58] We note that non-hermitian generalizations of the quantum geometric tensor have also recently been put forward [117–119], although not explicitly applied to the case of bosonic BdG systems.
- [59] D. T. Tran, A. Dauphin, A. G. Grushin, P. Zoller, and N. Goldman, *Science Advances* **3**, e1701207 (2017).
- [60] T. Ozawa and N. Goldman, *Phys. Rev. B* **97**, 201117 (2018).
- [61] D. Xiao, M.-C. Chang, and Q. Niu, *Rev. Mod. Phys.* **82**, 1959 (2010).
- [62] G. Sundaram and Q. Niu, *Phys. Rev. B* **59**, 14915 (1999).
- [63] M.-C. Chang and Q. Niu, *Phys. Rev. Lett.* **75**, 1348 (1995).
- [64] Throughout this letter, we assume a (thermodynamically) stable system, i.e., the coefficient matrix \mathcal{M} is positive definite such that a diagonalization of the BdG Hamiltonian (1) satisfying the bosonic commutation relations with strictly positive eigenvalues is always possible [2, 65, 120] as dictated by Williamson’s theorem [121, 122].
- [65] V. P. Flynn, E. Cobanera, and L. Viola, *New J. Phys.* **22**, 083004 (2020).
- [66] See supplementary material, which includes Refs. [2, 7, 12, 17, 20, 21, 28, 45, 56, 57, 59–63, 65, 67, 69, 92, 95, 97, 102, 121–136], where we provide details regarding: the paraunitary Bogoliubov transformation and the associated indefinite inner product space, the derivation of alternative representations of the SQGT, the relation between the symplectic quantum metric and the infinitesimal distance measure, the local conservation law for the symplectic Berry curvature, the derivation of the equation of motion for Bogoliubov modes and its solution using time-dependent perturbation theory, the application of the measurement protocol to BBdG lattice systems, the multi-quasiparticle case in Fock space, adiabatic perturbation theory of the equation of motion, the derivation of the symplectic anomalous velocity, the Bogoliubov-Haldane model, and observable signatures for the original particles.
- [67] J. Dereziński, *Journal of Mathematical Physics* **58**, 121101 (2017).
- [68] J.-P. Blaizot and G. Ripka, *Quantum Theory of Finite Systems* (MIT Press, Cambridge, Mass., 1986).
- [69] H. Schulz-Baldes and C. Villegas-Blas, *Mathematische Nachrichten* **290**, 1840 (2017).
- [70] Y. Aharonov and J. Anandan, *Phys. Rev. Lett.* **58**, 1593 (1987).
- [71] J. Anandan and Y. Aharonov, *Phys. Rev. Lett.* **65**, 1697 (1990).
- [72] G. Fubini, *Sulle Metriche Definite Da Una Forma Hermitiana: Nota* (Office graf. C. Ferrari, 1904).
- [73] E. Study, *Math. Ann.* **60**, 321 (1905).
- [74] P. Zanardi, P. Giorda, and M. Cozzini, *Phys. Rev. Lett.* **99**, 100603 (2007).
- [75] L. Campos Venuti and P. Zanardi, *Phys. Rev. Lett.* **99**, 095701 (2007).
- [76] A. T. Rezakhani, D. F. Abasto, D. A. Lidar, and P. Zanardi, *Phys. Rev. A* **82**, 012321 (2010).
- [77] A. Carollo, D. Valenti, and B. Spagnolo, *Physics Reports* **838**, 1 (2020).
- [78] J. Liu, H. Yuan, X.-M. Lu, and X. Wang, *J. Phys. A:*

- Math. Theor.* **53**, 023001 (2020).
- [79] Y.-Q. Ma, S.-J. Gu, S. Chen, H. Fan, and W.-M. Liu, *EPL* **103**, 10008 (2013).
- [80] P. Hauke, M. Heyl, L. Tagliacozzo, and P. Zoller, *Nature Phys* **12**, 778 (2016).
- [81] T. Ozawa and N. Goldman, *Phys. Rev. Res.* **1**, 032019 (2019).
- [82] C.-R. Yi, J. Yu, H. Yuan, R.-H. Jiao, Y.-M. Yang, X. Jiang, J.-Y. Zhang, S. Chen, and J.-W. Pan, *Phys. Rev. Res.* **5**, L032016 (2023).
- [83] M. Yu, Y. Liu, P. Yang, M. Gong, Q. Cao, S. Zhang, H. Liu, M. Heyl, T. Ozawa, N. Goldman, and J. Cai, *npj Quantum Inf* **8**, 1 (2022).
- [84] M. Yu, X. Li, Y. Chu, B. Mera, F. N. Ünal, P. Yang, Y. Liu, N. Goldman, and J. Cai, *National Science Review*, nwae065 (2024).
- [85] X. Tan, D.-W. Zhang, Z. Yang, J. Chu, Y.-Q. Zhu, D. Li, X. Yang, S. Song, Z. Han, Z. Li, Y. Dong, H.-F. Yu, H. Yan, S.-L. Zhu, and Y. Yu, *Phys. Rev. Lett.* **122**, 210401 (2019).
- [86] M. Yu, P. Yang, M. Gong, Q. Cao, Q. Lu, H. Liu, S. Zhang, M. B. Plenio, F. Jelezko, T. Ozawa, N. Goldman, and J. Cai, *National Science Review* **7**, 254 (2020).
- [87] A. Gianfrate, O. Bleu, L. Dominici, V. Ardizzone, M. De Giorgi, D. Ballarini, G. Lerario, K. W. West, L. N. Pfeiffer, D. D. Solnyshkov, D. Sanvitto, and G. Malpuech, *Nature* **578**, 381 (2020).
- [88] W. Zheng, J. Xu, Z. Ma, Y. Li, Y. Dong, Y. Zhang, X. Wang, G. Sun, P. Wu, J. Zhao, S. Li, D. Lan, X. Tan, and Y. Yu, *Chinese Phys. Lett.* **39**, 100202 (2022).
- [89] J. Cuerda, J. M. Taskinen, N. Källman, L. Grabitz, and P. Törmä, *Phys. Rev. Res.* **6**, L022020 (2024).
- [90] L. Asteria, D. T. Tran, T. Ozawa, M. Tarnowski, B. S. Rem, N. Fläschner, K. Sengstock, N. Goldman, and C. Weitenberg, *Nat. Phys.* **15**, 449 (2019).
- [91] D. M. Stamper-Kurn, A. P. Chikkatur, A. Görlitz, S. Inouye, S. Gupta, D. E. Pritchard, and W. Ketterle, *Phys. Rev. Lett.* **83**, 2876 (1999).
- [92] F. D. M. Haldane, *Phys. Rev. Lett.* **61**, 2015 (1988).
- [93] G. Jotzu, M. Messer, R. Desbuquois, M. Lebrat, T. Uehlinger, D. Greif, and T. Esslinger, *Nature* **515**, 237 (2014).
- [94] M. Tarnowski, F. N. Ünal, N. Fläschner, B. S. Rem, A. Eckardt, K. Sengstock, and C. Weitenberg, *Nat Commun* **10**, 1728 (2019).
- [95] J. M. Vogels, K. Xu, C. Raman, J. R. Abo-Shaeer, and W. Ketterle, *Phys. Rev. Lett.* **88**, 060402 (2002).
- [96] A. Brunello, F. Dalfovo, L. Pitaevskii, and S. Stringari, *Phys. Rev. Lett.* **85**, 4422 (2000).
- [97] C. Zhang, A. M. Dudarev, and Q. Niu, *Phys. Rev. Lett.* **97**, 040401 (2006).
- [98] We note that the dynamics of Bogoliubov Bloch wave packets have also been studied using a time-dependent variational principle in Ref. [97], but no connection to the symplectic Berry curvature has been made therein.
- [99] N. R. Cooper, J. Dalibard, and I. B. Spielman, *Rev. Mod. Phys.* **91**, 015005 (2019).
- [100] C. Gross and I. Bloch, *Science* **357**, 995 (2017).
- [101] F. Schäfer, T. Fukuhara, S. Sugawa, Y. Takasu, and Y. Takahashi, *Nat Rev Phys* **2**, 411 (2020).
- [102] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
- [103] M. Notomi, E. Kuramochi, and T. Tanabe, *Nature Photon* **2**, 741 (2008).
- [104] S. Mookherjea and A. Yariv, *IEEE Journal of Selected Topics in Quantum Electronics* **8**, 448 (2002).
- [105] B. J. Eggleton, B. Luther-Davies, and K. Richardson, *Nature Photon* **5**, 141 (2011).
- [106] J. Dahdah, M. Pilar-Bernal, N. Courjal, G. Ulliac, and F. Baida, *Journal of Applied Physics* **110**, 074318 (2011).
- [107] D. Smirnova, D. Leykam, Y. Chong, and Y. Kivshar, *Applied Physics Reviews* **7**, 021306 (2020).
- [108] T. Shi, H. J. Kimble, and J. I. Cirac, *Proceedings of the National Academy of Sciences* **114**, E8967 (2017).
- [109] C.-E. Bardyn, L. Wawer, A. Altland, M. Fleischhauer, and S. Diehl, *Phys. Rev. X* **8**, 011035 (2018).
- [110] L. Wawer and M. Fleischhauer, *Phys. Rev. B* **104**, 094104 (2021).
- [111] J. C. Budich and S. Diehl, *Phys. Rev. B* **91**, 165140 (2015).
- [112] A. Bouhon, Q. Wu, R.-J. Slager, H. Weng, O. V. Yazyev, and T. Bzdušek, *Nat. Phys.* **16**, 1137 (2020).
- [113] F. N. Ünal, A. Bouhon, and R.-J. Slager, *Phys. Rev. Lett.* **125**, 053601 (2020).
- [114] J. Ahn, S. Park, and B.-J. Yang, *Phys. Rev. X* **9**, 021013 (2019).
- [115] P. Weinberg and M. Bukov, *SciPost Physics* **2**, 003 (2017).
- [116] P. Weinberg and M. Bukov, *SciPost Physics* **7**, 020 (2019).
- [117] D.-J. Zhang, Q.-h. Wang, and J. Gong, *Phys. Rev. A* **99**, 042104 (2019).
- [118] Y.-Q. Zhu, W. Zheng, S.-L. Zhu, and G. Palumbo, *Phys. Rev. B* **104**, 205103 (2021).
- [119] C. Chen Ye, W. L. Vleeshouwers, S. Heatley, V. Gritsev, and C. Morais Smith, *Phys. Rev. Res.* **6**, 023202 (2024).
- [120] R. Rossignoli and A. M. Kowalski, *Phys. Rev. A* **72**, 032101 (2005).
- [121] J. Williamson, *American Journal of Mathematics* **58**, 141 (1936), 2371062.
- [122] F. Nicacio, *American Journal of Physics* **89**, 1139 (2021).
- [123] M. Ueda, *Fundamentals and New Frontiers of Bose-Einstein Condensation* (WORLD SCIENTIFIC, 2010).
- [124] C. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases*, 2nd ed. (Cambridge University Press, 2008).
- [125] J. Bognár, *Indefinite Inner Product Spaces* (Springer Berlin Heidelberg, Berlin, Heidelberg, 1974).
- [126] R. S. MacKay, in *Hamiltonian Dynamical Systems* (CRC Press, 1987).
- [127] B. Diu, F. Laloe, and C. Cohen-Tannoudji, *Quantum Mechanics: Volume II: Angular Momentum, Spin, and Approximation Methods*, 2nd ed. (Wiley-VCH, Weinheim, Germany, 2019).
- [128] R. Karplus and J. M. Luttinger, *Phys. Rev.* **95**, 1154 (1954).
- [129] G. Rigolin, G. Ortiz, and V. H. Ponce, *Phys. Rev. A* **78**, 052508 (2008).
- [130] C. De Grandi and A. Polkovnikov, in *Quantum Quenching, Annealing and Computation*, edited by A. K. Chandra, A. Das, and B. K. Chakrabarti (Springer, Berlin, Heidelberg, 2010) pp. 75–114.
- [131] M. V. Berry, *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences* **392**, 45

- (1984).
- [132] M. Kolodrubetz, V. Gritsev, and A. Polkovnikov, *Phys. Rev. B* **88**, 064304 (2013).
 - [133] H. M. Price, O. Zeitler, T. Ozawa, I. Carusotto, and N. Goldman, *Phys. Rev. B* **93**, 245113 (2016).
 - [134] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Rev. Mod. Phys.* **71**, 463 (1999).
 - [135] Y.-J. Wu, W.-Y. Zhou, and S.-P. Kou, *Phys. Rev. A* **95**, 023620 (2017).
 - [136] P. Hauke, M. Lewenstein, and A. Eckardt, *Phys. Rev. Lett.* **113**, 045303 (2014).
 - [137] H. Schulz-Baldes, *Integr. Equ. Oper. Theory* **78**, 323 (2014).

Supplementary Material for “Quantum geometry of bosonic Bogoliubov quasiparticles”

Isaac Tesfaye^{1,*} and André Eckardt^{1,†}

¹*Institut für Theoretische Physik, Technische Universität Berlin, Hardenbergstr. 36, D-10623 Berlin, Germany*

(Dated: July 17, 2024)

CONTENTS

Acknowledgments	5
References	5
S1. Paraunitary Bogoliubov transformation from bosonic commutation relations	1
S2. Indefinite Bogoliubov inner product space	3
S3. Symplectic quantum geometric tensor	4
S3A. Symplectic quantum metric	5
S3B. Local conservation law for symplectic Berry curvature	6
S4. Time-dependent Schrödinger equation for Bogoliubov modes	7
S5. Time-dependent perturbation theory for the effective Schrödinger equation	8
S6. Measuring the symplectic quantum geometry in lattice systems	9
S7. Time-dependent perturbation theory for many-body Bogoliubov Fock states	11
S7A. Transitions between Bogoliubov quasiparticle excitations lead to SQGT	11
S8. Adiabatic perturbation theory for the effective Schrödinger equation	13
S9. Symplectic anomalous velocity	15
S10. Bogoliubov Haldane model	16
S10A. Algorithm for constructing paraunitary Bogoliubov transformation matrix	20
S10B. SQGT from integrated Bogoliubov excitation rates	21
S10C. Signatures for the distribution of the original particles	22

S1. PARAUNITARY BOGOLIUBOV TRANSFORMATION FROM BOSONIC COMMUTATION RELATIONS

First, for ease of reference, we restate the bosonic BdG Hamiltonian [Eq. (1) in the main text] here

$$\hat{H} = \sum_{i,j=1}^N K_{ij} \hat{a}_i^\dagger \hat{a}_j + \frac{1}{2} \left(G_{ij} \hat{a}_i^\dagger \hat{a}_j^\dagger + G_{ji}^* \hat{a}_i \hat{a}_j \right) = \frac{1}{2} \hat{\Psi}^\dagger \mathcal{M} \hat{\Psi} - \frac{1}{2} \text{Tr} \{K\}, \quad \mathcal{M} = \begin{pmatrix} K & G \\ G^* & K^* \end{pmatrix}. \quad (\text{S1})$$

* i.tesfaye@tu-berlin.de

† eckardt@tu-berlin.de

The hermiticity of $\hat{H} = \hat{H}^\dagger$ imposes K to be hermitian, i.e., $K_{ij} = K_{ji}^* \in \mathbb{C}$, and the bosonic commutation relations for $\hat{a}_i^{(\dagger)}$ lead to G being symmetric, i.e., $G_{ij} = G_{ji} \in \mathbb{C}$ [2, 12, 65]. In the second equality, the BdG Hamiltonian is reformulated in terms of the bosonic Nambu spinors $\hat{\Psi}^{(\dagger)}$,

$$\hat{\Psi}^\dagger = (\hat{a}_1^\dagger \cdots \hat{a}_N^\dagger \hat{a}_1 \cdots \hat{a}_N), \quad \hat{\Psi} = (\hat{a}_1 \cdots \hat{a}_N \hat{a}_1^\dagger \cdots \hat{a}_N^\dagger)^T, \quad (\text{S2})$$

and the $2N \times 2N$ coefficient matrix \mathcal{M} consisting of $N \times N$ matrix blocks K and G . The bosonic operators $\hat{a}_i^{(\dagger)}$ in (S1) preserve the bosonic commutation relations (BCR),

$$[\hat{a}_i, \hat{a}_j] = 0, \quad [\hat{a}_i, \hat{a}_j^\dagger] = \delta_{ij}, \Leftrightarrow [\hat{\Psi}_i, \hat{\Psi}_j^\dagger] = (\tau_z)_{ij}, \quad \tau_z = \begin{pmatrix} \mathbb{1}_N & 0_N \\ 0_N & -\mathbb{1}_N \end{pmatrix}, \quad (\text{S3})$$

where in the last equation, we have reformulated the bosonic commutation relations via the Nambu spinors (S2) with $\tau_z = \sigma_z \otimes \mathbb{1}_N$, where $\sigma_z = \text{diag}(1, -1)$ is the Pauli-z matrix and $\mathbb{1}_N$ (0_N) is the $N \times N$ identity (zero) matrix. Now, our goal is to find a (Bogoliubov) transformation from the original set of bosonic operators $\hat{a}_i^{(\dagger)}$ to a new set of bosonic operators $\hat{b}_n^{(\dagger)}$ that diagonalizes the Hamiltonian (S1).

Let us first discuss the constraints on this transformation. Importantly, this transformation must preserve the BCR (S3), i.e., the BCR must also hold for the new bosonic operators $\hat{b}_n^{(\dagger)}$,

$$[\hat{b}_n, \hat{b}_m] = 0, \quad [\hat{b}_n, \hat{b}_m^\dagger] = \delta_{nm}, \Leftrightarrow [\hat{\Phi}_n, \hat{\Phi}_m^\dagger] = (\tau_z)_{nm}, \quad (\text{S4})$$

where we defined $\hat{\Phi} = (\hat{b}_1 \cdots \hat{b}_N \hat{b}_1^\dagger \cdots \hat{b}_N^\dagger)^T$ and $\hat{\Phi}^\dagger = (\hat{b}_1^\dagger \cdots \hat{b}_N^\dagger \hat{b}_1 \cdots \hat{b}_N)$ as the bosonic Nambu spinors (S2) for the new bosonic operators $\hat{b}_n^{(\dagger)}$.

We parametrize such a general Bogoliubov transformation from $\hat{a}_i^{(\dagger)}$ to $\hat{b}_n^{(\dagger)}$ by [2, 3, 68]

$$\hat{a}_i = \sum_n U_{in} \hat{b}_n + (V^*)_{in} \hat{b}_n^\dagger, \quad \hat{a}_i^\dagger = \sum_n \hat{b}_n^\dagger (U^\dagger)_{ni} + \hat{b}_n (V^T)_{ni}, \quad (\text{S5})$$

$$\hat{\Psi} = W \hat{\Phi}, \quad \hat{\Psi}^\dagger = \hat{\Phi}^\dagger W^\dagger, \quad W = \begin{pmatrix} U & V^* \\ V & U^* \end{pmatrix}, \quad W^\dagger = \begin{pmatrix} U^\dagger & V^\dagger \\ V^T & U^T \end{pmatrix}, \quad (\text{S6})$$

where in the second line we have reformulated the Bogoliubov transformation via a transformation between the original and new Nambu spinors, $\hat{\Psi}$ and $\hat{\Phi}$ respectively.

The requirement that also the $\hat{b}_n^{(\dagger)}$ are bosonic operators (S4) will impose constraints on the complex $2N \times 2N$ transformation matrix W , or equivalently on its constituents, the $N \times N$ complex matrices U and V . Using (S5) these constraints from the BCR (S3)-(S4) are given by

$$\delta_{ij} = [\hat{a}_i, \hat{a}_j^\dagger] = \sum_{n, n'} [(U_{in} \hat{b}_n + (V^*)_{in} \hat{b}_n^\dagger), (\hat{b}_{n'}^\dagger (U^\dagger)_{n'j} + \hat{b}_{n'} (V^T)_{n'j})] = \sum_n (U_{in} (U^\dagger)_{nj} - (V^*)_{in} (V^T)_{nj}), \quad (\text{S7})$$

$$0 = [\hat{a}_i, \hat{a}_j] = \sum_{n, n'} [(U_{in} \hat{b}_n + (V^*)_{in} \hat{b}_n^\dagger), (U_{jn'} \hat{b}_{n'} + (V^*)_{jn'} \hat{b}_{n'}^\dagger)] = \sum_n (U_{in} (V^*)_{jn} - (V^*)_{in} (U)_{jn}). \quad (\text{S8})$$

In matrix form, these conditions read

$$UU^\dagger - (V^*)V^T = \mathbb{1}_N, \quad V^*U^T - UV^\dagger = 0_N, \quad (\text{S9})$$

where $\mathbb{1}_N$ (0_N) is again the $N \times N$ identity (zero) matrix. We can also formulate the conditions for the entire transformation matrix W (S6) by using BCR in terms of the Nambu spinors (S3)-(S4)

$$(\tau_z)_{ij} = [\hat{\Psi}_i, \hat{\Psi}_j^\dagger] = \sum_{m, m'} [(W_{im} \hat{\Phi}_m), (\hat{\Phi}_{m'}^\dagger (W^\dagger)_{m'j})] = \sum_{m, m'} W_{im} (W^\dagger)_{m'j} [\hat{\Phi}_m, \hat{\Phi}_{m'}^\dagger] = \sum_{m, m'} W_{im} (\tau_z)_{mm'} (W^\dagger)_{m'j}. \quad (\text{S10})$$

In matrix form these conditions for the full transformation matrix W read

$$\tau_z = W \tau_z W^\dagger = W^\dagger \tau_z W, \quad (\text{S11})$$

which lead to Eq. (3) in the main text. Note that we have added a second equality in (S11) in addition to the first from (S10). To see that the second equality holds, note that the first equality, $\tau_z = W \tau_z W^\dagger$, is equivalent to

$W^\dagger = \tau_z W^{-1} \tau_z$. Then we immediately see that $W^\dagger \tau_z W = (\tau_z W^{-1} \tau_z) \tau_z W = \tau_z$ proving the second equality in (S11). The Bogoliubov transformation matrices W which fulfill the constraints of Eq. (S11) are called *paraunitary* matrices [2].

Conditions (S11) further allow us to obtain the useful relations for the inverse transformation of (S6), from the new bosonic operators $\hat{b}_n^{(\dagger)}$ to the original bosonic operators $\hat{a}_i^{(\dagger)}$ via

$$\hat{b}_n = \sum_i (U^\dagger)_{ni} \hat{a}_i - (V^\dagger)_{ni} \hat{a}_i^\dagger, \quad \hat{b}_n^\dagger = \sum_i \hat{a}_i^\dagger U_{in} - \hat{a}_i V_{in}, \quad (\text{S12})$$

$$\hat{\Phi} = W^{-1} \hat{\Psi}, \quad \hat{\Phi}^\dagger = \hat{\Psi}^\dagger (W^\dagger)^{-1} = \hat{\Psi}^\dagger \tau_z W \tau_z, \quad (\text{S13})$$

$$W^{-1} = \tau_z W^\dagger \tau_z = \begin{pmatrix} U^\dagger & -V^\dagger \\ -V^T & U^T \end{pmatrix}, \quad (W^\dagger)^{-1} = \tau_z W \tau_z = \begin{pmatrix} U & -V^* \\ -V & U^* \end{pmatrix}. \quad (\text{S14})$$

If we denote the n -th column of $W = (\mathbf{w}^1, \dots, \mathbf{w}^{2N})$ by \mathbf{w}^n , we can directly see from (S12) that the new bosonic quasiparticle operators $\hat{b}_n^{(\dagger)}$ can be compactly expressed as

$$\hat{b}_n = (\mathbf{w}^n)^\dagger \tau_z \hat{\Psi}, \quad \hat{b}_n^\dagger = \hat{\Psi}^\dagger \tau_z \mathbf{w}^n, \quad \text{for } n \leq N, \text{ or equivalently} \quad (\text{S15})$$

$$\hat{b}_n = -\hat{\Psi}^\dagger \tau_z \mathbf{w}^{N+n}, \quad \hat{b}_n^\dagger = -(\mathbf{w}^{N+n})^\dagger \tau_z \hat{\Psi}, \quad \text{for } n > N. \quad (\text{S16})$$

Having discussed the constraints on the Bogoliubov transformation matrix W , we turn to the diagonalization of the BdG Hamiltonian (S1). With the Bogoliubov transformation (S6) the Hamiltonian (S1) can be written in terms of the new bosonic quasiparticle operators $\hat{b}_n^{(\dagger)}$,

$$\hat{H} = \frac{1}{2} \hat{\Phi}^\dagger W^\dagger \mathcal{M} W \hat{\Phi} - \frac{1}{2} \text{Tr} \{K\}. \quad (\text{S17})$$

For the Hamiltonian (S1) to become diagonal in $\hat{b}_n^{(\dagger)}$, the coefficient matrix \mathcal{M} must become diagonal via the Bogoliubov transformation,

$$W^\dagger \mathcal{M} W = \Omega = \begin{pmatrix} \boldsymbol{\omega} & 0 \\ 0 & \boldsymbol{\omega} \end{pmatrix}, \quad \boldsymbol{\omega} = \text{diag}(\omega_1, \dots, \omega_N), \quad (\text{S18})$$

where Ω is the $2N \times 2N$ diagonal matrix consisting of the $N \times N$ diagonal matrix $\boldsymbol{\omega}$ with Bogoliubov excitation energies ω_n . As dictated by Williamson's theorem [121, 122] such a diagonalization (S18) via Bogoliubov matrices W (S5) is always possible for a positive definite coefficient matrix \mathcal{M} [2, 68]. This then eventually leads to the diagonal form of the Hamiltonian (S1)

$$\hat{H} = \frac{1}{2} \hat{\Phi}^\dagger \Omega \hat{\Phi} - \frac{1}{2} \text{Tr} \{K\} = \sum_{n=1}^N \omega_n \left(\hat{b}_n^\dagger \hat{b}_n + \frac{1}{2} \right) - \frac{1}{2} \text{Tr} \{K\} = \sum_{n=1}^N \omega_n \hat{b}_n^\dagger \hat{b}_n + C, \quad (\text{S19})$$

with $C \equiv \sum_n \omega_n / 2 - \text{Tr} \{K\} / 2$. The ground state $|\psi_0\rangle$ of this Hamiltonian (S19) is generally given by a multi-mode squeezed state with a non-zero number of original $\hat{a}_i^{(\dagger)}$ bosons, which is also called the Bogoliubov vacuum. This Bogoliubov vacuum is defined by the condition that it contains no Bogoliubov quasiparticle excitations, i.e., $\hat{b}_n |\psi_0\rangle = 0$ for all n [123, 124].

S2. INDEFINITE BOGOLIUBOV INNER PRODUCT SPACE

We collect some useful relations for the indefinite inner product space spanned by the Bogoliubov modes \mathbf{w}^n . For a more rigorous introduction into these concepts pertaining indefinite inner product spaces, we refer to Ref. [125]. Given a Bogoliubov transformation matrix $W \in \mathbb{C}^{2N \times 2N}$ which satisfies (S11)

$$W \tau_z W^\dagger = W^\dagger \tau_z W = \tau_z \quad (\text{S20})$$

and whose n -th column corresponds to the Bogoliubov mode \mathbf{w}^n , we can define a projector P^n onto the n -th non-degenerate Bogoliubov eigenspace (and its complement Q^n) (5)

$$P^n = W \Gamma^n W^{-1} = W \Gamma^n \tau_z W^\dagger \tau_z = s_n \mathbf{w}^n (\mathbf{w}^n)^\dagger \tau_z, \quad Q^n = \mathbb{1}_{2N} - P^n, \quad (\text{S21})$$

with $(\Gamma^n)_{ml} = \delta_{nm}\delta_{ml}$ being the $2N \times 2N$ matrix whose only non-zero element is the n -th diagonal element being equal to $+1$ and where $s_n = 1$ (-1) for $n \leq N$ ($n > N$) for the so-called particle (hole) states. We note that these projectors X^n are necessarily invariant under (local) $U(1)$ gauge transformations for the Bogoliubov modes of the form $\mathbf{w}^n \rightarrow e^{i\phi} \mathbf{w}^n$, i.e., $X^n \rightarrow X^n$ for $X^n \in \{P^n, Q^n\}$ (S21).

Both P^n and Q^n are Krein-hermitian (pseudo-hermitian) $(X^n)^\dagger = \tau_z X^n \tau_z$ for $X^n \in \{P^n, Q^n\}$ [28, 69], which directly follows from (S20). Moreover, the projectors are idempotent $P^n P^k = \delta_{nk} P^n$ and span the entire inner product space $\sum_n P^n = \mathbb{1}_{2N}$. This allows for a resolution of the identity in the indefinite Bogoliubov inner product space,

$$\mathbb{1}_{2N} = \sum_{n=1}^{2N} P^n = \sum_{n=1}^{2N} s_n \mathbf{w}^n (\mathbf{w}^n)^\dagger \tau_z = \sum_{l=1}^N (\mathbf{w}_+^l (\mathbf{w}_+^l)^\dagger - \mathbf{w}_-^l (\mathbf{w}_-^l)^\dagger) \tau_z, \quad (\text{S22})$$

where in the second equality we have separated the Bogoliubov modes into so-called particle \mathbf{w}_+^l and hole parts \mathbf{w}_-^l [45], which correspond to the first N and last N columns of the Bogoliubov transformation matrix W respectively. Furthermore, note that the particle and hole parts are not independent but are directly related via $\mathbf{w}_-^l = \tau_x (\mathbf{w}_+^l)^*$, with $\tau_x = \sigma_x \otimes \mathbb{1}_N$, where σ_x is the Pauli-x matrix, which can also be seen from the parametrization of W (S6).

The Bogoliubov modes \mathbf{w}^n adopt a generalized orthonormalization condition [2, 7, 12, 45] following from (S20),

$$(\mathbf{w}^n)^\dagger \tau_z \mathbf{w}^m = s_n \delta_{nm}, \Leftrightarrow (\mathbf{w}_\pm^l)^\dagger \tau_z \mathbf{w}_\pm^k = \pm \delta_{lk}, \quad (\mathbf{w}_\pm^l)^\dagger \tau_z \mathbf{w}_\mp^k = 0. \quad (\text{S23})$$

Next, note that the trace of a matrix $A \in \mathbb{C}^{2N \times 2N}$ can be evaluated in any basis which spans the entire \mathbb{C}^{2N} . The Bogoliubov modes are such a basis which can be seen from the resolution of the identity (S22). Thus, the trace of a matrix A can be expressed in terms of the Bogoliubov modes as

$$\begin{aligned} \text{Tr}\{A\} &= \sum_{n,m} \text{Tr}\{P^n A P^m\} = \sum_{n,m,i} (\mathbf{e}_i)^\dagger (P^n A P^m) \mathbf{e}_i = \sum_{n,m,i} s_n s_m [(\mathbf{e}_i)^\dagger \mathbf{w}^n (\mathbf{w}^n)^\dagger \tau_z A \mathbf{w}^m (\mathbf{w}^m)^\dagger \tau_z \mathbf{e}_i] \\ &= \sum_{n,m} s_n s_m [(\mathbf{w}^n)^\dagger \tau_z A \mathbf{w}^m (\mathbf{w}^m)^\dagger \tau_z (\sum_i \mathbf{e}_i (\mathbf{e}_i)^\dagger) \mathbf{w}^n] = \sum_n s_n [(\mathbf{w}^n)^\dagger \tau_z A \mathbf{w}^n], \end{aligned} \quad (\text{S24})$$

where \mathbf{e}_i are the standard basis vectors in \mathbb{C}^{2N} . In the last equality of (S24) we have further used that $\mathbb{1}_{2N} = \sum_i \mathbf{e}_i (\mathbf{e}_i)^\dagger$ and the orthonormalization condition of the Bogoliubov modes (S23).

Any vector $\psi \in \mathbb{C}^{2N}$ can be expanded in the basis of the Bogoliubov modes by making use of the resolution of identity (S22), such that

$$\psi = \sum_{j=1}^{2N} P^j \psi = \sum_{n=1}^{2N} s_n \mathbf{w}^n (\mathbf{w}^n)^\dagger \tau_z \psi \equiv \sum_n c_n^\psi \mathbf{w}^n, \quad (\text{S25})$$

where $c_n^\psi = s_n (\mathbf{w}^n)^\dagger \tau_z \psi$. Moreover, since ψ lives in the indefinite inner product space, its norm is only correctly defined via the generalized inner product $(\psi)^\dagger \tau_z \psi = \pm 1$, which is also called Krein signature [126, 137]. Using the generalized orthonormality condition (S23) this norm can consequently also be expressed as

$$(\psi)^\dagger \tau_z \psi = \sum_{nm} (c_n^\psi)^* c_m^\psi (\mathbf{w}^n)^\dagger \tau_z \mathbf{w}^m = \sum_{nm} (c_n^\psi)^* c_m^\psi s_m \delta_{nm} = \sum_{n=1}^{2N} s_n |c_n^\psi|^2 = \pm 1, \quad (\text{S26})$$

which may be viewed as a generalization of Parseval's identity for this indefinite inner product space.

S3. SYMPLECTIC QUANTUM GEOMETRIC TENSOR

In addition to the manifestly gauge-invariant form of the symplectic quantum geometric tensor $\eta_{\mu\nu}^n = \text{Tr}\{\partial_\mu P^n Q^n \partial_\nu P^n\}$ [Eq. (6) in the main text], we derive two equivalent representations of the SQGT, which are more suitable for practical calculations. We again assume implicitly that the BdG Hamiltonian (S1) depends on a set of λ parameters and all derivatives should be understood as derivatives with respect to these parameters, i.e., $\partial_\mu \equiv \frac{\partial}{\partial \lambda^\mu}$.

First, note that the diagonalization of the BdG Hamiltonian (S1) via the paraunitary matrix W (S6) leads to the diagonalization of the positive definite coefficient matrix \mathcal{M} (S18), $W^\dagger \mathcal{M} W = \Omega$. This in turn corresponds to the

generalized eigenvalue equation [2, 12],

$$\mathcal{M}W = (W^\dagger)^{-1}\Omega = \tau_z W \tau_z \Omega = \tau_z W \tilde{\Omega}, \text{ with } \tilde{\Omega} \equiv \tau_z \Omega = \begin{pmatrix} \omega & 0 \\ 0 & -\omega \end{pmatrix}, \quad (\text{S27})$$

$$\Leftrightarrow DW = W\tilde{\Omega}, \text{ with } D \equiv \tau_z \mathcal{M}, \quad (\text{S28})$$

which rewritten in terms of the Bogoliubov modes \mathbf{w}^n (n -th column of W) reads

$$D\mathbf{w}^n = \tilde{\Omega}_n \mathbf{w}^n, \text{ with } D = \tau_z \mathcal{M} = \begin{pmatrix} K & G \\ -G^* & -K^* \end{pmatrix}, \quad (\text{S29})$$

where $D \equiv \tau_z \mathcal{M}$ is the so-called dynamical matrix of the system and \mathbf{w}^n is the n -th Bogoliubov mode corresponding to the excitation energy $\tilde{\Omega}_n = s_n \omega_n \bmod N$ [7, 12]. Even though D is generally non-hermitian, the positive definiteness of \mathcal{M} ensures that the eigenvalues $\tilde{\Omega}_n$ are all real [2, 68].

Using the definition of the trace via the Bogoliubov modes (S24), we see that the SQGT $\eta_{\mu\nu}^n = \text{Tr} \{ \partial_\mu P^n Q^n \partial_\nu P^n \}$ can equivalently be expressed as

$$\begin{aligned} \eta_{\mu\nu}^n &= \sum_m s_m [(\mathbf{w}^m)^\dagger \tau_z (\partial_\mu P^n Q^n \partial_\nu P^n) \mathbf{w}^m] = \sum_m s_m [(\mathbf{w}^m)^\dagger \tau_z \mathbf{w}^n (\partial_\mu \mathbf{w}^n)^\dagger \tau_z Q^n (\partial_\nu \mathbf{w}^n) (\mathbf{w}^n)^\dagger \tau_z \mathbf{w}^m] \\ &= s_n (\partial_\mu \mathbf{w}^n)^\dagger \tau_z Q^n (\partial_\nu \mathbf{w}^n). \end{aligned} \quad (\text{S30})$$

Here, in the second equality we have used that $Q^n \mathbf{w}^n = (\mathbf{w}^n)^\dagger \tau_z Q^n = 0$ which directly follows from (S21). In the final equality, the orthonormality conditions (S23) has been used. Eq. (S30) is the first equivalent representation of the SQGT.

To obtain the spectral representation of the SQGT, we first derive a symplectic version of the Hellmann-Feynman theorem. To this end, we first observe from eigenvalue equation (S29) that

$$(\partial_\mu D)\mathbf{w}^n + D\partial_\mu \mathbf{w}^n = (\partial_\mu \tilde{\Omega}_n)\mathbf{w}^n + \tilde{\Omega}_n \partial_\mu \mathbf{w}^n, \quad (\text{S31})$$

Then we project a distinct ($m \neq n$) Bogoliubov mode $(\mathbf{w}^m)^\dagger \tau_z$ from the left onto (S31) and obtain with use of the orthonormality conditions (S23) the symplectic version of the Hellmann-Feynman theorem,

$$(\mathbf{w}^m)^\dagger (\partial_\mu \mathcal{M}) \mathbf{w}^n = (\tilde{\Omega}_n - \tilde{\Omega}_m) (\mathbf{w}^m)^\dagger \tau_z (\partial_\mu \mathbf{w}^n). \quad (\text{S32})$$

Next, by inserting two resolutions of the identity (S22) into Eq. (S30) and using the idempotence property of the projectors P^n , i.e., $P^n P^m = \delta_{nm} P^n$ (S21), one obtains the spectral representation of the SQGT,

$$\begin{aligned} \eta_{\mu\nu}^n &= s_n (\partial_\mu \mathbf{w}^n)^\dagger \tau_z Q^n (\partial_\nu \mathbf{w}^n) = \sum_{m,m'} s_n (\partial_\mu \mathbf{w}^n)^\dagger \tau_z P^m Q^n P^{m'} (\partial_\nu \mathbf{w}^n) \\ &= \sum_{m,m'} s_n (\partial_\mu \mathbf{w}^n)^\dagger \tau_z P^m (\mathbb{1}_{2N} - P^n) P^{m'} (\partial_\nu \mathbf{w}^n) = \sum_{m,m'} s_n (\partial_\mu \mathbf{w}^n)^\dagger \tau_z (\delta_{mm'} P^m - \delta_{nm'} \delta_{nm} P^m) (\partial_\nu \mathbf{w}^n) \\ &= s_n \sum_m [(\partial_\mu \mathbf{w}^n)^\dagger \tau_z (P^m) (\partial_\nu \mathbf{w}^n)] - s_n (\partial_\mu \mathbf{w}^n)^\dagger \tau_z (P^n) (\partial_\nu \mathbf{w}^n) \\ &= s_n \sum_{\substack{m \neq n \\ m=1}}^{2N} [(\partial_\mu \mathbf{w}^n)^\dagger \tau_z (P^m) (\partial_\nu \mathbf{w}^n)] = s_n \sum_{\substack{m \neq n \\ m=1}}^{2N} s_m \frac{(\mathbf{w}^n)^\dagger (\partial_\mu \mathcal{M}) \mathbf{w}^m \times (\mathbf{w}^m)^\dagger (\partial_\nu \mathcal{M}) \mathbf{w}^n}{(\tilde{\Omega}_n - \tilde{\Omega}_m)^2}, \end{aligned} \quad (\text{S33})$$

where in the last step, the symplectic Hellmann-Feynman theorem (S32) has been used. Eq. (S33) is the spectral representation of the SQGT.

S3A. Symplectic quantum metric

We show that the symplectic quantum metric naturally arises when considering the distance ds^2 between two infinitesimally close Bogoliubov modes. In analogy to Refs. [56, 57], we define the distance ds^2 between nearby states $\mathbf{w}^n(\boldsymbol{\lambda})$ and $\mathbf{w}^n(\boldsymbol{\lambda} + d\boldsymbol{\lambda})$ (omitting the index n in the following) via

$$ds^2 = 1 - f^2 = 1 - |(\mathbf{w}(\boldsymbol{\lambda}))^\dagger \tau_z \mathbf{w}(\boldsymbol{\lambda} + d\boldsymbol{\lambda})|^2, \quad (\text{S34})$$

with the fidelity $f = |(\mathbf{w}(\boldsymbol{\lambda}))^\dagger \tau_z \mathbf{w}(\boldsymbol{\lambda} + d\boldsymbol{\lambda})|$. We first expand $\mathbf{w}(\boldsymbol{\lambda} + d\boldsymbol{\lambda})$ to second order in $d\boldsymbol{\lambda}$,

$$\mathbf{w}(\boldsymbol{\lambda} + d\boldsymbol{\lambda}) = \mathbf{w}(\boldsymbol{\lambda}) + \partial_\mu \mathbf{w}(\boldsymbol{\lambda}) d\lambda^\mu + \frac{1}{2} \partial_\mu \partial_\nu \mathbf{w}(\boldsymbol{\lambda}) d\lambda^\mu d\lambda^\nu, \quad (\text{S35})$$

where we use the Einstein summation convention over Greek indices in the following. We then insert this expression (S35) into (S34), using orthonormalization conditions (S23). Collecting only terms up to second order in $d\boldsymbol{\lambda}$ then yields

$$\begin{aligned} ds^2 &= 1 - \left\{ (\mathbf{w}^\dagger \tau_z \mathbf{w})^2 + (\mathbf{w}^\dagger \tau_z \mathbf{w}) [((\partial_\mu \mathbf{w})^\dagger \tau_z \mathbf{w}) + (\mathbf{w}^\dagger \tau_z (\partial_\mu \mathbf{w}))] d\lambda^\mu + [((\partial_\mu \mathbf{w})^\dagger \tau_z \mathbf{w}) (\mathbf{w}^\dagger \tau_z (\partial_\nu \mathbf{w}))] d\lambda^\mu d\lambda^\nu \right. \\ &\quad \left. + \frac{1}{2} (\mathbf{w}^\dagger \tau_z \mathbf{w}) [((\partial_\nu \partial_\mu \mathbf{w})^\dagger \tau_z \mathbf{w}) + (\mathbf{w}^\dagger \tau_z (\partial_\nu \partial_\mu \mathbf{w}))] d\lambda^\mu d\lambda^\nu \right\} + \mathcal{O}(|d\boldsymbol{\lambda}|^3) \\ &= \{ (\mathbf{w}^\dagger \tau_z \mathbf{w}) [(\partial_\mu \mathbf{w})^\dagger \tau_z (\partial_\nu \mathbf{w})] - (\partial_\mu \mathbf{w})^\dagger \tau_z \mathbf{w} \mathbf{w}^\dagger \tau_z (\partial_\nu \mathbf{w}) \} d\lambda^\mu d\lambda^\nu + \mathcal{O}(|d\boldsymbol{\lambda}|^3) \\ &= [(\mathbf{w}^\dagger \tau_z \mathbf{w}) (\partial_\mu \mathbf{w})^\dagger \tau_z Q (\partial_\nu \mathbf{w})] d\lambda^\mu d\lambda^\nu + \mathcal{O}(|d\boldsymbol{\lambda}|^3) \end{aligned} \quad (\text{S36})$$

where the explicit parameter dependence of the states has been omitted for clarity. In the second equality above, we have on the one hand used $(\mathbf{w}^\dagger \tau_z \mathbf{w})^2 = 1$ (S23) and on the other hand that the linear terms ($\sim d\lambda^\mu$) vanish, which follows from the relation $(\mathbf{w}^\dagger \tau_z (\partial_\mu \mathbf{w})) = -(\partial_\mu \mathbf{w})^\dagger \tau_z (\mathbf{w})$ (S23). From this relation, it moreover follows that $((\partial_\nu \partial_\mu \mathbf{w})^\dagger \tau_z \mathbf{w}) + (\mathbf{w}^\dagger \tau_z (\partial_\nu \partial_\mu \mathbf{w})) = -[(\partial_\mu \mathbf{w})^\dagger \tau_z (\partial_\nu \mathbf{w}) + (\partial_\nu \mathbf{w})^\dagger \tau_z (\partial_\mu \mathbf{w})]$ which has also been used in the second equality above (S36). Finally, comparing this expression to the definition of the symplectic quantum geometric tensor in Eq. (S30), we see that the symplectic quantum geometric tensor (SQGT) naturally emerges as a distance measure in the space of the Bogoliubov modes

$$ds^2 = \eta_{\mu\nu} d\lambda^\mu d\lambda^\nu + \mathcal{O}(|d\boldsymbol{\lambda}|^3) = \frac{1}{2} (\eta_{\mu\nu} + \eta_{\nu\mu}) d\lambda^\mu d\lambda^\nu + \mathcal{O}(|d\boldsymbol{\lambda}|^3) = g_{\mu\nu} d\lambda^\mu d\lambda^\nu + \mathcal{O}(|d\boldsymbol{\lambda}|^3). \quad (\text{S37})$$

More precisely, since the distance measure ds^2 (S36) is a symmetric bilinear form, the antisymmetric part of the SQGT does not contribute to ds^2 , i.e., only the symmetric part, i.e., the symplectic quantum metric $g_{\mu\nu} = \frac{1}{2} (\eta_{\mu\nu} + \eta_{\nu\mu}) = \text{Re}[\eta_{\mu\nu}]$, determines the distance measure between two infinitesimally close Bogoliubov states, hence, the second equality in Eq. (S37).

S3B. Local conservation law for symplectic Berry curvature

We show that the symplectic Berry curvature, admits a local conservation law over all subspaces (Bogoliubov subspaces), including both the particle and hole subspaces [7]. For this, note that using Eq. (S30), the symplectic Berry curvature of the subspace n , $B_{\mu\nu}^n = -2\text{Im}[\eta_{\mu\nu}^n]$ can be written as

$$B_{\mu\nu}^n = i s_n ((\partial_\mu \mathbf{w}^n)^\dagger \tau_z (\partial_\nu \mathbf{w}^n) - (\partial_\nu \mathbf{w}^n)^\dagger \tau_z (\partial_\mu \mathbf{w}^n)), \quad (\text{S38})$$

where we omit the explicit parameter dependence for clarity. Next, inserting the resolution of the identity (S22) into (S38), we see that we may also write $B_{\mu\nu}^n$ as

$$\begin{aligned} B_{\mu\nu}^n &= i s_n \left\{ \sum_m s_m (\partial_\mu \mathbf{w}^n)^\dagger \tau_z \mathbf{w}^m (\mathbf{w}^m)^\dagger \tau_z (\partial_\nu \mathbf{w}^n) - (\partial_\nu \mathbf{w}^n)^\dagger \tau_z (\partial_\mu \mathbf{w}^n) \right\} \\ &= i s_n \left\{ \sum_m [s_m (\partial_\nu \mathbf{w}^m)^\dagger \tau_z \mathbf{w}^n (\mathbf{w}^n)^\dagger \tau_z (\partial_\mu \mathbf{w}^m)] - (\partial_\nu \mathbf{w}^n)^\dagger \tau_z (\partial_\mu \mathbf{w}^n) \right\}, \end{aligned} \quad (\text{S39})$$

where in the second equality we used $(\mathbf{w}^m)^\dagger \tau_z (\partial_\nu \mathbf{w}^n) = -(\partial_\nu \mathbf{w}^m)^\dagger \tau_z (\mathbf{w}^n)$ twice, which itself simply follows from Eq. (S23). Now, by summing Eq. (S39) over all subspaces n ,

$$\begin{aligned} \sum_n B_{\mu\nu}^n &= i \left\{ \sum_m \left[s_m (\partial_\nu \mathbf{w}^m)^\dagger \tau_z \left(\sum_j s_n \mathbf{w}^n (\mathbf{w}^n)^\dagger \tau_z \right) (\partial_\mu \mathbf{w}^m) \right] - \sum_j s_n (\partial_\nu \mathbf{w}^n)^\dagger \tau_z (\partial_\mu \mathbf{w}^n) \right\}, \\ &= i \left\{ \sum_m [s_m (\partial_\nu \mathbf{w}^m)^\dagger \tau_z (\partial_\mu \mathbf{w}^m)] - \sum_j [s_n (\partial_\nu \mathbf{w}^n)^\dagger \tau_z (\partial_\mu \mathbf{w}^n)] \right\} = 0, \end{aligned} \quad (\text{S40})$$

we immediately see that the symplectic Berry curvature admits a local conservation law for all parameter values $\boldsymbol{\lambda}$,

$$\sum_n B_{\mu\nu}^n(\boldsymbol{\lambda}) = 0, \quad \forall \boldsymbol{\lambda}. \quad (\text{S41})$$

S4. TIME-DEPENDENT SCHRÖDINGER EQUATION FOR BOGOLIUBOV MODES

Starting from the Heisenberg equations of motions (EOMs) for the original bosonic operators, we derive an effective time-dependent single-particle Schrödinger equation for the Bogoliubov modes \boldsymbol{w}^n . Using Eq. (S1), we see that the Heisenberg EOM ($\hbar \equiv 1$ throughout) for the original bosonic operators read

$$i\partial_t \hat{a}_m = [\hat{a}_m, \hat{H}] = \sum_j K_{mj} \hat{a}_j + \frac{1}{2} [\hat{a}_j^\dagger G_{jm} + \hat{a}_j^\dagger G_{jm}], \quad (\text{S42})$$

$$i\partial_t \hat{a}_m^\dagger = [\hat{a}_m^\dagger, \hat{H}] = -\sum_j \hat{a}_j^\dagger K_{jm} - \frac{1}{2} [G_{mj}^* \hat{a}_j + G_{mj}^* \hat{a}_j]. \quad (\text{S43})$$

These EOMs can also be compactly written in terms of the bosonic Nambu spinor $\hat{\Psi} = (\hat{a}_1, \dots, \hat{a}_N, \hat{a}_1^\dagger, \dots, \hat{a}_N^\dagger)^T$ [12, 65]

$$i\partial_t \hat{\Psi} = [\hat{\Psi}, \hat{H}] = D\hat{\Psi}, \quad \text{with } D = \tau_z \mathcal{M}, \quad (\text{S44})$$

where D is again the dynamical matrix (S29) and \mathcal{M} is the coefficient matrix of the BdG Hamiltonian consisting of matrix blocks K and G (S1). Eq. (S44) is easily solved by

$$\hat{\Psi}(t) = e^{-iDt} \hat{\Psi}(0) \quad (\text{S45})$$

which itself leads to the following time-dependence for the Nambu spinor of the new bosonic operators $\hat{\Phi}(t) = W^{-1} \hat{\Psi}(t)$ (S6)

$$\hat{\Phi}(t) = e^{-i\tilde{\Omega}t} \hat{\Phi}(0) = e^{-i\tilde{\Omega}t} W^{-1} \hat{\Psi}(0), \quad (\text{S46})$$

where we have utilized that $W^{-1}DW = \tau_z W^\dagger \mathcal{M} W = \tau_z \tilde{\Omega} \equiv \tilde{\Omega}$ which follows from Eq. (S28). Thus, each new bosonic operator acquires a phase factor associated to eigenenergy $\hat{b}_n(t) = e^{-i\omega_n t} \hat{b}_n(0)$ [12, 65].

Next, we note that we may also absorb the time-dependence of the original operators $\hat{\Psi}(t)$ into the (inverse) Bogoliubov matrix, $\hat{\Phi}(t) = W^{-1} \hat{\Psi}(t) \equiv W^{-1}(t) \hat{\Psi}(0)$, i.e., in (S46) $W^{-1}(t) = e^{-i\tilde{\Omega}t} W^{-1}(0) = W^{-1}(0) e^{-iDt}$. This leads to the immediate question, what kind of equation of motion $W^{-1}(t)$ fulfills. In compact matrix form, the answer is given by the following set of equations for the Bogoliubov modes,

$$\partial_t W^{-1}(t) = -iW^{-1}(t)D \Leftrightarrow \partial_t W(t) = iDW(t). \quad (\text{S47})$$

Since we know that $W^{-1}D = \tilde{\Omega}W^{-1}$, we see that (S47) is solved by $W^{-1}(t) = e^{-i\tilde{\Omega}t} W^{-1}(0)$, which is in agreement with (S46). This may also be expressed in as an effective Schrödinger equation for the individual Bogoliubov modes $\boldsymbol{w}^n(t)$ [65, 97]

$$-i\partial_t \boldsymbol{w}^n(t) = D\boldsymbol{w}^n(t), \quad (\text{S48})$$

which is the sought after effective time-dependent Schrödinger equation. From Eq. (S48), for a time-independent dynamical matrix D , we can define the paraunitary time evolution operator $P(t)$,

$$P(t) = \exp(iDt), \quad (\text{S49})$$

such that $\boldsymbol{w}^n(t) = P(t)\boldsymbol{w}^n(0)$. For a time-dependent dynamical matrix $D(t)$ in (S48), the paraunitary time evolution operator reads

$$P(t) = \mathcal{T} \exp\left(i \int_0^t dt' D(t')\right), \quad (\text{S50})$$

where \mathcal{T} denotes the time-ordering operator. Note that the paraunitarity of $P(t)$, i.e., $\tau_z P^\dagger(t) \tau_z = P^{-1}(t)$, directly follows from the pseudo-hermiticity of the dynamical matrix, $D^\dagger(t) = \tau_z D(t) \tau_z$. Thus, the time evolution of the Bogoliubov modes \mathbf{w}^n (S48) directly specifies the time-dependence of the Heisenberg-picture quasiparticle operators $\hat{b}_n^\dagger(t) = \hat{\Psi}^\dagger(t) \tau_z \mathbf{w}^n$ (S16),

$$\hat{b}_n^\dagger(t) = \hat{\Psi}^\dagger(t) \tau_z \mathbf{w}^n = \hat{\Psi}^\dagger(0) e^{i(D^\dagger)t} \tau_z \mathbf{w}^n = \hat{\Psi}^\dagger(0) \tau_z e^{iDt} \mathbf{w}^n = \hat{\Psi}^\dagger(0) \tau_z \mathbf{w}^n(t), \quad (\text{S51})$$

where we have used (S45) and the pseudo-hermiticity of the dynamical matrix $D^\dagger = \tau_z D \tau_z$.

S5. TIME-DEPENDENT PERTURBATION THEORY FOR THE EFFECTIVE SCHRÖDINGER EQUATION

Consider the effective time-dependent single-particle Schrödinger equation (S48) for some state $\psi(t) \in \mathbb{C}^{2N}$ in the indefinite inner product space spanned by the Bogoliubov modes $\mathbf{w}^n(t)$ with the addition of a small time-dependent perturbation $V(t)$ added to the dynamical matrix, i.e.,

$$-i\partial_t \psi(t) = D(t) \psi(t) = [D + V(t)] \psi(t) \quad (\text{S52})$$

where $V(t)$ shall only be present for $t > 0$. Adapting standard time-dependent perturbation theory [127] to the space of Bogoliubov modes, we now derive the effect of the perturbation $V(t)$ on the population of Bogoliubov modes.

At $t = 0$, where $V(0) = 0$, we consider the initial state $\psi(0)$ to be given by the Bogoliubov eigenstate \mathbf{w}^n , i.e., $D\psi(0) = \tilde{\Omega}_n \mathbf{w}^n$ (S29). At this point, \mathbf{w}^n adheres to the effective time-dependent Schrödinger equation without the perturbation (S48), which is easily solved by $\mathbf{w}^n(t) = e^{iDt} \mathbf{w}^n(0) = e^{i\tilde{\Omega}_n t} \mathbf{w}^n(0)$ (S48). To first order in the perturbation $V(t)$, we now compute the resulting excitation probability to other distinct Bogoliubov modes $\mathbf{w}^m(t)$ at time t .

First, we introduce the interaction picture

$$\psi_I(t) = \tilde{P}(t) \psi(t) = e^{-iDt} \psi(t). \quad (\text{S53})$$

The transformation is explicitly given by $\tilde{P}(t) = e^{-iDt}$ where D is the static part in Eq. (S52). Due to the pseudo-hermiticity of the dynamical matrix, $D^\dagger = \tau_z D \tau_z$, in the equation of motion (S48), the transformation $\tilde{P}(t)$ must be paraunitary, i.e., $\tilde{P}^\dagger(t) = \tau_z \tilde{P}^{-1}(t) \tau_z$. This directly follows from $\tau_z \tilde{P}^\dagger(t) \tau_z = \tau_z e^{iD^\dagger t} \tau_z = e^{iDt} = \tilde{P}^{-1}(t)$. The effective time-dependent Schrödinger equation in the interaction picture then reads

$$-i\partial_t \psi_I(t) = D_I(t) \psi_I(t), \quad (\text{S54})$$

where we used Eq. (S48) and introduced $D_I(t)$, the dynamical matrix in the interaction picture given by

$$D_I(t) = \tilde{P}(t) D(t) \tilde{P}^{-1}(t) - i(\partial_t \tilde{P}(t)) \tilde{P}^{-1}(t). \quad (\text{S55})$$

With $\tilde{P}(t) = e^{-iDt}$, Eq. (S54) then becomes

$$-i\partial_t \psi_I(t) = V_I(t) \psi_I(t), \quad (\text{S56})$$

where $V_I(t) \equiv \tilde{P}(t) V(t) \tilde{P}^{-1}(t) = e^{-iDt} V(t) e^{iDt}$ is the perturbation in the interaction picture. The solution to (S56) is then given by a Dyson-type series,

$$\psi_I(t) = \psi_I(0) + i \int_0^t dt' V_I(t') \psi_I(0) + i^2 \int_0^t dt' \int_0^{t'} dt'' V_I(t') V_I(t'') \psi_I(0) + \dots, \quad (\text{S57})$$

where we are only interested in the first order contribution. Now, given an initial eigenstate $\psi_I(0) = \mathbf{w}^n(0)$, we compute the transition amplitude $c_n^\psi(t)$ to some other arbitrary mode $\mathbf{w}^m(t)$ to first order at time t by first transforming back to the original picture (S53)

$$\psi(t) = e^{iDt} \psi_I(t) = e^{i\tilde{\Omega}_n t} \mathbf{w}^n(0) + e^{iDt} i \int_0^t dt' V_I(t') \mathbf{w}^n \quad (\text{S58})$$

and then projecting onto $s_m(\mathbf{w}^m)^\dagger \tau_z$ [cf. Eq. (S25)],

$$c_m^\psi(t) = s_m(\mathbf{w}^m)^\dagger \tau_z \psi(t) = s_m s_n \delta_{nm} e^{i(\tilde{\Omega}_n - \tilde{\Omega}_m)t} + i s_m \int_0^t dt' (\mathbf{w}^m)^\dagger \tau_z V(t') \mathbf{w}^n e^{i(\tilde{\Omega}_n - \tilde{\Omega}_m)t'}, \quad (\text{S59})$$

where we have utilized that the dynamical matrix is pseudo-hermitian, $D^\dagger = \tau_z D \tau_z$, which implies that $\exp(iDt)^\dagger = \tau_z \exp(-iDt) \tau_z$. Eventually, the probability for the system to transition into some distinct orthogonal state $m \neq n$ is given by

$$p_{mn} \equiv |c_m^\psi(t)|^2 = \left| \int_0^t dt' ((\mathbf{w}^m)^\dagger \tau_z V(t') \mathbf{w}^n) e^{i(\tilde{\Omega}_n - \tilde{\Omega}_m)t'} \right|^2. \quad (\text{S60})$$

Next, we consider a periodic perturbation of the form $V(t) = F \cos(\omega t - \varphi) = \frac{1}{2} F (e^{i(\omega t - \varphi)} + e^{-i(\omega t - \varphi)})$, where F is some constant real-valued amplitude. Thus, using standard assumptions of $t \gg 1/\omega$ and $\omega \simeq (\tilde{\Omega}_m - \tilde{\Omega}_n)$ [127], Eq. (S60) becomes

$$p_{mn}(\omega, t) = \frac{2\pi}{4} t |((\mathbf{w}^m)^\dagger \tau_z F e^{i\varphi} \mathbf{w}^n)|^2 \delta_t(\tilde{\Omega}_m - \tilde{\Omega}_n - \omega), \quad (\text{S61})$$

where $\delta_t(\alpha) = \frac{\sin^2(\alpha t/2)}{\pi \alpha^2 t}$ approaches a Dirac-Delta distribution in the limit of large observation times t , which then yields Eq. (13) in the main text.

S6. MEASURING THE SYMPLECTIC QUANTUM GEOMETRY IN LATTICE SYSTEMS

In this appendix, we specifically apply the general formalism above to the case of two-dimensional lattice BBdG systems, described by a translationally invariant dynamical matrix D , which is inspired by Refs. [59, 60]. For such a system, the Bogoliubov eigenmodes of the dynamical matrix D have Bloch function form, $\psi^n(\mathbf{q}) = e^{i\mathbf{q}\cdot\mathbf{r}} \mathbf{w}^n(\mathbf{q})$ [20, 97], such that the Bogoliubov eigenvalue equation in the quasimomentum space representation (S29) is given by $D(\mathbf{q}) \mathbf{w}^n(\mathbf{q}) = \tilde{\Omega}_n(\mathbf{q}) \mathbf{w}^n(\mathbf{q})$. Here, $\mathbf{w}^n(\mathbf{q})$ is the cell-periodic part of the Bogoliubov Bloch function for the n -th band. Next, in direct analogy to Refs. [59, 60], we consider periodically shaking the underlying lattice along x -direction of the lattice, resulting in the following time-dependent dynamical matrix,

$$D_x(t) = D + 2A \cos(\omega t) \mathcal{X} = D + V(t), \quad V(t) = 2A \cos(\omega t) \mathcal{X}, \quad (\text{S62})$$

where D is the initial dynamical matrix in position space and $\mathcal{X} = \sigma_z \otimes x$ the (pseudo-hermitian) position operator for the direction x (extended to Nambu space), and with x being the usual position operator.

We note that such periodically shaken systems described via $D_x(t)$ are in direct analogy to the general discussion following Eq. (12) in the main text. To see this, we first note that $D_x(t)$ becomes translationally invariant in the co-moving frame of the lattice [59, 60], when introducing a paraunitary rotation

$$\tilde{P}(t) = \exp\left(-i \int dt' V(t')\right) = \exp(-i(2A/\omega) \sin(\omega t) \mathcal{X}), \quad (\text{S63})$$

such that the dynamical matrix in the co-moving frame (S55) becomes

$$\begin{aligned} D'_x(t) &= \tilde{P} D_x(t) \tilde{P}^{-1} - i(\partial_t \tilde{P}) \tilde{P}^{-1} = \tilde{P} [D + V(t)] \tilde{P}^{-1} - i(-iV(t) \tilde{P}) \tilde{P}^{-1} \\ &= \tilde{P} D \tilde{P}^{-1} = D(\mathbf{q}) + \frac{2A}{\omega} \sin(\omega t) \partial_{q_x} D(\mathbf{q}) + \mathcal{O}((A/\omega)^2), \end{aligned} \quad (\text{S64})$$

which agrees with the formalism of Eq. (12) in the main text.

As in the general discussion in the main text, given an initially populated Bogoliubov mode $\psi^n = e^{i\mathbf{q}_0 \cdot \mathbf{r}} \mathbf{w}^n(\mathbf{q}_0)$, we compute the total excitation rate into all other available Bogoliubov modes ψ^m ,

$$\Gamma_x^n(\omega) = \frac{1}{t} \sum_{m \neq n} s_m p_{mn}(\omega, t), \quad (\text{S65})$$

with $p_{mn}(\omega, t)$ being given by (S61). We note that the factor s_m comes from the indefinite inner product structure of the Bogoliubov modes [cf. Eq. (S26)]. The total excitation rate can then be expressed as

$$\begin{aligned}\Gamma_x^n(\omega) &= 2\pi A^2 \sum_{m \neq n} s_m |(\mathbf{w}^m(\mathbf{q}))^\dagger e^{-i\mathbf{q}\cdot\mathbf{r}} \tau_z \mathcal{X} e^{i\mathbf{q}_0\cdot\mathbf{r}} \mathbf{w}^n(\mathbf{q}_0)|^2 \times \delta(\tilde{\Omega}_m(\mathbf{q}) - \tilde{\Omega}_n(\mathbf{q}_0) - \omega) \\ &= 2\pi A^2 \sum_{m \neq n} s_m |(\mathbf{w}^m(\mathbf{q}_0))^\dagger \tau_z i\partial_{q_x} \mathbf{w}^n(\mathbf{q}_0)|^2 \times \delta(\tilde{\Omega}_m(\mathbf{q}_0) - \tilde{\Omega}_n(\mathbf{q}_0) - \omega).\end{aligned}\tag{S66}$$

Here, we have used that the position operator acts as a derivative in the quasimomentum space representation, i.e., $(\mathbf{w}^m(\mathbf{q}))^\dagger e^{-i\mathbf{q}\cdot\mathbf{r}} \tau_z \mathcal{X} e^{i\mathbf{q}_0\cdot\mathbf{r}} \mathbf{w}^n(\mathbf{q}_0) = i\delta_{\mathbf{q},\mathbf{q}_0} (\mathbf{w}^m(\mathbf{q}))^\dagger \tau_z \partial_{q_x} \mathbf{w}^n(\mathbf{q})$ [60, 128]. By integrating the excitation rate (S66) over ω , we arrive at

$$\begin{aligned}\Gamma_{x,\text{int}}^n &= \int d\omega \Gamma_x^n(\omega) = 2\pi A^2 \sum_{m \neq n} s_m |(\mathbf{w}^m(\mathbf{q}_0))^\dagger \tau_z \partial_{q_x} \mathbf{w}^n(\mathbf{q}_0)|^2 \\ &= \sum_{m \neq n} [(\partial_{q_x} \mathbf{w}^n(\mathbf{q}_0))^\dagger \tau_z (P^m) (\partial_{q_x} \mathbf{w}^n(\mathbf{q}_0))] = 2\pi A^2 g_{xx}^n(\mathbf{q}_0),\end{aligned}\tag{S67}$$

where in the last step we have used (S33) to identify the symplectic quantum metric. Thus, the integrated excitation rate after shaking the lattice along the x direction is directly proportional to the diagonal component $g_{xx}^n(\mathbf{q}_0)$ of the symplectic quantum metric tensor. By similarly shaking the lattice along the y direction, the diagonal component $g_{yy}^n(\mathbf{q}_0)$ is obtained.

To obtain the remaining off-diagonal components of the symplectic quantum geometric tensor, i.e., the symplectic quantum metric $g_{xy}^n(\mathbf{q}_0)t$ and the symplectic Berry curvature $B_{xy}^n(\mathbf{q}_0)$, we now imagine shaking the lattice along both the x and y directions [59, 60], which is described by

$$D_{x\pm y}(t) = D + 2A(\cos(\omega t)\mathcal{X} \pm \cos(\omega t - \varphi)\mathcal{Y}) = D + V(t), \quad V(t) = 2A(\cos(\omega t)\mathcal{X} \pm \cos(\omega t - \varphi)\mathcal{Y}),\tag{S68}$$

where here $\mathcal{Y} = \sigma_z \otimes y$ is the (pseudo-hermitian) y -position operator (extended to Nambu space) with y being the usual position operator.

For completeness, we also provide the paraunitarity rotation to the co-moving frame for this case (S68)

$$P(t) = \exp\left(-i \int dt' V(t')\right) = \exp\{-i(A/\omega)(\sin(\omega t)\mathcal{X} \pm \sin(\omega t - \varphi)\mathcal{Y})\},\tag{S69}$$

which then leads to the following dynamical matrix in the co-moving frame (S55)

$$D'_{x\pm y}(t) = D(\mathbf{q}) + \frac{2A}{\omega} (\sin(\omega t)\partial_{q_x} D(\mathbf{q}) \pm \sin(\omega t - \varphi)\partial_{q_y} D(\mathbf{q})) + \mathcal{O}((A/\omega)^2).\tag{S70}$$

This allows us again to see the connection to the general discussion following Eq. (12) in the main text. We emphasize that Eq. (S70) is in direct analogy to the results for the conventional case in Ref. [59].

Then, in the same fashion as before (S65)-(S67), computing the integrated excitation rate leads to the emergence of the off-diagonal components of the SQGT depending on the phase offset φ ,

$$\Gamma_{\text{int}}^{n,\pm} = 2\pi A^2 \times \begin{cases} (g_{xx}^n(\mathbf{q}_0) \pm 2g_{xy}^n(\mathbf{q}_0) + g_{yy}^n(\mathbf{q}_0)), & \text{for } \varphi = 0, \\ (g_{xx}^n(\mathbf{q}_0) \mp B_{xy}^n(\mathbf{q}_0) + g_{yy}^n(\mathbf{q}_0)), & \text{for } \varphi = \pi/2. \end{cases}\tag{S71}$$

By taking the differential integrated rate [59] $\Delta\Gamma_{\text{int}}^n = \Gamma_{\text{int}}^{n,+} - \Gamma_{\text{int}}^{n,-}$ of the two protocols with the opposite sign in $D_{x\pm y}(t)$ (S68), we obtain

$$\Delta\Gamma_{\text{int}}^n = \begin{cases} 8\pi A^2 g_{xy}^n(\mathbf{q}_0), & \text{for } \varphi = 0, \\ -4\pi A^2 B_{xy}^n(\mathbf{q}_0), & \text{for } \varphi = \pi/2. \end{cases}\tag{S72}$$

Depending on the phase offset φ of the periodic modulation in (S68), we see that the differential integrated rate is directly proportional to the off-diagonal symplectic quantum metric $g_{xy}^n(\mathbf{q}_0)$ or the symplectic Berry curvature $B_{xy}^n(\mathbf{q}_0)$.

All the prior formalism, which assumed a single populated Bogoliubov mode at some quasimomentum \mathbf{q}_0 , may also be generalized to an initial Bogoliubov Bloch wave packet, e.g., a weighted average of multiple Bogoliubov (particle)

modes within the same band n , i.e., $\phi^n = \sum_{\mathbf{q}} c(\mathbf{q}) e^{i\mathbf{q}\cdot\mathbf{r}} w^n(\mathbf{q})$, where the wave packet ϕ^n is normalized with respect to the Bogoliubov norm $(\phi^n)^\dagger \tau_z \mathbf{r} \phi^n = \sum_{\mathbf{q}} |c(\mathbf{q})|^2 = 1$ [97]. One may further assume that the wave packet is localized around some central quasimomentum mode $\mathbf{q}_0 = \sum_{\mathbf{q}} |c(\mathbf{q})|^2 \mathbf{q}$, as well as around some central position \mathbf{r}_c , which is given by $\mathbf{r}_c = (\phi^n)^\dagger \tau_z \mathbf{r} \phi^n$ [97]. This then simply leads to the integrated rates being proportional to the weighted average of the symplectic quantum geometric quantities, e.g., for the linear shaking protocol of Eq. (S62), the integrated rate becomes

$$\Gamma_{x,\text{int}}^n = 2\pi A^2 \sum_{\mathbf{q}} |c(\mathbf{q})|^2 g_{xx}^n(\mathbf{q}), \quad (\text{S73})$$

which is in direct analogy to the proposals for probing the quantum geometry in particle-number-conserving systems [60].

S7. TIME-DEPENDENT PERTURBATION THEORY FOR MANY-BODY BOGOLIUBOV FOCK STATES

In the main text we have shown that the symplectic quantum geometric tensor can be extracted from integrated excitation rates to other Bogoliubov modes in response to weak periodic modulations of bosonic BdG systems Eq. (12). For this, we have used the results of first-order perturbation theory for the effective Schrödinger equation (S61). In this appendix, we now show that this derivation is equivalent to applying standard first-order time-dependent perturbation theory (Fermi's Golden rule) to treat transitions between many-body Bogoliubov Fock states with quasiparticle excitations present.

Let us first consider the general case, where, in addition to the already solved (unperturbed) problem of (S1), we apply a generic time-dependent particle-number-conserving perturbation of the form $V(t) = \sum_{i,j} \hat{a}_i^\dagger A_{ij}(t) \hat{a}_j$ to the system (S1). To understand how this perturbation affects the population of Bogoliubov quasiparticle excitations, we express $V(t)$ via the diagonal (quasiparticle) basis of the unperturbed problem using (S5)

$$\begin{aligned} V(t) &= \sum_{i,j} \hat{a}_i^\dagger A_{ij}(t) \hat{a}_j = \sum_{i,j} A_{ij}(t) \sum_{m,m'} \left(\hat{b}_m^\dagger (U^\dagger)_{mi} + \hat{b}_m (V^T)_{mi} \right) A_{ij}(t) \left(U_{jm'} \hat{b}_{m'} + (V^*)_{jm'} \hat{b}_{m'}^\dagger \right) \\ &= \sum_{m,m'} [\hat{b}_m^\dagger \hat{b}_{m'} \sum_{ij} (U^\dagger)_{mi} A_{ij}(t) U_{jm'} + \hat{b}_m \hat{b}_{m'}^\dagger \sum_{ij} (V^T)_{mi} A_{ij}(t) (V^*)_{jm'} \\ &\quad + \hat{b}_m^\dagger \hat{b}_{m'}^\dagger \sum_{ij} (U^\dagger)_{mi} A_{ij}(t) (V^*)_{jm'} + \hat{b}_m \hat{b}_{m'} \sum_{ij} (V^T)_{mi} A_{ij}(t) U_{jm'}]. \end{aligned} \quad (\text{S74})$$

Note that the ground state of the original BdG Hamiltonian (S19) contains zero Bogoliubov quasiparticle excitations. In the main text, we have argued that such a generic perturbation (S74), can be used to controllably create pairs of quasiparticle excitations in desired Bogoliubov energy bands via terms $\propto \hat{b}_m^\dagger \hat{b}_{m'}^\dagger$ or $\propto \hat{b}_m^\dagger$ [cf. Fig. 2(a) in the main text]. The presence of such Bogoliubov quasiparticle excitations is the starting point for the protocol to extract the symplectic quantum geometric tensor from excitation rates between Bogoliubov modes, as will show in the following [cf. Eq. (13) in the main text].

S7A. Transitions between Bogoliubov quasiparticle excitations lead to SQGT

Now, as a first simple case, we consider the transition of single Bogoliubov quasiparticle excitation from some occupied mode β to some unoccupied mode α via the perturbation $V(t)$ (S74). As an initial state $|\phi_\beta\rangle$, we then choose the Bogoliubov Fock state with a single quasiparticle excitation in some mode β and, i.e., $|\phi_\beta\rangle = |\dots, (0)_\alpha, (1)_\beta, \dots\rangle$ and as a final state the state with an excitation in some mode α , i.e., $|\psi_\alpha\rangle = |\dots, (1)_\alpha, (0)_\beta, \dots\rangle$. Standard first-order time-dependent perturbation theory [127], tells us that we have to look at matrix elements $\langle \psi_\alpha | V(t) | \phi_\beta \rangle$ to analyze the transitions between the initial and final states. So the only matrix elements of the perturbation $\langle \psi_\alpha | V(t) | \phi_\beta \rangle$ that contribute to the transition are only the ones which are number-conserving in the Bogoliubov basis, such that

using (S74) we obtain

$$\begin{aligned}
\langle \psi_\alpha | V(t) | \phi_\beta \rangle &= \langle \psi_\alpha | \sum_{m,m'} \left[\delta_{\alpha,m} \delta_{\beta,m'} \hat{b}_m^\dagger \hat{b}_{m'} \sum_{ij} (U^\dagger)_{mi} A_{ij}(t) U_{jm'} + \delta_{\alpha,m'} \delta_{\beta,m} \hat{b}_m \hat{b}_{m'}^\dagger \sum_{ij} (V^T)_{mi} A_{ij}(t) (V^*)_{jm'} \right] | \phi_\beta \rangle \\
&= \sum_{ij} (U^\dagger)_{\alpha i} A_{ij}(t) U_{j\beta} \langle \psi_\alpha | \hat{b}_\alpha^\dagger \hat{b}_\beta | \phi_\beta \rangle + (V^T)_{\beta i} A_{ij}(t) (V^*)_{j\alpha} \langle \psi_\alpha | \hat{b}_\beta \hat{b}_\alpha^\dagger | \phi_\beta \rangle \\
&= \sum_{ij} (U^\dagger)_{\alpha i} A_{ij}(t) U_{j\beta} + (V^\dagger)_{\alpha j} (A^*)_{ji}(t) V_{i\beta} = \sum_{ij} \begin{pmatrix} U^\dagger & V^\dagger \\ V^T & U^T \end{pmatrix}_{\alpha i} \begin{pmatrix} A(t) & 0 \\ 0 & A^*(t) \end{pmatrix}_{ij} \begin{pmatrix} U & V^* \\ V & U^* \end{pmatrix}_{j\beta} \\
&= \sum_{ij} (W^\dagger)_{\alpha i} \tilde{\mathcal{M}}_{ij}(t) W_{j\beta} = \mathbf{w}_\alpha^\dagger \tilde{\mathcal{M}}(t) \mathbf{w}_\beta, \text{ with } \tilde{\mathcal{M}}(t) = \begin{pmatrix} A(t) & 0 \\ 0 & A^*(t) \end{pmatrix}, \tag{S75}
\end{aligned}$$

where in the second to last step we have introduced the $2N \times 2N$ matrix $\tilde{\mathcal{M}}(t)$ and reformulated the Bogoliubov transformation in terms of the paraunitary matrix W (S6), where both the α and β indices are restricted to the particle sector, i.e., $1 \leq \alpha, \beta \leq N$ for the equality to hold. In the last step of (S75), we have introduced the column vector \mathbf{w}_β given by the β -th column of W and the row vector $\mathbf{w}_\alpha^\dagger$ given by the α -th row of W^\dagger , where again $1 \leq \alpha, \beta \leq N$ (restriction to the particle modes).

We can generalize this result to the case of an initial eigenstate of (S19) with n_β quasiparticle excitations in mode β and n_α excitations in mode α , i.e., $|\phi_\beta\rangle = |\dots, (n_\alpha)_\alpha, (n_\beta)_\beta, \dots\rangle$, and a final state with $n_\beta - 1$ quasiparticle excitations in mode β , and $n_\alpha + 1$ excitation in mode α , i.e., $|\psi_\alpha\rangle = |\dots, (n_\alpha + 1)_\alpha, (n_\beta - 1)_\beta, \dots\rangle$. This will yield a similar result as in (S75) but with an additional bosonic enhancement factor of due to $\hat{b}_\alpha^\dagger \hat{b}_\beta |\phi_\beta\rangle = \sqrt{n_\beta(n_\alpha + 1)} |\phi_\alpha\rangle$,

$$\langle \psi_\alpha | \mathcal{A}(t) | \phi_\beta \rangle = \sqrt{n_\beta(n_\alpha + 1)} \mathbf{w}_\alpha^\dagger \tilde{\mathcal{M}}(t) \mathbf{w}_\beta. \tag{S76}$$

Now we explicitly specify the explicit form of the perturbation. Inspired by Ref. [60], we assume an original parameter-dependent coefficient matrix $\mathcal{M}(\boldsymbol{\lambda}(t))$, where $\boldsymbol{\lambda}(t) = (\lambda^1(t), \dots, \lambda^r(t))$ is a set of time-dependent parameters. We first restrict ourselves only to have parameter dependence in the diagonal (particle-number-conserving) blocks of the coefficient matrix $\mathcal{M}(\boldsymbol{\lambda}(t))$, i.e., $K(\boldsymbol{\lambda}(t))$ and $G = \text{const.}$ (cf. Eq. (S1)). The system shall initially be prepared at some parameter values $\boldsymbol{\lambda} = \boldsymbol{\lambda}_0$. Let us then, for simplicity, consider a single parameter modulation $\lambda^1(t)$ of the form $\lambda^1(t) = \lambda_0^1 + 2(A/\omega) \cos(\omega t)$ with $A/\omega \ll 1$. The rest of the parameters are assumed to be constant, i.e., $\lambda^\nu(t) = \lambda_0^\nu$ for $\nu \neq 1$. We then Taylor-expand the relevant particle-number-conserving perturbation $K(\boldsymbol{\lambda}(t))$ to first order

$$\begin{aligned}
K(\boldsymbol{\lambda}(t)) &= K(\boldsymbol{\lambda}_0) + \partial_1 K(\boldsymbol{\lambda}_0) \times (2A/\omega) \cos(\omega t) = K(\boldsymbol{\lambda}_0) + \tilde{K}(t), \tag{S77} \\
\text{with } \tilde{K}(t) &= f(t) \tilde{K}, \quad f(t) \equiv (2A/\omega) \cos(\omega t), \quad \tilde{K} \equiv \partial_1 K(\boldsymbol{\lambda}_0).
\end{aligned}$$

Here, the matrix $(\tilde{K})_{ij}(t)$ plays the role of the matrix $A_{ij}(t)$ in the general perturbation considered in Eq. (S74). We note that this explicit form of the parameter modulations leading to (S77) is not arbitrary but readily appears in the co-moving frame of periodically shaken systems (S62) [59, 60].

On the level of the Hamiltonian (S1) the perturbation provides an additional term of the form

$$V(t) = \sum_{ij} \hat{a}_i^\dagger \tilde{K}_{ij}(t) \hat{a}_j = \left(\sum_{ij} \hat{a}_i^\dagger \tilde{K}_{ij} \hat{a}_j \right) f(t) \equiv \tilde{V} f(t), \tag{S78}$$

Now, initially, we assume the system to be in some eigenstate $|\phi_\beta\rangle$ of (S19). As before, we let us first discuss the case of a single excitation and generalize to multiple excitations later. Thus, we start with a state with a single Bogoliubov quasiparticle excitation in some mode β , i.e., $|\phi_\beta\rangle = |\dots, (0)_\alpha, (1)_\beta, \dots\rangle$ with the corresponding energy eigenvalue equation $\hat{H}|\phi_\beta\rangle = \omega_\beta |\phi_\beta\rangle$ (S19). Our goal is to compute the transition probability to some final eigenstate $|\psi_\alpha\rangle$, which for now shall consist of a single Bogoliubov quasiparticle excitation in some other mode α , i.e., $|\psi_\alpha\rangle = |\dots, (1)_\alpha, (0)_\beta, \dots\rangle$ with corresponding energy eigenvalue equation $\hat{H}|\psi_\alpha\rangle = \omega_\alpha |\psi_\alpha\rangle$.

Then, given the periodic perturbation of type (S77)-(S78), we can use Fermi's Golden rule (first-order time-dependent perturbation theory) [127] to compute this transition probability at time t with via

$$n_{\alpha\beta}(\omega, t) = 2\pi t \left| \langle \psi_\alpha | \tilde{V} | \phi_\beta \rangle \right|^2 \delta(\omega_\alpha - \omega_\beta - \omega) = 2\pi t \left| \langle \psi_\alpha | \sum_{ij} \hat{a}_i^\dagger \tilde{K}_{ij} \hat{a}_j | \phi_\beta \rangle \right|^2 \delta(\omega_\alpha - \omega_\beta - \omega), \tag{S79}$$

where $\hbar \equiv 1$ in the following. Into Eq. (S79) we can insert the explicit expansion of the perturbation \tilde{V} (S78) in terms of the Bogoliubov quasiparticle operators (S74)-(S75) leading to

$$\begin{aligned} n_{\alpha\beta}(\omega, t) &= 2\pi t \left| \sum_{ij} [(U^\dagger)_{\alpha i} \tilde{K}_{ij} U_{j\beta} \langle \psi_\alpha | \hat{b}_\alpha^\dagger \hat{b}_\beta | \phi_\beta \rangle + (V^T)_{\beta i} \tilde{K}_{ij} (V^*)_{j\alpha} \langle \psi_\alpha | \hat{b}_\beta \hat{b}_\alpha^\dagger | \phi_\beta \rangle] \right|^2 \delta(\omega_\alpha - \omega_\beta - \omega) \\ &= 2\pi t \frac{A^2}{\omega^2} |\mathbf{w}_\alpha^\dagger(\partial_1 \mathcal{M}) \mathbf{w}_\beta|^2 \delta(\omega_\alpha - \omega_\beta - \omega). \end{aligned} \quad (\text{S80})$$

Next, we compute the total excitation rate by summing Eq. (S80) over all possible final states $|\psi_\alpha\rangle$ with a single excitation in some mode α ,

$$\Gamma^\alpha(\omega) = \frac{1}{t} \sum_{\alpha \neq \beta} n_{\alpha\beta}(\omega, t) = \sum_{\alpha \neq \beta} 2\pi \frac{A^2}{\omega^2} |\mathbf{w}_\alpha^\dagger(\partial_1 \mathcal{M}) \mathbf{w}_\beta|^2 \delta(\omega_\alpha - \omega_\beta - \omega) \quad (\text{S81})$$

Then, inspired by Ref. [59], we introduce the integrated excitation rate $\Gamma_{\text{int}}^\alpha$

$$\Gamma_{\text{int}}^\alpha = \int d\omega \Gamma^\alpha(\omega) = 2\pi A^2 \sum_{\alpha \neq \beta, \alpha=1}^N \frac{|\mathbf{w}_\alpha^\dagger(\partial_1 \mathcal{M}) \mathbf{w}_\beta|^2}{(\omega_\alpha - \omega_\beta)^2}. \quad (\text{S82})$$

The last expression already resembles the spectral representation of the symplectic quantum metric (S33), although not exactly, since the sum over α in (S82) is only restricted to the particle Bogoliubov modes $(\mathbf{w}_\alpha^\dagger)_i = (W^\dagger)_{\alpha i}$ with $1 \leq \alpha \leq N$. However, since the spectral representation of the symplectic quantum metric is defined via the sum over all Bogoliubov modes, including the hole modes with $N+1 \leq \alpha \leq 2N$, we need to extend the sum in (S82). This is allowed by the fact that the operator $\partial_1 \mathcal{M}$ (with only particle-number-conserving parts) does not couple the particle and hole Bogoliubov modes. By this, we arrive at

$$\Gamma_{\text{int}} = \int d\omega \sum_\alpha \Gamma^\alpha(\omega) = 2\pi A^2 \sum_{\alpha \neq \beta, \alpha=1}^{2N} s_\alpha \frac{|\mathbf{w}_\alpha^\dagger(\partial_1 \mathcal{M}) \mathbf{w}_\beta|^2}{(\tilde{\omega}_\alpha - \omega_\beta)^2} = 2\pi A^2 g_{11}^\beta. \quad (\text{S83})$$

We observe that the integrated excitation rates between states $|\phi_\beta\rangle$ and $|\psi_\alpha\rangle$, with single Bogoliubov quasiparticle excitations, give rise to the symplectic quantum metric (S33). This agrees with the result derived from first-order perturbation theory for the effective Schrödinger equation (S61) in Eq. (14) in the main text.

As before this result can be generalized, to the case of an initial Bogoliubov Fock state with n_β quasiparticle excitations in mode β and n_α excitations in mode α , i.e., $|\phi_\beta\rangle = |\dots, (n_\alpha)_\alpha, (n_\beta)_\beta, \dots\rangle$, and a final state with $n_\beta - 1$ quasiparticle excitations in mode β , and n_β excitation in mode α , i.e., $|\psi_\alpha\rangle = |\dots, (n_\alpha + 1)_\alpha, (n_\beta - 1)_\beta, \dots\rangle$. This will lead to an additional bosonic enhancement factor of $\sqrt{n_\beta}$ in the transition matrix elements as in (S76). The energy difference between the initial and final state is still given by $[(n_\alpha + 1)\omega_\alpha + \omega_\beta(n_\beta - 1)] - [\omega_\beta n_\beta - \omega_\alpha n_\alpha] = \omega_\alpha - \omega_\beta$ such that we can directly generalize the Fermi's Golden rule result of Eq. (S83) to

$$\Gamma_{\text{int}} = \int d\omega \sum_\alpha \Gamma^\alpha(\omega) = n_\beta(n_\alpha + 1) \times 2\pi A^2 g_{11}^\beta. \quad (\text{S84})$$

S8. ADIABATIC PERTURBATION THEORY FOR THE EFFECTIVE SCHRÖDINGER EQUATION

In this appendix, we derive the first order correction to the adiabatic theorem for the time evolution of Bogoliubov modes (S48). This is a prerequisite for deriving the symplectic anomalous velocity term in the following Section S9. To this end, we adapt the standard theory of adiabatic perturbation theory for conventional systems [129, 130] to the symplectic Bogoliubov structure here.

We consider a time-dependent dynamical matrix $D(t) \equiv D(\boldsymbol{\lambda}(t))$, where the time dependence is encoded via some dimensionless parameters $\boldsymbol{\lambda}(t) = (\lambda^1(t), \dots, \lambda^r(t))$. At each point in time t , we assume that the Bogoliubov system is thermodynamically stable (the coefficient matrix $\mathcal{M}(\boldsymbol{\lambda}(t))$ is positive definite $\forall t$ [65]), such that an instantaneous eigenvalue equation of the following form holds (S29)

$$D(t) \mathbf{w}^n(t) = \tilde{\Omega}_n(t) \mathbf{w}^n(t), \quad (\text{S85})$$

where the time-dependence of the Bogoliubov modes $\mathbf{w}^n(t)$ is again specified through the parameters $\boldsymbol{\lambda}(t)$. In what follows, we assume a non-degenerate spectrum $\tilde{\Omega}_n \neq \tilde{\Omega}_m$ for $n \neq m$ for all times t . This assumption can be relaxed both regarding states that are not coupled to \mathbf{w}^n as a result of symmetry and when considering the adiabatic theorem with respect to a degenerate eigenspace, which is however beyond the scope of this work. We are interested in the time-evolution an arbitrary state $\boldsymbol{\psi}(t)$ in the space spanned by the Bogoliubov modes (S25),

$$\boldsymbol{\psi}(t) = \sum_n c_n(t) \mathbf{w}^n(t), \quad (\text{S86})$$

which is governed by the effective time-dependent Schrödinger equation (S48),

$$-i\partial_t \boldsymbol{\psi}(t) = D(t) \boldsymbol{\psi}(t). \quad (\text{S87})$$

Now, under the assumption of slow parameter variations of $\boldsymbol{\lambda}(t)$ (adiabatic limit), we would like to solve Eq. (S87) perturbatively. To this end, we first insert the instantaneous expansion of $\boldsymbol{\psi}(t)$ (S86) into the effective Schrödinger equation (S87), then project onto a specific mode via multiplying with $s_m(\mathbf{w}^m)^\dagger \tau_z$ from the left, by which we arrive

$$\dot{c}_m + \sum_n c_n s_m(\mathbf{w}^m)^\dagger \tau_z \partial_t \mathbf{w}^n = i\tilde{\Omega}_m c_m. \quad (\text{S88})$$

The explicit time-dependence will be omitted for brevity in the following. Next, we perform a gauge transformation $c_m(t) \rightarrow c_m(t) e^{i\xi_m(t)}$, where $\xi_m(t) = \int_0^t dt' \tilde{\Omega}_m(t')$. This removes the dynamical phase factor from the equations, which will, however, obviously be reintroduced at the end. This leads to the following differential equation for the time-dependent coefficients $c_m(t)$

$$\dot{c}_m = - \sum_n c_n s_m(\mathbf{w}^m)^\dagger \tau_z \partial_t \mathbf{w}^n e^{i\xi_{nm}(t)} \quad (\text{S89})$$

with $\xi_{nm}(t) \equiv \xi_n(t) - \xi_m(t)$. We formally integrate (S89) and obtain

$$c_m(t) = c_m(0) - \int_0^t dt' \sum_n c_n(t') s_m(\mathbf{w}^m)^\dagger \tau_z \partial_{\lambda^\mu} \mathbf{w}^n \dot{\lambda}^\mu e^{i\xi_{nm}(t')} \quad (\text{S90})$$

where we have used that $\partial_t \mathbf{w}^n(\boldsymbol{\lambda}(t)) = \sum_\mu \partial_\mu \mathbf{w}^n \dot{\lambda}^\mu \equiv \partial_\mu \mathbf{w}^n \dot{\lambda}^\mu$ with $\partial_\mu \equiv \frac{\partial}{\partial \lambda^\mu}$.

Let us now consider the situation, where the system is initially occupying a single Bogoliubov mode, i.e., $\boldsymbol{\psi}(0) = \mathbf{w}^m$ in (S86), such that $c_m(0) = 1$ and $c_n(0) = 0$ for $n \neq m$. In the limit of $\dot{\lambda}^\mu \rightarrow 0$ all transition amplitudes to other modes are highly suppressed, and the only leading order contribution in (S90) stems from the diagonal $n = m$ term in the sum [130]. Thus, to zeroth order, the solution for $c_m(t)$ is given by

$$c_m(t) = c_m(0) e^{i\gamma^m(t)}. \quad (\text{S91})$$

Here, γ^m denotes the symplectic version of the Berry phase [131]

$$\gamma^m(t) = i \int_0^t dt' s_m(\mathbf{w}^m)^\dagger \tau_z \partial_\mu \mathbf{w}^m \dot{\lambda}^\mu = \int_{\boldsymbol{\lambda}(0)}^{\boldsymbol{\lambda}(t)} A_\mu^m(\boldsymbol{\lambda}) d\lambda^\mu = \int A^m, \quad (\text{S92})$$

which may also written as an integral over the symplectic Berry connection $A^m = A_\mu^m d\lambda^\mu = [i s_m(\mathbf{w}^m)^\dagger \tau_z \partial_\mu \mathbf{w}^m] d\lambda^\mu$ [7, 17, 45, 97]. The result (S91) represents the conventional adiabatic theorem (adiabatic limit) adapted to the time evolution Bogoliubov modes, i.e., under slow parameter variations the system will remain in the initially occupied mode while acquiring an additional symplectic Berry phase $\gamma^m(t)$. We stress that this represents the 0-th order contribution, since Berry phase is independent of the rate of change in the parameters $\dot{\lambda}^\mu$ [cf. Eq. (S92)].

Next, we aim to solve (S90) perturbatively, to self-consistently obtain corrections to first order in $\dot{\lambda}^\mu$. To this end, we insert the 0-th order solution $c_m(t)$ (S91) into the right-hand side of the Eq. (S90), by which we obtain

$$c_m(t) \approx c_m(0) - \int_0^t dt' \sum_n c_n(0) e^{i\gamma^n(t')} s_m(\mathbf{w}^m)^\dagger \tau_z \partial_\mu \mathbf{w}^n \dot{\lambda}^\mu e^{i\xi_{nm}(t')} \quad (\text{S93})$$

Now for all other coefficients $c_k(t)$ with $k \neq m$, we only need to consider the $n = m$ term within the sum in (S93), where m is again the index of the initially occupied mode. To first order, this leads to the following equation for $c_{k \neq m}(t)$

$$\begin{aligned} c_k(t) &= - \int_0^t dt' c_m(0) e^{i\gamma^m(t')} s_k(\mathbf{w}^k)^\dagger \tau_z \partial_\mu \mathbf{w}^m \dot{\lambda}^\mu e^{i\xi_{km}(t')} \\ &= i \int_0^t dt' e^{i\gamma^m(t')} s_k \frac{(\mathbf{w}^k)^\dagger \tau_z \partial_\mu \mathbf{w}^m}{\tilde{\Omega}_{km}} \dot{\lambda}^\mu \frac{d}{dt'} e^{i\xi_{km}(t')}, \end{aligned} \quad (\text{S94})$$

with $\tilde{\Omega}_{km} \equiv \tilde{\Omega}_k - \tilde{\Omega}_m$ and where we have used that $\frac{d}{dt'} e^{i\xi_{km}(t')} = \tilde{\Omega}_{km} e^{i\xi_{km}(t')}$. Due to the fast oscillating exponential factor in Eq. (S94), we can integrate this expression by parts [130] and obtain the following solution for the off-diagonal coefficients

$$c_k(t) = i s_k e^{i\gamma^m(t)} \frac{(\mathbf{w}^k)^\dagger \tau_z \partial_\mu \mathbf{w}^m}{\tilde{\Omega}_{km}} \dot{\lambda}^\mu e^{i\xi_{km}(t)}. \quad (\text{S95})$$

Thus, to first order in $\dot{\lambda}^\mu$, the state $\psi(t)$ (S86) is finally given by

$$\begin{aligned} \psi(t) &= e^{-i(\xi_m(t) - \gamma^m(t))} \left[\mathbf{w}^m + i \sum_{k \neq m} \mathbf{w}^k s_k \frac{(\mathbf{w}^k)^\dagger \tau_z \partial_\mu \mathbf{w}^m}{(\tilde{\Omega}_{km})} \dot{\lambda}^\mu \right] \\ &= e^{-i(\xi_m(t) - \gamma^m(t))} \left[\mathbf{w}^m + i \sum_{k \neq m} \mathbf{w}^k s_k \frac{(\mathbf{w}^k)^\dagger \partial_\mu \mathcal{M} \mathbf{w}^m}{(\tilde{\Omega}_{km})^2} \dot{\lambda}^\mu \right], \end{aligned} \quad (\text{S96})$$

where we have reintroduced the dynamical phase $c_n(t) \rightarrow c_n(t) e^{-i\xi_n(t)}$ and utilized the symplectic version of the Hellmann-Feynman theorem (S32) in the second equality. Eq. (S96) constitutes the first order correction (in $\dot{\lambda}^\mu$) to the adiabatic theorem for the time evolution of Bogoliubov modes.

If required, one may also normalize the state $\psi(t)$ (S96) up to second order [132], i.e., up to $(\dot{\lambda}^\mu)^2 \equiv (v^\mu)^2$, which leads to

$$\psi(t) = e^{-i(\xi_m(t) - \gamma^m(t))} \left[(1 + \beta_\mu^m (v^\mu)^2) \mathbf{w}^m + i v^\mu \sum_{k \neq m} \alpha_\mu^{km} \mathbf{w}^k \right], \quad (\text{S97})$$

where we have defined $\alpha_\mu^{km} \equiv s_k \frac{(\mathbf{w}^k)^\dagger \partial_\mu \mathcal{M} \mathbf{w}^m}{(\tilde{\Omega}_{km})^2}$ with $\beta_\mu^m \equiv -\frac{1}{2} s_m \sum_{k \neq m} |\alpha_\mu^{km}|^2$.

S9. SYMPLECTIC ANOMALOUS VELOCITY

In this appendix, we provide a more detailed derivation of the symplectic anomalous velocity term, which is proportional to the symplectic Berry curvature. Consider a translationally invariant BdG system such that the Bogoliubov eigenmodes of the dynamical matrix D admit Bloch function type solution, i.e., $D\psi^n(\mathbf{q}) = D e^{i\mathbf{q}\cdot\mathbf{r}} \mathbf{w}^n(\mathbf{q}) = \tilde{\Omega}_n(\mathbf{q}) e^{i\mathbf{q}\cdot\mathbf{r}} \mathbf{w}^n(\mathbf{q})$. Here, $\mathbf{w}^n(\mathbf{q})$ is the cell-periodic part of the Bogoliubov Bloch function for the n -th band, which fulfill the Bogoliubov eigenvalue equation for the quasimomentum transformed dynamical matrix $D(\mathbf{q})\mathbf{w}^n(\mathbf{q}) = \tilde{\Omega}_n(\mathbf{q})\mathbf{w}^n(\mathbf{q})$. To make connections to the conventional anomalous velocity, we consider a two-dimensional time-dependent quasimomentum parameter space $\mathbf{q}(t) = (q_x(t), q_y(t))$. We imagine a uniform force \mathbf{F} to be applied to the Bogoliubov system of interest, such that quasimomenta satisfy $\dot{\mathbf{q}} = \mathbf{F}$, leading to $\mathbf{q}(t) = \mathbf{q}(0) + \mathbf{F}t$.

Next, we consider an initially populated Bogoliubov particle mode $\psi(0) = \mathbf{w}^n(\mathbf{q})$ and let the applied field point in the x -direction, i.e., $\mathbf{F} = F_x \mathbf{e}_x$. Using the results of Eq. (S96), we see that the solution to the time-dependent effective Schrödinger equation for the Bogoliubov modes is, to first order in F_x , given by

$$\psi(\mathbf{q}) = \mathbf{w}^n(\mathbf{q}) + i F_x \sum_{m \neq n} \mathbf{w}^m(\mathbf{q}) s_n \frac{(\mathbf{w}^m(\mathbf{q}))^\dagger \partial_{q_x} \mathcal{M} \mathbf{w}^n(\mathbf{q})}{(\tilde{\Omega}_n(\mathbf{q}) - \tilde{\Omega}_m(\mathbf{q}))^2} + \mathcal{O}(F_x^2), \quad (\text{S98})$$

where the irrelevant phase factors have been omitted for clarity. To obtain the velocity in the y -direction to first order in F_x , we take the expectation value of the quasiparticle velocity operator $v_y(\mathbf{q}) = \partial_{q_y} D(\mathbf{q})$ with respect to the

state of Eq. (S98), where $D(\mathbf{q})$ is again the dynamical matrix in the quasimomentum space representation. Note that $v_y(\mathbf{q}) = \partial_{q_y} D(\mathbf{q})$ just follows from the Heisenberg equations of motion for the position operator \mathbf{r} transformed into the quasimomentum space [61] and then extended to Nambu space. The velocity then equates to

$$\begin{aligned} \langle v_y(\mathbf{q}) \rangle &= (\boldsymbol{\psi}(\mathbf{q}))^\dagger \tau_z \partial_{q_y} D(\mathbf{q}) \boldsymbol{\psi}(\mathbf{q}) \\ &= (\mathbf{w}^n)^\dagger \partial_{q_y} \mathcal{M}(\mathbf{q}) \mathbf{w}^n + \left(i F_x \sum_{m \neq n} s_n \frac{\mathbf{w}^n \partial_{q_y} \mathcal{M} \mathbf{w}^m \times (\mathbf{w}^m)^\dagger \partial_{q_x} \mathcal{M} \mathbf{w}^n}{(\tilde{\Omega}_n - \tilde{\Omega}_m)^2} + \text{c.c.} \right) + \mathcal{O}(F_x^2). \end{aligned} \quad (\text{S99})$$

By making use of the eigenvalue equation for the Bogoliubov modes, $D(\mathbf{q}) \mathbf{w}^n(\mathbf{q}) = \tilde{\Omega}_n(\mathbf{q}) \mathbf{w}^n(\mathbf{q})$ (S29), as well as the spectral representation of the symplectic Berry curvature $B_{\mu\nu}^n = -2 \text{Im}[\eta_{\mu\nu}^n]$ (S33), we see that the velocity v_y of the Bogoliubov state in n -th band is given by

$$\langle v_y(\mathbf{q}) \rangle = \frac{\partial \tilde{\Omega}_n(\mathbf{q})}{\partial q_y} + F_x B_{xy}^n(\mathbf{q}). \quad (\text{S100})$$

By symmetry, we can immediately generalize this result (S100) to general directions of the applied force \mathbf{F}

$$\langle \mathbf{v}(\mathbf{q}) \rangle = \nabla_{\mathbf{q}} \tilde{\Omega}_n(\mathbf{q}) + \mathbf{B}_n(\mathbf{q}) \times \mathbf{F}, \quad (\text{S101})$$

with $\mathbf{B}_n(\mathbf{q}) \equiv \epsilon_{\mu\nu\kappa} B_{\mu\nu}^n(\mathbf{q}) \mathbf{e}_\kappa$. Here, the first term corresponds to the slope of the occupied Bogoliubov energy band $\tilde{\Omega}_n(\mathbf{q})$ and the second term is proportional to the symplectic Berry curvature $\mathbf{B}_n(\mathbf{q})$. The second term describes a transverse velocity component and may be viewed as a symplectic generalization of the conventional anomalous velocity term [61–63].

The results above may easily be generalized to the case of an initial Bogoliubov wave packet consisting of a superposition (particle) Bogoliubov modes $\boldsymbol{\psi}^n(\mathbf{q}) = e^{i\mathbf{q}\cdot\mathbf{r}} \mathbf{w}^n(\mathbf{q})$ of a given band n ,

$$\phi^n = \sum_{\mathbf{q}} c(\mathbf{q}) e^{i\mathbf{q}\cdot\mathbf{r}} \mathbf{w}^n(\mathbf{q}), \quad (\text{S102})$$

which is normalized with respect to the Bogoliubov norm, i.e., $\langle \phi^n | \tau_z | \phi^n \rangle = \sum_{\mathbf{q}} |c(\mathbf{q})|^2 = 1$ [97]. The velocity of such a Bogoliubov wave packet may then be related to a bulk observable, such as the total current density j_y in real space

$$j_y = \frac{1}{L^2} \sum_{\mathbf{q}} \rho(\mathbf{q}) \langle v_y(\mathbf{q}) \rangle = \int_{\text{1. BZ}} \frac{d^2 \mathbf{q}}{(2\pi)^2} \rho(\mathbf{q}) \langle v_y(\mathbf{q}) \rangle, \quad (\text{S103})$$

where L^2 is the area of the system and $\rho(\mathbf{q}) = |c(\mathbf{q})|^2$ the distribution function of Bogoliubov quasiparticles in the respective band. Although j_y is not quantized by the (symplectic) Chern number of the respective Bogoliubov band (for the typical case of a non-uniformly populated Bogoliubov band), the current density still gives rise to a finite transverse velocity contribution, which is directly related to the anomalous term governed by the underlying symplectic Berry curvature of the Bogoliubov band. We note in passing that special care may be required when switching to the continuum of quasimomentum states in Eq. (S103), which is however beyond the scope of this work [61, 133].

S10. BOGOLIUBOV HALDANE MODEL

In this appendix, we provide a brief derivation of the Bogoliubov-Haldane model considered as an example in the main text, which mainly follows Ref. [21]. Adding to this, we also review the numerical algorithm described in Refs. [2, 7] for obtaining the Bogoliubov matrices (S6) with the correct paraunitarity conditions (S20).

The starting point is a Bose-Hubbard model on a two-dimensional lattice with two sublattice sites,

$$\hat{H} = \sum_{\ell\ell', ss'} h_{\ell s, \ell' s'} \hat{a}_{\ell s}^\dagger \hat{a}_{\ell' s'} + \frac{U}{2} \sum_{\ell, s} \hat{n}_{\ell s} (\hat{n}_{\ell s} - 1) \quad (\text{S104})$$

where the elementary unit cells are labeled by ℓ , the sublattice sites by $s \in \{A, B\}$. The non-interacting (quadratic) part is governed by the matrix elements $h_{\ell s, \ell' s'}$ in position space and the interacting part by the on-site interaction strength U .

The first summand is the non-interacting part, which may diagonalized by first moving into quasimomentum space via

$$\mathcal{H}_{ss'}(\mathbf{q}) \equiv \sum_{\ell\ell'} h_{\ell s, \ell' s'} e^{-i(\mathbf{r}_{\ell s} - \mathbf{r}_{\ell' s'}) \cdot \mathbf{q}} = (h_0(\mathbf{q}) \mathbb{1} + \mathbf{h}(\mathbf{q}) \cdot \hat{\boldsymbol{\sigma}})_{ss'}, \quad (\text{S105})$$

where $\hat{\boldsymbol{\sigma}} = (\sigma_x, \sigma_y, \sigma_z)^T$ is the vector of Pauli matrices. At each quasimomentum point, \mathbf{q} , the Hamiltonian $\mathcal{H}_{ss'}(\mathbf{q})$ describes an operator acting on a two-dimensional Hilbert space, whose elements can be represented on the Bloch sphere. The north and south pole are denoted by $|\mathbf{q}A\rangle$ and $|\mathbf{q}B\rangle$ respectively. Due to the non-interacting nature of the system, the model can be further diagonalized by rotating into the eigenbasis $|\mathbf{q}\pm\rangle$,

$$|\mathbf{q}-\rangle = \sin(\theta_{\mathbf{q}}/2)|\mathbf{q}A\rangle - \cos(\theta_{\mathbf{q}}/2)e^{i\varphi_{\mathbf{q}}}|\mathbf{q}B\rangle, \quad |\mathbf{q}+\rangle = \cos(\theta_{\mathbf{q}}/2)|\mathbf{q}A\rangle + \sin(\theta_{\mathbf{q}}/2)e^{i\varphi_{\mathbf{q}}}|\mathbf{q}B\rangle. \quad (\text{S106})$$

which reside at $\pm\hat{\mathbf{h}}(\mathbf{q})$ on the Bloch sphere, with $\hat{\mathbf{h}}(\mathbf{q}) = \mathbf{h}(\mathbf{q})/|\mathbf{h}(\mathbf{q})| = (\cos(\varphi_{\mathbf{q}})\sin(\theta_{\mathbf{q}}), \sin(\varphi_{\mathbf{q}})\sin(\theta_{\mathbf{q}}), \cos(\theta_{\mathbf{q}}))^T$. The corresponding eigenenergies $\epsilon_{\pm}(\mathbf{q})$ of $|\mathbf{q}\pm\rangle$ are given by $\epsilon_{\pm}(\mathbf{q}) = h_0(\mathbf{q}) \pm |\mathbf{h}(\mathbf{q})|$ which define the band structure of the lattice. Now, an ideal Bose gas at zero temperature [with $U = 0$ in (S104)] forms a perfect Bose-Einstein condensate (BEC) characterized by all bosons condensing into the lowest energy eigenstate $|\mathbf{q}_0, -\rangle$ at some quasimomentum \mathbf{q}_0 in the lower band $\epsilon_-(\mathbf{q})$, with the ground state given by $|\psi_B\rangle = (\sqrt{N!})^{-1}(\hat{a}_{\mathbf{q}_0, -}^\dagger)^N|\text{vac}\rangle$. Here, N shall denote the total number of bosons in the system and $\hat{a}_{\mathbf{q}, \pm}^\dagger$ the creation operator for a boson at quasimomentum \mathbf{q} in the lower or upper band, respectively.

In this work, we consider the Haldane model [92] on a hexagonal lattice for which the non-interacting part (S105) reads,

$$\begin{aligned} h_x(\mathbf{q}) &= J_1 \sum_i \cos(\mathbf{q} \cdot \mathbf{d}_i), \quad h_y(\mathbf{q}) = J_1 \sum_i \sin(\mathbf{q} \cdot \mathbf{d}_i), \\ h_z(\mathbf{q}) &= \Delta + 2J_2 \sin(\theta) \sum_i \sin(\mathbf{q} \cdot \mathbf{a}_i), \quad h_0(\mathbf{q}) = 2J_2 \cos(\theta) \sum_i \cos(\mathbf{q} \cdot \mathbf{a}_i), \end{aligned} \quad (\text{S107})$$

where $J_1 (J_2 e^{i\theta})$ is the real nearest neighbor (complex next-nearest neighbor) hopping amplitude and Δ the mass term (sublattice offset). The complex next-nearest neighbor hopping amplitude breaks the time reversal symmetry of the system, and the mass term Δ breaks the inversion symmetry of the lattice. On the hexagonal lattice, translation vectors connecting the nearest neighbors are denoted by \mathbf{d}_i , while the vectors \mathbf{a}_i connect the next-nearest neighbors [Fig. S1],

$$\mathbf{d}_3 = \frac{a}{2} \begin{pmatrix} \sqrt{3} \\ -1 \end{pmatrix}, \quad \mathbf{d}_2 = \frac{a}{2} \begin{pmatrix} -\sqrt{3} \\ -1 \end{pmatrix}, \quad \mathbf{d}_1 = -(\mathbf{d}_2 + \mathbf{d}_3) = \mathbf{a} \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (\text{S108})$$

$$\mathbf{a}_3 = \mathbf{d}_3 - \mathbf{d}_2, \quad \mathbf{a}_2 = \mathbf{d}_1 - \mathbf{d}_3, \quad \mathbf{a}_1 = \mathbf{d}_1 - \mathbf{d}_2, \quad (\text{S109})$$

where a is the lattice constant. The reciprocal lattice (quasimomentum space) is spanned by the reciprocal lattice vectors \mathbf{b}_i

$$\mathbf{b}_1 = \frac{2\pi}{3a} \begin{pmatrix} \sqrt{3} \\ 1 \end{pmatrix}, \quad \mathbf{b}_2 = \frac{2\pi}{3a} \begin{pmatrix} -\sqrt{3} \\ 1 \end{pmatrix}, \quad \mathbf{b}_3 = \mathbf{b}_1 + \mathbf{b}_2, \quad (\text{S110})$$

such that $\mathbf{a}_i \cdot \mathbf{b}_j = 2\pi\delta_{ij}$ for $i, j \in \{1, 2\}$ [see Fig. S1].

Having discussed the non-interacting part, we turn to the Bogoliubov theory, which accounts for additional quantum (and thermal) fluctuations on top of a mean-field solution for a weakly interacting bosonic system [123, 124, 134]. For this, we first obtain a mean-field solution via a minimization of an associated Gross-Pitaevskii (GP) energy functional. The stationary solutions gained by minimizing the GP energy functional will then yield the mean-field ground state, which is assumed to be macroscopically occupied. Then, we will describe quantum fluctuations on top of the condensate, by considering non-condensed deviations from this mean-field solution. This procedure can be formally understood by replacing the operators within the Hamiltonian \hat{H} (S104) via $\hat{a}_{\ell s} \rightarrow \sqrt{n_0}\zeta_{\ell s} + \hat{a}_{\ell s}$, where the first term denotes the complex-valued mean-field solution, $\zeta_{\ell s} \in \mathbb{C}$, and the second term the additional quantum fluctuations where n_0 denotes condensate density of the system. Expanding the Bose-Hubbard Hamiltonian (S104) in orders of the condensate density n_0 then yields [20]

$$\hat{H} = n_0 E_{\text{GP}} + n_0^{1/2} \hat{H}^{(L)} + \hat{H}^{(B)} + \mathcal{O}(n_0^{-1/2}). \quad (\text{S111})$$




FIG. S1. (a) Illustration of the Bogoliubov-Haldane model (S122) on a hexagonal lattice with two sublattice sites A (B) with potential energy offset Δ ($-\Delta$), real (complex-valued) tunneling matrix element J_1 ($J_2e^{i\theta}$) for the nearest (next-) nearest neighbors, and on-site interaction strength U . The translation vectors connecting the (next-) nearest neighbors (\mathbf{d}_i) \mathbf{a}_i (S109) are shown. (b) Illustration of reciprocal lattice of the hexagonal lattice structure of (a) including the reciprocal lattice vectors \mathbf{b}_i (S110). In the 1st Brillouin zone, we depict a high-symmetry path along the high-symmetry points Γ , K , M , and K' .

The first term denotes the GP energy functional E_{GP} which is a function of the mean-field solutions $\zeta_{\ell s}$. The next two terms $\hat{H}^{(L)}$ and $\hat{H}^{(B)}$ are linear and quadratic in the bosonic operators $\hat{a}_{\ell s}^{(\dagger)}$, where the latter is known as the Bogoliubov Hamiltonian, which is sought after in the following.

First, the energy functional E_{GP} is found by replacing the operators $\hat{a}_{\ell s}^{(\dagger)}$ in \hat{H} by complex-valued functions $\sqrt{n}\zeta_{\ell s}^{(*)}$, where it is assumed that almost all bosons are condensed, i.e., $n_0 \approx n = N/V$, with $V = 2N_{uc}$ and N denoting the total number of bosons in the system and N_{uc} the number of unit cells in the lattice [20]. For the parameters of the bosonic Haldane model considered here, the condensate is formed at the single zero quasimomentum mode $\mathbf{q} = \mathbf{0}$ in the lower band such that a suitable ansatz for the ground state is given by a spatially homogeneous function $\zeta_{\ell s} = \chi_s$, $s \in \{A, B\}$. This leads to the following energy functional in quasimomentum space

$$E_{GP} = (\chi_A^* \chi_B^*) (\mathcal{H}(\mathbf{q} = \mathbf{0}) - \mu_{\text{eff}} \mathbb{1}_2) \begin{pmatrix} \chi_A \\ \chi_B \end{pmatrix} + \frac{U}{2} n (|\chi_A|^4 + |\chi_B|^4), \quad (\text{S112})$$

where we have introduced μ_{eff} as a Lagrange multiplier for satisfying the conservation of the total number of bosons in the system, i.e., $\sum_{\ell s} |\zeta_{\ell s}|^2 = N$. We minimize E_{GP} by taking the variational derivative of E_{GP} with respect to χ_s^* which leads to the so-called GP equations,

$$(\mathcal{H}(\mathbf{q} = \mathbf{0}) - \mu_{\text{eff}} \mathbb{1}_2) \begin{pmatrix} \chi_A \\ \chi_B \end{pmatrix} + nU \begin{pmatrix} |\chi_A|^2 \chi_A \\ |\chi_B|^2 \chi_B \end{pmatrix} = 0. \quad (\text{S113})$$

Since the non-interacting condensate is formed at the $|\mathbf{q} = \mathbf{0}, -\rangle$ mode we parametrize χ_s via the Bloch sphere representation $|\mathbf{q} = \mathbf{0}, -\rangle$ (S106) [21]

$$\begin{pmatrix} \chi_A \\ \chi_B \end{pmatrix} = \begin{pmatrix} \sin(\theta/2) \\ -\cos(\theta/2)e^{i\varphi} \end{pmatrix}. \quad (\text{S114})$$

Multiplying (S113) by $(\chi_A^* \chi_B^*)$ and $(-\chi_B^* \chi_A^*)$ from the left yields three equations [21, 135]

$$(h_0(\mathbf{0}) - \mu_{\text{eff}}) - h(\mathbf{0})[\cos(\theta_0)\cos(\theta) + \sin(\theta_0)\sin(\theta)\cos(\varphi - \varphi_0)] + 2Un(|\chi_A|^4 + |\chi_B|^4) = 0, \quad (\text{S115})$$

$$h(\mathbf{0})[\cos(\theta_0)\sin(\theta)\cos(\varphi) - \cos(\varphi_0)\sin(\theta_0)\cos(\theta)] - Un\sin(\theta)\cos(\theta)\cos(\varphi) = 0, \quad (\text{S116})$$

$$(h_0(\mathbf{0}) - \mu_{\text{eff}})\sin(\theta)\sin(\varphi) + h(\mathbf{0})\sin(\theta_0)\sin(\varphi_0) + Un\sin(\theta)\sin(\varphi) = 0, \quad (\text{S117})$$

where $\mathbf{h}(\mathbf{0}) = h(\mathbf{0}) (\cos(\varphi_0)\sin(\theta_0), \sin(\varphi_0)\sin(\theta_0), \cos(\theta_0))^T$ and $h_0(\mathbf{0})$ stem from (S105). For a given interaction energy U and density n , we numerically minimize the Eqs. (S115) to (S117) and gain the stationary solutions $(\theta, \varphi, \mu_{\text{eff}})$ specifying the mean-field ground state in (S114).

Since the BEC occurs in the lowest band at $\mathbf{q} = \mathbf{0}$, we further specify the unitary rotation (provided by the eigenstates of the non-interacting system (S106)) for the obtained solution [21]

$$\begin{pmatrix} \hat{a}_{\mathbf{0},A} \\ \hat{a}_{\mathbf{0},B} \end{pmatrix} = U(\theta, \varphi) \begin{pmatrix} \hat{a}_{\mathbf{0},+} \\ \hat{a}_{\mathbf{0},-} \end{pmatrix} = \begin{pmatrix} f_{A,+} & f_{A,-} \\ f_{B,+} & f_{B,-} \end{pmatrix} \begin{pmatrix} \hat{a}_{\mathbf{0},+} \\ \hat{a}_{\mathbf{0},-} \end{pmatrix}, \quad (\text{S118})$$

$$\text{with } U(\theta, \varphi) = \begin{pmatrix} \cos(\theta/2) & \sin(\theta/2) \\ \sin(\theta/2)e^{i\varphi} & -\cos(\theta/2)e^{i\varphi} \end{pmatrix}, \quad (\text{S119})$$

where $f_{s,\pm}(\theta, \varphi)$ with $s \in \{A, B\}$ denote the stationary solutions of the GP functional (S112).

The next step is to go beyond this mean-field description and derive the corresponding Bogoliubov Hamiltonian $\hat{H}^{(B)}$ (S111). The starting point is the quasimomentum space representation of the Bose-Hubbard Hamiltonian (S104)

$$\hat{H} = \sum_{\mathbf{q}} \sum_{s,s'} \hat{a}_{\mathbf{q},s}^\dagger \mathcal{H}_{s,s'}(\mathbf{q}) \hat{a}_{\mathbf{q},s'} + \frac{U}{V} \sum_{\mathbf{q},\mathbf{q}',\mathbf{q}''} \sum_s \hat{a}_{\mathbf{q}+\mathbf{q}'',s}^\dagger \hat{a}_{\mathbf{q}'-\mathbf{q}'',s} \hat{a}_{\mathbf{q},s} \hat{a}_{\mathbf{q}',s}, \quad (\text{S120})$$

where $\hat{a}_{\mathbf{q},s} = \sum_{\ell} \langle \mathbf{q}s | \ell s \rangle \hat{a}_{\ell s}$ denote the bosonic operators transformed into quasimomentum space. Utilizing another rotation into the eigenbasis (S106) the bosonic operators in quasimomentum space are decomposed into a condensed and a non-condensed part via [21]

$$\sum_{\mathbf{q}} \hat{a}_{\mathbf{q},s} = \hat{a}_{\mathbf{0},s} + \sum_{\mathbf{q} \neq \mathbf{0}} \hat{a}_{\mathbf{q},s} = f_{s,-} \hat{a}_{\mathbf{0},-} + f_{s,+} \hat{a}_{\mathbf{0},+} + \sum_{\mathbf{q} \neq \mathbf{0}} \hat{a}_{\mathbf{q},s}, \quad (\text{S121})$$

where the condensation mode is assumed to be located at $\mathbf{q} = \mathbf{0}$ in the lowest band and $f_{s,\pm} \equiv f_{s,\pm}(\varphi, \theta)$ are functions depending on the stationary solutions of the energy functional E_{GP} (S112).

Next, the decomposition (S121) is inserted into \hat{H} (S120) and the Hamiltonian is separated into parts of $\mathbf{q} = \mathbf{0}$ and $\mathbf{q} \neq \mathbf{0}$ while only terms up to second order in $\hat{a}_{\mathbf{q} \neq \mathbf{0},s}^\dagger$ are retained, leading to an approximate Hamiltonian. For the actual Bogoliubov approximation, the condensation mode is then replaced via $\hat{a}_{\mathbf{0},-} \rightarrow \sqrt{N_0}$, where N_0 denotes the number of particles in the condensate mode. Performing all these steps will eventually lead to the following Bogoliubov Hamiltonian in quasimomentum space

$$\begin{aligned} \hat{H}^{(B)} = & \frac{1}{2} \sum_{\mathbf{q} \neq \mathbf{0}} \left(\hat{a}_{\mathbf{q},A}^\dagger \hat{a}_{\mathbf{q},B}^\dagger \hat{a}_{-\mathbf{q},A}, \hat{a}_{-\mathbf{q},B} \right) \begin{pmatrix} \mathcal{H}(\mathbf{q}) + \mathcal{H}_1 & 2\mathcal{H}_2 \\ 2\mathcal{H}_2^* & (\mathcal{H}(-\mathbf{q}) + \mathcal{H}_1)^* \end{pmatrix} \begin{pmatrix} \hat{a}_{\mathbf{q},A} \\ \hat{a}_{\mathbf{q},B} \\ \hat{a}_{-\mathbf{q},A}^\dagger \\ \hat{a}_{-\mathbf{q},B}^\dagger \end{pmatrix} \\ & + \frac{1}{2} \left(\hat{a}_{\mathbf{0},+}^\dagger, \hat{a}_{\mathbf{0},+} \right) \begin{pmatrix} h_3 & 2h_4 \\ 2h_4^* & h_3^* \end{pmatrix} \begin{pmatrix} \hat{a}_{\mathbf{0},+} \\ \hat{a}_{\mathbf{0},+}^\dagger \end{pmatrix} + \text{const.} \end{aligned} \quad (\text{S122})$$

We can also compactly write this Bogoliubov Hamiltonian (S122) in terms of the Nambu spinor representation

$$\hat{H}^{(B)} = \frac{1}{2} \sum_{\mathbf{q} \neq \mathbf{0}} \Psi_{\mathbf{q}}^\dagger \mathcal{M}_{\mathbf{q}} \Psi_{\mathbf{q}} + \frac{1}{2} \Psi_+^\dagger \mathcal{M}_+ \Psi_+ + \text{const.}, \quad (\text{S123})$$

where $\Psi_{\mathbf{q}}^\dagger \equiv \left(\hat{a}_{\mathbf{q},A}^\dagger, \hat{a}_{\mathbf{q},B}^\dagger, \hat{a}_{-\mathbf{q},A}, \hat{a}_{-\mathbf{q},B} \right)$ and $\Psi_+^\dagger \equiv \left(\hat{a}_{\mathbf{0},+}^\dagger, \hat{a}_{\mathbf{0},+} \right)$. Here, \mathcal{M}_+ is the 2×2 matrix and $\mathcal{M}_{\mathbf{q}}$ the quasimomentum dependent 4×4 matrix from (S122), which are explicitly given by [21, 135]

$$\mathcal{M}_{\mathbf{q}} = \begin{pmatrix} \mathcal{H}(\mathbf{q}) + \mathcal{H}_1 & 2\mathcal{H}_2 \\ 2\mathcal{H}_2^* & (\mathcal{H}(-\mathbf{q}) + \mathcal{H}_1)^* \end{pmatrix}, \quad \mathcal{M}_+ = \begin{pmatrix} h_3 & 2h_4 \\ 2h_4^* & h_3^* \end{pmatrix}, \quad (\text{S124})$$

$$\text{with } \mathcal{H}_1 = 4Un |F_-|^2 - \mu_{\text{eff}} \mathbb{1}_2, \quad \mathcal{H}_2 = Un(F_-)^2,$$

where $F_- = \text{diag}(f_{A,-}, f_{B,-})$ and where the entries of $\mathcal{H}(\mathbf{q})$ are given by (S105) and (S107). The entries of \mathcal{M}_+ are given by

$$h_3 = 4Un \sum_s \left(|f_{s,-}|^2 |f_{s,+}|^2 \right) + \sum_{s,s'} f_{s,+}^* \mathcal{H}_{s,s'}(0) f_{s',+} - \mu_{\text{eff}}, \quad h_4 = Un \sum_s (f_{s,-})^2 (f_{s,+}^*)^2, \quad (\text{S125})$$

where μ_{eff} follows the stationary solutions of Eqs. (S115) to (S117).

The Bogoliubov Hamiltonian (S122) can now be diagonalized by applying the paraunitary (Bogoliubov transformation) $W_{\mathbf{q}} (W_+)$ [21]

$$\begin{pmatrix} \hat{a}_{\mathbf{q},A} \\ \hat{a}_{\mathbf{q},B} \\ \hat{a}_{-\mathbf{q},A}^\dagger \\ \hat{a}_{-\mathbf{q},B}^\dagger \end{pmatrix} = W_{\mathbf{q}} \begin{pmatrix} \hat{b}_{\mathbf{q},+} \\ \hat{b}_{\mathbf{q},-} \\ \hat{b}_{-\mathbf{q},+}^\dagger \\ \hat{b}_{-\mathbf{q},-}^\dagger \end{pmatrix} \Leftrightarrow \Psi_{\mathbf{q}} = W_{\mathbf{q}}\beta_{\mathbf{q}}, \quad \begin{pmatrix} \hat{a}_{\mathbf{0},+} \\ \hat{a}_{\mathbf{0},+}^\dagger \end{pmatrix} = W_+ \begin{pmatrix} \hat{b}_{\mathbf{0},+} \\ \hat{b}_{\mathbf{0},+}^\dagger \end{pmatrix} \Leftrightarrow \Psi_+ = W_+\beta_+ \quad (\text{S126})$$

with $\beta_{\mathbf{q}} \equiv \left(\hat{b}_{\mathbf{q},+}, \hat{b}_{\mathbf{q},-}, \hat{b}_{-\mathbf{q},+}^\dagger, \hat{b}_{-\mathbf{q},-}^\dagger \right)^T$ and $\beta_+ \equiv \left(\hat{b}_{\mathbf{0},+}, \hat{b}_{\mathbf{0},+}^\dagger \right)^T$ consisting of the new bosonic quasi-particle annihilation (creation) operators $\hat{b}_{\mathbf{q},\pm}^{(\dagger)}$ at quasimomentum \mathbf{q} in the upper (+) or lower Bogoliubov energy band (-). The 4×4 (2×2) Bogoliubov transformation matrix $W_{\mathbf{q}} (W_+)$ can be parametrized (S6) as

$$W_+ = \begin{pmatrix} u_{\mathbf{0}} & v_{\mathbf{0}}^* \\ v_{\mathbf{0}} & u_{\mathbf{0}}^* \end{pmatrix}, \quad W_{\mathbf{q}} = \begin{pmatrix} U_{\mathbf{q}} & V_{-\mathbf{q}}^* \\ V_{\mathbf{q}} & U_{-\mathbf{q}}^* \end{pmatrix} \quad (\text{S127})$$

$$\text{with } U_{\mathbf{q}} = \begin{pmatrix} u_{A,+}^{\mathbf{q}} & u_{A,-}^{\mathbf{q}} \\ u_{B,+}^{\mathbf{q}} & u_{B,-}^{\mathbf{q}} \end{pmatrix}, \quad V_{\mathbf{q}} = \begin{pmatrix} v_{A,+}^{\mathbf{q}} & v_{A,-}^{\mathbf{q}} \\ v_{B,+}^{\mathbf{q}} & v_{B,-}^{\mathbf{q}} \end{pmatrix}. \quad (\text{S128})$$

Since these quasi-particle operators still have to obey the bosonic commutation relations, the Bogoliubov transformation matrix is paraunitary [cf. Eq. (S20)], i.e., $W^\dagger \tau W = W \tau W^\dagger = \tau$, with $\tau = \sigma_z \otimes \mathbb{1}_4$ ($\tau = \sigma_z \otimes \mathbb{1}_2$) for $W = W_{\mathbf{q}}$ ($W = W_+$). These paraunitary matrices are specifically constructed [2, 7] such that

$$W_{\mathbf{q}}^\dagger \mathcal{M}_{\mathbf{q}} W_{\mathbf{q}} = \text{diag} (E_+(\mathbf{q}), E_-(\mathbf{q}), E_+(-\mathbf{q}), E_-(-\mathbf{q})), \quad W_+^\dagger \mathcal{M}_+ W_+ = \text{diag} (E_+(\mathbf{0}), E_+(\mathbf{0})), \quad (\text{S129})$$

holds, by which the Bogoliubov Hamiltonian is eventually diagonalized [21] to

$$\hat{H}^{(B)} = \sum_{\mathbf{q}} E_+(\mathbf{q}) \hat{b}_{\mathbf{q},+}^\dagger \hat{b}_{\mathbf{q},+} + \sum_{\mathbf{q} \neq \mathbf{0}} E_-(\mathbf{q}) \hat{b}_{\mathbf{q},-}^\dagger \hat{b}_{\mathbf{q},-} + \text{const.} \quad (\text{S130})$$

Here, $E_{\pm}(\mathbf{q})$ is the Bogoliubov energy spectrum of the upper (+) and lower (-) band.

The Bogoliubov energy spectrum can be obtained by (numerically) solving the energy eigenvalue problem (S29) of the (non-hermitian) dynamical matrix $D(\mathbf{q}) = \tau_z \mathcal{M}_{\mathbf{q}}$, i.e., $D(\mathbf{q}) \mathbf{w}^n(\mathbf{q}) = \tilde{\Omega}_j \mathbf{w}^n(\mathbf{q})$. However, the eigenmodes obtained from this diagonalization procedure do *not* lead to the correct paraunitary conditions for the Bogoliubov modes (S20).

As described in Refs. [2, 7], the correct paraunitary Bogoliubov matrices can be obtained via an algorithm, which includes a Cholesky decomposition of the coefficient matrices ($\mathcal{M}_{\mathbf{q}}, \mathcal{M}_+$). For completeness, we briefly review this algorithm in the next section.

S10A. Algorithm for constructing paraunitary Bogoliubov transformation matrix

Here, mainly following Refs. [2, 7], we briefly discuss how to (numerically) obtain a paraunitary Bogoliubov transformation matrix such as $W_{\mathbf{q}}$ and W_+ (S127) when given a bosonic BdG Hamiltonian.

We discuss the algorithm for a translationally invariant bosonic BdG Hamiltonian such that we have quasimomentum dependent $2N \times 2N$ (hermitian) coefficient matrix $\mathcal{M}_{\mathbf{q}}$ (S1) of the form

$$\mathcal{M}_{\mathbf{q}} = \begin{pmatrix} K_{\mathbf{q}} & G_{\mathbf{q}} \\ G_{-\mathbf{q}}^* & K_{-\mathbf{q}}^* \end{pmatrix}, \quad (\text{S131})$$

with $K_{\mathbf{q}}$ and $G_{\mathbf{q}}$ being quasimomentum dependent $N \times N$ matrices with N denoting the number internal degrees of freedom (e.g., the number sublattice sites within a unit cell). For each quasimomentum \mathbf{q} the bosonic system can then be diagonalized by a paraunitary matrix $W_{\mathbf{q}}$ (S18) such that

$$W_{\mathbf{q}}^\dagger \mathcal{M}_{\mathbf{q}} W_{\mathbf{q}} = \begin{pmatrix} \omega_{\mathbf{q}} & 0 \\ 0 & \omega_{-\mathbf{q}} \end{pmatrix}, \quad \text{with } \omega_{\mathbf{q}} = \text{diag} (\omega_{1,\mathbf{q}}, \dots, \omega_{N,\mathbf{q}}), \quad (\text{S132})$$

where $W_{\mathbf{q}}$ again fulfill the paraunitary conditions (S20),

$$W_{\mathbf{q}}^{\dagger} \tau_z W_{\mathbf{q}} = W_{\mathbf{q}} \tau_z W_{\mathbf{q}}^{\dagger} = \tau_z \text{ with } \tau_z = \sigma_z \otimes \mathbb{1}_N. \quad (\text{S133})$$

Now, given that $\mathcal{M}_{\mathbf{q}}$ is a positive-definite matrix (all eigenvalues of $\mathcal{M}_{\mathbf{q}}$ are strictly positive), the paraunitary matrix $W_{\mathbf{q}}$ can then be constructed via the following steps [2].

First, we apply a Cholesky decomposition to the coefficient matrix $\mathcal{M}_{\mathbf{q}}$ (which is possible since $\mathcal{M}_{\mathbf{q}}$ is positive-definite) (S131), which decomposes $\mathcal{M}_{\mathbf{q}}$ into a product between an upper triangular matrix $K_{\mathbf{q}}$ and its hermitian conjugate $K_{\mathbf{q}}^{\dagger}$, i.e., $\mathcal{M}_{\mathbf{q}} = K_{\mathbf{q}}^{\dagger} K_{\mathbf{q}}$. We note that the positive-definiteness of $\mathcal{M}_{\mathbf{q}}$ also ensures the existence of the inverse $K_{\mathbf{q}}^{-1}$.

Second, we introduce a hermitian matrix $T_{\mathbf{q}} \equiv K_{\mathbf{q}} \tau_z K_{\mathbf{q}}^{\dagger}$ from which we obtain the unitary matrix $U_{\mathbf{q}}$ via the eigendecomposition of $T_{\mathbf{q}}$,

$$U_{\mathbf{q}}^{\dagger} T_{\mathbf{q}} U_{\mathbf{q}} = \begin{pmatrix} \omega_{\mathbf{q}} & 0 \\ 0 & -\omega_{-\mathbf{q}} \end{pmatrix}. \quad (\text{S134})$$

These two diagonal matrices $\omega_{\mathbf{q}}$ and $\omega_{-\mathbf{q}}$ already provide the Bogoliubov energy spectrum of Eq. (S132) [7]. One can further show that both $\omega_{\mathbf{q}}$ and $\omega_{-\mathbf{q}}$ are real and strictly positive [2].

Third, the paraunitary matrix $W_{\mathbf{q}}$ is then constructed via

$$W_{\mathbf{q}} = K_{\mathbf{q}}^{-1} U_{\mathbf{q}} \begin{pmatrix} \omega_{\mathbf{q}}^{1/2} & 0 \\ 0 & \omega_{-\mathbf{q}}^{1/2} \end{pmatrix}, \quad (\text{S135})$$

which will diagonalize the coefficient matrix $\mathcal{M}_{\mathbf{q}}$ (S131) and importantly fulfill the paraunitary conditions (S133). This concludes the construction of the paraunitary matrix $W_{\mathbf{q}}$ for each quasimomentum \mathbf{q} .

One can then show the diagonalization of Eq. (S132) by the direct application of (S135)

$$W_{\mathbf{q}}^{\dagger} \mathcal{M}_{\mathbf{q}} W_{\mathbf{q}} = W_{\mathbf{q}}^{\dagger} K_{\mathbf{q}}^{\dagger} K_{\mathbf{q}} W_{\mathbf{q}} = \begin{pmatrix} \omega_{\mathbf{q}}^{1/2} & 0 \\ 0 & \omega_{-\mathbf{q}}^{1/2} \end{pmatrix} U_{\mathbf{q}}^{\dagger} (K_{\mathbf{q}}^{\dagger})^{-1} K_{\mathbf{q}}^{\dagger} K_{\mathbf{q}} K_{\mathbf{q}}^{-1} U_{\mathbf{q}} \begin{pmatrix} \omega_{\mathbf{q}}^{1/2} & 0 \\ 0 & \omega_{-\mathbf{q}}^{1/2} \end{pmatrix} = \begin{pmatrix} \omega_{\mathbf{q}} & 0 \\ 0 & \omega_{-\mathbf{q}} \end{pmatrix}. \quad (\text{S136})$$

Moreover, we can see that the construction of $W_{\mathbf{q}}$ (S135) also fulfills the paraunitary conditions (S133),

$$W_{\mathbf{q}}^{\dagger} \tau_z W_{\mathbf{q}} = \begin{pmatrix} \omega_{\mathbf{q}}^{1/2} & 0 \\ 0 & \omega_{-\mathbf{q}}^{1/2} \end{pmatrix} U_{\mathbf{q}}^{\dagger} (K_{\mathbf{q}}^{\dagger})^{-1} \tau_z K_{\mathbf{q}}^{-1} U_{\mathbf{q}} \begin{pmatrix} \omega_{\mathbf{q}}^{1/2} & 0 \\ 0 & \omega_{-\mathbf{q}}^{1/2} \end{pmatrix} = \begin{pmatrix} \omega_{\mathbf{q}}^{1/2} & 0 \\ 0 & \omega_{-\mathbf{q}}^{1/2} \end{pmatrix} U_{\mathbf{q}}^{\dagger} T_{\mathbf{q}}^{-1} U_{\mathbf{q}} \begin{pmatrix} \omega_{\mathbf{q}}^{1/2} & 0 \\ 0 & \omega_{-\mathbf{q}}^{1/2} \end{pmatrix} \quad (\text{S137})$$

$$= \begin{pmatrix} \omega_{\mathbf{q}}^{1/2} & 0 \\ 0 & \omega_{-\mathbf{q}}^{1/2} \end{pmatrix} \begin{pmatrix} \omega_{\mathbf{q}}^{-1} & 0 \\ 0 & -\omega_{-\mathbf{q}}^{-1} \end{pmatrix} \begin{pmatrix} \omega_{\mathbf{q}}^{1/2} & 0 \\ 0 & \omega_{-\mathbf{q}}^{1/2} \end{pmatrix} = \begin{pmatrix} \mathbb{1}_N & 0 \\ 0 & -\mathbb{1}_N \end{pmatrix} = \tau_z, \quad (\text{S138})$$

where in the first step we have just inserted the definition of $W_{\mathbf{q}}$ (S135). In the second equality, we used that $T_{\mathbf{q}}^{-1} = (K_{\mathbf{q}}^{\dagger})^{-1} \tau_z K_{\mathbf{q}}^{-1}$, which itself directly follows from $T_{\mathbf{q}} = K_{\mathbf{q}} \tau_z K_{\mathbf{q}}^{\dagger}$. In the second equality, following from (S134)

we used that $U_{\mathbf{q}}^{\dagger} T_{\mathbf{q}}^{-1} U_{\mathbf{q}} = \begin{pmatrix} \omega_{\mathbf{q}}^{-1} & 0 \\ 0 & -\omega_{-\mathbf{q}}^{-1} \end{pmatrix}$.

S10B. SQGT from integrated Bogoliubov excitation rates

Here, we briefly discuss how we numerically implement our proposed method to extract the SQGT from the integrated Bogoliubov excitation rate applied to the Bogoliubov-Haldane model (S122). We start with an initially populated Bogoliubov particle mode $\mathbf{w}^n(\mathbf{q})$ which is obtained from the paraunitary Bogoliubov transformation matrix (S126). The time evolution of the Bogoliubov particle mode $\mathbf{w}^n(\mathbf{q}, t)$ is performed via (S48) in the translationally invariant co-moving frame. Specifically, when the system is shaken along the x -direction the time-dependent dynamical matrix is given by Eq. (S64). The total excitation rate into other Bogoliubov in response to this perturbation is then computed via

$$\Gamma_{\mathbf{q}}^n(\omega) = \frac{1}{t} \sum_m s_m |(\mathbf{w}^m(\mathbf{q}))^{\dagger} \tau_z \mathbf{w}^n(\mathbf{q}, t)|^2, \quad (\text{S139})$$




FIG. S2. Integrated and ideal symplectic quantum metric component $g_{yy}(\mathbf{q})$ and $g_{xy}(\mathbf{q})$ for the upper particle band of the Bogoliubov-Haldane model (S122). (a) Integrated (left) and ideal (right) symplectic quantum metric component $g_{yy}(\mathbf{q})$ for $U_n/J_1 = 3$ shown in 2D quasimomentum space. (b) Same as in (a) but for the off-diagonal component $g_{xy}(\mathbf{q})$. (c) Ideal (solid line) and integrated (crosses) symplectic quantum metric component $g_{yy}(\mathbf{q})$ along a high-symmetry path in quasimomentum space for two interactions strengths $U_n/J_1 = 0$ (non-interacting Haldane model) and $U_n/J_1 = 3$. (d) Same as in (c) but for the off-diagonal component $g_{xy}(\mathbf{q})$. All the simulations have been performed for driving strengths $A/J_1 = 0.01$ up to a final time of $J_1 t/\hbar = 10$ for a frequency range of $\hbar\omega/J_1 \in [0.1(0.2), 5.5]$, for $U_n/J_1 = 3$ (0) with a spacing of $\delta\omega = 0.05$ over which the excitation rates have been integrated. The red-colored hexagon in (a) and (b) indicates the first Brillouin zone. Other parameters of the underlying Haldane model (S107) are the same as in Fig. 3 in the main text, i.e., $J_2 = 0.1J_1$, $\theta = 0.5\pi$, $\Delta/(3\sqrt{3}J_2) = 0.5$.

where t is the final integration time and ω the frequency of the perturbation (S64). Integrating (S139) over a range of probing frequencies ω , $\Gamma_{\text{int},\mathbf{q}}^n = \int d\omega \Gamma_{\mathbf{q}}^n(\omega)$, then allows us to obtain the symplectic quantum metric component $g_{xx}(\mathbf{q})$ via (S67). This similarly applies to the $g_{yy}(\mathbf{q})$ component of the SQGT when the system is shaken along the y -direction where the dynamical matrix in co-moving frame becomes $D'_y(t) = D(\mathbf{q}) + (2A/\omega) \sin(\omega t) \partial_{q_y} D(\mathbf{q}) + \mathcal{O}((A/\omega)^2)$. For the off-diagonal components the lattice is shaken along both directions (S68), the time-dependent dynamical matrix is given by Eq. (S70). The total excitation rates $\Gamma_{\mathbf{q}}^{n,\pm}(\omega)$ are obtained in the same way as in (S139). Taking the differential integrated rate, $\Delta\Gamma_{\text{int}}^n = \Gamma_{\text{int}}^{n,+} - \Gamma_{\text{int}}^{n,-} = \int d\omega (\Gamma_{\mathbf{q}}^{n,+}(\omega) - \Gamma_{\mathbf{q}}^{n,-}(\omega))$ over a range of probing frequencies ω eventually allows to obtain the off-diagonal component of the quantum metric $g_{xy}(\mathbf{q})$ (for $\varphi = 0$) and the Berry curvature $B_{xy}(\mathbf{q})$ (for $\varphi = \pi/2$) via (S72).

The results of this simulation for $g_{xx}(\mathbf{q})$ and $B_{xy}(\mathbf{q})$ are shown in Fig. 3 in the main text. For completeness, here we show the results for the remaining components of the SQGT, $g_{yy}(\mathbf{q})$ and $g_{xy}(\mathbf{q})$ in Fig. S2.

S10C. Signatures for the distribution of the original particles

Our measurement protocol for extracting the SQGT requires measuring the excitation fraction in specific Bogoliubov (particle) modes. Focusing on the Bogoliubov-Haldane model (S122), we discuss the signatures for the quasimomentum distribution of the original particles $c_{\mathbf{q},s}$ given the presence of the single or multiple quasiparticles excitations $\hat{b}_{\mathbf{q},\pm}^\dagger$ on top of the Bogoliubov vacuum, which can then be measured e.g., via time-of-flight techniques in optical lattice platforms [91, 95, 96, 102].

We start by noting that the ground state $|\psi_0\rangle$ of the Bogoliubov-Hamiltonian (S130) is given by the vacuum state of the Bogoliubov quasiparticles fulfilling the conditions,

$$\hat{b}_{\mathbf{q},+}|\psi_0\rangle = 0, \quad \forall \mathbf{q}, \quad \text{and} \quad \hat{b}_{\mathbf{q},-}|\psi_0\rangle = 0, \quad \forall \mathbf{q} \neq 0, \quad (\text{S140})$$

where in the second equality the condensation mode has to be explicitly excluded, as the Bogoliubov theory is only concerned with excitations outside the condensate mode at $\mathbf{q} = \mathbf{0}$ in the lower band. Even though the Bogoliubov ground state contains no quasiparticles, there is still a finite amount of original particles outside the condensation mode which are (virtually) excited by interparticle interactions. Since these excitations are only driven by quantum fluctuations, this result is called quantum depletion.

We now consider the more general situation where a finite number of l quasiparticles are excited on top of the Bogoliubov vacuum at quasimomentum \mathbf{q} in band $n \in \{\pm\}$, i.e., $(b_{\mathbf{q},n}^\dagger)^l / \sqrt{l!} |\psi_0\rangle \equiv |\phi_0\rangle$. For $l = 0$ this state is again the Bogoliubov ground state (S140). Now, for the state $|\phi_0\rangle$ with l quasiparticle excitations, we compute the original particle contributions outside the condensation mode $\langle \phi_0 | \hat{a}_{\mathbf{q},s}^\dagger \hat{a}_{\mathbf{q},s'} | \phi_0 \rangle$ where $s, s' \in \{A, B\}$ is again the sublattice index (S122). To this end, we explicitly use the Bogoliubov transformations for $\mathbf{q} \neq \mathbf{0}$ (S126)-(S128) and arrive at

$$\begin{aligned} \langle \hat{a}_{\mathbf{q},s}^\dagger \hat{a}_{\mathbf{q},s} \rangle &= l |u_{s,m}^{\mathbf{q}}|^2 \delta_{m,n} + |v_{s,+}^{-\mathbf{q}}|^2 + |v_{s,-}^{-\mathbf{q}}|^2, \quad \langle \hat{a}_{-\mathbf{q},s}^\dagger \hat{a}_{-\mathbf{q},s} \rangle = l |v_{s,m}^{\mathbf{q}}|^2 \delta_{m,n} + |v_{s,+}^{\mathbf{q}}|^2 + |v_{s,-}^{\mathbf{q}}|^2, \\ \langle \hat{a}_{\mathbf{q},A}^\dagger \hat{a}_{\mathbf{q},B} \rangle &= l \left[(u_{A,m}^{\mathbf{q}})^* u_{B,m}^{\mathbf{q}} \right] \delta_{m,n} + (v_{A,+}^{-\mathbf{q}})(v_{B,+}^{-\mathbf{q}})^* + (v_{A,-}^{-\mathbf{q}})(v_{B,-}^{-\mathbf{q}})^* = \langle \hat{a}_{\mathbf{q},B}^\dagger \hat{a}_{\mathbf{q},A} \rangle^*, \\ \langle \hat{a}_{-\mathbf{q},A}^\dagger \hat{a}_{-\mathbf{q},B} \rangle &= l \left[v_{A,m}^{\mathbf{q}} (v_{B,m}^{\mathbf{q}})^* \right] \delta_{m,n} + (v_{A,+}^{\mathbf{q}})(v_{B,+}^{\mathbf{q}})^* + (v_{A,-}^{\mathbf{q}})(v_{B,-}^{\mathbf{q}})^* = \langle \hat{a}_{-\mathbf{q},B}^\dagger \hat{a}_{-\mathbf{q},A} \rangle^*. \end{aligned} \quad (\text{S141})$$

For $l = 0$ initial quasiparticle excitation present in band n in (S141), we recover the quantum depletion result. As an example, let us consider l initial quasiparticle excitations in the lower Bogoliubov energy band at \mathbf{q} , i.e., $|\phi_0\rangle = (b_{\mathbf{q},-}^\dagger)^l / \sqrt{l!} |\psi_0\rangle$. Then, for the sublattice site $s = A$, we have $l \times |u_{A,-}^{\mathbf{q}}|^2 + |v_{A,+}^{-\mathbf{q}}|^2 + |v_{A,-}^{-\mathbf{q}}|^2$ original particles moving with quasimomentum $+\mathbf{q}$ and $(l+1) \times |v_{A,-}^{-\mathbf{q}}|^2 + |v_{A,+}^{\mathbf{q}}|^2$ original particles moving with quasimomentum $-\mathbf{q}$. These results (S141) directly relate the presence of initial Bogoliubov quasiparticle excitations with the quasimomentum distribution of the real bosonic particles.

In optical lattice platforms, the momentum distribution can be observed via time-of-flight (TOF) imaging techniques. The momentum distribution $N(\mathbf{k})$ of the system after TOF is given by [102, 136]

$$N(\mathbf{k}) = |\tilde{w}(\mathbf{k})|^2 \frac{1}{N_{\text{uc}}} \sum_{\ell\ell'} \sum_{ss'} e^{-i\mathbf{k} \cdot (\mathbf{r}_{\ell s} - \mathbf{r}_{\ell' s'})} \langle \hat{a}_{\ell s}^\dagger \hat{a}_{\ell' s'} \rangle, \quad (\text{S142})$$

where $|\tilde{w}(\mathbf{k})|^2$ is the Wannier function envelope with the Fourier-transformed Wannier function $\tilde{w}(\mathbf{k})$, N_{uc} the number of unit cells, and $\hat{a}_{\ell s} = \sum_{\mathbf{q}} \langle \ell s | \mathbf{q} s \rangle \hat{a}_{\mathbf{q},s} = (N_{\text{uc}})^{-1/2} \sum_{\mathbf{q}} e^{-i\mathbf{q} \cdot \mathbf{r}_{\ell s}} \hat{a}_{\mathbf{q},s}$ the real bosonic operators in the original real space basis with the unit cell index ℓ and sublattice index s (S104). Then, for the initial state with multiple Bogoliubov quasiparticle excitations in band n (S141) the TOF momentum distribution $N(\mathbf{k})$ (S142) becomes

$$\begin{aligned} N(\mathbf{k}) &= \sum_{s,s'} \langle \hat{a}_{\mathbf{k}s}^\dagger \hat{a}_{\mathbf{k}s'} \rangle = \langle \hat{a}_{\mathbf{k}A}^\dagger \hat{a}_{\mathbf{k}A} \rangle + \langle \hat{a}_{\mathbf{k}B}^\dagger \hat{a}_{\mathbf{k}B} \rangle + \left(\langle \hat{a}_{\mathbf{k}A}^\dagger \hat{a}_{\mathbf{k}B} \rangle + \text{c.c.} \right) \\ &= \sum_{s \in \{A,B\}} \left\{ l |u_{s,m}^{\mathbf{k}}|^2 \delta_{m,n} + |v_{s,+}^{-\mathbf{k}}|^2 + |v_{s,-}^{-\mathbf{k}}|^2 \right\} \\ &\quad + \left[\left(l \left[(u_{A,m}^{\mathbf{k}})^* u_{B,m}^{\mathbf{k}} \right] \delta_{m,n} + (v_{A,+}^{-\mathbf{k}})(v_{B,+}^{-\mathbf{k}})^* + (v_{A,-}^{-\mathbf{k}})(v_{B,-}^{-\mathbf{k}})^* \right) + \text{c.c.} \right], \end{aligned} \quad (\text{S143})$$

where we neglected the overall Wannier envelope function $|\tilde{w}(\mathbf{k})|^2$ for now. Thus, equipped with the quasimomentum distribution results obtained from Bogoliubov theory (S141) and the TOF images (S143), the distribution of Bogoliubov quasiparticle excitations can be inferred to extract the SQGT.

We believe that the following experimental protocol can be used to measure the Bogoliubov excitation fraction after periodic modulation the systems' parameters to extract the SQGT as described in the main text. First, one should measure the TOF momentum distribution without performing the periodic modulation and without exciting any quasiparticles to obtain a reference measurement. Then, as a second step, we imagine exciting Bogoliubov quasiparticles at a desired quasimomentum \mathbf{q} in band n on top of the Bogoliubov vacuum with the protocol described in the main text or the one of Refs. [91, 95], whose momentum distribution should then be measured via TOF. The difference of these first two measurements then provides a measure of the excited Bogoliubov quasiparticle distribution of step two. As a third step, after exciting initial Bogoliubov quasiparticles, one should perform the periodic parameter modulation protocol to induce scattering of Bogoliubov quasiparticles into the other band. Then, the momentum distribution of the system should be measured again via TOF. Eventually, the difference of this final momentum distribution and the one after step two gives us the induced Bogoliubov quasiparticle excitation fraction which can be used to extract the SQGT as described in the main text.