

Randomized Greedy Online Edge Coloring Succeeds for Dense and Randomly-Ordered Graphs

Aditi Dudeja^{*1}, Rashmika Goswami^{†2}, and Michael Saks^{‡2}

¹University of Salzburg

²Rutgers University

Abstract

Vizing’s theorem states that any graph of maximum degree Δ can be properly edge colored with at most $\Delta + 1$ colors. In the online setting, it has been a matter of interest to find an algorithm that can properly edge color any graph on n vertices with maximum degree $\Delta = \omega(\log n)$ using at most $(1 + o(1))\Delta$ colors. Here we study the naïve random greedy algorithm, which simply chooses a legal color uniformly at random for each edge upon arrival. We show that this algorithm can $(1 + \epsilon)\Delta$ -color the graph for arbitrary ϵ in two contexts: first, if the edges arrive in a uniformly random order, and second, if the edges arrive in an adversarial order but the graph is sufficiently dense, i.e., $n = O(\Delta)$. Prior to this work, the random greedy algorithm was only known to succeed in trees.

Our second result is applicable even when the adversary is *adaptive*, and therefore implies the existence of a deterministic edge coloring algorithm which $(1 + \epsilon)\Delta$ edge colors a dense graph. Prior to this, the best known deterministic algorithm for this problem was the simple greedy algorithm which utilized $2\Delta - 1$ colors.

1 Introduction

The edge coloring problems for graphs is to assign colors to the edges of a given graph so that any two edges meeting at a vertex are assigned different colors. Trivially, the number of colors needed is at least the maximum vertex degree Δ . A theorem of Vizing’s states that every graph can be properly edge colored using $\Delta + 1$ colors. Vizing’s proof is constructive, and gives an algorithm to $\Delta + 1$ color a graph with n vertices and m edges in $O(mn)$ time (see [MG92]).

Our focus is on the edge coloring problem in the online setting, which was first introduced by Bar-Noy, Motwani and Naor [BMN92]. Starting from the empty graph on vertex set V , edges of the graph are revealed one at a time, and the algorithm must irrevocably assign colors to the edges as they arrive. The simplest online edge coloring algorithms are greedy algorithms, which color each arriving edge by some previously used color whenever it is possible to do so. Since every arriving edge touches at most $2\Delta - 2$ existing edges, any greedy algorithm uses at most $2\Delta - 1$ colors. In their paper, [BMN92] showed that for $\Delta = O(\log n)$, no deterministic online algorithm can guarantee better than $2\Delta - 1$ coloring and thus greedy algorithms are optimal in this case. They also showed that for $\Delta = O(\sqrt{\log n})$, no randomized online algorithm can achieve a better than $2\Delta - 1$ coloring. Thus, most of the focus has been designing algorithms for $\Delta = \omega(\log n)$. There has been steady progress on this problem [CPW19, BGW21, SW21, KLS⁺22, BSVW23], culminating in a recent work of [BSVW24], which presents a randomized algorithm that edge colors an online graph using $\Delta + o(\Delta)$ colors.

In this paper, we investigate the *randomized greedy algorithm* which is a natural variation of the greedy algorithm. Given a set Γ of colors, the randomized greedy algorithm \mathcal{A} on input the online graph

*aditi.dudeja@salzburg.ac.at

†rg894@scarletmail.rutgers.edu

‡saks@math.rutgers.edu

G chooses the color of each arriving edge uniformly at random from the currently allowed colors for that edge, and leaves the edge uncolored if no colors are allowed. The algorithm is said to succeed if every edge is colored. The example in Figure 1 shows that the algorithm will fail if $|\Gamma| = \Delta + o(\sqrt{\Delta})$ (because the set of colors not used by the first $\Delta - 1$ edges is likely to be disjoint from the set of colors not used by the second $\Delta - 1$ edges.)




Figure 1: A simple example to illustrate that if $|\Gamma| = \Delta + o(\sqrt{\Delta})$, then \mathcal{A} likely fails.

In their paper, [BMN92] conjectured that given any online graph G with $\Delta = \omega(\log n)$ the randomized greedy algorithm \mathcal{A} succeeds in $\Delta + o(\Delta)$ edge coloring G with high probability. This algorithm is simple and natural, but subsequent researchers (e.g. [KLS⁺22, SW21, ASvS23, BMM10]) have noted that it seems difficult to analyze. Prior to the present paper, the analysis was only done for the special case of trees (see Theorem 5 below). Our first result gives an analysis of this algorithm for the case of random-order arrivals. In this setting, an adversary can pick a worst-case graph but the edges of the graph arrive in a random order. (Recall that a graph with no multiple edges is said to be *simple*.)

Theorem 1 (Informal version of Theorem 6). (**Random order case**) Let $\epsilon \in (0, 1)$ be a constant. The algorithm \mathcal{A} , when given any simple graph G of maximum degree $\Delta = \omega(\log n)$, whose edges are presented in a uniformly random order, edge colors G with $(1 + \epsilon)\Delta$ colors with high probability.

Our second result applies to the setting where the adversary is *adaptive*. This means that the adversary does not have to decide the graph *a priori*. They can instead decide the subsequent edges of graph based on the coloring of the prior edges. We call such a graph *adaptively chosen*.

Theorem 2 (Informal version of Theorem 7). (**Dense case**) Let $\epsilon \in (0, 1)$ be a constant. Suppose G is an *adaptively chosen* simple graph with maximum degree Δ and $n = O(\Delta)$. Then, \mathcal{A} succeeds in $(1 + \epsilon)\Delta$ -edge coloring G with high probability.

To clarify the setting of Theorem 2, we note that prior results (as far as we know) assumed an *oblivious adversary* that chooses the graph G and the edge-arrival order but must fix *both* of these before their online algorithm receives any input. In contrast, an *adaptive* adversary builds the input as \mathcal{A} runs and may choose each edge depending on the coloring of the graph so far.

For the case of online algorithms, [BBK⁺94] established a connection between randomized algorithms against an adaptive adversary and deterministic algorithms. Their result states that for a given online problem, if there is an α -competitive randomized algorithm against an adaptive adversary, then there exists a α -competitive deterministic algorithm. Exploiting this connection, we get the following corollary:

Corollary 3. Let $\epsilon \in (0, 1)$ be a constant. For Δ sufficiently large and $n = O(\Delta)$ there is a deterministic online algorithm that given any *adaptively chosen* simple graph with n vertices and maximum degree Δ will produce a valid coloring using $(1 + \epsilon)\Delta$ colors.

Since the above-mentioned previous algorithms for online edge-coloring were applicable against an *oblivious adversary*, they did not have any implications to the deterministic setting. As far as we know, Corollary 3 is the first result for a general class of graphs that gives a deterministic online algorithm using fewer than $2\Delta - 1$ colors.

Edge Coloring in other computational models. The edge coloring problem has been considered in numerous settings. In the standard (offline) computational model, it is NP-hard to distinguish whether a graph is Δ or $\Delta + 1$ colorable [Hol81], so attention has focused on algorithms that color in at most

$\Delta + 1$ colors. As mentioned before, Vizing's proof gives such an algorithm that runs in $O(mn)$ time. Additionally, This was improved to $O(m \cdot \min \{\Delta \cdot \log n, \sqrt{n \cdot \log n}\})$ by [Arj82, GNK⁺85] and then to $O(m \cdot \min \{\Delta \cdot \log n, \sqrt{n}\})$ by [Sin19]. The current best known bounds are $O(n^2)$ for sparse graphs (due to [Ass24]), and $O(mn^{1/3})$ for dense graphs (due to [BCC⁺24]). There has also been a considerable effort to study edge coloring in other computational models: such as the dynamic model [BM17, BCHN18, DHZ19, Chr23, BCPS24, Chr24], the distributed model [PR01, EK24, FGK17, GKMU18, BBKO22, CHL⁺20, Ber22, Chr23, Dav23], and the streaming model [BDH⁺19, BS23, CMZ23, GS23].

Remark 4 (Note about ϵ). We only consider $\epsilon < 1$, since for $\epsilon \geq 1$, we have $\geq 2\Delta$ colors, and the randomized greedy algorithm must succeed.

1.1 Our Approach

We start with describing \mathcal{A} in more detail. Fix the color set Γ . The edges of $G = (V, E)$ arrive in some order e_1, e_2, \dots and so on. At all points during the algorithm, for each vertex v , the *free set at v* is the set of colors not yet used to color an edge incident on v . The free set of v when edge e_i arrives is denoted $F_{i-1}(v)$. When edge $e_i = (u, v)$ arrives, algorithm \mathcal{A} uniformly chooses a color c from $F_{i-1}(u) \cap F_{i-1}(v)$ and colors e_i with that color. If $F_{i-1}(u) \cap F_{i-1}(v) = \emptyset$ when e_i arrives then e_i is left uncolored and the algorithm continues. In this case the algorithm is said to *fail on e_i* .

As mentioned in the introduction, this algorithm was previously analyzed for trees. We give a proof of this for completeness, since the tree case gives intuition for our proof strategy for general graphs.

Theorem 5. [FMP] Suppose the online graph G is a tree and $\Delta = \omega(\log n)$. If $|\Gamma| = \Delta + 2\sqrt{\Delta \cdot \log n}$, then \mathcal{A} succeeds with probability at least $1 - 1/n$.

Proof. We prove an upper bound on the probability that the algorithm fails to color the i th edge $e_i = (u, v)$ conditioned on not having failed previously. Let $d_{i-1}(u)$, respectively $d_{i-1}(v)$, be the degree of u and v prior to the arrival of e and let $k(u) = (1+\epsilon)\Delta - d_{i-1}(u)$ and $k(v) = (1+\epsilon)\Delta - d_{i-1}(v)$. Conditioned on the algorithm having succeeded thus far, $|F_{i-1}(u)| = k(u)$ and $|F_{i-1}(v)| = k(v)$. By symmetry $F_{i-1}(u)$ and $F_{i-1}(v)$ are uniformly random from $\binom{\Gamma}{k(u)}$, $\binom{\Gamma}{k(v)}$, respectively. Since u and v are in separate components of the tree prior to the arrival of e_i , $F_{i-1}(u)$ and $F_{i-1}(v)$ are independently sampled from Γ , even if we condition on successful coloring prior to e_i . Therefore, conditioned on successful coloring prior to e_i ,

$$\begin{aligned}
\Pr[e_i \text{ not colored}] &\leq \Pr[F_{i-1}(u) \cap F_{i-1}(v) = \emptyset] \\
&= \sum_{S \in \binom{\Gamma}{k(u)}} \Pr[S \cap F_{i-1}(v) = \emptyset \mid F_{i-1}(u) = S] \cdot \Pr[F_{i-1}(u) = S] \\
&= \sum_{S \in \binom{\Gamma}{k(u)}} \Pr[S \cap F_{i-1}(v) = \emptyset] \cdot \Pr[F_{i-1}(u) = S] && \text{(Indep. of } F_{i-1}(v), F_{i-1}(u)) \\
&\leq \sum_{S \in \binom{\Gamma}{k(u)}} \left(1 - \frac{k(u)}{|\Gamma|}\right)^{k(v)} \cdot \Pr[F_{i-1}(u) = S] \\
&\leq \exp\left(-\frac{k(u)k(v)}{|\Gamma|}\right) \\
&\leq \exp\left(-\frac{\epsilon^2 \cdot \Delta}{(1+\epsilon)}\right) && (k(u), k(v) \geq \epsilon\Delta) \\
&= O(n^{-3}) && \left(\epsilon \geq 2\sqrt{\frac{\log n}{\Delta}}\right)
\end{aligned}$$

Taking a union bound over all edges e_i , \mathcal{A} fails with probability $O(n^{-1})$. \square

Taking inspiration from the proof for trees, we aim to show that for any edge (u, v) , the sets $F_{i-1}(u)$ and $F_{i-1}(v)$, though not independent, are close to independent. It is difficult to estimate the dependencies

of the sets $F_{i-1}(u) \cap F_{i-1}(v)$ directly. Instead we aim to show that for *any* fixed subset S of colors, and for any time step i , the set $F_{i-1}(v)$ “looks random” with respect to S :

$$|S \cap F_{i-1}(v)| \approx \frac{|S||F_{i-1}(v)|}{(1 + \epsilon)\Delta}. \quad (1)$$

The right hand side is the expected size of $|S \cap F_{i-1}(v)|$ if $F_{i-1}(v)$ was a uniformly random subset of Γ of size $|F_{i-1}(v)|$. It is fairly obvious that Equation (1) cannot hold for every set S , vertex v , and time step i . (for example, $S = \Gamma - F_{i-1}(v)$). What we will show is that for each S , and for all but constantly many vertices v , Equation (1) holds for all time steps i (up to some small error).

The intuition for our proof is as follows. The set $F_i(v)$ evolves over time as colors are removed one by one. If, at each step j , the color removed was chosen uniformly from $F_{j-1}(v)$, then at any time i , the set $F_i(v)$ would be a uniform set from Γ and would likely satisfy Equation (1). In fact, since we are only interested in the *size* of $F_i(v) \cap S$ appearing to be random, and not the set itself, in order to satisfy Equation (1) with high probability, it is enough that the probability the color chosen at step j is in S is the same as it would be if the color was chosen uniformly from $F_{j-1}(v)$. More explicitly, we would like to have

$$\frac{|S \cap F_{j-1}(v) \cap F_{j-1}(w)|}{|F_{j-1}(v) \cap F_{j-1}(w)|} \approx \frac{|S \cap F_{j-1}(v)|}{|F_{j-1}(v)|}, \quad (2)$$

where the left hand side is the probability that the color selected for $e_j = (v, w_j)$ belongs to S . Note that if we set $S' = S \cap F_{j-1}(v)$ and $S'' = F_{j-1}(v)$, and in addition we knew that Equation (1) was satisfied for vertex w_j and each of the color sets S' and S'' , then the above equation would hold (with a marginal increase in error.) This idea forms the basis of an inductive proof of our main lemma Lemma 33.

As mentioned, we will show that with high probability, for each set S , the number of vertices v for which Equation (1) fails is bounded by a constant. Ideally, we would like to use this to show that for all but constantly many $e_j = (v, w_j)$ adjacent to v , Equation (2) holds. Unfortunately, it could happen that w_j is one of the vertices for which Equation (1) fails for $S' = S \cap F_{j-1}(v)$ or $S'' = F_{j-1}(v)$, and this would prevent us from reasoning as above. To get around this we note that the sets $F_j(v)$ cannot be completely independent from each other - $F_j(v)$ will differ from $F_{j+t}(v)$ by at most t colors. So for each vertex v , we will partition the time steps into a (large) constant number of *v-phases* so that each *v-phase* contains only a small fraction of the edges incident on v . Then for each *v-phase* r we will approximate $F_{j-1}(v)$ for all time steps j of that *v-phase* by the set $A_{j-1}(v)$ which is defined to be the free set of v at the beginning of the *v-phase* that contains j . Since $F_{j-1}(v)$ will not differ too much from $A_{j-1}(v)$, we will argue that if we replace $F_{j-1}(v)$ with $A_{j-1}(v)$ in Equation (2) and the new equation holds for all but constantly many $e_j = (v, w_j)$, then the original equation must also hold for all but constantly many neighbors w_j . This allows us to carry out the reasoning of the previous paragraph.

To this end, our analysis will consider a modified version of the algorithm which we denote by \mathcal{A}' . This modified version produces the same distribution over colorings as \mathcal{A} but it has the advantage that it explicitly reflects the partition of time steps into *v-phases* for each v , and the approximation of $F_{j-1}(v)$ by $A_{j-1}(v)$ described above. The modified algorithm works as follows. When edge $e_j = (u, v)$ arrives we first sample a color uniformly from $A_{j-1}(u) \cap A_{j-1}(v)$ (rather than from $F_{j-1}(u) \cap F_{j-1}(v)$). If the selected color is valid (which means that it belongs to $F_{j-1}(u) \cap F_{j-1}(v)$) then we use it, otherwise we discard it and then sample uniformly from $F_{j-1}(u) \cap F_{j-1}(v)$. We will have the following chain of comparisons: first, that the number of times \mathcal{A}' initially selects and invalid color and resamples is small, and thus the colors chosen for \mathcal{A} are close to the colors initially chosen by \mathcal{A}' . Second, the inductive assumption that Equation (2) holds for all but constantly many edges $e_j = (v, w_j)$ adjacent to v , and thus the colors initially chosen by \mathcal{A}' hit any given set S about as often as a uniform choice from $F_{j-1}(v)$ would. Finally, we show that the uniform choice from $F_{j-1}(v)$ would likely satisfy Equation (1). Proposition 26 quantifies the error added in each step of this chain in terms of martingale sums defined in Section 5.1.

2 Preliminaries

2.1 Online Coloring

Online coloring can be described as a two player game between *Builder* and *Colorer*. The game $\Phi(n, \Delta, \epsilon)$ is parameterized by the number of vertices n , degree bound Δ and $\epsilon > 0$. The game starts with the empty graph on n vertices and a color set Γ of size $\lceil (1 + \epsilon)\Delta \rceil$. The game lasts for $m = \lfloor \Delta n/2 \rfloor$ steps. In each time step, Builder selects an edge to add to the graph, subject to the restriction that all vertex degrees remain below Δ . Colorer then assigns a color to the edge from Γ so that the overall coloring remains valid. If this is impossible (i.e. any color choice will invalidate the coloring) the edge is left uncolored. Colorer wins if every edge is successfully colored. For convenience, we also allow Builder to add *null edges*; such edges are not adjacent to any vertex in the graph and may be assigned any color without affecting whether or not the coloring is valid. This has the advantage that we can fix the number of steps to m , and if Builder gets stuck (is unable to add an edge without violating the degree bound) then he can add null edges for the remaining steps.

The *state* of the game after i steps, denoted \mathcal{S}_i , consists of the list of the first i edges chosen by Builder and the coloring chosen by Colorer. A strategy for Builder is a function which given any game state \mathcal{S}_i determines the next edge to be added or terminates the game. A strategy for Colorer is a function which given the game state \mathcal{S}_i and an additional edge e assigns a color to e (or leaves e uncolored if no color can be assigned.)

We say that a color c is *free for v at step i* if among the first $i - 1$ edges, no edge that touches v is colored by c . If the i th edge chosen by Builder is (u, v) then the set of colors that can be used to color (u, v) is the intersection of the free set for v at step i and the free set for u at step i . If that set is empty then the edge is necessarily left uncolored.

We are interested in analyzing the behavior of the *randomized greedy strategy* for Colorer, denoted \mathcal{A} : for each new edge $e_i = (u, v)$, if the intersection of the free set for u and the free set for v is nonempty then choose the color for e_i uniformly at random from that set.

During the game Builder produces a graph G together with an ordering of its edges which we view as a one-to-one function $\sigma : E(G) \rightarrow [m]$. Elements in $[m]$ to which no edge in $E(G)$ is mapped are interpreted as *null edges*. We refer to (G, σ) as an edge-ordered graph. In general the edge ordered graph produced by Builder may depend on the coloring of edges chosen by Colorer. A strategy of Builder is *oblivious* if the choice of edge to be added at each step depends only on the current edge set and not on the coloring. An oblivious strategy is fully described by the pair (G, σ) where the edge selected at step i is σ_i^{-1} (and is a null edge if that is undefined). We denote this strategy by $\mathbf{obl}(G, \sigma)$. We can now give a more precise formulation of our first main result.

Theorem 6. (random order case) For any constant $\epsilon \in (0, 1)$ there are constants $N = N(\epsilon)$ (sufficiently large), $\gamma_1 = \gamma_1(\epsilon)$ and $\gamma_2 = \gamma_2(\epsilon)$ (sufficiently small) such that the following holds. Suppose that n is sufficiently large, $\Delta > N \log(n)$ and consider the edge coloring game $\Phi(n, \Delta, \epsilon)$. For any G on n vertices with maximum degree Δ , for all but at most a $2^{-\gamma_1 \Delta}$ fraction of mappings σ of $E(G)$ to $[m]$, \mathcal{A} will defeat the oblivious strategy $\mathbf{obl}(G, \sigma)$ (i.e., produce an edge coloring for G) with probability at least $1 - 2^{-\gamma_2 \Delta}$.

Our second main result applies to arbitrary adaptive strategies of Builder. (Here we use the word *adaptive* to emphasize that Builder's choices may depend on the past coloring.)

Theorem 7. (dense case) For any constant $\epsilon \in (0, 1)$ and constant $M > 1$, there is a constant $\gamma = \gamma(\epsilon, M)$ so that the following holds. For sufficiently large n , and for $\Delta > n/M$, for any (possibly adaptive) strategy for the online coloring game $\Phi(n, \Delta, \epsilon)$, \mathcal{A} wins (produces an edge coloring of the resulting graph) with probability $1 - 2^{-\gamma \Delta}$.

2.2 The Algorithm \mathcal{A}'

In order to prove the above theorems we analyze a modified strategy \mathcal{A}' that against any given Builder strategy produces exactly the same distribution over colorings as \mathcal{A} , but will be easier to analyze. For

each vertex v , the modified algorithm will keep track of a partition of the time steps into at most $b = b(\epsilon)$ contiguous v -phases. The value of $b(\epsilon)$ is specified in Section 5.2.

For each v , the v -phases are numbered from 1 to b . The partition into v -phases is represented by a *phase-partition counter* $\{\phi_i(v) : i \in \{1, \dots, t\}\}$ where $\phi_i(v)$ is the number of the v -phase that contains time step i . For each $i > 1$ we have either $\phi_i(v) = \phi_{i-1}(v)$ (if $i - 1$ and i are in the same phase) or $\phi_i(v) = 1 + \phi_{i-1}(v)$ (if i starts a new v -phase.) This function is determined online, so that $\phi_i(v)$ is determined after step $i - 1$ of the game.

The phase-partition counters that we use are defined formally in Definition 11. In the dense case (Theorem 7) for each vertex v , the v -phases are determined by the number of edges incident to v that have arrived. For $r \geq 2$, the r^{th} v -phase starts with the time step where the number of edges incident on v first exceeds $(r - 1)\Delta/b$. (Thus the number of edges incident on v in each v -phase is within 1 of Δ/b .) The phase-partition counters in this case are denoted by ϕ^D . In the random case (Theorem 6), the v -phase partition is the same for every vertex. For $r \geq 2$, the r^{th} v -phase starts with the time step where the total number of edges arrived first exceeds $(r - 1)m/b$. The phase-partition counters in this case are denoted ϕ^R .

Algorithm Description In \mathcal{A}' , Colorer maintains for each vertex v , a color set $A_i(v)$ that approximates $F_i(v)$ but remains constant during each v -phase. We call $A_i(v)$ the *palette* of v at the end of time step i . For vertex v and time i , we define $A_i(v) = \Gamma$ if $i + 1$ belongs to the first v -phase and otherwise:

$$\begin{aligned} A_i(v) &= F_{i'}(v) \text{ where } i' \text{ is the final step of the } v\text{-phase prior to the } v\text{-phase that contains time step } i + 1. \\ &= \text{the set of available (free) colors } v \text{ at the completion of } v\text{-phase } \phi_{i+1}(v) - 1. \end{aligned}$$

Note that $A_i(v) \supset F_i(v)$ for all i , and we think of $A_i(v)$ as an (over-)approximation to $F_i(v)$.

Definition 8 (Algorithm \mathcal{A}'). Start with $A_0(v) = F_0(v) = \Gamma$ for all v . When edge $e_i = (u, v)$ arrives:

- (a) Choose c uniformly at random from $A_{i-1}(u) \cap A_{i-1}(v)$. This is the *preliminary color* for e_i . If $A_{i-1}(u) \cap A_{i-1}(v) = \emptyset$, e_i is left uncolored.
- (b) Next, choose the final color for e_i :
 - i) If $c \in F_{i-1}(u) \cap F_{i-1}(v)$, color edge e_i with c .
 - ii) Otherwise, $c \notin F_{i-1}(u) \cap F_{i-1}(v)$. (We refer to this as a *collision* at e_i .) In this case, choose c' uniformly from $F_{i-1}(u) \cap F_{i-1}(v)$ for edge e_i . If $F_{i-1}(u) \cap F_{i-1}(v) = \emptyset$, e_i is left uncolored.
- (c) For all vertices w , if $\phi_i(w) < \phi_{i+1}(w)$ (i completes the current w -phase) then $A_i(w)$ is set to $F_i(w)$, otherwise $A_i(w) = A_{i-1}(w)$.

It is obvious from the algorithm that the final color selected for e is uniformly random over $F_{i-1}(u) \cap F_{i-1}(v)$, so the distribution over colorings produced by \mathcal{A}' is the same as \mathcal{A} . To prove Theorems 7 and 6 it suffices to prove the corresponding statements with \mathcal{A} replaced by \mathcal{A}' and this is what we'll do.

2.3 The probability space

Fix any (possibly adaptive) strategy for Builder and consider the coloring game against \mathcal{A}' . Recall that the state of the game \mathcal{S}_i after i steps consists of the sequence e_1, \dots, e_i of edges and the coloring of those edges. For analysis purposes we augment \mathcal{S}_i so that it includes the preliminary color that \mathcal{A}' chooses for each edge. The random choices of \mathcal{A}' determine a discrete probability space over the space of final states of the game; we view this space as a stochastic process that evolves over time.

Notation. For a step $1 \leq i \leq m$ we define the following random variables depending only on \mathcal{S}_i , the transcript after the i^{th} edge is processed:

$$e_{i+1} = \text{the } (i + 1)^{\text{th}} \text{ edge to arrive.}$$

For vertex v and time step i , we define the following random variables that are determined by \mathcal{S}_i :

$$\begin{aligned} F_i(v) &= \text{the set of free colors of } v \text{ at the end of step } i. \\ \phi_{i+1}(v) &= \text{the } v\text{-phase for step } i+1. \\ A_i(v) &= \text{the palette of free colors of } v \text{ at the end of step } i. \end{aligned}$$

We will track the above variables as the algorithm progresses, and they will form the basis for the martingale difference sequences we analyze in Section 5. Each of these variables is subscripted by a step i , at which time they are fixed. This is what allows us to apply our concentration lemmas and make probabilistic claims about such variables.

In contrast, the variables defined below will be superscripted by a phase number r , if at all, and are defined only after the last edge in phase r of v arrives and is colored (or fails to be colored.) However, since the choice of edges is not fixed ahead of time, we do not know a priori the step i at which they will be defined. Thus, we must be careful when dealing with such quantities. In the rest of the paper, outside of Section 5, we will assume the algorithm has concluded, and make deterministic statements about the quantities defined below, conditioned on the results from Section 5.

For a vertex v the following set depends on the final state \mathcal{S}_m :

$$T(v) = \text{the set of arrival times of edges adjacent to } v.$$

Also for vertex v and v -phase r , we define the following sets depending on \mathcal{S}_m :

$$\begin{aligned} A^r(v) &= \text{the set of free colors of } v \text{ at the end of the } r^{\text{th}} \text{ } v\text{-phase.} \\ U^r(v) &= A^{r-1}(v) \setminus A^r(v) \\ &= \text{the set of colors used to color edges incident on } v \text{ during the } r^{\text{th}} \text{ } v\text{-phase.} \\ T^r(v) &= \{i \in T(v) : \phi_i(v) = r\} \\ &= \text{the set of arrival times of edges incident on } v \text{ during the } r^{\text{th}} \text{ } v\text{-phase.} \\ T^{\leq r}(v) &= \cup_{r' \leq r} T^{r'}(v). \\ \mathbf{last}^r(v) &= \max \{i : i \in T^r(v)\} \\ &= \text{the time at which the last edge of } v\text{-phase } r \text{ that is adjacent to } v \text{ arrives.} \end{aligned}$$

We remark that by this notation, for each time i , $A_{i-1}(v) = A^{\phi_i(v)-1}(v)$.

Finally, we define the following quantity, called the *error of vertex v with respect to S after phase r* :

$$\delta^r(v, S) := \frac{|A^r(v) \cap S|}{|A^r(v)|} - \frac{|S|}{(1 + \epsilon)\Delta}. \quad (3)$$

In our inductive arguments, we will be proving statements about quantities of the form $Q^r(v)$ based on the relative ordering of the times $\mathbf{last}^r(v)$. Thus, the times $\mathbf{last}^r(v)$ are important to note down. More formally, we define the following partial order on vertex-phase pairs $(v, r) \in V \times [b]$.

Definition 9 (Vertex-Phase and Edge Arrival Ordering, \prec). Given pairs $(v, r), (u, s) \in V(G) \times [b]$, we say that $(v, r) \prec (u, s)$ if $\mathbf{last}^r(v) < \mathbf{last}^s(u)$.

We make one final note about the variables defined in this section. Above we have defined two families of variables: those indexed by times steps i that are defined at a known time step i , and those that are not defined until the algorithms concludes. Note that although the sets in the second family depend on \mathcal{S}_m , the indicator variable for a particular index i belonging to one of these sets only depends on \mathcal{S}_{i-1} , since the identity of e_i and $\phi_i(v)$ are determined at that point. For instance, $T^r(v)$ depends on \mathcal{S}_m , but the indicator $\mathbb{1}_{\{i \in T^r(v)\}}$ only depends on \mathcal{S}_{i-1} . Thus, indicator variables of this form will also belong to the first family of variables and will be used in Section 5.

2.4 Proof Idea

As mentioned above, for each set $S \subseteq \Gamma$ and vertex $v \in V$, we would like to track $|F_i(v) \cap S|$ as \mathcal{A} progresses. We will do this indirectly by tracking $\frac{|A^r(v) \cap S|}{|A^r(v)|}$ with the goal of bounding $\delta^r(v, S)$. Observe that $\delta^0(v, S) = 0$, so our goal will be to bound how much $\delta^r(v, S)$ increases in each phase. Note that if $U^r(v)$ was chosen uniformly from $A^{r-1}(v)$, we would expect that

$$|U^r(v) \cap S| \approx |U^r(v)| \cdot \frac{|A^{r-1}(v) \cap S|}{|A^{r-1}(v)|}, \quad (4)$$

which would imply that the error with respect to any set S would not increase too much when the phase of v changes:

$$\begin{aligned} \delta^{r-1}(v, S) - \delta^r(v, S) &= \frac{|A^{r-1}(v) \cap S|}{|A^{r-1}(v)|} - \frac{|A^r(v) \cap S|}{|A^r(v)|} \\ &= \frac{|A^{r-1}(v) \cap S|}{|A^{r-1}(v)|} - \frac{|A^{r-1}(v) \cap S| - |U^r(v) \cap S|}{|A^r(v)|} \\ &= \frac{1}{|A^r(v)|} \left(|U^r(v) \cap S| + |A^{r-1}(v) \cap S| \left(\frac{|A^r(v)|}{|A^{r-1}(v)|} - 1 \right) \right) \\ &= \frac{1}{|A^r(v)|} \left(|U^r(v) \cap S| + |A^{r-1}(v) \cap S| \cdot \frac{|A^r(v)| - |A^{r-1}(v)|}{|A^{r-1}(v)|} \right) \\ &= \frac{1}{|A^r(v)|} \left(|U^r(v) \cap S| - |A^{r-1}(v) \cap S| \cdot \frac{|U^r(v)|}{|A^{r-1}(v)|} \right). \end{aligned} \quad (5)$$

$$(6)$$

Our goal is to show that Equation (4) is never too far from the truth, and therefore $\delta^r(v, S)$ does not grow too large in any given phase. As we previously noted, this cannot hold for every set S and every vertex v , but we can show that there exists a constant C depending on ϵ such that with high probability, for each set S , for all but at most C vertices v and all phases r , we have

$$|\delta^r(v, S)| \leq \frac{\epsilon^3 \Delta}{10|A^r(v)|}. \quad (7)$$

Proposition 26 allows us to bound the amount $\delta^r(v, S)$ grows during a phase of v in terms of three main sources of error: the error from the collisions at each phase, the error inherent to a locally independent algorithm, and the error in the palettes of the neighbors of v for that phase.

To that end, for all $e_i = (u, v)$, we define the indicator variables $X_i(S)$ to track whether the preliminary color chosen for e_i from $A_{i-1}(u) \cap A_{i-1}(v)$ hits S , $Y_i(S)$ to track whether the final color chosen for e_i from $A_{i-1}(u) \cap A_{i-1}(v)$ hits S , and the collision indicator variables Z_i to track whether the preliminary color chosen for e_i needs to be resampled. Furthermore, we let

$$p_i(S) := \frac{|A_{i-1}(u) \cap A_{i-1}(v) \cap S|}{|A_{i-1}(u) \cap A_{i-1}(v)|} \quad (8)$$

be the probability that $X_i(S) = 1$, conditioned on the partial coloring of edges before e_i arrives, and let

$$D_i(S) = X_i(S) - p_i(S). \quad (9)$$

Note that,

$$\left| |U^r(v) \cap S| - \sum_{j \in T^r(v)} X_j(S) \right| = \left| \sum_{j \in T^r(v)} Y_j(S) - X_j(S) \right| \leq \sum_{j \in T^r(v)} Z_j \quad (10)$$

since the colors used in the preliminary coloring and the final coloring differ only in the edges which experience collisions. This produces the first source of error. Similarly,

$$\left| \sum_{j \in T^r(v)} X_j(S) - \sum_{j \in T^r(v)} p_j(S) \right| = \left| \sum_{j \in T^r(v)} D_j(S) \right| \quad (11)$$

models the inherent error of the local algorithm on phase r of v . Therefore, if we can show that for most $i \in T^r(v)$,

$$p_i(S) \approx \frac{|A^{r-1}(v) \cap S|}{|A^{r-1}(v)|} \quad (12)$$

and bound the quantities on the right hand sides of Equation (10) and Equation (11), we will have

$$|U^{r+1}(v) \cap S| \approx |U^{r+1}(v)| \cdot \frac{|A^r(v) \cap S|}{|A^r(v)|}, \quad (13)$$

as desired. Our main tool for bounding the sums above will be martingale concentration inequalities.

2.5 Technical parameters

For easy reference we collect the parameters that are used in the analysis.

Definition 10 (Technical parameters).

- $\epsilon \in (0, 1)$ is the parameter appearing in the statements of Theorem 7 and Theorem 6.
- n always represents the number of vertices in the graph.
- $m = \lfloor n\Delta/2 \rfloor$ is the total number of time steps.

In the dense case, there is a *density parameter* $M > 1$ that is an upper bound on n/Δ . For notational convenience we will say that $M = 0$ in the random-order case.

There are several parameters given below that arise in the analysis. All of the parameters depend on ϵ and on M . As indicated above, the value $M = 0$ is used to refer to the random case.

- $\zeta = \zeta(\epsilon, M)$ is an *error-control parameter*. In the random-order case, $\zeta(\epsilon, 0) = e^{-20/\epsilon^2} \frac{\epsilon^3}{10}$. In the dense case, for $M > 1$, $\zeta(\epsilon, M) = \frac{\epsilon^5}{100M} e^{-(5M/\epsilon^2)^2}$.
- $\alpha(\epsilon, M) = \zeta(\epsilon, M) = \zeta \frac{\epsilon^3}{5}$. In Section 5.2 we define certain bad events for the run of the algorithm. These bad events are that certain “error” quantities associated with the algorithm grow too large. The margin of error for these quantities is $\alpha\Delta$.
- $b = b(\epsilon, M) = \frac{40}{\alpha\epsilon^2}$. This is the number of phases in the phase-partition for each vertex.
- $C = C(\epsilon, M) = \frac{2000}{\alpha^4}$. For each subset S of colors we say that a vertex v is S -atypical (Definition 28) if (very roughly) at some point in the algorithm the fraction of free colors at v that belong to S differs significantly from, $\frac{|S|}{(1+\epsilon)\Delta}$, the overall fraction of colors that belong to S . One of the bad events is that for some color set S , the number of S -atypical vertices is at least C .

The final parameter is only relevant for the random case:

- $N = N(\epsilon) = \max(400C(\epsilon, 0), 50b(\epsilon, 0))$. The theorem for the random case requires that $\Delta = \Omega(\log n)$. The parameter N is the lower bound on $\Delta/\log(n)$ for which the result holds.

3 Properties of the phase partitions

As described above, the algorithm \mathcal{A}' makes use of the phase-counter sequences $\phi_i(v)$ for each vertex. We use a different choice of ϕ for the dense case and the random-order case. The parameters m and b are as in Definition 10.

Definition 11 (Phase Partition Counters). We use the following phase partition counter sequences:

- (a) **Phase Partition Counters** $\phi^D = \{\phi_i^D\}_{i \in [m]}$ **for the dense case:** For each vertex v , $\phi_0^D(v) = 0$, and for $i \in [m]$

$$\phi_i^D(v) = \left\lceil \frac{|T(v) \cap \{1, \dots, i\}| \cdot b}{\Delta} \right\rceil$$

Less formally, the counter is the number of edges so far incident to v times b/Δ rounded up to the nearest integer.

- (b) **Phase Partition Counters** $\phi^R = \{\phi_i^R\}_{i \in [m]}$ **for the random-order case:** For every vertex v , $\phi_i^R(v) = \lceil \frac{i \cdot b}{m} \rceil$ for $i \in [m]$.

Our proof that \mathcal{A}^l succeeds relies on two properties of the phase-partition counters, which we call *balance* and *controlled error*. Here we will state these properties and establish conditions under which ϕ^R and ϕ^D satisfy them. The properties are determined by the edge-ordered graph (G, σ) produced by Builder.

3.1 The balance property

Definition 12 (Balance). A family $\phi(v)$ of phase counter sequences is said to be *balanced* with respect to (G, σ) provided that for all v and all phases r , every v -phase contains at most $2\Delta/b$ edges incident on v .

The definition of ϕ^D immediately gives that it is balanced with respect to (G, σ) for any graph G of maximum degree Δ and ordering σ . On the other hand, ϕ^R does not satisfy balance for all (G, σ) but for any fixed graph G it satisfies the property for almost all orderings σ , as we now show. We use the following concentration bounds.

Lemma 13. [JLR00, Theorem 2.10] Let X be a hypergeometric random variable with parameters m, d and k , where $\mu = \mathbb{E}[X] = \frac{kd}{m}$, then,

$$\Pr[X \geq \mu + t] \leq \exp\left(-\frac{t^2}{2(\mu + \frac{t}{3})}\right).$$

The following lemma shows that for a random ordering, ϕ^R is almost certainly balanced.

Lemma 14 (Probability that ϕ^R is unbalanced). Let Δ be sufficiently large and suppose G is a graph on $n \leq 2^{\Delta/N}$ vertices, where N is given in Definition 10, and m edges, where some of the edges may be null edges. Then the fraction of orderings σ of edges of G such that ϕ^R is not balanced with respect to (G, σ) is at most $\exp(-\frac{\Delta}{20b})$.

Proof. Consider σ chosen uniformly at random from all orderings. Let E^r be the (multi)-set of edges in phase r (which is the same for all v by the definition of ϕ^R). We say that E^r is a multi-set because many of the edges can be multi-edges. Let $E^r(v)$ be the set of edges in E^r that are incident to v . Our goal is to show that for all $v \in V$ and $r \in [b]$, $|E^r(v)| \leq \frac{2\Delta}{b}$ with high probability. E^r is a uniformly random subset of the edges of size $k_r \in \{\lfloor \frac{m}{b} \rfloor, \lceil \frac{m}{b} \rceil\}$ and therefore $|E^r(v)|$ is a hypergeometric random variable with expectation $\deg(v) \cdot \frac{k_r}{m} \leq \frac{\Delta}{b} \cdot (1 + \frac{b}{m}) \leq \frac{\Delta}{b} \cdot \frac{3}{2}$, for Δ sufficiently large. Thus, by Lemma 13, we have

$$\begin{aligned} \Pr\left[|E^r(v)| \geq \frac{2\Delta}{b}\right] &\leq \exp\left(-\frac{(\Delta/b)^2/4}{3(\Delta/b) + (\Delta/b)/3}\right) \\ &\leq \exp\left(-\frac{3\Delta}{40b}\right). \end{aligned}$$

Taking a union bound over all choices of $v \in V$ and $r \in [b]$ gives us that the probability that ϕ^R is not balanced is at most

$$b \cdot 2^{\frac{\Delta}{N}} \cdot \exp\left(-\frac{3\Delta}{40b}\right) \leq \exp\left(-\frac{3\Delta}{40b} + \frac{\Delta}{50b} + \log(b)\right) \leq \exp\left(-\frac{\Delta}{20b}\right)$$

by Definition 10 for Δ sufficiently large. □

3.2 The controlled error property

The second, and more significant property that we want from our phase counter sequences is *controlled error*. As indicated in the overview, a key part of the proof is to establish for each color set S , an upper bound on $\delta^r(v, S)$ that holds for all but constantly many vertices. The upper bound will be expressed in terms of the error function $\widehat{\epsilon}^r(v)$, which we now define.

The following notation will be useful. For edge e incident on v , $e - v$ denotes the other vertex of e . Also if $j \in T(v)$ (so that $v \in e_j$) then $s_j(v) = \phi_j(e_j - v)$, the $(e_j - v)$ -phase that contains step j . Also it is useful to recall the partial order on vertex-phase pairs from Definition 9 : $(u, s) \prec (v, r)$ if the last edge in u -phase s comes before the last edge of v -phase r .

Definition 15 (Error Bounds). Let $\zeta > 0$. Given a phase-partition counter ϕ for the edge-ordered graph (G, σ) , the error function $\widehat{\epsilon} = \widehat{\epsilon}_\zeta$ is defined on vertex-phase pairs (v, r) inductively as follows:

$$\begin{aligned}\widehat{\epsilon}^0(v) &= 0 \\ \widehat{\epsilon}^r(v) &= \zeta + \frac{5}{\Delta\epsilon^2} \sum_{j \in T^{\leq r}(v)} \widehat{\epsilon}^{s_j(v)-1}(e_j - v).\end{aligned}$$

The recurrence is well-defined because the value for (v, r) only depends on the value for vertex-phase pairs $(u, s) \prec (v, r)$. To see this note that the summation is over time-steps $j \in T^{\leq r}(v)$. The $(e_j - v)$ -phase of j is $s_j(v)$, so u -phase $s_j(v) - 1$ ends before j which is before the end of the v -phase r .

Definition 16. For $\zeta > 0$, a phase partition counter $\{\phi_i\}_{i=1}^m$ has ζ -controlled error with respect to the ordered graph (G, σ) provided that for all vertices v phases r :

$$\widehat{\epsilon}_\zeta^r(v) \leq \frac{\epsilon^3}{10}.$$

In what follows we show that for any (G, σ) , (1) if $n \leq \Delta M$ (dense case), the phase partition counter ϕ^D has $\zeta(\epsilon, M)$ -controlled error with respect to (G, σ) , and (2) if $\Delta \geq N(\epsilon, 0)\Delta$, ϕ^R has $\zeta(\epsilon, 0)$ -controlled error with respect to any (G, σ) for which ϕ^R is balanced. We start by formulating a bound on $\widehat{\epsilon}$ by unwinding the recursive definition.

Definition 17 (Valid Paths from v). Let ϕ be a phase partition function. Let $\mathcal{P}^r(v)$ be the set of paths $(x_0, x_1, \dots, x_t) = (e_{i_1}, e_{i_2}, \dots, e_{i_t})$ such that, $x_0 = v$, $i_1 \in T^{\leq r}(v)$, and for all $1 \leq k < t$, $\phi_{i_{k+1}}(x_k) < \phi_{i_k}(x_k)$. That is, $e_{i_{k+1}}$ arrives in an earlier phase of x_k than edge e_{i_k} .

Proposition 18. For any $\zeta > 0$ and phase-vertex pair (v, r) ,

$$\widehat{\epsilon}_\zeta^r(v) \leq \zeta \sum_{P \in \mathcal{P}^r(v)} \left(\frac{5}{\Delta\epsilon^2} \right)^{l(P)}$$

where $l(P)$ is the length of the path P .

Proof. The proof is by induction with respect to the vertex-phase partial order. The base case where the phase number is 0 is trivial since $\widehat{\epsilon}^0(v) = 0$ for all v . As noted earlier, for any $j \in T^{\leq r}(v)$, $(e_j - v, s_j(v) - 1) \prec (v, r)$. Furthermore, for $i_0 \in T^{\leq r}(v)$ and for any valid path $(e_{i_1}, \dots, e_{i_t}) \in \mathcal{P}^{s_{i_0}(v)-1}(e_{i_0} - v)$ with $\phi_{i_0}(x_1) > \phi_{i_1}(x_1)$, the path $(e_{i_0}, e_{i_1}, \dots, e_{i_t}) \in \mathcal{P}^r(v)$. Thus, we have

$$\begin{aligned}\widehat{\epsilon}^r(v) &= \zeta + \frac{5}{\Delta\epsilon^2} \sum_{j \in T^{\leq r}(v)} \widehat{\epsilon}^{s_j(v)-1}(e_j - v) \\ &\leq \zeta + \zeta \cdot \frac{5}{\Delta\epsilon^2} \sum_{j \in T^{\leq r}(v)} \sum_{P \in \mathcal{P}^{s_j(v)-1}(e_j - v)} \left(\frac{5}{\Delta\epsilon^2} \right)^{l(P)} \\ &\leq \zeta \sum_{P \in \mathcal{P}^r(v)} \left(\frac{5}{\Delta\epsilon^2} \right)^{l(P)}\end{aligned}$$

where we include the path of length 0 starting at v . □

We now show that the phase counter functions ϕ^R and ϕ^D are ζ -error controlled under suitable conditions.

Proposition 19 (Bound on $\widehat{\epsilon}$: Dense Case). For any $\epsilon > 0$ and $M > 1$, let $\zeta = \zeta(\epsilon, M)$. Then for any graph G with degree Δ (sufficiently large) and $n \leq M\Delta$ and any edge ordering σ the phase partition function ϕ^D has $\zeta(\epsilon, M)$ -controlled error.

Proof. We first show that there are at most $\frac{n^t}{(\lfloor t/2 \rfloor)!}$ valid paths of length t starting from v . Note that it is enough to show this for t even since if there are at most r valid paths of length t , then there are at most rn valid paths of length $t+1$. Let $P = (e_{i_1}, \dots, e_{i_t})$ be any valid path where $v \in e_{i_1}$. Note that the edges in even positions $e_{i_2}, e_{i_4}, \dots, e_{i_t}$ determine the path. Furthermore, these edges can only be placed in reverse arrival order for the path to be valid, so choosing the set of edges determines the path. Therefore, there can be at most $\binom{n^2}{\lfloor t/2 \rfloor} \leq \frac{n^t}{\lfloor t/2 \rfloor!}$ such paths. Thus, if $n \leq M\Delta$, then, by Proposition 18,

$$\begin{aligned} \widehat{\epsilon}^r(v) &\leq \zeta \sum_{t=0}^{\infty} \left(\frac{5}{\Delta\epsilon^2} \right)^t \left(\frac{n^t}{(\lfloor t/2 \rfloor)!} \right) \\ &\leq \zeta \sum_{t'=0}^{\infty} \frac{2(5M/\epsilon^2)^{2t'+1}}{t'!} \\ &\leq \frac{10M}{\epsilon^2} \cdot \zeta \sum_{t'=0}^{\infty} \frac{((5M/\epsilon^2)^2)^{t'}}{t'!} \\ &\leq \frac{10M}{\epsilon^2} \cdot \zeta e^{(5M/\epsilon^2)^2} \leq \frac{\epsilon^3}{10} \end{aligned}$$

by our choice of $\zeta = \zeta(\epsilon, M)$ from Definition 10. □

We remark that there is a trivial bound of $\binom{m}{t}$ on the number of paths. The more careful bound of $\frac{n^t}{\lfloor t/2 \rfloor!}$ is crucial in the above analysis.

Proposition 20 (Bound on $\widehat{\epsilon}$: Random Order Setting). Let Δ be sufficiently large and suppose G is a graph on $n \leq 2^{\Delta/N}$ vertices, where N is given in Definition 10. Let $\epsilon > 0$ and suppose that ϕ^R is balanced with respect to (G, σ) . Then the phase counter function ϕ^R is ζ -error controlled provided that $\zeta \leq \zeta(\epsilon, 0)$.

Proof. Recall that under ϕ^R , the phase counters $\phi_i^R(v)$ for all $v \in V$ are updated in lockstep. Consequently, for any edge $e_j = (u, v)$, $\phi_j^R(u) = \phi_j^R(v)$. As before, we bound $\widehat{\epsilon}^r(v)$ by bounding $|\mathcal{P}^r(v)|$ in Proposition 18. Consider a valid path $P \in \mathcal{P}^r(v)$, where $P = (e_{i_1}, \dots, e_{i_t})$. Let (x_0, x_1, \dots, x_t) be the sequence of vertices such that $e_{i_j} = (x_{j-1}, x_j)$. Recall P has the property that $v = x_0$ and $e_{i_1} \in T^{\leq r}(v)$ and for all $1 \leq k < t$, we have, $\phi_{i_k}^R(x_{i_k}) > \phi_{i_{k+1}}^R(x_{i_k})$. In other words, each $e_{i_{k+1}}$ arrives in an earlier phase of x_k than e_{i_k} . Under ϕ^R the phase-partition for all vertices is the same so there is an associated unique phase, r_k associated to e_{i_k} and $r_1 > r_2 > \dots > r_t$. So, we count the number of paths by first picking r_i 's and then fixing the edges themselves. The number of ways of picking r_i 's is at most $\binom{b}{t}$. Now we show how to inductively choose e_{i_k} 's. The number of ways of choosing e_{i_1} after one has fixed r_1 , is $\frac{2\Delta}{b}$ (by the definition of balance property in Definition 12). Having fixed edge e_{i_j} and r_{j+1} the number of ways of picking $e_{i_{j+1}}$ is at most $\frac{2\Delta}{b}$. Thus, we have, by Proposition 18,

$$\begin{aligned} \widehat{\epsilon}^r(v) &\leq \zeta \sum_{t=0}^b \left(\frac{5}{\Delta\epsilon^2} \right)^t \binom{b}{t} \left(\frac{2\Delta}{b} \right)^t \\ &\leq \zeta \sum_{t=0}^b \binom{b}{t} \left(\frac{10}{b\epsilon^2} \right)^t \\ &= \zeta \left(1 + \frac{10}{b\epsilon^2} \right)^b \\ &\leq \zeta e^{\frac{10}{\epsilon^2}} \end{aligned}$$

$$\leq \frac{\epsilon^3}{10}$$

where the last inequality follows from our choice of $\zeta(\epsilon, 0)$ in Definition 10 and using the fact that $\zeta \leq \zeta(\epsilon, 0)$. \square

4 Background on martingales and supermartingales

In this section we review needed definitions and facts about martingales and supermartingales including Freedman's concentration inequality. We also state a Lemma that bounds the probability that several related supermartingales all deviate significantly from their expectation, which we prove using Freedman's inequality.

As in the previous section, we consider a random process which determines a sequence $\{\mathcal{S}_i\}$ where \mathcal{S}_i represents the history of the process through time i . Suppose $\{Y_i\}$ is a sequence of random variables such that Y_i is determined by \mathcal{S}_i .

- $\{Y_i\}$ is a *martingale* with respect to $\{\mathcal{S}_i\}$ provided that

$$\mathbb{E}[Y_i \mid \mathcal{S}_{i-1}] = Y_{i-1}.$$

- $\{Y_i\}$ is a *supermartingale* with respect to $\{\mathcal{S}_i\}$ provided that

$$\mathbb{E}[Y_i \mid \mathcal{S}_{i-1}] \leq Y_{i-1}.$$

Suppose $\{D_i\}$ is also a sequence of random variables where D_i is determined by \mathcal{S}_i .

- $\{D_i\}$ is a *martingale difference sequence* with respect to $\{\mathcal{S}_i\}$ provided that

$$\mathbb{E}[D_i \mid \mathcal{S}_{i-1}] = 0.$$

- $\{D_i\}$ is a *supermartingale difference sequence* with respect to $\{\mathcal{S}_i\}$ provided that

$$\mathbb{E}[D_i \mid \mathcal{S}_{i-1}] \leq 0.$$

For any martingale (resp. supermartingale) difference sequence $\{D_i\}$, the sequence of sums $\{Y_i = \sum_{j=1}^i D_j\}$ form a martingale (resp. supermartingale). Conversely, for any martingale (resp. supermartingale) $\{Y_i\}$, the differences $\{D_i = Y_i - Y_{i-1}\}$ form a martingale (resp. supermartingale) difference sequence.

We make the following observation about properties of difference sequences that will be useful later on:

Observation 21. Let $\{D_i\}$ be a martingale difference sequence with respect to $\{\mathcal{S}_i\}$.

- (a) For any $\ell < i$, conditioning on \mathcal{S}_{i-1} fixes D_ℓ , so we have

$$\mathbb{E}[D_\ell D_i \mid \mathcal{S}_{i-1}] = D_\ell(\mathcal{S}_{i-1}) \mathbb{E}[D_i \mid \mathcal{S}_{i-1}] = 0.$$

- (b) More generally, let $\{\beta_i\}$ be a sequence of random variables where β_i is determined by \mathcal{S}_{i-1} . Then the sequence $\{\beta_i D_i\}$ is also a martingale difference sequence with respect to $\{\mathcal{S}_i\}$:

$$\mathbb{E}[\beta_i D_i \mid \mathcal{S}_{i-1}] = \beta_i(\mathcal{S}_{i-1}) \mathbb{E}[D_i \mid \mathcal{S}_{i-1}] = 0.$$

We say that $\{\beta_i D_i\}$ is *derived from* the martingale $\{D_i\}$ and refer to β_i as the *coefficient sequence* for the derived martingale. We emphasize that the coefficients here are themselves random variables.

In this paper, the difference sequences that we consider are indexed by the time steps of the process (corresponding to the edges of the graph) and we will associate a β sequence to each vertex, where the

β -sequence associated to v is nonzero only on the edges that touch v . For our results, we will need to show that certain derived supermartingales associated with particular vertices are unlikely to exceed their expected value by too much. The well-known Azuma-Hoeffding bound provides bounds of this type where the bound (on the probability of too large a value) degrades inversely with $\sum_i \lambda_i^2$ where λ_i is a bound on the i th difference D_i . Unfortunately, in our applications the bounds on the individual D_i are not sufficient to get a good bound from Azuma-Hoeffding, which does not allow us to take advantage of the situation we have where most of the D_i are 0, but we don't know which ones in advance. Instead we use a variant of Azuma-Hoeffding due to Freedman that is well-suited for analyzing processes whose evolution is partially controlled by an adversary (which for us is Builder). In Freedman's theorem one considers the auxiliary sequence $V_i = \text{Var}(D_i \mid \mathcal{S}_{i-1})$, which is the variance of D_i conditioned on \mathcal{S}_{i-1} . Note that V_i is itself a random variable and Freedman's theorem applies in situations when $\sum_i V_i$ can be bounded above with probability 1. (Actually, Freedman's theorem applies even if the bound on $\sum_i V_i$ fails with small probability, but we only need a simplified version where $\sum_i V_i$ is always bounded.)

Lemma 22. [Fre75, Theorem 4.1] Suppose $\{Y_i\}$ is a supermartingale with respect to $\{\mathcal{S}_i\}$ and its corresponding difference sequence $\{D_i\}$ satisfies $|D_i| \leq D$ for all i . Let $V_i = \text{Var}(D_i \mid \mathcal{S}_{i-1})$ and $W_i = \sum_{j \leq i} V_j$, and suppose $W_m \leq b$ with probability 1. Then

$$\Pr[Y_m \geq Y_0 + \delta] \leq \exp\left(-\frac{\delta^2}{2(D \cdot \delta/3 + b)}\right).$$

In the next section we will use Lemma 22 to bound the probability that the derived supermartingale associated to a vertex gets too large. This bound is stated in the first part of the Lemma below. Additionally, there will be times we want to show that for a set of C vertices, v_1, \dots, v_C , the derived supermartingales associated to each of those vertices cannot *all* become too large at once. In particular, we would like to show that the probability of this occurring decays exponentially in C . Note that if we were guaranteed that the edges adjacent to each of the v_k were disjoint and arrived contiguously - that is, if we could partition the difference sequence $\{D_i\}_{i=1}^m$ into C sequences $\{D_i\}_{i=1}^{m_1}, \dots, \{D_i\}_{i=m_{C-1}+1}^{m_C}$ such that the derived supermartingale associated to v_k was nonzero only on $\{D_i\}_{i=m_{k-1}+1}^{m_k}$ - then we could get this result by iteratively applying Lemma 22, since conditioned on the value of $\{D_i\}_{i=1}^{m_{k-1}}$, the sequence $\{D_i\}_{i=m_{k-1}+1}^{m_k}$ is still a difference sequence. However, in our case, the arrival times of the edges for the different vertices can be interleaved, and even intersect. Nevertheless, we manage to provide a general sufficient condition for a similar conclusion to hold.

Lemma 23. Suppose $\{D_i\}_{i=1}^t$ is a martingale difference sequence with respect to $\{\mathcal{S}_i\}$ such that $|D_i| \leq 1$ for all i and let α, a, Δ be positive real numbers where Δ is sufficiently large (depending on a and α).

- (a) Suppose $\{\beta_i\}_{i=1}^t$ is a coefficient sequence where β_i depends only on \mathcal{S}_{i-1} and such that with probability 1, $\sum_{i=1}^t |\beta_i| \leq \Delta$ and $|\beta_i| \leq a$ for all i . Then:

$$\Pr\left[\sum_i \beta_i D_i \geq \alpha \Delta\right] \leq \exp\left(-\frac{\alpha^4 \Delta}{128a}\right).$$

- (b) Suppose that for each $k \in \{1, \dots, C\}$, $\{\beta_i^k\}_{i=1}^t$ is a coefficient sequence where β_i^k depends only on \mathcal{S}_{i-1} and such that with probability 1, $\sum_{i=1}^t |\beta_i^k| \leq \Delta$ for all k . Suppose further that with probability 1, for all $i \in \{1, \dots, t\}$, $\sum_{k=1}^C |\beta_i^k| \leq a$. Then:

$$\Pr\left[\forall k, \sum_i \beta_i^k D_i \geq \alpha \Delta\right] \leq \exp\left(-\frac{C \alpha^4 \Delta}{128a}\right).$$

The key thing to note about the conclusion is that for α, Δ and a fixed, the probability upper bound shrinks exponentially with C . The first part of the Lemma is just the case $C = 1$ of the second part; we stated it separately to help the reader to digest the lemma statement, and also because the special case $C = 1$ will be applied twice in what follows.

Proof of Lemma 23. The proof is obtained by applying Lemma 22 to a single random sequence $\{Y_j\}$ that is constructed from $\{D_i\}$ and all C coefficient sequences. Let $Y_0 = 0$, and $j \in \{1, \dots, t\}$ let:

$$Y_j = \sum_{k=1}^C \left[\left(\sum_{i \leq j} \beta_i^k D_i \right)^2 - \sum_{i \leq j} (\beta_i^k)^2 \right].$$

If it is the case that for all k , $|\sum_{i=1}^t \beta_i^k D_i| \geq \alpha \Delta$ then:

$$Y_t = \sum_{k=1}^C \left[\left(\sum_{i \leq t} \beta_i^k D_i \right)^2 - \sum_{i \leq t} (\beta_i^k)^2 \right] \geq C\alpha^2 \Delta^2 - C\Delta a \geq \frac{C\alpha^2 \Delta^2}{2},$$

for Δ sufficiently large and therefore:

$$\Pr \left[\forall k, \sum_i \beta_i^k D_i \geq \alpha \Delta \right] \leq \Pr \left[Y_t \geq \frac{C\alpha^2 \Delta^2}{2} \right],$$

so it suffices to bound the probability on the right. We first show that $\{Y_j\}$ is a supermartingale. Defining $\{Z_j\}$ to be the difference sequence associated to $\{Y_j\}$, we have

$$Z_j = \sum_{k=1}^C \left[(\beta_j^k)^2 D_j^2 + 2(\beta_j^k) D_j \sum_{i < j} \beta_i^k D_i - (\beta_j^k)^2 \right].$$

To see that $\{Y_j\}$ is a supermartingale, note that by Observation 21, for any $\ell < i$,

$$\mathbb{E}[D_\ell D_i | \mathcal{S}_{i-1}] = D_\ell \mathbb{E}[D_i | \mathcal{S}_{i-1}] = 0.$$

Furthermore, since $|D_i| \leq 1$ for all i , we have $\mathbb{E}[D_i^2 | \mathcal{S}_{i-1}] \leq 1$. Therefore, for any $1 \leq j \leq t$, we have

$$\begin{aligned} \mathbb{E}[Y_j - Y_{j-1} | \mathcal{S}_{j-1}] &= \mathbb{E}[Z_j | \mathcal{S}_{j-1}] = \sum_{k=1}^C \left[(\beta_j^k)^2 \mathbb{E}[D_j^2 | \mathcal{S}_{j-1}] + 2\beta_j^k \sum_{i < j} \beta_i^k \mathbb{E}[D_j D_i | \mathcal{S}_{j-1}] - (\beta_j^k)^2 \right] \\ &= \sum_{k=1}^C (\beta_j^k)^2 (\mathbb{E}[D_j^2 | \mathcal{S}_{j-1}] - 1) \\ &\leq 0, \end{aligned}$$

which shows that $\{Y_j\}$ is indeed a supermartingale. We now will use Lemma 22 to upper bound the indicated probability. For this, we must bound the variance sums $\{W_j\}$ of $\{Y_j\}$ and the absolute values of the associated difference sequences $\{Z_i\}$. Note:

$$\begin{aligned} |Z_j| &\leq \sum_{k=1}^C \left| (\beta_j^k)^2 D_j^2 + 2\beta_j^k D_j \sum_{i < j} \beta_i^k D_i - (\beta_j^k)^2 \right| \\ &\leq \sum_{k=1}^C \left(2|\beta_j^k|^2 + 2|\beta_j^k| \sum_{i < j} |\beta_i^k| \right) && \text{(since } |D_i| \leq 1) \\ &= \sum_{k=1}^C 2|\beta_j^k| \left(\sum_{i \leq j} |\beta_i^k| \right) \\ &\leq 2\Delta \sum_{k=1}^C |\beta_j^k| && \text{(since } \sum_i |\beta_i^k| \leq \Delta) \\ &\leq 2\Delta a && \text{(since } \sum_k |\beta_i^k| \leq a) \end{aligned}$$

Thus,

$$V_j = \text{Var}(Z_j \mid \mathcal{S}_{j-1}) \leq \mathbb{E}[Z_j^2 \mid \mathcal{S}_{j-1}] \leq 2\Delta a \sum_{k=1}^C 4\Delta |\beta_j^k| = 8\Delta^2 a \sum_{k=1}^C |\beta_j^k|$$

which tells us

$$W_t \leq \sum_j V_j \leq \sum_j 8\Delta^2 a \sum_{k=1}^C |\beta_j^k| = 8\Delta^2 a \sum_{k=1}^C \sum_j |\beta_j^k| \leq 8\Delta^2 a \sum_{k=1}^C \Delta \leq 8aC\Delta^3$$

Then Lemma 22 tells us that,

$$\begin{aligned} \Pr \left[Y_t \geq \frac{C\alpha^2\Delta^2}{2} \right] &\leq \exp \left(-\frac{C^2\alpha^4\Delta^4}{4 \cdot (C\alpha^2a\Delta^3 + 16aC\Delta^3)} \right) \\ &\leq \exp \left(-\frac{C\alpha^4\Delta}{128a} \right). \end{aligned}$$

□

5 Well-Behaved Colorings

5.1 Some Martingales Difference Sequences

Our main goal in this section will be to define the martingale difference sequences we will be considering. Recall that we are viewing the progression of the algorithm as a filtered probability space with \mathcal{S}_i representing the space of partial colorings of the first i edges to arrive. We first introduce the random variables which will form the basis for the difference sequences we track throughout the course of the algorithm. All of the quantities defined below for an edge e_i will be set to 0 by default if e_i is null or we are unable to color e_i .

Definition 24 (Collision Variables). The following random variables relate to the collisions experienced by algorithm \mathcal{A}' . For a non-null edge $e_i = (u, v)$

- Z_i is defined to be 1 if e_i is in a collision and then successfully colored, and is 0 otherwise. Thus $Z_i = 1$ provided that there is a collision at e_i (i.e., the preliminary color is not valid) and $F_{i-1}(u) \cap F_{i-1}(v) \neq \emptyset$.
- $q_i = \mathbb{E}[Z_i \mid \mathcal{S}_{i-1}]$. If $F_{i-1}(u) \cap F_{i-1}(v) = \emptyset$ this is 0. Otherwise:

$$q_i := 1 - \frac{|F_{i-1}(u) \cap F_{i-1}(v)|}{|A_{i-1}(u) \cap A_{i-1}(v)|} = \frac{|(A_{i-1}(u) \cap A_{i-1}(v)) \setminus (F_{i-1}(u) \cap F_{i-1}(v))|}{|A_{i-1}(u) \cap A_{i-1}(v)|}.$$

Since the phase counter functions we use are balanced, at most $\frac{4\Delta}{b}$ colors from $A_{i-1}(u) \cap A_{i-1}(v)$ could have been used by the time we color (u, v) , which gives us:

$$q_i \leq \frac{|(A_{i-1}(u) \setminus F_{i-1}(u)) \cup (A_{i-1}(v) \setminus F_{i-1}(v))|}{|A_{i-1}(u) \cap A_{i-1}(v)|} \leq \frac{4\Delta}{b \cdot |A_{i-1}(u) \cap A_{i-1}(v)|}. \quad (14)$$

- $\{Z_i - q_i\}_{i=1}^m$ is a martingale difference sequence with respect to $\{\mathcal{S}_i\}$, since $\mathbb{E}[Z_i - q_i \mid \mathcal{S}_{i-1}] = 0$. Additionally, since $Z_i, q_i \in [0, 1]$, we have for all i ,

$$|Z_i - q_i| \leq 1.$$

The significance of the next set of variables is little more subtle. Recall that our goal is to bound

$$\delta^r(v, S) := \frac{|A^r(v) \cap S|}{|A^r(v)|} - \frac{|S|}{(1 + \epsilon)\Delta},$$

the error of vertex v with respect to color set S after its r^{th} phase. A natural way to do this would be to track how often the colors chosen for edges incident to v hit S . However, the probability that the color of

an edge (u, v) hits S is highly dependent on the palette of u , which makes it difficult to control $\delta^r(v, S)$ on its own. Instead we consider a related, and easier to control, quantity: the difference between the probability of the color of an edge (u, v) hitting S and the indicator for the event. This doesn't directly bound $\delta^r(v, S)$, but it does allow us to approximate $|A^r(v) \cap S|$ in terms of intersections of the form $|A^{r'}(v') \cap S'|$ for neighbors v' of v which - crucially - complete their phase r' before v completes its phase r . This will allow us to use an inductive argument to bound $\delta^r(v, S)$ in terms of such $\delta^{r'}(v', S')$.

Definition 25 (Difference Variables). Let $e_i = (u, v)$ be a non-null edge and $S \subseteq \Gamma$.

- $X_i(S)$ is 1 if the preliminary color for e_i belongs to S and 0 otherwise.
- $Y_i(S)$ is 1 if the final color chosen for e_i belongs to S and is 0 otherwise. Note that

$$|X_i(S) - Y_i(S)| \leq Z_i,$$

since the final color chosen for e_i differs from the preliminary color only if there is a collision.

- $p_i(S)$ is the probability that the preliminary color chosen for edge e_i is in S , conditioned on the coloring of all previous edges:

$$p_i(S) := \mathbb{E}[X_i(S) \mid \mathcal{S}_{i-1}] = \frac{|A_{i-1}(u) \cap A_{i-1}(v) \cap S|}{|A_{i-1}(u) \cap A_{i-1}(v)|}.$$

- $D_i(S) = X_i(S) - p_i(S)$. This is a martingale difference sequence with respect to \mathcal{S}_i since:

$$\mathbb{E}[D_i(S) \mid \mathcal{S}_{i-1}] = \mathbb{E}[X_i(S) - p_i(S) \mid \mathcal{S}_{i-1}] = 0,$$

- Furthermore if S_1, S_2, \dots is a sequence of color sets where S_i is determined by \mathcal{S}_{i-1} then $\{D_i(S_i)\}$ is also a martingale difference sequence, satisfying

$$|D_i(S_i)| = |X_i(S_i) - p_i(S_i)| \leq 1$$

for all i .

The following proposition relates the variables above to the error $\delta^r(v, S)$ and motivates the difference sequences we will define. For a vertex v and phase ℓ , let

$$\tilde{T}^\ell(v) = \{j : j \in T^\ell(v) \text{ and } e_j \text{ is successfully colored}\}.$$

Note that $|\tilde{T}^\ell(v)| = |U^\ell(v)|$ and recall that if e_i was not colored, then by definition $Z_i = Y_i(S) = X_i(S) = D_i(S) = p_i(S) = 0$ for all S .

Proposition 26. For any vertex v , subset of colors S , and phase r of v ,

$$|\delta^r(v, S)| \leq \frac{1}{|A^r(v)|} \sum_{i \in T^{\leq r}(v)} Z_i + \left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in T^\ell(v)} D_i(S) \right| + \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in \tilde{T}^\ell(v)} \left| p_i(S) - \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right|$$

Proof. From Equation (6),

$$\delta^\ell(v, S) - \delta^{\ell+1}(v, S) = \frac{1}{|A^{\ell+1}(v)|} \left(|U^{\ell+1}(v) \cap S| - |A^\ell(v) \cap S| \cdot \frac{|U^{\ell+1}(v)|}{|A^\ell(v)|} \right).$$

By definition $\delta^0(v, S) = 0$ and so:

$$\begin{aligned} |\delta^r(v, S)| &= \left| \sum_{\ell=1}^r \delta^{\ell-1}(v, S) - \delta^\ell(v, S) \right| \\ &= \left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \left(|U^\ell(v) \cap S| - |U^{\ell-1}(v)| \cdot \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right) \right| \end{aligned}$$

$$\begin{aligned}
&= \left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \left(\sum_{i \in T^\ell(v)} Y_i(S) - |U^\ell(v)| \cdot \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right) \right| \\
&= \left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \left(\sum_{i \in T^\ell(v)} (Y_i(S) - X_i(S)) + \sum_{i \in T^\ell(v)} X_i(S) - |U^\ell(v)| \cdot \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right) \right| \\
&\leq \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in T^\ell(v)} |Y_i(S) - X_i(S)| + \left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \left(\sum_{i \in T^\ell(v)} X_i(S) - |U^\ell(v)| \cdot \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right) \right| \\
&\leq \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in T^\ell(v)} Z_i + \left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \left(\sum_{i \in T^\ell(v)} (X_i(S) - p_i(S)) + \sum_{i \in T^\ell(v)} p_i(S) - |U^\ell(v)| \cdot \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right) \right| \\
&= \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in T^\ell(v)} Z_i + \left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \left(\sum_{i \in T^\ell(v)} D_i(S) + \sum_{i \in \tilde{T}^\ell(v)} \left(p_i(S) - \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right) \right) \right| \\
&\leq \frac{1}{|A^r(v)|} \sum_{i \in T^{\leq r}(v)} Z_i + \left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in T^\ell(v)} D_i(S) \right| + \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in \tilde{T}^\ell(v)} \left| p_i(S) - \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right|.
\end{aligned}$$

□

5.2 Bad Events

In this section, we will define certain bad events for the run of the algorithm. These bad events are that some “error quantities” associated with the algorithm grow too large. These bad events, and their likelihood of occurring will be defined in terms of parameters in Definition 10. Note that our parameters vary depending on the random-order or dense case. In particular, in the dense case they depend on M .

We now identify three bad events, each associated with one of the three summands in Proposition 26. If none of them occur, we say that the resulting coloring is *well-behaved*. In this section we show that the coloring is very likely to be well-behaved. In the next section we show that in the two situations (an oblivious strategy that uses an arbitrary graph and random order, or an adaptively chosen dense graph) a well-behaved coloring will not have any uncolored edges.

The reader is reminded that various technical parameters are collected in Definition 10. The key parameter in this section is α .

The first type of bad event will occur if there are too many collisions at a particular vertex. This event corresponds directly to the first summand in Proposition 26.

Definition 27 (Too Many Collisions, $\mathcal{W}(v)$). Given a vertex v , the bad event $\mathcal{W}(v)$ occurs if there exists a $j \in \{1, \dots, m\}$ such that:

$$\sum_{i \in T(v), i \leq j} (Z_i - q_i) > \alpha \Delta.$$

The second type of bad event relates to the second summation in Proposition 26. As mentioned earlier, we can't hope to say that the summation is suitably small for all choices of S and v but it will be enough that for all S it is small for all but constantly many v .

Definition 28 (Too many S -atypical vertices, $\mathcal{B}(S)$). For a vertex v and color set S , we say that v is *S -atypical* if there is a v -phase $1 \leq r \leq b$ such that:

$$\left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in T^\ell(v)} D_i(S) \right| > \frac{\alpha}{\epsilon}$$

Let $B(S)$ be the set of S -atypical vertices. We say that the bad event $\mathcal{B}(S)$ occurs if $|B(S)| \geq C$. (Here α and C are as given in Definition 10.)

The final family of bad events helps track $|F_i(u) \cap F_i(v)|$ for an edge $e = (u, v)$. This will ultimately be used to show that no edge runs out of colors.

Definition 29 (Too Much Drift at a pair of vertices, $\mathcal{D}(u, v)$). Given a pair of vertices u, v , let $S_i = F_i(u) \cap F_i(v)$ be the set of colors free at both u and v at time i . Then, the bad event $\mathcal{D}(u, v)$ occurs if,

$$\left| \sum_{\substack{i \in T(u) \cup T(v) \\ j_1 \leq i \leq j_2}} D_i(S_i) \right| > \alpha \Delta$$

for any $1 \leq j_1 \leq j_2 \leq m$.

Next we will bound the probability of too many bad events occurring to show that the algorithm succeeds with high probability.

Definition 30 (Well-behaved Coloring). We say that a coloring is well-behaved if:

- (a) There are no vertices v such that $\mathcal{W}(v)$ occurs.
- (b) There are no sets S such that $\mathcal{B}(S)$ occurs.
- (c) There are no pairs of vertices u, v such that $\mathcal{D}(u, v)$ occurs.

Lemma 31. If $n \leq 2^{\frac{\Delta}{N}}$, then with probability at least $1 - \exp\left(-\frac{\alpha^4 \Delta}{1000}\right)$, the events Definition 30(a)-(c) do not occur, and consequently, the coloring is well-behaved.

We emphasize that this Lemma applies even for adaptive adversaries, and to sparse graphs, provided that $\Delta \geq N \log(n)$.

Proof. We show that the coloring is well-behaved by enumerating over each of the conditions (a)-(c), and bounding the probability they fail.

- (a) Fix a vertex v and time $1 \leq j \leq m$. Apply the first part of Lemma 23 with

$$\beta_i = \begin{cases} 1 & i \in T(v), i \leq j \\ 0 & \text{otherwise} \end{cases}.$$

Since the event that $i \in T(v)$ depends only on \mathcal{S}_{i-1} , the same holds for β_i . We have $|\beta_i| \leq 1$ and $\sum |\beta_i| \leq |T(v)| \leq \Delta$. Applying the first part of Lemma 23 we obtain:

$$\Pr \left[\sum_{i \in T(v), i \leq j} (Z_i - q_i) > \alpha \Delta \right] = \Pr \left[\sum_{i=1}^m \beta_i (Z_i - q_i) > \alpha \Delta \right] \leq \exp \left(-\frac{\alpha^4 \Delta}{128} \right).$$

Taking a union bound over at most $m \leq n \Delta \leq \Delta \cdot 2^{\frac{\Delta}{N}}$ choices for j , we see that

$$\Pr [\mathcal{W}(v) \text{ occurs}] \leq \Delta \cdot 2^{\frac{\Delta}{N}} \cdot \exp \left(-\frac{\alpha^4 \Delta}{128} \right)$$

for any vertex v . Then taking a union bound over at most $2^{\frac{\Delta}{N}}$ vertices, we get

$$\begin{aligned} \Pr [\exists v \text{ s.t. } \mathcal{W}(v) \text{ occurs}] &\leq \Delta \cdot 2^{\frac{2\Delta}{N}} \cdot \exp \left(-\frac{\alpha^4 \Delta}{128} \right) \\ &\leq \exp \left(-\frac{\alpha^4 \Delta}{128} + \frac{2\Delta}{N} + \ln \Delta \right) \\ &\leq \exp \left(-\frac{\alpha^4 \Delta}{500} \right), \end{aligned}$$

for Δ sufficiently large, using $N \geq C = \frac{2000}{\alpha^4}$ in the last line.

- (b) Fix a set S of colors and set v_1, \dots, v_C of C vertices. By definition if v_1, \dots, v_C are all S -atypical, then for each $k \in \{1, \dots, C\}$ there is a v_k -phase $r_k \in \{1, \dots, b\}$ such that:

$$\left| \sum_{\ell=1}^{r_k} \frac{1}{|A^\ell(v_k)|} \sum_{i \in T^\ell(v_k)} D_i(S) \right| > \frac{\alpha}{\epsilon}$$

We can think of this sum as having the form $\sum_{i \geq 1} \beta_i^k D_i(S)$ where

$$\beta_i^k = \begin{cases} \frac{1}{|A^{\phi_i(v_k)}(v_k)|} & \text{if } i \in T^{\leq r_k}(v_k) \\ 0 & \text{otherwise,} \end{cases}$$

and then we might hope to apply Lemma 23. However, the lemma requires that β_i^k be determined by \mathcal{S}_{i-1} and that is not the case here because $|A^{\phi_i(v)}(v)|$ depends on the number of edges incident on v through the end of v -phase $\phi_i(v)$ (and whether they are colored or not) and this is not determined by \mathcal{S}_{i-1} .

We address this by constructing a family of *fixed coefficient sequences* which is large enough that one of them agrees with the above coefficient sequence. Now for each choice of C fixed coefficient sequences (one for each vertex) we will apply Lemma 23, and then take a union bound over all such choices.

We note that all of the above coefficients are of the form $1/|A^\ell(v)|$ where $|A^\ell(v)|$ is an integer between $\epsilon\Delta$ and $(1+\epsilon)\Delta$. Thus if v_1, \dots, v_C are all S -atypical, then for each $k \in \{1, \dots, C\}$ there is a v_k -phase $1 \leq r_k \leq b$ and for each ℓ between 1 and b there is an integer $s_k^\ell \in [\epsilon\Delta, (1+\epsilon)\Delta]$ such that:

$$\left| \sum_{\ell=1}^{r_k} \frac{\Delta\epsilon}{s_k^\ell} \sum_{i \in T^\ell(v_k)} D_i(S) \right| > \alpha\Delta$$

Consider a fixed choice of r_k and $s_k^\ell : 1 \leq \ell \leq b, k \in \{1, \dots, C\}$.

For each $k \in \{1, \dots, C\}$, define the coefficient sequence β^k by

$$\beta_i^k = \begin{cases} \frac{\epsilon\Delta}{s_k^\ell} & i \in T^\ell(v_k) \text{ with } \ell \leq r_k \\ 0 & \text{otherwise} \end{cases}.$$

As before, $\{i \in T^\ell(v_k)\}$ is determined by $\phi_i(v)$, which is determined by \mathcal{S}_{i-1} , so the same holds for β_i^k . Then, since for all ℓ, k , $\left| \frac{\epsilon\Delta}{s_k^\ell} \right| \leq 1$, for all i , $\sum_k |\beta_i^k| \leq |\{v_k : i \in T(v_k)\}| \leq 2$, and for all k , $\sum_i |\beta_i^k| \leq |T(v_k)| \leq \Delta$, taking $a = 2$ in Lemma 23 gives us

$$\Pr \left[\forall k \in \{1, \dots, C\} \sum_i \beta_i^k D_i > \alpha\Delta \right] \leq \exp \left(-\frac{C\alpha^4\Delta}{256} \right).$$

This time we take a union bound over at most b^C choices of r_k for each vertex and at most $(\Delta+1)^{bC} \leq 2^{bC} \Delta^{bC}$ choices of $\{s_k^\ell\}$ to get

$$\Pr [\text{For all } k, v_k \text{ is } S\text{-atypical}] \leq b^C \cdot \Delta^{bC} \cdot 2^{bC} \cdot \exp \left(-\frac{C\alpha^4\Delta}{256} \right).$$

Taking another union bound over at most $\binom{n}{C} \leq 2^{\frac{cn}{N}}$ sets of C vertices and $2^{(1+\epsilon)\Delta}$ sets S gives us

$$\begin{aligned} \Pr [\exists S, v_1, \dots, v_C \text{ s.t. } v_k \text{ is } S\text{-atypical } \forall k] &\leq 2^{(1+\epsilon)\Delta} \cdot 2^{\frac{cn}{N} + bC} \cdot b^C \cdot \Delta^{bC} \cdot \exp \left(-\frac{C\alpha^4\Delta}{256} \right) \\ &\leq \exp \left(-\frac{C\alpha^4\Delta}{128} + 2\Delta + \frac{C\Delta}{N} + bC + bC \ln \Delta + C \ln b \right) \\ &\leq \exp(-\Delta), \end{aligned}$$

for Δ sufficiently large, since $C = \frac{2000}{\alpha^4}$ and $N \geq 400C$.

(c) Fix $u, v \in V$ and $1 \leq j_1 \leq j_2 \leq m$. Define the sequence β by:

$$\beta_i = \begin{cases} 1 & i \in T(u) \cup T(v), j_1 \leq i \leq j_2 \\ 0 & \text{otherwise} \end{cases}.$$

Then, since the event $\{i \in T(u) \cup T(v)\}$ depends only on \mathcal{S}_{i-1} , $|\beta_i| \leq 1$, and $\sum |\beta_i| \leq |T(u) \cup T(v)| \leq 2\Delta$, the first part of Lemma 23 with $a = 1$ gives us:

$$\Pr \left[\left| \sum_{\substack{i \in T(u) \cup T(v) \\ j_1 \leq i \leq j_2}} D_i(S_i) \right| > \alpha\Delta \right] \leq \exp \left(-\frac{(\alpha/2)^4 \Delta}{128} \right).$$

Taking a union bound over at most $(n\Delta)^2 \leq n^4 \leq 2^{\frac{4\Delta}{N}}$ choices for j_1, j_2 gives us

$$\Pr [\mathcal{D}(u, v) \text{ occurs}] \leq 2^{\frac{4\Delta}{N}} \cdot \exp \left(-\frac{\alpha^4 \Delta}{512} \right).$$

Taking another union bound over at most $n^2 \leq 2^{\frac{2\Delta}{N}}$ vertex pairs gives us

$$\begin{aligned} \Pr [\exists u, v \text{ s.t. } \mathcal{D}(u, v) \text{ occurs}] &\leq 2^{\frac{6\Delta}{N}} \cdot \exp \left(-\frac{\alpha^4 \Delta}{512} \right) \\ &\leq \exp \left(-\frac{\alpha^4 \Delta}{512} + \frac{4\Delta}{N} \right) \\ &\leq \exp \left(-\frac{\alpha^2 \Delta}{800} \right) \end{aligned}$$

for Δ sufficiently large, where in the last line we used $N \geq 400C = \frac{800000}{\alpha^4}$.

Thus, for Δ sufficiently large, our total probability of a bad event occurring is at most

$$\exp \left(-\frac{\alpha^2 \Delta}{500} \right) + \exp(-\Delta) + \exp \left(-\frac{\alpha^4 \Delta}{800} \right) \leq \exp \left(-\frac{\alpha^4 \Delta}{1000} \right).$$

□

6 Main Lemma

We begin with a brief summary of what we've shown so far and what remains to be shown. For the dense case with a fixed adaptive adversary, we are given a constant M such that $n \leq \frac{\Delta}{M}$. We run the game \mathcal{A}' using the phase-counter ϕ^D , which, by Proposition 19 is $\zeta(\epsilon, M)$ -error controlled. Since $n \leq \frac{\Delta}{M} \leq 2^{\Delta/N}$ (for Δ sufficiently large), Lemma 31 implies that the coloring produced by \mathcal{A}' is well-behaved with high probability.

Similarly, if $n \leq 2^{\frac{\Delta}{N}}$, then by Lemma 14, with high probability, for uniformly chosen ordering σ , ϕ^R is balanced with respect to (G, σ) . Conditioned on ϕ^R being balanced with respect to (G, σ) , guarantees us that by Proposition 20, ϕ^R is $\zeta(\epsilon, 0)$ -controlled. Finally, in this case also Lemma 31 tells us that the coloring produced by \mathcal{A}' is well-behaved with high probability.

In this final section, we will show that if the coloring produced by \mathcal{A}' is well-behaved and ϕ is a phase partition counter with ζ -controlled error, then \mathcal{A}' must have successfully produced a proper coloring of G . We do this by inductively showing that vertices are *good* according to the following definition. In this definition, and all following definitions in this section, we will assume that we are given (G, σ) with its corresponding phase counter ϕ and parameter ζ as defined above, we will denote $\hat{\epsilon}_\zeta$ simply as $\hat{\epsilon}$.

Definition 32 (Good Vertices). A vertex v is *good* for S during its r^{th} phase if

$$|\delta^{r-1}(v, S)| \leq \frac{\hat{\epsilon}^{r-1}(v) \cdot \Delta}{|A^{r-1}(v)|}.$$

Note that in this definition we say v is good with respect to S during phase r rather than $r - 1$, because the palette for v used during phase r is $A^{r-1}(v)$.

Lemma 33 (Main Lemma). Let $\zeta > 0$, let phase partition counter $\{\phi_i\}_{i=1}^m$ be balanced and have ζ -controlled error with respect to the ordered graph (G, σ) . If the coloring is well-behaved and for all vertex phase pairs (v, r) , for all color sets S , if v is an S -typical vertex, then v is good for S during its r^{th} phase.

We will prove this lemma by induction on the pairs (v, r) according to the order \prec and by bounding each of the three summands in Proposition 26. The next two propositions relate these terms to the error terms.

Proposition 34. For any set $S \subseteq \Gamma$, and any i with $e_i = (u, v)$, $\phi_i(u) = s$, and $\phi_i(v) = r$, the preliminary color set $A_{i-1}(u) \cap A_{i-1}(v) = A^{s-1}(u) \cap A^{r-1}(v)$, satisfies

$$|A^{s-1}(u) \cap A^{r-1}(v)| \geq \frac{\epsilon^2 \Delta}{1 + \epsilon} - |\delta^{s-1}(u, A^{r-1}(v))| |A^{s-1}(u)|.$$

Proof. By the definition of $\delta^{s-1}(u)$ in (3) and the fact that $|A^{r-1}(v)|$ is always at least $\epsilon \Delta$,

$$\begin{aligned} |A^{s-1}(u) \cap A^{r-1}(v)| &= \frac{|A^{s-1}(u)| |A^{r-1}(v)|}{(1 + \epsilon) \Delta} + \delta^{s-1}(u, A^{r-1}(v)) \cdot |A^{s-1}(u)| \\ &\geq \frac{\epsilon^2 \Delta}{1 + \epsilon} - |\delta^{s-1}(u, A^{r-1}(v))| |A^{s-1}(u)|, \end{aligned}$$

□

Note that this bounds the preliminary colors available to $e_i = (u, v)$ in terms of $\delta^{s-1}(u, S)$. Next we establish bounds on q_i and $p_i(S)$ if we know that u is good with respect to $A^{r-1}(v)$ and $A^{r-1}(v) \cap S$ during its phase s .

Proposition 35. For $\zeta > 0$, let phase partition counter $\{\phi_i\}_{i=1}^m$ be balanced and have ζ -controlled error with respect to the ordered graph (G, σ) . Let $e_i = (u, v)$, $\phi_i(u) = s$, and $\phi_i(v) = r$. For any set S , if u is good for both $A^{r-1}(v)$ and $A^{r-1}(v) \cap S$ during its s^{th} phase, then:

(a) The number of colors available to edge e_i is not too low, that is,

$$|A^{s-1}(u) \cap A^{r-1}(v)| \geq \frac{2\epsilon^2 \Delta}{5}$$

(b) The probability of a collision is low:

$$q_i \leq \frac{\alpha}{4}$$

(c) The probability that the preliminary color chosen for edge e_i hits S is close to what we would expect if the color was chosen randomly from $A^{r-1}(v)$:

$$\left| p_i(S) - \frac{|A^{r-1}(v) \cap S|}{|A^{r-1}(v)|} \right| \leq \frac{5\widehat{\epsilon}^{s-1}(u)}{\epsilon^2}.$$

Proof. For the first part:

$$\begin{aligned} |A^{s-1}(u) \cap A^{r-1}(v)| &\geq \frac{\epsilon^2 \Delta}{1 + \epsilon} - |\delta^{s-1}(u, A^{r-1}(v))| |A^{s-1}(u)| && \text{(From Proposition 34)} \\ &\geq \frac{\epsilon^2 \Delta}{1 + \epsilon} - \widehat{\epsilon}^{s-1}(u) \Delta && \text{(Since } u \text{ is good for } A^{r-1}(v) \text{ during phase } s) \\ &\geq \frac{\epsilon^2 \Delta}{1 + \epsilon} - \frac{\epsilon^3 \Delta}{10} && \text{(By Definition 16 since } \{\phi_i\}_{i=1}^m \text{ is error-controlled)} \\ &\geq \frac{2\epsilon^2 \Delta}{5} && \text{(Since } \epsilon < 1.) \end{aligned}$$

Using this, (14), and the definition of b from Definition 10:

$$q_i \leq \frac{4\Delta}{b \cdot |A^{s-1}(u) \cap A^{r-1}(v)|} \leq \frac{10}{b\epsilon^2} \leq \frac{\alpha}{4},$$

For the last part, again by (3), for any color set T :

$$|A^{s-1}(u) \cap T| = \delta^{s-1}(u, T) \cdot |A^{s-1}(u)| + \frac{|A^{s-1}(u)||T|}{(1+\epsilon)\Delta},$$

so

$$\begin{aligned} p_i(S) - \frac{|A^{r-1}(v) \cap S|}{|A^{r-1}(v)|} &= \frac{|A^{s-1}(u) \cap A^{r-1}(v) \cap S|}{|A^{s-1}(u) \cap A^{r-1}(v)|} - \frac{|A^{r-1}(v) \cap S|}{|A^{r-1}(v)|} \\ &= \frac{|A^{s-1}(u) \cap A^{r-1}(v) \cap S| |A^{r-1}(v)| - |A^{r-1}(v) \cap S| |A^{s-1}(u) \cap A^{r-1}(v)|}{|A^{s-1}(u) \cap A^{r-1}(v)| |A^{r-1}(v)|} \\ &= \frac{|A^{s-1}(u)| \left[\left(\frac{|A^{r-1}(v) \cap S|}{(1+\epsilon)\Delta} + \delta^{s-1}(u, A^{r-1}(v) \cap S) \right) |A^{r-1}(v)| - |A^{r-1}(v) \cap S| \left(\frac{|A^{s-1}(u)|}{(1+\epsilon)\Delta} + \delta^{s-1}(u, A^{r-1}(v)) \right) \right]}{|A^{s-1}(u) \cap A^{r-1}(v)| |A^{r-1}(v)|} \\ &= \frac{|A^{s-1}(u)| (\delta^{s-1}(u, A^{r-1}(v) \cap S) |A^{r-1}(v)| - |A^{r-1}(v) \cap S| \delta^{s-1}(u, A^{r-1}(v)))}{|A^{s-1}(u) \cap A^{r-1}(v)| |A^{r-1}(v)|}. \end{aligned}$$

Taking the absolute value, we get,

$$\begin{aligned} \left| p_i(S) - \frac{|A^{r-1}(v) \cap S|}{|A^{r-1}(v)|} \right| &= \left| \frac{|A^{s-1}(u)| (\delta^{s-1}(u, A^{r-1}(v) \cap S) \cdot |A^{r-1}(v)| - |A^{r-1}(v) \cap S| \cdot \delta^{s-1}(u, A^{r-1}(v)))}{|A^{s-1}(u) \cap A^{r-1}(v)| |A^{r-1}(v)|} \right| \\ &\leq \frac{|A^{s-1}(u)| (|\delta^{s-1}(u, A^{r-1}(v) \cap S)| \cdot |A^{r-1}(v)| + |A^{r-1}(v) \cap S| \cdot |\delta^{s-1}(u, A^{r-1}(v))|)}{|A^{s-1}(u) \cap A^{r-1}(v)| \cdot |A^{r-1}(v)|} \\ &\leq \frac{|A^{s-1}(u)| (|\delta^{s-1}(u, A^{r-1}(v) \cap S)| + |\delta^{s-1}(u, A^{r-1}(v))|)}{|A^{s-1}(u) \cap A^{r-1}(v)|} \\ &\leq \frac{2\widehat{\epsilon}^{s-1}(u)\Delta}{2\epsilon^2\Delta/5} = \frac{5\widehat{\epsilon}^{s-1}(u)}{\epsilon^2}, \end{aligned}$$

where the final inequality uses the first part and the assumption that u is good for $A^{r-1}(v)$ and $A^{r-1}(v) \cap S$ during phase r . \square

We now turn to the proof of the main lemma.

Proof of Main Lemma. Assume the coloring is well-behaved. We proceed by induction on vertex-phase pairs. First note that since $\delta^0(v, S) = 0$ by definition, we know that for all vertices v and sets S , v is good for S during its 1^{st} phase. Now, for any pair (v, r) with $1 \leq r < b$, we would like to show that for any set S such that v is S -typical, v is good for S during its $(r+1)^{th}$ phase.

Since we proceed to by strong induction on the vertex-phase pair, we fix (v, r) and suppose that for any $(u, s) \prec (v, r)$ and any set S' such that u is S' -typical. Then our induction hypothesis states that u is good for S' during its $(s+1)^{th}$ phase. Consider any S such that v is S -typical. Since we know the coloring is well-behaved, we know that for any phase $\ell \leq r$ of v , $|B(A^{\ell-1}(v))|, |B(A^{\ell-1}(v) \cap S)| \leq C$ (recall Definition 28). Let

$$T_B^\ell(v) = \{i \in T^\ell(v) : e_i - v \in B(A^{\ell-1}(v)) \cup B(A^{\ell-1}(v) \cap S)\},$$

so that $|T_B^\ell(v)| \leq 2C$. Then, for any $i \in T^\ell(v) \setminus T_B^\ell(v)$, with $u = e_i - v$ and $\phi_i(u) = s+1$, $\mathbf{last}^s(u) < i \leq \mathbf{last}^r(v)$, so $(u, s) \prec (v, r)$. Thus, by the inductive hypothesis, since u is both $A^{\ell-1}(v)$ -typical and $(A^{\ell-1}(v) \cap S)$ -typical, u must be good for both $A^{\ell-1}(v)$ and $A^{\ell-1}(v) \cap S$ during its $(s+1)^{th}$ phase.

Recall that by Proposition 26,

$$|\delta^r(v, S)| \leq \frac{1}{|A^r(v)|} \sum_{i \in T^{\leq r}(v)} Z_i + \left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in T^\ell(v)} D_i(S) \right| + \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in \widetilde{T}^\ell(v)} \left| p_i(S) - \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right|.$$

Since the coloring is well-behaved, we know that $\mathcal{W}(v)$ does not occur, and therefore,

$$\sum_{i \in T^{\leq r}(v)} Z_i - q_i \leq \alpha \Delta.$$

By Proposition 35,

$$\begin{aligned} \sum_{i \in T^{\leq r}(v)} Z_i &\leq \alpha \Delta + \sum_{\ell \leq r} \sum_{i \in T^\ell(v) \setminus T_B^\ell(v)} q_i + \sum_{\ell \leq r} \sum_{i \in T_B^\ell(v)} q_i \\ &\leq \alpha \Delta + \sum_{\ell \leq r} \sum_{i \in T^\ell(v) \setminus T_B^\ell(v)} \frac{\alpha}{4} + \sum_{\ell \leq r} \sum_{i \in T_B^\ell(v)} 1 \\ &\leq \alpha \Delta + \frac{\alpha \Delta}{4} + 2bC \\ &\leq 2\alpha \Delta. \end{aligned} \tag{15}$$

Furthermore, since v is S -typical, we are guaranteed that

$$\left| \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in T^\ell(v)} D_i(S) \right| \leq \frac{\alpha}{\epsilon} \leq \frac{\alpha \Delta (1 + \epsilon)}{\epsilon \cdot |A^r(v)|}.$$

Finally, again by Proposition 35,

$$\begin{aligned} &\sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in \tilde{T}^\ell(v)} \left| p_i(S) - \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right| \\ &= \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in \tilde{T}^\ell(v) \setminus T_B^\ell(v)} \left| p_i(S) - \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right| + \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in \tilde{T}^\ell(v) \cap T_B^\ell(v)} \left| p_i(S) - \frac{|A^{\ell-1}(v) \cap S|}{|A^{\ell-1}(v)|} \right| \\ &= \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{\substack{i \in \tilde{T}^\ell(v) \setminus T_B^\ell(v) \\ s = \phi(e_i - v) - 1}} \frac{5\hat{\epsilon}^s(e_i - v)}{\epsilon^2} + \sum_{\ell=1}^r \frac{1}{|A^\ell(v)|} \sum_{i \in \tilde{T}^\ell(v) \cap T_B^\ell(v)} 1 \\ &\leq \frac{1}{|A^r(v)|} \left(2bC + \sum_{\substack{i \in T^{\leq r}(v) \\ s = \phi(e_i - v) - 1}} \frac{5\hat{\epsilon}^s(e_i - v)}{\epsilon^2} \right) \end{aligned}$$

Combining this gives us

$$\begin{aligned} |\delta^r(v, S)| &\leq \frac{1}{|A^r(v)|} \left(2\alpha \Delta + \frac{1 + \epsilon}{\epsilon} \alpha \Delta + 2bC + \frac{5}{\epsilon^2} \sum_{\substack{i \in T^{\leq r}(v) \\ s = \phi(e_i - v) - 1}} \hat{\epsilon}^s(e_i - v) \right) \\ &\leq \frac{\Delta}{|A^r(v)|} \left(\zeta + \frac{5}{\epsilon^2 \Delta} \sum_{\substack{i \in T^{\leq r}(v) \\ s = \phi(e_i - v) - 1}} \hat{\epsilon}^s(e_i - v) \right) = \frac{\hat{\epsilon}^r(v) \Delta}{|A^r(v)|}, \end{aligned}$$

where the second inequality holds because $2\alpha + \frac{1 + \epsilon}{\epsilon} \alpha + \frac{2bC}{\Delta} \leq \frac{5}{\epsilon} \alpha \leq \zeta$, for Δ sufficiently large. \square

Finally, we use the main lemma to show that in a well-behaved coloring, no edge could have been left uncolored.

Corollary 36. For $\zeta > 0$, let phase partition counter $\phi = \{\phi_i\}_{i=1}^m$ have ζ -controlled error with respect to the ordered graph (G, σ) . If ϕ is balanced with respect to (G, σ) and the coloring produced by the \mathcal{A} is well-behaved, then no edge is left uncolored.

Proof. Suppose for contradiction that there is an edge $e = (u, v)$ and time i such that $F_i(u) \cap F_i(v) = \emptyset$. Let $S_j = F_{j-1}(u) \cap F_{j-1}(v)$ and e_t be the edge adjacent to u or v that uses up the last color in $F_{t-1}(u) \cap F_{t-1}(v) = S_t$. This gives us $|S_1| \geq |S_2| \geq \dots \geq |S_t| = 1$ and $|S_{t+1}| = 0$. Let $R = \{k \in T(u) \cup T(v) : k \leq t\}$ be the set of arrival times of edges adjacent to either u or v and define the partition $\{R_j\}$ of R to be $R_0 = \{k \in R : |S_k| \geq \frac{\epsilon^2 \Delta}{2}\}$ and for $j \geq 1$,

$$R_j = \left\{ k \in R : \frac{\epsilon^2 \Delta}{2^{j+1}} \leq |S_k| < \frac{\epsilon^2 \Delta}{2^j} \right\}$$

Note that $|\bigcup_j R_j| \leq |T(u) \cup T(v)| \leq 2\Delta$. We will derive a contradiction by establishing lower bounds on the size of each R_j , and show their sum exceeds 2Δ . Since $S_1 = (1 + \epsilon)\Delta$, $|S_i|$ can drop below $\epsilon^2 \Delta / 2$ only after at least $(1 + \epsilon - \epsilon^2 / 2)\Delta$ edges incident on u or v have arrived, we have

$$|R_0| \geq (1 + \epsilon - \epsilon^2 / 2)\Delta. \quad (16)$$

Claim 37. For each $j \geq 1$, $|R_j| \geq \frac{\epsilon^2 \Delta}{8} - 7\alpha\Delta 2^{j-2}$

Proof. Since the coloring is well-behaved, for each $1 \leq r \leq b$, $|B(A^r(u))|, |B(A^r(v))| \leq C$, so if we let

$$R^B = \left\{ k \in R : k \in \bigcup_{r=1}^b (B(A^r(u)) \cup B(A^r(v))) \right\},$$

we have $|R^B| \leq 2bC$. Then, since $0 \leq Z_i, q_i \leq 1$ for all i and $\mathcal{W}(v), \mathcal{W}(u)$ didn't occur, we have by Proposition 35

$$\begin{aligned} \sum_{i \in R} Z_i &\leq \sum_{i \in T(u), i \leq t} Z_i + \sum_{i \in T(v), i \leq t} Z_i \\ &\leq 2\alpha\Delta + \sum_{i \in T(u), i \leq t} q_i + \sum_{i \in T(v), i \leq t} q_i && \text{(From Definition 30(a) and Lemma 31)} \\ &\leq 2\alpha\Delta + 2 \sum_{i \in R} q_i && \text{(From definition of } R) \\ &\leq 2\alpha\Delta + 2 \sum_{i \in R \setminus R^B} q_i + 2 \sum_{i \in R^B} q_i \\ &\leq 2\alpha\Delta + 4\Delta \cdot \frac{\alpha}{4} + 4bC && \text{(From Proposition 35(b))} \\ &\leq 4\alpha\Delta. \end{aligned}$$

Note that since $Z_i \in [0, 1]$ for all i , this means that for any $R' \subseteq R$, $\sum_{i \in R'} Z_i \leq 4\alpha\Delta$.

Similarly, since $\mathcal{D}(u, v)$ did not occur, and the sets R_j are intersections of $T(u) \cup T(v)$ with intervals we have for all j ,

$$\begin{aligned} \sum_{i \in R_j} X_i(S_i) &\leq \left| \sum_{i \in R_j} p_i(S_i) \right| + \left| \sum_{i \in R_j} D_i(S_i) \right| \leq \sum_{i \in R_j} p_i(S_i) + 2\alpha\Delta \\ &\leq \sum_{i \in R_j \setminus R^B} p_i(S_i) + 2\alpha\Delta + 2bC \\ &\leq \sum_{i \in R_j \setminus R^B} p_i(S_i) + 3\alpha\Delta \end{aligned}$$

for all j . Finally, for $i \in R_j \setminus R^B$, suppose without loss of generality that $i \in T^r(v)$, with $x = e_i - v$ and $\phi_i(x) = s$. Then by Proposition 35,

$$p_i(S_i) = \frac{|S_i|}{|A^{r-1}(v) \cap A^{s-1}(x)|} < \frac{\epsilon^2 \Delta / 2^j}{2\epsilon^2 \Delta / 5} \leq 5 \cdot 2^{-j-1} \leq 2^{-j+2},$$

so

$$\sum_{i \in R_j} X_i(S_i) \leq |R_j|2^{-j+2} + 3\alpha\Delta$$

On the other hand, since the colors of at least $\frac{\epsilon^2\Delta}{2^{j+1}}$ edges e_i with $i \in R_j$ must hit S_i , we have

$$\frac{\epsilon^2\Delta}{2^{j+1}} \leq \sum_{i \in R_j} Y_i(S_i) \leq \sum_{i \in R_j} X_i(S_i) + \sum_{i \in R_j} Z_i \leq \sum_{i \in R_j} X_i(S_i) + 4\alpha\Delta.$$

Therefore,

$$\frac{\epsilon^2\Delta}{2^{j+1}} - 4\alpha\Delta \leq \sum_{i \in R_j} X_i(S_i) \leq |R_j|2^{-j+2} + 3\alpha\Delta$$

so

$$|R_j| \geq \frac{\epsilon^2\Delta}{8} - 7\alpha\Delta \cdot 2^{j-2},$$

to complete the proof of the claim. \square

Let $h = \lceil 8/\epsilon^2 \rceil$. Summing the lower bounds on R_j for $1 \leq j \leq h$ yields:

$$\left| \bigcup_{j=1}^h R_j \right| \geq \Delta - 7\alpha\Delta \cdot 2^h \geq \left(1 - \frac{\epsilon^3}{10}\right) \Delta$$

since $7\alpha 2^{8/\epsilon^2+1} < 14\alpha e^{8/\epsilon^2} \leq 14\zeta e^{8/\epsilon^2} \leq \frac{\epsilon^3}{10}$ by our choice of ζ from Definition 10. Combining with (Equation (16)) and the fact that $\epsilon \leq 1$, yields the desired contradiction:

$$\left| \bigcup_{j=0}^h R_j \right| > 2\Delta.$$

\square

We can now complete the proofs of Theorem 7 and Theorem 6. According to Corollary 36, \mathcal{A}' succeeds provided the partition function is ζ -controlled with respect to (G, σ) and the coloring is well-behaved.

In the dense case of Theorem 7, Proposition 19 ensures that there is a partition function ϕ^D that is balanced and ζ -error controlled for (G, σ) . Note that for any M , if Δ is sufficiently large, then $n \leq M\Delta$ implies $n \leq 2^{\frac{\Delta}{M}}$. Thus, in this case Lemma 31 implies that the coloring is well-behaved with probability at least $1 - 2^{-\alpha^4\Delta/1000}$, so this upper bounds the probability that \mathcal{A}' (and also \mathcal{A}) fails to color (G, σ) . Thus, letting $\gamma = \alpha^4$, we have our proof for Theorem 7.

In the random case of Theorem 6, from Lemma 14, the phase counter ϕ^R is balanced with respect to (G, σ) with probability at least $1 - 2^{-\Delta/(20b)}$, and combined with Proposition 20 this ensures that the partition function is ζ -controlled with respect to (G, σ) . As in the dense case the probability that the coloring is not well behaved is at most $2^{-\alpha^4\Delta/1000}$. Therefore for all but at most a $2^{-\frac{\Delta}{20b}}$ fraction of edge orderings, the probability that \mathcal{A} succeeds against $\mathbf{obl}(G, \sigma)$ is at least $1 - 2^{-\alpha^4\Delta/1000}$. Thus, letting $\gamma_1 = \frac{1}{20b}$ and $\gamma_2 = \frac{\alpha^4}{1000}$, we have our claim.

Acknowledgements

We would like to thank Jeff Kahn for helpful discussions.

References

- [Arj82] Eshrat Arjomandi. An efficient algorithm for colouring the edges of a graph with delta + 1 colours. *INFOR: Information Systems and Operational Research*, 20(2):82–101, 1982.

- [Ass24] Sepehr Assadi. Faster vizing and near-vizing edge coloring algorithms. *arXiv*, 2024. <https://arxiv.org/abs/2405.13371>.
- [ASvS23] Aflatoun Amouzandeh, Soren Schmitt, and Rob van Stee. Sigact news online algorithms column 40: 2022 in review. *ACM SIGACT News*, 54(1):90–104, 2023. <https://dl.acm.org/doi/abs/10.1145/3586165.3586177>.
- [BBK⁺94] Shai Ben-David, Allan Borodin, Richard M. Karp, Gábor Tardos, and Avi Wigderson. On the power of randomization in on-line algorithms. *Algorithmica*, 11(1):2–14, 1994. <https://doi.org/10.1007/BF01294260>.
- [BBKO22] Alkida Balliu, Sebastian Brandt, Fabian Kuhn, and Dennis Olivetti. Distributed edge coloring in time polylogarithmic in Δ . In *Proceedings of the 2022 ACM Symposium on Principles of Distributed Computing*, pages 15–25, 2022.
- [BCC⁺24] Sayan Bhattacharya, Din Carmon, Martin Costa, Shay Solomon, and Zhang Tianyi. Faster $\delta + 1$ -edge coloring: Breaking the $m\sqrt{n}$ time barrier. *arXiv*, 2024. <https://arxiv.org/abs/2405.15449>.
- [BCHN18] Sayan Bhattacharya, Deeparnab Chakrabarty, Monika Henzinger, and Danupon Nanongkai. Dynamic algorithms for graph coloring. In *Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1–20. SIAM, 2018.
- [BCPS24] Sayan Bhattacharya, Martin Costa, Nadav Panski, and Shay Solomon. Nibbling at long cycles: Dynamic (and static) edge coloring in optimal time. In *Proceedings of the ACM-SIAM Symposium on Discrete Algorithms (SODA)*. SIAM, 2024.
- [BDH⁺19] Soheil Behnezhad, Mahsa Derakhshan, MohammadTaghi Hajiaghayi, Marina Knittel, and Hamed Saleh. Streaming and massively parallel algorithms for edge coloring. In *27th Annual European Symposium on Algorithms (ESA)*, volume 144 of *LIPICs*, pages 15:1–15:14, 2019.
- [Ber22] Anton Bernshteyn. A fast distributed algorithm for $(\delta + 1)$ -edge-coloring. *J. Comb. Theory, Ser. B*, 152:319–352, 2022.
- [BGW21] Sayan Bhattacharya, Fabrizio Grandoni, and David Wajc. Online edge coloring algorithms via the nibble method. In Dániel Marx, editor, *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms, SODA 2021, Virtual Conference, January 10 - 13, 2021*, pages 2830–2842. SIAM, 2021.
- [BM17] Leonid Barenboim and Tzalik Maimon. Fully-dynamic graph algorithms with sublinear time inspired by distributed computing. In *International Conference on Computational Science (ICCS)*, volume 108 of *Procedia Computer Science*, pages 89–98. Elsevier, 2017.
- [BMM10] Bahman Bahmani, Aranyak Mehta, and Rajeev Motwani. A 1.43-competitive online graph edge coloring algorithm in the random order arrival model. In Moses Charikar, editor, *Proceedings of the Twenty-First Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2010, Austin, Texas, USA, January 17-19, 2010*, pages 31–39. SIAM, 2010.
- [BMN92] Amotz Bar-Noy, Rajeev Motwani, and Joseph Naor. The greedy algorithm is optimal for on-line edge coloring. *Inf. Process. Lett.*, 44(5):251–253, 1992.
- [BS23] Soheil Behnezhad and Mohammad Saneian. Streaming edge coloring with asymptotically optimal colors. *arXiv preprint arXiv:2305.01714*, 2023.
- [BSVW23] Joakim Blikstad, Ola Svensson, Radu Vintan, and David Wajc. Simple and asymptotically optimal online bipartite edge coloring. *CoRR*, abs/2311.04574, 2023.
- [BSVW24] Joakim Blikstad, Ola Svensson, Radu Vintan, and David Wajc. Online edge coloring is (nearly) as easy as offline. *CoRR*, abs/2402.18339, 2024.

- [CHL⁺20] Yi-Jun Chang, Qizheng He, Wenzheng Li, Seth Pettie, and Jara Uitto. Distributed edge coloring and a special case of the constructive lovász local lemma. *ACM Trans. Algorithms*, 16(1):8:1–8:51, 2020.
- [Chr23] Aleksander Bjørn Grodt Christiansen. The power of multi-step vizing chains. In *Proceedings of the 55th Annual ACM Symposium on Theory of Computing (STOC)*, pages 1013–1026. ACM, 2023.
- [Chr24] Aleksander B. G. Christiansen. Deterministic dynamic edge-colouring. *CoRR*, abs/2402.13139, 2024.
- [CMZ23] Shiri Chechik, Doron Mukhtar, and Tianyi Zhang. Streaming edge coloring with subquadratic palette size. *arXiv preprint arXiv:2305.07090*, 2023.
- [CPW19] Ilan Reuven Cohen, Binghui Peng, and David Wajc. Tight bounds for online edge coloring. In David Zuckerman, editor, *60th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2019, Baltimore, Maryland, USA, November 9-12, 2019*, pages 1–25. IEEE Computer Society, 2019.
- [Dav23] Peter Davies. Improved distributed algorithms for the lovász local lemma and edge coloring. In *Proceedings of the ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 4273–4295. SIAM, 2023.
- [DHZ19] Ran Duan, Haoqing He, and Tianyi Zhang. Dynamic edge coloring with improved approximation. In Timothy M. Chan, editor, *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2019, San Diego, California, USA, January 6-9, 2019*, pages 1937–1945. SIAM, 2019.
- [EK24] Michael Elkin and Ariel Khuzman. Deterministic simple $(1+\epsilon)\Delta$ -edge-coloring in near-linear time. *CoRR*, abs/2401.10538, 2024.
- [FGK17] Manuela Fischer, Mohsen Ghaffari, and Fabian Kuhn. Deterministic distributed edge-coloring via hypergraph maximal matching. In *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 180–191. IEEE, 2017.
- [FMP] Tomás Feder, Rajeev Motwani, and Rina Panigrahy. Online algorithms for edge coloring. <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=e300284ac75ed317c6550bea907>
- [Fre75] David A. Freedman. On tail probabilities for martingales. *The Annals of Probability*, 3(1):100–118, 1975.
- [GKMU18] Mohsen Ghaffari, Fabian Kuhn, Yannic Maus, and Jara Uitto. Deterministic distributed edge-coloring with fewer colors. In *Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing*, pages 418–430, 2018.
- [GNK⁺85] Harold N Gabow, Takao Nishizeki, Oded Kariv, Daneil Leven, and Osamu Terada. Algorithms for edge coloring. *Technical Report*, 1985.
- [GS23] Prantar Ghosh and Manuel Stoeckl. Low-memory algorithms for online and w-streaming edge coloring. *arXiv preprint arXiv:2304.12285*, 2023.
- [Hol81] Ian Holyer. The np-completeness of edge-coloring. *SIAM J. Comput.*, 10(4):718–720, 1981.
- [JLR00] Svante Janson, Tomasz Luczak, and Andrzej Rucinski. *Random graphs*. Wiley-Interscience series in discrete mathematics and optimization. Wiley, 2000.
- [KLS⁺22] Janardhan Kulkarni, Yang P. Liu, Ashwin Sah, Mehtaab Sawhney, and Jakub Tarnawski. Online edge coloring via tree recurrences and correlation decay. In Stefano Leonardi and Anupam Gupta, editors, *STOC '22: 54th Annual ACM SIGACT Symposium on Theory of Computing, Rome, Italy, June 20 - 24, 2022*, pages 104–116. ACM, 2022.

- [MG92] Jayadev Misra and David Gries. A constructive proof of vizing’s theorem. *Inf. Process. Lett.*, 41(3):131–133, 1992.
- [PR01] Alessandro Panconesi and Romeo Rizzi. Some simple distributed algorithms for sparse networks. *Distributed computing*, 14(2):97–100, 2001.
- [Sin19] Corwin Sinnamon. A randomized algorithm for edge-colouring graphs in $o(m\sqrt{n})$ time. *CoRR*, abs/1907.03201, 2019.
- [SW21] Amin Saberi and David Wajc. The greedy algorithm is not optimal for on-line edge coloring. In Nikhil Bansal, Emanuela Merelli, and James Worrell, editors, *48th International Colloquium on Automata, Languages, and Programming, ICALP 2021, July 12-16, 2021, Glasgow, Scotland (Virtual Conference)*, volume 198 of *LIPICs*, pages 109:1–109:18. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.