

AN UNIFORM LOWER BOUND FOR CLASSICAL KLOOSTERMAN SUMS AND AN APPLICATION

JEWEL MAHAJAN, JISHU DAS, AND STEPHAN BAIER

ABSTRACT. We present an elementary uniform lower bound for the classical Kloosterman sum $S(a, b; c)$ under the condition of its non-vanishing and $(ab, c) = 1$, with c being an odd integer. We then apply this lower bound for Kloosterman sums to derive an explicit lower bound in the Petersson's trace formula, subject to a pertinent condition. Consequently, we achieve a modified version of [JS20, Theorem 1.7], wherein the parameters k and N are permitted to vary independently.

1. INTRODUCTION

The classical Kloosterman sum $S(a, b; c)$ is given by

$$S(a, b; c) = \sum_{\substack{x \pmod{c} \\ \gcd(x, c) = 1}} e^{2\pi i \left(\frac{ax + b\overline{\mathfrak{X}}(c)}{c} \right)},$$

where a, b are integers, c is a natural number and $\overline{\mathfrak{X}}(c)$ denotes the multiplicative inverse of x modulo c . In 1926, Kloosterman sums were introduced to solve the following problem: Given natural numbers u, v, w, z and n , Kloosterman obtained an asymptotic formula in [Klo27] for the number of representations of a natural number n in the form $ux^2 + vy^2 + wz^2 + zt^2$. The sums for the special case of $a = 0$ or $b = 0$ are called Ramanujan sums. Kloosterman sums appear in various problems in number theory. One of the important places of its occurrence is in the Kuznetsov trace formula [Kuz80, Theorem 1]. Recently the non-vanishing aspect of Kloosterman sums has been used in [BDS23, Theorem 2] and [Das23] to obtain discrepancy results for the distribution of eigenvalues of Hecke operators. Whenever we have non-vanishing of a sum, a natural question that arises is whether we can obtain a lower bound for the same. In the next section, we answer this question for $S(a, b; c)$, where $(ab, c) = 1$. First, we state our main result on the obtained lower bound of classical Kloosterman sums.

Theorem 1. *Let c be an odd natural number such that $c = df$ where f is squarefree and d is a powerful number with $(d, f) = 1$. Let a, b integers such that $(ab, c) = 1$ and ab is a quadratic residue modulo each prime dividing d (and hence, ab is a square modulo d by Hensel's lemma and Chinese remainder theorem). Then*

$$|S(a, b; c)| \geq \left(\prod_{p|d} \frac{2}{\sqrt{p}} \right) \left(\frac{1}{(\tau(f)\sqrt{f})^{\phi(f)/\tau(f)-1}} \right),$$

where $\tau(f)$ is the number of positive divisors of f . If ab is a quadratic non-residue modulo a prime dividing d (and hence ab is not a square modulo d), then $S(a, b; c) = 0$.

Let $S_k(N)$ denote the space of cusp forms of even weight $k \geq 2$ and level N . The discrepancy between two probability measures involving eigenvalues of cusp forms on a given closed interval has been a well-studied topic in number theory. The upper bound (see [MS09, Theorem 2], [TW16, Theorem 1]) and the lower bound estimates (see [JS20, Theorem 1.1, Theorem 1.6], [Das23, Theorem 1, Theorem 2]) of the discrepancy of two probability measures are of particular interest due to its various applications to other problems. For upper bounds, the key technique is the Eichler-Selberg trace formula ([Sel56]) and for lower bounds, one

Date: June 21, 2024.

2020 Mathematics Subject Classification. Primary 11L05, 11F72.

Key words and phrases. Kloosterman sums, Petersson trace formula.

of the key ingredients is Petersson's trace formula ([Pet32]). We obtain the following version of Petersson's trace formula for the space $S_k(N)$ as an application to Theorem 1.

Theorem 2. *Let m, n, N be natural numbers with $(mn, N) = 1$ and N odd. Let $H_0 = 1 - \frac{4}{9} - \log(\frac{9}{5})$, $D_0 \in (e^{H_0}, 1)$ be a real number. Let k_0 be the least positive even integer greater than or equal to 22 satisfying the inequality $(D_0 e^{-H_0})^{k-1} \geq 7$. Let $N = df$ where d is powerful and f is squarefree, where $(d, f) = 1$. Let $\frac{4\pi\sqrt{mn}}{N} \in [D_0(k-1), (k-1)]$ and assume that mn is a quadratic residue modulo each prime dividing d . Let $k_0(N) = 2 + \left\lfloor \frac{\log 7 - \log H(N)}{\log A_0} \right\rfloor$, where $A_0 = D_0 e^{-H_0}$, and $H(N) = 2^{\omega(N)} \frac{G(N)}{N^{5/4}}$. Then, for all $k \geq k_0(N)$, and with $\Delta_{k,N}(m, n)$ as defined in equation (3.1), we have*

$$|\Delta_{k,N}(m, n) - \delta(m, n)| \geq 7.99\pi J_{k-1}(k-1) e^{(k-1)(1 - \frac{4}{9} - \log(\frac{9}{5}))},$$

where $G(N) = \frac{f^{3/4}}{(\tau(f)^2 f)^{\phi(f)/2\tau(f)}}$.

Some key features of Theorem 2 are as follows:

- The main results that help us to achieve Theorem 2 are as follows. Theorem 1 gives us an explicit lower bound for a Kloosterman sum appearing in $\Delta_{k,N}(m, n)$. We then use explicit bounds for the J -Bessel function as given in Lemma 3.1 and Lemma 3.2. We emphasise that Lemma 3.2, which follows from a direct application of [Kra14, Theorem 1], is crucial for deriving an explicit lower bound for $|\Delta_{k,N}(m, n) - \delta(m, n)|$. One of the main purposes of [Kra14] is to derive an explicit expression for $J_k(x)$ for a transition range of x and for any $k \geq 2$.
- The theorems [JS20, Theorem 1.7] and [Das23, Theorem 4], assume N to be fixed and provide an asymptotic for $|\Delta_{k,N}(m, n) - \delta(m, n)|$ as $k \rightarrow \infty$. However, Theorem 2 is applicable for every k and N , provided they satisfy certain hypotheses. For a fixed level N , [ILS00, Corollary 2.3], gives an asymptotic for $\Delta_{k,N}(m, n)$ with the condition $\frac{4\pi\sqrt{mn}}{N} \leq \frac{k}{3}$, whereas Theorem 2 gives us an asymptotic for the condition $D_0(k-1) \leq \frac{4\pi\sqrt{mn}}{N} \leq (k-1)$.
- In comparison to [JS20, Theorem 1.7], we increase the size of the interval in which $\frac{4\pi\sqrt{mn}}{N}$ lies to $(1 - D_0)k$ instead of $2k^{\frac{1}{3}}$, which weakens the lower bound to $e^{(k-1)(1 - \frac{4}{9} - \log(\frac{9}{5}))}$ as compared to $k^{-\frac{1}{3}}$. However, for a shorter range given by $[(k-1), (k-1) + (k-1)^{\frac{1}{3}}]$, we obtain back the lower bound of order $(k-1)^{-\frac{1}{3}}$, which is given by the following theorem.

Theorem 3. *Let m, n, N be natural numbers with $(mn, N) = 1$ and N odd. Let $N = df$ where d is powerful and f is squarefree, where $(d, f) = 1$. Let $k \geq 180$, $\frac{4\pi\sqrt{mn}}{N} \in [(k-1), (k-1) + (k-1)^{\frac{1}{3}}]$ and assume that mn is a quadratic residue modulo each prime that divides the powerful part of N . Let E_0 be a positive constant less than $0.327H(N)$ and $k_1(N) := 2 + \left\lfloor \frac{\log(1.632) - \log(0.327H(N) - E_0)}{-H_0} \right\rfloor$. Then, for all $k \geq \max(180, k_1(N))$, we have*

$$|\Delta_{k,N}(m, n) - \delta(m, n)| \geq \frac{2\pi E_0}{(k-1)^{\frac{1}{3}}}.$$

Organization of the article. The article is organised as follows. In Section 2, we discuss the proof of the lower bound of classical Kloosterman sums. In Sections 3, we prove Theorems 2 and 3.

2. LOWER BOUND FOR KLOOSTERMAN SUMS

In this section, we primarily prove Theorem 1 by considering squarefree and odd prime power cases separately and we consider the general case in section 2.3.

2.1. Squarefree case.

Proposition 1. *Let c be a squarefree positive integer such that $(a, c) = 1$. Then*

$$|S(a, b; c)| \geq \frac{1}{(\tau(c)\sqrt{c})^{\phi(c)/\tau(c)-1}},$$

Proof. Let $\alpha = S(a, b; c) \in \mathbb{Q}(\zeta_c)$, where $\zeta_c = e^{2\pi i/c}$. An element of $\text{Gal}(\mathbb{Q}(\zeta_c)/\mathbb{Q})$ sends ζ_c to ζ_c^n where $(n, c) = 1$. Hence, the conjugates of α are the Kloosterman sums $S(an, bn; c)$, where $(n, c) = 1$. In fact, we have $S(an, bn; c) = S(an^2, b; c) = S(a, bn^2; c)$. For c prime, $S(a, b; c)$ is a linear combination of at most $c - 1$ c -th roots of unity, but they are necessarily linearly independent over \mathbb{Q} . So if the linear combination becomes 0, all the coefficients need to be 0 which is not the case. Hence $S(a, b; c) \neq 0$ for c prime. Using the multiplicativity of Kloosterman sums (see Lemma 4), $S(a, b; c) \neq 0$ for squarefree c . Let $\mathcal{N}(\alpha)$ denote the norm of $\alpha \in \mathbb{Q}(\zeta_c)$ over \mathbb{Q} . Since $\alpha = S(a, b; c)$ is a non-zero algebraic integer in $\mathbb{Q}(\zeta_c)$, its norm $\mathcal{N}(\alpha)$ is a non-zero integer and therefore,

$$1 \leq |\mathcal{N}(\alpha)| = |\mathcal{N}(S(a, b; c))| = \prod_{\substack{(n,c)=1 \\ 1 \leq n \leq c}} |S(an^2, b; c)|.$$

Hence,

$$\prod_{\substack{(n,c)=1 \\ 1 \leq n \leq c}} |S(an^2, b; c)| \geq 1.$$

Let $\rho(c)$ be the number of solutions modulo c of $x^2 \equiv 1 \pmod{c}$. We have $\rho(c) = \tau(c)$ for squarefree c . Using Weil's bound together with $(a, c) = 1$, we have

$$\frac{1}{|S(a, b; c)|^{\tau(c)}} \leq \prod_{\substack{(n,c)=1 \\ 1 < n \leq c \\ n^2 \not\equiv 1(c)}} |S(an^2, b; c)| \leq \prod_{\substack{(n,c)=1 \\ 1 < n \leq c \\ n^2 \not\equiv 1(c)}} (\tau(c)\sqrt{c}) = (\tau(c)\sqrt{c})^{\phi(c) - \tau(c)}.$$

Hence we obtain

$$|S(a, b; c)| \geq \frac{1}{(\tau(c)\sqrt{c})^{\phi(c)/\tau(c) - 1}}.$$

□

2.2. Odd prime power case. Let q be odd, and ϵ_q be 1 or i depending on whether $q \equiv 1$ or -1 modulo 4. For odd prime power moduli q , we have the following explicit evaluation of $S(a, b; q)$.

Proposition 2. [IK04, Chapter 12, Exercise 1] *Let $q = p^\beta$ for some prime p and $\beta \geq 2$. Let p does not divide $2ab$. If ab is a quadratic residue modulo p , then*

$$(2.1) \quad S(a, b; q) = 2 \left(\frac{l}{q} \right) \sqrt{q} \operatorname{Re} \left(\epsilon_q e \left(\frac{2l}{q} \right) \right),$$

where $l^2 \equiv ab \pmod{q}$. If ab is a quadratic non-residue modulo p , then we have $S(a, b; q) = 0$.

From the above Proposition, we deduce the following lower bound for $|S(a, b; q)|$, if q is an odd prime power.

Proposition 3. *Let $q = p^\beta$ for some prime p and $\beta \geq 2$. Suppose p does not divide $2ab$. If ab is a quadratic residue modulo p , then*

$$|S(a, b; q)| \geq \frac{2}{\sqrt{q}}.$$

If ab is a quadratic non-residue modulo p , then $S(a, b; q) = 0$.

Proof. Assume that ab is a quadratic residue modulo p . Using Proposition 2, we have

$$S(a, b; q) = 2 \left(\frac{l}{q} \right) \sqrt{q} \operatorname{Re} \left(\epsilon_q e \left(\frac{2l}{q} \right) \right),$$

where

$$(2.2) \quad l^2 \equiv ab \pmod{q}.$$

We start by noting that q does not divide l and hence $\left| \left(\frac{l}{q} \right) \right| = 1$. Let $l = qt + r$ where $0 \leq r \leq q - 1$. Note that $p \nmid 2ab$ implies that p is odd and so is q . We consider the following two cases depending on whether $q \equiv 1$ or -1 modulo 4.

Case 1: $q \equiv 1 \pmod{4}$

In this case, we have $\epsilon_q = 1$. For $x \in \mathbb{R}$, let $\|x\|$ denote the distance from x to the nearest integer. Using $|\sin \pi x| \geq 2\|x\|$ for all real x , we have

$$\begin{aligned} |S(a, b; q)| &= 2 \left| \left(\frac{l}{q} \right) \sqrt{q} \operatorname{Re} \left(e \left(\frac{2l}{q} \right) \right) \right| = 2\sqrt{q} \left| \cos \left(\frac{4\pi r}{q} \right) \right| \\ &= 2\sqrt{q} \left| \sin \left(\frac{\pi}{2} - \frac{4\pi r}{q} \right) \right| \geq 4\sqrt{q} \left\| \frac{1}{2} - \frac{4r}{q} \right\|. \end{aligned}$$

Case 2: $q \equiv -1 \pmod{4}$

In this case, we have $\epsilon_q = i$ and hence, we have

$$\begin{aligned} |S(a, b; q)| &= 2 \left| \left(\frac{l}{q} \right) \sqrt{q} \operatorname{Re} \left(ie \left(\frac{2l}{q} \right) \right) \right| \\ &= 2\sqrt{q} \left| \sin \left(\frac{4\pi r}{q} \right) \right| \geq 4\sqrt{q} \left\| \frac{4r}{q} \right\|. \end{aligned}$$

Combining both cases, we obtain

$$(2.3) \quad |S(a, b; q)| \geq 4\sqrt{q} \min \left(\left\| \frac{q-8r}{2q} \right\|, \left\| \frac{4r}{q} \right\| \right).$$

□

We first show that $\left\| \frac{q-8r}{2q} \right\| \neq 0$. Suppose not, then $\frac{q-8r}{2q} = t$ for some $t \in \{0, -1, -2, -3\}$ as $0 \leq r \leq q-1$. We have $p \mid r^2 - ab$ which implies that p divides $\frac{(1-2t)^2 q^2 - 64ab}{64}$. This is an impossibility since p does not divide $2ab$. A similar argument yields $\left\| \frac{4r}{q} \right\| \neq 0$. Proceeding now from the equation (2.3), we conclude that

$$|S(a, b; q)| \geq 4\sqrt{q} \min \left(\left\| \frac{q-8r}{2q} \right\|, \left\| \frac{4r}{q} \right\| \right) \geq 4\sqrt{q} \min \left(\frac{1}{2q}, \frac{1}{q} \right) \geq \frac{2}{\sqrt{q}}.$$

2.3. General Case. In this section, we apply the multiplicativity property of Kloosterman sums. For $m, n \in \mathbb{N}$, $(m, n) = 1$, let $\overline{m}(n)$ denote the multiplicative inverse of m in $(\mathbb{Z}/n\mathbb{Z})^*$ i.e. $m\overline{m}(n) \equiv 1 \pmod{n}$. We have the following factorization of Kloosterman sums into Kloosterman sums to prime power moduli.

Proposition 4. [Das23, Lemma 3.1] *Let $c = q_1^{\beta_1} \cdots q_t^{\beta_t}$, where q_1, \dots, q_t are distinct prime numbers, $q_i^{\beta_i} = l_i$, and*

$$c_i = \frac{c}{\prod_{j=1}^i l_j}$$

for $1 \leq i \leq t$. Further let $m_0 = 1$, $m_i = \prod_{j=1}^i \overline{l_j}(c_j)$ for $1 \leq i \leq t-1$. Then

$$S(a, b; c) = \prod_{i=1}^t S(am_{i-1}\overline{c_i}(l_i), bm_{i-1}\overline{c_i}(l_i); l_i).$$

Proof. The proof uses equation (1.59) of [IK04] repetitively which is given by

$$S(a, b; ll') = S(a\overline{l'}(l), b\overline{l'}(l); l) S(a\overline{l}(l'), b\overline{l}(l'); l')$$

where $(l, l') = 1$. □

Remark 1. *Assume the hypotheses in the above Proposition 4 are satisfied. We note that ab is a quadratic residue modulo q_i if and only if $a'b'$ with $a' = am_{i-1}\overline{c_i}(l_i)$ and $b' = bm_{i-1}\overline{c_i}(l_i)$ is a quadratic residue modulo q_i .*

Now we give a proof of Theorem 1.

Proof of Theorem 1. We first factorize $S(a, b; c)$ using Proposition 4 into a Kloosterman sum with modulus f and a product of Kloosterman sums with prime power moduli l_i dividing d . Then we apply Proposition 1 to the Kloosterman sum with modulus f and Proposition 3 to the Kloosterman sums with moduli l_i , taking the condition $(ab, c) = 1$ and Remark 1 into account. □

Corollary 1. *Let $a, b \in \mathbb{Z}$ and c be an odd natural number such that $c = df$, where f is squarefree and d is a powerful number with $(d, f) = 1$, and $(ab, c) = 1$. Assume that ab is a quadratic residue modulo each prime dividing d . Then*

$$|S(a, b; c)| \geq \left(\frac{f^3}{c}\right)^{1/4} \frac{2^{\omega(c)}}{(\tau(f)^2 f)^{\phi(f)/2\tau(f)}},$$

where $\phi(\cdot)$ denotes the Euler's phi function and $\tau(\cdot)$ denotes the divisor counting function. In particular, when c is squarefree, we put $d = 1$ and when c is powerful we put $f = 1$ in the above inequality.

Proof. Let $\omega(n)$ denote the number of distinct prime divisors of n . Let $\nu_p(n)$ denote the highest exponent of p appearing in the factorisation of n . The statement follows from the following observation:

$$\begin{aligned} & \left(\prod_{p|d} \frac{2}{\sqrt{p}}\right) \left(\frac{1}{(\tau(f)\sqrt{f})^{\phi(f)/\tau(f)-1}}\right) \\ & \geq \left(\prod_{p|d} 2\right) \left(\left(\prod_{p|d} \frac{1}{p^{\nu_p(c)/2}}\right) \frac{1}{\sqrt{f}}\right)^{1/2} \left(\frac{1}{(\tau(f))^{\phi(f)/\tau(f)-1}}\right) \left(\frac{1}{(\sqrt{f})^{\phi(f)/\tau(f)-3/2}}\right) \\ & \geq 2^{\omega(d)+\omega(f)} \frac{1}{c^{1/4}} \frac{1}{\tau(f)^{\phi(f)/\tau(f)}} \frac{1}{f^{\phi(f)/2\tau(f)-3/4}} = 2^{\omega(c)} \frac{1}{c^{1/4}} \frac{1}{\tau(f)^{\phi(f)/\tau(f)}} \frac{1}{f^{\phi(f)/2\tau(f)-3/4}} \\ & = \left(\frac{f^3}{c}\right)^{1/4} \frac{2^{\omega(c)}}{(\tau(f)^2 f)^{\phi(f)/2\tau(f)}}. \end{aligned}$$

□

3. AN APPLICATION

Let k and N be positive integers with k even. Recall that $S_k(N)$ denotes the vector space of cusp forms of weight k and level N . For $(n, N) = 1$, the n -th (normalised) Hecke operator acting on $S_k(N)$ is given by

$$T_n(f)(z) = n^{\frac{k-1}{2}} \sum_{ad=n, d>0} \frac{1}{d^k} \sum_{b \pmod{d}} f\left(\frac{az+b}{d}\right).$$

Let $\mathcal{F}_k(N)$ be an orthonormal basis of $S_k(N)$ consisting only of joint eigenfunctions of the Hecke operators T_n . Since $S_k(N)$ is finite dimensional, $|\mathcal{F}_k(N)|$ is finite. For $f \in S_k(N)$, the Fourier expansion of f at the cusp ∞ is given by

$$f(z) = \sum_{n=1}^{\infty} a_n(f) n^{\frac{k-1}{2}} e^{2\pi inz}.$$

We denote $\lambda_n(f)$ to be the n -th normalised Hecke eigenvalue of f , i.e. $T_n(f) = \lambda_n(f)f$. The Fourier coefficient $a_n(f)$ and $\lambda_n(f)$ are related by the condition $\overline{a_m(f)} a_n(f) = a_1(f) \lambda_n(f)$.

Petersson (see [Pet32]) expressed a weighted sum of $\overline{a_m(f)} a_n(f)$ over f with $f \in \mathcal{F}_k(N)$ in terms of Bessel function of the first kind and Kloosterman sums. Let us consider the Fourier coefficient $a_n(f)$ with harmonic weights

$$\rho_n(f) = \left(\frac{\Gamma(k-1)}{(4\pi)^{k-1}}\right)^{\frac{1}{2}} a_n(f).$$

The sum under consideration is

$$(3.1) \quad \Delta_{k,N}(m, n) = \sum_{f \in \mathcal{F}_k(N)} \overline{\rho_m(f)} \rho_n(f).$$

Putting $m = 1$ in the above equation, we have

$$\begin{aligned}\Delta_{k,N}(1, n) &= \sum_{f \in \mathcal{F}_k(N)} \overline{\rho_1(f)} \rho_n(f) \\ &= \sum_{f \in \mathcal{F}_k(N)} \left(\frac{\Gamma(k-1)}{(4\pi)^{k-1}} \right) \overline{a_1(f)} a_n(f) = \sum_{f \in \mathcal{F}_k(N)} \left(\frac{\Gamma(k-1)}{(4\pi)^{k-1}} \right) |a_1(f)|^2 \lambda_n(f),\end{aligned}$$

since $a_n(f) = a_1(f) \lambda_n(f)$. Hence,

$$(3.2) \quad \Delta_{k,N}(1, n) = \sum_{f \in \mathcal{F}_k(N)} \left(\frac{\Gamma(k-1)}{(4\pi)^{k-1}} \right) |a_1(f)|^2 \lambda_n(f).$$

The above equation demonstrates how Petersson's trace formula is indeed a weighted trace formula for Hecke operators. We use the following version of Petersson's trace formula given in [ILS00, Proposition 2.1].

Proposition 5 ([ILS00, Proposition 2.1]). *Let k, m, n, N be natural numbers with k even. We have*

$$\Delta_{k,N}(m, n) = \delta(m, n) + 2\pi i^k \sum_{N|c, c>0} \frac{S(m, n; c)}{c} J_{k-1} \left(\frac{4\pi \sqrt{mn}}{c} \right),$$

where $\delta(m, n)$ is the Kronecker diagonal symbol, $J_{k-1}(x)$ is the Bessel function of the first kind, and $S(m, n; c)$ is the classical Kloosterman sum.

We will use the following two lemmas concerning the J -Bessel function.

Lemma 3.1. *If $\nu > 0$ and $0 < x \leq 1$, we have*

$$1 \leq \frac{J_\nu(\nu x)}{x^\nu J_\nu(\nu)} \leq e^{\nu(1-x)}.$$

Proof. We refer to [Par84] for a proof. □

Lemma 3.2. *Let $\epsilon \in (0, C_0)$ be arbitrary and $A(\epsilon) := C_0 - \epsilon$, where $C_0 = \frac{2^{1/3}}{3^{2/3}\Gamma(\frac{2}{3})} = 0.4473\dots$. Then, for all $\nu \geq \left(\frac{3}{\epsilon}\right)^{3/2}$, we have*

$$(3.3) \quad \frac{A(\epsilon)}{\nu^{1/3}} \leq J_\nu(\nu) \leq \frac{C_0}{\nu^{1/3}} + \frac{3}{\nu}.$$

In particular, when $\epsilon = 0.4470$, equation (3.3) holds for all $\nu \geq 18$.

Proof. The proof follows from a direct application of [Kra14, Theorem 1] with $z = 0$, which implies that

$$(3.4) \quad J_\nu(\nu) = \frac{2^{1/3}}{\nu^{1/3}} Ai(0) + \frac{3\theta}{\nu},$$

where θ is such that $|\theta| \leq 1$, and $Ai(x)$ is the Airy function. We have $Ai(0) = \frac{1}{3^{2/3}\Gamma(\frac{2}{3})}$. Therefore,

$$-\frac{3}{\nu} \leq J_\nu(\nu) - \frac{C_0}{\nu^{1/3}} \leq \frac{3}{\nu}.$$

For all $\nu \geq \left(\frac{3}{\epsilon}\right)^{3/2}$, we have $\nu^{2/3} \geq \frac{3}{C_0 - A(\epsilon)}$, i.e., $\frac{C_0}{\nu^{1/3}} - \frac{3}{\nu} \geq \frac{A(\epsilon)}{\nu^{1/3}}$ and this proves the lower bound for $J_\nu(\nu)$. □

We give a detailed proof of the Theorem 2 after an outline of the same.

Outline of the proof of Theorem 2. We first break the infinite sum over c in Proposition 5 into two parts given by $c = N$ and $c > N$. For $c > N$, applying the integral test, we give an upper bound using the exponential bound for the J-Bessel function. This is achieved in equation (3.10). In the next step, we use Lemma 3.2 and Theorem 1 to get a lower bound for $|\Delta_{k,N}(m, n) - \delta(m, n)|$.

Proof of Theorem 2. Applying Proposition 5, breaking the c -sum into the parts with $c = N$ and $c > N$, and writing $c = bN$, we have

$$\Delta_{k,N}(m, n) = \delta(m, n) + 2\pi i^k \frac{S(m, n; N)}{N} J_{k-1}\left(\frac{4\pi\sqrt{mn}}{N}\right) + 2\pi i^k \sum_{b \geq 2} \frac{S(m, n; bN)}{bN} J_{k-1}\left(\frac{4\pi\sqrt{mn}}{bN}\right).$$

Using Lemma 3.1, we get

$$\left| J_{k-1}\left(\frac{4\pi\sqrt{mn}}{bN}\right) \right| = \left| J_{k-1}\left((k-1)\frac{4\pi\sqrt{mn}}{(k-1)bN}\right) \right| \leq e^{\nu(1-x)} x^\nu |J_\nu(\nu)|,$$

where $\nu = k-1$ and $x = \frac{4\pi\sqrt{mn}}{(k-1)bN}$. For $b \geq 2$, and $k \geq 28$, we have

$$(3.5) \quad \frac{8}{9b} \leq \frac{4\pi\sqrt{mn}}{(k-1)bN} \leq \frac{10}{9b},$$

which implies

$$e^{\nu(1-x)} x^\nu = e^{(k-1)(1-x+\log x)} \leq e^{(k-1)\left(1-\frac{8}{9b}+\log\frac{10}{9b}\right)}.$$

Therefore, using a trivial bound $|S(m, n; bN)| \leq \phi(bN)$, we obtain

$$(3.6) \quad \begin{aligned} & \left| \sum_{b \geq 2} \frac{S(m, n; bN)}{bN} J_{k-1}\left(\frac{4\pi\sqrt{mn}}{bN}\right) \right| \\ & \leq \sum_{b \geq 2} \left| J_{k-1}\left(\frac{4\pi\sqrt{mn}}{bN}\right) \right| \leq |J_{k-1}(k-1)| e^{(k-1)} \sum_{b \geq 2} e^{-(k-1)\left(\frac{8}{9b}+\log\frac{9b}{10}\right)} \\ & = |J_{k-1}(k-1)| e^{(k-1)} \left(\sum_{b=2}^4 e^{-(k-1)\left(\frac{8}{9b}+\log\frac{9b}{10}\right)} \right) + |J_{k-1}(k-1)| e^{(k-1)} \left(\sum_{b \geq 5} e^{-(k-1)\left(\frac{8}{9b}+\log\frac{9b}{10}\right)} \right). \end{aligned}$$

Since

$$(3.7) \quad e^{(k-1)} \left(\sum_{b=2}^4 e^{-(k-1)\left(\frac{8}{9b}+\log\frac{9b}{10}\right)} \right) \leq 3e^{(k-1)\left(1-\frac{4}{9}-\log\left(\frac{9}{5}\right)\right)},$$

we focus on $\sum_{b \geq 5} e^{-(k-1)\left(\frac{8}{9b}+\log\frac{9b}{10}\right)}$ only.

Let $f(x) = e^{-(k-1)\left(\frac{8}{9x}+\log\frac{9x}{10}\right)}$. Then f is decreasing for $x > 2$. Using integral test, we obtain

$$(3.8) \quad \begin{aligned} \sum_{b \geq 5} e^{-(k-1)\left(\frac{8}{9b}+\log\frac{9b}{10}\right)} & \leq \int_4^\infty e^{-(k-1)\left(\frac{8}{9x}+\log\frac{9x}{10}\right)} dx \\ & = \frac{25}{18} \int_0^{\frac{2}{9}} e^{-(k-1)y} \left(\frac{5y}{4}\right)^{k-3} dy \\ & \leq \frac{25}{18} \int_0^{\frac{2}{9}} \left(\frac{5}{18}\right)^{k-3} e^{-(k-1)y} dy \\ & = \frac{25}{18} \left(\frac{5}{18}\right)^{k-3} \left(\frac{1 - e^{-\frac{2(k-1)}{9}}}{(k-1)}\right) \leq \frac{25}{18} \left(\frac{5}{18}\right)^{k-3} \frac{1}{(k-1)}. \end{aligned}$$

Combining equations (3.6), (3.7), and (3.8), we have

$$(3.9) \quad \sum_{b \geq 2} \left| J_{k-1} \left(\frac{4\pi\sqrt{mn}}{bN} \right) \right| \leq |J_{k-1}(k-1)| \left(3e^{(k-1)(1-\frac{4}{9}-\log(\frac{9}{5}))} + \frac{25}{18}e^{(k-1)} \left(\frac{5}{18} \right)^{k-3} \frac{1}{(k-1)} \right) \\ \leq |J_{k-1}(k-1)| \left(3e^{(k-1)(1-\frac{4}{9}-\log(\frac{9}{5}))} + \frac{25}{18} \frac{e^2}{(k-1)} \left(\frac{5e}{18} \right)^{k-3} \right).$$

On considering equation (3.9) and equation (3.6), we obtain

$$\Delta_{k,N}(m, n) = \delta(m, n) + 2\pi i^k \frac{S(m, n; N)}{N} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{N} \right) + B,$$

where

$$B = 2\pi i^k \sum_{b \geq 2} \frac{S(m, n; bN)}{bN} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{bN} \right),$$

with

$$(3.10) \quad |B| \leq 2\pi |J_{k-1}(k-1)| \left(3e^{H_0(k-1)} + \frac{25}{18} \frac{e^2}{(k-1)} \left(\frac{5e}{18} \right)^{k-3} \right) \\ \leq 6.01\pi |J_{k-1}(k-1)| e^{H_0(k-1)},$$

where we use the fact that

$$\frac{25}{18} \frac{e^2}{(k-1)} \left(\frac{5e}{18} \right)^{k-3} \leq 0.005e^{H_0(k-1)},$$

for all $k \geq 22$. Since $\frac{4\pi\sqrt{mn}}{N(k-1)} \leq 1$, using Lemma 3.1 together with the fact that $J_{k-1}(k-1) > 0$ (this follows from Lemma 3.2), we obtain

$$(3.11) \quad J_{k-1} \left(\frac{4\pi\sqrt{mn}}{N} \right) \geq J_{k-1}(k-1) \left(\frac{4\pi\sqrt{mn}}{N(k-1)} \right)^{k-1} \geq D_0^{k-1} J_{k-1}(k-1).$$

Let $G(N) = \frac{f^{3/4}}{(\tau(f)^2 f)^{\phi(f)/2\tau(f)}}$, for $N = df$. Then, Corollary 1 gives

$$|S(m, n; N)| \geq 2^{\omega(N)} \frac{G(N)}{N^{1/4}},$$

Combining the previous equation with equation 3.11, we get

$$(3.12) \quad \left| \frac{S(m, n; N)}{N} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{N} \right) \right| \geq 2^{\omega(N)} \frac{G(N)}{N^{5/4}} D_0^{k-1} J_{k-1}(k-1).$$

We now find a lower bound for $|\Delta_{k,N}(m, n) - \delta(m, n)|$. For all $k \geq k_0(N)$, we have

$$k-1 \geq 1 + \left\lfloor \frac{\log 7 - \log H(N)}{\log A_0} \right\rfloor \geq \frac{\log 7 - \log H(N)}{\log A_0},$$

i.e., $(D_0 e^{-H_0})^{k-1} \geq \frac{7}{H(N)}$, and this condition implies that

$$(3.13) \quad 7e^{H_0(k-1)} J_{k-1}(k-1) \leq H(N) D_0^{k-1} J_{k-1}(k-1),$$

holds. Combining equations (3.10), (3.12) and (3.13), we obtain the following inequality for all $k \geq k_0(N)$,

$$\frac{|B|}{2\pi} \leq 3.005e^{H_0(k-1)} J_{k-1}(k-1) \leq H(N) D_0^{k-1} J_{k-1}(k-1) \leq \left| \frac{S(m, n; N)}{N} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{N} \right) \right|.$$

Therefore, for all $k \geq k_0(N)$, we have

$$\begin{aligned}
& |\Delta_{k,N}(m, n) - \delta(m, n)| \\
&= \left| 2\pi i^k \frac{S(m, n; N)}{N} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{N} \right) + B \right| \\
&\geq 2\pi \left| \left| \frac{S(m, n; N)}{N} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{N} \right) \right| - \frac{|B|}{2\pi} \right| \\
&\geq 2\pi J_{k-1}(k-1) \left| H(N) D_0^{k-1} - 3.005 e^{H_0(k-1)} \right| \\
&\geq 7.99\pi J_{k-1}(k-1) e^{H_0(k-1)},
\end{aligned}$$

since $(D_0 e^{-H_0})^{k-1} \geq \frac{7}{H(N)}$ is true for all $k \geq k_0(N)$. \square

Remark 2. We recall that Theorem 2 is a variant of [JS20, Theorem 1.7]. The bounds have been made explicit so that one can use these results for weights at least $k_0(N)$. The size of the interval is increased to size $(1 - D_0)(k - 1)$ from $2k^{\frac{1}{3}}$ of [JS20, Theorem 1.7], for a fixed N and this weakens the lower bound to $e^{(k-1)(1-\frac{4}{9}-\log(\frac{9}{5}))}$.

Corollary 2. Let $0 < \epsilon < 0.4473$. Let m, n, N be natural numbers with $(mn, N) = 1$ and N odd. Let $H_0, D_0, k_0(N)$ be as defined in Theorem 2 satisfying $(D_0 e^{-H_0})^{k-1} \geq \frac{7}{H(N)}$. Let $N = df$ where d is powerful and f is squarefree where $(d, f) = 1$. If $\frac{4\pi\sqrt{mn}}{N} \in [D_0(k-1), (k-1)]$ and mn is a quadratic residue modulo each prime dividing d , then for all $k \geq \max\left(k_0(N), 1 + \left(\frac{3}{\epsilon}\right)^{3/2}\right)$, we have

$$|\Delta_{k,N}(m, n) - \delta(m, n)| \geq 7.99\pi \frac{(0.4473 - \epsilon)}{(k-1)^{1/3}} e^{(k-1)(1-\frac{4}{9}-\log(\frac{9}{5}))}.$$

Proof. The statement follows from Lemma 3.2 and Theorem 2. \square

Remark 3. Applying Corollary 2 with its hypotheses, $\epsilon = 0.4470$, $D_0 = 0.999$, $k \geq \max(22, k_0(N))$ and $\frac{4\pi\sqrt{mn}}{N} \in [0.999(k-1), (k-1)]$, we get

$$|\Delta_{k,N}(m, n) - \delta(m, n)| \geq \frac{0.002397\pi}{(k-1)^{1/3}} e^{(k-1)(1-\frac{4}{9}-\log(\frac{9}{5}))}.$$

Corollary 3. Let k, N be natural numbers with $k \geq \max(22, k_0(N))$, k even and N odd. Let $n \geq 2$ be a natural number such that $n \in \left[\left(\frac{0.999(k-1)N}{4\pi} \right)^2, \left(\frac{(k-1)N}{4\pi} \right)^2 \right]$ with $(n, N) = 1$, and n is a quadratic residue modulo each prime dividing the powerful part of N . Then we have

$$|\Delta_{k,N}(1, n)| \geq \frac{0.002397\pi}{(k-1)^{1/3}} e^{(k-1)(1-\frac{4}{9}-\log(\frac{9}{5}))}.$$

Remark 4. Keeping equation (3.2) in mind, we note that Corollary 3 gives us a lower bound for a weighted trace of Hecke operator T_n .

Now we give a proof of the Theorem 3.

Proof of Theorem 3. The proof is similar to the proof of Theorem 2. We briefly outline the key points that need to be addressed. Since $k \geq 28$, we see that equation (3.5) is satisfied. Now using [Kra14, Theorem 1], we get

$$J_{k-1} \left((k-1) + t(k-1)^{\frac{1}{3}} \right) = \frac{2^{\frac{1}{3}} Ai(-2^{\frac{1}{3}} t)}{(k-1)^{\frac{1}{3}}} + \theta \frac{(4t^{\frac{9}{4}} + 21)}{7(k-1)},$$

where $|\theta| \leq 1$ and $0 \leq t \leq 1$. Note that for $k \geq 180$,

$$\left| \frac{\theta(4t^{\frac{9}{4}} + 21)}{7(k-1)} \right| = \left| \frac{\theta(4t^{\frac{9}{4}} + 21)}{7(k-1)^{1/3}(k-1)^{2/3}} \right| \leq \frac{25}{7 \times 179^{2/3}(k-1)^{1/3}} \leq \frac{0.113}{(k-1)^{1/3}}$$

and $2^{\frac{1}{3}}Ai(-2^{\frac{1}{3}}t) \geq 2^{\frac{1}{3}}Ai(0) \geq 0.44$, so that

$$J_{k-1} \left((k-1) + t(k-1)^{\frac{1}{3}} \right) \geq \frac{0.327}{(k-1)^{\frac{1}{3}}}.$$

The above inequality plays the role of equation (3.11) in the proof of Theorem 2. Using Lemma 3.2, $J_{k-1}(k-1) \leq \frac{0.448}{(k-1)^{\frac{1}{3}}} + \frac{0.095}{(k-1)^{\frac{1}{3}}} = \frac{0.543}{(k-1)^{\frac{1}{3}}}$. Hence, for all $k \geq k_1(N)$, we have

$$\frac{|B|}{2\pi} \leq 3.005e^{H_0(k-1)}J_{k-1}(k-1) \leq e^{H_0(k-1)}\frac{1.632}{(k-1)^{\frac{1}{3}}} \leq \frac{0.327H(N)}{(k-1)^{\frac{1}{3}}} \leq \left| \frac{S(m, n; N)}{N} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{N} \right) \right|.$$

Using equation (3.10), we obtain the following for $k \geq k_1(N)$.

$$\begin{aligned} & |\Delta_{k,N}(m, n) - \delta(m, n)| \\ &= \left| 2\pi i^k \frac{S(m, n; N)}{N} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{N} \right) + B \right| \\ &\geq 2\pi \left| \left| \frac{S(m, n; N)}{N} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{N} \right) \right| - \frac{|B|}{2\pi} \right| \\ &\geq 2\pi \left| \frac{0.327H(N)}{(k-1)^{\frac{1}{3}}} - e^{H_0(k-1)}\frac{1.632}{(k-1)^{\frac{1}{3}}} \right| \\ &\geq \frac{2\pi E_0}{(k-1)^{\frac{1}{3}}}. \end{aligned}$$

The last inequality is true if we have the following for some positive constant E_0 :

$$\frac{0.327H(N)}{(k-1)^{\frac{1}{3}}} - e^{H_0(k-1)}\frac{1.632}{(k-1)^{\frac{1}{3}}} \geq \frac{E_0}{(k-1)^{\frac{1}{3}}},$$

i.e,

$$0.327H(N) - E_0 \geq e^{H_0(k-1)}1.632,$$

i.e.

$$e^{-H_0(k-1)} \geq \frac{1.632}{0.327H(N) - E_0},$$

which is true for $k \geq k_1(N)$, where

$$k_1(N) = 2 + \left\lceil \frac{\log(1.632) - \log(0.327H(N) - E_0)}{-H_0} \right\rceil.$$

Remark 5. Note that the above theorem can be extended similarly to $\left[(k-1), (k-1) + \gamma(k-1)^{\frac{1}{3}} \right]$ with $-2^{\frac{1}{3}}\gamma > a$, where $a = -2.33811 \dots$ is the largest zero of the airy function. This forces $\gamma < 1.86$. Furthermore, it is important to note that we can not apply [Kra14, Theorem 1] directly for the interval $[2, (k-1))$ as [Kra14, Theorem 1] is valid for $t \geq 0$. □

Acknowledgements. The first named author acknowledges the support from Bhaskaracharya Pratisthan and IISER Pune. The second named author acknowledges the support of IISER Pune. The first and second named authors would like to thank Kaneenika Sinha for her valuable input. All the authors are thankful to Harish-Chandra Research Institute, Prayagraj for the hospitality while the parts of the work were carried out.

REFERENCES

- [BDS23] Baskar Balasubramanyam, Jishu Das, and Kaneenika Sinha. A discrepancy result for Hilbert modular forms. *arXiv:2307.16736v2*, pages 1–23, 2023.
- [Das23] Jishu Das. A lower bound for the discrepancy in a Sato-Tate type measure. *arXiv:2311.18798(to appear in The Ramanujan Journal)*, pages 1–14, 2023.
- [IK04] Henryk Iwaniec and Emmanuel Kowalski. *Analytic number theory*, volume 53 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2004.

- [ILS00] Henryk Iwaniec, Wenzhi Luo, and Peter Sarnak. Low lying zeros of families of L -functions. *Inst. Hautes Études Sci. Publ. Math.*, (91):55–131 (2001), 2000.
- [JS20] Junehyuk Jung and Naser Talebizadeh Sardari. Asymptotic trace formula for the Hecke operators. *Math. Ann.*, 378(1-2):513–557, 2020. With an appendix by Simon Marshall.
- [Klo27] H. D. Kloosterman. On the representation of numbers in the form $ax^2 + by^2 + cz^2 + dt^2$. *Acta Math.*, 49(3-4):407–464, 1927.
- [Kra14] Ilia Krasikov. On the Bessel function $J_\nu(x)$ in the transition region. *LMS J. Comput. Math.*, 17(1):273–281, 2014.
- [Kuz80] N. V. Kuznecov. The Petersson conjecture for cusp forms of weight zero and the Linnik conjecture. Sums of Kloosterman sums. *Mat. Sb. (N.S.)*, 111(153)(3):334–383, 479, 1980.
- [MS09] M. Ram Murty and Kaneenika Sinha. Effective equidistribution of eigenvalues of Hecke operators. *J. Number Theory*, 129(3):681–714, 2009.
- [Par84] R. B. Paris. An inequality for the Bessel function $J_\nu(\nu x)$. *SIAM J. Math. Anal.*, 15(1):203–205, 1984.
- [Pet32] Hans Petersson. Über die Entwicklungskoeffizienten der automorphen Formen. *Acta Math.*, 58(1):169–215, 1932.
- [Sel56] A. Selberg. Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series. *J. Indian Math. Soc. (N.S.)*, 20:47–87, 1956.
- [TW16] Hengcai Tang and Yingnan Wang. Quantitative versions of the joint distributions of Hecke eigenvalues. *J. Number Theory*, 169:295–314, 2016.

BHASKARACHARYA PRATISHTHANA, 56/14, OFF LAW COLLEGE ROAD, DAMLE ROAD, ERANDWANE, PUNE 411004
Email address: jewel.mahajan@students.iiserpune.ac.in

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH PUNE, DR. HOMI BHABHA ROAD, PUNE 411008
Email address: jishu.das@students.iiserpune.ac.in

RAMAKRISHNA MISSION VIVEKANANDA UNIVERSITY, BELUR MATH, HOWRAH, WEST BENGAL 711202
Email address: stephanbaier2017@gmail.com