

## An extension of Gauss's arithmetic-geometric mean (AGM) to three variables iteration scheme

Kiyoshi Sogo<sup>1, a)</sup>

*Institute of Computational Fluid Dynamics, 1-16-5, Haramachi, Meguro, Tokyo, 152-0011, Japan*

Gauss's arithmetic-geometric mean (AGM) which is described by two variables iteration  $(a_n, b_n) \rightarrow (a_{n+1}, b_{n+1})$  by  $a_{n+1} = (a_n + b_n)/2$ ,  $b_{n+1} = \sqrt{a_n b_n}$ . We extend it to three variables iteration  $(a_n, b_n, c_n) \rightarrow (a_{n+1}, b_{n+1}, c_{n+1})$  which reduces to Gauss's AGM when  $c_0 = 0$ . Our iteration starting from  $a_0 > b_0 > c_0 > 0$  with further restriction  $a_0 > b_0 + c_0$  converges to  $a_\infty = b_\infty = M(a_0, b_0, c_0)$  and  $c_\infty = 0$ . The limit  $M(a_0, b_0, c_0)$  is expressed by Appell's hyper-geometric function  $F_1(1/2, \{1/2, 1/2\}, 1; \kappa, \lambda)$  of two variables  $(\kappa, \lambda)$  which are determined by  $(a_0, b_0, c_0)$ . A relation between two hyper-geometric functions (Gauss's and Appell's) is found as a by-product.

Keywords: Gauss's arithmetic-geometric mean, three variables iteration, Appell's hyper-geometric function

---

<sup>a)</sup>EMail: sogo@icfd.co.jp

## 1. INTRODUCTION

The theory of arithmetic-geometric mean (AGM) by Gauss was published posthumously in his Werke.<sup>1</sup> Gauss observed that the iteration

$$a_{n+1} = \frac{1}{2}(a_n + b_n), \quad b_{n+1} = \sqrt{a_n b_n}, \quad (a_0 > b_0 > 0) \quad (1.1)$$

gives converging series  $a_0 > a_1 > \dots > a_\infty = b_\infty > \dots > b_1 > b_0$ . We write the limit  $a_\infty = b_\infty = \text{agM}(a_0, b_0)$  and call it arithmetic-geometric mean (AGM), which satisfies a functional equation

$$\text{agM}(a, b) = \text{agM}((a + b)/2, \sqrt{ab}). \quad (1.2)$$

The amazing discovery<sup>2</sup> by Gauss is the equality

$$\frac{a_0}{\text{agM}(a_0, b_0)} = \frac{1}{\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa u)}}, \quad \kappa = 1 - \left(\frac{b_0}{a_0}\right)^2 \quad (1.3)$$

where the right hand side integral ( $0 \leq \kappa < 1$ ) is a special case of Gauss's hyper-geometric function

$$\frac{1}{\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa u)}} = {}_2F_1\left(\frac{1}{2}, \frac{1}{2}, 1; \kappa\right), \quad (1.4)$$

which is generally defined by

$${}_2F_1(\alpha, \beta, \gamma; \kappa) = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\gamma-\alpha)} \int_0^1 u^{\alpha-1}(1-u)^{\gamma-\alpha-1}(1-\kappa u)^{-\beta} du, \quad (1.5)$$

$$= \sum_{n=0}^{\infty} \frac{(\alpha)_n(\beta)_n}{(\gamma)_n} \cdot \frac{\kappa^n}{n!}, \quad (|\kappa| < 1) \quad (1.6)$$

where  $\gamma > \alpha > 0$  is assumed, and Pochhammer's symbol  $(\alpha)_0 = 1$ ,  $(\alpha)_1 = \alpha$ ,  $(\alpha)_n = \alpha(\alpha+1)\dots(\alpha+n-1) = \Gamma(\alpha+n)/\Gamma(\alpha)$  is used. To derive this series expansion, the following Newton's expansion formula is used

$$(1-\kappa u)^{-\beta} = \sum_{m=0}^{\infty} \left(\frac{(\beta)_m}{m!} \kappa^m\right) \cdot u^m. \quad (1.7)$$

To extend the above results, we start by considering the integral with an additional new parameter  $\lambda$

$$\frac{1}{\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa u)(1-\lambda u)}}, \quad (0 < \lambda < \kappa < 1) \quad (1.8)$$

which reduces to (1.4) when  $\lambda = 0$ . In the next section we construct a model with the iteration rule  $(a_n, b_n, c_n) \rightarrow (a_{n+1}, b_{n+1}, c_{n+1})$  and show the converging limit  $M(a_0, b_0, c_0)$  is given by

$$\frac{a_0}{M(a_0, b_0, c_0)} = \frac{1}{\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa u)(1-\lambda u)}}, \quad (1.9)$$

where the relations of  $(\kappa, \lambda)$  with  $(a_0, b_0, c_0)$  are given by

$$\left(\frac{b_0}{a_0}\right)^2 = (1-\kappa)(1-\lambda), \quad \left(\frac{c_0}{a_0}\right)^2 = \kappa\lambda, \quad (1.10)$$

which will be derived in the next section.

## 2. EXTENSION OF GAUSS'S AGM

### A. Iteration rule

Let us begin to observe that the integral (1.4) of Gauss is rewritten, by setting  $u = \sin^2 \theta$ ,

$$\frac{2a_0}{\pi} \int_0^{\pi/2} \frac{d\theta}{\sqrt{a_0^2 \cos^2 \theta + b_0^2 \sin^2 \theta}} = \frac{1}{\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa u)}}, \quad \kappa = 1 - \left(\frac{b_0}{a_0}\right)^2, \quad (2.1)$$

which can be extended to

$$\begin{aligned} \frac{2a_0}{\pi} \int_0^{\pi/2} \frac{d\theta}{\sqrt{a_0^2 \cos^2 \theta + b_0^2 \sin^2 \theta - c_0^2 \cos^2 \theta \sin^2 \theta}} \\ = \frac{1}{\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa u)(1-\lambda u)}}, \end{aligned} \quad (2.2)$$

by introducing a new variable  $c_0$ . Here we have, by substituting  $\sin^2 \theta = u$ ,

$$\begin{aligned} a_0^2 \cos^2 \theta + b_0^2 \sin^2 \theta - c_0^2 \cos^2 \theta \sin^2 \theta &= a_0^2 - (a_0^2 - b_0^2 + c_0^2)u + c_0^2 u^2 \\ &= a_0^2(1-\kappa u)(1-\lambda u) \implies (1-\kappa)(1-\lambda) = \left(\frac{b_0}{a_0}\right)^2, \quad \kappa\lambda = \left(\frac{c_0}{a_0}\right)^2, \end{aligned} \quad (2.3)$$

from which we can express  $\kappa, \lambda$  in terms of  $a_0, b_0, c_0$  as follows

$$\begin{aligned} \kappa &= \frac{1}{2} \left\{ 1 - \frac{b_0^2 - c_0^2}{a_0^2} + \sqrt{\left(1 - \frac{(b_0 + c_0)^2}{a_0^2}\right) \left(1 - \frac{(b_0 - c_0)^2}{a_0^2}\right)} \right\}, \\ \lambda &= \frac{1}{2} \left\{ 1 - \frac{b_0^2 - c_0^2}{a_0^2} - \sqrt{\left(1 - \frac{(b_0 + c_0)^2}{a_0^2}\right) \left(1 - \frac{(b_0 - c_0)^2}{a_0^2}\right)} \right\}, \end{aligned} \quad (2.4)$$

and

$$\begin{aligned} 1 - \kappa &= \frac{1}{2} \left\{ 1 + \frac{b_0^2 - c_0^2}{a_0^2} - \sqrt{\left(1 - \frac{(b_0 + c_0)^2}{a_0^2}\right) \left(1 - \frac{(b_0 - c_0)^2}{a_0^2}\right)} \right\}, \\ 1 - \lambda &= \frac{1}{2} \left\{ 1 + \frac{b_0^2 - c_0^2}{a_0^2} + \sqrt{\left(1 - \frac{(b_0 + c_0)^2}{a_0^2}\right) \left(1 - \frac{(b_0 - c_0)^2}{a_0^2}\right)} \right\}, \end{aligned} \quad (2.5)$$

where conditions  $a_0 > b_0 > c_0 > 0$  and  $a_0 > b_0 + c_0$  are assumed to guarantee  $\kappa$  and  $\lambda$  take real values ( $0 < \lambda < \kappa < 1$ ). If we introduce  $\xi, \eta$  by

$$\xi = \frac{b_0 + c_0}{a_0}, \quad \eta = \frac{b_0 - c_0}{a_0}, \quad (2.6)$$

eqs. (2.4) and (2.5) are written in much simpler forms by

$$\kappa = \frac{1}{2} \left( 1 - \xi\eta + \sqrt{(1 - \xi^2)(1 - \eta^2)} \right), \quad (2.7)$$

$$\lambda = \frac{1}{2} \left( 1 - \xi\eta - \sqrt{(1 - \xi^2)(1 - \eta^2)} \right),$$

$$1 - \kappa = \frac{1}{2} \left( 1 + \xi\eta - \sqrt{(1 - \xi^2)(1 - \eta^2)} \right), \quad (2.8)$$

$$1 - \lambda = \frac{1}{2} \left( 1 + \xi\eta + \sqrt{(1 - \xi^2)(1 - \eta^2)} \right).$$

Therefore  $\kappa, \lambda$ , which are  $\kappa_0, \lambda_0$  actually, are determined by  $(a_0, b_0, c_0)$  through  $\xi, \eta$ .

To construct the iteration rule, we notice, when  $0 \leq \lambda < \kappa < 1$ ,

$$\frac{1}{\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa u)(1-\lambda u)}} = \frac{2}{\pi\sqrt{1-\lambda}} \cdot K(k_0), \quad k_0^2 = 1 - \frac{1-\kappa}{1-\lambda} \quad (2.9)$$

where  $K(k)$  is the complete elliptic integral of the first kind

$$K(k) = \int_0^{\pi/2} \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} = \frac{1}{2} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-k^2 u)}}. \quad (2.10)$$

We use here the Landen transformation formula<sup>2</sup>

$$K(k_0) = (1 + k_1) \cdot K(k_1), \quad k_1 = \frac{1 - \sqrt{1 - k_0^2}}{1 + \sqrt{1 - k_0^2}}, \quad (2.11)$$

which can be rewritten such as

$$K(k_1) = \frac{1 + \sqrt{1 - k_0^2}}{2} \cdot K(k_0), \quad (2.12)$$

because we have

$$1 + k_1 = \frac{2}{1 + \sqrt{1 - k_0^2}}. \quad (2.13)$$

Therefore our iteration rule,  $(\kappa_0, \lambda_0) \rightarrow (\kappa_1, \lambda_1)$ , is

$$\begin{aligned}
& \frac{1}{a_0\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa_0u)(1-\lambda_0u)}} = \frac{1}{a_1\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa_1u)(1-\lambda_1u)}} \\
& \implies \frac{1}{a_0\sqrt{1-\lambda_0}} \cdot K\left(k_0 = \sqrt{1 - \frac{1-\kappa_0}{1-\lambda_0}}\right) = \frac{1}{a_1\sqrt{1-\lambda_1}} \cdot K\left(k_1 = \sqrt{1 - \frac{1-\kappa_1}{1-\lambda_1}}\right) \\
& \implies \frac{1}{a_0\sqrt{1-\lambda_0}} \cdot \frac{2}{1 + \sqrt{1-k_0^2}} = \frac{1}{a_1\sqrt{1-\lambda_1}}, \tag{2.14}
\end{aligned}$$

which is rewritten into

$$\begin{aligned}
\frac{a_1}{a_0} &= \frac{1 + \sqrt{1-k_0^2}}{2} \cdot \sqrt{\frac{1-\lambda_0}{1-\lambda_1}}, & k_0^2 &= 1 - \frac{1-\kappa_0}{1-\lambda_0}, \\
&= \frac{\sqrt{1-\lambda_0} + \sqrt{1-\kappa_0}}{2\sqrt{1-\lambda_1}}. \tag{2.15}
\end{aligned}$$

Now it is convenient to introduce parameters  $\alpha$  and  $\varepsilon$  ( $0 < \varepsilon < \alpha < \pi/2$ ) by

$$\kappa = \sin^2 \alpha, \quad \lambda = \sin^2 \varepsilon, \tag{2.16}$$

which imply

$$\begin{aligned}
\frac{b}{a} &= \cos \alpha \cos \varepsilon, & \frac{c}{a} &= \sin \alpha \sin \varepsilon, \\
\frac{b+c}{a} &= \cos(\alpha - \varepsilon), & \frac{b-c}{a} &= \cos(\alpha + \varepsilon). \tag{2.17}
\end{aligned}$$

By use of these parameters, we have

$$k_0^2 = 1 - \frac{1-\kappa_0}{1-\lambda_0} = 1 - \frac{\cos^2 \alpha_0}{\cos^2 \varepsilon_0} = \frac{\sin(\alpha_0 + \varepsilon_0) \sin(\alpha_0 - \varepsilon_0)}{\cos^2 \varepsilon_0}, \tag{2.18}$$

and similarly

$$k_1^2 = 1 - \frac{1-\kappa_1}{1-\lambda_1} = \frac{\sin(\alpha_1 + \varepsilon_1) \sin(\alpha_1 - \varepsilon_1)}{\cos^2 \varepsilon_1}. \tag{2.19}$$

Substituting these into (2.13), we have

$$\begin{aligned}
\tan^2\left(\frac{\alpha_0 + \varepsilon_0}{2}\right) &= \frac{\sin(\alpha_1 + \varepsilon_1)}{\cos \varepsilon_1} = \sin \alpha_1 + \cos \alpha_1 \tan \varepsilon_1, \\
\tan^2\left(\frac{\alpha_0 - \varepsilon_0}{2}\right) &= \frac{\sin(\alpha_1 - \varepsilon_1)}{\cos \varepsilon_1} = \sin \alpha_1 - \cos \alpha_1 \tan \varepsilon_1, \tag{2.20}
\end{aligned}$$

from which we obtain

$$\begin{aligned}
\sin \alpha_1 &= \frac{1}{2} \left\{ \tan^2\left(\frac{\alpha_0 + \varepsilon_0}{2}\right) + \tan^2\left(\frac{\alpha_0 - \varepsilon_0}{2}\right) \right\}, \\
\tan^2 \varepsilon_1 &= \tan^2 \alpha_1 - \sec^2 \alpha_1 \cdot \tan^2\left(\frac{\alpha_0 + \varepsilon_0}{2}\right) \cdot \tan^2\left(\frac{\alpha_0 - \varepsilon_0}{2}\right), \tag{2.21}
\end{aligned}$$

which gives  $\kappa_1 = \sin^2 \alpha_1$  and  $\lambda_1 = \sin^2 \varepsilon_1$  in terms of  $(\alpha_0, \varepsilon_0)$ . These are the iteration scheme what we are looking for.

Here we can rewrite these quantities in terms of  $\xi, \eta$ . Since

$$\cos(\alpha_0 - \varepsilon_0) = \frac{b_0 + c_0}{a_0} = \xi, \quad \cos(\alpha_0 + \varepsilon_0) = \frac{b_0 - c_0}{a_0} = \eta, \quad (2.22)$$

we have at first

$$\tan^2 \left( \frac{\alpha_0 - \varepsilon_0}{2} \right) = \frac{1 - \xi}{1 + \xi}, \quad \tan^2 \left( \frac{\alpha_0 + \varepsilon_0}{2} \right) = \frac{1 - \eta}{1 + \eta} \quad (2.23)$$

$$\implies \sin \alpha_1 = \frac{1}{2} \left( \frac{1 - \xi}{1 + \xi} + \frac{1 - \eta}{1 + \eta} \right) = \frac{1 - \xi\eta}{(1 + \xi)(1 + \eta)}, \quad (2.24)$$

therefore we obtain

$$\kappa_1 = \sin^2 \alpha_1 = \left( \frac{1 - \xi\eta}{(1 + \xi)(1 + \eta)} \right)^2. \quad (2.25)$$

We have next, after some calculations

$$\begin{aligned} \tan^2 \varepsilon_1 &= \frac{\sin^2 \alpha_1}{1 - \sin^2 \alpha_1} - \frac{1}{1 - \sin^2 \alpha_1} \cdot \frac{(1 - \xi)(1 - \eta)}{(1 + \xi)(1 + \eta)} \\ &= \frac{(\xi - \eta)^2}{(2 + \xi + \eta)(\xi + \eta + 2\xi\eta)}, \end{aligned} \quad (2.26)$$

therefore we obtain

$$\lambda_1 = \sin^2 \varepsilon_1 = \frac{\tan^2 \varepsilon_1}{1 + \tan^2 \varepsilon_1} = \frac{(\xi - \eta)^2}{2(1 + \xi)(1 + \eta)(\xi + \eta)}. \quad (2.27)$$

In conclusion, the procedure of iteration is described as follows.

- (1) Compute  $\kappa_0, \lambda_0$  by using  $\xi = (b_0 + c_0)/a_0, \eta = (b_0 - c_0)/a_0$ ,

$$\begin{aligned} \kappa_0 &= \frac{1}{2} \left( 1 - \xi\eta + \sqrt{(1 - \xi^2)(1 - \eta^2)} \right), \\ \lambda_0 &= \frac{1}{2} \left( 1 - \xi\eta - \sqrt{(1 - \xi^2)(1 - \eta^2)} \right). \end{aligned} \quad (2.28)$$

- (2) Compute  $\kappa_1, \lambda_1$  by using the same  $\xi, \eta$ ,

$$\begin{aligned} \kappa_1 &= \left( \frac{1 - \xi\eta}{(1 + \xi)(1 + \eta)} \right)^2, \\ \lambda_1 &= \frac{(\xi - \eta)^2}{2(1 + \xi)(1 + \eta)(\xi + \eta)}. \end{aligned} \quad (2.29)$$

- (3) Compute  $(a_1, b_1, c_1)$  by using  $\kappa_0, \lambda_0$  and  $\kappa_1, \lambda_1$

$$\begin{aligned} a_1 &= a_0 (\sqrt{1 - \lambda_0} + \sqrt{1 - \kappa_0})/2\sqrt{1 - \lambda_1}, \\ b_1 &= a_1 \sqrt{(1 - \kappa_1)(1 - \lambda_1)}, \\ c_1 &= a_1 \sqrt{\kappa_1 \lambda_1}. \end{aligned} \quad (2.30)$$

We perform these sequences repeatedly, regarding the sub-indices  $(0, 1)$  as  $(n, n + 1)$ . The result of iterations is  $a_\infty = b_\infty = M(a_0, b_0, c_0)$  and  $c_\infty = 0$  as was announced before.

Let us give an example of numerical computation by Python, whose code is given in Appendix. For the case of  $a_0 = 1.0$ ,  $b_0 = 0.5$ ,  $c_0 = 0.2$ , after ten iterations we get  $a_{10} = b_{10} = 0.7250921711406717$  and  $c_{10} = 0.0$ , as is expected. On the other hand, the numerical integration of (1.9) gives  $M(a_0, b_0, c_0) = 0.725092$ , which can be regarded as the same value.

## B. Appell's hyper-geometric function

Our integration (1.8) is expressed by

$$\frac{1}{\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa u)(1-\lambda u)}} = F_1 \left( \frac{1}{2}, \left\{ \frac{1}{2}, \frac{1}{2} \right\}, 1; \kappa, \lambda \right), \quad (2.31)$$

which is a special case of Appell's hyper-geometric function of two variables defined by<sup>3</sup>

$$\begin{aligned} F_1(\alpha, \{\beta, \beta'\}, \gamma; x, y) &= \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\gamma-\alpha)} \int_0^1 u^{\alpha-1} (1-u)^{\gamma-\alpha-1} (1-xu)^{-\beta} (1-yu)^{-\beta'} du \\ &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(\alpha)_{m+n} (\beta)_m (\beta')_n}{(\gamma)_{m+n}} \cdot \frac{x^m y^n}{m! n!}, \end{aligned} \quad (2.32)$$

where is used Newton's expansion formula (1.7) twice. Appell's function  $F_1$  satisfies the differential equations<sup>3</sup>

$$\begin{aligned} x(1-x) \frac{\partial^2 F_1}{\partial x^2} + y(1-x) \frac{\partial^2 F_1}{\partial x \partial y} + [\gamma - (\alpha + \beta + 1)x] \frac{\partial F_1}{\partial x} - \beta y \frac{\partial F_1}{\partial y} - \alpha \beta F_1 &= 0, \\ y(1-y) \frac{\partial^2 F_1}{\partial y^2} + x(1-y) \frac{\partial^2 F_1}{\partial y \partial x} + [\gamma - (\alpha + \beta' + 1)y] \frac{\partial F_1}{\partial y} - \beta' x \frac{\partial F_1}{\partial x} - \alpha \beta' F_1 &= 0. \end{aligned} \quad (2.33)$$

From the equality (2.9) we have an interesting relation

$$F_1 \left( \frac{1}{2}, \left\{ \frac{1}{2}, \frac{1}{2} \right\}, 1; x, y \right) = \begin{cases} \frac{1}{\sqrt{1-y}} \cdot {}_2F_1 \left( \frac{1}{2}, \frac{1}{2}, 1; \frac{x-y}{1-y} \right), & (0 < y \leq x < 1) \\ \frac{1}{\sqrt{1-x}} \cdot {}_2F_1 \left( \frac{1}{2}, \frac{1}{2}, 1; \frac{y-x}{1-x} \right), & (0 < x \leq y < 1) \end{cases} \quad (2.34)$$

between Appell's and Gauss's hyper-geometric functions. Especially when we set  $y = x$ , the right hand side becomes  $1/\sqrt{1-x}$ , which is verified by calculating the left hand side integral.

In general such kind of relationship can be derived at least under the condition  $\gamma = \beta + \beta'$ , which is satisfied in our case ( $1 = 1/2 + 1/2$ ). Such formula is given by

$$F_1(\alpha, \{\beta, \beta'\}, \gamma; x, y) = \begin{cases} (1-y)^{-\alpha} \cdot {}_2F_1\left(\alpha, \beta, \gamma; \frac{x-y}{1-y}\right), & (0 < y \leq x < 1) \\ (1-x)^{-\alpha} \cdot {}_2F_1\left(\alpha, \beta', \gamma; \frac{y-x}{1-x}\right), & (0 < x \leq y < 1) \end{cases} \quad (2.35)$$

which is derived for  $0 < y \leq x < 1$  case by the change of integration variable  $u \rightarrow v = (1-y)u/(1-yu)$ , under the condition  $\gamma = \beta + \beta'$ .

It should be noted here that general transformation formulas are known<sup>4</sup> for much wider class of multi-variable hyper-geometric functions, where the above relation (2.35) is a special case of such formulas.

### 3. SUMMARY

The arithmetic-geometric mean by Gauss is extended to three variables iteration  $(a_n, b_n, c_n) \rightarrow (a_{n+1}, b_{n+1}, c_{n+1})$ , with  $a_\infty = b_\infty = M(a_0, b_0, c_0)$ ,  $c_\infty = 0$ . Our main result is expressed by the equalities such as

$$\frac{a_0}{M(a_0, b_0, c_0)} = \frac{2a_0}{\pi} \int_0^{\pi/2} \frac{d\theta}{\sqrt{a_0^2 \cos^2 \theta + b_0^2 \sin^2 \theta - c_0^2 \cos^2 \theta \sin^2 \theta}}, \quad (3.1)$$

$$= \frac{1}{\pi} \int_0^1 \frac{du}{\sqrt{u(1-u)(1-\kappa u)(1-\lambda u)}}, \quad (3.2)$$

$$= F_1\left(\frac{1}{2}, \left\{\frac{1}{2}, \frac{1}{2}\right\}, 1; \kappa, \lambda\right) \quad (3.3)$$

$$= \frac{1}{\sqrt{1-\lambda}} \cdot {}_2F_1\left(\frac{1}{2}, \frac{1}{2}, 1; \frac{\kappa-\lambda}{1-\lambda}\right), \quad (0 < \lambda < \kappa < 1) \quad (3.4)$$

where the relations among  $(\kappa, \lambda)$  and  $(a_0, b_0, c_0)$  are given by

$$(1-\kappa)(1-\lambda) = \left(\frac{b_0}{a_0}\right)^2, \quad \kappa\lambda = \left(\frac{c_0}{a_0}\right)^2. \quad (3.5)$$

It should be noted that the conditions  $a_0 > b_0 > c_0 > 0$  and  $a_0 > b_0 + c_0$  are assumed.



## REFERENCES

- <sup>1</sup>C.F. Gauss, *Fragmente zur Theorie des arithmetisch-geometrischen Mittels aus den Jahren 1797-1799*, edited by F. Klein, M. Brendel and L. Schlesinger, (Teubner, 1912).
- <sup>2</sup>J.M. Borwein and P.B. Borwein, *Pi and the AGM*, (Wiley, 1987).
- <sup>3</sup>E.T. Whittaker and G.N. Watson, *A Course of Modern Analysis*, (Cambridge Univ. Press, 1969) p.300.
- <sup>4</sup>T.H. Koornwinder and J.V. Stokman edit. *Encyclopedia of Special Functions: The Askey-Bateman Project, Vol. 2: Multivariable Special Functions*, (Cambridge Univ. Press, 2021) p.90.

## APPENDIX

```
# Python code of extended Gauss-agM
import math
# initial variables
a0, b0, c0=1.0, 0.5, 0.2
# iterations
for n in range(0, 10):
    print (a0, b0, c0)
    x, y=(b0+c0)/a0, (b0-c0)/a0
    dum1=1-x*y
    dum2=math.sqrt((1-x*x)*(1-y*y))
    K0=(dum1+dum2)/2
    L0=(dum1-dum2)/2
#
    dum1=1-x*y
    dum2=(1+x)*(1+y)
    K1=math.pow(dum1/dum2, 2)
    dum1=(x-y)*(x-y)
    dum2=2*(1+x)*(1+y)*(x+y)
    L1=dum1/dum2
#
    dum1=math.sqrt(1-L0)+math.sqrt(1-K0)
    dum2=2*math.sqrt(1-L1)
    a1=a0*dum1/dum2
    b1=a1*math.sqrt((1-K1)*(1-L1))
    c1=a1*math.sqrt(K1*L1)
#
    a0, b0, c0=a1, b1, c1
# end of code
```