

A different approach to positive traces on generalized q -Weyl algebras

Daniil Klyuev

Abstract Positive twisted traces are mathematical objects that could be useful in computing certain parameters of superconformal field theories. The case when \mathcal{A} is a q -Weyl algebra and ρ is a certain antilinear automorphism of \mathcal{A} was considered in [4]. Here we consider more general choices of ρ . In particular, we show that for ρ corresponding to a standard Schur index of a four-dimensional gauge theory a positive trace is unique.

1 Introduction

Let \mathcal{A} be a generalized q -Weyl algebra, it is generated by u, v, Z, Z^{-1} with relations $ZuZ^{-1} = q^2u$, $ZvZ^{-1} = q^{-2}v$, $uv = P(q^{-1}Z)$, $vu = P(qZ)$, where P is a Laurent polynomial. Let ρ be a conjugation on \mathcal{A} . We are interested in positive traces on \mathcal{A} . We assume that $0 < q < 1$.

Positive traces are defined as follows. Let \mathcal{A} be a noncommutative algebra over \mathbb{C} , ρ be an antilinear automorphism of \mathcal{A} . Let $g = \rho^2$. We say that a linear map $T: \mathcal{A} \rightarrow \mathbb{C}$ is a g -twisted trace if $T(ab) = T(bg(a))$ for all $a, b \in \mathcal{A}$. A g -twisted trace is *positive* if $T(a\rho(a)) > 0$ for all nonzero $a \in \mathcal{A}$.

Positive traces for q -Weyl algebras appear in the study of the Coulomb branch of 4-dimensional superconformal field theories [1, 3].

We will consider g -twisted traces for automorphisms g such that $g(Z) = Z$. In this case we have $T(ZaZ^{-1}) = T(a)$ for all $a \in \mathcal{A}$. This means that T is zero for any expression $u^k R(z)$ and $v^k R(z)$, where R is a Laurent polynomial and k is a positive integer. Hence T is uniquely defined by its values on Laurent polynomials.

Daniil Klyuev
Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA
02139, USA e-mail: klyuev@mit.edu

More precisely, we will consider $g = g_k$ such that $g(Z) = Z$, $g(u) = q^k Z^{-k} u$, $g(v) = q^k Z^k v$ for some integer k .

We will identify a trace T on \mathcal{A} with its restriction to $\mathbb{C}[Z, Z^{-1}] \subset \mathcal{A}$ below.

2 Traces via formal integral

Let P be a Laurent polynomial that has n nonzero roots. For simplicity, we will consider the case when P has no roots with absolute value $q^{\pm 1}$, but we expect that our methods could be modified to work in the general case.

Similarly to [4], $T: \mathbb{C}[Z, Z^{-1}] \rightarrow \mathbb{C}$ is a g_k -twisted trace if and only if $T(vuR(q^{-1}Z) - vR(Z)q^k Z^{-k}u) = 0$ for all $R \in \mathbb{C}[z, z^{-1}]$. We have

$$vR(Z)q^k Z^{-k}u = vuR(qZ)Z^{-k} = P(qZ)R(qZ)Z^{-k}.$$

Hence T is a twisted trace if and only if

$$T(P(q^{-1}Z)R(q^{-1}Z) - P(qZ)R(qZ)Z^{-k}) = 0$$

for all Laurent polynomials R .

2.1 Properties of the vector space of formal power series.

Let $V = \mathbb{C}[[z, z^{-1}]]$ be the linear space of two-sided formal power series $\sum_{i=-\infty}^{\infty} a_i z^i$. Note that V is a $\mathbb{C}[z]$ -module. Let CT be the constant term map $\sum a_i z^i \mapsto a_0$.

For a nonzero Laurent polynomial P let $P_r^{-1} \in \mathbb{C}((z))$ be its inverse in Laurent series and $P_l^{-1} \in \mathbb{C}((z^{-1}))$ be its inverse in Laurent series in the opposite direction.

Lemma 1. *The map $w(z) \mapsto w(z)P(z)$ is surjective and has kernel of dimension n . Moreover, if P has distinct roots $\alpha_1, \dots, \alpha_k$ with multiplicity m_1, \dots, m_k respectively, then the kernel is linearly spanned by the elements $f_{i,j} = \sum_{l \in \mathbb{Z}} l(l-1) \cdots (l-i) \alpha_j^{-l} z^l$, where $j = 1, \dots, k$, $i = 0, \dots, m_j - 1$.*

Proof. Let $w = \sum a_i z^i$ be an element of V . Let $w_+ = \sum_{i \geq 0} a_i z^i$, $w_- = \sum_{i < 0} a_i z^i$. Then $w_+ = (w_+ P_r^{-1})P$ and $w_- = (w_- P_l^{-1})P$, so that $w = (w_+ P_r^{-1} + w_- P_l^{-1})P$. This proves surjectivity.

We turn to the statement on the dimension of the kernel. Using the surjectivity and induction on n it is enough to consider the case $P = z - a$

for $a \in \mathbb{C}^\times$. In this case direct computation gives that the kernel is one-dimensional and spanned by $\sum a^{-i} z^i$.

It is not hard to see that the n elements in the statement of the theorem belong to the kernel and are linearly independent, hence they span the kernel.

2.2 Growth condition on traces.

Let $T: \mathbb{C}[z, z^{-1}] \rightarrow \mathbb{C}$ be a trace. Let $T(z^i) = c_i$. Define an element w of $\mathbb{C}[[z, z^{-1}]]$ by $w = \sum c_i z^{-i}$. Then $T(R(z)) = CT(R(z)w(z))$ for any Laurent polynomial R .

Let T be a positive trace with respect to a conjugation ρ such that $\rho(Z) = Z^{-1}$. Assume that $T(1) = 1$, so that $c_0 = 1$.

Lemma 2. *For all integers k we have $|T(z^k)| < 1$. Hence $|c_i| < 1$ for all $i \neq 0$.*

Proof. Consider an element $a = 1 + \alpha Z^k$. By positivity of T we have $T(a\rho(a)) > 0$. Note that

$$a\rho(a) = (1 + \alpha Z^k)(1 + \bar{\alpha} Z^{-k}) = 1 + |\alpha|^2 + \alpha Z^k + \bar{\alpha} Z^{-k}.$$

Hence

$$T(a\rho(a)) = 1 + |\alpha|^2 + \alpha T(z^k) + \bar{\alpha} T(z^{-k}).$$

Since this expression is positive for all α , we get $|T(z^k)| < 1$.

Assume that $\rho^2 = g_l$, so that T is a positive g_l -twisted trace.

Theorem 1. *The element $w(z)$ coincides with the Laurent expansion of a function $f(z)$ holomorphic on a neighborhood of the circle $|z| = 1$. The function $f(z)P(qz)$ can be analytically continued to a meromorphic function on \mathbb{C}^\times holomorphic on $1 < |z| < q^{-2}$.*

Proof. For all Laurent polynomials R we have

$$T(P(qz)R(qz) - z^l P(q^{-1}z)R(q^{-1}z)) = 0.$$

Hence

$$CT(P(qz)R(qz)w(z)) = CT(z^l P(q^{-1}z)R(q^{-1}z)w(z)).$$

It follows that

$$CT(P(z)R(z)w(q^{-1}z)) = CT(q^l z^l P(z)R(z)w(qz)),$$

hence

$$CT\left(R(z)(P(z)w(q^{-1}z) - q^l z^l P(z)w(qz))\right) = 0.$$

Since this holds for any Laurent polynomial R , we get

$$P(z)w(q^{-1}z) = q^l z^l P(z)w(qz). \quad (1)$$

Hence $w(q^{-1}z) - q^l z^l w(qz)$ belongs to the kernel of multiplication by $P(z)$. Using Lemma 1, we get

$$w(q^{-1}z) - q^l z^l w(qz) = \sum_{i,j} r_{i,j} f_{i,j}.$$

We note that the coefficients on the left-hand side have growth at most q^{-N} when N tends to $+\infty$. On the right-hand side the coefficients have growth $|\alpha_j|^N N^a$, where α_j is a root with the largest absolute value such that $f_{i,j} \neq 0$ for some i and a is a nonnegative integer. In fact, a is the largest number such that for some k we have $f_{a,k} \neq 0$ and $|\alpha_k| = |\alpha_j|$. Since there are no roots with absolute value q^{-1} , the right-hand side grows as $\alpha^N N^a$, where $|\alpha| < q^{-1}$. Using that the coefficient of z^N in $w(q^{-1}z) - q^l z^l w(qz)$ equals to $q^{-N} c_N + O(q^N)$, we get that $c_N = O(\kappa_+^N N^a)$, where $0 < \kappa_+ < 1$. Similarly, $c_{-N} = \kappa_-^N N^b$, where $0 < \kappa_- < 1$ and b is a nonnegative integer. This proves the first statement.

We turn to the second statement. Consider the expansion $P(qz)w(z)$ for the function $P(qz)f(z)$ in a neighborhood of the circle $|z| = 1$. The coefficients of $P(qz)w(z)$ decay exponentially in the negative direction by the reasoning above. It follows from (1) that $P(qz)w(z) = sq^{2l} z^l P(qz)w(q^2 z)$. Using Lemma 2 again, we see that the coefficients of z^N in $P(qz)w(z)$ is $O(q^{2N} \kappa_+^N N^a)$ when N tends to $+\infty$. It follows that $P(qz)w(z)$ gives a function holomorphic in the annulus $r_1 < |z| < r_2$, where $r_1 < 1 < q^{-2} < r_2$. Hence $f(z)$ can be analytically continued to a meromorphic function in the annulus $r_1 < |z| < r_2$. Using (1) again, we see that $f(z)$ can be analytically continued to a meromorphic function on \mathbb{C}^\times .

3 The positivity condition

Consider the antilinear automorphism ρ such that $\rho(Z) = Z^{-1}$, $\rho(u) = Z^k q^k v$, $\rho(v) = Z^{-k} q^k u$. Similarly to [4] ρ is well-defined when $P(z) = \overline{P}(z^{-1})$. We have $\rho^2 = g_{2k}$. Since we have a full classification for the case $k = 0$, assume that $k \neq 0$. The case when $k = \frac{n}{2}$ is of particular interest, since it corresponds to the ‘‘standard’’ Schur index of a four-dimensional gauge theory [2].

As in [4], Lemma 3.2, it is enough to check the positivity condition $T(a\rho(a)) > 0$ when $a \in \mathbb{C}[Z, Z^{-1}]$ or $a \in u\mathbb{C}[Z, Z^{-1}]$. The first condition translates to $T(R(z)\overline{R}(z^{-1})) > 0$ for all nonzero Laurent polynomials R , whereas the second one translates to

$$T(uR(qZ)Z^k q^k v \overline{R}(qZ^{-1})) > 0,$$

which is equivalent to

$$T(z^k P(q^{-1}z)R(q^{-1}z)\overline{R}(qz^{-1})) > 0.$$

Reasoning as in [4], Theorem 3.3, this is equivalent to $w(z)$ and $z^k P(z)w(qz)$ being nonnegative on the unit circle $|z| = 1$.

From (1) we have

$$w(q^{-1}z) = q^{2k} z^{2k} w(qz). \quad (2)$$

Let $\vartheta(z) = \vartheta_{11}(\frac{\log z}{2\pi i})$, where ϑ_{11} is a Jacobi theta function for period $\tau = \frac{\log q}{\pi i}$. Hence $\vartheta(1) = 0$. It is well-known that any meromorphic function satisfying (2) can be expressed as $w = cz^l \frac{\prod_i \vartheta(\frac{z}{\alpha_i})}{\prod_j \vartheta(\frac{z}{\beta_j})}$, where $\alpha_1, \dots, \alpha_N$ and β_1, \dots, β_M are the zeroes and poles of w . Since $w(z)P(qz)$ is holomorphic on $1 < |z| < q^{-2}$, elements $q\beta_1, \dots, q\beta_M$ are roots of P . Moreover, in order to cancel out poles, $q\beta_1, \dots, q\beta_M$ should satisfy $q < |q\beta_i| < q^{-1}$. Hence we can assume that M equals to the number of zeroes of P in the annulus $q < |z| < q^{-1}$.

We have $\vartheta(q^2 z) = q^{-1} z^{-1} \vartheta(z)$. In particular, multiplying α_1 or β_1 by a power of q^2 , we can assume that $l = 0$. Also

$$\vartheta(q^2 \frac{z}{\alpha}) = q^{-1} \alpha z^{-1} \vartheta(\frac{z}{\alpha}).$$

It follows that

$$w(q^2 z) = w(z) \frac{\prod_i (q^{-1} \alpha_i z^{-1})}{\prod_j (q^{-1} \beta_j z^{-1})} = w(z) q^{-N+M} z^{M-N} \frac{\prod_i \alpha_i}{\prod_j \beta_j}.$$

Comparing with (2) we get $N = M - 2k$, $\prod_i \alpha_i = \prod_j \beta_j q^{-N+M}$. Since $P(z) = \overline{P}(z^{-1})$ the product $\prod_j \beta_j$ has absolute value q^{-M} . Multiplying Z by a complex number with absolute value one if necessary, we can assume that $\prod_j \beta_j = q^{-M}$. We get a classification similar to [4]: possible w are parametrized by $n - 2k$ parameters α_i with fixed product q^N , and c . Note that for $k = \frac{M}{2}$ we have no numerator and the function w is unique up to scaling.

Similarly to [4], Section 3.2 the denominator of w is positive on S^1 and qS^1 . Also reasoning similarly to [4], Section 3.2, we see that when c is positive and $\alpha_1, \dots, \alpha_N$ are divided into pairs α_i and $\alpha_j = q^2 \overline{\alpha_i^{-1}}$, the numerator is positive on S^1 and qS^1 . It follows that the cone of positive traces has maximal possible real dimension $N = M - 2k$ (one if $N = 0$).

Hence we obtain the following

Theorem 2. *Let \mathcal{A} , ρ be as above. Then the dimension of the cone of positive traces is $N = M - 2k$, one if $N = 0$, and there are no positive traces if $N < 0$. In particular, when $k = \frac{n}{2}$, a positive trace exists if and only if all roots of P belong to the annulus $q < |z| < q^{-1}$, and if it exists, it is unique up to scaling.*

In particular, we confirm a conjecture of Gaiotto and Teschner [3] in the case of abelian gauge theories: if A is a K -theoretic Coulomb branch and ρ corresponds to the “standard” Schur index, then the positive trace on A , if it exists, is unique up to scaling.

Acknowledgements. I would like to thank the organizers of the LT-15 conference for the opportunity to listen to interesting talks, give a talk and write this contribution. I am thankful to Davide Gaiotto for explaining to me the physical meaning of different choices of the conjugation ρ .

References

1. M. Dedushenko and D. Gaiotto, “Algebras, traces, and boundary correlators in $\mathcal{N} = 4$ SYM,” *JHEP* **12**, 050 (2021)
2. D. Gaiotto, personal correspondence.
3. D. Gaiotto, J. Teschner, “Schur Quantization and Complex Chern-Simons theory”, arXiv:2406.09171 [hep-th]
4. D. Klyuev, “Twisted Traces and Positive Forms on Generalized q -Weyl Algebras”, *SIGMA*, 18 (2022), 009, 28 pp.