# Testing for Underpowered Literatures

Stefan Faridani\*

June 21, 2024

#### Abstract

How many experimental studies would have come to different conclusions had they been run on larger samples? I show how to estimate the expected number of statistically significant results that a set of experiments would have reported had their sample sizes all been counterfactually increased by a chosen factor. The estimator is consistent and asymptotically normal. Unlike existing methods, my approach requires no assumptions about the distribution of true effects of the interventions being studied other than continuity. This method includes an adjustment for publication bias in the reported *t*-scores. An application to randomized controlled trials (RCTs) published in top economics journals finds that doubling every experiment's sample size would only increase the power of two-sided *t*-tests by 7.2 percentage points on average. This effect is small and is comparable to the effect for systematic replication projects in laboratory psychology where previous studies enabled accurate power calculations ex ante. These effects are both smaller than for non-RCTs. This comparison suggests that RCTs are on average relatively insensitive to sample size increases. The policy implication is that grant givers should generally fund more experiments rather than fewer, larger ones.

#### Keywords: Experiments, Meta-Analysis, Statistical Power

**JEL Codes:** C01, C90

<sup>&</sup>lt;sup>\*</sup>UC San Diego. sfaridan@ucsd.edu. I thank Graham Elliott, Paul Niehaus, Xinwei Ma, Kaspar Wüthrich, Davide Viviano, Andrés Shahidinejad, Jordan van Rijn, Sabareesh Ramachandran, and Nikolay Kudrin for very helpful comments at various stages of this project.

# 1 Introduction

A key tradeoff in experimental design is balancing the risk of drawing an erroneous conclusion with project cost. Nearly all experiments published in top economics journals use *t*-tests to interpret their results. Sampling variance means that the *t*-test can fail to reject the null hypothesis of zero treatment effect even when there is in fact an effect of meaningful magnitude. Every experiment therefore runs the risk of a false negative. Increasing the sample size raises statistical power and reduces the probability of a false negative but also requires additional funding. A central question faced by funders and researchers is whether to concentrate resources into a small number of experiments or to spread them across many.

This paper provides meta-analysts and research funding organizations with a statistical procedure to estimate how much larger the expected fraction of statistically significant t-scores would have been had every experiment in some collection been counterfactually run with  $c^2$  times the sample size where c > 1. This power gain is a number between zero and one that I call  $\Delta_c$ . If this quantity is large within a given scientific literature or funding initiative, then the rejection decisions of t-tests reported by that population of experiments are sensitive on average to sample size choices. In this case, the returns to concentrating resources into fewer and larger experiments could be relatively high.

To estimate  $\Delta_c$  with the proposed method, the meta-analyst needs a dataset of n t-scores reported by a collection of experiments. The statistical procedure works by first finding the distribution of true intervention treatment effects that best fits a smoothed version of the empirical distribution of t-scores. The fitted distribution of true effects is then integrated to calculate a point estimate of  $\Delta_c$ . I show that this procedure is consistent, asymptotically normal, and converges in a power of n.

The main contribution of this paper is that it removes the reliance on functional form assumptions made by related meta-analyses that study power. The methods of Ioannidies et al. (2017), Brunner and Schimmack (2020), and others are only consistent when the population distribution of true intervention treatment effects has a specific shape. Functional form restrictions of this kind need not hold in practice. In contrast, the only assumption made by this paper on the distribution of true treatment effects is that it has a probability density function with bounded height.

A second contribution of this paper is to make its method robust to simple forms of publication bias. Recent empirical evidence shows that statistically insignificant *t*-scores are less likely to be published in academic journals (Franco et al., 2014; Andrews and Kasy, 2019). Selective reporting can distort the distribution of reported *t*-scores and potentially confound estimation. This paper addresses selective reporting by parameterizing a simple model of publication bias, estimating the model, and then reweighting the observed *t*-scores to remove publication bias. Accommodating this first stage requires that I use a particular smoothing method which results in a different estimator than those used to solve mathematically related *deconvolution* problems like Carrasco and Florens (2011).

An empirical estimate of  $\Delta_c$  is useful to meta-analysts and funders because they may otherwise lack a way to evaluate how efficiently sample sizes are being chosen in practice. Sample size decisions are typically made under high uncertainty. If funders and researchers knew ex ante how quickly the statistical power of a proposed experiment responded to its sample size, they could optimally balance power and cost. But power also depends on how effective the treatment under study actually is, a quantity which is ex ante unknown. Grant-giving organizations try to address this challenge by requiring experimenters to collect samples large enough to guarantee at least 80% power to detect a true effect larger than some chosen threshold (Doyle and Feeney, 2021). The aim of these *power calculations* is to direct resources to projects that will only fail to detect an effect when that effect is in fact small.

Power calculations may fall short of this goal in practice because they involve a great deal of guesswork. Statistical power depends on many unknown parameters besides the true effectiveness of the intervention, e.g. the degree of heteroskedasticity of the outcome variable or how heterogeneous treatment effects are across individuals. If the power calculations are inaccurate, then the choice of sample size made on their basis may trade off power and cost sub-optimally.

Even ex post it is difficult to determine what the consequences of a larger sample size would have been for a single experiment. This problem arises because it is not in general possible to infer how powerful an individual experiment was. Plugging the estimated treatment effect into the power function is known to be highly misleading (Hoenig and Heisey, 2001). Experimenters and funders therefore lack a rigorous way to assess how efficiently sample sizes are being chosen in practice. The method proposed in this paper aims to address this need.

This paper applies its method to an empirical question of broad interest: how sensitive are randomized controlled trials (RCTs) published in top economics journals to counterfactual sample size increases? I use the data from Brodeur et al. (2020) which contains t-tests of main hypotheses reported by articles published in 25 top economics journals during 2015-2018. I estimate that counterfactually doubling the sample sizes of every RCT in that set would only increase the expected number of t-scores clearing the critical value of 1.96 by 7.2 percentage points with a standard error of 2.5.

This power gain is small in comparison to several benchmarks. First, I estimate that doubling the sample size of all non-RCTs in the same dataset would increase average power by 17.3 percentage points which is significantly larger. As a second benchmark, consider a hypothetical literature where every experiment is adequately powered to detect the true effect. By conventional standards, a literature where every experiment were powered at exactly 80% power would meet this criterion (Doyle and Feeney, 2021). For such a hypothetical literature, we can calculate that doubling every sample size would increase power by 17.8 percentage points, which is much larger than the empirical estimate.

I construct a third benchmark using data from the Many Labs systematic replication project (Klein et al., 2014). In this project, 36 laboratories each independently attempted to replicate 11 published effects from laboratory psychology. We would expect these replication experiments to be very well powered because they were designed using previous experimental data and the consequences of failure to detect an already published effect could be great. Yet I cannot reject the null hypothesis that  $\Delta_c$  is the same for both RCTs in economics and the Many Labs replications. This means that RCTs are on average about as sensitive to sample size as replication experiments that were designed using a great deal of prior knowledge. The non-rejection is not simply from high uncertainty because we can indeed reject the null that  $\Delta_c$  is

the same for non-RCTs in economics vs replications from Many Labs. I also leverage the fact that the Many Labs experiments are replications to estimate power gain conditional on sample true effects. The conditional Many Labs power gain is precisely estimated at 7.8 percentage points with a standard error of 0.6 percentage points and is also statistically indistinguishable from the  $\Delta_c$  for economics RCTs.

I conclude from this analysis that randomized trials in economics are on average relatively insensitive to counterfactual sample size increases. The policy implication is that funders and researchers in economics should broadly consider running more randomized trials, not fewer, larger ones.

This paper is organized as follows. Section 2 discusses the contributions of this paper to existing applied and theoretical literatures. Section 3 sets up the estimation problem without publication bias. Section 4 shows identification, proposes an estimator, and derives its rate of consistency. Section 5 introduces publication bias. Section 6 shows asymptotic normality and discusses inference. Section 7 uses simulations to recommend tuning parameters. Section 8 presents two empirical applications and Section 9 concludes.

# 2 Related Literature

This paper contributes to two literatures. The first is an applied literature that empirically studies statistical power under strict assumptions. Ioannidies et al. (2017) estimates median power in empirical economics under the assumption that papers can be sorted into groups within which every study is estimating the same true effect. A similar assumption is used by DellaVigna and Linos (2022), Arel-Bundock et al. (2022), and Ferraro and Shukla (2023). Other methods instead assume that the distribution of true effects has a specific shape (Brunner and Schimmack, 2020; Sotola, 2023; Lang, 2023). Assumptions like these need not hold in practice and are difficult to check. This paper contributes to this literature by assuming only that the population distribution of true effects has a probability density function with bounded height.

The second related literature is technical. The estimation problem studied in this paper belongs to a class of problems called *deconvolutions* which aim to "de-blur" a smooth probability density. The theoretical econometrics literature has proposed a number of solution methods for deconvolutions (Fan, 1991; Carrasco et al., 2007; Carrasco and Florens, 2011; Racine et al., 2014). But in the presence of publication bias none of these methods will be consistent. There is growing evidence that statistically insignificant *t*-scores are less likely to be reported in economics publications (Andrews and Kasy, 2019; Elliott et al., 2022; Brodeur et al., 2020, 2016). Such omissions can create a discontinuity in the probability density of *t*-scores at the significance threshold (Gerber and Malhotra, 2008; Kudrin, 2023).

Addressing this problem is not as simple as concatenating some existing deconvolution method with publication bias removal. This paper proposes a new deconvolution step in order to manage the interactions between the two stages. The method begins with the same singular value decomposition as Carrasco and Florens (2011) but uses a different kind of regularization called *spectral cutoff* that prevents uncertainty about the extent of selective reporting from magnifying the regularization bias. This choice of smoothing method results in a new estimator that converges at a different rate.<sup>1</sup>

# 3 Setup

First define a key piece of notation. Let  $\varphi(z)$  denote the probability density function of the standard normal distribution. Adding a subscript  $\varphi_{\sigma^2}(z)$  denotes the density of the normal distribution with variance  $\sigma^2$ . The absence of the subscript means that the variance is unity.

#### 3.1 Estimand

Consider a population of experiments. Each experiment studies a unique intervention with its own treatment effect  $b \in \mathbb{R}$ . The treatment effect b is unobserved, but the experimenter estimates it with an estimator  $\hat{b}$  that is unbiased and normally distributed. This means that:  $\hat{b} \mid b \sim N(b, \sigma^2)$ . The experimenter knows the standard error  $\sigma$  and summarizes the evidence against the null hypothesis of zero treatment effect by reporting the *t*-score:  $T = \frac{\hat{b}}{\sigma}$ . Defining  $h \equiv \frac{b}{\sigma}$ , makes it possible to express the conditional distribution of T in terms of h only. I call hthe "true effect." The ratio T is conditionally normally distributed centered on h with variance 1, i.e.  $T \mid h \sim N(h, 1)$ . Its conditional probability density function  $f_T(t \mid h)$  is the following:

(1) 
$$f_T(t \mid h) = \varphi(t - h)$$

T is interpreted as the test statistic of a two-sided t-test of the null hypothesis that h = 0. The power of a size- $\alpha$  test run by an individual experiment is the conditional probability that |T| exceeds the critical value  $cv(\alpha)$ . I suppress the dependence of the critical value on  $\alpha$  for ease of notation. I call this conditional probability the *conditional power*. Conditional power can be written in terms of an integral over the normal density.

(2) 
$$\underbrace{\Pr\left(|T| > cv \mid h\right)}_{\text{Conditional Power}} = 1 - \int_{-cv}^{cv} \varphi(t-h) dt$$

Conditional power is always unknown. Since h is unobserved, the meta-analyst never gets to condition on it. In practice this means that it is not possible to recover the power of any *individual* experiment using only the *t*-score that it reported (Hoenig and Heisey, 2001). To see why, notice that Jensen's Inequality guarantees that even though E[T | h] = h, nevertheless  $E[\varphi(t-T) | h] \neq \varphi(t-h)$ . Invoking consistency and the continuous mapping theorem does not help since the limit of conditional power as the experimental sample size goes to infinity is either  $\alpha$  or 1.

The meta-analyst does not need to know the true power of any individual experiment. Instead the meta-analyst wishes to know the expected statistical power of an experiment randomly drawn from a population of experiments. I call this expectation the "unconditional power." Unconditional power depends on the distribution of h. Assumption 1, states that the population

<sup>&</sup>lt;sup>1</sup>Spectral cutoff regularization is used by Florens et al. (2017) in related but distinct setting. They estimate an inner product, assume more smoothness than we do, and equip their domain and codomain with the same inner product to construct eigenfunctions.

distribution of h is continuous with probability density function  $\pi_0$ . This density must have finite height. Assumption 1 is not as restrictive as it seems because the distribution of t-scores under a discrete  $\pi_0$  can be arbitrarily well approximated by the distribution of t-scores under a continuous  $\pi_0$ .

Assumption 1. The distribution of h is continuous with bounded probability density function  $\pi_0(h)$ .

Unconditional power is defined as the expectation of conditional power over the distribution of h. By Fubini's Theorem, the order of integration can be exchanged and unconditional power can be expressed in the following way.

(3) 
$$\underbrace{\Pr\left(|T| > cv\right)}_{\text{Unconditional Power}} = 1 - \int_{-cv}^{cv} \int_{-\infty}^{\infty} \varphi(t-h)\pi_0(h)dh \, dt$$

Experimenters have some control over the power of their experiments even though they do not know h because they can choose the sample size of the experiment. Increasing the sample size of an experiment will increase its power whenever  $h \neq 0$ . The meta-analyst wishes to learn how much larger unconditional power would have been had the sample sizes of every experiment been counterfactually increased while holding the distribution of true effects constant.

Define the random variable  $T_c$  as a *t*-score randomly drawn from a counterfactual population where every experiment has been run at  $c^2$  times the actual sample size while holding  $\pi_0$ constant. Since it is counterfactual, no draw of  $T_c$  is ever actually observed. Multiplying the sample size by  $c^2$  shrinks the standard error and grows *h* by a factor of *c*. Conditional on the true effect,  $T_c$  is normally distributed but with a larger mean, i.e.  $T_c \mid h \sim N(ch, 1)$ . The unconditional counterfactual power is defined as the probability that  $T_c$  exceeds the critical value. This can be expressed as the following double integral.

(4) 
$$\underbrace{\Pr\left(|T_c| > cv\right)}_{\text{Unconditional Counterfactual Power}} = 1 - \int_{-cv}^{cv} \int_{-\infty}^{\infty} \varphi(t - ch) \pi_0(h) dh \, dt$$

The aim of this paper is to determine whether the unconditional power of a given population of experiments is sensitive to sample size increases. This represents an "opportunity missed" because there are in expectation many false negatives that could have been avoided had the experiments been somewhat larger. The next step is to quantify this idea. Define the estimand  $\Delta_c$  as the power gain resulting from increasing every sample size in the population of experiments by a factor of  $c^2$ .

(5) 
$$\Delta_c \equiv \Pr\left(|T_c| > cv\right) - \Pr\left(|T| > cv\right)$$

 $\Delta_c$  will be the estimated throughout this paper. Example 1 illustrates why I interpret a small value of  $\Delta_c$  as an indicator of an underpowered literature. The subsequent remarks build further intuition for the estimated.

**Example 1.** The difference between status quo and counterfactual power depends on the distribution of true effects  $\pi_0$ . When a true effect is extremely small or large, power conditional

on that effect is unresponsive to sample size. So literatures that contain many extreme true effects will be less responsive than those with few extreme true effects. The following example illustrates. Consider two different literatures each with their own distribution of true effects. For both literatures, the standard error and sample size are the same with  $\sigma = 1$  and  $\sqrt{n} = 100$ . So for both literatures h = 10b. In literature 1, the intervention impact is distributed  $b \sim N(0.17, 0.1)$ . In literature 2, the intervention impact b is a 50-50 mixture of two normals, one centered on 0.02 and the other centered on 2, both with standard deviation 0.1. Figure 1 shows how unconditional power (y-axis) responds to counterfactual sample size increases (x-axis). Increasing n leads to rapid gains for literature 1 but not literature 2 because literature 2 is composed almost entirely of extremely large or small true effects. The task of this paper is to use a sample of t-scores to determine whether they were drawn from a population more like literature 1 or literature 2.

**Remark 1.** The estimand  $\Delta_c$  considers a counterfactual where the sample sizes increase but the distribution of true effects  $\pi_0$  remains constant. This framework is flexible but does have limitations. For example, if increasing the sample size of a field experiment would entail increasing spillover effects then  $\pi_0$  is different under that counterfactual and  $\Delta_c$  would not be the estimand of interest.

### 3.2 Convolution Operators

It is possible to think of the random variable T as the sum of h plus an independent "noise" random variable that has the normal distribution, i.e. T = h + Z. The sum of two independent random variables is called a *convolution* and an operator that maps the distribution of h into the distribution of h plus an noise is called a *convolution operator*. It will be useful to express the distributions of T and  $T_c$  in terms of convolution operators. The first step is to write down the densities of T and  $T_c$  as integrals over the distribution of true effects.

$$f_T(t) = \int_{-\infty}^{\infty} \varphi(t-h)\pi_0(h)dh$$
$$f_{T_c}(t) = \int_{-\infty}^{\infty} \varphi(t-ch)\pi_0(h)dh$$

Both densities are the outcomes of closely related mappings of  $\pi_0$ . This type of mapping can be generalized. Consider the operator  $K_{\sigma^2}$  below that maps the density of h to the density of  $h + \sigma Z$  where Z is a standard normal random variable independent of h where  $\sigma > 0$ . The domain of this operator will be defined later on. For now it is enough to point out that  $K_{\sigma^2}$ maps any probability density into another probability density.

$$(K_{\sigma^2}\pi)[t] \equiv \int_{-\infty}^{\infty} \varphi_{\sigma^2}(t-h)\pi(h)dh$$

A key fact about normal convolutions is that they can be decomposed. Lemma 1 says that adding normal noise of variance 1 is equivalent to adding normal noise of variance  $c^{-2}$  and then adding further independent normal noise of variance  $1 - c^{-2}$ . **Lemma 1.** For any function  $\pi$  in the domain of  $K_{c^{-2}}$ , and c > 0:

$$K_1 \pi = K_{1-c^{-2}} K_{c^{-2}} \pi$$

Proof: Appendix D.1.

This decomposition will be very useful. It is immediate to see that  $f_T = K_{1-c^{-2}}K_{c^{-2}}\pi_0$ . Lemma 2 expresses the estimand  $\Delta_c$  in terms of  $K_{c^{-2}}\pi_0$  as well. The motivation for this decomposition is that  $K_{c^{-2}}\pi_0$  turns out to be much easier to estimate than  $\pi_0$  itself because it has already been smoothed out. The next subsection sets up the theory to show why this is the case.

Lemma 2. If Assumption 1 holds, then:

$$\Delta_c = \int_{-cv}^{cv} (K_{1-c^{-2}} K_{c^{-2}} \pi_0)[t] dt - \int_{-cv/c}^{cv/c} (K_{c^{-2}} \pi_0)[t] dt$$

Proof: Appendix D.2.

#### 3.3 Singular Value Decomposition

The meta-analyst observes draws from  $f_T$  and wishes to estimate  $\Delta_c$ . To do this it is sufficient to estimate  $\pi_0$ . The problem of recovering  $\pi_0$  from  $f_T$  is severely ill-posed because very large changes in  $\pi_0$  can result in very small changes in  $f_T$ . The intuition is that applying a convolution operator K produces smoothed-out version of the argument. Since  $\pi_0$  is not necessarily smooth, many of its "high-frequency" features are destroyed by convolution and are therefore difficult to recover from  $f_T$ . However, the problem of recovering  $K_{c^{-2}}\pi_0$  from  $f_T$  is much better-posed. This target density doesn't have many high frequency features because  $K_{c^{-2}}$  has already destroyed them.

This intuition can be formalized using the singular value decomposition. Intuitively, the singular value decomposition expresses  $K_{c^{-2}}\pi_0$  as an infinite weighted sum of components of  $\pi_0$  where the weights are guaranteed to decay geometrically fast. This decomposition reveals which features of  $\pi_0$  are preserved by  $K_{c^{-2}}$  and which cannot be recovered. The singular value decompositions presented in this section are modified versions of the decompositions in Carrasco and Florens (2011) and Wand and Jones (1995).

The first task is to precisely define the domain and range of the two convolution operators  $K_{c^{-2}}$  and  $K_{1-c^{-2}}$ . The meta-analyst must start by choosing the scalar  $\sigma_Y^2 > 0$ . This could be considered a tuning parameter but in this paper I always choose  $\sigma_Y^2 = 1$  and never deviate from this choice. Define the following three Hilbert spaces of functions:  $\mathcal{L}_Y, \mathcal{L}_X, \mathcal{L}_W$ .

$$\mathcal{L}_{Y} \equiv \left\{ \phi(x) \text{ such that } \int_{-\infty}^{\infty} \phi(x)^{2} \varphi_{\sigma_{Y}^{2}}(x) dx < \infty \right\}$$
$$\mathcal{L}_{X} \equiv \left\{ \phi(x) \text{ such that } \int_{-\infty}^{\infty} \phi(x)^{2} \varphi_{\sigma_{Y}^{2}+1-c^{-2}}(x) dx < \infty \right\}$$
$$\mathcal{L}_{W} \equiv \left\{ \phi(x) \text{ such that } \int_{-\infty}^{\infty} \phi(x)^{2} \varphi_{\sigma_{Y}^{2}+1}(x) dx < \infty \right\}$$

These spaces are all very large. Each contains every function in  $\mathcal{L}^2(\mathbb{R})$  and every probability density function of a real-valued random variable. Moreover,  $\mathcal{L}_Y \subset \mathcal{L}_X \subset \mathcal{L}_W$ . Equip each space with the following integral inner products respectively:

$$\begin{split} \langle \phi_1, \phi_2 \rangle_Y &\equiv \int_{-\infty}^{\infty} \phi_1(x) \phi_2(x) \varphi_{\sigma_Y^2}(x) dx \\ \langle \phi_1, \phi_2 \rangle_X &\equiv \int_{-\infty}^{\infty} \phi_1(x) \phi_2(x) \varphi_{\sigma_Y^2 + 1 - c^{-2}}(x) dx \\ \langle \phi_1, \phi_2 \rangle_W &\equiv \int_{-\infty}^{\infty} \phi_1(x) \phi_2(x) \varphi_{\sigma_Y^2 + 1}(x) dx \end{split}$$

These inner products induce norms. So  $||\phi||_Y^2 \equiv \langle \phi, \phi \rangle_Y$ . Now the convolution operators can be fully defined by specifying their domains and the spaces that they map into.

$$K_{c^{-2}} : \mathcal{L}_W \to \mathcal{L}_X$$
  
 $K_{1-c^{-2}} : \mathcal{L}_X \to \mathcal{L}_Y$ 

Both  $K_{c^{-2}}, K_{1-c^{-2}}$  are compact linear operators. This means that they have singular value decompositions. In this case the singular value decomposition will express the outcome of a convolution as a weighted sum of known orthonormal polynomials where the weights are known to decay at an geometric rate. These decompositions are expressed below.

$$K_{c^{-2}}\pi = \sum_{j=0}^{\infty} \underbrace{\eta_{j}}_{\text{scalar}} \langle \chi_{j}, \pi \rangle_{W} \underbrace{\phi_{j}}_{\text{polynomial}}$$
$$K_{1-c^{-2}}g = \sum_{j=0}^{\infty} \underbrace{\lambda_{j}}_{\text{scalar}} \langle \phi_{j}, g \rangle_{X} \underbrace{\psi_{j}}_{\text{polynomial}}$$

The singular values  $\eta_j$ ,  $\lambda_j$  are the sequences of scalars defined below. These decay geometrically fast to zero. The fast rate of decay implies that the problem of recovering  $\pi_0$  from  $K_1\pi_0$  is ill-posed because components of  $\pi_0$  with large j play a very small role in  $K_1\pi_0$ .

$$\eta_{j} = \left(\frac{1 + \sigma_{Y}^{2} - c^{-2}}{1 + \sigma_{Y}^{2}}\right)^{j/2}$$
$$\lambda_{j} = \left(\frac{\sigma_{Y}^{2}}{\sigma_{Y}^{2} + 1 - c^{-2}}\right)^{j/2}$$

The singular functions  $\chi_j, \psi_j, \phi_i$  are the generalized probabilist's Hermite Polynomials. Since the expressions for the polynomials are very long, they are relegated to Appendix C.1. There are four important properties of the Hermite polynomials that the reader should take note of. First, the polynomials are normalized so that, for example  $\langle \chi_j, \chi_j \rangle_W = 1$ . Second, each set forms a complete basis for its corresponding Hilbert space, all of which are weighted  $\mathcal{L}^2$  spaces (Johnston, 2014). This means that for any two probability densities  $\pi_1, \pi_2$ , if  $\langle \pi_1 - \pi_2, \chi_j \rangle_X = 0$  for all j, then  $\pi_1 = \pi_2$  almost everywhere. Third, the polynomials are orthogonal, so  $\langle \chi_j, \chi_k \rangle_W = 0$  when  $j \neq k$ . Fourth, while these polynomials are themselves unbounded, they are uniformly bounded over all t, j when multiplied by the kernels from their respective inner products (Indritz, 1961). Formally this means that:  $\sup_{t,j} |\varphi_{\sigma_Y^2+1}(t)\chi_j(t)| < \infty$ .

Equation 6 below expresses the density of T as a linear combination of the Hermite polynomial. Notice that the singular values  $\lambda_j$  and  $\eta_j$  are forcing higher order polynomials to play a small role. The implication for the meta-analyst is that high-frequency information about  $\pi_0$  is not easy to recover from  $f_T$ .

(6) 
$$f_T = \sum_{j=0}^{\infty} \underbrace{\lambda_j \eta_j}_{\text{scalars}} \langle \chi_j, \pi_0 \rangle_W \underbrace{\psi_j}_{\text{polynomials}}$$

Lemma 3 expresses the target estimand  $\Delta_c$  in terms of the singular values and Hermite polynomials. This gives insight into the ill-posedness. The sequence of coefficients  $\langle \chi_j, \pi_0 \rangle_W$  is sufficient for both the distribution of the data and for the estimand  $\Delta_c$ . Equation 6 shows that the larger j is, the more difficult it is to recover  $\langle \chi_j, \pi_0 \rangle_W$  from  $f_T$  since it is damped away by the geometrically decaying  $\eta_j \lambda_j$ . But on the other hand, Lemma 3 shows that the larger j is, the smaller the weight that  $\langle \chi_j, \pi_0 \rangle_W$  is given in the estimand because  $\eta_j$  is decaying as well. This illustrates why  $\Delta_c$  is fundamentally easier to recover from  $f_T$  than  $\pi_0$ .

#### Lemma 3.

$$\Delta_c = \sum_{j=0}^{\infty} \eta_j \langle \chi_j, \pi_0 \rangle_W \left( \lambda_j \int_{-cv}^{cv} \phi_j(t) dt - \int_{-cv/c}^{cv/c} \phi_j(t) dt \right)$$

Proof: Appendix D.3.

### 4 Identification and Estimation without Publication Bias

This section shows that  $\pi_0$  is identified and proposes a consistent estimator of  $\Delta_c$  under the assumption that there is no publication bias. Econometric deconvolution problems are well studied (Fan, 1991; Carrasco et al., 2007; Carrasco and Florens, 2011; Racine et al., 2014). My estimator has an unusual motivation in that its real goal is to recover  $\Delta_c$  and it must be set up in order to be compatible with an adjustment for publication bias later on. This motivates my choice of the spectral cutoff regularization method. Spectral cutoff has two advantages in this context. First, it eliminates dependence of the smoothing parameter on c. Second, when publication bias is introduced in the next section, spectral cutoff will allow me to concentrate the objective function and estimate publication bias and  $\pi_0$  one at a time. But to avoid introducing too many ideas at once, I first show how my deconvolution method works when every t-score is reported.

#### 4.1 Identification

I prove that  $\pi_0$  is identified in the standard way by showing that it is the unique minimizer of a population objective function (Newey and McFadden, 1994). I specify the objective function  $Q_0(\pi)$  to be the integral of the squared difference between the true density  $f_T$  and the density of T implied by the argument distribution  $\pi$ . The difference is weighted by the normal density. Intuitively, minimizing this objective function is the continuous analogue of a weighted least squares regression.

(7) 
$$Q_0(\pi) \equiv \int_{-\infty}^{\infty} \left( f_T(t) - \left( \int_{-\infty}^{\infty} \varphi(t-h)\pi(h) dh \right) \right)^2 \varphi_{\sigma_Y^2}(t) dt$$

By inspection the objective is non-negative and  $Q_0(\pi_0) = 0$ . So  $\pi_0$  is a minimizer. There are several ways to show that this minimum is unique. The following paragraphs provide an argument using the singular value decomposition.<sup>2</sup> This argument is analogous to checking whether a regression matrix in a least squares problem is full rank, which can also be done using the matrix singular value decomposition.

Since the integrand is weighted by the normal density, the objective function is actually the inner product of the difference in densities with itself. This is the norm induced by the inner product  $\langle \cdot, \cdot \rangle_Y$ . The expression below makes the analogy to regression even more plain.

$$Q_0(\pi) = ||f_T - K_1\pi||_Y^2$$

This new expression says that  $\pi_0$  is the unique minimizer of  $Q_0$  if and only if convolution cannot map two different densities into the same density. The singular value decomposition reveals why this is the case. Just like in Euclidean space, norms induced by inner products can be expressed as squared sums over orthonormal basis vectors. The singular functions  $\psi_j$  form a complete orthonormal basis for  $\mathcal{L}_Y$ . The population objective function can now be written as the sum of squared differences in basis coefficients.

(8) 
$$Q_0(\pi) = \sum_{j=0}^{\infty} \left( \langle f_T, \psi_j \rangle_Y - \lambda_j \eta_j \langle \pi, \chi_j \rangle_W \right)^2$$

The final step is to substitute  $f_T = K_1 \pi_0$  and then substitute in the singular value decomposition in for  $K_1$ , keeping in mind that the  $\psi_j$  are orthonormal. The population objective function can now be expressed as the following series.

(9) 
$$Q_0(\pi) = \sum_{j=0}^{\infty} \eta_j^2 \lambda_j^2 \langle \chi_j, \pi_0 - \pi \rangle_W^2$$

The singular values  $\eta_j$ ,  $\lambda_j$  are all strictly positive even though they decay. Since all of the summands are squared real numbers,  $Q_0(\pi) = 0$  if and only if  $\langle \chi_j, \pi_0 - \pi \rangle_W = 0$  for all j. But, since the generalized Hermite polynomials  $\chi_j$  form a complete basis, any function in  $\mathcal{L}_W$  that is orthogonal to all of the  $\chi_j$  must be equal zero almost everywhere (Johnston, 2014). This means that any  $\pi$  that satisfies  $Q_0(\pi) = 0$  must equal  $\pi_0$  almost everywhere. If two probability density functions agree almost everywhere, then they are the same density. This gives identification of  $\pi_0$ , which is stated formally in Proposition 1. Its proof formalizes the argument sketched in the preceding paragraphs.

**Proposition 1.** If Assumption 1 holds, then  $Q_0(\pi)$  is uniquely minimized over  $\mathcal{L}_Y$  by  $\pi_0$ . Proof: Appendix D.4.

 $<sup>^{2}</sup>$ An alternative argument using characteristic functions instead of the singular value decomposition is provided in the subsequent Remark 2 for interested readers.

Remark 2 provides an alternative proof of identification that does not use the singular value decomposition. While the argument in the remark is more traditional and accessible, its technique does not get us much farther than identification.

**Remark 2.** Identification can also be shown with characteristic functions instead of the singular value decomposition. Briefly,  $Q_0(\pi)$  will be zero if and only if the following unweighted integral is zero.

$$\int_{-\infty}^{\infty} \left( f_T(t) - \left( \int_{-\infty}^{\infty} \varphi(t-h)\pi(h)dh \right) \right)^2 dt = 0 \iff Q_0(\pi) = 0$$

By the Plancherel Theorem, the integrand can be replaced with its Fourier transform, i.e. the difference in characteristic functions of the two distributions. By the Convolution Theorem, the characteristic function of a convolution of two densities is the pointwise product of the characteristic functions. This yields the following integral in terms of the characteristic functions  $\zeta_{\pi_0}, \zeta_{\pi}$ 

$$\int_{-\infty}^{\infty} e^{-\omega^2} \left(\zeta_{\pi_0}(\omega) - \zeta_{\pi}(\omega)\right)^2 d\omega = 0 \iff Q_0(\pi) = 0$$

In order for the integral on the left-hand side to be zero, the characteristic functions must be equal almost everywhere, which means that  $\pi_0, \pi$  are the same distribution.

#### 4.2 Estimation

The meta analyst observes a sample of n t-scores  $t_{i,k}$  indexed by  $i \in \{1, \dots n\}$  reported by m studies indexed by  $k \in \{1, \dots m\}$ . Let the number of t-scores per study be uniformly bounded by C > 0. t-scores are independent across studies, but not necessarily independent within a study. The k subscript will sometimes be suppressed. For now there is no selective reporting. The meta-analyst wishes to construct an estimate  $\hat{\pi}_n$  by computing and minimizing a sample analogue  $\hat{Q}_n(\pi)$  of the population objective  $Q_0(\pi)$  as in Newey and McFadden (1994). They then wish to use  $\hat{\pi}_n$  to compute  $\Delta_c$ .

Even though  $\pi_0$  is identified, estimating it is a severely ill-posed inverse problem because the singular values  $\lambda_j \eta_j$  decay to zero exponentially fast. This means that information about the high-frequency components of  $\pi_0$  is severely attenuated in the distribution of T. The rate at which is possible to estimate  $\pi_0$  depends on how we measure the difference between the estimated density and the true density. If we take this difference to be the  $\mathcal{L}_{\infty}$  norm, then the minimax rate is known to be logorithmic in n (Fan, 1991). This is extremely slow.

Fortunately the  $\mathcal{L}_{\infty}$  norm is too high a standard for the meta-analyst's purposes because they only need to estimate the features of  $\pi_0$  that matter for counterfactual power. I now propose a new metric that weights up the features of  $\pi_0$  that matter for the estimand and weights down features that do not matter. Let  $\rho_c(\cdot, \cdot) : \mathcal{L}_W \times \mathcal{L}_W \to \mathbb{R}$  be defined as the following:

(10) 
$$\rho_c(\pi_1, \pi_2) \equiv \sum_{j=0}^{\infty} \eta_j \left| \langle \chi_j, \pi_1 \rangle_W - \langle \chi_j, \pi_2 \rangle_W \right|$$

This is connected to  $\Delta_c$ . Lemma 4 below says that if some sequence  $\pi_n$  is converging to  $\pi_0$  at a certain rate in  $\rho_c(\cdot, \cdot)$ , then the meta-analyst can compute an estimate of  $\Delta_c$  that converges

at the same rate by plugging  $\pi_n$  into Lemma 3.

**Lemma 4.** For any random sequence  $\pi_n \subseteq \mathcal{L}_W$ , if  $\rho_c(\pi_0, \pi_n) = \mathcal{O}_p(r_n)$ , then

$$\Delta_c - \sum_{j=0}^{\infty} \eta_j \langle \chi_j, \pi_n \rangle_W \left( \lambda_j \int_{-cv}^{cv} \phi_j(t) dt - \int_{-cv/c}^{cv/c} \phi_j(t) dt \right) = \mathcal{O}_p(r_n)$$

Proof: Appendix D.5.

The next step is to propose an estimator  $\hat{\pi}_n$  for  $\pi_0$ . My estimator will be the minimizer of a sample analogue to the population objective function. There are many ways to construct the sample version of  $Q_0$ . The following method draws a direct connection to the singular value decomposition, and is easy to regularize and compute. Starting from Equation 8, notice that the inner products  $\langle f_T, \psi_j \rangle_Y$  are integrals over the probability density  $f_T$ . Integrals over probability densities are expectations. So the population objective can be expressed in terms of population moments:

(11) 
$$Q_0(\pi) = \sum_{j=0}^{\infty} \left( E\left[\psi_j(T)\varphi_{\sigma_Y^2}(T)\right] - \lambda_j \eta_j \langle \pi, \chi_j \rangle_W \right)^2$$

This immediately suggests a sample analogue to the population objective where sample averages are plugged in place of the expectations. But some care is needed. Instead of summing over all j I will regularize the sample objective by summing only up to the integer  $J_n$  which grows with n. This smoothing method is called "spectral cutoff." The sample objective  $\hat{Q}_n(\pi)$ can now be expressed as:

(12) 
$$\hat{Q}_n(\pi) = \sum_{j=0}^{J_n} \left( \frac{1}{n} \sum_{i=1}^n \psi_j(t_i) \varphi_{\sigma_Y^2}(t_i) - \lambda_j \eta_j \langle \pi, \chi_j \rangle_W \right)^2$$

The fact that the  $\chi_j$  form an orthonormal basis for  $\mathcal{L}_W$  guarantees that  $\hat{Q}_n$  can be minimized at zero. Strictly speaking, spectral cutoff guarantees that there are infinitely many sample minimizers, but this does not matter. I define my estimator  $\hat{\pi}_n$  as the minimizer of smallest norm. This is easily computed because it is a linear combination of known Hermite polynomials where the weights are sample averages.

(13) 
$$\hat{\pi}_n(h) = \sum_{j=0}^{J_n} \frac{\frac{1}{n} \sum_{i=1}^n \psi_j(t_i) \varphi_{\sigma_Y^2}(t_i)}{\eta_j \lambda_j} \chi_j(h)$$

The estimation error of  $\hat{\pi}_n$  in the metric  $\rho_c(\cdot, \cdot)$  is straightforward to compute. It is composed of two sums. The first sum is random sampling error which comes from the difference between the sample means and their expectations. The second sum is the deterministic "regularization bias" that is the price paid for cutting the sum off at  $J_n$ .

(14) 
$$\rho_c(\pi_0, \hat{\pi}_n) = \underbrace{\sum_{j=0}^{J_n} \frac{1}{\lambda_j} \left| \frac{1}{n} \sum_{i=1}^n \psi_j(t_i) \varphi_{\sigma_Y^2}(t_i) - E\left[\psi_j(T) \varphi_{\sigma_Y^2}(T)\right] \right|}_{\text{Sampling Error}} + \underbrace{\sum_{j=J_n+1}^\infty \eta_j \left| \langle \pi_0, \chi_j \rangle_W \right|}_{\text{Reg. Bias}}$$

To see why cutting the sum off at  $J_n$  is necessary, consider the variance of the terms inside the first sum. Since  $\lambda_j$  is decaying to zero, if  $J_n$  grows too quickly (or is infinite) then the variance of  $\hat{\pi}_n$  can explode. The expectation of the square of the sampling error can be bounded by studying the rate of decay of the  $\lambda_j$  and by uniformly bounding the functions  $\psi_j(t)\varphi_{\sigma_Y^2}(t)$  over all j, t (Indritz, 1961). This argument yields a bound on the sampling error.

$$\underbrace{\sum_{j=0}^{J_n} \frac{1}{\lambda_j} \left| \frac{1}{n} \sum_{i=1}^n \psi_j(t_i) \varphi_{\sigma_Y^2}(t_i) - E\left[ \psi_j(T) \varphi_{\sigma_Y^2}(T) \right] \right|}_{\text{Sampling From}} = \mathcal{O}_p\left( n^{-\frac{1}{2}} \lambda_{J_n}^{-1} \right)$$

Sampling Error

The cost of cutting the sum off at  $J_n$  is a (vanishing) regularization bias term. This bias term can be bounded using the fact that since  $\pi_0$  is a probability density function with finite height, all of its inner products can be uniformly bounded. This means that the sum of regularization bias terms is bounded by a geometric series which is the same order of its first summand.

$$\underbrace{\sum_{j=J_n+1}^{\infty} \eta_j \left| \langle \pi_0, \chi_j \rangle_W \right|}_{\text{Reg. Bias}} = \mathcal{O}\left(\eta_{J_n}\right)$$

The meta-analyst wishes to tighten the regularization parameter at a rate that maximizes the speed at which  $\rho_c(\pi_0, \hat{\pi}_n)$  converges in probability to zero. This means balancing bias and variance. Surprisingly, the rate-optimal choice of smoothing parameter does not depend on c. This is a consequence of the spectral cutoff regularization method. Theorem 1 gives the optimized rate. At this optimized rate the regularization bias is possibly of the same order as the sampling error.

**Theorem 1.** Let Assumption 1 hold and assume there is no publication bias. Then, for any constant  $\sigma_Y^2 > 0$  and any sequence  $J_n \subseteq \mathbb{N}$  chosen by the meta-analyst,

$$\rho_c(\pi_0, \hat{\pi}_n) = \mathcal{O}_p\left(n^{-\frac{1}{2}}\lambda_{J_n}^{-1} + \eta_{J_n}\right)$$

If the meta-analyst chooses  $J_n, \sigma_Y^2$  such that  $n^{1/2} \left(\frac{\sigma_Y^2}{\sigma_Y^2+1}\right)^{J_n/2}$  converges to a positive number, then:

$$\rho_c(\pi_0, \hat{\pi}_n) = \mathcal{O}_p\left(n^{-\frac{q}{2}}\right), \qquad \text{where } q \equiv \log\left(\frac{1 + \sigma_Y^2}{1 + \sigma_Y^2 - c^{-2}}\right) / \log\left(\frac{1 + \sigma_Y^2}{\sigma_Y^2}\right)$$

Proof: Appendix D.6.

The rate of convergence may seem complicated but it contains several insights. If the metaanalyst wishes to estimate  $\Delta_c$  without any sample size change, then they set c = 1. Then q = 1and the estimator achieves the parameteric rate  $n^{-1/2}$  in  $\rho_c$ . As the meta-analyst increases c, the rate of convergence slows down. This illustrates that counterfactual power resulting from a larger sample size increase is harder to estimate. The intuition is the following. The larger c is, the greater the difference between h = 0 and h close to zero. Therefore for large c, the high-frequency components of  $\pi_0$  matter more. Since the high-frequency components are harder to estimate, the rate of convergence slows down.

The meta-analyst now wishes to estimate  $\Delta_c$  by plugging  $\hat{\pi}_n$  directly into Lemma 3. I call this estimator  $\hat{\Delta}_{c,n}$ . It can be expressed compactly as a weighted sum of sample means which makes  $\hat{\Delta}_{c,n}$  fast to compute.

$$\hat{\Delta}_{c,n} = \sum_{j=0}^{J_n} \left( \frac{1}{n} \sum_{i=1}^n \psi_j(t_i) \varphi_{\sigma_Y^2}(t_i) \right) \left( \int_{-cv}^{cv} \phi_j(t) dt - \frac{1}{\lambda_j} \int_{-cv/c}^{cv/c} \phi_j(t) dt \right)$$

Combining Theorem 1 with Lemma 4, it is immediate that  $\hat{\Delta}_{c,n}$  converges to  $\Delta_c$  at the same rate as  $\hat{\pi}_n$  converges to  $\pi_0$  in the metric  $\rho_c$ .

Corollary 1. Under the same conditions as Theorem 1,

$$\Delta_c - \hat{\Delta}_{c,n} = \mathcal{O}_p \left( n^{-\frac{1}{2}} \lambda_{J_n}^{-1} + \eta_{J_n} \right)$$

**Remark 3.** One virtue of  $\hat{\Delta}_{c,n}$  is that the signs of the observations  $t_j$  do not matter. This is useful because in many meta-data sets the *t*-scores have all been reported as positive. To see why the estimator is unaffected by signs, look at the polynomial definitions in Appendix C.1 and notice that if *j* is odd then the polynomial  $\phi_j(t), \psi_j(t)$  are odd and if *j* is even then  $\phi_j(t), \psi_j(t)$  are even. Since the integrals in  $\hat{\Delta}_{c,n}$  over  $\phi_j(t)$  are symmetrical about zero, then the summands where *j* is odd are zero. For the rest of the summands, the functions  $\psi_j(t)$  are even so the signs of the data points  $t_i$  do not matter.

**Remark 4.** The researcher may wish to optimize the rate of convergence by choosing the tuning parameter  $\sigma_Y^2$  that makes q as large as possible. By L'Hopital's Rule, as  $\sigma_Y^2$  goes to infinity, qconverges to  $c^{-2}$ , making the rate approach  $n^{-\frac{1}{2c^2}}$ . But, taking  $\sigma_Y^2$  to infinity also causes the optimal  $J_n$  to grow to infinity for a fixed n, which imposes computational burdens. The choice of  $\sigma_Y^2$  is therefore a balance between rate of convergence and computability. In my simulations and empirical application I always choose  $\sigma_Y^2 = 1$  for reproducibility.

**Remark 5.** There are other methods of regularizing deconvolutions. For example Carrasco and Florens (2011) use the Tikhonov or  $||\cdot|||_2$  regularization which is the same technique used in ridge regression. My choice of the "spectral cutoff" regularization method is useful for this particular problem because when I add publication bias later on, it will allow me to concentrate the sample objective function. This would not be so straightforward for Tikhonov or other methods. Moreover, in this particular problem the smoothing parameter for spectral cutoff does not depend on *c*. This reduces the meta-analyst's researcher degrees of freedom.

# 5 Publication Bias

It is not realistic in practice to assume that every t-score computed by an experimenter is reported with equal probability. After t-scores are computed, only a subset may be published. There is growing evidence that t-scores in economics and the social sciences are selected for publication partially on the basis of whether they cross certain significance thresholds (Brodeur

et al., 2016, 2020; Andrews and Kasey, 2019; Franco et al., 2014). In this section I add publication bias to the problem. I show that  $\pi_0$  is still identified and consistently estimable by adding a term to the objective function that penalizes discontinuities in the t-curve.

#### 5.1 Identification under Publication Bias

This paper approaches publication bias by specifying a parametric model of selective reporting. Let R be the event indicating that the *t*-score T was reported. Let the conditional probability of reporting be equal to:

(15) 
$$\Pr\left(R \mid T=t\right) = w_{\theta_0}(t)$$

where the form of the function  $w_{\theta}(t)$  is known to the meta-analyst and the parameter  $\theta_0$  is an unknown member of the compact set  $\theta_0 \in C \subseteq \mathbb{R}^K$ . I make several assumptions about the publication bias model. Assumption 2 below requires that  $w_{\theta}(t)$  be bounded away from zero and from infinity. Without an assumption like this,  $\pi_0$  is not necessarily identified.

**Assumption 2.** There is a constant  $\overline{\theta} > 1$  such that  $\frac{1}{\overline{\theta}} \le w_{\theta}(t) \le \overline{\theta}$  for all  $t \in \mathbb{R}$  and all  $\theta \in C$ .

Under publication bias the meta-analyst observes only T conditional on R = 1. This is problematic because the expectations  $E[\psi_j(T)\varphi_{\sigma_Y}(T)]$  from Equation 8 are no longer directly available in population. Instead the population distribution available to the meta-analyst is the conditional distribution  $T \mid R$ . If the meta-analyst knew  $\theta_0$ , then to recover an expectation over T, they could weight the  $T \mid R$  and take the weighted expectation. Lemma 5 computes this weighting.

Lemma 5. If Assumptions 1 and 2 hold, then:

$$\mathbb{E}[\psi_{j}(T)\varphi_{\sigma_{Y}}(T)] = \mathbb{E}\left[\frac{\psi_{j}(T)\varphi_{\sigma_{Y}}(T)}{w_{\theta_{0}}(T)} \mid R\right] / E\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]$$

Proof: Appendix D.7.

The meta-analyst must estimate  $\theta_0$ . For  $\theta_0$  to be identified, publication bias must always distort the distribution of t-scores in a way that could never happen "naturally." I now add an assumption on the shape of  $w_{\theta}(t)$  to this effect. In the absence of publication bias, T must always have a continuous probability density function. Most models of publication bias specify that reporting decisions are made on the basis of statistical significance, i.e. on whether Tcrosses certain thresholds. If publication bias introduces discontinuities into the density  $f_T$ , then it is possible to identify it. Assumption 3 stipulates that publication bias "reveals itself" through discontinuities at a finite set of known points. These discontinuities are sometimes called "Caliper Gaps" and they have been well studied by others (Gerber and Malhotra, 2008; Elliott et al., 2022; Kudrin, 2023).

**Assumption 3.** There is a finite set  $\mathcal{X} \subseteq \mathbb{R}$  and a constant L > 0 such that for all  $\theta_1, \theta_2 \in C$ 

the function  $\frac{w_{\theta_1}(t)}{w_{\theta_2}(t)}$  is uniformly continuous in t on  $\mathcal{X}^c$  and

$$\sup_{t \in \mathbb{R}} \frac{1}{L} \left| \frac{1}{w_{\theta_1}(t)} - \frac{1}{w_{\theta_2}(t)} \right| \le \left| |\theta_1 - \theta_2| \right|_{\infty} \le \max_{x \in \mathcal{X}} L \lim_{\epsilon \to 0} \left| \frac{w_{\theta_1}(x+\epsilon)}{w_{\theta_2}(x+\epsilon)} - \frac{w_{\theta_1}(x-\epsilon)}{w_{\theta_2}(x-\epsilon)} \right|$$

Assumption 3 is useful for the following reason. Suppose that the meta analyst has a guess  $\theta$  for  $\theta_0$  and attempts to remove the publication bias by reweighting the density  $f_{T|R}$  by  $1/w_{\theta}$ . Then the meta-analyst can tell how close they got to  $f_T$  by checking the magnitude of the largest discontinuity left in the reweighted density. If the meta-analyst has successfully removed every discontinuity, then  $1/w_{\theta} = 1/w_{\theta_0}$  and they have recovered  $f_T$ . Example 2 shows how the "illustrative example" of publication bias from Andrews and Kasey (2019) satisfies Assumption 3. This will be the model of publication bias that I use in all of the simulations and empirical applications later on.

**Example 2.** Here is a simple parametric model of publication bias that satisfies all of the assumptions in this paper. Suppose that the probability of publication depends only on whether |T| falls above the critical value 1.96. The conditional probability ratio is equal to the scalar  $\theta_0 = \frac{Pr(R \mid |T| < 1.96)}{Pr(R \mid |T| \geq 1.96)} \in \left(\frac{1}{\overline{\theta}}, \overline{\theta}\right)$ . This means that the weighting function is:

$$w_{\theta_0}(t) = \theta_0 \mathbf{1} \{ |t| < 1.96 \} + \mathbf{1} \{ |t| \ge 1.96 \}$$

In this case the set  $\mathcal{X}$  contains just {1.96, -1.96}, the set C is  $\left(\frac{1}{\overline{\theta}}, \overline{\theta}\right)$  and the Lipschitz constant L is  $\overline{\theta}$ . This model is used by Andrews and Kasey (2019).

These assumptions are enough to identify  $\pi_0$ , and  $\theta_0$ . To show this, consider the following publication-bias-aware population objective function  $Q_0(\pi, \theta)$ . This is similar to the objective in the absence of publication bias with two differences. First, the weighted expectations conditional on R have been substituted for the unweighted unconditional expectations. Second there is now a second sum that penalizes Caliper discontinuities in the weighted density.

$$Q_{0}(\pi,\theta) \equiv \sum_{j=0}^{\infty} \left( E\left[ \frac{\psi_{j}(T)\varphi_{\sigma_{Y}}(T)}{w_{\theta_{0}}(T)} \mid R \right] / E\left[ \frac{1}{w_{\theta_{0}}(T)} \mid R \right] - \lambda_{j}\eta_{j}\langle \pi, \chi_{j} \rangle_{W} \right)^{2} + \sum_{x \in \mathcal{X}} \lim_{\epsilon \to 0} \left( \frac{f_{T \mid R}(x+\epsilon)}{w_{\theta}(x+\epsilon)} - \frac{f_{T \mid R}(x-\epsilon)}{w_{\theta}(x-\epsilon)} \right)^{2}$$

Proposition 2 says that  $\pi_0, \theta_0$  are identified when the publication bias is correctly specified. To see why, notice that Assumption 3 guarantees that the second sum is zero if and only if  $\theta = \theta_0$ . Concentrating at the true  $\theta_0$  yields  $Q_0(\pi, \theta_0)$  which is uniquely minimized over  $\pi$  by  $\pi_0$  by Proposition 1. So  $\pi_0$  and  $\theta_0$  are identified.

**Proposition 2.** If Assumptions 1, 2, and 3 hold and publication bias follows Equation 15, then  $Q_0(\pi, \theta)$  is uniquely minimized over  $\mathcal{L}_Y$  by  $\pi_0, \theta_0$ .

**Remark 6.** Andrews and Kasey (2019) identify publication bias by assuming that each study's standard error and estimand are independent. This is defensible in the context of a specific and narrow collection of studies that are all estimating similar estimands with similar methods. But

for the applications in this paper this assumption is not defensible because I examine literatures that contain highly heterogeneous studies. For such literatures, correlation between standard errors and estimands can occur in many ways, e.g. when different studies report results using different units.

**Remark 7.** The term penalizing discontinuities in derivatives can actually be dropped from  $Q_0(\pi, \theta)$  and identification can still hold. However, the regularized plug-in estimator suggested by this alternative population objective seems to perform poorly in simulation so I do not study it in this paper.

#### 5.2 Estimation under Publication Bias

The meta-analyst estimates  $\pi_0, \theta_0$  jointly by minimizing a sample analogue of the population objective function. As in the case without publication bias, the meta-analyst obtains the sample objective by plugging sample averages in place of expectations and regularizes using the spectral cutoff method by summing only over  $J_n$  terms.

The sample analogue to the penalty for Caliper discontinuities is constructed in the following way. The meta-analyst first chooses a bin width  $\epsilon_n > 0$  and computes a histogram from the sample of published *t*-scores. The height of the histogram bars is divided by  $\epsilon_n$  so that the histogram approximates the density  $f_{T|R}$ . Then the heights of the histogram bars are weighted by  $w_{\theta}(t)$  such that weighted histogram would approach continuous function as  $\epsilon_n \to 0$  if  $\theta$  were equal to  $\theta_0$ . Finally, the differences between pairs of histogram bars straddling the members of  $\mathcal{X}$  are penalized in the objective. This yields the following sample objective:

$$\hat{Q}_n(\pi,\theta) \equiv \sum_{j=0}^{J_n} \left( \frac{\frac{1}{n} \sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\theta}(t_i)}}{\frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\theta}(t_i)}} - \eta_j \lambda_j \langle \chi_j, \pi \rangle_W \right)^2 + \sum_{x \in \mathcal{X}} \left( \frac{\frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbf{1} \left\{ t_i \in (x, x + \epsilon_n] \right\}}{w_{\theta}(x + \epsilon_n)} - \frac{\frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbf{1} \left\{ t_i \in (x - \epsilon_n, x] \right\}}{w_{\theta}(x - \epsilon_n)} \right)^2$$

There is now a third tuning parameter  $\epsilon_n$  to choose. If  $\epsilon_n$  is large, then the penalty is likely to be large even when we input the correct parameter  $\theta = \theta_0$ . This may lead to substantial bias. If  $\epsilon_n$  is small, then the penalty term will have high variance and be unreliable. The optimal rate at which to decrease  $\epsilon_n$  as *n* increases turns out to be  $n^{-1/3}$  which is the same as the optimal rate for pointwise convergence of the histogram. In Section 7, I provide more specific recommendations.

The key to estimation under publication bias is that  $\pi$  can easily be concentrated out of the sample objective function. To avoid confusion with the previous estimator I call the estimator under publication bias  $\hat{\pi}_n^{pb}$ . Once again because there are infinitely many  $\pi$  that set the sample objective to zero, I define  $\hat{\pi}_n^{pb}$  as the function of smallest norm which is expressed below in terms of  $\hat{\theta}_n$ .

$$\hat{\pi}_{n}^{pb} = \frac{1}{\frac{1}{n}\sum_{i=1}^{n}\frac{1}{w_{\hat{\theta}_{n}}(t_{i})}} \sum_{j=0}^{J_{n}} \frac{\frac{1}{n}\sum_{i=1}^{n}\frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})}}{\eta_{j}\lambda_{j}}\chi_{j}$$

Since  $\pi$  has been concentrated out of the sample objective and everything but the Caliper penalty sum has been set to zero, minimization proceeds by minimizing the penalty over  $\theta$ . The sample minimizer  $\hat{\theta}_n$  expressed below can be plugged back into the previous equation to obtain  $\hat{\pi}_n^{pb}$ .

$$\hat{\theta}_n = \min_{\theta} \sum_{x \in \mathcal{X}} \left( \frac{\frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbf{1} \left\{ t_i \in (x, x + \epsilon_n] \right\}}{w_{\theta}(x + \epsilon_n)} - \frac{\frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbf{1} \left\{ t_i \in (x - \epsilon_n, x] \right\}}{w_{\theta}(x - \epsilon_n)} \right)^2$$

Theorem 2 shows the consistency of  $\hat{\pi}_n^{pb}$ . The rate of convergence has slowed down to  $n^{-q/3}$  compared to the case without publication bias. This is because a small change in the histogram near a point in x can change the weight that every  $t_i$  gets in each of the sample averages in the sample objective. The previous remarks 3, 4, and 5 also apply here.

**Theorem 2.** Let Assumptions 1, 2, and 3 hold and assume that publication bias follows Equation 15. If the meta-analyst chooses  $\epsilon_n \propto n^{-\frac{1}{3}}$ 

$$\rho_c\left(\pi_0, \hat{\pi}_n^{pb}\right) = \mathcal{O}_p\left(\lambda_{J_n}^{-1} n^{-\frac{1}{3}} + \eta_{J_n}\right)$$

If the meta-analyst also chooses  $J_n, \sigma_Y^2$  such that  $n^{1/3} \left(\frac{\sigma_Y^2}{\sigma_Y^2+1}\right)^{J_n/2}$  converges to a positive number, then:

$$\rho_c\left(\pi_0, \hat{\pi}_n^{pb}\right) = \mathcal{O}_p\left(n^{-\frac{q}{3}}\right), \qquad \text{where } q \equiv \log\left(\frac{1+\sigma_Y^2}{1+\sigma_Y^2-c^{-2}}\right) / \log\left(\frac{1+\sigma_Y^2}{\sigma_Y^2}\right)$$

Proof: Appendix D.9.

Just as in the previous section, the meta-analyst can proceed to estimate the unconditional counterfactual power by plugging  $\hat{\pi}_n$  into Lemma 3. Since the sample objective function can be concentrated, it is possible to express the estimator  $\hat{\Delta}_{c,n}^{pb}$  in terms of sample averages just as in the case without publication bias.

$$\hat{\Delta}_{c,n}^{pb} = \frac{\sum_{j=0}^{J_n} \left( \int_{-cv}^{cv} \phi_j(t) dt - \frac{1}{\lambda_j} \int_{-cv/c}^{cv/c} \phi_j(t) dt \right) \frac{1}{n} \sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\hat{\theta}_n}(t_i)}}{\frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\hat{\theta}_n}(t_i)}}$$

Combining Theorem 2 and Lemma 4, gives us the rate of consistency for  $\Delta_c$ .

**Corollary 2.** Under the same conditions as Theorem 2,

$$\Delta_c - \hat{\Delta}_{c,n}^{pb} = \mathcal{O}_p\left(n^{-\frac{q}{3}}\right)$$

### 6 Inference

Next I show asymptotic normality of  $\hat{\Delta}_{n,c}^{pb}$  and derive a consistent variance estimator. Asymptotic normality is shown using several Taylor approximations. I will need to assume that the estimator does not converge so fast that the Taylor residuals become leading terms. Since the

regularization bias can be of the same order as the sampling error, I will also need to center  $\hat{\Delta}_{n,c}^{pb}$  on a deterministic sequence  $\tilde{\Delta}_{n,c}^{pb}$  that converges to  $\Delta_c$  but is not necessarily equal to it.<sup>3</sup>

#### 6.1 Asymptotic Normality

Publication bias makes the estimator nonlinear. This is dealt with in the usual way using Taylor approximations. In order for Taylor approximations to be valid,  $w_{\theta}(t)$  needs to be sufficiently smooth in  $\theta$ . This motivates Assumption 4. It says that the gradient of  $w_{\theta}(t)$  in  $\theta$  is Lipschitz continuous in  $\theta$ .

**Assumption 4.** There is an L > 0 such that for all  $\theta_1, \theta_2 \in C$ , the  $K \times 1$  gradients of  $w_{\theta}(t)$  satisfy:

$$\left| \left| \nabla_{\theta} \frac{1}{w_{\theta_1}(t)} - \nabla_{\theta} \frac{1}{w_{\theta_2}(t)} \right| \right|_{\infty} \le L ||\theta_1 - \theta_2||_{\infty}$$

Regularization bias will affect the centering of the estimator. The centering term is nonnegligible because  $\hat{\Delta}_{n,c}^{pb}$  contains smoothing bias from two different sources. First, since only the first  $J_n$  terms of the singular value decomposition are used, the set the rest of the terms are set to zero and this incurs regularization bias. Second, since the meta-analyst only observes a sample of *t*-scores, they estimate the discontinuities in  $f_{T|R}$  by looking in a window of width  $\epsilon_n$  on either side of each possible point of discontinuity. Since the PDF can change over this interval, smoothing bias is incurred here as well.

I call the centering term  $\tilde{\Delta}_{c,n}^{pb}$ . Its expression is lengthy and is relegated to Appendix C.2. The important point for now is that  $\tilde{\Delta}_{c,n}^{pb}$  converges to  $\Delta_c$  at least as fast as  $\hat{\Delta}_{c,n}^{pb}$  converges. Lemma 6 proves this.

Lemma 6. If Assumptions 1-4 hold, then:

$$\Delta_c - \tilde{\Delta}_{c,n}^{pb} = \mathcal{O}\left(\eta_{J_n} + \epsilon_n\right)$$

If the meta-analyst chooses  $\epsilon_n = \mathcal{O}\left(n^{-1/3}\right)$  and  $n^{1/3} \left(\frac{\sigma_Y^2}{\sigma_Y^2 + 1}\right)^{J_n/2} = O(1)$ , then

$$\Delta_c - \tilde{\Delta}_{c,n}^{pb} = \mathcal{O}\left(n^{-q/3}\right)$$

Proof: Appendix D.11.

After centering properly, it is possible to linearize the estimator  $\hat{\Delta}_{c,n}^{pb}$ . Our goal is to construct a triangular array of sample means. Lemma 7 below uses Taylor's Theorem several times to rewrite the estimator as a sample mean plus a vanishing term. The function  $Z_{n,J_n}(t)$  is deterministic. Since it is lengthy to write down, its expression is relegated to Appendix C.3.

**Lemma 7.** If Assumptions 1-4 hold, and the meta-analyst chooses  $\epsilon_n \propto n^{-1/3}$ , then there exist deterministic functions  $Z_{n,J_n}$  :  $\mathbb{R} \to \mathbb{R}$  such that (i)  $\max_n \sup_{t \in \mathbb{R}} n^{-1/3} \lambda_{J_n} |Z_{n,J_n}(t)| < \infty$  and

 $<sup>^{3}</sup>$ Ongoing work aims to lower bound the variance of the estimator, allow for undersmoothing, and remove these caveats.

(ii)  $\max_n n^{-1/3} \lambda_{J_n}^2 \operatorname{Var}\left(Z_{n,J_n}(T) \mid R\right) < \infty$  and

$$\hat{\Delta}_{c,n}^{pb} - \tilde{\Delta}_{c,n} = \frac{1}{n} \sum_{i=1}^{n} Z_{n,J_n}(t_i) - \mathbb{E}\left[Z_{n,J_n}(T)\right] + \mathcal{O}_p\left(n^{-2/3}\lambda_{J_n}\right)$$

Proof: Appendix D.12.

Next I show that the triangular array of sample means  $\frac{1}{n} \sum_{i=1}^{n} Z_{n,J_n}(t_i)$  is asymptotically normal. In order for this to guarantee that the estimator  $\hat{\Delta}_{c,n}^{pb}$  is itself also normal, the sample means must dominate the Taylor residuals. To guarantee this I add the next assumption. Assumption 5 says that the estimator does not converge much faster than its guaranteed rate.

#### Assumption 5.

$$\lim\inf_{n\to\infty} n^{5/6} \lambda_{J_n}^2 \operatorname{Var}\left(\hat{\Delta}_{c,n}\right) = \infty$$

To show asymptotic normality of the triangular array of sample sums  $\frac{1}{n} \sum_{i=1}^{n} Z_{n,J_n}(t_i)$  I use a Lindeberg-Feller type Central Limit Theorem. The key step is to verify the Lyapunov Condition. While verifying this condition is a common tactic in ill-posed problems, my argument is quite different than Carrasco and Florens (2011). Using the bounds on the magnitude and variance of the  $Z_{n,J_n}$  from Lemma 7, we can verify that Assumption 5 is sufficient to guarantee the following Lyapunov Condition that uses the fourth moments.

$$\frac{\mathbb{E}\left[\left(Z_{n,J_n}(T)\right)^4\right]}{nE\left[\left(Z_{n,J_n}(T)\right)^2\right]^2} \to 0$$

Since each observation of T|R is identically distributed and independent of all but at most C other observations, the dependence is weak and the sample means are asymptotically normal. Theorem 3 shows that the assumptions so far are enough to satisfy all of the hypotheses of the central limit theorem in Theorem 2.1 of Neumann (2013).

**Theorem 3.** If Assumptions 1-5 hold and the meta-analyst chooses  $\epsilon_n \propto n^{-1/3}$ , then:

$$\frac{\hat{\Delta}_{c,n} - \tilde{\Delta}_{c,n}}{\sqrt{\operatorname{Var}(\hat{\Delta}_{c,n})}} \to_d N(0,1)$$

Proof: Appendix D.10.

#### 6.2 Variance Estimation

The next step is to derive an estimator consistent for the variance of  $\frac{1}{n} \sum_{i=1}^{n} Z_{n,J_n}(t_i)$ . Under Assumption 5, this sample mean dominates the other random terms in  $\hat{\Delta}_{n,c}^{pb}$  and it is enough to estimate its variance. In other words, Assumption 5 guarantees that the ratio of the variances converges to one:

$$\frac{\operatorname{Var}(\Delta_{n,c})}{\operatorname{Var}\left(\frac{1}{n}\sum_{i=1}^{n}Z_{n,J_n}(t_i)\right)} \to 1$$

The variance of the sample mean in the denominator is straightforward to derive. The variance of the sum is the sum of the covariances. Since two *t*-scores drawn from different studies are independent, the covariances are zero across studies. Let the symmetric  $n \times n$  adjacency matrix  $\Lambda$  equal 1 when two *t*-scores are from the same study. This gives us the following expression for the variance:

$$\operatorname{Var}\left(\frac{1}{n}\sum_{i=1}^{n} Z_{n,J_{n}}(t_{i})\right) = \frac{1}{n^{2}}\sum_{i=1}^{n}\sum_{k=1}^{n} \Lambda_{ik}\operatorname{Cov}\left(Z_{n,J_{n}}(t_{i}), Z_{nJ_{n}}(t_{k})\right)$$

Since the functions  $Z_{n,J_n}$  depend on the true  $\theta_0$  and  $\pi_0$ , the meta-analyst does not know them. But there is a sample version  $\hat{Z}_{n,J_n}$  that the meta-analyst does observe and can be used for variance estimation. Its expression is once again very lengthy and is relegated to Appendix C.3. Theorem 4 guarantees that variance estimation is consistent by proving that  $\hat{Z}_{n,J_n}(t)$ converge uniformly to  $Z_{n,J_n}(t)$  fast enough for a variance estimator to converge faster than the variance itself decays to zero.

**Theorem 4.** If Assumptions 1-5 hold and the meta-analyst chooses  $\epsilon_n \propto n^{-1/3}$  then:

$$\frac{\hat{\Delta}_{c,n} - \tilde{\Delta}_{c,n}}{\sqrt{\frac{1}{n^2} \sum_{i=1}^n \sum_{k=1}^n \Lambda_{ik} \hat{Z}_{n,J_n}(t_i) \hat{Z}_{nJ_n}(t_k)}} \to_d N(0,1)$$

Proof: Appendix D.13.

Theorem 4 says that the meta-analyst can construct valid confidence intervals covering the centering sequence  $\tilde{\Delta}_{c,n}$ . Lemma 3 showed that the centering sequence converges to  $\Delta_c$  at the same rate as the variance decays. In theory, this could affect the coverage of the confidence intervals for  $\Delta_c$  itself. A natural solution is to "undersmooth" or to change  $J_n, \epsilon_n$  more quickly than the optimal rate in order to force  $\tilde{\Delta}_{c,n}$  to converge to  $\Delta_c$  faster than the variance decays. Ongoing work aims to lower bound the variance of the estimator to allow for undersmoothing. The simulations and empirical applications in this paper choose not to undersmooth.

# 7 Simulations

To compute  $\hat{\Delta}_{n,c}^{pb}$ , the meta-analyst must choose three tuning parameters:  $\epsilon_n$ ,  $J_n$ , and  $\sigma_Y$ . To guarantee the optimized rates of convergence in Theorems 1 and 2 the meta-analyst sets  $\epsilon_n = Cn^{-1/3}$  and  $J_n = \log(Dn^{-1/3}) / \log(\sigma_Y^2 / (1 + \sigma_Y^2))$  where C, D > 0. This means that the meta-analyst's choice is actually over the triple of constants  $\{C, D, \sigma_Y\}$ . The theorems in the preceding sections provide no specific guidance on how these constants are to be chosen and we must turn to simulations.

I recommend that the meta-analyst should always at least disclose results using the following tuning parameters: C = 2,  $D = 10^{-4}$ , and  $\sigma_Y = 1$ . These choices of tuning parameters yield good confidence interval coverage of  $\Delta_c$  in simulation for a very wide variety of data generating processes. I have found that using a larger D sometimes leads to reduced coverage and that decreasing D widens confidence sets with no gain. Choosing a larger  $\sigma_Y$  increases computational costs with no obvious gain in simulation. I generate simulated data in the following way. First I use the simple model of publication bias from Example 2 where t-scores are reported with probability  $\theta_0$  if they do not clear 1.96 and are reported with certainty otherwise. This matches the illustrative example from Andrews and Kasey (2019). In the simulations in this section I set  $\theta_0 = 0.9$  but the results are not sensitive to this.

The small sample performance of the confidence intervals could depend on the distribution of true effects  $\pi_0$ . I report simulations for data generating processes where I expect coverage to be poor. Theory predicts that coverage could be low in three situations. First, if the distribution  $\pi_0$  is not very smooth, then the coefficients  $\langle \chi_j, \pi_0 \rangle$  decay slowly in j and regularization bias is large. Similarly, if  $\pi_0$  has large maximum height, then  $||\pi_0||_W$  is large which could also increase regularization bias. Third, if the density  $f_T$  is close to zero or has steep slope at the critical threshold 1.96 then  $\hat{\theta}_n$  is poor and this could in theory affect coverage of  $\Delta_c$ . I report simulations against five different distributions  $\pi_0$ , four of which meet some or all of these criteria.

- First consider a situation where every true effect is very close to zero. I call this the "Null" density and specify it to be  $h \sim \text{Unif}(-10^{-3}, 10^{-3})$ . This density is tall and non-smooth so regularization bias may be high. In addition, the density is small at 1.96.
- Second consider a case where the distribution of true effects has thick tails. Outliers are common in my empirical applications and it is useful to confirm that the confidence sets are robust to extreme t-scores. I specify h ~ Cauchy. This has low density at 1.96.
- Third consider a case where the distribution of h is bimodal. Brodeur et al. (2020) documents bimodality in the empirical distribution of t-scores in economics. For the bimodal distribution I specify that h is a 50% mixture between two normals. One normal is centered at zero and another at 2.8 and both have variance one. I choose this centering because when h = 2.8 then conditional power is close to 80% for a size 5% test which is the standard target for power calculations. This bimodal distribution models a situation where conditional power is either close to 80% or close to 5%.
- Fourth consider the distribution that brings  $\Delta_c$  close to its maximum. The "Large" distribution is  $h \sim \text{Unif}(1.96 10^{-3}, 1.96 + 10^{-3})$  and makes the estimand  $\Delta_c = 0.28$  which is large and different from the rest of the simulations. This density is tall and non-smooth so regularization bias could be high.
- The fifth distribution puts the confidence sets to a severe test. For the "Slope" distribution I set  $h \sim \text{Unif} \left(0.96 10^{-3}, 0.96 + 10^{-3}\right)$ . This increases the bias of  $\hat{\theta}_n$  by maximizing the slope of the density  $f_T$  near the critical value 1.96.

I compare performance for these five choices of  $\pi_0$  under two different meta-sample sizes, a small meta-sample of 50 *t*-scores versus a large meta-sample of 1000 *t*-scores.

The simulation results are reported in Table 1. The first column reports the meta-sample size n. The second column reports the distribution of true effects  $\pi_0$  that was used to generate the data. The third and fourth columns report unconditional power and the power gain  $\Delta_c$  from a counterfactual doubling of every sample size in the population of experiments. This

means that  $c = \sqrt{2}$ . The fifth column reports the mean over 2000 simulation repetitions of the estimate  $\hat{\Delta}_{c,n}$ . The sixth and seventh columns compare the standard deviation of the estimator and the average standard error. The final column reports coverage of  $\Delta_c$  by the 95% confidence set centered on  $\hat{\Delta}_{c,n}$ . Coverage is close to 95% for all of the  $\pi_0$  for both large and small n. This is not necessarily the case for other choices of D and C.

# 8 Empirical Applications

I apply  $\hat{\Delta}_{c,n}^{pb}$  to two real world settings. First I present an empirical exercise using data from a multi-site replication study in laboratory psychology (Klein et al., 2014). This setting is special because counterfactual power can be credibly and precisely estimated by other means. I find that the confidence sets produced by my method contain a credible alternative point estimate. This exercise provides evidence that this novel method can perform well with real data. Second I present an empirical application where I use my method to answer a broad question of practical interest: are randomized controlled trials published in top economics journals underpowered? My method provides evidence of the opposite: the power of *t*-tests reported by RCTs published in these outlets are on average quite insensitive to sample size increases.

#### 8.1 Replication Studies in Laboratory Psychology

While the simulations in the previous section show good coverage for the confidence intervals against a variety of artificial data generating processes, I cannot know from those results how realistic the simulations are or how well the method performs when it is used on real world data. In this section I present an exercise that takes place in a special empirical environment where it is possible to construct a credible and precisely estimated alternative point estimate for  $\Delta_c$  that I can benchmark my confidence sets against.

The Many Labs systematic replication project is an ideal environment for this exercise. Klein et al. (2014) recruited 36 independent research teams who each attempted to replicate 13 effects from experimental psychology. Each team collected their own independent data and ran some or all of the thirteen experiments on their respective samples. Every sample contained at least 80 participants and many contained far more for a total of 6344. Eleven of the experiments were analyzed using t-tests of the equality in means between a treatment group and a control group and I will limit my analysis to these. This yields a meta-sample of 385 t-scores. The aim of all of the experiments run by Many Labs was to replicate effects that had been published in top psychology journals and to investigate how consistently effects could be replicated across study sites.

This environment is special because it is plausible to assume that all of the research teams were investigating the same (or very similar) true effects. A key conclusion reported by Klein et al. (2014) is that variation in the true effect size across study sites was found to be very small compared to the variation in the effect sizes across the experimental treatments. This finding is plausible because the experiments took place in controlled laboratory environments, the researchers were all using the same protocols, and the primary aim of every experiment was to replicate existing results consistently. Given that the research teams had no incentive

to selectively report their results, it is also reasonable to assume that there was no publication bias in this setting.

Assuming that the true effects did not vary across study sites makes it possible to estimate  $\Delta_c^{COND}$ , which is the power gain conditional on sample realizations of h.

(16) 
$$\Delta_c^{(n)} \equiv \frac{1}{n} \sum_{i=1}^n \left[ \Pr\left( |T_c| > 1.96 \mid h = h_i \right) - \Pr\left( |T| > 1.96 \mid h = h_i \right) \right]$$

Notice that  $E\left[\Delta_c^{(n)}\right] = \Delta_c$  and in large samples,  $\Delta_c^{(n)} \to \Delta_c$ . Even though  $\Delta_c^{(n)}$  is not identical to  $\Delta_c$  in every sample, its meaning and interpretation are similar enough to use as a benchmark.

To estimate  $\Delta_c^{(n)}$  we proceed as follows. For each of the 11 experimental treatments, I take the mean of the reported effect sizes across the 36 study sites to estimate the true effect b for that treatment. Then I replace each of the 386 *t*-scores with the ratio  $\frac{b}{s_i}$  where  $s_i$  is the standard error used to compute the *i*th *t*-score. To compute unconditional power, I plug each ratio  $\frac{b}{s_i}$ into the power function for the size 5% *t*-test and take the mean. To compute power were the sample sizes all to have been counterfactually doubled, I plug  $\sqrt{2}\frac{b}{s_i}$  into the power function instead. Taking the average yields the point estimate:  $\hat{\Delta}_c^{(n)} = 0.078$ . This estimation technique is a simplified and unweighted version of those used by Ioannidies et al. (2017); Arel-Bundock et al. (2022) and employs the same basic assumption. Computing the standard error under the conservative worst-case assumption that two experiments in the same lab have covariance 1 yields standard error 0.006. The alternative point estimate  $\hat{\Delta}_c^{(n)}$  is quite precise.

I test the coverage of the confidence sets proposed by this paper by checking whether they contain  $\hat{\Delta}_c^{(n)}$ . Table 2 presents the results. The third row of Table 2 reports the 95% confidence interval in square brackets. The confidence interval contains the benchmark value 0.078. This exercise complements the simulations by arguing that inference is likely to work reasonably well in a real-world environment.

To show that the results are not sensitive to reasonable choices of the tuning parameters, I present four specifications. In columns 1 and 2 I scale  $J_n$ ,  $\epsilon_n$  by the number of experimental study sites. In columns 3 and 4 I instead scale the tuning parameters by the total number of *t*-scores. In columns 1 and 3 I use the recommended values of the constants C, D from Table 1. In columns 2 and 4 I vary these choices to show that the confidence sets do not change much even when the changes to C, D are large.

 $\Delta_c^{(n)}$  is not identical to  $\Delta_c$  except in large samples. Nevertheless I argue that the benchmark is still very useful because it demonstrates that the confidence intervals produce plausible results in a well-understood empirical environment.

### 8.2 Randomized Controlled Trials in Economics

I apply  $\hat{\Delta}_{c,n}^{pb}$  to answer an empirical question with policy implications: Are randomized controlled trials published in top economics journals underpowered? Experiments are lauded as the "gold standard" of empirical evidence in social science because their design allows them to credibly control the rate of type-I errors. But what about type-II errors? If the conclusions reported by influential RCTs are sensitive to reasonable increases in sample size, then funders and researchers should consider running experiments with fewer treatment arms and allocate more resources toward data collection.

This is an empirical question and the answer will depend on the population of experiments under study. Here I study the experiments that have the most influence in academic economics: those published in top journals. The data source is Brodeur et al. (2020). This meta-study examined the universe of 684 articles published in top economics journals during 2015-2018. The data contain 21,740 test statistics of which 20,419 were t-scores that I can use. All test statistics corresponded to main hypotheses of interest and excluded regression controls, robustness checks, placebo tests, and the like. Every t-score was produced using one of four empirical methods: Randomized Controlled Trials (RCTs), Difference in Differences (DID), Discontinuity Designs (DD), and Instrumental Variables (IV).

I estimate the sensitivity of tests conducted by RCTs to hypothetical sample size increases in Table 3. The estimate in column 1 says that counterfactually doubling the sample sizes of every RCT published in top economics journals would only increase the expected number of *t*-scores clearing the critical value of 1.96 by 7.2 percentage points with standard error 2.5.

This result can be interpreted by comparing it to other empirical estimates of  $\Delta_c$ . First I return to the Many Labs result from the previous section as our first benchmark. The Many Labs  $\Delta_c$  should be very low because Many Labs consisted of deliberate laboratory replications of previously published results and used samples chosen to ensure adequate power. Since experiments published in top economics journals are usually novel in some way, the researcher knows less in the design phase and the sample size choice is less likely to be optimal.

Table 3 shows that I fail to reject the null hypothesis that  $\Delta_c$  is equal across economics RCTs and the Many Labs replications (p = 0.18). This is not simply a consequence of high uncertainty because the confidence intervals involved are small enough to make other meaningful comparisons. Column 2 of Table 3 provides an example. It reports that among tests conducted by non-RCTs, doubling sample sizes would increase power by 17.3 percentage points which is significantly different from Many Labs (p = 0.00).

These results are not very sensitive to the choice of tuning parameters. Columns 3 and 4 of Table 3 show a robustness check where the tuning parameters  $J, \epsilon$  are scaled by the number of articles instead of the number of t-scores. These confidence intervals are smaller but at greater risk of bias. The results are largely unchanged. The only comparison that changes is that RCTs and the Many Labs  $\Delta_c$  become marginally significantly different (p = 0.04). This may be because scaling by number of articles makes confidence intervals less likely to cover. Otherwise the results are qualitatively unchanged.

Figures 2, 3, and 4 visualize these comparisons over many possible sample size increases  $c^2$ . Figure 2 shows how power climbs much faster for non-experiments than RCTs in the Brodeur et al. (2016) sample even when we increase sample sizes by less than double. Figures 3 and 4 show that the Many Labs experiments respond at a pace indistinguishable from the economics RCTs. These figures reinforce and visualize the results from the tables.

As a final benchmark, consider a literature where every experiment is run at exactly 80% power. Then we can calculate that doubling every sample size would increase power by 17.8 percentage points. We can easily reject the null hypothesis that RCTs in the Brodeur et al.

(2016) data are this sensitive.

I conclude from this exercise that RCTs published in top economics journals are relatively insensitive to counterfactual sample size increases. The policy implication is that funders and researchers should generally consider devoting more resources to running more experiments or adding treatment arms rather than raising sample size standards.

There is one important caveat to mention. The estimand  $\Delta_c$  is the change in unconditional power and is therefore an average over many highly heterogeneous studies. This estimand likely masks important heterogeneity. Future research could use the method in this paper to identify sub-populations of experiments that are relatively sensitive to counterfactual sample size increases.

# 9 Conclusion

This paper proposed an estimator consistent for the fraction of t-scores that would have been statistically significant had every experiment in a given collection had its sample size counterfactually increased by a chosen factor c > 1. The only assumption imposed on the distribution of true effects was that it had a probability density function with bounded height. The proposed estimator is asymptotically normal and robust to simple forms of publication bias. The key insight of the paper was that even though it is not feasible to estimate the distribution of true effects  $\pi_0$  it is feasible to estimate enough components of  $\pi_0$  to recover counterfactual power. The key technical step was to use the spectral cutoff regularization method to concentrate the publication bias out of the sample objective function.

I provided tuning parameter recommendations that yielded good small sample coverage for the confidence sets across many simulations. An empirical exercise using the Many Labs dataset confirmed that the confidence interval contained a point estimate computed using an alternative method that the Many Labs environment made possible. In a second empirical application I found that the power of randomized trials in economics would only increase by 7.2 percentage points on average if every sample size had been doubled. I argued that this number is small. The policy implication is that funding organizations should in general fund more randomized trials rather than fewer, larger ones.

# References

- Andrews, Isaiah and Maximilian Kasey, "Identification of and Correction for Publication Bias," American Economic Review, August 2019, 109 (8), 2766–94.
- and Maximilian Kasy, "Identification of and Correction for Publication Bias," American Economic Review, August 2019, 109 (8), 2766–94.
- Arel-Bundock, Vincent, Ryan C. Briggs, Hristos Doucouliagos, Marco Mendoza Aviña, and Tom D. Stanley, "Quantitative Political Science Research is Greatly Underpowered," I4R Discussion Paper Series 6, The Institute for Replication (I4R) 2022.
- **Bateman, Harry**, *Higher Transcendental Functions [Volumes I-III]*, McGraw-Hill Book Company, 1953. Funding by Office of Naval Research (ONR).
- Brodeur, Abel, Mathias Lé, Marc Sangnier, and Yanos Zylberberg, "Star Wars: The Empirics Strike Back," American Economic Journal: Applied Economics, January 2016, 8 (1), 1–32.
- \_ , Nikolai Cook, and Anthony Heyes, "Methods Matter: p-Hacking and Publication Bias in Causal Analysis in Economics," *American Economic Review*, November 2020, 110 (11), 3634–60.
- Brunner, Jerry and Ulrich Schimmack, "Estimating Population Mean Power Under Conditions of Heterogeneity and Selection for Significance," *Meta-Psychology*, 05 2020, 4.
- Carrasco, Marine and Jean-Pierre Florens, "A spectral method for deconvolving a density," *Econometric Theory*, 2011, 27 (3), 546–581.
- \_ , \_ , and Eric Renault, in James J. Heckman and Edward E. Leamer, eds., Chapter 77 Linear Inverse Problems in Structural Econometrics Estimation Based on Spectral Decomposition and Regularization, Vol. 6 of Handbook of Econometrics, Elsevier, 2007, pp. 5633–5751.
- **DellaVigna, Stefano and Elizabeth Linos**, "RCTs to Scale: Comprehensive Evidence From Two Nudge Units," *Econometrica*, 2022, *90* (1), 81–116.
- Doyle, M and L Feeney, "Quick guide to power calculations," 2021.
- Elliott, Graham, Nikolay Kudrin, and Kaspar Wüthrich, "Detecting p-Hacking," *Econometrica*, 2022, 90 (2), 887–906.
- Fan, Jianqing, "On the Optimal Rates of Convergence for Nonparametric Deconvolution Problems," The Annals of Statistics, 1991, 19 (3), 1257 1272.
- Ferraro, Paul J. and Pallavi Shukla, "Credibility crisis in agricultural economics," Applied Economic Perspectives and Policy, 2023, 45 (3), 1275–1291.
- Florens, Jean-Pierre, Joël L. Horowitz, and Ingrid Van Keilegom, "Bias-Corrected Confidence Intervals in a Class of Linear Inverse Problems," Annals of Economics and Statistics, 2017, (128), 203–228.

- Franco, Annie, Neil Malhotra, and Gabor Simonovits, "Publication bias in the social sciences: Unlocking the file drawer," *Science*, 2014, 345 (6203), 1502–1505.
- Gerber, Alan and Neil Malhotra, "Do Statistical Reporting Standards Affect What Is Published? Publication Bias in Two Leading Political Science Journals," *Quarterly Journal* of Political Science, 2008, 3 (3), 313–326.
- Hoenig, John M and Dennis M Heisey, "The Abuse of Power," *The American Statistician*, 2001, 55 (1), 19–24.
- Indritz, Jack, "An Inequality for Hermite Polynomials," Proceedings of the American Mathematical Society, 1961, 12 (6), 981–983.
- Ioannidies, John P. A., T. D. Stanley, and Hristos Doucouliagos, "The Power of Bias in Economics Research," *The Economic Journal*, October 2017, 127, F236–F265.
- Johnston, William, "The Weighted Hermite Polynomials Form a Basis for L2(R)," *The American Mathematical Monthly*, 2014, 121 (3), pp. 249–253.
- Klein, Richard A., Kate A. Ratliff, Michelangelo Vianello, Reginald B. Adams, Štěpán Bahník, Michael J. Bernstein, Konrad Bocian, Mark J. Brandt, Beach Brooks, Claudia Chloe Brumbaugh, Zeynep Cemalcilar, Jesse Chandler, Winnee Cheong, William E. Davis, Thierry Devos, Matthew Eisner, Natalia Frankowska, David Furrow, Elisa Maria Galliani, Fred Hasselman, Joshua A. Hicks, James F. Hovermale, S. Jane Hunt, Jeffrey R. Huntsinger, Hans IJzerman, Melissa-Sue John, Jennifer A. Joy-Gaba, Heather Barry Kappes, Lacy E. Krueger, Jaime Kurtz, Carmel A. Levitan, Robyn K. Mallett, Wendy L. Morris, Anthony J. Nelson, Jason A. Nier, Grant Packard, Ronaldo Pilati, Abraham M. Rutchick, Kathleen Schmidt, Jeanine L. Skorinko, Robert Smith, Troy G. Steiner, Justin Storbeck, Lyn M. Van Swol, Donna Thompson, A. E. van 't Veer, Leigh Ann Vaughn, Marek Vranka, Aaron L. Wichman, Julie A. Woodzicka, and Brian A. Nosek, "Investigating Variation in Replicability," Social Psychology, 2014, 45 (3), 142–152.
- Kudrin, Nikolay, "Robust Caliper Tests," Working Paper, 2023.
- Lang, Kevin, "How Credible is the Credibility Revolution?," Working Paper 31666, National Bureau of Economic Research September 2023.
- Neumann, Michael H., "A central limit theorem for triangular arrays of weakly dependent random variables, with applications in statistics," *ESAIM: Probability and Statistics*, 2013, 17, 120–134.
- Newey, Whitney K. and Daniel McFadden, in "Chapter 36 Large sample estimation and hypothesis testing," Vol. 4 of *Handbook of Econometrics*, Elsevier, 1994, pp. 2111–2245.
- Racine, Jeffrey S., Liangjun Su, and Aman Ullah, The Oxford Handbook of Applied Nonparametric and Semiparametric Econometrics and Statistics, Oxford University Press, 02 2014.

Sotola, Lukas, "How Can I Study from Below, that which Is Above?: Comparing Replicability Estimated by Z-Curve to Real Large-Scale Replication Attempts," *Meta-Psychology*, 09 2023, 7.

Wand, Matt P. and M. Chris Jones, Kernel Smoothing 1995.

# A Tables

Table 1: Simulations										
Parameters					Results					
n	$\pi_0$	Unc. Pwr.	$\Delta_c$	Mean $\hat{\Delta}_{c,n}$	$\mathrm{SD}\ \hat{\Delta}_{c,n}$	Mean SE $\hat{\Delta}_{c,n}$	Coverage			
50	Null	0.05	0.00	0.01	0.18	0.18	0.95			
50	Cauchy	0.45	0.09	0.07	0.15	0.15	0.94			
50	Bimodal	0.57	0.12	0.09	0.16	0.15	0.94			
50	Large	0.50	0.28	0.26	0.15	0.15	0.95			
50	Slope	0.27	0.11	0.11	0.17	0.17	0.94			
1000	Null	0.05	0.00	0.00	0.04	0.04	0.95			
1000	Cauchy	0.45	0.09	0.08	0.03	0.03	0.96			
1000	Bimodal	0.57	0.12	0.12	0.03	0.03	0.95			
1000	Large	0.50	0.28	0.28	0.03	0.04	0.96			
1000	Slope	0.27	0.11	0.12	0.04	0.04	0.95			

Notes: Table showing simulation results. Each row is a parameterization. Each parameterization was run for 2000 repetitions. All simulations used the tuning parameters: C = 2,  $D = 10^{-4}$ , and  $\sigma_Y = 1$ . The first column in the table is the number of *t*-scores. The second is the distribution of true effects and each of these is described in Section 7. The third column is unconditional (mean) statistical power at the status quo. The fourth column is the increase in unconditional power resulting from doubling the sample size of every experiment, so  $c = \sqrt{2}$ . The Fifth and sixth columns are the mean and standard devition of the point estimates. The seventh column is the mean of the estimated standard errors and the final column is the fraction of confidence intervals that contained the true  $\Delta_c$ .

### **B** Figures

	By Number of t-scores		By Nun	nber of Sites
	Main	Rob. Check	Main	Rob. Check
Â	.005	.033	.005	.033
-c	(.042)	(.026)	(.039)	(.030)
	[0, .088]	[0, .083]	[0, .082]	[0, .092]
$\hat{ heta}$	.92	1.20	1.03	0.86
	(.70)	(.73)	(.30)	(.67)
D	1e-4	3e-4	1e-4	3e-4
С	2	1	2	1
J	16	15	15	13
$\epsilon$	.27	.07	.61	.15
$\hat{\Delta}_{c}^{(n)}$	0.078	0.078	0.078	0.078
	0.006	0.006	0.006	0.006
Unconditional Power	.61	.61	.61	.61
t-scores	385	385	385	385
Sites	36	36	36	36
Treatments	11	11	11	11

Table 2: Empirical Exercise: Many Labs Replication Project

Notes: Table reports estimates  $\hat{\Delta}$  of gain in unconditional statistical power of a two-sided size 5% t-test resulting from counterfactually doubling every study's sample size from the status quo. Columns 1 and 2 report results where  $J, \epsilon$  are scaled based on the number of t-scores while in columns 3 and 4 these tuning parameters are scaled based on the number of study sites. The confidence intervals in columns 1 and 2 are more conservative. Columns 1 and 3 use the preferred choice of C, D while columns 2 and 4 present a robustness check where we have significantly altered C, D. We always set  $\sigma_Y = 1$ . Standard errors are reported in round brackets and 95% confidence intervals are in square brackets. Standard errors clustered by study site and treatment type. The bottom five rows report the estimate and standard error of  $\hat{\Delta}^{(n)}$  using the method of means, the fraction of t-scores larger than 1.96, the number of t-scores in total, the number of unique experimental research sites, and the number of unique experimental treatments. Data from Klein et al. (2014).

	By Number of t-scores		By Number of Articles	
	RCT	non-RCT	RCT	non-RCT
$\hat{\Delta}$	.072	.173	.095	.164
	(.025)	(.028)	(.020)	(.021)
	[.02, .12]	[.12, .23]	[.06, .13]	[.12, .20]
$\hat{ heta}$	1.04	.93	.73	.59
	(0.11)	(.10)	(.20)	(.17)
J	18	18	15	16
$\epsilon$	.10	.08	.46	.38
= RCT (p-value)		0.01		0.02
$= \Delta_c$ Many labs (p-value)	0.18	0.00	0.04	0.02
$=\Delta_c^{(n)}$ Many labs (p-value)	0.81	0.00	0.40	0.00
Status Quo Power	.38	.55	.38	.55
t-scores	7569	14171	7569	14171
Articles	80	145	80	145

Table 3: Empirical Application: Randomized Trials in Economics

Notes: Reports estimates  $\hat{\Delta}$  of the gain in unconditional statistical power of a two-sided size 5% *t*-test resulting from counterfactually doubling every study's sample size from the status quo  $(c=\sqrt{2})$ . Compares *t*-scores testing main hypotheses from randomized trials published in top economics journals versus those reported by studies using difference in difference, regression discontinuity or instrumental variables. Tuning parameters are  $D = 10^{-4}$ , C = 2, and  $\sigma_Y = 1$ . Table compares results when  $J, \epsilon$  are chosen based on the number of articles (less conservative) vs the number of *t*-scores (more conservative) in the meta-sample. Standard errors are reported in round brackets and 95% confidence intervals are in square brackets. Standard errors clustered by article. The bottom four rows report the outcome of a test for equality of the RCT vs non-RCT values of  $\Delta$ , the fraction of *t*-scores larger than 1.96 in each sample. Data from Brodeur et al. (2020).




Figure 1: Underpowered literature vs well-powered literature.




Figure 2: Compares power gain (y axis) of Economics RCTs vs non-RCTs over many  $c^2$  (x-axis). Data from Brodeur et al. (2016).




Figure 3: Compares power gain (y axis) of Many Labs vs Economics RCTs over many  $c^2$  (x-axis). Data from Brodeur et al. (2016) and Klein et al. (2014).




Figure 4: Compares power gain (y axis) of Many Labs vs Economics RCTs over many  $c^2$  (x-axis). Unlike Figure 3, this Many Labs estimate increases precision by conditioning on sample values of h and using the assumption that interventions are identical across study sites. Data from Brodeur et al. (2016) and Klein et al. (2014).

# C Detailed expressions

This section contains explicit expressions for terms that are too lengthy and involved for the main text.

#### C.1 Hermite Polynomials

The generalized and scaled probabalist's Hermite Polynomials are written explicitly below:

$$\chi_j(t) = \frac{1}{\sqrt{j!}} \sum_{l=0}^{[j/2]} (-1)^l \frac{(2l)!}{2^l l!} {j \choose 2l} \left(\frac{x}{\sqrt{1+\sigma_Y^2}}\right)^{j-2l}$$
$$\phi_j(t) = \frac{1}{\sqrt{j!}} \sum_{l=0}^{[j/2]} (-1)^l \frac{(2l)!}{2^l l!} {j \choose 2l} \left(\frac{x}{\sqrt{1+\sigma_Y^2-c^{-2}}}\right)^{j-2l}$$
$$\psi_j(t) = \frac{1}{\sqrt{j!}} \sum_{l=0}^{[j/2]} (-1)^l \frac{(2l)!}{2^l l!} {j \choose 2l} \left(\frac{x}{\sigma_Y}\right)^{j-2l}$$

# C.2 Centering term $\tilde{\Delta}_{c,n}$

When we do inference we must center the estimator on a sequence  $\tilde{\Delta}_{c,n}$ . While  $\tilde{\Delta}_{c,n}$  does converge to  $\Delta_c$ , it may still be large enough to matter for inference. The full expression is below.

$$\begin{split} \tilde{\Delta}_{c,n} &\equiv \Delta_c + \sum_{j=J_n+1}^{\infty} \eta_j \langle \psi_j, \pi_0 \rangle_W a_j \\ &+ \sum_{j=0}^{J_n} a_j \frac{\mathbb{E} \left[ \psi_j(T) \varphi_{\sigma_Y}(T) \hat{B}'_{\theta_0}(T) \mid R \right] \left[ G(\theta_0)' G(\theta_0) \right]^{-1} G(\theta_0)' E[\mathbf{v}(T) \mid R]}{\lambda_j \mathbb{E} \left[ \frac{1}{w_{\theta_0}(T)} \mid R \right]^2} \\ &+ \frac{\Delta_c}{\mathbb{E} \left[ \frac{1}{w_{\theta_0}(T)} \mid R \right]^2} B'_{\theta_0} \left[ G(\theta_0)' G(\theta_0) \right]^{-1} G(\theta_0)' E[\mathbf{v}(T) \mid R] \end{split}$$

where  $a_j$  is shorthand for:

$$a_j \equiv \lambda_j \int_{-cv}^{cv} \phi_j(s) ds - \int_{-cv/c}^{cv/c} \phi_j(s) ds$$

The expression references the gradients  $\hat{B}$  and B which are defined as the  $K \times 1$  vectors:

$$\hat{B}_{\theta}(t) \equiv \nabla_{\theta} \frac{1}{w_{\theta}(t)}$$
$$B_{\theta} \equiv \mathbb{E} \left[ \hat{B}_{\theta_0}(T) \mid R \right]$$

The  $|\mathcal{X}| \times K$  population gradient matrix G is defined as:

$$G(\theta)_{x,k} = -\lim_{\epsilon \to 0} \frac{f_{T|R}(x+\epsilon)}{w_{\theta}(x+\epsilon_n)^2} \frac{d}{d\theta} w_{\theta}(x+\epsilon) + \frac{f_{T|R}(x-\epsilon)}{w_{\theta}(x-\epsilon)^2} \frac{d}{d\theta} w_{\theta}(x-\epsilon)$$

 $\mathbf{v}_n(t)$  is a function that maps  $t \in \mathbb{R}$  to a  $|\mathcal{X}| \times 1$  vector with components:

$$\mathbf{v}_{n,x}(t) = \frac{1}{\epsilon_n} \left( \frac{\mathbf{1} \left\{ t \in (x, x + \epsilon_n] \right\}}{w_{\theta_0}(x + \epsilon_n)} - \frac{\mathbf{1} \left\{ t \in (x - \epsilon_n, x] \right\}}{w_{\theta_0}(x - \epsilon_n)} \right)$$

# C.3 Linearized Functions $Z_{n,J_n}(t)$

Lemma 7 expresses the estimation error in terms of a sample average of  $\frac{1}{n} \sum_{i=1}^{n} Z_{n,J_n}(t_i)$ . The functional form is written down below:

$$\begin{split} Z_{n,J_n}(t) &= \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]} \sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} \frac{\psi_j(t)\varphi_{\sigma_Y}(t)}{w_{\theta_0}(t)} \\ &+ \frac{\sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\theta_0}(T) \mid R\right] \left[G(\theta_0)'G_n(\theta_0)\right]^{-1} G(\theta_0)'\mathbf{v}(t)}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} \\ &+ \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} \frac{1}{w_{\theta_0}(t)} \\ &+ \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} B'_{\theta_0} \left[G(\theta_0)'G_n(\theta_0)\right]^{-1} G(\theta_0)'\mathbf{v}(t) \end{split}$$

where  $a_j$  is shorthand for:

$$a_j \equiv \lambda_j \int_{-cv}^{cv} \phi_j(s) ds - \int_{-cv/c}^{cv/c} \phi_j(s) ds$$

The meta-analyst doesn't know  $Z_n$  but in Theorem 4 they use an estimate  $\hat{Z}_n$  to do inference. The function  $\hat{Z}_n$  is written down below:

$$\begin{split} \hat{Z}_{n,J_{n}}(t) &= \frac{1}{\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\hat{\theta}_{n}}(t_{i})}} \sum_{j=0}^{J_{n}} \frac{a_{j}}{\lambda_{j}} \frac{\psi_{j}(t)\varphi_{\sigma_{Y}}(t)}{w_{\hat{\theta}_{n}}(t)} \\ &+ \frac{\sum_{j=0}^{J_{n}} \frac{a_{j}}{\lambda_{j}} E\left[\psi_{j}(T)\varphi_{\sigma_{Y}}(T)\hat{B}'_{\hat{\theta}_{n}}(T) \mid R\right] \left[G(\hat{\theta}_{n})'G_{n}(\hat{\theta}_{n})\right]^{-1} G(\hat{\theta}_{n})'\mathbf{v}_{\hat{\theta}_{n}}(t)}{\left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\hat{\theta}_{n}}(t_{i})}\right)^{2}} \\ &+ \frac{\hat{\Delta}_{c,n}}{\left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\hat{\theta}_{n}}(t_{i})}\right)^{2}} \frac{1}{w_{\hat{\theta}_{n}}(t)} \\ &+ \frac{\hat{\Delta}_{c,n}}{\left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\hat{\theta}_{n}}(t_{i})}\right)^{2}} B'_{\hat{\theta}_{n}} \left[G(\hat{\theta}_{n})'G_{n}(\hat{\theta}_{n})\right]^{-1} G(\hat{\theta}_{n})'\mathbf{v}_{\hat{\theta}_{n}}(t) \end{split}$$

# D Proofs

## D.1 Proof of Lemma 1

We show this using characteristic functions.  $K_1\pi$  is the PDF of the random variable h + Zwhere  $Z \sim N(0,1)$  and Z is independent of h. By the Convolution Theorem, the characteristic function of the sum of any two independent random variables is the pointwise product of the characteristic functions of the summands. The characteristic function of a normal random variable with mean zero and variance  $\sigma^2$  is  $e^{\omega^2/2\sigma^2}$ . Let  $\zeta_h(\omega)$  denote the characteristic function of h. The characteristic function of h + Z is  $e^{-\omega^2/2}\zeta_h$ .

Now consider  $K_{1-c^{-2}}K_{c^{-2}}\pi$ . This is the PDF of the random variable  $h+c^{-1}Z_1+\sqrt{1-c^{-2}}Z_2$ where  $h, Z_1, Z_2$  are all independent and  $Z_1, Z_2 \sim N(0, 1)$ . To find the characteristic function of the sum, we take the pointwise product of the chacteristic functions of the summands which is:  $e^{-\omega^2/2c^{-2}}e^{-\omega^2/2(1-c^{-2})}\zeta_h = e^{-\omega^2/2}\zeta_h$ . So  $K_1\pi$  and  $K_{1-c^{-2}}K_{c^{-2}}\pi$  are PDFs that share the same characteristic function. So they must agree almost everywhere and be the same distribution.

# D.2 Proof of Lemma 2

It is easier to decompose  $\Delta_c$  into the difference of the type-II error probabilities:  $\Delta_c = \beta_1 - \beta_c$ . Here  $\beta_1$  is the type-II error probability under the factual sample sizes:  $\beta_1 \equiv \int_{-cv}^{cv} f_T(t) dt$  and  $\beta_c$  is the error probability under counterfactual sample sizes:  $\beta_c \equiv \int_{-cv}^{cv} f_{T_c}(t) dt$ . First we use a change of variables of integration to rewrite  $\beta_c$ . Then we substitute in the definition of  $K_{c^{-2}}$ . These steps yield:

$$\beta_{c} = \int_{-cv}^{cv} f_{T_{c}}(t)dt = \int_{-cv}^{cv} \int_{-\infty}^{\infty} \varphi(t-ch)\pi_{0}(h)dhdt = \int_{-cv/c}^{cv/c} \int_{-\infty}^{\infty} \varphi(ct-ch)\pi_{0}(h)dhdt = \int_{-cv/c}^{cv/c} \int_{-\infty}^{\infty} \varphi_{c^{-2}}(t-h)\pi_{0}(h)dhdt = \int_{-cv/c}^{cv/c} K_{c^{-2}}\pi_{0}(h)dt$$

By an identical argument:  $\beta_1 = \int_{-cv}^{cv} K_1 \pi_0(h) dt$ . Then taking the difference:

$$\Delta_c = \int_{-cv}^{cv} (K_{1-c^{-2}} K_{c^{-2}} \pi_0)[t] dt - \int_{-cv/c}^{cv/c} (K_{c^{-2}} \pi_0)[t] dt$$

#### D.3 Proof of Lemma 3

As in Lemma 2, is easier to first decompose  $\Delta_c$  into the difference of the type-II error probabilities:  $\Delta_c = \beta_1 - \beta_c$ . Here  $\beta_1$  is the type-II error probability under the factual sample sizes:  $\beta_1 \equiv \int_{-cv}^{cv} f_T(t) dt$  and  $\beta_c$  is the error probability under counterfactual sample sizes:  $\beta_c \equiv \int_{-cv}^{cv} f_{T_c}(t) dt$ . Using the same argument as in Lemma 2, we do a change of variables of integration. Then we substitute in the definition of  $K_{c^{-2}}$ . Then we substitute in the singular value decomposition from (Carrasco and Florens, 2011). These steps yield:

$$\int_{-cv}^{cv} f_{T_c}(t)dt = \int_{-cv/c}^{cv/c} K_{c^{-2}}\pi_0 dt = \int_{-cv/c}^{cv/c} \sum_{j=0}^{\infty} \eta_j \langle \chi_j, \pi_0 \rangle_W \phi_j(t) dt$$

We would like to exchange the sum and the integral with Fubini's Theorem. To do this it is sufficient to show that:  $\sum_{j=0}^{\infty} \eta_j |\langle \chi_j, \pi_0 \rangle_W| \int_{-cv/c}^{cv/c} |\phi_j(t)| dt < \infty$ . To see that this condition is satisfied we first use the Cauchy Schwarz Inequality:

$$\int_{-cv/c}^{cv/c} \phi_j(t) dt \le \int_{-cv/c}^{cv/c} |\phi_j(t)| dt \le 2\frac{cv}{c} \int_{-cv/c}^{cv/c} \phi_j(t)^2 dt$$

The singular functions are normalized in their respective Hilbert spaces so  $||\phi_j||_X = 1$ . This is the same as taking the inner product with itself:  $||\phi_j||_X \equiv \langle \phi_j, \phi_j \rangle_X = \int_{-\infty}^{\infty} \phi_j(t)^2 \varphi_{1+\sigma_Y^2+c^{-2}}(t) dt = 1$ . Since the integrand is non-negative, this implies that bounded integral is less than one.  $\int_{-cv/c}^{cv/c} \phi_j(t)^2 \varphi_{1+\sigma_Y^2+c^{-2}}(t) dt \leq 1$ . Now we use the fact that the normal density is bounded below on this compact set:

$$\min_{t \in -cv/c, cv/c} \varphi_{1+\sigma_Y^2+c^{-2}}(t) = \varphi_{1+\sigma_Y^2+c^{-2}}(cv/c) = \exp\left(\frac{-cv^2}{c^2(1+\sigma_Y^2+c^{-2})}\right) / \sqrt{2\pi(1+\sigma_Y^2+c^{-2})}$$

Since the minimum of  $\varphi_{1+\sigma_Y^2+c^{-2}}(t)$  is bounded below, the integral without the normal kernel can be bounded above:

$$\int_{-cv/c}^{cv/c} \phi_j(t)^2 dt \le \exp\left(\frac{-cv^2}{c^2(1+\sigma_Y^2+c^{-2})}\right) / \sqrt{2\pi(1+\sigma_Y^2+c^{-2})}$$

Combining this with the Cauchy Schwarz bound from before:

$$\int_{-cv/c}^{cv/c} |\phi_j(t)| dt \le 2\frac{cv}{c} \exp\left(\frac{-cv^2}{c^2(1+\sigma_Y^2+c^{-2})}\right) / \sqrt{2\pi(1+\sigma_Y^2+c^{-2})}$$

This bound applies uniformly to all of the integrals for all j. This gives us the following bound on the absolute sum of the absolute values.

$$\sum_{j=0}^{\infty} \eta_j \left| \langle \chi_j, \pi_0 \rangle_W \right| \int_{-cv/c}^{cv/c} \left| \phi_j(t) \right| dt \le \left( 2 \frac{cv}{c} \exp\left( \frac{-cv^2}{c^2(1+\sigma_Y^2+c^{-2})} \right) \right) / \sqrt{2\pi(1+\sigma_Y^2+c^{-2})} \right) \sum_{j=0}^{\infty} \eta_j \left| \langle \chi_j, \pi_0 \rangle_W \right|$$

Since  $\pi_0$  is a PDF:  $|\langle \chi_j, \pi_0 \rangle_W| \leq 1$  by Holder's Inequality for all j. Since the  $\eta_j$  are a positive power series, their sum converges. So we have proven that  $\sum_{j=0}^{\infty} \eta_j |\langle \chi_j, \pi_0 \rangle_W| \int_{-cv/c}^{cv/c} |\phi_j(t)| dt < \infty$ . Since the absolute sum of the absolute integrals is finite we can use Fubini's Theorem and exchange the sum and the integral to write the  $\beta_c$  as a sum of intetrals.

$$\beta_c = \sum_{j=0}^{\infty} \eta_j \langle \chi_j, \pi_0 \rangle_W \int_{-cv/c}^{cv/c} \phi_j(t) dt$$

An identical argument with c = 1 lets use compute  $\beta_1$  and convert back into  $\Delta_c$  to conclude the Lemma:

$$\Delta_c = \sum_{j=0}^{\infty} \eta_j \langle \chi_j, \pi_0 \rangle_W \left( \lambda_j \int_{-cv}^{cv} \phi_j(t) dt - \int_{-cv/c}^{cv/c} \phi_j(t) dt \right)$$

#### D.4 Proof of Proposition 1

This proof formalizes the argument in Section 4.1. We start with the definition of  $Q_0$ . We then rewrite this in terms of the inner product.

$$Q_0(\pi) = ||K_1(\pi_0 - \pi)||_Y$$

Next we substitue in  $K_1 = K_{1-c^{-1}}K_{c^{-2}}$  from Lemma 1:

$$||K_1(\pi_0 - \pi)||_Y = ||K_{1-c^{-1}}K_{c^{-2}}(\pi_0 - \pi)||_Y$$

Recall the singular value decompositions of  $K_{1-c^{-1}}$  and  $K_{c^{-2}}$  from Section 3.3:  $K_{c^{-2}}\pi = \sum_{j=0}^{\infty} \eta_j \langle \chi_j, \pi \rangle_W \phi_j$  and  $K_{1-c^{-2}}g = \sum_{j=0}^{\infty} \lambda_j \langle \phi_j, g \rangle_X \psi_j$ . Substituting in the expressions for the singular value decomposition and then using the fact that the  $\phi_j$  are orthonormal in  $\langle \cdot, \cdot \rangle_X$ :

$$K_{1-c^{-1}}K_{c^{-2}}\pi = \sum_{j=0}^{\infty} \lambda_j \langle \phi_j, \ K_{c^{-2}}\pi \rangle_X \psi_j = \sum_{j=0}^{\infty} \lambda_j \eta_j \langle \chi_k, \pi \rangle_W \psi_j$$

Substituting the SVD into the objective function:

$$||K_{1-c^{-1}}K_{c^{-2}}(\pi_0 - \pi)||_Y = \left\| \sum_{j=0}^{\infty} \lambda_j \eta_j \langle \chi_k, \pi_0 - \pi \rangle_W \psi_j \right\|_Y$$

Since  $\psi_j$  are orthogonal in  $\langle \cdot, \cdot \rangle_Y$  we can use the Pythagorean Formula to write the norm as the sum of inner products. Then we can use the orthonormality of  $\psi_j$  a second time to eliminate the cross terms:

$$\left\| \left\| \sum_{j=0}^{\infty} \lambda_j \eta_j \langle \chi_k, \pi_0 - \pi \rangle_W \psi_j \right\|_Y = \sum_{j=0}^{\infty} \lambda_j^2 \eta_j^2 \langle \chi_j, \pi_0 - \pi \rangle_W^2$$

The singular values  $\eta_j$ ,  $\lambda_j$  are all strictly positive. Since the generalized Hermite polynomials  $\chi_j$  form a complete basis, any function in  $\mathcal{L}_W$  that is orthogonal to all of the  $\chi_j$  must be equal zero almost everywhere (Johnston, 2014). This means that any  $\pi$  that satisfies  $Q_0(\pi) = 0$  must equal  $\pi_0$  almost everywhere. If two probability density functions agree almost everywhere, then they are the same density. So  $Q_0(\pi) = 0$  if and only if  $\pi_0 = 0$ . Since  $Q_0$  is everywhere non-negative this gives us identification of  $\pi_0$ .

### D.5 Proof of Lemma 4

As in Lemma 2, is easier to first decompose  $\Delta_c$  into the difference of the type-II error probabilities:  $\Delta_c = \beta_1 - \beta_c$ . Here  $\beta_1$  is the type-II error probability under the factual sample sizes:  $\beta_1 \equiv \int_{-cv}^{cv} f_T(t) dt$  and  $\beta_c$  is the error probability under counterfactual sample sizes:  $\beta_c \equiv \int_{-cv}^{cv} f_{T_c}(t) dt$ . Recall that in the proof of Lemma 3 we showed that:

$$\beta_c = \sum_{j=0}^{\infty} \eta_j \langle \chi_j, \pi_0 \rangle_W \int_{-cv/c}^{cv/c} \phi_j(t) dt$$
$$\beta_1 = \sum_{j=0}^{\infty} \eta_j \lambda_j \langle \chi_j, \pi_0 \rangle_W \int_{-cv}^{cv} \phi_j(t) dt$$
$$\int_{-cv/c}^{cv/c} |\phi_j(t)| dt \le 2 \frac{cv}{c} \exp\left(\frac{-cv^2}{c^2(1+\sigma_Y^2+c^{-2})}\right) \Big/ \sqrt{2\pi(1+\sigma_Y^2+c^{-2})}$$

This means that for any  $\pi \in \mathcal{L}_W$ ,

$$\begin{aligned} \left| \beta_c - \sum_{j=0}^{\infty} \eta_j \langle \chi_j, \pi \rangle_W \int_{-cv/c}^{cv/c} \phi_j(t) dt \right| &= \left| \sum_{j=0}^{\infty} \eta_j \langle \chi_j, \pi_0 - \pi \rangle_W \int_{-cv/c}^{cv/c} \phi_j(t) dt \right| \\ &\leq \sum_{j=0}^{\infty} \eta_j \left| \langle \chi_j, \pi_0 - \pi \rangle_W \right| \left| \int_{-cv/c}^{cv/c} \phi_j(t) dt \right| \\ &\leq \left( 2\frac{cv}{c} \exp\left(\frac{-cv^2}{c^2(1+\sigma_Y^2+c^{-2})}\right) \right/ \sqrt{2\pi(1+\sigma_Y^2+c^{-2})} \right) \rho_c(\pi_0, \pi) \end{aligned}$$

By an identical argument:

$$\left|\beta_1 - \sum_{j=0}^{\infty} \eta_j \lambda_j \langle \chi_j, \pi \rangle_W \int_{-cv}^{cv} \phi_j(t) dt \right| \le \left(2cv \frac{e^{\frac{cv^2}{(1+\sigma_Y^2 + c^{-2})}}}{\sqrt{2\pi(1+\sigma_Y^2 + 1)}}\right) \rho_c(\pi_0, \pi)$$

Since the constant in front does not depend on  $\pi$ , for any sequence of  $\pi_n$ :

$$\rho_c(\pi_0, \pi_n) = \mathcal{O}_p(r_b) \implies \Delta_c - \sum_{j=0}^{\infty} \eta_j \langle \chi_j, \pi_n \rangle_W \left( \lambda_j \int_{-cv}^{cv} \phi_j(t) dt - \int_{-cv/c}^{cv/c} \phi_j(t) dt \right) = \mathcal{O}_p(r_n)$$

# D.6 Proof of Theorem 1

From definition of  $\rho_c$ :  $\rho_c(\pi_0, \hat{\pi}_n) = \sum_{j=0}^{\infty} \eta_j |\langle \chi_j, \pi_0 \rangle_W - \langle \chi_j, \hat{\pi}_n \rangle_W|$ . The spectral cutoff regularization of  $\hat{\pi}_n$  sets:  $\langle \chi_j, \hat{\pi}_n \rangle_W = 0 \forall_{j>J_n}$ . So we can split  $\rho_c$  into the sampling error and regularization bias parts:

$$\rho_c(\pi_0, \hat{\pi}_n) = \sum_{j=0}^{J_n} \eta_j \left| \langle \chi_j, \pi_0 \rangle_W - \langle \chi_j, \hat{\pi}_n \rangle_W \right| + \sum_{J_n+1}^{\infty} \eta_j \left| \langle \chi_j, \pi_0 \rangle_W \right|$$

Now we rewrite the first sum which comes from sampling error. We do this by substituting in  $\hat{\pi}_n$  from Equation 13 and then using the linearity of the inner product and the fact that the  $\chi_j$  are orthonormal in  $\langle \cdot, \cdot \rangle_W$ .

$$\sum_{j=0}^{J_n} \eta_j \left| \langle \chi_j, \pi_0 \rangle_W - \langle \chi_j, \hat{\pi}_n \rangle_W \right| = \sum_{j=0}^{J_n} \left| \eta_j \langle \chi_j, \pi_0 \rangle_W - \frac{\frac{1}{n} \sum_{i=1}^n \psi_j(t_i) \varphi_{\sigma_Y^2}(t_i)}{\lambda_j} \right|$$

We now wish to substitute in an expectation over T for  $\eta_j \langle \chi_j, \pi_0 \rangle_W$ . We can do this with equation 6 which says:  $f_T = \sum_{j=1}^{\infty} \lambda_j \eta_j \langle \chi_j, \pi_0 \rangle_W \psi_j$ . Since the  $\psi_j$  are orthonormal:

$$\langle f_T, \psi_j \rangle_Y = \lambda_j \eta_j \langle \chi_j, \pi_0 \rangle_W$$

The inner  $\langle \cdot, \cdot \rangle_Y$  product is an integral and  $f_T$  is a PDF. The integral over a PDF is an expectation. So we have:

$$\frac{\mathbb{E}\left[\psi_j(T)\varphi_{\sigma_Y^2}(T)\right]}{\lambda_j} = \eta_j \langle \chi_j, \pi_0 \rangle_W$$

Substituting this into our expression for the sampling error:

$$\sum_{j=0}^{J_n} \left| \eta_j \langle \chi_j, \pi_0 \rangle_W - \frac{\frac{1}{n} \sum_{i=1}^n \psi_j(t_i) \varphi_{\sigma_Y^2}(t_i)}{\lambda_j} \right| = \sum_{j=0}^{J_n} \frac{1}{\lambda_j} \left| \mathbb{E} \left[ \psi_j(T) \varphi_{\sigma_Y^2}(T) \right] - \frac{1}{n} \sum_{i=1}^n \psi_j(t_i) \varphi_{\sigma_Y^2}(t_i) \right|$$

The observations  $t_i$  are identically distributed draws of T. While they are not all independent, each observation is independent of at least n - C other observations. So, the variance is bounded by:

$$\begin{aligned} \operatorname{Var}\left(\frac{1}{n}\sum_{i=1}^{n}\psi_{j}(t_{i})\varphi_{\sigma_{Y}^{2}}(t_{i})\right) &= \frac{1}{n^{2}}\sum_{i=1}^{n}\sum_{l=1}^{n}Cov\left(\psi_{j}(t_{i})\varphi_{\sigma_{Y}^{2}}(t_{i}),\psi_{j}(t_{l})\varphi_{\sigma_{Y}^{2}}(t_{l})\right) \\ &\leq C\frac{1}{n^{2}}\sum_{i=1}^{n}\sum_{l=1}^{n}\operatorname{Var}\left(\psi_{j}(t_{i})\varphi_{\sigma_{Y}^{2}}(t_{i})\right) \\ &= \frac{C}{n}\operatorname{Var}\left(\psi_{j}(T)\varphi_{\sigma_{Y}^{2}}(T)\right) \end{aligned}$$

The distribution of T is not changing. Furthermore, we can upper bound  $\operatorname{Var}\left(\psi_j(T)\varphi_{\sigma_Y^2}(T)\right)$  by 1 using Holder's Inequality and the fact that  $||\psi||_Y = 1$ .

$$\operatorname{Var}\left(\psi_j(T)\varphi_{\sigma_Y^2}(T)\right) \le \mathbb{E}\left[\left(\psi_j(T)\varphi_{\sigma_Y^2}(T)\right)^2\right] \le \int_{-\infty}^{\infty} \psi_j(t)^2 \varphi_{\sigma_Y^2}(t) dt = 1$$

This lets us upper bound the variance of the sums:

$$\operatorname{Var}\left(\frac{1}{n}\sum_{i=1}^{n}\psi_{j}(t_{i})\varphi_{\sigma_{Y}^{2}}(t_{i})\right) \leq \frac{C}{n}$$

By Chebyshev's Inequality:

$$\frac{1}{n}\sum_{i=1}^n\psi_j(t_i)\varphi_{\sigma_Y^2}(t_i) = \mathcal{O}_p(n^{-1/2})$$

Now we want to bound the sum of the centered sample means over j in the sampling error.

use the fact that  $\lambda_j$  is a power sequence, i.e.  $\lambda_j = \lambda_0^{j/2}$ . This gives us the identity:

$$\lambda_{J_n} \sum_{j=0}^{J_n} \frac{1}{\lambda_j} = \sum_{j=0}^{J_n} \lambda_0^{J_n - j/2} = \sum_{j=0}^{J_n} \lambda_0^{j/2} = \sum_{j=0}^{J_n} \lambda_j \to \frac{1}{1 - \lambda_0}$$

This means that we can control the order of the sum:  $\sum_{j=0}^{J_n} \frac{1}{\lambda_j} = \mathcal{O}(\lambda_{J_n}^{-1})$ . Combining this with the order of the centered sample means we can control the order of the sampling error part:

$$\sum_{j=0}^{J_n} \frac{1}{\lambda_j} \left| \mathbb{E} \left[ \psi_j(T) \varphi_{\sigma_Y^2}(T) \right] - \frac{1}{n} \sum_{i=1}^n \psi_j(t_i) \varphi_{\sigma_Y^2}(t_i) \right| = \mathcal{O}_p \left( \lambda_{J_n}^{-1} n^{-1/2} \right)$$

Next we bound the second sum that comes from regularization bias. Since  $\pi_0$  is a PDF, its  $\mathcal{L}_1$  norm is 1. Since the normal density with variance greater than 1 is always itself less than 1 the  $|| \cdot ||_Y$ -norm is less than the  $\mathcal{L}_1$  norm and  $||\pi_0||_W \leq 1$ . Using the Cauchy Schwarz Indequality and the fact that the  $\eta_j = \eta_2^{j/2}$ :

$$\sum_{J_n+1}^{\infty} \eta_j \left| \langle \chi_j, \pi_0 \rangle_W \right| \le \sqrt{\sum_{J_n+1}^{\infty} \eta_j^2} \sqrt{\sum_{J_n+1}^{\infty} \langle \chi_j, \pi_0 \rangle_W^2} = \mathcal{O}\left( \sqrt{\sum_{J_n+1}^{\infty} \eta_j^2} \right) = \mathcal{O}\left(\eta_{J_n}\right)$$

Combining the bounds on the sampling error and regularization bias we have proven the first claim in Theorem 1:

$$\rho_c(\pi_0, \hat{\pi}_n) = \mathcal{O}_p\left(\lambda_{J_n}^{-1} n^{-1/2} + \eta_{J_n}\right)$$

Now we prove the second claim in Theorem 1. Our goal is to choose a growth rate for  $J_n$  given  $\sigma_Y^2 > 0$  such that the order  $\lambda_{J_n}^{-1} n^{-1/2} + \eta_{J_n}$  is minimized. To do this it is sufficient to set the orders of the two summands equal to each other. A sufficient condition for this is that for some d > 0,

$$\frac{\eta_{J_n}}{\lambda_{J_n}^{-1}n^{-1/2}} \to c$$

Recall the definitions of the singular values  $\lambda_j$  and  $\eta_j$  from Section 3.3:  $\eta_j = \left(\frac{1+\sigma_Y^2-c^{-2}}{1+\sigma_Y^2}\right)^{j/2}$ and  $\lambda_j = \left(\frac{\sigma_Y^2}{\sigma_Y^2+1-c^{-2}}\right)^{j/2}$ . Notice that we can rewrite  $\eta_j$  as:  $\eta_j = \lambda_j^{-1} \left(\frac{\sigma_Y^2}{1+\sigma_Y^2}\right)^{j/2}$ . This makes their ratio:

$$\frac{\eta_{J_n}}{\lambda_{J_n}^{-1} n^{-1/2}} = \left(\frac{\sigma_Y^2}{1 + \sigma_Y^2}\right)^{j/2} n^{1/2}$$

So if the meta-analyst chooses  $J_n$  such that for some c > 0,  $n^{1/2} \left( \frac{\sigma_Y^2}{\sigma_Y^2 + 1} \right)^{J_n/2} \to c$ , then:

$$\frac{\eta_{J_n}}{\lambda_{J_n}^{-1}n^{-1/2}} \to c$$

So the order of  $\eta_{J_n}$  and  $n^{-1/2}\lambda_{J_n}^{-1}$  are the same and the order of the sum is minimized. Next we compute this order exactly. To do this we compute:

$$\left(\frac{\sigma_Y^2}{\sigma_Y^2+1}\right)^{J_n/2} = \mathcal{O}\left(n^{-1/2}\right)$$
$$J_n/2 = \mathcal{O}\left(\log_{\frac{\sigma_Y^2}{\sigma_Y^2+1}}\left(n^{-1/2}\right)\right)$$
$$\eta_{J_n} = \mathcal{O}\left(\left(\frac{1+\sigma_Y^2-c^{-2}}{1+\sigma_Y^2}\right)^{\log_{\frac{\sigma_Y^2}{\sigma_Y^2+1}}\left(n^{-1/2}\right)}\right)$$

Using the change of log base:

$$\log_{\frac{\sigma_Y^2}{\sigma_Y^2 + 1}} \left( n^{-1/2} \right) = \frac{\log_{\frac{1 + \sigma_Y^2 - c^{-2}}{1 + \sigma_Y^2}} \left( n^{-1/2} \right)}{\log_{\frac{1 + \sigma_Y^2 - c^{-2}}{1 + \sigma_Y^2}} \left( \frac{\sigma_Y^2}{\sigma_Y^2 + 1} \right)}$$

Some algebra lets us rewrite the rate to obtain the claim of Theorem 1.

$$\left(\frac{1+\sigma_Y^2-c^{-2}}{1+\sigma_Y^2}\right)^{\log_{\frac{\sigma_Y^2}{\sigma_Y^2+1}}(n^{-1/2})} = n^{-q/2}$$

# D.7 Proof of Lemma 5

By applying some algebra to Bayes' Rule we get:  $f_T(t) = \frac{f_T(t|R)\mathbb{E}[w_{\theta_0}(T)]}{w_{\theta_0}(t)}$ . For any measurable function  $\gamma : \mathbb{R} \to \mathbb{R}$ ,

$$E[\gamma(T)] = \int_{-\infty}^{\infty} \gamma(t) f_T(t) dt = \int_{-\infty}^{\infty} \gamma(t) \frac{f_T(t) | R}{w_{\theta_0}(t)} dt \mathbb{E}[w_{\theta_0}(T)] = \mathbb{E}\left[\frac{\gamma(T)}{w_{\theta_0}(T)} \mid R\right] \mathbb{E}[w_{\theta_0}(T)]$$

To compute  $\mathbb{E}[w_{\theta_0}(T)]$  by taking an expectation of over T|R, we use the following trick:

$$\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)}|R\right] = \int_{-\infty}^{\infty} \frac{1}{w_{\theta_0}(T)} f_T(t|R) dt = \int_{-\infty}^{\infty} \frac{1}{w_{\theta_0}(T)} \frac{f_T(t)w_{\theta_0}(t)}{E[w_{\theta_0}(T)]} dt = \frac{1}{E[w_{\theta_0}(T)]}$$

Setting  $\gamma(t) = \psi_j(T)\varphi_{\sigma_Y}(T)$  yields:

$$\mathbb{E}[\psi_j(T)\varphi_{\sigma_Y}(T)] = \frac{\mathbb{E}\left[\frac{\psi_j(T)\varphi_{\sigma_Y}(T)}{w_{\theta_0}(T)} \mid R\right]}{\mathbb{E}[w_{\theta_0}(T) \mid R]}$$

#### D.8 Proof of Proposition 2

Recall the population objective function:

$$Q_{0}(\pi,\theta) \equiv \sum_{j=0}^{\infty} \left( E\left[ \frac{\psi_{j}(T)\varphi_{\sigma_{Y}}(T)}{w_{\theta_{0}}(T)} \mid R \right] / E\left[ \frac{1}{w_{\theta_{0}}(T)} \mid R \right] - \lambda_{j}\eta_{j}\langle\pi,\chi_{j}\rangle_{W} \right)^{2} + \sum_{x \in \mathcal{X}} \lim_{\epsilon \to 0} \left( \frac{f_{T \mid R}(x+\epsilon)}{w_{\theta}(x+\epsilon)} - \frac{f_{T \mid R}(x-\epsilon)}{w_{\theta}(x-\epsilon)} \right)^{2}$$

As in the case without publication bias,  $Q_0(\pi_0, \theta_0) = 0$  and  $Q_0(\pi, \theta)$  is everywhere nonnegative. To show identification we must show that the objective cannot equal zero for any other arguments. First we show that if  $\theta \neq \theta_0$  then  $Q_0(\pi, \theta_0) > 0$  for any  $\pi$ . Using Bayes' Rule:  $f_T(t|R) = \frac{f_T(t)w_{\theta_0}(t)}{E[w_{\theta_0}(T)]}$ . Since  $f_T$  must be continuous,

$$\sum_{x \in \mathcal{X}} \lim_{\epsilon \to 0} \left( \frac{f_{T \mid R}(x+\epsilon)}{w_{\theta}(x+\epsilon)} - \frac{f_{T \mid R}(x-\epsilon)}{w_{\theta}(x-\epsilon)} \right)^2 = \frac{1}{E[w_{\theta_0}(T)]} \sum_{x \in \mathcal{X}} \lim_{\epsilon \to 0} f_T(x) \left( \frac{w_{\theta_0}(x+\epsilon)}{w_{\theta}(x+\epsilon)} - \frac{w_{\theta_0}(x+\epsilon)}{w_{\theta}(x-\epsilon)} \right)^2$$
$$\geq \frac{1}{E[w_{\theta_0}(T)]} \max_{x \in \mathcal{X}} \lim_{\epsilon \to 0} f_T(x) \left( \frac{w_{\theta_0}(x+\epsilon)}{w_{\theta}(x+\epsilon)} - \frac{w_{\theta_0}(x+\epsilon)}{w_{\theta}(x-\epsilon)} \right)^2$$

Now recall the inequality form Assumption 3:

$$\max_{x \in \mathcal{X}} \lim_{\epsilon \to 0} L \left| \frac{w_{\theta_0}(x+\epsilon)}{w_{\theta}(x+\epsilon)} - \frac{w_{\theta_0}(x-\epsilon)}{w_{\theta}(x-\epsilon)} \right| \ge ||\theta - \theta_0||_{\infty}$$

Substituting this in:

$$\frac{1}{E[w_{\theta_0}(T)]} \max_{x \in \mathcal{X}} \lim_{\epsilon \to 0} f_T(x) \left( \frac{w_{\theta_0}(x+\epsilon)}{w_{\theta}(x+\epsilon)} - \frac{w_{\theta_0}(x+\epsilon)}{w_{\theta}(x-\epsilon)} \right)^2 \ge \frac{1}{E[w_{\theta_0}(T)]} ||\theta - \theta_0||_{\infty}^2 \min_{x \in \mathcal{X}} f_T(x)$$

Since  $\mathcal{X}$  is finite and  $f_T$  is supported everywhere,  $\min_{x \in \mathcal{X}} f_T(x) \ge 0$ . Moreover, Assumption 2 guarantees that  $0 < \frac{1}{E[w_{\theta_0}(T)]} < \infty$ . So, the second sum in the objective function is zero if and only if  $\theta = \theta_0$ .

$$\sum_{x \in \mathcal{X}} \lim_{\epsilon \to 0} \left( \frac{f_{T \mid R}(x+\epsilon)}{w_{\theta}(x+\epsilon)} - \frac{f_{T \mid R}(x-\epsilon)}{w_{\theta}(x-\epsilon)} \right)^2 = 0 \iff ||\theta - \theta_0||_{\infty} = 0$$

This means that in order for  $Q_0(\pi, \theta) = 0$ , it must be  $Q(\pi, \theta_0) = 0$ . Concentrating the objective function gives:

$$Q_0(\pi,\theta_0) = \sum_{j=0}^{\infty} \left( E\left[ \frac{\psi_j(T) \varphi_{\sigma_Y}(T)}{w_{\theta_0}(T)} \mid R \right] / E\left[ \frac{1}{w_{\theta_0}(T)} \mid R \right] - \lambda_j \eta_j \langle \pi, \chi_j \rangle_W \right)^2$$

Now we use Lemma 5 to substitute in the unconditional expectations:

$$Q_0(\pi, \theta_0) = \sum_{j=0}^{\infty} \left( E\left[ \psi_j(T) \,\varphi_{\sigma_Y}(T) \right] - \lambda_j \eta_j \langle \pi, \chi_j \rangle_W \right)^2$$

By Proposition 1, this is uniquely minimized at  $\pi_0$ . So  $\pi_0, \theta_0$  are the unique minimizers of  $Q_0(\pi, \theta)$  and are therefore identified.

# D.9 Proof of Theorem 2

This proof consists of two steps. In the first step we show the consistency of  $\hat{\theta}_n$ . In the second step we use this find the rate of convergence of  $\rho_c(\pi_0, \hat{\pi}_n^{pb})$ .

## Step 1

 $\hat{\theta}_n$  is the minimizer of the following sample objective function (which is  $\hat{Q}_n(\pi, \theta)$  with the  $\pi$  concentrated out).

$$\hat{\theta}_n = \min_{\theta} \hat{Q}_n(\hat{\pi}_n, \theta)$$

$$= \min_{\theta} \sum_{x \in \mathcal{X}} \left( \frac{\frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbf{1} \{ t_i \in (x, x + \epsilon_n] \}}{w_{\theta}(x + \epsilon_n)} - \frac{\frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbf{1} \{ t_i \in (x - \epsilon_n, x] \}}{w_{\theta}(x - \epsilon_n)} \right)^2$$

This is the analogue of the concentrated population objective function.

$$\theta_0 = \min_{\theta} Q_0(\pi_0, \theta)$$
  
= 
$$\min_{\theta} \sum_{x \in \mathcal{X}} \left( \frac{\frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbf{1} \{ t_i \in (x, x + \epsilon_n] \}}{w_{\theta}(x + \epsilon_n)} - \frac{\frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbf{1} \{ t_i \in (x - \epsilon_n, x] \}}{w_{\theta}(x - \epsilon_n)} \right)^2$$

Next we show that the objective functions converge.  $n^{2/3} \sup_{\theta} \left| \hat{Q}_n(\hat{\pi}_n, \theta) - Q_0(\pi_0, \theta) \right| = \mathcal{O}_p(1)$ . We will call this fact (i).

By Assumption 3,  $||\theta_0 - \hat{\theta}_n||_{\infty}^2 \leq LQ_0(\pi_0, \hat{\theta}_n)$ . Call this fact (ii) Moreover, it is clear by inspection that  $Q_0(\pi_0, \theta_0) = 0$ . Call this fact (iii).

Now we will successively apply facts (i), (iii), the definition of  $\hat{\theta}_n$  as the minimizer, (iii), and (ii):

$$\begin{aligned} n^{2/3} ||\theta_0 - \hat{\theta}_n||_{\infty}^2 &\leq n^{2/3} L Q_0(\pi_0, \hat{\theta}_n) \\ &= n^{2/3} L \hat{Q}_n(\pi_0, \hat{\theta}_n) + \mathcal{O}_p(1) \\ &\leq n^{2/3} L \hat{Q}_n(\pi_0, \theta_0) + \mathcal{O}_p(1) \\ &= n^{2/3} L Q_0(\pi_0, \theta_0) + \mathcal{O}_p(1) \\ &= \mathcal{O}_p(1) \end{aligned}$$

So we conclude:

$$||\theta_0 - \hat{\theta}_n||_{\infty} = \mathcal{O}_p\left(n^{-1/3}\right)$$

#### Step 2

Using an argument identical to the proof of Theorem 1 we can obtain:

$$\rho_c(\pi_0, \hat{\pi}_n^{pb}) = \sum_{j=0}^{J_n} \frac{1}{\lambda_j} \left| \frac{1}{n \frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\hat{\theta}_n}(t_i)}} \sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y^2}(t_i)}{w_{\hat{\theta}_n}(t_i)} - E\left[\psi_j(T)\varphi_{\sigma_Y^2}(T)\right] \right| + \sum_{j=J_n+1}^\infty \eta_j \left| \langle \pi_0, \chi_j \rangle_W \right|$$

We can further break up the first sum into:

$$\begin{split} &\sum_{j=0}^{J_n} \frac{1}{\lambda_j} \left| \frac{1}{n \frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\hat{\theta}_n}(t_i)}} \sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y^2}(t_i)}{w_{\hat{\theta}_n}(t_i)} - E\left[\psi_j(T)\varphi_{\sigma_Y^2}(T)\right] \right| \\ &\leq \sum_{j=0}^{J_n} \frac{1}{\lambda_j} \left| \frac{1}{n \frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\hat{\theta}_n}(t_i)}} \sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y^2}(t_i)}{w_{\theta_0}(t_i)} - E\left[\psi_j(T)\varphi_{\sigma_Y^2}(T)\right] \right| \\ &+ \sum_{j=0}^{J_n} \frac{1}{\lambda_j} \left| \frac{1}{n \frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\hat{\theta}_n}(t_i)}} \sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y^2}(t_i)}{w_{\hat{\theta}_n}(t_i)} - \frac{1}{n \frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\hat{\theta}_n}(t_i)}} \sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y^2}(t_i)}{w_{\hat{\theta}_n}(t_i)} \right| \\ &= [A] + [B] \end{split}$$

First we bound sum [A]. Using Lemma 5 we can replace the unconditional expectation with the ratio of conditional expectations:

$$\begin{split} &\sum_{j=0}^{J_n} \frac{1}{\lambda_j} \left| \frac{1}{n_n^{\frac{1}{n}} \sum_{i=1}^n \frac{1}{w_{\hat{\theta}_n}(t_i)}} \sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y^2}(t_i)}{w_{\theta_0}(t_i)} - E\left[\psi_j(T)\varphi_{\sigma_Y^2}(T)\right] \right| \\ &= \sum_{j=0}^{J_n} \frac{1}{\lambda_j} \left| \frac{1}{n_n^{\frac{1}{n}} \sum_{i=1}^n \frac{1}{w_{\hat{\theta}_n}(t_i)}} \sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y^2}(t_i)}{w_{\theta_0}(t_i)} - \frac{\mathbb{E}\left[\frac{\psi_j(T)\varphi_{\sigma_Y}(T)}{w_{\theta_0}(T)} \mid R\right]}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]} \right| \end{split}$$

Assumption 2 guarantees that  $w_{\theta_0}(t)$  is bounded away from zero. By the same arguments as Theorem 1,

$$\frac{1}{n}\sum_{i=1}^{n}\frac{1}{w_{\hat{\theta}_{n}}(t_{i})} - \mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right] = \mathcal{O}_{p}\left(n^{-1/2}\right)$$
$$\frac{1}{n}\sum_{i=1}^{n}\frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}^{2}}(t_{i})}{w_{\theta_{0}}(t_{i})} - \mathbb{E}\left[\frac{\psi_{j}\left(T\right)\varphi_{\sigma_{Y}}\left(T\right)}{w_{\theta_{0}}(T)} \mid R\right] = \mathcal{O}_{p}\left(n^{-1/2}\right)$$

In the proof of Theorem 1 we showed that  $\lambda_{J_n} \sum_{j=0}^{J_n} \frac{1}{\lambda_j} \to \frac{1}{1-\lambda_0}$ . So we have:

$$[A] = \mathcal{O}_p\left(\lambda_{J_n}^{-1} n^{-1/2}\right)$$

Bounding term [B] requires bounding the convergence of  $\hat{\theta}_n$ . We already proved in step 1 that  $||\hat{\theta}_n - \theta_0||_{\infty} = \mathcal{O}_p(n^{-1/3})$ . Now by Assumption 3,  $\sup_t \left| \frac{1}{w_{\hat{\theta}_n}(t)} - \frac{1}{w_{\theta_0}(t)} \right| = \mathcal{O}_p(n^{-1/3})$ . This implies:

$$\frac{1}{n\frac{1}{n}\sum_{i=1}^{n}\frac{1}{w_{\hat{\theta}_{n}}(t_{i})}}\sum_{i=1}^{n}\frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}^{2}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})} - \frac{1}{n\frac{1}{n}\sum_{i=1}^{n}\frac{1}{w_{\hat{\theta}_{n}}(t_{i})}}\sum_{i=1}^{n}\frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}^{2}}(t_{i})}{w_{\theta_{0}}(t_{i})} = \mathcal{O}_{p}\left(n^{-1/3}\right)$$

Again using the fact that  $\lambda_{J_n} \sum_{j=0}^{J_n} \frac{1}{\lambda_j} \to \frac{1}{1-\lambda_0}$ ,

$$[B] = \mathcal{O}_p\left(\lambda_{J_n}^{-1} n^{-1/3}\right)$$

The regularization bias identical to the one in Theorem 1. So we have:

$$\rho_c(\pi_0, \hat{\pi}_n^{pb}) = \mathcal{O}_p\left(\lambda_{J_n}^{-1} n^{-1/3} + \eta_{J_n}\right)$$

If the meta-analyst chooses  $J_n$  such that for some c > 0,  $n^{1/3} \left(\frac{\sigma_Y^2}{\sigma_Y^2 + 1}\right)^{J_n/2} \to c$  then again by an identical argument to the one in the proof of Theorem 1, with  $\frac{1}{2}$  exchanged for  $\frac{1}{3}$ 

$$\rho_c(\pi_0, \hat{\pi}_n^{pb}) = \mathcal{O}_p \left( n^{-\frac{1}{2} \frac{\log\left(\frac{1+\sigma_Y^2}{1+\sigma_Y^2-c^{-2}}\right)}{\log\left(\frac{\sigma_Y^2+1}{\sigma_Y^2}\right)}} \right)$$

# D.10 Proof of Theorem 3

We want to invoke Theorem 2.1 from Neumann (2013). To do this we need to show (i) weak dependence and the (ii) Lindebergh condition. Weak dependence is immediate because each observation is independent of all but C other observations and C is fixed. Next we check the Lindebergh condition. The following Lyapunov condition is sufficient for the Lindebergh condition:

$$\frac{\mathbb{E}\left[\left(\sum_{j=0}^{J_n} \frac{1}{\lambda_j} Z_j(T,\theta_0)\right)^4\right]}{nE\left[\left(\sum_{j=0}^{J_n} \frac{1}{\lambda_j} Z_j(T,\theta_0)\right)^2\right]^2} \to 0$$

Next we show that the Lyapunov condition holds. Since  $\lambda_j = \left(\frac{\sigma_Y^2}{1+\sigma_Y^2+c^{-2}}\right)^{j/2}$ , we can bound the sum:  $\lambda_{J_n} \sum_j^{J_n} \frac{1}{\lambda_j} = \sum_{j=0}^{J_n} \lambda_j < \frac{1}{1-\frac{\sigma_Y^2}{1+\sigma_Y^2+c^{-2}}}$ . Combining this fact with (i) of Theorem 7, there is an M > 0 such that  $\left|\sum_j \frac{1}{\lambda_j} Z_j(T, \theta_0)\right| / (Mn^{1/3}\lambda_{J_n}) < 1$ . If we multiply the numerator and denominator by  $M^{-4}\lambda_{J_n}^4 n^{-4/3}$ , we obtain  $\frac{\mathbb{E}\left[\left(\sum_{j=0}^{J_n} \frac{\lambda_{J_n}}{\lambda_j} n^{-1/3} Z_j(T, \theta_0)/M\right)^4\right]}{nE\left[\left(\sum_{j=0}^{J_n} \frac{\lambda_{J_n}}{\lambda_j} n^{-1/3} Z_j(T, \theta_0)/M\right)^2\right]^2}$ . Since

 $\begin{aligned} \left| \sum_{j=0}^{J_n} \frac{\lambda_{J_n}}{\lambda_j} n^{-1/3} Z_j(T,\theta_0) / M \right| < 1, \text{ the fourth moment is smaller than the second moment. So,} \\ \frac{\mathbb{E} \left[ \left( \sum_{j=0}^{J_n} \frac{1}{\lambda_j} Z_j(T,\theta_0) \right)^4 \right]}{n E \left[ \left( \sum_{j=0}^{J_n} \frac{1}{\lambda_j} Z_j(T,\theta_0) \right)^2 \right]^2} \le \frac{1}{n^{1/3} \lambda_{J_n}^2 E \left[ \left( \sum_{j=0}^{J_n} \frac{1}{\lambda_j} Z_j(T,\theta_0) \right)^2 \right]}. \end{aligned}$ By Assumption 5, the sum of  $Z_j$  dominates the other terms and

$$\liminf n^{4/3} \lambda_{J_n}^2 n^{-1} \mathbb{E}\left[\left(\sum_{j=0}^{J_n} \frac{1}{\lambda_j} Z_j(T,\theta_0)\right)^2\right] = \liminf n^{1/3} \lambda_{J_n}^2 E\left[\left(\sum_{j=0}^{J_n} \frac{1}{\lambda_j} Z_j(T,\theta_0)\right)^2\right] = \infty$$

So the Lyapunov condition holds.

### D.11 Proof of Lemma 6

Using the statement from Appendix C.2 for  $\tilde{\Delta}_{c,n}$ :

$$\begin{split} \tilde{\Delta}_{c,n} - \Delta_c &= \sum_{j=J_n+1}^{\infty} \eta_j \langle \psi_j, \pi_0 \rangle_W a_j \\ &+ \sum_{j=0}^{J_n} a_j \frac{\mathbb{E}\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\theta_0}(T) \mid R\right] \left[G(\theta_0)'G(\theta_0)\right]^{-1} G(\theta_0)' E[\mathbf{v}(T) \mid R]}{\lambda_j \mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} \\ &+ \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} B'_{\theta_0} \left[G(\theta_0)'G(\theta_0)\right]^{-1} G(\theta_0)' E[\mathbf{v}(T) \mid R] \end{split}$$

where  $a_j$  is shorthand for:

$$a_j \equiv \lambda_j \int_{-cv}^{cv} \phi_j(s) ds - \int_{-cv/c}^{cv/c} \phi_j(s) ds$$

First we bound the first sum. We showed in the Proof of Lemma 3 that the integrals  $\int_{-cv/c}^{cv/c} \phi_j(s) ds$  and  $\int_{-cv}^{cv} \phi_j(s) ds$  are bounded uniformly in j which means that  $\sup_j |a_j| < \infty$ . We also showed in the proof of Lemma 3 that  $|\langle \psi_j, \pi_0 \rangle_W| \leq 1$  for all j. We also showed in the proof of Theorem 1 that  $\sum_{j=J_n+1}^{\infty} \frac{|\langle \psi_j, \pi_0 \rangle_W|}{\lambda_j} \leq \eta_{J_n}$ . By Holder's Inequality and the properties of geometric series, the regularization bias term is:

$$\sum_{j=J_n+1}^{\infty} \eta_j \langle \psi_j, \pi_0 \rangle_W a_j = \mathcal{O}\left(\sum_{j=J_n+1}^{\infty} \eta_j\right) = \mathcal{O}\left(\eta_{J_n}\right)$$

Next we bound the second and third sum. This involves bounding the  $|\mathcal{X}| \times 1$  vector  $E[\mathbf{v}(T) | R]$ . It will be enough to bound its infinity norm since the number of element does not change. Recall from Appendix C.3 that  $\mathbf{v}_n(t)$  is a function that maps  $t \in \mathbb{R}$  to a  $|\mathcal{X}| \times 1$  vector with components:

$$\mathbf{v}_{n,x}(t) = \frac{1}{\epsilon_n} \left( \frac{\mathbf{1} \left\{ t \in (x, x + \epsilon_n] \right\}}{w_{\theta_0}(x + \epsilon_n)} - \frac{\mathbf{1} \left\{ t \in (x - \epsilon_n, x] \right\}}{w_{\theta_0}(x - \epsilon_n)} \right)$$

For each x,

$$\mathbb{E}[\mathbf{v}_{n,x}(T) \mid R] = \frac{1}{\epsilon_n} \left( \frac{\Pr\left(T \in (x, x + \epsilon_n] \mid R\right)}{w_{\theta_0}(x + \epsilon_n)} - \frac{\Pr\left(T \in (x - \epsilon_n, x] \mid R\right)}{w_{\theta_0}(x - \epsilon_n)} \right)$$

Using the same argument as the proof in Lemma 5,

$$\begin{aligned} \frac{1}{\epsilon_n} \Pr\left(T \in (x, x + \epsilon_n] \mid R\right) &= \operatorname{E}\left[\frac{\mathbf{1}\left\{T \in (x, x + \epsilon_n]\right\}}{\epsilon_n} w_{\theta_0}(T)\right] \frac{1}{E[w_{\theta_0}(T)]} \\ &= \operatorname{E}\left[\frac{\mathbf{1}\left\{T \in (x, x + \epsilon_n]\right\}}{\epsilon_n}\right] \frac{w_{\theta_0}(x + \epsilon_n)}{E[w_{\theta_0}(T)]} \\ &+ \operatorname{E}\left[\frac{\mathbf{1}\left\{T \in (x, x + \epsilon_n]\right\}}{\epsilon_n}(w_{\theta_0}(T) - w_{\theta_0}(x + \epsilon_n))\right] \frac{1}{E[w_{\theta_0}(T)]} \end{aligned}$$

Since  $f_T$  is continuous,

$$\mathbf{E}\left[\frac{\mathbf{1}\left\{T\in(x,x+\epsilon_n]\right\}}{\epsilon_n}\right]\frac{1}{E[w_{\theta_0}(T)]} - \mathbf{E}\left[\frac{\mathbf{1}\left\{T\in[x-\epsilon_n,x)\right\}}{\epsilon_n}\right]\frac{1}{E[w_{\theta_0}(T)]} = O(\epsilon_n)$$

The leaves the term:  $\operatorname{E}\left[\frac{\mathbf{1}\{T \in (x, x+\epsilon_n]\}}{\epsilon_n}(w_{\theta_0}(T) - w_{\theta_0}(x+\epsilon_n))\right] \frac{1}{E[w_{\theta_0}(T)]}$ Since  $w_{\theta}(t)$  is by Assumption 3 uniformly continous except on a finite set,

$$\operatorname{E}\left[\frac{\mathbf{1}\left\{T \in (x, x + \epsilon_n]\right\}}{\epsilon_n} (w_{\theta_0}(T) - w_{\theta_0}(x + \epsilon_n))\right] \frac{1}{E[w_{\theta_0}(T)]} = O(\epsilon_n)$$

So we have:  $\mathbb{E}[\mathbf{v}_{n,x}(T) \mid R] = O(\epsilon_n)$  and since the number of elements isn't changing  $||\mathbb{E}[\mathbf{v}_n(T) \mid R]||_{\infty} = O(\epsilon_n).$ 

Now we bound the vectors that  $\mathbb{E}[\mathbf{v}_n(T) \mid R]$  is multiplied by. In the second sum, first recall that the integrals over  $\phi$  are uniformly bounded over j. The matrix  $[G(\theta_0)'G(\theta_0)]^{-1}G(\theta_0)'$  is not changing so we don't need to bound it. It remains to bound the sum of vector norms:

$$\sum_{j=0}^{J_n} \left| \left| E\left[ \psi_j(T) \varphi_{\sigma_Y}(T) \hat{B}'_{\theta_0}(T) \mid R \right] \right| \right|_{\infty}$$

We can write this in terms of inner products:

$$\sum_{j=0}^{J_n} \left| \left| E\left[ \psi_j(T) \varphi_{\sigma_Y}(T) \hat{B}'_{\theta_0}(T) \mid R \right] \right| \right|_{\infty} = \sum_{j=0}^{J_n} \left| \left| \left\langle \psi_j, f_{T|T} \frac{d}{d\theta_k} \frac{1}{w_{\theta}(t)} \right\rangle_Y \right| \right|_{\infty}$$

Assumption 3 says that  $w_{\theta}(t)$  is everywhere differentiable in  $\theta$  and that the derivatives are uniformly bounded. So  $||f_{T|R}\frac{d}{d\theta_k}\frac{1}{w_{\theta}(t)}||_Y \leq ||f_{T|R}||_Y \leq 1$ . So,

$$\sum_{j=0}^{J_n} \left\| E\left[ \psi_j(T) \varphi_{\sigma_Y}(T) \hat{B}'_{\theta_0}(T) \mid R \right] \right\|_{\infty} \le |\mathcal{X}|$$

In the third sum,  $\frac{\beta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)}|R\right]^2}B'_{\theta_0}\left[G(\theta_0)'G(\theta_0)\right]^{-1}G(\theta_0)'$  is a fixed vector.

Since all three terms are now bounded:

$$\tilde{\Delta}_{c,n} - \Delta_c = \mathcal{O}\left(\eta_{J_n} + \epsilon_n\right)$$

## D.12 Proof of Lemma 7

Since it minimizes the sample objective at zero,  $\hat{\pi}_n$  must satisfy:

$$\eta_j \langle \chi_j, \hat{\pi}_n \rangle_W = \frac{1}{\lambda_j} \frac{\frac{1}{n} \sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\hat{\theta}_n}(t_i)}}{\frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\hat{\theta}_n}(t_i)}}$$

Let the deterministic sequence  $r_{1,n}$  be:

$$r_{1,n} \equiv \sum_{j=J_n+1}^{\infty} \eta_j \langle \psi_j, \pi_0 \rangle_W \left( \lambda_j \int_{-cv}^{cv} \phi_j(s) ds - \int_{-cv/c}^{cv/c} \phi_j(s) ds \right)$$

Using Taylor's Theorem for some  $w_{\bar{\theta}_n}$  such that  $E\left[\frac{1}{w_{\bar{\theta}}(T)}\right]$  is between  $E\left[\frac{1}{w_{\hat{\theta}_n}(T)}\right]$  and  $E\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]$ :

$$\begin{split} \frac{1}{\lambda_{j}} \frac{\frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})}}{\frac{1}{n} \sum_{i=1}^{n} \frac{1}{\lambda_{j}} \frac{\frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})}}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]} \\ &- \frac{1}{\mathbb{E}\left[\frac{1}{w_{\bar{\theta}}(T)}\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\hat{\theta}_{n}}(t_{i})} - E\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]\right) \frac{1}{\lambda_{j}} \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})} \\ &= \frac{1}{\lambda_{j}} \frac{\frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})}}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]} \\ &- \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})} - E\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]\right) \frac{1}{\lambda_{j}} \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})} \\ &- \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})} - \frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})}\right) \frac{1}{\lambda_{j}} \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})} \\ &- \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})} - \frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})}\right) \frac{1}{\lambda_{j}} \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})} \\ &- \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})} - \frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})}\right) \frac{1}{\lambda_{j}} \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})} \\ &- \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})}\right) \frac{1}{\lambda_{j}} \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})}} \\ &- \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})}\right) \frac{1}{\lambda_{j}} \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\theta_{n}}(t_{i})}} \\ &- \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})}\right) \frac{1}{\lambda_{j}} \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\theta_{n}}(t_{i})}} \\ &- \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_$$

Since  $\left|\frac{1}{w_{\hat{\theta}_n}(t)} - \frac{1}{w_{\theta_0}(t)}\right| = \mathcal{O}_p\left(n^{-1/3}\right)$  and  $w_{\theta}(t)$  is uniformly bounded away from zero and uniformly bounded above by Assumption 2,  $\frac{1}{\mathbb{E}\left[\frac{1}{w_{\bar{\theta}}(T)}\right]^2} - \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)}|R\right]^2} = \mathcal{O}\left(n^{-1/3}\right)$ . We have already shown that  $\operatorname{Var}(\psi_j(T)\varphi_{\sigma_Y}(T)) \leq 1$  so  $\frac{1}{n}\sum_{i=1}^n \frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\hat{\theta}_n}(t_i)} - \mathbb{E}\left[\frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\theta_0}(t_i)}\right] = \mathcal{O}_p(n^{-1/3})$ . So we have:

$$\begin{split} \eta_{j} \langle \chi_{j}, \hat{\pi}_{n} \rangle_{W} &- \eta_{j} \langle \chi_{j}, \pi_{0} \rangle_{W} = \frac{1}{\lambda_{j}} \frac{\frac{1}{n} \sum_{i=1}^{n} \frac{\psi_{j}(t_{i})\varphi_{\sigma_{Y}}(t_{i})}{w_{\hat{\theta}_{n}}(t_{i})} - E\left[\frac{\psi_{j}(T)\varphi_{\sigma_{Y}}(T)}{w_{\theta_{0}}(T)}\right]}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)}\right]} \\ &- \frac{\mathbb{E}\left[\frac{\psi_{j}(T)\varphi_{\sigma_{Y}}(T)}{w_{\theta_{0}}(T)}\right]}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\theta_{0}}(t_{i})} - E\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]\right) \frac{1}{\lambda_{j}} \\ &- \frac{\mathbb{E}\left[\frac{\psi_{j}(T)\varphi_{\sigma_{Y}}(T)}{w_{\theta_{0}}(T)}\right]}{\mathbb{E}\left[\frac{1}{w_{\theta_{0}}(T)} \mid R\right]^{2}} \left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\hat{\theta}_{n}}(t_{i})} - \frac{1}{w_{\theta_{0}}(t_{i})}\right) \frac{1}{\lambda_{j}} \\ &+ \mathcal{O}_{p}\left(\frac{1}{\lambda_{j}n^{1/3}}\right) \\ &= [D] + [E] + [F] + \mathcal{O}_{p}\left(\frac{1}{\lambda_{j}n^{2/3}}\right) \end{split}$$

Term [E] can already be written as a sample mean minus its expectation. For the other two terms, we need to establish the following facts. We will use the Taylor Theorem. To do so, first define the following two  $K \times 1$  gradient vectors:

$$\hat{B}_{\theta}(t_i) \equiv \nabla_{\theta} \frac{1}{w_{\theta}(t_i)}$$
$$B_{\theta} \equiv \mathbb{E} \left[ \hat{B}_{\theta_0}(T) \mid R \right]$$

Since the gradients and  $\theta_0$  are  $K \times 1$ ,  $B'_{\theta_0} \theta_0$  is a scalar. With the gradients defined, we can use the Taylor Theorem:

$$\frac{1}{w_{\hat{\theta}_n}(t_i)} - \frac{1}{w_{\theta_0}(t_i)} = B'_{\theta_0}(t_i) \left(\hat{\theta}_n - \theta_0\right) + \mathcal{O}_p\left(\left(\hat{\theta}_n - \theta_0\right)^2\right)$$

Since we assumed that each element of  $B_{\theta}$  is uniformly bounded,  $\left\| \frac{1}{n} \sum_{i=1}^{n} \hat{B}_{\theta}(t_i) - \mathbb{E} \left[ \hat{B}_{\theta_0}(T) \mid R \right] \right\|_{\infty} = \mathcal{O}_p(n^{-1/2})$ . This gives us:

$$[F] = \frac{1}{\lambda_j} \frac{\mathbb{E}\left[\frac{\psi_j(T)\varphi_{\sigma_Y}(T)}{w_{\theta_0}(T)} \mid R\right]}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} B'_{\theta_0}\left(\hat{\theta}_n - \theta_0\right) + \mathcal{O}_p\left(\frac{1}{n^{2/3}\lambda_j}\right)$$

To linearize term [D]. First consider its numerator.

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\hat{\theta}_n}(t_i)} - E\left[\frac{\psi_j(T)\varphi_{\sigma_Y}(T)}{w_{\theta_0}(T)} \mid R\right] &= \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\theta_0}(t_i)} - E\left[\frac{\psi_j(T)\varphi_{\sigma_Y}(T)}{w_{\theta_0}(T)} \mid R\right] \\ &+ \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\hat{\theta}_n}(t_i)} - \frac{1}{n} \sum_{i=1}^{n} \frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\theta_0}(t_i)} \\ &= [D1] + [D2] \end{aligned}$$

Term [D1] is a sample mean minus its expectation. Term [D2] can be written as:

$$[D2] = \frac{1}{n} \sum_{i=1}^{n} \psi_j(t_i) \varphi_{\sigma_Y}(t_i) \left( \frac{1}{w_{\hat{\theta}_n}(t_i)} - \frac{1}{w_{\theta_0}(t_i)} \right)$$
$$= \frac{1}{n} \sum_{i=1}^{n} \psi_j(t_i) \varphi_{\sigma_Y}(t_i) \hat{B}'_{\theta_0}(t_i) (\hat{\theta}_n - \theta_0) + \mathcal{O}_p \left( (\hat{\theta}_n - \theta_0)^2 \right)$$
$$= \mathbb{E} \left[ \psi_j(T) \varphi_{\sigma_Y}(T) \hat{B}'_{\theta_0}(T) \mid R \right] (\hat{\theta}_n - \theta_0) + \mathcal{O}_p \left( n^{-2/3} \right)$$

To keep things concise we will need to define the following shorthand. We showed in the proof of Lemma 3 that the integrals are uniformly bounded over all j. Since the  $\lambda_j$  are decreasing this means that  $\sup_j |a_j| < \infty$ .

$$a_j \equiv \lambda_j \int_{-cv}^{cv} \phi_j(s) ds - \int_{-cv/c}^{cv/c} \phi_j(s) ds$$

No we can write the entire esimation error as:

$$\begin{split} \hat{\Delta}_{c,n} - \Delta_c + r_{1,n} &= \frac{1}{\mathbb{E}\left[\frac{1}{|w_{\theta_0}(T)| | R\right]}} \left( \frac{1}{n} \sum_{i=1}^n \sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} \frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\theta_0}(t_i)} - \sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\frac{\psi_j(T)\varphi_{\sigma_Y}(T)}{w_{\theta_0}(T)| | R\right]} \right) \\ &+ \frac{\sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\theta_0}(T) | R\right]}{\mathbb{E}\left[\frac{1}{|w_{\theta_0}(T)| | R\right]^2} (\hat{\theta}_n - \theta_0) \\ &+ \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{|w_{\theta_0}(T)| | R\right]^2} \left(\frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\theta_0}(t_i)} - E\left[\frac{1}{|w_{\theta_0}(T)| | R\right]}\right) \\ &+ \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{|w_{\theta_0}(T)| | R\right]^2} B'_{\theta_0} \left(\hat{\theta}_n - \theta_0\right) \\ &+ \mathcal{O}_p\left(\frac{1}{|\lambda_{J_n}n^{2/3}|}\right) \end{split}$$

To proceed we must now linearize  $\hat{\theta}_n - \theta_0$ . Using the result from section D.12.1 below, where the  $|\mathcal{X}| \times K$  matrix G and  $K \times 1$  vector  $\hat{g}_n$  are defined there:

$$\hat{\theta}_n - \theta_0 = \left[ G(\theta_0)' G_n(\theta_0) \right]^{-1} G(\theta_0)' n^{1/3} \hat{g}_n(\theta_0) + \mathcal{O}_p\left( n^{-2/3} \right)$$

Substituting this into the first term and using the facts that  $\sup_j |a_j| < \infty$  and  $\lambda_J \sum_{j=0}^J \lambda_j^{-1} = O(1)$ :

$$\frac{\sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\theta_0}(T) \mid R\right]}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} (\hat{\theta}_n - \theta_0) \\
= \frac{\sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\theta_0}(T) \mid R\right] [G(\theta_0)'G_n(\theta_0)]^{-1} G(\theta_0)'\hat{g}_n(\theta_0)}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} + \mathcal{O}_p\left(\frac{1}{\lambda_{J_n}n^{2/3}}\right)$$

Linearizing the second term:

$$\frac{\Delta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} B'_{\theta_0} \left(\hat{\theta}_n - \theta_0\right) = \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} B'_{\theta_0} \left[G(\theta_0)'G_n(\theta_0)\right]^{-1} G(\theta_0)'\hat{g}_n(\theta_0) + \mathcal{O}_p\left(n^{-2/3}\right)$$

Putting all of these pieces together:

$$\begin{split} \hat{\Delta}_{c,n} &- \Delta_c + r_{1,n} \\ &= \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]} \left( \frac{1}{n} \sum_{i=1}^n \sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} \frac{\psi_j(t_i)\varphi_{\sigma_Y}(t_i)}{w_{\theta_0}(t_i)} - \sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\frac{\psi_j(T)\varphi_{\sigma_Y}(T)}{w_{\theta_0}(T)} \mid R\right] \right) \\ &+ \frac{\sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\theta_0}(T) \mid R\right] \left[G(\theta_0)'G_n(\theta_0)\right]^{-1} G(\theta_0)'\hat{g}_n(\theta_0)}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} \\ &+ \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} \left(\frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\theta_0}(t_i)} - E\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]\right) \\ &+ \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} B'_{\theta_0} \left[G(\theta_0)'G_n(\theta_0)\right]^{-1} G(\theta_0)'\hat{g}_n(\theta_0) \\ &+ \mathcal{O}_p\left(\frac{1}{\lambda_{J_n}n^{2/3}}\right) \end{split}$$

Now we write down the function  $Z_{n,J_n}$  explicitly:

$$\begin{split} Z_{n,J_n}(t) &= \frac{1}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]} \sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} \frac{\psi_j(t)\varphi_{\sigma_Y}(t)}{w_{\theta_0}(t)} \\ &+ \frac{\sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\theta_0}(T) \mid R\right] [G(\theta_0)'G_n(\theta_0)]^{-1} G(\theta_0)'\mathbf{v}(t)}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} \\ &+ \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} \frac{1}{w_{\theta_0}(t)} \\ &+ \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} B'_{\theta_0} \left[G(\theta_0)'G_n(\theta_0)\right]^{-1} G(\theta_0)'\mathbf{v}(t) \\ &= [G] + [H] + [I] + [J] \end{split}$$

Where  $\mathbf{v}_n(t)$  is an  $|\mathcal{X}| \times 1$  vector with components:

$$\mathbf{v}_{n,x}(t) = \frac{1}{\epsilon_n} \left( \frac{\mathbf{1} \left\{ t \in (x, x + \epsilon_n] \right\}}{w_{\theta_0}(x + \epsilon_n)} - \frac{\mathbf{1} \left\{ t \in (x - \epsilon_n, x] \right\}}{w_{\theta_0}(x - \epsilon_n)} \right)$$

We will establish two facts:

•  $\sup_{t,n} n^{-1/3} \lambda_{J_n} |Z_{n,J_n}(t)| < \infty.$ 

•  $\sup_n \lambda_{J_n}^{-2} n^{-1/3} Var(Z_{n,J_n}(T)) < \infty$ 

First we show that  $\sup_{t,n} n^{-1/3} \lambda_{J_n} |Z_{n,J_n}(t)| < \infty$ . First, consider [G]. The functions  $\psi_j(t)\varphi_{\sigma_Y}(t)$  are called the "Hermite Functions." They are uniformly bounded by Cramér's Inequality which can be found in Indritz (1961) and Bateman (1953):

$$\sup_{j,t} |\psi_j(t)\varphi_{\sigma_Y}(t)| < \pi^{1/2}$$

We have already shown that  $\lambda_{J_n} \sum_{j=0}^{J_n} \frac{1}{\lambda_j} = O(1)$ . So we can conclude:

$$[G] \lesssim \frac{1}{\lambda_{J_n}}$$

For the same reasons:

$$\sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\theta_0}(T) \mid R\right] \left[G(\theta_0)'G_n(\theta_0)\right]^{-1} G(\theta_0)' \lesssim \frac{1}{\lambda_{J_n}}$$

By Assumption 2,

$$\mathbf{v}_{n,x}(t) \lesssim \frac{1}{\epsilon_n} \lesssim n^{1/3}$$

So we conclude that  $\sup_t |[H]| \leq \frac{n^{1/3}}{\lambda_{J_n}}$ . Assumption 2 guarantees that  $\sup_t |[I]| \leq 1$ . Finally, using the same arguments as the for the other parts:  $\sup_t |[J]| \leq n^{1/3}$ . Putting all three parts together,

$$\sup_{t \in \mathbb{R}} \max_{j \in 1, \dots, J_n} |Z_{j, J_n}(t)| \lesssim \frac{n^{1/3}}{\lambda_{J_n}}$$

Now to show  $\sup_n \lambda_{J_n}^{-2} n^{-1/3} Var(Z_{n,J_n}(T)) < \infty$ . The variance of the sum of four terms is no greater than sixteen times the largest of their variances. So we need only show that [G], [H], [I], and [J] all have variances of order  $\lambda_{J_n}^{-2} n^{-1/3}$ . For [G] and [I], the upper bounds already established suffice since the variance cannot be greater than the square of the maximum. For [H] and [J], it is sufficient to establish that  $\max_x Var(\mathbf{v}_{n,x}(T)) \leq n^{1/3}$ . To do this, first use Assumption 2 to bound  $1/w_{\theta_0}$ . Then notice that  $\mathbf{v}_{n,x}(T)$  is  $n^{1/3}$  times a bounded number times the difference of two Bernoulli random variables. The variance of a Bernoulli is p(1-p). Since T must have a continuous distribution  $P(T \in (x, x + \epsilon_n]) \leq \epsilon_n \leq \frac{1}{n^{1/3}}$ . So  $\operatorname{Var}(\mathbf{v}_{n,x}(T)) \leq \frac{n^{2/3}}{n^{1/3}} = n^{1/3}$ . Since there are only finitely many x,  $\max_x Var(\mathbf{v}_{n,x}(T)) \leq n^{1/3}$  and we have established that [H],[J] have variance of order  $\lambda_{J_n}^{-2} n^{-1/3}$ , which is enough to guarantee that  $\sup_n \lambda_{J_n}^{-2} n^{-1/3} Var(Z_{n,J_n}(T)) < \infty$ .

Notice that  $\frac{1}{n} \sum_{i=1}^{n} Z_{n,J_n}(t_i) - \mathbb{E}[Z_{n,J_n}(T)]$  is not asymptotically equivalent to  $\hat{\beta}_{c,n} - \tilde{\beta}_{c,n}$  because  $E[\mathbf{v}_{n,x}(t)] \neq 0$ . We need to subtract this bias term off on the left-hand side.

$$\mathbb{E}\left[\mathbf{v}_{n,x}(t)\right] = \frac{1}{\epsilon_n} \left( \frac{\Pr\left(t \in (x, x + \epsilon_n]\right)}{w_{\theta_0}(x + \epsilon_n)} - \frac{\Pr\left(t \in (x - \epsilon_n, x]\right)}{w_{\theta_0}(x - \epsilon_n)} \right)$$

Now define the bias terms as:

$$r_{2,n} \equiv \frac{\sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\theta_0}(T) \mid R\right] [G(\theta_0)'G_n(\theta_0)]^{-1} G(\theta_0)'E[\mathbf{v}(T) \mid R]}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} + \frac{\Delta_c}{\mathbb{E}\left[\frac{1}{w_{\theta_0}(T)} \mid R\right]^2} B'_{\theta_0} \left[G(\theta_0)'G_n(\theta_0)\right]^{-1} G(\theta_0)'E[\mathbf{v}(T) \mid R]$$

This expectation is  $E[||\mathbf{v}(T)||_{\infty}] = O(\epsilon_n) = \mathcal{O}(n^{-1/3})$ . So we can conclude:

$$\hat{\Delta}_{c,n} - \Delta_0 - r_{1,n} - r_{2,n} = \frac{1}{n} \sum_{i=1}^n Z_{n,J_n}(t_i) - \mathbb{E}\left[Z_{n,J_n}(T)\right] + \mathcal{O}_p\left(n^{-2/3}\lambda_{J_n}\right)$$

# **D.12.1** Linearization of $\hat{\theta}_n - \tilde{\theta}_n$

Next we study the convergence of  $\hat{\theta}_n$ . Since the first term of the sample objective is zero at its minimizer with probability approaching 1,  $\hat{\theta}_n$  satisfies

$$\hat{\theta}_n = \arg\min_{\theta} \sum_{x \in \mathcal{X}} \left( \frac{\frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbf{1} \{ t_i \in (x, x + \epsilon_n] \}}{w_{\theta}(x + \epsilon_n)} - \frac{\frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbf{1} \{ t_i \in (x - \epsilon_n, x] \}}{w_{\theta}(x - \epsilon_n)} \right)^2$$

This form coincides with a classical minimum distance estimator and we can show its asymptotic normality in the standard way from Newey and McFadden (1994). Let  $\hat{g}_n(\theta)$  be a  $|\mathcal{X}| \times 1$  vector with components equal to the summands  $\left(\frac{\frac{1}{n\epsilon_n}\sum_{i=1}^n \mathbf{1}\{t_i \in (x,x+\epsilon_n)\}}{w_{\theta}(x+\epsilon_n)} - \frac{\frac{1}{n\epsilon_n}\sum_{i=1}^n \mathbf{1}\{t_i \in (x-\epsilon_n,x]\}}{w_{\theta}(x-\epsilon_n)}\right)$ . So,

$$\hat{\theta}_n = \arg\min_{\theta} \hat{g}_n(\theta)' \hat{g}_n(\theta)$$

Since  $w_{\theta}(t)$  is assumed to be everywhere continuously differentiable in  $\theta$ ,  $\hat{g}_n(\theta)$  is also everywhere continuously differentiable in  $\theta$ . Let  $\hat{G}(\theta)$  be the  $|\mathcal{X}| \times K$  matrix of first derivatives of  $\hat{g}_n(\theta)$  with respect to the K components of  $\theta$ .

$$\hat{G}(\theta)_{x,k} = -\frac{\frac{1}{n\epsilon_n}\sum_{i=1}^n \mathbf{1}\left\{t_i \in (x, x+\epsilon_n)\right\}}{w_\theta(x+\epsilon_n)^2} \frac{d}{d\theta} w_\theta(x+\epsilon_n) + \frac{\frac{1}{n\epsilon_n}\sum_{i=1}^n \mathbf{1}\left\{t_i \in (x-\epsilon_n, x]\right\}}{w_\theta(x-\epsilon_n)^2} \frac{d}{d\theta} w_\theta(x-\epsilon_n)$$

The population gradient  $G(\theta)$  is:

$$G(\theta)_{x,k} = -\lim_{\epsilon \to 0} \frac{f_{T|R}(x+\epsilon)}{w_{\theta}(x+\epsilon_n)^2} \frac{d}{d\theta} w_{\theta}(x+\epsilon) + \frac{f_{T|R}(x-\epsilon)}{w_{\theta}(x-\epsilon)^2} \frac{d}{d\theta} w_{\theta}(x-\epsilon)$$

The sample gradients converge uniformly to the population gradient:

$$\max_{x \in \mathcal{X}, k \in \{1, \dots K\}} \sup_{\theta} n^{1/3} \left| \hat{G}(\theta)_{x,k} - G(\theta)_{x,k} \right| = \mathcal{O}_p(1)$$

So at the solution the FOC is satisfied:

$$\mathbf{0} = \hat{G}(\hat{\theta}_n)'\hat{g}_n(\hat{\theta}_n)$$

Taking the Taylor expansion about  $\tilde{\theta}$ :

$$\hat{g}_n(\hat{\theta}_n) - \hat{g}_n(\theta_0) = \hat{G}_n(\bar{\theta}_n)(\hat{\theta}_n - \tilde{\theta}_n)$$

Substituting this into the FOC:

$$\mathbf{0} = \hat{G}(\hat{\theta}_n)'\hat{g}_n(\theta_0) + \hat{G}(\hat{\theta}_n)'\hat{G}_n(\bar{\theta}_n)(\hat{\theta}_n - \tilde{\theta}_n)$$

Solving and using the nonsingularity of G,

$$(\hat{\theta}_n - \tilde{\theta}_n) = \left[\hat{G}(\hat{\theta}_n)'\hat{G}_n(\bar{\theta}_n)\right]^{-1}\hat{G}(\hat{\theta}_n)'\hat{g}_n(\theta_0)$$

Since  $(\tilde{\theta}_n - \theta_0) \to 0$  and  $\bar{\theta}$  is a mean value and the derivatives of  $w_{\theta}$  in  $\theta$  are continuous and the fact that matrix inverses are continuous in the 2-norm at invertible matrices:

$$\left\| \left[ G(\theta_0)'G_n(\theta_0) \right]^{-1} G(\theta_0)' - \left[ \hat{G}(\hat{\theta}_n)'\hat{G}_n(\bar{\theta}_n) \right]^{-1} \hat{G}(\hat{\theta}_n)' \right\|_2 \to 0$$

Since the dimensions of G are fixed and each dimension is converging with  $n^{-1/3}$ ,  $n^{1/3}||\hat{G}(\hat{\theta}_n) - G(\theta_0)||_F = \mathcal{O}_p(1)$ . Since the Frobenius norm upper bounds the 2-norm,  $n^{1/3}||\hat{G}(\hat{\theta}_n) - G(\theta_0)||_2 = \mathcal{O}_p(1)$ . Since matrix multiplication and inversion are Lipschitz continuous with respect to the 2-norm:

$$n^{1/3} \left\| \left[ G(\theta_0)'G_n(\theta_0) \right]^{-1} G(\theta_0)' - \left[ \hat{G}(\hat{\theta}_n)'\hat{G}_n(\bar{\theta}_n) \right]^{-1} \hat{G}(\hat{\theta}_n)' \right\|_2 = \mathcal{O}_p(1)$$

Since  $n^{1/3}\hat{g}_n(\theta_0) = \mathcal{O}_p(1),$ 

$$n^{1/3}(\hat{\theta}_n - \tilde{\theta}_n) = \left[ G(\theta_0)' G_n(\theta_0) \right]^{-1} G(\theta_0)' n^{1/3} \hat{g}_n(\theta_0) + \mathcal{O}_p\left( n^{-1/3} \right)$$

But  $\hat{g}_n(\theta_0)$  is a vector of sample averages.

## D.13 Proof of Theorem 4

Define:

$$\begin{split} \hat{Z}_{n,J_n}(t) &= \frac{1}{\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\hat{\theta}_n}(t_i)}} \sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} \frac{\psi_j(t)\varphi_{\sigma_Y}(t)}{w_{\hat{\theta}_n}(t)} \\ &+ \frac{\sum_{j=0}^{J_n} \frac{a_j}{\lambda_j} E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\hat{\theta}_n}(T) \mid R\right] \left[G(\hat{\theta}_n)'G_n(\hat{\theta}_n)\right]^{-1} G(\hat{\theta}_n)' \mathbf{v}_{\hat{\theta}_n}(t)}{\left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\hat{\theta}_n}(t_i)}\right)^2} \\ &+ \frac{\hat{\Delta}_{c,n}}{\left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\hat{\theta}_n}(t_i)}\right)^2} \frac{1}{w_{\hat{\theta}_n}(t)} \\ &+ \frac{\hat{\Delta}_{c,n}}{\left(\frac{1}{n} \sum_{i=1}^{n} \frac{1}{w_{\hat{\theta}_n}(t_i)}\right)^2} B'_{\hat{\theta}_n} \left[G(\hat{\theta}_n)'G_n(\hat{\theta}_n)\right]^{-1} G(\hat{\theta}_n)' \mathbf{v}_{\hat{\theta}_n}(t) \end{split}$$

The proof proceeds in two steps. First we show that  $\sup_{t \in \mathbb{R}} n^{1/3} \lambda_{J_n} |\hat{Z}_{n,J_n}(t) - Z_{n,J_n}(t)| = \mathcal{O}_p(1)$ . Then we show that the variance estimator converges faster than the variance decays. Step 1

First we want to show that:  $\sup_{t \in \mathbb{R}} n^{1/3} \lambda_{J_n} |\hat{Z}_{n,J_n}(t) - Z_{n,J_n}(t)| = \mathcal{O}_p(1)$ . We will show that each term above converges at this rate one by one.

First, we have already shown in Theorem 1 that  $\hat{\Delta}_{c,n} - \Delta_c = \mathcal{O}_p\left(\lambda_{J_n}n^{-1/3}\right)$ . We showed in step 1 of the proof of Theorem 2 that  $\sup_t n^{1/3} \left| \frac{1}{w_{\theta_n}(t)} - \frac{1}{w_{\theta_0}(t)} \right| = \mathcal{O}_p(1)$ . This implies that  $\sup_t n^{1/3} ||\mathbf{v}_{\theta_n}(t) - \mathbf{v}_{\theta_0}(t)||_{\infty} = \mathcal{O}_p(1)$ . By the law of large numbers,  $\frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\theta_n}(t_i)} - E\left[\frac{1}{w_{\theta_0}(T)}\right] = \mathcal{O}_p\left(n^{-1/2}\right)$ . By Assumption 2,  $\frac{1}{w_{\theta(t)}} > \overline{\theta}$  for all  $t, \theta$ , so  $\left| \frac{1}{\frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\theta_n}(t_i)}} - E\left[\frac{1}{w_{\theta_0}(T)}\right] \right| = \mathcal{O}_p(n^{-1/3})$ . Since all quantities are uniformly bounded by  $\overline{\theta}$ ,  $\left| \frac{1}{\frac{1}{n} \sum_{i=1}^n \frac{1}{w_{\theta_n}(t_i)}} - E\left[\frac{1}{w_{\theta_0}(T)}\right] \right|^2 = \mathcal{O}_p(n^{-1/3})$ . Next, we showed in the proof of Lemma 7 that  $n^{1/3} \left| \left| \left[ G(\hat{\theta}_n)'G_n(\hat{\theta}_n) \right]^{-1} G(\hat{\theta}_n)' \right| \right|_2 = \mathcal{O}_p(1)$ . Finally, since we assumed that  $\nabla_{\theta} \frac{1}{w_{\theta(t)}}$  is Lipschitz-continuous in  $\theta$ ,  $\sup_t n^{1/3} \left| \left| \hat{B}_{\theta_n}(t) - \right| \right|_1 < \infty$ . Combining this with the fact that  $\psi_j(t)\varphi_{\sigma_Y}(t) < \pi^{1/4}$  from Indritz (1961),

$$\left\| E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\hat{\theta}_n}(T) \mid R\right] - E\left[\psi_j(T)\varphi_{\sigma_Y}(T)\hat{B}'_{\theta_0}(T) \mid R\right] \right\|_1 = \mathcal{O}\left(n^{-1/3}\right)$$

Combining all of these facts, we conclude that  $\sup_{t \in \mathbb{R}} n^{1/3} \lambda_{J_n} |\hat{Z}_{n,J_n}(t) - Z_{n,J_n}(t)| = \mathcal{O}_p(1).$ Step 2

# Now we show the main result. $n^{5/6} \lambda_{J_n}^2$ times the difference between variance estimator and the variance estimator if we knew $Z_{n,J_n}$ is bounded by:

$$n^{5/6} \lambda_{J_n}^2 \left| \frac{1}{n^2} \sum_{i=1}^n \sum_{k=1}^n \Lambda_{ik} Z_{n,J_n}(t_i) Z_{n,J_n}(t_k) - \frac{1}{n^2} \sum_{i=1}^n \sum_{k=1}^n \Lambda_{ik} \hat{Z}_{n,J_n}(t_i) \hat{Z}_{n,J_n}(t_k) \right|$$
$$= \mathcal{O}_p \left( n^{5/6} \lambda_{J_n}^2 \frac{1}{n} \right) = \mathcal{O}_p(1)$$

But by Assumption 5,  $\liminf n^{5/6} \lambda_{J_n}^2 Var(\hat{\Delta}_{n,c}) = \infty$ , so this difference must be dominated by the variance itself. Now consider the estimation error. Since  $\operatorname{Var}(Z_{n,J_n}(t_i)) = O(\lambda_{J_n}^{-2}n^{1/3})$ , then  $\operatorname{Var}(Z_{n,J_n}(t_i)^2) = O(\lambda^{-4}n^{4/3})$ .

$$\frac{1}{n}\sum_{i=1}^{n}\sum_{k=1}^{n}\Lambda_{ik}Z_{n,J_n}(t_i)Z_{n,J_n}(t_k) - \frac{1}{n}\sum_{i=1}^{n}\sum_{k=1}^{n}\Lambda_{ik}\mathbb{E}[Z_{n,J_n}(t_i)Z_{n,J_n}(t_k)] = \mathcal{O}_p\left(\frac{n^{2/3}\lambda_{J_n}^{-2}}{n^{1/2}}\right)$$
$$= \mathcal{O}_p\left(n^{1/6}\lambda_{J_n}^{-2}\right)$$

So dividing by a further n yields:

$$\frac{1}{n^2} \sum_{i=1}^n \sum_{k=1}^n \Lambda_{ik} Z_{n,J_n}(t_i) Z_{n,J_n}(t_k) - \frac{1}{n^2} \sum_{i=1}^n \sum_{k=1}^n \Lambda_{ik} \mathbb{E}[Z_{n,J_n}(t_i) Z_{n,J_n}(t_k)] = \mathcal{O}_p\left(n^{-5/6} \lambda_{J_n}^{-2}\right)$$

Multiplying by  $n^{5/6}\lambda_{J_n}^2$ :

$$n^{5/6}\lambda_{J_n}^2\left(\frac{1}{n^2}\sum_{i=1}^n\sum_{k=1}^n\Lambda_{ik}Z_{n,J_n}(t_i)Z_{n,J_n}(t_k) - \frac{1}{n^2}\sum_{i=1}^n\sum_{k=1}^n\Lambda_{ik}\mathbb{E}[Z_{n,J_n}(t_i)Z_{n,J_n}(t_k)]\right) = \mathcal{O}_p(1)$$

Again by Assumption 5,  $\liminf n^{5/6} \lambda_{J_n}^2 Var(\hat{\Delta}_{n,c}) = \infty$ , so this difference is also dominated by the variance itself. So we can conclude,

$$\frac{\frac{1}{n^2} \sum_{i=1}^n \sum_{k=1}^n \Lambda_{ik} \hat{Z}_{n,J_n}(t_i) \hat{Z}_{nJ_n}(t_k)}{\operatorname{Var}(\hat{\Delta}_{n,c})} \to 1$$