Free energy equivalence between mean-field models and nonsparsely diluted mean-field models

Manaka Okuyama¹ and Masayuki Ohzeki^{1,2,3}

¹Graduate School of Information Sciences, Tohoku University, Sendai 980-8579, Japan ²Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan and ³Sigma-i Co., Ltd., Tokyo 108-0075, Japan

(Dated: June 25, 2024)

Abstract

We studied nonsparsely diluted mean-field models that differ from sparsely diluted mean-field models, such as the Viana–Bray model. We prove that the free energy of nonsparsely diluted mean-field models coincides exactly with that of the corresponding mean-field models with different parameters in ferromagnetic and spin-glass models composed of any discrete spin *S* in the thermodynamic limit. Our results are a broad generalization of the results of a previous study [Bovier and Gayrard, J. Stat. Phys. 72, 643 (1993)], where the densely diluted mean-field ferromagnetic Ising model (diluted Curie–Weiss model) was analyzed rigorously, and it was proven that its free energy was exactly equivalent to that of the corresponding mean-field model (Curie–Weiss model) with different parameters.

I. INTRODUCTION

The importance of understanding the effects of randomness on the system properties has increased over the past few decades. The spin-glass model is a typical example of randomness in statistical mechanics. The concept of replica symmetry breaking in mean-field models [1] has played an important role in statistical mechanics and information science.

In statistical mechanics, randomness can be introduced not only by randomizing the interaction strength, as in spin-glass models, but also by diluting the interaction. Diluted mean-field models can be defined in three ways: The first definition is sparsely diluted mean-field models, where the strength of the interaction is O(1) and the existence probability of each edge is $O(N^{-1})$. A concrete example of sparsely diluted mean-field models is the spin model on the Erdős-Rényi random graph (such as the Viana–Bray model [2]). The sparsely diluted mean-field models are closely related to information science problems [3–6].

The second definition of diluted mean-field models considers a densely diluted mean-field model, where the strength of the interaction is $O(N^{-1})$ and the existence probability of each edge is O(1). Densely diluted mean-field models are naturally defined from a statistical mechanics perspective. As the number of interactions is proportional to $O(N^2)$, the properties of these models are expected to be similar to those of the corresponding mean-field model. Bovier and Gayrard [7] proved that the free energy of the densely diluted Curie–Weiss model coincides exactly with that of a Curie–Weiss model with different parameters in the thermodynamic limit. Densely diluted mean-field models have not been studied well [8–10], but progress has been made recently [11–17]. The zero-temperature properties of the densely diluted Sherrington–Kirkpatrick (SK) model (not the Viana–Bray model) were numerically investigated in Refs. [16]. It was revealed that the ground-state energy coincides with that of the SK model and depends neither on the distribution of interactions nor on the concentration of dilution. Interestingly, this universal behavior appears to be within the limit of the dilution concentration $\alpha \rightarrow 0$. However, the thermodynamic properties have not been clarified at finite temperatures.

The third definition of diluted mean-field models considers an intermediate regime between sparse and dense dilution, where the strength of the interaction is $O(N^{-b})$ and the existence probability of each edge is $O(N^{b-1})$ with 0 < b < 1. Recent studies [18, 19] showed that the thermodynamic properties of this intermediate regime are equivalent to those of the corresponding mean-field models if the distribution of the existence probability of each edge is trivial such as a Bernoulli distribution. In addition, if the distribution of the existence probability of each edge is nontrivial, it was shown that the thermodynamic properties of the intermediate regime differ from those of sparsely and densely diluted mean-field models.

In the present study, we prove that the free energy of nonsparsely diluted mean-field models $(0 < b \le 1)$ is exactly equal to that of the corresponding mean-field models for both ferromagnetic and spin-glass models composed of an arbitrary discrete spin in the thermodynamic limit. Thus, it is sufficient to analyze the corresponding mean-field models to investigate the thermodynamic properties of nonsparsely diluted mean-field models. Our result is a broad generalization of a previous study by Bovier and Gayrard for the densely diluted Curie–Weiss model [7]. The present proof is based on the free energy equivalence between sparsely diluted mean-field models in the infinite connectivity limit and the corresponding mean-field models using the interpolation method [20, 21].

The remainder of this paper is organized as follows. In Sec. II, we define the model and present the main results (Theorem 1). Section III presents the proof of Theorem 1. Finally, a discussion is presented in Sec. IV.

II. MODELS AND RESULT

We define the nonsparsely diluted mean-field ferromagnetic model as

$$H_{\rm dMF, F} = -\frac{p!}{2\alpha N^{b(p-1)}} \sum_{i_1 < \dots < i_p} K_{i_1 \cdots i_p} S_{i_1} \cdots S_{i_p}, \qquad (1)$$

where $0 < \alpha \le 1$, $0 < b \le 1$, *p* is any positive integer, S_i takes any bounded discrete value with $|S_i| \le C < \infty$, and $K_{i_1 \dots i_p}$ are independent and identically distributed (i.i.d.) random variables following a Bernoulli distribution, with

$$\mathbb{E}[K_{i_1\cdots i_n}] = \alpha N^{(b-1)(p-1)},\tag{2}$$

where $\mathbb{E}[\cdots]$ denotes the expectation with respect to all the random variables. Note that b = 1 and 0 < b < 1 correspond to the densely diluted mean-field model and intermediately diluted mean-field model, respectively (the case b = 0 corresponds to sparsely diluted mean-field models but is not treated in the present study). Similarly, the nonsparsely diluted mean-field spin-glass model is defined as follows:

$$H_{\text{dMF, SG}} = -\sqrt{\frac{p!}{2\alpha N^{b(p-1)}}} \sum_{i_1 < \dots < i_p} K_{i_1 \cdots i_p} J_{d, i_1 \cdots i_p} S_{i_1} \cdots S_{i_p}, \qquad (3)$$

where we consider the following two probability distribution cases of $J_{d,i_1\cdots i_p}$: (i) a Gaussian distribution $N\left(J_0\sqrt{p!/(2\alpha N^{b(p-1)})}, 1\right)$ and (ii) any bounded discrete probability distribution with

$$\mathbb{E}[J_{d,i_1\cdots i_p}] = J_0 \sqrt{\frac{p!}{2\alpha N^{b(p-1)}}},\tag{4}$$

$$\mathbb{E}[J_{d,i_1\cdots i_p}^2] = 1, \tag{5}$$

$$\mathbb{E}[J^n_{d,i_1\cdots i_p}] < \infty \quad (n \ge 3).$$
(6)

The partition functions of the nonsparsely diluted mean-field models are given by

$$Z_{\rm dMF, F} = {\rm Tr}(e^{-\beta H_{\rm dMF, F}}), \qquad (7)$$

$$Z_{\rm dMF, SG} = {\rm Tr}(e^{-\beta H_{\rm dMF, SG}}), \qquad (8)$$

where $Tr(\dots)$ denotes the summation with respect to all spin variables, and β is the inverse temperature. The quenched free energies of the nonsparsely diluted mean-field models are defined as

$$f_{\rm dMF, F} = -\lim_{N \to \infty} \frac{1}{N\beta} \mathbb{E}[\log Z_{\rm dMF, F}], \qquad (9)$$

$$f_{\rm dMF, SG} = -\lim_{N \to \infty} \frac{1}{N\beta} \mathbb{E}[\log Z_{\rm dMF, SG}].$$
(10)

The Hamiltonians of the corresponding mean-field ferromagnetic and spin-glass models are defined as

$$H_{\rm MF, F} = -\frac{p!}{2N^{p-1}} \sum_{i_1 < \dots < i_p} S_{i_1} \cdots S_{i_p}, \qquad (11)$$

$$H_{\rm MF, SG} = -\sqrt{\frac{p!}{2N^{p-1}}} \sum_{i_1 < \dots < i_p} J_{i_1 \cdots i_p} S_{i_1} \cdots S_{i_p}, \qquad (12)$$

where $J_{i_1 \cdots i_p}$ are i.i.d. random variables following a Gaussian distribution, $N(J_0 \sqrt{p!/(2N^{p-1})}, 1)$.

The quenched free energies of the corresponding mean-field models are defined as follows:

$$f_{\rm MF, F} = -\lim_{N \to \infty} \frac{1}{N\beta} \mathbb{E}[\log Z_{\rm MF, F}], \qquad (13)$$

$$f_{\rm MF, SG} = -\lim_{N \to \infty} \frac{1}{N\beta} \mathbb{E}[\log Z_{\rm MF, SG}].$$
(14)

Our results are as follows:

Theorem 1. Let $\alpha \in (0, 1]$ and $b \in (0, 1]$ be functions of N such that $\alpha N^{b(p-1)} \to \infty$ is $N \to \infty$. Then,

$$f_{MF,F} = f_{dMF,F}, \tag{15}$$

$$f_{MF,SG} = f_{dMF,SG}.$$
 (16)

Remark 2. The free energy of the densely diluted ferromagnetic Ising model (b = 1 and p = 2) coincides exactly with that of the Curie–Weiss model [7]. Theorem 1 extends this result to any bounded discrete spin, p-body interaction, intermediately diluted regime (0 < b < 1), and spin-glass model.

III. PROOF OF THEOREM 1

We provide a proof only for the spin-glass model (a similar proof also applies to the ferromagnetic model). The interpolating pressure function is defined as

$$A_N(t) = \frac{1}{N} \mathbb{E}\left[\log \operatorname{Tr}\left(e^{\beta \sqrt{\frac{p!}{2aN^{b(p-1)}}\sum_{i_1 < \dots < i_p} K_{i_1 \cdots i_p}(t) J_{d,i_1 \cdots i_p} S_{i_1} \cdots S_{i_p} + \beta \sqrt{\frac{p!}{2N^{p-1}}\sum_{i_1 < \dots < i_p} J_{i_1 \cdots i_p}(t) S_{i_1} \cdots S_{i_p}}\right)\right], \quad (17)$$

where $K_{i_1\cdots i_p}(t)$ follows a Bernoulli distribution with $\mathbb{E}[K_{i_1\cdots i_p}(t)] = t\alpha N^{(b-1)(p-1)}$, and $J_{i_1\cdots i_p}(t)$ follows a Gaussian distribution $N\left((1-t)J_0\sqrt{p!/(2\alpha N^{p-1})}, (1-t)\right)$. Note that

$$\lim_{N \to \infty} A_N(0) = -\beta f_{\text{MF, SG}},\tag{18}$$

$$\lim_{N \to \infty} A_N(1) = -\beta f_{\text{dMF, SG}}.$$
(19)

The following relationship is useful for any function f(x)

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}[f(K_{i_1\cdots i_p}(t))] = \alpha N^{(b-1)(p-1)}\left(f(1) - f(0)\right).$$
(20)

Using Eq. (20) and integration by parts, we obtain

$$\frac{\mathrm{d}A_{N}(t)}{\mathrm{d}t} = \frac{1}{N} \mathbb{E} \left[\alpha N^{(b-1)(p-1)} \sum_{i_{1} < \dots < i_{p}} \log \left\langle e^{\beta \sqrt{\frac{p!}{2aN^{b(p-1)}}} J_{d,i_{1} \cdots i_{p}} S_{i_{1}} \cdots S_{i_{p}}} \right\rangle_{t,-(i_{1} \cdots i_{p})} \right] \\
- \frac{1}{N} \mathbb{E} \left[\frac{p!}{2N^{p-1}} J_{0}\beta \sum_{i_{1} < \dots < i_{p}} \langle S_{i_{1}} \cdots S_{i_{p}} \rangle_{t} + \frac{\beta^{2} p!}{4N^{p-1}} \sum_{i_{1} < \dots < i_{p}} \left(\langle S_{i_{1}}^{2} \cdots S_{i_{p}}^{2} \rangle_{t} - \langle S_{i_{1}} \cdots S_{i_{p}} \rangle_{t}^{2} \right) \right],$$
(21)

where $\langle \cdots \rangle_t$ is the thermal average with respect to the interpolating Hamiltonian (17), and $\langle \cdots \rangle_{t,-(i_1\cdots i_p)}$ is the thermal average with respect to the interpolating Hamiltonian (17), except for the interaction $\sqrt{p!/(2\alpha N^{b(p-1)})}K_{i_1\cdots i_p}(t)J_{d,i_1\cdots i_p}S_{i_1}\cdots S_{i_p}$.

A. Case where $J_{d,i_1\cdots i_p}$ follows any bounded discrete probability distribution

As the values of $J_{d,i_1\cdots i_p}$ and S_{i_j} are bounded, the following inequality holds by choosing N to be sufficiently large:

$$|e^{\beta\sqrt{\frac{p!}{2\alpha N^{b(p-1)}}}J_{d,i_1\cdots i_p}|S_{i_1}\cdots S_{i_p}|} - 1| < 1.$$
(22)

We can then expand the logarithmic function as

$$\log \left\langle e^{\beta \sqrt{\frac{p!}{2\alpha N^{b(p-1)}} J_{d,i_{1}\cdots i_{p}} S_{i_{1}}\cdots S_{i_{p}}}} \right\rangle_{t,-(i_{1}\cdots i_{p})}$$
(23)
= $\log \left(1 + \sum_{n=1}^{\infty} \frac{1}{n!} \left(\beta \sqrt{\frac{p!}{2\alpha N^{b(p-1)}} J_{d,i_{1}\cdots i_{p}}} \right)^{n} \langle S_{i_{1}}^{n} \cdots S_{i_{p}}^{n} \rangle_{t,-(i_{1}\cdots i_{p})} \right)$
= $\sum_{l=1}^{\infty} \frac{(-1)^{l-1}}{l} \left(\sum_{n=1}^{\infty} \frac{1}{n!} \left(\beta \sqrt{\frac{p!}{2\alpha N^{b(p-1)}} J_{d,i_{1}\cdots i_{p}}} \right)^{n} \langle S_{i_{1}}^{n} \cdots S_{i_{p}}^{n} \rangle_{t,-(i_{1}\cdots i_{p})} \right)^{l}.$ (24)

From Eqs. (4), (5), and (6), the leading-order term of Eq. (24) is as follows: l = n = 1

$$\mathbb{E}\left[\beta\sqrt{\frac{p!}{2\alpha N^{b(p-1)}}}J_{d,i_{1}\cdots i_{p}}\langle S_{i_{1}}\cdots S_{i_{p}}\rangle_{t,-(i_{1}\cdots i_{p})}\right] = \mathbb{E}\left[\beta\frac{p!}{2\alpha N^{b(p-1)}}J_{0}\langle S_{i_{1}}\cdots S_{i_{p}}\rangle_{t,-(i_{1}\cdots i_{p})}\right], \quad (25)$$

$$l = 1, n = 2$$

$$\mathbb{E}\left[\frac{1}{2}\left(\beta\sqrt{\frac{p!}{2\alpha N^{b(p-1)}}}J_{d,i_{1}\cdots i_{p}}\right)^{2}\langle S_{i_{1}}^{2}\cdots S_{i_{p}}^{2}\rangle_{t,-(i_{1}\cdots i_{p})}\right] = \mathbb{E}\left[\frac{\beta^{2}p!}{4\alpha N^{b(p-1)}}\langle S_{i_{1}}^{2}\cdots S_{i_{p}}^{2}\rangle_{t,-(i_{1}\cdots i_{p})}\right], \quad (26)$$

and l = 2, n = 1

$$\mathbb{E}\left[\frac{-1}{2}\left(\beta\sqrt{\frac{p!}{2\alpha N^{b(p-1)}}}J_{d,i_1\cdots i_p}\langle S_{i_1}\cdots S_{i_p}\rangle_{t,-(i_1\cdots i_p)}\right)^2\right] = \mathbb{E}\left[-\frac{\beta^2 p!}{4\alpha N^{b(p-1)}}\langle S_{i_1}\cdots S_{i_p}\rangle_{t,-(i_1\cdots i_p)}^2\right].$$
(27)

Thus, we can rewrite Eq. (21) as follows:

$$\frac{dA_{N}(t)}{dt} = \frac{1}{N} \mathbb{E} \left[\frac{p!}{2N^{p-1}} J_{0}\beta \sum_{i_{1} < \dots < i_{p}} (\langle S_{i_{1}} \cdots S_{i_{p}} \rangle_{t, -(i_{1} \cdots i_{p})} - \langle S_{i_{1}} \cdots S_{i_{p}} \rangle_{t}) \right] \\
+ \frac{1}{N} \mathbb{E} \left[\frac{\beta^{2} p!}{4N^{p-1}} \sum_{i_{1} < \dots < i_{p}} (\langle S_{i_{1}}^{2} \cdots S_{i_{p}}^{2} \rangle_{t, -(i_{1} \cdots i_{p})} - \langle S_{i_{1}} \cdots S_{i_{p}} \rangle_{t, -(i_{1} \cdots i_{p})}^{2} - \langle S_{i_{1}}^{2} \cdots S_{i_{p}}^{2} \rangle_{t} + \langle S_{i_{1}} \cdots S_{i_{p}} \rangle_{t}^{2} \right] \\
+ O(\alpha^{-1/2} N^{-b(p-1)/2}).$$
(28)

Furthermore, it is easy to verify the following:

$$\mathbb{E}[\langle \cdots \rangle_t] = \mathbb{E}[\langle \cdots \rangle_{t, -(i_1 \cdots i_p)}] + O(\alpha^{-1/2} N^{-b(p-1)/2}).$$
⁽²⁹⁾

Consequently, we obtain

$$\frac{\mathrm{d}A_N(t)}{\mathrm{d}t} = O(\alpha^{-1/2} N^{-b(p-1)/2}). \tag{30}$$

Finally, taking the thermodynamic limit $N \to \infty$ such that $\alpha N^{b(p-1)} \to \infty$, we obtain Finally, taking the thermodynamic limit $N \to \infty$ such that $\alpha N^{b(p-1)} \to \infty$, we obtain

$$\lim_{N \to \infty} A_N(0) = \lim_{N \to \infty} A_N(1).$$
(31)

This is the proof of Theorem 1.

B. Case where $J_{d,i_1\cdots i_p}$ follows the Gaussian distribution

The procedure of the proof is almost the same but we have to pay attention to expanding the logarithmic function (23) as $J_{d,i_1\cdots i_p}$ is defined on $(-\infty,\infty)$. For the Gaussian distribution, $N\left(J_0\sqrt{p!/(2\alpha N^{b(p-1)})},1\right)$, Mills' inequality [22] enables us to evaluate whether the probability of $J_{d,i_1\cdots i_p}$ is extremely large, such that

$$|e^{\beta \sqrt{\frac{p!}{2\alpha N^{b(p-1)}} J_{d,i_1 \cdots i_p} |S_{i_1} \cdots S_{i_p}|} - 1| \ge 1,$$
(32)

is exponentially small $O\left(e^{-\left(\log 2/(\beta C^p)\right)^2 \alpha N^{b(p-1)}/p!}\right)$. Thus, we can expand the logarithmic function (23) as in the case of bounded discrete probability distribution, and the calculations are the same.

IV. DISCUSSIONS

We rigorously proved that the free energies of the densely diluted mean-field models (b = 1) and intermediately diluted mean-field models (0 < b < 1) exactly coincide with that of the corresponding mean-field models with different parameters in both ferromagnetic and spin-glass models composed of any discrete spin *S* in the thermodynamic limit.

Note that the value of α is allowed to be close to zero, as long as the condition $\alpha N^{b(p-1)} \rightarrow \infty$ is satisfied within the thermodynamic limit. This explains why the ground-state energy of the densely diluted SK model (b = 1 and p = 2) coincides with that of the SK model, and depends neither on the distribution of interactions nor on the dilution concentration, even within the limit $\alpha \rightarrow 0$ [16]. Furthermore, our results determine the thermodynamic properties of intermediately diluted mean-field models (0 < b < 1) at finite temperatures. This rigorously confirms recent

studies [18, 19], where the thermodynamic properties of the intermediate regime coincide with those of the corresponding mean-field models if the distribution of the existence probability of each edge is trivial such as a Bernoulli distribution.

ACKNOWLEDGMENT

The authors are grateful to Fernando Lucas Metz for useful comments. This study was supported by JSPS KAKENHI Grant Nos. 24K16973 and 23H01432. Our study received financial support from the Public\Private R&D Investment Strategic Expansion PrograM (PRISM) and programs for bridging the gap between R&D and IDeal society (Society 5.0) and Generating Economic and social value (BRIDGE) from the Cabinet Office.

- [1] G. Parisi, Infinite Number of Order Parameters for Spin-Glasses, Phys. Rev. Lett. 43, 1754 (1979).
- [2] L. Viana and A. J. Bray, Phase diagrams for dilute spin-glasses, J. Phys. C, 18, 3037 (1985).
- [3] R. Monasson, R. Zecchina, S. Kirkpatrick, B. Selman, and L. Troyansky, Determining computational complexity from characteristic 'phase transitions, Nature 400, 133 (1999).
- [4] S. Franz, M. Leone, F. R. Tersenghi, and R. Zecchina, Exact Solutions for Diluted Spin Glasses and Optimization Problems, Phys. Rev. Lett. 87, 127209 (2001).
- [5] M. Mezard, G. Parisi, and R. Zecchina, Analytic and algorithmic solution of random satisfiability problems, Science 297, 812 (2002).
- [6] A. Montanari and M. Mézard, Information, Physics and Computation (Oxford Univ. Press, 2009).
- [7] A. Bovier and V. Gayrard, The thermodynamics of the Curie-Weiss model with random couplings, J. Stat. Phys. 72, 643 (1993).
- [8] H. Sompolinsky, Neural networks with nonlinear synapses and a static noise, Phys. Rev. A 34, 2571 (1986).
- [9] A. Canning and J-P. Naef, Phase diagrams and the instability of the spin glass states for the diluted Hopfield neural network model, J. Phys. I France 2, 1791 (1992).
- [10] F.L. Metz and W.K. Theumann, Feed-forward chains of recurrent attractor neural networks with finite dilution near saturation, Physica A 368, 273 (2006).

- [11] J. Barré, A. Ciani, D. Fanelli, F. Bagnoli, and S. Ruffo, Finite size effects for the Ising model on random graphs with varying dilution, Phys. A 388, 3413 (2009).
- [12] Z. Kabluchko, M. Löwe, and K. Schubert, Fluctuations of the Magnetization for Ising Models on Dense Erdös-Rényi Random Graphs, J. Stat. Phys. 177, 78 (2019).
- [13] Z. Kabluchko, M. Löwe, and K. Schubert, Fluctuations of the magnetization for Ising models on Erdös-Rényi graphs-the regimes of small *p* and the critical temperature, J. Phys. A 53, 355004 (2020).
- [14] Z. Kabluchko, M. Löwe, and K. Schubert, Fluctuations for the partition function of Ising models on Erdös-Rényi random graphs, Ann. Inst. H. Poincaré Probab. Statist. 57, 2017 (2021).
- [15] A. Bovier, S. Marello, and E. Pulvirenti, Metastability for the dilute Curie-Weiss model with Glauber dynamics, Electron. J. Probab. 26, 1 (2021),
- [16] S. Boettcher, Ground State Properties of the Diluted Sherrington-Kirkpatrick Spin Glass, Phys. Rev. Lett. 124, 177202 (2020).
- [17] W. Wang, Ground-state interface exponents of the diluted Sherrington-Kirkpatrick spin glass, Phys. Rev. B 106, 134209 (2022).
- [18] F. L. Metz and T. Peron, Mean-field theory of vector spin models on networks with arbitrary degree distributions, J. Phys. Complex. 3, 015008 (2022).
- [19] L. S. Ferreira and F. L. Metz, Nonequilibrium dynamics of the Ising model on heterogeneous networks with an arbitrary distribution of threshold noise, arXiv:2212.09424.
- [20] F. Guerra and F. L. Toninelli, The high temperature region of the Viana-Bray diluted spin glass model, J. Stat. Phys. 115, 531 (2004).
- [21] L. D. Sanctis and F. Guerra, Mean field dilute ferromagnet I. High temperature and zero temperature behavior, J. Stat. Phys. 132, 759 (2008).
- [22] G. Grimmett, Probability and Random Processes (Oxford University, New York, 2001).