

ON AREA-MINIMIZING SUBGRAPHS IN INTEGER LATTICES

ZUNWU HE AND BOBO HUA

ABSTRACT. We introduce area-minimizing subgraphs in an infinite graph via the formulation of functions of bounded variations initiated by De Giorgi. We classify area-minimizing subgraphs in the two-dimensional integer lattice up to isomorphisms, and prove general geometric properties for those in high-dimensional cases.

Mathematics Subject Classification 2010: 05C10, 28A75, 52C99.

1. INTRODUCTION

Area-minimizing submanifolds in the Euclidean space \mathbb{R}^n are important concepts in geometric measure theory. For the co-dimensional one case in \mathbb{R}^n , De Giorgi [DG61] initiated a formulation using functions of bounded variations, called an area-minimizing hypersurface or a boundary of a subset of least perimeter, see [Giu84]. A celebrated result is as follows.

Theorem 1.1 (Simons [Sim68]). *Every area minimizing hypersurface in \mathbb{R}^n for $3 \leq n \leq 7$ is flat, i.e. a boundary of a half space.*

The following equation is called the minimal surface equation

$$(1) \quad \operatorname{div} \left(\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right) = 0, \quad \mathbb{R}^k.$$

Bernstein first proved that any entire solution of the minimal surface equation on \mathbb{R}^2 is affine, see [Ber27]. Such a statement on the triviality of entire solutions is now called the Bernstein theorem. By the observations of Fleming [Fle62] and De Giorgi [DG65], the Bernstein theorem of the minimal surface equation is reduced to the classification of area-minimizing hypersurfaces or area-minimizing cones in the Euclidean space. This leads to the following Bernstein theorem: any entire solution of the minimal surface equation on \mathbb{R}^k is affine if and only if $k \leq 7$, for which the sharpness follows from the construction of a non-affine solution on \mathbb{R}^8 by Bombieri, De Giorgi, and Giusti [BDGG69].

We recall the basic setting in \mathbb{R}^n . A function f is called of locally bounded variations in \mathbb{R}^n if $f \in L^1_{loc}(\mathbb{R}^n)$ and whose distributional derivative is a Radon measure in \mathbb{R}^n . We denote by $BV_{loc}(\mathbb{R}^n)$ the space of functions of locally bounded variations in \mathbb{R}^n . A Lebesgue measurable set $F \subset \mathbb{R}^n$ is

called a Caccioppoli set if $1_F \in BV_{loc}(\mathbb{R}^n)$, where 1_F is the indicator function on F . A Caccioppoli set F is called area-minimizing in \mathbb{R}^n if for any open set $\Omega \subset \subset \mathbb{R}^n$, any Caccioppoli set W with $L^n((W \Delta F) \setminus \Omega) = 0$,

$$\int_{\Omega} |\nabla 1_F| \leq \int_{\Omega} |\nabla 1_W|,$$

where $W \Delta F := (W \setminus F) \cup (F \setminus W)$, L^n is the Lebesgue measure, and $\int |\nabla \cdot|$ is the BV seminorm. This in fact means that ∂F is an area-minimizing hypersurface. This is equivalent to that for any open set $\Omega \subset \subset \mathbb{R}^n$, and $g \in BV_{loc}(\mathbb{R}^n)$ with $g - 1_F = 0$ a.e. on $\mathbb{R}^n \setminus \Omega$,

$$\int_{\Omega} |\nabla 1_F| \leq \int_{\Omega} |\nabla g|.$$

Such a function 1_F is called of least gradient in \mathbb{R}^n , which has been extensively studied in the literature, see e.g. [BDGG69, Mir67, SWZ92, Juu05, MRSdL14, Mor17, JMN18, G18, Mor18, FM19, Zun19].

Let $\mathcal{G} = (V, E)$ be a simple undirected graph with the set of vertices V and the set of edges E . Two vertices x, y are called neighbors, denoted by $x \sim y$, if there is an edge connecting x and y , i.e. $\{x, y\} \in E$. For any subset $\Omega \subset V$, denote by $\Omega^c := V \setminus \Omega$ the complement of Ω , by $\delta\Omega := \{x \in \Omega : \exists y \in \Omega^c, x \sim y\}$ the vertex boundary of Ω and by $\tau\Omega := \{z \in \Omega^c : \exists w \in \Omega, z \sim w\}$ the exterior vertex boundary of Ω . We write $\bar{\Omega} := \Omega \cup \tau\Omega$. Given any $A, B \subset V$, we set

$$E(A, B) := \{\{x, y\} \in E : x \in A, y \in B\}.$$

We say $\partial\Omega := E(\Omega, \Omega^c)$ is the edge boundary of Ω , and set $E_{\Omega} := E(\Omega, \bar{\Omega})$. We introduce discrete analogs of a subset of least perimeter and an area-minimizing subset.

Definition 1.2. For a finite subset $U \subset V$, a subset $K \subset \bar{U}$ is called of least perimeter in U if for any $\hat{K} \subset \bar{U}$ with $\hat{K} \cap \tau U = K \cap \tau U$, we have

$$|\partial K \cap E_U| \leq |\partial \hat{K} \cap E_U|,$$

where $|\cdot|$ denotes the cardinality of a set.

Definition 1.3. A proper nonempty subset $A \subset V$ is called area-minimizing in V if for any finite $U \subset V$, $A \cap \bar{U}$ is of least perimeter in U . In this case, we identify the subset A with its induced subgraph \mathcal{G}_A on A , and call A a minimal subgraph in V for convenience.

Remark 1.4. The subset $A \subset V$ is called minimal if and only if for any finite $\Omega, \hat{F} \subset V$ with $\hat{F} \Delta F \subset \Omega$, then

$$|\partial F \cap E_{\Omega}| \leq |\partial \hat{F} \cap E_{\Omega}|.$$

For any $x \sim y$, we define

$$\nabla_{(x,y)} f := f(y) - f(x).$$

For finite $U \subset V$, the 1-Dirichlet energy on U is given by, for any $f \in \mathbb{R}^{\bar{U}}$,

$$J_U(f) := \frac{1}{2} \sum_{\{x,y\} \in E_U} |\nabla_{(x,y)} f|.$$

For a subset $\Omega \subset V$, any antisymmetric function a on E_Ω , i.e. for any x, y with $\{x, y\} \in E_\Omega$, $a_{xy} = -a_{yx}$, is called a *current* on Ω . For a function $f \in \mathbb{R}^{\bar{\Omega}}$, we call the current a is a *current associated with f on Ω* if

$$a_{xy} \in \text{Sgn}(f(x) - f(y)), \quad \forall x, y \text{ with } \{x, y\} \in E_\Omega,$$

where

$$\text{Sgn}(t) := \begin{cases} 1, & x > 0, \\ [-1, 1], & x = 0, \\ -1, & x < 0. \end{cases}$$

Let $\mathcal{C}_\Omega(f)$ be the set of currents associated with f on Ω . The 1-Laplacian is defined as a set-valued mapping $\Delta_1 : \mathbb{R}^\Omega \rightarrow 2^{\mathbb{R}^\Omega}$,

$$(2) \quad \Delta_1^\Omega f = \{g \in \mathbb{R}^\Omega : g(x) = \sum_{y \sim x} a_{xy}, a \in \mathcal{C}_\Omega(f)\}.$$

A current $a \in \mathcal{C}_\Omega(f)$ is called *minimal* in Ω if $\sum_{y \sim x} a_{xy} = 0$ for all $x \in \Omega$. The following is well-known using the results in convex analysis [Roc70, RW98], see e.g. Chang [Cha16] and Hein-Bühler [HB10].

Proposition 1.5. *For finite $U \subset V$ and a given function $\varphi \in \mathbb{R}^{\tau U}$, the subdifferential of the functional J_U at f in $\{f \in \mathbb{R}^{\bar{U}} : f|_{\tau U} = \varphi\}$ is given by $\Delta_1^U f$.*

The discrete analog of functions of least gradient was introduced in metric random walk spaces including graphs by Mazón, Pérez-Llanos, Rossi, and Toledo [MPLRT16], see also [GM21, MT23, MSDTM23].

Definition 1.6. *For finite $U \subset V$, a function $f \in \mathbb{R}^{\bar{U}}$ is called of least gradient in U if for any $g \in \mathbb{R}^{\bar{U}}$ with $g|_{\tau U} = f|_{\tau U}$, then*

$$J_U(g) \geq J_U(f).$$

The following are the characterizations of minimal subgraphs.

Theorem 1.7. *For $A \subset V$, the following are equivalent:*

- (1) *A is a minimal subgraph in V .*
- (2) *For any finite $U \subset V$, $1_{A \cap \bar{U}}$ is of least gradient in U .*
- (3) *$0 \in \Delta_1^V(1_A)$.*

Now we turn to minimal subgraphs in the integer lattices. We denote by \mathbb{Z}^n the graph of the n -dimensional integer lattice consisting of the set of vertices $\mathbb{Z}^n = \{x \in \mathbb{R}^n : x_i \in \mathbb{Z}, \forall 1 \leq i \leq n\}$ and the set of edges $\{\{x, y\} : |x - y| = 1, x, y \in \mathbb{Z}^n\}$.

The first main result is the following statement.

Theorem 1.8. Any minimal subgraph in \mathbb{Z}^2 is exactly one of the following subgraphs up to isomorphisms:

- (1) 7 families of minimal subgraphs with non-geodesic boundary (see Section 2 for the definition); see Fig.1-Fig. 7.
- (2) 3 families of minimal subgraphs with geodesic non-simple boundary (see Section 2 for the definition); see Fig.8-Fig.10.
- (3) 19 families of minimal subgraphs with geodesic simple boundary. More precisely, there are 7 families of connected minimal subgraphs: 5 families all have exactly one connected component of boundaries; see Fig.11-Fig.15; 2 families both have exactly two connected of boundaries; see Fig.16, Fig.17. There are 12 families of disconnected minimal subgraphs: they are all complementary subgraphs of connected minimal subgraphs.




FIGURE 1. $h \leq 2$.




FIGURE 2. $h = d = 1$.




FIGURE 3. $h = d = 1$.




FIGURE 4. $h = d = 1$.




FIGURE 5. $h = d = 1, a \geq 2, b \geq 2$.




FIGURE 6. $h = d = 1$.




FIGURE 7. $h = d = 1$.




FIGURE 8. $h \leq 2$.




FIGURE 9. $h = d = 1$.




FIGURE 10. $h = d = 1$.




FIGURE 11.




FIGURE 12.




FIGURE 13.




FIGURE 14.




FIGURE 15.




FIGURE 16. $0 \leq d \leq h + 2$




FIGURE 17. $d \geq 0, h \geq 0$

The classification of minimal subgraphs is quite complicated. The subtleties mainly lie in two aspects: 1. The boundaries of minimal subgraphs may not be geodesic. 2. The boundary of minimal subgraphs may not be simple. For the first aspect, we find that all boundaries of minimal subgraphs consist of geodesic rays instead of geodesic lines; for example see Claim 3.21. For the second aspect, the boundary of any minimal subgraph contains at most one unit square and distributes diagonally in some way; see Corollary 3.20 and Lemma 3.12. The arguments are based on topological, combinatorial and coarsely geometric methods.

The proof strategies of Theorem 1.8 are as follows. There are three key observations for the proof: Corollary 3.5, Lemma 3.7, and Lemma 3.9. Corollary 3.5 yields some geodesic convexity of minimal subgraphs, and implies that the boundary of any minimal subgraph determines this minimal subgraph itself in some sense. So the crucial point is to characterize the boundary of any minimal subgraph. Lemma 3.7 is a useful tool to describe the geometry of oriented boundary of any minimal subgraph, which suggests that the boundary behaves like a simple path in general. Lemma 3.9 gives a strong geometric restriction of the boundary of any minimal subgraph, which applies to deduce a strong property, Corollary 3.19, that the number of connected components of a minimal subgraph and its boundary is at most two. By the results in Lemma 3.1 and Lemma 3.19, it suffices to consider connected minimal subgraphs with at most two connected boundary components. There are two steps: First, we use several important properties to describe the geometry and topology of the boundary of any minimal subgraph (whether the boundary is geodesic or simple). Second, we use the arguments of currents to confirm the precise structure of boundary and the minimal subgraph; see Corollary 2.3 and Lemma 3.15.

On the other hand, the classification of high-dimensional minimal subgraphs (i.e. in $\mathbb{Z}^n, n \geq 3$) seems much more complicated. The reasons are as follows: The result as in Lemma 3.7 fails in higher dimension, see Fig.18; the analogs of Corollary 3.5 and Lemma 3.9 in higher dimension don't provide enough information about the local geometry.




FIGURE 18.




FIGURE 19

Nevertheless, we can decompose any three dimensional minimal subgraph \mathcal{M} into a minimal sub-subgraph $\mathcal{M}^3 \cup \mathcal{M}^2$ (union of three skeleton and two skeleton) and its one skeleton.

Theorem 1.9. *For a minimal subgraph $\mathcal{M} \subset \mathbb{Z}^3$, $\mathcal{M}^3 \cup \mathcal{M}^2$ is minimal.*

Remark 1.10. \mathcal{M}^3 is not minimal in general, see Fig. 19 for a counterexample by observing $0 \notin \Delta_1(1_{\mathcal{M}})(x)$.

Recall that $\phi : (X, d_X) \rightarrow (Y, d_Y)$ is called a rough isometry if

$$d_X(x_1, x_2) = d_Y(\phi(x_1), \phi(x_2)), d_Y(Y, \phi(X)) \leq c,$$

for any x_1, x_2 in X and some positive constant c . Note that this is stronger than usual definitions in [Woe00, BBI01]. For a minimal subgraph $\mathcal{M} \subset \mathbb{Z}^n$, we know \mathcal{M}^n does not always inherit the minimality of \mathcal{M} . But it shares the coarse geometry of \mathcal{M} and its boundary. We prove the following result.

Theorem 1.11. *For a minimal subgraph $\mathcal{M} \subset \mathbb{Z}^n$, the natural embedding \mathcal{M}^n (with the induced metric) $\subset \mathcal{M}$ is a rough isometry between metric spaces. Moreover, there exists a positive constant $c(n)$, depending only on*

n , such that for sufficiently large r and any $x \in \mathcal{M}$,

$$(3) \quad \frac{|\mathcal{M}^n \cap \hat{B}_r(x)|}{|\mathcal{M} \cap \hat{B}_r(x)|} \geq c(n),$$

$$(4) \quad \frac{|\delta\mathcal{M}^n \cap \hat{B}_r(x)|}{|\delta\mathcal{M} \cap \hat{B}_r(x)|} \geq \frac{1}{1+2n},$$

where $\hat{B}_r(x) := \{y \in \mathbb{R}^n : \max_{1 \leq i \leq n} |y_i - x_i| \leq r\}$ denotes the ∞ -normed ball centered at x of radius r .

We expect that \mathcal{M}^n determines the asymptotic geometry of \mathcal{M} and its boundary, so that the inequalities (3), (4) in Theorem 1.11 could be possibly improved, see Problem 5.1.

Moreover, we prove some restriction on the geometry of n dimensional minimal subgraphs.

Theorem 1.12. *If $\mathcal{M} \subset \mathbb{Z}^n$ is minimal, then there exist no two parallel hyperplanes bounding \mathcal{M} .*

We remark that the conclusions in Section 4 are independent of those of Section 3, and note that the maximum principle holds for minimal subgraphs in \mathbb{Z}^n in some sense; see Proposition 4.8.

The organization of this paper is as follows. In Section 2, we introduce some basic concepts and properties of minimal subgraphs. In Section 3, we focus on two dimensional minimal subgraphs and prove Theorem 1.8. In Section 4, we study the geometry of high-dimensional minimal subgraphs and prove Theorem 1.9. In Section 5, we list some open problems on geometry and topology of high-dimensional minimal subgraphs.

2. PRELIMINARIES

Let $\mathcal{G} = (V, E)$ be a simple, undirected graph. For each edge $\{x, y\} \in E$, we write (x, y) and (y, x) for associated directed edges. We say \mathcal{G} is connected if for any $x, y \in V$, there is a path $x = x_0 \sim x_1 \sim \cdots \sim x_n = y$ connecting x and y for some $n \in \mathbb{N}$. For $x \in V$, we denote by $\deg(x)$ the vertex degree of x in V . Usually, we consider the subgraph induced on a subset \mathcal{M} , for which the degree of a vertex refers to that in \mathcal{M} . The combinatorial distance $d_{\mathcal{G}}$ or simply d unless specially stated on the graph is defined as, for any $x, y \in V$ and $x \neq y$,

$$d(x, y) := \inf\{n \in \mathbb{N} : \exists \{x_i\}_{i=1}^{n-1} \subset V, x \sim x_1 \sim \cdots \sim x_{n-1} \sim y\}.$$

For any function $\varphi \in \mathbb{R}^{\tau\Omega}$, we consider the functional $J_{\Omega, \varphi} : \mathbb{R}^{\Omega} \rightarrow \mathbb{R}$ with Dirichlet boundary condition φ given by

$$J_{\Omega, \varphi}(g) = J_{\Omega}(\tilde{g}),$$

where $\tilde{g} \in \mathbb{R}^{\bar{\Omega}}$ such that $\tilde{g}|_{\Omega} = g, \tilde{g}|_{\tau\Omega} = \varphi$.

Note that $f \in \mathbb{R}^{\bar{\Omega}}$ is of least gradient if and only if $f|_{\Omega}$ is the minimizer of $J_{\Omega, f|_{\tau\Omega}}$. One readily sees that if $\Omega' \subset \Omega$, f is of least gradient in Ω , then f is of least gradient in Ω' . A similar result holds for sets of least perimeter.

The following is the discrete co-area formula, see [Bar17].

Proposition 2.1. *For any function $f \in \mathbb{R}^{\bar{\Omega}}$,*

$$J_{\Omega}(f) = \int_{-\infty}^{\infty} |\partial\{f > t\} \cap E_{\Omega}| dt,$$

where $\{f > t\} := \{x \in \bar{\Omega} : f(x) > t\}$.

Now we prove the following proposition.

Proposition 2.2. *For finite $\Omega \subset V$ and $K \subset \bar{\Omega}$, the following are equivalent:*

- (1) K is of least perimeter in Ω .
- (2) 1_K is of least gradient in Ω .
- (3) $0 \in \Delta_1^{\Omega}(1_K)$.

Proof. (1) \implies (2): Consider any $g \in \mathbb{R}^{\bar{\Omega}}$ with $g|_{\tau\Omega} = 1_K|_{\tau\Omega}$. For any $t \in (0, 1)$, $\{g > t\} \cap \tau\Omega = K \cap \tau\Omega$. Since K is of least perimeter in Ω ,

$$|\partial\{g > t\} \cap E_{\Omega}| \geq |\partial K \cap E_{\Omega}|.$$

By the co-area formula,

$$\begin{aligned} J_{\Omega}(g) &= \int_{-\infty}^{\infty} |\partial\{g > t\} \cap E_{\Omega}| dt \geq \int_0^1 |\partial\{g > t\} \cap E_{\Omega}| dt \\ &\geq |\partial K \cap E_{\Omega}| = J_{\Omega}(1_K). \end{aligned}$$

(2) \implies (1): For any \tilde{K} with $K \cap \tau\Omega = \tilde{K} \cap \tau\Omega$, the result follows from the least gradient property of 1_K by choosing $g = 1_{\tilde{K}}$.

(2) \iff (3): $1_K \in \mathbb{R}^{\bar{\Omega}}$ is of least gradient in Ω if and only if $1_K|_{\Omega}$ is the minimizer of $J_{\Omega, 1_K|_{\tau\Omega}}$. Since $J_{\Omega, 1_K|_{\tau\Omega}}$ is a convex function on $\mathbb{R}^{\bar{\Omega}}$, $1_K|_{\Omega}$ is the minimizer if and only if 0 is in the subdifferential of $J_{\Omega, 1_K|_{\tau\Omega}}$ at $1_K|_{\Omega}$, which is $0 \in \Delta_1^{\Omega}(1_K)$ by Proposition 1.5. \square

We write $B_r(x) := \{y \in V : d(y, x) \leq r\}$ for the ball of radius r centered at x .

Proof of Theorem 1.7. (1) \iff (2): This follows from Proposition 2.2.

(2) \implies (3): Consider a sequence of balls $\{B_r\}_{r=1}^{\infty}$ where $B_r := B_r(p)$ for some $p \in V$. For $U = B_r$, $r \geq 1$, it follows from Proposition 2.2,

$$0 \in \Delta_1^{B_r}(1_{A \cap \bar{B}_r}).$$

That is, for each $r \geq 1$, there is a minimal current a^r associated with $1_{A \cap \bar{B}_r}$ on B_r . Since there are countable edges in E and

$$\sup_{\{x, y\} \in E_{B_r}} |a_{xy}^r| \leq 1, \quad \forall r \geq 1,$$

there is a subsequence $r_i \rightarrow \infty$ and a current a^∞ on V such that

$$a_{xy}^{r_i} \rightarrow a_{xy}^\infty, \quad \forall \{x, y\} \in E.$$

One easily sees that $a^\infty \in \mathcal{C}_V(1_A)$ and a^∞ is minimal. Hence $0 \in \Delta_1^V(1_A)$.

(3) \implies (2): For any minimal $a \in \mathcal{C}_V(1_A)$, one easily verifies that for any finite $U \subset V$, $a|_{E_U} \in \mathcal{C}_U(1_{A \cap \bar{U}})$, which is also minimal in U .

This proves the theorem. \square

Now we introduce some notions on \mathbb{Z}^2 . For $x_1, x_2, \dots, x_k \in \mathbb{Z}^2$, we say these vertices are *horizontal (vertical, resp.)* or x_1, x_2, \dots, x_i is *horizontal (vertical, resp.)* to x_{i+1}, \dots, x_k if they are in a horizontal (vertical, resp.) line. We say x is *left-horizontal (right-horizontal, up-vertical, down-vertical, resp.)* to or a *left-horizontal (right-horizontal, up-vertical, down-vertical, resp.)* neighbor of y if x is horizontal (vertical, resp.) to y and is on the left (right, up, down, resp.) of y .

Given any two paths $\alpha \supset \alpha_1 := x_1 \sim x_2 \sim x_3$, α is called *flat* if its vertices are all horizontal or vertical. The vertex x_2 is called *a corner* of α if the subpath α_1 is not flat. We say a path is *simple* if all vertices have one or two neighbors in the path.

Given any path α containing x , x is called a *projective horizontal (vertical, resp.) interior point* if there is a horizontal (vertical, resp.) line β such that the distance projection (with respect to β) image of x lies in the interior of that of α ; see Fig. 20. Assume the path $\alpha := x_0 \sim x_1 \sim x_2 \cdots \sim x_k \sim x_{k+1} \subset \mathcal{M}$, we call α is *an isolated path* in \mathcal{M} if $\deg(x_i) = 2$ for $1 \leq i \leq k$ for the induced subgraph \mathcal{M} ; see Fig. 21.

By Theorem 1.7, we have the following.

Corollary 2.3. $\mathcal{M} \subset \mathbb{Z}^n$ is minimal if and only if $0 \in \Delta_1(1_{\mathcal{M}})$.

Remark 2.4. Corollary 2.3 provides an effective way to check the minimality of subgraphs by the equation (2). This plays an important role to classify all minimal subgraphs in \mathbb{Z}^2 .

Given a minimal graph $\mathcal{G} = (V, E) \subset \mathbb{Z}^2$, we associate it with a geometric space $X(\mathcal{G}) \subset \mathbb{R}^2$. To be precise, we identify the graph structure with the natural corresponding 1-skeleton in \mathbb{R}^2 , i.e. identify $(x, y) \in E$ with $[x, y] \subset \mathbb{R}^2$, and associate any loop $x \sim y \sim z \sim w \sim x$ with a unit square enclosed by the loop. The boundary $\delta\mathcal{G}$ is called *simple* if for any $x \in \delta\mathcal{G}$, there are at most two neighbors of x in $\delta\mathcal{G}$. $\delta\mathcal{G}$ is called *geodesic* if $d_{\delta\mathcal{G}}(y, z) = d_{\mathcal{G}}(y, z)$ holds for all $y, z \in \delta\mathcal{G}$.

For $(x, y) \in E$, the edge (x, y) is called a *boundary edge* of \mathcal{G} if there is at most one unit square in $X(\mathcal{G})$ containing $[x, y]$. A vertex $z \in v$ is called a *boundary vertex* if it is contained by a boundary edge. A boundary path $\alpha := x_0 \sim x_1 \cdots \sim x_k$ is called *oriented in \mathcal{G}* , oriented in short, if $k \geq 2$ and \mathcal{G} contains one of the following subgraphs for any subpath $\hat{\alpha} := u \sim v \sim w \subset \alpha$; see Fig. 22. If we equip α with an orientation, α is called *right (left, resp.)*




FIGURE 20. a, b are horizontal interior points, but c is not an interior point.




FIGURE 21. The horizontal path $[x, w]$ is an isolated path in \mathcal{M} .

oriented if these subgraphs for the subpaths lie on the right (left, resp.) hand side of α ; see Fig. 23.




FIGURE 22. There are three local subgraphs for black bold boundary subpath of length two.




FIGURE 23. The black bold boundary path is right oriented.

3. GEOMETRY OF TWO DIMENSIONAL MINIMAL SUBGRAPHS AND PROOF OF THEOREM 1.8

In this section, we write \mathcal{M} for a minimal subgraph in \mathbb{Z}^2 .

Lemma 3.1. *If \mathcal{M} is minimal, then so is the complementary subgraph \mathcal{M}^c .*

Proof. This is direct by definition and Corollary 2.3. □

Lemma 3.2. *$\delta\mathcal{M}$ can not contain an isolated geodesic path with length $L \geq 3$.*

Proof. If not, we may assume $\alpha := x_0 \sim x_1 \sim \cdots \sim x_k$ is an isolated geodesic path in $\delta\mathcal{M}$ with $k \geq 3$. Then one can remove this path except x_0, x_k , and obtain the new subgraph \mathcal{M}_1 . Taking $\Omega = \alpha$, one easily sees that

$$|\partial\mathcal{M} \cap E_\Omega| \geq |\partial\mathcal{M}_1 \cap E_\Omega| + 2.$$

This is impossible since \mathcal{M} is minimal. □

Lemma 3.3. *\mathcal{M} can not contain an isolated point (i.e. it has only one neighbor in \mathcal{M}). As a consequence, $\delta\mathcal{M}$ is locally one of the three subgraphs and any boundary vertex has at least two boundary neighbors in $\delta\mathcal{M}$; see Fig. 24.*




FIGURE 24. There are three local subgraphs for the bigger black boundary vertex.

Proof. If not, we get another subgraph \mathcal{M}_1 by deleting the isolated point x . It is clear that $|\partial\mathcal{M} \cap E_\Omega| = |\partial\mathcal{M}_1 \cap E_\Omega| + 2$, where $\Omega := \{x\}$. This contradicts the minimality of \mathcal{M} . □

Lemma 3.4. *If there is a finite non-closed simple path in \mathcal{M} lying on one side of a horizontal/vertical line and the line contains two endpoints of the path, then the domain enclosed by the line and the path is contained in \mathcal{M} .*

Proof. We argue by contradiction.

We may assume the path $\alpha := x_0 \sim x_1 \sim \cdots \sim x_n$ lies above on the horizontal line l such that $x_0, x_1 \in l$. Denote by \mathcal{A} the domain enclosed by the line and the path.

If \mathcal{A} is not in \mathcal{M} , then one can find $y_0 \sim y_1 \sim \cdots \sim y_m$ such that the domain enclosed by the horizontal line l' through y_0, y_m (except the line l') is contained in \mathcal{M} , and $y_0, y_m \in \mathcal{M}, y_1, \cdots, y_{m-1} \notin \mathcal{M}$. One can get another subgraph \mathcal{M}_1 by adding y_1, \cdots, y_{m-1} to \mathcal{M} , then one easily deduces that

$$|\partial\mathcal{M} \cap E_\Omega| \geq |\partial\mathcal{M}_1 \cap E_\Omega| + 1 + m - (m - 1) = |\partial\mathcal{M}_1 \cap E_\Omega| + 2.$$

This contradicts the minimality of \mathcal{M} . □

Corollary 3.5. *If $x, y \in \mathcal{M}$ lie in a horizontal/vertical line and in the same connected component of \mathcal{M} , then the horizontal/vertical segment between x and y is contained in \mathcal{M} .*

Proof. Since x, y are in the same connected component of \mathcal{M} , then there is a finite simple path $\alpha := x = x_0 \sim x_1 \cdots \sim x_n = y$ between x and y . Let l be the line on which x, y lie, and $\{x = z_0, z_1, \cdots, z_k = y\} := l \cap \alpha$. Note that each subpath between z_i and z_{i+1} of α satisfies the condition of Lemma 3.4. Thus we prove the result by Lemma 3.4. □

Corollary 3.6. *Any horizontal/vertical line cannot intersect two boundary edges and one other edge consecutively in a connected component of \mathcal{M} , i.e. there cannot exist two boundary edges $(x_1, x_2), (y_1, y_2)$ and one other edge (z_1, z_2) satisfying that x_1, y_1, z_1 and x_2, y_2, z_2 lie in two horizontal/vertical lines in the given order.*

Proof. If not, one can deduce that the rectangle $\mathcal{R}(x_1, x_2, z_1, z_2)$ is contained in \mathcal{M} by Corollary 3.5. This is impossible, since (y_1, y_2) is a boundary edge. □

Lemma 3.7. *If there is a finite simple right (left, resp.) oriented boundary path $\alpha := x_0 \sim x_1 \sim \cdots \sim x_n$ in $\delta\mathcal{M}$, and x_k is a projective horizontal (vertical, resp.) interior point in α with $1 \leq k \leq n - 1$. Then there is no vertex in \mathcal{M} on the left(right, resp.) hand side of α such that it is adjacent and vertical (horizontal, resp.) to x_k in \mathcal{M} .*

Proof. Otherwise, we may assume there is a vertex $y \sim x_k \in \mathcal{M}$ on the left side of α with $y \notin \alpha$. By Lemma 3.3, there is a vertex $y_1 (\neq x_k) \sim y \in \mathcal{M}$.

Case 1. If x_{k-1}, x_k, x_{k+1} are horizontal/vertical, then it is clear y, x_k are vertical/horizontal. Furthermore, y_1, x_k are also vertical/horizontal. If not, we can assume y_1, x_{k+1} are vertical/horizontal. Combining Corollary 3.5 and the condition that α is left oriented, one can deduce that (x_k, x_{k+1}) is contained in two unit squares in \mathcal{M} and it is not a boundary edge of \mathcal{M} . It is a contradiction. Applying Lemma 3.3 again, there is a vertex $y_2 (\neq y_1) \sim y_1 \in \mathcal{M}$. By the same argument, we have that y_2, x_k are vertical/horizontal. Continuing the process, we finally get an isolated vertical/horizontal ray

$x_k \sim y \sim y_1 \sim y_2 \sim \dots$ in \mathcal{M} . This contradicts the minimality of \mathcal{M} by Lemma 3.2.

Case 2. If x_{k-1}, x_k, x_{k+1} are not horizontal/vertical, we may assume that x_{k-1}, x_k are horizontal and x_k, x_{k+1} are vertical. Recall that x_k is projective interior vertex in α , so that we may assume x_k, x_s are vertical and x_s, x_{s+1} are horizontal for some $k+1 \leq s \leq n-1$. If x_k, y are horizontal, then x_k, x_s and y, x_{s+1} are both vertical. Combining Corollary 3.5 with the condition that α is left oriented, one can deduce that (x_s, x_{s+1}) is not a boundary edge of \mathcal{M} . It is a contradiction. Hence x_k, y are vertical. Using same arguments in Case 1, one can get a vertical ray $x_k \sim y \sim y_1 \sim y_2 \sim \dots$ in \mathcal{M} . This contradicts the minimality of \mathcal{M} by Lemma 3.2. \square

Remark 3.8. *The proof of Lemma 3.7 is valid for more general case and it is used frequently throughout this section.*

Lemma 3.9. *$\delta\mathcal{M}$ can not contain two parallel horizontal or vertical rays.*

Proof. If not, we may assume that $l_1 := x_1 \sim x_2 \sim \dots, l_2 := y_1 \sim y_2 \sim \dots \subset \delta\mathcal{M}$ are two horizontal rays originating from two vertical vertices x_1, y_1 respectively with $c := d(x_1, y_1)$. Denote by \mathcal{S} the strip bounded by l_1, l_2 . So one can find $x_i \in l_1, y_i \in l_2$ subjected to $\hat{c} := d(x_1, x_i) = d(y_1, y_i) \geq c+3$, where i is some positive integer. There are two cases.

Case 1. l_1, l_2 are in the same connected component of \mathcal{M} . Then \mathcal{M} contains \mathcal{S} by Corollary 3.5. One can obtain another graph \mathcal{M}_1 via removing the rectangle $\mathcal{R}(x_2, x_{i-1}, y_2, y_{i-1})$. Note that $x_2, y_2, x_3, y_3, \dots, x_{i-1}, y_{i-1}$ have edges not in \mathcal{M} by Lemma 3.7. Therefore, this yields that

$$|\partial\mathcal{M} \cap E_\Omega| \geq |\partial\mathcal{M}_1 \cap E_\Omega| + 2(\hat{c} - 2) - 2c \geq |\partial\mathcal{M}_1 \cap E_\Omega| + 2,$$

where $\Omega = \mathcal{R}(x_1, x_i, y_1, y_i)$. This is a contradiction by the minimality of \mathcal{M} .

Case 2. l_1, l_2 are in different connected components of \mathcal{M} . We may assume that l_1 is above l_2 . For each x_k, y_k with integer $k \geq 1$, there exist $[\hat{x}_k, \hat{y}_k] \subset [x_k, y_k]$ with

$$(5) \quad [\hat{x}_k, \hat{y}_k] \cap \mathcal{M} = \{\hat{x}_k, \hat{y}_k\}, d(\hat{x}_k, \hat{y}_k) \geq 2.$$

Let $\alpha_1 := \{\hat{x}_1, \hat{x}_2, \dots\}$ ($\alpha_2 := \{\hat{y}_1, \hat{y}_2, \dots\}$, *resp.*). Then we get another graph \mathcal{M}_1 by filling a rectangle $\mathcal{R}(x_1, x_i, y_1, y_i)$. Note that these \hat{x}_k, \hat{y}_k are boundary vertices of \mathcal{M} . Therefore, by using (5) one can show that

$$|\partial\mathcal{M} \cap E_\Omega| \geq |\partial\mathcal{M}_1 \cap E_\Omega| + 2\hat{c} - 2c \geq |\partial\mathcal{M}_1 \cap E_\Omega| + 4,$$

where $\Omega = \mathcal{R}(x_1, x_j, y_1, y_j)$. This contradicts the minimality of \mathcal{M} . \square

Corollary 3.10. *Any two disjoint infinite simple boundary paths in $\delta\mathcal{M}$ can not be contained in some unbounded infinite strip isometric to $[0, d] \times [0, \infty)$, where d is some positive constant.*

Proof. The proof is similar to that of Lemma 3.9. Assume that l_j is the line through x_j, y_j . We may assume $\beta_1 := z_1 \sim z_2 \sim \dots, \beta_2 := w_1 \sim w_2 \sim \dots \subset \delta\mathcal{M}$ are two disjoint infinite boundary simple paths bounded by two

horizontal rays $\alpha_1 := x_1 \sim x_2 \sim \dots$, $\alpha_2 := y_1 \sim y_2 \sim \dots$ and x_1, z_1, w_1, y_1 are vertical. Let γ_j be the vertical line through x_j, y_j , $c := d(x_1, y_1)$, $\hat{c} := d(x_1, x_i) \geq c + 3$ for some integer i .

We only prove the case that β_1, β_2 are in the same connected component of \mathcal{M} , and the others are similar. It follows that the strip $\hat{\mathcal{S}}$ bounded by β_1, β_2 is contained in \mathcal{M} by Corollary 3.5. Note that there is at least one edge not in \mathcal{M} for $\gamma_j \cap \beta_i (i = 1, 2)$ with $j \geq 2$ by Corollary 3.7. Therefore, we have

$$|\partial\mathcal{M} \cap E_\Omega| \geq |\partial\mathcal{M}_1 \cap E_\Omega| + 2(\hat{c} - 2) - 2c \geq |\partial\mathcal{M}_1 \cap E_\Omega| + 2,$$

where $\Omega = \hat{\mathcal{S}} \cap \mathcal{R}(x_2, x_{i-1}, y_{i-1}, y_2)$ and \mathcal{M}_1 is obtained by removing Ω from \mathcal{M} . This contradicts the minimality of \mathcal{M} . \square

Lemma 3.11. \mathcal{M} doesn't contain one of three subgraphs with $[u, y] \cup [y, z] \cup [z, w] \subset \delta\mathcal{M}$; see Fig. 25.




FIGURE 25. $x \sim y \sim u, z \sim w \sim v$.

Proof. By Corollary 3.5, we have that for the first subgraph in Fig. 25 the rectangle $R(x, y, z, w) \subset \mathcal{M}$ and then $[y, z]$ is not a boundary path. It is a contradiction. For the second one in Fig. 25, applying Lemma 3.7 to the projective vertical interior point y in the oriented path $[u, y] \cup [y, z] \cup [z, w]$, this yields that $[y, z]$ or $[u, y]$ is not a boundary path. It is a contradiction.

For the third one in Fig. 25, we have

Claim 3.12. The connected component \mathcal{C} containing x of \mathcal{M} and the interior of the domain II (or III, IV) are disjoint; see Fig. 26.




FIGURE 26.

Proof of Claim 3.12. By Corollary 3.5 and Corollary 3.7, we deduce that \mathcal{C} and the interior of the domain IV are disjoint. By symmetry of the domain II and III , one can assume that there is a vertex $s \in \mathcal{C}$ lying in the interior of the domain II . By definition of \mathcal{C} , it is easy to obtain that there is a vertex p (distinct to x or u respectively) vertical or horizontal to x or u respectively. Then using Corollary 3.5 again, we get that $[p, x] \subset \mathcal{M}$ or $[p, u] \subset \mathcal{M}$ and therefore \mathcal{M} contains a unit square containing x, y, u . This is impossible since $[y, u]$ is a boundary edge. \square

Assume that \mathcal{C}_1 is the connected component of \mathcal{M} in the domain I . Now we consider the boundary $\delta\mathcal{C}_1$.

Case 1. $\delta\mathcal{C}_1$ is finite. Using Lemma 3.3, we can find a closed simple path $\alpha \subset \delta\mathcal{C}_1$ to enclose \mathcal{C}_1 . By Corollary 3.5, we have $\alpha = \delta\mathcal{C}_1$ and that \mathcal{C}_1 is the domain enclosed by α . Taking $\Omega = \mathcal{C}$, by comparing the edge boundary of \mathcal{C}_1 with that of $[y, z]$, we get that

$$|\partial\mathcal{M} \cap E_\Omega| > |\partial\mathcal{M}_1 \cap E_\Omega| + 2,$$

where \mathcal{M}_1 is obtained by removing \mathcal{C} except $[y, z]$ from \mathcal{M} . This is impossible by the minimality of \mathcal{M} .

Case 2. $\delta\mathcal{C}_1$ is infinite. Applying Corollary 3.10, we have that $\delta\mathcal{C}$ contains at most one infinite half simple path. If $\delta\mathcal{C}$ contains one infinite half simple path β , then by Corollary 3.6 one deduces that β only have finite backtracks in the direction of the line containing x, y . Hence one can get that the subpath β_1 which is sufficiently far away from x of β is a geodesic ray. This contradicts the minimality of \mathcal{M} by Lemma 3.2. \square

Lemma 3.13. *If there is a simple loop in $\delta\mathcal{M}$, then the loop is a unit square.*

Proof. Note that the domain enclosed by any closed simple loop in $\delta\mathcal{M}$ is contained in \mathcal{M} by Corollary 3.5. Let α be a simple loop in $\delta\mathcal{M}$ and $\mathcal{D} \subset \mathcal{M}$ be the domain enclosed by α . We may assume that \mathcal{A} is the smallest rectangle containing \mathcal{D} . Using Corollary 3.5, we have $\mathcal{A} \cap \alpha = [a, d'] \cup [a', b] \cup [b', c] \cup [c', d]$, where $[a, d']$, $[a', b]$, $[b', c]$, $[c', d]$ are the highest, the leftmost, the lowest, the rightmost subpath of α .

Since α is a simple loop and $\mathcal{A} \cap \alpha = [a, d'] \cup [a', b] \cup [b', c] \cup [c', d]$ for the smallest rectangle \mathcal{A} containing \mathcal{D} , we can assume that $a, a', b, b', c, c', d, d'$ are arranged counterclockwise on the loop α . Let $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ be the boundary subpaths connecting a and a' , b and b' , c and c' , d and d' in α , respectively.

We claim that $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ are geodesic. It suffices to prove the result for $\alpha_1 := a = a_1 \sim a_2 \sim \cdots \sim a_i = a'$. Suppose it is not true, we may assume that a_j, a_{j+1} and a_{j+2}, a_{j+3} are horizontal, a_j, a_{j+3} and a_{j+1}, a_{j+2} are vertical. If a_{j+1} is on the left of a_j , then by the fact that $[a', b]$ is the leftmost, there is a boundary edge (a_k, a_{k+1}) for some integer $j+3 < k < i-1$ such that a_j, a_{j+3}, a_k and $a_{j+1}, a_{j+2}, a_{k+1}$ are vertical. This is impossible by Corollary 3.6. If a_{j+1} is on the right of a_j , then the square $R(a_j, a_{j+1}, a_{j+2}, a_{j+3})$ is

contained in \mathcal{M} by Corollary 3.5. This implies (a_{j+1}, a_{j+2}) is not a boundary edge and it is a contradiction. Thus we prove the claim.

It follows from the above claim that $a = a', b = b', c = c', d = d'$ if and only if $a, a'; b, b'; c, c'; d, d'$ are pairwise horizontal or vertical. Note that \mathcal{D} have other neighbors in \mathcal{M} exactly adjacent to the subset $\mathcal{B} \subset \{a, a', b, b', c, c', d, d'\}$ by Lemma 3.7.

Case 1. At least three of $a \neq a', b \neq b', c \neq c', d \neq d'$ hold. It is clear that $|\mathcal{B}| \leq 1$ by Lemma 3.7. This obviously yields that

$$|\partial\mathcal{M} \cap E_\Omega| - |\partial\mathcal{M}_1 \cap E_\Omega| \geq 6 \cdot 2 - 2 > 0,$$

where $\Omega := \mathcal{D}$ and the new subgraph \mathcal{M}_1 is obtained by removing \mathcal{D} from \mathcal{M} except \mathcal{B} . This contradicts the minimality of \mathcal{M} .

Case 2. Two or one of $a \neq a', b \neq b', c \neq c', d \neq d'$ hold. We only need to prove the result for the subcase that one of $a \neq a', b \neq b', c \neq c', d \neq d'$ holds, since the other case is similar to those of Case 1 and the previous subcase. Now we may assume $a \neq a'$ and $b = b', c = c', d = d'$. Applying Lemma 3.7 and Lemma 3.11, we have $|\mathcal{B}| \leq 2$ and $\mathcal{B} = \{b, d\}$ if $|\mathcal{B}| = 2$. Therefore we get

$$|\partial\mathcal{M} \cap E_\Omega| - |\partial\mathcal{M}_1 \cap E_\Omega| \geq 3 \cdot 2 - 2 \cdot 2 > 0,$$

where $\Omega := \mathcal{D}$ and the new subgraph \mathcal{M}_1 is obtained by removing \mathcal{D} from \mathcal{M} except \mathcal{B} . This contradicts the minimality of \mathcal{M} .

Case 3. None of $a \neq a', b \neq b', c \neq c', d \neq d'$ holds, i.e. $a = a', b = b', c = c', d = d'$. Then $\mathcal{D} = \mathcal{R}(a, b, c, d)$ is a rectangle with $d(a, b) = m \geq 1, d(a, d) = n \geq 1$. By Lemma 3.7, Lemma 3.11, $|\mathcal{B}| \leq 2$ and $\mathcal{B} = \{a, c\}$ or $\mathcal{B} = \{b, d\}$. Observing that \mathcal{M} is minimal, we deduce that

$$|\partial\mathcal{M}_1 \cap E_\Omega| \geq |\partial\mathcal{M} \cap E_\Omega| \geq |\partial\mathcal{M}_1 \cap E_\Omega| + 2(m + n) - 4,$$

where $\Omega := \mathcal{D}$ and the new subgraph \mathcal{M}_1 is obtained by removing \mathcal{D} from \mathcal{M} except \mathcal{B} . This implies $m = n = 1$ and the result follows. \square

Corollary 3.14. *Given any connected component \mathcal{C} of \mathcal{M} , if $\delta\mathcal{C}$ contains some loops, then these loops are exactly one unit square.*

Proof. Otherwise, we may assume that there are two unit distinct squares $\mathcal{R}(x, y, z, w), \mathcal{R}(\hat{x}, \hat{y}, \hat{z}, \hat{w})$ in $\delta\mathcal{C}$ by Lemma 3.13. Suppose that x, y, z, w and $\hat{x}, \hat{y}, \hat{z}, \hat{w}$ are arranged anticlockwise on $\mathcal{R}(x, y, z, w), \mathcal{R}(\hat{x}, \hat{y}, \hat{z}, \hat{w})$ respectively, and $(x, w)((\hat{x}, \hat{w}), (x, y), (\hat{x}, \hat{y}), \text{resp.})$ are the leftmost (rightmost, lowest, highest, resp.) in $\mathcal{R}(x, y, z, w), \mathcal{R}(\hat{x}, \hat{y}, \hat{z}, \hat{w})$ respectively.

Recall that $\mathcal{R}(x, y, z, w), \mathcal{R}(\hat{x}, \hat{y}, \hat{z}, \hat{w})$ are in the connected component \mathcal{C} . Applying Corollary 3.6, we may assume that $d(z, \hat{z}) = d(\mathcal{R}(x, y, z, w), \mathcal{R}(\hat{x}, \hat{y}, \hat{z}, \hat{w}))$ and there is a simple boundary path α realizing $d(z, \hat{z})$ in the domain I ; see Fig. 27.

Observe that w, \hat{y} both have exactly two neighbors in \mathcal{M} , by the proof of Lemma 3.7 for the simple boundary path $x \sim w \sim z \cup \alpha \cup \hat{z} \sim \hat{y} \sim \hat{x}$ with




FIGURE 29. α has finitely many corners and the current-determined region \mathcal{D} is a quadrant region.




FIGURE 30. α has infinitely many corners in one direction and the current-determined region \mathcal{D} is a half plane.




FIGURE 31. α has bi-infinitely many corners and the current-determined region \mathcal{D} is the whole plane or \mathbb{Z}^2 .

Proof. The first statement can be deduced by the equation (2) and Corollary 2.3. The currents in \mathcal{D} yields that $0 \in \Delta_1(1_{\mathcal{D}})$. Using Corollary 2.3 again, we have that $\mathcal{M}' := \mathcal{M} \cup \mathcal{D}$ is minimal. Applying Lemma 3.7, one can obtain that α_1 is an oriented boundary geodesic line (ray, line, resp.) in Fig. 28 (29, 30, resp.), and α_2 is an oriented boundary geodesic ray in Fig. 29. Thus, we get that $\alpha' := \alpha_1 (\alpha_1 \cup \alpha_2, \alpha_1, \text{resp.})$ is an oriented boundary geodesic line in Fig. 28 (Fig. 29, Fig. 30, resp.). \square

Corollary 3.16. *If the minimal subgraph \mathcal{M} contains two oriented boundary geodesic lines α, β in different connected components of $\delta\mathcal{M}$, then α, β are both of the form in Fig. 29 in Lemma 3.15.*

Proof. We argue by contradiction. Note that $\alpha \cap \beta = \emptyset$.

Case 1. α is of the form in Fig. 28 or Fig. 30. By Lemma 3.15 we may assume α' is a horizontal line.

If β is of the form in Fig. 28 or Fig. 30, and we assume that β' is a vertical line. Then it is easy to deduce that $\alpha \cap \beta \neq \emptyset$, which is impossible. If β' is also a horizontal line, then we can get a new minimal subgraph containing two boundary horizontal lines α', β' by Lemma 3.15. This is impossible by Lemma 3.9.

If β is of the form in Fig. 29, then we can get a new minimal subgraph containing two boundary horizontal rays by Lemma 3.15. This is impossible by Lemma 3.9.

If β is of the form in Fig. 31, then it always holds that $\alpha \cap \beta \neq \emptyset$. This is impossible.

Case 2. α, β are both of the form in Fig. 31. Since $\alpha \cap \beta = \emptyset$, α, β divide the plane into three domains $\mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3$, where \mathcal{D}_2 is bounded by α and β ; see Fig. 32. Thus the currents determined by α are inconsistent with these




FIGURE 32.

determined by β by Lemma 3.15, i.e. $0 \notin \Delta_1(1_{\mathcal{M}})$. □

Lemma 3.17. *Assume $\delta\mathcal{M}$ contains no loops. Let α be a boundary geodesic line in $\delta\mathcal{M}$ with exactly one corner $x \in \alpha$. Then the connected component \mathcal{D} of \mathcal{M} containing x in a quadrant region \mathcal{E} enclosed by α is α or \mathcal{E} .*

Proof. If not, then one can find a vertex $p \in \mathcal{E} \setminus \mathcal{D}$. We may assume that $\alpha = \alpha_1 \cup \alpha_2$, where α_1, α_2 are flat geodesic rays from x ; see Fig. 33. By Corollary 3.5, two flat geodesic rays α'_1, α'_2 from p , which don't intersect with \mathcal{D} ; see Fig. 33. Then $\mathcal{D} \cap \mathcal{E}$ is bounded by $\alpha_1 \cup \alpha_2$ and $\alpha'_1 \cup \alpha'_2$.

Since $\mathcal{E} \setminus \mathcal{D} \neq \emptyset$, there is another bi-infinite simple boundary path $\beta \subset \delta\mathcal{M} \cap \mathcal{D}$ with $\beta \neq \alpha_1 \cup \alpha_2$. Since $\delta\mathcal{M}$ has no loops, one can find an infinite simple subpath $\beta_1 \subset \beta$ such that $\beta_1 \cap (\alpha_1 \cup \alpha_2) = \emptyset$. Thus we can find two disjoint infinite simple boundary paths of \mathcal{D} in the unbounded strip enclosed by α'_1 and α_1 or enclosed by α'_2 and α_2 . This is impossible by Corollary 3.10.




FIGURE 33.

□

Corollary 3.18. *Assume $\delta\mathcal{M}$ contains no loops. Let α be a boundary geodesic line in $\delta\mathcal{M}$ with finitely many corners. Then the connected component of \mathcal{M} containing α in a quadrant region \mathcal{D} enclosed by α is α or \mathcal{D} .*

Proof. The argument of Lemma 3.17 still works with minor modifications.

□

Lemma 3.19. *Assume that $\delta\mathcal{M}$ is geodesic and contains no loops. Then both $\delta\mathcal{M}$ and \mathcal{M} have at most two connected components.*

Proof. If $\delta\mathcal{M}$ has more than two connected components, then there are three oriented disjoint boundary geodesic lines by Lemma 3.3 and Lemma 4.3. Using Corollary 3.16, one easily sees that there are two disjoint parallel boundary geodesic rays in a bounded strip. It is a contradiction by Lemma 3.9.

□

Now we can prove Theorem 1.8.

Proof of Theorem 1.8. We divide it into three cases.

Case 1. $\delta\mathcal{M}$ is non-geodesic.

We may assume that $\delta\mathcal{M}$ contains a path $x \sim z \cup [z, w] \cup w \sim y$ such that $[x, y]$ is not a boundary path with $d(x, y) = d(z, w)$, and z, w are horizontal; see Fig. 34. Then the rectangle determined by x, y, w, z is contained in \mathcal{M} by Corollary 3.5. If $\deg(z) \geq 3$ and $\deg(w) \geq 3$, then this contradicts




FIGURE 34.

Lemma 3.11 by symmetry of z and w .

If $\deg(z) = 2$ and $\deg(w) = 2$, then we get the new subgraph \mathcal{M}_1 by removing the rectangle $\mathcal{R}(x, y, z, w)$ except $[x, y]$. By Lemma 3.7, we have

$$|\partial\mathcal{M} \cap E_\Omega| = |\partial\mathcal{M}_1 \cap E_\Omega| + 2 > |\partial\mathcal{M}_1 \cap E_\Omega|,$$

where $\Omega := \mathcal{R}(x, y, z, w)$. This contradicts the minimality of \mathcal{M} .

Thus, either $\deg(z) \geq 3, \deg(w) = 2$ or $\deg(w) \geq 3, \deg(z) = 2$ occurs. So that we can assume $\deg(w) \geq 3$ and $\deg(z) = 2$. Denote by \mathcal{C} the connected component of \mathcal{M} containing w and $\mathcal{C}_1(\mathcal{C}_2, \text{resp.})$, the part of \mathcal{C} in the domain $I(II, \text{resp.})$; see Fig. 35.




FIGURE 35.

Step 1 of Case 1. $\mathcal{C}_1(\mathcal{C}_2, \text{resp.})$ is enclosed by two geodesic rays from w in the domain $I(II, \text{resp.})$.

Claim 3.20. $\delta\mathcal{C}_1$ contains no loops.

Proof of Claim 3.20. Otherwise, $\delta\mathcal{C}_1$ contains exactly one unit square $\mathcal{R}(\hat{x}, \hat{y}, \hat{z}, \hat{w})$ by Corollary 3.14. We can assume that $\hat{x}, \hat{y}, \hat{w}, \hat{z}$ are arranged anticlockwise, \hat{x}, \hat{y} are horizontal, and \hat{x} is on the left of \hat{y} . By Lemma 3.6, there is a boundary simple path α connecting w, \hat{x} in the domain I . Applying Lemma 3.7 to the boundary simple paths $\alpha \cup \hat{x} \sim \hat{y} \sim \hat{w}$ and $\alpha \cup \hat{x} \sim \hat{z} \sim \hat{w}$, we have $\deg(\hat{y}) = \deg(\hat{z}) = 2$.

Consider the new subgraph \mathcal{M}_1 by removing the part \mathcal{S} in the rectangle determined by $x, y, z, \hat{w}, \hat{y}, \hat{z}$ from \mathcal{M} except $[x, y], [\hat{x}, \hat{y}]$. Since $\deg(z) = \deg(\hat{y}) = \deg(\hat{z}) = 2$, we have

$$|\partial\mathcal{M} \cap E_\Omega| \geq |\partial\mathcal{M}_1 \cap E_\Omega| + 2 \cdot 2 - 2 > |\partial\mathcal{M}_1 \cap E_\Omega|,$$

where $\Omega := \mathcal{S}$. This contradicts the minimality of \mathcal{M} . \square

Using Claim 3.20 and Lemma 3.2, we may assume that $\hat{\beta}_1 := \beta \cup \beta_1 = (\hat{\beta}_2 := \beta \cup \beta_2, \text{resp.})$ is the leftmost(rightmost, resp.) simple infinite path in \mathcal{C}_1 such that $\beta = \hat{\beta}_1 \cap \hat{\beta}_2$ is a path connecting w and \hat{w} . Let $l_1(l_2, \text{resp.})$ be a vertical(horizontal, resp.) line through w ; see Fig. 36. Applying Corollary 3.10, the distance projection of $\beta_1(\beta_2, \text{resp.})$ on $l_1(l_2, \text{resp.})$ is an infinite ray.

Claim 3.21. β_1, β_2 are geodesic, and the length of β is at most one.




FIGURE 36.

Proof of Claim 3.21. If β_1 is not geodesic, then there are four cases for some non-geodesic boundary subpath $\hat{w} \sim \dots \sim x_1 \sim x_2 \sim \dots \sim x_3 \sim x_4 \subset \beta_1$. If $x_1(x_4, \text{ resp.})$ is right-horizontal to $x_2(x_3, \text{ resp.})$, x_4 is up-vertical to x_1 , then there is a boundary edge $(y_1, y_2) \subset \beta \cup \hat{w} \sim \dots \sim x_1$ such that x_1, x_4, y_1 and x_2, x_3, y_2 are vertical. This contradicts Corollary 3.6.

If $x_1(x_4, \text{ resp.})$ is down-vertical to $x_2(x_3, \text{ resp.})$, x_1 is left-horizontal to x_4 , then by that fact that the distance projection of β_1 on l_1 is an infinite ray, there is an edge $(y_1, y_2) \subset \beta_1$ such that x_1, x_4, y_1 and x_2, x_3, y_2 are horizontal. This contradicts Corollary 3.6.

If $x_1(x_4, \text{ resp.})$ is left-horizontal to $x_2(x_3, \text{ resp.})$, x_4 is up-vertical to x_1 , then we have $\mathcal{R}(x_1, x_2, x_3, x_4) \subset \mathcal{M}$ by Corollary 3.5. This contradicts that β_1 is leftmost.

If $x_1(x_4, \text{ resp.})$ is up-vertical to $x_2(x_3, \text{ resp.})$, x_1 is left-horizontal to x_4 , then we have $\mathcal{R}(x_1, x_2, x_3, x_4) \subset \mathcal{M}$ by Corollary 3.5. This contradicts that β_1 is leftmost.

Hence we have that β_1 is geodesic. The same argument yields that β_2 is geodesic.

Denote by $L(\beta)$ the length of β . Note that β is geodesic by Corollary 3.5 and Claim 3.20. Now consider a new subgraph \mathcal{M}_1 obtained by removing the part of \mathcal{C} in the rectangle \mathcal{R}' determined by x, y, z, \hat{w} , from \mathcal{M} except $[x, y]$ and \hat{w} . Combining Lemma 3.7 with the minimality of \mathcal{M} , we have

$$|\partial \mathcal{M}_1 \cap E_\Omega| \geq |\partial \mathcal{M} \cap E_\Omega| \geq |\partial \mathcal{M}_1 \cap E_\Omega| + 2(L(\beta) - 1),$$

where $\Omega := \mathcal{C} \cap \mathcal{R}'$. This implies $L(\beta) \leq 1$. \square

Claim 3.22. *The domain \mathcal{D}_1 enclosed by β_1, β_2 in domain I is contained in \mathcal{C}_1 . As a consequence, we have $\mathcal{C}_1 = \mathcal{D}_1 \cup \beta$.*

Proof of Claim 3.22. By same arguments in Lemma 3.17, one can show that $\mathcal{C}_1 - \beta = \mathcal{D}_1$ or $\mathcal{C}_1 - \beta = \beta_1 \cup \beta_2$. By definition, $\mathcal{C}_1 - \beta = \beta_1 \cup \beta_2$ implies that $\beta_1 \cup \beta_2$ is an isolated path. It is impossible by Lemma 3.2. Thus we prove the result.

□

Similar arguments yield that \mathcal{C}_2 is the domain enclosed by two geodesic rays β'_1 and β'_2 from w , where the subpath $w \sim \dots \sim z \sim x \subset \beta'_1$, the subpath $w \sim y \subset \beta'_2$, and $\beta'_1 \cap \beta'_2 = \{w\}$.

Step 2 of Case 1. The classification of \mathcal{M} of Case 1 is indeed as listed in Theorem 1.8 by Lemma 3.15.

Claim 3.23. *At most one of β_1 and β_2 has corners. If β_1 (β_2 , resp.) has corners, then $\beta_1 = \hat{\beta}_1$ ($\beta_2 = \hat{\beta}_2$, resp.) contains exactly two corners and the two corners are adjacent.*

Proof of Claim 3.23. We argue via Corollary 2.3 and the equation (2). We give only some arguments and the remaining are similar. If both β_1 and β_2 have corners, then the current flowing out of $w \geq 3 - 1 = 2 > 0$; see the first picture in Fig. 37 for $L(\beta) = 0$. This is a contradiction by Corollary 2.3. Note that if β_1 has corners, then it has at least two corners by Lemma 3.9. On the other hand, β_1 has at most two corners by Corollary 2.3; see Fig. 38. The remaining claims are easy to verify by Corollary 2.3; see the last two pictures in Fig. 37 and Fig. 39.




FIGURE 37. The second picture shows $\Delta_1(1_{\mathcal{M}})(w) \neq 0$ or $\Delta_1(1_{\mathcal{M}})(w_3) \neq 0$ if two corners of β_1 are not adjacent.




FIGURE 38. The first picture shows $\Delta_1(1_{\mathcal{M}})(w) \neq 0$ or $\Delta_1(1_{\mathcal{M}})(w_2) \neq 0$ if β_1 has more than two corners.

□




FIGURE 39.

Claim 3.24. For β'_1 and β'_2 , denote by $\mathcal{R}_1, \mathcal{R}_2$ the rectangles containing w and p_1 , w and p_1 respectively, and let l_1 and h_1 , l_2 and h_2 represent sides of $\mathcal{R}_1, \mathcal{R}_2$ respectively. Let d_1, d_2 be the length of h_1, h_2 respectively; see the graph (a) in Fig. 40. Then we have that $d_1 \leq 2, d_2 \leq 1$. In particular, β'_1 (β'_2 , resp.) has at most four (two, resp.) corners. If β'_2 has two corners, then two corners in β'_1 and β'_2 are both adjacent.




FIGURE 40. In (a): The oriented paths γ_1, γ_2 from w to p_1, p_2 are finite subpaths in β'_1, β'_2 respectively. C_1, C_2 denote the currents determined by the oriented paths γ_1, γ_2 in the interiors of the rectangles containing w and p_1 , w and p_2 respectively. In (b): $w \sim y \sim y_1$. In (c): $w \sim w_1$ and $w_1 \in [z, w]$.

Proof of Claim 3.24. If $d_1 \geq 3$, then this yields (b) in Fig. 40 by the currents indicated by C_1 in Fig. 40 and Corollary 2.3. This implies that $\Delta_1(1_{\mathcal{M}})(y) \neq 0$ or $\Delta_1(1_{\mathcal{M}})(y_1) \neq 0$, which is impossible by Corollary 2.3. Similarly, if $d_2 \geq 2$, then we have (c) in Fig. 40 by the currents indicated by C_2 in Fig. 40. This implies that $\Delta_1(1_{\mathcal{M}})(w_1) \neq 0$, which is impossible by Corollary 2.3. \square

Claim 3.25. \mathcal{M} is connected.

Proof of Claim 3.25. If not, let $\hat{\mathcal{C}}$ be another connected component of \mathcal{M} and $\hat{\mathcal{C}} \cap \mathcal{C} = \emptyset$. Then $\delta\hat{\mathcal{C}}$ is geodesic. Otherwise, $\hat{\mathcal{C}}$ contains two quadrants located diagonally such that $\hat{\mathcal{C}} \cap \mathcal{C} \neq \emptyset$ by Claim 3.23 and Claim 3.24, which is impossible. So that one can find a geodesic line γ in $\delta\hat{\mathcal{C}}$ by Lemma 3.3 and

where $\Omega := R(x, y, z, w)$. This implies that $d \leq h + 2$.

On the other hand, we shall prove that \mathcal{M}_1 is minimal if $d \leq h + 2$. We argue it by induction on h . If $h = 0$, then we get the result by Corollary 2.3; see Fig. 42. To be precise, for $d = 0$, the currents of α_1, α_2 can be




FIGURE 42. $\alpha_1(\alpha_2, \text{resp.})$ are horizontal(vertical, resp.) lines through x and they divide the plane into four quadrant regions. The crossed arrows denote the currents in the interiors of respective quadrant regions.

arbitrarily given (for example, zero currents) whenever they are of sum zero along respective lines; see (a) in Fig. 42. For $d = 1$, the currents of α_1 can be arbitrarily given whenever they are of sum zero along respective lines. The currents of α_2 are indicated in Fig. 42; see (b) in Fig. 42. For $d = 2$, one can see (c) in Fig. 42 to find that the minimal currents of α_1, α_2 are indicated.

Now let the current from t to x_1 be as in Fig. 41. Then we obtain the currents in Fig. 41 by Corollary 2.3 and the currents in the interior of the upper half plane bounded by the horizontal line through x and t , which are indicated by crossed arrows in Fig. 41. Denote by α'_1 the vertical ray from t in α_1 , and α'_2 the horizontal ray from t through w . Then one gets a new connected graph \mathcal{M}_2 bounded by $\alpha_1 \cup \alpha_2$ and β ; see Fig. 41. Note that $d(x_1, y_1) = h - 1, d(x_1, w_1) = d - 1$, and one can reverse the currents of the geodesic line $\alpha'_1 \cup \alpha'_2$. One can show that this preserves that $0 \in \Delta_1(1_{\mathcal{M}_2})$. Thus, by applying induction assumption we finish the proof. \square

Subcase 2-1-2-2 If \mathcal{M} is of the form in Fig. 17, then \mathcal{M} is minimal if and only if the new subgraph \mathcal{M}_1 in Fig. 43 is minimal by Lemma 3.15. Now we show that \mathcal{M}_1 in Fig. 43 is indeed minimal by Corollary 2.3. More precisely, α, β divide the plane into three regions $\mathcal{D}_1, \mathcal{D}_2$ and \mathcal{M}_1 ; see Fig. 43. The currents in the interiors of $\mathcal{D}_1, \mathcal{D}_2$ are indicated as the crossed arrows in Fig. 43, and the currents for the horizontal lines or vertical lines through the




FIGURE 43. \mathcal{M}_1 is a connected subgraph bounded by two geodesic lines α and β .

rectangle $R(x, y, z, w)$ in Fig. 43 only need to be of sum zero along these lines. Hence, this yields that $0 \in \Delta_1(1_{\mathcal{M}_1})$.

Subcase 2-2 The number of connected components of \mathcal{M} is two.

One can show that $\delta\mathcal{M}$ has exactly two connected components by Lemma 3.19. Recalling that $\delta\mathcal{M}$ is geodesic and simple, one obtains that the complementary subgraph \mathcal{M}^c of \mathcal{M} is connected. Lemma 3.1 yields that \mathcal{M}^c is also minimal.

Case 3. $\delta\mathcal{M}$ is geodesic and non-simple.

Subcase 3-1 $\delta\mathcal{M}$ contains some loops. Then one of connected components of \mathcal{M} contains exactly one unit square by Corollary 3.14.

We may assume that there is a square $R(x, y, z, w) \subset \delta\mathcal{M} \cap \mathcal{C}$, where \mathcal{C} is the connected component of \mathcal{M} containing x . Using Lemma 3.11 and symmetry of x, z and y, w , one can assume that \mathcal{C} is in the following shadow region; see Fig. 44.




FIGURE 44.

One easily gets that the minimal currents for the square $R(x, y, z, w)$; see Fig. 44. Then by $\Delta_1(1_{\mathcal{M}})(x) = \Delta_1(1_{\mathcal{M}})(z) = 0$, we deduce that there are two neighbors z_1, z_2 (distinct from w, y) of z and two neighbors x_1, x_2 (distinct from w, y) of x . Applying Lemma 3.3 and Corollary 3.14 and recalling that $\delta\mathcal{M}$ is geodesic, we have two flat boundary geodesic rays α_1, α_2

from z through z_1, z_2 respectively and two flat boundary geodesic rays α_3, α_4 from x through x_1, x_2 respectively. Using Lemma 3.17, it implies that \mathcal{C} is isometric to the subgraph in Fig. 9. Similar arguments as in the proof of Claim 3.25 yield that $\mathcal{M} = \mathcal{C}$.

Subcase 3-2 $\delta\mathcal{M}$ contains no loops.

Denote by $\deg(p, \delta\mathcal{M})$ the number of neighbors of p in $\delta\mathcal{M}$ for any $p \in \mathcal{M}$, and by \mathcal{C} the connected component of \mathcal{M} containing x for some $x \in \delta\mathcal{M}$. Recalling that $\delta\mathcal{M}$ is non-simple, we may assume that $\deg(x, \delta\mathcal{M}) \geq 3$.

Subcase 3-2-1 Assume $\deg(x, \delta\mathcal{M}) = 4$ and y, z, v, w are neighbors of x .

By Lemma 3.3, Lemma 4.3 and the fact that $\delta\mathcal{M}$ is geodesic, there are four boundary geodesic rays $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ from x through y, z, v, w respectively. By Corollary 3.5 and recalling that $\delta\mathcal{M}$ is geodesic and contains no loops, we obtain that these geodesic rays are all flat or have no corners. The four geodesic rays divide the plane into four quadrant regions. Applying Lemma 3.17 to these four quadrant regions, we have that \mathcal{C} is isometric to the subgraph with $h = 0$ in Fig. 8. By similar arguments to Claim 3.25, we get $\mathcal{C} = \mathcal{M}$.

Subcase 3-2-2 Assume $\deg(x, \delta\mathcal{M}) = 3$ and $\deg(p, \delta\mathcal{M}) \leq 3$ for any $p \in \delta\mathcal{M} \cap \mathcal{C}$. Let y, z, w be neighbors of x in $\delta\mathcal{M}$, and $y(z, w, \text{ resp.})$ be up-vertical(left-horizontal, right-horizontal, resp.) to x .

Subcase 3-2-2-1 Assume $\deg(q, \delta\mathcal{M}) = 3$ for some $(x \neq)q \in \delta\mathcal{M} \cap \mathcal{C}$.

We may assume $q \in \alpha_3$ with boundary neighbors p, s, t , and p is in the geodesic subpath $\hat{\alpha}$ between x and q . By similar arguments of Subcase 3-2-1, we get two flat boundary geodesic rays α_1, α_2 from x through y, z respectively, two flat boundary geodesic rays α_3, α_4 from q through s, t respectively, and $s(t, \text{ resp.})$ is right-horizontal(down-vertical, resp.) to q . Using Corollary 3.18 for geodesic boundary line $\alpha_1 \cup \alpha_2, \alpha_3 \cup \alpha_4, \alpha_1 \cup \hat{\alpha} \cup \alpha_3, \alpha_2 \cup \hat{\alpha} \cup \alpha_4$, we have $\mathcal{C} = \mathcal{C}_1 \cup \hat{\alpha} \cup \mathcal{C}_2$, where \mathcal{C}_1 is the quadrant domain enclosed by $\alpha_1 \cup \alpha_2$ and \mathcal{C}_2 is the quadrant domain enclosed by $\alpha_3 \cup \alpha_4$. This yields that $\hat{\alpha}$ is of length at most two by Lemma 3.2. By similar arguments in Claim 3.25, we have $\mathcal{M} = \mathcal{C}$ and it is isometric to the subgraph in Fig. 8 with $1 \leq h \leq 2$, or Fig. 10.

Subcase 3-2-2-2 Assume $\deg(q, \delta\mathcal{M}) = 2$ for any $x \neq q \in \delta\mathcal{M}$.

By similar arguments in Subcase 3-2-1, we get three boundary geodesic rays $\alpha_1, \alpha_2, \alpha_3$ from x through y, z, w respectively and α_1 is flat or vertical. If α_3 has corners, then α_2 is horizontal by Corollary 3.5 and the facts that $\delta\mathcal{M}$ is geodesic and contains no loops. So that we can assume α_2 is horizontal by symmetry of α_2, α_3 . Let $\mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3$ be regions enclosed by $\alpha_1 \cup \alpha_2, \alpha_1 \cup \alpha_3, \alpha_2 \cup \alpha_3$. We get $\mathcal{C} \cap \mathcal{D}_1 = \mathcal{D}_1$ or $\mathcal{C} \cap \mathcal{D}_1 = \alpha_1 \cup \alpha_2$ by Lemma 3.17.

We claim $\mathcal{C} \cap \mathcal{D}_3 = \mathcal{D}_3$ or $\mathcal{C} \cap \mathcal{D}_3 = \alpha_2 \cup \alpha_3$. Otherwise, recalling $\deg(q, \delta\mathcal{M}) = 2$ for any $x \neq q \in \delta\mathcal{M}$ and $\delta\mathcal{M}$ contains no loops, we have another boundary geodesic line β with $\beta \cap (\alpha_1 \cup \alpha_2 \cup \alpha_3) = \emptyset$. By Lemma 3.9, there is a vertical edge (a, b) in β such that a is up-vertical to b . Then one can find one geodesic ray β_1 in β from a , which doesn't contain b , is above

the horizontal line \hat{l} through b . Then we get a contradiction by Corollary 3.10 for two disjoint geodesic rays α_2 and β_1 , which proves the claim.

Similarly, we have $\mathcal{C} \cap \mathcal{D}_2 = \mathcal{D}_2$ or $\mathcal{C} \cap \mathcal{D}_3 = \alpha_1 \cup \alpha_3$. Note that $\alpha_1, \alpha_2, \alpha_3 \subset \delta\mathcal{M}$. Thus one of them is isolated and it contradicts Lemma 3.2.

Finally, we prove the result. \square

4. GEOMETRY OF HIGH-DIMENSIONAL MINIMAL SUBGRAPHS

For $x \in \mathbb{R}^n$ and $r > 0$, denote by $\hat{S}_r(x) = \{y \in \mathbb{R}^n : \max_{1 \leq i \leq n} |y_i - x_i| = r\}$

the ∞ -normed sphere centered at x of radius r . We say $\mathcal{I}^k \subset \mathbb{R}^n$ is a unit k -cube if it is a k dimensional cube with all edges of length one. Similar to the setting for \mathbb{Z}^2 , we identify the set $\{v_1, v_2, \dots, v_{2^k}\}$ with some k -cube \mathcal{I}^k if $\{v_1, v_2, \dots, v_{2^k}\}$ are exactly the vertices of \mathcal{I}^k , where $k = 0, 1, \dots, n$. So that any subgraph \mathcal{G} in \mathbb{Z}^n can be identified with a n -dimensional cell complex or a geometric space in \mathbb{R}^n , called the *geometric realization* of \mathcal{G} , which includes all cells whose vertices are contained in \mathcal{G} . Given any subgraph $\mathcal{G} \subset \mathbb{Z}^n$, let \mathcal{G}^k be the k -skeleton of \mathcal{G} (i.e. the union of k -cubes of \mathcal{G}) for $k = 0, 1, \dots, n$.

Proposition 4.1. *For any minimal subgraph \mathcal{M} in \mathbb{Z}^n , every connected component of \mathcal{M} is also minimal.*

Proof. Assume \mathcal{M}_1 is one of connected components of \mathcal{M} . One can restrict the minimal recurrents of \mathcal{M} to \mathcal{M}_1 , which yields the result by Corollary 2.3. \square

Lemma 4.2. *Given any minimal subgraph $\mathcal{M} \subset \mathbb{Z}^n$ and any $x \in \mathcal{M}$, then $\deg(x) \geq n$. Moreover, it never occurs that $\deg(x) = \deg(y) = n$ for any $y \sim x$ in \mathcal{M} .*

Proof. If $\deg(x) < n$, then the currents flowing out of x are at least $2n - \deg(x) > n$, but the currents flowing into x are at most $\deg(x) < n$. This is impossible by Lemma 2.3.

If $\deg(x) = \deg(y) = n$, then applying Lemma 2.3 to x , one can get that the current from y to x is one. By symmetry of x and y , one get the current from x to y is one. This is impossible. \square

Lemma 4.3. *If $\mathcal{M} \subset \mathbb{Z}^n$ is minimal, then both \mathcal{M} and $\delta\mathcal{M}$ are infinite.*

Proof. If not, then $\delta\mathcal{M} \subset B_r$ with some $r \in \mathbb{N}$. Hence $\mathcal{M} \subset B_r$ or $\mathcal{M}^c \subset B_r$. One can get a new graph \mathcal{M}' by removing B_r from \mathcal{M} or adding B_r to \mathcal{M} . Comparing \mathcal{M} with \mathcal{M}' , one easily sees that \mathcal{M} is not minimal. \square

Proposition 4.4. *If $\mathcal{M} \subset \mathbb{Z}^n$ is minimal, then we have $|\delta\mathcal{M} \cap \hat{B}_r(x)| \leq c(n)r^{n-1}$, where $c(n)$ is a constant depending only on n .*

Proof. One can get a new graph \mathcal{M}' by removing $\hat{B}_r(x)$ from \mathcal{M} for any $x \in \mathcal{M}$ and $r \in \mathbb{N}$. Comparing $|\partial\mathcal{M} \cap E_{\hat{B}_r(x)}|$ with $|\partial\mathcal{M}' \cap E_{\hat{B}_r(x)}|$ and using

the minimality of \mathcal{M} , one can get

$$\begin{aligned} c(n)|\delta\mathcal{M} \cap \hat{B}_r(x)| &\leq |\partial\mathcal{M} \cap E_{\hat{B}_r(x)}| \\ &\leq |\partial\mathcal{M}' \cap E_{\hat{B}_r(x)}| \\ &\leq 2n|\delta\hat{B}_r(x) \cap \mathcal{M}| \\ &\leq 4n^2(2r+1)^{n-1}. \end{aligned}$$

Thus we complete the proof. \square

Now we are ready to prove Theorem 1.9.

Proof of Theorem 1.9. It suffices to show the restriction of some minimal currents I of \mathcal{M} to $\mathcal{M}^3 \cup \mathcal{M}^2$, denoted by \hat{I} , are well defined, and then they are minimal currents, i.e. the sum of the restrictive currents \hat{I} for any vertex of \mathbb{Z}^n is zero; see Lemma 2.3. Note that \mathcal{M}^1 may be nonempty. So that it suffices to show the currents I on $\mathcal{M}^3 \cup \mathcal{M}^2 \cap \mathcal{M}^1$ agree with those induced naturally by $\mathcal{M}^3 \cup \mathcal{M}^2$. Precisely speaking, the minimal current on any (x, y) agrees with that induced by $\mathcal{M}^3 \cup \mathcal{M}^2$, where $x \sim y$ for $x \in \mathcal{M}^3 \cup \mathcal{M}^2$ and $y \in \mathcal{M} - \mathcal{M}^3 \cup \mathcal{M}^2$. This yields that $\deg(x) \geq 3$, $\deg(y) \geq 3$ by Lemma 4.2.

If $\deg(y) \geq 5$, then $y \in \mathcal{M}^3 \cup \mathcal{M}^2$. If $\deg(y) = 3$, then the result is true.

Thus we only need to consider the case $\deg(y) = 4$. Note that if $\deg(x) \geq 5$, one easily deduces that there exists a square in \mathcal{M} containing the edge (x, y) . Hence $y \in \mathcal{M}^3 \cup \mathcal{M}^2$. We have the following two cases.

Case 1. $\deg(x) = 3$, $\deg(y) = 4$.

Subcase 1-1. The neighbors of x and y in \mathcal{M} are as in Fig. 45. Using Lemma 2.3 to x and y , we have that the currents on the oriented edges (y, x) , (p, y) , (u, y) , (v, y) are all one, indicated by arrows in Fig. 45. Applying Lemma 2.3 to p , we obtain that the current on (w, z) is one, and hence $w \in \mathcal{M}$. By symmetry, we get $s, t \in \mathcal{M}$. Then the currents on the oriented edges (t, z) , (y, z) , (t, z) , (w, z) are all one and the sum of the currents flowing into $z \geq 4 - 2 = 2 > 0$. This is a contradiction.

Subcase 1-2. The neighbors of x and y in \mathcal{M} are as in Fig. 46. Using Lemma 2.3 to x and y , we have that the currents on the oriented edges (y, x) , (z, y) are both one. Since $y \in \mathcal{M}^1$, $u, v, w \notin \mathcal{M}$ and the sum of the currents flowing out of $z \geq 4 - 2 = 2 > 0$. This is a contradiction.

Subcase 1-3. The neighbors of x and y in \mathcal{M} are as in Fig. 47. Using Lemma 2.3 to x and y , we have that the currents on the oriented edges (y, x) , (u, y) , (v, y) are all one, indicated by arrows in Fig. 47. Using Lemma 2.3 for u, v , one can get the currents on the oriented edges (s, u) , (t, v) are both one, and hence $s, t \in \mathcal{M}$. Then the currents on the oriented edges (w, p) , (y, p) , (s, p) , (t, p) are all one. Thus the sum of the currents flowing in of $p \geq 4 - 2 = 2 > 0$. This is a contradiction.

Subcase 1-4. The neighbors of x and y in \mathcal{M} are as in Fig. 48 or Fig. 49. Using Lemma 2.3 to x and y , we deduce that the currents on oriented edges




FIGURE 45.




FIGURE 46.




FIGURE 47.




FIGURE 48.




FIGURE 49.

must be indicated by arrows in Fig. 48 or Fig. 49. Noting that $y \notin \mathcal{M}^3 \cup \mathcal{M}^2$, we have that the sum of currents flowing out of $z \geq 4 - 2 = 2 > 0$. This is a contradiction.

Case 2. $\deg(x) = \deg(y) = 4$.

Subcase 2-1. The neighbors of x in \mathcal{M} are as in Fig. 50. Then it is obvious that $y \in \mathcal{M}^3 \cup \mathcal{M}^2$, which is impossible.

Subcase 2-2. The neighbors of x and y in \mathcal{M} are as in Fig. 51. If $p \notin \mathcal{M}$, then using Lemma 2.3 to z , we have that the current on the oriented edge




FIGURE 50.




FIGURE 51.




FIGURE 52.

(y, z) is one. So that we get the current on the oriented edge (x, y) is one by applying Lemma 2.3 to y , which agrees with that induced by $\mathcal{M}^3 \cup \mathcal{M}^2$, and we finish the proof in this setting. By symmetry of z, w, u , the remaining setting in this subcase is that $p, q, v \in \mathcal{M}$. In this case, by direct computation the currents flowing into $t \geq 4 - 2 = 2 > 0$, which is impossible.

Subcase 2-3. The neighbors of x and y in \mathcal{M} are as in Fig. 52. It follows that $u, v, w \notin \mathcal{M}$ by $y \in \mathcal{M}^1$. By Lemma 2.3 for z , we have that the current on the oriented edge (y, z) is one. Recalling that $\deg(y) = 4$ and using Lemma 2.3 to y , we deduce that the current on the oriented edge (x, y) is one, which agrees with that induced by $\mathcal{M}^3 \cup \mathcal{M}^2$. Hence we finish the proof in this subcase. \square

Lemma 4.5. *For any minimal subgraph $\mathcal{M} \subset \mathbb{Z}^n$ and any point $x \in \mathcal{M}$, $\mathcal{M}^n \neq \emptyset$ and there exists a positive constant $c_1 = c_1(n)$, depending only on n , such that*

$$d(x, \mathcal{M}^n) \leq c_1.$$

Proof. For any $x \in \mathcal{M}$, let r be a positive integer with $\hat{B}_r(x) \cap \mathcal{M}^n = \emptyset$. Then $\hat{B}_r(x) \cap \mathcal{M} \subset \delta \mathcal{M}$. We can get a new graph \mathcal{M}' by removing $\hat{B}_{r-1}(x) \cap \mathcal{M}$ from \mathcal{M} .

Comparing $E_{\hat{B}_r(x)} \cap \partial\mathcal{M}$ with $E_{\hat{B}_r(x)} \cap \partial\mathcal{M}'$, one can deduce that by the minimality of \mathcal{M}

$$\begin{aligned} 2n|\delta\hat{B}_r(x) \cap \mathcal{M}| &= 2n(|\hat{B}_r(x) \cap \mathcal{M}| - |\hat{B}_{r-1}(x) \cap \mathcal{M}|) \\ &\geq |E_{\hat{B}_r(x)} \cap \partial\mathcal{M}'| \\ &\geq |E_{\hat{B}_r(x)} \cap \partial\mathcal{M}| \\ &\geq |\hat{B}_{r-1}(x) \cap \mathcal{M}|. \end{aligned}$$

This gives

$$|\hat{B}_r(x) \cap \mathcal{M}| \geq (1 + \frac{1}{2n})|\hat{B}_{r-1}(x) \cap \mathcal{M}|.$$

Therefore, we obtain that

$$|\hat{B}_r(x) \cap \mathcal{M}| \geq (1 + \frac{1}{2n})^r.$$

On the other hand, since $\mathcal{M} \subset \mathbb{Z}^n$,

$$|\hat{B}_r(x) \cap \mathcal{M}| \leq (1 + 2r)^n.$$

Hence we have

$$r \leq c_1(n),$$

for some positive constant $c_1(n)$ depending only on n . □

Corollary 4.6. *Given any minimal subgraph $\mathcal{M} \subset \mathbb{Z}^n$ and any point $x \in \mathcal{M}$, let $\mathcal{C}(\mathcal{M}^n \cap \hat{B}_r(x))$ be the set of connected components of $\mathcal{M}^n \cap \hat{B}_r(x)$. Then*

$$\liminf_{r \rightarrow \infty, A \in \mathcal{C}(\mathcal{M}^n \cap \hat{B}_r(x))} \frac{|\delta A|}{|A|} = 0.$$

In particular,

$$\limsup_{r \rightarrow \infty, A \in \mathcal{C}(\mathcal{M}^n \cap \hat{B}_r(x))} |A| = \infty.$$

Proof. The idea of the proof is similar to that of Lemma 4.5.

We argue by contradiction. Then this yields

$$\frac{|\delta A|}{|A|} \geq c(n) > 0,$$

for any $A \in \mathcal{C}(\mathcal{M}^n \cap \hat{B}_r(x))$, any $r \in \mathbb{N}$ and some positive constant $c(n)$ depending only on n . Hence we have

$$(6) \quad c(n)|\mathcal{M} \cap \hat{B}_r(x)| \leq |\delta\mathcal{M} \cap \hat{B}_r(x)|.$$

We can get a new graph \mathcal{M}' by removing $\hat{B}_{r-1}(x) \cap \mathcal{M}$ from \mathcal{M} . Comparing \mathcal{M} with \mathcal{M}' , using (6) and applying the minimality of \mathcal{M} , we have that

$$\begin{aligned} 0 &\geq |E_{\hat{B}_r(x)} \cap \partial\mathcal{M}| - |E_{\hat{B}_r(x)} \cap \partial\mathcal{M}'| \\ &\geq |\delta\mathcal{M} \cap \hat{B}_{r-1}(x)| - 2n|\mathcal{M} \cap \delta\hat{B}_r(x)|, \end{aligned}$$

and hence

$$(7) \quad 2n|\mathcal{M} \cap \delta \hat{B}_r(x)| \geq |\delta \mathcal{M} \cap \hat{B}_{r-1}(x)| \geq c(n)|\mathcal{M} \cap \hat{B}_{r-1}(x)|.$$

Note that

$$|\mathcal{M} \cap \delta \hat{B}_r(x)| = |\mathcal{M} \cap \hat{B}_r(x)| - |\mathcal{M} \cap \hat{B}_{r-1}(x)|,$$

which yields

$$|\mathcal{M} \cap \hat{B}_r(x)| \geq \left(1 + \frac{c(n)}{2n}\right)r,$$

for any $r \in \mathbb{N}$. But recall that

$$|\mathcal{M} \cap \hat{B}_r(x)| \leq (2r + 1)^n.$$

This is impossible for sufficiently large r . \square

Lemma 4.7. *For any minimal subgraph $\mathcal{M} \subset \mathbb{Z}^n$ and any point $x \in \mathcal{M}$, we have*

$$|(\mathcal{M} - \mathcal{M}^n) \cap \hat{B}_r(x)| \leq 2n|\mathcal{M}^n \cap N_1(\mathcal{M} - \mathcal{M}^n) \cap \hat{B}_r(x)|,$$

where $N_1(\mathcal{M} - \mathcal{M}^n) := \{x \in \mathbb{Z}^n | d(x, \mathcal{M} - \mathcal{M}^n) \leq 1\}$.

Proof. Observe the new graph \mathcal{M}' obtained by removing $\mathcal{M} - \mathcal{M}^n$ from $\mathcal{M} \cap \hat{B}_r(x)$ and note that $\mathcal{M} - \mathcal{M}^n \subset \delta \mathcal{M}$. Comparing $E_{\hat{B}_r(x)} \cap \partial \mathcal{M}'$ with $E_{\hat{B}_r(x)} \cap \partial \mathcal{M}$ and recalling that \mathcal{M} is minimal, we get that

$$\begin{aligned} |(\mathcal{M} - \mathcal{M}^n) \cap \hat{B}_r(x)| &\leq |E_{\hat{B}_r(x)} \cap \partial \mathcal{M}| \\ &\leq |E_{\hat{B}_r(x)} \cap \partial \mathcal{M}'| \\ &\leq 2n|\mathcal{M}^n \cap N_1(\mathcal{M} - \mathcal{M}^n) \cap \hat{B}_r(x)|. \end{aligned}$$

\square

Next, we prove Theorem 1.11 and Theorem 1.12.

Proof of Theorem 1.11. The natural embedding $\mathcal{M}^n \subset \mathcal{M}$ is surely a rough isometry by Lemma 4.5. Recall that $\deg(x) \leq 2n$, $|\hat{B}_{c_1}(x)| \leq (2c_1 + 1)^n$, and one has $|\hat{B}_{c_1}(x) \cap \mathcal{M}^n| \geq 1$ for any given $x \in \mathcal{M}$ by Lemma 4.5. One can show the inequality (3) in Theorem 1.11 for sufficiently large r .

It is obvious that $\mathcal{M}^n \cap N_1(\mathcal{M} - \mathcal{M}^n) \subset \delta \mathcal{M}^n$.

Hence we have

$$\begin{aligned} |(\mathcal{M} - \mathcal{M}^n) \cap \hat{B}_r(x)| &\leq 2n|\mathcal{M}^n \cap N_1(\mathcal{M} - \mathcal{M}^n) \cap \hat{B}_r(x)| \\ &\leq 2n|\delta \mathcal{M}^n \cap \hat{B}_r(x)|. \end{aligned}$$

By $\delta \mathcal{M} = \delta \mathcal{M}^n \cup (\mathcal{M} - \mathcal{M}^n)$, direct computation yields that

$$\frac{|\delta \mathcal{M}^n \cap \hat{B}_r(x)|}{|\delta \mathcal{M} \cap \hat{B}_r(x)|} \geq \frac{1}{1 + 2n}.$$

\square

Proof of Theorem 1.12. Suppose it is not true, let P be one of two parallel hyperplanes and $\bar{\mathcal{M}}$ be the usual distance projection image from \mathcal{M} to P . Assume the distance of two parallel hyperplanes is l and \mathcal{M}' is the new graph obtained by removing $\hat{B}_{r-1}(x)$ from \mathcal{M} for any given $x \in \mathcal{M}$. Then one can show that

$$(8) \quad 2|\bar{\mathcal{M}} \cap \hat{B}_r(x)| \leq |\delta\mathcal{M} \cap \hat{B}_r(x)| \leq l|\bar{\mathcal{M}} \cap \hat{B}_r(x)|.$$

$$(9) \quad |\bar{\mathcal{M}} \cap \delta\hat{B}_r(x)| \leq |\mathcal{M} \cap \delta\hat{B}_r(x)| \leq l|\bar{\mathcal{M}} \cap \delta\hat{B}_r(x)|.$$

On the other hand, using (8), (9) and the minimality of \mathcal{M} , we have

$$\begin{aligned} 0 &\geq |E_{\hat{B}_r(x)} \cap \partial\mathcal{M}| - |E_{\hat{B}_r(x)} \cap \partial\mathcal{M}'| \\ &\geq 2|\bar{\mathcal{M}} \cap \hat{B}_{r-1}(x)| - 2n|\mathcal{M} \cap \delta\hat{B}_r(x)| \\ &\geq 2|\bar{\mathcal{M}} \cap \hat{B}_{r-1}(x)| - 2nl|\bar{\mathcal{M}} \cap \delta\hat{B}_r(x)|, \end{aligned}$$

which yields that

$$|\bar{\mathcal{M}} \cap \hat{B}_r(x)| \geq (1 + \frac{1}{nl})^r.$$

But since $\bar{\mathcal{M}} \subset \mathbb{Z}^{n-1}$,

$$|\bar{\mathcal{M}} \cap \hat{B}_r(x)| \leq (2r + 1)^{n-1}.$$

This is a contradiction when r is sufficiently large. \square

Before introducing a maximum principle property of minimal subgraphs, we adopt some notations.

Let $p_i : \mathbb{Z}^n \rightarrow \mathbb{Z}^{n-1}$ be the i -th projection for any integer $1 \leq i \leq n$, where

$$p_i(x_1, x_2, \dots, x_n) = (x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$$

for any $x = (x_1, x_2, \dots, x_n) \in \mathbb{Z}^n$.

For any subgraph $\mathcal{M} \subset \mathbb{Z}^n$ and any $\hat{x} = (x_1, x_2, \dots, x_{n-1}) \in p_i(\mathcal{M})$, we define

$$H_i(\hat{x}, \mathcal{M}) := \sup_{y=(y_1, \dots, y_n) \in \mathcal{M}, p_i(y)=\hat{x}} y_i$$

and

$$h_i(x, \mathcal{M}) := \inf_{y=(y_1, \dots, y_n) \in \mathcal{M}, p_i(y)=\hat{x}} y_i.$$

Note that $H_i(\hat{x}, \mathcal{M}), h_i(\hat{x}, \mathcal{M})$ may be infinite.

Proposition 4.8. *Suppose that $\mathcal{M} \subset \mathbb{Z}^n$ is minimal and given two finite subsets $\Omega_1, \Omega_2 := \{x \in \mathbb{Z}^{n-1} | d(x, \Omega_1) \leq 1\} \subset p_i(\mathcal{M}) \subset \mathbb{Z}^{n-1}$, such that the geometric realization of Ω_1 in \mathbb{R}^{n-1} is the closure of a bounded open domain. Then neither*

$$(10) \quad \max_{\hat{x} \in \Omega_2 - \Omega_1} H_i(\hat{x}, \mathcal{M}) < H := \max_{\hat{x} \in \Omega_1} H_i(\hat{x}, \mathcal{M}) < +\infty,$$

nor

$$(11) \quad \min_{\hat{x} \in \Omega_2 - \Omega_1} h_i(\hat{x}, \mathcal{M}) > h := \min_{\hat{x} \in \Omega_1} h_i(\hat{x}, \mathcal{M}) > -\infty.$$

holds for any integer $1 \leq i \leq n$.

Proof. It suffices to prove (10) fails for $i = 1$. We argue by contradiction.

Consider the nonempty set $\Omega := \{\hat{x} \in \Omega_2 : H(\hat{x}, \mathcal{M}) = H\}$ and the set $\tilde{\Omega} := \{x \in \mathbb{Z}^n : p_1(x) \in \Omega, x_1 = H\} \subset \{H\} \times \mathbb{Z}^{n-1}$. Note that $\Omega \subset \Omega_1$ and $\tilde{\Omega} \subset \mathcal{M}$, by the assumption (10). Now one can obtain a new graph \mathcal{M}' from \mathcal{M} by deleting the finite subset $\tilde{\Omega}$.

Using (10) again, one easily deduces that

$$|E_B \cap \partial\mathcal{M}| - |E_B \cap \partial\mathcal{M}'| \geq |\hat{\partial}\tilde{\Omega}| > 0,$$

where B is any finite ball containing $\tilde{\Omega}$ and $\hat{\partial}\tilde{\Omega}$ is the boundary edge of $\tilde{\Omega}$ in $\{H\} \times \mathbb{Z}^{n-1}$. This is impossible by the minimality of \mathcal{M} . \square

5. OPEN PROBLEMS

In this section, let \mathcal{M} always be a minimal subgraph in \mathbb{Z}^n . We propose some open problems on geometry and topology of high-dimensional minimal subgraphs.

The first problem is the stronger version of Theorem 1.11.

Problem 5.1. *Are the following true:*

$$\lim_{r \rightarrow \infty} \frac{|\mathcal{M}^n \cap \hat{B}_r(x)|}{|\mathcal{M} \cap \hat{B}_r(x)|} = 1 \quad \text{and} \quad \lim_{r \rightarrow \infty} \frac{|\delta\mathcal{M}^n \cap \hat{B}_r(x)|}{|\delta\mathcal{M} \cap \hat{B}_r(x)|} = 1?$$

The second problem is about the number of connected components and some “big” connected component of any minimal subgraph.

Problem 5.2. *Do \mathcal{M}^n and \mathcal{M} have only finitely many connected components? Does at least one of these connected components contains a subgraph isometric to $\underbrace{\mathbb{Z}_{\geq 0} \times \cdots \times \mathbb{Z}_{\geq 0}}_{n \text{ times}}$?*

Motivated by positive density at infinity of any minimal hypersurface in \mathbb{R}^n , it is natural to ask the growth rates of any minimal subgraph and its boundary.

Problem 5.3. *Whether do there exist positive constants $c_2 = c_2(n), c_3 = c_3(n), c_4 = c_4(n)$ and $c_5 = c_5(n)$ depending only on n such that*

$$c_2 r^n \leq |\mathcal{M} \cap \hat{B}_r(x)| \leq c_3 r^n \quad \text{and} \quad c_4 r^{n-1} \leq |\delta\mathcal{M} \cap \hat{B}_r(x)| \leq c_5 r^{n-1}?$$

Remark 5.4. \mathcal{M} can be replaced by \mathcal{M}^n by Lemma 4.5 and Lemma 4.7.

Note that the monotonicity formula of any minimal subgraph or its boundary does not hold in general; see Figure 8. Proposition 4.4 gives the upper bounds in Problem 5.3. So that it suffices to consider the lower bounds in Problem 5.3.

The last question is as follows.

Problem 5.5. For a minimal subgraph $\mathcal{M} \subset \mathbb{Z}^n$, do bounded connected components of \mathcal{M}^n exist? How to characterize the bounded connected components of \mathcal{M}^n if they exist?

Acknowledgements. We thank Florentin Münch for helpful discussions and suggestions on the 1-Laplacian and the definition of area-minimizing subgraphs in \mathbb{Z}^n . B.H. is supported by NSFC, no. 12371056 and Shanghai Science and Technology Program [Project No. 22JC1400100]. Z.H. is supported by NSFC, no.12301094 and Guangzhou Basic and Applied Basic Research Foundation No. 2024A04J3483.

Data Available Statement. The data used to support the findings of this study are available from the corresponding author upon request.

REFERENCES

- [Bar17] Martin T. Barlow. *Random walks and heat kernels on graphs*, volume 438 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2017.
- [BBI01] D. Burago, Yu. Burago, and S. Ivanov. *A course in metric geometry*, volume 33 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001.
- [BDGG69] E. Bombieri, E. De Giorgi, and E. Giusti. Minimal cones and the Bernstein problem. *Invent. Math.*, 7:243–268, 1969.
- [Ber27] Serge Bernstein. Über ein geometrisches Theorem und seine Anwendung auf die partiellen Differentialgleichungen vom elliptischen Typus. *Math. Z.*, 26(1):551–558, 1927.
- [Cha16] K. C. Chang. Spectrum of the 1-Laplacian and Cheeger’s constant on graphs. *J. Graph Theory*, 81(2):167–207, 2016.
- [DG61] Ennio De Giorgi. *Frontiere orientate di misura minima*. Editrice Tecnico Scientifica, Pisa, 1961. Seminario di Matematica della Scuola Normale Superiore di Pisa, 1960–61.
- [DG65] Ennio De Giorgi. Una estensione del teorema di Bernstein. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)*, 19:79–85, 1965.
- [Fle62] Wendell H. Fleming. On the oriented Plateau problem. *Rend. Circ. Mat. Palermo (2)*, 11:69–90, 1962.
- [FM19] Morteza Fotouhi and Amir Moradifam. General least gradient problems with obstacle. *Calc. Var. Partial Differential Equations*, 58(5):Paper No. 182, 19, 2019.
- [Gí18] Wojciech Górny. Planar least gradient problem: existence, regularity and anisotropic case. *Calc. Var. Partial Differential Equations*, 57(4):Paper No. 98, 27, 2018.
- [Giu84] Enrico Giusti. *Minimal surfaces and functions of bounded variation*, volume 80 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1984.
- [GM21] Wojciech Górny and José M. Mazón. Least gradient functions in metric random walk spaces. *ESAIM Control Optim. Calc. Var.*, 27:Paper No. S28, 32, 2021.
- [HB10] M. Hein and T. Bühler. An inverse power method for nonlinear eigenproblems with applications in 1-spectral clustering and sparse pca. *Advances in Neural Information Processing Systems (NIPS)*, pages 847–855, 2010. MIT Press, Cambridge, MA.

- [JMN18] Robert L. Jerrard, Amir Moradifam, and Adrian I. Nachman. Existence and uniqueness of minimizers of general least gradient problems. *J. Reine Angew. Math.*, 734:71–97, 2018.
- [Juu05] Petri Juutinen. p -Harmonic approximation of functions of least gradient. *Indiana Univ. Math. J.*, 54(4):1015–1029, 2005.
- [Mir67] M. Miranda. Comportamento delle successioni convergenti di frontiere minimali. *Rend. Sem. Mat. Univ. Padova*, 38:238–257, 1967.
- [Mor17] Amir Moradifam. Least gradient problems with Neumann boundary condition. *J. Differential Equations*, 263(11):7900–7918, 2017.
- [Mor18] Amir Moradifam. Existence and structure of minimizers of least gradient problems. *Indiana Univ. Math. J.*, 67(3):1025–1037, 2018.
- [MPLRT16] José M. Mazón, Mayte Pérez-Llanos, Julio D. Rossi, and Julián Toledo. A nonlocal 1-Laplacian problem and median values. *Publ. Mat.*, 60(1):27–53, 2016.
- [MRSdL14] José M. Mazón, Julio D. Rossi, and Sergio Segura de León. Functions of least gradient and 1-harmonic functions. *Indiana Univ. Math. J.*, 63(4):1067–1084, 2014.
- [MSDTM23] José M. Mazón, Marcos Solera-Diana, and J. Julián Toledo-Melero. *Variational and diffusion problems in random walk spaces*, volume 103 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser/Springer, Cham, 2023.
- [MT23] José M. Mazón and Julián Toledo. Cahn-Hilliard equations on random walk spaces. *Anal. Appl. (Singap.)*, 21(4):959–1000, 2023.
- [Roc70] R. Tyrrell Rockafellar. *Convex analysis*. Princeton Mathematical Series, No. 28. Princeton University Press, Princeton, N.J., 1970.
- [RW98] R. Tyrrell Rockafellar and Roger J.-B. Wets. *Variational analysis*, volume 317 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1998.
- [Sim68] James Simons. Minimal varieties in riemannian manifolds. *Ann. of Math. (2)*, 88:62–105, 1968.
- [SWZ92] Peter Sternberg, Graham Williams, and William P. Ziemer. Existence, uniqueness, and regularity for functions of least gradient. *J. Reine Angew. Math.*, 430:35–60, 1992.
- [Woe00] W. Woess. *Random walks on infinite graphs and groups*. Number 138 in Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2000.
- [Zun19] Andres Zuniga. Continuity of minimizers to weighted least gradient problems. *Nonlinear Anal.*, 178:86–109, 2019.

ZUNWU HE: SCHOOL OF MATHEMATICS, SOUTH CHINA UNIVERSITY OF TECHNOLOGY, 510641, GUANGZHOU, CHINA

Email address: hzwmath789@scut.edu.cn

BOBO HUA: SCHOOL OF MATHEMATICAL SCIENCES, LMNS, FUDAN UNIVERSITY, SHANGHAI 200433, CHINA; SHANGHAI CENTER FOR MATHEMATICAL SCIENCES, JIANGWAN CAMPUS, FUDAN UNIVERSITY, NO. 2005 SONGHU ROAD, SHANGHAI 200438, CHINA.

Email address: bobohua@fudan.edu.cn