Existence of traveling wave solutions in continuous OV models

Kota Ikeda[∗] , Toru Kan† , Toshiyuki Ogawa‡

Abstract

In traffic flow, self-organized wave propagation, which characterizes congestion, has been reproduced in macroscopic and microscopic models. Hydrodynamic models, a subset of macroscopic models, can be derived from microscopic-level car-following models, and the relationship between these models has been investigated. However, most validations have relied on numerical methods and formal analyses; therefore, analytical approaches are necessary to rigorously ensure their validity. This study aims to investigate the relationship between macroscopic and microscopic models based on the properties of the solutions corresponding to congestion with sparse and dense waves. Specifically, we demonstrate the existence of traveling wave solutions in macroscopic models and investigate their properties.

1 Introduction

Various aspects of traffic dynamics and congestion formation present challenges for mathematicians and physicists, drawing on more than 80 years of engineering experience. In the early 1990s, traffic flow was recognized as a non-equilibrium system. Empirical evidence indicates multiple dynamic phases in traffic flow and dynamic phase transitions. Several mathematical models have been proposed to explain these empirical results, with some models qualitatively reproduce all known features of traffic flows, including localized and extended forms of congestion, self-organized propagation of stop-and-go waves, and observed hysteresis effects. These characteristics are criteria for good traffic models, as noted by Helbing [1]. However, many of these models have only been validated using numerical techniques or formal analyses and have not yet been rigorously proven.

Mathematical models can be categorized into macroscopic and microscopic models. These models are often interrelated through Taylor and mean-field approximations. Payne [2] developed a macroscopic model based on the compressible fluid equation and the dynamic velocity equation. It was demonstrated in [3] that the linear instability condition of the Payne model aligns precisely with those of well-known microscopic models, such as the car-following model or the optimal velocity model proposed by [4]. However, the Payne model produces shock-like waves that compromise numerical robustness. Hence, Kühne [5] and Kerner and Konhäuser [6] introduced models incorporating artificial viscosity terms to the Payne model. In these studies, uniform flows were destabilized based on density, and numerical calculations confirmed the stable formation of vehicle clusters. Lee et al. [7] attempted to derive a fluid-dynamic model from a car-following model using a coarse-graining procedure to elucidate the relationship between these models. The authors verified this through numerical simulations, demonstrating that the macroscopic model based on the mean-field method quantitatively approximates the microscopic model.

As noted by [1], many macroscopic models, including those mentioned above, can be expressed in the following general form:

$$
\begin{cases}\n\frac{\partial \rho}{\partial t} + \frac{\partial (\rho v)}{\partial x} = 0, \\
\frac{\partial v}{\partial t} + v \frac{\partial v}{\partial x} = -\frac{1}{\rho} \frac{d(P(\rho))}{dx} + \kappa(\rho) \frac{\partial^2 v}{\partial x^2} + \frac{1}{\tau} (V(\rho^{-1}) - v),\n\end{cases}
$$
\n(1.1)

[∗]School of Interdisciplinary Mathematical Sciences, Meiji University, ikeda@meiji.ac.jp

[†]Department of Mathematics, Osaka Metropolitan University

[‡]School of Interdisciplinary Mathematical Sciences, Meiji University

where $\rho = \rho(x, t)$ and $v = v(x, t)$ denote the density and velocity of the vehicles at x and t, respectively. The terms κ, τ, P and V represent viscosity-like quantities, relaxation time, traffic pressure, and equilibrium velocity, respectively. We neglected the diffusion effect of the density in the first equation and external forces, such as fluctuations, in (1.1). We assume that V and κ depend on ρ , and $P(\rho) \equiv -V(\rho^{-1})/2\tau$, where $\tau > 0$ is constant. In specific cases, V is often given by a sigmoid function

$$
V(u) \equiv V_0[\tanh(\beta(u - u_c)) + M], \qquad (1.2)
$$

where V_0, β, u_c, M are positive constants. The function V is known as the *optimal velocity function* [4]. The density-dependent function $\kappa = \kappa(\rho)$ has been assumed to be 0 [2], κ_0 [3, 5], κ_0/ρ [6], and $1/(6\tau\rho^2)$ [7], where κ_0 denotes a positive constant. We do not consider the case $\kappa(\rho) = 0$ in this paper. The derivations and analyses of continuous models that are not included in (1.1) have also been conducted (see for instance [8, 9]).

The microscopic optimal velocity model exhibits two typical types of collective motion depending on the density. In the low-density region, the distance between any two neighboring vehicles converges to a constant $t \to \infty$, known as *free flow*. When the density is relatively high, the distance oscillates over time, known as congestion or jamming. The transition between these two states occurs via a Hopf bifurcation, as shown in Fig. 12 of [10]. The characteristics of the global bifurcation diagram indicate that a congested state has only one congested region, and all periodic solutions on any other bifurcation branch associated with multiple congested regions are unstable. This implies that multiple congested regions merge as t increases and combine into a single lump. These observations in the microscopic model are valid for the macroscopic model. More precisely, the congestion phenomenon of vehicles in (1.1) can be considered as the dynamics of a traveling wave solution, which moves at a constant speed and forms a pulse shape (see Figure 1). If the initial state has a relatively high ρ , then all solutions transition to states with multiple pulses after a short time, as long as numerical calculations are feasible. Eventually, the multiple pulses merge into one. Therefore analyzing the single-pulse traveling wave solution in (1.1) is crucial to understand congestion phenomena in vehicles.

Generally, φ_a represents the (partial) derivative of φ with respect to a for a function φ that depends on variable a, i.e., $\varphi_a = \partial \varphi / \partial a$. If φ depends solely on one variable a, then we may use the symbol φ' , instead of φ_a . We make the following assumptions for the optimal velocity function V and viscosity coefficient κ throughout the study.

- (A1) $V \in C^1([0,\infty)) \cap C^2((0,\infty))$. $V'(u)$ is positive and bounded in $u > 0$, and converges to 0 as $u \to \infty$. Moreover, there is a global maximum point $U_M \in (0, \infty)$ of $V'(u)$ such that $V''(u) = 0$ attains a unique root at $u = U_M$.
- (A2) $\kappa \in C^1((0,\infty))$ and $\kappa > 0$ in $(0,\infty)$.

It follows from $(A1)$ that $V'(u)$ has no local minimum points. In some of our results, we additionally need the following assumption for the diffusion coefficient κ (see condition (C)):

$$
\int_{1}^{\infty} \frac{d\rho}{\rho^3 \kappa(\rho)} = \infty.
$$
 (H)

The traveling wave solution $(\rho(x, t), v(x, t)) = (\rho(z), v(z))$ in (1.1) is governed by

$$
\begin{cases} c\rho_z + (\rho v)_z = 0, \\ cv_z + vv_z = -\frac{1}{\rho} \frac{d(P(\rho))}{dz} + \kappa(\rho)v_{zz} + \frac{1}{\tau}(V(\rho^{-1}) - v), \end{cases}
$$
(1.3)

where c is the wave speed and $z = x + ct$ is the moving coordinate. As shown in Figure 1, the traveling wave solution to (1.1) approaches a constant outside the region where v is relatively small, implying the existence of a homoclinic orbit for a certain value of c.

Figure 1: Snapshots of solutions in (1.1) with the periodic boundary condition and in the microscopic optimal velocity model at (a) $t = 20$ and (b) $t = 880$. The optimal velocity function V is given by (1.2). Solid and dashed lines represent the graphs of v in (1.1) under $1/(6\tau\rho^2)$ (Lee et al. model) and $\kappa(\rho) = \kappa_0$ (Kühne model), respectively. The unfilled circles denote the pairs of $(x_i, dx_i/dt)$ in the microscopic optimal velocity model, where x_i represents the position of the *i*-th vehicle for $i = 1, \ldots, 77$. The numerical simulations were performed with parameters $L = 2.33$, $V_0 = 0.0168$, $M = 0.913$, $u_c = 0.025$, $\beta = 89.7$, $\tau = 0.5$, and $\kappa_0 = 1/7500$, where L denotes the length of the domain. The Fourier-spectral method was used in (1.1) for space discretization with a truncation wave number of 100, while the time discretization employed the classical Euler scheme with a time increment of 0.001. The classical fourth-order Runge-Kutta scheme was used for the microscopic optimal velocity model for time discretization with a time increment of 0.01.

We are interested in traveling wave solutions that connect constant steady states, imposing the condition:

$$
\lim_{z \to \pm \infty} (\rho(z), v(z)) = (\overline{\rho}_{\pm}, \overline{v}_{\pm})
$$
\n(1.4)

for certain $(\overline{\rho}_{\pm}, \overline{v}_{\pm}),$ where $\overline{v}_{\pm} = V((\overline{\rho}_{\pm})^{-1})$. If $\overline{\rho}_{+} > \overline{\rho}_{-}$ (resp. $\overline{\rho}_{+} < \overline{\rho}_{-}$), the corresponding solution is a traveling back solution (resp. traveling front solution). If $\overline{\rho}_+ = \overline{\rho}_-$, it is called a traveling pulse solution. We are also interested in a *periodic traveling wave solution*, which satisfies $(\rho(z+Z), v(z+Z)) = (\rho(z), v(z))$ for some $Z > 0$ instead of (1.4).

We obtain the following by integrating the first equation of (1.3) :

$$
\rho(z)(v(z) + c) = K \tag{1.5}
$$

for any z, where K is a constant. We assume $K \neq 0$ because of our interest in obtaining nonconstant solutions of (1.3). We obtain the following by substituting $u \equiv \rho^{-1} = (v+c)/K$ and $P(\rho) = -V(u)/2\tau$ into the second equation of (1.3):

$$
K^{2}uu_{z} = \frac{1}{2\tau}V'(u)uu_{z} + \kappa(u^{-1})Ku_{zz} + \frac{1}{\tau}(V(u) - Ku + c).
$$

This is equivalent to the following dynamical system:

$$
\begin{cases}\nu_z = w, \\
w_z = g_1(u)f(u) + g_2(u)h(u,\mu)w,\n\end{cases} (1.6)
$$

where $\mu = 2\tau K - 1$ and

$$
f(u) = \frac{Ku - V(u) - c}{K}, \quad h(u, \mu) = f'(u) + \mu, \quad g_1(u) = \frac{1}{\tau \kappa (u^{-1})}, \quad g_2(u) = \frac{u}{2\tau \kappa (u^{-1})}.
$$

The traveling back/front, traveling pulse, and periodic solutions of (1.3) correspond to a heteroclinic orbit, a homoclinic orbit, and a periodic orbit in (1.6), respectively. Hence, they are also referred to as a traveling back/front solution, a traveling pulse solution, and a periodic solution in (1.6). Our goal is to prove the existence of these solutions. We require that the solution (u, w) satisfies $u(z) > 0$ throughout the study because $u = \rho^{-1}$ and ρ must be positive. We note that c and K are unknown constants, which must be determined such that (1.6) has appropriate orbits. This task may be simplified by considering μ as a control parameter rather than addressing (K, c) directly. Specifically, we seek a particular value of μ such that the desired traveling wave solutions exist for a given pair (K, c) . The original problem can be then solved by determining (K, c) such that $\mu = 2\tau K - 1$. This study presents the first step toward addressing these issues. Our results identify all (K, c, μ) such that a homoclinic or periodic orbit exists, demonstrating the existence of particular triplets and the nonexistence of others.

We provide several definitions and notations to state the main results. Define $K_0 \equiv V'(0)$ and $K_M \equiv$ $V'(U_M) > K_0$. The assumption (A1) implies that the function $Ku - V(u)$ has a unique local maximum point $u_M = u_M(K) \in (0, U_M)$ for $K \in (K_0, K_M)$. Similarly, for $K \in (0, K_M)$, $Ku - V(u)$ has a unique local minimum point $u_m = u_m(K) \in (U_M, \infty)$. Set $c_0 \equiv -V(0)$ and define $c_M = c_M(K), c_m = c_m(K)$, and $c_1 = c_1(K)$ by

$$
c_M \equiv K u_M - V(u_M), \quad c_m \equiv K u_m - V(u_m), \quad c_1 \equiv \max\{c_0, c_m\}.
$$

It is elementary to show that some $K_1 \in (K_0, K_M)$ exists such that

$$
c_m(K) < c_0 \quad \text{if } 0 < K < K_1,
$$
\n
$$
c_m(K) = c_0 \quad \text{if } K = K_1,
$$
\n
$$
c_m(K) > c_0 \quad \text{if } K > K_1.
$$

After that, we define

$$
\mathcal{D}_1 \equiv \{ (K, c) \mid K_0 < K < K_M, \ c_1(K) < c < c_M(K) \},
$$
\n
$$
\mathcal{D}_2 \equiv \{ (K, c) \mid K_0 < K < K_1, \ c = c_0 \} \cup \{ (K, c) \mid 0 < K < K_1, \ c_m(K) < c < c_0 \}.
$$

If $(K, c) \in \mathcal{D}_1$, the function f has three zeros, denoted by u_0, u_1, u_2 with $0 < u_1 < u_0 < u_2$. Similarly, if $(K, c) \in \mathcal{D}_2$, f has two zeroes, u_0, u_2 , with $0 < u_0 < u_2$. More precise properties of f will be discussed in Lemma 2. Since we aim to find positive solutions in (1.6), we divide a region of (K, c) into \mathcal{D}_1 and \mathcal{D}_2 based on the number of zeroes of f for $u > 0$.

First, we study the existence of heteroclinic orbits in (1.6) that satisfy one of the following conditions:

$$
u(-\infty) = u_1 \quad u(\infty) = u_2, \quad w(\pm \infty) = 0, \quad w > 0,
$$
 (HE1)

$$
u(-\infty) = u_2, \quad u(\infty) = u_1, \quad w(\pm \infty) = 0, \quad w < 0.
$$
 (HE2)

Theorem 1. For $(K, c) \in \mathcal{D}_1$, there exists $\mu_b = \mu_b(K, c) \in \mathbb{R}$ (resp. $\mu_f = \mu_f(K, c) \in \mathbb{R}$) such that (1.6) with (HE1) (resp. (HE2)) has a solution if and only if $\mu = \mu_b$ (resp. $\mu = \mu_f$).

Subsequently, we state the existence and nonexistence of homoclinic orbits in (1.6), which satisfy

$$
(u(\pm\infty), w(\pm\infty)) = (\overline{u}, 0), \quad (u(z), w(z)) \neq (\overline{u}, 0),
$$
 (HO)

where \overline{u} is a positive number satisfying $f(\overline{u}) = 0$. The solution to (1.6) under condition (HO) is called a traveling pulse solution. As will be shown in Proposition 2, we can find $c_* = c_*(K)$ such that $\mu_* \equiv$ $\mu_b(K, c_*(K)) = \mu_f(K, c_*(K))$ under one of the following conditions:

$$
K \in (K_1, K_M) \quad \text{or} \quad (\text{H}) \text{ and } K \in (K_0, K_1]. \tag{C}
$$

This implies that (1.6) for $(c, \mu) = (c_*, \mu_*)$ has a *heteroclinic cycle* consisting of the equilibrium points $(u_1, 0)$ and $(u_2, 0)$ and two heteroclinic orbits that join them. After that, we divided \mathcal{D}_1 into

$$
\mathcal{D}_{1,1} \equiv \{ (K,c) \in \mathcal{D}_1 \mid c_*(K) < c < c_M(K) \},
$$
\n
$$
\mathcal{D}_{1,2} \equiv \{ (K,c) \in \mathcal{D}_1 \mid c_1(K) < c < c_*(K) \},
$$
\n
$$
\mathcal{D}_{1,3} \equiv \{ (K,c) \in \mathcal{D}_1 \mid c = c_*(K) \}.
$$

Theorem 2. The following statements hold:

- (i) If $(K, c) \notin \mathcal{D}_1 \cup \mathcal{D}_2$, then (1.6) has no solution satisfying (HO) for any \overline{u} and $\mu \in \mathbb{R}$.
- (ii) If $(K, c) \in \mathcal{D}_1 \cup \mathcal{D}_2$, then (1.6) has no solution satisfying (HO) with $\overline{u} = u_0$ for any $\mu \in \mathbb{R}$.
- (iii) Assume (C). If $(K, c) \in \mathcal{D}_{1,2} \cup \mathcal{D}_{1,3}$ (resp. $(K, c) \in \mathcal{D}_{1,1} \cup \mathcal{D}_{1,3}$), then (1.6) with (HO) has no solution for $\overline{u} = u_1$ (resp. $\overline{u} = u_2$) for any $\mu \in \mathbb{R}$.
- (iv) Assume (C). If $(K, c) \in \mathcal{D}_{1,1}$ (resp. $(K, c) \in \mathcal{D}_{1,2} \cup \mathcal{D}_2$), then there exists $\mu_{pul}^1 = \mu_{pul}^1(K, c)$ (resp. $\mu_{pul}^2 = \mu_{pul}^2(K, c)$ such that (1.6) with (HO) has a solution with $\overline{u} = u_1$ (resp. $\overline{u} = u_2$) if and only if $\mu = \mu_{pul}^1$ (resp. $\mu = \mu_{pul}^2$).

Finally, we discuss the existence or nonexistence of periodic orbits in (1.6) . Considering the Poincaré section $\{w = 0\}$, we obtain a periodic solution (u, w) to (1.6) satisfying the initial condition $(u(0), w(0)) =$ $(q, 0).$

Theorem 3. Assume (C) . Let q satisfy

$$
q \in \begin{cases} (u_1, u_0) & \text{if } (K, c) \in \mathcal{D}_{1,1} \cup \mathcal{D}_{1,3}, \\ (u_0, u_2) & \text{if } (K, c) \in \mathcal{D}_{1,2} \cup \mathcal{D}_2. \end{cases} \tag{1.7}
$$

Then there exists $\mu_{per} = \mu_{per}(K, c, q) > 0$ such that (1.6) has a periodic solution with the initial condition $(u(0), w(0)) = (q, 0)$ for $\mu = \mu_{per}$. Moreover, if $(K, c) \notin \mathcal{D}_1 \cup \mathcal{D}_2$ or $\mu \neq \mu_{per}$, there does not exist any periodic solution to (1.6).

As previously mentioned, (1.6) exhibits a heteroclinic cycle for $(K, c) \in \mathcal{D}_{1,3}$ under assumption (C). As discussed in [11], a global bifurcation can occur in systems with heteroclinic cycles, where a homoclinic orbit emerges from a heteroclinic cycle. Sections 7 and 8 demonstrate that such a bifurcation occurs in (1.6). In our case, a homoclinic orbit obtained in Theorem 2 (iv) bifurcates from the heteroclinic cycle. However, this study does not use the global bifurcation results established by [11].

All theorems are proven using phase-plane analysis. The key of the proofs is the monotonicity of solution trajectories with respect to μ (see Lemmas 6, 10 below). The shooting method facilitates us to directly study the behavior of the solution initiated from the equilibrium points. Such methods are widely used to demonstrate the existence of traveling wave solutions (see [12]).

The remainder of this paper is organized as follows. Section 2 discusses the investigation of the basic properties of (1.1) and (1.6) . Additionally, the properties of f are described. In Section 3, we consider the existence of the heteroclinic orbits in (1.6) and Theorem 1. Section 4 discusses the properties of μ_b and μ_f as given in Theorem 1 (see Proposition 2). Sections 5 and 6 address the existence of homoclinic and periodic orbits in (1.6) as stated in Theorems 2 and 3. Section 7 discusses the parameter dependence of heteroclinic, homoclinic, and periodic orbits. Additionally, we study the bifurcations that occur when these solutions appear. Section 8 presents a numerical investigation of the relationship among K, c, μ in the presence of traveling wave solutions, revealing the global bifurcation structure of (1.6) using AUTO [13]. One of our results, shown in Figure 4, is qualitatively equivalent to that in Figure 4 of [14], where the authors categorized the parameter space (K, c) based on the topological structure of the flow diagram and highlighted the existence of a heteroclinic cycle without using AUTO.

2 Preliminaries

2.1 Basic properties of (1.1)

To begin with, we consider the local existence and uniqueness of a solution in (1.1). If we consider (1.1) in the whole line $S \equiv \mathbb{R}$, we impose

$$
\lim_{x \to \pm \infty} (\rho(x, t), v(x, t)) = (\overline{\rho}_{\pm}, \overline{v}_{\pm}). \tag{2.1}
$$

If we are concerned with the bounded interval $S \equiv (0, L)$, we impose the periodic boundary conditions for ρ and v. Let $C^{k+\alpha}(S)$ for integer $k \geq 0$ and $0 < \alpha < 1$ be the set of kth-differentiable functions $\phi : S \to \mathbb{R}$ whose kth-derivatives are bounded and α -Hölder continuous [15]. We find a unique solution $(\rho, v) \in C^{1+\alpha}(S) \times C^{2+\alpha}(S)$ locally in time. The following proposition is standard so that we omit the details of the proof (see [16] and [17]).

Proposition 1. Give $(\rho_0, v_0) \in C^{1+\alpha}(S) \times C^{2+\alpha}(S)$ and assume that $T > 0$ is sufficiently small. Then the problem (1.1) with the initial condition $(\rho(z, 0), v(z, 0)) = (\rho_0(z), v_0(z))$ has a unique solution (ρ, v) in $C^{1+\alpha}(S) \times C^{2+\alpha}(S)$ in $(0,T)$.

As described in Introduction, congestion in the microscopic optimal velocity model proposed in [4] arises via Hopf bifurcation associated with the destabilization of the free flow [10]. This fact strongly implies the instability of a constant stationary solution (ρ_*, v_*) in (1.1), where $v_* = V(\rho_*^{-1})$. To examine the ρ_* -dependency of the stability, we set $(\rho, v) = (\rho_*, v_*) + (\phi, \psi)e^{\lambda t}e^{ikx}$ and study the linearized eigenvalue problem of (1.1), given by

$$
\begin{cases} \lambda \phi + ik(\rho_* \psi + v_* \phi) = 0, \\ \lambda \psi + ikv_* \psi = -\frac{ik}{2\tau \rho_*^3} V'(\rho_*^{-1}) \phi - \kappa(\rho_*) k^2 \psi - \frac{1}{\tau} \left(\frac{1}{\rho_*^2} V'(\rho_*^{-1}) \phi + \psi \right). \end{cases} (2.2)
$$

Then we obtain the characteristic equation

$$
(\lambda + ikv_*)\left(\lambda + ikv_* + \frac{1}{\tau} + \kappa(\rho_*)k^2\right) - \frac{ik}{\tau \rho_*}\left(\frac{ik}{2\rho_*} + 1\right)V'(\rho_*^{-1}) = 0.
$$

The eigenvalues $\lambda = \lambda_{\pm}(k)$ for each $k \in \mathbb{R}$ are explicitly given by

$$
\lambda_{\pm}(k) = -ikv_{*} + \frac{1}{2} \left(-\frac{1}{\tau} - \kappa(\rho_{*})k^{2} \pm \sqrt{z(k)} \right),
$$

$$
z(k) \equiv \left(\frac{1}{\tau} + \kappa(\rho_{*})k^{2} \right)^{2} + \frac{4ik}{\tau \rho_{*}} \left(\frac{ik}{2\rho_{*}} + 1 \right) V'(\rho_{*}^{-1}).
$$

Obviously, $Re\lambda_{-}(k) < 0$ for any ρ_{*} and k, where Re λ stands for the real part of complex number λ . To determine the stability of the stationary solution, it is sufficient to consider the sign of the real part of $\lambda_{+}(k)$ for each ρ_* and k.

Lemma 1. If $1 > 2\tau V'(\rho_*^{-1})$, then there are $k_1 < 0$ and $k_2 > 0$ such that $Re\lambda_+(k) > 0$ in $k_1 < k < k_2$ except for $k = 0$ and $Re\lambda_+(k) < 0$ in $k < k_1, k_2 < k$. On the other hand, if $1 \leq 2\tau V'(\rho_*^{-1})$, $Re\lambda_+(k) < 0$ in any $k \neq 0$.

Proof. Define $\lambda_r(k) = \text{Re}\lambda_+(k)$. It is clear that λ_r is smooth in $k \in \mathbb{R}$. It is easy to obtain $\lambda_r(0) = \lambda'_r(0) = 0$, and $\lambda''_r(0) = -V'(\rho_*^{-1})(1 - 2\tau V'(\rho_*^{-1}))/\rho_*^2$. Hence, $1 - 2\tau V'(\rho_*^{-1})$ determines the sign of $\lambda''_r(0)$ unless it is not equal to 0. If $1 = 2\tau V'(\rho_*^{-1})$, direct calculations imply $\lambda''_r(0) = \lambda_r^{(3)}(0) = 0$ and $\lambda_r^{(4)}(0) =$ $-24V'(\rho_*^{-1})\kappa(\rho_*)\tau/\rho_*^2 < 0$. Moreover, we easily calculate

$$
\lim_{k \to \pm \infty} \lambda_r(k) = -\frac{V'(\rho_*^{-1})}{2\tau \rho_*^2 \kappa(\rho_*)} < 0. \tag{2.3}
$$

Next we assume that there exist $k \neq 0$ and $a \in \mathbb{R}$ such that $\lambda_{+}(k) = i(a - kv_{*}),$ and calculate $\lambda'_{r}(k)$. Since

$$
\sqrt{z(k)} = 2ia + \frac{1}{\tau} + \kappa(\rho_*)k^2,
$$

we have

$$
a^{2} = \frac{k^{2}}{2\tau\rho_{*}^{2}} V'(\rho_{*}^{-1}), \quad a\left(\frac{1}{\tau} + \kappa(\rho_{*})k^{2}\right) = \frac{k}{\tau\rho_{*}} V'(\rho_{*}^{-1}),
$$

from which we have

$$
V'(\rho_*^{-1}) = \frac{\tau}{2} \left(\frac{1}{\tau} + \kappa(\rho_*) k^2 \right)^2.
$$

Then we find

$$
a = \frac{k}{2\rho_*} \left(\frac{1}{\tau} + \kappa(\rho_*) k^2 \right), \quad \sqrt{z(k)} = \left(\frac{ik}{\rho_*} + 1 \right) \left(\frac{1}{\tau} + \kappa(\rho_*) k^2 \right).
$$

Direct calculations give us

$$
z'(k) = 4\kappa(\rho_*)k\left(\frac{1}{\tau} + \kappa(\rho_*)k^2\right) + \frac{4i}{\tau \rho_*}\left(\frac{ik}{\rho_*} + 1\right)V'(\rho_*^{-1})
$$

= $\left(\frac{1}{\tau} + \kappa(\rho_*)k^2\right)\left(4\kappa(\rho_*)k - \frac{2k}{\rho_*^2}\left(\frac{1}{\tau} + \kappa(\rho_*)k^2\right)\right) + \frac{2i}{\rho_*}\left(\frac{1}{\tau} + \kappa(\rho_*)k^2\right)^2$

and

$$
\lambda'_{+}(k) + iv_{*} + \kappa(\rho_{*})k = \frac{z'(k)}{4\sqrt{z(k)}} = \frac{2\kappa(\rho_{*})k - \frac{k}{\rho_{*}^{2}}\left(\frac{1}{\tau} + \kappa(\rho_{*})k^{2}\right) + \frac{i}{\rho_{*}}\left(\frac{1}{\tau} + \kappa(\rho_{*})k^{2}\right)}{2\left(\frac{ik}{\rho_{*}} + 1\right)}.
$$

Picking up the real part of the above equality, we obtain

$$
\lambda'_{r}(k) = -\frac{k^{3}\kappa(\rho_{*})}{k^{2} + \rho_{*}^{2}},
$$
\n(2.4)

which shows that $\lambda'_r(k) < 0$ if $k > 0$, while $\lambda'_r(k) > 0$ if $k < 0$.

Combining the facts above, we show Lemma 1. Consider the case that $1 \leq 2\tau V'(\rho_*^{-1})$ and assume that there exists $k > 0$ such that $\lambda_r(k) > 0$. Then there must be some $\tilde{k} > 0$ such that $\lambda_r(\tilde{k}) = 0$ and $\lambda'_r(\tilde{k}) \ge 0$, which contradicts the sign of $\lambda'_r(k)$. On the other hand, if $1 > 2\tau V'(\rho_*^{-1})$, then $\lambda''_r(0) > 0$, (2.3) and (2.4) imply the statement of Lemma 1 by the same argument as above. \Box

2.2 Properties of f and flows near equilibria

We first summarize the sign and zeros of f .

Lemma 2. The following statements hold.

(i) If $(K, c) \in \mathcal{D}_1$, then f has three zeros $u_0 = u_0(K, c)$, $u_1 = u_1(K, c)$, $u_2 = u_2(K, c)$ with $0 < u_1 < u_0 <$ u_2 and satisfies

$$
\begin{cases}\nf(u) > 0 \quad \text{for} \quad u \in (u_1, u_0) \cup (u_2, \infty), \\
f(u) < 0 \quad \text{for} \quad u \in (0, u_1) \cup (u_0, u_2), \\
f'(u_1) > 0, \quad f'(u_2) > 0, \quad f'(u_0) < 0.\n\end{cases} \tag{2.5}
$$

(ii) If $(K, c) \in \mathcal{D}_2$, then f has two zeros $u_0 = u_0(K, c)$, $u_2 = u_2(K, c)$ with $0 < u_0 < u_2$ and satisfies

$$
\begin{cases}\nf(u) > 0 \quad \text{for} \quad u \in (0, u_0) \cup (u_2, \infty), \\
f(u) < 0 \quad \text{for} \quad u \in (u_0, u_2), \\
f'(u_2) > 0, \quad f'(u_0) < 0.\n\end{cases}
$$

Moreover,

$$
f(0) = 0, \ f'(0) > 0 \quad \text{if} \quad K_0 < K < K_1 \quad \text{and} \quad c = c_0,
$$
\n
$$
f(0) > 0 \quad \text{if} \quad 0 < K < K_1 \quad \text{and} \quad c_m < c < c_0.
$$

(iii) u_0 , u_1 and u_2 are of class C^1 and satisfy

$$
u_0(K, c), u_1(K, c) \to u_M(K) \quad as \quad c \to c_M(K), \tag{2.6}
$$

$$
u_0(K, c), u_2(K, c) \to u_m(K) \quad \text{as} \quad c \to c_m(K). \tag{2.7}
$$

We next study the flows of (1.6) near the equilibria $(u_0, 0)$, $(u_1, 0)$ and $(u_2, 0)$. Let $F(u, w)$ and $J(u)$ be the two-dimensional vector field associated with (1.6) and the Jacobian matrix of F at $(u, 0)$, respectively. They are explicitly given by

$$
F(u, w) \equiv \begin{pmatrix} w \\ g_1(u)f(u) + g_2(u)h(u, \mu)w \end{pmatrix}, \quad J(u) \equiv \begin{pmatrix} 0 & 1 \\ g_1(u)f'(u) & g_2(u)h(u, \mu) \end{pmatrix}.
$$
 (2.8)

The eigenvalues $\lambda_{\pm}(u)$ of $J(u)$ are explicitly given by

$$
\lambda_{\pm}(u) = \frac{1}{2} \left(g_2(u) h(u,\mu) \pm \sqrt{g_2(u)^2 h(u,\mu)^2 + 4g_1(u)f'(u)} \right).
$$

It is elementary to verify the following lemma by direct calculation.

Lemma 3. The following hold.

(i) For
$$
(K, c) \in \mathcal{D}_1
$$
 (resp. $(K, c) \in \mathcal{D}_1 \cup \mathcal{D}_2$), it holds that at $u = u_1$ (resp. $u = u_2$)
 $\lambda_+(u) > 0$, $\lambda_-(u) < 0$, $(\lambda_+)_{\mu}(u) > 0$, $(\lambda_-)_{\mu}(u) > 0$.

(ii) Assume $(K, c) \in \mathcal{D}_1 \cup \mathcal{D}_2$. If $\mu \neq -f'(u_0)$, then $Re\lambda_{\pm}(u_0)$ are either all positive or all negative, while if $\mu = -f'(u_0)$, then $\lambda_{\pm}(u_0) = \pm i\omega_0$, where $\omega_0 \equiv \sqrt{-g_1(u_0)f'(u_0)} > 0$.

2.3 Useful lemmas

We give some simple lemmas to be used throughout the subsequent sections.

Lemma 4. Let $A = A(s)$ and $B = B(s)$ be continuous integrable functions on a bounded interval (s_1, s_2) , and let $W \in C^1((s_1, s_2)) \cap C([s_1, s_2))$ be a solution of

$$
W' = A(s)W + B(s) \quad \text{in } (s_1, s_2).
$$

Then W can be extended continuously up to $s = s_2$. Moreover, if $B > 0$ in (s_1, s_2) and $W(s_1) \geq 0$, then $W > 0$ on $(s_1, s_2]$.

Proof. Solving the differential equation above, we have

$$
W(s) = \exp\left(\int_{s_1}^s A(\tau)d\tau\right)W(s_1) + \int_{s_1}^s \exp\left(\int_{\tau}^s A(\zeta)d\zeta\right)B(\tau)d\tau
$$

for $s \in [s_1, s_2)$. The lemma follows immediately from the above equality.

Lemma 5. Let A and B be continuous functions on a bounded interval $[s_1, s_2]$. Assume that $W \in$ $C^1((s_1, s_2)) \cap C([s_1, s_2])$ satisfies

$$
(W^2)' = A(s) + B(s)W \quad \text{in } (s_1, s_2).
$$

Then there hold

$$
e^{-s_2}W(s_2)^2 - e^{-s_1}W(s_1)^2 \le \int_{s_1}^{s_2} e^{-s} \left(A(s) + \frac{1}{4}B(s)^2\right)ds,
$$

$$
e^{s_2}W(s_2)^2 - e^{s_1}W(s_1)^2 \ge \int_{s_1}^{s_2} e^{s} \left(A(s) - \frac{1}{4}B(s)^2\right)ds.
$$

Proof. By the differential equation above and the inequality $|BW| \leq B^2/4 + W^2$, we have

$$
(W^2)' - W^2 \le A(s) + \frac{1}{4}B(s)^2
$$
, $(W^2)' + W^2 \ge A(s) - \frac{1}{4}B(s)^2$,

which concludes the lemma by Gronwall's inequality.

 \Box

 \Box

3 Existence of traveling back and front solutions

The goals of this section are to find monotone traveling back and front solutions and to prove Theorem 1. In order to obtain desired heteroclinic orbits, we consider stable and unstable manifolds emanating from $(u_1, 0)$ and $(u_2, 0)$. Let $(K, c) \in \mathcal{D}_1$. Lemma 3 shows that for each $j = 1, 2$, there is an orbit $\{(u_j^s(z), w_j^s(z)) \mid z \in \mathbb{R}\}\$ (resp. $\{(u_j^u(z), w_j^u(z)) \mid z \in \mathbb{R}\}\)$ which lies on the stable (resp. unstable) manifold of the equilibrium $(u_j, 0)$ and satisfies

$$
(u_1^{\rm s}(z),w_1^{\rm s}(z))\in S_-, \quad (u_2^{\rm s}(z),w_2^{\rm s}(z))\in S_+, \quad (u_1^{\rm u}(-z),w_1^{\rm u}(-z))\in S_+, \quad (u_2^{\rm u}(-z),w_2^{\rm u}(-z))\in S_-
$$

for large z, where $S_ = \{(u, w) | u_1 < u < u_2, w < 0\}$ and $S_ + \equiv \{(u, w) | u_1 < u < u_2, w > 0\}$. Hence we define z_j^s and z_j^u by

$$
z_1^s \equiv \inf\{z_0 \mid \{(u_1^s(z), w_1^s(z)) \mid z > z_0\} \subset S_{-}\} \in [-\infty, \infty),
$$

\n
$$
z_1^u \equiv \sup\{z_0 \mid \{(u_1^u(z), w_1^u(z)) \mid z < z_0\} \subset S_+\} \in (-\infty, \infty],
$$

\n
$$
z_2^s \equiv \inf\{z_0 \mid \{(u_2^s(z), w_2^s(z)) \mid z > z_0\} \subset S_+\} \in [-\infty, \infty),
$$

\n
$$
z_2^u \equiv \sup\{z_0 \mid \{(u_2^u(z), w_2^u(z)) \mid z < z_0\} \subset S_{-}\} \in (-\infty, \infty].
$$
\n(3.1)

It follows from Lemma 2 that

$$
(u_1^s(z_1^s), w_1^s(z_1^s)) \in \{(u, 0) \mid u_0 \le u \le u_2\} \cup \{(u_2, w) \mid w < 0\},(u_1^u(z_1^u), w_1^u(z_1^u)) \in \{(u, 0) \mid u_0 \le u \le u_2\} \cup \{(u_2, w) \mid w > 0\},(u_2^s(z_2^s), w_2^s(z_2^s)) \in \{(u, 0) \mid u_1 \le u \le u_0\} \cup \{(u_1, w) \mid w > 0\},(u_2^u(z_2^u), w_2^u(z_2^u)) \in \{(u, 0) \mid u_1 \le u \le u_0\} \cup \{(u_1, w) \mid w < 0\}.
$$
\n
$$
(3.2)
$$

As far as $u_z \neq 0$, we see from the inverse function theorem that each of orbits $\{(u_j^s(z), w_j^s(z)) \mid z > z_j^s\}$ and $\{(u_j^u(z), w_j^u(z)) \mid z < z_j^u\}$ is expressed as the graph of a function of u. More precisely, there are $u_j^{\pm} \in \mathbb{R}$ and functions $w_j^{\pm} = w_j^{\pm}(u)$ such that

$$
\mathcal{O}_1^+ \equiv \{ (u_1^u(z), w_1^u(z)) \mid z > z_1^u \} = \{ (u, w_1^+(u)) \mid u_1 < u < u_1^+ \},
$$
\n
$$
\mathcal{O}_1^- \equiv \{ (u_1^s(z), w_1^s(z)) \mid z < z_1^s \} = \{ (u, w_1^-(u)) \mid u_1 < u < u_1^- \},
$$
\n
$$
\mathcal{O}_2^+ \equiv \{ (u_2^s(z), w_2^s(z)) \mid z < z_2^s \} = \{ (u, w_2^+(u)) \mid u_2^+ < u < u_2 \},
$$
\n
$$
\mathcal{O}_2^- \equiv \{ (u_2^s(z), w_2^s(z)) \mid z > z_2^u \} = \{ (u, w_2^-(u)) \mid u_2^- < u < u_2 \}.
$$
\n
$$
(3.3)
$$

To emphasize the dependency of the parameters, we may write $u_j^{\pm} = u_j^{\pm}(K, c, \mu)$ and $w_j^{\pm}(u) = w_j^{\pm}(u; K, c, \mu)$. In the case of $u_j^{\pm} \neq u_1, u_0, u_2$, we see that u_j^{\pm} is C^1 with respect to (K, c, μ) by the implicit function theorem. By (3.2), we have $u_0 \leq u_1^{\pm} \leq u_2$ and $u_1 \leq u_2^{\pm} \leq u_0$.

We see from (1.6) that w_j^{\pm} is a solution of the equation

$$
w_u = \frac{g_1(u)f(u)}{w} + g_2(u)h(u,\mu),
$$
\n(3.4)

which is also written as

$$
\frac{1}{2}(w^2)_u = g_1(u)f(u) + g_2(u)h(u,\mu)w.
$$
\n(3.5)

By Lemma 3, we infer that w_j^{\pm} can be extended smoothly up to $u = u_j$ and

$$
w_1^{\pm} = 0, \quad (w_1^+)_{u} = \lambda_+(u_1), \quad (w_1^-)_{u} = \lambda_-(u_1) \quad \text{at } u = u_1,
$$

\n
$$
w_2^{\pm} = 0, \quad (w_2^+)_{u} = \lambda_-(u_2), \quad (w_2^-)_{u} = \lambda_+(u_2) \quad \text{at } u = u_2.
$$
\n(3.6)

We also infer that w_j^{\pm} can be extended continuously up to $u = u_j^{\pm}$ and

$$
w_1^{\pm} = 0 \text{ at } u = u_1^{\pm} \text{ if } u_0 \le u_1^{\pm} < u_2,
$$

\n
$$
w_2^{\pm} = 0 \text{ at } u = u_2^{\pm} \text{ if } u_1 < u_2^{\pm} \le u_0.
$$
\n
$$
(3.7)
$$

Moreover, w_j^{\pm} are continuously differentiable with respect to (K, μ, c) . Also, u_2^{\pm} and w_2^{\pm} are defined for $(K, c) \in \mathcal{D}_2$ by setting $u_1 = 0$.

We show the monotonicity of w_j^{\pm} with respect to μ .

Lemma 6. For $(K, c) \in \mathcal{D}_1$, there holds $(w_1^{\pm})_{\mu} > 0$ in $u \in (u_1, u_1^{\pm})$. Similarly, for $(K, c) \in \mathcal{D}_1 \cup \mathcal{D}_2$, there $holds (w_2^{\pm})_{\mu} < 0 \in (u_2^{\pm}, u_2).$

Proof. We put $W \equiv (w_1^+)_\mu$. By (3.6) and Lemma 3, we have $W = 0$, $W_u > 0$ at $u = u_1$. Hence we can pick a point $\hat{u} \in (u_1, u_1^+)$ such that $W > 0$ on $(u_1, \hat{u}]$. Differentiating (3.4) with respect to μ , we see that W satisfies

$$
W_u = -\frac{g_1(u)f(u)}{(w_1^+)^2}W + g_2(u)
$$

in (u_1, u_1^+) . By applying Lemma 4, $W > 0$ in (\hat{u}, u_1^+) . Thus we conclude that $(w_1^+)_\mu > 0$ in (u_1, u_1^+) . By a similar argument, we obtain $(w_1^-)_{\mu} > 0$ and $(w_2^{\pm})_{\mu} < 0$. Therefore the lemma follows.

Lemma 7. Let $(K, c) \in \mathcal{D}_1$ and $a \in (u_1, u_2)$. If μ (resp. $-\mu$) is large enough, then $u_1^+ > a$ and $u_2^- < a$ (resp. $u_1^- > a$ and $u_2^+ < a$). Moreover, there hold

$$
w_{1}^{\pm}(a; K, c, \mu) \to \pm \infty \quad as \quad \mu \to \pm \infty,
$$

\n
$$
w_{2}^{\pm}(a; K, c, \mu) \to \pm \infty \quad as \quad \mu \to \mp \infty.
$$
\n(3.8)

Proof. To estimate w_1^+ , we first integrate (3.4). Then, by Lemma 2, we have

$$
w_1^+(u) \ge \int_{u_1}^u g_2(s)h(s,\mu)ds
$$

for $u \in [u_1, u_0]$. Since the right-hand side goes to ∞ as $\mu \to \infty$, we find

$$
w_1^+(u;K,c,\mu) \to \infty \tag{3.9}
$$

locally uniformly for $u \in (u_1, u_0]$ as $\mu \to \infty$.

Next we integrate (3.5). We then have

$$
w_1^+(u)^2 = w_1^+(u_0)^2 + 2 \int_{u_0}^u g_1(s)f(s)ds + 2 \int_{u_0}^u g_2(s)h(s,\mu)w_1^+(s)ds.
$$

It follows from (3.9) that the right-hand side goes to ∞ uniformly for $u \in [u_0, u_1^+]$ as $\mu \to \infty$. From this and (3.7), we conclude that $u_1^+ = u_2$ for large μ and $w_1^+(u; K, c, \mu) \to \infty$ uniformly for $u \in [u_0, u_2]$ as $\mu \to \infty$. Thus the assertion for w_1^+ is proved. The others can be shown in a similar way.

We are now in a position to prove Theorem 1.

Proof of Theorem 1. We fix $a \in (u_0, u_2)$ and consider the behavior of $\phi(\mu) \equiv (w_1^+ - w_2^+)|_{u=a}$. From Lemma 6, we deduce that u_1^+ is nondecreasing with respect to μ . Since $u_1^+ > a$ for large μ from Lemma 7, there is some $b \in [-\infty, \infty)$ such that $\{\mu \in \mathbb{R} \mid u_1^+ > a\} = (b, \infty)$. Using Lemmas 6 and 7 again, we infer that $\phi_{\mu} > 0$, $\lim_{\mu\to\infty}\phi(\mu)=\infty$, and

$$
\lim_{\mu \to b} \phi(\mu) = \begin{cases} -w_2^+(a; K, c, b) < 0 & \text{if } b \neq -\infty, \\ -\infty & \text{if } b = -\infty \end{cases}
$$

because $u_1^+ = a$ and $w_1^+(a) = 0$ for $\mu = b$ when $b \neq -\infty$. Therefore there exists a unique zero μ_b of ϕ . We have thus proved the assertion for (1.6) with (HE1). Since the same argument is valid for (1.6) with (HE2), we conclude Theorem 1. \Box

Remark 1. By the implicit function theorem, μ_b and μ_f are of class C^1 with respect to $(K, c) \in \mathcal{D}_1$.

4 Behavior of μ_b and μ_f

This section is devoted to the proof of the following proposition. We examine the behaviors of μ_b and μ_f as c runs from c_1 to c_M and prove the existence of a heteroclinic cycle in (1.6).

Proposition 2. Assume (C). For $K \in (K_0, K_M)$, there is a unique number $c_* = c_*(K) \in (c_1, c_M)$ such that

$$
\mu_b < \mu_f \quad \text{if} \quad c_1 < c < c_*, \\
\mu_b > \mu_f \quad \text{if} \quad c_* < c < c_M, \\
\mu_b = \mu_f \quad \text{if} \quad c = c_*.\n\tag{4.1}
$$

We begin by showing the monotonicity of w_j^{\pm} with respect to c.

Lemma 8. For $(K, c) \in \mathcal{D}_1$ and $\mu \in \mathbb{R}$, there hold $\mp (w_1^{\pm})_c > 0$ in $u \in (u_1, u_1^{\pm})$ and $\pm (w_2^{\pm})_c > 0$ in $u\in(u_2^{\pm},u_2).$

Proof. We only estimate $W \equiv -(w_1^+)_c$. The other inequalities can be obtained in the same way. Differentiating the equalities $w_1^+(u_1) = 0$ and $f(u_1) = 0$ with respect to c gives $W(u_1) = \lambda_+(u_1)/(Kf'(u_1))$, where $\lambda_+(u_1)$ was given in Lemma 3. Since the right-hand side of this equality is positive, we infer that $W > 0$ in a neighborhood of u_1 . Differentiating (3.4) yields

$$
W_u = -\frac{g_1(u)f(u)}{(w_1^+)^2}W + \frac{g_1(u)}{Kw_1^+}.
$$

Applying Lemma 4, we conclude that $W > 0$ in (u_1, u_1^+) . This completes the proof.

Next we show the monotonicity of μ_b and μ_f with respect to $c \in (c_1, c_M)$.

Lemma 9. The inequalities $(\mu_b)_c > 0$ and $(\mu_f)_c < 0$ hold.

Proof. We recall that the equality $w_1^+(u; K, c, \mu_b) = w_2^+(u; K, c, \mu_b)$ holds. Differentiating this with respect to c, we have

$$
(w_1^+)_c + (w_1^+)_\mu (\mu_b)_c = (w_2^+)_c + (w_2^+)_\mu (\mu_b)_c.
$$

This with Lemmas 6 and 8 yields

$$
(\mu_b)_c = \frac{(w_2^+)_c - (w_1^+)_c}{(w_1^+)_\mu - (w_2^+)_\mu} > 0.
$$

In a similar way, we have

$$
(\mu_f)_c = -\frac{(w_1^-)_c - (w_2^-)_c}{(w_1^-)_\mu - (w_2^-)_\mu} < 0.
$$

Therefore the lemma follows.

Let us prove Proposition 2.

Proof of Proposition 2. For simplicity of notation, we ignore the dependence on K and write $\mu_b(c)$ instead of $\mu_b(K, c)$, for instance. By Lemma 9, it suffices to show that

$$
\lim_{c \to c_M} \mu_b(c) > \lim_{c \to c_M} \mu_f(c), \quad \lim_{c \to c_1} \mu_b(c) < \lim_{c \to c_1} \mu_f(c). \tag{4.2}
$$

To derive the former inequality of (4.2), we prove that

$$
\lim_{c \to c_M} \mu_b(c) > \mu^*,\tag{4.3}
$$

 \Box

 \Box

where μ^* is determined by the relation

$$
\int_{u_M}^{u_2^*} g_2(u)h(u, \mu^*) du = 0 \tag{4.4}
$$

for $u_2^* \equiv \lim_{c \to c_M} u_2(c) > 0$. To obtain a contradiction, suppose that (4.3) is false. Then Lemma 9 yields $\mu_b(c) < \mu^*$ for all $c \in (c_1, c_M)$. Set $w_b(u) \equiv w_1^+(u; c, \mu_b(c)) = w_2^+(u; c, \mu_b(c))$. Since w_b satisfies (3.5) for $\mu = \mu_b$, we can apply Lemma 5 with $W = w_b$, $A = 2g_1f - 2(\mu^* - \mu_b)g_2w_b$, $B = 2g_2h(\cdot, \mu^*)$, $s_1 = u_1$ and $s_2 = u \in [u_1, u_2]$ to obtain

$$
w_b(u)^2 \le \int_{u_1}^u e^{u-s} (2g_1(s)f(s) + g_2(s)^2 h(s, \mu^*)^2) ds - 2(\mu^* - \mu_b) \int_{u_1}^u e^{u-s} g_2(s) w_b(s) ds
$$

$$
\le \int_{u_1}^u e^{u-s} (2g_1(s)f(s) + g_2(s)^2 h(s, \mu^*)^2) ds.
$$
 (4.5)

If $u = u_0$, we find from (2.6) that the right-hand side approaches 0 as $c \to c_M$, which implies that $\lim_{c \to c_M} w_b(u_0(c); c, \mu_b(c)) = 0.$ Moreover, it follows from (4.5) that

$$
\limsup_{c \to c_M} \max_{u \in [u_1, u_2]} w_b(u) < \infty. \tag{4.6}
$$

We now integrate (3.4) over $[u_0, u_2]$ to obtain

$$
w_b(u_0) + \int_{u_0}^{u_2} g_2(u)h(u, \mu_b) du = \int_{u_0}^{u_2} \frac{-g_1(u)f(u)}{w_b(u)} du.
$$

By (4.4) and the assumption $\lim_{c\to c_M} \mu_b(c) \leq \mu^*$, we see that the left-hand side converges to some nonpositive number as $c \to c_M$. On the other hand, (4.6) implies that the right-hand side is bounded below by some positive constant for any c close to c_M . This is a contradiction, and therefore (4.3) holds. The inequality $\lim_{c \to c_M} \mu_f(c) < \mu^*$ can also be verified in a similar way. We have thus shown the former inequality of (4.2).

Let us prove the latter inequality of (4.2). In the case $K_1 < K < K_M$, we have $c_1 = c_m$ and put $u_1^* \equiv \lim_{c \to c_m} u_1(c) > 0$. Hence we can apply an argument similar to the proof of (4.3) to obtain

$$
\lim_{c \to c_m} \mu_b(c) < \bar{\mu}^* < \lim_{c \to c_m} \mu_f(c),
$$

where $\bar{\mu}^*$ is determined by the relation

$$
\int_{u_1^*}^{u_m} g_2(u)h(u,\bar{\mu}^*)du = 0.
$$

Next we assume (H) and $K_0 < K \leq K_1$. In this case, we have $c_1 = c_0$. We prove that

$$
\lim_{c \to c_0} \mu_b(c) \le \mu_0 < \lim_{c \to c_0} \mu_f(c),\tag{4.7}
$$

where $\mu_0 \equiv -f'(0)$. Note that μ_0 is given by $h(0, \mu_0) = 0$. Moreover, one can easily check that

$$
\lim_{c \to c_0} u_1(c) = 0, \quad \lim_{c \to c_0} u_2(c) > 0, \quad f'(u) > 0 \text{ for } u \in [0, u_M).
$$

We prove the first inequality of (4.7) by contradiction. Obviously, there is a positive constant δ_0 such that $h(u, \mu_b(c)) > 0$ for all $u \in [0, \delta_0]$ and $c \in (c_0, c_M)$. By integrating (3.5) over $[u_1, \delta_0]$, we deduce that

$$
\frac{1}{2}w_b(\delta_0)^2 \ge \int_{u_1}^{\delta_0} g_1(u)f(u)du.
$$
\n(4.8)

To estimate the left-hand side, we apply Lemma 5 with $W = w_b$, $A = 2g_1f - 2(\mu_0 - \mu_b)g_2w_b$, $B = 2g_2h(\cdot, \mu_0)$, $s_1 = \delta_0$ and $s_2 = u_2$. The result is

$$
w_b(\delta_0)^2 \leq \int_{\delta_0}^{u_2} e^{s-\delta_0} (-2g_1(u)f(u) + g_2(u)^2 h(u, \mu_0)^2) du + 2(\mu_0 - \mu_b) \int_{\delta_0}^{u_2} e^{s-\delta_0} g_2(u) w_b(u) du,
$$

which implies that $w_b(\delta_0)^2$ is bounded by some constant independent of c. On the other hand, the right-hand side of (4.8) is estimated as

$$
\int_{u_1}^{\delta_0} g_1(u) f(u) du \ge m_0 \int_{u_1}^{\delta_0} (u - u_1) g_1(u) du \ge \frac{m_0}{2} \int_{2u_1}^{\delta_0} u g_1(u) du,
$$

where we have applied $f(u) \geq m_0(u - u_1)$ in all $u \in [u_1, \delta_1]$ and $c \in [c_0, c_0 + \delta_2]$ for some positive constants m_0, δ_1, δ_2 . By the assumption (H), we conclude that the right-hand side of (4.8) diverges to ∞ as $c \to c_0$. This leads to a contradiction.

A similar argument works for the second inequality of (4.7). Thus we obtain the latter inequality of (4.2), and the proof is complete. П

5 Structure of traveling pulse solutions

Let us consider traveling pulse solutions of (1.6) with (HO) . For simplicity of notation, we let

$$
\tilde{u}_1 = \begin{cases} u_1 & \text{if } (K, c) \in \mathcal{D}_1, \\ 0 & \text{if } (K, c) \in \mathcal{D}_2. \end{cases}
$$
\n
$$
(5.1)
$$

To prove Theorem 2, we examine the properties of u_j^{\pm} . We note that the derivatives $(u_1^{\pm})_\mu$ (resp. $(u_2^{\pm})_\mu$) exist if $u_1^{\pm} \in (u_0, u_2)$ (resp. $u_2^{\pm} \in (\tilde{u}_1, u_0)$), thanks to the smooth dependence of stable and unstable manifolds of the equilibria $(u_1, 0)$ and $(u_2, 0)$ on parameters.

Lemma 10. It holds that $\pm (u_1^{\pm})_{\mu} > 0$ if $u_1^{\pm} \in (u_0, u_2)$ and $\pm (u_2^{\pm})_{\mu} > 0$ if $u_2^{\pm} \in (\tilde{u}_1, u_0)$.

Proof. We only prove the assertion for u_1^+ . The others can be handled in a similar way. For abbreviation, we write w instead of w_1^+ .

Assume $u_1^{\dagger} \in (u_0, u_2)$ and set $\zeta = \zeta(u) \equiv (w^2)_{\mu}$. We first show that ζ is continuously extended up to the point $u = u_1^+$ and $\zeta(u_1^+) > 0$. It is easy to see from Lemma 6 that ζ is positive if u is bigger than and close to u_1 . By differentiating (3.5) with respect to μ , we see that ζ satisfies

$$
\zeta_u = \frac{g_2(u)h(u,\mu)}{w}\zeta + 2g_2(u)w.
$$

Let us check that $g_2h(\cdot,\mu)/w$ is locally integrable on $(u_1, u_1^+]$. From (3.5) and (3.7), we have $\lim_{u\to u_1^+}ww_u$ $g_1(u_1^+)f(u_1^+)$. Since $f(u_1^+) < 0$, there is a constant $C > 0$ such that $w(u) \ge C\sqrt{u_1^+ - u}$ for u close to u_1^+ , which implies the local integrability of $g_2h(\cdot,\mu)/w$. Then it follows from Lemma 4 that ζ is continuously extended up to u_1^+ and positive in (u_1, u_1^+) .

Recall that u_1^{\dagger} is determined implicitly by $w(u_1^{\dagger})^2 = 0$. Differentiating this equality with respect μ and then using (3.5), we see that

$$
2g_1(u_1^+)f(u_1^+)(u_1^+)_\mu + \zeta(u_1^+) = 0.
$$

Therefore we obtain

$$
(u_1^+)_{\mu} = -\frac{\zeta(u_1^+)}{2g_1(u_1^+)f(u_1^+)} > 0,
$$

which completes the proof.

 \Box

Lemma 11. Define

$$
\mu_1^+ \equiv \inf \{ \mu \in \mathbb{R} \mid u_1^+ > u_0 \}, \quad \mu_1^- \equiv \sup \{ \mu \in \mathbb{R} \mid u_1^- > u_0 \}
$$

for $(K, c) \in \mathcal{D}_1$ and

$$
\mu_2^+ \equiv \sup \{ \mu \in \mathbb{R} \mid u_2^+ < u_0 \}, \quad \mu_2^- \equiv \inf \{ \mu \in \mathbb{R} \mid u_2^- < u_0 \}
$$

for $(K, c) \in \mathcal{D}_1 \cup \mathcal{D}_2$. Then $\mu_1^+, \mu_2^- \in [-\infty, \infty)$ and $\mu_1^-, \mu_2^+ \in (-\infty, \infty]$. Moreover, one has

$$
\mu_1^+ < \mu_1^-, \quad \mu_2^+ > \mu_2^-. \tag{5.2}
$$

Proof. Assume $(K, c) \in \mathcal{D}_1$. The first statement follows immediately from Lemma 7. We prove (5.2). On the contrary, suppose that $\mu_1^+ \geq \mu_1^-$. From Lemma 10, we see that u_1^+ and u_1^- are nondecreasing and nonincreasing with respect to μ , respectively. Hence $u_1^+ = u_1^- = u_0$ for all $\mu \in [\mu_1^-, \mu_1^+]$. By (3.4), we have

$$
(w_1^+ - w_1^-)_u = g_1(u) f(u) \left(\frac{1}{w_1^+} - \frac{1}{w_1^-}\right).
$$

Since the right-hand side is positive in (u_1, u_0) , we deduce that

$$
w_1^+(u_0) - w_1^-(u_0) > w_1^+(u_1) - w_1^-(u_1).
$$

This leads to a contradiction because $w_1^+(u_0) = w_1^-(u_0) = 0$ and $w_1^+(u_1) = w_1^-(u_1) = 0$. Therefore the former inequality of (5.2) holds. The latter inequality can be shown in a similar way.

Next we assume $(K, c) \in \mathcal{D}_2$. By the same argument as in the proof of Lemma 10, it follows that $u_2^- < a$ (resp. $u_2^+ < a$) if μ (resp. $-\mu$) is large enough. Then we easily verify (5.2) in the same manner as above.

We give sufficient conditions on μ for which u_2^{\pm} is positive in the case of $(K, c) \in \mathcal{D}_2$. We use the notation $\mu_0 = -f'(0)$, which has already been defined in the proof of Proposition 2.

Lemma 12. Let $(K, c) \in \mathcal{D}_2$ and assume (H). Then $u_2^+ > 0$ if $\mu \ge \mu_0$, while $u_2^- > 0$ if $\mu \le \mu_0$.

Proof. Let $\mu \geq \mu_0$. Then there is $\delta_0 > 0$ such that $h(u, \mu) \geq 0$ for $u \in [0, \delta_0]$. To obtain a contradiction, we assume that $w_2^+(u)$ is positive in $(0, u_2)$. For any $u \in (0, \delta_0)$, we integrate (3.5) over $[u, \delta_0]$ and then have

$$
w_2^+(\delta_0)^2 - w_2^+(u)^2 \ge 2 \int_u^{\delta_0} g_1(s) f(s) ds.
$$
\n(5.3)

□

From (ii) of Lemma 2, we can choose a constant $C > 0$ such that $f(u) \geq Cu$ for all u close to 0. Then it follows that the right-hand side of (5.3) diverges to ∞ as $u \to 0$ by the assumption (H). On the other hand, the left-hand side of (5.3) is bounded above as $u \to 0$, which is a contradiction. Therefore w_2^+ must vanish at some point in $(0, u_2)$, which implies $u_2^+ > 0$.

In a similar way, we can also show that $u_2^- > 0$ if $\mu \leq \mu_0$. Therefore the lemma follows.

Before proceeding to the proof of Theorem 2, we first give necessary conditions for the existence of solutions to the problem (1.6) with (HO). We will apply the following arguments not only for homoclinic orbits but also for periodic orbits discussed in the next section.

Lemma 13. If $(K, c) \notin \mathcal{D}_1 \cup \mathcal{D}_2$, then there is no solution (u, w) of (1.6) satisfying (HO) and $u(z) > 0$ in $-\infty < z < \infty$.

Proof. From the assumption, $(K, c) \in \mathcal{D}_3 \cup \mathcal{D}_4$, where

$$
\mathcal{D}_3 \equiv \{(K, c) \mid K < 0, \ c \ge c_0\} \cup \{(K, c) \mid 0 < K \le K_1, \ c \le c_m(K)\} \cup \{(K, c) \mid K_1 < K, \ c \le c_0\},
$$
\n
$$
\mathcal{D}_4 \equiv \{(K, c) \mid K < 0, \ c < c_0\} \cup \{(K, c) \mid 0 < K \le K_0, \ c \ge c_0\} \cup \{(K, c) \mid K_0 < K < K_M, \ c \ge c_M(K)\}
$$
\n
$$
\cup \{(K, c) \mid K_1 < K < K_M, \ c_0 < c \le c_m(K)\} \cup \{(K, c) \mid K \ge K_M, \ c > c_0\}.
$$

We easily see that $f(u) \geq 0$ in $u > 0$ if $(K, c) \in \mathcal{D}_3$. On the other hand, if $(K, c) \in \mathcal{D}_4$, then there is $u_* > 0$ such that $f(u) \leq 0$ in $0 \leq u \leq u_*$, while $f(u) \geq 0$ in $u \geq u_*$.

To obtain a contradiction, suppose that (1.6) has a solution (u, w) satisfying (HO) . We first consider the case $(K, c) \in \mathcal{D}_3$. Let H be a primitive of $g_2h(\cdot, \mu)$. Then we have

$$
(w - H(u))_z = w_z - g_2(u)h(u, \mu)u_z = g_1(u)f(u).
$$

Integrating this over $(-\infty, \infty)$ and using (HO), we deduce that

$$
0 = \int_{-\infty}^{\infty} g_1(u(z)) f(u(z)) dz.
$$
\n(5.4)

Then we obtain $f(u(z)) \equiv 0$, which contradicts the condition $(u(z), w(z)) \not\equiv (\overline{u}, 0)$.

Next we assume $(K, c) \in \mathcal{D}_4$. From (HO), we see that u has either a global maximum or a global minimum. Suppose that u has a global maximum at some z_0 . Then we have

$$
w(z_0) = u_z(z_0) = 0
$$
, $w_z(z_0) = u_{zz}(z_0) \le 0$.

Substituting these into the second equality of (1.6) yields $q_1(u(z_0))f(u(z_0)) \leq 0$. Since (u, w) is not an equilibrium, we see $f(u(z_0)) < 0$ and then $u(z) \le u(z_0) \le u_*$. However, it follows from the equality (5.4) that $f(u(z)) \equiv 0$, which is a contradiction. The other case can be derived in the same way as above. Thus the proof is complete. the proof is complete.

We are now in a position to show Theorem 2.

Proof of Theorem 2. The assertion (i) is a direct consequence of Lemma 13. We begin with the proof of (ii). On the contrary, suppose that there exists a solution (u, w) of (1.6) satisfying (HO) with $\overline{u} = u_0$. Then we must have $\mu = -f'(u_0)$ because (ii) of Lemma 3 shows that any solution of (1.6) cannot converge to $(u_0, 0)$ as either $z \to \infty$ or $z \to -\infty$ if $\mu \neq -f'(u_0)$. Let $z_* > 0$ be sufficiently large. Clearly, (u, w) is close to $(u_0, 0)$ in $|z| \geq z_*$. More precisely, there are z_0 and $r = r(z) > 0$ such that $r \to 0$ as $|z| \to \infty$ and (u, w) is approximated by

$$
(u_0,0) + r(\cos\omega_0(z-z_0), -\omega_0\sin\omega_0(z-z_0))
$$

in $|z| \geq z_*$ from (ii) of Lemma 3. Then the orbit of (u, w) must intersect with itself, which leads to the contradiction because of the uniqueness of a solution in ordinary differential equations.

Let us turn to the proofs of (iii) and (iv). First we consider the case $(K, c) \in \mathcal{D}_1$ and $\overline{u} = u_1$. It is sufficient to check the condition

$$
u_1^+ = u_1^- \in (u_0, u_2). \tag{5.5}
$$

Recall that

 $u_1^+ = u_2$ and $w_1^+(u_2) = 0$ if and only if $\mu = \mu_b$, $u_1^- = u_2$ and $w_1^-(u_2) = 0$ if and only if $\mu = \mu_f$.

These with Lemma 10 imply that

$$
u_1^+ \begin{cases} \in (u_0, u_2) & \text{if } \mu_1^+ < \mu < \mu_b, \\ = u_2 & \text{if } \mu \ge \mu_b, \end{cases}
$$
\n
$$
u_1^- \begin{cases} = u_2 & \text{if } \mu \le \mu_f, \\ \in (u_0, u_2) & \text{if } \mu_f < \mu < \mu_1^-.\end{cases}
$$
\n(5.6)

We also recall that

$$
\mu_b \le \mu_f \quad \text{if} \quad (K, c) \in \mathcal{D}_{1,2} \cup \mathcal{D}_{1,3},
$$
\n
$$
\mu_b > \mu_f \quad \text{if} \quad (K, c) \in \mathcal{D}_{1,1}.
$$
\n
$$
(5.7)
$$

Combining (5.2), (5.6), and (5.7), we conclude that (5.5) is never satisfied if $(K, c) \in \mathcal{D}_{1,2} \cup \mathcal{D}_{1,3}$, while (5.5) holds for some $\mu \in (\mu_f, \mu_b)$ if $(K, c) \in \mathcal{D}_{1,1}$. The uniqueness of μ satisfying (5.5) follows from Lemma 10. Therefore the assertion is proved in this case.

Next, we examine the case $\overline{u} = u_2$. By the same argument applied above, we can show the unique existence of $\mu_{pul}^2 \in (\mu_b, \mu_f)$ for $(K, c) \in \mathcal{D}_{1,2}$ and the nonexistence of solutions for $(K, c) \in \mathcal{D}_{1,1} \cup \mathcal{D}_{1,3}$. Hence we only need to consider the case of $(K, c) \in \mathcal{D}_2$ under (H). Put

$$
\underline{\mu}\equiv \inf\{\mu\in\mathbb{R}~|~u_2^+>0\},\quad \overline{\mu}\equiv \sup\{\mu\in\mathbb{R}~|~u_2^->0\}.
$$

From Lemmas 11 and 12, we see $\underline{\mu} < \mu_2^+$ and $\mu_2^- < \overline{\mu}$. We then have

$$
u_2^+ \in (0, u_0) \text{ if } \underline{\mu} < \mu < \mu_2^+, \quad u_2^- \in (0, u_0) \text{ if } \mu_2^- < \mu < \overline{\mu} \tag{5.8}
$$

by Lemma 10. Using Lemma 12 and the fact that $\{\mu \in \mathbb{R} \mid u_2^->0\}$ is open, we see that

$$
\underline{\mu} < \mu_0 < \overline{\mu}.\tag{5.9}
$$

Combining (5.2), (5.8) and (5.9), we obtain μ_{pul}^2 such that $u_2^+ = u_2^- \in (0, u_0)$ for $\mu = \mu_{pul}^2$. Thus the proof is complete. \Box

We conclude this section by deriving an estimate of μ_{pul}^j to be used in Section 7.

Lemma 14. For $\mu = \mu_{pul}^j$ $(j = 1, 2)$, let (u, w) be the traveling pulse solution obtained in (iv) of Theorem 2. Then

$$
-\sup_{u\in(\underline{m},\overline{m})}f'(u)<\mu_{pul}^j<-\inf_{u\in(\underline{m},\overline{m})}f'(u),
$$

where

$$
\underline{m} \equiv \inf_{z \in \mathbb{R}} u(z), \quad \overline{m} \equiv \sup_{z \in \mathbb{R}} u(z).
$$

Proof. Let $D \subset \mathbb{R}^2$ be the region enclosed by the closed curve $\{(u(z), w(z))\}_{z \in \mathbb{R}} \cup \{(u_j, 0)\}\)$ for $j = 1, 2$, and let $F(u, w)$ be the two-dimensional vector field defined in (2.8). It is well-known that

$$
\int_D \nabla \cdot F(u, w) du dw = 0,
$$

which follows easily from the divergence theorem and (1.6). Since $\nabla \cdot F(u, w) = g_2(u)h(u, \mu_{pul}^j)$, $h(\cdot, \mu_{pul}^j)$ must change its sign in (m, \overline{m}) . Therefore we obtain

$$
\inf_{u \in (\underline{m}, \overline{m})} h(u, \mu_{pul}^j) < 0 < \sup_{u \in (\underline{m}, \overline{m})} h(u, \mu_{pul}^j),
$$

which concludes the lemma.

6 Structure of periodic traveling wave solutions

In this section, we study periodic traveling wave solutions. The proof of Theorem 3 is similar to that of Theorem 2. Let (u, w) be a solution of (1.6) with the initial condition $(u(0), w(0)) = (q, 0)$. First, we assume that $(K, c) \in \mathcal{D}_1$ and $q \in (u_1, u_0)$. It is then seen from Lemma 2 that $(u(z), w(z)) \in S_+$ and $(u(-z), w(-z)) \in$ S− for small $z > 0$, where $S_-\equiv \{(u, w) \mid q < u < u_2, w < 0\}$ and $S_+\equiv \{(u, w) \mid q < u < u_2, w > 0\}$. Hence we define z^{\pm} by

$$
z^{+} \equiv \sup\{z_{0} \mid \{(u(z), w(z))\}_{0 < z < z_{0}} \subset S_{+}\} \in (0, \infty],
$$

\n
$$
z^{-} \equiv \inf\{z_{0} \mid \{(u(z), w(z))\}_{z_{0} < z < 0} \subset S_{-}\} \in [-\infty, 0).
$$
\n(6.1)

 \Box

Lemma 2 leads to

$$
(u(z^+), w(z^+)) \in \{(u, 0) \mid u_0 \le u \le u_2\} \cup \{(u_2, w) \mid w > 0\},\
$$

$$
(u(z^-), w(z^-)) \in \{(u, 0) \mid u_0 \le u \le u_2\} \cup \{(u_2, w) \mid w < 0\}.
$$

Since $u_z \neq 0$ in $z \in (0, z^+) \cup (z^-, 0)$, there are $u^{\pm} \in [u_0, u_2]$ and functions $w^{\pm} = w^{\pm}(u)$ such that

$$
\{(u(z), w(z)) \mid 0 < z < z^+\} = \{(u, w^+(u)) \mid q < u < u^+\},\
$$

$$
\{(u(z), w(z)) \mid z^- < z < 0\} = \{(u, w^-(u)) \mid q < u < u^-\}.
$$

To emphasize the dependency of the parameters, we may write $u^{\pm} = u^{\pm}(K, c, \mu)$ and $w^{\pm} = w^{\pm}(u; K, c, \mu)$. By definition, we see that w^{\pm} satisfy (3.4), (3.5) and $\lim_{u \to q} w^{\pm}(u) = 0$. Moreover, w^{\pm} are continuously

differentiable with respect to (K, c, μ) .

One can similarly define z^{\pm} , u^{\pm} and w^{\pm} for $(K, c) \in \mathcal{D}_1 \cup \mathcal{D}_2$ and $q \in (u_0, u_2)$. In this case, we have $\pm z^{\pm} < 0, \, \pm w^{\pm} > 0, \, u^{\pm} \in [\tilde{u}_1, u_0], \text{ and}$

$$
\{(u(z), w(z)) \mid z^+ < z < 0\} = \{(u, w^+(u)) \mid u^+ < u < q\},
$$

$$
\{(u(z), w(z)) \mid 0 < z < z^-\} = \{(u, w^-(u)) \mid u^- < u < q\},
$$

where \tilde{u}_1 was given in (5.1).

Proof of Theorem 3. In the same way as in the proof of Lemma 10, we easily verify

$$
\pm(u^{\pm})_{\mu} > 0. \tag{6.2}
$$

Let $(K, c) \in \mathcal{D}_{1,1} \cup \mathcal{D}_{1,3}$ and $q \in (u_1, u_0)$. It is clear that (u, w) is a periodic solution of (1.6) with (1.7) if and only if $u^+ = u^- \in (u_0, u_2)$. By the same argument as in the proof of Lemma 7, if μ (resp. $-\mu$) is large enough for arbitrarily fixed $a \in (q, u_2)$, $u^+ > a$ (resp. $u^- > a$). Define

$$
\underline{\mu}^{+} \equiv \inf \{ \mu \in \mathbb{R} \mid u^{+} > u_{0} \}, \quad \overline{\mu}^{-} \equiv \sup \{ \mu \in \mathbb{R} \mid u^{-} > u_{0} \},
$$

$$
\overline{\mu}^{+} \equiv \inf \{ \mu \in \mathbb{R} \mid u^{+} = u_{2} \}, \quad \underline{\mu}^{-} \equiv \sup \{ \mu \in \mathbb{R} \mid u^{-} = u_{2} \}.
$$

Then $-\infty \leq \underline{\mu}^+ < \overline{\mu}^+ < \infty$ and $-\infty < \underline{\mu}^- < \overline{\mu}^- \leq \infty$. By (6.2), we have

$$
u^{+}\begin{cases} \in (u_{0}, u_{2}) & \text{if } \underline{\mu}^{+} < \mu < \overline{\mu}^{+}, \\ = u_{2} & \text{if } \mu \geq \overline{\mu}^{+}, \end{cases}
$$

$$
u^{-}\begin{cases} = u_{2} & \text{if } \mu \leq \underline{\mu}^{-}, \\ \in (u_{0}, u_{2}) & \text{if } \underline{\mu}^{-} < \mu < \overline{\mu}^{-}. \end{cases}
$$

$$
\mu^{+} < \overline{\mu}^{-}, \quad \overline{\mu}^{+} > \mu^{-}. \tag{6.3}
$$

It is therefore sufficient to show that

The former inequality above is shown in the same way as the proof of Lemma 11. To derive the latter inequality, we observe that $u^+ < u_1^+$ when $u^+ \in (u_0, u_2)$, which follows from the fact that the orbits $\{(u, w_1^+(u)) \mid u_1 < u < u_1^+\}$ and $\{(u, w^+(u)) \mid q < u < u^+\}$ cannot intersect. In particular, we have $u^+ < u_1^+ = u_2$ if $\mu = \mu_b$. Hence it follows that $\overline{\mu}^+ > \mu_b$. In a similar manner, we also have $\underline{\mu}^- < \mu_f$. Since the inequality $\mu_b \geq \mu_f$ holds under the condition $(K, c) \in \mathcal{D}_{1,1} \cup \mathcal{D}_{1,3}$, we obtain the latter inequality of (6.3). Therefore we conclude that for each q satisfying (1.7), there is μ_{per} such that a periodic solution of (1.6) with $(u(0), w(0)) = (q, 0)$ exists for $\mu = \mu_{per}$. The uniqueness of μ_{per} follows from (6.2).

In the same way as in the proof of Lemma 13, we readily see that the condition $(K, c) \in \mathcal{D}_1 \cup \mathcal{D}_2$ is a necessary condition for the existence of a periodic solution of (1.6) with (1.7) . Finally we prove that there exists no periodic solutions if $\mu \neq \mu_{per}$ in (1.6) with (1.7). We first assume that $(K, c) \in \mathcal{D}_{1,1} \cup \mathcal{D}_{1,3}$.

Let (u, w) be a periodic solution of (1.6) with the period $Z > 0$. Then there exist q and z_0 such that $(u(z_0), w(z_0)) = (q, 0)$. If q is not in $[u_1, u_2]$, we see from Lemma 2 that

$$
\{(u(z), w(z)) \mid z > z_0\} \subset \begin{cases} \{(u, w) \mid 0 < u < q, w < 0\} & \text{if } q < u_1, \\ \{(u, w) \mid u > q, w > 0\} & \text{if } q > u_2, \end{cases}
$$

contrary to the fact that $(u(z_0+Z), w(z_0+Z)) = (q, 0)$. Hence we have $q \in (u_1, u_0) \cup (u_0, u_2)$ because (u, w) is not an equilibrium. If $q \in (u_1, u_0)$, then we must have $\mu = \mu_p e^p$ by the above argument. Hence a periodic solution exists only for $\mu = \mu_{per}$. If $q \in (u_0, u_2)$, the orbit $\{(u(z), w(z)) \mid z > z_0\}$ meets a point $(\tilde{q}, 0)$ with $\tilde{q} \in (u_1, u_0)$, since otherwise, one could show by Lemma 2 that

$$
\{(u(z), w(z)) \mid z > z_0\} \subset \{(u, w) \mid u < q, w < 0\},\
$$

contrary to the fact that $(u(z_0 + Z), w(z_0 + Z)) = (q, 0)$. Therefore we again have $\mu = \mu_{per}$.

We omit the discussion for the other cases because the same argument as above can be applied. Thus the proof is complete. 口

7 Bifurcations of traveling wave solutions

We have discussed several types of traveling wave solutions in Sections 3–6. It is then natural to investigate connections between them. In this section, we observe that some of the solutions converge to other solutions when parameters approach specific values. This study provides information on the structure of solutions in a bifurcation diagram.

To state the results of this section, we introduce some notation. Let (u_{pul}^j, w_{pul}^j) be the homoclinic orbit of (1.6) with (HO) for $\mu = \mu_{pul}^j$ and $\overline{u} = u_j$ in $j = 1, 2$, which is obtained in Theorem 2. Similarly, (u_{per}, w_{per}) denotes the solution of (1.6) with $(u_{per}(0), w_{per}(0)) = (q, 0)$ for q satisfying (1.7) and $\mu = \mu_{per}$ as seen in Theorem 3. We define Z_{per} to be the (fundamental) period of (u_{per}, w_{per}) . Moreover, set

$$
\mathcal{O}_{pul}^{j} = \{ (u_{pul}^{j}(z), w_{pul}^{j}(z)) \mid z \in \mathbb{R} \},
$$

$$
\mathcal{O}_{per} = \{ (u_{per}(z), w_{per}(z)) \mid 0 \le z < Z_{per} \},
$$

which represent the homoclinic and the periodic orbits in the phase plane, respectively. To emphasize the dependency of the parameters, we may write $\mathcal{O}_{pul}^j = \mathcal{O}_{pul}^j(K,c)$ and $\mathcal{O}_{per} = \mathcal{O}_{per}(K,c,q)$. Let \mathcal{O}_* be the heteroclinic cycle in (1.6) for $(c, \mu) = (c_*, \mu_*)$ consisting of two heteroclinic orbits connecting the equilibrium points $(u_1, 0)$, $(u_2, 0)$. Note that $\mathcal{O}_* = \mathcal{O}_1^+ \cup \mathcal{O}_1^ (= \mathcal{O}_2^+ \cup \mathcal{O}_2^-)$, where \mathcal{O}_j^{\pm} for $j = 1, 2$ were defined in Section 3.

We also introduce the notion of convergence for sets in \mathbb{R}^2 . Let $I \subset \mathbb{R}$ be an interval and let $\tau_0 \in \overline{I}$. For $A \subset \mathbb{R}^2$ and $\{A_\tau\}_{\tau \in I} \subset \mathbb{R}^2$, the notation $A_\tau \to A$ as $\tau \to \tau_0$ is used if $\{A_\tau\}$ converges to A with respect to the Hausdorff distance in \mathbb{R}^2 ([18]), that is,

$$
\max\left\{\sup_{a\in A}\inf_{b\in A_{\tau}}|a-b|,\sup_{b\in A_{\tau}}\inf_{a\in A}|a-b|\right\}\to 0 \text{ as } \tau\to\tau_0.
$$

We note that if A consists of a single point $(\overline{u}, \overline{w})$ and A_{τ} is an orbit $\{(u^{\tau}(z), w^{\tau}(z)) \mid z \in \mathbb{R}\},$ then the convergence of A_{τ} to A means that $(u^{\tau}(z), w^{\tau}(z)) \to (\overline{u}, \overline{w})$ uniformly for $z \in \mathbb{R}$ as $\tau \to \tau_0$.

The goal of this section is to present two propositions. First, we examine the relationship between the homoclinic orbits and the heteroclinic cycle (Proposition 3). Next, we show that the periodic orbit converges to the homoclinic orbit when q approaches the equilibrium point (Proposition 4).

Proposition 3. Assume (C). Then, for $j = 1, 2$,

$$
\mu_{pul}^j(K,c) \to \mu_*, \quad \mathcal{O}_{pul}^j(K,c) \to \mathcal{O}_*, \quad \text{as} \quad c \to c_*. \tag{7.1}
$$

Furthermore, there hold

$$
\mu_{pul}^1(K, c) \to 0, \quad \mathcal{O}_{pul}^1(K, c) \to \{(u_M, 0)\} \quad \text{as} \quad c \to c_M,\tag{7.2}
$$

$$
\mu_{pul}^2(K, c) \to 0, \quad \mathcal{O}_{pul}^2(K, c) \to \{(u_m, 0)\} \quad \text{as} \quad c \to c_m. \tag{7.3}
$$

Proposition 4. Assume the condition (C) . Then there holds

$$
\mu_{per} \rightarrow -f'(u_0), \ Z_{per} \rightarrow \frac{2\pi}{\omega_0}, \ \mathcal{O}_{per} \rightarrow \{(u_0, 0)\}\tag{7.4}
$$

as $q \to u_0$, where ω_0 was given in Lemma 3. Moreover,

$$
\mu_{per} \to \mu_{pul}^1, \ Z_{per} \to \infty, \ \mathcal{O}_{per} \to \mathcal{O}_{pul}^1 \quad as \ \ q \to u_1 \tag{7.5}
$$

if $(K, c) \in \mathcal{D}_{1,1}$,

$$
\mu_{per} \to \mu_{pul}^2, \ Z_{per} \to \infty, \ \mathcal{O}_{per} \to \mathcal{O}_{pul}^2 \quad as \ \ q \to u_2 \tag{7.6}
$$

if $(K, c) \in \mathcal{D}_{1,2} \cup \mathcal{D}_2$, and

$$
\mu_{per} \to \mu_*, \ Z_{per} \to \infty, \ \mathcal{O}_{per} \to \mathcal{O}_* \ \ as \ \ q \to u_1, u_2 \tag{7.7}
$$

if $(K, c) \in \mathcal{D}_{1,3}$.

- **Remark 2.** (i) Bifurcations of homoclinic and heteroclinic orbits are observed: (7.1) indicates that the homoclinic orbits \mathcal{O}^1_{pul} and \mathcal{O}^2_{pul} bifurcate from the heteroclinic cycle \mathcal{O}_* ; (7.5) indicates that the periodic orbit \mathcal{O}_{per} becomes the homoclinic orbit \mathcal{O}_{pul}^j or the heteroclinic cycle \mathcal{O}_* when the initial value q approaches u_j . A Hopf bifurcation is also observed: (7.4) shows that \mathcal{O}_{per} bifurcates from $(u_0, 0)$.
- (ii) We note that the equalities

$$
K = V'(u_M) = V'(u_m)
$$
\n(7.8)

hold since u_M and u_m are critical points of the function $Ku - V(u)$. These with (2.6) and (2.7) show that $f'(u_0) = -\lim_{q\to u_0} \mu_{per}$ converges to 0 as $c \to c_M, c_m$. Therefore (7.2) and (7.3) imply that a Bogdanov-Takens bifurcation ([19]) occurs at $(c, \mu) = (c_M, 0), (c_m, 0)$. We emphasize that the propositions yield information on not a local bifurcation diagram but a global one; the proofs will be done without using the theory of local bifurcations.

7.1 Proof of Proposition 3

In the following proofs of this subsection, we ignore the dependence on K in order to simplify notation. Before the proof of Proposition 3, we examine the behavior of u_1^{\pm} and w_1^{\pm} (resp. u_2^{\pm} and w_2^{\pm}) as $c \to c_M$ (resp. $c \to c_m$).

Lemma 15. Give $\mu_{\infty}^{\pm} \in [-\infty, 0]$ and $\mu_{\infty}^{-} \in [0, \infty]$ arbitrarily. Fix $K > 0$. If (c, μ) converges to $(c_M, \mu_{\infty}^{\pm})$ and satisfies $(K, c) \in \mathcal{D}_1$, then $u_1^{\pm}(K, c, \mu) \to u_M(K)$ and $w_1^{\pm}(u_1, K, c, \mu) \to 0$ uniformly in u under the limit. Similarly, if $(K, c) \in \mathcal{D}_1 \cup \mathcal{D}_2$ and $(c, \mu) \to (c_m, \mu^{\pm}_{\infty}),$ then $u_2^{\pm}(K, c, \mu) \to u_m(K)$ and $w_2^{\pm}(u; K, c, \mu) \to 0$ uniformly in u.

Proof. We may represent $w_1^+(u; c, \mu)$ by $w_1^+(u)$ for simplicity. In order to show the assertion for u_1^+ and w_1^+ , it is sufficient to consider only the case $\mu_{\infty}^{+} = 0$ because it follows from Lemmas 6 and 10 that

$$
u_0(c) \le u_1^+(c, \mu) \le u_1^+(c, 0), \quad 0 \le w_1^+(u; c, \mu) \le w_1^+(u; c, 0)
$$

if $\mu \leq 0$. Applying Lemma 5 with $W = w_1^+$, $A = 2g_1f$, $B = 2g_2h(\cdot, \mu)$, $s_1 = u_1$ and $s_2 = u$ gives

$$
w_1^+(u)^2 \le \int_{u_1}^u e^{u-s} \left(2g_1(s)f(s) + g_2(s)^2 h(s,\mu)^2\right) ds\tag{7.9}
$$

for $u \in [u_1, u_1^+]$. From this and (2.6), we particularly have

$$
w_1^+(u_0; c, \mu) \to 0 \text{ as } (c, \mu) \to (c_M, 0). \tag{7.10}
$$

Integrating (3.4) over $[u_0, u]$ and using the fact that $f \leq 0$ on $[u_0, u_2]$ yield

$$
w_1^+(u) \le w_1^+(u_0) + \int_{u_0}^u g_2(s)|h(s,\mu)|ds
$$

for $u \in [u_0, u_1^+]$. Furthermore, integrating (3.5) over $[u_0, u]$ and then plugging the above inequality into the result, we deduce that

$$
\frac{1}{2}w_1^+(u)^2 = \frac{1}{2}w_1^+(u_0)^2 + \int_{u_0}^u (g_1(s)f(s) + g_2(s)h(s,\mu)w_1^+(s))ds
$$
\n
$$
\leq \frac{1}{2}w_1^+(u_0)^2 + \int_{u_0}^u g_1(s)f(s)ds + \int_{u_0}^u g_2(s)|h(s,\mu)|\left(w_1^+(u_0) + \int_{u_0}^s g_2(\tau)|h(\tau,\mu)|d\tau\right)ds
$$
\n(7.11)

for $u \in [u_0, u_1^+]$.

Define $u^* \equiv \limsup_{(c,\mu)\to(c_M,0)} u_1^+(c,\mu)$. By (2.6) and (7.10), we see that the right-hand side of (7.11) converges to

$$
I(u) \equiv \int_{u_M}^u g_1(s) f(s; c_M) ds + \int_{u_M}^u g_2(s) |h(s, 0)| \left(\int_{u_M}^s g_2(\tau) |h(\tau, 0)| d\tau \right) ds
$$

for each $u \in [u_M, u^*]$ as $(c, \mu) \to (c_M, 0)$, where we used the notation $f(u; c)$ to emphasize c-dependency of f. Recall that $f(u_M; c_M) = f'(u_M; c_M) = 0$ and $h(u, 0) = f'(u; c) < 0$ for $u \in (u_M, u_m)$. Then $I(u)$ is estimated as

$$
I(u) \le \int_{u_M}^{u} (g_1(s) + Cf'(s; c_M)) f(s; c_M) ds \tag{7.12}
$$

for $u \in [u_M, \min\{u^*, u_m\}],$ where $C > 0$ is some constant. Then u^* must be equal to u_M . Otherwise, since $f'(u; c_M)$ is sufficiently small, the integral on the right-hand side of (7.12) is negative for u close to u_M , which contradicts the fact that the left-hand side of (7.11) is nonnegative. Therefore $u_1^+(c,\mu) \to u_M$ as $(c, \mu) \rightarrow (c_M, 0)$. From this and (7.9), we also obtain the uniform convergence of w_1^+ to 0.

The remainder of the lemma can be shown in the same way as above. So we omit the details of the proofs. \Box

Proof of Proposition 3. Recall that $\mu_f < \mu_{pul}^1 < \mu_b$ and $\mu_b < \mu_{pul}^2 < \mu_f$, which were shown in the proof of Theorem 2. We hence have $\mu_{pul}^j \to \mu_*$ as $c \to c_*$. This with the continuous dependence of stable and unstable manifolds on parameters gives the convergence of \mathcal{O}_{pul}^j to \mathcal{O}_* . We have therefore proved (7.1).

Let us show (7.2). It suffices to verify that $\mu_{pul}^1 \to 0$ as $c \to c_M$. Indeed, if this is true, it is shown that $\mathcal{O}_{pul}^1 \rightarrow \{(u_M, 0)\}$ from Lemma 15 and the fact that

$$
\mathcal{O}_{pul}^1 = \{(u, w_1^+(u; c, \mu_{pul}^1)) \mid u \in (u_1, u_1^+]\} \cup \{(u, w_1^-(u; c, \mu_{pul}^1)) \mid u \in (u_1, u_1^-]\}.
$$

Let ${c_n}_n$ be any sequence such that $\lim_{n\to\infty} c_n = c_M$ and

$$
\mu_{\infty} \equiv \lim_{n \to \infty} \mu_{pul}^1(c_n) \in [-\infty, \infty]
$$

exists. We prove $\mu_{\infty} = 0$. We apply Lemma 15 for $\mu_{\infty} = \mu_{\infty}^{+}$ if $\mu_{\infty} \le 0$ and for $\mu_{\infty} = \mu_{\infty}^{-}$ if $\mu_{\infty} \ge 0$. In either case, we have

$$
u_1^n \equiv u_1^+(c_n, \mu_{pul}^1(c_n)) = u_1^-(c_n, \mu_{pul}^1(c_n)) \to u_M
$$
\n(7.13)

as $n \to \infty$. We now use the inequalities

$$
-\sup_{u\in(u_1,u_1^n)}f'(u)<\mu_{pul}^1(c_n)<-\inf_{u\in(u_1,u_1^n)}f'(u),
$$

which follows from Lemma 14. From (2.6) , (7.8) , and (7.13) , we find that both the left-hand and the righthand sides converge to 0 as $n \to \infty$, which leads to $\mu_{\infty} = 0$. We can derive (7.3) in a similar way, and thus the proof is complete. the proof is complete.

7.2 Proof of Proposition 4

We define orbits \mathcal{O}^{\pm} by

$$
\mathcal{O}^{\pm} \equiv \begin{cases} \{ (u, w^{\pm}(u)) \mid u \in [q, u^{\pm}] \} & \text{if } q \in (u_1, u_0), \\ \{ (u, w^{\pm}(u)) \mid u \in [u^{\pm}, q] \} & \text{if } q \in (u_0, u_2). \end{cases}
$$

Lemma 16. For $j = 1, 2$, there hold $\mathcal{O}^{\pm} \to \overline{\mathcal{O}^{\pm}_j}$ locally uniformly in $\mu \in \mathbb{R}$ as $q \to u_j$, where \mathcal{O}^{\pm}_j were defined in (3.3). In particular, $u^{\pm} \rightarrow u_j^{\pm}$ as $q \rightarrow u_j$.

Proof. The convergence of \mathcal{O}^{\pm} to \mathcal{O}_j^{\pm} in a neighborhood of the equilibrium $(u_j, 0)$ follows from the Hartman-Grobman theorem. The proof for the convergence away from equilibria is then standard. \Box

Let us conclude this section by showing Proposition 4.

Proof of Proposition 4. From (ii) of Lemma 3, we see that no periodic orbit exists in a neighborhood of the equilibrium $(u_0, 0)$ provided $\mu \neq -f'(u_0)$. Hence μ_{per} must converge to $-f'(u_0)$ as $q \to u_0$. By the same argument as in the proof of Theorem 2 (ii), the behavior of (u_{per}, w_{per}) is approximated by

 $(u_0, 0) + (q - u_0)(\cos \omega_0 z, -\omega_0 \sin \omega_0 z)$

uniformly in z if q is close to u_0 . This implies that $(u_{per}, w_{per}) \to (u_0, 0)$ and $Z_{per} \to 2\pi/\omega_0$ as $q \to u_0$ from the representation above. Therefore (7.4) holds.

Let us proceed to the proof of (7.5). We consider the case $(K, c) \in \mathcal{D}_{1,1}$. To emphasize the dependency of q, we may write $\mu_{per} = \mu_{per}^q$ and $u^{\pm} = u^{q,\pm}$. On the other hand, we ignore the dependence of u_1^{\pm} and $u^{q,\pm}$ on (K, c) and write $u_1^{\pm}(\mu)$ and $u^{q,\pm}(\mu)$ instead of $u_1^{\pm}(K, c, \mu)$ and $u^{q,\pm}(K, c, \mu)$ for simplicity. We show that $\mu_{per}^q \to \mu_{pul}^1$ as $q \to u_1$ by contradiction. If this is false, we can take a constant $\varepsilon_0 > 0$ and a sequence $\{q_n\}_n$ which satisfies $q_n \to u_1$ as $n \to \infty$ and $|\mu_{per}^{q_n} - \mu_{pul}^{1}| \geq \varepsilon_0$ for all n. Let us consider the case that $\mu_{per}^{q_n} \ge \mu_{pul}^1 + \varepsilon_0$ for infinitely many n. By (6.2), we have

$$
u^{q_n,+}(\mu_{pul}^1+\varepsilon_0) \le u^{q_n,+}(\mu_{per}^{q_n}) = u^{q_n,-}(\mu_{per}^{q_n}) \le u^{q_n,-}(\mu_{pul}^1+\varepsilon_0).
$$

Hence letting $n \to \infty$ and using Lemma 16 yield $u_1^+(\mu_{pul}^1 + \varepsilon_0) \le u_1^-(\mu_{pul}^1 + \varepsilon_0)$. However it follows from Lemma 10 that

$$
u_1^+(\mu_{pul}^1 + \varepsilon_0) > u_1^+(\mu_{pul}^1) = u_1^-(\mu_{pul}^1) > u_1^-(\mu_{pul}^1 + \varepsilon_0),
$$

which is a contradiction. We can deal with the other case in the same way as above.

The convergence of \mathcal{O}_{per} to \mathcal{O}_{pul}^1 is verified by combining Lemma 16 and the fact that $\mu_{per}^q \to \mu_{pul}^1$ as $q \to u_1$. Furthermore, since the initial value $(u_{per}(0), w_{per}(0)) = (q, 0)$ approaches the equilibrium $(u_1, 0)$, it is easy to see that Z_{per} diverges to ∞ . We have thus proved (7.5), and the proof is complete. is easy to see that Z_{per} diverges to ∞ . We have thus proved (7.5), and the proof is complete.

8 Numerical continuation of traveling wave solutions

We illustrate all branches of the traveling wave solutions in (1.6) with the constants $V_0 = 0.0168, M =$ $0.913, u_c = 0.025, \beta = 89.7$, which are identical to those in Figure 1. These constants are fixed throughout this section. We used the numerical continuation package HomCont/AUTO [13] for heteroclinic, homoclinic, and periodic orbits. The numerical approximations of the heteroclinic and homoclinic orbits were achieved by solving a truncated problem using the projection boundary conditions. Refer to [20], [21], and [22] for the theoretical background.

We study the model proposed by Lee et al., characterized by the function $\kappa(\rho) = 1/(6\tau\rho^2)$. Notably, the system described by (1.6) remains independent of τ . We fixed $K = 1.25$, resulting in approximate values of $c_1 = c_m \approx 0.0151$ and $c_M \approx 0.0167$. Therefore, Theorem 1 stipulates the existence of traveling back and front solutions when $c_m < c < c_M$ and $\mu = \mu_b, \mu_f$. Furthermore, traveling pulse solutions emerge when $\mu = \mu_{pul}^1, \mu_{pul}^2.$

Figure 2: Numerical continuations for heteroclinic and homoclinic orbits, and Hopf bifurcation points. (a) Branches of traveling back (black), front (red) and pulse (blue) solutions. The green curve represents the graph of $\mu = -f'(u_0)$. (ii) of Lemma 3 shows that a Hopf bifurcation occurs in (1.6) at each point of the green curve. All the branches terminate at the values of $c = c_m$ or $c = c_M$ where two of the equilibriums among the three collide. (b), (c) Corresponding orbits for traveling pulse solutions at p_1, p_2, p_3, p_4 on the blue branches in (a). They are homoclinic orbits in the (u, w) phase plane, which are represented by black curves. Red curves indicate the heteroclinic cycle (HC), consisting of two heteroclinic orbits at the intersection (c_*, μ_*) of the black and red curves in (a). The black disks in (b), (c) are $(u_m, 0)$ and $(u_M, 0)$ corresponding to the endpoints $(c_m, 0)$ and $(c_M, 0)$ of the blue and green curves in (a), respectively.

Figure 2 illustrates the branches of heteroclinic and homoclinic orbits within the (c, μ) -parameter space, obtained through numerical continuations. Each point (c, μ) along the branches (black, red, or blue) in the upper-left plot corresponds to specific parameter settings conducive to heteroclinic or homoclinic orbits. These branches represent the loci of $\mu = \mu_b, \mu_f, \mu_{pul}^1, \mu_{pul}^2$. Notably, the crossing of two branches of heteroclinic orbits occurs at the point (c_*, μ_*) , indicating the presence of a heteroclinic cycle. Additionally, the two branches of the homoclinic orbits diverge from this point. These findings are consistent with Theorems 1 and 2, and Proposition 3.

The bifurcation of the homoclinic orbit from the heteroclinic cycle was discussed in [11]. While this previous study necessitated non-degeneracy hypotheses, we do not undertake such analytical investigations. However, numerical estimates of saddle quantity, computed as the sum of eigenvalues at saddle points O_1 = $(u_1, 0)$ and $O_2 = (u_2, 0)$ in (1.6), were performed. We estimated $u_1 = 0.0166$ and $u_2 = 0.0344$, along with the parameter values (c_*, μ_*) = (0.01611, 0.0734). Eigenvalue calculations for $\lambda_{\pm}(u_i)$, as defined in Lemma 3, yielded approximate values of $(\lambda_-(u_1), \lambda_+(u_1)) = (-52.88, 117.82)$ and $(\lambda_-(u_2), \lambda_+(u_2)) = (-27.90, 66.08)$. Consequently, positive saddle quantities were deduced at $(u_1, 0)$ and $(u_2, 0)$, signifying that the homoclinic orbit branches from O_1 to O_1 and from O_2 to O_2 are tangential to the heteroclinic orbit branches from O_1 to O_2 and from O_2 to O_1 at (c^*, μ^*) , respectively. This tangency is evident in Figure 2 (a).

The numerical continuation process relies on having approximate heteroclinic orbit as a starting point, which must be sufficiently accurate. While having an exact solution at a specific parameter value is advantageous, it is not feasible for (1.6). However, the Allen-Cahn-Nagumo equation

$$
\begin{cases} u_z = w, \\ w_z = -\mu w + u(u - a)(u - 1) \end{cases} \tag{8.1}
$$

offer exact families of heteroclinic solutions

$$
u(z) = \frac{1}{1 + e^{z/\sqrt{2}}}, \quad \mu = \sqrt{2} \left(\frac{1}{2} - a\right)
$$
 (8.2)

for $a \in (0,1)$, and homoclinic solutions

$$
u(z) = \frac{6a}{2(1+a) + \sqrt{2(2-a)(1-2a)}\cosh\sqrt{a}z}, \quad \mu = 0
$$
\n(8.3)

Figure 3: Continuation of periodic orbits from the Hopf to homoclinic bifurcation. (a), (c) Maximum of u versus μ . (b), (d) Orbits in the (u, w) phase plane. We set $c = 0.0163$ in the upper figures, whereas $c = 0.0155$ in the lower ones. The equilibrium point $(u_0, 0)$ is denoted by A, while B and C represent periodic orbits, and D corresponds to a homoclinic orbit.

for $a \in (0, 1/2)$. These exact solutions can serve as the seeds for homotopy continuation from (8.1) to (1.6) for $a \in (0, 1/2)$. Specifically, we scaled the nonlinear terms $u(u-a)(u-1)$ in (8.1) linearly, defining $f_{AC}(u; u_1, u_0, u_2) = (u - u_1)(u - u_0)(u - u_2)$, and performed homotopy continuation for

$$
\begin{cases}\nu_z = w, \\
w_z = -\mu w + (1 - \phi)f_{AC}(u; u_1, u_0, u_2) + \phi(g_1(u)f(u) + g_2(u)h(u, \mu)w),\n\end{cases} (8.4)
$$

where $\phi \in [0,1]$ represents a homotopic parameter. Note that $f_{AC}(u; u_1, u_0, u_2)$ and $f(u)$ have the same zeroes for $(K, c) \in \mathcal{D}_1$. At $\phi = 0$, the exact solutions of the heteroclinic and homoclinic orbits are as described earlier. Therefore, the continuation from $\phi = 0$ to 1 yields approximate solutions for (1.6).

Figure 4: (a) Continuation branches for the traveling back (black), traveling front (red), and traveling pulse (green, blue) solutions in the (K, c) -parameter space. (b) Orbits in the (u, w) phase plane. The curves A, B (blue) and C, D (green) correspond to the points along the branches of traveling pulse solutions in (a).

We delve into the bifurcation of periodic orbits, as outlined in Proposition 4, using the same parameters depicted in Figure 2. $\mu = -f'(u_0)$ serves as a critical value for each $c \in (c_m, c_M)$, where the middle

equilibrium $(u_0, 0)$ possesses purely imaginary eigenvalues, as denoted by the green curve in Figure 2. We can numerically trace the continuation of periodic orbits from the Hopf bifurcation point by considering μ as a control parameter while c remains fixed in (c_m, c_M) . The branches of these periodic orbits seem to culminate in the homoclinic orbit, represented by the blue curves in Figure 2 (a). Figure 3 showcases two typical situations: $(K, c) \in \mathcal{D}_{1,1}((a), (b))$ and $(K, c) \in \mathcal{D}_{1,2}((c), (d))$. We infer that periodic orbits exist within the parameter region delineated by the three curves in Figure 2 (a): the two blue curves (branches of homoclinic orbits) and the green curve (Hopf bifurcation points).

Furthermore, we delineated the branches of heteroclinic and homoclinic orbits in the (K, c) -parameter space illustrated in Figure 4, where we set $\mu = 2\tau K - 1$. As depicted in the figure, bifurcation branches for heteroclinic and homoclinic orbits correspond to the boundary conditions (HE1), (HE2), and (HO), considering a given u_1, u_2 . Obtaining a figure akin to Figure 4 rigorously is challenging because K and c feature in various locations in (1.6) . Hence, regarding μ as an independent parameter is apt. We could obtain qualitatively similar results for the Kühne model as shown in Figure 4 for Lee et al.'s model. However, we have not reported these findings.

We discuss the relationship between the traveling wave solutions for (1.6) and those of the original problem (1.1) . We numerically compute K and c using the data used from Figure 1 and confirm that the preceding discussion implies the existence of a traveling wave solution in (1.1). First, we estimate the time required for the pulse to traverse the region as $T = 178$. After that, $c = L/T$ is approximated to be 0.0130899. Subsequently, we determine $(\rho_-, v_-) = (26.516, 0.029025)$ using the data for $\rho(x, t)$ at $(x, t) = (0, 1000)$. Our observations indicate that $x = 0$ lies outside the congestion phase at $t = 1000$. We use (1.5) to find $K = \rho_-(v_- + c) \approx 1.11672$. Using the estimated values of (K, c) , and employing AUTO like in the case from which we obtained the results depicted in Figure 3, we derive a periodic traveling wave solution with a period of 2.33022, which closely aligns with $L = 2.33$. Moreover, we obtain $c = 0.013089412$ and $\mu_{per} = 0.11640802175$. We note that $(K, c) \in \mathcal{D}_1$, μ_{per} is approximately $2\tau K - 1 \approx 0.11672$, and (c, μ_{per}) is included in the parameter region bounded by the three curves related to the branches of homoclinic and Hopf bifurcation points, as shown in Figure 2 (a). Hence, we conclude that the solution illustrated in Figure 1 coincides with the periodic orbit verified numerically by AUTO.

9 Discussions

This study rigorously established the existence of various traveling wave solutions in the macroscopic traffic model (1.1). The emergence of congested states as time-periodic solutions in microscopic models is highlighted via Hopf bifurcation, which is a promising method for obtaining such solutions. However, obtaining a solution away from the bifurcation point is often infeasible. Alternatively, [23] employed the step function as an OV function and successfully formally constructed a solution corresponding to the congestion phase. Consequently, strong restrictions are usually necessary to rigorously obtain the congestion phase in microscopic models. Therefore, continuous models are useful in treating traveling wave solutions in a congestion phase.

In presenting Theorems 2 and 3, condition (H) was considered. If the viscosity coefficient $\kappa(\rho)$ exhibits a strong singularity at $\rho = 0$, as in Lee et al.'s model, all theorems in this study remain valid. However, (H) does not hold in the case where $\kappa(\rho)$ is constant, as in the Kühne model, or has a weak singularity, as in the Kerner and Konhäuser model. Actually, constructing a heteroclinic orbit is feasible in (1.6) protruding outside the region $\{u > 0\}$ even if $\kappa(\rho) = \kappa_0$. However, this solution is meaningless in (1.1) because $\rho = 1/u$ should be positive. Further analyses are necessary to obtain analogous results to Proposition 2 without (H).

References

[1] D. Helbing, Traffic and related self-driven many-particle systems, Reviews of modern physics 73 (4) (2001) 1067.

- [2] H. J. Payne, A critical review of a macroscopic freeway model, in: Research directions in computer control of urban traffic systems, ASCE, 1979, pp. 251–256.
- [3] R. Kühne, Macroscopic freeway model for dense traffic-stop-start waves and incident detection, Transportation and traffic theory 9 (1984) 21–42.
- [4] M. Bando, K. Hasebe, A. Nakayama, A. Shibata, Y. Sugiyama, Structure stability of congestion in traffic dynamics, Japan Journal of Industrial and Applied Mathematics 11 (1994) 203–223.
- [5] R. K¨uhne, Freeway speed distribution and acceleration noise calculations from a stochastic continuum theory and comparison with measurements, Transportation and traffic theory 12 (1987) 119–137.
- [6] B. S. Kerner, P. Konhäuser, Cluster effect in initially homogeneous traffic flow, Physical review E 48 (4) (1993) R2335.
- [7] H. Lee, H.-W. Lee, D. Kim, Macroscopic traffic models from microscopic car-following models, Physical Review E 64 (5) (2001) 056126.
- [8] P. Berg, A. Mason, A. Woods, Continuum approach to car-following models, Physical Review E 61 (2) (2000) 1056.
- [9] P. Berg, A. Woods, Traveling waves in an optimal velocity model of freeway traffic, Physical review E 63 (3) (2001) 036107.
- [10] I. Gasser, G. Sirito, B. Werner, Bifurcation analysis of a class of 'car following'traffic models, Physica D: Nonlinear Phenomena 197 (3-4) (2004) 222–241.
- [11] H. Kokubu, Homoclinic and heteroclinic bifurcations of vector fields, Japan Journal of Applied Mathematics 5 (1988) 455–501.
- [12] D. G. Aronson, H. F. Weinberger, Nonlinear diffusion in population genetics, combustion, and nerve pulse propagation, in: Partial Differential Equations and Related Topics: Ford Foundation Sponsored Program at Tulane University, January to May, 1974, Springer, 2006, pp. 5–49.
- [13] E. J. Doedel, A. R. Champneys, F. Dercole, T. F. Fairgrieve, Y. A. Kuznetsov, B. Oldeman, R. Paffenroth, B. Sandstede, X. Wang, C. Zhang, Auto-07p: Continuation and bifurcation software for ordinary differential equations (2007).
- [14] H. Lee, H.-W. Lee, D. Kim, Steady-state solutions of hydrodynamic traffic models, Physical Review E 69 (1) (2004) 016118.
- [15] D. Gilbarg, N. S. Trudinger, Elliptic partial differential equations of second order, Classics in Mathematics, Springer-Verlag, Berlin, 2001, reprint of the 1998 edition.
- [16] L. C. Evans, Partial differential equations, Vol. 19 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 1998. doi:10.1090/gsm/019.
- [17] A. Lunardi, Analytic semigroups and optimal regularity in parabolic problems, Vol. 16 of Progress in Nonlinear Differential Equations and their Applications, Birkhäuser Verlag, Basel, 1995. doi:10.1007/978-3-0348-9234-6.
- [18] T. Kato, Perturbation theory for linear operators, Classics in Mathematics, Springer-Verlag, Berlin, 1995, reprint of the 1980 edition.
- [19] Y. A. Kuznetsov, Elements of applied bifurcation theory, 3rd Edition, Vol. 112 of Applied Mathematical Sciences, Springer-Verlag, New York, 2004. doi:10.1007/978-1-4757-3978-7.
- [20] E. J. Doedel, M. J. Friedman, Numerical computation of heteroclinic orbits, Continuation Techniques and Bifurcation Problems (1990) 155–170.
- [21] M. J. Friedman, E. J. Doedel, Numerical computation and continuation of invariant manifolds connecting fixed points, SIAM Journal on Numerical Analysis 28 (3) (1991) 789–808.
- [22] W.-J. Beyn, The numerical computation of connecting orbits in dynamical systems, IMA Journal of Numerical Analysis 10 (3) (1990) 379–405.
- [23] Y. Sugiyama, H. Yamada, Simple and exactly solvable model for queue dynamics, Physical Review E 55 (6) (1997) 7749.