IBIS PRIMITIVE GROUPS OF ALMOST SIMPLE TYPE

FABIO MASTROGIACOMO AND PABLO SPIGA

ABSTRACT. Let G be a finite permutation group on Ω . An ordered sequence $(\omega_1 \ldots, \omega_\ell)$ of elements of Ω is an irredundant base for G if the pointwise stabilizer is trivial and no point is fixed by the stabilizer of its predecessors. The minimal cardinality of a base is said to be the base size of G. If all irredundant bases of G have the same cardinality, G is said to be an IBIS group.

In this paper, we classify the finite almost simple primitive IBIS groups whose base size is at least 6.

1. INTRODUCTION

A **base** for a permutation group G is a sequence $(\omega_1, \ldots, \omega_\ell)$ whose pointwise stabilizer in G is the identity. A base is said to be **irredundant** if no base point is fixed by the stabilizer its predecessors. Therefore, this gives rise to a decreasing sequence of stabilizers

$$G > G_{\omega_1} > G_{\omega_1,\omega_2} > \dots > G_{\omega_1,\omega_2,\dots,\omega_{\ell-1}} > G_{\omega_1,\omega_2,\dots,\omega_{\ell}} = 1.$$

The minimal cardinality of a base, called the **base size** of G, is usually denoted by b(G) and has played a key role in the investigation of primitive groups.

Cameron and Fon-der-Flaass [16] (see also [15, Section 4.14]) have proved that in a finite permutation group the following conditions are equivalent:

- all irredundant bases have the same size;
- the irredundant bases are invariant under re-ordering;
- the irredundant bases are the bases of a matroid.

A permutation group satisfying one, and hence all, of these conditions is said to be **IBIS**, that is, **Irredundant Bases of Invariant Size**. In particular, in an IBIS group G, all irredundant bases have cardinality b(G) and this value is sometimes referred to as the **rank** of the IBIS group, since it is the rank of the corresponding matroid. Aside from this introductory section, we avoid using the term rank with this meaning, because it can be easily confused with other properties of permutation groups that also bear this name.

Lucchini, Morigi and Moscatiello [34] made the first attempt of classifying finite primitive IBIS groups. Their approach is via the O'Nan-Scott classification of primitive groups (we use the terminology from [36]).

Theorem 1.1 (Theorem 1.1, [34]). Let G be a primitive IBIS group. Then one of the following holds:

²⁰¹⁰ Mathematics Subject Classification. primary 20B15.

Key words and phrases. IBIS, base size, irredundant base, almost simple, Lie group.

The authors are members of the GNSAGA INdAM research group and kindly acknowledge their support.

(1.1) G is of affine type,

(1.2) G is almost simple,

(1.3) G is of diagonal type.

Moreover, G is a primitive IBIS group of diagonal type if and only if it belongs to the infinite family of diagonal groups $\{PSL_2(2^f) \times PSL_2(2^f) \mid f \in \mathbb{N}, f \geq 2\}$ having degree $|PSL_2(2^f)| = 2^f (4^f - 1)$.

In light of this result, the problem of understanding finite primitive IBIS groups is reduced to affine groups and to almost simple groups. A first attempt of classifying finite primitive IBIS groups of affine type can be found in [35].

Lee and the third author [31] have classified the almost simple primitive IBIS groups having socle an alternating group. In the same paper, the authors have proposed a conjecture; indeed, they have conjectured that every almost simple primitive IBIS group is one of the groups in [31, Table 1]. However, we discovered two additional families of primitive IBIS groups that do not appear in the table of [31], namely $\Omega_d^{\pm}(q)$, with $q = 2^f$, acting on non-singular 1-dimensional subspaces. We have reported [31, Table 1] in Table 1 with these two new families.

Later, Lee [30] has classified the almost simple primitive IBIS groups having socle a sporadic simple group.

There are two special circumstances where IBIS groups arise. A permutation group G on Ω is said to be **sharply** *k*-**transitive** if, for any two *k*-tuples of distinct elements of Ω , there exists a unique element of G mapping the first *k*-tuple to the second. Observe that if G is sharply *k*-transitive, then b(G) = k and G is an IBIS group. Moreover, a transitive permutation group G is said to be **geometric** (or, base-transitive, see [1]) if G permutes its irredundant bases transitively. Clearly, every sharply *k*-transitive group is geometric and every geometric group is IBIS.^a

In this paper, we deal with almost simple primitive groups having socle a simple group of Lie type.

Theorem 1.2. Let G be an almost simple primitive IBIS group having socle a simple group of Lie type G_0 , with $b(G) \ge 6$ when G_0 is exceptional and $b(G) \ge 4$ when G_0 is classical. Then G is either $SL_d(2)$ or $Sp_d(2)$ in its natural action on the non-zero vectors of a d-dimensional vector space over the field with two elements, or $G = \Omega_d^{\pm}(q)$ with $d \ge 4$ and $q \ge 4$ acting on the non-singular 1-dimensional subspaces of the d-dimensional vector space over the field with q elements.

Our proof of Theorem 1.2 uses the astoneshing work on the Cameron-Kantor conjecture [14]. Most interest in the base size of primitive groups originated from Jordan's classic results [26], which bound the cardinality of a primitive group via its base size. This interest was spurred in the '90s by the Cameron-Kantor conjecture [14]: there exists an absolute constant b with $b(G) \leq b$, for every almost simple primitive group in a non-standard action. (We refer to [14] and to Section 1.1 for the definition of **standard action**). This conjecture was settled in the positive in [11]; however, the refinement of Cameron [15], asking whether one can take b = 7 has required considerable more effort. The detailed analysis of Burness [4, 5, 6, 7] on fixed point ratios has resulted in an answer to Cameron's question in [8, 10, 12].

 $\mathbf{2}$

^aIn Table 1, the third column indicates whether the group is sharply k-transitive, denoted by a \checkmark , and similarly, whether the group is geometric, marked with a \checkmark in the forth column.

From these papers, one can infer that "most" almost simple primitive groups in non-standard actions have base size 2.^b

Therefore, our proof of Theorem 1.2 uses deatailed information on the standard actions. Indeed, we prove this stronger result.

Theorem 1.3. Let G be an almost simple primitive IBIS group having socle a simple classical group in a standard action. Then, one of the following holds:

- (1.1) $G = SL_d(2)$ or $G = Sp_d(2)$ in its natural action on the non-zero vectors of an n-dimensional vector space over the field with two elements,
- (1.2) G has socle $PSL_2(q)$ and G is endowed of its natural action on the q + 1 points of the projective line,
- (1.3) the socle of G is $\text{Sp}_4(2)'$ in its natural action on the fifteen non-zero vectors of a 4-dimension vector space over the field with two elements.
- (1.4) $G = \Omega_d^{\pm}(q), d \ge 4, q \ge 4$ in its natural action on the non-singular 1dimensional subspaces.^c

While proving Theorem 1.3, we have delved deeply into the problem of classifying almost simple primitive IBIS groups. As a result, in the future, we intend to revisit this issue and examine the case of groups having base size at most 5. Indeed, in our opinion, there is some hope in dealing with groups having small base size. For instance, suppose that G is an almost simple primitive IBIS group with b(G) = 2. In particular, all irredundant bases of G have cardinality 2. Thus, for any two distinct points α and β in the domain of G we have $G_{\alpha} \cap G_{\beta} = 1$. Therefore G is a Frobenius group, contradicting the fact that G is almost simple. This simple argument shows that, if G is an almost simple primitive IBIS group, then $b(G) \geq 3$. Therefore, from Theorem 1.2, the cases that need to be considered are when $b(G) \in \{3, 4, 5\}$.^d

1.1. Standard and non-standard actions. To start with, we use well-known results bounding b(G) to deal with the so-called *non-standard actions*. The terminology below follows [8, Definition 2.1].^e

Definition 1.4. Let G be an almost simple classical group over \mathbb{F}_q , where $q = p^f$ and p is prime, with socle G_0 and associated natural module V. A subgroup H of G not containing G_0 is a **subspace subgroup** if, for each maximal subgroup M of G_0 containing $H \cap G_0$, one of the following holds:

- (1.1) M is the stabilizer in G_0 of a proper nonzero subspace U of V, where U is totally singular, non-degenerate, or, if G_0 is orthogonal and p = 2, a non-singular 1-space (U can be any subspace if $G_0 = \text{PSL}(V)$).
- (1.2) $G_0 = \tilde{\mathrm{Sp}}_{2m}(q)', \ p = 2, \text{ and } M = \mathrm{O}_{2m}^{\pm}(q).$

A subspace action of the classical group G is the action of G on the coset space [G : H], where H is a subspace subgroup of G. Note that the definition above amounts precisely to this: a maximal subgroup of G is a subspace subgroup if it lies in any C_1 -class, or is the even-characteristic symplectic case in the C_8 -class. This definition requires that we follow [28] in labeling the Aschbacher classes $C_1 - C_8$.

^bJan Saxl's pioneering work on primitive groups with base size not equal to 2 continues to inspire and drive research in this field. His legacy lives on as we pay tribute to his remarkable contributions.

^cIn this case, all bases have cardinality d - 1.

^dMost examples in Table 1 have $b(G) \in \{3, 4\}$.

^eOriginally, this notation was introduced by Liebeck and Shalev [33].

Group	Degree	b(G)	s. <i>k</i> -t.	g.	Comments
$\operatorname{Alt}(n)$	n	n-2	\checkmark	\checkmark	$n \ge 5$
$\operatorname{Sym}(n)$	n	n-1	\checkmark	\checkmark	$n \ge 5$
$PGL_2(q)$	q+1	3	\checkmark	\checkmark	
$\mathrm{PSL}_2(q)$	q+1	3			$q \mathrm{odd}$
$\mathrm{PSL}_2(q) \rtimes \langle \tau \rangle$	q+1	4			q odd, τ field automorphism
					of prime order
$\mathrm{PGL}_2(q) \rtimes \langle \tau \rangle$	q+1	4			$q > 4, \tau$ field automorphism
	-				of prime order
$\operatorname{Aut}(\operatorname{PSL}_2(4))$	5	4	\checkmark	\checkmark	-
$\mathrm{PSL}_2(q).\langle \tau \rangle$	q+1	3	\checkmark	\checkmark	$q = p^f$ odd, f even, τ not
2(1)()	1				field automorphism, $G \neq \mathrm{PGL}_2(q)$
M_{11}	11	4	\checkmark	\checkmark	
M_{12}^{11}	12	5	\checkmark	\checkmark	
M_{22}^{12}	22	5			
$M_{23}^{}$	23	6			
M_{24}	24	7			
$\operatorname{SL}_d(2)$	$2^{d} - 1$	n		\checkmark	$d \ge 3$
$\operatorname{Sp}_d(2)$	$2^{d} - 1$	n			$d \ge 4$
$\operatorname{Sp}_4(2)' \cong \operatorname{Alt}(6)$	15	3			
$^{2}B_{2}(q)$	$q^2 + 1$	3			$q = 2^{2f+1}, f \ge 1$
${}^{2}G_{2}(q)$	$q^3 + 1$	3			$q = 3^{2f+1}, f \ge 1$
$^{2}G_{2}(3)' \cong \mathrm{PSL}_{2}(8)$	28	3			
$SL_2(q)$	$\frac{(q-1)q}{2}$	3			$q = 2^f, f \ge 3$
$\operatorname{Aut}(\operatorname{SL}_2(q))$	$\frac{(q-1)q}{2}$	3			$q = 2^f, f > 3$ prime
$SL_2(4)$	$\ddot{6}$	3		\checkmark	
$P\Gamma L_2(q)$	$\frac{(q+1)q}{2}$	3			$q = 2^p, p \text{ odd prime}$
$\operatorname{SL}_2(q)$	$\sqrt{q}(q+1)$	3			$q = 2^f, f \ge 4, f$ even
$\operatorname{Alt}(7)$	15	3		\checkmark	
$\Omega_d^{\pm}(q)$	$q^{\frac{d}{2}-1}(q^{\frac{d}{2}} \mp 1)$	d-1		l	$q \ge 4, q$ even, $n \ge 4$
<i>u</i> (-/	- (1)				action on non-singular 1-spaces

TABLE 1. Results on sharply k-transitive and/or geometric almost simple IBIS groups.

A small extra collection of maximal subgroups arises due to the existence of peculiar outer automorphisms, when $G_0 = \text{Sp}_4(2^f)'$ and G contains a graph automorphism, or when $G_0 = P\Omega_8^+(q)$ and G contains a triality graph automorphism. We note that [28] explicitly excludes these cases, but they are included in [2].

Definition 1.5. A transitive action of G on a set Ω is said to be standard if, up to equivalence of actions, one of the following holds:

- (1.1) $G_0 = \operatorname{Alt}(m)$ and Ω is an orbit of subsets or uniform partitions of $\{1, \ldots, m\}$.
- (1.2) G is a classical group in a subspace action.

For an almost simple primitive permutation group in a non-standard action, the base size is bounded by an absolute constant. This was conjectured by Cameron and Kantor (see [17, 18]) and then settled in the affirmative by Liebeck and Shalev in [33, Theorem 1.3]. The constant was then made explicit in subsequent works [8, 9, 10, 11, 12]. The following theorem summarizes these results.

Theorem 1.6. Let G be a finite almost simple group in a primitive faithful nonstandard action with socle G_0 . Then, $b(G) \leq 7$, with equality if and only if G is the Mathieu group M_{24} in its natural action of degree 24. Moreover, b(G) = 6 if and only if one of the following holds:

- (1.1) $G = M_{23}$ in its natural action of degree 23,
- (1.2) $G = Co_3$ in its action on the right cosets of a maximal subgroup isomorphic to McL.2,
- (1.3) $G = Co_2$ in its action on the right cosets of a maximal subgroup isomorphic to U₄(3).2.2,
- (1.4) $G = Fi_{22}.2$ in its action on the right cosets of a maximal subgroup isomorphic to 2.U₆(2).2,
- (1.5) $G_0 = E_7(q)$ in its action on the cosets of a parabolic subgroup labeled P_7 ,
- (1.6) $G_0 = E_6(q)$ in its action on the cosets of a parabolic subgroup labeled P_1 or P_6 .

Moreover, if G_0 is a classical group, then either $b(G) \leq 4$ or $G = U_6(2).2$ with stabilizer $U_4(3).2^2$.

The labeling of the parabolic subgroups in part (1.5) and (1.6) follows [3].

1.2. Structure of the paper. Let G be an almost simple primitive group having socle a simple group of Lie type G_0 . When G_0 is classical, let V be the natural module of the covering group of G_0 and let p be the characteristic of the underlying field. From Definition 1.4^f and Theorem 1.6, the proof of Theorem 1.2 reduces to the following cases:

- (1.1) G_0 is a classical group acting on totally singular subspaces of V,
- (1.2) $G_0 = \text{PSL}_d(q)$ acting on pairs of subspaces of V, see (3.2) and (3.3) in Section 3.1 for more details,
- (1.3) $G_0 = \operatorname{Sp}_{2m}(q)', q$ is even and G_0 is acting on the right cosets of $\operatorname{O}_{2m}^{\pm}(q),$
- (1.4) G_0 is classical acting on non-degenerate subspaces of V,
- (1.5) G_0 is orthogonal of even characteristic and G_0 is acting on non-singular 1-subspaces of V,
- (1.6) $G_0 = \text{Sp}_4(2^a)$, G contains a graph automorphism and G_0 is acting on the cosets of a local subgroup of type $[q^4] : C_{q-1}^2$, see [2, Table 8.14],
- (1.7) $G_0 = P\Omega_8^+(q)$, G contains a triality graph automorphism and G_0 is acting on the cosets of a local subgroup of type

$$[q^{11}]: \left[\frac{q-1}{d}\right]^2 \cdot \frac{1}{d} \mathrm{GL}_2(q) \cdot d^2,$$

where $d = \gcd(2, q - 1)$, see [2, Table 8.50],

- (1.8) $G_0 = P\Omega_8^+(q)$, G contains a triality graph automorphism and G_0 is acting on the cosets of a subgroup isomorphic to $G_2(q)$, see [2, Table 8.50],
- (1.9) $G_0 = E_7(q)$ in its action on the cosets of a parabolic subgroup labeled P_7 ,
- (1.10) $G_0 = E_6(q)$ in its action on the cosets of a parabolic subgroup labeled P_1 or P_6 .

^fSee also the comment following Definition 1.4 and [8, page 549].

The structure of the paper is straightforward. In Section 2 we collect some basic remarks and observations. The proof of Theorem 1.2 is divided by cases (1.1) - (1.10) as follow.

We prove case (1.1) and case (1.2) in Section 3. We prove case (1.3) in Section 4. We prove case (1.4) in Section 5. We prove case (1.5) in Section 6. We prove case (1.6) in Section 7. We prove case (1.7) and (1.8) in Section 8. We prove cases (1.9) and (1.10) in Section 9.

2. Preliminaries

Let G be a permutation group on Ω . Given $\omega_1, \ldots, \omega_\ell \in \Omega$, we say that the sequence $(\omega_1, \ldots, \omega_\ell)$ is *irredundant* if

$$G > G_{\omega_1} > G_{\omega_1,\omega_2} > \cdots > G_{\omega_1,\omega_2,\dots,\omega_\ell}.$$

Lemma 2.1. Let G be a permutation group on Ω and let $(\omega_1, \ldots, \omega_\ell)$ and $(\omega'_1, \ldots, \omega'_{\ell'})$ be irredundant with

$$G_{\omega_1,\ldots,\omega_\ell} = G_{\omega'_1,\ldots,\omega'_{\ell'}}.$$

if $\ell \neq \ell'$, then G is not IBIS.

Proof. As $G_{\omega_1,\ldots,\omega_\ell} = G_{\omega'_1,\ldots,\omega'_{\ell'}}$, we may extend the irredundant sequences $(\omega_1,\ldots,\omega_\ell)$ and $(\omega'_1,\ldots,\omega'_{\ell'})$ using the same set of points to obtain irredundant bases for G. \Box

The following results are just simple observations.

Lemma 2.2. Let G be a permutation group on Ω and let H be a subgroup of G. If $(\omega_1, \ldots, \omega_\ell)$ is irredundant for H, then it is irredundant for G.

Lemma 2.3. Let A, G and B be permutation groups on Ω with $A \leq G \leq B$. If A has an irredundant base of length ℓ_A and B has an irredundant base of length ℓ_B with $\ell_A > \ell_B$, then G is not IBIS.

3. Primitive action on totally-singular subspaces: case (1.1) and case (1.2)

We divide the proof of Theorem 1.2 for case (1.1) depending on whether G_0 is linear, unitary, symplectic or orthogonal. There are strong similarities in the arguments we use to deal with each case. However, each family has its own peculiar behavior and hence we have preferred to divide the proofs in separate cases.

In this and in later sections, we denote the projective image of a matrix with square brackets.

3.1. Linear groups. Let $d \ge 2$ be a positive integer, let q be a prime power (that is, $q = p^f$ for some prime number p and some positive integer f), let $G_0 = \text{PSL}_d(q)$ and let G be an almost simple group with socle G_0 . In particular, $G_0 \le G \le \text{Aut}(G_0)$, where $\text{Aut}(G_0) = \text{P}\Gamma L_d(q)$ when d = 2 and

$$\operatorname{Aut}(G_0) = \operatorname{P}\Gamma \operatorname{L}_d(q) \rtimes \langle \iota \rangle = \operatorname{PSL}_d(q) \rtimes (\operatorname{Aut}(\mathbb{F}_q) \times \langle \iota \rangle),$$

when d > 2, where ι is a graph automorphism of G_0 and $\operatorname{Aut}(\mathbb{F}_q)$ is the group of field automorphisms of \mathbb{F}_q .

We now describe the primitive subspace actions. We let $V = \mathbb{F}_q^d$ be the *d*-dimensional vector space over the finite field \mathbb{F}_q of cardinality q and let $k \in \{1, \ldots, d-1\}$. Then,

IBIS LARGE BASE

- (3.1) we let Ω_k be the collection of all k-dimensional subspaces of V, here $\mathrm{PSL}_d(q) \trianglelefteq G \le \mathrm{P}\Gamma\mathrm{L}_d(q)$,
- (3.2) for $1 \leq k < d/2$, we let $\Omega_k^1 = \{\{W, U\} \mid W \in \Omega_k, U \in \Omega_{d-k}, V = W \oplus U\}$, here $\text{PSL}_d(q) \leq G \leq \text{P}\Gamma L_d(q)$,
- (3.3) for $1 \leq k < d/2$, we let $\Omega_k^2 = \{\{W, U\} \in W \in \Omega_k, U \in \Omega_{d-k}, W \leq U\},$ here $\text{PSL}_d(q) \leq G \leq \text{P}\Gamma L_d(q).$

Since our argument is inductive, we need to discuss in detail the cases when d is small. The case d = 2 is quite special and it was already discussed in [31]. Here, we recall some basic facts from [31, Example 3.1]. Let G be an almost simple group with socle $PSL_2(q)$ acting on the q + 1 points of the projective line. In particular,

$$\operatorname{PSL}_2(q) \trianglelefteq G \le \operatorname{P}\Gamma\operatorname{L}_2(q).$$

Set $r = |G: G \cap PGL_2(q)|$. Recall that $q = p^f$.

Lemma 3.1 (Example 3.1, [31]). The group G is IBIS if and only if r = 1, or r is prime and $r \mid f$.

Suppose now $d \ge 3$. Let $V = \mathbb{F}_q^d$ and let e_1, \ldots, e_d be an \mathbb{F}_q -basis of V.

3.1.1. Case (3.1). We first discuss the case k = 1.

Lemma 3.2. Let G be a group as in (3.1) acting on Ω_k with k = 1. Then G is IBIS if and only if q = 2.

Proof. When q = 2, $G = GL_d(2)$ and G is the group described in [31, Example 3.3]; hence G is IBIS. Therefore, we may suppose that q > 2.

Let $\omega_1 = \omega'_1 = \langle e_1 \rangle$, $\omega_2 = \omega'_2 = \langle e_2 \rangle$, $\omega_3 = \omega'_4 = \langle e_3 \rangle$, $\omega'_3 = \langle e_1 + e_2 \rangle$, $\omega_4 = \omega'_5 = \langle e_1 + e_2 + e_3 \rangle$. As q > 2, with a simple computation, we see that $(\omega_1, \ldots, \omega_4)$ and $(\omega'_1, \ldots, \omega'_5)$ are irredundant chains for $G_0 = \text{PSL}_d(q)$. Therefore, by Lemma 2.2, these are irredundant chains for G.

We show that

$$G_{\omega_1,\ldots,\omega_4} = G_{\omega_1',\ldots,\omega_5'},$$

from which the proof immediately follows from Lemma 2.1.

As $\{\omega_1, \ldots, \omega_4\} \subseteq \{\omega'_1, \ldots, \omega'_5\}$, we deduce $G_{\omega_1, \ldots, \omega_4} \geq G_{\omega'_1, \ldots, \omega'_5}$. Thus, let $g \in G_{\omega_1, \ldots, \omega_4}$. In particular, there exists a linear transformation $x \in \operatorname{GL}(V)$ and a Galois automorphism $\sigma \in \operatorname{Aut}(\mathbb{F}_q)$ such that g is the permutation induced by σx on Ω_1 . Thus

$$e_1^{\sigma} x = e_1 x \in \langle e_1 \rangle = \omega_1,$$

$$e_2^{\sigma} x = e_2 x \in \langle e_2 \rangle = \omega_2,$$

$$e_3^{\sigma} x = e_3 x \in \langle e_3 \rangle = \omega_3,$$

$$(e_1 + e_2 + e_3)^{\sigma} x = (e_1 + e_2 + e_3) x \in \langle e_1 + e_2 + e_3 \rangle = \omega_4.$$

In particular, there exist $\lambda_1, \lambda_2, \lambda_3, \lambda_4 \in \mathbb{F}_q$ with

$$e_1 x = \lambda_1 e_1,$$

$$e_2 x = \lambda_2 e_2,$$

$$e_3 x = \lambda_3 e_3,$$

$$(e_1 + e_2 + e_3) x = \lambda_4 (e_1 + e_2 + e_3).$$

This linear system of equations gives $\lambda_1 = \lambda_2 = \lambda_3 = \lambda_4$. This shows that x acts as a scalar on the subspace $W = \langle e_1, e_2, e_3 \rangle$ of V. Since $\omega'_1, \ldots, \omega'_5$ are 1-dimensional

subspaces of W and since $e_1, e_2, e_1 + e_2, e_3, e_1 + e_2 + e_3$ have coefficients in the prime field \mathbb{F}_p , we deduce $g \in G_{\omega'_1, \dots, \omega'_5}$.

Lemma 3.3. Let $G = GL_d(2)$ be acting on Ω_k with k = 2. Then G is not IBIS.

Proof. As k = 2, we have $d \ge 4$. Let $W = \langle e_1, e_2, e_3, e_4 \rangle$ and let G_W be the setwise stabilizer of W in G. We let

$$\begin{aligned} \omega_1 &= \langle e_1, e_2 \rangle, \, \omega_2 = \langle e_3, e_4 \rangle, \, \omega_3 = \langle e_1 + e_3, e_2 + e_4 \rangle, \, \omega_4 = \langle e_1, e_3 \rangle, \, \omega_5 = \langle e_2, e_4 \rangle, \\ \omega_1' &= \langle e_1, e_2 \rangle, \, \omega_2' = \langle e_1, e_3 \rangle, \, \omega_3' = \langle e_2, e_4 \rangle, \, \omega_4' = \langle e_3, e_4 \rangle. \end{aligned}$$

A computation shows that $(\omega_1, \omega_2, \omega_3, \omega_4, \omega_5)$ and $(\omega'_1, \omega'_2, \omega'_3, \omega'_4)$ are irredundant chains with

$$G_{\omega_1,\omega_2,\omega_3,\omega_4,\omega_5} = G_W = G_{\omega_1',\omega_2',\omega_3',\omega_4'}$$

Therefore, by Lemma 2.1, G in its action on Ω_2 is not IBIS.

$$\square$$

The action of $G = \operatorname{GL}_4(2)$ on the 2-subspaces of $V = \mathbb{F}_2^4$ is actually very interesting; Lemma 3.3 shows that G has irredundant bases of size 4 and 5, however, it can be verified with the auxiliary help of a computer that every minimal base^g of G has cardinality 4. In [20], these permutation groups were christianised with MiBIS, from *Minimal Bases of Invariant Sizes*.

Using Lemmas 3.2 and 3.3 we can now deal with the case $k \ge 2$.

Lemma 3.4. Let G be a group as in (3.1) acting on Ω_k with $k \ge 2$. Then G is not IBIS.

Proof. When q > 2, let $W = \langle e_1, \ldots, e_{k-1} \rangle$ and, when q = 2, let $W = \langle e_1, \ldots, e_{k-2} \rangle$. Now, let $\Delta = \{ \omega \in \Omega_k \mid W \leq \omega \}$, let G_W be the stabilizer of W in G, let G_Δ be the setwise stabilizer of Δ in G and let H be the permutation group induced by G_W on its action on Δ . We claim that

(1)
$$G_W = G_{\Delta}$$

Indeed, if $g \in G_W$, then g fixes setwise W and hence g permutes the k-dimensional subspaces of V containing W, that is, g fixes setwise Δ . Thus $g \in G_{\Delta}$. Conversely, if $g \in G_{\Delta}$, then g fixes setwise

$$\bigcap_{\delta \in \Delta} \delta = W$$

and hence $g \in G_W$.

Observe that when q = 2, from Lemma 3.3, we may assume that $k \ge 3$.

When q > 2, H is an almost simple group with socle $\text{PSL}_{d-k+1}(q)$ and the action of H on Δ is permutation isomorphic to the natural action of H on the 1-dimensional subspaces of $V/W \cong \mathbb{F}_q^{d-k+1}$; moreover, $d-k+1 \ge 3$ because $2 \le k \le d/2$. When q = 2, $H = \text{GL}_{d-k+2}(2)$ and the action of H on Δ is permutation isomorphic to the natural action of H on the 2-dimensional subspaces of $V/W \cong \mathbb{F}_q^{d-k+2}$; moreover, $d-k+2 \ge 4$ because $3 \le k \le d/2$.

Thus, from Lemmas 3.2 and 3.3, the action of H on Δ is not IBIS. Therefore, in its action on Δ the group H has two irredundant bases $(\omega_1, \ldots, \omega_\ell)$ and $(\omega'_1, \ldots, \omega'_{\ell'})$

^gRecall that a minimal base is a base for G, for which none of its proper subsets is again a base.

with $\ell' \neq \ell$. We claim that

(2)
$$\bigcap_{i=1}^{\ell} \omega_i = W = \bigcap_{i=1}^{\ell'} \omega'_i.$$

We only show that $W = \bigcap_{i=1}^{\ell} \omega_i$, because the argument for showing $W = \bigcap_{i=1}^{\ell'} \omega_i'$ is similar. As $\omega_i \in \Delta$ for each i and as $W \leq \omega_i$, we have $W \leq \bigcap_{i=1}^{\ell} \omega_i$. Suppose first q > 2. Since $\dim_{\mathbb{F}_q} W = k - 1$ and $\dim_{\mathbb{F}_q} \omega_i = k$, we have $W \neq \bigcap_{i=1}^{\ell} \omega_i$ only if $\ell = 1$. However, when $\ell = 1$, we get a contradiction because $\mathrm{PSL}_{d-k+1}(q)$ in its action on the projective points cannot have a base of size $\ell = 1$, and hence neither can H. Suppose now q = 2. Let $W' = \bigcap_{i=1}^{\ell} \omega_i$. Since $\dim_{\mathbb{F}_q} W = k - 2$ and $\dim_{\mathbb{F}_q} \omega_i = k$, we have $W \neq W'$ only when $\dim_{\mathbb{F}_q} W' = k$ or $\dim_{\mathbb{F}_q} W' = k - 1$. In the first case, we have $\ell = 1$ and we get a contradiction because $H = \mathrm{GL}_{d-k+2}(2)$ in its action on the 2-dimensional subspaces of \mathbb{F}_q^{d-k+2} cannot have a base of size $\ell = 1$. Assume then $\dim_{\mathbb{F}_q} W' = k - 1$. Without loss of generality, replacing the standard \mathbb{F}_q -basis of V if necessary, we have $W' = \langle W, e_{k-1} \rangle$. Let $\omega = \langle W, e_k, e_{k+1} \rangle$ and let $x \in \mathrm{GL}_d(q)$ with $e_k x = e_{k-1} + e_k$ and $e_i x = e_i, \forall i \neq k$. Since $W \leq \omega$ and $\dim_{\mathbb{F}_q} \omega = k$, we have $\omega \in \Delta$. Since x acts trivially on V/W' and since $W' \leq \omega_i$ for each i, we have $x \in G_{\omega_1, \dots, \omega_\ell}$. Since $\omega_1, \dots, \omega_\ell$ is an irredundant base for the action of H on the 2-dimensional subspaces of V/W, we deduce that x fixes each 2-dimensional subspace of V/W. However,

$$\omega^x = \langle Wx, e_k x, e_{k+1} x \rangle = \langle W, e_{k-1} + e_k, e_{k+1} \rangle \neq \langle e_1, \dots, e_{k-2}, e_k, e_{k+1} \rangle = \omega,$$

which is a contradiction. This concludes the proof of (2).

From (1) and (2), we have $G_{\omega_1,\ldots,\omega_{\ell}} \leq G_{\Delta}$ and $G_{\omega'_1,\ldots,\omega'_{\ell'}} \leq G_{\Delta}$. Moreover, since $\omega_1,\ldots,\omega_{\ell}$ and $\omega'_1,\ldots,\omega'_{\ell'}$ are irredundant bases for H, we get

$$G_{\omega_1,\dots,\omega_\ell} = G_{(\Delta)} = G_{\omega_1',\dots,\omega_{\ell'}'},$$

where as usual $G_{(\Delta)}$ is the pointwise stabiliser of Δ . Therefore, by Lemma 2.1, G is not IBIS in its action on Ω_k .

3.1.2. Case (3.2). We set some notation that we use in this section and in Section 3.1.3. As above, we let e_1, \ldots, e_d be the canonical basis of $V = \mathbb{F}_q^d$; we think of e_1, \ldots, e_d as d-tuples of row vectors with coefficients in \mathbb{F}_q . We let e_1^T, \ldots, e_d^T be the dual basis and we think of e_1^T, \ldots, e_d^T as d-tuples of column vectors with coefficients in \mathbb{F}_q . Given $x \in \operatorname{GL}_d(q), v \in V$ and $v^T \in V^*$, the matrix x acts on V and V^* by setting $v \mapsto vx$ and $w^T \mapsto x^{-1}w^T$. Moreover, we let ι be the graph automorphism acting on $V \cup V^*$ by setting $v^{\iota} = v^T$ and $(v^T)^{\iota} = v$. We have

$$v^{\iota x \iota} = (v^T)^{x \iota} = (x^{-1} v^T)^{\iota} = v(x^{-1})^T,$$

where x^T is the transposed of the matrix x. Since this holds for every $v \in V$, we have $\iota^2 = 1$ and $\iota x \iota = (x^{-1})^T$.

Let $v \in V$ and $w^T \in V^*$ with $v \neq 0 \neq w^T$. Using the notation above, $\langle w^T \rangle$ is an hyperplane of V and the vector v belongs to the hyperplane $\langle w^T \rangle$ if and only if $v \cdot w^T = 0$.

Lemma 3.5. Let G be a group as in (3.2) acting on Ω_k^1 . Then G is not IBIS.

Proof. Let $U = \langle e_4, \ldots, e_{k+2} \rangle$ and let $W = \langle e_{k+3}, \ldots, e_d \rangle$. There is a slight abuse of notation here: when d = 3, U = W = 0; when k = 1, U = 0.

Let

$$\begin{split} &\omega_1 = \omega_1' = \{ \langle e_1, U \rangle, \langle e_2, e_3, W \rangle \}, \\ &\omega_2 = \omega_2' = \{ \langle e_2, U \rangle, \langle e_1, e_3, W \rangle \}, \\ &\omega_3' = \{ \langle e_1 + e_2, U \rangle, \langle e_1, e_3, W \rangle \}, \\ &\omega_3 = \omega_4' = \{ \langle e_1 + e_2 + e_3, U \rangle, \langle e_1, e_2, W \rangle \}. \end{split}$$

When $q \neq 2$ and when $(d, q) \neq (3, 4)$, a simple computation shows that $(\omega_1, \omega_2, \omega_3)$ and $(\omega'_1, \omega'_2, \omega'_3, \omega'_4)$ are irredundant chains for $\text{PSL}_d(q)^{\text{h}}$. Moreover,

$$\operatorname{PSL}_{d}(q)_{\omega_{1},\omega_{2},\omega_{3}} = \operatorname{PSL}_{d}(q)_{\omega_{1}',\omega_{2}',\omega_{3}',\omega_{4}'} = \left\{ \begin{bmatrix} I_{3} & 0 & 0\\ 0 & X & 0\\ 0 & 0 & Y \end{bmatrix} \middle| \begin{array}{c} X \in \operatorname{GL}_{k-1}(q), \\ Y \in \operatorname{GL}_{d-k-2}(q), \\ \det(X)\det(Y) = 1 \end{array} \right\},$$

where I_3 is the 3×3 identity matrix. Therefore, by Lemma 2.2, $(\omega_1, \omega_2, \omega_3)$ and $(\omega'_1, \omega'_2, \omega'_3, \omega'_4)$ are irredundant chains for G.

We claim that

(3)
$$G_{\omega_1,\omega_2,\omega_3} = G_{\omega'_1,\omega'_2,\omega'_3,\omega'_4}$$

As $\{\omega_1, \omega_2, \omega_3\} = \{\omega'_1, \omega'_2, \omega'_4\}$, it suffices to show that $G_{\omega_1, \omega_2, \omega_3}$ fixes ω'_3 . Let $A = \Pr \Gamma_d(q) \rtimes \langle \iota \rangle$. Using the fact that ι and $\operatorname{Aut}(\mathbb{F}_q)$ fix ω_1 and ω_2 , it is easy to verify that

$$A_{\omega_1,\omega_2} = \left\{ \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & a & 0 & 0 & 0 \\ 0 & 0 & c & 0 & x \\ 0 & 0 & 0 & X & 0 \\ 0 & 0 & 0 & 0 & Y \end{bmatrix} x^{\varepsilon} | \begin{array}{c} a, b \in \mathbb{F}_q \setminus \{0\}, x \in \mathbb{F}_q^{d-k-2} \\ X \in \operatorname{GL}_{k-1}(q), Y \in \operatorname{GL}_{d-k-2}(q), \\ \tau \in \operatorname{Aut}(\mathbb{F}_q), \varepsilon \in \{0,1\} \end{array} \right\}.$$

From this, it follows that

$$A_{\omega_1,\omega_2,\omega_3} = \left\{ \begin{bmatrix} I_3 & 0 & 0\\ 0 & X & 0\\ 0 & 0 & Y \end{bmatrix} \mid \begin{array}{c} X \in \mathrm{GL}_{k-1}(q), \\ Y \in \mathrm{GL}_{d-k-2}(q) \end{array} \right\} \rtimes \mathrm{Aut}(\mathbb{F}_q).$$

From this explicit calculation, it is immediate that each element of $A_{\omega_1,\omega_2,\omega_3}$ fixes ω'_3 and hence, each element of $G_{\omega_1,\omega_2,\omega_3}$ fixes ω'_3 . By (3) and Lemma 2.1, G is not IBIS.

It remains to deal with the cases q = 2 and (d, q) = (3, 4). When (d, q) = (3, 4), PSL₃(4) $\leq G \leq PSL_3(4) \rtimes \langle \iota \rangle$ and $|\Omega| = 336$. It can be verified directly, or with the auxiliary help of a computer, that G is not IBIS.

Assume q = 2. Thus $G = \operatorname{GL}_d(2) \rtimes \langle \iota \rangle$. Let $U = \langle e_3, \ldots, e_{k+1} \rangle$ and $W = \langle e_{k+2}, \ldots, e_d \rangle$. As above, there is an abuse of notation here: when d = 3, U = 0. We define

$$\begin{split} \omega_1 &= \omega_1' = \{ \langle e_1, U \rangle, \langle e_2, W \rangle \}, \\ \omega_2 &= \omega_3' = \{ \langle e_2, U \rangle, \langle e_1 + e_2, W \rangle \}, \\ \omega_2' &= \{ \langle e_2, U \rangle, \langle e_1, W \rangle \}. \end{split}$$

We claim that

$$G_{\omega_1,\omega_2} = G_{\omega_1',\omega_2',\omega_3'},$$

10

^hThe condition $q \neq 2$ guarantees that $(\omega_1, \omega_2, \omega_3)$ is an irredundant chain and the condition $(d, q) \neq (3, 4)$ guarantees that $(\omega'_1, \omega'_2, \omega'_3, \omega'_4)$ is also an irredundant chain.

from which it follows, thanks to Lemma 2.1, that G is not IBIS. It is immediate to verify that

$$G_{\omega'_1,\omega'_2} = \left\{ \begin{bmatrix} I_2 & 0 & 0\\ 0 & X & 0\\ 0 & 0 & Y \end{bmatrix} \mid X \in \mathrm{GL}_{k-1}(2), Y \in \mathrm{GL}_{d-k-1}(2) \right\} \rtimes \langle \iota \rangle$$

and

$$G_{\omega_1',\omega_2',\omega_3'} = \left\{ \left[\begin{array}{ccc} I_2 & 0 & 0\\ 0 & X & 0\\ 0 & 0 & Y \end{array} \right] \mid X \in \mathrm{GL}_{k-1}(2), Y \in \mathrm{GL}_{d-k-1}(2) \right\}.$$

It is also clear that

$$\operatorname{GL}_{d}(2)_{\omega_{1},\omega_{2}} = \left\{ \begin{bmatrix} I_{2} & 0 & 0\\ 0 & X & 0\\ 0 & 0 & Y \end{bmatrix} \mid X \in \operatorname{GL}_{k-1}(2), Y \in \operatorname{GL}_{d-k-1}(2) \right\} = G_{\omega_{1}',\omega_{2}',\omega_{3}'}.$$

Suppose, arguing by contradiction, that $\iota g \in G_{\omega_1,\omega_2}$, for some $g \in \mathrm{GL}_d(2)$. Now,

$$\begin{split} \omega_1^\iota &= \{ \langle e_1, U \rangle, \langle e_2, W \rangle \}^\iota = \{ \langle e_2, W \rangle, \langle e_1, U \rangle \} = \omega_1, \\ \omega_2^\iota &= \{ \langle e_2, U \rangle, \langle e_1 + e_2, W \rangle \}^\iota = \{ \langle e_1, W \rangle, \langle e_1 + e_2, U \rangle \} \end{split}$$

From this and from the fact that ιg fixes ω_1, ω_2 , we deduce

$$\langle e_1, U \rangle^g = \langle e_1, U \rangle, \langle e_2, W \rangle^g = \langle e_2, W \rangle, \langle e_1 + e_2, U \rangle^g = \langle e_2, U \rangle, \langle e_1, W \rangle^g = \langle e_1 + e_2, W \rangle.$$

From this it follows that g fixes the subspaces $\langle e_1, e_2 \rangle$, U and W. Now, the first equality implies $e_1^g = e_1$ and the forth equality implies $e_1^g = e_2$, which is a contradiction.

3.1.3. Case (3.3). In this section, we deal with case (3.3). This case is easier to handle.

Lemma 3.6. Let G be a group as in (3.3) acting on Ω_k^2 . Then G is not IBIS.

Proof. Let $A = \Pr L_d(q) \rtimes \langle \iota \rangle$. Let $U = \langle e_4, \ldots, e_{k+2} \rangle$ and $W = \langle e_4, \ldots, e_{d-k+1} \rangle$. Observe that $U \leq W$, because $k+2 \leq d-k+1$ (recall k < d/2). Moreover, dim U = k-1 and dim W = d-k-2. As in the previous lemmas, there is a slight abuse of notation here. Indeed, when k = 1, U = 0. Now, let

$$\begin{split} \omega_1 &= \omega_1' = \{ \langle e_1, U \rangle, \langle e_1, e_2, W \rangle \}, \\ \omega_2 &= \omega_2' = \{ \langle e_1, U \rangle, \langle e_1, e_3, W \rangle \}, \\ \omega_3 &= \{ \langle e_2, U \rangle, \langle e_1, e_2, W \rangle \}, \\ \omega_4 &= \omega_3' = \{ \langle e_2, U \rangle, \langle e_2, e_3, W \rangle \}. \end{split}$$

Clearly, $G_{\omega_1,\omega_2} = G_{\omega'_1,\omega'_2} \leq \Pr \Gamma_d(q)$, because ω_1 and ω_2 share the same k-dimensional subspace. From $G_{\omega'_1,\omega'_2,\omega'_3} \leq \Pr \Gamma_d(q)$, it follows that $G_{\omega'_1,\omega'_2,\omega'_3}$ fixes ω_3 and hence

(4)
$$G_{\omega_1',\omega_2',\omega_3'} = G_{\omega_1,\omega_2,\omega_3,\omega_4}.$$

Now, consider the matrices

$$g_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & I_{d-3} \end{bmatrix}, g_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & I_{d-3} \end{bmatrix}, g_3 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & I_{d-3} \end{bmatrix}.$$

Observe $g_1, g_2, g_3 \in \text{PSL}_d(q) \leq G$. Moreover, $g_1 \in G_{\omega_1} \setminus G_{\omega_1, \omega_2}, g_2 \in G_{\omega_1, \omega_2} \setminus$ $G_{\omega_1,\omega_2,\omega_3}, g_3 \in G_{\omega_1,\omega_2,\omega_3} \setminus G_{\omega_1,\omega_2,\omega_3,\omega_4}$. Therefore, $(\omega_1,\omega_2,\omega_3,\omega_4)$ and $(\omega'_1,\omega'_2,\omega'_3)$ are irredundant chains for G. Now, from (4) and Lemma 2.1, this shows that G is not IBIS.

We conclude this section summing up the main result.

Proposition 3.7. Let G be an almost simple primitive group having socle $PSL_d(q)$ in a subspace action. If G is IBIS, then either

- q = 2 and $G = \operatorname{GL}_d(2)$ is acting on the non-zero vectors of \mathbb{F}_2^d , or
- d = 2 and $|G: G \cap PGL_2(q)|$ is either 1 or a prime number.

3.2. Unitary groups. Let d be a positive integer and $q = p^{f}$, with p a prime number. Let $G_0 = \text{PSU}_d(q)$ and let G be an almost simple group having socle G_0 . In particular, $G_0 \leq G \leq \operatorname{Aut}(G_0) = \operatorname{PGU}_d(q) \rtimes \operatorname{Aut}(\mathbb{F}_{q^2})$. In the following, we set $A = \operatorname{Aut}(\operatorname{PSU}_d(q)).$

Lemma 3.8. There exists $\alpha \in \mathbb{F}_{q^2}$ with $\alpha + \alpha^q + 1 = 0$ such that the orbit of α under the action of $\operatorname{Aut}(\mathbb{F}_{q^2})$ has cardinality 2f.

Proof. Let

$$S = \{ \alpha \in \mathbb{F}_{q^2} : \alpha + \alpha^q + 1 = 0 \}.$$

Consider the trace map $\operatorname{Tr} : \mathbb{F}_{q^2} \to \mathbb{F}_q$ defined by $x \mapsto x + x^q$. Since Tr is a surjective \mathbb{F}_q -linear mapping, we deduce that

$$(5) |S| = q.$$

Recall that $\operatorname{Aut}(\mathbb{F}_{q^2})$ is cyclic of order 2f. Let $2f = r_1^{l_1} \dots r_t^{l_t}$ be the factorization of 2f in prime numbers with $2 = r_1 < r_2 < \cdots < r_t$. The elements of S not lying on a regular orbit under the action of $\operatorname{Aut}(\mathbb{F}_{q^2})$ are fixed by some element of $\operatorname{Aut}(\mathbb{F}_{q^2})$ of prime order r_i , for some $i \in \{1, \ldots, t\}$. Actually, using the fact that this group is cyclic, we deduce that the elements of S fixed by some automorphism of order r_i are the elements of the set

$$S_i = S \cap \mathbb{F}_{q^{2/r_i}} = \{ \alpha \in S \mid \alpha \in \mathbb{F}_{q^{2/r_i}} \}.$$

Suppose i = 1 and let $\alpha \in S_1$. Then $\alpha^q = \alpha$ because $\alpha \in \mathbb{F}_q$. Therefore $0 = \alpha + \alpha^q + 1 = 2\alpha + 1$ and hence $|S_1| \leq 1^i$

Suppose i > 1 and let $\alpha \in S_i$. Since r_i is odd and $\alpha \in \mathbb{F}_{q^{2/r_i}}^{j}$, we deduce $\alpha^q = \alpha^{p^{f/r_i}}$. Therefore

$$0 = \alpha + \alpha^q + 1 = \alpha + \alpha^{p^{\frac{f}{r_i}}} + 1.$$

12

ⁱActually, $|S_1| = 0$ when q is even and $|S_1| = 1$ when q is odd. ^jSince r_i is odd, we may write $f = \frac{2f}{r_i} \cdot \frac{r_i - 1}{2} + \frac{f}{r_i}$. Now observe that, as $\alpha \in \mathbb{F}_{q^{2/r_i}} = \mathbb{F}_{p^{2f/r_i}}$, we get $\alpha^{p^{\frac{2f}{r_i} \cdot \frac{r_i - 1}{2}}} = \alpha$

Since this can be viewed as a polynomial equation in α of degree p^{f/r_i} , we deduce that it has at most $p^{f/r_i} = q^{1/r_i}$ solutions and hence $|S_i| \leq q^{1/r_i}$. Summing up,

(6)
$$\left| \bigcup_{i=1}^{t} S_{i} \right| \leq \sum_{i=1}^{t} |S_{i}| \leq 1 + \sum_{i=2}^{t} |S_{i}| \leq 1 + \sum_{i=2}^{t} q^{1/r_{i}} \leq 1 + (t-1)q^{1/3}$$
$$\leq 1 + (\log_{2} q - 1)q^{1/3}.$$

Using (5) and (6), we deduce that $\bigcup_i S_i$ is a proper subset of S as long as $q > 1 + (\log_2 q - 1)q^{1/3}$. Since this inequality is always satisfied, we deduce the existence of $\alpha \in S$ lying on an orbit of cardinality 2f under the action of $\operatorname{Aut}(\mathbb{F}_{q^2})$. \Box

We let $V = \mathbb{F}_{q^2}^d$ be the *d*-dimensional vector space over the finite field \mathbb{F}_{q^2} of cardinality q^2 and let $k \in \{1, \ldots, d-1\}$. Then, for $1 \leq k \leq d/2$, we let Ω_k be the collection of all *k*-dimensional totally singular subspaces of *V*.

Lemma 3.9. When d = 3 and $q \neq 2$, there exist two positive integers $\ell, \ell' \geq 2$ with $\ell \neq \ell'$ and there exist $\omega_1, \ldots, \omega_\ell, \omega'_1, \ldots, \omega'_{\ell'}$ elements of Ω_1 such that

- (ω₁,..., ω_ℓ) and (ω'₁,..., ω'_{ℓ'}) are irredudandant bases for the action of G on Ω₁,
- $\omega_1 + \cdots + \omega_\ell = V = \omega'_1 + \cdots + \omega'_{\ell'}, \ \omega_1 \cap \cdots \cap \omega_\ell = 0 = \omega'_1 \cap \cdots \cap \omega'_{\ell'}.$

In particular, the action of G on Ω_1 is not IBIS. When (d,q) = (3,2), the action of G on Ω_1 is IBIS.

Proof. Let e_1, e_2, e_3 be an \mathbb{F}_{q^2} -basis of V such that the Hermitian form defining G_0 is

$$\begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

When (d,q) = (3,2), the statement can be checked directly: G is IBIS for each choice of G with $PSU_3(2) \le G \le P\Gamma U_3(2)$. Therefore, for the rest of the proof we suppose $(d,q) \ne (3,2)$.

The set on which G acts is

$$\Omega = \{ \langle (1,0,0) \rangle \} \cup \{ (\alpha,\beta,1) : \alpha + \alpha^q + \beta \beta^q = 0 \}.$$

We give an irredundant base of size 4 for G_0 and an irredundant base of size 3 for A, hence by Lemma 2.3 G is not IBIS.

Take $\omega_1 = \langle e_1 \rangle$, $\omega_2 = \langle e_3 \rangle$ and $\omega_3 = \langle \alpha e_1 + e_3 \rangle$, where $\alpha \in \mathbb{F}_{q^2} \setminus \{0\}$ satisfies $\alpha + \alpha^q = 0$, and $\omega_4 = \langle \tilde{\alpha} e_1 + \beta e_2 + e_3 \rangle$, where $\tilde{\alpha}, \beta \in \mathbb{F}_{q^2}$ satisfy $\tilde{\alpha} + \tilde{\alpha}^q + \beta^{q+1} = 0$ and $\beta \neq 0$. A computation shows that

$$(G_0)_{\omega_1,\omega_2} = \left\{ \left[\begin{array}{rrr} a & 0 & 0 \\ 0 & a^{q-1} & 0 \\ 0 & 0 & a^{-q} \end{array} \right] \mid a \in \mathbb{F}_{q^2} \setminus \{0\} \right\}.$$

Observe that $(G_0)_{\omega_1,\omega_2} \neq 1$, because $q \neq 2$. From this description of $(G_0)_{\omega_1,\omega_2}$, we get

$$(G_0)_{\omega_1,\omega_2,\omega_3} = \left\{ \begin{bmatrix} a & 0 & 0 \\ 0 & a^{q-1} & 0 \\ 0 & 0 & a^{-q} \end{bmatrix} \mid a \in \mathbb{F}_{q^2}, a^{q+1} = 1 \right\}.$$

Observe that $(G_0)_{\omega_1,\omega_2,\omega_3}$ has order $(q+1)/\gcd(3, q+1)$. Thus $(G_0)_{\omega_1,\omega_2,\omega_3} \neq 1$, because $q \neq 2$. It is now easy to check that $(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4} = 1$ and hence $(\omega_1, \omega_2, \omega_3, \omega_4)$ is an irredundant base for G_0 . Finally, observe that $\omega_1 + \cdots + \omega_4 = V$ and $\omega_1 \cap \cdots \cap \omega_4 = 0$. We can now complete this sequence to construct an irredundant base $(\omega_1, \ldots, \omega_\ell)$ for G with $\ell \geq 4$ such that $\omega_1 + \cdots + \omega_\ell = V$ and $\omega_1 \cap \cdots \cap \omega_\ell = 0$.

Let $A = P\Gamma U_d(q)$. We now construct an irredundant base of cardinality 3 for A. From Lemma 3.8, let $\alpha \in \mathbb{F}_{q^2}$ with $\alpha + \alpha^q + 1 = 0$ and such that the orbit of α under the action of $\operatorname{Aut}(\mathbb{F}_{q^2})$ has cardinality 2f. Now take $\omega'_1 = \omega_1, \, \omega'_2 = \omega_2$ and $\omega'_3 = \langle \alpha e_1 + e_2 + e_3 \rangle$. The same computation as above shows that

$$A_{\omega_1',\omega_2'} = \left\langle \begin{bmatrix} \gamma & 0 & 0\\ 0 & \delta & 0\\ 0 & 0 & \gamma^{-q} \end{bmatrix}, \operatorname{Aut}(\mathbb{F}_{q^2} \setminus \mathbb{F}_p) : \gamma, \delta \in \mathbb{F}_{q^2} \setminus \{0\}, \delta^{q+1} = 1 \right\rangle.$$

Let $g \in A_{\omega'_1,\omega'_2}$ with g fixing ω'_3 . From the description above, we see that there exists $\gamma, \delta \in \mathbb{F}_{q^2} \setminus \{0\}$ with $\delta^{q+1} = 1$ and there exists $\sigma \in \operatorname{Aut}(\mathbb{F}_{q^2})$ with

 $\alpha^{\sigma}\gamma e_1 + \delta e_2 + \gamma^{-q} e_3 \in \omega_3' = \langle \alpha e_1 + e_2 + e_3 \rangle.$

By checking the second and the third coordinate, we deduce $\delta = \gamma^{-q}$ and hence $\gamma^{q+1} = 1$, because $\delta^{q+1} = 1$. Now, by checking the first and second coordinate, we deduce

$$\alpha^{\sigma}\gamma = \alpha\delta = \alpha\gamma^{-q} = \alpha\gamma.$$

Therefore $\alpha^{\sigma} = \sigma$. Since α lies in a regular orbit under the action of $\operatorname{Aut}(\mathbb{F}_{q^2})$, we get $\sigma = 1$. This shows that g = 1 and hence $A_{\omega'_1,\omega'_2,\omega'_3} = 1$. As before $\omega'_1 + \omega'_2 + \omega'_3 = V$, $\omega'_1 \cap \omega'_2 \cap \omega'_3 = 0$ and it is also a base for G.

We deal with the case d = 4 and k = 2.

Lemma 3.10. When d = 4, there exist two positive integers $\ell, \ell' \geq 2$ with $\ell \neq \ell'$ and there exist $\omega_1, \ldots, \omega_\ell, \omega'_1, \ldots, \omega'_{\ell'}$ elements of Ω_2 such that

- (ω₁,..., ω_ℓ) and (ω'₁,..., ω'_{ℓ'}) are irredudandant bases for the action of G on Ω₂,
- $\omega_1 + \cdots + \omega_\ell = V = \omega'_1 + \cdots + \omega'_{\ell'}, \ \omega_1 \cap \cdots \cap \omega_\ell = 0 = \omega'_1 \cap \cdots \cap \omega'_{\ell'}.$

Proof. We use the Hermitian form given by the matrix

$$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

and we use the basis e_1, e_2, f_1, f_2 of $V = \mathbb{F}_{q^2}^4$. Let G_0 be the socle of G.

Let $\omega_1 = \langle e_1, e_2 \rangle$, $\omega_2 = \langle f_1, f_2 \rangle$, $\omega_3 = \langle e_1, f_2 \rangle$, $\omega_4 = \langle e_2, f_1 \rangle$, $\omega_5 = \langle e_1 + e_2, f_1 - f_2 \rangle$ and $\omega_6 = \langle e_1 + f_2, e_2 - f_1 \rangle$. It is readily seen that $\omega_1, \ldots, \omega_6$ are 2-dimensional totally singular subspaces of V. An easy computation shows that

$$(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4} = \left\{ \begin{bmatrix} a & 0 & 0 & 0\\ 0 & b & 0 & 0\\ 0 & 0 & a^{-q} & 0\\ 0 & 0 & 0 & b^{-q} \end{bmatrix} \mid a, b \in \mathbb{F}_{q^2} \setminus \{0\} \text{ with } ab \in \mathbb{F}_q \right\}.$$

14

In particular, $|(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4}| = (q^2 - 1)(q - 1)/\gcd(4, q + 1) \neq 1$. We deduce

$$(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4,\omega_5} = \left\{ \begin{bmatrix} a & 0 & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & a^{-q} & 0 \\ 0 & 0 & 0 & a^{-q} \end{bmatrix} \mid a \in \mathbb{F}_{q^2} \setminus \{0\} \text{ with } a^2 \in \mathbb{F}_q \right\}.$$

From this it follows that,

$$|(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4,\omega_5}| = \begin{cases} q-1 & \text{if } q \text{ is even,} \\ \frac{2(q-1)}{\gcd(4,q+1)} & \text{if } q \text{ is odd.} \end{cases}$$

Thus $|(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4,\omega_5}| = 1$ if and only if $q \in \{2,3\}$. This show that $\omega_1, \ldots, \omega_5$ is an irredundant base for G_0 of cardinality 5 when $q \in \{2,3\}$, whereas $\omega_1, \ldots, \omega_6$ is an irredundant base for G_0 of cardinality 6 when $q \geq 4$.

The argument above shows that

$$A_{\omega_1,\omega_2,\omega_3,\omega_4} = \left\{ \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & b & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & b^{-q} \end{bmatrix} \mid b \in \mathbb{F}_{q^2} \setminus \{0\} \right\} \rtimes \operatorname{Aut}(\mathbb{F}_{q^2}).$$

Let $q = p^f$ and let α be a generator of the multiplicative group of the field \mathbb{F}_{q^2} and let $\omega'_5 = \langle e_1 + \alpha f_2, e_2 - \alpha f_1 \rangle$. Observe that ω'_5 is a 2-dimensional totally singular subspace of V. Let $g \in A_{\omega_1,\omega_2,\omega_3,\omega_4,\omega'_5}$. Then there exist $b \in \mathbb{F}_{q^2} \setminus \{0\}$ and $\eta \in \operatorname{Aut}(\mathbb{F}_{q^2})$ such that

$$\omega_5' = \omega_5'^g = \langle e_1 + \alpha^{\eta} b^{-q} f_2, b e_2 - \alpha^{\eta} f_1 \rangle.$$

This yields $\alpha = \alpha^{\eta} b^{-q}$ and $\alpha b = \alpha^{\eta}$. Therefore $b = b^q = \alpha^q \alpha^{-1}$ and hence $\alpha^{\eta} \alpha^{-1} \in \mathbb{F}_q$. Now, $\alpha^{\eta} = \alpha^{p^i}$, where *i* is a divisor of 2f. Since $\alpha^{\eta} \alpha^{-1} = \alpha^{p^i-1} \in \mathbb{F}_q$, we deduce that $\alpha^{(p^i-1)(p^f-1)} = 1$ and, as α is a generator of the multiplicative group of \mathbb{F}_{q^2} , we get that $p^{2f} - 1$ divides $(p^i - 1)(p^f - 1)$. If i = 2f, then α is the identity and b = 1 and hence g = 1. Assume that i < 2f. Then $i \leq f$ and hence $(p^i - 1)(p^f - 1) < p^{2f} - 1$, however this contradicts the fact that $p^{2f} - 1$ divides $(p^i - 1)(p^f - 1)$. This shows that $\omega_1, \omega_2, \omega_3, \omega_4, \omega'_5$ is an irredundant base for A (and hence for G) of cardinality 5. This implies that G is not IBIS when $q \geq 4$. Observe also that these two irredundant bases satisfy the additional requirements in the statement of the lemma.

Suppose q = 2. When $G = PSU_4(2)$, we have found irredundant bases of cardinality 4 and 5 (satisfying the requirements in the statement of this lemma) and hence G is not IBIS. Similarly, if $G \neq PSU_4(2)$, then $G = P\Gamma U_4(2)$ and we have found irredundant bases of cardinality 5 and 6 and hence G is not IBIS.

Suppose q = 3. We have found irredundant bases of cardinality 6 for $PSU_4(3)$ and hence we may use the argument above to deduce that G is not IBIS.

Lemma 3.11. When (d,q) = (5,2), there exist two positive integers $\ell, \ell' \geq 2$ with $\ell \neq \ell'$ and there exist $\omega_1, \ldots, \omega_\ell, \omega'_1, \ldots, \omega'_{\ell'}$ in Ω_2 such that

- (ω₁,..., ω_ℓ) and (ω'₁,..., ω'_{ℓ'}) are irredudandant bases for the action of G on Ω₂,
- $\omega_1 + \cdots + \omega_\ell = V = \omega'_1 + \cdots + \omega'_{\ell'}, \ \omega_1 \cap \cdots \cap \omega_\ell = 0 = \omega'_1 \cap \cdots \cap \omega'_{\ell'}.$

Proof. This follows by a direct inspection in $P\Gamma U_5(2)$ and in $PSU_5(2)$.

Proposition 3.12. When $(d,q) \neq (3,2)$, the action of G on Ω_k is not IBIS and $b(G) \geq 2$. When (d,q) = (3,2), the action of G on Ω_1 is IBIS.

Proof. We first deal with the case k = 1, this will set up the base case of an inductive argument.

When d = 3, the proof follows from Lemma 3.9; therefore, for the rest of the argument, we suppose $d \ge 4$. Assume $q \ne 2$. Let W be a non-degenerate 3-dimensional subspace of V and let

$$\Delta = \{ \omega \mid \omega \le W, \dim_{\mathbb{F}_{a^2}}(\omega) = 1 \text{ and } \omega \text{ sigular} \}.$$

Let G_{Δ} be the stabilizer of Δ in G and let H be the permutation group induced by G_{Δ} on Δ .

Observe that $G_{\Delta} = G_W$ because the span of the singular vectors of V contained in W is W itself. Moreover, H is an almost simple primitive group having socle $PSU_3(q)$ acting on the singular 1-dimensional subspaces of W. From Lemma 3.9, there exist two positive integers $\ell, \ell' \geq 2$ with $\ell \neq \ell'$ and there exist $\omega_1, \ldots, \omega_{\ell}, \omega'_1, \ldots, \omega'_{\ell'}$ totally singular 1-dimensional subspaces of W such that

- $(\omega_1, \ldots, \omega_\ell)$ and $(\omega'_1, \ldots, \omega'_{\ell'})$ are irreduced and ant bases for the action of H on Δ ,
- $\omega_1 + \cdots + \omega_\ell = W = \omega'_1 + \cdots + \omega'_{\ell'}$.

Since $\omega_1 + \cdots + \omega_\ell = W = \omega'_1 + \cdots + \omega'_{\ell'}$, we deduce that $G_{\omega_1,...,\omega_\ell}$ and $G_{\omega'_1,...,\omega'_{\ell'}}$ both stabilize W, that is, $G_{\omega_1,...,\omega_\ell}, G_{\omega'_1,...,\omega'_{\ell'}} \leq G_W = G_\Delta$. From this, since $H_{\omega_1,...,\omega_\ell} = H_{\omega'_1,...,\omega'_{\ell'}} = 1$, we deduce

$$G_{\omega_1,\dots,\omega_\ell} = G_{\omega_1',\dots,\omega_{\ell'}'}$$

As $\ell \neq \ell'$, Lemma 2.1 implies that G is not IBIS.

Finally, assume q = 2. Let $A = P\Gamma U_d(2)$. We first suppose d = 4. We use the Hermitian form given by the matrix

$$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

Let $\omega_1 = \langle e_1 \rangle$, $\omega_2 = \langle e_2 \rangle$, $\omega_3 = \langle e_3 \rangle$ and $\omega_4 = \langle e_4 \rangle$. A computation shows that $(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4}$ has order 3 and is generated by

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \alpha & 0 & 0 \\ 0 & 0 & \alpha^2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

where α is a generator of \mathbb{F}_4^* . This shows that G has an irredundant base $\omega_1, \ldots, \omega_\ell$, with $\ell = 5$ and with $V = \omega_1 + \cdots + \omega_\ell$. Similarly, let $\omega'_1 = \langle e_1 \rangle$, $\omega'_2 = \langle e_2 \rangle$, $\omega'_3 = \langle e_4 \rangle$ and $\omega'_4 = \langle e_1 + e_2 + \alpha e_3 + \alpha e_4 \rangle$. A computation shows that $A_{\omega'_1, \omega'_2, \omega'_3} = (G_0)_{\omega'_1, \omega'_2, \omega'_3} \rtimes \operatorname{Aut}(\mathbb{F}_4)$, where

$$(G_0)_{\omega_1',\omega_2',\omega_3'} = \left\{ \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & z & a^2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \mid a, z \in \mathbb{F}_4, a \neq 0, az + a^2 z^2 = 0 \right\}.$$

Let $g \in A_{\omega'_1,\omega'_2,\omega'_3,\omega'_4}$. From the description of $A_{\omega'_1,\omega'_2,\omega'_3}$, there exist $a, z \in \mathbb{F}_4$ (with $a \neq 0$ and $az + a^2 z^2 = 0$) and an automorphism η of \mathbb{F}_4 such that

$$\omega_4' = \langle e_1 + (a + \alpha^\eta z)e_2 + \alpha^\eta a^2 e_3 + \alpha^\eta e_4 \rangle.$$

This yields the linear system of equations

$$\begin{cases} 1 = a + \alpha^{\eta} z \\ \alpha = a^2 \alpha^{\eta}, \\ \alpha = \alpha^{\eta}. \end{cases}$$

Using the fact that α is a generator of \mathbb{F}_4^* , this gives a = 1, z = 0 and $\eta = 1$. This shows that $\omega'_1, \omega'_2, \omega'_3, \omega'_4$ is an irredundant base for G with $V = \omega'_1 + \omega'_2 + \omega'_3 + \omega'_4$. This has established that G is not IBIS. The case $d \ge 4$ can be deal with a similar argument as above (used for $q \ne 2$), by taking W a non-degenerate 4-dimensional subspace of V. This completes the case when k = 1.

Suppose that k > 1. Let U be a totally singular subspace of V with $\dim_{\mathbb{F}_{q^2}}(U) = k - 1$ and let

 $\Delta = \{ W \mid U \le W, \dim_{\mathbb{F}_{q^2}}(W) = k \text{ and } W \text{ totally singular} \}.$

Let H be the permutation group induced by G on Δ .

Observe that $G_{\Delta} = G_U$, where G_U is the setwise stabilizer of U in G. The Hermitial form κ defining the unitary group G_0 induces a form κ' on the \mathbb{F}_{q^2} -vector space U^{\perp}/U . Observe that $\dim_{\mathbb{F}_{q^2}}(U^{\perp}/U) = d - 2k + 2$ and that the form induced on U^{\perp}/U is non-degenerate and Hermitian. Let $\bar{V} = U^{\perp}/U$ and consider

 $\Omega = \{ W' \le \bar{V} \mid \dim_{\mathbf{F}_{q_0}}(W') = 1 \text{ and } W' \text{ is totally singular for } \kappa' \}.$

Clearly, there exists a one-to-one correspondence between the elements of Δ and the elements of Ω ; moreover, this correspondence allows to identify the action of H on Δ with the natural action on the 1-dimensional totally singular subspaces of \bar{V} .

Suppose $d-2k+2 \ge 3$ if $q \ge 3$ and $d-2k+2 \ge 4$ when q = 2. From Lemma 3.9, the action of H on Δ is not IBIS and hence there exist two irredundant sequences $(\delta_1, \ldots, \delta_\ell)$ and $(\delta'_1, \ldots, \delta'_{\ell'})$ of points of Δ such that $\ell, \ell' \ge 2, \ell \neq \ell'$ and

(7)
$$G_{\delta_1,\dots,\delta_\ell} \cap G_\Delta = G_{\delta'_1,\dots,\delta'_{\ell'}} \cap G_\Delta$$

Since $\ell \geq 2$, δ_1 and δ_2 are two distinct k-dimensional subspaces of V containing U and hence

$$\delta_1 \cap \delta_2 = U.$$

This shows that $G_{\delta_1,\delta_2} = G_{\delta_1} \cap G_{\delta_2} \leq G_U = G_{\Delta}$ and hence $G_{\delta_1,\ldots,\delta_\ell} \leq G_{\Delta}$. Since this argument applies also for ℓ' , from (7), we deduce

$$G_{\delta_1,\ldots,\delta_\ell} = G_{\delta'_1,\ldots,\delta'_{\ell'}}$$

As $\ell \neq \ell'$, from Lemma 2.1, we deduce that the action of G on Ω_k is not IBIS. It remains to consider the case $d-2k+2 \leq 2$ (that is, d=2k), and (d,q) = (2k+1,2).

Suppose d = 2k, or d = 2k + 1 and q = 2. Let U be a totally singular subspace of V with $\dim_{\mathbb{F}_{q^2}}(U) = k - 2$ and let

$$\Delta = \{ W \mid U \le W, \dim_{\mathbb{F}_{2}}(W) = k \text{ and } W \text{ totally singular} \}.$$

We may apply the argument above verbatim, with the old U replaced by this new U. Observe that $\dim(U^{\perp}/U) = 4$ when d = 2k and $\dim(U^{\perp}/U) = 5$ when

(d,q) = (2k+1,2). We deduce, using Lemma 3.10 when d = 2k and Lemma 3.11 when (d,q) = (2k+1,2), that G is not IBIS.

3.3. Symplectic groups. Let $d \ge 4$ be an even positive integer, let $q = p^f$ be a prime power with p a prime number, let $G_0 = \operatorname{PSp}_d(q)$ and let G be an almost simple group with socle G_0 . We fix a basis $e_1, \ldots, e_d, f_1, \ldots, f_d$ of $V = \mathbb{F}_q^d$ such that the symplectic form defining G_0 satisfies $(e_i, e_j) = (f_i, f_j) = 0$ and $(e_i, f_j) = 1$ if and only i = j. Let μ be a generator of the multiplicative group of the field \mathbb{F}_q and let δ be the projective image of diag $(\mu, \ldots, \mu, 1, \ldots, 1)$. In particular, $G_0 \leq G \leq \operatorname{Aut}(G_0)$ and

$$\operatorname{Aut}(G_0) = \begin{cases} \operatorname{Sp}_d(q) \rtimes \operatorname{Aut}(\mathbb{F}_q) & \text{if } q \text{ even,} \\ (\operatorname{PSp}_d(q) \rtimes \langle \delta \rangle) \rtimes \operatorname{Aut}(\mathbb{F}_q) & \text{if } q \text{ odd,} \end{cases}$$

see [28, Section 2.4]. In the following, we set $A = \operatorname{Aut}(\operatorname{PSp}_d(q))$.

For $1 \leq k \leq d/2$, we let Ω_k be the collection of all k-dimensional totally singular subspaces of V. Observe that, since we are in a symplectic space, every 1-dimensional subspace of V is singular, hence Ω_1 is the collection of all 1-dimensional subspaces.

Lemma 3.13. When d = 4 and $q \neq 2$, there exist two positive integers $\ell, \ell' \geq 2$ with $\ell \neq \ell'$ and there exist $\omega_1, \ldots, \omega_\ell, \omega'_1, \ldots, \omega'_{\ell'}$ elements of Ω_1 such that

- (ω₁,..., ω_ℓ) and (ω'₁,..., ω'_{ℓ'}) are irredudandant bases for the action of G on Ω₁,
- $\omega_1 + \dots + \omega_\ell = V = \omega'_1 + \dots + \omega'_{\ell'}, \ \omega_1 \cap \dots \cap \omega_\ell = 0 = \omega'_1 \cap \dots \cap \omega'_{\ell'}.$

In particular, the action of G on the set of the totally singular subspaces of dimension 1 is not IBIS.

When d = 4 and q = 2, G is IBIS.

Proof. When d = 4 and q = 2, the action of G is IBIS, as proved in [31]. Therefore, suppose $q \neq 2$.

For q = 3, the veracity of the lemma can be checked with the help of a computer. So we now suppose that q > 3. We recall that we are using the symplectic form given by

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}$$

and the base e_1, e_2, f_1, f_2 of $V = \mathbb{F}_q^4$.

We give an irredundant base of cardinality 6 for $G_0 = \text{PSp}_4(q)$ and an irredundant base of cardinality 5 for $A = \text{Aut}(G_0)$. From this, the result follows immediately from Lemma 2.3. Firstly, consider $\omega_1 = \langle e_1 \rangle$, $\omega_2 = \langle e_2 \rangle$, $\omega_3 = \langle f_1 \rangle$, $\omega_4 = \langle f_2 \rangle$, $\omega_5 = \langle e_1 + e_2 \rangle$, $\omega_6 = \langle e_1 + f_1 \rangle$. A computation shows that

$$(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4} = \left\{ \begin{bmatrix} a & 0 & 0 & 0 \\ 0 & b & 0 & 0 \\ 0 & 0 & a^{-1} & 0 \\ 0 & 0 & 0 & b^{-1} \end{bmatrix} \mid a, b \in \mathbb{F}_q \setminus \{0\} \right\}.$$

Now an element $g \in (G_0)_{\omega_1,\omega_2,\omega_3,\omega_4}$ fixes ω_5 if and only if $ae_1 + be_2 = \alpha(e_1 + e_2)$ for some $\alpha \in \mathbb{F}_q$. This implies that a = b.

Finally, an element $g \in (G_0)_{\omega_1,\omega_2,\omega_3,\omega_4,\omega_5}$ fixes ω_6 if and only if $ae_1 + a^{-1}f_1 =$

 $\alpha(e_1 + f_1)$ for some $\alpha \in \mathbb{F}_q$. This yields $a = a^{-1}$, that is, $a = \pm 1$. In conclusion, ω_1,\ldots,ω_6 is an irredundant base of cardinality 6 for G_0 . Moreover, the sum of these subspaces is equal to V and their intersection is equal to 0. This implies that there exists $\ell \geq 6$ and an irredundant base $(\omega_1, \ldots, \omega_\ell)$ for G with $\omega_1 + \cdots + \omega_\ell = V$ and $\omega_1 \cap \cdots \cap \omega_\ell = 0$.

We now take in account A. Let α be a generator of the multiplicative group of the field \mathbb{F}_q . Take $\omega'_1 = \omega_1, \ldots, \omega'_4 = \omega_4$ and $\omega'_5 = \langle \alpha e_1 + e_2 + \alpha f_1 + f_2 \rangle$. We deal simultaneously with the case q odd and q even, by defining $\delta = 1$ when q is even. A computation shows that

$$A_{\omega'_1,...,\omega'_4} = \left\{ \begin{bmatrix} a & 0 & 0 & 0 \\ 0 & b & 0 & 0 \\ 0 & 0 & a^{-1} & 0 \\ 0 & 0 & 0 & b^{-1} \end{bmatrix} \mid a, b \in \mathbb{F}_q \setminus \{0\} \right\} \rtimes (\langle \delta \rangle \times \operatorname{Aut}(\mathbb{F}_q)).$$

Let $g \in A_{\omega'_1,\ldots,\omega'_4}$ be an element fixing ω'_5 . Then, from the description of $A_{\omega'_1,\ldots,\omega'_4}$ above, we see that there exist $a, b \in \mathbb{F}_q \setminus \{0\}, \varepsilon \in \{0, 1\}$ and $\eta \in \operatorname{Aut}(\mathbb{F}_q)$ such that

$$\langle \alpha e_1 + e_2 + \alpha f_1 + f_2 \rangle = \omega_5' = \omega_5'^g = \langle \mu^{\varepsilon} \alpha^{\eta} a e_1 + \mu^{\varepsilon} b e_2 + \alpha^{\eta} a^{-1} f_1 + b^{-1} f_2 \rangle.$$

We obtain the following system

$$\begin{cases} \alpha c &= \mu^{\varepsilon} \alpha^{\eta} a \\ c &= \mu^{\varepsilon} b \\ \alpha c &= \alpha^{\eta} a^{-1} \\ c &= b^{-1}, \end{cases}$$

for some $c \in \mathbb{F}_q \setminus \{0\}$. The second and the forth equations give $\mu^{\varepsilon} = b^2$ and this implies $\varepsilon = 0$, because μ is a generator of the multiplicative group \mathbb{F}_q . Therefore, $b^2 = 1$ and hence $c = b \in \{1, -1\}$. The first and the third equations give $a = a^{-1}$ and hence $a \in \{1, -1\}$. Now, the first equation gives $ac\alpha = \alpha^{\eta}$ and hence $\alpha^2 = \alpha^{\eta}$ $(\alpha^2)^{\eta}$. Therefore, α^2 is invariant under η . Suppose, arguing by contradiction, that $\eta \neq 1$. Then the set of elements of \mathbb{F}_q invariant under η forms a proper subfield \mathbb{F} of \mathbb{F}_q , with $|\mathbb{F}| \leq \sqrt{q}$. On the other hand, since α is a generator of the multiplicative group of \mathbb{F}_q and since $\alpha^2 \in \mathbb{F}$, we get $|\mathbb{F}| \geq (q+1)/2$. However, the inequality $\sqrt{q} \ge (q+1)/2$ is not satisfied. This contradiction has shown that $\eta = 1$. Now, the linear system above gives $a = b = c \in \{1, -1\}$ and hence $A_{\omega'_1, \dots, \omega'_d, \omega'_5} = 1$.

We deal with the case d = 4 and k = 2. This is very similar to the proof of Lemma 3.10.

Lemma 3.14. Let d = 4. When $q \neq 2$, there exist two positive integers $\ell, \ell' > 2$ with $\ell \neq \ell'$ and there exist $\omega_1, \ldots, \omega_\ell, \omega'_1, \ldots, \omega'_{\ell'}$ elements of Ω_2 such that

- $(\omega_1, \ldots, \omega_\ell)$ and $(\omega'_1, \ldots, \omega'_{\ell'})$ are irreduced and and bases for the action of G on Ω_2 , • $\omega_1 + \dots + \omega_\ell = V = \omega'_1 + \dots + \omega'_{\ell'}, \ \omega_1 \cap \dots \cap \omega_\ell = 0 = \omega'_1 \cap \dots \cap \omega'_{\ell'}.$

In particular, G is not IBIS and $b(G) \geq 2$. When q = 2, the action of G on the totally singular 2-dimensional subspaces of V is IBIS and $b(G) \ge 2$.

Proof. We use the symplectic form given by the matrix

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}$$

and we use the basis e_1, e_2, f_1, f_2 of $V = \mathbb{F}_q^4$. Let G_0 be the socle of G and assume, for the time being, that $q \notin \{2,3\}$.

Let $\omega_1 = \langle e_1, e_2 \rangle$, $\omega_2 = \langle f_1, f_2 \rangle$, $\omega_3 = \langle e_1, f_2 \rangle$, $\omega_4 = \langle e_2, f_1 \rangle$, $\omega_5 = \langle e_1 + e_2, f_1 - f_2 \rangle$ and $\omega_6 = \langle e_1 + f_2, e_2 + f_1 \rangle$. It is readily seen that $\omega_1, \ldots, \omega_6$ are 2-dimensional totally singular subspaces of V. An easy computation shows that

$$(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4} = \left\{ \begin{bmatrix} a & 0 & 0 & 0 \\ 0 & b & 0 & 0 \\ 0 & 0 & a^{-1} & 0 \\ 0 & 0 & 0 & b^{-1} \end{bmatrix} \mid a, b \in \mathbb{F}_q \setminus \{0\} \right\}$$

In particular, $|(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4}| = (q-1)^2/\gcd(2,q-1) \neq 1$, because $q \neq 2$. We deduce

$$(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4,\omega_5} = \left\{ \begin{bmatrix} a & 0 & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & a^{-1} & 0 \\ 0 & 0 & 0 & a^{-1} \end{bmatrix} \mid a \in \mathbb{F}_q \setminus \{0\} \right\}$$

From this it follows that $|(G_0)_{\omega_1,\omega_2,\omega_3,\omega_4,\omega_5}| = (q-1)/\gcd(2,q-1) \neq 1$, because $q \neq 3$. This show that $\omega_1, \ldots, \omega_6$ is an irredundant base for G_0 of cardinality 6.

The argument above shows that

$$A_{\omega_1,\omega_2} = \left(\left\{ \begin{bmatrix} X & 0 \\ 0 & (X^{-1})^{tr} \end{bmatrix} \mid X \in \mathrm{GL}_2(q) \right\} \rtimes \langle \delta \rangle \right) \rtimes \mathrm{Aut}(\mathbb{F}_{q^2}).$$

Here by abuse of notation we are including the case q even by setting $\delta = 1$. Let $\omega'_3 = \langle e_1, e_2 + f_2 \rangle$ and observe that ω'_3 is a totally singular 2-dimensional subspace of V. Using the description of A_{ω_1,ω_2} and using the fact that δ is the projective image of diag $(\mu, \mu, 1, 1)$ with μ a generator of the multiplicative group of the field \mathbb{F}_q , it follows that

$$A_{\omega_1,\omega_2,\omega'_3} = \left\{ \begin{bmatrix} a & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & a^{-1} & 0\\ 0 & 0 & 0 & 1 \end{bmatrix} \mid a \in \mathbb{F}_q \setminus \{0\} \right\} \rtimes \operatorname{Aut}(\mathbb{F}_q).$$

Let $\omega'_4 = \langle e_1 + e_2, f_1 + f_2 \rangle$ and observe that ω'_4 is a totally singular 2-dimensional subspace of V. Clearly,

$$A_{\omega_1,\omega_2,\omega'_3,\omega'_4} = \operatorname{Aut}(\mathbb{F}_q).$$

Since $\operatorname{Aut}(\mathbb{F}_q)$ admits a regular orbit on its action on \mathbb{F}_q , the irredudant chain $\omega_1, \omega_2, \omega'_3, \omega'_4$ can be extended to an irredundant base of cardinality 5. This implies that G is not IBIS when $q \geq 4$.

Suppose q = 2. Here, $G = \text{Sp}_4(2)' \cong \text{Alt}(6)$ or $G = \text{Sp}_4(2) \cong \text{Sym}(6)$, and G has degree 15. We have verified with a computer that in both cases the group is IBIS, indeed, this example is in Table 1.

Suppose q = 3. Here, $G = PSp_4(3)$ or $G = PSp_4(3).2$, and the degree is 40. We have verified with a computer that in both cases the group is not IBIS and it admits

bases of different cardinality satisfying the additional conditions in the statement of the lemma. \square

Lemma 3.15. Let d = 6 and q = 2 and let $k \in \{2, 3\}$. There exist two positive integers $\ell, \ell' \geq 2$ with $\ell \neq \ell'$ and there exist $\omega_1, \ldots, \omega_\ell, \omega'_1, \ldots, \omega'_{\ell'}$ elements of Ω_k such that

- $(\omega_1, \ldots, \omega_\ell)$ and $(\omega'_1, \ldots, \omega'_{\ell'})$ are irreduced and and bases for the action of G on Ω_k , • $\omega_1 + \dots + \omega_\ell = V = \omega'_1 + \dots + \omega'_{\ell'}, \ \omega_1 \cap \dots \cap \omega_\ell = 0 = \omega'_1 \cap \dots \cap \omega'_{\ell'}.$

In particular, G is not IBIS on Ω_k and $b(G) \ge 2$.

Proof. This follows with a computer computation.

Proposition 3.16. When $(q,k) \neq (2,1)$ and when $(d,q,k) \neq (4,2,2)$, the action of G on Ω_k is not IBIS and $b(G) \geq 2$. When (q,k) = (2,1) and when (d,q,k) =(4, 2, 2), the action of G on Ω_{κ} is IBIS.

Proof. We first deal with the case k = 1, this will set up the base case of an inductive argument.

When d = 4, the proof follows from Lemma 3.13; therefore, for the rest of the argument, we suppose d > 4. Assume q > 2. Let W be a non-degenerate 4-dimensional subspace of V and let

$$\Delta = \{ \omega \mid \omega \leq W, \dim_{\mathbb{F}_a}(\omega) = 1 \text{ and } \omega \text{ sigular} \}.$$

Let G_{Δ} be the stabilizer of Δ in G and let H be the permutation group induced by G_{Δ} on Δ .

Observe that $G_{\Delta} = G_W$ because the span of the singular vectors of V contained in W is W itself. Moreover, H is an almost simple primitive group having socle $PSp_4(q)$ acting on the totally singular 1-dimensional subspaces of W. From Lemma 3.13, there exist two positive integers $\ell, \ell' \geq 2$ with $\ell \neq \ell'$ and $\omega_1, \ldots, \omega_\ell$, $\omega'_1, \ldots, \omega'_{\ell'}$ totally singular 1-dimensional subspaces of V such that

- $\omega_1 + \dots + \omega_\ell = W = \omega'_1 + \dots + \omega'_{\ell'}$
- $H_{\omega_1,...,\omega_\ell} = H_{\omega'_1,...,\omega'_{\ell'}}.$

Since $\omega_1 + \dots + \omega_\ell = W = \omega'_1 + \dots + \omega'_{\ell'}$, we deduce $G_{\omega_1,\dots,\omega_\ell}, G_{\omega'_1,\dots,\omega'_{\ell'}} \leq G_W = G_\Delta$. Now, since $H_{\omega_1,\ldots,\omega_\ell} = H_{\omega'_1,\ldots,\omega'_{\ell'}}$, we deduce

$$G_{\omega_1,\ldots,\omega_\ell} = G_{\omega'_1,\ldots,\omega'_{\ell'}}$$

As $\ell \neq \ell'$, Lemma 2.1 implies that G is not IBIS.

Assume now q = 2. Since each element of V is totally singular, the action of $G = \text{Sp}_d(2)$ on the totally singular 1-dimensional subspaces of V is the natural action of G on the non-zero vectors of V. This action is IBIS, see [31], and is reported in Table 1. This completes the case k = 1.

Suppose that k > 1. Let U be a totally singular subspace of V with $\dim_{\mathbb{F}_a}(U) =$ k-1 and let

$$\Delta = \{ W \mid U \le W, \dim_{\mathbb{F}_q}(W) = k \text{ and } W \text{ totally singular} \}.$$

Let H be the permutation group induced by G on Δ .

Observe that $G_{\Delta} = G_U$, where G_U is the setwise stabilizer of U in G. The symplectic form κ defining the symplectic group G_0 induces a form κ' on the \mathbb{F}_q -vector space U^{\perp}/U . Observe that $\dim_{\mathbb{F}_q}(U^{\perp}/U) = d - 2k + 2$ and that the form induced on U^{\perp}/U is non-degenerate and symplectic. Let $\overline{V} = U^{\perp}/U$ and consider

 $\Omega = \{ W' \le \bar{V} \mid \dim_{\mathbb{F}_a}(W') = 1 \text{ and } W' \text{ is totally singular for } \kappa' \}.$

Clearly, there exists a one-to-one correspondence between the elements of Δ and the elements of Ω ; moreover, this correspondence allows to identify the action of H on Δ with the natural action on the 1-dimensional totally singular subspaces of \bar{V} .

Suppose $d - 2k + 2 \ge 4$ and $q \ne 2$. From Lemma 3.13, the action of H on Δ is not IBIS and hence there exist two irredundant chains $(\delta_1, \ldots, \delta_\ell)$ and $(\delta'_1, \ldots, \delta'_{\ell'})$ of points of Δ such that $\ell, \ell' \ge 2, \ell \ne \ell'$ and

(8)
$$G_{\delta_1,\dots,\delta_\ell} \cap G_\Delta = G_{\delta'_1,\dots,\delta'_{\ell'}} \cap G_\Delta.$$

Since $\ell \geq 2$, δ_1 and δ_2 are two distinct k-dimensional subspaces of V containing U and hence

$$\delta_1 \cap \delta_2 = U$$

This shows that $G_{\delta_1,\delta_2} = G_{\delta_1} \cap G_{\delta_2} \leq G_U = G_{\Delta}$ and hence $G_{\delta_1,\ldots,\delta_\ell} \leq G_{\Delta}$. Since this argument applies also for ℓ' , from (8), we deduce

$$G_{\delta_1,\ldots,\delta_\ell} = G_{\delta'_1,\ldots,\delta'_{\ell'}}$$

As $\ell \neq \ell'$, from Lemma 2.1, we deduce that the action of G on Ω_k is not IBIS. It remains to consider the case $d - 2k + 2 \leq 2$ (that is, d = 2k), and the case q = 2.

Suppose d = 2k. Let U be a totally singular subspace of V with $\dim_{\mathbb{F}_q}(U) = k-2$ and let

$$\Delta = \{ W \mid U \le W, \dim_{\mathbb{F}_q}(W) = k \text{ and } W \text{ totally singular} \}.$$

We may apply the argument above verbatim, using Lemma 3.14 instead of Lemma 3.13, with the old U replaced by this new U. We deduce that G is not IBIS. It remains to deal with the case q = 2.

Suppose q = 2 and $k \in \{2, 3\}$. Clearly, we may suppose that $d \ge 6$. Let W be a non-degenerate 6-dimensional subspace of V and let

$$\Delta = \{ \omega \mid \omega \leq W, \dim_{\mathbb{F}_a}(\omega) = k \text{ and } \omega \text{ sigular} \}.$$

Let G_{Δ} be the stabilizer of Δ in G and let H be the permutation group induced by G_{Δ} on Δ .

Observe that $G_{\Delta} = G_W$ because the span of the k-dimensional singular subspaces of V of dimension k contained in W is W itself. Moreover, H is an almost simple primitive group having socle $PSp_4(q)$ acting on the totally singular k-dimensional subspaces of W. From Lemma 3.15, there exist two positive integers $\ell, \ell' \geq 2$ with $\ell \neq \ell'$ and $\omega_1, \ldots, \omega_\ell, \omega'_1, \ldots, \omega'_{\ell'}$ totally singular k-dimensional subspaces of V such that

• $\omega_1 + \dots + \omega_\ell = W = \omega'_1 + \dots + \omega'_{\ell'}$

• $H_{\omega_1,\ldots,\omega_\ell} = H_{\omega_1,\ldots,\omega'_{\ell'}}.$

Since $\omega_1 + \cdots + \omega_\ell = W = \omega'_1 + \cdots + \omega'_{\ell'}$, we deduce $G_{\omega_1,\dots,\omega_\ell}, G_{\omega'_1,\dots,\omega'_{\ell'}} \leq G_W = G_\Delta$. Now, since $H_{\omega_1,\dots,\omega_\ell} = H_{\omega_1,\dots,\omega'_{\ell'}}$, we deduce

$$G_{\omega_1,\ldots,\omega_\ell} = G_{\omega_1,\ldots,\omega'_{\ell'}}.$$

As $\ell \neq \ell'$, Lemma 2.1 implies that G is not IBIS.

IBIS LARGE BASE

Suppose now q = 2 and $k \ge 4$. Again, we may let U be a totally singular subspace of V with $\dim_{\mathbb{F}_q}(U) = k - 3$ and let

$$\Delta = \{ W \mid U \le W, \dim_{\mathbb{F}_q}(W) = k \text{ and } W \text{ totally singular} \}.$$

We may apply the argument above verbatim, with the old U replaced by this new U. We deduce that G is not IBIS.

3.4. The Klein correspondence. In this section we make a brief digression on the *Klein correspondence* in geometry. For this purpose, we follow [13, Chapter 8]. This will allow us to use the previous results on the unitary and symplectic groups to deal with the orthogonal groups.

Let W be the \mathbb{F}_q -vector space of all 4×4 skew-symmetric matrices. Clearly, dim $_{\mathbb{F}_q} W = 6$, because the entries of an element of W above the diagonal uniquely determine the matrix. Let $Q: W \to \mathbb{F}_q$ be the quadratic form defined by

$$X \mapsto Q(X) = \operatorname{Pf}(X)$$

where Pf(X) is the *Pfaffian* of the matrix X. In particular, if

$\begin{pmatrix} 0 \end{pmatrix}$	x_{12}	x_{13}	x_{14}	
$-x_{12}$	0	x_{23}	x_{24}	
$-x_{13}$	$-x_{23}$	0	x_{34}	,
$\sqrt{-x_{14}}$	$-x_{24}$	$-x_{34}$	0 /	

then

$$Pf(X) = x_{12}x_{34} - x_{13}x_{24} + x_{14}x_{23}$$

Clearly, Q has Witt defect 0. The group $GL_4(q)$ acts as a linear group on W by

 $X \mapsto P^T X P$,

where $X \in W$, $P \in GL_4(q)$ and P^T is the transposed of the matrix P. Using the properties of the Pfaffian, we have $Pf(P^TXP) = det(P)Pf(X)$. Therefore, Ppreserves the quadratic form Q if and only if det(P) = 1. Since $SL_4(q)$ equals its own derived subgroup, we deduce $SL_4(q) \leq \Omega_6^+(q)$. Observe that we are identifying here $SL_4(q)$ with a subgroup of $GL_6(q)$ using the action of $SL_4(q)$ on W. Moreover, from this embedding, we deduce $PSL_4(q) \leq P\Omega_6^+(q)$. Comparing the orders of $PSL_4(q)$ and of $P\Omega_6^+(q)$, we get $PSL_4(q) = P\Omega_6^+(q)$.

We now deduce some geometric facts. Let $0 \neq X \in W$ be singular with respect to Q. This means Q(X) = 0. Since det(X) is the square of Pf(X) = Q(X), we deduce that det(X) = 0. Since X is skew-symmetric, we deduce that the rank of X is even. As $X \neq 0$, we deduce that X has rank 2. It is not hard to verify that a skew-symmetric matrix of rank 2 is of the form

$$X = v^T w - w^T v,$$

for some vectors $v, w \in \mathbb{F}_q^4$. Observe that v and w are linearly independent over \mathbb{F}_q , because otherwise $X \neq 0$. This sets up a bijection between the points on the quadric Q and the 2-dimensional subspaces of \mathbb{F}_q^4 .

We summarise in the following proposition a few facts that we need later. A **plain pencil** in \mathbb{F}_q^4 is an incident point-plane pair, that is, a pair $\{A, B\}$ where A and B are subspaces of \mathbb{F}_q^4 with A < B, $\dim_{\mathbb{F}_q} A = 1$ and $\dim_{\mathbb{F}_q} B = 3$. Given a plain pencil $\{A, B\}$, the **plain pencil of lines** of $\{A, B\}$ is the collection of all the 2-dimensional subspaces of B containing A.

- **Proposition 3.17.** A: There exists a natural one-to-one correspondence between the 1-dimensional singular subspaces of $\mathbb{F}_q^6 \cong W$ with respect to Q and the 2-dimensional subspaces of \mathbb{F}_q^4 .
 - **B:** Under this correspondence, lines in Q, that is, 2-dimensional subspaces of $W = \mathbb{F}_q^6$ which are totally singular for Q, correspond to plane pencils of lines in \mathbb{F}_q^4 .
 - C: Under this correspondence, planes in Q, that is, 3-dimensional subspaces of $W = \mathbb{F}_q^6$ which are totally singular for Q, correspond to maximal families of pairwise intersecting lines in \mathbb{F}_q^4 . There are two types: all lines through a fixed point and all lines in a fixed plane.

3.5. Orthogonal groups of plus type. Let $d \ge 6$ be an even positive integer, let $q = p^f$ be a prime power with p a prime number, let $G_0 = P\Omega_d^+(q)$ and let Gbe an almost simple group with socle G_0 . We now describe the primitive subspace actions.

We let $V = \mathbb{F}_q^d$ be the *d*-dimensional vector space over the finite field \mathbb{F}_q of cardinality *q*. Then, for $1 \leq k \leq d/2$, we let Ω_k be the collection of all *k*-dimensional totally singular subspaces of *V*, with respect to the quadratic form *Q* preserved by G_0 . When k = d/2, the collection $\Omega_{d/2}$ of all maximal totally isotropic subspaces of *V* splits into two disjoint families, $\Omega_{d/2} = \Omega_{d/2,1} \cup \Omega_{d/2,2}$, namely the **greeks** and the **latins** where two elements *X* and *Y* of $\Omega_{d/2}$ are in the same family if and only if $X \cap Y$ has even codimension in either *X* or *Y*, for details we refer to [13].

Lemma 3.18. The action of G on Ω_k , or $\Omega_{d/2,1}$, $\Omega_{d/2,2}$, is IBIS if and only if (d, q, k) = (6, 2, 3).

Proof. When d = 6, using the Klein correspondence, we are reduced to the primitive actions of almost simple groups with socle $PSL_4(q)$ on 1-, 2-, 3-dimensional subspaces of \mathbb{F}_q^4 , and on pairs $\{A, B\}$ of subspaces with A < B and dim B = 3, dim A = 1. The result in this case follows from Lemmas 3.2, 3.4 and 3.6. Therefore, we suppose $d \geq 8$. Let W be a non-degenerate subspace of V with dim W = 6 and such that the quadratic form Q restricts to a quadratic form having Witt defect 0 on W. In particular, the quadratic form Q restricts also to W^{\perp} to a quadratic form having Witt defect 0.

Suppose that $k \leq d/2 - 2$. Let U be a totally isotropic subspace of W^{\perp} of dimension k - 1. Let

 $\Delta = \{ U' \oplus U \mid U' \le W, \dim(U' \oplus U) = k, U' \text{ totally isotropic} \}.$

Clearly, $G_{\Delta} = G_{W,U}$. Using the Klein correspondence and the results on the subspace actions of $PSL_4(q)$ (see Lemma 3.4 applied with k = 2), we deduce that G is not IBIS.

Suppose that k = d/2 - 1. Let U be a totally isotropic subspace of W^{\perp} of dimension k - 2, observe that U is a maximal totally isotropic subspace of W^{\perp} . Let

$$\Delta = \{ U' \oplus U \mid U' \le W, \dim(U' \oplus U) = k, U' \text{ totally isotropic} \}.$$

Clearly, $G_{\Delta} = G_{W,U}$. Before using the Klein correspondence we need to make an additional remark. Observe that the orthogonal decoposition $V = W \perp W^{\perp}$ gives rise to an embedding of $\mathrm{SO}_6^+(q) \times \mathrm{SO}_{d-6}^+(q)$ into $\mathrm{SO}_d^+(q)$. From [28], we see that $(\mathrm{SO}_6^+(q) \times \mathrm{SO}_{d-6}^+(q)) \cap \Omega_d^+(q)$ projects surjectively onto $\mathrm{SO}_6^+(q)$. In particular, the action of G_{Δ} on the 2-dimensional totally singular subspaces of W induces an almost simple group containing $PSO_6^+(q)$. Now, we can apply the Klein correspondence because the associated group in the action on pairs of subspaces contains $PSL_4(q)\langle \iota \rangle$, where ι is the inverse transposed automorphism. Now, using Lemma 3.6, we deduce that G is not IBIS.

Suppose that k = d/2. Without loss of generality, we may suppose that G is acting on $\Omega_{d/2,1}$. Let U be a totally isotropic subspace of W^{\perp} of dimension k-3 and let U'' be a 3-dimensional totally singular subspace of W with $U' \oplus U \in \Omega_{d/2,1}$. Let

 $\Delta = \{ U' \oplus U \mid U' \le W, \dim(U' \oplus U) = k,$

U' totally isotropic and dim $(U'/(U' \cap U''))$ even}.

Clearly, $G_{\Delta} = G_{W,U}$. What is more, the condition that $\dim(U'/(U' \cap U''))$ is even guarantees that every element of Δ is in $\Omega_{d/2,1}$. Using the Klein correspondence and Lemma 3.2, we deduce that G is not IBIS, except when q = 2.

Assume k = d/2 and q = 2. Let U be a totally isotropic subspace of V with $\dim U = k - 4$ and consider $\Delta = \{\omega \in \Omega \mid U \leq \omega\}$. We have a one-to-one correspondence between the elements of Δ and the 4-dimensional totally isotropic subspaces of U^{\perp}/U falling in the same family. Therefore, the result for an arbitrary value of d follows from the analogous result for d = 8. We have checked with a computer that when d = 8, the action of G on $\Omega_{4,1}$ and on $\Omega_{4,2}$ is not IBIS. \Box

3.6. Odd dimensional orthogonal groups. To deal with this case we continue our analysis on the Klein correspondence. Recall that we have an embedding of $SL_4(q)$ in $\Omega_6^+(q)$.

Let A be a non-singular skew-symmetric matrix. For the way that the action of $GL_4(q)$ was defined on W, the stabilizer of A in $SL_4(q)$ consists of the matrices P such that $P^T A P = A$. Therefore, this stabilizer is the symplectic group defined by the matrix A.

On the other hand, A can be viewed as a vector in W with $Q(A) \neq 0$ and so the stabilizer of A viewed in the orthogonal group $\Omega_6^+(q)$ is the 5-dimensional orthogonal group on A^{\perp} . Thus $\Omega_5(q) \cong PSp_4(q)$. Under this correspondence the orthogonal geometry and the symplectic geometry are linked together.

- **Proposition 3.19.** A: There exists a natural one-to-one correspondence between the 1-dimensional singular subspaces of $A^{\perp} \leq \mathbb{F}_q^6 \cong W$ with respect to Q and the 2-dimensional subspaces of \mathbb{F}_q^4 which are totally singular for $\operatorname{Sp}_4(q)$.
 - **B:** Under this correspondence, lines in Q contained in A^{\perp} correspond to 1-dimensional subspaces of \mathbb{F}_q^4 .

Let d be an odd integer with $d \geq 5$, let q be an odd prime power and let k be an integer with $1 \leq k \leq (d-1)/2$. We let G be an almost simple group with socle $\Omega_d(q) = P\Omega_d(q)$ and let Ω_k be the collection of all totally isotropic subspaces of \mathbb{F}_q^d of dimension k, with respect to the quadradic form preserved by $\Omega_d(q)$. We consider the primitive action of G on Ω_k .

Lemma 3.20. The action of G on Ω_k is not IBIS.

Proof. We dismiss the notation established to describe the Klein correspondence and we use Proposition 3.19 directly. Therefore, as usual, we let $V = \mathbb{F}_q^d$, G_0 be the socle of G and Q be the quadratic form on V preseved by G_0 . Let W be a non-degenerate subspace of V with dim W = 5 and such that the quadratic form Q restricts to a non-degenerate quadratic form having Witt defect 0 on W^{\perp} . Now, when k = (d-1)/2, let U be a totally isotropic subspace of W^{\perp} of dimension k - 2; when k < (d - 1)/2, let U be a totally isotropic subspace of W^{\perp} of dimension k - 1. Finally, let

$$\Delta = \{ U' \oplus U \mid U' \le W, \dim(U' \oplus U) = k, U' \text{ totally isotropic} \}.$$

Clearly, $G_{\Delta} = G_{W,U}$. Using the fact that q is odd and using the Klein correspondence and the results on the subspace actions of $PSp_4(q)$ (see Proposition 3.16), we deduce that G is not IBIS.

3.7. Orthogonal groups of minus type. The Klein correspondence allows to identify $P\Omega_6^-(q)$ with $PSU_4(q)$, details can be found in [13]. Under this identitication the orthogonal geometry and the Hermitian geometry are linked together. When $k \in \{1, 2\}$, the k-dimensional totally isotropic subspaces for the orthogonal group correspond to the k-dimensional totally singular subspaces for the Hermitian group.

Let d be an even integer with $d \ge 6$, let q be a prime power and let k be an integer with $1 \le k \le d/2 - 1$. We let G be an almost simple group with socle $P\Omega_d^-(q)$ and let Ω_k be the collection of all totally isotropic subspaces of \mathbb{F}_q^d of dimension k, with respect to the quadradic form Q preserved by $\Omega_d(q)$. We consider the primitive action of G on Ω_k .

Lemma 3.21. The action of G on Ω_k is not IBIS.

Proof. Let W be a non-degenerate subspace of V with dim W = 6 and such that the quadratic form Q restricts to a non-degenerate quadratic form having Witt defect 0 on W^{\perp} and to a non-degenerate quadratic form having Witt defect 1 on W. Now, when k = d/2 - 1, let U be a totally isotropic subspace of W^{\perp} of dimension k - 2; when k < d/2 - 1, let U be a totally isotropic subspace of W^{\perp} of dimension k - 1. Finally, let

$$\Delta = \{ U' \oplus U \mid U' \le W, \dim(U' \oplus U) = k, U' \text{ totally isotropic} \}.$$

Clearly, $G_{\Delta} = G_{W,U}$. Using the Klein correspondence and the results on the subspace actions of $PSU_4(q)$ (see Proposition 3.12), we deduce that G is not IBIS.

4. Even characteristic symplectic groups: case (1.3)

In this section we deal with the actions listed at item (1.2) of Definition 1.4. In particular G is almost simple group with socle $G_0 = \operatorname{Sp}_{2m}(q)'$ and q a power of 2. Consulting [2, 28], it is clear that if H is a maximal subgroup of G not containing G_0 , and $H \cap G_0$ is a subgroup of $M = O_{2m}^{\pm}(q)$, then $H \cap G_0 = M$. In light of this, we may first consider the action of $G_0 = \operatorname{Sp}_{2m}(q)$ on the set of right cosets of $M = O_{2m}^{\pm}(q)$.^k

Our treatment follows [24], which in turn was inspired by [21, §7.7] where the case of $\operatorname{Sp}_{2m}(2)$ was considered. Let e and m be positive integers, let $q = 2^e$, let \mathbb{F}_q be the finite field with q elements, and let $V = \mathbb{F}_q^{2m}$ be the 2*m*-dimensional vector

^kThe embedding of $O_{2m}^{\pm}(q)$ in $\operatorname{Sp}_{2m}(q)$ when q is even was discovered by Dye, see [22].

space of row vectors over \mathbb{F}_q . To start with we adjust notation slightly, and assume that $G = \operatorname{Sp}_{2m}(q)$ is the symplectic group defined by the symmetric matrix

$$f = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$$

where 0 and I are the zero and identity $m \times m$ -matrices, respectively. In particular, G is the group of invertible matrices preserving the bilinear form $\varphi: V \times V \to \mathbb{F}_q$ defined by

$$\varphi(u,v) = ufv^T,$$

for every $u, v \in V$, that is

$$G = \left\{ g \in \mathrm{GL}_{2m}(q) \mid gfg^T = f \right\}.$$

Note that the bilinear form φ is alternating, i.e. for all $u \in V$, we have

(9)
$$\varphi(u, u) = 0$$

Moreover, since \mathbb{F}_q is of characteristic 2, the form φ is symmetric, i.e. for all $u, v \in V$, we have

(10)
$$\varphi(u,v) = \varphi(v,u)$$

Now we let Ω be the set of quadratic forms $\theta: V \to \mathbb{F}_q$ polarising to φ . Recall that this means that $\theta: V \to \mathbb{F}_q$ is a function satisfying

- $\theta(u+v) \theta(u) \theta(v) = \varphi(u,v)$ for every $u, v \in V$, and
- $\theta(cu) = c^2 u$, for every $c \in \mathbb{F}_q$ and $u \in V$.

Next, consider the matrix

$$e = \begin{pmatrix} 0 & I \\ 0 & 0 \end{pmatrix}$$

and the quadratic form $\theta_0: V \to \mathbb{F}_q$ defined by

$$\theta_0(u) = ueu^T$$
,

for every $u \in V$. Then, for every $u, v \in V$, we have

(11)

$$\theta_{0}(u+v) - \theta_{0}(u) - \theta_{0}(v) = (u+v)e(u+v)^{T} - ueu^{T} - vev^{T}$$

$$= ueu^{T} + vev^{T} + uev^{T} + veu^{T} - ueu^{T} - vev^{T}$$

$$= uev^{T} + veu^{T} = uev^{T} + ue^{T}v^{T}$$

$$= u(e+e^{T})v^{T} = ufv^{T} = \varphi(u,v).$$

In particular, θ_0 is a quadratic form whose polarisation is the symplectic form φ . This shows that $\theta_0 \in \Omega$.

Let $\theta \in \Omega$ and define $\lambda = \theta - \theta_0$. We have

$$\begin{split} \lambda(u+v) =& \theta(u+v) - \theta_0(u+v) = \theta(u) + \theta(v) + \varphi(u,v) - \theta_0(u) - \theta_0(v) - \varphi(u,v) \\ =& \lambda(u) + \lambda(v), \\ \lambda(cu) =& \theta(cu) - \theta_0(cu) = c^2 \theta(u) - c^2 \theta_0(u) = c^2 \lambda(u), \end{split}$$

for every $u, v \in V$ and for every $c \in \mathbb{F}_q$. Therefore, since \mathbb{F}_q is of characteristic 2, the function $\lambda : V \to \mathbb{F}_q$ is semilinear and hence there exists a unique $b \in V$ such that

$$\lambda(u) = (u \cdot b^T)^2$$
, for every $u \in V$.

Since f is an invertible matrix, there exists a unique $a \in V$ with b = af and hence

$$\lambda(u) = (ufa^T)^2 = \varphi(u, a)^2$$

for every $u \in V$. Summing up, we have shown that an arbitrary element of Ω is of the form

$$u \mapsto \theta_0(u) + \varphi(u, a)^2$$

where $a \in V$. We denote this element of Ω simply by θ_a . Thus

(12)
$$\theta_a(u) = \theta_0(u) + \varphi(u, a)^2$$
, for every $u \in V$.

In particular, the elements of Ω are parametrised by the vectors of V. Moreover, if $\theta_a = \theta_{a'}$ for some $a, a' \in V$, then $\theta_a(u) = \theta_{a'}(u)$ for every $u \in V$ and this implies $\varphi(u, a) = \varphi(u, a')$ for every $u \in V$. Since φ is non-degenerate, we obtain a = a'. Hence, the set Ω is in one-to-one correspondence with V. This, in particular, yields that

$$|\Omega| = q^{2m}.$$

Lemma 4.1 (Lemma 6.8, [24]). For every $\theta \in \Omega$ and $x \in G$, the mapping

(13)
$$\begin{aligned} \theta^x \colon V \to \mathbb{F}_q \\ u \mapsto \theta(ux^{-1}) \end{aligned}$$

lies in Ω . In particular, this defines an action of the group G on the set Ω .

Before continuing our discussion, we gather some information on G. Let $a \in V$, we define the mapping

(14)
$$\begin{aligned} t_a \colon V \to V \\ u \mapsto u + \varphi(u, a)a. \end{aligned}$$

Such a function is called a *transvection*. For every $u, v \in V$ and $c \in \mathbb{F}_q$ we have $(u+v)t_a = (u+v) + \varphi(u+v, a)a = (u+\varphi(u, a)a) + (v+\varphi(v, a)a) = (u)t_a + (v)t_a;$ $(cu)t_a = cu + \varphi(cu, a)a = c(u+\varphi(u, a)a) = c(u)t_a.$ Hence t_a is linear. Moreover, for every $u \in V$ we have $(u)t^2 = (u+\varphi(u, a)a)t_a = u + \varphi(u, a)a + \varphi(u+\varphi(u, a)a, a)a$

$$\begin{aligned} (u)t_a^2 &= (u+\varphi(u,a)a)t_a = u+\varphi(u,a)a+\varphi(u+\varphi(u,a)a,a)a\\ &= u+\varphi(u,a)a+\varphi(u,a)a+\varphi(u,a)\varphi(a,a)a = u+2\varphi(u,a)a+\varphi(u,a)\varphi(a,a)a\\ &= u+\varphi(u,a)\varphi(a,a)a = u, \end{aligned}$$

where in the second-last equality we use the fact that the characteristic of \mathbb{F}_q is 2. This shows that t_a is an involution. Finally, for every $u, v \in V$, we have

$$\begin{aligned} \varphi((u)t_a,(v)t_a) &= \varphi(u + \varphi(u,a)a, v + \varphi(v,a)a) \\ &= \varphi(u,v) + \varphi(v,a)\varphi(u,a) + \varphi(u,a)\varphi(a,v) + \varphi(u,a)\varphi(v,a)\varphi(a,a) \\ &= \varphi(u,v) + 2\varphi(v,a)\varphi(u,a) + \varphi(u,a)\varphi(v,a)\varphi(a,a) \\ &= \varphi(u,v) + \varphi(u,a)\varphi(v,a)\varphi(a,a) \\ &= \varphi(u,v). \end{aligned}$$

Therefore t_a preserves φ and hence lies in the symplectic group G.

We are now interested in computing the image of θ_a under the transvection t_c . First recall that, in a field of characteristic 2, since x = -x for every $x \in \mathbb{F}_q$, the square root $\sqrt{\cdot} \colon \mathbb{F}_q \to \mathbb{F}_q$ is a well-defined map. Moreover, for every $a, b, x, y \in \mathbb{F}_q$ such that $x = a^2$ and $y = b^2$ we have $(\sqrt{x} + \sqrt{y})^2 = (a+b)^2 = a^2 + b^2 = x + y$, which implies $\sqrt{x} + \sqrt{y} = \sqrt{x+y}$. Moreover, recall that θ_a is a quadratic form polarising to φ and that t_a is an involution, in particular $t_c = t_c^{-1}$. By using these facts, given $v \in V$, we have

$$\begin{split} \theta_a^{t_c}(u) &= \theta_a(ut_c^{-1}) = \theta_a(ut_c) = \theta_a(u + \varphi(u, c)c) \\ &= \theta_a(u) + \theta_a(\varphi(u, c)c) + \varphi(u, \varphi(u, c)c) \\ &= \theta_a(u) + \varphi(u, c)^2 \theta_a(c) + \varphi(u, c)^2 \\ &= \theta_a(u) + (\theta_a(c) + 1)\varphi(u, c)^2 = \theta_a(u) + (\sqrt{\theta_a(c)} + 1\varphi(u, c))^2 \\ &= \theta_a(u) + \varphi(u, (\sqrt{\theta_a(c)} + 1)c)^2 = \theta_0(u) + \varphi(u, a)^2 + \varphi(u, (\sqrt{\theta_a(c)} + 1)c)^2 \\ &= \theta_0(u) + \varphi(u, a + (\sqrt{\theta_a(c)} + 1)c)^2 \\ &= \theta_{a+(\sqrt{\theta_a(c)} + 1)c}(u). \end{split}$$

From this, we deduce

(15)
$$\theta_a^{t_c} = \theta_{a+(\sqrt{\theta_a(c)}+1)c}.$$

We now recall some facts about Galois theory. For a reference see [29, Chapter VI]. The Frobenius mapping $\phi: x \mapsto x^2$ from \mathbb{F}_q to itself is a generator of the Galois group of \mathbb{F}_q over \mathbb{F}_2 . There exists a well-defined \mathbb{F}_2 -linear *trace* mapping $\operatorname{Tr}: \mathbb{F}_q \to \mathbb{F}_2$. In what follows, we need only two basic facts about Tr: first, Tr is surjective and second, from Hilbert's 90 Theorem, the kernel of Tr consists of the set $\{x^2 + x \mid x \in \mathbb{F}_q\}$ and has cardinality q/2.

Define

(16)
$$\Omega^{+} = \{\theta_{a} \mid \operatorname{Tr}(\theta_{0}(a)) = 0\},$$
$$\Omega^{-} = \{\theta_{a} \mid \operatorname{Tr}(\theta_{0}(a)) = 1\}.$$

Observe that the above definition is a generalization of the definition of Ω^+ and Ω^- in [21, Corollary 7.7 A]. Indeed, if q = 2, then the Galois group is the trivial group and hence the trace map is the identity.

Let $N = \langle t_a \mid a \in V \rangle$ be the subgroup of G generated by the transvections. Observe that, for all $a, u \in V$ and $x \in G$, we have

$$(u)x^{-1}t_ax = (ux^{-1} + \varphi(ux^{-1}, a)a)x = u + \varphi(u, ax)ax = (u)t_{ax}.$$

In particular, this shows that $N \trianglelefteq G$.

Lemma 4.2 (Lemma 6.8, [24]). The sets Ω^+ and Ω^- are N-orbits on Ω , with

$$|\Omega^+| = \frac{q^m(q^m+1)}{2}, \quad |\Omega^-| = \frac{q^m(q^m-1)}{2}$$

These are also orbits for G. Moreover, the action of $G = \text{Sp}_{2m}(q)$ on Ω^{\pm} is permutation equivalent to the action of G on the right cosets of $O_{2m}^{\pm}(q)$.

Now, we select a distinguished element of Ω^- . Let $\epsilon \in \mathbb{F}_q$ with $\operatorname{Tr}(\epsilon) = 1$ and set $\varepsilon = \epsilon e_1 + e_{m+1}$, where $(e_i)_{i \in \{1, \dots, 2m\}}$ is the standard basis of V. Since $\theta_0(\epsilon e_1) = 0 = \theta_0(e_{m+1})$, we have

$$\theta_0(\varepsilon) = \theta_0(\epsilon e_1) + \theta_0(e_{m+1}) + \varphi(\epsilon e_1, e_{m+1}) = \epsilon \varphi(e_1, e_{m+1}) = \epsilon,$$

and hence $\operatorname{Tr}(\theta_0(\varepsilon)) = \operatorname{Tr}(\epsilon) = 1$. Therefore, $\theta_{\varepsilon} \in \Omega^-$.

We now compute stabilizer chains in these two actions. Let $\circ \in \{+, -\}$ (we deal simultaneously with both cases). Let $\{\theta_{a_1}, \theta_{a_2}, \ldots, \theta_{a_k}\}$ be a subset of Ω° . Without loss of generality, we may suppose that $a_1 = 0$ when $\circ = +$ and $a_1 = \varepsilon$ when $\circ = -$.

Let us define

 $\mathbf{C}_{G_{\theta_{a_1}}}(a_1 + a_i) = \{ x \in G_{\theta_{a_1}} \mid (a_1 + a_i)x = a_1 + a_i \},\$

that is the set of matrices in $G_{\theta_{a_1}}$ fixing the vector $a_1 + a_i \in V$.

Lemma 4.3 (Lemma 6.10, [24]). For every $i \in \{2, ..., k\}$,

(17)
$$G_{\theta_{a_1}} \cap G_{\theta_{a_i}} = \mathbf{C}_{G_{\theta_{a_1}}}(a_1 + a_i),$$

We are now ready to prove our main result of this section.

Proposition 4.4. Let G be a permutation group as in Definition 1.4 (1.2). Then G is not IBIS, except in one of the folling cases:

- (4.1) $m = 1, G = \operatorname{Sp}_2(q)$ in its action on $\Omega^-, q = 2^f, f \ge 3$,
- (4.2) m = 1, $G = \operatorname{Sp}_2(q)$ or $G = \operatorname{Aut}(\operatorname{Sp}_2(q))$ in its action on Ω^- , q = 4,
- (4.3) $m = 1, G = \operatorname{Aut}(\operatorname{Sp}_2(q)) \cong \Gamma \operatorname{L}_2(q)$ in its action on $\Omega^+, q = 2^f, f$ odd prime,
- (4.4) m = q = 2.

Proof. Suppose first m = 1. Here, $G_0 = \operatorname{Sp}_2(q) = \operatorname{SL}_2(q)$. The stabilizer of an element of Ω^+ in G_0 is isomorphic to the dihedral group of order 2(q-1). This action was discussed in Example 3.8 in [31] and it was proven that it is IBIS only when $G = \Gamma L_2(q)$, $q = 2^f$ and f is an odd prime. In particular, we obtain the examples in (4.3). The stabilizer of an element of Ω^- in G_0 is isomorphic to the dihedral group of order 2(q+1). This action was discussed in Example 3.7 in [31] and it was proven to be IBIS only when $G = \operatorname{SL}_2(q)$, $q = 2^f$ and $f \ge 3$, or when $\operatorname{SL}_2(q) \le G \le \operatorname{Aut}(\operatorname{SL}_2(q))$ with q = 4. In particular, we obtain the examples in (4.1) and (4.2).

We now deal with the case m = q = 2. In this case $\text{Sp}_4(2)$ is not simple, since $\text{Sp}_4(2)' \cong \text{Alt}(6)$. Hence G is an almost simple group having socle Alt(6). Moreover, $|\Omega^+| = 10$ and $|\Omega^-| = 6$. The action of G on Ω^{\pm} is IBIS, see [31, Theorem 1.2]. In particular, we obtain the examples in (4.4). Therefore, for the rest of our argument, we may suppose that $(m, q) \neq (2, 2)$ and $m \geq 2$.

As usual, set $G_0 = \text{Sp}_{2m}(q)$. From Lemma 4.2, we may idenfity the action of G_0 on the right cosets of $O_{2m}^{\pm}(q)$ with the action of G_0 on Ω^{\pm} and we may also use the notation above.

Suppose first q = 2. This case is rather special because in this case the action of G_0 on both Ω^+ and Ω^- is 2-transitive.¹ We first consider Ω^+ and observe that $\theta_0, \theta_{e_1} \in \Omega^+$ because $\theta_0(e_1) = 0$, (that is, e_1 is singular for the quadratic form θ_0), see (16). From Lemma 4.3, $G_{\theta_0} \cap G_{\theta_{e_1}} = \mathbf{C}_{O_{2m}^+(q)}(e_1)$. Since the transitive action of G_{θ_0} on $\Omega^+ \setminus \{\theta_0\}$ is permutation equivalent to the action of $G_{\theta_0} = O_{2m}^+(q)$ on the right cosets of $G_{\theta_0} \cap G_{\theta_{e_1}} = \mathbf{C}_{O_{2m}^+(q)}(e_1)$, we deduce that the action of G_{θ_0} on $\Omega^+ \setminus \{\theta_0\}$ is permutation equivalent to the action of $O_{2m}^+(q)$ on the set of singular non-zero vectors of V, with respect to the quadratic form θ_0 . In the special case q = 2, this implies that the action of G_{θ_0} on $\Omega^+ \setminus \{\theta_0\}$ is permutation equivalent to the action of $O_{2m}^+(q)$ on the set of 1-dimensional singular subspaces of V, which

30

¹In general the action of G_0 on Ω^{\pm} has rank q, see for instance [25].

we have shown in Lemma 3.18 to be not IBIS. The argument for Ω^- is similar. Observe that $\theta_{\varepsilon} \in \Omega^-$. Hence, using (15) and using the fact that $\theta_{\varepsilon}(e_2) = 0$,

$$\theta_{\varepsilon}^{t_{e_2}} = \theta_{\varepsilon + (\sqrt{\theta_{\varepsilon}(e_2)} + 1)e_2} = \theta_{\varepsilon + e_2} \in \Omega^-.$$

From Lemma 4.3, $G_{\theta_{\varepsilon}} \cap G_{\theta_{\varepsilon+e_2}} = \mathbf{C}_{\mathcal{O}_{2m}^-(q)}(e_2)$. Since the transitive action of $G_{\theta_{\varepsilon}}$ on $\Omega^- \setminus \{\theta_{\varepsilon}\}$ is permutation equivalent to the action of $G_{\theta_{\varepsilon}} = \mathcal{O}_{2m}^-(q)$ on the right cosets of $G_{\theta_{\varepsilon}} \cap G_{\theta_{\varepsilon+e_2}} = \mathbf{C}_{\mathcal{O}_{2m}^-(q)}(e_2)$ and since $\theta_{\varepsilon}(e_2) = 0$, we deduce that the action of $G_{\theta_{\varepsilon}}$ on $\Omega^- \setminus \{\theta_{\varepsilon}\}$ is permutation equivalent to the action of $\mathcal{O}_{2m}^-(q)$ on the set of singular non-zero vectors of V, with respect to the quadratic form θ_{ε} . In the special case q = 2, this implies that the action of $G_{\theta_{\varepsilon}}$ on $\Omega^- \setminus \{\theta_{\varepsilon}\}$ is permutation equivalent to the action of $\mathcal{O}_{2m}^-(q)$ on the set of 1-dimensional singular subspaces of V, which we have shown in Lemma 3.21 to be not IBIS.

Suppose next q > 2. We first take in account Ω^+ . Consider

$$H = G_{\theta_0} \cap \bigcap_{i=3}^m (G_{\theta_{e_i}} \cap G_{\theta_{e_{i+m}}}).$$

By construction, $H \cong O_4^+(q)$ fixes pointwise the vector space $\langle e_i, e_{i+m} | 3 \le i \le m \rangle$ and induces the orthogonal group $O_4^+(q)$ on $\langle e_1, e_2, e_{1+m}, e_{2+m} \rangle$. This shows that we may deduce the fact that the action of G on Ω^+ is not IBIS for general values of m, arguing only on the case m = 2. Therefore, in what follows, we assume m = 2. A direct computation with the quadratic form θ_0 shows that

$$G_{\theta_0} \cap G_{\theta_{e_2}} = \left\{ \begin{pmatrix} a & ay & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & a^{-1}x & a^{-1} & 0\\ x & xy & y & 1 \end{pmatrix} \mid a, x, y \in \mathbb{F}_q, a \neq 0 \right\}$$
$$\cup \left\{ \begin{pmatrix} 0 & ax & a & 0\\ 0 & 1 & 0 & 0\\ a^{-1} & a^{-1}y & 0 & 0\\ x & xy & y & 1 \end{pmatrix} \mid a, x, y \in \mathbb{F}_q, a \neq 0 \right\}$$

In particular, $|G_{\theta_0} \cap G_{\theta_{e_2}}| = 2(q-1)q^2$. Similarly,

$$G_{\theta_0} \cap G_{\theta_{e_2}} \cap G_{\theta_{e_1}} = \left\{ \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & x & 1 & 0 \\ x & 0 & 0 & 1 \end{pmatrix} \mid x \in \mathbb{F}_q \right\}.$$

In particular, $|G_{\theta_0} \cap G_{\theta_{e_2}} \cap G_{\theta_{e_3}}| = q$. Moreover, from this description, it is clear that $G_{\theta_0} \cap G_{\theta_{e_2}} \cap G_{\theta_{e_3}} \cap G_{\theta_{e_4}} = 1$. Therefore $\operatorname{Sp}_4(q)$ on Ω^+ has a base of size 4. Now, let $\lambda \in \mathbb{F}_q$ with $\operatorname{Tr}(\lambda) = 0$ and $\lambda \neq 0$: observe that this is possible because $q \neq 2$. Let $a = e_1 + \lambda e_3$ and observe that $\theta_a \in \Omega^+$ because $\theta_0(a) = 0$. A direct

computation with the quadratic form θ_0 shows that

$$G_{\theta_{0}} \cap G_{\theta_{e_{2}}} \cap G_{\theta_{a}} = \left\{ \begin{pmatrix} 1 & \lambda x & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & ax & 1 & 0 \\ x & \lambda x^{2} & \lambda x & 1 \end{pmatrix} \mid x \in \mathbb{F}_{q} \right\}$$
$$\cup \left\{ \begin{pmatrix} 0 & \lambda x & \lambda & 0 \\ 0 & 1 & 0 & 0 \\ \lambda^{-1} & x & 0 & 0 \\ x & \lambda x^{2} & \lambda x & 1 \end{pmatrix} \mid x \in \mathbb{F}_{q} \right\}$$

In particular, $|G_{\theta_0} \cap G_{\theta_{e_2}} \cap G_{\theta_a}| = 2q$. Moreover, from this description, it is clear that

$$G_{\theta_0} \cap G_{\theta_{e_2}} \cap G_{\theta_a} \cap G_{\theta_{e_4}} = \left\langle \begin{pmatrix} 0 & 0 & \lambda & 0 \\ 0 & 1 & 0 & 0 \\ \lambda^{-1} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \right\rangle$$

has order 2. Now, $G_{\theta_0} \cap G_{\theta_{e_2}} \cap G_{\theta_a} \cap G_{\theta_{e_4}} \cap G_{\theta_{e_1}} = 1$. Therefore the action of $\operatorname{Sp}_4(q)$ on Ω^+ has a base of size 5. This concludes the proof that G on Ω^+ is not IBIS when q > 2.

We now consider Ω^- . Observe that $\theta_{\varepsilon}(e_i) = 0$ for each $i \in \{1, \ldots, 2m\} \setminus \{1, 1 + m\}$. In particular, for these values of i, we have

$$\theta_{\varepsilon}^{te_i} = \theta_{\varepsilon + (\sqrt{\theta_{\varepsilon}(e_i)} + 1)e_i} = \theta_{\varepsilon + e_i}.$$

Consider

$$H = G_{\theta_{\varepsilon}} \cap \bigcap_{i=3}^{m} (G_{\theta_{\varepsilon+e_i}} \cap G_{\theta_{\varepsilon+e_{i+m}}}) = \mathbf{C}_{G_{\theta_{\varepsilon}}}(\langle e_i, e_{i+1} \mid 3 \le i \le m \rangle).$$

By construction, $H \cong O_4^-(q)$ fixes pointwise the vector space $\langle e_i, e_{i+m} | 3 \le i \le m \rangle$ and induces the orthogonal group $O_4^-(q)$ on $\langle e_1, e_2, e_{1+m}, e_{2+m} \rangle$. This shows that we may deduce the fact that the action of G on Ω^- is not IBIS for general values of m, arguing only on the case m = 2. Therefore, in what follows, we assume m = 2. Observe that when m = 2, the quadratic form θ_{ε} has matrix

$$\begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & \epsilon^2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

As $\theta_{\varepsilon}(e_2) = 0$, we have

$$\boldsymbol{\theta}_{\varepsilon}^{t_{e_2}} = \boldsymbol{\theta}_{\varepsilon + (\sqrt{\theta_{\varepsilon}(e_2)} + 1)e_2} = \boldsymbol{\theta}_{\varepsilon + e_2}$$

A computation gives that the group

$$G_{\theta_{\varepsilon}} \cap G_{\theta_{\varepsilon}+e_2} = \mathbf{C}_{G_{\theta_{\varepsilon}}}(e_2) = \mathbf{C}_{\mathcal{O}_4^-(q)}(e_2)$$

consists of the matrices

$$\begin{pmatrix} a & cx_3 + ay_3 & c & 0\\ 0 & 1 & 0 & 0\\ x_1 & x_1y_3 + x_3y_1 & y_1 & 0\\ x_3 & x_3^2 + x_3y_3 + \epsilon^2 y_3^2 & y_3 & 1 \end{pmatrix} \text{ subject to } \begin{cases} ay_1 + cx_1 &= 1\\ a^2 + ac + \epsilon^2 c^2 &= 1\\ x_1^2 + x_1y_1 + \epsilon^2 y_1^2 &= \epsilon^2. \end{cases}$$

Moreover, as $\theta_{\varepsilon}(e_4) = 0$, we have

$$\theta_{\varepsilon}^{t_{e_4}} = \theta_{\varepsilon + (\sqrt{\theta_{\varepsilon}(e_4)} + 1)e_4} = \theta_{\varepsilon + e_4}.$$

A computation gives that the group

$$G_{\theta_{\varepsilon}} \cap G_{\theta_{\varepsilon+e_{2}}} \cap G_{\theta_{\varepsilon+e_{4}}} = \mathbf{C}_{G_{\theta_{\varepsilon}}}(\langle e_{2}, e_{4} \rangle) = \mathbf{C}_{\mathbf{O}_{4}^{-}(q)}(\langle e_{2}, e_{4} \rangle)$$

consists of the matrices

$$\begin{pmatrix} a & 0 & c & 0 \\ 0 & 1 & 0 & 0 \\ x_1 & 0 & y_1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \text{ subject to } \begin{cases} ay_1 + cx_1 &= 1 \\ a^2 + ac + \epsilon^2 c^2 &= 1 \\ x_1^2 + x_1 y_1 + \epsilon^2 y_1^2 &= \epsilon^2. \end{cases}$$

Moreover, as $\theta_{\varepsilon}(e_3) = \epsilon^2$, we have

$$\theta_{\varepsilon}^{t_{e_3}} = \theta_{\varepsilon + (\sqrt{\theta_{\varepsilon}(e_3)} + 1)e_3} = \theta_{\varepsilon + (1+\epsilon)e_3}$$

Without loss of generality, we may suppose that $\epsilon \neq 1$. A computation gives that the group

$$G_{\theta_{\varepsilon}} \cap G_{\theta_{\varepsilon+e_2}} \cap G_{\theta_{\varepsilon+e_4}} \cap G_{\theta_{\varepsilon+(1+\epsilon)e_3}}$$

has order 2 and is generated by

$$\begin{pmatrix} 1 & 0 & \epsilon^{-2} & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Therefore $\text{Sp}_4(q)$ in its action on Ω^- has a basis of size 5. Now, fix $\alpha \in \mathbb{F}_q \setminus \{0, 1\}$ and let $a = \alpha e_1$. We have

$$\theta_{\varepsilon}^{t_a} = \theta_{\varepsilon + (\sqrt{\theta_{\varepsilon}(a) + 1})a} = \theta_{\varepsilon + \alpha(\alpha + 1)e_1}.$$

A computation gives that the group

$$G_{\theta_{\varepsilon}} \cap G_{\theta_{\varepsilon+e_2}} \cap G_{\theta_{\varepsilon+(1+\alpha)\alpha e_1}}$$

consists of the matrices

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ x_1 & x_3 & 1 & 0 \\ x_3 & x_3^2 & 0 & 1 \end{pmatrix} \text{ subject to } x_1 \in \{0, 1\}.$$

Finally,

$$G_{\theta_{\varepsilon}} \cap G_{\theta_{\varepsilon+e_2}} \cap G_{\theta_{\varepsilon+(1+\alpha)\alpha e_1}} \cap G_{\theta_{\varepsilon+(1+\epsilon)e_3}} = 1$$

and hence $\operatorname{Sp}_4(q)$ on Ω^- has also a base of size 4. The case when $G_0 < G \leq \operatorname{Sp}_{2m}(q) \rtimes \operatorname{Aut}(\mathbb{F}_q)$ is analogous.

5. Primitive action on non-degenerate subspaces: Case (1.4)

Here, we deal with the action on the non-degenerate subspaces and hence with case (1.4). Let G an almost simple group with socle a classical group G_0 (either symplectic, unitary or orthogonal), and let V be its associated module of dimension d. We describe the primitive action of G on non-degenerate subspaces.

When G is symplectic or unitary, the action of G is on the collection of all non-degenerate subspaces of V of a given dimension k with k < d/2. When G is orthogonal, there are a few cases to consider. Indeed, the collection of nondegenerate k-subspaces of V is not always a G_0 -orbit. Indeed, it is a single orbit if and only if d is even and k is odd.

Lemma 5.1. Let V, d and k be as above and let W_1 be a non-degenerate k-dimensional subspace of V. Except when

- (q,k) = (2,2) and V is symplectic, or
- (q,k) = (2,1) and V is Hermitian, or
- (q,k) = (3,1) and V is orthogonal, or
- (q,k) = (2,2), V is orthogonal and W_1 is anisotropic,

there exist two non-degenerate subspaces W_2 and W_3 of V and an element g of G_0 with

- (5.1) $W_1 \perp W_2$,
- (5.2) $W_1 \oplus W_2 = W_1 \oplus W_3$ and $W_1 \cap W_3 = W_2 \cap W_3 = 0$, (5.3) $W_1^g = W_1, W_2^g = W_2$ and $W_3^g \neq W_3$.

Proof. We give complete details with V is symplectic, all other cases are similar. Let b be the symplectic form defining V. Fix a non-degenerate k-dimensional subspace W_1 of V and fix a non-degenerate k-dimensional subspace W_2 of W_1^{\perp} . Clearly, (5.1) is satisfied. Let $f: W_1 \to W_2$ be an arbitrary isomorphism such that $b(f(w_1), f(w_2)) \neq b(w_1, w_2)$, for some $w_1, w_2 \in W_1$. The existence of f is clear when q > 2 because we may take f the scalar multiplication by a scalar different form 0 and 1. When q = 2, the existence of f depends on the fact that $SL_{2k}(2) = Sp_{2k}(2)$ only when k = 2. Therefore, for the rest of the argument assume $(k,q) \neq (2,2)$.

Let $W_3 = \{w + f(w) \mid w \in W_1\}$. Clearly, W_3 satisfies (5.2). Let $w_1, w_1' \in W_1$ and let $w_3 = w_1 + f(w_1), w'_3 = w'_1 + f(w'_1)$. As

$$b(w_3, w'_3) = b(w_1, w'_1) + b(w_1, f(w'_1)) + b(f(w_1), w'_1) + b(f(w_1), f(w'_1))$$

= b(w_1, w_2) + b(f(w_1), f(w_2)),

we see that W_3 is non-degenerate from our assumption on f.

Finally, let $x: W_1 \to W_1$ be any non-identity symplectic isomorphism and, using Witt's lemma, extend x to a symplectic automorphism g of V fixing pointwise W_2 . Clearly, q satisfies (5.3).

We need an analogous result for the exceptional cases arising in Lemma 5.1.

Lemma 5.2. Assume that V, k, W_1 are one of the four exceptional cases in Lemma 5.1. Then, there exist three non-degenerate subspaces W_2 , W_3 and W_4 of V and an element g of G_0 with

- (5.1) $W_1 \perp W_2 \perp W_3$,
- (5.2) $W_1 \oplus W_2 \oplus W_3 = W_1 \oplus W_2 \oplus W_4$ and $W_1 \cap W_4 = W_2 \cap W_4 = W_3 \cap W_4 = 0$,

IBIS LARGE BASE

(5.3)
$$W_1^g = W_1, W_2^g = W_2, W_3^g = W_3 \text{ and } W_4^g \neq W_4.$$

Proof. The proof uses the same argument as the proof of Lemma 5.1. In this case, it suffices to take $W_4 = \{w + f_2(w) + f_3(w) \mid w \in W_1\}$, where $f_2 : W_1 \to W_2$ and $f_3 : W_1 \to W_3$ are suitable isomorphisms.

Using Lemmas 5.1 and 5.2 , we immediately deduce the following result, which concludes case 1.4.

Lemma 5.3. The primitive actions of G on non-degenerate subspaces are not IBIS.

Proof. Suppose first that we are not in one of the exceptional cases in Lemma 5.1 and let W_1, W_2, W_3 be as in the statement of Lemma 5.1. Clearly,

$$G_{W_1} > G_{W_1,W_2} > G_{W_1,W_2,W_3},$$

where the last strict inequality uses (5.3) of Lemma 5.1. Finally, observe that $G_{W_1,W_3} \leq G_{W_1,W_2}$ from (5.1) and (5.2) of Lemma 5.1.

Next, suppose that we are in one of the exceptional cases in Lemma 5.1 and let W_1, W_2, W_3, W_4 be as in the statement of Lemma 5.2. Clearly,

$$G_{W_1} > G_{W_1,W_2} > G_{W_1,W_2,W_3} > G_{W_1,W_2,W_3,W_4},$$

where the last strict inequality uses (5.3) of Lemma 5.2. Finally, observe that $G_{W_1,W_2,W_4} \leq G_{W_1,W_2,W_3}$ from (5.1) and (5.2) of Lemma 5.2.

6. ACTION ON NON-SINGULAR 1-SUBSPACES: CASE (1.5)

Here G is an almost simple group having socle $G_0 = \Omega_d^{\pm}(q)$ acting on 1-dimensional non-singular subspaces of V and q is even. Since $\Omega_2^{\pm}(q)$ is solvable, we may suppose that $d \ge 4$.

We look closely at the case d = 4 because this will serve as a base case of an induction.

Lemma 6.1. When d = 4, the action is IBIS if and only if $G = G_0 = \Omega_4^{\pm}(q)$ and $q \ge 4$.

Proof. We first deal with the hyperbolic case.

In this case, $G_0 = \Omega_4^+(q) = \mathrm{SL}_2(q) \times \mathrm{SL}_2(q)$. Moreover, the stabilizer of a nonsingular vector in G_0 is isomorphic to $\mathrm{Sp}_2(q) = \mathrm{SL}_2(q)$ diagonally embedded in $\mathrm{SL}_2(q) \times \mathrm{SL}_2(q)$. Therefore, in terms of the O'Nan-Scott classification of primitive groups, this action is diagonal. Therefore, using the main result in [34], we deduce that the action is IBIS exactly when $G = G_0$.

Suppose now we are in the elliptic case.

Let $q = 2^f$ for some integer f and let $\phi : \mathbb{F}_q \to \mathbb{F}_q$ be the field automorphism defined by $x^{\phi} = x^2, \forall x \in \mathbb{F}_q$. We use the quadratic form

$$X_1 X_3 + X_2^2 + X_2 X_4 + \mu X_4^2,$$

for some $\mu \in \mathbb{F}_q$.

Assume first that $G = G_0 = \Omega_4^-(q) = \text{PSL}_2(q^2)$. The stabilizer of the action of G is $\text{Sp}_2(q) = \text{SL}_2(q)$, so the action is the one described in Example 3.9 in [31], and it is IBIS if $q \ge 4$. Suppose then that

$$G_0 < G \leq \operatorname{Aut}(G_0) = \operatorname{SO}_4^-(q)\operatorname{Aut}(\mathbb{F}_q).$$

Let α be a generator of the multiplicative group of the field \mathbb{F}_q . Using the quadratic form above, we see that the stabilizer of $\langle e_2 \rangle$ and $\langle e_4 \rangle$ in the whole group $\mathrm{SO}_4^-(q)\mathrm{Aut}(\mathbb{F}_q)$ is

$$\left\{ \begin{pmatrix} a & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & a^{-1} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 0 & a & 0 \\ 0 & 1 & 0 & 0 \\ a^{-1} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \mid a \in \mathbb{F}_q \setminus \{0\} \right\} \rtimes \operatorname{Aut}(\mathbb{F}_q).$$

With this, it is easy to verify that

$$(\langle e_2 \rangle, \langle e_4 \rangle, \langle e_1 + \alpha e_2 + e_4 \rangle)$$

is an irredundant base^m for $\mathrm{SO}_{4}^{-}(q)\mathrm{Aut}(\mathbb{F}_{q})$ of cardinality 3, where α is a generator of the multiplicative group of the field \mathbb{F} . To conclude that G is not IBIS, we show that G admits an irredundant base of cardinality at least 4 and then we invoke Lemma 2.3. If $G < \mathrm{SO}_{4}^{-}(q)\mathrm{Aut}(\mathbb{F}_{q})$, then either

- $\operatorname{SO}_4^-(q) \le G$, or
- $G \leq \Omega_4^-(q) \rtimes \langle \phi^i \rangle$ where i|f, or
- $G \leq \Omega_4^-(q) \rtimes \langle \iota \phi^i \rangle$ where i | f, f/i is even, and ι is the matrix

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

In the first case, using the description of the stabilizer of $\langle e_2 \rangle$ and $\langle e_4 \rangle$, we get $\iota \in G_{\langle e_2 \rangle, \langle e_1 + e_3 \rangle}$ and hence $(\langle e_2 \rangle, \langle e_4 \rangle, \langle e_1 + e_3 \rangle)$ is an irredundant chain having non-identity stabilizer. In the second and third case, since $(\langle e_2 \rangle, \langle e_4 \rangle, \langle e_1 + e_3 \rangle)$ is an irredundant base for $\Omega_4^-(q)$, the stabilizer of these three points is $\langle \phi^i \rangle$ or $\langle \iota \phi^i \rangle$ respectively. Hence, in all cases G admits a base of cardinality at least 4, so it is not IBIS.

In the proof of Lemma 6.1, we have used [34] in the hyperbolic case and [31] in the elliptic case (in this latter case, only to show that G_0 is IBIS). However, in both cases we could also have argued geometrically.

Proposition 6.2. The action of G on non-singular subspaces is IBIS if and only if $G = G_0 = \Omega_d^{\pm}(q)$ and $q \ge 4$.

Proof. From Lemma 6.1, we may suppose that $d \ge 6$. We use the hyperbolic quadratic form given by

$$X_1X_2 + X_3X_4 + \dots + X_{d-1}X_d,$$

and the elliptic quadratic form given by

$$X_1 X_{\frac{d}{2}+1} + X_2 X_{\frac{d}{2}+2} + \dots + X_{\frac{d}{2}-1} X_{d-1} + X_{\frac{d}{2}}^2 + X_{\frac{d}{2}} X_d + \mu X_d^2$$

We first show that if $G = G_0$ with $q \ge 4$, then G is IBIS. We show that all irredundant bases of G have cardinality d - 1. Let $\omega_1 = \langle v_1 \rangle, \ldots, \omega_{\ell} = \langle v_{\ell} \rangle$ be an irredundant base for G. Since $G = \Omega_d^{\pm}(q)$, G fixes the vectors v_1, \ldots, v_{ℓ} and hence it fixes pointwise the subspace generated by v_1, \ldots, v_{ℓ} . This immediately shows that $\ell \le d$. Suppose $d = \ell$. Then v_1, \ldots, v_d is a basis of the vector space V.

36

^mObserve that $\langle e_1 + \alpha e_2 + e_4 \rangle$ is non-singular since μ is chosen in such a way that the polynomial $T^2 + T + \mu$ is irreducible over \mathbb{F}_q .

The pointwise stabilizer of the subspace $\langle v_1, \ldots, v_{d-1} \rangle$ must be the identity group because $G = \Omega_d^{\pm}(q)$ does not contain any transvection. Therefore, we contradict the fact that $\omega_1, \ldots, \omega_\ell$ is an irredundant base. Finally, suppose $\ell \leq d-2$ and let $W = \langle v_1, \ldots, v_\ell \rangle$. Choose any (d-2)-dimensional subspace U of V containing W. Suppose first U^{\perp} is non-degenerate. The pointwise stabiliser of U in G is either $\Omega_2^+(q)$ or $\Omega_2^-(q)$. In both cases, since $q \geq 4$, the pointwise stabiliser of U in G is not the identity and it does move some non-singular vector, contradicting the fact that $\omega_1, \ldots, \omega_\ell$ is a base. Suppose next that U^{\perp} is totally isotropic and hence $U^{\perp} \leq U$. In this case, the pointwise stabilizer of U in G contains an element of order 2 which is the product of two communing transvectionsⁿ and hence again we contradict the fact that $\omega_1, \ldots, \omega_\ell$ is a base.

We now show that, if G is IBIS, then $G = G_0$ and $q \ge 4$. Here, we divide the case depending on G_0 . Suppose firstly we are in the hyperbolic case. Assume $q \geq 4$ and let $\alpha \in \mathbb{F}_q \setminus \{0,1\}$. Let $\omega_1 = \langle e_1 + e_2 \rangle$ and $\omega_2 = \langle e_1 + \alpha e_2 \rangle$. Observe that G_{ω_1,ω_2} stabilizes the subspace $\langle e_1, e_2 \rangle$ and fixes two non-singular vectors in this subspace. Therefore, G_{ω_1,ω_2} fixes pointwise $\langle e_1, e_2 \rangle$. Thus, G_{ω_1,ω_2} is a (d - d)2)-dimensional orthogonal group acting on the non-singular 1-subspaces. Hence, arguing inductively and using Lemma 6.1 as the base case of the induction, we deduce $G = G_0$. Assume now q = 2. In this case, either $G = G_0 = \Omega_d^+(2)$ or $G = SO_d^+(2)$. Let $\omega_1 = \langle e_1 + e_2 \rangle$, $\omega_2 = \langle e_1 + e_2 + e_3 \rangle$, $\omega_3 = \langle e_1 + e_2 + e_4 \rangle$ and $\omega_4 = \langle e_1 + e_3 + e_4 \rangle$. Observe that $G_{\omega_1,\omega_2,\omega_3,\omega_4}$ fixes pointwise $\langle e_1, e_2, e_3, e_4 \rangle$. Therefore, $G_{\omega_1,\omega_2,\omega_3,\omega_4}$ is a d-4 dimensional orthogonal group acting on the nonsingular 1-subspaces. When $d \equiv 0 \pmod{4}$, arguing inductively, we reduce to the case d = 8 (observe that, when q = 2, $\Omega_4^+(2)$ is solvable and hence it is of no concern in this work). We have verified that both $\Omega_8^+(2)$ and $SO_8^+(2)$ are not IBIS. When $d \equiv 2 \pmod{4}$, arguing inductively, we reduce to the case d = 6. We have verified that both $\Omega_6^+(2)$ and $\mathrm{SO}_6^+(2)$ are not IBIS. For instance, when $G = \mathrm{SO}_6^+(2)$, it can be verified that

 $\begin{array}{l} (\langle e_1 + e_2 \rangle, \, \langle e_1 + e_2 + e_6 \rangle, \, \langle e_1 + e_2 + e_3 \rangle, \, \langle e_1 + e_2 + e_4 \rangle, \, \langle e_5 + e_6 \rangle, \, \langle e_2 + e_3 + e_4 \rangle), \\ (\langle e_1 + e_2 \rangle, \, \langle e_1 + e_2 + e_6 \rangle, \, \langle e_1 + e_2 + e_3 \rangle, \, \langle e_1 + e_2 + e_4 \rangle, \, \langle e_2 + e_3 + e_4 \rangle) \end{array}$

are irredundant bases of length 6 and 5.

The elliptic case is analogous.

7. $Sp_4(2^a)$ containing a graph automorphism: Case (1.6)

In this case, $G_0 = \operatorname{Sp}_4(q) \leq G$ and G contains a graph automorphism swapping the two types of parabolic subgroups: stabilizers of 1-dimensional subspaces and stabilizers of 2-dimensional totally singular subspaces. The action of G can be identified with the action on pairs $\{p, \ell\}$, where p is a 1-dimensional subspace and ℓ is a 2-dimensional totally singular subspace with $p \in \ell$. It is not hard to verify that this action is not IBIS. To prove that, we fix the symplectic form given by the matrix

$$\begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}.$$

ⁿIndeed, if we are in the hyperbolic case and if $U^{\perp} = \langle e_1, e_d \rangle$, then we may take the transvection with respect to e_1 and to e_d and then take their product. Recall that each element of $\Omega_d^+(q)$ is the product of an even number of transvections in $\mathrm{SO}_d^+(q)$. The same argument works for the elliptic case.

Where 0 and I are the zero and identity 2×2 matrices respectively. Let

$$\begin{split} \omega_1 &= \omega_1' = \{ \langle e_1 \rangle, \langle e_1, e_2 \rangle \}, \\ \omega_2 &= \omega_2' = \{ \langle e_2 \rangle, \langle e_1, e_2 \rangle \}, \\ \omega_3 &= \{ \langle e_3 \rangle, \langle e_3, e_4 \rangle \}, \\ \omega_3' &= \{ \langle e_3 \rangle, \langle e_2, e_3 \rangle \}, \\ \omega_4' &= \{ \langle e_4 \rangle, \langle e_1, e_4 \rangle \}, \\ \omega_4 &= \omega_5' = \{ \langle e_1 + e_2 \rangle, \langle e_1 + e_2, e_3 + e_4 \rangle \}, \\ \omega_5 &= \omega_6' = \{ \langle e_2 + e_3 \rangle, \langle e_2 + e_3, e_1 + e_4 \rangle \}. \end{split}$$

Observe that $\langle e_1, e_2 \rangle$ is in common to ω_1 and ω_2 , and hence every element of G_{ω_1,ω_2} fixes $\langle e_1, e_2 \rangle$. Thus, G_{ω_1,ω_2} does not contain graph automorphisms swapping the two subspaces. With this in hands, using the same argument that we have used in Lemma 3.2, it is not hard to verify that $(\omega_1, \ldots, \omega_5)$ and $(\omega'_1, \ldots, \omega'_6)$ are two irredundant bases for G, hence G is not IBIS.

8. $P\Omega_8^+(q)$ containing triality: Cases (1.7) and (1.8)

In this section we deal with the case $G_0 = P\Omega_8^+(q)$ and G contains a triality.

Lemma 8.1. If G is as in Case (1.7), then G is not IBIS.

Proof. We have a geometric description of the domain Ω of G. We briefly recall a few basic facts about trialities, see [13, Chapter 8]. Let $V = \mathbb{F}_q^8$ be the orthogonal spaces associated to G. The 4-dimensional totally singular subspaces fall into two families (the greeks and the latins), where two subspaces W_1, W_2 are in the same class if dim (W_1) -dim $(W_1 \cap W_2)$ is even. Now, Ω can be identified with the collection of all triples $\{p, \pi_1, \pi_2\}$, where p is a point (that is, a 1-dimensional totally singular subspace) and π_1 and π_2 are solids containing p and of different type. Without loss of generality we may suppose that the hyperbolic quadric is

$$X_1X_5 + X_2X_6 + X_3X_7 + X_4X_8$$

Let

$$\begin{split} \omega_1 &= \{ \langle e_1 \rangle, \langle e_1, e_2, e_3, e_4 \rangle, \langle e_1, e_2, e_3, e_8 \rangle \}, \\ \omega_2 &= \{ \langle e_2 \rangle, \langle e_1, e_2, e_3, e_4 \rangle, \langle e_1, e_2, e_3, e_8 \rangle \}, \\ \omega_3 &= \{ \langle e_3 \rangle, \langle e_1, e_2, e_3, e_4 \rangle, \langle e_1, e_2, e_3, e_8 \rangle \}, \\ \omega_4 &= \{ \langle e_1 + e_2 + e_3 \rangle, \langle e_1, e_2, e_3, e_4 \rangle, \langle e_1, e_2, e_3, e_8 \rangle \}, \\ \omega'_3 &= \{ \langle e_1 + e_2 \rangle, \langle e_1, e_2, e_3, e_4 \rangle, \langle e_1, e_2, e_3, e_8 \rangle \}, \\ \omega'_4 &= \omega_3, \\ \omega'_5 &= \omega_4. \end{split}$$

As we have done in 7, since $\langle e_1, e_2, e_3, e_4 \rangle$ is in common to ω_1 and ω_2 , every element of G_{ω_1,ω_2} fixes $\langle e_1, e_2, e_3, e_4 \rangle$. This implies that $G_{\omega_1,\omega_2} \leq \Pr GO_8^+(q)$, that is, Gdoes not contain triality. With this remark, it is elementary to show that, when q > 2, $(\omega_1, \omega_2, \omega_3, \omega_4)$ and $(\omega_1, \omega_2, \omega'_3, \omega'_4, \omega'_5)$ are irredundant chains with

$$G_{\omega_1,\omega_2,\omega_3,\omega_4} = G_{\omega_1,\omega_2,\omega_3',\omega_4',\omega_5'}.$$

Therefore G is not IBIS by Lemma 2.1.

When q = 2, we have verified with a computer that b(G) = 3 and that G also admits bases of cardinality 4.

Dealing with case 1.8 is much more delicate and in fact we need to start with a lemma dealing with certain non-subspace actions. Our proof is quite geometric and we believe that it is interesting in its own.

Lemma 8.2. Let q be even and $G_0 = \operatorname{Sp}_6(q)$ or let q be odd and $G_0 = \Omega_7(q)$. Let G be an almost simple primitive group with socle G_0 such that the stabilizer of a point in G_0 is maximal subgroup in the Aschbacher class C_9 isomorphic to $G_2(q)$. Then G is not IBIS.

Proof. The proofs of Lemma 3.1 (when $G_0 = \text{Sp}_6(q)$) and Proposition 3.2 (when $G_0 = \Omega_7(q)$) in [8] show that G has base size 4. The same proofs can also be used to show that G has irredundant bases of cardinality 5.

Using Lemma 8.2 we may now deal with (1.8).

Lemma 8.3. If G is as in Case (1.8), then G is not IBIS.

Proof. Here $G_0 = P\Omega_8^+(q)$ and the stabilizer of a point in G_0 is isomorphic to $G_2(q)$: the centralizer of a graph automorphism of order 3.

Let Ω be the domain of G and fix $\alpha \in \Omega$. Thus $(G_0)_{\alpha} = \mathbf{C}_{G_0}(\tau) \cong G_2(q)$, for some graph automorphism τ of order 3. From [27], we deduce that the lattice of subgroups of G_0 containing $(G_0)_{\alpha}$ consists of five subgroups: G_0 , $(G_0)_{\alpha}$ and three maximal subgroups M_1, M_2, M_3 of G_0 with $M_1 \cong M_2 \cong M_3 \cong \Omega_7(q)$ when q is odd and $M_1 \cong M_2 \cong M_3 \cong \mathrm{Sp}_6(q)$ when q is even. These embeddings arise from the spin representations of orthogonal groups.

The group G_0 is imprimitive on Ω and each subgroup M_i gives rise to a system of imprimitivity Σ_i for the action of G_0 on Ω . Clearly, G does not preserve Σ_i and in fact G acts transitively on the three systems of imprimitivity $\Sigma_1, \Sigma_2, \Sigma_3$ making the action of G primitive. Let $\theta_i : \Omega \to \Sigma_i$ be the natural projection, that is, $\beta_i(\beta)$ is the unique part in the partition Σ_i containing the point β . We now consider the map

$$\begin{aligned} \theta : \Omega \to \Sigma_1 \times \Sigma_2 \times \Sigma_3, \\ \beta \mapsto \theta(\beta) &= (\theta_1(\beta), \theta_2(\beta), \theta_3(\beta)). \end{aligned}$$

Since G is primitive on Ω , θ is injective. Therefore, when necessary, we identify Ω with its image under θ .

Let $i \in \{1, 2, 3\}$ and let $\{1, 2, 3\} \setminus \{i\} = \{j, k\}$. We claim that the mapping $\Omega \to \Sigma_j \times \Sigma_k$ defined by $\beta \mapsto (\theta_j(\beta), \theta_k(\beta))$ is a bijection. In other words, in the function θ , each coordinate is uniquely determined by the remaining two coordinates. Firstly, note that

$$\begin{aligned} |\Omega| &= |\mathbf{P}\Omega_8^+(q) : G_2(q)| = q^6 (q^4 - 1)^2, \\ |\Sigma_i| &= \begin{cases} |\mathbf{P}\Omega_8^+(q) : \Omega_7(q)| = q^3 (q^4 - 1), & \text{when } q \text{ is odd,} \\ |\mathbf{P}\Omega_8^+(q) : \mathbf{Sp}_6(q)| = q^3 (q^4 - 1), & \text{when } q \text{ is even.} \end{cases} \end{aligned}$$

Therefore, $|\Omega| = |\Sigma_j \times \Sigma_k|$. Hence, it suffices to show that $\beta \mapsto (\theta_j(\beta), \theta_k(\beta))$ is injective. Let $\beta, \beta' \in \Omega$ with $(\theta_j(\beta), \theta_k(\beta)) = (\theta_j(\beta'), \theta_k(\beta'))$. Set $\lambda = \theta_j(\beta)$ and

 $\lambda' = \theta_k(\beta)$. From the classification of the maximal factorizations of the simple group $P\Omega_8^+(q)$ (see first line of Table 4 in [32]), we deduce that

$$G_0 = (G_0)_{\lambda} (G_0)_{\lambda}$$

and hence simply checking the cardinality, we deduce that

$$(G_0)_{\beta} = (G_0)_{\lambda} \cap (G_0)_{\lambda'} = (G_0)_{\beta'}$$

The primitivity of G implies that $(G_0)_\beta$ can fix at most one point and hence $\beta = \beta'$.

Let $\theta(\alpha) = (\lambda_1, \lambda_2, \lambda_3)$ and let X be the setwise stabilizer of λ_1 in G. Then, X is an almost simple group with socle either $\text{Sp}_6(q)$ or $\Omega_7(q)$ and the action of X on λ_1 is the one investigated in Lemma 8.2. Therefore, there exist $\alpha = \alpha_1, \alpha_2, \alpha_3, \alpha_4 \in \lambda_1$ and $\alpha = \alpha'_1, \alpha'_2, \alpha'_3, \alpha'_4, \alpha'_5 \in \lambda_1$ forming irredundant bases of cardinality 4 and 5 for the action of X on λ_1 . We claim that these are also irredundant chains for the action of G on Ω with

$$G_{\alpha_1,\alpha_2,\alpha_3,\alpha_4} = G_{\alpha'_1,\alpha'_2,\alpha'_3,\alpha'_4,\alpha'_5} = G_{(\lambda_1)},$$

where $G_{(\lambda_1)}$ is the pointwise stabilizer of λ_1 in G. To prove this it suffices to show that $G_{\alpha_1,\alpha_2,\alpha_3,\alpha_4}$ and $G_{\alpha'_1,\alpha'_2,\alpha'_3,\alpha'_4,\alpha'_5}$ fix setwise λ_1 and then use the fact these points form two irredundant bases for the action of $G_{\{\lambda_1\}}$ on λ_1 .

Clearly,

$$G > G_{\alpha_1} > G_{\alpha_1,\alpha_2} > G_{\alpha_1,\alpha_2,\alpha_3} > G_{\alpha_1,\alpha_2,\alpha_3,\alpha_4},$$

because the analogous chain of stabilizers in X is strictly decreasing. Let $Y = G_{\alpha_1,\alpha_2,\alpha_3,\alpha_4}$. Observe that, for each *i*, the first coordinate of $\theta(\alpha_i)$ is λ_1 , that is,

$$\theta_1(\alpha_1) = \theta_1(\alpha_2) = \theta_1(\alpha_3) = \theta_1(\alpha_4)$$

Assume that Y contains some permutation g moving Σ_1 , say $\Sigma_1^g = \Sigma_2$. Then, since g fixes α_i for each i, we deduce

$$\theta_2(\alpha_1) = \theta_2(\alpha_2) = \theta_2(\alpha_3) = \theta_2(\alpha_4).$$

However, since the mapping $\beta \mapsto (\theta_1(\beta), \theta_2(\beta))$ is bijective, this implies $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4$, which is clearly a contradiction. This has shown that every element of Y fixes Σ_1 (and eventually permutes Σ_2 with Σ_3). Therefore, Σ_1 is a system of imprimitivity for the action of Y. Since $\alpha_1 \in \lambda_1$ and since Y fixes α_1 , we deduce that Y fixes setwise λ_1 . With the same argument, we can show that $G_{\alpha'_1,\alpha'_2,\alpha'_3,\alpha'_4,\alpha'_5}$ fixes setwise λ_1 . Therefore, G has two irredundant bases of cardinality 4 and 5 respectively, and hence it is not IBIS.

9. Almost simple groups having socle $E_6(q)$ or $E_7(q)$: case (1.9) AND (1.10)

In this case, $G_0 = E_6(q) \leq G$ or $G_0 = E_7(q) \leq G$, and the action is one of those described in cases (1.9) and (1.10).

Proposition 9.1. Let G be an almost simple group on Ω with socle $G_0 = E_6(q)$, where the action of G_0 on Ω is permutation equivalent to the action on the right cosets of a parabolic subgroup labeled P_1 or P_6 . Then G is not IBIS.

Proof. Observe that a graph automorphism of $G_0 = E_6(q)$ fuses the G_0 -conjugacy classes of parabolic subgroups labeled P_1 and P_6 .

By [9], b(G) = 6 and hence, by Lemma 2.2, it suffices to show that G_0 has an irredundant base of length at least 7.

40

Let V be the 27-dimensional natural module for the covering group of G_0 over the finite field \mathbb{F}_q and let \mathcal{P} be the collection of all 1-dimensional subspaces of V. From [19], the 1-dimensional subspaces of V are partitioned into three parts

$$\mathcal{P}=\mathcal{W}\cup\mathcal{G}\cup\mathcal{B},$$

namely the set of white points \mathcal{W} , the set of gray points \mathcal{G} and the set of black points \mathcal{B} . In [19], it is noticed that the action of G_0 on the right cosets of a maximal parabolic subgroup of type P_1 (or P_6) is permutation equivalent to the action of G_0 on the set of white points \mathcal{W} of V.^o In particular, without loss of generality, we may suppose that $\Omega = \mathcal{W}$.

Let \mathcal{W}_6 be the collection of all 6-dimensional subspaces W of V consisting only of white points, that is, each 1-dimensional subspace of W is white. From [19, (P.3), page 470], we see that $\mathcal{W}_6 \neq \emptyset$ and that G_0 acts transitively on \mathcal{W}_6 . Let $W \in \mathcal{W}_6$ and let $H = (G_0)_{\{\Delta\}}$ be the setwise stabilizer of

$$\Delta = \{ \omega \in \Omega \mid \omega \le W \},\$$

in other words, $\Delta \subseteq \Omega$ consists of the white points contained in W. Again from [19, (P.3), page 470], we deduce that the permutation group induced by H on Δ is permutation equivalent to an almost simple group X having socle $Y = \text{PSL}_6(q)$ in its natural action on the projective points of a 6-dimensional vector space \mathbb{F}_q^6 over \mathbb{F}_q . In particular, we may identify the action of $H = G_{\{\Delta\}}$ on Δ with the natural action of X on the projective points. We use this identification in what follows.

Now, let $\delta_1 = \langle v_1 \rangle, \ldots, \delta_6 = \langle v_6 \rangle$ be elements of Δ such that the vectors v_1, \ldots, v_6 are linearly independent in \mathbb{F}_q^6 . When q > 2, let $\delta_7 = \langle v_1 + \cdots + v_6 \rangle$. Moreover, let $\ell = 6$ when q = 2 and let $\ell = 7$ when q > 2. Now,

$$G_0 > (G_0)_{\delta_1} > (G_0)_{\delta_1, \delta_2} > \dots > (G_0)_{\delta_1, \delta_2, \dots, \delta_\ell}.$$

In particular, when q > 2, we may extend the sequence $(\delta_1, \ldots, \delta_\ell)$ to an irredundant base for G_0 of length at least 7. When q = 2, by definition $\ell = 6$ and, moreover, $(G_0)_{\delta_1, \delta_2, \ldots, \delta_\ell}$ is the pointwise stabilizer of Δ . Since this subgroup contains the unipotent radical of the parabolic subgroup $(G_0)_{\delta_1}{}^p$, we deduce that $(G_0)_{\delta_1, \delta_2, \ldots, \delta_\ell} \neq$ 1, hence we may extend the sequence $(\delta_1, \ldots, \delta_\ell)$ to an irredundant base for G_0 of length at least 7.^q

Proposition 9.2. Let G be an almost simple group on Ω with socle $G_0 = E_7(q)$, where the action of G_0 on Ω is permutation equivalent to the action on the right cosets of a parabolic subgroup labeled P_7 . Then G is not IBIS.

Proof. From [9], G has a base of cardinality 6. Therefore, from Lemma 2.2, it suffices to prove that G_0 has irredundant bases of cardinality larger than 6.

Let Ω be the set of cosets of P_7 in G_0 and let $\omega \in \Omega$. From $[37]^r$, we have

$$|\Omega| = \frac{(q^{14} - 1)(q^9 + 1)(q^5 + 1)}{q - 1}$$

^oIncidentally, this fact was first observed by Freudenthal in [23].

^pSee [19] or [23].

^qWe observe that the argument when q = 2 also applies to the case q > 2.

^rVasilev [37] employs a distinct labeling system for the parabolic subgroups compared to that

of [3]. Specifically, parabolic subgroups denoted as P_7 in [3] are identified as P_1 in [37].

Moreover, again from [37], we see that G_0 has (permutational) rank 4 in its action on Ω , and that G_0 has a suborbit of cardinality

(18)
$$n_2 = \frac{q(q^9 - 1)(q^8 + q^4 + 1)}{q - 1}$$

Therefore, $(G_0)_{\omega}$ has an orbit, Δ say, having cardinality n_2 . Moreover, from [37], we deduce that the action of $(G_0)_{\omega}$ on Δ is faithful. From this, we have that G_0 admits an irredundant base $(\omega_0 = \omega, \omega_1, \ldots, \omega_{\ell-1})$, where $\omega_1, \ldots, \omega_{\ell-1} \in \Delta$. This implies

$$n_2^{\ell-1} = |\Delta|^{\ell-1} > |(G_0)_{\omega}| = |P_7| = \frac{|G_0|}{|\Omega|}$$
$$= \frac{1}{d}q^{63}(q^9 - 1)(q^{12} - 1)(q^5 - 1)(q^8 - 1)(q^6 - 1)(q^2 - 1)(q - 1)$$

where $d = \gcd(2, q - 1)$. Using (18), it can be verified that this implies $\ell \geq 7$. \Box

References

- R. F. Bailey, Uncoverings-by-bases for base-transitive permutation groups, Des. Codes Cryptogr. 41 (2006), no. 2, 153–176.
- [2] J. N. Bray, D. F. Holt, C. M. Roney-Dougal, The maximal subgroups of the low-dimensional finite classical groups, Cambridge: Cambridge University Press, 2013.
- [3] N. Bourbaki, Groupes et Algebrès de Lie (Chapters 4, 5 and 6), Hermann, Paris, 1968.
- [4] T. C. Burness, Fixed point ratios in actions of finite classical groups. I, J. Algebra 309 (2007), 69–79.
- [5] T. C. Burness, Fixed point ratios in actions of finite classical groups. II, J. Algebra 309 (2007), 80–138.
- [6] T. C. Burness, Fixed point ratios in actions of finite classical groups. III, J. Algebra 309 (2007), 693-748.
- [7] T. C. Burness, Fixed point ratios in actions of finite classical groups. IV, J. Algebra 309 (2007), 749–788.
- [8] T. C. Burness, On base sizes for actions of finite classical groups, J. Lond. Math. Soc. (2) 75 (2007), 545–562.
- [9] T. C. Burness, On base sizes for almost simple primitive groups, J. Algebra **516** (2018), 38–74.
- [10] T. C. Burness, R. M. Guralnick, J. Saxl, On base sizes for symmetric groups, Bull. Lond. Math. Soc. 43 (2011), 386–391.
- [11] T. C. Burness, M. W. Liebeck, A. Shalev, Base sizes for simple groups and a conjecture of Cameron, Proc. Lond. Math. Soc. (3) 98 (2009), 116–162.
- [12] T. C. Burness, E. A. O'Brien, R. A. Wilson, Base sizes for sporadic simple groups, Israel J. Math. 177 (2010), 307–333.
- [13] P. J. Cameron, *Projective and Polar Spaces*, University of London, Queen Mary and Westfield college, 1992.
- [14] P. J. Cameron, W. M. Kantor, Random permutations: some group-theoretic aspects, Combin. Probab. Comput. 2 (1993) 257–262.
- [15] P. J. Cameron, *Permutation Groups*, London Mathematical Society Student Texts, Cambridge University Press, 1999.
- [16] P. J. Cameron, D. G. Fon-Der-Flaass, Bases for permutation groups and matroids, *Eur. J. Comb.* 16 (1995), 537–544.
- [17] P. J. Cameron, "Some open problems on permutation group" in Groups, Combinatorics and Geometry. Proceedings of the L.M.S. Durham Symposium, Held July 5–15, 1990 in Durham, UK, Cambridge University Press, Cambridge, MA, 1992, 340–350.
- [18] P. J. Cameron, W. M. Kantor, "Random permutations: Some group-theoretic aspect" in Combinatorics, Geometry and Probability. A Tribute to Paul Erdős. Proceedings of the Conference Dedicated to Paul Erdős on the Occasion of His 80th Birthday, Cambridge, UK, 26 March 1993, Cambridge University Press, Cambridge, MA, 1997, 139–144.

IBIS LARGE BASE

- [19] A. M. Cohen, B. N. Cooperstein, The 2-spaces of the standard $E_6(q)$ -module, Geom. Dedicata **25** (1988), 467–480.
- [20] F. Dalla Volta, F. Mastrogiacomo, P. Spiga, On the cardinality of irredundant and minimal bases of finite permutation groups, J. Algebr. Comb., to appear.
- [21] J. D. Dixon, B. Mortimer, *Permutation groups*, Graduate Texts in Mathematics 163, Springer-Verlag, New York, 1996.
- [22] R. H. Dye, Interrelations of Symplectic and Orthogonal Groups in Characteristic Two, J. Algebra 59 1979, 202–221.
- [23] H. Freudenthal, Beziehungen der E_7 und E_8 zur Oktavenebene I., Indagationes Math. 16 1954, 218–230.
- [24] N. Gill, B. Lodá, P. Spiga, On the height and relational complexity of a finite permutation group, Nagoya Math. J. 246 (2022), 372–411.
- [25] S. Guest, A. Previtali, P. Spiga, A remark on the permutation representations afforded by the embeddings of $O_{2m}^{\pm}(2^f)$ in $\operatorname{Sp}_{2m}(2^f)$, Bull. Aust. Math. Soc. 89 (2014), 331–336.
- [26] C. Jordan, Traité des Substitutions et des Équations Algébriques, Gauthier-Villars, Paris, 1870.
- [27] P. Kleidman, The maximal subgroups of the finite 8-dimensional orthogonal groups $P\Omega_8^+(q)$ and of their automorphism groups, J. Algebra **110** (1987), 173–242.
- [28] P. Kleidman, M. Liebeck, The Subgroup Structure of the Finite Classical Groups, Lond. Math. Soc. Lect. Note Ser. 129, Cambridge University Press, Cambridge, MA, 1990.
- [29] S. Lang, Algebra. 3rd revised ed., Graduate Texts in Mathematics 211, 2002, New York, NY: Springer.
- [30] M. Lee, Primitive almost simple IBIS groups with sporadic socle, arXiv:2302.01521.
- [31] M. Lee, P. Spiga, A classification of finite primitive IBIS groups with alternating socle, J. Group Theory 26 (2023), 915–930.
- [32] M. W. Liebeck, C. E. Praeger and J. Saxl, The maximal factorizations of the finite simple groups and their automorphism groups, *Mem. Amer. Math. Soc.* 86 (1990).
- [33] M. W. Liebeck, A. Shalev, Simple groups, permutation groups, and probability, J. Amer. Math. Soc. 12 (1999), 497–520.
- [34] A. Lucchini, M. Morigi, M. Moscatiello, Primitive permutation IBIS groups, J. Combin. Theory Ser. A 184 (2021), Paper No. 105516.
- [35] A. Lucchini, D. Malinin, IBIS soluble linear groups, Comm. Algebra 52 (2024), 457-465.
- [36] C. E. Praeger, The inclusion problem for finite primitive permutation groups, Proc. Lond. Math. Soc. (3) 60 (1990), 68–88.
- [37] A. V. Vasilyev, Minimal permutation representations of finite simple exceptional groups of type E₆, E₇ and E₈, Algebra and Logic 36 (1997), 302–3010.

FABIO MASTROGIACOMO, DIPARTIMENTO DI MATEMATICA "FELICE CASORATI", UNIVERSITY OF PAVIA, VIA FERRATA 5, 27100 PAVIA, ITALY *Email address*: fabio.mastrogiacomo01@universitadipavia.it

PABLO SPIGA, DIPARTIMENTO DI MATEMATICA PURA E APPLICATA, UNIVERSITY OF MILANO-BICOCCA, VIA COZZI 55, 20126 MILANO, ITALY Email address: pablo.spiga@unimib.it