

# High-probability minimax lower bounds

Tianyi Ma\*, Kabir A. Verchand\*,<sup>◦</sup> and Richard J. Samworth\*

\*Statistical Laboratory, University of Cambridge

<sup>◦</sup>Industrial and Systems Engineering, Georgia Institute of Technology

June 21, 2024

## Abstract

The minimax risk is often considered as a gold standard against which we can compare specific statistical procedures. Nevertheless, as has been observed recently in robust and heavy-tailed estimation problems, the inherent reduction of the (random) loss to its expectation may entail a significant loss of information regarding its tail behaviour. In an attempt to avoid such a loss, we introduce the notion of a minimax quantile, and seek to articulate its dependence on the quantile level. To this end, we develop high-probability variants of the classical Le Cam and Fano methods, as well as a technique to convert local minimax risk lower bounds to lower bounds on minimax quantiles. To illustrate the power of our framework, we deploy our techniques on several examples, recovering recent results in robust mean estimation and stochastic convex optimisation, as well as obtaining several new results in covariance matrix estimation, sparse linear regression, nonparametric density estimation and isotonic regression. Our overall goal is to argue that minimax quantiles can provide a finer-grained understanding of the difficulty of statistical problems, and that, in wide generality, lower bounds on these quantities can be obtained via user-friendly tools.

## 1 Introduction

Over the last 15 years or so, minimax lower bounds, which used to be regarded as primarily a branch of information theory, have entered the statistical mainstream. This transition has been brought about by an increased awareness of what they have to offer to statisticians, in terms of providing a benchmark against which we can compare the performance of different statistical procedures. Their increased prominence can be attributed in part to the surge of interest in high-dimensional statistical problems over the last 25 years, which has encouraged statisticians to move away from traditional diverging sample size asymptotics (with other problem parameters held fixed) and to focus instead on finite-sample performance bounds.

To fix the decision-theoretic foundations that underpin this work, suppose that we are interested in estimating a parameter  $\theta$  taking values in a parameter space  $\Theta$ , based on data  $X$  taking values in a measurable space  $\mathcal{X}$ . The notation here might suggest that  $\Theta$  is a subset of a Euclidean space, so it is worth emphasising that we may be interested in

estimating more complicated objects like density or regression functions, in which case  $\Theta$  would be an appropriate function space. Likewise, the data need not be a single observation, but could be a sample of size  $n$ , for example. Let  $\widehat{\Theta}$  denote the set of estimators of  $\theta \in \Theta$ , i.e. the set of measurable functions from  $\mathcal{X}$  to  $\Theta$ . Our loss function is a measurable function  $L : \Theta \times \Theta \rightarrow [0, \infty)$ .

Since  $L(\widehat{\theta}(X), \theta)$  is random when  $\widehat{\theta} \in \widehat{\Theta}$ , it is common to work with the *risk function*

$$R(\widehat{\theta}, \theta) := \mathbb{E}_\theta L(\widehat{\theta}(X), \theta),$$

a deterministic quantity. A typical ideal goal, then, is to find an estimator  $\widehat{\theta}$  that achieves the *minimax risk* in the sense that

$$\sup_{\theta \in \Theta} R(\widehat{\theta}, \theta) = \inf_{\widetilde{\theta} \in \widehat{\Theta}} \sup_{\theta \in \Theta} R(\widetilde{\theta}, \theta).$$

In this paper, we revisit one of the early choices in the development of this paradigm, namely the reduction of the random  $L(\widehat{\theta}(X), \theta)$  to the deterministic risk function. This simplification may fail to tell the whole story regarding the variability and tail behaviour of the loss function for different data realisations. For instance, consider the simple problem of estimating a univariate mean  $\theta$  with respect to squared error loss  $L(\widehat{\theta}, \theta) := (\widehat{\theta} - \theta)^2$ , based on a sample  $X_1, \dots, X_n$  from a distribution having variance at most  $\sigma^2$ . It is well-known that the minimax risk in this problem is  $\sigma^2/n$ , and is attained by the sample mean  $\bar{X} := n^{-1} \sum_{i=1}^n X_i$ . On the other hand, [Catoni \(2012, Proposition 6.2\)](#) proved that the sample mean has undesirable tail behaviour in the sense that for each  $\delta \leq 1/e$ , there exists a distribution with mean  $\theta$  and variance  $\sigma^2$  such that with probability at least  $\delta$ ,

$$L(\bar{X}, \theta) \geq \frac{\sigma^2}{n\delta} \left(1 - \frac{e\delta}{n}\right)^{n-1} > \frac{\sigma^2}{en\delta}.$$

This lower bound admits an interpretation as a lower bound on the  $(1 - \delta)$ th quantile of  $L(\bar{X}, \theta)$ . By contrast, the performance of the median-of-means estimator  $\widehat{\theta}_\delta^{\text{MoM}}$  ([Nemirovski and Yudin, 1983, p. 243](#)) can be controlled via an upper bound on the  $(1 - \delta)$ th quantile of  $L(\widehat{\theta}_\delta^{\text{MoM}}, \theta)$ : with probability at least  $1 - \delta$ ,

$$L(\widehat{\theta}_\delta^{\text{MoM}}, \theta) \leq C \cdot \frac{\sigma^2 \log(1/\delta)}{n},$$

for some universal constant  $C > 0$ .

This stark gap in the dependence on  $\delta$  in the above bounds, and hence the guarantees that can be made with high confidence for the two estimators, makes clear that the reduction to the minimax risk may entail significant information loss. We can retain this information by defining, for  $\delta \in (0, 1]$ , the *minimax  $(1 - \delta)$ th quantile*

$$\mathcal{M}(\delta) := \inf_{\widehat{\theta} \in \widehat{\Theta}} \sup_{\theta \in \Theta} \inf \left\{ r \in [0, \infty) : \mathbb{P}_\theta \{ L(\widehat{\theta}(X), \theta) > r \} \leq \delta \right\}.$$

It is often the case that upper bounds on the minimax risk are obtained by finding an estimator  $\widehat{\theta} \in \widehat{\Theta}$  and integrating a tail bound of the form

$$\sup_{\theta \in \Theta} \mathbb{P}_\theta \{ L(\widehat{\theta}(X), \theta) > r_\delta \} \leq \delta;$$

in such circumstances, we immediately have  $\mathcal{M}(\delta) \leq r_\delta$ , so upper bounds on the minimax quantile are frequently implicit in upper bound arguments for the minimax risk. The main aim of this work is to introduce user-friendly tools to facilitate lower bounds on  $\mathcal{M}(\delta)$ ; we give a variety of examples where these lower bounds match corresponding upper bounds up to constants, and thereby provide a finer-grained understanding of these statistical challenges.

As two further illustrations of the benefits of minimax quantiles, we first remark that the basis for the analysis of several estimators is a union bound over (potentially exponentially) many different events. Examples include the sample covariance matrix as an estimator of its population counterpart in operator norm (Vershynin, 2018, Theorem 4.6.1) and the Tukey median (Tukey, 1975; Chen, Gao and Ren, 2018), as well as tournament and skeleton estimators (Devroye and Lugosi (2001, Chapters 6 and 7) and Chen, Gao and Ren (2016)). In such cases, it is the behaviour of extreme quantiles of simpler estimators that govern ultimate performance. Second, consider the problem of robust mean estimation with respect to squared error loss in a Huber contamination model (Huber, 1964). There, the unboundedness of the loss function, combined with the fact that there is a positive probability with which all data are contaminated, ensures that the minimax risk is infinite, so uninformative from the point of view of comparing estimators; on the other hand, in Corollary 11 we will determine (the finite)  $\mathcal{M}(\delta)$  for a broad interval of quantile levels  $\delta$ .

In the high-probability minimax framework, an estimator achieving an upper bound on the minimax quantile  $\mathcal{M}(\delta)$  may in principle depend on  $\delta$ . Indeed, this is the case in the median-of-means example discussed above. Nevertheless, there is the potential for a single estimator  $\hat{\theta}$  to attain the minimax quantile (up to constants) simultaneously over a wide range of quantile levels. Such adaptation to the quantile level may be interpreted as a further justification for the use of  $\hat{\theta}$ .

To outline our contributions in greater detail, in Section 2, we formally define the minimax quantile  $\mathcal{M}(\delta)$ , and a closely-related minorant  $\mathcal{M}_-(\delta)$ , which we call the *lower minimax quantile*; see Definitions 1 and 3 below. This latter quantity turns out to be especially amenable to analysis via standard tools in the minimax lower bound literature, and in Lemma 5 and Lemma 7 respectively, we apply Le Cam’s two-point lemma and Fano’s inequality to obtain lower bounds on the lower minimax quantile. In fact, in Theorem 8, we provide a reduction from a local minimax lower bound on the risk to a lower bound on  $\mathcal{M}_-(\delta)$ , so the full range of minimax lower bound techniques, including Assouad’s lemma, can be applied.

In Section 3, we provide a suite of examples to illustrate the power of the framework and give a sense of the breadth of problems to which it may be applied. Indeed, the settings we consider include (robust) multivariate mean estimation, covariance matrix estimation, sparse linear regression, nonparametric density estimation, isotonic regression and stochastic convex optimisation. In all of these examples, we determine the minimax quantile up to multiplicative constants\*. One of the main messages we aim to convey through this selection of examples is that it is relatively straightforward to apply our results from Section 2, and thereby to enhance the strength of the justification for many statistical procedures through the minimax quantile paradigm.

Proofs of our main results in Section 2 are given in Section 4, with the exception of one

---

\*Strictly speaking, in sparse linear regression, these constants depend on a restricted eigenvalue condition, while in nonparametric density estimation, they depend on smoothness parameter of a Hölder class.

very short corollary. The Appendix comprises some standard minimax risk lower bounds, as well as the statements and proofs of several auxiliary lemmas.

## 1.1 Related work

To the best of our knowledge, our work represents the first systematic study of high-probability minimax lower bounds, though isolated results in certain settings have started to appear in recent years. The importance of the dependence on the quantile level of upper bounds on the minimax quantile has been emphasised in the literature on robust and heavy-tailed estimation (e.g. [Catoni, 2012](#); [Lugosi and Mendelson, 2019b](#); [Hopkins, 2020](#)). On the other hand, lower bounds are scarcer, though [Devroye et al. \(2016\)](#) established a lower bound on  $\mathcal{M}_-(\delta)$  in univariate mean estimation, and [Depersin and Lecué \(2022a\)](#) extended this result to the multivariate case using techniques developed by [Lecué and Mendelson \(2013\)](#) based on the Gaussian shift theorem. In particular, our Proposition 10 provides an alternative, streamlined proof of [Depersin and Lecué \(2022a, Theorem 4\)](#).

The quantity  $\mathcal{M}_-(\delta)$  was introduced by [Donoho \(1994\)](#) (albeit without our moniker) to study minimax confidence interval length in the setting of linear functional estimation. He showed that it could be expressed in terms of a modulus of continuity (see also [Cai, 2012, Section 5](#)). Recently, in a different yet still univariate setting, [Wang and Ramdas \(2024\)](#) and [Duchi and Haque \(2024\)](#) invoked information-theoretic techniques to obtain lower bounds on the minimax length of an anytime-valid  $(1 - \delta)$ -level confidence interval. [Tsybakov \(2009, Theorem 2.3\)](#) established a lower bound on  $\mathcal{M}_-(\delta)$  in the context of univariate nonparametric regression.

In stochastic convex optimisation, [Davis et al. \(2021\)](#) considered heavy-tailed stochastic gradients and design methods with logarithmic dependence on the quantile level. Very recently, [Carmon and Hinder \(2024\)](#) introduced the notion of a minimax quantile in this setting and applied information-theoretic tools to obtain lower bounds that match their upper bounds up to multiplicative universal constants. In Section 3.6, we provide a different proof of their lower bound based on our general theory developed in Section 2 in conjunction with Le Cam’s two-point method.

## 1.2 Notation

For  $n \in \mathbb{N}$ , we define  $[n] := \{1, \dots, n\}$ . We let  $\mathbb{S}^{d-1}$  denote the unit Euclidean sphere in  $\mathbb{R}^d$  and for  $R > 0$ , write  $\mathbb{B}_2^d(R)$  for the closed Euclidean ball in  $\mathbb{R}^d$  of radius  $R$ . For  $v = (v_1, \dots, v_d)^\top \in \mathbb{R}^d$  and  $q > 0$ , we let  $\|v\|_q := (\sum_{j=1}^d |v_j|^q)^{1/q}$ ,  $\|v\|_0 := \sum_{j=1}^d \mathbb{1}_{\{v_j \neq 0\}}$  and  $\|v\|_\infty := \max_{j \in [d]} |v_j|$ . We let  $\mathcal{S}_+^{d \times d}$  and  $\mathcal{S}_{++}^{d \times d}$  denote the cones of symmetric, positive semi-definite and symmetric, positive definite matrices in  $\mathbb{R}^{d \times d}$  respectively. For symmetric matrices  $A, B \in \mathbb{R}^{d \times d}$ , we write  $A \succeq B$  if  $A - B \in \mathcal{S}_+^{d \times d}$  and  $A \preceq B$  if  $B - A \in \mathcal{S}_+^{d \times d}$ . For a matrix  $A = (A_{ij})_{i \in [n], j \in [d]} \in \mathbb{R}^{n \times d}$ , we let  $\|A\|_\infty := \max_{i \in [n], j \in [d]} |A_{ij}|$  and  $\|A\|_{\text{op}} := \sup_{u \in \mathbb{S}^{n-1}, v \in \mathbb{S}^{d-1}} u^\top A v$ . Given  $(a_1, \dots, a_d)^\top \in \mathbb{R}^d$ , let  $\text{diag}(a_1, \dots, a_d) \in \mathbb{R}^{d \times d}$  denote the diagonal matrix with entries  $a_1, \dots, a_d$ . We let  $I_d := \text{diag}(1, \dots, 1)$  denote the identity matrix in  $\mathbb{R}^{d \times d}$  and, for  $j \in [d]$ , let  $e_j \in \mathbb{R}^d$  denote the  $j$ th standard basis vector. Given a measurable space  $(\mathcal{X}, \mathcal{A})$ , we will denote the space of all probability measures on  $\mathcal{X}$  as  $\mathcal{Q}(\mathcal{X})$ . For  $\omega = (\omega_1, \dots, \omega_d)^\top, \omega' = (\omega'_1, \dots, \omega'_d)^\top \in \{0, 1\}^d$ , we define their Hamming distance as  $d_H(\omega, \omega') :=$

$\sum_{j=1}^d \mathbb{1}_{\{\omega_j \neq \omega'_j\}}$ . We let  $\text{Pow}(S)$  denote the power set of an arbitrary set  $S$ .

For  $a, b \in \mathbb{R}$ , we write  $a \gtrsim b$  if there exists a universal constant  $c > 0$  such that  $a \geq cb$ , and write  $a \lesssim b$  if there exists a universal constant  $C > 0$  such that  $a \leq Cb$ . We write  $a \asymp b$  if both  $a \lesssim b$  and  $a \gtrsim b$ . Similarly, we write  $a \asymp_\alpha b$  if there exist  $c(\alpha), C(\alpha) > 0$ , depending only on  $\alpha$ , such that  $c(\alpha)b \leq a \leq C(\alpha)b$ .

## 2 Problem setup and main results

Let  $(\Theta, d)$  be a non-empty pseudo-metric<sup>†</sup> space, equipped with the topology induced by the open balls in  $d$  and the corresponding Borel  $\sigma$ -algebra. It will be convenient for some of our examples to consider a slightly more general setting than that outlined in the introduction. In particular, for  $\theta \in \Theta$ , let  $\mathcal{P}_\theta$  denote a family of probability measures on a measurable space  $(\mathcal{X}, \mathcal{A})$ , and define  $\mathcal{P}_\Theta := \{\mathcal{P}_\theta : \theta \in \Theta\}$ . Although in many applications, each  $\mathcal{P}_\theta$  will consist of a single distribution, the additional generality allows us to cover heavy-tailed and Huber contamination models with our theory.

Let  $g : [0, \infty) \rightarrow [0, \infty)$  be an increasing (i.e. non-decreasing) function and define a loss function  $L : \Theta \times \Theta \rightarrow [0, \infty)$  by

$$L(\theta_1, \theta_2) := g(d(\theta_1, \theta_2)),$$

for  $\theta_1, \theta_2 \in \Theta$ . Writing  $\widehat{\Theta}$  for the set of all measurable functions from  $\mathcal{X}$  to  $\Theta$ , for  $\delta \in (0, 1]$ ,  $\theta \in \Theta$ ,  $P_\theta \in \mathcal{P}_\theta$  and  $\widehat{\theta} \in \widehat{\Theta}$ , we then define the  $(1 - \delta)$ th quantile of  $L(\widehat{\theta}, \theta)$  under  $P_\theta$  as

$$\text{Quantile}(1 - \delta; P_\theta, L(\widehat{\theta}, \theta)) := \inf \left\{ r \in [0, \infty) : P_\theta \{L(\widehat{\theta}, \theta) \leq r\} \geq 1 - \delta \right\}.$$

In our more general setting, we have the following definition of minimax quantiles.

**Definition 1** (Minimax quantile). *For  $\delta \in (0, 1]$ , we define the minimax  $(1 - \delta)$ th quantile of the loss  $L$  for estimating  $\theta$  over  $\mathcal{P}_\Theta$  as*

$$\mathcal{M}(\delta) \equiv \mathcal{M}(\delta, \mathcal{P}_\Theta, L) := \inf_{\widehat{\theta}_\delta \in \widehat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} \text{Quantile}(1 - \delta; P_\theta, L(\widehat{\theta}_\delta, \theta)).$$

A high-probability upper bound for an estimator implies an upper bound on the minimax quantile. That is, if there exists  $\widehat{\theta}_\delta \in \widehat{\Theta}$  such that with  $P_\theta$ -probability at least  $1 - \delta$ , we have  $L(\widehat{\theta}_\delta, \theta) \leq \text{UB}(\delta)$  for all  $\theta \in \Theta$  and  $P_\theta \in \mathcal{P}_\theta$ , then  $\mathcal{M}(\delta) \leq \text{UB}(\delta)$ . The main goal of this section is to provide general lower bounds on minimax quantiles. The following basic result shows that a lower bound on the minimax quantile  $\mathcal{M}(\delta)$  implies a lower bound on the minimax risk.

**Proposition 2.** *For  $\delta \in (0, 1]$ , we have*

$$\inf_{\widehat{\theta} \in \widehat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} \mathbb{E}_{P_\theta} L(\widehat{\theta}, \theta) \geq \delta \mathcal{M}(\delta).$$

---

<sup>†</sup> $(\Theta, d)$  is a pseudo-metric space if  $d : \Theta \times \Theta \rightarrow [0, \infty)$  satisfies for all  $\theta_1, \theta_2, \theta_3 \in \Theta$  that  $d(\theta_1, \theta_1) = 0$ ,  $d(\theta_1, \theta_2) = d(\theta_2, \theta_1)$  and  $d(\theta_1, \theta_2) \leq d(\theta_1, \theta_3) + d(\theta_2, \theta_3)$ ; in contrast to a metric space, we may have  $d(\theta_1, \theta_2) = 0$  with  $\theta_1 \neq \theta_2$ .

Key to our development is the notion of a lower minimax quantile, which turns out to be more amenable to direct analysis.

**Definition 3** (Lower minimax quantile). *For  $\delta \in (0, 1]$ , we define the lower minimax  $(1-\delta)$ th quantile of the loss  $L$  for estimating  $\theta$  over  $\mathcal{P}_\Theta$  as*

$$\mathcal{M}_-(\delta) \equiv \mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) := \inf \left\{ r \in [0, \infty) : \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{ L(\hat{\theta}, \theta) > r \} \leq \delta \right\}.$$

The following theorem relates minimax quantiles to their lower analogues. In particular, it reveals that the lower minimax quantile is not only a minorant of the corresponding minimax quantile, but also that the two quantities coincide almost everywhere.

**Theorem 4.** *For all  $\delta \in (0, 1]$  and  $\xi \in (0, \delta)$ , we have*

$$\mathcal{M}_-(\delta) \leq \mathcal{M}(\delta) \leq \mathcal{M}_-(\delta - \xi). \quad (1)$$

Moreover,  $\mathcal{M}(\delta) = \mathcal{M}_-(\delta)$  at all but countably many points  $\delta \in (0, 1]$ .

The convenience of the lower bound in Theorem 4 arises from the fact that if

$$\inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{ L(\hat{\theta}, \theta) > r \} > \delta,$$

then  $\mathcal{M}(\delta) \geq \mathcal{M}_-(\delta) \geq r$ .

With these notions in place, we now establish high-probability versions of standard information-theoretic tools, beginning with a high-probability variant of Le Cam’s two-point lemma.

**Lemma 5** (High-probability Le Cam’s two-point lemma). *Let  $\delta \in (0, 1/2)$ , and suppose that  $\theta_1, \theta_2 \in \Theta$ ,  $P_1 \in \mathcal{P}_{\theta_1}$  and  $P_2 \in \mathcal{P}_{\theta_2}$  satisfy  $\text{TV}(P_1, P_2) < 1 - 2\delta$ . Then, writing  $\eta := d(\theta_1, \theta_2)/2$ , we have  $\mathcal{M}_-(\delta) \geq g(\eta)$ .*

In our applications in Section 3, we will typically apply this lemma to two product distributions for which the total variation distance may be difficult to compute. The following lemma, which is a straightforward consequence of the Bretagnolle–Huber inequality, provides a more user-friendly version of the above lemma, stated in terms of the Kullback–Leibler divergence.

**Corollary 6.** *Let  $\delta \in (0, 1/2)$ , and suppose that  $\theta_1, \theta_2 \in \Theta$ ,  $P_1 \in \mathcal{P}_{\theta_1}$  and  $P_2 \in \mathcal{P}_{\theta_2}$  satisfy  $\text{KL}(P_1, P_2) < \log\left(\frac{1}{4\delta(1-\delta)}\right)$ . Then, writing  $\eta := d(\theta_1, \theta_2)/2$ , we have  $\mathcal{M}_-(\delta) \geq g(\eta)$ .*

*Proof.* By the Bretagnolle–Huber inequality (Bretagnolle and Huber, 1979),

$$\text{TV}(P_1, P_2) \leq \sqrt{1 - \exp\{-\text{KL}(P_1, P_2)\}} < 1 - 2\delta.$$

The result then follows by Lemma 5. □

As we will see in Section 3, in several examples, the minimax quantile decomposes into a sum of a term reflecting the dependence on the quantile level  $\delta$ , and another capturing the minimax risk (that is independent of  $\delta$ ). Lemma 5 and Corollary 6 are effective for controlling the first of these terms. On the other hand, our next two results are well-suited for the minimax risk term, with the first providing lower bounds on the minimax quantile via a specialisation of Fano’s lemma.

**Lemma 7.** Let  $M \geq 2$ ,  $\theta_1, \dots, \theta_M \in \Theta$ , and  $P_j \in \mathcal{P}_{\theta_j}$  for  $j \in [M]$ . Suppose that there exists  $\epsilon \in (0, 1]$  such that

$$\frac{M^{-1} \inf_{Q \in \mathcal{Q}(\mathcal{X})} \sum_{j=1}^M \text{KL}(P_j, Q) + \log(2 - M^{-1})}{\log M} \leq 1 - \epsilon.$$

Then, writing  $\eta := \min_{1 \leq j < k \leq M} d(\theta_j, \theta_k)/2$ , we have  $\mathcal{M}_-(\delta) \geq g(\eta)$  for all  $\delta \in (0, \epsilon)$ .

The second of our results tailored for the minimax risk term obtains lower bounds on the lower minimax quantile  $\mathcal{M}_-$  from local minimax risk lower bounds, thereby facilitating the use of other tools such as Assouad's lemma. During the preparation of this manuscript, the authors became aware of the recent result of [Chhor, Klopp and Tsybakov \(2024, Lemma 6\)](#), which provides such a reduction for a bounded parameter space. The removal of this assumption turns out to be important, for instance, in the problem of multivariate mean estimation ([Proposition 10](#)).

**Theorem 8.** Let  $\Theta_0 \subseteq \Theta$  be non-empty and suppose that  $D := \sup_{\theta_1, \theta_2 \in \Theta_0} d(\theta_1, \theta_2)$  satisfies  $g(D) > 0$ . Let  $\Delta \in [0, \infty)$  be such that

$$\inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} \mathbb{E}_{P_{\theta_0}} L(\hat{\theta}, \theta_0) \geq \Delta.$$

Then for every  $\epsilon > 0$  and  $\delta \in (0, \frac{\Delta - g(\epsilon D)}{g((1+\epsilon)D)})$ , we have  $\mathcal{M}_-(\delta) \geq g(\epsilon D)$ .

In order to apply this theorem, we note that standard minimax lower bound techniques proceed by constructing a carefully-chosen finite subset of the parameter space, and hence automatically yield local minimax risk lower bounds; see, e.g., [Propositions 10 and 16](#). The upper bound on  $\delta$  in [Theorem 8](#) may be small in some cases; [Proposition 19](#) provides a way to relax this to any universal constant less than  $1/2$ . Recall that a *Polish space* is a separable topological space that may be equipped with a metric that generates the topology and under which the space is complete; for example, any open or closed subset of  $\mathbb{R}^d$  is a Polish space.

**Proposition 9.** For  $m \in \mathbb{N}$ , let  $\mathcal{P}_{\theta}^{\otimes m} := \{P_{\theta}^{\otimes m} \in \mathcal{Q}(\mathcal{X}^m) : P_{\theta} \in \mathcal{P}_{\theta}\}$  and  $\mathcal{P}_{\Theta}^{\otimes m} := \{P_{\theta}^{\otimes m} : \theta \in \Theta\}$ . Suppose that  $\Theta$  is a non-empty Polish space, that  $g$  is continuous and that there exists  $A > 0$  such that  $L(\theta_1, \theta_2) \leq A\{L(\theta_1, \theta_3) + L(\theta_2, \theta_3)\}$  for all  $\theta_1, \theta_2, \theta_3 \in \Theta$ . Define  $h : [0, 1] \rightarrow [0, 1]$  by  $h(x) := x^3 + 3x^2(1 - x)$  and for  $m \in \mathbb{N}$ , let  $h^{\circ m}$  denote the function obtained by composing  $h$  with itself  $m$  times. Let  $\delta_- \in (0, 1/2)$  and  $\delta_+ \in (\delta_-, 1/2)$ . Then there exists  $k \equiv k(\delta_-, \delta_+) \in \mathbb{N}$  such that  $h^{\circ k}(\delta_+) \leq \delta_-$ . Moreover, for any such  $k$ , we have for  $\delta \in (0, \delta_+]$  that

$$\mathcal{M}_-(\delta, \mathcal{P}_{\Theta}, L) \geq \frac{1}{(2A)^k} \cdot \mathcal{M}_-(\delta_-, \mathcal{P}_{\Theta}^{\otimes 3k}, L).$$

### 3 Examples

With the general techniques of [Section 2](#) now in place, the aim of this section is to determine expressions for the minimax quantiles of loss functions in several examples. In each of the statements to follow, we will consider quantile levels  $\delta \leq 1/4$ ; this value is chosen simply for convenience and could be relaxed to any constant less than  $1/2$  by, e.g., a further application of [Proposition 9](#). Throughout this section, we will slightly abuse notation when the loss is of the form  $L(\theta, \theta') = g(\|\theta - \theta'\|)$  for some norm  $\|\cdot\|$ , by writing  $\mathcal{M}(\delta, \mathcal{P}_{\Theta}, g(\|\cdot\|))$  in place of  $\mathcal{M}(\delta, \mathcal{P}_{\Theta}, L)$ .

### 3.1 Multivariate mean estimation

For this basic problem, we determine the minimax quantile in three settings, one for each of the next subsections.

#### 3.1.1 Gaussian mean estimation in squared Euclidean norm

Proposition 10 below recovers the minimax  $(1 - \delta)$ th quantile in multivariate Gaussian mean estimation, up to multiplicative universal constants. The matching upper bound is attained by the sample mean, which is a  $\delta$ -independent estimator. This result therefore demonstrates the optimality of the sample mean in Gaussian mean estimation in a much stronger manner than usual minimax risk lower bounds.

**Proposition 10.** *Let  $n, d \in \mathbb{N}$ ,  $\delta \in (0, 1/4]$ ,  $\Sigma \in \mathcal{S}_{++}^{d \times d}$ ,  $\Theta := \mathbb{R}^d$  and  $\mathcal{P}_\theta := \{\mathbf{N}_d(\theta, \Sigma)^{\otimes n}\}$  for  $\theta \in \Theta$ . Then*

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_2^2) \asymp \frac{\text{tr}(\Sigma)}{n} + \frac{\|\Sigma\|_{\text{op}} \log(1/\delta)}{n}.$$

*Proof.* Let  $v_1, \dots, v_d \in \mathbb{R}^d$  denote orthonormal eigenvectors of  $\Sigma$  with corresponding eigenvalues  $\lambda_1 \geq \dots \geq \lambda_d$ . We capture each of the two additive terms separately, starting with the  $\delta$ -dependent term. To this end, take  $\theta_1 := 0$ ,  $\theta_2 := \sqrt{\frac{\lambda_1}{n} \log\left(\frac{1}{4\delta(1-\delta)}\right)} v_1$  and set  $P_1 = \mathbf{N}_d(\theta_1, \Sigma)^{\otimes n}$  and  $P_2 = \mathbf{N}_d(\theta_2, \Sigma)^{\otimes n}$ . Then

$$\text{KL}(P_1, P_2) = n \text{KL}(\mathbf{N}_d(\theta_1, \Sigma), \mathbf{N}_d(\theta_2, \Sigma)) = \frac{1}{2} \log\left(\frac{1}{4\delta(1-\delta)}\right) < \log\left(\frac{1}{4\delta(1-\delta)}\right).$$

Thus, applying Theorem 4 in conjunction with Corollary 6, we deduce that for  $\delta \in (0, 1/4]$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_2^2) \geq \mathcal{M}_-(\delta, \mathcal{P}_\Theta, \|\cdot\|_2^2) \geq \frac{\|\Sigma\|_{\text{op}} \log\left(\frac{1}{4\delta(1-\delta)}\right)}{4n} \geq \frac{\|\Sigma\|_{\text{op}} \log(1/\delta)}{20n}. \quad (2)$$

We turn now to the other term in the lower bound. For  $\theta, \theta' \in \mathbb{R}^d$ , we have

$$\|\theta - \theta'\|_2^2 = \left\| \sum_{j=1}^d \{(v_j^\top \theta) v_j - (v_j^\top \theta') v_j\} \right\|_2^2 = \sum_{j=1}^d \{v_j^\top (\theta - \theta')\}^2 = \sum_{j=1}^d g(\rho_j(\theta, \theta')),$$

where we defined the pseudo-metrics  $\rho_j : \Theta \times \Theta \rightarrow [0, \infty)$  by  $\rho_j(\theta, \theta') := |v_j^\top (\theta - \theta')|$  for  $j \in [d]$  and the increasing function  $g : [0, \infty) \rightarrow [0, \infty)$  by  $g(x) := x^2$ . Now let  $\Phi := \{0, 1\}^d$  and for  $j \in [d]$  set  $a_j := \frac{4}{3} \sqrt{\frac{\lambda_j}{n}}$ . For  $\phi = (\phi_1, \dots, \phi_d)^\top \in \Phi$ , take  $\theta_\phi := \sum_{j=1}^d a_j \phi_j v_j$  and let  $P_\phi := \mathbf{N}_d(\theta_\phi, \Sigma)^{\otimes n}$ . Define  $\Theta_0 := \{\theta_\phi : \phi \in \Phi\} \subseteq \Theta$  with diameter  $D := \frac{4}{3} \sqrt{\frac{\text{tr}(\Sigma)}{n}}$ . For  $\phi, \phi' \in \Phi$  that differ only in the  $j$ th coordinate, we have  $\rho_j(\theta_\phi, \theta_{\phi'}) = |a_j|$  and, by Pinsker's inequality (Wainwright, 2019, Lemma 15.2),  $\text{TV}(P_\phi, P_{\phi'}) \leq \left\{ \frac{n}{2} \text{KL}(\mathbf{N}_d(\theta_\phi, \Sigma), \mathbf{N}_d(\theta_{\phi'}, \Sigma)) \right\}^{1/2} = 2/3$ . Thus, by Assouad's lemma (Lemma 23),

$$\inf_{\hat{\theta} \in \Theta} \max_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} \mathbb{E}_{P_{\theta_0}} (\|\hat{\theta} - \theta_0\|_2^2) \geq \frac{4 \text{tr}(\Sigma)}{27n}.$$



Taking  $\epsilon = 3/40$  in Theorem 8, we deduce for  $\delta \in (0, 1/15]$  that

$$\mathcal{M}_-(\delta, \mathcal{P}_\Theta, \|\cdot\|_2^2) \geq g(\epsilon D) = \frac{\text{tr}(\Sigma)}{100n}. \quad (3)$$

Now, recalling the function  $h$  from Proposition 9, since  $h(h(1/4)) < 1/15$ , we may apply Proposition 9 with  $k = 2$  and (3) to deduce that for any  $\delta \in (0, 1/4]$ ,

$$\mathcal{M}_-(\delta, \mathcal{P}_\Theta, \|\cdot\|_2^2) \geq \frac{\text{tr}(\Sigma)}{2^6 \cdot 3^2 \cdot 5^2 \cdot n}. \quad (4)$$

Combining Theorem 4 with (2) and (4) therefore yields that for  $\delta \in (0, 1/4]$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_2^2) \geq \mathcal{M}_-(\delta, \mathcal{P}_\Theta, \|\cdot\|_2^2) \geq \frac{\text{tr}(\Sigma)}{2^7 \cdot 3^2 \cdot 5^2 \cdot n} + \frac{\|\Sigma\|_{\text{op}} \log(1/\delta)}{40n}.$$

For the upper bound, consider the sample mean as an estimator. Set  $\bar{X} := n^{-1} \sum_{i=1}^n X_i$ , where  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \mathbf{N}(\theta, \Sigma)$ , and note that by Jensen's inequality,  $\mathbb{E}(\|\bar{X} - \theta\|_2) \leq \sqrt{\text{tr}(\Sigma)/n}$ . Moreover,  $\|\bar{X} - \theta\|_2 = \sup_{u \in \mathbb{S}^{d-1}} u^\top (\bar{X} - \theta)$ , so using the fact that  $2(a + b) \geq (a^{1/2} + b^{1/2})^2$  for  $a, b \geq 0$  by another application of Jensen's inequality, as well as the Borell-TIS inequality (see, e.g., Adler and Taylor, 2009, Theorem 2.1.1), we have for any  $\delta \in (0, 1]$  that

$$\begin{aligned} \mathbb{P}\left(\|\bar{X} - \theta\|_2^2 > \frac{2 \text{tr}(\Sigma)}{n} + \frac{4\|\Sigma\|_{\text{op}} \log(1/\delta)}{n}\right) \\ \leq \mathbb{P}\left(\|\bar{X} - \theta\|_2 > \mathbb{E}(\|\bar{X} - \theta\|_2) + \sqrt{\frac{2\|\Sigma\|_{\text{op}} \log(1/\delta)}{n}}\right) \leq \delta, \end{aligned}$$

as required.  $\square$

### 3.1.2 Robust mean estimation in squared Euclidean norm

Recent literature has studied high-probability upper bounds for mean estimation with heavy-tailed data (see, e.g., the review article of Lugosi and Mendelson (2019a) and the references therein), or more generally, data from the Huber  $\epsilon$ -contamination model (see, e.g., Lugosi and Mendelson, 2021; Depersin and Lecu e, 2022b). These upper bound are attained using  $\delta$ -dependent estimators. As a corollary of Proposition 10, we provide a matching lower bound on the minimax  $(1 - \delta)$ th quantile up to universal constants in these settings.

**Corollary 11.** *Let  $n, d \in \mathbb{N}$ ,  $\epsilon \in [0, 1)$ ,  $\Sigma \in \mathcal{S}_{++}^{d \times d}$ ,  $\Theta = \mathbb{R}^d$  and*

$$\mathcal{P}_\theta := \left\{ R^{\otimes n} : R \in \mathcal{Q}(\mathbb{R}^d), R = (1 - \epsilon)P + \epsilon Q, \mathbb{E}_P(X) = \theta, \text{Cov}_P(X) = \Sigma, Q \in \mathcal{Q}(\mathbb{R}^d) \right\}.$$

*There exists a universal constant  $c > 0$  such that if  $e^{-cn(1-300\epsilon)} \leq \delta \leq 1/4$ , then*

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_2^2) \asymp \frac{\text{tr}(\Sigma)}{n} + \|\Sigma\|_{\text{op}} \epsilon + \frac{\|\Sigma\|_{\text{op}} \log(1/\delta)}{n}.$$

*Proof.* We first consider the lower bound. When  $\varepsilon = 0$ , the claimed lower bound follows from Proposition 10, so without loss of generality we assume that  $\varepsilon > 0$ . First, since  $\mathbb{N}_d(\theta, \Sigma)^{\otimes n} \in \mathcal{P}_\theta$ , we have by Proposition 10 that for  $\delta \in (0, 1/4]$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\theta, \|\cdot\|_2^2) \gtrsim \frac{\text{tr}(\Sigma)}{n} + \frac{\|\Sigma\|_{\text{op}} \log(1/\delta)}{n}. \quad (5)$$

By the spectral theorem,  $\Sigma = V\Lambda V^\top \in \mathbb{R}^{d \times d}$  where  $\Lambda := \text{diag}(\lambda_1, \dots, \lambda_d) \in \mathbb{R}^{d \times d}$ ,  $V \in \mathbb{R}^{d \times d}$  is orthogonal and  $\lambda_1 \geq \dots \geq \lambda_d$ . Let  $a := (\varepsilon + \varepsilon^2)/2$  and  $b := (3\varepsilon + \varepsilon^2)/2$ . Define random vectors  $X = (X_1, \dots, X_d)^\top \sim P_X$  and  $Y = (Y_1, \dots, Y_d)^\top \sim P_Y$  with independent components satisfying

$$X_1 := \begin{cases} -\sqrt{\frac{\lambda_1}{2\varepsilon}} & \text{with probability } \varepsilon \\ 0 & \text{with probability } 1 - 2\varepsilon, \\ \sqrt{\frac{\lambda_1}{2\varepsilon}} & \text{with probability } \varepsilon \end{cases}, \quad Y_1 := \begin{cases} -\sqrt{\frac{\lambda_1}{2\varepsilon}} & \text{with probability } a \\ 0 & \text{with probability } 1 - a - b \\ \sqrt{\frac{\lambda_1}{2\varepsilon}} & \text{with probability } b, \end{cases}$$

and  $X_j \stackrel{d}{=} Y_j \sim \mathbb{N}(0, \lambda_j)$  for all  $j \in \{2, \dots, d\}$ . Then

$$\text{Var}(X_1) = \lambda_1, \quad \text{Var}(Y_1) = \frac{\lambda_1}{2\varepsilon} \{(a+b) - (b-a)^2\} = \lambda_1.$$

Thus  $\text{Cov}(X) = \text{Cov}(Y) = \Lambda$ , and

$$\text{TV}(P_X, P_Y) = \frac{1}{2} \left( \frac{\varepsilon - \varepsilon^2}{2} + \varepsilon^2 + \frac{\varepsilon + \varepsilon^2}{2} \right) = \frac{\varepsilon + \varepsilon^2}{2} \leq \varepsilon.$$

Define  $X' := VX$ ,  $Y' := VY$  and let  $P_{X'}$ ,  $P_{Y'}$  denote the distributions of  $X'$ ,  $Y'$  respectively. Then  $\text{Cov}(X') = \text{Cov}(Y') = \Sigma$ , and by the data processing inequality (e.g., Gerchinovitz, Ménard and Stoltz, 2020, Lemma 2.1),

$$\text{TV}(P_{X'}, P_{Y'}) \leq \text{TV}(P_X, P_Y) \leq \varepsilon \leq \frac{\varepsilon}{1 - \varepsilon}.$$

Then, let  $\mathcal{R}_\theta := \{R \in \mathcal{Q}(\mathbb{R}^d) : \mathbb{E}_R(X) = \theta, \text{Cov}_R(X) = \Sigma\}$  and  $\mathcal{R}_\Theta := \{\mathcal{R}_\theta : \theta \in \Theta\}$ . Then, writing  $\theta_1 := \mathbb{E}(X')$  and  $\theta_2 := \mathbb{E}(Y')$ , we have by Theorem 4 and Lemma 25 that

$$\mathcal{M}(\delta, \mathcal{P}_\theta, \|\cdot\|_2^2) \geq \mathcal{M}_-(\delta, \mathcal{P}_\theta, \|\cdot\|_2^2) \geq \frac{\|\theta_1 - \theta_2\|_2^2}{4} = \frac{(b-a)^2 \cdot \|\Sigma\|_{\text{op}}}{8\varepsilon} = \frac{\|\Sigma\|_{\text{op}} \varepsilon}{8}. \quad (6)$$

Combining (5) and (6) proves that the stated lower bound holds for all  $\delta \in (0, 1/4]$ .

The matching upper bound follows from Depersin and Lecué (2022b, Theorem 2.1), by taking  $u = \infty$  in their notation (note that the lower bound on  $\delta$ —which in turn implies an upper bound on  $\varepsilon$ —is implicit from their theorem).  $\square$

### 3.1.3 Gaussian mean estimation in $\ell_\infty$ -norm

In our next result, we switch to  $\ell_\infty$ -loss, and invoke Fano's lemma (Lemma 7) for the lower bound. As in Proposition 10, the matching upper bound is achieved by the sample mean.

**Proposition 12.** Let  $n, d \in \mathbb{N}$ ,  $\sigma > 0$ ,  $\delta \in (0, 1/4]$ ,  $\Theta := \mathbb{R}^d$  and  $\mathcal{P}_\theta := \{\mathbf{N}_d(\theta, \Sigma)^{\otimes n} : \Sigma \in \mathcal{S}_{++}^{d \times d}, \|\Sigma\|_\infty \leq \sigma^2\}$ . Then

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_\infty) \asymp \sigma \sqrt{\frac{\log(d/\delta)}{n}}.$$

**Remark.** In our lower bound construction below, we only use distributions whose covariance matrices are scalar multiples of the identity matrix. In that setting, we see from Proposition 12 that the term involving the quantile level  $\delta$  dominates when  $\delta \ll 1/d$ , whereas with squared Euclidean error, Proposition 10 reveals that this happens when  $\delta \ll e^{-d}$ .

*Proof.* Define  $\theta_1 := \sigma \sqrt{\frac{1}{n} \log\left(\frac{1}{4\delta(1-\delta)}\right)} e_1$ ,  $\theta_2 := 0$ , and  $P_\ell := \mathbf{N}_d(\theta_\ell, \sigma^2 I_d)^{\otimes n} \in \mathcal{P}_{\theta_\ell}$  for  $\ell \in \{1, 2\}$ . Then  $\text{KL}(P_1, P_2) = \frac{1}{2} \log\left(\frac{1}{4\delta(1-\delta)}\right) < \log\left(\frac{1}{4\delta(1-\delta)}\right)$ . Hence, by Theorem 4 and Corollary 6, we have for  $\delta \in (0, 1/4]$  that

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_\infty) \geq \mathcal{M}_-(\delta, \mathcal{P}_\Theta, \|\cdot\|_\infty) \geq \sigma \sqrt{\frac{1}{4n} \log\left(\frac{1}{4\delta(1-\delta)}\right)} \geq \sigma \sqrt{\frac{\log(1/\delta)}{20n}}. \quad (7)$$

Now assume for the time being that  $d \geq 4$ . For  $j \in [d]$ , define  $\theta^{(j)} := \sigma \sqrt{\frac{\log d}{2n}} e_j$ ,  $P^{(j)} := \mathbf{N}_d(\theta^{(j)}, \sigma^2 I_d)^{\otimes n} \in \mathcal{P}_{\theta^{(j)}}$ , and define  $Q := \mathbf{N}_d(0, \sigma^2 I)^{\otimes n} \in \mathcal{Q}((\mathbb{R}^d)^n)$ . Then  $\text{KL}(P^{(j)}, Q) = n \text{KL}(\mathbf{N}_d(\theta^{(j)}, \sigma^2 I_d), \mathbf{N}_d(0, \sigma^2 I_d)) = \frac{1}{4} \log d$  and thus

$$\frac{d^{-1} \sum_{j=1}^d \text{KL}(P^{(j)}, Q) + \log(2 - d^{-1})}{\log d} \leq \frac{1}{4} + \frac{\log(2 - d^{-1})}{\log d} \leq \frac{7}{10}.$$

Therefore, by applying Theorem 4, followed by Lemma 7 with  $\epsilon = 3/10$ , we deduce that for  $\delta \in (0, 1/4]$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_\infty) \geq \mathcal{M}_-(\delta, \mathcal{P}_\Theta, \|\cdot\|_\infty) \geq \sigma \sqrt{\frac{\log d}{8n}}. \quad (8)$$

Combining (7) and (8) yields that for  $\delta \in (0, 1/4]$  and  $d \geq 4$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_\infty) \geq \frac{\sigma}{2} \left( \sqrt{\frac{\log d}{8n}} + \sqrt{\frac{\log(1/\delta)}{20n}} \right) \geq \sigma \sqrt{\frac{\log(d/\delta)}{80n}}. \quad (9)$$

Now consider the case  $d \leq 3$ . By (7), we have for  $\delta \in (0, 1/4]$  that

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_\infty) \geq \sigma \sqrt{\frac{\log(1/\delta)}{20n}} \geq \sigma \sqrt{\frac{\log(d/\delta)}{40n}}. \quad (10)$$

Combining (9) and (10) proves the lower bound.

For the upper bound, let  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} P_\theta \in \mathcal{P}_\theta$  and  $\bar{X} := n^{-1} \sum_{i=1}^n X_i$ . Then for  $j \in [d]$ ,

$$\mathbb{P}\left(|\bar{X}_j - \theta_j| > \sigma \sqrt{\frac{2 \log(d/\delta)}{n}}\right) = 2 \int_{\sqrt{2 \log(d/\delta)}}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx \leq \frac{\delta}{d}.$$

Applying a union bound, we deduce that

$$\mathbb{P}\left(\|\bar{X} - \theta\|_\infty > \sigma \sqrt{\frac{2 \log(d/\delta)}{n}}\right) \leq \delta,$$

so  $\mathcal{M}_-(\delta, \mathcal{P}_\Theta, \|\cdot\|_\infty) \leq \mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_\infty) \leq \sigma \sqrt{\frac{2 \log(d/\delta)}{n}}$  for  $\delta \in (0, 1]$ .  $\square$

### 3.2 Covariance matrix estimation

Here we consider the estimation of a covariance matrix in operator norm based on a sample of size  $n$ , and determine the minimax quantile up to universal constants. As we show in the proof of Proposition 13 below, the upper bound is attained by the sample covariance matrix, applying the careful analysis of Koltchinskii and Lounici (2017) (see also Zhivotovskiy (2024)) when the quantile level  $\delta$  satisfies  $\delta \geq e^{-n}$ , and the trivial zero estimator when  $\delta < e^{-n}$ . By contrast, the matrix Bernstein inequality yields sub-optimal concentration as compared to the minimax quantile in Proposition 13; see Lemma 26 for details. Recall that the *effective rank* of  $\Sigma \in \mathcal{S}_+^{d \times d}$  is defined as  $\mathbf{r}(\Sigma) := \text{tr}(\Sigma)/\|\Sigma\|_{\text{op}}$ , with the convention that  $0/0 := 0$ .

**Proposition 13.** *Let  $n, d \in \mathbb{N}$ ,  $\delta \in (0, 1/4]$ ,  $\sigma > 0$ ,  $r \in [1, n \wedge d]$ ,  $\Theta := \{\Sigma \in \mathcal{S}_+^{d \times d} : \|\Sigma\|_{\text{op}} \leq \sigma^2, \mathbf{r}(\Sigma) \leq r\}$ , and for  $\Sigma \in \Theta$ , define  $\mathcal{P}_\Sigma := \{\mathbf{N}_d(0, \Sigma)^{\otimes n}\}$ . Then*

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_{\text{op}}) \asymp \sigma^2 \sqrt{\frac{r + \log(1/\delta)}{n}} \wedge \sigma^2.$$

*Proof.* Let  $r_0 := \lfloor r \rfloor$ , so that  $r_0 \geq (r/2) \vee 1$ . First consider the case  $\delta \in [e^{-n}/3, 1/4]$ , and set  $a := 1 - \sqrt{\frac{1}{4n} \log\left(\frac{1}{4\delta(1-\delta)}\right)} \in [1/2, 1]$ . Define  $\Sigma^{(1)}, \Sigma^{(2)} \in \mathcal{S}_+^{d \times d}$  by

$$\Sigma^{(1)} := \begin{pmatrix} a^2 \sigma^2 & 0 & 0 \\ 0 & \sigma^2 I_{r_0-1} & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad \Sigma^{(2)} := \begin{pmatrix} \sigma^2 I_{r_0} & 0 \\ 0 & 0 \end{pmatrix}.$$

Then, for  $\ell \in \{1, 2\}$ , we have  $\|\Sigma^{(\ell)}\|_{\text{op}} \leq \sigma^2$  and  $\mathbf{r}(\Sigma^{(\ell)}) \leq r_0 \leq r$ , so that  $\Sigma^{(\ell)} \in \Theta$ . Letting  $P_\ell := \mathbf{N}_d(0, \Sigma^{(\ell)})^{\otimes n}$  for  $\ell \in \{1, 2\}$ , we find by Taylor's theorem with the integral form of the remainder that

$$\begin{aligned} \text{KL}(P_1, P_2) &= n \cdot \frac{\log(1/a^2) - (1-a^2)}{2} = \frac{n}{2} \cdot \left\{ 2(1-a)^2 + 2 \int_a^1 \frac{1}{t} \left(1 - \frac{a}{t}\right)^2 dt \right\} \\ &\leq 3n(1-a)^2 = \frac{3}{4} \log\left(\frac{1}{4\delta(1-\delta)}\right) < \log\left(\frac{1}{4\delta(1-\delta)}\right). \end{aligned}$$

Therefore, combining Theorem 4 and Corollary 6, we find that

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_{\text{op}}) \geq \frac{\|\Sigma^{(1)} - \Sigma^{(2)}\|_{\text{op}}}{2} = \frac{(1-a^2)\sigma^2}{2} \geq \frac{\sigma^2}{4\sqrt{n}} \sqrt{\log\left(\frac{1}{4\delta(1-\delta)}\right)} \geq \frac{\sigma^2}{9} \sqrt{\frac{\log(1/\delta)}{n}}.$$

On the other hand, when  $\delta \in (0, e^{-n}/3)$ , since  $\mathcal{M}(\delta)$  is decreasing in  $\delta$ , we have

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_{\text{op}}) \geq \mathcal{M}(e^{-n}, \mathcal{P}_\Theta, \|\cdot\|_{\text{op}}) \geq \frac{\sigma^2}{9}.$$

Hence, for  $\delta \in (0, 1/4]$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_{\text{op}}) \gtrsim \sigma^2 \sqrt{\frac{\log(1/\delta)}{n}} \wedge \sigma^2. \tag{11}$$

Now assume that  $r_0 \geq 20$  and define  $\Phi := \{0, 1\}^{r_0}$ . By the Gilbert–Varshamov lemma (e.g., [Massart, 2007](#), Lemma 4.7), there exists  $\Phi_0 \subseteq \Phi$  with  $|\Phi_0| \geq e^{r_0/8}$  and  $d_{\text{H}}(\phi, \phi') > r_0/4$  for all distinct  $\phi, \phi' \in \Phi_0$ , where  $d_{\text{H}}$  denotes the Hamming distance. Let  $b := \frac{\sigma^2}{6\sqrt{nr_0}}$ , and for  $\phi \in \Phi_0$ , define  $\Sigma^{(\phi)} \in \mathcal{S}_+^{d \times d}$  as well as  $\Sigma^{(0)} \in \mathcal{S}_+^{d \times d}$  by

$$\Sigma^{(\phi)} := \begin{pmatrix} \frac{\sigma^2}{2}I_{r_0} + b\phi\phi^\top & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad \Sigma^{(0)} := \begin{pmatrix} \frac{\sigma^2}{2}I_{r_0} & 0 \\ 0 & 0 \end{pmatrix}.$$

Then  $\|\Sigma^{(\phi)}\|_{\text{op}} = \frac{\sigma^2}{2} + b\|\phi\|_2^2 \leq \frac{\sigma^2}{2} + br_0 \leq \sigma^2$  and  $\mathbf{r}(\Sigma^{(\phi)}) \leq r_0 \leq r$  for all  $\phi \in \Phi_0$ , so  $\Sigma^{(\phi)} \in \Theta$ . For  $\phi \in \Phi_0$ , define  $P^{(\phi)} := \mathbf{N}_d(0, \Sigma^{(\phi)})^{\otimes n}$  so that  $P^{(\phi)} \in \mathcal{P}_{\Sigma^{(\phi)}}$ . Then, taking  $Q := \mathbf{N}_d(0, \Sigma^{(0)})^{\otimes n}$ , we have for  $\phi \in \Phi_0$  that

$$\begin{aligned} \text{KL}(P^{(\phi)}, Q) &= \frac{n}{2} \left\{ \log \left( \frac{(\sigma^2/2)^{r_0}}{(b\|\phi\|_2^2 + \sigma^2/2)(\sigma^2/2)^{r_0-1}} \right) + \frac{2b\|\phi\|_2^2}{\sigma^2} \right\} \\ &= \frac{n}{2} \left\{ \log \left( \frac{1}{1 + 2b\|\phi\|_2^2/\sigma^2} \right) + \frac{2b\|\phi\|_2^2}{\sigma^2} \right\} \leq \frac{nb^2\|\phi\|_2^4}{\sigma^4} \leq \frac{r_0}{36}, \end{aligned}$$

where the penultimate inequality follows because  $\log(\frac{1}{1+x}) + x \leq x^2/2$  for  $x \geq 0$  and the final inequality follows since  $\|\phi\|_2^2 \leq r_0$ . Therefore,

$$\frac{|\Phi_0|^{-1} \sum_{\phi \in \Phi_0} \text{KL}(P^{(\phi)}, Q) + \log(2 - |\Phi_0|^{-1})}{\log |\Phi_0|} \leq \frac{r_0/36 + \log(2)}{r_0/8} \leq \frac{1}{2}, \quad (12)$$

where the final inequality holds under the assumption  $r_0 \geq 20$ . Now fix distinct  $\phi = (\phi_1, \dots, \phi_{r_0})^\top$ ,  $\phi' = (\phi'_1, \dots, \phi'_{r_0})^\top \in \Phi_0$ , and define  $S := \{j \in [r_0] : \phi_j = 1 \text{ and } \phi'_j = 0\}$ . By reversing the roles of  $\phi$  and  $\phi'$  if necessary, we may assume that  $|S| \geq r_0/8$  since  $d_{\text{H}}(\phi, \phi') > r_0/4$ . Define  $v = (v_1, \dots, v_{r_0})^\top \in \{0, 1\}^{r_0}$  by  $v_j := \mathbb{1}_{\{j \in S\}}$  for  $j \in [r_0]$ . Then

$$\|\Sigma^{(\phi)} - \Sigma^{(\phi')}\|_{\text{op}} \geq \frac{v^\top (\Sigma^{(\phi)} - \Sigma^{(\phi')}) v}{\|v\|_2^2} \geq \frac{b(r_0/8)^2}{r_0} \geq \frac{\sigma^2}{384} \sqrt{\frac{r_0}{n}}. \quad (13)$$

Combining (12) and (13) with Theorem 4 and Lemma 7, we deduce that for  $\delta \in (0, 1/2)$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_{\text{op}}) \geq \frac{\sigma^2}{768} \sqrt{\frac{r_0}{n}} \geq \frac{\sigma^2}{800} \sqrt{\frac{r}{n}}. \quad (14)$$

Hence, combining (11) and (14) yields that for  $r \geq 20$  and  $\delta \in (0, 1/4]$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_{\text{op}}) \gtrsim \sigma^2 \sqrt{\frac{r}{n}} + \left( \sigma^2 \sqrt{\frac{\log(1/\delta)}{n}} \wedge \sigma^2 \right) \gtrsim \sigma^2 \sqrt{\frac{r + \log(1/\delta)}{n}} \wedge \sigma^2.$$

On the other hand, for  $r < 20$  and  $\delta \in (0, 1/4]$ , we have by (11) that

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_{\text{op}}) \gtrsim \sigma^2 \sqrt{\frac{\log(1/\delta)}{n}} \wedge \sigma^2 \gtrsim \sigma^2 \sqrt{\frac{r + \log(1/\delta)}{n}} \wedge \sigma^2.$$

This proves the lower bound.

For the upper bound, let  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \mathbf{N}_d(0, \Sigma)$  where  $\Sigma \in \Theta$ . When  $\delta \in [e^{-n}, 1/4]$ , let  $\widehat{\Sigma} := n^{-1} \sum_{i=1}^n X_i X_i^\top$ . Then by [Koltchinskii and Lounici \(2017, Corollary 2\)](#), we have with probability at least  $1 - \delta$  that

$$\|\widehat{\Sigma} - \Sigma\|_{\text{op}} \lesssim \sigma^2 \sqrt{\frac{r + \log(1/\delta)}{n}}.$$

On the other hand, when  $\delta \in (0, e^{-n})$ , let  $\widehat{\Sigma} := 0$ , so that  $\|\widehat{\Sigma} - \Sigma\|_{\text{op}} = \|\Sigma\|_{\text{op}} \leq \sigma^2$ . Thus

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, \|\cdot\|_{\text{op}}) \lesssim \sigma^2 \sqrt{\frac{r + \log(1/\delta)}{n}} \wedge \sigma^2,$$

for  $\delta \in (0, 1/4]$ . □

### 3.3 Sparse linear regression

In this section, we consider sparse linear regression with fixed design  $X \in \mathbb{R}^{n \times d}$ . We will require the weighted restricted eigenvalue (WRE) condition of [Bellec, Lecué and Tsybakov \(2018, p. 3616\)](#). To recall this, for  $v = (v_1, \dots, v_d)^\top \in \mathbb{R}^d$ , let  $v^\# = (v_1^\#, \dots, v_d^\#)^\top \in [0, \infty)^d$  denote the vector obtained by ordering the absolute values of the components of  $v$  in decreasing order. For  $j \in [d]$  and some  $\sigma > 0$  (which will be taken to be the noise standard deviation), let  $\lambda_j := 6\sigma \sqrt{n^{-1} \log(2d/j)}$ . Now, for  $c_0 > 0$ ,  $s \in [d]$ , define the cone

$$\mathcal{C}_{\text{WRE}}(s, c_0) := \left\{ v = (v_1, \dots, v_d)^\top \in \mathbb{R}^d : \sum_{j=1}^d \lambda_j v_j^\# \leq (1 + c_0) \|v\|_2 \left( \sum_{j=1}^s \lambda_j^2 \right)^{1/2} \right\}$$

and let

$$\vartheta(s, c_0) := \min_{v \in \mathcal{C}_{\text{WRE}}(s, c_0) \setminus \{0\}} \frac{\|Xv\|_2}{n^{1/2} \|v\|_2}.$$

The following assumption imposes regularity conditions on our design matrix.

**Assumption 1.** *Each column of  $X \in \mathbb{R}^{n \times d}$  has Euclidean norm at most  $\sqrt{n}$ . Moreover, there exist  $c, C > 0$  such that when  $d = 1$ ,  $X$  satisfies  $\|X\|_2 \geq c\sqrt{n}$ ; when  $d \geq 2$ ,  $X$  satisfies  $\vartheta(s \vee 2, 3) \geq c$  and  $\max_{T \subseteq [d]: |T| \leq s \vee 2} \|X_T\|_{\text{op}} \leq C\sqrt{n}$ , where  $X_T \in \mathbb{R}^{n \times |T|}$  is the submatrix of  $X$  obtained by extracting the columns with indices in  $T$ .*

Existing results show that if  $X \in \mathbb{R}^{n \times d}$  has independent  $\mathbf{N}(0, 1/2)$  entries, then there exist universal constants  $c', C' > 0$  such that, when  $d \geq 2$ , Assumption 1 holds with  $c = 1/2$  and  $C = 2$  with probability at least  $1 - e^{-c'n}$  when  $n \geq C's \log(2ed/s)$ . Indeed, the verification of the condition  $\vartheta(s \vee 2, 3) \geq 1/2$  is given by [Bellec, Lecué and Tsybakov \(2018, Theorem 8.3\)](#), and verification of the condition  $\max_{T \subseteq [d]: |T| \leq s \vee 2} \|X_T\|_{\text{op}} / \sqrt{n} \leq 2$  can be obtained via concentration results for Gaussian ensembles (e.g., [Wainwright, 2019, Theorem 6.1](#)).

We note that the matching upper bound in Proposition 14 below is attained by the SLOPE estimator ([Bogdan et al., 2015; Bellec, Lecué and Tsybakov, 2018](#)), which is  $\delta$ -independent.

**Proposition 14.** Let  $n, d \in \mathbb{N}$ ,  $s \in [d]$ ,  $\delta \in (0, 1/4]$ ,  $\sigma > 0$  and  $\Theta := \{\theta \in \mathbb{R}^d : \|\theta\|_0 \leq s\}$ . Assume that  $X \in \mathbb{R}^{n \times d}$  satisfies Assumption 1 for some  $c, C > 0$ , and for  $\theta \in \Theta$ , let  $\mathcal{P}_\theta := \{\mathbf{N}_n(X\theta, \sigma^2 I_n)\}$ . Then, setting  $L(\theta, \theta') := n^{-1} \|X(\theta - \theta')\|_2^2$ , we have

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, L) \asymp_{c,C} \sigma^2 \cdot \frac{s \log(ed/s) + \log(1/\delta)}{n}.$$

*Proof.* Let  $\theta_1 := 0 \in \mathbb{R}^d$  and  $\theta_2 := e_1 \sqrt{\frac{\sigma^2}{n} \log\left(\frac{1}{4\delta(1-\delta)}\right)} \in \mathbb{R}^d$ , so that  $\theta_1, \theta_2 \in \Theta$ . For  $\ell \in \{1, 2\}$ , let  $P_\ell := \mathbf{N}_n(X\theta_\ell, \sigma^2 I_n)$ . Then  $\text{KL}(P_1, P_2) = \frac{1}{2\sigma^2} \|X\theta_2\|_2^2 \leq \frac{n}{2\sigma^2} \|\theta_2\|_2^2 < \log\left(\frac{1}{4\delta(1-\delta)}\right)$ . Therefore, combining Theorem 4, Assumption 1 (since  $\theta_1 - \theta_2 \in \mathcal{C}_{\text{WRE}}(s \vee 2, 3)$  when  $d \geq 2$ ) and Corollary 6, we have for  $\delta \in (0, 1/4]$  that

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, L) \geq \frac{1}{4n} \|X(\theta_1 - \theta_2)\|_2^2 \geq \frac{c^2}{4} \|\theta_1 - \theta_2\|_2^2 \geq \frac{c^2 \sigma^2 \log(1/\delta)}{20n}. \quad (15)$$

To capture the first term in the lower bound, let us first consider the case where  $d \geq 4e$ . By Lemma 27, there exist  $M \in \mathbb{N}$  with

$$\log M \geq \frac{3(1 \vee \lfloor s/2 \rfloor \wedge \lfloor d/(4e) \rfloor)}{4} \log\left(\frac{d}{4(1 \vee \lfloor s/2 \rfloor \wedge \lfloor d/(4e) \rfloor)}\right)$$

and  $\mathcal{V} := \{v^{(1)}, \dots, v^{(M)}\}$  such that  $\mathcal{V}$  forms a Euclidean norm  $(1/2)$ -packing of the set  $\{\theta \in \mathbb{R}^d : \|\theta\|_0 \leq 1 \vee \lfloor s/2 \rfloor \wedge \lfloor d/(4e) \rfloor, \|\theta\|_2 \leq 1\}$ . For  $m \in [M]$ , define  $\theta^{(m)} := v^{(m)} \sqrt{\frac{\sigma^2 \log M}{4C^2 n}}$ , so that  $\theta^{(m)} \in \Theta$ , and define  $P^{(m)} := \mathbf{N}_n(X\theta^{(m)}, \sigma^2 I_n)$ . Further define  $Q := \mathbf{N}_n(0, \sigma^2 I_n)$  so that  $\text{KL}(P^{(m)}, Q) = \frac{1}{2\sigma^2} \|X\theta^{(m)}\|_2^2 \leq \frac{\log M}{8}$  for  $m \in [M]$ . Since  $\log M \geq 3/4$  and since  $M \mapsto \log(2 - M^{-1})/\log M$  is decreasing for  $M \geq e^{3/4}$ , we have

$$\frac{M^{-1} \sum_{m=1}^M \text{KL}(P^{(m)}, Q) + \log(2 - M^{-1})}{\log M} \leq \frac{1}{8} + \frac{\log(2 - M^{-1})}{\log M} \leq \frac{7}{10}.$$

Since  $\theta^{(m)} - \theta^{(m')} \in \mathcal{C}_{\text{WRE}}(s \vee 2, 3)$  for distinct  $m, m' \in [M]$ , we have by Lemma 7, that for  $\delta \in (0, 1/4]$ ,

$$\begin{aligned} \mathcal{M}(\delta, \mathcal{P}_\Theta, L) &\geq \frac{1}{4n} \cdot \min_{m \neq m'} \|X(\theta^{(m)} - \theta^{(m')})\|_2^2 \geq \frac{c^2}{4} \min_{m \neq m'} \|\theta^{(m)} - \theta^{(m')}\|_2^2 \\ &> \frac{c^2 \sigma^2 \{s/4 \wedge 3d/(32e)\} \log\left(\frac{d}{4s}\right)}{2^6 C^2 n} \geq \frac{c^2 \sigma^2 s \log(ed/s)}{2^{13} C^2 n}. \end{aligned} \quad (16)$$

From (15) and (16), we deduce that for  $d \geq 4e$  and  $\delta \in (0, 1/4]$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, L) \geq \frac{c^2 \sigma^2 \log(1/\delta)}{40n} + \frac{c^2 \sigma^2 s \log(ed/s)}{2^{14} C^2 n}.$$

On the other hand, when  $d < 4e$ , we have by (15) that for  $\delta \in (0, 1/4]$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, L) \geq \frac{c^2 \sigma^2 \log(1/\delta)}{20n} \geq c^2 \sigma^2 \cdot \frac{s \log(ed/s) + \log(1/\delta)}{200n}.$$

This proves the lower bound.

The upper bound is achieved, for example, by the SLOPE estimator (Bogdan et al., 2015), with a  $\delta$ -independent tuning parameter. In particular, applying Bellec, Lecué and Tsybakov (2018, Eq. (6.2) and (6.5)) with  $\gamma = \tau = 1/4$  in their notation yields that for any  $\delta \in (0, 1)$ ,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, L) \leq 100(4 + \sqrt{2})\sigma^2 \left( \frac{s \log(ed/s)}{c^2 n} + \frac{\log(1/\delta)}{n} \right),$$

as required.  $\square$

### 3.4 Nonparametric density estimation

Turning to a nonparametric statistical problem, we now consider density estimation at a point. We remark that this is the only one of our examples in which the matching upper bound is not available in the literature, to the best of our knowledge. We provide, using Lepski's method, a  $\delta$ -independent estimator in Lemma 29.

To fix the setting, let  $\beta, \gamma > 0$  and  $m := \lceil \beta \rceil - 1$ . The Hölder class  $\mathcal{F}(\beta, \gamma)$  of density functions is the set of all Borel measurable functions  $f : \mathbb{R} \rightarrow [0, \infty)$  satisfying that  $\int_{-\infty}^{\infty} f(x) dx = 1$ , that  $f$  is  $m$ -times differentiable and that

$$|f^{(m)}(x) - f^{(m)}(x')| \leq \gamma |x - x'|^{\beta - m},$$

for all  $x, x' \in \mathbb{R}$ .

**Proposition 15.** *Let  $n \in \mathbb{N}$ ,  $\delta \in (0, 1/4]$ ,  $\Theta := \mathcal{F}(\beta, \gamma)$  be the Hölder class of density functions and for  $f \in \Theta$ , define  $\mathcal{P}_f := \{P_f^{\otimes n}\}$  where  $P_f$  is the distribution with density  $f$ . Fix  $x_0 \in \mathbb{R}$  and let  $L(f_1, f_2) := \{f_1(x_0) - f_2(x_0)\}^2$ . Then*

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, L) \asymp_\beta \gamma^{2/(\beta+1)} \cdot \left\{ \left( \frac{\log(1/\delta)}{n} \right)^{2\beta/(2\beta+1)} \wedge 1 \right\}.$$

*Proof.* By the translation invariance of  $\mathcal{F}(\beta, \gamma)$ , the minimax quantile  $\mathcal{M}(\delta, \mathcal{P}_\Theta, L)$  is the same for all  $x_0 \in \mathbb{R}$ , so assume without loss of generality that  $x_0 = 0$ . Let  $m := \lceil \beta \rceil - 1$  and take  $K : \mathbb{R} \rightarrow [0, \infty)$  to be the infinitely differentiable density given by  $K(x) \propto e^{-1/(1-4x^2)} \mathbb{1}_{\{|x| < 1/2\}}$ , which satisfies

$$\|K\|_{C^\beta} := \sup_{x \neq x'} \frac{|K^{(m)}(x) - K^{(m)}(x')|}{|x - x'|^{\beta - m}} < \infty.$$

Define  $\tilde{\gamma} := \gamma / \|K\|_{C^\beta}$ , and for  $\lambda \in (0, 1]$  that will be chosen later to depend on  $n, \beta$  and  $\delta$ , define  $g, h : \mathbb{R} \rightarrow [0, \infty)$  by

$$g(x) := \tilde{\gamma}^{1/(\beta+1)} K(\tilde{\gamma}^{1/(\beta+1)} x), \quad h(x) := \lambda \tilde{\gamma}^{1/(\beta+1)} K(\lambda^{-1/\beta} \tilde{\gamma}^{1/(\beta+1)} x). \quad (17)$$

Then  $g \in \mathcal{F}(\beta, \gamma)$  because it is  $m$ -times differentiable, and for  $x, x' \in \mathbb{R}$ ,

$$\begin{aligned} |g^{(m)}(x) - g^{(m)}(x')| &= \tilde{\gamma}^{\frac{m+1}{\beta+1}} |K^{(m)}(\tilde{\gamma}^{1/(\beta+1)} x) - K^{(m)}(\tilde{\gamma}^{1/(\beta+1)} x')| \\ &\leq \tilde{\gamma}^{\frac{m+1}{\beta+1}} \|K\|_{C^\beta} |\tilde{\gamma}^{1/(\beta+1)} x - \tilde{\gamma}^{1/(\beta+1)} x'|^{\beta - m} = \gamma |x - x'|^{\beta - m}. \end{aligned}$$






Figure 1: Illustration of the construction in Proposition 15. Left: The functions  $g$  (solid) and  $h$  (dot-dashed) from (17). Right: The functions  $f_0$  (solid) and  $f_1$  (dot-dashed) from (18) and (19).

Similarly,  $|h^{(m)}(x) - h^{(m)}(x')| \leq \gamma|x - x'|^{\beta-m}$  for  $x, x' \in \mathbb{R}$ , and  $\int_{-\infty}^{\infty} h = \lambda^{(\beta+1)/\beta}$ . Moreover,

$$h(x) \leq \lambda \tilde{\gamma}^{1/(\beta+1)} K(\tilde{\gamma}^{1/(\beta+1)} x) = \lambda g(x)$$

for all  $x \in \mathbb{R}$ . Now define  $f_0, f_1 : \mathbb{R} \rightarrow [0, \infty)$  by

$$f_0(x) := \frac{(g+h)(x) + g(2\tilde{\gamma}^{-1/(\beta+1)} - x)}{2 + \lambda^{(\beta+1)/\beta}}, \quad (18)$$

$$f_1(x) := \frac{g(x) + (g+h)(2\tilde{\gamma}^{-1/(\beta+1)} - x)}{2 + \lambda^{(\beta+1)/\beta}}; \quad (19)$$

see Figure 1. Since  $g, h$  are zero outside the interval  $[-\tilde{\gamma}^{-1/(\beta+1)}/2, \tilde{\gamma}^{-1/(\beta+1)}/2]$ , we deduce that  $f_0, f_1 \in \mathcal{F}(\beta, \gamma)$  and since  $K(0) \geq 1$ ,

$$f_0(0) - f_1(0) = \frac{h(0)}{2 + \lambda^{(\beta+1)/\beta}} = \frac{\lambda \tilde{\gamma}^{1/(\beta+1)} K(0)}{2 + \lambda^{(\beta+1)/\beta}} \geq \frac{\lambda \tilde{\gamma}^{1/(\beta+1)}}{3}.$$

In addition, writing  $I := [-\lambda^{1/\beta} \tilde{\gamma}^{-1/(\beta+1)}/2, \lambda^{1/\beta} \tilde{\gamma}^{-1/(\beta+1)}/2]$  for the support of  $h$  and  $I' := 2\tilde{\gamma}^{-1/(\beta+1)} + I$ , we have  $f_0 = f_1$  on  $\mathbb{R} \setminus (I \cup I')$ . Thus, since  $\log(1 + \lambda) \leq \lambda$ , it follows that the product densities  $\{f_j^n : j \in \{0, 1\}\}$  given by  $f_j^n(x_1, \dots, x_n) := \prod_{i=1}^n f_j(x_i)$  satisfy

$$\begin{aligned} \text{KL}(f_0^n, f_1^n) &= n \text{KL}(f_0, f_1) = n \int_{I \cup I'} f_0 \log\left(\frac{f_0}{f_1}\right) \\ &= \frac{n}{2 + \lambda^{(\beta+1)/\beta}} \int_I \left\{ (g+h) \log\left(\frac{g+h}{g}\right) + g \log\left(\frac{g}{g+h}\right) \right\} \\ &= \frac{n}{2 + \lambda^{(\beta+1)/\beta}} \int_I h \log\left(1 + \frac{h}{g}\right) \end{aligned}$$

$$\leq \frac{n}{2} \log(1 + \lambda) \int_I h \leq \frac{n}{2} \lambda^{(2\beta+1)/\beta}.$$

Taking  $\lambda = \left\{ \frac{1}{n} \log\left(\frac{1}{4\delta(1-\delta)}\right) \right\}^{\beta/(2\beta+1)} \wedge 1 \in (0, 1]$ , we have that  $\text{KL}(f_0^n, f_1^n) \leq \frac{1}{2} \log\left(\frac{1}{4\delta(1-\delta)}\right) < \log\left(\frac{1}{4\delta(1-\delta)}\right)$ , so by Theorem 4 and Corollary 6, for  $\delta \in (0, 1/4]$ ,

$$\begin{aligned} \mathcal{M}(\delta, \mathcal{P}_\Theta, L) &\geq \mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) \geq \left( \frac{\lambda \tilde{\gamma}^{1/(\beta+1)}}{2 \cdot 3} \right)^2 \\ &\geq \frac{\gamma^{2/(\beta+1)}}{36 \cdot 5^{2\beta/(2\beta+1)} \|K\|_{C^\beta}^{2/(\beta+1)}} \left\{ \left( \frac{\log(1/\delta)}{n} \right)^{2\beta/(2\beta+1)} \wedge 1 \right\}. \end{aligned}$$

This completes the proofs of the lower bound. The upper bound follows from Lemma 29, after fixing a bounded kernel supported on  $[-1, 1]$ .  $\square$

### 3.5 Isotonic regression

Other natural nonparametric estimation problems are provided by shape-constrained models (Groeneboom and Jongbloed, 2014; Samworth and Sen, 2018), and here we consider isotonic regression. The matching upper bound is attained by the projected isotonic least squares estimator, which is  $\delta$ -independent.

**Proposition 16.** *Let  $n \geq 2$ ,  $\delta \in (0, 1/4]$ ,  $\Theta := \{\theta = (\theta_1, \dots, \theta_n)^\top \in [0, 1]^n : \theta_1 \leq \dots \leq \theta_n\}$  and  $\mathcal{P}_\theta := \{\mathbf{N}_n(\theta, I_n)\}$  for  $\theta \in \Theta$ . Then*

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, n^{-1} \|\cdot\|_2^2) \asymp \left( \frac{1}{n^{2/3}} + \frac{\log(1/\delta)}{n} \right) \wedge 1.$$

*Proof.* First consider the case where  $\delta \in [e^{-n}/3, 1/4]$ . Let  $\theta^{(1)} := 0$ ,  $\theta^{(2)} := \sqrt{\frac{1}{n} \log\left(\frac{1}{4\delta(1-\delta)}\right)} \mathbf{1}_n$  and  $P^{(\ell)} := \mathbf{N}_n(\theta^{(\ell)}, I_n)$  for  $\ell \in \{1, 2\}$ . Then  $\theta^{(\ell)} \in \Theta$  and  $P^{(\ell)} \in \mathcal{P}_{\theta^{(\ell)}}$  for  $\ell \in \{1, 2\}$  and

$$\text{KL}(P^{(1)}, P^{(2)}) = \frac{1}{2} \log\left(\frac{1}{4\delta(1-\delta)}\right) < \log\left(\frac{1}{4\delta(1-\delta)}\right).$$

Therefore, by Corollary 6,

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, n^{-1} \|\cdot\|_2^2) \geq \frac{\|\theta^{(1)} - \theta^{(2)}\|_2^2}{4n} \geq \frac{\log(1/\delta)}{20n}, \quad (20)$$

for  $\delta \in [e^{-n}/3, 1/4]$ . Next, for the case where  $\delta \in (0, e^{-n}/3)$ , we have by (20) that

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, n^{-1} \|\cdot\|_2^2) \geq \mathcal{M}(e^{-n}, \mathcal{P}_\Theta, n^{-1} \|\cdot\|_2^2) \geq \frac{1}{20}. \quad (21)$$

Combining (20) and (21) yields

$$\mathcal{M}(\delta, \mathcal{P}_\Theta, n^{-1} \|\cdot\|_2^2) \geq \frac{1}{20} \left( \frac{\log(1/\delta)}{n} \wedge 1 \right), \quad (22)$$

for  $\delta \in (0, 1/4]$ .

For the second part of the lower bound, we let  $\Theta_0 \subseteq \Theta$  denote the finite set constructed in the proof of [Chatterjee, Guntuboyina and Sen \(2015, Theorem 5.3\)](#) to obtain

$$\inf_{\hat{\theta} \in \hat{\Theta}} \max_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} \mathbb{E}_{P_{\theta_0}} (n^{-1} \|\hat{\theta} - \theta_0\|_2^2) \geq \frac{1}{2^9 \cdot 3^{1/3}} \cdot n^{-2/3}.$$

In this construction,  $D := \sup_{\theta, \theta' \in \Theta_0} \|\theta - \theta'\|_2 \in [\frac{n^{1/6}}{2^3 \cdot 3^{1/6}}, n^{1/6}]$ . Hence by [Theorem 8](#) with  $\epsilon = 2^{-6}$ , we obtain that for  $\delta \in (0, 1/1000]$ ,

$$\mathcal{M}_-(\delta, \mathcal{P}_{\Theta}, n^{-1} \|\cdot\|_2^2) \geq \frac{1}{2^{18} \cdot 3^{1/3}} \cdot n^{-2/3}.$$

Then, applying [Proposition 9](#) with  $k = 4$  in conjunction with [Theorem 4](#), we deduce that for  $\delta \in (0, 1/4]$ ,

$$\mathcal{M}(\delta, \mathcal{P}_{\Theta}, n^{-1} \|\cdot\|_2^2) \geq \mathcal{M}_-(\delta, \mathcal{P}_{\Theta}, n^{-1} \|\cdot\|_2^2) \geq \frac{1}{2^{26} \cdot 3^3} \cdot n^{-2/3}. \quad (23)$$

Combining [\(22\)](#) and [\(23\)](#) proves the lower bound.

For the upper bound, let  $\hat{\theta}$  denote the isotonic least squares estimator. By [Bellec \(2018, Corollary 3.3\)](#), for every  $\delta \in (0, 1]$ , we have with probability at least  $1 - \delta$  that

$$n^{-1} \|\hat{\theta} - \theta\|_2^2 \lesssim \frac{1}{n^{2/3}} + \frac{\log(1/\delta)}{n}.$$

Now define  $\tilde{\theta} = (\tilde{\theta}_1, \dots, \tilde{\theta}_n)^\top \in \mathbb{R}^n$  by  $\tilde{\theta}_i := 0 \vee \hat{\theta}_i \wedge 1$  for  $i \in [n]$ . Then, since  $n^{-1} \|\tilde{\theta} - \theta\|_2^2 \leq n^{-1} \|\hat{\theta} - \theta\|_2^2$  and  $n^{-1} \|\tilde{\theta} - \theta\|_2^2 \leq 1$ , we deduce that

$$\mathcal{M}(\delta, \mathcal{P}_{\Theta}, n^{-1} \|\cdot\|_2^2) \lesssim \left( \frac{1}{n^{2/3}} + \frac{\log(1/\delta)}{n} \right) \wedge 1$$

for  $\delta \in (0, 1]$ . □

### 3.6 Stochastic convex optimisation

As our final example, we move away from the traditional setting of statistical estimation and consider stochastic convex optimisation ([Duchi, 2018](#)), where the goal is to understand the iteration complexity of minimising a convex function in expectation, given access only to queries at random observations. To do so, we require some preliminaries. Let  $R > 0$ , let  $\mathcal{X} \subseteq \mathbb{R}^d$  be closed and convex such that  $\mathbb{B}_2^d(R) \subseteq \mathcal{X}$ , and let  $\mathcal{Y}$  be a measurable space. For  $P \in \mathcal{Q}(\mathcal{Y})$  and  $\gamma > 0$ , let  $\mathcal{F}(P, \gamma)$  denote the set of functions  $f : \mathbb{R}^d \times \mathcal{Y} \rightarrow \mathbb{R}$  for which

- (i)  $f(x, \cdot) : \mathcal{Y} \rightarrow \mathbb{R}$  is  $P$ -integrable for each  $x \in \mathcal{X}$ ;
- (ii)  $f(\cdot, y) : \mathbb{R}^d \rightarrow \mathbb{R}$  is convex for each  $y \in \mathcal{Y}$  with subdifferential at  $x \in \mathbb{R}^d$  denoted by  $\partial_x f(x, y)$ ;
- (iii) for each  $x \in \mathcal{X}$ , define the function  $g_x : \mathcal{Y} \rightarrow \mathbb{R}^d$  by  $g_x(y) := \operatorname{argmin}_{x' \in \partial_x f(x, y)} \|x'\|_2$ , and assume that  $\sup_{x \in \mathcal{X}} \mathbb{E}_P \{\|g_x(Y)\|_2^2\} \leq \gamma^2$ .

The selected subgradient  $g_x$  in part (iii) of this definition is well-defined as the subdifferential is compact and convex (Rockafellar, 1997, Theorem 23.4 and p. 215), and moreover  $g_x$  is measurable on  $\mathcal{Y}$  for each  $x \in \mathcal{X}$ , by Aliprantis and Border (2007, Theorem 18.19). For  $f \in \mathcal{F}(P, \gamma)$ , define  $F_P : \mathcal{X} \rightarrow \mathbb{R}$  by  $F_P(x) := \mathbb{E}_P\{f(x; Y)\}$ , so that  $F_P$  is convex on  $\mathcal{X}$ . The *optimality gap*  $L^{\text{opt}}(\cdot; P, f) : \mathcal{X} \rightarrow [0, \infty)$  is defined to be  $L^{\text{opt}}(x; P, f) := F_P(x) - \inf_{x' \in \mathcal{X}} F_P(x')$ . For  $P_1, P_2 \in \mathcal{Q}(\mathcal{Y})$  and  $f \in \mathcal{F}(P_1, \gamma) \cap \mathcal{F}(P_2, \gamma)$ , the *f-separation* between  $P_1$  and  $P_2$  is defined as

$$\text{sep}(P_1, P_2; f) := \sup \left\{ \Delta \geq 0 : L^{\text{opt}}(x; P_2, f) > \Delta \text{ for all } x \in \mathcal{X} \text{ with } L^{\text{opt}}(x; P_1, f) \leq \Delta \right\}.$$

Now define the set

$$\mathcal{G}(\gamma, R) := \left\{ (P, f) : P \in \mathcal{Q}(\mathcal{Y}), f \in \mathcal{F}(P, \gamma) \text{ and } \|x^*\|_2 \leq R \text{ for some } x^* \in \underset{x' \in \mathcal{X}}{\text{argmin}} F_P(x') \right\}.$$

For each  $t \in \mathbb{N}$ , let  $\widehat{\mathcal{X}}_t$  denote the set of all measurable functions from  $\mathcal{Y}^t$  to  $\mathcal{X}$ ; in particular, this includes methods such as stochastic (sub)gradient descent (SGD) or stochastic versions of Newton's algorithm. For  $\delta \in (0, 1]$  and  $T \in \mathbb{N}$ , the minimax quantile  $\mathcal{M}^{\text{opt}}$  for the optimality gap (cf. Carmon and Hinder, 2024, Eq. (1)) is given by

$$\mathcal{M}^{\text{opt}}(\delta; T) := \inf_{\widehat{x}_\delta \in \widehat{\mathcal{X}}_T} \sup_{(P, f) \in \mathcal{G}(\gamma, R)} \text{Quantile}(1 - \delta; L^{\text{opt}}(\widehat{x}_\delta; P, f), P^{\otimes T}).$$

By a virtually identical argument to that given in the proof of Theorem 4,

$$\mathcal{M}^{\text{opt}}(\delta; T) \geq \inf \left\{ r \in [0, \infty) : \inf_{\widehat{x}_\delta \in \widehat{\mathcal{X}}_T} \sup_{(P, f) \in \mathcal{G}(\gamma, R)} P^{\otimes T} \{ L^{\text{opt}}(\widehat{x}_\delta; P, f) > r \} \leq \delta \right\} =: \mathcal{M}_-^{\text{opt}}(\delta; T). \quad (24)$$

We next present a variant of Lemma 5, specialised to the current optimisation setting. It can also be understood as a high-probability variant of, e.g., Duchi (2018, Eq. (5.2.3)).

**Lemma 17.** *Let  $T \in \mathbb{N}$ ,  $\gamma, R > 0$  and  $\delta \in (0, 1/2)$ . Suppose  $(P_1, f), (P_2, f) \in \mathcal{G}(\gamma, R)$  and  $\text{TV}(P_1^{\otimes T}, P_2^{\otimes T}) < 1 - 2\delta$ . Then  $\mathcal{M}_-^{\text{opt}}(\delta; T) \geq \text{sep}(P_1, P_2; f)$ .*

*Proof.* Given  $\epsilon > 0$ , choose  $r \in [0, \infty)$  such that  $r > \text{sep}(P_1, P_2; f) - \epsilon$  and such that for every  $x \in \mathcal{X}$  we have  $L^{\text{opt}}(x; P_2, f) > r$  whenever  $L^{\text{opt}}(x; P_1, f) \leq r$ . Then

$$\begin{aligned} & \inf_{\widehat{x}_\delta \in \widehat{\mathcal{X}}_T} \sup_{(P, f) \in \mathcal{G}(\gamma, R)} P^{\otimes T} \{ L^{\text{opt}}(\widehat{x}_\delta; P, f) > r \} \\ & \geq \frac{1}{2} \inf_{\widehat{x}_\delta \in \widehat{\mathcal{X}}_T} \left\{ P_1^{\otimes T} \{ L^{\text{opt}}(\widehat{x}_\delta; P_1, f) > r \} + P_2^{\otimes T} \{ L^{\text{opt}}(\widehat{x}_\delta; P_2, f) > r \} \right\} \\ & \geq \frac{1}{2} \inf_{\widehat{x}_\delta \in \widehat{\mathcal{X}}_T} \left\{ 1 - P_1^{\otimes T} \{ L^{\text{opt}}(\widehat{x}_\delta; P_2, f) > r \} + P_2^{\otimes T} \{ L^{\text{opt}}(\widehat{x}_\delta; P_2, f) > r \} \right\} \\ & \geq \frac{1}{2} \{ 1 - \text{TV}(P_1^{\otimes T}, P_2^{\otimes T}) \} > \delta. \end{aligned}$$

We deduce that  $\mathcal{M}_-^{\text{opt}}(\delta, T) \geq r > \text{sep}(P_1, P_2; f) - \epsilon$ . Since  $\epsilon > 0$  was arbitrary, the result follows.  $\square$




Figure 2: Illustration of the functions in the proof of Proposition 18 with  $f : \mathcal{X} \times \{-1, 1\}$  given by  $f(x, y) = \gamma|x + yR|$ .

The following result, first obtained by [Carmon and Hinder \(2024, Proposition 1\)](#), then provides tight bounds on the minimax  $(1 - \delta)$ th quantile over  $\mathcal{G}(\gamma, R)$ . The upper bound is attained by clipped SGD, where the truncation level is  $\delta$ -dependent. We provide here a short proof of the lower bound using our techniques in conjunction with the lower bound construction of [Duchi \(2018, Eq. \(5.2.7\)\)](#).

**Proposition 18.** *Let  $T \in \mathbb{N}$ ,  $\gamma, R > 0$  and  $\delta \in (0, 1/4]$ . Then*

$$\mathcal{M}^{\text{opt}}(\delta; T) \asymp \gamma R \cdot \left\{ \sqrt{\frac{\log(1/\delta)}{T}} \wedge 1 \right\}.$$

*Proof.* For the lower bound, we initially consider the case  $d = 1$  where  $\mathcal{X} \supseteq [-R, R]$ . Take  $\mathcal{Y} = \{-1, 1\}$  and consider the function  $f : \mathbb{R} \times \mathcal{Y} \rightarrow \mathbb{R}$  given by  $f(x, y) := \gamma|x + yR|$ . Let  $g : [0, 1) \rightarrow \mathbb{R}$  be the strictly increasing function given by  $g(x) := x \log\left(\frac{1+x}{1-x}\right)$ , and set  $s := g^{-1}\left(\frac{1}{2T} \log\left(\frac{1}{4\delta(1-\delta)}\right)\right) \in [0, 1)$ . Now let  $P_1 = \text{Ber}\left(\frac{1+s}{2}\right)$  and  $P_2 = \text{Ber}\left(\frac{1-s}{2}\right)$ , so that  $\text{KL}(P_1, P_2) = g(s)$ ,

$$\inf_{x \in \mathcal{X}} F_{P_1}(x) = F_{P_1}(-R) = \gamma R(1 - s) = F_{P_2}(R) = \inf_{x \in \mathcal{X}} F_{P_2}(x),$$

and  $(P_1, f), (P_2, f) \in \mathcal{G}(\gamma, R)$ . Further,  $\text{sep}(P_1, P_2; f) = F_{P_1}(0) - \gamma R(1 - s) = \gamma s R$ ; see [Figure 2](#). Applying the Bretagnolle–Huber inequality, we deduce that

$$\text{TV}(P_1^{\otimes T}, P_2^{\otimes T}) \leq \sqrt{1 - \exp\{-T \cdot \text{KL}(P_1, P_2)\}} = \{1 - \sqrt{4\delta(1 - \delta)}\}^{1/2} < 1 - 2\delta.$$

Thus, by Lemma 17,  $\mathcal{M}^{\text{opt}}(\delta; T) \geq \gamma s R$ . Now consider the case where  $\log(1/\delta) \leq T$ . Then, since  $\delta \in (0, 1/4]$  and  $g^{-1}$  is a strictly increasing function,

$$s \geq g^{-1}\left(\frac{\log(1/\delta)}{10T}\right) \geq \sqrt{\frac{\log(1/\delta)}{30T}},$$

where the final inequality follows since  $g(x) \leq 3x^2$  for  $x \in [0, 4/5]$ , so that  $g^{-1}(w) \geq (w/3)^{1/2}$  for  $w \in [0, 7/4]$ . On the other hand, when  $\log(1/\delta) > T$ , we have  $s > g^{-1}(1/10) \geq 1/5$ . Thus, by (24),

$$\mathcal{M}^{\text{opt}}(\delta; T) \geq \mathcal{M}_-^{\text{opt}}(\delta; T) \geq \frac{\gamma R}{30^{1/2}} \cdot \left\{ \sqrt{\frac{\log(1/\delta)}{T}} \wedge 1 \right\}.$$

Turning to the more general case where  $d > 1$  and  $\mathcal{X} \supseteq \mathbb{B}_2^d(R)$ , we consider the function  $\tilde{f} : \mathbb{R}^d \times \mathcal{Y}$  given by  $\tilde{f}(x, y) := f(x^\top e_1, y)$ , where  $f$  is as defined above. The same lower bound then follows.

To obtain a matching upper bound, we set  $\hat{x}_T = 0$  if  $\log(1/\delta) > T$  and apply clipped SGD with a  $\delta$ -dependent clipping parameter otherwise. The conclusion then follows from Carmon and Hinder (2024, Theorem 4).  $\square$

## 4 Proofs from Section 2

*Proof of Proposition 2.* For  $\theta \in \Theta$ ,  $P_\theta \in \mathcal{P}_\theta$  and  $\hat{\theta} \in \hat{\Theta}$ , define  $r(P_\theta, \hat{\theta}) := \text{Quantile}(1 - \delta; P_\theta, L(\hat{\theta}, \theta))$ . Then

$$\begin{aligned} \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} \mathbb{E}_{P_\theta} L(\hat{\theta}, \theta) &\geq \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{L(\hat{\theta}, \theta) \geq r(P_\theta, \hat{\theta})\} \cdot r(P_\theta, \hat{\theta}) \\ &\geq \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} \delta \cdot r(P_\theta, \hat{\theta}) = \delta \mathcal{M}(\delta), \end{aligned}$$

as desired.  $\square$

*Proof of Theorem 4.* We begin with the lower bound on  $\mathcal{M}(\delta)$ . For  $\theta \in \Theta$ ,  $P_\theta \in \mathcal{P}_\theta$  and  $\hat{\theta} \in \hat{\Theta}$ , define  $r(P_\theta, \hat{\theta}) := \text{Quantile}(1 - \delta; P_\theta, L(\hat{\theta}, \theta))$ , so that  $P_\theta \{L(\hat{\theta}, \theta) > r(P_\theta, \hat{\theta})\} \leq \delta$ . Then

$$P_\theta \left\{ L(\hat{\theta}, \theta) > \sup_{\theta' \in \Theta} \sup_{P_{\theta'} \in \mathcal{P}_{\theta'}} r(P_{\theta'}, \hat{\theta}) \right\} \leq \delta,$$

for all  $\theta \in \Theta$  and  $P_\theta \in \mathcal{P}_\theta$ , so that

$$\sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} r(P_\theta, \hat{\theta}) \geq \inf \left\{ r \in [0, \infty) : \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{L(\hat{\theta}, \theta) > r\} \leq \delta \right\} =: r(\hat{\theta}). \quad (25)$$

Further, if  $r > \inf_{\hat{\theta} \in \hat{\Theta}} r(\hat{\theta})$ , then there exists  $\tilde{\theta} \in \hat{\Theta}$  and  $\epsilon > 0$  such that  $r(\tilde{\theta}) \leq r - \epsilon$ , so

$$\begin{aligned} \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{L(\hat{\theta}, \theta) > r\} &\leq \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{L(\tilde{\theta}, \theta) > r\} \\ &\leq \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{L(\tilde{\theta}, \theta) > r(\tilde{\theta}) + \epsilon\} \leq \delta. \end{aligned} \quad (26)$$

Hence,

$$\begin{aligned}
\mathcal{M}_-(\delta) &= \inf \left\{ r \in [0, \infty) : \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{L(\hat{\theta}, \theta) > r\} \leq \delta \right\} \\
&\stackrel{(i)}{\leq} \inf \left\{ r \in [0, \infty) : r > \inf_{\hat{\theta} \in \hat{\Theta}} r(\hat{\theta}) \right\} \\
&= \inf_{\hat{\theta} \in \hat{\Theta}} r(\hat{\theta}) \stackrel{(ii)}{\leq} \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} r(P_\theta, \hat{\theta}) = \mathcal{M}(\delta),
\end{aligned}$$

where inequality (i) follows from (26) and inequality (ii) follows from (25).

For the upper bound, let  $\epsilon > 0$  and  $\xi \in (0, \delta)$ . There exists  $r \leq \mathcal{M}_-(\delta - \xi) + \epsilon$  such that

$$\inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{L(\hat{\theta}, \theta) > r\} \leq \delta - \xi.$$

It follows that there exists  $\tilde{\theta} \in \hat{\Theta}$  such that

$$\sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{L(\tilde{\theta}, \theta) > r\} \leq \delta.$$

Hence,  $\mathcal{M}(\delta) \leq r \leq \mathcal{M}_-(\delta - \xi) + \epsilon$ . Since  $\epsilon > 0$  was arbitrary, we deduce that  $\mathcal{M}(\delta) \leq \mathcal{M}_-(\delta - \xi)$ .

For the final claim, observe that  $\delta \mapsto \mathcal{M}_-(\delta)$  is a decreasing function on  $(0, 1]$ , so it has at most countably many discontinuities. But if  $\mathcal{M}_-(\cdot)$  is continuous at  $\delta$ , then we must have  $\mathcal{M}(\delta) = \mathcal{M}_-(\delta)$ , by (1).  $\square$

*Proof of Lemma 5.* There is nothing to prove in the case  $g(\eta) = 0$ , so we may assume without loss of generality that  $g(\eta) > 0$ . For  $a \in (0, 1)$ , define  $\tilde{g}_a : [0, \infty) \rightarrow \{0, 1\}$  by  $\tilde{g}_a(x) := \mathbb{1}_{\{g(x) > ag(\eta)\}}$  for  $x \in [0, \infty)$ , so that  $\tilde{g}_a$  is an increasing function. By applying Le Cam's two-point lemma (stated as Lemma 21 for convenience), we obtain

$$\begin{aligned}
\inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{L(\hat{\theta}, \theta) > ag(\eta)\} &= \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} \mathbb{E}_{P_\theta} \{\tilde{g}_a(d(\hat{\theta}, \theta))\} \\
&\geq \frac{\tilde{g}_a(\eta)}{2} (1 - \text{TV}(P_1, P_2)) > \delta.
\end{aligned}$$

We deduce that  $\mathcal{M}_-(\delta) \geq ag(\eta)$ . Since  $a \in (0, 1)$  was arbitrary, the conclusion follows.  $\square$

*Proof of Lemma 7.* Again, we may assume without loss of generality that  $g(\eta) > 0$ . For  $a \in (0, 1)$ , define  $\tilde{g}_a : [0, \infty) \rightarrow \{0, 1\}$  by  $\tilde{g}_a(x) := \mathbb{1}_{\{g(x) > ag(\eta)\}}$  for  $x \in [0, \infty)$ . By Fano's lemma (stated as Lemma 22 for convenience),

$$\begin{aligned}
\inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta \{L(\hat{\theta}, \theta) > ag(\eta)\} &= \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} \mathbb{E}_{P_\theta} \{\tilde{g}_a(d(\hat{\theta}, \theta))\} \\
&\geq \tilde{g}_a(\eta) \left\{ 1 - \frac{M^{-1} \inf_{Q \in \mathcal{Q}(X)} \sum_{j=1}^M \text{KL}(P_j, Q) + \log(2 - M^{-1})}{\log M} \right\} \geq \epsilon > \delta.
\end{aligned}$$

Since  $a \in (0, 1)$  was arbitrary, the conclusion follows.  $\square$




Figure 3: Illustration of the construction of  $\hat{\theta}^{(1)}$  in the proof of Theorem 8. If  $x \in S(\hat{\theta})$ , then we set  $\hat{\theta}^{(1)}(x) := \hat{\theta}(x)$ ; otherwise we set  $\hat{\theta}^{(1)}(x) := \theta^{(1)} \in \Theta_1$ .

*Proof of Theorem 8.* For  $\theta \in \Theta$  and  $r \geq 0$ , let  $B(\theta, r) := \{\theta' \in \Theta : d(\theta, \theta') < r\}$  be the open ball in  $\Theta$  centred at  $\theta$  with radius  $r$ . Fix  $\epsilon > 0$  and define  $\Theta_1 := \bigcup_{\theta_0 \in \Theta_0} B(\theta_0, \epsilon D)$ , so that  $d(\theta_0, \theta_1) < (1 + \epsilon)D$  for all  $\theta_0 \in \Theta_0$  and  $\theta_1 \in \Theta_1$ . Let  $\hat{\Theta}_1$  denote the set of all measurable functions from  $\mathcal{X}$  to  $\Theta_1$ . For any  $\tilde{\theta} \in \hat{\Theta}_1$ , we have

$$\begin{aligned} \Delta &\leq \sup_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} \mathbb{E}_{P_{\theta_0}} L(\tilde{\theta}, \theta_0) \\ &= \sup_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} \left\{ \mathbb{E}_{P_{\theta_0}} \left\{ L(\tilde{\theta}, \theta_0) \mathbb{1}_{\{L(\tilde{\theta}, \theta_0) \leq g(\epsilon D)\}} \right\} + \mathbb{E}_{P_{\theta_0}} \left\{ L(\tilde{\theta}, \theta_0) \mathbb{1}_{\{L(\tilde{\theta}, \theta_0) > g(\epsilon D)\}} \right\} \right\} \\ &\leq g(\epsilon D) + g((1 + \epsilon)D) \sup_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} P_{\theta_0}(L(\tilde{\theta}, \theta_0) > g(\epsilon D)). \end{aligned}$$

Hence,

$$\inf_{\tilde{\theta} \in \hat{\Theta}_1} \sup_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} P_{\theta_0}(L(\tilde{\theta}, \theta_0) > g(\epsilon D)) \geq \frac{\Delta - g(\epsilon D)}{g((1 + \epsilon)D)}. \quad (27)$$

Now for any  $\hat{\theta} \in \hat{\Theta}$ , let

$$S(\hat{\theta}) := \{x \in \mathcal{X} : \hat{\theta}(x) \in \Theta_1\} \in \mathcal{A}.$$

Fix  $\theta^{(1)} \in \Theta_1$  and  $\hat{\theta} \in \hat{\Theta}$ , and define a new estimator  $\hat{\theta}^{(1)} \in \hat{\Theta}_1$  by

$$\hat{\theta}^{(1)}(x) := \begin{cases} \hat{\theta}(x) & \text{if } x \in S(\hat{\theta}) \\ \theta^{(1)} & \text{if } x \notin S(\hat{\theta}); \end{cases}$$



see Figure 3. Then for any  $\xi > 0$ ,  $\theta_0 \in \Theta_0$  and  $P_{\theta_0} \in \mathcal{P}_{\theta_0}$ , we have

$$\begin{aligned} P_{\theta_0}(L(\widehat{\theta}, \theta_0) > g(\epsilon D) - \xi) &\stackrel{(i)}{=} P_{\theta_0}(S(\widehat{\theta})^c) + P_{\theta_0}(S(\widehat{\theta}) \cap \{L(\widehat{\theta}, \theta_0) > g(\epsilon D) - \xi\}) \\ &= P_{\theta_0}(S(\widehat{\theta})^c) + P_{\theta_0}(S(\widehat{\theta}) \cap \{L(\widehat{\theta}^{(1)}, \theta_0) > g(\epsilon D) - \xi\}) \\ &\geq P_{\theta_0}(L(\widehat{\theta}^{(1)}, \theta_0) > g(\epsilon D) - \xi), \end{aligned}$$

where (i) follows since  $g(d(\widehat{\theta}(x), \theta_0)) \geq g(\epsilon D) > g(\epsilon D) - \xi$  for all  $x \in S(\widehat{\theta})^c$ . Therefore,

$$\begin{aligned} &\inf_{\widehat{\theta} \in \widehat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} P_\theta(L(\widehat{\theta}, \theta) > g(\epsilon D) - \xi) \\ &\geq \inf_{\widehat{\theta} \in \widehat{\Theta}} \sup_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} P_{\theta_0}(L(\widehat{\theta}^{(1)}, \theta_0) > g(\epsilon D) - \xi) \\ &\geq \inf_{\widehat{\theta} \in \widehat{\Theta}_1} \sup_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} P_{\theta_0}(L(\widetilde{\theta}, \theta_0) > g(\epsilon D) - \xi) \geq \frac{\Delta - g(\epsilon D)}{g((1 + \epsilon)D)}, \end{aligned}$$

where the final inequality follows from (27). We conclude that  $\mathcal{M}_-(\delta) \geq g(\epsilon D) - \xi$  for all  $\delta \in (0, \frac{\Delta - g(\epsilon D)}{g((1 + \epsilon)D)})$ , and since  $\xi > 0$  was arbitrary, the result follows.  $\square$

We now turn to the proof of our boosting result, Proposition 9, which requires two preliminary lemmas. Let  $(\mathcal{Y}, \mathcal{B})$  be a measurable space, and suppose that  $\Theta$  and  $\mathcal{Z}$  are topological spaces. A function  $f : \mathcal{Y} \times \Theta \rightarrow \mathcal{Z}$  is a *Carathéodory function* if:

- (i) the function  $f(\cdot, \theta) : \mathcal{Y} \rightarrow \mathcal{Z}$  is measurable (when  $\mathcal{Z}$  is equipped with its Borel  $\sigma$ -algebra) for each  $\theta \in \Theta$ ;
- (ii) the function  $f(y, \cdot) : \Theta \rightarrow \mathcal{Z}$  is continuous for each  $y \in \mathcal{Y}$ .

**Lemma 19.** *For  $m \in \mathbb{N}$ , let  $\mathcal{P}_\theta^{\otimes m} := \{P_\theta^{\otimes m} \in \mathcal{Q}(\mathcal{X}^m) : P_\theta \in \mathcal{P}_\theta\}$  and  $\mathcal{P}_\Theta^{\otimes m} := \{P_\theta^{\otimes m} : \theta \in \Theta\}$ . Suppose that  $\Theta$  is a non-empty Polish space, that  $L : \Theta \times \Theta \rightarrow [0, \infty)$  is a Carathéodory function, and there exists  $A > 0$  such that  $L(\theta_1, \theta_2) \leq A\{L(\theta_1, \theta_3) + L(\theta_2, \theta_3)\}$  for all  $\theta_1, \theta_2, \theta_3 \in \Theta$ . Then, for any  $\delta \in (0, 1]$ , we have*

$$\mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) \geq \frac{1}{2A} \mathcal{M}_-(\delta^3 + 3\delta^2(1 - \delta), \mathcal{P}_\Theta^{\otimes 3}, L).$$

*Proof.* Fix  $\epsilon > 0$ , there exists a measurable function  $\widetilde{\theta} : \mathcal{X} \rightarrow \Theta$  such that for all  $\theta \in \Theta$ ,

$$P_\theta\{L(\widetilde{\theta}, \theta) < \mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) + \epsilon\} \geq 1 - \delta.$$

Now fix  $\theta \in \Theta$ , and define the event

$$E := \left\{ (x_1, x_2, x_3) \in \mathcal{X}^3 : \sum_{\ell=1}^3 \mathbb{1}_{\{L(\widetilde{\theta}(x_\ell), \theta) < \mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) + \epsilon\}} \geq 2 \right\}$$

so that  $P_\theta^{\otimes 3}(E) \geq 1 - \delta^3 - 3\delta^2(1 - \delta)$ . Further, define  $\psi : \mathcal{X}^3 \rightarrow \text{Pow}(\Theta)$  by

$$\psi(x_1, x_2, x_3) := \left\{ \theta' \in \Theta : \sum_{\ell=1}^3 \mathbb{1}_{\{L(\widetilde{\theta}(x_\ell), \theta') < \mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) + \epsilon\}} \geq 2 \right\}.$$

For  $(x_1, x_2, x_3) \in \mathcal{X}^3$ , let  $\bar{\psi}(x_1, x_2, x_3)$  denote the closure of  $\psi(x_1, x_2, x_3)$  in  $\Theta$ , and define  $S : \mathcal{X}^3 \rightarrow \text{Pow}(\Theta)$  by

$$S(x_1, x_2, x_3) := \begin{cases} \bar{\psi}(x_1, x_2, x_3) & \text{when } \psi(x_1, x_2, x_3) \neq \emptyset \\ \Theta & \text{otherwise.} \end{cases}$$

By Lemma 20, there exists a measurable function  $\hat{\theta} : \mathcal{X}^3 \rightarrow \Theta$  such that  $\hat{\theta}(x_1, x_2, x_3) \in S(x_1, x_2, x_3)$  for all  $(x_1, x_2, x_3) \in \mathcal{X}^3$ . For any  $(x_1, x_2, x_3) \in E$ , we have  $\theta \in \psi(x_1, x_2, x_3)$ , so  $\psi(x_1, x_2, x_3)$  is non-empty; for such  $(x_1, x_2, x_3)$ , it follows that  $\hat{\theta}(x_1, x_2, x_3) \in \bar{\psi}(x_1, x_2, x_3)$ , so

$$\sum_{\ell=1}^3 \mathbb{1}_{\{L(\tilde{\theta}(x_\ell), \hat{\theta}(x_1, x_2, x_3)) \leq \mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) + \epsilon\}} \geq 2.$$

Therefore, for all  $(x_1, x_2, x_3) \in E$ , there exists  $k \in \{1, 2, 3\}$  such that both  $L(\tilde{\theta}(x_k), \theta) < \mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) + \epsilon$  and  $L(\tilde{\theta}(x_k), \hat{\theta}(x_1, x_2, x_3)) \leq \mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) + \epsilon$ . Thus, for all  $(x_1, x_2, x_3) \in E$ ,

$$L(\hat{\theta}(x_1, x_2, x_3), \theta) \leq A\{L(\tilde{\theta}(x_k), \theta) + L(\tilde{\theta}(x_k), \hat{\theta}(x_1, x_2, x_3))\} \leq 2A\mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) + 2A\epsilon,$$

and we deduce that

$$P_\theta^{\otimes 3}\{L(\hat{\theta}, \theta) \leq 2A\mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) + 2A\epsilon\} \geq P_\theta^{\otimes 3}(E) \geq 1 - \delta^3 - 3\delta^2(1 - \delta).$$

This holds for all  $\theta \in \Theta$ , so

$$\mathcal{M}_-(\delta^3 + 3\delta^2(1 - \delta), \mathcal{P}_\Theta^{\otimes 3}, L) \leq 2A\mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) + 2A\epsilon,$$

and the result follows since  $\epsilon > 0$  was arbitrary.  $\square$

The following technical lemma on the existence of a measurable selector is based on the theory of correspondences, so we now present some basic definitions, following Aliprantis and Border (2007). A *correspondence*  $\varphi$  from a set  $\mathcal{Y}$  to a set  $\mathcal{Z}$ , written  $\varphi : \mathcal{Y} \rightrightarrows \mathcal{Z}$ , assigns to each  $y \in \mathcal{Y}$  a subset  $\varphi(y)$  of  $\mathcal{Z}$ ; in other words, it is a function from  $\mathcal{Y}$  to  $\text{Pow}(\mathcal{Z})$ . Its *lower inverse*  $\varphi^\text{L} : \text{Pow}(\mathcal{Z}) \rightarrow \text{Pow}(\mathcal{Y})$  is defined by

$$\varphi^\text{L}(B) := \{y \in \mathcal{Y} : \varphi(y) \cap B \neq \emptyset\}.$$

When  $\mathcal{Z}$  is a topological space, the *closure correspondence*  $\bar{\varphi} : \mathcal{Y} \rightrightarrows \mathcal{Z}$  of  $\varphi$  assigns to each  $y \in \mathcal{Y}$  the subset of  $\mathcal{Z}$  given by the closure of  $\varphi(y)$ . If, in addition,  $(\mathcal{Y}, \mathcal{B})$  is a measurable space, we say that the correspondence  $\varphi : \mathcal{Y} \rightrightarrows \mathcal{Z}$  is *measurable* if  $\varphi^\text{L}(F) \in \mathcal{B}$  for every closed subset  $F$  of  $\mathcal{Z}$ , and *weakly measurable* if  $\varphi^\text{L}(G) \in \mathcal{B}$  for every open subset  $G$  of  $\mathcal{Z}$ .

**Lemma 20.** *Let  $(\mathcal{X}, \mathcal{A})$  be a measurable space, let  $\Theta$  be a non-empty Polish space and suppose that  $L : \Theta \times \Theta \rightarrow [0, \infty)$  is a Carathéodory function. Let  $\tilde{\theta} : \mathcal{X} \rightarrow \Theta$  be measurable, let  $r \in (0, \infty)$  and define a correspondence  $\psi : \mathcal{X}^3 \rightrightarrows \Theta$  by*

$$\psi(x_1, x_2, x_3) := \left\{ \theta' \in \Theta : \sum_{\ell=1}^3 \mathbb{1}_{\{L(\tilde{\theta}(x_\ell), \theta') < r\}} \geq 2 \right\}.$$

Further, define a non-empty correspondence  $S : \mathcal{X}^3 \rightarrow \Theta$  by

$$S(x_1, x_2, x_3) := \begin{cases} \bar{\psi}(x_1, x_2, x_3) & \text{when } \psi(x_1, x_2, x_3) \neq \emptyset \\ \Theta & \text{otherwise.} \end{cases}$$

Then there exists a measurable function  $\hat{\theta} : \mathcal{X}^3 \rightarrow \Theta$  such that  $\hat{\theta}(x_1, x_2, x_3) \in S(x_1, x_2, x_3)$  for all  $(x_1, x_2, x_3) \in \mathcal{X}^3$ .

*Proof.* The map  $(x_1, x_2, x_3, \theta') \mapsto (L(\tilde{\theta}(x_1), \theta'), L(\tilde{\theta}(x_2), \theta'), L(\tilde{\theta}(x_3), \theta'))$  is a Carathéodory function from  $\mathcal{X}^3 \times \Theta$  to  $[0, \infty)^3$ . Define  $G := \{(v_1, v_2, v_3)^\top \in [0, \infty)^3 : \sum_{\ell=1}^3 \mathbb{1}_{\{v_\ell < r\}} \geq 2\}$ , which is an open subset of  $[0, \infty)^3$ . Then, by Aliprantis and Border (2007, Lemma 18.7), the correspondence  $\psi$ , which can be equivalently expressed as  $\psi(x_1, x_2, x_3) = \{\theta' \in \Theta : (L(\tilde{\theta}(x_\ell), \theta'))_{\ell=1}^3 \in G\}$ , is measurable, so by Aliprantis and Border (2007, Lemma 18.2) it is also weakly measurable. It follows that the lower inverse  $\psi^L : \text{Pow}(\Theta) \rightarrow \text{Pow}(\mathcal{X}^3)$  satisfies  $\psi^L(\Theta) = \{(x_1, x_2, x_3) \in \mathcal{X}^3 : \psi(x_1, x_2, x_3) \neq \emptyset\} \in \mathcal{A}^{\otimes 3}$ , so  $U := \psi^L(\Theta)^c \in \mathcal{A}^{\otimes 3}$ . Now define a non-empty correspondence  $\varphi : \mathcal{X}^3 \rightarrow \Theta$  by

$$\varphi(x_1, x_2, x_3) := \begin{cases} \psi(x_1, x_2, x_3) & \text{if } (x_1, x_2, x_3) \in U^c \\ \Theta & \text{if } (x_1, x_2, x_3) \in U. \end{cases}$$

Then, for any non-empty open set  $V \subseteq \Theta$ , we have  $\psi^L(V) \in \mathcal{A}^{\otimes 3}$ , so

$$\begin{aligned} \varphi^L(V) &= \{(x_1, x_2, x_3) \in \mathcal{X}^3 : \varphi(x_1, x_2, x_3) \cap V \neq \emptyset\} \\ &= \{(x_1, x_2, x_3) \in \mathcal{X}^3 : \psi(x_1, x_2, x_3) \cap V \neq \emptyset\} \cup U = \psi^L(V) \cup U \in \mathcal{A}^{\otimes 3}. \end{aligned}$$

Since we also have  $\varphi^L(\emptyset) = \emptyset \in \mathcal{A}^{\otimes 3}$ , we deduce that  $\varphi$  is weakly measurable, and by Aliprantis and Border (2007, Lemma 18.3), its closure correspondence  $\bar{\varphi}$  is also weakly measurable. Therefore, by Aliprantis and Border (2007, Theorem 18.13),  $\bar{\varphi}$  admits a measurable selector  $\hat{\theta} : \mathcal{X}^3 \rightarrow \Theta$ ; in other words,  $\hat{\theta}$  is measurable, with  $\hat{\theta}(x_1, x_2, x_3) \in \bar{\varphi}(x_1, x_2, x_3)$  for all  $(x_1, x_2, x_3) \in \mathcal{X}^3$ . Since  $\bar{\varphi} = S$ , the conclusion follows.  $\square$

We are now in a position to complete the proof of Proposition 9.

*Proof of Proposition 9.* The sequence  $x, h(x), h^{\circ 2}(x), h^{\circ 3}(x), \dots$  is strictly decreasing and converges to zero for  $x \in (0, 1/2)$ . It follows that there exists  $k \equiv k(\delta_-, \delta_+) \in \mathbb{N}$  such that  $h^{\circ k}(\delta_+) \leq \delta_-$ . Moreover, since  $h^{\circ k}(\cdot)$  is increasing on  $(0, 1/2)$ , we have  $h^{\circ k}(\delta) \leq h^{\circ k}(\delta_+) \leq \delta_-$  for all  $\delta \in (0, \delta_+]$ . Since  $d(\cdot, \theta)$  and  $d(\theta, \cdot)$  are continuous for all  $\theta \in \Theta$ , and since  $g$  is continuous, we have that  $L$  is a Carathéodory function. Hence, by iteratively applying Lemma 19  $k$  times, we deduce that

$$\mathcal{M}_-(\delta, \mathcal{P}_\Theta, L) \geq \frac{1}{(2A)^k} \cdot \mathcal{M}_-(h^{\circ k}(\delta), \mathcal{P}_\Theta^{\otimes 3k}, L) \geq \frac{1}{(2A)^k} \cdot \mathcal{M}_-(\delta_-, \mathcal{P}_\Theta^{\otimes 3k}, L),$$

as required.  $\square$

**Acknowledgement:** The research of all three authors was supported by the third author's European Research Council Advanced Grant 101019498.

## References

- Adler, R. J. and Taylor, J. E. (2009) *Random Fields and Geometry*. Springer.
- Aliprantis, C. D. and Border, K. C. (2007) *Infinite Dimensional Analysis: A Hitchhiker's Guide*. Springer.
- Bellec, P. C. (2018) Sharp oracle inequalities for least squares estimators in shape restricted regression. *The Annals of Statistics*, **46**, 745–780.
- Bellec, P. C., Lecué, G. and Tsybakov, A. B. (2018) Slope meets Lasso: improved oracle bounds and optimality. *The Annals of Statistics*, **46**, 3603–3642.
- Bogdan, M., Van Den Berg, E., Sabatti, C., Su, W. and Candès, E. J. (2015) SLOPE—Adaptive variable selection via convex optimization. *The Annals of Applied Statistics*, **9**, 1103.
- Bretagnolle, J. and Huber, C. (1979) Estimation des densités: Risque minimax. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, **47**, 119–137.
- Cai, T. T. (2012) Minimax and adaptive inference in nonparametric function estimation. *Statistical Science*, **27**, 31–50.
- Candès, E. J. and Davenport, M. A. (2013) How well can we estimate a sparse vector? *Applied and Computational Harmonic Analysis*, **34**, 317–323.
- Carmon, Y. and Hinder, O. (2024) The price of adaptivity in stochastic convex optimization. *arXiv preprint arXiv:2402.10898*.
- Catoni, O. (2012) Challenging the empirical mean and empirical variance: A deviation study. *Annales de l'Institut Henri Poincaré, Probabilités et Statistiques*, **48**, 1148–1185.
- Chatterjee, S., Guntuboyina, A. and Sen, B. (2015) On risk bounds in isotonic and other shape restricted regression problems. *The Annals of Statistics*, **43**, 1774–1800.
- Chen, M., Gao, C. and Ren, Z. (2016) A general decision theory for Huber's  $\epsilon$ -contamination model. *Electronic Journal of Statistics*, **10**, 3752–3774.
- Chen, M., Gao, C. and Ren, Z. (2018) Robust covariance and scatter matrix estimation under Huber's contamination model. *The Annals of Statistics*, **46**, 1932–1960.
- Chhor, J., Klopp, O. and Tsybakov, A. (2024) Generalized multi-view model: Adaptive density estimation under low-rank constraints. *arXiv preprint arXiv:2404.17209*.
- Davis, D., Drusvyatskiy, D., Xiao, L. and Zhang, J. (2021) From low probability to high confidence in stochastic convex optimization. *Journal of Machine Learning Research*, **22**, 1–38.
- Depersin, J. and Lecué, G. (2022a) Optimal robust mean and location estimation via convex programs with respect to any pseudo-norms. *Probability Theory and Related Fields*, **183**, 997–1025.

- Depersin, J. and Lecué, G. (2022b) Robust sub-Gaussian estimation of a mean vector in nearly linear time. *The Annals of Statistics*, **50**, 511–536.
- Devroye, L., Lerasle, M., Lugosi, G. and Oliveira, R. I. (2016) Sub-Gaussian mean estimators. *The Annals of Statistics*, **44**, 2695–2725.
- Devroye, L. and Lugosi, G. (2001) *Combinatorial Methods in Density Estimation*. Springer.
- Donoho, D. L. (1994) Statistical estimation and optimal recovery. *The Annals of Statistics*, **22**, 238–270.
- Duchi, J. C. (2018) Introductory Lectures on Stochastic Optimization. In *The Mathematics of Data*, 99–185, American Mathematical Society.
- Duchi, J. C. and Haque, S. (2024) An information-theoretic lower bound in time-uniform estimation. *arXiv preprint arXiv:2402.08794*.
- Gerchinovitz, S., Ménard, P. and Stoltz, G. (2020) Fano’s inequality for random variables. *Statistical Science*, **35**, 178–201.
- Groeneboom, P. and Jongbloed, G. (2014) *Nonparametric Estimation under Shape Constraints*. Cambridge University Press.
- Hopkins, S. B. (2020) Mean estimation with sub-Gaussian rates in polynomial time. *The Annals of Statistics*, **48**, 1193–1213.
- Huber, P. J. (1964) Robust estimation of a location parameter. *The Annals of Mathematical Statistics*, **35**, 73–101.
- Koltchinskii, V. and Lounici, K. (2017) Concentration inequalities and moment bounds for sample covariance operators. *Bernoulli*, **23**, 110–133.
- Lecué, G. and Mendelson, S. (2013) Learning sub-Gaussian classes: Upper and minimax bounds. In *Topics in Learning Theory*, Société Mathématique de France.
- Lugosi, G. and Mendelson, S. (2019a) Mean estimation and regression under heavy-tailed distributions: A survey. *Foundations of Computational Mathematics*, **19**, 1145–1190.
- Lugosi, G. and Mendelson, S. (2019b) Sub-Gaussian estimators of the mean of a random vector. *The Annals of Statistics*, **47**, 783–794.
- Lugosi, G. and Mendelson, S. (2021) Robust multivariate mean estimation: The optimality of trimmed mean. *The Annals of Statistics*, **49**, 393–410.
- Massart, P. (2007) *Concentration Inequalities and Model Selection*. Springer.
- Nemirovski, A. and Yudin, D. (1983) *Problem Complexity and Method Efficiency in Optimization*. Wiley-Interscience.
- Robbins, H. (1955) A remark on Stirling’s formula. *The American Mathematical Monthly*, **62**, 26–29.

- Rockafellar, R. T. (1997) *Convex Analysis*. Princeton University Press.
- Samworth, R. J. and Sen, B. (2018) Editorial: Special issue on “Nonparametric inference under shape constraints”. *Statistical Science*, **33**, 469–472.
- Samworth, R. J. and Shah, R. D. (2024+) *Modern Statistical Methods and Theory*. Cambridge University Press, to appear.
- Tropp, J. A. (2012) User-friendly tail bounds for sums of random matrices. *Foundations of computational mathematics*, **12**, 389–434.
- Tsybakov, A. B. (2009) *Introduction to Nonparametric Estimation*. Springer.
- Tukey, J. W. (1975) Mathematics and the picturing of data. In *Proceedings of the International Congress of Mathematicians*, vol. 2, 523–531.
- Vershynin, R. (2018) *High-Dimensional Probability: An Introduction with Applications in Data Science*. Cambridge University Press.
- Wainwright, M. J. (2019) *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge University Press.
- Wang, H. and Ramdas, A. (2024) Anytime-valid  $t$ -tests and confidence sequences for Gaussian means with unknown variance. *arXiv preprint arXiv:2310.03722*.
- Zhivotovskiy, N. (2024) Dimension-free bounds for sums of independent matrices and simple tensors via the variational principle. *Electronic Journal of Probability*, **29**, 1–28.
- Zhu, Z., Wang, T. and Samworth, R. J. (2022) High-dimensional principal component analysis with heterogeneous missingness. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, **84**, 2000–2031.

## A Minimax risk lower bounds

In this section, we state some general techniques for minimax risk lower bounds from [Samworth and Shah \(2024+\)](#); similar statements can be found in the books by [Tsybakov \(2009\)](#) and [Wainwright \(2019\)](#). We work in the setting of [Section 2](#).

**Lemma 21** (Le Cam’s two-point lemma). *Let  $\theta_1, \theta_2 \in \Theta$ ,  $P_1 \in \mathcal{P}_{\theta_1}$  and  $P_2 \in \mathcal{P}_{\theta_2}$ . Then, writing  $\eta := \frac{1}{2}d(\theta_1, \theta_2)$  and  $\Theta_0 := \{\theta_1, \theta_2\}$ , we have*

$$\inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} \mathbb{E}_{P_\theta} L(\hat{\theta}, \theta) \geq \inf_{\hat{\theta} \in \hat{\Theta}} \max_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} \mathbb{E}_{P_{\theta_0}} L(\hat{\theta}, \theta_0) \geq \frac{g(\eta)}{2} \{1 - \text{TV}(P_1, P_2)\}.$$

**Lemma 22** (Fano’s lemma). *Let  $M \geq 2$ ,  $\theta_1, \dots, \theta_M \in \Theta$ , and  $P_j \in \mathcal{P}_{\theta_j}$  for  $j \in [M]$ . Then, writing  $\eta := \frac{1}{2} \min_{1 \leq j < k \leq M} d(\theta_j, \theta_k)$  and  $\Theta_0 := \{\theta_j : j \in [M]\}$ , we have*

$$\begin{aligned} \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} \mathbb{E}_{P_\theta} L(\hat{\theta}, \theta) &\geq \inf_{\hat{\theta} \in \hat{\Theta}} \max_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} \mathbb{E}_{P_{\theta_0}} L(\hat{\theta}, \theta_0) \\ &\geq g(\eta) \left\{ 1 - \frac{M^{-1} \inf_{Q \in \mathcal{Q}(\mathcal{X})} \sum_{j=1}^M \text{KL}(P_j, Q) + \log(2 - M^{-1})}{\log M} \right\}. \end{aligned}$$

**Lemma 23** (Assouad's lemma). *Let  $m \in \mathbb{N}$ ,  $\Phi := \{0, 1\}^m$ . For  $\phi \in \Phi$ , let  $\theta_\phi \in \Theta$  and  $P_\phi \in \mathcal{P}_{\theta_\phi}$ . For  $\phi, \phi' \in \Phi$ , write  $\phi \sim \phi'$  whenever  $\phi$  and  $\phi'$  differ in precisely one coordinate, and  $\phi \sim_j \phi'$  when that coordinate is the  $j$ th. Suppose now that the loss function is of the form*

$$L(\theta_1, \theta_2) := \sum_{j=1}^m g(d_j(\theta_1, \theta_2)),$$

for  $\theta_1, \theta_2 \in \Theta$ , where  $d_1, \dots, d_m$  are pseudo-metrics on  $\Theta$  with  $d_j(\theta_\phi, \theta_{\phi'}) \geq \alpha_j$  whenever  $\phi \sim_j \phi'$ , and where  $g$  is an increasing function satisfying  $g(x + y) \leq A\{g(x) + g(y)\}$  for all  $x, y \in [0, \infty)$  and some  $A > 0$ . Then, writing  $\Theta_0 := \{\theta_\phi : \phi \in \Phi\}$ , we have

$$\begin{aligned} \inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P_\theta \in \mathcal{P}_\theta} \mathbb{E}_{P_\theta} L(\hat{\theta}, \theta) &\geq \inf_{\hat{\theta} \in \hat{\Theta}} \max_{\theta_0 \in \Theta_0} \sup_{P_{\theta_0} \in \mathcal{P}_{\theta_0}} \mathbb{E}_{P_{\theta_0}} L(\hat{\theta}, \theta_0) \\ &\geq \frac{1}{2A} \left\{ 1 - \max_{\phi, \phi' \in \Phi: \phi \sim \phi'} \text{TV}(P_\phi, P_{\phi'}) \right\} \sum_{j=1}^d g(\alpha_j). \end{aligned}$$

## B Auxiliary lemmas

Our first auxiliary lemma is a minor modification of [Chen, Gao and Ren \(2018, Theorem 5.1\)](#) that we apply in the proof of [Corollary 11](#). To introduce the setting, let  $\mathcal{Y}$  be a measurable space; for  $\theta \in \Theta$ , let  $\mathcal{R}_\theta \subseteq \mathcal{Q}(\mathcal{Y})$  and let  $\mathcal{R}_\Theta := \{\mathcal{R}_\theta : \theta \in \Theta\}$ .

**Definition 24.** *Let  $\varepsilon \in [0, 1)$ . We define the modulus of continuity of  $\mathcal{R}_\Theta$  with respect to  $\varepsilon$  and  $g$  to be*

$$\omega(\varepsilon, \mathcal{R}_\Theta, g) := \sup \left\{ g\left(\frac{d(\theta_1, \theta_2)}{2}\right) : \theta_1, \theta_2 \in \Theta \text{ and there exist } R_{\theta_1} \in \mathcal{R}_{\theta_1}, \right. \\ \left. R_{\theta_2} \in \mathcal{R}_{\theta_2} \text{ such that } \text{TV}(R_{\theta_1}, R_{\theta_2}) \leq \frac{\varepsilon}{1 - \varepsilon} \right\}.$$

**Lemma 25.** *For  $n \in \mathbb{N}$ ,  $\theta \in \Theta$  and  $\varepsilon \in [0, 1)$ , let  $\mathcal{X} = \mathcal{Y}^n$  and*

$$\mathcal{P}_\theta \equiv \mathcal{P}_{\theta, \varepsilon} := \{P^{\otimes n} \in \mathcal{Q}(\mathcal{X}) : P = (1 - \varepsilon)R_\theta + \varepsilon Q : R_\theta \in \mathcal{R}_\theta \text{ and } Q \in \mathcal{Q}(\mathcal{Y})\}$$

denote the class of contaminated distributions. Then, for all  $\delta \in (0, 1/2)$ ,

$$\mathcal{M}_-(\delta) \geq \omega(\varepsilon, \mathcal{R}_\Theta, g).$$

*Proof.* We apply the same proof strategy as [Chen, Gao and Ren \(2018, Theorem 5.1\)](#). Consider any  $R_1 \in \mathcal{R}_{\theta_1}$  and  $R_2 \in \mathcal{R}_{\theta_2}$  that satisfy  $\text{TV}(R_1, R_2) \leq \varepsilon/(1 - \varepsilon)$ . Then there exists  $\varepsilon' \in [0, \varepsilon]$  such that  $\text{TV}(R_1, R_2) = \varepsilon'/(1 - \varepsilon')$ . Next, for  $\ell \in \{1, 2\}$ , define densities

$$r_\ell := \frac{dR_\ell}{dR_1 + dR_2}.$$

We also define probability distributions  $Q_1$  and  $Q_2$  on  $\mathcal{X}$  through their densities

$$\frac{dQ_1}{dR_1 + dR_2} := \frac{(r_2 - r_1)\mathbb{1}_{\{r_2 \geq r_1\}}}{\text{TV}(R_1, R_2)} \quad \text{and} \quad \frac{dQ_2}{dR_1 + dR_2} := \frac{(r_1 - r_2)\mathbb{1}_{\{r_1 \geq r_2\}}}{\text{TV}(R_1, R_2)}.$$

Now define  $P_\ell := (1 - \varepsilon')R_\ell + \varepsilon'Q_\ell$  for  $\ell \in \{1, 2\}$ . Then  $P_\ell \in \mathcal{P}_{\theta_\ell}$  for  $\ell \in \{1, 2\}$ . Moreover, direct calculation yields

$$\begin{aligned} \frac{dP_1}{dR_1 + dR_2} &= (1 - \varepsilon')r_1 + \varepsilon' \frac{(r_2 - r_1)\mathbb{1}_{\{r_2 \geq r_1\}}}{\varepsilon'/(1 - \varepsilon')} = (1 - \varepsilon')\{r_1 + (r_2 - r_1)\mathbb{1}_{\{r_2 \geq r_1\}}\} \\ &= (1 - \varepsilon')\{r_2 + (r_1 - r_2)\mathbb{1}_{\{r_1 \geq r_2\}}\} = \frac{dP_2}{dR_1 + dR_2}, \end{aligned}$$

so  $P_1 = P_2$ . Thus, by Lemma 5,

$$\inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P \in \mathcal{P}_\theta} P \left\{ L(\hat{\theta}, \theta) \geq g \left( \frac{d(\theta_1, \theta_2)}{2} \right) \right\} \geq \frac{1}{2} \{1 - \text{TV}(P_1^{\otimes n}, P_2^{\otimes n})\} = \frac{1}{2},$$

where the inequality follows since  $P_1^{\otimes n}, P_2^{\otimes n} \in \mathcal{P}_\Theta$ . Further, since  $R_1, R_2 \in \mathcal{R}_\Theta$  with  $\text{TV}(R_1, R_2) \leq \varepsilon/(1 - \varepsilon)$  were arbitrary, for any  $t > 0$ , we can find  $\theta_1^{(t)}, \theta_2^{(t)} \in \Theta$ ,  $R_1^{(t)} \in \mathcal{R}_{\theta_1^{(t)}}$  and  $R_2^{(t)} \in \mathcal{R}_{\theta_2^{(t)}}$  such that  $\text{TV}(R_1^{(t)}, R_2^{(t)}) \leq \varepsilon/(1 - \varepsilon)$  and  $g(d(\theta_1^{(t)}, \theta_2^{(t)})/2) \geq \omega(\varepsilon, \mathcal{R}_\Theta, g) - t$ . Hence, by taking  $t \rightarrow 0$ , we deduce that

$$\inf_{\hat{\theta} \in \hat{\Theta}} \sup_{\theta \in \Theta} \sup_{P \in \mathcal{P}_\theta} P \left\{ L(\hat{\theta}, \theta) \geq \omega(\varepsilon, \mathcal{R}_\Theta, g) \right\} \geq \frac{1}{2},$$

so the result follows.  $\square$

Lemma 26 below applies a non-central version of the matrix Bernstein inequality (Tropp, 2012; Zhu, Wang and Samworth, 2022) to provide a high-probability upper bound on the operator norm of the difference between a sample covariance matrix and its population counterpart.

**Lemma 26.** *Let  $n, d \in \mathbb{N}$ ,  $\Sigma \in \mathcal{S}_+^{d \times d}$  and  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \mathbf{N}_d(0, \Sigma)$ . Then*

$$\left\| \frac{1}{n} \sum_{i=1}^n \mathbb{E}\{(X_i X_i^\top - \Sigma)^2\} \right\|_{\text{op}} \asymp \left\| \frac{1}{n} \sum_{i=1}^n \mathbb{E}\{(X_i X_i^\top)^2\} \right\|_{\text{op}} \asymp \|\Sigma\|_{\text{op}}^2 \mathbf{r}(\Sigma). \quad (28)$$

Moreover, writing  $\hat{\Sigma} := n^{-1} \sum_{i=1}^n X_i X_i^\top$ , if  $\frac{\mathbf{r}(\Sigma)\{\log(1/\delta) + \log(8d)\}}{n} \leq 1$ , then with probability at least  $1 - \delta$ , we have

$$\|\hat{\Sigma} - \Sigma\|_{\text{op}} \leq 513 \|\Sigma\|_{\text{op}} \sqrt{\frac{\mathbf{r}(\Sigma)\{\log(1/\delta) + \log(8d)\}}{n}}. \quad (29)$$

**Remark.** *From the bounds on the variance parameter in (28), we see that the upper bound in (29) is the tightest (up to multiplicative constants) that one can obtain from the matrix Bernstein inequality in the sub-Gaussian concentration regime (i.e. when  $\frac{\mathbf{r}(\Sigma)\{\log(1/\delta) + \log(8d)\}}{n} \leq 1$ ). In particular, the  $\log(1/\delta)$  term in (29) appears multiplicatively with the effective rank, in stark contrast with Proposition 13, where the  $\log(1/\delta)$  term appears only additively.*

*Proof.* Since the operator norm is invariant to orthogonal transformations, we may assume without loss of generality that  $\Sigma$  is diagonal, so that  $\Sigma = \text{diag}(\lambda_1, \dots, \lambda_d)$  where  $\lambda_1 \geq \dots \geq \lambda_d \geq 0$ . Then, writing  $X_1 = (X_{11}, \dots, X_{1d}) \in \mathbb{R}^d$ , we have

$$\mathbb{E}\{(X_1 X_1^\top - \Sigma)^2\} = \mathbb{E}\{(X_1 X_1^\top)^2\} - \Sigma^2$$



$$\begin{aligned}
&= \mathbb{E}(\|X_1\|_2^2 X_1 X_1^\top) - \Sigma^2 \\
&= \text{diag}(\mathbb{E}(\|X_1\|_2^2 X_{11}^2), \dots, \mathbb{E}(\|X_1\|_2^2 X_{1d}^2)) - \text{diag}(\lambda_1^2, \dots, \lambda_d^2) \\
&= \text{diag}\left(2\lambda_1^2 + \lambda_1 \sum_{j \neq 1} \lambda_j, \dots, 2\lambda_d^2 + \lambda_d \sum_{j \neq d} \lambda_j\right).
\end{aligned}$$

Therefore,

$$\|\Sigma\|_{\text{op}} \text{tr}(\Sigma) \leq \left\| \frac{1}{n} \sum_{i=1}^n \mathbb{E}\{(X_i X_i^\top - \Sigma)^2\} \right\|_{\text{op}} \leq 2\|\Sigma\|_{\text{op}} \text{tr}(\Sigma).$$

Similarly,

$$\|\Sigma\|_{\text{op}} \text{tr}(\Sigma) \leq \left\| \frac{1}{n} \sum_{i=1}^n \mathbb{E}\{(X_i X_i^\top)^2\} \right\|_{\text{op}} \leq 3\|\Sigma\|_{\text{op}} \text{tr}(\Sigma),$$

and (28) follows.

For the second claim, observe that for  $p \geq 2$ , we have

$$\begin{aligned}
\mathbb{E}\{(X_1 X_1^\top)^p\} &= \mathbb{E}(\|X_1\|_2^{2p-2} X_1 X_1^\top) \\
&= \text{diag}(\mathbb{E}(\|X_1\|_2^{2p-2} X_{11}^2), \dots, \mathbb{E}(\|X_1\|_2^{2p-2} X_{1d}^2)) \\
&\preceq \{\mathbb{E}(\|X_1\|_2^{4p-4})\}^{1/2} \text{diag}(\{\mathbb{E}(X_{11}^4)\}^{1/2}, \dots, \{\mathbb{E}(X_{1d}^4)\}^{1/2}) \\
&\stackrel{(i)}{\preceq} \left(\frac{8}{3}\right)^{p-1} \cdot (2p-2)^{p-1} \text{tr}(\Sigma)^{p-1} \cdot 3^{1/2} \Sigma \\
&\stackrel{(ii)}{\preceq} \left(\frac{16e \text{tr}(\Sigma)}{3}\right)^{p-1} \cdot \frac{1}{p\sqrt{p-1}} \cdot p! \cdot \Sigma \\
&\preceq \frac{p!}{2} \cdot (16 \text{tr}(\Sigma))^{p-2} \cdot 16 \text{tr}(\Sigma) \cdot \Sigma.
\end{aligned}$$

Here, Step (i) follows by recalling the sub-exponential norm of a random variable  $Z$ , given by  $\|Z\|_{\psi_1} := \inf\{t > 0 : \mathbb{E}(e^{|Z|/t}) \leq 2\}$ , so that  $\|\|X_1\|_2^2\|_{\psi_1} \leq \sum_{j=1}^d \|X_{1j}^2\|_{\psi_1} = 8 \text{tr}(\Sigma)/3$ . Moreover,  $(y/r)^r \leq e^y$  for all  $y, r \geq 0$ , so

$$\mathbb{E}\left\{\left(\frac{\|X_1\|_2^2}{(2p-2) \cdot 8 \text{tr}(\Sigma)/3}\right)^{2p-2}\right\} \leq \mathbb{E} \exp\left(\frac{\|X_1\|_2^2}{8 \text{tr}(\Sigma)/3}\right) \leq 2.$$

Step (ii) follows from Stirling's formula (e.g., [Robbins, 1955](#), Eq. (1)). Noting that  $X_1 X_1^\top$  is positive semi-definite, we may apply a non-central version of the matrix Bernstein inequality ([Zhu, Wang and Samworth, 2022](#), Lemma 3) to both  $\widehat{\Sigma}$  and  $-\widehat{\Sigma}$  to deduce that, for all  $t > 0$ ,

$$\mathbb{P}(\|\widehat{\Sigma} - \Sigma\|_{\text{op}} \geq t) \leq 8d \exp\left\{-\frac{nt^2}{2^9(\|\Sigma\|_{\text{op}} + t) \text{tr}(\Sigma)}\right\}.$$

The result then follows on setting  $t = (2^9 + 1)\|\Sigma\|_{\text{op}} \sqrt{\frac{\mathbf{r}(\Sigma)\{\log(1/\delta) + \log(8d)\}}{n}}$  and invoking the assumption  $\frac{\mathbf{r}(\Sigma)\{\log(1/\delta) + \log(8d)\}}{n} \leq 1$ .  $\square$

The following lemma provides a lower bound on the packing number of the set of  $s$ -sparse vectors in  $\mathbb{S}^{d-1}$ . Very similar results exist in the literature (e.g. [Candès and Davenport \(2013, Lemma 3\)](#) or [Wainwright \(2019, Exercise 5.8\)](#)); we present a short, constructive proof without imposing a parity restriction on the sparsity level  $s$ .

**Lemma 27.** *Fix  $d \in \mathbb{N}$  and  $s \in [d]$ , and let  $\mathcal{V} := \{v \in \mathbb{S}^{d-1} : \|v\|_0 \leq s\}$ . There exist  $\log M \geq \frac{3s}{4} \log\left(\frac{d}{4s}\right)$  and  $v_1, \dots, v_M \in \mathcal{V}$  such that  $\|v_j - v_k\|_2 > 1/2$  for all distinct  $j, k \in [M]$ .*

*Proof.* Let  $L := \lfloor s/4 \rfloor$  and let  $\Omega_1 := \{\omega \in \{0, 1\}^d : \|\omega\|_0 \leq s\}$ . For each  $j \in \mathbb{N}$ , if  $\Omega_j \neq \emptyset$ , fix an arbitrary  $\omega_j \in \Omega_j$ , let

$$\Omega_{j+1} := \{\omega \in \Omega_j : d_{\text{H}}(\omega, \omega_j) > s/4\},$$

and let  $M := \max\{j \in \mathbb{N} : \Omega_j \neq \emptyset\}$ . Then  $|\Omega_j \setminus \Omega_{j+1}| \leq \sum_{\ell=0}^L \binom{d}{\ell}$  for each  $j \in [M]$ . Since the sets  $\{\Omega_j \setminus \Omega_{j+1} : j \in [M]\}$  form a partition of  $\Omega_1$ , we deduce that

$$\sum_{\ell=0}^s \binom{d}{\ell} = \sum_{j=1}^M |\Omega_j \setminus \Omega_{j+1}| \leq M \sum_{\ell=0}^L \binom{d}{\ell}.$$

Hence

$$M \geq \frac{\sum_{\ell=0}^s \binom{d}{\ell}}{\sum_{\ell=0}^L \binom{d}{\ell}} \geq \frac{\binom{d}{s}}{(L+1)\binom{d}{L}} \geq \frac{(d/s)^s}{(L+1)(ed/L)^L}.$$

It follows that

$$\begin{aligned} \log M &\geq s \log\left(\frac{d}{s}\right) - L \log\left(\frac{ed}{L}\right) - \log(L+1) \geq s \log\left(\frac{d}{s}\right) - \frac{s}{4} \log\left(\frac{4ed}{s}\right) - \log\left(\frac{s}{4} + 1\right) \\ &\geq \frac{3s}{4} \log\left(\frac{d}{s}\right) - \frac{s}{4} \log(4e^2) = \frac{3s}{4} \log\left(\frac{d}{(4e^2)^{1/3}s}\right) \geq \frac{3s}{4} \log\left(\frac{d}{4s}\right). \end{aligned}$$

The points  $\omega_1, \dots, \omega_M$  satisfy  $d_{\text{H}}(\omega_j, \omega_k) > s/4$  for all distinct  $j, k \in [M]$ , and the result follows by taking,  $v_j = \omega_j / \sqrt{s}$  for  $j \in [M]$ , so that  $v_j \in \mathcal{V}$  for all  $j$ .  $\square$

Recall that in the context of kernel density estimation, a *kernel* is a Borel measurable function  $K : \mathbb{R} \rightarrow \mathbb{R}$  with  $\int_{-\infty}^{\infty} K(u) du = 1$ . It is of *order*  $\ell \in \mathbb{N}$  if  $\int_{-\infty}^{\infty} u^j K(u) du = 0$  for all  $j \in [\ell - 1]$ .

**Lemma 28.** *Let  $\beta, \gamma > 0$ , let  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} f \in \mathcal{F}(\beta, \gamma)$ , let  $x_0 \in \mathbb{R}$  and let  $K : \mathbb{R} \rightarrow \mathbb{R}$  be a kernel of order  $\ell := \lceil \beta \rceil$  satisfying  $\|K\|_{\infty} < \infty$  and*

$$\mu_{\beta}(K) := \int_{-\infty}^{\infty} |u|^{\beta} |K(u)| du < \infty.$$

*There exist  $c(\beta, K), C(\beta, K) > 0$ , depending only on  $\beta, K$ , and a choice of bandwidth  $h \equiv h_{n, \beta, \gamma, K, \delta}$  such that if  $\delta \in [2e^{-c(\beta, K)n}, 1]$ , then with probability at least  $1 - \delta$ ,*

$$\{\widehat{f}_n(x_0) - f(x_0)\}^2 \leq C(\beta, K) \cdot \gamma^{2/(\beta+1)} \left(\frac{\log(2/\delta)}{n}\right)^{2\beta/(2\beta+1)},$$

*where  $\widehat{f}_n \equiv \widehat{f}_{n, h, K}$  is the kernel density estimator with bandwidth  $h$  and kernel  $K$ .*

*Proof.* By [Tsybakov \(2009, Proposition 1.2\)](#), we have

$$|\mathbb{E}\widehat{f}_n(x_0) - f(x_0)| \leq \frac{\gamma}{(\ell - 1)!} \mu_\beta(K) h^\beta. \quad (30)$$

Moreover, by [Tsybakov \(2009, Proposition 1.1\)](#) and the first display in the proof of [Samworth and Shah \(2024+, Theorem 4.6\)](#), we have

$$\text{Var}(K_h(X_1 - x_0)) \leq \frac{1}{h} \|f\|_\infty R(K) \leq \frac{1}{h} C_1(\beta) \gamma^{1/(\beta+1)} R(K),$$

where  $R(K) := \int_{-\infty}^{\infty} K^2(u) du$  and where  $C_1(\beta) > 0$  depends only on  $\beta$ . Now by Bernstein's inequality, we have for any  $\delta \in (0, 1]$  that

$$\mathbb{P}\left\{ |\widehat{f}_n(x_0) - \mathbb{E}\widehat{f}_n(x_0)| > \left( \frac{2C_1(\beta) \gamma^{1/(\beta+1)} R(K) \log(2/\delta)}{nh} \right)^{1/2} + \frac{\|K\|_\infty \log(2/\delta)}{3nh} \right\} \leq \delta.$$

Hence, when

$$\log(2/\delta) \leq nh \left( \frac{18C_1(\beta) \gamma^{1/(\beta+1)} R(K)}{\|K\|_\infty^2} \right), \quad (31)$$

we have with probability at least  $1 - \delta$  that

$$|\widehat{f}_n(x_0) - \mathbb{E}\widehat{f}_n(x_0)| \leq 2 \sqrt{\frac{2C_1(\beta) \gamma^{1/(\beta+1)} R(K) \log(2/\delta)}{nh}}. \quad (32)$$

Combining (32) with (30), we deduce that when  $\delta$  satisfies (31), we have with probability at least  $1 - \delta$  that

$$\begin{aligned} |\widehat{f}_n(x_0) - f(x_0)| &\leq |\widehat{f}_n(x_0) - \mathbb{E}\widehat{f}_n(x_0)| + |\mathbb{E}\widehat{f}_n(x_0) - f(x_0)| \\ &\leq 2 \sqrt{\frac{2C_1(\beta) \gamma^{1/(\beta+1)} R(K) \log(2/\delta)}{nh}} + \frac{\gamma}{(\ell - 1)!} \mu_\beta(K) h^\beta. \end{aligned}$$

Now set

$$C_2(\beta, K) := \left( \frac{C_1(\beta) R(K) \{(\ell - 1)!\}^2}{\mu_\beta^2(K)} \right)^{1/(2\beta+1)},$$

and

$$h \equiv h_{n,\beta,\gamma,K,\delta} := C_2(\beta, K) \cdot \gamma^{-1/(\beta+1)} \left( \frac{\log(2/\delta)}{n} \right)^{1/(2\beta+1)}.$$

Then there exist  $c(\beta, K), C(\beta, K) > 0$  such that

$$\{\widehat{f}_n(x_0) - f(x_0)\}^2 \leq C(\beta, K) \cdot \gamma^{2/(\beta+1)} \left( \frac{\log(2/\delta)}{n} \right)^{2\beta/(2\beta+1)},$$

with probability at least  $1 - \delta$ , whenever  $\delta \geq 2e^{-c(\beta,K) \cdot n}$ . □

The bandwidth in Lemma 28 depends on  $\delta$ , which is undesirable. The following lemma uses a Lepski type method to obtain a  $\delta$ -independent estimator for  $f(x_0)$ . The same method was employed by Devroye et al. (2016, Theorem 4.2) for heavy-tailed mean estimators.

**Lemma 29.** *Let  $\beta, \gamma > 0$ , let  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} f \in \mathcal{F}(\beta, \gamma)$ , let  $x_0 \in \mathbb{R}$  and let  $K : \mathbb{R} \rightarrow \mathbb{R}$  be a kernel of order  $\ell := \lceil \beta \rceil$  satisfying  $\|K\|_\infty < \infty$  and*

$$\mu_\beta(K) := \int_{-\infty}^{\infty} |u|^\beta |K(u)| du < \infty.$$

*There exists a  $\delta$ -independent estimator  $\check{f}(x_0) \equiv \check{f}_{n,\beta,\gamma,K}(x_0)$  and  $\check{C}(\beta, K) > 0$  depending only on  $\beta, K$  such that for all  $\delta \in (0, 1]$ , we have with probability at least  $1 - \delta$  that*

$$\{\check{f}_n(x_0) - f(x_0)\}^2 \leq \check{C}(\beta, K) \cdot \gamma^{2/(\beta+1)} \cdot \left\{ \left( \frac{\log(8/\delta)}{n} \right)^{2\beta/(2\beta+1)} \wedge 1 \right\}.$$

*Proof.* Let  $c(\beta, K) > 0$  be as in Lemma 28. First consider the case  $n \leq 3 \log(2)/c(\beta, K)$ . We set  $\tilde{f}_n(x_0) = 0$ , and recall from the first display in the proof of Samworth and Shah (2024+, Theorem 4.6) that  $\|f\|_\infty \leq C_1(\beta) \gamma^{1/(\beta+1)}$  for some  $C_1(\beta) > 0$  depending only on  $\beta$ . Therefore, when  $n \leq 3 \log(2)/c(\beta, K)$ , we have

$$\{\tilde{f}_n(x_0) - f(x_0)\}^2 \leq C_1(\beta)^2 \gamma^{2/(\beta+1)} \leq C_2(\beta, K) \cdot \gamma^{2/(\beta+1)} \cdot \left\{ \left( \frac{\log(8/\delta)}{n} \right)^{2\beta/(2\beta+1)} \wedge 1 \right\}, \quad (33)$$

for some  $C_2(\beta, K) > 0$  depending only on  $\beta$  and  $K$ . This proves the claim for  $n \leq 3 \log(2)/c(\beta, K)$ .

Next consider the case where  $n > 3 \log(2)/c(\beta, K)$ . Let  $\hat{f}_{n,\delta}(x_0)$  denote the kernel density estimator of  $f(x_0)$  with bandwidth  $h_{n,\beta,\gamma,K,\delta}$  as in Lemma 28. Define  $k_+ := \lfloor c(\beta, K) \cdot n / \log(2) - 1 \rfloor \in \mathbb{N}$ , which by assumption, satisfies  $k_+ \geq c(\beta, K) \cdot n / (2 \log 2)$ . For  $k \in [k_+]$ , define the interval

$$\hat{I}_k := \left[ \hat{f}_{n,2^{-k}}(x_0) - C(\beta, K)^{1/2} \cdot \gamma^{1/(\beta+1)} \left( \frac{\log(2^{k+1})}{n} \right)^{\beta/(2\beta+1)}, \right. \\ \left. \hat{f}_{n,2^{-k}}(x_0) + C(\beta, K)^{1/2} \cdot \gamma^{1/(\beta+1)} \left( \frac{\log(2^{k+1})}{n} \right)^{\beta/(2\beta+1)} \right].$$

By Lemma 28, we have for each  $k \in [k_+]$  that  $f(x_0) \in \hat{I}_k$  with probability at least  $1 - 2^{-k}$ . Now define  $\hat{k} := \min\{k \in [k_+] : \bigcap_{j=k}^{k_+} \hat{I}_j \neq \emptyset\}$ . Define the estimator

$$\tilde{f}_n(x_0) := \begin{cases} \inf\{\bigcap_{j=\hat{k}}^{k_+} \hat{I}_j\} & \text{if } \bigcap_{j=\hat{k}}^{k_+} \hat{I}_j \neq \emptyset \\ 0 & \text{otherwise,} \end{cases}$$

and set  $\check{f}_n(x_0) := 0 \vee \tilde{f}_n(x_0) \wedge C_1(\beta) \gamma^{1/(\beta+1)}$ . For  $\delta \in [2^{-(k_+-1)}, 1]$ , we set  $k_\delta := \min\{k \in [k_+] : \delta \geq 2^{-(k-1)}\}$ , so that  $2^{k_\delta-2} < 1/\delta \leq 2^{k_\delta-1}$ . Then, by a union bound,

$$\mathbb{P}\left(f(x_0) \in \bigcap_{j=k_\delta}^{k_+} \hat{I}_j\right) \geq 1 - \sum_{j=k_\delta}^{k_+} 2^{-j} \geq 1 - 2^{-(k_\delta-1)} \geq 1 - \delta.$$

On this event,  $\bigcap_{j=k_\delta}^{k_+} \widehat{I}_j \neq \emptyset$ , so  $\widehat{k} \leq k_\delta$ . Hence, for all  $\delta \in [2^{-(k_+-1)}, 1]$ , we have with probability at least  $1 - \delta$  that  $\widetilde{f}_n(x_0) \in \widehat{I}_{k_\delta}$ , so that with probability at least  $1 - \delta$ ,

$$\begin{aligned} \{\widetilde{f}_n(x_0) - f(x_0)\}^2 &\leq 4C(\beta, K) \cdot \gamma^{2/(\beta+1)} \left( \frac{\log(2^{k_\delta+1})}{n} \right)^{2\beta/(2\beta+1)} \\ &\leq 4C(\beta, K) \cdot \gamma^{2/(\beta+1)} \left( \frac{\log(8/\delta)}{n} \right)^{2\beta/(2\beta+1)}. \end{aligned}$$

Since  $f(x_0) \in [0, C_1(\beta)\gamma^{1/(\beta+1)}]$ , it follows that when  $\delta \in [2^{-(k_+-1)}, 1]$ , we have with probability at least  $1 - \delta$  that

$$\{\check{f}_n(x_0) - f(x_0)\}^2 \leq \{\widetilde{f}_n(x_0) - f(x_0)\}^2 \leq 4C(\beta, K) \cdot \gamma^{2/(\beta+1)} \left( \frac{\log(8/\delta)}{n} \right)^{2\beta/(2\beta+1)}. \quad (34)$$

On the other hand, for  $\delta \in (0, 2^{-(k_+-1)})$ ,

$$\{\check{f}_n(x_0) - f(x_0)\}^2 \leq \|f\|_\infty^2 \leq C_1^2(\beta) \cdot \gamma^{2/(\beta+1)}. \quad (35)$$

Note that when  $\delta = 2^{-(k_+-1)}$  and  $n > 3 \log(2)/c(\beta, K)$ , we have that

$$\frac{\log(8/\delta)}{n} = \frac{1}{n} \log(2^{k_++2}) \in \left[ \frac{c(\beta, K)}{2}, \frac{4c(\beta, K)}{3} \right].$$

Thus, there exists  $C_3(\beta, K) \geq 1$  depending only on  $\beta, K$  such that when  $\delta \in (0, 2^{-(k_+-1)})$ , we have  $1 \leq C_3(\beta, K) \left\{ \left( \frac{\log(8/\delta)}{n} \right)^{2\beta/(2\beta+1)} \wedge 1 \right\}$ , and when  $\delta \in [2^{-(k_+-1)}, 1]$ , we have

$$\left( \frac{\log(8/\delta)}{n} \right)^{2\beta/(2\beta+1)} \leq C_3(\beta, K) \left\{ \left( \frac{\log(8/\delta)}{n} \right)^{2\beta/(2\beta+1)} \wedge 1 \right\}.$$

The final conclusion follows by combining (33), (34) and (35), and setting  $\check{C}(\beta, K) := C_2(\beta, K) \vee [C_3(\beta, K) \{4C(\beta, K) \vee C_1^2(\beta)\}]$ .  $\square$