

An extension of the low-rank Lyapunov ADI to non-zero initial values and its applications

Jonas Schulze * Jens Saak †

† *Max Planck Institute for Dynamics of Complex Technical Systems,
Sandtorstr. 1, 39106 Magdeburg, Germany.*

* *Corresponding author. Email: jschulze@mpi-magdeburg.mpg.de*

Abstract:

We derive the Alternating-Direction Implicit (ADI) method based on a commuting operator split and apply the results to the continuous time algebraic Lyapunov equation with low-rank constant term and approximate solution. Previously, it has been mandatory to start the low-rank ADI (LR-ADI) with an all-zero initial value. Our approach retains the known efficient iteration schemes of low-rank increments and residual to arbitrary low-rank initial values for the LR-ADI method. We further generalize some of the known properties of the LR-ADI for Lyapunov equations to larger classes of algorithms or problems.

We investigate the performance of arbitrary initial values using two outer iterations in which LR-ADI is typically called. First, we solve an algebraic Riccati equation with the Newton method. Second, we solve a differential Riccati equation with a first-order Rosenbrock method. Numerical experiments confirm that the proposed new initial value of the alternating-directions implicit (ADI) can lead to a significant reduction in the total number of ADI steps, while also showing a 17% and 8× speed-up over the zero initial value for the two equation types, respectively.

Keywords: commuting splitting scheme, commuting operator split, low-rank Lyapunov ADI, complex data, non-zero initial value

Mathematics subject classification: 15A24, 65F10, 65F45, 65F55

Novelty statement: We introduce the notion of fully commuting splitting schemes to solve arbitrary linear systems, and derive the ADI method in that context. This allows us to extend the low-rank Lyapunov ADI to non-zero initial values. Furthermore, we generalize the permutation invariance of ADI iterates to a more general class of algorithms, as well as the existence of a real-valued ADI double-step for complex-conjugated shifts to arbitrary linear systems.

1. Intro

We consider the numerical solution of the continuous-time algebraic Lyapunov equation (ALE)

$$AX + XA^H = -GSG^H \quad (1)$$

with large and sparse coefficient matrix $A \in \mathbb{C}^{n \times n}$ and a low-rank constant term comprised of the factors $G \in \mathbb{C}^{n \times g}$ and $S \in \mathbb{C}^{g \times g}$, where $g \ll n$. This type of equation arises in, e.g., optimal control and model order reduction. We refer to [2, 18, 42] and the references therein for a more detailed introduction.

Given the low rank of the right-hand side of (1), the solution can, at least numerically, be well approximated by a low-rank factorization [36]. We choose the symmetric indefinite factorization, $X \approx ZYZ^H$, with tall and skinny $Z \in \mathbb{C}^{n \times z}$ and Hermitian $Y \in \mathbb{C}^{z \times z}$, $z \ll n$, [15, 31]. For this kind of large-scale equations with low-rank solutions, the most successful algorithms used recently are Krylov subspace projection method, e.g., [28, 29, 34, 41], as well as the ADI method [12, 15, 32, 36, 46]. The latter will be the focus of this paper.

The low-rank Lyapunov ADI had been derived and analyzed using Butcher tableaus for ordinary differential equations (ODEs) [20, 21], as a Smith method [26], as a Krylov method [19, 47], or by Cayley transformations [30]. We revise the derivation of the ADI method using the language of linear splitting methods following [40, Section 5]. The ADI requires the solution of a linear system at every iteration, which is the dominant part of the run-time of the algorithm. Thus far, for every derivation of the low-rank Lyapunov ADI (LR-ADI), it has been mandatory to use an all zero initial guess, $X_0 = 0$, [10–13, 15, 16, 18, 31, 32, 36]. Our motivation is to reduce the number of ADI iterations by extending the method to arbitrary low-rank initial values, $X_0 \neq 0$. We generalize the iteration scheme of Li and White [32] and the residual formulation of Benner, Kürschner, and Saak [12] to arbitrary low-rank initial values for the LR-ADI method at the expense of an indefinite residual.

We investigate the performance of improved initial values using two outer iterations in which LR-ADI is typically called. First, we solve an algebraic Riccati equation with the Newton-ADI method; see e.g., Benner, Li, and Penzl [15]. Every Newton step requires the solution of one algebraic Lyapunov equation. Especially close to convergence of the Newton method, the solution of the previous Newton step is expected to be a good candidate to start the ADI with. We investigate how much an LR-ADI warm-start of the Newton-ADI method can close the performance gap compared to the Riccati ADI (RADI) method of Benner et al. [7].

Second, we solve a differential Riccati equation with a first-order Rosenbrock method. Each time step requires the solution of an algebraic Lyapunov equation; see e.g., Lang, Mena, and Saak [31]. Due to the smoothness of the solution, the solution at the previous time step is a natural candidate to start the next ADI with. Here, we are only concerned with an autonomous equation, and again we investigate how much an LR-ADI warm-start in each time step can close the performance gap to projection-based solvers for the DRE. We expect the real benefits for the non-autonomous case, where no alternative solvers exist, and will separately report the results together with similar approaches for BDF methods for the non-autonomous DRE along the lines of [4, 5].

Throughout the paper, $\|\cdot\|$ denotes the Frobenius norm. $\operatorname{Re}(a+bi) = a$ and $\operatorname{Im}(a+bi) = b$ denote the real and complex part of a complex scalar (or matrix), respectively; $i = \sqrt{-1}$ and $a, b \in \mathbb{R}$. We denote complex conjugation by $\bar{a} + \bar{b}i = a - bi$. $\mathbb{F}^{m \times n}$ denotes the space of m -by- n matrices with entries in the field \mathbb{F} . Transposition of a real matrix is denoted by $(\cdot)^T : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^{n \times m}$, Hermitian transposition of a complex matrix by $(\cdot)^H : \mathbb{C}^{m \times n} \rightarrow \mathbb{C}^{n \times m}$. I_q denotes the identity matrix of size $q \in \mathbb{N}$. Whenever we refer to general linear operators, or the matrix dimensions are evident from the context, we omit the subscript. To simplify notation in many places, we formulate the concatenation of linear operators as multiplication, i.e. $A(B(X)) = AB(X) = ABX$. Spectrum and spectral radius are denoted by $\Lambda(\cdot)$ and $\rho(\cdot)$, respectively.

The paper is structured as follows. In Section 2, we generalize the notion of splitting schemes and prove some of the known properties of the LR-ADI in this more general context. Afterwards, in Section 3, we derive the ADI for arbitrary linear systems using the framework described in Section 2, while generalizing some of the properties known in the Lyapunov case. Section 4 specializes the ADI for Lyapunov equations. In Section 5 we present the two applications mentioned above, as well as some numerical experiments. We conclude the paper in Section 6.

2. Instationary Splitting Schemes

Many iterative methods solving $Ax = b$ can be written in a one-step *splitting* form

$$Mx^{k+1} = Nx^k + b \quad (2)$$

where $M - N = A$ and systems $Mx = d$ are “easy” to solve [24, Section 11.2.3]. These methods are *consistent* by construction, i.e. every solution to $Ax = b$ is a fixed point of the iteration. Conversely,

every fixed point $x^{k+1} = x^k$ is a solution as well. Assuming that M^{-1} exists, the method converges if the spectral radius $\rho(G) < 1$ for $G := M^{-1}N$ [24, Theorem 11.2.1]. G is called *iteration matrix* of the scheme.

The structure above is called *first normal form* of the method [27, Chapter 2]. Consistency is equivalent to $M - N = A$. The *second normal form* of a consistent linear iteration method reads

$$x^{k+1} = M^{-1}((M - A)x^k + b) = x^k - M^{-1}(Ax^k - b). \quad (3)$$

Furthermore, a splitting method is called *nonstationary*¹ if the iteration matrix depends on the iteration, i.e. $A = M_k - N_k$ for $k \in \mathbb{N}$. The normal forms are thus given by

$$M_k x^{k+1} = N_k x^k + b, \quad (4a)$$

$$x^{k+1} = x^k - M_k^{-1}(Ax^k - b). \quad (4b)$$

Here, again, we assume that M_k^{-1} exists. We will see in the later sections that this is actually not a strong assumption in the context of this article.

2.1. Commuting Splitting Schemes

We call an operator split $A = M_k - N_k$ and the corresponding iterative method (4) *commuting* if M_k, N_k commute, i.e.

$$M_k N_k = N_k M_k \quad \forall k \in \mathbb{N}. \quad (5)$$

It is a well known fact that the iteration matrix G_k determines the evolution of the *error* $e^k := x^k - x^*$, where $Ax^* = b$. Due to the consistency of the method, $M_k x^* = N_k x^* + b$ holds. Subtract this from the first normal form (4a) to obtain $M_k e^{k+1} = N_k e^k$ or, equivalently,

$$e^{k+1} = G_k e^k. \quad (6)$$

That the same recursion holds for the *residual* $r^k := Ax^k - b$, is an interesting observation for commuting operator splits.

Proposition 2.1 (Residual Recursion [40, Proposition 5.2]). *Let $A = M_k - N_k$ define a commuting nonstationary splitting method (4). Then, the residual $r^k := Ax^k - b$ adheres to*

$$r^{k+1} = G_k r^k$$

where $G_k := M_k^{-1}N_k$ denotes the iteration matrix.

Proof. Due to $A = M_k - N_k$ it is $AM_k^{-1} = I - N_k M_k^{-1}$. Multiplying (5) with M_k^{-1} from both the left and the right, we find that $M_k^{-1}N_k = N_k M_k^{-1}$. Thus, since M_k and N_k commute, so do A , M_k^{-1} , and N_k . Hence, by substituting the first normal form (4a) into the definition of the residual, we obtain

$$\begin{aligned} r^{k+1} &= Ax^{k+1} - b \\ &= A(M_k^{-1}(N_k x^k + b)) - b \\ &= M_k^{-1}N_k \underbrace{Ax^k}_{r^k + b} + \underbrace{AM_k^{-1}b - b}_{I - N_k M_k^{-1}} \\ &= M_k^{-1}N_k r^k + \underbrace{(M_k^{-1}N_k - N_k M_k^{-1})}_{0} b. \quad \square \end{aligned}$$

This observation allows us to formulate a variant of the second normal form (4b) that iterates the residual r^k and *increment* $v^k := -M_k^{-1}r^k$ alongside the solution x^k . Although Proposition 2.1 may be seen as a special case of Hackbusch [27, Exercise 2.15], it allows us to give much simpler proofs of the known properties of the ADI method applied to Lyapunov equations. The details are covered in Section 4.

Using Proposition 2.1 together with commutation of M_k^{-1} and N_k , observed in its proof, as well as the definition of the increment $v^k := -M_k^{-1}r^k$, we conclude the following corollary.

¹Schulze [40] called this family of algorithms *parametrized* splitting schemes.

Algorithm 1: Commuting Splitting Scheme

Input: operator split $A = M_k - N_k$ with $M_k N_k = N_k M_k$ for $k \in \mathbb{N}$, initial guess x^0
Output: v^0, v^1, \dots such that $x \approx x^0 + v^0 + v^1 + \dots$ solves $Ax = b$

- 1 Assemble initial residual $r^0 \leftarrow Ax^0 - b$
- 2 **for** $k \in \{0, 1, \dots\}$ **do**
- 3 Compute increment $v^k \leftarrow -M_k^{-1} r^k$
- 4 Update residual $r^{k+1} \leftarrow -N_k v^k$ (alternatively, via $r^{k+1} \leftarrow M_k^{-1} N_k r^k$)
- 5 **if** converged **then** break
- 6 **end**
- 7 Assemble solution $x \leftarrow x^0 + v^0 + v^1 + \dots$ if needed

Corollary 2.2. *The residual r^{k+1} only depends on the increment v^k leading to the iterate x^{k+1} , that is*

$$r^{k+1} = -N_k v^k.$$

Combining this with the increment formula once more, gives:

Corollary 2.3. *The increment v^k only depends on the previous increment v^{k-1} , that is*

$$M_k v^k = N_{k-1} v^{k-1}.$$

It may seem that [Corollary 2.2](#) should always give the most efficient way to update the residual. However, [Remark 4.3](#) provides a counter example. At times, [Proposition 2.1](#) in combination with $v^k := -M_k^{-1} r^k$ yields a formula for the residual r^{k+1} that is more efficient to evaluate. In summary, this yields [Algorithm 1](#).

2.2. Fully Commuting Splitting Schemes

We call an operator split $A = M_k - N_k$ and the corresponding iterative method [\(4\)](#) *fully commuting* if $\forall i, j \in \mathbb{N}$ we have

$$M_i N_j = N_j M_i, \quad M_i M_j = M_j M_i, \quad \text{and} \quad N_i N_j = N_j N_i. \quad (7)$$

In the latter case, the order in which the steps defined by (M_0, N_0) to (M_k, N_k) are applied does not matter:

Proposition 2.4 (Permutation Invariance [[40](#), Proposition 5.1]). *Let $A = M_k - N_k$ be a fully commuting operator split. Let the initial iterate x^0 be fixed. Then, the value of x^{k+1} given by the nonstationary splitting method [\(4\)](#) does not depend on the order in which the steps are executed.*

Proof. Let x^* denote the unique solution to $Ax = b$. Observe that $\forall i, j \in \mathbb{N}$, all iteration matrices $G_i := M_i^{-1} N_i$ and G_j commute. Therefore, after applying G_0 through G_k , the error $e^{k+1} = G_k G_{k-1} \cdots G_0 e^0$ does not depend on the order of the steps, as does the iterate $x^{k+1} = e^{k+1} + x^*$. \square

[Proposition 2.4](#) generalizes Li and White [[32](#), Theorem 4.1].

3. ADI Method

The ADI method has originally been described by Peaceman and Rachford [[35](#)]. Suppose $A = H + V$ for some linear operators H and V and the system to be solved is, again, $Ax = b$ for given b . Select some parameters $\alpha_k, \beta_k \in \mathbb{C}$. Using the shorthand notation

$$\begin{aligned} H_k^+ &:= H + \alpha_k I & V_k^+ &:= V + \beta_k I \\ H_k^- &:= H - \beta_k I & V_k^- &:= V - \alpha_k I, \end{aligned} \quad (8)$$

the ADI method reads

$$\begin{aligned} H_k^+ x^{k+\frac{1}{2}} &= b - V_k^- x^k \\ V_k^+ x^{k+1} &= b - H_k^- x^{k+\frac{1}{2}}. \end{aligned} \quad (9)$$

As a splitting scheme (4), the ADI method yields the consistent operator split

$$\begin{aligned} M_k &:= (\alpha_k + \beta_k)^{-1} H_k^+ V_k^+ \\ N_k &:= (\alpha_k + \beta_k)^{-1} H_k^- V_k^- . \end{aligned} \quad (10)$$

If H and V commute, i.e. $HV = VH$, so do all the shorthands (8) as well as the split operators (10). We summarize these findings in the following theorem.

Theorem 3.1 (Fully Commuting Splitting Scheme). *The ADI method is a fully commuting splitting scheme (7) assuming that H and V commute.*

As we will motivate in the next section (Remark 4.4) the parameters may occur in conjugated pairs. That is, $\alpha_{k+1} = \overline{\alpha_k}$ and $\beta_{k+1} = \overline{\beta_k}$. It is therefore reasonable to look for a more efficient means to handle both steps k and $k+1$ at once, especially as the ADI method preserves real iterates.

Theorem 3.2 (Real-valued Double-Step). *Let H and V be commuting real operators, i.e. they map real elements onto real ones. Suppose that iterate x^k and residual r^k are real for some fixed $k \in \mathbb{N}$, and that the next ADI parameters fulfill*

$$\alpha_k \alpha_{k+1}, \quad \alpha_k + \alpha_{k+1}, \quad \beta_k \beta_{k+1}, \quad \beta_k + \beta_{k+1} \in \mathbb{R},$$

for example, $\alpha_{k+1} = \overline{\alpha_k}$ and $\beta_k, \beta_{k+1} \in \mathbb{R}$. Then, two ADI steps later, the iterate x^{k+2} and residual r^{k+2} are real again.

Proof. We show constructively how to compute iterate x^{k+2} and residual r^{k+2} using real arithmetic. The residual $r^{k+2} = G_{k+1} G_k r^k$ is real if the operator $G_{k+1} G_k$ of the combined step is real. Let the scalar

$$\sigma_{k+1,k} := (\alpha_{k+1} + \beta_{k+1})(\alpha_k + \beta_k) \in \mathbb{C} \setminus \{0\}. \quad (11)$$

Recall that H_k^+ and V_{k+1}^+ commute. We observe that the scaled

$$\begin{aligned} \sigma_{k+1,k} M_{k+1} M_k &= \underbrace{(H_{k+1}^+ H_k^+)}_{(H + \alpha_{k+1} I)(H + \alpha_k I)} (V_{k+1}^+ V_k^+), \\ &= H^2 + (\alpha_k + \alpha_{k+1}) H + \alpha_k \alpha_{k+1} I \end{aligned} \quad (12)$$

is a real operator, as is the scaled $\sigma_{k+1,k} N_{k+1} N_k$ and, consequently, $G_{k+1} G_k$. This proves the desired property of the residual r^{k+2} .

Moreover, Corollary 2.3 together with the definition of the increment $v^k := -M_k^{-1} r^k$ implies that $M_{k+1} v^{k+1} = N_k v^k = -N_k M_k^{-1} r^k = -M_k^{-1} N_k r^k$, as N_k and M_k commute. Therefore, by multiplying v^k and v^{k+1} with $M_{k+1} M_k$, we conclude that the increment of the combined step $x^{k+2} = x^k + v^k + v^{k+1}$ is given by

$$M_{k+1} M_k (v^k + v^{k+1}) = -(M_{k+1} + N_k) r^k. \quad (13)$$

Observe that the scaled

$$\sigma_{k+1,k} (M_{k+1} + N_k) = (\alpha_k + \beta_k) (H + \alpha_{k+1} I) (V + \beta_{k+1} I) \quad (14a)$$

$$\begin{aligned} &+ (\alpha_{k+1} + \beta_{k+1}) (H - \beta_k I) (V - \alpha_k I) \\ &= (\alpha_k + \alpha_{k+1} + \beta_k + \beta_{k+1}) HV \end{aligned} \quad (14b)$$

$$\begin{aligned} &+ (\alpha_k \beta_{k+1} + \beta_k \beta_{k+1} - \alpha_k \alpha_{k+1} - \alpha_k \beta_{k+1}) H \\ &+ (\alpha_k \alpha_{k+1} + \alpha_{k+1} \beta_k - \alpha_{k+1} \beta_k - \beta_k \beta_{k+1}) V \\ &+ ((\alpha_k + \alpha_{k+1}) \beta_k \beta_{k+1} + (\beta_k + \beta_{k+1}) \alpha_k \alpha_{k+1}) I \\ &= (\alpha_k + \alpha_{k+1} + \beta_k + \beta_{k+1}) HV + (\beta_k \beta_{k+1} - \alpha_k \alpha_{k+1}) (H - V) \\ &+ ((\alpha_k + \alpha_{k+1}) \beta_k \beta_{k+1} + \alpha_k \alpha_{k+1} (\beta_k + \beta_{k+1})) I \end{aligned} \quad (14c)$$

is a real operator as well. Therefore, given that the residual r^k is real, equation (13) can be solved for the combined increment $v^k + v^{k+1}$ using only real arithmetic. This proves the desired property for the iterate x^{k+2} . \square

Theorem 3.2 generalizes [13]. It motivates to look for an efficient formulation that handles complex shift parameters when applying the ADI method to *any* linear system, in particular, *any* linear matrix equation.

4. Low-rank Lyapunov ADI

In this section, we apply the ADI method to an ALE with low-rank right-hand side and solution. We show how to derive several low-rank variants of the ADI method [9, 15, 16, 31, 32], while extending all these algorithms to non-zero initial values $x^0 \neq 0$. Throughout the section, we use the SVD-type low-rank factorization of, e.g., [16], for the iterates x^k , increments v^k , residuals r^k , and right-hand sides b .

Consider the algebraic Lyapunov equation

$$\mathcal{L}(X) := GSG^H + AX + XA^H = 0 \quad (15)$$

for square matrices $A, X \in \mathbb{C}^{n \times n}$, $G \in \mathbb{C}^{n \times g}$, and $S \in \mathbb{C}^{g \times g}$. We assume that $\lambda_i + \overline{\lambda_j} \neq 0$ for any two eigenvalues $\lambda_i, \lambda_j \in \mathbb{C}$ of A ; that is, equation (15) permits a unique solution; see, e.g., [1, Corollary 1.1.4]. We are aiming to approximate the solution as $X \approx ZYZ^H$, where $Z \in \mathbb{C}^{n \times z}$, $Y \in \mathbb{C}^{z \times z}$. If the constant term has a low rank $g \ll n$, we can expect the solution to have a low numerical rank as well [3, 25, 37, 39, 45].

In terms of the previous section, $H(U) := AU$ and $V(U) := UA^H$, which obviously commute, $(HV)(U) = AUA^H = (VH)(U)$. For generalized equations, the situation is not as simple.

Remark 4.1 (Generalized Matrix Equations). *For a generalized Lyapunov equation*

$$AXE^T + EXA^T = -W$$

we can, in general, not expect A and E to commute. For non-singular E , however, it is still possible to symbolically transform the above equation into its simpler equivalent (15), in which case the restriction of Theorem 3.1 becomes trivial. Then, however, it is mandatory to rephrase the actual steps of the ADI individually along the lines of Saak [38, Section 5.2], to avoid inversion of E .

For this reason, we continue deriving the low-rank ADI for standard equations (15). The overall ADI operator split reads

$$\begin{aligned} M_k(U) &= (\alpha_k + \beta_k)^{-1} (A + \alpha_k I) U (A + \overline{\beta_k} I)^H, \\ N_k(U) &= (\alpha_k + \beta_k)^{-1} (A - \beta_k I) U (A - \overline{\alpha_k} I)^H, \end{aligned} \quad (16)$$

which is symmetry preserving if we choose $\beta_k := \overline{\alpha_k}$. We further require that $-\alpha_k$ is not an eigenvalue of A , such that M_k is invertible. As long as the initial guess $X_0 = Z_0 Y_0 Z_0^H$ is symmetric, the same will hold for the residuals and increments. The residual of a low-rank factorization can itself be expressed as a low-rank factorization, $\mathcal{L}(ZYZ^H) = RTR^H$, where

$$\begin{aligned} R &= \begin{bmatrix} G & Z & AZ \end{bmatrix} \in \mathbb{C}^{n \times (g+2z)} \\ T &= \begin{bmatrix} S & \cdot & \cdot \\ \cdot & \cdot & Y \\ \cdot & Y & \cdot \end{bmatrix} \in \mathbb{C}^{(g+2z) \times (g+2z)} \end{aligned} \quad (17)$$

have rank at most $g + 2z \ll n$. In particular, $\mathcal{L}(0) = GSG^H$ is a rank- g factorization, given both G and S have full rank.

Lemma 4.2 (Consistent Lyapunov Residual). *There exists a factorization of the Lyapunov ADI residual $r^k = R_k TR_k^H$ that is consistent with the initial residual $r^0 = R_0 TR_0^H$. That is, the inner factor T does not depend on the iteration $k \in \mathbb{N}$.*

Proof. Without loss of generality, suppose $r^k = R_k T_k R_k^H$ for $k \in \mathbb{N}$. Substitute the split operators (16) into the residual update of Proposition 2.1, $M_k r^{k+1} = N_k r^k$, to obtain

$$(A + \alpha_k I) R_{k+1} T_{k+1} R_{k+1}^H (A + \alpha_k I)^H = (A - \overline{\alpha_k} I) R_k T_k R_k^H (A - \overline{\alpha_k} I)^H. \quad (18)$$

By choosing the outer factor according to

$$(A + \alpha_k I) R_{k+1} = (A - \overline{\alpha_k} I) R_k, \quad (19)$$

the inner factor does not need to change; that is, $T_{k+1} = T_k = T$. \square

Moreover, we may utilize the definition of the increment v^k for a more efficient evaluation of the residual factors. The formulation $M_k(v^k) = -r^k$ directly reveals a factorization $v^k = V_k \hat{Y}_k V_k^H$, which we will now explain. Using the Lyapunov operator split (16),

$$(2 \operatorname{Re} \alpha_k)^{-1} (A + \alpha_k I) V_k \hat{Y}_k V_k^H (A + \alpha_k I)^H = -R_k T R_k^H, \quad (20)$$

which is fulfilled for

$$(A + \alpha_k I) V_k = R_k, \quad \hat{Y}_k = -2 \operatorname{Re}(\alpha_k) T. \quad (21)$$

Therefore, by simple algebraic reformulations of the Lyapunov residual formula (19), we observe

$$\begin{aligned} R_{k+1} &= (A + \alpha_k I)^{-1} (A - \overline{\alpha_k} I) R_k \\ &= R_k - 2 \operatorname{Re}(\alpha_k) (A + \alpha_k I)^{-1} R_k \\ &= R_k - 2 \operatorname{Re}(\alpha_k) V_k. \end{aligned} \quad (22)$$

This relation has previously been derived by Benner, Kürschner, and Saak [12] in the low-rank ADI context as well as Wolf and Panzer [47] interpreting the iteration as an implicit, in general oblique, Krylov subspace projection method.

Remark 4.3 (Efficient Residual Update). *Had we instead used Corollary 2.2, $r^{k+1} = -N_k v^k$, to derive the residual factors, we would have obtained*

$$R_{k+1} = (A - \overline{\alpha_k} I) V_k, \quad (23)$$

which can at best be evaluated with near-linear complexity if A is sparse. In contrast, formula (22) can be evaluated with linear complexity irrespective of A .

Remark 4.4 (Shifts and Eigenvalues). *Ideally, the residual $r^{k+1} = R_{k+1} T R_{k+1}^H = 0$ should vanish. By formula (23), that means $A V_k = \overline{\alpha_k} V_k$, which is only possible if $\overline{\alpha_k}$ is an eigenvalue of A , and V_k spans (a subspace of) the corresponding eigenspace. Relaxing this to a Ritz-Galerkin-type condition, multiplying with V_k^H from the left, and thus turning the eigenvalues into Ritz-values, motivates why both Penzl's heuristic shifts [36] and the self-generating projection shifts [14] usually perform very well. The fact that, in the case of a real matrix, eigenvalues occur in conjugated pairs motivates Theorem 3.2.*

The problem with the so-called $V(u)$ -shifts [14], however, is that α_k and V_k are not chosen at the same time. Instead, V_k is determined by α_k and the previous residual factor R_k via formula (21), while α_k is determined by some previous increments $\tilde{V} := [V_{k-\ell-1}, V_{k-\ell-2}, \dots, V_{k-\ell-u}]$ via $\alpha_k \in \Lambda(\tilde{V}^H A \tilde{V}, \tilde{V}^H E \tilde{V})$, where $\ell \in \mathbb{N}$ denotes the number of steps taken since the last shift computation. Recall that the solution X has a low (numerical) rank. Therefore, we do not expect the space spanned by the increment V_k to deviate much from \tilde{V} , such that the offset ℓ does not matter much.

The arguments above lead to the algorithm described by Lang, Mena, and Saak [31, Algorithm 3.1]. Recall, however, that our derivation does not require a zero matrix initial guess $X_0 = 0$.

Unfortunately, using the construction from the proof of Theorem 3.2 we obtain the algorithm described by Benner, Li, and Penzl [15, Algorithm 4], and not the more efficient formulation by Benner, Kürschner, and Saak [13, Algorithm 3].

However, the formulations by Lang, Mena, and Saak [31] to handle complex shifts straightforwardly apply to the case of a non-zero initial value in the ADI iteration. We summarize the

Algorithm 2: Frobenius norm of a low-rank factorization

Input: matrices $Z \in \mathbb{C}^{n \times z}$ and $Y \in \mathbb{C}^{z \times z}$, $z \ll n$
Output: $\rho = \|ZYZ^H\|$
1 Compute R from the economy-size QR factorization $Z = QR$, i.e. $R \in \mathbb{C}^{z \times z}$
2 $\rho \leftarrow \|RYR^H\|$

Algorithm 3: Low-rank Lyapunov ADI

Input: system matrices A, E, G , and S , initial value $X_0 = Z_0 Y_0 Z_0^H$,
parameters $\{\alpha_0, \alpha_1, \dots\}$
Output: $V_0, V_1, \dots \in \mathbb{C}^{n \times m}$ and $T \in \mathbb{C}^{m \times m}$ such that $X \approx ZYZ^H$ solves the Lyapunov
equation $AXE^H + EXA^H = -GSG^H$
1 Assemble initial residual factors:

$$R_0 \leftarrow [G \quad EZ_0 \quad AZ_0], \quad T \leftarrow \begin{bmatrix} S & \cdot & \cdot \\ \cdot & \cdot & Y_0 \\ \cdot & Y_0 & \cdot \end{bmatrix}$$

2 $k \leftarrow 0$
3 **repeat**
4 **if** α_k is real **then** // single step
5 Compute increment factor $V_k \leftarrow (A + \alpha_k E)^{-1} R_k$
6 Update residual factor $R_{k+1} \leftarrow R_k - 2 \operatorname{Re}(\alpha_k) E V_k$
7 $k \leftarrow k + 1$
8 **else** // double-step; $\alpha_{k+1} = \overline{\alpha_k}$ must hold
9 Solve complex-valued system $\hat{V}_k \leftarrow (A + \alpha_k E)^{-1} R_k$
10 Compute increment factors:
 $\delta_k \leftarrow \operatorname{Re}(\alpha_k) / \operatorname{Im}(\alpha_k)$
 $V_k \leftarrow \sqrt{2} (\operatorname{Re}(\hat{V}_k) + \delta_k \operatorname{Im}(\hat{V}_k))$
 $V_{k+1} \leftarrow \sqrt{2\delta_k + 2} \operatorname{Im}(\hat{V}_k)$
11 Update residual factor $R_{k+2} \leftarrow R_k - 2\sqrt{2} \operatorname{Re}(\alpha_k) E V_k$
12 $k \leftarrow k + 2$
13 **end**
14 **until** converged
15 Assemble solution factors, if needed:
 $Z \leftarrow [Z_0 \quad V_0 \quad V_1 \quad \dots]$
 $Y \leftarrow \operatorname{blockdiag}(Y_0, -2 \operatorname{Re}(\alpha_0) T, -2 \operatorname{Re}(\alpha_1) T, \dots)$

low-rank Lyapunov ADI as derived by [Proposition 2.1](#) and [Remark 4.1](#) in [Algorithm 3](#). For the convergence criterion in [line 14](#) we use

$$\|r^k\| = \|R_k T R_k^H\| \leq \begin{cases} \operatorname{abstol}_{\text{ADI}} \\ \operatorname{reitol}_{\text{ADI}} \|GSG^H\| = \operatorname{reitol}_{\text{ADI}} \|b\|, \end{cases} \quad (24)$$

depending on whether $\operatorname{abstol}_{\text{ADI}} \in \mathbb{R}$ or $\operatorname{reitol}_{\text{ADI}} \in \mathbb{R}$ has been provided. Either way, we evaluate all Frobenius norms of low-rank factorizations using [Algorithm 2](#) [[15](#), [36](#)]. Note that accumulation of Q can be avoided [[15](#)], e.g., using Householder QR or (shifted) Cholesky QR; see [[24](#)].

Remark 4.5 (Other formulations of the ADI iteration for Linear Matrix Equations). *Applying [Corollary 2.3](#), $M_k v^k = N_{k-1} v^{k-1}$, to the Lyapunov operator split (16) for Cholesky-type low-rank iterates $X_k = Z_k Z_k^H$ and data $W = BB^H$ yields the CF-ADI algorithm described by Li and White [[32](#)]. Applying the same reformulation as in equation (22) leads to the LR-ADI as described by Benner, Li, and Penzl [[15](#)]. Recall, however, that our derivation does not require a zero matrix initial guess; that is, $X_0 = 0$ or $Z_0 = []$.*

Applying [Corollary 2.3](#), $M_k v^k = N_{k-1} v^{k-1}$, to the Sylvester equation $AX - XB = W$ leads to the fADI method described by Benner, Li, and Truhar [[16](#)]. Specializing [Algorithm 1](#) to the

Sylvester equation leads to the G -fADI described by Benner and Kürschner [9], which iterates the residual alongside.

Remark 4.6 (Alternative Derivation). *Observe that $\mathcal{L}(X_0 + \hat{V}) = 0$ is a Lyapunov equation in the increment \hat{V} ,*

$$\mathcal{L}(X_0 + \hat{V}) = RTR^H + A\hat{V} + \hat{V}A^H = 0.$$

where R and T are the Lyapunov residual factors (17). Applying the ADI to solve for \hat{V} and starting with rank-zero $\hat{V}_0 = 0$, is equivalent to applying the ADI to solve for $X_0 + \hat{V}$ and starting with X_0 .

5. Applications and Numerical Experiments

In this section, we recall two iterative algorithms that solve an ALE at every step. If we solve these ALEs with another iterative method, like the ADI, it is natural to use the previous outer iterate as the initial guess for the inner method.

More specifically, let X_ℓ denote the outer iterates, $\ell \in \mathbb{N}$, and let $X_{\ell+1,k}$ denote the inner iterates, $k \in \mathbb{N}$, whose final iterate will become $X_{\ell+1}$. Then, it is natural to choose $X_{\ell+1,0} = X_\ell$ instead of, e.g., $X_{\ell+1,0} = 0$.

All computations in this section have been performed on an Intel Xeon Silver 4110. We evaluated the following shift strategies:

- `heur(10, 10, 10)`: Penzl's $\text{heur}(l_0, k_+, k_-)$ strategy, which first computes eigenvalues based on k_+ Arnoldi iterations of $E^{-1}A$ and k_- Arnoldi iterations of $A^{-1}E$, and selects l_0 of them based on a greedy heuristic [36]. If more than l_0 shifts are needed, we repeat these l_0 -many selected shifts cyclically.
- `heur(20, 30, 30)`
- `proj(2, heur)`: One of the self-generating projection strategies, which is called $V(u)$ -shifts in [30, Section 5.3.1]. In our case $u = 2$, to properly account for a potential ADI double-step. Here, we order these Ritz values based on Penzl's heuristic.
- `proj(2, dec)`: The same $V(2)$ -shifts ordered by decreasing real part. In the case of real-valued shifts, the shifts are equivalently ordered by increasing magnitude, since all such shifts must have a negative real part.
- `proj(2, inc)`: The same $V(2)$ -shifts ordered by increasing real part.

In general, we expect fewer ADI iterations for larger $l_0 \in \mathbb{N}$ in the case of $\text{heur}(l_0, k_+, k_-)$ shifts, and even fewer iterations for the `proj` shifts. In that notion, we call shifts to be better or worse than others.

Remark 5.1 (Accuracy of Ritz Values). *It is critical that Penzl's shifts are computed precisely as described in [36], since it is by no means clear that the largest or smallest eigenvalues and corresponding eigenspaces are the most relevant for the solution process. For example, using the k_+ largest and k_- smallest eigenvalues of $E^{-1}A$ as computed by `ArnoldiMethod.jl` [44] (version 0.2.0) compared to Penzl's procedure, limiting to only a few Arnoldi steps and Ritz values, being comparably rough approximations of the eigenvalues, leads to 5 to 10× more ADI iterations. These values are too accurate and ignore relevant eigenspaces still covered by Penzl's Ritz values. When computing highly accurate eigenvalues, choosing those from the dominant poles of the underlying dynamical system, for which the ALE is solved, can be beneficial [38, Section 8.1].*

Remark 5.2 (Order of Shifts and Eigenvalues). *Benner, Kürschner, and Saak require that complex conjugated pairs remain adjacent, but do otherwise not specify the order in which the V -shifts shall be used [14]. As we will see later, the ADI can be quite susceptible to the order of the shifts. Given that, for example, MATLAB's `eig` does not guarantee a particular order but appears to keep complex conjugated pairs adjacent, while Julia's `eigvals` sorts the eigenvalues lexicographically*

by real and imaginary parts (similar to the $\text{proj}(2, \text{inc})$ shifts), the unaware user may observe drastically different convergence behaviors. Try, for example,

$$A = \text{blockdiag} \left(\begin{bmatrix} -3 & 4 \\ -4 & 3 \end{bmatrix}, \begin{bmatrix} -4 & 3 \\ -3 & 4 \end{bmatrix}, \begin{bmatrix} -3 & 2 \\ -2 & 3 \end{bmatrix}, -5, -3 \right).$$

Julia's order is problematic insofar as complex conjugated pairs are not adjacent, which is required for an ADI double-step. Therefore, when sorting the eigenvalues based on their real part, we also sort by the absolute value of their imaginary parts, in order to keep complex conjugated pairs adjacent. Penzl's heuristic ensures this property by itself.

We conclude this section's introduction by mentioning some implementation details. The examples to follow will all lead to real-valued system matrices. We therefore expect a real-valued solution $ZYZ^T \in \mathbb{R}^{n \times n}$ as well; see [Theorem 3.2](#). All Frobenius norms of ZYZ^T factorizations are evaluated using [Algorithm 2](#). The systems arising in the ADI increment formulas (21), lines 4 and 8 of [Algorithm 3](#), may have a low-rank updated structure,

$$\underbrace{A + \alpha_k E}_{\text{sparse}} + \underbrace{UV^T}_{\text{low-rank}} \in \mathbb{C}^{n \times n}, \quad (25)$$

where U and V are real and have few columns. The only complex contribution will be from the ADI shifts $\alpha_k \in \mathbb{C}$. In this case, we apply the Sherman-Morrison-Woodbury formula; e.g., [\[24, Section 2.1.4\]](#).²

We perform a column compression of any factored representation $ZYZ^T \in \mathbb{R}^{n \times n}$, as described by Lang, Mena, and Saak [\[31\]](#), every 10 low-rank additions/subtractions,

$$Z_1 Y_1 Z_1^T \pm Z_2 Y_2 Z_2^T = [Z_1 \quad Z_2] \underbrace{\begin{bmatrix} Y_1 \\ \pm Y_2 \end{bmatrix}}_{Y \in \mathbb{R}^{k \times k}} [Z_1 \quad Z_2]^T, \quad (26)$$

or once the inner dimension reaches 50% of the outer dimension; $k \geq 0.5n$. The eigenvalues of the resulting inner matrix \hat{Y} are truncated such that

$$\Lambda(\hat{Y}) = \{ \lambda \in \Lambda(Y) : |\lambda| \geq \max\{1, \rho(Y)\} \cdot k \cdot u_{\text{mach}} \}, \quad (27)$$

where $\Lambda(\cdot)$ and $\rho(\cdot)$ denote the spectrum and spectral radius, respectively. We further compress every low-rank Lyapunov right-hand side W before solving $AX + XA^T = -W$, unless mentioned otherwise.

5.1. Algebraic Riccati Equation

We apply the Newton-Kleinman method to the algebraic Riccati equation (ARE)

$$\mathcal{R}(X) := C^T C + A^T X E + E^T X A - E^T X B B^T X E = 0, \quad (28)$$

with matrices

$$E, A \in \mathbb{R}^{n \times n}, \quad B \in \mathbb{R}^{n \times m}, \quad C \in \mathbb{R}^{q \times n}, \quad (29)$$

and $m, q \ll n$. Due to the low rank of $C^T C$ and $B B^T$, we can expect the solution to have a low numerical rank; see Benner and Bujanović [\[6\]](#). Hence, we factorize the Newton iterates according to $X_\ell = Z_\ell Y_\ell Z_\ell^T$ with $Z_\ell \in \mathbb{R}^{n \times z_\ell}$ and $Y_\ell \in \mathbb{R}^{z_\ell \times z_\ell}$. Adapted from Benner, Li, and Penzl [\[15\]](#), the ℓ th step ($\ell \in \mathbb{N}$) of the Kleinman formulation reads

$$0 = \mathcal{L}_\ell(X_{\ell+1}) := G_\ell S_\ell G_\ell^T + A_\ell^T X_{\ell+1} E + E^T X_{\ell+1} A_\ell, \quad (30)$$

²For more general examples, the structure of the low-rank part in (25) may be UDV^T for some $D \neq I$. In that case, the center terms S_ℓ of the Lyapunov equations to come (equations (30) and (38)) need to be modified slightly, and the Sherman-Morrison-Woodbury formula requires a term D . For ease of notation, we stick to the simpler case (25).

where

$$\begin{aligned} A_\ell &= A - BB^\top X_\ell E \in \mathbb{R}^{n \times n}, \\ G_\ell &= [C^\top \quad E^\top X_\ell B,] \in \mathbb{R}^{n \times (q+m)}, \\ S_\ell &= I \in \mathbb{R}^{(q+m) \times (q+m)}. \end{aligned} \quad (31)$$

Note that A_ℓ has the structure of a sparse matrix updated by a low-rank matrix product, using $U = B$ and $V^\top = B^\top X_\ell E$. Let $X_{\ell+1,k}$ for $k \in \mathbb{N}$ denote the ADI iterates during the solution of (30). The dimension of the right-hand side factor G_ℓ does not depend on the current iterate X_ℓ .³ Therefore, the ADI started with $X_{\ell+1,0} = 0$ will have an advantage, as its Lyapunov residuals $\mathcal{L}_\ell(X_{\ell+1,k})$ will always have (at most) $q + m$ columns, which determines the cost to solve the linear systems defining the ADI increments. The new ADI with $X_{\ell+1,0} = X_\ell = Z_\ell Y_\ell Z_\ell^\top$ having inner dimension z_ℓ will lead to Lyapunov residuals having up to $(q + m) + 2z_\ell$ columns; see formula (17). We do not yet use the common $E^\top Z_\ell$ block in the outer low-rank factor, which would allow factorization having inner dimension $q + 2z_\ell < q + m + 2z_\ell$ and, thus, slightly reduced compression times.

The Riccati residual of a low-rank factorization can naively be written in a low-rank form as $\mathcal{R}(Z_\ell Y_\ell Z_\ell^\top) = RTR^\top$, where

$$\begin{aligned} R &= [C^\top \quad A^\top Z_\ell \quad E^\top Z_\ell] \in \mathbb{R}^{n \times (q+2z_\ell)} \\ T &= \begin{bmatrix} I & \cdot & \cdot \\ \cdot & \cdot & Y_\ell \\ \cdot & Y_\ell & -Y_\ell Z_\ell^\top B B^\top Z_\ell Y_\ell \end{bmatrix} \in \mathbb{R}^{(q+2z_\ell) \times (q+2z_\ell)}, \end{aligned} \quad (32)$$

compare, e.g., [15]. We consider the Newton method to have converged once

$$\|\mathcal{R}(X_{\ell+1})\| < \text{reitol}_{\text{Newton}} \|C^\top C\| \quad (33)$$

for some user-provided $\text{reitol}_{\text{Newton}}$. Next, we describe how to configure the ADI convergence criterion (24) based on three different variations of the Newton method. First, for the classical Newton method we use

$$\text{reitol}_{\text{ADI}} = \text{reitol}_{\text{Newton}}/10. \quad (34a)$$

Second, to speed up the early Newton iterations, we use an inexact Newton method with variable forcing term $\eta \in \mathbb{R}$ described by Dembo, Eisenstat, and Steihaug [23], resulting in

$$\text{abstol}_{\text{ADI}} = \eta \|\mathcal{R}(X_\ell)\|. \quad (34b)$$

Third, to speed up the later Newton steps with already small $\|\mathcal{R}(X_\ell)\|$, we switch back to the classical Newton method if the inexact bound (34b) becomes smaller than the classical bound (34a). This ‘‘hybrid’’ Newton method results in

$$\text{abstol}_{\text{ADI}} = \max \left\{ \eta \|\mathcal{R}(X_\ell)\|, \frac{1}{10} \text{reitol}_{\text{Newton}} \|\mathcal{L}_\ell(0)\| \right\}. \quad (34c)$$

Furthermore, to keep the potential overshoot of the Riccati residual in control, which refers to an increasing Riccati residual, as usually observed in the first Newton iterations, we employ an Armijo line search.⁴ As described by Benner et al. [8], the line search is applied if $\|\mathcal{R}(\hat{X}_{\ell+1})\| > 0.9 \|\mathcal{R}(X_\ell)\|$, where $\hat{X}_{\ell+1}$ denotes the solution to the classical Newton step (30).

Example 5.3. *We apply the procedure described above to the Steel Profile benchmark [17, 33], which is based on a semi-discretized heat transfer problem. The corresponding matrices (36) have $m = 7$ inputs, $q = 6$ outputs, and are available for several sizes n . For this work, we focus on the configuration $n = 5177$. However, the input matrix $B = 1000\hat{B}$ is a scalar multiple of the original Steel Profile matrix \hat{B} . The factor 1000 controls the relative weighting of control and output costs of the corresponding linear quadratic regulator problem. In this setting it corresponds to a factor 10^{-6}*

³Except for an outer initial value $X_0 = 0$, which would allow $G_0 = C^\top$ which has only q columns. For simplicity, we did not implement this, nor do we perform a column-compression on the right-hand side.

⁴Our implementation is rather naive in that it does not utilize the connection between $\mathcal{R}(X)$ and $\mathcal{L}_\ell(X)$ described in [8, Equation (5.2b)]. However, the overall runtime cost of the line search was negligible (less than 1% of the total runtime).

on the control costs, decreasing the regularity of the optimal control problem and, thus, making the ARE harder to solve. Consequently, we observe a higher overshoot of the Riccati residual norm and more Newton steps. We use the quadratic forcing term $\eta = \min\{0.1, 0.9\|\mathcal{R}(X_\ell)\|\}$ also used by Benner et al. [8]. The Newton tolerance is $\text{reltol}_{\text{Newton}} = 10^{-10}$.

Table 1 gives an overview on the number of ADI and Newton iterations required to solve the ARE (28) arising from Example 5.3, as well as the accumulated run-time of the ZYZ^T columns compression, t_{compress} , the time to compute the ADI shifts, t_{shifts} , the time of all the linear solves of Algorithm 3, t_{solve} , as well as the total run-time of the Newton method. The ADI shift strategies are described at the beginning of this section. We observe that enabling the line search reduces the number of Newton steps, but only mildly reduces the number of ADI steps, as mentioned in [8]. Switching to the inexact or hybrid Newton method has a much larger effect on the number of ADI steps. Replacing the zero initial ADI value by the previous Newton iterate further reduces the number of ADI steps by up to $4\times$ (516 vs 121). This reduction is strongest for the simplest shift strategy, $\text{heur}(10, 10, 10)$. The old ADI outperforms the new one in most configurations of the Newton method, and its run-time is dominated by t_{solve} . Only for the inexact and hybrid Newton method with $\text{heur}(10, 10, 10)$ shifts, our new ADI has a slight run-time advantage. In these cases, however, t_{compress} is the most expensive part of the algorithm, indicating further potential for improvements especially for our new ADI, when only bad shifts are known.

We observe a certain dependence of the new ADI on the order of the (projection) shifts. By merely changing from the decreasing to the increasing order (see Remark 5.2), the number of ADI iterations more than doubles (595 vs 1302). The large number of ADI iterations for $\text{proj}(2, \text{inc})$ shifts is somewhat reasonable; refer to Appendix A for an explanation. Meanwhile, the old ADI seems barely affected. However, most configurations using inexact or hybrid Newton with projection shifts did not converge. Only the new ADI with $\text{proj}(2, \text{dec})$ shifts made the inexact methods without line search converge. Further inspection is needed to understand this phenomenon.

In terms of the number of ADI iterations, the optimal configuration for the old ADI is the hybrid Newton method with line search and $\text{heur}(20, 30, 30)$ shifts. The new ADI performed best for the same Newton method and the simpler $\text{heur}(10, 10, 10)$ shifts. Our present implementation of the heuristic shift strategy is not yet optimized, which skews the total run-time of the old ADI in favor of the simpler shifts. Thus, currently, we observe a $3\times$ reduction in the number of ADI steps (348 vs 121), and a modest speedup of about 17% in favor of the new ADI (18.3s vs 15.7s) when compared to the old ADI with the lowest number of ADI steps and run-time, respectively. However, for any configuration other than the inexact Newton methods with the simplest shift strategy, the old ADI outperforms our new ADI. More research is needed in how to select the optimal Newton configuration and ADI shifts.

For the remainder of this subsection, we focus on configuration with the lowest run-time for either ADI, namely the hybrid Newton method with line search using $\text{heur}(10, 10, 10)$ shifts. Figure 1 shows that the new ADI requires much fewer iterations per Newton step; as expected. We further see that the number of old ADI iterations per step does not increase by much beyond Newton step 10 due to the classical bound becoming active in formula (34c) for the last three Newton iterations. However, the linear systems to be solved for the ADI increments (21) have more columns for the non-zero initial guess; $6\times$ as many (78 vs 13) at Newton step 7. Despite this imbalance, the new ADI requires slightly fewer linear system solves overall (6457 vs 6708). In total, we observe a 6.0s reduction in t_{solve} at the cost of a 1.4s increase in t_{compress} . In the future, we need to optimize our implementations computing the heuristic shifts as well as the low-rank column compression.

If we plot the Riccati and Lyapunov residuals for all inner ADI iterations directly one after the other, we obtain Figure 2. When starting the next Newton step's ADI with the current Newton iterate, $X_{\ell+1,0} = X_\ell$, the corresponding Riccati residual obviously does not change. Meanwhile, when starting with the zero matrix, $X_{\ell+1,0} = 0$, the normalized Riccati residual will jump back to 1. Both effects are clearly visible in Figure 2; for the curve of the old ADI it marks the start of every Newton iteration. The curve of the new ADI jumps back to 1 only due to the single line search applied after the first Newton step. Furthermore, we observe a stagnation of the naive Riccati residual (32) during the ADI towards the end of every Newton step irrespective of its initial value (solid lines), even though the Lyapunov residual keeps getting smaller (dashed lines). Benner et al. describe a more efficient representation of the Riccati residual $\mathcal{R}(\cdot)$ in relation to the Lyapunov residual $\mathcal{L}_\ell(\cdot)$ and the change in the thin rectangular matrix $B^T X_\ell E \in \mathbb{R}^{m \times n}$ [8, Equation 5.2b].




Figure 1: Newton method (hybrid w/ line search) applied to solve ARE (28) arising from Example 5.3. ADI shifts: $\text{heur}(10, 10, 10)$. The ADI is started from a zero value (●) or with the previous Newton iterate (▲).

This representation allows to monitor the outer Riccati residual in the inner ADI iteration, which we plan to add to our implementation in the future. Then the ADI can stop early whenever a sufficient decrease condition for the Newton step is fulfilled.

We recommend to investigate the cause of the aforementioned stagnation of the Riccati residual. Furthermore, a hybrid ADI approach should be studied, that switches from the zero initial value to the non-zero one, once the hybrid Newton criterion (34c) resolves to the classical condition (34a).

5.2. Differential Riccati Equation

We apply a first-order Rosenbrock scheme, the implicit Euler method, to the differential Riccati equation (DRE)

$$E^T \dot{X} E = \mathcal{R}(X) := C^T C + A^T X E + E^T X A - E^T X B B^T X E, \quad E^T X(t_0) E = C^T C, \quad (35)$$

where, again,

$$E, A \in \mathbb{R}^{n \times n}, \quad B \in \mathbb{R}^{n \times m}, \quad C \in \mathbb{R}^{q \times n}, \quad (36)$$

and $m, q \ll n$. As $C^T C$ and $B B^T$ are positive semi-definite, equation (35) has a unique solution; see, e.g., [1, Theorem 4.1.6]. Motivated by Stillfjord [43] and following Lang, Mena, and Saak [31], we factorize the solution according to $X_\ell = Z_\ell Y_\ell Z_\ell^T \approx X(t_0 + \ell\tau)$ with

$$Z_\ell \in \mathbb{R}^{n \times z_\ell}, \quad Y_\ell \in \mathbb{R}^{z_\ell \times z_\ell}, \quad (37)$$

where $z_\ell \ll n$. Thus, every Rosenbrock step reads

$$\mathcal{L}_\ell(X_{\ell+1}) := G_\ell S_\ell G_\ell^T + A_\ell^T X_{\ell+1} E + E^T X_{\ell+1} A_\ell = 0 \quad (38)$$

where

$$\begin{aligned} A_\ell &= A - \frac{1}{2\tau} E - B B^T X_\ell E && \in \mathbb{R}^{n \times n} \\ G_\ell &= \begin{bmatrix} C^T & E^T Z_\ell \end{bmatrix} && \in \mathbb{R}^{n \times (q+z_\ell)} \\ S_\ell &= \begin{bmatrix} I & \\ & Y_\ell Z_\ell^T B B^T Z_\ell Y_\ell + \frac{1}{\tau} Y_\ell \end{bmatrix} && \in \mathbb{R}^{(q+z_\ell) \times (q+z_\ell)}. \end{aligned} \quad (39)$$




Figure 2: Normalized Riccati residuals $\|\mathcal{R}(X_{\ell+1,k})\|/\|\mathcal{R}(0)\|$ following the naive formula (32), and normalized Lyapunov residuals $\|\mathcal{L}_\ell(X_{\ell+1,k})\|/\|\mathcal{L}_\ell(0)\|$ following Algorithm 3, over the course of all ADI iterations k during all Newton steps ℓ . Newton method: hybrid w/ line search. ADI shifts: $\text{heur}(10, 10, 10)$. The ADI is started from a zero value (old) or with the previous Newton iterate (new).

Note, again, that A_ℓ has the structure of a sparse matrix updated by a low-rank matrix product. This time, however, the inner dimension of the right-hand side $G_\ell S_\ell G_\ell^\top$, and therefore the corresponding Lyapunov residual $\mathcal{L}_\ell(X_{\ell+1,k})$ both depend on the rank of the current Rosenbrock iterate X_ℓ . More precisely, the rank is at most $(q + z_\ell) + 2z_\ell$; see formula (17). Again, we do not yet utilize the common $E^\top Z_\ell$ in the outer low-rank factor, which would allow for a factorization having inner dimension $q + 2z_\ell < q + 3z_\ell$ and, thus, reduced compression times.

Example 5.4. We apply the procedure described above to the Steel Profile benchmark [17, 33], see Example 5.3. The time span is $[t_0, t_f] = [0, 4500]$, which we discretize in $\ell_{\max} = 45$ or 450 equidistant segments. That is, $\tau = 100$ or 10. The convergence criterion (24) is based on $\text{reitol}_{\text{ADI}} = 10^{-10}$;

Table 2 gives an overview on the number of ADI iterations required to solve the DRE (35) arising from Example 5.4. The timings are as described for Table 1. Refer to the beginning of this section for an explanation on the shift strategies chosen. Changing the initial ADI guess from the zero matrix to the previous Rosenbrock iterate $X_{\ell-1}$ always reduced the overall number of ADI iterations. In contrast to the previous section, the run-time is always dominated by t_{compress} . This motivates, again, to improve the implementation of the low-rank column compression.

For this application, both ADIs are susceptible to the order of the shifts. Focusing on the projection shifts for 450 Rosenbrock steps, the number of ADI iterations varies by more than $6 \times$ by merely changing the order of the shifts (old: 54 657 vs 8247, new: 9693 vs 1306). In terms of the number of iterations, the old ADI performs best for $\text{proj}(2, \text{heur})$ shifts and appears to degenerate for $\text{proj}(2, \text{dec})$ shifts. Meanwhile, the new ADI performs best for this order; taking less than 3 steps on average; and mediocre for $\text{proj}(2, \text{heur})$ shifts.

Figure 3 shows some more detail for one of the configurations of the table; the general shape is the same for all configurations. The DRE (35) is very stiff during early Rosenbrock steps $\ell \in \mathbb{N}$. Once the integrator reaches a more transient regime for large-enough ℓ , using the non-zero initial ADI value reduces the number of ADI steps drastically; as expected. Surprisingly, however, the number of columns comprising the linear systems to compute the ADI increments (21) decreases as well. Apparently, the previous Rosenbrock iterate X_ℓ contains enough information to cause the Lyapunov residual $\mathcal{L}_\ell(X_\ell)$ to vanish in many directions of the corresponding subspace. Overall, the new ADI has to solve fewer and cheaper linear systems. Consequently, the finer the temporal

Table 2: Total number of ADI iterations k_{total} performed, as well as several run-time metrics to solve the DRE (35) arising from Example 5.4. All timings are in seconds. The ADI is started from a zero value (old) or with the previous Rosenbrock iterate (new). A speedup >1 indicates that our new ADI is faster.

#Rosenbrock steps	ADI shifts	#ADI steps		t_{compress}		t_{shifts}		t_{solve}		t_{total}		Speedup
		old	new	old	new	old	new	old	new	old	new	
45	heur(10, 10, 10)	1395	671	211.8	41.2	7.6	7.6	53.5	18.8	314.7	89.5	3.52
45	heur(20, 30, 30)	1170	683	177.8	43.4	24.3	24.1	44.7	19.0	282.5	109.9	2.57
45	proj(2, dec)	5685	550	971.1	72.7	3.7	2.1	222.5	19.3	1357.7	119.6	11.35
45	proj(2, heur)	1266	821	201.0	49.3	3.9	1.6	49.9	22.2	293.7	97.3	3.02
45	proj(2, inc)	4247	2172	670.3	181.8	3.7	1.7	164.4	64.1	958.7	296.0	3.24
450	heur(10, 10, 10)	9450	3473	1572.3	127.9	172.5	133.8	562.4	93.3	2742.8	536.7	5.11
450	heur(20, 30, 30)	9000	3392	1626.2	139.6	243.6	285.9	487.2	93.8	2849.3	706.9	4.03
450	proj(2, dec)	54 657	1306	10 727.1	73.6	38.2	8.2	3525.8	54.0	16 405.3	313.2	52.38
450	proj(2, heur)	8247	3739	1511.3	129.8	38.6	10.1	538.5	119.0	2484.7	469.9	5.29
450	proj(2, inc)	25 487	9693	4677.9	396.3	37.3	12.4	1646.7	322.1	7401.2	1037.6	7.13




Figure 3: Rosenbrock method applied to solve DRE (35) arising from Example 5.4. ADI shifts: $\text{heur}(20, 30, 30)$. Rosenbrock step size $\tau = 10$. The ADI is started from a zero value (\bullet) or with the previous Rosenbrock iterate (\blacktriangle).

resolution, the larger the expected speedup of the new ADI.⁵ This notion is confirmed by Table 2. When increasing the number of Rosenbrock steps by $10\times$, the total number of iterations and run-time of the new ADI only increase by roughly 2.5 to $5\times$ and 2 to $6\times$, respectively, depending on the shifts. Meanwhile, the total number of iterations and run-time of the old ADI increase by roughly 6 to $10\times$ and 7.5 to $12\times$, respectively, depending on the shifts. Comparing the best shifts for each of the ADIs applied over 450 Rosenbrock steps, we observe a $6\times$ reduction in ADI steps (8247 vs 1306) and an $8\times$ reduction in run-time (2484.7s vs 313.2s) in favor of our new ADI.

Note that, although prescribed, the order of the heuristic shifts does not matter that much. The heuristic strategies compute only few (here: 10 or 20) shifts at once. Once all of them have been used, the iterates and residuals coincide; see Proposition 2.4. The projection strategies, however, compute twice the inner dimension of the Lyapunov residual many at once (here: approx. 80 to 250). For the vast majority of Rosenbrock steps, the ADI does converge after only a subset of these shifts has been used; that is, Proposition 2.4 does not apply. Further research is needed to understand the effect of the shift order in such a scenario, and how to select the optimal shift order (and subset of shifts).

6. Conclusion

We introduced the notion of fully commuting splitting schemes to solve arbitrary linear systems, and derived the ADI method in that context. This allowed us to extend the low-rank Lyapunov ADI for complex data to non-zero initial values. Furthermore, we generalized the permutation invariance of ADI iterates to arbitrary fully commuting splitting schemes (Proposition 2.4), as well as the existence of a real-valued ADI double-step for complex-conjugated shifts to arbitrary linear systems (Theorem 3.2).

We applied the extended low-rank Lyapunov ADI to a Newton and a Rosenbrock method to solve an algebraic and a differential Riccati equation, respectively. For the Newton method we observed a $4\times$ lower total number of ADI steps for heuristic Penzl shifts, but more expensive linear systems at every ADI step. Overall, for this application, our implementation only showed

⁵Adaptive time stepping may reduce this advantage of our new ADI, as regions allowing for large time steps are the ones that would benefit most from our initial values. Therefore, as a stopgap specifically for Example 5.4, we select temporal resolutions that are rather coarse from an engineer's perspective.

a modest 17% run-time improvement over the old ADI. Therefore, at the moment we doubt that our modifications to the Newton-ADI for AREs suffice to make the method competitive with the RADI [7]. For the Rosenbrock method, however, we observed a 2 to 6× lower total number of ADI steps, depending on the shifts, whose linear systems have fewer columns, resulting in an 8× speed-up in our implementation.

Code and Data Availability




The algorithms have been implemented using Julia [22] and are available at:

DOI [10.5281/zenodo.10650859](https://doi.org/10.5281/zenodo.10650859)

The datasets analyzed in this paper are available at:

Newton: DOI [10.5281/zenodo.10650872](https://doi.org/10.5281/zenodo.10650872)
 Rosenbrock: DOI [10.5281/zenodo.10651124](https://doi.org/10.5281/zenodo.10651124)

Acknowledgments

We would like to thank Daniel Szyld  for pointing out the name *nonstationary* splitting scheme. We further thank Fan Wang  and Martin Köhler  for their review of our codes and independent verification of the reproducibility of the numerical experiments.

A. Order of ADI shifts

As we have observed in Section 5, the low-rank Lyapunov ADI can be quite sensitive to the order of its shifts. In this section, we give some intuition on why the projection shifts ordered by increasing real part, $\text{proj}(2, \text{inc})$, sometimes lead to large numbers of ADI iterations.

Using the notation of Section 2, the iteration map of the symmetry-preserving low-rank Lyapunov ADI, as derived in Section 4, is given by

$$G_k(U) := (M_k^{-1}N_k)(U) = \mathcal{C}(A, \alpha_k) U \mathcal{C}(A, \alpha_k)^H, \quad (40)$$

using the operator split (16), $\beta_k := \overline{\alpha_k}$, and the Cayley transformation

$$\mathcal{C}(A, \alpha) := (A + \alpha I)^{-1}(A - \overline{\alpha}I). \quad (41)$$

If applied to a symmetric low-rank factorization $ZY Z^H$, the iteration map effectively only operates on the outer factors Z . Recall that a nonstationary splitting scheme converges iff $\rho(G_k \cdots G_0) \rightarrow 0$ as $k \rightarrow \infty$. A sufficient condition is thus

$$\rho(\mathcal{C}(A, \alpha_k) \cdots \mathcal{C}(A, \alpha_0)) \rightarrow 0 \quad (42)$$

as $k \rightarrow \infty$. As the spectral radius is sub-multiplicative, the upper bound

$$\rho(\mathcal{C}(A, \alpha_k) \cdots \mathcal{C}(A, \alpha_0)) \leq \rho(\mathcal{C}(A, \alpha_k)) \cdots \rho(\mathcal{C}(A, \alpha_0)) =: \hat{\rho}_k \quad (43)$$

holds. It can be computed as follows. By, e.g., [30, Proposition 2.16], the spectral radius of such a Cayley transformation is given as

$$\rho(\mathcal{C}(A, \alpha)) = \max \left\{ \frac{|\lambda - \overline{\alpha}|}{|\lambda + \alpha|} : \lambda \in \Lambda(A) \right\}. \quad (44)$$

If we chose the parameters $\{\alpha_0, \dots, \alpha_k\}$ to be (a subset of) the spectrum $\Lambda(A)$, it holds $\rho(\mathcal{C}(A, \alpha_k)) \leq 1$ for any k . Furthermore, every eigenvector v of A to the eigenvalue λ lives in the null space of $\mathcal{C}(A, \lambda)$. Consequently, if the whole $\Lambda(A)$ is chosen, the norm of the combined step $\rho(\mathcal{C}(A, \alpha_k) \cdots \mathcal{C}(A, \alpha_0))$ will be zero. On the other hand, any individual $\rho(\mathcal{C}(A, \alpha_k)) > 0$. That is, $\hat{\rho}_k$ is not a sharp upper bound.




Figure 4: Spectral radius of Cayley transformations associated to spectrum of A , and upper bound $\hat{\rho}_k$ on the norm of parts of the iteration map over the course of multiple ADI iterations k , for different permutations of the spectrum $\Lambda(A)$.

Figure 4 shows the aforementioned upper bound $\hat{\rho}_0 = \rho(\mathcal{C}(A, \alpha_0))$ and $\hat{\rho}_k$ for a smaller configuration of the Steel Profile [17, 33] benchmark having matrix dimension $n = 371$. The corresponding (generalized) spectrum is contained in the negative half plane, and the whole set is chosen as the parameter set $\{\alpha_0, \dots, \alpha_{n-1}\}$. These parameters may be ordered by increasing real part (similar to `proj(2, inc)`), by decreasing real part (similar to `proj(2, dec)`), or by Penzl's heuristic (similar to `proj(2, heur)`); see page 9. Note that Penzl's heuristic [36] chooses the first shift α_0 to be the one minimizing $\rho(\mathcal{C}(A, \cdot)) = \hat{\rho}_0$, and continues greedily to select α_k by minimizing

$$\prod_{i=0}^{k-1} \frac{|\alpha_k - \alpha_i|}{|\alpha_k + \alpha_i|} \quad (45)$$

for given $\{\alpha_0, \dots, \alpha_{k-1}\}$.⁶ Quantity (45) is related to $\hat{\rho}_k$, but not the same.⁷ Consequently, the heuristic is the only one for which the corresponding $\hat{\rho}_0$ is visibly smaller than 1.

Obviously, all upper bounds $\hat{\rho}_k$ coincide for $k = n - 1$, as the underlying product of spectral radii (43) is formed over the whole set of parameters. The decreasing order clearly yields a better a priori upper bound than the increasing order. Interestingly, the heuristic order starts out smallest, remains in between the other two orders for most iterations, but comes out largest towards the last iterations. Thus, for most cases, we expect the `proj(2, dec)` shifts to require the least number of ADI steps until convergence, followed by `proj(2, heur)` and `proj(2, inc)`, in that order. Unfortunately, however, this intuition can only be confirmed for the new ADI in Tables 1 and 2.

References

- [1] H. ABOU-KANDIL, G. FREILING, V. IONESCU, AND G. JANK, *Matrix Riccati Equations in Control and Systems Theory*, Systems & Control: Foundations & Applications, Birkhäuser Basel, Basel, 2003. DOI [10.1007/978-3-0348-8081-7](https://doi.org/10.1007/978-3-0348-8081-7).
- [2] A. C. ANTOUNAS, *Approximation of Large-Scale Dynamical Systems*, Society for Industrial and Applied Mathematics, 2005. DOI [10.1137/1.9780898718713](https://doi.org/10.1137/1.9780898718713).

⁶The heuristic includes $\alpha_{k+1} := \overline{\alpha_k}$ if $\alpha_k \in \mathbb{C} \setminus \mathbb{R}$. For the present example, however, this does not occur.

⁷Note the lack of conjugation in the numerators. Furthermore, each factor containing α_i is in general only a lower bound to $\rho(\mathcal{C}(A, \alpha_i))$, which renders quantity (45) smaller than $\hat{\rho}_{k-1}$.

-
- [3] A. C. ANTOULAS, D. C. SORENSSEN, AND Y. ZHOU, *On the Decay Rate of Hankel Singular Values and Related Issues*, *Systems & Control Letters*, 46 (2002), pp. 323–342. DOI [10.1016/S0167-6911\(02\)00147-0](https://doi.org/10.1016/S0167-6911(02)00147-0).
- [4] B. BARAN, P. BENNER, AND J. SAAK, *Riccati Feedback Control of a Two-dimensional Two-phase Stefan Problem*, e-print arXiv:2209.05476, arXiv, 2022. math.NA. DOI [10.48550/arXiv.2209.05476](https://doi.org/10.48550/arXiv.2209.05476).
- [5] B. BARAN, P. BENNER, J. SAAK, AND T. STILLFJORD, *Numerical methods for closed-loop systems with non-autonomous data*, e-print arXiv:2402.13656, arXiv, 2024. math.NA. DOI [10.48550/arXiv.2402.13656](https://doi.org/10.48550/arXiv.2402.13656).
- [6] P. BENNER AND Z. BUJANOVIĆ, *On the solution of large-scale algebraic Riccati equations by using low-dimensional invariant subspaces*, *Linear Algebra Appl.*, 488 (2016), pp. 430–459. DOI [10.1016/j.laa.2015.09.027](https://doi.org/10.1016/j.laa.2015.09.027).
- [7] P. BENNER, Z. BUJANOVIĆ, P. KÜRSCHNER, AND J. SAAK, *RADI: a low-rank ADI-type algorithm for large scale algebraic Riccati equations*, *Numerische Mathematik*, 138 (2018), pp. 301–330. DOI [10.1007/s00211-017-0907-5](https://doi.org/10.1007/s00211-017-0907-5).
- [8] P. BENNER, M. HEINKENSCHLOSS, J. SAAK, AND H. K. WEICHEL, *An inexact low-rank Newton-ADI method for large-scale algebraic Riccati equations*, *Appl. Numer. Math.*, 108 (2016), pp. 125–142. DOI [10.1016/j.apnum.2016.05.006](https://doi.org/10.1016/j.apnum.2016.05.006).
- [9] P. BENNER AND P. KÜRSCHNER, *Computing Real Low-Rank Solutions of Sylvester Equations by the Factored ADI Method*, *Computers & Mathematics with Applications*, 67 (2014), pp. 1656–1672. DOI [10.1016/j.camwa.2014.03.004](https://doi.org/10.1016/j.camwa.2014.03.004).
- [10] P. BENNER, P. KÜRSCHNER, AND J. SAAK, *Real versions of low-rank ADI methods with complex shifts*, Technical report MPIMD/12-11, Max Planck Institute Magdeburg, 2012. <https://csc.mpi-magdeburg.mpg.de/preprints/2012/11/>.
- [11] ———, *A Reformulated Low-Rank ADI Iteration with Explicit Residual Factors*, *PAMM*, 13 (2013), pp. 585–586. DOI [10.1002/pamm.201310273](https://doi.org/10.1002/pamm.201310273).
- [12] ———, *An improved numerical method for balanced truncation for symmetric second-order systems*, *Mathematical and Computer Modelling of Dynamical Systems*, 19 (2013), pp. 593–615. DOI [10.1080/13873954.2013.794363](https://doi.org/10.1080/13873954.2013.794363).
- [13] ———, *Efficient Handling of Complex Shift Parameters in the Low-Rank Cholesky Factor ADI Method*, *Numerical Algorithms*, 62 (2013), pp. 225–251. DOI [10.1007/s11075-012-9569-7](https://doi.org/10.1007/s11075-012-9569-7).
- [14] ———, *Self-Generating and Efficient Shift Parameters in ADI Methods for Large Lyapunov and Sylvester Equations*, *Electron. Trans. Numer. Anal.*, 43 (2014), pp. 142–162. <http://etna.mcs.kent.edu/volumes/2011-2020/vol143/abstract.php?vol=43&pages=142-162>.
- [15] P. BENNER, J.-R. LI, AND T. PENZL, *Numerical solution of large-scale Lyapunov equations, Riccati equations, and linear-quadratic optimal control problems*, *Numerical Linear Algebra with Applications*, 15 (2008), pp. 755–777. DOI [10.1002/nla.622](https://doi.org/10.1002/nla.622).
- [16] P. BENNER, R.-C. LI, AND N. TRUHAR, *On the ADI method for Sylvester equations*, *Journal of Computational and Applied Mathematics*, 233 (2009), pp. 1035–1045. DOI [10.1016/j.cam.2009.08.108](https://doi.org/10.1016/j.cam.2009.08.108).
- [17] P. BENNER AND J. SAAK, *Linear-Quadratic Regulator Design for Optimal Cooling of Steel Profiles*, techreport SFB393/05-05, Sonderforschungsbereich 393 *Parallele Numerische Simulation für Physik und Kontinuumsmechanik*, TU Chemnitz, D-09107 Chemnitz (Germany), 2005. <http://nbn-resolving.de/urn:nbn:de:swb:ch1-200601597>.
- [18] ———, *Numerical Solution of Large and Sparse Continuous Time Algebraic Matrix Riccati and Lyapunov Equations: A State of the Art Survey*, *GAMM-Mitteilungen*, 36 (2013), pp. 32–52. DOI [10.1002/gamm.201310003](https://doi.org/10.1002/gamm.201310003).

-
- [19] C. BERTRAM AND H. FASSBENDER, *Riccati ADI: Existence, Uniqueness and New Iterative Methods*, techreport arXiv:2004.11212, 2020. <http://arxiv.org/abs/2004.11212>.
- [20] ———, *A Link Between Gramian-Based Model Order Reduction and Moment Matching*, in *Model Reduction of Complex Dynamical Systems*, P. Benner, T. Breiten, H. Faßbender, M. Hinze, T. Stykel, and R. Zimmermann, eds., vol. 171, Springer International Publishing, Cham, 2021, pp. 119–139. DOI [10.1007/978-3-030-72983-7_6](https://doi.org/10.1007/978-3-030-72983-7_6).
- [21] ———, *A Quadrature Framework for Solving Lyapunov and Sylvester Equations*, *Linear Algebra and its Applications*, 622 (2021), pp. 66–103. DOI [10.1016/j.laa.2021.03.029](https://doi.org/10.1016/j.laa.2021.03.029).
- [22] J. BEZANSON, A. EDELMAN, S. KARPINSKI, AND V. B. SHAH, *Julia: A Fresh Approach to Numerical Computing*, *SIAM Review*, 59 (2017), pp. 65–98. DOI [10.1137/141000671](https://doi.org/10.1137/141000671).
- [23] R. S. DEMBO, S. C. EISENSTAT, AND T. STEIHAUG, *Inexact Newton Methods*, *SIAM Journal on Numerical Analysis*, 19 (1982), pp. 400–408. DOI [10.1137/0719025](https://doi.org/10.1137/0719025).
- [24] G. H. GOLUB AND C. F. VAN LOAN, *Matrix Computations*, Johns Hopkins University Press, 4 ed., 2013.
- [25] L. GRASEDYCK, *Existence of a Low Rank or \mathcal{H} -matrix Approximant to the Solution of a Sylvester Equation*, *Numerical Linear Algebra with Applications*, 11 (2004), pp. 371–389. DOI [10.1002/nla.366](https://doi.org/10.1002/nla.366).
- [26] S. GUGERCIN AND J.-R. LI, *Smith-Type Methods for Balanced Truncation of Large Sparse Systems*, in *Dimension Reduction of Large-Scale Systems*, P. Benner, D. C. Sorensen, and V. Mehrmann, eds., vol. 45, Springer-Verlag, Berlin/Heidelberg, 2005, pp. 49–82. DOI [10.1007/3-540-27909-1_2](https://doi.org/10.1007/3-540-27909-1_2).
- [27] W. HACKBUSCH, *Iterative Solution of Large Sparse Systems of Equations*, vol. 95 of *Applied Mathematical Sciences*, 2016. DOI [10.1007/978-3-319-28483-5](https://doi.org/10.1007/978-3-319-28483-5).
- [28] I. M. JAIMOUKHA AND E. M. KASENALLY, *Krylov Subspace Methods for Solving Large Lyapunov Equations*, *SIAM Journal on Numerical Analysis*, 31 (1994), pp. 227–251. DOI [10.1137/0731012](https://doi.org/10.1137/0731012).
- [29] D. KRESSNER, K. LUND, S. MASSEI, AND D. PALITTA, *Compress-and-restart Block Krylov Subspace Methods for Sylvester Matrix Equations*, *Numerical Linear Algebra with Applications*, 28 (2021), p. e2339. DOI [10.1002/nla.2339](https://doi.org/10.1002/nla.2339).
- [30] P. KÜRSCHNER, *Efficient low-rank solution of large-scale matrix equations*, Ph.D. thesis, Otto-von-Guericke-Universität Magdeburg, 2016. <https://hdl.handle.net/11858/00-001M-0000-0029-CE18-2>.
- [31] N. LANG, H. MENA, AND J. SAAK, *On the benefits of the LDL^T factorization for large-scale differential matrix equation solvers*, *Linear Algebra and its Applications*, 480 (2015), pp. 44–71. DOI [10.1016/j.laa.2015.04.006](https://doi.org/10.1016/j.laa.2015.04.006).
- [32] J.-R. LI AND J. WHITE, *Low rank solution of Lyapunov equations*, *SIAM Journal on Matrix Analysis and Applications*, 24 (2002), pp. 260–280. DOI [10.1137/S0895479801384937](https://doi.org/10.1137/S0895479801384937).
- [33] OBERWOLFACH BENCHMARK COLLECTION, *Steel Profile*. hosted at MORwiki – Model Order Reduction Wiki, 2005. http://modelreduction.org/index.php/Steel_Profile.
- [34] D. PALITTA AND V. SIMONCINI, *Computationally Enhanced Projection Methods for Symmetric Sylvester and Lyapunov Matrix Equations*, *Journal of Computational and Applied Mathematics*, 330 (2018), pp. 648–659. DOI [10.1016/j.cam.2017.08.011](https://doi.org/10.1016/j.cam.2017.08.011).
- [35] D. W. PEACEMAN AND H. H. RACHFORD, JR, *The numerical solution of parabolic and elliptic differential equations*, *Journal of the Society for industrial and Applied Mathematics*, 3 (1955), pp. 28–41. DOI [10.1137/0103003](https://doi.org/10.1137/0103003).

-
- [36] T. PENZL, *A Cyclic Low-Rank Smith Method for Large Sparse Lyapunov Equations*, SIAM Journal on Scientific Computing, 21 (1999), pp. 1401–1418. DOI [10.1137/S1064827598347666](https://doi.org/10.1137/S1064827598347666).
- [37] ———, *Eigenvalue Decay Bounds for Solutions of Lyapunov Equations: The Symmetric Case*, Systems & Control Letters, 40 (2000), pp. 139–144. DOI [10.1016/S0167-6911\(00\)00010-4](https://doi.org/10.1016/S0167-6911(00)00010-4).
- [38] J. SAAK, *Efficient Numerical Solution of Large Scale Algebraic Matrix Equations in PDE Control and Model Order Reduction*, 2009. <http://nbn-resolving.de/urn:nbn:de:bsz:ch1-200901642>.
- [39] J. SABINO, *Solution of Large-Scale Lyapunov Equations via the Block Modified Smith Method*, Ph.D. thesis, Rice University, 2007. <https://hdl.handle.net/1911/20641>.
- [40] J. SCHULZE, *A Low-Rank Parareal Solver for Differential Riccati Equations Written in Julia*, 2022. DOI [10.5281/zenodo.7843198](https://doi.org/10.5281/zenodo.7843198).
- [41] V. SIMONCINI, *A New Iterative Method for Solving Large-Scale Lyapunov Matrix Equations*, SIAM Journal on Scientific Computing, 29 (2007), pp. 1268–1288. DOI [10.1137/06066120X](https://doi.org/10.1137/06066120X).
- [42] ———, *Computational Methods for Linear Matrix Equations*, SIAM Review, 58 (2016), pp. 377–441. DOI [10.1137/130912839](https://doi.org/10.1137/130912839).
- [43] T. STILLFJORD, *Singular Value Decay of Operator-Valued Differential Lyapunov and Riccati Equations*, SIAM J. Control Optim., 56 (2018), pp. 3598–3618. DOI [10.1137/18M1178815](https://doi.org/10.1137/18M1178815).
- [44] H. T. STOPPELS AND L. NYMAN, *ArnoldiMethod.jl: Arnoldi Method with Krylov-Schur restart, natively in Julia*. <https://github.com/JuliaLinearAlgebra/ArnoldiMethod.jl>.
- [45] N. TRUHAR AND K. VESELIĆ, *Bounds on the Trace of a Solution to the Lyapunov Equation with a General Stable Matrix*, Systems & Control Letters, 56 (2007), pp. 493–503. DOI [10.1016/j.sysconle.2007.02.003](https://doi.org/10.1016/j.sysconle.2007.02.003).
- [46] E. WACHSPRESS, *The ADI Model Problem*, Springer New York, New York, NY, 2013. DOI [10.1007/978-1-4614-5122-8](https://doi.org/10.1007/978-1-4614-5122-8).
- [47] T. WOLF AND H. K. F. PANZER, *The ADI Iteration for Lyapunov Equations Implicitly Performs H2 Pseudo-Optimal Model Order Reduction*, International Journal of Control, 89 (2016), pp. 481–493. DOI [10.1080/00207179.2015.1081985](https://doi.org/10.1080/00207179.2015.1081985).