

Semiparametric Localized Principal Stratification Analysis with Continuous Strata

Yichi Zhang* and Shu Yang^{†‡}

Abstract

Principal stratification is essential for revealing causal mechanisms involving post-treatment intermediate variables. Principal stratification analysis with continuous intermediate variables is increasingly common but challenging due to the infinite principal strata and the nonidentifiability and nonregularity of principal causal effects. Inspired by recent research, we resolve these challenges by first using a flexible copula-based principal score model to identify principal causal effect under weak principal ignorability. We then target the local functional substitute of principal causal effect, which is statistically regular and can accurately approximate principal causal effect with vanishing bandwidth. We simplify the full efficient influence function of the local functional substitute by considering its oracle-scenario alternative. This leads to a computationally efficient and straightforward estimator for the local functional substitute and principal causal effect with vanishing bandwidth. We prove the double robustness and statistical optimality of our proposed estimator, and derive its asymptotic normality for inferential purposes. We illustrate the appealing statistical performance of our proposed estimator in simulations, and apply it to two real datasets with intriguing scientific discoveries.

Keywords: Causal effect predictiveness surface; Double robustness; Local functional; Semiparametric efficiency; Surrogate evaluation

1 Introduction

When an intermediate variable exists between treatment and outcome, identifying and inferring treatment effects becomes more challenging compared to the classical “covariate-treatment-outcome” mechanism. Principal strata, defined by the joint potential values of the intermediate variable under both treatment and control, offer a principled way to understand such underlying causal mechanisms (Frangakis and Rubin, 2002). Principal stratification involves comparing causal effects between key strata or groupings of the data, often referred to as principal causal effects (PCEs). PCEs have been

*Department of Biostatistics and Bioinformatics & Fuqua School of Business, Duke University, Durham, North Carolina 27708, U.S.A. Email: yz847@duke.edu

[†]Department of Statistics, North Carolina State University, Raleigh, North Carolina 27607, U.S.A. Email: syang24@ncsu.edu

[‡]Corresponding author.

useful in various applications, such as noncompliance and truncation-by-death problems in clinical trials (Frangakis and Rubin, 2002, Rubin, 2006), surrogate evaluation in vaccinology (Rubin, 2004, Gilbert and Hudgens, 2008), and causal mediation analysis (Kim et al., 2019).

Like potential outcomes, potential intermediate variable values suffer from inherent missing data because at most one potential value is observable. Therefore, additional assumptions are necessary to identify PCEs. In this paper, we consider *weak principal ignorability* (Jo and Stuart, 2009), an ignorability-type assumption for the intermediate variable analogous to the celebrated treatment ignorability assumption (Rosenbaum and Rubin, 1983). Principal ignorability has been widely adopted in various applications (Follmann, 2000, Hill et al., 2002, Jo and Stuart, 2009, Stuart et al., 2015, Jiang et al., 2022). For a binary intermediate variable, Jiang et al. (2022) establish various identification formulas for PCEs under principal ignorability and an additional monotonicity assumption and derive the semiparametric efficient influence function (EIF). The resulting PCE estimator exhibits desirable properties, e.g., triple robustness and semiparametric efficiency.

1.1 From a binary to continuous intermediate variable

In contrast to the noncompliance and truncation-by-death problems, one often encounters a continuous intermediate variable in surrogate evaluation and mediation analysis. For example, Follmann (2006) considers the immune response measured after randomization as the continuous surrogate outcome for infection in vaccine trials. Schwartz et al. (2011) assess the mediation effect of the continuous body mass index on cardiovascular disease. Zigler and Belin (2012) evaluate the surrogacy of the continuous short-term CD4 count for a long-term clinical endpoint in AIDS clinical trials. The identification and estimation of PCE with a continuous intermediate variable (Cont.PCE) are challenging due to infinitely many principal strata generated by a continuous intermediate variable. Early attempts identify and estimate Cont.PCE through parametric models (Gilbert and Hudgens, 2008, Jin and Rubin, 2008). Later works consider more flexible semiparametric models of continuous principal strata (see, e.g., Bartolucci and Grilli (2011), Schwartz et al. (2011), Zigler and Belin (2012)). However, these methods still depend on correctly specifying the entire underlying distribution (cf. Section 5).

This paper aims to develop an efficient and flexible semiparametric framework for the identification and estimation of Cont.PCEs under the weak principal ignorability. This work can be seen as an extension of Jiang et al. (2022), which, however, is non-trivial due to challenges with respect to estimand, identification, and estimation:

- (i). The Cont.PCE is a non-pathwise differentiable estimand, making classical semiparametric efficiency theory inapplicable for deriving a semiparametric efficient estimator.
- (ii). The monotonicity assumption used to identify PCE in Jiang and Ding (2021) fails to identify PCE when the intermediate variable is continuous and may be implausible in various applications such as surrogate analysis.

(iii). After addressing (i) and (ii) properly, the resulting full EIF (cf. Section 3) is still too complex to construct a computationally feasible estimator.

1.2 From a perfectly localized estimand to local estimand

The non-pathwise differentiability, as mentioned in (i) above, presents a significant obstacle in developing an efficient estimator for Cont.PCE. To resolve this type of challenge for efficient statistical estimation, Chernozhukov et al. (2022, 2023) focus on a specific non-pathwise differentiable estimand, namely, the *perfectly localized functional* ϑ_0 ,

$$\vartheta_0 = \lim_{h \rightarrow 0} \vartheta_h, \quad \vartheta_h = \mathbb{E}\{m(W, Y, \gamma^*, k_{h,d_0})\}, \quad (1)$$

where ϑ_h is the *local functional* with a fixed and sufficiently small $h > 0$, which is pathwise differentiable.¹ Here Y is the response, W includes other variables, $\gamma^*(\cdot) = \mathbb{E}(Y \mid W = \cdot)$, $m(w, y, \gamma^*, k_{h,d_0})$ is a linear functional of $\gamma^*(\cdot)$, and $k_{h,d_0}(\cdot)$ is a pre-specified smoothing function centered at a specific value $D = d_0$ in W with bandwidth $h > 0$. Examples of ϑ_0 include the conditional average treatment effect and continuous treatment effect. This paper generalizes the original definition of ϑ_0 by allowing W to contain unobserved variables and proves that Cont.PCE is also a perfectly localized functional under proper identification assumptions (cf. Proposition 1).

Instead of directly estimating the perfectly localized functional, Chernozhukov et al. (2022, 2023) alternatively estimate its substitute, the local functional ϑ_h in (1) with small $h > 0$. They further propose a doubly robust estimator using Riesz representation. Switching from ϑ_0 to ϑ_h as the target estimand has multiple advantages, as discussed in Chernozhukov et al. (2022, 2023): ϑ_h is pathwise differentiable and can be estimated semiparametrically efficiently; ϑ_h serves as a good approximation of ϑ_0 when h is small, thus efficient estimation of ϑ_h translates to efficient estimation of ϑ_0 up to a small localization bias; and inference on ϑ_h is more robust and does not rely on the localization bias. Following Chernozhukov et al. (2022, 2023), this paper aims to construct a computationally and statistically efficient estimator for the local functional substitute of Cont.PCE (Loc.PCE), and subsequently, estimate Cont.PCE efficiently and accurately as $h \rightarrow 0$ with growing sample size. However, the paper deals with a more complex setting than Chernozhukov et al. (2022, 2023) due to the unobserved variable in W and an additional nuisance function. To address these challenges, a new EIF-invoked estimation strategy is proposed for Loc.PCE. The recent work of Lu et al. (2023) adopts a different strategy by using a global approximation of the underlying Cont.PCE surface, focusing on a finite-dimensional parameter within a working model. This enables them to derive the corresponding EIF and doubly robust estimators. In contrast, we directly target the localized Cont.PCE and Cont.PCE (c.f. Section 5).

¹Notably, Chernozhukov et al. (2023) terms ϑ_0 as the “local functional” and ϑ_h as the “global functional”. This paper adopts the terminology in Chernozhukov et al. (2022, Definition 2.2) to ensure consistence.

1.3 Outline and contributions

The paper is organized as follows. Section 2.1 provides the background for PCE estimation. Section 2.2 motivates Cont.PCE estimation using a synthetic example in surrogate analysis, while Section 2.3 presents the identification assumptions and formula for Cont.PCE. Section 2.4 introduces Loc.PCE, the local functional substitute of Cont.PCE, as the target estimand for which we develop a semiparametric efficient estimator. The identification and estimation of Cont.PCE and Loc.PCE involve three nuisance models:

\mathcal{M}_{tp} : *treatment probability*, the distribution of treatment given covariates;

\mathcal{M}_{ps} : *principal score*, the distribution of intermediate variable given covariates & treatment;

\mathcal{M}_{om} : *outcome mean*, the conditional expectation of outcome given all other variables.

In Section 3, we derive the EIF of Loc.PCE under nonparametric nuisance functions. However, the resulting EIF is too complex for a computationally viable estimator. Thus, we derive simplified EIFs in two special scenarios: (i) the parametric scenario where \mathcal{M}_{ps} is parametrically modeled, and (ii) the oracle scenario where \mathcal{M}_{ps} is known. We construct our proposed estimator using the oracle-scenario EIF, which is computationally feasible. In Sections 4.1 and 4.2, we show that the proposed estimator is doubly robust and asymptotically normal for Loc.PCE and Cont.PCE. Interestingly, our estimator attains the semiparametric efficiency bound for Loc.PCE, under the more general condition when \mathcal{M}_{ps} is parametric, despite being derived using the EIF assuming \mathcal{M}_{ps} is known. We can also construct confidence intervals (CIs) for the inference of Loc.PCE and Cont.PCE. In Section 6, simulation results corroborate our theoretical results. In Section 7, we apply the proposed estimator to two real-world problems in surrogate analysis and causal pathway discovery, and in Section 8 we provide concluding remarks.

2 Principal stratification with continuous strata

2.1 Basic setups and Cont.PCE

Suppose we have n i.i.d. observations $\{V_i = (X_i, Z_i, M_i, Y_i)\}_{i=1}^n$. For each sample V_i , we denote the pre-treatment covariates by $X_i \in \mathbb{X} \subseteq \mathbb{R}^{d_x}$, treatment assignment by $Z_i \in \{0, 1\}$, intermediate variable by $M_i \in \mathbb{M} \subseteq \mathbb{R}$, and outcome by $Y_i \in \mathbb{Y} \subseteq \mathbb{R}$. Our data generation mechanism invokes the Stable Unit Treatment Value Assumption (Rubin, 1980). In particular, we assume

$$(X_i, Z_i, M_{1i}, M_{0i}, Y_{1i}, Y_{0i}) \stackrel{\text{i.i.d.}}{\sim} (X, Z, M_1, M_0, Y_1, Y_0),$$

where M_{1i} and M_{0i} are potential intermediate variables, and Y_{1i} and Y_{0i} are potential outcomes, such that $M_i = Z_i M_{1i} + (1 - Z_i) M_{0i}$ and $Y_i = Z_i Y_{1i} + (1 - Z_i) Y_{0i}$.

Frangakis and Rubin (2002) use the joint potential values of intermediate variables to define the principal stratum $U = (M_1, M_0)$. A stratum U captures the information on how the intermediate

variable is affected by the treatment. When M is a binary variable, principal stratum U takes values in $\{(0, 0), (1, 0), (0, 1), (1, 1)\}$, which in noncompliance problems (Frumento et al., 2012) are referred to the never-taker, complier, defier, and always-taker, respectively. When M is a continuous variable, stratum U takes values in \mathbb{R}^2 . In surrogate analysis, M_1 and M_0 are the counterfactual surrogate outcomes had treatments been set to 1 and 0, respectively, and $M_1 - M_0$ is the surrogate treatment effect.

For a specific principal stratum $u^* = (m_1^*, m_0^*) \in \mathbb{R}^2$, define PCE as the average causal effect on the primary outcome Y within the principal stratum $U = u^*$,

$$\tau_{u^*}^* = \mathbb{E}(Y_1 - Y_0 \mid U = u^*).$$

Referring to Jiang et al. (2022, § 2), $\tau_{u^*}^*$ can have different scientific meanings when M is binary. For continuous M , it is the Cont.PCE as we alluded earlier. In the next section, we demonstrate the practical utility of Cont.PCE in evaluating surrogate outcomes (Frangakis and Rubin, 2002).

2.2 Interpretation of Cont.PCE in surrogate analysis

In clinical trials, obtaining primary clinical outcomes can be time-consuming and expensive. As a result, there is considerable interest in identifying an easy-to-obtain surrogate that can reliably reflect treatment effects on the primary clinical outcome. Cont.PCE, also known as the causal effect predictiveness (CEP) surface (Gilbert and Hudgens, 2008), can provide a comprehensive evaluation of any tested surrogate. In particular, Frangakis and Rubin (2002) suggest evaluating the associative and dissociative effects of a surrogate outcome, both of which can be quantified through Cont.PCE.

- A good surrogate is required to meet *causal sufficiency*, indicating that any causal influence of the treatment on the surrogate must be accompanied by an effect on the primary outcome (Gilbert and Hudgens, 2008, § 3.1). The associative effect compares the average difference between Y_1 and Y_0 when $M_1 = m_1 \neq M_0 = m_0$, which is quantified by $\tau_{(m_1, m_0)}^*$ for any two distinct principal levels $m_1, m_0 \in \mathbb{R}$. Therefore, causal sufficiency is satisfied *if* all associative effects $\tau_{(m_1, m_0)}^* \neq 0$ when $m_1 \neq m_0$.
- Another criterion a good surrogate shall satisfy is *causal necessity*, which entails that the absence of a causal impact of the treatment on the surrogate outcome must also imply the absence of any effect on the primary outcome (Gilbert and Hudgens, 2008, § 3.1). The dissociative effect compares the average difference between Y_1 and Y_0 when $M_1 = M_0 = m$, which is quantified through $\tau_{(m, m)}^*$ for any principal level $m \in \mathbb{R}$. Causal necessity can be true *only if* dissociative effects $\tau_{(m, m)}^* = 0$ for all $m \in \mathbb{R}$. In another word, if one observes some $\tau_{(m, m)}^* \neq 0$, M is evaluated as a surrogate not satisfying causal necessity.

We use synthetic examples to demonstrate the usage of Cont.PCE, in which we denote $\Sigma_0 = \begin{pmatrix} 1 & 0.25 \\ 0.25 & 1 \end{pmatrix}$ and generate $X = (X_1, X_0) \sim \mathcal{N}\{(0, 0)^\top, \Sigma_0\}$ and $Z \sim \text{Bern}(1/2)$, independently. Given




Figure 1: Cont.PCEs under Procedure (i) and (ii). The X- and Y- axes represent m_1 and m_0 , respectively, and Z-axis represents the corresponding τ_u^* . The orange dashed line represents $\tau_u^* = 0$ when $m_1 = m_0$.

X , we consider two different generating procedures of (M_1, M_0) :

$$\text{P(i). } (M_1, M_0) \sim \mathcal{N} \left\{ (X_1, X_0)^\top, \Sigma_0 \right\}; \quad \text{P(ii). } (M_1, M_0) \sim \mathcal{N} \left\{ (0, 0)^\top, \Sigma_0 \right\}.$$

Finally, we generate the potential outcomes. To facilitate the mediation analysis, we generalize the single-indexed potential outcome notation Y_z to double-indexed $Y_{z,m}$. Under the background of surrogate analysis, in principle, the post-treatment surrogate outcome M can be regarded as an additional continuous treatment that has causal effect on the primary outcome Y . Conceptually, $Y_{z,m}$ is the potential clinical outcome if the treatment Z and surrogate outcome M were simultaneously manipulated to some levels z and m , respectively (Robins and Greenland, 1992, Mealli and Mattei, 2012, Baccini et al., 2017). The single-indexed and double-indexed potential outcome notations can be linked through $Y_z = Y_{z,M_z}$ for $z \in \{0, 1\}$ (Forastiere et al., 2018, § 2.1). For each $z \in \{0, 1\}$ and $m \in \mathbb{R}$, with given X , we generate the potential outcome $Y_{z,m}$ independently according to the following distributions:

$$\text{P(i). } Y_{z,m} \sim m/2 + \mathcal{N}(X_z + 0.5X_{1-z}, 1); \quad \text{P(ii). } Y_{z,m} \sim \mathcal{N}(X_z + 0.5X_{1-z}, 1).$$

Figure 1 visualizes the CEP surfaces in P(i) where $\tau_u^* = 0.75(m_1 - m_0)$ and in P(ii) where $\tau_u^* = 0$. Therefore, P(i) and P(ii) generate two types of surrogates: in P(i), $M_1 - M_0$ has a positive correlation with $Y_1 - Y_0$, and M is both causally necessary and sufficient to be a good surrogate for Y , while in P(ii) M has no association with Y and thus is uninformative.

Mediation analysis is another widely recognized approach for handling a continuous intermediate variable. Through our synthesized examples, we demonstrate that the conventional mediation analysis may not be suitable for evaluating the surrogate and, thus, highlight the need for Cont.PCE estimation. Mediation analysis focuses on how M plays a role in the causal pathway $Z \rightarrow M \rightarrow Y$. The natural indirect effect (NIE) and natural direct effect (NDE) are $\text{NIE} = \mathbb{E}(Y_{1,M_1} - Y_{1,M_0})$ and $\text{NDE} = \mathbb{E}(Y_{1,M_0} - Y_{0,M_0})$, respectively. By fixing Z at the treatment level ($Z = 1$), NIE

measures the average causal effect of counterfactual mediators. By fixing the mediator at the control level ($M = M_0$), NDE measures the average causal effect of treatments (Imai et al., 2010). In the synthetic examples, we can deduce that $Y_{1,M_1} - Y_{1,M_0} \sim \mathcal{N}\{(M_1 - M_0)/2, 2\}$ in P(i) and $Y_{1,M_1} - Y_{1,M_0} \sim \mathcal{N}(0, 2)$ in P(ii), while $Y_{1,M_0} - Y_{0,M_0} \sim \mathcal{N}\{(X_1 - X_0)/2, 2\}$ in both P(i) and P(ii). Then by the zero means of (X_1, X_0) and (M_1, M_0) , we have NIE = NDE = 0 in both P(i) and P(ii), which make NIE and NDE uninformative for the surrogate evaluation of M . However, the strong and positive correlation between $M_1 - M_0$ and $Y_1 - Y_0$ actually makes M a good surrogate in P(i), which can be captured by the Cont.PCE surface.

Remark 1 *The mediation analysis and principal stratification are generally not comparable due to their conceptual difference (Rubin, 2004, VanderWeele, 2011, Lipkovich et al., 2022). When the intermediate variable cannot be intervened, or causal pathway $Z \rightarrow M \rightarrow Y$ does not exist, the principal stratification is a more reasonable framework to deal with intermediate variables (Rubin, 2004, VanderWeele, 2011). In mediation analysis, Cont.PCE can also be used to evaluate the direct effect; Non-zero $\tau_{(m,m)}^*$ for some $m \in \mathbb{R}$, imply that Z affects Y through some causal pathways other than $Z \rightarrow M \rightarrow Y$ (Mattei and Mealli, 2011). See Section 7.2 for a real-data illustration.*

Remark 2 *Another branch of research analyzes the surrogate variable through the estimation of proportion of the treatment effect (PTE) (Wang and Taylor, 2002, Parast et al., 2016, Agniel and Parast, 2021, Wang et al., 2020, Han et al., 2022). PTE summarizes how much of the treatment effect on Y is explained by its effect on M , offering a different perspective on surrogacy. We compare the identification assumptions of these works with ours in Remark 3.*

Additional notations Before we present our identification results, we introduce some new notations for ease of exposition. Denote $f(V = v)$ the density function of $V = (X, Z, M, Y)$. Define $\pi_z(x) = \mathbb{P}(Z = z | X = x)$ as the treatment probability, also known as the propensity score, $e_u(x) = f(U = u | X = x)$ the generalized principal score, which is the density function of principal stratum u given the covariates, and $\mu_{zm}(x) = \mathbb{E}(Y | X = x, Z = z, M = m)$ the outcome mean. When there is no conflict of notations, we abbreviate $\pi_z(x)$ and $\mu_{zm}(x)$ as π and μ . Denote $\|\cdot\|_\infty$ as the functional sup-norm. Let $e_u = \mathbb{E}\{e_u(X)\} = f(U = u)$ denote the marginalized principal score. We use $f(\cdot)$ and $F(\cdot)$ to represent the probability density function (PDF) and the conditional cumulative distribution function (CDF), respectively. We also denote $f_{zm}(x) = f(M = m | X = x, Z = z)$ and $F_{zm}(x) = \mathbb{P}(M \leq m | X = x, Z = z)$.

2.3 Nonparametric identification

We first consider the treatment ignorability for both the intermediate variable and outcome, which is an extension of the classic treatment ignorability (Rubin, 1974).

Assumption 1 (Treatment ignorability) $Z \perp\!\!\!\perp (M_0, M_1, Y_0, Y_1) | X$.

Assumption 1 precludes unobservable confounders that affect both the treatment and intermediate variable and those that impact both the treatment and outcome. In observational studies, its

plausibility relies on whether or not observed covariates include all the confounders that affect the treatment as well as the outcome and intermediate variables. Next, we extend the weak principal ignorability assumption (Jiang et al., 2022) to the continuous- M scenario. Weak principal ignorability is an ignorability-type assumption for the principal strata similar to the treatment ignorability.

Assumption 2 (Weak principal ignorability) $\mathbb{E}(Y_z | X, U = u) = \mathbb{E}(Y_z | X, M_z = m_z)$ for any $z \in \{0, 1\}$ and $u = (m_1, m_0) \in \mathbb{R}^2$.

Assumption 2 is a relaxed, expectation-based version of the general weak principal ignorability (see, e.g., Mattei et al. (2023, Assumption 12)) that assumes $Y_z \perp\!\!\!\perp M_{1-z} | X, M_z, \forall z \in \{0, 1\}$. Briefly, weak principal ignorability implies that given X and M_1 (resp., M_0), the distribution of Y_1 (resp., Y_0) does not depend on the intermediate variable under the opposite treatment M_0 (resp., M_1) (Mattei et al., 2023). Assumption 2 is generally weaker than such implication. In particular, it requires that, for both $z = 0$ and $z = 1$, the expected value of the potential outcome Y_z should be the same across principal strata when the value of M_z is fixed at a specific level $m_z \in \mathbb{R}$, and M_{1-z} can vary over any level $m_{1-z} \in \mathbb{R}$, given observed covariates. Under Assumption 1, Assumption 2 is equivalent to

$$\mathbb{E}(Y_z | X, Z = z, U = u) = \mathbb{E}(Y_z | X, Z = z, M = m_z), \text{ for any } z \in \{0, 1\} \text{ and } (m_1, m_0) \in \mathbb{R}^2, \quad (2)$$

which further implies that $\mathbb{E}(Y_z | X, U = u)$ can be simplified to an observable conditional expectation $\mathbb{E}(Y | X, Z = z, M = m_z)$. The plausibility of weak principal ignorability depends on the problem context and whether X captures all the Z - Y and M - Y confounders. For the surrogate analysis, the surrogate outcome under the opposite treatment does not have a direct causal effect on the primary outcome (Parast et al., 2016), then weak principal ignorability is plausible if X collects all the confounders.

Remark 3 *In mediation analysis, the strong principal ignorability (Forastiere et al., 2018) and sequential ignorability (Imai et al., 2010) are two standard assumptions for the identification of causal mediation effects. Some other surrogate identifying conditions have been proposed in Parast et al. (2016), Wang et al. (2020) for PTE estimation, under the randomized clinical trial (RCT). All assumptions above, however, directly imply and thus are stronger than Assumption 2; see Section S2 in the Supplementary File for detailed discussions.*

By Assumption 1, the conditional density of M_z given X can be identified as

$$f(M_z = m_z | X = x) = f(M = m_z | X = x, Z = z),$$

for any $x \in \mathbb{R}^{d_x}, z \in \{0, 1\}, m_z \in \mathbb{R}$. However, the joint density $e_u(x) = f(M_1 = m_1, M_0 = m_0 | X = x)$ is still not identifiable due to the fundamental problem that M_1 and M_0 cannot be jointly observed. Moreover, the monotonicity assumption, as used in the binary- M scenario (Jiang et al., 2022), is inadequate to identify $e_u(x)$ in the continuous- M scenario. To overcome

this issue, we propose a copula-based approach (Bartolucci and Grilli, 2011, Kim et al., 2020) to model the association between M_1 and M_0 conditional on X . Our approach encompasses several popular association models, including the monotonicity and equipercentile equating model (Efron and Feldman, 1991, Jin and Rubin, 2008) as special cases; see, e.g., Jiang and Ding (2021, § 4.1) and Nelsen (2007).

Assumption 3 (Copula models of $e_u(x)$) Let $F(M_z = \cdot | X = \star)$ be the CDF of M_z given X , $c_\rho(\cdot, \cdot)$ be any specified copula function, and ρ be the correlation between M_1 and M_0 given X . Assume $e_u(x) = f(M_1 = m_1, M_0 = m_0 | X = x) = c_\rho\{F(M_1 = m_1 | X = x), F(M_0 = m_0 | X = x)\} \prod_{z \in \{0,1\}} f(M_z = m_z | X = x)$, for any $x \in \mathbb{R}^{d_x}$, $z \in \{0, 1\}$, $u = (m_1, m_0) \in \mathbb{R}^2$.

Sklar's theorem (Sklar, 1959) guarantees any regular $e_u(x)$ can be represented via Assumption 3 with correctly specified $c_\rho(\cdot, \cdot)$. Some popular models of $c_\rho(\cdot, \cdot)$ include Gaussian copula, Clayton copula, and Farli-Gumbel-Morgenstern copula (Jaworski et al., 2010). We assume the copula function $c_\rho(\cdot, \cdot)$ and correlation parameter ρ are correctly specified. In practice, one can conduct sensitivity analysis for various combinations of $c_\rho(\cdot, \cdot)$ and ρ .

The following theorem provides the identification result for Cont.PCE.

Theorem 1 (Identification of Cont.PCE) Under Assumptions 1 and 2, Cont.PCE can be expressed as

$$\tau_{u^*} = \mathbb{E}\{\mu_{1m_1^*}(X) - \mu_{0m_0^*}(X) | U = u^*\}. \quad (3)$$

Suppose Assumptions 1–3 hold. Given any principal level $u^* = (m_1^*, m_0^*) \in \mathbb{R}^2$ with $e_{u^*} > 0$, τ_{u^*} can be identified as

$$\tau_{u^*} = \frac{\mathbb{E}[\{\mu_{1m_1^*}(X) - \mu_{0m_0^*}(X)\}e_{u^*}(X)]}{e_{u^*}}, \quad (4)$$

where $e_{u^*}(X) = c_\rho\{F_{1m_1^*}(X), F_{0m_0^*}(X)\} \prod_{z \in \{0,1\}} f_{zm_z^*}(X)$.

In Theorem 1, we emphasize that τ_{u^*} in (3) is defined by conditioning on the unobserved U . This distinguishes our problem from other localized functional estimation problems discussed by Chernozhukov et al. (2022, 2023). For example, while CATE can be expressed through conditional outcome means similar to (3), it conditions on an observable variable X . This highlights the need to model the joint distribution of U in Assumption 3 for τ_{u^*} identification. In the identification formula (4), under the treatment and weak principal ignorability, we have $\mu_{1m_1^*}(X) - \mu_{0m_0^*}(X) = \mathbb{E}(Y_1 - Y_0 | X, U = u^*)$, which is the conditional Cont.PCE given X . The weight $e_{u^*}(X)/e_{u^*}$, by Bayes' Theorem, is the density ratio of X conditional and unconditional on $U = u^*$; Here e_{u^*} is the normalizing constant of $e_{u^*}(X)$, and they can be identified by Assumptions 1 and 3. Therefore, our identification formula marginalizes the conditional Cont.PCE over the distribution of X given $U = u^*$, and thereby giving Cont.PCE within stratum $U = u^*$.

2.4 Target estimand: from Cont.PCE to Loc.PCE

We show that Cont.PCE is a perfectly localized estimand (Chernozhukov et al., 2022). Specifically, there exists a sequence of local estimands indexed by $h > 0$, i.e., Loc.PCE (cf. (5)), that converges

to Cont.PCE as $h \rightarrow 0$. Define the normalized kernel function centered at $u^* \in \mathbb{R}^2$ as $k_{u^*}(u) = k_{h, \mathcal{K}(\cdot, \cdot), u^*}\{u = (m_1, m_0)\} = h^{-2} \mathcal{K}\{h^{-1}(m_1 - m_1^*), h^{-1}(m_0 - m_0^*)\}$, where $\mathcal{K}(\cdot, \cdot)$ is any pre-specified two-dimensional kernel function and $h > 0$ is a fixed bandwidth. Define Loc.PCE as

$$\tau_{u^*} = \tau_{u^*}\{\mathcal{K}(\cdot, \cdot), h\} = \frac{\mathbb{E}[\{\mu_{1M_1}(X) - \mu_{0M_0}(X)\}k_{u^*}(U)]}{\mathbb{E}\{k_{u^*}(U)\}}, \quad (5)$$

where we drop the dependence of τ_{u^*} on $\mathcal{K}(\cdot, \cdot)$ and h for brevity.

Proposition 1 verifies that Cont.PCE is a perfectly localized functional and Loc.PCE is its local functional substitute under the identification assumptions (c.f. Section 2.3). To enhance the understanding, we demonstrate how to verify the conditional average treatment effect as a perfectly localized functional in Section S3 in Supplementary File; see also Chernozhukov et al. (2023, Examples 3.1–3.3) for more examples.

Proposition 1 *Suppose Assumptions 1–3 and the regularity conditions in Section S4.1 in the Supplementary File hold. For any $u^* \in \mathbb{R}^2$ such that $e_{u^*} > 0$, we have,*

$$|\tau_{u^*} - \tau_{u^*}^*| = \mathcal{O}(h^2). \quad (6)$$

In (1), let $W = (X, Z, U)$, $\gamma^*(x, z, u) = \mu_{zmz}(x)$, $D = U$, $d_0 = u^*$, $m(w, y, \gamma^*, k_h) = \{\gamma^*(x, 1, u) - \gamma^*(x, 0, u)\} [\mathbb{E}\{k_h(U)\}]^{-1} k_h(u)$, $\vartheta_0 = \tau_{u^*}^*$, and $\vartheta_h = \tau_{u^*}$. Then (6) implies that $\tau_{u^*}^*$ is a perfectly localized functional, and τ_{u^*} is the associated local functional substitute.

Remark 4 *If Cont.PCE $\tau_{u^*}^*$ is non-linear yet smooth enough, Proposition 1 controls the smoothing bias between τ_u and $\tau_{u^*}^*$ and ensures that Loc.PCE with a small h is a good approximation of Cont.PCE. Moreover, if Cont.PCE surface $\tau_{u^*}^*$ is a linear function of $u \in \mathbb{R}^2$ (e.g., $\tau_{u^*}^*$ in Figure 1) and kernel function $\mathcal{K}(\cdot, \cdot)$ is symmetric, then Loc.PCE τ_u indeed is identical to Cont.PCE $\tau_{u^*}^*$.*

In light of Proposition 1 and following Chernozhukov et al. (2022), we will focus on Loc.PCE as our target estimand to develop an efficient estimator. We begin by deriving the identification result for Loc.PCE, similar to Theorem 1. For ease of exposition, given any function $h_u(\mathcal{T})$ with \mathcal{T} being any variables other than u , we define $[\![\cdot]\!]_{u^*}$ as a kernel smoother of $h_u(\mathcal{T})$ around $u = u^*$,

$$[\![h_{(\cdot, \star)}(\mathcal{T})]\!]_{u^*} = \int_{\mathbb{R}^2} h_u(\mathcal{T}) k_{u^*}(u) du. \quad (7)$$

Here (\cdot, \star) represents the two-dimensional vector u in $h_u(\mathcal{T})$ are integrated out.

Theorem 2 (Identification of Loc.PCE) *Suppose Assumptions 1–3 hold. Given $u^* \in \mathbb{R}^2$, kernel function $k_{u^*}(\cdot)$ and bandwidth $h > 0$, then τ_{u^*} can be identified as*

$$\tau_{u^*} = \frac{\mathbb{E}[\{\mu_{1\cdot}(X) - \mu_{0\star}(X)\}e_{\star}(X)]_{u^*}}{\mathbb{E}[e_{\star}(X)]_{u^*}}, \quad (8)$$

where $e_{u^*}(X) = c_\rho\{F_{1m_1^*}(X), F_{0m_0^*}(X)\} \prod_{z \in \{0,1\}} f_{zm_z^*}(X)$.

Comparing (4) and (8), it becomes clear that Loc.PCE acts as a locally kernel-smoothed approximation of Cont.PCE.

3 EIFs and an oracle EIF-invoked estimator

We adopt a semiparametric approach similar to the binary- M case in Jiang et al. (2022) to propose a principled estimator for Loc.PCE, and subsequently, for Cont.PCE when $h \rightarrow 0$ as $n \rightarrow \infty$. We begin by deriving EIFs under two scenarios, where \mathcal{M}_{tp} and \mathcal{M}_{om} are both nonparametric, and \mathcal{M}_{ps} is either nonparametric or follows a parametric model,

$$f_{zm}(x | \beta) = f(M = m | X = x, Z = z, \beta). \quad (9)$$

Here $\beta \in \mathbb{R}^{d_\beta}$ is the parameter and $f_{zm}(x) = f_{zm}(x | \beta^*)$. The resulting EIFs are complex, posing challenges in constructing a computationally viable estimator. We will elaborate on this issue later. To simplify the EIF and facilitate computation, we further derive the EIF under an oracle scenario, where the true \mathcal{M}_{ps} is known.

Theorem 3 *Suppose Assumptions 1–3 hold, and \mathcal{M}_{tp} and \mathcal{M}_{om} are nonparametric.*

(i) *Suppose \mathcal{M}_{ps} is known. The EIF of τ_{u^*} is*

$$\phi_{\text{kn}}(x, z, m, y) = \underbrace{\frac{\xi_{1,u^*}(x, z, m)\{y - \mu_{zm}(x)\}}{\mathbb{E}\llbracket e_{\cdot,\star}(X) \rrbracket_{u^*}}}_{\text{(I)}} + \underbrace{\frac{\llbracket \{\mu_{1\cdot}(x) - \mu_{0\star}(x)\}e_{\cdot,\star}(x) \rrbracket_{u^*} - \tau_{u^*}\llbracket e_{\cdot,\star}(x) \rrbracket_{u^*}}{\mathbb{E}\llbracket e_{\cdot,\star}(X) \rrbracket_{u^*}}}_{\text{(II)}}, \quad (10)$$

where $\xi_{1,u^*}(x, z, m) = \{\pi_z(x)f_{zm}(x)\}^{-1}(-1)^{z+1}\gamma_{1,u^*}(z)(x, m)$, $\gamma_{1,u^*}^{(1)}(x, m) = \int_{\mathbb{R}} k_{u^*}(m, m_0) e_{(m,m_0)}(x) dm_0$, and $\gamma_{1,u^*}^{(0)}(x, m) = \int_{\mathbb{R}} k_{u^*}(m_1, m) e_{(m_1,m)}(x) dm_1$.

(ii) *Suppose \mathcal{M}_{ps} is unknown but follows a parametric model (9). Let $s_{zm}(x | \beta^*) = \partial \log f_{zm}(x | \beta) / \partial \beta |_{\beta=\beta^*}$ be the score function of (9), and suppose information matrix $I_{\beta^*} = \mathbb{E}\{s_{ZM}(X | \beta^*)s_{ZM}^\top(X | \beta^*)\}$ is positive-definite. The EIF of τ_{u^*} is $\phi_{\text{p}}(x, z, m, y) = \phi_{\text{kn}}(x, z, m, y) + \tilde{\phi}_{\text{p}}(x, z, m)$, where $\tilde{\phi}_{\text{p}}(x, z, m)$ is defined in (S3).*

(iii) *Suppose \mathcal{M}_{ps} is unknown and nonparametric. The EIF of τ_{u^*} is $\phi_{\text{np}}(x, z, m, y) = \phi_{\text{kn}}(x, z, m, y) + \tilde{\phi}_{\text{np}}(x, z, m)$, where $\tilde{\phi}_{\text{np}}(x, z, m)$ is defined in (S4).*

The expressions for $\tilde{\phi}_{\text{p}}(x, z, m)$ and $\tilde{\phi}_{\text{np}}(x, z, m)$ are complex and do not provide additional insights and thus are deferred to (S3) and (S4) in the Supplement File. We discuss the connections and comparisons among the EIFs. In $\phi_{\text{np}}(\cdot)$, function $\phi_{\text{kn}}(\cdot)$ comprises two orthogonal functionals, (I) and (II) (cf. (10)), which belong to the orthogonal tangent spaces spanned by the scores of $f(y | x, z, m)$ and $f(x)$, respectively, and function $\tilde{\phi}_{\text{np}}(\cdot)$ belongs to the tangent space spanned by the score of $f(m | x, z)$. To derive EIFs $\phi_{\text{p}}(\cdot)$ and $\phi_{\text{kn}}(\cdot)$, we project $\phi_{\text{np}}(\cdot) = \phi_{\text{kn}}(\cdot) + \tilde{\phi}_{\text{np}}(\cdot)$

onto the corresponding tangent spaces under Scenarios (i) and (ii), respectively. Compared with Scenario (iii), under Scenarios (i) and (ii), only the assumption on $f(m | x, z)$ changes, and $\phi_{\text{kn}}(\cdot)$ is orthogonal to the corresponding tangent spaces spanned by the scores of $f(m | x, z)$. Therefore, $\phi_{\text{kn}}(\cdot)$ remains the same after projection. The projection of $\tilde{\phi}_{\text{np}}(\cdot)$ is zero under Scenario (i), because the tangent space spanned by a known $f(m | x, z)$ is null, and the projection of $\tilde{\phi}_{\text{np}}(\cdot)$ is $\tilde{\phi}_{\text{p}}(\cdot)$ under Scenario (ii). Under Scenarios (i)–(iii), the corresponding semiparametric efficiency bounds (SEBs) are $\text{SEB}_\star = \mathbb{E}\{\phi_\star^2(X, Z, M, Y)\}$, $\star \in \{\text{kn}, \text{p}, \text{np}\}$. The additional terms in $\phi_{\text{p}}(\cdot)$ and $\phi_{\text{np}}(\cdot)$ amplify the variances in the SEBs for Scenarios (ii) and (iii). This is because, under Scenario (i), we have complete information about \mathcal{M}_{ps} , which results in the smallest efficiency bound. In particular, as the knowledge about \mathcal{M}_{ps} reduces from Scenario (i) to Scenario (iii), we have $\text{SEB}_{\text{kn}} \leq \text{SEB}_{\text{p}} \leq \text{SEB}_{\text{np}}$.

The EIFs in Theorem 3 can be used to construct semiparametrically efficient estimators via classical semiparametric theory (Bickel et al., 1993), but the complexity of $\phi_{\text{p}}(\cdot)$ and $\phi_{\text{np}}(\cdot)$ makes it impractical to generate computationally feasible estimators. For instance, accurately computing $\tilde{\phi}_{\text{np}}(\cdot)$ and $\tilde{\phi}_{\text{p}}(\cdot)$ can be time-consuming due to the need for $2n$ times of 2D-numerical integrals with intricate integrands, and this can be further exacerbated when using Bootstrap to construct confidence intervals. Additionally, calculating derivatives such as $c_\rho^{(1)}(\cdot, \cdot)$ and $s_{zm}(x | \beta^*)$ in $\tilde{\phi}_{\text{p}}(\cdot)$ and $\tilde{\phi}_{\text{np}}(\cdot)$ requires additional effort from the researchers.

As a result, our proposed estimator $\hat{\tau}_{u^*}$ is based on the oracle EIF, which is the simplest among the three. Importantly, $\hat{\tau}_{u^*}$ only requires n times of 2D numerical integrals and does not require calculating any derivatives. Consider the empirical version of $\phi_{\text{kn}}(x, z, m, y)$,

$$\hat{\phi}_{\text{kn}}(x, z, m, y) = \frac{\hat{\xi}_{1,u^*}(x, z, m)\{y - \hat{\mu}_{zm}(x)\} + \left[\left\{ \hat{\mu}_1(x) - \hat{\mu}_{0\star}(x) \right\} \hat{e}_{\cdot\star}(x) \right]_{u^*} - \hat{\tau}_{u^*} \left[\hat{e}_{\cdot\star}(x) \right]_{u^*}}{\mathbb{P}_n \left[\left[\hat{e}_{\cdot\star}(X) \right]_{u^*} \right]},$$

where the nuisance functions $\{\hat{\pi}_z(x), \hat{f}_{zm}(x), \hat{\mu}_{zm}(x)\}$ are trained by any off-the-shelf estimation algorithms, and other quantities are approximated by the plug-in estimators:

- (i) $\hat{F}_{zm_z}(x) = \int_{-\infty}^{m_z} \hat{f}_{zm}(x) dm$, $\hat{e}_u(x) = c_\rho\{\hat{F}_{1m_1}(x), \hat{F}_{0m_0}(x)\} \prod_{z \in \{0,1\}} \hat{f}_{zm_z}(x)$;
- (ii) $\hat{\gamma}_{1,u^*}^{(1)}(m, x) = \int_{\mathbb{R}} k_{u^*}(m, m_0) \hat{e}_{(m,m_0)}(x) dm_0$, $\hat{\gamma}_{1,u^*}^{(0)}(m, x) = \int_{\mathbb{R}} k_{u^*}(m_1, m) \hat{e}_{(m_1,m)}(x) dm_1$;
- (iii) $\hat{\xi}_{1,u^*}(x, z, m) = (-1)^{z+1} \{\hat{\pi}_z(x) \hat{f}_{zm}(x)\}^{-1} \hat{\gamma}_{1,u^*}^{(z)}(m, x)$.

Solving $\hat{\tau}_{u^*}$ from the estimating equation $\mathbb{P}_n\{\hat{\phi}_{\text{kn}}(X, Z, M, Y)\} = 0$, we obtain the proposed estimator,

$$\hat{\tau}_{u^*} = \frac{\mathbb{P}_n \left[\hat{\xi}_{1,u^*}(X, Z, M) \{Y - \hat{\mu}_{ZM}(X)\} + \left[\left\{ \hat{\mu}_1(X) - \hat{\mu}_{0\star}(X) \right\} \hat{e}_{\cdot\star}(X) \right]_{u^*} \right]}{\mathbb{P}_n \left[\left[\hat{e}_{\cdot\star}(X) \right]_{u^*} \right]}, \quad (11)$$

for Loc.PCE, and subsequently, for Cont.PCE when $h \rightarrow 0$ as $n \rightarrow \infty$.

4 Theoretical properties

4.1 Limit double robustness

Denote the functional limits $\bar{\pi}_z(x)$ and $\bar{\mu}_{zm}(x)$, such that $\hat{\pi}_z(x) \rightarrow \bar{\pi}_z(x)$, $\hat{\mu}_{zm}(x) \rightarrow \bar{\mu}_{zm}(x)$ as $n \rightarrow \infty$. Recalling (11), the probabilistic limit of $\hat{\tau}_{u^*}$ is

$$\bar{\tau}_{u^*} = \frac{\mathbb{E}\left[\bar{\xi}_{1,u^*}(X, Z, M)\{Y - \bar{\mu}_{ZM}(X)\} + \left[\left\{\bar{\mu}_{1\cdot}(X) - \bar{\mu}_{0\cdot}(X)\right\}e_{\cdot\star}(X)\right]_{u^*}\right]}{\mathbb{E}\left[e_{\cdot\star}(X)\right]_{u^*}}, \quad (12)$$

where $\bar{\xi}_{1,u^*}(x, z, m) = \{\bar{\pi}_z(x)f_{zm}(x)\}^{-1}(-1)^{z+1}\gamma_{1,u^*}^{(z)}(x, m)$. Theorem 4 below justifies the limit double robustness of $\hat{\tau}_{u^*}$ when \mathcal{M}_{ps} is correctly specified.

Theorem 4 *Suppose Assumptions 1–3 hold, and \mathcal{M}_{ps} is correctly specified. If either $\bar{\pi} = \pi$ or $\bar{\mu} = \mu$, we have $\bar{\tau}_{u^*} = \tau_{u^*}$, and moreover, $\lim_{h \rightarrow 0} \bar{\tau}_{u^*} = \tau_{u^*}^*$ under the conditions of Proposition 1.*

Intuitively, with nonparametric \mathcal{M}_{tp} and \mathcal{M}_{om} , our EIF-invoked estimator enjoys the Neyman orthogonality (Chernozhukov et al., 2017) with respect to these two nuisance functions, which leads to the double robustness property as shown in Theorem 4.

Remark 5 *When M is binary, Jiang et al. (2022) illustrate the triple robustness of their EIF-based PCE estimator; i.e., their estimator is consistent as long as any pair of the three models in $(\mathcal{M}_{tp}, \mathcal{M}_{ps}, \mathcal{M}_{om})$ is correctly specified. The crucial reason why their estimator is triply robust while ours is doubly robust is that the underpinning identification assumptions are different. Their monotonicity assumption provides a neat identification PCE formula, enabling triple robustness. However, when M is continuous, the monotonicity fails to identify Cont.PCE and can be implausible in real-world problems such as surrogate evaluation. Due to the nonlinearity of copula $c_\rho(\cdot, \cdot)$, copula-based identification requires correct specification of \mathcal{M}_{ps} , losing robustness against its misspecification. If \mathcal{M}_{ps} is incorrectly specified in $e_u(x)$, the bias for $\bar{\tau}_{u^*}$ cannot be eliminated through correct specifications of $\mu_{zm}(x)$ and $\pi_z(x)$.*

4.2 Rate-double robustness and optimality

Desirable asymptotic properties of $\hat{\tau}_{u^*}$ are shown under the semiparametric setting that \mathcal{M}_{ps} is parametrically modeled through (9), while \mathcal{M}_{tp} and \mathcal{M}_{om} are nonparametrically modeled. Examples of (9) include the linear model, additive errors nonlinear model and generalized linear model (Boos et al., 2013). With fixed $h > 0$, we first derive the asymptotic results of $\hat{\tau}_{u^*}$ towards the Loc.PCE τ_{u^*} . Assumption 4 below imposes some typical regularity conditions for the nuisance functions and their estimators; see, e.g. Kennedy (2016).

Assumption 4 *(i) Either $\bar{\mu} = \mu$ or $\bar{\pi} = \pi$; (ii) $\|\hat{\mu} - \bar{\mu}\|_\infty, \|\hat{\pi} - \bar{\pi}\|_\infty = o_{\mathbb{P}}(1)$; (iii) $\mathbb{P}(|\hat{\pi} - \pi|^2) = \mathcal{O}(r_\pi^2)$ and $\mathbb{P}(|\hat{\mu} - \mu|^2) = \mathcal{O}(r_\mu^2)$ with $r_\pi, r_\mu \lesssim 1$ as $n \rightarrow \infty$; (iv) Functions like $\hat{\pi}(x), \bar{\pi}(x), \hat{\mu}_{zm}(x), \bar{\mu}_{zm}(x)$ and $\hat{\mu}_{am_a}(x), \bar{\mu}_{am_a}(x)$ for $a = 0, 1$, are contained in uniformly bounded function classes with finite*

uniform entropy integrals (see Section S6 in the Supplementary File for details), and $\hat{\pi}^{-1}$, $\bar{\pi}^{-1}$ are uniformly bounded.

If one exploits the cross-fitting strategy for $\hat{\tau}_{u^*}$, (iv) can be further dropped, and our theoretical results remain valid; see, e.g., [Robins et al. \(2008\)](#) and [Chernozhukov et al. \(2018\)](#) for details.

Theorem 5 (Loc.PCE inference) *Suppose Assumptions 1–4 and other regularity conditions in Section S5.5 hold. In addition, assume the parametric model of \mathcal{M}_{ps} in (9) is correctly specified and $\hat{\beta}$ is a regular \sqrt{n} -consistent estimator of β^* .*

I. (Rate-double robustness) *As $n \rightarrow \infty$, we have*

$$\hat{\tau}_{u^*} - \tau_{u^*} = \mathcal{O}_{\mathbb{P}}(n^{-1/2} + r_{\pi}r_{\mu}). \quad (13)$$

II. (Asymptotic normality) *Furthermore, suppose $r_{\pi}r_{\mu} = o(n^{-1/2})$, and $\hat{\beta}$ is asymptotically linear, i.e., there exists some $\psi(x, z, m)$ such that $\mathbb{E}\{\psi(X, Z, M)\} = 0$, $\mathbb{E}\|\psi(X, Z, M)\|^2 < +\infty$, and $\sqrt{n}(\hat{\beta} - \beta^*) = n^{-1/2} \sum_{i=1}^n \psi(X_i, Z_i, M_i) + o_{\mathbb{P}}(1)$. Then, with $\bar{\sigma}_{u^*}^2$ defined in (S5), we have*

$$\sqrt{n}(\hat{\tau}_{u^*} - \tau_{u^*}) \rightsquigarrow \mathcal{N}(0, \bar{\sigma}_{u^*}^2). \quad (14)$$

III. (Semiparametric efficiency) *Since h is fixed, τ_{u^*} is a regular estimand. Then we can use the classic semiparametric statistics theory to justify the optimality of $\hat{\tau}_{u^*}$ ([Bickel et al., 1993](#)). Suppose all conditions above hold. Further assume $\bar{\pi} = \pi$, $\bar{\mu} = \mu$, and $\hat{\beta}$ is a semiparametrically efficient estimator for β^* such that $\psi(x, z, m) = I_{\beta^*}^{-1} s_{zm}(x | \beta^*)$, where $s_{zm}(x | \beta^*)$ and I_{β^*} are defined in Theorem 3. Then we have $\sigma_{u^*}^2 = \text{SEB}_{\text{p}} = \mathbb{E}\{\phi_{\text{p}}^2(X, Z, M, Y)\}$. Thus, by Theorem 3(ii), $\hat{\tau}_{u^*}$ is semiparametrically efficient.*

The error rate (13) in Theorem 5(I) illustrates the rate-double robustness of our proposed estimator. The asymptotic linearity condition for $\hat{\beta}$ can be attained by commonly-used methods, e.g., the likelihood-based or quasi-likelihood-based estimators, and generalized method of moments ([Boos et al., 2013](#), [McCullagh and Nelder, 2019](#)). To attain the optimal root- n convergence of $\hat{\tau}_{u^*}$, we only require $\mu_{zm}(x)$ and $\pi_z(x)$ to be well estimated, e.g., $r_{\pi}, r_{\mu} \lesssim n^{-1/4}$. When one of $\mu_{zm}(x)$ and $\pi_z(x)$ can be estimated parametrically (i.e., r_{π} or $r_{\mu} \lesssim n^{-1/2}$), $\hat{\tau}_{u^*}$ is root- n consistent even the other one is not correctly specified. For asymptotic normality (14) in Theorem 5(II), the condition $r_{\pi}r_{\mu} = o_{\mathbb{P}}(n^{-1/2})$ is standard in the double machine learning literature ([Chernozhukov et al., 2018](#)). Under the condition of Theorem 5(II), we can construct the CI for Loc.PCE. A closed-form estimator for the asymptotic variance $\sigma_{u^*}^2$ is feasible, which however is complicated due to the complex form of $\sigma_{u^*}^2$ (c.f. (S5)). Thus a standard nonparametric bootstrap procedure ([Efron, 1981](#)) is recommended to estimate $\sigma_{u^*}^2$; see, e.g., [Imbens \(2004, § IV\)](#) and [Jiang et al. \(2022\)](#) for the use of bootstrap asymptotic variance estimation in different problems.

Theorem 5(III) further justifies the optimality of $\hat{\tau}_{u^*}$. It requires correct specification of both \mathcal{M}_{tp} and \mathcal{M}_{om} , and that $\hat{\beta}$ is semiparametrically efficient, which holds when $\hat{\beta}$ is a regular MLE.

These conditions are necessary for achieving semiparametric efficiency in doubly robust estimators, such as the augmented inverse propensity weighted estimator (Hahn, 1998).

Finally, we derive nonparametric convergence results of $\hat{\tau}_{u^*}$ for Cont.PCE estimation. In particular, the asymptotic rate for $\hat{\tau}_{u^*} - \tau_{u^*}$ in (13) can be generalized to include h , when we let $h \rightarrow 0$ as $n \rightarrow \infty$. Combing such results with the asymptotic bias of $\tau_{u^*} - \tau_{u^*}^*$ in Proposition 1, we show the convergence rate and rate-double robustness of $\hat{\tau}_{u^*} - \tau_{u^*}^*$. Similarly, we generalize the result in Theorem 5(II) and show the asymptotic normality of $\hat{\tau}_{u^*} - \tau_{u^*}^*$.

Theorem 6 (Cont.PCE inference) *Suppose Assumptions 1–4 and regularity conditions in Section S4.1 and Section S5.5 hold. We have*

$$\hat{\tau}_{u^*} - \tau_{u^*}^* = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2} + h^2 + h^{-1}r_{\pi}r_{\mu}). \quad (15)$$

Moreover, suppose the additional conditions in Theorem 5(II) hold and we choose any $h = o(n^{-1/6})$. Then with $\tilde{\sigma}_{u^*}^2$ defined in (S104) in the Supplementary File, we have

$$\sqrt{nh}(\hat{\tau}_{u^*} - \tau_{u^*}^*) \rightsquigarrow \mathcal{N}(0, \tilde{\sigma}_{u^*}^2). \quad (16)$$

In (15), we derive the nonparametric convergence rate of $\hat{\tau}_{u^*}$ towards $\tau_{u^*}^*$. The three terms on the right-hand side of (15) represent the nonparametric variance, nonparametric bias, and the effect of nuisance estimation errors, respectively. Specifically, under the rate-doubly robust condition $r_{\pi}r_{\mu} = \mathcal{O}(n^{-1/2})$, we have $\hat{\tau}_{u^*} - \tau_{u^*}^* = \mathcal{O}_{\mathbb{P}}(n^{-1/3})$ by choosing $h \asymp n^{-1/6}$, which is minimax optimal for the pointwise estimation of a two-dimensional and twice-differentiable nonparametric regression function $\tau_{u^*}^*$ (Tsybakov, 2008). See the regularity conditions for $\tau_{u^*}^*$ in Section S4.1. With an undersmoothing selection of $h > 0$ and well-estimated nuisance functions such that $r_{\pi}r_{\mu} = \mathcal{O}(n^{-1/2})$, we further achieve the asymptotic normality in (16). Similar to the Loc.PCE, we can use nonparametric bootstrap to estimate $\tilde{\sigma}_{u^*}^2$ and thereby construct CI for $\tau_{u^*}^*$.

Remark 6 (Nonparametric estimation of \mathcal{M}_{ps}) *In general, suppose $\hat{f}_{zm}(x)$ is nonparametrically estimated with $\mathbb{E}\{\hat{f}_{zm}(X) - f_{zm}(X)\}^2 = \mathcal{O}_{\mathbb{P}}(r_f^2) = o_{\mathbb{P}}(1)$ for $z = 0, 1$, and $r_{\pi}r_{\mu} \rightarrow 0$. Following the proofs of Theorem 5 and Theorem 6, we can show the rates in (13) and (15) become $\mathcal{O}_{\mathbb{P}}(r_f + r_{\pi}r_{\mu})$ and $\mathcal{O}_{\mathbb{P}}(h^{-1}r_f + h^2 + h^{-1}r_{\pi}r_{\mu})$, respectively, and thus $\hat{\tau}_{u^*}$ remains consistent for $\tau_{u^*}^*$, and furthermore, for $\tau_{u^*}^*$ when we select $r_f \prec h \prec 1$ as $n \rightarrow \infty$.*

5 Related works

Joint modeling of continuous principal strata. Pioneering works have proposed flexible methods to model the joint distribution of continuous M_1 and M_0 . Bartolucci and Grilli (2011) use copula to model the joint distribution of (M_1, M_0) with parametric outcome model for randomized experiment. They propose a likelihood approach to estimate and perform sensitivity analysis with respect to ρ . Schwartz et al. (2011) further adopt the Dirichlet process mixture model to model

$(M_1, M_0) \mid X$. [Zigler and Belin \(2012\)](#) develop a fully Bayesian approach, and assume $(M_1, M_0) \mid X$ follows a joint conditional Gaussian model with a known ρ , which corresponds to a Gaussian linear regression model for (9) with a Gaussian copula $c_\rho(\cdot, \cdot)$. Unlike previous works that rely on concrete modeling assumptions without assuming principal ignorability, our proposed estimator under weak principal ignorability is more robust and flexible. It does not necessarily require parametric assumptions for all nuisance functions and remains consistent even if \mathcal{M}_{tp} or \mathcal{M}_{om} is misspecified.

Semiparametric estimation of Cont.PCE under principal ignorability. During the revision of the current manuscript, another paper uploaded on arXiv ([Lu et al., 2023](#)) also studies the semiparametric estimation problem of Cont.PCE under principal ignorability. To handle the non-regularity of Cont.PCE, the authors propose a global approximation approach. They project the Cont.PCE surface onto a general parametric working model and then study the semiparametric efficient estimation for this parametric substitute of Cont.PCE. While their approach elegantly reduces the estimation complexity of Cont.PCE, its precision may depend on the approximation power of the parametric working model. In contrast, we focus on the semiparametric estimation for the local approximation of Cont.PCE, which further facilitates the nonparametric and doubly robust estimation for the exact Cont.PCE through a vanishing bandwidth.

Local and localized functional estimation. [Chernozhukov et al. \(2022, 2023\)](#) propose a general Riesz-representation-invoked method which estimates local functionals semiparametrically efficiently ([Chernozhukov et al., 2022](#), Theorem 4.2), and thereby approximates the corresponding perfectly localized functionals accurately. They consider the minimal Riesz representer and outcome mean model as nuisance functions, and demonstrate the double robustness and semiparametric efficiency of their estimator. However, their framework cannot be applied to estimate Loc.PCE and Cont.PCE, because they assume all variables in (W, Y) (cf. (1)) are observable, while under this paper’s setting, principal strata U in W contain one unobserved variable. Moreover, for identification, we need to deal with three nuisance functions instead of two, which introduces extra difficulties in estimation. To resolve these challenges while achieving the same desirable properties as [Chernozhukov et al. \(2022, 2023\)](#), we propose our estimator through a new oracle EIF-invoked strategy. Although $\hat{\tau}_{u^*}$ is derived through the EIF under the oracle \mathcal{M}_{ps} -known scenario, it can still attain the semiparametric efficient bound SEB_p when \mathcal{M}_{ps} follows a parametric model (9) (c.f. Theorem 5(III)).

6 Simulation study

We perform numerical experiments to assess the large-sample properties of $\hat{\tau}_u$ on estimating Cont.PCE. The data generation process involves the following steps: first, we generate $X = (X_1, X_2, X_3)^\top$ from the standard normal distribution $\mathcal{N}(0, I_3)$. Second, we generate Z from \mathcal{F}_{tp} given X . Third, we generate M from \mathcal{F}_{ps} given X and Z . Finally, we generate Y from \mathcal{F}_{om} given all the other variables. We use the standard Gaussian copula $c_\rho(\cdot, \cdot)$ with $\rho = 0.5$. We consider two comparative distributions for each of the three distributions $\mathcal{F}_{\text{tp}}, \mathcal{F}_{\text{ps}}, \mathcal{F}_{\text{om}}$ as follows:




Figure 2: The box-plots of $\hat{\tau}_u - \tau_u^*$ over 100 MC rounds when $n = 1000$. X-axis represents different data generation mechanisms. For example, (TP2,PS1,OM1) means data are generated from $(\mathcal{F}_{\text{tp}}^{(2)}, \mathcal{F}_{\text{ps}}^{(1)}, \mathcal{F}_{\text{ps}}^{(1)})$. Different colors represent different principal levels u . The top and bottom panels present the results when nuisance functions are estimated through parametric and semiparametric strategies, respectively.

- $\mathcal{F}_{\text{tp}}^{(1)} = \text{Bern}\{\text{logit}(X_1 + X_2 + X_3)\}$ and $\mathcal{F}_{\text{tp}}^{(2)} = \text{Bern}\{\text{logit}(-X_1^2 + X_2)\}$;
- $\mathcal{F}_{\text{ps}}^{(1)} = \mathcal{N}(X_1 + X_2 + X_3 + Z, 1)$ and $\mathcal{F}_{\text{ps}}^{(2)} = \mathcal{N}(X_1^2 + 2X_3Z, 1 + X_1^2Z - 0.5Z)$;
- $\mathcal{F}_{\text{om}}^{(1)} = \mathcal{N}(X_1 + X_2 + X_3 + Z + M, 1)$ and $\mathcal{F}_{\text{om}}^{(2)} = \mathcal{N}(X_1Z + X_2 + M^2, 1)$.

Based on the identification formula (4), we generate 10^6 samples, $\{(X_i, Z_i, M_i, Y_i)\}_{i=1}^{10^6}$, under the corresponding data generation setting, and approximate τ_u^* as $\tau_u^* \approx [\sum_{i=1}^{10^6} \{\mu_{1m_1}(X_i) - \mu_{0m_0}(X_i)\}e_u(X_i)] / [\sum_{i=1}^{10^6} e_u(X_i)]$, where $e_u(X)$ can be specified by Assumption 3.

6.1 Simulation: the consistency and double robustness

We simulate data through eight different data generation mechanisms, with $(\mathcal{F}_{\text{tp}}, \mathcal{F}_{\text{ps}}, \mathcal{F}_{\text{om}}) = (\mathcal{F}_{\text{tp}}^{(a)}, \mathcal{F}_{\text{ps}}^{(b)}, \mathcal{F}_{\text{om}}^{(c)})$, $a, b, c \in \{1, 2\}$. In addition, we choose $n \in \{500, 800, 1000\}$ and $h = 0.8n^{-1/6}$ following the optimal rate in Theorem 6. We consider both a parametric and a semiparametric strategy for estimating the nuisance functions.

The parametric strategy We estimate $\pi_z(x)$ and $\mu_{zm}(x)$ through logistic regression and linear regression, respectively. To estimate $f_{zm}(x)$, we specify the parametric model in (9) as $f_{zm}\{x \mid \beta = (\ell, \sigma^2)\} = \varphi\{m \mid \ell^\top(x, z), \sigma^2\}$, where ℓ is a vector in \mathbb{R}^4 and $\varphi(\cdot \mid \iota, \omega)$ denotes the Gaussian density function with mean ι and variance ω . We estimate ℓ and σ^2 through the linear regression $M \sim (X_1, X_2, X_3, Z)$, where $\hat{\ell}$ denotes the vector of regression coefficients and $\hat{\sigma}^2$ represents the empirical residual variance. Hence, \mathcal{M}_{ps} is correctly specified if and only if $\mathcal{F}_{\text{ps}} = \mathcal{F}_{\text{ps}}^{(1)}$.

The semiparametric strategy We estimate $\pi_z(x)$ and $\mu_{zm}(x)$ through an ensemble learning algorithm, namely SuperLearner (Van der Laan et al., 2007), which combines various machine

learning methods such as linear regression, multivariate adaptive regression splines, single-layer neural networks, and recursive regression trees. The estimation of \mathcal{M}_{ps} is the same as the parametric strategy, where we estimate ℓ and σ^2 through the linear regression $M \sim (X_1, X_2, X_3, Z)$, and $\hat{\ell}$ is the vector of regression coefficients while $\hat{\sigma}^2$ is the empirical residual variance.


We evaluate the performance of $\hat{\tau}_u$ on Cont.PCE estimation at six different levels of u , namely, $u = (0, 0), (0, 0.5), (0, 1), (0.5, 0.5), (0.5, 1), (1, 1)$, and demonstrate the estimation errors of $\hat{\tau}_u$ across 100 Monte Carlo (MC) iterations in Figure 2 when $n = 1000$. The complete simulation results with different sample sizes $n \in \{500, 800, 1000\}$ are contained in Figure S5 in the Supplementary File. Our numerical results indicate: (1) When employing a parametric nuisance strategy, the estimation biases of $\hat{\tau}_u - \tau_u^*$ are close to zero for all tested u , provided that $\mathcal{F}_{\text{ps}} = \mathcal{F}_{\text{ps}}^{(1)}$ and either $\mathcal{F}_{\text{tp}} = \mathcal{F}_{\text{tp}}^{(1)}$ or $\mathcal{F}_{\text{om}} = \mathcal{F}_{\text{om}}^{(1)}$, and the estimation variances are particularly small when all nuisance functions are correctly specified, i.e., $(\mathcal{F}_{\text{tp}}, \mathcal{F}_{\text{ps}}, \mathcal{F}_{\text{om}}) = (\mathcal{F}_{\text{tp}}^{(1)}, \mathcal{F}_{\text{ps}}^{(1)}, \mathcal{F}_{\text{om}}^{(1)})$; (2) When using a semiparametric nuisance strategy, the estimation biases are close to zero and the estimation variances are particularly small whenever $\mathcal{F}_{\text{ps}} = \mathcal{F}_{\text{ps}}^{(1)}$, as \mathcal{M}_{tp} and \mathcal{M}_{om} are always correctly specified by **SuperLearner**; (3) Under aforementioned settings, the estimation errors become smaller as n increases. These findings confirm the consistency and rate-double robustness of $\hat{\tau}_u$ as shown in Theorem 6.

6.2 Simulation: the limiting distribution


We verify the asymptotic normality of $\hat{\tau}_u$ by $n = 700$ samples and calculating the bootstrap standard deviation of $\hat{\tau}_u$ using 100 bootstrap samples, which serves as the nonparametric Bootstrap estimator of σ_u . To validate the asymptotic normality in Theorem 6, we choose undersmoothing $h = 0.8n^{-1/5}$. We obtain the 95% bootstrap CIs and calculate the empirical coverage rate (ECR) over 300 MC rounds, to check if the CIs capture the true τ_u^* . All nuisance functions are estimated via the parametric strategy (c.f. Section 6.1) for tested u . As shown in Table 1, when $(\mathcal{F}_{\text{tp}}, \mathcal{F}_{\text{ps}}, \mathcal{F}_{\text{om}}) = (\mathcal{F}_{\text{tp}}^{(1)}, \mathcal{F}_{\text{ps}}^{(1)}, \mathcal{F}_{\text{om}}^{(1)})$ and thus all nuisances can be estimated under the root- n rates, ECRs are close to 0.95 for all tested u because of the asymptotic normality of $\hat{\tau}_u$ (cf. (14)). When $(\mathcal{F}_{\text{tp}}, \mathcal{F}_{\text{ps}}, \mathcal{F}_{\text{om}}) = (\mathcal{F}_{\text{tp}}^{(2)}, \mathcal{F}_{\text{ps}}^{(1)}, \mathcal{F}_{\text{om}}^{(1)})$ or $(\mathcal{F}_{\text{tp}}^{(1)}, \mathcal{F}_{\text{ps}}^{(1)}, \mathcal{F}_{\text{om}}^{(2)})$, we have $r_{\pi} r_{\mu} \lesssim n^{-1/2}$, as one of $\mu_{zm}(x)$ and $\pi_z(x)$ is misspecified while the other can be estimated parametrically, which is of the bound-

Table 1: Empirical coverage rates (rounded to three decimal places) of $\hat{\tau}_u$'s 95% CIs over 300 Monte Carlo rounds, when all nuisance functions are trained via linear models. The triplet abc in the first row represents scenario $(\mathcal{F}_{\text{tp}}^{(a)}, \mathcal{F}_{\text{ps}}^{(b)}, \mathcal{F}_{\text{om}}^{(c)})$ for $a, b, c \in \{1, 2\}$.

u	111	211	112	212	121	221	122	222
(0, 0)	0.937	0.947	0.927	0.880	0.937	0.947	0.963	0.880
(0, 0.5)	0.957	0.937	0.967	0.910	0.940	0.947	0.930	0.963
(0, 1)	0.947	0.947	0.927	0.933	0.953	0.960	0.963	0.927
(0.5, 0.5)	0.920	0.937	0.947	0.923	0.927	0.957	0.933	0.950
(0.5, 1)	0.940	0.933	0.920	0.893	0.943	0.937	0.937	0.807
(1, 1)	0.930	0.930	0.943	0.900	0.920	0.953	0.877	0.837



(a) The surface of $\hat{\tau}_u$.



(b) 95% bootstrap CIs of $\hat{\tau}_u$ with varying u .

Figure 3: Panel (a) shows the surface of $\hat{\tau}_u$ with $\rho = 0.5, h = 0.25$ (for the standardized M), when $u \in [190 \text{ cells/mm}^3, 520 \text{ cells/mm}^3]^2$. The X-axis of (b) is different levels of u . The top panel of (b) shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying u when $m_1 = m_0$. The bottom panel of (b) shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying m_1 , while m_0 is fixed at 340 cells/mm^3 .

any case of condition $r_\pi r_\mu = o(n^{-1/2})$ for (14). Nevertheless, in these cases, the nuisance function estimation biases are still negligible enough, and ECRs are close to 0.95 for all tested u .

7 Real data applications

7.1 Surrogate analysis of short-term CD4 count in ACTG 175

ACTG 175 (Hammer et al., 1996) is one of the first randomized clinical trials to demonstrate the superiority of combination therapy with zidovudine and didanosine over monotherapy with zidovudine or didanosine for patients infected with human immunodeficiency virus type I (HIV-I). Each patient was randomly assigned a specific therapy. It is of essential interest to investigate whether a short-term endpoint measured several weeks after randomization can be a valid surrogate for a long-term clinical endpoint (Burzykowski et al., 2005). In this study, we explore this question using data from ACTG175 contained in the R package `speff2trial`, focusing on patients treated with zidovudine only ($Z = 0$) or zidovudine + didanosine ($Z = 1$). To ensure weak principal ignorability, we collect various covariates in X , including age, weight, hemophilia, homosexual activity, history of intravenous drug use, Karnofsky score, non-zidovudine antiretroviral therapy prior to initiation, zidovudine use in the 30 days prior to treatment initiation, race (white/non-white), gender, antiretroviral history, and symptomatic indicator. We select the short-term CD4 count at 20 ± 5 weeks as M and the long-term CD4 count at 96 ± 5 weeks as Y , both of which reflect patients' immune functions in different periods. To handle censoring and outliers, we exclude patients censored two years after randomization and patients with M_i larger than the 99th percentile or smaller than the 1st percentile among all M_i . Ultimately, our study focuses on 640 patients.


The parametric strategy for nuisance training, as outlined in Section 6.1, is employed in this study. The Q-Q plot of \mathcal{M}_{ps} estimation suggests that a linear regression model is appropriate, as demonstrated in Figure S6(a) in the Supplementary File. Our approach utilizes a Gaussian kernel, denoted by $\mathcal{K}(\cdot, \cdot)$, and a Gaussian copula, denoted by $c_\rho(\cdot, \cdot)$. To evaluate the estimator’s sensitivity under different scenarios of (M_1, M_0) ’s correlation, we investigate three values of $\rho = -0.5, 0, 0.5$. All continuous variables are standardized during the implementation of the proposed method, and are rescaled back when presenting the results. The standardized M_i values fall approximately within the range of -2 and 3 , as depicted in Figure S6(b). Thus, we have chosen two relatively small bandwidths $h = 0.25, 0.5$ to ensure that τ_u can approximate τ_u^* effectively. In practice, we recommend specifying h to be a small fraction of the standard deviation of the observed M_i values. For $\rho = 0.5$ and $h = 0.25$, the estimated $\hat{\tau}_u$ surface and 95% confidence intervals are reported in Figure 3, with both m_1 and m_0 ranging from 190 cells/mm³ to 520 cells/mm³, which roughly correspond to the 10th and 85th percentiles among all M_i . Our results are robust to different choices of ρ and h , as demonstrated in Figures S7 and S8.

The estimated Cont.PCE surface exhibits a pattern similar to the synthetic example in Figure 1(a) of an ideal surrogate, where the estimated dissociative effects are close to zero, and the estimated associative effects are positive (negative) when $m_1 > m_0$ ($m_1 < m_0$). We show the 95% Bootstrap CIs with 100 resamples of several different $\hat{\tau}_u$ with $m_1 = m_0$ in the top panel of Figure 3(b), none of which are significantly different from zero. However, the 95% CIs of $\hat{\tau}_u$ with different m_1 while fixing m_0 at 340 cells/mm³ in the bottom panel of Figure 3(b) reveal that τ_u^* becomes significantly larger as $m_1 - m_0$ increases.


To the best of authors’ knowledge, this is the first time that the short-term CD4 count is evaluated as a surrogate under the principal stratification framework. Previous works assess PTE (cf. Remark 2) and suggest not a large proportion of the treatment effect on the clinical endpoint can be explained by the short-term CD4 count (Hughes et al., 1998, Agniel and Parast, 2021). Nevertheless, our observations imply that, from the perspective of association strength and prediction power, the short-term CD4 count can still be a desirable principal surrogate of long-term CD4 for the considered population (Gilbert and Hudgens, 2008). Another reason why the short-term CD4 count might be a reasonable surrogate in our analysis is that we focus on a more healthy and cooperative population with no censoring due to death or lost-to-follow-up. This reduces the cases that can weaken the association between short-term and long-term CD4 counts, such as non-compliance, viral resistance, and changes of therapy in the face of disease progression (Hughes et al., 1998).

7.2 Causal effect of flooding on health

In this section, we extend the principal stratification analysis of the Bangladesh flooding dataset conducted by Jiang et al. (2022) to a continuous- M scenario. The dataset comprises data on 774 households during the 1998 Bangladesh flood. We investigate the causal effect of flooding (Z) on the days of children’s diarrhea (Y) through the variable of per capita calorie consumption of the household (M). In their study, Jiang et al. (2022) adapted the data to a binary- M framework by



(a) The surface of $\hat{\tau}_u$.



(b) 95% bootstrap CIs of $\hat{\tau}_u$ with varying u .

Figure 4: Panel (a) shows the surface of $\hat{\tau}_u$ with $\rho = 0.5, h = 0.5$ (for the standardized M), when $u \in [1400 \text{ calories}, 2500 \text{ calories}]^2$. The X-axis of (b) is different levels of u . The top panel of (b) shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying u when $m_1 = m_0$. The bottom panel of (b) shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying m_1 , while m_0 is fixed at 1900 calories.

truncating M to 1 if the per capita calorie consumption of the household was larger than 2000 calories and 0 otherwise. Our study does not require dichotomization and thus might conduct a more meticulous analysis. To ensure ignorability, we include gender, age, the size of the household, father’s education, mother’s education, father’s age, and mother’s age in covariates X .

The estimation of the nuisance functions and Loc.PCE follows a similar approach as outlined in Section 7.1. Additionally, we exclude several outliers where the M_i values are larger than the 99th percentile or smaller than the 1st percentile among all M_i . The standardized M_i values range approximately between -2.2 and 3.1 . Consequently, we select small bandwidths of $h = -0.25$ and 0.5 . Our model diagnosis indicates the validity of our preprocessing step, as depicted in Figure S9(a) and (b). To evaluate sensitivity, we examine ρ values of $-0.5, 0$, and 0.5 . We present the estimated $\hat{\tau}_u$ surface when $\rho = 0.5$ and $h = 0.5$ in Figure 4. Similar results with other ρ and h are referred to Figures S10 and S11.

Figure S9(a) illustrates that the Cont.PCE surface is positioned above the $\hat{\tau}_u = 0$ plane, indicating positive dissociative effects, i.e., $\tau_u^* > 0$ when $m_1 = m_0$. This observation suggests a positive direct effect (cf. Remark 1), where Z may affect Y through a causal pathway other than $Z \rightarrow M \rightarrow Y$, for instance, the pathway through mother’s health (Del Ninno, 2001). Thus, our findings support the conclusions of Jiang et al. (2022). From a principal stratification perspective, the τ_u^* surface also suggests negative associative effects, given that $\hat{\tau}_u$ increases as $m_1 - m_0$ decreases. This is also reasonable because for the households whose capita calorie consumption can be more severely affected by floods, they are more likely to have children who are strongly and persistently affected by diarrhea. Similar to Section 7.1, a thorough pointwise investigation of $\hat{\tau}_u$ with accompanying confidence intervals in Figure 4(b) quantifies the uncertainty and further supports our conclusions.

8 Discussion

To efficiently estimate the principal causal effect with continuous strata, we extend the local functional estimation framework in [Chernozhukov et al. \(2022, 2023\)](#). Our new oracle EIF-invoked estimator serves as a valuable complement to the existing literature for resolving computation and estimation difficulties in this scenario.

For future work, we suggest three directions. Firstly, it is interesting to examine the relationship between the EIF and the Riesz representation ([Chernozhukov et al., 2022, 2023](#)) of a target estimand for constructing estimators with desirable statistical properties. While the Riesz representation is identical to the EIF for some regular estimands such as the average treatment effect, it can be challenging to determine for more complex problems. We hypothesize that the EIF can offer insights into the Riesz representation in general, and plan to establish a formal connection in our future research. Secondly, some key identification assumptions, such as weak principal ignorability, are not verifiable based on the observed data, necessitating sensitivity analysis to assess the robustness of the proposed estimator against their violations. Similar to [Jiang et al. \(2022\)](#), sensitivity models can be introduced to handle possible breaches of these critical assumptions. Finally, when h is vanishing as n grows, non-asymptotic analysis of Cont.PCE estimation has been further shown in [Theorem 6](#). Thus, a data-driven algorithm for optimal and adaptive h selection, is also of interest for future research.

References

- D. Agniel and L. Parast. Evaluation of longitudinal surrogate markers. *Biometrics*, 77(2):477–489, 2021.
- D. W. Andrews. Empirical process methods in econometrics. *Handbook of econometrics*, 4:2247–2294, 1994.
- M. Baccini, A. Mattei, and F. Mealli. Bayesian inference for causal mechanisms with application to a randomized study for postoperative pain control. *Biostatistics*, 18(4):605–617, 2017.
- F. Bartolucci and L. Grilli. Modeling partial compliance through copulas in a principal stratification framework. *Journal of the American Statistical Association*, 106(494):469–479, 2011.
- P. J. Bickel, C. Klaassen, Y. Ritov, and J. Wellner. *Efficient and Adaptive Inference in Semiparametric Models*. Johns Hopkins University Press, Baltimore, 1993.
- D. D. Boos, L. A. Stefanski, et al. *Essential Statistical Inference*, volume 120. Springer, 2013.
- S. Boyd, S. P. Boyd, and L. Vandenberghe. *Convex Optimization*. Cambridge university press, 2004.
- T. Burzykowski, M. Buyse, and G. Molenberghs. *The Evaluation of Surrogate Endpoints*, volume 427. Springer, 2005.

- X. Chen and T. Christensen. Optimal sup-norm rates, adaptivity and inference in nonparametric instrumental variables estimation. *arXiv:1508:03365v1*, 2015.
- V. Chernozhukov, D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, and W. Newey. Double/debiased/neyman machine learning of treatment effects. *American Economic Review*, 107: 261–265, 2017.
- V. Chernozhukov, D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins. Double/debiased machine learning for treatment and structural parameters: Double/debiased machine learning. *The Econometrics Journal*, 21:C1–C68, 2018.
- V. Chernozhukov, W. K. Newey, and R. Singh. Debiased machine learning of global and local parameters using regularized riesz representers. *The Econometrics Journal*, 25:576–601, 2022.
- V. Chernozhukov, W. K. Newey, and R. Singh. A simple and general debiased machine learning theorem with finite-sample guarantees. *Biometrika*, 110:257–264, 2023.
- J. Chu, W. Lu, and S. Yang. Targeted Optimal Treatment Regime Learning Using Summary Statistics. *Biometrika*, 03 2023. ISSN 1464-3510. doi: 10.1093/biomet/asad020. URL <https://doi.org/10.1093/biomet/asad020>. In press.
- C. Del Ninno. *The 1998 Floods in Bangladesh: Disaster Impacts, Household Coping Strategies, and Response*, volume 122. The International Food Policy Research Institute, 2001.
- B. Efron. Nonparametric estimates of standard error: the jackknife, the bootstrap and other methods. *Biometrika*, 68:589–599, 1981.
- B. Efron and D. Feldman. Compliance as an explanatory variable in clinical trials. *Journal of the American Statistical Association*, 86:9–17, 1991.
- J. Fan. Local linear regression smoothers and their minimax efficiencies. *The Annals of Statistics*, pages 196–216, 1993.
- J. Fan and H. Peng. Nonconcave penalized likelihood with a diverging number of parameters. *The Annals of Statistics*, 32:928–961, 2004.
- D. Follmann. Augmented designs to assess immune response in vaccine trials. *Biometrics*, 62: 1161–1169, 2006.
- D. A. Follmann. On the effect of treatment among would-be treatment compliers: An analysis of the multiple risk factor intervention trial. *Journal of the American Statistical Association*, 95: 1101–1109, 2000.
- L. Forastiere, A. Mattei, and P. Ding. Principal ignorability in mediation analysis: through and beyond sequential ignorability. *Biometrika*, 105:979–986, 2018.

- C. E. Frangakis and D. B. Rubin. Principal stratification in causal inference. *Biometrics*, 58:21–29, 2002.
- P. Frumento, F. Mealli, B. Pacini, and D. B. Rubin. Evaluating the effect of training on wages in the presence of noncompliance, nonemployment, and missing outcome data. *Journal of the American Statistical Association*, 107:450–466, 2012.
- P. B. Gilbert and M. G. Hudgens. Evaluating candidate principal surrogate endpoints. *Biometrics*, 64(4):1146–1154, 2008.
- J. Hahn. On the role of the propensity score in efficient semiparametric estimation of average treatment effects. *Econometrica*, 66:315–331, 1998.
- S. M. Hammer, D. A. Katzenstein, M. D. Hughes, H. Gundacker, R. T. Schooley, R. H. Haubrich, W. K. Henry, M. M. Lederman, J. P. Phair, M. Niu, et al. A trial comparing nucleoside monotherapy with combination therapy in hiv-infected adults with cd4 cell counts from 200 to 500 per cubic millimeter. *The New England Journal of Medicine*, 335:1081–1090, 1996.
- L. Han, X. Wang, and T. Cai. Identifying surrogate markers in real-world comparative effectiveness research. *Statistics in Medicine*, 41(26):5290–5304, 2022.
- J. Hill, J. Waldfogel, and J. Brooks-Gunn. Differential effects of high-quality child care. *Journal of Policy Analysis and Management: The Journal of the Association for Public Policy Analysis and Management*, 21:601–627, 2002.
- M. D. Hughes, M. J. Daniels, M. A. Fischl, S. Kim, and R. T. Schooley. Cd4 cell count as a surrogate endpoint in hiv clinical trials: a meta-analysis of studies of the aids clinical trials group. *AIDS*, 12:1823–1832, 1998.
- K. Imai, L. Keele, and T. Yamamoto. Identification, inference and sensitivity analysis for causal mediation effects. *Statistical Science*, 25:51–71, 2010.
- G. W. Imbens. Nonparametric estimation of average treatment effects under exogeneity: A review. *Review of Economics and Statistics*, 86:4–29, 2004.
- P. Jaworski, F. Durante, W. K. Hardle, and T. Rychlik. *Copula Theory and Its Applications*, volume 198. Springer, 2010.
- Z. Jiang and P. Ding. Identification of causal effects within principal strata using auxiliary variables. *Statistical Science*, 36:493–508, 2021.
- Z. Jiang, S. Yang, P. Ding, et al. Multiply robust estimation of causal effects under principal ignorability. *Journal of the Royal Statistical Society: Series B (Methodological)*, 84:1423–1445, 2022.

- H. Jin and D. B. Rubin. Principal stratification for causal inference with extended partial compliance. *Journal of the American Statistical Association*, 103:101–111, 2008.
- B. Jo and E. A. Stuart. On the use of propensity scores in principal causal effect estimation. *Statistics in Medicine*, 28:2857–2875, 2009.
- E. H. Kennedy. Semiparametric theory and empirical processes in causal inference. In *Statistical Causal Inferences and Their Applications in Public Health Research*, pages 141–167. Springer, 2016.
- E. H. Kennedy, Z. Ma, M. D. McHugh, and D. S. Small. Non-parametric methods for doubly robust estimation of continuous treatment effects. *Journal of the Royal Statistical Society: Series B (Methodological)*, 79:1229–1245, 2017.
- C. Kim, M. J. Daniels, J. W. Hogan, C. Choirat, and C. M. Zigler. Bayesian methods for multiple mediators: Relating principal stratification and causal mediation in the analysis of power plant emission controls. *The Annals of Applied Statistics.*, 13:1927, 2019.
- C. Kim, L. R. Henneman, C. Choirat, C. M. Zigler, et al. Health effects of power plant emissions through ambient air quality. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 183:1677–1703, 2020.
- I. Lipkovich, B. Ratitch, Y. Qu, X. Zhang, M. Shan, and C. Mallinckrodt. Using principal stratification in analysis of clinical trials. *Statistics in Medicine*, 41:3837–3877, 2022.
- S. Lu, Z. Jiang, and P. Ding. Principal stratification with continuous post-treatment variables: Nonparametric identification and semiparametric estimation. *arXiv preprint arXiv:2309.12425*, 2023.
- A. R. Luedtke and M. J. Van Der Laan. Statistical inference for the mean outcome under a possibly non-unique optimal treatment strategy. *The Annals of statistics*, 44:713–742, 2016.
- A. Mattei and F. Mealli. Augmented designs to assess principal strata direct effects. *Journal of the Royal Statistical Society: Series B (Methodological)*, 73:729–752, 2011.
- A. Mattei, L. Forastiere, and F. Mealli. Assessing principal causal effects using principal score methods. In *Handbook of Matching and Weighting Adjustments for Causal Inference*, pages 313–348. Chapman and Hall/CRC, 2023.
- P. McCullagh and J. A. Nelder. *Generalized Linear Models*. Routledge, 2019.
- F. Mealli and A. Mattei. A refreshing account of principal stratification. *The international journal of biostatistics*, 8(1), 2012.
- R. B. Nelsen. *An Introduction to Copulas*. Springer science & business media, 2007.

- L. Parast, M. M. McDermott, and L. Tian. Robust estimation of the proportion of treatment effect explained by surrogate marker information. *Statistics in medicine*, 35(10):1637–1653, 2016.
- D. Pollard. *Convergence of Stochastic Processes*. Springer Science & Business Media, 2012.
- J. Robins, L. Li, E. Tchetgen, A. van der Vaart, et al. Higher order influence functions and minimax estimation of nonlinear functionals. *Probability and Statistics: Essays in Honor of David A. Freedman*, 66:335–421, 2008.
- J. M. Robins and S. Greenland. Identifiability and exchangeability for direct and indirect effects. *Epidemiology*, 3:143–155, 1992.
- P. R. Rosenbaum and D. B. Rubin. The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70:41–55, 1983.
- D. B. Rubin. Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of educational Psychology*, 66:688–701, 1974.
- D. B. Rubin. Randomization analysis of experimental data: The fisher randomization test comment. *Journal of the American statistical association*, 75(371):591–593, 1980.
- D. B. Rubin. Direct and indirect causal effects via potential outcomes. *Scandinavian Journal of Statistics*, 31:161–170, 2004.
- D. B. Rubin. Causal inference through potential outcomes and principal stratification: application to studies with “censoring” due to death. *Statistical Science*, 21:299–309, 2006.
- S. L. Schwartz, F. Li, and F. Mealli. A bayesian semiparametric approach to intermediate variables in causal inference. *Journal of the American Statistical Association*, 106:1331–1344, 2011.
- M. Sklar. Fonctions de repartition an dimensions et leurs marges. *Publications de L’Institut de Statistique de L’Universit  de Paris*, 8:229–231, 1959.
- E. A. Stuart, C. P. Bradshaw, and P. J. Leaf. Assessing the generalizability of randomized trial results to target populations. *Prevention Science*, 16:475–485, 2015.
- E. J. T. Tchetgen and I. Shpitser. Semiparametric theory for causal mediation analysis: efficiency bounds, multiple robustness, and sensitivity analysis. *The Annals of Statistics*, 40:1816–1845, 2012.
- A. B. Tsybakov. *Introduction to nonparametric estimation*. Springer, 2008.
- A. W. Vaart and J. A. Wellner. Weak convergence. In *Weak Convergence and Empirical Processes*, pages 16–28. Springer, 1996.
- M. J. Van der Laan, E. C. Polley, and A. E. Hubbard. Super learner. *Statistical Applications in Genetics and Molecular Biology*, 6:25, 2007.

- A. Van der Vaart and M. J. van der Laan. Estimating a survival distribution with current status data and high-dimensional covariates. *The International Journal of Biostatistics*, 2(1), 2006.
- A. W. Van der Vaart. *Asymptotic Statistics*, volume 3. Cambridge university press, 2000.
- T. J. VanderWeele. Principal stratification—uses and limitations. *The International Journal of Biostatistics*, 7:1–14, 2011.
- X. Wang, L. Parast, L. Tian, and T. Cai. Model-free approach to quantifying the proportion of treatment effect explained by a surrogate marker. *Biometrika*, 107(1):107–122, 2020.
- Y. Wang and J. M. Taylor. A measure of the proportion of treatment effect explained by a surrogate marker. *Biometrics*, 58(4):803–812, 2002.
- L. Wasserman. *All of Nonparametric Statistics*. Springer Science & Business Media, 2006.
- T. Westling, P. Gilbert, and M. Carone. Causal isotonic regression. *Journal of the Royal Statistical Society: Series B (Methodological)*, 82:719–747, 2020.
- C. M. Zigler and T. R. Belin. A bayesian approach to improved estimation of causal effect predictiveness for a principal surrogate endpoint. *Biometrics*, 68:922–932, 2012.

Supplementary Material for “Semiparametric Localized Principal Stratification Analysis with Continuous Strata”

The Supplementary File is organized as follows. Section S1 presents short-hand notations used throughout the Supplementary File. Section S2 contains technical details of Remark 3. An illustrative example demonstrating the verification of CATE as a perfectly localized estimand is presented in Section S3. Sections S4–S6 include lemmas and proofs. Section S8 reports additional results and figures for our real-data applications.

S1 Summary of notations

For simplicity of presentation, we summarize the short-hand notations used in the proofs.

- $\mathbb{I}(\mathcal{E})$ is the indicator function such that $\mathbb{I}(\mathcal{E}) = 1$ when \mathcal{E} happens and 0 otherwise.
- $\xi_{1,u^*}(x, z, m) = \{\pi_z(x) f_{zm}(x)\}^{-1} (-1)^{z+1} \gamma_1^{(z)}(x, m)$;
- $\gamma_{1,u^*}^{(1)}(x, m) = \int_{\mathbb{R}} k_{u^*}(m, m_0) e_{(m, m_0)}(x) dm_0$;
- $\gamma_{1,u^*}^{(0)}(x, m) = \int_{\mathbb{R}} k_{u^*}(m_1, m) e_{(m_1, m)}(x) dm_1$;
- $\tilde{\xi}_{a,u^*}(x, z, m) = \xi_{a,u^*}(x, z, m) - \mathbb{E}\{\xi_{a,u^*}(X, Z, M) \mid X = x, Z = z\}$ for $a = 2, 3$;
- $\xi_{2,u^*}(x, z, m) = \{\pi_z(x)\}^{-1} \int_{\mathbb{R}^2} \mathbb{1}(m \leq m_z) \{\mu_{1m_1}(x) - \mu_{0m_0}(x)\} k_{u^*}(u) c_\rho^{(z)} \{F_{1m_1}(x), F_{0m_0}(x)\} \{\prod_{z \in \{0,1\}} f_{zm_z}(x)\} du$;
- $\xi_{3,u^*}(x, z, m) = \{\pi_z(x)\}^{-1} \gamma_{2,u^*}^{(z)}(x, m)$;
- $\gamma_{2,u^*}^{(1)}(x, m) = \int_{\mathbb{R}} \{\mu_{1m}(x) - \mu_{0m_0}(x)\} k_{u^*}(m, m_0) c_\rho \{F_{1m}(x), F_{0m_0}(x)\} f_{0m_0}(x) dm_0$;
- $\gamma_{2,u^*}^{(0)}(x, m) = \int_{\mathbb{R}} \{\mu_{1m_1}(x) - \mu_{0m}(x)\} k_{u^*}(m_1, m) c_\rho \{F_{1m_1}(x), F_{0m}(x)\} f_{1m_1}(x) dm_1$;
- $\tilde{\xi}'_{a,u^*}(x, z, m) = \xi'_{a,u^*}(x, z, m) - \mathbb{E}\{\xi'_{a,u^*}(X, Z, M) \mid X = x, Z = z\}$ for $a = 2, 3$;
- $\xi'_{2,u^*}(x, z, m) = \{\pi_z(x)\}^{-1} \int_{\mathbb{R}^2} \mathbb{1}(m \leq m_z) k_{u^*}(u) c_\rho^{(z)} \{F_{1m_1}(x), F_{0m_0}(x)\} \{\prod_{z \in \{0,1\}} f_z(m_z \mid x)\} du$;
- $\xi'_{3,u^*}(x, z, m) = \{\pi_z(x)\}^{-1} \gamma'^{(z)}_{2,u^*}(x, m)$;
- $\gamma'^{(1)}_{2,u^*}(x, m) = \int_{\mathbb{R}} k_{u^*}(m, m_0) c_\rho \{F_{1m}(x), F_{0m_0}(x)\} f_{0m_0}(x) dm_0$;
- $\gamma'^{(0)}_{2,u^*}(x, m) = \int_{\mathbb{R}} k_{u^*}(m_1, m) c_\rho \{F_{1m_1}(x), F_{0m}(x)\} f_{1m_1}(x) dm_1$;

- \mathcal{A}_{1,u^*}^* and \mathcal{A}_{2,u^*}^* are defined as follows:

$$\begin{aligned}\mathcal{A}_{1,u^*}^* &= \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X) \{f'_{zm_z}(X)\}^\top du \right] \\ &\quad + \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[c_\rho^{(z)} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1m_1}(X) f_{0m_0}(X) \{F'_{zm_z}(X)\}^\top \right] du \right], \\ \mathcal{A}_{2,u^*}^* &= \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X) \{f'_{zm_z}(X)\}^\top du \right] \\ &\quad + \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} c_\rho^{(z)} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1m_1}(X) f_{0m_0}(X) \{F'_{zm_z}(X)\}^\top du \right];\end{aligned}$$

- \mathcal{A}_{1,u^*} and \mathcal{A}_{2,u^*} are defined as follows:

$$\begin{aligned}\mathcal{A}_{1,u^*} &= \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X) \{f'_{zm_z}(X)\}^\top du \right] \\ &\quad + \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[c_\rho^{(z)} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1m_1}(X) f_{0m_0}(X) \{F'_{zm_z}(X)\}^\top \right] du \right],\end{aligned}\tag{S1}$$

$$\begin{aligned}\mathcal{A}_{2,u^*} &= \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{ \bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X) \} c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X) \{f'_{zm_z}(X)\}^\top du \right] \\ &\quad + \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{ \bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X) \} c_\rho^{(z)} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1m_1}(X) f_{0m_0}(X) \{F'_{zm_z}(X)\}^\top du \right] \\ &\quad + \sum_{j=1,2,3} \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{Y - \bar{\mu}_{ZM}(X)\} \right] \delta_{X,Z,M}^{(j)} \right];\end{aligned}\tag{S2}$$

- $\tilde{\phi}_p(x, z, m)$ and $\tilde{\phi}_{np}(x, z, m)$ are defined as follows:

$$\tilde{\phi}_p(x, z, m) = \frac{\mathcal{A}_{2,u^*}^* - \tau_{u^*} \mathcal{A}_{1,u^*}^*}{\mathbb{E} \llbracket e_{\bullet\star}(X) \rrbracket_{u^*}} I_{\beta^*}^{-1} s_{zm}(x | \beta^*),\tag{S3}$$

$$\tilde{\phi}_{np}(x, z, m) = \frac{\tilde{\xi}_{2,u^*}(x, z, m) + \tilde{\xi}_{3,u^*}(x, z, m) - \tau_{u^*} \{ \tilde{\xi}_{2,u^*}^t(x, z, m) + \tilde{\xi}_{3,u^*}^t(x, z, m) \}}{\mathbb{E} \llbracket e_{\bullet\star}(X) \rrbracket_{u^*}};\tag{S4}$$

- $\bar{\sigma}_{u^*}^2$ is defined as follows:

$$\begin{aligned}\bar{\sigma}_{u^*}^2 &= \mathbb{E} \left[\frac{\tilde{\xi}_{1,u^*}(X, Z, M) \{Y - \bar{\mu}_{ZM}(X)\} + \llbracket \{ \bar{\mu}_{1\bullet}(X) - \bar{\mu}_{0\bullet}(X) \} e_{\bullet\star}(X) \rrbracket_{u^*} - \tau_{u^*} \llbracket e_{\bullet\star}(X) \rrbracket_{u^*}}{\llbracket e_{\bullet\star}(X) \rrbracket_{u^*}} \right. \\ &\quad \left. + \frac{\mathcal{A}_{2,u^*} - \tau_{u^*} \mathcal{A}_{1,u^*}}{\llbracket e_{\bullet\star}(X) \rrbracket_{u^*}} \psi(X, Z, M) \right]^2;\end{aligned}\tag{S5}$$

- $\delta_{X,1,M,u^*}^{(1)}$ and $\delta_{X,1,M,u^*}^{(3)}$ are defined as follows:

$$\delta_{X,1,M,u^*}^{(1)} = \{F'_{1M}(X | \beta^*)\}^\top \int_{\mathbb{R}} k_{u^*}(M, m_0) c_\rho^{(1)} \{F_{1M}(X), F_{0m_0}(X)\} f_{0m_0}(X) dm_0,$$

$$\begin{aligned}
\delta_{X,1,M,u^*}^{(2)} &= \int_{\mathbb{R}} k_{u^*}(M, m_0) c_{\rho}^{(0)} \{F_{1M}(X), F_{0m_0}(X)\} \{F'_{0m_0}(X | \beta^*)\}^{\top} f_{0m_0}(X) dm_0, \\
\delta_{X,1,M,u^*}^{(3)} &= \int_{\mathbb{R}} k_{u^*}(M, m_0) c_{\rho} \{F_{1M}(X), F_{0m_0}(X)\} \{f'_{0m_0}(X | \beta^*)\}^{\top} dm_0; \tag{S6}
\end{aligned}$$

- $\delta_{X,0,M,u^*}^{(1)} - \delta_{X,0,M,u^*}^{(3)}$ are defined as follows:

$$\begin{aligned}
\delta_{X,0,M,u^*}^{(1)} &= \{F'_{0M}(X | \beta^*)\}^{\top} \int_{\mathbb{R}} k_{u^*}(m_1, M) c_{\rho}^{(0)} \{F_{1m_1}(X), F_{0M}(X)\} f_{1m_1}(X) dm_1, \\
\delta_{X,0,M,u^*}^{(2)} &= \int_{\mathbb{R}} k_{u^*}(m_1, M) c_{\rho}^{(1)} \{F_{1m_1}(X), F_{0M}(X)\} \{F'_{1m_1}(X | \beta^*)\}^{\top} f_{1m_1}(X) dm_1, \\
\delta_{X,0,M,u^*}^{(3)} &= \int_{\mathbb{R}} k_{u^*}(m_1, M) c_{\rho} \{F_{1m_1}(X), F_{0M}(X)\} \{f'_{1m_1}(X | \beta^*)\}^{\top} dm_1. \tag{S7}
\end{aligned}$$

S2 Comparison with identification assumptions in mediation analysis

Recall the setting of mediation analysis in Section 2.2. In particular, the potential outcome notation Y_z and potential outcome notation $Y_{z,m}$ in mediation analysis are linked by

$$Y_z = Y_{z,M_z}, \quad z = 0, 1, \tag{S8}$$

with no conflict of settings; See Forastiere et al. (2018, §2.1). We show that Assumption 2 is weaker than the following two common identification assumptions in mediation analysis.

- The *strong principal ignorability* assumes

$$Y_{z,m} \perp\!\!\!\perp U | X \quad \text{for any } z = 0, 1 \tag{S9}$$

and $m \in \mathbb{R}$. When M is binary, Forastiere et al. (2018) show that this assumption is equivalent to the sequential ignorability under monotonicity. When M is continuous, it implies Assumption 2 as

$$\begin{aligned}
\mathbb{E}\{Y_z | X, U = (m_1, m_0)\} &= \mathbb{E}\{Y_{z,m_z} | X, U = (m_1, m_0)\} \quad (\text{by (S8) and } M_z = m_z) \\
&= \mathbb{E}(Y_{z,m_z} | X) \quad (\text{by (S9)}) \\
&= \mathbb{E}(Y_{z,m_z} | X, M_z = m_z) \quad (\text{by (S9)}) \\
&= \mathbb{E}(Y_{z,M_z} | X, M_z = m_z) = \mathbb{E}(Y_z | X, M_z = m_z) \quad (\text{by (S8)}).
\end{aligned}$$

- The *sequential ignorability* (Imai et al., 2010, Tchetgen and Shpitser, 2012) assumes for any $z, z' \in \{0, 1\}$, $m \in \mathbb{R}$:

$$(i). (Y_{z',m}, M_z) \perp\!\!\!\perp Z \mid X; \quad (\text{S10})$$

$$(ii). Y_{z',m} \perp\!\!\!\perp M \mid Z = z, X, \quad (\text{S11})$$

which imply Assumption 2. This is because for any $z \in \{0, 1\}$ and $(m_1, m_0) \in \mathbb{R}^2$,

$$\begin{aligned} \mathbb{E}(Y_z \mid X, M_z = m_z) &= \mathbb{E}(Y_{z,m_z} \mid X, Z = 1 - z, M_z = m_z) \quad (\text{by (S8) and (S10)}) \\ &= \mathbb{E}(Y_{z,m_z} \mid X, Z = 1 - z, M_z = m_z, M = m_{1-z}) \quad (\text{by (S11)}) \\ &= \mathbb{E}(Y_{z,m_z} \mid X, Z = 1 - z, M_z = m_z, M_{1-z} = m_{1-z}) \\ &= \mathbb{E}\{Y_{z,m_z} \mid X, U = (m_1, m_0)\} \quad (\text{by (S10)}) \\ &= \mathbb{E}\{Y_{z,M_z} \mid X, U = (m_1, m_0)\} = \mathbb{E}\{Y_z \mid X, U = (m_1, m_0)\}. \end{aligned}$$

- Parast et al. (2016) and Wang et al. (2020) consider the RCT setting where Z is randomly assigned and no covariates X are collected. The identification assumption of Parast et al. (2016) assumes

$$Y_1 \perp\!\!\!\perp M_0 \mid M_1 \text{ and } Y_0 \perp\!\!\!\perp M_1 \mid M_0. \quad (\text{S12})$$

The identification assumption of Wang et al. (2020) assumes

$$(Y_1, M_1) \perp\!\!\!\perp (Y_0, M_0). \quad (\text{S13})$$

Under either (S12) or (S13), one can see that $Y_z \mid M_z = m_z$ and $Y_z \mid U = u$ share the same distribution and thus $\mathbb{E}(Y_z \mid M_z = m_z) = \mathbb{E}\{Y_z \mid U = (m_1, m_0)\}$, which verifies Assumption 2.

S3 Verification of CATE as a perfectly localized functional

In (1), let $W = (X, Z)$, $D = X$, and x_0 be a specific individual level of X . Let $\gamma^*(x, z) = \mathbb{E}(Y \mid X = x, Z = z)$ and $k_h(x \mid x_0)$ be some regular kernel centered at x_0 . Denote

$$m(w, y, \gamma^*, k_h) = k_h(x \mid x_0)\{\gamma^*(x, 1) - \gamma^*(x, 0)\}.$$

We show that CATE with $X = x_0$, namely $\vartheta_0 = \mathbb{E}(Y_1 - Y_0 \mid X = x_0)$, can be represented through (1) and thus is a perfectly localized function. In particular, under the assumption of selection on observables, we have $\vartheta_0 = \mathbb{E}\{\gamma^*(X, 1) - \gamma^*(X, 0) \mid X = x_0\} = \lim_{h \rightarrow 0} \mathbb{E}[k_h(X \mid x_0)\{\gamma^*(X, 1) - \gamma^*(X, 0)\}] = \lim_{h \rightarrow 0} \vartheta_h$, which complies with (1).

S4 Proof of Proposition 1

S4.1 Regularity conditions

We list the three regularity conditions for Proposition 1 as follows.

Assumption S5 *The second-order partial derivative of $\mu_{zm}(x)$ with respect to m is uniformly bounded for all $x \in \mathbb{X}$, $m_z \in \mathbb{R}^2$ and $z = 0, 1$.*

Assumption S6 *All second-order partial derivatives of $f\{(M_1, M_0) = u \mid X = x\} = e_u(x)$ with respect to u , are uniformly bounded for all $x \in \mathbb{X}$, $u \in \mathbb{R}^2$ and $z = 0, 1$.*

Assumption S7 *$\int_{\mathbb{R}^2} \mathcal{K}(\delta_1, \delta_0) d\delta_1 d\delta_0 = 1$, $\int_{\mathbb{R}^2} \delta_z \mathcal{K}(\delta_1, \delta_0) d\delta_1 d\delta_0 = 0$, and $\int_{\mathbb{R}^2} |\delta_1|^{\alpha_1} |\delta_0|^{\alpha_0} \mathcal{K}(\delta_1, \delta_0) d\delta_1 d\delta_0 < +\infty$ for all integers α_1, α_0 such that $\alpha_1 + \alpha_0 \leq 4$ and $z = 0, 1$.*

The bounded second-order partial derivatives of $\mu_{zm}(x)$ and $e_u(x)$ in Assumptions S5 and S6 are standard in the nonparametric kernel regression literature. Assumption S7 for $\mathcal{K}(\cdot, \cdot)$ is also mild and can be typically satisfied by the two-dimensional Gaussian Kernel, box kernel, Epanechnikov kernel, among others. See e.g. Fan (1993) and Wasserman (2006).

S4.2 Main proof of Proposition 1

Some technical arguments are standard in the nonparametric kernel regression literature for bias control, and thus we will omit the corresponding details; See, e.g. Fan (1993) for more discussions. For the ease of exposition, denote $\Delta_u(x) = \mu_{1m_1}(x) - \mu_{0m_0}(x)$ for any $u = (m_1, m_0) \in \mathbb{R}^2$ and $x \in \mathbb{R}^{d_X}$.

By definition, one has

$$\begin{aligned} \tau_{u^*} &= \int_{\mathbb{R}^{d_X+2}} \Delta_u(x) \frac{k_{u^*}(u)}{\mathbb{E}\{k_{u^*}(U)\}} e_u(x) f(x) dm_1 dm_0 dx \\ &= \int_{\mathbb{R}^{d_X+2}} \Delta_{u^*+(\delta_1, \delta_0)}(x) \frac{k_{u^*}\{u^* + (\delta_1, \delta_0)\}}{\mathbb{E}\{k_{u^*}(U)\}} e_{u^*+(\delta_1, \delta_0)}(x) f(x) d\delta_1 d\delta_0 dx \end{aligned}$$

$$\begin{aligned}
&= \int_{\mathbb{R}^{d_X+2}} \Delta_{u^*+h\delta'}(x) \frac{k_{u^*}\{u^*+h\delta'\}}{\mathbb{E}\{k_{u^*}(U)\}} e_{u^*+h\delta'}(x) f(x) h^2 d\delta'_1 d\delta'_0 dx \\
&= \int_{\mathbb{R}^{d_X+2}} \Delta_{u^*+h\delta'}(x) \frac{\mathcal{K}(\delta'_1, \delta'_0)}{\mathbb{E}\{k_{u^*}(U)\}} e_{u^*+h\delta'}(x) f(x) d\delta'_1 d\delta'_0 dx \\
&= \int_{\mathbb{R}^{d_X}} \frac{f(x)}{\mathbb{E}\{k_{u^*}(U)\}} \int_{\mathbb{R}^2} \Delta_{u^*+h\delta'}(x) \mathcal{K}(\delta'_1, \delta'_0) e_{u^*+h\delta'}(x) d\delta'_1 d\delta'_0 dx, \tag{S14}
\end{aligned}$$

where we denote $\delta' = (\delta'_1, \delta'_0)$. By Taylor expansion, one has for any given $x \in \mathbb{X}$,

$$\begin{aligned}
\Delta_{u^*+h\delta'}(x) &= \mu_{1m_1^*}(x) + h\delta'_1 \frac{\partial \mu_{1m_1}(x)}{\partial m_1} \Big|_{m_1=m_1^*} + \frac{(h\delta'_1)^2}{2} \frac{\partial^2 \mu_{1m_1}(x)}{\partial^2 m_1} \Big|_{m_1=m_1^*+h_1\delta'_1} \\
&\quad - \mu_{0m_0^*}(x) - h\delta'_0 \frac{\partial \mu_{0m_0}(x)}{\partial m_0} \Big|_{m_0=m_0^*} - \frac{(h\delta'_0)^2}{2} \frac{\partial^2 \mu_{0m_0}(x)}{\partial^2 m_0} \Big|_{m_0=m_0^*+h_0\delta'_0} \tag{S15}
\end{aligned}$$

where $|h_z| \leq h$ and $\text{sgn}(h_z) = \text{sgn}(h)$ for $z = 0, 1$; $\text{sgn}(\cdot)$ is the sign function. Similarly, we have,

$$\begin{aligned}
e_{u^*+h\delta'}(x) &= e_{u^*}(x) + h\delta'_1 \frac{\partial e_u(x)}{\partial m_1} \Big|_{u=u^*} + h\delta'_0 \frac{\partial e_u(x)}{\partial m_0} \Big|_{u=u^*} \\
&\quad + \frac{(h\delta'_1)^2}{2} \frac{\partial^2 e_u(x)}{\partial^2 m_1} \Big|_{u=u^*+h_1\delta'} + \frac{(h\delta'_0)^2}{2} \frac{\partial^2 e_u(x)}{\partial^2 m_0} \Big|_{u=u^*+h_0\delta'} + \frac{h^2 \delta'_1 \delta'_0}{2} \frac{\partial^2 e_u(x)}{\partial m_1 \partial m_0} \Big|_{u=u^*+h_{10}\delta'}, \tag{S16}
\end{aligned}$$

for some $|h_{10}| \leq h$ and $\text{sgn}(h_{10}) = \text{sgn}(h)$. Substituting the right-hand side of (S14) with (S15) and (S16), we have

$$\begin{aligned}
\tau_{u^*} &= \int_{\mathbb{R}^{d_X}} \frac{f(x)}{\mathbb{E}\{k_{u^*}(U)\}} \Delta_{u^*}(x) e_{u^*}(x) \int_{\mathbb{R}^2} \mathcal{K}(\delta'_1, \delta'_0) d\delta'_1 d\delta'_0 dx \\
&\quad + h \cdot \left\{ \int_{\mathbb{R}^{d_X}} \frac{f(x)}{\mathbb{E}\{k_{u^*}(U)\}} \Delta_{u^*}(x) \frac{\partial e_u(x)}{\partial m_1} \Big|_{u=u^*} \underbrace{\int_{\mathbb{R}^2} \delta'_1 \mathcal{K}(\delta'_1, \delta'_0) d\delta'_1 d\delta'_0 dx}_{=0} + \dots \right. \\
&\quad \left. - \int_{\mathbb{R}^{d_X}} \frac{f(x)}{\mathbb{E}\{k_{u^*}(U)\}} e_{u^*}(x) \delta'_0 \frac{\partial \mu_{0m_0}(x)}{\partial m_0} \Big|_{m_0=m_0^*} \underbrace{\int_{\mathbb{R}^2} \delta'_0 \mathcal{K}(\delta'_1, \delta'_0) d\delta'_1 d\delta'_0 dx}_{=0} \right\} \\
&\quad + \left\{ h^2 \int_{\mathbb{R}^{d_X}} \frac{f(x)}{\mathbb{E}\{k_{u^*}(U)\}} \left\{ \frac{\partial e_u(x)}{\partial m_1} \frac{\partial \mu_{1m_1}(x)}{\partial m_1} \right\} \Big|_{u=u^*} \int_{\mathbb{R}^2} (\delta'_1)^2 \mathcal{K}(\delta'_1, \delta'_0) d\delta'_1 d\delta'_0 dx + \dots \right. \\
&\quad \left. - \frac{h^4}{4} \int_{\mathbb{R}^{d_X}} \frac{f(x)}{\mathbb{E}\{k_{u^*}(U)\}} \int_{\mathbb{R}^2} \frac{\partial^2 \mu_{0m_0}(x)}{\partial^2 m_0} \Big|_{m_0=m_0^*+h_0\delta'_0} \frac{\partial^2 e_u(x)}{\partial m_1 \partial m_0} \Big|_{u=u^*+h_{10}\delta'} \delta'_1 (\delta'_0)^3 \mathcal{K}(\delta'_1, \delta'_0) d\delta'_1 d\delta'_0 dx \right\} \\
&= \int_{\mathbb{R}^{d_X}} \frac{f(x)}{\mathbb{E}\{k_{u^*}(U)\}} \Delta_{u^*}(x) e_{u^*}(x) dx + hR_1 + h^2 R_2 \\
&= \int_{\mathbb{R}^{d_X}} \frac{f(x)}{\mathbb{E}\{k_{u^*}(U)\}} \Delta_{u^*}(x) e_{u^*}(x) dx + \mathcal{O}(h^2), \tag{S17}
\end{aligned}$$

where the last equality is because that, by the regularity conditions and the fact that $[\mathbb{E}\{k_{u^*}(U)\}]^{-1} = \mathcal{O}(1)$ (see (S21) in the following), we can derive $R_1 = 0$ and $R_2 = \mathcal{O}(1)$ when $h \rightarrow 0$ with standard arguments for controlling the bias of the kernel nonparametric regression (Fan, 1993).

Next, we show $|1/\mathbb{E}\{k_{u^*}(U)\} - 1/e_{u^*}| = \mathcal{O}(h^2)$. By the Taylor expansion, one has

$$\left| \frac{1}{\mathbb{E}\{k_{u^*}(U)\}} - \frac{1}{e_{u^*}} \right| = \frac{1}{\xi^2} |\mathbb{E}\{k_{u^*}(U)\} - e_{u^*}| \quad (\text{S18})$$

for some ξ between $\mathbb{E}\{k_{u^*}(U)\}$ and e_{u^*} . Similar to (S14)–(S16), one has

$$\begin{aligned} \mathbb{E}\{k_{u^*}(U)\} &= \mathbb{E}[\mathbb{E}\{k_{u^*}(U) \mid X\}] \\ &= \int_{\mathbb{R}^{d_X}} f(x) \int_{\mathbb{R}^2} \frac{1}{h^2} \mathcal{K}\left(\frac{m_1 - m_1^*}{h}, \frac{m_0 - m_0^*}{h}\right) e_u(x) dm_1 dm_0 dx \\ &= \int_{\mathbb{R}^{d_X}} f(x) \int_{\mathbb{R}^2} \mathcal{K}(\delta'_1, \delta'_0) e_{u^* + h\delta'}(x) d\delta'_1 d\delta'_0 dx \\ &= \int_{\mathbb{R}^{d_X}} f(x) e_{u^*}(x) \int_{\mathbb{R}^2} \mathcal{K}(\delta'_1, \delta'_0) d\delta'_1 d\delta'_0 dx \\ &\quad + h \cdot \left\{ \int_{\mathbb{R}^{d_X}} f(x) \frac{\partial e_u(x)}{\partial m_1} \Big|_{u=u^*} \underbrace{\int_{\mathbb{R}^2} \delta'_1 \mathcal{K}(\delta'_1, \delta'_0) d\delta'_1 d\delta'_0 dx}_{=0} \right. \\ &\quad \left. + \int_{\mathbb{R}^{d_X}} f(x) \frac{\partial e_u(x)}{\partial m_0} \Big|_{u=u^*} \underbrace{\int_{\mathbb{R}^2} \delta'_0 \mathcal{K}(\delta'_1, \delta'_0) d\delta'_1 d\delta'_0 dx}_{=0} \right\} \\ &\quad + h^2 \cdot \left\{ \int_{\mathbb{R}^{d_X}} \frac{f(x)}{2} \int_{\mathbb{R}^2} \frac{\partial^2 e_u(x)}{\partial m_1^2} \Big|_{u=u^* + h_{10}\delta'} (\delta'_1)^2 \mathcal{K}(\delta'_1, \delta'_0) d\delta'_1 d\delta'_0 dx + \dots \right. \\ &\quad \left. + \int_{\mathbb{R}^{d_X}} \frac{f(x)}{2} \int_{\mathbb{R}^2} \frac{\partial^2 e_u(x)}{\partial m_1 \partial m_0} \Big|_{u=u^* + h_{10}\delta'} \delta'_1 \delta'_0 \mathcal{K}(\delta'_1, \delta'_0) d\delta'_1 d\delta'_0 dx \right\} \\ &= \int_{\mathbb{R}^{d_X}} f(x) e_{u^*}(x) dx + \mathcal{O}(h^2) \\ &= e_{u^*} + \mathcal{O}(h^2). \end{aligned} \quad (\text{S19})$$

Thus $|\mathbb{E}\{k_{u^*}(U)\} - e_{u^*}| = \mathcal{O}(h^2) = o(1)$ when $h \rightarrow 0$, which implies $\xi \rightarrow e_{u^*}$. Thus when $h \rightarrow 0$, we have,

$$\xi^{-1} \lesssim 1 \quad \text{and} \quad [\mathbb{E}\{k_{u^*}(U)\}]^{-1} \lesssim 1, \quad (\text{S21})$$

as $e_{u^*} > 0$ is fixed. Then (S18)–(S21) imply,

$$\left| \frac{1}{\mathbb{E}\{k_{u^*}(U)\}} - \frac{1}{e_{u^*}} \right| = \mathcal{O}(h^2). \quad (\text{S22})$$

Recalling (S17), one has

$$\begin{aligned} \tau_{u^*} &= \int_{\mathbb{R}^{d_X}} \Delta_{u^*}(x) \{e_{u^*}(x)/e_{u^*}\} f(x) dx + \left[\frac{1}{\mathbb{E}\{k_{u^*}(U)\}} - \frac{1}{e_{u^*}} \right] \int_{\mathbb{R}^{d_X}} \Delta_{u^*}(x) e_{u^*}(x) f(x) dx + \mathcal{O}(h^2) \\ &= \int_{\mathbb{R}^{d_X}} \Delta_{u^*}(x) f(X = x | U = u^*) dx + \underbrace{\left[\frac{1}{\mathbb{E}\{k_{u^*}(U)\}} - \frac{1}{e_{u^*}} \right] e_{u^*} \int_{\mathbb{R}^{d_X}} \Delta_{u^*}(x) f(X = x | U = u^*) dx}_{=R_3} \\ &\quad + \mathcal{O}(h^2), \end{aligned} \quad (\text{S23})$$

where we note $\{e_{u^*}(x)/e_{u^*}\} f(x) = f\{(M_1, M_0) = u^* | X = x\} f(X = x) / f\{(M_1, M_0) = u^*\} = f\{X = x | U = u^*\}$. Furthermore, under Assumptions 1–2, one has

$$\begin{aligned} \int_{\mathbb{R}^{d_X}} \Delta_{u^*}(x) f\{X = x | (M_1, M_0) = u^*\} dx &= \mathbb{E}\{\mu_{1m_1^*}(X) - \mu_{0m_0^*}(X) | U = u^*\} \\ &= \tau_{u^*}^*, \end{aligned} \quad (\text{S24})$$

by an argument same as (S26) in the Proof of Theorem 1. Combining (S22)–(S24), one has

$$\begin{aligned} |R_3| &\leq \left| \frac{1}{\mathbb{E}\{k_{u^*}(U)\}} - \frac{1}{e_{u^*}} \right| \cdot e_{u^*} \int_{\mathbb{R}^{d_X}} \Delta_{u^*}(x) f\{X = x | U = u^*\} dx \\ &= \left| \frac{1}{\mathbb{E}\{k_{u^*}(U)\}} - \frac{1}{e_{u^*}} \right| \cdot e_{u^*} \tau_{u^*}^* \\ &= \mathcal{O}(h^2). \end{aligned} \quad (\text{S25})$$

Combining (S23)–(S25), we have $\tau_{u^*} = \tau_{u^*}^* + \mathcal{O}(h^2)$ and thus $|\tau_{u^*} - \tau_{u^*}^*| = \mathcal{O}(h^2)$ as desired. \square

S5 Proofs of main theorems

Throughout the proofs, we will use $f(\cdot)$ to denote the probability density function for continuous random variables. We will use the law of total expectation repeatedly and will mark the steps as

LOTE. We first recall some notations in the main text,

$$\begin{aligned}\pi_z(X) &= \mathbb{P}(Z = z | X), & e_u(X) &= f(U = u | X), \\ e_u &= \mathbb{E}\{e_u(X)\} = f(U = u), & \mu_{zm}(X) &= \mathbb{E}(Y | X, Z = z, M = m), \\ f_{zm}(X) &= f(M = m | X, Z = z), & F_{zm}(X) &= \mathbb{P}(M \leq m | X, Z = z),\end{aligned}$$

for $z = 0, 1$, $m \in \mathbb{R}$ and $u = (m_1, m_0)$. Under Assumptions 1 and 3, we have

$$\begin{aligned}e_u(X) &= c_\rho \{F(M_1 = m_1 | X), F(M_0 = m_0 | X)\} \prod_{z \in \{0,1\}} f(M_z = m_z | X) \\ &= c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} \prod_{z \in \{0,1\}} f_{zm_z}(X).\end{aligned}$$

□

S5.1 Proof of Theorem 1

We first introduce the following lemma from Jiang et al. (2022, Lemma S3).

Lemma S1 *Let A and B be two random variables with densities $f_A(\cdot)$ and $f_B(\cdot)$. For any function $h(\cdot)$ with $\mathbb{E}\{h(A)\} < \infty$, we have*

$$\mathbb{E}\{h(A)\} = \mathbb{E}\left\{\frac{f_A(B)}{f_B(B)}h(B)\right\}.$$

By the law of the expectation (LOTE), we have

$$\begin{aligned}\tau_{u^*}^* &= \mathbb{E}(Y_1 - Y_0 | U = u^*) \\ &= \mathbb{E}\left\{\mathbb{E}(Y_1 - Y_0 | X, U = u^*) | U = u^*\right\} \\ &= \mathbb{E}\left[\mathbb{E}\{Y_1 | X, U = u^*\} - \mathbb{E}\{Y_0 | X, U = u^*\} | U = u^*\right] \\ &= \mathbb{E}\left[\mathbb{E}\{Y_1 | X, M_1 = m_1^*\} - \mathbb{E}\{Y_0 | X, M_0 = m_0^*\} | U = u^*\right] \quad (\text{Assumption 2}) \\ &= \mathbb{E}\left[\mathbb{E}\{Y_1 | X, Z = 1, M_1 = m_1^*\} - \mathbb{E}\{Y_0 | X, Z = 0, M_0 = m_0^*\} | U = u^*\right] \quad (\text{Assumption 1}) \\ &= \mathbb{E}\left\{\mathbb{E}(Y | X, Z = 1, M = m_1^*) - \mathbb{E}(Y | X, Z = 0, M = m_0^*) | U = u^*\right\} \\ &= \mathbb{E}\{\mu_{1m_1^*}(X) - \mu_{0m_0^*}(X) | U = u^*\} \tag{S26}\end{aligned}$$

$$= \mathbb{E}\left[\frac{f(X | U = u^*)}{f(X)}\{\mu_{1m_1^*}(X) - \mu_{0m_0^*}(X)\}\right] \tag{S27}$$

$$\begin{aligned}
&= \mathbb{E}\left[\frac{f(U = u^* | X)f(X)}{f(U = u^*)f(X)}\{\mu_{1m_1^*}(X) - \mu_{0m_0^*}(X)\}\right] \\
&= \mathbb{E}\left[\frac{e_{u^*}(X)}{e_{u^*}}\{\mu_{1m_1^*}(X) - \mu_{0m_0^*}(X)\}\right],
\end{aligned}$$

where in (S27) we use Lemma S1 with $A = X | U = u^*$ and $B = X$. By Assumptions 1 and 3, $e_{u^*}(x)$ can be further identified as

$$\begin{aligned}
e_{u^*}(X) &= f(M_1 = m_1^*, M_0 = m_0^* | X) \\
&= c_\rho \{F(M_1 = m_1^* | X), F(M_0 = m_0^* | X)\} \prod_{z \in \{0,1\}} f(M_z = m_z^* | X) \\
&= c_\rho \{F(M_1 = m_1^* | X, Z = 1), F(M_0 = m_0^* | X, Z = 0)\} \prod_{z \in \{0,1\}} f(M_z = m_z^* | X, Z = z) \\
&\hspace{20em} \text{(Assumption 1)} \\
&= c_\rho \{F_{1m_1^*}(X), F_{0m_0^*}(X)\} \prod_{z \in \{0,1\}} f_{zm_z^*}(X). \tag{S28}
\end{aligned}$$

□

S5.2 Proof of Theorem 2

By the law of the expectation (LOTE), we have

$$\begin{aligned}
\tau_{u^*} &= \mathbb{E}\left[\{\mu_{1M_1}(X) - \mu_{0M_0}(X)\} \frac{k_{u^*}(U)}{\mathbb{E}\{k_{u^*}(U)\}}\right] \\
&= \frac{1}{\mathbb{E}\{k_{u^*}(U)\}} \cdot \mathbb{E}\left[\mathbb{E}\left[\{\mu_{1M_1}(X) - \mu_{1M_0}(X)\} k_{u^*}(U) \mid X\right]\right] \\
&= \frac{1}{\mathbb{E}\{k_{u^*}(U)\}} \cdot \mathbb{E}\left[\int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} k_{u^*}(u) e_u(X) du\right], \tag{S29}
\end{aligned}$$

We also have $\mathbb{E}\{k_{u^*}(U)\} = \mathbb{E}[\mathbb{E}\{k_{u^*}(U) \mid X\}] = \mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du\}$. Combining above results and recalling the definition of $\llbracket \cdot \rrbracket$ in (7), the identification formula follows, and $e_u(X)$ can be identified same as (S28).

□

S5.3 Proof of Theorem 3

S5.3.1 Preliminary and auxiliary lemmas

To obtain the EIFs, we consider some arbitrary one-dimensional parametric submodel of the whole data density, denoted by $\{f_\theta(V = v)\}_{\theta \in \mathbb{R}}$. Let $f_\theta(v)$ contain the truth $f(v)$ at $\theta = 0$, i.e., $f_0(v) = f(v)$. We use $s_\theta(\cdot)$ and $s(\cdot)$ to denote the corresponding score functions, e.g., $s_\theta(x) = \partial \log f_\theta(X = x) / \partial \theta$ and $s(x) = \partial \log f_\theta(X = x) / \partial \theta |_{\theta=0}$. The submodel likelihood is factorized as

$$\begin{aligned} & f_\theta(V = v) \\ = & f_\theta(X = x) f_\theta(Z = z | X = x) f_\theta(M = m | X = x, Z = z) f_\theta(Y = y | X = x, Z = z, M = m). \end{aligned} \tag{S30}$$

Under the scenario of nonparametric estimation, the score function under submodel can be decomposed as

$$s_\theta(v) = s_\theta(x) + s_\theta(z | x) + s_\theta(m | x, z) + s_\theta(y | x, z, m).$$

Then the tangent space is

$$\Lambda = H_1 \oplus H_2 \oplus H_3 \oplus H_4 \tag{S31}$$

where \oplus is the set sum and,

$$\begin{aligned} H_1 &= \{h(x) \mid \mathbb{E}\{h(X)\} = 0\}, \\ H_2 &= \{h(x, z) \mid \mathbb{E}\{h(X, Z) \mid X\} = 0\}, \\ H_3 &= \{h(x, z, m) \mid \mathbb{E}\{h(X, Z, M) \mid X, Z\} = 0\}, \\ H_4 &= \{h(x, z, m, y) \mid \mathbb{E}\{h(X, Z, M, Y) \mid X, Z, M\} = 0\}. \end{aligned}$$

Based on the semiparametric theory (see e.g. [Bickel et al. \(1993\)](#), [Jiang et al. \(2022\)](#)), we have that under the nonparametric scenario (Scenario (iii)) with a tangent space like (S31), EIF of target parameter τ_{u^*} is the unique function $\phi(\cdot)$ (up to the almost surely equivalence) satisfying,

$$\tau_{u^*, \theta} \Big|_{\theta=0} = \mathbb{E}\left\{\phi(V) \cdot s(V)\right\}. \tag{S32}$$

Under the scenario of semiparametric estimation with known or parametric \mathcal{M}_{ps} (Scenarios (i) and (ii)), due to the assumption that the distribution (or the parametric distribution model) of $M | X, T$ is already known, the corresponding tangent space is more restrictive in comparison with (S31). As a result, the $\phi(\cdot)$ satisfying (S32) is no longer unique. Any $\phi(\cdot)$ satisfying (S32) serves as an influence function (IF) of τ_{u^*} , but is not necessary to be an EIF. To show that the $\phi(\cdot)$ we derive under Scenarios (i) or (ii) are indeed EIFs, we further verify that the derived $\phi(\cdot)$ belong to the corresponding semiparametric tangent spaces; Technical details are deferred to Section S5.3.4. Different from the order of Theorem 3, during the proof, we first derive $\phi_{\text{np}}(x, z, m, y)$ under Scenario (iii) in Section S5.3.2, based on which we further derive the EIFs under Scenarios (ii) and (i) in Sections S5.3.3 and S5.3.4, respectively.

We list two technical lemmas that will be used in the main proof.

Lemma S2 *Suppose function $\xi = \xi(A_1, A_2)$, where A_1, A_2 are random variables belonging to V , and A_1, A_2 have no overlapping. We have*

$$\mathbb{E}\{\xi(A_1, A_2) \cdot s(A_1 | A_2)\} = \mathbb{E}[\{\xi(A_1, A_2) - \mathbb{E}\{\xi(A_1, A_2) | A_2\}\} \cdot s(V)].$$

Proof of Lemma S2. By property of score function, we have $s(A_1 | A_2) = s(A_1, A_2) - s(A_2)$. Then

$$\mathbb{E}\{\xi(A_1, A_2) \cdot s(A_1 | A_2)\} = \mathbb{E}\{\xi(A_1, A_2) \cdot s(A_1, A_2)\} - \mathbb{E}\{\xi(A_1, A_2) \cdot s(A_2)\}. \quad (\text{S33})$$

If (A_1, A_2) do not contain all the covariates in V , we let A_3 be the rest of the covariates in V such that $A_3 = V \setminus (A_1, A_2)$. For the first term of (S33), we note that

$$\begin{aligned} \mathbb{E}\{\xi(A_1, A_2) \cdot s(A_3 | A_1, A_2)\} &= \mathbb{E}\left[\mathbb{E}\{\xi(A_1, A_2) \cdot s(A_3 | A_1, A_2) | A_1, A_2\}\right] \quad (\text{S34}) \\ &= \mathbb{E}\left[\xi(A_1, A_2) \cdot \mathbb{E}\{s(A_3 | A_1, A_2) | A_1, A_2\}\right] \\ &= \mathbb{E}\{\xi(A_1, A_2) \cdot 0\} \\ &= 0, \end{aligned}$$

where the third equality is by the property of score function as $\mathbb{E}\{s(A_3 | A_1, A_2) | A_1, A_2\} = 0$.

Thus we have

$$\mathbb{E}\{\xi(A_1, A_2) \cdot s(A_1, A_2)\} = \mathbb{E}\{\xi(A_1, A_2) \cdot s(A_1, A_2)\} + \mathbb{E}\{\xi(A_1, A_2) \cdot s(A_3 | A_1, A_2)\}$$

$$\begin{aligned}
&= \mathbb{E}\left[\xi(A_1, A_2) \cdot \{s(A_1, A_2) + s(A_3 | A_1, A_2)\}\right] \\
&= \mathbb{E}\left\{\xi(A_1, A_2) \cdot s(V)\right\},
\end{aligned} \tag{S35}$$

where the first equality is by (S34). If (A_1, A_2) contain all covariates in V , then $\mathbb{E}\{\xi(A_1, A_2) \cdot s(A_1, A_2)\} = \mathbb{E}\{\xi \cdot s(V)\}$ trivially.

For the second term of (S33), we have

$$\begin{aligned}
\mathbb{E}\left\{\xi(A_1, A_2) \cdot s(A_2)\right\} &= \mathbb{E}\left[s(A_2) \cdot \mathbb{E}\{\xi(A_1, A_2) | A_2\}\right] \quad (\text{LOTE}) \\
&= \mathbb{E}\left[s(A_2) \cdot \mathbb{E}\{\xi(A_1, A_2) | A_2\}\right] + \mathbb{E}\left[s(A_1, A_3 | A_2) \cdot \mathbb{E}\{\xi(A_1, A_2) | A_2\}\right] \\
&= \mathbb{E}\left[\{s(A_2) + s(A_1, A_3 | A_2)\} \cdot \mathbb{E}(\xi | A_2)\right] \\
&= \mathbb{E}\left\{\mathbb{E}(\xi | A_2) \cdot s(V)\right\},
\end{aligned} \tag{S36}$$

where the second equality holds because

$$\begin{aligned}
\mathbb{E}\left[s(A_1, A_3 | A_2) \cdot \mathbb{E}\{\xi(A_1, A_2) | A_2\}\right] &= \mathbb{E}\left[\mathbb{E}\left[s(A_1, A_3 | A_2) \cdot \mathbb{E}\{\xi(A_1, A_2) | A_2\} | A_2\right]\right] \\
&= \mathbb{E}\left[\mathbb{E}\{\xi(A_1, A_2) | A_2\} \cdot \mathbb{E}\left\{s(A_1, A_3 | A_2) | A_2\right\}\right] \\
&= 0,
\end{aligned}$$

by the property of score function as $\mathbb{E}\{s(A_1, A_3 | A_2) | A_2\} = 0$. Note here if (A_1, A_2) contain all the covariates in V , A_3 can be empty. Combing (S33), (S35), (S36), the desired result follows immediately. □

Lemma S3 *Suppose N_θ and D_θ are differentiable with respect to $\theta \in \mathbb{R}$. If $\partial N_\theta / \partial \theta |_{\theta=0} = \mathbb{E}\{\phi_N(V)s(V)\}$ and $\partial D_\theta / \partial \theta |_{\theta=0} = \mathbb{E}\{\phi_D(V)s(V)\}$, then*

$$\frac{\partial}{\partial \theta} \frac{N_\theta}{D_\theta} \Big|_{\theta=0} = \mathbb{E}\left[\left\{\frac{1}{D_0} \phi_N(V) - \frac{N_0}{D_0^2} \phi_D(V)\right\} \cdot s(V)\right]. \tag{S37}$$

Proof of Lemma S3. See Lemma S2 in Jiang et al. (2022). □

S5.3.2 Derivation of $\phi_{\text{np}}(x, z, m, y)$ under Scenario (iii)

For the nonparametric scenario, by verifying condition (S32), we derive the EIFs of the denominator and numerator of τ_{u^*} 's identification formula in (8), respectively. We then combine these EIFs

together and derive the desired EIF of τ_{u^*} based on Lemma S3.

Step 1 (EIF of $\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\}$): Based on the submodel introduced before, we can write $\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\}$ as

$$\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\} = \mathbb{E}_\theta\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u^{(\theta)}(X)du\right\}\Big|_{\theta=0}. \quad (\text{S38})$$

In (S38), obtain the derivative with respect to θ ,

$$\begin{aligned} \frac{\partial}{\partial\theta}\mathbb{E}_\theta\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u^{(\theta)}(X)du\right\}\Big|_{\theta=0} &= \mathbb{E}_\theta\left\{\int_{\mathbb{R}^2} k_{u^*}(u)\frac{\partial}{\partial\theta}e_u^{(\theta)}(X)\Big|_{\theta=0}du\right\} \\ &\quad + \mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du \cdot s(X)\right\}. \end{aligned}$$

Now we take a careful look at $\partial e_u^{(\theta)}(x)/\partial\theta|_{\theta=0}$. By definition (3), we have

$$\begin{aligned} \frac{\partial}{\partial\theta}e_u^{(\theta)}(x)\Big|_{\theta=0} &= \frac{\partial}{\partial\theta}\left[c_\rho\left\{F_{1m_1}^{(\theta)}(x), F_{0m_0}^{(\theta)}(x)\right\} \prod_{z \in \{0,1\}} f_{zm_z}^{(\theta)}(x)\right]\Big|_{\theta=0} \\ &= \sum_{z'=1,0} c_\rho^{(z')}\left\{F_{1m_1}(x), F_{0m_0}(x)\right\}\left\{\prod_{z \in \{0,1\}} f_{zm_z}(x)\right\} \cdot \frac{\partial}{\partial\theta}F_{z'm_{z'}}^{(\theta)}(x)\Big|_{\theta=0} \\ &\quad + \sum_{z'=1,0} c_\rho\left\{F_{1m_1}(x), F_{0m_0}(x)\right\} \cdot \frac{\partial}{\partial\theta}f_{z'm_{z'}}^{(\theta)}(x)\Big|_{\theta=0} \cdot f_{1-z'm_{1-z'}}(x). \end{aligned}$$

For $z' = 0, 1$, since $F_{z'm_{z'}}^{(\theta)}(x) = \int_{-\infty}^{m_{z'}} f^{(\theta)}(M_{z'} = m | X = x)dm$, we have

$$\begin{aligned} &\frac{\partial}{\partial\theta}F_{z'm_{z'}}^{(\theta)}(x)\Big|_{\theta=0} \\ &= \int_{\mathbb{R}} \mathbb{I}(m \leq m_{z'}) \frac{\partial f^{(\theta)}(M_{z'} = m | X = x)}{\partial\theta}\Big|_{\theta=0} dm \\ &= \int_{\mathbb{R}} \mathbb{I}(m \leq m_{z'}) \frac{1}{f(M_{z'} = m | X = x)} \frac{\partial f^{(\theta)}(M_{z'} = m | X = x)}{\partial\theta}\Big|_{\theta=0} f(M_{z'} = m | X = x) dm \\ &= \int_{\mathbb{R}} \mathbb{I}(m \leq m_{z'}) \frac{1}{f(M = m | X, Z = z')} \frac{\partial f^{(\theta)}(M = m | X, Z = z')}{\partial\theta}\Big|_{\theta=0} f(M = m | X = x, Z = z') dm \\ &= \mathbb{E}\left\{\mathbb{I}(M \leq m_{z'}) s(M | X, Z) \Big| X, Z = z'\right\}, \end{aligned}$$

where the third equality is by $f(M_{z'} = m | X = x) = f(M = m | X = x, Z = z')$ under Assumption

1. And similarly,

$$\begin{aligned}
& \frac{\partial}{\partial \theta} f^{(\theta)}(M_{z'} = m_{z'} \mid X = x) \Big|_{\theta=0} \\
&= \frac{1}{f(M = m_{z'} \mid X = x, Z = z')} \left\{ \frac{\partial}{\partial \theta} f^{(\theta)}(M = m_{z'} \mid X = x, Z = z') \Big|_{\theta=0} \right\} f(M = m_{z'} \mid X = x, Z = z') \\
&= s(M = m_{z'} \mid X = x, Z = z') f(M = m_{z'} \mid X = x, Z = z').
\end{aligned}$$

Combining the results above, we have

$$\begin{aligned}
& \frac{\partial}{\partial \theta} \mathbb{E}_\theta \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u^{(\theta)}(X) du \right\} \Big|_{\theta=0} \\
&= \sum_{z'=1,0} \mathbb{E}_\theta \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho^{(z')} \{F_{1m_1}(X), F_{0m_0}(X)\} \left\{ \prod_{z \in \{0,1\}} f_{zm_z}(X) \right\} \cdot \frac{\partial}{\partial \theta} F_{z'm_{z'}}^{(\theta)}(X) \Big|_{\theta=0} du \right] \\
&\quad + \sum_{z'=1,0} \mathbb{E}_\theta \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} \cdot \frac{\partial}{\partial \theta} f_{z'm_{z'}}^{(\theta)}(X) \Big|_{\theta=0} \cdot f_{1-z'm_{1-z'}}(X) du \right] \\
&\quad + \mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \cdot s(X) \right\} \\
&= \sum_{z'=1,0} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho^{(z')} \{F_{1m_1}(X), F_{0m_0}(X)\} \left\{ \prod_{z \in \{0,1\}} f_{zm_z}(X) \right\} \right. \\
&\quad \cdot \mathbb{E} \left\{ \mathbb{I}(M \leq m_{z'}) s(M \mid X, Z = z') \mid X, Z = z' \right\} du \Big] \\
&\quad + \sum_{z'=1,0} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} \left\{ \prod_{z \in \{0,1\}} f_{zm_z}(X) \right\} s(M = m_{z'} \mid X, Z = z') du \right] \\
&\quad + \mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \cdot s(X) \right\}. \tag{S39}
\end{aligned}$$

By LOTE and Fubini's theorem, the first term on the right-hand side of (S39) equals to

$$\begin{aligned}
& \sum_{z'=1,0} \mathbb{E} \left[\mathbb{E} \left[s(M \mid X, Z = z') \int_{\mathbb{R}^2} \mathbb{1}(M \leq m_{z'}) k_{u^*}(u) c_\rho^{(z')} \{F_{1m_1}(X), F_{0m_0}(X)\} \right. \right. \\
&\quad \left. \left. \cdot \prod_{z \in \{0,1\}} f_{zm_z}(X) du \mid X, Z = z' \right] \right] \\
&= \sum_{z'=1,0} \mathbb{E} \left[\mathbb{E} \left[\frac{\mathbb{1}(Z = z') s(M \mid X, Z = z')}{\pi_{z'}(X)} \right. \right. \\
&\quad \left. \left. \cdot \int_{\mathbb{R}^2} \mathbb{1}(M \leq m_{z'}) k_{u^*}(u) c_\rho^{(z')} \{F_{1m_1}(X), F_{0m_0}(X)\} \prod_{z \in \{0,1\}} f_{zm_z}(X) du \mid X \right] \right] \\
&= \mathbb{E} \left[\sum_{z'=1,0} \frac{\mathbb{1}(Z = z') s(M \mid X, Z = z')}{\pi_{z'}(X)} \int_{\mathbb{R}^2} \mathbb{1}(M \leq m_{z'}) k_{u^*}(u) c_\rho^{(z')} \{F_{1m_1}(X), F_{0m_0}(X)\} \prod_{z \in \{0,1\}} f_{zm_z}(X) du \right]
\end{aligned}$$

$$\begin{aligned}
&= \mathbb{E} \left[s(M | X, Z) \cdot \frac{1}{\pi_Z(X)} \int_{\mathbb{R}^2} \mathbf{1}(M \leq m_Z) k_{u^*}(u) c_\rho^{(Z)} \{F_{1m_1}(X), F_{0m_0}(X)\} \prod_{z \in \{0,1\}} f_{zm_z}(X) du \right] \\
&= \mathbb{E} \left\{ \tilde{\xi}'_2(X, Z, M) \cdot s(V) \right\} \quad (\text{By Lemma S2 with } A_1 = M, A_2 = (X, Z)), \tag{S40}
\end{aligned}$$

recalling

$$\begin{aligned}
\xi'_2(X, Z, M) &= \frac{1}{\pi_Z(X)} \int_{\mathbb{R}^2} \mathbf{1}(M \leq m_Z) k_{u^*}(u) c_\rho^{(Z)} \{F_{1m_1}(X), F_{0m_0}(X)\} \prod_{z \in \{0,1\}} f_{zm_z}(X) du, \\
\tilde{\xi}'_2(X, Z, M) &= \xi_2(X, Z, M) - \mathbb{E} \{ \xi_2(X, Z, M) | X, Z \}.
\end{aligned}$$

Note here and in the following, for simplicity, we will omit the footnote u^* in $\xi_{2,u^*}(x, z, m)$, $\xi'_{2,u^*}(x, z, m)$, $\tilde{\xi}'_{2,u^*}(z, x, m)$, etc. By definition, for both $z' = 0$ or 1 we have

$$\gamma_2'^{(z')}(M = m_{z'}, X) = \int_{\mathbb{R}} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1-z'm_{1-z'}}(X) dm_{1-z'}.$$

The second term on the right-hand side of (S39) equals to

$$\begin{aligned}
&\sum_{z'=1,0} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} \prod_{z=0,1} f_{zm_z}(X) \cdot s(M = m_{z'} | X, Z = z') du \right] \\
&= \sum_{z'=1,0} \mathbb{E} \left[\int_{\mathbb{R}} \int_{\mathbb{R}} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1-z'm_{1-z'}}(X) dm_{1-z'} \right. \\
&\quad \left. \cdot s(M = m_{z'} | X, Z = z') f(M = m_{z'} | X, Z = z') dm_{z'} \right] \\
&= \sum_{z'=1,0} \mathbb{E} \left\{ \int_{\mathbb{R}} \gamma_2'^{(z')}(X, M = m_{z'}) s(M = m_{z'} | X, Z = z') f(M = m_{z'} | X, Z = z') dm_{z'} \right\} \\
&= \sum_{z'=1,0} \mathbb{E} \left[\mathbb{E} \left\{ \gamma_2'^{(z')}(X, M) s(M | X, Z = z') | X, Z = z' \right\} \right] \\
&= \sum_{z'=1,0} \mathbb{E} \left[\mathbb{E} \left\{ \frac{\mathbf{1}(Z = z')}{\pi_{z'}(X)} \gamma_2'^{(z')}(X, M) s(M | X, Z) | X \right\} \right] \\
&= \mathbb{E} \left[\left\{ \sum_{z'=1,0} \frac{\mathbf{1}(Z = z')}{\pi_{z'}(X)} \gamma_2'^{(z')}(X, M) \right\} s(M | X, Z) \right] \quad (\text{LOTE}) \\
&= \mathbb{E} \left\{ \frac{1}{\pi_Z(X)} \gamma_2'^{(Z)}(X, M) s(M | X, Z) \right\} \\
&= \mathbb{E} \{ \xi'_3(X, Z, M) s(M | X, Z) \} \\
&= \mathbb{E} \{ \tilde{\xi}'_3(X, Z, M) s(V) \} \quad (\text{Lemma S2 with } A_1 = M, A_2 = (X, Z)), \tag{S41}
\end{aligned}$$

recalling $\xi'_3(X, Z, M) = \{\pi_Z(X)\}^{-1} \gamma_2'^{(Z)}(X, M)$ and $\tilde{\xi}'_3(X, Z, M) = \xi'_3(X, Z, M) - \mathbb{E}[\xi'_3(X, Z, M) |$

$X, Z]$.

For the third term on the right-hand side of (S39), since $s(X) + s(Z, M, Y | X) = s(V)$, we have

$$\begin{aligned}
& \mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \cdot s(X) \right\} \\
&= \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \cdot \{s(V) - s(Z, M, Y | X)\} \right] \\
&= \mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \cdot s(V) \right\} - \underbrace{\mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \cdot \mathbb{E} \{s(Z, M, Y | X) | X\} \right]}_{=0} \quad (\text{LOTE}) \\
&= \mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \cdot s(V) \right\} \\
&= \mathbb{E} \left[\left[\int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du - \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right] \right] \cdot s(V) \right], \tag{S42}
\end{aligned}$$

where the last equality is by $\mathbb{E} \left[\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} s(V) \right] = \mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \mathbb{E} \{s(V)\} = 0$.

Combining (S39), (S40), (S41) and (S42), we have

$$\begin{aligned}
& \frac{\partial}{\partial \theta} \mathbb{E}_\theta \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u^{(\theta)}(X) du \right\} \Big|_{\theta=0} \\
&= \mathbb{E} \left[\left[\tilde{\xi}'_2(X, Z, M) + \tilde{\xi}'_3(X, Z, M) + \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du - \mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \right] s(V) \right],
\end{aligned}$$

and therefore

$$\tilde{\xi}'_2(x, z, m) + \tilde{\xi}'_3(x, z, m) + \int_{\mathbb{R}^2} k_{u^*}(u) e_u(x) du - \mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \tag{S43}$$

is the EIF of $\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}$.

Step 2 (EIF of $\mathbb{E} \left[\int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} k_{u^*}(u) e_u(X) du \right]$): We obtain the following derivatives,

$$\begin{aligned}
& \frac{\partial}{\partial \theta} \mathbb{E}_\theta \left[\int_{\mathbb{R}^2} \{\mu_{1m_1}^{(\theta)}(X) - \mu_{0m_0}^{(\theta)}(X)\} k_{u^*}(u) e_u^{(\theta)}(X) du \right] \Big|_{\theta=0} \\
&= \mathbb{E} \left[\int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} k_{u^*}(u) e_u(X) du \cdot s(X) \right] \\
&+ \mathbb{E} \left[\int_{\mathbb{R}^2} \left\{ \frac{\partial}{\partial \theta} \mu_{1m_1}^{(\theta)}(X) \Big|_{\theta=0} - \frac{\partial}{\partial \theta} \mu_{0m_0}^{(\theta)}(X) \Big|_{\theta=0} \right\} k_{u^*}(u) e_u(X) du \right] \\
&+ \mathbb{E} \left[\int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} k_{u^*}(u) \cdot \frac{\partial}{\partial \theta} e_u^{(\theta)}(X) \Big|_{\theta=0} du \right]. \tag{S44}
\end{aligned}$$

Similar to (S42), the first term on the right-hand side of (S44) equals to

$$\begin{aligned}
& \mathbb{E} \left[\int_{\mathbb{R}^2} \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} k_{u^*}(u) e_u(X) du \cdot s(V) \right] \\
&= \mathbb{E} \left[\left\{ \int_{\mathbb{R}^2} \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} k_{u^*}(u) e_u(X) du \right. \right. \\
&\quad \left. \left. - \mathbb{E} \int_{\mathbb{R}^2} \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} k_{u^*}(u) e_u(X) du \right\} \cdot s(V) \right]. \tag{S45}
\end{aligned}$$

For the second term of (S44), we first derive

$$\begin{aligned}
\frac{\partial}{\partial \theta} \mu_{zm_z}^{(\theta)}(x) \Big|_{\theta=0} &= \frac{\partial}{\partial \theta} \mathbb{E}_\theta(Y | X = x, Z = z, M = m_z) \Big|_{\theta=0} \\
&= \mathbb{E}\{Y s(Y | X = x, Z = z, M = m_z) | X = x, Z = z, M = m_z\} \\
&= \mathbb{E}\{Y s(Y | X, Z, M) | X = x, Z = z, M = m_z\}.
\end{aligned}$$

Then we have

$$\begin{aligned}
& \mathbb{E} \left\{ \int_{\mathbb{R}^2} \frac{\partial}{\partial \theta} \mu_{zm_z}^{(\theta)}(X) \Big|_{\theta=0} k_{u^*}(u) e_u(X) du \right\} \\
&= \mathbb{E} \left[\int_{\mathbb{R}} \int_{\mathbb{R}} \mathbb{E}\{Y s(Y | X, Z, M) | X, Z = z, M = m_z\} \cdot k_{u^*}(u) e_u(X) dm_{1-z} dm_z \right] \\
&= \mathbb{E} \left[\int_{\mathbb{R}} \int_{\mathbb{R}} \underbrace{k_{u^*}(u) e_{(m_1, m_0)}(X) dm_{1-z}}_{=\gamma_1^{(z)}(X, M=m_z)} \cdot \mathbb{E}\{Y s(Y | X, Z, M) | X, Z = z, M = m_z\} dm_z \right] \\
&= \mathbb{E} \left[\int_{\mathbb{R}} \frac{\gamma_1^{(z)}(X, m_z)}{f(M = m_z | X, Z = z)} \cdot \mathbb{E}\{Y s(Y | X, Z, M) | X, Z = z, M = m_z\} \cdot f(M = m_z | X, Z = z) dm_z \right] \\
&= \mathbb{E} \left[\mathbb{E} \left[\frac{\gamma_1^{(z)}(X, M)}{f(M | X, Z = z)} \cdot \mathbb{E}\{Y s(Y | X, Z = z, M) | X, Z = z, M\} | X, Z = z \right] \right] \\
&= \mathbb{E} \left[\mathbb{E} \left[\mathbb{E} \left\{ \frac{\gamma_1^{(z)}(X, M)}{f(M | X, Z = z)} \cdot Y s(Y | X, Z = z, M) | X, Z = z, M \right\} | X, Z = z \right] \right] \\
&= \mathbb{E} \left[\mathbb{E} \left[\frac{\gamma_1^{(z)}(X, M)}{f(M | X, Z = z)} \cdot Y s(Y | X, Z = z, M) | X, Z = z \right] \right] \quad (\text{LOTE}) \\
&= \mathbb{E} \left[\mathbb{E} \left\{ \frac{1(Z = z)}{\pi_z(X)} \frac{\gamma_1^{(Z)}(X, M)}{f(M | X, Z = z)} \cdot Y s(Y | X, Z = z, M) | X \right\} \right] \quad (\text{LOTE}) \\
&= \mathbb{E} \left[\mathbb{E} \left\{ \frac{\mathbb{I}(Z = z) \gamma_1^{(Z)}(X, M)}{\pi_Z(X) f(M | X, Z)} \cdot Y s(Y | X, Z, M) | X \right\} \right] \\
&= \mathbb{E} \left\{ \frac{1(Z = z) \gamma_1^{(Z)}(X, M)}{\pi_Z(X) f_{ZM}(X)} \cdot Y s(Y | X, Z, M) \right\} \quad (\text{LOTE}). \tag{S46}
\end{aligned}$$

By (S46), the second term on the right-hand side of (S44) is

$$\begin{aligned} & \mathbb{E} \left[\left\{ \frac{1(Z=1)\gamma_1^{(Z)}(X, M)}{\pi_Z(X)f_{ZM}(X)} Y - \frac{1(Z=0)\gamma_1^{(Z)}(X, M)}{\pi_Z(X)f_{ZM}(X)} Y \right\} s(Y | X, Z, M) \right] \\ &= \mathbb{E} \left\{ \frac{(-1)^{Z+1}\gamma_1^{(Z)}(X, M)}{\pi_Z(X)f_{ZM}(X)} Y s(Y | X, Z, M) \right\}, \end{aligned}$$

recalling that $\xi_1(x, z, m) = \{\pi_z(x)f_{zm}(x)\}^{-1}(-1)^{z+1}\gamma_1^{(z)}(x, m)$. By Lemma S2 with $A_1 = Y$ and $A_2 = (X, Z, M)$, the second term on the right-hand side of (S44) then is,

$$\mathbb{E} \left[\xi_1(X, Z, M) \{Y - \mu_{ZM}(X)\} \cdot s(V) \right]. \quad (\text{S47})$$

where $\tilde{\xi}_1(X, Z, M, Y) = \xi_1(X, Z, M, Y) - \mathbb{E}\{\xi_1(X, Z, M, Y) | X, Z, M\}$.

For the third term on the right-hand side of (S44), similar arguments to (S40) and (S41) can directly show

$$\begin{aligned} & \mathbb{E} \left[\int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} k_{u^*}(u) \cdot \frac{\partial}{\partial \theta} e_u^{(\theta)}(X) \Big|_{\theta=0} du \right] \\ &= \mathbb{E} \left[\{\tilde{\xi}_2(X, Z, M) + \tilde{\xi}_3(X, Z, M)\} \cdot s(V) \right], \end{aligned}$$

where recalling Section S1, we denote

- (i) $\tilde{\xi}_a(X, Z, M) = \xi_a(X, Z, M) - \mathbb{E}\{\xi_a(X, Z, M) | X, Z\}$ for $a = 2, 3$;
- (ii) $\xi_2(X, Z, M) = \frac{1}{\pi_Z(X)} \int_{\mathbb{R}^2} \mathbb{I}(M \leq m_Z) \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} k_{u^*}(u) c_\rho^{(Z)} \{F_{1m_1}(X), F_{0m_0}(X)\} \{\prod_{z \in \{0,1\}} f_{zm_z}(X)\} du$;
- (iii) $\xi_3(X, Z, M) = \frac{1}{\pi_Z(X)} \gamma_2^{(Z)}(X, M)$;
- (iv) $\gamma_2^{(1)}(X, M) = \int_{\mathbb{R}} \{\mu_{1M}(X) - \mu_{0m_0}(X)\} k_{u^*}(M, m_0) c_\rho \{F_{1M}(X), F_{0m_0}(X)\} f_{0m_0}(X) dm_0$;
- (v) $\gamma_2^{(0)}(X, M) = \int_{\mathbb{R}} \{\mu_{1m_1}(X) - \mu_{0M}(X)\} k_{u^*}(m_1, M) c_\rho \{F_{1m_1}(X), F_{0M}(X)\} f_{1m_1}(X) dm_1$.

Combining (S44), (S45), (S47) we have

$$\begin{aligned} & \frac{\partial}{\partial \theta} \mathbb{E}_\theta \left[\int_{\mathbb{R}^2} \{\mu_{1m_1}^{(\theta)}(X) - \mu_{0m_0}^{(\theta)}(X)\} k_{u^*}(u) e_u^{(\theta)}(X) du \right] \Big|_{\theta=0} \\ &= \mathbb{E} \left[\left[\tilde{\xi}_2(X, Z, M) + \tilde{\xi}_3(X, Z, M) + \xi_1(X, Z, M) \{Y - \mu_{ZM}(X)\} \right. \right. \\ & \quad \left. \left. + \int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} k_{u^*}(u) e_u(X) du \right] \right] \end{aligned}$$

$$-\mathbb{E} \int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du \Big] \cdot s(V) \Big].$$

Therefore the EIF of $\mathbb{E} \left[\int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du \right]$ is

$$\begin{aligned} & \tilde{\xi}_2(x, z, m) + \tilde{\xi}_3(x, z, m) + \xi_1(x, z, m) \{y - \mu_{zm}(x)\} + \int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(x) - \mu_{0m_0}(x) \right\} k_{u^*}(u) e_u(x) du \\ & - \mathbb{E} \left\{ \int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du \right\}. \end{aligned} \quad (\text{S48})$$

Step 3 (EIF of τ_{u^*}): Recalling identification formula (8) and applying Lemma S3 with derived EIFs in Step 1 and Step 2, we have

$$\begin{aligned} \frac{\partial}{\partial \theta} \tau_{u^*, \theta} &= \frac{1}{\mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right]} \frac{\partial}{\partial \theta} \mathbb{E}_\theta \left[\int_{\mathbb{R}^2} \left\{ \mu_{1m_1}^{(\theta)}(X) - \mu_{0m_0}^{(\theta)}(X) \right\} k_{u^*}(u) e_u^{(\theta)}(X) du \right] \Big|_{\theta=0} \\ & - \frac{\mathbb{E} \left[\int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du \right]}{\left\{ \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right] \right\}^2} \frac{\partial}{\partial \theta} \mathbb{E}_\theta \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u^{(\theta)}(X) du \right\} \Big|_{\theta=0} \\ &= \frac{1}{\mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right]} \cdot \mathbb{E} \left[\left[\tilde{\xi}_2(X, Z, M) + \tilde{\xi}_3(X, Z, M) + \xi_1(X, Z, M) \{Y - \mu_{ZM}(X)\} \right. \right. \\ & \left. \left. + \int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du - \mathbb{E} \int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du \right] \cdot s(V) \right] \\ & - \frac{\mathbb{E} \left[\int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du \right]}{\left[\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \right]^2} \cdot \mathbb{E} \left[\left\{ \tilde{\xi}'_2(X, Z, M) + \tilde{\xi}'_3(X, Z, M) \right. \right. \\ & \left. \left. + \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du - \mathbb{E} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \cdot s(V) \right] \\ &= \mathbb{E} \left[\left[\frac{1}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} \cdot \left[\tilde{\xi}_2(X, Z, M) + \tilde{\xi}_3(X, Z, M) + \xi_1(X, Z, M) \{Y - \mu_{ZM}(X)\} \right. \right. \right. \\ & \left. \left. + \int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du - \mathbb{E} \int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du \right] \right. \right. \\ & \left. \left. - \frac{\mathbb{E} \left[\int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du \right]}{\left[\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \right]^2} \cdot \left\{ \tilde{\xi}'_1(X, Z, M) + \tilde{\xi}'_2(X, Z, M) \right. \right. \right. \\ & \left. \left. + \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du - \mathbb{E} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \right] \cdot s(V) \right], \\ &= \mathbb{E} \left[\left[\frac{1}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} \cdot \left[\tilde{\xi}_2(X, Z, M) + \tilde{\xi}_3(X, Z, M) + \xi_1(X, Z, M) \{Y - \mu_{ZM}(X)\} \right. \right. \right. \\ & \left. \left. + \int_{\mathbb{R}^2} \left\{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \right\} k_{u^*}(u) e_u(X) du \right] \right. \right. \\ & \left. \left. - \frac{\tau_{u^*}}{\left[\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \right]} \cdot \left\{ \tilde{\xi}'_2(X, Z, M) + \tilde{\xi}'_3(X, Z, M) + \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \right] \cdot s(V) \right], \\ &= \mathbb{E} \left[\phi_{\text{np}}(X, Z, M, Y) s(V) \right], \end{aligned} \quad (\text{S49})$$

after rearranging the terms and recalling the definitions in Section S1 and (7). (S49) combining with (S32) proves that $\phi_{\text{np}}(x, z, m, y) = \phi_{\text{kn}}(x, z, m, y) + \tilde{\phi}_{\text{np}}(x, z, m)$ is the full EIF of τ_{u^*} .

S5.3.3 Derivation of $\phi_{\text{p}}(x, z, m, y)$ under Scenario (ii)

Under the model assumption that \mathcal{M}_{ps} is parametrically modeled and estimated, we first derive the corresponding tangent space Λ and then project $\phi_{\text{np}}(x, z, m, y)$ onto Λ to obtain the new EIF $\phi_{\text{p}}(x, z, m, y)$ under this specific scenario (Van der Vaart, 2000, § 25.3).

Consider any regular submodel with likelihood $f_{\theta, \beta}(V = v)$, parametrized with some one-dimensional parameter $\theta \in \mathbb{R}$ and the principal score model parameter $\beta \in \mathbb{R}^{d_\beta}$; the true model takes place when $f_{\theta=0, \beta=\beta^*}(v) = f(v)$. The likelihood is decomposed as

$$f_{\theta, \beta}(v) = f_\theta(X = x)f_\theta(Z = z | X = x)f(M = m | X = x, Z = z, \beta) \cdot f_\theta(Y = y | X = x, Z = z, M = m).$$

Thus the score function of submodel is,

$$s_{\beta^*}(v) = \begin{bmatrix} s(x) + s(z | x) + s(y | x, z, m) \\ s_{zm}(x | \beta^*) \end{bmatrix} \in \mathbb{R}^{1+d_\beta},$$

where we recall $s(x) = \partial \log f_\theta(X = x) / \partial \theta |_{\theta=0}$, $s_{zm}(x | \beta^*) = \partial \log f(M = m | X = x, Z = z, \beta) / \partial \beta |_{\beta=\beta^*}$, and similarly for others. The parametric submodel tangent space can be represented as

$$\Lambda_\theta = \underbrace{\left\{ \ell_1 \{s(x) + s(z | x) + s(y | x, z, m)\} \mid \ell_1 \in \mathbb{R} \right\}}_{\Lambda_1} \oplus \underbrace{\left\{ \ell_2^\top s_{zm}(x | \beta^*) \mid \ell_2 \in \mathbb{R}^{d_\beta} \right\}}_{\Lambda_2}.$$

One has $\Lambda_1 \perp \Lambda_2$, as by LOTE,

$$\begin{aligned} & \mathbb{E} \left[\ell_1 \{s(X) + s(Z | X) + s(Y | X, Z, M)\} \cdot \ell_2^\top s_{ZM}(X | \beta^*) \right] \\ &= \mathbb{E} \left[\ell_1 \{s(X) + s(Z | X)\} \cdot \mathbb{E} \{ \ell_2^\top s_{ZM}(X | \beta^*) \mid X, Z \} \right] \\ & \quad + \mathbb{E} \left[\ell_2^\top s_{ZM}(X | \beta^*) \cdot \mathbb{E} \{ s(Y | X, Z, M) \mid X, Z, M \} \right] \\ &= 0, \end{aligned} \tag{S50}$$

for any $\ell_1 \in \mathbb{R}$ and $\ell_2 \in \mathbb{R}^{d_\beta}$. Then the semiparametric tangent space Λ is the mean-square closure of Λ_θ , over all possible regular submodels associated with parameter θ . Note β is associated with a

known and fixed model, namely $f_{zm}(x | \beta)$. Thus we have $\Lambda = \bar{\Lambda}_1 + \Lambda_2$, where $\bar{\Lambda}_1$ is the mean-square closure of Λ_1 over all regular parametric submodels associated with θ . Recalling (S55)–(S56), we similarly have $\bar{\Lambda}_1 = H_1 \oplus H_2 \oplus H_4$ and thus

$$\Lambda = H_1 \oplus H_2 \oplus H_4 \oplus \Lambda_2, \quad (\text{S51})$$

where

$$\begin{aligned} H_1 &= \{h(x) \mid \mathbb{E}\{h(X)\} = 0\}, \\ H_2 &= \{h(x, z) \mid \mathbb{E}\{h(X, Z) \mid X\} = 0\}, \\ H_4 &= \{h(x, z, m, y) \mid \mathbb{E}\{h(X, Z, M, Y) \mid X, Z, M\} = 0\}, \\ \Lambda_2 &= \left\{ \ell_2^\top s_{zm}(x \mid \beta^*) \mid \ell_2 \in \mathbb{R}^{d_\beta} \right\}. \end{aligned}$$

Similar to (S50), we can check that

$$H_1 \perp H_2 \perp H_4 \perp \Lambda_2. \quad (\text{S52})$$

To derive the semiparametric EIF, we need to obtain the projection of $\phi_{\text{np}}(x, z, m, y)$ onto the semiparametric tangent space Λ . We first decompose $\phi_{\text{np}}(x, z, m, y)$ in (S49) as follows,

$$\begin{aligned} \phi_{\text{np}}(x, z, m, y) &= \underbrace{\frac{\tilde{\xi}_2(x, z, m) + \tilde{\xi}_3(x, z, m) - \tau_{u^*} \{ \tilde{\xi}'_2(x, z, m) + \tilde{\xi}'_3(x, z, m) \}}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}}}_{\phi_1(x, z, m)} + \underbrace{\frac{\xi_1(x, z, m) \{ y - \mu_{zm}(x) \}}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}}}_{\phi_2(x, z, m, y)} \\ &\quad + \underbrace{\frac{\int_{\mathbb{R}^2} \{ \mu_{1m_1}(x) - \mu_{0m_0}(x) \} k_{u^*}(u) e_u(x) du - \tau_{u^*} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(x) du}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}}}_{\phi_3(x)}. \end{aligned} \quad (\text{S53})$$

Since the orthogonal projection is a linear operator, we only need to derive the projections of ϕ_1 – ϕ_3 onto Λ , respectively, and sum them up to get the final projections. By definition, one has

$$\begin{aligned} \mathbb{E}\{ \phi_2(X, Z, M, Y) \mid X, Z, M \} &= c_k^{-1} \left[\xi_1(X, Z, M) \{ \mathbb{E}(Y \mid X, Z, M) - \mu_{ZM}(X) \} \right] \\ &= 0, \end{aligned}$$

recalling ξ_1 is defined in Section S1; we denote $c_k = \mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}$ for simplicity. Therefore $\phi_2(x, z, m, y) \in H_4$, and by (S52), we have the projection of $\phi_2(x, z, m, y)$ onto Λ is itself. Similarly,

we can check that $\mathbb{E}\{\phi_3(X)\} = \tau_{u^*} - \tau_{u^*} = 0$ by Theorem 2. Thus $\phi_3(x) \in H_1$. Similarly, the projection of $\phi_3(x)$ onto Λ is also itself.

It is left to find the projection of $\phi_1(x, z, m)$. Recalling the definitions of $\tilde{\xi}_2, \tilde{\xi}_3, \tilde{\xi}'_2, \tilde{\xi}'_3$ in Section S1, one can easily show that

$$\mathbb{E}\{\phi_1(X, Z, M) \mid X, Z\} = 0.$$

We claim that the projections of $\phi_1(X, Z, M)$ onto H_1, H_2 , and H_4 are all zero, because $\phi_1(X, Z, M) \perp H_j, j = 1, 2, 4$. For example, denote $h_1^{(1)}(x) \in H_1$ as the projection of $\phi_1(x, z, m)$ onto Λ . Then $h_1^{(1)}(x)$ minimizes the following function

$$\begin{aligned} & \mathbb{E}\{h_1^{(1)}(x) - \phi_1(X, Z, M)\}^2 \\ = & \mathbb{E}\{h_1^{(1)}(X)\}^2 + \mathbb{E}\{\phi_1^2(X, Z, M)\} - 2\mathbb{E}[h_1^{(1)}(X) \underbrace{\mathbb{E}\{\phi_1(X, Z, M) \mid X, Z\}}_{=0}] \\ = & \mathbb{E}\{h_1^{(1)}(X)\}^2 + \mathbb{E}\{\phi_1^2(X, Z, M)\}, \end{aligned}$$

among all functions in H_1 , which is clearly satisfied if and only if $h_1^{(1)}(x) = 0$, a.s. Thus $\phi_1(x, z, m) \perp H_1$. Observing that

$$\begin{aligned} \mathbb{E}\{h_2(X, Z)\phi_1(X, Z, M)\} &= \mathbb{E}[h_2(X, Z)\mathbb{E}\{\phi_1(X, Z, M) \mid X, Z\}] = 0, \\ \mathbb{E}\{h_4(X, Z, M, Y)\phi_1(X, Z, M)\} &= \mathbb{E}[\phi_1(X, Z, M)\mathbb{E}\{h_4(X, Z, M, Y) \mid X, Z, M\}] = 0, \end{aligned}$$

for any $h_2(x, z) \in H_2$ and $h_4(x, z, m, y) \in H_4$, we thus similarly have $\phi_1(x, z, m) \perp H_2$ and $\phi_1(x, z, m) \perp H_4$. So the projection of $\phi_1(x, z, m)$ onto Λ is only the projection of $\phi_1(x, z, m)$ onto Λ_2 .

Now we obtain the projections of $\tilde{\xi}_2, \tilde{\xi}'_2, \tilde{\xi}'_3$ onto Λ_2 one by one. For example, for $\tilde{\xi}_2$, this is equivalent to finding $\tilde{\ell}_1^\top s_{zm}(x \mid \beta^*) \in \Lambda_2$ solving the following optimization problem,

$$\begin{aligned} & \arg \min_{\tilde{\ell}_1 \in \mathbb{R}^{d_\beta}} \mathbb{E}\left\{\tilde{\xi}_2(X, Z, M) - \tilde{\ell}_1^\top s_{ZM}(X \mid \beta^*)\right\}^2 \\ = & \arg \min_{\tilde{\ell}_1 \in \mathbb{R}^{d_\beta}} \tilde{\ell}_1^\top I_{\beta^*} \tilde{\ell}_1 - 2\tilde{\ell}_1^\top \mathbb{E}\{\tilde{\xi}_2(X, Z, M) s_{ZM}(X \mid \beta^*)\} + \mathbb{E}\{\tilde{\xi}_2^2(X, Z, M)\}, \end{aligned}$$

which can be solved by standard quadratic programming (Boyd et al., 2004). In particular, the solution is

$$\tilde{\ell}_{1, \diamond} = I_{\beta^*}^{-1} \mathbb{E}\{\tilde{\xi}_2(X, Z, M) s_{ZM}(X \mid \beta^*)\}$$

$$\begin{aligned}
&= I_{\beta^*}^{-1} \mathbb{E} \{ \xi_2(X, Z, M) s_{ZM}(X | \beta^*) \} - I_{\beta^*}^{-1} \mathbb{E} \left[\mathbb{E}(\xi_2 | X, Z) \underbrace{\mathbb{E}\{s_{ZM}(X | \beta^*) | X, Z\}}_{=0} \right] \\
&= I_{\beta^*}^{-1} \mathbb{E} \{ \xi_2(X, Z, M) s_{ZM}(X | \beta^*) \} \\
&= I_{\beta^*}^{-1} \mathbb{E} \left[\frac{1}{\pi_Z(X)} \int_{\mathbb{R}^2} 1(M \leq m_z) \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} k_{u^*}(u) \cdot c_\rho^{(Z)} \{ F_{1m_1}(X), F_{0m_0}(X) \} \left\{ \prod_{z \in \{0,1\}} f_{zm_z}(X) \right\} du \cdot s_{ZM}(X | \beta^*) \right] \\
&= I_{\beta^*}^{-1} \sum_{z=0,1} \mathbb{E} \left[\mathbb{E} \left[\int_{\mathbb{R}^2} 1(M \leq m_z) \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} k_{u^*}(u) c_\rho^{(z)} \{ F_{1m_1}(X), F_{0m_0}(X) \} \right. \right. \\
&\quad \left. \left. \cdot \left\{ \prod_{z \in \{0,1\}} f_{zm_z}(X) \right\} du \cdot s_{ZM}(X | \beta^*) \mid X, Z = z \right] \right] \quad (\text{LOTE}) \\
&= I_{\beta^*}^{-1} \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} \int_{-\infty}^{m_z} \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} k_{u^*}(u) c_\rho^{(z)} \{ F_{1m_1}(X), F_{0m_0}(X) \} \right. \\
&\quad \left. \cdot \left\{ \prod_{z \in \{0,1\}} f_{zm_z}(X) \right\} du \cdot \underbrace{f'_{zm}(X) f_{zm}^{-1}(X) f_{zm}(X)}_{=1} dm du \right] \\
&= I_{\beta^*}^{-1} \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} c_\rho^{(z)} \{ F_{1m_1}(X), F_{0m_0}(X) \} f_{1m_1}(X) f_{0m_0}(X) F'_{zm_z}(X) du \right].
\end{aligned}$$

Thus $\tilde{\ell}_{1,\diamond}^\top s_{zm}(x | \beta^*)$ is the projection of $\tilde{\xi}_2$ onto Λ_1 , which, combing with the previous discussions, is also the projection of $\tilde{\xi}_2$ onto the whole Λ . With similar techniques, we have $\tilde{\ell}_{2,\diamond}^\top s_{zm}(x | \beta^*)$ is the projection of $\tilde{\xi}_3$ onto Λ , such that

$$\begin{aligned}
\tilde{\ell}_{2,\diamond} &= I_{\beta^*}^{-1} \mathbb{E} \{ \xi_3(X, Z, M) s_{ZM}(X | \beta^*) \} \\
&= \sum_{z=0,1} I_{\beta^*}^{-1} \mathbb{E} \left[\mathbb{E} \{ \gamma_2^{(z)}(X, M) s_{ZM}(X | \beta^*) \mid X, Z = z \} \right].
\end{aligned}$$

When $z = 1$, for the term on the right-hand side of above equation, we further have

$$\begin{aligned}
&I_{\beta^*}^{-1} \mathbb{E} \left[\mathbb{E} \{ \gamma_2^{(1)}(X, M) s_{ZM}(X | \beta^*) \mid X, Z = 1 \} \right] \\
&= I_{\beta^*}^{-1} \mathbb{E} \left[\mathbb{E} \left\{ \int_{\mathbb{R}} \{ \mu_{1M}(X) - \mu_{0m_0}(X) \} k_{u^*}(M, m_0) c_\rho \{ F_{1M}(X), F_{0m_0}(X) \} f_{0m_0}(X) dm_0 \right. \right. \\
&\quad \left. \left. \cdot f^{-1}(M | X, Z = 1) f'_{1M}(X) \mid X, Z = 1 \right\} \right] \\
&= I_{\beta^*}^{-1} \mathbb{E} \left[\int_{\mathbb{R}} \int_{\mathbb{R}} \{ \mu_{1m}(X) - \mu_{0m_0}(X) \} k_{u^*}(m, m_0) c_\rho \{ F_{1m}(X), F_{0m_0}(X) \} f_{0m_0}(X) f'_{1m}(X) dm_0 dm \right] \\
&= I_{\beta^*}^{-1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} c_\rho \{ F_{1m_1}(X), F_{0m_0}(X) \} f_{0m_0}(X) f'_{1m_1}(X) du \right]
\end{aligned}$$

We can similarly deal with the $z = 0$ case and deduce,

$$\tilde{\ell}_{2,\diamond} = I_{\beta^*}^{-1} \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} c_\rho \{ F_{1m_1}(X), F_{0m_0}(X) \} f_{(1-z)m_{1-z}}(X) f'_{zm_z}(X) du \right].$$

Notice that $\tilde{\xi}'_2, \tilde{\xi}'_3$ have very similar structures as $\tilde{\xi}_2, \tilde{\xi}_3$. Following the derivation strategy as above, we can show that $\tilde{\ell}'_{1,\diamond\top} s_{zm}(x | \beta^*), \tilde{\ell}'_{2,\diamond\top} s_{zm}(x | \beta^*)$ are the projections of $\tilde{\xi}'_2, \tilde{\xi}'_3$, respectively, such that

$$\begin{aligned}\tilde{\ell}'_{1,\diamond} &= I_{\beta^*}^{-1} \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_{\rho}^{(z)} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1m_1}(X) f_{0m_0}(X) F'_{zm_z}(X) du \right], \\ \tilde{\ell}'_{2,\diamond} &= I_{\beta^*}^{-1} \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_{\rho} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X) f'_{zm_z}(X) du \right].\end{aligned}$$

Recalling that $\phi_1(x, z, m)$ is a linear combination of $\tilde{\xi}_2, \tilde{\xi}_3, \tilde{\xi}'_2, \tilde{\xi}'_3$, we have the projection of $\phi_1(x, z, m)$ on Λ is the corresponding linear combination of $\tilde{\xi}_2, \tilde{\xi}_3, \tilde{\xi}'_2, \tilde{\xi}'_3$'s projections on Λ . In particular, the projection of $\phi_1(x, z, m)$ is

$$\frac{\mathcal{A}_2^* - \tau_{u^*} \mathcal{A}_1^*}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} I_{\beta^*}^{-1} s_{zm}(x | \beta^*), \quad (\text{S54})$$

where we recall that in Section S1,

$$\begin{aligned}\mathcal{A}_1^* &= \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_{\rho} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X) \{f'_{zm_z}(X)\}^{\top} du \right] \\ &\quad + \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[c_{\rho}^{(z)} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1m_1}(X) f_{0m_0}(X) \{F'_{zm_z}(X)\}^{\top} \right] du \right], \\ \mathcal{A}_2^* &= \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} c_{\rho} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X) \{f'_{zm_z}(X)\}^{\top} du \right] \\ &\quad + \sum_{z=0,1} \mathbb{E} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{ \mu_{1m_1}(X) - \mu_{0m_0}(X) \} c_{\rho}^{(z)} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1m_1}(X) f_{0m_0}(X) \{F'_{zm_z}(X)\}^{\top} du \right].\end{aligned}$$

Summarizing the projections of $\phi_1(\cdot) - \phi_3(\cdot)$ on Λ defined in (S51) as above, and recalling (S53), we conclude the projection of $\phi_{\text{np}}(x, z, m, y)$ on Λ , i.e., the semiparametric EIF of τ_{u^*} when the parametric model of \mathcal{M}_{ps} is known, is

$$\begin{aligned}\phi_{\text{p}}(x, z, m, y) &= \frac{\mathcal{A}_2^* - \tau_{u^*} \mathcal{A}_1^*}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} I_{\beta^*}^{-1} s_{zm}(x | \beta^*) + \frac{\xi_1(x, z, m) \{y - \mu_{zm}(x)\}}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} \\ &\quad + \frac{\int_{\mathbb{R}^2} \{ \mu_{1m_1}(x) - \mu_{0m_0}(x) \} k_{u^*}(u) e_u(x) du - \tau_{u^*} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(x) du}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}},\end{aligned}$$

which shows $\phi_{\text{p}}(x, z, m, y) = \phi_{\text{kn}}(x, z, m, y) + \tilde{\phi}_{\text{p}}(x, z, m)$ by recalling (7) and (S3).

S5.3.4 Derivation of $\phi_{\text{kn}}(x, z, m, y)$ under Scenario (i)

Now we assume the distribution of M given X, Z is known and thus we have some model restriction. Since the model assumption becomes more restrictive, the $\phi(\cdot)$ satisfying (S32) is not necessary to

be an EIF but must be an influence function (IF) of τ_{u^*} and the IF can be not unique (Bickel et al., 1993). Therefore, we can still derive an IF of τ_{u^*} in a similar yet simpler way as in Section S5.3.2. In particular, we find that $\phi(\cdot) = \phi_{\text{kn}}(\cdot)$ in (10) satisfying (S32) and thus $\phi_{\text{kn}}(\cdot)$ is an IF. Furthermore, based on the fundamental semiparametric statistics theory, we can verify that $\phi_{\text{kn}}(\cdot)$ is actually the EIF by showing that it belongs to the semiparametric tangent space and thereby finish our proof; See e.g. Van der Vaart (2000, § 25.3) for the relevant semiparametric statistics theory used here.

We show that $\phi_{\text{kn}}(\cdot)$ is an IF following the proof routine in Section S5.3.2 with some simplification. In particular, since $f_{zm}(x)$ and $F_{zm}(x)$ are all known, we do not need to parametrize $e_u(x)$ as $e_u^{(\theta)}(x)$ and take it into account when deriving the EIFs of

$$\mathbb{E}\left[\int_{\mathbb{R}^2}\{\mu_{1m_1}(X)-\mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right]$$

and $\mathbb{E}\left[\int_{\mathbb{R}^2}\{\mu_{1m_1}(X)-\mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right]$ in Section S5.3.2. Then we can show that the function

$$\int_{\mathbb{R}^2}k_{u^*}(u)e_u(x)du-\mathbb{E}\left\{\int_{\mathbb{R}^2}k_{u^*}(u)e_u(X)du\right\},$$

which drops the terms $\tilde{\xi}'_2, \tilde{\xi}'_3$ in (S43), is the IF of $\mathbb{E}\left[\int_{\mathbb{R}^2}\{\mu_{1m_1}(X)-\mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right]$ under Scenario (i). Similarly, the function

$$\tilde{\xi}_1(x, z, m, y)+\int_{\mathbb{R}^2}\{\mu_{1m_1}(x)-\mu_{0m_0}(x)\}k_{u^*}(u)e_u(x)du-\mathbb{E}\left\{\int_{\mathbb{R}^2}\{\mu_{1m_1}(X)-\mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right\},$$

which drops the terms $\tilde{\xi}_2, \tilde{\xi}_3$ in (S48), is the IF of

$$\mathbb{E}\left[\int_{\mathbb{R}^2}\{\mu_{1m_1}(X)-\mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right].$$

With simplified IFs of $\mathbb{E}\left[\int_{\mathbb{R}^2}\{\mu_{1m_1}(X)-\mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right]$ and $\mathbb{E}\left[\int_{\mathbb{R}^2}\{\mu_{1m_1}(X)-\mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right]$ as above and similar to (S49), we can use Lemma S3 to show that $\phi(\cdot) = \phi_{\text{kn}}(\cdot)$ in (10) satisfying (S32) and thus $\phi_{\text{kn}}(\cdot)$ is an IF of τ_{u^*} .

We now derive the semiparametric tangent space under Scenario (i). With known \mathcal{M}_{ps} , the density function decomposition similar to (S30) is now

$$\begin{aligned} & f_{\theta}(V=v) \\ &= f_{\theta}(X=x)f_{\theta}(Z=z|X=x)f(M=m|X=x,Z=z)f_{\theta}(Y=y|X=x,Z=z,M=m). \end{aligned}$$

Thus score function of some arbitrary parametric submodel is now decomposed as

$$s_\theta(v) = s_\theta(x) + s_\theta(z | x) + s_\theta(y | x, z, m). \quad (\text{S55})$$

By considering the mean-square closure of all functions $s_\theta(v)$ over all possible submodels, the tangent space is now degenerated to,

$$\Lambda = H_1 \oplus H_2 \oplus H_4, \quad (\text{S56})$$

where \oplus is the set sum and,

$$\begin{aligned} H_1 &= \{h(x) \mid \mathbb{E}\{h(X)\} = 0\}, \\ H_2 &= \{h(x, z) \mid \mathbb{E}\{h(X, Z) \mid X\} = 0\}, \\ H_4 &= \{h(x, z, m, y) \mid \mathbb{E}\{h(X, Z, M, Y) \mid X, Z, M\} = 0\}. \end{aligned}$$

First, observing that $\mathbb{E}[\xi_1(X, Z, M)\{Y - \mu_{ZM}(X)\}] = \mathbb{E}[\xi_1(X, Z, M)\mathbb{E}\{Y - \mu_{ZM}(X) \mid X, Z, M\}] = 0$, we have

$$\frac{\xi_1(x, z, m)\{y - \mu_{zm}(x)\}}{\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}} \in H_4.$$

Second, observe that

$$\begin{aligned} & \mathbb{E}\left[\frac{\int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du - \tau_{u^*}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}}{\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}}\right] \\ &= \frac{\mathbb{E}\left[\int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right]}{\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}} - \tau_{u^*} \\ &= 0, \end{aligned} \quad (\text{S57})$$

by Theorem 2. Recalling the form of $\phi_{\text{kn}}(\cdot)$ in (10), thus for $\psi_{\text{kn}}(x)$ defined as follows, we have

$$\begin{aligned} \psi_{\text{kn}}(x) &= \frac{\int_{\mathbb{R}^2} \{\mu_{1m_1}(x) - \mu_{0m_0}(x)\}k_{u^*}(u)e_u(x)du - \tau_{u^*}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(x)du\right\}}{\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}} \\ &= \phi_{\text{kn}}(x, z, m, y) - \frac{\xi_1(x, z, m)\{y - \mu_{zm}(x)\}}{\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}} \in H_1. \end{aligned}$$

Summarizing above results, we conclude that IF $\phi_{\text{kn}}(x, z, m, y)$ belongs to the semiparametric tan-

gent space, as

$$\begin{aligned}\phi_{\text{kn}}(x, z, m, y) &= \psi_{\text{kn}}(x) + \frac{\xi_1(x, z, m)\{y - \mu_{zm}(x)\}}{\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}} \\ &\in H_1 \oplus H_4 \subseteq \Lambda.\end{aligned}$$

This conclusion directly confirms that $\phi_{\text{kn}}(\cdot)$ is not only the IF, but also the EIF of τ_{u^*} under the semiparametric scenario with known \mathcal{M}_{ps} by the classic semiparametric statistics theory; See the discussions after Lemma 25.14 in [Van der Vaart \(2000, § 25.3\)](#).

S5.4 Proof of Theorem 4

With fixed u^* , we omit the footnote u^* in $\bar{\xi}_{1,u^*}(\cdot)$ and directly write $\bar{\xi}_1(\cdot)$ for simplicity. By definition, we further write,

$$\begin{aligned}& \bar{\xi}_1(X, Z, M)\{Y - \bar{\mu}_{ZM}(X)\} \\ = & \underbrace{\frac{\mathbb{I}(Z=1)\gamma_1^{(Z)}(X, M)}{\bar{\pi}_Z(X)f(M|X, Z)}Y}_{\bar{\xi}_1^{(1)}(X, Z, M, Y)} - \underbrace{\frac{\mathbb{I}(Z=0)\gamma_1^{(Z)}(X, M)}{\bar{\pi}_Z(X)f(M|X, Z)}Y}_{\bar{\xi}_1^{(0)}(X, Z, M, Y)} \\ & - \underbrace{\frac{\mathbb{I}(Z=1)\gamma_1^{(Z)}(X, M)}{\bar{\pi}_Z(X)f(M|X, Z)}\bar{\mu}_{ZM}(X)}_{\bar{E}_{\xi_1}^{(1)}(X, Z, M)} + \underbrace{\frac{\mathbb{I}(Z=0)\gamma_1^{(Z)}(X, M)}{\bar{\pi}_Z(X)f(M|X, Z)}\bar{\mu}_{ZM}(X)}_{\bar{E}_{\xi_1}^{(0)}(X, Z, M)}\end{aligned}$$

We first consider the scenario when $\pi = \bar{\pi}$. Applying LOTE, one has

$$\begin{aligned}\mathbb{E}\left\{\bar{\xi}_1^{(1)}(X, Z, M, Y)\right\} &= \mathbb{E}\left\{\frac{\mathbb{I}(Z=1)\gamma_1^{(Z)}(X, M)}{\pi_Z(X)f(M|X, Z)}Y\right\} \\ &= \mathbb{E}\left[\mathbb{E}\left\{\frac{\mathbb{I}(Z=1)\gamma_1^{(Z)}(X, M)}{\pi_Z(X)f(M|X, Z)}Y \mid X\right\}\right] \\ &= \mathbb{E}\left[\pi_1(X)\mathbb{E}\left\{\frac{\gamma_1^{(Z)}(X, M)}{\pi_1(X)f(M|X, Z)}Y \mid X, Z=1\right\} + \pi_0(X) \cdot 0\right] \\ &= \mathbb{E}\left[\mathbb{E}\left\{\frac{\gamma_1^{(1)}(X, M)}{f(M|X, Z=1)}Y \mid X, Z=1\right\}\right] \\ &= \mathbb{E}\left[\mathbb{E}\left\{\frac{\gamma_1^{(1)}(X, M)}{f(M|X, Z=1)}\mathbb{E}(Y \mid X, Z=1, M) \mid X, Z=1\right\}\right] \\ &= \mathbb{E}\left\{\int_{\mathbb{R}} \frac{\gamma_1^{(1)}(M=m_1, X)\mathbb{E}(Y \mid X, Z=1, M=m_1)}{f(M=m_1|X, Z=1)}f(M=m_1|X, Z=1)dm_1\right\}\end{aligned}$$

$$\begin{aligned}
&= \mathbb{E} \left\{ \int_{\mathbb{R}} \gamma_1^{(1)}(M = m_1, X) \mu_{1m_1}(X) dm_1 \right\} \\
&= \mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(m_1, m_0) e_{(m_1, m_0)}(X) \mu_{1m_1}(X) du \right\}.
\end{aligned} \tag{S58}$$

Similarly, one has $\mathbb{E}\{\bar{\xi}_1^{(0)}(X, Z, M, Y)\} = \mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(m_1, m_0) e_{(m_1, m_0)}(X) \mu_{0m_0}(X) du\}$. With same techniques as above, we can show

$$\begin{aligned}
\mathbb{E}\{\bar{E}_{\xi_1}^{(1)}(X, Z, M)\} &= \mathbb{E} \left\{ \frac{\mathbb{I}(Z = 1) \gamma_1^{(Z)}(X, M)}{\pi_Z(X) f(M | X, Z)} \bar{\mu}_{ZM}(X) \right\} \\
&= \mathbb{E} \left[\mathbb{E} \left\{ \frac{\gamma_1^{(1)}(X, M)}{f(M | X, Z = 1)} \bar{\mu}_{1M}(X) \mid X, Z = 1 \right\} \right] \\
&= \mathbb{E} \left\{ \int_{\mathbb{R}} \gamma_1^{(1)}(M = m_1, X) \bar{\mu}_{1m_1}(X) dm_1 \right\} \\
&= \mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) \bar{\mu}_{1m_1}(X) du \right\},
\end{aligned}$$

and, similarly $\mathbb{E}\{\bar{E}_{\xi_1}^{(0)}(X, Z, M)\} = \mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) \bar{\mu}_{0m_0}(X) du\}$. Summarizing all results above,

$$\begin{aligned}
\bar{\tau}_{u^*} &= \frac{1}{\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du\}} \left[\mathbb{E} \left[\int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} k_{u^*}(u) e_u(X) du \right] \right. \\
&\quad - \mathbb{E} \left[\int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} k_{u^*}(u) e_u(X) du \right] \\
&\quad \left. + \mathbb{E} \left[\int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} k_{u^*}(u) e_u(X) du \right] \right] \\
&= \tau_{u^*},
\end{aligned} \tag{S59}$$

where the last equality is from Theorem 8 and we recall (7).

Next we consider the scenario when $\bar{\mu} = \mu$. With same technique as (S58), we can show

$$\mathbb{E}\{\bar{\xi}_1^{(z)}(X, Z, M, Y)\} = \mathbb{E} \left\{ \frac{\pi_z(X)}{\bar{\pi}_z(X)} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) \mu_{zm_z}(X) du \right\}$$

and

$$\mathbb{E}\{\bar{E}_{\xi_1}^{(z)}(X, Z, M)\} = \mathbb{E} \left\{ \frac{\pi_z(X)}{\bar{\pi}_z(X)} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) \bar{\mu}_{zm_z}(X) du \right\}$$

$$= \mathbb{E} \left\{ \frac{\pi_z(X)}{\bar{\pi}_z(X)} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) \mu_{zm_z}(X) du \right\},$$

for both $z = 0, 1$. This implies

$$\mathbb{E}\{\tilde{\xi}_1(X, Z, M, Y)\} = 0 \quad (\text{S60})$$

Finally since $\bar{\mu} = \mu$, by Theorem 2 we have

$$\bar{\tau}_{u^*} = \frac{0 + \mathbb{E} \left[\int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} k_{u^*}(u) e_u(X) du \right]}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} = \tau_{u^*}.$$

Combining with Proposition 1, we further have $\lim_{h \rightarrow 0} \bar{\tau}_{u^*} = \lim_{h \rightarrow 0} \tau_{u^*} = \tau_{u^*}^*$ \square

S5.5 Regularity assumptions for Theorem 5

We first discuss Assumption 4 in the main paper. Assumption 4 (i) allows either one of the models \mathcal{M}_{tp} or \mathcal{M}_{om} , to be incorrectly estimated, while consistency and limiting distribution results in Theorem 5 will still be valid. Assumption 4 (ii)–(iii) are standard which assume that when the sample size grows, nuisance function estimators are consistent to their limits under \mathcal{L}_∞ norm and specify their \mathcal{L}_2 convergence rates to the truths. Assumption 4 (iv) restricts the functional space complexities of estimators $\hat{\pi}$, $\hat{\mu}$ and their limits, which is a usual minimal condition for the use of empirical process techniques in theoretical analysis; See e.g., Kennedy (2016), Westling et al. (2020).

Now we list other regularity conditions used for Theorem 5 as follows.

Assumption S8 *There exists some $\epsilon > 0$ such that $\pi_z(X), \bar{\pi}_z(X) \in (\epsilon, 1/\epsilon)$, uniformly for all $X \in \mathbb{R}$ and $z = 0, 1$.*

Assumption S9 *For some universal constant $C > 0$, we have*

$$\max \left\{ \sup_{h>0} \int_{\mathbb{R}^2} k_{u^*}(u) du, \sup_{z \in \{0,1\}, h>0, m_z \in \mathbb{R}} h \int_{\mathbb{R}} k_{u^*}(u) dm_{1-z}, \sup_{h>0} h^2 \int_{\mathbb{R}^2} k_{u^*}^2(u) du \right\} \leq C.$$

Assumption S10 *(1) $\mathbb{E}\{e_U(X)\} < +\infty$; (2) $c(u, v)$ and all its first and second-order partial derivatives are bounded away from $+\infty$, uniformly over $(u, v) \in [-\infty, \infty] \times [-\infty, \infty]$; (3) $c^{(10)}(u, v) = c^{(01)}(u, v)$.*

Assumption S11 *$\sup_{u \in \mathbb{R}^2} e_u = \sup_{(m_1, m_0) \in \mathbb{R}^2} f(M_1 = m_1, M_0 = m_0) < C_M$ for a universal constant $C_M > 0$.*

Assumption S12 \mathbb{Y} is compact and thus $\sup_{\substack{x,m \in \mathbb{R} \\ z \in \{0,1\}}} \text{Var}(Y | X = x, Z = z, M = m) < +\infty$.

Assumption S13 For any $g_{zm_z}(\cdot | \beta) \in \{f_{zm_z}(\cdot | \beta), F_{zm_z}(\cdot | \beta)\}$ and any $\tilde{\beta} \in \mathbb{R}^{d_\beta}$, denote $g'_{zm_z}(\cdot | \tilde{\beta}) = \frac{\partial}{\partial \tilde{\beta}} g_{zm_z}(\cdot | \beta) \Big|_{\beta=\tilde{\beta}}$ and $g''_{zm_z}(\cdot | \tilde{\beta}) = \frac{\partial^2}{\partial \tilde{\beta} \partial \tilde{\beta}^\top} g_{zm_z}(\cdot | \beta) \Big|_{\beta=\tilde{\beta}}$. There exist constants $c_0, C_0, \delta_0 > 0$ such that, $\inf_{\substack{m_z, x \in \mathbb{R} \\ z \in \{0,1\}}} f_{zm_z}(x | \beta^*) \geq c_0$ and

$$\max \left\{ \sup_{\substack{m_z, x \in \mathbb{R} \\ z \in \{0,1\}}} |g_{zm_z}(x | \beta^*)|, \sup_{\substack{m_z, x \in \mathbb{R} \\ z \in \{0,1\}}} \|g'_{zm_z}(x | \beta^*)\|, \sup_{\substack{m_z, x \in \mathbb{R} \\ z \in \{0,1\} \\ \|\tilde{\beta} - \beta^*\| \leq \delta_0}} \|g''_{zm_z}(x | \tilde{\beta})\|_2 \right\} \leq C_0.$$

Assumption S8 is the standard overlap condition (Rosenbaum and Rubin, 1983). Assumptions S9–S10 can be satisfied by most of the commonly-used kernel functions and copula functions (Jaworski et al., 2010, Fan, 1993, Wasserman, 2006). Assumptions S11–S12 are standard density and moment conditions for the data distribution (Chen and Christensen, 2015, Luedtke and Van Der Laan, 2016, Chu et al., 2023). The condition of a compact \mathbb{Y} in Assumption S12 is only for controlling the empirical process term (see Section S6), and can be further relaxed if one exploits the standard sample splitting technique. Assumption S13 is a standard regularity condition to bound the first and second-order derivatives of some parametrized likelihood (Van der Vaart, 2000, Fan and Peng, 2004).

S5.6 Proof of Theorem 5

Although h is fixed for Loc.PCE, to further enable our nonparametric theoretical analysis of Cont.PCE in Section S5.7, we consider a more general setting in the proof such that we allow $h \rightarrow 0$. Our results in Theorem 5 can be easily obtained by setting h as fixed in our general results. For the ease of exposition, in the following, we will omit the footnote u^* in $\hat{\xi}_{1,u^*}(\cdot)$, $\bar{\xi}_{1,u^*}(\cdot)$, etc, and directly write $\hat{\xi}_1(\cdot)$, $\bar{\xi}_1(\cdot)$, etc. We will also occasionally omit random variables in $\hat{\xi}_1(X, Z, M)$ and directly write $\hat{\xi}_1$ for simplicity. On the other hand, for better illustration, we expand the notation of $\llbracket \cdot \rrbracket$ through its definition in (7) during the proof, and only use $\llbracket \cdot \rrbracket$ when wrapping up results.

We first define

$$\begin{aligned} \Delta_{1,n} &= (\mathbb{P}_n - \mathbb{P}) \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} + \mathbb{P}_n \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) \hat{e}_u(X) du - \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}, \\ \Delta_{2,n} &= \mathbb{P}_n \left[\hat{\xi}_1(X, Z, M) \{Y - \hat{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\} k_{u^*}(u) \hat{e}_u(X) du \right] \\ &\quad - \mathbb{P}_n \left[\bar{\xi}_1(X, Z, M) \{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} k_{u^*}(u) e_u(X) du \right]. \end{aligned}$$

Then by Taylor expansion, we have for some $\delta_n \in [0, 1]$,

$$\begin{aligned}
\hat{\tau}_{u^*} - \tau_{u^*} &= -\tau_{u^*} + \frac{\mathbb{P}_n[\hat{\xi}_1\{Y - \hat{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\}k_{u^*}(u)\hat{e}_u(X)du]}{\mathbb{P}_n\left\{\int_{\mathbb{R}^2} k_{u^*}(u)\hat{e}_u(X)du\right\}} \\
&= -\tau_{u^*} + \frac{\mathbb{P}_n[\bar{\xi}_1\{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du] + \Delta_{2,n}}{\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\} + \Delta_{1,n}} \\
&= -\tau_{u^*} + \frac{\mathbb{P}_n[\bar{\xi}_1\{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du] + \Delta_{2,n}}{\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}} + \Delta_{2,n} \\
&\quad - \frac{\mathbb{P}_n[\bar{\xi}_1\{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du] + \Delta_{2,n}}{\left[\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}\right]^2} \Delta_{1,n} \\
&\quad + \frac{\mathbb{P}_n[\bar{\xi}_1\{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du] + \Delta_{2,n}}{\left[\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\} + \delta_n\Delta_{1,n}\right]^3} \Delta_{1,n}^2 \\
&= \frac{(\mathbb{P}_n - \mathbb{P})[\bar{\xi}_1\{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du] + \Delta_{2,n}}{\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}} \\
&\quad - \frac{\mathbb{P}[\bar{\xi}_1\{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du] \Delta_{1,n}}{\left[\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}\right]^2} \\
&\quad - \frac{(\mathbb{P}_n - \mathbb{P})[\bar{\xi}_1\{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du] + \Delta_{2,n}}{\left[\mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}\right]^2} \Delta_{1,n} \\
&\quad + \frac{\mathbb{P}_n[\bar{\xi}_1\{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du] + \Delta_{2,n}}{\left[\mathbb{P}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\} + \delta_n\Delta_{1,n}\right]^3} \Delta_{1,n}^2 \quad (\text{S61})
\end{aligned}$$

where the last equality is by the double robustness introduced in Theorem 4. Next, in Sections S5.6.1 and S5.6.2, we bound the relative error terms above, respectively.

S5.6.1 Bounding $\Delta_{1,n}$

For simplicity, we denote

$$\Delta_{1,n}^{(1)} = (\mathbb{P}_n - \mathbb{P})\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\} \quad \text{and} \quad \Delta_{1,n}^{(2)} = \mathbb{P}_n\left\{\int_{\mathbb{R}^2} k_{u^*}(u)\hat{e}_u(X)du - \int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}.$$

Thus we have $\Delta_{1,n} = \Delta_{1,n}^{(1)} + \Delta_{1,n}^{(2)}$.

Notice that,

$$\begin{aligned}
\mathbb{P}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\}^2 &\leq \mathbb{P}\left\{\int_{\mathbb{R}^2} k_{u^*}^2(u)du \cdot \int_{\mathbb{R}^2} e_u^2(X)du\right\} \quad (\text{Cauchy-Schwartz inequality}) \\
&\lesssim h^{-2} \cdot \mathbb{P}\left\{\int_{\mathbb{R}^2} e_u^2(X)du\right\} \quad (\text{Assumption S9}) \\
&= h^{-2} \cdot \int_{\mathbb{R}^3} f^2(U = u \mid X = x) \cdot f(X = x)dxdu
\end{aligned} \quad (\text{S62})$$

$$\begin{aligned}
&= h^{-2} \cdot \int_{\mathbb{R}^3} f(U = u \mid X = x) \cdot f(X = x, U = u) dx du \\
&= h^{-2} \cdot \mathbb{E}\{e_U(X)\} \lesssim h^{-2}. \quad (\text{Assumption S10}).
\end{aligned}$$

Then by a standard argument based on the Chebyshev's inequality, we conclude

$$\text{pr} \left[\left| \mathbb{P}_n \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} - \mathbb{P} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \right| \geq kh^{-1}n^{-1/2} \right] \lesssim k^{-2},$$

for any $k > 0$, which implies that,

$$\begin{aligned}
\Delta_{1,n}^{(1)} &= (\mathbb{P}_n - \mathbb{P}) \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \\
&= \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}).
\end{aligned} \tag{S63}$$

On the other hand, by Taylor expansion one has

$$\Delta_{1,n}^{(2)} = \mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[c_{\rho} \{ F_{1m_1}(X \mid \hat{\beta}), F_{0m_0}(X \mid \hat{\beta}) \} f_{1m_1}(X \mid \hat{\beta}) f_{0m_0}(X \mid \hat{\beta}) - e_u(X) \right] du \right] \tag{S64}$$

$$= \mathbb{P}_n \left[\underbrace{\int_{\mathbb{R}^2} k_{u^*}(u) \left[c_{\rho} \{ F_{1m_1}(X \mid \beta^*), F_{0m_0}(X \mid \beta^*) \} f_{1m_1}(X \mid \hat{\beta}) f_{0m_0}(X \mid \hat{\beta}) - e_u(X) \right] du}_{\Delta_{1,n}^{(2,1)}} \right] \tag{S65}$$

$$+ \underbrace{\sum_{z=0,1} \mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \cdot \left[c_{\rho}^{(z)} \{ F_{1m_1}(X \mid \beta^*), F_{0m_0}(X \mid \beta^*) \} f_{1m_1}(X \mid \hat{\beta}) f_{0m_0}(X \mid \hat{\beta}) \{ F_{zm_z}(X \mid \hat{\beta}) - F_{zm_z}(X \mid \beta^*) \} \right] du \right]}_{\Delta_{1,n}^{(2,2)}}$$

$$+ \underbrace{\frac{1}{2} \sum_{z=0,1} \mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[c_{\rho}^{(zz)} \{ \xi_{1,n}, \xi_{0,n} \} f_{1m_1}(X \mid \hat{\beta}) f_{0m_0}(X \mid \hat{\beta}) \{ F_{zm_z}(X \mid \hat{\beta}) - F_{zm_z}(X \mid \beta^*) \}^2 \right] du \right]}_{\Delta_{1,n}^{(2,3)}}$$

$$+ \underbrace{\mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[c_{\rho}^{(10)} \{ \xi_{1,n}, \xi_{0,n} \} f_{1m_1}(X \mid \hat{\beta}) f_{0m_0}(X \mid \hat{\beta}) \prod_{z=0,1} \{ F_{zm_z}(X \mid \hat{\beta}) - F_{zm_z}(X \mid \beta^*) \} \right] du \right]}_{\Delta_{1,n}^{(2,4)}}$$

where $\xi_{z,n} = \delta_{z,n} F_{zm_z}(X \mid \beta^*) + (1 - \delta_{z,n}) F_{zm_z}(X \mid \hat{\beta})$, for some $\delta_{z,n} \in [0, 1]$ and $z = 0, 1$. We recall that $c^{(1)}(a, b) = \frac{\partial c(a, b)}{\partial a}$, $c^{(11)}(a, b) = \frac{\partial^2 c(a, b)}{\partial^2 a}$ and analogously for other notations, and in (S65) we use the symmetric property of the copula function under Assumption S10. With straightforward algebra, we further have

$$\begin{aligned}
&f_{1m_1}(X \mid \hat{\beta}) f_{0m_0}(X \mid \hat{\beta}) - f_{1m_1}(X \mid \beta^*) f_{0m_0}(X \mid \beta^*) \\
&= \{ f_{1m_1}(X \mid \hat{\beta}) - f_{1m_1}(X \mid \beta^*) \} \{ f_{0m_0}(X \mid \hat{\beta}) - f_{0m_0}(X \mid \beta^*) \}
\end{aligned}$$

$$\begin{aligned}
& + \{f_{1m_1}(X | \hat{\beta}) - f_{1m_1}(X | \beta^*)\} f_{0m_0}(X | \beta^*) \\
& + \{f_{0m_0}(X | \hat{\beta}) - f_{0m_0}(X | \beta^*)\} f_{1m_1}(X | \beta^*),
\end{aligned} \tag{S66}$$

meanwhile by Taylor expansion we actually have for $z = 0, 1$,

$$f_{zm_z}(X | \hat{\beta}) - f_{zm_z}(X | \beta^*) = \{f'_{zm_z}(X | \beta^*)\}^\top (\hat{\beta} - \beta^*) + (\hat{\beta} - \beta^*)^\top f''_{zm_z}(X | \tilde{\beta}_{X,m_z})(\hat{\beta} - \beta^*), \tag{S67}$$

where $\tilde{\beta}_{X,m_z} = \delta_{X,m_z} \hat{\beta} + (1 - \delta_{X,m_z}) \beta^*$ for some $\delta_{X,m_z} \in [0, 1]$. Combining (S66) and (S67), the first term on the right-hand side of (S65) equals to

$$\begin{aligned}
\Delta_{1,n}^{(2,1)} &= \mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X | \beta^*), F_{0m_0}(X | \beta^*)\} \right. \\
&\quad \cdot \{f_{1m_1}(X | \hat{\beta}) f_{0m_0}(X | \hat{\beta}) - f_{1m_1}(X | \beta^*) f_{0m_0}(X | \beta^*)\} du \Big] \\
&= \mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X | \beta^*), F_{0m_0}(X | \beta^*)\} \right. \\
&\quad \cdot \left[(\hat{\beta} - \beta^*)^\top f'_{1m_1}(X | \beta^*) \{f'_{0m_0}(X | \beta^*)\}^\top (\hat{\beta} - \beta^*) \right. \\
&\quad + (\hat{\beta} - \beta^*)^\top f'_{1m_1}(X | \beta^*) \cdot (\hat{\beta} - \beta^*)^\top f''_{0m_0}(X | \tilde{\beta}_{X,m_0})(\hat{\beta} - \beta^*) \\
&\quad + (\hat{\beta} - \beta^*)^\top f'_{0m_0}(X | \beta^*) \cdot (\hat{\beta} - \beta^*)^\top f''_{1m_1}(X | \tilde{\beta}_{X,m_1})(\hat{\beta} - \beta^*) \\
&\quad \left. \left. + (\hat{\beta} - \beta^*)^\top f''_{0m_0}(X | \tilde{\beta}_{X,m_0})(\hat{\beta} - \beta^*) \cdot (\hat{\beta} - \beta^*)^\top f''_{1m_1}(X | \tilde{\beta}_{X,m_1})(\hat{\beta} - \beta^*) \right] du \right] \\
&\quad + \sum_{z=0,1} \mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X | \beta^*), F_{0m_0}(X | \beta^*)\} \right. \\
&\quad \cdot \left[f_{(1-z)m_{1-z}}(X | \beta^*) \{f'_{zm_z}(X | \beta^*)\}^\top (\hat{\beta} - \beta^*) \right. \\
&\quad \left. \left. + f_{(1-z)m_{1-z}}(X | \beta^*) \cdot (\hat{\beta} - \beta^*)^\top f''_{zm_z}(X | \tilde{\beta}_{X,m_z})(\hat{\beta} - \beta^*) \right] du \right].
\end{aligned} \tag{S68}$$

Since $\hat{\beta} - \beta^* = \mathcal{O}_{\mathbb{P}}(n^{-1/2})$ under a parametric model of \mathcal{M}_{ps} , we have with probability approaching 1 (wpa1), $\|\tilde{\beta}_{x,m_z} - \beta^*\| \leq \delta_0$ uniformly for all $x, m_z \in \mathbb{R}$ when n is sufficiently large. Denote $\|\cdot\|_2$ the matrix spectral norm. Then by Assumption S13, we have both $\sup_{m_z, x \in \mathbb{R}} \|f''_{zm_z}(x | \tilde{\beta}_{x,m_z})\|_2$ and $\sup_{m_z, x \in \mathbb{R}} \|f'_{zm_z}(x | \tilde{\beta}_{x,m_z})\|$ are $\mathcal{O}(1)$ for both $z = 0, 1$. By the bounds of $k_{u^*}(\cdot, \cdot)$ and $c(\cdot, \cdot)$ given in Assumptions S9 and S10, we can bound the first and second terms in (S68) as following,

$$\mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} (\hat{\beta} - \beta^*)^\top f'_{1m_1}(X | \beta^*) \{f'_{0m_0}(X | \beta^*)\}^\top (\hat{\beta} - \beta^*) du \right]$$

$$\begin{aligned}
&\leq \sup_{x \in \mathbb{R}} \left| (\hat{\beta} - \beta^*)^\top \int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(x), F_{0m_0}(x)\} f'_{1m_1}(x | \beta^*) \{f'_{0m_0}(x | \beta^*)\}^\top du \cdot (\hat{\beta} - \beta^*) \right| \\
&\leq \|\hat{\beta} - \beta^*\|^2 \cdot \sup_{(u,v) \in \mathbb{R}^2} |c(u,v)| \cdot \prod_{z=1,0} \sup_{m_z \in \mathbb{R}, x \in \mathbb{R}^{d_X}} \|f'_{zm_z}(x | \beta^*)\| \underbrace{\sqrt{\int_{\mathbb{R}^2} k_{u^*}^2(u) du}}_{\mathcal{O}(h^{-1}) \text{ by Assumption S9}} \\
&= \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1})
\end{aligned}$$

and similarly,

$$\begin{aligned}
&\mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} (\hat{\beta} - \beta^*)^\top f'_{1m_1}(X | \beta^*) \cdot (\hat{\beta} - \beta^*)^\top f''_{0m_0}(X | \tilde{\beta}_{X,m_0}) (\hat{\beta} - \beta^*) du \right] \\
&\leq \|\hat{\beta} - \beta^*\|^3 \sup_{(u,v) \in \mathbb{R}^2} |c(u,v)| \sup_{m_1 \in \mathbb{R}, x \in \mathbb{R}^{d_X}} \|f'_{1m_1}(x | \beta^*)\| \sup_{m_0 \in \mathbb{R}, x \in \mathbb{R}^{d_X}} \|f''_{0m_0}(X | \tilde{\beta}_{X,m_0})\|_2 \cdot \sqrt{\int_{\mathbb{R}^2} k_{u^*}^2(u) du} \\
&= \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-3/2}).
\end{aligned}$$

With similar argument, we can finally show the first term in (S65) is

$$\begin{aligned}
\Delta_{1,n}^{(2,1)} &= \sum_{z=0,1} \mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X | \beta^*) \{f'_{zm_z}(X | \beta^*)\}^\top du \right] (\hat{\beta} - \beta^*) \\
&\quad + \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}) \\
&= \sum_{z=0,1} \mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X | \beta^*) \{f'_{zm_z}(X | \beta^*)\}^\top du \right] (\hat{\beta} - \beta^*) \\
&\quad + \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2});
\end{aligned}$$

Here in the last equality, by Chebyshev's inequality, we can easily show that,

$$\begin{aligned}
&(\mathbb{P}_n - \mathbb{P}) \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X | \beta^*) \{f'_{zm_z}(X | \beta^*)\}^\top du \right] \cdot (\hat{\beta} - \beta^*) \\
&= \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}) \cdot \mathcal{O}_{\mathbb{P}}(n^{-1/2}) = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1}).
\end{aligned}$$

Note here $\int_{\mathbb{R}^2} k_{u^*}(u) c_\rho \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X | \beta^*) \{f'_{zm_z}(X | \beta^*)\}^\top du$ is a random variable with finite variance under Assumptions S9–S10. With similar argument, we can finally show the second term on the right-hand side of (S65) is

$$\begin{aligned}
\Delta_{1,n}^{(2,2)} &= \sum_{z=0,1} \mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[c_\rho^{(z)} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1m_1}(X | \beta^*) f_{0m_0}(X | \beta^*) \{f'_{zm_z}(X | \beta^*)\}^\top \right] du \right] (\hat{\beta} - \beta^*) \\
&\quad + \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}), \tag{S69}
\end{aligned}$$

and $\Delta_{1,n}^{(2,3)} - \Delta_{1,n}^{(2,4)}$ on the right-hand side of (S65) are both $o_{\mathbb{P}}(h^{-1}n^{-1/2})$. Summarizing the results above, we finally conclude,

$$\begin{aligned}
\Delta_{1,n} &= (\mathbb{P}_n - \mathbb{P}) \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \\
&+ \sum_{z=0,1} \mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_{\rho} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X | \beta^*) \{f'_{zm_z}(X | \beta^*)\}^{\top} du \right] (\hat{\beta} - \beta^*) \\
&+ \sum_{z=0,1} \mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[c_{\rho}^{(z)} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1m_1}(X | \beta^*) f_{0m_0}(X | \beta^*) \{F'_{zm_z}(X | \beta^*)\}^{\top} \right] du \right] (\hat{\beta} - \beta^*) \\
&+ o_{\mathbb{P}}(h^{-1}n^{-1/2}) \\
&= \underbrace{\Gamma_{1,n}}_{\mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2})} + o_{\mathbb{P}}(h^{-1}n^{-1/2}) = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}), \tag{S70}
\end{aligned}$$

where the last equality follows from (S63), $\|\hat{\beta} - \beta^*\| = \mathcal{O}_{\mathbb{P}}(n^{-1/2})$, and the facts that for $z \in \{0, 1\}$,

$$\begin{aligned}
&\left\| \mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}(u) c_{\rho} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{(1-z)m_{1-z}}(X | \beta^*) \{f'_{zm_z}(X | \beta^*)\}^{\top} du \right] \right\| \\
&\leq \sup_{(u,v) \in \mathbb{R}^2} c_{\rho}(u, v) \sup_{m_{1-z} \in \mathbb{R}, x \in \mathbb{R}^{d_X}} \left| f_{(1-z)m_{1-z}}(x | \beta^*) \right| \sup_{m_z \in \mathbb{R}, x \in \mathbb{R}^{d_X}} \|f'_{zm_z}(x | \beta^*)\| \cdot \sqrt{\int_{\mathbb{R}^2} k_{u^*}^2(u) du} \\
&\lesssim h^{-1}. \tag{S71}
\end{aligned}$$

Under Assumptions S10 and S13, we similarly have,

$$\left\| \mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[c_{\rho}^{(z)} \{F_{1m_1}(X), F_{0m_0}(X)\} f_{1m_1}(X | \beta^*) f_{0m_0}(X | \beta^*) \{F'_{zm_z}(X | \beta^*)\}^{\top} \right] du \right] \right\| \lesssim h^{-1}.$$

S5.6.2 Bounding $\Delta_{2,n}$

We now bound $\Delta_{2,n}$. Decompose $\Delta_{2,n}$ as

$$\begin{aligned}
\Delta_{2,n} &= \underbrace{\mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[\{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\} \{\hat{e}_u(X) - e_u(X)\} \right] du \right]}_{\Delta_{2,n}^{(1)}} \\
&+ \underbrace{\mathbb{P}_n \left[\hat{\xi}_1 \{Y - \hat{\mu}_{ZM}(X)\} - \bar{\xi}_1 \{Y - \bar{\mu}_{ZM}(X)\} \right] + \mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[\{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\} - \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} \right] e_u(X) du \right]}_{\Delta_{2,n}^{(2)}}.
\end{aligned}$$

We bound the two terms above respectively. For $\Delta_{2,n}^{(1)}$, we further decompose it as

$$\begin{aligned} \Delta_{2,n}^{(1)} &= \underbrace{\mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[\{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} \{\hat{e}_u(X) - e_u(X)\} \right] du \right]}_{\Delta_{2,n}^{(1,1)}} \\ &\quad + \underbrace{\mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \left[\{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\} - \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} \right] \{\hat{e}_u(X) - e_u(X)\} du \right]}_{\Delta_{2,n}^{(1,2)}}. \end{aligned}$$

A similar argument to (S64)–(S69) that handles $\mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{\hat{e}_u(X) - e_u(X)\} du \right]$ can similarly show that

$$\begin{aligned} &\Delta_{2,n}^{(1,1)} \\ &= \sum_{z=0,1} \mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} c_\rho \{F_{1m_1}(X | \beta^*), F_{0m_0}(X | \beta^*)\} f_{(1-z)m_{1-z}}(X | \beta^*) \{f'_{zm_z}(X | \beta^*)\}^\top du \right] (\hat{\beta} - \beta^*) \\ &\quad + \sum_{z=0,1} \mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} c_\rho^{(a)} \{F_{1m_1}(X | \beta^*), F_{0m_0}(X | \beta^*)\} f_{1m_1}(X | \beta^*) f_{0m_0}(X | \beta^*) \{F'_{zm_z}(X | \beta^*)\}^\top du \right] (\hat{\beta} - \beta^*) \\ &\quad + o_{\mathbb{P}}(h^{-1}n^{-1/2}) \\ &= \underbrace{\Gamma_{2,n}^{(1)}}_{\mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2})} + o_{\mathbb{P}}(h^{-1}n^{-1/2}), \end{aligned}$$

under Assumptions 4 and S9. Note $\Delta_{2,n}^{(1,2)}$ is built by $\mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{\hat{e}_u(X) - e_u(X)\} du \right]$ with an additional factor $\{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\} - \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}$ in the integrand. Thus we can similarly use the Taylor expansion technique shown in (S64)–(S69) to deduce

$$\mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{\hat{e}_u(X) - e_u(X)\} du \right] = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}),$$

with a uniform bound such that

$$\sup_{x \in \mathbb{R}^{d_X}} \left| \{\hat{\mu}_{1m_1}(x) - \hat{\mu}_{0m_0}(x)\} - \{\bar{\mu}_{1m_1}(x) - \bar{\mu}_{0m_0}(x)\} \right| = o_{\mathbb{P}}(1),$$

under Assumption 4, thus we have $\Delta_{2,n}^{(1,2)} = o_{\mathbb{P}}(h^{-1}n^{-1/2})$. In summary, one has

$$\Delta_{2,n}^{(1)} = \Gamma_{2,n}^{(1)} + o_{\mathbb{P}}(h^{-1}n^{-1/2}).$$

For $\Delta_{2,n}^{(2)}$, we decompose

$$\begin{aligned}
\Delta_{2,n}^{(2)} &= \mathbb{P}_n \left[\frac{(-1)^{Z+1} \hat{\gamma}_1^{(Z)}(X, M)}{\hat{\pi}_Z(X) f_{ZM}(X | \hat{\beta})} \{Y - \hat{\mu}_{ZM}(X)\} \right] - \mathbb{P}_n \left[\frac{(-1)^{Z+1} \gamma_1^{(Z)}(X, M)}{\hat{\pi}_Z(X) f_{ZM}(X | \beta^*)} \{Y - \hat{\mu}_{ZM}(X)\} \right] \\
&+ \mathbb{P}_n \left[\frac{(-1)^{Z+1} \gamma_1^{(Z)}(X, M)}{\hat{\pi}_Z(X) f_{ZM}(X | \beta^*)} \{Y - \hat{\mu}_{ZM}(X)\} \right] - \mathbb{P}_n \left[\frac{(-1)^{Z+1} \gamma_1^{(Z)}(X, M)}{\bar{\pi}_Z(X) f_{ZM}(X | \beta^*)} \{Y - \bar{\mu}_{ZM}(X)\} \right] \\
&+ \mathbb{P}_n \left[\int_{\mathbb{R}^2} k_{u^*}(u) [\{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\} - \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}] e_u(X) du \right] \\
&= \mathbb{P}_n \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \left\{ \frac{\hat{\gamma}_1^{(Z)}(X, M)}{f_{ZM}(X | \hat{\beta})} - \frac{\gamma_1^{(Z)}(X, M)}{f_{ZM}(X | \beta^*)} \right\} \right] \\
&+ \mathbb{P}_n \left[\left[\frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} \{Y - \hat{\mu}_{ZM}(X)\} - \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \right] \left\{ \frac{\hat{\gamma}_1^{(Z)}(X, M)}{f_{ZM}(X | \hat{\beta})} - \frac{\gamma_1^{(Z)}(X, M)}{f_{ZM}(X | \beta^*)} \right\} \right] \\
&+ \mathbb{P}_n \left[\frac{(-1)^{Z+1} \gamma_1^{(Z)}(X, M)}{\hat{\pi}_Z(X) f_{ZM}(X | \beta^*)} \{Y - \hat{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} k_{u^*}(u) \{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\} e_u(X) du \right] \\
&- \mathbb{P}_n \left[\frac{(-1)^{Z+1} \gamma_1^{(Z)}(X, M)}{\bar{\pi}_Z(X) f_{ZM}(X | \beta^*)} \{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} k_{u^*}(u) \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} e_u(X) du \right] \\
&= \underbrace{\mathbb{P}_n \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \left\{ \frac{\hat{\gamma}_1^{(Z)}(X, M)}{f_{ZM}(X | \hat{\beta})} - \frac{\gamma_1^{(Z)}(X, M)}{f_{ZM}(X | \beta^*)} \right\} \right]}_{\Delta_{2,n}^{(2,1)}} \tag{S72} \\
&+ \underbrace{\mathbb{P}_n \left[\left[\frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} \{Y - \bar{\mu}_{ZM}(X)\} - \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \right] \left\{ \frac{\hat{\gamma}_1^{(Z)}(X, M)}{f_{ZM}(X | \hat{\beta})} - \frac{\gamma_1^{(Z)}(X, M)}{f_{ZM}(X | \beta^*)} \right\} \right]}_{\Delta_{2,n}^{(2,2)}} \\
&+ \underbrace{\mathbb{P}_n \left[\left[\frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} \{Y - \hat{\mu}_{ZM}(X)\} - \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{Y - \bar{\mu}_{ZM}(X)\} \right] \left\{ \frac{\hat{\gamma}_1^{(Z)}(X, M)}{f_{ZM}(X | \hat{\beta})} - \frac{\gamma_1^{(Z)}(X, M)}{f_{ZM}(X | \beta^*)} \right\} \right]}_{\Delta_{2,n}^{(2,3)}} \\
&+ \mathbb{P}_n \left[\frac{(-1)^{Z+1} \gamma_1^{(Z)}(X, M)}{\hat{\pi}_Z(X) f_{ZM}(X | \beta^*)} \{Y - \hat{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} k_{u^*}(u) [\{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\} e_u(X)] du \right] \\
&- \mathbb{P}_n \left[\frac{(-1)^{Z+1} \gamma_1^{(Z)}(X, M)}{\bar{\pi}_Z(X) f_{ZM}(X | \beta^*)} \{Y - \bar{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} k_{u^*}(u) [\{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} e_u(X)] du \right] \\
&= \Delta_{2,n}^{(2,1)} + \Delta_{2,n}^{(2,2)} + \Delta_{2,n}^{(2,3)} + \mathbb{P}_n \{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\},
\end{aligned}$$

where we denote

$$h(V | \pi, \mu) = \frac{(-1)^{Z+1} \gamma_1^{(Z)}(X, M)}{\pi_Z(X) f_{ZM}(X)} \{Y - \mu_{ZM}(X)\} + \int_{\mathbb{R}^2} k_{u^*}(u) \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} e_u(X) du.$$

(S73)

First, we study $\Delta_{2,n}^{(2,1)}$. By Taylor expansion, one has

$$\begin{aligned}
& \frac{\hat{\gamma}_1^{(1)}(X, M)}{f_{1M}(X | \hat{\beta})} - \frac{\gamma_1^{(1)}(X, M)}{f_{1M}(X | \beta^*)} \\
&= \int_{\mathbb{R}} k_{u^*}(M, m_0) c_\rho \{F_{1M}(X | \hat{\beta}), F_{0m_0}(X | \hat{\beta})\} f_{0m_0}(X | \hat{\beta}) dm_0 \\
&\quad - \int_{\mathbb{R}} k_{u^*}(M, m_0) c_\rho \{F_{1M}(X | \beta^*), F_{0m_0}(X | \beta^*)\} f_{0m_0}(X | \beta^*) dm_0 \\
&= \underbrace{\left[\{F'_{1M}(X | \beta^*)\}^\top \int_{\mathbb{R}} k_{u^*}(M, m_0) c_\rho^{(1)} \{F_{1M}(X), F_{0X}(m_0)\} f_{0m_0}(X) dm_0 \right.}_{\delta_{X,1,M}^{(1)}} \\
&\quad + \underbrace{\int_{\mathbb{R}} k_{u^*}(M, m_0) c_\rho^{(0)} \{F_{1M}(X), F_{0m_0}(X)\} \{F'_{0m_0}(X | \beta^*)\}^\top f_{0m_0}(X) dm_0}_{\delta_{X,1,M}^{(2)}} \\
&\quad + \left. \underbrace{\int_{\mathbb{R}} k_{u^*}(M, m_0) c_\rho \{F_{1M}(X), F_{0m_0}(X)\} \{f'_{0m_0}(X | \beta^*)\}^\top dm_0 \right] \cdot (\hat{\beta} - \beta^*)}_{\delta_{X,1,M}^{(3)}} \\
&\quad + (\tilde{\beta}_{X,1,M} - \beta^*)^\top I_{X,1,M} (\tilde{\beta}_{X,1,M} - \beta^*) \\
&= (\delta_{X,1,M}^{(1)} + \delta_{X,1,M}^{(2)} + \delta_{X,1,M}^{(3)}) (\hat{\beta} - \beta^*) + (\tilde{\beta}_{X,1,M} - \beta^*)^\top I_{X,1,M} (\tilde{\beta}_{X,1,M} - \beta^*),
\end{aligned} \tag{S74}$$

for some $\tilde{\beta}_{X,1,M} = \delta_{X,1,M} \hat{\beta} + (1 - \delta_{X,1,M}) \beta^*$ for some $\delta_{X,1,M} \in (0, 1)$. Here $\delta_{X,1,M}^{(1)}(\hat{\beta} - \beta^*)$, $\delta_{X,1,M}^{(2)}(\hat{\beta} - \beta^*)$, $\delta_{X,1,M}^{(3)}(\hat{\beta} - \beta^*)$ are the first three terms of (S74), and $I_{X,1,M}$ is the second order derivative matrix of $\gamma_1^{(1)}(X, M)/f_1(M | X, \beta)$ with respect to β , localized at $\beta = \tilde{\beta}_{X,1,M}$. In particular, one has

$$\begin{aligned}
I_{X,1,M} &= F_1''(M | X, \tilde{\beta}_{X,1,M}) \\
&\quad \cdot \int_{\mathbb{R}} k_{u^*}(M, m_0) c_\rho^{(1)} \{F_1(M | X, \tilde{\beta}_{X,1,M}), F_0(m_0 | X, \tilde{\beta}_{X,1,M})\} f_0(m_0 | X, \tilde{\beta}_{X,1,M}) dm_0 \\
&\quad + F_1'(M | X, \tilde{\beta}_{X,1,M}) \{F_1'(M | X, \tilde{\beta}_{X,1,M})\}^\top \\
&\quad \cdot \int_{\mathbb{R}} k_{u^*}(M, m_0) c_\rho^{(11)} \{F_1(M | X, \tilde{\beta}_{X,1,M}), F_0(m_0 | X, \tilde{\beta}_{X,1,M})\} f_0(m_0 | X, \tilde{\beta}_{X,1,M}) dm_0 \\
&\quad + \dots \\
&\quad + \int_{\mathbb{R}} k_{u^*}(M, m_0) c_\rho \{F_1(M | X, \tilde{\beta}_{X,1,M}), F_0(m_0 | X, \tilde{\beta}_{X,1,M})\} f_0''(m_0 | X, \tilde{\beta}_{X,1,M}) dm_0.
\end{aligned}$$

We can define $\delta_{X,0,M}^{(1)}(\hat{\beta} - \beta^*)$, $\delta_{X,0,M}^{(2)}(\hat{\beta} - \beta^*)$, $\delta_{X,0,M}^{(3)}(\hat{\beta} - \beta^*)$ and $I_{X,0,M}$ in a similar manner. We then write,

$$\begin{aligned} \Delta_{2,n}^{(2,1)} &= \sum_{j=1,2,3} \mathbb{P}_n \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \delta_{X,Z,M}^{(j)} \right] (\hat{\beta} - \beta^*) \\ &\quad + \mathbb{P}_n \left[(\tilde{\beta}_{X,1,M} - \beta^*)^\top \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} I_{X,Z,M} (\tilde{\beta}_{X,1,M} - \beta^*) \right]. \end{aligned} \quad (\text{S75})$$

For $j = 1, 2, 3$, similar to (S71), we can show that uniformly,

$$\sup_{x,z,m} \|\delta_{x,z,m}^{(j)}\| = \max \left\{ \sup_{x,m} \|\delta_{x,1,m}^{(j)}\|, \sup_{x,m} \|\delta_{x,0,m}^{(j)}\| \right\} \lesssim h^{-1}.$$

To illustrate this, we use $\sup_{x,m} \|\delta_{x,1,m}^{(1)}\|$ as an example. In particular, we have

$$\begin{aligned} &\sup_{x,m} \|\delta_{x,1,m}^{(1)}\| \\ &= \sup_{x,m} \left[\left\| \{F'_{1m}(x | \beta^*)\}^\top \int_{\mathbb{R}} k_{u^*}(m, m_0) c_\rho^{(1)} \{F_{1m}(x), F_{0x}(m_0)\} f_{0m_0}(x) dm_0 \right\| \right] \\ &\leq \sup_{x,m} \left[\left\| F'_{1m}(x | \beta^*) \right\| \cdot \left| \int_{\mathbb{R}} k_{u^*}(m, m_0) c_\rho^{(1)} \{F_{1m}(x), F_{0x}(m_0)\} f_{0m_0}(x) dm_0 \right| \right] \\ &\leq \sup_{u,v} |c_\rho^{(1)}(u, v)| \cdot \sup_{x,m} |f_{0m}(x)| \cdot \sup_{x,m} \left[\left\| F'_{1m}(x | \beta^*) \right\| \cdot \int_{\mathbb{R}} k_{u^*}(m, m_0) dm_0 \right] \\ &\lesssim h^{-1}, \end{aligned}$$

by Assumptions S9-S13. Then we have

$$\begin{aligned} &\mathbb{E} \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \delta_{X,Z,M,k}^{(j)} \right]^2 \lesssim h^{-1} \cdot \mathbb{E} \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \right]^2 \\ &\lesssim \mathbb{E} \left[\{Y - \mu_{ZM}(X)\}^2 \right] \quad (\text{Assumption S8}) \\ &\lesssim h^{-1}. \quad (\text{Assumption S12}). \end{aligned} \quad (\text{S76})$$

Here $\delta_{X,Z,M,k}^{(j)}$ represents the k th coordinate of $\delta_{X,Z,M}^{(j)}$. Finally by a similar argument as (S62)–(S63), one can show the following by Chebyshev's inequality,

$$(\mathbb{P}_n - \mathbb{P}) \left[\left\| \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \delta_{X,Z,M,k}^{(j)} \right\| \right] = \mathcal{O}_{\mathbb{P}}(h^{-1} n^{-1/2}),$$

and by an entry-wise argument, we have,

$$\left\| \left(\mathbb{P}_n - \mathbb{P} \right) \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \delta_{X,Z,M}^{(j)} \right] \right\|_2 = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}). \quad (\text{S77})$$

In addition, with an entry-wise argument, we can also show that,

$$\mathbb{E} \left[\left\| \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} I_{X,Z,M} \right\|_2 \right] \leq \mathbb{E} \left[\sum_{1 \leq l_1, l_2 \leq d_{\beta}} \left\| \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} I_{X,Z,M} \right\|_{l_1 l_2} \right] \lesssim h^{-1}, \quad (\text{S78})$$

and similarly as (S76)–(S77) we have

$$\left(\mathbb{P}_n - \mathbb{P} \right) \left[\left\| \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} I_{X,Z,M} \right\|_2 \right] = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}).$$

Summarizing the results above and by the fact that $\sup_{x,z,m} \|\tilde{\beta}_{x,z,m} - \beta^*\| = \sup_{x,z,m} \delta_{x,z,m} \|\hat{\beta} - \beta^*\| \leq \|\hat{\beta} - \beta^*\| = \mathcal{O}_{\mathbb{P}}(n^{-1/2})$, we have

$$\begin{aligned} \Delta_{2,n}^{(2,1)} &= \sum_{j=1,2,3} \mathbb{P} \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \delta_{X,Z,M}^{(j)} \right] (\hat{\beta} - \beta^*) \\ &\quad + \sum_{j=1,2,3} (\mathbb{P}_n - \mathbb{P}) \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \delta_{X,Z,M}^{(j)} \right] (\hat{\beta} - \beta^*) \\ &\quad + \left\{ (\mathbb{P}_n - \mathbb{P}) + \mathbb{P} \right\} \left[\|\tilde{\beta}_{X,1,M} - \beta^*\|^2 \left(\frac{\tilde{\beta}_{X,1,M} - \beta^*}{\|\tilde{\beta}_{X,1,M} - \beta^*\|} \right)^\top \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \cdot I_{X,Z,M} \left(\frac{\tilde{\beta}_{X,1,M} - \beta^*}{\|\tilde{\beta}_{X,1,M} - \beta^*\|} \right) \right] \\ &= \underbrace{\sum_{j=1,2,3} \mathbb{P} \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \delta_{X,Z,M}^{(j)} \right] (\hat{\beta} - \beta^*)}_{\Gamma_{2,n}^{(2,1)}} + o_{\mathbb{P}}(h^{-1}n^{-1/2}) \end{aligned}$$

where in the last equality we use the property of spectral norm such that by (S78),

$$\begin{aligned} &\left\| \left\{ (\mathbb{P}_n - \mathbb{P}) + \mathbb{P} \right\} \left[\|\tilde{\beta}_{X,1,M} - \beta^*\|^2 \left(\frac{\tilde{\beta}_{X,1,M} - \beta^*}{\|\tilde{\beta}_{X,1,M} - \beta^*\|} \right)^\top \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \cdot I_{X,Z,M} \left(\frac{\tilde{\beta}_{X,1,M} - \beta^*}{\|\tilde{\beta}_{X,1,M} - \beta^*\|} \right) \right] \right\| \\ &\leq \|\hat{\beta} - \beta^*\|^2 \left\{ (\mathbb{P}_n - \mathbb{P}) + \mathbb{P} \right\} \left[\sup_{\|v\|=1} \left| v^\top \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} I_{X,Z,M} \right| \right] \\ &= \|\hat{\beta} - \beta^*\|^2 \left\{ (\mathbb{P}_n - \mathbb{P}) + \mathbb{P} \right\} \left[\left\| \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} I_{X,Z,M} \right\|_2 \right] \\ &= \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1}), \quad (\text{S79}) \end{aligned}$$

and also for $j = 1, 2, 3$, by (S77),

$$\begin{aligned}
& \left\| (\mathbb{P}_n - \mathbb{P}) \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \delta_{X,Z,M}^{(j)} \right] (\hat{\beta} - \beta^*) \right\| \\
& \leq \left\| (\mathbb{P}_n - \mathbb{P}) \left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \delta_{X,Z,M}^{(j)} \right] \right\|_2 \|\hat{\beta} - \beta^*\| \\
& = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1}).
\end{aligned} \tag{S80}$$

Next we study $\Delta_{2,n}^{(2,2)}$. When $\bar{\pi} = \pi$ and $\bar{\mu} = \mu$, it is easy to see $\Delta_{2,n}^{(2,2)} = 0$. When either $\bar{\pi} \neq \pi$ and $\bar{\mu} \neq \mu$, similar to (S75), we have

$$\begin{aligned}
& \Delta_{2,n}^{(2,2)} \\
& = \sum_{j=1,2,3} \mathbb{P}_n \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{Y - \bar{\mu}_{ZM}(X)\} - \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \right] \delta_{X,Z,M}^{(j)} \right] (\hat{\beta} - \beta^*) \\
& \quad + \mathbb{P}_n \left[(\tilde{\beta}_{X,1,M} - \beta^*)^\top \left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{Y - \bar{\mu}_{ZM}(X)\} - \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \right] I_{X,Z,M}(\tilde{\beta}_{X,1,M} - \beta^*) \right].
\end{aligned}$$

We then exploit the same technical argument as for $\Delta_{2,n}^{(2,1)}$ to bound $\Delta_{2,n}^{(2,2)}$. In particular, recalling $\|\delta_{X,Z,M}^{(j)}\|$ can be uniformly bounded by $\mathcal{O}(h^{-1})$, we have for any $j = 1, 2, 3$ and $k \in [d_\beta]$,

$$\begin{aligned}
& \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{Y - \bar{\mu}_{ZM}(X)\} - \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \right] \delta_{X,Z,M}^{(j)} \right]^2 \\
& \lesssim h^{-2} \cdot \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{Y - \bar{\mu}_{ZM}(X)\} - \frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \right]^2 \right] \\
& \leq 2h^{-2} \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{Y - \bar{\mu}_{ZM}(X)\} \right]^2 \right] + 2h^{-2} \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \right]^2 \right] \\
& \leq 4h^{-2} \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{Y - \mu_{ZM}(X)\} \right]^2 \right] + 4h^{-2} \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{\mu_{ZM}(X) - \bar{\mu}_{ZM}(X)\} \right]^2 \right] \\
& \quad + 2h^{-2} \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \right]^2 \right] \\
& \lesssim h^{-2},
\end{aligned} \tag{S81}$$

where the first and third term in (S81) can be bounded similar to (S76), while the second term in (S81) can be bounded by Assumptions S8 and 4,

$$\begin{aligned} \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{ \mu_{ZM}(X) - \bar{\mu}_{ZM}(X) \} \right]^2 \right] &\leq \epsilon^{-2} \cdot \{ \mathbb{E} | \mu_{ZM}(X) |^2 + \mathbb{E} | \bar{\mu}_{ZM}(X) |^2 \} \\ &= \mathcal{O}(1). \end{aligned}$$

Similar to (S77), (S81) combining with Chebyshev's inequality and an entry-wise argument implies that

$$\left\| (\mathbb{P}_n - \mathbb{P}) \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{ Y - \bar{\mu}_{ZM}(X) \} - \frac{(-1)^{Z+1}}{\pi_Z(X)} \{ Y - \mu_{ZM}(X) \} \right] \delta_{X,Z,M}^{(j)} \right] \right\|_2 = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}).$$

Then, a similar argument to (S76)–(S80) can show that

$$\begin{aligned} \Delta_{2,n}^{(2,2)} &= \underbrace{\sum_{j=1,2,3} \mathbb{P} \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{ Y - \bar{\mu}_{ZM}(X) \} - \frac{(-1)^{Z+1}}{\pi_Z(X)} \{ Y - \mu_{ZM}(X) \} \right] \delta_{X,Z,M}^{(j)} \right]}_{\Gamma_{2,n}^{(2,2)}} (\hat{\beta} - \beta^*) \\ &\quad + o_{\mathbb{P}}(h^{-1}n^{-1/2}) \end{aligned}$$

Next we bound $\Delta_{2,n}^{(2,3)}$. Based on (S6), we can show

$$\begin{aligned} &\Delta_{2,n}^{(2,3)} \\ &= \mathbb{P}_n \left[\left[\frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} \{ Y - \hat{\mu}_{ZM}(X) \} - \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{ Y - \bar{\mu}_{ZM}(X) \} \right] (\delta_{X,1,M}^{(1)} + \delta_{X,1,M}^{(2)} + \delta_{X,1,M}^{(3)}) \right] (\hat{\beta} - \beta^*) \\ &\quad + \mathbb{P}_n \left[\left[\frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} \{ Y - \hat{\mu}_{ZM}(X) \} - \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{ Y - \bar{\mu}_{ZM}(X) \} \right] (\tilde{\beta}_{X,1,M} - \beta^*)^\top I_{X,1,M} (\tilde{\beta}_{X,1,M} - \beta^*) \right] \\ &= \mathbb{P}_n \left\{ (\kappa_{X,Z,Y}^{(1)} + \kappa_{X,Z,M}^{(2)}) (\delta_{X,1,M}^{(1)} + \delta_{X,1,M}^{(2)} + \delta_{X,1,M}^{(3)}) \right\} (\hat{\beta} - \beta^*) \tag{S82} \\ &\quad + \mathbb{P}_n \left\{ (\kappa_{X,Z,Y}^{(1)} + \kappa_{X,Z,M}^{(2)}) (\tilde{\beta}_{X,1,M} - \beta^*)^\top I_{X,1,M} (\tilde{\beta}_{X,1,M} - \beta^*) \right\}, \end{aligned}$$

where we further define,

$$\begin{aligned} \kappa_{X,Z,Y}^{(1)} &= \frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} Y - \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} Y, \\ \kappa_{X,Z,M}^{(2)} &= \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \bar{\mu}_{ZM}(X) - \frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} \hat{\mu}_{ZM}(X). \end{aligned}$$

We argue that all terms in (S82) are $o_{\mathbb{P}}(h^{-1}n^{-1/2})$ and thus

$$\Delta_{2,n}^{(2,3)} = o_{\mathbb{P}}(h^{-1}n^{-1/2}), \quad (\text{S83})$$

is negligible. For example, we have

$$\begin{aligned} \left| \mathbb{P}_n(\kappa_{X,Z,Y}^{(1)} \delta_{X,1,M}^{(1)})(\hat{\beta} - \beta^*) \right| &\leq \|\hat{\beta} - \beta^*\| \cdot \mathbb{P}_n(\|\kappa_{X,Z,Y}^{(1)} \delta_{X,1,M}^{(1)}\|) \\ &= o_{\mathbb{P}}(h^{-1}n^{-1/2}), \end{aligned} \quad (\text{S84})$$

where $\|\hat{\beta} - \beta^*\| = \mathcal{O}_{\mathbb{P}}(n^{-1/2})$ and

$$\begin{aligned} &\mathbb{P}_n(\|\kappa_{X,Z,Y}^{(1)} \delta_{X,1,M}^{(1)}\|) \\ = &\mathbb{P}_n \left[\left\| \left\{ \frac{\bar{\pi}_Z(X) - \hat{\pi}_Z(X)}{\hat{\pi}_Z(X) \bar{\pi}_Z(X)} \right\} Y F'_{1M}(X | \beta^*) \int_{\mathbb{R}} k_{u^*}(M, m_0) c_{\rho}^{(1)} \{F_{1M}(X), F_{0X}(m_0)\} f_{0m_0}(X) dm_0 \right\| \right] \\ \leq &\|\bar{\pi}_Z(X) - \hat{\pi}_Z(X)\|_{\infty} \|\hat{\pi}_Z^{-1}(X)\|_{\infty} \|\bar{\pi}_Z^{-1}(X)\|_{\infty} \cdot \sup_{m_1, x} \|F'_{1m_1}(x | \beta^*)\| \cdot \sup_{u, v} |c_{\rho}^{(1)}(u, v)| \cdot \sup_{m_0, x} f_{0m_0}(x) \\ &\cdot \left\{ \sup_{m_1} \int_{\mathbb{R}} k_{u^*}(u) dm_0 \right\} \mathbb{P}_n(|Y|) \end{aligned} \quad (\text{S85})$$

$$= o_{\mathbb{P}}(1) \cdot \mathcal{O}_{\mathbb{P}}(1) \cdots \mathcal{O}(1) \cdot \mathcal{O}(h^{-1}) \cdot \{(\mathbb{P}_n - \mathbb{E})(|Y|) + \mathbb{E}(|Y|)\} \quad (\text{S86})$$

$$= o_{\mathbb{P}}(h^{-1}),$$

where $\|\cdot\|_{\infty}$ denotes the sup-norm for function and random variable. For (S85), we have $\|\bar{\pi}_Z(X) - \hat{\pi}_Z(X)\|_{\infty} = o_{\mathbb{P}}(1)$ under Assumption 4, $\|\bar{\pi}_Z^{-1}(X)\|_{\infty} \geq \epsilon^{-1}$ under Assumption S8 and thus $\|\hat{\pi}_Z^{-1}(X)\|_{\infty}$ is also bounded away from zero with probability approaching 1, meanwhile all other terms are $\mathcal{O}_{\mathbb{P}}(1)$ under Assumptions S9–S13. For (S86), $\sup_{m_1} \int_{\mathbb{R}} k_{u^*}(u) dm_0 = \mathcal{O}(h^{-1})$ under Assumption S9, $\mathbb{E}(|Y|) \leq \sqrt{\mathbb{E}(|Y|^2)} < +\infty$ under Assumption S12 and then $(\mathbb{P}_n - \mathbb{E})(|Y|) = \mathcal{O}_{\mathbb{P}}(n^{-1/2})$ follows by an application of the Markov's inequality. Similarly, we can show

$$\left| \mathbb{P}_n(\kappa_{X,Z,Y}^{(1)} \delta_{X,1,M}^{(2)})(\hat{\beta} - \beta^*) \right| = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}) \quad \text{and} \quad \left| \mathbb{P}_n(\kappa_{X,Z,Y}^{(1)} \delta_{X,1,M}^{(3)})(\hat{\beta} - \beta^*) \right| = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2}).$$

We can also show that

$$\begin{aligned} \left| \mathbb{P}_n(\kappa_{X,Z,M}^{(2)} \delta_{X,1,M}^{(1)})(\hat{\beta} - \beta^*) \right| &\leq \|\hat{\beta} - \beta^*\| \mathbb{P}_n(\|\kappa_{X,Z,Y}^{(2)} \delta_{X,1,M}^{(1)}\|) \\ &= o_{\mathbb{P}}(h^{-1}n^{-1/2}), \end{aligned} \quad (\text{S87})$$

where $\|\hat{\beta} - \beta^*\| = \mathcal{O}_{\mathbb{P}}(n^{-1/2})$, and similar to (S84),

$$\begin{aligned}
& \mathbb{P}_n(\|\kappa_{X,Z,Y}^{(2)} \delta_{X,1,M}^{(1)}\|) \\
= & \mathbb{P}_n \left[\left\| \left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \bar{\mu}_{ZM}(X) - \frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} \hat{\mu}_{ZM}(X) \right] F'_{1M}(X | \beta^*) \int_{\mathbb{R}} k_{u^*}(M, m_0) \right. \right. \\
& \left. \left. \cdot c_{\rho}^{(1)} \{F_{1M}(X), F_{0X}(m_0)\} f_{0m_0}(X) dm_0 \right\| \right] \\
\leq & \left\| \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{\bar{\mu}_{ZM}(X) - \hat{\mu}_{ZM}(X)\} + \left\{ \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} - \frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} \right\} \hat{\mu}_{ZM}(X) \right\|_{\infty} \quad (\text{S88}) \\
& \cdot \sup_{m_1, x} \|F'_{1m_1}(x | \beta^*)\| \cdot \sup_{u, v} |c_{\rho}^{(1)}(u, v)| \cdot \sup_{m_0, x} f_{0m_0}(x) \cdot \sup_{m_1} \int_{\mathbb{R}} k_{u^*}(u) dm_0 \cdot \mathbb{P}_n(|Y|) \\
= & o_{\mathbb{P}}(h^{-1}).
\end{aligned}$$

For (S88), based on Assumptions S8 and 4, we bound

$$\begin{aligned}
& \left\| \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{\bar{\mu}_{ZM}(X) - \hat{\mu}_{ZM}(X)\} + \left\{ \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} - \frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} \right\} \hat{\mu}_{ZM}(X) \right\|_{\infty} \\
\leq & \left\| \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{\bar{\mu}_{ZM}(X) - \hat{\mu}_{ZM}(X)\} \right\|_{\infty} + \left\| \left\{ \frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} - \frac{(-1)^{Z+1}}{\hat{\pi}_Z(X)} \right\} \hat{\mu}_{ZM}(X) \right\|_{\infty} \\
\leq & \epsilon^{-1} \|\bar{\mu}_{ZM}(X) - \hat{\mu}_{ZM}(X)\|_{\infty} + \left\| \frac{\hat{\pi}_Z(X) - \bar{\pi}_Z(X)}{\bar{\pi}_Z(X) \hat{\pi}_Z(X)} \right\|_{\infty} \|\hat{\mu}_{ZM}(X)\|_{\infty} \\
\leq & \epsilon^{-1} \|\bar{\mu}_{ZM}(X) - \hat{\mu}_{ZM}(X)\|_{\infty} + \|\bar{\pi}_Z(X) - \hat{\pi}_Z(X)\|_{\infty} \|\hat{\pi}_Z^{-1}(X)\|_{\infty} \|\bar{\pi}_Z^{-1}(X)\|_{\infty} \\
& \cdot \left\{ \|\bar{\mu}_{ZM}(X)\|_{\infty} + \|\hat{\mu}_{ZM}(X) - \bar{\mu}_{ZM}(X)\|_{\infty} \right\} \\
= & o_{\mathbb{P}}(1).
\end{aligned}$$

By applying the similar technical arguments as above to each entry of $\mathbb{P}_n(\kappa_{X,Z,Y}^{(1)} I_{X,1,M})$ and $\mathbb{P}_n(\kappa_{X,Z,M}^{(2)} I_{X,1,M})$, we can show that $\mathbb{P}_n(\|\kappa_{X,Z,Y}^{(1)} I_{X,1,M}\|_2)$, $\mathbb{P}_n(\|\kappa_{X,Z,M}^{(2)} I_{X,1,M}\|_2) = \mathcal{O}_{\mathbb{P}}(h^{-1})$ similar to (S78). Then by the basic property of spectral norm and similar to (S79), one has

$$\begin{aligned}
& \left| \mathbb{P}_n \left\{ (\kappa_{X,Z,Y}^{(1)} + \kappa_{X,Z,M}^{(2)}) (\tilde{\beta}_{X,1,M} - \beta^*)^{\top} I_{X,1,M} (\tilde{\beta}_{X,1,M} - \beta^*) \right\} \right| \\
\leq & \|\hat{\beta} - \beta\|^2 \left\{ \mathbb{P}_n(\|\kappa_{X,Z,Y}^{(1)} I_{X,1,M}\|_2) + \mathbb{P}_n(\|\kappa_{X,Z,M}^{(2)} I_{X,1,M}\|_2) \right\} \\
= & \mathcal{O}_{\mathbb{P}}(h^{-1} n^{-1}). \quad (\text{S89})
\end{aligned}$$

Combining (S82)–(S89), (S83) is thus shown.

Summarizing the results above, we have

$$\Delta_{2,n} = \Gamma_{2,n}^{(1)} + \Gamma_{2,n}^{(2,1)} + \Gamma_{2,n}^{(2,2)} + o_{\mathbb{P}}(h^{-1}n^{-1/2}) + \mathbb{P}_n\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\} \quad (\text{S90})$$

We now study the last term on the right-hand side above. We decompose

$$\begin{aligned} & \mathbb{P}_n\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\} \\ &= (\mathbb{P}_n - \mathbb{P})\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\} + \mathbb{P}\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\}. \end{aligned}$$

We first bound $(\mathbb{P}_n - \mathbb{P})\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\}$ through Lemma S4 using empirical process theory. We defer a brief introduction of empirical process theory as well as the technical proof of Lemma S4 to Section S6.

Lemma S4 *Recall (S73). Under the conditions of Theorem 5, we have*

$$(\mathbb{P}_n - \mathbb{P})\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\} = o_{\mathbb{P}}(h^{-1}n^{-1/2}). \quad (\text{S91})$$

By Lemma S4, we have

$$\mathbb{P}_n\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\} = o_{\mathbb{P}}(h^{-1}n^{-1/2}) + \mathbb{P}\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\}.$$

In addition, if either $\bar{\mu} = \mu$ or $\bar{\pi} = \pi$, one has

$$\begin{aligned} \mathbb{P}\{h(V | \bar{\pi}, \bar{\mu})\} &= \mathbb{E}\left[\bar{\xi}_1(X, Z, M)\{Y - \mu_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right] \\ &= \bar{\tau}_{u^*} \cdot \mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\} \\ &= \tau_{u^*} \cdot \mathbb{E}\left\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\right\} \quad (\text{Theorem 4}) \\ &= \mathbb{E}\left[\int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right] \\ &= \mathbb{E}\left[\xi_1(X, Z, M)\{Y - \mu_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\}k_{u^*}(u)e_u(X)du\right] \\ &= \mathbb{P}\{h(V | \pi, \mu)\}, \end{aligned}$$

where the fourth equality is by Theorem 2, the fifth equality is by $\mathbb{E}\tilde{\xi}_1 = 0$ when $\tilde{\mu} = \mu$; See (S60).

Therefore we have,

$$\begin{aligned}
& \mathbb{P}\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\} \tag{S92} \\
&= \mathbb{P}\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \pi, \mu)\} \\
&= \mathbb{P}\left[\frac{(-1)^{Z+1}\gamma_1^{(Z)}(X, M)}{\hat{\pi}_Z(X)f_Z(M | X, \beta^*)} \{Y - \hat{\mu}_{ZM}(X)\} + \int_{\mathbb{R}^2} k_{u^*}(u) \{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\} e_u(X) du \right. \\
&\quad \left. - \frac{(-1)^{Z+1}\gamma_1^{(Z)}(X, M)}{\pi_Z(X)f_Z(M | X, \beta^*)} \{Y - \mu_{ZM}(X)\} - \int_{\mathbb{R}^2} k_{u^*}(u) \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} e_u(X) du \right] \\
&= \mathbb{P}\left[\frac{(-1)^{Z+1}\gamma_1^{(Z)}(X, M)}{\hat{\pi}_Z(X)f_Z(M | X, \beta^*)} \{Y - \hat{\mu}_{ZM}(X)\} - \frac{(-1)^{Z+1}\gamma_1^{(Z)}(X, M)}{\pi_Z(X)f_Z(M | X, \beta^*)} \{Y - \mu_{ZM}(X)\} \right] \\
&\quad + \mathbb{P}\left[\int_{\mathbb{R}^2} k_{u^*}(u) \{\hat{\mu}_{1m_1}(X) - \hat{\mu}_{0m_0}(X)\} e_u(X) du - \int_{\mathbb{R}^2} k_{u^*}(u) \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} e_u(X) du \right]. \tag{S93}
\end{aligned}$$

By LOTE, the first term in (S93) equals to,

$$\begin{aligned}
& \sum_{z=0,1} \mathbb{P}\left[\mathbb{P}\left[\pi_z(X) \frac{(-1)^{z+1}\gamma_1^{(z)}(X, M)}{\hat{\pi}_z(X)f_z(M | X, \beta^*)} \{Y - \hat{\mu}_{zM}(X)\} \right. \right. \\
&\quad \left. \left. - \frac{(-1)^{z+1}\gamma_1^{(z)}(X, M)}{f_z(M | X, \beta^*)} \{Y - \mu_{zM}(X)\} \mid X, Z = z \right] \right] \\
&= \sum_{z=0,1} \mathbb{P}\left[\mathbb{P}\left[\mathbb{P}\left[\pi_z(X) \frac{(-1)^{z+1}\gamma_1^{(z)}(X, M)}{\hat{\pi}_z(X)f_z(M | X, \beta^*)} \{Y - \hat{\mu}_{zM}(X)\} \right. \right. \right. \\
&\quad \left. \left. - \frac{(-1)^{z+1}\gamma_1^{(z)}(X, M)}{f_z(M | X, \beta^*)} \{Y - \mu_{zM}(X)\} \mid X, Z = z, M \right] \mid X, Z = z \right] \\
&= \sum_{z=0,1} \mathbb{P}\left[\mathbb{P}\left[\pi_z(X) \frac{(-1)^{z+1}\gamma_1^{(z)}(X, M)}{\hat{\pi}_z(X)f_z(M | X, \beta^*)} \{\mu_{zM}(X) - \hat{\mu}_{zM}(X)\} \right. \right. \\
&\quad \left. \left. - \frac{(-1)^{z+1}\gamma_1^{(z)}(X, M)}{f_z(M | X, \beta^*)} \underbrace{\{\mu_{zM}(X) - \mu_{zM}(X)\}}_{=0} \mid X, Z = z \right] \right] \\
&= \sum_{z=0,1} \mathbb{P}\left[\frac{\pi_z(X)(-1)^{z+1}}{\hat{\pi}_z(X)} \mathbb{P}\left[\frac{\gamma_1^{(z)}(X, M)}{f_z(M | X, \beta^*)} \{\mu_{zM}(X) - \hat{\mu}_{zM}(X)\} \mid X, Z = z \right] \right] \\
&= \sum_{z=0,1} \mathbb{P}\left[\frac{\pi_z(X)(-1)^{z+1}}{\hat{\pi}_z(X)} \int_{\mathbb{R}^2} k_{u^*}(u) \{\mu_{zm_z} - \hat{\mu}_{zm_z}(X)\} e_u(X) du \right],
\end{aligned}$$

which combining with (S92) yields

$$\begin{aligned} & \mathbb{P}\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\} \\ &= \sum_{z=0,1} (-1)^{z+1} \cdot \mathbb{P} \left[\left\{ \frac{\pi_z(X)}{\hat{\pi}_z(X)} - 1 \right\} \int_{\mathbb{R}^2} k_{u^*}(u) \{\mu_{zm_z}(X) - \hat{\mu}_{zm_z}(X)\} e_u(X) du \right]. \end{aligned}$$

For either $z = 0$ or 1 , by Cauchy–Schwartz inequality, we have

$$\begin{aligned} & \left| \mathbb{P} \left[\left\{ \frac{\pi_z(X)}{\hat{\pi}_z(X)} - 1 \right\} \int_{\mathbb{R}^2} k_{u^*}(u) \{\mu_{zm_z}(X) - \hat{\mu}_{zm_z}(X)\} e_u(X) du \right] \right| \\ & \leq \left| \mathbb{P} \left\{ \frac{\pi_z(X) - \hat{\pi}_z(X)}{\hat{\pi}_z(X)} \right\}^2 \right|^{1/2} \cdot \sqrt{\int_{\mathbb{R}^2} k_{u^*}^2(u) du} \cdot \left| \mathbb{P} \left[\int_{\mathbb{R}^2} \{\mu_{zm_z}(X) - \hat{\mu}_{zm_z}(X)\}^2 e_u^2(X) du \right] \right|^{1/2} \\ & = \mathcal{O}_{\mathbb{P}}(h^{-1} r_{\pi} r_{\mu}). \end{aligned} \tag{S94}$$

To see (S94), firstly we note by Assumptions S8 and 4 and LOTE,

$$\begin{aligned} \mathbb{P}[\{\pi_z(X) - \hat{\pi}_z(X)\}^2] &= \mathbb{P}[\mathbb{P}[\{\pi_Z(X) - \hat{\pi}_Z(X)\}^2 | X, Z = z]] \\ &\leq \pi_z(X)^{-1} \cdot \mathbb{P}[\pi_z(X) \mathbb{P}[\{\pi_Z(X) - \hat{\pi}_Z(X)\}^2 | X, Z = z] \\ &\quad + \pi_{1-z}(X) \mathbb{P}[\{\pi_Z(X) - \hat{\pi}_Z(X)\}^2 | X, Z = 1 - z]] \\ &= \pi_z(X)^{-1} \cdot \mathbb{P}[\mathbb{P}[\{\pi_Z(X) - \hat{\pi}_Z(X)\}^2 | X]] \\ &= \pi_z(X)^{-1} \cdot \mathbb{P}[\{\pi_Z(X) - \hat{\pi}_Z(X)\}^2] \\ &= \mathcal{O}_{\mathbb{P}}(r_{\pi}^2). \end{aligned}$$

and thus

$$\left| \mathbb{P} \left\{ \frac{\pi_z(X) - \hat{\pi}_z(X)}{\hat{\pi}_z(X)} \right\}^2 \right| \leq \|\hat{\pi}^{-1}\|_{\infty}^2 \cdot |\mathbb{P}\{\pi_z(X) - \hat{\pi}_z(X)\}^2| = \mathcal{O}_{\mathbb{P}}(r_{\pi}^2),$$

because $\|\hat{\pi}^{-1}\|_{\infty} \lesssim 1$ with probability approaching 1 under Assumptions S8 and 4, where $\bar{\pi}_z(X) \lesssim 1$ and $\|\hat{\pi} - \bar{\pi}\|_{\infty} = o_{\mathbb{P}}(1)$. Secondly, we have

$$\mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}^2(u) du \int_{\mathbb{R}^2} \{\mu_{zm_z}(X) - \hat{\mu}_{zm_z}(X)\}^2 e_u^2(X) du \right]$$

$$\begin{aligned}
&\lesssim h^{-2} \cdot \mathbb{P} \left[\int_{\mathbb{R}^2} \{\mu_{zm_z}(X) - \hat{\mu}_{zm_z}(X)\}^2 e_u^2(X) du \right] \quad (\text{Assumption S9}) \\
&\lesssim h^{-2} \cdot \mathbb{P} \left[\int_{\mathbb{R}^2} \{\mu_{zm_z}(X) - \hat{\mu}_{zm_z}(X)\}^2 e_u(X) du \right] \\
&= h^{-2} \cdot \mathbb{P} \left[\mathbb{P} \left[\{\mu_{zM_z}(X) - \hat{\mu}_{zM_z}(X)\}^2 \mid X \right] \right] \\
&= h^{-2} \cdot \mathbb{P} \left[\mathbb{P} \left[\{\mu_{ZM}(X) - \hat{\mu}_{ZM}(X)\}^2 \mid X, Z = z \right] \right] \quad (\text{Assumption 1}) \\
&\leq h^{-2} \cdot \pi_z^{-1}(X) \cdot \mathbb{P} \left[\pi_z(X) \mathbb{P} \left[\{\mu_{ZM}(X) - \hat{\mu}_{ZM}(X)\}^2 \mid X, Z = z \right] \right. \\
&\quad \left. + \pi_{1-z}(X) \mathbb{P} \left[\{\mu_{ZM}(X) - \hat{\mu}_{ZM}(X)\}^2 \mid X, Z = 1 - z \right] \right] \\
&= h^{-2} \cdot \pi_z^{-1}(X) \cdot \mathbb{P} \left[\mathbb{P} \left[\{\mu_{ZM}(X) - \hat{\mu}_{ZM}(X)\}^2 \mid X \right] \right] \\
&= h^{-2} \cdot \pi_z^{-1}(X) \cdot \mathbb{P} \left[\{\mu_{ZM}(X) - \hat{\mu}_{ZM}(X)\}^2 \right] \\
&= \mathcal{O}(h^{-2} r_\mu^2) \quad (\text{Assumptions S8 and 4}),
\end{aligned}$$

where the second inequality is by $e_u(x)$ is uniformly bounded by some constants for all $x \in \mathbb{X}$ and $u \in \mathbb{R}^2$, under Assumptions S10 and S13. Summarizing results above, (S94) is thus shown. We finally conclude,

$$\begin{aligned}
&\mathbb{P}\{h(V \mid \hat{\pi}, \hat{\mu}) - h(V \mid \bar{\pi}, \bar{\mu})\} = \mathcal{O}_{\mathbb{P}}(h^{-1} r_\pi r_\mu) \\
&\text{and } \mathbb{P}_n\{h(V \mid \hat{\pi}, \hat{\mu}) - h(V \mid \bar{\pi}, \bar{\mu})\} = \mathcal{O}_{\mathbb{P}}(h^{-1} r_\pi r_\mu) + o_{\mathbb{P}}(h^{-1} n^{-1/2}),
\end{aligned}$$

which combining with (S90) yields

$$\begin{aligned}
\Delta_{2,n} &= \Gamma_{2,n}^{(1)} + \Gamma_{2,n}^{(2,1)} + \Gamma_{2,n}^{(2,2)} + \mathcal{O}_{\mathbb{P}}(h^{-1} r_\pi r_\mu) + o_{\mathbb{P}}(h^{-1} n^{-1/2}) \\
&= \underbrace{\Gamma_{2,n}}_{\mathcal{O}_{\mathbb{P}}(h^{-1} n^{-1/2})} + \mathcal{O}_{\mathbb{P}}(h^{-1} r_\pi r_\mu) + o_{\mathbb{P}}(h^{-1} n^{-1/2}), \tag{S95}
\end{aligned}$$

where in summary, we define,

$$\begin{aligned}
&\Gamma_{2,n} \\
&= \sum_{z=0,1} \mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} c_\rho \{F_{1m_1}(X \mid \beta^*), F_{0m_0}(X \mid \beta^*)\} f_{(1-z)m_{1-z}}(X \mid \beta^*) \{f'_{zm_z}(X \mid \beta^*)\}^\top du \right] (\hat{\beta} - \beta^*)
\end{aligned}$$

$$\begin{aligned}
& + \sum_{z=0,1} \mathbb{P} \left[\int_{\mathbb{R}^2} k_{u^*}(u) \{ \bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X) \} c_\rho^{(z)} \{ F_{1m_1}(X | \beta^*), F_{0m_0}(X | \beta^*) \} \right. \\
& \quad \left. f_{1m_1}(X | \beta^*) f_{0m_0}(X | \beta^*) \{ F'_{zm_z}(X | \beta^*) \}^\top du \right] \cdot (\hat{\beta} - \beta^*) \\
& + \sum_{j=1,2,3} \mathbb{P} \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{ Y - \bar{\mu}_{ZM}(X) \} \right] \delta_{X,Z,M}^{(j)} \right] (\hat{\beta} - \beta^*).
\end{aligned}$$

S5.6.3 Putting all pieces together

Combining (S61), (S70) and (S95), we have that, in general,

$$\begin{aligned}
& \hat{\tau}_{u^*} - \tau_{u^*} \\
& = (\mathbb{P}_n - \mathbb{P}) \left[\frac{\bar{\xi}_1 \{ Y - \mu_{ZM}(X) \} + \int_{\mathbb{R}^2} \{ \bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X) \} k_{u^*}(u) e_u(X) du}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} \right] \\
& \quad - \frac{\mathbb{E} \left[\bar{\xi}_1 \{ Y - \mu_{ZM}(X) \} + \int_{\mathbb{R}^2} \{ \bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X) \} k_{u^*}(u) e_u(X) du \right]}{\left[\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} \right]^2} \left[(\mathbb{P}_n - \mathbb{P}) \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} + \mathcal{A}_1(\hat{\beta} - \beta^*) \right] \\
& \quad + \frac{\mathcal{A}_2}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} (\hat{\beta} - \beta^*) + \mathcal{O}_{\mathbb{P}}(h^{-1} r_\pi r_\mu) + o_{\mathbb{P}}(h^{-1} n^{-1/2}) \\
& = (\mathbb{P}_n - \mathbb{P}) \left[\frac{\bar{\xi}_1 \{ Y - \mu_{ZM}(X) \} + \int_{\mathbb{R}^2} \{ \bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X) \} k_{u^*}(u) e_u(X) du - \tau_{u^*} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} \right] \tag{S96}
\end{aligned}$$

$$\begin{aligned}
& \quad + \frac{\mathcal{A}_2 - \tau_{u^*} \mathcal{A}_1}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} (\hat{\beta} - \beta^*) + \mathcal{O}_{\mathbb{P}}(h^{-1} r_\pi r_\mu) + o_{\mathbb{P}}(h^{-1} n^{-1/2}) \\
& = \mathcal{O}_{\mathbb{P}}(h^{-1} n^{-1/2} + h^{-1} r_\pi r_\mu), \tag{S97}
\end{aligned}$$

where the second equality is by Theorem 4 under Assumption 4(1); The first two terms in (S96) are $\mathcal{O}_{\mathbb{P}}(h^{-1} n^{-1/2})$ with a standard argument similar as before. We note that when h is fixed, we have $\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} > 0$ is fixed, and if $h \rightarrow 0$,

$$\begin{aligned}
\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\} & = \mathbb{E} \left[\left\{ k_{u^*}(U) \mid X \right\} \right] \\
& = \mathbb{E} [k_{u^*}(U)] \\
& = e_{u^*} + \mathcal{O}(h^2) \lesssim 1 \tag{S98}
\end{aligned}$$

by (S20), which is also bounded away from 0. We define \mathcal{A}_1 and \mathcal{A}_2 in (S1) and (S2), $\delta_{X,1,M}^{(1)} - \delta_{X,1,M}^{(3)}$ in (S6), and $\delta_{X,0,M}^{(1)} - \delta_{X,0,M}^{(3)}$ in (S7).

In the following, we focus on the fixed- h setting for the asymptotic analysis of Loc.PCE. Then (S97) becomes

$$\hat{\tau}_{u^*} - \tau_{u^*} = \mathcal{O}_{\mathbb{P}}(n^{-1/2} + r_\pi r_\mu),$$

which implies (13). If moreover, $r_\pi r_\mu = o(n^{-1/2})$ and $\hat{\beta}$ is asymptotically linear, we further have,

$$\begin{aligned}
& \sqrt{n}(\hat{\tau}_{u^*} - \tau_{u^*}) \\
= & \sqrt{n}(\mathbb{P}_n - \mathbb{P}) \left[\frac{\bar{\xi}_1\{Y - \mu_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} k_{u^*}(u) e_u(X) du - \tau_{u^*} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}} \right] \\
& + \frac{\mathcal{A}_2 - \tau_{u^*} \mathcal{A}_1}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}} (\hat{\beta} - \beta^*) + o_{\mathbb{P}}(1) \\
= & \sqrt{n}(\mathbb{P}_n - \mathbb{P}) \left[\frac{\bar{\xi}_1\{Y - \mu_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} k_{u^*}(u) e_u(X) du - \tau_{u^*} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}} \right. \\
& \left. + \frac{\mathcal{A}_2 - \tau_{u^*} \mathcal{A}_1}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}} \psi(X, Z, M) \right] + o_{\mathbb{P}}(1), \tag{S99}
\end{aligned}$$

$$\rightsquigarrow \mathcal{N}(0, \bar{\sigma}_{u^*}^2) \tag{S100}$$

where the second equality is by that $\psi(X, Z, M)$ is mean-zero and $\|\{\mathcal{A}_2 - \tau_{u^*} \mathcal{A}_1\} [\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}]^{-1}\|$ is bounded by some constant, which can be easily shown with a similar argument as the previous proof. Finally, (S100) follows from the central limit theorem (CLT) and Slutsky's theorem, as the random variable in $\sqrt{n}(\mathbb{P}_n - \mathbb{P})[\cdot]$ in (S99) is mean-zero and has bounded variance under the conditions of Theorem 4. In particular, we have

$$\begin{aligned}
\bar{\sigma}_{u^*}^2 &= \text{Var} \left[\frac{\bar{\xi}_1\{Y - \mu_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} k_{u^*}(u) e_u(X) du - \tau_{u^*} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}} \right. \\
& \left. + \frac{\mathcal{A}_2 - \tau_{u^*} \mathcal{A}_1}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}} \psi(X, Z, M) \right] \\
&= \mathbb{E} \left[\frac{\mathcal{A}_2 - \tau_{u^*} \mathcal{A}_1}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}} \psi(X, Z, M) + \frac{\bar{\xi}_1\{Y - \mu_{ZM}(X)\}}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}} \right. \\
& \left. + \frac{\int_{\mathbb{R}^2} \{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\} k_{u^*}(u) e_u(X) du - \tau_{u^*} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}} \right]^2.
\end{aligned}$$

In Theorem 3 (ii), we have derived the semiparametric EIF under the setting that the parametric model \mathcal{M}_{ps} in (9) is specified and $\hat{\beta}$ is estimated. By the classic semiparametric theory (see e.g. [Bickel et al. \(1993\)](#)), the corresponding semiparametric efficiency bound is

$$\begin{aligned}
\sigma_{u^*, \text{eff}}^2 &= \mathbb{E}\{\phi_{\text{p}}^2(X, Z, M, Y)\} \\
&= \mathbb{E} \left[\frac{\bar{\xi}_1\{Y - \mu_{ZM}(X)\} + \int_{\mathbb{R}^2} \{\mu_{1m_1}(X) - \mu_{0m_0}(X)\} k_{u^*}(u) e_u(X) du - \tau_{u^*} \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du}{\mathbb{E}\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \}} \right]^2
\end{aligned}$$

$$+ \frac{\mathcal{A}_2^* - \tau_{u^*} \mathcal{A}_1^*}{\mathbb{E} \left\{ \int_{\mathbb{R}^2} k_{u^*}(u) e_u(X) du \right\}} I_{\beta^*}^{-1} s_{ZM}(X | \beta^*) \Big]^2. \quad (\text{S101})$$

We now show that, under the setting of Theorem 5, the asymptotic variance of our proposed estimator given in (S5) achieves such bound and thus is semiparaetric efficient. Under the conditions of Theorem 5(III), we can check the following facts through definitions:

- (i) $\bar{\xi}_1 = \tilde{\xi}_1$ because $\bar{\mu} = \mu$, $\bar{\pi} = \pi$;
- (ii) $\mathcal{A}_1 = \mathcal{A}_1^*$, $\mathcal{A}_2 = \mathcal{A}_2^*$, noting that the third term of \mathcal{A}_2 in (S2) under our scenario becomes,

$$\begin{aligned} & \sum_{j=1,2,3} \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \{Y - \mu_{ZM}(X)\} \right] \delta_{X,Z,M}^{(j)} \right] \\ &= \sum_{j=1,2,3} \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\pi_Z(X)} \underbrace{\mathbb{E}\{Y - \mu_{ZM}(X) | X, Z, M\}}_{=0} \right] \delta_{X,Z,M}^{(j)} \right] = 0; \end{aligned}$$

- (iii) $\psi(X, Z, M) = I_{\beta^*}^{-1} s_{ZM}(X | \beta^*)$ by $\psi(x, z, m) = I_{\beta^*}^{-1} s_{zm}(x | \beta^*)$.

Based on above facts and comparing (S5) with (S101), we can observe that $\bar{\sigma}_{u^*}^2 = \sigma_{u^*,\text{eff}}^2$, which implies that $\hat{\tau}_{u^*}$ is semiparametric efficient. □

S5.7 Proof of Theorem 6

Combining (S97) and Proposition 1, we have,

$$\begin{aligned} \hat{\tau}_{u^*} - \tau_{u^*}^* &= (\hat{\tau}_{u^*} - \tau_{u^*}) + (\tau_{u^*} - \tau_{u^*}^*) \\ &= \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2} + h^{-1}r_{\pi}r_{\mu} + h^2). \end{aligned}$$

If $r_{\pi}r_{\mu} = \mathcal{O}(n^{-1/2})$, one further has $\hat{\tau}_{u^*} - \tau_{u^*}^* = \mathcal{O}_{\mathbb{P}}(h^{-1}n^{-1/2} + h^2)$. When choosing $h \asymp n^{-1/6}$, we then deduce the minimax optimal convergence rate $\hat{\tau}_{u^*} - \tau_{u^*}^* = \mathcal{O}_{\mathbb{P}}(n^{-1/3})$.

By (S96) and Proposition 1, when $h = o(n^{-1/6})$, $r_{\mu}r_{\pi} = o(n^{-1/2})$ and $\hat{\beta}$ is asymptotically linear, we further have,

$$\begin{aligned} & \sqrt{\bar{n}h}(\hat{\tau}_{u^*} - \tau_{u^*}^*) \\ &= \sqrt{\bar{n}h}(\hat{\tau}_{u^*} - \tau_{u^*}) + \sqrt{\bar{n}h}(\tau_{u^*} - \tau_{u^*}^*) \end{aligned}$$

$$= \sqrt{n}(\mathbb{P}_n - \mathbb{P}) \left[\frac{h\bar{\xi}_1\{Y - \mu_{ZM}(X)\}}{\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\}} + \frac{h\int_{\mathbb{R}^2}\{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du}{\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\}} - \frac{h\tau_{u^*}\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du}{\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\}} \right] \quad (\text{S102})$$

$$+ \frac{h\mathcal{A}_2 - h\tau_{u^*}\mathcal{A}_1}{\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\}} \psi(X, Z, M) \Big] + \underbrace{\mathcal{O}_{\mathbb{P}}(\sqrt{nr}\pi r\mu) + o_{\mathbb{P}}(1) + \mathcal{O}(\sqrt{nh^3})}_{o_{\mathbb{P}}(1)}. \quad (\text{S103})$$

To apply central limit theorem, under regularity conditions, we check the L_2 -norm bounds of all terms in (S102) and (S103). For example, for the first term in (S102), we have

$$\begin{aligned} \mathbb{E} \left[\frac{h\bar{\xi}_1\{Y - \mu_{ZM}(X)\}}{\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\}} \right]^2 &= \mathbb{E} \left[\frac{h^2\{\bar{\pi}_Z(X)f_{ZM}(X)\}^{-2}\{\gamma_1^{(Z)}(Z, M)\}^2\{Y - \mu_{ZM}(X)\}^2}{[\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\}]^2} \right] \\ &\lesssim \mathbb{E} \left[\{h\gamma_1^{(Z)}(Z, M)\}^2\{Y - \mu_{ZM}(X)\}^2 \right] \\ &\lesssim \mathbb{E} \left[\{Y - \mu_{ZM}(X)\}^2 \right] = \mathcal{O}(1), \end{aligned}$$

where the first inequality is by (S98), Assumption S8 and Assumption S13, and the second inequality is by $\sup_{z,m} h\gamma_1^{(z)}(z, m) = \mathcal{O}(1)$ under Assumption S9. For the second term in (S102), we have

$$\begin{aligned} \mathbb{E} \left[\frac{h\int_{\mathbb{R}^2}\{\bar{\mu}_{1m_1}(X) - \bar{\mu}_{0m_0}(X)\}k_{u^*}(u)e_u(X)du}{\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\}} \right]^2 &\lesssim \mathbb{E} \left[h\int_{\mathbb{R}^2} k_{u^*}(u)du \right]^2 \\ &= h^2 \rightarrow 0. \end{aligned}$$

Thus the second term in (S102) is negligible, i.e., it does not contribute to the asymptotic variance of $\sqrt{nh}(\hat{\tau}_{u^*} - \tau_{u^*}^*)$. We can similarly analyze all terms in (S102) and (S103) and show all terms have either bounded or vanishing L_2 norms; We omit details for simplicity. Finally by CLT, we can finally show that

$$\sqrt{nh}(\hat{\tau}_{u^*} - \tau_{u^*}^*) \rightsquigarrow \mathcal{N}(0, \tilde{\sigma}_{u^*}^2),$$

where

$$\begin{aligned} \tilde{\sigma}_{u^*}^2 &= \frac{\mathbb{E}[\mathcal{C}\psi(X, Z, M) + h\bar{\xi}_1\{Y - \mu_{ZM}(X)\}]^2}{[\mathbb{E}\{\int_{\mathbb{R}^2} k_{u^*}(u)e_u(X)du\}]^2}, \quad (\text{S104}) \\ \mathcal{C} &= \sum_{j=1,2,3} \mathbb{E} \left[\left[\frac{(-1)^{Z+1}}{\bar{\pi}_Z(X)} \{Y - \bar{\mu}_{ZM}(X)\} \right] h\delta_{X,Z,M}^{(j)} \right]. \end{aligned}$$

S6 Proof of Lemma S4

S6.1 Preliminary

We first introduce some basic concepts and results in the empirical process theory, e.g., the stochastic equicontinuity, covering number, entropy integral; See [Vaart and Wellner \(1996\)](#) for a more detailed introduction. Based on the application of empirical process theory, we prove Lemma S4. Denote $\mathbb{G}_n = \sqrt{n}(\mathbb{P}_n - \mathbb{P})$. Let $\mathcal{E}(g)$ be an operator of function g where g is in some metric space \mathcal{G} equipped with semimetric $d(\cdot, \cdot)$. In this paper, we typically consider the uniform difference metric, i.e., for any $g_1, g_2 \in \mathcal{G}$,

$$d(g_1, g_2) = \sup_{v \in \mathbb{V}} |g_1(v) - g_2(v)|.$$

We now introduce some notations to measure the complexity of the functional space \mathcal{G} . Let G be an envelope function for \mathcal{G} such that $G(v) \geq |g(v)|$ for any $g \in \mathcal{G}$ and $v \in \mathbb{V}$. Let $N(\epsilon, \mathcal{G}, \|\cdot\|)$ denote the covering number of \mathcal{G} , which is the minimal number of ϵ -balls needed to cover \mathcal{G} ; the ϵ -ball is under the distance induced by norm $\|\cdot\|$. We then define the uniform entropy integral,

$$J(\delta, \mathcal{G}, \mathcal{L}_2) = \sup_Q \int_0^\delta \sqrt{\log N\{\epsilon \|G\|_{Q,2}, \mathcal{G}, \|\cdot\|_{Q,2}\}} d\epsilon,$$

where Q can be any random measure, $\|\cdot\|_{Q,2}$ is the \mathcal{L}_2 -norm under Q , such that $\|g\|_{Q,2} = \sqrt{\int_{\mathbb{V}} g^2 dQ}$. A series of empirical process $\{\mathbb{G}_n \mathcal{E}(g) \mid n \geq 1, g \in \mathcal{G}\}$ is *stochastically equicontinuous* if for every $\epsilon, \zeta > 0$, there exists a $\delta > 0$ such that

$$\limsup_{n \rightarrow +\infty} \mathbb{P} \left\{ \sup_{\substack{d(g_1, g_2) < \delta \\ g_1, g_2 \in \mathcal{G}}} \left| \mathbb{G}_n \mathcal{E}(g_1) - \mathbb{G}_n \mathcal{E}(g_2) \right| > \epsilon \right\} < \zeta.$$

Above property implies that for any stochastically equicontinuous $\{\mathbb{G}_n \mathcal{E}(g) \mid n \geq 1, g \in \mathcal{G}\}$,

$$\mathbb{G}_n \{\mathcal{E}(g_1) - \mathcal{E}(g_2)\} = o_{\mathbb{P}}(1) \tag{S105}$$

whenever $d(g_1, g_2) = o_{\mathbb{P}}(1)$ and $g_1, g_2 \in \mathcal{G}$ ([Pollard, 2012](#)).

We also denote $\|\mathcal{E}\|_{\mathcal{G}} = \sup_{g \in \mathcal{G}} |\mathcal{E}\{g(V)\}|$. It can be seen that $\|\mathcal{E}\|_{\mathcal{G}}$ is generally a random variable as it depends on V . To show a series of empirical process $\{\mathbb{G}_n \mathcal{E}(g) \mid n \geq 1, g \in \mathcal{G}\}$ is stochastically equicontinuous, one can use Theorem 2.11.1 in [Vaart and Wellner \(1996\)](#), which boils down the

verification of stochastic equicontinuity, to checking the following two Lindeberg conditions and one condition of \mathcal{G} 's complexity,

- (1). $\mathbb{E}\{\|\mathcal{E}\|_{\mathcal{G}}^2 \cdot 1(\|\mathcal{E}\|_{\mathcal{G}} > \epsilon\sqrt{n})\} \rightarrow 0$ for every $\epsilon > 0$,
- (2). $\sup_{d(g_1, g_2) < \delta_n} \mathbb{E}[\{\mathcal{E}(g_1) - \mathcal{E}(g_2)\}^2] \rightarrow 0$ for every sequence $\delta_n \rightarrow 0$,
- (3). $\int_0^{\delta_n} \sqrt{\log N(\epsilon, \mathcal{G}, \|\cdot\|_{\mathbb{P}_n, 2})} d\epsilon \rightarrow 0$ in probability, for every sequence $\delta_n \rightarrow 0$.

Now we prove Lemma S4. Using the new notation, we decompose our target term to bound, namely $(\mathbb{P}_n - \mathbb{P})\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\}$, as

$$\begin{aligned} (\mathbb{P}_n - \mathbb{P})\{h(V | \hat{\pi}, \hat{\mu}) - h(V | \bar{\pi}, \bar{\mu})\} &= h^{-1}n^{-1/2}\mathbb{G}_n\{h_1(V | \hat{\pi}, \hat{\mu}) - h_1(V | \bar{\pi}, \bar{\mu})\} \\ &\quad + n^{-1/2}\mathbb{G}_n\{h_2(V | \hat{\mu}) - h_2(V | \bar{\mu})\} \\ &\quad + n^{-1/2}\mathbb{G}_n\{h_3(V | \hat{\mu}) - h_3(V | \bar{\mu})\}, \end{aligned} \tag{S106}$$

where we define

$$\begin{aligned} h_1(V | \pi, \mu) &= \frac{(-1)^{Z+1}h\gamma_1^{(Z)}(X, M)}{\pi_Z(X)f_{ZM}(X)}\{Y - \mu_{ZM}(X)\}, \\ h_2(V | \mu) &= \int_{\mathbb{R}^2} k_{u^*}(u)\mu_{1m_1}(X)e_u(X)du, \\ h_3(V | \mu) &= -\int_{\mathbb{R}^2} k_{u^*}(u)\mu_{0m_0}(X)e_u(X)du. \end{aligned}$$

In the following sections, we show the root- n -rates of all three terms on the right-hand side of (S106), and thereby show the desired result (S91).

S6.2 Bounding $\mathbb{G}_n\{h_1(V | \hat{\pi}, \hat{\mu}) - h_1(V | \bar{\pi}, \bar{\mu})\}$

We use a similar proof strategy as Kennedy et al. (2017, Appendix 5) to bound the first term of (S106). Denote

$$\mathcal{E}_1(g) = \frac{(-1)^{Z+1}h\gamma_1^{(Z)}(X, M)}{f_{ZM}(X)}g(V).$$

The following lemma gives a sufficient condition, under which $\{\mathbb{G}_n\mathcal{E}_1(g) | n \geq 1\}$ is stochastically equicontinuous.

Lemma S5 Suppose Assumptions 1–4 and other regularity conditions in Section S5.5 hold. Suppose g belongs to \mathcal{G} , such that \mathcal{G} has:

- (i). A finite uniform entropy integral, i.e., $J(\delta, \mathcal{G}, \mathcal{L}_2) < +\infty$ for every $\delta > 0$;
- (ii). A uniformly bounded envelope function $G(v)$ such that $\sup_{v \in \mathbb{V}} G(v) < C_G$ with some universal constant $C_G > 0$, and $G(v) > |g(v)|$ for any $g \in \mathcal{G}$ and any $v \in \mathbb{V}$.

Then $\{\mathbb{G}_n \mathcal{E}_1(g) \mid n \geq 1\}$ is stochastically equicontinuous.

Proof of Lemma S5. To show $\{\mathbb{G}_n \mathcal{E}_1(g) \mid n \geq 1\}$ is stochastically equicontinuous, we only need to verify aforementioned Conditions (1)–(3) in Section S6.1. We first verify Lindeberg Condition (1). We have for any $c > 0$,

$$\begin{aligned} \lim_{n \rightarrow +\infty} \text{pr} \left\{ \|\mathcal{E}_1\|_{\mathcal{G}}^2 \cdot \mathbf{1}(\|\mathcal{E}_1\|_{\mathcal{G}} > \epsilon\sqrt{n}) \geq c \right\} &\leq \lim_{n \rightarrow +\infty} \text{pr} \left(\|\mathcal{E}_1\|_{\mathcal{G}} > c\sqrt{n} \right) \\ &\leq \lim_{n \rightarrow +\infty} \text{pr} \left\{ \frac{h\gamma_1^{(Z)}(X, M)}{f_{ZM}(X)} G(V) > c\sqrt{n} \right\} \\ &= 0, \end{aligned} \tag{S107}$$

where the last equality is for some universal constant $C_{\mathcal{E}_1} > 0$, we have a.s.,

$$\left| \frac{h\gamma_1^{(Z)}(X, M)}{f_{ZM}(X)} G(V) \right| < C_{\mathcal{E}_1},$$

where $G(V)$ is bounded by C_G , $f_{ZM}(X)$ is bounded away from 0 under Assumption S13, and when $Z = 0$,

$$\begin{aligned} h\gamma_1^{(Z)}(X, M) &= h \int_{\mathbb{R}} k_{u^*}(m_1, M) e_{(m_1, M)}(X) dm_1 \\ &\lesssim h \int_{\mathbb{R}} k_{u^*}(m_1, M) dm_1 \quad (\text{Assumptions S10 and S13}) \\ &\leq \sup_{m_0 \in \mathbb{R}, h > 0} h \int_{\mathbb{R}} k_{u^*}(m_1, m_0) dm_1 \\ &\lesssim 1 \quad (\text{Assumption S9}), \end{aligned}$$

similar for $Z = 1$ and thus $h\gamma_1^{(Z)}(X, M)$ is uniformly bounded by some constant. By (S107),

$$\|\mathcal{E}_1\|_{\mathcal{G}}^2 \cdot \mathbf{1}(\|\mathcal{E}_1\|_{\mathcal{G}} > \epsilon\sqrt{n}) = o_{\mathbb{P}}(1). \tag{S108}$$

On the other hand, under the regularity conditions,

$$\begin{aligned}
\mathbb{E}\|\mathcal{E}_1\|_{\mathcal{G}}^2 &\leq \mathbb{E}\left\{\frac{h\gamma_1^{(Z)}(X, M)}{f_{ZM}(X)}G(V)\right\}^2 \\
&\leq C_G^2 c_0^{-2} h^2 \mathbb{E}\left\{\gamma_1^{(Z)}(X, M)\right\}^2 \\
&= \sum_{z=0,1} C_G^2 c_0^{-2} h^2 \mathbb{E}\left[\pi_z(X) \cdot \mathbb{E}\left[\left\{\gamma_1^{(z)}(X, M)\right\}^2 \mid X, Z = z\right]\right] \\
&\lesssim 1,
\end{aligned} \tag{S109}$$

where the last inequality follows because we have when $z = 0$,

$$\begin{aligned}
\mathbb{E}\left[\pi_z(X) \cdot \mathbb{E}\left[\left\{\gamma_1^{(z)}(X, M)\right\}^2 \mid X, Z = z\right]\right] &\leq \epsilon \cdot \mathbb{E}\left[\mathbb{E}\left[\left\{\gamma_1^{(0)}(X, M)\right\}^2 \mid X, Z = 0\right]\right] \\
&\leq \epsilon \cdot \mathbb{E}\left[\mathbb{E}\left\{\int_{\mathbb{R}} k_{u^*}^2(m_1, M) e_{(m_1, M)}^2(X) dm_1 \mid X, Z = 0\right\}\right] \\
&\lesssim \mathbb{E}\left[\mathbb{E}\left\{\int_{\mathbb{R}} k_{u^*}^2(m_1, M) dm_1 \mid X, Z = 0\right\}\right] \\
&= \mathbb{E}\left[\int_{\mathbb{R}^2} k_{u^*}^2(u) f_{0m_0}(X) du\right] \\
&\lesssim h^{-2},
\end{aligned} \tag{S110}$$

where the third inequality is because that $e_u(x)$ is uniformly bounded under regularity conditions and the last inequality is by $h^2 \int_{\mathbb{R}^2} k_{u^*}^2(u) du \lesssim 1$ and $f_{0m_0}(x)$ is uniformly upper bounded; Similar result also holds for $z = 1$. Combining (S108) and (S110), and noting $\|\mathcal{E}_1\|_{\mathcal{G}}^2 \cdot 1(\|\mathcal{E}_1\|_{\mathcal{G}} > \epsilon\sqrt{n}) \leq \|\mathcal{E}_1\|_{\mathcal{G}}^2$, we can invoke dominated convergence theorem and show Lindeberg Condition (1) holds for $\mathcal{E}_1(g)$.

We next verify Lindeberg Condition (2) when $d(g_1, g_2) = \|g_1 - g_2\|_{\infty}$, where $\|\cdot\|_{\infty}$ is the sup-norm. We have,

$$\begin{aligned}
\sup_{d(g_1, g_2) < \delta_n} \mathbb{E}\left[\left\{\mathcal{E}_1(g_1) - \mathcal{E}_1(g_2)\right\}^2\right] &= \mathbb{E}\left[\frac{\{h\gamma_1^{(Z)}(X, M)\}^2}{f_{ZM}^2(X)} \left\{g_1(V) - g_2(V)\right\}^2\right] \\
&\leq c_0^{-2} h^2 \|g_1 - g_2\|_{\infty}^2 \mathbb{E}\left[\left\{\gamma_1^{(Z)}(X, M)\right\}^2\right] \\
&\lesssim \delta_n \rightarrow 0,
\end{aligned}$$

where the last inequality follows from $\|g_1 - g_2\|_{\infty} \leq \delta_n$ and $\mathbb{E}\left[\left\{\gamma_1^{(Z)}(X, M)\right\}^2\right] \lesssim h^{-2}$ as shown in

(S109).

We finally verify Condition (3). By [Vaart and Wellner \(1996, § 2.11.1.1 & Lemma 2.11.6\)](#), Condition (3) is satisfied if \mathcal{G} has a finite uniform entropy integral and for some random measure ν_n ,

$$\frac{1}{n} \left\{ \mathcal{E}_1(g_1) - \mathcal{E}_1(g_2) \right\}^2 \leq \int_{\mathbb{V}} \{g_1(v) - g_2(v)\}^2 d\nu_n; \quad (\text{S111})$$

see also the proof of Lemma 1 in [Kennedy et al. \(2017\)](#). Noticing that

$$\frac{1}{n} \left\{ \mathcal{E}_1(g_1) - \mathcal{E}_1(g_2) \right\}^2 = \frac{\{h\gamma_1^{(Z)}(X, M)\}^2}{nf_{ZM}^2(X)} \left\{ g_1(V) - g_2(V) \right\}^2,$$

(S111) is satisfied by the following random measure $\nu_n = \frac{\{h\gamma_1^{(Z)}(X, M)\}^2}{nf_{ZM}^2(X)} \delta_V$, where δ_V is a (random) Dirac measure centered at V . We thus have verified Condition (3). With Conditions (1)–(3) being satisfied, we have $\{\mathbb{G}_n \mathcal{E}_1(g) \mid n \geq 1\}$ is stochastically equicontinuous. □

Define

$$g^{\natural}(V \mid \pi, \mu) = \frac{Y - \mu_{ZM}(X)}{\pi_Z(X)}.$$

We then have

$$\mathbb{G}_n \{h_1(V \mid \hat{\pi}, \hat{\mu}) - h_1(V \mid \bar{\pi}, \bar{\mu})\} = \mathbb{G}_n \left[\mathcal{E}_1 \{g^{\natural}(V \mid \hat{\pi}, \hat{\mu})\} - \mathcal{E}_1 \{g^{\natural}(V \mid \bar{\pi}, \bar{\mu})\} \right]. \quad (\text{S112})$$

For the series of empirical process $\{\mathbb{G}_n \mathcal{E}_1(g^{\natural}) \mid n \geq 1, g^{\natural} \in \mathcal{G}^{\natural}\}$ where g^{\natural} can be all possible $g^{\natural}(\cdot \mid \hat{\pi}, \hat{\mu})$ and $g^{\natural}(\cdot \mid \bar{\pi}, \bar{\mu})$ under Assumption 4. We first prove that there exists some metric space \mathcal{G}^{\natural} satisfying the condition in Lemma S5, and thus $\{\mathbb{G}_n \mathcal{E}_1(g^{\natural}) \mid n \geq 1, g^{\natural} \in \mathcal{G}^{\natural}\}$ is stochastically equicontinuous. Under Assumption S13, we have functional classes \mathcal{G}_{μ} and \mathcal{G}_{π} containing $\bar{\mu}, \hat{\mu}$ and $\bar{\pi}, \hat{\pi}$, respectively, and both \mathcal{G}_{μ} and \mathcal{G}_{π} have uniformly bounded envelop functions and finite uniform entropy integrals. We then construct

$$\mathcal{G}^{\natural} = (\mathcal{Y} \ominus \mathcal{G}_{\mu}) \circ \mathcal{G}_{\pi}^{-1},$$

where we denote \mathcal{Y} is the set of a single function outputting Y from Z , and for two functional sets, e.g., \mathcal{G} and \mathcal{G}' , we define the operations, $\mathcal{G} \ominus \mathcal{G}' = \{g - g' \mid g \in \mathcal{G}, g' \in \mathcal{G}'\}$, $\mathcal{G}^{-1} = \{g^{-1} \mid g \in \mathcal{G}\}$,

and $\mathcal{G} \circ \mathcal{G}' = \{gg' \mid g \in \mathcal{G}, g' \in \mathcal{G}'\}$. Under Assumptions S12 and S13 such that $\mu_{zm}(x)$ and $\pi_z(x)^{-1}$ are uniformly bounded by some constant and \mathbb{Y} is compact, it can be seen that \mathcal{G}^{\natural} has an envelope function with a uniform constant upper bound; thus Condition (ii) in Lemma S5 is verified. The Condition (i) in Lemma S5 follows from the basic property of uniform entropy integral. Under Assumption S13, both \mathcal{G}_μ and \mathcal{G}_π^{-1} have bounded entropy integrals and bounded envelop functions and so does the single-function space \mathcal{Y} . Therefore, by Andrews (1994, Theorem 3), one has the uniform entropy integral of \mathcal{G}^{\natural} is bounded. Thus Lemma S5 verifies $\{\mathbb{G}_n \mathcal{E}_1(g^{\natural}) \mid n \geq 1, g^{\natural} \in \mathcal{G}^{\natural}\}$ is stochastically equicontinuous.

On the other hand, recalling we define $d(\cdot, \cdot)$ as the uniform difference metric, we have

$$\begin{aligned}
d\left\{g^{\natural}(\cdot \mid \hat{\pi}, \hat{\mu}), g^{\natural}(\cdot \mid \bar{\pi}, \bar{\mu})\right\} &= \sup_{v \in \mathbb{V}} \left| \frac{y - \hat{\mu}_{zm}(x)}{\hat{\pi}_z(x)} - \frac{y - \bar{\mu}_{zm}(x)}{\bar{\pi}_z(x)} \right| \\
&\leq \sup_{v \in \mathbb{V}} \left| \frac{y}{\hat{\pi}_z(x)\bar{\pi}_z(x)} \{\bar{\pi}_z(x) - \hat{\pi}_z(x)\} \right| + \sup_{v \in \mathbb{V}} \left| \frac{1}{\hat{\pi}_z(x)} \{\hat{\mu}_{zm}(x) - \bar{\mu}_{zm}(x)\} \right| \\
&\quad + \sup_{v \in \mathbb{V}} \left| \frac{\bar{\mu}_{zm}(x)}{\hat{\pi}_z(x)\bar{\pi}_z(x)} \{\bar{\pi}_z(x) - \hat{\pi}_z(x)\} \right| \\
&= o_{\mathbb{P}}(\|\hat{\pi} - \bar{\pi}\|_{\infty} + \|\hat{\mu} - \bar{\mu}\|_{\infty}) \\
&= o_{\mathbb{P}}(1),
\end{aligned} \tag{S113}$$

under Assumptions S12 and S13. Finally, by (S105), (S112) and (S113), we conclude $h^{-1}n^{-1/2}\mathbb{G}_n\{h_1(V \mid \hat{\pi}, \hat{\mu}) - h_1(V \mid \bar{\pi}, \bar{\mu})\} = o_{\mathbb{P}}(h^{-1}n^{-1/2})$.

S6.3 Bounding $\mathbb{G}_n\{h_j(V \mid \hat{\mu}) - h_j(V \mid \bar{\mu})\}$ for $j = 2, 3$

When bounding $\mathbb{G}_n\{h_2(V \mid \hat{\mu}) - h_2(V \mid \bar{\mu})\}$ and $\mathbb{G}_n\{h_3(V \mid \hat{\mu}) - h_3(V \mid \bar{\mu})\}$, the arguments will be similar. Therefore, we focus on bounding the former for brevity. Define,

$$\mathcal{E}_2(\mu) = \int_{\mathbb{R}^2} k_{u^*}(u) \mu_{1m_1}(X) e_u(X) du.$$

We then write

$$n^{-1/2}\mathbb{G}_n\{h_2(V \mid \hat{\mu}) - h_2(V \mid \bar{\mu})\} = n^{-1/2}\mathbb{G}_n\{\mathcal{E}_2(\hat{\mu}) - \mathcal{E}_2(\bar{\mu})\}. \tag{S114}$$

By Van der Vaart (2000, Theorem 19.14) and Assumption 4, we have the functional space \mathcal{G}_{μ_1} which contains all possible functions $\hat{\mu}_{1m_1}(x), \bar{\mu}_{1m_1}(x)$, is P -Donsker, because \mathcal{G}_{μ_1} has a finite uniform entropy integral when $\delta = 1$ and a uniformly bounded (and thus also mean-squared-average bounded)

envelop function; See the definition of P -Donsker in [Van der Vaart \(2000, § 19.2\)](#). Next, we consider the functional space,

$$\mathcal{E}_2(\mathcal{G}_{\mu_1}) = \left\{ \mathcal{E}_2(\mu)(x) = \int_{\mathbb{R}^2} k_{u^*}(u) \mu_{1m_1}(x) e_u(x) du \mid \mu_{1m_1}(x) \in \mathcal{G}_{\mu_1} \right\},$$

which contains functions for all possible $\mathcal{E}_2(\hat{\mu})$ and $\mathcal{E}_2(\bar{\mu})$.

Now we show that $\mathcal{E}_2(\mathcal{G}_{\mu_1})$ is P -Donsker. Since $e_u(x)$ is a fixed function and uniformly bounded (Assumptions [S10](#) and [S13](#)), we have $\mathcal{G}_{\mu_1 e} = \{\mu_{1m_1}(x) e_u(x) \mid \mu_{1m_1}(x) \in \mathcal{G}_{\mu_1}\}$ is also P -Donsker by [Van der Vaart \(2000, Theorem 19.14\)](#). Define $R(u)$ as a fixed probability measure over u with probability density function $k_{u^*}(u) / \int_{\mathbb{R}^2} k_{u^*}(u) du$ and clearly $C_{\mathcal{E}} = \int_{\mathbb{R}^2} k_{u^*}(u) du < +\infty$ is a fixed constant under our regularity conditions. Then noticing that

$$\int_{\mathbb{R}^2} k_{u^*}(u) \mu_{1m_1}(x) e_u(x) du = C_{\mathcal{E}} \int_{\mathbb{R}^2} \mu_{1m_1}(x) e_u(x) R(du),$$

we can apply [Van der Vaart and van der Laan \(2006, Lemma 5.2\)](#) with $r = s = 2$ and deduce,

$$\sup_Q N\{\epsilon \|G_{\mathcal{E}}\|_{Q,2}, \mathcal{E}_2(\mathcal{G}_{\mu_1}), \|\cdot\|_{Q,2}\} \leq \sup_Q N\{\epsilon \|G_{\mu_1 e}\|_{Q,2}/2, \mathcal{G}_{\mu_1 e}, \|\cdot\|_{Q,2}\}, \quad (\text{S115})$$

where $G_{\mu_1 e}(x, u)$ can be any envelope function of $\mathcal{G}_{\mu_1 e}$ and, correspondingly, define

$$G_{\mathcal{E}}(x) = C_{\mathcal{E}} \left\{ \int_{\mathbb{R}^2} G_{\mu_1 e}^2(x, u) dR(du) \right\}^{1/2}, \quad (\text{S116})$$

which serves as an envelope function of $\mathcal{E}_2(\mathcal{G}_{\mu_1})$ by Hölder's inequality.

On the other hand, noticing that for any $\nu(u, x) = \mu_{1m_1}(x) e_u(x)$, $\nu'(v) = \mu'_{1m_1}(x) e_u(x)$ for any $\mu, \mu' \in \mathcal{G}_{\mu_1}$, we have

$$\begin{aligned} \|\nu - \nu'\|_{Q,2} &\leq \left\{ \sup_{x,u} e_u(x) \right\} \cdot \|\mu_{1\cdot}(\cdot) - \mu'_{1\cdot}(\cdot)\|_{Q,2} \\ &\leq C_e \|\mu_{1\cdot}(\cdot) - \mu'_{1\cdot}(\cdot)\|_{Q,2} \end{aligned}$$

for some fixed constant $C_e > 0$ such that

$$\sup_{x,u} e_u(x) \leq C_e, \quad (\text{S117})$$

as $e_u(x)$ is uniformly bounded. Thus for any $(C_e^{-1}c)$ -balls covering \mathcal{G}_{μ_1} with functional centers,

$\mu_{1m_1}^{(1)}(x), \mu_{1m_1}^{(2)}(x)$, etc, the corresponding c -balls with functional centers, $e_u(x)\mu_{1m_1}^{(1)}(x), e_u(x)\mu_{1m_1}^{(2)}(x)$, etc, cover $\mathcal{G}_{\mu_1 e}$ for any $c > 0$. Thus we have

$$N\{c, \mathcal{G}_{\mu_1 e}, \|\cdot\|_{Q,2}\} \leq N\{C_e^{-1}c, \mathcal{G}_{\mu_1}, \|\cdot\|_{Q,2}\}, \quad (\text{S118})$$

for any $c > 0$ and any random measure Q . Let $G_{\mu_1}(x, m_1)$ be an envelope function of \mathcal{G}_{μ_1} satisfying Assumption 4. By (S117), we can take $G_{\mu_1 e}(x, u) = C_e G_{\mu_1}(x, m_1)$ as the envelop function of $\mathcal{G}_{\mu_1 e}$. Taking $c = \epsilon \|G_{\mu_1 e}\|_{Q,2}/2 = \epsilon C_e \|G_{\mu_1}\|_{Q,2}/2$ in (S118) and considering the supremum over all possible Q , we have

$$\sup_Q N\{\epsilon \|G_{\mu_1 e}\|_{Q,2}/2, \mathcal{G}_{\mu_1 e}, \|\cdot\|_{Q,2}\} \leq \sup_Q N\{\epsilon \|G_{\mu_1}\|_{Q,2}/2, \mathcal{G}_{\mu_1}, \|\cdot\|_{Q,2}\}. \quad (\text{S119})$$

Combing (S115) and (S119), we have

$$\sup_Q N\{\epsilon \|G_{\mathcal{E}}\|_{Q,2}, \mathcal{E}_2(\mathcal{G}_{\mu_1}), \|\cdot\|_{Q,2}\} \leq \sup_Q N\{\epsilon \|G_{\mu_1}\|_{Q,2}/2, \mathcal{G}_{\mu_1}, \|\cdot\|_{Q,2}\}$$

and therefore,

$$\begin{aligned} J\{1, \mathcal{E}_2(\mathcal{G}_{\mu_1}), \mathcal{L}_2\} &= \sup_Q \int_0^1 \sqrt{\log N\{\epsilon \|G_{\mathcal{E}}\|_{Q,2}, \mathcal{E}_2(\mathcal{G}_{\mu_1}), \|\cdot\|_{Q,2}\}} d\epsilon \\ &\leq \sup_Q \int_0^1 \sqrt{\log N\{\epsilon \|G_{\mu_1}\|_{Q,2}/2, \mathcal{G}_{\mu_1}, \|\cdot\|_{Q,2}\}} d\epsilon \\ &\leq 2 \sup_Q \int_0^{1/2} \sqrt{\log N\{\epsilon \|G_{\mu_1}\|_{Q,2}, \mathcal{G}_{\mu_1}, \|\cdot\|_{Q,2}\}} d\epsilon \\ &< +\infty, \end{aligned} \quad (\text{S120})$$

under Assumption 4. In addition, we have

$$\begin{aligned} \mathbb{E}\{G_{\mathcal{E}}^2(X)\} &= \mathbb{E}\left\{C_{\mathcal{E}}^2 C_e^2 \int_{\mathbb{R}^2} G_{\mu_1}^2(m_1, X) dR(du)\right\} \\ &\leq C_{\mathcal{E}}^2 C_e^2 \left\{\sup_{m_1, x} G_{\mu_1}(m_1, x)\right\}^2 \\ &< +\infty, \end{aligned} \quad (\text{S121})$$

as $G_{\mu_1}(m_1, x)$ is uniformly bounded. Summarizing (S120)–(S121) and by Van der Vaart (2000, Theorem 19.14), we conclude that $\mathcal{E}_2(\mathcal{G}_{\mu_1})$ is P -Donsker.

With the Donsker property, a standard application of the continuous mapping theorem (Van der

Vaart, 2000, Theorem 19.24) implies that

$$\mathbb{G}_n\{\mathcal{E}_2(\hat{\mu}) - \mathcal{E}_2(\bar{\mu})\} = o_{\mathbb{P}}(1) \quad (\text{S122})$$

for any $\mathcal{E}_2(\hat{\mu})$ and $\mathcal{E}_2(\bar{\mu})$ in $\mathcal{E}_2(\mathcal{G}_{\mu_1})$, as long as $\|\mathcal{E}_2(\hat{\mu}) - \mathcal{E}_2(\bar{\mu})\|_{\infty} = o_{\mathbb{P}}(1)$; we omit this standard argument for simplicity and refer the interested reader to Kennedy (2016, § 4.2) for details. Thus (S122) gives the desired bound for (S114). Therefore, to finish the proof, it is only left to show $\|\mathcal{E}_2(\hat{\mu}) - \mathcal{E}_2(\bar{\mu})\|_{\infty} = o_{\mathbb{P}}(1)$ which can be bounded as follows under Assumption 4,

$$\begin{aligned} \|\mathcal{E}_2(\hat{\mu}) - \mathcal{E}_2(\bar{\mu})\|_{\infty} &= \sup_{x \in \mathbb{X}} \left| \int_{\mathbb{R}^2} k_{u^*}(u) \{\hat{\mu}_{1m_1}(x) - \bar{\mu}_{1m_1}(x)\} e_u(x) du \right| \\ &\leq \|\hat{\mu}_{1m_1} - \bar{\mu}_{1m_1}\|_{\infty} C_e C_{\mathcal{E}} \\ &= o_{\mathbb{P}}(1). \end{aligned}$$

Summarizing above results, we have $n^{-1/2}\mathbb{G}_n\{h_2(V | \hat{\mu}) - h_2(V | \bar{\mu})\} = o_{\mathbb{P}}(n^{-1/2})$, and symmetrically we can show $n^{-1/2}\mathbb{G}_n\{h_3(V | \hat{\mu}) - h_3(V | \bar{\mu})\} = o_{\mathbb{P}}(n^{-1/2})$. Combining the bounds in Sections S6.2 and S6.3 along with (S106), Lemma S4 follows. \square

S7 Additional Figures for simulation in Section 6.1

In Figure S5, we report the complete simulation results for our numerical experiments in Section 6.1. In Figures S5(a)–(c), we present the histograms of $\hat{\tau}_u - \tau_u^*$ over 100 Monte Carlo iterations under different data generation settings, with $n = 500, 800, \text{ and } 1000$, respectively. In each subfigure, the top and bottom panels present the results when nuisance functions are estimated through parametric and semiparametric strategies, respectively. X-axis represents different data generation mechanisms. For example, (TP2,PS1,OM1) means data are generated from $(\mathcal{F}_{\text{tp}}, \mathcal{F}_{\text{ps}}, \mathcal{F}_{\text{ps}}) = (\mathcal{F}_{\text{tp}}^{(2)}, \mathcal{F}_{\text{ps}}^{(1)}, \mathcal{F}_{\text{ps}}^{(1)})$. Different colors represent different principal levels u .

Under the data generation settings when $\hat{\tau}_u$ can consistently estimate τ_u^* as indicated by Theorem 6, i.e., the data generation settings are (TP1,PS1,OM1), (TP2,PS1,OM1), (TP1,PS1, OM2) when nuisance functions are trained by the parametric strategy, and the data generation settings are (TP1,PS1,OM1), (TP2,PS1,OM1), (TP1,PS1,OM2), (TP2,PS1,OM2) when nuisance functions are trained by the semiparametric strategy, we can observe the corresponding estimation biases of $\hat{\tau}_u - \tau_u^*$ are close to zero for all n , and in general, the estimation variances and biases become

smaller as n increases, which verifies the nonparametric consistency of $\hat{\tau}_u$ (cf. Theorem 6). In addition, when all nuisance functions can be consistently estimated, i.e., the data generation setting is (TP1,PS1,OM1) with parametric nuisance training and the data generation settings are (TP1,PS1,OM1), (TP2,PS1,OM1), (TP1,PS1,OM2), (TP2,PS1,OM2) with semiparametric nuisance training, the estimation errors of $\hat{\tau}_u$ become particularly small due to the rate-double robustness (cf. Theorem 6).

S8 Additional Figures and results for the real-data applications in Section 7

This section contains additional figures and results for our real-data applications in Section 7. In particular, Figures S6–S8 contain additional results for the surrogate analysis of ACTG 175 data in Section 7.1. Figures S6–S8 contain additional results for the causal pathway discovery of Bangladesh flooding data in Section 7.1.

Figure S6(a) shows the Q-Q plot of linear regression residuals when training \mathcal{M}_{ps} in the ACTG 175 data analysis. The approximate linear pattern implies that our regular linear regression-type modeling of $M \mid X, Z$ is reasonable. Figure S6(b) is the histogram of all standardized M_i , which roughly range from -2 to 3 . This implies that our selected bandwidths $h = 0.25$ and 0.5 are small, and Loc.PCE surface can approximate Cont.PCE surface reasonably well. In Figures S7 and S8, we report the $\hat{\tau}_u$ surface as well as the 95% Bootstrap confidence intervals for $\hat{\tau}_u$ with 100 resamples, varying ρ and h with different values. Our proposed estimators with different ρ and h show similar patterns as Figure 3, and imply similar conclusions as we discussed in Section 7.1. Therefore, additional results in Figure S7 and S8 show that our proposed estimator is not sensitive with different ρ and h for ACTG 175 data analysis, and further confirm our conclusions in Section 7.1.

Similar to Figure S6, Figure S9 show that our linear model for \mathcal{M}_{ps} and the selection of h are reasonable for Bangladesh flooding data analysis. The additional results with different ρ and h in Figures S10 and S11 show that our proposed estimator is not sensitive to the variability of ρ and h , and therefore further validate our conclusions in Section 7.2.






Figure S5: The box-plots of $\hat{\tau}_u - \tau_u^*$ over 100 MC rounds when $n = 500, 800$ and 1000 . X-axis represents different data generation mechanisms. The top and bottom panels in each subfigure present the results when nuisance functions are estimated through parametric and semiparametric strategies, respectively.




(a) The Q-Q plot of the residuals for linear regression $M \sim X, Z$.




(b) The histogram of standardized M_i among all samples.


Figure S6: Model diagnosis for ACTG 175 data in Section 7.1



(a) $\rho = 0.5, h = 0.25$




(b) $\rho = 0, h = 0.25$




(c) $\rho = -0.5, h = 0.25$


Figure S7: Additional results for ACTG 175 data with $h = 0.25$ and $\rho = 0.5, 0, -0.5$. For each subfigure, the left panel shows the surface of $\hat{\tau}_u$ with specific h and ρ , when $u \in [190 \text{ cells/mm}^3, 520 \text{ cells/mm}^3]^2$. The X-axis of right panel is different levels of u . The top right panel shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying u when $m_1 = m_0$. The bottom right panel shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying m_1 , while m_0 is fixed at 340 cells/mm^3 .



(a) $\rho = 0.5, h = 0.5$




(b) $\rho = 0, h = 0.5$




(c) $\rho = -0.5, h = 0.5$

Figure S8: Additional results for ACTG 175 data with $h = 0.5$ and $\rho = 0.5, 0, -0.5$. For each subfigure, the left panel shows the surface of $\hat{\tau}_u$ with specific h and ρ , when $u \in [190 \text{ cells/mm}^3, 520 \text{ cells/mm}^3]^2$. The X-axis of right panel is different levels of u . The top right panel shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying u when $m_1 = m_0$. The bottom right panel shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying m_1 , while m_0 is fixed at 340 cells/mm^3 .




(a) The Q-Q plot of the residuals for linear regression $M \sim X, Z$.




(b) The histogram of standardized M_i among all samples.


Figure S9: Model diagnosis for Bangladesh flooding data in Section 7.2



(a) $\rho = 0.5, h = 0.25$




(b) $\rho = 0, h = 0.25$




(c) $\rho = -0.5, h = 0.25$


Figure S10: Additional results for Bangladesh flooding data with $h = 0.25$ and $\rho = 0.5, 0, -0.5$. For each subfigure, the left panel shows the surface of $\hat{\tau}_u$ with specific h and ρ , when $u \in [1400 \text{ calories}, 2500 \text{ calories}]^2$. The X-axis of the right panel is different levels of u . The top right panel shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying u when $m_1 = m_0$. The bottom right panel shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying m_1 , while m_0 is fixed at 1900 calories.



(a) $\rho = 0.5, h = 0.5$



(b) $\rho = 0, h = 0.5$



(c) $\rho = -0.5, h = 0.5$

Figure S11: Additional results for Bangladesh flooding data with $h = 0.5$ and $\rho = 0.5, 0, -0.5$. For each subfigure, the left panel shows the surface of $\hat{\tau}_u$ with specific h and ρ , when $u \in [1400 \text{ calories}, 2500 \text{ calories}]^2$. The X-axis of the right panel is different levels of u . The top right panel shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying u when $m_1 = m_0$. The bottom right panel shows the 95% bootstrap CIs of $\hat{\tau}_u$ with varying m_1 , while m_0 is fixed at 1900 calories.