## CONIC BUNDLE THREEFOLDS DIFFERING BY A CONSTANT BRAUER CLASS AND CONNECTIONS TO RATIONALITY

SARAH FREI, LENA JI, SOUMYA SANKAR, BIANCA VIRAY, AND ISABEL VOGT

ABSTRACT. A double cover Y of  $\mathbb{P}^1 \times \mathbb{P}^2$  ramified over a general (2,2)-divisor will have the structure of a geometrically standard conic bundle ramified over a smooth plane quartic  $\Delta \subset \mathbb{P}^2$  via the second projection. These threefolds are rational over algebraically closed fields, but over nonclosed fields, including over  $\mathbb{R}$ , their rationality is an open problem. In this paper, we characterize rationality over  $\mathbb{R}$  when  $\Delta(\mathbb{R})$  has at least two connected components (extending work of M. Ji and the second author) and over local fields when all odd degree fibers of the first projection have nonsquare discriminant.

We obtain these applications by proving general results comparing the conic bundle structure on Y with the conic bundle structure on a well-chosen intersection of two quadrics. The difference between these two conic bundles is encoded by a constant Brauer class, and we prove that this class measures a certain failure of Galois descent for the codimension 2 Chow group of Y.

### 1. INTRODUCTION

For threefolds over algebraically closed fields, much is known about the rationality problem (i.e., the property of being birational to projective space), see e.g. [IP99]. However, the classical rationality obstructions cannot always detect irrationality over nonclosed fields. In this direction, Hassett–Tschinkel and Benoist–Wittenberg introduced a rationality obstruction over nonclosed fields k that considers restrictions on certain torsors under the intermediate Jacobian [HT21b, BW23, HT21a], refining the classical Clemens–Griffiths intermediate Jacobian obstruction [CG72]. Subsequent works show that this refined intermediate Jacobian torsor (IJT) obstruction is strong enough to characterize rationality for many classes of Fano threefolds with k-points [HT21b, BW23, KP23].

In an earlier paper [FJS<sup>+</sup>24], the authors studied the IJT obstruction for geometrically standard conic bundle threefolds over  $\mathbb{P}^2$  with singular fibers over a smooth plane quartic  $\Delta$ , and proved that although the IJT obstruction characterizes rationality for these degree 4 conic bundles over fields k with  $\operatorname{Br} k[2] = 0$ , the IJT obstruction fails to characterize rationality over any subfield of  $\mathbb{R}$ . The root cause of this discrepancy is that the intermediate Jacobian torsors of a conic bundle  $\pi: X \to \mathbb{P}^2$  depend only on the discriminant cover  $\tilde{\Delta} \to \Delta$ of  $\pi$ , whereas the isomorphism class of the generic fiber  $X_{\eta}$  depends on the discriminant cover and the constant 2-torsion Brauer class given by a smooth fiber at any point  $w \in (\mathbb{P}^2 \setminus \Delta)(k)$ .

Thus, one may ask whether the vanishing of the IJT obstruction for a degree 4 conic bundle  $X \to \mathbb{P}^2$  implies the rationality of *some* degree 4 conic bundle  $X' \to \mathbb{P}^2$  with the

<sup>2020</sup> Mathematics Subject Classification. Primary: 14C25. Secondary: 14E08, 14G27, 14H40, 14K30.

Key words and phrases. Conic bundles, the Brauer group, rationality, intermediate Jacobians.

same discriminant cover  $\hat{\Delta}/\Delta$  as X. We prove that this indeed occurs when the so-called Prym curve<sup>1</sup>  $\Gamma_{\tilde{\Delta}/\Delta}$  has a k-point (see Corollary 3.2).

The question then becomes: what geometric information is captured by the constant Brauer class  $[X_w] - [X'_w]$ ? The primary goal of this paper is to answer this question for an important class of Fano threefolds with a degree 4 conic bundle structure. Let  $\Delta \subset \mathbb{P}^2$  be a smooth plane quartic and let  $\varpi : \tilde{\Delta} \to \Delta$  be a smooth projective geometrically integral étale double cover. Then there exists a unique double cover  $Y = Y_{\tilde{\Delta}/\Delta} \to \mathbb{P}^1 \times \mathbb{P}^2$  branched over a (2, 2)-divisor such that the second projection has the structure of a (geometrically standard) conic bundle with discriminant cover  $\tilde{\Delta} \to \Delta$ . The first projection  $\pi_1$  realizes  $Y_{\tilde{\Delta}/\Delta}$  as a quadric surface fibration; in particular,  $Y_{\tilde{\Delta}/\Delta}$  is rational as soon as  $\pi_1$  has a section.

**Theorem 1.1.** Over a field k of characteristic different from 2, let  $\tilde{\Delta}/\Delta$ ,  $Y_{\tilde{\Delta}/\Delta}$ , and  $\Gamma_{\tilde{\Delta}/\Delta}$ be as above and set  $X := Y_{\tilde{\Delta}/\Delta}$ . Assume that  $\Gamma_{\tilde{\Delta}/\Delta}(k) \neq \emptyset$  and that the IJT obstruction vanishes. Then there is a conic bundle  $\psi \colon X' \to \mathbb{P}^2$  with discriminant cover  $\tilde{\Delta}/\Delta$ , such that

- (1) The threefold X' is rational, and
- (2)  $[X_{\eta}] [X'_{\eta}] \in \operatorname{Br} k[2]$  is trivial if and only if  $\pi_1$  has a section.

Theorem 1.1 raises the question of whether conic bundles  $X \to \mathbb{P}^2$ ,  $X' \to \mathbb{P}^2$  that have the same discriminant cover, but whose generic fibers are not isomorphic, can *both* be rational. In other words, can  $\mathbb{P}^3$  have more than one isomorphism class of conic bundle structure with the same discriminant cover? If not, then Theorem 1.1 would imply that  $Y_{\tilde{\Delta}/\Delta} \to \mathbb{P}^1 \times \mathbb{P}^2$ is rational if and only if  $\pi_1$  has a section. In general, this question seems difficult to answer, but we provide two contexts in which there is a positive answer. First, in Proposition 2.1, we show that when  $\tilde{\Delta}$  has index 1, any two conic bundles with discriminant cover  $\tilde{\Delta}/\Delta$ necessarily have isomorphic generic fibers. When  $\Delta$  is a smooth plane quartic, to the best of our knowledge the condition that  $\tilde{\Delta}$  has index 1 is the unique sufficient rationality criteria that depends *only* on the discriminant cover (see [FJS<sup>+</sup>24, Section 8.1] and Corollary 2.2).

On the other hand, over  $\mathbb{R}$ , topological obstructions can shed light on whether two conic bundles with the same discriminant cover can both be rational. Exploiting this extra information, we extend the characterization of rationality of real degree 4 conic bundles in [JJ24] to cover one of the two remaining isotopy classes of the quartic  $\Delta(\mathbb{R})$ .

**Theorem 1.2.** Let  $k = \mathbb{R}$  and let  $\Delta/\Delta$  be a geometrically integral étale double cover of a smooth plane quartic. Assume that  $\Delta(\mathbb{R})$  is not a single oval. Then  $Y_{\bar{\Delta}/\Delta}$  is rational if and only if  $\pi_1$  has a section.

For further equivalent rationality conditions over  $\mathbb{R}$ , see Theorem 7.2 (which includes Theorem 1.2 as a special case).

The topological arguments over  $\mathbb{R}$  do not extend to nonarchimedean local fields. However, we can use our methods to show that if the Prym curve  $\Gamma_{\tilde{\Delta}/\Delta}$  has index 2, then the IJT obstruction characterizes rationality.

**Theorem 1.3** (Corollary 5.4). Let k be a local field of characteristic different from 2 and let  $\tilde{\Delta}/\Delta$  be a geometrically integral étale double cover of a smooth plane quartic. If  $\Gamma_{\tilde{\Delta}/\Delta}$  has index 2, then  $Y_{\tilde{\lambda}/\Delta}$  is rational if and only if the IJT obstruction vanishes.

<sup>&</sup>lt;sup>1</sup>The intermediate Jacobian of X is the Prym variety of  $\tilde{\Delta}/\Delta$ , which, in the case deg  $\Delta = 4$ , is isomorphic to the Jacobian of an explicit genus 2 curve  $\Gamma_{\tilde{\Delta}/\Delta}$  with equation given in (2.4) [FJS<sup>+</sup>24, Definition 4.6].

1.1. **Outline.** We begin in Section 2 with a brief review of conic bundles, focusing on the discriminant cover  $\varpi: \tilde{\Delta} \to \Delta$  of a conic bundle and the possible isomorphism classes of the generic fiber. In this section we prove Proposition 2.1: if  $\tilde{\Delta}$  has index 1, then any two conic bundles with discriminant cover  $\tilde{\Delta}/\Delta$  necessarily have isomorphic generic fibers. In Section 2.2 we review results from [FJS<sup>+</sup>24] that rephrase the IJT obstruction for conic bundles whose discriminant cover is a geometrically irreducible étale double cover of a smooth plane quartic in terms of torsors under the Prym variety of  $\tilde{\Delta} \to \Delta$ ; in doing so, we also give equations for the Prym curve  $\Gamma_{\tilde{\Delta}/\Delta}$ . In Section 3, we show that under the assumption that the Prym curve has a rational point, there exists a conic bundle with discriminant cover  $\tilde{\Delta} \to \Delta$  that is the blow-up along a smooth conic of an intersection of two quadrics. In Section 4, we construct a conic bundle over one of the Prym torsors from Section 2.2.1 and show that if the torsor is trivial, then this conic bundle is constant. This construction is used in Section 5 to study the obstruction of a section to  $\pi_1: Y_{\tilde{\Delta}/\Delta} \to \mathbb{P}^1$ . These ingredients are combined in Section 6 to prove Theorem 1.1. We end in Section 7 by specializing to the case  $k = \mathbb{R}$  and proving Theorem 1.2.

### Acknowledgements

The authors thank David Saltman for an intriguing question and helpful follow-up discussion that motivated this project, and Asher Auel for helpful conversations.

This material is based partially upon work supported by National Science Foundation grant DMS-1928930 while the last two authors were in residence at the Simons Laufer Mathematical Sciences Institute in Berkeley, California, during the Spring 2023 semester. S.F. was supported in part by the Hausdorff Research Institute for Mathematics, funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2047/1 – 390685813. L.J. was supported in part by NSF MSPRF grant DMS-2202444. S.S. was supported by the Dutch Research Council (NWO) grant OCENW.XL21.XL21.011. B.V. was supported in part by an AMS Birman Fellowship and NSF grant DMS-2101434. I.V. was supported in part by NSF grant DMS-2200655. For the purpose of open access, a CC BY public copyright license is applied to any Author Accepted Manuscript version arising from this submission.

**Notation.** Throughout, k denotes a field of characteristic different from 2. A variety over k is a separated k-scheme of finite type. If Y is an integral k-variety,  $\mathbf{k}(Y)$  denotes its function field. More generally, if Y is a finite union of integral k-varieties  $Y_i$ , then  $\mathbf{k}(Y) := \prod_i \mathbf{k}(Y_i)$  is the ring of global sections of the sheaf of total quotient rings.

For a smooth variety Y over a field k, we write Br Y for the (cohomological) Brauer group  $\operatorname{H}^2_{\operatorname{et}}(Y, \mathbb{G}_m)$ . If  $Y = \operatorname{Spec} R$  is affine we write Br  $R := \operatorname{Br} \operatorname{Spec} R$ . For a field k, Br k can also be viewed as describing the isomorphism classes of Severi–Brauer varieties. In particular, any rank 3 quadric defines an isomorphism class of a smooth plane conic which has an associated class in the 2-torsion part of the Brauer group. Given a rank 2 quadric Q, we may associate the Brauer class given by the conic  $V(z^2 - Q) \subset \mathbb{P}^2$ . For a quadratic form  $Q \in k[\mathbf{x}]$ , the symmetric matrix M associated to Q is defined to be the matrix such that  $\mathbf{x}M\mathbf{x}^T = Q$ .

The Brauer group  $\mathrm{H}^2_{\mathrm{\acute{e}t}}(-, \mathbb{G}_m)$  is a contravariant functor, and given a morphism of smooth varieties  $f: Y \to Z$  we write  $\alpha_Y := f^* \alpha$  for  $\alpha \in \mathrm{Br} Z$ . Given any point  $z \in Z$  we write  $Y_z$  for the fiber of f over z.

For a curve C, i.e., a smooth projective variety of dimension 1, we write  $\operatorname{Pic}_{C}$  for the Picard scheme of C and write  $\operatorname{Pic}_{C}^{i}$  for the connected component that parametrizes degree i line bundles.

Throughout the paper, we will write  $\eta$  for the generic point of  $\mathbb{P}^2$ .

### 2. Conic bundles and discriminant covers

The central objects of interest in this paper are **conic bundles** over  $\mathbb{P}^2$ , i.e., the class of smooth projective varieties X with a flat morphism  $\pi: X \to \mathbb{P}^2$  of relative dimension 1 such that  $\omega_X$  is relatively anti-ample. This implies that locally over  $\mathbb{P}^2$ , X is defined by a quadratic form and that the fiber of  $\pi$  over any  $w \in \mathbb{P}^2$  is a conic in  $\mathbb{P}^2$  (possibly of rank less than 3 if w is not the generic point).

There is a curve  $\Delta \subset \mathbb{P}^2$  with the property that the fiber over  $w \in \mathbb{P}^2$  is a smooth conic if and only if  $w \notin \Delta$ . Furthermore, there is a curve  $\tilde{\Delta}$  that is a double cover of  $\Delta$  and that parametrizes the irreducible components of the fibers of  $\pi: X_{\Delta} \to \Delta$ . The curve  $\tilde{\Delta}$  can be obtained by considering the Stein factorization of the relative variety  $\mathcal{F}_1(X/\mathbb{P}^2) \to \Delta \subset \mathbb{P}^2$  of lines in the fibers of  $\pi$ . We call  $\Delta$  the discriminant locus of  $\pi$  and  $\varpi: \tilde{\Delta} \to \Delta$  the discriminant cover of  $X \to \mathbb{P}^2$ . If  $\Delta$  is smooth, then  $\varpi$  is étale.

A conic bundle is standard if  $\operatorname{Pic} X = \pi^* \operatorname{Pic} \mathbb{P}^2 \oplus \mathbb{Z}$ ; this is equivalent to  $\tilde{\Delta}$  and  $\Delta$  having the same number of irreducible components. Throughout the paper, we reserve  $\pi \colon X \to \mathbb{P}^2$ for a geometrically standard conic bundle, i.e., the base change  $X_{\overline{k}} \to \mathbb{P}^2_{\overline{k}}$  remains standard.

2.1. Discriminant covers and isomorphism classes of conic bundles. Discriminant covers are crucial to the study of conic bundles and control much of their geometry. Indeed, over fields with trivial 2-torsion part of the Brauer group, the discriminant cover determines the isomorphism class of  $X_{\eta}$ . Over arbitrary fields, we study the isomorphism class of  $X_{\eta}$  via its class in Br  $\mathbf{k}(\mathbb{P}^2)[2]$  by considering the purity exact sequence [CTS21, Theorems 3.7.3 and 6.1.3]

$$0 \longrightarrow \operatorname{Br} k[2] \longrightarrow \operatorname{Br} \mathbf{k}(\mathbb{P}^2)[2] \xrightarrow{(\partial_z)_z} \bigoplus_{z \in (\mathbb{P}^2)^{(1)}} H^1(\mathbf{k}(z), \mathbb{Z}/2\mathbb{Z}).$$

By definition of the residue maps  $\partial_z$  [CTS21, Definition 1.4.11],  $\partial_z([X_\eta])$  is trivial for every z outside of the support of  $\Delta$ . Furthermore, if  $\mathbf{k}(\tilde{\Delta})_z$  denotes the maximal subfield of  $\mathbf{k}(\tilde{\Delta})$  that is an extension of  $\mathbf{k}(z)$ , then  $\partial_z([X_\eta]) = \mathbf{k}(\tilde{\Delta})_z$  for every  $z \in (\mathbb{P}^2)^{(1)}$  in the support of  $\Delta$ . Thus, if  $\pi: X \to \mathbb{P}^2$  and  $\pi': X' \to \mathbb{P}^2$  are two standard conic bundles with the same discriminant cover, then


$$[X_{\eta}] - [X'_{\eta}] \in \operatorname{im}(\operatorname{Br} k[2] \to \operatorname{Br} \mathbf{k}(\mathbb{P}^2)[2]).$$
(2.1)

Further, this difference agrees with the difference  $[X_w] - [X'_w]$  for any  $(w \in \mathbb{P}^2 \setminus \Delta)(k)$ . However, as the following proposition shows, not every element of Br k[2] can be realized as a difference of two standard conic bundles with a fixed discriminant cover.

**Proposition 2.1.** Let  $\varpi: \Delta \to \Delta$  be an étale double cover of smooth, projective, geometrically integral curves with  $\Delta$  a plane curve over a field k of characteristic different from 2. If  $\tilde{\Delta}$  has index 1, i.e., if  $\tilde{\Delta}(L) \neq \emptyset$  for some odd degree extension L/k, then any conic bundles  $\pi: X \to \mathbb{P}^2, \ \tilde{\pi}: \ \tilde{X} \to \mathbb{P}^2$  with discriminant cover  $\tilde{\Delta}/\Delta$  must have isomorphic generic fibers. That is,  $[X_{\eta}] - [\ \tilde{X}_{\eta}] = 0 \in \operatorname{Br} \mathbf{k}(\mathbb{P}^2)$ . Proof. Let  $\pi: X \to \mathbb{P}^2$ ,  $\tilde{\pi}: \tilde{X} \to \mathbb{P}^2$  be standard conic bundles with discriminant cover  $\tilde{\Delta}/\Delta$ . Then  $[X_\eta] - [\tilde{X}_\eta] \in \operatorname{im}(\operatorname{Br} k[2] \to \operatorname{Br} \mathbf{k}(\mathbb{P}^2)[2])$  by (2.1). We will show that for any nontrivial  $\alpha \in \operatorname{Br} k[2], [X_\eta] + \alpha$  has no quadratic splitting field, which implies that  $[X_\eta] + \alpha$  is not represented by a conic. Thus,  $[X_\eta] - [\tilde{X}_\eta]$  must equal  $0 \in \operatorname{Br} \mathbf{k}(\mathbb{P}^2)$ , as desired.

The statement that  $[X_{\eta}] + \alpha$  has no quadratic splitting field is equivalent to the statement that the index of  $[X_{\eta}] + \alpha$  is  $2^n$  for some n > 1. Since the index of a 2-primary Brauer class is unchanged under odd degree extensions [GS17, Corollary 4.5.11], we may reduce to the case that  $\tilde{\Delta}(k) \neq \emptyset$ .

We now return to proving that  $[X_{\eta}] + \alpha$  has no quadratic splitting field, assuming that  $\tilde{\Delta}(k) \neq \emptyset$ . By a result of Albert [Lam05, Chapter III, Theorem 4.8],  $[X_{\eta}] + \alpha$  has a quadratic splitting field if and only if  $\alpha$  and  $[X_{\eta}]$  have a common quadratic splitting field  $L/\mathbf{k}(\mathbb{P}^2)$ . Assume that there exists such an L. Let R denote the local ring  $\mathcal{O}_{\mathbb{P}^2,\Delta}$  and identify  $\mathbf{k}(\mathbb{P}^2) = \operatorname{Frac}(R)$ ; let S denote the integral closure of R in L. Since L splits  $X_{\eta}$  and  $\pi \colon X \to \mathbb{P}^2$  is proper, the valuative criterion for properness gives a morphism  $\operatorname{Spec} S \to X_R$ . If  $R \hookrightarrow S$  is unramified, then the image of  $\operatorname{Spec} S \to X_R$  cannot meet the singular locus of the special fiber of  $X_R \to \operatorname{Spec} R$ . In particular, since  $X_{\mathbf{k}(\Delta)}$  is birational to  $\mathbb{P}^1_{\mathbf{k}(\tilde{\Delta})}$  (by definition of the discriminant cover), if  $R \hookrightarrow S$  is an unramified extension, then the extension of residue fields agrees with  $\mathbf{k}(\tilde{\Delta})/\mathbf{k}(\Delta)$ . Thus, we have the following commutative diagrams:



By assumption L splits  $\alpha$ , so the diagram on the right shows that  $\alpha$  becomes trivial when pulled back to  $\mathbf{k}(\tilde{\Delta})$ . The pullback map factors as  $\operatorname{Br} k \to \operatorname{Br} \tilde{\Delta} \hookrightarrow \operatorname{Br} \mathbf{k}(\tilde{\Delta})$ . Since  $\tilde{\Delta}(k) \neq \emptyset$ ,  $\operatorname{Br} k \to \operatorname{Br} \tilde{\Delta}$  is injective [CTS21, Proposition 5.4.2 and Remark 5.4.3(1)], so  $\alpha = 0 \in \operatorname{Br} k$ .

In the case that  $R \hookrightarrow S$  is ramified, then the residue field of S is  $\mathbf{k}(\Delta)$ . Since  $\tilde{\Delta}$  is a cover of  $\Delta$ , we have  $\Delta(k) \neq \emptyset$  and so we can repeat the above argument with  $\Delta$  replacing  $\tilde{\Delta}$ .  $\Box$ 

In the case that  $\Delta$  has degree 4, we prove the following strengthening of [FJS<sup>+</sup>24, Proposition 8.1(ii)].

**Corollary 2.2.** In the setting of Proposition 2.1, assume  $\Delta$  is a smooth plane quartic and  $\tilde{\Delta}$  has index 1. Then any conic bundle  $X \to \mathbb{P}^2$  with discriminant cover  $\tilde{\Delta}/\Delta$  is rational.

Proof. By  $[FJS^+24$ , Section 6] there exists a threefold Y with the structure of a conic bundle with discriminant cover  $\tilde{\Delta}/\Delta$  and the structure of a quadric fibration  $\pi_1: Y \to \mathbb{P}^1$ .  $[FJS^+24$ , Proposition 6.1(v)] and the assumption on  $\tilde{\Delta}$  imply that there is an odd degree extension L/k over which  $(\pi_1)_L: Y_L \to \mathbb{P}^1_L$  has a section. Then  $\pi_1$  has a section over k by Springer's theorem [Spr52], so Y is rational. Next, let  $X \to \mathbb{P}^2$  be any conic bundle with discriminant cover  $\tilde{\Delta}/\Delta$ . Then Proposition 2.1 implies that X and Y are birationally equivalent, and therefore X is rational. 2.2. Prym varieties and the IJT obstruction to rationality. The primary conic bundles  $\pi: X \to \mathbb{P}^2$  of interest in this paper are those whose discriminant curve  $\Delta \subset \mathbb{P}^2$  is a smooth plane curve (often of degree 4) and whose discriminant cover  $\varpi: \tilde{\Delta} \to \Delta$  is geometrically integral. The smoothness of X and  $\Delta$  imply that  $\varpi$  is an étale double cover.

The Prym variety of  $\Delta/\Delta$ , which we denote by P, is the connected component of the identity of ker( $\varpi_*$ :  $\operatorname{Pic}_{\tilde{\Delta}/k} \to \operatorname{Pic}_{\Delta/k}$ ); it is a principally polarized abelian variety (see [Mum74]). The Prym variety is the intermediate Jacobian of X, and can be thought of as a parameter space for algebraically trivial curve classes on X [Bea77, Proposition 3.3, FJS<sup>+</sup>24, Theorem 5.8]. The kernel of  $\varpi_*$ :  $\operatorname{Pic}_{\tilde{\Delta}/k} \to \operatorname{Pic}_{\Delta/k}$  has a unique additional connected component, which we denote  $\tilde{P}$ , and it is a torsor under P.

2.2.1. Étale double covers of smooth plane quartics. If  $\Delta \subset \mathbb{P}^2$  is a smooth plane quartic, then any geometrically integral étale double cover  $\tilde{\Delta}$  is a smooth genus 5 curve. Bruin [Bru08] proved that the Prym variety of a such a double cover is always isomorphic to the Jacobian of a smooth genus 2 curve  $\Gamma = \Gamma_{\tilde{\Delta}/\Delta}$ . This implies that on the threefold X, the Clemens–Griffiths intermediate Jacobian obstruction to rationality [CG72] vanishes. Defining equations of  $\Delta, \tilde{\Delta}, \Gamma$  can be given in terms of 3 quadrics  $Q_1, Q_2, Q_3 \in k[u, v, w]$  as follows:

$$\Delta = V(Q_2^2 - Q_1 Q_3), \tag{2.2}$$

$$\tilde{\Delta} = V(Q_1 - r^2, Q_2 - rs, Q_3 - s^2) \subset \mathbb{P}^4, \text{and}$$
(2.3)

$$\Gamma = V(y^2 - \det(t_0^2 M_1 + 2t_0 t_1 M_2 + t_1^2 M_3)) \subset \mathbb{P}(1, 1, 3), \tag{2.4}$$

(here  $M_i$  is the symmetric matrix associated to  $Q_i$ ); see, for example, [Bru08]. In addition,  $Q_1, Q_2, Q_3$  are uniquely determined up to a PGL<sub>2</sub>-action [FJS<sup>+</sup>24, Theorem 4.5(i)] which agrees with changing coordinates of  $t_0, t_1$  on  $\Gamma$ .

The intermediate Jacobian torsor (IJT) obstruction for X, introduced by [HT21b] over  $\mathbb{R}$  and [BW23] over arbitrary k (see also [HT21a] for  $k \subset \mathbb{C}$ ), is an obstruction to rationality of X that is encoded by certain torsors under P. In the case where  $\Delta \subset \mathbb{P}^2$  is degree 4, the obstruction is encoded by  $\tilde{P}$  and the following two P-torsors:

$$P^{(1)} := \{ D \in \mathbf{Pic}_{\tilde{\Delta}/k}^4 \colon \varpi_* D \text{ lies on a line in } \mathbb{P}^2, \mathbf{h}^0(D) \equiv 0 \mod 2 \}, \text{ and}$$
(2.5)  
$$\tilde{P}^{(1)} := \{ D \in \mathbf{Pic}_{\tilde{\Delta}/k}^4 \colon \varpi_* D \text{ lies on a line in } \mathbb{P}^2, \mathbf{h}^0(D) = 1 \}.$$
(2.6)

**Proposition 2.3.** Let k be a field of characteristic different from 2, and let  $\pi: X \to \mathbb{P}^2$  be a conic bundle over k with discriminant cover  $\tilde{\Delta} \to \Delta$  such that  $\Delta$  is a smooth quartic and  $\tilde{\Delta}$  is geometrically integral.

- (1) [Bru08, Section 5, Case 4] As P-torsors,  $P^{(1)} \simeq \operatorname{Pic}_{\Gamma}^{1}$ .
- (2)  $[FJS^+24, Corollary of Theorem 6.4, Section 6.4]$  The IJT obstruction vanishes for X if and only if either  $\tilde{P}(k) \neq \emptyset$  or  $\tilde{P}^{(1)}(k) \neq \emptyset$ . Moreover, if  $\operatorname{Pic}_{\Gamma}^{1}(k) \neq \emptyset$ , then  $\tilde{P}^{(1)}(k) \neq \emptyset$  if and only if  $\tilde{P}(k) \neq \emptyset$ , so the vanishing of the IJT obstruction for X is equivalent to  $\tilde{P}^{(1)}(k) \neq \emptyset$ .

*Remark* 2.4. The torsors  $\tilde{P}, P^{(1)}, \tilde{P}^{(1)}$  are part of a larger family of torsors, given by the polarized Prym scheme, which can be defined for arbitrary conic bundles; see [FJS<sup>+</sup>24, Section 4] for more details.

2.3. Relation to curve classes. The intermediate Jacobian torsors are indexed by Galoisinvariant algebraic curve classes  $\gamma$  on X. For conic bundles, the vanishing of the IJT obstruction implies the existence of certain Galois-invariant geometric curve classes up to *rational* equivalence, as we now explain. Let  $\operatorname{CH}^2(X)$  denote the group of rational equivalence classes of codimension 2 cycles on X, and for a fixed class  $\gamma$ , let  $\operatorname{CH}^2(X)^{\gamma}$  denote the classes that are algebraically equivalent to  $\gamma$ . In the degree 4 case,  $\tilde{P}^{(1)}(k) \cong (\operatorname{CH}^2(X_{\bar{k}})^{\tilde{\gamma}_1})^{G_k}$  for some algebraic curve class  $\tilde{\gamma}_1$ . Proposition 2.3 shows that if the IJT obstruction vanishes and  $\operatorname{Pic}^1_{\Gamma_{\tilde{\Delta}/\Delta}}(k) \neq \emptyset$ , then there is a cycle on  $X_{\bar{k}}$  algebraically equivalent to  $\tilde{\gamma}_1$  whose rational equivalence class is Galois invariant. (See [FJS<sup>+</sup>24, Theorem 6.4] for a construction of  $\tilde{\gamma}_1$ .)

## 3. Constructing an intersection of quadrics with a conic bundle structure

The goal of this section is to prove the following proposition.

## **Proposition 3.1.** Let k be a field of characteristic different from 2.

Given quadrics  $Q_1, Q_2, Q_3 \in k[u, v, w]$  such that  $V(Q_2^2 - Q_1Q_3)$  is a smooth quartic and such that  $\operatorname{disc}(Q_1)$  is a square, there exist two quadrics  $q_0, q_\infty \in k[x_0, x_1, \ldots, x_5]$  such that

- (1) The intersection  $Z := V(q_0, q_\infty) \subset \mathbb{P}^5$  is a smooth projective threefold;
- (2) The linear section  $Z \cap V(x_3, x_4, x_5)$  is a smooth conic C; and
- (3) Projection away from the plane  $V(x_3, x_4, x_5)$  induces a morphism  $\operatorname{Bl}_C Z \to \mathbb{P}^2$ , endowing  $\operatorname{Bl}_C Z$  with a geometrically standard conic bundle structure whose generic fiber equals

 $[(\operatorname{Bl}_C Z)_{\eta}] = (Q_1, Q_2^2 - Q_1 Q_3) + [Q_1] \in \operatorname{Br} \mathbf{k}(\mathbb{P}^2),$ 

where  $[Q_1]$  denotes the pullback of the class in Br k[2] corresponding to the rank 2 or 3 quadric  $Q_1$ . In particular, the discriminant cover of the conic bundle  $\operatorname{Bl}_C Z \to \mathbb{P}^2$  is  $\tilde{\Delta}/\Delta$ .

Using this proposition together with work of Hassett–Tschinkel and Benoist–Wittenberg, we can deduce the assertion from the introduction.

**Corollary 3.2** (Corollary of Proposition 3.1 & [HT21b, Theorem 36, BW23, Theorem A]). Let k be a field of characteristic different from 2, and let  $\pi: X \to \mathbb{P}^2$  be a conic bundle over k with discriminant cover  $\varpi: \tilde{\Delta} \to \Delta$  where  $\Delta \subset \mathbb{P}^2$  is a smooth plane quartic and  $\tilde{\Delta}$  is geometrically irreducible. Assume that  $\Gamma_{\tilde{\Delta}/\Delta}(k) \neq \emptyset$ .

There exists a conic bundle  $\pi': X' \to \mathbb{P}^2$  with discriminant cover  $\tilde{\Delta}/\Delta$  (namely  $X' = \operatorname{Bl}_C Z$  with Z, C as in Proposition 3.1) such that X' is rational if and only if the IJT obstruction vanishes for X (equivalently for X').

Proof. The assumption that  $\Gamma_{\tilde{\Delta}/\Delta}(k) \neq \emptyset$  means that there exists some  $[t_0:t_1] \in \mathbb{P}^1(k)$  such that the discriminant of  $t_0^2 Q_1 + 2t_0 t_1 Q_2 + t_1^2 Q_3$  is a square. By a change of coordinates on  $\mathbb{P}^1(k)$ , we may assume that this point is [1:0] so  $\operatorname{disc}(Q_1)$  is a square. Then we may apply Proposition 3.1 to obtain an intersection of quadrics Z containing a conic C such that the conic bundle morphism  $\pi' \colon \operatorname{Bl}_C Z \to \mathbb{P}^2$  has discriminant cover  $\varpi \colon \tilde{\Delta} \to \Delta$ . Then [HT21b, Theorem 36, BW23, Theorem A] imply that the IJT obstruction characterizes rationality for  $\operatorname{Bl}_C Z$  (see [FJS<sup>+</sup>24, Corollary 8.3] for details).

Remark 3.3. If k is algebraically closed, then the proposition is suggested by a dimension count of the relevant moduli spaces. Over nonclosed fields k, we must show that the intersection of quadrics Z has the same field of definition as the quadrics  $Q_1, Q_2, Q_3$ . We do so by constructing the quadrics  $q_0, q_\infty$  (see (3.1) and (3.2)) and then we verify that Z has the desired properties.

Proof of Proposition 3.1. The assumption that  $V(Q_2^2 - Q_1Q_3)$  is a smooth quartic implies that the quadrics  $Q_1, Q_3$  both have rank at least 2. This furthermore implies that the curve  $\tilde{\Delta}$  defined in (2.3) is geometrically integral. We consider two cases for the rank of  $Q_1$ .

**Case 1**: disc $(Q_1) \neq 0$ , i.e.,  $Q_1$  has rank 3. We change coordinates on  $\mathbb{P}^2$  to assume that  $Q_1 = au^2 + bv^2 - abw^2$  for some  $a, b \in k^{\times}$ . Note that then  $[Q_1] = (a, b) \in \text{Br } k[2]$ . Consider the following  $6 \times 6$  symmetric matrices defined in terms of the symmetric matrices  $M_i$  associated to  $Q_i$ :

$$A_0 := \begin{pmatrix} abM_1^{-1} & -M_1^{-1}M_2 \\ -M_2M_1^{-1} & \frac{-1}{ab}(M_3 - M_2M_1^{-1}M_2) \end{pmatrix}, \qquad A_\infty := \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}.$$
(3.1)

Let  $q_0, q_\infty$  be the two quadratic forms defined by these matrices, i.e.,  $q_i = \mathbf{x} A_i \mathbf{x}^T$ . We claim that  $Z := V(q_0, q_\infty) \subset \mathbb{P}^5$  has the desired properties.

An intersection of two quadrics in six variables is smooth and three-dimensional if and only if the discriminant polynomial  $\det(A_0 - TA_\infty)$  is separable [Rei72, Proposition 2.1]. Using the Schur complement, we compute that up to a nonzero scalar,  $\det(A_0 - TA_\infty)$  is equal to the determinant of the following  $3 \times 3$  matrix

$$M_3 - M_2 M_1^{-1} M_2 + (M_2 M_1^{-1} + TI) M_1 (M_1^{-1} M_2 + TI) = M_3 + 2TM_2 + T^2 M_1.$$

Thus, by (2.4), the discriminant polynomial  $det(A_0 - TA_\infty)$  is a scalar multiple of the polynomial describing the Weierstrass points of  $\Gamma$ , which is separable. Hence, Z is smooth and three-dimensional, as desired.

Since  $q_{\infty}$  is identically zero when restricted to  $V(x_3, x_4, x_5)$ , the linear section of Z given by  $V(x_3, x_4, x_5)$  is equal to  $V(q_0, x_3, x_4, x_5)$  so by (3.1) is given by the vanishing of the rank 3 form corresponding to  $M_1^{-1}$ . Thus we have proved (2).

To prove (3), we must further describe the conic bundle structure on  $\operatorname{Bl}_C Z$ . Let  $\Lambda := V(x_3, x_4, x_5)$ , and let  $\phi: Z \dashrightarrow \mathbb{P}^2$  be the projection away from  $C := V(q_0) \cap \Lambda$ . Fix a point  $p = [u : v : w] \in \mathbb{P}^2$ . We obtain the fiber of  $\phi$  above p by taking the intersection  $\operatorname{Span}(\Lambda, [0:0:0:u:v:w]) \cap Z \subset \mathbb{P}^5$ , removing  $\Lambda$ , and then taking the Zariski closure.

Note that the restriction of  $q_{\infty}$  to  $\text{Span}(\Lambda, [0:0:v:w])$  is a rank 2 quadratic form, and one can compute that the vanishing of this quadratic form is the union of  $\Lambda$  and the 3-dimensional column span of the following matrix

$$B_p = \begin{pmatrix} -v & -w & 0 & 0\\ u & 0 & -w & 0\\ 0 & u & v & 0\\ & \mathbf{0}_{3,3} & & [u, v, w]^T \end{pmatrix};$$

we denote the column span of this matrix  $\Lambda_p$ . Thus the fiber of  $\phi$  above p is  $\Lambda_p \cap V(q_0)$ , which is given by the rank 3 quadratic form associated to the  $4 \times 4$  symmetric matrix  $B_p^T A_0 B_p$ . Let **M** denote the bottom left  $3 \times 3$  submatrix of  $B_p^T A_0 B_p$ . One can check<sup>2</sup> that the determinants of the top right  $i \times i$  submatrices of **M** are:

$$-(u^2 - bw^2), \quad -w^2Q_1, \quad \frac{-w^2}{ab}(Q_2^2 - Q_1Q_3), \quad \text{for } i = 1, 2, 3, \text{ respectively.}$$

<sup>&</sup>lt;sup>2</sup>This, and all following computational claims, can be verified with the Magma [BCP97] code in the Github repository. This submatrix  $\mathbf{M}$  is denoted 'bM' in the code.

 $(bv)^2 - (-ab(u^2 - bw^2))$ , we see that  $(-Q_1, Q_1)$  is trivial and that

$$(-ab(u^2 - bw^2), Q_1) = (-ab(u^2 - bw^2), b) = (-ab, b) = (a, b),$$

which completes the proof.

**Case 2**:  $Q_1$  has rank 2. The proof will proceed similarly as in the rank 3 case with the specific equations modified slightly, so we will omit more of the algebraic details. As in the previous case, verification of all steps can be found in the Magma [BCP97] code in the Github repository associated to this paper.

We change coordinates on  $\mathbb{P}^2$  to assume that  $Q_1 = av^2 + bw^2$  for some  $a, b \in k^{\times}$ . Note that then  $[Q_1] = (a, b) \in \operatorname{Br} k[2]$ . Since  $\Delta$  is smooth,  $\lambda := Q_2(1, 0, 0) \neq 0$  and so by replacing  $Q_2$  and  $Q_3$  with  $-ab\lambda^{-1}Q_2$  and  $(-ab\lambda^{-1})^2Q_3$ , we may assume that  $Q_2(1, 0, 0) = -ab$ . We will again consider  $6 \times 6$  symmetric matrices defined in terms of the symmetric matrices  $M_i$ associated to  $Q_i$ , but we need some additional notation. Let  $T_2$  denote the unique upper triangular matrix such that  $T_2 + T_2^T = 2M_2$ . For any  $2 \times 2$  matrix N and any  $c \in k$ , we write  $c \oplus N$  for the  $3 \times 3$  block diagonal matrix with c in the top left entry and N in the bottom right entry. Let  $N_1$  denote the  $2 \times 2$  diagonal matrix with entries a, b so that  $M_1 = 0 \oplus N_1$ . Now define the  $6 \times 6$  symmetric matrices

$$A_{0} := \begin{pmatrix} 1 \oplus -abN_{1}^{-1} & -(0 \oplus N_{1}^{-1})T_{2}^{T} \\ -T_{2}(0 \oplus N_{1}^{-1}) & \frac{1}{ab}(M_{3} + T_{2}(0 \oplus -N_{1}^{-1})T_{2}^{T}) \end{pmatrix}, \quad A_{\infty} := \begin{pmatrix} \mathbf{0}_{3,3} & 0 \oplus I_{2} \\ 0 \oplus I_{2} & 2 \oplus \mathbf{0}_{2,2} \end{pmatrix}.$$
(3.2)

Let  $q_0, q_\infty$  be the two quadratic forms defined by these matrices, i.e.,  $q_i = \mathbf{x} A_i \mathbf{x}^T$ ; we will show that  $Z := V(q_0, q_\infty) \subset \mathbb{P}^5$  has the desired properties.

We again have  $\det(A_0 - TA_\infty) = \det(M_3 + 2TM_2 + T^2M_1)$  so Z is smooth and threedimensional, as desired. Since  $q_\infty$  is identically zero when restricted to  $V(x_3, x_4, x_5)$ , the linear section of Z given by  $V(x_3, x_4, x_5)$  is equal to  $V(q_0, x_3, x_4, x_5)$  and so is given by the rank 3 quadratic form associated to the top left  $3 \times 3$  submatrix of  $A_0$ . This proves (2).

To prove (3), we must further describe the conic bundle structure on  $\operatorname{Bl}_C Z$ . We again let  $\Lambda := V(x_3, x_4, x_5)$ , and let  $\phi \colon Z \dashrightarrow \mathbb{P}^2$  be the projection away from  $C := V(q_0) \cap \Lambda$ . Fix a point  $p = [u : v : w] \in \mathbb{P}^2$ . As above, the fiber of  $\phi$  above p is obtained by taking the Zariski closure of  $(\operatorname{Span}(\Lambda, [0 : 0 : 0 : u : v : w]) \cap Z) \smallsetminus \Lambda \subset \mathbb{P}^5$ . Note that the restriction of  $q_{\infty}$  to  $\operatorname{Span}(\Lambda, [0 : 0 : 0 : u : v : w])$  is a rank 2 quadratic form, and one can compute that vanishing of this quadratic form is the union of  $\Lambda$  and the 3-dimensional column span of the following matrix

$$B_p = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -w & -u^2 & 0 \\ 0 & v & 0 & -u^2 \\ \hline \mathbf{0}_{3,2} & [u, v, w]^T [v, w] \end{pmatrix};$$

we denote the column span of this matrix  $\Lambda_p$ . Thus the fiber of  $\phi$  above p is  $\Lambda_p \cap V(q_0)$ , which is given by the rank 3 quadratic form associated to the  $4 \times 4$  symmetric matrix  $B_p^T A_0 B_p$ . Let **M** denote the negative of the middle  $2 \times 2$  submatrix of  $B_p^T A_0 B_p$ ; **M** has rank 2. One can check by inspection of  $B_p$  and  $A_0$  that  $B_p^T A_0 B_p$  is block diagonal with the top left entry 1 and that the bottom right  $3 \times 3$  block has rank 2. Thus, the Brauer class of the generic fiber is given by the class of the rank 2 quadratic form associated to **M**. One can also check that the top left entry of **M** is  $Q_1$  and that the discriminant of the associated quadratic form, which is  $-\det(\mathbf{M})$ , is equal to  $-(ab)^{-1}v^2(Q_2^2 - Q_1Q_3)$ . Therefore,

$$[(\operatorname{Bl}_C Z)_\eta] = (-(ab)^{-1}v^2(Q_2^2 - Q_1Q_3), Q_1) = (-ab(Q_2^2 - Q_1Q_3), Q_1)$$
$$= (Q_2^2 - Q_1Q_3, Q_1) + (-ab, Q_1).$$

Furthermore, the Brauer relations show  $(-ab, av^2 + bw^2) = (-ab, a) = (b, a)$ , as desired.  $\Box$ 

# 4. A constant Brauer class from k-points of $\tilde{P}^{(1)}$

Let  $\pi: X \to \mathbb{P}^2$  be a geometrically standard conic bundle whose discriminant curve is a smooth quartic, as in Section 2.2.1. In this section, we pull back  $\pi: X \to \mathbb{P}^2$  to obtain a conic bundle over an open subset of  $\tilde{P}^{(1)}$  and then show that if  $\tilde{P}^{(1)}(k) \neq \emptyset$  then this conic bundle is constant, i.e., its Brauer class is in the image of the map  $\operatorname{Br} k \to \operatorname{Br} \mathbf{k}(\tilde{P}^{(1)})$ . This construction can be viewed as a globalization of the argument in [FJS<sup>+</sup>24, Section 6.4], and, as such, the resulting Brauer class can be viewed as an obstruction to descending representatives of curve classes to k (see Corollary 4.5 for details).

By Equation (2.6), a point  $x \in \tilde{P}^{(1)}$  corresponds to an effective divisor  $D_x$  on  $\tilde{\Delta}$ , whose image under the pushforward by  $\varpi \colon \tilde{\Delta} \to \Delta$  is contained in a line in  $\mathbb{P}^2$  (defined over  $\mathbf{k}(x)$ ). We write  $\ell_x = \operatorname{Span}(\varpi_*D_x)$  for this line. Let I be the universal line over  $\tilde{P}^{(1)}$ :


$$I := \{ (p, x) \in \mathbb{P}^2 \times \tilde{P}^{(1)} : p \in \ell_x \}$$

This comes equipped with projections  $p_1: I \to \mathbb{P}^2$  and  $p_2: I \to \tilde{P}^{(1)}$ .

**Lemma 4.1.** The second projection  $p_2: I \to \tilde{P}^{(1)}$  is a  $\mathbb{P}^1$ -bundle, and any line  $\ell \subset \mathbb{P}^2$  gives a rational section  $I_\ell := p_1^{-1}(\ell) \subset I$ . Further,  $p_2^*: \operatorname{Br} \tilde{P}^{(1)}[2] \to \operatorname{Br} I[2]$  is an isomorphism.

Proof. Fix a line  $\ell \subset \mathbb{P}^2$ . Over  $\tilde{P}^{(1)} \smallsetminus \varpi_*^{-1}(\ell \cap \Delta)$ , the second projection has an inverse morphism that sends a point  $D \in \tilde{P}^{(1)} \backsim \varpi_*^{-1}(\ell \cap \Delta)$  to the point  $(\ell_D \cap \ell, D) \in I$ . Thus  $p_2: I \to \tilde{P}^{(1)}$  is a  $\mathbb{P}^1$ -bundle, and  $I_\ell$  is a rational section. In particular, the generic fiber of  $p_2$  is  $\mathbb{P}^1_{\mathbf{k}(\tilde{P}^{(1)})}$  [CTS21, Definition 11.1.1]. Furthermore,  $p_2$  is smooth, so, since we assumed 2 is invertible in k, we have an isomorphism on the 2-primary torsion of  $\operatorname{Br}_{\operatorname{vert}}(I/S)$  and the pullback of  $\operatorname{Br}(S)$  by [CTS21, Proof of Corollary 11.1.6(ii)] (noting that [CTS21, Equation (11.3)] holds for p-primary torsion when  $p \in k^{\times}$ ).

Let  $x \in \tilde{P}^{(1)}(k)$ . We have the following commutative diagrams of varieties and Brauer groups:



**Theorem 4.2** (Strengthening of [FJS<sup>+</sup>24, Proposition 6.5]). Let k be a field of characteristic different from 2, and let  $x \in \tilde{P}^{(1)}(k)$  and let  $w \in (\ell_x \smallsetminus (\Delta \cap \ell_x))(k)$ . Then

$$[X_w] = [X_{\ell_x}] \in \operatorname{Br} \mathbf{k}(\ell_x) \quad \text{and} \quad [X_w] = p_1^*[X_\eta] \in \operatorname{Br} \mathbf{k}(I).$$

In particular,  $p_1^*[X_\eta]$  and  $[X_{\ell_x}]$  are unramified and constant.

**Definition 4.3.** Assume that  $\tilde{P}^{(1)}(k) \neq \emptyset$ . Let  $x \in \tilde{P}^{(1)}(k)$  and define

$$\tilde{\alpha}^{(1)} := x^* (p_2^*)^{-1} (p_1^*[X_\eta]) \in \operatorname{Br} k[2].$$

In particular,  $\tilde{\alpha}^{(1)} = [X_w]$  for any  $w \in (\ell_x \smallsetminus (\Delta \cap \ell_x))(k)$ .

Remarks 4.4.

- (1) Theorem 4.2 implies that  $\tilde{\alpha}^{(1)}$  is independent of the choice of  $x \in \tilde{P}^{(1)}(k)$ . However, the definition of  $\tilde{\alpha}^{(1)} \in \operatorname{Br} k$  does require that  $\tilde{P}^{(1)}(k) \neq \emptyset$ , which motivates our choice of notation.
- (2) It is possible to prove that  $p_1^*[X_\eta] \in \operatorname{Br} \mathbf{k}(I)$  is unramified without assuming that  $\tilde{P}^{(1)}(k) \neq \emptyset$ . However, in the paper we will only invoke Theorem 4.2 in situations when  $\tilde{P}^{(1)}(k) \neq \emptyset$ , so we chose to present the shorter proof with this assumption.

Proof of Theorem 4.2. By [FJS<sup>+</sup>24, Proposition 6.5], the conic bundle  $\pi|_{\ell_x} \colon X_{\ell_x} \to \ell_x$  is unramified and so, since  $\ell_x \simeq \mathbb{P}^1$ ,  $[X_{\ell_x}] = [X_w]$ . Then, the commutative diagrams (4.1) show that  $p_1^*[X_\eta]$  is equal to the image of  $[X_{\ell_x}] = [X_w]$ , as desired.

4.1. Viewing  $\tilde{\alpha}^{(1)}$  as an obstruction to the existence of cycles. One of the primary reasons that vanishing of the IJT obstruction fails to imply rationality for conic bundle threefolds is that it fails to guarantee the existence of cycles defined over k (see [FJS<sup>+</sup>24, Prop. 6.5, Theorem 6.7, and Theorem 1.4(i)]). The class  $\tilde{\alpha}^{(1)}$  measures this failure in the following precise way.

As discussed in Section 2.2.1, the k-points of  $\tilde{P}^{(1)}$  are in bijection with  $(\operatorname{CH}^2(X_{\overline{k}})^{\tilde{\gamma}_1})^{G_k}$ . The classes represented by k-rational codimension 2 cycles form a subgroup  $\operatorname{CH}^2(X)^{\tilde{\gamma}_1}$  of  $(\operatorname{CH}^2(X_{\overline{k}})^{\tilde{\gamma}_1})^{G_k}$  [CT15, Corollaire 4.2], and in general the containment can be strict.

**Corollary 4.5.** Let k be a field of characteristic different from 2. Assume that  $X(k) \neq \emptyset$ and  $\tilde{P}^{(1)}(k) \neq \emptyset$ . If  $\tilde{\alpha}^{(1)} = 0$ , then  $\operatorname{CH}^2(X)^{\tilde{\gamma}_1} = (\operatorname{CH}^2(X_{\overline{k}})^{\tilde{\gamma}_1})^{G_k}$ .

Proof. Let  $x \in \tilde{P}^{(1)}(k)$ . By  $[FJS^+24$ , Theorem 6.4] the bijection  $(CH^2(X_{\overline{k}})^{\tilde{\gamma}_1})^{G_k} \xrightarrow{\sim} \tilde{P}^{(1)}(k)$ can be realized explicitly as  $p_*j^*\varepsilon^*$  where p, j and  $\varepsilon$  are defined in  $[FJS^+24, (5.3)]$ . Thus, we will show that there exists a curve  $\gamma \subset X$ , defined over k such that  $p_*j^*\varepsilon^*[\gamma] = x \in \tilde{P}^{(1)}(k)$ . If  $\tilde{\alpha}^{(1)} = 0$  then, by Theorem 4.2,  $\pi|_{\ell_x} : X_{\ell_x} \to \ell_x$  has a section defined over k. In particular  $X_{\ell_x}(k) \neq \emptyset$ , so by  $[FJS^+24$ , Proposition 6.5] there is a curve  $\gamma$ , defined over k, such that  $p_*j^*\varepsilon^*[\gamma] = x \in \tilde{P}^{(1)}(k)$ .

5. Double covers of  $\mathbb{P}^1 \times \mathbb{P}^2$  and obstructions to sections of  $\pi_1$ 

Any smooth double cover of  $\mathbb{P}^1_{[t_0:t_1]} \times \mathbb{P}^2_{[u:v:w]}$  branched over a (2, 2)-divisor has equation

$$z^2 = t_0^2 Q_1 + 2t_0 t_1 Q_2 + t_1^2 Q_3, (5.1)$$

for a tuple of quadratic forms  $Q_1, Q_2, Q_3 \in k[u, v, w]$ . Further the quadratic forms are welldefined up to an action of Aut( $\mathbb{P}^1$ ), and this action agrees with the action of PGL<sub>2</sub> on the defining equations of  $\tilde{\Delta}$  and  $\Delta$  described in Section 2.2, so the isomorphism class of the double cover depends only on  $\tilde{\Delta}/\Delta$ . We write  $\tilde{\pi}: Y_{\tilde{\Delta}/\Delta} \to \mathbb{P}^1_{[t_0:t_1]} \times \mathbb{P}^2_{[u:v:w]}$  for this isomorphism class.

If  $\tilde{\Delta}$  and  $\Delta$  are smooth, then so is  $Y_{\tilde{\Delta}/\Delta}$  [FJS<sup>+</sup>24, Theorem 6.1(a)]. The second projection  $\pi: Y_{\tilde{\Delta}/\Delta} \to \mathbb{P}^2$  gives  $Y = Y_{\tilde{\Delta}/\Delta}$  the structure of a standard conic bundle with discriminant cover  $\tilde{\Delta}/\Delta$ , and the first projection  $\pi_1: Y \to \mathbb{P}^1$  gives Y the structure of a quadric surface bundle. By [FJS<sup>+</sup>24, Theorem 4.5(ii) and Proposition 6.3(iii)], the Prym curve  $\Gamma = \Gamma_{\tilde{\Delta}/\Delta}$  arises from the Stein factorization of  $\mathcal{F}_1(Y/\mathbb{P}^1) \to \mathbb{P}^1$ , so  $[\mathcal{F}_1(Y/\mathbb{P}^1)]$ gives a Brauer class on  $\Gamma$ . Furthermore,  $\pi_1$  has a section if and only if  $[\mathcal{F}_1(Y/\mathbb{P}^1)] = 0 \in \operatorname{Br} \Gamma^3$ .

**Theorem 5.1.** Let k be a field of characteristic different from 2, let  $\varpi \colon \tilde{\Delta} \to \Delta$  be a geometrically irreducible étale double cover of a smooth plane quartic over k, and let  $Y = Y_{\tilde{\Delta}/\Delta} \to \mathbb{P}^1 \times \mathbb{P}^2$  be the double cover associated to  $\tilde{\Delta}/\Delta$ . Then for any  $x \in \tilde{P}^{(1)}(k)$  and any  $w \in (\ell_x \smallsetminus (\ell_x \cap \Delta))(k)$ ,

$$[\mathcal{F}_1(Y/\mathbb{P}^1)] = [Y_w]_{\Gamma} = (\tilde{\alpha}^{(1)})_{\Gamma} \in \operatorname{Br} \Gamma.$$

*Proof.* The second equality holds by Definition 4.3. For the first equality, it suffices to show equality in Br  $\mathbf{k}(\Gamma)$ . By [Ami55, Corollary 9.5], if two finite-dimensional central simple algebras over  $\mathbf{k}(\Gamma)$  have the same splitting fields over  $\mathbf{k}(\Gamma)$ , then they generate the same subgroup in Br  $\mathbf{k}(\Gamma)$ . Since  $[Y_w]$  and  $[\mathcal{F}_1(Y/\mathbb{P}^1)]$  are both 2-torsion, showing they have the same splitting fields will imply the desired equality.

Let k' be an extension of  $\mathbf{k}(\Gamma)$  which is a splitting field for  $[\mathcal{F}_1(Y/\mathbb{P}^1)]$ . Then the generic fiber of  $Y \to \mathbb{P}^1$  contains lines over k', so, over k', the generic fiber becomes isomorphic to  $\mathbb{P}^1_{k'} \times \mathbb{P}^1_{k'}$ . In particular, every hyperplane section of the generic fiber contains k' points. Note that  $Y_{\ell_x}$  is a hyperplane section of the generic fiber, so then  $Y_{\ell_x}(k') \neq \emptyset$ . By Theorem 4.2,  $Y_{\ell_x}$  is Brauer equivalent to  $Y_w$ , so  $Y_w(k') \neq \emptyset$ .

Now assume that k' is an extension of  $\mathbf{k}(\Gamma)$  which is a splitting field for  $Y_w$ . Then since  $Y_w \subset Y_{\ell_x}, Y_{\ell_x}(k') \neq \emptyset$ , so [FJS<sup>+</sup>24, Prop. 6.5] implies that  $Y \to \mathbb{P}^1$  has a section over an odd degree extension of k'. By Springer's theorem [Spr52],  $Y \to \mathbb{P}^1$  then has a section over k' so  $[\mathcal{F}_1(Y/\mathbb{P}^1)]$  is trivial in Br k'.  $\square$ 

**Corollary 5.2.** Let  $\tilde{\Delta}, \Delta$  and  $Y = Y_{\tilde{\Delta}/\Delta}$  be as in Theorem 5.1. Consider the following conditions:

- (1)  $\tilde{P}^{(1)}(k) \neq \emptyset$  and  $\tilde{\alpha}^{(1)} \in \ker(\operatorname{Br} k \to \operatorname{Br} \Gamma)$  (*i.e.*,  $(\tilde{\alpha}^{(1)})_{\Gamma} = 0 \in \operatorname{Br} \Gamma$ ).
- (2)  $\pi_1: Y \to \mathbb{P}^1$  has a section (over k).
- (3) Y is rational (over k).

Then  $(1) \Rightarrow (2) \Rightarrow (3)$ .

*Proof.* (1) $\Rightarrow$ (2): By Theorem 5.1,  $(\tilde{\alpha}^{(1)})_{\Gamma} = [\mathcal{F}_1(Y/\mathbb{P}^1)] \in \operatorname{Br} \Gamma$ , and hence  $[\mathcal{F}_1(Y/\mathbb{P}^1)] = 0 \in$ Br  $\Gamma$ . Therefore  $\pi_1$  has a section (see Footnote 3).

(2) $\Rightarrow$ (3): The generic fiber of  $\pi_1$  is a quadric, which is rational as soon as it has a point.  $\Box$ 

Remark 5.3. By Proposition 2.3, the vanishing of the IJT obstruction for Y is equivalent to either  $\hat{P}^{(1)}(k) \neq \emptyset$  or  $\hat{P}(k) \neq \emptyset$ . Thus, the condition (1) in Corollary 5.2 is stronger than assuming only that the IJT obstruction vanishes.

<sup>&</sup>lt;sup>3</sup>Any quadric surface has a point if and only if it has lines defined over the discriminant extension. Indeed, if the quadric surface has a point, then slicing the surface with the tangent plane at that point will give a geometrically reducible conic whose geometric components are defined over the discriminant extension. Conversely, if there are lines defined over the discriminant extension, then the intersection of a line and its Galois conjugate will give a rational point.

**Corollary 5.4.** Let k be a local field of characteristic different from 2, let  $\hat{\Delta}, \Delta$  be as above, and let  $Y = Y_{\tilde{\Delta}/\Delta} \to \mathbb{P}^1 \times \mathbb{P}^2$  be the double cover associated to  $\tilde{\Delta}/\Delta$ . If  $\Gamma$  has index 2, i.e., if  $\Gamma(L) = \emptyset$  for all extensions L/k of odd degree, then Y is rational if and only if the IJT obstruction vanishes.

Proof. We will first show that the vanishing of the IJT obstruction is equivalent to (1) in Corollary 5.2. First,  $\operatorname{Pic}_{\Gamma}^{1}(k) \neq \emptyset$  since k is a local field [Lic69] (see Remark 7.3(i) for  $k = \mathbb{R}$ ). So by Proposition 2.3, the vanishing of the IJT obstruction is equivalent to  $\tilde{P}^{(1)}(k) \neq \emptyset$ , and in this setting we may define  $\tilde{\alpha}^{(1)} \in \operatorname{Br} k[2]$  (see Definition 4.3). Furthermore,  $\operatorname{Pic}^{1}(\Gamma) = \emptyset$  since  $\Gamma$  has index 2, so the map  $\operatorname{Pic}(\Gamma) \to \operatorname{Pic}_{\Gamma}(k)$  has a nontrivial element of order 2 in its cokernel. Since  $\operatorname{Br} k[2] \simeq \mathbb{Z}/2\mathbb{Z}$ , the low degree terms of the Hochschild–Serre spectral sequence imply that  $\operatorname{Br} k[2] \to \operatorname{Br} \Gamma[2]$  is identically 0. In particular, the condition  $\tilde{\alpha}^{(1)} \in \ker(\operatorname{Br} k \to \operatorname{Br} \Gamma)$ always holds. This shows that (1) holds if and only if the IJT obstruction vanishes.

Corollary 5.2 shows (1) $\Rightarrow$ (3). Conversely, if (3) holds, then the IJT obstruction vanishes by [BW23, Theorem 3.11].

## 6. Proof of Theorem 1.1

Let  $Q_1, Q_2, Q_3 \in k[u, v, w]$  be homogeneous forms defining a smooth plane quartic  $\Delta$  and a geometrically integral étale double cover  $\tilde{\Delta} \to \Delta$  as in (2.2)–(2.3). By assumption, there exists  $y \in \Gamma_{\tilde{\Delta}/\Delta}(k)$  which, without loss of generality, we assume lies over  $[1:0] \in \mathbb{P}^1(k)$ so that  $Q_1$  has square discriminant. By Proposition 3.1, there exists an intersection of quadrics  $Z \subset \mathbb{P}^5$  containing a conic C such that  $\operatorname{Bl}_C Z \to \mathbb{P}^2$  has discriminant cover  $\tilde{\Delta}/\Delta$ . Furthermore, the generic fiber of the conic bundle has Brauer class

$$[(\operatorname{Bl}_C Z)_{\eta}] = (Q_1, Q_2^2 - Q_1 Q_3) + [Q_1] \in \operatorname{Br} \mathbf{k}(\mathbb{P}^2).$$

Since Y is defined by the equation  $z^2 = Q_1(u, v, w)t_0^2 + 2Q_2(u, v, w)t_0t_1 + Q_3(u, v, w)t_1^2$ , completing the square shows that the Brauer class of generic fiber of the conic bundle  $\pi_2 \colon Y \to \mathbb{P}^2$  is equal to the quaternion algebra  $(Q_1, Q_2^2 - Q_1Q_3)$ . In particular,

$$[(\operatorname{Bl}_C Z)_{\eta}] - [Y_{\eta}] = [Q_1] \in \operatorname{Br} k[2].$$

On the other hand, the fiber of  $\pi_1: Y \to \mathbb{P}^1$  above [1:0] is given by  $z^2 = Q_1$ , so  $[Q_1]$  is equal to the evaluation of  $[\mathcal{F}_1(Y/\mathbb{P}^1)]$  at y. By Theorem 5.1,  $\mathcal{F}_1(Y/\mathbb{P}^1)$  is a constant algebra, so it is equal to its evaluation at y. Furthermore, the class of the relative Fano variety of lines  $[\mathcal{F}_1(Y/\mathbb{P}^1)]$  is exactly the obstruction to  $\pi_1$  having a section (see Footnote 3). Finally, we deduce that  $\operatorname{Bl}_C Z$  is rational by Corollary 3.2.

## 7. Real topological consequences of Theorem 1.1

Throughout this section we work over  $\mathbb{R}$ , where the real topology gives us additional tools to attack rationality. In particular, we are able to show that a necessary condition for rationality is that  $\tilde{\alpha}^{(1)} \in \ker(\operatorname{Br} \mathbb{R} \to \operatorname{Br} \Gamma)$  for all but one isotopy class of  $\Delta(\mathbb{R})$ .

Remark 7.1. The rigid isotopy classes of smooth quartic plane curves over  $\mathbb{R}$  are  $\emptyset$ , one oval, two nested ovals, two non-nested ovals, three ovals, and four ovals [Zeu74], where an oval is a homotopically trivial simple closed curve in  $\mathbb{P}^2(\mathbb{R})$  [Man20, Definition 2.7.7].

**Theorem 7.2.** Let  $\varpi \colon \hat{\Delta} \to \Delta$  be an étale double cover of a smooth plane quartic over  $\mathbb{R}$ , and let  $Y_{\tilde{\Delta}/\Delta}$  be the double cover defined in (5.1). Assume that  $\Delta(\mathbb{R})$  is not a single oval. Then the following are equivalent.

- (1) The IJT obstruction vanishes and  $\tilde{\alpha}^{(1)} \in \ker(\operatorname{Br} \mathbb{R} \to \operatorname{Br} \Gamma)$ .
- (2)  $\pi_1: Y \to \mathbb{P}^1$  has a section.
- (3) Y is rational.
- (4) The IJT obstruction vanishes and  $Y(\mathbb{R})$  is connected.

### Remarks 7.3.

- (i) If the IJT obstruction vanishes, then the class  $\tilde{\alpha}^{(1)}$  is defined. Indeed,  $\operatorname{Pic}_{\Gamma}^{1} \simeq \operatorname{Pic}_{\Gamma}^{3}$  has an  $\mathbb{R}$ -point because there exists a Galois-invariant partition of the Weierstrass points of  $\Gamma$  into two sets of cardinality 3. Then Proposition 2.3 shows that the vanishing of the IJT obstruction is equivalent to  $\tilde{P}^{(1)}(\mathbb{R}) \neq \emptyset$ .
- (ii) The implications  $(1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4)$  hold without any assumptions on  $\Delta(\mathbb{R})$ .
- (iii) When  $\Delta(\mathbb{R})$  is not two nested ovals (nor a single oval), the equivalences  $(2) \Leftrightarrow (3) \Leftrightarrow (4)$  were proven in [JJ24, Theorem 1.2].
- (iv) There are examples where Y is rational over  $\mathbb{R}$  and  $\tilde{\alpha}^{(1)} \in \operatorname{Br} \mathbb{R}$  is the nontrivial element; see [JJ24, Example 4.10(1)]. If  $\Delta(\mathbb{R})$  is not a single oval, such examples necessarily have that  $\operatorname{Br} \mathbb{R} \to \operatorname{Br} \Gamma$  is identically zero, which is equivalent to  $\Gamma(\mathbb{R}) = \emptyset$ .

*Proof.*  $(1) \Rightarrow (2) \Rightarrow (3)$  by Corollary 5.2. The implication  $(3) \Rightarrow (4)$  follows because the number of connected components of  $Y(\mathbb{R})$  is a birational invariant [DK81, Theorem 13.3] and the IJT obstruction is an obstruction to rationality (see Section 2.2.1).

(4) $\Rightarrow$ (1): If  $\Gamma(\mathbb{R}) = \emptyset$ , then this follows from Corollary 5.4. Now assume that  $\Gamma(\mathbb{R}) \neq \emptyset$ . Since the IJT obstruction vanishes,  $\tilde{P}^{(1)}(\mathbb{R}) \neq \emptyset$  and so by Theorem 4.2,  $\tilde{\alpha}^{(1)}$  is defined. In this case Br  $\mathbb{R} \to$  Br  $\Gamma$  is injective (evaluating at a point of  $\Gamma(\mathbb{R})$  gives a section), so we must show that  $\tilde{\alpha}^{(1)} = 0$ . Assume not. By Theorem 1.1, there is a rational threefold X' and a standard conic bundle  $\psi: X' \to \mathbb{P}^2$  such that  $[Y_\eta] - [X'_\eta] \in \operatorname{im}(\operatorname{Br} \mathbb{R} \to \operatorname{Br} \mathbf{k}(\mathbb{P}^2))$  is the obstruction to the existence of a section to  $\pi_1$ , thus is equal to  $[\mathcal{F}_1(Y/\mathbb{P}^1)] \in \operatorname{im}(\operatorname{Br} \mathbb{R} \to \operatorname{Br} \Gamma)$ . Further, Theorems 4.2 and 5.1 imply that  $[Y_\eta] - [X'_\eta] = \tilde{\alpha}^{(1)}$ . Since we assumed  $\tilde{\alpha}^{(1)}$  is nontrivial,  $\pi(Y(\mathbb{R}) - Y_\Delta(\mathbb{R})) = \overline{\mathbb{P}^2(\mathbb{R}) \setminus (\psi(X'(\mathbb{R})) \cup \Delta(\mathbb{R}))}$ . Additionally,  $\pi|_{Y_\Delta(\mathbb{R})}: Y_\Delta(\mathbb{R}) \to \Delta(\mathbb{R})$  is surjective, so  $\pi(Y(\mathbb{R})) = \overline{\mathbb{P}^2(\mathbb{R}) \setminus \psi(X'(\mathbb{R}))}$ . Furthermore, both  $\pi(Y(\mathbb{R}))$  and  $\psi(X'(\mathbb{R}))$  are nonempty and connected so Corollary 7.5(2) below implies that  $\Delta(\mathbb{R})$  is a single oval and we arrive at a contradiction.

## 7.1. Real points on conic bundles.

**Proposition 7.4.** Let  $\varpi : \tilde{\Delta} \to \Delta$  be a geometrically integral étale double cover of a smooth plane curve over  $\mathbb{R}$ , and let  $\pi : X \to \mathbb{P}^2_{\mathbb{R}}$  be a conic bundle with discriminant cover  $\tilde{\Delta}/\Delta$ . Then  $\partial(\pi(X(\mathbb{R}))) = \Delta(\mathbb{R}) \smallsetminus \varpi(\tilde{\Delta}(\mathbb{R}))$ .

**Corollary 7.5.** Let  $\varpi \colon \tilde{\Delta} \to \Delta$  be a geometrically integral étale double cover of a smooth plane curve over  $\mathbb{R}$ , and let  $\pi \colon X \to \mathbb{P}^2_{\mathbb{R}}$  be a conic bundle with discriminant cover  $\varpi \colon \tilde{\Delta} \to \Delta$ .

- (1) Then  $\pi(X(\mathbb{R}))$  and  $(\mathbb{P}^2(\mathbb{R}) \setminus \pi(X(\mathbb{R})))$  are both nonempty and connected if and only if  $\Delta(\mathbb{R}) \setminus \varpi(\tilde{\Delta}(\mathbb{R}))$  is a simple closed curve.
- (2) Then  $\pi(X(\mathbb{R}))$  and  $(\mathbb{P}^2(\mathbb{R}) \setminus \pi(X(\mathbb{R}))) \cup \Delta(\mathbb{R})$  are both nonempty and connected if and only if  $\Delta(\mathbb{R})$  is a simple closed curve.

*Remark* 7.6. Note that we make no assumption on the degree of  $\Delta$ .

Proof of Corollary 7.5. The loci  $\pi(X(\mathbb{R}))$  and  $(\mathbb{P}^2(\mathbb{R}) \setminus \pi(X(\mathbb{R})))$  are both nonempty and connected if and only if the boundary  $\partial(\pi(X(\mathbb{R})))$  is connected. Thus, part (1) follows

from Proposition 7.4, the Jordan curve theorem, and the classification of the real loci of smooth plane curves [Man20, Section 2.7]. Part (2) follows from part (1) because  $(\mathbb{P}^2(\mathbb{R}) \setminus \pi(X(\mathbb{R}))) \cup \Delta(\mathbb{R})$  is equal to  $(\mathbb{P}^2(\mathbb{R}) \setminus \pi(X(\mathbb{R}))) \sqcup \varpi(\tilde{\Delta}(\mathbb{R}))$ .

Proof of Proposition 7.4. Fix a Zariski affine open  $U = \operatorname{Spec} A \subset \mathbb{P}^2$  where  $X_U$  can be viewed as a subvariety of  $\mathbb{P}^2_A$ , and fix a defining quadratic equation F for  $X_U$ . Since  $\pi$  is smooth away from  $\Delta$ , the signature of F is constant on real connected components of  $U(\mathbb{R}) \setminus (\Delta \cap U)(\mathbb{R})$ . Note that a connected component B of  $U(\mathbb{R}) \setminus (\Delta \cap U)(\mathbb{R})$  is in the image of  $X_U(\mathbb{R})$  if and only if  $F|_B$  is an indefinite quadratic form.

Fix a real connected component c of  $(\Delta \cap U)(\mathbb{R})$  and let  $B_1, B_2$  be the two distinct connected components of  $U(\mathbb{R}) \setminus (\Delta \cap U)(\mathbb{R})$  such that  $B_1 \cup B_2 \cup c$  becomes connected, i.e.,  $B_1, B_2$  are the immediate interior and immediate exterior of c. The proposition now follows from the following two claims.

- (1) The signatures of  $F|_{B_1}$  and  $F|_{B_2}$  differ by exactly one sign. In particular, we may assume that  $F|_{B_1}$  is indefinite.
- (2) The quadratic form  $F|_{B_2}$  is also indefinite if and only if  $c \subset \varpi(\tilde{\Delta}(\mathbb{R}))$ .

By continuity, it suffices to prove both claims in a (Euclidean) open neighborhood around a point  $x_0 \in c \subset (\Delta \cap U)(\mathbb{R})$ . In an open neighborhood of  $x_0$ , we may change coordinates so that F is diagonal and two of the coefficients are nowhere vanishing (see [IOOV17, Lemma 6.3]). Thus, the sign of those coefficients are equal on  $B_1$  and  $B_2$ . Since the vanishing of the discriminant of F exactly agrees with  $\Delta$ , the third coefficient must vanish with multiplicity 1 exactly along  $\Delta$ . In particular, it has a different sign on  $B_1$  than on  $B_2$ . This completes the proof of the first claim.

To prove the second claim, we assume without loss of generality that the signature of  $F|_{B_1}$  is (+, +, -). By the first claim, the signature of  $F|_{B_2}$  is either (+, +, +) or (+, -, -). Furthermore, the proof of the first claim above shows that the signature of  $F|_{B_2}$  is (+, -, -) if and only if the signature of  $F|_{x_0}$  is (+, 0, -). Finally, since  $\Delta$  is smooth and  $\tilde{\Delta}$  is geometrically integral,  $F|_{x_0}$  is a rank 2 quadratic form, and its vanishing has a smooth  $\mathbb{R}$ -point if and only if  $x_0 \in \varpi(\tilde{\Delta}(\mathbb{R}))$ . This completes the proof of the second claim.

The outside of  $\Delta(\mathbb{R})$  is the unique non-orientable component of  $\mathbb{P}^2(\mathbb{R}) \setminus \Delta(\mathbb{R})$  (which exists by [Man20, Section 2.7]).

**Proposition 7.7.** Let  $\varpi : \tilde{\Delta} \to \Delta$  be an étale double cover of a smooth plane quartic over  $\mathbb{R}$ , and let  $Y_{\tilde{\Delta}/\Delta}$  be the double cover defined in (5.1). Assume that  $\tilde{P}^{(1)}(\mathbb{R}) \neq \emptyset$ . Then the class  $\tilde{\alpha}^{(1)}$  is trivial if and only if the image of  $Y(\mathbb{R}) \to \mathbb{P}^2(\mathbb{R})$  contains the outside of  $\Delta(\mathbb{R})$ .

Proof. Let  $x \in \tilde{P}^{(1)}(\mathbb{R})$ ; write  $D_x \in \operatorname{Pic}_{\tilde{\Delta}}^4$  for the associated effective divisor (see (2.6)) and  $\ell_x$  for the line in  $\mathbb{P}^2$  spanned by  $\varpi_* D_x$ . The key observation is that  $\ell_x(\mathbb{R}) \smallsetminus (\ell_x \cap \Delta)(\mathbb{R})$  meets the outside of  $\Delta(\mathbb{R})$ . If  $\tilde{\alpha}^{(1)} = 0$ , then  $Y_{\ell_x}(\mathbb{R}) \to \ell_x(\mathbb{R})$  is surjective by Theorem 4.2, and hence the image of  $Y(\mathbb{R}) \to \mathbb{P}^2(\mathbb{R})$  meets (and consequently contains) the outside of  $\Delta(\mathbb{R})$ . Conversely, if the image of  $Y(\mathbb{R}) \to \mathbb{P}^2(\mathbb{R})$  contains the outside of  $\Delta(\mathbb{R})$ , then there exists  $w \in \ell_x(\mathbb{R}) \smallsetminus (\ell_x \cap \Delta)(\mathbb{R})$  such that  $Y_w(\mathbb{R}) \neq \emptyset$ . Thus  $\tilde{\alpha}^{(1)} = 0$  by Theorem 4.2.

7.2. Cases when the IJT obstruction or topological obstruction individually characterize rationality. Theorem 7.2 allows us to extend results of [JJ24] and classify the cases from Theorem 7.2 when the IJT obstruction or the real topological obstruction individually characterize rationality. **Proposition 7.8.** Let  $\varpi \colon \dot{\Delta} \to \Delta$  be an étale double cover of a smooth plane quartic over  $\mathbb{R}$ , and let  $Y_{\dot{\Delta}/\Delta}$  be the double cover defined in (5.1). Assume that  $\Delta(\mathbb{R})$  is not one oval.

If  $\Delta(\mathbb{R})$  is two nested ovals, three ovals, or four ovals, then the four conditions of Theorem 7.2 are also equivalent to

(5)  $Y(\mathbb{R})$  is connected.

If  $\Delta(\mathbb{R})$  is two nested ovals or four ovals then (5) is additionally equivalent to

(6) The IJT obstruction vanishes for Y.

Moreover, the assumptions above are the strongest possible, i.e., if  $\Delta(\mathbb{R})$  is empty or two non-nested ovals then connectedness is not equivalent to rationality, and if  $\Delta(\mathbb{R})$  is empty, two non-nested ovals, or three ovals then vanishing of the IJT obstruction is not equivalent to rationality.

Remark 7.9. The part of the proposition that is new is the case of two nested ovals; the rest follows from prior work. If  $\Delta(\mathbb{R})$  is four ovals, then Y is always rational [JJ24, Theorem 1.2(4)] so the theorem vacuously holds. If  $\Delta(\mathbb{R})$  is three ovals, then the result that connectedness of  $Y(\mathbb{R})$  characterizes rationality is [JJ24, Theorem 1.2(3)]. That connectedness fails to characterize rationality if  $\Delta(\mathbb{R})$  is empty or two non-nested ovals is [JJ24, Examples 4.2 and 4.4]. That the IJT obstruction fails to characterize rationality if  $\Delta(\mathbb{R})$  is two non-nested ovals is shown by [FJS<sup>+</sup>24, Theorem 1.4(i)]; if  $\Delta(\mathbb{R})$  is empty or three ovals, this is shown by [JJ24, Examples 4.1 and 4.6].

*Proof.* Given the remark above, we restrict ourselves to the case that  $\Delta(\mathbb{R})$  is two nested ovals. In this case, (6) $\Rightarrow$ (5) by [JJ24, Corollary 3.9]. This shows that (6) is equivalent to (4).

It remains to show that  $(5) \Rightarrow (6)$ . Assume for the sake of contradiction that (5) holds and that Y has an IJT obstruction to rationality over  $\mathbb{R}$ . Then  $\pi_1$  does not have a section by Theorem 7.2, so  $\pi_1$  is not surjective on  $\mathbb{R}$ -points by [Wit37, Satz 22]. Thus there exists a point  $t \in \mathbb{P}^1(\mathbb{R})$  such that  $Y_t(\mathbb{R}) = \emptyset$ , and so is defined by a definite quadratic form. In particular, there exists  $z \in \Gamma_t(\mathbb{R})$  with  $\mathcal{F}(Y/\mathbb{P}^1)_z = \mathbb{H}$ . By Proposition 3.1 and Corollary 3.2, there exists an irrational complete intersection  $Z \subset \mathbb{P}^5$  of two quadrics, containing a conic C, such that  $\psi: \operatorname{Bl}_C Z \to \mathbb{P}^2$  is a conic bundle with

$$[(\operatorname{Bl}_C Z)_\eta] = [Y_\eta] + \mathbb{H} \in \operatorname{Br} \mathbf{k}(\mathbb{P}^2),$$

so the image  $\psi((\operatorname{Bl}_C Z)(\mathbb{R}))$  is equal to  $\overline{\mathbb{P}^2(\mathbb{R}) \setminus \psi(Y(\mathbb{R}))} \cup \Delta(\mathbb{R})$ . Since  $Y(\mathbb{R})$  and  $(\operatorname{Bl}_C Z)(\mathbb{R})$ are both 3-manifolds, the images  $\pi(Y(\mathbb{R}))$  and  $\psi((\operatorname{Bl}_C Z)(\mathbb{R}))$  do not have any connected components that are 1-manifolds. Thus, since  $Y(\mathbb{R})$  is connected and  $\Delta(\mathbb{R})$  is two nested ovals, the image  $\pi(Y(\mathbb{R}))$  must be the region (diffeomorphic to an annulus) enclosed by the two ovals of  $\Delta(\mathbb{R})$ . This implies  $Z(\mathbb{R})$  is disconnected, so Krasnov's real isotopy classification of 3-dimensional intersections of two quadrics [Kra18, Theorem 5.4] (see also [HT21b, Sections 11.2 and 11.4]) implies that  $Z(\mathbb{R})$  is diffeomorphic to  $\mathbb{S}^3 \sqcup \mathbb{S}^3$  and that the Prym curve  $\Gamma$ has 6 real Weierstrass points. On the other hand, since we've assumed that  $\pi_1(Y(\mathbb{R})) \subseteq \mathbb{P}^1(\mathbb{R})$ is connected, the fact that  $\Gamma$  has 6 real Weierstrass points implies that  $\pi_1: Y \to \mathbb{P}^1$  must have a smooth fiber with signature (+, +, -, -). After a coordinate change on  $\mathbb{P}^1$  we may assume that the fiber of  $\pi_1$  over [1:0] has signature (+,+,-,-) [FJS<sup>+</sup>24, Theorem 4.5], i.e. that the quadratic form  $Q_1$  has signature (+, +, -). Then  $(Q_1 \ge 0) = \mathbb{P}^2(\mathbb{R}) \setminus \mathbb{D}^2 \subset \mathbb{P}^2(\mathbb{R})$ , and this region is contained in  $\pi(Y(\mathbb{R}))$  by [JJ24, Lemma 2.8]. This contradicts our earlier conclusion that  $Y(\mathbb{R})$  is the region enclosed by the two ovals of  $\Delta(\mathbb{R})$ , so we have shown (5) implies (6). 

#### References

- [Ami55] S. A. Amitsur, Generic splitting fields of central simple algebras, Ann. of Math. (2) 62 (1955), 8–43. MR70624 ↑5
- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust, The Magma algebra system. I. The user language, J. Symbolic Comput. 24 (1997), no. 3-4, 235–265. Computational algebra and number theory (London, 1993). MR1484478 ↑2, 3
- [Bea77] Arnaud Beauville, Variétés de Prym et jacobiennes intermédiaires, Ann. Sci. École Norm. Sup. (4) 10 (1977), no. 3, 309–391. MR472843 <sup>↑</sup>2.2
- [Bru08] Nils Bruin, The arithmetic of Prym varieties in genus 3, Compos. Math. 144 (2008), no. 2, 317–338. MR2406115 ↑2.2.1, 2.2.1, 1
- [BW23] Olivier Benoist and Olivier Wittenberg, Intermediate Jacobians and rationality over arbitrary fields, Ann. Sci. Éc. Norm. Supér. (4) 56 (2023), no. 4, 1029–1086. MR4650157 ↑1, 2.2.1, 3.2, 3, 5
- [CG72] C. Herbert Clemens and Phillip A. Griffiths, The Intermediate Jacobian of the cubic threefold, Ann. of Math. 95 (1972), no. 2, 281–356. MR0302652 ↑1, 2.2.1
- [CT15] Jean-Louis Colliot-Thélène, Descente galoisienne sur le second groupe de Chow: mise au point et applications, Doc. Math. (2015), 195–220. MR3404380 ↑4.1
- [CTS21] Jean-Louis Colliot-Thélène and Alexei N. Skorobogatov, The Brauer-Grothendieck group, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 71, Springer, Cham, 2021. MR4304038 ↑2.1, 2.1, 3, 4
- [DK81] Hans Delfs and Manfred Knebusch, Semialgebraic topology over a real closed field. II. Basic theory of semialgebraic spaces, Math. Z. 178 (1981), no. 2, 175–213. MR631628 ↑7
- [FJS<sup>+</sup>24] Sarah Frei, Lena Ji, Soumya Sankar, Bianca Viray, and Isabel Vogt, Curve classes on conic bundle threefolds and applications to rationality, Algebr. Geom. 11 (2024), no. 3, 421–459. ↑1, 1, 1, 1, 1, 2.1, 2.1, 2.2, 2.2.1, 2, 2.4, 2.3, 3, 4, 4.2, 4, 4.1, 4.1, 5, 5, 7.9, 7.2
  - [GS17] Philippe Gille and Tamás Szamuely, Central simple algebras and Galois cohomology, Second, Cambridge Studies in Advanced Mathematics, vol. 165, Cambridge University Press, Cambridge, 2017. MR3727161 ↑2.1
- [HT21a] Brendan Hassett and Yuri Tschinkel, Cycle class maps and birational invariants, Comm. Pure Appl. Math. 74 (2021), no. 12, 2675–2698. MR4373165 ↑1, 2.2.1
- [HT21b] \_\_\_\_\_, Rationality of complete intersections of two quadrics over nonclosed fields, Enseign. Math. 67 (2021), no. 1-2, 1-44. With an appendix by Jean-Louis Colliot-Thélène. MR4323991 ↑1, 2.2.1, 3.2, 3, 7.2
- [IOOV17] Colin Ingalls, Andrew Obus, Ekin Ozman, and Bianca Viray, Unramified Brauer classes on cyclic covers of the projective plane, Brauer groups and obstruction problems, 2017, pp. 115–153. With an appendix by Hugh Thomas. MR3616009 ↑7.1
  - [IP99] V. A. Iskovskikh and Yu. G. Prokhorov, Fano varieties, Algebraic geometry, V, 1999, pp. 1–247. MR1668579 ↑1
  - [JJ24] Lena Ji and Mattie Ji, Rationality of real conic bundles with quartic discriminant curve, Int. Math. Res. Not. IMRN 1 (2024), 115–151. MR4686647 ↑1, (iii), (iv), 7.2, 7.9, 7.2
  - [KP23] Alexander Kuznetsov and Yuri Prokhorov, Rationality of Fano threefolds over non-closed fields, Amer. J. Math. 145 (2023), no. 2, 335–411. MR4570985 ↑1
  - [Kra18] V. A. Krasnov, On the intersection of two real quadrics, Izv. Ross. Akad. Nauk Ser. Mat. 82 (2018), no. 1, 97–150. MR3749598 ↑7.2
  - [Lam05] T. Y. Lam, Introduction to quadratic forms over fields, Graduate Studies in Mathematics, vol. 67, American Mathematical Society, Providence, RI, 2005. MR2104929 ↑2.1
  - [Lic69] Stephen Lichtenbaum, Duality theorems for curves over p-adic fields, Invent. Math. 7 (1969), 120–136. MR242831 ↑5
- [Man20] Frédéric Mangolte, Real algebraic varieties, Springer Monographs in Mathematics, Springer, Cham, 2020. Translated from the 2017 French original [3727103] by Catriona Maclean. MR4179588 ↑7.1, 7.1, 7.1
- [Mum74] David Mumford, Prym varieties. I, Contributions to analysis (a collection of papers dedicated to Lipman Bers), 1974, pp. 325–350. MR0379510 ↑2.2

17

- [Rei72] Miles Reid, The complete intersection of two or more quadrics, 1972. Ph.D thesis (Trinity College, Cambridge). ↑3
- [Spr52] Tonny Albert Springer, Sur les formes quadratiques d'indice zéro, C. R. Acad. Sci. Paris 234 (1952), 1517–1519. MR47021 ↑2.1, 5
- [Wit37] Ernst Witt, Theorie der quadratischen Formen in beliebigen Körpern, J. Reine Angew. Math. 176 (1937), 31–44. MR1581519 ↑7.2
- [Zeu74] Hieronymus Georg Zeuthen, Sur les différentes formes des courbes planes du quatrième ordre, Mathematische Annalen 7 (1874), 410–432 (fre). ↑7.1

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, 27 N. MAIN STREET, HANOVER, NH 03755 Email address: sarah.frei@dartmouth.edu URL: https://math.dartmouth.edu/~sfrei

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 530 CHURCH STREET, ANN ARBOR, MI 48109-1043

Email address: lenaji.math@gmail.com URL: https://www-personal.umich.edu/~lenaji

MATHEMATICAL INSTITUTE, UTRECHT UNIVERSITY, HANS FREUDENTHAL BUILDING, BUDAPEST 6, 3584 CD UTRECHT, THE NETHERLANDS

Email address: s.sankar@uu.nl

URL: https://sites.google.com/site/soumya3sankar/

University of Washington, Department of Mathematics, Box 354350, Seattle, WA 98195, USA

*Email address*: bviray@uw.edu *URL*: http://math.washington.edu/~bviray

BROWN UNIVERSITY, DEPARTMENT OF MATHEMATICS, BOX 1917, 151 THAYER STREET, PROVIDENCE, RI 02912, USA

Email address: ivogt.math@gmail.com URL: https://www.math.brown.edu/ivogt/