

DUAL OF THE GEOMETRIC LEMMA AND THE SECOND ADJOINTNESS THEOREM FOR p -ADIC REDUCTIVE GROUPS

KEI YUEN CHAN

ABSTRACT. Let P, Q be standard parabolic subgroups of a p -adic reductive group G . We study the smooth dual of the filtration on a parabolically induced module arising from the geometric lemma associated to the cosets $P \backslash G/Q$. We prove that the dual filtration coincides with the filtration associated to the cosets $P \backslash G/Q^-$ via the Bernstein-Casselman canonical pairing from the second adjointness of parabolic induction. This result generalizes a result of Bezrukavnikov-Kazhdan on the explicit description in the second adjointness. Along the way, we also study some group theoretic results.

1. INTRODUCTION

Let G be a connected reductive group over a non-Archimedean local field F . Let $\text{Alg}(G)$ be the category of smooth representations of G over \mathbb{C} . The parabolic induction and Jacquet functor form a pair of adjoint functors and are fundamental tools in the representation theory of reductive p -adic groups. The Bernstein geometric lemma describes how to decompose the Jacquet functors of parabolically induced modules, an analog of Mackey decomposition in finite groups.

It follows from definitions that the parabolic induction admits the Jacquet functor as the natural left adjoint functor. Casselman (see [Ca93]) discovered the opposite Jacquet functor is right adjoint to the parabolic induction for admissible representations. Generalizing to all smooth representations is highly non-trivial and is first shown by Bernstein [Be92] using his stabilization theorem. The later adjointness is usually referred to the second adjointness theorem. The geometric viewpoint of the second adjointness is studied in [BK15].

This paper studies the interplay between the second adjointness and the geometric lemma. To state results more precisely, we need more notations. Let P be a parabolic subgroup of G with Levi decomposition MU (where M is the Levi subgroup of P and U is the unipotent radical of P). For a smooth representation σ of M , define $\text{Ind}_P^G \sigma$ to be the normalized parabolic induction; and for a smooth representation π of G , define π_U to be the normalized Jacquet functor.

Let Q be another parabolic subgroup of G with the Levi decomposition NV , where N is the Levi subgroup of Q while V is the unipotent subgroup of Q . The geometric lemma describes a decomposition for

$$(\text{Ind}_P^G \sigma)_V$$

as N -representations. For a smooth representation π of G , we denote by π^\vee the smooth dual of π . By taking the exact contravariant dual smooth functor (as N -representations), we then obtain one filtration on $((\text{Ind}_P^G \sigma)_V)^\vee$.

On the other hand, the second adjointness gives the natural isomorphism:

$$((\text{Ind}_P^G \sigma)_V)^\vee \cong ((\text{Ind}_P^G \sigma)^\vee)_{V^-},$$

where V^- is the unipotent radical of the parabolic subgroup opposite to Q . The work of Bernstein and Casselman shows a more explicit non-degenerate N -equivariant pairing for

$(\mathrm{Ind}_P^G \sigma)_V$ and $((\mathrm{Ind}_P^G \sigma)^\vee)_{V^-}$, which we shall refer to the Bernstein-Casselman canonical pairing. There is a natural isomorphism between $(\mathrm{Ind}_P^G \sigma)^\vee$ and $\mathrm{Ind}_P^G \sigma^\vee$ and so one can also obtain a filtration on the space $((\mathrm{Ind}_P^G \sigma)^\vee)_{V^-}$, using the orbits in $P \backslash G/Q^-$. Here Q^- is the parabolic subgroup opposite to Q . Our main result is the following:

Theorem 1.1. *The two filtrations above coincide via the Bernstein-Casselman canonical pairing.*

The statement seems to be natural, but we are not aware of a reference. The proof of our approach involves some technicalities which may be of independent interest.

More precise definitions for the filtrations are given in Sections 6.1 and 6.3. We remark that in order to match two filtrations, we need the compatibility of the Bruhat orderings in Section 2.

We now describe how to show Theorem 1.1. The Bernstein-Casselman canonical pairing for $(\mathrm{Ind}_P^G \sigma)_V$ and $((\mathrm{Ind}_P^G \sigma)^\vee)_{V^-}$ is expressed in terms of the original pairing of $\mathrm{Ind}_P^G \sigma$ and its dual. One crucial ingredient is the so-called canonical lifting defined from the stabilization theorem. Using the group-theoretic properties in Section 3, we show that the canonical lift is well-behaved under the filtration arising from the geometry (Proposition 7.3).

We now give a motivation for Theorem 1.1. Let P be a parabolic subgroup of G and let P^- be its opposite parabolic subgroup. Let $P = MU$ be its Levi decomposition with M to be the Levi subgroup and U to be the unipotent radical of P . Let σ be an irreducible representation of M and let π be a quotient of $\mathrm{Ind}_P^G \sigma$. We have the following composition of maps:

$$C_c^\infty(PP^-, \sigma) \hookrightarrow \mathrm{Ind}_P^G \sigma \twoheadrightarrow \pi,$$

where $C_c^\infty(PP^-, \sigma)$ is the subspace of functions in $\mathrm{Ind}_P^G \sigma$ compactly supported in the open subspace $P \backslash PP^-$. Taking the Jacquet functor on U^- , it gives a map $\sigma \rightarrow \pi_{U^-}$, accounting for the adjoint map in the second adjointness. This map is explicitly studied before in the work of Bezrukavnikov-Kazhdan [BK15] from geometric viewpoint. Theorem 1.1 recovers such map (see Section 8.2) and so can be viewed as a refinement of the description of the adjoint map. We are originally motivated from some questions of the study of derivatives and related discussions are in Section 8.

We finally remark that it completely makes sense to ask similar questions in the setting for affine Hecke algebras or graded Hecke algebras. Those settings are less technical and can be mainly deduced from Section 2. We shall not go into the details. One may see [BC15] for stating the second adjointness in those relevant settings.

This paper is organized as follows. Section 2 reviews some properties of Weyl groups, including the Bruhat ordering. Section 3 studies intersections of Bruhat cells. Section 4 reviews the Bernstein-Casselman pairing. Section 5 performs a support computation for a canonical lift of an element in a parabolically induced module, which uses results in Section 3. Section 6 describes filtrations from the geometric lemma. Section 7 proves our main result on matching filtrations in Theorem 7.7. Section 8 studies some consequences.

1.1. Acknowledgements. The author would like to thank Xuhua He and Jianghua Lu for discussions on Richardson varieties. The author would also like to thank Caihua Luo for helpful discussions. This project is supported in part by the Research Grants Council of the Hong Kong Special Administrative Region, China (Project No: 17305223) and the National Natural Science Foundation of China (Project No. 12322120).

2. SOME RESULTS ON WEYL GROUPS

2.1. Basic setup and notations. Let R be a root system and let Φ be a choice of simple roots. Let W be the Weyl group associated to R . For a subset $S \subset \Phi$, let $W(S)$ be the subgroup of W generated by the simple reflections associated to roots in S . For $w \in W$, the S -part, denoted by w_S , of w is the unique element such that $w w_S^{-1} \in W(S)$. Let w_0 be the longest element in W . In particular, we write $w_{0,S}$ for the S -part of w_0 . We let $\theta(\alpha) = -w_0(\alpha)$ for $\alpha \in R$.

Let $w \in {}_F W$ with a reduced expression $s_r \dots s_1$. A *subexpression* of w is a sequence $\{w_0, w_1, \dots, w_r\}$ of elements in ${}_F W$ with the conditions:

- (i) $w_0 = \text{Id}$;
- (ii) $w_i w_{i-1}^{-1} \in \{\text{Id}, s_i\}$ for all $1 \leq i \leq r$.

By abuse of notations, we also say w_r is a subexpression of w . Let \leq be the Bruhat ordering on W i.e. $w' \leq w$ if w' is a subexpression of w .

One basic result is the compatibility of the Bruhat ordering with S parts:

Lemma 2.1. [De77, Lemmas 3.5,3.6] *Let $w, w' \in W$. Let $S \subset \Phi$. If $w \leq w'$, then $w_S \leq w'_S$.*

2.2. Order reversing operation. For $S, T \subset \Phi$, let $W_{S,T}$ be the set of minimal representatives of the double cosets in $W(S) \setminus W/W(T)$. We also have the following version of Lemma 2.1:

Lemma 2.2. *Let $w, w' \in W$. Let u, u' be the respective minimal representatives in $W(S)wW(T)$ and $W(S)w'W(T)$. If $w \leq w'$, then $u \leq u'$.*

Proof. Suppose $w \leq w'$. Let v, v' be the respective minimal representatives in $wW(T)$ and $w'W(T)$. Then, by Lemma 2.1, $v \leq v'$. Note that u, u' are minimal representatives of the double cosets in $W(S)v$ and $W(S)v'$ respectively. Then, by the left version of Lemma 2.1, we have $u \leq u'$. □

We shall need the following lemma:

Lemma 2.3. *Let $T \subset \Phi$. Then $w_{0,T}^{-1}W(T)w_{0,T} = W(\theta(T))$.*

Proof. It is well-known that $-w_0(\Phi) = \Phi$ and so $\theta(T) \subset \Phi$. Let x be the longest element in $W(T)$. Then $xw_{0,T} = w_0$. Then

$$w_{0,T}^{-1}W(T)w_{0,T} = w_{0,T}^{-1}x^{-1}W(T)xw_{0,T} = w_0^{-1}W(T)w_0 = W(\theta(T)).$$

□

We now define a map:

$$(2.1) \quad \Omega_{S,T} : W_{S,T} \rightarrow W_{S,\theta(T)}$$

such that $\Omega_{S,T}(u)$ is the minimal representative in the coset

$$W(S)uw_{0,T}W(\theta(T)).$$

We may simply write Ω for $\Omega_{S,T}$ if it is clear from the context.

Proposition 2.4. (c.f. [Ri92, Lemma 3.6]) *The map Ω is an order-reversing bijection.*

Proof. We first show it is a bijection. Note that, by Lemma 2.3,

$$W(S)uW(T)w_{0,T} = W(S)uw_{0,T}W(\theta(T)).$$

Thus $\Omega(u) = \Omega(u')$ if and only if $u = u'$, which shows that Ω is a bijection.

We now show it is order-reversing. Let $u, u' \in W_{S,T}$ with $u \leq u'$. It is standard that:

$$(2.2) \quad uw_0 \geq u'w_0.$$

Note that the $\theta(T)$ -part of uw_0 (resp. $u'w_0$) coincides with that of $uw_{0,T}$ (since uw_0 can be obtained from $uw_{0,T}$ by multiplying an element in $W(T)$ on the left). Thus, by Lemma 2.1 and (2.2), $\Omega(u) \geq \Omega(u')$. \square

3. SOME RESULTS ON BRUHAT CELLS

3.1. Notations. We shall follow the notions of structure of linear algebraic groups in [Bo91], which will be used in the rest of the article. Let \mathbb{G} be a connected reductive group defined over a non-Archimedean local field F . One may see [CGP15] for a modern account on the subject. Let $G = \mathbb{G}(F)$ be the group of F -rational points of \mathbb{G} . Fix a maximal split torus T in G . Let ${}_F R = R(G, T)$ be the relative root system of G with respect to T ([Bo91, 8.17]). Let $Z_G(T)$ be the subgroup of G centralizing T and let ${}_F W = N_G(T)/Z_G(T)$ be the relative Weyl group of G . Let R_{nd} be the subset of non-divisible roots in ${}_F R$. We fix a set Φ of simple roots in ${}_F R$, which determines the set of positive roots, denoted R^+ . Let $R_{nd}^+ = R^+ \cap R_{nd}$. For $S \subset \Phi$, let $R_{nd,S}$ be the root sub-system of $R_{nd} \cap (\mathbb{Z}S)$.

For each root $\alpha \in {}_F R$, we associate to a root subgroup \mathbb{U}_α defined over F and write $U_\alpha = \mathbb{U}_\alpha(F)$ [Bo91, Proposition 21.9]. The group U is directly spanned by $U_{(\alpha)}$ ($\alpha \in R_{nd}^+$) taken in any order i.e.

$$\prod_{\alpha \in R_{nd}^+} U_{(\alpha)} \rightarrow U$$

is an isomorphism (in any order taken) [Bo91, Proposition 21.9]. For $\alpha \in R_{nd}$, let $U_{(\alpha)} = U_\alpha$ if $2\alpha \notin R$ and let $U_{(\alpha)}$ be the subgroup generated by U_α and $U_{2\alpha}$ otherwise.

Let \mathbb{B} be the minimal parabolic F -subgroup generated by $Z_G(T)$ and \mathbb{U}_α ($\alpha \in R$) and let $B = \mathbb{B}(F)$. A *standard parabolic subgroup* of G is a subgroup generated by $Z_G(T)$ and $U_{(\alpha)}$ for $\alpha \in R_{nd,S}$ for some subset $S \subset \Phi$ ([Bo91, Section 21.11]). For a standard parabolic subgroup P of G , we also write M_P to be the Levi subgroup in P , write U_P to be the unipotent subgroup in P , and write U_P^- to be the unipotent subgroup of the opposite parabolic subgroup of P .

3.2. Bruhat cells. We keep using the above notations. For each $w \in {}_F W$, we choose a representative \dot{w} of w in $N_G(T)$. Define

$$C(w) := B\dot{w}B,$$

which is independent of a choice of \dot{w} .

According to [Bo91, Theorem 21.26], the Zariski closure

$$\overline{C(w)} = \bigsqcup_{w' \leq w} C(w'),$$

which is also closed under the topology of an l -group.

Let B^- be the opposite minimal parabolic subgroup of B . Let U be the unipotent radical of B . Define

$$U_{w,-} = U \cap \dot{w}^{-1}U^-\dot{w}, \quad U_{w,+} = U \cap \dot{w}^{-1}U\dot{w}.$$

3.3. Open compact subgroups. It is well-known that G is a l -group i.e. Hausdorff topological group such that the identity has a basis of neighborhoods which are open and compact (see [Be92, Ch II, Section 2.1]). Let K_0 be the hyperspecial open compact subgroup in G . For an open compact subgroup K and $\alpha \in R_{nd}$, let $K_{(\alpha)} = K \cap U_{(\alpha)}$ and $K' = K \cap T$. An open compact subgroup K of G is said to be *in a good position* if the natural map

$$K' \times \prod_{\alpha \in R_{nd}} K_{(\alpha)} \rightarrow K$$

is a bijection, for any ordering on α 's.

In the context below, all chosen open compact subgroups are assumed to be in K_0 and are normal in K_0 . We can always choose such sufficiently small open compact subgroup which is also in a good position (see Bruhat's theorem [Be92, Page 41], also see [BD84]), which we shall use several times. In particular, this also implies one can always find a sufficiently small open compact subgroup K with respect to any parabolic subgroup P i.e.

$$K = (K \cap U_P) \cdot (K \cap M_P) \cdot (K \cap U_{\bar{P}}).$$

3.4. Intersection of Bruhat cells. The intersection $B\dot{w}B \cap B\dot{w}'B^-$ typically refers to the Richardson variety when the underlying field is over \mathbb{C} . The study of the intersection appears in [De86], and some refinement is obtained in [Ri92]. We shall explain the intersection in our content and we first explain some combinatorial ingredients.

Definition 3.1. Let $w \in {}_F W$. A subexpression of w is said to be *distinguished* if for each $1 \leq i \leq r$, $w_i \leq w_{i-1} s_i$.

Note that for the distinguished condition, if it is a strict inequality, then $w_{i-1} = w_i$.

Theorem 3.2. [De86] *The set of distinguished subexpressions of w coincides with the set of subexpressions.*

Proposition 3.3. (c.f. [De86]) *Let $w, w' \in {}_F W$. Then $B\dot{w}B \cap B\dot{w}'B^- \neq \emptyset$ if and only if $w' \leq w$.*

Proof. Since [De86, Theorem 1.1] is stated for over \mathbb{C} , we shall give some details following his proof in our case. We make some simplifications and hence we do not have (and do not need) a concrete description on the intersection.

We first prove the only if direction. Suppose $B\dot{w}B \cap B\dot{w}'B^- \neq \emptyset$. Write the reduced expression for w as $s_r \dots s_1$. Then we have:

$$\dot{s}_r \dots \dot{s}_1 u \in B\dot{w}'B^-.$$

for some $u \in U$. Let $z'_i \in W$ be the unique element such that $\dot{s}_i \dots \dot{s}_1 u \in Bz'_i B^-$. Now, one begins with u and inductively applies \dot{s}_i on the left. Then

$$\dot{s}_i \dots \dot{s}_1 u \in \dot{s}_i Bz'_{i-1} B^-.$$

By using $\dot{s}_i Bz'_{i-1} B^- \subset B\dot{s}_i z'_{i-1} B^- \cup Bz'_{i-1} B^-$, we have that $z'_i = s_i z'_{i-1}$ or z'_{i-1} . Note that $u \in Bz_r B^-$ and so $z'_0 = \text{Id}$. In other words, by the definition of the Bruhat ordering, z'_r is a subexpression of w . This proves the only if direction.

We now consider the if direction. Let $z_0 = \text{Id}, z_1, \dots, z_r$ be a sequence of elements in ${}_F W$ representing the subexpression w' of w . We now construct elements inductively in the intersection:

$$B\dot{s}_q \dots \dot{s}_1 B \cap Bz_q B^-.$$

When $q = r$, we shall have the desired element in the intersection.

When $q = 0$, it is clear that $B \cap BB^- \neq \emptyset$ and we can pick a particular element in the intersection. Suppose we have constructed an element in

$$(3.3) \quad B\dot{s}_q \dots \dot{s}_1 B \cap B\dot{z}_q B^-.$$

We now consider

$$B\dot{s}_{q+1} \dots \dot{s}_1 B \cap B\dot{z}_{q+1} B^-$$

and consider the following cases:

- Case 1: $z_{q+1} \neq z_q$ i.e. $z_{q+1} = s_{q+1}z_q$. By multiplying an element in B on the left in (3.3), we may assume the intersection $B\dot{s}_q \dots \dot{s}_1 B \cap B\dot{z}_q B^-$ contains an element in $B\dot{z}_q B^-$ of the form:

$$\dot{z}_q b'$$

for some $b' \in B^-$. Then $\dot{s}_{q+1}\dot{z}_q b' \in B\dot{z}_{q+1} B^-$ and is also in $\dot{s}_{q+1}B\dot{z}_q B \subset B\dot{s}_{q+1}\dot{z}_q B$ ([Bo91, Proposition 21.22]), as desired.

- Case 2: $z_{q+1} = z_q$. Let α_{q+1} be the simple root associated to the simple reflection s_{q+1} . Let $v \in U_{(-\alpha_{q+1})} = \dot{s}_{q+1}U_{(\alpha_{q+1})}\dot{s}_{q+1}^{-1}$ be the non-identity element. Note that $v \in \dot{s}_{q+1}B\dot{s}_{q+1}^{-1} \subset B \cup B\dot{s}_{q+1}B$ by [Bo91, Theorem 21.15]. Then v can be written in the form

$$x' \dot{s}_{q+1} x$$

for some $x \in U_{\alpha_{q+1}}$ and $x' \in U$. Thus,

$$x' \dot{s}_{q+1} x \dot{z}_q b',$$

where $b' \in B^-$. Then, we have:

$$\dot{s}_{q+1} x \dot{z}_q b' \in \dot{s}_{q+1} B \dot{s}_q \dots \dot{s}_1 B \subset B \dot{s}_{q+1} \dots \dot{s}_1 B.$$

Hence, $v \dot{z}_q b' \in B \dot{s}_{q+1} \dots \dot{s}_1 B$.

On the other hand, the distinguishedness above implies that $z_q^{-1}(\alpha_q) > 0$, and so $\dot{z}_q^{-1} v \dot{z}_q \in U_{\alpha_q}^- \subset B^-$. Hence, the element $v \dot{z}_q b'$ is also in $B \dot{z}_{q+1} B^- = B \dot{z}_q B^-$. This proves this case. □

3.5. Consequences on other intersections. Let P, Q be standard parabolic subgroups of G . Let S and T be the corresponding subset of Φ determining P and Q respectively. Then

$$G = \bigsqcup_{w \in W_{S,T}} P \dot{w} Q.$$

Indeed, we can also use $W_{S,T}$ to parametrize double cosets of $P \setminus G/Q^-$:

Lemma 3.4. *We use the notations above. Let $w, w' \in W_{S,T}$. Then*

$$P \dot{w} Q^- = P \dot{w}' Q^- \iff w = w'.$$

Moreover,

$$G = \bigsqcup_{w \in W_{S,T}} P \dot{w} Q^-.$$

Proof. Let $\bar{Q} = \dot{w}_{0,T}^{-1}(Q^-)\dot{w}_{0,T}$. Since $w_{0,T}^{-1}$ sends all negative roots in the root subsystem generated by T to positive roots, \bar{Q} is still a standard parabolic subgroup of G . Moreover, the subset of Φ determining \bar{Q} is $\theta(T)$. Then, we have:

$$P \dot{w} Q^- = P \dot{w}_{0,T} \bar{Q} \dot{w}_{0,T}^{-1} = P \dot{v} \bar{Q} \dot{w}_{0,T}^{-1},$$

where $v = \Omega_{S,T}(w)$ ($\Omega_{S,T}$ is defined in (2.1)). Then the two assertions follow from Proposition 2.4. □

Lemma 3.5. *Let $\bar{Q} = \dot{w}_0 Q^- \dot{w}_0^{-1}$. Let $w \in W_{S,T}$. Let $u = \Omega_{S,T}(w)$. Then*

$$P\dot{w}Q = \bigsqcup_{x \in W(S), y \in W(T)} B\dot{x}\dot{w}\dot{y}B, \quad P\dot{w}Q^- = \bigsqcup_{x \in W(S), y \in W(T)} B\dot{x}\dot{w}\dot{y}B^-.$$

Proof. We only prove the second one and the first one is simpler. The second assertion follows from:

$$\begin{aligned} P\dot{w}Q^- &= P\dot{w}\dot{w}_0\bar{Q}\dot{w}_0^{-1} \\ &= \bigsqcup_{x \in W(S), y \in W(\theta(T))} B\dot{x}B\dot{w}\dot{w}_0B\dot{y}B\dot{w}_0^{-1} \\ &= \bigsqcup_{x \in W(S), y \in W(\theta(T))} B\dot{x}B\dot{w}\dot{w}_0\dot{y}B\dot{w}_0^{-1} \\ &= \bigsqcup_{x \in W(S), y \in W(\theta(T))} B\dot{x}\dot{w}\dot{w}_0\dot{y}B\dot{w}_0^{-1} \\ &= \bigsqcup_{x \in W(S), y \in W(\theta(T))} B\dot{x}\dot{w}\dot{w}_0\dot{y}\dot{w}_0^{-1}B^- \\ &= \bigsqcup_{x \in W(S), z \in W(T)} B\dot{x}\dot{w}\dot{z}B^-, \end{aligned}$$

where the third equality follows that for $v \in {}_F W$ and a simple reflection s ,

$$B\dot{v}\dot{s}B \subset B\dot{v}B\dot{s}B \subset B\dot{v}\dot{s}B \cup B\dot{v}B$$

and subexpression of an element of $W(\theta(T))$ is still in $W(\theta(T))$, and the fourth equality is similar. \square

Corollary 3.6. *We use the notations above. Let $w, w' \in W_{S,T}$. Then,*

$$P\dot{w}Q \cap P\dot{w}'Q^- \neq \emptyset,$$

if and only if $w' \leq w$.

Proof. We first prove the if direction. Suppose $w \leq w'$. Then $B\dot{w}B \cap B\dot{w}'B^- \neq \emptyset$ by Proposition 3.3 and so $P\dot{w}Q \cap P\dot{w}'Q^- \neq \emptyset$.

We now prove the only if direction. Suppose $P\dot{w}Q \cap P\dot{w}'Q^- \neq \emptyset$. Then, by $B\dot{x}\dot{w}\dot{y}B \cap B\dot{x}'\dot{w}'\dot{y}'B^- \neq \emptyset$ for some $x, x' \in W(S)$ and $y, y' \in W(T)$. Hence, $x'w'y' \leq xwy$ by Proposition 3.3. By Lemma 2.2, we have $w' \leq w$. \square

4. BERNSTEIN-CASSELMAN CANONICAL PAIRING

4.1. Stable operator. For a vector space V with an operator $\omega \in \text{End}(V)$, we say that ω is *stable* if ω is invertible on $\text{im } \omega$ and

$$V = \ker \omega \oplus \text{im } \omega.$$

Proposition 4.1. *Let V be a smooth representation of G and let $\omega \in \text{End}(V)$ be stable. Let V^\vee be a smooth dual of V . Define ω^\vee to be an operator on V^\vee given by $(\omega^\vee \cdot f)(v) = f(\omega \cdot v)$, where $f \in V^\vee$ and $v \in V$. Then ω^\vee is also stable.*

Proof. Let X be the subspace of functions in V^\vee which vanishes on $\ker(\omega)$. We claim that $\text{im}(\omega^\vee) = X$.

Let $f \in \text{im}(\omega^\vee)$. Then $f = \omega^\vee \cdot \tilde{f}$ for some $\tilde{f} \in V^\vee$. For $x \in \ker(\omega)$, $f(x) = (\omega^\vee \cdot \tilde{f})(x) = \tilde{f}(\omega \cdot x) = 0$. Hence, $f \in X$. This proves $\text{im}(\omega^\vee) \subset X$.

Let $f \in X$. For each $x \in \text{im}(\omega)$, we can write $x = \omega^{-1}.\tilde{x}$ for some $\tilde{x} \in \text{im}(\omega)$. Then, define $\tilde{f}(x) = f(\tilde{x})$ if $x \in \text{im}(\omega)$ and $\tilde{f}(x) = 0$ if $x \in \ker(\omega)$. This then implies that $(\omega^\vee.\tilde{f})(x) = f(x)$ for all x . Since f is smooth, \tilde{f} is also smooth and in V^\vee . This shows that $X \subset \text{im}(\omega')$. \square

4.2. Stabilization Theorem. For a representation π of G and an open compact subgroup K of G , we define

$$\pi_K(g).v = \int_G \text{ch}_{KgK}(h) \cdot \pi(h)v \, dh = \int_{KgK} \pi(h)v \, dh,$$

where ch_{KgK} is the characteristic function of KgK , and dh is the normalized Haar measure on G .

Let P be a standard parabolic subgroup of G . Let $M = M_P$, let $U = U_P$ and let $Z(M)$ be the center of M . An element $a \in Z(M)$ is said to be *strictly dominant* for M if for any two open compact subgroups H_1, H_2 in U , there exists an integer $n \geq 0$ such that $a^n H_1 a^{-n} \subset H_2$. The existence of such element is shown in [BK98, Lemma 6.14] and allows one to find an 'inverse' of the projection $\pi^K \rightarrow (\pi_U)^{K_M}$ as follows.

Theorem 4.2 (Jacquet's lemma, Bernstein's stabilization theorem). [BD84, Be92] *Let π be a smooth representation of G . Let P be a parabolic subgroup, let $M = M_P$ and let $U = U_P$. Let K be an open compact subgroup of G admitting an Iwahori decomposition. Let $K_M = M \cap K$. Then*

- (1) [Re10, Proposition IV.6.1] *For a strictly dominant element a for M , $\pi_K(a^i)$ is stable for large i . Let $W = \pi_K(a^i)V^K$. Then the natural surjection above induces a bijection from W to $V_U^{K_M}$.*
- (2) [Re10, Théorème IV.6.1] *The natural surjection $\pi^K \rightarrow (\pi_U)^{K_M}$ has a natural inverse, as vector spaces.*
- (3) [Re10, Pages 261, 262] *The projection sends $\ker \pi(a^i)$ to zero.*

4.3. Bernstein-Casselman canonical pairing. We now state a canonical pairing, which is due to Casselman [Ca93] for admissible representations. The stabilization theorem of Bernstein above removes the admissibility condition.

Theorem 4.3 (Bernstein, Casselman). (see [Re10, Proposition VI 9.6]) *Let π be a smooth representation of G . Let P be a standard parabolic subgroup of G . Let $M = M_P$, let $U = U_P$ and let $U^- = U_P^-$. Then the followings hold:*

- (1) *We have a functorial isomorphism $(\pi_U)^\vee \cong (\pi^\vee)_{U^-}$, as M -representations.*
- (2) *Moreover, the isomorphism in (1) is given explicitly as follows. For $x \in \pi_U$ and $f \in (\pi^\vee)_{U^-}$, let K be an open compact subgroup of G in a good position. By choosing sufficiently small K , we may assume $x \in \pi_U^{K_M}$ and $f \in (\pi^\vee)_{U^-}^{K_M}$. Let \tilde{x} and \tilde{f} be the lifts of $x \in \pi^K$ and $f \in (\pi^\vee)^K$ via the isomorphism in Theorem 4.2. Then the isomorphism is given by the pairing $\langle x, f \rangle_M := \tilde{f}(\tilde{x})$, which is independent of a choice of K .*
- (3) *Let a be a strictly dominant element for M . Let i be a sufficiently large integer such that $\pi_K(a^i)$ and $\pi_K^\vee(a^{-i})$ are stable. Let $p : \text{im } \pi_K(a^i) \rightarrow \pi_U$ be the isomorphism from the natural projection in Theorem 4.2. Similarly, we have an isomorphism $p^\vee : \text{im } \pi_K^\vee(a^{-i}) \rightarrow \pi_{U^-}$. Then, for any $\tilde{x} \in \pi^K$ and $\tilde{f} \in (\pi^\vee)^K$,*

$$\langle \pi_K(a^i)\tilde{x}, \tilde{f} \rangle = \langle \pi_U(a^i)p(\tilde{x}), p^\vee(\tilde{f}) \rangle_M.$$

- (4) Let $x \in \pi_U$ and let $f \in (\pi^\vee)_{U^-}$. Let \tilde{x} be the canonical lift of x and let f' be any lift of f in π^\vee . Then,

$$\langle x, f \rangle_M = \langle \tilde{x}, f' \rangle.$$

Similarly, let x' be any lift of x and let \tilde{f} be the canonical lift of f in π^\vee . Then,

$$\langle x, f \rangle_M = \langle x', \tilde{f} \rangle.$$

Definition 4.4. Let π be a smooth representation of G . Let P be a standard parabolic subgroup of G . Let $x \in \pi_{U_P}$. We shall say that an element \tilde{x} in π is a *lift* of x if the image of \tilde{x} under the natural projection map $\pi \rightarrow \pi_{U_P}$ is x .

Definition 4.5. [Ca93, Be92] Let π be a smooth representation of G with the underlying space V . Let V^\vee be the underlying space of π^\vee . Let P be a standard parabolic subgroup of G . Let $M = M_P$.

- (1) Let $x \in \pi_{U_P}$. Let K be an open compact subgroup of G in a good position such that $x \in \pi_{U_P}^{K_M}$. The unique element $\tilde{x} \in \pi_K(a^i)V^K$ such that \tilde{x} mapped to x in Theorem 4.2(3) is called the *canonical lift* of x . Note that the canonical lift is independent of K by Theorem 4.3(2).
- (2) Let $y \in (\pi^\vee)_{U_P^-}$. Let K be an open compact subgroup of G in a good position such that $y \in (\pi^\vee)_{U_P^-}^{K_M}$. The unique element $\tilde{y} \in \pi_K^\vee(a^{-1})(V^\vee)^K$ mapped to y in Theorem 4.2(3) is called the *canonical lift* of y .

5. CANONICAL LIFT ON SOME PARABOLICALLY INDUCED MODULES

5.1. Parabolic induction. Let P be a standard parabolic subgroup of G . Let σ be a smooth representation of M_P , inflated to a P -representation. Define

$$\text{Ind}_P^G \sigma := \{f : G \rightarrow \sigma \mid f \text{ is smooth, } f(pg) = p.f(g) \text{ for } p \in P\}.$$

We shall regard $\text{Ind}_P^G \sigma$ as a G -representation via the left translation i.e.

$$(g.f)(h) = f(hg)$$

for any $g, h \in G$.

For $f \in \text{Ind}_P^G \sigma$, define

$$\text{supp}(f) = \{x \in G : f(x) \neq 0\},$$

called the *support* of f .

5.2. Support of a function under actions of a strictly dominant element. We shall first show two lemmas on the group structure, continuing Section 3.

Lemma 5.1. *Let K be an open compact subgroup of G in a good position and let Q be a standard parabolic subgroup of G . Let $M = M_Q$, $U = U_Q$ and $U^- = U_Q^-$. Let $g \in G$ and let $m \in M$. Let a be a strictly dominant element for (G, M) and let i be a sufficiently large integer such that $a^i(K \cap U)a^{-i} \subset K \cap U \subset K$. Then $Ka^iK \subset QU^-$.*

Proof. Since K is in a good position, we have the Iwahori decomposition of K :

$$K_U \cdot K_M \cdot K_{U^-},$$

where $K_* = * \cap K$ for $* = M, U, U^-$. For a sufficiently large integer i , we have $a^i K_U a^{-i} \subset K_U$.

Let $g \in Ka^iK$. Write $g = k_1 a^i k_2$ for $k_1, k_2 \in K$. By the Iwahori decomposition, write $k_2 = umu'$ for $u \in K_U$, $m \in K_M$ and $u' \in K_{U^-}$. Then

$$g = k_1 a^i k_2 = k_1 a^i (umu') = k_1 (a^i u a^{-i}) a^i m u' \in KQ^-.$$

which follows from $a^i u a^{-i} \in K$ and $a^i m u' \in Q^-$.

On the other hand, by the Iwahori decomposition again, $K \subset K_U \cdot K_M \cdot K_{U^-} \subset Q Q^-$. Thus, we also have $g \in Q Q^- = Q U^-$. \square

Lemma 5.2. *We use the notations in Lemma 5.1. Let P be another standard parabolic subgroup of G . Let S and T be the subsets of Φ determining P and Q respectively. Let $u, w \in W_{S,T}$. Suppose $g k a^{-i} \in P \dot{w} m K$ for some $k \in K$. If $u \not\leq w$, then*

$$P g K \cap P \dot{u} Q^- = \emptyset.$$

Proof. The hypothesis gives that $g \in P \dot{w} m K a^i K$. Thus

$$P g K \subset P \dot{w} m K a^i K$$

Now, by Lemma 5.1, we then have:

$$P g K \subset P \dot{w} Q U_Q^-.$$

On the other hand, since $U_Q^- \subset Q^-$, we also have:

$$P \dot{w} Q U_Q^- \cap P \dot{u} Q^- \neq \emptyset \iff P \dot{w} Q \cap P \dot{u} Q^- \neq \emptyset.$$

Now the lemma follows from Corollary 3.6. \square

As seen in Theorem 4.3, computing the canonical pairing may involve computing the action of $\pi_K(a^i)$ for some strictly dominant a for some Levi subgroup. We shall need the following related lemma later.

Proposition 5.3. *Let K be an open compact subgroup of G in a good position. Let P, Q be standard parabolic subgroups of G . Let S, T be the subsets of Φ determining P and Q respectively. Let σ be a smooth representation of M_P . Let $w \in {}_F W$. Let*

$$\mathcal{O}_w^- = \bigsqcup_{w' \in {}_F W : w' \leq w} P \dot{w}' Q^-.$$

Let f be a function in $\text{Ind}_P^G \sigma$ such that $\text{supp}(f) \subset P \dot{w} m K$ for some $m \in M_Q$. For any strictly dominant element a for M_Q , $\pi_K(a^{-k})f$ has support in \mathcal{O}_w .

Proof. Let $\tilde{f} = \pi_K(a^{-i})f$. Let $g \in G$ such that $\tilde{f}(g) \neq 0$. By definition,

$$\tilde{f}(g) = \int_{K a^{-i} K} f(gh) dh$$

and so $g k a^{-i} \in P \dot{w} K$ for some $k \in K$. By Lemma 5.2, $g \in P \dot{w}' Q^-$ for some $w' \leq w$. This implies the proposition. \square

6. GEOMETRIC LEMMA AND ITS DUAL

6.1. Geometric lemma. We use the notations as in Section 3.1. Let P and Q be standard parabolic subgroups of G . Let S and T be the subsets of Φ determining P and Q respectively. Then, according to [Bo91, Section 21.16], there is a one-to-one correspondence between

$$W(S) \setminus W/W(T) \longleftrightarrow P \setminus G/Q.$$

Recall that $W_{S,T}$ is the set of minimal representatives in $W(S) \setminus W/W(T)$. Let $r = |W|$. We enumerate the elements u_1, \dots, u_r in $W_{S,T}$ such that $i < j$ implies $l(u_i) \leq l(u_j)$. This, in particular, gives that $u_i \leq u_j$ implies $i \leq j$.

Let

$$\mathcal{O}_i = \bigsqcup_{i' \geq i} P \dot{u}_{i'} Q.$$

Then, by [Bo91, Theorem 21.26], \mathcal{O}_i is open in G (in Zariski topology) and so is open in G as an l -group. Let $\mathcal{I}_i := \mathcal{I}_i(Q, \sigma)$ be the subspace of $\text{Ind}_P^{G^n} \sigma$ with functions supported in \mathcal{O}_i . By [BZ76, Proposition 1.8], we have:

$$\mathcal{I}_i / \mathcal{I}_{i-1} \cong C_c^\infty(P\dot{u}_i Q, \sigma),$$

where $C_c^\infty(P\dot{u}_i Q, \sigma)$ is the space of smooth σ -valued functions on $P\dot{u}_i Q$ satisfying $f(pg) = p.f(g)$ with the support to be compactly supported module P .

Let $\mathcal{J}_i := \mathcal{J}_i(Q, \sigma)$ be the projection from \mathcal{I}_i to π_{U_Q} . This then induces a filtration on $(\text{Ind}_P^G \sigma)_{U_Q}$:

$$0 \subseteq \mathcal{J}_r \subseteq \dots \subseteq \mathcal{J}_1 = (\text{Ind}_P^G \sigma)_{U_Q}.$$

As shown in [BZ77], there is an isomorphism between

$$(6.4) \quad C_c^\infty(P\dot{w}Q, \sigma)_{U_Q} \cong \text{ind}_{P^w \cap M_Q}^{M_Q} (\sigma_{P \cap \dot{w}U_Q \dot{w}^{-1}})^w,$$

where $P^w = \dot{w}^{-1}P\dot{w}$, given by:

$$(6.5) \quad f \mapsto \left(m \mapsto \int_{U^w} \text{pr}(f(\dot{w}um)) \, du \right),$$

where $m \in M_Q$ and pr is the projection from σ to $\sigma_{P \cap \dot{w}U_Q \dot{w}^{-1}}$.

6.2. Dual filtration I: smooth duals. We keep using notations in Section 6.1. This then induces a filtration on $((\text{Ind}_P^G \sigma)_{U_Q})^\vee$. More precisely, let $\mathcal{J}_i^\vee := \mathcal{J}_i^\vee(Q, \sigma)$ be the subspace of $((\text{Ind}_P^G \sigma)_{U_Q})^\vee$ precisely containing all functions from $(\text{Ind}_P^G \sigma)_{U_Q}$ to \mathbb{C} which vanish on $\mathcal{J}_{i+1}(Q, \sigma)$. Then we have a filtration on $((\text{Ind}_P^G \sigma)_{U_Q})^\vee$ given by:

$$0 \subseteq \mathcal{J}_1^\vee \subseteq \dots \subseteq \mathcal{J}_r^\vee = ((\text{Ind}_P^G \sigma)_{U_Q})^\vee$$

6.3. Dual filtration II: Bernstein-Casselman pairings. We keep using notations in Section 6.1. Let Q^- be the opposite parabolic subgroup of Q and let $\bar{Q} = \dot{w}_{0,T}^{-1}(Q^-)\dot{w}_{0,T}$, which is a standard parabolic subgroup of G . We now construct a filtration for $(\text{Ind}_P^G \sigma^\vee)_{U_{\bar{Q}}}$, as M_Q -representations.

To do so, by Lemma 3.4,

$$G = \bigsqcup_{u \in W_{S,T}} P\dot{u}Q^- = \bigsqcup_{u \in W_{S,T}} P\dot{u}\dot{w}_{0,T}\bar{Q}\dot{w}_{0,T}^{-1}.$$

We keep using the enumeration u_1, \dots, u_r of $W_{S,T}$ in Section 6.1.

Let

$$\mathcal{O}_i^- = \bigsqcup_{i' \leq i} P\dot{u}_{i'}Q^- = \bigsqcup_{i' \leq i} P\dot{v}_{i'}\bar{Q}\dot{w}_T^{-1},$$

where $v_{i'} = \Omega_{S,T}(u_{i'})$. Now, by Proposition 2.4, \mathcal{O}_i^- is open in G .

We now define a filtration for $\text{Ind}_P^G \sigma^\vee$. Let $\mathcal{I}_i^- := \mathcal{I}_i^-(Q^-, \sigma^\vee)$ be the space of functions in $\text{Ind}_P^G(\sigma^\vee)$ whose support lies in \mathcal{O}_i^- . Then we have a filtration on $\text{Ind}_P^G \sigma^\vee$ given by:

$$0 \subset \mathcal{I}_1^- \subset \dots \subset \mathcal{I}_r^- = \text{Ind}_P^G(\sigma^\vee).$$

Let pr be the projection from $\text{Ind}_P^G(\sigma^\vee)$ to $(\text{Ind}_P^G \sigma^\vee)_{U_{\bar{Q}}}$. Let $\mathcal{J}_i^- := \mathcal{J}_i(Q^-, \sigma^\vee)$ be the image of pr on $\mathcal{I}_i^-(\sigma^\vee)$. This gives another filtration for $(\text{Ind}_P^G \sigma^\vee)_{U_{\bar{Q}}}$:

$$0 \subseteq \mathcal{J}_1^- \subseteq \dots \subseteq \mathcal{J}_r^- = (\text{Ind}_P^G \sigma^\vee)_{U_{\bar{Q}}}.$$

7. MATCHING FILTRATIONS

Let P be a standard parabolic subgroup of G . Let σ be a smooth representation of M_P . Define a pairing $\langle \cdot, \cdot \rangle_\sigma : \sigma \times \sigma^\vee \rightarrow \mathbb{C}$ given by:

$$\langle x, f \rangle_\sigma = f(x)$$

for $x \in \sigma$ and $f \in \sigma^\vee$.

There is a natural isomorphism [BZ76]:

$$(\text{Ind}_P^G \sigma)^\vee \cong \text{Ind}_P^G(\sigma^\vee)$$

determined by the non-degenerate pairing on $\text{Ind}_P^G \sigma \times \text{Ind}_P^G(\sigma^\vee)$ given by:

$$\langle f, f' \rangle = \int_{P \backslash G} \langle f(g), f'(g) \rangle_\sigma dg,$$

where dg is the normalized Haar measure on $P \backslash G$.

On the other hand, via Theorem 4.3(1), we have:

$$((\text{Ind}_P^G \sigma)_{U_Q})^\vee \cong ((\text{Ind}_P^G \sigma)^\vee)_{U_Q^-}.$$

Combining two isomorphisms, we have:

$$((\text{Ind}_P^G \sigma)_{U_Q})^\vee \cong (\text{Ind}_P^G(\sigma^\vee))_{U_Q^-}$$

Hence, we have a non-degenerate M_Q -equivariant pairing $\langle \cdot, \cdot \rangle_{M_Q} : (\text{Ind}_P^G \sigma)_{U_Q} \times (\text{Ind}_P^G \sigma^\vee)_{U_Q^-}$ given by:

$$\langle f, f' \rangle_{M_Q} = \langle \tilde{f}, \tilde{f}' \rangle,$$

where \tilde{f} is a lift of f and \tilde{f}' is the canonical lift of f' .

The goal of this section is to show that the filtrations in Sections 6.2 and 6.3 coincide.

7.1. Canonical lift under a filtration. Let $\pi = \text{Ind}_P^G(\sigma^\vee)$. We first construct a canonical lift of an element in $\pi_{U_Q^-}$. Let $f \in \text{Ind}_P^G(\sigma^\vee)$. Fix an open compact subgroup K of G in a good position such that $f \in (\text{Ind}_P^G \sigma)^K$. Let pr be the natural projection from $\text{Ind}_P^G(\sigma^\vee)$ to $(\text{Ind}_P^G \sigma^\vee)_{U_Q^-}$.

Let a be a strictly dominant element for M_P . Let k be a sufficiently large integer such that the action of a^k on $\text{Ind}_P^G \sigma$ is stable and so the action of a^{-k} on $(\text{Ind}_P^G \sigma)^\vee$ is also stable (see Proposition 4.1).

Now one defines the filtrations $\mathcal{I}_i = \mathcal{I}_i(Q, \sigma)$, $\mathcal{J}_i = \mathcal{J}_i(Q, \sigma)$, $\mathcal{I}_i^- = \mathcal{I}_i^-(Q^-, \sigma^\vee)$, $\mathcal{J}_i^- = \mathcal{J}_i^-(Q^-, \sigma^\vee)$, $\mathcal{J}_i^\vee = \mathcal{J}_i^\vee(Q, \sigma)$ as in Section 6.

Lemma 7.1. *Let $f \in \mathcal{I}_i^-$. Then there exists $\tilde{f} \in \mathcal{J}_i^-$ such that $\text{pr}(\tilde{f}) = f$.*

Proof. This follows from the definition of \mathcal{I}_i^- and \mathcal{J}_i^- . □

We shall need the following useful formula:

Lemma 7.2. [Be87, Proposition 5.2] $((\pi^\vee)_{U_Q^-})(a^{-k}) \circ \text{pr} = \text{pr} \circ \pi_K^\vee(a^{-k})$.

Proof. By the Iwahori decomposition of K ,

$$K = (U_Q^- \cap K) \cdot (M_Q \cap K) \cdot (U_Q \cap K).$$

Note that $a^k(U_Q \cap K)a^{-k} \subset K$ by the strict dominance of a . We also have $a^k(M_Q \cap K)a^{-k} \subset K$ and so:

$$\pi_K^\vee(a^{-k}) = e_{U_Q^- \cap K} \circ \pi^\vee(a^{-k}) \circ e_K,$$

where $e_{K'}(f)(g) = \int_{h \in K'} f(gh) dh$ for $K' = U_{\bar{Q}} \cap K$ or $M_P \cap K$. (Here dh is the normalized Haar measure.) We also remark that $\pi^\vee(a^{-k})$ is simply the group action of a^{-1} on π^\vee . Since $e_{U_{\bar{Q}} \cap K}$ coincides with taking the $U_{\bar{Q}} \cap K$ -coinvariants and pr coincides with taking $U_{\bar{Q}}$ -coinvariants, we have that $\text{pr} \circ e_{U_{\bar{Q}} \cap K} = \text{pr}$. Now the lemma follows by the commutation between pr and $\pi(a^{-k})$. \square

Proposition 7.3. *Let $f \in \mathcal{J}_i^-$. Then the canonical lift of f is in \mathcal{I}_i^- .*

Proof. Let $f' = a^k \cdot f$. By Lemma 7.1, there exists $\tilde{f}' \in \mathcal{J}_i^-$ such that $\text{pr}(\tilde{f}') = f'$. By Lemma 7.2, $f = \text{pr}(\pi_K(a^{-k})\tilde{f}')$. Thus $\pi_K(a^{-k})\tilde{f}'$ is the canonical lift of f by Proposition 4.1 and the definition. Thus, it remains to see that $\pi_K(a^{-k})\tilde{f}'$ is in \mathcal{I}_i^- , but this follows from Proposition 5.3. \square

7.2. Constructing a lift for \mathcal{J}_{i+1} .

Lemma 7.4. *Let $f \in \mathcal{J}_i$. There exists a lift of f lying in \mathcal{I}_i .*

Proof. This again follows from definitions. \square

Lemma 7.5. *Let $\tilde{f} \in \mathcal{I}_i$ such that $\text{supp}(\tilde{f}) \subset PgK$ for some $g \in Q$ and some open compact subgroup K . Then $\text{supp}(u^{-1} \cdot \tilde{f}) \subset PgKu^{-1}$.*

Proof. This also follows from definitions. \square

We now have the following key technical lemma for constructing a lift of an element in \mathcal{I}_i .

Lemma 7.6. *Let $f \in \mathcal{J}_i$. There exists $\tilde{f} \in \mathcal{I}_i$ and an open compact subgroup K of G such that $\text{pr}(\tilde{f}) = f$ and*

$$\text{supp}(\tilde{f}) \subset \bigsqcup_{i' \geq i} P\dot{w}_{i'} M_Q K$$

and

$$\mathcal{O}_i^- \cap \text{supp}(\tilde{f}) = \emptyset.$$

Proof. Let \tilde{f} be a lift of f in \mathcal{I}_i by Lemma 7.4 and so $\text{supp}(\tilde{f}) \subset \mathcal{O}_{i+1}$.

For each $g \in \text{supp}(\tilde{f}) \subset \mathcal{O}_{i+1}$, let K'_g be an open compact subgroup of G such that

$$\tilde{f}|_{gK'_g}$$

is a constant function and $gK'_g \cap \mathcal{O}_i^- = \emptyset$ (and so $PgK'_g \cap \mathcal{O}_i^- = \emptyset$). The last property is guaranteed by the Hausdorffness property of G .

Write $g = pw_jq$ for some $p \in P$, $j \leq i$ and $q \in Q$. Write $q = mu$ for $m \in M_Q$ and $u \in U_Q$. We can also choose an open compact subgroup of the form $uK''_g u^{-1}$ such that

$$(7.6) \quad PgK''_g u^{-1} = P\dot{w}_j m (uK''_g u^{-1}) \cap \mathcal{O}_i^- = \emptyset.$$

Let $K_g = K''_g \cap K'_g$.

Now by using the compactness of $P \setminus G$, we have finitely many g_1, \dots, g_r such that $g_1 K_{g_1}, \dots, g_r K_{g_r}$ cover $\text{supp}(\tilde{f})$. We can now let $K = K_{g_1} \cap \dots \cap K_{g_r}$. Then we partition $\text{supp}(f)$ by

$$Pg_1 k_{1,1} K, g_1 k_{1,2} K, \dots, g_1 k_{1,s_x} K, g_2 k_{2,1} K, g_2 k_{2,2} K, \dots, g_2 k_{2,s_2} K, \dots$$

for each $k_{x,y} \in K_{g_x}$ such that all those cosets are disjoint.

Thus, we have

$$\tilde{f} = \sum_{x=1}^r \sum_{y=1}^{s_x} c_{x,y},$$

where $c_{x,y}$ is a function uniquely determined by the support exactly equal to $Pg_x k_{x,y} K$.

Now by the definition of the Jacquet functor, we can replace each function $c_{x,y}$ by $u_x^{-1} \cdot c_{x,y}$, where u_x is the element that $g_x = pw_j m u_x$ for $p \in P$, $m \in M_Q$ and $u_x \in U_Q$.

Let such function by \bar{f} . Then \bar{f} is also a lift of f . It remains to see

$$\text{supp}(u_x^{-1} \cdot c_{x,y}) \cap \mathcal{O}_i^- = \emptyset.$$

To this end, from Lemma 7.5,

$$\text{supp}(u_x^{-1} c_{x,y}) = Pg_x(k_{x,y} K) u_x^{-1}$$

and so

$$\text{supp}(u_x^{-1} c_{x,y}) \subset Pg_x k_{x,y} K u_x^{-1} \subset Pg_x K_{g_x} u_x^{-1}.$$

Hence, by (7.6),

$$\text{supp}(u_x^{-1} c_{x,y}) \cap \mathcal{O}_i^- = \emptyset$$

as desired. \square

7.3. Coincidence of filtrations.

Theorem 7.7. *Let P, Q be standard parabolic subgroups of G . Let σ be a smooth representation of M_P . We write $\mathcal{I}_i^\vee(Q, \sigma)$ for the filtration on $((\text{Ind}_P^G \sigma)_{U_Q})^\vee$ as in Section 6.1 with a fixed ordering on $W_{S,T}$. Write $\mathcal{I}_i^-(Q^-, \sigma^\vee)$ for the filtration on $(\text{Ind}_P^G \sigma^\vee)_{U_Q^-}$ as in Section 6.3. Then the two filtrations coincide via the Bernstein-Casselman isomorphism in Theorem 4.3(1).*

The proof of Theorem 7.7 is divided into two steps:

(1) We first show that $\langle \cdot, \cdot \rangle_{M_Q}$ is identically zero on $\mathcal{J}_{i+1} \times \mathcal{J}_i^-$.

(2) We next show the non-degeneracy of $\langle \cdot, \cdot \rangle_{M_Q}$ when restricted to $(\mathcal{J}_i / \mathcal{J}_{i+1}) \times \mathcal{J}_i^-$.

We first prove Step 1.

Lemma 7.8. *For $f' \in \mathcal{J}_{i+1}^-$ and $\tilde{f} \in \mathcal{I}_i$, there exists a lift \tilde{f}' of f' such that $\langle \tilde{f}, \tilde{f}' \rangle = 0$.*

Proof. By Lemma 7.6, we can find a lift \tilde{f} of f such that $\text{supp}(\tilde{f}) \cap \text{supp}(\tilde{f}') = \emptyset$. Now, we have

$$\begin{aligned} \langle \tilde{f}, \tilde{f}' \rangle &= \int_{P \backslash G} \langle \tilde{f}(g), \tilde{f}'(g) \rangle \sigma dg \\ &= 0 \end{aligned}$$

where dg is a Haar measure on $P \backslash G$, the first equality follows from Theorem 4.3(4), the second equality follows from the definition of the pairing [BZ76] and the third equality follows from the disjointness of support. \square

Lemma 7.9. *Let $f \in \mathcal{J}_{i+1}(Q, \sigma)$ and let $f' \in \mathcal{J}_i^-(Q, \sigma^\vee)$. Then $\langle f, f' \rangle_{M_Q} = 0$.*

Proof. This follows from Theorem 4.3(4), Proposition 7.3 and Lemma 7.8. \square

We now prove Step 2.

Lemma 7.10. *Let $\tilde{f}' \in \mathcal{J}_i^-$, which does not lie in \mathcal{J}_{i-1}^- . Then there exists $\tilde{f} \in \mathcal{J}_i$ such that $\langle \tilde{f}, \tilde{f}' \rangle \neq 0$.*

Proof. Since $\tilde{f}' \in \mathcal{J}_i^- - \mathcal{J}_{i-1}^-$, we have $\tilde{f}'(\dot{w}_j q) \neq 0$ for some $q \in Q$. We find a sufficiently small open compact subgroup K' such that \tilde{f} is constant in $\dot{w}_j q K'$.

Now we let $y = \tilde{f}(\dot{w}_j q)$ and let x be an element in σ^\vee such that $\langle x, y \rangle_\sigma = 1$. Let \tilde{f} be the function in $\text{Ind}_P^G \sigma$ with $\text{supp}(\tilde{f}) = P\dot{w}_j q K'$ and $f(\dot{w}_j q K') = x$. Then

$$\langle \tilde{f}, \tilde{f}' \rangle = \text{vol}(P \setminus P\dot{w}_j q K') \neq 0.$$

□

Proof of Theorem 7.7. By Lemma 7.8, $\mathcal{I}_i^- \subset \mathcal{I}_i^\vee$. We now prove another inclusion. Let $f' \in \mathcal{I}_i^\vee$. If $f' \notin \mathcal{I}_i^-$, then $f' \in \mathcal{I}_j^-$ for some $j > i$ and does not lie in \mathcal{I}_i^- . Let \tilde{f}' be the canonical lift of f' . By Proposition 7.3, $\tilde{f}' \in \mathcal{J}_j^-$. Since \tilde{f}' does not lie in \mathcal{I}_i^- , its canonical lift also does not lie in \mathcal{J}_i^- . By Lemma 7.10, there exists $\tilde{f} \in \mathcal{J}_k$ for $j \geq k > i$ such that $\langle \tilde{f}, \tilde{f}' \rangle \neq 0$. Thus, $\langle f, f' \rangle_{M_Q} = \langle \tilde{f}, \tilde{f}' \rangle \neq 0$, where f is the projection of \tilde{f} from $\text{Ind}_P^G \sigma$ onto $(\text{Ind}_P^G \sigma)_{U_Q}$. By definition, f is also in $\mathcal{I}_k \subset \mathcal{I}_i$. This contradicts that $f' \in \mathcal{I}_i^\vee$. Hence, $f' \in \mathcal{I}_i^-$ as desired. □

Remark 7.11. Let p be the residual characteristics of F . Let R be a noetherian $\mathbb{Z}[\frac{1}{p}]$ -ring. A recent paper of Dat-Helm-Kurinczuk-Moss [DHKM23] establishes the second adjointness theorem for the category of smooth R -representations. Indeed, they also established the stabilization theorem in [DHKM23, Corollary 4.9]. One may expect that such analogue can also be generalized to such category.

8. APPLICATIONS

8.1. Quotients of induced representations.

Corollary 8.1. *Let P be a standard parabolic subgroup of G . Let σ be a smooth representation of M_P . Let π be a non-zero quotient of $\text{Ind}_P^G \sigma$. Let $\lambda = C_c^\infty(PQ^-, \sigma)$ be the subspace of $\text{Ind}_P^G \sigma$ containing all functions supported in PQ^- . Then*

- (1) *the natural composition $\lambda_{U_Q^-} \hookrightarrow (\text{Ind}_P^G \sigma)_{U_Q^-} \twoheadrightarrow \pi_{U_Q^-}$ is non-zero;*
- (2) *if $P = Q$, then the map in (a) takes the form:*

$$\sigma \hookrightarrow (\text{Ind}_P^G \sigma)_{U_Q^-} \twoheadrightarrow \pi_{U_Q^-}.$$

Proof. We first prove (1). By taking the dual on the surjection $\text{Ind}_P^G \sigma \twoheadrightarrow \pi$, we have an embedding:

$$\pi^\vee \hookrightarrow \text{Ind}_P^G \sigma^\vee$$

Taking the Jacquet functor gives:

$$(8.7) \quad (\pi^\vee)_{U_Q} \hookrightarrow (\text{Ind}_P^G \sigma^\vee)_{U_Q} \twoheadrightarrow C_c^\infty(PQ, \sigma^\vee)_{U_Q}$$

which is non-zero. (The last module does not necessarily give a M_Q -representation, but makes sense as a space.)

Since the map in Theorem 4.3(1) is functorial, the first injection in (8.7) takes the following form under the isomorphism of Theorem 4.3(1):

$$(\pi_{U_Q^-})^\vee \hookrightarrow ((\text{Ind}_P^G \sigma)_{U_Q^-})^\vee$$

and is obtained by first taking the Jacquet functor with respect to Q^- on the projection $\text{Ind}_P^G \sigma$ and then taking the smooth dual.

By Theorem 7.7, the second map coincides with the embedding $\lambda_{U_Q^-} \hookrightarrow (\text{Ind}_P^G \sigma)_{U_Q^-}$. Thus (8.7) is obtained by taking the smooth dual on $\lambda_{U_Q^-} \hookrightarrow (\text{Ind}_P^G \sigma)_{U_Q^-} \twoheadrightarrow \pi_{U_Q^-}$. Then we must have the non-zero composition in the first assertion.

We now prove (2). Note that $C_c^\infty(PP^-, \sigma)$ is naturally isomorphic to the space of smooth functions from U_P^- to σ . Taking the Jacquet functor by U_P^- is isomorphic to σ . One checks the M_P -action is isomorphic to σ . \square

8.2. The adjoint map.

Lemma 8.2. *Let π_1, π_2 be smooth representations of G . Let $f : \pi_1 \rightarrow \pi_2^\vee$. Let $f' : \pi_2 \rightarrow \pi_1^\vee$ be the map given by $f'(x)(y) = f(y)(x)$ for $x \in \pi_2$ and $y \in \pi_1$. Let $f^\vee : (\pi_2^\vee)^\vee \rightarrow \pi_1^\vee$ given by $f^\vee(h)(y) = h(f(y))$ for $h \in (\pi_2^\vee)^\vee$ and $y \in \pi_1$. Let $\iota : \pi_2 \rightarrow (\pi_2^\vee)^\vee$ be the embedding given by: for $x \in \pi_2$ and $f \in \pi_2^\vee$, $\iota(x)(f) = f(x)$. Then $f' = f^\vee \circ \iota$.*

The above lemma is straightforward and we omit the details.

Lemma 8.3. *Let σ be a smooth representation of M_P . Let $f \in \text{Hom}_G(\pi, \text{Ind}_P^G \sigma)$. Via Frobenius reciprocity, the corresponding map in $\text{Hom}_{M_P}(\pi_{U_P}, \sigma)$ is given by the composition:*

$$\pi_{U_P} \xrightarrow{\tilde{f}} (\text{Ind}_P^G \sigma)_{U_P} \twoheadrightarrow C_c^\infty(P, \sigma) \cong \sigma,$$

where \tilde{f} is descended from f .

Proof. Recall that the adjoint map F of f is given by

$$F(x) = f(x)(1),$$

where 1 is the identity element in G [BZ76]. Note that the projection $\text{Ind}_P^G \sigma$ to $C_c^\infty(P, \sigma)$ coincides with the evaluation map of a function of $\text{Ind}_P^G \sigma$ at the identity. Thus the two maps coincide. \square

We now recall the relation between the second adjointness theorem and the Bernstein-Casselman pairing. Let π be a smooth representation of G and let σ be a smooth representation of M_P . It follows from [Be92, Page 63] that

$$\begin{aligned} \text{Hom}_G(\text{Ind}_P^G \sigma, \pi^\vee) &\cong \text{Hom}_G(\pi, (\text{Ind}_P^G \sigma)^\vee) \\ &\cong \text{Hom}_G(\pi, \text{Ind}_P^G(\sigma^\vee)) \\ &\cong \text{Hom}_G(\pi_{U_P}, \sigma^\vee) \\ &\cong \text{Hom}_G(\sigma, (\pi_{U_P})^\vee) \\ &\cong \text{Hom}_G(\sigma, (\pi^\vee)_{U_P^-}), \end{aligned}$$

where the first isomorphism follows by taking dual maps, the second isomorphism follows from [BZ76], the third isomorphism follows from Frobenius reciprocity, the fourth isomorphism follows from taking dual maps again and the last isomorphism follows from Theorem 4.3(1). The above isomorphism determines the second adjointness theorem.

Corollary 8.4. *(c.f. [BK15]) Let π be a smooth representation of G . Let P be a standard parabolic subgroup of G and let σ be a smooth representation of M_P . Let $f \in \text{Hom}_G(\text{Ind}_P^G \sigma, \pi)$. Let $U^- = U_P^-$. Then, the map $\sigma \twoheadrightarrow \pi_{U^-}$ arising from the adjointness coincides from taking the Jacquet functor U^- on the following composition of maps:*

$$C_c^\infty(PP^-, \sigma) \hookrightarrow \text{Ind}_P^G \sigma \twoheadrightarrow \pi.$$

with the isomorphism $C_c^\infty(PP^-, \sigma)_{U^-} \cong \sigma$.

Proof. Let $U = U_P$. We begin with a non-zero map f in $\text{Hom}_G(\text{Ind}_P^G \sigma, \pi^\vee)$. Then let f^\vee be the corresponding dual map in $\text{Hom}_G(\pi, (\text{Ind}_P^G \sigma)^\vee)$. By Lemma 8.3, the adjoint map of f in $\text{Hom}_G(\pi_U, \sigma)$ is given by:

$$\pi_U \hookrightarrow (\text{Ind}_P^G \sigma^\vee)_U \twoheadrightarrow C_c^\infty(P, \sigma^\vee)_U \cong \sigma,$$

where the maps are natural maps in Lemma 8.3.

Taking the dual and the Bernstein-Casselman isomorphism gives the following non-zero composition:

$$(8.8) \quad (\pi^\vee)_{U^-} \leftarrow ((\text{Ind}_P^G \sigma^\vee)^\vee)_{U^-} \leftarrow C_c^\infty(PP^-, \sigma^\vee)_{U^-}^\vee \cong (\sigma^\vee)^\vee,$$

where the first map follows from the functorial isomorphism of Bernstein-Casselman and the second map follows from our matching of filtrations in Theorem 7.7.

In order to obtain the adjoint map in $\text{Hom}_{M_P}(\sigma, (\pi^\vee)_{U^-})$, we then compose (8.8) with the natural embedding $\sigma \hookrightarrow \sigma^{\vee\vee}$ by Lemma 8.2. By functoriality, we also have the following commutative diagram:

$$\begin{array}{ccc} (\pi^\vee)_{U^-} & \xleftarrow{h} & (\text{Ind}_P^G(\sigma^{\vee\vee}))_{U^-} \xrightarrow{\quad} C_c^\infty(PP^-, \sigma^{\vee\vee})_{U^-} \cong \sigma^{\vee\vee}, \\ & & \uparrow s \qquad \qquad \qquad \uparrow \\ (\text{Ind}_P^G \sigma)_{U^-} & \xleftarrow{i} & C_c^\infty(PP^-, \sigma)_{U^-} \cong \sigma \end{array}$$

where h is the natural map obtained from f via taking dual, Jacquet functors and the Bernstein-Casselman pairing, and i is the map induced from the embedding as a subspace.

The natural map from $\text{Ind}_P^G \sigma$ to $\text{Ind}_P^G(\sigma^{\vee\vee})$ coincides with the natural embedding $\text{Ind}_P^G \sigma$ to $(\text{Ind}_P^G \sigma)^{\vee\vee}$ under the isomorphism between $\text{Ind}_P^G(\sigma^{\vee\vee})$ and $(\text{Ind}_P^G \sigma)^{\vee\vee}$. Thus, by Lemma 8.3, $h \circ s$ is obtained from f via taking the Jacquet functor. Using the composition $h \circ s \circ i$, we obtain the desired description of the adjoint map of f in the case that f is in $\text{Hom}_G(\text{Ind}_P^G \sigma, \pi^\vee)$.

We now consider general π (not necessarily of the form π^\vee). Let $f \in \text{Hom}_G(\text{Ind}_P^G \sigma, \pi)$. One considers the following embedding:

$$0 \rightarrow \pi \rightarrow (\pi^\vee)^\vee.$$

Hence, we have an injection from $\text{Hom}_G(\text{Ind}_P^G \sigma, \pi)$ to $\text{Hom}_G(\text{Ind}_P^G \sigma, (\pi^\vee)^\vee)$. Hence, f descends to a map, denoted by \bar{f} , in $\text{Hom}_G(\text{Ind}_P^G \sigma, \pi)$. By functoriality of the second adjointness, we then have the following commutativity diagram

$$\begin{array}{ccc} \pi_{U^-} & \xleftarrow{\bar{f}} & \sigma \\ \downarrow & & \parallel \\ (\pi^{\vee\vee})_{U^-} & \xleftarrow{\quad} & (\text{Ind}_P^G \sigma)_{U^-} \xleftarrow{\quad} C_c^\infty(PP^-, \sigma)_{U^-} \end{array},$$

where the bottom horizontal maps are the maps whose composition gives the adjoint map of \bar{f} shown above, the left vertical map is induced from the natural embedding π to $\pi^{\vee\vee}$. This then gives the desired description of the map \bar{f} in general case. \square

8.3. Applications on derivatives for GL. Let $G_n = \text{GL}_n(F)$ be the general linear group over a non-Archimedean local field F . Let P_{n_1, n_2} be the parabolic subgroup containing $\text{diag}(g_1, g_2)$ for $g_1 \in G_{n_1}$ and $g_2 \in G_{n_2}$. For smooth representations π_1 and π_2 of G_{n_1} and G_{n_2} respectively, we shall denote by $\pi_1 \times \pi_2$ the normalized parabolic induction $\text{Ind}_{P_{n_1, n_2}}^{G_{n_1+n_2}} \pi_1 \boxtimes \pi_2$.

Corollary 8.5. *Let τ be an irreducible representation of G_n and let σ be a smooth representation of G_k . Let π be an irreducible quotient of $\sigma \times \tau$. Suppose the level of π is equal to the level of τ . Then ${}^- \pi$ is a quotient of $\sigma \times {}^- \tau$.*

Proof. Let l be the level of π . Let U^- be the opposite unipotent radical in $P_{k,n}$. It follows from Corollary 8.1 that we have a non-zero map $\sigma \boxtimes \tau \rightarrow \pi_{U^-}$ from the second adjointness theorem on the quotient map from $\sigma \times \tau$ to π , and so we have a non-zero map

$$\sigma \boxtimes \tau \rightarrow \sigma' \boxtimes \tau \hookrightarrow \pi_{U^-}$$

for some quotient σ' of σ . On the factor G_n in $G_k \times G_n$, we consider the unipotent subgroup R containing the elements:

$$(8.9) \quad \begin{pmatrix} I_{n-l} & \\ & u \end{pmatrix} \subset G_n,$$

where $m \in \text{Mat}_{l,n-l}$ and $u \in U_l^-$. Here U_l^- is the subgroup containing all unipotent lower triangular matrices. Let $\psi : R \rightarrow \mathbb{C}$ be given by $\psi\left(\begin{pmatrix} g & \\ m & u \end{pmatrix}\right) = \bar{\psi}(u)$, where $\bar{\psi}$ is a non-degenerate character on U_l^- . Thus, by exactness, we obtain a non-zero map:

$$\sigma \boxtimes \tau_{R,\psi} \rightarrow \sigma' \boxtimes \tau_{R,\psi} \hookrightarrow (\pi_{N^-})_{R,\psi}.$$

Now, one defines similarly R' in G_{n+k} by replacing I_{n-l} with I_{n+k-l} and replacing $m \in \text{Mat}_{l,n-l}$ with elements in $\text{Mat}_{l,n+k-l}$. We similarly have a character ψ' on R' .

Now let \tilde{R} (resp. \tilde{R}') be the subgroup containing matrices of the form

$$\begin{pmatrix} g_1 & & & \\ * & g_2 & & \\ * & * & & \\ * & * & & u \end{pmatrix} \quad (\text{resp.} \quad \begin{pmatrix} g & \\ * & u \end{pmatrix}),$$

where $g_1 \in G_k$, $g_2 \in G_{n-l}$ and $u \in U_l^-$ (resp. $g \in G_{n+k-l}$ and $u \in U_l^-$). By taking Jacquet functors in stage, we have a non-zero map

$$C_c^\infty(P\tilde{R}, \sigma \boxtimes \tau)_{R',\psi'} \hookrightarrow (\sigma \times \tau)_{R',\psi'} \rightarrow \pi_{R',\psi}.$$

Since we have a surjection from $C_c^\infty(P\tilde{R}, \sigma \boxtimes \tau)$ to $C_c^\infty(P\tilde{R}', \sigma \boxtimes \tau)$ and $\sigma \times \tau$, we then also have a non-zero map:

$$C_c^\infty(P\tilde{R}', \sigma \boxtimes \tau)_{R',\psi'} \rightarrow \pi_{R',\psi'}.$$

But, the first space is isomorphic to $\sigma \times {}^- \tau$ (here we have to use that the level of τ is also l) and the second one is isomorphic to ${}^- \pi$. This implies the corollary. \square

The above example can be regarded as an instance on how the two operations – Bernstein-Zelevinsky derivatives and parabolic inductions – commute. More extensive studies and applications appear in [Ch22+b] and [Ch22+].

8.4. Connection to commutative triples and its dual version.

Definition 8.6. A *segment* is a combinatorial datum of the form $[a, b]_\rho$ for some cuspidal representation ρ of some G_k and for some $a, b \in \mathbb{Z}$ with $b - a \in \mathbb{Z}_{\geq 0}$.

For each segment Δ , we associate an essentially square-integrable representation denoted by $\text{St}(\Delta)$ [Ze80] (see [Ch22+c] for more details of the notion). We now introduce a notion of derivatives, which can be seen as a computable replacement for some structure of Bernstein-Zelevinsky derivatives (see [Ch22+c]).

Definition 8.7. Let Δ be a segment. Let π be an irreducible representation of G_n .

- Define $D_\Delta(\pi)$ to be the unique (if exists) irreducible representation of some G_k such that

$$\pi \hookrightarrow D_\Delta(\pi) \times \text{St}(\Delta).$$

Set $D_\Delta(\pi) = 0$ if such module does not exist.

- Define $I_\Delta(\pi)$ to be the unique irreducible representation of $\text{St}(\Delta) \times \pi$.

Definition 8.8. [Ch22+b] Let ρ_1 and ρ_2 be cuspidal representations of G_{n_1} and G_{n_2} respectively. Let $\Delta_1 = [a_1, b_1]_{\rho_1}$, $\Delta_2 = [a_2, b_2]_{\rho_2}$ be segments. Let $l_k = (b_k - a_k + 1)n_k$ ($k = 1, 2$). Let π be an irreducible representation of G_n . For $k = 1, 2$, let P_k be the parabolic subgroup of G_{l_2+n} containing upper triangular matrices a let N_k be the unipotent radicals of P_k respectively. We say that $(\Delta_1, \Delta_2, \pi)$ is a *pre-commutative triple* if the following composition:

$$D_{\Delta_1} \circ I_{\Delta_2}(\pi) \boxtimes \text{St}(\Delta_1) \hookrightarrow (I_{\Delta_2}(\pi))_{N_1} \hookrightarrow (\text{St}(\Delta_2) \times \pi)_{N_1} \twoheadrightarrow C_c^\infty(P_{n_2}P_{n_1}, \text{St}(\Delta_2) \boxtimes \pi)_{N_1},$$

where

- the first map is the unique embedding in Definition 8.7;
- the second map is induced from the unique embedding $I_{\Delta_2}(\pi) \hookrightarrow \text{St}(\Delta_2) \times \pi$;
- the third map is induced from the natural surjection from $\text{St}(\Delta_2) \times \pi$ to $C_c^\infty(P_{n_2}P_{n_1}, \text{St}(\Delta_2) \boxtimes \pi)$ in the geometric lemma.

We now define a dual version:

Definition 8.9. Let Δ be a segment. Let π be an irreducible representation of G_n .

- Define $D_\Delta^\vee(\pi)$ to be the unique irreducible representation such that

$$D_\Delta^\vee(\pi) \times \text{St}(\Delta) \twoheadrightarrow \pi.$$

- Define $I_\Delta^\vee(\pi)$ to be the unique quotient of $\text{St}(\Delta) \times \pi$.

Definition 8.10. We use the notations in Definition 8.8. Let U_1^- be the opposite unipotent radical for P_1 . We say that $(\Delta_1, \Delta_2, \pi)$ is a *dual commutative triple* if the following composition:

$$C_c^\infty(P_{n_2}P_{n_1}^-, \text{St}(\Delta_2) \boxtimes \pi)_{U_{n_1}^-} \hookrightarrow (\text{St}(\Delta_2) \times \pi)_{U_1^-} \twoheadrightarrow I_{\Delta_2}^\vee(\pi)_{U_1^-} \twoheadrightarrow D_{\Delta_1}^\vee \circ I_{\Delta_2}^\vee(\pi) \boxtimes \text{St}(\Delta_2)$$

is non-zero, where

- the first map is induced from the natural embedding from $C_c^\infty(P_{n_2}P_{n_1}^-, \text{St}(\Delta_2) \boxtimes \pi)$ to $\text{St}(\Delta_2) \times \pi$;
- the second map is induced from the natural map from $\text{St}(\Delta) \times \pi$ to $I_\Delta^\vee(\pi)$;
- the third map is the map from the first bullet of Definition 8.9.

Proposition 8.11. *Let Δ_1, Δ_2 be segments. Let π be an irreducible representation of G_n . Then $(\Delta_1, \Delta_2, \pi)$ is a commutative triple if and only if $(\Delta_1^\vee, \Delta_2^\vee, \pi^\vee)$ is a dual commutative triple.*

Proof. This follows from Theorem 7.7 and the following standard facts: for a segment Δ , and for irreducible representations π and π' ,

- (1) $(\pi \times \pi')^\vee \cong \pi^\vee \times \pi'^\vee$;
- (2) $I_\Delta(\pi)^\vee \cong I_{\Delta^\vee}^\vee(\pi^\vee)$;
- (3) $D_\Delta(\pi)^\vee \cong D_{\Delta^\vee}^\vee(\pi^\vee)$.

We remark that (2) and (3) can also be deduced from (1) in a quite straightforward manner, \square

Proposition 8.11 implies that the notion of dual commutative triples coincides with another notion called LdRi-commutative triples in [Ch22+b].

REFERENCES

- [BC15] D. Barbasch and D. Ciubotaru, Hermitian forms for affine Hecke algebras, arXiv:1312.3316
- [BD84] J. N. Bernstein, Le centre de Bernstein, edited by P. Deligne. Travaux en Cours, Representations of reductive groups over a local field, 1-32, Hermann, Paris, 1984.
- [BZ76] I. N. Bernstein and A. V. Zelevinsky, Representations of the group GL_n ; F where F is a local non-Archimedean field, Russian Math. Surveys 31 (1976), no. 3, 1-68.
- [BZ77] I. N. Bernstein and A. V. Zelevinsky, *Induced representations of reductive p -adic groups*, I, Ann. Sci. Ecole Norm. Sup. **10** (1977), 441-472.
- [Be87] J. Bernstein, Second Adjointness Theorem for Representations of p -adic Groups, 1987
- [Be92] J. Bernstein. Representations of p -adic groups. Harvard University, 1992. Lectures by Joseph Bernstein. Written by Karl E. Rumelhart.
- [Bo91] A. Borel, Linear Algebraic Groups, GTM 1991.
- [BK15] R. Bezrukavnikov and D. Kazhdan, Geometry of Second Adjointness for p -adic groups, with an Appendix by Yakov Varshavsky, Roman Bezrukavnikov, and David Kazhdan, Represent. Theory 19 (2015), 299-332, <https://doi.org/10.1090/ert/471>
- [BK98] C.J. Bushnell, P.C. Kutzko, Smooth representations of reductive p -ADIC groups: structure theory via types, Proceedings of the London Mathematical Society, Volume 77, Issue 3, November 1998, Pages 582-634, <https://doi.org/10.1112/S0024611598000574>
- [Ca93] W. Casselman, Introduction to the theory of admissible representations of p -adic reductive groups. Preprint written 1974-1993
- [Ch22+] K.Y. Chan, Quotient branching law for p -adic (GL_{n+1}, GL_n) I: Generalized Gan-Gross-Prasad relevant pairs, arXiv:2212.05919, preprint.
- [Ch22+b] K.Y. Chan, Duality for generalized GGP relevant pairs for p -adic GL , arXiv:2210.17249, preprint
- [Ch22+c] K.Y. Chan, Construction of simple quotients of Bernstein-Zelevinsky derivatives and highest derivative multisegments I: reduction to combinatorics, preprint (2022)
- [CGP15] B. Conrad, O. Gabber, G. Prasad, Pseudo-reductive groups (2nd edition), new mathematical monographs: 26 (2015).
- [DHKM23] J.F. Dat, D. Helm, R. Kurinczuk, G. Moss, Finiteness for Hecke algebras of p -adic groups, Journal of AMS (2023), DOI:10.1090/jams/1034.
- [De77] V.V. Deodhar, Some characterizations of Bruhat ordering on a Coxeter group and determination of the relative Möbius function. Invent Math 39, 187-198 (1977). <https://doi.org/10.1007/BF01390109>
- [De86] V.V. Deodhar, On some geometric aspects of Bruhat orderings. I. A finer decomposition of Bruhat cells. Invent Math 79, 499-511 (1985). <https://doi.org/10.1007/BF01388520>
- [Re10] D. Renard, Représentations des groupes réductifs p -adiques, Cours Spécialisés, 17. Société Mathématique de France, Paris, 2010.
- [Ri92] R.W. Richardson, Intersections of double cosets in algebraic groups, Indagationes Mathematicae, Volume 3, Issue 1, 1992, Pages 69-77, [doi.org/10.1016/0019-3577\(92\)90028-J](https://doi.org/10.1016/0019-3577(92)90028-J).
- [Ze80] A. Zelevinsky, Induced representations of reductive p -adic groups II, Ann. Sci. Ecole Norm. Sup. **13** (1980), 154-210.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG,
Email address: kychan1@hku.hk