Generative Modeling by Minimizing the Wasserstein-2 Loss

Yu-Jui Huang^{*}

Zachariah Malik[†]

July 16, 2024

Abstract

This paper approaches the unsupervised learning problem by minimizing the second-order Wasserstein loss (the W_2 loss) through a *distribution-dependent* ordinary differential equation (ODE), whose dynamics involves the Kantorovich potential associated with the true data distribution and a current estimate of it. A main result shows that the time-marginal laws of the ODE form a gradient flow for the W_2 loss, which converges exponentially to the true data distribution. An Euler scheme for the ODE is proposed and it is shown to recover the gradient flow for the W_2 loss in the limit. An algorithm is designed by following the scheme and applying *persistent training*, which naturally fits our gradient-flow approach. In both low- and high-dimensional experiments, our algorithm outperforms Wasserstein generative adversarial networks by increasing the level of persistent training appropriately.

MSC (2020): 34A06, 49Q22, 68T01

Keywords: generative modeling, distribution-dependent ODEs, gradient flows, nonlinear Fokker-Planck equations, persistent training, generative adversarial networks

1 Introduction

From observed data points, a generative model aims at learning the unknown data distribution μ_d , a probability measure on \mathbb{R}^d , so as to generate new data points indistinguishable from the observed ones. The success of it demands a rigorous underlying theory.

In this paper, starting with any initial guess μ_0 of μ_d , we measure the discrepancy between μ_0 and μ_d by the second-order Wasserstein distance (i.e., $W_2(\cdot, \cdot)$ in (2.1) below) and strive to minimize the discrepancy efficiently. As the map $\mu \mapsto W_2^2(\mu, \mu_d)$ is convex, it is natural to ask if this minimization (or, learning of μ_d) can be done by *gradient descent*, the traditional wisdom for convex minimization. The crucial question is how we should define "gradient" in the present setting.

Thanks to subdifferential calculus for probability measures, widely studied under the W_2 distance in Ambrosio et al. [1], the Fréchet subdifferential of $\mu \mapsto J(\mu) := \frac{1}{2}W_2^2(\mu, \mu_d)$, denoted by $\partial J(\mu)$, always exists; see Definition 2.2 and the discussion below it. Hence, we can take the "gradient of J" evaluated at μ to be an element of $\partial J(\mu)$ and the resulting gradient flow converges exponentially to μ_d under W_2 (Proposition 3.1). What is *not* answered by the full-fledged W_2 -theory is how this gradient flow can actually be computed, for μ_d to be found realistically; see Remark 3.5.

To materialize the gradient flow, we plan to simulate a gradient-descent ordinary differential equation (ODE) with distribution dependence. Let Y be an \mathbb{R}^d -valued process satisfying

$$\frac{dY_t}{dt} \in -\partial J(\mu^{Y_t}), \quad \mu^{Y_0} = \mu_0, \tag{1.1}$$

^{*}Department of Applied Mathematics, University of Colorado, Boulder, CO 80309-0526, USA, email: yujui.huang@colorado.edu. Partially supported by National Science Foundation (DMS-2109002).

[†]Department of Applied Mathematics, University of Colorado, Boulder, CO 80309-0526, USA, email: zachariah.malik@colorado.edu.

where μ^{Y_t} denotes the law of Y_t at $t \ge 0$. This stipulates that Y_t evolves instantaneously along a "negative gradient" of J, evaluated at the current distribution μ^{Y_t} . Note that (1.1) is an inclusion, but not an equation, as $\partial J(\mu^{Y_t})$ in general may contain multiple elements. Notably, for the special case where μ^{Y_t} is absolutely continuous with respect to (w.r.t.) the Lebesgue measure (written $\mu^{Y_t} \ll \mathcal{L}^d$), $\partial J(\mu^{Y_t})$ reduces to a singleton containing only $\nabla \phi_{\mu^{Y_t}}^{\mu_d}$ (Lemma 2.1), where $\phi_{\mu^{Y_t}}^{\mu_d} : \mathbb{R}^d \to \mathbb{R}^d$ is a Kantorovich potential from μ^{Y_t} to μ_d (which, by definition, maximizes the dual formulation of $W_2(\mu^{Y_t}, \mu_d)$). Hence, if " $\mu^{Y_t} \ll \mathcal{L}^d$ for all $t \ge 0$ " holds, (1.1) will take the concrete form

$$dY_t = -\nabla \phi^{\mu_d}_{\mu^{Y_t}}(Y_t) dt, \quad \mu^{Y_0} = \mu_0 \ll \mathcal{L}^d.$$
 (1.2)

This is a distribution-dependent ODE. At time 0, the law of Y_0 is given by μ_0 , an initial distribution satisfying $\mu_0 \ll \mathcal{L}^d$. This initial randomness trickles through the ODE dynamics in (1.2), such that Y_t remains an \mathbb{R}^d -valued random variable for each t > 0. How the ODE evolves is determined jointly by $\phi_{\mu_{Y_t}}^{\mu_d}$ (depending on the current law $\mu_{Y_t}^{Y_t}$ and μ_d) and the actual realization of Y_t .

Solving ODE (1.2) is nontrivial. Compared with typical stochastic differential equations (SDEs) with distribution dependence (i.e., McKean-Vlasov SDEs), how (1.2) depends on distributions—via the Kantorovich potential $\phi^{\mu_d}_{\mu}$ —is quite unusual. As $\phi^{\mu_d}_{\mu}$ is only known to exist generally (without any tractable formula), the regularity of $(y, \mu) \mapsto \nabla \phi^{\mu_d}_{\mu}(y)$ is largely obscure, which prevents direct application of standard results for McKean-Vlasov SDEs.

In view of this, we will *not* deal with ODE (1.2) directly, but focus on the nonlinear Fokker-Planck equation associated with it. By finding a solution ν_t to the Fokker-Planck equation, we can in turn construct a solution Y to (1.2), with $\mu^{Y_t} = \nu_t$, by Trevisan's superposition principle [16]. This strategy was introduced in Barbu and Röckner [3] and recently used in Huang and Zhang [9] for generative modeling under the Jensen-Shannon divergence (JSD).

Our first observation is that if the gradient flow $\boldsymbol{\mu} = \{\boldsymbol{\mu}_t\}_{t\geq 0}$ for $J(\cdot) := \frac{1}{2}W_2^2(\cdot,\mu_d)$ satisfies " $\boldsymbol{\mu}_t \ll \mathcal{L}^d$ for all $t \geq 0$," it will automatically solve the nonlinear Fokker-Planck equation (i.e., (2.10) below). That is, the solvability of (2.10) hinges on the claim: "if we start with $\mu_0 \ll \mathcal{L}^d$, the resulting gradient flow $\boldsymbol{\mu}$ for J must fulfill $\boldsymbol{\mu}_t \ll \mathcal{L}^d$ for all t > 0." This will be proved in a constructive, yet indirect, way. First, we construct a curve $\boldsymbol{\beta} = \{\boldsymbol{\beta}_s\}_{s\in[0,1]}$ that moves $\boldsymbol{\beta}_0 := \mu_0$ to $\boldsymbol{\beta}_1 := \mu_d$ at a constant speed; see (4.1) below. By the definition of $\boldsymbol{\beta}$, Lemma 4.2 finds that " $\mu_0 \ll \mathcal{L}^d$ " entails " $\boldsymbol{\beta}_s \ll \mathcal{L}^d$ for all $s \in [0,1]$ " (excluding s = 1). By a change of speed from constant to exponentially decreasing, $\{\boldsymbol{\beta}_s\}_{s\in[0,1)}$ is rescaled into $\boldsymbol{\mu}^* = \{\boldsymbol{\mu}_t^*\}_{t\geq 0}$ (i.e., (4.3) below), so that $\boldsymbol{\mu}_t^* \ll \mathcal{L}^d$ for all $t \geq 0$. As $\boldsymbol{\mu}^*$ is by definition a time-changed constant-speed geodesic, the dynamics of $t \mapsto J(\boldsymbol{\mu}_t^*)$ can be decomposed and explicitly computed (Lemma 4.1), which reveals that $J(\boldsymbol{\mu}_t^*)$ always decreases in time with a largest possible slope. This "maximal slope" behavior in turn implies that $\boldsymbol{\mu}^*$ simply coincides with $\boldsymbol{\mu}$, the gradient flow for J (Theorem 4.1). The claim is thus proved and the explicitly constructed $\boldsymbol{\mu}^*$ solves the nonlinear Fokker-Planck equation (2.10); see Corollary 4.1.

By substituting $\boldsymbol{\mu}_t^*$ for $\boldsymbol{\mu}_t^{Y_t}$ in (1.2), we obtain a standard ODE without distribution dependence, i.e., $dY_t = f(t, Y_t)dt$ with $f(t, y) := -\nabla \phi_{\boldsymbol{\mu}_t^*}^{\mu_d}(y)$. To construct a solution to (1.2), we aim at finding a process Y such that (i) $t \mapsto Y_t$ satisfies the above standard ODE, and (ii) the law of Y_t coincides with $\boldsymbol{\mu}_t^*$ (i.e., $\boldsymbol{\mu}_t^{Y_t} = \boldsymbol{\mu}_t^*$) for all $t \ge 0$. Thanks to Trevisan's superposition principle [16], we can choose a probability measure \mathbb{P} on the canonical path space, so that the canonical process $Y_t(\omega) := \omega(t)$ fulfills (i) and (ii) above. This gives a solution Y to (1.2) (Proposition 5.1) and it is in fact unique up to time-marginal laws under suitable regularity conditions (Proposition 5.2). As $\boldsymbol{\mu}^*$ coincides with the gradient flow for $J(\cdot) := \frac{1}{2}W_2^2(\cdot, \mu_d)$, we conclude that $\boldsymbol{\mu}_t^{Y_t} = \boldsymbol{\mu}_t^* \to \mu_d$ exponentially under W_2 as $t \to \infty$; see Theorem 5.1, the main theoretic result of this paper.

Theorem 5.1 suggests that we can uncover μ_d by simulating ODE (1.2). To facilitate the actual implementation, we propose a forward Euler scheme (i.e., (5.7) below), which, for any fixed time step $0 < \varepsilon < 1$, simulates a sequence of random variables $\{Y_n^{\varepsilon}\}_{n \in \mathbb{N} \cup \{0\}}$ at time points $\{n\varepsilon\}_{n \in \mathbb{N} \cup \{0\}}$. Interestingly, under our scheme, how the distribution $\mu^{Y_n^{\varepsilon}}$ is transformed into $\mu^{Y_{n+1}^{\varepsilon}}$ is optimal in the sense of W_2 (Lemma 5.1). This crucially implies that the laws of $\{Y_n^{\varepsilon}\}_{n \in \mathbb{N} \cup \{0\}}$ all lie on the curve μ^* , which consists of the time-marginal laws of ODE (1.2) (Proposition 5.3). As $\varepsilon \downarrow 0$, Theorem 5.2 shows that the laws of $\{Y_n^{\varepsilon}\}_{n \in \mathbb{N} \cup \{0\}}$ gradually cover the entire curve μ^* . That is, our Euler scheme does correctly recover the evolving distributions induced by ODE (1.2) in the limit.

Based on this Euler scheme, Algorithm 1 (called W2-FE, where FE means "forward Euler") is designed to simulate ODE (1.2). In particular, a "generator" $G_{\theta} : \mathbb{R}^d \to \mathbb{R}^d$, modeled by a deep neural network, is updated at each time point to keep track of the changing distributions along the (discretized) ODE. Remarkably, in updating G_{θ} , we reuse the same minibatch of points newly generated by the ODE in $K \in \mathbb{N}$ consecutive iterations of stochastic gradient descent (SGD). This is in line with *persistent training* in Fischetti et al. [5], distinct from standard SGD implementation that takes K = 1 by default. Note that persistent training fits our ODE approach naturally: as the goal of updating G_{θ} is to learn the new distribution induced by the ODE at the next time point, keeping the newly generated points unchanged in SGD iterations allows G_{θ} to more accurately represent the new distribution, so that the ODE can be more closely followed.

By a min-max game perspective, generative adversarial networks (GANs) also aim at minimizing the discrepancy between an initial estimate μ_0 and the true data distribution μ_d . In measuring the discrepancy, the original GAN [6] uses JSD; Wasserstein GAN (WGAN) [2], a popular version of GANs known for its stability, uses the first-order Wasserstein distance; W2-GAN [11], less heeded in the literature, uses the W_2 distance (the same as this paper). Intriguingly, while our algorithm is designed by a gradient-flow perspective (instead of the min-max game idea), it covers W2-GAN as a special case with K = 1 (i.e., no persistent training); see Proposition 6.1.

We implement our algorithm W2-FE with varying persistency levels (i.e., K values) and compare its performance with that of the refined WGAN algorithm in [13] (called W1-LP), which is arguably one of the most well-performing and stable WGAN algorithm. In both low- and high-dimensional experiments, by increasing the persistency level appropriately, W2-FE converges significantly faster than W1-LP and achieves similar or better outcomes; see Figures 1, 2, and 3.

The rest of the paper is organized as follows. Section 1.1 collects fundamental notation. Section 2 motivates the gradient-descent ODE to be studied. In Section 3, the gradient flow theory under W_2 is briefly reviewed and tailored to the needs of our study. Section 4 finds a solution to the nonlinear Fokker-Planck equation associated with the gradient-descent ODE. Based on this, Section 5 builds a solution to the ODE, whose time-marginal laws converge to μ_d exponentially. A discretized scheme for the ODE is proposed and analyzed in detail in Section 5.1. Section 6 designs an algorithm based on the discretized scheme. Its performance is illustrated by numerical experiments in Section 7.

1.1 Notation

For any Polish space \mathcal{X} , let $\mathcal{B}(\mathcal{X})$ be the Borel σ -algebra of \mathcal{X} and $\mathcal{P}(\mathcal{X})$ be the set of all probability measures on $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$. Given two Polish spaces \mathcal{X} and $\mathcal{Y}, \mu \in \mathcal{P}(\mathcal{X})$, and a Borel $f : \mathcal{X} \to \mathcal{Y}$, we define $f_{\#}\mu \in \mathcal{P}(\mathcal{Y})$, called the pushforward of μ through f, by $f_{\#}\mu(B) := \mu(f^{-1}(B))$ for $B \in \mathcal{B}(\mathcal{Y})$. On the product space $\mathcal{X} \times \mathcal{Y}$, consider the projection operators $\pi^1(x, y) := x$ and $\pi^2(x, y) := y$ for all $(x, y) \in \mathcal{X} \times \mathcal{Y}$. Given $\mu \in \mathcal{P}(\mathcal{X})$ and $\nu \in \mathcal{P}(\mathcal{Y})$, a Borel $\mathbf{t} : \mathcal{X} \to \mathcal{Y}$ is called a transport map from μ to ν if $\mathbf{t}_{\#}\mu = \nu$. Also, a probability $\gamma \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ is called a transport plan from μ to ν if $\pi^1_{\#}\gamma = \mu$ and $\pi^2_{\#}\gamma = \nu$. Let $\Gamma(\mu, \nu)$ denote the set of all transport plans from μ to ν .

Fix any $d \in \mathbb{N}$. Let **i** denote the identity map on \mathbb{R}^d (i.e., $\mathbf{i}(y) := y$ for all $y \in \mathbb{R}^d$) and \mathcal{L}^d denote the Lebesgue measure on \mathbb{R}^d . For any measure μ on \mathbb{R}^d , we write $\mu \ll \mathcal{L}^d$ if it is absolutely continuous w.r.t. \mathcal{L}^d . Given $p \ge 1$ and $S \subseteq \mathbb{R}^d$, let $L^p(S, \mu)$ be the set of all $f : \mathbb{R}^d \to \mathbb{R}$ with $\int_S |f(y)|^p d\mu(y) < \infty$. For the case $\mu = \mathcal{L}^d$, we will simply write $L^p(S)$ for $L^p(S, \mu)$.

$\mathbf{2}$ **Preliminaries**

Fix $d \in \mathbb{N}$ and denote by $\mathcal{P}_2(\mathbb{R}^d)$ the set of elements in $\mathcal{P}(\mathbb{R}^d)$ with finite second moments, i.e.,

$$\mathcal{P}_2(\mathbb{R}^d) := \bigg\{ \mu \in \mathcal{P}(\mathbb{R}^d) : \int_{\mathbb{R}^d} |y|^2 d\mu(y) < \infty \bigg\}.$$

The second-order Wasserstein distance, a metric on $\mathcal{P}_2(\mathbb{R}^d)$, is defined by

$$W_2(\mu,\nu) := \inf_{\gamma \in \Gamma(\mu,\nu)} \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} |x-y|^2 \, d\gamma(x,y) \right)^{1/2}, \quad \forall \mu, \nu \in \mathcal{P}_2(\mathbb{R}^d).$$
(2.1)

For $\{\mu_n\}_{n\in\mathbb{N}}$ and μ in $\mathcal{P}_2(\mathbb{R}^d)$ with $W_2(\mu_n, \mu) \to 0$, we will often write $\mu_n \to \mu$ for brevity. To state the duality formula for the W_2 distance, we set $c(x, y) := \frac{1}{2}|x - y|^2$ for $x, y \in \mathbb{R}^d$ and define the *c*-transform of any $\varphi : \mathbb{R}^d \to \mathbb{R}$ by $\varphi^c(y) := \inf_{x \in \mathbb{R}^d} \{c(x,y) - \varphi(x)\}, y \in \mathbb{R}^d$. Also, we say φ is *c*-concave if $\varphi = \psi^c$ for some $\psi : \mathbb{R}^d \to \mathbb{R}$. By [17, Theorem 5.10 (i)], the W_2 distance fulfills

$$W_2(\mu,\nu) = \sup\left\{\int_{\mathbb{R}^d} \phi(x) \, d\mu(x) + \int_{\mathbb{R}^d} \phi^c(y) \, d\nu(y) : \phi \in L^1(\mathbb{R}^d,\mu) \text{ is } c\text{-concave}\right\}.$$
(2.2)

Thanks to $|x-y|^2 \leq 2(|x|^2+|y|^2)$ for $x, y \in \mathbb{R}^d$, [17, Theorem 5.10 (iii)] asserts that a minimizer of (2.1) and a maximizer of (2.2) always exist.

Definition 2.1. Given $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$, a $\gamma \in \Gamma(\mu, \nu)$ that minimizes (2.1) is called an optimal transport plan from μ to ν . We denote by $\Gamma_0(\mu,\nu)$ the set of all such plans. A c-concave $\phi \in$ $L^1(\mathbb{R}^d,\mu)$ that maximizes (2.2) is called a Kantorovich potential from μ to ν and denoted by ϕ^{ν}_{μ} .

Remark 2.1. The c-concavity of ϕ^{ν}_{μ} implies that $f(x) := \frac{1}{2}|x|^2 - \phi^{\nu}_{\mu}(x)$, $x \in \mathbb{R}^d$, is convex; see [1, Section 6.2.3]. It follows that $\nabla f(x)$ (and accordingly $\nabla \phi_{\mu}^{\nu}(x)$) exists for \mathcal{L}^{d} -a.e. $x \in \mathbb{R}^{d}$.

Remark 2.2. For any $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$, suppose additionally that μ belongs to

$$\mathcal{P}_2^r(\mathbb{R}^d) := \{ \mu \in \mathcal{P}_2(\mathbb{R}^d) : \mu \ll \mathcal{L}^d \}.$$

Then, by [1, Theorem 6.2.4 and Remark 6.2.5 a)], $\Gamma_0(\mu,\nu)$ is a singleton and the unique optimal transport plan $\gamma \in \mathcal{P}(\mathbb{R}^d \times \mathbb{R}^d)$ takes the form

$$\gamma = (\mathbf{i} \times \mathbf{t}^{\nu}_{\mu})_{\#} \mu \tag{2.3}$$

for some transport map $\mathbf{t}^{\nu}_{\mu}: \mathbb{R}^d \to \mathbb{R}^d$ from μ to ν , and the map \mathbf{t}^{ν}_{μ} can be expressed as

$$\mathbf{t}^{\nu}_{\mu}(x) = x - \nabla \phi^{\nu}_{\mu}(x) \quad \text{for } \mu\text{-a.e. } x \in \mathbb{R}^d, \tag{2.4}$$

for any Kantorovich potential ϕ^{ν}_{μ} . Note that $\nabla \phi^{\nu}_{\mu}$ is well-defined μ -a.e. by Remark 2.1 and $\mu \ll \mathcal{L}^d$.

Remark 2.3. We will call $\mathbf{t}^{\nu}_{\mu} : \mathbb{R}^d \to \mathbb{R}^d$ in (2.3) an optimal transport map from μ to ν . Indeed, as $\gamma = (\mathbf{i} \times \mathbf{t}^{\nu}_{\mu})_{\#}\mu$ is an optimal transport plan from μ to ν , we deduce from (2.1) that

$$W_{2}(\mu,\nu) = \left(\int_{\mathbb{R}^{d}} |\mathbf{t}_{\mu}^{\nu}(x) - x|^{2} d\mu(x)\right)^{1/2} = \inf_{\mathbf{t}:\mathbb{R}^{d}\to\mathbb{R}^{d},\mathbf{t}_{\#}\mu=\nu} \left(\int_{\mathbb{R}^{d}} |\mathbf{t}(x) - x|^{2} d\mu(x)\right)^{1/2}.$$

2.1 Problem Formulation

Let $\mu_{d} \in \mathcal{P}_{2}(\mathbb{R}^{d})$ denote the (unknown) data distribution. Starting with an arbitrary initial guess $\mu_{0} \in \mathcal{P}_{2}(\mathbb{R}^{d})$ of μ_{d} , we aim at solving the problem

$$\min_{\mu \in \mathcal{P}_2(\mathbb{R}^d)} W_2^2(\mu, \mu_d) \tag{2.5}$$

efficiently. By (2.1), it can be checked directly that $\mu \mapsto W_2^2(\mu, \mu_d)$ is convex on $\mathcal{P}_2(\mathbb{R}^d)$. Hence, it is natural to ask if (2.5) can be solved simply by *gradient descent*. The challenge here is how we should define the geadient of a function on $\mathcal{P}_2(\mathbb{R}^d)$. By subdifferential calculus in $\mathcal{P}_2(\mathbb{R}^d)$ developed by [1], for a general $G : \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$, we can take the "gradient of G" to be an element of its Fréchet subdifferential, defined below following (10.3.12)-(10.3.13) and Definition 10.3.1 in [1].

Definition 2.2. Given $G : \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ and $\mu \in \mathcal{P}_2(\mathbb{R}^d)$, the Fréchet subdifferential of G at μ , denoted by $\partial G(\mu)$, is the collection of all $\xi \in L^2(\mathbb{R}^d, \mu)$ such that

$$G(\nu) - G(\mu) \ge \inf_{\gamma \in \Gamma_0(\mu,\nu)} \int_{\mathbb{R}^d} \xi(x) \cdot (y-x) \, d\gamma(x,y) + o(W_2(\mu,\nu)) \quad \forall \nu \in \mathcal{P}_2(\mathbb{R}^d).$$
(2.6)

For the function $J: \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ defined by

$$J(\mu) := \frac{1}{2} W_2^2(\mu, \mu_{\rm d}), \tag{2.7}$$

the proof of [1, Theorem 10.4.12], particularly (10.4.53) therein, shows that $\xi \in \partial J(\mu)$ generally exists. The construction of ξ , however, relies on an optimal transport plan $\gamma \in \Gamma_0(\mu, \mu_d)$, which is not unique and does not admit concrete characterizations in general. Fortunately, for the special case where $\mu \in \mathcal{P}_2^r(\mathbb{R}^d)$, $\partial J(\mu)$ becomes much more tractable.

Lemma 2.1. At any $\mu \in \mathcal{P}_2^r(\mathbb{R}^d)$, the Fréchet subdifferential of $J : \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ in (2.7) is a singleton. Specifically, $\partial J(\mu) = \{-(\mathbf{t}_{\mu}^{\mu_d} - \mathbf{i})\} = \{\nabla \phi_{\mu}^{\mu_d}\}$, where $\mathbf{t}_{\mu}^{\mu_d}$ (resp. $\phi_{\mu}^{\mu_d}$) is an optimal transport map (resp. a Kantorovich potential) from μ to μ_d .

Proof. For any $\nu \in \mathcal{P}_2(\mathbb{R}^d)$, by [1, Corollary 10.2.7] (with $\mu_1 \in \mathcal{P}_2^r(\mathbb{R}^d)$ and $\mu_2, \mu_3 \in \mathcal{P}_2(\mathbb{R}^d)$ therein taken to be $\mu \in \mathcal{P}_2^r(\mathbb{R}^d)$ and $\mu_d, \nu \in \mathcal{P}_2(\mathbb{R}^d)$, respectively), we have

$$J(\nu) - J(\mu) = -\int_{\mathbb{R}^d} (\mathbf{t}_{\mu}^{\mu_d}(x) - x) \cdot (\mathbf{t}_{\mu}^{\nu}(x) - x) \, d\mu(x) + o(W_2(\mu, \nu)).$$

In addition, as $\mu \in \mathcal{P}_2^r(\mathbb{R}^d)$, $\Gamma_0(\mu, \nu)$ is a singleton that contains only the transport plan in (2.3) (see Remark 2.2). Hence, the previous equality can be equivalently written as

$$J(\nu) - J(\mu) = \inf_{\gamma \in \Gamma_0(\mu,\nu)} \int_{\mathbb{R}^d} -(\mathbf{t}_{\mu}^{\mu_d}(x) - x) \cdot (y - x) \, d\gamma(x,y) + o(W_2(\mu,\nu)).$$
(2.8)

That is, J fulfills (2.6) with $\xi = -(\mathbf{t}_{\mu}^{\mu_d} - \mathbf{i}) \in \partial J(\mu)$. By (2.8) and $\Gamma_0(\mu, \nu)$ being a singleton, we may follow the same argument in the proof of [4, Proposition 5.63] to show that any $\xi' \in \partial J(\mu)$ must coincide with $-(\mathbf{t}_{\mu}^{\mu_d} - \mathbf{i}) \mu$ -a.e.. That is, $-(\mathbf{t}_{\mu}^{\mu_d} - \mathbf{i})$ is the unique element in $\partial J(\mu)$. The fact $-(\mathbf{t}_{\mu}^{\mu_d} - \mathbf{i}) = \nabla \phi_{\mu}^{\mu_d}$ follows directly from (2.4).

Lemma 2.1 suggests that (2.5) can potentially be solved by the gradient-descent ODE

$$dY_t = -\nabla \phi_{\mu Y_t}^{\mu_d}(Y_t) dt, \quad \mu^{Y_0} = \mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d).$$
 (2.9)

This ODE is distribution-dependent in nontrivial ways. At time 0, Y_0 is an \mathbb{R}^d -valued random variable whose law is given by $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, an arbitrarily specified initial distribution. This initial randomness trickles through the ODE dynamics, such that Y_t remains an \mathbb{R}^d -valued random variable, with its law denoted by $\mu^{Y_t} \in \mathcal{P}_2(\mathbb{R}^d)$, at each t > 0. The evolution of the ODE is determined jointly by a Kantorovich potential from the present distribution μ^{Y_t} to μ_d (i.e., the function $\phi_{\mu^{Y_t}}^{\mu_d}$) and the actual realization of Y_t (which is plugged into the map $y \mapsto \nabla \phi_{\mu^{Y_t}}^{\mu_d}(y)$).

Our goal is to show that (i) there is a (unique) solution Y to ODE (2.9) and (ii) the law of Y_t will ultimately converge to μ_d (i.e., $W_2(\mu^{Y_t}, \mu_d) \to 0$ as $t \to \infty$), thereby solving (2.5). How (2.9) depends on distributions—via the Kantorovich potential $\phi^{\mu}_{\mu^{Y_t}}$ —poses nontrivial challenges.

- (a) For the existence of a unique solution to a McKean-Vlasov SDE (i.e., an SDE with dependence on the present law μ^{Y_t}), one needs the SDE's coefficients to be sufficiently regular in both the state variable and μ^{Y_t} . As $\phi_{\mu^{Y_t}}^{\mu_d}$ in (2.9) is only known to exist (without any tractable formula), the regularity of $(y, \mu) \mapsto \nabla \phi_{\mu}^{\mu_d}(y)$ is left largely unexplored.
- (b) The coefficient $-\nabla \varphi_{\mu^{Y_t}}^{\mu_d}(\cdot)$ in (2.9) is only well-defined \mathcal{L}^d -a.e. (Remark 2.1). If the random variable Y_t takes a value where $-\nabla \varphi_{\mu^{Y_t}}^{\mu_d}(\cdot)$ is undefined, the ODE dynamics will simply break down. To avoid this, we need " $\mu^{Y_t} \ll \mathcal{L}^d$ for all $t \ge 0$," for every Y_t to concentrate on where $-\nabla \varphi_{\mu^{Y_t}}^{\mu_d}(\cdot)$ is well-defined. At t = 0, this is achieved by choosing μ_0 directly from $\mathcal{P}_2^r(\mathbb{R}^d)$. The question is whether subsequent distributions $\{\mu^{Y_t}\}_{t>0}$ will remain in $\mathcal{P}_2^r(\mathbb{R}^d)$.

In view of the above challenges, we will *not* work with ODE (2.9) directly, but instead focus on the corresponding nonlinear Fokker-Planck equation (or, continuity equation)

$$\partial_t \boldsymbol{\mu}_t - \operatorname{div}(\nabla \varphi_{\boldsymbol{\mu}_t}^{\mu_d} \boldsymbol{\mu}_t) = 0, \quad \lim_{t \ge 0} \boldsymbol{\mu}_t = \mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d).$$
(2.10)

By first finding a solution $\{\mu_t\}_{t\geq 0}$ to (2.10), we can in turn construct a solution to ODE (2.9), made specifically from $\{\mu_t\}_{t\geq 0}$. This way of solving a distribution-dependent ODE (or SDE) was introduced in Barbu and Röckner [3] and recently used in Huang and Zhang [9] for a learning problem similar to (2.5), with the W_2 distance replaced by the Jensen-Shannon divergence (JSD). To actually find a solution to (2.10), we will rely on "gradient flows" in $\mathcal{P}_2(\mathbb{R}^d)$, which are now introduced.

3 Gradient Flows in $\mathcal{P}_2(\mathbb{R}^d)$

In this section, key aspects of the gradient flow theory for probability measures will be introduced from [1] and tailored to the needs of Sections 4 and 5 below. To begin with, we introduce absolutely continuous curves in $\mathcal{P}_2(\mathbb{R}^d)$ in line with [1, Definition 1.1.1].

Definition 3.1. Fix an open interval $I \subseteq (0, \infty)$. For any $p \ge 1$, we say $\boldsymbol{\mu} : I \to \mathcal{P}_2(\mathbb{R}^d)$ belongs to $AC^p(I; \mathcal{P}_2(\mathbb{R}^d))$ (resp. $AC^p_{loc}(I; \mathcal{P}_2(\mathbb{R}^d))$) if there exists $m \in L^p(I)$ (resp. $m \in L^p_{loc}(I)$) such that

$$W_2(\boldsymbol{\mu}_{t_1}, \boldsymbol{\mu}_{t_2}) \le \int_{t_1}^{t_2} m(t) \, dt, \quad \forall t_1, t_2 \in I \text{ with } t_1 \le t_2.$$
(3.1)

For the case p = 1, we call μ an absolutely continuous (resp. locally absolutely continuous) curve and simply write $\mu \in AC(I; \mathcal{P}_2(\mathbb{R}^d))$ (resp. $\mu \in AC_{loc}(I; \mathcal{P}_2(\mathbb{R}^d)))$.

Remark 3.1. Fix any $p \ge 1$ and open interval $I \subseteq (0, \infty)$.

(i) If $\mathcal{L}^1(I) < \infty$, since $L^p(I) \subseteq L^1(I)$, the inclusion $AC^p(I; \mathcal{P}_2(\mathbb{R}^d)) \subseteq AC(I; \mathcal{P}_2(\mathbb{R}^d))$ follows; similarly, we also have $AC^p_{\text{loc}}(I; \mathcal{P}_2(\mathbb{R}^d)) \subseteq AC_{\text{loc}}(I; \mathcal{P}_2(\mathbb{R}^d))$.

(ii) If $\mathcal{L}^1(I) = \infty$, the inclusions in (i) no longer hold in general. Nonetheless, for any open interval $I' \subseteq I$ with $\mathcal{L}^1(I') < \infty$, since $L^p_{\text{loc}}(I) \subseteq L^p(I') \subseteq L^1(I')$, we have $AC^p_{\text{loc}}(I; \mathcal{P}_2(\mathbb{R}^d)) \subseteq AC^p(I'; \mathcal{P}_2(\mathbb{R}^d)) \subseteq AC(I'; \mathcal{P}_2(\mathbb{R}^d))$.

Remark 3.2. Fix any $p \ge 1$ and open interval $I \subseteq (0,\infty)$. For any $\mu \in AC^p(I; \mathcal{P}_2(\mathbb{R}^d))$ (resp. $\mu \in AC_{loc}^p(I; \mathcal{P}_2(\mathbb{R}^d))$), as a direct consequence of (3.1), the so-called "metric derivative"

$$|\boldsymbol{\mu}'|(t) := \lim_{s \to t} \frac{W_2(\boldsymbol{\mu}_s, \boldsymbol{\mu}_t)}{|s - t|}$$
(3.2)

exists with $|\boldsymbol{\mu}'|(t) \leq m(t)$ for \mathcal{L}^1 -a.e. $t \in I$, which in turn implies $|\boldsymbol{\mu}'| \in L^p(I)$ (resp. $|\boldsymbol{\mu}'| \in L^p_{loc}(I)$); see [1, Theorem 1.1.2] for detailed arguments.

The next result shows that every $\boldsymbol{\mu} \in AC^2_{loc}((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$ is inherently associated with a vector field $v: (0,\infty) \times \mathbb{R}^d \to \mathbb{R}^d$, which will be understood as the "velocity" of $\boldsymbol{\mu}$.

Lemma 3.1. For any $\boldsymbol{\mu} \in AC^2_{loc}((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$, there exists a unique Borel vector field $v : (0,\infty) \times \mathbb{R}^d \to \mathbb{R}^d$, written $(t,x) \mapsto v_t(x)$, such that

$$v_t \in L^2(\mathbb{R}^d, \mu_t) \quad and \quad \|v_t\|_{L^2(\mathbb{R}^d, \mu_t)} = |\mu'|(t) \quad for \ \mathcal{L}^1 \text{-} a.e. \ t > 0$$

$$(3.3)$$

and the resulting Fokker-Planck equation

$$\partial_t \boldsymbol{\mu}_t + \operatorname{div}(v_t \boldsymbol{\mu}_t) = 0 \quad on \quad (0, \infty) \times \mathbb{R}^d$$
(3.4)

holds in the sense of distributions, i.e.,

$$\int_0^\infty \int_{\mathbb{R}^d} \left(\partial_t \varphi(t, x) + v_t(x) \cdot \nabla \varphi(t, x) \right) d\boldsymbol{\mu}_t(x) dt = 0 \quad \forall \varphi \in C_c^\infty((0, \infty) \times \mathbb{R}^d).$$
(3.5)

Proof. Take an increasing sequence $\{I_n\}_{n\in\mathbb{N}}$ of open intervals such that $\mathcal{L}^1(I_n) < \infty$ and $\bigcup_{n\in\mathbb{N}} I_n = (0,\infty)$. With $\boldsymbol{\mu} \in AC_{\text{loc}}^2((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$, Remark 3.1 (ii) implies $\boldsymbol{\mu} \in AC(I_n; \mathcal{P}_2(\mathbb{R}^d))$ for all $n \in \mathbb{N}$. We can then apply [1, Theorem 8.3.1] to $\boldsymbol{\mu}$ on each interval I_n , which asserts the existence of a Borel vector field $v^n : I_n \times \mathbb{R}^d \to \mathbb{R}^d$ such that

$$v_t^n \in L^2(\mathbb{R}^d, \boldsymbol{\mu}_t) \quad \text{and} \quad \|v_t^n\|_{L^2(\mathbb{R}^d, \boldsymbol{\mu}_t)} = |\boldsymbol{\mu}'|(t) \quad \text{for } \mathcal{L}^1\text{-a.e. } t \in I_n$$

$$(3.6)$$

and $\partial_t \boldsymbol{\mu}_t - \operatorname{div}(v_t^n \boldsymbol{\mu}_t) = 0$ holds on $I_n \times \mathbb{R}^d$ in the sense of distributions, i.e.,

$$\int_{I_n} \int_{\mathbb{R}^d} \left(\partial_t \varphi(t, x) + v_t^n(x) \cdot \nabla \varphi(t, x) \right) d\boldsymbol{\mu}_t(x) dt = 0 \quad \forall \varphi \in C_c^\infty(I_n \times \mathbb{R}^d).$$
(3.7)

Moreover, by [1, Proposition 8.4.5], $v_t^n : \mathbb{R}^d \to \mathbb{R}^d$ is uniquely determined \mathcal{L}^1 -a.e. in I_n by (3.6) and (3.7). By such uniqueness on each I_n , the function $v_t(x) := v_t^n(x)$, for $(t, x) \in I_n \times \mathbb{R}^d$ and $n \in \mathbb{N}$, is well-defined and Borel. Then, (3.3) and (3.5) follow directly from (3.6) and (3.7).

Definition 3.2. Given $\boldsymbol{\mu} \in AC^2_{loc}((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$, the unique Borel vector field $v : (0,\infty) \times \mathbb{R}^d \to \mathbb{R}^d$ that fulfills (3.3) and (3.4) is called the velocity field of $\boldsymbol{\mu}$.

Remark 3.3. Note that (3.3) and (3.4) allow the use of Trevisan's superposition principle [16], which gives a weak solution to the following ODE: $dY_t = v_t(Y_t)dt$ with $\mu^{Y_t} = \mu_t$ for all $t \in (0, \infty)$; see the precise arguments in the proof of Theorem 5.1 below. That is, at every t > 0, the vector $v_t(Y_t)$ dictates how Y_t moves in \mathbb{R}^d instantaneously, and the resulting laws of the random variables $\{Y_t\}_{t>0}$ recover $\mu \in AC^2_{loc}((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$. This explains why we call v the velocity field of μ . If the velocity field v always guides μ forward along the "negative gradient" of some function $G: \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$, we naturally call μ a "gradient flow."

Definition 3.3. We say $\boldsymbol{\mu} \in AC^2_{\text{loc}}((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$ is a gradient flow for $G : \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ if its velocity field $v : (0,\infty) \times \mathbb{R}^d \to \mathbb{R}^d$ satisfies $v_t \in -\partial G(\boldsymbol{\mu}_t)$ for \mathcal{L}^1 -a.e. t > 0.

Definition 3.3 is in line with [1, Definition 11.1.1]. While the latter seems to require additionally that v_t lies in the tangent bundle to $\mathcal{P}_2(\mathbb{R}^d)$ at μ_t , defined as the $L^2(\mathbb{R}^d, \mu_t)$ -closure of $\{\nabla \psi : \psi \in C_c^{\infty}(\mathbb{R}^d)\}$, v_t readily fulfills this by [1, Proposition 8.4.5] and the proof of Lemma 3.1 above.

For $J : \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ in (2.7), a gradient flow μ does exist and is unique once an initial measure $\mu_0 \in \mathcal{P}_2(\mathbb{R}^d)$ is specified. Moreover, it diminishes J exponentially fast.

Proposition 3.1. For any $\mu_0 \in \mathcal{P}_2(\mathbb{R}^d)$, there exists a gradient flow $\boldsymbol{\mu} \in AC^2_{loc}((0,\infty);\mathcal{P}_2(\mathbb{R}^d))$ for $J:\mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ in (2.7) with $\lim_{t\downarrow 0} \boldsymbol{\mu}_t = \mu_0$. Moreover,

$$J(\boldsymbol{\mu}_t) \le e^{-2t} J(\boldsymbol{\mu}_0) \quad \forall t > 0;$$
(3.8)

$$J(\boldsymbol{\mu}_t) - J(\boldsymbol{\mu}_s) = -\int_s^t \int_{\mathbb{R}^d} |v_u(x)|^2 d\boldsymbol{\mu}_u(x) du \le 0 \quad \forall 0 \le s < t,$$
(3.9)

where $v: (0, \infty) \times \mathbb{R}^d$ is the velocity field of $\boldsymbol{\mu}$. If $\bar{\boldsymbol{\mu}} \in AC^2_{loc}((0, \infty); \mathcal{P}_2(\mathbb{R}^d))$ is also a gradient flow for J with $\lim_{t\downarrow 0} \bar{\boldsymbol{\mu}}_t = \mu_0$, then $W_2(\boldsymbol{\mu}_t, \bar{\boldsymbol{\mu}}_t) = 0$ for all t > 0.

Proof. By [1, Lemma 9.2.1], $J : \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ in (2.7) is 1-convex along generalized geodesics. Hence, we can apply [1, Theorem 11.2.1] to conclude that there exists a gradient flow for J with $\lim_{t \to 0} \mu_t = \mu_0$ and it fulfills (3.8) and (3.9) (which corresponds to (11.2.5) and (11.2.4) in [1], respectively). The uniqueness result follows from [1, Theorem 11.1.4].

Remark 3.4. Proposition 3.1 shows the advantage of working with W_2 in (2.5): not only does a unique gradient flow exists, it converges to μ_d exponentially fast, thanks to the full-fledged gradient flow theory under W_2 . This is in contrast to Huang and Zhang [9], where (2.5) is studied with the W_2 distance replaced by JSD. As no gradient flow theory under JSD is known, a substantial effort is spent in [9] to define a proper "gradient" notion and show that the associated gradient flow does exist and converge to μ_d , relying on the theory of differential equations on Banach spaces.

Remark 3.5. Despite the desirable properties in Proposition 3.1, it is unclear how the gradient flow for J in (2.7) can actually be computed. The construction of the gradient flow in [1, Theorem 11.2.1] is based on a discretized scheme that requires an optimization problem to be solved at each time step. Although [1, Corollary 2.2.2] asserts that solutions to these problems generally exist, it provides no clear clues for actually finding the solutions and thereby implementing the discretized scheme.

A gradient flow for $J : \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ in (2.7) can be characterized equivalently as a curve of "maximal slope." Specifically, consider the local slope of J at $\nu \in \mathcal{P}_2(\mathbb{R}^d)$, defined by

$$|\partial J|(\nu) := \limsup_{\rho \to \nu} \frac{(J(\nu) - J(\rho))^+}{W_2(\nu, \rho)}.$$
(3.10)

Now, for any $\boldsymbol{\mu} \in AC^2_{\text{loc}}((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$, observe that

$$\frac{d}{dt}J(\boldsymbol{\mu}_t) = \lim_{h \to 0} \frac{J(\boldsymbol{\mu}_{t+h}) - J(\boldsymbol{\mu}_t)}{W_2(\boldsymbol{\mu}_{t+h}, \boldsymbol{\mu}_t)} \frac{W_2(\boldsymbol{\mu}_{t+h}, \boldsymbol{\mu}_t)}{h} \ge -|\partial J|(\boldsymbol{\mu}_t)|\boldsymbol{\mu}'|(t) \quad \forall t > 0,$$

where the inequality follows directly from the definitions of $|\partial J|$ and $|\mu'|$ in (3.10) and (3.2), respectively. That is, $|\partial J|(\mu_t)|\mu'|(t)$ is the largest possible slope (in magnitude) of $s \mapsto J(\mu_s)$ at time t. By Young's inequality, the above relation yields

$$\frac{d}{dt}J(\boldsymbol{\mu}_t) \ge -|\partial J|(\boldsymbol{\mu}_t)|\boldsymbol{\mu}'|(t) \ge -\frac{1}{2}\left(|\partial J|^2(\boldsymbol{\mu}_t) + |\boldsymbol{\mu}'|^2(t)\right) \quad \forall t > 0.$$

Hence, if it can be shown that

$$\frac{d}{dt}J(\boldsymbol{\mu}_t) \le -\frac{1}{2}\left(|\partial J|^2(\boldsymbol{\mu}_t) + |\boldsymbol{\mu}'|^2(t)\right) \quad \text{for } \mathcal{L}^1\text{-a.e. } t > 0,$$
(3.11)

we get $\frac{d}{dt}J(\boldsymbol{\mu}_t) = -|\partial J|(\boldsymbol{\mu}_t)|\boldsymbol{\mu}'|(t)$ for \mathcal{L}^d -a.e. t > 0, i.e., $s \mapsto J(\boldsymbol{\mu}_s)$ almost always moves with the largest possible slope. This behavior characterizes gradient flows for J.

Proposition 3.2. Given $\mu \in AC^2_{loc}((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$, μ satisfies (3.11) if and only if μ is a gradient flow for J as in Definition 3.3.

Proof. By [1, Lemma 9.2.1 and Remark 9.2.8], $J : \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ in (2.7) is 1-geodesically convex per [1, Definition 9.1.1]. This gives two important properties. First, it allows the use of [1, Lemma 10.3.8], which asserts that J is a regular functional per [1, Definition 10.3.9]. Second, it also indicates that [1, Theorem 4.1.2 (i)] is applicable, which asserts that a discretized minimization problem has a solution. With these two properties of J, we conclude from [1, Theorem 11.1.3] that μ satisfies (3.11) if and only if it is a gradient flow for J and $t \mapsto J(\mu_t)$ equals a function of bounded variation for \mathcal{L}^1 -a.e. t > 0. Finally, by Proposition 3.1, for any gradient flow μ for $J, t \mapsto J(\mu_t)$ is nonincreasing (see (3.9)) and thus of bounded variation. This then gives us the desired result.

4 Solving the Fokker-Planck Equation (2.10)

For any $\mu_0 \in \mathcal{P}_2(\mathbb{R}^d)$, in view of Proposition 3.1 and Definitions 3.3 and 3.2, there is a solution $\{\boldsymbol{\mu}_t\}_{t>0}$ to the Fokker-Planck equation (3.4) with $\lim_{t\downarrow 0} \boldsymbol{\mu}_t = \mu_0$ and its velocity field v_t lies in the negative Fréchat subdifferential $-\partial J(\mu_t)$ for all t > 0. A key observation is that if we additionally have " $\mu_t \ll \mathcal{L}^d$ for all $t \ge 0$," as $\partial J(\mu_t)$ in this case reduces to a singleton containing only $\nabla \phi_{\mu_t}^{\mu_d}$ for all $t \ge 0$ (Lemma 2.1), the velocity field v_t must equal $-\nabla \phi_{\mu_t}^{\mu_d}$. This turns the general Fokker-Planck equation (3.4) into the specific one (2.10) we aim to solve. Hence, the problem boils down to: If we take $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, does the gradient flow $\boldsymbol{\mu}$ in Proposition 3.1 fulfill $\boldsymbol{\mu}_t \in \mathcal{P}_2^r(\mathbb{R}^d)$ for all t > 0?

In this section, we will provide an affirmative answer to this. First, for any $\mu_0 \in \mathcal{P}_2(\mathbb{R}^d)$, we will construct a curve in $\mathcal{P}_2(\mathbb{R}^d)$ that moves μ_0 to μ_d at a constant speed. By its explicit construction, we find that as long as $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, the entire curve will remain in $\mathcal{P}_2^r(\mathbb{R}^d)$, except possibly the end point $\mu_d \in \mathcal{P}_2(\mathbb{R}^d)$; see Lemma 4.2. Next, by a suitable change of speed, the rescaled curve will exactly coincide with the gradient flow in Proposition 3.1; see Theorem 4.1.

To begin with, for any $\mu_0 \in \mathcal{P}_2(\mathbb{R}^d)$, take $\gamma \in \Gamma_0(\mu_0, \mu_d)$ and consider

$$\boldsymbol{\beta}_{s} := \left((1-s)\pi^{1} + s\pi^{2} \right)_{\#} \gamma \in \mathcal{P}_{2}(\mathbb{R}^{d}), \quad s \in [0,1].$$
(4.1)

Remark 4.1. By [1, Theorem 7.2.2], $\{\beta_s\}_{s \in [0,1]}$ in (4.1) is a constant-speed geodesic in $\mathcal{P}_2(\mathbb{R}^d)$ connecting μ_0 and μ_d , i.e., it satisfies, per [1, Definition 2.4.2],

$$W_2(\boldsymbol{\beta}_{s_1}, \boldsymbol{\beta}_{s_2}) = (s_2 - s_1)W_2(\mu_0, \mu_d), \quad \forall 0 \le s_1 \le s_2 \le 1.$$
(4.2)

To change the speed of β from constant to exponentially decreasing, we consider

$$\boldsymbol{\mu}_t^* := \boldsymbol{\beta}_{1-e^{-t}} = \left(e^{-t}\pi^1 + (1-e^{-t})\pi^2\right)_{\#}\gamma, \quad t \in [0,\infty).$$
(4.3)

Lemma 4.1. For any $\mu_0 \in \mathcal{P}_2(\mathbb{R}^d)$, $\mu^* \in AC^p((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$ for all $p \ge 1$. Moreover,

$$|(\boldsymbol{\mu}^*)'|(t) = e^{-t}W_2(\mu_0, \mu_d)$$
 and $\frac{d}{dt}J(\boldsymbol{\mu}^*_t) = -e^{-2t}W_2^2(\mu_0, \mu_d)$ $\forall t > 0$

where $|(\boldsymbol{\mu}^*)'|(t)$ is the metric derivative defined in (3.2) and $J: \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}^d$ is given by (2.7).

Proof. For any $0 < t_1 \le t_2 < \infty$, by the definition of μ^* and (4.2),

$$W_{2}(\boldsymbol{\mu}_{t_{1}}^{*}, \boldsymbol{\mu}_{t_{2}}^{*}) = W_{2}\left(\boldsymbol{\beta}_{1-e^{-t_{1}}}, \boldsymbol{\beta}_{1-e^{-t_{2}}}\right)$$

= $-(e^{-t_{2}} - e^{-t_{1}})W_{2}(\mu_{0}, \mu_{d}) = \int_{t_{1}}^{t_{2}} e^{-t}W_{2}(\mu_{0}, \mu^{d}) dt.$ (4.4)

As $e^{-t} \in L^p(0,\infty)$ for all $p \geq 1$, we conclude $\mu^* \in AC^p((0,\infty);\mathcal{P}_2(\mathbb{R}^d))$ for all $p \geq 1$; recall Definition 3.1. For any t > 0, by (3.2) and using (4.2) in a similar way,

$$|(\boldsymbol{\mu}^*)'|(t) = \lim_{t' \to t} \frac{W_2(\boldsymbol{\mu}_{t'}^*, \boldsymbol{\mu}_t^*)}{|t' - t|} = \lim_{t' \to t} \frac{|e^{-t'} - e^{-t}|W_2(\mu_0, \mu_d)}{|t' - t|}$$
$$= W_2(\mu_0, \mu_d) \left| \left(\frac{d}{ds} e^{-s}\right) \right|_{s=t} \right| = e^{-t} W_2(\mu_0, \mu_d).$$

On the other hand,

$$\frac{d}{dt}J(\boldsymbol{\mu}_{t}^{*}) = \frac{1}{2}\frac{d}{dt}W_{2}^{2}(\boldsymbol{\mu}_{t}^{*},\boldsymbol{\mu}_{d}) = \frac{1}{2}\lim_{h\to 0}\frac{W_{2}^{2}(\boldsymbol{\mu}_{t+h}^{*},\boldsymbol{\mu}_{d}) - W_{2}^{2}(\boldsymbol{\mu}_{t}^{*},\boldsymbol{\mu}_{d})}{h} \\
= \frac{1}{2}\lim_{h\to 0}\frac{e^{-2(t+h)}W_{2}^{2}(\boldsymbol{\mu}_{0},\boldsymbol{\mu}_{d}) - e^{-2t}W_{2}^{2}(\boldsymbol{\mu}_{0},\boldsymbol{\mu}_{d})}{h} \\
= \frac{1}{2}W_{2}^{2}(\boldsymbol{\mu}_{0},\boldsymbol{\mu}_{d})\left(\frac{d}{ds}e^{-2s}\right)\Big|_{s=t} = -e^{-2t}W_{2}^{2}(\boldsymbol{\mu}_{0},\boldsymbol{\mu}_{d}), \quad (4.5)$$
second line follows from (4.3), $\boldsymbol{\beta}_{1} = \boldsymbol{\mu}_{d}$, and (4.2).

where the second line follows from (4.3), $\beta_1 = \mu_d$, and (4.2).

Now, let us assume additionally that $\mu_0 \ll \mathcal{L}^d$ (i.e., $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$). By Remarks 2.2 and 2.3, an optimal transport map $\mathbf{t}_{\mu_0}^{\mu_d} : \mathbb{R}^d \to \mathbb{R}^d$ from μ_0 to μ_d exists and $\gamma = (\mathbf{i} \times \mathbf{t}_{\mu_0}^{\mu_d})_{\#} \mu_0$ is the unique element in $\Gamma_0(\mu_0, \mu_d)$. Hence, $\{\boldsymbol{\beta}_s\}_{s \in [0,1]}$ in (4.1) now takes the more concrete form

$$\boldsymbol{\beta}_{s} = \left((1-s)\mathbf{i} + s\mathbf{t}_{\mu_{0}}^{\mu_{d}} \right)_{\#} \mu_{0} \in \mathcal{P}_{2}(\mathbb{R}^{d}), \quad s \in [0,1].$$
(4.6)

Furthermore, the entire curve β now lies in $\mathcal{P}_2^r(\mathbb{R}^d)$, except possibly the right endpoint.

Lemma 4.2. As long as $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, we have $\beta_s \in \mathcal{P}_2^r(\mathbb{R}^d)$ for all $s \in [0, 1)$.

Proof. For any fixed $s \in (0,1)$, thanks to (4.6) and (2.4), β_s can be equivalently expressed as

$$\boldsymbol{\beta}_s = (\mathbf{i} - s \nabla \phi_{\mu_0}^{\mu_d})_{\#} \mu_0. \tag{4.7}$$

Recall from Remark 2.1 that $f(x) := \frac{1}{2}|x|^2 - \phi_{\mu_0}^{\mu_d}(x), x \in \mathbb{R}^d$, is convex. It follows that $f_s(x) := \frac{1}{2}|x|^2 - \frac{1}{2}|x|^2 - \frac{1}{2}|x|^2 - \frac{1}{2}|x|^2 + \frac{1}$ $\frac{1}{2}|x|^2 - s\phi^{\mu_d}_{\mu_0}(x), x \in \mathbb{R}^d$, is strictly (in fact, uniformly) convex. Indeed, f_s can be expressed as

$$f_s(x) = \frac{1}{2}|x|^2 - s\left(\frac{1}{2}|x|^2 - f(x)\right) = \frac{1}{2}(1-s)|x|^2 + sf(x).$$

where $\frac{1}{2}(1-s)|x|^2$ is uniformly convex and sf(x) is convex. Hence, by [1, Theorem 5.5.4], there exists a \mathcal{L}^d -negligible $S \subset \mathbb{R}^d$ such that ∇f_s exists and is injective with $|\det \nabla^2 f_s| > 0$ on $\mathbb{R}^d \setminus S$. We can then apply [1, Lemma 5.5.3] to conclude that $(\nabla f_s)_{\#}\mu_0 \ll \mathcal{L}^d$. Finally, by the definition of f_s and Remark 2.1, $\nabla f_s = \mathbf{i} - s \nabla \phi_{\mu_0}^{\mu_d} \mu_0$ -a.s. This, together with (4.7), implies $\beta_s = (\nabla f_s)_{\#} \mu_0 \ll \mathcal{L}^d$. \Box

For any $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, thanks to the concrete form of β in (4.6), μ^* defined in (4.3) becomes

$$\boldsymbol{\mu}_{t}^{*} = \boldsymbol{\beta}_{1-e^{-t}} = \left(e^{-t}\mathbf{i} + (1-e^{-t})\mathbf{t}_{\mu_{0}}^{\mu_{d}}\right)_{\#}\mu_{0}, \quad t \in [0,\infty).$$
(4.8)

In fact, this exactly coincides with the unique gradient flow for $J: \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ in (2.7) with $\lim_{t \to 0} \mu_t = \mu_0$. While the general existence of such a gradient flow is known from Proposition 3.1, we show here that it can be explicitly constructed as a time-changed constant-speed geodesic.

Theorem 4.1. For any $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, μ^* defined in (4.3) satisfies the following:

(i) $\boldsymbol{\mu}^* \in \mathcal{P}_2^r(\mathbb{R}^d)$ for all $t \ge 0$;

(ii) μ^* is the unique gradient flow for $J: \mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ in (2.7) with $\lim_{t \downarrow 0} \mu_t = \mu_0$.

Proof. From Lemma 4.2, we immediately see that $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$ implies $\boldsymbol{\mu}_t^* = \boldsymbol{\beta}_{1-e^{-t}} \in \mathcal{P}_2^r(\mathbb{R}^d)$ for all $t \geq 0$. By the definitions of $\boldsymbol{\mu}^*$ and $\boldsymbol{\beta}$,

$$\lim_{t \downarrow 0} W_2(\boldsymbol{\mu}_t^*, \mu_0) = \lim_{t \downarrow 0} W_2(\boldsymbol{\beta}_{1-e^{-t}}, \boldsymbol{\beta}_0) = \lim_{t \downarrow 0} (1-e^{-t}) W_2(\mu_0, \mu_d) = 0,$$

where the second equality follows from (4.2). That is, $\lim_{t\downarrow 0} \mu_t^* = \mu_0$. Hence, it remains to show that μ^* is a gradient flow for J in (2.7). As $\mu^* \in AC^2((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$ (recall Lemma 4.1), it suffices to prove that μ^* satisfies (3.11), thanks to Proposition 3.2. For any t > 0, as $\frac{d}{dt}J(\mu_t^*)$ and $|(\mu^*)'|(t)$ have been derived explicitly in Lemma 4.1, it remains to compute $|\partial J|(\mu_t^*)$. By [1, Theorem 10.4.12],

$$|\partial J|(\boldsymbol{\mu}_t^*) = \min\left\{\int_{\mathbb{R}^d} |\bar{\gamma} - \mathbf{i}|^2 d\boldsymbol{\mu}_t^* : \gamma \in \Gamma_0(\boldsymbol{\mu}_t^*, \boldsymbol{\mu}_d)\right\}^{1/2},$$

where $\bar{\gamma} : \mathbb{R}^d \to \mathbb{R}^d$ is the barycentric projection of $\gamma \in \Gamma_0(\boldsymbol{\mu}_t^*, \mu_d)$, defined by

$$\bar{\gamma}(x) := \int_{\mathbb{R}^d} y d\gamma (dy \mid x) \quad \text{for } \mu_t^* \text{-a.e. } x \in \mathbb{R}^d,$$

with $\gamma(dy \mid x)$ denoting the conditional probability of $\gamma(\cdot, \cdot) \in \mathcal{P}(\mathbb{R}^d \times \mathbb{R}^d)$ given that the first entry is known to be some $x \in \mathbb{R}^d$. As $\mu_t^* \in \mathcal{P}_2^r(\mathbb{R}^d)$, $\Gamma_0(\mu_t^*, \mu_d)$ is a singleton containing only $\gamma = (\mathbf{i} \times \mathbf{t}_{\mu_t^*}^{\mu_d})_{\#} \mu_t^*$ (Remark 2.2). It follows that $\gamma(dy \mid x)$ is the dirac measure concentrated on the point $\mathbf{t}_{\mu_t^*}^{\mu_d}(x)$, such that $\bar{\gamma}(x) = \mathbf{t}_{\mu_t^*}^{\mu_d}(x)$, for μ_t^* -a.e. $x \in \mathbb{R}^d$. This in turn implies that

$$|\partial J|(\boldsymbol{\mu}_t^*) = \left(\int_{\mathbb{R}^d} |\mathbf{t}_{\boldsymbol{\mu}_t^*}^{\mu_{\rm d}} - \mathbf{i}|^2 d\boldsymbol{\mu}_t^*\right)^{1/2} = W_2(\boldsymbol{\mu}_t^*, \mu_{\rm d}) = e^{-t} W_2(\mu_0, \mu_{\rm d}), \tag{4.9}$$

where the last equality follows from (4.3), $\beta_1 = \mu_d$, and (4.2). By Lemma 4.1 and (4.9),

$$\frac{d}{dt}J(\boldsymbol{\mu}_t^*) = -e^{-2t}W_2^2(\mu_0,\mu_d) = -(e^{-t}W_2(\mu_0,\mu_d))^2 = -\frac{1}{2}\left(|\partial J|^2(\boldsymbol{\mu}_t^*) + |(\boldsymbol{\mu}^*)'|^2(t)\right).$$

That is, (3.11) is fulfilled, as desired.

Now, we are ready to show that a solution to the Fokker-Planck equation (2.10) exists and it is unique up to time-marginal laws.

Corollary 4.1. For any $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, μ^* defined in (4.3) satisfies

$$\nabla \phi_{\boldsymbol{\mu}_{t}^{*}}^{\mu_{d}} \in L^{2}(\mathbb{R}^{d}, \boldsymbol{\mu}_{t}^{*}) \quad and \quad \|\nabla \phi_{\boldsymbol{\mu}_{t}^{*}}^{\mu_{d}}\|_{L^{2}(\mathbb{R}^{d}, \boldsymbol{\mu}_{t}^{*})} = |(\boldsymbol{\mu}^{*})'|(t) \quad for \ \mathcal{L}^{1}\text{-}a.e. \ t > 0$$
(4.10)

and it is a solution to the Fokker-Planck equation (2.10). For any $\boldsymbol{\mu} \in AC^2_{loc}((0,\infty), \mathcal{P}_2(\mathbb{R}^d))$ with $\boldsymbol{\mu}_t \in \mathcal{P}_2^r(\mathbb{R}^d)$ for all t > 0, if it satisfies (4.10) (with $\boldsymbol{\mu}$ in place of $\boldsymbol{\mu}^*$) and is a solution to (2.10), then $W_2(\boldsymbol{\mu}_t, \boldsymbol{\mu}_t^*) = 0$ for all $t \ge 0$.

Proof. By Theorem 4.1 (ii), $\boldsymbol{\mu}^*$ is the unique gradient flow for J with $\lim_{t\downarrow 0} \boldsymbol{\mu}^*_t = \mu_0$. In view of Definitions 3.3 and 3.2, this implies that $\boldsymbol{\mu}^*$ is a solution to the Fokker-Planck equation (3.4), where $v: (0, \infty) \times \mathbb{R}^d \to \mathbb{R}^d$ therein is the velocity field of $\boldsymbol{\mu}^*$ and it satisfies (3.3) and $v_t \in -\partial J(\boldsymbol{\mu}^*_t)$ for all t > 0. As $\boldsymbol{\mu}^*_t \in \mathcal{P}^r_2(\mathbb{R}^d)$ for all t > 0 (by Theorem 4.1 (i)), we know from Lemma 2.1 that $\partial J(\boldsymbol{\mu}^*_t)$ is a

singleton containing only $\nabla \phi_{\mu_t}^{\mu_d}$ for all t > 0. Hence, $v_t = -\nabla \phi_{\mu_t}^{\mu_d}$ must hold for all t > 0, such that (3.3) and (3.4) become (4.10) and (2.10), respectively. Conversely, given $\boldsymbol{\mu} \in AC_{\text{loc}}^2((0,\infty), \mathcal{P}_2(\mathbb{R}^d))$ with $\boldsymbol{\mu}_t \in \mathcal{P}_2^r(\mathbb{R}^d)$ for all t > 0, we can argue as above that the velocity field v of $\boldsymbol{\mu}$ must satisfy $v_t = -\nabla \phi_{\mu_t}^{\mu_d}$ and $\nabla \phi_{\mu_t}^{\mu_d}$ is the unique element in $\partial J(\boldsymbol{\mu}_t)$ for all t > 0. Hence, if $\boldsymbol{\mu}$ satisfies (4.10) (with $\boldsymbol{\mu}$ in place of $\boldsymbol{\mu}^*$) and is a solution to (2.10), it automatically fulfills (3.3) and (3.4). This means, by Definitions 3.3 and 3.2, that $\boldsymbol{\mu}$ is a gradient flow for J in (2.7) with $\lim_{t \downarrow 0} \boldsymbol{\mu}_t = \mu_0$. The uniqueness part of Proposition 3.1 then yields $W_2(\boldsymbol{\mu}_t, \boldsymbol{\mu}_t^*) = 0$ for all $t \ge 0$.

5 Solving the Gradient-descent ODE (2.9)

In this section, relying on the (unique) solution μ^* to the Fokker-Planck equation (2.10) (recall Corollary 4.1), we will first construct a (unique) solution Y to ODE (2.9) on the strength of Trevisan's superposition principle [16]. Next, to facilitate numerical simulation of ODE (2.9), we propose an Euler scheme and show that it does converge to the laws of $\{Y_t\}_{t\geq 0}$ (i.e., μ^*) in the limit.

To state precisely what forms a solution to ODE (2.9), consider the space of continuous paths

 $(\Omega, \mathcal{F}) := (C([0, \infty); \mathbb{R}^d), \mathcal{B}(C([0, \infty); \mathbb{R}^d)).$

Definition 5.1. A process $Y : [0, \infty) \times \Omega \to \mathbb{R}^d$ is said to be a solution to (2.9) if $Y_t(\omega) = \omega(t)$ for all $(t, \omega) \in [0, \infty) \times \Omega$ and there exists a probability measure \mathbb{P} on (Ω, \mathcal{F}) under which

- (i) $\boldsymbol{\mu}_t := \boldsymbol{\mu}^{Y_t}$, the law of $Y_t : \Omega \to \mathbb{R}^d$, belongs to $\mathcal{P}_2^r(\mathbb{R}^d)$ for all $t \ge 0$ with $\boldsymbol{\mu}_0 = \boldsymbol{\mu}_0$;
- (i) the collection of paths $\left\{\omega \in \Omega: \omega(t) = \omega(0) \int_0^t \nabla \phi_{\mu_s}^{\mu_d}(\omega(s)) ds, t \ge 0 \right\}$ has probability one.

Proposition 5.1. For any $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, there exists a solution Y to ODE (2.9) such that $\mu^{Y_t} = \mu_t^*$ for all $t \ge 0$, with μ^* defined as in (4.3).

Proof. In view of (4.3), as $\mu_t^* = \beta_{1-e^{-t}}$ for all $t \ge 0$, where β is a constant-speed geodesic (i.e., satisfying (4.2)), we immediately conclude that $t \mapsto \mu_t^*$ is continuous under the W_2 distance, and thus narrowly continuous. Also, by Corollary 4.1, μ^* is a solution to the Fokker-Planck equation (2.10). Hence, to apply the "superposition principle" (i.e., [16, Theorem 2.5]), it remains to verify equation (2.3) in [16], which in our case takes the form

$$\int_0^T \int_{\mathbb{R}^d} |\nabla \phi_{\boldsymbol{\mu}_t^*}^{\mu_d}(y)| d\boldsymbol{\mu}_t^*(y) dt < \infty \quad \forall T > 0,$$
(5.1)

where $\phi_{\boldsymbol{\mu}_t^*}^{\mu_{\mathrm{d}}}$ is a Kantorovich potential from $\boldsymbol{\mu}_t^*$ to μ_{d} . For any T > 0, note that

$$\int_0^T \int_{\mathbb{R}^d} |\nabla \phi_{\boldsymbol{\mu}_t^*}^{\mu_d}(y)| d\boldsymbol{\mu}_t^*(y) dt = \int_0^T \|\nabla \phi_{\boldsymbol{\mu}_t^*}^{\mu_d}\|_{L^1(\mathbb{R}^d, \boldsymbol{\mu}_t^*)} dt \le \int_0^T \|\nabla \phi_{\boldsymbol{\mu}_t^*}^{\mu_d}\|_{L^2(\mathbb{R}^d, \boldsymbol{\mu}_t^*)} dt \le \int_0^T |(\boldsymbol{\mu}^*)'|(t) dt,$$

where the last inequality stems from (4.10). As $\mu^* \in AC((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$ by Lemma 4.1, Remark 3.2 indicates $|(\mu^*)'| \in L^1((0,\infty))$. This, along with the previous inequality, implies that (5.1) holds.

Consider $Y_t(\omega) := \omega(t)$ for all $(t, \omega) \in [0, \infty) \times \Omega$. By [16, Theorem 2.5], there is a probability \mathbb{P} on (Ω, \mathcal{F}) such that (i) $\mathbb{P} \circ (Y_t)^{-1} = \boldsymbol{\mu}_t^*$ for all $t \ge 0$, and (ii) for any $f \in C^{1,2}((0, \infty) \times \mathbb{R}^d)$,

$$t \mapsto f(t,\omega(t)) - f(0,\omega(0)) - \int_0^t \left(\partial_s f - \nabla \phi_{\boldsymbol{\mu}_s^*}^{\mu_d} \cdot \nabla f\right)(s,\omega(s))ds, \quad t \ge 0$$
(5.2)

is a local martingale under \mathbb{P} . As $\mu_t^* \in \mathcal{P}_2^r(\mathbb{R}^d)$ for all $t \ge 0$ (by Theorem 4.1), (i) above already implies that Definition 5.1 (i) is satisfied. For each $i \in \{1, 2, ..., d\}$, by taking $f(t, y) = y_i$ for

 $y = (y_1, y_2, ..., y_d) \in \mathbb{R}^d$ in (5.2), we see that $t \mapsto M_t^{(i)}(\omega) := (\omega(t))_i - (\omega(0))_i + \int_0^t \partial_{y_i} \phi_{\mu_s^*}^{\mu_d}(\omega(s)) ds$, $t \ge 0$, is a local martingale under \mathbb{P} . By the same argument on p. 315 of [10], we find that the quadratic variation of $M^{(i)}$ is constantly zero, i.e., $\langle M^{(i)} \rangle_t = 0$ for all $t \ge 0$. As $M^{(i)}$ is a local martingale with zero quadratic variation under \mathbb{P} , $M_t^{(i)} = 0$ for all $t \ge 0$ P-a.s. That is,

$$(\omega(t))_i = (\omega(0))_i - \int_0^t \partial_{y_i} \phi_{\boldsymbol{\mu}_s^*}^{\mu_d}(\omega(s)) ds \quad t \ge 0, \qquad \text{for } \mathbb{P}\text{-a.e. } \omega \in \Omega.$$

Because this holds for all $i \in \{1, 2, ..., d\}$, we conclude that

$$\omega(t)=\omega(0)-\int_0^t \nabla \phi_{\boldsymbol{\mu}^s_s}^{\mu_{\mathrm{d}}}(\omega(s)) ds \quad t\geq 0, \quad \text{for \mathbb{P}-a.e. $\omega\in\Omega$},$$

which shows that Definition 5.1 (ii) is fulfilled. Hence, Y is a solution to (2.9).

In fact, under appropriate regularity conditions, the time-marginal laws of any solution Y to ODE (2.9) must coincide with μ^* in (4.3).

Proposition 5.2. Given $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, let Y be a solution to (2.9) and consider $\boldsymbol{\mu}_t := \mu^{Y_t} \in \mathcal{P}_2^r(\mathbb{R}^d)$ for all $t \ge 0$. If $\boldsymbol{\mu}$ belongs to $AC((0,\infty); \mathcal{P}_2(\mathbb{R}^d)) \cap AC^2_{loc}((0,\infty); \mathcal{P}_2(\mathbb{R}^d))$ and satisfies (4.10) (with $\boldsymbol{\mu}$ in place of $\boldsymbol{\mu}^*$), then $W_2(\boldsymbol{\mu}_t, \boldsymbol{\mu}_t^*) = 0$ for all $t \ge 0$, with $\boldsymbol{\mu}^*$ given by (4.3).

Proof. Let \mathbb{P} be the probability measure on (Ω, \mathcal{F}) associated with the solution Y to (2.9); recall Definition 5.1. As $\boldsymbol{\mu}$ satisfies (4.10) (with $\boldsymbol{\mu}$ in place of $\boldsymbol{\mu}^*$) and belongs to $AC((0, \infty); \mathcal{P}_2(\mathbb{R}^d))$, we may follow the same argument below (5.1) to show that

$$\int_0^T \int_{\mathbb{R}^d} |\nabla \phi_{\boldsymbol{\mu}_t}^{\mu_{\mathrm{d}}}(y)| d\boldsymbol{\mu}_t(y) dt < \infty \quad \forall T > 0.$$
(5.3)

For any $\psi \in C_c^{1,2}((0,\infty) \times \mathbb{R}^d)$, take T > 0 large such that $\psi(T,y) = \psi(0,y) = 0$ for all $y \in \mathbb{R}^d$. With $Y_t(\omega) = \omega(t)$ for all $(t,\omega) \in [0,\infty) \times \Omega$, we deduce from Definition 5.1 (ii) that $0 = \psi(T,Y_T) - \psi(0,Y_0) = \int_0^T \left(\partial_t \psi - \nabla \phi_{\mu_t}^{\mu_d} \cdot \nabla \psi\right)(t,Y_t) dt$ P-a.s. Taking expectation under \mathbb{P} on both sides gives

$$0 = \mathbb{E}\bigg[\int_0^T \left(\partial_t \psi - \nabla \phi_{\mu_t}^{\mu_d} \cdot \nabla \psi\right)(t, Y_t) dt\bigg] = \int_0^T \int_{\mathbb{R}^d} \left(\partial_t \psi - \nabla \phi_{\mu_t}^{\mu_d} \cdot \nabla \psi\right)(t, y) d\mu_t(y) dt,$$

where the second equality follows from Fubini's theorem, which is applicable here thanks to (5.3). As $T \to \infty$, we see that $\boldsymbol{\mu}$ fulfills the Fokker-Planck equation (2.10). Given that $\boldsymbol{\mu}$ belongs to $AC_{loc}^2((0,\infty);\mathcal{P}_2(\mathbb{R}^d))$, satisfies (4.10) (with $\boldsymbol{\mu}$ in place of $\boldsymbol{\mu}^*$), and solves (2.10), we conclude from Corollary 4.1 that $W_2(\boldsymbol{\mu}_t, \boldsymbol{\mu}_t^*) = 0$ for all $t \geq 0$.

Now, we are ready to present the main theoretic result of this paper.

Theorem 5.1. For any $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, there exists a solution Y to ODE (2.9) such that $\boldsymbol{\mu}_t := \mu^{Y_t}$ belongs to $\mathcal{P}_2^r(\mathbb{R}^d)$ for all $t \ge 0$ and satisfies

$$\boldsymbol{\mu} \in AC((0,\infty); \mathcal{P}_2(\mathbb{R}^d)) \cap AC^2_{\text{loc}}((0,\infty); \mathcal{P}_2(\mathbb{R}^d)),$$
(5.4)

$$\|\nabla \phi_{\mu_t}^{\mu_d}\|_{L^2(\mathbb{R}^d,\mu_t)} = |\mu'|(t) \quad \text{for } \mathcal{L}^1 \text{-a.e. } t > 0.$$
(5.5)

Moreover, for any such a solution Y to ODE (2.9), $W_2(\mu^{Y_t}, \mu_d) \downarrow 0$ as $t \to \infty$; specifically,

$$W_2(\mu^{Y_t}, \mu_d) \le e^{-t} W_2(\mu_0, \mu_d) \quad \forall t \ge 0.$$

Proof. Recall from Lemma 4.1 and Corollary 4.1 that μ^* defined in (4.3) fulfills (5.4) and (5.5) (with μ^* in place of μ). Hence, the solution Y to ODE (2.9) constructed in Proposition 5.1 readily satisfies all the conditions in the theorem. As μ^* is the unique gradient flow for J in (2.7) with $\lim_{t \to 0} \mu_t^* = \mu_0$ (by Theorem 4.1), we know from Proposition 3.1 that

$$J(\boldsymbol{\mu}_t^*) \downarrow 0 \text{ as } t \to \infty \quad \text{and} \quad J(\boldsymbol{\mu}_t^*) \le e^{-2t} J(\mu_0) \quad \forall t \ge 0.$$
 (5.6)

For any solution Y to ODE (2.9) that fulfills all the conditions in the theorem, Proposition 5.2 asserts that $\mu^{Y_t} = \boldsymbol{\mu}_t^*$ for all $t \ge 0$. By (2.7), this implies $W_2(\mu^{Y_t}, \mu_d) = W_2(\boldsymbol{\mu}_t^*, \mu_d) = (2J(\boldsymbol{\mu}_t^*))^{1/2}$ for all $t \ge 0$. Thus, (5.6) implies $W_2(\mu^{Y_t}, \mu_d) \downarrow 0$ and $W_2(\mu^{Y_t}, \mu_d) \le e^{-t}W_2(\mu_0, \mu_d)$ for all $t \ge 0$. \Box

Theorem 5.1 suggests that one can uncover the true data distribution $\mu_d \in \mathcal{P}_2(\mathbb{R}^d)$ by simulating the distribution-dependent ODE (2.9). To facilitate the numerical implementation of this, we now propose a discretization of (2.9) and closely study its convergence as the time step diminishes.

5.1 An Euler Scheme

As ODE (2.9) is distribution-dependent, when we discretize it, we need to simulate a random variable and keep track of its law at every discrete time point. Specifically, fix a time step $0 < \varepsilon < 1$. At time 0, we simulate a random variable Y_0^{ε} according to a given initial distribution, i.e., $\mu^{Y_0^{\varepsilon}} = \mu_{0,\varepsilon} :=$ $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$. At time ε , we simulate the random variable

$$Y_1^{\varepsilon} := Y_0^{\varepsilon} - \varepsilon \nabla \phi_{\mu_0}^{\mu_d}(Y_0^{\varepsilon}), \quad \text{such that} \quad \mu^{Y_1^{\varepsilon}} = \mu_{1,\varepsilon} := (\mathbf{i} - \varepsilon \nabla \phi_{\mu_0}^{\mu_d})_{\#} \mu_0,$$

where $\phi_{\mu_0}^{\mu_d}$ is a Kantorovich potential from μ_0 to μ_d . It is crucial that μ_0 lies in $\mathcal{P}_2^r(\mathbb{R}^d)$, instead of the more general $\mathcal{P}_2(\mathbb{R}^d)$. Indeed, as $\nabla \phi_{\mu_0}^{\mu_d}$ is only well-defined \mathcal{L}^d -a.e. (Remark 2.1), $\mu_0 \ll \mathcal{L}^d$ ensures that Y_1^{ε} above is well-defined. Moreover, with $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, we may apply the same arguments below (4.7) and find that $\mu_{1,\varepsilon}$ remains in $\mathcal{P}_2^r(\mathbb{R}^d)$. At time 2ε , we simulate the random variable

$$Y_2^{\varepsilon} := Y_1^{\varepsilon} - \varepsilon \nabla \phi_{\mu_{1,\varepsilon}}^{\mu_{\rm d}}(Y_1^{\varepsilon}), \quad \text{such that} \quad \mu_{Y_2^{\varepsilon}}^{Y_2^{\varepsilon}} = \mu_{2,\varepsilon} := (\mathbf{i} - \varepsilon \nabla \phi_{\mu_{1,\varepsilon}}^{\mu_{\rm d}})_{\#} \mu_{1,\varepsilon},$$

where $\phi_{\mu_{1,\varepsilon}}^{\mu_{d}}$ is a Kantorovich potential from $\mu_{1,\varepsilon}$ to μ_{d} . Again, $\mu_{1,\varepsilon} \in \mathcal{P}_{2}^{r}(\mathbb{R}^{d})$ ensures that Y_{2}^{ε} above is well-defined and allows the use of the arguments below (4.7), which in turn guarantees $\mu_{2,\varepsilon} \in \mathcal{P}_{2}^{r}(\mathbb{R}^{d})$. Continuing this procedure, we simulate for all $n \in \mathbb{N}$ the random variables

$$Y_n^{\varepsilon} := Y_{n-1}^{\varepsilon} - \varepsilon \nabla \phi_{\mu_{n-1,\varepsilon}}^{\mu_{\mathrm{d}}}(Y_{n-1}^{\varepsilon}), \quad \text{such that} \quad \mu_n^{Y_n^{\varepsilon}} = \mu_{n,\varepsilon} := (\mathbf{i} - \varepsilon \nabla \phi_{\mu_{n-1,\varepsilon}}^{\mu_{\mathrm{d}}})_{\#} \mu_{n-1,\varepsilon}, \tag{5.7}$$

and we know $\mu_{n,\varepsilon} \in \mathcal{P}_2^r(\mathbb{R}^d)$ for all $n \in \mathbb{N}$. This allows us to invoke Remark 2.2, particularly (2.4), to express $\{\mu_{n,\varepsilon}\}_{n\in\mathbb{N}}$ equivalently as

$$\mu_{n,\varepsilon} = \left(\mathbf{i} + \varepsilon (\mathbf{t}^{\mu_{\mathrm{d}}}_{\mu_{n-1,\varepsilon}} - \mathbf{i})\right)_{\#} \mu_{n-1,\varepsilon} \quad \forall n \in \mathbb{N},$$
(5.8)

using their associated optimal transport maps. Interestingly, how $\mu_{n-1,\varepsilon}$ is transported to $\mu_{n,\varepsilon}$ in (5.8) is already optimal, as the next result shows.

Lemma 5.1. Fix $0 < \varepsilon < 1$. For any $n \in \mathbb{N}$, $\mathbf{i} + \varepsilon(\mathbf{t}_{\mu_{n-1,\varepsilon}}^{\mu_d} - \mathbf{i}) : \mathbb{R}^d \to \mathbb{R}^d$ is an optimal transport map from $\mu_{n-1,\varepsilon}$ to $\mu_{n,\varepsilon}$. That is, $\mathbf{t}_{\mu_{n-1,\varepsilon}}^{\mu_{n,\varepsilon}} = \mathbf{i} + \varepsilon(\mathbf{t}_{\mu_{n-1,\varepsilon}}^{\mu_d} - \mathbf{i})$ for all $n \in \mathbb{N}$.

Proof. For any $n \in \mathbb{N}$, by [1, Theorem 7.2.2], $\boldsymbol{\nu}_s := ((1-s)\mathbf{i} + s\mathbf{t}_{\mu_{n-1,\varepsilon}}^{\mu_d})_{\#}\mu_{n-1,\varepsilon}$, $s \in [0,1]$, is a constant-speed geodesic connecting $\mu_{n-1,\varepsilon}$ to μ_d , i.e., it satisfies, per [1, Definition 2.4.2],

$$W_2(\boldsymbol{\nu}_{s_1}, \boldsymbol{\nu}_{s_2}) = (s_2 - s_1) W_2(\mu_{n-1,\varepsilon}, \mu_d), \quad \forall 0 \le s_1 \le s_2 \le 1.$$
(5.9)

Observe from (5.8) that $\mu_{n,\varepsilon} = \nu_{\varepsilon}$. Hence,

$$W_2(\mu_{n-1,\varepsilon},\mu_{n,\varepsilon}) = W_2(\boldsymbol{\nu}_0,\boldsymbol{\nu}_\varepsilon) = \varepsilon W_2(\mu_{n-1,\varepsilon},\mu_d), \qquad (5.10)$$

where the last equality follows from (5.9). Now, set $\mathbf{t} := \mathbf{i} + \varepsilon(\mathbf{t}_{\mu_{n-1,\varepsilon}}^{\mu_d} - \mathbf{i})$ and note that

$$\int_{\mathbb{R}^d} |\mathbf{t} - \mathbf{i}|^2 d\mu_{n-1,\varepsilon} = \varepsilon^2 \int_{\mathbb{R}^d} |\mathbf{t}_{\mu_{n-1,\varepsilon}}^{\mu_{\mathrm{d}}} - \mathbf{i}|^2 d\mu_{n-1,\varepsilon} = \varepsilon^2 W_2^2(\mu_{n-1,\varepsilon},\mu_{\mathrm{d}}) = W_2^2(\mu_{n-1,\varepsilon},\mu_{n,\varepsilon}),$$

where the second equality follows from the fact that $\mathbf{t}_{\mu_{n-1,\varepsilon}}^{\mu_{d}}$ is an optimal transport map from $\mu_{n-1,\varepsilon}$ to μ_{d} and the third equality stems from (5.10). This, together with (5.8), readily shows that $\mathbf{t} := \mathbf{i} + \varepsilon(\mathbf{t}_{\mu_{n-1,\varepsilon}}^{\mu_{d}} - \mathbf{i})$ is an optimal transport map from $\mu_{n-1,\varepsilon}$ to $\mu_{n,\varepsilon}$.

An important consequence of Lemma 5.1 is that the measures $\{\mu_{n,\varepsilon}\}_{n\in\mathbb{N}}$ all lie on the constantspeed geodesic connecting μ_0 and μ_d , i.e., β given in (4.6), or equivalently, on the curve μ^* defined in (4.3). To see this, we consider, for each $0 < \varepsilon < 1$, the unique $t_{\varepsilon} > 0$ such that

$$\varepsilon = 1 - e^{-t_{\varepsilon}}.\tag{5.11}$$

Remark 5.1. It can be checked directly from (5.11) that (i) $t_{\varepsilon} > \varepsilon$ for any $0 < \varepsilon < 1$; (ii) as $\varepsilon \downarrow 0$, we have $t_{\varepsilon} \downarrow 0$ and $\frac{t_{\varepsilon}}{\varepsilon} \to 1$.

Proposition 5.3. For any $0 < \varepsilon < 1$, we have $\mu_{n,\varepsilon} = \beta_{1-e^{-nt_{\varepsilon}}} = \mu_{nt_{\varepsilon}}^*$ for all $n \in \mathbb{N}$, where $t_{\varepsilon} > 0$ is given by (5.11).

Proof. We will prove by induction that

$$\mu_{n,\varepsilon} = \boldsymbol{\beta}_{1-e^{-nt_{\varepsilon}}} = \boldsymbol{\mu}_{nt_{\varepsilon}}^{*} \quad \text{and} \quad \mathbf{t}_{\mu_{0}}^{\mu_{n,\varepsilon}} = \mathbf{i} + (1 - e^{-nt_{\varepsilon}})(\mathbf{t}_{\mu_{0}}^{\mu_{d}} - \mathbf{i}) \qquad \forall n \in \mathbb{N},$$
(5.12)

By (5.8) with n = 1, $\mu_{0,\varepsilon} = \mu_0$, and the definitions of $\boldsymbol{\beta}$ and $\boldsymbol{\mu}^*$, we see that $\mu_{1,\varepsilon} = \boldsymbol{\beta}_{\varepsilon} = \boldsymbol{\beta}_{1-e^{-t\varepsilon}} = \boldsymbol{\mu}_{t_{\varepsilon}}^*$. Also, by Lemma 5.1 and $\mu_{0,\varepsilon} = \mu_0$, we have $\mathbf{t}_{\mu_0}^{\mu_{1,\varepsilon}} = \mathbf{i} + \varepsilon(\mathbf{t}_{\mu_0}^{\mu_d} - \mathbf{i}) = \mathbf{i} + (1 - e^{-t_{\varepsilon}})(\mathbf{t}_{\mu_0}^{\mu_d} - \mathbf{i})$. That is, (5.12) holds for n = 1. Now, suppose that (5.12) holds for some $n = k - 1 \in \mathbb{N}$, i.e.,

$$\mu_{k-1,\varepsilon} = \beta_{1-e^{-(k-1)t_{\varepsilon}}} = \mu^*_{(k-1)t_{\varepsilon}} \quad \text{and} \quad \mathbf{t}^{\mu_{k-1,\varepsilon}}_{\mu_0} = \mathbf{i} + (1 - e^{-(k-1)t_{\varepsilon}})(\mathbf{t}^{\mu_{\mathrm{d}}}_{\mu_0} - \mathbf{i}), \tag{5.13}$$

and we aim to show that (5.12) remains true for n = k. By (5.8), $\mu_{k-1,\varepsilon} = \mathbf{t}_{\mu_0}^{\mu_{k-1,\varepsilon}} \# \mu_0$, and the composition rule for pushforward measures (see e.g., [1, (5.2.4)]),

$$\mu_{k,\varepsilon} = \left(\mathbf{i} + \varepsilon (\mathbf{t}_{\mu_{k-1,\varepsilon}}^{\mu_{d}} - \mathbf{i})\right)_{\#} \mu_{k-1,\varepsilon} = \left(\mathbf{i} + \varepsilon (\mathbf{t}_{\mu_{k-1,\varepsilon}}^{\mu_{d}} - \mathbf{i})\right)_{\#} (\mathbf{t}_{0}^{\mu_{k-1,\varepsilon}}_{\#} + \mu_{0})$$
$$= \left(\mathbf{t}_{\mu_{0}}^{\mu_{k-1,\varepsilon}} + \varepsilon (\mathbf{t}_{\mu_{k-1,\varepsilon}}^{\mu_{d}} \circ \mathbf{t}_{\mu_{0}}^{\mu_{k-1,\varepsilon}} - \mathbf{t}_{\mu_{0}}^{\mu_{k-1,\varepsilon}})\right)_{\#} \mu_{0}.$$

As $\mu_{k-1,\varepsilon}$ lies on the constant-speed geodesic β from μ_0 to μ_d (by the first relation in (5.13)), [1, Lemma 7.2.1] implies that $\mathbf{t}_{\mu_0}^{\mu_d} = \mathbf{t}_{\mu_{k-1,\varepsilon}}^{\mu_d} \circ \mathbf{t}_{\mu_0}^{\mu_{k-1,\varepsilon}}$. Plugging the second relation in (5.13), $\varepsilon = 1 - e^{-t_{\varepsilon}}$, and $\mathbf{t}_{\mu_0}^{\mu_d} = \mathbf{t}_{\mu_{k-1,\varepsilon}}^{\mu_d} \circ \mathbf{t}_{\mu_0}^{\mu_{k-1,\varepsilon}}$ into the previous equality, we get

$$\mu_{k,\varepsilon} = \left(\mathbf{i} + (1 - e^{-(k-1)t_{\varepsilon}})(\mathbf{t}_{\mu_{0}}^{\mu_{d}} - \mathbf{i}) + (1 - e^{-t_{\varepsilon}})(\mathbf{t}_{\mu_{0}}^{\mu_{d}} - \left(\mathbf{i} + (1 - e^{-(k-1)t_{\varepsilon}})(\mathbf{t}_{\mu_{0}}^{\mu_{d}} - \mathbf{i})\right)\right)\right)_{\#} \mu_{0}$$
$$= \left(e^{-kt_{\varepsilon}}\mathbf{i} + (1 - e^{-kt_{\varepsilon}})\mathbf{t}_{\mu_{0}}^{\mu_{d}}\right)_{\#} \mu_{0} = \boldsymbol{\mu}_{kt_{\varepsilon}}^{*} = \boldsymbol{\beta}_{1-e^{-kt_{\varepsilon}}}, \tag{5.14}$$

where the second last equality is due to (4.8). Now, we see that μ_0 , $\mu_{k-1,\varepsilon}$, and $\mu_{k,\varepsilon}$ are all on the the constant-speed geodesic β connecting μ_0 and μ_d , simply at the different time points 0, $1 - e^{-(k-1)t_{\varepsilon}}$, and $1 - e^{-kt_{\varepsilon}}$, respectively. Hence, by a proper change of time, there exists a constantspeed geodesic that connects μ_0 and $\mu_{k,\varepsilon}$ and passes through $\mu_{k-1,\varepsilon}$. By applying [1, Lemma 7.2.1] to this constant-speed geodesic, we have

$$\mathbf{t}_{\mu_{0}}^{\mu_{k,\varepsilon}} = \mathbf{t}_{\mu_{k-1,\varepsilon}}^{\mu_{k,\varepsilon}} \circ \mathbf{t}_{\mu_{0}}^{\mu_{k-1,\varepsilon}} = \mathbf{t}_{\mu_{0}}^{\mu_{k-1,\varepsilon}} + \varepsilon (\mathbf{t}_{\mu_{k-1,\varepsilon}}^{\mu_{d}} \circ \mathbf{t}_{\mu_{0}}^{\mu_{k-1,\varepsilon}} - \mathbf{t}_{\mu_{0}}^{\mu_{k-1,\varepsilon}})
= \mathbf{i} + (1 - e^{-(k-1)t_{\varepsilon}}) (\mathbf{t}_{\mu_{0}}^{\mu_{d}} - \mathbf{i}) + (1 - e^{-t_{\varepsilon}}) (\mathbf{t}_{\mu_{0}}^{\mu_{d}} - \mathbf{i} - (1 - e^{-(k-1)t_{\varepsilon}}) (\mathbf{t}_{\mu_{0}}^{\mu_{d}} - \mathbf{i}))
= e^{-kt_{\varepsilon}} \mathbf{i} + (1 - e^{-kt_{\varepsilon}}) \mathbf{t}_{\mu_{0}}^{\mu_{d}} = \mathbf{i} + (1 - e^{-kt_{\varepsilon}}) (\mathbf{t}_{\mu_{0}}^{\mu_{d}} - \mathbf{i}),$$
(5.15)

where the second equality is due to $\mathbf{t}_{\mu_{k-1,\varepsilon}}^{\mu_{k,\varepsilon}} = \mathbf{i} + \varepsilon(\mathbf{t}_{\mu_{k-1,\varepsilon}}^{\mu_{d}} - \mathbf{i})$ (Lemma 5.1) and the third equality follows from plugging the second relation in (5.13), $\varepsilon = 1 - e^{-t_{\varepsilon}}$, and $\mathbf{t}_{\mu_{0}}^{\mu_{d}} = \mathbf{t}_{\mu_{k-1,\varepsilon}}^{\mu_{d}} \circ \mathbf{t}_{\mu_{0}}^{\mu_{k-1,\varepsilon}}$ into the first line. By (5.14) and (5.15), we see that (5.12) holds for n = k, as desired.

Given $0 < \varepsilon < 1$, we now define a piecewise-constant flow of measures $\mu^{\varepsilon} : [0, \infty) \to \mathcal{P}_2(\mathbb{R}^d)$ by

$$\boldsymbol{\mu}_t^{\varepsilon} := \mu_{n,\varepsilon} \quad \text{if} \quad t \in [n\varepsilon, (n+1)\varepsilon), \qquad \forall n \in \mathbb{N} \cup \{0\}.$$
(5.16)

That is, μ^{ε} is the flow of measures obtained under our Euler scheme (5.7). As $\varepsilon \downarrow 0$, the next result shows that μ^{ε} converges to μ^* defined in (4.8).

Theorem 5.2. For any $\mu_0 \in \mathcal{P}_2^r(\mathbb{R}^d)$, $\lim_{\varepsilon \downarrow 0} W_2(\boldsymbol{\mu}_t^{\varepsilon}, \boldsymbol{\mu}_t^*) = 0$ for all $t \in [0, \infty)$.

Proof. For any $t \in [0, \infty)$, there exists some $n \in \mathbb{N} \cup \{0\}$ such that $t \in [n\varepsilon, (n+1)\varepsilon)$. Then,

$$W_2(\boldsymbol{\mu}_t^{\varepsilon}, \boldsymbol{\mu}_t^*) = W_2(\mu_{n,\varepsilon}, \boldsymbol{\mu}_t^*) = W_2(\boldsymbol{\mu}_{nt_{\varepsilon}}^*, \boldsymbol{\mu}_t^*) = \left(\int_{\min\{nt_{\varepsilon}, t\}}^{\max\{nt_{\varepsilon}, t\}} e^{-s} ds\right) W_2(\mu_0, \mu_d)$$

where the second equality follows from Proposition 5.3 (with $t_{\varepsilon} > 0$ defined as in (5.11)) and the last equality results from a calculation similar to (4.4). As $t_{\varepsilon} > \varepsilon$ (by Remark 5.1) and $t \in [n\varepsilon, (n+1)\varepsilon)$, we have $n\varepsilon \leq \min\{nt_{\varepsilon}, t\} \leq \max\{nt_{\varepsilon}, t\} \leq (n+1)t_{\varepsilon}$. Hence, the previous equality implies

$$W_{2}(\boldsymbol{\mu}_{t}^{\varepsilon},\boldsymbol{\mu}_{t}^{*}) \leq \left(\int_{n\varepsilon}^{(n+1)t_{\varepsilon}} e^{-s} ds\right) W_{2}(\mu_{0},\mu_{d}) = \left(e^{-n\varepsilon} - e^{-(n+1)t_{\varepsilon}}\right) W_{2}(\mu_{0},\mu_{d})$$
$$= e^{-(n+1)t_{\varepsilon}} \left(e^{n(t_{\varepsilon}-\varepsilon)+t_{\varepsilon}} - 1\right) W_{2}(\mu_{0},\mu_{d})$$
$$\leq \left(e^{n\varepsilon\left(\frac{t_{\varepsilon}}{\varepsilon}-1\right)+t_{\varepsilon}} - 1\right) W_{2}(\mu_{0},\mu_{d}) \leq \left(e^{t\left(\frac{t_{\varepsilon}}{\varepsilon}-1\right)+t_{\varepsilon}} - 1\right) W_{2}(\mu_{0},\mu_{d}),$$

where the last inequality follows from $t \ge n\varepsilon$ and $t_{\varepsilon} > \varepsilon$. As $\varepsilon \downarrow 0$, since $t_{\varepsilon} \downarrow 0$ and $\frac{t_{\varepsilon}}{\varepsilon} \to 1$ (by Remark 5.1), the right-hand side above vanishes, which entails $W_2(\boldsymbol{\mu}_t^{\varepsilon}, \boldsymbol{\mu}_t^*) \to 0$.

Recall from Proposition 5.2 that μ^* is the unique flow of probability measures induced by any solution Y to ODE (2.9) under suitable regularity. Hence, Theorem 5.2 stipulates that our Euler scheme (5.7) does converge to the right limit, recovering the time-marginal laws of ODE (2.9).

6 Algorithms

To simulate the distribution-dependent ODE (2.9), we follow the Euler scheme (5.7) in designing Algorithm 1 (called W2-FE, where "FE" means "forward Euler"). The algorithm uses three deep neural networks. One network trains a "generator" $G_{\theta} : \mathbb{R}^d \to \mathbb{R}^d$ to properly represent the evolving time-marginal laws of ODE (2.9). The other two networks, by following [15, Algorithm 1], estimate the Kantorovich potential from the distribution induced by G_{θ} to the data distribution $\mu_{\rm d}$ (denoted by ϕ_w) as well as its *c*-transform (denoted by ψ_v).

How G_{θ} is updated demands a closer examination. Based on the estimated Kantorovich potential ϕ_w, G_{θ} is trained via the Euler scheme (5.7). Given a sample z_i from a prior distribution, the point $y_i = G_{\theta}(z_i)$ represents a sample from $\mu^{Y_n^{\varepsilon}}$. A forward Euler step yields a new point ζ_i , which represents a sample from $\mu^{Y_{n+1}^{\varepsilon}}$. The generator's task is to produce samples indistinguishable from the new points $\{\zeta_i\}$ —or more precisely, to learn the distribution $\mu^{Y_{n+1}^{\varepsilon}}$, represented by the points $\{\zeta_i\}$. To this end, we fix the points $\{\zeta_i\}$ and update G_{θ} by descending the mean square error (MSE) between $\{G_{\theta}(z_i)\}$ and $\{\zeta_i\}$ up to $K \in \mathbb{N}$ times. Let us stress that throughout the K updates of G_{θ} , the points $\{\zeta_i\}$ are kept unchanged. This sets us apart from the standard implementation of stochastic gradient descent (SGD), but for a good reason: as our goal is to learn the distribution represented by $\{\zeta_i\}$, it is important to keep $\{\zeta_i\}$ unchanged for the eventual G_{θ} to more accurately represent $\mu^{Y_{n+1}^{\varepsilon}}$, such that the (discretized) ODE (5.7) is more closely followed.

How we update G_{θ} is reminiscent of *persistent training* in Fischetti et al. [5], a technique consisting of reusing the same minibatch for K consecutive SGD iterations. As shown in [5, Figure 1], a persistency level of five (i.e., K = 5) allows much faster convergence on the CIFAR-10 dataset. Our numerical examples in Section 7 also indicate a similar effect of an enlarged persistency level.

Algorithm 1 W2-FE, our proposed algorithm

Require: Input measures μ^Z, μ_d , batch sizes m, learning rates γ_g, γ_d , regularizer λ , time step Δt , persistency level K, Kantorovich potential pair (ψ_v, ϕ_w) parameterized as deep neural networks, and generator G_θ parameterized as a deep neural network.

for Number of training epochs do for Number of updates do Sample a batch { x_i } from μ_d Sample a batch { z_i } from μ^Z $L_D \leftarrow \frac{1}{m} \sum_{i=1}^{m} \left[\phi_w(G_{\theta}(z_i)) + \psi_v(x_i) - \lambda \left(\phi_w(G_{\theta}(z_i)) + \psi_v(x_i) - \frac{1}{2} ||G_{\theta}(z_i) - x_i||_2^2 \right)_+ \right]$ $w \leftarrow w + \gamma_d \nabla_w L_D$ $v \leftarrow v + \gamma_d \nabla_v L_D$ end for Sample a batch { z_i } from μ^Z Generate $y_i \leftarrow G_{\theta}(z_i)$ Generate $\zeta_i \leftarrow y_i - \Delta t \nabla \phi_w(y_i)$ for K generator updates do $\theta \leftarrow \theta - \frac{\gamma_g}{m} \nabla_{\theta} \sum_{i=1}^{m} |\zeta_i - G_{\theta}(z_i)|^2$ end for end for

6.1 Connections to Generative Adversarial Networks (GANs)

GANs of Goodfellow et al. [6] also approach (2.5)—yet with the W_2 distance replaced by the Jensen-Shannon divergence (JSD). The key idea of [6] is to minimize the JSD loss by a min-max game between a generator and a discriminator. While the GAN algorithm had been impactful on artificial intelligence, it was well-known for its instability; see e.g., [7] and [12]. A popular solution is to replace the distance function JSD by the first-order Wasserstein distance (W_1). This yields the Wasserstein GAN (WGAN) algorithm [2], which minimizes the W_1 loss by the same min-max game idea in [6]. As shown in [2, Figure 2], WGAN performs well where early versions of GANs fare poorly.

If one further replaces the distance function W_1 by W_2 , the problem becomes exactly (2.5). Note

that (2.5) has received much less attention in the literature (relative to the W_1 counterpart), with only a few exceptions (e.g., [11] and [8]). In particular, to solve (2.5), [11] develops an algorithm (called W2-GAN) by following the same min-max game idea in [6]. While we solve (2.5) by a different approach (i.e., the gradient-flow perspective), it is interesting to note that our algorithm W2-FE covers W2-GAN as a special case. For K = 1 in W2-FE, the generator update reduces to standard SGD without persistent training, which in turn equates W2-FE with W2-GAN.

Proposition 6.1. W2-GAN (i.e., [11, Algorithm 1]) is equivalent to W2-FE (i.e., Algorithm 1 above) with K = 1, up to an adjustment of learning rates.

Proof. As W2-GAN and W2-FE update the Kantorovich potential pair (ϕ_w, ψ_v) in the same way, it suffices to show that their generator updates are equivalent under K = 1. Observe that with K = 1,

$$\nabla_{\theta} \frac{1}{m} \sum_{i=1}^{m} |\zeta_i - G_{\theta}(z_i)|^2 = -\frac{2}{m} \sum_{i=1}^{m} (\zeta_i - G_{\theta}(z_i)) \nabla_{\theta} G_{\theta}(z_i)$$

$$= \frac{2}{m} \sum_{i=1}^{m} \Delta t \nabla \phi_w(G_{\theta}(z_i)) \nabla_{\theta} G_{\theta}(z_i) = 2\Delta t \nabla_{\theta} \left(\frac{1}{m} \sum_{i=1}^{m} \phi_w(G_{\theta}(z_i))\right),$$
(6.1)

where the second equality stems from $\zeta_i = G_{\theta}(z_i) - \Delta t \nabla \phi_w(G_{\theta}(z_i))$, due to the two lines above the generator update in W2-FE. Thus, the generator update in W2-FE takes the form $\theta \leftarrow \theta - \gamma_g 2\Delta t \nabla_{\theta} \frac{1}{m} \sum_{i=1}^m \phi_w(G_{\theta}(z_i))$, the same as that in W2-GAN with learning rate $\gamma_g 2\Delta t$.

It is tempting to think that if we carry out the generator update K > 1 consecutive times in W2-GAN, W2-GAN will be equivalent to W2-FE with the same K > 1. In fact, when G_{θ} is updated for the second time in W2-FE, the second equality in (6.1) no longer holds, as " $\zeta_i = G_{\theta}(z_i) - \Delta t \nabla \phi_w(G_{\theta}(z_i))$ " is true only when G_{θ} has not been updated. That is, the generator updates in W2-FE and W2-GAN coincide only when K = 1, and can be quite different for K > 1.

7 Numerical Experiments

First, as in [12, Section 3.1], we consider learning a two-dimensional mixture of Gaussians arranged on a circle from an initially given Gaussian distribution. We carry out this task using Algorithm 1 (W2-FE) with varying K values and compare its performance with that of the refined WGAN algorithm in [13] (called W1-LP), which is arguably one of the most well-performing and stable WGAN algorithms. Figure 1 shows the qualitative evolution of the models through time.

Figure 2, on the other hand, presents the actual W_1 and W_2 losses through time. We see that W1-LP converges to the same loss level as W2-FE and achieves this faster than W2-FE with K = 1 (i.e., without persistent training). However, W2-FE with a higher K value converges much faster than W1-LP in both number of epochs and wallclock time. With K = 10, W2-FE achieves convergence within 100 training epochs, while W1-LP needs more than 600 training epochs.

Next, we consider domain adaptation from the USPS dataset to the MNIST dataset. We carry out this task using Algorithm 1 (W2-FE) with varying K values and W1-LP in [13], respectively, and evaluate their performance every 100 training epochs using a 1-nearest neighbor (1-NN) classifier.¹ Figure 3 presents the results. It is clear that W2-FE with K = 1 already outperforms W1-LP: while their performance are similar before epoch 3000, W2-FE with K = 1 consistently achieves a higher accuracy rate after epoch 3000, with a margin of 5%. When we raise the persistency level to K = 3, W2-FE converges significantly faster and consistently achieves an even higher accuracy rate. Indeed, it takes W2-FE with K = 1 8000 epochs to attain its ultimate accuracy rate, which is achieved by W2-FE with K = 3 by epoch 3000; W2-FE with K = 3 continues to improve after epoch 3000, yielding the best accuracy rate among all the models. Raising persistency level further to K = 5 worsens the training quality, which might result from overfitting.

¹This is the same performance metric as in [15], but [15] evaluate their models only once at the end of training.




Figure 1: Qualitative evolution of learning a mixture of Gaussians on a circle (in green) from an initial Gaussian at the center (in magenta). Each grey arrow indicates how a sample from the initial Gaussian is transported toward the target distribution.




Figure 2: The first (resp. second) row plots the Wasserstein-1 (resp. Wasserstein-2) loss against training epoch and wallclock time, respectively, for learning a mixture of Gaussians on a circle from an initial Gaussian at the center.




Figure 3: 1-NN classifier accuracy against training epoch for domain adaptation from the USPS dataset to the MNIST dataset.

References

- L. AMBROSIO, N. GIGLI, AND G. SAVARÉ, Gradient flows in metric spaces and in the space of probability measures, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, second ed., 2008.
- [2] M. ARJOVSKY, S. CHINTALA, AND L. BOTTOU, Wasserstein generative adversarial networks, in Proceedings of the 34th International Conference on Machine Learning, D. Precup and Y. W. Teh, eds., vol. 70 of Proceedings of Machine Learning Research, PMLR, 06–11 Aug 2017, pp. 214–223.
- [3] V. BARBU AND M. RÖCKNER, From nonlinear Fokker-Planck equations to solutions of distribution dependent SDE, Ann. Probab., 48 (2020), pp. 1902–1920.
- [4] R. CARMONA AND F. DELARUE, Probabilistic theory of mean field games with applications. I, Springer, Cham, 2018.
- [5] M. FISCHETTI, I. MANDATELLI, AND D. SALVAGNIN, Faster SGD training by minibatch persistency, CoRR, abs/1806.07353 (2018).
- [6] I. GOODFELLOW, J. POUGET-ABADIE, M. MIRZA, B. XU, D. WARDE-FARLEY, S. OZAIR, A. COURVILLE, AND Y. BENGIO, *Generative adversarial nets*, in Advances in Neural Information Processing Systems 27, 2014, pp. 2672–2680.
- [7] I. J. GOODFELLOW, NIPS 2016 tutorial: Generative adversarial networks, (2016). Available at https://arxiv.org/abs/1701.00160.
- [8] Y.-J. HUANG, S.-C. LIN, Y.-C. HUANG, K.-H. LYU, H.-H. SHEN, AND W.-Y. LIN, On characterizing optimal Wasserstein GAN solutions for non-Gaussian data, in 2023 IEEE International Symposium on Information Theory (ISIT), 2023, pp. 909–914.
- [9] Y.-J. HUANG AND Y. ZHANG, GANs as gradient flows that converge, Journal of Machine Learning Research, 24 (2023), pp. 1–40.
- [10] I. KARATZAS AND S. E. SHREVE, Brownian motion and stochastic calculus, vol. 113 of Graduate Texts in Mathematics, Springer-Verlag, New York, second ed., 1991.

- [11] J. LEYGONIE, J. SHE, A. ALMAHAIRI, S. RAJESWAR, AND A. C. COURVILLE, Adversarial computation of optimal transport maps, CoRR, abs/1906.09691 (2019).
- [12] L. METZ, B. POOLE, D. PFAU, AND J. SOHL-DICKSTEIN, Unrolled generative adversarial networks, in International Conference on Learning Representations, 2017.
- [13] H. PETZKA, A. FISCHER, AND D. LUKOVNIKOV, On the regularization of wasserstein GANs, in International Conference on Learning Representations, 2018.
- [14] F. SANTAMBROGIO, Optimal transport for applied mathematicians, vol. 87 of Progress in Nonlinear Differential Equations and their Applications, Birkhäuser/Springer, Cham, 2015. Calculus of variations, PDEs, and modeling.
- [15] V. SEGUY, B. B. DAMODARAN, R. FLAMARY, N. COURTY, A. ROLET, AND M. BLONDEL, Large scale optimal transport and mapping estimation, in International Conference on Learning Representations, 2018.
- [16] D. TREVISAN, Well-posedness of multidimensional diffusion processes with weakly differentiable coefficients, Electron. J. Probab., 21 (2016), pp. Paper No. 22, 41.
- [17] C. VILLANI, Optimal transport, vol. 338 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Springer-Verlag, Berlin, 2009. Old and new.