DIFFRACTION OF THE PRIMES AND OTHER SETS OF ZERO DENSITY

ADAM HUMENIUK, CHRISTOPHER RAMSEY, AND NICOLAE STRUNGARU

ABSTRACT. In this paper, we show that the diffraction of the primes is absolutely continuous, showing no bright spots (Bragg peaks). We introduce the notion of counting diffraction, extending the classical notion of (density) diffraction to sets of density zero. We develop the counting diffraction theory and give many examples of sets of zero density of all possible spectral types.

1. INTRODUCTION

In 2018, much interest was given to a group of papers [28, 26, 27] that studied the prime numbers through diffraction experiments. They discovered that the primes in certain large intervals possess a pattern of Bragg-like peaks that is evocative of the diffraction pattern one sees in quasicrystals, aperiodic solids with diffraction that is crystal-like. The main conjecture of these papers was that this discovered pattern showed deep structural results about the prime numbers and that these approximate diffraction pictures suggest that the whole set of primes have a pure point diffraction spectrum.

In this paper, we establish in Theorem 5.12 that the diffraction measure of the prime numbers is the Lebesgue measure, therefore, absolutely continuous. This means that any perceived Bragglike peak structure observed in a physical diffraction experiment of a finite portion of the primes is an artifact of the experiment's finiteness and disappears when taking the limit. As always, the effectiveness of an approximation comes down to what form of convergence is being used and how fast that convergence happens. We argue here that the correct setting is the convergence of measures in the vague topology, which is the foundation of mathematical diffraction theory, which was developed with its feet firmly planted in physics [8, 13, 10, 4]. It is of note that the diffraction of the finite approximations converge extremely slowly for many highly ordered structures, and are nearly impossible to pick up convincingly in a physical experiment, notably the circular symmetry of the pinwheel tiling (see for example [12, Section 4]). As we discuss in Remark 5.8, this also seems to be the case for the diffraction of the primes, which explains why the simulations in [28, 26, 27] do not show the real picture for the infinite set of primes.

The diffraction of the mathematical idealization of quasicrystals and other nice structures has been studied so far by using the structure's uniform, positive density [13, 4]. However, the prime numbers have zero density in the positive real line so a new approach is needed. Here, we define the counting diffraction, which is studied through its counting version of the autocorrelation (or twopoint correlation) measure. The great advantage to this is that the counting and usual diffraction measures correspond on positive density sets, Theorem 3.6, while on sets of zero density, the former measure gives a sensible extension of the Patterson formula to an infinite set, Theorem 3.8.

Given a finite set $F \subseteq \mathbb{R}^d$, its diffraction is given by the Patterson formula:

$$I_F(y) \coloneqq \frac{1}{\operatorname{card}(F)} \left| \sum_{x \in F} e^{2\pi i x \cdot y} \right|^2$$

A simple computation shows that I_F is the Fourier transform of the finite measure

$$\gamma_F \coloneqq \frac{1}{\operatorname{card}(F)} \delta_F * \widetilde{\delta_F} = \frac{1}{\operatorname{card}(F)} \sum_{x, y \in F} \delta_{x-y} \,.$$

The measure γ_F is called the autocorrelation, or 2-point correlation, measure of the finite sample F.

Next, consider an infinite uniformly discrete set $\Lambda \subseteq \mathbb{R}^d$ and some nice averaging sequence $A_n \subseteq \mathbb{R}^d$, such as $A_n = [-n, n]^d$. The diffraction of Λ is defined as the limit in a suitable topology of the diffraction measures I_{F_n} of the finite sets $F_n \coloneqq \Lambda \cap A_n$. For uniformly discrete sets Λ of positive density, it is more advantageous to define the autocorrelation and diffraction of Λ by dividing by the volume, $\operatorname{vol}(A_n)$, of the averaging sequence instead of the cardinality, $\operatorname{card}(F_n)$. As introduced by Hof [13] in \mathbb{R}^d , the sequence

$$\gamma_n \coloneqq \frac{1}{\operatorname{vol}(A_n)} \delta_{F_n} * \widetilde{\delta_{F_n}}$$

of autocorrelation measures of the finite samples F_n has a subsequence converging to some positive definite measure γ . The measure γ is Fourier transformable, and, by [18, Lemma 4.11.10] or [7, Theorem 4.5], its Fourier transform $\widehat{\gamma}$ is a positive measure called the (density) diffraction of Λ .

The choice of averaging by $vol(A_n)$ has the great advantage that one can often use spectral theory of dynamical systems via the so called Dworkin argument (see [10, 4, 11], just to name a few). The relationship

$$\gamma_n = \frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)} \gamma_{F_n}$$

shows that this change from γ_{F_n} to γ_n simply multiplies the diffraction measure by the positive density

$$d = \lim_{n} \frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)},$$

of the point set. In particular, for sets of positive density, the two approaches should be equivalent, and they are indeed equivalent (see Theorem 3.6). On the other hand, the same relation shows that, for sets of density zero, defining the autocorrelation as the limit of γ_n is wrong, as this would always give a zero diffraction measure.

Over the last few years, there has grown interest in the field of aperiodic order towards the study of point sets that are not relatively dense, such as for example maximal density weak model sets (see [3, 14, 15]). So far, they have been studied using certain averaging sequences with respect to which they have positive density. It is worth emphasizing that a point set Λ is not relatively dense if and only if there exists a van Hove sequence with respect to which the set has a density of zero¹. These examples, as well as the diffraction of primes we mention above, suggest that it would be good to properly define the diffraction for sets of density zero, which we do in this paper.

The paper is organized as follows: in Section 2 we collect some basic definitions and properties needed in the paper, and define the density and counting autocorrelations and diffractions. We study the existence and basic properties of the autocorrelation/diffraction measures, as well as their relationship in Section 3. In particular, we show that for uniformly discrete point sets, both the density and counting autocorrelations exist along subsequences of a given van Hove sequence (Proposition 3.3). Furthermore, for sets of positive density, the counting and density diffraction

¹In fact, this is equivalent to the seemingly stronger statement that for each van Hove sequence A_n there exists a sequence $t_n \in G$ such that Λ has zero density with respect to the van Hove sequence $t_n + A_n$.

measures are proportional (Theorem 3.6), while for sets of density zero, the density diffraction is always null while the counting diffraction is non-zero for non-trivial point sets. In Section 4 we show that for sets which are very sparse, such as the Fibonacci numbers or any fast-growing sequence, the diffraction is the Lebesgue measure. In Section 5 we prove in Theorem 5.12 that the diffraction of the primes is the Lebesgue measure. This is one of the main results of the paper, and shows that the primes have absolutely continuous diffraction spectrum. We show that this holds when the averaging sequence is any reasonable van Hove sequence of intervals. We also study the diffraction of some related sets, such as the diffraction of prime powers and the diffraction of twin primes. We complete the paper by showing in Section 6 that there exist sets of density zero with diffraction of any spectral type and showing in Section 7 that the counting diffraction is preserved when a point set is embedded into higher-dimensional space.

2. Preliminaries

We begin with recalling a number of definitions. It should be noted that everything before the definition of counting autocorrelation and counting diffraction for infinite sets is standard.

Throughout this paper, G is a second countable locally compact abelian group (LCAG), and any such group is metrisable. While in most examples G will simply be \mathbb{R} , we want to set up the theory of counting diffraction in general settings.

Definition 2.1. A set $\Lambda \subseteq G$ is called **uniformly discrete** if for any choice of metric d that generates the topology on G, there exists a constant r > 0 such that for all $x, y \in \Lambda$, if $x \neq y$, then $d(x,y) \geq r$.

The set $\Lambda \subseteq G$ is called **locally finite** if, for all compact sets $K \subseteq G$, the set $\Lambda \cap K$ is finite.

It is easy to see that $\Lambda \subset G$ is uniformly discrete if and only if there exists some non-empty open set U such that, any translate t + U of U meets Λ in at most one point. On another hand, Λ is locally finite exactly when Λ is closed and discrete in G.

Finally, a set Λ is locally finite exactly when the Dirac comb

$$\delta_\Lambda\coloneqq \sum_{x\in\Lambda}\delta_x$$

is a locally finite measure on G.

Next, let us recall the definition of FLC.

Definition 2.2. A point set Λ has finite local complexity (FLC) if $\Lambda - \Lambda$ is locally finite.

For a finite set $F \subseteq \mathbb{R}^d$, define the **autocorrelation measure** of F as

$$\gamma_F := \begin{cases} \frac{1}{\operatorname{card}(F)} \delta_F * \widetilde{\delta_F} & \text{if } F \neq \emptyset \\ 0 & \text{if } F = \emptyset \end{cases}$$

where \star denotes convolution of finite measures, and $\widetilde{\delta_F}$ is the reflected measure $\widetilde{\delta_F}(U) = \delta_F(-U)$, i.e.

$$\widetilde{\delta_F} = \sum_{x \in F} \delta_{-x}$$

Recall that the **Fourier transform** of a finite Radon measure μ on a LCAG G is the extended complex function $\hat{\mu}$ on the dual group \hat{G} defined by

$$\widehat{\mu}(\chi) = \int_G \overline{\chi(\mu)} \,\mathrm{d}\mu(\chi)$$

Definition 2.3. The counting diffraction of a finite set $F \subseteq G$ is the Fourier transform

$$\widehat{\gamma_F}(\chi) = \frac{1}{\operatorname{card}(F)} \widehat{\delta_F * \delta_F}(\chi) = \frac{1}{\operatorname{card}(F)} \left| \sum_{x \in F} \chi(x) \right|^2$$

If Λ is not finite, defining the autocorrelation γ_{Λ} , and therefore the diffraction of Λ , is more difficult as the convolution $\delta_{\Lambda} * \widetilde{\delta_{\Lambda}}$ is ill-defined. If Λ is locally finite, then we can avoid this issue by restricting to a sequence of finite sets $F_n = A_n \cap \Lambda$, where $\mathcal{A} = (A_n)$ is an appropriate averaging sequence of compact sets in G. As it was noted by Schlottmann [24], when dealing with point sets and measures, one has to work with so-called van Hove sequences. Let us recall here the definition.

Definition 2.4. A van Hove sequence $\mathcal{A} = (A_n)_n$ in a second countable LCAG G is a sequence of compact sets of positive measure such that for every compact subset $K \subseteq G$,

$$\lim_{n \to \infty} \frac{\operatorname{vol}(\partial^K(A_n))}{\operatorname{vol}(A_n)},$$

where "vol" denotes Haar measure, and

$$\partial^{K}(A) = ((A+K) \setminus A^{\circ}) \cup ((\overline{G \setminus A} - K) \cap A)$$

is the K-boundary of a set A.

The K boundary $\partial^{K}(A)$ is the set of points which are K-translates of points in A that land outside A° , or which are points in A which are translated outside of A° by a point in K. If $K = \{g\}$ is a singleton, the symmetric difference $(g + A) \bigtriangleup A$ satisfies

$$(g+A) \bigtriangleup A \subseteq \partial^{\{g,-g\}}(A),$$

Therefore if A_n is a van Hove sequence, and g is in G, then

$$\lim_{n \to \infty} \frac{\operatorname{vol}((g + A_n) \bigtriangleup A_n)}{\operatorname{vol}(A_n)} = 0$$

That is, every van Hove sequence is a Følner sequence.

Remark 2.5.

- (a) Every second countable LCAG admits van Hove sequences [24].
- (b) Some authors define van Hove sequences to be precompact. While this seems to allow more general averaging sequences, it typically makes no difference to calculations. Indeed, it is easy to see that a sequence A_n of precompact sets is a van Hove sequence if and only if A_n is a van Hove sequence. Moreover, [24, Lemma 1.1] implies that for every translation bounded measure μ (as defined below) we have

$$\lim_{n} \left| \frac{1}{\operatorname{vol}(A_n)} \mu(A_n) - \frac{1}{\operatorname{vol}(\overline{A_n})} \mu(\overline{A_n}) \right| = 0.$$

Because of this, restricting to compact van Hove sequences is not a restriction.

(c) All the examples below are examples of point sets $\Lambda \subseteq \mathbb{R}$. For those examples, we will always use van Hove sequences $A_n = [a_n, b_n]$ of intervals. It is easy to see that a sequence $A_n = [a_n, b_n]$ of intervals is a van Hove sequence if and only if $\lim_{n \to \infty} b_n - a_n = \infty$.

We will sometimes prefer to work with van Hove sequences which are **nested** in the sense that $A_n \subseteq A_{n+1}$ for every n. We will also say that (A_n) is **exhausting** if $\bigcup_n A_n = G$. The most typical van Hove sequences on \mathbb{R}^d are the collection of cubes $[-n, n]^d$ or balls $A_n = \{x \in \mathbb{R}^d \mid ||x|| \le n\}$.

When proving the existence of the autocorrelation, we will need the following measure theoretic background.

By the Riesz Representation Theorem [9, 22], a **Radon measure** (or simply measure) on G can be seen as a linear function μ on the space $C_{c}(G)$ of compactly supported continuous functions on G, which is continuous with respect to the so-called "inductive" topology on $C_{c}(G)$, obtained by writing

$$C_{\mathsf{c}}(G) = \bigcup_{\substack{K \subseteq G \\ K \text{ compact}}} C(G:K),$$

where

$$C(G:K) \coloneqq \{f \in C_{\mathsf{c}}(G) \mid \operatorname{supp}(f) \subseteq K\}.$$

The **vague topology** for measure is defined as the weak-* topology on the space $\mathcal{M}(G)$ of measures on G viewed as the dual space of $C_{\mathsf{c}}(G)$. More precisely, a net μ_{α} of measures converges vaguely to a measure μ precisely when

$$\mu(f) = \lim_{\alpha} \mu_{\alpha}(f) \quad \text{for all } f \in C_{\mathsf{c}}(G).$$

Give a Radon measure μ on G and a precompact Borel set B we can define

$$\|\mu\|_B \coloneqq \sup_{t \in G} |\mu|(t+B)|$$

where $|\mu|$ denotes the total variation measure of μ (see [19, Sect. 6.5.5] for the definition and properties).

The set of translation bounded measures is defined as

$$\mathcal{M}^{\infty}(G) = \{ \mu \in \mathcal{M}(G) \mid \|\mu\|_{K} < \infty \text{ for all compact } K \subseteq G \}.$$

In fact, if U and V are any precompact sets with nonempty interior, then $\|\cdot\|_U$ and $\|\cdot\|_V$ define norms (not just seminorms) on $\mathcal{M}^{\infty}(G)$, and these norms are equivalent [5].

Let us note here that for any uniformly discrete point set Λ we have $\delta_{\Lambda} \in \mathcal{M}^{\infty}(G)$ [4], which we use below.

For any constant C > 0 and open precompact set $U \subseteq G$, write

$$\mathcal{M}_{C,U}(G) \coloneqq \{ \mu \in \mathcal{M}^{\infty}(G) \mid \|\mu\|_U \leq C \}.$$

Since the vague topology is a weak-* topology, the Banach-Alaoglu Theorem implies compactness of $\mathcal{M}_{C,U}(G)$.

Lemma 2.6. [4, Theorem 2] Let G be a LCAG. Then, the set $\mathcal{M}_{C,U}(G)$ is vaguely compact. If G is second countable, then the vague topology is metrisable on $\mathcal{M}_{C,U}(G)$.

Let us now introduce the following two definitions, and discuss their relevance. Note again that the infinite version of the counting autocorrelation and diffraction are novel definitions and these have not appeared elsewhere to the best of our knowledge. **Definition 2.7.** Let $\Lambda \subseteq G$ be a uniformly discrete point set, and let $\mathcal{A} = (A_n)$ be a van Hove sequence. Define $F_n \coloneqq \Lambda \cap A_n$ for each n, and set

$$\gamma_{F_n} := \begin{cases} \frac{1}{\operatorname{card}(F_n)} \delta_{F_n} * \widetilde{\delta_{F_n}} & \text{if } F_n \neq \emptyset \\ 0 & \text{if } F_n = \emptyset \end{cases}$$

and

$$\gamma_n \coloneqq \frac{1}{\operatorname{vol}(A_n)} \delta_{F_n} * \tilde{\delta}_{F_n},$$

where vol denotes Haar measure.

We say that the **density autocorrelation** γ_{dens} of Λ exists with respect to \mathcal{A} if the limit

$$\gamma_{dens} = \lim_{n} \gamma_n$$

exists in the vague topology. In the same way, we say that the **counting autocorrelation** γ_{count} of Λ exists with respect to \mathcal{A} if the limit

$$\gamma_{count} = \lim_{n} \gamma_{F_n}$$

exists in the vague topology.

As limits of positive definite measures, $\gamma_{count}, \gamma_{dens}$ are positive definite measures [18, Lemma 4.11.10] and hence Fourier transformable. We refer to the positive measures $\overline{\gamma_{dens}}$ and $\overline{\gamma_{count}}$ as the **density diffraction** and the **counting diffraction** of Λ with respect to \mathcal{A} .

Let us note here the following simple relation which we will often use:

(2.1)
$$\gamma_n = \frac{\operatorname{card}(\Lambda \cap A_n)}{\operatorname{vol}(A_n)} \gamma_{F_n} \,.$$

Using a standard argument, we explain below in Proposition 3.3 that if Λ is a uniformly discrete point set in G, and $\mathcal{A} = (A_n)$ is any van Hove sequence, then there is a subsequence of \mathcal{A} with respect to which the counting and density autocorrelations of Λ both exist.

Remark 2.8. Since the supports of each measure satisfy $\operatorname{supp}(\gamma_n), \operatorname{supp}(\gamma_{F_n}) \subseteq \Lambda - \Lambda$ we have

$$\operatorname{supp}(\gamma_{dens}), \operatorname{supp}(\gamma_{count}) \subseteq \Lambda - \Lambda.$$

In the next section, we will show that the density and counting diffractions are related via a relation of the type

$$\widehat{\gamma_{\text{dens}}} = \text{dens}(\Lambda)\widehat{\gamma_{\text{count}}}$$

In particular, in Theorem 3.6 and Theorem 3.8 we will show the following:

- For point sets of positive density, the counting and density diffraction coincide up to multiplication by non-zero constants. In this case, the two approaches to the diffraction spectrum are equivalent, and either one can be used.
- For point sets of zero density, the density diffraction is zero and the counting diffraction is non-zero. In this case, the counting diffraction must be used.

3. Density and autocorrelation

In this section, we study the basic properties of the density and counting autocorrelations.

3.1. Density and counting autocorrelations.

Definition 3.1. Let $\Lambda \subseteq G$ be uniformly discrete and $\mathcal{A} = (A_n)$ a van Hove sequence. We define the lower and upper density of Λ with respect to \mathcal{A} by

$$\underline{\operatorname{dens}}_{\mathcal{A}}(\Lambda) \coloneqq \liminf_{n} \frac{\operatorname{card}(\Lambda \cap A_{n})}{\operatorname{vol}(A_{n})} \quad and$$
$$\overline{\operatorname{dens}}_{\mathcal{A}}(\Lambda) \coloneqq \limsup_{n} \frac{\operatorname{card}(\Lambda \cap A_{n})}{\operatorname{vol}(A_{n})},$$

respectively. We say that Λ has **positive density** with respect to \mathcal{A} if

 $\underline{\operatorname{dens}}_{\mathcal{A}}(\Lambda) > 0.$

We say that the density of Λ exists with respect to \mathcal{A} if the limit

dens_{$$\mathcal{A}$$} $(\Lambda) \coloneqq \lim_{n} \frac{\operatorname{card}(\Lambda \cap A_n)}{\operatorname{vol}(A_n)}$ exists.

By definition, the density of Λ exists with respect to \mathcal{A} if and only if $\underline{\mathrm{dens}}_{\mathcal{A}}(\Lambda) = \overline{\mathrm{dens}}_{\mathcal{A}}(\Lambda)$, and if this is the case, then we have

$$\underline{\operatorname{dens}}_{\mathcal{A}}(\Lambda) = \overline{\operatorname{dens}}_{\mathcal{A}}(\Lambda) = \operatorname{dens}_{\mathcal{A}}(\Lambda).$$

The following result follows immediately from [24, Lemma 1.1(2)].

Lemma 3.2. If Λ is uniformly discrete, then the sequence $\frac{\operatorname{card}(\Lambda \cap A_n)}{\operatorname{vol}(A_n)}$ is bounded. In particular, $\overline{\operatorname{dens}}_{\mathcal{A}}(\Lambda) < \infty$.

Proof. Since Λ is uniformly discrete, the measure δ_{Λ} is translation bounded. Therefore, by [24, Lemma 1.1(2)], the sequence

$$\frac{\operatorname{card}(\Lambda \cap A_n)}{\operatorname{vol}(A_n)} = \frac{\delta_{\Lambda}(A_n)}{\operatorname{vol}(A_n)} = \frac{|\delta_{\Lambda}|(A_n)}{\operatorname{vol}(A_n)}$$

is bounded.

As promised in Section 2, we can now prove that the counting and density diffractions exist for a uniformly discrete set and any van Hove sequence, perhaps upon passing to a subsequence.

The following lemma is standard, but we include the proof for completeness.

Proposition 3.3. Let $\Lambda \subseteq G$ be any set such that $\delta_{\Lambda} \in \mathcal{M}^{\infty}(G)$ and let $\mathcal{A} = \{A_n\}$ be a van Hove sequence. Then, for each open precompact set U, there exists some C > 0 such that

$$\gamma_n, \gamma_{F_n} \in \mathcal{M}_{C,U}(G)$$
 for all n .

In particular, there exists a subsequence \mathcal{B} of \mathcal{A} such that $\gamma_{dens}, \gamma_{count}, \operatorname{dens}_{\mathcal{B}}(\Lambda)$ exist and

$$\gamma_{dens} = \operatorname{dens}_{\mathcal{B}}(\Lambda) \gamma_{count}.$$

In particular, the conclusion holds whenever Λ is uniformly discrete.

Proof. Pick any open pre-compact set V such that $\overline{U} \subseteq V$. Since $\widetilde{\delta_{F_n}}/\text{card}(F_n)$ is a finite measure of total mass 1, by [25, Lemma 6.1] we have

$$\|\gamma_{F_n}\|_U \le \|\delta_{F_n}\|_V \left(\frac{\widetilde{\delta_{F_n}}}{\operatorname{card}(F_n)}(G)\right) \le \|\delta_\Lambda\|_V =: C_1$$

where $C_1 < \infty$ since $\delta_{\Lambda} \in \mathcal{M}^{\infty}(G)$.

Next, since $\frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)} = \frac{\delta_{\Lambda}(A_n)}{\operatorname{vol}(A_n)}$ is bounded above by some constant *D*, from Lemma 3.2, by (2.1) we have

$$\|\gamma_n\|_U = \left\|\frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)}\gamma_{F_n}\right\|_U \le DC_1.$$

The claim follows by picking $C = \max\{C_1, DC_1\}$. The existence of the limit along subsequences follows now from 2.6 and the boundedness of $\frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)}$. Finally, by definition, Λ is uniformly discrete if and only if there exists a precompact open set W

Finally, by definition, Λ is uniformly discrete if and only if there exists a precompact open set W such that

 $\|\delta_{\Lambda}\|_W \leq 1$.

The claim follows immediately.

Let us now look at an example which shows why translation boundedness is an important assumption, even for sets of density zero. We construct a (non-uniformly discrete) set $\Lambda \subseteq \mathbb{R}$ with zero density such that $\delta_{\Lambda} \notin \mathcal{M}^{\infty}(\mathbb{R})$ and γ_{F_n} has no vaguely convergent subsequence.

Example 3.4. Define

$$\Lambda_m \coloneqq \left\{ 4^m - \frac{j}{2^{m+1}} : 1 \le j \le 2^m \right\}, \qquad \Lambda \coloneqq \bigcup_{m=1}^\infty \Lambda_m.$$

Let us note here in passing that we start at $4^m - \frac{1}{2^{m+1}}$ and not 4^m to avoid having integers in Λ_m , as, in that case, the F_n we introduce below would have an extra point whenever $n = 4^m$. This extra point would not change any conclusion, but would make the proof slightly more technical.

By construction we have $\operatorname{card}(\Lambda_m) = 2^m$ and $\Lambda_m \subset [4^m - \frac{1}{2}, 4^m)$. This immediately implies that

$$F_n = \Lambda \cap \big[-n, n \big] = \bigcup_{j=1}^m \Lambda_j$$

where $m = \lfloor \log_4 n \rfloor$ is the last positive integer satisfying $4^m \le n$. Note here we are using $A_n = [-n, n]$ as our usual averaging sequence. In particular,

$$\operatorname{card}(F_n) = 2 + \dots + 2^m = 2^{m+1} - 2 = 2(2^{\lfloor \log_4 n \rfloor} - 1).$$

This implies that

$$\operatorname{dens}(\Lambda) = \lim_{n} \frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)} = \lim_{n} \frac{2(2^{\lfloor \log_4 n \rfloor} - 1)}{2n} \le \lim_{n} \frac{2^{\log_4 n} - 1}{n} = \lim_{n} \frac{\sqrt{n} - 1}{n} = 0$$

as claimed.

Now, fix some non-negative function $f \in C_{\mathsf{c}}(\mathbb{R})$ so that $f(x) \ge 1$ for all $x \in [-\frac{1}{2}, \frac{1}{2}]$. Let $n \in \mathbb{N}$ be arbitrary, and set

$$m = \lfloor \log_4 n \rfloor$$
.

Then,

$$\gamma_{F_n}(f) = \frac{1}{2(2^{\lfloor \log_4 n \rfloor} - 1)} \sum_{x, y \in F_n} f(x - y) \ge \frac{1}{2(2^{\log_4 n})} \sum_{x, y \in \Lambda_m} f(x - y).$$

Recall that $\Lambda_m \subseteq [4^m - \frac{1}{2}, 4^m)$. Therefore, for all $x, y \in \Lambda_m$ we have $x - y \in [-\frac{1}{2}, \frac{1}{2}]$ and hence $f(x - y) \ge 1$. It follows that

$$\gamma_{F_n}(f) \ge \frac{1}{2\sqrt{n}} \sum_{x,y \in \Lambda_m} 1 = \frac{1}{2\sqrt{n}} \left(\operatorname{card}(\Lambda_m) \right)^2$$
$$= \frac{1}{2\sqrt{n}} 2^{2m} = \frac{1}{2\sqrt{n}} 4^{\lfloor \log_4 n \rfloor} \ge \frac{1}{2\sqrt{n}} 4^{\log_4 n - 1} = \frac{1}{8} \sqrt{n}$$

Therefore,

$$\gamma_{F_n}(f) \ge \frac{1}{8}\sqrt{n}$$
 for all n

which implies that γ_{F_n} has no vaguely convergent subsequence.

Let us next prove the following result.

Lemma 3.5. Let $\Lambda \subseteq G$ be uniformly discrete.

(a) If γ_{dens} exists with respect to $\mathcal{A} = (A_n)_n$ then dens_{\mathcal{A}}(Λ) exists and

$$\gamma_{dens}(\{0\}) = \operatorname{dens}_{\mathcal{A}}(\Lambda)$$

- (b) If γ_{count} exists with respect to \mathcal{A} , then the following are equivalent:
 - (i) $\gamma_{count}(\{0\}) = 1$.
 - (ii) $F_n = \Lambda \cap A_n$ contains a subsequence of non-empty sets.
 - (iii) There exists some N such that for all n > N we have $F_n \neq \emptyset$.
 - (iv) $\gamma_{count} \neq 0$.

Proof. First, let us note that since Λ is uniformly discrete, 0 is an isolated point of $\overline{\Lambda - \Lambda}$. Therefore, there exists some open set $U \subseteq G$ containing 0 such that

$$\left(\overline{\Lambda-\Lambda}\right)\cap U=\left\{0\right\}.$$

For the entire proof, we fix some $f \in C_{\mathsf{c}}(G)$ such that $f(0) = 1, 0 \le f \le 1$, and $\operatorname{supp}(f) \subseteq U$.

(a) Then, Remark 2.8 implies that

$$\gamma(f) = f(0)\gamma(\{0\}) = \gamma(\{0\})$$
$$\gamma_n(f) = f(0)\gamma_n(\{0\}) = \frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)}.$$

Therefore,

$$\gamma(\{0\}) = \gamma(f) = \lim_{n} \gamma_n(f),$$

and so (a) follows.

(b) Using Remark 2.8 again, we have

$$\gamma_{F_n}(f) = \gamma_{F_n}(0) = \begin{cases} 1 & \text{if } F_n \neq \emptyset \\ 0 & \text{if } F_n = \emptyset. \end{cases}$$

$$\gamma_{\text{count}}(f) = \gamma_{\text{count}}(\{0\}).$$

Since γ_{F_n} converges vaguely to γ_{count} , we get

$$\gamma_{\text{count}}(\{0\}) = \gamma_{\text{count}}(f) = \lim_{n} \gamma_{F_n}(f) = \lim_{n} \begin{cases} 1 & \text{if } F_n \neq \emptyset, \\ 0 & \text{if } F_n = \emptyset. \end{cases}$$

Now, the implications (iii) \iff (ii) \iff (i) \implies (iv) are immediate. We will complete the proof by showing that (iv) \implies (iii). If we assume by contradiction that (iii) fails, then we can find a subsequence $F_{k_n} = \emptyset$. But then

$$\gamma_{\text{count}} = \lim_{n} \gamma_{F_n} = \lim_{n} \gamma_{F_{k_n}} = \lim_{n} \gamma_{\varnothing} = \lim_{n} 0 = 0,$$

a contradiction.

We can now prove that for sets of positive density, there is no difference between working with density or counting autocorrelation. The same is not true in the case of zero-density sets.

Theorem 3.6. Let $\Lambda \subseteq G$ be uniformly discrete and \mathcal{A} be a van Hove sequence. Suppose Λ has a positive density with respect to \mathcal{A} .

(a) If γ_{count} is any cluster point of γ_{F_n} then there exists some $C \in (0, \infty)$ and a cluster point γ_{dens} of γ_n such that

$$\gamma_{dens} = C \gamma_{count}$$
 .

(b) If γ_{dens} is any cluster point of γ_n then there exists some $D \in (0, \infty)$ and a cluster point γ_{count} of γ_{F_n} such that

 $\gamma_{count} = D\gamma_{dens}$.

Proof. (a) Let \mathcal{A}' be the subsequence of \mathcal{A} along which γ_{count} is the limit. By Lemma 3.3 there exists a subsequence \mathcal{B} of \mathcal{A}' along which the density autocorrelation and the density of Λ exist and

$$\gamma_{\rm dens} = C \gamma_{\rm count}$$

for $C = \text{dens}_{\mathcal{B}}(\Lambda)$. Positive density of Λ ensures that C > 0.

(b) The proof of (b) is a symmetrical argument, in which $D = 1/\text{dens}_{\mathcal{B}}(\Lambda)$.

Remark 3.7. The relationship between the cluster point(s) γ_{dens} and the cluster point(s) γ_{count} is explained by (2.1):

$$\gamma_n = \frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)} \gamma_{F_n} \,.$$

This relationship implies that, as long as $\frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)}$ is bounded away from 0, if two of γ_n , $\frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)}$, γ_{F_n} exist along \mathcal{A} then so does the third. Furthermore, by Lemma 3.5, if γ_n is convergent, then so is $\frac{\operatorname{card}(F_n)}{\operatorname{vol}(A_n)}$.

This means that for point sets of positive density we have the following implications:

- (a) If γ_{dens} exists along \mathcal{A} then, dens_{\mathcal{A}}(Λ) and γ_{count} exist along \mathcal{A} .
- (b) Assume that γ_{count} exists along \mathcal{A} . Then, γ_{dens} exists along \mathcal{A} if and only if dens_{\mathcal{A}}(Λ) exists along \mathcal{A} .

In contrast, in the case of zero density, the following result shows that γ_{dens} and γ_{count} are not proportional.

Theorem 3.8. Let $\Lambda \subseteq G$ be uniformly discrete and $\mathcal{A} = (A_n)$ be a van Hove sequence for which $F_n = \Lambda \cap A_n$ are not eventually all empty. Suppose dens_{\mathcal{A}}(Λ) = 0.

(a) The autocorrelation γ_{dens} of Λ exists with respect to \mathcal{A} and

$$\gamma_{dens} = 0$$

(b) If γ_{count} is any cluster point of γ_{F_n} then

 $\gamma_{count} \neq 0$.

Proof. (a) By Lemma 3.3 we have

$$\gamma_n \in \mathcal{M}_{C,U}(G)$$

for some C > 0 and precompact open U. Since $\mathcal{M}_{C,U}(G)$ is vaguely compact and metrisable, showing that $\gamma_n \to 0$ is equivalent to showing that if η is a cluster point of γ_n then $\eta = 0$.

Let η be a cluster point of γ_n calculated along some subsequence \mathcal{A}' of \mathcal{A} . By Lemma 3.3, there exists some subsequence \mathcal{B} of \mathcal{A}' such that

$$\eta = \mathrm{dens}_\mathcal{B}(\Lambda) \gamma_\mathrm{count}$$
 .

The zero density property gives dens_{\mathcal{B}}(Λ) = 0 and hence η = 0. This proves (a).

(b) By Lemma 3.5 and (a),

$$\gamma_{\text{count}}(\{0\}) = 1 \neq 0 = \gamma_{\text{dens}}(\{0\}).$$

3.2. Point sets with finite local complexity. If Λ has finite local complexity (FLC), then the following lemma shows how to write the autocorrelation as a Dirac comb.

Lemma 3.9. Let $\Lambda \subseteq G$ have FLC and let \mathcal{A} be a van Hove sequence. Then γ_{dens} exists if and only if for all $t \in \Lambda - \Lambda$ the limit

$$\eta_{dens}(t) \coloneqq \lim_{n} \frac{1}{\operatorname{vol}(A_n)} \operatorname{card}(F_n \cap (F_n - t)) = \operatorname{dens}_{\mathcal{A}}(\Lambda \cap (\Lambda - t))$$

exists. Moreover, in this case we have

$$\gamma_{dens} = \sum_{t \in \Lambda - \Lambda} \eta_{dens}(t) \delta_t \,.$$

Proof. Let $f \in C_c(G)$, and then for any finite n, rearranging the sum shows that

$$\begin{split} \gamma_n(f) &= \frac{1}{\operatorname{vol}(A_n)} \sum_{x,y \in F_n} f(y - x) \\ &= \frac{1}{\operatorname{vol}(A_n)} \sum_{t \in \Lambda - \Lambda} \sum_{\substack{x \in F_n \\ x + t \in F_n}} f(t) \\ &= \sum_{t \in \Lambda} \left(\frac{\operatorname{card}(F_n \cap (F_n - t))}{\operatorname{vol}(A_n)} \right) f(t) \\ &= \left(\sum_{t \in \Lambda} \left(\frac{\operatorname{card}(F_n \cap (F_n - t))}{\operatorname{vol}(A_n)} \right) \delta_t \right) (f), \end{split}$$

where this sum contains at most finitely many nonzero terms, since f has compact support and $\Lambda - \Lambda$ is locally finite.

If each limit

$$\eta_{\text{dens}}(t) = \lim_{n} \frac{\operatorname{card}(F_n \cap (F_n - t))}{\operatorname{vol}(A_n)}$$

exists, then this shows that γ_n vaguely converges to

$$\gamma_{\text{dens}} = \sum_{t \in \Lambda - \Lambda} \eta_{\text{dens}}(t) \delta_t$$

Conversely, suppose the density diffraction $\gamma_{\text{dens}} = \lim_n \gamma_n$ exists. Let $t \in \Lambda - \Lambda$. Since $\Lambda - \Lambda$ is locally finite, we can find a function $f \in C_c(G)$ for which f(t) = 1 and f(s) = 0 for all other $s \in \Lambda - \Lambda$. In this case, we have

$$\frac{\operatorname{card}(F_n \cap (F_n - t))}{\operatorname{vol}(A_n)} = \gamma_n(f).$$

which converges to $\gamma_{\text{dens}}(f)$, showing that the limit $\eta_{\text{dens}}(t)$ exists.

Finally, it remains to show that $\eta_{dens}(t) = dens_{\mathcal{A}}(\Lambda \cap (\Lambda - t))$, i.e.

(3.1)
$$\lim_{n} \frac{\operatorname{card}(F_n \cap (F_n - t))}{\operatorname{vol}(A_n)} = \lim_{n} \frac{\operatorname{card}(\Lambda \cap (\Lambda - t) \cap A_n)}{\operatorname{vol}(A_n)},$$

when either limit exists. Here,

$$F_n \cap (F_n - t) \subseteq \Lambda \cap (\Lambda - t) \cap A_n,$$

with set difference

$$(\Lambda \cap (\Lambda - t) \cap A_n) \setminus (F_n \cap (F_n - t)) \subseteq \Lambda \cap (A_n \setminus (A_n - t)).$$

Therefore, we can bound

$$\left|\frac{\operatorname{card}(\Lambda \cap (\Lambda - t) \cap A_n)}{\operatorname{vol}(A_n)} - \frac{\operatorname{card}(F_n \cap (F_n - t))}{\operatorname{vol}(A_n)}\right| \le \frac{\operatorname{card}(\Lambda \cap (A_n \smallsetminus (A_n - t)))}{\operatorname{vol}(A_n)}$$
$$= \frac{\delta_\Lambda(A_n \smallsetminus (A_n - t))}{\operatorname{vol}(A_n)} \le \frac{\delta_\Lambda(\partial^{\{t.-t\}}(A_n))}{\operatorname{vol}(A_n)} \to 0$$

by [24, Lemma 1.1]. The claim follows.

The next result is proven in exactly the same way, and so we will skip the proof.

Lemma 3.10. Let $\Lambda \subseteq G$ have FLC and let \mathcal{A} be a van Hove sequence such that $F_n = A_n \cap \Lambda$ is eventually always nonempty. Then γ_{count} exists if and only if for all $t \in \Lambda - \Lambda$ the limit

$$\eta_{count}(t) \coloneqq \lim_{n} \frac{\operatorname{card} \left(F_n \cap \left(-t + F_n \right) \right)}{\operatorname{card} \left(F_n \right)}$$

exists. Moreover, in this case we have

$$\gamma_{count} = \sum_{t \in \Lambda - \Lambda} \eta_{count}(t) \delta_t \, .$$

Comparing to Lemma 3.9, we may not be able to say that the limit $\eta_{\text{count}}(t)$ is the same as the limit $\ldots \operatorname{card}(\Lambda \cap (-t + \Lambda) \cap A_n)$

$$\lim_{n} \frac{\operatorname{card}(\Lambda \cap (-t + \Lambda) \cap A_n)}{\operatorname{card}(F_n)}$$

for general Λ and \mathcal{A} . But for van Hove sequences of intervals in \mathbb{R} , and non-trivial uniformly discrete sets $\Lambda \subseteq \mathbb{R}$, we can do so.

Lemma 3.11. Let Λ be a uniformly discrete point set in \mathbb{R} , and let $\mathcal{A} = (A_n)$ be a van Hove sequence of intervals for which $\operatorname{card}(F_n) = \operatorname{card}(\Lambda \cap A_n) \to \infty$ as $n \to \infty$.

Then for all $t \in \Lambda - \Lambda$ we have

$$\lim_{n} \frac{\operatorname{card}(\Lambda \cap (-t + \Lambda) \cap A_n) - \operatorname{card}(F_n \cap (-t + F_n))}{\operatorname{card}(F_n)} = 0$$

In particular, if Λ has FLC, then η_{count} exists if and only if for all $t \in \mathbb{R}$ the following limit exists

$$\eta_{count}(t) \coloneqq \lim_{n} \frac{\operatorname{card} \left(F_n \cap \left(-t + \Lambda \right) \right)}{\operatorname{card} \left(F_n \right)}$$

Moreover, in this case,

$$\gamma_{count} = \sum_{t \in \Lambda - \Lambda} \eta_{count}(t) \delta_t$$

Proof. Let $t \in \mathbb{R}$. Then the sets

$$-t + \Lambda \cap A_n$$
 and $(-t + \Lambda) \cap A_n$

have symmetric difference contained in

$$(-t+A_n) \bigtriangleup A_n$$

Because A_n is an interval, this symmetric difference is no larger than two intervals of width |t|. Since Λ is uniformly discrete, there is some minimum distance r > 0 between points of Λ , and so $-t + \Lambda \cap A_n$ and $(-t + \Lambda) \cap A_n$ can differ by at most 2|t|/r points. Intersecting with $F_n = \Lambda \cap A_n$ shows that

$$F_n \cap (-t + F_n)$$
 and $F_n \cap (-t + F)_n = \Lambda \cap (-t + \Lambda) \cap A_n$

differ still by at most 2|t|/r elements.

By assumption, $\operatorname{card}(F_n) \to \infty$, and so $\lim_n \frac{2|t|/r}{\operatorname{card}(F_n)} = 0$, and therefore

$$\lim_{n} \frac{\operatorname{card}(F_n \cap (-t + F_n)) - \operatorname{card}(\Lambda \cap (-t + \Lambda) \cap A_n)}{\operatorname{card}(F_n)} = 0.$$

The remaining claims follow from Lemma 3.10.

Remark 3.12. In \mathbb{R}^d the same proof fails, since the number of points in $\Lambda \cap ((-t + A_n) \bigtriangleup A_n)$ can be unbounded as $n \to \infty$. To draw a similar conclusion for uniformly discrete pointsets in \mathbb{R}^d using $A_n = [-n, n]^d$ we would need the stronger assumption

$$\lim_{n} \frac{\operatorname{card}(F_n)}{n^{d-1}} = \infty$$

Let us now cover a simple consequence of this, which we will use later in the paper.

Proposition 3.13. Let $\mathcal{A} = (A_n)$ be a van Hove sequence consisting of intervals in \mathbb{R} . If $\Lambda \subseteq \mathbb{R}$ is a uniformly discrete point set with counting autocorrelation δ_0 with respect to \mathcal{A} , and for which $\lim_n \operatorname{card}(\Lambda \cap A_n) = \infty$, then for any nonzero integer k, the set $\Lambda \cup (k+\Lambda)$ has counting autocorrelation and diffraction

$$\gamma_{count} = \delta_0 + \frac{\delta_k + \delta_{-k}}{2}$$
$$\widehat{\gamma_{count}} = (1 + \cos(2\pi kx))\lambda$$

with respect to \mathcal{A} , where λ is the Lebesgue measure.

Proof. Write $\Gamma = \Lambda \cup (k + \Lambda)$. By Lemma 3.11, we have that $\gamma_{\text{count}} = \sum_t \eta_{\text{count}}(t)\delta_t$ where for $t \in \mathbb{Z}$,

$$\eta_{\text{count}}(t) = \lim_{n \to \infty} \frac{\operatorname{card}(\Gamma \cap (-t + \Gamma) \cap A_n)}{\operatorname{card}(\Gamma \cap A_n)}$$

Let $\Lambda_n = \Lambda \cap A_n$ and similarly $\Gamma_n = \Gamma \cap A_n$, et cetera.

The assumption that Λ has counting autocorrelation δ_0 means that whenever $s \in \mathbb{R}$ is nonzero, card $(\Lambda \cap (-s + \Lambda) \cap A_n)$ is $o(\operatorname{card}(\Lambda_n))$ as $n \to \infty$. As in the proof of Lemma 3.11, because Λ is uniformly discrete and card $(\Lambda_n) \to \infty$, we have that card $((-s + \Lambda)_n) \sim \operatorname{card}(-s + \Lambda_n)$ as $n \to \infty$, and similarly for Γ . Using inclusion-exclusion,

$$\operatorname{card}(\Gamma_n) = \operatorname{card}(\Lambda_n) + \operatorname{card}((k+\Lambda)_n) - \operatorname{card}(\Lambda \cap (k+\Lambda) \cap A_n)$$
$$\sim \operatorname{card}(\Lambda_n) + \operatorname{card}(k+\Lambda_n) - \operatorname{card}(\Lambda \cap (k+\Lambda) \cap A_n)$$
$$= 2\operatorname{card}(\Lambda_n) - \operatorname{card}(\Lambda \cap (k+\Lambda) \cap A_n),$$

and the last term is $o(\operatorname{card}(\Lambda_n))$, so $\operatorname{card}(\Gamma_n) \sim 2\operatorname{card}(\Lambda_n)$ as $n \to \infty$.

Now, let $t \in \mathbb{R}$. The autocorrelation of Γ considers the intersection

$$\Gamma \cap (-t + \Gamma) = (\Lambda \cap (-t + \Lambda)) \cup (\Lambda \cap (k - t + \Lambda))$$
$$\cup ((k + \Lambda) \cap (-t + \Lambda)) \cup ((k + \Lambda) \cap (k - t + \Lambda)).$$

Because $k \neq 0$, the autocorrelation of Λ being δ_0 implies that the intersection of any two or more sets in this union with A_n has cardinality that is $o(\operatorname{card}(\Lambda_n))$ as $n \to \infty$. So, after using the inclusion-exclusion formula and discarding terms that go to zero, we find that

$$\eta_{\text{count}}(t) = \lim_{n} \frac{\operatorname{card}(\Gamma \cap (-t+\Gamma) \cap A_{n})}{\operatorname{card}(\Gamma_{n})} = \lim_{n} \frac{\operatorname{card}(\Gamma \cap (-t+\Gamma) \cap A_{n})}{2\operatorname{card}(\Lambda_{n})}$$
$$= \frac{1}{2} \lim_{n} \left(\frac{\operatorname{card}((\Lambda \cap (-t+\Lambda))_{n})}{\operatorname{card}(\Lambda_{n})} + \frac{\operatorname{card}((\Lambda \cap (k-t+\Lambda))_{n})}{\operatorname{card}(\Lambda_{n})} + \frac{\operatorname{card}(((k+\Lambda) \cap (k-t+\Lambda))_{n})}{\operatorname{card}(\Lambda_{n})} \right)$$
$$= \frac{1}{2} \left(\delta_{0}(t) + \delta_{0}(t-k) + \delta_{0}(t+k) + \delta_{0}(t) \right)$$
$$= \left(\delta_{0} + \frac{\delta_{k} + \delta_{-k}}{2} \right)(t).$$

Therefore Lemma 3.11 implies that $\gamma_{\text{count}} = \delta_0 + (\delta_k + \delta_{-k})/2$ for Γ .

We complete this subsection by showing that the addition of very few points does not change the counting autocorrelation.

It is well known that if two sets differ by a set of density zero, then the density autocorrelation of one exists if and only if the density autocorrelation of the other one exists. Moreover, in this case, the two autocorrelations are equal. We prove the corresponding result for counting autocorrelations.

Lemma 3.14. Let $\Lambda \subseteq \Gamma \subseteq G$ be uniformly discrete point sets and A_n be a van Hove sequence with the following properties:

- (a) $\Lambda \cap A_n$ are eventually non-empty.
- (b) The autocorrelation $\gamma_{count,\Lambda}$ exists with respect to A_n .
- (c)

$$\lim_{n} \frac{\operatorname{card}(\Lambda \cap A_n)}{\operatorname{card}(\Gamma \cap A_n)} = 1.$$

Then, the autocorrelation $\gamma_{count,\Gamma}$ exists with respect to A_n and

$$\gamma_{count,\Gamma}$$
 = $\gamma_{count,\Lambda}$.

Proof. Let $F_n \coloneqq \Lambda \cap A_n$ and $E_n \coloneqq \Gamma \cap A_n$. We will show that

$$\frac{1}{\operatorname{card}(E_n)}\delta_{E_n} * \widetilde{\delta_{E_n}} - \frac{1}{\operatorname{card}(F_n)}\delta_{F_n} * \widetilde{\delta_{F_n}} \to 0$$

vaguely, from which the result follows.

First let us note that

$$\frac{1}{\operatorname{card}(E_n)}\delta_{F_n} * \widetilde{\delta_{F_n}} - \frac{1}{\operatorname{card}(F_n)}\delta_{F_n} * \widetilde{\delta_{F_n}} = \left(\frac{\operatorname{card}(F_n) - \operatorname{card}(E_n)}{\operatorname{card}(E_n)}\right) \frac{1}{\operatorname{card}(F_n)} \left(\delta_{F_n} * \widetilde{\delta_{F_n}}\right)$$

converges to $0\cdot\gamma_{{\rm count},\Lambda}=0.$ Therefore, we need to show that

$$\frac{1}{\operatorname{card}(E_n)}\delta_{E_n} * \widetilde{\delta_{E_n}} - \frac{1}{\operatorname{card}(E_n)}\delta_{F_n} * \widetilde{\delta_{F_n}} \to 0.$$

Let $f \in C_{\mathsf{c}}(G)$ be arbitrary. Then

$$\begin{aligned} \left| \frac{1}{\operatorname{card}(E_n)} \delta_{E_n} * \widetilde{\delta_{E_n}}(f) - \frac{1}{\operatorname{card}(E_n)} \delta_{F_n} * \widetilde{\delta_{F_n}}(f) \right| \\ \leq \left| \frac{1}{\operatorname{card}(E_n)} \delta_{E_n} * \widetilde{\delta_{E_n}}(f) - \frac{1}{\operatorname{card}(E_n)} \delta_{F_n} * \widetilde{\delta_{E_n}}(f) \right| \\ + \left| \frac{1}{\operatorname{card}(E_n)} \delta_{F_n} * \widetilde{\delta_{E_n}}(f) - \frac{1}{\operatorname{card}(E_n)} \delta_{F_n} * \widetilde{\delta_{F_n}}(f) \right| \\ \leq \frac{1}{\operatorname{card}(E_n)} \left| (\delta_{E_n} - \delta_{F_n}) * \widetilde{\delta_{E_n}}(f) \right| + \frac{1}{\operatorname{card}(E_n)} \left| \delta_{F_n} * \left(\widetilde{\delta_{E_n}} - \delta_{F_n} \right)(f) \right| \\ = \frac{1}{\operatorname{card}(E_n)} \left| \int_G \int_G f(x+y) \mathrm{d} \widetilde{\delta_{E_n}}(x) \mathrm{d} \left(\delta_{E_n} - \delta_{F_n} \right)(y) \right| \\ + \frac{1}{\operatorname{card}(E_n)} \left| \int_G \int_G f(x+y) \mathrm{d} \widetilde{\delta_{E_n}}(x) \mathrm{d} \left(\delta_{E_n} - \delta_{F_n} \right)(y) \right| \\ \leq \frac{1}{\operatorname{card}(E_n)} \int_G \int_G |f(x+y)| \mathrm{d} \widetilde{\delta_{E_n}}(x) \mathrm{d} \left(\delta_{E_n \setminus F_n} \right)(y) \\ + \frac{1}{\operatorname{card}(E_n)} \int_G \int_G |f(x+y)| \mathrm{d} \widetilde{\delta_{F_n}}(x) \mathrm{d} \left(\widetilde{\delta_{E_n \setminus F_n}} \right)(y) \\ \leq \frac{1}{\operatorname{card}(E_n)} \int_G \int_G |f(x+y)| \mathrm{d} \widetilde{\delta_{F_n}}(x) \mathrm{d} \left(\delta_{E_n \setminus F_n} \right)(y) \\ + \frac{1}{\operatorname{card}(E_n)} \int_G \int_G |f(x+y)| \mathrm{d} \delta_{F_n}(x) \mathrm{d} \left(\delta_{E_n \setminus F_n} \right)(y) \\ \end{cases}$$

Now, since $f \in C_{\mathsf{c}}(G)$ and Γ is uniformly discrete, all measures involved are translation bounded and so there exists a constant C such that for all $y \in G$ we have

$$\begin{split} &\int_{G} |f(x+y)| \, \mathrm{d}\delta_{\Lambda}(x) \leq C \quad \text{and} \\ &\int_{G} |f(x+y)| \, \mathrm{d}\widetilde{\delta_{\Gamma}}(x) \leq C. \end{split}$$

It follows that

$$\left| \frac{1}{\operatorname{card}(E_n)} \delta_{E_n} * \widetilde{\delta_{E_n}}(f) - \frac{1}{\operatorname{card}(E_n)} \delta_{F_n} * \widetilde{\delta_{F_n}}(f) \right|$$

$$\leq \frac{1}{\operatorname{card}(E_n)} \int_G Cd\left(\delta_{E_n \setminus F_n}\right)(y) + \frac{1}{\operatorname{card}(E_n)} \int_G Cd\left(\widetilde{\delta_{E_n \setminus F_n}}\right)(y)$$

$$= \frac{C}{\operatorname{card}(E_n)} 2\operatorname{card}(E_n \setminus F_n) \to 0,$$

completing the proof.

3.3. Diffraction of subsets of integers. As mentioned before, many examples we will consider below are subsets Λ of the integers, considered as point sets in $G = \mathbb{R}$. The averaging van Hove sequences A_n will be of the form $A_n = [a_n, b_n]$, where $\lim_n b_n - a_n = \infty$.

Let us now fix $\Lambda \subseteq \mathbb{Z}$ and A_n a van Hove sequence in \mathbb{R} . For each n, let $f_n, g_n : \mathbb{Z} \to \mathbb{C}$ be defined as

$$f_n(d) \coloneqq \frac{1}{\operatorname{vol}(A_n)} \operatorname{card}(\Lambda \cap (d + \Lambda) \cap A_n)$$
$$g_n(d) \coloneqq \frac{\operatorname{card}(\Lambda \cap (d + \Lambda) \cap A_n)}{\operatorname{card}(\Lambda \cap A_n)} = \frac{f_n(d)}{f_n(0)}.$$

We then have:

Proposition 3.15. Let $\Lambda \subseteq \mathbb{Z}$ and let A_n be a van Hove sequence in \mathbb{R} . Then, γ_{dens} exists with respect to A_n if and only if f_n converges pointwise to a function $f : \mathbb{Z} \to \mathbb{C}$.

Moreover, in this case $\gamma_{dens} = \sum_{m \in \mathbb{Z}} f(m) \delta_m$ and f is positive definite. Finally, if σ is the positive measure on \mathbb{R}/\mathbb{Z} corresponding to f via Bochner's theorem, then σ and the 1-periodic measure $\overline{\gamma_{dens}}$ are related via the so-called Weil formula:

$$\int_{\mathbb{R}} f(x) d\overline{\gamma_{dens}}(x) = \int_{\mathbb{R}/\mathbb{Z}} \mathcal{W}f(x+\mathbb{Z}) d\sigma(x) \qquad \forall f \in C_{\mathsf{c}}(G) \,.$$

where

$$\mathcal{W}f(x+\mathbb{Z}) \coloneqq \sum_{y\in\mathbb{Z}} f(x+y) \qquad \forall x\in\mathbb{R}.$$

Proof. The first part follows trivially from Proposition 3.9. The last part follows from [21]. \Box

Let us note here in passing that intuitively, the Weil formula says that $\overline{\gamma_{\text{dens}}}$ is the 1-periodic measure obtained by "periodizing" σ . More precisely, σ is the measure obtained by restricting $\overline{\gamma_{\text{dens}}}$ to [0, 1) and identifying this interval with \mathbb{R}/\mathbb{Z} .

Now, exactly the same result holds for the counting autocorrelation. Since the proof is similar, we skip it.

Proposition 3.16. Let $\Lambda \subseteq \mathbb{Z}$ and let A_n be a van Hove sequence in \mathbb{R} . Then, γ_{count} exists with respect to A_n if and only if g_n converges pointwise to a function $g:\mathbb{Z} \to \mathbb{C}$.

Moreover, in this case $\gamma_{count} = \sum_{m \in \mathbb{Z}} g(m) \delta_m$ and g is positive definite. Finally, if σ is the positive measure on \mathbb{R}/\mathbb{Z} corresponding to g via Bochner's theorem, then σ and the 1-periodic measure $\overline{\gamma_{count}}$ are related via the Weil formula.

4. DIFFRACTION OF SETS WITH NO INFINITE TRANSLATED SUBSETS

As we saw in the last section, the autocorrelation encodes how often an element $t \in G$ appears as the difference of two points of Λ . The easiest examples to study, then, are those sets with periodic subsets or those where Λ and $t + \Lambda$ have little agreement. In this section we look at the former situation.

Lemma 4.1. Let $\Lambda \subseteq G$ be a set of FLC and let $\mathcal{A} = (A_n)$ be a van Hove sequence. Assume that for $F_n = \Lambda \cap A_n$

- (a) $\lim_{n \to \infty} \operatorname{card}(F_n) = \infty$.
- (b) For all $0 \neq t \in \Lambda \Lambda$ we have

$$\operatorname{card}(\Lambda \cap (-t + \Lambda)) < \infty$$
 .

Then, the counting autocorrelation is $\gamma_{count} = \delta_0$ and it exists with respect to \mathcal{A} .

Proof. For any nonzere $0 \neq t \in \Lambda - \Lambda$, we can bound

$$\frac{\operatorname{card}(F_n \cap (-t + F_n))}{\operatorname{card}(F_n)} \leq \frac{\operatorname{card}(\Lambda \cap (-t + \Lambda))}{\operatorname{card}(F_n)},$$

where assumptions (a) and (b) imply that the right-hand side tends to zero. So, for nonzero t, in the notation of Lemma 3.10, we have

$$\eta_{\text{count}}(t) = \lim_{n} \frac{\operatorname{card}(F_n \cap (-t + F_n))}{\operatorname{card}(F_n)} = 0.$$

For t = 0, $\eta_{\text{count}}(0) = 1$, because F_n is eventually nonempty. So, each limit $\eta_{\text{count}}(t)$ exists, and so by Lemma 3.10, the autocorrelation γ_{count} exists with respect to \mathcal{A} and it is

$$\gamma_{\text{count}} = \sum_{t \in \Lambda - \Lambda} \eta_{\text{count}}(t) \delta_t = \delta_0.$$

Remark 4.2.

(a) Let $\Lambda \subseteq G$ be locally finite and let $K \subseteq G$ be compact. A short computation shows that

$$\begin{aligned} |\gamma_{F_n} - \delta_0|(K) &= |\gamma_{F_n} - \delta_0|(\{0\}) + |\gamma_{F_n}|(K \setminus \{0\}) \\ &= |\gamma_{F_n}(\{0\}) - 1| + \frac{1}{\operatorname{card}(F_n)} \sum_{x,y \in F_n} \delta_{x-y}(K \setminus \{0\}) \\ &= \left| \frac{1}{\operatorname{card}(F_n)} \delta_{F_n} * \delta_{F_n}(\{0\}) - 1 \right| + \frac{1}{\operatorname{card}(F_n)} \sum_{\substack{t \in (F_n - F_n) \cap K \\ t \neq 0}} \operatorname{card}(F_n \cap (-t + F_n)) \\ &\leq \left| \frac{1}{\operatorname{card}(F_n)} \sum_{x,y \in F_n} \delta_{x-y}(\{0\}) \right| + \frac{1}{\operatorname{card}(F_n)} \sum_{\substack{t \in (\Lambda - \Lambda) \cap K \\ t \neq 0}} \operatorname{card}(\Lambda \cap (-t + \Lambda)). \end{aligned}$$

Now, if Λ has FLC, then the set $(\Lambda - \Lambda) \cap K$ is finite. Therefore, if the assumptions in Lemma 4.1 hold, then we get the stronger conclusion that

$$\left|\gamma_{F_n} - \delta_0\right|(K) \to 0$$

for all compact sets K.

(b) The conclusion of Lemma 4.1 does not hold for sets without FLC. For example,

$$\Lambda = \bigcup_{n} \left\{ n! + \frac{1}{e^n}, \dots, n! + \frac{\lfloor e^n \rfloor}{e^n} \right\}$$

has density zero, meets any of its translates in at most one point, but one can show that

$$\gamma_{count} = \lambda|_{[-1,1]}$$

As an immediate consequence of Lemma 4.1 we get:

Lemma 4.3. Let $\Lambda = \{a_n : n \in \mathbb{N}\} \subset \mathbb{R}$ be any sequence satisfying

$$\lim (a_{n+1} - a_n) = \infty$$

Then, Λ has FLC, and for any van Hove sequence A_n such that $\lim_{n \to \infty} (A_n \cap \Lambda) = \infty$ we have

(4.1)
$$\gamma_{count} = \delta_0, \quad \widehat{\gamma_{count}} = \lambda.$$

In particular, (4.1) holds for all van Hove sequences which are nested and exhausting.

Proof. First note that for each R > 0 there exists some N such that, for all n > N we have

$$\Lambda \cap [a_n - R, a_n + R] = \{a_n\}.$$

It follows immediately that Λ has FLC.

Note that the given condition implies that for all $t \neq 0$ the set

 $\Lambda \cap (t + \Lambda)$

is finite. Therefore, for any van Hove sequence for which $\lim_n \operatorname{card}(F_n) = \infty$, we get $\gamma_{\text{count}} = \delta_0$ by Lemma 4.1.

Finally, if A_n is exhausting, $\bigcup_n A_n = \mathbb{R}$ and so $\bigcup_n F_n = \Lambda$. If (A_n) is nested, so is the sequence (F_n) , and so since Λ is infinite, we must have $\lim_n \operatorname{card}(F_n) = \infty$.

The following two-sided version is proven in exactly the same way, and we skip the details.

Lemma 4.4. Let $\Lambda = \{a_n : n \in \mathbb{Z}\} \subset \mathbb{R}$ so that

$$\lim_{n \to \infty} a_{n+1} - a_n = \infty \quad and$$
$$\lim_{n \to \infty} a_{n-1} - a_n = -\infty.$$

Then, Λ has FLC, and for any van Hove sequence A_n such that $\lim_{n \to \infty} (A_n \cap \Lambda) = \infty$ we have

(4.2)
$$\gamma_{count} = \delta_0, \quad \widehat{\gamma_{count}} = \lambda.$$

In particular, (4.2) holds for all van Hove sequences which are nested and exhausting.

The next three examples follow immediately from these results.

Example 4.5. Choose any a > 1, let

$$\Lambda = \{a^n : n \in \mathbb{N}\},\$$

and let $A_n = [-n, n]$. Then $\gamma_{count} = \delta_0$ and $\widehat{\gamma_{count}} = \lambda$.

Example 4.6. Let

$$\Lambda = \{n! : n \in \mathbb{N}\}$$

and let $A_n = [-n, n]$. Then $\gamma_{count} = \delta_0$ and $\widehat{\gamma_{count}} = \lambda$.

Example 4.7. Let

$$\Lambda = \{ f_n : n \in \mathbb{N} \}$$

where f_n is the Fibonacci sequence $f_0 = 0, f_1 = 1$ and $f_{n+1} = f_n + f_{n-1}$. Let $A_n = [-n, n]$. Then $\gamma_{count} = \delta_0$ and $\widehat{\gamma_{count}} = \lambda$.

More generally, the same behaviour is exhibited by many point sets arising from linear recurrence relations, in particular when the largest root in modulus of the characteristic polynomial is strictly greater than 1. The point being that all of these have approximately exponential growth.

It should be noted that these examples can be put into Proposition 3.13 to get a point set with diffraction and autocorrelation

$$\gamma_{\text{count}} = \delta_0 + \frac{\delta_k + \delta_{-k}}{2} \quad \text{and} \quad \widehat{\gamma_{\text{count}}} = (1 + \cos(2\pi kx))\lambda.$$

5. DIFFRACTION OF THE PRIMES

In this section, we discuss the diffraction of primes $\mathbb{P} \subseteq \mathbb{Z}$ (including negative primes) for van Hove sequences of intervals. We will see that the density autocorrelation is always 0, while the counting autocorrelation is, under weak conditions, δ_0 .

Throughout this entire section, for $d \in \mathbb{N}$, let us recall the notation

$$\pi_d(x) = \operatorname{card}\{p \in \mathbb{P} \cap [0, x] : p + d \in \mathbb{P}\}.$$

Of course, $\pi_0(x)$ is just the prime counting function $\pi(x)$. Since there is only one even prime, for odd d we have

$$\pi_d(x) \le 1$$

for any x.

5.1. Density autocorrelation with respect to van Hove sequences of intervals. Consider a van Hove sequence $A_n = [a_n, b_n]$ of intervals. By eventually replacing a_n, b_n by nearby integers, we can assume without loss of generality that $a_n, b_n \in \mathbb{Z}$.

A simple consequence of the Brun-Titchmarsh theorem is that the density of the primes with respect to A_n is zero. Indeed, let us recall the following version of this theorem:

Theorem 5.1. [17, Corollary 2] Let m, n be positive integers. Then

$$\pi(m+n) - \pi(m) \le 2\pi(n).$$

As an immediate consequence, we get:

Theorem 5.2. Let $A_n = [a_n, b_n]$ be any van Hove sequence of intervals. Then,

$$\operatorname{card}(\mathbb{P} \cap A_n) \le \frac{2\pi(b_n - a_n)}{b_n - a_n}$$

In particular, with respect to A_n we have dens(\mathbb{P}) = 0 and so the density diffraction is

$$\gamma_{dens}$$
 = 0.

Proof. Let $r_n = b_n - a_n$. Then, since A_n is a van Hove sequence, we have $\lim_n r_n = \infty$. Let $F_n := \mathbb{P} \cap A_n$.

We split the problem into three cases:

<u>Case 1:</u> Suppose $0 < a_n$. Then, by Theorem 5.1 we have

$$\frac{\operatorname{card}(F_n)}{r_n} = \frac{\pi(b_n) - \pi(a_n)}{r_n} \le 2\frac{\pi(r_n)}{r_n}$$

<u>Case 2:</u> Suppose $b_n < 0$. Then, by Theorem 5.1 we have

$$\frac{\operatorname{card}(F_n)}{r_n} = \frac{\pi(|a_n|) - \pi(|b_n|)}{r_n} \le 2\frac{\pi(r_n)}{r_n} \,.$$

<u>Case 3:</u> Suppose $a_n \leq 0 \leq b_n$. Then, $|a_n| < r_n$ and $b_n < r_n$, and hence

$$\frac{\operatorname{card}(F_n)}{r_n} = \frac{\pi(b_n) + \pi(|a_n|)}{r_n} \le 2\frac{\pi(r_n)}{r_n} \,.$$

Therefore, for all n we have

$$\frac{\operatorname{card}(F_n)}{r_n} \le 2\frac{\pi(r_n)}{r_n} \,.$$

Since $r_n \to \infty$, the last claim follows from the Prime number Theorem.

Remark 5.3. Let $\mathbb{X}(\mathbb{P})$ be the dynamical system generated by \mathbb{P} under the translation action of \mathbb{R} , see [4] for details. Then, we have $\emptyset \in \mathbb{X}(\mathbb{P})$.

Now, for each $\varphi_1, \ldots, \varphi_n \in C_{\mathsf{c}}(\mathbb{R})$ define $F_{\varphi_1, \ldots, \varphi_n} : \mathbb{X}(\mathbb{P}) \to \mathbb{C}$ via

$$F_{\varphi_1,\ldots,\varphi_n}(\Lambda) \coloneqq \prod_{j=1}^n \left(\sum_{x \in \Lambda} \varphi_j(x)\right).$$

Let

$$\mathbb{A} := \{ F_{\varphi_1, \dots, \varphi_n}(\Lambda) + c1_{\mathbb{X}(\mathbb{P})} : 0 \le n, \varphi_1, \dots, \varphi_n \in C_{\mathsf{c}}(\mathbb{R}), c \in \mathbb{R} \}$$

Then, $A \subseteq C(X(\mathbb{P}))$ is a subalgebra separating the points and hence dense in $C(X(\mathbb{P}))$ by Stone–Weierstrass.

Now, it follows immediately from Theorem 5.2 that

$$\lim_{n} \frac{1}{2n} \int_{t-n}^{t+n} F_{\varphi_1,\dots,\varphi_n}(s+\mathbb{P}) ds = 0 = \int_{\mathbb{X}(\mathbb{P})} F_{\varphi_1,\dots,\varphi_n}(\Gamma) d\delta_{\emptyset}(\Gamma)$$

uniformly in t. Also, we trivially have

$$\lim_{n} \frac{1}{2n} \int_{t-n}^{t+n} c \mathbb{1}_{\mathbb{X}(\mathbb{P})}(s+\mathbb{P}) ds = c = \int_{\mathbb{X}(\mathbb{P})} c \mathbb{1}_{\mathbb{X}(\mathbb{P})}(\Gamma) d\delta_{\emptyset}(\Gamma) ds$$

The density of \mathbb{A} in $C(\mathbb{X}(\mathbb{P}))$ immediately implies that $(\mathbb{X}(\mathbb{P}), \mathbb{R})$ is uniquely ergodic, with unique ergodic measure $m = \delta_0$. Therefore, by [4], the primes must have density and autocorrelation zero with respect to any van Hove sequence (not necessarily of intervals).

The same conclusion holds if we replace the translation action of \mathbb{R} by the translation action of \mathbb{Z} .

5.2. Counting diffraction with respect to van Hove sequences of intervals. Numerical estimates have reported that the diffraction of primes is pure point, meaning that the primes are a model for quasicrystals [27] (compare [26, 28]). This would mean that the (counting) diffraction would have the form

$$\widehat{\gamma_{\text{count}}} = \sum_{y \in B} I(y) \delta_y$$
.

This seems too nice to be true. Indeed, for a set $\Lambda \subseteq \mathbb{Z}$, the much weaker statement that $\widehat{\gamma_{\text{count}}} \neq \lambda$ is equivalent to the fact that there exists some $d \in \mathbb{Z} \setminus \{0\}$ such that

$$\limsup_{n} \frac{\operatorname{card}(\Lambda \cap (d+\Lambda) \cap [-n,n])}{\operatorname{card}(\Lambda \cap [-n,n])} > 0.$$

When $\Lambda = \mathbb{P}$, this would imply that there exists a positive integer d > 0 such that the set $\{p \in \mathbb{P} : p + d \in \mathbb{P}\}$ would have positive density among the primes.

As mentioned above, such a result is too nice to be true. It would give some results which are much much stronger than the latest developments in number theory ([29, 20, 16]). In fact, it turns out that this would contradict some known results in this area, and hence cannot be true. This means that the diffraction of the primes must be the Lebesgue measure

$$\widehat{\gamma_{\mathrm{count}}} = \lambda$$

We now prove that this is indeed the case. We show that the counting autocorrelation γ_{count} exists along many van Hove sequences of intervals and it is $\gamma_{\text{count}} = \delta_0$. This implies that the diffraction is the Lebesgue measure and hence absolutely continuous.

First, we cover the following generalisation of Brun's Theorem.

Proposition 5.4. For all $d \ge 1$ we have

$$\lim_{x \to \infty} \frac{\pi_d(x)}{\pi_0(x)} = 0.$$

Proof. For even d, it is well known using sieve methods in number theory that there exists a constant C > 0 such that

$$\pi_d(x) \le C \frac{x}{\log^2 x}, \quad \forall x \ge 1.$$

Since we are not experts, we do not know precisely who came up with this argument but it can be found in Bateman and Stemmler [6, Lemma 3]. Therefore, by the prime number theorem,

$$0 \le \limsup_{x \to \infty} \frac{\pi_d(x)}{\pi_0(x)} \le \lim_{x \to \infty} \frac{Cx/\log^2 x}{x/\log x} = 0,$$

so $\lim_{x\to\infty} \pi_d(x)/\pi_0(x) = 0$.

Now note that if $F \subseteq \mathbb{R}$ is any finite set we have

$$\gamma_{-F}$$
 = γ_{F}

Since $-\mathbb{P} = \mathbb{P}$, for any set A we have

$$(5.1) \qquad \qquad -(\mathbb{P} \cap A) = \mathbb{P} \cap (-A).$$

Therefore, we immediately get :

Lemma 5.5. Let A_n be any van Hove sequence, let $\epsilon_n \in \{\pm 1\}$ and let $B_n = \epsilon_n A_n$. Let $\mathcal{A} = \{A_n\}, \mathcal{B} = \{B_n\}$. Then, the counting autocorrelation $\gamma_{count,\mathcal{A}}$ of \mathbb{P} exists with respect to \mathcal{A} if and only if the counting autocorrelation $\gamma_{count,\mathcal{B}}$ of \mathbb{P} exists with respect to \mathcal{B} . Moreover, in this case $\gamma_{count,\mathcal{A}} = \gamma_{count,\mathcal{B}}$.

Whenever when we are given a van Hove sequence $A_n = [a_n, b_n]$ of intervals, Lemma 5.5 will allow us to flip some of those intervals so that $|b_n| \ge |a_n|$. Since, after the flip, we also have $a_n < b_n$, we get that $b_n \ge 0$ and must have

$$\lim_{n} b_n = \infty$$

because the width of the intervals goes to infinity. We will often make this extra assumption.

Next, let us cover the following two results.

Lemma 5.6. Let $A_n = [a_n, b_n]$ be a van Hove sequence of intervals such that for all n we have $0 \in A_n$. Then, with respect to A_n we have $\gamma_{count} = \delta_0$.

Proof. By (5.1) and Lemma 5.5 we can assume without loss of generality that $|b_n| \ge |a_n|$. This implies that $b_n > 0$ and $\lim_n b_n = \infty$. As usual, let $F_n = \mathbb{P} \cap A_n$.

For each $t \neq 0$ we have

$$\operatorname{card}(F_n \cap (t + \mathbb{P})) \le 2\pi_{|t|}(b_n)$$
$$\operatorname{card}(F_n) \ge \pi(b_n).$$

The claim now follows from Proposition 5.4 and Lemma 3.11.

As a consequence we get:

Theorem 5.7. The primes \mathbb{P} have counting autocorrelation and diffraction with respect to the van Hove sequence $A_n = [-n, n]$

7

$$\gamma_{count} = \delta_0,$$

 $\widehat{\gamma_{count}} = \lambda.$

Remark 5.8. Combining the proofs of Lemma 5.6, Proposition 5.4, Lemma 3.11 and the Prime Number Theorem, it follows that, for each van Hove sequence A_n of intervals such that $0 \in A_n$ there exists a constant $C_1 > 0$ such that for all $m \in \mathbb{Z}$ we have

$$|\gamma_{F_n} - \delta_0|(m) \le \frac{2\pi_{|m|}(n) + |m|}{\pi_0(n)} \le C_1\left(\frac{1}{\log(n)} + \frac{|m|\log(n)}{n}\right).$$

This seem to suggest that γ_{F_n} converges very slowly to γ_{count} , meaning that numerical estimates are useless when studying the diffraction of primes.

Lemma 5.9. Let $A_n = [a_n, b_n]$ be a van Hove sequence of intervals such that for all n we have $0 < a_n < b_n$. Assume that there exists some c > 1 and N such that for all n > N we have $b_n \ge ca_n$.

Then, with respect to A_n the primes have counting autocorrelation $\gamma_{count} = \delta_0$.

Proof. For each $t \neq 0$ we have

$$\operatorname{card}(F_n \cap (t + \mathbb{P})) \le \pi_{|t|}(b_n)$$
$$\operatorname{card}(F_n) = \pi(b_n) - \pi(a_n) \ge \pi(b_n) - \pi(b_n/c)$$

Now, by Proposition 5.4 and the prime number theorem we have

$$\lim_{n} \frac{\pi_{|t|}(b_n)}{\frac{b_n}{\ln(b_n)}} = 0 \quad \text{and}$$
$$\lim_{n} \frac{\pi(b_n) - \pi(b_n/c)}{\frac{b_n}{\ln(b_n)}} = 1 - \frac{1}{c} > 0.$$

It follows that for all $t \neq 0$ we have

$$\lim_{n} \frac{\operatorname{card}(F_n \cap (t + \mathbb{P}))}{\operatorname{card}(F_n)} = \lim_{n} \frac{\pi_{|t|}(b_n)}{\pi(b_n) - \pi(b_n/c)} = 0.$$

The claim follows from Lemma 3.11.

Remark 5.10. The condition that $b_n \ge ca_n$ is important and cannot be dropped. For instance, $A_n = [n!, n! + n]$ is a van Hove sequence that intersects \mathbb{P} in at most one point, because n! + k is always divisible by k. With respect to this van Hove sequence we have

$$\gamma_{count} = 0$$

Let us now state a few immediate consequences of the previous result.

Corollary 5.11. Let a_n, b_n be any sequences of integers such that $b_n > a_n > 0$ and $\lim_n b_n = \infty$. If

$$\lim_{n} \frac{b_n}{a_n} = L > 1 \,.$$

Then, $A_n = [a_n, b_n]$ is a van Hove sequence and with respect to this van Hove sequence we have

$$\gamma_{count} = \delta_0 , \quad \widehat{\gamma_{count}} = \lambda$$

We can now prove our most general version of these results.

Theorem 5.12. Let $A_n = [a_n, b_n]$ be any van Hove sequence of intervals. Assume that there exists some constant d > 0 such that

$$a_n | < d(b_n - a_n)$$

Then, with respect to A_n we have $\gamma_{count} = \delta_0$ and $\widehat{\gamma_{count}} = \lambda$.

Proof. As usual, let $r_n = b_n - a_n$.

First let us note that

$$|b_n| \le |a_n| + |b_n - a_n| \le (d+1)r_n$$
.

In particular, for all n we have

$$\max\{|a_n|, |b_n|\} \le (d+1)r_n$$

Define

$$B_n = \begin{cases} \begin{bmatrix} a_n, b_n \end{bmatrix} & \text{if } |a_n| \le |b_n| \\ \begin{bmatrix} -b_n, -a_n \end{bmatrix} & \text{if } |a_n| \ge |b_n| \end{cases}$$

Then, by Lemma 5.5, it suffices to prove that our conclusion holds with respect to B_n . Note that if $B_n = [a'_n, b'_n]$ we have

$$|b'_n| \le (d+1)(b'_n - a'_n).$$

Therefore, without loss of generality, we can assume that $|b_n| \ge |a_n|$ for all n. As before, this implies that $b_n > 0$ and $b_n \to \infty$.

Define

$$A \coloneqq \{n : a_n \le 0\}$$
$$B \coloneqq \{n : a_n > 0\}$$

We split the proof into three cases.

<u>Case 1:</u> A is finite. Then, by eventually erasing the first few terms of A_n , we can assume that $a_n > 0$ for all n.

Next, we have

$$\frac{b_n}{a_n} = \frac{b_n - a_n}{a_n} + 1 \ge d + 1 > 1 \,.$$

The conclusion follows from Lemma 5.9.

<u>Case 2:</u> B is finite. Then, by eventually erasing the first few terms of A_n , we can assume that $a_n \leq 0$ for all n. The conclusion then follows from Lemma 5.6.

<u>Case 3:</u> A, B are infinite. Then write A as the sequence $k_1, k_2, \ldots, k_n \ldots$ and B as the sequence $m_1, m_2, \ldots, m_n \ldots$

By Lemma 5.6, as in Case 2, we have

$$\lim_n \gamma_{F_{k_n}} = \delta_0 \,.$$

Also, exactly as in Case 1, we have

$$\frac{b_{m_n}}{a_{m_n}} \ge d+1 > 1$$

Therefore, by Lemma 5.9 we have

$$\lim_n \gamma_{F_{m_n}} = \delta_0 \,.$$

 $\lim_n \gamma_{F_n} = \delta_0 \,,$

Since $A \cup B = \mathbb{N}$ we conclude that

as claimed.

Corollary 5.13. Let A_n be a van Hove sequence of intervals which have a common point c. Then, with respect to A_n we have $\gamma_{count} = \delta_0$ and $\widehat{\gamma_{count}} = \lambda$.

In particular, the conclusion holds for any nested van Hove sequence of intervals.

Proof. Let $A_n = [a_n, b_n]$ and let $r_n = b_n - a_n$. Then

$$\frac{a_n}{r_n} < \frac{c}{r_n}$$

which is bounded from above since $\lim_{n \to \infty} \frac{c}{r_n} = 0$. Moreover

$$\frac{-a_n}{r_n} = 1 - \frac{b_n}{r_n} \le 1 - \frac{c}{r_n}$$

which again is bounded from above.

Therefore, we are in the situation of Theorem 5.12.

Remark 5.14. If we do not work with van Hove sequences of intervals, there are many other auto-correlations which are possible. Indeed, let $Q \subseteq \mathbb{P}$ be any finite set.

24

Then, $A_n = [n!, n! + n] \cup Q$ is a van Hove sequence and $F_n = A_n \cap \mathbb{P} = Q$. Comparing to Remark 5.10, it follows that

$$\gamma_{count} = \frac{1}{\operatorname{card}(Q)} \sum_{p,q \in Q} \delta_{p-q}, \qquad \widehat{\gamma_{count}}(x) = \frac{1}{\operatorname{card}(Q)} \sum_{p \in Q} \left| e^{2\pi i (q-p)x} \right|^2.$$

By combining Proposition 3.13 and Theorem 5.7 we also get the following example:

Example 5.15. Let k be a nonzero integer and let

$$\Lambda = \{p, p+k \mid p \in \mathbb{P}\} = \mathbb{P} \cup (k+\mathbb{P}).$$

Then, with respect to the van Hove sequence [-n,n], $\gamma_{count} = \delta_0 + \frac{1}{2}(\delta_k + \delta_{-k})$ and so

 $\widehat{\gamma_{count}} = (1 + \cos(2\pi kx))\lambda$.

5.3. Diffraction of prime powers. Consider now the set

$$\Lambda = \{\pm p^n : p \in \mathbb{P}, n \in \mathbb{N}\}$$

of positive and negative prime powers. The point sets $\mathbb{P} \subseteq \Lambda \subseteq \mathbb{R}$ and $A_n = [-n, n]$ satisfy the conditions of Lemma 3.14.

Indeed, a standard computation (see for example [23]) shows that for all n we have

$$\operatorname{card}(\Lambda \cap [-n, n]) = 2\operatorname{card}(\Lambda \cap [0, n]) = 2(\pi_0(n) + \pi_0(\sqrt{n}) + \pi_0(\sqrt[3]{n}) + \ldots + \pi_0(\sqrt[m]{n})).$$

where $m = \lfloor \log_2(n) \rfloor + 1$ since $n < 2^{m+1}$. This implies that

$$0 \leq \frac{\operatorname{card}(\Lambda \cap [-n, n])}{\operatorname{card}(\mathbb{P} \cap [-n, n])} - 1 \leq \frac{\pi_0(\sqrt{n}) + \pi_0(\sqrt[3]{n}) + \dots + \pi_0(\sqrt[m]{n})}{\pi_0(n)} \leq \frac{\sqrt{n} + \sqrt[3]{n} + \dots + \sqrt[m]{n}}{\pi_0(n)} \leq \frac{(\lfloor \log_2(n) \rfloor + 1) \cdot \sqrt{n}}{\pi_0(n)}$$

Therefore, by the Prime Number Theorem, we get

$$\lim_{n} \frac{\operatorname{card}(\Lambda \cap [-n, n])}{\operatorname{card}(\mathbb{P} \cap [-n, n])} = 1$$

Lemma 3.14 and Theorem 5.12 then give:

Proposition 5.16. Let $A_n = [a_n, b_n]$ be any van Hove sequence of intervals. Assume that there exists some constant d > 0 such that

$$|a_n| < d(b_n - a_n).$$

Then, with respect to A_n the set Λ of powers of primes has counting autocorrelation $\gamma_{count} = \delta_0$ and diffraction $\widehat{\gamma_{count}} = \lambda$.

5.4. **Primes with fixed distance.** We complete the section on primes by discussing the diffraction of twin primes and other similar sets. While we cannot explicitly calculate the diffraction of the twin primes, as the answer would need us first to settle the twin prime conjecture, we can show that the diffraction of the twin primes is different from the diffraction of the primes.

Fix a positive integer $d \ge 1$. Define

$$\mathbb{P}_d \coloneqq \{p, p+d \colon p, p+d \in \mathbb{P}\}.$$

When d = 2, the set \mathbb{P}_2 is exactly the set of twin primes.

Proposition 5.17. Let $d \ge 1$ be any positive integer. Then,

(a) If \mathbb{P}_d is non-empty and finite then the autocorrelation γ_{count} exists with respect to $A_n = [-n,n]$ and

$$\begin{split} \gamma_{count} &= \frac{1}{\mathrm{card}(\mathbb{P}_d)} \sum_{p,q \in \mathbb{P}_d} \delta_{p-q} \neq \delta_0, \\ \widehat{\gamma_{count}} &= \frac{1}{\mathrm{card}(\mathbb{P}_d)} \sum_{p,q \in \mathbb{P}_d} e^{2\pi i (p-q)x} \neq \lambda. \end{split}$$

Moreover,

 $\frac{1}{2} \le \gamma_{count}(\{d\})$

with equality if and only if p, p + d, p + 2d cannot be prime at the same time.

(b) Assume that \mathbb{P}_d is infinite and d is not divisible by 3. Let γ_{count} be the counting autocorrelation of \mathbb{P}_d with respect to a subsequence of [-n, n]. Then,

$$\gamma_{count}(\{d\}) = \frac{1}{2},$$
$$\widehat{\gamma_{count}} \neq \lambda.$$

Proof. Let us define

$$\Lambda \coloneqq \{ p \in \mathbb{P}_d : p + d \in \mathbb{P}_d \}.$$

and note that

(5.2) $\mathbb{P}_d = \Lambda \cup (d + \Lambda).$

(a) Since \mathbb{P}_d is finite, we have

$$F_n = \mathbb{P}_d \cap [-n, n] = \mathbb{P}_d$$

for n large enough. The two formulas are then immediate.

Next, (5.2) implies that

$$\operatorname{card}(\mathbb{P}_d) \leq 2\operatorname{card}(\Lambda)$$

with equality if and only if $\Lambda \cap (d + \Lambda) = \emptyset$. Now,

$$\gamma_{\text{count}}(\{d\}) = \frac{1}{\text{card}(\mathbb{P}_d)} \sum_{p,q \in \mathbb{P}_d} \delta_{p-q}(d)$$
$$= \frac{1}{\text{card}(\mathbb{P}_d)} \sum_{q,q+d \in \mathbb{P}_d} 1 = \frac{\text{card}(\Lambda)}{\text{card}(\mathbb{P}_d)} \ge \frac{1}{2}$$

with equality if and only if $\Lambda \cap (p + \Lambda) = \emptyset$. This is in turn equivalent to saying that p, p + d, p + 2d are never primes at the same time.

(b) Next, let us note that if

 $p \in \Lambda \cap (d + \Lambda)$

then $p, p + d, p + 2d \in \mathbb{P}$. Since $3 \neq d$, one of p, p + d, p + 2d must be divisible by 3. This shows that

$$\Lambda \cap (d + \Lambda) \subseteq \{-3 - 2d, -3 - d, -3, 3, 3 - d, 3 - 2d\}$$

and hence $\Lambda \cap (d + \Lambda)$ is finite.

A short computation similar to the proof of Proposition 3.13 shows that

$$\gamma_{\text{count}}(\{d\}) = \frac{1}{2}$$

The claim follows.

Since $3, 5, 7 \in \mathbb{P}$ we obtain:

Corollary 5.18. Let γ_{count} be any counting autocorrelation of the twin primes \mathbb{P}_2 with respect to a subsequence of [-n,n]. Then, $\gamma_{count}(\{2\}) = \frac{1}{2}$ if and only if there are infinitely many twin primes.

Remark 5.19.

- (a) The diffraction of the set \mathbb{P}_2 of twin primes is not λ .
- (b) When \mathbb{P}_d is finite then the diffraction is absolutely continuous. It is unclear to us what happens when \mathbb{P}_d is infinite, for example, for the values of d from [29, 20, 16].

The above result shows that, as long as d is not a multiple of 3, if \mathbb{P}_d is infinite, then \mathbb{P}_d and \mathbb{P} have different diffraction. It is reasonable to expect that for infinite \mathbb{P}_d , the set \mathbb{P}_d and \mathbb{P} have different diffraction even when 3|d, but the above proof does not work anymore. To make the proof work, one would need to show that the set of primes p for which both p + d and p + 2d are also prime has asymptotic density zero inside \mathbb{P}_d , which looks reasonable but extremely hard to show.

6. Subsets of zero density with non-trivial Bragg spectrum

All the examples we have seen so far have continuous counting diffraction spectrum. We show below that this is not always the case. The following result allows us to construct sets of density zero whose counting diffraction spectrum can be the density diffraction of any relatively dense set that we want.

Theorem 6.1. Let $\Lambda \subseteq \mathbb{Z}$ be relatively dense, and assume that the density autocorrelation $\gamma_{dens} = \gamma$ of Λ exists with respect to $A_n = [-n, n]$.

For each $n \ge 2$, set

$$\Gamma_n = n! + (\Lambda \cap [-n, n])$$

and define

$$\Gamma \coloneqq \bigcup_{n>2} \Gamma_n,$$

which is a disjoint union. Then, with respect to $A_n = [-n, n]$ we have

- (a) dens(Γ) = 0, and
- (b) $\gamma_{count} = C\gamma$ where

$$C = \frac{1}{\gamma(\{0\})} = \frac{1}{\operatorname{dens}(\Lambda)} \neq 0.$$

Proof. Let $\Lambda_n = \Lambda \cap A_n = \Lambda \cap [-n, n]$, so that $\Gamma_n = n! + \Lambda_n$ and $\operatorname{card}(\Gamma_n) = \operatorname{card}(\Lambda_n) \le 2n + 1$. Denote $F_n = \Gamma \cap A_n$.

(a) Let $n \in \mathbb{N}$, and let m be the unique natural number for which

$$m! \le n < (m+1)!.$$

Then $F_n \subseteq \bigcup_{k=2}^{m+1} \Gamma_k$, and so

$$\frac{\operatorname{card}(F_n)}{2n} = \frac{1}{2n} \sum_{k=2}^{m+1} \operatorname{card}(\Gamma_k \cap [-n, n])$$
$$\leq \frac{1}{2n} \sum_{k=2}^{m+1} \operatorname{card}(\Gamma_k) \leq \frac{m(2m+1)}{2n} \leq \frac{m(2m+1)}{2m!} \to 0$$

as $m \to \infty$. As $n \to \infty$, we also have $m \to \infty$, and so therefore

dens_{$$\mathcal{A}$$}(Γ) = $\lim_{n} \frac{\operatorname{card}(F_n)}{2n} = 0$

(b) Let $t \in \mathbb{Z}$. For readability, introduce the notation

$$\ell_n = \operatorname{card}(\Lambda_n) = \operatorname{card}(\Gamma_n), \text{ and}$$

 $c_n = \operatorname{card}(\Lambda_n \cap (-t + \Lambda)).$

Then, by Lemma 3.9, we know that

$$\gamma(\{0\}) = \operatorname{dens}_{\mathcal{A}}(\Lambda) = \lim_{n} \frac{\ell_n}{2n} \neq 0 \quad \text{and} \quad \gamma(\{t\}) = \lim_{n} \frac{c_n}{2n}.$$

Our goal is to show that

$$\lim_{n} \frac{\operatorname{card}(F_n \cap (-t + \Gamma))}{\operatorname{card}(F_n)} = \frac{\gamma(\{t\})}{\gamma(\{0\})} = \lim_{n} \frac{c_n}{\ell_n}$$

and then the result follows from Lemma 3.10. We will accomplish this by showing that $\frac{\operatorname{card}(F_n \cap (-t+\Gamma))}{\operatorname{card}(F_n)}$ is asymptotically a ratio of the Césaro means for (c_n) and (ℓ_n) .

As in the proof of (a), let $n \in \mathbb{N}$ and let $m \in \mathbb{N}$ be the unique number satisfying $m! \le n \le (m+1)!$. Then

$$\bigcup_{k=2}^{m} \Gamma_k \subseteq F_n \subseteq \bigcup_{k=2}^{m+1} \Gamma_k,$$

with these unions disjoint, and so

$$\sum_{k=2}^{m} \ell_k \le \operatorname{card}(F_n) \le \sum_{k=2}^{m+1} \ell_k.$$

Eventually, as k becomes large, k! is large enough that $t + \Gamma_k$ intersects Γ only in Γ_k . That is, for some constant N depending only on t, if $k \ge N$, then

$$\Gamma_k \cap (-t + \Gamma) = \Gamma_k \cap (-t + \Gamma_k),$$

which has cardinality $\operatorname{card}(\Gamma_k \cap (-t + \Gamma_k)) = \operatorname{card}(\Lambda_k \cap (-t + \Lambda_k)) = c_k$. (In fact, we could take N = |t|, though a smaller number will typically work.) Now, assume that n is large enough that $m \ge N$, and estimate

$$\operatorname{card}(F_n \cap (-t + \Gamma)) \ge \sum_{k=2}^m \operatorname{card}(\Gamma_k \cap (-t + \Gamma))$$
$$\ge \sum_{k=N}^m \operatorname{card}(\Gamma_k \cap (-t + \Gamma))$$
$$= \sum_{k=N}^m c_k = \sum_{k=2}^m c_k - C,$$

where $C = \sum_{k=2}^{N-1} c_k$ is a constant not depending on n (only on Λ and t). And, for an upper bound,

$$\operatorname{card}(F_n \cap (-t + \Gamma)) \leq \sum_{k=2}^{m+1} \operatorname{card}(\Gamma_k \cap (-t + \Gamma))$$
$$= \sum_{k=2}^{N-1} \operatorname{card}(\Gamma_k \cap (-t + \Gamma)) + \sum_{k=N}^{m+1} \operatorname{card}(\Gamma_k \cap (-t + \Gamma))$$
$$\leq \sum_{k=2}^{N-1} \operatorname{card}(\Gamma_k) + \sum_{k=N}^{m+1} \operatorname{card}(\Gamma_k \cap (-t + \Gamma_k))$$
$$= \sum_{k=2}^{N-1} \ell_k + \sum_{k=N}^{m+1} c_k \leq \sum_{k=2}^{m+1} c_k + D,$$

where $D = \sum_{k=2}^{N-1} (\ell_k - c_k)$ is a constant not depending on n. Combining all of our bounds so far shows that for large enough n,

(6.1)
$$\frac{\sum_{k=2}^{m} c_k - C}{\sum_{k=2}^{m+1} \ell_k} \le \frac{\operatorname{card}(F_n \cap (-t + \Gamma))}{\operatorname{card}(F_n)} \le \frac{\sum_{k=2}^{m+1} c_k + D}{\sum_{k=2}^{m} \ell_k}$$

Now, because Λ_n is relatively dense, there must be a constant R > 0 so that

 $Rn \leq \ell_n \leq 2n+1$

for all n. So,

$$\frac{\sum_{k=2}^{m+1} \ell_k}{\sum_{k=2}^m \ell_k} - 1 = \frac{\ell_{m+1}}{\sum_{k=2}^m \ell_k} \le \frac{2m+1}{\sum_{k=2}^m Rk} = \frac{2m+1}{R\frac{m(m+1)}{2} - R} \to 0$$

as $m \to \infty$, so

(6.2)
$$\lim_{m \to \infty} \frac{\sum_{k=2}^{m+1} \ell_k}{\sum_{k=2}^m \ell_k} = 1.$$

Now, as $n \to \infty$, $m \to \infty$, and Equation (6.2) gives

$$\lim_{m \to \infty} \frac{\sum_{k=2}^{m} c_k - C}{\sum_{k=2}^{m+1} \ell_k} = \lim_{m \to \infty} \frac{\sum_{k=2}^{m} c_k - C}{\sum_{k=2}^{m} \ell_k} = \lim_{m \to \infty} \frac{c_m}{\ell_m}$$

by the Stolz-Césaro Theorem. Similarly,

$$\lim_{m \to \infty} \frac{\sum_{k=2}^{m+1} c_k + D}{\sum_{k=2}^m \ell_k} = \lim_{m \to \infty} \frac{\sum_{k=2}^{m+1} c_k + D}{\sum_{k=2}^{m+1} \ell_k} = \lim_{m \to \infty} \frac{c_{m+1}}{\ell_{m+1}} = \lim_{m \to \infty} \frac{c_m}{\ell_m}.$$

Now, the Squeeze Theorem applied to Equation (6.1) yields

$$\eta_{\text{count}}(t) = \lim_{n \to \infty} \frac{\operatorname{card}(F_n \cap (-t + \Gamma))}{\operatorname{card}(F_n)} = \lim_{m \to \infty} \frac{c_m}{\ell_m} = \frac{\gamma(\{t\})}{\gamma(\{0\})}.$$

Lemma 3.10 now shows that γ_{count} exists for Γ with respect to \mathcal{A} , and it is $\gamma_{\text{count}} = \frac{1}{\gamma(\{0\})}\gamma$.

Remark 6.2. The conclusion of Theorem 6.1 holds if in the definition of Γ_n , the sequence n! is replaced by any increasing sequence a_n of natural numbers with the property that

$$\lim_{n} \frac{a_{n+1} - a_n}{n^2} = \infty \,.$$

The proof in this more general situation is a straightforward modification of the proof of Theorem 6.1.

By applying this result to various known examples, we get a list of subsets of \mathbb{Z} of density zero covering most spectral types. Note that the primes already provide an example of a set of density zero with a purely absolutely continuous spectrum; the only spectral type currently missing is purely singular continuous. Such an example cannot be produced by the method of this section, as a relatively dense subset of \mathbb{Z} always has a trivial Bragg peak at the origin.

Example 6.3. Let $\Lambda = \mathbb{Z}$ and let

$$\Gamma = \bigcup_{n=2}^{\infty} \{n! - n, n! - n + 1, \dots, n! + n\}$$

Then Γ has

$$\begin{aligned} \gamma_{count} &= \delta_{\mathbb{Z}} \quad and \\ \widehat{\gamma_{count}} &= \delta_{\mathbb{Z}}, \end{aligned}$$

which are the density autocorrelation and diffraction of \mathbb{Z} , respectively.

Example 6.4. Let Λ be the positions of a's in the Thue-Morse substitution (see [1, Sect. 4.6]) and let

$$\Gamma = \bigcup_{n=2}^{\infty} n! + (\Lambda \cap [-n, n])$$

Then

$$\widehat{\gamma_{count}} = \delta_{\mathbb{Z}} + \omega$$

where ω is a \mathbb{Z} -periodic singular continuous measure (see [1, Remark 10.3]).

Example 6.5. Let Λ be the positions of a's in the Rudin-Shapiro substitution (see [1, Sect. 4.7]) and let

$$\Gamma = \bigcup_{n=2}^{\infty} n! + (\Lambda \cap [-n, n])$$

Then

$$\widehat{\gamma_{count}} = \frac{1}{2}\delta_{\mathbb{Z}} + \lambda$$

where λ is the Lebesgue measure on \mathbb{R} (see [1, Remark 10.5]).

Example 6.6. Let Λ' be the Thue-Morse sequence, and let Λ be a generic element for the Bernoulisation process on Λ' [1, Subsection 11.2.2]. Then, $\Lambda \subseteq \mathbb{Z}$ almost surely has diffraction

$$\widehat{\gamma_{dens}} = \frac{1}{4}\delta_{\mathbb{Z}} + \frac{1}{4}\omega + \frac{1}{4}\lambda$$

where ω is the singular continuous measure from Example 6.4 (see [1, Remark 11.3]). Thm. 6.1 then produces a set Γ of density zero with diffraction spectrum containing all three components.

7. Embedding point sets in higher dimension

Let us now briefly look at a different situation where sets of density zero appear naturally. Let $\Lambda \subseteq \mathbb{R}^d$ be an uniformly discrete point set. Then we can identify Λ with $\Lambda \times \{0\} \subseteq \mathbb{R}^{d+m}$. Using the standard van Hove sequence $A_n = [-n, n]^{d+m}$ we have

$$\operatorname{dens}(\Lambda) = 0$$
.

This means that the classical definition of (density) diffraction would always give a diffraction

 $\widehat{\gamma_{dens}} = 0$.

But now consider an (idealized) multiple-slit interference experiment. This can simply be described by punching tiny holes into a wall at locations $\Lambda \subseteq \mathbb{R}$ and doing a 2-dimensional diffraction experiment. The outcome of the experiment is usually a mixture of lights and shadows on a wall, which are periodic in one direction and change in the opposite direction.

We will see below that the counting autocorrelation precisely describes this phenomena. Recall first that given two measures μ, ν on G and H respectively, there exists a unique product measure $\mu \times \nu$ on $G \times H$ with the property that

$$\int_{G \times H} f(x)g(y) \mathrm{d}\mu \times \nu(x,y) = \left(\int_G f(x) \mathrm{d}\mu(x)\right) \left(\int_H g(y) \mathrm{d}\nu(y)\right), \qquad \forall f \in C_{\mathsf{c}}(G), g \in C_{\mathsf{c}}(H)$$

With this notation, we trivially have

$$\delta_x \times \delta_y = \delta_{(x,y)} \, .$$

Let us now show how the counting autocorrelation behaves when we increase the dimension of the underlying space.

Theorem 7.1. Let $\Lambda \subseteq G$ be any point set, and assume that the autocorrelation $\gamma_{count,\Lambda}$ exists with respect to some van Hove sequence A_n .

Let H be any second countable LCAG and B_n any van Hove sequence in H, such that $0 \in B_n$ for all n. Let $C_n = A_n \times B_n \subseteq G \times H$.

Define

$$\Gamma = \Lambda \times \{0\} \subseteq G \times H$$

Then, the counting autocorrelation $\gamma_{count,\Gamma}$ of $\Gamma \subseteq G \times H$ exists with respect to C_n and

$$\gamma_{count,\Gamma} = \gamma_{count,\Lambda} \times \delta_0$$
.

In particular,

$$\widehat{\gamma_{count,\Gamma}} = \widehat{\gamma_{count,\Lambda}} \times \theta_{\hat{H}}$$

where $\theta_{\hat{H}}$ is a Haar measure on \hat{H} .

Proof. Define

$$F_n \coloneqq \Lambda \cap A_n$$
$$E_n = \Gamma \cap C_n$$

Then, since $0 \in B_n$, we have

$$E_n = F_n \times \{0\}$$

and hence

 $\delta_{E_n} = \delta_{F_n} \times \delta_0$.

It follows immediately that

$$\gamma_{E_n} = \frac{1}{\operatorname{card}(E_n)} \delta_{E_n} * \widetilde{\delta_{E_n}} = \frac{1}{\operatorname{card}(F_n)} \delta_{F_n \times \{0\}} * \widetilde{\delta_{F_n \times \{0\}}}$$
$$= \left(\frac{1}{\operatorname{card}(F_n)} \delta_{F_n} * \widetilde{\delta_{F_n}}\right) \times \delta_0 = \gamma_{F_n} \times \delta_0,$$

from which the rest of the proof follows.

Example 7.2. Consider $\mathbb{Z} \times \{0\} \subseteq \mathbb{R}^d$. Then, its counting diffraction is

$$\widehat{\gamma_{count}} = \delta_{\mathbb{Z}} \times \lambda$$

Example 7.3. Let $\Lambda \subseteq \mathbb{R}$ be the Silver mean model set (see [1, Example 4.5]) and let $\Gamma = \Lambda \times \{0\} \subseteq \mathbb{R}^{1+m}$. Then, the diffraction of Γ is

$$\widehat{\gamma_{count}} = \sigma \times \lambda$$
,

where

$$\sigma = \sum_{k \in \frac{1}{2}\mathbb{Z} + \frac{1}{2\sqrt{2}}\mathbb{Z}} A_k \delta_k$$

is the pure point measure from [1, Theorem 9.3].

Acknowledgements

A.H. was partially supported by the NSERC Discovery grant 2024-03883, C.R. was supported by the NSERC Discovery grant 2019-05430, and N.S. was supported by the NSERC Discovery grants 2020-00038 and 2024-04853.

References

- M. Baake and U. Grimm, Aperiodic Order. Vol. 1. A Mathematical Invitation, Encyclopedia of Mathematics and its Applications 149, Cambridge University Press, Cambridge, 2013.
- [2] M. Baake and U. Grimm (eds.), Aperiodic Order. Vol. 2: Crystallography and Almost Periodicity, Encyclopedia of Mathematics and its Applications 166, Cambridge University Press, Cambridge, 2017.
- [3] M. Baake, C. Huck, and N. Strungaru, On weak model sets of extremal density, Indag. Math. 28 (2017), 3-31.
- [4] M. Baake and D. Lenz, Dynamical systems on translation bounded measures: pure point dynamical and diffraction spectra, Ergod. Th. Dynam. Sys. 24 (2004), 1867–1893.
- [5] M. Baake and R.V. Moody, Weighted Dirac combs with pure point diffraction, J. Reine Angew. Math. 573 (2004), 61–94.
- [6] P. Bateman and R. Stemmler, Waring's problem for algebraic number fields and primes of the form $(p^r-1)/(p^d-1)$, Illinois J. Math. 6 (1962), 142-156.
- [7] C. Berg and G. Forst, Potential Theory on Locally Compact Abelian Groups, Springer, Berlin, 1975.
- [8] J.M. Cowley, Diffraction Physics, 3rd ed., North-Holland, Amsterdam, 1995.
- [9] J. Dieudonné, Treatise on analysis, vol. 2, Academic Press, 1970.
- [10] S. Dworkin, Spectral theory and X-ray diffraction, J. Math. Phys. 34 (1993), 2965–2967.
- [11] J.-B. Gouéré, Diffraction and Palm measure of point processes, C. R. Acad. Sci. Paris 336 (2003), 57–62.
- [12] U. Grimm and X. Deng, Some comments on pinwheel tilings and their diffraction, J. Phys.: Conf. Ser. 284 (2011).
- [13] A. Hof, On diffraction by aperiodic structures, Commun. Math. Phys. 169 (1995), 25–43.
- [14] G. Keller and C. Richard, Dynamics on the graph of the torus parametrisation, Ergod. Th. Dynam. Sys. 38 (2018), 1048–1085.
- [15] G. Keller and C. Richard, Periods and factors of weak model sets, Israel J. Math. 229 (2019), 85–132.
- [16] J. Maynard, Small gaps between primes, Ann. Math. **181** (2015), 383–413.
- [17] H. L. Montgomery, R. C. Vaughan, The large sieve, Mathematika 20, 119–134, 1973.
- [18] R.V. Moody and N. Strungaru, Almost Periodic Measures and their Fourier Transforms, In: [2], pp. 173–270, 2017.
- [19] G.K. Pedersen, Analysis Now, Springer, New York, 1989; Revised printing 1995.
- [20] D.H.J. Polymath, Variants of the Selberg sieve, and bounded intervals containing many primes, Res. Math. Sci. 1 (2014).
- [21] C. Richard and N. Strungaru, Fourier analysis of measures with lattice support, in preperation (2024).
- [22] W. Rudin, Real and Complex Analysis, McGraw-Hill, New York, 1987.
- [23] Samuel, What's the asymptotic distribution of pⁿ (powers of primes)?, Mathstackexchange, q/396665, 2016. https://math.stackexchange.com/q/396665
- [24] M. Schlottmann, Generalised model sets and dynamical systems. In: Directions in Mathematical Quasicrystals, eds. M. Baake, R.V. Moody, CRM Monogr. Ser., AMS, Providence, RI, pp. 143–159, 2000.
- [25] N. Strungaru, On the Fourier analysis of measures with Meyer set support, J. Funct. Anal. 278, (2020).

- [26] S. Torquato, G. Zhang, and M. de Courcy-Ireland, Hidden multiscale order in the primes, J. Physics A 52 (2018).
- [27] S. Torquato, G. Zhang, and M. de Courcy-Ireland, Uncovering multiscale order in the prime numbers via scattering, J. Statistical Mechanics (2018).
- [28] G. Zhang, F. Martelli, and S. Torquato, The structure factor of primes, J. Physics A 51 (2018).
- [29] Y. Zhang, Bounded gaps between primes, Ann. Math. 179 (2014), 1121–1174.

DEPARTMENT OF MATHEMATICS AND COMPUTING, MOUNT ROYAL UNIVERSITY CALGARY, ALBERTA, CANADA *Email address:* ahumeniuk@mtroyal.ca

DEPARTMENT OF MATHEMATICS AND STATISTICS, MACEWAN UNIVERSITY, EDMONTON, ALBERTA, CANADA Email address: ramseyc5@macewan.ca URL: https://sites.google.com/macewan.ca/chrisramsey/

DEPARTMENT OF MATHEMATICS AND STATISTICS, MACEWAN UNIVERSITY, EDMONTON, ALBERTA, CANADA, AND INSTITUTE OF MATHEMATICS "SIMON STOILOW", BUCHAREST, ROMANIA Email address: strungarun@macewan.ca

 $\mathit{URL:}\ \texttt{https://sites.google.com/macewan.ca/nicolae-strungaru/home}$