

A nonlocal approximation of the area in codimension two

Michele Caselli, Mattia Freguglia and Nicola Picenni

Abstract

For $s \in (0, 1)$ we introduce a notion of fractional s -mass on $(n - 2)$ -dimensional closed, orientable surfaces in \mathbb{R}^n . Moreover, we prove its Γ -convergence, with respect to the flat topology, and pointwise convergence to the $(n - 2)$ -dimensional area.

Contents

1	Introduction	2
1.1	Main result	3
1.2	Motivation and related works	4
1.3	Future directions	6
2	Definitions and notations	7
2.1	Linking number between submanifolds	7
2.2	Fractional Sobolev spaces	9
2.3	Slicing with subspaces	11
2.4	Flat convergence of boundaries	14
3	Prescribing singularities and weakly linking maps	15
3.1	Existence of a competitor	15
3.2	Existence of a minimizer	18
3.3	Linking with squares	18
3.4	Positivity of the fractional mass	20
4	Proof of the main result	23
4.1	Lower bound (proof of Proposition 1.2)	24
4.2	Upper bound (proof of Proposition 1.3)	34
5	Possible generalizations	40
5.1	Ambient Riemannian manifolds	40
5.2	Higher codimension	40
A	Appendix: energy asymptotics of the planar vortex	42

Mathematics Subject Classification (2020): 49J45, 28A75, 49Q20.

Keywords: Fractional mass, fractional Sobolev spaces, Gamma-convergence, flat convergence, codimension two submanifolds, Ginzburg-Landau energy.

1 Introduction

In this paper, for every $s \in (0, 1)$, we consider a notion of fractional s -mass for surfaces of codimension two in \mathbb{R}^n , with $n \geq 3$, and we prove that it suitably converges to the $(n - 2)$ -dimensional Hausdorff measure.

In order to define this quantity, let $\Sigma \subset \mathbb{R}^n$ be a closed (compact without boundary), oriented, $(n - 2)$ -dimensional surface of class C^2 , with locally constant integer multiplicity. More precisely, we consider

$$\Sigma := d_1 \Sigma_1 \cup \dots \cup d_m \Sigma_m, \quad (1.1)$$

where $m \geq 1$ is a positive integer, $d = (d_1, \dots, d_m) \in \mathbb{N}_+^m$, and $\Sigma_1, \dots, \Sigma_m$ are closed, connected, oriented, $(n - 2)$ -dimensional surfaces of class C^2 . With a little abuse of notation, we use the same symbol Σ to denote the submanifold with multiplicities (1.1), the associated integral current (see subsection 2.4) and the set $\Sigma_1 \cup \dots \cup \Sigma_m \subset \mathbb{R}^n$.

Let us consider the following set of curves

$$L(\Sigma) := \{\gamma \subset \mathbb{R}^n \setminus \Sigma : \gamma \text{ is a bi-Lipschitz image of } \mathbb{S}^1\}. \quad (1.2)$$

For every $s \in (0, 1)$ we consider the following class of maps

$$\mathfrak{F}_s(\Sigma) := \left\{ u \in C^1(\mathbb{R}^n \setminus \Sigma; \mathbb{S}^1) : \mathcal{E}_{\frac{1+s}{2}}(u, \mathbb{R}^n) < +\infty \text{ and} \right. \\ \left. |\deg(u, \gamma)| = |d_1 \text{link}(\gamma, \Sigma_1) + \dots + d_m \text{link}(\gamma, \Sigma_m)| \text{ for any } \gamma \in L(\Sigma) \right\}, \quad (1.3)$$

where $|\deg(u, \gamma)|$ denotes the absolute value of the degree of the restriction of u to γ , $\text{link}(\gamma, \Sigma_i)$ denotes the linking number between γ and Σ_i , that is how many times γ winds around Σ_i (see Subsection 2.1 for a precise definition), and

$$\mathcal{E}_\alpha(u, \Omega) := \iint_{\mathbb{R}^n \times \mathbb{R}^n \setminus \Omega^c \times \Omega^c} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2\alpha}} dx dy$$

is defined more in general for any $\alpha \in (0, 1)$ and every open set $\Omega \subset \mathbb{R}^n$.

Moreover, if $u \in \mathfrak{F}_s(\Sigma)$ we say that u *links with* Σ . For our purposes, it is useful to extend this definition of linking between a function and a surface to less regular functions. To this aim, we consider the following class of maps

$$\mathfrak{F}_s^w(\Sigma) := \left\{ u : \mathbb{R}^n \rightarrow \mathbb{S}^1 : \exists \{u_k\} \subset \mathfrak{F}_s(\Sigma), u_k \rightarrow u \text{ in } H_{\text{loc}}^\alpha(\mathbb{R}^n; \mathbb{S}^1) \forall \alpha \in \left(0, \frac{1+s}{2}\right) \right\}. \quad (1.4)$$

In this case, if $u \in \mathfrak{F}_s^w(\Sigma)$ we say that u *weakly links with* Σ .

We can now introduce the s -fractional mass of Σ as the least possible fractional energy of a map that weakly links with Σ , namely

$$\mathbb{M}_s(\Sigma) := \min_{u \in \mathfrak{F}_s^w(\Sigma)} \mathcal{E}_{\frac{1+s}{2}}(u, \mathbb{R}^n). \quad (1.5)$$

Here the use of the exponent $(1 + s)/2$ in the fractional space is motivated by the fact that this choice yields the expected scaling property of \mathbb{M}_s (in analogy with the fractional perimeter), namely

$$\mathbb{M}_s(\lambda \cdot \Sigma) = \lambda^{n-1-s} \cdot \mathbb{M}_s(\Sigma) \quad \forall \lambda > 0,$$

where $\lambda \cdot \Sigma$ denotes Σ rescaled by a factor λ .

We prove in Section 3 that for every $s \in (0, 1)$ and every Σ as above, the class $\mathfrak{F}_s(\Sigma)$ is non-empty, and that the minimum in (1.5) is strictly positive and is achieved by a minimizer in $\mathfrak{F}_s^w(\Sigma)$ that, in some weak sense, still enjoys the geometric property that defines the class $\mathfrak{F}_s(\Sigma)$.

1.1 Main result

Our main result concerns the asymptotic behavior of $\mathbb{M}_s(\Sigma)$ in the limit as $s \rightarrow 1^-$. More precisely, we prove the pointwise convergence and the Γ -convergence with respect to the flat norm of boundaries (see Subsection 2.4) of $\mathbb{M}_s(\Sigma)$ to the $(n-2)$ -Hausdorff measure with multiplicity, after multiplication by a suitable rescaling factor.

Let ω_{n-1} be the area of \mathbb{S}^{n-1} , then the precise statement is the following.

Theorem 1.1. *Let $\Sigma \subset \mathbb{R}^n$ be as in (1.1).*

Then it holds that

$$\Gamma\text{-}\lim_{s \rightarrow 1^-} (1-s)^2 \mathbb{M}_s(\Sigma) = \lim_{s \rightarrow 1^-} (1-s)^2 \mathbb{M}_s(\Sigma) = \frac{2\pi\omega_{n-1}}{n} \sum_{i=1}^m d_i \mathcal{H}^{n-2}(\Sigma_i),$$

where the Γ -limit is intended with respect to the flat convergence of boundaries in \mathbb{R}^n .

More precisely, we obtain Theorem 1.1 as a consequence of the following two estimates, which actually provide a stronger local result involving the fractional seminorm, that is

$$[u]_{H^\alpha(\Omega)}^2 := \iint_{\Omega \times \Omega} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2\alpha}} dx dy,$$

for any $\alpha \in (0, 1)$ and every open set $\Omega \subset \mathbb{R}^n$. We point out that $\mathcal{E}_\alpha(u, \Omega) \geq [u]_{H^\alpha(\Omega)}^2$.

Proposition 1.2 (Lower bound). *Let $\{\Sigma_s\}_{s \in (0,1)} \subset \mathbb{R}^n$ and $\Sigma \subset \mathbb{R}^n$ be as in (1.1), and let us assume that $\Sigma_s \rightarrow \Sigma$ as $s \rightarrow 1^-$ with respect to the flat convergence of boundaries. For every $s \in (0, 1)$, let also $u_s \in \mathfrak{F}_s^w(\Sigma_s)$ be any map that weakly links with Σ_s .*

Then it holds that

$$\liminf_{s \rightarrow 1^-} (1-s)^2 [u_s]_{H^{\frac{1+s}{2}}(\Omega)}^2 \geq \frac{2\pi\omega_{n-1}}{n} \sum_{i=1}^m d_i \mathcal{H}^{n-2}(\Sigma_i \cap \Omega), \quad (1.6)$$

for every open set $\Omega \subseteq \mathbb{R}^n$.

Proposition 1.3 (Upper bound). *Let $\Sigma \subset \mathbb{R}^n$ be as in (1.1). Then there exists a function $u: \mathbb{R}^n \rightarrow \mathbb{S}^1$, such that $u \in \mathfrak{F}_s(\Sigma)$ for every $s \in (0, 1)$ and it holds that*

$$\limsup_{s \rightarrow 1^-} (1-s)^2 \mathcal{E}_{\frac{1+s}{2}}(u, \Omega) \leq \frac{2\pi\omega_{n-1}}{n} \sum_{i=1}^m d_i \mathcal{H}^{n-2}(\Sigma_i \cap \bar{\Omega}). \quad (1.7)$$

for every open set $\Omega \subseteq \mathbb{R}^n$.

We observe that in codimension one, that is for the classical fractional perimeter, the correct rescaling factor to get a meaningful (nonzero, not identically infinite) Γ -limit is $(1-s)$ (see [ADPM11]). The reason why $(1-s)^2$ is the correct factor in the codimension two setting will be clear from our proofs, but the reader can see Lemma A.1 in the Appendix for a model case that enlightens this feature.

Remark 1.4. The combination of Proposition 1.2 and Proposition 1.3 shows that our convergence result can be localized, and any reasonable definition of fractional mass inside an open set $\Omega \subset \mathbb{R}^n$, which agrees with (1.5) when $\Omega = \mathbb{R}^n$, would converge to the classical area as $s \rightarrow 1^-$.

In the codimension one case, the fractional perimeter inside Ω of a set $E \subset \mathbb{R}^n$ is usually defined (up to constant factors) by

$$\text{Per}_s(E, \Omega) := \iint_{E \times E^c \setminus \Omega^c \times \Omega^c} \frac{dx dy}{|y - x|^{n+s}} = [\mathbb{1}_E]_{H^{s/2}(\mathbb{R}^n)} - [\mathbb{1}_E]_{H^{s/2}(\Omega^c)},$$

where $\mathbb{1}_E$ denotes the characteristic function of the set E (see [CRS10]).

In our case, a few different ways to define the local fractional mass exist. For example, one could just take (1.5) and replace $\mathbb{R}^n \times \mathbb{R}^n$ with $\mathbb{R}^n \times \mathbb{R}^n \setminus \Omega^c \times \Omega^c$.

Alternatively, one could take a minimizer for (1.5) and compute its fractional energy on $\mathbb{R}^n \times \mathbb{R}^n \setminus \Omega^c \times \Omega^c$. This notion could, in principle, depend on the choice of the minimizer. Nevertheless, it would be the most natural notion if the minimizer for (1.5) turned out to be unique (up to rotations in the target \mathbb{S}^1). Indeed, in this case this definition would be completely analogous to the one of the localized fractional perimeter above, with the unique minimizer playing the role of the characteristic function.

Remark 1.5. It can be checked that the (equivalent) notions of linking number, described in Subsection 2.1, used in this work are also equivalent to the one given by Serra in [Ser23, Section 5] in the multiplicity one case. Indeed, in [Ser23] it is said that $u \in C^1(\mathbb{R}^n \setminus \Sigma; \mathbb{S}^1)$ is linked with Σ if

- (i) For every small 2-dimensional disk $D \subset \mathbb{R}^n$ orthogonal to Σ and with center on Σ we have $|\deg(u, \partial D)| = 1$.
- (ii) For every embedded disk $D \subset \mathbb{R}^n \setminus \Sigma$ not intersecting Σ we have $|\deg(u, \partial D)| = 0$.

Note that (i) only is not sufficient for our purposes. Asking only that (i) holds for tiny disks, say contained in a δ -neighborhood N_δ of Σ , would give a notion of s -fractional mass in (1.5) that depends on δ , and tends to zero as $\delta \rightarrow 0$. Roughly speaking, the minimizer u_\circ would tend to create a vortex of opposite degree just outside N_δ to make the energy small. This issue is resolved by adding (ii).

1.2 Motivation and related works

One of the central questions in modern Riemannian geometry is the existence and regularity of minimal surfaces in various codimension.

Nevertheless, with the exception of the case of geodesics on surfaces that can be treated by parametrization effectively, the existence of such objects turned out to be a challenging problem since regular surfaces (of any codimension) with bounded area is a very non-compact set. To overcome this difficulty, one must either consider weak topologies on larger spaces or approximate the area functional with a better-behaved functional, and recover minimal surfaces after a limit procedure of this relaxation. This approach of relaxing the functional has been successful in many great works such as [Riv17, Gua18, GG18, CM20, PS21].

In codimension one, many different relaxations of the area functional were successful in the construction of minimal hypersurfaces in \mathbb{R}^n or in ambient Riemannian manifolds. For example, building on the seminal works by Modica-Mortola [MM77a, MM77b] and Modica [Mod87] on the Γ -convergence of the Allen-Cahn energy to the classical perimeter, many works succeeded in proving the existence of (two-sided) minimal hypersurfaces in Riemannian manifolds.

The Allen-Cahn energy, provides a PDE-based alternative to the classical min-max theory of existence by Almgren-Pitts [Alm62, Alm65, Pit81]. Using this approach, together with some profound results on the regularity of stable solutions of the Allen-Cahn equation by Tonegawa-Wickramasekera [TW12], Gaspar-Guaraco [GG18] and Guaraco [Gua18] managed to prove the existence of smooth embedded minimal hypersurfaces in ambient Riemannian manifolds. The effectiveness of this approach culminated in the work by Chodosh-Mantoulidis [CM20] where the authors prove the multiplicity-one conjecture and, with it, give a new proof of Yau's conjecture for generic metrics, that was recently shown by Irie-Marques-Neves in [IMN18].

Very recently, a new possible approach to the approximation of minimal surfaces appeared: the one of fractional minimal surfaces. Since their first precise definition and study in the

seminal work [CRS10] by Caffarelli, Roquejoffre and Savin, much interest has been devoted to their study and the study of the fractional perimeter in general. These objects, at first sight, less “natural” than other approximations of minimal surfaces, enjoy a long list of properties utterly analogous to the ones for classical minimal surfaces. We refer to [Dip20, Ser23] for a complete discussion of the similarities with the classical world.

Apart from a long list of similarities, the literature has recently pointed out a few striking differences with the world of classical minimal surfaces.

A first instance of these differences appeared in a work by Cinti-Serra-Valdinoci [CSV19]. In this paper, the authors prove that stable fractional minimal surfaces (say, in a ball of radius one) enjoy an interior uniform bound on the classical perimeter of the surfaces. This is in clear contrast with classical minimal surfaces since many parallel hyperplanes (arbitrarily close) are a stable configuration for the perimeter with, clearly, no uniform bound on the area in any ball.

This feature has an analog for the fractional Allen-Cahn equation proved in [CCS21], where the authors show that stable solutions of the fractional Allen-Cahn equation enjoy a uniform BV -estimate.

Another remarkable result for the fractional Allen-Cahn equation with no parallel in the classical world is the improvement of flatness theorem proved by Dipierro-Serra-Valdinoci in [DSV20]. In this work, the authors show that entire solutions to the fractional Allen-Cahn equation with asymptotically flat level sets (say, in Hausdorff distance) are one-dimensional. This is in contrast with the classical Allen-Cahn equation since the solutions constructed in [dPKW13] concentrate on catenoids, which have arbitrary flat blow-downs in Hausdorff distance but are clearly not one-dimensional.

This feature represents a remarkable departure from the classical world and is in accordance with the fact that nonlocal catenoids have conical, nontrivial blow-downs. This is to say the blow-down of the nonlocal catenoids is not a single plane with multiplicity two. These nonlocal catenoids were constructed in [DdPW18], where the authors prove that in \mathbb{R}^3 there exists a connected, embedded s -minimal surface of revolution whose blow-down is a nontrivial cone. Moreover, in the same work [DdPW18] it is also proved that these nonlocal catenoids have infinite Morse index, in any reasonable sense.

These features suggest that sequences of s -minimal surfaces with uniformly bounded index should enjoy stronger compactness properties than classical minimal surfaces with bounded index. This turns out to be true. Indeed, in the recent paper [CFS23], the first author, Florit and Serra, obtain a uniform BV -estimate for finite index solutions of the fractional Allen-Cahn equation, extending the one of [CCS21] for stable solutions to the case of finite index.

These surprisingly strong estimates for finite Morse index nonlocal minimal surfaces confer exceptional compactness and regularity properties to these objects. Thanks to this feature and a classical min-max method, the authors in [CFS23] establish far-reaching existence result of infinitely many s -minimal surfaces in every n -dimensional closed Riemannian manifold. In the same work, they are also proved to be smooth for $n = 3, 4$ and to have a small singular set for $n \geq 5$. This work suggests that s -minimal surfaces are an ideal class of objects on which to apply min-max methods, as they seem to prevent almost every pathology that arises for classical minimal surfaces (as multiplicity and pinching).

From here, a natural question arises: Can one take advantage of these exceptional compactness and regularity properties of s -minimal surfaces and send $s \rightarrow 1^-$ afterward to recover classical minimal surfaces? Surprisingly, the answer seems to be affirmative.

In the recent work [CDSV23], Chan, Dipierro, Serra, and Valdinoci obtain robust curvature estimates and optimal sheet separation estimates, as $s \rightarrow 1^-$, for stable s -minimal surfaces in three dimensions. This result allows to send $s \rightarrow 1^-$ and to pass to the limit *the supports* of the s -minimal surfaces to obtain standard minimal surfaces.

In codimension two, the goal of finding a diffuse approximation of the codimension two

area that is well-behaved in the limit is an extremely challenging problem. For example, it is known that with the classical complex valued Ginzburg-Landau model, one can produce nontrivial, stationary $(n - 2)$ -dimensional varifolds in ambient Riemannian manifolds. Nevertheless, it has been shown in [PS23] that, in general, these varifolds are not integral and that every density $\theta \in \{1\} \cup [2, \infty)$ on an $(n - 2)$ -plane can be realized as a limit of complex Ginzburg–Landau critical points.

In the groundbreaking work [PS21] by Pigati and Stern, the authors substitute the complex Ginzburg-Landau energy with the Yang-Mills–Higgs energy and can produce stationary, integral $(n - 2)$ -varifolds in the limit. As a byproduct, in codimension two, they obtain a new proof of Almgren’s existence result of nontrivial, stationary integral $(n - 2)$ -dimensional varifolds in closed Riemannian manifolds.

At this point, it is natural to ask if there is a natural codimension two analog of the fractional perimeter that reflects similar properties for stationary objects as the ones recently discovered for s -minimal surfaces. To the authors’ knowledge, tentative notions of codimension ≥ 2 fractional masses appeared just in two very recent preprints: [CHK⁺24] for codimension $(n - 1)$ and [MS23] in any codimension. In the same works, both notions are proved to converge pointwise, as $s \rightarrow 1^-$, to the standard Hausdorff measure of the correct dimension.

The convergence of the fractional area to the classical one in the limit $s \rightarrow 1^-$ is indeed the first basic property that any possible notion need to satisfy if one wants to exploit the fractional approximation to obtain results on classical minimal surface. In the case of the codimension one fractional perimeter, the pointwise convergence is just a particular case of the so-called BBM formula (see [BBM01, Dá02]). It has also been showed in [ADPM11] (see also [Pon04]) that the fractional perimeter converges to the classical perimeter also in the sense of Γ -convergence with respect to the L^1 convergence of sets (that basically corresponds to the flat convergence of their boundaries).

Recently, Serra in [Ser23, Section 5] suggested a notion of s -fractional mass for codimension $k \in \{1, 2, \dots, n - 1\}$ smooth, multiplicity one submanifolds of \mathbb{R}^n that are level sets of (regular values of) maps from \mathbb{R}^n to \mathbb{R}^k .

In this work, we study the case of codimension two in detail and full generality. We prove that this notion is well-defined for closed, oriented (not necessarily connected) codimension two surfaces of locally constant multiplicity. Moreover, we prove its Γ -convergence with respect to the flat topology, and pointwise convergence (as $s \rightarrow 1^-$) to the $(n - 2)$ -dimensional Hausdorff measure with multiplicity.

Our study is robust and suited to be extended naturally to ambient Riemannian manifolds and every codimension. We refer to Section 5 for more details on these generalizations. In particular, on Riemannian manifolds, our notion is well-defined for $(n - 2)$ -dimensional oriented boundaries with integer multiplicity, and this is the natural class to apply min-max methods on ambient Riemannian manifolds.

Compared to their local counterparts, we expect stationary sets for this notion to exhibit enhanced regularity and compactness properties, much like the codimension one scenario of the s -minimal surfaces described above.

1.3 Future directions

This work focuses mainly on the definition and study of the fractional s -mass for codimension two surfaces in \mathbb{R}^n and its convergence to the standard \mathcal{H}^{n-2} Hausdorff measure.

Because of this focus, a few basic questions related but not strictly needed to prove this convergence were left unanswered. In particular, we expect to explore the following few (related) questions regarding the minimization in (1.5) and the characterization of the minimizer, whose structure is left almost entirely open in this work.

(P1) Prove that there is no energy gap taking the infimum on $\mathfrak{F}_s(\Sigma)$ instead of $\mathfrak{F}_s^w(\Sigma)$, that is

$$\inf_{u \in \mathfrak{F}_s(\Sigma)} \mathcal{E}_{\frac{1+s}{2}}(u, \mathbb{R}^n) = \min_{u \in \mathfrak{F}_s^w(\Sigma)} \mathcal{E}_{\frac{1+s}{2}}(u, \mathbb{R}^n).$$

(P2) Prove that every minimizer u_o in (1.5) is smooth in $\mathbb{R}^n \setminus \Sigma$ and therefore $u_o \in \mathfrak{F}_s(\Sigma)$.

Moreover, on an ambient Riemannian manifold the s -mass \mathbb{M}_s is well-defined on $(n - 2)$ -dimensional boundaries with integer coefficients (see Subsection 5.1 on the generalization of our notion to Riemannian manifolds), hence it is natural to wonder whether many critical points of \mathbb{M}_s exist. This question was already posed by Serra in [Ser23, Section 5, Open problem 3] in general.

In particular, since $H_{n-2}(\mathbb{S}^n) = 0$ for all $n \geq 3$, on Riemannian spheres (\mathbb{S}^n, g) there holds that every $(n - 2)$ -dimensional integral cycle is an $(n - 2)$ -dimensional integral boundary. Moreover, see for example the end of Section 6 in [MN16], the space of $(n - 2)$ -dimensional integral cycles endowed with the flat topology is weakly homotopic to $\mathbb{C}\mathbb{P}^\infty$, and $\mathbb{C}\mathbb{P}^\infty$ contains a nontrivial class on every even dimensional homology group. Hence, in the space of $(n - 2)$ -dimensional integral cycles, one could perform min-max for \mathbb{M}_s over these infinitely many nontrivial families. This suggests that the answer to the following open question should be affirmative.

(Q1) For $n \geq 3$, do there exist infinitely critical points of \mathbb{M}_s on every sphere (\mathbb{S}^n, g) ?

Lastly, other than existence of critical points, it is natural to wonder whether local minimizers of \mathbb{M}_s are regular, outside a singular set of suitable dimension. Precisely

(Q2) If Σ is a local minimizer (in some suitable class) of \mathbb{M}_s , is Σ smooth/analytic outside a singular set at most $(n - 4)$ -dimensional?

Organization. In Section 2, we deal with all the preliminary tools that we need to define the fractional mass \mathbb{M}_s . These include the linking number, fractional Sobolev spaces, and the notion of flat convergence.

Section 3 deals with the existence of a competitor in $\mathfrak{F}_s(\Sigma)$ for the minimum problem (1.5) (see Theorem 3.1), the existence of a minimizer for this problem, and some properties of this minimizer. In particular, we prove that any minimizer still links with the surface Σ in a suitable weak sense, and we provide a lower bound for the energy for fixed $s \in (0, 1)$.

Then, Section 4 deals with the most technical part of this work and the proof of the Γ -convergence result of Theorem 1.1.

Lastly, in Section 5, we briefly describe possible generalizations (and variants) of our notion to higher codimension or to ambient Riemannian manifolds.

2 Definitions and notations

2.1 Linking number between submanifolds

Let $n \geq 3$ and $n_1, n_2 \geq 1$ be integers with $n_1 + n_2 = n - 1$. Let M_1 and M_2 be disjoint, closed, oriented, connected C^2 submanifolds of \mathbb{R}^n , with dimensions n_1 and n_2 . We denote by $H_k(X)$ and $H^k(X)$ respectively the k -th singular homology and cohomology groups of the space X , with integer coefficients.

There exist several equivalent definitions of linking number between M_1 and M_2 , and each of them enlightens different properties of this quantity that we need in the sequel of the paper. For this reason, we now consider three of these definitions.

Definition 2.1. Let M_1 and M_2 be as above. The Gaussian linking number $\text{link}_G(M_1, M_2)$ is the degree of the map $G : M_1 \times M_2 \rightarrow \mathbb{S}^{n-1}$ defined by

$$G(x, y) := \frac{x - y}{|x - y|}.$$

Definition 2.2. Let M_1 and M_2 be as above, and let us assume that there exists an oriented $(n_1 + 1)$ -submanifold $N_1 \subset \mathbb{R}^n$ with $\partial N_1 = M_1$ intersecting M_2 transversally. Then the intersection linking number is defined as

$$\text{link}_I(M_1, M_2) := \sum_{p \in N_1 \cap M_2} s(p),$$

where $s(p) = 1$ if the orientation on $T_p N_1 \oplus T_p M_2 = \mathbb{R}^n$ induced by the orientations on N_1 and M_2 agrees with the orientation of \mathbb{R}^n , and $s(p) = -1$ otherwise.

Definition 2.3. Let M_1 and M_2 be as above. Let $[M_1] \in H_{n_1}(M_1)$ be the fundamental class of M_1 and let $[M_1]^* \in H^{n-n_1-1}(\mathbb{R}^n \setminus M_1) = H^{n_2}(\mathbb{R}^n \setminus M_1)$ be its Alexander dual. Let also $[M_2]$ be (the push-forward of) the fundamental class of M_2 in $H_{n_2}(\mathbb{R}^n \setminus M_1)$.

Then the homological linking number is defined as

$$\text{link}_H(M_1, M_2) := [M_1]^*([M_2]).$$

The following proposition shows that these definitions actually agree up to a sign.

Proposition 2.4. Let M_1 and M_2 be as above, and let us assume that there exists N_1 as in Definition 2.2. Then it turns out that

$$\text{link}_G(M_1, M_2) = (-1)^{n_2} \text{link}_I(M_1, M_2) = (-1)^{n_2} \text{link}_H(M_1, M_2). \quad (2.1)$$

Proof. The first equality is basically proved in [ABO03, Section 2.11]. The idea consists in considering a function $F : N_1 \times M_2 \rightarrow \mathbb{R}^n$ such that $F(x, y) = (x - y)/|x - y|$ when $(x, y) \in M_1 \times M_2$, with $F(x, y) = x - y$ when $|x - y|$ is small, and no other zeros outside the diagonal $\{x = y\}$. Then it turns out that

$$\text{link}_G(M_1, M_2) = \deg(F, M_1 \times M_2) = \deg(F, N_1 \times M_2, \mathbb{R}^n, 0),$$

where the expression in the right-hand side is the degree of F as a map defined on the manifold with boundary $N_1 \times M_2$ with values in \mathbb{R}^n , computed with respect to the origin. By the construction of F , this coincides with $(-1)^{n_2} \text{link}_I(M_1, M_2)$, because $F^{-1}(0) = N_1 \cap M_2$ and $\text{sign}(\det dF(p)) = (-1)^{n_2} s(p)$ for every $p \in N_1 \cap M_2$.

The second equality can be proved by analyzing the construction of the Alexander duality. Indeed, if νM_1 is a tubular neighborhood of M_1 that does not intersect M_2 , the Alexander duality can be obtained by composing the following isomorphisms

$$H_{n_1}(M_1) \simeq H_{n_1+1}(\mathbb{R}^n, M_1) \simeq H_{n_1+1}(\mathbb{R}^n \setminus \nu M_1, \partial(\mathbb{R}^n \setminus \nu M_1)) \simeq H^{n_2}(\mathbb{R}^n \setminus M_1).$$

The first isomorphism is given by the inverse of the boundary operator in the long exact sequence of the pair (\mathbb{R}^n, M_1) , so it sends $[M_1]$ to $[N_1]$. The second isomorphism is given by the excision theorem, together with some deformation retraction, and is just restricting $[N_1]$ to $\mathbb{R}^n \setminus \nu M_1$. Finally, the last isomorphism is given by the Poincaré-Lefschetz duality (to be precise, one should apply it in \mathbb{S}^n), again together with a deformation.

This means that $[M_1]^*$ is the Poincaré-Lefschetz dual of $[N_1]$, hence its action on $[M_2]$ is given by the algebraic intersection number (see [Bre93, Section VI.11]), which is exactly $\text{link}_I(M_1, M_2)$. \square

This result allows us to define the linking number $\text{link}(M_1, M_2)$ as any of the three equal quantities appearing in (2.1).

The equivalence of the previous definitions shows that the linking number satisfies several useful properties. For example, Definition 2.1 shows that the linking number is commutative, up to a sign, while from Definition 2.3 it follows that $\text{link}(M_1, M_2) = \text{link}(M_1, M'_2)$ if $[M_2] = [M'_2]$ in $H_{n-2}(\mathbb{R}^n \setminus M_1)$. However, Definition 2.2 provides the simpler way to compute the linking number in concrete situations.

Remark 2.5. Actually, the linking number can be defined for much more general couples of disjoint sets and, since the algebraic intersection number can be defined also for singular cycles, in Proposition 2.4 it is not necessary to assume the existence of a regular submanifold N_1 with $\partial N_1 = M_1$.

In particular, the linking number can be defined also if one of the two submanifolds is a bi-Lipschitz curve in the class $L(\Sigma)$ defined in (1.2). In this paper, however, the only non-smooth objects that we consider are boundaries of two-dimensional squares, in which case extending the three definitions of the linking number and Proposition 2.4 is straightforward.

Remark 2.6. We observe that the choice of the orientations on M_1 and M_2 affects the linking number at most by a sign, and more precisely

$$\text{link}(M_1, M_2) = -\text{link}(-M_1, M_2) = -\text{link}(M_1, -M_2) = \text{link}(-M_1, -M_2),$$

where $-M_1$ and $-M_2$ denote, respectively, the manifolds M_1 and M_2 with opposite orientation.

As a consequence, if Σ is as in (1.1) and $\gamma \in L(\Sigma)$, then

$$|d_1 \text{link}(\gamma, \Sigma_1) + \cdots + d_m \text{link}(\gamma, \Sigma_m)|$$

is independent of the choice of the orientation for γ .

Similarly, we point out that if $u : \mathbb{R}^n \setminus \Sigma \rightarrow \mathbb{S}^1$ is continuous, then $|\deg(u, \gamma)|$ is independent of the choice of an orientation (or a parameterization) for γ , hence the class (1.3) is well-defined.

2.2 Fractional Sobolev spaces

We briefly recall here the function spaces we will work with.

Let $\Omega \subset \mathbb{R}^n$ be an open set. For $\alpha \in (0, 1)$ we denote by

$$H^\alpha(\Omega; \mathbb{R}^2) = \left\{ u \in L^2(\Omega; \mathbb{R}^2) : [u]_{H^\alpha(\Omega)}^2 = \iint_{\Omega \times \Omega} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2\alpha}} dx dy < +\infty \right\},$$

the Sobolev-Slobodeckij space of order s . This is a separable Hilbert space endowed with the norm $\|u\|_{H^\alpha(\Omega)}^2 := \|u\|_{L^2(\Omega)}^2 + [u]_{H^\alpha(\Omega)}^2$. We also set

$$H^\alpha(\Omega; \mathbb{S}^1) := \left\{ u \in H^\alpha(\Omega; \mathbb{R}^2) : |u(x)| = 1 \text{ for almost every } x \in \Omega \right\},$$

As usual, we denote by $H_{\text{loc}}^\alpha(\mathbb{R}^n; \mathbb{S}^1)$ the space of functions $u \in L_{\text{loc}}^2(\mathbb{R}^n; \mathbb{S}^1)$ that belong to $H^\alpha(B_R, \mathbb{S}^1)$ for every ball $B_R \subset \mathbb{R}^n$ with radius $R > 0$ centered in the origin.

For every open set $\Omega \subset \mathbb{R}^n$ with Lipschitz (or empty) boundary, let us consider the following space of functions

$$\mathbb{X}_s(\Omega) := \left\{ u : \Omega \rightarrow \mathbb{S}^1 : u \in H^\alpha(\Omega \cap B_R) \text{ for every } R > 0 \text{ and every } \alpha \in \left(0, \frac{1+s}{2}\right) \right\},$$

endowed with the distance

$$d_{\mathbb{X}_s(\Omega)}(u_1, u_2) := \sum_{k=2}^{+\infty} 2^{-k} \min \left\{ \|u_1 - u_2\|_{H^{\frac{1+s}{2} - \frac{1}{k}}(\Omega \cap B_k)}, 1 \right\}.$$

Since the inclusion $H^\alpha(\Omega \cap B_R; \mathbb{S}^1) \hookrightarrow H^{\alpha'}(\Omega \cap B_R; \mathbb{S}^1)$ is continuous (and actually compact, see e.g. [Leo23, Theorem 8.5]) for every $R > 0$ and every $\alpha' < \alpha$, we deduce that a sequence $\{u_k\} \subset \mathbb{X}_s(\Omega)$ converges to some $u \in \mathbb{X}_s(\Omega)$ with respect to this distance if and only if $u_k \rightarrow u$ in $H^\alpha(\Omega \cap B_R; \mathbb{S}^1)$ for every $R > 0$ and every $\alpha \in (0, (1+s)/2)$. As a consequence, the completeness of the spaces $H^\alpha(\Omega \cap B_R; \mathbb{S}^1)$ implies that also $(\mathbb{X}_s(\Omega), d_{\mathbb{X}_s(\Omega)})$ is a complete metric space.

Remark 2.7. We observe that the class $\mathfrak{F}_s^w(\Sigma)$ defined in (1.4) is actually the closure in $\mathbb{X}_s(\mathbb{R}^n)$ of the class $\mathfrak{F}_s(\Sigma)$ defined in (1.3). Moreover, if a sequence $\{u_k\} \subset H_{\text{loc}}^{\frac{1+s}{2}}(\mathbb{R}^n; \mathbb{S}^1)$ converges weakly in $H_{\text{loc}}^{\frac{1+s}{2}}(\mathbb{R}^n; \mathbb{S}^1)$, then it also converges in the sense of $\mathbb{X}_s(\mathbb{R}^n)$. In particular, any sequence of functions $\{u_k\} \subset \mathbb{X}_s(\mathbb{R}^n)$ for which

$$\sup_{k \in \mathbb{N}} \mathcal{E}_{\frac{1+s}{2}}(u_k, \mathbb{R}^n) < +\infty,$$

is relatively compact in $\mathbb{X}_s(\mathbb{R}^n)$. This follows by a standard weak compactness (see, for example, [MPS21, Remark 2.2]) and diagonal argument. However, the weak topology is not metrizable, and this is the reason for which it is convenient to consider the spaces \mathbb{X}_s .

We conclude this subsection by recalling a few results about fractional Sobolev spaces. While all these facts are well-known, we need the explicit dependence on the parameter s of all the constants, so here state them in the form that we need, and we also include short proofs.

The first result is the following version of the fractional Sobolev embedding in the one-dimensional case.

Lemma 2.8 (Fractional Sobolev embedding). *For every $s_0 \in (0, 1)$ there exists a positive constant $C_S = C_S(s_0) > 0$ such that for every $s \in [s_0, 1)$, every interval $(a, b) \subset \mathbb{R}$, and every function $u : (a, b) \rightarrow \mathbb{R}^2$ it holds that*

$$\text{osc}(u, (a, b))^2 \leq C_S (b-a)^s (1-s) [u]_{H^{\frac{1+s}{2}}((a,b))}^2,$$

where $\text{osc}(u, (a, b))$ denotes the (essential) oscillation of u in (a, b) , namely

$$\text{osc}(u, (a, b)) := \| |u(y) - u(x)| \|_{L^\infty((a,b) \times (a,b))}.$$

In particular, if $[u]_{H^{\frac{1+s}{2}}((a,b))}^2 < +\infty$, then u agrees almost everywhere with a continuous function.

Proof. Let us consider first the case in which $(a, b) = (0, 1)$ and u is scalar-valued, and let us assume by contradiction that there exists $s_0 \in (0, 1)$ such that for every $k \in \mathbb{N}^+$ there exist $s_k \in [s_0, 1)$ and a function $u_k : (0, 1) \rightarrow \mathbb{R}$ such that

$$\text{osc}(u_k, (0, 1)) = 1 \quad \text{and} \quad (1-s_k) [u_k]_{H^{\frac{1+s_k}{2}}((0,1))}^2 < \frac{1}{k}. \quad (2.2)$$

Up to a translation, we can also assume that $\{u_k\}$ is bounded in $L^\infty((0, 1))$.

From [BBM01, Remark 5] we deduce that there exists a universal constant $C > 0$ such that

$$\lim_{k \rightarrow +\infty} [u_k]_{H^{\frac{1+s_0}{2}}((0,1))}^2 \leq \lim_{k \rightarrow +\infty} C \cdot \frac{1-s_k}{2} \cdot [u_k]_{H^{\frac{1+s_k}{2}}((0,1))}^2 = 0,$$

and hence [DNPV12, Theorem 7.1] implies that, up to a subsequence, $\{u_k\}$ converges to some constant function in $H^{\frac{1+s_0}{2}}((0, 1))$.

As a consequence, from [DNPV12, Theorem 8.2] we deduce that $\{u_k\}$ converges to a constant also in $C^{0,\alpha}((0, 1))$, with $\alpha = s_0/2$, but this contradicts the first condition in (2.2).

At this point, the case of a general interval (a, b) follows just by scaling, and the vector-valued case just by applying the inequality component-wise. \square

The second result is a version of the Poincaré inequality. This type of results are well-known, even in a wider generality than what we need for our purposes, see e.g. [BBM02, Theorem 1] or [Pon03, Theorem 1.1]. However, in our case, the proof is simpler and with the same strategy used in the previous lemma (with the L^2 -norm instead of the oscillation) we can prove the following result as well.

Lemma 2.9 (Poincaré inequality). *Let $\alpha_0 \in (0, 1)$ and let $B_1 \subset \mathbb{R}^n$ be the unit ball. Then, there exists a constant $C_P = C_P(\alpha_0, n) > 0$ such that for every $\alpha \in [\alpha_0, 1)$ and for every $u \in H^\alpha(B_1, \mathbb{R}^2)$ it holds that*

$$\|u - \bar{u}\|_{L^2(B_1)}^2 \leq C_P(1 - \alpha)[u]_{H^\alpha(B_1)}^2, \quad (2.3)$$

where $\bar{u} := \int_{B_1} u$ is the average of u in B_1 .

Finally, we need the following extension result.

Lemma 2.10 (Extension theorem). *Let $\alpha_0 \in (0, 1)$ and let $B_R \subset \mathbb{R}^n$ be the ball of radius R . Then there exists a constant $C_E = C_E(\alpha_0, n) > 0$, depending only on α_0 and n , such that for every $\alpha \in [\alpha_0, 1)$ and every $u \in H^\alpha(B_R; \mathbb{R}^2)$ with $\int_{B_R} u = 0$ there exists a function $E(u) \in H^\alpha(\mathbb{R}^n; \mathbb{R}^2)$ with $E(u) = u$ in B_R and*

$$[E(u)]_{H^\alpha(\mathbb{R}^n)}^2 \leq C_E[u]_{H^\alpha(B_R)}^2. \quad (2.4)$$

Proof. We start with the case $R = 1$. A careful reading of [DNPV12, Section 5] shows that there exists a constant $C(n) > 0$ such for any $u \in H^\alpha(B_1)$ there exists $E(u) \in H^\alpha(\mathbb{R}^n)$ such that $E(u) = u$ in B_1 and

$$[E(u)]_{H^\alpha(\mathbb{R}^n)}^2 \leq C(n) \left(\alpha^{-1}[u]_{H^\alpha(B_1)}^2 + \alpha^{-1}(1 - \alpha)^{-1}\|u\|_{L^2(B_1)}^2 \right). \quad (2.5)$$

Indeed, the constant appearing in [DNPV12, Theorem 5.4] depends on the domain Ω (via a covering, a partition of unity and the bi-Lipschitz maps locally parameterizing the boundary), and on the constants coming from [DNPV12, Lemma 5.1 and Lemma 5.3].

Now we observe that the constant $C(n, \alpha, p, K, \Omega)$ in [DNPV12, Lemma 5.1] could actually be written as $C(n, p, K, \Omega)\alpha^{-1}$, while in [DNPV12, Lemma 5.3] the dependence of the constant on α appears only in the estimate

$$\int_{\Omega} dx \int_{\Omega \cap |x-y| < 1} \frac{|u(x)|^p}{|y-x|^{n+(\alpha-1)p}} + \int_{\Omega} dx \int_{\Omega \cap |x-y| > 1} \frac{|u(x)|^p}{|y-x|^{n+\alpha p}} \leq C(n, p, \alpha)\|u\|_{L^p(\Omega)}^p,$$

and in this case $C(n, p, \alpha)$ could be written as $C(n, p)\alpha^{-1}(1 - \alpha)^{-1}$, and it multiplies only the L^p part of the norm. Hence the estimate (5.6) in [DNPV12] in the case $p = 2$ could actually be written as

$$[\psi u]_{H^\alpha(\Omega)} \leq C(n, \Omega, \psi) \left([u]_{H^\alpha(\Omega)}^2 + \alpha^{-1}(1 - \alpha)^{-1}\|u\|_{L^2(\Omega)}^2 \right).$$

In the end, the arguments in [DNPV12, Section 5], in the case $p = 2$ and $\Omega = B_1$, actually yield (2.5). Then, since u has zero average by hypothesis, the Poincaré inequality (2.3) yields (2.4) as desired.

Finally, reasoning component-wise we obtain the same result also when u takes values in \mathbb{R}^2 , and, by scaling, we also deduce that the same holds for every $R > 0$. \square

2.3 Slicing with subspaces

We denote by \mathcal{G}_k^n the Grassmanian of k -dimensional linear subspaces in \mathbb{R}^n . This comes naturally endowed with a unit measure invariant under the action of the orthogonal group $O(n)$.

In the case $k = 1$, it can be easily proved using polar coordinates that

$$\begin{aligned} \iint_{\mathbb{R}^n \times \mathbb{R}^n} f(x, y) dx dy &= \frac{1}{2} \int_{\mathbb{S}^{n-1}} d\theta \int_{\theta^\perp} d\mathcal{H}^{n-1}(h) \iint_{\mathbb{R} \times \mathbb{R}} f(h + \theta x, h + \theta y) |x - y|^{n-1} dx dy \\ &= \frac{\omega_{n-1}}{2} \int_{\mathcal{G}_1^n} dL \int_{L^\perp} d\mathcal{H}^{n-1}(h) \iint_{(L+h) \times (L+h)} f(x, y) |x - y|^{n-1} dx dy, \end{aligned}$$

for every measurable function f . In particular, choosing f of the form

$$f(x, y) = \frac{|u(x) - u(y)|^2}{|x - y|^{n+2\alpha}} \mathbb{1}_{\Omega \times \Omega}(x, y)$$

gives

$$\begin{aligned} [u]_{H^\alpha(\Omega)}^2 &= \iint_{\Omega \times \Omega} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2\alpha}} dx dy \\ &= \frac{\omega_{n-1}}{2} \int_{\mathcal{G}_1^n} dL \int_{L^\perp} d\mathcal{H}^{n-1}(h) \iint_{((L+h) \cap \Omega)^2} \frac{|u(x) - u(y)|^2}{|x - y|^{1+2\alpha}} dx dy \\ &= \frac{\omega_{n-1}}{2} \int_{\mathcal{G}_1^n} dL \int_{L^\perp} [u]_{H^\alpha((L+h) \cap \Omega)}^2 d\mathcal{H}^{n-1}(h). \end{aligned} \quad (2.6)$$

In the case in which $n = 2$ and $\Omega = D_r(x)$ is a disk, from this formula and the invariance by rotations of the measure on $\mathcal{G}_1^n \simeq \mathbb{S}^1$ we deduce that

$$\begin{aligned} [u]_{H^\alpha(D_r(x))}^2 &= \frac{1}{2} \int_{\mathbb{S}^1} d\theta \int_{-r}^r [u]_{H^\alpha(S_{\theta,t}(x))}^2 dt \\ &= \frac{1}{4} \int_{\mathbb{S}^1} d\theta \int_0^r \left([u]_{H^\alpha(S_{\theta,t}(x))}^2 + [u]_{H^\alpha(S_{\theta^\perp,t}(x))}^2 + [u]_{H^\alpha(S_{\theta,-t}(x))}^2 + [u]_{H^\alpha(S_{\theta^\perp,-t}(x))}^2 \right) dt, \end{aligned} \quad (2.7)$$

where θ^\perp denotes the counterclockwise rotation of θ by an angle equal to $\pi/2$, and the set $S_{\theta,t}(x) := \{x + t\theta^\perp + \xi\theta : \xi \in (-\sqrt{r^2 - t^2}, \sqrt{r^2 - t^2})\}$ is the intersection of the line $t\theta^\perp + \theta\mathbb{R}$ with the disk $D_r(x)$. We point out that when $t < r/\sqrt{2}$ the four segments in the right-hand side contain the boundary of a square centered at x .

The next result of this section is a generalization of (2.6) with k -dimensional subspaces instead of lines. This is the content of Theorem 2.12 below.

Before proving it, we recall a classical fact about expressing Sobolev seminorms in terms of the Fourier transform. We adopt the convention of the normalized Fourier transform, that is

$$\mathcal{F}u(\xi) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-ix \cdot \xi} u(x) dx.$$

Proposition 2.11 ([DNPV12, Proposition 3.4]). *Let $\alpha \in (0, 1)$. Then there exists a positive constant $C_F(n, \alpha) > 0$ such that for every $u \in H^\alpha(\mathbb{R}^n)$ it holds that*

$$[u]_{H^\alpha(\mathbb{R}^n)}^2 = C_F(n, \alpha) \int_{\mathbb{R}^n} |\xi|^{2\alpha} |\mathcal{F}u(\xi)|^2 d\xi.$$

Theorem 2.12. *Let $\alpha \in (0, 1)$ and $k \in \{1, \dots, n-1\}$. Then there exists a constant $C(n, k, \alpha) > 0$ such that for every $u \in H^\alpha(\mathbb{R}^n)$ it holds that*

$$[u]_{H^\alpha(\mathbb{R}^n)}^2 = C(n, k, \alpha) \int_{\mathcal{G}_k^n} dL \int_{L^\perp} [u]_{H^\alpha(L+h)}^2 d\mathcal{H}^{n-k}(h),$$

where

$$C(n, k, \alpha) = \frac{C_F(n, \alpha)}{C_F(k, \alpha)} \left(\int_{\mathcal{G}_k^n} |P_L(e_1)|^{2\alpha} dL \right)^{-1},$$

and $P_L : \mathbb{R}^n \rightarrow L$ is the orthogonal projection on the subspace L .

Proof. Fix $u \in H^\alpha(\mathbb{R}^n)$ and denote $(x, y) \in \mathbb{R}^n$ with $x \in \mathbb{R}^k$ and $y \in \mathbb{R}^{n-k}$. By Proposition 2.11 on the x -variables of u (that is, with k in place of n) we have for every fixed $y \in \mathbb{R}^{n-k}$:

$$[u(\cdot, y)]_{H^\alpha(\mathbb{R}^k)}^2 = C_F(k, \alpha) \int_{\mathbb{R}^k} |\xi|^{2\alpha} |\mathcal{F}_x u(\xi, y)|^2 d\mathcal{H}^k(\xi).$$

Completely analogously, we can do the same for every $L \in \mathcal{G}_k^n$. If we denote by $(x_L, y_L) \in L \times L^\perp = \mathbb{R}^n$ the coordinates on \mathbb{R}^n , we get

$$[u(\cdot, y_L)]_{H^\alpha(L)}^2 = C_F(k, \alpha) \int_L |\xi_L|^{2\alpha} |\mathcal{F}_L u(\xi_L, y_L)|^2 d\mathcal{H}^k(\xi_L).$$

Integrating this formula over $y_L \in L^\perp$ and using Plancherel's identity on the y_L -variables gives

$$\begin{aligned} \int_{L^\perp} [u(\cdot, y_L)]_{H^\alpha(L)}^2 d\mathcal{H}^{n-k}(y_L) &= C_F(k, \alpha) \int_{L^\perp} \int_L |\xi_L|^{2\alpha} |\mathcal{F}_L u(\xi_L, y_L)|^2 d\mathcal{H}^k(\xi_L) d\mathcal{H}^{n-k}(y_L) \\ &= C_F(k, \alpha) \int_L |\xi_L|^{2\alpha} \left(\int_{L^\perp} |\mathcal{F}_L u(\xi_L, y_L)|^2 d\mathcal{H}^{n-k}(y_L) \right) d\mathcal{H}^k(\xi_L) \\ &= C_F(k, \alpha) \int_L \int_{L^\perp} |\xi_L|^{2\alpha} |(\mathcal{F}_{L^\perp} \circ \mathcal{F}_L) u(\xi_L, \xi_{L^\perp})|^2 d\mathcal{H}^{n-k}(\xi_{L^\perp}) d\mathcal{H}^k(\xi_L) \\ &= C_F(k, \alpha) \int_{\mathbb{R}^n} |P_L(\xi)|^{2\alpha} |\mathcal{F}u|^2 d\mathcal{H}^n(\xi), \end{aligned}$$

since $P_L(\xi) = \xi_L$.

Since this formula holds for every $L \in \mathcal{G}_k^n$, we can integrate it over the entire Grassmanian \mathcal{G}_k^n to get

$$\int_{\mathcal{G}_k^n} dL \int_{L^\perp} [u(\cdot, y_L)]_{H^\alpha(L)}^2 d\mathcal{H}^{n-k}(y_L) = C_F(k, \alpha) \int_{\mathbb{R}^n} \left(\int_{\mathcal{G}_k^n} |P_L(\xi)|^{2\alpha} dL \right) |\mathcal{F}u|^2 d\mathcal{H}^n(\xi).$$

Moreover, just by scaling and rotational invariance, we see that

$$\int_{\mathcal{G}_k^n} |P_L(\xi)|^{2\alpha} dL = |\xi|^{2\alpha} \int_{\mathcal{G}_k^n} |P_L(\xi/|\xi|)|^{2\alpha} dL = |\xi|^{2\alpha} \int_{\mathcal{G}_k^n} |P_L(e_1)|^{2\alpha} dL,$$

and the last integrals depends only on n, k and α . Putting everything together

$$\begin{aligned} \int_{\mathcal{G}_k^n} dL \int_{L^\perp} [u(\cdot, y_L)]_{H^\alpha(L)}^2 d\mathcal{H}^{n-k}(y_L) &= C_F(k, \alpha) \left(\int_{\mathcal{G}_k^n} |P_L(e_1)|^{2\alpha} dL \right) \int_{\mathbb{R}^n} |\xi|^{2\alpha} |\mathcal{F}u|^2 d\mathcal{H}^n(\xi) \\ &= \frac{C_F(k, \alpha)}{C_F(n, \alpha)} \left(\int_{\mathcal{G}_k^n} |P_L(e_1)|^{2\alpha} dL \right) [u]_{H^\alpha(\mathbb{R}^n)}^2, \end{aligned}$$

where, in the last line, we have used Proposition 2.11 again.

Lastly, since

$$\int_{\mathcal{G}_k^n} dL \int_{L^\perp} [u(\cdot, y_L)]_{H^\alpha(L)}^2 d\mathcal{H}^{n-k}(y_L) = \int_{\mathcal{G}_k^n} dL \int_{L^\perp} [u]_{H^\alpha(L+y_L)}^2 d\mathcal{H}^{n-k}(y_L),$$

this concludes the proof with

$$C(n, k, \alpha) = \frac{C_F(n, \alpha)}{C_F(k, \alpha)} \left(\int_{\mathcal{G}_k^n} |P_L(e_1)|^{2\alpha} dL \right)^{-1} > 0.$$

□

Remark 2.13. Since by [DNPV12, Corollary 4.2] we have

$$\lim_{\alpha \rightarrow 1^-} (1 - \alpha) C_F(n, \alpha) = \frac{\omega_{n-1}}{2n},$$

one can easily see that when $\alpha \geq \alpha_0 \in (0, 1)$ the constant $C(n, k, \alpha)$ in Theorem 2.12 is bounded, above and below, by constants depending only on n and α_0 .

Corollary 2.14. *Let $\alpha_0 \in (0, 1)$ and let $\alpha \in [\alpha_0, 1)$ and $R > 0$. Then there exists a constant $c(\alpha_0, n)$ depending only on α_0 and the space dimension n , such that for every $u \in H^\alpha(B_R)$ and every $k \in \{1, \dots, n-1\}$ it holds that*

$$[u]_{H^\alpha(B_R)}^2 \geq c(\alpha_0, n) \int_{\mathcal{G}_k^n} dL \int_{L^\perp} [u]_{H^\alpha((L+h) \cap B_R)}^2 d\mathcal{H}^{n-k}(h).$$

Proof. Let $\bar{u} := \mathcal{f}_{B_R} u$. By Lemma 2.10 there exists a constant $c(\alpha_0, n) = C_E(\alpha_0, n)^{-1}$ and an extension $E(u - \bar{u})$ such that

$$[u]_{H^\alpha(B_R)} = [u - \bar{u}]_{H^\alpha(B_R)} \geq c(\alpha_0, n) [E(u - \bar{u})]_{H^\alpha(\mathbb{R}^n)}.$$

From Theorem 2.12 we deduce that

$$\begin{aligned} [E(u - \bar{u})]_{H^\alpha(\mathbb{R}^n)} &= C(n, k, \alpha) \int_{\mathcal{G}_k^n} dL \int_{L^\perp} [E(u - \bar{u})]_{H^\alpha((L+h))}^2 d\mathcal{H}^{n-k}(h) \\ &\geq C(n, k, \alpha) \int_{\mathcal{G}_k^n} dL \int_{L^\perp} [E(u - \bar{u})]_{H^\alpha((L+h) \cap B_R)}^2 d\mathcal{H}^{n-k}(h) \\ &= C(n, k, \alpha) \int_{\mathcal{G}_k^n} dL \int_{L^\perp} [u]_{H^\alpha((L+h) \cap B_R)}^2 d\mathcal{H}^{n-k}(h), \end{aligned}$$

where in the last line we have used that $E(u - \bar{u}) = u - \bar{u}$ in B_R and that $[u - \bar{u}]_{H^\alpha((L+h) \cap B_R)}^2 = [u]_{H^\alpha((L+h) \cap B_R)}^2$ by the translation invariance of the seminorm. By Remark 2.13, this concludes the proof. \square

Finally, we recall one of the Crofton formulas for rectifiable sets in \mathbb{R}^n .

Theorem 2.15 (Crofton formula, [SW08, Theorem 5.4.3]). *Let $n \geq 2$, $k \in \{1, \dots, n-1\}$ be integers, and let $M \subset \mathbb{R}^n$ be a q -dimensional rectifiable set with $k + q \geq n$. Then*

$$\int_{\mathcal{G}_k^n} dL \int_{L^\perp} \mathcal{H}^{k+q-n}(M \cap (L+h)) d\mathcal{H}^{n-k}(h) = \beta_{k+q-n, n}^{k, q} \mathcal{H}^q(M),$$

where

$$\beta_{i, j}^{k, \ell} := \frac{\Gamma\left(\frac{k+1}{2}\right) \Gamma\left(\frac{\ell+1}{2}\right)}{\Gamma\left(\frac{i+1}{2}\right) \Gamma\left(\frac{j+1}{2}\right)}.$$

2.4 Flat convergence of boundaries

In this paper, we use integral currents just as a tool to define a distance, and hence a notion of convergence, for submanifolds of \mathbb{R}^n .

For this reason, here we recall the few basic notions we need to define the flat norm, and we refer to [ABO03, ABO05, FF60] for more details.

For $k \in \{1, \dots, n\}$, the space of k -currents in \mathbb{R}^n is defined to be the dual of the space of smooth k -forms with compact support in \mathbb{R}^n .

For every $k \in \{1, \dots, n\}$, we have a boundary operator that associates to a k -current T the $(k-1)$ -current ∂T defined as $\langle \partial T, \omega \rangle := \langle T, d\omega \rangle$ for every $(k-1)$ -form ω .

A k -current T is called *rectifiable* if it can be represented by integration on a rectifiable set, up to an integer multiplicity, namely if

$$\langle T, \omega \rangle = \int_M \theta(x) \omega_x(\tau(x)) d\mathcal{H}^k(x),$$

where M is a k -rectifiable set, $\theta : M \rightarrow \mathbb{Z} \setminus \{0\}$ is an integer-valued measurable function, and $\tau(x)$ is a measurable choice of a unitary k -vector spanning $T_x M$ at \mathcal{H}^k -almost every $x \in M$. In this case we write $T = [M, \theta, \tau]$.

The mass of a rectifiable current $T = [M, \theta, \tau]$ is then defined as

$$\mathbb{M}(T) := \int_M |\theta(x)| d\mathcal{H}^k(x).$$

A k -current is called *integral* if both T and ∂T are rectifiable.

To any submanifold $\Sigma \subset \mathbb{R}^n$ as in (1.1) we can associate the integral $(n - 2)$ -current

$$[\Sigma_1, d_1, \tau_{\Sigma_1}] + \cdots + [\Sigma_m, d_m, \tau_{\Sigma_m}],$$

where $\tau_{\Sigma_i}(x)$ is the unitary $(n - 2)$ -vector given by the orientation of $T_x \Sigma_i$. Stokes' theorem ensures that the boundary operator agrees with the classical notion of boundary on currents coming from submanifolds.

Another special class of integral k -currents that we need in the sequel is the class of *integral Lipschitz chains*, namely Lipschitz images of linear combinations (with integer coefficients) of currents of the form $[P, 1, \tau_P]$, where P is a k -dimensional convex polyhedron. We point out that, through a triangulation, the current associated to any oriented compact manifold can be written as an integral Lipschitz chain.

In [FF60, Theorem 5.11], it was proved that on reasonable subsets of \mathbb{R}^n , and in particular in the whole Euclidean space, the boundary operator on integral currents generates a homology that is equivalent to the singular homology with integer coefficient. In particular, any integral k -current T with $\partial T = 0$ can be obtained as the boundary ∂S of an integral $(k + 1)$ -current S .

Therefore, for an integral k -current without boundary T , we can define its *flat norm of boundaries* (see [ABO05, section 2.5]) as

$$\mathbf{F}(T) := \inf\{\mathbb{M}(S) : S \text{ is an integral } (k + 1)\text{-current with } \partial S = T\}.$$

Consequently, we say that a sequence $\{\Sigma_i\}$ of integral k -currents with null boundary converge to an integral k -current Σ with respect to the flat distance of boundaries if $\mathbf{F}(\Sigma - \Sigma_i) \rightarrow 0$ as $i \rightarrow +\infty$.

Remark 2.16. If $\partial S = T$ is an integral Lipschitz chain, by [FF60, Lemma 5.7] we can approximate S with an integral Lipschitz chain S' with the same boundary $\partial S' = T$. Therefore, in this case, and in particular when T is the difference between two closed oriented submanifolds with multiplicities, we have

$$\mathbf{F}(T) := \inf\{\mathbb{M}(S) : S \text{ is an integral Lipschitz chain with } \partial S = T\}.$$

3 Prescribing singularities and weakly linking maps

3.1 Existence of a competitor

The following result shows that the family $\mathfrak{F}_s(\Sigma)$ is always not empty. This might not be the case for ambient Riemannian manifolds, as there could be topological obstructions to the existence of such a map. See Subsection 5.1 for a short discussion on this feature.

Proposition 3.1. *Let $\Sigma \subset \mathbb{R}^n$ be as in (1.1). Then there exists a function $u : \mathbb{R}^n \rightarrow \mathbb{S}^1$, such that $u \in \mathfrak{F}_s(\Sigma)$ for every $s \in (0, 1)$. Moreover, such map is constant outside a compact set containing Σ .*

There are many results in the literature, similar to our Proposition 3.1, regarding functions with prescribed singularities.

In [ABO03], it has been shown (see in particular [ABO03, Theorem 4.4]) a similar existence result of a map $u : \mathbb{R}^n \rightarrow \mathbb{S}^1$ with prescribed Jacobian equal to (a multiple of) a codimension two integral current with finite mass and no boundary in \mathbb{R}^n . Nevertheless, on the one hand, the result in [ABO03, Theorem 4.4] is more general than ours since it deals with general integral

currents. On the other hand it only shows $u \in W_{\text{loc}}^{1,p}(\mathbb{R}^n; \mathbb{S}^1)$ for every $1 < p < n/(n-1)$, and this range of p is not enough even to conclude that $u \in H_{\text{loc}}^\alpha(\mathbb{R}^n; \mathbb{S}^1)$ for α close to 1.

More recently, in [BM21, Theorem 3.3] the authors proved a similar existence result in the smooth setting and for $n = 3$ in $W_{\text{loc}}^{1,p}(\mathbb{R}^3; \mathbb{S}^1) \cap C^\infty(\mathbb{R}^3 \setminus \Sigma; \mathbb{S}^1)$, for every $1 < p < 2$.

However, for our purposes, it is essential that the map u can be taken constant outside a compact set since this implies that $[u]_{H^\alpha(\mathbb{R}^n)}^2 < +\infty$, and also that it looks like a vortex in each fiber of a tubular neighborhood of Σ , in order to establish Proposition 1.3.

For this reason, we present here a different proof, which relies on the following result about codimension two submanifolds (see [Kir89, Chapter VIII]).

Theorem 3.2. *Let N be an oriented n -dimensional manifold and let $\Sigma \subset N$ be a closed, oriented, connected, $(n-2)$ -dimensional submanifold such that $[\Sigma] = [0] \in H_{n-2}(N)$. Then the normal bundle of Σ in N is trivial, and there exists an oriented hypersurface $M \subset N$ such that $\Sigma = \partial M$.*

In our setting $N = \mathbb{R}^n$, so the assumption of Theorem 3.2 is always trivially satisfied. Hence, in the following proof, for any $i \in \{1, \dots, m\}$, we let $M_i \subset \mathbb{R}^n$ be a so-called ‘‘Seifert hypersurface’’ for Σ_i , namely an oriented hypersurface such that $\partial M_i = \Sigma_i$.

Proof of Proposition 3.1. We consider first the case of a connected surface $\Sigma = \partial M$ with multiplicity one, that is $m = d_1 = 1$ in (1.1). We parameterize the normal bundles of M and Σ in the following compatible way.

Let $\delta > 0$ be a small positive number. First, we take $\Phi_M : M \times [-\delta, \delta] \rightarrow \mathbb{R}^n$ defined by

$$\Phi_M(y, t) := y + tn_M(y),$$

where $n_M \in C^1(M; \mathbb{S}^{n-1})$ is normal to M and such that $(n_M(y), v_1(y), \dots, v_{n-1}(y))$ is a positive basis of \mathbb{R}^n whenever $(v_1(y), \dots, v_{n-1}(y))$ is a positive basis of $T_y M$.

Now we construct a parameterization $\Phi_\Sigma : \Sigma \times [-\delta, \delta]^2 \rightarrow \mathbb{R}^n$ of a tubular neighborhood for Σ in the following way. We start by parameterizing a collar neighborhood of Σ inside M with the map $\Phi_\Sigma : \Sigma \times [0, \delta] \times \{0\} \rightarrow M$ defined as

$$\Phi_\Sigma(\sigma, p_1, 0) := \text{Exp}_\sigma^M(p_1 n_\Sigma(\sigma)),$$

where $n_\Sigma(\sigma) \in T_\sigma M$ is the unit normal vector to Σ at σ pointing inward.

We observe that this definition implies that

$$\Phi_\Sigma(\sigma, 0, 0) = \sigma, \quad \text{and} \quad \frac{\partial \Phi_\Sigma}{\partial p_1}(\sigma, 0, 0) = n_\Sigma(\sigma), \quad \forall \sigma \in \Sigma. \quad (3.1)$$

Then we extend it to $\Phi_\Sigma : \Sigma \times [-\delta, \delta]^2 \rightarrow \mathbb{R}^n$ by setting

$$\Phi_\Sigma(\sigma, p_1, p_2) := \begin{cases} \Phi_M(\Phi_\Sigma(\sigma, p_1, 0), p_2) & \text{if } p_1 \geq 0, \\ \sigma + p_1 n_\Sigma(\sigma) + p_2 n_M(\sigma) & \text{if } p_1 \leq 0. \end{cases} \quad (3.2)$$

By computing the partial derivatives, it is easy to check that $\Phi_\Sigma \in C^1(\Sigma \times [-\delta, \delta]^2; \mathbb{R}^n)$. We also observe that $d\Phi_\Sigma(\sigma, 0, 0)$ is an isometry because of (3.1) and

$$\frac{\partial \Phi_\Sigma}{\partial p_2}(\sigma, 0, 0) = n_M(\sigma). \quad (3.3)$$

Therefore there exists a small positive real number $\delta_0 > 0$ such that both the maps

$$\Phi_\Sigma : \Sigma \times [-\delta_0, \delta_0]^2 \rightarrow \Phi_\Sigma(\Sigma \times [-\delta_0, \delta_0]^2) \quad \text{and} \quad \Phi_M : M \times [-\delta_0, \delta_0] \rightarrow \Phi_M(M \times [-\delta_0, \delta_0]) \quad (3.4)$$

are diffeomorphisms of class C^1 , namely they parameterize some tubular neighborhoods of Σ and M in \mathbb{R}^n . We also observe that $\Phi_\Sigma^{-1}(M) = \Sigma \times [0, \delta_0] \times \{0\}$.

Now let us fix a smooth function $\vartheta : [-1, 1] \rightarrow [0, 2\pi]$ such that $\vartheta(t) = 0$ for $t \in [-1, -1/2]$ and $\vartheta(t) = 2\pi$ for $t \in [1/2, 1]$.

Then we fix a smooth function $u_* : [-1, 1]^2 \rightarrow \mathbb{S}^1$ such that

$$u_*(p) = \begin{cases} p/|p| & \text{if } p \in [-1/2, 1/2]^2, \\ (1, 0) & \text{if } p \in [-1, 3/4] \times [-1, 1] \setminus [-3/4, 3/4]^2, \\ (\cos(\vartheta(p_2)), \sin(\vartheta(p_2))) & \text{if } p \in [3/4, 1] \times [-1, 1]. \end{cases}$$

We observe that such a function exists because

$$\deg(u_*, \partial([-1/2, 1/2]^2)) = \deg(u_*, \partial([-3/4, 3/4]^2)),$$

and any two functions with the same degree on the circle are homotopic (and in this case the homotopy can be made smooth).

Now we define a function $u_\Sigma : \mathbb{R}^n \setminus \Sigma \rightarrow \mathbb{S}^1$ as follows.

$$u_\Sigma(x) := \begin{cases} u_*(p/\delta_0) & \text{if } x = \Phi_\Sigma(\sigma, p) \text{ with } (\sigma, p) \in \Sigma \times [-\delta_0, \delta_0]^2, \\ u_*(1, t/\delta_0) & \text{if } x = \Phi_M(y, t) \text{ with } (y, t) \in M \times [-\delta_0, \delta_0] \setminus \Phi_M^{-1}(\Phi_\Sigma(\Sigma \times [-\delta_0, \delta_0]^2)), \\ (1, 0) & \text{if } x \in \mathbb{R}^n \setminus (\Phi_\Sigma(\Sigma \times [-\delta_0, \delta_0]^2) \cup \Phi_M(M \times [-\delta_0, \delta_0])) \end{cases} \quad (3.5)$$

We point out that the definitions of the parameterizations Φ_M , Φ_Σ , and of the function u_* ensure that the junction in $\Phi_\Sigma(\Sigma \times \{\delta_0\} \times [-\delta_0, \delta_0])$ is of class C^1 .

We also observe that the function u_Σ defined in this way is identically equal to $(1, 0)$ near the boundary of $\Phi_\Sigma(\Sigma \times [-\delta_0, \delta_0]^2) \cup \Phi_M(M \times [-\delta_0, \delta_0])$, hence also the junction at this boundary is smooth.

Hence we have defined a function $u_\Sigma \in C^1(\mathbb{R}^n \setminus \Sigma; \mathbb{S}^1)$ that coincides with the planar vortex on each fiber of a small tubular neighborhood of Σ , and is constant outside a neighborhood of the compact submanifold M .

We claim that the function u_Σ that we have just defined actually links with Σ .

Let $\gamma \in L(\Sigma)$. Up to a homotopy, we can assume that $\gamma \subset \mathbb{R}^n \setminus \Phi_\Sigma(\Sigma \times [-\delta_0, \delta_0]^2)$, and that γ intersects M transversally, and more precisely that $\gamma \cap \Phi_M(M \times [-\delta_0, \delta_0])$ consists of finitely many segments of type $\Phi_M(\{y\} \times [-\delta_0, \delta_0])$, for some $y \in M$. Since the restriction of u_Σ to γ is constant outside these segments, if we fix any parameterization (or any orientation) of γ we obtain that

$$\deg(u_\Sigma, \gamma) = \sum_{y \in M \cap \gamma} s(y) = (-1)^{n-2} \text{link}(\gamma, \Sigma) \quad (3.6)$$

where $s(y) \in \{\pm 1\}$ depends on whether the segment $\Phi_M(\{y\} \times [-\delta_0, \delta_0])$ is crossed from $\Phi_M(y, -\delta_0)$ to $\Phi_M(y, \delta_0)$, or vice versa.

It remains to prove that $\mathcal{E}_\alpha(u_\Sigma, \mathbb{R}^n) < +\infty$ for any $\alpha \in (0, 1)$. This is a consequence of the fact that the function $p/|p| \in H_{\text{loc}}^\alpha(\mathbb{R}^2; \mathbb{S}^1)$ for any $\alpha \in (0, 1)$. However, we will perform this computation in detail in the proof of Proposition 1.3.

Now, we treat the general case. Let Σ be as in (1.1), then there exists $\delta_0 > 0$ such that for any $i \in \{1, \dots, m\}$ the maps Φ_{Σ_i} and Φ_{M_i} in (3.4) are diffeomorphism of class C^1 . In addition, we can assume that the sets $\Phi_{\Sigma_i}(\Sigma_i \times [-\delta_0, \delta_0]^2)$ are pairwise disjoint. For any $i \in \{1, \dots, m\}$, we consider the function u_{Σ_i} defined by (3.5), then we claim that the function

$$u_\Sigma(x) := u_{\Sigma_1}^{d_1}(x) \cdot \dots \cdot u_{\Sigma_m}^{d_m}(x), \quad \forall x \in \mathbb{R}^n \setminus \Sigma, \quad (3.7)$$

has the desired properties, where the product has to be understood in the sense of complex numbers identifying \mathbb{S}^1 as the unit circle in \mathbb{C} .

From the preceding discussion we know that for any $i \in \{1, \dots, m\}$ we have $u_{\Sigma_i} \in C^1(\mathbb{R}^n \setminus \Sigma_i; \mathbb{S}^1)$, therefore $u_\Sigma \in C^1(\mathbb{R}^n \setminus \Sigma; \mathbb{S}^1)$. We prove that u_Σ links with Σ . It is well-known that the

degree is additive with respect to complex multiplication, hence combining this fact with (3.6) and (3.7), we deduce that for any $\gamma \in L(\Sigma)$ we have

$$\deg(u_\Sigma, \gamma) = d_1 \deg(u_{\Sigma_1}, \gamma) + \cdots + d_m \deg(u_{\Sigma_m}, \gamma) = (-1)^{n-2} (\text{link}(\gamma, \Sigma_1) + \cdots + \text{link}(\gamma, \Sigma_m)), \quad (3.8)$$

which in particular implies that u_Σ links with Σ .

At this point, iterating the elementary inequality $|ab - cd| \leq |a - c| + |b - d|$ which is valid for any $a, b, c, d \in \mathbb{S}^1 \subset \mathbb{C}$, we derive that for any $\alpha \in (0, 1)$ it holds that

$$\mathcal{E}_\alpha(u_\Sigma, \mathbb{R}^n) \leq d_1 \mathcal{E}_\alpha(u_{\Sigma_1}, \mathbb{R}^n) + \cdots + d_m \mathcal{E}_\alpha(u_{\Sigma_m}, \mathbb{R}^n), \quad (3.9)$$

and since each addendum on the right-hand side is finite this completes the proof. \square

3.2 Existence of a minimizer

We can now show that there always exists a minimizer in the problem defining the fractional mass (1.5).

Lemma 3.3. *Let $\Sigma \subset \mathbb{R}^n$ be as in (1.1). Then the minimum in (1.5) is achieved by a map $u_\circ \in \mathfrak{F}_s^w(\Sigma)$.*

Proof. Let $m_0 \geq 0$ be the infimum in (1.5) and $\{u_k\} \subset \mathfrak{F}_s^w(\Sigma)$ be a minimizing sequence, that is $\mathcal{E}_{\frac{1+s}{2}}(u_k, \mathbb{R}^n) \downarrow m_0$. The fact that $m_0 < +\infty$ is clear by considering the explicit competitor given in Proposition 3.1.

In particular $\mathcal{E}_{\frac{1+s}{2}}(u_k, \mathbb{R}^n) \leq C$ for some $C > 0$ independent of k . Hence by Remark 2.7 there exists a function $u_\circ \in H_{\text{loc}}^{\frac{1+s}{2}}(\mathbb{R}^n; \mathbb{S}^1)$ such that $u_k \rightarrow u_\circ$ in $\mathbb{X}_s(\mathbb{R}^n)$. Moreover, there holds

$$\mathcal{E}_{\frac{1+s}{2}}(u_\circ, B_R) \leq \liminf_{k \rightarrow \infty} \mathcal{E}_{\frac{1+s}{2}}(u_k, B_R),$$

for every $R \geq 1$. Therefore

$$\mathcal{E}_{\frac{1+s}{2}}(u_\circ, \mathbb{R}^n) = \sup_{R \geq 1} \mathcal{E}_{\frac{1+s}{2}}(u_\circ, B_R) \leq \sup_{R \geq 1} \liminf_{k \rightarrow \infty} \mathcal{E}_{\frac{1+s}{2}}(u_k, B_R) \leq \liminf_{k \rightarrow \infty} \mathcal{E}_{\frac{1+s}{2}}(u_k, \mathbb{R}^n) = m_0.$$

To show that u_\circ is a minimizer we are left to prove that $u_\circ \in \mathfrak{F}_s^w(\Sigma)$. Since $\{u_k\} \subset \mathfrak{F}_s^w(\Sigma)$ there is $\{u_{k,i}\} \subset \mathfrak{F}_s(\Sigma)$ such that $u_{k,i} \rightarrow u_k$ in $\mathbb{X}_s(\mathbb{R}^n)$ as $i \rightarrow \infty$ (see Remark 2.7). Moreover, by the first part of this proof, $u_k \rightarrow u_\circ$ in $\mathbb{X}_s(\mathbb{R}^n)$. Hence, as $\mathbb{X}_s(\mathbb{R}^n)$ is a complete metric space there exists a diagonal subsequence $\{i(k)\} \subset \mathbb{N}^+$ such that $u_{k,i(k)} \rightarrow u_\circ$ in $\mathbb{X}_s(\mathbb{R}^n)$ as $k \rightarrow \infty$. Since $\mathfrak{F}_s^w(\Sigma)$ is the closure of $\mathfrak{F}_s(\Sigma)$ with respect to the distance $d_{\mathbb{X}_s(\mathbb{R}^n)}$ of $\mathbb{X}_s(\mathbb{R}^n)$, we have that $u_\circ \in \mathfrak{F}_s^w(\Sigma)$. \square

3.3 Linking with squares

In this section we show that all weakly linking maps, and in particular the minimizers for (1.5) whose existence is proved in Lemma 3.3, still satisfy in a weak sense the geometric property that defines the class (1.3).

To this end, for $n \geq 2$, we need to parameterize the space \mathcal{S}_n of planar squares in \mathbb{R}^n , and more precisely of their boundaries. Indeed, we remark that we will call "square" the union of the four segments forming the boundary of a planar square.

Let us first consider the case $n = 2$. In this case we can describe the set of squares in \mathbb{R}^2 as

$$\mathcal{S}_2 := \left\{ Q_{\theta,t}(x) : \theta \in \mathbb{S}^1, x \in \mathbb{R}^2, t > 0 \right\},$$

where $Q_{\theta,t}(x)$ is (the boundary of) the square centered at x , with sides of length $2t$, two of which are parallel to θ .

We observe that actually each square corresponds to four different choices of θ . Anyway, we can exploit this description to endow \mathcal{S}_2 with a measure Q_2 defined by

$$\int_{\mathcal{S}_2} \varphi dQ_2 := \int_{\mathbb{S}^1} d\theta \int_{\mathbb{R}^2} dx \int_0^{+\infty} \varphi(Q_{\theta,t}(x)) dt$$

for every $\varphi \in C_c(\mathcal{S}_2)$.

In the higher dimensional case $n \geq 3$, we also need to fix the plane in which the square lies and, since we do not have a canonical set of directions in each plane, the parameter θ has to be chosen a bit more carefully. So we write

$$\mathcal{S}_n := \left\{ Q_{\theta,t}^{L+h}(x) : L \in \mathcal{G}_2^n, h \in L^\perp, x \in L+h, \theta \in \partial D_1^{L+h}(x), t > 0 \right\},$$

where $D_1^{L+h}(x)$ is the unit disk with center x in the affine 2-plane $L+h$ and $Q_{\theta,t}^{L+h}(x)$ is (the boundary of) the square contained $L+h$, centered at x , with sides of length $2t$, two of which are parallel to $\theta - x$. We endow this space with the measure Q_n defined by

$$\int_{\mathcal{S}_n} \varphi dQ_n := \int_{\mathcal{G}_2^n} dL \int_{L^\perp} d\mathcal{H}^{n-2}(h) \int_{L+h} d\mathcal{H}^2(x) \int_{\partial D_1^{L+h}(x)} d\theta \int_0^\infty \varphi(Q_{\theta,t}^{L+h}(x)) dt$$

for every $\varphi \in C_c(\mathcal{S}_n)$.

In what follows, we say that some property holds for *almost every square* $Q \in S$, for some subset $S \subset \mathcal{S}_n$, meaning that the set of squares $Q \in S$ for which that property fails is negligible with respect to the measure Q_n .

Lemma 3.4. *Let $\Sigma \subset \mathbb{R}^n$ be as in (1.1), let $s \in (0, 1)$ and $u \in \mathfrak{F}_s^w(\Sigma)$. Then u has continuous trace on almost every square $Q \in \mathcal{S}_n$ and*

$$|\deg(u, Q)| = |d_1 \text{link}(Q, \Sigma_1) + \dots + d_m \text{link}(Q, \Sigma_m)|, \quad (3.10)$$

for almost every square $Q \in \mathcal{S}_n \cap L(\Sigma)$.

Proof. Let us fix $s_0 \in (0, s)$. In what follows, let c denote a constant that depends only on n and s_0 , and whose value may vary in each line.

Let us fix $R > 0$. By Corollary 2.14 with $k = 2$, for every $v : B_R \rightarrow \mathbb{R}^2$ we have

$$[v]_{H^{\frac{1+s_0}{2}}(B_R)}^2 \geq c \int_{\mathcal{G}_2^n} dL \int_{L^\perp} [v]_{H^{\frac{1+s_0}{2}}((L+h) \cap B_R)}^2 d\mathcal{H}^{n-2}(h). \quad (3.11)$$

We observe that $(L+h) \cap B_R$ is a two-dimensional disk of radius $\sqrt{R^2 - |h|^2}$ and center h in the plane $L+h$, if $|h| \leq R$, and is empty otherwise.

Hence, for every $L \in \mathcal{G}_2^n$ and for every $h \in L^\perp \cap \{|h| \leq R/4\}$, from (2.7) we deduce that

$$\begin{aligned} [v]_{H^{\frac{1+s_0}{2}}((L+h) \cap B_R)}^2 &\geq \int_{(L+h) \cap B_{R/4}(h)} [u]_{H^{\frac{1+s_0}{2}}(D_{R/4}^{L+h}(x))}^2 d\mathcal{H}^2(x) \\ &\geq \int_{(L+h) \cap B_{R/4}(h)} d\mathcal{H}^2(x) \int_{\partial D_1^{L+h}(x)} d\theta \\ &\int_0^{R/4} \left([u]_{H^{\frac{1+s_0}{2}}(S_{\theta,t}^{L+h}(x))}^2 + [u]_{H^{\frac{1+s_0}{2}}(S_{\theta^\perp,t}^{L+h}(x))}^2 + [u]_{H^{\frac{1+s_0}{2}}(S_{\theta,-t}^{L+h}(x))}^2 + [u]_{H^{\frac{1+s_0}{2}}(S_{\theta^\perp,-t}^{L+h}(x))}^2 \right) dt, \end{aligned}$$

where $S_{\theta,t}^{L+h}(x) := \{x + t(\theta - x)^\perp + \xi(\theta - x) : \xi \in (-\sqrt{R/16 - t^2}, \sqrt{R/16 - t^2})\}$.

Applying Lemma 2.8 on the four sides of $Q_{\theta,t}^{L+h}(x)$, since the sum of the oscillation of u on these sides is larger than or equal to the oscillation of u on $Q_{\theta,t}^{L+h}(x)$, we obtain that

$$[v]_{H^{\frac{1+s_0}{2}}((L+h) \cap B_R)}^2 \geq c \int_{(L+h) \cap B_{R/4}(h)} d\mathcal{H}^2(x) \int_{\partial D_1^{L+h}(x)} d\theta \int_0^{R/8} t^{-s_0} \text{osc}(v, Q_{\theta,t}^{L+h}(x))^2 dt.$$

Hence, from (3.11) we get

$$[v]_{H^{\frac{1+s_0}{2}}(B_R)}^2 \geq c \int_{\mathcal{G}_2^n} dL \int_{L^\perp \cap \{|h| \leq R/4\}} d\mathcal{H}^{n-2}(h) \int_{(L+h) \cap B_{R/4}(h)} d\mathcal{H}^2(x) \int_{\partial D_1^{L+h}(x)} d\theta \int_0^{R/8} t^{-s_0} \text{osc}(v, Q_{\theta,t}^{L+h}(x))^2 dt. \quad (3.12)$$

Since $u \in \mathfrak{F}_s^w(\Sigma)$, there exists a sequence $\{u_k\} \subset \mathfrak{F}_s(\Sigma)$ such that $u_k \rightarrow u$ in $\mathbb{X}_s(\mathbb{R}^n)$. Then, applying (3.12) with $v = u - u_k$, we get

$$0 = \lim_{k \rightarrow +\infty} [u - u_k]_{H^{\frac{1+s_0}{2}}(B_R)}^2 \geq \lim_{k \rightarrow +\infty} c \int_{\mathcal{G}_2^n} dL \int_{L^\perp \cap \{|h| \leq R/4\}} d\mathcal{H}^{n-2}(h) \int_{(L+h) \cap B_{R/4}(h)} d\mathcal{H}^2(x) \int_{\partial D_1^{L+h}(x)} d\theta \int_0^{R/8} t^{-s_0} \text{osc}(u - u_k, Q_{\theta,t}^{L+h}(x))^2 dt.$$

Hence, since the oscillation is bounded by 2, the dominated convergence theorem implies that $u_k \rightarrow u$ uniformly on almost every square $Q_{\theta,t}^{L+h}(x) \in \mathcal{S}_n$ with $|h| < R/4$, $|x - h| < R/4$, and $t < R/8$.

Since all squares in \mathcal{S}_n satisfy these conditions for R large enough, we deduce that $u_k \rightarrow u$ in $L^\infty(Q)$ for almost every square $Q \in \mathcal{S}_n$, and in particular u has continuous trace on almost every $Q \in \mathcal{S}_n$.

Moreover, since $u_k \in \mathfrak{F}_s(\Sigma)$, we know that

$$|\deg(u_k, Q)| = |d_1 \text{link}(Q, \Sigma_1) + \dots + d_m \text{link}(Q, \Sigma_m)|$$

for all $Q \in \mathcal{S}_n \cap L(\Sigma)$. Since the degree is continuous under uniform convergence, this implies that (3.10) holds for almost every $Q \in \mathcal{S}_n \cap L(\Sigma)$. \square

3.4 Positivity of the fractional mass

Building on the previous results about weakly linking functions, in this subsection we prove that the fractional mass is always positive. Actually, we prove a stronger result, namely that the mass is locally positive around any point in Σ .

Before stating our result, we prove a few preliminary estimates in the planar case.

Lemma 3.5. *Let $Q = Q_{\theta,t}(x) \in \mathcal{S}_2$ be a square in \mathbb{R}^2 and let L_1, L_2, L_3, L_4 denote its four sides. Let $u : Q \rightarrow \mathbb{S}^1$ be continuous and $d = |\deg(u, Q)|$. Then, for every $s_0 \in (0, 1)$ there exists a constant $c = c(s_0) > 0$ depending only on s_0 such that for every $s \in (s_0, 1)$ we have*

$$[u]_{H^{\frac{1+s}{2}}(L_1)}^2 + [u]_{H^{\frac{1+s}{2}}(L_2)}^2 + [u]_{H^{\frac{1+s}{2}}(L_3)}^2 + [u]_{H^{\frac{1+s}{2}}(L_4)}^2 \geq c \frac{d}{1-s} t^{-s}$$

Proof. Let us assume that $d > 0$, otherwise the thesis is trivial.

Then we can write Q as the union of d connected curves $\gamma_1, \dots, \gamma_d$, intersecting at most at their endpoints, such that $u : \gamma_i \rightarrow \mathbb{S}^1$ is surjective for every i .

Then we can write

$$\sum_{j=1}^4 [u]_{H^{\frac{1+s}{2}}(L_j)}^2 \geq \sum_{i=1}^d \sum_{j=1}^4 [u]_{H^{\frac{1+s}{2}}(L_j \cap \gamma_i)}^2,$$

hence, by Lemma 2.8, we obtain that

$$\sum_{j=1}^4 [u]_{H^{\frac{1+s}{2}}(L_j)}^2 \geq \frac{C_S^{-1}}{1-s} \sum_{i=1}^d \sum_{j=1}^4 \mathcal{H}^1(L_j \cap \gamma_i)^{-s} \text{osc}(u, L_j \cap \gamma_i)^2 \geq \frac{C_S^{-1} (2t)^{-s}}{1-s} \sum_{i=1}^d \sum_{j=1}^4 \text{osc}(u, L_j \cap \gamma_i)^2,$$

and we conclude by observing that

$$\sum_{i=1}^d \sum_{j=1}^4 \text{osc}(u, L_j \cap \gamma_i)^2 \geq \sum_{i=1}^d \frac{1}{4} \left(\sum_{j=1}^4 \text{osc}(u, L_j \cap \gamma_i) \right)^2 \geq \sum_{i=1}^d \frac{1}{4} \text{osc}(u, \gamma_i)^2 = d,$$

because the surjectivity of u on γ_i implies that $\text{osc}(u, \gamma_i) = 2$ for every $i \in \{1, \dots, d\}$. \square

Lemma 3.6. *In the case $n = 2$, let $D_r(x_0)$ denote the disk of center $x_0 \in \mathbb{R}^2$ and radius $r > 0$. Let $d \in \mathbb{N}_+$, let $s_0 \in (0, 1)$ and $s \in [s_0, 1)$, and let $u \in H^{\frac{1+s}{2}}(D_r(x_0); \mathbb{S}^1)$ be a map with $|\deg(u, Q)| = d$ for almost every square $Q \in \mathcal{S}_2$ that is contained in $D_r(x_0)$ for which x_0 lies inside Q , namely in the bounded connected component of $\mathbb{R}^2 \setminus Q$. Then there exists a constant $c = c(s_0) > 0$ such that*

$$[u]_{H^{\frac{1+s}{2}}(D_r(x_0))}^2 \geq c \frac{d}{(1-s)^2} r^{1-s}.$$

Proof. By scaling and translating, we can assume that $r = 1$ and $x_0 = 0$. Let us fix $x \in D_{1/4}(0)$, and let us set $S_{\theta,t}(x) = \{x + t\theta^\perp + \xi\theta : \xi \in (-\sqrt{1/4 - t^2}, \sqrt{1/4 - t^2})\}$, where θ^\perp is the counter-clockwise rotation of θ by an angle equal to $\pi/2$.

By the slicing formula (2.7), we obtain that

$$\begin{aligned} [u]_{H^{\frac{1+s}{2}}(D_{1/2}(x))}^2 &= \\ \frac{1}{4} \int_{\mathbb{S}^1} d\theta \int_0^{1/2} &\left([u]_{H^{\frac{1+s}{2}}(S_{\theta,t}(x))}^2 + [u]_{H^{\frac{1+s}{2}}(S_{\theta,-t}(x))}^2 + [u]_{H^{\frac{1+s}{2}}(S_{\theta^\perp,t}(x))}^2 + [u]_{H^{\frac{1+s}{2}}(S_{\theta^\perp,-t}(x))}^2 \right) dt. \end{aligned}$$

We observe that when $t < \sqrt{2}/4$ the four segments $S_{\theta,t}(x)$, $S_{\theta,-t}(x)$, $S_{\theta^\perp,t}(x)$ and $S_{\theta^\perp,-t}(x)$ appearing in the right-hand side contain the four sides of the square $Q_{\theta,t}(x)$.

By assumption we know that for almost every $x \in D_{1/4}(0)$, for almost every $\theta \in \mathbb{S}^1$ and for almost every $t \in (|x|, \sqrt{2}/4)$, we have that $|\deg(u, Q_{\theta,t}(x))| = d$. Therefore, from Lemma 3.5 we deduce that for almost every $x \in D_{1/4}(0)$ it holds that

$$[u]_{H^{\frac{1+s}{2}}(D_{1/2}(x))}^2 \geq \frac{1}{4} \int_{\mathbb{S}^1} d\theta \int_{|x|}^{\sqrt{2}/4} \frac{cd}{1-s} t^{-s} dt = \frac{cd}{(1-s)^2} (2^{-\frac{3}{2}(1-s)} - |x|^{1-s}),$$

for some constants $c_1, c_2 > 0$ depending only on s_0 .

Now let us fix $\lambda \in (0, 1/4)$. Then we have that

$$\begin{aligned} [u]_{H^{\frac{1+s}{2}}(D_1(0))}^2 &= \int_{D_\lambda(0)} [u]_{H^{\frac{1+s}{2}}(D_1(0))}^2 dx \geq \int_{D_\lambda(0)} [u]_{H^{\frac{1+s}{2}}(D_{1/2}(x))}^2 dx \\ &\geq \frac{cd}{(1-s)^2} \int_{D_\lambda} (2^{-\frac{3}{2}(1-s)} - |x|^{1-s}) dx \geq \frac{cd}{(1-s)^2} (2^{-\frac{3}{2}(1-s)} - \lambda^{1-s}). \end{aligned}$$

Letting $\lambda \rightarrow 0^+$, we conclude the proof. \square

Exploiting the slicing formulas proved in subsection 2.3, we can pass from the planar case to our case of general dimension. To this end, let us introduce some notation and prove another preliminary geometric lemma.

We endow \mathcal{G}_2^n with the following metric structure. For $L, L' \in \mathcal{G}_2^n$ we set

$$d_G(L, L') := \|P_L - P_{L'}\| = \sup_{x \in \mathbb{S}^{n-1}} |(P_L - P_{L'})(x)|,$$

where $P_L, P_{L'}$ are the orthogonal projections on L, L' respectively. This metric is invariant under the action of $O(n)$. Moreover, for every $L \in \mathcal{G}_2^n$ and $x \in \mathbb{R}^n$, let us denote by x_{L^\perp} the orthogonal projection of x onto L^\perp , namely $x_{L^\perp} := x - P_L(x)$.

Lemma 3.7. *Let Σ be as in (1.1). Then there exists a real number $r_0 = r_0(\Sigma) > 0$, depending only on Σ , with the following property.*

For every $(r, x) \in (0, r_0) \times \Sigma$, every $L \in \mathcal{G}_2^n$ with $\text{d}_{\mathcal{G}}(L, (T_x \Sigma)^\perp) \leq 1/4$ and every $h \in L^\perp$ with $|h - x_{L^\perp}| \leq r/8$, there exists a unique point $y \in B_{r/4}(x)$ such that

$$(L + h) \cap \Sigma \cap B_r(x) = \{y\},$$

and the intersection between $L + h$ and Σ in $B_r(x)$ is transversal.

Proof. Let us assume by contradiction that there exists Σ as in (1.1) such that there is no $r_0 > 0$ with the required property.

Then we can find a sequence of numbers $r_k \rightarrow 0^+$, a sequence of points $\{x_k\} \subset \Sigma$, a sequence of 2-planes $\{L_k\} \subset \mathcal{G}_2^n$ and a sequence $\{h_k\} \subset \mathbb{R}^n$ with $h_k \in L_k^\perp$ for every $k \in \mathbb{N}$ such that $|h_k - (x_k)_{L_k^\perp}| \leq r_k/8$ and $\text{d}_{\mathcal{G}}(L_k, (T_{x_k} \Sigma)^\perp) \leq 1/4$, and one of the following three options holds

- $(L_k + h_k) \cap \Sigma \cap B_{r_k/4}(x_k)$ is empty,
- $(L_k + h_k) \cap \Sigma \cap B_{r_k}(x_k)$ contains (at least) two distinct points,
- $L_k + h_k$ is not transversal to Σ in $B_{r_k}(x_k)$.

Up to subsequences (not relabelled), we can also assume that all the points x_k belong to the same connected component Σ_i and that, as $k \rightarrow +\infty$, it holds that

$$\frac{h_k - (x_k)_{L_k^\perp}}{r_k} \rightarrow h_*,$$

for some $h_* \in \mathbb{R}^n$ with $|h_*| \leq 1/8$, that $x_k \rightarrow x_*$ for some $x_* \in \Sigma_i$, and that $L_k \rightarrow L_*$ for some $L_* \in \mathcal{G}_2^n$ with $\text{d}_{\mathcal{G}}(L_*, (T_{x_*} \Sigma_i)^\perp) \leq 1/4$.

We observe that

$$\frac{L_k + h_k - x_k}{r_k} \cap B_1(0) = \left(L_k + \frac{h_k - (x_k)_{L_k^\perp}}{r_k} \right) \cap B_1(0),$$

is a sequence of disks converging to $(L_* + h_*) \cap B_1(0)$, while, taking into account the C^2 regularity of Σ_i , we deduce that

$$\frac{\Sigma - x_k}{r_k} \cap B_1(0)$$

is a sequence of C^2 graphs over $T_{x_k} \Sigma_i$, and hence eventually also over $T_{x_*} \Sigma_i$, converging in C^2 to the disk $T_{x_*} \Sigma_i \cap B_1(0)$ (that is the graph of the null function).

Now we observe that for every $(n-2)$ -plane $T \in \mathcal{G}_{n-2}^n$, every 2-plane $L \in \mathcal{G}_2^n$ with $\text{d}_{\mathcal{G}}(L, T^\perp) \leq 1/4$ and every $h \in \mathbb{R}^n$ with $|h| \leq 1/8$, it holds that $L + h$ intersects T transversally in a single point z , and $|z| < 1/4$. Indeed, since $z \in T \cap (L + h)$, we have that

$$|z| = |z - P_{T^\perp}(z)| \leq |z - P_L(z)| + |P_L(z) - P_{T^\perp}(z)| \leq 1/8 + |z|/4,$$

from which we deduce that $|z| \leq 1/6 < 1/4$.

Therefore, we deduce that eventually

$$\frac{L_k + h_k - x_k}{r_k} \cap \frac{\Sigma - x_k}{r_k} \cap B_1(0)$$

contains exactly one point z_k , at which $(L_k + h_k - x_k)/r_k$ and $(\Sigma - x_k)/r_k$ intersect transversally, and, moreover, $|z_k| < 1/4$.

As a consequence, when k is large enough, $(L_k + h_k) \cap \Sigma \cap B_{r_k}(x_k)$ contains exactly one point $y_k = x_k + r_k z_k$, which also belongs to $B_{r_k/4}(x_k)$ and at which $L_k + h_k$ and Σ are transversal, so none of the previous three options can be true. This concludes the proof. \square

We are finally ready to prove that the fractional mass is locally positive around Σ . The precise statement is the following.

Proposition 3.8. *Let $\Sigma \subset \mathbb{R}^n$ be as in (1.1), let $s_0 \in (0, 1)$, let $s \in [s_0, 1)$ and let $u \in \mathfrak{F}_s^w(\Sigma)$. Then there exists a radius $r_0 = r_0(\Sigma) > 0$ and a constant $c = c(n, s_0) > 0$ such that, for every $i \in \{1, \dots, m\}$, every $x \in \Sigma_i$ and $r \leq r_0$ there holds*

$$[u]_{H^{\frac{1+s}{2}}(B_r(x))}^2 \geq \frac{c}{(1-s)^2} d_i r^{n-1-s}.$$

Proof. By Corollary 2.14 with $k = 2$ and $\alpha_0 = 1/2$ there exists a constant $c(n) > 0$, depending only on n such that for every $r > 0$ it holds that

$$[u]_{H^{\frac{1+s}{2}}(B_r(x))}^2 \geq c(n) \int_{\mathcal{G}_2^n} dL \int_{L^\perp} [u]_{H^{\frac{1+s}{2}}((L+h) \cap B_r(x))}^2 d\mathcal{H}^{n-2}(h). \quad (3.13)$$

Let r_0 be as in Lemma 3.7. Then, for every (r, x, L, h) as in the Lemma 3.7, with $x \in \Sigma_i$, we know that $(L+h) \cap B_r(x)$ is a two-dimensional disk of radius $\sqrt{r^2 - |h - x_{L^\perp}|^2} \geq r\sqrt{1 - 1/64} > r/2$ intersecting Σ transversally in a single point $y = y(r, x, L, h) \in B_{r/4}(x) \cap \Sigma_i$.

In particular, a two-dimensional disk $D_{r/4}^{L+h}(y)$ in the plane $L+h$, with radius equal to $r/4$ and center at y , is contained in $(L+h) \cap B_r(x)$. Using Definition 2.2, we see that all the squares $Q_{\theta,t}^{L+h}(y') \in \mathcal{S}_n$ that are contained in $D_{r/4}^{L+h}(y)$ and for which y lies inside $Q_{\theta,t}^{L+h}(y')$ in the plane $L+h$, satisfy $|\text{link}(Q_{\theta,t}^{L+h}(y'), \Sigma_i)| = 1$ and $\text{link}(Q_{\theta,t}^{L+h}(y'), \Sigma_j) = 0$ for $j \neq i$.

Therefore, since $u \in \mathfrak{F}_s^w(\Sigma)$, from Lemma 3.4 we deduce that $|\deg(u, Q_{\theta,t}^{L+h}(y'))| = d_i$ for almost every square $Q_{\theta,t}^{L+h}(y') \in \mathcal{S}_n$ as above. Hence, for every $(r, x) \in (0, r_0) \times \Sigma$, for almost every $L \in \mathcal{G}_2^n$ and almost every $h \in L^\perp$ as in Lemma 3.7, the disk $D_{r/4}^{L+h}(y)$ fits into the framework of Lemma 3.6, which implies that

$$[u]_{H^{\frac{1+s}{2}}((L+h) \cap B_r(x))}^2 \geq [u]_{H^{\frac{1+s}{2}}(D_{r/4}^{L+h}(y))}^2 \geq \frac{c(s_0)}{(1-s)^2} d_i r^{1-s}$$

for every $s \in (0, s_0)$.

Then, we can continue from (3.13) to get

$$\begin{aligned} [u]_{H^{\frac{1+s}{2}}(B_r(x))}^2 &\geq c(n) \int_{\mathcal{G}_2^n \cap \{d_{\mathcal{G}}(L, (T_x \Sigma_i)^\perp) \leq 1/4\}} dL \int_{L^\perp \cap \{|h - x_{L^\perp}| \leq r/8\}} \frac{c(s_0)}{(1-s)^2} d_i r^{1-s} d\mathcal{H}^{n-2}(h) \\ &\geq \frac{c(n)c(s_0)}{(1-s)^2} d_i r^{1-s} \cdot \text{Vol}_{\mathcal{G}_2^n}(\{d_{\mathcal{G}}(L, (T_x \Sigma_i)^\perp) \leq 1/4\}) \cdot \mathcal{H}^{n-2}(B_{r/8}(x_{L^\perp}) \cap L^\perp) \\ &= \frac{c(n, s_0)}{(1-s)^2} d_i r^{n-1-s} \end{aligned}$$

where $c(n, s_0)$ is a positive constant depending only on n and s_0 . □

4 Proof of the main result

Let $\Phi_i := \Phi_{\Sigma_i}: \Sigma \times [-\delta_0, \delta_0]^2 \rightarrow \mathbb{R}^n$ be the map defined by (3.2), with $\Sigma = \Sigma_i$. For any $\delta \leq \delta_0$, we define the open set $\mathcal{U}_{i,\delta} := \Phi(\Sigma_i \times (-\delta, \delta)^2)$. Of course, we can assume that δ_0 is sufficiently small to ensure that both (3.4) holds and $\mathcal{U}_{i,\delta} \cap \mathcal{U}_{j,\delta} = \emptyset$ for every $i \neq j$ and every $\delta \leq \delta_0$. Moreover, we will use the symbol $\Psi_i = (\Psi_{i,1}, \Psi_{i,2})$ to denote the inverse map of Φ_i , which is defined on $\overline{\mathcal{U}_{i,\delta_0}}$.

As we already observed, $d\Phi_i(\sigma, 0)$ is an isometry for any $\sigma \in \Sigma_i$, see (3.1) and (3.3), therefore for any $\eta > 0$, there exists $\delta(\eta) \in (0, \delta_0)$ such that for any $W \subset \Sigma_i$ with $\text{diam}(W) \leq \delta(\eta)$ it holds

$$(1 - \eta)^2 (d_{\Sigma_i}(\sigma, \tau)^2 + |p - q|^2) \leq |\Phi_i(\sigma, p) - \Phi_i(\tau, q)|^2 \leq (1 + \eta)^2 (d_{\Sigma_i}(\sigma, \tau)^2 + |p - q|^2), \quad (4.1)$$

for any $\sigma, \tau \in W$, for any $p, q \in (-\delta(\eta), \delta(\eta))^2$ and . From the previous estimates we deduce that

$$(1 - \eta)^n \leq |\det d\Phi_i(\sigma, p)| \leq (1 + \eta)^n, \quad \forall (\sigma, p) \in \Sigma_i \times (-\delta(\eta), \delta(\eta))^2. \quad (4.2)$$

Now, for any $\eta > 0$ and for any $\delta > 0$, there exist $N = N(\eta, \delta)$ and a finite disjoint collection $\mathcal{W}_i = \{W_{i,1}, \dots, W_{i,N}\}$ of Lipschitz open subsets of Σ_i with the following properties:

- a) for any $i = 1, \dots, N$, it holds $\text{diam}(W_i) \leq \delta$, and $\bigcup_{i=1}^N \overline{W}_i = \Sigma_i$,
- b) for any $j = 1, \dots, N$, there exists a Lipschitz (open) domain $A_{i,j} \subset \mathbb{R}^{n-2}$ and a bi-Lipschitz diffeomorphism $\varphi_{i,j}: A_{i,j} \rightarrow W_{i,j}$ such that

$$(1 - \eta)|z - w| \leq d_{\Sigma_i}(\varphi_{i,j}(z), \varphi_{i,j}(w)) \leq (1 + \eta)|z - w|, \quad \forall z, w \in A_{i,j}, \quad (4.3)$$

and, as a consequence,

$$(1 - \eta)^{n-2} \leq |\det d\varphi_{i,j}(z)| \leq (1 + \eta)^{n-2}, \quad \forall z \in A_{i,j}. \quad (4.4)$$

For the existence of such a collection, see for example [LMN18, Subsection 4.2].

4.1 Lower bound (proof of Proposition 1.2)

This section is devoted to the proof of Proposition 1.2, which provides the liminf inequality for Theorem 1.1. The main ingredient for the proof is the following flat version of the estimate, for maps weakly linking with a $(n - 2)$ -plane in a bounded region.

Proposition 4.1. *Let $n \geq 3$ be an integer. Let $E \subset \mathbb{R}^{n-2}$ be a bounded open set with Lipschitz boundary, and let $\delta > 0$. Let us set*

$$F := E \times (-\delta, \delta)^2 \quad \text{and} \quad F^* := E \times ((-\delta, \delta)^2 \setminus \{(0, 0)\}).$$

For every $\ell \in (0, \delta)$ and every $\sigma \in E$ we consider the closed curves

$$\gamma_{\sigma, \ell} := \{\sigma\} \times \partial([-\ell, \ell]^2) \subset F^*.$$

For every $s \in (0, 1)$ let $M_s \subset F$ be a $(n - 1)$ -rectifiable set, and let $w_s \in H^{\frac{1+s}{2}}(F; \mathbb{S}^1)$ be a function.

Let also $d \in \mathbb{N}_+$ be a positive integer, and let us assume that for every $s \in (0, 1)$ there exists a sequence of functions $\{w_{s,k}\} \subset C^1(F^*; \mathbb{S}^1) \cap H^{\frac{1+s}{2}}(F; \mathbb{S}^1)$ such that $w_{s,k} \rightarrow w_s$ in $\mathbb{X}_s(F)$ and $|\deg(w_{s,k}, \gamma)| = d$ for every $k \in \mathbb{N}$ and for every curve $\gamma \subset F^* \setminus M_s$ that is homotopic to some $\gamma_{\sigma, \ell}$ in F^* . Let us assume also that $\mathcal{H}^{n-1}(M_s) \rightarrow 0$ as $s \rightarrow 1^-$. Then it turns out that

$$\liminf_{s \rightarrow 1^-} (1 - s)^2 [w_s]_{H^{\frac{1+s}{2}}(F)}^2 \geq \frac{2\pi\omega_{n-1}}{n} d \cdot \mathcal{H}^{n-2}(E). \quad (4.5)$$

Let us first show that Proposition 1.2 follows from Proposition 4.1.

Proof of Proposition 1.2. For every $i \in \{1, \dots, m\}$ let us set $\Omega_i := \Omega \cap \mathcal{U}_{i, \delta_0}$. Then we have that

$$[u]_{H^{\frac{1+s}{2}}(\Omega)} \geq \sum_{i=1}^m [u]_{H^{\frac{1+s}{2}}(\Omega_i)},$$

so (1.6) follows if we prove that

$$\liminf_{s \rightarrow 1^-} [u]_{H^{\frac{1+s}{2}}(\Omega_i)} \geq \frac{2\pi\omega_{n-1}}{n} d_i \mathcal{H}^{n-2}(\Sigma_i \cap \Omega), \quad (4.6)$$

for every $i \in \{1, \dots, m\}$.

For every $s \in (0, 1)$ let $\{u_{s,k}\} \subset \mathfrak{F}_s(\Sigma_s)$ be a sequence converging to u_s in $\mathbb{X}_s(\mathbb{R}^n)$ as $k \rightarrow +\infty$.

Let us fix $i \in \{1, \dots, m\}$ and $\eta > 0$ and let $W_{i,1}, \dots, W_{i,N}$ be as above, with $\delta(\eta)$ as in (4.1). Let us fix $\delta \in (0, \delta(\eta))$, and for every $j \in \{1, \dots, N\}$ let us consider the sets

$$W_{i,j}^\delta := \{x \in W_{i,j} : \Phi_i(\{x\} \times (-\delta, \delta)^2) \subset \Omega_i\}$$

It is clear that

$$[u_s]_{H^{\frac{1+s}{2}}(\Omega_i)}^2 \geq \sum_{j=1}^N [u_s]_{H^{\frac{1+s}{2}}(\Phi_i(W_{i,j}^\delta \times (-\delta, \delta)^2))}^2, \quad (4.7)$$

while, from (4.1), (4.2), (4.3) and (4.4), we deduce that

$$\begin{aligned} [u_s]_{H^{\frac{1+s}{2}}(\Phi(W_{i,j}^\delta \times (-\delta, \delta)^2))}^2 &\geq \frac{(1-\eta)^{2n}}{(1+\eta)^{n+1+s}} \iint_{(W_{i,j}^\delta \times (-\delta, \delta)^2)^2} \frac{|u_s(\Phi_i(\sigma, p)) - u_s(\Phi_i(\tau, q))|^2}{(d_\Sigma(\sigma, \tau)^2 + |p - q|^2)^{\frac{n+1+s}{2}}} \\ &\geq \frac{(1-\eta)^{2n+2(n-2)}}{(1+\eta)^{2(n+1+s)}} [\tilde{u}_{s,i,j}]_{H^{\frac{1+s}{2}}(A_{i,j}^\delta \times (-\delta, \delta)^2)}^2, \end{aligned} \quad (4.8)$$

where $\tilde{u}_{s,i,j} = u_s \circ \Phi_i \circ (\varphi_{i,j} \times \text{Id})$, and $A_{i,j}^\delta := \varphi_{i,j}^{-1}(W_{i,j}^\delta)$.

We also observe that (4.4) yields

$$\mathcal{H}^{n-2}(A_{i,j}^\delta) \geq \frac{1}{(1+\eta)^{n-2}} \mathcal{H}^{n-2}(W_{i,j}^\delta) \quad \forall j \in \{1, \dots, N\}, \quad (4.9)$$

and that

$$\lim_{\delta \rightarrow 0^+} \sum_{j=1}^N \mathcal{H}^{n-2}(W_{i,j}^\delta) = \sum_{j=1}^N \lim_{\delta \rightarrow 0^+} \mathcal{H}^{n-2}(W_{i,j}^\delta) = \sum_{j=1}^N \mathcal{H}^{n-2}(W_{i,j} \cap \Omega_i) = \mathcal{H}^{n-2}(\Sigma_i \cap \Omega). \quad (4.10)$$

Now, by Remark 2.16, for every $s \in (0, 1)$ we can find an integral Lipschitz chain T_s such that $\partial T_s = \Sigma_s - \Sigma$ in the sense of integral currents and

$$\mathbb{M}(T_s) \leq 2\mathbf{F}(\Sigma_s - \Sigma).$$

Let N_s be the support of T_s , which is a $(n-1)$ -rectifiable set, because it is the Lipschitz image of a finite union of $(n-1)$ -dimensional polyhedra. Moreover, we have that

$$\mathcal{H}^{n-1}(N_s) \leq \mathbb{M}(T_s) \leq 2\mathbf{F}(\Sigma_s - \Sigma),$$

and hence $\mathcal{H}^{n-1}(N_s) \rightarrow 0$ as $s \rightarrow 1^-$.

Let us set $N'_s := N_s \cup \Sigma \cup \Sigma_s$. Since Σ and Σ_s are $(n-2)$ -dimensional, we have that N'_s is still $(n-1)$ -rectifiable and also that $\mathcal{H}^{n-1}(N'_s) \rightarrow 0$ as $s \rightarrow 1^-$.

We claim that for every closed curve $\gamma \subset \mathbb{R}^n \setminus N'_s$ it turns out that

$$d_1 \text{link}(\gamma, \Sigma_1) + \dots + d_m \text{link}(\gamma, \Sigma_m) = d_{s,1} \text{link}(\gamma, \Sigma_{s,1}) + \dots + d_{s,m_s} \text{link}(\gamma, \Sigma_{s,m_s}).$$

Indeed we know that $\Sigma_s = \Sigma + \partial T_s$ as currents and the support of T_s is $N_s \subset \mathbb{R}^n \setminus \gamma$, so $[\Sigma] = [\Sigma_s]$ in $H_{n-2}(\mathbb{R}^n \setminus \gamma)$, because the homology of integral currents in $\mathbb{R}^n \setminus \gamma$ is isomorphic to the singular homology (see [FF60, Theorem 5.11]). Therefore our claim follows using Definition 2.3 to compute the linking numbers, because

$$[\gamma]^*(d_1[\Sigma_1] + \dots + d_m[\Sigma_m]) = [\gamma]^*([\Sigma]) = [\gamma]^*([\Sigma_s]) = [\gamma]^*(d_{s,1}[\Sigma_{s,1}] + \dots + d_{s,m_s}[\Sigma_{s,m_s}]).$$

As a consequence, for every $i \in \{1, \dots, m\}$ and every $j \in \{1, \dots, N\}$ we are in the framework of Proposition 4.1 with

$$E := A_{i,j}^\delta, \quad w_s = \tilde{u}_{s,i,j}, \quad w_{s,k} = u_{s,k} \circ \Phi_i \circ (\varphi_{i,j} \times \text{Id}), \quad M_s := (\Phi_i \circ (\varphi_{i,j} \times \text{Id}))^{-1}(N'_s), \quad d = d_i.$$

Indeed, let us check that the assumptions are satisfied.

First of all, we observe that $\mathcal{H}^{n-1}(M_s) \rightarrow 0$ because $\mathcal{H}^{n-1}(N'_s) \rightarrow 0$, and that $w_{s,k} \rightarrow w_s$ in $\mathbb{X}_s(A_{i,j}^\delta \times (-\delta, \delta)^2)$, because $u_{s,k} \rightarrow u_s$ in $\mathbb{X}_s(\mathbb{R}^n)$ as $k \rightarrow +\infty$.

Moreover, if $\gamma \subset (A_{i,j}^\delta \times ((-\delta, \delta)^2 \setminus \{(0, 0)\})) \setminus M_s$ is a curve that is homotopic to some $\gamma_{\sigma,\ell}$ in $A_{i,j}^\delta \times ((-\delta, \delta)^2 \setminus \{(0, 0)\})$, then we have that $\hat{\gamma} := \Phi_i \circ (\varphi_{i,j} \times \text{Id})(\gamma) \subset \mathbb{R}^n \setminus N'_s$ satisfies

$$\begin{aligned} |d_{s,1} \text{link}(\hat{\gamma}, \Sigma_{s,1}) + \cdots + d_{s,m_s} \text{link}(\hat{\gamma}, \Sigma_{s,m_s})| &= |d_1 \text{link}(\hat{\gamma}, \Sigma_1) + \cdots + d_m \text{link}(\hat{\gamma}, \Sigma_m)| \\ &= |d_1 \text{link}(\Phi_i \circ (\varphi_{i,j} \times \text{Id})(\gamma_{\sigma,\ell}), \Sigma_1) + \cdots + d_m \text{link}(\Phi_i \circ (\varphi_{i,j} \times \text{Id})(\gamma_{\sigma,\ell}), \Sigma_m)| = d_i. \end{aligned}$$

Here the first equality follows from our previous claim, the second one from the homotopy invariance of the linking number, and the third one from Definition 2.2, because we know that $\Phi_i \circ (\varphi_{i,j} \times \text{Id})(\{\sigma\} \times [-\ell, \ell]^2)$ intersects Σ_i transversally in the point $\Phi_i \circ (\varphi_{i,j} \times \text{Id})(\sigma, 0)$, and does not intersect $\Sigma_{i'}$ for any $i' \neq i$.

Therefore, since $u_{s,k} \in \mathfrak{F}_s(\Sigma_s)$, we deduce that for every such curve γ it holds that

$$|\deg(w_{s,k}, \gamma)| = |\deg(u_{s,k}, \hat{\gamma})| = d_i,$$

and hence all the assumptions of Proposition 4.1 hold true.

Thus, (4.5) yields

$$\liminf_{s \rightarrow 1^-} (1-s)^2 [\tilde{u}_{s,i,j}]_{H^{\frac{1+s}{2}}(A_{i,j}^\delta \times (-\delta, \delta)^2)}^2 \geq \frac{2\pi\omega_{n-1}}{n} d_i \mathcal{H}^{n-2}(A_{i,j}^\delta).$$

Combining this estimate with (4.7), (4.8) and (4.9) we obtain that

$$\liminf_{s \rightarrow 1^-} (1-s)^2 [u_s]_{H^{\frac{1+s}{2}}(\Omega_i)}^2 \geq \frac{(1-\eta)^{4n-4}}{(1+\eta)^{3n+2}} \sum_{j=1}^N \frac{2\pi\omega_{n-1}}{n} d_i \mathcal{H}^{n-2}(W_{i,j}^\delta).$$

Letting $\delta \rightarrow 0^+$ and $\eta \rightarrow 0^+$, and recalling (4.10), we obtain (4.6). \square

We now focus on the proof of Proposition 4.1, which is quite involved and requires several preliminary results. The first step consists in writing our fractional energy in terms of the discrete functionals studied in [AC09], as it was done in [Sol24] in a similar context.

To this end, let us introduce some notation. Following [Sol24, Section 4] we consider the set V_n of all orthonormal bases of \mathbb{R}^n , namely the set of $\nu \in (\mathbb{S}^{n-1})^n$ such that $\nu_i \cdot \nu_j = 0$ for every $i \neq j$. This set can be identified with $O(n)$ (which is a Lie group of dimension $n(n-1)/2$) by associating to every basis $\nu \in V_n$ the rotation $R_\nu \in O(n)$ bringing the canonical basis into ν .

Therefore it is natural to endow V_n with the measure $\mathcal{H}^{n(n-1)/2}$. For every $\theta \in \mathbb{S}^{n-1}$ and every $i \in \{1, \dots, n\}$ we also consider the set $V_n^{i,\theta}$ of all $\nu \in V_n$ with $\nu_i = \theta$. This set can be identified with $O(n-1)$, and its volume is equal to (see [Sol24, equation (4.3)])

$$\mathcal{H}^{(n-1)(n-2)/2}(V_n^{i,\theta}) = \frac{\mathcal{H}^{n(n-1)/2}(V_n)}{\omega_{n-1}}.$$

As a consequence, for every non-negative measurable function $f : \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ we have the following formula

$$\int_{\mathbb{S}^{n-1}} f(\theta) d\theta = \frac{\omega_{n-1}}{n} \int_{\mathbb{S}^{n-1}} \left(\sum_{i=1}^n \int_{V_n^{i,\theta}} f(\theta) d\nu \right) d\theta = \frac{\omega_{n-1}}{n} \int_{V_n} \sum_{i=1}^n f(\nu_i) d\nu. \quad (4.11)$$

Now let us denote by $\mathcal{Q} := (0, 1)^n$ the unit cube in \mathbb{R}^n . For an open set $\Omega \subset \mathbb{R}^n$ and a triple $(\varepsilon, \nu, z) \in (0, 1) \times V_n \times \mathcal{Q}$, we consider the lattice

$$\mathbb{Z}_{\varepsilon, \nu, z}^n(\Omega) := R_\nu(\varepsilon \mathbb{Z}^n + \varepsilon z) \cap \Omega,$$

the set of cubes into which this lattice divides the domain

$$\mathcal{C}_{\varepsilon,\nu,z}(\Omega) := \{j + R_\nu(\varepsilon\mathcal{Q}) : j \in \mathbb{Z}_{r_s,\nu,z}^n(\Omega) \text{ and } j + R_\nu(\varepsilon\mathcal{Q}) \subset \Omega\},$$

and the set of all their one-dimensional edges $\mathcal{E}_{\varepsilon,\nu,z}(\Omega)$.

For every function $w : \mathbb{Z}_{\varepsilon,\nu,z}^n(\Omega) \rightarrow \mathbb{S}^1$ we consider the following discrete energy

$$E_{\varepsilon,\nu,z}(w, \Omega) := \frac{\varepsilon^{n-2}}{2|\log \varepsilon|} \sum_{\substack{j,k \in \mathbb{Z}_{\varepsilon,\nu,z}^n(\Omega) \\ |j-k|=\varepsilon}} |w(j) - w(k)|^2.$$

For every $r \in (0, 1)$ and every $s \in (0, 1)$, we set $r_s := r^{\frac{2}{1-s}}$, and for every $\eta \in (0, \min\{1, \delta/4\})$ we fix $s_0(\eta) \in (0, 1)$ in such a way that

$$(3\sqrt{n} + 1)r_s \leq \eta \quad \forall (r, s) \in (0, 1 - \eta) \times (s_0(\eta), 1). \quad (4.12)$$

Finally, for every small real number $\varepsilon > 0$, we set

$$E_\varepsilon := \{x \in E : \text{dist}(x, \mathbb{R}^{n-2} \setminus E) > \varepsilon\}, \quad \text{and} \quad F_\varepsilon := \{x \in F : \text{dist}(x, \mathbb{R}^n \setminus F) > \varepsilon\}.$$

The next lemma shows that we can estimate the fractional energy of a function $w : F \rightarrow \mathbb{S}^1$ in terms of an average of the discrete functional computed on the restriction of w to some lattice, as the step, the orientation and the position of the lattice vary.

Lemma 4.2. *For every $\eta \in (0, 1)$, every $s \in (s_0(\eta), 1)$ and every $w \in H^{\frac{1+s}{2}}(F; \mathbb{S}^1)$ it turns out that*

$$(1-s)^2 [w]_{H^{\frac{1+s}{2}}(F)}^2 \geq \frac{\omega_{n-1}}{n} \int_0^{1-\eta} dr \, 8r |\log r| \int_{V_n} d\nu \int_{\mathcal{Q}} E_{r_s,\nu,z}(w^{r_s,\nu,z}, F_{\eta-\sqrt{n}r_s}) dz, \quad (4.13)$$

where for every $(\varepsilon, \nu, z) \in (0, 1) \times V_n \times \mathcal{Q}$ the function $w^{\varepsilon,\nu,z} : \mathbb{Z}_{\varepsilon,\nu,z}^n(F) \rightarrow \mathbb{S}^1$ is the restriction of w to $\mathbb{Z}_{\varepsilon,\nu,z}^n(F)$.

Proof. With the change of variable $y = x + r_s\theta$, we obtain that

$$(1-s)^2 [w]_{H^{\frac{1+s}{2}}(F)}^2 \geq \int_0^{1-\eta} 2(1-s)r \, dr \int_{\mathbb{S}^{n-1}} d\theta \int_{F_{\eta-3\sqrt{n}r_s}} \frac{|w(x+r_s\theta) - w(x)|^2}{r_s^2} dx,$$

where the inequality follows just from the fact that we are reducing the domain of integration, because of (4.12). Observing that $1-s = 2|\log r|/|\log r_s|$ and exploiting (4.11), we can rewrite the right-hand side as

$$\frac{\omega_{n-1}}{n} \int_0^{1-\eta} 8r |\log r| \, dr \int_{V_n} d\nu \frac{1}{2|\log r_s|} \int_{F_{\eta-3\sqrt{n}r_s}} \sum_{i=1}^n \frac{|w(x+r_s\nu_i) - w(x)|^2}{r_s^2} dx.$$

Now we observe that

$$\begin{aligned} & \frac{1}{2|\log r_s|} \int_{F_{\eta-3\sqrt{n}r_s}} \sum_{i=1}^n \frac{|w(x+r_s\nu_i) - w(x)|^2}{r_s^2} dx \\ & \geq \frac{1}{2|\log r_s|} \sum_{j \in \mathbb{Z}_{r_s,\nu,0}^n(F_{\eta-2\sqrt{n}r_s})} \int_{j+r_sR_\nu(\mathcal{Q})} \sum_{i=1}^n \frac{|w(x+r_s\nu_i) - w(x)|^2}{r_s^2} dx \\ & = \frac{r_s^{n-2}}{2|\log r_s|} \sum_{j \in \mathbb{Z}_{r_s,\nu,0}^n(F_{\eta-2\sqrt{n}r_s})} \int_{\mathcal{Q}} \sum_{i=1}^n |w(j+r_sR_\nu(z)+r_s\nu_i) - w(j+R_\nu(z))|^2 dz \\ & \geq \frac{r_s^{n-2}}{2|\log r_s|} \int_{\mathcal{Q}} \sum_{j \in \mathbb{Z}_{r_s,\nu,z}^n(F_{\eta-\sqrt{n}r_s})} \sum_{i=1}^n |w(j+r_s\nu_i) - w(j)|^2 dz \\ & \geq \int_{\mathcal{Q}} E_{r_s,\nu,z}(w^{r_s,\nu,z}, F_{\eta-\sqrt{n}r_s}) dz. \end{aligned}$$

Putting everything together we obtain (4.13). \square

At this point, following [Sol24], one might try to exploit the compactness statement and the Γ -liminf estimate of [AC09, Theorem 3] to deduce that for almost every (r, ν, z) , up to subsequences, it holds that $w_s^{r, \nu, z} \rightarrow \Sigma^{r, \nu, z}$ (in the sense used in [AC09]) for some $(n-2)$ -dimensional integral current $\Sigma^{r, \nu, z}$ and

$$\liminf_{s \rightarrow 1^-} E_{r_s, \nu, z}(w_s^{r_s, \nu, z}, F_{\eta - \sqrt{n}r_s}) \geq \pi \mathbb{M}(\Sigma^{r, \nu, z}).$$

At this point, one might be tempted to infer the assumption on the degree implies that the rectifiable set supporting the current $\Sigma^{r, \nu, z}$ must contain the set $E_\eta \times \{(0, 0)\}$, and hence $\mathbb{M}(\Sigma^{r, \nu, z}) \geq \mathcal{H}^{n-2}(E_\eta)$.

Unfortunately, this is not simple, and actually it is not even true that in general $\mathbb{M}(\Sigma^{r, \nu, z})$ is comparable with $\mathcal{H}^{n-2}(E)$ for almost every choice of the parameters. Indeed, in our case, the argument that we have just outlined does not work for two reasons.

- The first one is that the notion of convergence used in [AC09, Sol24] involves the Jacobian of some piecewise affine interpolations of the discrete functions, and in general it is not true that these interpolations still satisfy some condition on the degree. Actually, the interpolations take values in the unit disk, so they could also vanish, in which case the degree might not even be defined, or it could also happen that w_s is constant on some of the lattices $\mathbb{Z}_{r_s, \nu, z}^n(F)$, in which case the interpolation would be a constant function, and its degree on any curve would be equal to zero.
- The second issue is that, even if one knows that a sequence of maps satisfies our assumption on the degree, it is not immediate to deduce that any limit point (in the sense of currents) of their Jacobians has positive mass in F , because the convergence allows cancellations, and some mass could also concentrate on the boundary.

We overcome the first issue by exploiting the one-dimensional fractional Sobolev embedding of Lemma 2.8 on the one-dimensional edges in $\mathcal{E}_{r_s, \nu, z}^n(F)$, in order to find sufficiently many of these edges on which we can control the oscillation of w_s , and hence the difference between w_s and its interpolations.

As for the second one, we first pass from the discrete energy to the Ginzburg-Landau energy of the interpolations, as in [AC09], then we exploit an estimate of [Jer99, ABO05] that relates the Ginzburg-Landau energy of a function defined in a two-dimensional square directly to its degree on the boundary, instead of the mass of the weak limits, up to a boundary remainder term.

In the end, for sufficiently many values of the parameters (r, ν, z) we manage to find sufficiently many points $\sigma \in E$ such that there exists $\ell_\sigma \in (0, \delta)$ for which we can apply the estimate of [ABO05] on $\{\sigma\} \times [-\ell_\sigma, \ell_\sigma]^2$ and also control the remainder term.

In order to make this reasoning precise, the first step consists in slicing the fractional norm along the lattices $\mathbb{Z}_{\varepsilon, \nu, z}^n(\Omega)$, and applying Lemma 2.8 in each edge $I \in \mathcal{E}_{\varepsilon, \nu, z}(\Omega)$.

Lemma 4.3. *For every $s_0 \in (0, 1)$ there exists a positive constant $c_{n, s_0} > 0$ such that for every open set $\Omega \subset \mathbb{R}^n$, every $s \in (s_0, 1)$, every function $w \in H^{\frac{1+s}{2}}(\Omega)$ and every $\varepsilon > 0$ it holds that*

$$\int_{V_n} d\nu \int_{\mathcal{Q}} dz \sum_{I \in \mathcal{E}_{\varepsilon, \nu, z}(\Omega)} \varepsilon^{n-1-s} \text{osc}(w, I)^2 \leq c_{n, s_0} (1-s) [w]_{H^{\frac{1+s}{2}}(\Omega)}^2$$

Proof. From the usual slicing formula (2.6) we obtain that

$$[w]_{H^{\frac{1+s}{2}}(\Omega)}^2 = \frac{1}{2} \int_{\mathbb{S}^{n-1}} d\theta \int_{\theta^\perp} [w]_{H^{\frac{1+s}{2}}(\Omega_{\theta, z'})}^2 dz',$$

where $\Omega_{\theta, z'}$ denotes the intersection of the line $z' + \mathbb{R}\theta$ with Ω . Then from (4.11) we deduce that

$$[w]_{H^{\frac{1+s}{2}}(\Omega)}^2 = \frac{\omega_{n-1}}{2n} \int_{V_n} d\nu \sum_{i=1}^n \int_{\nu_i^\perp} [w]_{H^{\frac{1+s}{2}}(\Omega_{\nu_i, z'})}^2 dz'.$$

Now let $\{e_1, \dots, e_n\}$ be the canonical basis of \mathbb{R}^n , so $\nu_i = R_\nu(e_i)$ for every $\nu \in V_n$ and every $i \in \{1, \dots, n\}$. Then we can rewrite the previous equality as

$$\begin{aligned}
[w]_{H^{\frac{1+s}{2}}(\Omega)}^2 &= \frac{\omega_{n-1}}{2n} \int_{V_n} d\nu \sum_{i=1}^n \int_{R_\nu(e_i^\perp)} [w]_{H^{\frac{1+s}{2}}(\Omega_{\nu_i, z'})}^2 dz' \\
&= \frac{\omega_{n-1}}{2n} \int_{V_n} d\nu \sum_{i=1}^n \int_{e_i^\perp} [w]_{H^{\frac{1+s}{2}}(\Omega_{\nu_i, R_\nu(z')})}^2 dz' \\
&= \frac{\omega_{n-1}}{2n} \int_{V_n} d\nu \sum_{i=1}^n \frac{1}{\varepsilon} \int_{e_i^\perp \times [0, \varepsilon]} [w]_{H^{\frac{1+s}{2}}(\Omega_{\nu_i, R_\nu(z)})}^2 dz \\
&= \frac{\omega_{n-1}}{2n} \int_{V_n} d\nu \sum_{i=1}^n \sum_{j \in \mathbb{Z}^n \cap e_i^\perp} \frac{1}{\varepsilon} \int_{\varepsilon Q} [w]_{H^{\frac{1+s}{2}}(\Omega_{\nu_i, R_\nu(j+z)})}^2 dz \\
&= \frac{\omega_{n-1}}{2n} \int_{V_n} d\nu \int_Q dz \sum_{i=1}^n \sum_{j \in \mathbb{Z}^n \cap e_i^\perp} \varepsilon^{n-1} [w]_{H^{\frac{1+s}{2}}(\Omega_{\nu_i, R_\nu(j+\varepsilon z)})}^2. \tag{4.14}
\end{aligned}$$

Since in the last expression for every $(\nu, z) \in V_n \times Q$ we are summing the fractional semi-norms on a collection of lines that contains all the edges of the cubes in $\mathcal{C}_{\varepsilon, \nu, z}(\Omega)$, and the fractional norm of a function on an interval is larger than the sum of the fractional norms on any family of disjoint subintervals, we finally obtain that

$$[w]_{H^{\frac{1+s}{2}}(\Omega)}^2 \geq \frac{\omega_{n-1}}{2n} \int_{V_n} d\nu \int_Q dz \sum_{I \in \mathcal{E}_{\varepsilon, \nu, z}(\Omega)} \varepsilon^{n-1} [w]_{H^{\frac{1+s}{2}}(I)}^2.$$

Since the length of each edge $I \in \mathcal{E}_{\varepsilon, \nu, z}(\Omega)$ is equal to ε , the conclusion follows from Lemma 2.8. \square

A simple consequence of the previous lemma is the following.

Lemma 4.4. *Let $s \in (0, 1)$, let $\Omega \subset \mathbb{R}^n$ be a bounded open set with Lipschitz boundary, let $w \in H^{\frac{1+s}{2}}(\Omega; \mathbb{S}^1)$ be a function, and let $\{w_k\} \subset C^0(\Omega; \mathbb{S}^1) \cap H^{\frac{1+s}{2}}(\Omega; \mathbb{S}^1)$ be a sequence such that $w_k \rightarrow w$ in $\mathbb{X}_s(\Omega)$.*

Then for every $\varepsilon > 0$ there exists a negligible set $N_\varepsilon \subset V_n \times Q$ such that for every $(\nu, z) \notin N_\varepsilon$ it turns out that w is continuous (namely it agrees \mathcal{H}^1 almost everywhere with a continuous function) on every $I \in \mathcal{E}_{\varepsilon, \nu, z}(\Omega)$ and, moreover,

$$\liminf_{k \rightarrow +\infty} |\deg(w_k, \gamma)| \leq |\deg(w, \gamma)| \leq \limsup_{k \rightarrow +\infty} |\deg(w_k, \gamma)|, \tag{4.15}$$

for every closed curve γ that is contained in the union of all $I \in \mathcal{E}_{\varepsilon, \nu, z}(\Omega)$.

Proof. Let us fix $s_0 \in (0, s)$. By assumption, we know that $w_k \rightarrow w$ strongly in $H^{\frac{1+s_0}{2}}(\Omega; \mathbb{S}^1)$. Hence, from Lemma 4.3 and Fatou's lemma we deduce that

$$\int_{V_n} d\nu \int_Q dz \liminf_{k \rightarrow +\infty} \sum_{I \in \mathcal{E}_{\varepsilon, \nu, z}(\Omega)} \varepsilon^{n-1-s_0} \text{osc}(w - w_k, I)^2 \leq \liminf_{k \rightarrow +\infty} c_n (1 - s_0) [w - w_k]_{H^{\frac{1+s_0}{2}}(\Omega)}^2 = 0.$$

Therefore, there exists a negligible set $N_\varepsilon \subset V_n \times Q$ such that for every $(\nu, z) \notin N_\varepsilon$ there is a subsequence $\{w_{k_h}\}$, which depends on (ν, z) , that converges to w in $L^\infty(I)$ for every edge $I \in \mathcal{E}_{\varepsilon, \nu, z}$.

As a consequence, w is continuous on each of these edges and

$$|\deg(w, \gamma)| = \lim_{h \rightarrow +\infty} |\deg(w_{k_h}, \gamma)|$$

for every closed curve γ contained in the union of all $I \in \mathcal{E}_{\varepsilon, \nu, z}$. This implies (4.15). \square

With the same slicing technique that we used in Lemma 4.3, we also obtain the following estimate, that allows us to control the number of cubes with an edge intersecting M_s .

Lemma 4.5. *There exists a positive constant $c_n > 0$ such that for every open set $\Omega \subset \mathbb{R}^n$, every $(n-1)$ -rectifiable set $M \subset \Omega$ with $\mathcal{H}^{n-1}(M) < +\infty$, and every $\varepsilon > 0$ it turns out that*

$$\int_{V_n} d\nu \int_{\mathcal{Q}} dz \sum_{I \in \mathcal{C}_{\varepsilon, \nu, z}(\Omega)} \varepsilon^{n-1} \mathcal{H}^0(I \cap M) \leq c_n \mathcal{H}^{n-1}(M).$$

Proof. From Theorem 2.15 we obtain that

$$\mathcal{H}^{n-1}(M) = \frac{\beta_{0,n}^{n-1,1}}{\omega_{n-1}} \int_{\mathbb{S}^{n-1}} d\theta \int_{\theta^\perp} \mathcal{H}^0(\Omega_{\theta, z'} \cap M) dz',$$

where $\Omega_{\theta, z'}$ denotes the intersection of the line $z' + \mathbb{R}\theta$ with Ω .

We conclude arguing as in (4.14). □

As we anticipated above, the next step consists in relating the discrete energies

$$E_{r_s, \nu, z}(w_s^{r_s, \nu, z}, F_{\eta - \sqrt{n}r_s})$$

to the Ginzburg-Landau energy of a piecewise affine interpolation of $w_s^{r_s, \nu, z}$. This was done in [AC09], so we just have to introduce some notation to recall the results that we need.

For every $(r, \nu, z) \in (0, \eta) \times V_n \times \mathcal{Q}$ and every $s \in (s_0(\eta), 1)$ we denote by $v_s^{r_s, \nu, z}$ the piecewise affine interpolation of w_s on the simplices of the Kuhn decomposition of each cube in $\mathcal{C}_{r_s, \nu, z}(F)$.

We refer to [AC09] for the detailed construction of this interpolation, and we limit to describe it in words. Basically, the idea is to divide the cube \mathcal{Q} into the $n!$ simplices $T_\pi := \{x_{\pi(1)} < \dots < x_{\pi(n)}\}$, where π is a permutation of $\{1, \dots, n\}$. We observe that the vertices of all these simplices are also vertices of \mathcal{Q} .

Thus, we can divide each of the cubes in $\mathcal{C}_{r_s, \nu, z}(F)$ into $n!$ simplices whose vertices lie in the lattice $\mathbb{Z}_{r_s, \nu, z}^n(F)$. In each of these simplices we take the affine function coinciding with w_s (and hence with $w_s^{r_s, \nu, z}$) on its vertices. What we obtain is a Lipschitz continuous piecewise affine function $v_s^{r_s, \nu, z}$ with values in the unit ball of \mathbb{R}^2 , that coincides with $w_s^{r_s, \nu, z}$ on $\mathbb{Z}_{r_s, \nu, z}^n(F)$. We point out that $v_s^{r_s, \nu, z}$ is not defined on the whole F , but only on a union of cubes contained in F , which anyway contains $F_{\sqrt{n}r_s}$, and hence also $F_{\eta - \sqrt{n}r_s}$, because of (4.12).

From [AC09, equation (4.21)], we deduce that

$$E_{r_s, \nu, z}(w_s^{r_s, \nu, z}, F_{\eta - \sqrt{n}r_s}) \geq \frac{1}{2|\log r_s|} \int_{F_\eta} |\nabla v_s^{r_s, \nu, z}(x)|^2 dx,$$

while from [AC09, Lemma 2] we deduce that there exists a constant $c_0 > 0$ such that

$$E_{r_s, \nu, z}(w_s^{r_s, \nu, z}, F_{\eta - \sqrt{n}r_s}) \geq \frac{c_0}{r_s^2 |\log r_s|} \int_{F_\eta} (1 - |v_s^{r_s, \nu, z}(x)|^2)^2 dx.$$

As a consequence, if for every $(\varepsilon, \beta) \in (0, 1)^2$ we consider the Ginzburg-Landau density

$$e_{\varepsilon, \beta}(v) := \frac{1 - \beta}{2|\log \varepsilon|} |\nabla v|^2 + \frac{\beta c_0}{\varepsilon^2 |\log \varepsilon|} (1 - |v|^2)^2,$$

and, for every open set $\Omega \subset \mathbb{R}^n$, the corresponding energy

$$GL_{\varepsilon, \beta}(v, \Omega) := \int_{\Omega} e_{\varepsilon, \beta}(v) dx,$$

then we obtain that

$$E_{r_s, \nu, z}(w_s^{r_s, \nu, z}, F_{\eta - \sqrt{n}r_s}) \geq GL_{r_s, \beta}(v_s^{r_s, \nu, z}, F_\eta), \quad (4.16)$$

for every $\eta \in (0, 1)$, every $(r, \nu, z) \in (0, 1 - \eta) \times V_n \times \mathcal{Q}$, every $s \in (s_0(\eta), 1)$, and every $\beta \in (0, 1)$.

Now we recall the following result from [ABO05], which is a refinement of a previous estimate in [Jer99].

Lemma 4.6 (Lemma 3.10 in [ABO05]). *There exists a universal constant $K > 0$ such that for every $\beta \in (0, 1)$ and every $\rho > 0$ there exist two constants $D_0, D_1 > 0$, depending only on β and ρ , such that for every $d \in \mathbb{N}_+$ and every smooth function $v : [-\ell, \ell]^2 \rightarrow \mathbb{R}^2$ with $|v| > 1/2$ on $\partial[-\ell, \ell]^2$ and $|\deg(v/|v|, \partial[-\ell, \ell]^2)| = d$ it holds that*

$$GL_{\varepsilon, \beta}(v, [-\ell, \ell]^2) + K\rho\ell \int_{\partial[-\ell, \ell]^2} e_{\varepsilon, \beta}(v) d\mathcal{H}^1 \geq (1 - \beta)\pi d \left(1 - \frac{\log(2\ell) + D_1(1 + \log d)}{|\log \varepsilon|}\right), \quad (4.17)$$

for every $\varepsilon \in (0, 2\ell)$ such that $(\varepsilon/2\ell)|\log(\varepsilon/2\ell)| < D_0/d$.

Remark 4.7. We will need to apply Lemma 4.6 to (the restriction to some planar domain of) the functions $v_s^{r_s, \nu, z}$, which are not smooth, but Lipschitz continuous and piecewise affine. However, if we approximate $v_s^{r_s, \nu, z}$ with smooth functions, every term in (4.17) passes to the limit, provided the boundary of the domain intersects the set where $v_s^{r_s, \nu, z}$ is not smooth (which is a finite union of $(n - 1)$ -dimensional simplices) transversally.

Now we are ready to prove Proposition 4.1.

Proof of Proposition 4.1. First of all, we observe that without loss of generality we can assume that the liminf in (4.5) is finite, and up to extracting a subsequence (that we do not relabel), we can also assume that it is a limit, so that in particular

$$\limsup_{s \rightarrow 1^-} (1 - s)^2 [w_s]_{H^{\frac{1+s}{2}}(F)}^2 < +\infty. \quad (4.18)$$

Let us fix $\eta \in (0, \min\{1, \delta/4\})$ and $s_0 = s_0(\eta)$ such that (4.12) holds.

Let us fix also $r \in (0, 1 - \eta)$ and $s \in (s_0, 1)$, and let $\mathcal{N}_{s,r} \subset V_n \times \mathcal{Q}$ be the negligible set provided by Lemma 4.4 applied with

$$\Omega = F, \quad w = w_s, \quad w_k = w_{s,k}, \quad \varepsilon = r_s.$$

Now we need to select the "good" cubes in $\mathcal{C}_{r_s, \nu, z}(F_{\eta - \sqrt{n}r_s})$, namely those cubes for which we can find a suitable estimate on the oscillation of w_s on each of their one-dimensional edge, and such that M_s does not intersect any of these edges.

To this end, for every $(\nu, z) \in (V_n \times \mathcal{Q}) \setminus \mathcal{N}_{s,r}$, we consider the set

$$\mathcal{O}_{s,r,\nu,z} := \left\{ Q \in \mathcal{C}_{r_s, \nu, z}(F_{\eta - \sqrt{n}r_s}) : \text{osc}(w_s, I) > \frac{\sqrt{2}}{n} \text{ for some edge } I \text{ of } Q \right\},$$

of cubes for which we do not have a good oscillation estimate on all their edges, and the set

$$\mathcal{M}_{s,r,\nu,z} := \left\{ Q \in \mathcal{C}_{r_s, \nu, z}(F_{\eta - \sqrt{n}r_s}) : I \cap M_s \neq \emptyset \text{ for some edge } I \text{ of } Q \right\},$$

of cubes with at least one edge intersecting M_s .

Now we fix a small positive real number $\lambda > 0$ and we consider the set

$$\mathcal{B}_{s,r} := \mathcal{N}_{s,r} \cup \left\{ (\nu, z) \in (V_n \times \mathcal{Q}) \setminus \mathcal{N}_{s,r} : \mathcal{H}^0(\mathcal{O}_{s,r,\nu,z} \cup \mathcal{M}_{s,r,\nu,z}) > \frac{2\lambda^2\delta}{r_s^{n-1}} \right\}.$$

Since each edge belongs to at most 2^{n-1} cubes, from Lemma 4.3 we deduce that

$$\begin{aligned} & \frac{\mathcal{H}^{n(n+1)/2} \left(\left\{ (\nu, z) \in (V_n \times \mathcal{Q}) \setminus \mathcal{N}_{s,r} : \mathcal{H}^0(\mathcal{O}_{s,r,\nu,z}) > \frac{\lambda^2\delta}{r_s^{n-1}} \right\} \right)}{\mathcal{H}^{n(n-1)/2}(V_n)} \\ & \leq \frac{r_s^{n-1}}{\lambda^2\delta} \cdot 2^{n-1} \cdot \frac{n^2}{2} \cdot \frac{c_{n,s_0}}{r_s^{n-1-s}} (1-s) [w_s]_{H^{\frac{1+s}{2}}(F)}^2 = \frac{c_{n,s_0} 2^{n-2} n^2}{\lambda^2\delta} r_s^s (1-s) [w_s]_{H^{\frac{1+s}{2}}(F)}^2, \end{aligned}$$

while from Lemma 4.5 we deduce that

$$\frac{\mathcal{H}^{n(n+1)/2} \left(\left\{ (\nu, z) \in (V_n \times \mathcal{Q}) \setminus \mathcal{N}_{s,r} : \mathcal{H}^0(\mathcal{M}_{s,r,\nu,z}) > \frac{\lambda^2 \delta}{r_s^{n-1}} \right\} \right)}{\mathcal{H}^{n(n-1)/2}(V_n)} \leq \frac{c_n 2^{n-1}}{\lambda^2 \delta} \mathcal{H}^{n-1}(M_s).$$

Since $\mathcal{N}_{s,r}$ is negligible, combining these two estimates we obtain that

$$\frac{\mathcal{H}^{n(n+1)/2}(\mathcal{B}_{s,r})}{\mathcal{H}^{n(n-1)/2}(V_n)} \leq \frac{c_{n,s_0} 2^{n-2} n^2}{\lambda^2 \delta} \frac{r_s^s}{(1-s)} (1-s)^2 [w_s]_{H^{\frac{1+s}{2}}(\Omega_{\delta,L})}^2 + \frac{c_n 2^{n-1}}{\lambda^2 \delta} \mathcal{H}^{n-1}(M_s),$$

which in turn, recalling (4.18), yields

$$\lim_{s \rightarrow 1^-} \mathcal{H}^{n(n+1)/2}(\mathcal{B}_{s,r}) = 0 \quad \forall r \in (0, 1 - \eta). \quad (4.19)$$

Let $\pi_1 : \mathbb{R}^2 \times \mathbb{R}^{n-2} \rightarrow \mathbb{R}^2$ and $\pi_2 : \mathbb{R}^2 \times \mathbb{R}^{n-2} \rightarrow \mathbb{R}^{n-2}$ be the orthogonal projections onto the two factors, and let consider the set

$$\mathfrak{S}_{s,r,\nu,z} := \left\{ \sigma \in E_\eta : \mathcal{H}^0(\{Q \in \mathcal{O}_{s,r,\nu,z} \cup \mathcal{M}_{s,r,\nu,z} : \sigma \in \pi_2(Q)\}) > \frac{\lambda \delta}{r_s} \right\},$$

that consists of the points $\sigma \in E_\eta$ for which the number of "bad" cubes intersecting $\{\sigma\} \times (-\delta, \delta)^2$ is not small enough.

We observe that, if χ_A denotes the indicator function of the set A , then

$$\begin{aligned} \frac{\lambda \delta}{r_s} \mathcal{H}^{n-2}(\mathfrak{S}_{s,r,\nu,z}) &\leq \int_{E_\eta} \sum_{Q \in \mathcal{O}_{s,r,\nu,z} \cup \mathcal{M}_{s,r,\nu,z}} \chi_{\pi_2(Q)}(\sigma) d\sigma \\ &= \sum_{Q \in \mathcal{O}_{s,r,\nu,z} \cup \mathcal{M}_{s,r,\nu,z}} \mathcal{H}^{n-2}(\pi_2(Q)) \\ &\leq p_n r_s^{n-2} \mathcal{H}^0(\mathcal{O}_{s,r,\nu,z} \cup \mathcal{M}_{s,r,\nu,z}), \end{aligned}$$

where $p_n := \sup\{\mathcal{H}^{n-2}(\pi_2(R_\nu(Q))) : \nu \in V_n\}$ is a constant depending only on the dimension n .

As a consequence, if $(\nu, z) \in (V_n \times \mathcal{Q}) \setminus \mathcal{B}_{s,r}$, then we have that

$$\mathcal{H}^{n-2}(\mathfrak{S}_{s,r,\nu,z}) \leq 2\lambda p_n. \quad (4.20)$$

Finally, for every $\sigma \in E_\eta$ let us consider the set

$$\mathcal{L}_{s,r,\nu,z}(\sigma) := \{\ell \in (\delta/2, \delta - \eta) : \exists Q \in \mathcal{O}_{s,r,\nu,z} \cup \mathcal{M}_{s,r,\nu,z} \text{ such that } \gamma_{\sigma,\ell} \cap Q \neq \emptyset\},$$

namely the set of $\ell \in (\delta/2, \delta - \eta)$ for which $\gamma_{\sigma,\ell}$ intersects some of the "bad" cubes. We recall that we have chosen $\eta < \delta/4$, so $(\delta/2, \delta - \eta)$ is not empty.

We observe that for every $\sigma \in E_\eta \setminus \mathfrak{S}_{s,r,\nu,z}$ it holds that

$$\mathcal{H}^1(\mathcal{L}_{s,r,\nu,z}(\sigma)) \leq \sum_{Q \in \mathcal{O}_{s,r,\nu,z} \cup \mathcal{M}_{s,r,\nu,z}} \text{diam}(Q \cap (\{\sigma\} \times (-\delta, \delta)^2)) \leq \frac{\lambda \delta}{r_s} \sqrt{n} r_s = \lambda \sqrt{n} \delta. \quad (4.21)$$

As a consequence, if we take λ small enough, and more precisely such that $\lambda \sqrt{n} \delta < \delta/2 - \eta$, then for every $(\nu, z) \in (V_n \times \mathcal{Q}) \setminus \mathcal{B}_{s,r}$ and every $\sigma \in E_\eta \setminus \mathfrak{S}_{s,r,\nu,z}$, we can find many curves $\gamma_{\sigma,\ell}$ that do not intersect any "bad" cube.

For every such curve, namely for every $\gamma_{\sigma,\ell}$ with $\ell \notin \mathcal{L}_{s,r,\nu,z}(\sigma)$, let us denote by $U_{\sigma,\ell}$ the union of the cubes in $\mathcal{O}_{r_s,\nu,z}(F_{\eta-\sqrt{n}r_s})$ intersected by $\gamma_{\sigma,\ell}$ and with $S_{\sigma,\ell}$ the union of their edges.

We claim that for every $\ell \notin \mathcal{L}_{s,r,\nu,z}(\sigma)$, with $(\nu, z) \notin \mathcal{N}_{s,r}$, it holds that

$$|v_s^{r_s,\nu,z}(x)| > 1/2 \quad \forall x \in U_{\sigma,\ell} \quad \text{and} \quad |\deg(v_s^{r_s,\nu,z}/|v_s^{r_s,\nu,z}|, \gamma_{\sigma,\ell})| = 1. \quad (4.22)$$

Indeed, if $Q \notin \mathcal{O}_{s,r,\nu,z}$, then from the estimate on the oscillation of w_s we deduce that $|w_s(x) - w_s(y)| \leq \sqrt{2}/n$ for every couple (x, y) of vertices of Q that are connected by an edge. Since every couple of vertices of a n -dimensional cube can be connected by a path made of at most n edges, by the triangle inequality we deduce that $|w_s(x) - w_s(y)| \leq \sqrt{2}$ for every couple (x, y) of vertices of Q , hence w_s sends the vertices of Q in a quarter of the circle. Since by construction $v_s^{r_s,\nu,z}(Q)$ is contained in the convex hull of the image of the vertices of Q , and on these vertices $v_s^{r_s,\nu,z}$ coincides with w_s , we deduce that actually $|v_s^{r_s,\nu,z}| \geq \sqrt{2}/2 > 1/2$ on each cube $Q \notin \mathcal{O}_{s,r,\nu,z}$, and hence on $U_{\sigma,\ell}$.

Moreover, $\gamma_{\sigma,\ell}$ can be continuously deformed inside $U_{\sigma,\ell}$ (and in particular inside F^*) onto a curve γ contained in $S_{\sigma,\ell}$, and hence not intersecting M_s , because the cubes in $U_{\sigma,\ell}$ do not belong to $\mathcal{M}_{s,r,\nu,z}$.

Therefore our hypothesis implies that $|\deg(w_{k,s}, \gamma)| = 1$ for every $k \in \mathbb{N}$, so (4.15) yields $|\deg(w_s, \gamma)| = 1$.

Finally, we observe that the estimate on the oscillation implies that $v_s^{r_s,\nu,z}/|v_s^{r_s,\nu,z}|$ is homotopic to w_s on each of the edges in $S_{\sigma,\ell}$, because in these edges w_s take values in an arc of length smaller than $\pi/2$ and $v_s^{r_s,\nu,z}$ in its convex hull, which does not contain the origin. Hence $v_s^{r_s,\nu,z}/|v_s^{r_s,\nu,z}|$ is homotopic to w_s on γ , so we deduce that

$$|\deg(v_s^{r_s,\nu,z}/|v_s^{r_s,\nu,z}|, \gamma_{\sigma,\ell})| = |\deg(v_s^{r_s,\nu,z}/|v_s^{r_s,\nu,z}|, \gamma)| = |\deg(w_s, \gamma)| = 1,$$

and this concludes the proof of (4.22).

The last thing we need to do before being able to exploit Lemma 4.6 is to make sure that we can control the boundary term. To this end it is enough to notice that for every fixed $\beta \in (0, 1)$ it holds that

$$\int_0^{\delta-\eta} d\ell \int_{\gamma_{\sigma,\ell}} e_{r_s,\beta}(v_s^{r_s,\nu,z}) d\mathcal{H}^1 = \int_{\{\sigma\} \times [-\delta+\eta, \delta-\eta]^2} e_{r_s,\beta}(v_s^{r_s,\nu,z}) d\mathcal{H}^2,$$

and hence, thanks to (4.21), for almost every $\sigma \in E_\eta \setminus \mathfrak{S}_{s,r,\nu,z}$ there exists $\ell_\sigma \in (\delta/2, \delta - \eta) \setminus \mathcal{L}_{s,r,\nu,z}(\sigma)$ such that

$$\int_{\gamma_{\sigma,\ell_\sigma}} e_{r_s,\beta}(v_s^{r_s,\nu,z}) d\mathcal{H}^1 \leq \frac{1}{\delta/2 - \eta - \lambda\sqrt{n}\delta} \int_{\{\sigma\} \times [-\delta+\eta, \delta-\eta]^2} e_{r_s,\beta}(v_s^{r_s,\nu,z}) d\mathcal{H}^2,$$

and such that both $\{\sigma\} \times [-\delta, \delta]^2$ and $\gamma_{\sigma,\ell_\sigma}$ intersect the singular set of $v_s^{r_s,\nu,z}$ transversally (see Remark 4.7). We point out that for almost every $\sigma \in E_\eta$, this transversality condition may fail only for a negligible set of $\ell \in (0, \delta)$. We also remark that actually ℓ_σ depends also on all the other parameters, namely on $\eta, \beta, s, r, \nu, z$.

Now we fix $\rho > 0$ and $s_1 = s_1(\eta, \delta, \beta, \rho, d) \in [s_0, 1)$ such that

$$r_s < 2\ell \quad \text{and} \quad \frac{r_s}{2\ell} \left| \log \frac{r_s}{2\ell} \right| < \frac{D_0}{d}$$

for every $s \in (s_1, 1)$ and every $\ell \in (\delta/2, \delta)$, where D_0 is the constant in Lemma 4.6. Then from (4.17) we deduce that

$$\begin{aligned} & \left(1 + \frac{K\rho\delta}{\delta/2 - \eta - \lambda\sqrt{n}\delta} \right) \int_{\{\sigma\} \times [-\delta+\eta, \delta-\eta]^2} e_{r_s,\beta}(v_s^{r_s,\nu,z}) d\mathcal{H}^2 \\ & \geq \int_{\{\sigma\} \times [-\ell_\sigma, \ell_\sigma]^2} e_{r_s,\beta}(v_s^{r_s,\nu,z}) d\mathcal{H}^2 + K\rho\ell_\sigma \int_{\gamma_{\sigma,\ell_\sigma}} e_{r_s,\beta}(v_s^{r_s,\nu,z}) d\mathcal{H}^1 \\ & \geq \int_{\{\sigma\} \times [-\ell_\sigma, \ell_\sigma]^2} \tilde{e}_{r_s,\beta}(v_s^{r_s,\nu,z}) d\mathcal{H}^2 + K\rho\ell_\sigma \int_{\gamma_{\sigma,\ell_\sigma}} \tilde{e}_{r_s,\beta}(v_s^{r_s,\nu,z}) d\mathcal{H}^1 \\ & \geq (1 - \beta)\pi d \left(1 - \frac{\log(2\ell_\sigma) + D_1(1 + \log d)}{|\log r_s|} \right), \end{aligned}$$

for every r_s such that $s \in (s_1, 1)$, where $\tilde{e}_{r_s, \beta}(v_s^{r_s, \nu, z})$ denote the Ginzburg-Landau energy density of $v_s^{r_s, \nu, z}$ as a function defined on the two dimensional set $[-\ell_\sigma, \ell_\sigma]^2 \times \{y\}$ or, equivalently, the energy density obtained by replacing the full gradient of $v_s^{r_s, \nu, z}$ with its "horizontal" component.

We are now ready to put all the estimates together and conclude the proof. First, we observe that for every $(r, \nu, z) \in (0, \eta) \times V_n \times \mathcal{Q}$ and every $s \in (s_1, 1)$ it holds that

$$\begin{aligned} GL_{r_s, \beta}(v_s^{r_s, \nu, z}, F_\eta) &\geq \int_{E_\eta \setminus \mathfrak{S}_{s, r, \nu, z}} d\sigma \int_{\{\sigma\} \times [-\eta\delta, \eta\delta]^2} e_{r_s, \beta}(v_s^{r_s, \nu, z}) d\mathcal{H}^2 \\ &\geq \mathcal{H}^{n-2}(E_\eta \setminus \mathfrak{S}_{s, r, \nu, z})(1 - \beta)\pi d \left(1 - \frac{\log(2\delta) + D_1(1 + \log d)}{|\log r_s|}\right) \left(1 + \frac{K\rho\delta}{\delta/2 - \eta - \lambda\sqrt{n}\delta}\right)^{-1}, \end{aligned}$$

and hence, if $(\nu, z) \notin \mathcal{B}_{s, r}$, from (4.20) we deduce that

$$\begin{aligned} GL_{r_s, \beta}(v_s^{r_s, \nu, z}, F_\eta) &\geq (\mathcal{H}^{n-2}(E_\eta) - 2\lambda p_n)(1 - \beta)\pi d \left(1 - \frac{\log(2\delta) + D_1(1 + \log d)}{|\log r_s|}\right) \left(1 + \frac{K\rho\delta}{\delta/2 - \eta - \lambda\sqrt{n}\delta}\right)^{-1}. \end{aligned}$$

Combining this estimate with (4.13), (4.16), Fatou's lemma and (4.19) we obtain that

$$\begin{aligned} \liminf_{s \rightarrow 1^-} (1 - s)^2 [w_s]_{H^{\frac{1+s}{2}}(F)}^2 &\geq \frac{\omega_{n-1}}{n} \int_0^{1-\eta} 8r |\log r| dr (\mathcal{H}^{n-2}(E_\eta) - 2\lambda p_n)(1 - \beta)\pi d \left(1 + \frac{K\rho\delta}{\delta/2 - \eta - \lambda\sqrt{n}\delta}\right)^{-1} \end{aligned}$$

Letting $\rho \rightarrow 0^+$, $\beta \rightarrow 0^+$, $\lambda \rightarrow 0^+$, and $\eta \rightarrow 0^+$, and computing the integral

$$\int_0^1 8r |\log r| dr = 2,$$

we obtain exactly (4.5). □

4.2 Upper bound (proof of Proposition 1.3)

This section is devoted to the proof of Proposition 1.3, which provides the limsup inequality for Theorem 1.1.

As for the lower bound, we first need a flat version of the estimate, namely the computation of the energy of the vortex around a $(n - 2)$ -plane. More precisely, we consider the vortex-type map $u_\star: \mathbb{R}^{n-2} \times (\mathbb{R}^2 \setminus \{0\}) \rightarrow \mathbb{S}^1$ defined by

$$u_\star(\sigma, p) := \frac{p}{|p|} \quad \forall (\sigma, p) \in \mathbb{R}^{n-2} \times (\mathbb{R}^2 \setminus \{0\}).$$

In the following lemma we compute the asymptotic limit of the rescaled fractional semi-norms of u_\star on bounded domains. This computation will be crucial to treat the general case.

Lemma 4.8. *Let $F \subset \mathbb{R}^n$ be a bounded open set such that $\mathcal{H}^{n-2}(\partial F \cap (\mathbb{R}^{n-2} \times \{0\})) = 0$. Then,*

$$\lim_{s \rightarrow 1^-} (1 - s)^2 [u_\star]_{H^{\frac{1+s}{2}}(F)}^2 = \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(F \cap (\mathbb{R}^{n-2} \times \{0\})), \quad (4.23)$$

and

$$\lim_{s \rightarrow 1^-} (1 - s)^2 \iint_{F \times F^c} \frac{|u_\star(x) - u_\star(y)|^2}{|x - y|^{n+1+s}} dx dy = 0. \quad (4.24)$$

Proof. For any $x = (\sigma, p) \in \mathbb{R}^{n-2} \times \mathbb{R}^2$ and for any $\delta > 0$ we define the quantities

$$\mathbf{I}_{s,\delta}(x) := \int_{B_\delta} \frac{|u_\star(x+z) - u_\star(x)|^2}{|z|^{n+1+s}} dz, \quad \mathbf{II}_{s,\delta}(x) := \int_{B_\delta^c} \frac{|u_\star(x+z) - u_\star(x)|^2}{|z|^{n+1+s}} dz, \quad (4.25)$$

and we claim that, if $p \neq 0$, then for any $\delta < |p|$, we have

$$\mathbf{I}_{s,\delta}(x) \leq \frac{\omega_{n-1}}{n(1-s)} \cdot \frac{\delta^{1-s}}{(|p| - \delta)^2}, \quad (4.26)$$

$$\mathbf{I}_{s,\delta}(x) \geq \frac{\omega_{n-1}}{n(1-s)} \cdot \frac{\delta^{1-s}}{(|p| + \delta)^2} - \frac{3\omega_{n-1}}{8} \cdot \frac{\delta^{3-s}}{(|p| - \delta)^4}, \quad (4.27)$$

and

$$\mathbf{II}_{s,\delta}(x) \leq 4\omega_{n-1}\delta^{-(1+s)}. \quad (4.28)$$

The estimate (4.28) is an easy consequence of the fact that u_\star takes value in the circle and hence it is bounded. We turn our attention to the bounds (4.26) and (4.27). We preliminarily observe that $\mathbf{I}_{s,\delta}(x)$ depends only on $|p|$, where $x = (\sigma, p)$. Indeed, let $\lambda = |p|$ and let $A \in O(2)$ such that $p = A(\lambda e_1)$, where $e_1 = (1, 0) \in \mathbb{R}^2$, we define $B \in O(n)$ by the formula $B(\tau, q) = (\tau, A(q))$.

Combining the elementary identity $|a - b|^2 = 2(1 - a \cdot b)$, which is valid for any $a, b \in S^1$, with the change of variable $z = B(\tau, q)$, we obtain

$$\begin{aligned} \mathbf{I}_{s,\delta}(x) &= 2 \int_{B_\delta} \frac{1 - u_\star(B(\sigma + \tau, \lambda e_1 + q)) \cdot u_\star(B(\sigma, \lambda e_1))}{(|\tau|^2 + |q|^2)^{\frac{n+1+s}{2}}} d\tau dq \\ &= 2 \int_{B_\delta} \left(1 - \frac{A(\lambda e_1 + q)}{|A(\lambda e_1 + q)|} \cdot \frac{A(\lambda e_1)}{|A(\lambda e_1)|} \right) \frac{d\tau dq}{(|\tau|^2 + |q|^2)^{\frac{n+1+s}{2}}} \\ &= 2 \int_{B_\delta} \left(1 - \frac{(\lambda e_1 + q) \cdot e_1}{|\lambda e_1 + q|} \right) \frac{d\tau dq}{(|\tau|^2 + |q|^2)^{\frac{n+1+s}{2}}} \\ &= 2 \int_{B_\delta} \left(1 - \frac{\lambda + q_1}{\sqrt{(\lambda + q_1)^2 + q_2^2}} \right) \frac{d\tau dq}{(|\tau|^2 + |q|^2)^{\frac{n+1+s}{2}}}, \end{aligned} \quad (4.29)$$

and the last expression depends only on $\lambda = |p|$. From (4.29) and the elementary inequality

$$\frac{1}{2}t^2 - \frac{3}{8}t^4 \leq 1 - \frac{1}{\sqrt{1+t^2}} \leq \frac{1}{2}t^2, \quad \forall t \in \mathbb{R},$$

applied with $t = q_2/(\lambda + q_1)$, we find that

$$\mathbf{I}_{s,\delta}(x) \leq \int_{B_\delta} \frac{q_2^2}{(\lambda + q_1)^2(|\tau|^2 + |q|^2)^{\frac{n+1+s}{2}}} d\tau dq, \quad (4.30)$$

and

$$\mathbf{I}_{s,\delta}(x) \geq \int_{B_\delta} \frac{q_2^2}{(\lambda + q_1)^2(|\tau|^2 + |q|^2)^{\frac{n+1+s}{2}}} d\tau dq - \frac{3}{4} \int_{B_\delta} \frac{q_2^4}{(\lambda + q_1)^4(|\tau|^2 + |q|^2)^{\frac{n+1+s}{2}}} d\tau dq. \quad (4.31)$$

Finally, the estimates (4.26) and (4.27) follow easily from (4.30) and (4.31) after noticing that $\lambda - \delta \leq \lambda + q_1 \leq \lambda + \delta$ for any $(\tau, q) \in B_\delta$ and that the following computations hold

$$\begin{aligned} \int_{B_\delta} \frac{q_2^2}{(|\tau|^2 + |q|^2)^{\frac{n+1+s}{2}}} d\tau dq &= \frac{1}{n} \int_{B_\delta} \frac{d\tau dq}{(|\tau|^2 + |q|^2)^{\frac{n-1+s}{2}}} = \frac{\omega_{n-1}}{n} \frac{\delta^{1-s}}{1-s}, \\ \int_{B_\delta} \frac{q_2^4}{(|\tau|^2 + |q|^2)^{\frac{n+1+s}{2}}} d\tau dq &\leq \int_{B_\delta} \frac{d\tau dq}{(|\tau|^2 + |q|^2)^{\frac{n-3+s}{2}}} = \omega_{n-1} \frac{\delta^{3-s}}{3-s} \leq \frac{1}{2} \omega_{n-1} \delta^{3-s}, \end{aligned}$$

which can be proved using polar coordinates.

We are ready to prove (4.23), we start by proving that for any bounded open set $F \subset \mathbb{R}^n$ it holds

$$\limsup_{s \rightarrow 1^-} (1-s)^2 [u_\star]_{H^{\frac{1+s}{2}}(F)}^2 \leq \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(\overline{F} \cap (\mathbb{R}^{n-2} \times \{0\})). \quad (4.32)$$

Let $A \subset \mathbb{R}^n$ be a Borel set, for any $\sigma \in \mathbb{R}^{n-2}$ we define $A^\sigma := \{p \in \mathbb{R}^2 : (\sigma, p) \in A\}$. Now, for any $s \in (0, 1)$ we introduce the nonnegative function

$$f_s(\sigma) := (1-s)^2 \int_{F^\sigma} \text{I}_{s,\delta}(p) + \text{II}_{s,\delta}(p) dp, \quad (4.33)$$

and we recall that both $\text{I}_{s,\delta}$ and $\text{II}_{s,\delta}$ depend only on $|p|$ as a consequence of (4.29). Moreover, their sum does not depend on δ . We fix $R > 0$ such that $F \subset B_R$. We claim that for every $\theta \in (0, 1)$ we have the estimate

$$f_s(\sigma) \leq \frac{2\pi\omega_{n-1}}{n} \cdot \frac{\theta^{1-s}}{(1-\theta)^2} R^{1-s} + 8\pi\omega_{n-1}(1-s)\theta^{-(1+s)} R^{1-s}, \quad (4.34)$$

in particular the family $\{f_s\}_{s \in (0,1)}$ is bounded in $L^\infty(\mathbb{R}^{n-2})$.

We consider $\delta = \theta|p|$, then from (4.26) and (4.28) we obtain

$$\text{I}_{s,\delta}(p) \leq \frac{\omega_{n-1}}{n(1-s)} \cdot \frac{\theta^{1-s}}{(1-\theta)^2} |p|^{-(1+s)}, \quad \text{and} \quad \text{II}_{s,\delta}(p) \leq 4\omega_{n-1}\theta^{-(1+s)} |p|^{-(1+s)}.$$

Integrating the previous estimates in D_R , we derive

$$(1-s)^2 \int_{D_R} \text{I}_{s,\delta}(p) + \text{II}_{s,\delta}(p) dp \leq \frac{2\pi\omega_{n-1}}{n} \cdot \frac{\theta^{1-s}}{(1-\theta)^2} R^{1-s} + 8\pi\omega_{n-1}(1-s)\theta^{-(1+s)} R^{1-s},$$

and using the fact that $F^\sigma \subset D_R$ for any $\sigma \in \mathbb{R}^{n-2}$ we immediately obtain (4.34).

From (4.33), Fubini's theorem and the definition of the fractional seminorm we have that

$$(1-s)^2 [u_\star]_{H^{\frac{1+s}{2}}(F)}^2 \leq (1-s)^2 \iint_{F \times \mathbb{R}^n} \frac{|u_\star(x) - u_\star(y)|^2}{|x-y|^{n+1+s}} dx dy = \int_{\{|\sigma| \leq R\}} f_s(\sigma) d\sigma, \quad (4.35)$$

where to deduce the last identity we used again that $F \subset B_R$. Therefore, the uniform estimate (4.34) implies that

$$\limsup_{s \rightarrow 1^-} (1-s)^2 \iint_{F \times \mathbb{R}^n} \frac{|u_\star(x) - u_\star(y)|^2}{|x-y|^{n+1+s}} dx dy \leq \int_{\{|\sigma| \leq R\}} \limsup_{s \rightarrow 1^-} f_s(\sigma) d\sigma. \quad (4.36)$$

We distinguish two cases:

i) if $0 \notin \overline{F}^\sigma$, then $\delta_\sigma := \min\{|p|/2 : p \in \overline{F}^\sigma\}$ is positive. Therefore, from (4.26) and (4.28) with $\delta = \delta_\sigma$ we clearly have

$$\lim_{s \rightarrow 1^-} f_s(\sigma) = 0, \quad (4.37)$$

where we used again the fact that $F^\sigma \subset D_R$ for any $\sigma \in \mathbb{R}^{n-2}$.

ii) if $0 \in \overline{F}^\sigma$, letting first $s \rightarrow 1^-$ and then $\theta \rightarrow 0^+$ in (4.34), we obtain

$$\limsup_{s \rightarrow 1^-} f_s(\sigma) \leq \frac{2\pi\omega_{n-1}}{n}. \quad (4.38)$$

Finally, from (4.36), (4.37) and (4.38) we have

$$\begin{aligned} \limsup_{s \rightarrow 1^-} (1-s)^2 \iint_{F \times \mathbb{R}^n} \frac{|u_\star(x) - u_\star(y)|^2}{|x-y|^{n+1+s}} dx dy &\leq \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(\{\sigma : 0 \in \overline{F}^\sigma\}) \\ &\leq \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(\overline{F} \cap (\mathbb{R}^{n-2} \times \{0\})), \end{aligned} \quad (4.39)$$

where to deduce the last inequality we used the fact that if $0 \in \partial F^\sigma$, then $(\sigma, 0) \in \partial F$. Combining (4.35) and (4.39) we get (4.32).

Now we prove that

$$\liminf_{s \rightarrow 1^-} (1-s)^2 [u_\star]_{H^{\frac{1+s}{2}}(F)}^2 \geq \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(F \cap (\mathbb{R}^{n-2} \times \{0\})). \quad (4.40)$$

We observe that if σ has the property that $0 \in F^\sigma$, then $r_\sigma := \text{dist}((\sigma, 0), \partial F)/2$ is positive and for any $|p| \leq r_\sigma$ we have $B_{r_\sigma}((\sigma, p)) \subset F$. In particular, from (4.25), Fubini's theorem and the definition of the fractional seminorm, we have that

$$[u_\star]_{H^{\frac{1+s}{2}}(F)}^2 \geq \int_{\{\sigma: 0 \in F^\sigma\}} d\sigma \int_{D_{r_\sigma}} I_{s,r_\sigma}(p) dp. \quad (4.41)$$

We fix $\theta \in (0, 1)$ and we notice that $I_{s,r_\sigma}(p) \geq I_{s,\theta|p|}(p)$ for any $|p| \leq r_\sigma$. Therefore, from (4.27) we obtain

$$\int_{D_{r_\sigma}} I_{s,r_\sigma}(p) dp \geq \int_{D_{r_\sigma}} I_{s,\theta|p|}(p) dp \geq \frac{2\pi\omega_{n-1}}{n(1-s)^2} \cdot \frac{\theta^{1-s}}{(1+\theta)^2} r_\sigma^{1-s} - \frac{3\pi\omega_{n-1}}{4(1-s)} \cdot \frac{\theta^{3-s}}{(1-\theta)^4} r_\sigma^{1-s}.$$

Moreover, letting first $s \rightarrow 1^-$ and then $\theta \rightarrow 0^+$ in the previous estimate, we deduce that

$$\liminf_{s \rightarrow 1^-} (1-s)^2 \int_{D_{r_\sigma}} I_{s,r_\sigma}(p) dp \geq \frac{2\pi\omega_{n-1}}{n},$$

which implies, together with (4.41) and Fatou's lemma, that

$$\liminf_{s \rightarrow 1^-} (1-s)^2 [u_\star]_{H^{\frac{1+s}{2}}(F)}^2 \geq \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(\{\sigma: 0 \in F^\sigma\}) = \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(F \cap (\mathbb{R}^{n-2} \times \{0\})).$$

This gives (4.40), and together with (4.32) and the assumption $\mathcal{H}^{n-2}(\partial F \cap (\mathbb{R}^{n-2} \times \{0\})) = 0$, it concludes the proof of (4.23).

It remains to prove (4.24). Combining (4.23), (4.39) and the fact that

$$\iint_{F \times F^c} \frac{|u_\star(x) - u_\star(y)|^2}{|x-y|^{n+1+s}} dx dy = \iint_{F \times \mathbb{R}^n} \frac{|u_\star(x) - u_\star(y)|^2}{|x-y|^{n+1+s}} dx dy - [u_\star]_{H^{\frac{1+s}{2}}(F)}^2,$$

we easily obtain (4.24). \square

We are ready to conclude the proof of the upper bound.

Proof of Proposition 1.3. Let u_Σ be defined by (3.7). From the proof of Theorem 3.1, see (3.8), we know that $u_\Sigma \in \mathfrak{F}_s(\Sigma)$ for any $s \in (0, 1)$. This is our candidate to show the validity of (1.7).

In view of (3.9), it is enough to prove the proposition separately for each u_{Σ_i} . Therefore, we can assume without loss of generality that Σ is connected and with multiplicity one. For the rest of the proof we set $\Phi := \Phi_\Sigma$, where Φ_Σ is the parameterization defined by (3.2).

We start with the case $\Omega = \mathbb{R}^n$. From the definition of the fractional seminorm, for any $\delta \leq \delta_0$, we have

$$[u_\Sigma]_{H^{\frac{1+s}{2}}(\mathbb{R}^n)}^2 = [u_\Sigma]_{H^{\frac{1+s}{2}}(\mathcal{U}_\delta)}^2 + [u_\Sigma]_{H^{\frac{1+s}{2}}(\mathcal{U}_\delta^c)}^2 + 2 \iint_{\mathcal{U}_\delta \times \mathcal{U}_\delta^c} \frac{|u_\Sigma(x) - u_\Sigma(y)|^2}{|x-y|^{n+1+s}} dx dy. \quad (4.42)$$

Moreover, if $\delta \leq \delta_0/2$ it holds that $u_\Sigma(\Phi(\sigma, p)) = p/|p|$. We start by proving that the estimate (1.7) holds if we replace the fractional seminorm on the whole \mathbb{R}^n with the fractional seminorm on \mathcal{U}_δ .

Combining the change of variable $(x, y) = (\Phi(\sigma, p), \Phi(\tau, q))$ with the definition of u_Σ we obtain

$$[u_\Sigma]_{H^{\frac{1+s}{2}}(\mathcal{U}_\delta)}^2 = \iint_{\mathcal{V}_\delta \times \mathcal{V}_\delta} \frac{|u_\star(p) - u_\star(q)|^2}{|\Phi(\sigma, p) - \Phi(\tau, q)|^{n+1+s}} J\Phi(\sigma, p) J\Phi(\tau, q) d\sigma d\tau dp dq, \quad (4.43)$$

where $\mathcal{V}_\delta := \Sigma \times (-\delta, \delta)^2$ and $J\Phi(\sigma, p) := |\det d\Phi(\sigma, p)|$ is the Jacobian of Φ evaluated at the point (σ, p) .

We fix $\eta > 0$ and a positive real number $\delta(\eta) \leq \delta_0/2$ for which (4.1) holds. We fix $\delta \leq \delta(\eta)$ and we consider a collection $\mathcal{W} = \{W_i, \dots, W_N\}$ associated to the parameters η and δ such that properties a) and b) stated at the beginning of Section 4 hold. For any $i = 1, \dots, N$ we define the set $\mathcal{V}_{i,\delta} := W_i \times (-\delta, \delta)^2$. From (4.43) and (4.2) we deduce that

$$[u_\Sigma]_{H^{\frac{1+s}{2}}(\mathcal{U}_\delta)}^2 \leq (1 + \eta)^{2n} \sum_{i,j=1}^N \iint_{\mathcal{V}_{i,\delta} \times \mathcal{V}_{j,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{|\Phi(\sigma, p) - \Phi(\tau, q)|^{n+1+s}} d\sigma d\tau dp dq. \quad (4.44)$$

We distinguish three cases:

i) if $i \neq j$ and the distance between W_i and W_j is greater than or equal to a positive number d_0 , then from (4.1) and the fact that u_\star has values in the circle we have

$$\begin{aligned} \iint_{\mathcal{V}_{i,\delta} \times \mathcal{V}_{j,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{|\Phi(\sigma, p) - \Phi(\tau, q)|^{n+1+s}} &\leq \frac{4}{(1 - \eta)^{n+1+s}} \iint_{\mathcal{V}_{i,\delta} \times \mathcal{V}_{j,\delta}} \frac{1}{(d_0^2 + |p - q|^2)^{\frac{n+1+s}{2}}} \\ &\leq \frac{(8\delta^2 \mathcal{H}^{n-2}(\Sigma))^2}{((1 - \eta)d_0)^{n+1+s}}. \end{aligned}$$

In particular, from the previous estimate it follows immediately that

$$\lim_{s \rightarrow 1^-} (1 - s)^2 \iint_{\mathcal{V}_{i,\delta} \times \mathcal{V}_{j,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{|\Phi(\sigma, p) - \Phi(\tau, q)|^{n+1+s}} = 0. \quad (4.45)$$

ii) if $i \neq j$ and W_i and W_j are at zero distance, we can find an open set $U \subset \mathbb{R}^{n-2}$ and a bi-Lipschitz parameterization $\psi: U \rightarrow B$, where B is some ball in Σ which contains both W_i and W_j . It is always possible to find such a map if δ is smaller than the injectivity radius of Σ .

We define the set $G_{i,\delta} := \psi^{-1}(W_i) \times (-\delta, \delta)^2 \subset \mathbb{R}^{n-2} \times \mathbb{R}^2$ and we consider the change of variable $(\sigma, \tau) = (\psi(z), \psi(w))$, then it is clear that there exists $L > 0$, depending on the Lipschitz constants of ψ , Φ^{-1} and ψ^{-1} , such that

$$\iint_{\mathcal{V}_{i,\delta} \times \mathcal{V}_{j,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{|\Phi(\sigma, p) - \Phi(\tau, q)|^{n+1+s}} \leq L \iint_{G_{i,\delta} \times G_{j,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{(|z - w|^2 + |p - q|^2)^{\frac{n+1+s}{2}}}. \quad (4.46)$$

We observe that $G_{j,\delta} \subset G_{i,\delta}^c$, therefore from (4.46) and (4.24) we have

$$\lim_{s \rightarrow 1^-} (1 - s)^2 \iint_{\mathcal{V}_{i,\delta} \times \mathcal{V}_{j,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{|\Phi(\sigma, p) - \Phi(\tau, q)|^{n+1+s}} = 0. \quad (4.47)$$

iii) if $i = j$, we consider the map $\varphi_i \times \text{Id}: A_i \times \mathbb{R}^2 \rightarrow W_i \times \mathbb{R}^2$ and the open set

$$F_{i,\delta} := (\varphi_i \times \text{Id})^{-1}(\mathcal{V}_{i,\delta}) = A_i \times (-\delta, \delta)^2, \quad (4.48)$$

where φ_i is the bi-Lipschitz diffeomorphism appearing in (4.3).

From (4.43) and (4.4) and the change of variables $(\sigma, \tau) = (\varphi_i(z), \varphi_j(w))$, we obtain

$$\iint_{\mathcal{V}_{i,\delta} \times \mathcal{V}_{i,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{|\Phi(\sigma, p) - \Phi(\tau, q)|^{n+1+s}} \leq (1 + \eta)^{2(n-2)} \iint_{F_{i,\delta} \times F_{i,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{|\Phi(\varphi_i(z), p) - \Phi(\varphi_j(w), q)|^{n+1+s}}. \quad (4.49)$$

Moreover, from (4.1) and (4.3) we have

$$\iint_{F_{i,\delta} \times F_{i,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{|\Phi(\varphi_i(z), p) - \Phi(\varphi_i(w), q)|^{n+1+s}} \leq \frac{1}{(1-\eta)^{2(n+1+s)}} \iint_{F_{i,\delta} \times F_{i,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{(|z-w|^2 + |p-q|^2)^{\frac{n+1+s}{2}}}. \quad (4.50)$$

It follows immediately from (4.48) that

$$\Phi((\varphi_i \times \text{Id})(\overline{F}_{i,\delta} \cap \mathbb{R}^{n-2} \times \{0\})) = \overline{W}_i. \quad (4.51)$$

Therefore, combining (4.49), (4.50) and (4.32) we deduce that

$$\begin{aligned} \limsup_{s \rightarrow 1^-} (1-s)^2 \iint_{\mathcal{V}_{i,\delta} \times \mathcal{V}_{i,\delta}} \frac{|u_\star(p) - u_\star(q)|^2}{|\Phi(\sigma, p) - \Phi(\tau, q)|^{n+1+s}} &\leq \frac{(1+\eta)^{2(n-2)}}{(1-\eta)^{2(n+2)}} \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(\overline{F}_{i,\delta} \cap \{p=0\}) \\ &\leq \frac{(1+\eta)^{2(n-2)}}{(1-\eta)^{3n+2}} \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(\overline{W}_i), \end{aligned} \quad (4.52)$$

where we used (4.51) and again the estimates (4.1) and (4.3).

Finally, from (4.44), (4.45), (4.47) and (4.52) we obtain

$$\begin{aligned} \limsup_{s \rightarrow 1^-} (1-s)^2 [u_\Sigma]_{H^{\frac{1+s}{2}}(\mathcal{U}_\delta)}^2 &\leq \frac{(1+\eta)^{4(n-1)}}{(1-\eta)^{3n+2}} \frac{2\pi\omega_{n-1}}{n} \sum_{i=1}^N \mathcal{H}^{n-2}(\overline{W}_i) \\ &= \frac{(1+\eta)^{4(n-1)}}{(1-\eta)^{3n+2}} \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(\Sigma). \end{aligned} \quad (4.53)$$

Now, we deal with the remaining terms in (4.42). We notice that $u_\Sigma \in H_{\text{loc}}^1(\mathcal{U}_\delta^c; \mathbb{S}^1)$ and it is constant outside a large ball, therefore we can use [DNPV12, Remark 4.3], to infer that

$$\limsup_{s \rightarrow 1^-} (1-s)^2 [u_\Sigma]_{H^{\frac{1+s}{2}}(\mathcal{U}_\delta^c)}^2 = 0. \quad (4.54)$$

For what concerns the last integral in (4.42), we observe that

$$\begin{aligned} \iint_{\mathcal{U}_\delta \times \mathcal{U}_\delta^c} \frac{|u_\Sigma(x) - u_\Sigma(y)|^2}{|x-y|^{n+1+s}} dx dy &\leq \iint_{\mathcal{U}_{\delta/2} \times \mathcal{U}_\delta^c} \frac{|u_\Sigma(x) - u_\Sigma(y)|^2}{|x-y|^{n+1+s}} dx dy \\ &\quad + \iint_{\mathcal{U}_{\delta/2}^c \times \mathcal{U}_{\delta/2}^c} \frac{|u_\Sigma(x) - u_\Sigma(y)|^2}{|x-y|^{n+1+s}} dx dy. \end{aligned} \quad (4.55)$$

Since $\mathcal{U}_{\delta/2}$ and \mathcal{U}_δ^c are at positive distance and $|u_\Sigma| = 1$, clearly

$$\lim_{s \rightarrow 1^-} (1-s)^2 \iint_{\mathcal{U}_{\delta/2} \times \mathcal{U}_\delta^c} \frac{|u_\Sigma(x) - u_\Sigma(y)|^2}{|x-y|^{n+1+s}} dx dy = 0. \quad (4.56)$$

Moreover, after multiplying the second term in (4.55) by $(1-s)^2$, it goes to zero for the same reason used in (4.54). Finally, putting together (4.42) and (4.53)-(4.56) we obtain

$$\limsup_{s \rightarrow 1^-} (1-s)^2 [u_\Sigma]_{H^{\frac{1+s}{2}}(\mathbb{R}^n)}^2 \leq \frac{(1+\eta)^{4(n-1)}}{(1-\eta)^{3n+2}} \frac{2\pi\omega_{n-1}}{n} \mathcal{H}^{n-2}(\Sigma).$$

Letting $\eta \rightarrow 0^+$ the conclusion follows.

We are left with the case of a general open set Ω . By definition, we have

$$\mathcal{E}_{\frac{1+s}{2}}(u_\Sigma, \Omega) \leq [u_\Sigma]_{H^{\frac{1+s}{2}}(\mathbb{R}^n)}^2 - [u_\Sigma]_{H^{\frac{1+s}{2}}(\mathbb{R}^n \setminus \overline{\Omega})}^2.$$

Therefore, by combining this simple observation with the estimate on the whole space that we have just proved and the lower bound (1.6), we obtain (1.7). \square

5 Possible generalizations

In order to enlighten the main ideas in our proofs, avoiding as much as possible inessential technicalities, we stated and proved our main result in the case of codimension two submanifolds of the Euclidean space.

However, our method is quite robust, and could be generalized to more general settings. In this section, we briefly describe two possible generalizations, pointing out the what can be extended verbatim, and what requires some carefulness.

5.1 Ambient Riemannian manifolds

The convergence results presented in this work (see Proposition 1.2 and Proposition 1.3) are local. Our main result Theorem 1.1 is stated globally essentially just because we do not know that the minimizer in (1.5) is unique. Then, the localized fractional s -mass could depend on the choice of the minimizer, as explained in Remark 1.4.

Apart from this technical point, our results are completely local. This means that our notion is well-suited to be extended to ambient Riemannian manifolds. The only point that requires care in doing so is that in an ambient n -dimensional Riemannian manifold N , even for a single $(n - 2)$ -dimensional closed, oriented surface (with multiplicity one) $\Sigma \subset N$, the family $\mathfrak{F}_s(\Sigma)$ defined in (1.3) could be empty because of topological obstructions.

Indeed, in our proof of the existence of a competitor in \mathbb{R}^n (that is, Proposition 3.1), we exploit Theorem 3.2, for which it is essential that $H_{n-2}(\mathbb{R}^n) = \{0\}$ to say that every cycle is a boundary. On an ambient Riemannian manifold N with $H_{n-2}(N) \neq \{0\}$, it is not true in general that every such cycle (the connected components of Σ) is a boundary, so a competitor might fail to exist.

For example, take the flat three torus T^3 , which we consider as $[-1, 1]^3$ with suitably identified faces. Then, for the closed curve $\Sigma_\circ = \{(0, 0, t) : t \in [-1, 1]\}$ no such $u \in \mathfrak{F}_s(\Sigma_\circ)$ exists. If such u would exist then some level set of u would be an immersed surface spanning Σ_\circ , and this would imply $[\Sigma_\circ] = 0$ in $H_1(T^3)$, which is not the case.

This phenomenon is due to the fact that Σ_\circ above does not lie in the natural class on which the s -mass is well-defined on ambient Riemannian manifolds. This natural class is the family of $(n - 2)$ -dimensional boundaries with integer coefficients.

Precisely, for $m \geq 1$ positive integer and $d = (d_1, \dots, d_m) \in \mathbb{N}_+^m$ one can consider

$$\Sigma = \partial(d_1 M_1 \cup \dots \cup d_m M_m), \quad (5.1)$$

where $\{M_i\}_{i=1}^m \subset \mathbb{R}^n$ are C^2 , connected, oriented hypersurfaces with boundary. Note that being an $(n - 2)$ -dimensional boundary puts some restrictions on the support, multiplicity and orientation of the connected components of Σ .

On a closed Riemannian manifold, everything done in this manuscript works for $(n - 2)$ -dimensional boundaries instead of $(n - 2)$ -dimensional cycles. In particular, on every Riemannian manifold with trivial $(n - 2)$ -dimensional homology (where every $(n - 2)$ -dimensional cycle with integer coefficients is an $(n - 2)$ -dimensional boundary), our proofs apply verbatim. That is, on these manifolds, our notion of s -mass is well-defined and, with similar proofs, it Γ -converges and pointwise converges to the $(n - 2)$ -Hausdorff measure with multiplicity. This is the case, for example, for Riemannian spheres (\mathbb{S}^n, g) for $n \geq 3$.

5.2 Higher codimension

Apart from being extended to ambient Riemannian manifolds, our notion is suited to be extended to higher codimension too. In what follows, we work in ambient \mathbb{R}^n and $k \in \{2, \dots, n - 1\}$ denotes the codimension of our surfaces.

Similarly to (5.1), the natural domain for the s -mass in codimension k is the space of k -dimensional boundaries with integer coefficients and trivial normal bundle. That is, for $m \geq 1$ positive integer and $d = (d_1, \dots, d_m) \in \mathbb{N}_+^m$ let

$$\Sigma = \partial(d_1 M_1 \cup \dots \cup d_m M_m) = \bigcup_{i=1}^m \bigcup_{j=1}^{m_i} d_i \Sigma_i^j, \quad (5.2)$$

where $\{M_i\}_{i=1}^m \subset \mathbb{R}^n$ are C^2 , connected, oriented $(n - k + 1)$ -dimensional submanifolds with $\partial M_i = \cup_{j=1}^{m_i} \Sigma_i^j$, and each Σ_i^j is connected and has trivial normal bundle (which is always true when $k = 2$, but not necessarily if $k \geq 3$). Here, for every $i \in \{1, \dots, m\}$ and $j \in \{1, \dots, m_i\}$, each Σ_i^j comes with the induced orientation from M_i .

Denote also

$$L_k(\Sigma) := \{S \subset \mathbb{R}^n \setminus \Sigma : S \text{ is a bi-Lipschitz image of } \mathbb{S}^{k-1}\}.$$

With this notation, we can consider the notion of fractional s -mass of codimension $k \in \{2, \dots, n - 1\}$, for surfaces Σ as in (5.2), defined by

$$\mathbb{M}_{k,s}(\Sigma) := \min \left\{ [u]_{W^{\frac{k-1+s}{k}, k}(\mathbb{R}^n)}^k : u \in \mathfrak{F}_{s,k}^w(\Sigma) \right\},$$

where

$$\begin{aligned} \mathfrak{F}_{s,k}(\Sigma) &:= \left\{ u \in C^1(\mathbb{R}^n \setminus \Sigma; \mathbb{S}^{k-1}) \cap W^{\frac{k-1+s}{k}, k}(\mathbb{R}^n; \mathbb{S}^{k-1}) \text{ such that} \right. \\ &\quad \left. |\deg(u, S)| = \left| \sum_{i=1}^m \sum_{j=1}^{m_i} d_i \text{link}(S, \Sigma_i^j) \right| \text{ for every } S \in L_k(\Sigma) \right\}, \quad (5.3) \end{aligned}$$

and

$$\mathfrak{F}_{s,k}^w(\Sigma) := \left\{ u : \mathbb{R}^n \rightarrow \mathbb{S}^{k-1} : \exists \{u_h\} \subset \mathfrak{F}_{s,k}(\Sigma), u_h \rightarrow u \text{ in } W_{\text{loc}}^{\alpha, k}(\mathbb{R}^n; \mathbb{S}^{k-1}) \forall \alpha \in \left(0, \frac{k-1+s}{2}\right) \right\}.$$

With this definition, it should be possible to extend Theorem 1.1 to any codimension.

Indeed, for the upper bound, at least in the case in which $m = 1$, since the normal bundle of each Σ_1^j is trivial and all the multiplicities agree with the one of M_1 , one can still use the standard vortex of \mathbb{R}^k (or a fixed map of higher degree) in each fiber of a tubular neighborhood of Σ_1^j , and extend this map to an admissible competitor that is defined on \mathbb{R}^n and is constant outside a neighborhood of M_1 . The case in which $m > 1$ here is more delicate, since we cannot use the complex multiplication to build a single map out of the ones relative to each component. Nevertheless, this can still be done trivially when the manifolds M_i are disjoint and probably, with some more effort, also if they are not.

On the other hand, in order to obtain the lower bound, one can still reduce the fractional energy to a discrete one and bound it from below with the following version of the Ginzburg-Landau energy

$$GL_{k,\varepsilon,\beta}(v, \Omega) := \int_{\Omega} \left(\frac{1-\beta}{k|\log \varepsilon|} |\nabla v|^k + \frac{\beta c_k}{\varepsilon^2 |\log \varepsilon|} (1 - |v|^2)^k \right) dx,$$

for which we still have an analogue of Lemma 4.6 (see [ABO05, Lemma 3.10]) that provides an estimate for the energy on k -dimensional cubes. Moreover, for every $s \in (0, 1)$ we have the embedding $W^{\frac{k-1+s}{k}, k}(\mathbb{R}^{k-1}) \hookrightarrow C^0(\mathbb{R}^{k-1})$, which provides an analogue of Lemma 2.8 that we can apply on each $(k - 1)$ -face of the cubes in $\mathcal{C}_{\varepsilon, \nu, z}(F)$.

Very similarly, Serra in [Ser23, Section 5] proposed a notion of $(n - k)$ -dimensional fractional volume $V_{n-k,s}$ (we keep the notation $V_{d,s}$ to match the one in [Ser23], where $d = n - k$) for

boundaries that are level sets of (regular values of) smooth functions. Precisely, for $s \in (0, 1)$ Serra defined the $(n - k)$ -dimensional fractional volume as

$$V_{n-k,s}(\Sigma) := \inf \left\{ [u]_{H^{\frac{k-1+s}{2}}(\mathbb{R}^n)}^2 : u \in \mathfrak{F}_{s,k}(\Sigma) \right\},$$

where the family $\mathfrak{F}_{s,k}(\Sigma)$ is defined as in (5.3) without the multiplicities, and with the space $H^{\frac{k-1+s}{2}}(\mathbb{R}^n; \mathbb{S}^{k-1})$ in place of $W^{\frac{k-1+s}{k},k}(\mathbb{R}^n; \mathbb{S}^{k-1})$.

We remark that on a closed Riemannian manifold M , the spaces $H^\alpha(M)$ for $\alpha \geq 1$ still have few (all equivalent) canonical definitions corresponding to the natural extensions of the ones in [Ser23, Section 4.1]. For example, see [Yan13] for the Caffarelli-Silvestre extension of higher-order fractional Laplacian. Hence, even in this setting of general codimension, this notion of fractional mass seems to be canonical also if \mathbb{R}^n is replaced by a closed Riemannian manifold without boundary.

Note that the family of level-set surfaces in [Ser23] satisfies our general hypothesis above. Indeed, in [Ser23] the $(n - k)$ -dimensional fractional volume $V_{n-k,s}$ is defined for surfaces $\Sigma = F^{-1}(y)$, where $F : \mathbb{R}^n \rightarrow \mathbb{R}^k$ is smooth and y is a regular value of F . These surfaces have trivial normal bundle, since in this setting, being y a regular value of F , $\{\nabla F^i(y)\}_{i=1}^k$ are k independent, nonzero sections of the normal bundle of Σ . Hence, they trivialize each connected component of Σ . Moreover, $\Sigma = F^{-1}(y)$ is the boundary of $M = (F/|F|)^{-1}(\theta)$, which is an embedded $(n - k + 1)$ -surface when $\theta \in \mathbb{S}^{k-1}$ is a regular value of $F/|F|$. This means that the class (5.2) contains all regular level sets of smooth functions.

A Appendix: energy asymptotics of the planar vortex

In the following result, we show the correct asymptotic of the fractional energy in the model case of the standard vortex in \mathbb{R}^2 . This has already been done in Lemma 4.8 in the case $n \geq 3$.

Here, using a slicing procedure, we give a simple proof of this fact for $n = 2$ and when the domain is a ball. Since this result is not essential for our purpose, we just sketch the proof, leaving some details to the reader.

Lemma A.1. *Let $u_\star : D_1 \setminus \{0\} \subset \mathbb{R}^2 \rightarrow \mathbb{S}^1$ be the standard vortex in two dimensions $u_\star(p) = p/|p|$. Then*

$$\lim_{s \rightarrow 1^-} (1 - s)^2 [u_\star]_{H^{\frac{1+s}{2}}(D_1)}^2 = 2\pi^2.$$

Proof. By the slicing formulas (2.6)-(2.7) we have

$$[u_\star]_{H^{\frac{1+s}{2}}(D_1)}^2 = \frac{1}{2} \int_{\mathbb{S}^1} d\theta \int_{-1}^1 [u_\star]_{H^{\frac{1+s}{2}}(S_{\theta,r})}^2 dr,$$

where $S_{\theta,r} := \{r\theta^\perp + \xi\theta : \xi \in (-\sqrt{1-r^2}, \sqrt{1+r^2})\}$. By the rotational invariance of the vortex we see that $[u_\star]_{H^{\frac{1+s}{2}}(S_{\theta,r})}^2$ does not depend on $\theta \in \mathbb{S}^1$, but only on $|r|$. Hence

$$[u_\star]_{H^{\frac{1+s}{2}}(B_1)}^2 = 2\pi \int_0^1 [u_\star]_{H^{\frac{1+s}{2}}(S_{e_1,r})}^2 dr,$$

where $e_1 = (1, 0)$ and $S_{e_1,r} = \{(x, r) \in B_1 \mid -\sqrt{1-r^2} \leq x \leq \sqrt{1-r^2}\}$. Moreover

$$\begin{aligned} [u_\star]_{H^{\frac{1+s}{2}}(S_{e_1,r})}^2 &= \int_{-\sqrt{1-r^2}}^{\sqrt{1-r^2}} \int_{-\sqrt{1-r^2}}^{\sqrt{1-r^2}} \frac{\left| \frac{(x,r)}{\sqrt{x^2+r^2}} - \frac{(y,r)}{\sqrt{y^2+r^2}} \right|^2}{|x-y|^{2+s}} dx dy \\ &= r^{-s} \iint_{I_r \times I_r} \frac{\left| \frac{(x,1)}{\sqrt{x^2+1}} - \frac{(y,1)}{\sqrt{y^2+1}} \right|^2}{|x-y|^{2+s}} dx dy, \end{aligned}$$

where we have substituted in the integral $x = r\tilde{x}, y = r\tilde{y}$ (without renaming the variables) and we have set $I_r = \{|x| \leq \sqrt{r^{-2} - 1}\}$. Note that $I_r \nearrow \mathbb{R}$ as $r \rightarrow 0^+$. Hence, setting $u_1(x) := \frac{(x,1)}{\sqrt{x^2+1}} : \mathbb{R} \rightarrow \mathbb{R}^2$ we can summarize the above computations as

$$[u_\star]_{H^{\frac{1+s}{2}}(\mathbb{D}_1)}^2 = 2\pi \int_0^1 r^{-s} [u_1]_{H^{\frac{1+s}{2}}(I_r)}^2 dr.$$

From [DNPV12, Remark 4.3] we deduce that

$$\lim_{s \rightarrow 1^-} (1-s) [u_1]_{H^{\frac{1+s}{2}}(I_r)}^2 = 2 \int_{I_r} |u_1'(x)|^2 dx,$$

and it is not difficult to check that the convergence is uniform with respect to $r \in (0, 1)$.

Since $(1-s)r^{-s} \mathbf{L}(0, 1] \rightarrow \delta_{\{0\}}$ in duality with $C^0([0, 1])$ and

$$\int_{\mathbb{R}} |u_1'(x)|^2 dx = \int_{\mathbb{R}} \left| \partial_x \left(\frac{x}{\sqrt{1+x^2}} \right) \right|^2 + \left| \partial_x \left(\frac{1}{\sqrt{1+x^2}} \right) \right|^2 dx = \int_{\mathbb{R}} \frac{1}{(1+x^2)^2} dx = \frac{\pi}{2},$$

we conclude that

$$\lim_{s \rightarrow 1^-} (1-s)^2 [u_\star]_{H^{\frac{1+s}{2}}(\mathbb{D}_1)}^2 = 2\pi \int_0^1 (1-s)r^{-s} \cdot (1-s) [u_1]_{H^{\frac{1+s}{2}}(I_r)}^2 dr = 2\pi \cdot 2 \cdot \frac{\pi}{2} = 2\pi^2.$$

□

Remark A.2. Note that the constant $2\pi^2$ is in accordance with the one in Theorem 1.1 for $n = 2$.

Acknowledgements. We are very grateful to Joaquim Serra for proposing this problem to us, and to Andrea Malchiodi and Daniel Stern for many valuable comments on a preliminary version of this work.

We also thank Clara Antonucci, Federica Bertolotti, Giovanni Italiano, Matteo Migliorini, and Diego Santoro for several fruitful discussions.

M. C. is funded by NSF grant DMS-2304432 and is grateful to Stanford University for its kind hospitality during the realization of part of this work.

M. F. and N. P. are members of the “Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni” (GNAMPA) of the “Istituto Nazionale di Alta Matematica” (INdAM).

N. P. acknowledges the MIUR Excellence Department Project awarded to the Department of Mathematics, University of Pisa, CUP I57G22000700001.

M. F. is a member of the PRIN Project 2022AKNSE4 *Variational and Analytical aspects of Geometric PDEs*.

References

- [ABO03] G. Alberti, S. Baldo, and G. Orlandi. Functions with prescribed singularities. *J. Eur. Math. Soc. (JEMS)*, 5(3):275–311, 2003.
- [ABO05] G. Alberti, S. Baldo, and G. Orlandi. Variational convergence for functionals of Ginzburg-Landau type. *Indiana Univ. Math. J.*, 54(5):1411–1472, 2005.
- [AC09] R. Alicandro and M. Cicalese. Variational analysis of the asymptotics of the XY model. *Arch. Ration. Mech. Anal.*, 192(3):501–536, 2009.
- [ADPM11] L. Ambrosio, G. De Philippis, and L. Martinazzi. Gamma-convergence of nonlocal perimeter functionals. *Manuscripta Math.*, 134(3-4):377–403, 2011.

- [Alm62] Frederick Justin Almgren, Jr. The homotopy groups of the integral cycle groups. *Topology*, 1:257–299, 1962.
- [Alm65] F. J. Almgren. The theory of varifolds. *mimeographed notes, Princeton*, 1965.
- [BBM01] J. Bourgain, H. Brezis, and P. Mironescu. Another look at Sobolev spaces. In *Optimal control and partial differential equations*, pages 439–455. IOS, Amsterdam, 2001.
- [BBM02] J. Bourgain, H. Brezis, and P. Mironescu. Limiting embedding theorems for $W^{s,p}$ when $s \uparrow 1$ and applications. *J. Anal. Math.*, 87:77–101, 2002.
- [BM21] H. Brezis and P. Mironescu. *Sobolev maps to the circle—from the perspective of analysis, geometry, and topology*, volume 96 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser/Springer, New York, 2021.
- [Bre93] G. E. Bredon. *Topology and geometry*, volume 139 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1993.
- [CCS21] X. Cabré, E. Cinti, and J. Serra. Stable solutions to the fractional allen-cahn equation in the nonlocal perimeter regime. *to appear in Amer. J. Math.*, 2021.
- [CDSV23] H. Chan, S. Dipierro, J. Serra, and E. Valdinoci. Nonlocal approximation of minimal surfaces: optimal estimates from stability. *arXiv:2308.06328*, 2023.
- [CFS23] M. Caselli, E. Florit, and J. Serra. Yau’s conjecture for nonlocal minimal surfaces. *arXiv:2306.07100*, 2023.
- [CHK⁺24] M. Cicalese, T. Heilmann, A. Kubin, F. Onoue, and M. Ponsiglione. A notion of s -fractional mass for 1-currents in higher codimension. *arXiv:2403.03901*, 2024.
- [CM20] O. Chodosh and C. Mantoulidis. Minimal surfaces and the Allen-Cahn equation on 3-manifolds: index, multiplicity, and curvature estimates. *Ann. of Math. (2)*, 191(1):213–328, 2020.
- [CRS10] L. Caffarelli, J.-M. Roquejoffre, and O. Savin. Nonlocal minimal surfaces. *Comm. Pure Appl. Math.*, 63(9):1111–1144, 2010.
- [CSV19] E. Cinti, J. Serra, and E. Valdinoci. Quantitative flatness results and BV -estimates for stable nonlocal minimal surfaces. *J. Differential Geom.*, 112(3):447–504, 2019.
- [DdPW18] J. Dávila, M. del Pino, and J. Wei. Nonlocal s -minimal surfaces and Lawson cones. *J. Differential Geom.*, 109(1):111–175, 2018.
- [Dip20] Serena Dipierro. A comparison between the nonlocal and the classical worlds: minimal surfaces, phase transitions, and geometric flows. *Notices Amer. Math. Soc.*, 67(9):1324–1335, 2020.
- [DNPV12] E. Di Nezza, G. Palatucci, and E. Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bull. Sci. Math.*, 136(5):521–573, 2012.
- [dPKW13] M. del Pino, M. Kowalczyk, and J. Wei. Entire solutions of the Allen-Cahn equation and complete embedded minimal surfaces of finite total curvature in \mathbb{R}^3 . *J. Differential Geom.*, 93(1):67–131, 2013.
- [DSV20] S. Dipierro, J. Serra, and E. Valdinoci. Improvement of flatness for nonlocal phase transitions. *Amer. J. Math.*, 142(4):1083–1160, 2020.
- [Dá02] J. Dávila. On an open question about functions of bounded variation. *Calc. Var. Partial Differential Equations*, 15(4):519–527, 2002.
- [FF60] H. Federer and W. H. Fleming. Normal and integral currents. *Ann. of Math. (2)*, 72:458–520, 1960.
- [GG18] P. Gaspar and M. A. M. Guaraco. The Allen-Cahn equation on closed manifolds. *Calc. Var. Partial Differential Equations*, 57(4):Paper No. 101, 42, 2018.

- [Gua18] M. A. M. Guaraco. Min-max for phase transitions and the existence of embedded minimal hypersurfaces. *J. Differential Geom.*, 108(1):91–133, 2018.
- [IMN18] K. Irie, F. C. Marques, and A. Neves. Density of minimal hypersurfaces for generic metrics. *Ann. of Math. (2)*, 187(3):963–972, 2018.
- [Jer99] R. L. Jerrard. Lower bounds for generalized Ginzburg-Landau functionals. *SIAM J. Math. Anal.*, 30(4):721–746, 1999.
- [Kir89] R. C. Kirby. *The topology of 4-manifolds*, volume 1374 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1989.
- [Leo23] G. Leoni. *A first course in fractional Sobolev spaces*, volume 229 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, [2023] ©2023.
- [LMN18] Y. Liokumovich, F. C. Marques, and A. Neves. Weyl law for the volume spectrum. *Ann. of Math. (2)*, 187(3):933–961, 2018.
- [MM77a] L. Modica and S. Mortola. Il limite nella Γ -convergenza di una famiglia di funzionali ellittici. *Boll. Un. Mat. Ital. A (5)*, 14(3):526–529, 1977.
- [MM77b] L. Modica and S. Mortola. Un esempio di Γ^- -convergenza. *Boll. Un. Mat. Ital. B (5)*, 14(1):285–299, 1977.
- [MN16] Fernando C. Marques and André Neves. Topology of the space of cycles and existence of minimal varieties. In *Surveys in differential geometry 2016. Advances in geometry and mathematical physics*, volume 21 of *Surv. Differ. Geom.*, pages 165–177. Int. Press, Somerville, MA, 2016.
- [Mod87] L. Modica. The gradient theory of phase transitions and the minimal interface criterion. *Arch. Ration. Mech. Anal.*, 98(2):123–142, 1987.
- [MPS21] V. Millot, M. Pegon, and A. Schikorra. Partial regularity for fractional harmonic maps into spheres. *Arch. Ration. Mech. Anal.*, 242(2):747–825, 2021.
- [MS23] C. Mihaila and B. Seguin. A definition of fractional k -dimensional measure: bridging the gap between fractional length and fractional area. *arXiv:2303.11542*, 2023.
- [Pit81] J. T. Pitts. *Existence and regularity of minimal surfaces on Riemannian manifolds*, volume 27 of *Mathematical Notes*. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1981.
- [Pon03] A. C. Ponce. A variant of Poincaré’s inequality. *C. R. Math. Acad. Sci. Paris*, 337(4):253–257, 2003.
- [Pon04] A. C. Ponce. A new approach to Sobolev spaces and connections to Γ -convergence. *Calc. Var. Partial Differential Equations*, 19(3):229–255, 2004.
- [PS21] A. Pigati and D. Stern. Minimal submanifolds from the abelian Higgs model. *Invent. Math.*, 223(3):1027–1095, 2021.
- [PS23] A. Pigati and D. Stern. Quantization and non-quantization of energy for higher-dimensional Ginzburg-Landau vortices. *Ars Inven. Anal.*, pages Paper No. 3, 55, 2023.
- [Riv17] T. Rivière. A viscosity method in the min-max theory of minimal surfaces. *Publ. Math. Inst. Hautes Études Sci.*, 126:177–246, 2017.
- [Ser23] J. Serra. Nonlocal minimal surfaces: recent developments, applications, and future directions. *SeMA Journal*, 2023.
- [Sol24] M. Solci. Nonlocal-Interaction Vortices. *SIAM J. Math. Anal.*, 56(3):3430–3451, 2024.
- [SW08] R. Schneider and W. Weil. *Stochastic and integral geometry*. Probability and its Applications (New York). Springer-Verlag, Berlin, 2008.

- [TW12] Y. Tonegawa and N. Wickramasekera. Stable phase interfaces in the van der Waals–Cahn–Hilliard theory. *J. Reine Angew. Math.*, 668:191–210, 2012.
- [Yan13] R. Yang. On higher order extensions for the fractional laplacian. *arXiv:1302.4413*, 2013.

MICHELE CASELLI

Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy
Stanford University, 450 Jane Stanford Way, Building 380, 94305 Stanford, CA
E-mail address: michele.caselli@sns.it

MATTIA FREGUGLIA

Scuola Normale Superiore
Piazza dei Cavalieri 7, 56126 Pisa, Italy
E-mail address: mattia.freguglia@sns.it

NICOLA PICENNI

University of Pisa, Department of Mathematics
Largo Bruno Pontecorvo 5, 56127 Pisa, Italy
E-mail address: nicola.picenni@dm.unipi.it