

12th Workshop on the CKM Unitarity Triangle  
Santiago de Compostela, 18-22 September 2023

## Working group 1 summary: $V_{ud}$ , $V_{us}$ , $V_{cd}$ , $V_{cs}$ and semileptonic/leptonic $D$ decays

Bipasha Chakraborty<sup>\*1</sup>, Alex Gilman<sup>†2</sup>,  
Martin Hoferichter<sup>‡3</sup>, and Michal Koval<sup>§4</sup>

<sup>1</sup>University of Southampton  
University Road, SO17 IBJ, United Kingdom

<sup>2</sup>University of Oxford

Denys Wilkinson Building

Keble Road, OX1 3RH, United Kingdom

<sup>3</sup>Albert Einstein Center for Fundamental Physics,  
Institute for Theoretical Physics, University of Bern,  
Sidlerstrasse 5, 3012 Bern, Switzerland

<sup>4</sup>Institute of Particle and Nuclear Physics,  
Faculty of Mathematics and Physics, Charles University,  
V Holesovickach 2, 180 00 Prague 8, Czech Republic

### Abstract

We summarize the program of working group 1 at the 12th Workshop on the CKM Unitarity Triangle, whose main subjects covered  $V_{ud}$ ,  $V_{us}$ , and first-row unitarity as well as  $V_{cd}$ ,  $V_{cs}$ , and (semi-)leptonic  $D$  decays.

---

\*b.chakraborty@soton.ac.uk

†alex.gilman@physics.ox.ac.uk

‡hoferichter@itp.unibe.ch

§michal.koval@matfyz.cuni.cz

# 1 Introduction

While the program of the CKM 2023 conference focused on precision tests of the CKM paradigm via unitarity triangles, a separate class of precision tests only involves the moduli of the CKM matrix elements  $|V_{ij}|$ , since unitarity demands that

$$\sum_{j=d,s,b} |V_{ij}|^2 = 1, \quad \sum_{i=u,c,t} |V_{ij}|^2 = 1, \quad (1.1)$$

for fixed  $i = u, c, t$  and  $j = d, s, b$ , respectively. The corresponding unitarity tests thus amount to unitarity circles, referred to as “first-row” and “first-column” unitarity. The primary goal of the work presented in WG 1 constitutes improvements in the unitarity test involving  $V_{ud}$ ,  $V_{us}$ ,  $V_{cd}$ ,  $V_{cs}$ , i.e., first/second row/column, given that the third-generation elements therein are negligible at the current level of precision. Looking at the status as quoted in Ref. [1]:

$$\begin{aligned} \text{first row:} & \quad |V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9984(7), \\ \text{second row:} & \quad |V_{cd}|^2 + |V_{cs}|^2 + |V_{cb}|^2 = 1.001(12), \\ \text{first column:} & \quad |V_{ud}|^2 + |V_{cd}|^2 + |V_{td}|^2 = 0.9971(20), \\ \text{second column:} & \quad |V_{us}|^2 + |V_{cs}|^2 + |V_{ts}|^2 = 1.003(12), \end{aligned} \quad (1.2)$$

one observes that there are hints for a deficit in the first-row and first-column unitarity tests, motivating increased efforts both in experiment and theory to corroborate the current results and further increase the precision. Here, we provide a summary of the corresponding working group 1 program, while for some presentations, separate write-ups are available [2–6].

## 2 $V_{ud}$ , $V_{us}$ , and first-row unitarity

The current status of the first-row unitarity test is shown in Fig. 1, including constraints from  $K_{\ell 3} = K \rightarrow \pi \ell \nu_\ell$  decays,  $\pi_{\ell 2}/K_{\ell 2}$ , and  $\beta$  decays. Tensions arise both between the global fit and with unitarity, but each pair of constraints by itself leads to a unitarity deficit  $\Delta_{\text{CKM}} \equiv |V_{ud}|^2 + |V_{us}|^2 - 1 < 0$  [7]

$$\begin{aligned} \Delta_{\text{CKM}}^{K_{\ell 2}-K_{\ell 3}} &= -0.016(6) [2.6\sigma], \\ \Delta_{\text{CKM}}^{K_{\ell 2}-\beta} &= -0.0010(6) [1.7\sigma], \\ \Delta_{\text{CKM}}^{K_{\ell 3}-\beta} &= -0.0018(6) [3.1\sigma], \\ \Delta_{\text{CKM}}^{\text{global}} &= -0.0018(6) [2.8\sigma], \end{aligned} \quad (2.1)$$

suggesting a tension even just within the kaon sector. Accordingly, it is clear that both  $V_{ud}$  and  $V_{us}$  need to be improved.




Figure 1: Current status of first-row CKM unitarity, figure taken from Ref. [7]. The (green) horizontal band represents  $V_{us}$  from  $K_{\ell 3}$  decays, the (blue) diagonal band  $V_{us}/V_{ud}$  from  $K_{\ell 2}/\pi_{\ell 2}$  decays, and the vertical bands to  $V_{ud}$  from superallowed  $\beta$  decays (left, red) as well as neutron decay (right, violet), respectively. The unitarity circle is displayed by the black solid line, the global fit by the yellow ellipse.

## 2.1 $V_{ud}$

Sources for  $V_{ud}$  determinations are summarized in Table 1, together with their respective advantages and disadvantages. In the following, we discuss recent developments for each method in more detail.

For superallowed  $\beta$  decays, one argument for the reliability of nuclear corrections brought forward in Ref. [8] is that once all corrections are applied, one observes good consistency among a large number of nuclear transitions. To test the nuclear models that were employed traditionally, a new method to compute nuclear-structure corrections using dispersion relations was developed [9, 10], which was recently applied to the  $^{10}\text{C} \rightarrow ^{10}\text{B}$  transition in combination with modern ab-initio nuclear-structure methods [11]. Moreover, measurements of nuclear charge radii could help constrain nuclear-structure calculations [12], and such measurements are possible, e.g., at PSI, FRIB, and ISOLDE.

A complementary approach using effective field theory (EFT) is being developed to establish a framework that allows one to explicitly take advantage of the separation of scales, resum large logarithms, and systematically improve

	+	-
Superaligned $\beta$ decays	many isotopes to average	nuclear uncertainties
Neutron decay	no nuclear uncertainties	need high precision for $\tau_n$ and $g_A$
Pion $\beta$ decay	theoretically pristine	experimentally challenging

Table 1: Sources for  $V_{ud}$  determinations, indicating some advantages and disadvantages of each method.

calculations by including subleading orders in the various EFT expansions. The application to neutron decay [13] was discussed in detail, demonstrating how the different scales ranging from the mass of the  $W$ -boson down to the electron mass and small  $Q$  value of the decay have to be evolved and at which places the non-perturbative input enters. Recently, this EFT approach was extended to nuclear decays, including a first ab-initio application to  $^{14}\text{O} \rightarrow ^{14}\text{N}$  [14, 15].

For neutron decay, the uncertainty in the resulting  $V_{ud}$  value is dominated by the experimental input, since both for the neutron lifetime  $\tau_n$  and the asymmetry parameter  $\lambda = g_A/g_V$  very high precision is required. For  $\tau_n$ , the precision necessary for a determination competitive with superallowed  $\beta$  decays is already within reach [16], but  $\lambda$  needs to be further improved, and, crucially, the tension between PERKEO III [17] and aSPECT [18] must be resolved. At the future PERC experiment, a precision of  $\Delta\lambda/\lambda \simeq 10^{-4}$  is anticipated, which, combined with ongoing improvements in  $\tau_n$  measurements, would allow for an independent, competitive  $V_{ud}$  determination free of nuclear uncertainties.

Third,  $V_{ud}$  can be determined from pion  $\beta$  decay,  $\pi^+ \rightarrow \pi^0 e^+ \bar{\nu}_e$ , but its measurement at the required level of precision is extremely challenging. PIONEER is a next-generation rare pion decay experiment at PSI [19] that will pursue such a measurement in Phases II+III of its physics program, while concentrating on lepton flavor universality in the ratio  $R_{e/\mu} = \Gamma[\pi^+ \rightarrow e^+ \nu_e(\gamma)]/\Gamma[\pi^+ \rightarrow \mu^+ \nu_\mu(\gamma)]$  in Phase I (which could be related to CKM unitarity [20, 21]). For both measurements, improvements by a factor 10 are possible before theory uncertainties become relevant, presenting unique opportunities to explore uncharted parameter space.

## 2.2 $V_{us}$

For  $V_{us}$ , the results of an updated global fit to kaon decays were presented, using the methodology from Refs. [22, 23], and leading to the situation depicted in Fig. 1, see Ref. [7]. While new data for  $K_S \rightarrow \pi e \nu$  have become available [24]

$$\text{Br}[K_S \rightarrow \pi e \nu] = 7.153(37)_{\text{stat}}(44)_{\text{syst}} \times 10^{-4}, \quad (2.2)$$

the tension between  $K_{\ell 2}$  and  $K_{\ell 3}$  determinations remains, motivating a new measurement of  $K_{\mu 3}/K_{\mu 2}$  at NA62 or elsewhere [25] to either resolve the tension or corroborate the need for a BSM explanation.




Figure 2: Updated Fermilab-MILC determination of  $|V_{cs}|$  compared to the 2021 HFLAV average results [37] from leptonic charm decays. Figure taken from Ref. [38].

The updated fit to  $K_{\ell 2}$  and  $K_{\ell 3}$  data also includes improvements in critical radiative corrections. For  $K_{\ell 3}$  decays these were evaluated in a dispersive framework [26–28], in agreement with earlier calculations in chiral perturbation theory (ChPT) [29, 30], but reducing the uncertainty due to low-energy constants, mainly by evaluating Born-term contributions in terms of known form factors and using input from lattice QCD. For  $K_{\ell 2}$  decays, the radiative corrections from ChPT [31, 32] have been crosschecked by direct lattice-QCD calculations [33, 34], and further improvements are being developed [35]. In the context of lattice-QCD calculations for rare kaon decays, also the prospects for  $\Sigma^+ \rightarrow p\ell^+\ell^-$  were discussed [36].

### 3 $V_{cd}$ , $V_{cs}$ , and (semi-)leptonic $D$ decays

Precision on the magnitudes  $|V_{cd}|$  and  $|V_{cs}|$  is driven by the study of purely leptonic and semileptonic decays of charmed hadrons. The magnitude of the CKM elements can be determined from the measured decay rates with external measurements of the charmed hadron lifetime and an estimation of QCD decay constants or form factors.

#### 3.1 Updated precision on $|V_{cs}|$

Significantly improved precision has been achieved on  $|V_{cs}|$  since the last iteration of the CKM conference due to new work from experimental collaborations and




Figure 3: Different kinematical contributions to the inclusive  $D_s^+ \rightarrow X_s \ell^+ \nu_e$  decay rate compared to exclusive lattice data [51].

new lattice determinations of QCD form factors. In the two intervening years, the BESIII collaboration published updated measurements of  $D_s^+ \rightarrow \mu^+ \nu_\mu$  [39] and  $D_s^+ \rightarrow \tau^+ \nu_\tau$  with multiple  $\tau^+$  final states [40–42]. In combination, these updated measurements determine  $|V_{cs}| = 0.9774(56)(72)$  with inputs of particle masses [1], the  $D_s^+$  lifetime [1], and the  $D_s^+$  decay constant [43]. New calculations of  $D$  to pseudoscalar form factors from both the Fermilab-MILC [38] and HPQCD collaborations [44] provide significantly improved determinations of  $|V_{cs}|$  from the BESIII measurements of  $D^0 \rightarrow K^- \ell^+ \nu_\ell$  [45, 46]. The impacts of the new Fermilab-MILC calculation are shown in Fig. 2, compared to an average determination from leptonic charm decays from HFLAV 2021 [37], which does not include the recent BESIII results.

The Belle II collaboration measured the lifetimes of multiple charmed mesons with significantly improved precision [47–50]. The updated measurements of the  $D^0$ ,  $D^+$ ,  $D_s^+$ , and  $\Lambda_c^+$  lifetimes reduce uncertainties in the determination of  $|V_{cd}|$  and  $|V_{cs}|$ . Prospects for competitive measurements of leptonic and semileptonic charm decays at Belle II in future datasets were also presented.

### 3.2 Progress on inclusive semileptonic charm decays

A joint session with working group 2 was held on inclusive semileptonic decays of heavy hadrons. New measurements of the inclusive  $D_s^+ \rightarrow X e^+ \nu_e$  and  $\Lambda_c^+ \rightarrow X e^+ \nu_e$  branching fractions and lepton momentum distributions were reported by the BESIII collaboration, both improving the precision on observables by roughly a factor of three [52, 53]. These measurements, in comparison to the sum of the known exclusive branching fractions [1], can be used to infer the

existence of unobserved semileptonic decay channels. In the  $D_s^+$  case, the difference between the measured inclusive branching fraction and the sum of exclusive branching fractions is  $-0.04(13)(20)\%$ , providing no evidence for unobserved  $D_s^+$  semileptonic modes within the measured precision. However, a similar comparison for the  $\Lambda_c^+$  case indicates the unobserved semileptonic  $\Lambda_c^+$  decays [54].

Due to theoretical challenges,  $|V_{cd}|$  and  $|V_{cs}|$  have solely been determined through exclusive semileptonic decays in contrast to the beauty sector. Recent progress in both lattice QCD and analytic theory presents exciting prospects for the extraction of  $|V_{cd}|$  and  $|V_{cs}|$  from inclusive semileptonic charm decays. Significant progress has been made in the study of inclusive semileptonic decays on the lattice, demonstrated in the study of  $D_s^+ \rightarrow X_s \ell^+ \nu_e$  [51]. The determinations of the different kinematical contributions to the inclusive  $D_s^+ \rightarrow X_s \ell^+ \nu_e$  decay rate as determined by Ref. [51] is shown in Fig. 3. Progress in the heavy quark effective theory treatment of inclusive charm decays were also discussed [55], similar to that employed in beauty decays as, e.g., in Ref. [56]. This progress from the theory and the lattice indicates exciting prospects for the study of inclusive semileptonic charm decays and encourages detailed experimental analysis of these processes.

## 4 Conclusions

Intriguing tensions of almost  $3\sigma$  persist in the first-row test of CKM unitarity, both of the global fit with the unitarity circle and among different constraints on  $V_{ud}$  and  $V_{us}$ . Several ways to corroborate or refute these tensions were discussed, for  $V_{ud}$  ranging from improved theory for radiative corrections to superallowed  $\beta$  decays and neutron decay, over experimental prospects for competitive measurements of neutron lifetime and decay asymmetry, to a long-term vision for a precision measurement of pion  $\beta$  decay. For  $V_{us}$  also improved radiative corrections for both  $K_{\ell 2}$  and  $K_{\ell 3}$  decays have become available, as well as new data on  $K_S \rightarrow \pi e \nu$ , leading to a situation that urgently calls for a new measurement of  $K_{\mu 2}/K_{\mu 3}$  at NA62 or elsewhere.

The total uncertainty on determinations of  $|V_{cs}|$  from direct measurements was reduced by a factor of two, achieving sub-percent precision since the last iteration of the CKM conference. Promising prospects for further precision on  $|V_{cs}|$  and  $|V_{cd}|$  were presented, including the growing data sample at Belle II, large new datasets collected near the  $\psi(3770)$  resonance by BESIII, and work from the lattice and analytic theory communities in analyzing charm decays. These could lead to competitive tests of unitarity from the second row and the first and second columns of the CKM matrix.

## Acknowledgments

We would like to thank the local organizers for putting together a great conference, and all the speakers in our working group for their contributions. Financial support by the SNSF (Project No. PCEFP2\_181117), the Czech Science Foundation (Project No. 23-06770S), Charles University (Project PRIMUS 23/SCI/025), Leverhulme Trust (ECF-2019-223 G100820), STFC (Grant no. ST/X000583/1), STFC (Grant no. ST/W006251/1), and EPSRC (Grant no. EP/W032635/1) is gratefully acknowledged.

## References

- [1] PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *PTEP* **2022** (2022) 083C01.
- [2] M. Hoferichter, *Prospects for PIONEER*, in *12th International Workshop on the CKM Unitarity Triangle*, 2024 [2403.18889].
- [3] BESIII collaboration, *Experimental status of  $|V_{cd}|$  and  $|V_{cs}|$* , in *12th International Workshop on the CKM Unitarity Triangle*, 2024 [2404.00592].
- [4] KLOE-2 collaboration, *A new  $K_S \rightarrow \pi e \nu$  branching fraction measurement from KLOE-2*, in *12th International Workshop on the CKM Unitarity Triangle*, 2024 [2404.01826].
- [5] A.J. Schwartz, *Charm lifetime measurements at Belle II*, in *12th International Workshop on the CKM Unitarity Triangle*, 2024 [2405.10394].
- [6] R. Kellermann, A. Barone, S. Hashimoto, A. Jüttner and T. Kaneko, *Updates on inclusive charmed and bottomed meson decays from the lattice*, in *12th International Workshop on the CKM Unitarity Triangle*, 2024 [2405.06152].
- [7] V. Cirigliano, A. Crivellin, M. Hoferichter and M. Moulson, *Scrutinizing CKM unitarity with a new measurement of the  $K_{\mu 3}/K_{\mu 2}$  branching fraction*, *Phys. Lett. B* **838** (2023) 137748 [2208.11707].
- [8] J.C. Hardy and I.S. Towner, *Superallowed  $0^+ \rightarrow 0^+$  nuclear  $\beta$  decays: 2020 critical survey, with implications for  $V_{ud}$  and CKM unitarity*, *Phys. Rev. C* **102** (2020) 045501.
- [9] C.-Y. Seng and M. Gorchtein, *Dispersive formalism for the nuclear structure correction  $\delta_{NS}$  to the  $\beta$  decay rate*, *Phys. Rev. C* **107** (2023) 035503 [2211.10214].

- [10] M. Gorchtein and C.Y. Seng, *Superaligned nuclear beta decays and precision tests of the Standard Model*, *Ann. Rev. Nucl. Part. Sci.* **74** (2024) 23 [2311.00044].
- [11] M. Gennari, M. Drissi, M. Gorchtein, P. Navratil and C.-Y. Seng, *An ab initio recipe for taming nuclear-structure dependence of  $V_{ud}$ : the  $^{10}\text{C} \rightarrow ^{10}\text{B}$  superallowed transition*, 2405.19281.
- [12] C.-Y. Seng and M. Gorchtein, *Electroweak nuclear radii constrain the isospin breaking correction to  $V_{ud}$* , *Phys. Lett. B* **838** (2023) 137654 [2208.03037].
- [13] V. Cirigliano, W. Dekens, E. Mereghetti and O. Tomalak, *Effective field theory for radiative corrections to charged-current processes: Vector coupling*, *Phys. Rev. D* **108** (2023) 053003 [2306.03138].
- [14] V. Cirigliano, W. Dekens, J. de Vries, S. Gandolfi, M. Hoferichter and E. Mereghetti, *Ab-initio electroweak corrections to superallowed  $\beta$  decays and their impact on  $V_{ud}$* , 2405.18464.
- [15] V. Cirigliano, W. Dekens, J. de Vries, S. Gandolfi, M. Hoferichter and E. Mereghetti, *Radiative corrections to superallowed  $\beta$  decays in effective field theory*, 2405.18469.
- [16] UCN $\tau$  collaboration, *Improved neutron lifetime measurement with UCN $\tau$* , *Phys. Rev. Lett.* **127** (2021) 162501 [2106.10375].
- [17] B. Märkisch et al., *Measurement of the Weak Axial-Vector Coupling Constant in the Decay of Free Neutrons Using a Pulsed Cold Neutron Beam*, *Phys. Rev. Lett.* **122** (2019) 242501 [1812.04666].
- [18] M. Beck et al., *Improved determination of the  $\beta$ - $\bar{\nu}_e$  angular correlation coefficient  $a$  in free neutron decay with the aSPECT spectrometer*, *Phys. Rev. C* **101** (2020) 055506 [1908.04785].
- [19] PIONEER collaboration, *PIONEER: Studies of Rare Pion Decays*, 2203.01981.
- [20] A. Crivellin and M. Hoferichter,  *$\beta$  Decays as Sensitive Probes of Lepton Flavor Universality*, *Phys. Rev. Lett.* **125** (2020) 111801 [2002.07184].
- [21] A. Crivellin, M. Hoferichter and C.A. Manzari, *Fermi Constant from Muon Decay Versus Electroweak Fits and Cabibbo-Kobayashi-Maskawa Unitarity*, *Phys. Rev. Lett.* **127** (2021) 071801 [2102.02825].
- [22] FLAVIANET WORKING GROUP ON KAON DECAYS collaboration, *An Evaluation of  $|V_{us}|$  and precise tests of the Standard Model from world data on leptonic and semileptonic kaon decays*, *Eur. Phys. J. C* **69** (2010) 399 [1005.2323].

- [23] M. Moulson, *Experimental determination of  $V_{us}$  from kaon decays*, *PoS CKM2016* (2017) 033 [[1704.04104](#)].
- [24] KLOE-2 collaboration, *Measurement of the  $K_S \rightarrow \pi e \nu$  branching fraction with the KLOE experiment*, *JHEP* **02** (2023) 098 [[2208.04872](#)].
- [25] G. Anzivino et al., *Workshop summary: Kaons@CERN 2023*, *Eur. Phys. J. C* **84** (2024) 377 [[2311.02923](#)].
- [26] C.-Y. Seng, D. Galviz, M. Gorchtein and U.-G. Meißner, *High-precision determination of the  $K_{e3}$  radiative corrections*, *Phys. Lett. B* **820** (2021) 136522 [[2103.00975](#)].
- [27] C.-Y. Seng, D. Galviz, M. Gorchtein and U.-G. Meißner, *Improved  $K_{e3}$  radiative corrections sharpen the  $K_{\mu 2}$ - $K_{l 3}$  discrepancy*, *JHEP* **11** (2021) 172 [[2103.04843](#)].
- [28] C.-Y. Seng, D. Galviz, M. Gorchtein and U.-G. Meißner, *Complete theory of radiative corrections to  $K_{e3}$  decays and the  $V_{us}$  update*, *JHEP* **07** (2022) 071 [[2203.05217](#)].
- [29] V. Cirigliano, M. Knecht, H. Neufeld, H. Rupertsberger and P. Talavera, *Radiative corrections to  $K_{l3}$  decays*, *Eur. Phys. J. C* **23** (2002) 121 [[hep-ph/0110153](#)].
- [30] V. Cirigliano, M. Giannotti and H. Neufeld, *Electromagnetic effects in  $K_{l3}$  decays*, *JHEP* **11** (2008) 006 [[0807.4507](#)].
- [31] M. Knecht, H. Neufeld, H. Rupertsberger and P. Talavera, *Chiral perturbation theory with virtual photons and leptons*, *Eur. Phys. J. C* **12** (2000) 469 [[hep-ph/9909284](#)].
- [32] V. Cirigliano and H. Neufeld, *A note on isospin violation in  $P_{l2(\gamma)}$  decays*, *Phys. Lett. B* **700** (2011) 7 [[1102.0563](#)].
- [33] M. Di Carlo et al., *Light-meson leptonic decay rates in lattice QCD+QED*, *Phys. Rev. D* **100** (2019) 034514 [[1904.08731](#)].
- [34] P. Boyle et al., *Isospin-breaking corrections to light-meson leptonic decays from lattice simulations at physical quark masses*, *JHEP* **02** (2023) 242 [[2211.12865](#)].
- [35] M. Di Carlo, *Isospin-breaking corrections to weak decays: the current status and a new infrared improvement*, *PoS LATTICE2023* (2024) 120 [[2401.07666](#)].
- [36] F. Erben, V. Gülpers, M.T. Hansen, R. Hodgson and A. Portelli, *Prospects for a lattice calculation of the rare decay  $\Sigma^+ \rightarrow p l^+ l^-$* , *JHEP* **04** (2023) 108 [[2209.15460](#)].

- [37] HFLAV collaboration, *Averages of b-hadron, c-hadron, and  $\tau$ -lepton properties as of 2021*, *Phys. Rev. D* **107** (2023) 052008 [2206.07501].
- [38] FERMILAB LATTICE, MILC collaboration, *D-meson semileptonic decays to pseudoscalars from four-flavor lattice QCD*, *Phys. Rev. D* **107** (2023) 094516 [2212.12648].
- [39] BESIII collaboration, *Improved measurement of the branching fraction of  $D_s^+ \rightarrow \mu^+ \nu_\mu$* , *Phys. Rev. D* **108** (2023) 112001 [2307.14585].
- [40] BESIII collaboration, *Measurement of the Absolute Branching Fraction of  $D_s^+ \rightarrow \tau^+ \nu_\tau$  via  $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$* , *Phys. Rev. Lett.* **127** (2021) 171801 [2106.02218].
- [41] BESIII collaboration, *Measurement of the absolute branching fractions for purely leptonic  $D_s^+$  decays*, *Phys. Rev. D* **104** (2021) 052009 [2102.11734].
- [42] BESIII collaboration, *Measurement of the branching fraction of leptonic decay  $D_s^+ \rightarrow \tau^+ \nu_\tau$  via  $\tau^+ \rightarrow \pi^+ \pi^0 \bar{\nu}_\tau$* , *Phys. Rev. D* **104** (2021) 032001 [2105.07178].
- [43] FLAVOUR LATTICE AVERAGING GROUP (FLAG) collaboration, *FLAG Review 2021*, *Eur. Phys. J. C* **82** (2022) 869 [2111.09849].
- [44] HPQCD collaboration, *Improved  $V_{cs}$  determination using precise lattice QCD form factors for  $D \rightarrow K \ell \nu$* , *Phys. Rev. D* **104** (2021) 034505 [2104.09883].
- [45] BESIII collaboration, *Study of Dynamics of  $D^0 \rightarrow K^- e^+ \nu_e$  and  $D^0 \rightarrow \pi^- e^+ \nu_e$  Decays*, *Phys. Rev. D* **92** (2015) 072012 [1508.07560].
- [46] BESIII collaboration, *Study of the  $D^0 \rightarrow K^- \mu^+ \nu_\mu$  dynamics and test of lepton flavor universality with  $D^0 \rightarrow K^- \ell^+ \nu_\ell$  decays*, *Phys. Rev. Lett.* **122** (2019) 011804 [1810.03127].
- [47] BELLE-II collaboration, *Precise measurement of the  $D^0$  and  $D^+$  lifetimes at Belle II*, *Phys. Rev. Lett.* **127** (2021) 211801 [2108.03216].
- [48] BELLE-II collaboration, *Precise Measurement of the  $D_s^+$  Lifetime at Belle II*, *Phys. Rev. Lett.* **131** (2023) 171803 [2306.00365].
- [49] BELLE-II collaboration, *Measurement of the  $\Lambda_c^+$  Lifetime*, *Phys. Rev. Lett.* **130** (2023) 071802 [2206.15227].
- [50] BELLE-II collaboration, *Measurement of the  $\Omega_c^0$  lifetime at Belle II*, *Phys. Rev. D* **107** (2023) L031103 [2208.08573].

- [51] R. Kellermann, A. Barone, S. Hashimoto, A. Jüttner and T. Kaneko, *Inclusive semi-leptonic decays of charmed mesons with Möbius domain wall fermions*, *PoS LATTICE2022* (2023) 414 [[2211.16830](#)].
- [52] BESIII collaboration, *Measurement of the absolute branching fraction of inclusive semielectronic  $D_s^+$  decays*, *Phys. Rev. D* **104** (2021) 012003 [[2104.07311](#)].
- [53] BESIII collaboration, *Improved measurement of the absolute branching fraction of inclusive semileptonic  $\Lambda_c^+$  decay*, *Phys. Rev. D* **107** (2023) 052005 [[2212.03753](#)].
- [54] BESIII collaboration, *First observation of the semileptonic decay  $\Lambda_c^+ \rightarrow pK^-e^+\nu_e$* , *Phys. Rev. D* **106** (2022) 112010 [[2207.11483](#)].
- [55] M. Fael, T. Mannel and K.K. Vos, *The Heavy Quark Expansion for Inclusive Semileptonic Charm Decays Revisited*, *JHEP* **12** (2019) 067 [[1910.05234](#)].
- [56] F. Bernlochner, M. Fael, K. Olschewsky, E. Persson, R. van Tonder, K.K. Vos et al., *First extraction of inclusive  $V_{cb}$  from  $q^2$  moments*, *JHEP* **10** (2022) 068 [[2205.10274](#)].