

# Neutrino Tomography of the Earth: the Earth Total Mass, Moment of Inertia and Hydrostatic Equilibrium Constraints

S. T. Petcov<sup>a,b)</sup> <sup>1</sup>

<sup>a</sup> *INFN/SISSA, Via Bonomea 265, 34136 Trieste, Italy*

<sup>b</sup> *Kavli IPMU (WPI), University of Tokyo, 5-1-5 Kashiwanoha, 277-8583 Kashiwa, Japan*

## Abstract

We investigate the implications of the constraints following from the precise knowledge of the total Earth mass,  $M_{\oplus}$ , and moment of inertia,  $I_{\oplus}$ , and from the requirement that Earth be in hydrostatic equilibrium (EHE), in the neutrino tomography studies of the Earth density structure. In order to estimate the sensitivity of a given neutrino detector to possible deviations of the inner core (IC), outer core (OC), core (IC + OC) and mantle Earth densities from those obtained using geophysical and seismological data and described by the preliminary reference Earth model (PREM), in the statistical analyses performed within the neutrino tomography studies one typically varies the density of each of these structures. These variations, however, must respect the  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints. Working with PREM average densities we derive the  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints on the possible density variations when one approximates the Earth density structure with i) three layers - mantle, outer core and inner core, and ii) four layers - upper mantle, lower mantle, outer core and inner core. We get drastically different results in the two cases.

---

<sup>1</sup>Also at: Institute of Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, 1784 Sofia, Bulgaria.

# 1 Introduction

At present our knowledge about the interior composition of the Earth and its density structure is based primarily on seismological and geophysical data (see, e.g., [1–9]). These data were used to construct the Preliminary Reference Earth Model (PREM) [10] of the density distribution of the Earth. In the PREM model, the Earth density distribution  $\rho_E$  is assumed to be spherically symmetric,  $\rho_E = \rho_E(r)$ ,  $r$  being the distance from the Earth center, and there are three major density structures (or layers)- the inner core, outer core and the mantle, and a certain number of substructures (shells). In addition there are three shells below, but very close to, the Earth surface - the ocean and the two crust “surface” layers. The mantle has five shells in the model. The mean Earth radius is  $R_\oplus = 6371$  km; the Earth core has a radius of  $R_c = 3480$  km, with the IC and OC extending respectively from  $r = 0$  to  $r = 1221.5$  km, and from  $r = 1221.5$  km to  $r = 3480$  km. The mantle layer is located between the outer core and the lower crust layer, i.e., for  $r$  in the interval  $R_c \leq r \leq R_{\text{man}}$ , with  $R_{\text{man}} = 6346.6$  km. The mean densities of the mantle, the inner and out core are respectively  $\bar{\rho}_{\text{man}} = 4.66$  g/cm<sup>3</sup>,  $\bar{\rho}_{\text{IC}} = 12.89$  g/cm<sup>3</sup> and  $\bar{\rho}_{\text{OC}} = 10.90$  g/cm<sup>3</sup>. The change of density between the mantle and the outer core is described by a step function.

The determination of the radial density distributions in the mantle, outer core and inner core (or core)  $\rho_{\text{man}}(r)$ ,  $\rho_{\text{OC}}(r)$  and  $\rho_{\text{IC}}(r)$  (or  $\rho_c(r)$ ) from seismological and geophysical data is not direct and suffers from uncertainties [2–5]. An approximate and perhaps rather conservative estimate of this uncertainty for  $\rho_{\text{man}}(r)$  is  $\sim 5\%$ ; for the core density  $\rho_c(r)$  it is larger and can be significantly larger [2–5]. It was concluded in [4] (see also [5]), in particular, that the density increase across the inner core - outer core boundary is known with an uncertainty of about 20%.

A precise knowledge of the Earth’s density distribution and of the average densities of the Earth’s three different major structures - the mantle, outer core and inner core - is essential for understanding the physical conditions and fundamental aspects of the structure and properties of the Earth’s interior (including the dynamics of mantle and core, the bulk composition of the Earth’s three structures, the generation, properties and evolution of the Earth’s magnetic field and the gravity field of the Earth) [1, 2, 11, 12]. The thermal evolution of the Earth’s core, in particular, depends critically on the density change across the inner core - outer core boundary (see, e.g., [13]).

A unique alternative method of obtaining information about the density profile of the Earth is the neutrino tomography of the Earth [14–27]. The propagation of the active flavour neutrinos and antineutrinos  $\nu_\alpha$  and  $\bar{\nu}_\alpha$ ,  $\alpha = e, \mu, \tau$ , in the Earth is affected by the Earth matter. The original idea of neutrino Earth tomography is based on the observation that the cross section of the neutrino-nucleon interaction rises with energy. For neutrinos with energies  $E_\nu \gtrsim$  a few TeV, the inelastic scattering off protons and neutrons leads to absorption of neutrinos and thus to attenuation of the initial neutrino flux. The magnitude of the attenuation depends on the Earth matter density along the neutrino path. Attenuation data for neutrinos with different path-lengths in the Earth carry information about the matter density distribution in the Earth interior. The absorption method of Earth tomography with accelerator neutrino beams, which is difficult (if not impossible) to realise in practice was

discussed first in [14, 15] and later in grater detail in [16–27].

The oscillations between the active flavour neutrinos and antineutrinos,  $\nu_\alpha \leftrightarrow \nu_\beta$  and  $\bar{\nu}_\alpha \leftrightarrow \bar{\nu}_\beta$ ,  $\alpha, \beta = e, \mu$  having energies in the range  $E \sim (0.1 - 15.0)$  GeV and traversing the Earth can be strongly modified by the Earth matter effects (see, e.g., [28]). These modifications depend on the Earth matter density along the path of the neutrinos <sup>1</sup>. Thus, by studying the effects of Earth matter on the oscillations of, e.g.,  $\nu_\mu$  and  $\nu_e$  ( $\bar{\nu}_\mu$  and  $\bar{\nu}_e$ ) neutrinos traversing the Earth along different trajectories it is possible to obtain information about the Earth density distribution.

Atmospheric neutrinos (see, e.g., [36, 37]) are a perfect tool for performing Earth tomography. Consisting of significant fluxes of muon and electron neutrinos and antineutrinos,  $\nu_\mu$ ,  $\nu_e$ ,  $\bar{\nu}_\mu$  and  $\bar{\nu}_e$ , produced in the interactions of cosmic rays with the Earth atmosphere, they have a wide range of energies spanning the interval from a few MeV to multi-GeV to multi-TeV. Being produced isotropically in the upper part of the Earth atmosphere at a height of  $\sim 15$  km, they travel distances from  $\sim 15$  km to 12742 km before reaching detectors located on the Earth surface, crossing the Earth along all possible directions and thus “scanning” the Earth interior. The interaction rates that allow to get information about the Earth density distribution can be obtained in the currently taking data IceCube experiment [38, 39] and its planned upgrade [39], in the ORCA [40], Hyper Kamiokande [41] and DUNE [42] experiments which are under construction, and in the planned INO [43] experiment.

The idea of using the absorption method of Earth tomography with atmospheric neutrinos was discussed first, to our knowledge, in [27]. In 2018 in [33] the authors used the data of the IceCube experiment on multi-TeV atmospheric  $\nu_\mu$  and  $\bar{\nu}_\mu$  with sufficiently long paths in the Earth and obtained information about the Earth density distribution, which, although not very precise, broadly agrees with the PREM model. More specifically, in [33] it is assumed that the Earth density distribution is spherically symmetric. The analysis is performed with a five layer Earth model: the inner core, two equal width layers of the outer core and two equal width layers of the mantle. The densities in each of the five layers are varied independently. The external constraints on the Earth total mass  $M_\oplus$  and moment of inertia  $I_\oplus$ , which are known with a remarkable high precision [44–46], as well as the Earth hydrostatic equilibrium (EHE) constraint, were not applied. The results are obtained with the IceCube data on the zenith angle dependence of the fluxes of up-going atmospheric  $\nu_\mu$  and  $\bar{\nu}_\mu$  with energies in the interval  $E = (1.5 - 20.0)$  TeV [47]. Four different models of the initial fluxes of atmospheric  $\nu_\mu$  and  $\bar{\nu}_\mu$  were used in the analysis. The value of the Earth mass found in [33],  $M_\oplus^\nu = (6.0_{-1.3}^{+1.6}) \times 10^{24}$  kg, is in good agreement with gravitationally determined value [44, 45],  $M_\oplus = (5.9722 \pm 0.0006) \times 10^{24}$  kg. Thus, the Earth was “weighted” with neutrinos. The results obtained in [33] contain evidence at  $2\sigma$  C.L. that the core is denser than the mantle:  $\bar{\rho}_c^\nu(3\text{layer}) - \bar{\rho}_{man}^\nu(2\text{layer}) = (13.1_{-6.3}^{+5.8})$  g/cm<sup>3</sup>, where  $\bar{\rho}_c^\nu(3\text{layer})$  and  $\bar{\rho}_{man}^\nu(2\text{layer})$  are the

---

<sup>1</sup>More precisely, they depend on the electron number density  $N_e(r)$ ,  $N_e(r) = \rho_{rmE}(r)Y_e(r)/m_N$ ,  $Y_e(r)$  and  $m_N$  being the electron fractions number and nucleon mass. It follows from the studies of the Earth matter composition that  $Y_e \cong (0.490 - 0.0496)$  in the mantle and  $Y_e \cong (0.466 - 0.471)$  in the core (see, e.g., [29–32]). Taking into account these uncertainties has no effect on the results on the densities of the mantle and the core obtained in neutrino tomography studies (see, e.g., [33]). The relation between  $N_e(r)$  and  $\rho_{rmE}(r)$  can be used to obtain information about  $Y_e(r)$ , i.e., the composition of the core and mantle layers of the Earth (see, e.g., [34, 35]).

values of the average core and mantle densities determined in [33]. This was the first time the study of neutrinos traversing the Earth provided information of the Earth interior and marked the beginning of real experimental data driven neutrino tomography of the Earth.

The neutrino oscillation tomography of the Earth with IceCube, ORCA, DUNE and INO detectors is discussed in [34,35,48–58] (the early studies include [59,60] briefly described in [52]). In order to estimate the sensitivity of a given neutrino detector to possible deviations of the inner core (IC), outer core (OC), core (IC + OC) and mantle Earth densities from those obtained using geophysical and seismological data and described by PREM, in the statistical analyses performed within the neutrino tomography studies one typically varies the density of each of these structures. These variations, however, must respect the  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints. In the studies performed so far only in [53] all three  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints were taken into account. The  $M_{\oplus}$  and EHE constraints, but not the  $I_{\oplus}$  one, were implemented in the analyses performed in [52, 57].

The existing studies provide very limited information about the implications, and thus on the relevance, if any, of the three external constraints -  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE - in the neutrino tomography studies of the Earth density distribution. The aim of the present letter is to assess the importance of the three constraints in the neutrino tomography studies of the Earth interior. Working with PREM average densities we derive the  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints on the possible density variations in two illustrative and widely considered cases, namely, when one approximates the Earth density structure with i) three layers - mantle, outer core and inner core, and ii) four layers - upper mantle, lower mantle, outer core and inner core.

The article is organised as follows. In Section 2 we derive the contributions of the “surface” layers, the mantle, outer core and inner core to the Earth total mass and moment of inertia in terms of average densities of these layers calculated using the corresponding PREM density distributions. In Section 3 we implement the  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints considering uniform ( $r$ -independent) deviations of the average densities of the Earth layers from their respective PREM values. In sub-sections 3.1 and 3.2 We consider the cases when one approximates the Earth density structure with i) three layers - mantle, outer core and inner core, and ii) four layers - upper mantle, lower mantle, outer core and inner core. We obtain the equations which the deviation parameters should satisfy and provide the analytic solutions of these equations. We further comment on the implications of the results obtained. Section 4 contains a brief summary.

## 2 Earth Total Mass and Moment of Inertia: the Mantle, Outer Core, Inner Core and “Surface” Layers Contributions

In order to estimate the sensitivity of any detector to the IC, OC, core (IC + OC) and mantle densities, one typically varies the density in one of these structures - the structure of interest. These variations, however, must respect the constraints following from the precise

knowledge of the total Earth mass,  $M_{\oplus}$ , and momentum of inertia,  $I_{\oplus}$  [44–46]:

$$M_{\oplus} = (5.9722 \pm 0.0006) \times 10^{24} \text{ kg}, \quad (1)$$

$$I_{\oplus} = (8.01736 \pm 0.00097) \times 10^{37} \text{ kg m}^2. \quad (2)$$

The mean value of  $I_{\oplus}$  can be related to the Earth mean total mass and radius  $R_{\oplus}$ :

$$I_{\oplus} = C_{G(\text{PREM})} M_{\oplus} R_{\oplus}^2. \quad (3)$$

where  $C_G$  ( $C_{\text{PREM}}$ ) is a constant determined in gravity experiments (the PREM model):

$$C_G = 0.330745, \quad C_{\text{PREM}} = 0.33080. \quad (4)$$

One can implement the total Earth mass and moment of inertia constraints by compensating the variation of the density in the structure of interest by corresponding change of the densities in at least two of the other structures.

Following PREM, we will assume that the Earth density distribution  $\rho_E$  is spherically symmetric,  $\rho_E = \rho_E(r)$ , where  $r$  is the distance from the Earth center. The Earth density structure consists of inner core, outer core, mantle and three relatively thin layers close to the Earth surface - the ocean and two crust “surface” layers, having masses  $M_{\text{IC}}$ ,  $M_{\text{OC}}$ ,  $M_{\text{man}}$  and  $M_{\text{o+c1+c2}}$ , and moments of inertia  $I_{\text{IC}}$ ,  $I_{\text{OC}}$ ,  $I_{\text{man}}$  and  $I_{\text{o+c1+c2}}$ , respectively. Correspondingly, the total Earth mass in PREM is given by:

$$\begin{aligned} M_{\oplus} &= M_{\text{IC}} + M_{\text{OC}} + M_{\text{man}} + M_{\text{o+c1+c2}} = \int_0^{R_{\oplus}} 4\pi \rho_E(r) r^2 dr \\ &= 4\pi \left[ \int_0^{R_{\text{IC}}} \rho_{\text{IC}}(r) r^2 dr + \int_{R_{\text{IC}}}^{R_{\text{C}}} \rho_{\text{OC}}(r) r^2 dr + \int_{R_{\text{C}}}^{R_{\text{man}}} \rho_{\text{man}}(r) r^2 dr + \int_{R_{\text{man}}}^{R_{\oplus}} \rho_{\text{o+c1+c2}}(r) r^2 dr \right] \end{aligned} \quad (5)$$

where  $\rho_{\text{IC}}(r)$ ,  $\rho_{\text{OC}}(r)$ ,  $\rho_{\text{mant}}(r)$  and  $\rho_{\text{o+c1+c2}}(r)$  are the inner core (IC), outer core (OC), the mantle and the “surface” layers densities as a function of  $r$  and, according to PREM,

$$R_{\text{IC}} = 1221.5 \text{ km}, \quad R_{\text{OC}} = R_{\text{C}} = 3480 \text{ km}, \quad R_{\text{man}} = 6346.6 \text{ km}, \quad R_{\oplus} = 6371 \text{ km}. \quad (6)$$

Note that  $R_{\text{man}}$  is smaller than the mean radius of the Earth  $R_{\oplus}$  due to the presence in PREM of an “ocean” and two crust “surface” layers.

Similarly, the Earth total moment of inertia can be written as:

$$\begin{aligned} I_{\oplus} &= I_{\text{IC}} + I_{\text{OC}} + I_{\text{man}} + I_{\text{o+c1+c2}} = \int_0^{R_{\oplus}} \frac{8\pi}{3} \rho_E(r) r^4 dr \\ &= \frac{8\pi}{3} \left[ \int_0^{R_{\text{IC}}} \rho_{\text{IC}}(r) r^4 dr + \int_{R_{\text{IC}}}^{R_{\text{C}}} \rho_{\text{OC}}(r) r^4 dr + \int_{R_{\text{C}}}^{R_{\text{man}}} \rho_{\text{man}}(r) r^4 dr + \int_{R_{\text{man}}}^{R_{\oplus}} \rho_{\text{o+c1+c2}}(r) r^4 dr \right] \end{aligned} \quad (7)$$

It follows from the seismological and geophysical data that in order for Earth to be in hydrostatic equilibrium the following inequalities should always hold:

$$\rho_{\text{man}} < \rho_{\text{OC}} < \rho_{\text{IC}}. \quad (8)$$

We will refer to these inequalities as Earth hydrostatic equilibrium (EHE) constraint. The EHE constraint should also be implemented in the studies of the Earth density structure using the method of neutrino tomography.

In our further analysis we will use the the IC, OC, mantle and the “surface” layers density distributions given by PREM. In PREM, the mantle is represented by five layers with different distributions in each layer. In addition, as we have already indicated, there are three thin layers with constant density close to the Earth’s surface: one “ocean” and two crust “surface” layers.

In PREM the IC and OC density distributions have the form:

$$\rho_{\text{IC}}(r) = 13.0885 - 8.8381 \frac{r^2}{R_{\oplus}^2}, \quad r = (0 - 1221.5) \text{ km}; \quad (9)$$

$$\rho_{\text{OC}}(r) = 12.5815 - 1.2638 \frac{r}{R_{\oplus}} - 3.6426 \frac{r^2}{R_{\oplus}^2} - 5.5281 \frac{r^3}{R_{\oplus}^3}, \quad r = (1221.5 - 3480.0) \text{ km} \quad (10)$$

where  $\rho_{\text{IC}}(r)$  and  $\rho_{\text{OC}}(r)$  are in units of  $\text{g/cm}^3$  and we have given also for convenience the “widths” of the IC and OC layers. Similarly, the densities (in units of  $\text{g/cm}^3$ ) and the widths of the ocean, the two crust and the five mantle layers read [10]:

$$\rho_{\text{ocean}}(r) = 1.020, \quad r = (6368 - 6371) \text{ km}; \quad (11)$$

$$\rho_{\text{crust1}}(r) = 2.600, \quad r = (6356 - 6368) \text{ km};$$

$$\rho_{\text{crust2}}(r) = 2.900, \quad r = (6346.6 - 6356.0) \text{ km}; \quad (12)$$

$$\rho_{\text{m1}}(r) = 2.6910 + 0.6924 \frac{r}{R_{\oplus}}, \quad r = (6151 - 6346.6) \text{ km};$$

$$\rho_{\text{m2}}(r) = 7.1089 - 3.8045 \frac{r}{R_{\oplus}}, \quad r = (5971 - 6151) \text{ km};$$

$$\rho_{\text{m3}}(r) = 11.2494 - 8.0298 \frac{r}{R_{\oplus}}, \quad r = (5771 - 5971) \text{ km};$$

$$\rho_{\text{m4}}(r) = 5.3197 - 1.4836 \frac{r}{R_{\oplus}}, \quad r = (5701 - 5771) \text{ km};$$

$$\rho_{\text{m5}}(r) = 7.9565 - 6.4761 \frac{r}{R_{\oplus}} + 5.5283 \frac{r^2}{R_{\oplus}^2} - 3.0807 \frac{r^3}{R_{\oplus}^3}, \quad r = (3480 - 5701) \text{ km}. \quad (13)$$

Using Eqs. (5) and (7) - (13), it is not difficult to obtain the contributions of the IC, OC, the mantle and “surface” layers in  $M_{\oplus}$  and  $I_{\oplus}$ . For the contributions of the IC and OC

layers,  $M_{IC}$ ,  $I_{IC}$  and  $M_{OC}$ ,  $I_{OC}$ , we get:

$$M_{IC} = \frac{4\pi}{3} \bar{\rho}_{IC} R_{IC}^3, \quad (14)$$

$$I_{IC} = \frac{2}{5} \frac{4\pi}{3} \bar{\rho}_{IC}^I R_{IC}^5 = \frac{2}{5} \frac{\bar{\rho}_{IC}^I}{\bar{\rho}_{IC}} M_{IC} R_{IC}^2, \quad (15)$$

$$M_{OC} = \frac{4\pi}{3} \bar{\rho}_{OC} (R_C^3 - R_{IC}^3), \quad R_C = R_{OC}, \quad (16)$$

$$I_{OC} = \frac{2}{5} \frac{4\pi}{3} \bar{\rho}_{OC}^I (R_C^5 - R_{IC}^5) = \frac{2}{5} \frac{\bar{\rho}_{OC}^I}{\bar{\rho}_{OC}} M_{OC} \frac{R_C^5 - R_{IC}^5}{R_C^3 - R_{IC}^3}, \quad (17)$$

where  $\bar{\rho}_{IC}$ ,  $\bar{\rho}_{IC}^I$ ,  $\bar{\rho}_{OC}$  and  $\bar{\rho}_{OC}^I$  are average densities,

$$\bar{\rho}_{IC} = \frac{4\pi \int_0^{R_{IC}} \rho_{IC}(r) r^2 dr}{4\pi \int_0^{R_{IC}} r^2 dr} = 13.0885 - 8.8381 \frac{3R_{IO}^2}{5R_{\oplus}^2}, \quad (18)$$

$$\bar{\rho}_{IC}^I = \frac{\frac{8\pi}{3} \int_0^{R_{IC}} \rho_{IC}(r) r^4 dr}{\frac{8\pi}{3} \int_0^{R_{IC}} r^4 dr} = 13.0885 - 8.8381 \frac{5R_{IC}^2}{7R_{\oplus}^2}, \quad (19)$$

$$\begin{aligned} \bar{\rho}_{OC} &= \frac{4\pi \int_{R_{IC}}^{R_C} \rho_{OC}(r) r^2 dr}{4\pi \int_{R_{IC}}^{R_C} r^2 dr} \\ &= 12.5815 - 1.2638 \frac{3(R_{OC}^4 - R_{IO}^4)}{4R_{\oplus}(R_{OC}^3 - R_{IO}^3)} - 3.6426 \frac{3(R_{OC}^5 - R_{IO}^5)}{5R_{\oplus}^2(R_{OC}^3 - R_{IO}^3)} - 5.5281 \frac{3(R_{OC}^6 - R_{IO}^6)}{6R_{\oplus}^3(R_{OC}^3 - R_{IO}^3)} \end{aligned} \quad (20)$$

$$\begin{aligned} \bar{\rho}_{OC}^I &= \frac{\frac{8\pi}{3} \int_{R_{IC}}^{R_C} \rho_{OC}(r) r^4 dr}{\frac{8\pi}{3} \int_{R_{IC}}^{R_C} r^4 dr} \\ &= 12.5815 - 1.2638 \frac{5(R_{OC}^6 - R_{IO}^6)}{6R_{\oplus}(R_{OC}^5 - R_{IO}^5)} - 3.6426 \frac{5(R_{OC}^7 - R_{IO}^7)}{7R_{\oplus}^2(R_{OC}^5 - R_{IO}^5)} - 5.5281 \frac{5(R_{OC}^8 - R_{IO}^8)}{8R_{\oplus}^3(R_{OC}^5 - R_{IO}^5)} \end{aligned} \quad (21)$$

Using the values of  $R_{IC}$ ,  $R_{OC}$  and  $R_{\oplus}$  we find:

$$\bar{\rho}_{IC} = 12.8936 \frac{g}{cm^3}, \quad \bar{\rho}_{IC}^I = 12.8564 \frac{g}{cm^3} = 0.9971 \bar{\rho}_{IC}, \quad (22)$$

$$\bar{\rho}_{OC} = 10.90070 \frac{g}{cm^3}, \quad \bar{\rho}_{OC}^I = 10.6583 \frac{g}{cm^3} = 0.9778 \bar{\rho}_{OC}. \quad (23)$$

Note that  $\bar{\rho}_{IC}^I$  and  $\bar{\rho}_{OC}^I$  differ respectively from  $\bar{\rho}_{IC}$  and  $\bar{\rho}_{OC}$ , being somewhat smaller.

We can consider also the the total mass and moment of inertia of the Earth core:

$$M_C = M_{IC} + M_{OC} = \frac{4\pi}{3} \bar{\rho}_C R_C^3, \quad (24)$$

$$I_C = I_{IC} + I_{OC} = \frac{2}{5} \frac{4\pi}{3} \bar{\rho}_C^I R_C^5 = \frac{2}{5} \frac{\bar{\rho}_C^I}{\bar{\rho}_C} M_C R_C^2, \quad (25)$$

where

$$\bar{\rho}_C = \bar{\rho}_{IC} \frac{R_{IC}^3}{R_C^3} + \bar{\rho}_{OC} \left( 1 - \frac{R_{IC}^3}{R_C^3} \right) = 10.9869 \frac{g}{cm^3}, \quad (26)$$

$$\bar{\rho}_C^I = \bar{\rho}_{IC}^I \frac{R_{IC}^5}{R_C^5} + \bar{\rho}_{OC}^I \left( 1 - \frac{R_{IC}^5}{R_C^5} \right) = 10.6700 \frac{g}{cm^3} = 0.9712 \bar{\rho}_C. \quad (27)$$

It is not difficult to calculate in a similar way the contributions of the five mantle layers to  $M_\oplus$  and  $I_\oplus$ . They can be written in the form:

$$M_{\text{man}} = \frac{4\pi}{3} \bar{\rho}_{\text{man}} (R_{\text{man}}^3 - R_C^3), \quad \bar{\rho}_{\text{man}} = \frac{4\pi \int_{R_C}^{R_{\text{man}}} \rho_{\text{man}}(r) r^2 dr}{4\pi \int_{R_C}^{R_{\text{man}}} r^2 dr} \quad (28)$$

$$I_{\text{man}} = \frac{8\pi}{15} \bar{\rho}_{\text{man}}^I (R_{\text{man}}^5 - R_C^5) = \frac{2}{5} \frac{\bar{\rho}_{\text{man}}^I}{\bar{\rho}_{\text{man}}} M_{\text{man}} \frac{R_{\text{man}}^5 - R_C^5}{R_{\text{man}}^3 - R_C^3}, \quad \bar{\rho}_{\text{man}}^I = \frac{\frac{8\pi}{3} \int_{R_C}^{R_{\text{man}}} \rho_{\text{man}}(r) r^4 dr}{\frac{8\pi}{3} \int_{R_C}^{R_{\text{man}}} r^4 dr} \quad (29)$$

For the average densities  $\bar{\rho}_{\text{man}}$  and  $\bar{\rho}_{\text{man}}^I$  we get:

$$\bar{\rho}_{\text{man}} = 4.6651 \frac{g}{cm^3}, \quad \bar{\rho}_{\text{man}}^I = 4.2833 \frac{g}{cm^3} = 0.9182 \bar{\rho}_{\text{man}}. \quad (30)$$

We remark that if we include the ‘‘surface’’ layers in the mantle, integrating from  $R_C = 3480$  km to  $R_\oplus = 6371$  km we obtain:  $\bar{\rho}_{\text{man}} = 4.4551$  g/cm<sup>3</sup>,  $\bar{\rho}_{\text{man}}^I = 4.2481$  g/cm<sup>3</sup> = 0.9535  $\bar{\rho}_{\text{man}}$ .

All results derived above are within the PREM framework. For the Earth total mass and moment on inertia in this framework we have:  $M_\oplus = M_{o+c1+c2} + M_{\text{man}} + M_C$  and  $I_\oplus = I_{o+c1+c2} + I_{\text{man}} + I_C$ , where  $M_C = M_{IC} + M_{OC}$ ,  $I_C = I_{IC} + I_{OC}$  and  $M_{o+c1+c2}$  and  $I_{o+c1+c2}$  are the contributions from the ocean and the two crust ‘‘surface’’ layers. Using the PREM results shown in Eqs. (9) - (13), we find  $M_{o+c1+c2} \ll M_{\text{man}} + M_C$  and  $I_{o+c1+c2} \ll I_{\text{man}} + I_C$ :  $M_{o+c1+c2}/(M_{\text{man}} + M_C) \cong 5 \times 10^{-3}$  and  $I_{o+c1+c2}/(I_{\text{man}} + I_C) \cong 0.01060$ .

We note that the ‘‘surface’’ layers play subleading (if not negligible) role in neutrino tomography of the Earth due to their relatively small widths and contributions to  $M_\oplus$  and  $I_\oplus$ . As far as  $M_{o+c1+c2}$  and  $I_{o+c1+c2}$  are kept constant, they do not play any role in our analysis and the results we will obtain are independent of the values of  $M_{o+c1+c2}$  and  $I_{o+c1+c2}$ , as it will become clear further. We will keep them for consistency. For this purpose we introduce



their average densities  $\bar{\rho}_{oc1c2}$  and  $\bar{\rho}_{oc1c2}^I$ :

$$M_{o+c1+c2} = \frac{4\pi}{3} \bar{\rho}_{oc1c2} (R_{\oplus}^3 - R_{\text{man}}^3),$$

$$\bar{\rho}_{oc1c2} = \frac{4\pi \left[ \int_{R_{\text{ocean}}}^{R_{\oplus}} \rho_{\text{ocean}}(r) r^2 dr + \int_{R_{\text{crust1}}}^{R_{\text{ocean}}} \rho_{\text{crust1}}(r) r^2 dr + \int_{R_{\text{crust2}}}^{R_{\text{crust1}}} \rho_{\text{crust2}}(r) r^2 dr \right]}{4\pi \int_{R_{\text{man}}}^{R_{\oplus}} r^2 dr}, \quad (31)$$

$$I_{o+c1+c2} = \frac{8\pi}{15} \bar{\rho}_{oc1c2}^I (R_{\oplus}^5 - R_{\text{man}}^5),$$

$$\bar{\rho}_{oc1c2}^I = \frac{\frac{8\pi}{3} \left[ \int_{R_{\text{ocean}}}^{R_{\oplus}} \rho_{\text{ocean}}(r) r^4 dr + \int_{R_{\text{crust1}}}^{R_{\text{ocean}}} \rho_{\text{crust1}}(r) r^4 dr + \int_{R_{\text{crust2}}}^{R_{\text{crust1}}} \rho_{\text{crust2}}(r) r^4 dr \right]}{\frac{8\pi}{3} \int_{R_{\text{man}}}^{R_{\oplus}} r^4 dr}, \quad (32)$$

where  $R_{\text{ocean}} = 6368$  km,  $R_{\text{crust1}} = 6356$  km and  $R_{\text{crust2}} \equiv R_{\text{man}} = 6346.6$  km. For  $\bar{\rho}_{oc1c2}$  and  $\bar{\rho}_{oc1c2}^I$  we get:

$$\bar{\rho}_{oc1c2} = 2.5204 \frac{g}{\text{cm}^3}, \quad \bar{\rho}_{oc1c2}^I = 2.5194 \frac{g}{\text{cm}^3} = 0.9996 \bar{\rho}_{oc1c2}. \quad (33)$$

Summarising the results obtained so far we have:

$$M_{\oplus} = M_{\text{man}} + M_{\text{OC}} + M_{\text{IC}} + M_{o+c1+c2}$$

$$= \frac{4\pi}{3} R_{\oplus}^3 \left[ \bar{\rho}_{\text{man}} \frac{R_{\text{man}}^3}{R_{\oplus}^3} \left( 1 - \frac{R_{\text{C}}^3}{R_{\text{man}}^3} \right) + \bar{\rho}_{\text{OC}} \frac{R_{\text{C}}^3}{R_{\oplus}^3} \left( 1 - \frac{R_{\text{IC}}^3}{R_{\text{OC}}^3} \right) + \bar{\rho}_{\text{IC}} \frac{R_{\text{IC}}^3}{R_{\oplus}^3} + \bar{\rho}_{oc1c2} \left( 1 - \frac{R_{\text{man}}^3}{R_{\oplus}^3} \right) \right] \quad (34)$$

$$I_{\oplus} = I_{\text{man}} + I_{\text{OC}} + I_{\text{IC}} + I_{o+c1+c2}$$

$$= \frac{2}{5} \frac{4\pi}{3} R_{\oplus}^5 \left[ \bar{\rho}_{\text{man}}^I \frac{R_{\text{man}}^5}{R_{\oplus}^5} \left( 1 - \frac{R_{\text{C}}^5}{R_{\text{man}}^5} \right) + \bar{\rho}_{\text{OC}}^I \frac{R_{\text{C}}^5}{R_{\oplus}^5} \left( 1 - \frac{R_{\text{IC}}^5}{R_{\text{OC}}^5} \right) + \bar{\rho}_{\text{IC}}^I \frac{R_{\text{IC}}^5}{R_{\oplus}^5} + \bar{\rho}_{oc1c2}^I \left( 1 - \frac{R_{\text{man}}^5}{R_{\oplus}^5} \right) \right] \quad (35)$$

In terms of mantle and total core densities the expressions for  $M_{\oplus}$  and  $I_{\oplus}$  read:

$$M_{\oplus} = M_{\text{man}} + M_{\text{C}} + M_{o+c1+c2}$$

$$= \frac{4\pi}{3} R_{\oplus}^3 \left[ \bar{\rho}_{\text{man}} \frac{R_{\text{man}}^3}{R_{\oplus}^3} \left( 1 - \frac{R_{\text{C}}^3}{R_{\text{man}}^3} \right) + \bar{\rho}_{\text{C}} \frac{R_{\text{C}}^3}{R_{\oplus}^3} + \bar{\rho}_{oc1c2} \left( 1 - \frac{R_{\text{man}}^3}{R_{\oplus}^3} \right) \right], \quad (36)$$

$$I_{\oplus} = I_{\text{man}} + I_{\text{C}} + I_{o+c1+c2}$$

$$= \frac{2}{5} \frac{4\pi}{3} R_{\oplus}^5 \left[ \bar{\rho}_{\text{man}}^I \frac{R_{\text{man}}^5}{R_{\oplus}^5} \left( 1 - \frac{R_{\text{C}}^5}{R_{\text{man}}^5} \right) + \bar{\rho}_{\text{C}}^I \frac{R_{\text{C}}^5}{R_{\oplus}^5} + \bar{\rho}_{oc1c2}^I \left( 1 - \frac{R_{\text{man}}^5}{R_{\oplus}^5} \right) \right], \quad (37)$$

where we have used Eqs. (24) - (27) and (31) - (32).

It proves convenient to express the constants  $\bar{\rho}_i^{(1)}$  in Eqs. (35) and (37),  $i = \text{IC}, \text{OC}, \text{C}, \text{man}, \text{oc1c2}$ , as  $\bar{\rho}_i^{(1)} = r_i \bar{\rho}_i$ . For the ratios  $r_i$  we have:

$$\begin{aligned} r_{\text{man}} &= \frac{\bar{\rho}_{\text{man}}^{\text{I}}}{\bar{\rho}_{\text{man}}} = 0.9182, \quad r_{\text{OC}} = \frac{\bar{\rho}_{\text{OC}}^{\text{I}}}{\bar{\rho}_{\text{OC}}} = 0.9778, \quad r_{\text{IC}} = \frac{\bar{\rho}_{\text{IC}}^{\text{I}}}{\bar{\rho}_{\text{IC}}} = 0.9971, \\ r_{\text{C}} &= \frac{\bar{\rho}_{\text{C}}^{\text{I}}}{\bar{\rho}_{\text{C}}} = 0.9712, \quad r_{\text{oc1c2}} = \frac{\bar{\rho}_{\text{oc1c2}}^{\text{I}}}{\bar{\rho}_{\text{oc1c2}}} = 0.9996. \end{aligned} \quad (38)$$

The values of these ratios are given in Eqs. (22), (23), (27), (30) and (33).

It also proves convenient to introduce the average Earth density:

$$\bar{\rho}_{\oplus} \equiv \frac{3}{4\pi} \frac{M_{\oplus}}{R_{\oplus}^3} = a_{\text{man}} \bar{\rho}_{\text{man}} + a_{\text{OC}} \bar{\rho}_{\text{OC}} + a_{\text{IC}} \bar{\rho}_{\text{IC}} + a_{\text{oc1c2}} \bar{\rho}_{\text{oc1c2}} \quad (39)$$

$$= a_{\text{man}} \bar{\rho}_{\text{man}} + a_{\text{C}} \bar{\rho}_{\text{C}} + a_{\text{oc1c2}} \bar{\rho}_{\text{oc1c2}}. \quad (40)$$

where

$$\begin{aligned} a_{\text{man}} &= \frac{R_{\text{man}}^3}{R_{\oplus}^3} \left(1 - \frac{R_{\text{C}}^3}{R_{\text{man}}^3}\right) \cong 0.8256, \quad a_{\text{OC}} = \frac{R_{\text{C}}^3}{R_{\oplus}^3} \left(1 - \frac{R_{\text{IC}}^3}{R_{\text{C}}^3}\right) \cong 0.15592, \quad a_{\text{C}} = \frac{R_{\text{C}}^3}{R_{\oplus}^3} \cong 0.1639, \\ a_{\text{IC}} &= \frac{R_{\text{IC}}^3}{R_{\oplus}^3} \cong 7.0479 \times 10^{-3}. \quad a_{\text{oc1c2}} = \left(1 - \frac{R_{\text{man}}^3}{R_{\oplus}^3}\right) \cong 0.0114. \end{aligned} \quad (41)$$

Replacing  $\bar{\rho}_i^{(1)}$  with  $r_i \bar{\rho}_i$  in Eqs. (35) and (37),  $i = \text{IC}, \text{OC}, \text{C}, \text{man}, \text{oc1c2}$ , and using Eq. (39) we get  $I_{\oplus} = C M_{\oplus} R_{\oplus}^2$ , which implies:

$$\frac{I_{\oplus}}{R_{\oplus}^5} \frac{5}{2} \frac{3}{4\pi} = C' \bar{\rho}_{\oplus} = b_{\text{man}} \bar{\rho}_{\text{man}} + b_{\text{OC}} \bar{\rho}_{\text{OC}} + b_{\text{IC}} \bar{\rho}_{\text{IC}} + b_{\text{oc1c2}} \bar{\rho}_{\text{oc1c2}} \quad (42)$$

$$= b_{\text{man}} \bar{\rho}_{\text{man}} + b_{\text{C}} \bar{\rho}_{\text{C}} + b_{\text{oc1c2}} \bar{\rho}_{\text{oc1c2}} \quad (43)$$

where  $C'$  is a constant,  $C' = 5C/2 \cong 0.8270$  and

$$\begin{aligned} b_{\text{man}} &= r_{\text{man}} \frac{R_{\text{man}}^5}{R_{\oplus}^5} \left(1 - \frac{R_{\text{C}}^5}{R_{\text{man}}^5}\right), \quad b_{\text{OC}} = r_{\text{OC}} \frac{R_{\text{C}}^5}{R_{\oplus}^5} \left(1 - \frac{R_{\text{IC}}^5}{R_{\text{C}}^5}\right), \quad b_{\text{C}} = r_{\text{C}} \frac{R_{\text{C}}^5}{R_{\oplus}^5}, \\ b_{\text{IC}} &= r_{\text{IC}} \frac{R_{\text{IC}}^5}{R_{\oplus}^5}, \quad b_{\text{oc1c2}} = r_{\text{oc1c2}} \left(1 - \frac{R_{\text{man}}^5}{R_{\oplus}^5}\right). \end{aligned} \quad (44)$$

The constant parameters  $r_{\text{man}}, r_{\text{OC}}, r_{\text{IC}}$  and  $r_{\text{oc1c2}}$  are defined and their values are given in Eq. (38). They are all close to, but different from, 1. The numerical values of  $b_{\text{man}}, b_{\text{OC}}, b_{\text{C}}, b_{\text{IC}}$  and  $b_{\text{oc1c2}}$  read:

$$b_{\text{man}} \cong 0.8561, \quad b_{\text{OC}} \cong 0.0473, \quad b_{\text{C}} \cong 0.0472, \quad b_{\text{IC}} \cong 2.58 \times 10^{-4}, \quad b_{\text{oc1c2}} \cong 0.0190. \quad (45)$$

The left-hand sides of Eqs. (39), (40), (42) and (43) are fixed since  $M_{\oplus}, I_{\oplus}$  and  $R_{\oplus}$  are known with relatively high precision. Thus, within the aspects of neutrino tomography of

the Earth we are considering, Eqs. (39), (40) and (42), (43) represent the constraints on the variations of  $\bar{\rho}_{\text{man}}$ , and  $\bar{\rho}_{\text{OC}}$ ,  $\bar{\rho}_{\text{IC}}$  or  $\bar{\rho}_C$ , from the knowledge of  $M_{\oplus}$  and  $I_{\oplus}$ .

We will assume that  $R_{\text{man}}$ ,  $R_{\text{OC}}$  and  $R_{\text{IC}}$  and, correspondingly, the values of the constants  $r_{\text{man}}$ ,  $r_{\text{OC}}$  and  $r_{\text{IC}}$  are correctly given by PREM. These are standard assumptions made in the neutrino tomography of the Earth studies of possible deviations of  $\bar{\rho}_{\text{man}}$ ,  $\bar{\rho}_{\text{OC}}$  and  $\bar{\rho}_{\text{IC}}$  from their PREM values. In our further analyses we will consider the ‘‘surface’’ terms in Eqs. (39) and (42),  $a_{\text{oc1c2}} \bar{\rho}_{\text{oc1c2}}$  and  $b_{\text{oc1c2}} \bar{\rho}_{\text{oc1c2}}$ , as fixed. Under these conditions, Eqs. (39) and (42) represent two equations for the three densities of interest:  $\bar{\rho}_{\text{man}}$ ,  $\bar{\rho}_{\text{OC}}$  and  $\bar{\rho}_{\text{IC}}$ .

### 3 Implementing the $M_{\oplus}$ , $I_{\oplus}$ and EHE Constraints

#### 3.1 The Mantle, Outer Core and Inner Core Case

Suppose one is interested in obtaining information about possible deviations of the densities of the mantle  $\rho_{\text{man}}$ , the outer core,  $\rho_{\text{OC}}$ , and the inner core,  $\rho_{\text{IC}}$  from their PREM values. In the corresponding statistical analysis one varies the density in the chosen layer of interest, say  $\rho_{\text{OC}}$ . In order to satisfy the  $M_{\oplus}$  and  $I_{\oplus}$  constraints one has to compensate the variation of  $\rho_{\text{OC}}$  by an appropriate change of the densities in the other two layers,  $\rho_{\text{man}}$  and  $\rho_{\text{IC}}$ . In the analyses performed so far it is assumed that the indicated variations and changes of interest are by constant  $r$ -independent factors  $\kappa_i$ , which can be positive, negative or zero:

$$\rho'_i = (1 + \kappa_i) \rho_i, \quad i = \text{man}, \text{OC}, \text{IC}. \quad (46)$$

We will analyse this possibility below.

The total Earth mass and momentum of inertia constraints in the simple case we are considering of average densities imply that  $\rho'_i$  should satisfy Eqs. (39) and (42) and the following two equations:

$$\begin{aligned} a_{\text{man}} (1 + \kappa_{\text{man}}) \bar{\rho}_{\text{man}} + a_{\text{OC}} (1 + \kappa_{\text{OC}}) \bar{\rho}_{\text{OC}} + a_{\text{IC}} (1 + \kappa_{\text{IC}}) \bar{\rho}_{\text{IC}} + a_{\text{oc1c2}} \bar{\rho}_{\text{oc1c2}} &= \bar{\rho}_{\oplus}, \\ b_{\text{man}} (1 + \kappa_{\text{man}}) \bar{\rho}_{\text{man}} + b_{\text{OC}} (1 + \kappa_{\text{OC}}) \bar{\rho}_{\text{OC}} + b_{\text{IC}} (1 + \kappa_{\text{IC}}) \bar{\rho}_{\text{IC}} + b_{\text{oc1c2}} \bar{\rho}_{\text{oc1c2}} &= C' \bar{\rho}_{\oplus}. \end{aligned} \quad (47)$$

Combining these equations with Eqs. (39) and (42) and taking into account that the terms  $a_{\text{oc1c2}} \bar{\rho}_{\text{oc1c2}}$  and  $b_{\text{oc1c2}} \bar{\rho}_{\text{oc1c2}}$  are fixed, we get neglecting the uncertainties in the determination of  $M_{\oplus}$  and  $I_{\oplus}$ :

$$\begin{aligned} a_{\text{man}} \kappa_{\text{man}} \bar{\rho}_{\text{man}} + a_{\text{OC}} \kappa_{\text{OC}} \bar{\rho}_{\text{OC}} + a_{\text{IC}} \kappa_{\text{IC}} \bar{\rho}_{\text{IC}} &= 0, \\ b_{\text{man}} \kappa_{\text{man}} \bar{\rho}_{\text{man}} + b_{\text{OC}} \kappa_{\text{OC}} \bar{\rho}_{\text{OC}} + b_{\text{IC}} \kappa_{\text{IC}} \bar{\rho}_{\text{IC}} &= 0. \end{aligned} \quad (48)$$

Suppose we vary  $\rho_{\text{OC}}$  and are interested in the sensitivity of a given detector to possible deviations of  $\rho'_{\text{OC}}$  from the PREM prediction, i.e., to values of  $\kappa_{\text{OC}} \neq 0$ . In the simple case of average densities we are considering we can express  $\kappa_{\text{man}}$  and  $\kappa_{\text{IC}}$  in terms of  $\kappa_{\text{OC}}$  using

Eq. (48):

$$\begin{aligned}\kappa_{\text{man}} &= \frac{\kappa_{\text{OC}}}{D_{\text{OC}}} (a'_{\text{IC}} b_{\text{OC}} - a_{\text{OC}} b'_{\text{IC}}) , \\ \kappa_{\text{IC}} &= \frac{\kappa_{\text{OC}}}{D_{\text{OC}}} (a_{\text{OC}} b'_{\text{man}} - a'_{\text{man}} b_{\text{OC}}) .\end{aligned}\quad (49)$$

Here

$$D_{\text{OC}} \equiv a'_{\text{man}} b'_{\text{IC}} - a'_{\text{IC}} b'_{\text{man}} , \quad (50)$$

and

$$a'_j \equiv \frac{\bar{\rho}_j}{\bar{\rho}_{\text{OC}}} a_j , \quad b'_j \equiv \frac{\bar{\rho}_j}{\bar{\rho}_{\text{OC}}} b_j , \quad j = \text{man}, \text{IC} . \quad (51)$$

It follows, in particular, from Eq. (49) that in order to satisfy the Earth total mass and moment of inertia constraints, the variations of  $\bar{\rho}_{\text{OC}}$  have to be compensated by changes of the densities of at least two other Earth layers. Using the PREM values of  $\bar{\rho}_{\text{IC}}$ ,  $\bar{\rho}_{\text{OC}}$  and  $\bar{\rho}_{\text{man}}$  given in Eqs. (22), (23) and (30), as well as the values of  $a_i$  (in Eq. (41)) and  $b_i$  (in Eq. (44)) we find:

$$\kappa_{\text{man}} \cong -0.1104 \kappa_{\text{OC}} , \quad \kappa_{\text{IC}} \cong -13.9558 \kappa_{\text{OC}} . \quad (52)$$

In the case of  $\kappa_{\text{OC}} > 0$  the relevant Earth hydrostatic equilibrium (EHE) constraint is  $\bar{\rho}_{\text{OC}}(1 + \kappa_{\text{OC}}) < \bar{\rho}_{\text{IC}}(1 + \kappa_{\text{IC}})$ , with  $\kappa_{\text{IC}} < 0$ . Using the relation between  $\kappa_{\text{IC}}$  and  $\kappa_{\text{OC}}$  given in Eq. (52), we find:

$$\kappa_{\text{OC}} < \frac{\bar{\rho}_{\text{IC}} - \bar{\rho}_{\text{OC}}}{\bar{\rho}_{\text{OC}} + 13.9558 \bar{\rho}_{\text{IC}}} \cong 0.0104 , \quad (53)$$

where we have used the PREM values of  $\bar{\rho}_{\text{IC}}$  and  $\bar{\rho}_{\text{OC}}$  (see Eq. (22)).

For  $\kappa_{\text{OC}} < 0$  the relevant EHE constraint is instead  $\bar{\rho}_{\text{man}}(1 + \kappa_{\text{man}}) < \bar{\rho}_{\text{OC}}(1 - |\kappa_{\text{OC}}|)$ . This inequality together with the relation between  $\kappa_{\text{man}}$  and  $\kappa_{\text{OC}}$  we have derived and the PREM values of  $\bar{\rho}_{\text{man}}$  and  $\bar{\rho}_{\text{OC}}$  leads to the following lower limit:

$$\kappa_{\text{OC}} > -\frac{\bar{\rho}_{\text{OC}} - \bar{\rho}_{\text{man}}}{\bar{\rho}_{\text{OC}} + 0.1104 \bar{\rho}_{\text{man}}} \cong -0.546 , \quad \kappa_{\text{OC}} < 0 . \quad (54)$$

However, the obtained lower limit on  $\kappa_{\text{OC}} < 0$  allows excessively large values of  $\kappa_{\text{IC}} > 0$  and thus of  $\bar{\rho}_{\text{IC}}(1 + \kappa_{\text{IC}})$ , as large as, e.g., 7.6 times bigger than the PREM value of  $\bar{\rho}_{\text{IC}}$ . The existing seismological data on  $\bar{\rho}_{\text{IC}}$  is claimed to have an uncertainty not exceeding  $\sim 10\%$ . Even if we allow a rather large maximum deviation of  $\bar{\rho}_{\text{IC}}$  from its PREM value of 25% (50%), which corresponds to  $\kappa_{\text{IC}} \leq 0.25$  (0.50), this leads to the following lower limit on  $\kappa_{\text{OC}} < 0$ :  $\kappa_{\text{OC}} \gtrsim -0.018$  (0.036).

Thus, we see that in the considered case of three layer Earth density distribution, the Earth mass and moment of inertia constraints together with the requirement of the Earth being in hydrostatic equilibrium and the value of the inner core average density lead to severe constraints on the possible uniform deviations of  $\bar{\rho}_{\text{OC}}$  from its PREM value:

$$-0.018 \text{ (0.036)} \lesssim \kappa_{\text{OC}} < 0.010 . \quad (55)$$

These very restrictive constraints can be traced to the relatively small contributions of the inner core to  $M_\oplus$  and especially to  $I_\oplus$ .

We note that we get practically the same results if we consider the “surface” layers as part of the mantle. In this case the mantle extends from  $R_C = 3480$  km to  $R_\oplus = 6371$  km (instead of  $R_{\text{man}} = 6346.6$  km).

In this analysis we did not take into account the uncertainties in the determination of  $M_\oplus$  and  $I_\oplus$ . As it follows from Eqs. (1) and (2), the relative  $1\sigma$  uncertainties in the values of  $M_\oplus$  and  $I_\oplus$  are  $\sim 10^{-4}$ . These uncertainties are too small to have any significant effect, if taken into account, on the results we have obtained.

### 3.2 The case of Two Mantle Layers, Outer Core and Inner Core

We consider next the case of two different mantle layers and the already discussed outer core, inner core and “surface” layers. The mantle is divided into i) lower mantle, extending from  $R_c = 3480$  km to  $R_{\text{lman}} = 5701$  km, with densities  $\bar{\rho}_{\text{lman}}$  and  $\bar{\rho}_{\text{lman}}^I$ , and ii) upper mantle, extending from  $R_{\text{lman}} = 5701$  km to  $R_{\text{man}} = 6346.6$  km, with densities  $\bar{\rho}_{\text{uman}}$  and  $\bar{\rho}_{\text{uman}}^I$ . It is not difficult to calculate the PREM values of  $\bar{\rho}_{\text{lman}}$ ,  $\bar{\rho}_{\text{lman}}^I$ ,  $\bar{\rho}_{\text{uman}}$  and  $\bar{\rho}_{\text{uman}}^I$ :

$$\bar{\rho}_{\text{lman}} = 4.9035 \frac{g}{\text{cm}^3}, \quad \bar{\rho}_{\text{lman}}^I = r_{\text{lman}} \bar{\rho}_{\text{lman}}, \quad r_{\text{lman}} = 0.9771, \quad (56)$$

$$\bar{\rho}_{\text{uman}} = 3.6046 \frac{g}{\text{cm}^3}, \quad \bar{\rho}_{\text{uman}}^I = r_{\text{uman}} \bar{\rho}_{\text{uman}}, \quad r_{\text{uman}} = 0.9425. \quad (57)$$

As in the preceding subsection we will assume that the variations and changes of  $\bar{\rho}_{\text{IC}}$ ,  $\bar{\rho}_{\text{OC}}$ ,  $\bar{\rho}_{\text{lman}}$  and  $\bar{\rho}_{\text{uman}}$  are by constant  $r$ -independent factors  $\kappa_{\text{IC}}$ ,  $\kappa_{\text{OC}}$ ,  $\kappa_{\text{lman}}$  and  $\kappa_{\text{uman}}$ :

$$\rho'_i = (1 + \kappa_i) \rho_i, \quad i = \text{IC}, \text{OC}, \text{lman}, \text{uman}. \quad (58)$$

The analogs of the equations in Eq. (48) now read:

$$\begin{aligned} a_{\text{lman}} \kappa_{\text{lman}} \bar{\rho}_{\text{lman}} + a_{\text{uman}} \kappa_{\text{uman}} \bar{\rho}_{\text{uman}} + a_{\text{OC}} \kappa_{\text{OC}} \bar{\rho}_{\text{OC}} + a_{\text{IC}} \kappa_{\text{IC}} \bar{\rho}_{\text{IC}} &= 0, \\ b_{\text{lman}} \kappa_{\text{lman}} \bar{\rho}_{\text{lman}} + b_{\text{uman}} \kappa_{\text{uman}} \bar{\rho}_{\text{uman}} + b_{\text{OC}} \kappa_{\text{OC}} \bar{\rho}_{\text{OC}} + b_{\text{IC}} \kappa_{\text{IC}} \bar{\rho}_{\text{IC}} &= 0. \end{aligned} \quad (59)$$

For the new parameters in these equations we find:

$$\begin{aligned} a_{\text{lman}} &= \frac{R_{\text{lman}}^3}{R_\oplus^3} \left(1 - \frac{R_C^3}{R_{\text{lman}}^3}\right) \cong 0.5535, \quad a_{\text{uman}} = \frac{R_{\text{man}}^3}{R_\oplus^3} \left(1 - \frac{R_{\text{lman}}^3}{R_{\text{man}}^3}\right) \cong 0.2720, \quad (60) \\ b_{\text{lman}} &= r_{\text{lman}} \frac{R_{\text{lman}}^5}{R_\oplus^5} \left(1 - \frac{R_c^5}{R_{\text{lman}}^5}\right) \cong 0.5131, \quad b_{\text{uman}} = r_{\text{uman}} \frac{R_{\text{man}}^5}{R_\oplus^5} \left(1 - \frac{R_{\text{lman}}^5}{R_{\text{man}}^5}\right) \cong 0.3838 \end{aligned} \quad (61)$$

where we have used the values of  $R_C$ ,  $R_{\text{lman}}$ ,  $R_{\text{man}}$ ,  $R_\oplus$ ,  $r_{\text{lman}}$  and  $r_{\text{uman}}$ .

Equations (59) represent the constraints following from the knowledge of  $M_\oplus$  and  $I_\oplus$  on the the possible deviations of  $\bar{\rho}_{\text{IC}}$ ,  $\bar{\rho}_{\text{OC}}$ ,  $\bar{\rho}_{\text{lman}}$  and  $\bar{\rho}_{\text{uman}}$  from their PREM values.

Suppose we vary  $\rho_{\text{OC}}$  and are interested in the sensitivity of a given detector to possible deviations  $\kappa_{\text{OC}} \neq 0$  of the PREM prediction for  $\rho_{\text{OC}}$ . Consider the simple case of  $\kappa_{\text{IC}} = 0$ .

Working with the average densities and using Eq. (59) we can express  $\kappa_{lman}$  and  $\kappa_{uman}$  in terms of  $\kappa_{OC}$ :

$$\begin{aligned}\kappa_{lman} &= \frac{\kappa_{OC}}{D_{OC}^{(m)}} (a'_{uman} b_{OC} - a_{OC} b'_{uman}) , \\ \kappa_{uman} &= \frac{\kappa_{OC}}{D_{OC}^{(m)}} (a_{OC} b'_{lman} - a'_{lman} b_{OC}) .\end{aligned}\quad (62)$$

Here

$$D_{OC}^{(m)} \equiv a'_{lman} b'_{uman} - a'_{uman} b'_{lman} , \quad (63)$$

and

$$a'_j \equiv \frac{\bar{\rho}_j}{\bar{\rho}_{OC}} a_j , \quad b'_j \equiv \frac{\bar{\rho}_j}{\bar{\rho}_{OC}} b_j , \quad j = lman, uman . \quad (64)$$

Using the PREM values of  $\bar{\rho}_{OC}$ ,  $\bar{\rho}_{lman}$  and  $\bar{\rho}_{uman}$  given in Eqs. (23) and (56) and (57), as well as the values of  $a_i$  and  $b_i$ ,  $i = OC, lman, uman$ , from Eqs. (41), (45), (60) and (61), we get:

$$\kappa_{lman} = f_{lman} \kappa_{OC} , \quad f_{lman} \cong -1.4327; \quad \kappa_{uman} = f_{uman} \kappa_{OC} , \quad f_{uman} \cong 2.2329 . \quad (65)$$

The EHE constraint for  $\kappa_{OC} > 0$  is  $\bar{\rho}_{OC}(1 + \kappa_{OC}) < \bar{\rho}_{IC}$  (recall that we set  $\kappa_{IC} = 0$ ). Using the PREM values of  $\bar{\rho}_{IC}$  and  $\bar{\rho}_{OC}$  (see Eq. (22)) we find:

$$\kappa_{OC} < \frac{\bar{\rho}_{IC} - \bar{\rho}_{OC}}{\bar{\rho}_{OC}} \cong 0.18 . \quad (66)$$

In the case of  $\kappa_{OC} < 0$  the EHE constraint reads:  $\bar{\rho}'_{man} < \bar{\rho}_{OC}(1 - |\kappa_{OC}|)$ , where  $\bar{\rho}'_{man}$  is the modified mantle density due to the changes of the lower and upper mantle densities  $\bar{\rho}_{lman}(1 + \kappa_{lman})$  and  $\bar{\rho}_{uman}(1 + \kappa_{uman})$ . It is not difficult to show that

$$\bar{\rho}'_{man} = \bar{\rho}_{lman}(1 + \kappa_{lman}) C_{lman} + \bar{\rho}_{uman}(1 + \kappa_{uman}) C_{uman} , \quad (67)$$

where

$$C_{lman} = \frac{R_{lman}^3}{R_{man}^3} \left( 1 - \frac{R_C^3}{R_{man}^3} \right) \cong 0.6705 , \quad C_{uman} = \frac{1 - \frac{R_{lman}^3}{R_{man}^3}}{1 - \frac{R_C^3}{R_{man}^3}} \cong 0.3295 , \quad (68)$$

where we have used the values of  $R_{lman}$ ,  $R_{man}$  and  $R_C$ . The EHE constraint in the considered case has the form:

$$\kappa_{OC} > - \frac{\bar{\rho}_{OC} - \bar{\rho}_{lman} C_{lman} - \bar{\rho}_{uman} C_{uman}}{\bar{\rho}_{OC} + \bar{\rho}_{lman} C_{lman} f_{lman} + \bar{\rho}_{uman} C_{uman} f_{uman}} , \quad \kappa_{OC} < 0 . \quad (69)$$

Using the values of  $\bar{\rho}_{OC}$ ,  $\bar{\rho}_{lman}$ ,  $\bar{\rho}_{uman}$ ,  $f_{lman}$ ,  $f_{uman}$ ,  $C_{lman}$  and  $C_{uman}$  given in Eqs. (23), (56), (57), (65) and (68) we get:

$$\kappa_{OC} > -0.73 . \quad (70)$$

Combining the two limits we have:

$$-0.73 < \kappa_{OC} < 0.18. \quad (71)$$

Practically the same results are obtained if we include the “surface” layers in the upper mantle. In this case the “surface” layers and the upper mantle form one “upper mantle” layer, which extends from  $R_{\text{lman}} = 3701$  km to  $R_{\oplus} = 6371$  km (instead of  $R_{\text{man}} = 6346.6$  km). Similarly, the results change insignificantly if one joins the inner core and the outer core in one core layer.

Thus, we see that in the case of two mantle, outer and inner core (or one core) layers, the  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints on the possible variation of  $\bar{\rho}_{OC}$  are much less restrictive than in the case of one mantle, outer and inner core layers, considered in the preceding subsection. The main reason for this difference is that the contributions of the inner core to  $M_{\oplus}$  and  $I_{\oplus}$  are not crucial for satisfying the  $M_{\oplus}$  and  $I_{\oplus}$  constraints when there are two mantle layers, while, being relatively small, they are critical in the one mantle layer case.

One may consider other layer structures of the Earth density distribution, for example, inner core, two equal width outer core layers and mantle layer. We did not performed the numerical analysis in this case but expect the  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints to be relevant but much less restrictive than in the case of the three layer case we have considered.

## 4 Summary and Conclusions

Summarising, we have shown that the impact of the Earth total mass,  $M_{\oplus}$ , moment of inertia,  $I_{\oplus}$ , and the Earth hydrostatic equilibrium (EHE) constraints in the neutrino tomography studies of the Earth density structure depends crucially on the Earth layer structure assumed in the studies. More specifically, we find that if one approximates the Earth density distribution with three layers - inner core, outer core and mantle, the  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints lead to severe limits on the possible uniform deviations of the outer core and mantle densities from their PREM values even if one assumes that the inner core density, which plays a crucial role in satisfying the constraints, can be by 50% larger than its PREM density. We have shown also that, in contrast, the  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints on the possible deviations of the outer core and mantle densities from their PREM value are much less restrictive if one assumes a four layer Earth density structure consisting of inner core, outer core, and two - upper and lower - mantle layers. In this case the inner core density plays an insignificant (or no) role in satisfying the constraints.

It follows from our results that the implementation of  $M_{\oplus}$ ,  $I_{\oplus}$  and EHE constraints in the neutrino tomography studies of the Earth density structure can have a very strong impact on the conclusions reached in the studies.

## Acknowledgements

This work was supported in part by the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 860881-HIDDeN,

by the INFN program on Theoretical Astroparticle Physics and by the World Premier International Research Center Initiative (WPI Initiative, MEXT), Japan.

## References

- [1] W.F. McDonough, “Treatise on Geochemistry: The Mantle and Core”, vol. 2 (ed. R. W. Carlson, Elsevier-Pergamon, Oxford, 2003), p. 547.
- [2] B. A. Bolt, Q. J. R. Astron. Soc. **32**, 367 (1991).
- [3] B. L. N. Kennett, Geophys. J. Int. **132** (1998) 374.
- [4] G. Masters and D. Gubbins, Phys. Earth Planet. Inter. **140** (2003) 159.
- [5] F. Nimmo, talk given at the International Workshop “Multi-Messenger Tomography of the Earth”, Snowbird, Utah, U.S.A., July 30-31, 2022.
- [6] K.D. Koper, talk given at the International Workshop “Multi-Messenger Tomography of the Earth”, Snowbird, Utah, U.S.A., July 30-31, 2022.
- [7] L. Waszek, talk given at the International Workshop “Multi-Messenger Tomography of the Earth”, APC, Paris, July 4-7, 2023.
- [8] W.F. McDonough, talk given at the International Workshop “Multi-Messenger Tomography of the Earth”, APC, Paris, July 4-7, 2023.
- [9] S. Noe, talk given at the International Workshop “Multi-Messenger Tomography of the Earth”, APC, Paris, July 4-7, 2023.
- [10] A. M. Dziewonski and D. L. Anderson, Phys. Earth Planet. Interiors **25** (1981) 297.
- [11] C. F. Yoder, “Global Earth Physics”, vol. 1, (ed. T. J. Ahrens, American Geophysical Union, Washington DC, 1995), p. 1.
- [12] W. F. McDonough and R. Arevalo, J. Phys. Conf. Ser. **136** (2008), 022006 (doi:10.1088/1742-6596/136/2/022006).
- [13] B.A. Baffet, H.E. Huppert, J.R. Lister and A.W. Woods, Submitted to Nature 1991.
- [14] A. Placci and E. Zavattini, “On the possibility of using high-energy neutrinos to study the Earth’s interior”, <https://cds.cern.ch/record/2258764> (1973), CERN Report.
- [15] L. V. Volkova and G. T. Zatsepin, Izv. Akad. Nauk Ser. Fiz. **38N5** (1974) 1060.
- [16] I. P. Nedyalkov, preprint JINR (Dubna), JINR-P2-81-645, 1981.
- [17] I. P. Nedyalkov, Bolgarskaia Akademiia Nauk **34** (1981) 177.



- [18] A. De Rujula, S. L. Glashow, R. R. Wilson, G. Charpak, Phys. Rept. **99** (1983) 341.
- [19] T. L. Wilson, Nature **309** (1984) 38.
- [20] G. A. Askar'yan, Usp. Fiz. Nauk **144** (1984) 523 [Sov. Phys. Usp. **27** (1984) 896].
- [21] A. B. Borisov, B. A. Dolgoshein, A. N. Kalinovsky, Yad. Fiz. **44** (1986) 681.
- [22] A. B. Borisov, B. A. Dolgoshein, Phys. Atom. Nucl. **56** (1993) 755.
- [23] W. Winter, Earth Moon Planets **99** (2006) 285.
- [24] C. Kuo, *et al.*, Earth Plan. Sci. Lett. **133** (1995) 95.
- [25] P. Jain, J. P. Ralston, G. M. Frichter, Astropart. Phys. **12** (1999) 193.
- [26] M. M. Reynoso, O. A. Sampayo, Astropart. Phys. **21** (2004) 315.
- [27] M. C. Gonzalez-Garcia, F. Halzen, M. Maltoni and H. K. M. Tanaka, Phys. Rev. Lett. **100** (2008) 061802 [arXiv:0711.0745 [hep-ph]].
- [28] M. Tanabashi *et al.* [Particle Data Group], Phys. Rev. D **98** (2018) 030001 (doi:10.1103/PhysRevD.98.030001). See therein the review “Neutrino Masses, Mixing and Oscillations” by K. Nakamura and S.T. Petcov.
- [29] J. Badro *et al.*, Proc. Natl. Acad. Sci. U. S. A. **112(40)** (2015) 12310-12314.
- [30] E. Kaminski and M. Javoy, Earth Plan. Sci. Lett. **365** (2013) 97-107.
- [31] T. Sakamaki *et al.*, Earth and Planetary Science Letters **287** (2009) 293-297.
- [32] <https://earthref.org/GERMRD/datamodel/> (cit. on pp. 114, 115).
- [33] A. Donini, S. Palomares-Ruiz and J. Salvado, Nature Phys. **15** (2019) 37 [arXiv:1803.05901 [hep-ph]].
- [34] C. Rott, A. Taketa and D. Bose, Sci. Rep. **5** (2015) 15225 [arXiv:1502.04930 [physics.geo-ph]].
- [35] S. Bourret, J. Coelho, E. Kaminski and V. Van Elewyck, PoS **ICRC2019** (2020), 1024 doi:10.22323/1.358.1024
- [36] T. K. Gaisser and M. Honda, Ann. Rev. Nucl. Part. Sci. **52** (2002) 153 [arXiv:hep-ph/0203272].
- [37] M. Honda, M. Sajjad Athar, T. Kajita, K. Kasahara and S. Midorikawa, Phys. Rev. D **92** (2015) 023004 [arXiv:1502.03916].
- [38] R. Abbasi *et al.* [IceCube Collab.], Astropart. Phys. **35** (2012) 615; M.G. Aartsen *et al.* [IceCube Collab.], Phys. Rev. Lett. **120** (2018) 071801; P. Eller *et al.* [IceCube], PoS **ICRC2023** (2023) 1036 [arXiv:2307.15295].

- [39] J.A. Sanchez [IceCube Collab], talk given at the International Conference on Neutrino Physics and Astrophysics, June 17-22, 2024, Milano, Italy.
- [40] S. Adrian-Martinez *et al.* [KM3Net], “Letter of intent for KM3NeT 2.0,” *J. Phys. G* **43** (2016) 084001 [arXiv:1601.07459].
- [41] K. Abe *et al.* [Hyper-Kamiokande], “Hyper-Kamiokande Design Report,” arXiv:1805.04163;  
M. Ishitsuka [on behalf of the Hyper-Kamiokande Proto-Collab.], talk given at the XXIX Int. Conference on Neutrino Physics and Astrophysics, Chicago, June 22 - July 2, 2020 (virtual conference), DOI:10.5281/zenodo.3959585.
- [42] B. Abi *et al.* [DUNE], [arXiv:1807.10334 [physics.ins-det]].
- [43] S. Ahmed *et al.* [ICAL], *Pramana* **88** (2017) 79 [arXiv:1505.07380].
- [44] B. Luzum *et al.*, *Celest. Mech. Phys.* **110** (2011) 110.
- [45] H.USAO, USNO, UKHO, “The Astronomical Almanac”, <http://asa.usno.navy.mil/>.
- [46] W. Chen, J. Ray, W.B. Shen and C.L. Huang, *J. Geod.* **89** (2015) 179.
- [47] M. G. Aartsen *et al.* [IceCube], *Phys. Rev. Lett.* **117** (2016) 071801 [arXiv:1605.01990].
- [48] S. K. Agarwalla, T. Li, O. Mena and S. Palomares-Ruiz, [arXiv:1212.2238].
- [49] W. Winter, *Nucl. Phys. B* **908** (2016), 250-267 doi:10.1016/j.nuclphysb.2016.03.033 [arXiv:1511.05154 [hep-ph]].
- [50] S. Bourret *et al.* [KM3NeT], *J. Phys. Conf. Ser.* **888** (2017) no.1, 012114 doi:10.1088/1742-6596/888/1/012114 [arXiv:1702.03723 [physics.ins-det]].
- [51] A. Kumar and S.Kumar Agarwalla, [arXiv:2104.11740].
- [52] F. Capozzi and S.T. Petcov, *Eur. Phys. J. C* **82** (2022) 461 [arXiv:2111.13048].
- [53] K.J. Kelly, P.A.N. Machado, I. Martinez-Soler and Y.F. Perez-Gonzalez, “DUNE atmospheric neutrinos: Earth tomography,” *JHEP* **05** (2022) 187 [arXiv:2110.00003].
- [54] P. B. Denton and R. Pestes, *Phys. Rev. D* **104** (2021) 113007 [arXiv:2110.01148].
- [55] L. Maderer, E. Kaminski, J.A.B. Coelho, S. Bourret and V. Van Elewyck, *Front. Earth Sci.* **11** (2023) 1008396 [arXiv:2208.00532].
- [56] A.K. Upadhyay, A. Kumar, S.K. Agarwalla and A. Dighe, *JHEP* **04** (2023) 068 [arXiv:2211.08688].
- [57] D. Raikwal and S. Choubey, *Phys. Rev. D* **109** (2024) 073011 [arXiv:2309.12573].
- [58] A.K. Upadhyay, A. Kumar, S.K. Agarwalla and A. Dighe, [arXiv:2405.04986].

- [59] S. Choubey, P. Ghoshal and S.T. Petcov, studies performed in the period 2008 - 2011 for the 100 kt liquid Argon and 1 Mt water Cerenkov (Hyper-Kamiokande) detectors, unpublished.
- [60] S. Choubey and S.T. Petcov, studies performed in 2014 for the PINGU detector, unpublished.