

Effects of quantum corrections to Lorentzian vacuum transitions in the presence of gravity

H. García-Compeán^{a,1}, J. Hernández-Aguilar^{b,2}, D. Mata-Pacheco^{a,3},
C. Ramírez^{b4}

^a*Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN
P.O. Box 14-740, CP. 07000, Ciudad de México, México*

^b*Facultad de Ciencias Físico Matemáticas,
Benemérita Universidad Autónoma de Puebla
4 Sur 104, Puebla 72000, Puebla, México*

Abstract

We present a study of the vacuum transition probabilities taking into account quantum corrections. We first introduce a general method that expand previous works employing the Lorentzian formalism of the Wheeler-De Witt equation by considering higher order terms in the semiclassical expansion. The method presented is applicable in principle to any model in the superspace and up to any desired order in the quantum correction terms. Then, we apply this method to obtain analytical solutions for the probabilities up to second quantum corrections for homogeneous isotropic and anisotropic universes. We use the Friedmann-Lemaitre-Robertson-Walker with positive and zero curvature for the isotropic case and the Bianchi III and Kantowski-Sachs metrics for the anisotropic case. Interpreting the results as distribution probabilities of creating universes by vacuum decay with a given size, we found that the general behaviour is that considering up to the second quantum correction leads to an avoidance of the initial singularity. However, we show that this result can only be achieved for the isotropic universe. Furthermore, we also study the effect of anisotropy on the transition probabilities.

¹e-mail address: hugo.compean@cinvestav.mx

²e-mail address: jorge.hernandezag@alumno.buap.mx

³e-mail address: daniel.mata@cinvestav.mx

⁴e-mail address: cramirez@fcfm.buap.mx

1 Introduction

Quantum tunnelling represents one of the first phenomenon that challenged the classical ideas and required a quantum theory for its proper understanding. It is a key process in many situations, in particular, in the canonical approach to quantum gravity it is thought to provide the description for the birth of our universe. Let us consider a general scenario where an scalar field is present with its corresponding potential with the general behaviour of containing two local minima of different value separated by a hill. Then we can always expect that a transition between the two minima can occur via quantum tunnelling. In field theory this is called a vacuum transition (analogously in the absence of an scalar field, the two minima can also be described in a simpler context with two different values of the cosmological constant). Since this setup appears in many scenarios, it has attracted a lot of attention over the decades. In particular, for field theory the study of such type of transitions began with the work of Sidney Coleman et. al. [1, 2] and it is described by the nucleation of true vacuum bubbles on a false vacuum background (the true vacuum in this context is located at the global minimum). Later on, by using a proposal for a quantum theory of gravity employing the path integral approach, these transition were studied including the gravitational field by Coleman and De Luccia [3] and then by Parke [4]. Over the years, many new results have been explored using these Euclidean techniques (see for example [5–14]).

On the other hand, it is well known that an alternative description to the path integral approach of quantum gravity can be described by employing a Hamiltonian formalism. In this way, employing the Arnowitt, Deser and Misner (ADM) formalism [15, 16], instead of a path integral, the classical Hamiltonian and momentum constraints of a gravitational system are quantized using the Dirac quantization procedure, leading to a set of constraints on a wave function for the universe (for a general introduction to the subject of quantum cosmology see [17–21]). In particular, when we are dealing with homogeneous metrics, all the information is encoded in the equation resulting from the Hamiltonian constraint, that states that the Hamiltonian operator annihilates such wave function, this is called the Wheeler-De Witt equation [22, 23]. One important feature of this formalism is that it does not rely on a Wick rotation, thus it is a purely Lorentzian method. The interpretation of this wave function turned out to be very troublesome, since there can not be a notion of an external observer (see for example [24, 25] for recent proposals and discussions on this regard). However, there are many ways in which we can use the solutions of the WDW equation to explore physical properties of the gravitational systems. In particular, the squared ratio of two solutions can be interpreted as the conditional probability between both configurations. Using this interpretation, the vacuum transition probabilities have also been explored in a Lorentzian formalism. First of all, Fischler, Morgan and Polchinski studied transition between two values of the cosmological constant in [26, 27]. Recently, such results were generalized in [28], one important feature of the Lorentzian formalism is that the end result can be a closed universe, contrary to the results obtained with the Euclidean approach. Later on, the inclusion of an scalar field was studied in [29]. One limitation of this formalism when dealing with the scalar field is that there is not a concrete description of bubble nucleation, however the transition probabilities

obtained can be interpreted as probability distributions of creating universes with a given size by vacuum decay, as a generalization of the standard tunnelling from nothing scenario. Based on these works, in [30] it was proposed a general method to compute such transition probabilities for any model in the superspace at the semiclassical level. In [31] such method was used to study the transitions in the Hořava-Lifshitz theory of gravity, then in [32] it was generalized to consider lineal terms in the momenta in order to take into account the effect of a Generalized Uncertainty Principle.

Let us remark that all these studies provide results only at the semiclassical level. The incorporation of quantum corrections to the transition probabilities in the Euclidean formalism is troublesome since there appears a negative mode problem that may spoil the semiclassical approximation [33]. This issue has been studied rigorously over the years [34–38]. On the other hand, the Lorentzian formalism employs a semiclassical expansion in the form of a WKB proposal. In this way, the quantum corrections are incorporated by considering the higher order terms in the \hbar expansion. It has been shown for field theory that the first quantum correction computed in this form indeed coincides with the 1-loop contribution of the Euclidean formalism [39]. Thus, the main purpose of this article is to expand the general method presented in [29, 30] to incorporate quantum corrections to the transition probabilities with gravity in a general setup, in this way avoiding the problems presented in the Euclidean formalism.

Let us also remark that recently, a new method has been proposed (at the semiclassical level) in the form of a tunnelling potential in which many of the Euclidean results can be derived [40–44]. Furthermore, holographic interpretation of these transitions have also been explored [45–48]. In this way, our results may be relevant for these proposals as well.

Moreover, although the Friedmann-Lemaître-Robertson-Walker (FLRW) metric describing a homogeneous and isotropic universe is the the natural and most used choice to study these transitions given its cosmological importance, it is also relevant to study homogeneous but anisotropic metrics. This is justified by the fact that inflation is thought to erase any signal of anisotropy on the early universe, but since we will be dealing in this regime, such metrics should be relevant. In addition, there are some recent experimental studies that suggest that such metrics may be important for the description of our universe at the present epoch as well [49, 50]. Therefore, in this article we will apply the general method to study both isotropic and anisotropic homogeneous metrics.

The outline of this article is as follows: In Section 2 we will present the general method to compute the transition probabilities with quantum corrections for any model on the superspace starting with a generic form of the Hamiltonian constraint. We will apply the method to study the transition probabilities with quantum corrections terms (up to second order in powers of \hbar) in the following sections. First, we will study the isotropy case with the closed FLRW metric in Section 3 and the flat FLRW metric in Section 4. Then, we will consider anisotropic metrics, in Section 5 we will consider the Bianchi III metric and compare the results with the flat FLRW result, which corresponds to its isotropy limit. Then, in Section 6 we will use the Kantowski-Sachs metric. Finally, in Section 7 we will present our Final Remarks and in Appendix A, we will discuss the dependence of the results on the choice of factorization for the Bianchi

III and flat FLRW metrics which is an important subject to remark.

2 Transition probabilities with quantum corrections

In this section, we begin by expanding the general method proposed in [29,30] to compute transition probabilities in a general scenario. We will consider higher order terms in the WKB expansion for the WDW equation to compute the quantum corrections to the transition probabilities. We will also consider transitions between a false and a true vacuum of a scalar field potential. We employ the notation and conventions of these references.

Let us begin by considering a general form for the Hamiltonian constraint in the context of the ADM formulation of general relativity given by

$$\mathcal{H} = \frac{1}{2}G^{MN}(\Phi)\pi_M\pi_N + f[\Phi] \simeq 0, \quad (1)$$

where G^{MN} denotes the inverse metric in Wheeler's superspace, Φ encompasses all superspace coordinates Φ^M (there can be an infinite number of them), these include the degrees of freedom of the three-dimensional metric, variables of the matter fields, etc., along with their respective canonically conjugate momenta π_M . Additionally, $f[\Phi]$ refers to all remaining terms related to the three-curvature, potential terms of matter fields or any other term that does not depend on the momenta. The WDW equation is obtained by performing a canonical quantization of the Hamiltonian constraint, specifically by promoting $\pi_M \rightarrow -i\hbar\frac{\delta}{\delta\Phi^M}$ in the last expression, leading to

$$\mathcal{H}\Psi(\Phi) = \left[-\frac{\hbar^2}{2}G^{MN}(\Phi)\frac{\delta}{\delta\Phi^M}\frac{\delta}{\delta\Phi^N} + f[\Phi] \right] \Psi[\Phi] = 0, \quad (2)$$

up to ordering ambiguities, where $\Psi[\Phi]$ is the wave functional in superspace. In order to obtain a semiclassical outcome and quantum corrections from this equation, we employ the general proposal of the WKB type

$$\Psi[\Phi] = \exp \left\{ \frac{i}{\hbar}S[\Phi] \right\} \quad (3)$$

with the \hbar -expansion

$$S[\Phi] = S_0[\Phi] + \hbar S_1[\Phi] + \hbar^2 S_2[\Phi] + \mathcal{O}(\hbar^3), \quad (4)$$

where S_0 is the classical action and S_1 and S_2 are the first and second quantum corrections respectively. Substituting this ansatz in the WDW equation (2) we obtain for the first three orders in \hbar

$$\frac{1}{2}G^{MN}\frac{\delta S_0}{\delta\Phi^M}\frac{\delta S_0}{\delta\Phi^N} + f[\Phi] = 0, \quad (5)$$

$$2G^{MN}\frac{\delta S_0}{\delta\Phi^M}\frac{\delta S_1}{\delta\Phi^N} = iG^{MN}\frac{\delta^2}{\delta\Phi^M\delta\Phi^N}S_0, \quad (6)$$

$$2G^{MN} \frac{\delta S_0}{\delta \Phi^M} \frac{\delta S_2}{\delta \Phi^N} + G^{MN} \frac{\delta S_1}{\delta \Phi^M} \frac{\delta S_1}{\delta \Phi^N} = iG^{MN} \frac{\delta^2 S_1}{\delta \Phi^M \delta \Phi^N}. \quad (7)$$

Let us consider now a set of integral curves on a particular slice on the space of fields defined with a parameter s in the form

$$C(s) \frac{d\Phi^M}{ds} = G^{MN} \frac{\delta S_0}{\delta \Phi^N}, \quad (8)$$

where $C(s)$ is an auxiliary function. Then, employing eqs. (5) and (8) the classical action can be written as

$$S_0[\Phi] = -2 \int^s \frac{ds'}{C(s')} \int_X f[\Phi(s')]. \quad (9)$$

Then, using (8), eq. (5) leads to the following form

$$G_{MN} \frac{d\Phi^M}{ds} \frac{d\Phi^N}{ds} = -\frac{2f[\Phi_s]}{C^2(s)}, \quad (10)$$

where G_{MN} is the inverse of G^{MN} . As we pointed out in [30] eqs. (8) and (10) form a system of $n+1$ equations for the $n+1$ variables: $\left(\frac{d\Phi^M}{ds}, C^2(s)\right)$ which in principle can be solved for any model in the superspace in order to obtain a solution for S_0 . In the following, we will assume that the fields Φ^M depend only on the time variable, then using eq. (9), the variational derivative in (8) can be expressed in terms of a partial derivative of the function $f[\Phi]$ and the integrals on the spatial slice will only give a factor of the volume of such slice $\text{Vol}(X)$.

In this way, considering the next order in the WKB expansion, we note that we can write

$$\frac{dS_1}{ds} = \int_X \frac{d\Phi^M}{ds} \frac{\delta S_1}{\partial \Phi^M}, \quad (11)$$

then, using (8), from eq. (6) we obtain

$$S_1 = \frac{i}{2} \int_s \int_X \frac{ds}{C(s)} G^{MN} \frac{\delta^2 S_0}{\delta \Phi^M \delta \Phi^N}. \quad (12)$$

Furthermore, from (9), it is clear that we can write

$$\frac{\delta S_0}{\delta \Phi^M} = -\frac{2 \text{Vol}(X)}{C(s)} \frac{\partial f}{\partial \Phi^M}, \quad (13)$$

and

$$\frac{\delta^2 S_0}{\delta \Phi^M \delta \Phi^N} = -\frac{2 \text{Vol}(X)}{C(s)} \frac{\partial^2 f}{\partial \Phi^M \partial \Phi^N}. \quad (14)$$

Thus it is found that the first quantum correction can be written as

$$S_1 = -i \text{Vol}^2(X) \int \frac{ds}{C^2(s)} \nabla^2 f, \quad (15)$$

where the laplacian is defined on the superspace, that is

$$\nabla^2 f = G^{MN} \frac{\partial^2 f}{\partial \Phi^M \partial \Phi^N}. \quad (16)$$

Likewise, following the same procedure using (7) we obtain that the second quantum correction can be written as

$$S_2 = \frac{1}{2} \int_s ds \frac{\text{Vol}^3(X)}{C^3(s)} \nabla^2 (\nabla^2 f) + \frac{1}{2} \int_s ds \frac{\text{Vol}^5(X)}{C^5(s)} (\nabla (\nabla^2 f))^2, \quad (17)$$

where we have denoted

$$(\nabla f)^2 = G^{MN} \frac{\partial f}{\partial \Phi^M} \frac{\partial f}{\partial \Phi^N}. \quad (18)$$

Thus, we note that once we have solved the system of equations, using (9), (15) and (17) we can have a solution to the WDW equation up to second order in the WKB expansion in general. Let us also remark that each term in the WKB expansion leads to an independent equation from the WDW equation (such as eqs. (5)-(7) for the first three terms), thus the system of equations will always have the same number of variables as equations. Then, in principle we can compute the transition probabilities up to any desired order. However, for simplicity, in the following we will keep only up to the second quantum correction.

As we remarked in [30] the system of equations can be solved in general as long as $\Phi^M = \Phi^M(s)$ is satisfied, yielding the solutions

$$C^2(s) = -\frac{2\text{Vol}(X)}{f} (\nabla f)^2, \quad \frac{d\Phi^M}{ds} = \frac{f}{\text{Vol}(X)} \frac{\nabla^M f}{(\nabla f)^2}, \quad (19)$$

where we rise indices with the superspace metric, that is

$$\nabla^M f = G^{MN} \frac{\partial f}{\partial \Phi^N}. \quad (20)$$

From eq. (19) we see that in general the various fields will be related by

$$\frac{d\Phi^M}{d\Phi^N} = \frac{\nabla^M f}{\nabla^N f}, \quad (21)$$

which is valid for every value of M and N such that $d\Phi^{M,N} \neq 0$. Therefore as long as all the expressions are different from zero, we will always be able to reduce the number of independent degrees of freedom to just one and change the integration variable from s to such field in (9), (15) and (17).

So far, we have only studied the way in which we can obtain a general solution from the WDW equation with up to second quantum correction terms. In order to compute the transition probabilities, following [29, 30] next it will be deemed a wave functional corresponding to a path in field space in which the scalar field evolves from the false minimum to the true one. We also discuss a system which is kept at the false minimum during the path, so we can consider that the squared ratio of the absolute

value of these wave functionals can be interpreted as the transition probability for the system to tunnel from the false to the true vacuum. Therefore, the transition tunnelling probability for going from the false vacuum at ϕ_A to the true vacuum at ϕ_B is the decay rate which is given by

$$P(A \rightarrow B) = \left| \frac{\Psi(\varphi_0^I, \phi_B; \varphi_m^I, \phi_A)}{\Psi(\varphi_0^I, \phi_A; \varphi_m^I, \phi_A)} \right|^2 = |e^{-\Gamma}|^2 = \exp[-2 \operatorname{Re}(\Gamma)], \quad (22)$$

where we have denoted as φ^I the remaining fields on superspace apart from the scalar field degree of freedom, $\Psi(\varphi_0^I, \phi_B; \varphi_m^I, \phi_A)$ is the wave functional corresponding to the path that starts in $\varphi^I(s=0) = \varphi_0^I$ with the scalar field ϕ_B and ends in $\varphi^I(s=s_M) = \varphi_m^I$ with the scalar field ϕ_A . Therefore in the WKB approach considered here we can write

$$\pm\Gamma = \frac{i}{\hbar} [S(\varphi_0^m, \phi_B; \varphi_M^m, \phi_A) - S(\varphi_0^m, \phi_A; \varphi_M^m, \phi_A)], \quad (23)$$

where the sign ambiguity appears due to the fact that the general solution of the wave functionals will be a linear superposition of exponential terms, however, we will keep only the dominant terms. Thus, using the expansion (4) we can write up to second order

$$\pm\Gamma = \Gamma_0 + \Gamma_1 + \Gamma_2, \quad (24)$$

where

$$\begin{aligned} \Gamma_0 &= \frac{i}{\hbar} [S_0(\varphi_0^m, \phi_B; \varphi_M^m, \phi_A) - S_0(\varphi_0^m, \phi_A; \varphi_M^m, \phi_A)], \\ \Gamma_1 &= i [S_1(\varphi_0^m, \phi_B; \varphi_M^m, \phi_A) - S_1(\varphi_0^m, \phi_A; \varphi_M^m, \phi_A)], \\ \Gamma_2 &= i\hbar [S_2(\varphi_0^m, \phi_B; \varphi_M^m, \phi_A) - S_2(\varphi_0^m, \phi_A; \varphi_M^m, \phi_A)]. \end{aligned} \quad (25)$$

In this case Γ_0 stands for the value that is calculated with the classical action, and it will be referred to as the semiclassical contribution, Γ_1 is the first quantum correction to the transition probabilities, and Γ_2 is the second quantum correction.

Let us remark that the concrete expression for the transition probabilities will depend on the specific choice of s . In this work we will choose the parameter s such that for the interval $[0, \bar{s} - \delta s]$, where $s=0$ is the initial value, the field remains close to its value at the true minimum ϕ_B , and for the interval $[\bar{s} + \delta s, s_M]$ the field remains very close to its value at the false minimum ϕ_A , then the scalar field takes the values

$$\phi(s) \approx \begin{cases} \phi_B, & 0 < s < \bar{s} - \delta s \\ \phi_A, & \bar{s} + \delta s < s < s_M. \end{cases} \quad (26)$$

Thus, we establish a configuration in which there is a bubble of true vacuum in the background of a false vacuum separated by a wall. Given the limitations of the minisuperspace approach in describing these transitions, this method remains our sole option for implementing this configuration. Furthermore, we will always consider the thin wall approximation in which $\delta s \rightarrow 0$. This choice for the s parameter allows to obtain the solutions from the semiclassical computations on the Euclidean approach as was shown in [29, 30].

With this choice, from (9) we obtain that the semiclassical contribution takes the form

$$\Gamma_0 = -\frac{2\text{Vol}(X)i}{\hbar} \left\{ \int_{s_0}^{\bar{s}-\delta s} \frac{ds}{C(s)} f \Big|_{\phi=\phi_B} - \int_{s_0}^{s-\delta s} \frac{ds}{C(s)} f \Big|_{\phi=\phi_A} + \int_{\bar{s}-\delta s}^{\bar{s}+\delta s} ds \left[\frac{f}{C(s)} - \frac{f}{C(s)} \Big|_{\phi=\phi_A} \right] \right\}. \quad (27)$$

From (15) the first quantum correction is given by

$$\Gamma_1 = \text{Vol}^2(X) \left[\int_{s_0}^{\bar{s}-\delta s} \frac{ds}{C^2(s)} \nabla^2 f \Big|_{\phi=\phi_B} - \int_{s_0}^{s-\delta s} \frac{ds}{C^2(s)} \nabla^2 f \Big|_{\phi=\phi_A} + \int_{\bar{s}-\delta s}^{\bar{s}+\delta s} ds \frac{1}{C^2(s)} \left(\nabla^2 f - \nabla^2 f \Big|_{\phi=\phi_A} \right) \right]. \quad (28)$$

Moreover from (17) the second quantum correction is written as

$$\begin{aligned} \Gamma_2 = & \frac{i\hbar}{2} \left[\int_{s_0}^{s-\delta s} ds \frac{\text{Vol}^3(X)}{C^3(s)} \nabla^2 (\nabla^2 f) \Big|_{\phi=\phi_B} + \int_{s_0}^{s-\delta s} ds \frac{\text{Vol}^5(X)}{C^5(s)} (\nabla (\nabla^2 f))^2 \Big|_{\phi=\phi_B} \right. \\ & - \int_{s_0}^{\bar{s}-\delta s} ds \frac{\text{Vol}^3(x)}{C^3(s)} \nabla^2 (\nabla^2 f) \Big|_{\phi=\phi_A} - \int_{s_0}^{\bar{s}-\delta s} ds \frac{\text{Vol}^5(x)}{C^5(s)} (\nabla (\nabla^2 f))^2 \Big|_{\phi=\phi_A} \\ & + \int_{\bar{s}-\delta s}^{\bar{s}+\delta s} ds \frac{\text{Vol}^3(X)}{C^3(s)} \left(\nabla^2 (\nabla^2 f) - \nabla^2 (\nabla^2 f) \Big|_{\phi=\phi_A} \right) \\ & \left. + \int_{\bar{s}-\delta s}^{\bar{s}+\delta s} ds \frac{\text{Vol}^5(x)}{C^5(s)} \left((\nabla (\nabla^2 f))^2 - (\nabla (\nabla^2 f))^2 \Big|_{\phi=\phi_A} \right) \right]. \quad (29) \end{aligned}$$

Therefore, with these expressions we can compute the transition probabilities up to second quantum correction for any model on superspace. Let us remark that we have only assumed a generic form of the Hamiltonian constraint (1), thus these expressions will be valid in principle for any metric and for any gravity theory that leads to a Hamiltonian constraint of this form, in the same way that the semiclassical treatment was valid for Hořava-Lifshitz gravity for example as shown in [31].

However, let us make some assumptions that will be relevant for this article in order to describe some general features of the quantum correction terms. In this work we will only consider General Relativity and the matter content of the system to be a scalar field canonically coupled to gravity with a potential $V(\phi)$ that has a false and a true minimum, thus the complete action is⁵

$$S = \frac{1}{2} \int d^4x \sqrt{-g} R - \int d^4x \sqrt{-g} \left[\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi + V(\phi) \right]. \quad (30)$$

This action will lead to a structure of the the Hamiltonian constraint which will allow us to recast the function f as

$$f = H(\varphi^I) + F(\varphi^I) V(\phi), \quad (31)$$

⁵Using natural units in which $c = 1$ and $8\pi G = 1$.

Furthermore, since the scalar field is canonically coupled, the superspace metric will satisfy $G^{M\phi} = 0$ for $M \neq \phi$. Moreover, for the same reason we can assume that $G^{\phi\phi} = G^{\phi\phi}(\varphi^I)$. In addition, as we pointed out earlier, due to the fact that all fields in superspace can be related through eq. (21), for the regions where the scalar field is constant we can always write $f = f(\Phi^i)$ for some fixed i . In this way, the general solutions (19) can be cast as

$$C(s) = \pm i \text{Vol}(X) \sqrt{\frac{2(\nabla f)^2}{f}}, \quad ds = \frac{\text{Vol}(X)(\nabla f)^2}{f \nabla^i f} d\Phi^i, \quad (32)$$

then we can change the integrals in s in the region of constant scalar field in (27), (28) and (29) to integrals in the chosen degree of freedom Φ^i .

Moreover, for the third term in (27) we note that on the region of the path where the scalar field is not a constant, we have the structure of a function that depends on the degrees of freedom of the three-metric times a function of the scalar field minus the same function evaluated at the false minimum. When dealing with the FLRW metric this term coincides with the tension term in the bubble as found in [29]. Moreover, in general we have dealt with these type of terms by defining constants (that we will call tension terms) accompanying the function that depends on the degrees of freedom of the metric evaluated at \bar{s} , as previously done in [30–32]. This will be valid only on the thin wall limit. Therefore, for the semiclassical contribution we define the tension term as

$$\text{Vol}(X) F|_{\bar{\Phi}^i} T_0 = -2 \text{Vol}(X) i \int_{\bar{s}-\delta s}^{\bar{s}+\delta s} \frac{F}{C(s)} (V - V_A). \quad (33)$$

Thus, employing the thin wall limit, we finally obtain that the semiclassical contribution is given by

$$\begin{aligned} \Gamma_0 = \mp \frac{\sqrt{2} \text{Vol}(X)}{\hbar} & \left[\int_{\Phi_0^i}^{\bar{\Phi}^i} \frac{\sqrt{f(\nabla f)^2}}{\nabla^i f} \Big|_{\phi=\phi_B} d\Phi^i - \int_{\Phi_0^i}^{\bar{\Phi}^i} \frac{\sqrt{f(\nabla f)^2}}{\nabla^i f} \Big|_{\phi=\phi_A} d\Phi^i \right] \\ & + \frac{\text{Vol}(X)}{\hbar} F \Big|_{\bar{\Phi}^i} T_0, \end{aligned} \quad (34)$$

where the sign ambiguity in the right hand side appears because of the general solution in (32) and will be independent of the sign ambiguity in the left hand side of the expression (24). This expression gives the correct semiclassical results for all the metrics considered in [30]. Furthermore, we note that in general, the transition probability depends on two parameters, in this case $\bar{\Phi}^i$ and T_0 that we will consider as independent. Moreover, we note that the volume of the spatial slice is an overall constant. Thus for metrics that do not have a finite value for this volume, we can always compactified the spatial slice and consider appropriate constant values.

Following the same procedure, for the first quantum correction we obtain in the

thin wall limit

$$\Gamma_1 = -\frac{\text{Vol}(X)}{2} \left[\int_{\Phi_0^i}^{\bar{\Phi}^i} \frac{\nabla^2 f}{\nabla^i f} \Big|_{\phi=\phi_B} d\Phi^i - \int_{\Phi_0^i}^{\bar{\Phi}^i} \frac{\nabla^2 f}{\nabla^i f} \Big|_{\phi=\phi_A} d\Phi^i \right] \quad (35)$$

$$+ \text{Vol}(X) \left[\nabla_R^2 F \Big|_{\bar{\Phi}^i} T_{1,1} + FG^{\phi\phi} \Big|_{\bar{\Phi}^i} T_{1,2} \right],$$

where $T_{1,1}$ and $T_{1,2}$ are tension terms defined analogously as in (33) but with their corresponding function of the scalar field. Furthermore, we have denoted the restricted laplacian as

$$\nabla_R^2 F = G^{IJ} \frac{\partial^2}{\partial \varphi^I \partial \varphi^J} F, \quad (36)$$

i.e. the laplacian restricted only to the fields that are not the scalar field. We note that if $\nabla_R^2 F = \zeta FG^{\phi\phi}$ for some constant ζ , or one of these functions is zero, the tension terms can be reduced to only one term. Therefore, we note that in general the second quantum correction will depend on three parameters, one coming from the metric, namely $\bar{\Phi}^i$ and two tension terms at most. Furthermore, we note that once again we obtain the global term $\text{Vol}(X)$, thus there is not any problem with metrics that contain noncompact spatial slices as well.

Finally, following the same procedure we obtain for the second quantum correction

$$\Gamma_2 = \mp \frac{\text{Vol}(X)\hbar}{4\sqrt{2}} \left[\int_{\Phi_0^i}^{\bar{\Phi}^i} \frac{\nabla^2(\nabla^2 f)}{\nabla^i f} \sqrt{\frac{f}{(\nabla f)^2}} \Big|_{\phi=\phi_B} d\Phi^i - \int_{\Phi_0^i}^{\bar{\Phi}^i} \frac{\nabla^2(\nabla^2 f)}{\nabla^i f} \sqrt{\frac{f}{(\nabla f)^2}} \Big|_{\phi=\phi_A} d\Phi^i \right]$$

$$\pm \frac{\text{Vol}(X)\hbar}{8\sqrt{2}} \left[\int_{\Phi_0^i}^{\bar{\Phi}^i} \frac{(\nabla(\nabla^2 f))^2}{\nabla^i f} \sqrt{\left(\frac{f}{(\nabla f)^2}\right)^3} \Big|_{\phi=\phi_B} d\Phi^i - \int_{\Phi_0^i}^{\bar{\Phi}^i} \frac{(\nabla(\nabla^2 f))^2}{\nabla^i f} \sqrt{\left(\frac{f}{(\nabla f)^2}\right)^3} \Big|_{\phi=\phi_A} d\Phi^i \right]$$

$$+ \text{Vol}(X)\hbar \left\{ \nabla_R^2(\nabla_R^2 F) \Big|_{\bar{\Phi}^i} T_{2,1} + [(\nabla_R^2 F)G^{\phi\phi} + \nabla_R^2(FG^{\phi\phi})] \Big|_{\bar{\Phi}^i} T_{2,2} + F(G^{\phi\phi})^2 \Big|_{\bar{\Phi}^i} T_{2,3} \right.$$

$$+ 2\nabla_R(\nabla_R^2 H) \cdot \nabla_R(\nabla_R^2 F) \Big|_{\bar{\Phi}^i} T_{2,4} + [\nabla(\nabla_R^2 F)]^2 \Big|_{\bar{\Phi}^i} T_{2,5} + G^{\phi\phi}(\nabla_R^2 F)^2 \Big|_{\bar{\Phi}^i} T_{2,6}$$

$$+ 2F(G^{\phi\phi})^2 \nabla_R^2 F \Big|_{\bar{\Phi}^i} T_{2,7} + \nabla_R(\nabla_R^2 F) \cdot \nabla_R(FG^{\phi\phi}) \Big|_{\bar{\Phi}^i} T_{2,8}$$

$$\left. + 2\nabla_R(\nabla_R^2 H) \cdot \nabla_R(FG^{\phi\phi}) \Big|_{\bar{\Phi}^i} T_{2,9} + [\nabla_R(FG^{\phi\phi})]^2 \Big|_{\bar{\Phi}^i} T_{2,10} + F^2(G^{\phi\phi})^3 \Big|_{\bar{\Phi}^i} T_{2,11} \right\}, \quad (37)$$

where we have defined the tension terms $T_{2,i}$ with $i = 1, \dots, 11$ in the same general form as above in (33) with their corresponding functions of the scalar field, and the sign ambiguity is the same as the one in eq. (34) for both terms. Furthermore, we have defined the dot product in the superspace as

$$\nabla f \cdot \nabla g = G^{MN} \frac{\partial f}{\partial \Phi^M} \frac{\partial g}{\partial \Phi^N}. \quad (38)$$

Therefore we note that the second quantum correction is described by $\bar{\Phi}^i$ one more time but we have many new tension terms, at most we can have 11 new independent constants (this is the maximum value, it can be reduced if some terms are zero or

equivalent). Furthermore, we obtain once again that the spatial slice volume $\text{Vol}(X)$ appears as a overall term. Thus we can compute the transition probabilities up to second quantum correction for metrics with a noncompact spatial slice.

Let us remark that the number of tension terms increase when the quantum corrections are taken into account. However, as we will see, at least for the metrics that will be of interest to us in the present article, most of the terms will be related or will vanish in such a way that in almost all cases we will only have one tension term per order in the WKB expansion.

Furthermore, let us also point out that in the Euclidean approach the loop contributions are taken into account by writing

$$\Phi^i = \Phi_{Cl}^i + \Phi_Q^i, \quad (39)$$

where Φ_{Cl}^i fulfils the equations of motion and Φ_Q^i are the quantum fluctuations. In the Lorentzian formalism the results of the quantum corrections are described with the same variable $\bar{\Phi}^i$. However, the quantum fluctuations of the degrees of freedom of the three metric are the responsible for adding all the integrals to Γ_1 and Γ_2 . Furthermore, the quantum fluctuations of the scalar field on the thin wall are the ones responsible of creating all these new tension terms. Thus, these quantum corrections should correspond to loop corrections in the Euclidean formalism as was proven at one loop for field theory in [39].

3 Transitions for a closed FLRW metric

Now that we have described in detail the general method to compute the transition probabilities, let us apply this method for metrics of cosmological interest. Let us begin with the FLRW closed metric that describes a closed homogeneous isotropic universe with positive spatial curvature.

The FLRW metric with positive curvature in 3+1 dimensions is written as

$$ds^2 = -N^2(t)dt^2 + a^2(t) (dr^2 + \sin^2 r d\Omega_2^2), \quad (40)$$

where $r \in [0, \pi]$, $d\Omega_2^2$ the metric of the sphere, $a(t)$ is the scale factor and $N(t)$ is the lapse function. Considering that the scalar field coupled in (30) depends only on the time variable, i.e., $\phi = \phi(t)$, we obtain the Lagrangian for this system as

$$\mathcal{L} = 3N(t)a(t) - \frac{3a(t)\dot{a}(t)^2}{N(t)} + \frac{a^3(t)\dot{\phi}^2}{2N(t)} - N(t)a^3(t)V(\phi) \quad (41)$$

where $\dot{a}(t)$ stands for the derivative of $a(t)$ with respect to the time variable t and $V(\phi)$ is the scalar field potential. In a standard way, we obtain the conjugate canonical momenta as

$$\pi_N = 0, \quad \pi_a = -\frac{6a\dot{a}}{N}, \quad \pi_\phi = \frac{a^3\dot{\phi}}{N} \quad (42)$$

which leads to the Hamiltonian constraint

$$H = N \left[\frac{\pi_\phi^2}{2a^3} - \frac{\pi_a^2}{12a} - 3a + a^3V \right] \simeq 0. \quad (43)$$

As always, since the canonical momentum with respect to N vanishes, we can ignore the prefactor and focus only on the term inside brackets in the last expression. Then by comparing it with the general form (1), we see that for this metric the set of minisuperspace variables is $\{\Phi^M\} = \{a, \phi\}$, while the metric in minisuperspace results in

$$(G^{MN}) = \begin{pmatrix} -1/6a & 0 \\ 0 & 1/a^3 \end{pmatrix}. \quad (44)$$

We identify the remaining terms as

$$f(a, \phi) = -3a + a^3 V(\phi). \quad (45)$$

We also note that for this metric the volume of the spatial slice is given by

$$\text{Vol}(X) = \int_{\phi=0}^{2\pi} \int_{\theta=0}^{\pi} \int_{r=0}^{\pi} \sin^2 r \sin \theta dr d\theta d\phi = 2\pi^2 \quad (46)$$

which is finite. In this case following the general method we can only choose $\Phi^i = a$ since there is only one degree of freedom coming from the metric, furthermore we can choose $a_0 = a(s=0) = 0$ since the expressions are well behaved for this value. Thus in this case the semiclassical contribution to the transition probability (34) takes the form

$$\begin{aligned} \Gamma_0 = & \pm \frac{12\pi^2}{\hbar} \left\{ \frac{1}{V_B} \left[\left(1 - \frac{V_B}{3} \bar{a}^2\right)^{3/2} - 1 \right] - \frac{1}{V_A} \left[\left(1 - \frac{V_A}{3} \bar{a}^2\right)^{3/2} - 1 \right] \right\} \\ & + \frac{2\pi^2}{\hbar} \bar{a}^3 T_0. \end{aligned} \quad (47)$$

This result was derived in [29, 30] and coincides with the one obtained with the Euclidean approach. We note from this result that $\lim_{\bar{a} \rightarrow 0} \Gamma_0 = 0$, then taking only this semiclassical contribution we will obtain that the transition probability (22) fulfils $\lim_{\bar{a} \rightarrow 0} P(A \rightarrow B) = 1$. Therefore, interpreting the transition probabilities as probability distributions of creating universes with a given size \bar{a} as it has been previously done in [29–32], we note that the semiclassical contribution implies that the most possible scenario is that the universe is created at the spatial singularity.

Let us continue with the first quantum correction, that is the term of order $\mathcal{O}(\hbar)$ in the WKB expansion. The general expression for this term is written in (35). For this metric we find that

$$\nabla^2 f = V^{(2)} - V, \quad (48)$$

where $V^{(n)}$ denotes the n -th derivative of the potential with respect to the scalar field. Therefore, substituting in (35) we obtain

$$\Gamma_1 = \pi^2 \left[\left(\frac{V_B^{(2)}}{V_B} - 1 \right) \ln(1 - \bar{a}^2 V_B) - \left(\frac{V_A^{(2)}}{V_A} - 1 \right) \ln(1 - \bar{a}^2 V_A) \right] + 2\pi^2 T_1. \quad (49)$$

We note that in this case there is only one tension term, since both expressions for the tension terms in the general form are equivalent, thus the first quantum correction only

adds one independent parameter to the transition probability. Moreover we note that $\lim_{\bar{a} \rightarrow 0} \Gamma_1 = 2\pi^2 T_1$ which is a constant. Therefore, the inclusion of the first quantum correction will not alter the fact that the point with the biggest probability is the spatial singularity ($\bar{a} = 0$) but it reduces the probability in that point.

Let us now move on to the second quantum correction. In this case we obtain

$$\begin{aligned}\nabla^2 (\nabla^2 f) &= \frac{1}{a^3} (V^{(4)} - V^{(2)}), \\ (\nabla (\nabla^2 f))^2 &= \frac{1}{a^3} (V^{(3)} - V^{(1)})^2.\end{aligned}\tag{50}$$

Then, for this metric we have

$$\frac{\text{Vol}(X)\hbar}{4\sqrt{2}} \int_{\Phi_0^i}^{\bar{\Phi}^i} \frac{\nabla^2 (\nabla^2 f)}{\nabla^i f} \sqrt{\frac{f}{(\nabla f)^2}} \Big|_{\phi=\phi_B} d\Phi^i = \pi^2 \hbar (V^{(4)} - V^{(2)}) F(V_B, a) \Big|_{a_0}^{\bar{a}},\tag{51}$$

$$\frac{\text{Vol}(X)\hbar}{8\sqrt{2}} \int_{\Phi_0^i}^{\bar{\Phi}^i} \frac{(\nabla (\nabla^2 f))^2}{\nabla^i f} \sqrt{\left(\frac{f}{(\nabla f)^2}\right)^3} \Big|_{\phi=\phi_B} = \pi^2 \hbar (V^{(3)})^2 G(V_B, a) \Big|_{a_0}^{\bar{a}},\tag{52}$$

where we have defined the functions

$$\begin{aligned}F(V, a) &= \int \frac{\left(1 - \frac{V}{3}\bar{a}^2\right)^{1/2}}{a(1 - V\bar{a}^2)^2} da = -\frac{\sqrt{1 - \frac{V}{3}a^2}}{2(-1 + Va^2)} + \frac{5}{2\sqrt{6}} \operatorname{arctanh}\left(\frac{\sqrt{3 - Va^2}}{\sqrt{2}}\right) \\ &\quad - \operatorname{arctanh}\left[\sqrt{1 - \frac{V}{3}a^2}\right],\end{aligned}\tag{53}$$

$$\begin{aligned}G(V, a) &= \int \frac{a\left(1 - \frac{V}{3}\bar{a}^2\right)^{3/2}}{(1 - V\bar{a}^2)^4} da = \frac{1}{576\sqrt{3}V} \left[-\frac{2\sqrt{3 - Va^2}(63 - 34Va^2 + 3V^2a^4)}{(-1 + Va^2)^3} \right. \\ &\quad \left. - 3\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{3 - Va^2}}{\sqrt{2}}\right) \right].\end{aligned}\tag{54}$$

We note that the $G(V, a)$ function is well behaved in the limit $a \rightarrow 0$. On the other hand the third term of the $F(V, a)$ function diverges in this limit. However, we note that this limit is independent of V , therefore we can eliminate this divergence in the transition probability by imposing the condition

$$V_B^{(4)} - V_B^{(2)} = V_A^{(4)} - V_A^{(2)}.\tag{55}$$

In this way we can still choose $a_0 = 0$ in order to have access to the initial spatial singularity. Furthermore for this metric from the eleven possible tension terms only one remains since all the nonzero terms have the same dependence on \bar{a} , thus we finally obtain that the second quantum correction is written as

$$\begin{aligned}\Gamma_2 &= \pm \pi^2 \hbar \left\{ \left(V_B^{(4)} - V_B^{(2)} \right) [F(V_B, \bar{a}) - F(V_A, \bar{a})] - \left(V_B^{(3)} \right)^2 [G(V_B, \bar{a}) - G(V_B, 0)] \right. \\ &\quad \left. + \left(V_A^{(3)} \right)^2 [G(V_A, \bar{a}) - G(V_A, 0)] \right\} + \frac{2\pi^2 \hbar}{\bar{a}^3} T_2.\end{aligned}\tag{56}$$

We note that once again, the second quantum correction only adds one more independent parameter tension term. From this result we also note that the limit $\lim_{\bar{a} \rightarrow 0} \Gamma_2$ will diverge because of the tension term. Thus, choosing appropriately the sign of (24) we will obtain that taking into account up to second order in the quantum corrections the transition probability fulfils $\lim_{\bar{a} \rightarrow 0} P(A \rightarrow B) = 0$. Therefore, the second order quantum correction implies that the most probable size of the universe to be created is at a finite nonzero value of the scale factor. In this way, the quantum correction terms leads to a result that avoids the spatial singularity.

Let us remark from (47) that the semiclassical contribution Γ_0 is well behaved for all values of the potentials, but it can be complex for positive values of the potential minima. However this not a problem since we always take only the real part. Nevertheless we see from (49) that Γ_1 have a divergence in $\bar{a}^2 = V_{A,B}^{-1}$ which can occur for positive values of the potential minima. Furthermore from (56) we note that Γ_2 will also have a divergence appearing in $\bar{a}^2 = \frac{5}{V_{A,B}}$, which can also appear for positive values of the potential minima. In these cases, the transition probability will only be well defined up to an upper limit. On the other hand, we note that for negative potential minima everything is well defined without any issue. Therefore positive values of the potential minima will imply some additional restrictions for the validity of the transition probabilities. Moreover, in order to have a well defined second quantum correction term we need to impose additional restrictions on the higher derivatives of the potential.

In order to explicitly visualize the effect of the first and second order quantum corrections for all values of the scale factor, we plot in Figure 1 the transition probability (22) using the results for this metric, that is using eq. (47) for the semiclassical contribution, eq. (49) for the first quantum correction and eq. (56) for the second order quantum correction. We choose units in which $\hbar = 0.1$, we also choose $V_B = -2$, $V_A = -1$, $\frac{V_B^{(2)}}{V_B} - 1 = \frac{V_A^{(2)}}{V_A} - 1 = 0.5$, $V_B^{(4)} - V_B^{(2)} = -0.05$, $V_B^{(3)} = V_A^{(3)} = 0.1$, $T_0 = 1$, $T_1 = 5 \times 10^{-4}$ and $T_2 = 10^{-6}$. In order to have a well defined transition probability we choose the plus sign in the left hand side of (24) and the minus sign in the right of (47) which corresponds to the minus sign in (56) as well. We note that the result with the semiclassical contribution only has a maximum value on the spatial singularity and as \bar{a} increases, the probability decreases. The result including the first quantum correction does not change the overall shape of the curve, it only reduces the overall probability and make it fall faster. However, as we have previously seen, when the second quantum correction is taken into account as well, the shape of the curve is changed in the ultraviolet (small values of the scale factor), in such a way that the maximum value of the transition probability is no longer in the spatial singularity, it is instead located at a small nonzero value of the scale factor. From this point, the same behaviour is encountered as in the other two results in the infrared, that is as the scale factor increases, the probability decreases. Thus, the quantum corrections are only relevant on the ultraviolet as expected.




Figure 1: Transition probability for the positive FLRW metric in units such that $\hbar = 0.1$, choosing $V_B = -2$, $V_A = -1$, $\frac{V_B^{(2)}}{V_B} - 1 = \frac{V_A^{(2)}}{V_A} - 1 = 0.5$, $V_B^{(4)} - V_B^{(2)} = -0.05$, $V_B^{(3)} = V_A^{(3)} = 0.1$, $T_0 = 1$, $T_1 = 5 \times 10^{-4}$, $T_2 = 10^{-6}$. We plot the results with the semiclassical contribution Γ_0 (red line), including the first quantum correction $\Gamma_0 + \Gamma_1$ (blue line) and including the second quantum correction $\Gamma_0 + \Gamma_1 + \Gamma_2$ (purple line).

4 Transitions for the flat FLRW metric

Let us consider now the FLRW metric with zero spatial curvature. This metric will be important in the following since it represents the isotropic limit of the Bianchi III metric. We can write this metric in cartesian coordinates as follows

$$ds^2 = -N(t)dt^2 + a^2(t) [dx^2 + dy^2 + dz^2]. \quad (57)$$

As stated above, the gravitational and scalar fields depend only on the time variable t . Thus the corresponding Lagrangian is given by

$$\mathcal{L} = -\frac{3a(t)\dot{a}(t)^2}{N(t)} + \frac{a^3(t)\dot{\phi}^2}{2N(t)} - N(t)a^3(t)V(\phi) \quad (58)$$

and the Hamiltonian constraint is

$$H = N \left[\frac{\pi_\phi^2}{2a^3} - \frac{\pi_a^2}{12a} + a^3V \right] \simeq 0. \quad (59)$$

As always the N function is not dynamical, so we can focus on the terms within brackets and proceed to identify the metric in minisuperspace and the function f . Comparing it with the general form (2) we see that once again the coordinates in minisuperspace are $\{\Phi^M\} = \{a, \phi\}$ with the metric

$$G^{MN} = \begin{pmatrix} -\frac{1}{6a} & 0 \\ 0 & \frac{1}{a^3} \end{pmatrix} \quad (60)$$

and the function f reads

$$f(a, \phi) = a^3 V(\phi). \quad (61)$$

We also note that in this case the spatial volume takes the form

$$\text{Vol}(X) = \iiint dx dy dz. \quad (62)$$

In a strict sense this volume is divergent as a result of the noncompact nature of the 3-space of the 3+1 decomposition. However, as we pointed out earlier we can constrain the variables to a finite interval in order to obtain a finite value for this term. Since it will be an overall constant we will not worry about its value and we will maintain it as an arbitrary constant that will not modify the behaviour of the transition probabilities.

Then, substituting the above identifications for this metric into eq. (34) we obtain

$$\Gamma_0 = \pm \frac{2i \text{Vol}(X)}{\sqrt{3}\hbar} \left(\sqrt{V_B} - \sqrt{V_A} \right) \bar{a}^3 + \frac{\text{Vol}(X)}{\hbar} \bar{a}^3 T_0. \quad (63)$$

We note that $\lim_{\bar{a} \rightarrow 0} \Gamma_0 = 0$, then considering only the semiclassical contribution we would obtain for the transition probability $\lim_{\alpha=0} P(A \rightarrow B) = 1$. Thus, once again, the semiclassical contribution leads to a maximum value for probability at the initial singularity.

Proceeding with the first quantum correction, we have also in this case

$$\nabla^2 f = V'' - V, \quad (64)$$

that substituting back in (35) we get

$$\Gamma_1 = \frac{\text{Vol}(X)}{2} \left(\frac{V_B^{(2)}}{V_B} - \frac{V_A^{(2)}}{V_A} \right) \ln(a) \Big|_{a_0}^{\bar{a}} + \frac{\text{Vol}(X)}{2} T_1. \quad (65)$$

Moreover, as in the previous case, there only remains a single tension term, thus the first quantum correction only adds one independent constant. From this expression we also notice that the choice $a_0 = 0$ may lead to a divergence due to the logarithmic term. Therefore, in order to obtain a consistent probability we are forced to impose the condition on the second derivatives of the potential

$$\frac{V_B^{(2)}}{V_B} = \frac{V_A^{(2)}}{V_A}. \quad (66)$$

Thus the first quantum correction is written finally as

$$\Gamma_1 = \frac{\text{Vol}(x)}{2} T_1, \quad (67)$$

which is just a constant. Consequently, as was discussed in the previous example, the first quantum correction does not alter the overall behaviour, in particular the point with maximum probability.

Let us move on to the computation of the second quantum correction. For this metric we also have

$$\nabla^2 (\nabla^2 f) = \frac{V^{(4)} - V^{(2)}}{a^3}, \quad (68)$$

$$(\nabla (\nabla^2 f))^2 = \frac{(V^{(3)} - V^{(1)})^2}{a^3}. \quad (69)$$

Therefore, substituting back these equations into (37) and computing all the remaining terms, we finally obtain

$$\Gamma_2 = \pm \frac{i\hbar \text{Vol}(X)}{2} \left[(W_B - W_A) \frac{1}{3a^3} \Big|_{a_0}^{\bar{a}} \right] + \frac{\hbar \text{Vol}(X)}{\bar{a}^3} T_2, \quad (70)$$

where we have defined

$$W = \frac{V^{(2)} - V^{(4)}}{\sqrt{3}V^{3/2}} - \frac{(V^{(3)})^2}{3\sqrt{3}V^{5/2}}. \quad (71)$$

For this metric we obtain once again that from the eleven possible different tension terms, only one dependence on \bar{a} appears, leading to only one tension term. In this way, the second quantum correction also adds only one independent constant. Here we also note that the choice $a_0 = 0$ will lead to a divergence in the first term of (70). However, for positive values of the potential minima we note that the first term never contributes to the transition probability after taking the real part. Therefore, for positive potentials the second quantum correction takes simply the form

$$\Gamma_2 = \frac{\text{Vol}(X)\hbar}{\bar{a}^3} T_2. \quad (72)$$

On the other hand, for negative potential minima, this divergence can be eliminated by imposing the constraint

$$W_A = W_B. \quad (73)$$

Then, the negative potential minima leads to an additional constraint on the higher derivative terms of the potential, but it takes the same form as in the previous example. In any case, we note that $\lim_{\alpha=0} \Gamma_2$ will diverge. Thus by choosing the proper signs, we will obtain that $\lim_{\bar{a} \rightarrow 0} P(A \rightarrow B) = 0$ as in the previous section.

We can plot the transition probability for this metric considering the semiclassical contribution (63), the first quantum correction (67) and the second quantum correction (72). In this case, we obtain the same general behaviour as the one encountered for the positive FLRW metric in Figure 1 in the previous section. Therefore, for this metric the second quantum correction will also lead to an avoidance of the initial singularity.

5 Transitions for the Bianchi III metric

Now that we have completed the study of the FLRW metrics, let us move on to consider metrics describing homogeneous but anisotropic universes.

We will begin by considering the Bianchi III metric in the present section. In cartesian coordinates this metric has the following form

$$ds^2 = -N^2(t)dt^2 + A^2(t)dx^2 + B^2(t)e^{-2\alpha x}dy^2 + C^2(t)dz^2, \quad (74)$$

where $\alpha \neq 0$ is a constant measuring the amount of anisotropy. We see that the isotropy limit is performed by taking $A(t) = B(t) = C(t)$ and $\alpha \rightarrow 0$ and results in the flat FLRW metric. Furthermore, if we consider different values for the scale factors $A(t)$, $B(t)$ and $C(t)$ and take $\alpha \rightarrow 0$ we will obtain the anisotropic Bianchi I metric.

Let us consider now as usual a homogeneous scalar field coupled to gravity with the above metric, the Lagrangian takes the form

$$\begin{aligned} \mathcal{L} = & -\frac{1}{N(t)}(\dot{A}(t)\dot{B}(t)C(t) + \dot{A}(t)\dot{C}(t)B(t) + \dot{B}(t)\dot{C}(t)A(t)) - \frac{\alpha^2}{A(t)}N(t)B(t)C(t) \\ & + \frac{\dot{\phi}^2}{2N(t)}A(t)B(t)C(t) - V(\phi)N(t)A(t)B(t)C(t). \end{aligned} \quad (75)$$

The canonical momenta are given by

$$\begin{aligned} \pi_N = 0, \quad \pi_A = & -\frac{1}{N}(\dot{B}C + \dot{C}B), \quad \pi_B = -\frac{1}{N}(\dot{A}C + \dot{C}A) \\ \pi_C = & -\frac{1}{N}(\dot{A}B + \dot{B}A), \quad \pi_\phi = \frac{\dot{\phi}}{N}ABC. \end{aligned} \quad (76)$$

Then, the Hamiltonian constraint takes the form

$$\begin{aligned} H = N \left[\frac{A}{4BC}\pi_A^2 + \frac{B}{4AC}\pi_B^2 + \frac{C}{4AB}\pi_C^2 - \frac{1}{2C}\pi_A\pi_B - \frac{1}{2B}\pi_A\pi_C - \frac{1}{2A}\pi_B\pi_C \right. \\ \left. + \frac{\pi_\phi^2}{2ABC} + \frac{\alpha^2 BC}{A} + V(\phi)ABC \right] \simeq 0. \end{aligned} \quad (77)$$

Comparing with the general form (1), we see that the coordinates on minisuperspace are $\{\Phi^M\} = \{A, B, C, \phi\}$. Moreover the metric is written as

$$(G^{MN}) = \frac{1}{2ABC} \begin{pmatrix} A^2 & -AB & -AC & 0 \\ -AB & B^2 & -BC & 0 \\ -AC & -BC & C^2 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix} \quad (78)$$

and the function f reads

$$f(A, B, C, \phi) = ABC \left(V(\phi) + \frac{\alpha^2}{A^2} \right). \quad (79)$$

We also note that in this case the volume of X is given by

$$\text{Vol}(X) = \iiint e^{-\alpha x} dx dy dz \quad (80)$$

which again is finite only if we restrict to a finite interval in y and z but we will consider it as an arbitrary constant. In [30] the semiclassical contribution to the transition probability was studied for this metric. However, in that work a convenient factorization was used to obtain a result consistent in the case of a vanishing potential. In the present case it will be more convenient to consider the Hamiltonian constraint without any factorization. The dependence of the results on a particular factorization will be explored in the Appendix A.

We note that for this metric we have more than one variable on the minisuperspace apart from the scalar field. Thus, as it was stated in the general method in Section 2 these variables will be related as a consequence of the semiclassical expansion. In particular, in the region where the scalar field is constant eq. (21) leads to

$$\frac{dB}{dC} = \frac{B}{C}, \quad \frac{dA}{dB} = \frac{A \left(V + \frac{3\alpha^2}{A^2} \right)}{B \left(B - \frac{\alpha^2}{A^2} \right)}, \quad (81)$$

which can be integrated to obtain

$$B = b_0 C = b_1 \frac{\left(A^2 + \frac{3\alpha^2}{V} \right)^{2/3}}{A^{1/3}}, \quad (82)$$

where b_0 and b_1 are integration constants. However, we can set these constants to 1 by rescaling the variables on the metric in a suitable way for each of the solutions and demanding continuity on the wall at \bar{B} . Therefore we obtain for the region where the scalar field is constant

$$B = C = \frac{\left(A^2 + \frac{3\alpha^2}{V} \right)^{2/3}}{A^{1/3}}. \quad (83)$$

From this relation we note that the isotropy limit leads to $\lim_{\alpha \rightarrow 0} B = A$. Therefore, all the anisotropy will be measured by the constant α . Moreover as we pointed out previously the limit $\alpha \rightarrow 0$ corresponds to the Bianchi I metric, thus from these relations we note that if we study the Bianchi I metric, we will be forced to obtain the flat FLRW metric and its corresponding result. Thus, the Bianchi III metric is more suitable to study the effect of anisotropy.

Furthermore, we note that given $\alpha \neq 0$, the variable B will diverge as $A \rightarrow 0$. On the other hand for big values of A , B increases with A . Thus, in this case the cosmological behaviour does not describe a spatial singularity at the beginning ($A = 0$). Instead we have in the early universe (understood as small values of A) a behaviour in which B and C start with big values (infinite in principle), then as A grows, they decrease until they reach a minimum value, from where they start growing. For positive potential minima this minimum value is nonzero, whereas for a negative potential minimum it is zero but always with a nonzero value of A . Thus, we have a phase of contraction on the B and C direction, until they reach a point where they start an expansion as A expands, that is, similar to a bounce scenario as the dimension described by A emerges⁶. We note that the size of the contraction phase before the bounce depends

⁶The relation between the coordinates can change if we consider a different factorization. This point will be discussed in the Appendix A.

on the size of the values of $\frac{3\alpha^2}{V}$ and it disappears in the isotropy limit. Let us remark that since we have two different degrees of freedom coming from the metric, namely A and B we can write the transition probabilities in terms of any of these two, however only A is single-valued in time, whereas the other two can have the same value twice given the bounce nature. For all these reasons we will consider only positive values of the potential minima and from now on write the transition probabilities in terms of the A function.

With the identifications made above for this metric, substituting back in (27) we obtain for the semiclassical contribution

$$\begin{aligned} \Gamma_0 = & \pm \frac{2i\text{Vol}(X)}{\hbar} \left[\sqrt{V_B} F_{III}(V_B, A) \Big|_{A_0}^{\bar{A}} - \sqrt{V_A} F_{III}(V_A, A) \Big|_{A_0}^{\bar{A}} \right] \\ & + \frac{\text{Vol}(X)}{\hbar} \bar{A}^{1/3} \left(\bar{A}^2 + \frac{3\alpha^2}{V_B} \right)^{4/3} T_0, \end{aligned} \quad (84)$$

where we have defined the function

$$F_{III}(V, A) = \int \frac{\left(A^2 + \frac{3\alpha^2}{V} \right)^{1/3}}{A^{5/3}} \sqrt{\left(A^2 + \frac{\alpha^2}{V} \right) \left(A^2 - \frac{\alpha^2}{V} \right) \left(3A^2 + \frac{5\alpha^2}{V} \right)} dA. \quad (85)$$

It can be shown that the Bianchi III results (84) coincides with the flat FLRW result (63) in the isotropy limit $\alpha \rightarrow 0$ as we expected.

Taking $A_0 = 0$ we note from the last expression that $\lim_{\bar{A} \rightarrow 0} \Gamma_0 = 0$. Thus we obtain once again that considering only the semiclassical contribution the probability will have its biggest value in $\bar{A} = 0$. However, let us remark that in this case this point does not represent a spatial singularity. Furthermore, we plot the last expression varying with respect to α to obtain Figure 2, where in order to have a well-behaved probability we have chosen the minus sign in both sign ambiguities. Additionally, we choose positive values for the potential minima, $V_B = 1$ and $V_A = 5$, and for the tension term $T_0 = 1$. We can see that the form of the semiclassical contribution coincides with the general form of the FLRW results. Moreover, the effect of anisotropy is to decrease the value of the transition probability as the value of α increases, and the case $\alpha \rightarrow 0$ corresponds to the FLRW result, in this form our result is consistent with [30].

Let us continue now with the computation of the first quantum correction, in this case we obtain

$$\nabla^2 f = V^{(2)} - 3V + \frac{2\alpha^2}{A^2}. \quad (86)$$

Then, computing the remaining terms and substituting back in eq. (35) we obtain for the first quantum correction

$$\begin{aligned} \Gamma_1 = & \text{Vol}(X) \left[\left(\frac{V_B^{(2)}}{V_B} - \frac{11}{3} \right) \ln \sqrt{\frac{\bar{A}^2 + \frac{3\alpha^2}{V_B}}{A_0^2 + \frac{3\alpha^2}{V_B}}} - \left(\frac{V_A^{(2)}}{V_A} - \frac{11}{3} \right) \ln \sqrt{\frac{\bar{A}^2 + \frac{3\alpha^2}{V_A}}{A_0^2 + \frac{3\alpha^2}{V_A}}} \right] \\ & + \text{Vol}(X) T_1. \end{aligned} \quad (87)$$




Figure 2: Transition probabilities for the Bianchi III metric in units such that $\hbar = 1$, choosing $V_B = 1$, $V_A = 5$, $T_0 = 1$. We plot the results of the semiclassical contribution Γ_0 varying the value of α . We choose the cases, $\alpha = 0$ that corresponds to the flat FLRW case (red line), $\alpha = 0.075$ (blue line), $\alpha = 0.1$ (yellow line), $\alpha = 0.15$ (green line) and $\alpha = 0.2$ (purple line).

We note that taking the isotropic limit $\alpha \rightarrow 0$ in the last expression we obtain the flat FLRW result of eq. (67). However, contrary to that case, we will not have any divergence if we take $A_0 = 0$ as long as $\alpha \neq 0$. Moreover, in order to obtain the flat FLRW result in the limit $\alpha \rightarrow 0$ we impose the same condition (66), thus the first quantum correction takes the form

$$\Gamma_1 = \text{Vol}(X) \left\{ \left(\frac{V_B^{(2)}}{V_B} - \frac{11}{3} \right) \ln \left[\sqrt{\frac{\bar{A}^2 + \frac{3\alpha^2}{V_B}}{A_0^2 + \frac{3\alpha^2}{V_B}}} \sqrt{\frac{\bar{A}^2 + \frac{3\alpha^2}{V_A}}{A_0^2 + \frac{3\alpha^2}{V_A}}} \right] \right\} + \text{Vol}(X)T_1. \quad (88)$$

Now choose $A_0 = 0$, then the latter simplifies to

$$\Gamma_1 = \text{Vol}(X) \left(\frac{V_B^{(2)}}{V_B} - \frac{11}{3} \right) \ln \sqrt{\frac{V_B}{V_A} \left(\frac{\bar{A}^2 + \frac{3\alpha^2}{V_B}}{\bar{A}^2 + \frac{3\alpha^2}{V_A}} \right)} + \text{Vol}(X)T_1. \quad (89)$$

However, we note that once we have chosen $A_0 = 0$ we can not longer recover the flat FLRW result (67) in the isotropy limit. In the isotropy limit the first quantum correction will be a constant, but the first term will contribute with a logarithmic constant term as well. Thus, the first quantum correction does not lead exactly to the flat FLRW result in the isotropy limit. Furthermore, we also note that we only obtain one tension term, thus the first quantum correction adds only one independent constant. Moreover, we note that in order to have a well defined probability we should choose only positive values of the potential minima as we have remarked before. In addition, we also note that $\lim_{\bar{A} \rightarrow 0} \Gamma_1 = \text{Vol}(X)T_1$ which is a constant, thus as in the

previous cases the first quantum correction will not alter the behaviour near the value $\bar{A} = 0$. In Figure 3 we show a plot of this result taking different values of α , we have considered the same values for the potential minima, and choose $\frac{V_B^{(2)}}{V_B} - \frac{11}{3} = -0.005$, $T_0 = 1$, $T_1 = 0.1$. We see that the general behaviour is the same that the FLRW result. That is, for all values of α , the first quantum correction does not change the overall behaviour, it only reduces the probability. The effect of anisotropy is to reduce the probability once again. Moreover, in this case the isotropy limit does not correspond exactly to the flat FLRW result.




Figure 3: Transition probabilities for the flat FLRW (orange line) and Bianchi III metrics in units such that $\hbar = 1$, choosing $V_B = 1$, $V_A = 5$, $\frac{V_B^{(2)}}{V_B} - \frac{11}{3} = -0.005$, $T_0 = 1$, $T_1 = 0.1$. We plot the results of the semiclassical contribution plus the first quantum correction, $\Gamma_0 + \Gamma_1$, varying the value of α ; we choose the cases $\alpha = 0$ (purple line), $\alpha = 0.075$ (blue line), $\alpha = 0.1$ (yellow line), $\alpha = 0.15$ (green line) and $\alpha = 0.2$ (red line). The isotropy limit of the Bianchi III metric $\alpha \rightarrow 0$, does not correspond exactly with the flat FLRW result

Finally, for the second correction we obtain in this case

$$\nabla^2 (\nabla^2 f) = \frac{V^{(4)} - 3V^{(2)} + \frac{6\alpha^2}{A^2}}{ABC}, \quad (90)$$

$$[\nabla (\nabla^2 f)]^2 = \frac{(V^{(3)} - 3V^{(1)})^2 + \frac{8\alpha^4}{A^4}}{ABC}. \quad (91)$$

Then, computing the remaining terms in eq. (37) we obtain for the second quantum correction

$$\begin{aligned} \Gamma_2 = & \pm \frac{i\hbar \text{Vol}(X)}{2} \left[\frac{1}{V_B^{3/2}} H_{III}(V_B, A) \Big|_{A_0}^{\bar{A}} - \frac{1}{V_A^{3/2}} H_{III}(V_A, A) \Big|_{A_0}^{\bar{A}} \right] \\ & + \frac{\text{Vol}(X)\hbar}{\bar{A}^{1/3} \left(\bar{A}^2 + \frac{3\alpha^2}{V_B} \right)^{4/3}} T_2, \end{aligned} \quad (92)$$

where the H_{III} function is defined as

$$H_{III}(V, A) = \int \frac{dA}{A^{1/3} \left(A^2 + \frac{3\alpha^2}{V}\right)^{7/3}} \sqrt{\frac{A^2 + \frac{\alpha^2}{V}}{\left(A^2 - \frac{\alpha^2}{V}\right)\left(3A^2 + \frac{5\alpha^2}{V}\right)}} \times \left[6\alpha^2 + (V^{(4)} - 3V^{(2)}) A^4 + \frac{((V^{(3)})^2 A^4 + 8\alpha^2) \left(A^2 + \frac{\alpha^2}{V}\right)}{V \left(A^2 - \frac{\alpha^2}{V}\right) \left(3A^2 + \frac{5\alpha^2}{V}\right)} \right]. \quad (93)$$

As in the previous case we note that the first term of eq. (92) has an overall i , thus if the function is always real for positive potentials, this term will not contribute. Furthermore we note that once again only one tension term is added from the eleven possible because the nonzero terms have the same depend with A . However, if we take the isotropy limit for this result we obtain

$$\lim_{\alpha=0} \Gamma_2 = \pm \frac{i\hbar \text{Vol}(X)}{2} \left[(\bar{W}_B - \bar{W}_A) \frac{1}{3A^3} \Big|_{A_0}^{\bar{A}} \right] + \frac{\hbar \text{Vol}(X)}{A^3} T_2, \quad (94)$$

where

$$\bar{W} = \frac{(3V^{(2)} - V^{(4)})}{\sqrt{3}V^{3/2}} - \frac{(V^{(3)})^2}{3\sqrt{3}V^{5/2}}, \quad (95)$$

which is very similar to the result for the FLRW metric in eq. (70). The only difference appears in the constant accompanying the second derivative on the W function, in this case it is a factor of 3, whereas for the flat FLRW metric it was just a factor of 1. The differences that appear at first and second quantum corrections come from the fact that although the FLRW metric follows from the isotropy limit $\alpha \rightarrow 0$, if we look for the definition of the canonical momenta on this case in (76), we note that the isotropy limit does not lead to the canonical momentum of the single scale factor of the FLRW metric in (42). The same occurs with the metric on superspace, thus the structure of the minisuperspace is not the same after taking the isotropy limit and we should not expect the same results in all scenarios. In particular we note that the laplacian in the Bianchi III metric (86) does not lead to the laplacian for the FLRW metric (86) in the isotropy limit. It is interesting that this discrepancy appears when we take into account the quantum corrections and it only changes by constants, it does not change the general behaviour. Furthermore, we note that in the general case we have for this metric that $\lim_{\bar{A} \rightarrow 0} \Gamma_2$ will diverge, thus, making the probability vanish on this limit after choosing the appropriate signs.

Now, by considering different values of α , and arduously exploring the expression (93) we can find values of the parameters in which H_{III} only takes real values. In this way, the first term in (92) does not contribute to the transition probability, we only have the tension term. This can be done for example by choosing $V_B = 1$, $V_A = 5$, $\frac{V_B^{(2)}}{V_B} - \frac{11}{3} = -0.005$, $T_0 = 1$, $T_1 = 0.1$, $T_2 = 0.05$, $(V_A^{(3)})^2 = (V_B^{(3)})^2 = 0.1$, $V_A^{(4)} - 3V_A^{(2)} = V_B^{(4)} - 3V_B^{(2)} = 0.1$. The resulting transition probabilities are shown in Figure 4 for different values of α . It is shown that the general form corresponds to the behaviour encountered in the FLRW metrics. The effect of anisotropy is once again to

reduce the probability and move the maximum value towards smaller values of \bar{A} as anisotropy is increased.

In summary, we have obtained that the behaviour of the transition probabilities for the Bianchi III metric is in general the same as for the FLRW metrics, that is, the semiclassical contribution leads to a maximum value at the origin of \bar{A} , the first quantum correction reduces the probability but does not change the overall shape and the second quantum correction leads to a vanishing probability in this point. However, as we pointed out earlier for this metric this point does not represent a spatial singularity. What is being avoided in this case is the situation where one dimension is absent and the other two are infinite in size. We could try different factorizations looking for a scenario that allows us to have access to a spatial singularity in this anisotropic metric looking for a description where the second quantum correction avoids such singularity. However, as we will show in the Appendix A this scenario does not exist. Therefore, we conclude that the avoidance of the initial singularity due to quantum corrections is only possible in the isotropic case. Moreover, we have shown that the effect of the anisotropy is to reduce the probability in all scenarios, making it fall faster. In particular, when considering up to second quantum correction terms, the maximum point of the probability is moved towards smaller values of \bar{A} .




Figure 4: Transition probabilities for the flat FLRW (orange line) and Bianchi III metrics in units such that $\hbar = 1$, choosing $V_B = 1$, $V_A = 5$, $\frac{V_B^{(2)}}{V_B} - \frac{11}{3} = -0.005$, $T_0 = 1$, $T_1 = 0.1$, $T_2 = 0.05$, $(V_A^{(3)})^2 = (V_B^{(3)})^2 = 0.1$, $V_A^{(4)} - 3V_A^{(2)} = V_B^{(4)} - 3V_B^{(2)} = 0.1$. We plot the results of the semiclassical contribution plus the first and second quantum corrections, $\Gamma_0 + \Gamma_1 + \Gamma_2$, varying the value of α ; we choose the cases $\alpha = 0$ (purple line), $\alpha = 0.075$ (blue line), $\alpha = 0.1$ (yellow line), $\alpha = 0.15$ (green line) and $\alpha = 0.2$ (red line). The isotropy limit of the Bianchi III metric $\alpha \rightarrow 0$, does not correspond exactly with the flat FLRW result

6 Transitions for the Kantowski-Sachs metric

Finally, we will consider the Kantowski-Sachs metric. This metric describes an anisotropic universe or the interior of a Schwarzschild black hole. Following the parametrization proposed by Misner [51] we can write down this metric in the form

$$ds^2 = -N^2(t)dt^2 + e^{2\sqrt{3}\beta(t)}dr^2 + e^{-2\sqrt{3}(\beta(t)+\Omega(t))} [d\theta^2 + \sin^2(\theta)d\psi^2], \quad (96)$$

with $0 \leq \theta \leq \pi$ and $0 \leq \psi \leq 2\pi$. Let us define for simplicity the functions $\gamma(t) = e^{\sqrt{3}\beta(t)}$ and $\sigma(t) = e^{-\sqrt{3}\Omega(t)}$, then the metric is written as

$$ds^2 = -N^2(t)dt^2 + \gamma^2(t)dr^2 + \frac{\sigma^2(t)}{\gamma^2(t)} [d\theta^2 + \sin^2\theta d\psi^2] \quad (97)$$

Considering once again a homogeneous scalar field the Lagrangian takes the form

$$\mathcal{L} = \frac{\sigma^2(t)}{\gamma^3(t)N(t)}\dot{\gamma}^2 - \frac{\dot{\sigma}^2}{\gamma(t)N(t)} + N(t)\gamma(t) + \left[\frac{\dot{\phi}^2}{2N(t)} - N(t)V(\phi) \right] \frac{\sigma^2(t)}{\gamma(t)} \quad (98)$$

and the Hamiltonian constraint turns out to be

$$H = N \left[\frac{\gamma^3}{4\sigma^2}\pi_\gamma^2 - \frac{\gamma}{4}\pi_\sigma^2 + \frac{\gamma}{2\sigma^2}\pi_\phi^2 + \frac{\sigma^2}{\gamma}V(\phi) - \gamma \right] \simeq 0. \quad (99)$$

Once more, we can focus only on the term inside brackets, and comparing it with the general form (1), we observe that the coordinates defining the minisuperspace are $\{\Phi^M\} = \{\gamma, \sigma, \phi\}$. Thus, the metric is

$$G^{MN} = \begin{pmatrix} \frac{\gamma^3}{2\sigma^2} & 0 & 0 \\ 0 & -\frac{\gamma}{2} & 0 \\ 0 & 0 & \frac{\gamma}{\sigma^2} \end{pmatrix} \quad (100)$$

and the function f is given by

$$f(\gamma, \sigma, \phi) = \frac{\sigma^2}{\gamma}V(\phi) - \gamma. \quad (101)$$

We also have in this case that the volume of the spatial slice X reads

$$\text{Vol}(X) = \int_r \int_{\theta=0}^{\pi} \int_{\psi=0}^{2\pi} \sin\theta dr d\theta d\psi = 4\pi \int dr, \quad (102)$$

which is finite only if we restrict it to a finite interval in r , since the spatial slice is again noncompact in this scenario. Thus we will consider it as an overall arbitrary constant as in the previous cases.

In order to obtain a consistent system of equations we need to consider nonzero values for the potential minima. Therefore for the region of constant scalar field, the relations (21) leads in this case to the differential equation

$$2\sigma V d\gamma - \gamma \left(\frac{\gamma^2}{\sigma^2} + V \right) d\sigma = 0. \quad (103)$$

Integration of this expression leads to the relation

$$\gamma^2 = \frac{\sigma^2 V}{1 - c\sigma}, \quad (104)$$

where c is an integration constant. We note that by definition γ and σ are positive functions, therefore c has to fulfill the condition

$$\frac{V}{1 - c\sigma} > 0, \quad (105)$$

which depends on the value of V (either V_A or V_B) and the possible values that σ can have. From this expression we note that $\lim_{\sigma \rightarrow 0} \gamma = 0$ but $\lim_{\sigma \rightarrow 0} \frac{\sigma^2}{\gamma^2} = \frac{1}{V}$. Therefore, we note from the form of the metric (97) that the limit $\sigma \rightarrow 0$ leads to the vanishing of the radial component but not of the angular part, thus once again we will not have access to a spatial singularity.

Computing the remaining terms and substituting back in (34) we obtain

$$\Gamma_0 = \pm \frac{\text{Vol}(X)i}{\hbar} \left[c_B F_{KS}[c_B, \sigma] \Big|_{\sigma_0}^{\bar{\sigma}} - c_A F_{KS}[c_A, \sigma] \Big|_{\sigma_0}^{\bar{\sigma}} \right] + \frac{\text{Vol}(X)}{\hbar} \frac{\bar{\sigma} T}{\sqrt{V_B}} \sqrt{1 - c_B \bar{\sigma}}, \quad (106)$$

where we have defined the function

$$\begin{aligned} F_{KS}[c, x] &= \int \frac{\sqrt{4 - 3cx}}{1 - cx} x dx \\ &= -\frac{2}{9c^2} [\sqrt{4 - 3cx}(5 + 3cx) - 9 \operatorname{arctanh}(\sqrt{4 - 3cx})]. \end{aligned} \quad (107)$$

This result was first derived in [30]. We note that $\lim_{\bar{\sigma} \rightarrow 0} \Gamma_0 = 0$. Thus once again the semiclassical contribution leads to a maximum value for the probability located at $\bar{\sigma} = 0$. Furthermore, we note that in this semiclassical contribution the tension term will be the only one that contributes after taking the real part since the integral with a constant scalar field gives always imaginary numbers.

Moving further, for the first quantum correction we obtain

$$\nabla^2 f = V''. \quad (108)$$

Then, computing the remaining terms and substituting back in (35) we obtain for the first quantum correction

$$\Gamma_1 = \frac{\text{Vol}(X)}{2} \left[\left(\frac{V_B^{(2)}}{V_B} - \frac{V_A^{(2)}}{V_A} \right) \ln(\sigma) \Big|_{\sigma_0}^{\bar{\sigma} - \delta\sigma} \right] + \text{Vol}(X) T_1, \quad (109)$$

where we note again that there only appears one tension term, thus it only adds one extra parameter. Furthermore, we note that the divergence that appeared in the flat FLRW result in (65) appears again here and we are led to impose also the condition (66). Thus, we finally obtain

$$\Gamma_1 = \text{Vol}(X) T_1, \quad (110)$$

which is only a constant. Therefore, the general behaviour is unaltered by the first quantum correction.

Carrying out with the analysis to the second quantum correction, we note that for this metric

$$\nabla^2 (\nabla^2 f) = \frac{\gamma}{\sigma^2} V^{(4)}, \quad (111)$$

$$(\nabla (\nabla^2 f))^2 = \frac{\gamma}{\sigma^2} (V^{(3)})^2. \quad (112)$$

Therefore, computing the remaining necessary terms we obtain from (37) that the second quantum correction term is written as

$$\begin{aligned} \Gamma_2 = \pm \frac{\text{Vol}(X) i \hbar}{4} & \left\{ \frac{V_B^{(4)}}{V_B} H_{KS}(c_B, \sigma) \Big|_{\sigma_0}^{\bar{\sigma}} + \left(\frac{V_B^{(3)}}{V_B} \right)^2 I_{KS}(c_B, \sigma) \Big|_{\sigma_0}^{\bar{\sigma}} - \frac{V_A^{(4)}}{V_A} H_{KS}(c_A, \sigma) \Big|_{\sigma_0}^{\bar{\sigma}} \right. \\ & \left. - \left(\frac{V_A^{(3)}}{V_A} \right)^2 I_{KS}(c_A, \sigma) \Big|_{\sigma_0}^{\bar{\sigma}} \right\} + \frac{\text{Vol}(X) \hbar}{\bar{\sigma}} \sqrt{\frac{V_B}{1 - c_B \bar{\sigma}}} T_2, \end{aligned} \quad (113)$$

where we have defined the functions

$$H_{KS}(c, \sigma) = \int \frac{d\sigma}{\sigma^2 \sqrt{4 - 3c\sigma}} = -\frac{\sqrt{4 - 3c\sigma}}{4\sigma} - \frac{3c}{8} \operatorname{arctanh} \left(\frac{\sqrt{4 - 3c\sigma}}{2} \right), \quad (114)$$

$$I_{KS}(c, \sigma) = \int \frac{1 - c\sigma}{\sigma^2 (4 - 3c\sigma)^{3/2}} d\sigma = -\frac{1}{16} \left[\frac{4 - c\sigma}{\sigma \sqrt{4 - 3c\sigma}} + \frac{c}{2} \operatorname{arctanh} \left(\frac{\sqrt{4 - 3c\sigma}}{2} \right) \right]. \quad (115)$$

Furthermore, we note from this result that the functions H_{KS} and I_{KS} have a divergence if we choose $\sigma_0 = 0$, but this divergence is independent of c . Therefore, in order to have a well defined probability we can eliminate those divergences by imposing the conditions

$$\frac{V_B^4}{V_B} = \frac{V_A^4}{V_A}, \quad \left(\frac{V_B^3}{V_B} \right)^2 = \left(\frac{V_A^3}{V_A} \right)^2. \quad (116)$$

We note that as in all the previous cases all the possible tension terms reduce to only one and that it is written in such a way that $\lim_{\bar{\sigma} \rightarrow 0} \Gamma_2$ diverges, therefore choosing the appropriate signs the probability of having a universe created at $\bar{\sigma} = 0$ vanishes. However, we note that as in the Bianchi III result we are not avoiding the singularity, we are just preventing this point which has a different physical meaning. Furthermore, when we plot the transition probabilities for the semiclassical contribution in (106), the first quantum correction in (110) and the second quantum correction in (113) we obtain the same qualitative behaviour as the one encountered for the FLRW metric in Figure 1.

7 Final Remarks

In this article we have studied the quantum corrections to the vacuum transitions probabilities between two minima of a scalar field potential employing a Lorentzian

formalism in the presence of gravity. First of all, we expanded a method previously proposed in [29,30] to describe these probabilities by computing solutions of the WDW equation using a semiclassical expansion of the WKB type. We have shown how to take into account the quantum corrections to the general method for any model in the superspace. We note that each quantum correction corresponds to considering one higher-order term in the \hbar expansion of the WKB proposal. Then, the WDW equation provides one equation for every term in the semiclassical expansion considered. Therefore, in principle, the system of equations will always be solvable. Thus this method can account for any desired order in the quantum corrections. Moreover, by choosing the parameter defining the integral curves in the superspace in an appropriate form as has been used in previous works, and applying some general considerations, we have obtained explicit analytical expressions for the semiclassical contributions and up to second order quantum corrections in (34), (35) and (37). One important feature of these general expressions is that all of them have the volume of the spatial slice as an overall constant, a feature that we expect will be maintained regardless of the order of the quantum correction that we wish to compute and that allows to compute the transition probabilities for metrics with a noncompact spatial slice. Furthermore, all expressions are described in terms of one degree of freedom coming from the metric evaluated at the point where the transition is made. Thus, the interpretation of the probability functions obtained can still be stated as describing probability distributions of creating universes with a given size as was done for the semiclassical contributions on previous works. However, taking into account the quantum corrections, new tension terms were added in increasing numbers. These terms arrived from the quantum fluctuations of the scalar field and may not be independent if the functions on the metric accompanying them are proportional or vanish. For example, on all the metrics considered in the main text only one tension term was added for each order in the expansion.

Then, we applied our method to some metrics of cosmological interest using General Relativity. We found analytical expressions for the transition amplitudes for homogeneous isotropic and anisotropic metrics. These expressions include the semiclassical contributions and two quantum corrections and are described by the potential minima and its derivatives, tension terms, and only one variable from the metric that represents the size of the universe at the time of nucleation.

For an isotropic universe we studied the FLRW metric with positive and zero curvature. In order to obtain well defined probabilities, the quantum corrections implied extra restrictions on the derivatives of the potential evaluated at the minima. The general behaviour of the transition probabilities was shown in Figure 1. It was shown that the semiclassical contribution leads to a probability that starts at its maximum value on the spatial singularity and falls to zero as the scale factor increases. The first quantum correction does not change the overall shape, it only reduces the probability. On the other hand, the second quantum correction changes the behaviour in the singularity, leading to the prediction that the maximum probability is located at small size of the scale factor but different from zero, in this way the quantum corrections leads to an avoidance of the initial singularity. We also noted that the quantum correction terms are relevant on the ultraviolet described by small values of the scale factor but

they do not change the behaviour in the infrared as we expected.

For an anisotropic universe we studied the Bianchi III and the Kantowski-Sachs metrics. For the Bianchi metric we encountered the same general form of the transition probability as in the isotropic case. However, the point of maximum probability in the semiclassical contribution that is avoided in the second quantum correction in this case does not represent an initial singularity. The different scale factors are related as a consequence of the integral curves on superspace in such a way that such point describes a scenario where two cartesian directions have infinite value for their scale factor whereas the third direction has a zero scale factor. However, if we factorize some terms in the Hamiltonian constraint we can obtain new Hamiltonians that are equivalent at the classical level but at the quantum level they lead to different forms of the WDW equation and different relations of the scale factors. This ambiguity on the correct choice for the Hamiltonian constraint is equivalent to the ordering ambiguity in the quantization procedure for the squared momentum. However, as we showed in the Appendix A, once we obtain a proper factorization that allows a initial spatial singularity, the avoidance of such point is lost and we only obtain transition probabilities that leads to a maximum value at the initial singularity. Thus, the avoidance of the spatial singularity due to quantum corrections to the transition probabilities can only be obtained in the isotropic universe. Furthermore, at the classical level the limit $\alpha \rightarrow 0$ of the Bianchi III metric results in the flat FLRW metric. We encountered that the results from the semiclassical contribution indeed lead to the flat FLRW result in such isotropy limit. However, the quantum corrections lead to a very similar result but differs from the flat FLRW result by some constants. Thus, the different structure of the superspace from these two metrics is revealed only after considering quantum corrections. Furthermore, by varying the parameter α we found that the effect of anisotropy, even adding quantum corrections, is to decrease the transition probability, preserving the form that was observed for the FLRW case of positive curvature. On the other hand, for the Kantowski-Sachs metric we also encountered the same general behaviour for the transition probabilities as for the previous cases. But once again the point that gets avoided represents a point where the radial part of the metric is set to zero but not the angular part, thus it is not an spatial singularity.

Let us remark, that in the Euclidean formalism the computation of the quantum corrections to the transition probabilities is troublesome. However, in the Lorentzian formalism we could compute such corrections in a fairly simple form by considering higher order terms in the semiclassical expansion. It will be interesting to investigate if some of the properties outlined in this article such as the avoidance of the initial singularity for the isotropic universe can also be encountered on the Euclidean formalism or if this represent a relevant departure from both approaches.

Acknowledgments

J. H. Aguilar (No. 933342) would like to thank CONAHCyT for a grant. He also wants to thank Cinvestav for its hospitality.

A Effects of a given factorization to the transition probabilities

In order to determine the effect of choosing a different factorization for the Hamiltonian constraint in the Bianchi III and flat FLRW metrics, let us consider a general factorization of the Hamiltonian of the Bianchi III metric (77) in the form

$$\begin{aligned}
H = & NA^{-\gamma}B^{-\delta}C^{-\zeta} \left[\frac{A^{\gamma+1}B^{\delta-1}C^{\zeta-1}}{4}\pi_A^2 + \frac{B^{\delta+1}A^{\gamma-1}C^{\zeta-1}}{4}\pi_B^2 + \frac{C^{\zeta+1}A^{\gamma-1}B^{\delta-1}}{4}\pi_C^2 \right. \\
& - \frac{A^\gamma B^\delta C^{\zeta-1}}{2}\pi_A\pi_B - \frac{A^\gamma C^\zeta B^{\delta-1}}{2}\pi_A\pi_C - \frac{A^{\gamma-1}B^\delta C^\zeta}{2}\pi_B\pi_C + \frac{A^{\gamma-1}B^{\delta-1}C^{\zeta-1}}{2}\pi_\phi^2 \\
& \left. + \alpha^2 A^{\gamma-1}B^{\delta+1}C^{\zeta+1} + V(\phi)A^{\gamma+1}B^{\delta+1}C^{\zeta+1} \right] \simeq 0,
\end{aligned} \tag{117}$$

where γ , δ and ζ are integers. We now consider the Hamiltonian constraint to be the terms within brackets, then after identifying the metric and the corresponding f function we can proceed to relate the degrees of freedom of the metric when the scalar field is constant by employing eq. (21). In this case we obtain

$$\frac{dA}{dB} = \left(\frac{A}{B} \right) \frac{(\gamma - 1 - \delta - \zeta)V + \frac{\alpha^2}{A^2}(\gamma - 3 - \delta - \zeta)}{(\delta - 1 - \gamma - \zeta)V + \frac{\alpha^2}{A^2}(\delta + 1 - \gamma - \zeta)}, \tag{118}$$

and

$$\frac{dB}{dC} = \left(\frac{B}{C} \right) \frac{(\delta - 1 - \gamma - \zeta)V + \frac{\alpha^2}{A^2}(\delta + 1 - \gamma - \zeta)}{(\zeta - 1 - \gamma - \delta)V + \frac{\alpha^2}{A^2}(\zeta + 1 - \gamma - \delta)}. \tag{119}$$

The second relation is very difficult to solve because of the appearance of the A function. We can simplify this expression by choosing $\delta = \zeta$. Then we obtain

$$B = b_0 C \tag{120}$$

where b_0 is an integration constant that will be taken as 1. In the same way, it is found that

$$B = a_0 A^{\phi_1} \left[A^2 + \phi_2 \frac{\alpha^2}{V} \right]^{\phi_3} \tag{121}$$

where a_0 is another integration constant that will be taken as 1 and

$$\phi_1 = \frac{1 - \gamma}{\gamma - 3 - 2\zeta}, \quad \phi_2 = \frac{\gamma - 3 - 2\zeta}{\gamma - 1 - 2\zeta}, \quad \phi_3 = \frac{2(\zeta + 1)}{(2\zeta + 1 - \gamma)(2\zeta + 3 - \gamma)}. \tag{122}$$

Nevertheless, we want to obtain an isotropy limit that leads to the flat FLRW metric in order to correctly study the effect of anisotropy, that is we require $\lim_{\alpha=0} B = A$. This leads to the condition

$$\phi_1 + 2\phi_3 = 1, \tag{123}$$

which gives us the solution $\zeta = \gamma$. In this way, the most general factorization requires $\gamma = \zeta = \delta$. So the above definitions are simplified as

$$\phi_1 = \frac{\gamma - 1}{\gamma + 3}, \quad \phi_2 = \frac{\gamma + 3}{\gamma + 1}, \quad \phi_3 = \frac{2}{\gamma + 3}. \tag{124}$$

However, as we remarked in the main text the motivation to consider a different factorization is to have access to an initial singularity for the Bianchi III metric, that is that we can have $\lim_{A \rightarrow 0} B = 0$. We note from eq. (121) that this behaviour can be obtained by requiring that $\phi_1 > 0$, which leads to

$$\gamma > 1, \quad \text{or} \quad \gamma < -3. \quad (125)$$

Furthermore, the general form of the metric in minisuperspace is

$$(G^{MN}) = \frac{1}{2} \begin{pmatrix} A^{\gamma+1}(BC)^{\gamma-1} & -(AB)^\gamma C^{\gamma-1} & -(AC)^\gamma B^{\gamma-1} & 0 \\ -(AB)^\gamma C^{\gamma-1} & B^{\gamma+1}(AC)^{\gamma-1} & -(BC)^\gamma A^{\gamma-1} & 0 \\ -(AC)^\gamma B^{\gamma-1} & -(BC)^\gamma A^{\gamma-1} & C^{\gamma+1}(AB)^{\gamma-1} & 0 \\ 0 & 0 & 0 & 2(ABC)^{\gamma-1} \end{pmatrix} \quad (126)$$

and we can identify

$$f = H(A, B, C) + F(A, B, C)V(\phi), \quad (127)$$

with

$$H(A, B, C) = \alpha^2 A^{\gamma-1} (BC)^{\gamma+1}, \quad F(A, B, C) = (ABC)^{\gamma+1}. \quad (128)$$

With these considerations we can proceed to compute the transition probabilities with up to second quantum corrections using the general expressions (34), (35) and (37). However, we note that all the integrals on the three expressions will vanish in the limit $\bar{A} \rightarrow A_0 = 0$. Therefore, the behaviour of the transition probabilities in the singularity is dictated only by the tension terms. Thus, let us analyse the resulting tension terms due the three contributions.

Following (34) the tension term for the semiclassical contribution takes the form

$$\frac{\text{Vol}(X)}{\hbar} \bar{A}^{-\frac{(\gamma+1)(3\gamma+1)}{\gamma+3}} \left[\bar{A}^2 + \frac{\gamma+3}{\gamma+1} \frac{\alpha^2}{V_B} \right]^{\frac{4(\gamma+1)}{\gamma+3}} T_0. \quad (129)$$

From this expression we note that the important quantity to keep track in order to study the behaviour in the initial singularity is given by

$$\beta_0 = \frac{(\gamma+1)(3\gamma+1)}{\gamma+3}. \quad (130)$$

Furthermore, taking the limit $\alpha \rightarrow 0$ of (129) we are lead consistently to the tension term of the flat FLRW metric

$$\frac{\text{Vol}(X)}{\hbar} \bar{a}^{3(\gamma+1)} T_0, \quad (131)$$

from where we note that relevant quantity is

$$\eta_0 = 3(\gamma+1). \quad (132)$$

In the same way, following (35) for the first quantum correction Γ_1 we obtain only one tension term written as

$$\text{Vol}(X) \bar{A}^{-\frac{2\gamma(3\gamma+1)}{\gamma+3}} \left[\bar{A}^2 + \frac{\gamma+3}{\gamma+1} \frac{\alpha^2}{V_B} \right]^{\frac{8\gamma}{\gamma+3}} T_1. \quad (133)$$

In the previous equation (133) we see that the relevant quantity to study the initial singularity is

$$\beta_1 = \frac{2\gamma(3\gamma + 1)}{\gamma + 3}. \quad (134)$$

One again taking the limit $\alpha \rightarrow 0$ we obtain the corresponding tension term for the flat FLRW metric

$$\text{Vol}(X)\bar{a}^{6\gamma}T_1, \quad (135)$$

where the relevant quantity is

$$\eta_1 = 6\gamma. \quad (136)$$

Finally, for the second quantum correction (37), we obtain only 3 different tension terms in the form

$$\begin{aligned} \text{Vol}(X)\hbar \left\{ \bar{A}^{-\frac{9\gamma^2-1}{\gamma+3}} \left[\bar{A}^2 + \frac{\gamma+3}{\gamma+1} \frac{\alpha^2}{V_B} \right]^{\frac{4(3\gamma-1)}{\gamma+3}} T_{2,1} + \bar{A}^{-\frac{15\gamma^2-7}{\gamma+3}} \left[\bar{A}^2 + \frac{\gamma+3}{\gamma+1} \frac{\alpha^2}{V_B} \right]^{\frac{4(5\gamma-1)}{\gamma+3}} T_{2,2} \right. \\ \left. + \bar{A}^{-\frac{(5\gamma-1)(3\gamma+1)}{\gamma+3}} \left[\bar{A}^2 + \frac{\gamma+3}{\gamma+1} \frac{\alpha^2}{V_B} \right]^{\frac{4(5\gamma-1)}{\gamma+3}} T_{2,3} \right\}. \end{aligned} \quad (137)$$

Therefore, the relevant quantities for the second correction are

$$\beta_2 = \frac{9\gamma^2 - 1}{\gamma + 3}, \quad \beta_3 = \frac{15\gamma^2 - 7}{\gamma + 3}, \quad \beta_4 = \frac{(5\gamma - 1)(3\gamma + 1)}{\gamma + 3}. \quad (138)$$

In this case in order to obtain correctly the FLRW flat result we need to take $\alpha \rightarrow 0$ and eliminate the second tension term since the functions depending on the scalar field vanish in this limit. Thus we are led to

$$\text{Vol}(X)a^{3(3\gamma-1)}T_{2,1} + \text{Vol}(X)a^{3(5\gamma-1)}T_{2,3}, \quad (139)$$

from where we identify

$$\eta_2 = 3(3\gamma - 1), \quad \eta_3 = 3(5\gamma - 1). \quad (140)$$

Therefore, the singularity will be the point of maximum probability for β_i and η_i positive, the general form of the probability distribution will not change for zero values of these parameters, whereas the singularity will be avoided if at least one of the β_i (for the Bianchi III metric) or one of the η_i (for the flat FLRW metric) is negative.

As stated before we are interested in exploring these behaviours for the regions where we can explore the initial singularity, that is $\gamma > 1$ or $\gamma < -3$. However, it can be shown that all β_i and η_i are positive for $\gamma > 1$ whereas all of these coefficients are negative for $\gamma < -3$. Since taking negative values for η_0 will change drastically our semiclassical result, we will not consider the region $\gamma < -3$. Therefore, focusing on $\gamma > 1$ we note that all coefficients are positive, thus the singularity will remain as the point with maximum probability even after taking up to second quantum correction terms. Therefore, we conclude that the avoidance of the initial singularity by taking quantum corrections can only be performed in the isotropic universe.

References

- [1] S. R. Coleman, “The Fate of the False Vacuum. 1. Semiclassical Theory,” Phys. Rev. D **15** (1977), 2929-2936 [erratum: Phys. Rev. D **16** (1977), 1248] doi:10.1103/PhysRevD.16.1248
- [2] C. G. Callan, Jr. and S. R. Coleman, “The Fate of the False Vacuum. 2. First Quantum Corrections,” Phys. Rev. D **16** (1977), 1762-1768 doi:10.1103/PhysRevD.16.1762
- [3] S. R. Coleman and F. De Luccia, “Gravitational Effects on and of Vacuum Decay,” Phys. Rev. D **21** (1980), 3305 doi:10.1103/PhysRevD.21.3305
- [4] S. J. Parke, “Gravity, the Decay of the False Vacuum and the New Inflationary Universe Scenario,” Phys. Lett. B **121** (1983), 313-315 doi:10.1016/0370-2693(83)91376-X
- [5] L. G. Jensen and P. J. Ruback, “Bubble Formation in Anisotropic Cosmologies,” Nucl. Phys. B **325** (1989), 660-686 doi:10.1016/0550-3213(89)90502-6
- [6] D. A. Samuel and W. A. Hiscock, “Effect of gravity on false vacuum decay rates for O(4) symmetric bubble nucleation,” Phys. Rev. D **44** (1991), 3052-3061 doi:10.1103/PhysRevD.44.3052
- [7] A. q. Wu, “Tunneling in the presence of gravity,” Phys. Rev. D **46** (1992), 5321-5330 doi:10.1103/PhysRevD.46.5321
- [8] R. Mansouri and M. Mohazzab, “Tunneling in anisotropic cosmological models,” Class. Quant. Grav. **10** (1993), 1353-1359 doi:10.1088/0264-9381/10/7/011 [arXiv:gr-qc/9211024 [gr-qc]].
- [9] U. Gen and M. Sasaki, “False vacuum decay with gravity in nonthin wall limit,” Phys. Rev. D **61** (2000), 103508 doi:10.1103/PhysRevD.61.103508 [arXiv:gr-qc/9912096 [gr-qc]].
- [10] R. G. Cai, B. Hu and S. Koh, “Gauss-Bonnet Term on Vacuum Decay,” Phys. Lett. B **671** (2009), 181-186 doi:10.1016/j.physletb.2008.11.053 [arXiv:0806.2508 [hep-th]].
- [11] S. Kanno and J. Soda, “Exact Coleman-de Luccia Instantons,” Int. J. Mod. Phys. D **21** (2012), 1250040 doi:10.1142/S021827181250040X [arXiv:1111.0720 [hep-th]].
- [12] B. Salehian and H. Firouzjahi, “Vacuum decay and bubble nucleation in $f(R)$ gravity,” Phys. Rev. D **99** (2019) no.2, 025002 doi:10.1103/PhysRevD.99.025002 [arXiv:1810.01391 [hep-th]].
- [13] N. Oshita, Y. Shoji and M. Yamaguchi, “Polychronic tunneling: New tunneling processes experiencing Euclidean and Lorentzian evolution simultaneously,” Phys. Rev. D **107** (2023) no.4, 045007 doi:10.1103/PhysRevD.107.045007 [arXiv:2112.10736 [hep-th]].

- [14] S. Vicentini, “Vacuum decay and quadratic gravity,” doi:10.15168/11572_335510
- [15] R. L. Arnowitt, S. Deser and C. W. Misner, “The Dynamics of general relativity,” *Gen. Rel. Grav.* **40** (2008), 1997-2027 doi:10.1007/s10714-008-0661-1 [[arXiv:gr-qc/0405109](#) [gr-qc]].
- [16] A. Corichi and D. Núñez, “Introduction to the ADM formalism,” *Rev. Mex. Fis.* **37** (1991), 720-747 [[arXiv:2210.10103](#) [gr-qc]].
- [17] J. J. Halliwell, “Introductory Lectures on Quantum Cosmology,” [[arXiv:0909.2566](#)].
- [18] D. L. Wiltshire, “An Introduction to quantum cosmology,” [[arXiv:gr-qc/0101003](#) [gr-qc]].
- [19] S. Carlip, “Quantum gravity: A Progress report,” *Rept. Prog. Phys.* **64** (2001), 885 doi:10.1088/0034-4885/64/8/301 [[arXiv:gr-qc/0108040](#) [gr-qc]].
- [20] C. Kiefer and B. Sandhoefer, “Quantum cosmology,” *Z. Naturforsch. A* **77** (2022) no.6, 543-559 doi:10.1515/zna-2021-0384 [[arXiv:0804.0672](#) [gr-qc]].
- [21] C. J. Isham, “Canonical quantum gravity and the problem of time,” *NATO Sci. Ser. C* **409** (1993), 157-287 [[arXiv:gr-qc/9210011](#) [gr-qc]].
- [22] J.A. Wheeler, “Superspace and the nature of quantum geometrodynamics,” pp 615-724 of *Topics in Nonlinear Physics*, (ed) N.J. Zabusky, Springer-Verlag NY, Inc. (1969).
- [23] B.S. DeWitt, “Quantum theory of gravity I, The canonical theory,” *Phys. Rev.* **160** (5) (1967) 1113.
- [24] D. A. Craig and P. Singh, “Consistent Probabilities in Wheeler-DeWitt Quantum Cosmology,” *Phys. Rev. D* **82** (2010), 123526 doi:10.1103/PhysRevD.82.123526 [[arXiv:1006.3837](#) [gr-qc]].
- [25] L. Chataignier, C. Kiefer and P. Moniz, “Observations in quantum cosmology,” *Class. Quant. Grav.* **40** (2023) no.22 , 223001 doi:10.1088/1361-6382/acfa5b [[arXiv:2306.14948](#) [gr-qc]].
- [26] W. Fischler, D. Morgan and J. Polchinski, “Quantum Nucleation of False Vacuum Bubbles,” *Phys. Rev. D* **41** (1990), 2638 doi:10.1103/PhysRevD.41.2638
- [27] W. Fischler, D. Morgan and J. Polchinski, “Quantization of False Vacuum Bubbles: A Hamiltonian Treatment of Gravitational Tunneling,” *Phys. Rev. D* **42** (1990), 4042-4055 doi:10.1103/PhysRevD.42.4042
- [28] S. P. De Alwis, F. Muia, V. Pasquarella and F. Quevedo, “Quantum Transitions Between Minkowski and de Sitter Spacetimes,” *Fortsch. Phys.* **68** (2020) no.9 , 2000069 doi:10.1002/prop.202000069 [[arXiv:1909.01975](#) [hep-th]].

- [29] S. Cespedes, S. P. de Alwis, F. Muia and F. Quevedo, “Lorentzian vacuum transitions: Open or closed universes?,” *Phys. Rev. D* **104** (2021) no.2, 026013 doi:10.1103/PhysRevD.104.026013 [[arXiv:2011.13936](#) [hep-th]].
- [30] H. García-Compeán and D. Mata-Pacheco, “Lorentzian vacuum transitions for anisotropic universes,” *Phys. Rev. D* **104** (2021) no.10, 106014 doi:10.1103/PhysRevD.104.106014 [[arXiv:2107.07035](#) [hep-th]].
- [31] H. García-Compeán and D. Mata-Pacheco, “Lorentzian Vacuum Transitions in Hořava–Lifshitz Gravity,” *Universe* **8** (2022) no.4, 237 doi:10.3390/universe8040237 [[arXiv:2111.11571](#) [gr-qc]].
- [32] H. García-Compeán and D. Mata-Pacheco, “Lorentzian vacuum transitions with a generalized uncertainty principle,” *Class. Quant. Grav.* **39** (2022) no.23, 235011 doi:10.1088/1361-6382/ac9efc [[arXiv:2206.06534](#) [gr-qc]].
- [33] G. V. Lavrelashvili, V. A. Rubakov and P. G. Tinyakov, “Tunnelling Transitions with Gravitation: Breaking of the Quasiclassical Approximation,” *Phys. Lett. B* **161** (1985), 280-284 doi:10.1016/0370-2693(85)90761-0
- [34] L. F. Abbott, D. Harari and Q. H. Park, “Vacuum Decay in Curved Backgrounds,” *Class. Quant. Grav.* **4** (1987), L201-L204 doi:10.1088/0264-9381/4/6/001
- [35] G. V. Lavrelashvili, “Negative mode problem in false vacuum decay with gravity,” *Nucl. Phys. B Proc. Suppl.* **88** (2000), 75-82 doi:10.1016/S0920-5632(00)00756-8 [[arXiv:gr-qc/0004025](#) [gr-qc]].
- [36] M. Koehn, G. Lavrelashvili and J. L. Lehners, “Towards a Solution of the Negative Mode Problem in Quantum Tunnelling with Gravity,” *Phys. Rev. D* **92** (2015) no.2, 023506 doi:10.1103/PhysRevD.92.023506 [[arXiv:1504.04334](#) [hep-th]].
- [37] S. F. Bramberger, M. Chitishvili and G. Lavrelashvili, “Aspects of the negative mode problem in quantum tunneling with gravity,” *Phys. Rev. D* **100** (2019) no.12, 125006 doi:10.1103/PhysRevD.100.125006 [[arXiv:1906.07033](#) [gr-qc]].
- [38] R. Jinno and R. Sato, “Negative mode problem of false vacuum decay revisited,” *Phys. Rev. D* **104** (2021) no.9, 096009 doi:10.1103/PhysRevD.104.096009 [[arXiv:2010.04462](#) [hep-th]].
- [39] S. P. de Alwis, “Revisiting Vacuum decay in Field Theory,” [[arXiv:2305.11786](#) [hep-th]].
- [40] J. R. Espinosa, “Fresh look at the calculation of tunneling actions including gravitational effects,” *Phys. Rev. D* **100** (2019) no.10, 104007 doi:10.1103/PhysRevD.100.104007 [[arXiv:1808.00420](#) [hep-th]].
- [41] J. R. Espinosa, J. F. Fortin and J. Huertas, “Exactly solvable vacuum decays with gravity,” *Phys. Rev. D* **104** (2021) no.6, 065007 doi:10.1103/PhysRevD.104.065007 [[arXiv:2106.15505](#) [hep-th]].

- [42] G. Calcagni, M. Frasca and A. Ghoshal, “Fate of false vacuum in non-perturbative regimes: Gravity effects,” [[arXiv:2206.09965](#) [hep-th]].
- [43] J. R. Espinosa, R. Jinno and T. Konstandin, “Tunneling potential actions from canonical transformations,” JCAP **02** (2023), 021 doi:10.1088/1475-7516/2023/02/021 [[arXiv:2209.03293](#) [hep-th]].
- [44] J. R. Espinosa and J. F. Fortin, “Vacuum decay actions from tunneling potentials for general spacetime dimension,” JCAP **02** (2023), 023 doi:10.1088/1475-7516/2023/02/023 [[arXiv:2211.13667](#) [hep-th]].
- [45] B. Freivogel, Y. Sekino, L. Susskind and C. P. Yeh, “A Holographic framework for eternal inflation,” Phys. Rev. D **74** (2006), 086003 doi:10.1103/PhysRevD.74.086003 [[arXiv:hep-th/0606204](#) [hep-th]].
- [46] J. L. F. Barbon and E. Rabinovici, “Holography of AdS vacuum bubbles,” JHEP **04** (2010), 123 doi:10.1007/JHEP04(2010)123 [[arXiv:1003.4966](#) [hep-th]].
- [47] J. K. Ghosh, E. Kiritsis, F. Nitti and L. T. Witkowski, “Revisiting Coleman-de Luccia transitions in the AdS regime using holography,” JHEP **09** (2021), 065 doi:10.1007/JHEP09(2021)065 [[arXiv:2102.11881](#) [hep-th]].
- [48] V. Pasquarella and F. Quevedo, “Vacuum transitions in two-dimensions and their holographic interpretation,” JHEP **05** (2023), 192 doi:10.1007/JHEP05(2023)192 [[arXiv:2211.07664](#) [hep-th]].
- [49] J. Colin, R. Mohayaee, M. Rameez and S. Sarkar, “Evidence for anisotropy of cosmic acceleration,” Astron. Astrophys. **631** (2019), L13 doi:10.1051/0004-6361/201936373 [[arXiv:1808.04597](#) [astro-ph.CO]].
- [50] K. Migkas, G. Schellenberger, T. H. Reiprich, F. Pacaud, M. E. Ramos-Ceja and L. Lovisari, “Probing cosmic isotropy with a new x-ray galaxy cluster sample through the $L_X - T$ scaling relation,” Astron. Astrophys. **636** (2020), A15 doi:10.1051/0004-6361/201936602 [[arXiv:2004.03305](#) [astro-ph.CO]].
- [51] C. W. Misner, “Minisuperspace,” edited by J.R. Klauder, *Magic Without Magic* John Archibald Wheeler, A collection in honor of his 60th birthday, San Francisco: W. H. Freeman, pages 441-473 (1972).