

Semi-supervised Regression Analysis with Model Misspecification and High-dimensional Data

Ye Tian,¹ Peng Wu,² and Zhiqiang Tan³

June 21, 2024

Abstract. The accessibility of vast volumes of unlabeled data has sparked growing interest in semi-supervised learning (SSL) and covariate shift transfer learning (CSTL). In this paper, we present an inference framework for estimating regression coefficients in conditional mean models within both SSL and CSTL settings, while allowing for the misspecification of conditional mean models. We develop an augmented inverse probability weighted (AIPW) method, employing regularized calibrated estimators for both propensity score (PS) and outcome regression (OR) nuisance models, with PS and OR models being sequentially dependent. We show that when the PS model is correctly specified, the proposed estimator achieves consistency, asymptotic normality, and valid confidence intervals, even with possible OR model misspecification and high-dimensional data. Moreover, by suppressing detailed technical choices, we demonstrate that previous methods can be unified within our AIPW framework. Our theoretical findings are verified through extensive simulation studies and a real-world data application.

Key words and phrases: Augmented inverse probability weighted estimator; Covariate shift transfer learning; High-dimensional data; Semi-supervised learning.

¹Department of Statistics, Rutgers University, Piscataway, NJ 08854, USA (E-mail yt334@stat.rutgers.edu).

²Department of Applied Statistics, Beijing Technology and Business University, Beijing, 102488, China
(E-mail: pengwu@btbu.edu.cn).

³Department of Statistics, Rutgers University, Piscataway, NJ 08854, USA (E-mail: ztan@stat.rutgers.edu).

1 Introduction

In recent years, vast volumes of unlabeled data have become increasingly accessible, sparking growing interest in how to leverage these data in both academic research and industry applications. One of the active areas of research is semi-supervised learning (SSL). In addition, covariate shift transfer learning (CSTL) also exploits information from unlabeled target data. Both have various application scenarios like computer vision (Sohn et al., 2020; Zhou and Levine, 2021; Zheng et al., 2022), natural language process (Chen and Huang, 2016; Ruder et al., 2019; Zhao et al., 2022), causal inference (Alvari et al., 2019; Aloui et al., 2023; Zhang et al., 2023), healthcare data analysis (Castro et al., 2020; Liu et al., 2023; Tang et al., 2024), etc.

In both the SSL and CSTL settings, we have access to a labeled dataset \mathcal{L} and an unlabeled dataset \mathcal{U} , where the labeled dataset \mathcal{L} contains observations with both the covariates \mathbf{X} and the outcome Y , while the unlabeled dataset \mathcal{U} consists solely of observations with the covariates \mathbf{X} . The training set \mathcal{T} is the union of \mathcal{L} and \mathcal{U} . Nevertheless, there is a key distinction between the classical SSL and CSTL setups (Chapelle et al., 2006; Liu et al., 2023). In CSTL, the conditional distributions of Y given \mathbf{X} in the labeled and unlabeled datasets are assumed to be the same, whereas the marginal distributions of \mathbf{X} are different (hence the term covariate shift), and the estimator is ultimately evaluated on unlabeled data. However, under the classical SSL setup, it is assumed that the distributions of labeled data, unlabeled data, and population are the same, making no difference in evaluating the estimator on which distribution. To accommodate both SSL and CSTL, we consider a more general setting. We only assume the conditional distribution of Y given \mathbf{X} is the same in labeled and unlabeled data, while marginal distributions of \mathbf{X} are permitted to be different. Estimators evaluated on the population and unlabeled data are both considered.

While there is a long history of SSL (Chapelle et al., 2006; Zhu, 2008) and CSTL (Quiñonero-Candela et al., 2009), a growing literature has considered inference procedures only recently. Notable advancements have been made in estimating the population mean $\mathbb{E}(Y)$ (Zhang et al., 2019; Zhang and Bradic, 2021) and regression coefficients in (generalized) linear models (Chakraborty, 2016; Chakraborty and Cai, 2018; Liu et al., 2023). Inferences of quantile regression (Chakraborty et al., 2022), explained variance (Cai and Guo, 2020), and model performance metrics such as true and false positive rates (Gronsbell and Cai, 2017) have also attracted interest.

In this article, we focus on the inference of regression coefficients in (conditional) mean models in SSL settings, hence called semi-supervised regression analysis. We demonstrate a unified framework

for estimating and inferring these coefficients, particularly in cases where the (conditional) mean model and outcome regression (OR) model $E[Y|\mathbf{X}]$ may be misspecified. Previous SSL and CSTL methods that considered the same goal, such as Chakraborty (2016), Chakraborty and Cai (2018), Zhang et al. (2019), Zhang and Bradic (2021) and Liu et al. (2023), can largely be accommodated in the augmented inverse probability weighting (AIPW) framework (Robins et al., 1994; Tan, 2020a; Wu et al., 2024). See Section 5 and Section 8 for further details.

Despite significant advancements made, there remain limitations in the previous AIPW methods. Methods developed in SSL settings usually treat the problem as the one where data are missing completely at random (MCAR) (Chakraborty, 2016; Chakraborty and Cai, 2018; Zhang et al., 2019; Zhang and Bradic, 2021). This restricts their application scenarios and overlooks the significance of constructing the propensity score (PS) model. In contrast, our setting is in general a missing-at-random (MAR) problem (Little and Rubin, 2019), and the estimation of the PS model is no longer negligible. In the setting of MCAR, the PS remains a constant, whereas in the setting of MAR, the PS varies with the covariates \mathbf{X} . In MAR problems, similarly as in Tan (2020a), the estimation of PS and OR models needs to be carefully handled in a way different from regularized least squares or maximum likelihood as in previous papers, so that \sqrt{N} -consistent estimation can be achieved with possible misspecification of the OR model, where N is the sample size of \mathcal{T} .

In summary, we mainly make the following two contributions. First, we present an inference framework that consolidates several previous settings, including the estimation of population mean and regression coefficients in conditional mean models of Y given any sub-vector of \mathbf{X} under both SSL and CSTL setups. Second, we propose a novel AIPW method that enables \sqrt{N} -consistent and asymptotically normal estimation and achieves valid confidence intervals (CIs), when the PS model is correctly specified but the OR model may be misspecified. This robustness to model misspecification is achieved by carefully exploiting the connection between PS and OR models and designing estimating equations for nuisance parameters, differently from regularized least-squares or maximum-likelihood estimation. The desirable properties of our method, \sqrt{N} -consistency, asymptotical normality, and valid confidence intervals, are formally established in the setting of sparse high-dimensional PS and OR models, while allowing that the estimator of PS model has convergence rates slower than $1/\sqrt{N}$ and the OR model may be misspecified.

This work is also related to the causal inference problem under the strong ignorability assumption (Rosenbaum and Rubin, 1983). Specifically, the SSL problem is similar to estimating the average treatment effect (ATE) and the conditional ATE (CATE) (Wager and Athey, 2018; Athey

et al., 2019; Zimmert and Lechner, 2019; Fan et al., 2022; Wu et al., 2024). The CSTL problem can be viewed as an analog to the estimation of the average treatment effect on the treated (ATT).

The rest of this paper is organized as follows. In section 2, we present our setup and define the target parameters of interest. In Section 3, we construct a novel AIPW estimator for the target parameter under the SSL setting. We show theoretical properties of the proposed estimator in Section 4, and compare them with the previous literature in Section 5. Simulation results are shown in Section 6, and an application to crime study is presented in Section 7. Extension of proposed methods to the CSTL setting is given in Section 8 followed by concluding discussions in Section 9. Proofs of all theoretical results and associated technical materials are presented in the Supplementary Material.

2 Setup and preliminaries

2.1 Data and target parameters

Let $Y \in \mathbb{R}$ be a response variable and $\mathbf{X} = (1, X_1, \dots, X_d)^\top \in \mathbb{R}^{d+1}$ be a covariate vector with the first element being the constant 1. In addition, let $R \in \{0, 1\}$ be the indicator of whether Y is observed: $R = 1$ if observed and $R = 0$ if missing. Assume that $\{(\mathbf{X}_i, Y_i, R_i) : i = 1, \dots, N\}$ is an independent and identically distributed (i.i.d.) sample from a joint distribution of (\mathbf{X}, Y, R) , denoted as \mathbb{P} . The observed dataset, $\{(\mathbf{X}_i, R_i Y_i, R_i) : i = 1, \dots, N\}$, can be split into a labeled dataset and an unlabeled dataset as follows:

$$\mathcal{L} = \{(\mathbf{X}_i, Y_i, R_i = 1), i = 1, \dots, n\}, \quad \mathcal{U} = \{(\mathbf{X}_i, R_i = 0), i = n + 1, \dots, N\}.$$

For \mathbf{Z} a sub-vector of \mathbf{X} , it is of interest to fit a regression model for the conditional mean $\mathbb{E}(Y|\mathbf{Z})$:

$$\mathbb{E}(Y|\mathbf{Z}) = \psi(\beta^{*\top} \mathbf{Z}), \tag{1}$$

where $\mathbb{E}(\cdot)$ denotes the expectation under \mathbb{P} , β^* is a parameter vector, and $\psi(\cdot)$ is an (increasing) inverse link function, such as the identity function $\psi(u) = u$ and logit function $\psi(u) = 1/\{1 + \exp(-u)\}$. The model (1) is allowed to be misspecified, that is, $\mathbb{E}(Y|\mathbf{Z})$ may not be put in the form $\psi(\beta^\top \mathbf{Z})$. With possible model misspecification, β^* is defined as the solution to the estimating equations:

$$\mathbb{E}[\{Y - \psi(\beta^\top \mathbf{Z})\} \mathbf{Z}] = 0. \tag{2}$$

For a generalized linear model with $\psi(\cdot)$ as the canonical inverse link, the estimating equation (2) leads to maximum likelihood estimation, so that $\psi(\beta^{*\text{T}} \mathbf{Z})$ can be interpreted as the best likelihood-based approximation to $\mathbb{E}(Y|\mathbf{Z})$ using model (1).

The regression model (1) is flexible. The target parameter β^* accommodates a variety of estimands in the previous literature.

- (a) If we set $\mathbf{Z} = 1$ and $\psi(u) = u$, then $\beta^* = \mathbb{E}(Y)$. The problem corresponds to the semi-supervised estimation of the population mean (Zhang et al., 2019; Zhang and Bradic, 2021).
- (b) If we set \mathbf{Z} to be a univariate covariate, for example, $\mathbf{Z} = X_1$, then β^* corresponds to the regression coefficient in the regression model of Y given the particular covariate X_1 . This problem was studied by Liu et al. (2023) and Wu et al. (2024).
- (c) If we set $\mathbf{Z} = \mathbf{X}$, then β^* corresponds to the coefficient vector in the regression model of Y given the full covariate vector \mathbf{X} (Chakraborty, 2016; Chakraborty and Cai, 2018).

In addition to the parameter vector β^* , it may also be of interest to consider target parameters defined within unlabeled data corresponding to the CSTL setting, as studied in several recent papers (Liu et al., 2023; He et al., 2024). To incorporate this setting, we consider a regression model for the conditional mean in the unlabeled data:

$$\mathbb{E}(Y|R = 0, \mathbf{Z}) = \psi(\beta^{0*\text{T}} \mathbf{Z}). \quad (3)$$

With possible misspecification of model (3), the parameter β^{0*} is defined as the solution to

$$\mathbb{E}[(1 - R) \{Y - \psi(\beta^{\text{T}} \mathbf{Z})\} \mathbf{Z}] = 0. \quad (4)$$

To illustrate the main ideas, we focus on the estimation of β^* , and defer the associated results for the estimation of β^{0*} to Section 8.

2.2 General assumptions

Without imposing any assumption, we cannot obtain a consistent estimator of β^* due to the missingness of Y in the unlabeled data. Below, we introduce the identifiability assumption.

Assumption 1. $Y \perp\!\!\!\perp R \mid \mathbf{X}$, *i.e.*, Y and R are conditionally independent given \mathbf{X} .

Assumption 1 is crucial for the identifiability of β^* and β^{0*} . It ensures $\mathbb{E}(Y|\mathbf{X}, R = 1) = \mathbb{E}(Y|\mathbf{X}, R = 0)$, indicating that the conditional mean of Y is the same for both unlabeled and

labeled data after accounting for the full covariates \mathbf{X} . This establishes the connection between the labeled and unlabeled data. Moreover, Assumption 1 implies that $\mathbb{P}(R = 1|\mathbf{X}, Y) = \mathbb{P}(R = 1|\mathbf{X})$, meaning that the label indicator R depends solely on the covariates \mathbf{X} , i.e., R is missing at random (Molenberghs et al., 2015; Imbens and Rubin, 2015). It should be noted that Assumption 1 does not imply $Y \perp\!\!\!\perp R | \mathbf{Z}$ when $\mathbf{Z} \neq \mathbf{X}$. If $Y \perp\!\!\!\perp R | \mathbf{Z}$ does hold and models (1) and (3) are correctly specified, then $\beta^* = \beta^{0*}$. Otherwise, β^* and β^{0*} may differ from each other.

Despite bearing many similarities, Assumption 1 differs from the classic SSL setup in the missing mechanism as represented by the probabilistic behavior of R . SSL assumes the missing-completely-at-random mechanism (MCAR) (Chakraborty, 2016; Chakraborty and Cai, 2018; Zhang et al., 2019; Zhang and Bradic, 2021), that is, $R \perp\!\!\!\perp (\mathbf{X}, Y)$ and thus $\mathbb{P}(R = 1|\mathbf{X})$ is a constant, independent of \mathbf{X} . In contrast, we allow R to probabilistically depend on the covariates \mathbf{X} . In addition, we make the following technical assumption.

Assumption 2. $\pi^*(\mathbf{X}) = \mathbb{P}(R = 1|\mathbf{X}) > 0$.

This condition is introduced to ensure that each unit has a positive probability of belonging to \mathcal{L} . Then the labeled dataset \mathcal{L} is of a non-negligible size compared with N .

By Assumption 2, as $N \rightarrow \infty$, the ratio n/N may randomly fluctuate but converges to a constant within the interval $(0, 1)$, equal to $\mathbb{E}(R) = \pi^*(\mathbf{X})$. This distinguishes our sampling process from the stratified sampling process widely used in the previous literature (Chakraborty, 2016; Chakraborty and Cai, 2018; Zhang et al., 2019; Zhang and Bradic, 2021), where the sizes of labeled and unlabeled datasets, n and $N - n$, are deterministic. For the asymptotic analysis, they assume that both n and N tend to infinity such that n/N converges to a constant in $[0, 1)$, including zero. For details, see the discussion in Section 5.

2.3 AIPW estimating equations

For estimating β^* , we introduce the augmented inverse probability weighting (AIPW) estimating equations. Essentially the same AIPW estimating equation has been used in the previous literature, albeit under somewhat different settings than ours. See Section 5 for a connection and comparison of our method with the previous methods.

With the true PS $\pi^*(\mathbf{X}) = \mathbb{P}(R = 1|\mathbf{X})$, then under Assumption 1, we have

$$\mathbb{E} \left[\frac{R}{\pi^*(\mathbf{X})} \{Y - \psi(\beta^T \mathbf{Z})\} \mathbf{Z} \right] = \mathbb{E} [\{Y - \psi(\beta^T \mathbf{Z})\} \mathbf{Z}].$$

Then a sample estimating equation for β^* is

$$\tilde{\mathbb{E}} \left[\frac{R}{\hat{\pi}(\mathbf{X})} \{Y - \psi(\beta^{\text{T}} \mathbf{Z})\} \mathbf{Z} \right] = 0, \quad (5)$$

where $\hat{\pi}(\mathbf{X})$ is an estimator of $\pi^*(\mathbf{X})$ and $\tilde{\mathbb{E}}$ denotes the sample mean, defined as $\tilde{\mathbb{E}}(U) = N^{-1} \sum_{i=1}^N U_i$ for a variable U .

Let $\hat{\beta}_{\text{IPW}}$ be a solution to equations (5). If the PS model is correctly specified, then under certain regularity conditions, $\hat{\pi}(\mathbf{X}) \xrightarrow{\mathbb{P}} \pi^*(\mathbf{X})$ and $\hat{\beta}_{\text{IPW}} \xrightarrow{\mathbb{P}} \beta^*$. However, if the PS model is misspecified, then $\hat{\pi}(\mathbf{X}) \not\xrightarrow{\mathbb{P}} \pi^*(\mathbf{X})$ and $\hat{\beta}_{\text{IPW}} \not\xrightarrow{\mathbb{P}} \beta^*$. To mitigate the possible inconsistency of $\hat{\beta}_{\text{IPW}}$, the AIPW method introduces an augmented term. Specifically, let $\phi^*(X) = \mathbb{E}[Y|\mathbf{X}]$ be the true OR function and $\hat{\phi}(\mathbf{X})$ be a corresponding estimator. Then the AIPW estimating equation is

$$\tilde{\mathbb{E}} \left[\frac{R}{\hat{\pi}(\mathbf{X})} \{Y - \psi(\beta^{\text{T}} \mathbf{Z})\} \mathbf{Z} + \left\{ 1 - \frac{R}{\hat{\pi}(\mathbf{X})} \right\} \left\{ \hat{\phi}(\mathbf{X}) - \psi(\beta^{\text{T}} \mathbf{Z}) \right\} \mathbf{Z} \right] = 0. \quad (6)$$

Let $\hat{\beta}_{\text{AIPW}}$ be the solution to equation (6). If the PS model is misspecified, the augmented term

$$\tilde{\mathbb{E}} \left[\left\{ 1 - \frac{R}{\hat{\pi}(\mathbf{X})} \right\} \left\{ \hat{\phi}(\mathbf{X}) - \psi(\beta^{\text{T}} \mathbf{Z}) \right\} \mathbf{Z} \right]$$

corrects the bias of $\tilde{\mathbb{E}} \left[\frac{R}{\hat{\pi}(\mathbf{X})} \{Y - \psi(\beta^{\text{T}} \mathbf{Z})\} \mathbf{Z} \right]$ by introducing the estimator $\hat{\phi}(\mathbf{X})$. In addition, if the PS model is correctly specified, the augmented term improves the estimation efficiency of β^* by leveraging the association between \mathbf{X} and Y . It can be shown that the left side of equation (6) converges in probability to that of equation (2), if either $\hat{\pi}(\mathbf{X}) \xrightarrow{\mathbb{P}} \pi^*(\mathbf{X})$ or $\hat{\phi}(\mathbf{X}) \xrightarrow{\mathbb{P}} \phi^*(\mathbf{X})$, which is the property of double robustness.

In the classic SSL setup, estimating β^* is considered to be an MCAR problem, where $\hat{\pi}(\mathbf{X})$ is a constant, independent of \mathbf{X} , and the estimator $\hat{\phi}(\mathbf{X})$ is usually defined using an OR model by (unweighted) least squares, maximum likelihood, or variations (Chakraborty, 2016; Chakraborty and Cai, 2018; Zhang et al., 2019). However, our semi-supervised regression is formulated as a MAR problem, where $\hat{\pi}(\mathbf{X})$ depends on \mathbf{X} . In such a scenario, as shown in the next section, the estimators $\hat{\pi}(\mathbf{X})$ and $\hat{\phi}(\mathbf{X})$ for the PS and OR functions can be defined in a sequential manner, different from least squares or maximum likelihood, in order to obtain desirable properties with possible model misspecification.

3 Method

We develop a novel AIPW method that achieves \sqrt{N} -consistency in the setting of sparse high-dimensional PS and OR models, even if the estimation of the PS model exhibits convergence rates

slower than $1/\sqrt{N}$ and the OR model is misspecified.

3.1 Model specification for nuisance parameters

AIPW estimation based on the estimating equation (6) requires constructing the estimators $\hat{\pi}(\mathbf{X})$ and $\hat{\phi}(\mathbf{X})$ for $\pi^*(\mathbf{X})$ and $\phi^*(\mathbf{X})$, using some PS and OR models. In contrast with the previous literature, we introduce a dependency between $\hat{\pi}(\mathbf{X})$ and $\hat{\phi}(\mathbf{X})$ by carefully specifying basis functions and incorporating weighted estimation.

Specifically, let $\mathbf{F}(\mathbf{X}) = \{1, f_1(\mathbf{X}), \dots, f_p(\mathbf{X})\}^\top$ be a vector of known functions of \mathbf{X} . We allow p to be high-dimensional, tending to infinity as N increases. As in Tan (2020a), we propose using logistic regression as a working model for the PS function $\pi^*(\mathbf{X})$,

$$\mathbb{P}(R = 1|\mathbf{X}) = \pi(\mathbf{X}; \gamma) = [1 + \exp\{-\gamma^\top \mathbf{F}(\mathbf{X})\}]^{-1}, \quad (7)$$

where γ is an unknown coefficient parameter.

Remark 1. In several related works on classic SSL and stratified sampling setups, making efforts to estimate the PS may not be necessary. Firstly, in the classic SSL setup, the true PS is a constant, leading to a constant PS model. Secondly, in the stratified sampling setup, the proportion n/N is fixed and known, which corresponds to a known PS function. Thus, researchers concentrate on specifying OR models. However, for our setup, both the PS model and the OR model must be carefully formulated to achieve desirable properties.

Next, we turn to modeling the OR function $\phi^*(\mathbf{X})$. The working model for $\phi^*(\mathbf{X})$ is specified as

$$\mathbb{E}[Y|\mathbf{X}] = \phi(\mathbf{X}; \alpha) = \psi\{\alpha^\top \mathbf{G}(\mathbf{X})\}, \quad (8)$$

where $\mathbf{G}(\mathbf{X}) = \{1, g_1(\mathbf{X}), \dots, g_q(\mathbf{X})\}^\top$ be a vector of known functions of \mathbf{X} and q can be high-dimensional. In contrast to the previous literature, to ensure valid inference even when the OR model is misspecified, we carefully specify a choice of $\mathbf{G}(\mathbf{X})$ as follows:

$$\mathbf{G}(\mathbf{X}) = [\mathbf{F}(\mathbf{X})^\top, \{\mathbf{Z} \otimes \mathbf{F}(\mathbf{X})\}^\top]^\top, \quad (9)$$

where $\mathbf{Z} \otimes \mathbf{F}(\mathbf{X})$ consists of all interactions between \mathbf{Z} and $\mathbf{F}(\mathbf{X})$ (i.e., all products of individual components from \mathbf{Z} and $\mathbf{F}(\mathbf{X})$). Equation (9) represents the minimal choice for $\mathbf{G}(\mathbf{X})$, and additional covariates can also be incorporated, such as nonlinear terms of \mathbf{Z} and $\mathbf{F}(\mathbf{X})$. Under sparsity conditions, these additional terms can be readily accommodated.

Remark 2. Under the classic SSL setups, various OR working models are used for estimating $\phi^*(\mathbf{X})$. For example, Chakrabortty (2016) recommended using a partial linear OR model. In addition, Zhang and Bradic (2021) proposed using flexible OR working models such as Lasso, elastic net, etc., provided that they meet certain rates of convergence.

3.2 Estimation procedures

The proposed method consists of the following three steps: (a) estimating the parameter γ in the PS model (7); (b) estimating the parameter α in the OR model (8); (c) estimating the target parameter β . Below, we give details for the three steps.

For estimating γ , rather than employing a regularized maximum likelihood estimator, we utilize a regularized calibrated estimator (Tan, 2020b), defined as

$$\hat{\gamma} = \underset{\gamma \in \mathbb{R}^{p+1}}{\operatorname{argmin}} L_{\text{RCAL}}(\gamma) = \underset{\gamma \in \mathbb{R}^{p+1}}{\operatorname{argmin}} \{\ell_{\text{CAL}}(\gamma) + \lambda_\gamma \|\gamma_{1:p}\|_1\}, \quad (10)$$

where

$$\ell_{\text{CAL}}(\gamma) = \tilde{\mathbb{E}}[R \exp\{-\gamma^T \mathbf{F}(\mathbf{X})\} + (1 - R)\gamma^T \mathbf{F}(\mathbf{X})],$$

λ_γ is a pre-specified tuning parameter, $\|\cdot\|_1$ denotes the L_1 -norm, and for any vector ν , $\nu_{i:j}$ is the sub-vector of ν consisting of its i -th to j -th elements (both ends included).

For a possibly misspecified model $\pi(\mathbf{X}; \gamma)$, under suitable regularity conditions, $\hat{\gamma}$ converges in probability to its target value $\bar{\gamma}$ defined by

$$\bar{\gamma} = \underset{\gamma \in \mathbb{R}^{p+1}}{\operatorname{argmin}} \mathbb{E}[R \exp\{-\gamma^T \mathbf{F}(\mathbf{X})\} + (1 - R)\gamma^T \mathbf{F}(\mathbf{X})]. \quad (11)$$

For estimating α , we adopt a regularized weighted maximum likelihood estimator (Tan, 2020a), which is defined as

$$\hat{\alpha} = \underset{\alpha \in \mathbb{R}^{q+1}}{\operatorname{argmin}} L_{\text{RWL}}(\alpha; \hat{\gamma}) = \underset{\alpha \in \mathbb{R}^{q+1}}{\operatorname{argmin}} \{\ell_{\text{WL}}(\alpha; \hat{\gamma}) + \lambda_\alpha \|\alpha_{1:q}\|_1\}, \quad (12)$$

where

$$\ell_{\text{WL}}(\alpha; \hat{\gamma}) = \tilde{\mathbb{E}}(Rw(\mathbf{X}; \hat{\gamma})[-Y\alpha^T \mathbf{G}(\mathbf{X}) + \Psi\{\alpha^T \mathbf{G}(\mathbf{X})\}]),$$

$w(X, \hat{\gamma}) = \{1 - \pi(\mathbf{X}, \hat{\gamma})\}/\pi(\mathbf{X}, \hat{\gamma})$, $\Psi(u) = \int_0^u \psi(t)dt$ is the antiderivative of ψ , λ_α is a pre-specified tuning parameter. Similarly to the target value $\bar{\gamma}$, we define the target value of α as follows

$$\bar{\alpha} = \underset{\alpha \in \mathbb{R}^{q+1}}{\operatorname{argmin}} \mathbb{E}(Rw(\mathbf{X}; \bar{\gamma})[-Y\alpha^T \mathbf{G}(\mathbf{X}) + \Psi\{\alpha^T \mathbf{G}(\mathbf{X})\}]). \quad (13)$$

Similarly as in Tan (2020a), the construction of the loss function $L_{\text{RWL}}(\alpha; \hat{\gamma})$ differs from the regularized maximum likelihood estimator in two aspects: (a) it utilizes a weight function $w(X, \hat{\gamma})$ for each labeled observation; (b) the weight function depends on the fitted PS $\pi(\mathbf{X}, \hat{\gamma})$. This design is crucial for the proposed method to achieve desirable properties with possible misspecification of the OR model. A subtle difference from Tan (2020a) is that the OR basis functions \mathbf{G} are here explicitly allowed to differ from the PS basis functions \mathbf{F} .

After obtaining the estimators of γ and α , the proposed calibrated AIPW estimator of β , denoted as $\hat{\beta}$, is the solution to the following estimating equation

$$\tilde{\mathbb{E}}\{\tau(\mathbf{O}, \hat{\alpha}, \beta, \hat{\gamma})\} = 0, \quad (14)$$

where $\mathbf{O} = (\mathbf{X}, \mathbf{Z}, Y, R)$ and

$$\begin{aligned} \tau(\mathbf{O}, \alpha, \beta, \gamma) &= \frac{R}{\pi(\mathbf{X}; \gamma)} \{Y - \psi(\beta^T \mathbf{Z})\} \mathbf{Z} + \left\{1 - \frac{R}{\pi(\mathbf{X}; \gamma)}\right\} \{\psi(\alpha^T \mathbf{G}) - \psi(\beta^T \mathbf{Z})\} \mathbf{Z} \\ &= \frac{R}{\pi(\mathbf{X}; \gamma)} \{Y - \psi(\alpha^T \mathbf{G})\} \mathbf{Z} + \{\psi(\alpha^T \mathbf{G}) - \psi(\beta^T \mathbf{Z})\} \mathbf{Z} \end{aligned} \quad (15)$$

Our estimating equations (14) share the same form as the AIPW estimating equations (6). However, there exists a crucial distinction between our method and previous related methods based on (6). In previous methods, the PS and OR models are specified and fitted independently of each other, typically both by least squares, maximum likelihood, or variations. In our method, the PS and OR models are specified and fitted in a sequentially dependent manner. This design allows our estimator $\hat{\beta}$ to achieve \sqrt{N} -consistency in the presence of misspecified OR models.

4 Theoretical analysis

In this section, we present the theoretical analysis of the proposed estimator $\hat{\beta}$. In Section 4.1, we examine the theoretical properties of estimators $\hat{\gamma}$ and $\hat{\alpha}$ in the PS and OR models. Then, we study the asymptotic properties of the proposed estimator $\hat{\beta}$ in Section 4.2. Finally, in Section 4.3, we extend our analysis to the classical SSL setting (stratified sampling with constant PS).

4.1 Properties of the estimators for nuisance parameters

We analyze the theoretical properties of $\hat{\gamma}$ and $\hat{\alpha}$, which provide the basis for investigating properties of $\hat{\beta}$. For ease of presentation, we denote $\mathbf{F}(\mathbf{X})$ and $\mathbf{G}(\mathbf{X})$ as \mathbf{F} and \mathbf{G} , respectively.

We first present the properties of $\hat{\gamma}$ based on Tan (2020a) [Theorem 1 and Theorem 3]. The following assumptions are taken from Tan (2020a).

Assumption 3 (Regularity conditions for $\hat{\gamma}$). *Suppose that the following conditions are satisfied:*

- (i) $\max_{j=0,\dots,p} |f_j(\mathbf{X})| \leq C_0$ a.s. for a constant $C_0 \geq 1$;
- (ii) $\bar{\gamma}^\top \mathbf{F} \geq B_0$ a.s. for a constant $B_0 \in \mathbb{R}$, that is, $\pi(\mathbf{X}; \bar{\gamma})$ is bounded from below by $\{1 + \exp(-B_0)\}^{-1}$;
- (iii) the compatibility condition holds for $\Sigma_{\bar{\gamma}}$ with the subset $S_{\bar{\gamma}} = \{0\} \cup \{j : \bar{\gamma}_j \neq 0, j = 1, \dots, p\}$ and some constants $\nu_0 > 0$ and $\xi_0 > 1$, where $\Sigma_{\bar{\gamma}} = \mathbb{E}\{Rw(\mathbf{X}; \bar{\gamma})\mathbf{F}\mathbf{F}^\top\}$ is the Hessian of $\mathbb{E}\{\ell_{\text{CAL}}(\gamma)\}$ at $\gamma = \bar{\gamma}$;
- (iv) $|S_{\bar{\gamma}}|\lambda_0 \leq \zeta_0$ for a sufficiently small constant $\zeta_0 > 0$, depending only on (A_0, C_0, ξ_0, ν_0) , where $|\cdot|$ denotes the cardinality of a set, $\lambda_0 = c_\gamma \sqrt{\ln\{(1+p)/\epsilon\}/N}$, c_γ is a constant only depending on (B_0, C_0) and $A_0 > (\xi_0 + 1)/(\xi_0 - 1)$ is a constant.¹

The conditions in Assumption 3 are plausible as discussed in Tan (2020a). Based on Assumption 3, we have the following Proposition 1.

Proposition 1. *Suppose that Assumption 3 is satisfied, and λ_γ in (10) is specified by $\lambda_\gamma = A_0\lambda_0$. Then, with probability at least $1 - 8\epsilon$,*

$$D_{\text{CAL}}^\dagger(\hat{\gamma}^\top \mathbf{F}, \bar{\gamma}^\top \mathbf{F}) + (A_0 - 1)\lambda_0 \|\hat{\gamma} - \bar{\gamma}\|_1 \leq M_0 |S_{\bar{\gamma}}| \lambda_0^2, \quad (16)$$

where $M_0 > 0$ is a constant depending only on $(A_0, C_0, B_0, \xi_0, \nu_0, \zeta_0)$, and $D_{\text{CAL}}^\dagger(\hat{\gamma}^\top \mathbf{F}, \bar{\gamma}^\top \mathbf{F})$ is the symmetrized Bregman Divergence w.r.t $\ell_{\text{CAL}}(\gamma)$, i.e.

$$D_{\text{CAL}}^\dagger(\hat{\gamma}^\top \mathbf{F}, \bar{\gamma}^\top \mathbf{F}) = -\tilde{\mathbb{E}}[R\{\exp(-\hat{\gamma}^\top \mathbf{F}) - \exp(-\bar{\gamma}^\top \mathbf{F})\}(\hat{\gamma}^\top \mathbf{F} - \bar{\gamma}^\top \mathbf{F})].$$

Note that $D_{\text{CAL}}^\dagger(\hat{\gamma}^\top \mathbf{F}, \bar{\gamma}^\top \mathbf{F}) \geq 0$, then equation (16) implies that

$$\|\hat{\gamma} - \bar{\gamma}\|_1 \leq \frac{M_0}{A_0 - 1} |S_{\bar{\gamma}}| \lambda_0,$$

which indicates that the L_1 -convergence rate of the proposed regularized calibrated estimator $\hat{\gamma}$ is $|S_{\bar{\gamma}}|\lambda_0$, where $|S_{\bar{\gamma}}|$ is the nonzero size of $\bar{\gamma}$ and $\lambda_0 = c_\gamma \sqrt{\ln\{(1+p)/\epsilon\}/N}$. For example, taking $\epsilon = 1/(1+p)$ gives $\lambda_0 = c_\gamma \sqrt{2 \ln(1+p)/N}$, which leads to $\|\hat{\gamma} - \bar{\gamma}\|_1 = O(|S_{\bar{\gamma}}| \sqrt{\ln(1+p)/N})$.

For studying the properties of $\hat{\alpha}$, we make the following Assumption 4.

Assumption 4 (Regularity conditions for $\hat{\alpha}$). *Let $\psi_1(u)$ denote the derivative of $\psi(u)$. Suppose that the following conditions are satisfied:*

¹ c_γ is defined in Section III.1 of Supplement.

- (i) $C_1 \leq \psi_1(\bar{\alpha}^\top \mathbf{G}) \leq C_2$ a.s. for positive constants (C_1, C_2) ;
- (ii) $\psi_1(u) \leq \psi_1(u') \exp(C_3|u - u'|)$ for any (u, u') and certain constant $C_3 \geq 0$;
- (iii) $\max_{j=0, \dots, q} |g_j(\mathbf{X})| \leq C_4$ a.s. for a constant $C_4 \geq 1$;
- (iv) $Y - \psi(\bar{\alpha}^\top \mathbf{G})$ is uniformly sub-Gaussian given \mathbf{X} : $D_0^2 \mathbb{E}(\exp[\{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 / D_0^2] - 1 | \mathbf{X}) \leq D_1^2$ for some positive constants (D_0, D_1) ;
- (v) the compatibility condition holds for $\Sigma_{\bar{\alpha}}$ with the subset $S_{\bar{\alpha}} = \{0\} \cup \{j : \bar{\alpha}_j \neq 0, j = 0, \dots, q\}$ and some constant $\nu_1 > 0$ and $\xi_1 > 1$, where $\Sigma_{\bar{\alpha}} = \mathbb{E}[Rw(\mathbf{X}; \bar{\gamma})\psi_1(\bar{\alpha}^\top \mathbf{G})\mathbf{G}\mathbf{G}^\top]$;
- (vi) $(1 + \xi_1)^2 \nu_1^{-2} |S_{\bar{\alpha}}| \lambda_1 \leq \zeta_1$ for a sufficiently small constant $\zeta_1 > 0$, where $\lambda_1 = \max[\lambda_0, c_\alpha \sqrt{\ln\{(1+q)/\epsilon\}/N}]$, c_α is a constant depending on $(B_0, C_2, C_4, D_0, D_1)$;²
- (vii) let $A_1 > (\xi_1 + 1)/(\xi_1 - 1)$ be a constant. There exist $0 \leq \eta_2, \eta_3 < 1$, such that $\tilde{c}_\alpha |S_{\bar{\alpha}}| \lambda_1 \leq \eta_2$ and $\tilde{c}_\gamma |S_{\bar{\gamma}}| \lambda_0 \leq \eta_3$, where \tilde{c}_α and \tilde{c}_γ are both constants.³

Assumptions 4(i)–(ii) are mild conditions on the smoothness of the inverse link function ψ . Commonly used functions like the identity and logit functions satisfy these requirements. Assumptions 4(iii)–(vii) are similar to those used in related analysis by Tan (2020a). A subtle difference is that the compatibility condition in Tan (2020a) [Assumption 2 (ii)] is assumed for $\Sigma_{\bar{\gamma}}$ with $\mathbf{F} = \mathbf{G}$, whereas our compatibility condition is assumed for $\Sigma_{\bar{\alpha}}$. In our setting, \mathbf{G} has a higher dimension than \mathbf{F} (except in the case where $\mathbf{Z} = 1$).

Proposition 2. *Suppose Assumptions 3 and 4 are satisfied. If $\ln\{(1+p)/\epsilon\}/N < 1$ and λ_α in (12) is specified as $A_1 \lambda_1$, then with probability at least $1 - 10\epsilon$,*

$$D_{\text{WL}}^\dagger(\hat{\alpha}^\top \mathbf{G}, \bar{\alpha}^\top \mathbf{G}, \bar{\gamma}) + \exp(\eta_{01})(A_1 - 1)\lambda_1 \|\hat{\alpha} - \bar{\alpha}\|_1 \leq M_{11} |S_{\bar{\gamma}}| \lambda_0^2 + M_{12} |S_{\bar{\alpha}}| \lambda_1^2, \quad (17)$$

where $\eta_{01} = (A_0 - 1)^{-1} M_0 C_0 \zeta_0$, M_{11} and M_{12} are also constants;⁴ $D_{\text{WL}}^\dagger(\hat{\alpha}^\top \mathbf{G}, \bar{\alpha}^\top \mathbf{G}, \bar{\gamma})$ is the symmetrized Bregman divergence with respect to $\ell_{\text{WL}}(\alpha; \bar{\gamma})$, which is given by

$$D_{\text{WL}}^\dagger(\hat{\alpha}^\top \mathbf{G}, \bar{\alpha}^\top \mathbf{G}, \bar{\gamma}) = \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma})\{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\bar{\alpha}^\top \mathbf{G})\}(\hat{\alpha}^\top \mathbf{G} - \bar{\alpha}^\top \mathbf{G})]. \quad (18)$$

² c_α is defined in Section III.2 of Supplement.

³ \tilde{c}_α depends on $(A_0, A_1, C_0, C_3, C_4, M_0, \nu_1, \xi_1, \zeta_0, \zeta_1)$ and \tilde{c}_γ depends on $(A_0, A_1, C_0, C_1, C_3, C_4, D_0, D_1, M_0, \xi_1, \zeta_0)$.

They are defined in Section III.2 of Supplement.

⁴Constant M_{11} depends on $(A_0, C_0, C_1, D_0, D_1, M_0, \eta_3, \zeta_0)$ and M_{12} depends on $(A_0, A_1, C_0, M_0, \nu_1, \zeta_0, \zeta_1, \xi_1, \eta_2)$.

They are defined in Section III.2 of Supplement.

Proposition 2 gives the convergence rate of $\hat{\alpha}$. Since $D_{\text{WL}}^\dagger(\hat{\alpha}^\top \mathbf{G}, \bar{\alpha}^\top \mathbf{G}, \bar{\gamma}) \geq 0$ and $\lambda_1 \geq \lambda_0$ (Assumption 4(vi)),

$$\|\hat{\alpha} - \bar{\alpha}\|_1 \leq c_m(|S_{\bar{\gamma}}| + |S_{\bar{\alpha}}|)\lambda_1,$$

for some constant c_m . Compared with an unweighted lasso estimator, the convergence rate of our estimator $\hat{\alpha}$ not only depends on the non-sparsity size of $\phi(\mathbf{X}; \bar{\alpha})$ itself but also on that of $\pi(\mathbf{X}; \bar{\gamma})$.

4.2 Large sample properties of the proposed estimator

In this subsection, we present our main result: the properties of the proposed estimator $\hat{\beta}$. We show that the proposed estimator is consistent if either the PS model or the OR model is correctly specified. In addition, the proposed confidence intervals are valid when the PS model is correctly specified, irrespective of whether the OR model is correctly specified or not.

Assumption 5. *Suppose that the following conditions are satisfied:*

- (i) *assume $\mathbf{Z} = \{Z_0, \dots, Z_{m-1}\} \in \mathbb{R}^m$, and $\max_{j=0, \dots, m-1} |Z_j| \leq C_5$ almost surely for a constant $C_5 \geq 1$, where m is fixed as N increases;*
- (ii) *assume that $\beta \in \Theta_\beta \subset \mathbb{R}^m$, and for all $\epsilon > 0$, $\inf_{\beta \in \Theta_\beta: \|\beta - \beta^*\|_1 \geq \epsilon} \mathbb{E}\{\|\tau(\bar{\alpha}, \beta, \bar{\gamma})\|_1\} > 0$;*
- (iii) $\mathbb{E}\{\sup_{\beta \in \Theta_\beta} \|\tau(\bar{\alpha}, \beta, \bar{\gamma})\|_1\} < \infty$;
- (iv) $\psi_1(\beta^{*\top} \mathbf{Z}) \leq C_6$ a.e., for some constant $C_6 > 0$;
- (v) $|S_{\bar{\gamma}}| \sqrt{\ln(1+p) \ln(1+q)} = o(N^{1/2})$ and $|S_{\bar{\alpha}}| \ln(1+q) = o(N^{1/2})$.

Assumptions 5(ii)–(iii) are standard conditions in the asymptotic theory of estimating equations. Assumption 5(iv) is a mild condition on the smoothness of the inverse link function at the value of $\beta^{*\top} \mathbf{Z}$. Assumption 5(v) is comparable to the sparsity requirement in Tan (2020a).

Theorem 1. *Suppose Assumptions 1–5 are satisfied, and the PS model (7) is correctly specified with $\pi(\cdot; \bar{\gamma}) = \pi^*(\cdot)$. If $\ln((1+p)/\epsilon)/N < 1$, then the following results hold.*

- (i) *The estimator $\hat{\beta}$ is consistent and asymptotically normal, and*

$$\sqrt{N}(\hat{\beta} - \beta^*) \xrightarrow{d} \text{N}(0, \boldsymbol{\Sigma}),$$

where \xrightarrow{d} denotes convergence in distribution, $\boldsymbol{\Sigma} = \boldsymbol{\Gamma}^{-1} \boldsymbol{\Lambda} \boldsymbol{\Gamma}^{-1}$ with $\boldsymbol{\Gamma} = \mathbb{E}\{\psi_1(\beta^{*\top} \mathbf{Z}) \mathbf{Z} \mathbf{Z}^\top\}$, and

$$\boldsymbol{\Lambda} = \mathbb{E}\{\tau(\mathbf{O}, \bar{\alpha}, \beta^*, \bar{\gamma}) \tau(\mathbf{O}, \bar{\alpha}, \beta^*, \bar{\gamma})^\top\}$$

$$\begin{aligned}
&= \mathbb{E} \left(\left[\frac{1}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2 + \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*\top} \mathbf{Z})\}^2 \right] \mathbf{Z} \mathbf{Z}^T \right) \\
&\quad + 2\mathbb{E}[\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*\top} \mathbf{Z})\} \mathbf{Z} \mathbf{Z}^T].
\end{aligned} \tag{19}$$

(ii) A consistent estimator of Σ is $\hat{\Sigma} = \hat{\Gamma}^{-1} \hat{\Lambda} \hat{\Gamma}^{-1}$, where $\hat{\Gamma} = \tilde{\mathbb{E}}\{\psi_1(\hat{\beta}^T \mathbf{Z}) \mathbf{Z} \mathbf{Z}^T\}$ and

$$\hat{\Lambda} = \tilde{\mathbb{E}} \left\{ \tau(\mathbf{O}, \hat{\alpha}, \hat{\beta}, \hat{\gamma}) \tau(\mathbf{O}, \hat{\alpha}, \hat{\beta}, \hat{\gamma})^T \right\}.$$

Thus, for a constant vector \mathbf{c} with the same dimension of β , an asymptotic $(1 - \eta)$ confidence interval for $\mathbf{c}^T \beta^*$ is $\mathbf{c}^T \hat{\beta} \pm z_{\eta/2} \sqrt{\mathbf{c}^T \hat{\Sigma} \mathbf{c} / N}$, where $z_{\eta/2}$ is the $(1 - \eta/2)$ quantile of the standard normal distribution.

Theorem 1 shows that if the PS model is correct, regardless of whether the OR working model is correct or not, the proposed estimator $\hat{\beta}$ is consistent and asymptotically normal, and the proposed confidence intervals based on $\hat{\Sigma}$ are valid. In contrast, the estimator $\hat{\beta}_{\text{IPW}}$ defined by equations (5) is not \sqrt{N} -consistent in general, even with a correctly specified PS model. This is because in high-dimensional settings, the convergence rate of $\hat{\pi}(\mathbf{X})$ is typically slower than $1/\sqrt{N}$, leading to a slower convergence rate of $\hat{\beta}_{\text{IPW}}$. Similarly, when the OR working model is misspecified and the estimator $\hat{\phi}(\mathbf{X})$ is inconsistent, the convergence rate of the AIPW estimator with nuisance parameters being estimated using conventional regularized maximum likelihood may still be slower than $1/\sqrt{N}$.

In Theorem 1, we present our results in high-dimensional settings. The conclusions in Theorem 1 also hold in low-dimensional settings with a reduced form of Assumptions 3, 4 and 5.

4.3 Extension to stratified sampling with constant PS

To facilitate comparison with existing methods described in Section 5, we extend the theoretical analysis of the proposed estimator to the classical SSL setting (stratified sampling with constant PS), where the sizes of labeled and unlabeled datasets, n and $N - n$, are deterministic. For fixed n and N , the observed data are generated as follows:

- The labeled dataset $(\mathbf{X}_1, Y_1), \dots, (\mathbf{X}_n, Y_n) \stackrel{\text{i.i.d.}}{\sim} \mathbb{P}(\mathbf{X}, Y | R = 1)$.
- The unlabeled dataset $(\mathbf{X}_{n+1}, \dots, \mathbf{X}_N) \stackrel{\text{i.i.d.}}{\sim} \mathbb{P}(\mathbf{X}, Y | R = 0)$.

Moreover, by letting $\mathbf{F} = 1$, $\hat{\pi}(\mathbf{X}) = \pi(\mathbf{X}; \hat{\gamma}) = n/N$ (constant PS) and allowing a general choice of \mathbf{G} instead of (9), our AIPW estimator, denoted as $\hat{\beta}^s$, can be rewritten as a solution to the

following estimating equation:

$$\frac{1}{n} \sum_{i=1}^n \{Y_i - \psi(\hat{\alpha}^T \mathbf{G}_i)\} \mathbf{Z}_i + \frac{1}{N} \sum_{i=1}^N \{\psi(\hat{\alpha}^T \mathbf{G}_i) - \psi(\beta^T \mathbf{Z}_i)\} \mathbf{Z}_i = 0, \quad (20)$$

where \mathbf{Z}_i denotes the i -th observation of \mathbf{Z} and \mathbf{G}_i is the abbreviation of $\mathbf{G}(\mathbf{X}_i)$. Due to the constant $\hat{\pi}(\mathbf{X})$, our estimator $\hat{\alpha}$ reduces to the regularized unweighted maximum likelihood or least squares estimator. The following proposition for $\hat{\beta}^s$ can be readily derived.

Proposition 3. *Suppose that the conditions of Theorem 1 are satisfied with $\mathbf{F} = 1$, $\pi^*(\mathbf{X}) \equiv n/N$, and a general choice of \mathbf{G} , where Assumption 5(v) reduces to $|S_{\bar{\alpha}}| \ln(q+1) = o_p(\sqrt{n})$. Then we have*

$$\sqrt{n}(\hat{\beta}^s - \beta^*) \xrightarrow{d} \mathbf{N}(0, \boldsymbol{\Sigma}^s),$$

where $\boldsymbol{\Sigma}^s = \boldsymbol{\Gamma}^{-1} \boldsymbol{\Lambda}^s \boldsymbol{\Gamma}^{-1}$, with $\boldsymbol{\Gamma}$ being defined as in Theorem 1 and

$$\begin{aligned} \boldsymbol{\Lambda}^s = & \mathbb{E} \left[\left\{ Y - \frac{N-n}{N} \psi(\bar{\alpha}^T \mathbf{G}) - \frac{n}{N} \psi(\beta^{*T} \mathbf{Z}) \right\}^2 \mathbf{Z} \mathbf{Z}^T \right] \\ & + \frac{(N-n)n}{N^2} \mathbb{E} [\{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\}^2 \mathbf{Z} \mathbf{Z}^T]. \end{aligned}$$

For comparison, by letting $\mathbf{F} = 1$ and $\pi^*(\mathbf{X}) \equiv n/N$ in Theorem 1, the variance matrix $\boldsymbol{\Sigma}$ for $\hat{\beta}$ such that $\sqrt{N}(\hat{\beta} - \beta^*) \xrightarrow{d} \mathbf{N}(0, \boldsymbol{\Sigma})$ is $\boldsymbol{\Sigma} = \boldsymbol{\Gamma}^{-1} \boldsymbol{\Lambda} \boldsymbol{\Gamma}^{-1}$, where $\boldsymbol{\Lambda}$ is simplified as

$$\begin{aligned} \boldsymbol{\Lambda} = & \mathbb{E} \left(\left[\frac{N}{n} \{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2 + \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\}^2 \right] \mathbf{Z} \mathbf{Z}^T \right) \\ & + 2 \left(\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\} \right) \mathbf{Z} \mathbf{Z}^T \end{aligned}$$

By direct calculation (see Section V of Supplement), it is shown that $\boldsymbol{\Sigma}/N = \boldsymbol{\Sigma}^s/n$, which means the unscaled asymptotic variances of $\hat{\beta}^s$ and $\hat{\beta}$ are the same. Hence, in the classical SSL with constant PS, the asymptotic variances of our estimators, $\hat{\beta}$ under random sampling or $\hat{\beta}^s$ under stratified sampling, are equivalent to each other.

5 Comparison with previous methods

In this subsection, we first briefly summarize characteristics of previous related methods. All of them can be integrated into the AIPW estimation framework. We also compare asymptotic variances of our methods with previous methods.

5.1 Unified framework

Various methods have been proposed in the classical SSL setting, i.e., stratified sampling with constant PS ($F = 1$) as in Section 4.3 (Chakrabortty, 2016; Chakrabortty and Cai, 2018; Zhang et al., 2019; Zhang and Bradic, 2021). From an AIPW point of view, the major difference among previous methods lies in the choices of OR working models. For example, Zhang et al. (2019) and Zhang and Bradic (2021) proposed linear OR working models for the estimation of $\mathbb{E}(Y)$, i.e., with $\mathbf{Z} = 1$. Chakrabortty (2016) and Chakrabortty and Cai (2018) proposed using non-parametric or semi-parametric OR working models, such as kernel smoothing or partially linear model, for regression analysis with \mathbf{Z} a sub-vector of \mathbf{X} .

If we disregard the specific choice of the OR working model, the previous methods can be incorporated into the AIPW estimating framework. In our notation, the estimators of Chakrabortty (2016), Zhang et al. (2019) and Zhang and Bradic (2021) can be reformulated as solutions of the following equations:

$$\frac{1}{n} \sum_{i=1}^n \{Y_i - \psi(\hat{\alpha}^T \mathbf{G}_i)\} \mathbf{Z}_i + \frac{1}{N} \sum_{i=1}^N \{\psi(\hat{\alpha}^T \mathbf{G}_i) - \psi(\beta^T \mathbf{Z}_i)\} \mathbf{Z}_i = 0, \quad (21)$$

for different choices of \mathbf{Z}_i and $\psi(\cdot)$. Specifically, Zhang et al. (2019) and Zhang and Bradic (2021) correspond to the case of $\mathbf{Z} = 1$ and $\psi(\cdot)$ is the identity function in equation (21), while Chakrabortty (2016) corresponds to the case where \mathbf{Z} is any sub-vector of \mathbf{X} and $\psi(\cdot)$ is an arbitrary inverse link function. Suppose a constant PS model (7) is used with $\pi^*(\mathbf{X}) \equiv n/N$ and a general choice of \mathbf{G} is used in the OR model (8). Then the AIPW estimating equation (14) or, in the simplified form, (20) in Section 4.3, coincides with (21).

In addition, Chakrabortty and Cai (2018) adopted a variation of AIPW estimating equations. By the assumption $\lim_{n, N \rightarrow \infty} n/N \rightarrow 0$ and controlling kernel smoothing in fitting OR working models, they made it possible to drop the labeled part and only retain the augmented term of unlabeled data in (21). Their estimating equations can be reformulated in our notation as

$$\frac{1}{N-n} \sum_{i=n+1}^N \{\psi(\hat{\alpha}^T \mathbf{G}_i) - \psi(\beta^T \mathbf{Z}_i)\} \mathbf{Z}_i = 0,$$

with $\psi(\cdot)$ to be identity function and $\mathbf{Z} = \mathbf{X}$, corresponding to full linear regression.

5.2 Variance comparison

Under stratified sampling with constant PS, both estimators of Zhang et al. (2019) and Zhang and Bradic (2021) of $\mathbb{E}(Y)$ achieve asymptotic normality and their asymptotic variance is $\text{Var}(Y - \bar{\alpha}^T \mathbf{G})$

+ $(n/N)\text{Var}(\bar{\alpha}^T \mathbf{G})$. Under this setting, by Proposition 3 with $\mathbf{Z} = 1$, our AIPW estimator has the asymptotic variance

$$\mathbb{E} \left\{ (Y - \bar{\alpha}^T \mathbf{G})^2 + \frac{n}{N} (\bar{\alpha}^T \mathbf{G} - \beta^*)^2 + 2 \frac{n}{N} (Y - \bar{\alpha}^T \mathbf{G})(\bar{\alpha}^T \mathbf{G} - \beta^*) \right\}, \quad (22)$$

where $\beta^* = \mathbb{E}(Y) = \mathbb{E}(\bar{\alpha}^T \mathbf{G})$ and $\mathbb{E}\{(Y - \bar{\alpha}^T \mathbf{G})\mathbf{G}\} = 0$ by definition of $\bar{\alpha}$ and the fact that \mathbf{G} includes 1. Then

$$\mathbb{E} \{(Y - \bar{\alpha}^T \mathbf{G})(\bar{\alpha}^T \mathbf{G} - \beta^*)\} = \mathbb{E}\{(Y - \bar{\alpha}^T \mathbf{G})(\bar{\alpha}^T \mathbf{G})\} = 0,$$

and (22) reduces to $\text{Var}(Y - \bar{\alpha}^T \mathbf{G}) + (n/N)\text{Var}(\bar{\alpha}^T \mathbf{G})$ matching results in Zhang et al. (2019) and Zhang and Bradic (2021).

Under stratified sampling with constant PS, the estimators of regression coefficients in conditional mean models proposed by Chakraborty (2016) and Chakraborty and Cai (2018) achieve asymptotic normality under the assumption that $\lim_{n, N \rightarrow \infty} n/N \rightarrow 0$. Their asymptotic variances satisfy the following formula:

$$\mathbf{\Gamma}^{-1} \text{Var}[\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \mathbf{Z}] \mathbf{\Gamma}^{-1}. \quad (23)$$

Under this setting, by Proposition 3, our AIPW estimator has the asymptotic variance

$$\begin{aligned} & \mathbf{\Gamma}^{-1} \text{Var}[\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \mathbf{Z}] \mathbf{\Gamma}^{-1} + \frac{n}{N} \mathbf{\Gamma}^{-1} \mathbb{E}[\{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\}^2 \mathbf{Z} \mathbf{Z}^T] \mathbf{\Gamma}^{-1} \\ & + 2 \frac{n}{N} \mathbf{\Gamma}^{-1} \mathbb{E}[\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\} \mathbf{Z} \mathbf{Z}^T] \mathbf{\Gamma}^{-1}, \end{aligned}$$

which, compared with (23), in general has additional term

$$\frac{n}{N} \mathbf{\Gamma}^{-1} \left\{ \mathbb{E}[\{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\}^2 + 2\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\}] \mathbf{Z} \mathbf{Z}^T \right\} \mathbf{\Gamma}^{-1}.$$

The additional term reduces to 0 under the condition $\lim_{n, N \rightarrow \infty} n/N \rightarrow 0$, implying that our result aligns with those of Chakraborty (2016) and Chakraborty and Cai (2018) with the same condition.

6 Simulation study

In this section, we design experiments to evaluate the finite-sample performance of the proposed method and compare it with the competing IPW and AIPW methods. We consider the estimators of population mean for $\mathbf{Z} = 1$, regression coefficients in the mean model for $\mathbf{Z} = X_1$ and $\mathbf{Z} = \mathbf{X}$, respectively, and $\psi(\cdot)$ is assumed to be the identity function.

6.1 Data generating process

Throughout the simulation, we generate the covariates \mathbf{X} as follows. We first generate a random vector from $N(0, \Sigma)$, where the variance matrix $\Sigma \in \mathbb{R}^{3 \times 3}$ has elements $\Sigma_{i,j}$ defined as $2^{-|i-j|}$ for $i, j = 1, 2, 3$. Then we clamp each of its coordinates within $[-3, 3]$ to obtain (X_1, X_2, X_3) and $\mathbf{X} = (1, X_1, X_2, X_3)$. In addition, the data source indicator R follows a Bernoulli distribution with success probability $\pi(\mathbf{X})$, where $\pi(\mathbf{X}) = \{1 + \exp(-\gamma^T \mathbf{F})\}^{-1}$, the parameter $\gamma = (-1.5, -0.8, -0.2, 0.3, 0, \dots, 0)^T$ and the basis functions \mathbf{F} are described in Section 6.2.

Study I. We first focus on the estimation of the population mean and consider two data-generating mechanisms:

- **Case 1.** The outcome $Y = -0.2 + 0.1\tilde{X}_1 + 0.4\tilde{X}_2 + 0.7\tilde{X}_3 + \epsilon$, where $\epsilon \sim \mathcal{N}(0, 0.1)$ and $\tilde{X}_j = X_j \cdot |X_j|^{0.1} + X_j \cdot |X_j|^{0.3} + X_j \cdot |X_j|^{0.5}$, for $j = 1, 2, 3$. We set $\mathbf{Z} = 1$.
- **Case 2.** The outcome $Y = -0.2 + 0.1\tilde{X}_1 + 0.4\tilde{X}_2 + 0.7\tilde{X}_3 + \epsilon$, where $\epsilon \sim \mathcal{N}(0, 0.1)$ and $\tilde{X}_j = |X_j| \exp(|X_j|^{0.1} + |X_j|^{0.3})$ for $j = 1, 2, 3$. We set $\mathbf{Z} = 1$.

In many scenarios, estimating the conditional mean given a subset of variables in \mathbf{X} garners statistical interest. Accordingly, we design experiments in Study II to evaluate the performance of the proposed estimator in such setups.

Study II. We further consider three additional cases for estimating regression coefficients in the conditional mean outcome model.

- **Case 3.** The outcome $Y = 0.4\tilde{X}_1 + 0.2\tilde{X}_2 + \epsilon$, where $\epsilon \sim \mathcal{N}(0, 0.1)$, $\tilde{X}_1 = X_1 + X_1^2$ and $\tilde{X}_2 = \cos(\pi/9 \cdot X_1 \cdot X_3)$. We set $\mathbf{Z} = X_1$.
- **Case 4.** The outcome $Y = 0.4\tilde{X}_1 + 0.2\tilde{X}_2 + \epsilon$, where $\epsilon \sim \mathcal{N}(0, 0.1)$, $\tilde{X}_1 = X_1 \cdot \mathbb{I}\{X_1 > 0\} \sqrt{|X_1|}$ and $\tilde{X}_2 = X_1 \cdot X_2$. We set $\mathbf{Z} = X_1$.
- **Case 5.** The outcome $Y = -0.2 + 0.1\tilde{X}_1 + 0.4\tilde{X}_2 + 0.7\tilde{X}_3 + \epsilon$, where $\epsilon \sim \mathcal{N}(0, 0.1)$, $\tilde{X}_1 = X_1 \cdot X_2$, $\tilde{X}_2 = X_2 \cdot X_3$ and $\tilde{X}_3 = X_1 \cdot X_3$. We set $\mathbf{Z} = \mathbf{X}$.

Cases 3 and 4 involve \mathbf{Z} as a specific covariate X_1 , while Case 5 involves \mathbf{Z} as the full set of covariates \mathbf{X} . In addition, for all Cases 1–5, OR models are misspecified.

6.2 Implementation details and competing methods

We compare the proposed method (AIPW_{RCAL}) with several alternative methods: the IPW method with Lasso-regularized maximum likelihood estimation for the PS model, AIPW methods with

Lasso-regularized maximum likelihood estimation for both the PS and OR models without cross-fitting as in Tan (2020a),⁵ and AIPW methods with cross-fitting and Lasso-regularized maximum likelihood estimation for both the PS and OR models (Chernozhukov et al., 2018; Zhang and Bradic, 2021). These competing estimators are denoted as IPW, AIPW_{RML}, and AIPW_{CF}, respectively. For the PS and OR models, the basis functions \mathbf{F} and \mathbf{G} are specified as follows.

- AIPW_{RCAL}: Given $\mathbf{X} = (1, X_1, \dots, X_d)^\top$ in Section 6.1, let $\{\xi_i\}_{i=1}^k$ be the k points equally spaced within $(-a, a)$, where $d = 3$, $k = 49$, and $a = 3$. Let $f_{ij}(\mathbf{X}) = (X_i - \xi_j)_+$, $i = 1, \dots, d, j = 1, \dots, k$. Let $\mathbf{F} = \{1, f_{11}(\mathbf{X}), \dots, f_{1n_k}(\mathbf{X}), \dots, f_{d1}(\mathbf{X}), \dots, f_{dn_k}(\mathbf{X})\}^\top$ be the basis functions in the PS model, and $\mathbf{G} = \{\mathbf{F}^\top, (\mathbf{Z} \otimes \mathbf{F})^\top\}^\top$ be the basis functions in the OR model. Then the dimension of \mathbf{F} is 148. For $\mathbf{Z} = 1, X_1, \mathbf{X}$, the dimensions of \mathbf{G} are 148, 285 and 589, respectively.
- IPW: Let \mathbf{F} be the basis functions for the PS model.
- AIPW_{RML}: Let \mathbf{F} and $\mathbf{G} = \mathbf{F}$ be the basis functions for both PS and OR models, respectively.
- AIPW_{CF}: Let \mathbf{F} and $\mathbf{G} = \mathbf{F}$ be the basis functions for both PS and OR models, respectively.

Both the Lasso-regularized calibrated and maximum likelihood estimators for the PS and OR models can be implemented using the R package RCAL (Tan and Sun, 2020). We employ 5-fold cross-fitting to select the optimal tuning parameters. In addition, by equation (2), $\beta^* = \mathbb{E}(\mathbf{Z}\mathbf{Z}^\top)^{-1}\mathbb{E}(Y\mathbf{Z})$. Thus the true value of β^* is calculated as $\tilde{\mathbb{E}}(\mathbf{Z}\mathbf{Z}^\top)^{-1}\tilde{\mathbb{E}}(Y\mathbf{Z})$ through a simulation with a sample size of 100,000. For $\mathbf{Z} = X_1$ and $\mathbf{Z} = \mathbf{X}$, we denote $\beta^* = \beta_1$ and $\beta^* = (\beta_0, \beta_1, \beta_2, \beta_3)^\top$, respectively.

6.3 Summary of results

We present and analyze the simulation results for Study I and Study II. In the following tables, we compare various methods in terms of five metrics: Bias (Monte Carlo bias), $\sqrt{\text{Var}}$ (Monte Carlo standard deviation), $\sqrt{\text{EVar}}$ (square root of the mean of variance estimates), CP90 (coverage proportions of the 90% CIs), and CP95 (coverage proportions of the 95% CIs). As discussed in the paragraph below Theorem 1, under high-dimensional settings, the IPW estimator is not \sqrt{N} -consistent, and its asymptotic normality is not well established. Therefore, we do not report its numerical results for $\sqrt{\text{EVar}}$, CP90, and CP95.

⁵For Lasso-regularized maximum likelihood estimation, the loss functions in fitting PS models and OR models are $-\tilde{\mathbb{E}}[Y\gamma^\top\mathbf{F} - \ln\{1 + \exp(\gamma^\top\mathbf{F})\}] + \lambda_\gamma\|\gamma_{1:p}\|_1$ and $\tilde{\mathbb{E}}[(Y - \alpha^\top\mathbf{G})^2] + \lambda_\alpha\|\alpha_{1:q}\|_1$, respectively.

Results for Study I. Table 1 shows the numerical results for the estimation of population mean $\mathbb{E}(Y)$. From the table, the proposed method $\text{AIPW}_{\text{RCAL}}$ has the smallest $\sqrt{\text{Var}}$ and $\sqrt{\text{EVar}}$, and Bias. Moreover, CP90 and CP95 of the proposed method are more aligned with their nominal values of 0.90 and 0.95, respectively. This indicates the effectiveness of the proposed method in terms of estimating the population mean.

Table 1: Summary of estimates of population mean for Study I.

	Case 1			Case 2		
	$\text{AIPW}_{\text{RCAL}}$	AIPW_{RML}	AIPW_{CF}	$\text{AIPW}_{\text{RCAL}}$	AIPW_{RML}	AIPW_{CF}
Bias	0.004	0.004	0.006	0.002	-0.006	0.031
$\sqrt{\text{Var}}$	0.078	0.078	0.079	0.139	0.143	0.166
$\sqrt{\text{EVar}}$	0.079	0.079	0.082	0.140	0.143	0.161
CP90	0.904	0.908	0.918	0.898	0.886	0.896
CP95	0.948	0.954	0.956	0.948	0.946	0.956

Figure 1 depicts the box plots of the estimates for Case 1 and Case 2, where the blue horizontal line indicates the true value. In both cases, our method $\text{AIPW}_{\text{RCAL}}$ exhibits the smallest biases, interquartile ranges, and whiskers, indicating the smallest variances compared to the other methods. In addition, AIPW_{CF} shows more outliers than the other two methods, which is more apparent in the results of Study II, suggesting that cross-fitting may cause instability for the estimates.




Figure 1: Box plots for estimates of population mean.

Results for Study II. The simulation results for Cases 3–5 are presented in Tables 2–3, and the corresponding box plots are displayed in Figures 2–3. We observe similar patterns as those in


Cases 1 and 2: the proposed method performs well in terms of all metrics.

Table 2: Summary of estimates of β_1 in Cases 3 and 4.


	Case 3			Case 4		
	AIPW _{RCAL}	AIPW _{RML}	AIPW _{CF}	AIPW _{RCAL}	AIPW _{RML}	AIPW _{CF}
Bias	0.000	-0.011	-0.004	0.002	-0.010	-0.002
$\sqrt{\text{Var}}$	0.036	0.043	0.058	0.023	0.031	0.042
$\sqrt{\text{EVar}}$	0.036	0.037	0.056	0.021	0.025	0.042
CP90	0.884	0.824	0.840	0.866	0.774	0.838
CP95	0.946	0.886	0.900	0.930	0.834	0.906

Table 3: Summary of estimates of $\beta_0, \beta_1, \beta_2, \beta_3$ in Case 5

	β_0			β_1		
	AIPW _{RCAL}	AIPW _{RML}	AIPW _{CF}	AIPW _{RCAL}	AIPW _{RML}	AIPW _{CF}
Bias	0.001	0.001	0.013	-0.007	-0.004	0.000
$\sqrt{\text{Var}}$	0.024	0.068	0.084	0.036	0.114	0.139
$\sqrt{\text{EVar}}$	0.025	0.055	0.077	0.036	0.089	0.131
CP90	0.910	0.800	0.846	0.886	0.774	0.830
CP95	0.958	0.874	0.914	0.942	0.842	0.900
	β_2			β_3		
	AIPW _{RCAL}	AIPW _{RML}	AIPW _{CF}	AIPW _{RCAL}	AIPW _{RML}	AIPW _{CF}
Bias	-0.001	0.002	0.001	-0.006	0.114	-0.060
$\sqrt{\text{Var}}$	0.033	0.086	0.111	0.040	0.095	0.132
$\sqrt{\text{EVar}}$	0.034	0.075	0.108	0.041	0.095	0.133
CP90	0.920	0.818	0.866	0.898	0.592	0.774
CP95	0.958	0.902	0.938	0.940	0.682	0.842




(a) Case 3




(b) Case 4


Figure 2: Box plots of estimates of β_1 in Cases 3 and 4.




(a) $\hat{\beta}_0$



(b) $\hat{\beta}_1$



(c) $\hat{\beta}_2$



(d) $\hat{\beta}_3$

Figure 3: Box plots of estimates of $\beta_0, \beta_1, \beta_2, \beta_3$ in Case 5

7 Application

7.1 Data description

The Communities and Crime dataset comprises 1994 records of crime-related information from communities in the United States, which combine socio-economic data from the 1990 US Census, law enforcement data from the 1990 US LEMAS survey, and crime data from the 1995 FBI UCR. Each record includes a response variable `ViolentCrimesPerPop`⁶ and 127 covariates, encompassing both location information (such as `state` and `county`) and socio-economic factors (such as `PctTeen2Par`⁷, `HousVacant`⁸, etc.). In this study, we are interested in examining the influence of univariate covariates on the response. We consider the case where $\mathbf{Z} = (1, X_i)^T$ for a particular univariate covariate X_i as discussed in Section 2.1, and we denote $\beta^* = (\beta_0, \beta_1)$.

Due to the presence of numerous missing values in high-dimensional covariates, we eliminate covariates with high missing ratios. See details of the pre-processing procedure in Section VI.1 of Supplement. After pre-processing, the analytical dataset consists of 1993 observations and 26 covariates (i.e., $d = 26$). The shift in covariates \mathbf{X} is naturally introduced by the different states where the communities are located.⁹ We set label indicators R for communities in New Jersey (Code 34) to be 1 and those for communities in other states to be 0 and remove the associated response data if $R = 0$, resulting in 211 labeled observations and 1782 unlabeled observations. The covariate shift of the joint distribution of \mathbf{X} was confirmed to exist using a Gaussian kernel two-sample test with maximum mean discrepancy (You, 2023). Additionally, we assess the shift of each individual covariate by a bootstrap version of the Kolmogorov–Smirnov test (Sekhon, 2011). For results of those tests, please see Section VI.2 of Supplement.

We randomly take 90% of labeled data and 90% of unlabeled data to form the training set with the remaining data used for the testing set. From the remaining 26 covariates, we select four representative ones: `PctTeen2Par`, `HousVacant`, `PctHousNoPhone`¹⁰ and `PopDens`¹¹, which illustrate different aspects of the socio-economic characteristics of communities. Notice that the covariate shift exists in all four covariates.

We compare the proposed method with IPW, AIPW_{RML} and AIPW_{CF} methods with piecewise

⁶total number of violent crimes per 100,000 population

⁷percent of kids age 12-17 in two parent households

⁸number of vacant households

⁹Notice that no covariate related to locations of communities is included in \mathbf{X} .

¹⁰percent of occupied housing units without phone

¹¹population density in persons per square mile

linear basis functions introduced in Section 6.2. The PS and OR working models are estimated the same way as described there. For details of the procedures for designing basis functions, please see Section VI.3 of Supplement.

7.2 Results

Table 4 presents the estimates of the regression coefficients $\hat{\beta}_1$ along with the prediction mean squared error (MSE), which are calculated using the test data. It reveals that the point estimates of the regression coefficient are similar across the different methods. Notably, our estimators achieve the lowest prediction MSE except `PctTeen2Par`, highlighting the superior performance of our methods in minimizing predictive errors.


Table 4: Summary of $\hat{\beta}_1$ and prediction MSE

	$\hat{\beta}_1$				prediction MSE			
	AIPW _{RCAL}	IPW	AIPW _{RML}	AIPW _{CF}	AIPW _{RCAL}	IPW	AIPW _{RML}	AIPW _{CF}
<code>PctTeen2Par</code>	-0.137	-0.172	-0.147	-0.256	0.034	0.032	0.034	0.044
<code>HousVacant</code>	0.107	0.261	0.073	0.133	0.046	0.085	0.046	0.047
<code>PctHousNoPhone</code>	0.123	0.245	0.103	0.050	0.036	0.058	0.039	0.047
<code>PopDens</code>	0.045	0.069	0.048	0.039	0.050	0.052	0.052	0.055


Moreover, signs of estimates of coefficients are the same among different methods for each covariate Z of interest. and they coincide with common sense and previous studies. For example, the coefficients of `PctTeen2Par` are negative, since it is believed to have protective effects in assaults (Luo and Qi, 2017); the coefficients of `HousVacant` is positive, and criminological theories predict a positive association between vacancy and crime since empty structures of houses could provide locations for some crimes (e.g., prostitution, drug dealing), and the absence of residents may prevent social organization and reduce guardianship (Roth, 2019). Moreover, AIPW_{RML} and ours are close in all cases, while IPW estimators and AIPW_{CF} estimators are far from others in some cases.

In Figure 4, we compare the 95% CIs of AIPW_{RCAL}, AIPW_{RML} and AIPW_{CF}. From the CIs, we see that for AIPW_{RML} and our estimators all four single effects are significant. CIs of our estimators and of AIPW_{RML}'s have similar lengths and are overlapped, except `HousVacant`. The reason of the small difference is that the estimates of β_0 are a bit different. CIs of AIPW_{CF} are much longer; for `PctHousNoPhone` and `PopDens`, the estimates are not significant. Both phenomena show that


AIPW_{CF} is not as efficient as other two methods.




(a) PctTeen2Par



(b) HousVacant



(c) PctHousNoPhone



(d) PopDens

Figure 4: Comparison of 95% CIs of AIPW_{RCAL}, AIPW_{RML} and AIPW_{CF}

8 Extension to estimation of β^{0*}

Consider the estimation of β^{0*} , defined as a solution to estimating equation (4). Under Assumption 1, $\mathbb{E}[R\{1 - \pi^*(\mathbf{X})\}/\pi^*(\mathbf{X})\{Y - \psi(\beta^T \mathbf{Z})\} \mathbf{Z}] = \mathbb{E}[\{1 - R\}\{Y - \psi(\beta^T \mathbf{Z})\} \mathbf{Z}]$. Then a natural sample estimating equation for β^{0*} is $\tilde{\mathbb{E}}[R\{1 - \hat{\pi}(\mathbf{X})\}/\hat{\pi}(\mathbf{X})\{Y - \psi(\beta^T \mathbf{Z})\} \mathbf{Z}] = 0$. We augment the estimating equations similarly as described in Section 2.3 and obtain the sample AIPW estimating equations:

$$\tilde{\mathbb{E}} \left[\frac{R\{1 - \hat{\pi}(\mathbf{X})\}}{\hat{\pi}(\mathbf{X})} \{Y - \psi(\beta^T \mathbf{Z})\} \mathbf{Z} + \left\{ 1 - \frac{R}{\hat{\pi}(\mathbf{X})} \right\} \left\{ \hat{\phi}(\mathbf{X}) - \psi(\beta^T \mathbf{Z}) \right\} \mathbf{Z} \right] = 0. \quad (24)$$

For the PS and OR models, we adopt a similar construction as in Section 3. Our AIPW estimator for β^{0*} , $\hat{\beta}^0$, is defined as the solution to the following estimating equations:

$$\tilde{\mathbb{E}}\{\tau^0(\mathbf{O}, \hat{\alpha}, \beta, \hat{\gamma})\} = 0, \quad (25)$$

where

$$\tau^0(\mathbf{O}, \alpha, \beta, \gamma) = \left[\frac{R\{1 - \pi(\mathbf{X}; \gamma)\}}{\pi(\mathbf{X}; \gamma)} \{Y - \psi(\beta^T \mathbf{Z})\} + \left\{ 1 - \frac{R}{\pi(\mathbf{X}; \gamma)} \right\} \{\psi(\alpha^T \mathbf{G}) - \psi(\beta^T \mathbf{Z})\} \right] \mathbf{Z}.$$

The following result is an extension of Theorem 1 to the estimation of β^{0*} .

Theorem 2. *Under Assumptions 1–5, if the PS model (7) is correctly specified with $\pi(\cdot; \bar{\gamma}) = \pi^*(\cdot)$, and $\ln\{(1+p)/\epsilon\}/N < 1$, then the following results hold.*

(i) *The estimator $\hat{\beta}^0$ is consistent and asymptotically normal, and*

$$\sqrt{N}(\hat{\beta}^0 - \beta^{0*}) \xrightarrow{d} \mathbf{N}(0, \boldsymbol{\Sigma}^0),$$

where $\boldsymbol{\Sigma}^0 = \boldsymbol{\Gamma}^{0-1} \boldsymbol{\Lambda}^0 \boldsymbol{\Gamma}^{0-1}$ with $\boldsymbol{\Gamma}^0 = \mathbb{E}\{[1 - \pi(\mathbf{X}; \bar{\gamma})] \psi_1(\beta^{0*\top} \mathbf{Z}) \mathbf{Z} \mathbf{Z}^\top\}$ and

$$\boldsymbol{\Lambda}^0 = \mathbb{E}\{\tau^0(\mathbf{O}, \bar{\alpha}, \beta^*, \bar{\gamma}) \tau^0(\mathbf{O}, \bar{\alpha}, \beta^*, \bar{\gamma})^\top\}.$$

(ii) *A consistent estimator of $\boldsymbol{\Sigma}^0$ is $\hat{\boldsymbol{\Sigma}}^0 = \hat{\boldsymbol{\Gamma}}^{0-1} \hat{\boldsymbol{\Lambda}}^0 \hat{\boldsymbol{\Gamma}}^{0-1}$, where $\hat{\boldsymbol{\Gamma}}^0 = \tilde{\mathbb{E}}\{\psi_1(\hat{\beta}^{0\top} \mathbf{Z}) \mathbf{Z} \mathbf{Z}^\top\}$ and*

$$\hat{\boldsymbol{\Lambda}}^0 = \tilde{\mathbb{E}}\left\{\tau^0(\mathbf{O}, \hat{\alpha}, \hat{\beta}^{0\top}, \hat{\gamma}) \tau^0(\mathbf{O}, \hat{\alpha}, \hat{\beta}^{0\top}, \hat{\gamma})^\top\right\}.$$

Thus, for a constant vector \mathbf{c} with the same dimension of β , an asymptotic $(1 - \eta)$ confidence interval for $\mathbf{c}^\top \beta^{0*}$ is $\mathbf{c}^\top \hat{\beta}^{0\top} \pm z_{\eta/2} \sqrt{\mathbf{c}^\top \hat{\boldsymbol{\Sigma}}^0 \mathbf{c} / N}$.

Theorem 2 shows that if the PS model is correct, regardless of the correctness of the OR working model, the proposed estimator $\hat{\beta}^0$ is consistent and asymptotically normal, and the proposed CIs based on $\hat{\boldsymbol{\Sigma}}^0$ are valid. Similarly to the estimation of β^* , the conclusions in Theorem 2 also hold in low-dimensional settings with a reduced form of Assumptions 3, 4 and 5.

We point out that the method of Liu et al. (2023) for CSTL can be viewed as an AIPW estimator of β^{0*} under the stratified sampling setting, where the labeled and unlabeled datasets \mathcal{L} and \mathcal{U} are treated as two independent samples of fixed sizes n and $N - n$. They employed partial linear models for both PS and OR working models. By replacing their choices of semi-parametric nuisance models with our parametric models, the estimator of β^{0*} in Liu et al. (2023) can be reformulated as the solution to the following estimating equations:

$$\frac{1}{n} \sum_{i=1}^n w(\mathbf{X}_i; \hat{\gamma}^s) [\{Y_i - \psi(\hat{\alpha}^\top \mathbf{G}_i)\} \mathbf{Z}_i] + \frac{1}{N - n} \sum_{i=n+1}^N [\{\psi(\hat{\alpha}^\top \mathbf{G}_i) - \psi(\beta^T \mathbf{Z}_i)\} \mathbf{Z}_i] = 0. \quad (26)$$

where $w(\mathbf{X}_i; \hat{\gamma}^s) = \exp(-\hat{\gamma}^{s\top} \mathbf{F}_i)$ and \mathbf{F}_i is the abbreviation of $\mathbf{F}(\mathbf{X}_i)$; $\hat{\gamma}^s = (\hat{\gamma}_0^s, \hat{\gamma}_{1:p}^{s\top})^\top$ is an estimator of the parameter γ^s in an exponential tilt model, defined as

$$dG_1 = \exp(\gamma_0^s + \gamma_{1:p}^{s\top} \mathbf{F}_{1:p}) dG_0, \quad (27)$$

where G_0 and G_1 are two probability distributions for the unlabeled and labeled data in $\mathbf{F}_{1:p}$ and $\gamma_0^s = -\log \left\{ \int \exp(\gamma_{1:p}^{s\top} \mathbf{F}_{1:p}) dG_0 \right\}$ to ensure that $\int dG_1 = 1$. The exponential tilt model (27) can be shown to be equivalent to the logistic PS model (7), where the coefficients are related as follows (Prentice and Pyke, 1979; Qin, 1998; Tian et al., 2023):

$$\gamma_0 = \gamma_0^s + \ln \left(\frac{\rho_m}{1 - \rho_m} \right), \quad \gamma_{1:p} = \gamma_{1:p}^s, \quad (28)$$

where $\rho_m = \mathbb{P}(R = 1)$, the true value of the proportion of missing data. When analyzing the asymptotic property in stratified sampling settings, we assume n/N to be constant and, consequently, assume that $\rho_m = n/N$. On the other hand, our estimating equations (24) can be rewritten as

$$\frac{1}{N} \sum_{i=1}^n w(\mathbf{X}_i; \hat{\gamma}) [\{Y_i - \psi(\hat{\alpha}^\top \mathbf{G}_i)\} \mathbf{Z}_i] + \frac{1}{N} \sum_{i=n+1}^N [\{\psi(\hat{\alpha}^\top \mathbf{G}_i) - \psi(\beta^\top \mathbf{Z}_i)\} \mathbf{Z}_i] = 0. \quad (29)$$

where $\hat{\gamma} = (\hat{\gamma}_0, \hat{\gamma}_{1:p})^\top$ is an estimator of the parameter γ in logistic PS model (7). Suppose that the estimators $\hat{\gamma}^s$ and $\hat{\gamma}$ satisfy the same relationship as (28), i.e., $\hat{\gamma}_0 = \hat{\gamma}_0^s + \ln \left(\frac{n}{N-n} \right)$ and $\hat{\gamma}_{1:p} = \hat{\gamma}_{1:p}^s$. Then, it is easily seen that $(N-n)/n \cdot w(\mathbf{X}_i; \hat{\gamma}^s) = \exp(-\hat{\gamma}^\top \mathbf{F}_i) = w(\mathbf{X}_i; \hat{\gamma})$, and the two equations (26) and (29) match each other. Therefore, the different forms of (26) and (29) can be explained by the relationship of the coefficient estimates between the exponential tilt model (27) and the logistic regression model (7).

9 Summary

We present a new AIPW method for the inference of regression coefficients in (conditional) mean models in SSL and CSTL settings. We demonstrate that various previous methods can be unified in our AIPW framework by suppressing detailed technical choices. By carefully exploiting the dependence of PS and OR models and designing estimating equations of nuisance parameters, our AIPW estimator achieves asymptotic normality, and valid CIs can be obtained, whether or not the OR working model is correctly specified, with high-dimensional data. Finite sample performances of the proposed method are confirmed by a simulation study and an application to a real-world dataset.

Currently, the proposed CIs can only achieve single robustness to the misspecification of the OR model. Doubly robust CIs can be developed using the approach of Ghosh and Tan (2022), albeit at the cost of increasing technical and numerical complexities. In addition, how to handle the case where $\lim_{n,N \rightarrow \infty} n/N \rightarrow 0$ under the random sampling process is also technically challenging, since the “positivity assumption” (Assumption 2) typical in missing data theory is violated. New analysis needs to be developed to address the problem.

References

- Aloui, A., Dong, J., Le, C. P., and Tarokh, V. (2023). Transfer learning for individual treatment effect estimation. In *Proceedings of the Thirty-Ninth Conference on Uncertainty in Artificial Intelligence*, pages 56–66.
- Alvari, H., Shaabani, E., Sarkar, S., Beigi, G., and Shakarian, P. (2019). Less is more: Semi-supervised causal inference for detecting pathogenic users in social media. In *Companion Proceedings of The 2019 World Wide Web Conference*, pages 154–161.
- Athey, S., Tibshirani, J., and Wager, S. (2019). Generalized random forests. *The Annals of Statistics*, 47:1148–1178.
- Bühlmann, P. and Van De Geer, S. (2011). *Statistics for High-dimensional Data: Methods, Theory and Applications*. Springer Science & Business Media.
- Cai, T. T. and Guo, Z. (2020). Semisupervised inference for explained variance in high dimensional linear regression and its applications. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 82:391–419.
- Castro, D. C., Walker, I., and Glocker, B. (2020). Causality matters in medical imaging. *Nature Communications*, 11:3673.
- Chakraborty, A. (2016). *Robust Semi-Parametric Inference in Semi-Supervised Settings*. PhD thesis, Harvard University.
- Chakraborty, A. and Cai, T. (2018). Efficient and adaptive linear regression in semi-supervised settings. *The Annals of Statistics*, 46:1541–1572.
- Chakraborty, A., Dai, G., and Carroll, R. J. (2022). Semi-supervised quantile estimation: Robust and efficient inference in high dimensional settings. *arXiv preprint arXiv:2201.10208*.
- Chapelle, O., Schölkopf, B., and Zien, A. (2006). *Semi-Supervised Learning*. The MIT Press.

- Chen, B. and Huang, F. (2016). Semi-supervised convolutional networks for translation adaptation with tiny amount of in-domain data. In *Proceedings of the 20th SIGNLL Conference on Computational Natural Language Learning*, pages 314–323.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21:C1–C68.
- Fan, Q., Hsu, Y.-C., Lieli, R. P., and Zhang, Y. (2022). Estimation of conditional average treatment effects with high-dimensional data. *Journal of Business & Economic Statistics*, 40:313–327.
- Ghosh, S. and Tan, Z. (2022). Doubly robust semiparametric inference using regularized calibrated estimation with high-dimensional data. *Bernoulli*, 28:1675–1703.
- Gronsbell, J. L. and Cai, T. (2017). Semi-supervised approaches to efficient evaluation of model prediction performance. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 80:579–594.
- He, Z., Sun, Y., and Li, R. (2024). Transfusion: Covariate-shift robust transfer learning for high-dimensional regression. In *International Conference on Artificial Intelligence and Statistics*, pages 703–711.
- Imbens, G. W. and Rubin, D. B. (2015). *Causal Inference for Statistics Social and Biomedical Science*. Cambridge University Press.
- Little, R. J. and Rubin, D. B. (2019). *Statistical Analysis with Missing Data*. John Wiley & Sons.
- Liu, M., Zhang, Y., Liao, K. P., and Cai, T. (2023). Augmented transfer regression learning with semi-non-parametric nuisance models. *Journal of Machine Learning Research*, 24:1–50.
- Luo, R. and Qi, X. (2017). Signal extraction approach for sparse multivariate response regression. *Journal of Multivariate Analysis*, 153:83–97.
- Molenberghs, G., Fitzmaurice, G., Kenward, M. G., Tsiatis, A., and Verbeke, G. (2015). *Handbook of Missing Data Methodology*. Chapman & Hall/CRC.
- Prentice, R. L. and Pyke, R. (1979). Logistic disease incidence models and case-control studies. *Biometrika*, 66:403–411.
- Qin, J. (1998). Inferences for case-control and semiparametric two-sample density ratio models. *Biometrika*, 85:619–630.
- Quiñonero-Candela, J., Sugiyama, M., Schwaighofer, A., and Lawrence, N. D. (2009). *Dataset Shift*

- in Machine Learning*. The MIT Press.
- Robins, J. M., Rotnitzky, A., and Zhao, L. P. (1994). Estimation of regression coefficients when some regressors are not always observed. *Journal of the American statistical Association*, 89:846–866.
- Rosenbaum, P. R. and Rubin, D. B. (1983). The central role of the propensity score in observational studies for causal. *Biometric*, 70:41–55.
- Roth, J. J. (2019). Empty homes and acquisitive crime: Does vacancy type matter? *American Journal of Criminal Justice*, 44:770–787.
- Ruder, S., Peters, M. E., Swayamdipta, S., and Wolf, T. (2019). Transfer learning in natural language processing. In *Proceedings of the 2019 Conference of the North American Chapter of the Association for Computational Linguistics: Tutorials*, pages 15–18.
- Sekhon, J. S. (2011). Multivariate and propensity score matching software with automated balance optimization: The Matching package for R. *Journal of Statistical Software*, 42:1–52.
- Sohn, K., Berthelot, D., Carlini, N., Zhang, Z., Zhang, H., Raffel, C. A., Cubuk, E. D., Kurakin, A., and Li, C.-L. (2020). Fixmatch: Simplifying semi-supervised learning with consistency and confidence. In *Advances in Neural Information Processing Systems*, pages 596–608.
- Tan, Z. (2020a). Model-assisted inference for treatment effects using regularized calibrated estimation with high-dimensional data. *The Annals of Statistics*, 48:811–837.
- Tan, Z. (2020b). Regularized calibrated estimation of propensity scores with model misspecification and high-dimensional data. *Biometrika*, 107:137–158.
- Tan, Z. and Sun, B. (2020). *RCAL: Regularized calibrated estimation*. R package version 2.0.
- Tang, C., Zeng, X., Zhou, L., Zhou, Q., Wang, P., Wu, X., Ren, H., Zhou, J., and Wang, Y. (2024). Semi-supervised medical image segmentation via hard positives oriented contrastive learning. *Pattern Recognition*, 146:110020.
- Tian, Y., Zhang, X., and Tan, Z. (2023). On semi-supervised estimation using exponential tilt mixture models. *arXiv preprint arXiv:2311.08504*.
- Van der Vaart, A. W. (2000). *Asymptotic Statistics*. Cambridge University Press.
- Wager, S. and Athey, S. (2018). Estimation and inference of heterogeneous treatment effects using random forests. *Journal of the American Statistical Association*, 113:1228–1242.

- Wu, P., Tan, Z., Hu, W., and Zhou, X. (2024). Model-assisted inference for covariate-specific treatment effects with high-dimensional data. *Statistica Sinica*, 34:459–479.
- You, K. (2023). *maotai: Tools for Matrix Algebra, Optimization and Inference*. R package version 0.2.5.
- Zhang, A., Brown, L. D., and Cai, T. T. (2019). Semi-supervised inference: General theory and estimation of means. *The Annals of Statistics*, 47:2538 – 2566.
- Zhang, Y. and Bradic, J. (2021). High-dimensional semi-supervised learning: in search of optimal inference of the mean. *Biometrika*, 109:387–403.
- Zhang, Y., Chakraborty, A., and Bradic, J. (2023). Semi-supervised causal inference: Generalizable and double robust inference for average treatment effects under selection bias with decaying overlap. *arXiv preprint arXiv:2305.12789*.
- Zhao, Y., Zheng, Y., Yu, B., Tian, Z., Lee, D., Sun, J., Li, Y., and Zhang, N. L. (2022). Semi-supervised lifelong language learning. In *Findings of the Association for Computational Linguistics: EMNLP 2022*, pages 3937–3951.
- Zheng, M., You, S., Huang, L., Wang, F., Qian, C., and Xu, C. (2022). Simmatch: Semi-supervised learning with similarity matching. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pages 14471–14481.
- Zhou, A. and Levine, S. (2021). Bayesian adaptation for covariate shift. In *Advances in Neural Information Processing Systems*, pages 914–927.
- Zhu, X. (2008). Semi-supervised learning literature survey. Technical Report No. 1530, Department of Computer Sciences, University of Wisconsin-Madison, USA.
- Zimmert, M. and Lechner, M. (2019). Nonparametric estimation of causal heterogeneity under high-dimensional confounding. *arXiv preprint arXiv:1908.08779*.

Supplementary Material for “Semi-supervised Regression Analysis with Model Misspecification and High-dimensional Data”

Ye Tian, Peng Wu and Zhiqiang Tan

This Supplementary Material consists of Sections I–VI, where Section I contains technical tools used in proofs of lemmas in Section II. Section II presents lemmas used in proofs of Proposition 2 and Theorem 1. Section III gives the technical proofs of Proposition 1 and Proposition 2. Section IV provides the proof of Theorem 1. Section V includes contents of the extension to stratified sampling settings, including the proof of Proposition 3 and the variance comparison. Section VI presents details of the application.

I Technical tools

We state the following concentration inequalities, to facilitate proofs of lemmas in Section II, which can be obtained from Bühlmann and Van De Geer (2011) [Lemmas 14.11, 14.16 & 14.9].

Lemma S1. *Let (Y_1, \dots, Y_N) be independent variables such that $\mathbb{E}(Y_i) = 0$ for $i = 1, \dots, n$ and $\max_{i=1, \dots, n} |Y_i| \leq c_0$ for some constant c_0 . Then for any $t > 0$,*

$$\mathbb{P} \left(\left| \frac{1}{N} \sum_{i=1}^N Y_i \right| > t \right) \leq 2 \exp \left(-\frac{nt^2}{2c_0^2} \right).$$

Lemma S2. *Let (Y_1, \dots, Y_N) be independent variables such that $\mathbb{E}(Y_i) = 0$ for $i = 1, \dots, n$ and (Y_1, \dots, Y_N) are uniformly sub-gaussian: $\max_{i=1, \dots, n} c_1^2 \mathbb{E}\{\exp(Y_i^2/c_1^2) - 1\} \leq c_2^2$. Then for any $t > 0$,*

$$\mathbb{P} \left(\left| \frac{1}{N} \sum_{i=1}^N Y_i \right| > t \right) \leq 2 \exp \left\{ -\frac{nt^2}{8(c_1^2 + c_2^2)} \right\}.$$

Lemma S3. *Let (Y_1, \dots, Y_N) be independent variables such that $\mathbb{E}(Y_i) = 0$ for $i = 1, \dots, n$ and,*

$$\frac{1}{N} \sum_{i=1}^N \mathbb{E}(|Y_i|^k) \leq \frac{k!}{2} c_3^{k-2} c_4^2, \quad k = 2, 3, \dots,$$

for some constants (c_3, c_4) . Then for any $t > 0$,

$$\mathbb{P} \left(\left| \frac{1}{N} \sum_{i=1}^N Y_i \right| > c_3 t + c_4 \sqrt{2t} \right) \leq 2 \exp(-nt).$$

Lemma S4. *Suppose that $|X| \leq c_5$, c_5 is some constant, and Y is sub-gaussian: $c_1^2 \mathbb{E}\{\exp\{(X^2/c_1^2)\}$*

$-1\} \leq c_2^2$ for some constants (c_1, c_2) . Then $Z = XY^2$ satisfies

$$\mathbb{E}\{|Z - \mathbb{E}(Z)|^k\} \leq \frac{k!}{2} c_6^{k-2} c_7^2, \quad k = 2, 3, \dots,$$

for $c_6 = 2c_5c_1^2$ and $c_7 = 2c_5c_1c_2$.

Lemma S5. Suppose that Y is sub-gaussian: $c_1^2 \mathbb{E}\{\exp\{(Y^2/c_1^2)\} - 1\} \leq c_2^2$ for some constants (c_1, c_2) . Then

$$\mathbb{E}(|Y|^k) \leq \Gamma\left(\frac{k}{2} + 1\right)(c_1^2 + c_2^2)c_1^{k-2}, \quad k = 2, 3, \dots$$

II Technical lemmas

II.1 Lemmas for the parameter in the PS model

The following Lemmas S6–S8 will be used in proofs of Proposition 1 and Theorem 1. Lemma S9 would be used in proofs of lemmas in Section II.2 and Theorem 1.

Lemma S6. Under Assumptions 3(i) and 3(ii), the following statements hold:

(i) Denoted by Ω_{00} the event that

$$\sup_{j=0, \dots, p} \left| \tilde{\mathbb{E}}[\{-Rw(\mathbf{X}; \bar{\gamma}) + (1 - R)\} f_j(\mathbf{X})] \right| \leq \lambda_0.$$

If $\lambda_0 \geq \sqrt{2}\{1 + \exp(-B_0)\}C_0\sqrt{\ln\{(1+p)/\epsilon\}/N}$, then $\mathbb{P}(\Omega_{00}) \geq 1 - 2\epsilon$.

(ii) Denote by Ω_{01} the event that

$$\sup_{j,k=0, \dots, p} \left| (\tilde{\Sigma}_\gamma)_{jk} - (\Sigma_\gamma)_{jk} \right| \leq \lambda_0, \tag{S1}$$

where $\tilde{\Sigma}_\gamma$ is the empirical version of Σ_γ . If $\lambda_0 \geq 4 \exp(-B_0)C_0^2\sqrt{\ln\{(1+p)/\epsilon\}/N}$, then $\mathbb{P}(\Omega_{01}) \geq 1 - 2\epsilon^2$.

The result of Lemma S6(ii) is taken from Tan (2020b)[Lemma 1(ii)], the result of Lemma S6(i) can be shown similarly using Lemma S1 in Section I and the union bound.

Let $\Sigma_{m^1} = \mathbb{E}\{Rw(\mathbf{X}; \bar{\gamma})|Y - \psi(\bar{\alpha}^T \mathbf{G})|\mathbf{F}\mathbf{F}^T\}$, and $\tilde{\Sigma}_{m^1}$ be the sample version of Σ_{m^1} .

Lemma S7. Let Ω_{02} denote the event that

$$\sup_{j,k=0, \dots, p} |(\Sigma_{m^1})_{jk} - (\tilde{\Sigma}_{m^1})_{jk}| \leq \sqrt{D_0^2 + D_1^2}\lambda_0. \tag{S2}$$

Under Assumptions 3(i), 3(ii) and 4(iv), if

$$\lambda_0 \geq 4 \exp(-B_0) C_0^2 \sqrt{\frac{\ln \left\{ \frac{(1+p)}{\epsilon} \right\}}{N}},$$

then $\mathbb{P}(\Omega_{02}) \geq 1 - 2\epsilon^2$.

Proof. Since $|Rw(\mathbf{X}; \bar{\gamma}) f_j(\mathbf{X}) f_k(\mathbf{X})| \leq \exp(-B_0) C_0^2$ for $j, k = 0, \dots, p$ by Assumptions 3(i), and 3(ii), and $|Y - \psi(\bar{\alpha}^\top \mathbf{G})|$ is uniformly sub-gaussian by Assumption 4(iv), $Rw(\mathbf{X}; \bar{\gamma}) |Y - \psi(\bar{\alpha}^\top \mathbf{G})| f_j(\mathbf{X}) f_k(\mathbf{X})$ is uniformly sub-gaussian. By Lemma S2, we have

$$\mathbb{P} \left\{ |(\boldsymbol{\Sigma}_{m^1})_{jk} - (\tilde{\boldsymbol{\Sigma}}_{m^1})_{jk}| > t \right\} \leq \frac{2\epsilon^2}{(1+p)^2},$$

for $j, k = 0, \dots, p$, where $t = \exp(-B_0) C_0^2 \sqrt{8(D_0^2 + D_1^2)} \sqrt{\ln\{(1+p)^2/\epsilon^2\}}/N$. By union bounds, (S2) holds. \square

Let $\boldsymbol{\Sigma}_{m^2} = \mathbb{E}[Rw(\mathbf{X}; \bar{\gamma}) \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 \mathbf{F} \mathbf{F}^\top]$, the sample version of $\boldsymbol{\Sigma}_{m^2}$, $\tilde{\boldsymbol{\Sigma}}_{m^2} = \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma}) \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 \mathbf{F} \mathbf{F}^\top]$.

Lemma S8. Denote by Ω_{03} the event that

$$\sup_{j,k=0,\dots,p} |(\boldsymbol{\Sigma}_{m^2})_{jk} - (\tilde{\boldsymbol{\Sigma}}_{m^2})_{jk}| \leq (D_0^2 + D_0 D_1) \lambda_0, \quad (\text{S3})$$

Under Assumptions 3(i), 3(ii) and 4(iv), if

$$(D_0^2 + D_0 D_1) \lambda_0 \geq 4 \exp(-B_0) C_0^2 [D_0^2 \ln\{(1+p)/\epsilon\}/N + D_0 D_1 \sqrt{\ln\{(1+p)/\epsilon\}}/N],$$

then, $\mathbb{P}(\Omega_{03}) \geq 1 - 2\epsilon^2$. Furthermore, if we assume that $\ln\{(1+p)/\epsilon\}/N < 1$, the above condition reduces to $\lambda_0 \geq 4 \exp(-B_0) C_0^2 \sqrt{\ln\{(1+p)/\epsilon\}}/N$.

Proof. For $j, k = 0, \dots, p$, the variable $Rw(\mathbf{X}; \bar{\gamma}) \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 f_j(\mathbf{X}) f_k(\mathbf{X})$ is the product of $Rw(\mathbf{X}; \bar{\gamma}) f_j(\mathbf{X}) f_k(\mathbf{X})$ and $\{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}^2$, where $|Rw(\mathbf{X}; \bar{\gamma}) f_j(\mathbf{X}) f_k(\mathbf{X})| \leq \exp(-B_0) C_0^2$ by Assumptions 3(i) and 3(ii); and $\{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}$ is sub-gaussian by Assumption 4(iv). By Lemmas S3 and S4 in Section I, we have

$$\mathbb{P} \left\{ |(\tilde{\boldsymbol{\Sigma}}_{m^2})_{j,k} - \boldsymbol{\Sigma}_{m^2}_{j,k}| > 2e^{(-B_0)} C_0^2 D_0^2 t + 2e^{(-B_0)} C_0^2 D_0 D_1 \sqrt{2t} \right\} \leq 2 \frac{\epsilon^2}{(1+p)^2},$$

for $j, k = 0, \dots, p$, where $t = \ln\{(1+p)^2/\epsilon^2\}/N$. The result then follows from the union bound. \square

Lemma S9. *Under the conditions of Proposition 1, in the event $\Omega_{00} \cap \Omega_{01}$, we have*

$$\tilde{\mathbb{E}} [Rw(\mathbf{X}; \bar{\gamma})\{(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}\}^2] \leq \exp(\eta_{01})M_0|S_{\bar{\gamma}}|\lambda_0^2, \quad (\text{S4})$$

where $\eta_{01} = (A_0 - 1)^{-1}M_0C_0\zeta_0$.

Proof. By Lemma S6 and Proposition 1, in the event $\Omega_{00} \cap \Omega_{01}$, (16) holds, we obtain

$$\|\hat{\gamma} - \bar{\gamma}\|_1 \leq (A_0 - 1)^{-1}M_0|S_{\bar{\gamma}}|\lambda_0 \leq (A_0 - 1)^{-1}M_0\zeta_0, \quad (\text{S5})$$

the second inequality holds due to Assumption 3(iv). By the definition of $D_{\text{CAL}}^\dagger(\cdot, \cdot)$, we obtain

$$\begin{aligned} D_{\text{CAL}}^\dagger(\hat{\gamma}^\top \mathbf{F}, \bar{\gamma}^\top \mathbf{F}) &= -\tilde{\mathbb{E}}[R\{\exp(-\hat{\gamma}^\top \mathbf{F}) - \exp(-\bar{\gamma}^\top \mathbf{F})\}\{(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}\}] \\ &= \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma}) \exp[-u\{(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}\}]\{(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}\}^2] \\ &\geq \exp(-C_0\|\hat{\gamma} - \bar{\gamma}\|_1)\tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma})\{(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}\}^2] \\ &\geq \exp(-\eta_{01})\tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma})\{(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}\}^2]. \end{aligned}$$

The second equality holds by the mean value theorem and u is some scalar in $(0, 1)$. Combining the inequality with (16), we obtain

$$\tilde{\mathbb{E}} [Rw(\mathbf{X}; \bar{\gamma})\{(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}\}^2] \leq \exp(\eta_{01})D_{\text{CAL}}^\dagger(\hat{\gamma}^\top \mathbf{F}, \bar{\gamma}^\top \mathbf{F}) \leq \exp(\eta_{01})M_0|S_{\bar{\gamma}}|\lambda_0^2.$$

□

II.2 Lemmas for the parameter in the OR model

The following Lemmas S10–S17 will be used in proofs of Proposition 2 and Theorem 1.

Lemma S10. *Let Ω_{10} denote the event that*

$$\sup_{j=0, \dots, q} \left| \tilde{\mathbb{E}} [Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}g_j(\mathbf{X})] \right| \leq \lambda_1. \quad (\text{S6})$$

Under Assumptions 3, 4(iii) and 4(iv), if

$$\lambda_1 \geq \exp(-B_0)C_4\sqrt{8(D_0^2 + D_1^2)}\sqrt{\ln\{(1+q)/\epsilon\}/N},$$

then $\mathbb{P}(\Omega_{10}) \geq 1 - 2\epsilon$.

Proof. Let $S_j = Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}g_j(\mathbf{X})$ for $j = 0, \dots, q$. Then, $\mathbb{E}(S_j) = 0$ by the definition of $\bar{\alpha}$. Under Assumptions 3 and 4(iii), $|S_j| \leq \exp(-B_0)C_4|R\{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}|$. By Assumption 4(iv),

the variables (S_0, \dots, S_q) are uniformly sub-gaussian: $\max_{j=0, \dots, q} D_2^2 \mathbb{E}\{\exp(S_j^2/D_2^2) - 1\} \leq D_3^2$, with $D_2 = \exp(-B_0)C_4D_0$, and $D_3 = e^{(-B_0)}C_4D_1$. Therefore, by Lemma S2 in Section I and the union bound, $\mathbb{P}(\Omega_{10}) \geq 1 - 2\epsilon$, if $\lambda_1 \geq \exp(-B_0)C_4\sqrt{8(D_0^2 + D_1^2)}\sqrt{\ln\{(1+q)/\epsilon\}/N}$. \square

Lemma S11. Denote by Ω_{11} the event that

$$\sup_{j,k=0, \dots, q} |(\boldsymbol{\Sigma}_{\bar{\alpha}})_{jk} - (\tilde{\boldsymbol{\Sigma}}_{\bar{\alpha}})_{jk}| \leq \lambda_1, \quad (\text{S7})$$

where $\tilde{\boldsymbol{\Sigma}}_{\bar{\alpha}}$ is the empirical version of $\boldsymbol{\Sigma}_{\bar{\alpha}}$. Under Assumptions 3(ii), 4(i) and 4(iii), if $\lambda_1 \geq 4\exp(-B_0)C_2C_4^2\sqrt{\ln\{(1+q)/\epsilon\}/N}$, then $\mathbb{P}(\Omega_{11}) \geq 1 - 2\epsilon^2$.

Proof. Notice that $|Rw(\mathbf{X}; \bar{\gamma})\psi_1(\bar{\alpha}^T \mathbf{G})g_j(\mathbf{X})g_k(\mathbf{X})| \leq e^{(-B_0)}C_2C_4^2$ for $j, k = 0, 1, \dots, q$, by Assumptions 3(ii), 4(i), and 4(iii). Thus,

$$\begin{aligned} & |Rw(\mathbf{X}; \bar{\gamma})\psi_1(\bar{\alpha}^T \mathbf{G})g_j(\mathbf{X})g_k(\mathbf{X}) - \mathbb{E}\{Rw(\mathbf{X}; \bar{\gamma})\psi_1(\bar{\alpha}^T \mathbf{G})g_j(\mathbf{X})g_k(\mathbf{X})\}| \\ & \leq 2\exp(-B_0)C_2C_4^2. \end{aligned}$$

By union bounds and applying Lemma S1 yields (S7). \square

Lemma S12. For any $\alpha \in \mathbb{R}^{q+1}$, we have

$$\begin{aligned} & D_{\text{WL}}^\dagger(\hat{\alpha}^T \mathbf{G}, \alpha^T \mathbf{G}; \hat{\gamma}) + \lambda_\alpha \|\hat{\alpha}_{1:q}\|_1 \\ & \leq (\hat{\alpha} - \alpha)^T \tilde{\mathbb{E}}[Rw(\mathbf{X}; \hat{\gamma})\{Y - \psi(\alpha^T \mathbf{G})\}\mathbf{G}] + \lambda_\alpha \|\alpha_{1:q}\|_1. \end{aligned} \quad (\text{S8})$$

Proof. For any $u \in (0, 1]$, by definition of $\hat{\alpha}$, we have

$$\ell_{\text{WL}}(\hat{\alpha}, \hat{\gamma}) + \lambda_\alpha \|\hat{\alpha}_{1:q}\|_1 \leq \ell_{\text{WL}}\{(1-u)\hat{\alpha} + u\alpha; \hat{\gamma}\} + \lambda_\alpha \|(1-u)\hat{\alpha}_{1:q} + u\alpha_{1:q}\|_1,$$

which implies

$$\ell_{\text{WL}}(\hat{\alpha}, \hat{\gamma}) - \ell_{\text{WL}}\{(1-u)\hat{\alpha} + u\alpha; \hat{\gamma}\} + \lambda_\alpha u \|\hat{\alpha}_{1:q}\|_1 \leq \lambda_\alpha u \|\alpha_{1:q}\|_1,$$

by the convexity of L_1 -norm. Dividing both sides of the preceding inequality by u and letting $u \rightarrow 0_+$ leads to

$$-\tilde{\mathbb{E}}[Rw(\mathbf{X}; \hat{\gamma})\{Y - \psi(\hat{\alpha}^T \mathbf{G})\}\{(\hat{\alpha} - \alpha)^T \mathbf{G}\}] + \lambda_\alpha \|\hat{\alpha}_{1:q}\|_1 \leq \lambda_\alpha \|\alpha_{1:q}\|_1,$$

which yields (S8) after rearranging using (18). \square

Lemma S13. For any function $h(\mathbf{X})$, under the conditions of Proposition 1, in the event $\Omega_{00} \cap \Omega_{01}$,

$$D_{\text{WL}}^\dagger\{\hat{\alpha}^\text{T}\mathbf{G}, h(\mathbf{X}); \hat{\gamma}\} \geq \exp(-\eta_{01})D_{\text{WL}}^\dagger\{\hat{\alpha}^\text{T}\mathbf{G}, h(\mathbf{X}); \bar{\gamma}\}. \quad (\text{S9})$$

Proof. By the definition of $D_{\text{WL}}^\dagger(\cdot, \cdot; \cdot)$,

$$\begin{aligned} & D_{\text{WL}}^\dagger\{\hat{\alpha}^\text{T}\mathbf{G}, h(\mathbf{X}); \hat{\gamma}\} \\ &= \tilde{\mathbb{E}}(Rw(\mathbf{X}; \hat{\gamma})[\psi(\hat{\alpha}^\text{T}\mathbf{G}) - \psi\{h(\mathbf{X})\}]\{\hat{\alpha}^\text{T}\mathbf{G} - h(\mathbf{X})\}) \\ &= \tilde{\mathbb{E}}(Rw(\mathbf{X}; \bar{\gamma}) \exp\{-(\hat{\gamma} - \bar{\gamma})^\text{T}\mathbf{F}\}[\psi(\hat{\alpha}^\text{T}\mathbf{G}) - \psi\{h(\mathbf{X})\}]\{\hat{\alpha}^\text{T}\mathbf{G} - h(\mathbf{X})\}) \\ &\geq \tilde{\mathbb{E}}(Rw(\mathbf{X}; \bar{\gamma}) \exp(-\eta_{01})[\psi(\hat{\alpha}^\text{T}\mathbf{G}) - \psi\{h(\mathbf{X})\}]\{\hat{\alpha}^\text{T}\mathbf{G} - h(\mathbf{X})\}) \\ &= \exp(-\eta_{01})\tilde{\mathbb{E}}(Rw(\mathbf{X}; \bar{\gamma})[\psi(\hat{\alpha}^\text{T}\mathbf{G}) - \psi\{h(\mathbf{X})\}]\{\hat{\alpha}^\text{T}\mathbf{G} - h(\mathbf{X})\}) \\ &= \exp(-\eta_{01})D_{\text{WL}}^\dagger\{\hat{\alpha}^\text{T}\mathbf{G}, h(\mathbf{X}); \bar{\gamma}\}. \end{aligned}$$

The inequality holds since in the event $\Omega_{00} \cap \Omega_{01}$, (16) holds. \square

For functions $h(\mathbf{X})$ and $h'(\mathbf{X})$, let $Q_{\text{WL}}\{h(\mathbf{X}), h'(\mathbf{X}); \cdot\} = \tilde{\mathbb{E}}[Rw(\mathbf{X}; \cdot)\{h(\mathbf{X}) - h'(\mathbf{X})\}^2]$.

Lemma S14. Suppose Assumption 4(iv) holds, in the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03}$, (S8) implies

$$\begin{aligned} & \exp(-\eta_{01})D_{\text{WL}}^\dagger(\hat{\alpha}^\text{T}\mathbf{G}, \bar{\alpha}^\text{T}\mathbf{G}; \bar{\gamma}) + \lambda_\alpha \|\hat{\alpha}_{1:q}\|_1 \\ &\leq (\hat{\alpha} - \bar{\alpha})^\text{T} \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^\text{T}\mathbf{G})\}\mathbf{G}] + \lambda_\alpha \|\bar{\alpha}_{1:q}\|_1 \\ &\quad + \exp(\eta_{01})\sqrt{M_1|S_{\bar{\gamma}}|\lambda_0^2}\{Q_{\text{WL}}(\hat{\alpha}^\text{T}\mathbf{G}, \bar{\alpha}^\text{T}\mathbf{G}; \bar{\gamma})\}^{1/2}, \end{aligned}$$

where $M_1 = \left[(2D_0^2 + D_1^2 + D_0D_1) \left\{ \frac{M_0^2\zeta_0}{(A_0-1)^2} \right\} + (D_0^2 + D_1^2) \exp(\eta_{01})M_0 \right]$.

Proof. Consider the following decomposition,

$$\begin{aligned} & (\hat{\alpha} - \bar{\alpha})^\text{T} \tilde{\mathbb{E}}[Rw(\mathbf{X}; \hat{\gamma})\{Y - \psi(\bar{\alpha}^\text{T}\mathbf{G})\}\mathbf{G}] \\ &= (\hat{\alpha} - \bar{\alpha})^\text{T} \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^\text{T}\mathbf{G})\}\mathbf{G}] \\ &\quad + \tilde{\mathbb{E}}[R\{w(\mathbf{X}; \hat{\gamma}) - w(\mathbf{X}; \bar{\gamma})\}\{Y - \psi(\bar{\alpha}^\text{T}\mathbf{G})\}\{(\hat{\alpha} - \bar{\alpha})^\text{T}\mathbf{G}\}], \end{aligned} \quad (\text{S10})$$

denoted as $\Delta_0^0 + \Delta_1^0$. By the mean value theorem and Cauchy-Schwartz inequality,

$$\begin{aligned} \Delta_1^0 &\leq \exp(C_0\|\hat{\gamma} - \bar{\gamma}\|_1)\tilde{\mathbb{E}}^{1/2}[Rw(\mathbf{X}; \bar{\gamma})\{(\hat{\alpha} - \bar{\alpha})^\text{T}\mathbf{G}\}^2] \\ &\quad \times \tilde{\mathbb{E}}^{1/2}[Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^\text{T}\mathbf{G})\}^2\{(\hat{\gamma} - \bar{\gamma})^\text{T}\mathbf{F}\}^2] \\ &\leq \exp(\eta_{01})\{Q_{\text{WL}}(\hat{\alpha}^\text{T}\mathbf{G}, \bar{\alpha}^\text{T}\mathbf{G}; \bar{\gamma})\}^{1/2} \\ &\quad \times \tilde{\mathbb{E}}^{1/2}[Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^\text{T}\mathbf{G})\}^2\{(\hat{\gamma} - \bar{\gamma})^\text{T}\mathbf{F}\}^2]. \end{aligned} \quad (\text{S11})$$

We bound the third term in (S11). By Assumption 4(iv) and Lemma S5, we have

$$\mathbb{E}[\{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2 | \mathbf{X}] \leq D_0^2 + D_1^2.$$

Therefore,

$$\mathbb{E}[Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2 \{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2] \leq (D_0^2 + D_1^2) \mathbb{E}[Rw(\mathbf{X}; \bar{\gamma})\{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2].$$

Let $(\tilde{\mathbb{E}} - \mathbb{E})(\mathbf{U})$ denote $\tilde{\mathbb{E}}\{\mathbf{U} - \mathbb{E}(\mathbf{U})\}$ for \mathbf{U} , which is a function of (\mathbf{X}, R, Y) . Then in the event Ω_{01} , by (S1), we have

$$(\mathbb{E} - \tilde{\mathbb{E}})[Rw(\mathbf{X}; \bar{\gamma})\{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2] \leq \lambda_0 \|\hat{\gamma} - \bar{\gamma}\|_1^2.$$

In the event Ω_{03} , by S7, we have

$$(\tilde{\mathbb{E}} - \mathbb{E})[Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2 \{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2] \leq (D_0^2 + D_0 D_1) \lambda_0 \|\hat{\gamma} - \bar{\gamma}\|_1^2.$$

Combining preceding inequalities, we obtain in $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03}$,

$$\begin{aligned} & \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2 \{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2] \\ & \leq (D_0^2 + D_0 D_1) \lambda_0 \|\hat{\gamma} - \bar{\gamma}\|_1^2 \\ & + (D_0^2 + D_1^2) \left\{ \lambda_0 \|\hat{\gamma} - \bar{\gamma}\|_1^2 + \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma})\{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2] \right\} \\ & \leq (2D_0^2 + D_1^2 + D_0 D_1) \left\{ \frac{M_0^2 \zeta_0}{(A_0 - 1)^2} \right\} |S_{\bar{\gamma}}| \lambda_0^2 + (D_0^2 + D_1^2) \exp(\eta_{01}) M_0 |S_{\bar{\gamma}}| \lambda_0^2 \\ & = M_1 |S_{\bar{\gamma}}| \lambda_0^2, \end{aligned} \tag{S12}$$

where the last inequality holds due to (16) and (S4). Combining (S10) – (S12), we obtain

$$\begin{aligned} & (\hat{\alpha} - \bar{\alpha})^T \tilde{\mathbb{E}}[Rw(\mathbf{X}; \hat{\gamma})\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \mathbf{G}] \\ & \leq (\hat{\alpha} - \bar{\alpha})^T \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \mathbf{G}] \\ & + \exp(\eta_{01}) (M_1 |S_{\bar{\gamma}}| \lambda_0^2)^{1/2} \{Q_{\text{WL}}(\hat{\alpha}^T \mathbf{G}, \bar{\alpha}^T \mathbf{G}; \bar{\gamma})\}^{1/2}. \end{aligned} \tag{S13}$$

The desired result follows by combining (S8), (S9) and (S13) in the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03}$. \square

Lemma S15. Denote $b = \hat{\alpha} - \bar{\alpha}$. Suppose Assumption 4(iv) holds. In the event, $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03} \cap \Omega_{10}$, we have

$$\begin{aligned} & \exp(-\eta_{01}) D_{\text{WL}}^\dagger(\hat{\alpha}^T \mathbf{G}, \bar{\alpha}^T \mathbf{G}; \bar{\gamma}) + (A_1 - 1) \lambda_1 \|b\|_1 \\ & \leq 2A_1 \lambda_1 \sum_{j \in S_{\hat{\alpha}}} |b_j| + \exp(\eta_{01}) \sqrt{M_1 |S_{\bar{\gamma}}| \lambda_0^2} \{Q_{\text{WL}}(\hat{\alpha}^T \mathbf{G}, \bar{\alpha}^T \mathbf{G}; \bar{\gamma})\}^{1/2}. \end{aligned} \tag{S14}$$

Proof. In the event Ω_{10} , we have

$$b^T \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma})\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \mathbf{G}] \leq \lambda_1 \|b\|_1,$$

by which and Lemma S14, we have in the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03} \cap \Omega_{10}$,

$$\begin{aligned} & \exp(-\eta_{01}) D_{\text{WL}}^\dagger(\hat{\alpha}^T \mathbf{G}, \bar{\alpha}^T \mathbf{G}; \bar{\gamma}) + A_1 \lambda_1 \|\hat{\alpha}_{1:q}\|_1 \\ & \leq \lambda_1 \|b\|_1 + A_1 \lambda_1 \|\bar{\alpha}_{1:q}\|_1 + \exp(\eta_{01}) \sqrt{M_1 |S_{\bar{\gamma}}| \lambda_0^2} \{Q_{\text{WL}}(\hat{\alpha}^T \mathbf{G}, \bar{\alpha}^T \mathbf{G}; \bar{\gamma})\}^{1/2}. \end{aligned}$$

Applying to the preceding inequality the identity $\hat{\alpha}_j = |\hat{\alpha}_j - \bar{\alpha}_j|$ for $j \notin S_{\bar{\alpha}}$ and the triangle inequality

$$|\hat{\alpha}_j| \geq |\bar{\alpha}_j| - |\hat{\alpha}_j - \bar{\alpha}_j|, \quad j \in S_{\bar{\alpha}} \setminus \{0\},$$

and rearranging the result gives

$$\begin{aligned} & \exp(-\eta_{01}) D_{\text{WL}}^\dagger(\hat{\alpha}^T \mathbf{G}, \bar{\alpha}^T \mathbf{G}; \bar{\gamma}) + (A_1 - 1) \lambda_1 \|b_{1:q}\|_1 \\ & \leq \lambda_1 |b_0| + 2A_1 \lambda_1 \sum_{j \in S_{\bar{\alpha}} \setminus \{0\}} |b_j| + \exp(\eta_{01}) \sqrt{M_1 |S_{\bar{\gamma}}| \lambda_0^2} \{Q_{\text{WL}}(\hat{\alpha}^T \mathbf{G}, \bar{\alpha}^T \mathbf{G}; \bar{\gamma})\}^{1/2} \\ & \leq (A_1 + 1) \lambda_1 |b_0| + 2A_1 \lambda_1 \sum_{j \in S_{\bar{\alpha}} \setminus \{0\}} |b_j| + \exp(\eta_{01}) \sqrt{M_1 |S_{\bar{\gamma}}| \lambda_0^2} \{Q_{\text{WL}}(\hat{\alpha}^T \mathbf{G}, \bar{\alpha}^T \mathbf{G}; \bar{\gamma})\}^{1/2}. \end{aligned}$$

By adding $(A_1 - 1) \lambda_1 |b_0|$ on both sides of the previous inequality, the conclusion follows. \square

Lemma S16. *Suppose Assumptions 4(ii) and 4(iii) hold. Then, for any $\alpha, \tilde{\alpha} \in \mathbb{R}^{q+1}$,*

$$D_{\text{WL}}^\dagger(\alpha^T \mathbf{G}, \tilde{\alpha}^T \mathbf{G}; \bar{\gamma}) \geq \{b^T \tilde{\Sigma}_\alpha(\tilde{\alpha}) b\} \frac{1 - \exp(-C_{40} \|b\|_1)}{C_{40} \|b\|_1},$$

where $b = \alpha - \tilde{\alpha}$, $C_{40} = C_3 C_4$ and $\tilde{\Sigma}_\alpha(\cdot) = \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma}) \psi_1\{(\cdot)^T \mathbf{G}\} \mathbf{G} \mathbf{G}^T]$. Throughout, set $\{1 - \exp(-c)\}/c = 1$, for $c = 0$.

Proof. By the definition of $D_{\text{WL}}^\dagger(\cdot, \cdot; \cdot)$,

$$\begin{aligned} & D_{\text{WL}}^\dagger(\alpha^T \mathbf{G}, \tilde{\alpha}^T \mathbf{G}; \bar{\gamma}) \\ & = \tilde{\mathbb{E}}[Rw(\mathbf{X}; \bar{\gamma})\{\psi(\alpha^T \mathbf{G}) - \psi(\tilde{\alpha}^T \mathbf{G})\}(\alpha^T \mathbf{G} - \tilde{\alpha}^T \mathbf{G})] \\ & = \tilde{\mathbb{E}}\left(Rw(\mathbf{X}; \bar{\gamma}) \left[\int_0^1 \psi_1\{\tilde{\alpha}^T \mathbf{G} + u(\alpha^T \mathbf{G} - \tilde{\alpha}^T \mathbf{G})\} du \right] \{\alpha^T \mathbf{G} - \tilde{\alpha}^T \mathbf{G}\}^2\right). \end{aligned}$$

By Assumptions 4(ii), 4(iii), and the fact that $|\alpha^T \mathbf{G} - \tilde{\alpha}^T \mathbf{G}| \leq \{\sup_{j=0, \dots, q} |g_j(\mathbf{X})|\} \|\alpha - \tilde{\alpha}\|_1 \leq C_4 \|\alpha - \tilde{\alpha}\|_1$, it follows that

$$D_{\text{WL}}^\dagger(\alpha^T \mathbf{G}, \tilde{\alpha}^T \mathbf{G}; \bar{\gamma})$$

$$\begin{aligned}
&\geq \tilde{\mathbb{E}} \left[R w(\mathbf{X}; \bar{\gamma}) \left\{ \int_0^1 \psi_1(\tilde{\alpha}^\top \mathbf{G}) \exp(-u C_3 |\alpha^\top \mathbf{G} - \tilde{\alpha}^\top \mathbf{G}|) du \right\} (\alpha^\top \mathbf{G} - \tilde{\alpha}^\top \mathbf{G})^2 \right] \\
&\geq \tilde{\mathbb{E}} \{ R w(\mathbf{X}; \bar{\gamma}) \psi_1(\tilde{\alpha}^\top \mathbf{G}) (\alpha^\top \mathbf{G} - \tilde{\alpha}^\top \mathbf{G})^2 \} \left\{ \int_0^1 \exp(-u C_{40} \|\tilde{\alpha} - \alpha\|_1) du \right\},
\end{aligned}$$

which gives the desired result since $\int_0^1 \exp(-cu) du = \{1 - \exp(-c)\}/c$ for $c \geq 0$. \square

Lemma S17. *Suppose that Assumption 4(v) holds. In the event Ω_{11} , Assumption 4(vi) implies a compatibility condition for $\tilde{\Sigma}_{\bar{\alpha}}$: for any vector $b = (b_0, \dots, b_q)^\top$ such that $\sum_{j \notin S_{\bar{\alpha}}} |b_j| \leq \xi_1 \sum_{j \in S_{\bar{\alpha}}} |b_j|$, we have*

$$(1 - \zeta_1) \nu_1^2 \left(\sum_{j \in S_{\bar{\alpha}}} |b_j| \right)^2 \leq |S_{\bar{\alpha}}| (b^\top \tilde{\Sigma}_{\bar{\alpha}} b). \quad (\text{S15})$$

Proof. In the event Ω_{11} , we have $|b^\top (\tilde{\Sigma}_{\bar{\alpha}} - \Sigma_{\bar{\alpha}}) b| \leq \lambda_1 \|b\|_1^2$. Then Assumption 4(v) implies that for any $b = (b_0, \dots, b_q)^\top$ satisfying $\sum_{j \notin S_{\bar{\alpha}}} |b_j| \leq \xi_1 \sum_{j \in S_{\bar{\alpha}}} |b_j|$,

$$\begin{aligned}
\nu_1^2 \|b_{S_{\bar{\alpha}}}\|_1^2 &\leq |S_{\bar{\alpha}}| (b^\top \Sigma_{\bar{\alpha}} b) \leq |S_{\bar{\alpha}}| (b^\top \tilde{\Sigma}_{\bar{\alpha}} b + \lambda_1 \|b\|_1^2) \\
&\leq |S_{\bar{\alpha}}| (b^\top \tilde{\Sigma}_{\bar{\alpha}} b) + |S_{\bar{\alpha}}| \lambda_1 (1 + \xi_1)^2 \|b_{S_{\bar{\alpha}}}\|_1^2 \\
&\leq |S_{\bar{\alpha}}| (b^\top \tilde{\Sigma}_{\bar{\alpha}} b) + \zeta_1 \nu_1^2 \|b_{S_{\bar{\alpha}}}\|_1^2
\end{aligned}$$

where $\|b_{S_{\bar{\alpha}}}\|_1 = \sum_{j \in S_{\bar{\alpha}}} |b_j|$; and the last inequality holds due to Assumption 4(vi), $(1 + \xi_1)^2 \nu_1^{-2} |S_{\bar{\alpha}}| \lambda_1 \leq \zeta_1$. Thus (S15) follows by rearrangement. \square

III Proofs of Propositions 1 and 2

III.1 Proof of Proposition 1

Proof. Let c_γ in Assumption 4(iv) = $\max[\sqrt{2}\{1 + \exp(-B_0)\}C_0, 4\exp(-B_0)C_0^2]$. This can be shown similarly to Tan (2020a) [Theorem 1] by Lemmas S6–S8 and Lemmas similar to S10–S17. The small difference in probability is due to extra constraints of λ_0 on Ω_{02} and Ω_{03} from the sequential estimate, which is also demonstrated in Tan (2020a) [Theorem 5]. \square

III.2 Proof of Proposition 2

Proof. To facilitate the proof, we first define some constants. Let $\nu_2 = \nu_1(1 - \zeta_1)^{1/2}$, $\xi_2 = 1 - 2A_1/\{(\xi_1 + 1)(A_1 - 1)\}$, $\xi_3 = (\xi_1 + 1)(A_1 - 1)$, c_α in Assumption 4(vi) equals to $\max\{C_4\sqrt{8(D_0^2 + D_1^2)},$

$4C_2C_4^2\} \exp(-B_0)$, \tilde{c}_α and \tilde{c}_γ in Assumption 4(vii) equal to $C_3C_4 \exp(\eta_{01})(A_1 - 1)^{-1}\xi_3^2\nu_2^{-2}$ and $C_1^{-1}C_3C_4(A_1 - 1)^{-1}\xi_2^{-2} \exp(3\eta_{01})M_1$, respectively.

Denote $b = \hat{\alpha} - \bar{\alpha}$, $D_{\text{WL}}^\dagger = D_{\text{WL}}^\dagger(\hat{\alpha}^\top \mathbf{G}, \bar{\alpha}^\top \mathbf{G}; \bar{\gamma})$, $Q_{\text{WL}} = Q_{\text{WL}}(\hat{\alpha}^\top \mathbf{G}, \bar{\alpha}^\top \mathbf{G}; \bar{\gamma})$ and $D_{\text{WL}}^\dagger = \exp(-\eta_{01})D_{\text{WL}}^\dagger + (A_1 - 1)\lambda_1\|b\|_1$.

In the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03} \cap \Omega_{10} \cap \Omega_{11}$, (S14) in Lemma S15 leads to two possible cases: either

$$\xi_2 D_{\text{WL}}^\dagger \leq \exp(\eta_{01})(M_1 |S_{\bar{\gamma}}| \lambda_0^2 Q_{\text{WL}})^{1/2}, \quad (\text{S16})$$

or $(1 - \xi_2)D_{\text{WL}}^\dagger \leq 2A_1\lambda_1 \sum_{j \in S_{\bar{\alpha}}} |b_j|$, i.e.,

$$D_{\text{WL}}^\dagger \leq (\xi_1 + 1)(A_1 - 1)\lambda_1 \sum_{j \in S_{\bar{\alpha}}} |b_j| = \xi_3 \lambda_1 \sum_{j \in S_{\bar{\alpha}}} |b_j|. \quad (\text{S17})$$

By Lemma S16, we have

$$D_{\text{WL}}^\dagger \geq \{b^\top \tilde{\Sigma}_\alpha(\bar{\alpha})b\} \frac{1 - \exp(-C_{40}\|b\|_1)}{C_{40}\|b\|_1} = (b^\top \tilde{\Sigma}_{\bar{\alpha}}b) \frac{1 - \exp(-C_{40}\|b\|_1)}{C_{40}\|b\|_1}. \quad (\text{S18})$$

If (S16) holds, notice that $D_{\text{WL}}^\dagger \leq \exp(\eta_{01})D_{\text{WL}}^\dagger$ and by Assumption 4(i), $Q_{\text{WL}} < C_1^{-1}(b^\top \tilde{\Sigma}_\alpha b)$, which together with (S18) yields

$$D_{\text{WL}}^\dagger \leq \exp(3\eta_{01})\xi_2^{-2}C_1^{-1}(M_1 |S_{\bar{\gamma}}| \lambda_0^2) \frac{C_{40}\|b\|_1}{1 - \exp(-C_{40}\|b\|_1)}. \quad (\text{S19})$$

Since $(A_1 - 1)\lambda_1\|b\|_1 \leq D_{\text{WL}}^\dagger$ and Assumption 4(vii) holds, (S19) implies that

$$1 - \exp(-C_{40}\|b\|_1) \leq (A_1 - 1)^{-1} \exp(3\eta_{01})\xi_2^{-2}C_1^{-1}(M_1 |S_{\bar{\gamma}}| \lambda_0)C_{40} \leq \eta_3 < 1.$$

As a result, $C_{40}\|b\|_1 \leq -\ln(1 - \eta_3)$, which leads to

$$\frac{1 - \exp(-C_{40}\|b\|_1)}{C_{40}\|b\|_1} = \int_0^1 \exp(-C_{40}\|b\|_1 u) du \geq \exp(-C_{40}\|b\|_1) \geq 1 - \eta_3. \quad (\text{S20})$$

Combining the inequality with (S19), we obtain

$$D_{\text{WL}}^\dagger \leq \exp(3\eta_{01})\xi_2^{-2}\{C_1(1 - \eta_3)\}^{-1}(M_1 |S_{\bar{\gamma}}| \lambda_0^2)$$

If (S17) holds, then $\sum_{j \notin S_{\bar{\alpha}}} |b_j| \leq \xi_1 \sum_{j \in S_{\bar{\alpha}}} |b_j|$, which, together with Assumptions 4(v)–4(vi), implies (S15) in Lemma S17, that is,

$$\sum_{j \in S_{\bar{\alpha}}} |b_j| \leq (1 - \zeta_1)^{-1/2} \nu_1^{-1} |S_{\bar{\alpha}}|^{1/2} \left(b^\top \tilde{\Sigma}_{\bar{\alpha}} b \right)^{1/2}. \quad (\text{S21})$$

Since $D_{\text{WL}}^\dagger \leq \exp(\eta_{01})D_{\text{WL}}^\dagger$, combining (S17), (S18) and (S21) yields

$$D_{\text{WL}}^\dagger \leq \exp(\eta_{01})\xi_3^2(1 - \zeta_1)^{-1}\nu_1^{-2}|S_{\bar{\alpha}}|\lambda_1^2 \frac{C_{40}\|b\|_1}{1 - \exp(-C_{40}\|b\|_1)}. \quad (\text{S22})$$

Since $(A_1 - 1)\lambda_1\|b\|_1 \leq D_{\text{WL}}^\dagger$ and Assumption 4(ii) holds, (S22) implies that

$$1 - \exp(-C_{40}\|b\|_1) \leq (A_1 - 1)^{-1} \exp(\eta_{01})\xi_3^2(1 - \zeta_1)^{-1}\nu_1^{-2}|S_{\bar{\alpha}}|\lambda_1 C_{40} \leq \eta_2 < 1.$$

As a result, $C_{40}\|b\|_1 \leq -\ln(1 - \eta_2)$, which leads to

$$\frac{1 - \exp(-C_{40}\|b\|_1)}{C_{40}\|b\|_1} = \int_0^1 \exp(-C_{40}\|b\|_1 u) du \geq \exp(-C_{40}\|b\|_1) \geq 1 - \eta_2.$$

Combining the inequality with (S22), we obtain $D_{\text{WL}}^\dagger \leq \exp(\eta_{01})\xi_3^2\nu_2^{-2}(1 - \eta_2)^{-2}|S_{\bar{\alpha}}|\lambda_1^2$.

Therefore, we obtain

$$\begin{aligned} & D_{\text{WL}}^\dagger(\hat{\alpha}^\top \mathbf{G}, \bar{\alpha}^\top \mathbf{G}, \bar{\gamma}) + \exp(\eta_{01})(A_1 - 1)\lambda_1\|\hat{\alpha} - \bar{\alpha}\|_1 \\ & \leq \exp(4\eta_{01})\xi_2^{-2}\{C_1(1 - \eta_3)\}^{-1}(M_1|S_{\bar{\gamma}}|\lambda_0^2) + \exp(2\eta_{01})\xi_3^2\nu_2^{-2}(1 - \eta_2)^{-2}|S_{\bar{\alpha}}|\lambda_1^2, \end{aligned}$$

Let $M_{11} = \exp(4\eta_{01})\xi_2^{-2}\{C_1(1 - \eta_3)\}^{-1}M_1$ and $M_{12} = \exp(2\eta_{01})\xi_3^2\nu_2^{-2}(1 - \eta_2)^{-2}$, then (17) holds. \square

IV Proof of Theorem 1

IV.1 Lemmas for the proposed estimator

Lemma S18. *Suppose that Assumptions 3(i), 4(i), 4(ii), and 4(vi) hold, if*

$$\lambda_0 \geq \sqrt{2}\{1 + \exp(-B_0)\}C_0\sqrt{\frac{\ln(\frac{1+P}{\epsilon})}{N}},$$

then, $\mathbb{P}(\Omega_{20}) \geq 1 - 2\epsilon$ for any $r \geq 0$, where Ω_{20} denotes the event

$$\sup_{\|\alpha - \bar{\alpha}\|_1 \leq r; j=0, \dots, d} \left| \left(\tilde{\mathbb{E}} - \mathbb{E} \right) \left[\left\{ \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} - 1 \right\} \{ \psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\bar{\alpha}^\top \mathbf{G}) \} Z_j \right] \right| \leq B_1 \lambda_0 r,$$

where B_1 is a positive constant, depending on (C_0, C_2, C_3, C_5) .

Proof. This can be shown similarly to Lemma 13 in the Supplement of Tan (2020a). \square

Lemma S19. *Suppose Assumptions 3, 4 and 5 hold, if a function $h(\cdot)$ on a set of samples satisfying $h[\{\mathbf{X}_i\}_{i=1}^N] \leq M\{|S_{\bar{\gamma}}|\lambda_0(\epsilon) + |S_{\bar{\alpha}}|\lambda_1(\epsilon)\}$ for some constant M with probability $1 - c\epsilon$ for some constant $c > 0$ and any $\epsilon > 0$, then, $h[\{\mathbf{X}_i\}_{i=1}^N] = o_p(1)$.¹²*

¹²We write λ_0, λ_1 as $\lambda_0(\epsilon), \lambda_1(\epsilon)$, since we treat them as functions of ϵ .

Proof. For $\forall \epsilon > 0$, let Ω_ϵ be the event where $h[\{\mathbf{X}_i\}_{i=1}^N] \leq M\{|S_{\bar{\gamma}}|\lambda_0(\epsilon/c) + |S_{\bar{\alpha}}|\lambda_1(\epsilon/c)\}$ holds. Suppose $|S_{\bar{\gamma}}|\lambda_0(\epsilon/c) < |S_{\bar{\alpha}}|\lambda_1(\epsilon/c)$, then, on Ω_ϵ ,

$$\frac{h[\{\mathbf{X}_i\}_{i=1}^N]\sqrt{N}}{|S_{\bar{\alpha}}|\sqrt{\ln\{e(q+1)\}}} \leq 2M\sqrt{\frac{\ln\left(\frac{q+1}{\epsilon/c}\right)}{\ln\{e(q+1)\}}} \leq 2M\sqrt{\frac{\ln(q+1)}{\ln(q+1)} + \ln\left(\frac{c}{\epsilon}\right)} \leq 2M\sqrt{1 + \ln\left(\frac{c}{\epsilon}\right)}, \quad (\text{S23})$$

which implies that

$$\mathbb{P}\left[\frac{h[\{\mathbf{X}_i\}_{i=1}^N]\sqrt{N}}{|S_{\bar{\alpha}}|\sqrt{\ln\{e(q+1)\}}} \leq 2M\sqrt{1 + \ln\left(\frac{c}{\epsilon}\right)}\right] \geq \mathbb{P}(\Omega_\epsilon) = 1 - \epsilon.$$

Similarly, suppose $|S_{\bar{\alpha}}|\lambda_1 < |S_{\bar{\gamma}}|\lambda_0$, then, on Ω_ϵ ,

$$\mathbb{P}\left[\frac{h[\{\mathbf{X}_i\}_{i=1}^N]\sqrt{N}}{|S_{\bar{\gamma}}|\sqrt{\ln\{e(p+1)\}}} \leq 2M\sqrt{1 + \ln\left(\frac{c}{\epsilon}\right)}\right] \geq \mathbb{P}(\Omega_\epsilon) = 1 - \epsilon.$$

For $\forall \epsilon > 0$, we have

$$\mathbb{P}\left[\frac{h[\{\mathbf{X}_i\}_{i=1}^N]\sqrt{N}}{|S_{\bar{\alpha}}|\sqrt{\ln\{e(q+1)\}} + |S_{\bar{\gamma}}|\sqrt{\ln\{e(p+1)\}}} \leq 2M\sqrt{1 + \ln\left(\frac{c}{\epsilon}\right)}\right] \geq 1 - \epsilon,$$

thus, $\frac{h[\{\mathbf{X}_i\}_{i=1}^N]\sqrt{N}}{|S_{\bar{\alpha}}|\sqrt{\ln\{e(q+1)\}} + |S_{\bar{\gamma}}|\sqrt{\ln\{e(p+1)\}}} = O_p(1)$, and by Assumption 5(vi), it follows that

$$h[\{\mathbf{X}_i\}_{i=1}^N] = o_p\left[\frac{|S_{\bar{\alpha}}|\sqrt{\ln\{e(q+1)\}} + |S_{\bar{\gamma}}|\sqrt{\ln\{e(p+1)\}}}{\sqrt{N}}\right] = o_p(1).$$

□

Lemma S20. *Suppose Assumptions 3, 4 and 5 hold, if a function $h(\cdot)$ on a set of samples satisfying $h[\{\mathbf{X}_i\}_{i=1}^N] \leq M(|S_{\bar{\gamma}}| + |S_{\bar{\alpha}}|)\lambda_0(\epsilon)\lambda_1(\epsilon)$ for some constant M with probability $1 - c\epsilon$ for some constant c and any $\epsilon > 0$, then, $h[\{\mathbf{X}_i\}_{i=1}^N] = o_p(1/\sqrt{N})$.*

Proof. By similar trick used in the proof of Lemma S19, it can be easily shown that $h[\{\mathbf{X}_i\}_{i=1}^N] = o_p\left[\frac{(|S_{\bar{\gamma}}| + |S_{\bar{\alpha}}|)\sqrt{\ln\{e(q+1)\}}\sqrt{\ln\{e(p+1)\}}}{N}\right] = o_p(1/\sqrt{N})$ by Assumption 5(vi). □

Lemma S21. *Under Assumptions 3, 4 and 5, suppose either the PS model (7) is correct or the OR model (8) is correct, the AIPW estimator $\hat{\beta} \xrightarrow{\mathbb{P}} \beta^*$.*

Proof. First, we notice that when $\pi(\mathbf{X}; \gamma)$ is correct or $\phi(\mathbf{X}; \alpha) = \psi(\alpha^T \mathbf{G})$ is correct, β^* is the unique solution to $\mathbb{E}\{\tau(\bar{\alpha}, \beta, \bar{\gamma})\} = 0$. If $\pi(\mathbf{X}; \gamma)$ is correct, we obtain,

$$\mathbb{E}\{\tau(\bar{\alpha}, \beta^*, \bar{\gamma})\}$$

$$\begin{aligned}
&= \mathbb{E} \left[\frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\beta^{*\text{T}} \mathbf{Z})\} \mathbf{Z} + \left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \{\psi(\bar{\alpha}^{\text{T}} \mathbf{G}) - \psi(\beta^{*\text{T}} \mathbf{Z})\} \mathbf{Z} \right] \\
&= \mathbb{E}_{\mathbf{X}} \left(\mathbb{E} \left[\frac{R\{Y - \psi(\beta^{*\text{T}} \mathbf{Z})\}}{\pi(\mathbf{X}; \bar{\gamma})} + \left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \{\psi(\bar{\alpha}^{\text{T}} \mathbf{G}) - \psi(\beta^{*\text{T}} \mathbf{Z})\} \right] \mathbf{Z} \middle| \mathbf{X} \right) \\
&= \mathbb{E} [\{Y - \psi(\beta^{*\text{T}} \mathbf{Z})\} \mathbf{Z}] \\
&= 0.
\end{aligned}$$

If $\psi(\alpha^{\text{T}} \mathbf{G})$ is correct, we obtain,

$$\begin{aligned}
&\mathbb{E}\{\tau(\bar{\alpha}, \beta^*, \bar{\gamma})\} \\
&= \mathbb{E} \left[\frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\beta^{*\text{T}} \mathbf{Z})\} \mathbf{Z} + \left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \{\psi(\bar{\alpha}^{\text{T}} \mathbf{G}) - \psi(\beta^{*\text{T}} \mathbf{Z})\} \mathbf{Z} \right] \\
&= \mathbb{E} [\{\psi(\bar{\alpha}^{\text{T}} \mathbf{G}) - \psi(\beta^{*\text{T}} \mathbf{Z})\} \mathbf{Z}] \\
&= \mathbb{E} [\{Y - \psi(\beta^{*\text{T}} \mathbf{Z})\} \mathbf{Z}] \\
&= 0.
\end{aligned}$$

The uniqueness is determined by the uniqueness of β^* .

Since Assumptions 5(ii) and 5(iii) hold, by standard argument of consistency (e.g. Van der Vaart (2000)), it suffices to show that

$$\tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \hat{\beta}, \bar{\gamma})\} = o_p(1). \quad (\text{S24})$$

We consider the following decomposition,

$$\begin{aligned}
\tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \hat{\beta}, \bar{\gamma})\} &= \tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \hat{\beta}, \bar{\gamma})\} - \tilde{\mathbb{E}}\{\tau(\hat{\alpha}, \hat{\beta}, \hat{\gamma})\} \\
&= \tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \hat{\beta}, \hat{\gamma})\} - \tilde{\mathbb{E}}\{\tau(\hat{\alpha}, \hat{\beta}, \hat{\gamma})\} \\
&\quad + \tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \hat{\beta}, \bar{\gamma})\} - \tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \hat{\beta}, \hat{\gamma})\} \\
&= \Delta_0^1 + \Delta_1^1,
\end{aligned}$$

where

$$\begin{aligned}
\Delta_0^1 &= \tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \hat{\beta}, \hat{\gamma})\} - \tilde{\mathbb{E}}\{\tau(\hat{\alpha}, \hat{\beta}, \hat{\gamma})\} \\
&= \tilde{\mathbb{E}} \left[\left\{ \frac{R}{\pi(\mathbf{X}; \hat{\gamma})} - 1 \right\} \{\psi(\hat{\alpha}^{\text{T}} \mathbf{G}) - \psi(\bar{\alpha}^{\text{T}} \mathbf{G})\} \mathbf{Z} \right], \\
\Delta_1^1 &= \tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \hat{\beta}, \bar{\gamma})\} - \tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \hat{\beta}, \hat{\gamma})\} \\
&= \tilde{\mathbb{E}} \left[R \left\{ \frac{1}{\pi(\mathbf{X}; \bar{\gamma})} - \frac{1}{\pi(\mathbf{X}; \hat{\gamma})} \right\} \{Y - \psi(\bar{\alpha}^{\text{T}} \mathbf{G})\} \mathbf{Z} \right].
\end{aligned}$$

By Assumptions 3(iv) and 4(vi), we know that $\exists \zeta_{10} < 0$ a constant, such that $|S_{\bar{\gamma}}|\lambda_0 + |S_{\bar{\alpha}}|\lambda_1 \leq \zeta_{10}$. By (17) in Proposition 2, we know that $\exists M_0^0 > 0$ a constant, such that in the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03} \cap \Omega_{10} \cap \Omega_{11}$, $\|\hat{\alpha} - \bar{\alpha}\|_1 \leq M_0^0(|S_{\bar{\gamma}}|\lambda_0 + |S_{\bar{\alpha}}|\lambda_1)$. Consider the j -th coordinate of Δ_0^1 ,

$$\begin{aligned}
|\Delta_{0,j}^1| &= \left| \tilde{\mathbb{E}} \left[\left\{ \frac{R}{\pi(\mathbf{X}; \hat{\gamma})} - 1 \right\} \{\psi(\hat{\alpha}^T \mathbf{G}) - \psi(\bar{\alpha}^T \mathbf{G})\} Z_j \right] \right| \\
&\leq \tilde{\mathbb{E}}^{1/2} \left[\left\{ \frac{R}{\pi(\mathbf{X}; \hat{\gamma})} - 1 \right\}^2 Z_j^2 \right] \times \tilde{\mathbb{E}}^{1/2} \{\psi(\hat{\alpha}^T \mathbf{G}) - \psi(\bar{\alpha}^T \mathbf{G})\}^2 \\
&\leq \tilde{\mathbb{E}}^{1/2} \left([R - 1 + R w(\mathbf{X}; \bar{\gamma}) \exp\{-(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}]^2 Z_j^2 \right) \\
&\quad \times \sqrt{2} \tilde{\mathbb{E}}^{1/2} [\{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\bar{\alpha}^T \mathbf{G})\} \psi_1(\bar{\alpha}^T \mathbf{G}) (\hat{\alpha} - \bar{\alpha})^T \mathbf{G}] \\
&\leq \sqrt{2} C_5 \tilde{\mathbb{E}}^{1/2} [(R - 1)^2 + R w^2(\mathbf{X}; \bar{\gamma}) \exp\{-2(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}] \\
&\quad \times \tilde{\mathbb{E}}^{1/2} \left[\int_0^1 \psi_1\{\bar{\alpha}^T \mathbf{G} + u(\bar{\alpha} - \bar{\alpha})^T \mathbf{G}\} du \psi_1(\bar{\alpha}^T \mathbf{G}) \{(\hat{\alpha} - \bar{\alpha})^T \mathbf{G}\}^2 \right] \\
&\leq \sqrt{2} C_5 \{1 + \exp(-2B_0 + 2C_0 \|\hat{\gamma} - \bar{\gamma}\|_1)\}^{\frac{1}{2}} \\
&\quad \times \tilde{\mathbb{E}}^{1/2} [\psi_1^2(\bar{\alpha}^T \mathbf{G}) \exp(2C_3 |\bar{\alpha}^T \mathbf{G} - \bar{\alpha}^T \mathbf{G}|) \{(\hat{\alpha} - \bar{\alpha})^T \mathbf{G}\}^2] \\
&\leq 2C_5 \{1 + \exp(-2B_0 + 2\eta_{01})\}^{\frac{1}{2}} \times C_2 C_4 \exp(C_3 C_4 \|\hat{\alpha} - \bar{\alpha}\|_1) \|\hat{\alpha} - \bar{\alpha}\|_1 \\
&\leq \sqrt{2} C_2 C_4 C_5 \{1 + \exp(-2B_0 + 2\eta_{01})\}^{\frac{1}{2}} \times \exp(C_3 C_4 M_0^0 \zeta_{10}) \|\hat{\alpha} - \bar{\alpha}\|_1 \\
&= M_0^1 (|S_{\bar{\gamma}}|\lambda_0 + |S_{\bar{\alpha}}|\lambda_1) \\
&= o_p(1)
\end{aligned}$$

where M_0^1 is a constant, $\tilde{\alpha} = u\hat{\alpha} + (1-u)\bar{\alpha}$ for some constant $u \in (0, 1)$ and the last equality holds by Lemma S19.

We consider the j -th coordinate of Δ_1^1 . By Cauchy-Schwarz inequality, Lemma S8, and Lemma S5, since Assumption 4(iv) holds, we obtain

$$\begin{aligned}
|\Delta_{1,j}^1| &= \left| \tilde{\mathbb{E}} \left[R \left\{ \frac{1}{\pi(\mathbf{X}; \bar{\gamma})} - \frac{1}{\pi(\mathbf{X}; \hat{\gamma})} \right\} \{Y - \psi(\bar{\alpha}^T \mathbf{G})\} Z_j \right] \right| \\
&= \left| \tilde{\mathbb{E}} [R w(\mathbf{X}; \bar{\gamma}) \exp\{-u(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\} (\bar{\gamma} - \hat{\gamma})^T \mathbf{F} \{Y - \psi(\bar{\alpha}^T \mathbf{G})\} Z_j] \right| \\
&\leq \tilde{\mathbb{E}}^{1/2} [R w(\mathbf{X}; \bar{\gamma}) \exp\{-2u(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\} \{(\bar{\gamma} - \hat{\gamma})^T \mathbf{F}\}^2 Z_j^2] \\
&\quad \times \tilde{\mathbb{E}}^{1/2} [R w(\mathbf{X}; \bar{\gamma}) \{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2] \\
&\leq \exp(-B_0/2 + C_0 \|\hat{\gamma} - \bar{\gamma}\|_1) C_0 C_5 \|\hat{\gamma} - \bar{\gamma}\|_1 \\
&\quad \times \sqrt{\mathbb{E}[R w(\mathbf{X}; \bar{\gamma}) \{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2] + (D_0^2 + D_0 D_1) \lambda_0} \\
&\leq \exp(-B_0/2 + \eta_{01}) C_0 C_5 \sqrt{(D_0^2 + D_1^2) \{\lambda_0 + \exp(-B_0)\}} \|\hat{\gamma} - \bar{\gamma}\|_1 \\
&\leq M_1^0 (|S_{\bar{\gamma}}|\lambda_0 + |S_{\bar{\alpha}}|\lambda_1)
\end{aligned}$$

$$=o_p(1),$$

where $u \in (0, 1)$ and M_1^0 are both constants. Therefore, $\|\tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \hat{\beta}, \bar{\gamma})\}\|_\infty \leq \|\Delta_0^1\|_\infty + \|\Delta_1^1\|_\infty = o_p(1)$. Hence, $\hat{\beta} \xrightarrow{\mathbb{P}} \beta^*$. \square

IV.2 Proof of Theorem 1(i)

We show the asymptotic normality of $\sqrt{N}(\hat{\beta} - \beta^*)$. First, we consider the following decomposition,

$$\begin{aligned} & \tau(\hat{\alpha}, \hat{\beta}, \hat{\gamma}) - \tau(\bar{\alpha}, \beta^*, \bar{\gamma}) \\ &= \left[\frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\hat{\beta}^\top \mathbf{Z})\} - \left\{ \frac{R}{\pi(\mathbf{X}; \hat{\gamma})} - 1 \right\} \{ \psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\hat{\beta}^\top \mathbf{Z}) \} \right] \mathbf{Z} \\ & \quad - \left[\frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\beta^{*\top} \mathbf{Z})\} - \left\{ \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} - 1 \right\} \{ \psi(\bar{\alpha}^\top \mathbf{G}) - \psi(\beta^{*\top} \mathbf{Z}) \} \right] \mathbf{Z} \\ &= \{ \psi(\beta^{*\top} \mathbf{Z}) - \psi(\hat{\beta}^\top \mathbf{Z}) \} \mathbf{Z} + R \left\{ \frac{1}{\pi(\mathbf{X}; \hat{\gamma})} - \frac{1}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} \mathbf{Z} \\ & \quad + \left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \{ \psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\bar{\alpha}^\top \mathbf{G}) \} \mathbf{Z} \\ & \quad + \left\{ \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} - \frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \right\} \{ \psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\bar{\alpha}^\top \mathbf{G}) \} \mathbf{Z}, \end{aligned}$$

denoted as $\delta_0^0 + \delta_1^0 + \delta_2^0 + \delta_3^0$. Then, $-\tilde{\mathbb{E}}\tau(\bar{\alpha}, \beta^*, \bar{\gamma}) = \Delta_0^2 + \Delta_1^2 + \Delta_2^2 + \Delta_3^2$, with $\Delta_i^2 = \tilde{\mathbb{E}}(\delta_i^0)$, $i = 0, 1, 2, 3$. First, we show that $\Delta_1^2 + \Delta_2^2 + \Delta_3^2 = o_p(1/\sqrt{N})$. To upper-bound Δ_1^2 , consider $\Delta_{1,j}^2$, the j -th coordinate of Δ_1^2 . By Taylor expansion in a neighborhood of $\bar{\gamma}$,

$$\begin{aligned} \Delta_{1,j}^2 &= \tilde{\mathbb{E}} \left[\left\{ \frac{R}{\pi(\mathbf{X}; \hat{\gamma})} - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} Z_j \right] \\ &= -(\hat{\gamma} - \bar{\gamma})^\top \tilde{\mathbb{E}} [\mathbf{F} R w(\mathbf{X}; \bar{\gamma}) \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} Z_j] \\ & \quad + \frac{1}{2} (\hat{\gamma} - \bar{\gamma})^\top \tilde{\mathbb{E}} [\mathbf{F} R w(\mathbf{X}; \tilde{\gamma}_j) \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} Z_j \mathbf{F}^\top] (\hat{\gamma} - \bar{\gamma}), \end{aligned}$$

denoted as $\Delta_{10,j}^2 + \Delta_{11,j}^2$ where $\tilde{\gamma}_j = u_j \hat{\gamma} + (1 - u_j) \bar{\gamma}$ for some $u_j \in (0, 1)$.

In the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{10}$, by (16) and Lemma S10, we obtain

$$|\Delta_{10,j}^2| \leq \|\hat{\gamma} - \bar{\gamma}\|_1 \|\tilde{\mathbb{E}} [\mathbf{F} R w(\mathbf{X}; \bar{\gamma}) \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} Z_j]\|_\infty \leq M_{10,j}^1 S_{\bar{\gamma}} |\lambda_0 \lambda_1|,$$

for some constant $M_{10,j}^1 > 0$. In the event $\Omega_{00} \cap \Omega_{01}$, by (S4), we obtain

$$\begin{aligned} |\Delta_{11,j}^2| &= \frac{1}{2} |(\hat{\gamma} - \bar{\gamma})^\top \tilde{\mathbb{E}} [\mathbf{F} R w(\mathbf{X}; \tilde{\gamma}_j) \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} Z_j \mathbf{F}^\top] (\hat{\gamma} - \bar{\gamma})| \\ &\leq \frac{1}{2} C_5 \exp(\|\hat{\gamma} - \bar{\gamma}\|_1 C_0) \tilde{\mathbb{E}} \{ R w(\mathbf{X}; \bar{\gamma}) |Y - \psi(\bar{\alpha}^\top \mathbf{G})| |(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}|^2 \}. \end{aligned}$$

We bound $\tilde{\mathbb{E}} \{Rw(\mathbf{X}; \bar{\gamma})|Y - \psi(\bar{\alpha}^T \mathbf{G})|(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}|^2\}$ by following steps. First, by Lemma S7, in the event Ω_{02} ,

$$(\tilde{\mathbb{E}} - \mathbb{E}) [Rw(\mathbf{X}; \bar{\gamma})|Y - \psi(\bar{\alpha}^T \mathbf{G})|(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}|^2] \leq \sqrt{D_0^2 + D_1^2} \|\hat{\gamma} - \bar{\gamma}\|_1^2 \lambda_0.$$

Second, by Assumption 4(iv) and Lemma S5, we have

$$\mathbb{E}[\{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2 | \mathbf{X}] \leq D_0^2 + D_1^2.$$

Therefore,

$$\begin{aligned} & \mathbb{E} [Rw(\mathbf{X}; \bar{\gamma})|Y - \psi(\bar{\alpha}^T \mathbf{G})|(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}|^2] \\ & \leq \mathbb{E}^{1/2} [Rw(\mathbf{X}; \bar{\gamma})|Y - \psi(\bar{\alpha}^T \mathbf{G})|^2 \{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2] \times \mathbb{E}^{1/2} [Rw(\mathbf{X}; \bar{\gamma}) \{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2] \\ & \leq \sqrt{D_0^2 + D_1^2} \mathbb{E} [Rw(\mathbf{X}; \bar{\gamma}) \{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2]. \end{aligned}$$

Third, in the event Ω_{01} , by (S1),

$$(\mathbb{E} - \tilde{\mathbb{E}}) [Rw(\mathbf{X}; \bar{\gamma}) \{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2] \leq \lambda_0 \|\bar{\gamma} - \hat{\gamma}\|_1^2.$$

Combining preceding inequalities and (S4) in LemmaS9, in the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03} \cap \Omega_{02}$,

$$\begin{aligned} & \tilde{\mathbb{E}} \{Rw(\mathbf{X}; \bar{\gamma})|Y - \psi(\bar{\alpha}^T \mathbf{G})|(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}|^2\} \\ & \leq \sqrt{D_0^2 + D_1^2} \|\hat{\gamma} - \bar{\gamma}\|_1^2 \lambda_0 + \sqrt{D_0^2 + D_1^2} \{\lambda_0 \|\bar{\gamma} - \hat{\gamma}\|_1^2 + \exp(\eta_{01}) M_0 |S_{\bar{\gamma}}| \lambda_0^2\} \\ & \leq 2\sqrt{D_0^2 + D_1^2} \left(\frac{M_0}{A_0 - 1}\right)^2 |S_{\bar{\gamma}}|^2 \lambda_0^3 + \sqrt{D_0^2 + D_1^2} \exp(\eta_{01}) M_0 |S_{\bar{\gamma}}| \lambda_0^2 \\ & \leq 2\zeta_0 \sqrt{D_0^2 + D_1^2} \left(\frac{M_0}{A_0 - 1}\right)^2 |S_{\bar{\gamma}}| \lambda_0^2 + \sqrt{D_0^2 + D_1^2} \exp(\eta_{01}) M_0 |S_{\bar{\gamma}}| \lambda_0^2. \end{aligned}$$

Therefore,

$$\begin{aligned} |\Delta_{11,j}^2| & \leq \frac{1}{2} C_5 \exp(\|\hat{\gamma} - \bar{\gamma}\|_1 C_0) \tilde{\mathbb{E}} \{Rw(\mathbf{X}; \bar{\gamma})|Y - \psi(\bar{\alpha}^T \mathbf{G})|(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}|^2\} \\ & \leq \frac{1}{2} C_5 \exp(\eta_{01}) \\ & \quad \times \left\{ 2\zeta_0 \sqrt{D_0^2 + D_1^2} \left(\frac{M_0}{A_0 - 1}\right)^2 |S_{\bar{\gamma}}| \lambda_0^2 + \sqrt{D_0^2 + D_1^2} \exp(\eta_{01}) M_0 |S_{\bar{\gamma}}| \lambda_0^2 \right\} \\ & = M_{11}^1 |S_{\bar{\gamma}}| \lambda_0^2, \end{aligned}$$

for some constant $M_{11}^1 > 0$. Hence, in the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03} \cap \Omega_{02} \cap \Omega_{10}$,

$$\begin{aligned}
& \sup_{j=0, \dots, m-1} |\Delta_{1,j}^2| \\
& \leq \sup_{j=0, \dots, m-1} (|\Delta_{10,j}^2| + |\Delta_{11,j}^2|) \\
& \leq \sup_{j=0, \dots, m-1} M_{10}^1 |S_{\bar{\gamma}}| \lambda_0 \lambda_1 + \sup_{j=0, \dots, m-1} M_{11}^1 |S_{\bar{\gamma}}| \lambda_0^2 \\
& = M_{10}^1 |S_{\bar{\gamma}}| \lambda_0 \lambda_1 + M_{11}^1 |S_{\bar{\gamma}}| \lambda_0^2 \\
& \leq M_1^1 (|S_{\bar{\gamma}}| \lambda_0 \lambda_1 + |S_{\bar{\gamma}}| \lambda_0^2),
\end{aligned} \tag{S25}$$

where $M_1^1 = \max(M_{10}^1, M_{11}^1)$.

To bound Δ_2^2 , consider $\Delta_{2,j}^2$, the j -th coordinate of Δ_2^2 , $\Delta_{2,j}^2$ can be decomposed as

$$\begin{aligned}
\Delta_{2,j}^2 = & (\tilde{\mathbb{E}} - \mathbb{E}) \left[\left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \{ \psi(\hat{\alpha}^T \mathbf{G}) - \psi(\bar{\alpha}^T \mathbf{G}) \} Z_j \right] \\
& + \mathbb{E} \left[\left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \{ \psi(\hat{\alpha}^T \mathbf{G}) - \psi(\bar{\alpha}^T \mathbf{G}) \} Z_j \right],
\end{aligned}$$

denoted as $\Delta_{20,j}^2 + \Delta_{21,j}^2$. Since $\pi(\mathbf{X}; \bar{\gamma})$ is correct, $\Delta_{21,j}^2 = 0$. By (17), $\exists M_\alpha > 0$, such that $\|\hat{\alpha} - \bar{\alpha}\|_1 \leq M_\alpha (|S_{\bar{\gamma}}| \lambda_0 + |S_{\bar{\alpha}}| \lambda_1)^{13}$. Take $r_\alpha = M_\alpha (|S_{\bar{\gamma}}| \lambda_0 + |S_{\bar{\alpha}}| \lambda_1)$ in Lemma S18, then in the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03} \cap \Omega_{10} \cap \Omega_{11} \cap \Omega_{20}$, we have $\|\hat{\alpha} - \bar{\alpha}\|_1 \leq r_\alpha$ and hence

$$|\Delta_{20,j}^2| \leq B_1 M_\alpha (|S_{\bar{\gamma}}| \lambda_0 + |S_{\bar{\alpha}}| \lambda_1) \lambda_0. \tag{S26}$$

Thus, in the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03} \cap \Omega_{10} \cap \Omega_{11} \cap \Omega_{20}$,

$$\sup_{j=0, \dots, m-1} |\Delta_{2,j}^2| \leq B_1 M_\alpha (|S_{\bar{\gamma}}| \lambda_0 + |S_{\bar{\alpha}}| \lambda_1) \lambda_0 = M_2^1 (|S_{\bar{\gamma}}| \lambda_0 + |S_{\bar{\alpha}}| \lambda_1) \lambda_0, \tag{S27}$$

for some positive constant M_2^1 .

To deal with Δ_3^2 , first, by mean value theorem, we obtain for some $u \in (0, 1)$,

$$\begin{aligned}
\frac{1}{\pi(\mathbf{X}; \hat{\gamma})} - \frac{1}{\pi(\mathbf{X}; \bar{\gamma})} & = - \exp\{-u \hat{\gamma}^T \mathbf{F} - (1-u) \bar{\gamma}^T \mathbf{F}\} (\hat{\gamma} - \bar{\gamma})^T \mathbf{F} \\
& = - w(\mathbf{X}; \bar{\gamma}) \exp\{-u(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\} (\hat{\gamma} - \bar{\gamma})^T \mathbf{F}
\end{aligned} \tag{S28}$$

and for some $\tilde{\alpha}$ lies between $\bar{\alpha}$ and $\hat{\alpha}$,

$$\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\hat{\alpha}^T \mathbf{G}) = -\psi_1(\tilde{\alpha}^T \mathbf{G}) (\hat{\alpha} - \bar{\alpha})^T \mathbf{G}. \tag{S29}$$

¹³Note that $\lambda_0 \leq \lambda_1$.

Combining (S28) and (S29) and applying Cauchy-Schwartz inequality to j -th coordinate of Δ_3^2 in the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03} \cap \Omega_{10} \cap \Omega_{11} \cap \Omega_{20}$, we get

$$\begin{aligned}
|\Delta_{3,j}^2| &= \left| \tilde{\mathbb{E}} \left[\left\{ \frac{R}{\pi(\mathbf{X}; \hat{\gamma})} - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \{ \psi(\bar{\alpha}^T \mathbf{G}) - \psi(\hat{\alpha}^T \mathbf{G}) \} Z_j \right] \right| \\
&= |\tilde{\mathbb{E}}([Rw(\mathbf{X}; \bar{\gamma}) \exp\{-u_j(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}] \{ \psi_1(\tilde{\alpha}_j^T \mathbf{G})(\hat{\alpha} - \bar{\alpha})^T \mathbf{G} \}) Z_j| \\
&\leq C_5 \exp(\eta_{01}) \tilde{\mathbb{E}}^{1/2} |Rw(\mathbf{X}; \bar{\gamma}) \{(\hat{\gamma} - \bar{\gamma})^T \mathbf{F}\}^2| \\
&\quad \times \tilde{\mathbb{E}}^{1/2} |Rw(\mathbf{X}; \bar{\gamma}) \psi_1^2(\tilde{\alpha}_j^T \mathbf{G}) \{(\hat{\alpha} - \bar{\alpha})^T \mathbf{G}\}^2| \\
&\leq C_5 \exp(\eta_{01}) \{ \exp(\eta_{01}) M_0 |S_{\bar{\gamma}}| \lambda_0^2 \}^{1/2} \\
&\quad \times \tilde{\mathbb{E}}^{1/2} (Rw(\mathbf{X}; \bar{\gamma}) [\psi_1(\tilde{\alpha}_j^T \mathbf{G}) \exp\{C_3 |(\hat{\alpha} - \bar{\alpha})^T \mathbf{G}|\}]^2 \{(\hat{\alpha} - \bar{\alpha})^T \mathbf{G}\}^2) \\
&\leq C_5 \exp(\eta_{01}) \{ \exp(\eta_{01}) M_0 |S_{\bar{\gamma}}| \lambda_0^2 \}^{1/2} C_2^{\frac{1}{2}} \exp(C_3 C_4 r_\alpha) \left\{ \frac{M^\dagger (|S_{\bar{\gamma}}| \lambda_0^2 + |S_{\bar{\alpha}}| \lambda_1^2)}{1 - \eta_3} \right\}^{\frac{1}{2}} \\
&\leq M_3^1 \{ |S_{\bar{\gamma}}| \lambda_0^2 + (|S_{\bar{\gamma}}| |S_{\bar{\alpha}}|)^{1/2} \lambda_0 \lambda_1 \}.
\end{aligned} \tag{S30}$$

The second inequality holds due to (S4) in Lemma S9. The third inequality holds by Assumption 4(ii), (S20), the facts that $\tilde{\mathbb{E}} [Rw(\mathbf{X}; \bar{\gamma}) \psi_1(\tilde{\alpha}_j^T \mathbf{G}) \{(\hat{\alpha} - \bar{\alpha})^T \mathbf{G}\}^2] = b^T \tilde{\Sigma}_\alpha b$ in (S18) and that by (17) in Proposition 2, $\exists M^\dagger > 0$ a constant, such that $D_{\text{WL}}^\dagger \leq M^\dagger (|S_{\bar{\gamma}}| \lambda_0^2 + |S_{\bar{\alpha}}| \lambda_1^2)$. Therefore,

$$\sup_{j=0, \dots, m-1} |\Delta_{3,j}^2| \leq M_3^1 \{ |S_{\bar{\gamma}}| \lambda_0^2 + (|S_{\bar{\gamma}}| |S_{\bar{\alpha}}|)^{1/2} \lambda_0 \lambda_1 \}, \tag{S31}$$

for some constant M_3^1 . Thus, on the event $\Omega_{00} \cap \Omega_{01} \cap \Omega_{03} \cap \Omega_{02} \cap \Omega_{10} \cap \Omega_{11} \cap \Omega_{20}$,

$$\begin{aligned}
\|\Delta_1^2 + \Delta_2^2 + \Delta_3^2\|_\infty &\leq \sup_{j=0, \dots, m-1} |\Delta_{1,j}^2| + \sup_{j=0, \dots, m-1} |\Delta_{2,j}^2| + \sup_{j=0, \dots, m-1} |\Delta_{3,j}^2| \\
&\leq M_1^1 (|S_{\bar{\gamma}}| \lambda_0 \lambda_1 + |S_{\bar{\gamma}}| \lambda_0^2) + M_2^1 (|S_{\bar{\gamma}}| \lambda_0 + |S_{\bar{\alpha}}| \lambda_1) \lambda_0 \\
&\quad + M_3^1 \{ |S_{\bar{\gamma}}| \lambda_0^2 + (|S_{\bar{\gamma}}| |S_{\bar{\alpha}}|)^{1/2} \lambda_0 \lambda_1 \} \\
&\leq M^1 (|S_{\bar{\gamma}}| + |S_{\bar{\alpha}}|) \lambda_0 \lambda_1.
\end{aligned}$$

By Lemma S20, $\|\Delta_1^2 + \Delta_2^2 + \Delta_3^2\|_\infty = o_p(1/\sqrt{N})$.

Then, we deal with Δ_0^2 . For the j -th coordinate of Δ_0^2 , $\Delta_{0,j}^2$, by mean value theorem,

$$\Delta_{0,j}^2 = \tilde{\mathbb{E}}[\{ \psi(\beta^{*\text{T}} \mathbf{Z}) - \psi(\hat{\beta}^T \mathbf{Z}) \} Z_j] = -\tilde{\mathbb{E}}\{ \psi_1(\tilde{\beta}_j^T \mathbf{Z}) Z_j \mathbf{Z}^T (\hat{\beta} - \beta^*) \},$$

where $\tilde{\beta}_j = (1 - u_j) \beta^* + u_j \hat{\beta}$ for some $u_j \in (0, 1)$. We first show that $\tilde{\mathbb{E}}\{ \psi_1(\beta^{*\text{T}} \mathbf{Z}) Z_j \mathbf{Z} \} - \tilde{\mathbb{E}}\{ \psi_1(\tilde{\beta}_j^T \mathbf{Z}) Z_j \mathbf{Z} \} \xrightarrow{\mathbb{P}} 0$. By Assumption 4(ii), we know that for $\forall u, u'$, if $\psi_1(u) > \psi_1(u')$, since $\psi_1(u') \geq \psi_1(u) \exp(-C_3 |u - u'|)$, then $|\psi_1(u) - \psi_1(u')| \leq \psi_1(u) \{1 - \exp(-C_3 |u - u'|)\}$; if $\psi_1(u) <$

$\psi_1(u')$, $\psi_1(u') \leq \psi_1(u) \exp(C_3|u - u'|)$, $|\psi_1(u) - \psi_1(u')| \leq \psi_1(u)\{\exp(C_3|u - u'|) - 1\}$; therefore, $|\psi_1(u) - \psi_1(u')| \leq \psi_1(u) \max\{1 - \exp(-C_3|u - u'|), \exp(C_3|u - u'|) - 1\}$. Consider the i -th element of the difference, if $C_3 = 0$, ψ_1 is a constant, then

$$|\tilde{\mathbb{E}}\{\psi_1(\beta^{*\text{T}}\mathbf{X})Z_iZ_j\} - \tilde{\mathbb{E}}\{\psi_1(\tilde{\beta}_j^{\text{T}}\mathbf{Z})Z_iZ_j\}| = 0.$$

Otherwise,

$$\begin{aligned} & |\tilde{\mathbb{E}}\{\psi_1(\beta^{*\text{T}}\mathbf{X})Z_iZ_j\} - \tilde{\mathbb{E}}\{\psi_1(\tilde{\beta}_j^{\text{T}}\mathbf{Z})Z_iZ_j\}| \leq \tilde{\mathbb{E}}\{|\psi_1(\beta^{*\text{T}}\mathbf{Z}) - \psi_1(\tilde{\beta}_j^{\text{T}}\mathbf{Z})||Z_iZ_j|\} \\ & \leq C_5^2 \tilde{\mathbb{E}}[\psi_1(\beta^{*\text{T}}\mathbf{Z}) \max\{1 - \exp(-C_3|(\hat{\beta} - \beta^*)^{\text{T}}\mathbf{Z}|), \exp(C_3|(\hat{\beta} - \beta^*)^{\text{T}}\mathbf{Z}|) - 1\}] \\ & = C_5^2 C_6 \tilde{\mathbb{E}}\{C_3|(\hat{\beta} - \beta^*)^{\text{T}}\mathbf{Z}| + o_p(|(\hat{\beta} - \beta^*)^{\text{T}}\mathbf{Z}|)\} \\ & \leq C_3 C_5^3 C_6 \|\hat{\beta} - \beta^*\|_1 + o_p(\|\hat{\beta} - \beta^*\|_1) \\ & = o_p(1), \end{aligned}$$

which leads to $-\tilde{\mathbb{E}}\{\psi_1(\tilde{\beta}_j^{\text{T}}\mathbf{Z})Z_j\mathbf{Z}^{\text{T}}\} + \tilde{\mathbb{E}}\{\psi_1(\beta^{*\text{T}}\mathbf{Z})Z_j\mathbf{Z}^{\text{T}}\} \xrightarrow{\mathbb{P}} 0$. Therefore, we consider the following decomposition,

$$\begin{aligned} & -\tilde{\mathbb{E}}\{\psi_1(\tilde{\beta}_j^{\text{T}}\mathbf{Z})Z_j\mathbf{Z}\} + \mathbb{E}\{\psi_1(\beta^{*\text{T}}\mathbf{Z})Z_j\mathbf{Z}\} \\ & = -\tilde{\mathbb{E}}\{\psi_1(\tilde{\beta}_j^{\text{T}}\mathbf{Z})Z_j\mathbf{Z}\} + \tilde{\mathbb{E}}\{\psi_1(\beta^{*\text{T}}\mathbf{X})Z_j\mathbf{Z}\} \\ & \quad - \tilde{\mathbb{E}}\{\psi_1(\beta^{*\text{T}}\mathbf{Z})Z_j\mathbf{Z}\} + E\{\psi_1(\beta^{*\text{T}}\mathbf{Z})Z_j\mathbf{Z}\} \\ & \xrightarrow{\mathbb{P}} 0. \end{aligned}$$

Hence,

$$\tilde{\mathbb{E}}\{\psi_1(\tilde{\beta}_j^{\text{T}}\mathbf{Z})Z_j\mathbf{Z}\} \xrightarrow{\mathbb{P}} \mathbb{E}\{\psi_1(\beta^{*\text{T}}\mathbf{Z})Z_j\mathbf{Z}\} = \mathbf{\Gamma}_j, \quad (\text{S32})$$

where $\mathbf{\Gamma}_j$ is the j -th row of $\mathbf{\Gamma}$, and (S32) holds for $j = 0, \dots, m-1$. Hence, $\Delta_0^2 = \tilde{\mathbb{E}}(\tilde{\mathbf{Z}}\mathbf{Z}^{\text{T}})(\hat{\beta} - \beta^*)$, where $\tilde{\mathbf{Z}}_j = -\psi_1(\tilde{\beta}_j^{\text{T}}\mathbf{Z})Z_j$, and $\tilde{\mathbb{E}}(\tilde{\mathbf{Z}}\mathbf{Z}^{\text{T}}) \xrightarrow{\mathbb{P}} -\mathbf{\Gamma}$. Suppose $\sqrt{N}(\hat{\beta} - \beta^*) \xrightarrow{d} G_2$, by continuous mapping theorem,

$$\sqrt{N}\Delta_0^2 \xrightarrow{d} -\mathbf{\Gamma}G_2. \quad (\text{S33})$$

Besides, by central limit theorem,

$$\sqrt{N}\Delta_0^2 = \sqrt{N}\tilde{\mathbb{E}}\{\tau(\mathbf{O}, \bar{\alpha}, \beta^*, \bar{\gamma})\} + o_p(1) \xrightarrow{d} \text{N}(0, \mathbf{\Lambda}). \quad (\text{S34})$$

Therefore,

$$\sqrt{N}(\hat{\beta} - \beta^*) \xrightarrow{d} G_2 \sim -\mathbf{\Gamma}^{-1}\text{N}(0, \mathbf{\Lambda}) \sim \text{N}(0, \mathbf{\Sigma}),$$

where \sim denotes "distributed as", i.e., for any two distributions G_0 and G_1 , $G_0 \sim G_1$ means the two distributions are the same.

IV.3 Proof of Theorem 1(ii)

We show the consistency of $\hat{\Sigma}$. First, if we let $\hat{\Gamma}_j = \tilde{\mathbb{E}}\{\psi_1(\hat{\beta}^\top \mathbf{Z})Z_j\mathbf{Z}\}$, then

$$\hat{\Gamma}_j \xrightarrow{\mathbb{P}} \mathbb{E}\{\psi_1(\beta^{*\top} \mathbf{Z})Z_j\mathbf{Z}\},$$

i.e., $\hat{\Gamma}_j \xrightarrow{\mathbb{P}} \Gamma_j$, which can be shown in the way similar to the proof of (S32). Then we get $\hat{\Gamma} \xrightarrow{\mathbb{P}} \Gamma$.

Next, we want to show that $\hat{\Lambda} \xrightarrow{\mathbb{P}} \Lambda$. Since $\Lambda = \mathbb{E}\{\tau(\bar{\alpha}, \beta^*, \bar{\gamma})\tau(\bar{\alpha}, \beta^*, \bar{\gamma})^\top\}$ and $\hat{\Lambda} = \tilde{\mathbb{E}}\{\tau(\hat{\alpha}, \hat{\beta}, \hat{\gamma})\tau(\hat{\alpha}, \hat{\beta}, \hat{\gamma})^\top\}$, it suffices to show that

$$\tilde{\mathbb{E}}\{\tau(\hat{\alpha}, \hat{\beta}, \hat{\gamma})\tau(\hat{\alpha}, \hat{\beta}, \hat{\gamma})^\top\} - \tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \beta^*, \bar{\gamma})\tau(\bar{\alpha}, \beta^*, \bar{\gamma})^\top\} = o_p(1).$$

We consider the i, j -th element of the difference above:

$$\begin{aligned} & \left| \tilde{\mathbb{E}}\{\tau(\hat{\alpha}, \hat{\beta}, \hat{\gamma})\tau(\hat{\alpha}, \hat{\beta}, \hat{\gamma})^\top\} - \tilde{\mathbb{E}}\{\tau(\bar{\alpha}, \beta^*, \bar{\gamma})\tau(\bar{\alpha}, \beta^*, \bar{\gamma})^\top\} \right|_{ij} \\ &= \left| \tilde{\mathbb{E}} \left(\left[\frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\hat{\alpha}^\top \mathbf{G})\} + \{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\hat{\beta}^\top \mathbf{Z})\} \right]^2 Z_i Z_j \right. \right. \\ & \quad \left. \left. - \left[\frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} + \{\psi(\bar{\alpha}^\top \mathbf{G}) - \psi(\beta^{*\top} \mathbf{Z})\} \right]^2 Z_i Z_j \right) \right| \\ &\leq \tilde{\mathbb{E}} \left(\left| \left[\frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\hat{\alpha}^\top \mathbf{G})\} + \{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\hat{\beta}^\top \mathbf{Z})\} \right]^2 \right. \right. \\ & \quad \left. \left. - \left[\frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} + \{\psi(\bar{\alpha}^\top \mathbf{G}) - \psi(\beta^{*\top} \mathbf{Z})\} \right]^2 \right| \left| Z_i Z_j \right| \right) \\ &\leq C_5^2 \tilde{\mathbb{E}} \left(\left| \left[\frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\hat{\alpha}^\top \mathbf{G})\} + \{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\hat{\beta}^\top \mathbf{Z})\} \right]^2 \right. \right. \\ & \quad \left. \left. - \left[\frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} + \{\psi(\bar{\alpha}^\top \mathbf{G}) - \psi(\beta^{*\top} \mathbf{Z})\} \right]^2 \right| \right) \\ &\leq C_5^2 \tilde{\mathbb{E}} \left(\left[\frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\hat{\alpha}^\top \mathbf{G})\} + \{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\hat{\beta}^\top \mathbf{Z})\} \right. \right. \\ & \quad \left. \left. - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} - \{\psi(\bar{\alpha}^\top \mathbf{G}) - \psi(\beta^{*\top} \mathbf{Z})\} \right]^2 \right) \\ & \quad + 2C_5^2 \tilde{\mathbb{E}} \left(\left[\frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\hat{\alpha}^\top \mathbf{G})\} + \{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\hat{\beta}^\top \mathbf{Z})\} \right. \right. \\ & \quad \left. \left. - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} - \{\psi(\bar{\alpha}^\top \mathbf{G}) - \psi(\beta^{*\top} \mathbf{Z})\} \right] \right. \\ & \quad \left. \times \left[\frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} + \{\psi(\bar{\alpha}^\top \mathbf{G}) - \psi(\beta^{*\top} \mathbf{Z})\} \right] \right) \\ &\leq C_5^2 \tilde{\mathbb{E}} \left(\left[\frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\hat{\alpha}^\top \mathbf{G})\} + \{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\hat{\beta}^\top \mathbf{Z})\} \right. \right. \\ & \quad \left. \left. - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\} - \{\psi(\bar{\alpha}^\top \mathbf{G}) - \psi(\beta^{*\top} \mathbf{Z})\} \right]^2 \right) \end{aligned}$$

$$\begin{aligned}
& + 2C_5^2 \tilde{\mathbb{E}}^{\frac{1}{2}} \left(\left[\frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\hat{\alpha}^T \mathbf{G})\} + \{\psi(\hat{\alpha}^T \mathbf{G}) - \psi(\hat{\beta}^T \mathbf{Z})\} \right. \right. \\
& \left. \left. - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^T \mathbf{G})\} - \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\} \right]^2 \right) \\
& \times \tilde{\mathbb{E}}^{\frac{1}{2}} \left(\left[\frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^T \mathbf{G})\} + \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\} \right]^2 \right),
\end{aligned}$$

therefore, we only need to show that

$$\begin{aligned}
& \tilde{\mathbb{E}} \left(\left[\frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\hat{\alpha}^T \mathbf{G})\} + \{\psi(\hat{\alpha}^T \mathbf{G}) - \psi(\hat{\beta}^T \mathbf{Z})\} \right. \right. \\
& \left. \left. - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^T \mathbf{G})\} - \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\} \right]^2 \right) = o_p(1).
\end{aligned}$$

Consider the following decomposition:

$$\begin{aligned}
& \frac{R}{\pi(\mathbf{X}; \hat{\gamma})} \{Y - \psi(\hat{\alpha}^T \mathbf{G})\} + \{\psi(\hat{\alpha}^T \mathbf{G}) - \psi(\hat{\beta}^T \mathbf{Z})\} \\
& - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \{Y - \psi(\bar{\alpha}^T \mathbf{G})\} - \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\} \\
& = \{\psi(\hat{\alpha}^T \mathbf{G}) - \psi(\bar{\alpha}^T \mathbf{G})\} \left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \\
& + R \{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \left\{ \frac{1}{\pi(\mathbf{X}; \hat{\gamma})} - \frac{1}{\pi(\mathbf{X}; \bar{\gamma})} \right\} \\
& + R \{\psi(\hat{\alpha}^T \mathbf{G}) - \psi(\bar{\alpha}^T \mathbf{G})\} \left\{ \frac{1}{\pi(\mathbf{X}; \bar{\gamma})} - \frac{1}{\pi(\mathbf{X}; \hat{\gamma})} \right\} \\
& + \psi(\beta^{*T} \mathbf{Z}) - \psi(\hat{\beta}^T \mathbf{Z}),
\end{aligned}$$

denoted as $\delta_0^1 + \delta_1^1 + \delta_2^1 + \delta_3^1$. Let $\Delta_i^3 = \tilde{\mathbb{E}}\{(\delta_i^1)^2\}$, $i = 0, \dots, 3$, we only need to show that $\Delta_i^3 = o_p(1)$, $i = 0, \dots, 3$.

By mean value theorem,

$$\begin{aligned}
\Delta_0^3 & = \tilde{\mathbb{E}} \left[\{\psi(\hat{\alpha}^T \mathbf{G}) - \psi(\bar{\alpha}^T \mathbf{G})\}^2 \left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\}^2 \right] \\
& = 2\tilde{\mathbb{E}} \left[\psi_1(\tilde{\alpha}^T \mathbf{G}) \{\psi(\tilde{\alpha}^T \mathbf{G}) - \psi(\bar{\alpha}^T \mathbf{G})\} (\hat{\alpha} - \bar{\alpha})^T \mathbf{G} \left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\}^2 \right] \\
& = 2\tilde{\mathbb{E}} \left(\psi_1(\tilde{\alpha}^T \mathbf{G}) \left[\int_0^1 \psi_1\{\bar{\alpha}^T \mathbf{G} + u(\tilde{\alpha} - \bar{\alpha})^T \mathbf{G}\} du \right] \right. \\
& \quad \left. \{(\hat{\alpha} - \bar{\alpha})^T \mathbf{G}\}^2 \left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\}^2 \right) \\
& \leq 2\tilde{\mathbb{E}} \left[\psi_1(\bar{\alpha}^T \mathbf{G}) \exp(C_3 C_4 r_\alpha) \left\{ \int_0^1 \psi_1(\bar{\alpha}^T \mathbf{G}) \exp(C_3 C_4 r_\alpha) du \right\} \right. \\
& \quad \left. \{(\hat{\alpha} - \bar{\alpha})^T \mathbf{G}\}^2 \left\{ 1 - \frac{R}{\pi(\mathbf{X}; \bar{\gamma})} \right\}^2 \right]
\end{aligned}$$

$$\begin{aligned}
&\leq 2C_2^2 C_4^2 \exp(2C_3 C_4 r_\alpha) \{1 + \exp(-B_0)\}^2 \|\hat{\alpha} - \bar{\alpha}\|_1^2 \\
&= o_p(1).
\end{aligned}$$

$$\begin{aligned}
\Delta_1^3 &= \tilde{\mathbb{E}} \left[R \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 \left\{ \frac{1}{\pi(\mathbf{X}; \hat{\gamma})} - \frac{1}{\pi(\mathbf{X}; \bar{\gamma})} \right\}^2 \right] \\
&= -2\tilde{\mathbb{E}} \left(R w(\mathbf{X}; \bar{\gamma})^2 [\exp\{-(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}\} - 1] \exp\{-(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}\} \right. \\
&\quad \left. \times \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 (\hat{\gamma} - \bar{\gamma})^\top \mathbf{G} \right) \\
&\leq 2\{1 + \exp(\eta_{01})\}^2 \exp(-B_0 + \eta_{01}) C_4 \|\hat{\gamma} - \bar{\gamma}\|_1 \tilde{\mathbb{E}} \left[R w(\mathbf{X}; \bar{\gamma}) \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 \right] \\
&\leq 2\{1 + \exp(\eta_{01})\}^2 \exp(-B_0 + \eta_{01}) C_4 \|\hat{\gamma} - \bar{\gamma}\|_1 \left(\mathbb{E} \left[R w(\mathbf{X}; \bar{\gamma}) \{Y - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 \right] \right. \\
&\quad \left. + (D_0^2 + D_0 D_1) \lambda_0 \right) \\
&= o_p(1).
\end{aligned}$$

By mean value theorem,

$$\begin{aligned}
\Delta_2^3 &= \tilde{\mathbb{E}} \left[R \{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 \left\{ \frac{1}{\pi(\mathbf{X}; \bar{\gamma})} - \frac{1}{\pi(\mathbf{X}; \hat{\gamma})} \right\}^2 \right] \\
&= \tilde{\mathbb{E}} \left(R \{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 w(\mathbf{X}; \bar{\gamma})^2 [1 - \exp\{-(\hat{\gamma} - \bar{\gamma})^\top \mathbf{F}\}]^2 \right) \\
&\leq \{1 + \exp(\eta_{01})\}^2 \exp(-2B_0) \tilde{\mathbb{E}} \left[\{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\bar{\alpha}^\top \mathbf{G})\}^2 \right] \\
&= 2\{1 + \exp(\eta_{01})\}^2 \exp(-2B_0) \tilde{\mathbb{E}} \left[\{\psi(\hat{\alpha}^\top \mathbf{G}) - \psi(\bar{\alpha}^\top \mathbf{G})\} \psi_1(\hat{\alpha}^\top \mathbf{G}) (\hat{\alpha} - \bar{\alpha})^\top \mathbf{G} \right] \\
&= 2\{1 + \exp(\eta_{01})\}^2 \exp(-2B_0) \\
&\quad \times \tilde{\mathbb{E}} \left(\psi_1(\bar{\alpha}^\top \mathbf{G}) \left[\int_0^1 \psi_1 \{\bar{\alpha}^\top \mathbf{G} + u(\hat{\alpha} - \bar{\alpha})^\top \mathbf{G}\} du \right] \{(\hat{\alpha} - \bar{\alpha})^\top \mathbf{G}\}^2 \right) \\
&\leq 2C_2^2 C_4^2 \{1 + \exp(\eta_{01})\}^2 \exp(2C_3 C_4 r_\alpha - 2B_0) \|\hat{\alpha} - \bar{\alpha}\|_1^2 \\
&= o_p(1).
\end{aligned}$$

$$\begin{aligned}
\Delta_3^3 &= \tilde{\mathbb{E}} [\{\psi(\beta^{*\top} \mathbf{Z}) - \psi(\hat{\beta}^\top \mathbf{Z})\}^2] \\
&= 2\tilde{\mathbb{E}} \left[\{\psi(\hat{\beta}^\top \mathbf{Z}) - \psi(\beta^{*\top} \mathbf{Z})\} \psi_1(\hat{\beta}^\top \mathbf{Z}) (\hat{\beta} - \beta^*)^\top \mathbf{Z} \right] \\
&= 2\tilde{\mathbb{E}} \left(\psi_1(\hat{\beta}^\top \mathbf{Z}) \left[\int_0^1 \psi_1 \{\beta^{*\top} \mathbf{Z} + u(\hat{\beta} - \beta^*)^\top \mathbf{Z}\} du \right] \{(\hat{\beta} - \beta^*)^\top \mathbf{Z}\}^2 \right) \\
&\leq 2C_5^2 C_6^2 \{1 + \exp(\eta_{01})\}^2 \exp(2C_3 C_5 \|\hat{\beta} - \beta^*\|_1) \|\hat{\beta} - \beta^*\|_1^2
\end{aligned}$$

$$=o_p(1).$$

Therefore, $\hat{\Lambda} \xrightarrow{\mathbb{P}} \Lambda$. Then, by continuous mapping theorem, $\hat{\Gamma}^{-1} \xrightarrow{\mathbb{P}} \Gamma^{-1}$. Thus, by continuous mapping theorem again, $\hat{\Sigma} \xrightarrow{\mathbb{P}} \Sigma$.

V Extension to the setting of stratified sampling

V.1 Proof of Proposition 3

Let

$$\tau^s(\alpha, \beta, \cdot) = \frac{1}{N} \sum_{i=1}^n \frac{1}{(\cdot)} \{Y_i - \psi(\alpha^T \mathbf{G}_i)\} \mathbf{Z}_i + \frac{1}{N} \sum_{i=1}^N \{\psi(\alpha^T \mathbf{G}_i) - \psi(\beta^T \mathbf{Z}_i)\} \mathbf{Z}_i. \quad (\text{S35})$$

We have

$$\begin{aligned} & -\tau^s\{\bar{\alpha}, \beta^*, \pi(\mathbf{X}; \bar{\gamma})\} = -\tau^s\{\bar{\alpha}, \beta^*, \pi^*(\mathbf{X})\} = \tau^s\{\hat{\alpha}, \hat{\beta}^s, \hat{\pi}(\mathbf{X})\} - \tau^s\{\bar{\alpha}, \beta^*, \pi^*(\mathbf{X})\} \\ &= \frac{1}{N} \sum_{i=1}^N [\{\psi(\hat{\alpha}^T \mathbf{G}_i) - \psi(\hat{\beta}^{sT} \mathbf{Z}_i)\} \mathbf{Z}_i] - \frac{1}{N} \sum_{i=1}^N [\{\psi(\hat{\alpha}^T \mathbf{G}_i) - \psi(\beta^{*T} \mathbf{Z}_i)\} \mathbf{Z}_i] \\ &+ \frac{1}{N} \sum_{i=1}^N [\{\psi(\hat{\alpha}^T \mathbf{G}_i) - \psi(\beta^{*T} \mathbf{Z}_i)\} \mathbf{Z}_i] - \frac{1}{N} \sum_{i=1}^N [\{\psi(\bar{\alpha}^T \mathbf{G}_i) - \psi(\beta^{*T} \mathbf{Z}_i)\} \mathbf{Z}_i] \\ &+ \frac{1}{n} \sum_{i=1}^n \{Y_i - \psi(\hat{\alpha}^T \mathbf{G}_i)\} \mathbf{Z}_i - \frac{1}{n} \sum_{i=1}^n \{Y_i - \psi(\bar{\alpha}^T \mathbf{G}_i)\} \mathbf{Z}_i, \end{aligned}$$

denoted as $\Delta_5^0 + \Delta_5^1 + \Delta_5^2$.

We first deal with Δ_5^0 . Let Z_{ij} denote the j -th co-ordinate of the i -th sample \mathbf{Z}_i . Then, we consider the j -th coordinate of Δ_5^0 , $\Delta_{5,j}^0$

$$\begin{aligned} \Delta_{5,j}^0 &= \frac{1}{N} \sum_{i=1}^N [\{\psi(\hat{\alpha}^T \mathbf{G}_i) - \psi(\hat{\beta}^{sT} \mathbf{Z}_i)\} Z_{ij}] - \frac{1}{N} \sum_{i=1}^N [\{\psi(\hat{\alpha}^T \mathbf{G}_i) - \psi(\beta^{*T} \mathbf{Z}_i)\} Z_{ij}] \\ &= \frac{1}{N} \sum_{i=1}^N [\{\psi(\beta^{*T} \mathbf{Z}_i) - \psi(\hat{\beta}^{sT} \mathbf{Z}_i)\} Z_{ij}] \\ &= \frac{1}{N} \sum_{i=1}^N \{\psi_1(\tilde{\beta}_j^T \mathbf{Z}_i) Z_{ij} (\beta^* - \hat{\beta}^s)^T \mathbf{Z}_i\} \\ &= \frac{1}{N} \sum_{i=1}^N \{\psi_1(\beta^{*T} \mathbf{Z}_i) Z_{ij} (\beta^* - \hat{\beta}^s)^T \mathbf{Z}_i\} + O_p(\|\beta^* - \hat{\beta}^s\|_1^2), \end{aligned}$$

where $\tilde{\beta}_j = u_j \beta^* + (1 - u_j) \hat{\beta}^s$ for some $u_j \in (0, 1)$. The last equality holds since $\psi_1(\beta^{*T} \mathbf{Z}_i) \exp(-C_3 C_5 \|\beta^* - \hat{\beta}^s\|_1) \leq \psi_1(\tilde{\beta}_j^T \mathbf{Z}_i) \leq \psi_1(\beta^{*T} \mathbf{Z}_i) \exp(C_3 C_5 \|\beta^* - \hat{\beta}^s\|_1)$, which leads to $\psi_1(\tilde{\beta}_j^T \mathbf{Z}_i)$

$$= \psi_1(\beta^{*\top} \mathbf{Z}_i) + O_p(\|\beta^* - \hat{\beta}^s\|_1).$$

Then we deal with $\Delta_{5,j}^1$ and $\Delta_{5,j}^2$ together. Consider the j -th coordinate of $\Delta_{5,j}^1 + \Delta_{5,j}^2$:

$$\begin{aligned} & |\Delta_{5,j}^1 + \Delta_{5,j}^2| \\ &= \left| \frac{1}{n} \sum_{i=1}^n [\{\psi(\bar{\alpha}^\top \mathbf{G}_i) - \psi(\hat{\alpha}^\top \mathbf{G}_i)\} Z_{ij}] \right. \\ &\quad \left. + \frac{1}{N} \sum_{i=1}^N [\{\psi(\hat{\alpha}^\top \mathbf{G}_i) - \psi(\bar{\alpha}^\top \mathbf{G}_i)\} Z_{ij}] \right| \\ &= \left| \frac{1}{n} \sum_{i=1}^n \{\psi_1(\tilde{\alpha}_j^\top \mathbf{G}_i)(\bar{\alpha} - \hat{\alpha})^\top \mathbf{G}_i Z_{ij}\} \right. \\ &\quad \left. + \frac{1}{N} \sum_{i=1}^N \{\psi_1(\tilde{\alpha}_j^\top \mathbf{G}_i)(\hat{\alpha} - \bar{\alpha})^\top \mathbf{G}_i Z_{ij}\} \right| \\ &\leq \|\hat{\alpha} - \bar{\alpha}\|_1 \left\| \frac{1}{n} \sum_{i=1}^n \{\psi_1(\tilde{\alpha}_j^\top \mathbf{G}_i) \mathbf{G}_i Z_{ij}\} - \frac{1}{N} \sum_{i=1}^N \{\psi_1(\tilde{\alpha}_j^\top \mathbf{G}_i) \mathbf{G}_i Z_{ij}\} \right\|_\infty, \end{aligned}$$

where $\tilde{\alpha}_j = u_j \hat{\alpha} + (1 - u_j) \bar{\alpha}$, for some $0 < u_j < 1$, $j = 0, \dots, m-1$. Consider the k -th coordinate of \mathbf{G}_i . For technical convenience, we assume that n is divisible by N . Let

$$V_{ijk} \triangleq \left\{ \frac{n}{N} - 1 \right\} \psi_1(\tilde{\alpha}_j^\top \mathbf{G}_i) G_{ik} Z_{ij} + \frac{n}{N} \sum_{s=n+(N/n-1)(i+1)+1}^{n+i(N/n-1)} \psi_1(\tilde{\alpha}_j^\top \mathbf{G}_s) G_{sk} Z_{sj},$$

then, \exists constant $C_7 > 0$, such that $|V_{ijk}| \leq C_7$. Moreover, we have $\mathbb{E}(V_{ijk}) = 0$; therefore, $\mathbb{E}(V_{ijk}^2) = \text{Var}(V_{ijk}) = \{(n - N)^2/N^2 + n^2/N^2(N/n - 1)\} \text{Var}\{\psi_1(\tilde{\alpha}_j^\top \mathbf{G}) G_k Z_j\} \leq C_8$ for some constant $C_8 > 0$. Let $t = \sqrt{C_8 \ln\{(q+1)/\epsilon\}/n}$, since $\ln(q+1) = o(n)$, $\exists n$ large enough, such that $t^2/C_8 \leq 3t/C_7$, then by Bernstein's inequality, we have

$$\mathbb{P}\left(\left|\frac{1}{n} \sum_{i=1}^n V_{ijk}\right| \geq t\right) = \mathbb{P}\left(\left|\sum_{i=1}^n V_{ijk}\right| \geq nt\right) \leq \exp\left(-\frac{nt^2}{C_8}\right) = \frac{\epsilon}{q+1}. \quad (\text{S36})$$

Let V_j denote $\sup_{k=0, \dots, q} |1/n \sum_{i=1}^n V_{ijk}|$, and

$$V_j < t \Rightarrow \frac{V_j \sqrt{n}}{\sqrt{C_8 \ln\{e(q+1)\}}} \leq \sqrt{1 - \ln(\epsilon)}.$$

It follows that

$$\begin{aligned} & \mathbb{P}\left[\frac{V_j \sqrt{n}}{\sqrt{C_8 \ln\{e(q+1)\}}} \leq \sqrt{1 - \ln(\epsilon)}\right] \geq \mathbb{P}(V_j < t) = \\ & 1 - \mathbb{P}(V_j \geq t) \geq 1 - \sum_{k=0}^q \mathbb{P}\left(\left|\frac{1}{n} \sum_{i=1}^n V_{ijk}\right| \geq t\right) = 1 - \epsilon. \end{aligned}$$

Therefore, $\frac{V_j\sqrt{n}}{\sqrt{C_8 \ln\{e(q+1)\}}} = O_p(1)$, for $j = 0, \dots, m-1$, it follows that

$$\left\| \frac{1}{n} \sum_{i=1}^n \left\{ \frac{1}{\pi^s(\bar{\gamma})} \psi_1(\tilde{\alpha}_j^T \mathbf{G}_i) \mathbf{G}_i Z_{ij} \right\} - \frac{1}{N} \sum_{i=1}^N \left\{ \psi_1(\tilde{\alpha}_j^T \mathbf{G}_i) \mathbf{G}_i Z_{ij} \right\} \right\|_{\infty} = V_j = O_p(\sqrt{\ln(q+1)/n}).$$

Therefore,

$$|\Delta_{5,j}^1 + \Delta_{5,j}^2| \leq O_p(\sqrt{\ln(q+1)/n}) O_p(|S_{\bar{\alpha}}| \sqrt{\ln(q+1)/n}) = o_p(1/\sqrt{n}),$$

it follows that $\Delta_{5,j}^1 + \Delta_{5,j}^2 = o_p(1/\sqrt{n})$. Hence, by central limit theorem and continuous mapping theorem, we have

$$\sqrt{n}(\hat{\beta}^s - \beta^*) \xrightarrow{d} \sqrt{n} \Gamma^{-1} \tau^s \{\bar{\alpha}, \beta^*, \pi^*(\mathbf{X})\} \sim N(0, \Sigma^s).$$

V.2 Variance comparison

Because the following relationship holds,

$$\begin{aligned} \frac{1}{n} \mathbf{\Lambda}^s &= \frac{1}{n} \mathbb{E} \left[\left\{ Y - \frac{N-n}{N} \psi(\bar{\alpha}^T \mathbf{G}) - \frac{n}{N} \psi(\beta^{*T} \mathbf{Z}) \right\}^2 \mathbf{Z} \mathbf{Z}^T \right] \\ &\quad + \frac{(N-n)}{N^2} \mathbb{E} [\{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\}^2 \mathbf{Z} \mathbf{Z}^T] \\ &= \frac{1}{n} \mathbb{E} \left[\{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2 \mathbf{Z} \mathbf{Z}^T + \left(\frac{n}{N}\right)^2 \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\}^2 \mathbf{Z} \mathbf{Z}^T \right] \\ &\quad + \frac{2}{N} \mathbb{E} [\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\} \mathbf{Z} \mathbf{Z}^T] \\ &\quad + \frac{N-n}{N^2} \mathbb{E} [\{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\}^2 \mathbf{Z} \mathbf{Z}^T] \\ &= \frac{1}{n} \mathbb{E} [\{Y - \psi(\bar{\alpha}^T \mathbf{G})\}^2 \mathbf{Z} \mathbf{Z}^T] + \frac{2}{N} \mathbb{E} [\{Y - \psi(\bar{\alpha}^T \mathbf{G})\} \{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\} \mathbf{Z} \mathbf{Z}^T] \\ &\quad + \frac{1}{N} \mathbb{E} [\{\psi(\bar{\alpha}^T \mathbf{G}) - \psi(\beta^{*T} \mathbf{Z})\}^2 \mathbf{Z} \mathbf{Z}^T] \\ &= \frac{1}{N} \mathbf{\Lambda}, \end{aligned}$$

we obtain

$$\Sigma^s/n = \Gamma^{-1} \left(\frac{1}{n} \mathbf{\Lambda}^s \right) \Gamma^{-1} = \Gamma^{-1} \left(\frac{1}{N} \mathbf{\Lambda} \right) \Gamma^{-1} = \Sigma/N.$$

VI Details of the application

VI.1 Pre-processing details of the community and crime dataset

We pre-process the data in following steps:

Step 1. remove 22 covariates missing 84% of data and 2 variables missing roughly 59% of data;

Step 2. remove covariates with weak linear relationships to the response `ViolentCrimesPerPop` based on their correlation coefficients.

Step 3. remove covariates that exhibit multi-collinearity based on their values of variance inflation factors.

After the process, we obtain 1993 observations of 26 covariates.

VI.2 Test results of the covariate shift

VI.2.1 Kernel two-sample test with maximum mean discrepancy

- Kernel: $\exp(-\|\cdot\|_2^2)$
- MMD: 0.39227
- P-value: 0.001

VI.2.2 Bootstrap KS-tests for univariate covariates

Covariate	Bootstrap-KS P-value	KS-test Statistic	KS-test Approximate P-value
<code>racePctHisp</code>	0.000	0.335	0.000
<code>pctWWage</code>	0.000	0.230	0.000
<code>pctWInvInc</code>	0.000	0.330	0.000
<code>blackPerCap</code>	0.000	0.421	0.000
<code>PctLess9thGrade</code>	0.010	0.119	0.010
<code>PctUnemployed</code>	0.000	0.231	0.000
<code>PctOccupManu</code>	0.000	0.205	0.000
<code>MalePctDivorce</code>	0.000	0.373	0.000
<code>MalePctNevMarr</code>	0.000	0.202	0.000
<code>PctTeen2Par</code>	0.000	0.289	0.000
<code>PctIlleg</code>	0.000	0.200	0.000
<code>NumImmig</code>	0.000	0.267	0.000
<code>PctImmigRec10</code>	0.001	0.141	0.001
<code>PctHousLess3BR</code>	0.000	0.218	0.000

MedNumBR	0.000	0.138	0.000
HousVacant	0.000	0.184	0.000
PctHousOccup	0.000	0.195	0.000
PctHousOwnOcc	0.000	0.285	0.000
PctVacantBoarded	0.797	0.040	0.923
PctHousNoPhone	0.000	0.373	0.000
PctWOFullPlumb	0.000	0.146	0.000
RentLowQ	0.000	0.529	0.000
MedRentPctHousInc	0.062	0.089	0.099
NumInShelters	0.022	0.087	0.113
NumStreet	0.001	0.101	0.043
PopDens	0.000	0.266	0.000

Table S1: Bootstrap KS-tests for univariate covariates

VI.3 Design of basis functions

In this application, we design basis functions in the following way: Given $\{X_{ij}\}_{j=1}^N$, i.e., N samples of the i -th coordinate of \mathbf{X} , let $\{\xi_{ij}\}_{j=1}^{n_k}$ be the n_k points equally spaced within the $[-a_i, b_i]$, where $a_i = \min_{j=1, \dots, N} X_{ij}$ and $b_i = \max_{j=1, \dots, N} X_{ij}$. Let $f_{ij}(\mathbf{X})$ denote $(X_i - \xi_{ij})_+$, $i = 1, \dots, d; j = 1, \dots, n_k$. Let $\mathbf{F} = \{1, f_{11}(\mathbf{X}), \dots, f_{1n_k}(\mathbf{X}), \dots, f_{d1}(\mathbf{X}), \dots, f_{dn_k}(\mathbf{X})\}^T$ be the basis functions of the PS model, and $\mathbf{G} = \{\mathbf{F}^T, (\mathbf{Z} \otimes \mathbf{F})^T\}^T$. We choose $n_k = 4$ in this application.