

The inviscid fixed point of the multi-dimensional Burgers-KPZ equation

Liubov Gosteva¹, Malo Tarpin², Nicolás Wschebor³, Léonie Canet^{1,4}

¹ *Université Grenoble Alpes, CNRS, LPMMC, 38000 Grenoble, France*

² *Ecole Centrale de Lyon, CNRS, Université Claude Bernard Lyon 1, INSA Lyon, LMFA, 69130, Ecully, France*

³ *Instituto de Física, Facultad de Ingeniería, Universidad de la República, J.H.y Reissig 565, 11000 Montevideo, Uruguay*

⁴ *Institut Universitaire de France, 5 rue Descartes, 75005 Paris*

A new scaling regime characterized by a $z = 1$ dynamical critical exponent has been reported in several numerical simulations of the one-dimensional Kardar-Parisi-Zhang and noisy Burgers equations. This scaling was found to emerge in the tensionless limit for the interface and in the inviscid limit for the fluid. Based on functional renormalization group, the origin of this scaling has been elucidated. It was shown to be controlled by a yet unpredicted fixed point of the one-dimensional Burgers-KPZ equation, termed inviscid Burgers fixed point. The associated universal properties, including the scaling function, were calculated. Here, we generalize this analysis to the multi-dimensional Burgers-KPZ equation. We show that the inviscid-Burgers fixed point exists in all dimensions d , and that it controls the large momentum behavior of the correlation functions in the inviscid limit. It turns out that it yields in all d the same super-universal value $z = 1$ for the dynamical exponent.

I. INTRODUCTION

The Kardar-Parisi-Zhang (KPZ) universality class is impressive by its broadness, which keeps extending. Beyond the realm of stochastically growing interfaces – for which the KPZ equation was originally derived [1] – the KPZ scaling has been both unveiled theoretically and observed experimentally in many quantum systems ranging from integrable systems, such as Heisenberg quantum spin chains [2, 3], to driven-dissipative systems, such as exciton-polariton condensates [4]. It has also emerged in quantum information in the scaling of entanglement growth [5] or in Anderson localization in the scaling of wave-packets density fluctuations [6]. This broadness was first recognized for classical systems [7, 8], since the KPZ equation maps to the noisy Burgers equation for randomly stirred fluid [9], as well as to the model of directed polymers in random media [10]. All of these problems are whole research fields in themselves, and a lot of efforts have been devoted to understanding the rich physics of each.

The hallmark of the KPZ universality class in dimension $d = 1$ is the superdiffusive scaling $z = 3/2$ [1], which was first reported for the noisy Burgers equation in [11, 12]. This scaling differs from the diffusive equilibrium one $z = 2$, which corresponds to the scaling of the linear Edwards-Wilkinson (EW) equation [13]. In the KPZ equation, the noise source is microscopic, in the sense that it predominantly acts at small scales. In this case, one is interested in the large-distance long-time behavior of the interface. In contrast, in the context of the Burgers equation, one rather considers a large-scale stochastic forcing, and the regime of interest is the inertial range, which corresponds to the small-scale short-time behavior. In this regime, the system is characterized by a kinetic energy spectrum $E(k) \sim k^{-2}$, and strong intermittency effects [14]. Intermittency effects are usually evidenced through static equal-time statistics. In particular, a prominent example are the structure functions,

whose scaling reflects a characteristic bi-fractal distribution of anomalous exponents [15].

The temporal properties are much less discussed. In this respect, a pioneering work was carried out in Ref. [16] using the mapping from the noisy Burgers equation to the disordered problem of directed polymers, and applying the replica method to compute averages over the disorder. It led to a solution obtained from a variational ansatz featuring a $z = 1$ dynamical exponent, which was interpreted as a convective scaling associated with the formation of cellular patterns. This variational solution is expected to become exact in the limit of infinite dimension $d \rightarrow \infty$ and infinite Reynolds number (inviscid limit). Whereas this result was obtained in the case of a large-scale forcing, a similar $z = 1$ dynamical exponent was also uncovered in different simulations of Burgers [17, 18] or KPZ [19] equations with short-scale microscopic noise in $d = 1$ and in the inviscid (respectively, tensionless) limit. The same scaling was found in a strongly interacting one-dimensional quantum bosonic system [20]. Using functional renormalization group (FRG) [21], this scaling was shown to result from the existence of a yet unpredicted fixed point of the KPZ equation, termed the inviscid Burgers fixed point.

In this work, we show that the inviscid Burgers fixed point exists in all dimensions, and independently of the scale of the forcing, as long as the latter remains Gaussian. This bridges the result of [16] valid for $d \rightarrow \infty$ and large-scale forcing with the ones of [17, 19–21] obtained for $d = 1$ and small-scale noise. This fixed point controls the large momentum behavior (UV scales, short distances) of the correlation functions in the limit $\nu \rightarrow 0$, and it yields a $z = 1$ dynamical exponent in all dimensions. In fact, a similar $z = 1$ scaling was also unveiled in the large-momentum behavior of correlation functions in Navier-Stokes turbulence [22, 23], and was related in this context to the random sweeping effect, that is, the sweeping of small-scale velocities by the large scale eddies of the turbulent flow. We show in this paper that this common

scaling is ultimately rooted in the Galilean invariance and the existence of a non-Gaussian fixed point. Our derivation of the $z = 1$ dynamical exponent in the multi-dimensional Burgers-KPZ equation stems from FRG, and deeply exploits the Ward identities associated with the symmetries of the underlying dynamics.

The remaining of the paper is organized as follows. We construct in Sec. II A an appropriate action for the d -dimensional Burgers equation, and discuss its symmetries and the associated Ward identities in Sec. II B. We introduce in Sec. III the FRG formalism, and elucidate in Sec. IV, based on a simple approximation, the fixed-point structure of the Burgers-KPZ equation, showing in particular the existence of the inviscid Burgers fixed point in all d . In Sec. V, we show that the flow equation for the two-point velocity correlation function can be closed in the large momentum p limit, and this closure is asymptotically exact for $p \rightarrow \infty$. We finally present the fixed-point solution of this flow equation and show that it features the $z = 1$ dynamical exponent in all d .

II. FIELD THEORY FORMULATION OF THE STOCHASTIC BURGERS-KPZ EQUATION

A. Action for the d -dimensional Burgers equation

The Burgers equation [9] is a simplified version of the Navier-Stokes equations for a potential flow – which satisfies $\nabla \times \mathbf{v} = 0$ – without pressure terms, and stirred by a stochastic force \mathbf{f} . It reads as

$$\partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} = \nu \nabla^2 \mathbf{v} + \mathbf{f}, \quad (1)$$

with ν the kinematic viscosity of the fluid. The random force is chosen to be Gaussian distributed with zero mean and covariance

$$\langle f_\alpha(t, \mathbf{x}) f_\beta(t', \mathbf{x}') \rangle = 2\delta(t - t') N_{\alpha\beta}(\mathbf{x} - \mathbf{x}'), \quad (2)$$

where N is centered around a large scale L to generate turbulence. Since the velocity is irrotational, one can define a scalar function h such that $\mathbf{v} = -\lambda \nabla h$ with λ a real parameter. The dynamics of this field, interpreted as the height of an interface, is then given by the KPZ equation

$$\partial_t h = \nu \nabla^2 h + \frac{\lambda}{2} (\nabla h)^2 + \eta. \quad (3)$$

In the original KPZ equation [1], η is a Gaussian noise of zero mean and covariance

$$\langle \eta(t, \mathbf{x}) \eta(t', \mathbf{x}') \rangle = 2D\delta(t - t') \delta^d(\mathbf{x} - \mathbf{x}'). \quad (4)$$

Through the mapping to the Burgers equation, one obtains that the stochastic forcing is related to the KPZ noise as $\mathbf{f} = -\lambda \nabla \eta$. Thus, the covariance (4) corresponds for the Burgers equation to \mathbf{f} being the thermal noise – implying in particular that it acts at small scales.

In order to apply renormalization group techniques, one can cast the stochastic equation into a path integral formulation which encompasses all the trajectories emanating from different noise realizations, following the Martin-Siggia-Rose-Janssen-de Dominicis formalism [24–26]. For the one-dimensional Burgers equation, this is straightforward and one obtains [21]

$$\begin{aligned} \mathcal{Z}[j, \bar{j}] &= \int Dv D\bar{v} e^{-\mathcal{S}[v, \bar{v}] + \int_{t,x} \{jv + \bar{j}\bar{v}\}}, \\ \mathcal{S}[v, \bar{v}] &= \int_{t,x} \left\{ \bar{v} \left[\partial_t v + v \partial_x v - \nu \partial_x^2 v \right] - \mathcal{D}(\partial_x \bar{v})^2 \right\}, \end{aligned} \quad (5)$$

where $\mathcal{D} = D\lambda^2$, and the conservative form of the noise term is inherited from the mapping with the KPZ equation.

In higher dimension d , one should further impose the irrotationality constraint. In the seminal work of Forster-Nelson-Stephen [12], the one-dimensional Burgers equation is formally continued to arbitrary dimension by using the identity $\mathbf{v} \cdot \nabla \mathbf{v} = \frac{1}{2} \nabla \mathbf{v}^2 - \mathbf{v} \times (\nabla \times \mathbf{v})$ and deleting the last term. However, in practice, only the one-dimensional case is studied in Ref. [12]. Since the FRG calculations presented in the following involve arbitrary n -point correlation functions, the irrotationality constraint must be explicitly encoded in the action. This can be simply achieved by introducing another response field $\bar{\mathbf{w}}$ which acts as a Lagrange multiplier for this constraint. This procedure is similar to the one used to impose incompressibility in the Navier-Stokes action [27, 28]. Furthermore, one can check from the KPZ mapping that the response velocity field $\bar{\mathbf{v}}$ should also be irrotational. We hence introduce another response field $\bar{\mathbf{w}}$ to impose this constraint. However, it turns out that the action resulting from adding these two auxiliary Lagrange multipliers is ill-defined as its Hessian is not invertible, which implies that the propagator does not exist, preventing any field-theoretical calculation. This is due to the presence of zero modes, which originate in the fact that the $\bar{\mathbf{w}}$ and $\bar{\mathbf{w}}$ fields are coupled only through their transverse parts. To fix their longitudinal parts and thereby remove the zero modes, one can introduce the additional scalar Lagrange multiplier fields $\theta, \bar{\theta}$. Thus, we obtain the following action for the Burgers-KPZ equation

$$\mathcal{Z}[\mathcal{J}] = \int D\Phi e^{-\mathcal{S}[\Phi] + \int_{t,x} \mathcal{J}\Phi} \quad (6)$$

$$\begin{aligned} \mathcal{S}[\Phi] &= \int_{t,x} \left\{ \bar{\mathbf{v}} \cdot [\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} - \nu \nabla^2 \mathbf{v}] - \mathcal{D}(\nabla \cdot \bar{\mathbf{v}})^2 \right. \\ &\quad \left. + \bar{\mathbf{v}} \cdot (\nabla \times \bar{\mathbf{w}}) + \bar{\mathbf{w}} \cdot (\nabla \times \mathbf{v}) + \bar{\theta} \nabla \cdot \bar{\mathbf{w}} + \bar{\mathbf{w}} \cdot \nabla \theta \right\}. \end{aligned} \quad (7)$$

Here and in the following, Φ denotes the multiplet of fields $(\mathbf{v}, \bar{\mathbf{v}}, \mathbf{w}, \bar{\mathbf{w}}, \theta, \bar{\theta})$, \mathcal{J} the multiplet of associated sources $(\mathbf{j}, \bar{\mathbf{j}}, \mathbf{L}, \bar{\mathbf{L}}, K, \bar{K})$, and summation over all fields is implicit in the notation $\mathcal{J}\Phi$. Note that, in the presence of a large-scale forcing (2) instead of the KPZ-

inherited noise term, the latter can be simply replaced with $\int_{t,\mathbf{x},\mathbf{x}'} \bar{v}_\alpha(t,\mathbf{x}) N_{\alpha\beta}(\mathbf{x}-\mathbf{x}') \bar{v}_\beta(t,\mathbf{x}')$.

Let us note that we have conveniently used vectorial product and mixed product notations, which are appropriate in $d=3$ (or $d=2$). However, this can be extended to arbitrary dimensions by expressing any product of the form $\mathbf{c} \cdot (\mathbf{a} \times \mathbf{b})$ in an anti-symmetrized form, such as $c_{\alpha\beta}(a_\alpha b_\beta - b_\alpha a_\beta)$ where c is now a rank-two tensor. For dimensions higher than $d=3$, it requires to introduce supplementary constraints to keep the Hessian of the action invertible. We show in Appendix A how this can be achieved without changing the discussion below. For simplicity, we keep using the notation of (7) in the following, and also express vectorial products using the Levi-Civita tensor $\epsilon_{\alpha\beta\gamma}$, bearing in mind that they can be replaced by anti-symmetric products in arbitrary dimensions $d > 1$. All the results presented in the following are thus valid in any d .

B. Extended symmetries and Ward identities

In order to constrain the FRG approximations, we are interested not only in symmetries in the strict sense, but also in extended symmetries. Extended symmetries are transformations that do not leave the action invariant, but induce a variation linear in the fields. These extended symmetries allow one to derive exact identities on correlation functions, under the form of Ward identities, which are typically more constraining than their non-extended counter-parts. The Ward identities follow from the requirement that the symmetry transformations do not affect the measure of the path integral (6) [28, 29]. This leads to functional identities satisfied by $\mathcal{W} = \ln \mathcal{Z}$, the generating functional of connected correlation functions. Equivalently, Ward identities can be derived for the Legendre transform of \mathcal{W} , which is called the effective action and denoted Γ . The effective action is the generating functional of one-particle irreducible correlation functions, also termed vertex functions. It is defined by

$$\Gamma[\Psi] = \sup_{\mathcal{J}} \left[\int_{t,\mathbf{x}} \mathcal{J}\Psi - \mathcal{W}[\mathcal{J}] \right] \quad (8)$$

where $\Psi \equiv \langle \Phi \rangle_{\mathcal{J}}$ are the expectation values of the fields in the presence of the sources \mathcal{J} . Since the FRG is primarily formulated in terms of vertex functions, we express in the following the Ward identities on the effective action.

The Burgers action (7) possesses three extended symmetries, in close analogy with the Navier-Stokes action [28, 29], that we now expound. Note that the form of these extended symmetries is independent of the precise profile of the noise as long as it is Gaussian and white in time. As a consequence, all the Ward identities derived in the following for n -point vertices ($n > 2$) hold both for a short-scale noise and for a large-scale forcing.

1. Fully-gauged shift symmetries of the auxiliary fields.

The four transformations

$$\varphi(t,\mathbf{x}) \rightarrow \varphi(t,\mathbf{x}) + \varepsilon_\varphi(t,\mathbf{x}), \quad \varphi \in \{\mathbf{w}, \bar{\mathbf{w}}, \theta, \bar{\theta}\},$$

where ε_φ is either a scalar or a vectorial infinitesimal field, are extended symmetries of the Burgers action. Indeed, as the terms involving the auxiliary fields are quadratic, their variation under a shift is automatically at most linear. They yield the functional Ward identities

$$\frac{\delta\Gamma[\Psi]}{\delta\Psi_i} = \frac{\delta S[\Psi]}{\delta\Psi_i}, \quad \Psi_i \in \left\{ \langle \mathbf{w} \rangle_{\mathcal{J}}, \langle \bar{\mathbf{w}} \rangle_{\mathcal{J}}, \langle \theta \rangle_{\mathcal{J}}, \langle \bar{\theta} \rangle_{\mathcal{J}} \right\}. \quad (9)$$

Note that the corresponding global (non-infinitesimal) constant shifts are exact symmetries of the action, and the associated Ward identities have the same form as (9), but integrated over space and time. Thus, the identities (9), which are local both in space and time, are more constraining. The identities (9) imply that the full dependence of Γ on the mean values of the auxiliary fields $\mathbf{w}, \bar{\mathbf{w}}, \theta, \bar{\theta}$ is known explicitly and keeps the bare form determined by the action. This means that this whole sector is not renormalized, and hence we use in the following the same notation for the fields $\varphi \in \{\mathbf{w}, \bar{\mathbf{w}}, \theta, \bar{\theta}\}$ and their averages $\langle \varphi \rangle_{\mathcal{J}}$. In particular, these fields do not enter any n -point vertex function with $n > 2$. As a consequence, these vertex functions only involve velocity or response velocity fields, so that we set the notation

$$\Gamma_{\alpha_1 \dots \alpha_{m+n}}^{(m,n)}(t_1, \mathbf{x}_1; \dots; t_{m+n}, \mathbf{x}_{m+n}) \equiv \frac{\delta^{m+n} \Gamma}{\delta u_{\alpha_1}(t_1, \mathbf{x}_1) \dots \delta \bar{u}_{\alpha_{m+n}}(t_{m+n}, \mathbf{x}_{m+n})} \Big|_{\mathbf{u}, \bar{\mathbf{u}}=0}, \quad (10)$$

where the first m fields are average velocities $\mathbf{u} = \langle \mathbf{v} \rangle_{\mathcal{J}}$ and the n last are average response velocities $\bar{\mathbf{u}} = \langle \bar{\mathbf{v}} \rangle_{\mathcal{J}}$. Because of translational invariance in space and time, their Fourier transforms take the form

$$\Gamma_{\alpha_1 \dots \alpha_{m+n}}^{(m,n)}(\omega_1, \mathbf{q}_1; \dots; \omega_{m+n}, \mathbf{q}_{m+n}) = (2\pi)^{d+1} \delta \left(\sum_{\ell=1}^{m+n} \omega_\ell \right) \delta^d \left(\sum_{\ell=1}^{m+n} \mathbf{q}_\ell \right) \times \bar{\Gamma}_{\alpha_1 \dots \alpha_{m+n}}^{(m,n)}(\omega_1, \mathbf{q}_1; \dots; \omega_{m+n-1}, \mathbf{q}_{m+n-1}), \quad (11)$$

where the last frequency and momentum of $\bar{\Gamma}^{(m,n)}$ are implicit as they are fixed by the conservation of the total frequency and momentum.

2. Time-gauged shift symmetry of the response field.

The transformation

$$\begin{cases} \bar{v}_\alpha(t,\mathbf{x}) \rightarrow \bar{v}_\alpha(t,\mathbf{x}) + \bar{\varepsilon}_\alpha(t) \\ \bar{w}_\alpha(t,\mathbf{x}) \rightarrow \bar{w}_\alpha(t,\mathbf{x}) + \epsilon_{\alpha\beta\gamma} \bar{\varepsilon}_\beta(t) v_\gamma(t,\mathbf{x}) - \bar{\varepsilon}_\alpha(t) \theta(t,\mathbf{x}), \end{cases}$$

where $\bar{\varepsilon}$ is a vectorial infinitesimal time-dependent shift is an extended symmetry of the Burgers action. It leads to the functional Ward identity

$$\int_{\mathbf{x}} \left(\frac{\delta \Gamma}{\delta \bar{u}_\alpha} - \epsilon_{\alpha\beta\gamma} \frac{\delta \Gamma}{\delta \bar{w}_\beta} u_\gamma - \theta \frac{\delta \Gamma}{\delta \bar{w}_\alpha} \right) = \int_{\mathbf{x}} (\partial_t u_\alpha + \epsilon_{\alpha\beta\gamma} \partial_\beta w_\gamma). \quad (12)$$

Taking m functional derivatives of this identity with respect to \mathbf{u} and n with respect to $\bar{\mathbf{u}}$, and evaluating them at vanishing fields, one obtains an identity for $\Gamma^{(m,n+1)}$ from the first term in the left-hand-side. Because of the integration over space, this yields in Fourier space that one of the momentum associated with a response field is zero, and one finds that the corresponding vertex vanishes:

$$\bar{\Gamma}_{\alpha_1 \dots \alpha_{m+n+1}}^{(m,n+1)}(\omega_1, \mathbf{p}_1; \omega_2, \mathbf{p}_2; \dots, \underbrace{\omega_{k>m}, 0}_{\bar{\mathbf{u}}}; \dots) = 0, \quad (13)$$

for any m, n except the lowest ones, which are given by:

$$\begin{aligned} \bar{\Gamma}_{\alpha_1 \alpha}^{(1,1)}(\omega_1, 0) &= i\omega_1 \delta_{\alpha\alpha_1}, \\ \bar{\Gamma}_{\alpha_1 \alpha_2 \alpha}^{(2,1)}(\omega_1, \mathbf{p}; \omega_2, -\mathbf{p}) &= \delta_{\alpha\alpha_2} i p_{\alpha_1} - \delta_{\alpha\alpha_1} i p_{\alpha_2}, \end{aligned}$$

where the identity (9) was used in the second expression.

3. Time-gauged Galilean symmetry.

The transformation

$$\begin{cases} v_\alpha(t, \mathbf{x}) \rightarrow v_\alpha(t, \mathbf{x}) - \partial_t \varepsilon_\alpha(t) + \varepsilon_\beta(t) \partial_\beta v_\alpha(t, \mathbf{x}) \\ \varphi(t, \mathbf{x}) \rightarrow \varphi(t, \mathbf{x}) + \varepsilon_\beta(t) \partial_\beta \varphi(t, \mathbf{x}), \quad \varphi \in \{\bar{v}, \mathbf{w}, \bar{\mathbf{w}}, \theta, \bar{\theta}\} \end{cases}$$

where ε is a vectorial infinitesimal time-dependent function is an extended symmetry of the Burgers action. This reduces to the standard Galilean symmetry when ε is linear in t . This yields the functional identity

$$- \int_{\mathbf{x}} \partial_t^2 \bar{u}_\alpha = \int_{\mathbf{x}} \partial_t \frac{\delta \Gamma}{\delta \bar{u}_\alpha} + \sum_{\varphi \in \{\mathbf{u}, \bar{\mathbf{u}}, \mathbf{w}, \bar{\mathbf{w}}, \theta, \bar{\theta}\}} \int_{\mathbf{x}} \partial_\alpha \varphi \frac{\delta \Gamma}{\delta \varphi}. \quad (14)$$

Taking functional derivatives of this identity with respect to velocity and response velocity fields and evaluating them at zero fields provides exact relations for vertex functions with a zero momentum associated with a velocity field. In Fourier space, they read as

$$\begin{aligned} \bar{\Gamma}_{\alpha_1 \dots \alpha_{m+n}}^{(m+1,n)}(\omega, 0; \underbrace{\omega_1, \mathbf{p}_1; \dots; \omega_{m+n}, \mathbf{p}_{m+n}}_{\mathbf{u}}) = \\ - \sum_{j=1}^{m+n-1} \frac{p_j \alpha}{\omega} \left[\bar{\Gamma}_{\alpha_1 \dots \alpha_{m+n}}^{(m,n)}(\omega_1, \mathbf{p}_1; \dots; \omega_j + \omega, \mathbf{p}_j; \dots) - \right. \\ \left. \bar{\Gamma}_{\alpha_1 \dots \alpha_{m+n}}^{(m,n)}(\omega_1, \mathbf{p}_1; \dots; \omega_j, \mathbf{p}_j; \dots) \right], \quad (15) \end{aligned}$$

for any $m, n \geq 1$ except the lowest one:

$$\bar{\Gamma}_{\alpha\alpha_1}^{(1,1)}(\omega, 0) = i\omega \delta_{\alpha\alpha_1}. \quad (16)$$

As an example, these identities consist in two non-trivial relations for the 3-point vertex functions, which are

$$\begin{aligned} \bar{\Gamma}_{\alpha\alpha_1\alpha_2}^{(2,1)}(\omega, 0; \omega_1, \mathbf{p}_1) = \\ - \frac{p_{1\alpha}}{\omega} \left[\bar{\Gamma}_{\alpha_1\alpha_2}^{(1,1)}(\omega_1 + \omega, \mathbf{p}_1) - \bar{\Gamma}_{\alpha_1\alpha_2}^{(1,1)}(\omega_1, \mathbf{p}_1) \right], \quad (17) \end{aligned}$$

$$\begin{aligned} \bar{\Gamma}_{\alpha\alpha_1\alpha_2}^{(1,2)}(\omega, 0; \omega_1, \mathbf{p}_1) = \\ - \frac{p_{1\alpha}}{\omega} \left[\bar{\Gamma}_{\alpha_1\alpha_2}^{(0,2)}(\omega_1 + \omega, \mathbf{p}_1) - \bar{\Gamma}_{\alpha_1\alpha_2}^{(0,2)}(\omega_1, \mathbf{p}_1) \right], \quad (18) \end{aligned}$$

since any $\Gamma^{(m,0)}$ vanishes for causality reasons [30], and Galilean invariance provides no constraint on $\Gamma^{(0,3)}$.

III. FUNCTIONAL RENORMALIZATION GROUP FORMALISM FOR THE BURGERS-KPZ EQUATION

A. The Wetterich equation

We briefly set the basis of the FRG formalism, referring the reader to comprehensive introductions, such as [31–34], for more details. The path integral (6) encompasses the integration over all realizations of the fluctuating fields Φ , which evolve according to the stochastic equation (1) with different realizations of the noise, and over all scales (or all modes) of these fields. The idea of the functional renormalization group is based on the Wilsonian RG. Instead of integrating over all scales simultaneously, one integrates up to a certain RG scale κ^{-1} (or, in momentum space, κ) and derives the evolution of the theory with the change of the RG scale. From the technical point of view, this is achieved in FRG through the introduction of a scale-dependent ‘weight’ $\exp(-\Delta\mathcal{S}_\kappa)$ in the functional integral to suppress fluctuations below the RG scale κ :

$$\mathcal{Z}_\kappa = \int D\Phi \exp \left(-\mathcal{S}[\Phi] - \Delta\mathcal{S}_\kappa[\Phi] + \int_{t,\mathbf{x}} \mathcal{J}\Phi \right), \quad (19)$$

where $\Delta\mathcal{S}_\kappa[\Phi] \equiv \frac{1}{2} \int_{t,\mathbf{x},\mathbf{x}'} \Phi(t, \mathbf{x}) \mathcal{R}_\kappa(|\mathbf{x} - \mathbf{x}'|) \Phi(t, \mathbf{x}')$. The kernel \mathcal{R}_κ is called ‘regulator’ and must satisfy the following properties (in Fourier space):

1. $\mathcal{R}_\kappa(\mathbf{p}) \xrightarrow{\kappa \rightarrow \infty} \infty$: all fluctuations are frozen;
2. $\mathcal{R}_\kappa(\mathbf{p}) \xrightarrow{\kappa \rightarrow 0} 0$: all fluctuations are averaged over;
3. $\mathcal{R}_\kappa(\mathbf{p}) \stackrel{|\mathbf{p}| \ll \kappa}{\sim} \kappa^2$: the contribution of the ‘slow’ modes is suppressed;
4. $\mathcal{R}_\kappa(\mathbf{p}) \stackrel{|\mathbf{p}| \gg \kappa}{\rightarrow} 0$: ‘fast’ modes are unaltered and averaged over.

The central object within the FRG formalism is the effective average action Γ_κ , defined as the following modified Legendre transform of $\mathcal{W}_\kappa = \ln \mathcal{Z}_\kappa$

$$\Gamma_\kappa[\Psi] + \Delta \mathcal{S}_\kappa[\Psi] = \sup_{\mathcal{J}} \left[\int_{t,\mathbf{x}} \mathcal{J}\Psi - \mathcal{W}_\kappa[\mathcal{J}] \right], \quad (20)$$

where as before $\Psi \equiv \langle \Phi \rangle_{\mathcal{J}}$. In the large- κ limit, the microscopic action $\Gamma_\infty \equiv \mathcal{S}$ is recovered since all the fluctuations are frozen, while in the small- κ limit, the full effective action $\Gamma_0 \equiv \Gamma$, encompassing all the fluctuations at all scales, is obtained [33, 34]. For the Burgers-KPZ equation, the macroscopic scale is set by the size of the system, or by the integral scale if the forcing acts at large scales. This scale can be denoted by L (or $k_L = L^{-1}$ for momentum), and it embodies the scale beyond which (*i.e.* for $\kappa \lesssim k_L$) the RG flow essentially stops, because fluctuations are negligible or inexistent beyond this scale. The microscopic (momentum) scale, denoted Λ , defines the maximal momentum accessible in the system – Λ^{-1} can be thought of as the scale at which the continuous fluid description is valid.

The evolution of Γ_κ with the RG scale in between these two limits is given by the exact Wetterich equation [35–37]

$$\partial_\kappa \Gamma_\kappa = \frac{1}{2} \text{tr} \int_{\omega,\mathbf{q}} \partial_\kappa \mathcal{R}_\kappa \mathcal{G}_\kappa, \quad (21)$$

where the trace means summation over all fields, and

$$\mathcal{G}_\kappa \equiv \left(\Gamma_\kappa^{(2)} + \mathcal{R}_\kappa \right)^{-1} \quad (22)$$

is the propagator matrix, where the Hessian

$$\Gamma_\kappa^{(2)}(t_1, \mathbf{x}_1; t_2, \mathbf{x}_2) \equiv \left. \frac{\delta^2 \Gamma}{\delta \Psi(t_1, \mathbf{x}_1) \delta \Psi(t_2, \mathbf{x}_2)} \right|_{\Psi=0} \quad (23)$$

is the matrix of two-point vertex functions. Their exact flow can be obtained by taking functional derivatives of (21), yielding in Fourier space:

$$\begin{aligned} \partial_s \bar{\Gamma}_{\kappa,AB}^{(2)}(\varpi, \mathbf{p}) &= \frac{1}{2} \text{tr} \left[\tilde{\partial}_s \int_{\omega,\mathbf{q}} \bar{\mathcal{G}}_\kappa(\omega, \mathbf{q}) \right. \\ &\cdot \left\{ \bar{\Gamma}_{\kappa,AB}^{(4)}(\varpi, \mathbf{p}, -\varpi, -\mathbf{p}, \omega, \mathbf{q}) - \bar{\Gamma}_{\kappa,A}^{(3)}(\varpi, \mathbf{p}, \omega, \mathbf{q}) \right. \\ &\cdot \left. \left. \bar{\mathcal{G}}_\kappa(\varpi + \omega, \mathbf{p} + \mathbf{q}) \cdot \bar{\Gamma}_{\kappa,B}^{(3)}(-\varpi, -\mathbf{p}, \varpi + \omega, \mathbf{p} + \mathbf{q}) \right\} \right], \end{aligned} \quad (24)$$

where the operator $\tilde{\partial}_s$ only acts on the regulators, *i.e.* $\tilde{\partial}_s \equiv \partial_s \mathcal{R}_{\kappa,AB} \frac{\partial}{\partial \mathcal{R}_{\kappa,AB}}$, with $s = \ln(\kappa/\Lambda)$ the RG ‘time’.

The 3- and 4-point vertices $\bar{\Gamma}_{\kappa,A}^{(3)}$ and $\bar{\Gamma}_{\kappa,AB}^{(4)}$ are written as 6×6 matrices, the remaining fields being fixed to the external ones (labeled by the indices A and B from the set Ψ). The summation is carried over the implicit field indices (matrix product), and integration over the internal frequency and momentum (ω, \mathbf{q}) .

B. Structure of the propagator matrix

One can infer from the set of Ward identities (9) the general form of the effective average action as

$$\Gamma_\kappa[\mathbf{u}, \bar{\mathbf{u}}, \mathbf{w}, \bar{\mathbf{w}}, \theta, \bar{\theta}] = \tilde{\Gamma}_\kappa[\mathbf{u}, \bar{\mathbf{u}}] + \int_{t,\mathbf{x}} \left\{ \bar{\mathbf{u}} \cdot (\nabla \times \mathbf{w}) + \bar{\mathbf{w}} \cdot (\nabla \times \mathbf{u}) + \bar{\theta} \nabla \cdot \mathbf{w} + \bar{\mathbf{w}} \cdot \nabla \theta \right\}, \quad (25)$$

where the explicit part of Γ_κ containing the auxiliary fields is not renormalized. One can then deduce from it the general form of the Hessian of Γ_κ , and of the propagator matrix \mathcal{G}_κ , which are both given in Appendix B. Since the auxiliary-field sector does not renormalize, the RG procedure of progressive integration of fluctuation modes only needs to be carried out on the velocity and response velocity fields. Thus, one can choose a regulator matrix \mathcal{R}_κ with only three non-zero elements $\mathcal{R}_{\kappa,\mathbf{u}\bar{\mathbf{u}}} = \mathcal{R}_{\kappa,\bar{\mathbf{u}}\mathbf{u}} \equiv \mathcal{M}_\kappa$ and $\mathcal{R}_{\kappa,\bar{\mathbf{u}}\bar{\mathbf{u}}} \equiv \mathcal{N}_\kappa$. Moreover, since the auxiliary fields do not enter any $n > 2$ vertex function, they completely decouple from the flow equations (24). One can thus focus on the sole velocity and response velocity sector and consider the reduced vertex functions (10), and the reduced propagator matrix $\bar{\mathcal{G}}_\kappa$ given by (see Appendix B)

$$\bar{\mathcal{G}}_\kappa(\omega, \mathbf{p}) \equiv \begin{pmatrix} u_\beta & \bar{u}_\beta \\ \bar{u}_\alpha & 0 \end{pmatrix} \begin{pmatrix} \bar{\mathcal{G}}_{\kappa,\alpha\beta}^{(2,0)}(\omega, \mathbf{p}) & \bar{\mathcal{G}}_{\kappa,\alpha\beta}^{(1,1)}(\omega, \mathbf{p}) \\ \bar{\mathcal{G}}_{\kappa,\alpha\beta}^{(1,1)}(-\omega, \mathbf{p}) & 0 \end{pmatrix} \quad (26)$$

where

$$\bar{\mathcal{G}}_{\kappa,\alpha\beta}^{(1,1)}(\omega, \mathbf{p}) \equiv \frac{1}{\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\omega, \mathbf{p}) + \mathcal{M}_\kappa(\mathbf{p})} P_{\alpha\beta}^{\parallel}(\mathbf{p}), \quad (27)$$

$$\bar{\mathcal{G}}_{\kappa,\alpha\beta}^{(2,0)}(\omega, \mathbf{p}) \equiv \frac{-\left(\bar{\Gamma}_{\kappa,\parallel}^{(0,2)}(\omega, \mathbf{p}) + \mathcal{N}_\kappa(\mathbf{p})\right)}{\left|\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\omega, \mathbf{p}) + \mathcal{M}_\kappa(\mathbf{p})\right|^2} P_{\alpha\beta}^{\parallel}(\mathbf{p}), \quad (28)$$

with the longitudinal projector

$$P_{\alpha\beta}^{\parallel}(\mathbf{p}) = \frac{p_\alpha p_\beta}{p^2}. \quad (29)$$

The reduced propagator matrix is purely longitudinal for the Burgers-KPZ equation as a consequence of irrotationality, whereas it is purely transverse for the Navier-Stokes equation as a consequence of incompressibility [23, 27].

The exact flow equations (24) can be projected onto the reduced $(\mathbf{u}, \bar{\mathbf{u}})$ sector. The different terms involved in these flow equations can be written as diagrams, representing the field \mathbf{u} by a line carrying an ingoing-into-a-vertex arrow and the response field $\bar{\mathbf{u}}$ by a line carrying an outgoing-of-a-vertex arrow. Performing the matrix products and the trace for $A = B = \bar{\mathbf{u}}$ leads to the exact flow equation for $\bar{\Gamma}_\kappa^{(0,2)}$ represented diagrammatically on Fig. 1 (without the combinatorial factors).

Similarly, performing the matrix products and the trace in (24) for $A = \mathbf{u}$ and $B = \bar{\mathbf{u}}$ leads to the exact




FIG. 1. Diagrammatic representation of the flow equation of $\Gamma_\kappa^{(0,2)}$, which is given by the component $A = B = \bar{\mathbf{u}}$ of the exact flow equation (24).

flow equation for $\Gamma_\kappa^{(1,1)}$ represented diagrammatically on Fig. 2.




FIG. 2. Diagrammatic representation of the flow equation of $\Gamma_\kappa^{(1,1)}$, which is given by the component $A = \mathbf{u}$, $B = \bar{\mathbf{u}}$ of the exact flow equation (24).

These flow equations are exact, but they are not closed in the sense that they involve higher-order 3- and 4-point vertices. We first solve them in Sec. IV within a simple approximation, in order to determine the fixed-point structure of the Burgers-KPZ equation. We then stress in Sec. V that in the inviscid limit, the regime of interest is the large momentum regime, and we show that the FRG flow equations can be closed exactly in the limit of large momentum thanks to the set of Ward identities (13) and (15).

IV. FIXED POINT STRUCTURE OF THE BURGERS-KPZ EQUATION

The FRG flow equations are exact but they are functional partial differential equations which cannot be solved exactly in most cases (see, however, [38]). Several approximation schemes to exploit them have been developed and thoroughly studied. We refer the readers to Refs. [31, 33, 34] for a detailed account of the standard approximations. In this section, we use a very simple approximation, which consists in keeping the bare form of the action, but with effective flowing parameters, which means that the original parameters \mathcal{D} and ν of the Burgers-KPZ action (7) are promoted to κ -dependent effective parameters \mathcal{D}_κ and ν_κ . This amounts to defining

the following ansatz for $\tilde{\Gamma}_\kappa$ defined in Eq. (25):

$$\tilde{\Gamma}_\kappa[\mathbf{u}, \bar{\mathbf{u}}] = \int_{t,\mathbf{x}} \left\{ \bar{\mathbf{u}} \cdot [\partial_t \mathbf{u} + \lambda_\kappa (\mathbf{u} \cdot \nabla) \mathbf{u} - \nu_\kappa \nabla^2 \mathbf{u}] - \mathcal{D}_\kappa (\nabla \cdot \bar{\mathbf{u}})^2 \right\}, \quad (30)$$

where we also introduced a parameter λ_κ since the corresponding term could in principle be renormalized.

Despite its simplicity, it was shown that this ansatz enables one to capture the KPZ strong-coupling fixed point in all dimensions d [39], and also the inviscid Burgers fixed point in $d = 1$ [21]. It is thus sufficient to provide a *qualitative* picture of the phase diagram. Of course, in order to obtain *quantitative* estimates of the critical exponents and scaling functions, one needs to resort to improved levels of approximation, such as the ones introduced in [40–42].

The advantage of the approximation (30) is that it leads to very simple flow equations. They are obtained by projecting the exact flow equations (24) onto the reduced functional space defined by the ansatz (30). In particular, the associated two-point vertex functions are simply given by

$$\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\omega, \mathbf{q}) = i\omega + \nu_\kappa \mathbf{q}^2, \quad \bar{\Gamma}_{\kappa,\parallel}^{(0,2)}(\omega, \mathbf{q}) = -2\mathbf{q}^2 \mathcal{D}_\kappa, \quad (31)$$

and the only non-vanishing n -point vertex function with $n > 2$ is the one present in the original action which reads as

$$\bar{\Gamma}_{\kappa,\alpha\beta\gamma}^{(2,1)}(\omega_1, \mathbf{q}_1, \omega_2, \mathbf{q}_2) = -i\lambda_\kappa (q_1^\beta \delta_{\alpha\gamma} + q_2^\alpha \delta_{\beta\gamma}). \quad (32)$$

One can show that the Galilean invariance (through the Ward identity (17)) imposes that λ_κ is not renormalized, *i.e.* $\lambda_\kappa \equiv \lambda = 1$ at all scales [41, 43]. The flows of ν_κ and \mathcal{D}_κ can be obtained as

$$\begin{aligned} \partial_s \mathcal{D}_\kappa &= -\frac{1}{4d} \partial_{p_\alpha} \partial_{p_\alpha} \partial_s \bar{\Gamma}_{\kappa,\parallel}^{(0,2)}(\varpi, \mathbf{p}) \Big|_{\varpi=0, \mathbf{p}=0} \\ \partial_s \nu_\kappa &= \frac{1}{2d} \partial_{p_\alpha} \partial_{p_\alpha} \Re \left[\partial_s \bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi, \mathbf{p}) \Big|_{\varpi=0, \mathbf{p}=0} \right]. \end{aligned} \quad (33)$$

The calculation is detailed in Appendix C.

In order to study the fixed points, one introduces dimensionless quantities, denoted with a hat symbol. Momenta are measured in units of κ , *eg* $\hat{q} = q/\kappa$, and frequencies in units of $\kappa^2 \nu_\kappa$, *eg* $\hat{\omega} = \omega/(\kappa^2 \nu_\kappa)$. One defines the two anomalous dimensions

$$\eta_\kappa^\nu = -\partial_s \ln \nu_\kappa, \quad \eta_\kappa^\mathcal{D} = -\partial_s \ln \mathcal{D}_\kappa, \quad (34)$$

where $s \equiv \ln(\kappa/\Lambda)$ and $\partial_s = \kappa \partial_\kappa$. The critical roughness and dynamical exponents χ and z are obtained from the fixed-point values of these anomalous dimensions as [42]

$$z = 2 - \eta_\kappa^\nu, \quad \chi = \frac{1}{2}(2 - d + \eta_\kappa^\mathcal{D} - \eta_\kappa^\nu). \quad (35)$$

The dimensionless effective coupling is defined as $\hat{g}_\kappa = \kappa^{d-2} \lambda^2 \mathcal{D}_\kappa / \nu_\kappa^3$. Its flow equation is thus given by

$$\partial_s \hat{g}_\kappa = \hat{g}_\kappa (d - 2 - \eta_\kappa^\mathcal{D} + 3\eta_\kappa^\nu). \quad (36)$$




FIG. 3. FRG flows of \hat{w}_κ in $d = 2$ and in $d = 3$: plain lines are IR flows from $s = 0$ ($\kappa = \Lambda$) towards an IR stable fixed point (KPZ and EW) reached as $s \rightarrow -\infty$ ($\kappa \rightarrow 0$); dashed lines are UV flows from a large negative value of $s_*(d)$ towards an UV fixed point (IB and EW or RT) reached as $s = 0$ ($\kappa = \Lambda$).

In fact, in order to study the inviscid Burgers fixed point, which corresponds to $\hat{g}_* \rightarrow \infty$, it is convenient to change variables to $\hat{w}_\kappa = \hat{g}_\kappa / (1 + \hat{g}_\kappa)$, whose flow equation reads as

$$\partial_s \hat{w}_\kappa = \hat{w}_\kappa (1 - \hat{w}_\kappa) (d - 2 - \eta_\kappa^D + 3\eta_\kappa^\nu). \quad (37)$$

Let us discuss the solutions of the fixed-point equation $\partial_s \hat{w}_\kappa = 0$. There are three ways to satisfy this equation: $\hat{w}_* = 0$, $\hat{w}_* = 1$, or $(d - 2 - \eta_*^D + 3\eta_*^\nu) = 0$. This last equation yields, using the definition (35) of the critical exponents, the identity $z + \chi = 2$ valid in any d . There are two such solutions with $0 < \hat{w}_* < 1$ and $z + \chi = 2$, which are the KPZ fixed point and the roughening transition (RT) fixed point. In contrast, this relation is not imposed neither for $\hat{w}_*^{\text{EW}} = 0$, which is the EW fixed point, nor for $\hat{w}_*^{\text{IB}} = 1$, which is the IB fixed point (although Galilean symmetry is preserved at both these fixed points). In most cases, the actual fixed-point values of the anomalous dimensions η_*^ν and η_*^D cannot be read off from the fixed point equation for \hat{w}_κ alone, but one has to solve the full set of equations (37), (C4).

We numerically solved these equations for dimensions $0 \leq d \leq 6$. Note that in the numerics we absorbed the factor $v_d = (2^{d-1} \pi^{d/2} \Gamma(d/2))^{-1}$ coming from the d -dimensional angular integration into the coupling, defining $\tilde{g}_\kappa \equiv \hat{g}_\kappa v_d$ and $\tilde{w}_\kappa = \tilde{g}_\kappa / (1 + \tilde{g}_\kappa)$. The plots are displayed using either \tilde{w} or \tilde{g} for clarity purpose. In




FIG. 4. Flow diagram of the Burgers-KPZ equation: IR and UV fixed point values of \tilde{g}_* as a function of the dimension d . The arrows indicate IR flows.

each dimension, we first integrated the flow from $s = 0$ ($\kappa = \Lambda$) with different initial conditions \hat{w}_Λ towards the IR $s \rightarrow -\infty$ ($\kappa \rightarrow 0$) until a fixed point was reached, stopping at a value that we denote $s_*(d)$. This is always the KPZ fixed point for $d \leq 2$. In $d > 2$, this is either the KPZ fixed point, when $\hat{w}_\Lambda > \hat{w}_\Lambda^c$, or the EW fixed point, when $\hat{w}_\Lambda < \hat{w}_\Lambda^c$, where \hat{w}_Λ^c is the critical value for the roughening transition. We refer to this flow from small scales to large scales as the IR flow.

We then ran the flow backwards, starting at $s_*(d)$ from initial conditions $\hat{w}_{s_*(d)} \lesssim \hat{w}_*^{\text{KPZ}}$ and $\hat{w}_{s_*(d)} \gtrsim \hat{w}_*^{\text{KPZ}}$ and letting $s \rightarrow 0$, in order to evidence the UV stable fixed points. We refer to this flow from large scales to small scales as the UV flow. We found, as UV fixed points, the IB and EW fixed points for $d \leq 2$, and the IB and RT fixed points for $d > 2$. We show in Fig. 3 the IR and UV flows in $d = 2$ as an illustration of the $d \leq 2$ case, and in $d = 3$ as an illustration of the $d > 2$ case.

To summarize, there are four fixed points:

- EW fixed point: $\hat{w}_*^{\text{EW}} = 0$,
IR stable for $d > 2$, and UV stable for $d \leq 2$.
- IB fixed point: $\hat{w}_*^{\text{IB}} = 1$, always UV stable.
- KPZ fixed point: $0 < \hat{w}_*^{\text{KPZ}} < 1$, always IR stable.
- RT fixed point: $0 < \hat{w}_*^{\text{RT}} < 1$, always UV stable
when it exists, *ie* for $d > 2$.

The complete flow diagram as a function of the dimension is displayed in Fig. 4.

The values of the critical exponents are trivially determined for the EW fixed point in all d because $\eta_*^D = \eta_*^\nu = 0$ for $\hat{w}_*^{\text{EW}} = 0$ (see (C4)) and thus $z = 2$ and $\chi = (2 - d)/2$ from (35). They are also determined for the KPZ fixed point in $d = 1$ since the accidental time reversal symmetry imposes $\eta_*^\nu = \eta_*^D \equiv \eta_*$ [40], which yields $\eta_* = 1/2$, and thus $z = 3/2$ and $\chi = 1/2$. In all other cases, the values of η_*^ν and η_*^D are to be calculated from




FIG. 5. Sketch of the fixed-point structure of the Burgers-KPZ equation in $d \leq 2$ and in $d > 2$. The fixed points are represented by the dots, the red-shaded area represents the region which controls the IR modes, while the blue-shaded area represents the region which controls the UV modes. The crossover regions correspond to scales κ around κ_{IR} introduced in the text.

the numerical solution of the full equations (37), (C4). At the simple level of approximation presented in this section, the estimates of the exponents are poor. Obtaining accurate estimates requires much richer approximations [40–42]. However, it is worth emphasizing that the simple one used here already encompasses the two strong-coupling KPZ and IB fixed points in all dimensions. In contrast, the perturbative RG equations, even resummed to all orders of perturbation theory, only contain the EW and RT fixed points [44]. The perturbative RG flow just runs away to infinity for $d \geq 2$ and completely fails to capture the strong-coupling fixed points.

We show in the next section that the dynamical exponent at the IB fixed point can be determined using another approach to solve the FRG equations, which yields the exact result $z = 1$.

V. EXACT FLOW EQUATIONS IN THE LARGE-MOMENTUM LIMIT

A. Relevance of the IB fixed point

We showed in the previous section that the IB fixed point exists in all dimensions and that it is always UV stable (IR unstable). Thus, this fixed point controls UV (large momentum) modes. Let us explain why the associated behavior is observable in the inviscid limit.

The fixed point structure of the Burgers-KPZ equation is schematically depicted in Fig. 5. The RG flow tends in the IR to the KPZ fixed point, for any initial condition g_Λ in $d \leq 2$, or for any $g_\Lambda > g_\Lambda^c$ in $d > 2$. Within the simple approximation (30), one simply has $g_\Lambda^c = \Lambda^{2-d} \hat{g}_*^{\text{RT}}$, but this does not hold in general when the effective average action is not constrained to keep the form of the bare action. In any dimension, the KPZ fixed point \hat{g}_*^{KPZ} is fully (or locally) IR attractive. Let us define $g_* \equiv \hat{g}_*^{\text{KPZ}} \Lambda^{2-d}$, which corresponds to the value of the microscopic coupling g_Λ for which the system would already be at the IR fixed point, and thus all modes p are controlled by this fixed point [45]. If $g_\Lambda \neq g_*$, then \hat{g}_κ renormalizes, and one can consider that it reaches the vicinity of the IR fixed point for a RG time s_{IR} such that

$\hat{g}_{s_{\text{IR}}} = \alpha \hat{g}_*$ (if $g_\Lambda > g_*$) or $\hat{g}_{s_{\text{IR}}} = \hat{g}_*/\alpha$ (if $g_\Lambda < g_*$) where $\alpha > 1$ is of order one, *e.g.* $\alpha = 2$. The larger $|g_\Lambda - g_*|$ is, the longer the time $|s_{\text{IR}}|$, or equivalently the smaller $\kappa_{\text{IR}} = \Lambda e^{s_{\text{IR}}}$. Thus, if a mode p is such that $p \lesssim \kappa_{\text{IR}}$, the corresponding \hat{g}_p is closed to \hat{g}_* and it is controlled by the IR fixed point. This corresponds to the red-shaded area in Fig. 5. In contrast, the large momentum modes $p \gtrsim \kappa_{\text{IR}}$ are controlled by the associated UV fixed point, which is the IB fixed point for any $g_\Lambda > g_*$ (and otherwise the EW ($d \leq 2$) or the RT ($d > 2$) fixed point). The larger $|g_\Lambda - g_*|$ is, the smaller the RG scale κ_{IR} which delineates IR and UV modes, and thus the larger the extent of the UV modes exhibiting the IB scaling. This situation is favored by the limit $\nu \rightarrow 0$, since $g_\Lambda = \lambda^2 D/\nu^3$. This explains why this scaling arises in the inviscid limit. We now show that this scaling is $z = 1$, by focusing on the large p behavior of the FRG flow equations.

B. Closure at large momentum

In this section, we aim at calculating the dynamical exponent at the IB fixed point. As explained in the previous section, this amounts to studying the large momentum $|\mathbf{p}| \gtrsim \kappa_{\text{IR}}$ behavior of the flow equations, where $\kappa_{\text{IR}} \rightarrow 0$ as $\nu \rightarrow 0$. An approximation scheme [46–48] turns out to be particularly appropriate to study this large momentum limit. It consists in an expansion of the vertices entering a given flow equation in powers of the internal loop momentum. This expansion is fully justified by the presence of the (scale derivative of the) regulator $\partial_s \mathcal{R}_\kappa(\mathbf{q})$ in the loop integral. Indeed, the general property 3- of the regulator implies that the integral over the internal momentum \mathbf{q} is cut to values $|\mathbf{q}| \lesssim \kappa$. Thus, in the limit of large external momentum $\mathbf{p} \gg \kappa$, one has $|\mathbf{q}| \ll |\mathbf{p}|$. Because analyticity of the vertices is guaranteed at all scale κ by construction [34], one can Taylor expand them in powers of $|\mathbf{q}|$, since \mathbf{q} is compared to \mathbf{p} and $|\mathbf{q}|/|\mathbf{p}| \rightarrow 0$. This expansion becomes exact in the limit of infinite \mathbf{p} [46–48]. Let us stress that this analysis holds at or in the vicinity of a fixed point of the FRG flow equations. This ensures that, in the large $|\mathbf{p}|$ limit, only the ratio $|\mathbf{q}|/|\mathbf{p}| \rightarrow 0$ can appear, and not independent occurrences of $|\mathbf{q}|/\kappa$.

It was realized that, in the context of turbulence, this approximation scheme leads to exact results for generic n -point correlation functions in the limit of large momentum. Their derivation is grounded in the existence of extended symmetries and the related Ward identities, which allows one to close the corresponding flow equations in the large-momentum limit. This closure was first achieved for the incompressible Navier-Stokes equation in three, and then two dimensions [22, 23, 27, 49, 50], and it was recently extended for the one-dimensional Burgers-KPZ equation [21]. We now show that it can be generalized to the d -dimensional Burgers equation as well.

The derivation is strictly analogous to Refs. [21, 27], with the main difference that the transverse projectors

(for Navier-Stokes equation) are replaced with longitudinal ones (for Burgers equation). We report this derivation for the two-point functions in Appendix D for completeness. We emphasize that it can also be applied to arbitrary n -point correlation function, following the same proof as in [23]. We here only discuss the result for the two-point connected correlation function of the velocity $C_\kappa(\varpi, \mathbf{p}) \equiv G_{\kappa, \parallel}^{(2,0)}(\varpi, \mathbf{p})$, whose flow equation is derived in Appendix D and given by (D8). Fourier transforming this equation back in time delay, one obtains

$$\partial_\kappa C_\kappa(t, \mathbf{p}) = \frac{1}{d} p^2 C_\kappa(t, \mathbf{p}) \int_\omega \frac{\cos(\omega t) - 1}{\omega^2} \tilde{\partial}_s \int_{\mathbf{q}} C_\kappa(\omega, \mathbf{q}). \quad (38)$$

Note that this flow equation is identical to the one obtained for the Navier-Stokes equation, apart from the $1/d$ pre-factor which arises because the longitudinal rather than the transverse part of the correlation function matters. In fact, Appendix D shows that the only ingredients entering the derivation of this equation are the Ward identities associated with the response-shift and Galilean time-gauged symmetries. Thus, any system sharing these basic symmetries will display similar flow equations, leading in turn to a $z = 1$ dynamical exponent at large momentum (as shown below). This $z = 1$ is thus purely the imprint of the Galilean and shift symmetries in a scaling regime.

C. Solution at the fixed point

We showed in Sec. IV that the FRG flow always reaches an IR fixed point. At this fixed point, the (dimensionless) term $\tilde{\partial}_s \int_{\hat{\mathbf{q}}} \hat{C}_\kappa(\hat{\omega}, \hat{\mathbf{q}})$ becomes a simple κ -independent function $\hat{J}(\hat{\omega})$. Hence, the flow equation (38) becomes linear, and can be solved analytically in both limits of large and small time-delays t where the remaining integral over ω can be simplified (we refer the interested reader to [21, 23] for the details of the calculation). The solution reads as

$$C(t, \mathbf{p}) = C(0, \mathbf{p}) \times \begin{cases} \exp(-\mu_0(pt)^2), & t \ll \tau_c \\ \exp(-\mu_\infty p^2 |t|), & t \gg \tau_c \end{cases} \quad (39)$$

where μ_0, μ_∞ are non-universal constants and τ_c is a characteristic time scale. This solution shows that the short-time behavior of the correlation function is a scaling form with the variable $pt^{1/z} \equiv pt$, which proves that the associated dynamical exponent is $z = 1$. The large momentum and short time regime of the Burgers equation, thus exhibits the $z = 1$ dynamical exponent, independently of the dimension d , and independently of the range of the forcing, as long as it is Gaussian and white in time. This regime arises for small ν , and it extends as $\nu \rightarrow 0$.

The solution (39) also predicts a crossover to another regime, with $z = 2$, at large time delays. This crossover was also predicted for the Navier-Stokes equation and

for the one-dimensional Burgers equation. It turns out that it is challenging to observe this regime in numerical simulations, since the short-time Gaussian decay rapidly leads to noise-level signal, which prevents from resolving the long-time behavior [51]. A hint of this crossover appears in the simulation of the one-dimensional Burgers equation of Ref. [17] as a lack of collapse of the data at large time when the scaling variable pt is used. There is one case, reported in Ref. [52], where it has been unambiguously observed, which is the case of passive scalars transported by a turbulent Gaussian velocity field with finite time correlations (generalization of the Kraichnan model). We expect that this large-time regime is also difficult to observe in simulations of the Burgers-KPZ equation in $d > 1$.

VI. CONCLUSION AND OUTLOOK

In this work, we have proposed an action for the d -dimensional Burgers-KPZ equation which properly enforces the irrotationality of the flow, through the introduction of auxiliary response fields imposing dynamically the corresponding constraints, and preventing the appearance of unphysical zero modes. We have shown that this action possesses extended symmetries which play a similar role to the ones of the Navier-Stokes action. In the latter case, these symmetries were exploited to achieve the exact closure of the flow equations for any n -point correlation functions in the limit of large momentum. We have thus transposed this closure to the d -dimensional Burgers equation. The resulting flow equation for the two-point correlation function of the velocity can be solved exactly at the fixed point and the solution shows that it features at short times a scaling regime with the dynamical exponent $z = 1$. This scaling is deeply rooted in the Galilean and shift extended symmetries of the system, together with the existence of a non-Gaussian fixed point of the FRG equations. As a consequence, it is super-universal, in the sense that it does not depend on the dimension d , and it does not depend either on the scale of the stirring force/noise. It thus fully bridges the result of [16] in $d \rightarrow \infty$ with large-scale forcing and of [21] in $d = 1$ with small-scale noise.

The $z = 1$ scaling describes the temporal behavior at large momentum of the velocity correlation function. It gives no information on the equal-time statistics, in particular on the structure functions, since the solution (39) becomes trivial at equal time delays $t = 0$. In contrast with the large momentum $z = 1$ regime, the equal-time properties are sensitive to both the dimension and the scale of the forcing. For a large-scale forcing, the multi-dimensional Burgers equation leads to turbulence with strong intermittency, and it would be very interesting to calculate the corresponding structure functions using FRG. For a small-scale noise, the behavior of the system is controlled by the KPZ fixed point, and calculating the associated universal scaling functions in all d would also

be very desirable and is work in progress.

ACKNOWLEDGMENTS

L.C. acknowledges support from IUF (Institut Universitaire de France). L.G. acknowledges support by the MSCA Cofund QuanG (Grant Number : 101081458) funded by the European Union. Views and opinions ex-

pressed are however those of the authors only and do not necessarily reflect those of the European Union or Université Grenoble Alpes. Neither the European Union nor the granting authority can be held responsible for them. N.W. thanks the ‘Programa de Desarrollo de las Ciencias Básicas’ for its support and the LPMMC for a scientific invitation during which part of the present work was done. This work received the support of the French-Uruguayan Institute of Physics (IFUΦ).

Appendix A: Burgers-KPZ action for arbitrary dimensions

In this appendix, the action presented in Eq. (7) is generalized for arbitrary dimensions. As mentioned in Sec. II A, in any dimension, the rotational of a vector does not correspond to a (pseudo-)vector, but must be viewed as a rank-two antisymmetric tensor. This leads to the fact that the action given by Eq. (7) must be slightly modified. As shown here, once these modifications are implemented, the results of the three-dimensional case generalize to arbitrary d . In particular, once the equations of motion of the auxiliary fields (necessary to impose the irrotationality of the velocity and response velocity fields) are imposed, the velocity and response velocity sector is decoupled from the rest, the only imprint resulting from these constraints being a longitudinal projector in the reduced propagator.

Firstly, the case $d = 1$ has already been discussed in Sec. II A. The case $d = 2$ is a trivial particular case of the $d = 3$ one since in $d = 2$, the w_β and \bar{w}_β fields only have non-zero components with $\beta = 3$, and obviously only depend on the x_1 and x_2 coordinates, so the divergence condition becomes trivial. That is, the action given by Eq. (7) also applies to the case $d = 2$ except for the fact that the terms $\propto \theta$ or $\propto \bar{\theta}$ can be omitted.

We proceed to analyze the nontrivial case corresponding to $d > 3$. In such a case, the term for imposing the irrotationality of the velocity field must be replaced by (and similarly for the response velocity field):

$$\bar{\mathbf{w}} \cdot (\nabla \times \mathbf{v}) \rightarrow \frac{1}{2} \bar{w}_{\alpha\beta} (\partial_\alpha v_\beta - \partial_\beta v_\alpha) = \bar{w}_{\alpha\beta} \partial_\alpha v_\beta. \quad (\text{A1})$$

As in the three-dimensional case, when $d > 3$, it is necessary to cancel the longitudinal components of the field $\bar{\mathbf{w}}$. In $d = 3$, this is achieved by introducing a (pseudo-)scalar Lagrange multiplier θ . For generic dimension, it is necessary to introduce an antisymmetric tensor of rank $d - 3$, thereby including in the density of the action a term of type

$$\bar{w}_{\alpha\beta} \epsilon_{\alpha\beta\gamma\mu_1\mu_2\dots} \partial_\gamma \theta_{\mu_1\mu_2\dots} \quad (\text{A2})$$

where $\epsilon_{\alpha\beta\gamma\mu_1\mu_2\dots}$ is the completely antisymmetric pseudo-tensor of rank d existing in dimension d . This expression naturally reduces to the term included in (7) in $d = 3$ but is valid in generic dimension. One proceeds similarly for the \mathbf{w} field.

The problem that arises at this point is that the included terms do not involve the longitudinal components of the θ and $\bar{\theta}$ fields which, in contrast with the three-dimensional case, appear for higher dimensions. The solution to this difficulty is to iterate the procedure of imposing the cancellation of the longitudinal components by new Lagrange multipliers in the same way as described for the \mathbf{w} and $\bar{\mathbf{w}}$ fields. For example, in dimension $d = 4$ the θ and $\bar{\theta}$ fields are vectors. In such a case it is sufficient to add to the density of the action a term of type

$$\bar{\chi} \partial_\alpha \theta_\alpha \quad (\text{A3})$$

(and a similar one for $\bar{\theta}$). In general dimension d , this iterative procedure stops after adding $d - 1$ pairs of Lagrange multipliers, the last pair being scalars.

Since all the added auxiliary fields enter through quadratic terms in the action, they lead to similar extended symmetries (fully gauged shifts) as in $d = 3$. It follows that the auxiliary field sector is not renormalized, or otherwise stated that the effective action Γ keeps the same form as the bare action \mathcal{S} in this sector. Moreover, we have checked that in any dimension, this procedure respects the extended shift and Galilean symmetries, and that the Hessian of the action is invertible, resulting in a well-defined propagator. Finally, once the equations of motion of the auxiliary fields are employed, one can restrict to the reduced velocity and response velocity sector. The only remnant of the irrotationality constraints is the longitudinal projector in the reduced propagator.

Appendix B: General structure of the propagator matrix

In this section, the derivation is presented in the more intuitive $d = 3$ notation, but it can be generalized for arbitrary d following the procedure described in Appendix A. Most importantly, one obtains that the general form of the reduced propagator matrix in the $(\mathbf{u}, \bar{\mathbf{u}})$ sector is given by (B1), (B2) in any $d > 1$.

The general form of the Hessian of the effective average action Γ_κ can be inferred from its general structure given by (25). One obtains

$$\bar{\Gamma}_\kappa^{(2)}(\omega, \mathbf{p}) = \begin{matrix} & u_\beta & \bar{u}_\beta & w_\beta & \bar{w}_\beta & \theta & \bar{\theta} \\ \begin{matrix} u_\alpha \\ \bar{u}_\alpha \\ w_\alpha \\ \bar{w}_\alpha \\ \theta \\ \bar{\theta} \end{matrix} & \begin{pmatrix} 0 & \bar{\Gamma}_{\kappa,\alpha\beta}^{(1,1)}(-\omega, \mathbf{p}) & 0 & i\epsilon_{\alpha\gamma\beta}p_\gamma & 0 & 0 & 0 \\ \bar{\Gamma}_{\kappa,\alpha\beta}^{(1,1)}(\omega, \mathbf{p}) & \bar{\Gamma}_{\kappa,\alpha\beta}^{(0,2)}(\omega, \mathbf{p}) & i\epsilon_{\alpha\gamma\beta}p_\gamma & 0 & 0 & 0 & 0 \\ 0 & i\epsilon_{\alpha\gamma\beta}p_\gamma & 0 & 0 & 0 & 0 & -ip_\alpha \\ i\epsilon_{\alpha\gamma\beta}p_\gamma & 0 & 0 & 0 & 0 & ip_\alpha & 0 \\ 0 & 0 & 0 & 0 & -ip_\beta & 0 & 0 \\ 0 & 0 & 0 & ip_\beta & 0 & 0 & 0 \end{pmatrix} \end{matrix}$$

using the general notation (10) and (11) for the vertex functions. The regulator matrix is chosen as

$$\mathcal{R}_\kappa(\omega, \mathbf{p}) = \begin{matrix} & u_\beta & \bar{u}_\beta & w_\beta & \bar{w}_\beta & \theta & \bar{\theta} \\ \begin{matrix} u_\alpha \\ \bar{u}_\alpha \\ w_\alpha \\ \bar{w}_\alpha \\ \theta \\ \bar{\theta} \end{matrix} & \begin{pmatrix} 0 & \mathcal{M}_\kappa(\mathbf{p}) & 0 & 0 & 0 & 0 & 0 \\ \mathcal{M}_\kappa(\mathbf{p}) & \mathcal{N}_\kappa(\mathbf{p}) & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} P_{\alpha\beta}^{\parallel} \end{matrix}$$

The propagator is defined as the inverse of the matrix $\bar{\Gamma}_\kappa^{(2)} + \mathcal{R}_\kappa$, which is given by

$$\bar{\mathcal{G}}_\kappa(\omega, \mathbf{p}) = \begin{matrix} & u_\beta & \bar{u}_\beta & w_\beta & \bar{w}_\beta & \theta & \bar{\theta} \\ \begin{matrix} u_\alpha \\ \bar{u}_\alpha \\ w_\alpha \\ \bar{w}_\alpha \\ \theta \\ \bar{\theta} \end{matrix} & \begin{pmatrix} \bar{G}_{\kappa,\alpha\beta}^{(2,0)}(\omega, \mathbf{p}) & \bar{G}_{\kappa,\alpha\beta}^{(1,1)}(\omega, \mathbf{p}) & 0 & \frac{i\epsilon_{\alpha\gamma\beta}p_\gamma}{p^2} & 0 & 0 & 0 \\ \bar{G}_{\kappa,\alpha\beta}^{(1,1)}(-\omega, \mathbf{p}) & 0 & \frac{i\epsilon_{\alpha\gamma\beta}p_\gamma}{p^2} & 0 & 0 & 0 & 0 \\ 0 & \frac{i\epsilon_{\alpha\gamma\beta}p_\gamma}{p^2} & -\frac{1}{p^2}\bar{\Gamma}_{\kappa,\perp}^{(0,2)}(\omega, \mathbf{p})P_{\alpha\beta}^\perp & -\frac{1}{p^2}\bar{\Gamma}_{\kappa,\perp}^{(1,1)}(\omega, \mathbf{p})P_{\alpha\beta}^\perp & 0 & 0 & -\frac{ip_\alpha}{p^2} \\ \frac{i\epsilon_{\alpha\gamma\beta}p_\gamma}{p^2} & 0 & -\frac{1}{p^2}\bar{\Gamma}_{\kappa,\perp}^{(1,1)}(-\omega, \mathbf{p})P_{\alpha\beta}^\perp & 0 & 0 & \frac{ip_\alpha}{p^2} & 0 \\ 0 & 0 & 0 & 0 & -\frac{ip_\beta}{p^2} & 0 & 0 \\ 0 & 0 & \frac{ip_\beta}{p^2} & 0 & 0 & 0 & 0 \end{pmatrix} \end{matrix}$$

where

$$\bar{G}_{\kappa,\alpha\beta}^{(1,1)}(\omega, \mathbf{p}) \equiv \frac{1}{\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\omega, \mathbf{p}) + \mathcal{M}_\kappa(\mathbf{p})} P_{\alpha\beta}^{\parallel}(\mathbf{p}), \quad (\text{B1})$$

$$\bar{G}_{\kappa,\alpha\beta}^{(2,0)}(\omega, \mathbf{p}) \equiv \frac{-(\bar{\Gamma}_{\kappa,\parallel}^{(0,2)}(\omega, \mathbf{p}) + \mathcal{N}_\kappa(\mathbf{p}))}{\left|\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\omega, \mathbf{p}) + \mathcal{M}_\kappa(\mathbf{p})\right|^2} P_{\alpha\beta}^{\parallel}(\mathbf{p}), \quad (\text{B2})$$

with the longitudinal and transverse projectors defined as

$$P_{\alpha\beta}^{\parallel}(\mathbf{p}) = \frac{p_\alpha p_\beta}{p^2}, \quad P_{\alpha\beta}^\perp(\mathbf{p}) = \delta_{\alpha\beta} - \frac{p_\alpha p_\beta}{p^2}. \quad (\text{B3})$$

Appendix C: Flow equations within the approximation (30)

The flow equations for ν_κ and \mathcal{D}_κ can be obtained following the definitions (33) from the flows of $\Gamma_{\kappa,||}^{(1,1)}$ and $\Gamma_{\kappa,||}^{(0,2)}$, given by the exact equations (24). Within the simple approximation corresponding to the ansatz (30), only the vertex $\Gamma_\kappa^{(2,1)}$ is non-zero, such that only the diagrams (c) of Fig. 2 and 1 contribute. They are given by

$$\begin{aligned} \left[\partial_s \bar{\Gamma}_{\kappa,||}^{(1,1)}(\varpi, \mathbf{p}) \right]_{(c)} &= -\tilde{\partial}_s \left[P_{\alpha\beta}^\parallel(\mathbf{p}) \int_{\omega, \mathbf{q}} \bar{\Gamma}_{\kappa, \alpha ij}^{(2,1)}(\varpi, \mathbf{p}, \omega, \mathbf{q}) \bar{G}_{\kappa, jk}^{(1,1)}(\omega + \varpi, \mathbf{p} + \mathbf{q}) \bar{\Gamma}_{\kappa, kl\beta}^{(2,1)}(\omega + \varpi, \mathbf{p} + \mathbf{q}, -\omega, -\mathbf{q}) \bar{G}_{\kappa, li}^{(2,0)}(\omega, \mathbf{q}) \right] \\ \left[\partial_s \bar{\Gamma}_{\kappa,||}^{(0,2)}(\varpi, \mathbf{p}) \right]_{(c)} &= -\frac{1}{2} \tilde{\partial}_s \left[P_{\alpha\beta}^\parallel(\mathbf{p}) \int_{\omega, \mathbf{q}} \bar{\Gamma}_{\kappa, ij\alpha}^{(2,1)}(\omega, \mathbf{q}, -\omega - \varpi, -\mathbf{p} - \mathbf{q}) \bar{G}_{\kappa, jk}^{(2,0)}(\omega + \varpi, \mathbf{p} + \mathbf{q}) \right. \\ &\quad \left. \times \bar{\Gamma}_{\kappa, kl\beta}^{(2,1)}(\omega + \varpi, \mathbf{p} + \mathbf{q}, -\omega, -\mathbf{q}) \bar{G}_{\kappa, li}^{(2,0)}(\omega, \mathbf{q}) \right]. \end{aligned} \quad (C1)$$

One then replaces the propagators $\bar{G}_\kappa^{(2,0)}$ and $\bar{G}_\kappa^{(1,1)}$ with their explicit expressions (B1) and (B2), and inserts the simplified expressions (31) and (32) for the vertex functions $\bar{\Gamma}_\kappa^{(1,1)}$, $\bar{\Gamma}_\kappa^{(0,2)}$, and $\bar{\Gamma}_\kappa^{(2,1)}$. Applying the $\tilde{\partial}_s$ derivative, and the $\partial_{p_\alpha}^2$ derivative, one obtains

$$\begin{aligned} \partial_s \nu_\kappa &= \frac{\lambda_\kappa^2}{d} \int \frac{d\omega}{2\pi} \frac{d^d \mathbf{q}}{(2\pi)^d} \frac{y}{(\ell_\kappa^2(y) + \omega^2)^3} \times \\ &\quad \left\{ 2\partial_s \mathcal{M}_\kappa(y) \left[k_\kappa(y) \left((d-1)\ell_\kappa^2(y) - (1+d)\omega^2 \right) + 2y\partial_y k_\kappa(y) (\ell_\kappa^2(y) - \omega^2) \right] \right. \\ &\quad \left. + \partial_s \mathcal{N}_\kappa(y) \left[-\ell_\kappa(y) (\ell_\kappa^2(y) + \omega^2) + 2y\partial_y \ell_\kappa(y) (\ell_\kappa^2(y) - \omega^2) \right] \right\}, \\ \partial_s \mathcal{D}_\kappa &= -\lambda_\kappa^2 \int \frac{d\omega}{2\pi} \frac{d^d \mathbf{q}}{(2\pi)^d} \frac{y^2 k_\kappa(y)}{(\ell_\kappa^2(y) + \omega^2)^3} \left[4k_\kappa(y) \ell_\kappa(y) \partial_s \mathcal{M}_\kappa(y) + (\ell_\kappa^2(y) + \omega^2) \partial_s \mathcal{N}_\kappa(y) \right], \end{aligned}$$

where $y \equiv q^2$, $\ell_\kappa(y) \equiv y\nu_\kappa + \mathcal{M}_\kappa(y)$, $k_\kappa(y) \equiv \mathcal{D}_\kappa - \frac{1}{2}\mathcal{N}_\kappa(y)$. We assumed the cut-off functions \mathcal{M}_κ , \mathcal{N}_κ to depend only on $y = q^2$, and we omitted terms which are odd in ω since they vanish upon integration. The frequency integrals can be explicitly carried out, yielding

$$\begin{aligned} \partial_s \nu_\kappa &= \frac{\lambda_\kappa^2}{4d} \int \frac{d^d \mathbf{q}}{(2\pi)^d} \frac{y}{\ell_\kappa^3(y)} \left[\left((d-2)k_\kappa(y) + 2y\partial_y k_\kappa(y) \right) \partial_s \mathcal{M}_\kappa(y) + \left(-\ell_\kappa(y) + y\partial_y \ell_\kappa(y) \right) \partial_s \mathcal{N}_\kappa(y) \right], \\ \partial_s \mathcal{D}_\kappa &= -\frac{\lambda_\kappa^2}{4} \int \frac{d^d \mathbf{q}}{(2\pi)^d} \frac{y^2 k_\kappa(y)}{\ell_\kappa^4(y)} \left[3k_\kappa(y) \partial_s \mathcal{M}_\kappa(y) + \ell_\kappa(y) \partial_s \mathcal{N}_\kappa(y) \right]. \end{aligned}$$

We then introduce dimensionless variables, and write the cut-off functions in the form $\mathcal{M}_\kappa(y) = \nu_\kappa y \hat{m}(\hat{y})$ and $\mathcal{N}_\kappa(y) = -2\mathcal{D}_\kappa \hat{n}(\hat{y})$ with $\hat{y} = y/\kappa^2$, such that one has

$$\begin{aligned} \partial_s \mathcal{M}_\kappa(y) &= -\nu_\kappa y (\eta_\kappa^\nu \hat{m}(\hat{y}) + 2\hat{y} \hat{m}'(\hat{y})) \equiv -\nu_\kappa y s_{\mathcal{M}}(\hat{y}) \\ \partial_s \mathcal{N}_\kappa(y) &= 2\mathcal{D}_\kappa (\eta_\kappa^{\mathcal{D}} \hat{n}(\hat{y}) + 2\hat{y} \hat{n}'(\hat{y})) \equiv 2\mathcal{D}_\kappa s_{\mathcal{N}}(\hat{y}). \end{aligned} \quad (C2)$$

The dimensionless flows of the running anomalous dimensions, defined as $\eta_\kappa^\nu = -\partial_s \ln \nu_\kappa$ and $\eta_\kappa^{\mathcal{D}} = -\partial_s \ln \mathcal{D}_\kappa$, are given by

$$\begin{aligned} \eta_\kappa^\nu &= \hat{g}_\kappa \frac{\nu_d}{8d} \int_0^\infty d\hat{y} \frac{\hat{y}^{d/2-2}}{(1+\hat{m}(\hat{y}))^3} \left[s_{\mathcal{M}}(\hat{y}) \left((d-2)(1+\hat{n}(\hat{y})) + 2\hat{y} \hat{n}'(\hat{y}) \right) - 2s_{\mathcal{N}}(\hat{y}) \hat{y} \hat{m}'(\hat{y}) \right], \\ \eta_\kappa^{\mathcal{D}} &= -\hat{g}_\kappa \frac{\nu_d}{8} \int_0^\infty d\hat{y} \frac{\hat{y}^{d/2-2}}{(1+\hat{m}(\hat{y}))^4} \left[(1+\hat{n}(\hat{y})) \left(3(1+\hat{n}(\hat{y})) s_{\mathcal{M}}(\hat{y}) - 2(1+\hat{m}(\hat{y})) s_{\mathcal{N}}(\hat{y}) \right) \right], \end{aligned} \quad (C3)$$

with $\hat{n}'(\hat{y}) \equiv \partial_{\hat{y}}\hat{n}(\hat{y})$ and $\hat{m}'(\hat{y}) \equiv \partial_{\hat{y}}\hat{m}(\hat{y})$, and where we defined $v_d = (2^{d-1}\pi^{d/2}\Gamma(d/2))^{-1}$. Solving this linear system for η_κ^ν and $\eta_\kappa^{\mathcal{D}}$ then yields

$$\begin{aligned}\eta_\kappa^\nu &= \frac{\mathcal{I}_0^\nu - \mathcal{I}_{\mathcal{D}}^\nu \mathcal{I}_0^\nu + \mathcal{I}_0^\nu \mathcal{I}_{\mathcal{D}}^\nu}{1 - \mathcal{I}_{\mathcal{D}}^\nu - \mathcal{I}_\nu^\nu - \mathcal{I}_\nu^\nu \mathcal{I}_{\mathcal{D}}^\nu + \mathcal{I}_{\mathcal{D}}^\nu \mathcal{I}_\nu^\nu}, \\ \eta_\kappa^{\mathcal{D}} &= \frac{\mathcal{I}_0^{\mathcal{D}} - \mathcal{I}_\nu^{\mathcal{D}} \mathcal{I}_0^{\mathcal{D}} + \mathcal{I}_0^{\mathcal{D}} \mathcal{I}_\nu^{\mathcal{D}}}{1 - \mathcal{I}_{\mathcal{D}}^{\mathcal{D}} - \mathcal{I}_\nu^{\mathcal{D}} - \mathcal{I}_\nu^{\mathcal{D}} \mathcal{I}_{\mathcal{D}}^{\mathcal{D}} + \mathcal{I}_{\mathcal{D}}^{\mathcal{D}} \mathcal{I}_\nu^{\mathcal{D}}},\end{aligned}\tag{C4}$$

where

$$\begin{aligned}\mathcal{I}_0^\nu &\equiv \hat{g}_\kappa \frac{v_d}{4d} \int_0^\infty d\hat{y} \hat{y}^{d/2-1} \frac{(d-2)(1+\hat{n}(\hat{y}))\hat{m}'(\hat{y})}{(1+\hat{m}(\hat{y}))^3}, \\ \mathcal{I}_\nu^\nu &\equiv \hat{g}_\kappa \frac{v_d}{8d} \int_0^\infty d\hat{y} \hat{y}^{d/2-2} \frac{[(d-2)(1+\hat{n}(\hat{y})) + 2\hat{y}\hat{n}'(\hat{y})]\hat{m}(\hat{y})}{(1+\hat{m}(\hat{y}))^3}, \\ \mathcal{I}_\nu^{\mathcal{D}} &\equiv -\hat{g}_\kappa \frac{v_d}{4d} \int_0^\infty d\hat{y} \hat{y}^{d/2-1} \frac{\hat{n}(\hat{y})\hat{m}'(\hat{y})}{(1+\hat{m}(\hat{y}))^3}, \\ \mathcal{I}_0^{\mathcal{D}} &\equiv -\hat{g}_\kappa \frac{v_d}{4} \int_0^\infty d\hat{y} \hat{y}^{d/2-1} \frac{(1+\hat{n}(\hat{y}))}{(1+\hat{m}(\hat{y}))^4} \left[3(1+\hat{n}(\hat{y}))\hat{m}'(\hat{y}) - 2(1+\hat{m}(\hat{y}))\hat{n}'(\hat{y}) \right], \\ \mathcal{I}_\nu^{\mathcal{D}} &\equiv -\hat{g}_\kappa \frac{3v_d}{8} \int_0^\infty d\hat{y} \hat{y}^{d/2-2} \frac{(1+\hat{n}(\hat{y}))^2\hat{m}(\hat{y})}{(1+\hat{m}(\hat{y}))^4}, \\ \mathcal{I}_{\mathcal{D}}^{\mathcal{D}} &\equiv \hat{g}_\kappa \frac{v_d}{4} \int_0^\infty d\hat{y} \hat{y}^{d/2-2} \frac{(1+\hat{n}(\hat{y}))\hat{n}(\hat{y})}{(1+\hat{m}(\hat{y}))^3}.\end{aligned}$$

It is clear that the integrals \mathcal{I}_0^ν , \mathcal{I}_ν^ν , $\mathcal{I}_\nu^{\mathcal{D}}$, $\mathcal{I}_0^{\mathcal{D}}$, $\mathcal{I}_\nu^{\mathcal{D}}$, $\mathcal{I}_{\mathcal{D}}^{\mathcal{D}}$ strongly depend on the choice of the cutoff functions $\hat{m}(y)$ and $\hat{n}(y)$. Since with this simple approximation we are only aiming at a qualitative picture of the phase diagram, we conveniently choose the regulator $\hat{m}(y) = \hat{n}(y) = (1/y - 1)\theta(1 - y)$, which allows one to analytically perform the momentum integrals. Defining $\tilde{g}_\kappa = v_d \hat{g}_\kappa$ One obtains

$$\begin{aligned}\eta_\kappa^\nu &= -\frac{(d+2)\tilde{g}_\kappa (d^3 - d(\tilde{g}_\kappa + 4) + \tilde{g}_\kappa)}{2d^5 + 8d^4 + d^3(8 - 3\tilde{g}_\kappa) - 2d^2\tilde{g}_\kappa + d\tilde{g}_\kappa(\tilde{g}_\kappa + 8) - \tilde{g}_\kappa^2}, \\ \eta_\kappa^{\mathcal{D}} &= \frac{(d+2)\tilde{g}_\kappa (d^3 + 2d^2 + \tilde{g}_\kappa(d-1))}{2d^5 + 8d^4 + d^3(8 - 3\tilde{g}_\kappa) - 2d^2\tilde{g}_\kappa + d\tilde{g}_\kappa(\tilde{g}_\kappa + 8) - \tilde{g}_\kappa^2}.\end{aligned}\tag{C5}$$

Appendix D: Calculation of the FRG flow equations in the large momentum limit

We consider the exact flow equations for the two-point functions $\bar{\Gamma}_\kappa^{(1,1)}(\varpi, \mathbf{p})$ and $\bar{\Gamma}_\kappa^{(0,2)}(\varpi, \mathbf{p})$, given by (24), and represented diagrammatically in Fig. 1 and 2. These equations can be closed (expressed in terms of two-point functions only) in the limit of large momentum $p = |\mathbf{p}|$, using the Ward identities. Indeed, as explained in Sec. V, in the large- p limit, one can set $\mathbf{q} = 0$ in all the vertices, where \mathbf{q} is the internal momentum, circulating in the loop. If this momentum is carried by a $\bar{\mathbf{u}}$ leg (represented by an outgoing arrow), then the corresponding vertex is zero because of the Ward identity (13), and the whole diagram vanishes. This occurs for the diagrams (b), (d) and (e) both in Fig. 1 and in Fig. 2, which can hence be neglected in the large p limit.

To evaluate the remaining diagrams, one first distribute the $\tilde{\partial}_s$ operator. This yields either a $\tilde{\partial}_s \bar{G}_{\kappa,ij}^{(2,0)}(\mathbf{Q})$ or a $\tilde{\partial}_s \bar{G}_{\kappa,ij}^{(1,1)}(\mathbf{Q})$, where $\mathbf{Q} = \pm \mathbf{q}$ or $\mathbf{Q} = \pm(\mathbf{p} + \mathbf{q})$. When $\mathbf{Q} = \pm(\mathbf{p} + \mathbf{q})$, one can change variables in the integrals to $\mathbf{q}' = \mp(\mathbf{p} + \mathbf{q})$ (and $\omega' = \mp(\varpi + \omega)$) such that it is always the momentum \mathbf{q} which is cutoff in the loop. After this operation, some terms turn out to vanish again because of (13) since the \mathbf{q} momentum appears to be carried by a $\bar{\mathbf{u}}$ leg. For the remaining non-zero terms, the $\mathbf{q} = 0$ momentum is carried by a \mathbf{u} leg (represented by an ingoing arrow). In fact, the corresponding diagrams always have the $\tilde{\partial}$ operator acting on a line with a black dot ($\bar{G}^{(2,0)}$ line). The vertices in these diagrams can be expressed in terms of two-point functions using the Ward identity (15) related to the extended Galilean symmetry, as we detail below.

1. Flow equation of $\Gamma_{\kappa,\parallel}^{(1,1)}(\varpi, \mathbf{p})$ in the large p limit

The diagram (a) of Fig. 2 can be expressed as

$$\begin{aligned}
\left[\partial_s \bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi, p) \right]_{(a)} &= \frac{1}{2} P_{\alpha\beta}^{\parallel}(\mathbf{p}) \int_{\omega, \mathbf{q}} \bar{\Gamma}_{\kappa,ij\alpha\beta}^{(3,1)}(\omega, \mathbf{q}, -\omega, -\mathbf{q}, \varpi, \mathbf{p}) \tilde{\partial}_s \bar{G}_{\kappa,ji}^{(2,0)}(\omega, \mathbf{q}) \\
&\stackrel{p \rightarrow \infty}{=} \frac{1}{2} P_{\alpha\beta}^{\parallel}(\mathbf{p}) \int_{\omega, \mathbf{q}} \bar{\Gamma}_{\kappa,ij\alpha\beta}^{(3,1)}(\omega, 0, -\omega, 0, \varpi, \mathbf{p}) \tilde{\partial}_s \bar{G}_{\kappa,ji}^{(2,0)}(\omega, \mathbf{q}) \\
&= \frac{1}{2} \frac{p^2}{d} \int_{\omega} \frac{1}{\omega^2} \left[\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\omega + \varpi, \mathbf{p}) - 2\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi, \mathbf{p}) + \bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(-\omega + \varpi, \mathbf{p}) \right] \tilde{\partial}_s \int_{\mathbf{q}} \bar{G}_{\kappa,\parallel}^{(2,0)}(\omega, \mathbf{q}), \quad (D1)
\end{aligned}$$

where the large p limit is taken in the second line, and the identity (15) is used twice to obtain the third line. The non-zero contribution of diagram (c) can be written as

$$\begin{aligned}
\left[\partial_s \bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi, \mathbf{p}) \right]_{(c)} &= -P_{\alpha\beta}^{\parallel}(\mathbf{p}) \int_{\omega, \mathbf{q}} \bar{\Gamma}_{\kappa,\alpha ij}^{(2,1)}(\varpi, \mathbf{p}, \omega, \mathbf{q}) \bar{G}_{\kappa,jk}^{(1,1)}(\omega + \varpi, \mathbf{p} + \mathbf{q}) \bar{\Gamma}_{\kappa,kl\beta}^{(2,1)}(\omega + \varpi, \mathbf{p} + \mathbf{q}, -\omega, -\mathbf{q}) \tilde{\partial}_s \bar{G}_{\kappa,li}^{(2,0)}(\omega, \mathbf{q}) \\
&\stackrel{p \rightarrow \infty}{=} -\frac{p^2}{d} \int_{\omega} \left[\frac{\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\omega + \varpi, \mathbf{p}) - \bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi, \mathbf{p})}{\omega} \right]^2 \bar{G}_{\kappa,\parallel}^{(1,1)}(\omega + \varpi, \mathbf{p}) \tilde{\partial}_s \int_{\mathbf{q}} \bar{G}_{\kappa,\parallel}^{(2,0)}(\omega, \mathbf{q}), \quad (D2)
\end{aligned}$$

where the three-point vertices in the first line are evaluated in their large p limit (setting $\mathbf{q} = 0$) and expressed using the identity (15). Summing up the two contributions (D1) and (D2), one obtains

$$\begin{aligned}
\partial_s \bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi, \mathbf{p}) &= \frac{p^2}{d} \int_{\omega} \left\{ - \left[\frac{\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\omega + \varpi, \mathbf{p}) - \bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi, \mathbf{p})}{\omega} \right]^2 \bar{G}_{\kappa,\parallel}^{(1,1)}(\omega + \varpi, \mathbf{p}) \right. \\
&\quad \left. + \frac{1}{2\omega^2} \left[\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi + \omega, \mathbf{p}) - 2\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi, \mathbf{p}) + \bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi - \omega, \mathbf{p}) \right] \right\} \times \tilde{\partial}_s \int_{\mathbf{q}} \bar{G}_{\kappa,\parallel}^{(2,0)}(\omega, \mathbf{q}). \quad (D3)
\end{aligned}$$

2. Flow equation of $\Gamma_{\kappa,\parallel}^{(0,2)}(\varpi, \mathbf{p})$ in the large p limit

Similarly, the diagram (a) of Fig. 1 can be expressed, in the large p limit and using twice (15), as

$$\begin{aligned}
\left[\partial_s \bar{\Gamma}_{\kappa,\parallel}^{(0,2)}(\varpi, \mathbf{p}) \right]_{(a)} &= \frac{1}{2} P_{\alpha\beta}^{\parallel}(\mathbf{p}) \int_{\omega} \bar{\Gamma}_{\kappa,ij\alpha\beta}^{(2,2)}(\omega, \mathbf{q}, -\omega, -\mathbf{q}, \varpi, \mathbf{p}) \int_{\omega, \mathbf{q}} \tilde{\partial}_s \bar{G}_{\kappa,ji}^{(2,0)}(\omega, \mathbf{q}) \\
&\stackrel{p \rightarrow \infty}{=} \frac{1}{2} \frac{p^2}{d} \int_{\omega} \frac{1}{\omega^2} \left[\bar{\Gamma}_{\kappa,\parallel}^{(0,2)}(\omega + \varpi, \mathbf{p}) - 2\bar{\Gamma}_{\kappa,\parallel}^{(0,2)}(\varpi, \mathbf{p}) + \bar{\Gamma}_{\kappa,\parallel}^{(0,2)}(-\omega + \varpi, \mathbf{p}) \right] \tilde{\partial}_s \int_{\mathbf{q}} \bar{G}_{\kappa,\parallel}^{(2,0)}(\omega, \mathbf{q}). \quad (D4)
\end{aligned}$$

The expression for the diagram (c) is obtained using (15) for the two vertices, and reads as

$$\begin{aligned}
\left[\partial_s \bar{\Gamma}_{\kappa,\parallel}^{(0,2)}(\varpi, \mathbf{p}) \right]_{(c)} &= -P_{\alpha\beta}^{\parallel}(\mathbf{p}) \int_{\omega, \mathbf{q}} \bar{\Gamma}_{\kappa,ij\alpha}^{(2,1)}(\omega, \mathbf{q}, -\omega - \varpi, -\mathbf{p} - \mathbf{q}) \bar{G}_{\kappa,jk}^{(2,0)}(\omega + \varpi, \mathbf{p} + \mathbf{q}) \\
&\quad \times \bar{\Gamma}_{\kappa,kl\beta}^{(2,1)}(\omega + \varpi, \mathbf{p} + \mathbf{q}, -\omega, -\mathbf{q}) \tilde{\partial}_s \bar{G}_{\kappa,li}^{(2,0)}(\omega, \mathbf{q}) \\
&\stackrel{p \rightarrow \infty}{=} -\frac{p^2}{d} \int_{\omega} \frac{1}{\omega^2} \left(\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(-\varpi, \mathbf{p}) - \bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(-\varpi - \omega, \mathbf{p}) \right) \\
&\quad \times \left(\bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi, \mathbf{p}) - \bar{\Gamma}_{\kappa,\parallel}^{(1,1)}(\varpi + \omega, \mathbf{p}) \right) \bar{G}_{\kappa,\parallel}^{(2,0)}(\omega + \varpi, \mathbf{p}) \tilde{\partial}_s \int_{\mathbf{q}} \bar{G}_{\kappa,\parallel}^{(2,0)}(\omega, \mathbf{q}). \quad (D5)
\end{aligned}$$

Finally, the expression for the diagram (f) is obtained in the same way yielding

$$\begin{aligned}
\left[\partial_s \bar{\Gamma}_{\kappa, \parallel}^{(0,2)}(\varpi, \mathbf{p}) \right]_{(f)} &= -P_{\alpha\beta}^{\parallel}(\mathbf{p}) \int_{\omega, \mathbf{q}} \bar{\Gamma}_{\kappa, ij\alpha}^{(1,2)}(\omega, \mathbf{q}, -\omega - \varpi, -\mathbf{p} - \mathbf{q}) \bar{G}_{\kappa, jk}^{(1,1)}(\omega + \varpi, \mathbf{p} + \mathbf{q}) \\
&\quad \times \bar{\Gamma}_{\kappa, kl\beta}^{(2,1)}(\omega + \varpi, \mathbf{p} + \mathbf{q}, -\omega, -\mathbf{q}) \tilde{\partial}_s \bar{G}_{\kappa, li}^{(2,0)}(\omega, \mathbf{q}) + c.c. \\
&\stackrel{p \rightarrow \infty}{=} -\frac{p^2}{d} \int_{\omega} \left[\frac{\bar{\Gamma}_{\kappa, \parallel}^{(0,2)}(\omega + \varpi, \mathbf{p}) - \bar{\Gamma}_{\kappa, \parallel}^{(0,2)}(\varpi, \mathbf{p})}{\omega} \right] \times \left[\frac{\bar{\Gamma}_{\kappa, \parallel}^{(1,1)}(\omega + \varpi, \mathbf{p}) - \bar{\Gamma}_{\kappa, \parallel}^{(1,1)}(\varpi, \mathbf{p})}{\omega} \right] \\
&\quad \times \bar{G}_{\kappa, \parallel}^{(1,1)}(\omega + \varpi, \mathbf{p}) \tilde{\partial}_s \int_{\mathbf{q}} \bar{G}_{\kappa, \parallel}^{(2,0)}(\omega, \mathbf{q}) + c.c., \tag{D6}
\end{aligned}$$

where $c.c.$ denotes the complex conjugate. Adding the three contributions (D4), (D5) and (D6) finally leads to

$$\begin{aligned}
\partial_s \bar{\Gamma}_{\kappa, \parallel}^{(0,2)}(\varpi, \mathbf{p}) &= \frac{p^2}{d} \int_{\omega} \left\{ - \left| \frac{\bar{\Gamma}_{\kappa, \parallel}^{(1,1)}(\varpi, \mathbf{p}) - \bar{\Gamma}_{\kappa, \parallel}^{(1,1)}(\varpi + \omega, \mathbf{p})}{\omega} \right|^2 \bar{G}_{\kappa, \parallel}^{(2,0)}(\omega + \varpi, \mathbf{p}) \right. \\
&\quad \left. - 2 \left[\frac{\bar{\Gamma}_{\kappa, \parallel}^{(0,2)}(\omega + \varpi, \mathbf{p}) - \bar{\Gamma}_{\kappa, \parallel}^{(0,2)}(\varpi, \mathbf{p})}{\omega} \right] \times \Re \left\{ \left[\frac{\bar{\Gamma}_{\kappa, \parallel}^{(1,1)}(\omega + \varpi, \mathbf{p}) - \bar{\Gamma}_{\kappa, \parallel}^{(1,1)}(\varpi, \mathbf{p})}{\omega} \right] \bar{G}_{\kappa, \parallel}^{(1,1)}(\omega + \varpi, \mathbf{p}) \right\} \right. \\
&\quad \left. + \frac{1}{2\omega^2} \left[\bar{\Gamma}_{\kappa, \parallel}^{(0,2)}(\omega + \varpi, \mathbf{p}) - 2\bar{\Gamma}_{\kappa, \parallel}^{(0,2)}(\varpi, \mathbf{p}) + \bar{\Gamma}_{\kappa, \parallel}^{(0,2)}(-\omega + \varpi, \mathbf{p}) \right] \right\} \times \tilde{\partial}_s \int_{\mathbf{q}} \bar{G}_{\kappa, \parallel}^{(2,0)}(\omega, \mathbf{q}). \tag{D7}
\end{aligned}$$

The two flow equations (D3) and (D7) for the two-point functions are thus closed in the limit of large p . However, they still bare a complicated nonlinear form. It turns out that they endow a much simpler form when expressed for the correlation functions $\bar{G}_{\kappa}^{(2,0)}$ and $\bar{G}_{\kappa}^{(1,1)}$ rather than for the vertex functions $\bar{\Gamma}_{\kappa}^{(0,2)}$ and $\bar{\Gamma}_{\kappa}^{(1,1)}$ [23, 27]. Indeed, using the definitions (B1) and (B2), one can calculate the flow equations for $\bar{G}_{\kappa}^{(2,0)}$ and $\bar{G}_{\kappa}^{(1,1)}$ from (D3) and (D7), and one finds, using parity and after some algebra

$$\partial_s \bar{G}_{\kappa, \parallel}^{(2,0)}(\varpi, \mathbf{p}) = \frac{p^2}{d} \int_{\omega} \frac{1}{2\omega^2} \left[\bar{G}_{\kappa, \parallel}^{(2,0)}(\omega + \varpi, \mathbf{p}) - 2\bar{G}_{\kappa, \parallel}^{(2,0)}(\varpi, \mathbf{p}) + \bar{G}_{\kappa, \parallel}^{(2,0)}(-\omega + \varpi, \mathbf{p}) \right] \tilde{\partial}_s \int_{\mathbf{q}} \bar{G}_{\kappa, \parallel}^{(2,0)}(\omega, \mathbf{q}) \tag{D8}$$

$$\partial_s \bar{G}_{\kappa, \parallel}^{(1,1)}(\varpi, \mathbf{p}) = \frac{p^2}{d} \int_{\omega} \frac{1}{2\omega^2} \left[\bar{G}_{\kappa, \parallel}^{(1,1)}(\omega + \varpi, \mathbf{p}) - 2\bar{G}_{\kappa, \parallel}^{(1,1)}(\varpi, \mathbf{p}) + \bar{G}_{\kappa, \parallel}^{(1,1)}(-\omega + \varpi, \mathbf{p}) \right] \tilde{\partial}_s \int_{\mathbf{q}} \bar{G}_{\kappa, \parallel}^{(2,0)}(\omega, \mathbf{q}). \tag{D9}$$

-
- [1] M. Kardar, G. Parisi, and Y.-C. Zhang, Dynamic scaling of growing interfaces, *Phys. Rev. Lett.* **56**, 889 (1986).
 - [2] D. Wei, A. Rubio-Abadal, B. Ye, F. Machado, J. Kemp, K. Srakaew, S. Hollerith, J. Rui, S. Gopalakrishnan, N. Y. Yao, I. Bloch, and J. Zeiher, Quantum gas microscopy of Kardar-Parisi-Zhang superdiffusion, *Science* **376**, 716 (2022).
 - [3] A. Scheie, N. E. Sherman, M. Dupont, S. E. Nagler, M. B. Stone, G. E. Granroth, J. E. Moore, and D. A. Tennant, Detection of Kardar-Parisi-Zhang hydrodynamics in a quantum Heisenberg spin-1/2 chain, *Nature Physics* **17**, 726 (2022).
 - [4] Q. Fontaine, D. Squizzato, F. Baboux, I. Amelio, A. Lemaître, M. Morassi, I. Sagnes, L. Le Gratiet, A. Harouri, M. Wouters, I. Carusotto, A. Amo, M. Richard, A. Minguzzi, L. Canet, S. Ravets, and J. Bloch, Kardar-Parisi-Zhang universality in a one-dimensional polariton condensate, *Nature* **608**, 687 (2022).
 - [5] A. Nahum, J. Ruhman, S. Vijay, and J. Haah, Quantum entanglement growth under random unitary dynamics, *Phys. Rev. X* **7**, 031016 (2017).
 - [6] S. Mu, J. Gong, and G. Lemarié, Kardar-parisi-zhang physics in the density fluctuations of localized two-dimensional wave packets, *Phys. Rev. Lett.* **132**, 046301 (2024).
 - [7] T. Halpin-Healy and Y.-C. Zhang, Kinetic roughening phenomena, stochastic growth, directed polymers and all that. Aspects of multidisciplinary statistical mechanics, *Phys. Rep.* **254**, 215 (1995).
 - [8] J. Krug, Origins of scale invariance in growth processes, *Adv. Phys.* **46**, 139 (1997).
 - [9] J. Burgers, A mathematical model illustrating the theory of turbulence (Elsevier, 1948) pp. 171–199.
 - [10] M. Kardar, Replica Bethe ansatz studies of two-dimensional interfaces with quenched random impurities,

- Nucl. Phys. B **290**, 582 (1987).
- [11] D. Forster, D. R. Nelson, and M. J. Stephen, Long-time tails and the large-eddy behavior of a randomly stirred fluid, *Phys. Rev. Lett.* **36**, 867 (1976).
- [12] D. Forster, D. R. Nelson, and M. J. Stephen, Large-distance and long-time properties of a randomly stirred fluid, *Phys. Rev. A* **16**, 732 (1977).
- [13] S. F. Edwards and D. R. Wilkinson, The Surface Statistics of a Granular Aggregate, *Proc. R. Soc. Lon. A* **381**, 17 (1982).
- [14] M. K. Verma, Intermittency exponents and energy spectrum of the Burgers and KPZ equations with correlated noise, *Physica A* **277**, 359 (2000).
- [15] J. Bec and K. Khanin, Burgers turbulence, *Physics Reports* **447**, 1 (2007).
- [16] J. P. Bouchaud, M. Mézard, and G. Parisi, Scaling and intermittency in burgers turbulence, *Phys. Rev. E* **52**, 3656 (1995).
- [17] C. Cartes, E. Tirapegui, R. Pandit, and M. Brachet, The Galerkin-truncated Burgers equation: crossover from inviscid-thermalized to Kardar–Parisi–Zhang scaling, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **380**, 20210090 (2022).
- [18] A. J. Majda and I. Timofeyev, Remarkable statistical behavior for truncated Burgers–Hopf dynamics, *Proceedings of the National Academy of Sciences* **97**, 12413 (2000).
- [19] E. Rodríguez-Fernández, S. N. Santalla, M. Castro, and R. Cuerno, Anomalous ballistic scaling in the tensionless or inviscid Kardar–Parisi–Zhang equation, *Phys. Rev. E* **106**, 024802 (2022).
- [20] K. Fujimoto, R. Hamazaki, and Y. Kawaguchi, Family-Vicsek scaling of roughness growth in a strongly interacting bose gas, *Phys. Rev. Lett.* **124**, 210604 (2020).
- [21] C. Fontaine, F. Vercesi, M. Brachet, and L. Canet, Unpredicted scaling of the one-dimensional kardar-parisi-zhang equation, *Phys. Rev. Lett.* **131**, 247101 (2023).
- [22] L. Canet, V. Rossetto, N. Wschebor, and G. Balarac, Spatiotemporal velocity-velocity correlation function in fully developed turbulence, *Phys. Rev. E* **95**, 023107 (2017).
- [23] M. Tarpin, L. Canet, and N. Wschebor, Breaking of scale invariance in the time dependence of correlation functions in isotropic and homogeneous turbulence, *Physics of Fluids* **30**, 055102 (2018).
- [24] P. C. Martin, E. D. Siggia, and H. A. Rose, Statistical dynamics of classical systems, *Phys. Rev. A* **8**, 423 (1973).
- [25] H.-K. Janssen, On a Lagrangean for classical field dynamics and Renormalization Group calculations of dynamical critical properties, *Z. Phys. B* **23**, 377 (1976).
- [26] C. de Dominicis, Techniques de renormalisation de la théorie des champs et dynamique des phénomènes critiques, *J. Phys. (Paris) Colloq.* **37**, 247 (1976).
- [27] L. Canet, B. Delamotte, and N. Wschebor, Fully developed isotropic turbulence: Nonperturbative renormalization group formalism and fixed-point solution, *Phys. Rev. E* **93**, 063101 (2016).
- [28] L. Canet, Functional renormalisation group for turbulence, *Journal of Fluid Mechanics* **950**, P1 (2022).
- [29] L. Canet, B. Delamotte, and N. Wschebor, Fully developed isotropic turbulence: Symmetries and exact identities, *Phys. Rev. E* **91**, 053004 (2015).
- [30] L. Canet, H. Chaté, and B. Delamotte, General framework of the non-perturbative Renormalization Group for non-equilibrium steady states, *J. Phys. A* **44**, 495001 (2011).
- [31] J. Berges, N. Tetradis, and C. Wetterich, Non-perturbative Renormalization flow in quantum field theory and statistical physics, *Phys. Rep.* **363**, 223 (2002).
- [32] P. Kopietz, L. Bartosch, and F. Schütz, *Introduction to the Functional Renormalization Group*, *Lecture Notes in Physics*, Vol. 798 (Springer, Berlin, 2010).
- [33] B. Delamotte, An Introduction to the nonperturbative renormalization group, *Lect. Notes Phys.* **852**, 49 (2012).
- [34] N. Dupuis, L. Canet, A. Eichhorn, W. Metzner, J. Pawłowski, M. Tissier, and N. Wschebor, The nonperturbative functional renormalization group and its applications, *Physics Reports* **910**, 1 (2021), the nonperturbative functional renormalization group and its applications.
- [35] C. Wetterich, Exact evolution equation for the effective potential, *Phys. Lett. B* **301**, 90 (1993).
- [36] U. Ellwanger, Flow equations and BRS invariance for Yang-Mills theories, *Physics Letters B* **335**, 364 (1994).
- [37] T. R. Morris, The exact renormalization group and approximate solutions, *International Journal of Modern Physics A* **09**, 2411 (1994).
- [38] F. Benitez and N. Wschebor, Branching and annihilating random walks: Exact results at low branching rate, *Phys. Rev. E* **87**, 052132 (2013).
- [39] L. Canet, H. Chaté, B. Delamotte, I. Dornic, and M. A. Muñoz, Nonperturbative fixed point in a nonequilibrium phase transition, *Phys. Rev. Lett.* **95**, 100601 (2005).
- [40] L. Canet, H. Chaté, B. Delamotte, and N. Wschebor, Nonperturbative renormalization group for the Kardar–Parisi–Zhang equation, *Phys. Rev. Lett.* **104**, 150601 (2010).
- [41] L. Canet, H. Chaté, B. Delamotte, and N. Wschebor, Nonperturbative Renormalization Group for the Kardar–Parisi–Zhang equation: General framework and first applications, *Phys. Rev. E* **84**, 061128 (2011).
- [42] T. Kloss, L. Canet, and N. Wschebor, Nonperturbative Renormalization Group for the stationary Kardar–Parisi–Zhang equation: Scaling functions and amplitude ratios in 1+1, 2+1, and 3+1 dimensions, *Phys. Rev. E* **86**, 051124 (2012).
- [43] E. Frey and U. C. Täuber, Two-loop Renormalization-Group analysis of the Burgers–Kardar–Parisi–Zhang equation, *Phys. Rev. E* **50**, 1024 (1994).
- [44] K. J. Wiese, On the perturbation expansion of the KPZ equation, *J. Stat. Phys.* **93**, 143 (1998).
- [45] Again, the statement that the system is already at the KPZ fixed point for $g_\Lambda = g_*$ only holds within the simplest approximation (30) where Γ_κ is restricted to the form of the bare action. In general, the fixed point effective action is more complicated, and g_* is defined as the initial value g_Λ which minimizes the RG time to reach the IR fixed point.
- [46] J.-P. Blaizot, R. Méndez-Galain, and N. Wschebor, A new method to solve the non-perturbative Renormalization Group equations, *Phys. Lett. B* **632**, 571 (2006).
- [47] J.-P. Blaizot, R. Méndez-Galain, and N. Wschebor, Non-perturbative Renormalization Group calculation of the scalar self-energy, *Eur. Phys. J. B* **58**, 297 (2007).
- [48] F. Benitez, J.-P. Blaizot, H. Chaté, B. Delamotte, R. Méndez-Galain, and N. Wschebor, Nonperturbative

- Renormalization Group preserving full-momentum dependence: Implementation and quantitative evaluation, *Phys. Rev. E* **85**, 026707 (2012).
- [49] M. Tarpin, L. Canet, C. Pagani, and N. Wschebor, Stationary, isotropic and homogeneous two-dimensional turbulence: a first non-perturbative renormalization group approach, *Journal of Physics A: Mathematical and Theoretical* **52**, 085501 (2019).
- [50] C. Pagani and L. Canet, Spatio-temporal correlation functions in scalar turbulence from functional renormalization group, *Physics of Fluids* **33**, 065109 (2021).
- [51] A. Gorbunova, G. Balarac, L. Canet, G. Eyink, and V. Rossetto, Spatio-temporal correlations in three-dimensional homogeneous and isotropic turbulence, *Physics of Fluids* **33**, 045114 (2021).
- [52] A. Gorbunova, C. Pagani, G. Balarac, L. Canet, and V. Rossetto, Eulerian spatiotemporal correlations in passive scalar turbulence, *Phys. Rev. Fluids* **6**, 124606 (2021).