

Estimating Time-Varying Parameters of Various Smoothness in Linear Models via Kernel Regression

Mikihito Nishi*¹

¹Graduate School of Economics, Hitotsubashi University

June 21, 2024

Abstract

We consider estimating nonparametric time-varying parameters in linear models using kernel regression. Our contributions are twofold. First, We consider a broad class of time-varying parameters including deterministic smooth functions, the rescaled random walk, structural breaks, the threshold model and their mixtures. We show that those time-varying parameters can be consistently estimated by kernel regression. Our analysis exploits the smoothness of time-varying parameters rather than their specific form. The second contribution is to reveal that the bandwidth used in kernel regression determines the trade-off between the rate of convergence and the size of the class of time-varying parameters that can be estimated. An implication from our result is that the bandwidth should be proportional to $T^{-1/2}$ if the time-varying parameter follows the rescaled random walk, where T is the sample size. We propose a specific choice of the bandwidth that accommodates a wide range of time-varying parameter models. An empirical application shows that the kernel-based estimator with this choice can capture the random-walk dynamics in time-varying parameters.

Keywords: Bandwidth, kernel estimation, random walk, structural break, time-varying parameter

JEL Codes: C14, C22

*I am deeply grateful to Atsushi Inoue for his constructive comments on the earlier version of this article. All errors are mine. Address correspondence to: Graduate School of Economics, Hitotsubashi University, 2-1 Naka, Kunitachi, Tokyo 186-8601, Japan; e-mail: ed225007@g.hit-u.ac.jp

1 Introduction

In this article, we consider estimating the following linear model with time-varying coefficients:

$$y_t = x_t' \beta_{T,t} + \varepsilon_t, \quad t = 1, 2, \dots, T, \quad (1)$$

where T is the sample size, $p \times 1$ vector x_t is the regressor, $p \times 1$ triangular array $\beta_{T,t}$ is the time-varying coefficient, and ε_t is the disturbance. Unlike the usual constant-coefficient model where $\beta_{T,t} = \beta$ for all t and T , model (1) allows the coefficient to vary over time.

In the literature, time-varying parameters are often estimated via rolling regression (Cushman, 1988; Pivetta and Reis, 2007; Inoue, Jin and Rossi, 2017), or more generally, via kernel regression where observations are weighted by some kernel function. Starting from Robinson (1989), researchers have developed estimation and inferential theory for the kernel-based estimation of time-varying parameters; Cai (2007), Chen and Hong (2012), and Friedrich and Lin (2022) to name a few. In this article, we follow this strand of literature and consider estimating $\beta_{T,t}$ by using kernel regression.

Our contributions are twofold. First, we consider estimating a broader class of time-varying parameters than the class considered in the literature. The most common assumption adopted in the literature is that $\beta_{T,t}$ is a smooth function, and usually, smooth enough that it is continuously differentiable (e.g., Cai, 2007; Inoue et al., 2017). However, smooth functions are not the only model for parameter instability that is popular in economics and statistics. For instance, the (rescaled) random walk model is a popular one, in which $\beta_{T,t}$ is modeled as $\beta_{T,t} = (1/\sqrt{T}) \sum_{i=1}^t u_i$ with u_i being a transitory process (e.g., Nyblom, 1989; Cogley and Sargent, 2005). Another example is (abrupt) structural breaks in $\beta_{T,t}$ (Andrews, 1993; Bai and Perron, 1998). These two modeling schemes have received less attention in the literature on kernel-based estimation, and therefore it is largely unknown what the consequence is if one applies kernel regression to these types of time-varying parameters.^{1,2} In this article, we develop the kernel-based estimation theory accommodating a wide class of time-varying parameters, including smooth functions, the rescaled random walk, and struc-

¹Giraitis, Kapetanios and Yates (2014) and Giraitis, Kapetanios and Marcellino (2021) are among few exceptions. They show that random walk type parameters can be consistently estimated via the kernel-based method.

²Pesaran and Timmermann (2007) and Pesaran, Pick and Pranovich (2013) apply kernel-based approaches for random walk and structural break type parameter instabilities, but their focus is on optimal forecasting, rather than estimation of $\beta_{T,t}$.

tural breaks. Moreover, this class encompasses the threshold regression model proposed by Hansen (2000), which has rarely been considered in the context of kernel regression.

Let us emphasize that the class of time-varying parameters considered in this article also includes the mixtures of the aforementioned models. The relationship between y_t and x_t , for example, can smoothly change over time but may experience discontinuous changes at some points because of global financial crises or pandemics. If this is the case, it is reasonable to model $\beta_{T,t}$ as a smooth function with some discontinuous changes at some points. Our class accommodates such types of parameter instability. The literature has acknowledged the importance of taking into account several types of parameter instability. For instance, Müller and Petalas (2010) consider inference in models with time-varying parameters approximated by Gaussian processes and continuous functions possibly with finitely many jumps. Although their framework allows for nonlinear models and thus is more general than ours in this respect, they focus on small parameter instabilities (relative to those considered in this work). Therefore, large instabilities are not allowed in their model. Chen and Hong (2012) construct tests for smooth structural changes possibly with a finite number of discontinuous breaks. However, they do not provide a result for the estimation of time-varying parameters of this type. Kristensen (2012) proposes a nonparametric estimation method for time-varying coefficients by developing a framework that might allow for smooth functions, structural breaks, and the rescaled random walk. However, his analysis is restricted to smooth functional parameters only, and his results do not directly apply to the other specifications. Giraitis et al. (2021) develop an IV estimation method for a class of time-varying parameters, which includes smooth deterministic functions, the rescaled random walk, and their mixture. However, (large) structural breaks are not allowed in their model. Unlike these earlier studies, we employ a general framework that accommodates all the aforementioned models and their mixtures.

This general framework is developed by exploiting the smoothness of nonparametric time-varying parameter $\beta_{T,t}$ rather than its specific form. In this article, we quantify the smoothness of time-varying parameters by a single parameter $\alpha > 0$. Our definition of smoothness, roughly speaking, extends the Hölder condition for real-valued functions to the case where the function of interest (i.e., $\beta_{T,t}$) may be random. For example, we will show that continuously differentiable functions have smoothness of $\alpha = 1$, and the random walk divided by \sqrt{T} has $\alpha = 1/2$. As with the Hölder condition, a smaller α means (possibly) more roughness of the path of the time-varying parameter. This implies that the random walk divided by \sqrt{T} is

less smooth than continuously differentiable functions.

Our second contribution is to reveal the role of the bandwidth in the above general setting. Within a textbook framework, it is well-known that the bandwidth determines the bias-variance trade-off inherent in the kernel-based estimation. The result shown in this article goes beyond that: The bandwidth determines the trade-off between the rate of convergence and the size of the class of time-varying parameters that can be estimated. In other words, if one wants to estimate $\beta_{T,t}$ at a fast rate of convergence, only a tiny class of $\beta_{T,t}$ can be estimated via kernel regression, while they have to accept a slower rate of convergence if they are to estimate a broader class of time-varying parameters. This indicates that there is a trade-off between efficiency (in terms of the rate of convergence) and robustness (in terms of the size of the class of time-varying parameters that can be estimated) in kernel-based estimation. We will reveal that the bandwidth determines the balance between them.

Two implications are obtained from the above result. First, we will show that the conventional choice of a $T^{-1/5}$ -rate bandwidth, which is specialized to continuously differentiable time-varying parameters, is invalid if $\beta_{T,t}$ is less smooth (e.g. if $\beta_{T,t}$ is the random walk divided by \sqrt{T}). We will show that the bandwidth should be proportional to $T^{-1/2}$ if $\beta_{T,t}$ follows the rescaled random walk. Second, we will show that the kernel-based estimator is biased if the time-varying parameter has a break (discontinuity) of a certain magnitude. We quantify the degree of the bias and show that the bandwidth determines the magnitude that yields the bias.

Based on the aforementioned findings, we also discuss how to choose the bandwidth. Given the fact that various models for parameter instability have been proposed (e.g., smooth functions, the random walk, structural breaks and the threshold model), a robust kernel-based method that is universally applicable to a wide class of time-varying parameters will be valuable. However, as stated above and as is the case in every aspect of econometrics, pursuing robustness is necessarily accompanied by inefficiency. Therefore, how to balance them is a crucial issue in econometric applications, and this question reduces to how to choose the bandwidth in our context. We propose a specific choice of the bandwidth that leads to a robust estimation method accommodating many important time-varying parameters, including continuously differentiable functions, the rescaled random walk, small to moderate abrupt breaks and threshold effects, and their mixtures. Although this choice leads to a slower rate of convergence than that achieved by the conventional bandwidth choice, our empirical application demonstrates that the kernel-based estimation with our bandwidth choice can

capture random walk type parameter instabilities and thus is useful in many applications.

The remainder of this paper is organized as follows. In Section 2, we propose a definition of the smoothness of (possibly random) time-varying parameters and derive the asymptotic properties of the kernel-based estimator. In Section 3, we discuss the consequence of an improper choice of bandwidth and propose a specific choice to accommodate a wide variety of time-varying parameters. Section 4 conducts Monte Carlo experiments, and Section 5 gives an empirical application. Section 6 concludes the article. All mathematical proofs of the main results are relegated to Appendix A.

Notation: For any matrix A , $\|A\| = \text{tr}(A'A)^{1/2}$ denotes the Frobenius norm of A . For any positive number b , $\lfloor b \rfloor$ denotes the integer part of b . \xrightarrow{p} and \xrightarrow{d} signify convergence in probability and convergence in distribution as $T \rightarrow \infty$, respectively. \Rightarrow signifies weak convergence of the associated probability measures.

2 Assumptions and Main Result

2.1 Assumptions

Throughout the article, we impose the following assumption on model (1).³

Assumption 1.

(a) $\{(x'_t, \varepsilon_t)\}_t$ is L_2 -NED of size $-(r-1)/(r-2)$ on an α -mixing sequence of size $-r/(r-2)$ for some $r > 2$, with respect to some positive constants d_t satisfying $\sup_t d_t < \infty$. Moreover, $\sup_t E[\|x_t\|^{2r}] + \sup_t E[|\varepsilon_t|^{2r}] < \infty$.

(b) $\{x_t \varepsilon_t\}_t$ has mean zero and is serially uncorrelated.

(c) For each $t = \lfloor Tr \rfloor$, $r \in (0, 1)$, there exist nonrandom symmetric matrices $\Omega(r) > 0$ and $\Sigma(r) > 0$ such that $(1/Th) \sum_{i=1}^T K((t-i)/Th) x_i x'_i \xrightarrow{p} \Omega(r)$ and $\text{Var}\left(\frac{1}{\sqrt{Th}} \sum_{i=1}^T K((t-i)/Th) x_i \varepsilon_i\right) \rightarrow \Sigma(r)$.

Assumption 1(a) allows the regressor and disturbance to be weakly serially dependent. The NED assumption is more general than mixing conditions, which have been commonly assumed in the literature (Cai, 2007; Chen and Hong, 2012; Giraitis et al., 2021; Friedrich and Lin, 2022). Also note that we do not impose stationarity unlike earlier studies, and thus our

³For the definition of near epoch dependence (NED), see, e.g., Davidson (1994).

framework allows for heteroskedasticity in ε_t . Assumption 1(b) requires that the product of regressors and disturbance be serially uncorrelated, which is satisfied when, for example, x_t is $F_{T,t-1}$ -measurable, where $\mathcal{F}_{T,t} := \sigma(\{x'_s, \varepsilon_s, \beta_{T,s}\}_{s \leq t})$, and ε_t is a martingale difference sequence (m.d.s.). The assumption of no serial correlation or m.d.s. is common in the literature (Chen and Hong, 2012; Kristensen, 2012; Giraitis et al., 2021). Assumption 1(c) holds under Assumptions 1(a)-(b) if x_t and $x_t \varepsilon_t$ are covariance-stationary (see Corollary 1).

We consider estimating $\beta_{T,t}$ by using the local constant (Nadaraya-Watson) estimator:

$$\hat{\beta}_t := \left(\sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' \right)^{-1} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i y_i,$$

where $K(\cdot)$ is a kernel function and h is the bandwidth parameter satisfying $h \rightarrow 0$ and $Th \rightarrow \infty$ as $T \rightarrow \infty$. We suppose the kernel function $K(\cdot)$ satisfies the following condition:

Assumption 2.

- (a) $K(x) \geq 0$, $x \in \mathbb{R}$, is Lipschitz continuous and has compact support $[-1, 1]$.
- (b) $\int_{-1}^1 K(x) dx = 1$.

Commonly used kernels such as the flat kernel (uniform density on $[-1, 1]$) and Epanechnikov kernel satisfy Assumption 2. Following the arguments of Giraitis et al. (2014, 2021), kernels with non-compact support such as the Gaussian kernel are permitted under some stronger condition. We focus on kernels with a compact support as specified in condition (a) to avoid unessential complications. Note that $2Th$ is the effective sample size of the kernel-based estimation, since $K((t-i)/Th) = 0$ for i such that $|t-i| > Th$ under Assumption 2.

2.2 Time-varying parameters with smoothness α

In discussing the consistency and asymptotic normality of $\hat{\beta}_t$, the smoothness of the path of $\beta_{T,t}$ has a decisive effect. In the following definition, we focus on the degree of smoothness of $\beta_{T,t}$ quantified by a single parameter α , rather than its stochastic/deterministic nature or specific form.

Definition 1. Triangular array $\beta_{T,t}$ such that $\beta_{T,t} = O_p(1)$ as $T \rightarrow \infty$ for all t is said to belong to the class type-a TVP(α) or type-b TVP(α), if the following condition (a) or (b) holds, respectively:

- (a) There exists some real $\alpha > 0$ such that for any sequence $\{a_T\}$ of positive integers satisfying $a_T \leq T$ and $a_T \rightarrow \infty$ as $T \rightarrow \infty$, and for any t ,

$$\max_{j:|t-j|\leq a_T} \|\beta_{T,t} - \beta_{T,j}\| = O_p\left(\left(\frac{a_T}{T}\right)^\alpha\right) \text{ as } T \rightarrow \infty.$$

- (b) There exists some real $\alpha > 0$ such that for any sequence $\{a_T\}$ of positive integers satisfying $a_T \leq T$ and $a_T \rightarrow \infty$ as $T \rightarrow \infty$, and for any t ,

$$\max_{j:|t-j|\leq a_T} \|\beta_{T,t} - \beta_{T,j}\| = O_p\left(\frac{1}{T^\alpha}\right) \text{ as } T \rightarrow \infty.$$

Definition 1 essentially controls by α the smoothness of the path of $\beta_{T,t}$ on any interval of any length (up to T). In typical applications, a_T will be set $a_T = \lfloor Th \rfloor$. Definition 1(a) allows the difference between the values of $\beta_{T,t}$ at distinct time points to grow as the time points gets further apart, while Definition 1(b) does not.⁴

Because $a_T/T \leq 1$ and $\alpha > 0$, smaller α permits larger differences $\|\beta_{T,t} - \beta_{T,j}\|$, resulting in $\beta_{T,t}$ possibly having a rougher path. Note that triangular arrays that are unbounded in probability are excluded from Definition 1. We emphasize that Definition 1 does not impose any parametric assumption on $\beta_{T,t}$, and moreover, $\beta_{T,t}$ may be deterministic or stochastic under this definition. In addition, $\beta_{T,t}$ is allowed to have arbitrary correlation with x_t and ε_t . Definition 1 is quite general and accommodates many important time-varying parameters, as shown below.

Remark 1. Giraitis et al. (2021) develop a kernel-based instrumental variable method to estimate time-varying parameters. The classes of time-varying parameters they consider are essentially type-a TVP(1) and TVP(1/2), albeit with slightly different definitions. They do not consider time-varying parameters belonging to type-a TVP(α) with $\alpha \neq 1/2, 1$ or type-b TVP(α). Our definition is more general than theirs.

Example 1 (Continuously differentiable functions). A popular model for time-varying parameters is deterministic smooth functions, accompanied by the formulation $\beta_{T,t} = \beta(t/T)$ for some continuously differentiable function β on $[0, 1]$ (e.g., Cai, 2007; Zhang and Wu, 2012; Chen and Hong, 2012). Under this formulation, the fact that $\sup_{0 \leq r \leq 1} \|\beta'(r)\| \leq C$ for some constant $C > 0$ implies that for any $s, t = 1, 2, \dots, T$,

$$\|\beta_{T,t} - \beta_{T,s}\| = \|\beta(t/T) - \beta(s/T)\| \leq C \frac{|t-s|}{T}$$

⁴Therefore, $\beta_{T,t}$ belongs to the type-a TVP(α) if it belongs to the type-b TVP(α).

by the mean value theorem. Therefore, we have $\max_{j:|t-j|\leq a_T} \|\beta_{T,t} - \beta_{T,j}\| \leq Ca_T/T = O(a_T/T)$, which implies continuously differentiable $\beta_{T,t}$ belongs to the type-a TVP(1) class. More generally, $\beta_{T,t}$ belongs to the type-a TVP(α) class if it is Hölder continuous with exponent α .

Example 2 (The random walk). It is often assumed in the time series literature that the parameters of interest follow the random walk (Cogley and Sargent, 2005; Giraitis et al., 2014; Georgiev, Harvey, Leybourne and Taylor, 2018). Suppose $\beta_{T,t}$ satisfies $\beta_{T,0} = \mu$ and $\beta_{T,t} = \mu + (1/\sqrt{T}) \sum_{i=1}^t u_i$, $t = 1, 2, \dots, T$, where μ is a constant and $\{u_i\}$ is a stationary sequence with $E[u_i] = 0$ and $V[u_i] = \Sigma_u > 0$. Also suppose that the functional central limit theorem (FCLT) applies to $\{u_i\}$, that is,

$$\beta_{T, \lfloor T \cdot \rfloor} = \mu + \frac{1}{\sqrt{T}} \sum_{i=1}^{\lfloor T \cdot \rfloor} u_i \Rightarrow \mu + \Sigma_u^{1/2} B_1(\cdot),$$

in the Skorokhod space $D_{[0,1]}^p$, where B_1 is a p -dimensional vector standard Brownian motion. Then, the following result holds: for any $t > a_T$,

$$\begin{aligned} \max_{j:|t-j|\leq a_T} \|\beta_{T,t} - \beta_{T,j}\| &\leq \max_{j:|t-j|\leq a_T} (\|\beta_{T,t} - \beta_{T,t-a_T}\| + \|\beta_{T,j} - \beta_{T,t-a_T}\|) \\ &\leq 2 \max_{1 \leq j \leq 2a_T} \|\beta_{T,t-a_T+j} - \beta_{T,t-a_T}\| \\ &= 2 \max_{1 \leq j \leq 2a_T} \left\| \frac{1}{\sqrt{T}} \sum_{i=t-a_T+1}^{t-a_T+j} u_i \right\| \\ &\stackrel{d}{=} 2 \max_{1 \leq j \leq 2a_T} \left\| \frac{1}{\sqrt{T}} \sum_{i=1}^j u_i \right\| \\ &= \sqrt{\frac{2a_T}{T}} 2 \sup_{0 \leq r \leq 1} \left\| \frac{1}{\sqrt{2a_T}} \sum_{i=1}^{\lfloor 2a_T r \rfloor} u_i \right\| = O_p \left(\sqrt{\frac{a_T}{T}} \right), \end{aligned}$$

by the stationarity of $\{u_t\}$ and the continuous mapping theorem (CMT). The same bound holds for $t \leq a_T$. Hence, the random walk divided by \sqrt{T} belongs to the type-a TVP(1/2). More generally, the random walk divided by T^α belongs to the type-a TVP(α) class for $\alpha \geq 1/2$. Note that the random walk divided by T^α with $\alpha < 1/2$ is excluded from Definition 1 because it is unbounded in probability.

Because a smaller α implies a (possibly) rougher path, the random walk divided by \sqrt{T} is rougher than continuously differentiable functions on $[0, 1]$. This is intuitively because

the random walk divided by \sqrt{T} weakly converges to Brownian motion, which is nowhere differentiable almost surely.

Remark 2. Müller and Petalas (2010) study an inferential problem concerning time-varying parameters approximated by Gaussian processes and piece-wise continuous functions scaled by a factor of $T^{-1/2}$. Leading examples are $T^{-1/2}\beta(t/T)$ with $\beta(\cdot)$ continuous on $[0, 1]$ and $T^{-1/2}B_1(t/T)$, which is approximately equivalent (in distribution) to $T^{-1}\sum_{i=1}^t u_i = O_p(1/\sqrt{T})$. Therefore, non-vanishing smooth functions and random walks are not considered in their framework.

Example 3 (Structural breaks). Structural breaks in parameters have been long studied in econometrics and statistics as they capture discontinuous changes (Casini and Perron, 2018, provide a recent survey on this topic). Suppose time-varying coefficient $\beta_{T,t}$ experiences one abrupt break during the sample period:

$$\beta_{T,t} = \begin{cases} \beta_1 & \text{for } t = 1, 2, \dots, T_B \\ \beta_2 & \text{for } t = T_B + 1, T_B + 2, \dots, T \end{cases}, \quad (2)$$

where $T_B = \lfloor \tau_B T \rfloor$, $\tau_B \in (0, 1)$, and $\|\beta_1 - \beta_2\| = \delta/T^\alpha$ for some $\delta > 0$ and $\alpha > 0$. Then, $\beta_{T,t}$ belongs to the type-b TVP(α) class. Specifically, we have, for any $t \in \{1, \dots, T_B - a_T\} \cup \{T_B + a_T + 1, \dots, T\}$, $\max_{j:|t-j|\leq a_T} \|\beta_{T,t} - \beta_{T,j}\| = 0$, and for any $t \in \{T_B - a_T + 1, \dots, T_B + a_T\}$, $\max_{j:|t-j|\leq a_T} \|\beta_{T,t} - \beta_{T,j}\| = \delta T^{-\alpha}$. Note that the asymptotically non-negligible discontinuity given by $\alpha = 0$ is excluded from Definition 1.

Example 4 (Threshold models). Hansen (2000) considers the threshold regression model, which is obtained by letting $\beta_{T,t} = \theta_1 + \delta_T 1\{q_t > \eta\}$, where q_t is the threshold variable that determines the regime at time t , depending on whether it exceeds threshold parameter η . δ_T , which Hansen (2000) refers to as the threshold effect, expresses the magnitude of discontinuous changes in $\beta_{T,t}$. To obtain the asymptotic distribution of the estimator for η that is free from nuisance parameters, Hansen (2000) imposes $\delta_T = c/T^\alpha$.⁵ Under this formulation, $\beta_{T,t}$ clearly satisfies $\|\beta_{T,t} - \beta_{T,j}\| \leq \|\delta_T\| = O_p(1/T^\alpha)$, for all t and j , which implies $\beta_{T,t}$ belongs to type-b TVP(α).

Example 5 (Mixed model). Suppose that $\beta_{T,t}$ is expressed as $\beta_{T,t} = \beta_{1,T,t} + \beta_{2,T,t}$, where $\beta_{1,T,t}$ is continuously differentiable and $\beta_{2,T,t} = \mu + (1/\sqrt{T})\sum_{i=1}^t u_i$ with u_i defined as in

⁵Hansen (2000) also imposes $0 < \alpha < 1/2$, but this restriction is not necessary in our framework.

Example 2. Then, it holds that

$$\begin{aligned} \max_{j:|t-j|\leq a_T} \|\beta_{T,t} - \beta_{T,j}\| &\leq \max_{j:|t-j|\leq a_T} \|\beta_{1,T,t} - \beta_{1,T,j}\| + \max_{j:|t-j|\leq a_T} \|\beta_{2,T,t} - \beta_{2,T,j}\| \\ &= O_p\left(\frac{a_t}{T}\right) + O_p\left(\sqrt{\frac{a_T}{T}}\right) = O_p\left(\sqrt{\frac{a_T}{T}}\right), \end{aligned}$$

since $a_T/T \leq 1$. Therefore, $\beta_{T,t}$ belongs to type-a TVP(1/2). More generally, for any finite positive integer S , if $\beta_{T,t}$ is expressed as the sum of S time-varying parameters each of which belongs to the type-a TVP(α_s) class ($s = 1, \dots, S$), then $\beta_{T,t}$ belongs to type-a TVP($\min\{\alpha_1, \dots, \alpha_S\}$).

2.3 Asymptotic properties of $\hat{\beta}_t$

In the following theorem, we establish the consistency and asymptotic normality of the kernel-based estimator, $\hat{\beta}_t$.

Theorem 1. *Suppose Assumptions 1 and 2 hold. Then, we have, for model (1) and $t = \lfloor Tr \rfloor$, $r \in (0, 1)$,*

$$\sqrt{Th}(\hat{\beta}_t - \beta_{T,t} - R_{T,t}) \xrightarrow{d} N(0, \Omega(r)^{-1}\Sigma(r)\Omega(r)^{-1}), \quad (3)$$

where

$$R_{T,t} = \begin{cases} O_p(h^\alpha) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(a)} \\ O_p(T^{-\alpha}) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(b)} \end{cases}. \quad (4)$$

In particular, for $h = cT^\gamma$, $c > 0, \gamma \in (-1, 0)$, we have

$$\sqrt{cT^{1+\gamma}}(\hat{\beta}_t - \beta_{T,t}) \xrightarrow{d} N(0, \Omega(r)^{-1}\Sigma(r)\Omega(r)^{-1}), \quad (5)$$

for $\gamma \in \Gamma(\alpha)$, where

$$\Gamma(\alpha) = \begin{cases} (-1, -\frac{1}{2\alpha+1}) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(a)} \\ (-1, 2\alpha - 1) \cap (-1, 0) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(b)} \end{cases}. \quad (6)$$

Proof. See Appendix A. □

Remark 3. We do not derive the asymptotic distribution of $\hat{\beta}_t$ at boundary points (near $t = 0$ and $t = T$), but the derivation will proceed along the lines of Cai (2007). As shown

by Cai (2007), the local constant (Nadaraya-Watson) estimator suffers from a larger bias at boundary points than the local linear estimator if $\beta_{T,t}$ is continuously differentiable. However, as revealed soon later (in Example 1 below), the local linear estimator is available only when $\beta_{T,t}$ is (continuously) differentiable and is not applicable to nondifferentiable time-varying parameters such as the random walk. In view of the prevalence of nondifferentiable time-varying parameters in applied economics, an estimator that only applies to differentiable time-varying parameters is not appealing. For this reason, we will continue to use the local constant estimator rather than the local linear one.

Corollary 1. *Suppose Assumptions 1(a)-(b) and 2 hold. Suppose also that $\{x_t\}_t$ and $\{x_t\varepsilon_t\}_t$ are covariance-stationary. Then, (3)-(6) hold with $\Omega(r)$ and $\Sigma(r)$ replaced by $\Omega := E[x_1x_1']$ and $\Sigma := \int_{-1}^1 K(x)^2 dx E[\varepsilon_1^2 x_1 x_1']$, respectively.*

Proof. See Appendix A. □

In what follows, we will set $h = cT^\gamma$ and call γ (as well as h) the bandwidth parameter. The effect of smoothness parameter α is twofold. First, the asymptotic order of the bias term, $R_{T,t}$, depends on α . The smaller α is, the larger the order of $R_{T,t}$ is, because the path of $\beta_{T,t}$ (possibly) becomes rougher. And it is through this bias term that $\Gamma(\alpha)$, the set of bandwidth parameter γ that yields \sqrt{Th} -consistency and asymptotic normality,⁶ depends on α , which in turn makes the rate of convergence $T^{(1+\gamma)/2}$ dependent on α . This is the second effect. Letting $\alpha \rightarrow 0$, the kernel-based estimation can accommodate time-varying parameters of arbitrary smoothness, but this is accompanied by $\Gamma(\alpha) \rightarrow -1$, resulting in the rate of convergence $T^{(1+\gamma)/2} \rightarrow 1$. In contrast, if we let $\alpha \rightarrow \infty$, then $\Gamma(\alpha)$ tends to $(-1, 0)$, and the choice $\gamma \approx 0$ yields a nearly \sqrt{T} -rate convergence, but only highly smooth parameters can be estimated. This observation reveals that there is a trade-off between efficiency (the rate of convergence) and robustness (the range of allowable smoothness α of the time-varying parameter) in the kernel-based estimation.

Because $\Gamma(\alpha)$ is the set of bandwidths that yield \sqrt{Th} -consistency and asymptotic normality under given α , we can obtain the set of α that leads to \sqrt{Th} -consistency and asymptotic normality of $\hat{\beta}_t$ under given γ , by inverting the expression of $\Gamma(\alpha)$. Letting $A(\gamma)$ denote such a set, we can say that $\hat{\beta}_t$ calculated using given γ is \sqrt{Th} -consistent and asymptotically

⁶Note that $\Gamma(\alpha)$ is the set of bandwidths such that $\sqrt{Th}R_{T,t} = o_p(1)$ and therefore should be labeled as the set of bandwidths that yield \sqrt{Th} -consistency (not mere consistency) and asymptotic normality. From (3), it holds that $\hat{\beta}_t - \beta_{T,t} = o_p(1)$ for any $\gamma < 0$ and $\alpha > 0$ because $R_{T,t} = O_p(\max\{T^{\gamma\alpha}, T^{-\alpha}\}) = o_p(1)$, but $\hat{\beta}_t$ is neither \sqrt{Th} -consistent nor asymptotically normal if $\gamma \notin \Gamma(\alpha)$ in general.

normal for time-varying parameters with smoothness $\alpha \in A(\gamma)$, where

$$A(\gamma) = \begin{cases} (-\frac{1+1/\gamma}{2}, \infty) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(a)} \\ (\frac{1+\gamma}{2}, \infty) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(b)} \end{cases}. \quad (7)$$

Letting $\gamma \rightarrow -1$, $A(\gamma)$ tends to $(0, \infty)$, which implies that time-varying parameters with any smoothness $\alpha > 0$ can be accommodated, but the rate of convergence becomes $T^{(1+\gamma)/2} \rightarrow 1$. On the other hand, if we let $\gamma \uparrow 0$, then the rate of convergence is as fast as \sqrt{T} , but $A(\gamma) \rightarrow \infty$ (the smoothness of constant parameters) in the type-a case. Hence, the bandwidth parameter γ determines the trade-off between efficiency and robustness.

Example 1 (Continued). Because continuously differentiable $\beta_{T,t}$ belongs to the type-a TVP(1) class, for any $\gamma \in \Gamma(1) = (-1, -1/3)$, we have $\sqrt{cT^{1+\gamma}}(\hat{\beta}_t - \beta_{T,t}) \xrightarrow{d} N(0, \Omega(r)^{-1}\Sigma(r)\Omega(r)^{-1})$. Setting $\gamma \approx -1/3$ gives the fastest rate of convergence of $T^{1/3}$. If we take advantage of the continuous differentiability of $\beta_{T,t}$, and the kernel is symmetric, then the set of the admissible bandwidths, $\Gamma(\alpha)$, widens to $(-1, -1/5)$, giving the faster rate of convergence of $T^{2/5}$ (see Cai, 2007). In general, we will be able to enlarge $\Gamma(\alpha)$ to $(-1, -1/(4\alpha + 1))$ in the type-a case if the following additional condition (mimicking the Taylor expansion) holds:

$$\max_{j:|t-j|\leq a_T} \|\beta_{T,t} - \beta_{T,j} - c_t(\frac{t}{T} - \frac{j}{T})\| = O_p\left(\left(\frac{a_T}{T}\right)^{2\alpha}\right), \quad (8)$$

for some (possibly random) bounded vector c_t . This condition is satisfied by twice continuously differentiable functions. Condition (8), however, essentially requires the differentiability of $\beta_{T,t}$ with respect to time, which is not satisfied by, e.g., the random walk divided by \sqrt{T} , so that the enlarged version of $\Gamma(\alpha)$ is only available to a limited class of time-varying parameters. For the same reason, the local linear estimator, which is based on the Taylor expansion of $\beta_{T,t}$, is not applicable to nondifferentiable time-varying parameters.

Example 2 (Continued). If $\beta_{T,t}$ is the random walk divided by \sqrt{T} , which belongs to the type-a TVP(1/2) class, then $\Gamma(1/2) = (-1, -1/2)$, and thus the fastest rate of convergence given by $\gamma \approx -1/2$ is $T^{1/4}$, which is slower than in the continuously differentiable case. The same set of admissible bandwidths is derived by Giraitis et al. (2014), who considers a random walk type time-varying coefficient in the context of univariate AR(1) models.

Furthermore, we show in Appendix B that the bandwidth minimizing the MSE of $\hat{\beta}_t$ is proportional to $T^{-1/2}$ when $\beta_{T,t}$ is the random walk divided by \sqrt{T} . We prove this result under a condition that is stronger than Assumption 1 but still retains generality to some extent. Therefore, the choice of $\gamma = -1/2$ may also be justified as the minimizer of the MSE of the kernel-based estimator.

Example 3 (Continued). Suppose $\beta_{T,t}$ is defined as in (2). Because $\beta_{T,t}$ belongs to the type-b TVP(α) class, arbitrary γ in $(-1, 0)$ yields the \sqrt{Th} -consistency and asymptotic normality of $\hat{\beta}_t$ as long as $\alpha \geq 1/2$. In particular, setting $\gamma \approx 0$ gives a near \sqrt{T} -consistency.⁷

For the case of $\alpha \in (0, 1/2]$, however, a smaller α leads to a larger discontinuity in $\beta_{T,t}$ and thus a slower rate of convergence (through a narrower $\Gamma(\alpha)$). Therefore, if $\beta_{T,t}$ experiences large structural breaks given by $\alpha < 1/2$ (and there is no other source of instability in the path of $\beta_{T,t}$), a conventional structural-break approach (e.g., the sequential procedure proposed by Bai and Perron, 1998) will be more suitable than the kernel-based estimation.

Example 4 (Continued). The argument given in Example 3 also applies to the threshold model: When $\delta_T = O_p(1/T^\alpha)$ with $\alpha \geq 1/2$, the kernel-based method delivers a \sqrt{Th} -consistent, asymptotically normal estimation of $\beta_{T,t}$, whereas Hansen's (2000) method should be used when $\alpha < 1/2$.

3 Consequence of Improper choice of the bandwidth

In Theorem 1, we showed the range of bandwidth that yields \sqrt{Th} -consistency and asymptotic normality depends on the smoothness α of $\beta_{T,t}$. This implies that an improper choice of the bandwidth (given by $\gamma \notin \Gamma(\alpha)$) leads to misleading inference. In this section, we illustrate this implication through some examples where the evolutionary mechanism of $\beta_{T,t}$ is misspecified, which leads to an improper bandwidth selection. We also discuss how to choose bandwidth parameter γ when there is no prior information on smooth parameter α .

3.1 When random-walk $\beta_{T,t}$ is mistakenly assumed to be continuously differentiable

Suppose one assumes $\beta_{T,t}$ is a continuously differentiable function and sets $\gamma \approx -1/3$, but the fact is that $\beta_{T,t}$ follows the random walk divided by \sqrt{T} . Recall from the analysis given in Example 2 that $\gamma \approx -1/3$ is invalid in this case. The kernel-based estimator satisfies $\sqrt{cT^{1+\gamma}}(\hat{\beta}_t - \beta_{T,t}) = S_{T,t} + O_p(T^{1/2+\gamma})$, where $S_{T,t} \xrightarrow{d} N(0, \Omega(r)^{-1}\Sigma(r)\Omega(r)^{-1})$. Since the bias term is of order $O_p(T^{1/2+\gamma})$ and $\gamma \approx -1/3$, the difference $\hat{\beta}_t - \beta_{T,t}$ is dominated by the bias term. This is the case for any $\gamma > -1/2$. Because the bias term, which is a function

⁷In fact, setting exactly $\gamma = 0$ yields \sqrt{T} -consistency and asymptotic normality if $\alpha > 1/2$. In this case, each $\beta_{T,t}$, $t = 1, \dots, T$ is estimated by using the full sample, but $\hat{\beta}_t$ and $\hat{\beta}_s$ ($t \neq s$) may take different values. This is because the weighting scheme (based on the kernel, $K(\cdot)$) is different for different time points. If $K(\cdot)$ is the uniform kernel, $\hat{\beta}_t$ equals the full-sample OLS estimator for all t .

of random-walk $\beta_{T,t}$, is not normal in general, confidence intervals based on the normal approximation given in Theorem 1 will perform poorly.

In the literature on smooth (differentiable) time-varying parameters, researchers have often used a rule-of-thumb bandwidth $h = \text{constant} \times T^{-1/5}$, or picked the minimizer of the cross validation criterion over $h \in [c_1 T^{-1/5}, c_2 T^{-1/5}]$ for some $0 < c_1 < c_2$ (Chen and Hong, 2012; Sun, Hong, Wang and Zhang, 2023). Although these selection rules lead to an efficient estimation of $\beta_{T,t}$ as long as it is correctly specified as a continuously differentiable function, they will yield a biased estimation if $\beta_{T,t}$ is a random walk, or more generally, if $\beta_{T,t}$ belongs to the type-a TVP(α) class with $\alpha < 1$ but does not belong to type-a TVP(1).

3.2 The effect of neglected breaks

Suppose $\beta_{T,t} = \mu_{T,t} + (1/\sqrt{T}) \sum_{i=1}^t u_i$, where u_t is specified as in Example 2, and $\mu_{T,t}$ satisfies

$$\mu_{T,t} = \begin{cases} \mu_1 & \text{for } t = 1, 2, \dots, T_B \\ \mu_2 & \text{for } t = T_B + 1, T_B + 2, \dots, T \end{cases}, \quad (9)$$

with $T_B = \lfloor \tau_B T \rfloor$ and $\mu_2 - \mu_1 = \delta/T^\alpha$. Then, we can show

$$R_{T,t} = \begin{cases} O_p(T^{\gamma/2}) & \text{for } t \in [1, T_B - \lfloor Th \rfloor] \cup [T_B + 1 + \lfloor Th \rfloor, T] \\ O_p(\max\{T^{\gamma/2}, T^{-\alpha}\}) & \text{for } t \in [T_B - \lfloor Th \rfloor + 1, T_B + \lfloor Th \rfloor] \end{cases},$$

where $R_{T,t}$ is defined in (3) and (4). The asymptotic order of the bias term, $R_{T,t}$, is $O_p(T^{\gamma/2})$ for t outside the $\lfloor Th \rfloor$ -neighborhood of break point T_B . On the $\lfloor Th \rfloor$ -neighborhood of T_B , it is $O_p(T^{\gamma/2})$ if $\alpha \geq -\gamma/2$, while it is $O_p(T^{-\alpha})$ if $0 < \alpha < -\gamma/2$.

Suppose we estimate $\beta_{T,t}$ by $\hat{\beta}_t$ assuming $\mu_{T,t} = \mu$, that is, the parameter instability is purely due to the zero-mean random walk. In this case, the (misleading) optimal rate of convergence is achieved by the choice of $\gamma \approx -1/2$, yielding

$$R_{T,t} = \begin{cases} O_p(T^{\gamma/2}) & \text{if } \alpha \geq 1/4 \\ O_p(T^{-\alpha}) & \text{if } 0 < \alpha < 1/4 \end{cases}.$$

When $\alpha \geq 1/4$, the asymptotic order of $R_{T,t}$ is $O_p(T^{\gamma/2})$, which is the same order as in the pure random walk case (see (4) and set $\alpha = 1/2$, $h = cT^\gamma$), so that the choice $\gamma \approx -1/2$ is valid and leads to the fastest rate of convergence. In contrast, if $0 < \alpha < 1/4$, $R_{T,t} = O_p(T^{-\alpha})$. Because the asymptotically normal component of the decomposition of $\hat{\beta}_t - \beta_{T,t}$ is $O_p(T^{-(1+\gamma)/2}) \approx O_p(T^{-1/4})$, the asymptotic behavior of $\hat{\beta}_t - \beta_{T,t}$ is dominated by the bias

term. Therefore, the structural break induces a severe bias in the kernel-based estimation if the break is of order $T^{-\alpha}$ with $0 < \alpha < 1/4$.

One faces this bias when estimating $\beta_{T,t}$ with $t = T_B \pm [rTh]$, $r \in [0, 1]$. Consider, for example, $t = T_B$. Then, the following decomposition holds for the kernel-based estimator:

$$\sqrt{Th}(\hat{\beta}_{T_B} - \beta_{T,T_B}) = S_{T,t} + \left(\frac{1}{Th} \sum_{i=T_B+1}^{T_B+[Th]} K\left(\frac{T_B-i}{Th}\right)x_i x_i' \right)^{-1} \frac{1}{\sqrt{Th}} \sum_{i=T_B+1}^{T_B+[Th]} K\left(\frac{T_B-i}{Th}\right)x_i x_i' \frac{\delta}{T^\alpha},$$

where $S_{T,t} \xrightarrow{d} N(0, \Omega(r)^{-1}\Sigma(r)\Omega(r)^{-1})$. The second term is of order $T^{(1+\gamma)/2-\alpha}$ (as long as $(Th)^{-1} \sum_{i=T_B+1}^{T_B+[Th]} K((T_B-i)/(Th))x_i x_i'$ converges in probability to some nonsingular matrix) and diverges as $T \rightarrow \infty$, since $\gamma \approx -1/2$ and $0 < \alpha < 1/4$. A bias of the same order appears for any $t = T_B \pm [rTh]$ with $r \in [0, 1]$. Therefore, the kernel-based estimator is severely biased on the $[Th]$ -neighborhood of the discontinuity point $t = T_B$.

A more general result can be derived if we invoke $\Gamma(\alpha)$ and $A(\gamma)$ defined in (6) and (7), respectively. Suppose that $\beta_{T,t}$ can be expressed as $\beta_{T,t}^1 + \mu_{T,t}$, where $\beta_{T,t}^1$ belongs to type-a TVP(α_1) with $\alpha_1 \in (0, \infty)$, and $\mu_{T,t}$ is defined as in (9) but the magnitude of the break is $\mu_2 - \mu_1 = \delta/T^{\alpha_2}$. Then, $\hat{\beta}_t$ is \sqrt{Th} -consistent and asymptotically normal under any $\gamma \in \Gamma(\alpha_1) = (-1, -1/(2\alpha_1 + 1))$ for $t \in [1, T_B - [Th]] \cup [T_B + 1 + [Th], T]$. On $[T_B - [Th] + 1, T_B + [Th]]$, the same γ leads to \sqrt{Th} -consistency and asymptotic normality if $\alpha_2 \in A(\gamma) = ((1+\gamma)/2, \infty)$, while the bias term dominates the asymptotically normal term in the decomposition of $\hat{\beta}_t$ if $\alpha_2 < (1+\gamma)/2$.

3.3 The choice of the bandwidth

In the previous (sub)sections, we have observed that there is a trade-off in kernel regression between the rate of convergence and the size of the class of time-varying parameters that can be estimated. On the one hand, if we set bandwidth parameter γ near 0, then the rate of convergence is nearly as fast as \sqrt{T} , but only a tiny class of time-varying parameters is accommodated. On the other hand, if we set γ near -1 , then the kernel-based estimator accommodates a large class of time-varying parameters, but the rate of convergence is nearly as slow as T^0 . If one can correctly specify the evolutionary mechanism of $\beta_{T,t}$, or there is some information on the smoothness of $\beta_{T,t}$, one may select γ appropriately. However, if there is no such information, how should γ be selected?

Given the popularity of the random walk coefficient model in the time series literature,

setting $\gamma \approx -1/2$ (or $\gamma = -1/2$ in view of the MSE) would be appealing. Note that this choice accommodates not only the rescaled random walk but also continuously differentiable coefficients and small to moderate breaks and threshold effects (given by $\alpha \geq 1/4$; recall Section 3.2). Moreover, mixtures of these specifications can also be estimated under this choice of γ . The price of this choice is that the rate of convergence of the kernel-based estimator is $T^{1/4}$, slower than that given by the conventional choice of $\gamma = -1/5$.

Note that time-varying parameters that belong to type-a TVP(α) with $\alpha < 1/2$ but do not belong to type-a TVP(α) with $\alpha \geq 1/2$ are not accommodated by the choice of $\gamma \approx -1/2$ (see the definition of $A(\gamma)$ given in (7)). Leading examples are functions that are Hölder continuous with exponent $\alpha < 1/2$ but are not for $\alpha \geq 1/2$. Thus, the validity of the choice $\gamma \approx -1/2$ should be evaluated under the exclusion of this class. Time-varying parameters that belong to type-a TVP(α) with $\alpha < 1/2$ possibly have a rougher and uncontrolled path than the rescaled random walk. Therefore, such models will be relevant only if one is interested in highly unstable economic relationships. Our empirical application will show that random walk type parameter instabilities can be well captured by the choice of $\gamma = -1/2$ (see Section 5). Therefore, this choice will be adequate in many applications.

Remark 4. Although the choice $\gamma \approx -1/2$ can accommodate many important time-varying parameter models, it still yields a biased estimation if $\beta_{T,t}$ experiences large structural breaks or threshold effects (given by $\alpha < 1/4$). A possible way to avoid facing bias around the discontinuous points would be to apply the sample splitting approach based on some test for structural breaks (e.g., as proposed by Bai and Perron, 1998) and then apply kernel regression within each subsample. If large threshold effects are suspected, Hansen’s (2000) sample-splitting method may be used.

A natural question about the use of those sample-splitting approaches is whether they can correctly identify latent discontinuous changes in the presence of another source of parameter instability (e.g., smooth functions or the random walk). Because theoretically analyzing the behavior of these methods in such a setting will be quite complicated, we investigate it via simulation. In Appendix C, we provide simulation results for the behavior of several structural break tests. According to the results, conventional structural break tests can both underestimate and overestimate the number of breaks with a nonnegligible (or quite large in some cases) probability. The underestimation of the number of breaks implies that some latent abrupt breaks are overlooked, and the overestimation implies that spurious abrupt breaks are detected. Therefore, conventional structural break tests probably are not suitable

for detecting abrupt breaks if they are mixed with smooth parameter instabilities. The development of a test that detects latent breaks with high probability and spurious breaks with small probability in such a setting is important, but is beyond the scope of the present work.

4 Monte Carlo Simulation

In this section, we conduct two Monte Carlo experiments to verify the implications provided in Section 3. We use the following data generating process (DGP) $y_t = \beta_{T,t}x_t + \varepsilon_t$, $t = 1, \dots, T$, where $x_t = 0.5x_{t-1} + \varepsilon_{x,t}$ with $\varepsilon_{x,t} \sim \text{i.i.d. } N(0, 1)$, and $\beta_{T,t}$ is defined differently in different experiments. For the specification of ε_t , we consider two cases: $\varepsilon_t = u_t$, where $u_t \sim \text{i.i.d. } N(0, 1)$ (i.i.d. case) and $\varepsilon_t = \sigma_t u_t$ with $\sigma_t^2 = 0.1 + 0.3\varepsilon_{y,t-1}^2 + 0.6\sigma_{t-1}^2$ (GARCH case). To obtain $\hat{\beta}_t$, we use the Epanechnikov kernel $K(x) = 0.75(1 - x^2)I(|x| \leq 1)$.

4.1 Simulation for Section 3.1

The first experiment is related to Section 3.1. In this simulation, $\beta_{T,t}$ is generated as the rescaled random walk: $\beta_{T,t} = T^{-1/2} \sum_{i=1}^t v_i$. We consider two DGPs for the driver process v_t : (i) $v_t \sim \text{i.i.d. } N(0, 1)$ and (ii) $v_t \sim \text{i.i.d. log normal}$ with parameters $\mu = 0$ and $\sigma = 1$.⁸ Four sample sizes are used: $T \in \{100, 200, 400, 800\}$. To evaluate the global performance of $\hat{\beta}_t$, we calculate $\text{MSE} = T^{-1} \sum_{t=1}^T (\hat{\beta}_t - \beta_{T,t})^2$ (the reported MSE is the mean MSE over 2000 replications). To evaluate how well the normal approximation works for $\hat{\beta}_t$, we consider $\beta_{T,t}$ with $t = 0.5T$ (the middle point of the sample). We construct the 95% confidence interval for $\beta_{T,0.5T}$ based on the normal approximation given in Corollary 1. The variance estimators are $\hat{\Omega} := T^{-1} \sum_{i=1}^T x_i^2$ and $\hat{\Sigma} := \int_{-1}^1 K(x)^2 dx \times T^{-1} \sum_{i=1}^T \hat{\varepsilon}_i^2 x_i^2$, where $\hat{\varepsilon}_i = y_i - \hat{\beta}_i x_i$. We experiment with bandwidth parameter $h = T^\gamma$ and $\gamma \in \{-0.2, -0.33, -0.5, -0.55, -0.6, -0.7\}$, and evaluate the performance for each pair (γ, T) . The results are presented in Tables 1 and 2.

Because $\beta_{T,t}$ is the random walk divided by \sqrt{T} , an appropriate bandwidth is $\gamma \approx -1/2$. According to the analysis given in Section 3.1, the kernel-based estimator leads to poor inference when $\gamma > -1/2$. Our simulation result corroborates this analysis. First, consider the case where $\varepsilon_t \sim \text{i.i.d. } N(0, 1)$ (Table 1). In case (i) (Gaussian random-walk $\beta_{T,t}$), when $\gamma = -0.2$, which is a common choice in the literature on models with differentiable time-

⁸Specifically, X follows a log normal distribution if $X = \exp(Z)$, where $Z \sim N(\mu, \sigma^2)$.

varying parameters, the coverage probability is far below the 95% confidence level. What is worse, it deviates from 0.95 as T increases. Note that the MSE is relatively large except when T is small. When $\gamma = -1/3$, the MSE takes the smallest value for all T considered, but the coverage probability is still too small. This result warns researchers against using these bandwidths unless they are confident that $\beta_{T,t}$ can be well approximated by smooth functions with smoothness parameter $\alpha = 1$. For $\gamma \leq -1/2$, the interval estimation performs well with coverage probability being 85-90% and getting better as T increases. However, $\gamma = -0.7$ leads to undercoverage when T is small and the largest MSE for all T . $\gamma = -0.6$ also gives large MSEs. The choices $\gamma \approx -1/2$ lead to good coverage and small MSE, so that these choices are recommended for random walk type parameters, or more generally, for parameters with $\alpha = 1/2$. The result for case (ii) (non-Gaussian random-walk $\beta_{T,t}$) is similar to that for case (i), so the same comment applies.

Results for the case where ε_t is GARCH (Table 2) are similar to those for the i.i.d case. Hence, we do not repeat the same analysis here.

4.2 Simulation for Section 3.2

The second experiment is for verifying the implication provided in Section 3.2. In this simulation, we analyze the effect of (neglected) structural breaks. For this purpose, we generate $\beta_{T,t}$ according to $\beta_{T,t} = \mu_{T,t} + T^{-1/2} \sum_{i=1}^t v_i$ (rescaled random walk plus intercept), where $v_i \sim \text{i.i.d. } N(0,1)$ and $\mu_{T,t}$ is an intercept term experiencing a break at $t = 0.5T$. Specifically, we let

$$\mu_{T,t} = \begin{cases} 0 & \text{for } t = 1, \dots, 0.5T \\ 2/T^\alpha & \text{for } t = 0.5T + 1, \dots, T \end{cases},$$

where $\alpha \in \{0.1, 0.2, 0.3, 0.4, 0.5\}$. A smaller α yields a larger break. We consider estimating $\beta_{T,t}$ with the choice $\gamma = -1/2$, reflecting the ignorance of the break. According to our theoretical analysis, the kernel-based estimator has a severe bias around $t = 0.5T$ when $\alpha < 0.25$, while breaks given by $\alpha > 0.25$ have no effect asymptotically. To confirm this implication, we calculate the MSE and coverage probability of $\hat{\beta}_t$ for $t = \tau T$ with $\tau = 0.4, 0.45, 0.5, 0.55, 0.6$. The MSE is calculated for each τ as the mean squared error over 2000 replications, that is, $\text{MSE}(\tau) = 2000^{-1} \sum_{i=1}^{2000} (\hat{\beta}_{\tau T}^{(i)} - \beta_{T,\tau T}^{(i)})^2$, where superscript i signifies $\hat{\beta}_{\tau T}^{(i)}$ and $\beta_{T,\tau T}^{(i)}$ are obtained in the i th replication. We consider four sample sizes; (i) $T = 100$, (ii) $T = 200$, (iii) $T = 400$, and (iv) $T = 800$. As the variance estimators, we use

$\hat{\Omega}_t = (Th)^{-1} \sum_{i=1}^T K((t-i)/Th)x_i^2$ and $\hat{\Sigma}_t = (Th)^{-1} \sum_{i=1}^T K((t-i)/Th)^2 \hat{\varepsilon}_i^2 x_i^2$ to evaluate the normal approximation given in Theorem 1. Results are reported in Tables 3 and 4.

First, let us see the case of ε_t being i.i.d. and $T = 100$ (Table 3, the row labeled as (i)). The MSEs and coverage probabilities for $\tau = 0.4$ and 0.6 are stable across α . This is because the break only affects estimation around the discontinuity point $t = 0.5T$. However, the break has a severe effect on $\hat{\beta}_{\tau T}$ with $\tau = 0.45, 0.5, 0.55$, both in terms of MSE and coverage. The smaller α is (i.e. the larger the break is), the worse the performance gets. Moreover, this effect is more profound for τ closer to 0.5 . For example, the MSE for $(\alpha, \tau) = (0.1, 0.5)$ is about six times as large as the MSE for $\alpha = 0.1$ and $\tau = 0.4, 0.6$. In terms of the coverage probability, smaller breaks given by $\alpha \geq 0.25$ affect the kernel-based estimation. This indicates that, although breaks of these magnitudes asymptotically have no impact, they do have nontrivial effects in finite samples.

Note that the MSE of $\hat{\beta}_t$ affects the precision of variance estimator $\hat{\Sigma}_t = (Th)^{-1} \sum_{i=1}^T K((t-i)/Th)^2 \hat{\varepsilon}_i^2 x_i^2$ through $\hat{\varepsilon}_t = \varepsilon_t - (\hat{\beta}_t - \beta_{T,t})x_t$. Therefore, a larger MSE of $\hat{\beta}_t$ leads to a wider confidence interval.⁹ To confirm this implication, we calculate the length of the confidence interval for each τ . Table 3 also reports the mean of the interval length taken over the 2000 replications. When $T = 100$, the means of the interval length take similar values across τ for $\alpha \geq 0.25$. However, larger breaks given by $\alpha < 0.25$ lead to wider confidence intervals around the break point. For example, the confidence interval for $\alpha = 0.1$ and $\tau = 0.5$ is about 10-13% longer than that for $\alpha = 0.1$ and $\tau = 0.4, 0.6$ on average.

For case (ii) ($T = 200$), MSEs for $\tau = 0.5$ and $\alpha < 0.25$ are still large. Note that MSEs for $\tau = 0.45, 0.55$ are comparable with those for $\tau = 0.4, 0.6$. This is because the abrupt break affects $\hat{\beta}_t$ on the Th -neighborhood of the break date. Because $t = 0.45T$ and $t = 0.55T$ are outside the Th -neighborhood of $0.5T$, the performance of $\hat{\beta}_{\tau T}$ improves as T increases for $\tau = 0.45, 0.55$. In terms of the interval length, the same comment as in case (i) applies. For example, the confidence interval for $\alpha = 0.1$ and $\tau = 0.5$ is 15% longer than that for $\tau = 0.4, 0.6$ on average. $\hat{\beta}_{0.5T}$ also suffers from poor coverage. In particular, the coverage probabilities for $\alpha < 0.25$ get worse as T increases. For the cases with $T = 400, 800$ (cases (iii) and (iv)), a similar comment applies.

Examining the case with ε_t being GARCH (see Table 4), the same conclusion is drawn, so the detail is omitted.

⁹Note that the width of the confidence interval for $\beta_{T,t}$ is identical across t if $\hat{\Omega}$ and $\hat{\Sigma}$ are used as the variance estimators.

5 Empirical Application

To illustrate the use of our results in empirical applications, we consider estimating the time-varying capital asset pricing model (CAPM).¹⁰ Parameter instabilities are widely observed in the CAPM literature (see Ghysels, 1998; Lewellen and Nagel, 2006; Fama and French, 2006; Ang and Chen, 2007; Ang and Kristensen, 2012; Guo, Wu and Yu, 2017, and references therein). We consider estimating the following factor model:

$$R_{j,t} = \alpha_{j,t} + \beta_{j,t}R_{M,t} + \varepsilon_t,$$

where $R_{j,t}$ denotes the excess return of portfolio j at time t , and $R_{M,t}$ is the market excess return. The coefficients alpha and beta are allowed to be time-varying.

5.1 Background

In the CAPM literature, a popular approach to modeling parameter instability is to assume that the parameters depend on observable instrumental variables. But results drawn from this approach tend to be sensitive to the choice of instruments (Ghysels, 1998). To overcome this problem, researchers have proposed time-varying parameter models that do not depend on the use of exogenous information.

Some assume that parameters experience abrupt changes at some time points (Lewellen and Nagel, 2006; Fama and French, 2006), and others model parameter instability via the (near) random walk (Ang and Chen, 2007) or smooth function of time (Li and Yang, 2011; Ang and Kristensen, 2012). However, these results can be misleading if the evolutionary mechanism of parameters is misspecified. For example, Fama and French (2006) and Lewellen and Nagel (2006) assume that changes in parameter value occur based on calendar time (e.g., monthly or yearly), and estimate parameters by the OLS within each subsample. However, estimates obtained in this fashion suffer from bias if the timing of structural breaks is misspecified. Ang and Chen (2007) estimate time-varying alpha and beta assuming that they are the (near) random walk and using Bayesian methods. However, it is unclear whether their estimates are reliable if the parameters have a different mechanism of evolution or experience abrupt breaks in their path. Li and Yang (2011) and Ang and Kristensen (2012) estimate time-varying parameters assuming that they are deterministic smooth functions of time, but their studies are

¹⁰The R code used for the empirical application is available on the author's website (<https://sites.google.com/view/mikihito-nishi/home>).

restricted to smooth functional parameters that belong to type-a TVP(1). In contrast, the kernel-based method with $\gamma = -1/2$ accommodates many important models of time-varying parameters, including smooth functions of time, the rescaled random walk, abrupt breaks and their mixtures.

5.2 Data

All data are extracted from Kenneth French’s website (<https://mba.tuck.dartmouth.edu/pages/faculty/ken.french>). Following Li, Su and Xu (2015), we form three portfolios denoted by G, V, and G-V, respectively, from the 25 size-B/M portfolios. G is the average of the five portfolios in the lowest B/M quintile, V is the average of the five portfolios in the highest B/M quintile, and V-G is simply their difference. All the data are monthly, spanning 1952:1-2019:12 ($T = 816$).

5.3 Results

We apply kernel regression with $\gamma = -1/2$ to the three portfolios. We use the Epanechnikov kernel and $\hat{\Omega}_t = (Th)^{-1} \sum_{i=1}^T K((t-i)/Th)x_i x_i'$ and $\hat{\Sigma}_t = (Th)^{-1} \sum_{i=1}^T K((t-i)/Th)^2 \hat{\varepsilon}_i^2 x_i x_i'$ as the variance estimators, where $x_t = (1, R_{M,t})'$. Bandwidth h is set to $h = T^{-1/2}$. To save space, we only discuss the result for portfolio V-G. The results for portfolios G and V are given in Appendix D.

5.3.1 Comparison with the Bayesian estimate

Recall that our choice of the bandwidth ($\gamma = -1/2$) is intended to accommodate parameters following the (rescaled) random walk. To evaluate to what extent this choice accommodates random walk dynamics in parameters, we compare the estimates obtained from the kernel-based method with those obtained from a Bayesian procedure in which parameters are assumed to be the random walk.

Let $\theta_t := (\alpha_t, \beta_t)'$. In the Bayesian method, we estimate the time-varying alpha and beta by using the Markov Chain Monte Carlo algorithm, assuming that $\theta_t = \theta_{t-1} + u_t$, where $u_t \sim N(0, D^2)$ with $D^2 = \text{diag}(D_1^2, D_2^2)$.¹¹ As the prior distributions for parameters θ_0 , D and $\text{Var}(\varepsilon_t) = \sigma_\varepsilon^2$, we suppose $\theta_0 \sim N(\mu \mathbf{1}_2, \sigma^2 \mathbf{I}_2)$, $D_i \sim \text{Gamma}(v_1, v_2)$, $i = 1, 2$, and $\sigma_\varepsilon \sim \text{Gamma}(\nu_1, \nu_2)$. Hyperparameters are set to $(\mu, \sigma, v_1, v_2, \nu_1, \nu_2) = (0, 32, 2, 10^{-4}, 2, 10^{-4})$.¹²

¹¹For computation, we use the R package `walker` developed by Helske (2023).

¹²The parameter values used for (v_1, v_2, ν_1, ν_2) are the default values in the R package `walker`.

In Figure 1, we compare the estimates obtained from this Bayesian method with those from the kernel method with $h = T^{-1/2}$. For the estimated alpha (Figure 1a), the trajectory obtained from the kernel method is more volatile (with a larger amplitude) than that obtained from the Bayesian algorithm, but the two trajectories seem to share the same frequency. More striking is the similarity between the estimates of the time-varying beta. The estimated trajectories obtained from the two distinct methods are similar on the whole line. In particular, it is remarkable that the peaks and troughs are similar between the two trajectories.

As a robustness check, we also perform the Bayesian estimation with a different set of hyperparameters. Specifically, we experiment with $(v_1, v_2) = (1, 10^{-4}), (4, 10^{-4})$ for the priors of D_i , $i = 1, 2$, and the other hyperparameters are unchanged.¹³ Figure 1 shows the results. For the time-varying alpha (Figure 1a), the Bayesian estimates with $v_1 = 4$ are more volatile and similar to the trajectory obtained from the kernel method, while the estimates are smoother with $v_1 = 1$. For the time-varying beta, the kernel and Bayesian methods continue to produce similar trajectories irrespective of the value of v_1 .

5.3.2 Interval estimation

Based on the normal approximation given in Theorem 1, we calculate the confidence band for the kernel-based estimator.¹⁴

In Figure 2a, we plot the estimated time-varying alpha and its confidence band. The estimated alpha fluctuates around the value zero on the whole line, and the confidence band includes zero at almost all time points. Figure 2b depicts the estimated time-varying beta. It starts with a positive value that is significantly different from zero and then fluctuates around zero up to about $t = 300$. Then, it starts to decrease, and, from about $t = 340$ to $t = 610$, it stays below zero, and the confidence band does not include the value zero. From $t = 580$, it starts to increase and fluctuates around zero from $t = 630$ till the end of the sample.

5.3.3 Estimates obtained from the conventional $T^{-1/5}$ -rate bandwidth

We also apply kernel regression using the conventional bandwidth given by $\gamma = -1/5$. Figure 3 shows the estimates obtained from this method (with $h = T^{-1/5}$). For both the time-varying alpha and beta, the estimates are smoother than those obtained from the kernel

¹³We also changed the values for $(\mu, \sigma, \nu_1, \nu_2)$, but the estimates were insensitive to the values of these parameters.

¹⁴This confidence band is obtained by sequentially calculating the pointwise 95% confidence intervals and is not the uniform 95% confidence band.

method with $h = T^{-1/2}$ and the Bayesian method. In the estimation with $h = T^{-1/5}$, the parameter value at each time point is estimated by using a wider range of observations than in the kernel method with $h = T^{-1/2}$. Specifically, the number of observations used for the estimation at each time point is $\lfloor Th \rfloor = 213$ for the case of $h = T^{-1/5}$, while it is $\lfloor Th \rfloor = 28$ for the case of $h = T^{-1/2}$. This results in smoothing out short-term fluctuations in the parameter path and obtaining smoother estimates. Those estimates, however, are neither \sqrt{Th} -consistent nor asymptotically normal if the true time-varying parameters follow the random walk, or experience abrupt breaks or threshold effects given by $\alpha \leq 2/5$; see the definition of $A(\gamma)$ in (7). The kernel-based estimator with $\gamma = -1/5$ is more prone to the misspecification of the time-varying parameter than that with $\gamma = -1/2$.

6 Conclusion

We considered the estimation of linear models with time-varying parameters, where the time-varying parameters are allowed to have a wide range of smoothness. To do this, we set up a general framework in which the smoothness of nonparametric time-varying parameters is quantified by a single parameter α . We showed that the kernel-based estimator can consistently estimate many important time-varying parameter models, including continuously differentiable functions, the rescaled random walk, abrupt structural breaks, the threshold regression model, and their mixtures. Our analysis revealed that a trade-off between efficiency and robustness is inherent in kernel-based estimation, and the choice of the bandwidth determines the balance between them. Within a textbook framework where the (infinite-dimensional) parameter of interest is (continuously) differentiable, the role of the bandwidth pertaining to the bias-variance trade-off has been repeatedly emphasized. When the parameter is possibly nondifferentiable, which is a popular assumption in the time series setting, the bandwidth determines the rate of convergence and the size of the class of time-varying parameters that can be estimated, between which a trade-off does exist.

Using this result, we demonstrated through theoretical and numerical analyses that the kernel-based estimation delivers invalid inference if an improper bandwidth is selected. In particular, the conventional choice of a $T^{-1/5}$ -rate bandwidth yields misleading inference in the case of nondifferentiable time-varying parameters such as the random walk. Another important implication from our result is that abrupt breaks of a certain magnitude (that depends on the bandwidth used) lead to biased estimation.

Given the diversity of existing time-varying parameter models (differentiable, continuous but nondifferentiable, or discontinuous) and the possibility that time-varying coefficients are a mixture of several models, it is often difficult to specify the evolutionary mechanism of time-varying parameters correctly. Therefore, we proposed a specific choice of the bandwidth that accommodates many important models for time-varying parameters, including smooth functions, the rescaled random walk, small to moderate abrupt breaks and threshold effects, and their mixtures. Our empirical application showed that the kernel-based estimator with this choice can capture the random walk type dynamics in time-varying parameters.

References

- Andrews, D. W. K. (1993) Tests for Parameter Instability and Structural Change With Unknown Change Point. *Econometrica*, 61 (4), 821–856.
- Ang, A. and J. Chen (2007) CAPM over the Long Run: 1926–2001. *Journal of Empirical Finance*, 14 (1), 1–40.
- Ang, A. and D. Kristensen (2012) Testing Conditional Factor Models. *Journal of Financial Economics*, 106 (1), 132–156.
- Bai, J. and P. Perron (1998) Estimating and Testing Linear Models with Multiple Structural Changes. *Econometrica*, 66 (1), 47–78.
- Cai, Z. (2007) Trending Time-Varying Coefficient Time Series Models with Serially Correlated Errors. *Journal of Econometrics*, 136 (1), 163–188.
- Casini, A. and P. Perron (2018) Structural Breaks in Time Series. May.
- Chen, B. and Y. Hong (2012) Testing for Smooth Structural Changes in Time Series Models via Nonparametric Regression. *Econometrica*, 80 (3), 1157–1183.
- Cogley, T. and T. J. Sargent (2005) Drifts and Volatilities: Monetary Policies and Outcomes in the Post WWII US. *Review of Economic Dynamics*, 8 (2), 262–302.
- Cushman, D. O. (1988) U.S. Bilateral Trade Flows and Exchange Risk during the Floating Period. *Journal of International Economics*, 24 (3), 317–330.
- Davidson, J. (1994) *Stochastic Limit Theory: An Introduction for Econometricians* Oxford University Press.

- Fama, E. F. and K. R. French (2006) The Value Premium and the CAPM. *The Journal of Finance*, 61 (5), 2163–2185.
- Friedrich, M. and Y. Lin (2022) Sieve Bootstrap Inference for Linear Time-Varying Coefficient Models. *Journal of Econometrics*.
- Georgiev, I., D. I. Harvey, S. J. Leybourne, and A. M. R. Taylor (2018) Testing for Parameter Instability in Predictive Regression Models. *Journal of Econometrics*, 204 (1), 101–118.
- Ghysels, E. (1998) On Stable Factor Structures in the Pricing of Risk: Do Time-Varying Betas Help or Hurt? *The Journal of Finance*, 53 (2), 549–573.
- Giraitis, L., G. Kapetanios, and T. Yates (2014) Inference on Stochastic Time-Varying Coefficient Models. *Journal of Econometrics*, 179 (1), 46–65.
- Giraitis, L., G. Kapetanios, and M. Marcellino (2021) Time-Varying Instrumental Variable Estimation. *Journal of Econometrics*, 224 (2), 394–415.
- Guo, H., C. Wu, and Y. Yu (2017) Time-Varying Beta and the Value Premium. *The Journal of Financial and Quantitative Analysis*, 52 (4), 1551–1576.
- Hansen, B. E. (2000) Sample Splitting and Threshold Estimation. *Econometrica*, 68 (3), 575–603.
- Helske, J. (2023) Walker: Bayesian Generalized Linear Models with Time-Varying Coefficients.
- Inoue, A., L. Jin, and B. Rossi (2017) Rolling Window Selection for Out-of-Sample Forecasting with Time-Varying Parameters. *Journal of Econometrics*, 196 (1), 55–67.
- Kristensen, D. (2012) Non-Parametric Detection and Estimation of Structural Change. *The Econometrics Journal*, 15 (3), 420–461.
- Lewellen, J. and S. Nagel (2006) The Conditional CAPM Does Not Explain Asset-Pricing Anomalies. *Journal of Financial Economics*, 82 (2), 289–314.
- Li, Y., L. Su, and Y. Xu (2015) A Combined Approach to the Inference of Conditional Factor Models. *Journal of Business & Economic Statistics*, 33 (2), 203–220.
- Li, Y. and L. Yang (2011) Testing Conditional Factor Models: A Nonparametric Approach. *Journal of Empirical Finance*, 18 (5), 972–992.

- Müller, U. K. and P.-E. Petalas (2010) Efficient Estimation of the Parameter Path in Unstable Time Series Models. *The Review of Economic Studies*, 77 (4), 1508–1539.
- Nyblom, J. (1989) Testing for the Constancy of Parameters Over Time. *Journal of the American Statistical Association*, 84 (405), 223–230.
- Pesaran, M. H., A. Pick, and M. Pranovich (2013) Optimal Forecasts in the Presence of Structural Breaks. *Journal of Econometrics*, 177 (2), 134–152.
- Pesaran, M. H. and A. Timmermann (2007) Selection of Estimation Window in the Presence of Breaks. *Journal of Econometrics*, 137 (1), 134–161.
- Pivetta, F. and R. Reis (2007) The Persistence of Inflation in the United States. *Journal of Economic Dynamics and Control*, 31 (4), 1326–1358.
- Robinson, P. M. (1989) Nonparametric Estimation of Time-Varying Parameters. in Hackl, P. ed. *Statistical Analysis and Forecasting of Economic Structural Change*, 253–264, Berlin, Heidelberg Springer.
- Sun, Y., Y. Hong, S. Wang, and X. Zhang (2023) Penalized Time-Varying Model Averaging. *Journal of Econometrics*, 235 (2), 1355–1377.
- Zhang, T. and W. B. Wu (2012) Inference of Time-Varying Regression Models. *The Annals of Statistics*, 40 (3), 1376–1402.

Table 1: Mean MSE and coverage probability (CP) when ε_t is i.i.d.

γ	MSE				CP ($t = 0.5T$)				
	T				T				
	100	200	400	800	100	200	400	800	
(i)	-0.2	0.069	0.055	0.043	0.036	0.626	0.538	0.461	0.395
	-0.33	0.056	0.039	0.027	0.019	0.777	0.746	0.734	0.709
	-0.5	0.073	0.048	0.032	0.022	0.850	0.853	0.874	0.899
	-0.55	0.087	0.058	0.040	0.028	0.842	0.876	0.886	0.914
	-0.6	0.107	0.074	0.053	0.038	0.837	0.866	0.884	0.910
	-0.7	0.198	0.138	0.103	0.077	0.792	0.835	0.848	0.872
(ii)	-0.2	0.070	0.054	0.044	0.036	0.628	0.553	0.460	0.373
	-0.33	0.056	0.039	0.027	0.020	0.790	0.771	0.736	0.696
	-0.5	0.073	0.048	0.032	0.022	0.853	0.865	0.877	0.906
	-0.55	0.087	0.058	0.040	0.028	0.852	0.874	0.881	0.906
	-0.6	0.107	0.074	0.053	0.038	0.843	0.875	0.882	0.903
	-0.7	0.198	0.138	0.103	0.077	0.791	0.828	0.850	0.865

Note: $\beta_{T,t} = T^{-1/2} \sum_{i=1}^T v_i$, where $v_i \sim$ i.i.d. $N(0, 1)$ for case (i) and v_i is log-normally distributed with $\mu = 0$, $\sigma = 1$ for case (ii). $\hat{\beta}_t$ is calculated using bandwidth parameter $h = T^\gamma$.

Table 2: Mean MSE and coverage probability (CP) when ε_t is GARCH

	γ	MSE				CP ($t = 0.5T$)			
		T				T			
		100	200	400	800	100	200	400	800
(i)	-0.2	0.070	0.054	0.043	0.036	0.603	0.529	0.460	0.394
	-0.33	0.057	0.039	0.027	0.019	0.753	0.736	0.727	0.702
	-0.5	0.074	0.048	0.032	0.022	0.847	0.866	0.886	0.900
	-0.55	0.090	0.059	0.040	0.028	0.847	0.878	0.893	0.912
	-0.6	0.111	0.075	0.053	0.038	0.844	0.878	0.889	0.918
	-0.7	0.206	0.141	0.104	0.077	0.819	0.855	0.868	0.890
(ii)	-0.2	0.070	0.054	0.044	0.036	0.604	0.525	0.459	0.376
	-0.33	0.057	0.039	0.027	0.019	0.770	0.759	0.725	0.695
	-0.5	0.074	0.048	0.032	0.022	0.862	0.874	0.881	0.903
	-0.55	0.089	0.059	0.040	0.028	0.858	0.879	0.892	0.906
	-0.6	0.111	0.075	0.053	0.038	0.855	0.885	0.887	0.909
	-0.7	0.206	0.141	0.104	0.077	0.815	0.850	0.863	0.884

Note: $\beta_{T,t} = T^{-1/2} \sum_{i=1}^T v_i$, where $v_i \sim$ i.i.d. $N(0, 1)$ for case (i) and v_i is log-normally distributed with $\mu = 0$, $\sigma = 1$ for case (ii). $\hat{\beta}_t$ is calculated using bandwidth parameter $h = T^\gamma$.

Table 3: MSE, coverage probability and mean interval length when ε_t is i.i.d


α	MSE					Coverage Probability					Interval Length							
	τ	0.4	0.45	0.5	0.55	0.6	τ	0.4	0.45	0.5	0.55	0.6	τ	0.4	0.45	0.5	0.55	0.6
(i)	0.1	0.07	0.11	0.44	0.16	0.07	0.84	0.80	0.36	0.72	0.85	0.79	0.85	0.89	0.88	0.81		
	0.2	0.07	0.08	0.22	0.10	0.07	0.84	0.82	0.54	0.77	0.84	0.77	0.80	0.82	0.82	0.79		
	0.3	0.07	0.07	0.13	0.08	0.07	0.83	0.84	0.69	0.80	0.84	0.77	0.78	0.79	0.79	0.78		
	0.4	0.07	0.07	0.09	0.07	0.07	0.83	0.84	0.77	0.81	0.83	0.77	0.77	0.78	0.78	0.78		
	0.5	0.07	0.07	0.08	0.07	0.07	0.83	0.84	0.80	0.83	0.83	0.77	0.77	0.77	0.78	0.78		
(ii)	0.1	0.04	0.05	0.39	0.06	0.05	0.86	0.86	0.30	0.83	0.85	0.66	0.71	0.76	0.72	0.66		
	0.2	0.04	0.04	0.17	0.05	0.05	0.86	0.86	0.52	0.84	0.85	0.66	0.67	0.69	0.68	0.66		
	0.3	0.04	0.04	0.09	0.05	0.05	0.86	0.87	0.70	0.85	0.85	0.66	0.66	0.67	0.67	0.65		
	0.4	0.04	0.04	0.06	0.05	0.05	0.86	0.87	0.79	0.85	0.85	0.66	0.66	0.66	0.66	0.65		
	0.5	0.04	0.04	0.05	0.05	0.05	0.86	0.87	0.83	0.85	0.85	0.66	0.66	0.66	0.66	0.65		
(iii)	0.1	0.03	0.03	0.33	0.03	0.03	0.87	0.87	0.21	0.88	0.86	0.57	0.58	0.65	0.58	0.56		
	0.2	0.03	0.03	0.12	0.03	0.03	0.87	0.86	0.51	0.87	0.86	0.57	0.57	0.59	0.57	0.56		
	0.3	0.03	0.03	0.06	0.03	0.03	0.87	0.86	0.74	0.87	0.86	0.57	0.57	0.57	0.56	0.56		
	0.4	0.03	0.03	0.04	0.03	0.03	0.87	0.86	0.82	0.87	0.86	0.57	0.56	0.57	0.56	0.56		
	0.5	0.03	0.03	0.03	0.03	0.03	0.87	0.86	0.86	0.87	0.86	0.57	0.56	0.57	0.56	0.56		
(iv)	0.1	0.02	0.02	0.29	0.02	0.02	0.88	0.89	0.13	0.87	0.88	0.48	0.48	0.55	0.48	0.48		
	0.2	0.02	0.02	0.09	0.02	0.02	0.88	0.89	0.48	0.87	0.88	0.48	0.48	0.50	0.48	0.48		
	0.3	0.02	0.02	0.04	0.02	0.02	0.88	0.89	0.75	0.87	0.88	0.48	0.48	0.48	0.48	0.48		
	0.4	0.02	0.02	0.03	0.02	0.02	0.88	0.89	0.85	0.87	0.88	0.48	0.48	0.48	0.48	0.48		
	0.5	0.02	0.02	0.02	0.02	0.02	0.88	0.89	0.88	0.87	0.88	0.48	0.48	0.48	0.48	0.48		

Note: $\beta_{T,t} = \mu_{T,t} + T^{-1/2} \sum_{i=1}^t v_i$, where $\mu_{T,t} = 0$ for $t \leq 0.5T$ and $\mu_{T,t} = 2/T^\alpha$ for $t > 0.5T$, and $v_i \sim$ i.i.d. $N(0, 1)$. $\beta_{T,t}$ with $t = \tau T$ is estimated using bandwidth $h = T^{-0.5}$. The sample size is $T = 100$ for case (i), $T = 200$ for case (ii) $T = 400$ for case (iii), and $T = 800$ for case (iv).

Table 4: MSE, coverage probability and mean interval length when ε_t is GARCH

α	MSE					Coverage Probability					Interval Length							
	τ	0.4	0.45	0.5	0.55	0.6	τ	0.4	0.45	0.5	0.55	0.6	τ	0.4	0.45	0.5	0.55	0.6
(i)	0.1	0.07	0.11	0.44	0.15	0.06	0.82	0.77	0.32	0.68	0.84	0.84	0.72	0.79	0.83	0.82	0.74	0.74
	0.2	0.07	0.09	0.22	0.10	0.06	0.81	0.79	0.48	0.74	0.82	0.82	0.71	0.74	0.75	0.75	0.75	0.72
	0.3	0.07	0.08	0.13	0.08	0.06	0.80	0.80	0.63	0.79	0.82	0.82	0.71	0.71	0.72	0.72	0.72	0.71
	0.4	0.07	0.07	0.09	0.07	0.06	0.80	0.81	0.73	0.79	0.82	0.82	0.70	0.70	0.70	0.70	0.70	0.71
	0.5	0.07	0.07	0.07	0.07	0.06	0.80	0.81	0.78	0.80	0.82	0.82	0.70	0.70	0.70	0.70	0.70	0.71
(ii)	0.1	0.05	0.05	0.40	0.06	0.04	0.84	0.85	0.26	0.82	0.84	0.84	0.62	0.67	0.72	0.67	0.61	0.61
	0.2	0.05	0.04	0.17	0.05	0.04	0.84	0.84	0.46	0.83	0.84	0.84	0.62	0.63	0.65	0.63	0.60	0.60
	0.3	0.05	0.04	0.09	0.05	0.04	0.84	0.84	0.65	0.83	0.84	0.84	0.61	0.62	0.62	0.61	0.60	0.60
	0.4	0.05	0.04	0.07	0.05	0.04	0.84	0.85	0.75	0.83	0.84	0.84	0.61	0.62	0.61	0.61	0.60	0.60
	0.5	0.05	0.04	0.06	0.05	0.04	0.84	0.84	0.80	0.83	0.84	0.84	0.61	0.61	0.61	0.61	0.60	0.60
(iii)	0.1	0.03	0.03	0.34	0.03	0.03	0.84	0.86	0.18	0.87	0.85	0.85	0.52	0.54	0.63	0.54	0.52	0.52
	0.2	0.03	0.03	0.13	0.03	0.03	0.84	0.85	0.44	0.86	0.85	0.85	0.52	0.53	0.56	0.53	0.52	0.52
	0.3	0.03	0.03	0.06	0.03	0.03	0.84	0.85	0.68	0.86	0.85	0.85	0.52	0.53	0.54	0.53	0.52	0.52
	0.4	0.03	0.03	0.04	0.03	0.03	0.84	0.85	0.79	0.86	0.85	0.85	0.52	0.53	0.53	0.52	0.52	0.52
	0.5	0.03	0.03	0.04	0.03	0.03	0.84	0.85	0.83	0.86	0.85	0.85	0.52	0.53	0.53	0.52	0.52	0.52
(iv)	0.1	0.02	0.02	0.29	0.02	0.02	0.86	0.88	0.12	0.86	0.86	0.86	0.45	0.45	0.53	0.45	0.45	0.45
	0.2	0.02	0.02	0.09	0.02	0.02	0.86	0.88	0.41	0.86	0.86	0.86	0.45	0.45	0.47	0.45	0.45	0.45
	0.3	0.02	0.02	0.04	0.02	0.02	0.86	0.88	0.69	0.86	0.86	0.86	0.45	0.45	0.45	0.45	0.45	0.45
	0.4	0.02	0.02	0.02	0.02	0.02	0.86	0.88	0.82	0.86	0.86	0.86	0.45	0.45	0.45	0.45	0.45	0.45
	0.5	0.02	0.02	0.02	0.02	0.02	0.86	0.88	0.85	0.86	0.86	0.86	0.45	0.45	0.45	0.45	0.45	0.45

Note: $\beta_{T,t} = \mu_{T,t} + T^{-1/2} \sum_{i=1}^t v_i$, where $\mu_{T,t} = 0$ for $t \leq 0.5T$ and $\mu_{T,t} = 2/T^\alpha$ for $t > 0.5T$, and $v_i \sim \text{i.i.d. } N(0, 1)$. $\beta_{T,t}$ with $t = \tau T$ is estimated using bandwidth $h = T^{-0.5}$. The sample size is $T = 100$ for case (i), $T = 200$ for case (ii) $T = 400$ for case (iii), and $T = 800$ for case (iv).




(a) Plot of the time-varying alpha




(b) Plot of the time-varying beta

Figure 1: Estimates from the kernel ($h = T^{-1/2}$) and Bayesian methods

—: Kernel, ·····: Bayesian ($v_1 = 1$), - - -: Bayesian ($v_1 = 2$), - · -: Bayesian ($v_1 = 4$)




(a) Plot of the time-varying alpha




(b) Plot of the time-varying beta

Figure 2: Estimates and 95% confidence band from the kernel-based method ($h = T^{-1/2}$) for V-G

(Horizontal lines in (a) and (b) indicate the value zero.)



(a) Plot of the time-varying alpha



(b) Plot of the time-varying beta

Figure 3: Estimates and 95% confidence band obtained from the conventional bandwidth ($h = T^{-1/5}$) for V-G

Appendix to “Estimating Time-Varying Parameters of Various Smoothness in Linear Models via Kernel Regression” by M.Nishi

Appendix A: Proofs of the Main Results

Lemma A.1. Under Assumptions 1 and 2, for each $t = \lfloor Tr \rfloor$, $r \in (0, 1)$, we have

$$\left\| \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' (\beta_{T,i} - \beta_{T,t}) \right\| = \begin{cases} O_p(Th^{1+\alpha}) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(a)} \\ O_p(T^{1-\alpha}h) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(b)} \end{cases}.$$

Proof. First, we have

$$\begin{aligned} \left\| \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' (\beta_{T,i} - \beta_{T,t}) \right\| &= \left\| \sum_{i=t-\lfloor Th \rfloor}^{t+\lfloor Th \rfloor} K\left(\frac{t-i}{Th}\right) x_i x_i' (\beta_{T,i} - \beta_{T,t}) \right\| \\ &\leq \max_{t-\lfloor Th \rfloor \leq i \leq t+\lfloor Th \rfloor} \|\beta_{T,i} - \beta_{T,t}\| \\ &\quad \times \sum_{i=t-\lfloor Th \rfloor}^{t+\lfloor Th \rfloor} K\left(\frac{t-i}{Th}\right) \|x_i x_i'\|, \end{aligned} \quad (\text{A.1})$$

because the support of K is $[-1, 1]$ under Assumption 2. Note that

$$\sum_{i=t-\lfloor Th \rfloor}^{t+\lfloor Th \rfloor} K\left(\frac{t-i}{Th}\right) \|x_i x_i'\| = O_p(Th), \quad (\text{A.2})$$

because

$$\begin{aligned} E \left[\sum_{i=t-\lfloor Th \rfloor}^{t+\lfloor Th \rfloor} K\left(\frac{t-i}{Th}\right) \|x_i x_i'\| \right] &\leq \max_i E[\|x_i x_i'\|] \sum_{i=t-\lfloor Th \rfloor}^{t+\lfloor Th \rfloor} K\left(\frac{t-i}{Th}\right) \\ &\leq \sup_t E[\|x_t\|^2] \times Th \times \frac{1}{Th} \sum_{i=-\lfloor Th \rfloor}^{\lfloor Th \rfloor} K\left(\frac{i}{Th}\right) \\ &= O(1) \times Th \sum_{i=-\lfloor Th \rfloor}^{\lfloor Th \rfloor} \int_{(i-1)/Th}^{i/Th} K\left(\frac{i}{Th}\right) dr \\ &= O(1) \times Th \sum_{i=-\lfloor Th \rfloor}^{\lfloor Th \rfloor} \int_{(i-1)/Th}^{i/Th} \left\{ K\left(\frac{i}{Th}\right) - K(r) + K(r) \right\} dr \\ &= O(1) \times Th \left(\int_{-\lfloor Th \rfloor/Th}^{\lfloor Th \rfloor/Th} K(r) dr + O(1/Th) \right) = O(Th), \end{aligned}$$

because $\sup_t E[|x_t|^2] < \infty$ under Assumption 1, K is Lipschitz continuous and $\int_{-1}^1 K(x)dx = 1$ under Assumption 2. We also have

$$\max_{t-[Th] \leq i \leq t+[Th]} \|\beta_{T,i} - \beta_{T,t}\| = \begin{cases} O_p(h^\alpha) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(a)} \\ O_p(T^{-\alpha}) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(b)} \end{cases}. \quad (\text{A.3})$$

Substituting (A.2) and (A.3) into (A.1), we deduce

$$\left\| \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' (\beta_{T,i} - \beta_{T,t}) \right\| = \begin{cases} O_p(Th^{1+\alpha}) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(a)} \\ O_p(T^{1-\alpha}h) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(b)} \end{cases}.$$

□

Lemma A.2. Under Assumptions 1 and 2, for each $t = \lfloor Tr \rfloor$, $r \in (0, 1)$, we have

$$\frac{1}{\sqrt{Th}} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i \varepsilon_i \xrightarrow{d} N(0, \Sigma(r)).$$

Proof. To prove this result, we use the Cramer-Wold device. Define $z_{T,i}^* := \lambda' K((t-i)/Th) x_i \varepsilon_i$, where $\lambda \neq 0$ is any $p \times 1$ vector, $\sigma_T^2 := \text{Var}(\sum_{i=1}^T z_{T,i}^*)$, and $z_{T,i} := z_{T,i}^*/\sigma_T$. Note that $\sigma_T^2/Th \rightarrow \lambda' \Sigma(r) \lambda$ by Assumption 1(c). Moreover, define positive constant array $\{c_{T,i}\}$ as

$$c_{T,i} = \begin{cases} \max\left\{\sqrt{\text{Var}(z_{T,i}^*)}, 1\right\}/\sigma_T & \text{for } i \in [t - \lfloor Th \rfloor, t + \lfloor Th \rfloor] \\ 1/\sqrt{T} & \text{otherwise} \end{cases}.$$

To show Lemma A.2, we rely on Theorem 2 of de Jong (1997), which requires that the following conditions hold for $\{z_{T,i}, c_{T,i}\}$:

- (i) $z_{T,i}$ has mean zero, and $\text{Var}(\sum_{i=1}^T z_{T,i}) = 1$.
- (ii) $z_{T,i}/c_{T,i}$ is L_r -bounded for some $r > 2$ uniformly in i and T .
- (iii) $z_{T,i}$ is L_2 -NED of size $-1/2$ on an α -mixing array of size $-r/(r-2)$, with respect to some constants $d_{T,i}$. Moreover, $d_{T,i}/c_{T,i}$ is bounded uniformly in i and T .
- (iv) Let b_T be a positive non-decreasing integer-valued sequence such that $b_T \leq T$, $b_T \rightarrow \infty$, and $b_T/T \rightarrow 0$ as $T \rightarrow \infty$. Also let $r_T := \lfloor T/b_T \rfloor$. Define $M_{T,j} := \max_{(j-1)b_T+1 \leq i \leq j b_T} c_{T,i}$, $j = 1, \dots, r_T$, and $M_{T,r_T+1} := \max_{r_T b_T+1 \leq i \leq T} c_{T,i}$. Then, we have $\max_{1 \leq j \leq r_T+1} M_{T,j} = o(b_T^{-1/2})$ and $\sum_{j=1}^{r_T} M_{T,j}^2 = O(b_T^{-1})$.

Conditions (i)-(iv) imply that $\sum_{i=1}^T z_{t,i} \xrightarrow{d} N(0,1)$. We show that the above four conditions hold.

(i) This condition trivially follows from Assumption 1(b) and the definition of $z_{T,i}$.

(ii) Noting that $z_{T,i}^* = 0$ for $i < t - \lfloor Th \rfloor$ and $i > t + \lfloor Th \rfloor$, we have

$$z_{T,i}/c_{T,i} = \begin{cases} z_{T,i}^*/\max\left\{\sqrt{\text{Var}(z_{T,i}^*)}, 1\right\} & \text{for } i \in [t - \lfloor Th \rfloor, t + \lfloor Th \rfloor] \\ 0 & \text{otherwise} \end{cases}. \quad (\text{A.4})$$

Because (x'_i, ε_i) is uniformly L_{2r} -bounded for $r > 2$ by Assumption 1(a), $z_{T,i}^*$ is L_r -bounded uniformly in i and T since kernel $K(\cdot)$ is bounded. This implies that $z_{T,i}/c_{T,i}$ is also L_r -bounded uniformly in i and T in view of (A.4).

(iii) Note that (x'_i, ε_i) is L_{2r} -bounded and L_2 -NED of size $-(r-1)/(r-2)$ on an α -mixing sequence of size $-r/(r-2)$. Thus, following the argument of Example 17.17 of Davidson (1994), we can show that $z_{T,i}$ is L_2 -NED of size $-1/2$ on the same α -mixing sequence, with respect to positive constant array $d_{T,i}$ satisfying

$$\sup_{t-\lfloor Th \rfloor \leq i \leq t+\lfloor Th \rfloor} d_{T,i} \leq \frac{C}{\sigma_T} = O\left(\frac{1}{\sqrt{Th}}\right),$$

for some positive constant $C < \infty$ independent of T , and $d_{T,i} = 0$ for $i \notin [t - \lfloor Th \rfloor, t + \lfloor Th \rfloor]$. This follows from the fact that $z_{T,i}^*$ is L_2 -NED of size $-1/2$ with respect to positive constant array $d_{T,i}^*$ satisfying $\sup_{T,i} d_{T,i}^* < \infty$ due to $\sup_t d_t < \infty$ by Assumption 1(a), $\sigma_T^2 = O(Th)$, and $K((t-i)/Th) = 0$ for $i < t - \lfloor Th \rfloor$ and $i > t + \lfloor Th \rfloor$. This implies that $d_{T,i}/c_{T,i}$ is bounded uniformly in i and T .

(iv) Let $b_T = \sqrt{Th}$. Then, by the definition of $c_{T,i}$ and the fact that $\sigma_T = O(\sqrt{Th})$ and $\text{Var}(z_{T,i}^*) < \infty$ uniformly in i and T by Assumption 1(a), we get

$$\max_{1 \leq j \leq r_{T+1}} M_{T,j} = O((Th)^{-1/2}) = o(b_T^{-1/2}).$$

Furthermore, letting $j_1 := \lfloor (t - \lfloor Th \rfloor)/b_T \rfloor$ and $j_2 := \lfloor (t + \lfloor Th \rfloor)/b_T \rfloor$, we obtain

$$\begin{aligned} \sum_{j=1}^{r_T} M_{T,j}^2 &= \sum_{j=1}^{j_1} M_{T,j}^2 + \sum_{j=j_1+1}^{j_2} M_{T,j}^2 + \sum_{j=j_2+1}^{r_T} M_{T,j}^2 \\ &= \frac{j_1}{T} + O\left(\frac{j_2 - j_1}{Th}\right) + \frac{r_T - j_2}{T} = O(b_T^{-1}). \end{aligned}$$

Now that conditions (i)-(iv) are shown to hold, we obtain

$$\sum_{i=1}^T z_{T,i} = \lambda' \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i \varepsilon_i / \sigma_T \xrightarrow{d} N(0, 1).$$

Moreover, we have

$$\frac{1}{Th} \sigma_T^2 = \frac{1}{Th} \lambda' \text{Var} \left(\sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i \varepsilon_i \right) \lambda \rightarrow \lambda' \Sigma(r) \lambda,$$

by Assumption 1(c). This implies that

$$\lambda' \frac{1}{\sqrt{Th}} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i \varepsilon_i \xrightarrow{d} N(0, \lambda' \Sigma(r) \lambda).$$

By the Cramer-Wold device, we deduce

$$\frac{1}{\sqrt{Th}} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i \varepsilon_i \xrightarrow{d} N(0, \Sigma(r)).$$

□

Proof of Theorem 1. Since

$$\begin{aligned} \hat{\beta}_t &= \left(\frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' \right)^{-1} \frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' \beta_{T,i} \\ &\quad + \left(\frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' \right)^{-1} \frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i \varepsilon_i, \end{aligned}$$

we have

$$\sqrt{Th}(\hat{\beta}_t - \beta_{T,t} - R_{T,t}) = \left(\frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' \right)^{-1} \frac{1}{\sqrt{Th}} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i \varepsilon_i,$$

where $R_{T,t} := \left(\frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' \right)^{-1} \frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' (\beta_{T,i} - \beta_{T,t})$. It follows from Assumption 1(c) and Lemma A.2 that

$$\begin{aligned} \left(\frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' \right)^{-1} \frac{1}{\sqrt{Th}} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i \varepsilon_i &\xrightarrow{d} \Omega(r)^{-1} \times N(0, \Sigma(r)) \\ &= N(0, \Omega(r)^{-1} \Sigma(r) \Omega(r)^{-1}). \end{aligned}$$

The bias term, $R_{T,t}$, satisfies

$$R_{T,t} = \begin{cases} O_p(h^\alpha) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(a)} \\ O_p(T^{-\alpha}) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(b)} \end{cases},$$

by Assumption 1(c) and Lemma A.1.

Set $h = cT^\gamma$ for some $c > 0$ and $\gamma \in (-1, 0)$. Because $\sqrt{Th}R_{T,t} = O_p(T^{1/2+\gamma(1/2+\alpha)})$ for the type-a TVP(α) case and $\sqrt{Th}R_{T,t} = O_p(T^{1/2-\alpha+\gamma/2})$ for the type-b TVP(α) case, $\sqrt{Th}R_{T,t} = o_p(1)$ if

$$\gamma \in \begin{cases} (-1, -\frac{1}{2\alpha+1}) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(a)} \\ (-1, 2\alpha - 1) \cap (-1, 0) & \text{if } \beta_{T,t} \text{ satisfies Definition 1(b)} \end{cases},$$

under which choice we obtain

$$\begin{aligned} \sqrt{cT^{1+\gamma}}(\hat{\beta}_t - \beta_{T,t}) &= \left(\frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' \right)^{-1} \frac{1}{\sqrt{Th}} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i \varepsilon_i + o_p(1) \\ &\xrightarrow{d} N(0, \Omega(r)^{-1} \Sigma(r) \Omega(r)^{-1}). \end{aligned}$$

□

Proof of Corollary 1. We show that Assumption 1(c) holds with $\Omega(r) = \Omega$ and $\Sigma(r) = \Sigma$ under Assumptions 1(a)-(b) and 2 and covariance-stationarity.

First, decompose $(1/Th) \sum_{i=1}^T K((t-i)/Th) x_i x_i'$ as

$$\begin{aligned} \frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i x_i' &= \frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) E[x_1 x_1'] + \frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) (x_i x_i' - E[x_1 x_1']) \\ &=: A_{T,1} + A_{T,2}. \end{aligned}$$

We show $A_{T,1} \xrightarrow{p} E[x_1 x_1']$ and $A_{T,2} = o_p(1)$. For $A_{T,1}$, we have

$$\begin{aligned} A_{T,1} &= E[x_1 x_1'] \frac{1}{Th} \sum_{i=-\lfloor Th \rfloor}^{\lfloor Th \rfloor} K\left(\frac{i}{Th}\right) \\ &= E[x_1 x_1'] \int_{-\lfloor Th \rfloor/Th}^{\lfloor Th \rfloor/Th} K(r) dr + O(1/Th) \\ &\rightarrow E[x_1 x_1'] \int_{-1}^1 K(r) dr = E[x_1 x_1'], \end{aligned}$$

by Assumption 2. To show $A_{T,2} = o_p(1)$, note that $\{K((t-i)/Th)(x_i x'_i - E[x_1 x'_1])\}_i$ is a L_r -bounded ($r > 2$), mean-zero L_2 -NED triangular array by Assumption 1(b) and covariance-stationarity of x_i , and thus is a uniformly integrable L_2 -mixingale (see Andrews, 1988). This result allows us to apply the law of large numbers (see Andrews (1988), p.464) and obtain

$$A_{T,2} = \frac{1}{Th} \sum_{i=0}^{\lfloor Th \rfloor} K\left(\frac{i}{Th}\right) (x_{t-i} x'_{t-i} - E[x_1 x'_1]) + \frac{1}{Th} \sum_{i=1}^{\lfloor Th \rfloor} K\left(\frac{-i}{Th}\right) (x_{t+i} x'_{t+i} - E[x_1 x'_1])$$

$$\xrightarrow{p} 0,$$

which, together with $A_{T,1} \xrightarrow{p} \Omega$, shows that $(1/Th) \sum_{i=1}^T K((t-i)/Th) x_i x'_i \xrightarrow{p} \Omega$. Similarly, noting that $x_t \varepsilon_t$ is serially uncorrelated under Assumption 1, we have

$$\text{Var}\left(\frac{1}{\sqrt{Th}} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right) x_i \varepsilon_i\right) = \frac{1}{Th} \sum_{i=1}^T K\left(\frac{t-i}{Th}\right)^2 E[\varepsilon_1^2 x_1 x'_1]$$

$$\rightarrow \int_{-1}^1 K(x)^2 dx E[\varepsilon_1^2 x_1 x'_1],$$

since $x_t \varepsilon_t$ is covariance-stationary. □

Appendix B: MSE-Minimizing Bandwidth in the Case of Rescaled Random Walk Coefficients

In this appendix, we show that the bandwidth that minimizes the MSE of the kernel-based estimator is proportional to $T^{-1/2}$. In what follows, we will assume that Th is an integer for simplicity.

B.1 A simple case

To gain some insight, we begin with the following local-level model:

$$y_t = \beta_{T,t} + \varepsilon_t, \tag{B.1}$$

where $\beta_{T,t} = T^{-1/2} \sum_{i=1}^t u_i$.

Assumption B.1. (ε_t, u_t) is an *i.i.d.* sequence with mean zero and variance $\Sigma = \text{diag}(\sigma_\varepsilon^2, \sigma_u^2)$. Moreover, ε_t and u_t are independent.

We estimate $\beta_{T,t}$ using $\hat{\beta}_t$ with $K(\cdot)$ being the uniform kernel, that is, $\hat{\beta}_t = (2Th + 1)^{-1} \sum_{i=t-Th}^{t+Th} y_i$. Let $\text{MSE}(h) := E[(\hat{\beta}_t - \beta_{T,t})^2]$ denote the MSE of $\hat{\beta}_t$ as a function of bandwidth parameter h .

From model (B.1), $\hat{\beta}_t - \beta_{T,t}$ admits the following decomposition:

$$\begin{aligned} \hat{\beta}_t - \beta_{T,t} &= \frac{1}{2Th+1} \sum_{i=t-Th}^{t+Th} (\beta_{T,i} - \beta_{T,t}) + \frac{1}{2Th+1} \sum_{i=t-Th}^{t+Th} \varepsilon_i \\ &= -\frac{1}{2Th+1} \sum_{i=t-Th}^{t-1} \left(\frac{1}{\sqrt{T}} \sum_{k=i+1}^t u_k \right) + \frac{1}{2Th+1} \sum_{i=t+1}^{t+Th} \left(\frac{1}{\sqrt{T}} \sum_{k=t+1}^i u_k \right) + \frac{1}{2Th+1} \sum_{i=t-Th}^{t+Th} \varepsilon_i. \end{aligned}$$

Given that (ε_t, u_t) and (ε_s, u_s) ($t \neq s$) are independent, and also ε_t and u_t are independent, we have

$$\begin{aligned} \text{MSE}(h) &= \left(\frac{1}{2Th+1} \right)^2 \left\{ E \left[\left(\sum_{i=t-Th}^{t-1} \frac{1}{\sqrt{T}} \sum_{k=i+1}^t u_k \right)^2 \right] + E \left[\left(\sum_{i=t+1}^{t+Th} \frac{1}{\sqrt{T}} \sum_{k=t+1}^i u_k \right)^2 \right] + (2Th+1) \sigma_\varepsilon^2 \right\} \\ &= \left(\frac{1}{2Th+1} \right)^2 \left\{ \frac{1}{T} E \left[\left(\sum_{i=1}^{Th} (Th-i+1) u_{t-i+1} \right)^2 \right] + \frac{1}{T} E \left[\left(\sum_{i=1}^{Th} (Th-i+1) u_{t+i} \right)^2 \right] \right\} + \frac{\sigma_\varepsilon^2}{2Th+1} \\ &= \frac{2\sigma_u^2}{(2Th+1)^2 T} \frac{Th(Th+1)(2Th+1)}{6} + \frac{\sigma_\varepsilon^2}{2Th+1} \\ &= \frac{\sigma_u^2 h(1+o(1))}{6(1+o(1))} + \frac{\sigma_\varepsilon^2}{2Th(1+o(1))}. \end{aligned}$$

Ignoring the $1+o(1)$ terms, the MSE of $\hat{\beta}_t$ is asymptotically

$$\text{MSE}(h) = \frac{\sigma_u^2}{6} h + \frac{\sigma_\varepsilon^2}{2T} h^{-1}. \quad (\text{B.2})$$

Letting h_{\min} denote the minimizer of (B.2), it can be easily shown that

$$h_{\min} = \left(\frac{3\sigma_\varepsilon^2}{\sigma_u^2} \right)^{1/2} T^{-1/2}.$$

Therefore, the MSE-minimizing bandwidth is proportional to $T^{-1/2}$.

B.2 A general case

The argument above can be extended to the multiple regression. Suppose we are interested in the following model:

$$y_t = x_t' \beta_{T,t} + \varepsilon_t, \quad (\text{B.3})$$

where $\beta_{T,t} = T^{-1/2} \sum_{i=1}^t u_i$ is a p -dimensional rescaled random walk driven by $u_t = (u_{t,1}, \dots, u_{t,p})'$. For model (B.3), we impose the following assumption.

Assumption B.2. (a) $\{x_t\}_t$ is a p -dimensional stationary sequence with $E[x_1 x_1'] > 0$.

(b) $\{(\varepsilon_t, u_t')\}_t$ is a $(p+1)$ -dimensional i.i.d. sequence that is independent of $\{x_t\}_t$ and has mean zero and variance Σ with

$$\Sigma = \begin{pmatrix} \sigma_\varepsilon^2 & 0' \\ 0 & \sigma_u^2 I_p \end{pmatrix}.$$

Moreover, $\varepsilon_t, u_{t,1}, \dots, u_{t,p}$ are mutually independent.

(c) There exist nonrandom matrices $\Omega > 0$, Λ , $\bar{\Lambda}$, and Ξ such that $\Lambda \bar{\Lambda} + \bar{\Lambda} \Lambda - 2\Xi > 0$, $(2Th)^{-1} \sum_{i=t-Th}^{t+Th} x_i x_i' \xrightarrow{L_1} \Omega$, $(Th)^{-1} \sum_{i=t-Th}^{t-1} x_i x_i' \xrightarrow{L_1} \Lambda$, $(Th)^{-1} \sum_{i=1}^{Th} \left(\frac{i}{Th}\right) x_{t-i} x_{t-i}' \xrightarrow{L_1} \bar{\Lambda}$, and $(Th)^{-1} \sum_{i=t-Th}^{t-1} x_i x_i' (Th)^{-1} \sum_{j=t-Th}^i \frac{i-j}{Th} x_j x_j' \xrightarrow{L_1} \Xi$ as $T \rightarrow \infty$.

Assumptions B.2(a)-(b) extend Assumption B.1 to the case of the multiple regression. Assumption B.2(c) will hold if $x_t x_t' - E[x_t x_t']$ satisfies the condition of the L_1 law of large numbers of Andrews (1988). In particular, if $x_t = 1$ for all t , then we have $\Omega = 1$, $\Lambda = 1$, $\bar{\Lambda} = 1/2$, and $\Xi = 1/6$, and all the convergence results hold deterministically. The L_1 convergence assumed in (c) facilitates the derivation of the asymptotic expression of the MSE of $\hat{\beta}_t$.

The estimator of $\beta_{T,t}$ is $\hat{\beta}_t = (\sum_{i=t-Th}^{t+Th} x_i x_i')^{-1} \sum_{i=t-Th}^{t+Th} x_i y_i$, and the associated MSE is $\text{MSE}(h) = E[|\hat{\beta}_t - \beta_{T,t}|^2]$.

Proposition B.1. Consider model (B.3). Under Assumption B.2, we have

$$\begin{aligned} \text{MSE}(h) &= \frac{\sigma_u^2 h}{4} \text{tr}[\Omega^{-1}(\Lambda \bar{\Lambda} + \bar{\Lambda} \Lambda - 2\Xi)\Omega^{-1}](1 + o(1)) \\ &\quad + \frac{\sigma_\varepsilon^2}{2Th} \text{tr}[\Omega^{-1}](1 + o(1)). \end{aligned}$$

Checking the first and second order conditions, one can easily verify that the MSE-minimizing h is proportional to $T^{-1/2}$.

Proof. Note that

$$\hat{\beta}_t - \beta_{T,t} = \left(\sum_{i=t-Th}^{t+Th} x_i x_i' \right)^{-1} \left\{ \sum_{i=t-Th}^{t+Th} x_i x_i' (\beta_{T,i} - \beta_{T,t}) + \sum_{i=t-Th}^{t+Th} x_i \varepsilon_i \right\}.$$

The conditional MSE given $X_T := \{x_t\}_{t=1}^T$ is

$$\begin{aligned} & E[|\hat{\beta}_t - \beta_{T,t}|^2 | X_T] \\ &= E[\text{tr}[(\hat{\beta}_t - \beta_{T,t})(\hat{\beta}_t - \beta_{T,t})' | X_T]] \\ &= \text{tr} \left[\left(\sum_{i=t-Th}^{t+Th} x_i x_i' \right)^{-1} E \left[\left\{ - \sum_{i=t-Th}^{t-1} x_i x_i' \frac{1}{\sqrt{T}} \sum_{k=i+1}^t u_k + \sum_{i=t+1}^{t+Th} x_i x_i' \frac{1}{\sqrt{T}} \sum_{k=t+1}^i u_k + \sum_{i=t-Th}^{t+Th} x_i \varepsilon_i \right\} \right. \right. \\ &\quad \times \left. \left\{ - \sum_{i=t-Th}^{t-1} x_i x_i' \frac{1}{\sqrt{T}} \sum_{k=i+1}^t u_k + \sum_{i=t+1}^{t+Th} x_i x_i' \frac{1}{\sqrt{T}} \sum_{k=t+1}^i u_k + \sum_{i=t-Th}^{t+Th} x_i \varepsilon_i \right\}' | X_T \right] \\ &\quad \times \left. \left(\sum_{i=t-Th}^{t+Th} x_i x_i' \right)^{-1} \right] \\ &= \text{tr} \left[\left(\sum_{i=t-Th}^{t+Th} x_i x_i' \right)^{-1} E \left[\frac{1}{T} \left(\sum_{i=t-Th}^{t-1} x_i x_i' \sum_{k=i+1}^t u_k \right) \left(\sum_{i=t-Th}^{t-1} x_i x_i' \sum_{k=i+1}^t u_k \right)' \right. \right. \\ &\quad + \frac{1}{T} \left(\sum_{i=t+1}^{t+th} x_i x_i' \sum_{k=t+1}^i u_k \right) \left(\sum_{i=t+1}^{t+th} x_i x_i' \sum_{k=t+1}^i u_k \right)' \\ &\quad \left. \left. + \left(\sum_{i=t-Th}^{t+Th} x_i \varepsilon_i \right) \left(\sum_{i=t-Th}^{t+Th} x_i \varepsilon_i \right)' | X_T \right] \left(\sum_{i=t-Th}^{t+Th} x_i x_i' \right)^{-1} \right], \quad (\text{B.4}) \end{aligned}$$

where the last equality follows from Assumption B.2(b).

Consider each of the three terms in the conditional expectation in (B.4).

$$\begin{aligned}
& E \left[\frac{1}{T} \left(\sum_{i=t-Th}^{t-1} x_i x'_i \sum_{k=i+1}^t u_k \right) \left(\sum_{i=t-Th}^{t-1} x_i x'_i \sum_{k=i+1}^t u_k \right)' \middle| X_T \right] \\
&= \frac{1}{T} \sum_{i=t-Th}^{t-1} \sum_{j=t-Th}^{t-1} x_i x'_i E \left[\sum_{k=i+1}^t u_k \sum_{l=j+1}^t u'_l \right] x_j x'_j \\
&= \frac{1}{T} \sum_{i=t-Th}^{t-1} \sum_{j=t-Th}^i x_i x'_i \sum_{k=i+1}^t E[u_k u'_k] x_j x'_j + \frac{1}{T} \sum_{i=t-Th}^{t-1} \sum_{j=i+1}^{t-1} x_i x'_i \sum_{k=j+1}^t E[u_k u'_k] x_j x'_j \\
&= \frac{\sigma_u^2}{T} \sum_{i=t-Th}^{t-1} x_i x'_i \left(\sum_{j=t-Th}^i (t-j+j-i) x_j x'_j + \sum_{j=i+1}^{t-1} (t-j) x_j x'_j \right) \\
&= \frac{\sigma_u^2}{T} \left(\sum_{i=t-Th}^{t-1} x_i x'_i \sum_{j=t-Th}^{t-1} (t-j) x_j x'_j - \sum_{i=t-Th}^{t-1} x_i x'_i \sum_{j=t-Th}^i (i-j) x_j x'_j \right), \tag{B.5}
\end{aligned}$$

where we used the independence between $\{u_t\}$ and $\{x_t\}$, the serial independence of $\{u_t\}$, and $E[u_t u'_t] = \sigma_u^2 I_p$. Similarly, the second term becomes

$$\begin{aligned}
& E \left[\frac{1}{T} \left(\sum_{i=t+1}^{t+Th} x_i x'_i \sum_{k=t+1}^i u_k \right) \left(\sum_{i=t+1}^{t+Th} x_i x'_i \sum_{k=t+1}^i u_k \right)' \middle| X_T \right] \\
&= \frac{1}{T} \sum_{i=t+1}^{t+Th} \sum_{j=t+1}^{t+Th} x_i x'_i E \left[\sum_{k=t+1}^i u_k \sum_{l=t+1}^j u'_l \right] x_j x'_j \\
&= \frac{1}{T} \sum_{i=t+1}^{t+Th} \left(x_i x'_i \sum_{j=t+1}^i \sum_{k=t+1}^j E[u_k u'_k] x_j x'_j + x_i x'_i \sum_{j=i+1}^{t+Th} \sum_{k=t+1}^i E[u_k u'_k] x_j x'_j \right) \\
&= \frac{\sigma_u^2}{T} \sum_{i=t+1}^{t+Th} x_i x'_i \left(\sum_{j=t+1}^i (j-i+i-t) x_j x'_j + \sum_{j=i+1}^{t+Th} (i-t) x_j x'_j \right) \\
&= \frac{\sigma_u^2}{T} \left(\sum_{i=t+1}^{t+Th} (i-t) x_i x'_i \sum_{j=t+1}^{t+Th} x_j x'_j - \sum_{i=t+1}^{t+Th} x_i x'_i \sum_{j=t+1}^i (i-j) x_j x'_j \right). \tag{B.6}
\end{aligned}$$

The last term in the conditional expectation in (B.4) is

$$\begin{aligned}
E \left[\left(\sum_{i=t-Th}^{t+Th} x_i \varepsilon_i \right) \left(\sum_{i=t-Th}^{t+Th} x_i \varepsilon_i \right)' \middle| X_T \right] &= \sum_{i=t-Th}^{t+Th} \sum_{j=t-Th}^{t+Th} x_i x'_j E[\varepsilon_i \varepsilon_j] \\
&= \sigma_\varepsilon^2 \sum_{i=t-Th}^{t+Th} x_i x'_i. \tag{B.7}
\end{aligned}$$

Substituting (B.5), (B.6), and (B.7) into (B.4) yields

$$\begin{aligned}
& E[|\hat{\beta}_t - \beta_{T,t}|^2 | X_T] \\
&= \text{tr} \left[\left(\sum_{i=t-Th}^{t+Th} x_i x_i' \right)^{-1} \left\{ \frac{\sigma_u^2}{T} \left(\sum_{i=t-Th}^{t-1} x_i x_i' \sum_{j=t-Th}^{t-1} (t-j) x_j x_j' - \sum_{i=t-Th}^{t-1} x_i x_i' \sum_{j=t-Th}^i (i-j) x_j x_j' \right. \right. \right. \\
&\quad \left. \left. \left. + \sum_{i=t+1}^{t+Th} (i-t) x_i x_i' \sum_{j=t+1}^{t+Th} x_j x_j' - \sum_{i=t+1}^{t+Th} x_i x_i' \sum_{j=t+1}^i (i-j) x_j x_j' \right) + \sigma_\varepsilon^2 \sum_{i=t-Th}^{t+Th} x_i x_i' \right\} \left(\sum_{i=t-Th}^{t+Th} x_i x_i' \right)^{-1} \right] \\
&= \text{tr} \left[\left(\frac{1}{Th} \sum_{i=t-Th}^{t+Th} x_i x_i' \right)^{-1} \left\{ \sigma_u^2 h \left(\frac{1}{Th} \sum_{i=t-Th}^{t-1} x_i x_i' \frac{1}{Th} \sum_{j=t-Th}^{t-1} \frac{t-j}{Th} x_j x_j' \right. \right. \right. \\
&\quad - \frac{1}{Th} \sum_{i=t-Th}^{t-1} x_i x_i' \frac{1}{Th} \sum_{j=t-Th}^i \frac{i-j}{Th} x_j x_j' + \frac{1}{Th} \sum_{i=t+1}^{t+Th} \frac{i-t}{Th} x_i x_i' \frac{1}{Th} \sum_{j=t+1}^{t+Th} x_j x_j' \\
&\quad \left. \left. - \frac{1}{Th} \sum_{i=t+1}^{t+Th} x_i x_i' \frac{1}{Th} \sum_{j=t+1}^i \frac{i-j}{Th} x_j x_j' \right) + \frac{\sigma_\varepsilon^2}{Th} \frac{1}{Th} \sum_{i=t-Th}^{t+Th} x_i x_i' \right\} \left(\frac{1}{Th} \sum_{i=t-Th}^{t+Th} x_i x_i' \right)^{-1} \right] \\
&= \frac{\sigma_u^2 h}{4} \text{tr} \left[\Omega^{-1} (\Lambda \bar{\Lambda} + \bar{\Lambda} \Lambda - 2\Xi) \Omega^{-1} \right] (1 + o(1)) + \frac{\sigma_\varepsilon^2}{2Th} \text{tr} \left[\Omega^{-1} \right] (1 + o(1))
\end{aligned}$$

in the sense of L_1 convergence. Note that we used the stationarity of x_t to derive the final expression. Therefore, the MSE of $\hat{\beta}_t$ satisfies

$$\begin{aligned}
\text{MSE}(h) &= E[E[|\hat{\beta}_t - \beta_{T,t}|^2 | X_T]] \\
&= \frac{\sigma_u^2 h}{4} \text{tr} \left[\Omega^{-1} (\Lambda \bar{\Lambda} + \bar{\Lambda} \Lambda - 2\Xi) \Omega^{-1} \right] (1 + o(1)) + \frac{\sigma_\varepsilon^2}{2Th} \text{tr} \left[\Omega^{-1} \right] (1 + o(1)),
\end{aligned}$$

where we interchanged the order of expectation and L_1 limit operator. \square

Appendix C: Performance of Structural Break Tests

In this appendix, we investigate the behavior of conventional structural break tests. Our focus is on whether the tests for structural breaks can discover latent breaks correctly even in the presence of another source of instability such as smooth functions and the random walk. We verify this via (limited) Monte Carlo experiments. The data is generated as $y_t = \beta_{T,t} x_t + \varepsilon_t$, $t = 1, \dots, T$, where $\varepsilon_t \sim \text{i.i.d. } N(0, 1)$, and $x_t = 0.5x_{t-1} + \varepsilon_{x,t}$ with $\varepsilon_{x,t} \sim \text{i.i.d. } N(0, 1)$. $\beta_{T,t}$ is defined as a smooth function or rescaled random walk with two abrupt breaks. Specifically, we let $\beta_{T,t} = \mu_{T,t} + h_{T,t}$, where $\mu_{T,t} = \sum_{i=1}^3 T^{-\alpha} \mu_i 1\{\lfloor \tau_{i-1} T \rfloor + 1 \leq$

$t \leq \lfloor \tau_i T \rfloor$ with $\tau_0 = 0$, $\tau_1 = 0.3$, $\tau_2 = 0.7$, and $\tau_3 = 1$. $h_{T,t}$ is specified as either a deterministic smooth function $f(t/T)$ or rescaled random walk $g_{T,t}$. The function f is equal to $f(u) = 2u + \exp(-16(u - 0.5)^2)$ or $f(u) = \{\sin(\pi u) + \cos(2\pi u) + \sin(3\pi u) + \cos(4\pi u)\}/4$. $g_{T,t}$ is generated as $g_{T,t} = T^{-1/2} \sum_{i=1}^t v_i$, where $v_i \sim \text{i.i.d. } N(0, 1)$ or $v_i \sim \text{i.i.d. log normal}$ with parameters $\mu = 0$ and $\sigma = 1$.

When $h_{T,t} = f(t/T)$, $\beta_{T,t}$ evolves smoothly and deterministically over time but experiences two abrupt breaks at the 30% and 70% points of the sample period. The magnitude of the breaks is determined by μ_i and α . We let $\mu_1 = 0$, $\mu_2 = 4$, $\mu_3 = -2$, and $\alpha \in \{0.1, 0.2\}$. When $h_{T,t} = g_{T,t}$, $\beta_{T,t}$ follows a rescaled random walk with two discontinuous jumps.

To identify abrupt breaks, we use the comprehensive estimation procedure developed by Nguyen, Perron and Yamamoto (2023). In this procedure, the number of breaks and break dates are estimated by the sequential method (SEQ) proposed by Bai and Perron (1998), the BIC suggested by Yao (1988), the modified SIC (LWZ) of Liu, Wu and Zidek (1997) or the modified BIC (KT) of Kurozumi and Tuvaandorj (2011) (see Nguyen et al. (2023) for the detailed description of the procedure and the associated R package). We investigate the performance of these four methods through 2000 replications with the sample size being (i) $T = 100$, (ii) $T = 200$, (iii) $T = 400$ and (iv) $T = 800$.

We calculate the frequency of particular numbers of breaks (up to 5) being selected and the estimated break date fraction (\hat{T}_B/T) being in the $1/25$ -neighborhood of the true one.¹⁵

Let us start with the case of $h_{T,t} = f(t/T)$ with $f(u) = 2u + \exp(-16(u - 0.5)^2)$ (Table C.1). When $\alpha = 0.1$ and $T = 100$ (case (i)), the SEQ method estimates no break with a probability of 13%, while it overestimates the number of breaks in about 25% of the 2000 replications. The estimate of the break date fraction falls in the $1/25$ -neighborhood of the true brake date fraction with a probability of 80%-85%. As T gets larger, the frequency of underestimating the number of breaks decreases, and the true break points are detected more frequently, but the number of breaks gets more likely to be overestimated. In particular, the estimated number of breaks is more than two in more than 90% of the 2000 replications when $T = 800$. The same tendency to overestimate the number of breaks is shared by the BIC and KT methods, although they can identify the true break points with a high probability even when $T = 100$. This implies that BIC and KT often detect spurious breaks in addition to the true ones. LWZ is the most successful in this case, identifying the true breaks in almost all

¹⁵We check the behavior of the estimate for the break date fraction, T_B/T , rather than break date T_B itself. This is because T_B/T can be consistently estimated but T_B cannot; see Casini and Perron (2018).

replications for $T \geq 200$ without detecting an additional spurious break. However, LWZ is more likely to overestimate the number of breaks as T gets larger, as the other methods are. When $\alpha = 0.2$, the tendency to overestimate the number of breaks is greater for all the four tests than in the case with $\alpha = 0.1$, and the probability of the true breaks being identified decreases. The LWZ method still performs well, estimating the number of breaks to be two with a probability of not less than 94%.

Next, we consider the case of $h_{T,t} = f(t/T)$ with $f(u) = \{\sin(\pi u) + \cos(2\pi u) + \sin(3\pi u) + \cos(4\pi u)\}/4$ (Table C.2). When $\alpha = 0.1$, the behaviors of the four methods are similar to those in the case of $f(u) = 2u + \exp(-16(u - 0.5)^2)$ with $\alpha = 0.1$, but the SEQ procedure estimates the number of breaks to be not less than two in almost all replications and detects the true breaks with a high probability even when $T = 100$. In this case, the LWZ procedure is the most successful one, identifying the true breaks without detecting a spurious in all replications for $T = 400, 800$. When $\alpha = 0.2$, there are several differences. First, LWZ is much more likely to underestimate the number of breaks than the other tests. For example, it estimates the number of breaks to be less than two with probabilities of 38% and 26% for $T = 100$ and $T = 200$, respectively. The probability of the underestimation is still nonnegligible even when $T = 400, 800$, under which sample size the other tests estimate the number of breaks to be not less than two in almost all replications. This causes the true breaks (in particular, the first one) to be overlooked by LWZ. For the other tests (SEQ, BIC, KT), the tendency to overestimate the number of breaks gets stronger as T increases. These tests are the most successful procedures in terms of identifying the true breaks.

We turn to the case with $h_{T,t} = g_{T,t} = T^{-1/2} \sum_{i=1}^t v_i$ where $v_i \sim \text{i.i.d. } N(0, 1)$ (Table C.3). When $\alpha = 0.1$, the behaviors of the four procedures are similar to those in the preceding cases: SEQ, BIC and KT not only identify true breaks but also detect spurious ones, with this tendency being greater for larger T , and LWZ identifies true breaks without detecting spurious one with a large probability. However, the probability of overestimating the number of breaks is nonnegligible for LWZ, and this probability gets larger as T increases. When $\alpha = 0.2$, SEQ and LWZ are more likely to underestimate the number of breaks than the other two methods. In particular, LWZ underestimates the number of breaks with a nonnegligible probability even when $T = 800$ and thus is more likely to overlook the latent breaks than the other tests. SEQ, BIC and KT can identify latent breaks with a high probability but tend to detect spurious breaks. This tendency is stronger for larger T , as in the preceding cases. The results for the case with $v_i \sim \text{i.i.d. log normal}$ are similar, so the same comment applies.

Table C.1: Results of structural break tests for $h_{T,t} = f(t/T)$ with $f(u) = 2u + \exp(-16(u - 0.5)^2)$

		# of estimated breaks						Frequency of $\hat{T}_B/T \in$	
		0	1	2	3	4	5	$[0.3 \pm 1/25]$	$[0.7 \pm 1/25]$
$\alpha = 0.1$									
(i)	SEQ	0.126	0	0.613	0.239	0.022	0	0.831	0.868
	BIC	0	0	0.870	0.126	0.005	0	0.966	0.995
	LWZ	0	0	0.996	0.004	0	0	0.979	0.995
	KT	0	0	0.802	0.184	0.015	0	0.964	0.994
(ii)	SEQ	0.004	0	0.618	0.351	0.027	0	0.977	0.996
	BIC	0	0	0.789	0.198	0.013	0	0.996	0.999
	LWZ	0	0	0.995	0.006	0	0	0.998	0.999
	KT	0	0	0.756	0.225	0.020	0	0.996	0.999
(iii)	SEQ	0	0	0.358	0.554	0.089	0	0.994	1
	BIC	0	0	0.518	0.411	0.072	0	1	1
	LWZ	0	0	0.995	0.006	0	0	1	1
	KT	0	0	0.515	0.430	0.056	0	1	1
(iv)	SEQ	0	0	0.069	0.620	0.311	0.001	0.999	1
	BIC	0	0	0.105	0.453	0.443	0	1	1
	LWZ	0	0	0.966	0.034	0	0	1	1
	KT	0	0	0.127	0.502	0.371	0	1	1
$\alpha = 0.2$									
(i)	SEQ	0.044	0	0.657	0.280	0.020	0	0.841	0.935
	BIC	0	0	0.847	0.149	0.005	0	0.883	0.981
	LWZ	0	0	0.993	0.007	0	0	0.920	0.981
	KT	0	0	0.775	0.211	0.015	0	0.876	0.980
(ii)	SEQ	0	0	0.588	0.392	0.021	0.001	0.923	0.995
	BIC	0	0	0.750	0.241	0.010	0	0.938	0.995
	LWZ	0	0	0.992	0.008	0	0	0.983	0.995
	KT	0	0	0.722	0.264	0.015	0	0.936	0.995
(iii)	SEQ	0	0	0.343	0.590	0.068	0	0.968	1
	BIC	0	0	0.463	0.481	0.056	0	0.949	1
	LWZ	0	0	0.987	0.014	0	0	0.996	1
	KT	0	0	0.473	0.481	0.047	0	0.950	1
(iv)	SEQ	0	0	0.067	0.682	0.251	0	0.992	1
	BIC	0	0	0.091	0.556	0.354	0	0.980	1
	LWZ	0	0	0.938	0.063	0	0	0.992	1
	KT	0	0	0.119	0.583	0.298	0	0.977	1

Note: $\beta_{T,t} = \mu_{T,t} + f(t/T)$, where $\mu_{T,t} = \sum_{i=1}^3 \mu_i T^{-\alpha} 1\{\lfloor \tau_{i-1} T \rfloor + 1 \leq t \leq \lfloor \tau_i T \rfloor\}$ and $f(u) = 2u + \exp(-16(u - 0.5)^2)$. The number of replications is 2000. The sample size is $T = 100$ for case (i), $T = 200$ for case (ii), $T = 400$ for case (iii), and $T = 800$ for case (iv).

Table C.2: Results of structural break tests for $h_{T,t} = f(t/T)$ with $f(u) = \{\sin(\pi u) + \cos(2\pi u) + \sin(3\pi u) + \cos(4\pi u)\}/4$

		# of estimated breaks						Frequency of $\hat{T}_B/T \in$	
		0	1	2	3	4	5	$[0.3 \pm 1/25]$	$[0.7 \pm 1/25]$
$\alpha = 0.1$									
(i)	SEQ	0.001	0	0.829	0.162	0.009	0	0.962	0.995
	BIC	0	0	0.972	0.028	0.001	0	0.962	0.995
	LWZ	0	0.002	0.999	0	0	0	0.961	0.995
	KT	0	0	0.924	0.073	0.004	0	0.962	0.995
(ii)	SEQ	0	0	0.845	0.151	0.005	0	0.997	0.999
	BIC	0	0	0.967	0.033	0	0	0.997	1
	LWZ	0	0	1	0	0	0	0.997	1
	KT	0	0	0.947	0.054	0	0	0.997	1
(iii)	SEQ	0	0	0.773	0.219	0.009	0	1	1
	BIC	0	0	0.929	0.070	0.001	0	1	1
	LWZ	0	0	1	0	0	0	1	1
	KT	0	0	0.921	0.077	0.003	0	1	1
(iv)	SEQ	0	0	0.522	0.425	0.053	0.001	1	1
	BIC	0	0	0.789	0.201	0.011	0	1	1
	LWZ	0	0	1	0	0	0	1	1
	KT	0	0	0.800	0.190	0.011	0	1	1
$\alpha = 0.2$									
(i)	SEQ	0.025	0.047	0.782	0.139	0.009	0	0.765	0.950
	BIC	0	0.044	0.924	0.032	0.001	0	0.798	0.970
	LWZ	0.078	0.303	0.620	0	0	0	0.529	0.898
	KT	0.001	0.047	0.877	0.073	0.004	0	0.793	0.971
(ii)	SEQ	0	0.006	0.845	0.145	0.005	0	0.912	0.994
	BIC	0	0.007	0.957	0.036	0	0	0.911	0.985
	LWZ	0.009	0.248	0.744	0	0	0	0.694	0.980
	KT	0	0.011	0.928	0.061	0.001	0	0.905	0.985
(iii)	SEQ	0	0.001	0.768	0.224	0.008	0	0.966	1
	BIC	0	0.001	0.923	0.076	0.001	0	0.968	0.997
	LWZ	0	0.134	0.866	0	0	0	0.842	0.997
	KT	0	0.001	0.917	0.079	0.004	0	0.967	0.997
(iv)	SEQ	0	0	0.528	0.413	0.059	0.001	0.979	1
	BIC	0	0	0.767	0.220	0.013	0	0.979	1
	LWZ	0	0.074	0.926	0	0	0	0.907	1
	KT	0	0.001	0.782	0.206	0.012	0	0.979	1

Note: $\beta_{T,t} = \mu_{T,t} + f(t/T)$, where $\mu_{T,t} = \sum_{i=1}^3 \mu_i T^{-\alpha} \mathbf{1}\{[\tau_{i-1}T] + 1 \leq t \leq [\tau_i T]\}$ and $f(u) = \{\sin(\pi u) + \cos(2\pi u) + \sin(3\pi u) + \cos(4\pi u)\}/4$. The number of replications is 2000. The sample size is $T = 100$ for case (i), $T = 200$ for case (ii), $T = 400$ for case (iii), and $T = 800$ for case (iv).

Table C.3: Results of structural break tests for $h_{T,t} = g_{T,t} = T^{-1/2} \sum_{i=1}^t v_i$ with $v_i \sim$ i.i.d. $N(0,1)$

		# of estimated breaks						Frequency of $\hat{T}_B/T \in$	
		0	1	2	3	4	5	$[0.3 \pm 1/25]$	$[0.7 \pm 1/25]$
$\alpha = 0.1$									
(i)	SEQ	0.025	0.002	0.737	0.223	0.014	0	0.932	0.967
	BIC	0	0.001	0.897	0.100	0.003	0	0.957	0.994
	LWZ	0	0.007	0.984	0.010	0	0	0.953	0.994
	KT	0	0.001	0.838	0.152	0.010	0	0.957	0.994
(ii)	SEQ	0.001	0.001	0.718	0.262	0.019	0.001	0.990	0.998
	BIC	0	0	0.812	0.181	0.008	0	0.996	1
	LWZ	0	0.003	0.979	0.019	0	0	0.991	1
	KT	0	0	0.779	0.214	0.008	0	0.996	1
(iii)	SEQ	0	0	0.581	0.369	0.050	0.001	0.999	1
	BIC	0	0	0.682	0.289	0.029	0.001	0.999	1
	LWZ	0	0.001	0.959	0.040	0.001	0	0.998	1
	KT	0	0	0.677	0.294	0.029	0.002	0.999	1
(iv)	SEQ	0	0.001	0.378	0.501	0.116	0.006	0.999	1
	BIC	0	0	0.470	0.423	0.103	0.005	1	1
	LWZ	0	0.001	0.881	0.116	0.003	0	0.999	1
	KT	0	0	0.487	0.416	0.093	0.005	1	1
$\alpha = 0.2$									
(i)	SEQ	0.056	0.065	0.685	0.183	0.011	0	0.730	0.896
	BIC	0.001	0.059	0.849	0.089	0.002	0	0.792	0.969
	LWZ	0.025	0.187	0.784	0.004	0	0	0.680	0.938
	KT	0.001	0.064	0.790	0.136	0.010	0	0.789	0.967
(ii)	SEQ	0.006	0.050	0.718	0.214	0.012	0.001	0.845	0.969
	BIC	0	0.046	0.792	0.157	0.006	0	0.867	0.985
	LWZ	0.010	0.170	0.812	0.009	0	0	0.762	0.968
	KT	0	0.044	0.760	0.188	0.009	0	0.865	0.984
(iii)	SEQ	0	0.047	0.605	0.317	0.031	0	0.863	0.982
	BIC	0	0.041	0.671	0.265	0.024	0.001	0.895	0.996
	LWZ	0.003	0.169	0.804	0.025	0	0	0.777	0.982
	KT	0	0.045	0.671	0.259	0.025	0.001	0.888	0.995
(iv)	SEQ	0.001	0.027	0.431	0.464	0.076	0.003	0.870	0.979
	BIC	0	0.025	0.482	0.402	0.088	0.004	0.906	0.998
	LWZ	0.005	0.163	0.761	0.069	0.003	0	0.775	0.980
	KT	0	0.031	0.495	0.388	0.084	0.003	0.902	0.997

Note: $\beta_{T,t} = \mu_{T,t} + g_{T,t}$, where $\mu_{T,t} = \sum_{i=1}^3 \mu_i T^{-\alpha} 1\{[\tau_{i-1}T] + 1 \leq t \leq [\tau_i T]\}$ and $g_{T,t} = T^{-1/2} \sum_{i=1}^t v_i$ with $v_i \sim$ i.i.d. $N(0,1)$. The sample size is $T = 100$ for case (i), $T = 200$ for case (ii), $T = 400$ for case (iii), and $T = 800$ for case (iv).

Table C.4: Results of structural break tests for $h_{T,t} = g_{T,t} = T^{-1/2} \sum_{i=1}^t v_i$ with $v_i \sim$ i.i.d. log normal

		# of estimated breaks					Frequency of $\hat{T}_B/T \in$		
		0	1	2	3	4	5	$[0.3 \pm 1/25]$	$[0.7 \pm 1/25]$
$\alpha = 0.1$									
(i)	SEQ	0.026	0.003	0.739	0.216	0.017	0.001	0.930	0.963
	BIC	0	0.002	0.899	0.095	0.005	0	0.958	0.994
	LWZ	0	0.008	0.986	0.007	0	0	0.954	0.995
	KT	0	0.002	0.864	0.126	0.009	0	0.956	0.994
(ii)	SEQ	0	0.001	0.728	0.259	0.014	0	0.992	0.997
	BIC	0	0	0.830	0.162	0.009	0	0.992	0.998
	LWZ	0	0.004	0.978	0.019	0	0	0.991	0.998
	KT	0	0	0.813	0.177	0.010	0	0.992	0.999
(iii)	SEQ	0	0	0.592	0.358	0.050	0.001	0.999	1
	BIC	0	0	0.703	0.266	0.031	0.001	1	1
	LWZ	0	0.002	0.954	0.044	0.001	0	0.998	1
	KT	0	0	0.705	0.261	0.033	0.001	1	1
(iv)	SEQ	0	0	0.378	0.495	0.124	0.003	0.999	1
	BIC	0	0	0.475	0.420	0.100	0.006	1	1
	LWZ	0	0.001	0.878	0.118	0.004	0	0.999	1
	KT	0	0	0.492	0.411	0.092	0.006	1	1
$\alpha = 0.2$									
(i)	SEQ	0.057	0.078	0.680	0.175	0.011	0	0.719	0.898
	BIC	0	0.067	0.843	0.088	0.003	0	0.780	0.968
	LWZ	0.026	0.198	0.772	0.004	0	0	0.665	0.935
	KT	0	0.070	0.800	0.122	0.009	0	0.775	0.966
(ii)	SEQ	0.004	0.059	0.710	0.219	0.010	0	0.846	0.966
	BIC	0	0.053	0.799	0.142	0.007	0	0.856	0.985
	LWZ	0.006	0.189	0.795	0.011	0	0	0.742	0.970
	KT	0	0.057	0.784	0.152	0.008	0	0.856	0.985
(iii)	SEQ	0.001	0.043	0.617	0.306	0.033	0.001	0.870	0.978
	BIC	0	0.042	0.688	0.247	0.023	0.001	0.892	0.994
	LWZ	0.004	0.170	0.803	0.023	0.001	0	0.776	0.978
	KT	0	0.045	0.688	0.240	0.027	0.001	0.889	0.993
(iv)	SEQ	0	0.034	0.428	0.448	0.089	0.002	0.872	0.986
	BIC	0	0.032	0.476	0.404	0.086	0.004	0.908	0.998
	LWZ	0.003	0.144	0.763	0.090	0.002	0	0.791	0.981
	KT	0	0.037	0.487	0.395	0.078	0.004	0.903	0.997

Note: $\beta_{T,t} = \mu_{T,t} + g_{T,t}$, where $\mu_{T,t} = \sum_{i=1}^3 \mu_i T^{-\alpha} 1\{\lfloor \tau_{i-1} T \rfloor + 1 \leq t \leq \lfloor \tau_i T \rfloor\}$ and $g_{T,t} = T^{-1/2} \sum_{i=1}^t v_i$ with $v_i \sim$ i.i.d. log normal with parameters $\mu = 0$ and $\sigma = 1$. The sample size is $T = 100$ for case (i), $T = 200$ for case (ii) $T = 400$ for case (iii), and $T = 800$ for case (iv).

In general, the tests for structural breaks can identify latent breaks in the presence of another source of parameter instability but tend to detect additional spurious breaks. This tendency is stronger for larger T . Investigating the behavior of each test, LWZ identifies latent breaks without estimating spurious breaks in some situations, but it underestimates the number of breaks and overlooks latent breaks in other situations. SEQ is likely to both underestimate and overestimate the number of breaks. BIC and KT can identify true breaks irrespective of the DGP, but they tend to detect additional spurious breaks.

Appendix D: Additional Results for the Empirical Application

In this appendix, we discuss the estimation results for portfolios G and V (see Section 5 for details). Figures D.1 and D.2 show the estimates for G, and Figures D.3 and D.4 for V.


D.1 Estimates for G

First, we compare the estimates obtained from the kernel method ($h = T^{-1/2}$) with those obtained from the Bayesian method explained in Section 5.3.1. See Figure D.1. For the time-varying alpha, the kernel estimates are more volatile than the Bayesian estimates, but both estimates share the same frequency. For the time-varying beta, the estimates obtained from the two procedures are similar, irrespective of the value of the hyperparameter used in the Bayesian method.


In Figure D.2a, we plot the estimated time-varying alpha and its confidence band obtained from the kernel method with $h = T^{-1/2}$. The estimated alpha stays around zero as a whole, but there are troughs around $t = 400$ and $t = 550$. Around these troughs, the confidence band does not include the value zero. Figure D.2b shows the estimates for the time-varying beta. It starts with the value of 0.8 and starts to increase soon later. From $t = 100$ till the end of the sample, it stays between 1.2 and 1.5. The confidence band does not include the value zero on the whole line.

D.2 Estimates for V

In Figure D.3, we compare the estimates obtained from the kernel ($h = T^{-1/2}$) and Bayesian approaches. For the time-varying alpha, the estimates are similar and share the same frequency. In particular, the kernel and Bayesian (with $v_1 = 4$) estimates have similar




(a) Plot of the time-varying alpha




(b) Plot of the time-varying beta

Figure D.1: Estimates from the kernel ($h = T^{-1/2}$) and Bayesian ($D \sim \text{Gamma}(2, 10^{-4})$) methods for G

—: Kernel, ·····: Bayesian ($v_1 = 1$), - - - : Bayesian ($v_1 = 2$), - · - : Bayesian ($v_1 = 4$)



(a) Plot of the time-varying alpha




(b) Plot of the time-varying beta

Figure D.2: Estimates and 95% confidence band from the kernel-based method ($h = T^{-1/2}$) for G


(The horizontal line in (a) indicates the value zero.)

peaks and troughs. For the time-varying beta, both estimates have a similar general pattern; the estimates fluctuate between 0.75 and 1.5 up to $t = 300$, then drop to and fluctuate at a lower level up to $t = 600$, and then go upward and stay at a higher level until the end of the sample. The two estimates also share a similar frequency, but the Bayesian estimate is more volatile (irrespective of the value of v_1) and has more peaks and troughs than the kernel-based estimate.

Figure D.4a shows the estimated time-varying alpha and its confidence band obtained from the kernel method with $h = T^{-1/2}$. Although the confidence band for the estimated alpha includes zero as a whole, there are several periods when the value zero is excluded from the band. The estimated alpha is significantly different from zero and positive during those periods. Figure D.4b depicts the estimated time-varying beta. From $t = 1$ to $t = 320$, it stays around 1.2 and then sharply drops to 0.8. From $t = 370$, it gently increases but sharply drops from $t = 510$ to reach the lowest level of 0.6 at $t = 580$. Then, the time-varying beta dramatically increases and re-enter the phase where it fluctuates around 1.2 from $t = 620$.




(a) Plot of the time-varying alpha



(b) Plot of the time-varying beta

Figure D.3: Estimates from the kernel ($h = T^{-1/2}$) and Bayesian ($D \sim \text{Gamma}(2, 10^{-4})$) methods for V

—: Kernel, ·····: Bayesian ($v_1 = 1$), - - - : Bayesian ($v_1 = 2$), - · - : Bayesian ($v_1 = 4$)



(a) Plot of the time-varying alpha



(b) Plot of the time-varying beta

Figure D.4: Estimates and 95% confidence band from the kernel-based method ($h = T^{-1/2}$) for V

(The horizontal line in (a) indicates the value zero.)

References to Appendices

- Andrews, D. W. K. (1988) Laws of Large Numbers for Dependent Non-Identically Distributed Random Variables. *Econometric Theory*, 4 (3), 458–467.
- Bai, J. and P. Perron (1998) Estimating and Testing Linear Models with Multiple Structural Changes. *Econometrica*, 66 (1), 47–78.
- Davidson, J. (1994) *Stochastic Limit Theory: An Introduction for Econometricians* Oxford University Press.
- de Jong, R. M. (1997) Central Limit Theorems for Dependent Heterogeneous Random Variables. *Econometric Theory*, 13 (3), 353–367.
- Kurozumi, E. and P. Tuvaandorj (2011) Model Selection Criteria in Multivariate Models with Multiple Structural Changes. *Journal of Econometrics*, 164 (2), 218–238.
- Liu, J., S. Wu, and J. V. Zidek (1997) On Segmented Multivariate Regression. *Statistica Sinica*, 7 (2), 497–525.
- Nguyen, L., P. Perron, and Y. Yamamoto (2023) Mbreaks: Estimation and Inference for Structural Breaks in Linear Regression Models.
- Yao, Y.-C. (1988) Estimating the Number of Change-Points via Schwarz' Criterion. *Statistics & Probability Letters*, 6 (3), 181–189.