

On one-step numerical schemes of weak convergence for SDEs with super-linear coefficients [†]

Yuying Zhao^{a,b}, Xiaojie Wang^a, and Zhongqiang Zhang ^{*b}

^a *School of Mathematics and Statistics, HNP-LAMA, Central South University, Changsha, Hunan, P. R. China*

^b *Department of Mathematical Sciences, Worcester Polytechnic Institute, Worcester, MA 01609 USA*

June 21, 2024

Abstract

We consider weak convergence of one-step schemes for solving stochastic differential equations (SDEs) with one-sided Lipschitz conditions. It is known that the super-linear coefficients may lead to a blowup of moments of solutions and their numerical solutions. When solutions to SDEs have all finite moments, weak convergence of numerical schemes has been investigated in [Wang et al (2023), Weak error analysis for strong approximation schemes of SDEs with super-linear coefficients, IMA Journal numerical analysis]. Some modified Euler schemes have been analyzed for weak convergence. In this work, we present a family of explicit schemes of first and second-order weak convergence based on classical schemes for SDEs. We explore the effects of limited moments on these schemes. We provide a systematic but simple way to establish weak convergence orders for schemes based on approximations/modifications of drift and diffusion coefficients. We present several numerical examples of these schemes and show their weak convergence orders.

AMS subject classification: 60H35, 60H15, 65C30.

Keywords: Non-globally Lipschitz coefficients, second order, weak approximation, modified scheme, sigmoid functions, truncation functions

1 Introduction

We consider the weak convergence of first- and second-order explicit schemes for stochastic differential equations (SDEs) with non-globally Lipschitz coefficients of superlinear growth. In [31], a theorem is established on the weak convergence orders of one-step numerical schemes with *infinity* moments. The weak convergence orders of the several tamed and balanced schemes are examined therein. However, there are several limitations to be addressed for further applications. The limitations include but are not limited to 1) requiring infinitely many moments of solutions to SDEs and numerical solutions, 2) schemes of at most first-order weak convergence, and 3) not structure-preserving from the schemes.

We will address the first two limitations by discussing 1) relaxing requirements of all finite moments, which necessitate the moments of derivatives of solutions with respect to the initial condition; 2) second-order schemes obtained from modifying classical second-order schemes for SDEs with Lipschitz coefficients. We will not focus on 3) while we discuss some possibilities in Remark 3.11. For 1), we will reduce the requirements of all finite moments of both exact solutions and numerical ones from the infinite order to a finite order for a q -th order numerical scheme

[†]This work was supported by the Natural Science Foundation of China (12071488, 12371417, 11971488), the Natural Science Foundation of Hunan Province (2020JJ2040), and the innovative project of graduate students of Central South University (2021zzts0040). E-mail addresses: zhaoyuying78@gmail.com, x.j.wang7@csu.edu.cn, zzhang7@wpi.edu

*Corresponding author

($0 < q \leq 2$). The reduction allows the design of numerical schemes of weak convergence when solutions have only a few moments. Also, we show the required moments of finite orders are critical for the desired convergence order. For 2), we modify schemes of weak order two for SDEs with Lipschitz coefficients and show that the weak convergence order of the modified schemes may degenerate to one for nonlinear SDEs with only limited moments. It can achieve second-order when numerical solutions have enough high-order moments. See Example 3.16 for a representative example.

We briefly discuss several related works on analyzing weak convergence using one-step numerical approximation. For a thorough review of schemes of weak convergence for SDEs with non-Lipschitz coefficients, we refer to [31]. The idea of analyzing the weak convergence by one-step numerical approximation dates back to [21, 22]. These typically establish the weak convergence theorem for general one-step numerical approximations of SDEs with global Lipschitz coefficients. See also discussions in [14, 23]. However, in practical applications, the coefficients of most SDEs are not Lipschitz continuous and exhibit super-linear or sublinear growth. In [31], the authors develop a weak error analysis for general one-step approximation schemes of SDEs with non-globally Lipschitz coefficients of superlinear growth. Under the same conditions on the coefficients of SDEs, the Euler-Maruyama scheme fails to have bounded moments and thus does not converge in the strong and weak sense; see e.g., in [7, 10, 20, 24]. For numerical schemes solving these SDEs, several types of methods have been proposed: 1) *Tamed explicit schemes*, such as in [9, 11, 12, 25–27, 29–31], where coefficients are approximated with a function of the form $\frac{f(x)}{1+h^\beta|f(x)|}$ ($0 \leq \beta \leq 1$), effectively controlling the superlinear growth; 2) *Explicit truncation schemes*, where coefficients are set to be constants when solutions reach a certain threshold of magnitude, as exemplified in prior studies [1, 2, 15, 17, 24, 27]; 3) *Projection methods*: trajectories with tremendous values are projected back onto a ball of radius inversely proportional to the step size. For further information, refer to the works of [1, 2, 27]; 4) *Implicit methods*: the drift and/or diffusion coefficients are treated implicitly, see e.g. [6, 8, 18, 31]. Also, the authors in [24] apply the schemes of weak convergence for SDEs with Lipschitz continuous coefficients and reject all the computed trajectories outside a ball centered at the origin with a certain radius. Therein the resulting scheme may be volume-preserving for stochastic Hamiltonian systems while there is no mathematical proof.

Except the implicit schemes, all the above schemes are established by modifying the working schemes for SDEs with Lipschitz coefficients. Along this avenue of modifying classical schemes, we extend the theorem of weak convergence for one-step numerical schemes in [31], using relaxed assumptions. Also, we discuss numerical schemes with second-order weak convergence, which haven't been explored for SDEs with non-Lipschitz coefficients. For example, only first-order schemes are considered in [31]. The main novelty and contributions of the work are summarized as follows.

- First, we relax the assumptions in the key theorem in [31]. Therein all moments of solutions to SDEs and numerical solutions are assumed. However, this assumption is invalid for many SDEs and their numerical schemes; see Example 3.16. The relaxation requires investigating the solutions and their derivatives in the initial data. After thoroughly investigating the moments of the derivatives, we present a simpler proof (see Appendix A and B) with relaxed assumptions than those in [4, 31].
- Second, we present several modified schemes (Examples 3.10-3.14) from classical second-order schemes and discuss the weak convergence orders of these schemes for nonlinear SDEs. We show in Examples 3.16 (theoretically) and 4.2 (numerically) that the convergence order of the schemes of second-order weak convergence become one when the moment conditions are not satisfied.

In Figure 1, we present the key components for verifying the weak convergence. The bold texts are requirements in this work. Here we consider modified Euler or Milstein-Talay schemes and thus we only need to check the properties of the modification maps to obtain the moment bounds and one-step approximation error to obtain a weak convergence order. In other words, users may compute the weak convergence order by checking these assumptions and conditions in Theorems 3.3 and 3.8 but they are not required to check the moment bounds and local approximation error. To illustrate how to apply our results, we present several examples in Section 3.4.




Figure 1: Sketch of components for proving weak convergence order of one-step schemes for SDE (2.2)

The rest of this paper is organized as follows. In Section 2, we first introduce some notations and assumptions. Then we present the general convergence theorem of one-step schemes for SDEs with assumptions weaker than those in [31]. The proof of this improvement is presented in Appendix C. We then present second-order weak convergence candidate schemes in Section 3. We also present necessary technical assumptions to obtain second-order convergence, especially on how to modify the classical second-order weakly convergent schemes. We present some numerical results in Section 4. The essential proofs of the main convergence results are presented in Section 5 while more details can be found in the appendix.

2 Preliminaries and assumptions

We introduce some notations used throughout this paper.

We use $|\cdot|$ to denote the l^2 -norm of vectors and matrices. We denote by $\mathbf{C}^k(\mathbb{R}^d, \mathbb{R}^m)$ the space consisting of k -th continuously differentiable functions from \mathbb{R}^d to \mathbb{R}^m , and denote $x \vee y := \max\{x, y\}$ and $x \wedge y := \min\{x, y\}$. For a multi-index α , we define the partial derivatives of $v: \mathbb{R}^d \rightarrow \mathbb{R}^l$ as

$$D^\alpha v(x) := \frac{\partial^{|\alpha|_1} v(x)}{\partial x_1^{\alpha_1} \cdots \partial x_d^{\alpha_d}} = \partial_{x_1}^{\alpha_1} \cdots \partial_{x_d}^{\alpha_d} v(x), \quad |\alpha|_1 = \sum_{i=1}^d \alpha_i. \quad (2.1)$$

For $l = 1$, We use $Dv := (\frac{\partial v}{\partial x_1}, \frac{\partial v}{\partial x_2}, \dots, \frac{\partial v}{\partial x_d})$ (gradient vector) and $D^2v := (\frac{\partial^2 v}{\partial x_i \partial x_j})_{d \times d}$ (Hessian matrix). For simplicity, we use the letter C to denote a generic positive constant independent of the step size h .

For a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$, we use \mathbb{E} to denote expectation and $L^r(\Omega; \mathbb{R}^{d \times m})$, $r \in \mathbb{N}$, to denote the family of $\mathbb{R}^{d \times m}$ -valued variables with the norm defined by $\|\xi\|_{L^r(\Omega; \mathbb{R}^{d \times m})} = (\mathbb{E}[|\xi|^r])^{\frac{1}{r}} < \infty$. Let \mathcal{F}_t^W be an increasing family of σ -subalgebras of \mathcal{F} induced by $W(t)$ for $0 \leq t \leq T$, where $(W(t), \mathcal{F}_t^W) = ((W_1(t), \dots, W_m(t))^\top, \mathcal{F}_t^W)$ is an m -dimensional standard Wiener process. Consider the following Itô stochastic differential equation:

$$dX(t) = f(X(t)) dt + g(X(t)) dW(t), \quad t \in (0, T], \quad X(0) = x_0 \in \mathbb{R}^d, \quad (2.2)$$

where $f = (f^1, f^2, \dots, f^d)^\top: \mathbb{R}^d \rightarrow \mathbb{R}^d$ is the drift coefficient and $f^i: \mathbb{R}^d \rightarrow \mathbb{R}$, and $g = (g^{i,j})_{d \times m}: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times m}$ is the diffusion coefficient and $g^{i,j}: \mathbb{R}^d \rightarrow \mathbb{R}$. The initial data $x_0 = (x_0^{(1)}, x_0^{(2)}, \dots, x_0^{(d)}) \in \mathbb{R}^d$ is deterministic for simplicity.

2.1 Conditions on coefficients of SDEs

In this subsection, we present assumptions on the drift and diffusion coefficients. These assumptions are crucial for establishing the existence, uniqueness, and moment estimates of SDEs.

Assumption 2.1. (A1) The drift coefficient function $f \in \mathbf{C}^{2q+2}(\mathbb{R}^d; \mathbb{R}^d)$ and for $r \geq 0$,

$$\sup_{x \in \mathbb{R}^d} \frac{|D^\alpha f(x)|}{1 + |x|^{(2r+1-j)\nu_0}} < \infty, \quad |\alpha|_1 = j, \quad j \in \{0, 1, \dots, 2q+2\}; \quad (2.3)$$

(A2) The diffusion coefficient function $g \in \mathbf{C}^{2q+2}(\mathbb{R}^d; \mathbb{R}^{d \times m})$ and there exists $\rho \leq r+1$ such that

$$\sup_{x \in \mathbb{R}^d} \frac{|D^\alpha g(x)|}{1 + |x|^{(\rho-j)\nu_0}} < \infty, \quad |\alpha|_1 = j, \quad j \in \{0, 1, \dots, 2q+2\}; \quad (2.4)$$

(A3) Let $p_0 \geq 2$ be a sufficiently large number and there exists $c_{p_0} \in \mathbb{R}$ such that

$$\langle Df(x)y, y \rangle + \frac{p_0 - 1}{2} |Dg(x)y|^2 \leq c_{p_0} |y|^2, \quad x, y \in \mathbb{R}^d; \quad (2.5)$$

Remark 2.2. Compared to [31], the conditions are relaxed as follows.

1. In (A2), the condition $\rho \leq r$ is relaxed to $\rho \leq r+1$;
2. In (A3), we only require a large $p_0 \geq 2$ instead of all $p_0 \geq 2$;
3. We remove the following condition (the condition (A4) in [31]), which is cited from [4]: there exist $a_1 > 0$ and $\gamma, c_1 \geq 0$ such that for any $x, y \in \mathbb{R}^d$ it holds

$$\langle f(x+y) - f(x), y \rangle \leq -a_1 |y|^{2r+2} + c_1 (|x|^\gamma + 1). \quad (2.6)$$

In the proof, we don't require $-a_1 < 0$, which is required when $t \rightarrow \infty$ in [4]. The polynomial growth condition (2.10) implies the above condition with $a_1 \geq 0$.

We also assume the following one-sided Lipschitz condition (globally monotone condition), e.g. in [26, 29],

- (A3') $\langle x - y, f(x) - f(y) \rangle + \frac{p'_0 - 1}{2} |g(x) - g(y)|^2 \leq c_{p'_0} |x - y|^2, \quad \forall x, y \in \mathbb{R}^d$, where $p'_0 > 1$.

By utilizing Young's inequality with arbitrarily small $\varepsilon > 0$, we have

$$\langle x, f(x) \rangle + \frac{p'_0 - 1 - \varepsilon}{2} |g(x)|^2 \leq c_0 + c'_{p'_0} |x|^2, \quad \forall x \in \mathbb{R}^d, \quad (2.7)$$

where $\varepsilon > 0$ is arbitrarily small if $g(0) \neq 0$ and $c_0 = \frac{|f(0)|^2}{2} + \frac{(p'_0 - 1)(p'_0 - 1 - \varepsilon)}{2\varepsilon} |g(0)|^2$; and $c'_{p'_0} = c_{p'_0} + \frac{1}{2}$. When $g(0) = 0$, we have $\varepsilon = 0$ and $c_0 = \frac{|f(0)|^2}{2}$. By Itô's formula, the inequality (2.7) is sufficient to ensure bounded moments [13, 16]: there is $C > 0$,

$$\mathbb{E}[|X^{0, x_0}(t)|^{p_1}] \leq C(1 + |x_0|^{p_1}), \quad 2 \leq p_1 \leq p'_0 - \varepsilon, \quad t \in [0, T]. \quad (2.8)$$

We remark that assumption (A3) in Assumption 2.1 implies this condition with $p'_0 \geq p_0$. In fact, the inequality may be proved by using Taylor's expansion and applying the inequality (2.5):

$$\langle x - y, \int_0^1 Df(x + s(y - x))(x - y) dt \rangle + \frac{p_0 - 1}{2} \left| \int_0^1 Dg(x + s(y - x))(x - y) ds \right|^2 \leq c_{p_0} |x - y|^2,$$

where p_0 can be arbitrarily large, e.g. in [4, 31], $p'_0 = p_0 < \infty$. However, it may be true that $p'_0 > p_0$, which is the case in Example 3.16. Even it is redundant when $p'_0 = p_0$, we assume (A3') in addition to Assumption 2.1.

Assumption (A1) immediately implies

$$|f(x) - f(y)| \leq C(1 + |x|^{2r} + |y|^{2r})|x - y|, \quad x, y \in \mathbb{R}^d, \quad (2.9)$$

which implies the polynomial growth at infinity

$$|f(x)| \leq C(1 + |x|^{2r+1}), \quad x \in \mathbb{R}^d. \quad (2.10)$$

Moreover, from (2.7) and (2.9), we may derive

$$|g(x)| \leq C(1 + |x|^{r+1}). \quad (2.11)$$

Under the inequalities (2.9), (2.4), and (2.7), it holds that $r+1 \geq \rho$. If $f(x)$ is a scalar polynomial and satisfies Assumption 2.1, then it is of the form $f(x) = -c_{2k+1}x^{2k+1} + \sum_{i=0}^{2k} c_i x^i$, where $c_{2k+1} > 0$, $c_i \in \mathbb{R}$, $0 \leq i \leq 2k$, and $r \geq k$.

2.2 A theorem on weak convergence order for one-step approximations

To approximate solutions to (2.2), we construct a uniform mesh on $[0, T]$ with $h = \frac{T}{N}$ being the step size, for any $N \in \mathbb{N}$. For $x \in \mathbb{R}^d$, $t \in [0, T]$, $h > 0$, $0 < t + h \leq T$, we introduce the one-step approximation $Y(t, x; t + h)$ for the solution $X(t, x; t + h)$ to (2.2) in the form of

$$Y(t, x; t + h) = x + \Phi(t, x, h; \nu_t), \quad (2.12)$$

where ν_t is a random vector defined on $(\Omega, \mathcal{F}, \mathbb{P})$ with moments of a sufficiently high order and Φ is a function from $[0, T] \times \mathbb{R}^d \times (0, T] \times \mathbb{R}^m$ to \mathbb{R}^d . Using the one-step approximation (2.12), we recurrently construct numerical approximations $\{Y_n\}_{0 \leq n \leq N}$ on the uniform mesh grid $\{t_n = nh, n = 0, 1, \dots, N\}$, given by

$$Y_0 = X_0, \quad Y_{n+1} = Y_n + \Phi(t_n, Y_n, h; \nu_n), \quad n = 0, 1, \dots, N-1, \quad (2.13)$$

where ν_n is independent of $Y_0, Y_1, \dots, Y_n, \nu_0, \nu_1, \dots, \nu_{n-1}$ for all $n \geq 1$.

We will consider the convergence of $\mathbb{E}[\varphi(Y_n)]$ to $\mathbb{E}[\varphi(X_{t_n})]$ where φ and its derivatives have at most a polynomial growth at infinity.

Assumption 2.3. Let $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}^m$ ($m \geq 1$). Assume that $\varphi \in \mathbb{G}^{2q+2}$, i.e., there exist constants $L > 0$ and $\kappa \geq 1$ such that

$$|D^j \varphi(x)| \leq L(1 + |x|^\kappa), \quad j \in \{j = 0, 1, 2, \dots, 2q + 2\}. \quad (2.14)$$

The following theorem has been known if the coefficients are Lipschitz continuous [21, 22] or p_0 being arbitrarily large in (A3) [31]. Here we remove the last constraint and extra constraints as stated in Remark 2.2.

Theorem 2.4 (Fundamental theorem of one-step schemes for weak convergence, c.f. [31]). Let Assumption 2.3 hold. Suppose

- (i) (Coefficients of SDEs) Assumptions 2.1 and (A3') hold;
- (ii) (Local approximation error) For some constants $C > 0$ and $\varkappa \geq 0$, and for any $i_j \in \{1, 2, \dots, d\}$ and $x \in \mathbb{R}^d$, the one-step approximation $Y(t, x; t + h)$ has the following orders of accuracy:

$$\left| \mathbb{E} \left[\prod_{j=1}^s (\delta_{X,x})^{i_j} \right] - \mathbb{E} \left[\prod_{j=1}^s (\delta_{Y,x})^{i_j} \right] \right| \leq C(1 + |x|^\varkappa) h^{q+1}, \quad s = 1, \dots, 2q + 1, \quad (2.15)$$

$$\left\| \prod_{j=1}^{2q+2} (\delta_{X,x})^{i_j} \right\|_{L^2(\Omega; \mathbb{R})} \leq C(1 + |x|^\varkappa) h^{q+1}, \quad (2.16)$$

$$\left\| \prod_{j=1}^{2q+2} (\delta_{Y,x})^{i_j} \right\|_{L^2(\Omega; \mathbb{R})} \leq C(1 + |x|^\varkappa) h^{q+1}, \quad (2.17)$$

where we denote $(\delta_{X,x})^{i_j} := X^{i_j}(t, x; t + h) - x^{i_j}$, $(\delta_{Y,x})^{i_j} := Y^{i_j}(t, x; t + h) - x^{i_j}$.

- (iii) (Moment bounds) There exist constants $\beta \geq 1$ and $C > 0$ such that for $p \geq 2\kappa \vee (\beta\kappa + \varkappa)$, the moments of the approximation $\{Y_n\}_{0 \leq n \leq N}$ are bounded by

$$\sup_{N \in \mathbb{N}} \sup_{0 \leq n \leq N} \mathbb{E}[|Y_n|^p] \leq C(1 + |Y_0|^{\beta p}). \quad (2.18)$$

Then we obtain a global weak convergence order of order q , i.e., for some $C > 0$ independent of h ,

$$\left| \mathbb{E}[\varphi(X(t_0, X_0; T))] - \mathbb{E}[\varphi(Y(t_0, Y_0; t_N))] \right| \leq C(1 + |X_0|^{\beta(\beta\kappa + \varkappa)}) h^q. \quad (2.19)$$

The proof of this theorem is presented in Appendix C and it uses the following lemma.

Lemma 2.5. (c.f. [4, Theorem 1.3.6]) Let Assumption 2.1 and inequality (A3') be fulfilled. Then, the solution to SDE (2.2) $X(t, x; s), 0 \leq t \leq s \leq T$ is $2q + 2$ times differentiable with respect to the initial data $x \in \mathbb{R}^d$ and for any $q \geq 1$ and $j = 2, \dots, 2q + 2$ it holds that

$$\sup_{x \in \mathbb{R}^d} \mathbb{E}[|D^j X(t, x; s)|^p] \leq C(T, \mathbb{P}, j), \quad \text{for all } \mathbb{P} \in \left[1, \frac{p_0}{((2r + 1 - j) \vee 0) + j}\right], \quad (2.20)$$

where $C(T, \mathbb{P}, j)$ is a constant that depends on T, \mathbb{P}, j . Here D^j 's refer to derivatives in x .

This lemma is proved in Appendix B. In fact, we need $\mathbb{P} \geq 4q + 4$ to prove Theorem 2.4.

Remark 2.6. As pointed out in [31], it is often subtle to verify the condition (2.18) on moments of a method $\{Y_n\}_{0 \leq n \leq N}$ in the case of the nonglobal Lipschitz coefficients. Usually, each scheme and each equation require special considerations (see Examples 3.10-3.16). However, it is not difficult to verify the conditions (2.15)-(2.17) following the calculations in [23] where the coefficients are Lipschitz continuous.

3 Schemes of weak convergence based on the Itô-Taylor expansion

Here we recall the following one-step approximation based on the Itô-Taylor expansion, known as the Milstein-Talay method [28]:

$$\begin{aligned} X_{n+1} &= Y_{MT}(t_n, X_n; t_n + h), \\ Y_{MT}(t, x; t + h) &= x + f(x)h + \sum_{r=1}^m \int_t^{t+h} g^r(x) dW_r(s) \\ &\quad + \sum_{r=1}^m \sum_{r_1=1}^m \int_t^{t+h} \int_t^s \Lambda_{r_1} g^r(x) dW_{r_1}(s_1) dW_r(s) + \sum_{r=1}^m \int_t^{t+h} \mathcal{L}g^r(x)h dW_r(s) \\ &\quad + \sum_{r=1}^m \int_t^{t+h} \int_t^s (\Lambda_r f(x) - \mathcal{L}g^r(x)) dW_r(s_1) ds + \frac{h^2}{2} \mathcal{L}f(x). \end{aligned} \quad (3.1)$$

Here $\Lambda_r = \sum_{i=1}^d g^{i,r} \frac{\partial}{\partial x_i}$ and $\mathcal{L} = f^\top \frac{\partial}{\partial x} + \frac{1}{2} \sum_{r=1}^m \sum_{i,j=1}^d g^{i,r} g^{j,r} \frac{\partial^2}{\partial x_i \partial x_j}$. The method is shown to have weak convergence of order two when f and g are Lipschitz continuous, see e.g. [23]. When Lipschitz continuity conditions are violated and coefficients have polynomial growth instead of linear growth, the Milstein-Talay method can explode at a certain time as in the Euler scheme [10]. Inspired by modified Euler schemes in the literature, we propose the following scheme

$$\begin{aligned} Y_{n+1} &= Y_n + \mathcal{T}_1(f(Y_n), h)h + \sum_{r=1}^m \mathcal{T}_2(g^r(Y_n), h) \Delta W_r(n) \\ &\quad + \sum_{r=1}^m \sum_{r_1=1}^m \int_{t_n}^{t_{n+1}} \int_{t_n}^s \mathcal{T}_3(\Lambda_{r_1} g^r(Y_n), h) dW_{r_1}(s_1) dW_r(s) + \sum_{r=1}^m \int_{t_n}^{t_{n+1}} \mathcal{T}_4(\mathcal{L}g^r(Y_n), h)h dW_r(s) \\ &\quad + \sum_{r=1}^m \int_{t_n}^{t_{n+1}} \int_{t_n}^s \mathcal{T}_5((\Lambda_r f(Y_n) - \mathcal{L}g^r(Y_n)), h) dW_r(s_1) ds + \int_{t_n}^{t_{n+1}} \mathcal{T}_6(\mathcal{L}f(Y_n), h) \frac{h}{2} ds, \end{aligned} \quad (3.2)$$

where the maps $\mathcal{T}_i(\cdot)$ ($i = 1, \dots, 6$) are approximations of ' \cdot ' or zero and will be specified in Section 3.1 and $\Delta W_r(n) = W_r(t_{n+1}) - W_r(t_n), n = 0, 1, 2, \dots, N - 1$.

3.1 Assumptions on the maps for bounded moments

To obtain moment boundedness of solutions to the scheme (3.2), we assume the following conditions:

Assumption 3.1. (H1) (Growth conditions on the maps \mathcal{T}_1 and \mathcal{T}_2) There exist positive constants γ_1, γ_2 and C , such that

$$|\mathcal{T}_i(z, h)| \leq Ch^{-\gamma_i} \wedge |z|, \quad i = 1, 2, \quad z \in \mathbb{R}^d. \quad (3.3)$$

(H2) (Approximation condition) There exist $C, \tau, l_1 > 0$ such that for all $z \in \mathbb{R}^d$,

$$|\mathcal{T}_1(z, h) - z| \leq Ch^\tau |z|^{l_1}. \quad (3.4)$$

(H3) (Growth conditions on the $\mathcal{T}_3 - \mathcal{T}_6$) There exist $\gamma_3 > \frac{1}{2}, \gamma_4 > 1$ and $C > 0$ such that for all $z \in \mathbb{R}^d$,

$$|\mathcal{T}_3(z, h)| \leq Ch^{-\gamma_1} \wedge |z|, \quad |\mathcal{T}_4(z, h)| + |\mathcal{T}_5(z, h)| \leq Ch^{-\gamma_3} \wedge |z|, \quad |\mathcal{T}_6(z, h)| \leq Ch^{-\gamma_4} \wedge |z|, \quad (3.5)$$

where γ_1 is from (3.3).

The condition (H2) in Assumption 3.1 may be replaced by the following condition:

(H2') (One-sided Lipschitz condition) For a sufficiently large $p_T \geq 2$, there is a constant $C > 0$ such that,

$$\langle x, \mathcal{T}_1(f(x), h) \rangle + \frac{p_T - 1}{2} \left| \sum_{r=1}^m \mathcal{T}_2(g^r(x), h) \right|^2 \leq C(1 + |x|^2), \quad x \in \mathbb{R}^d. \quad (3.6)$$

It is shown in Section 3.3 that Assumption 3.1 is sufficient to obtain moment bounds of the scheme (3.2).

3.2 Modified Euler schemes and the maps \mathcal{T}_i

In this section, we revisit schemes of at most first-order weak convergence when $\mathcal{T}_i = 0, i = 3, 4, 5, 6$ in (3.2). These choices lead to the modified Euler scheme:

$$Y_{n+1} = Y_n + \mathcal{T}_1(f(Y_n), h)h + \sum_{r=1}^m \mathcal{T}_2(g^r(Y_n), h) \Delta W_r(n), \quad (3.7)$$

where $\Delta W_r(n) := W_r(t_{n+1}) - W_r(t_n), n \in \{0, 1, 2, \dots, N-1\}$. Moreover, the maps $\mathcal{T}_1, \mathcal{T}_2$ satisfy $\mathcal{T}_1: \mathbb{R}^d \times (0, 1) \rightarrow \mathbb{R}^d$ and $\mathcal{T}_2: \mathbb{R}^d \times (0, 1) \rightarrow \mathbb{R}^d$. Recall that the maps $\mathcal{T}_i(z, h), i = 1, 2$ represent an approximation of z when values of $|z|$ are not excessively large. Moreover, $\mathcal{T}_i(z, h)$ is bounded in z and is bounded above by a negative power of the time step size h . In Figure 2, we sketch some popular choices of the map $\mathcal{T}_i(z, h)$ for modified Euler schemes which have at most first-order weak convergence [31].




Figure 2: Illustration of approximations of truncation functions $x\mathbf{1}_{|x| \leq h^{-1}} + \text{sign}(x)h^{-1}\mathbf{1}_{|x| \geq h^{-1}}$, $\frac{x}{1+h|x|}$, and $h^{-1} \tanh(hx)$.

To apply Theorem 2.4, the first step is to discuss the moment bounds of the numerical approximation (3.7). We assume that $\mathcal{T}_1(z, h)$ and $\mathcal{T}_2(z, h)$ are deterministic* and we assume either (H1), (H2) in Assumption 3.1 or (H1), (H2') hold, depending on the specific choices of \mathcal{T}_1 and \mathcal{T}_2 .

*It is possible to apply randomized functions but we only consider deterministic functions for brevity.

Lemma 3.2. Suppose Assumption 2.1, (A3'), and (H1) and (H2) in Assumption 3.1 hold. Let $\{Y_n\}_{0 \leq n \leq N}$ be given by (3.7) and let p'_0 coming from (2.7) be sufficiently large. Then there exist $\beta_1 \geq 1$ and $C > 0$ independent of h such that for all $n = 0, 1, \dots, N$

$$\mathbb{E}[|Y_n|^p] \leq C(1 + |X_0|^{p\beta_1}), \quad \text{for all } p \in [1, \mathbb{B}_1], \quad (3.8)$$

where $\beta_1 = 1 + \frac{(p\gamma_1+1)\mathbb{G}_1}{p} \wedge \frac{(1+\frac{1}{2}p+p\gamma_2)\mathbb{G}_1}{p}$ and $\mathbb{B}_1 = \frac{p'_0-\varepsilon-\mathbb{G}_1}{1+\gamma_1\mathbb{G}_1} \wedge \frac{p'_0-\varepsilon-\mathbb{G}_1}{1+(\frac{1}{2}+\gamma_2)\mathbb{G}_1}$ with γ_1, γ_2 from (H1) in Assumption 3.1. Here $\mathbb{G}_1 = 6r \vee \frac{(2r+1)l_1-1}{\tau}$ with $r \geq 0$ from (A1) and $\tau, l_1 > 0$ from (H2).

If (H2) is replaced by (H2'), then $\mathbb{G}_1 = 6r$ and $\mathbb{B}_1 = \frac{p\tau-\mathbb{G}_1}{1+\gamma_1\mathbb{G}_1} \wedge \frac{p\tau-\mathbb{G}_1}{1+(\frac{1}{2}+\gamma_2)\mathbb{G}_1}$.

This lemma can be proved by extending the results in [5, 32]. See Appendix D for details.

Note that β_1 plays the role of β in Theorem 2.4 (iii). Applying Theorem 2.4, with the moment bounds for the modified Euler scheme (3.7) in Lemma 3.2, we obtain the following theorem.

Theorem 3.3 (Weak convergence order for the modified Euler scheme (3.7)). Let Assumption 2.3 on the test function φ hold. Let Assumptions 2.1 and (A3') be satisfied. Additionally, assume that (H1), (H2) (or (H2')) in Assumption 3.1 hold. Assume further that there exist $C > 0, q_0 > 0$ and $\eta_{q_0} \geq 0$ such that

$$|z - \mathcal{T}_i(z, h)| \leq C(1 + |z|^{\eta_{q_0}})h^{q_0}, \quad z \in \mathbb{R}^d, \quad i = 1, 2. \quad (3.9)$$

If \mathbb{P} in Lemma 2.5 is no less than $4q + 4$ and \mathbb{B}_1 in Lemma 3.2 is no less than $2\kappa \vee (\beta_1\kappa + \varkappa)$, the scheme (3.7) has a global weak convergence of order $q \wedge 1 \leq q_0$:

$$|\mathbb{E}[\varphi(X(t_0, X_0; T))] - \mathbb{E}[\varphi(Y(t_0, Y_0; T))]| \leq C(1 + |X_0|^{\beta_1(\beta_1\kappa + \varkappa)})h^{q \wedge 1},$$

where β_1 is from (3.8), $\varkappa = (2q + 2)(2r + 1)(1 \vee \eta_{q_0})$ and r comes from (2.3).

The proof is similar to that in [31] and we omit the proof. Here we use \varkappa from [31], which can be verified as in Lemmas 5.1 and 5.2. The parameters in Table 1 are essential for weak convergence order of numerical schemes (3.7).

γ_1, γ_2	$\mathbb{G}_1, \mathbb{B}_1, \beta_1$	η_{q_0}, q_0	κ	\varkappa
approximation	moments bound	approximation	test function	minimum moments
Assumption 3.1	Lemma 3.2	Theorem 3.3	Assumption 2.3	Theorem 3.3

Table 1: Essential parameters for weak convergence orders of the modified Euler scheme (3.7)

Remark 3.4. The condition (3.9) is a sufficient but not a necessary condition. For example, in some schemes with $\mathcal{T}_i(z, h; w)$, one could also by $|z - \mathcal{T}_i(z, h; w)|$ to derive similar result as Theorem 3.3.

Below we list some examples of modified Euler scheme (3.7), where \mathcal{T}_1 and \mathcal{T}_2 are explicitly given. Furthermore, Table 2 gives the corresponding parameters for the numerical schemes.

- fully tamed Euler (c.f. [26]), for $\forall z \in \mathbb{R}^d$ and $0 < \alpha_1 \leq 1$,

$$\mathcal{T}_1(z, h; w) = \frac{z}{1 + h^{\alpha_1} |z| + h^{\alpha_1} |w|}, \quad \mathcal{T}_2(w, h; z) = \frac{w}{1 + h^{\alpha_1} |z| + h^{\alpha_1} |w|}. \quad (3.10)$$

- balanced scheme [32]

$$\mathcal{T}_i(z, h) = h^{-1} \tanh(hz), \quad i = 1, 2, \quad z \in \mathbb{R}^d. \quad (3.11)$$

- tamed Euler scheme [5]

$$\mathcal{T}_i(z, h) = \frac{z}{1 + h|z|}, \quad i = 1, 2, \quad z \in \mathbb{R}^d. \quad (3.12)$$

scheme	γ_1	γ_2	\mathbf{G}_1	\mathbf{B}_1	\varkappa	η_{q_0}	q_0
fully tamed (3.10)	α_1	α_1	$6r$	$\frac{p_T - \mathbf{G}_1}{1 + (\frac{1+\alpha_1}{2})\mathbf{G}_1}$	$(2q+2)(2r+1)\eta_{q_0}$	$1 + \varsigma$	$\varsigma\alpha_1$
balanced (3.11)	1	1	$6r \vee \frac{(3-2\varsigma)(2r+1)-1}{2-2\varsigma}$	$\frac{p_0 - \varepsilon - \mathbf{G}_1}{1 + \frac{3}{2}\mathbf{G}_1}$	$(2q+2)(2r+1)\eta_{q_0}$	$3 - 2\varsigma$	$2 - 2\varsigma$
tamed Euler (3.12)	1	1	$6r \vee \frac{(2r+1)(1+\varsigma)-1}{\varsigma}$	$\frac{p_0 - \varepsilon - \mathbf{G}_1}{1 + \frac{3}{2}\mathbf{G}_1}$	$(2q+2)(2r+1)\eta_{q_0}$	$1 + \varsigma$	ς
truncation (3.14)	α	α	$6r \vee \frac{(2r+1)(1+\frac{\varepsilon}{\alpha})-1}{\varepsilon}$	$\frac{p_0 - \varepsilon - \mathbf{G}_1}{1 + (\frac{3}{2} + \alpha)\mathbf{G}_1}$	$(2q+2)(2r+1)\eta_{q_0}$	$1 + \frac{\varepsilon}{\alpha}$	ε

Table 2: Essential parameters for weak convergence orders of the scheme (3.2). Here $\varsigma \in [0, 1]$ and $\varepsilon > 0$.

- modified Euler scheme [31]

$$\mathcal{T}_1(z, h) = \frac{z}{1 + h|z|^2}, \quad \mathcal{T}_2(w, h; z) = \frac{w}{1 + h|z|^2}, \quad z, w \in \mathbb{R}^d. \quad (3.13)$$

- truncation scheme

$$\mathcal{T}_i(z, h) = z \mathbb{1}_{|z| \leq h^{-\alpha}} + \vartheta \operatorname{sgn}(z) h^{-\alpha} \mathbb{1}_{|z| > h^{-\alpha}}, \quad \vartheta = 0 \text{ or } 1, \quad i = 1, 2. \quad (3.14)$$

The fully tamed Euler (3.10) satisfies condition (H1) from Assumption 3.1 with $\gamma_1 = \alpha_1$ and $\gamma_2 = \frac{\alpha_1}{2}$. Due to (2.7), $\mathcal{T}_1(f(x), h; g(x))$ and $\mathcal{T}_2(g(x), h; f(x))$ satisfy (H2'), by Lemma 3.2, $\mathbf{G}_1 = 6r$. Furthermore, due to $|z - \mathcal{T}_i(z, h; w)| = h^{\alpha_1} |z| \frac{|z| + |w|}{1 + h^{\alpha_1}|z| + h^{\alpha_1}|w|} \leq h^{\alpha_1} |z| (|z|^\varsigma + |w|^\varsigma)$, condition (3.9) holds with $q_0 = \varsigma\alpha_1$ and $\eta_{q_0} = 1 + \varsigma$.

For the balanced scheme (3.11), we need to verify the condition (H2) from Assumption 3.1. In fact, by (3.22), we have

$$|h^{-1} \tanh(hz) - z| \leq h^{-1} |\tanh(hz) - hz| \leq h^{-1} |hz|^{3-2\varsigma} = h^{2-2\varsigma} |z|^{3-2\varsigma}, \quad 0 \leq \varsigma \leq 1, \quad (3.15)$$

i.e., $\tau = 2 - 2\varsigma$ and $l_1 = 3 - 2\varsigma$. Then, $\mathbf{G}_1 = 6r \vee \frac{(3-2\varsigma)(2r+1)-1}{2-2\varsigma}$, $\eta_{q_0} = 3 - 2\varsigma$ and $q_0 = 2 - 2\varsigma$.

For the tamed Euler (3.12), condition (H1) is satisfied with $\gamma_1 = \gamma_2 = 1$. Furthermore, we have $|z - \mathcal{T}_1(z, h)| = h |z| \frac{|z|}{1 + h^2|z|} \leq h^\varsigma |z|^{1+\varsigma}$, $0 \leq \varsigma \leq 1$. Then, condition (H2) from Assumption 3.1 hold with $\tau = \varsigma$ and $l_1 = 1 + \varsigma$. So, we get $\mathbf{G}_1 = 6r \vee \frac{(2r+1)(1+\varsigma)-1}{\varsigma}$, $\eta_{q_0} = 1 + \varsigma$ and $q_0 = \varsigma$.

In the modified Euler scheme (3.13), $\mathcal{T}_2(g(x), h; f(x))$ does not satisfy (H1) in Assumption 3.1. We refer interested readers to [31] for the proof of bounded moments when p_0 is arbitrarily large.

It's straightforward to check that for the truncation scheme (3.14), $|\mathcal{T}_i(z, h)| \leq h^{-\alpha} \wedge |z|$, and thus (H1) in Assumption 3.1 hold with $\alpha = \gamma_1 = \gamma_2$. For any $\varepsilon > 0$, we also have

$$|\mathcal{T}_1(z, h) - z| = |\mathcal{T}_1(z, h) - z| \mathbb{1}_{|z| > h^{-\alpha}} \leq |z - \vartheta \operatorname{sgn}(z) h^{-\alpha}| \mathbb{1}_{|z| > h^{-\alpha}} \leq (1 + \vartheta) |z|^{1+\frac{\varepsilon}{\alpha}} h^\varepsilon. \quad (3.16)$$

Subsequently, (H2) in Assumption 3.1 is satisfied with $\tau = \varepsilon$ and $l_1 = 1 + \frac{\varepsilon}{\alpha}$. One can show $\mathbf{G}_1 = 6r \vee \frac{(2r+1)(1+\frac{\varepsilon}{\alpha})-1}{\varepsilon}$, $\eta_{q_0} = 1 + \frac{\varepsilon}{\alpha}$ and $q_0 = \varepsilon$.

Remark 3.5. The moment bounds in Table 2 are consistent with those in the literature. The balanced scheme (3.11) and the tamed Euler scheme (3.12), as discussed in [32] and [5] exhibit the same moment bound for $1 \leq p \leq \mathbf{B}_1$.

3.3 Weak convergence order for the scheme (3.2)

Next, we consider the weak convergence order of the scheme (3.2). According to Theorem 2.4, we need to show the moment bounds of numerical solutions.

Lemma 3.6 (Moment bounds of numerical solutions (3.2)). Let Assumptions 2.1, (A3'), and 3.1 hold. Let p'_0 from (2.7) be sufficiently large. Then there exist $\beta_2 \geq 1$ and $C > 0$ independent of h such that for all $n = 0, \dots, N$

$$\mathbb{E}[|Y_n|^p] \leq C(1 + |X_0|^{\beta_2 p}), \quad \text{for all } p \in [1, \mathbf{B}_2], \quad (3.17)$$

where $\beta_2 = 1 + (\beta_1 - 1) \wedge \frac{(1 - \frac{1}{2}p + p\gamma_3)\mathbf{G}_1}{p} \wedge \frac{(1 - p + p\gamma_4)\mathbf{G}_1}{p}$, and $\mathbf{B}_2 = \mathbf{B}_1 \wedge \frac{p'_0 - \varepsilon - \mathbf{G}_1}{1 + (\gamma_3 - \frac{1}{2})\mathbf{G}_1} \wedge \frac{p'_0 - \varepsilon - \mathbf{G}_1}{1 + (\gamma_4 - 1)\mathbf{G}_1}$. Also, γ_3, γ_4 are from Assumption 3.1. Here \mathbf{G}_1, β_1 and \mathbf{B}_1 come from Theorem 3.3.

If (H2) in Assumption 3.1 is replaced by (H2'), then $\mathbf{G}_1 = 6r$, and $\mathbf{B}_2 = \mathbf{B}_1 \wedge \frac{p_T - \mathbf{G}_1}{1 + (\gamma_3 - \frac{1}{2})\mathbf{G}_1} \wedge \frac{p_T - \mathbf{G}_1}{1 + (\gamma_4 - 1)\mathbf{G}_1}$.

Note that β_2 plays the role of β in Theorem 2.4 (iii). The proof for the above lemma is presented in Section 5.1.

By Theorem 2.4, we need the one-step approximation of the scheme (3.2) to be of certain order. To this end, we assume the following on the maps \mathcal{T}_i 's.

Assumption 3.7. For $\mathcal{T}_i: \mathbb{R}^d \times (0, 1) \rightarrow \mathbb{R}^d$, there exist $C > 0$, $q_0 > 0$ and $\eta_{q_0} \geq 0$ such that for all $z \in \mathbb{R}^d$,

$$\begin{aligned} |z - \mathcal{T}_i(z, h)| &\leq C(1 + |z|^{\eta_{q_0}})h^{q_0}, \quad i = 1, 2, 3, \\ |z - \mathcal{T}_j(z, h)| &\leq C(1 + |z|^{\eta_{q_0}})h^{q_0-1}, \quad j = 4, 5, 6. \end{aligned} \quad (3.18)$$

Under the above assumptions, by Theorem 2.4, our main result is summarized as follows.

Theorem 3.8 (Weak convergence order for the scheme (3.2)). Let Assumption 2.3 hold. Let Assumptions 2.1 and (A3') be satisfied. Additionally, assume that (H1), (H2) (or (H2')), (H3) in Assumption 3.1, and Assumption 3.7 hold. If \mathbb{P} in Lemma 2.5 is no less than $4q + 4$ and $\mathbb{B}_2 \geq 2\kappa \vee (\beta_2\kappa + \varkappa)$, the scheme (3.2) has a global weak convergence order $q \wedge 2 \leq q_0$:

$$|\mathbb{E}[\varphi(X(t_0, X_0; T))] - \mathbb{E}[\varphi(Y(t_0, Y_0; T))]| \leq C(1 + |X_0|^{\beta_2(\beta_2\kappa + \varkappa)})h^{q \wedge 2},$$

where β_2 is from (3.17), $\varkappa = (2q + 2)r_1 \vee r_2$ and $r_1 = (4r + 1)\eta_{q_0}$, $r_2 = (2q + 1)(6r + 1)$ come from Lemmas 5.1 and 5.2. Also, r is from (2.3).

We summarize the key parameters in Table 3 for the weak convergence order of the scheme (3.2).

γ_i 's	$\mathbb{G}_1, \mathbb{B}_2, \beta_2$	η_{q_0}, q_0	κ	\varkappa
approximation	moments bound	approximation	test function	minimum moments
Assumption 3.1	Lemma 3.6	Assumption 3.7	Assumption 2.3	Theorem 3.8

Table 3: Essential parameters for the weak convergence order of the scheme (3.2).

Remark 3.9. The Assumption 3.7 is a sufficient but not a necessary condition. For example, in some schemes with $\mathcal{T}_i(z, h; w)$, one could also by $|z - \mathcal{T}_i(z, h; w)|$ to derive similar result as Theorem 3.8.

3.3.1 Examples of the maps \mathcal{T}_i 's

We now present three examples, where \mathcal{T}_i 's are specified, and present the key parameters for weak convergence orders.

Example 3.10 (Truncation scheme). In the scheme (3.2), we take

$$\mathcal{T}_i(z, h) = z\mathbb{1}_{|z| \leq h^{-\alpha}} + \vartheta \operatorname{sgn}(z)h^{-\alpha}\mathbb{1}_{|z| > h^{-\alpha}}, \quad \vartheta = 0 \text{ or } 1, \quad i = 1, 2, 3, 4, 5, 6. \quad (3.19)$$

It's straightforward to check that $|\mathcal{T}_i(z, h)| \leq h^{-\alpha} \wedge |z|$, and thus (H1) and (H3) in Assumption 3.1 hold with $\alpha = \gamma_1 = \gamma_2 = \gamma_3 = \gamma_4$. As shown in (3.16), for any $\epsilon > 0$, (H2) in Assumption 3.1 is satisfied with $\tau = \epsilon$ and $l_1 = 1 + \frac{\epsilon}{\alpha}$. Additionally, Assumption 3.7 is fulfilled with $\eta_{q_0} = 1 + \frac{\epsilon}{\alpha}$ and $q_0 = \epsilon$, by (3.16). Thus, by Lemmas 3.6 and 5.2, we can show that $\mathbb{G}_1 = 6r \vee \frac{(2r+1)(1+\frac{\epsilon}{\alpha})-1}{\epsilon}$, $\beta_2 = 1 + \frac{(1-p+p\alpha)\mathbb{G}_1}{p}$, $p \in [1, \mathbb{B}_2]$, $\mathbb{B}_2 = \frac{p'_0 - \epsilon - \mathbb{G}_1}{1 + (\frac{1}{2} + \alpha)\mathbb{G}_1}$, and $\varkappa = r_2 \vee (2q + 2)r_1$, $r_1 = (4r + 1)\epsilon$, $r_2 = (2q + 1)(6r + 1)$. Let $\epsilon = \alpha = 2$. If $\mathbb{P} \geq 4q + 4$ and $\mathbb{B}_2 \geq 2\kappa \vee (\beta_2\kappa + \varkappa)$, the weak convergence order is $q \leq q_0 = \alpha = 2$, by Theorem 3.8. The second-order weak convergence is confirmed numerically in Examples 4.1, where enough high order moments of numerical solutions are valid.

scheme (3.2)	γ_i 's	\mathbb{G}_1	\mathbb{B}_2	β_2	\varkappa	η_{q_0}	q_0
truncation (3.19)	α	$6r \vee \frac{(2r+1)(1+\frac{\epsilon}{\alpha})-1}{\epsilon}$	$\frac{p'_0 - \epsilon - \mathbb{G}_1}{1 + (\frac{1}{2} + \alpha)\mathbb{G}_1}$	$1 + \frac{(1-p+p\alpha)\mathbb{G}_1}{p}$	$r_2 \vee (2q + 2)r_1$	$1 + \frac{\epsilon}{\alpha}$	ϵ

Table 4: Essential parameters for the weak convergence order of the scheme in Example 3.10

Remark 3.11 (Structure preserving schemes). Taking $\vartheta = 0$ in (3.19) and applying structure-preserving schemes for SDEs with Lipschitz continuous coefficients may lead to structure-preserving schemes for SDEs with non-Lipschitz coefficients. In this case, we may also stop the trajectories where truncation is needed and thus we have an acceptance-rejection stordergy similar to that in [24]. Therein, the authors discard the approximate trajectories that leave a sufficiently large ball $S_R := \{x : |x| < R\}$. This strategy has been tested on quasi-symplectic schemes for nonlinear stochastic Hamiltonian systems. A similar truncation strategy has been considered in [15], which may be structure-preserving while the scheme requires small time sizes for stability. We will not discuss this research direction for brevity as verifying that the resulting schemes admit enough high-order moments for convergence order is necessary.

Example 3.12. In the scheme (3.2), we propose the balanced scheme as follows[†]: for all $z \in \mathbb{R}^d$,

$$\mathcal{T}_i(z, h) = h^{-2} \tanh(h^2 z), \quad i = 1, 2, 3, 4, 5, 6. \quad (3.20)$$

It's clear that (H1) and (H3) from Assumption 3.1 are fulfilled when $\gamma_1 = \gamma_2 = \gamma_3 = \gamma_4 = 2$. Also, based on (3.22), it holds for any $\varsigma \in [0, 1]$ that

$$|\mathcal{T}_i(z, h) - z| \leq h^{-2} (h^2 |z|)^{3-2\varsigma} = h^{4-4\varsigma} |z|^{3-2\varsigma}, \quad i = 1, 2, 3, 4, 5, 6. \quad (3.21)$$

This implies that (H2) in Assumption 3.1 is valid with $\tau = 4 - 4\varsigma$ and $l_1 = 3 - 2\varsigma$, and Assumption 3.7 holds with $q_0 = 4 - 4\varsigma$ and $\eta_{q_0} = 3 - 2\varsigma$. By Lemmas 3.6 and 5.2, $\mathbb{G}_1 = 6r \vee \frac{(2r+1)(3-2\varsigma)-1}{4-4\varsigma}$, $\mathbb{B}_2 = \frac{p'_0 - \varepsilon - \mathbb{G}_1}{1 + \frac{1}{2}\mathbb{G}_1}$, $\beta_2 = 1 + \frac{(p+1)\mathbb{G}_1}{p}$, $p \in [1, \mathbb{B}_2]$, and $\varkappa = r_2 \vee (2q+2)r_1$, $r_1 = (4r+1)\eta_{q_0}$, $r_2 = (2q+1)(6r+1)$. By Theorem 3.8, the weak convergence order is $q \wedge 2 \leq q_0 = 2$, if $\mathbb{P} \geq 4q+4$ and $\mathbb{B}_2 \geq 2\kappa \vee (\beta_2\kappa + \varkappa)$. We also present numerical result (Example 4.1) and verify the second-order weak convergence.

scheme (3.2)	γ_i 's	\mathbb{G}_1	\mathbb{B}_2	β_2	\varkappa	η_{q_0}	q_0
truncation (3.20)	2	$6r \vee \frac{(2r+1)(3-2\varsigma)-1}{4-4\varsigma}$	$\frac{p'_0 - \varepsilon - \mathbb{G}_1}{1 + \frac{1}{2}\mathbb{G}_1}$	$1 + \frac{(p+1)\mathbb{G}_1}{p}$	$r_2 \vee (2q+2)r_1$	$3 - 2\varsigma$	$4 - 4\varsigma$

Table 5: Essential parameters for the weak convergence order of the scheme in Example 3.12

Remark 3.13. The choices of the maps can be made more general as long as the maps satisfy Assumption 3.7. We use the hyperbolic tangent function since it has the following properties:

$$\begin{aligned} |\tanh(y)| &\leq 1, \quad |\tanh(y)| \leq |y|, \\ |y - \tanh(y)| &\leq \tanh^2(\theta y) |y|, \quad \text{for some } 0 \leq \theta \leq 1, \\ |y - \tanh y| &\leq |y|^{3-2\varsigma}, \quad \text{for any } 0 \leq \varsigma \leq 1. \end{aligned} \quad (3.22)$$

Under some certain conditions, Assumption 3.7 can be extended when the mappings \mathcal{T}_i depend on some $w \in \mathbb{R}^d$, written as $\mathcal{T}_i = \mathcal{T}_i(z, h; w)$.

Example 3.14. In the scheme (3.2), we apply the following modification maps, where

$$\mathcal{T}_i = \mathcal{T}_i(z, h; w) = \frac{z}{1 + h^2 |z| + h^2 |w|}, \quad i = 1, 2; \quad \mathcal{T}_j = \mathcal{T}_j(z, h) = \frac{z}{1 + h^2 |z|}, \quad j = 3, 4, 5, 6. \quad (3.23)$$

Evidently, (H1) and (H3) in Assumption 3.1 are fulfilled with $\gamma_1 = \gamma_2 = \gamma_3 = \gamma_4 = 2$. Moreover, for any $\varsigma \in [0, 1]$, we have

$$|z - \mathcal{T}_i(z, h; w)| = h^2 |z| \frac{|z| + |w|}{1 + h^2 |z| + h^2 |w|} \leq h^{2\varsigma} |z| (|z|^\varsigma + |w|^\varsigma), \quad i = 1, 2,$$

and

$$|z - \mathcal{T}_j(z, h)| = h^2 |z| \frac{|z|}{1 + h^2 |z|} \leq h^{2\varsigma} |z|^{1+\varsigma}, \quad j = 3, 4, 5, 6.$$

[†]For convenience all \mathcal{T}_i are the same. We can take $\mathcal{T}_2(z, h) = h^{-\frac{3}{2}} \tanh(h^{-\frac{3}{2}} z)$ which may lead to a smaller \mathbb{B}_2 .

i.e. (H2) in Assumption 3.1 and Assumption 3.7 are satisfied with $\tau = 2\varsigma$, $l_1 = 1 + \varsigma$, $\eta_{q_0} = 1 + \varsigma$, and $q_0 = 2\varsigma$. It can be readily verified that (H2') is satisfied with $p\tau = p'_0 - \varepsilon$ if we take

$$\mathcal{T}_1(f(x), h) = \mathcal{T}_1(f(x), h; g(x)), \quad \mathcal{T}_2(g(x), h) = \mathcal{T}_2(g(x), h; f(x)). \quad (3.24)$$

Subsequently, applying Lemmas 3.6 and 5.2 yields that $\mathbb{G}_1 = 6r$, $\mathbb{B}_2 = \frac{p\tau - \mathbb{G}_1}{1 + \frac{\varsigma}{2}\mathbb{G}_1}$, $\beta_2 = 1 + \frac{(p+1)\mathbb{G}_1}{p}$, $p \in [1, \mathbb{B}_2]$, and $\varkappa = r_2 \vee (2q + 2)r_1$, $r_1 = (4r + 1)(1 + \varsigma)$, $r_2 = (2q + 1)(6r + 1)$. The weak convergence order becomes $q \leq q_0 = 2$ when $\mathbb{P} \geq 4q + 4$ and $\mathbb{B}_2 \geq 2\kappa \vee (\beta_2\kappa + \varkappa)$, according to Theorem 3.8. We numerically verify this scheme (3.24) in Example 4.1, confirming the second-order weak convergence if enough moments of numerical solutions exist.

scheme (3.2)	γ_i 's	\mathbb{G}_1	\mathbb{B}_2	β_2	\varkappa	η_{q_0}	q_0
modified (3.23)	2	$6r$	$\frac{p\tau - \mathbb{G}_1}{1 + \frac{\varsigma}{2}\mathbb{G}_1}$	$1 + \frac{(p+1)\mathbb{G}_1}{p}$	$r_2 \vee (2q + 2)r_1$	$1 + \varsigma$	2ς

Table 6: Essential parameters for the weak convergence order of the scheme in Example 3.14. Here $\varsigma \in [0, 1]$.

3.4 Examples of weak convergence order depending on moments bounds

Example 3.15. Consider the following SDE with Lipschitz diffusion coefficient

$$dX(t) = (X(t) - X^3(t)) dt + X(t) dW(t), \quad t \in (0, T], \quad X(0) = x. \quad (3.25)$$

Assumption 2.1 is fulfilled with $r = 1$, $\rho = 1$ while $p_0 < +\infty$ and $c_{p_0} = \frac{p_0+1}{2}$ in (2.5). Also, the inequality (A3') holds with $p'_0 = p_0 < +\infty$. As p'_0 can be arbitrarily large, \mathbb{B}_1 and \mathbb{B}_2 are arbitrarily large and thus both are larger than $2\kappa \vee (\beta_2\kappa + \varkappa)$. Recall that $\varkappa = (2q + 2)(4r + 1)\eta_{q_0} \vee (2q + 1)(6r + 1)$ for the scheme (3.2) and $\varkappa = (2q + 2)(2r + 1)(1 \vee \eta_{q_0})$ for the scheme (3.7). By Theorems 3.3 and 3.8, the weak convergence orders are $q_0 \wedge 2$ and $q_0 \wedge 1$.

Applying Theorem 3.8, we obtain the second-order weak convergence of the truncation scheme (3.19) with $\alpha = 2$, the balanced scheme (3.20), and the modified scheme (3.24) with $\varphi(x) = \cos x$, x^2 . By Theorem 3.3, we

scheme (3.2)	\mathbb{P}	\mathbb{G}_1	\mathbb{B}_2	\varkappa	$2\kappa \vee (\beta_2\kappa + \varkappa)$	η_{q_0}	q_0	q (order)
truncation (3.19) ($\alpha = 2$)	12	6	$\frac{p'_0 - \varepsilon - 6}{16}$	60	60	2	$= \varepsilon = 2$	2
balanced (3.20)	12	6	$\frac{p'_0 - \varepsilon - 6}{16}$	60	60	2	2	2
modified (3.23)	12	6	$\frac{p'_0 - \varepsilon - 6}{16}$	60	60	2	2	2
scheme (3.7)	\mathbb{P}	\mathbb{G}_1	\mathbb{B}_1	\varkappa	$2\kappa \vee (\beta_2\kappa + \varkappa)$	η_{q_0}	q_0	q (order)
fully tamed (3.10) ($\alpha_1 = 1$)	6	6	$\frac{p'_0 - \varepsilon - 6}{7}$	18	18	2	$= \alpha_1$	α_1
balanced (3.11)	8	6	$\frac{p'_0 - \varepsilon - 6}{10}$	24	24	2	1	1
tamed Euler (3.12)	8	6	$\frac{p'_0 - \varepsilon - 6}{10}$	24	24	2	1	1
truncation (3.14)	8	6	$\frac{p'_0 - \varepsilon - 6}{10}$	24	24	2	$= \alpha = 1$	1

Table 7: Essential parameters for weak convergence order of the schemes in Example 3.15, $\varphi(x) = \cos x$

may show first-order weak convergence of the fully tamed Euler (3.10) with $\alpha_1 = 1$, the balanced scheme (3.11), the tamed Euler scheme (3.12), the modified Euler scheme (3.13), and the truncation scheme (3.19) with $\alpha = 1$. In Table 7, we present the essential parameters in case of $\varphi(x) = \cos x$, where $\kappa = 0$. The weak convergence orders are numerically verified in Figure 3 in Section 4.

The next example shows that the weak convergence order deteriorates if the numerical solutions don't have enough high-order moments.

Example 3.16. Consider the following stochastic differential equations

$$dX(t) = (X(t) - X^3(t)) dt + \sigma X^2(t) dW(t), \quad t \in (0, T], \quad X(0) = x_0. \quad (3.26)$$

Assumption 2.1 is fulfilled with $r = 1$ and $\rho = 2$ while $p_0 = 3/(2\sigma^2) + 1$ and $c_{p_0} = 1$ in (2.5). Also, we may check that for all $p'_0 \leq \frac{2}{\sigma^2} + 1$, it holds that $x(x - x^3) + \frac{p'_0 - 1}{2}\sigma^2 x^4 \leq x^2$. Take $\varphi(x) = \cos x$ and two sets of model parameters: **Case I:** $\sigma = 0.1$, **Case II:** $\sigma = 0.5$. In the calculations, we may observe that q is not an integer while the order q in some previous theorems is required to be an integer. For simplicity, we keep q as is for the discussion while q in assumptions and lemmas should be replaced by the rounding of q to its nearest integer.

When $\sigma = 0.1$, $p'_0 \leq 201$. For the balanced scheme (3.20), $\mathbb{B}_2 = \frac{p'_0 - \epsilon - 6}{16} \approx 12.2$, $q_0 = 4 - 4\varsigma$, $\eta_{q_0} = 3 - 2\varsigma$, $\varsigma \in [0, 1]$. According to Theorem 3.8, we need $\mathbb{B}_2 \geq 2\kappa \vee (\beta_2 \kappa + \varkappa) = \varkappa$, i.e. $12.2 \geq \varkappa = (2q + 2)(4r + 1)\eta_{q_0} \vee (2q + 1)(6r + 1) = 10(q + 1)\eta_{q_0} \vee 7(2q + 1)$. Since $q \leq q_0 = 4 - 4\varsigma$, we take $\varsigma = 0.964970$ and thus $\eta_{q_0} \approx 1$ and $q \leq 0.14$. For the scheme (3.11), $\mathbb{B}_1 = \frac{p'_0 - \epsilon - 6}{10} \approx 19.5$, $q_0 = 2 - 2\varsigma$, $\eta_{q_0} = 3 - 2\varsigma$, $\varsigma \in [0, 1]$. By Theorem 3.3, we need $\mathbb{B}_1 \geq 2\kappa \vee (\beta_1 \kappa + \varkappa) = \varkappa$, i.e. $19.5 \geq \varkappa = (2q + 2)(2r + 1)(1 \vee \eta_{q_0}) = 6(q + 1)(1 \vee (3 - 2\varsigma))$. As $q \leq q_0 = 2 - 2\varsigma$, we may take $\varsigma \approx 0.965$ and thus $q \leq q_0 = 2 - 2\varsigma \approx 0.803$. For the rest of schemes, calculations are similar and we present results in Table 8.

Observe that when $\sigma = 0.5$, $p'_0 \leq 9$ while Theorem 3.8 requires $\varkappa = (2q + 2)(4r + 1)\eta_{q_0} \vee (2q + 1)(6r + 1)$ -th moments of numerical solutions. In this case, we cannot draw any conclusion on the convergence order. For the scheme (3.19) with $\alpha = 2$, the schemes (3.20) and (3.24), we present numerical results in Example 4.2, where we observe a weak convergence order around one.

scheme (3.2)	\mathbb{G}_1	\mathbb{B}_2	$\varkappa = 10(q + 1)\eta_{q_0} \vee 7(2q + 1)$	q_0	q (order)
Truncation (3.19)($\alpha = 2$)	6	$\frac{p'_0 - \epsilon - 6}{16} \approx 12.2$	$10(q + 1)(1 + \frac{\epsilon}{\alpha}) \vee 7(2q + 1)$	ϵ	≈ 0.14
Balanced (3.20)	6	$\frac{p'_0 - \epsilon - 6}{16} \approx 12.2$	$10(q + 1)(3 - 2\varsigma) \vee 7(2q + 1)$	$4 - 4\varsigma$	≈ 0.14
Modified (3.23)	6	$\frac{p'_0 - \epsilon - 6}{16} \approx 12.2$	$10(q + 1)(1 + \varsigma) \vee 7(2q + 1)$	2ς	≈ 0.14
scheme (3.7)	\mathbb{G}_1	\mathbb{B}_1	$\varkappa = (2q + 2)(2r + 1)\eta_{q_0}$	q_0	q (order)
fully tamed (3.10) ($\alpha_1 = 1$)	6	$\frac{p'_0 - \epsilon - 6}{7} \approx 27.86$	$3(2q + 2)(1 + \varsigma)$	$= \varsigma = 1$	$= \varsigma = 1$
balanced (3.11)	6	$\frac{p'_0 - \epsilon - 6}{10} \approx 19.5$	$3(2q + 2)(3 - 2\varsigma)$	$2 - 2\varsigma$	≈ 0.803
tamed Euler (3.12)	6	$\frac{p'_0 - \epsilon - 6}{10} \approx 19.5$	$3(2q + 2)(1 + \varsigma)$	ς	≈ 0.803
truncation (3.14) ($\alpha = 1$)	6	$\frac{p'_0 - \epsilon - 6}{10} \approx 19.5$	$3(2q + 2)(1 + \frac{\epsilon}{\alpha})$	ϵ	≈ 0.803

Table 8: **Case I:** $\sigma = 0.1$. Essential parameters for checking the weak convergence orders of schemes for (3.26) with $\varphi(x) = \cos x$. Here $2\kappa \vee (\beta\kappa + \varkappa) = \varkappa$, $\epsilon > 0$, and $\varsigma \in [0, 1]$.

4 Numerical Experiments

In this section, we will test the scheme (3.2) for three examples with different maps \mathcal{T}_i 's: the truncation scheme (3.19) with $\alpha = 2$ (denoted by **TS2**); the balanced scheme (3.20) (**BS2**); the modified Euler scheme (3.24) (**MS2**). In Example 4.2, we also compare these schemes with those of at most first-order weak convergence from Section 3.2: the balanced scheme (3.11) (**BS1**), the modified Euler scheme (3.13) (**MS1**) and the truncation scheme (3.14) with $\alpha = 1$ (**TS1**).

We present numerical results for schemes in three examples: the first has one-sided Lipschitz drift and Lipschitz diffusion; the second has locally Lipschitz drift and diffusion, where the solution has limited moments; the third is two-dimensional.

In the experiments, we compare their weak error and computational costs. We consider different test functions $\varphi(x) = x, x^2, \cos x$ in Assumption 2.3 and we use a large number of Monte Carlo samples with a small enough time step size h_{ref} to compute references: $\mathbb{E}[\varphi(X_T)] \approx \frac{1}{M} \sum_{i=1}^M \varphi(X_T(\omega_i, h_{ref}))$. Here M is sufficiently large to have negligible statistical errors with the 95% confidence interval. In simulations, we use the Mersenne twister algorithm [19] to generate pseudorandom numbers. The experiments are performed using Matlab R2022a on a desktop (31.7 GB RAM, 12th Gen Intel(R) Core(TM) i9-12900H CPU at 2.50 GHz) with 64-bit Windows 11 operating system.

Example 4.1. Consider the following SDE in Example 3.15

$$dX(t) = (X(t) - X^3(t)) dt + X(t) dW(t), \quad t \in (0, T], \quad X(0) = x. \quad (4.1)$$

In this example, we fix $T = 2$, the initial condition $x = 0.5$, and take $h_{ref} = 10^{-3}$ and $M = 2 \times 10^6$. In Figure 3, we present weak errors of three different numerical schemes (**TS2**, **MS2** and **BS2**) with the test functions $\varphi(x) = x^2, \cos x$ against different step sizes 0.002, 0.003, 0.004 on a log-log scale. As predicted in Example 3.15, the weak convergence orders of the proposed schemes are close to 2.




Figure 3: Example 4.1: weak convergence orders with $\varphi(x) = x^2$ (Left) and $\varphi(x) = \cos(x)$ (Right).

Example 4.2. Consider the equation in Example 3.16:

$$dX(t) = (X(t) - X^3(t)) dt + \sigma X^2(t) dW(t), \quad t \in (0, T], \quad X(0) = x_0 = 0.1.$$

Consider **Case I** ($\sigma = 0.1$) and **Case II** ($\sigma = 0.5$) in Example 3.16. We first investigate the weak convergence order when $\sigma = 0.1$. For the reference solution, we take $h_{ref} = 2^{-13}$ and $M = 3 \times 10^6$. We present in Table 9 weak errors of the selected schemes for different step sizes $h = 2^{-2}, 2^{-3}, 2^{-4}, 2^{-5}, 2^{-6}, 2^{-7}$ where we use the test function $\varphi(x) = \cos x$. The number of trajectories $M = 3 \times 10^6$ yields statistical errors that are 10 times smaller than the reported weak errors. We omit the statistical errors for brevity. We underline errors at the order of $2 \times 10^{-4} \sim 4 \times 10^{-4}$ in Table 9 and their corresponding computational time at the bottom of the table. We observe that schemes in the second to fourth columns are more efficient than their corresponding schemes in the last three columns. The empirical convergence orders in Table 9 are higher than the theoretical predictions. According to Theorems 3.8 and 3.3, the weak convergence order with $\varphi(x) = \cos x$ should be less than 1 for the TS2, BS2, MS2, and approximately 1 for TS1, BS1, MS1 schemes. Here we conjecture that the step sizes are not small enough to apply Theorems 3.3 and 3.8. To verify our conjecture, we further take $h_{ref} = 10^{-6}$ and $M = 10^8$ and present results in Table 10 weak errors of the schemes with $\varphi(x) = \cos x$ for smaller step-size $h = 5 \times 10^{-4}, 10^{-3}, 2 \times 10^{-3}, 4 \times 10^{-3}$. We observe that the convergence orders of scheme (3.2) (**TS2**, **BS2**, **MS2**) decrease to almost one as time step-sizes become smaller and smaller. We could expect that the convergence order should be less than one if we take even smaller time step sizes. We also observe that the schemes of **TS2**, **BS2** and **MS2** have better accuracy than schemes of **TS1**, **BS1** and **MS1**.

For **Case II**, where $\sigma = 0.5$, we do not expect the numerical schemes (**TS2**, **MS2** and **BS2**) to have weak convergence order of 1 or 2 (see discussions in Example 3.16). However, we observe a first order weak convergence in Figure 4, where we present the weak errors of three different numerical schemes (**TS2**, **MS2** and **BS2**) with the test functions $\varphi(x) = x^2, \cos x$. In this case, the empirical observation does not match the theoretical prediction. We conjecture that the mismatch may disappear when much smaller step sizes are used. However, we cannot perform such experiments due to limited computational resources.

Example 4.3. Consider the stochastic FitzHugh-Nagumo (FHN) model [3] in the form of

$$\begin{pmatrix} dX_1(t) \\ dX_2(t) \end{pmatrix} = \begin{pmatrix} X_1(t) - X_1^3(t) - X_2(t) \\ X_1(t) - X_2(t) + 1 \end{pmatrix} dt + \begin{pmatrix} X_1(t) + 1 & 0 \\ 0 & X_2(t) + 1 \end{pmatrix} dW(t), \quad (4.2)$$

for $t \in (0, T]$ and $X(0) = (0.8, 0.8)^\top$, with solution $X(t) := (X_1(t), X_2(t))^\top$ for $t \in [0, T]$, where $W(t) := (W_1(t), W_2(t))^\top$ is a two-dimensional Brownian motion.

h	TS2	order	BS2	order	MS2	order	TS1	order	BS1	order	MS1	order
2^{-7}	8.11e-07	-	8.11e-07	-	1.46e-06	-	<u>2.21e-04</u>	-	<u>2.22e-04</u>	-	<u>3.00e-04</u>	-
2^{-6}	2.45e-06	1.59	2.45e-06	1.59	5.02e-06	1.78	4.43e-04	1.00	4.44e-04	1.00	5.99e-04	1.00
2^{-5}	8.21e-06	1.74	8.21e-06	1.75	1.83e-05	1.87	8.78e-04	0.99	8.79e-04	0.99	1.18e-03	0.98
2^{-4}	2.93e-05	1.83	2.93e-05	1.83	6.84e-05	1.90	1.71e-03	0.96	1.71e-03	0.96	2.27e-03	0.94
2^{-3}	1.08e-04	1.88	1.08e-04	1.88	<u>2.54e-04</u>	1.89	3.25e-03	0.92	3.26e-03	0.93	4.20e-03	0.89
2^{-2}	<u>3.96e-04</u>	1.88	<u>3.97e-04</u>	1.88	9.04e-04	1.83	5.89e-03	0.86	5.91e-03	0.86	7.33e-03	0.80
	7.36e+02		7.53e+02		6.52e+02		7.48e+02		7.52e+02		7.16e+02	

Table 9: **Case I** in Example 4.2: weak errors with the test function $\varphi(x) = \cos x$ of the selected schemes at $T = 1$. CPU time (in seconds) is listed at the bottom for schemes with underlined errors.

h	TS2	order	BS2	order	MS2	order	TS1	order	BS1	order	MS1	order
5×10^{-4}	2.66e-08	-	2.66e-08	-	2.93e-08	-	1.45e-05	-	1.45e-05	-	1.97e-05	-
1×10^{-3}	5.91e-08	1.15	5.91e-08	1.15	6.98e-08	1.25	2.90e-05	1.00	2.90e-05	1.00	3.94e-05	1.00
2×10^{-3}	1.32e-07	1.16	1.32e-07	1.16	1.74e-07	1.32	5.79e-05	1.00	5.79e-05	1.00	7.87e-05	1.00
4×10^{-3}	3.13e-07	1.25	3.13e-07	1.25	4.83e-07	1.47	1.16e-04	1.00	1.16e-04	1.00	1.57e-04	1.00

Table 10: **Case I** in Example 4.2: weak errors with $\varphi(x) = \cos x$ for smaller step sizes of the selected schemes at $T = 1$.




Figure 4: **Case II** in Example 4.2: weak convergence orders with $\varphi(x) = x^2$ (Left) and $\varphi(x) = \cos(x)$ (Right).

We fix $T = 2$ and take $h_{ref} = 10^{-3}$ and $M = 2 \times 10^6$ to generate a reference solution.

In this example, Assumption 2.1 is satisfied with $r = 1$ and $\rho = 1$ and $p_0 < +\infty$. Also, $p'_0 = p_0 < \infty$ in the assumption (A3'), according to the discussions in Section 3.3.1, all the schemes of **TS2**, **MS2** and **BS2** have weak convergence of order two. In Figure 5, we present the weak errors of these three numerical schemes with the test functions ($\varphi(x) = x, \cos x$) and step sizes $h = 0.002, 0.003, 0.004$. We observe that the weak convergence orders of these schemes are close to 2 in agreement with the theoretical prediction.

5 Weak convergence of second-order numerical schemes

In this section, we prove the second-order weak convergence for the proposed scheme (3.2) under the assumptions in Sections 2 and 3.




Figure 5: Example 4.3: weak convergence orders of the scheme (3.2) with $\varphi(x) = x$ (Left) and $\varphi(x) = \cos(x)$ (Right).

5.1 Moment bound of second-order numerical schemes

We first prove the moment bound of second-order numerical schemes (3.2).

Proof of Lemma 3.6. To prove the boundedness of moments, we adopt a proof strategy similar to that of Lemma 3.2. However, we provide only the necessary details to streamline our presentation. The key to proving the boundedness of moments is to estimate the growth of the solution under some events

$$\Omega_{\mathcal{R},n} := \left\{ \omega \in \Omega : \sup_{0 \leq i \leq n} |Y_i(\omega)| \leq \mathcal{R} \right\}, \quad n = 0, 1, \dots, N, \quad N \in \mathbb{N}.$$

For integer $\bar{p} \geq 1$, We have

$$\begin{aligned} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n+1}} |Y_{n+1}|^{\bar{p}}] &\leq \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}] + \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} (\bar{p} \langle Y_n, Y_{n+1} - Y_n \rangle + \frac{\bar{p}(\bar{p}-1)}{2} |Y_{n+1} - Y_n|^2)] \\ &\quad + C \sum_{l=3}^{\bar{p}} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-l} |Y_{n+1} - Y_n|^l] \\ &:= \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}] + I_1 + I_2. \end{aligned} \quad (5.1)$$

Compared to the proof of bounded moments for the scheme (3.7), it is essential to provide a proper upper bound for I_1 . Similar to the proof of the upper bound for I_1 in Lemma 3.2, we have

$$\begin{aligned} I_1 &\leq \bar{p} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} \langle Y_n, \mathcal{T}_1(f(Y_n), h)h \rangle] + \bar{p} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} \langle Y_n, \int_{t_n}^{t_{n+1}} \mathcal{T}_6(\mathcal{L}f(Y_n), h) \frac{h}{2} ds \rangle] \\ &\quad + \frac{\bar{p}(\bar{p}-1)}{2} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} |\mathcal{T}_1(f(Y_n), h)h|^2] + \frac{\bar{p}(\bar{p}-1)}{2} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} |\sum_{r=1}^m \mathcal{T}_2(g^r(Y_n), h) \Delta W_r(n)|^2] \\ &\quad + \frac{\bar{p}(\bar{p}-1)}{2} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} |\sum_{r=1}^m \sum_{r_1=1}^m \int_{t_n}^{t_{n+1}} \int_{t_n}^s \mathcal{T}_3(\Lambda_{r_1} g^r(Y_n), h) dW_{r_1}(s_1) dW_r(s)|^2] \\ &\quad + \bar{p}(\bar{p}-1) \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} |\sum_{r=1}^m \int_{t_n}^{t_{n+1}} \mathcal{T}_4(\mathcal{L}g^r(Y_n), h) h dW_r(s)|^2] \\ &\quad + \bar{p}(\bar{p}-1) \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} |\sum_{r=1}^m \int_{t_n}^{t_{n+1}} \int_{t_n}^s \mathcal{T}_5((\Lambda_r f(Y_n) - \mathcal{L}g^r(Y_n)), h) dW_r(s_1) ds|^2] \\ &\quad + \frac{\bar{p}(\bar{p}-1)}{2} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} |\int_{t_n}^{t_{n+1}} \mathcal{T}_6(\mathcal{L}f(Y_n), h) \frac{h}{2} ds|^2] \end{aligned}$$

$$\begin{aligned}
& +\bar{p}(\bar{p}-1)\mathbb{E}\left[\mathbf{1}_{\Omega_{\mathcal{R},n}}|Y_n|^{\bar{p}-2}\left\langle\sum_{r=1}^m\mathcal{T}_2(g^r(Y_n),h)\Delta W_r(n),\sum_{r=1}^m\int_{t_n}^{t_{n+1}}\mathcal{T}_4(\mathcal{L}g^r(Y_n),h)h\,dW_r(s)\right.\right. \\
& \left.\left.+\sum_{r=1}^m\int_{t_n}^{t_{n+1}}\int_{t_n}^s\mathcal{T}_5((\Lambda_r f(Y_n)-\mathcal{L}g^r(Y_n)),h)\,dW_r(s_1)ds\right\rangle\right]. \tag{5.2}
\end{aligned}$$

By Schwarz inequality, (D.2), (H3), (A1) and (A2), we have

$$\begin{aligned}
I_1 & \leq Ch + Ch\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^{\bar{p}}] + Ch^2\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^{4r+\bar{p}}] + Ch^{\tau+1}\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^{(2r+1)l_1+\bar{p}-1}] \\
& \quad + Ch^3\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^{4r+2\rho+\bar{p}-2}] + Ch^4\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^{8r+\bar{p}}]. \tag{5.3}
\end{aligned}$$

The scheme for estimating I_2 in (5.1) is analogous to the approach used to estimate I_2 in (D.3). By leveraging Assumption (H3) and an elementary inequality, we can show that

$$\begin{aligned}
I_2 & \leq C\sum_{l=3}^{\bar{p}}\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}}|Y_n|^{\bar{p}-l}(h^l|\mathcal{T}_1(f(Y_n),h)|^l + h^{\frac{l}{2}}|\sum_{r=1}^m\mathcal{T}_2(g^r(Y_n),h)|^l + h^l|\Lambda_{r_1}g^r(Y_n)|^l + h^{\frac{3l}{2}}|\mathcal{L}g^r(Y_n)|^l \\
& \quad + h^{2l}|\Lambda_r f(Y_n) - \mathcal{L}g^r(Y_n)|^l + h^{\frac{3l}{2}}|\mathcal{L}f(Y_n)|^l)] \\
& \leq Ch + C\sum_{l=3}^{\bar{p}}h^l\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}}|Y_n|^{2rl+\bar{p}}] + C\sum_{l=3}^{\bar{p}}h^{\frac{l}{2}}\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}}|Y_n|^{rl+\bar{p}}] \\
& \quad + C\sum_{l=3}^{\bar{p}}h^{\frac{3l}{2}}\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}}|Y_n|^{3rl+\bar{p}}] + C\sum_{l=3}^{\bar{p}}h^{2l}\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}}|Y_n|^{4rl+\bar{p}}]. \tag{5.4}
\end{aligned}$$

Similar to the proof of Lemma 3.2, we set $\mathcal{R} = \mathcal{R}(h) = h^{-1/G}$ with $G_1 = 6r \vee \frac{(2r+1)l_1-1}{\tau}$ to show that

$$\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}}|Y_n|^{\bar{p}}] \leq C(1 + |Y_0|^{\bar{p}}). \tag{5.5}$$

It remains to estimate $\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}^c}|Y_n|^{\bar{p}}]$. It follows from (3.2), (D.6) and Assumption (H3) that

$$\begin{aligned}
|Y_{n+1}| & \leq |Y_n| + Ch^{1-\gamma_1} + \sum_{r=1}^m Ch^{-\gamma_2}|W_r(t_{n+1}) - W_r(t_n)| + \sum_{r=1}^m \sum_{r_1=1}^m Ch^{-\gamma_1} \left| \int_{t_n}^{t_{n+1}} \int_{t_n}^s dW_{r_1}(s_1)dW_r(s) \right| \\
& \quad + \sum_{r=1}^m Ch^{1-\gamma_3}|W_r(t_{n+1}) - W_r(t_n)| + \sum_{r=1}^m Ch^{-\gamma_3} \left| \int_{t_n}^{t_{n+1}} \int_{t_n}^s dW_r(s_1)ds \right| + Ch^{2-\gamma_4} \\
& \leq |Y_0| + (n+1)Ch^{1-\gamma_1} + \sum_{k=0}^n \sum_{r=1}^m Ch^{-\gamma_2}|W_r(t_{k+1}) - W_r(t_k)| \\
& \quad + \sum_{k=0}^n \sum_{r=1}^m Ch^{-\gamma_3} \left| \int_{t_k}^{t_{k+1}} \int_{t_k}^s dW_r(s_1)ds \right| + \sum_{k=0}^n \sum_{r=1}^m \sum_{r_1=1}^m Ch^{-\gamma_1} \left| \int_{t_k}^{t_{k+1}} \int_{t_k}^s dW_{r_1}(s_1)dW_r(s) \right| \\
& \quad + \sum_{k=0}^n \sum_{r=1}^m Ch^{1-\gamma_3}|W_r(t_{k+1}) - W_r(t_k)| + (n+1)Ch^{2-\gamma_4}. \tag{5.6}
\end{aligned}$$

Similar to arguments in the proof of (D.7)-(D.9), using the Hölder inequality with $\frac{1}{p'} + \frac{1}{q'} = 1$ for $q' = \frac{\bar{p}}{(p\gamma_1+1)G_1} \vee \frac{\bar{p}}{(1+(\frac{1}{2}+\gamma_2)p)G_1} \vee \frac{\bar{p}}{(1-\frac{1}{2}p+p\gamma_3)G_1} \vee \frac{\bar{p}}{(1-p+p\gamma_4)G_1} > 1$ due to $p \leq \frac{\bar{p}-G_1}{1+\gamma_1 G_1} \wedge \frac{\bar{p}-G_1}{1+(\frac{1}{2}+\gamma_2)G_1} \wedge \frac{\bar{p}-G_1}{1+(\gamma_3-\frac{1}{2})G_1} \wedge \frac{\bar{p}-G_1}{1+(\gamma_4-1)G_1}$ and the Chebyshev inequality gives

$$\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}^c}|Y_n|^{\bar{p}}] \leq (\mathbb{E}[|Y_n|^{pp'}])^{\frac{1}{p'}} \sum_{i=0}^n \frac{(\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),i-1}}|Y_i|^{\bar{p}}])^{\frac{1}{q'}}}{\mathcal{R}(h)^{\bar{p}/q'}}.$$

Using the Hölder inequality, (5.6) and the elementary inequality implies

$$(\mathbb{E}[|Y_n|^{pp'}])^{\frac{1}{p'}} \leq C(|Y_0|^{\bar{p}} + n^{\bar{p}}h^{\bar{p}(1-\gamma_1)} + n^{\bar{p}}h^{\bar{p}(\frac{1}{2}-\gamma_2)} + n^{\bar{p}}h^{\bar{p}(\frac{3}{2}-\gamma_3)} + n^{\bar{p}}h^{\bar{p}(2-\gamma_4)})^{\frac{p}{p'}}$$

$$\leq Ch^{-p\gamma_1} + Ch^{-\frac{p}{2}-p\gamma_2} + Ch^{\frac{p}{2}-p\gamma_3} + Ch^{p-p\gamma_4} + C(1 + |X_0|^{\bar{p}})^{\frac{p}{\bar{p}}}. \quad (5.7)$$

Notice that arguments similar to those in the proof of Lemma 3.2 yield that

$$\begin{aligned} \mathbb{E}[|Y_n|^p] &= \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^p] + \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}^c}|Y_n|^p] \\ &\leq (\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^{\bar{p}}])^{\frac{p}{\bar{p}}} + \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}^c}|Y_n|^p] \\ &\leq C(1 + |X_0|^p)^\beta \end{aligned} \quad (5.8)$$

for all integer $p \in [1, \frac{\bar{p}-G_1}{1+\gamma_1 G_1} \wedge \frac{\bar{p}-G_1}{1+(\frac{1}{2}+\gamma_2)G_1} \wedge \frac{\bar{p}-G_1}{1+(\gamma_3-\frac{1}{2})G_1} \wedge \frac{\bar{p}-G_1}{1+(\gamma_4-1)G_1}]$. Here $\beta = 1 + \frac{\bar{p}}{pq'} = 1 + \frac{(p\gamma_1+1)G_1}{p} \wedge \frac{(1+\frac{1}{2}p+p\gamma_2)G_1}{p} \wedge \frac{(1-\frac{1}{2}p+p\gamma_3)G_1}{p} \wedge \frac{(1-p+p\gamma_4)G_1}{p}$. Then, by Jensen's inequality, (3.17) holds for non-integer p as well.

Except for the second term I_1 on the right-hand side of (5.1), the proof of bounded moments for the scheme (3.2) under Assumption (H2') is identical to that under Assumption (H2). Applying the elementary inequality and Assumption (H2'), we get

$$\begin{aligned} I_1 &\leq Ch + Ch\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^{\bar{p}}] + Ch^2\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^{4r+\bar{p}}] \\ &\quad + Ch^3\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^{4r+2\rho+\bar{p}-2}] + Ch^4\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}}|Y_n|^{8r+\bar{p}}]. \end{aligned}$$

Similar to the proof above, we choose $\mathcal{R} = h^{-1/G_1}$ and obtain a similar conclusion. \square

5.2 Weak convergence of second-order numerical schemes

To verify the one-step weak convergence rates (2.15)-(2.17) in Theorem 2.4, we will examine the one-step approximation of the numerical scheme (3.2), which is expressed as follows:

$$\begin{aligned} Y(t, x; t+h) &= x + \mathcal{T}_1(f(x), h)h + \sum_{r=1}^m \int_t^{t+h} \mathcal{T}_2(g^r(x), h) dW_r(s) \\ &\quad + \sum_{r=1}^m \sum_{r_1=1}^m \int_t^{t+h} \int_t^s \mathcal{T}_3(\Lambda_{r_1} g^r(x), h) dW_{r_1}(s_1) dW_r(s) + \sum_{r=1}^m \int_t^{t+h} \mathcal{T}_4(\mathcal{L}g^r(x), h)h dW_r(s) \\ &\quad + \sum_{r=1}^m \int_t^{t+h} \int_t^s \mathcal{T}_5((\Lambda_r f(x) - \mathcal{L}g^r(x)), h) dW_r(s_1) ds + \int_t^{t+h} \mathcal{T}_6(\mathcal{L}f(x), h) \frac{h}{2} ds. \end{aligned} \quad (5.9)$$

In the context of analyzing the weak error of the second-order numerical schemes, we adopt Assumption 3.7 to characterize the difference between f, g and $\mathcal{T}_1 - \mathcal{T}_6$. Based on this assumption, we can derive accurate estimates of the strong errors of the one-step approximations, which are essential for obtaining the desired estimates of the weak errors.

Lemma 5.1. Let Assumptions (A1), (A2) and Assumption 3.7 hold, then for any $p \geq 1$ it holds that

$$\|\delta_{Y_{MT},x} - \delta_{Y,x}\|_{L^{2p}(\Omega;\mathbb{R}^d)} \leq C(1 + |x|^{r_1})h^{q_0+\frac{1}{2}}, \quad (5.10)$$

$$\|\delta_{Y,x}\|_{L^{2p}(\Omega;\mathbb{R}^d)} \leq C(1 + |x|^{r_1})h^{\frac{1}{2}}, \quad (5.11)$$

$$\|\delta_{Y_{MT},x}\|_{L^{2p}(\Omega;\mathbb{R}^d)} \leq C(1 + |x|^{4r+1})h^{\frac{1}{2}}, \quad (5.12)$$

where $r_1 = (4r+1)\eta_{q_0}$, r is from (2.10) and η_{q_0} is from Assumption 3.7.

Proof. Using the Minkowski inequality, moment inequality [16, Theorem 7.1] and Assumption 3.7 gives

$$\begin{aligned} \|\delta_{Y_{MT},x} - \delta_{Y,x}\|_{L^{2p}(\Omega;\mathbb{R}^d)} &\leq Ch\|f(x) - \mathcal{T}_1(f(x), h)\|_{L^{2p}(\Omega;\mathbb{R}^d)} + Ch^{\frac{1}{2}}\|g^r(x) - \mathcal{T}_2(g^r(x), h)\|_{L^{2p}(\Omega;\mathbb{R}^d)} \\ &\quad + Ch\|\Lambda_{r_1} g^r(x) - \mathcal{T}_3(\Lambda_{r_1} g^r(x), h)\|_{L^{2p}(\Omega;\mathbb{R}^d)} \\ &\quad + Ch^{\frac{3}{2}}\|\mathcal{L}g^r(x) - \mathcal{T}_4(\mathcal{L}g^r(x), h)\|_{L^{2p}(\Omega;\mathbb{R}^d)} + Ch^2\|\mathcal{L}f(x) - \mathcal{T}_6(\mathcal{L}f(x), h)\|_{L^{2p}(\Omega;\mathbb{R}^d)} \end{aligned}$$

$$\begin{aligned}
& +Ch^{\frac{3}{2}}\|(\Lambda_r f(x) - \mathcal{L}g^r(x)) - \mathcal{T}_5((\Lambda_r f(x) - \mathcal{L}g^r(x)), h)\|_{L^{2p}(\Omega, \mathbb{R}^d)} \\
\leq & C(1 + |x|^{r_1})h^{q_0 + \frac{1}{2}},
\end{aligned} \tag{5.13}$$

which implies the first assertion. Similar arguments lead to

$$\begin{aligned}
\|\delta_{Y,x}\|_{L^{2p}(\Omega; \mathbb{R}^d)} & \leq Ch\|\mathcal{T}_1(f(x), h)\|_{L^{2p}(\Omega, \mathbb{R}^d)} + Ch^{\frac{1}{2}}\|\mathcal{T}_2(g^r(x), h)\|_{L^{2p}(\Omega, \mathbb{R}^d)} \\
& + Ch\|\mathcal{T}_3(\Lambda_{r_1}g^r(x), h)\|_{L^{2p}(\Omega, \mathbb{R}^d)} \\
& + Ch^{\frac{3}{2}}\|\mathcal{T}_4(\mathcal{L}g^r(x), h)\|_{L^{2p}(\Omega, \mathbb{R}^d)} + Ch^2\|\mathcal{T}_6(\mathcal{L}f(x), h)\|_{L^{2p}(\Omega, \mathbb{R}^d)} \\
& + Ch^{\frac{3}{2}}\|\mathcal{T}_5((\Lambda_r f(x) - \mathcal{L}g^r(x)), h)\|_{L^{2p}(\Omega, \mathbb{R}^d)} \\
& \leq C(1 + |x|^{r_1})h^{\frac{1}{2}}.
\end{aligned} \tag{5.14}$$

The estimate of $\|\delta_{Y_{MT},x}\|_{L^{2p}(\Omega; \mathbb{R}^d)}$ can be done similarly. \square

Lemma 5.2 (One-step weak error estimates). Under the same conditions of Lemma 5.1, the one-step approximation (5.9) of the numerical scheme (3.2) satisfies

$$\begin{aligned}
\left| \mathbb{E} \left[\prod_{j=1}^s (\delta_{X,x})^{i_j} - \prod_{j=1}^s (\delta_{Y,x})^{i_j} \right] \right| & \leq C(1 + |x|^{r_2} + |x|^{(2q+1)r_1})h^{3 \wedge (q_0+1)}, \quad s = 1, \dots, 2q+1, \\
\left\| \prod_{j=1}^{2q+2} (\delta_{Y,x})^{i_j} \right\|_{L^2(\Omega; \mathbb{R})} & \leq C(1 + |x|^{(2q+2)r_1})h^{q+1}, \\
\left\| \prod_{j=1}^{2q+2} (\delta_{X,x})^{i_j} \right\|_{L^2(\Omega; \mathbb{R})} & \leq C(1 + |x|^{(2q+2)(2r+1)})h^{q+1},
\end{aligned}$$

Here r_1 comes from Lemma 5.1 and $r_2 = (2q+1)(6r+1)$.

Proof. Thanks to the triangle inequality, we have

$$\begin{aligned}
& \left| \mathbb{E} \left[\prod_{j=1}^s (\delta_{X,x})^{i_j} - \prod_{j=1}^s (\delta_{Y,x})^{i_j} \right] \right| \\
\leq & \left| \mathbb{E} \left[\prod_{j=1}^s (\delta_{X,x})^{i_j} - \prod_{j=1}^s (\delta_{Y_{MT},x})^{i_j} \right] \right| + \left| \mathbb{E} \left[\prod_{j=1}^s (\delta_{Y_{MT},x})^{i_j} - \prod_{j=1}^s (\delta_{Y,x})^{i_j} \right] \right| \\
=: & J_1 + J_2, \quad s = 1, \dots, 2q+1.
\end{aligned} \tag{5.15}$$

Building on the proof presented in [23, Lemma 1.4, Chapter 2], we establish the inequality $J_1 \leq C(1 + |x|^{r_2})h^3$. Now, our focus shifts to the estimation of J_2 . Specifically, when $s = 1$, we can estimate the second term J_2 in the following manner:

$$\begin{aligned}
|\mathbb{E}[(\delta_{Y_{MT},x})^{i_1} - (\delta_{Y,x})^{i_1}]| & \leq |\mathbb{E}[(f(x) - \mathcal{T}_1(f(x), h))h + \frac{h^2}{2}(\mathcal{L}f(x) - \mathcal{T}_6(\mathcal{L}f(x), h)) \\
& + \sum_{r=1}^m \sum_{r_1=1}^m \int_t^{t+h} \int_t^s (\Lambda_{r_1}g^r(x) - \mathcal{T}_3(\Lambda_{r_1}g^r(x), h)) dW_{r_1}(s_1)dW_r(s)]| \\
& \leq C(1 + |x|^{r_1})h^{q_0+1}.
\end{aligned} \tag{5.16}$$

With the Hölder inequality and Lemma 5.1 available, one can readily obtain the result for $s = 2$, yielding

$$\begin{aligned}
& \left| \mathbb{E} \left[\prod_{j=1}^2 (\delta_{Y_{MT},x})^{i_j} - \prod_{j=1}^2 (\delta_{Y,x})^{i_j} \right] \right| \\
\leq & \|\delta_{Y_{MT},x} - \delta_{Y,x}\|_{L^2(\Omega; \mathbb{R}^d)} \|\delta_{Y_{MT},x}\|_{L^2(\Omega; \mathbb{R}^d)} + \|\delta_{Y,x}\|_{L^2(\Omega; \mathbb{R}^d)} \|\delta_{Y_{MT},x} - \delta_{Y,x}\|_{L^2(\Omega; \mathbb{R}^d)}
\end{aligned}$$

$$\leq C(1 + |x|^{2r_1})h^{q_0+1}. \quad (5.17)$$

Utilizing a technique analogous to the one demonstrated in Equation (5.17), we obtain the following result

$$\left| \mathbb{E} \left[\prod_{j=1}^3 (\delta_{Y_{MT}, x})^{i_j} - \prod_{j=1}^3 (\delta_{Y, x})^{i_j} \right] \right| \leq C(1 + |x|^{3r_1})h^{q_0 + \frac{3}{2}}. \quad (5.18)$$

Similar to (5.18), we have

$$J_2 \leq C(1 + |x|^{(2q+1)r_1})h^{q_0+1}, \quad s = 1, \dots, 2q + 1.$$

Using the Hölder inequality and (5.11) yields

$$\left\| \prod_{j=1}^{2q+2} (\delta_{Y, x})^{i_j} \right\|_{L^2(\Omega; \mathbb{R})} \leq \prod_{j=1}^{2q+2} \left\| (\delta_{Y, x})^{i_j} \right\|_{L^{4q+4}(\Omega; \mathbb{R})} \leq C(1 + |x|^{(2q+2)r_1})h^{q+1}. \quad \square$$

By Lemmas 3.6 and 5.2 and Theorem 2.4, we can readily derive the weak convergence order of the numerical scheme (3.2) in Theorem 3.8.

6 Discussion

We have discussed the weak convergence orders of modified Euler and Milstein-Talay schemes. When solutions and numerical solutions have finite moments, it is crucial to find the highest moment of numerical solutions to determine the convergence. We show how the modification maps in the modified schemes determine the highest order of the moments. We investigate the effects of limited moments on weak convergence orders theoretically and numerically. However, the theoretical orders of weak convergence seem to be lower than the observed orders. We conjecture that the highest order of moments may be underestimated in Theorems 3.3 and 3.8. Further exploration along this direction is needed.

References

- [1] W.-J. Beyn, E. Isaak, and R. Kruse. Stochastic C-stability and B-consistency of explicit and implicit Euler-type schemes. *Journal of Scientific Computing*, 67:955–987, 2016.
- [2] W.-J. Beyn, E. Isaak, and R. Kruse. Stochastic c-stability and b-consistency of explicit and implicit milstein-type schemes. *Journal of Scientific Computing*, 70:1042–1077, 2017.
- [3] E. Buckwar, A. Samson, M. Tamborrino, and I. Tubikanec. A splitting method for SDEs with locally Lipschitz drift: Illustration on the FitzHugh-Nagumo model. *Applied Numerical Mathematics*, 179:191–220, 2022.
- [4] S. Cerrai. *Second order PDE's in finite and infinite dimension: A probabilistic approach*, volume 1762 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2001.
- [5] Z. Chen, S. Gan, and X. Wang. Mean-square approximations of lévy noise driven sdes with super-linearly growing diffusion and jump coefficients. *Discrete & Continuous Dynamical Systems-Series B*, 24(8), 2019.
- [6] D. J. Higham, X. Mao, and A. M. Stuart. Strong convergence of Euler-type methods for non-linear stochastic differential equations. *SIAM Journal on Numerical Analysis*, 40:1041–1063, 2002.
- [7] D. J. Higham, X. Mao, and C. Yuan. Almost sure and moment exponential stability in the numerical simulation of stochastic differential equations. *SIAM Journal on Numerical Analysis*, 45(2):592–609, 2007.
- [8] Y. Hu. Semi-implicit Euler-Maruyama scheme for stiff stochastic equations. In *Stochastic analysis and related topics, V (Silivri, 1994)*, volume 38 of *Progr. Probab.*, pages 183–202. Birkhäuser Boston, Boston, MA, 1996.
- [9] M. Hutzenthaler and A. Jentzen. On a perturbation theory and on strong convergence rates for stochastic ordinary and partial differential equations with nonglobally monotone coefficients. *Ann. Probab.*, 48(1):53–93, 2020.

- [10] M. Hutzenthaler, A. Jentzen, and P. E. Kloeden. Strong and weak divergence in finite time of Euler’s method for stochastic differential equations with non-globally Lipschitz continuous coefficients. *Proceedings of The Royal Society of London. Series A. Mathematical, Physical and Engineering Sciences*, 467(2130):1563–1576, 2011.
- [11] M. Hutzenthaler, A. Jentzen, and P. E. Kloeden. Strong convergence of an explicit numerical method for SDEs with non-globally Lipschitz continuous coefficients. *The Annals of Applied Probability*, 22(4):1611–1641, 2012.
- [12] M. Hutzenthaler, A. Jentzen, and X. Wang. Exponential integrability properties of numerical approximation processes for nonlinear stochastic differential equations. *Math. Comp.*, 87(311):1353–1413, 2018.
- [13] R. Khasminskii. *Stochastic stability of differential equations*, volume 66. Springer Science & Business Media, 2011.
- [14] P. E. Kloeden and E. Platen. *Numerical Solution of Stochastic Differential Equations*. Springer, 1992.
- [15] W. Liu and X. Mao. Strong convergence of the stopped Euler-Maruyama method for nonlinear stochastic differential equations. *Applied Mathematics and Computation*, 223:389–400, 2013.
- [16] X. Mao. *Stochastic differential equations and applications*. Horwood Publishing Limited, Chichester, second edition, 2008.
- [17] X. Mao. The truncated Euler- Maruyama method for stochastic differential equations. *Journal of Computational and Applied Mathematics*, 290:370–384, 2015.
- [18] X. Mao and L. Szpruch. Strong convergence rates for backward Euler-Maruyama method for non-linear dissipative-type stochastic differential equations with super-linear diffusion coefficients. *Stochastics*, 85(1):144–171, 2013.
- [19] M. Matsumoto and T. Nishimura. Mersenne twister: a 623-dimensionally equidistributed uniform pseudo-random number generator. *ACM Transactions on Modeling and Computer Simulation (TOMACS)*, 8(1):3–30, 1998.
- [20] J. C. Mattingly, A. M. Stuart, and D. J. Higham. Ergodicity for SDEs and approximations: locally Lipschitz vector fields and degenerate noise. *Stochastic processes and their applications*, 101(2):185–232, 2002.
- [21] G. N. Milstein. A method with second order accuracy for the integration of stochastic differential equations. *Theory of Probability & Its Applications*, 23(2):396–401, 1978.
- [22] G. N. Milstein. Weak approximation of solutions of systems of stochastic differential equations. *Theory of Probability & Its Applications*, 30(4):706–721, 1985.
- [23] G. N. Milstein and M. V. Tretyakov. *Stochastic numerics for mathematical physics*. Scientific Computation. Springer-Verlag, Berlin, 2004.
- [24] G. N. Milstein and M. V. Tretyakov. Numerical integration of stochastic differential equations with nonglobally Lipschitz coefficients. *SIAM Journal on Numerical Analysis*, 43(3):1139–1154, 2005.
- [25] S. Sabanis. A note on tamed Euler approximations. *Electron. Commun. Probab.*, 18:no. 47, 10, 2013.
- [26] S. Sabanis. Euler approximations with varying coefficients: the case of super-linearly growing diffusion coefficients. *The Annals of Applied Probability*, 26(4):2083–2105, 2016.
- [27] L. Szpruch. V-stable tamed euler schemes. *arXiv preprint arXiv:1310.0785*, 2013.
- [28] D. Talay. Efficient numerical schemes for the approximation of expectations of functionals of the solution of a SDE and applications. In *Filtering and control of random processes (Paris, 1983)*, volume 61 of *Lect. Notes Control Inf. Sci.*, pages 294–313. Springer, Berlin, 1984.
- [29] M. V. Tretyakov and Z. Zhang. A fundamental mean-square convergence theorem for SDEs with locally Lipschitz coefficients and its applications. *SIAM Journal on Numerical Analysis*, 51(6):3135–3162, 2013.

- [30] X. Wang and S. Gan. The tamed Milstein method for commutative stochastic differential equations with non-globally Lipschitz continuous coefficients. *Journal of Difference Equations and Applications*, 19(3):466–490, 2013.
- [31] X. Wang, Y. Zhao, and Z. Zhang. Weak error analysis for strong approximation schemes of SDEs with super-linear coefficients. *IMA Journal of Numerical Analysis*, page drad083, 2023.
- [32] Z. Zhang and H. Ma. Order-preserving strong schemes for SDEs with locally Lipschitz coefficients. *Applied Numerical Mathematics*, 112:1–16, 2017.

A Moment estimates for an SDE with random forcing

In this section, we establish general moment estimates for the SDE (2.2), which will be used in the analysis in Appendix B.

Let $\mathcal{H}_1, \mathcal{H}_2$ be random functions defined on $\Omega \times [0, T]$ and be adapted to the natural filtration generated by $W(t)$.

Assumption A.1. There exists a constant $K \geq 0$ and $q_1 \geq 1, q_2 \geq 2$, such that for all $t \in [0, T]$,

$$\mathbb{E}\left[\int_0^t |\mathcal{H}_1(\omega, s)|^{q_1} ds\right] \leq K, \quad \mathbb{E}\left[\int_0^t |\mathcal{H}_2(\omega, s)|^{q_2} ds\right] \leq K.$$

Lemma A.2. Under Assumption 2.1 with sufficiently large $p_0 \geq 2$ and Assumption A.1, the stochastic differential equation

$$Z(t) = Z(0) + \int_0^t (Df(X(s))Z(s) + \mathcal{H}_1(\omega, s)) ds + \int_0^t (Dg(X(s))Z(s) + \mathcal{H}_2(\omega, s)) dW(s), \quad (\text{A.1})$$

admits a unique strong solution $\{Z(t) : t \in [0, T]\}$ such that for any $p_2 \in [1, q_1 \wedge q_2 \wedge \frac{p_0+1}{2}]$,

$$\mathbb{E}[|Z(t)|^{p_2}] \leq C_{p_2} \mathbb{E}\left[(|Z(0)|^2)^{p_2} + \int_0^t |\mathcal{H}_1(\omega, s)|^{p_2} ds + \int_0^t |\mathcal{H}_2(\omega, s)|^{p_2} ds \right] \exp(C_{p_2} T), \quad t \in [0, T], \quad (\text{A.2})$$

where the constant $C > 0$ depends only on K, p_2 and T . Here q_1 and q_2 are from Assumption A.1.

The existence and uniqueness of the solution can be proved by applying a truncation argument and Assumption 2.1, see e.g., in [16, Chapter 2]. Applying Itô's formula and Assumption (A3) leads to the moment bound (A.2).

If $\mathcal{H}_1(\omega, t, \epsilon)$ and $\mathcal{H}_2(\omega, t, \epsilon)$ depend on ϵ and have limits in a certain sense when $\epsilon \rightarrow 0$, then we can show that the solution to the resulting equation, denoted by $\mathcal{S}_\epsilon(t)$, also has a limit.

Corollary A.3. Under the same conditions in Lemma A.2, if there exists $p_3 \geq 1$ and $\bar{p}_3 \geq 2$ such that

$$\lim_{\epsilon \rightarrow 0} \mathbb{E}\left[\int_0^t |\mathcal{H}_1(\omega, s, \epsilon)|^{p_3} ds\right] = 0, \quad \lim_{\epsilon \rightarrow 0} \mathbb{E}\left[\int_0^t |\mathcal{H}_2(\omega, s, \epsilon)|^{\bar{p}_3} ds\right] = 0, \quad (\text{A.3})$$

then it holds that

$$\lim_{\epsilon \rightarrow 0} \mathbb{E}[|\mathcal{S}_\epsilon(t)|^{\tilde{p}_3}] \leq C_{\tilde{p}_3} \lim_{\epsilon \rightarrow 0} \mathbb{E}[|\mathcal{S}_\epsilon(0)|^{\tilde{p}_3}], \quad \tilde{p}_3 \in [1, p_3 \wedge \bar{p}_3 \wedge \frac{p_0+1}{2}],$$

where $C_{\tilde{p}_3}$ is independent of ϵ but dependent on \tilde{p}_3 and T .

B Derivatives of the solutions in the initial condition

In this section, we focus on the SDEs that characterize the derivatives of X^{0, x_0} with respect to the initial condition x_0 . We adopt the notation used in [4]. For random functions, we refer to Page 166 of [13] and below for the definition of differentiability in the $L^p(\Omega)$ ($p \geq 1$) sense. For simplicity, we present the proofs for the derivatives up to order two.

Definition B.1 (L^p differentiable). Let $\Psi: \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}$ and $\phi_i: \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}$ be random functions and suppose for each $i \in \{1, 2, \dots, d\}$

$$\lim_{\tau \rightarrow 0} \mathbb{E} \left[\left| \frac{1}{\tau} [\Psi(x + \tau e_i) - \Psi(x)] - \phi_i(x) \right|^p \right] = 0, \quad p \geq 1, \quad (\text{B.1})$$

where $e_i \in \mathbb{R}^d$ is a unit vector in \mathbb{R}^d , with the i -th element being 1. Then Ψ is called to be differentiable, with $\phi = (\phi_1, \phi_2, \dots, \phi_d)$ being the derivative (in the $L^p(\Omega)$ sense) of Ψ and we write $\partial_{(i)} \Psi = \phi_i$.

This definition can be extended to the case of vector-valued functions by considering their components separately.

We now apply conclusions in Appendix A to analyze the derivatives of the solution to the SDE (2.2) concerning the initial conditions in the sense of $L^p(\Omega)$ differentiability. To do so, we first examine the limit behavior of solutions for the SDEs with perturbed initial conditions.

Proposition B.2. Let Assumption 2.1 and the inequality (A3') hold. Then the solution $X^{0,x_0}(t)$ to (2.2) satisfies, for any $\epsilon > 0$,

$$\mathbb{E} [|X^{0,x_0+\epsilon e_i}(t) - X^{0,x_0}(t)|^{p_4}] \leq |\epsilon|^{p_4} \exp(C_{p_4} T), \quad p_4 \in [1, p'_0], \quad (\text{B.2})$$

where $e_i \in \mathbb{R}^d$ is defined in Definition B.1 and C_{p_4} is independent of ϵ but dependent on p_4 . For brevity, we denote

$$\chi_{x_0, \epsilon}(s) := X^{0,x_0+\epsilon e_i}(s) - X^{0,x_0}(s), \quad \Delta f^{x_0+\epsilon e_i, x_0}(s) := f(X^{0,x_0+\epsilon e_i}(s)) - f(X^{0,x_0}(s)), \quad (\text{B.3})$$

$$\Delta g^{x_0+\epsilon e_i, x_0}(s) := g(X^{0,x_0+\epsilon e_i}(s)) - g(X^{0,x_0}(s)). \quad (\text{B.4})$$

Proof. By the definition of $X^{0,x_0+\epsilon e_i}(t)$,

$$X^{0,x_0+\epsilon e_i}(t) = x_0 + \epsilon e_i + \int_0^t f(X^{0,x_0+\epsilon e_i}(s)) ds + \int_0^t g(X^{0,x_0+\epsilon e_i}(s)) dW(s), \quad 0 < t \leq T.$$

Using the Itô formula on the functional $|x|^{p_4}$ for $x \in \mathbb{R}^d$, $p_4 \geq 2$ and the Schwarz inequality, we obtain that

$$\begin{aligned} |\chi_{x_0, \epsilon}(t)|^{p_4} &\leq |\epsilon e_i|^{p_4} + p_4 \int_0^t |\chi_{x_0, \epsilon}(s)|^{p_4-2} \langle \chi_{x_0, \epsilon}(s), \Delta f^{x_0+\epsilon e_i, x_0}(s) \rangle ds \\ &\quad + \frac{p_4(p_4-1)}{2} \int_0^t |\chi_{x_0, \epsilon}(s)|^{p_4-2} |\Delta g^{x_0+\epsilon e_i, x_0}(s)|^2 ds \\ &\quad + p_4 \int_0^t |\chi_{x_0, \epsilon}(s)|^{p_4-2} \langle \chi_{x_0, \epsilon}(s), \Delta g^{x_0+\epsilon e_i, x_0}(s) dW(s) \rangle. \end{aligned}$$

Applying the inequality (A3') and the properties of Itô integrals implies

$$\mathbb{E} [|\chi_{x_0, \epsilon}(t)|^{p_4}] \leq |\epsilon|^{p_4} + C_{p_4} \int_0^t \mathbb{E} [|\chi_{x_0, \epsilon}(s)|^{p_4}] ds, \quad p_4 \in [2, p'_0], \quad (\text{B.5})$$

where C_{p_4} is independent of ϵ but dependent on p_4 . The Gronwall inequality shows that for all $t \in [0, T]$, $\mathbb{E} [|X^{0,x_0+\epsilon e_i}(t) - X^{0,x_0}(t)|^{p_4}] \leq |\epsilon|^{p_4} \exp(C_{p_4} T)$, $p_4 \in [2, p'_0]$. For the case $p_4 \in [1, 2)$, applying the Hölder inequality immediately suggests (B.2). Thus we complete the proof. \square

Corollary B.3. Under the same assumptions of Proposition B.2 with $p_0 \geq 2r + 1$ (r is defined in (2.3)), for each $i \in \{1, \dots, d\}$, $\xi_i(t) := \frac{\partial X^{0,x_0}(t)}{\partial x^{(i)}} \in \mathbb{R}^d$ exists and satisfies the following stochastic differential equation

$$\begin{cases} d\xi_i(t) = Df(X(t))\xi_i(t) dt + Dg(X(t))\xi_i(t) dW(t), & t \in (0, T], \\ \xi_i(0) = e_i, \end{cases} \quad (\text{B.6})$$

where $e_i \in \mathbb{R}^d$ comes from Definition B.1. Furthermore, there exists a constant $C > 0$, such that

$$\mathbb{E} [|\xi_i(t)|^{p_5}] \leq C \exp(C_{p_5} t), \quad 1 \leq p_5 \leq p_0, \quad \text{for all } t \in [0, T], \quad (\text{B.7})$$

$$\text{and } \lim_{\epsilon \rightarrow 0} \mathbb{E} \left[\left| \frac{\chi_{x_0, \epsilon}(t)}{\epsilon} - \xi_i(t) \right|^{p_6} \right] = 0, \quad 1 \leq p_6 \leq \frac{p_0}{(2r \vee 0) + 1}, \quad \text{for all } t \in [0, T], \quad (\text{B.8})$$

where $\chi_{x_0, \epsilon}(t) := X^{0,x_0+\epsilon e_i}(t) - X^{0,x_0}(t)$.

Proof. Following the lines similar to those in Lemma A.2, (B.7) can be derived by observing that $\xi_i(t)$ satisfies the equation (A.2) with $\mathcal{H}_1 = \mathcal{H}_2 = 0$, $Z(0) = e_i$. Below, we apply Corollary A.3 to derive (B.8). First of all, $\frac{\chi_{x_0, \epsilon}(t)}{\epsilon} - \xi_i(t)$ satisfies the equation (A.1) with zero initial condition and \mathcal{H}_1 and \mathcal{H}_2 defined by

$$\begin{aligned}\mathcal{H}_1(\omega, s, \epsilon) &= \frac{f(X^{0, x_0 + \epsilon e_i}(s)) - f(X^{0, x_0}(s))}{\epsilon} - Df(X^{0, x_0}(s)) \frac{\chi_{x_0, \epsilon}(s)}{\epsilon}, \\ \mathcal{H}_2(s)(\omega, s, \epsilon) &= \frac{g(X^{0, x_0 + \epsilon e_i}(s)) - g(X^{0, x_0}(s))}{\epsilon} - Dg(X^{0, x_0}(s)) \frac{\chi_{x_0, \epsilon}(s)}{\epsilon}.\end{aligned}$$

It only requires to verify that the corresponding \mathcal{H}_1 and \mathcal{H}_2 satisfy (A.3). In fact, using the Taylor expansion, the Hölder inequality with $\frac{1}{\theta_1} + \frac{1}{\theta_2} = 1$, for $\theta_2 = \frac{p_0}{p_6}$ and Proposition B.2, we have

$$\begin{aligned}\mathbb{E}\left[\int_0^t |\mathcal{H}_1|^{p_6} ds\right] &\leq \int_0^t \left(\mathbb{E}\left[\left|\int_0^1 Df(X^{0, x_0}(s) + \lambda(X^{0, x_0 + \epsilon e_i}(s) - X^{0, x_0}(s))) - Df(X^{0, x_0}(s)) d\lambda\right|^{p_6 \theta_1}\right]\right)^{1/\theta_1} \\ &\quad \left(\mathbb{E}\left[\left|\frac{\chi_{x_0, \epsilon}(s)}{\epsilon}\right|^{p_6 \theta_2}\right]\right)^{1/\theta_2} ds \\ &\leq C \int_0^t \left(\mathbb{E}\left[\left|\int_0^1 Df(X^{0, x_0}(s) + \lambda(X^{0, x_0 + \epsilon e_i}(s) - X^{0, x_0}(s))) - Df(X^{0, x_0}(s)) d\lambda\right|^{p_6 \theta_1}\right]\right)^{1/\theta_1} ds.\end{aligned}$$

By the Assumption (A1), (2.8) and the dominated convergence theorem, we get

$$\lim_{\epsilon \rightarrow 0} \mathbb{E}\left[\int_0^t |\mathcal{H}_1|^{p_6} ds\right] = 0, \quad 1 \leq p_6 \leq \frac{p_0}{(2r \vee 0) + 1}, \quad \text{for all } t \in [0, T]. \quad (\text{B.9})$$

Following a similar way as (B.9) and Assumption (A2), we have

$$\lim_{\epsilon \rightarrow 0} \mathbb{E}\left[\int_0^t |\mathcal{H}_2|^{p_6 \vee 2} ds\right] = 0, \quad 1 \leq p_6 \leq \frac{p_0}{((\rho - 1) \vee 0) + 1}, \quad \text{for all } t \in [0, T].$$

The proof is thus complete. \square

We now extend the above result to higher-order derivatives.

Proposition B.4. Under the same assumptions of Proposition B.2 with $p_0 \geq 2r + 1$, the solution $\xi_i(t)$ of (B.6) satisfy

$$\lim_{\epsilon \rightarrow 0} \mathbb{E}\left[|\xi_i^{0, x_0 + \epsilon_1 e_j}(t) - \xi_i^{0, x_0}(t)|^{p_7}\right] = 0, \quad p_7 \in \left[1, \frac{p_0}{(2r \vee 0) + 1}\right], \quad (\text{B.10})$$

where $e_j \in \mathbb{R}^d$ comes from Definition B.1.

Proof. We apply Corollary A.3 to derive (B.10), which requires us to verify that the corresponding \mathcal{H}_1 and \mathcal{H}_2 satisfy (A.3). Here

$$\begin{aligned}\mathcal{H}_1 &= Df(X^{0, x_0 + \epsilon_1 e_j}(s)) \xi_i^{0, x_0 + \epsilon_1 e_j}(s) - Df(X^{0, x_0}(s)) \xi_i^{0, x_0 + \epsilon_1 e_j}(s), \\ \mathcal{H}_2 &= Dg(X^{0, x_0 + \epsilon_1 e_j}(s)) \xi_i^{0, x_0 + \epsilon_1 e_j}(s) - Dg(X^{0, x_0}(s)) \xi_i^{0, x_0 + \epsilon_1 e_j}(s), \quad \mathcal{S}_\epsilon(0) = 0.\end{aligned}$$

Using the Hölder inequality with $\frac{1}{\theta_1} + \frac{1}{\theta_2} = 1$, for $\theta_2 = \frac{p_0}{p_7}$ and (B.7), we have

$$\begin{aligned}\mathbb{E}\left[\int_0^t |\mathcal{H}_1|^{p_7} ds\right] &\leq \int_0^t \left(\mathbb{E}\left[\left|Df(X^{0, x_0 + \epsilon_1 e_j}(s)) - Df(X^{0, x_0}(s))\right|^{p_7 \theta_1}\right]\right)^{1/\theta_1} \left(\mathbb{E}\left[\left|\xi_i^{0, x_0 + \epsilon_1 e_j}(s)\right|^{p_7 \theta_2}\right]\right)^{1/\theta_2} ds \\ &\leq C \int_0^t \left(\mathbb{E}\left[\left|Df(X^{0, x_0 + \epsilon_1 e_j}(s)) - Df(X^{0, x_0}(s))\right|^{p_7 \theta_1}\right]\right)^{1/\theta_1} ds.\end{aligned}$$

Therefore, from the Assumption (A1), (2.8) and dominated convergence theorem, we get

$$\lim_{\epsilon \rightarrow 0} \mathbb{E}\left[\int_0^t |\mathcal{H}_1|^{p_7} ds\right] = 0, \quad 1 \leq p_7 \leq \frac{p_0}{(2r \vee 0) + 1}, \quad \text{for all } t \in [0, T].$$

Similarly, one can show that

$$\lim_{\epsilon \rightarrow 0} \mathbb{E} \left[\int_0^t |\mathcal{H}_2|^{p_7 \vee 2} ds \right] = 0, \quad 1 \leq p_7 \leq \frac{p_0}{((\rho - 1) \vee 0) + 1}, \quad \text{for all } t \in [0, T].$$

Using the Corollary B.3, we complete the proof. \square

Corollary B.5. Under the same assumptions of Proposition B.2 with $p_0 \geq 4r + 4$, for each $i, j \in \{1, \dots, d\}$, $\zeta_{i,j}(t) := \frac{\partial^2 X^{0,x_0}(t)}{\partial x_0^{(i)} \partial x_0^{(j)}}$ exists and satisfies the following equation

$$\begin{cases} d\zeta_{i,j}(t) = Df(X(t))\zeta_{i,j}(t) dt + Dg(X(t))\zeta_{i,j}(t) dW(t) + d\mathfrak{S}^{i,j}(t, x_0), & t \in (0, T], \\ \zeta_{i,j}(0) = 0, \end{cases} \quad (\text{B.11})$$

where $\mathfrak{S}^{i,j}(t, x_0)$ is the process defined by

$$\mathfrak{S}^{i,j}(t, x_0) = \int_0^t D^2 f(X(s))(\xi_i(s), \xi_j(s)) ds + \int_0^t D^2 g(X(s))(\xi_i(s), \xi_j(s)) dW(s).$$

Furthermore, there exists a constant $C > 0$, such that for all $t \in [0, T]$,

$$\mathbb{E}[|\zeta_{i,j}(t)|^{p_8}] \leq C \exp(Cp_8 T) (1 + |x_0|^{((2r-1) \vee 0)p_8}), \quad 1 \leq p_8 \leq \frac{p_0}{((2r-1) \vee 0) + 2}, \quad (\text{B.12})$$

$$\text{and } \lim_{\epsilon_1 \rightarrow 0} \mathbb{E} \left[\left| \frac{\xi_i^{0, x_0 + \epsilon_1 e_j}(t) - \xi_i^{0, x_0}(t)}{\epsilon_1} - \zeta_{i,j}(t) \right|^{p_9} \right] = 0, \quad 1 \leq p_9 \leq \frac{p_0}{((2r-1) \vee 0) + (2r \vee 0) + 2}. \quad (\text{B.13})$$

Proof. We apply Lemma A.2 to derive (B.12), which requires us to verify that the corresponding \mathcal{H}_1 and \mathcal{H}_2 satisfy Assumption A.1. Here

$$\mathcal{H}_1 = D^2 f(X(s))(\xi_i(s), \xi_j(s)), \quad \mathcal{H}_2 = D^2 g(X(s))(\xi_i(s), \xi_j(s)), \quad Z(0) = 0.$$

It follows from the Hölder inequality, Assumption 2.1, (2.8) and (B.7) that for $p_0 \geq (2r + 1) \vee 2$

$$\mathbb{E} \left[\int_0^t |\mathcal{H}_1(\omega, s)|^{p_8} ds \right] \leq C(1 + |x_0|^{((2r-1) \vee 0)p_8}), \quad 1 \leq p_8 \leq \frac{p_0}{((2r-1) \vee 0) + 2}. \quad (\text{B.14})$$

By similar arguments in the proof of (B.14), we obtain

$$\mathbb{E} \left[\int_0^t |\mathcal{H}_2(\omega, s)|^{p_8 \vee 2} ds \right] \leq C(1 + |x_0|^{((\rho-2) \vee 0)p_8}), \quad 1 \leq p_8 \leq \frac{p_0}{((2r-1) \vee 0) + 2}.$$

Next, based on the Corollary A.3, we are in the position to prove (B.13). Here

$$\begin{aligned} \mathcal{H}_1 &= \frac{(Df(X^{0, x_0 + \epsilon_1 e_j}(s)) - Df(X^{0, x_0}(s)))\xi_i^{0, x_0 + \epsilon_1 e_j}(s)}{\epsilon_1} - D^2 f(X^{0, x_0}(s))(\xi_i(s), \xi_j(s)), \\ \mathcal{H}_2 &= \frac{(Dg(X^{0, x_0 + \epsilon_1 e_j}(s)) - Dg(X^{0, x_0}(s)))\xi_i^{0, x_0 + \epsilon_1 e_j}(s)}{\epsilon_1} - D^2 g(X^{0, x_0}(s))(\xi_i(s), \xi_j(s)), \quad \mathcal{S}_\epsilon(0) = 0. \end{aligned}$$

Using the Taylor expansion and the elementary inequality, we have

$$\begin{aligned} \mathbb{E} \left[\int_0^t |\mathcal{H}_1|^{p_9} ds \right] &\leq C \mathbb{E} \left[\int_0^t \left| \int_0^1 D^2 f(\lambda(s)) d\lambda \left(\frac{X^{0, x_0 + \epsilon_1 e_j}(s) - X^{0, x_0}(s)}{\epsilon_1} - \xi_i^{0, x_0 + \epsilon_1 e_j}(s) \right) \xi_i^{0, x_0 + \epsilon_1 e_j}(s) \right|^{p_9} ds \right] \\ &\quad + C \mathbb{E} \left[\int_0^t \left| \int_0^1 D^2 f(\lambda(s)) d\lambda (\xi_i^{0, x_0 + \epsilon_1 e_j}(s) - \xi_i^{0, x_0}(s)) \xi_i^{0, x_0}(s) \right|^{p_9} ds \right] \\ &\quad + C \mathbb{E} \left[\int_0^t |(D^2 f(\lambda(s)) - D^2 f(X^{0, x_0}(s))) (\xi_j^{0, x_0}(s), \xi_i^{0, x_0}(s))|^{p_9} ds \right] \end{aligned}$$

$$=: I_1 + I_2 + I_3,$$

where $\lambda(s) = \lambda X^{0,x_0}(s) + (1 - \lambda)X^{0,x_0+\epsilon_1 e_j}(s)$. By the Hölder inequality, Assumption 2.1, Corollary B.3 and the dominated convergence theorem, we have that for $p_0 \geq 4r + 4$

$$\lim_{\epsilon_1 \rightarrow 0} I_1 = 0, \quad 1 \leq p_9 \leq \frac{p_0}{((2r-1) \vee 0) + (2r \vee 0) + 2}, \quad \text{for all } t \in [0, T]. \quad (\text{B.15})$$

Similar arguments lead to $\lim_{\epsilon_1 \rightarrow 0} I_2 = 0$ and $\lim_{\epsilon_1 \rightarrow 0} I_3 = 0$, which indicates that $\lim_{\epsilon_1 \rightarrow 0} \mathbb{E}[\int_0^t |\mathcal{H}_1|^{p_9} ds] = 0$. Similarly, we have

$$\mathbb{E}[\int_0^t |\mathcal{H}_2|^{p_9 \vee 2} ds] = 0, \quad 1 \leq p_9 \leq \frac{p_0}{((\rho-2) \vee 0) + (2r \vee 0) + 2}, \quad \text{for all } t \in [0, T],$$

which, together with Corollary B.3, completes the proof. \square

Proof of Lemma 2.5. The proof is conducted by induction with the $j = 1$ case in Corollary B.3 as the base step, and $j = 2$ case in Corollary B.5 as the first induction step. We omit the details here. \square

Remark B.6. As discussed in Remark 2.2, notice we prove Lemma 2.5 under the relax condition compared to [4].

C Proof of fundamental theorem for weak convergence

To carry out the weak error analysis, we introduce the function $u: [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ defined by $u(t, x) = \mathbb{E}[\varphi(X(t, x; T))]$. We first prove a useful result.

Lemma C.1 (One-step error). Under the same condition of Theorem 2.4, function u fulfills

$$\sup_{t \in [0, T-h]} |\mathbb{E}[u(t+h, X(t, x; t+h))] - \mathbb{E}[u(t+h, Y(t, x; t+h))]| \leq C(1 + |x|^{\beta\kappa+\varkappa})h^{q+1}, \quad (\text{C.1})$$

where β, \varkappa come from assumption (ii) and (iii) in Theorem 2.4, $h > 0$ and $t+h \leq T$.

Proof. We recall that $\varphi \in \mathbb{G}^{2q+2}$. By the results in Corollary B.3 and Lemma 2.5 together with chain rule we could show that $u(t, x)$ is $2q+2$ times differentiable with respect to x and

$$\left| \frac{\partial^k u(t+h, x)}{\partial x_{i_1} \dots \partial x_{i_k}} \right| \leq C(1 + |x|^\kappa), \quad k = 1, 2, \dots, 2q+2. \quad (\text{C.2})$$

By the Taylor expansion, we get

$$\begin{aligned} & \mathbb{E}[u(t+h, X(t, x; t+h))] - \mathbb{E}[u(t+h, Y(t, x; t+h))] \\ &= \sum_{k=1}^{2q+1} \sum_{i_1, \dots, i_k=1}^d \frac{1}{k!} \frac{\partial^k u(t+h, x)}{\partial x_{i_1} \dots \partial x_{i_k}} \mathbb{E}\left[\prod_{j=1}^k (\delta_{X,x})^{i_j} - \prod_{j=1}^k (\delta_{Y,x})^{i_j} \right] + \mathbb{E}[\mathcal{R}_{2q+2}], \end{aligned} \quad (\text{C.3})$$

where we denote $(\delta_{X,x})^\alpha := \prod_{j=1}^{2q+2} (\delta_{X,x})^{i_j}$, $(\delta_{Y,x})^\alpha := \prod_{j=1}^{2q+2} (\delta_{Y,x})^{i_j}$, $\alpha = (i_1, i_2, \dots, i_{2q+2})$ and

$$\begin{aligned} \mathcal{R}_{2q+2} &:= \sum_{|\alpha|=2q+2} \frac{|\alpha|}{\alpha!} \left(\int_0^1 (1-s)^{|\alpha|-1} D^\alpha u(t+h, x + s\delta_{X,x}) ds (\delta_{X,x})^\alpha \right. \\ &\quad \left. - \int_0^1 (1-s)^{|\alpha|-1} D^\alpha u(t+h, x + s\delta_{Y,x}) ds (\delta_{Y,x})^\alpha \right). \end{aligned}$$

Combining the assumption (ii) with (C.2) yields

$$\sum_{k=1}^{2q+1} \sum_{i_1, \dots, i_k=1}^d \frac{1}{k!} \left| \frac{\partial^k u(t+h, x)}{\partial x_{i_1} \dots \partial x_{i_k}} \right| \cdot \left| \mathbb{E}\left[\prod_{j=1}^k (\delta_{X,x})^{i_j} - \prod_{j=1}^k (\delta_{Y,x})^{i_j} \right] \right| \leq C(1 + |x|^{\kappa+\varkappa})h^{p+1}. \quad (\text{C.4})$$

Using the Hölder inequality, (2.8), assumption (iii) and (C.2), one can derive

$$\begin{aligned}
|\mathbb{E}[\mathcal{R}_{2q+2}]| &\leq \sum_{|\alpha|=2q+2} C \int_0^1 \|D^\alpha u(t+h, (1-s)x + sX(t, x; t+h))\|_{L^2(\Omega; \mathbb{R})} ds \|(\delta_{X,x})^\alpha\|_{L^2(\Omega; \mathbb{R})} \\
&\quad + \sum_{|\alpha|=2q+2} C \int_0^1 \|D^\alpha u(t+h, (1-s)x + sY(t, x; t+h))\|_{L^2(\Omega; \mathbb{R})} ds \|(\delta_{Y,x})^\alpha\|_{L^2(\Omega; \mathbb{R})} \\
&\leq C(1 + |x|^{\beta\kappa+\varkappa})h^{p+1},
\end{aligned} \tag{C.5}$$

where we use the assumption $p_1 = 2\kappa$, $p = 2\kappa$, and $\mathbb{P} = 4q + 4$. Plugging (C.4) and (C.5) into (C.3), we get

$$|\mathbb{E}[u(t+h, X(t, x; t+h))] - \mathbb{E}[u(t+h, Y(t, x; t+h))]| \leq C(1 + |x|^{\beta\kappa+\varkappa})h^{p+1}.$$

The proof of the lemma is thus complete. \square

Proof of Theorem 2.4. The proof follows the main ideas from [22, 23, 31]. Here, we show the key steps of the proof. Fix $\varphi \in \mathbb{G}^{2q+2}$ and define $u(t, x) = \mathbb{E}[\varphi(X(t, x; T))]$. Then, we find that

$$\mathbb{E}[\varphi(X(t_0, X_0; T))] - \mathbb{E}[\varphi(Y(t_0, Y_0; t_N))] = \mathbb{E}[u(t_0, X_0)] - \mathbb{E}[u(t_N, Y_N)],$$

and

$$\mathbb{E}[u(t_0, X_0)] = \mathbb{E}[u(t_1, X(t_1))] - \mathbb{E}[u(t_1, Y_1)] + \mathbb{E}[u(t_2, X(t_1, Y_1; t_2))].$$

Continuing this process and subtracting $\mathbb{E}[u(t_N, Y_N)]$, we get

$$\mathbb{E}[u(t_0, X_0)] - \mathbb{E}[u(t_N, Y_N)] = \sum_{i=0}^{N-1} \left(\mathbb{E}[u(t_{i+1}, X(t_i, Y_i; t_{i+1}))] - \mathbb{E}[u(t_{i+1}, Y(t_i, Y_i; t_{i+1}))] \right), \tag{C.6}$$

and so

$$\begin{aligned}
&|\mathbb{E}[\varphi(X(t_0, X_0; T))] - \mathbb{E}[\varphi(Y(t_0, Y_0; t_N))]| \\
&\leq \sum_{i=0}^{N-1} \left| \mathbb{E}[u(t_{i+1}, X(t_i, Y_i; t_{i+1}))] - \mathbb{E}[u(t_{i+1}, Y(t_i, Y_i; t_{i+1}))] \right| \\
&= \sum_{i=0}^{N-1} \left| \mathbb{E} \left[\mathbb{E}[u(t_{i+1}, X(t_i, Y_i; t_{i+1})) - u(t_{i+1}, Y(t_i, Y_i; t_{i+1})) | \sigma(Y_i)] \right] \right|.
\end{aligned} \tag{C.7}$$

Finally, combining a conditional version of (C.1) with (C.7) yields

$$|\mathbb{E}[\varphi(X(t_0, X_0; T))] - \mathbb{E}[\varphi(Y(t_0, Y_0; t_N))]| \leq C(1 + |X_0|^{\beta(\beta\kappa+\varkappa)})Th^q, \tag{C.8}$$

which validates the desired assertion (2.19). \square

D Proofs of moment bound of modified Euler schemes

Proof of Lemma 3.2. In the proof, we shall use the letter C to denote various constants independent of h and n . Let $\mathcal{R} > 0$ be sufficiently large and define a sequence of decreasing subevents

$$\Omega_{\mathcal{R},n} := \left\{ \omega \in \Omega : \sup_{0 \leq i \leq n} |Y_i(\omega)| \leq \mathcal{R} \right\}, \quad n = 0, 1, \dots, N, \quad N \in \mathbb{N},$$

and their compliments $\Omega_{\mathcal{R},n}^c$. At first, we show that the boundedness of high-order moments is valid within a family of appropriate subevents. For integer $\bar{p} \geq 1$, We have

$$\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n+1}} |Y_{n+1}|^{\bar{p}}] \leq \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}] + \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} (\bar{p}(Y_n, Y_{n+1} - Y_n) + \frac{\bar{p}(\bar{p}-1)}{2} |Y_{n+1} - Y_n|^2)]$$

$$+C \sum_{l=3}^{\bar{p}} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-l} |Y_{n+1} - Y_n|^l]. \quad (\text{D.1})$$

Consider the second term in the right-hand side of (D.1):

$$\begin{aligned} I_1 &= \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} (\bar{p} \langle Y_n, Y_{n+1} - Y_n \rangle + \frac{\bar{p}(\bar{p}-1)}{2} |Y_{n+1} - Y_n|^2)] \\ &= \bar{p} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} \langle Y_n, \mathcal{T}_1(f(Y_n), h)h - f(Y_n)h \rangle] + \frac{\bar{p}(\bar{p}-1)}{2} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} |\mathcal{T}_1(f(Y_n), h)h|^2] \\ &\quad + \bar{p} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} (\langle Y_n, f(Y_n)h \rangle + \frac{(\bar{p}-1)}{2} |\sum_{r=1}^m \mathcal{T}_2(g^r(Y_n), h) \Delta W_r(n)|^2)]. \end{aligned}$$

Using the Schwarz inequality, (H1), (H2) in Assumption 3.1, (2.7) and (2.10) yields

$$\begin{aligned} I_1 &\leq Ch + Ch \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}} |Y_n|^{\bar{p}}] \\ &\quad + \bar{p} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{2\bar{p}-1} |\mathcal{T}_1(f(Y_n), h) - f(Y_n)h|] + \frac{\bar{p}(\bar{p}-1)}{2} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{2\bar{p}-2} |\mathcal{T}_1(f(Y_n), h)h|^2] \\ &\leq Ch + Ch \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}} |Y_n|^{\bar{p}}] + Ch^2 \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}} |Y_n|^{4r+\bar{p}}] \\ &\quad + Ch^{\tau+1} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}} |Y_n|^{(2r+1)l_1+\bar{p}-1}], \end{aligned} \quad (\text{D.2})$$

where $\bar{p} \leq p'_0 - \varepsilon$. Now consider the last term in (D.1)

$$\begin{aligned} I_2 &= C \sum_{l=3}^{\bar{p}} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-l} |Y_{n+1} - Y_n|^l] \\ &\leq C \sum_{l=3}^{\bar{p}} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-l} (h^l |\mathcal{T}_1(f(Y_n), h)|^l + h^{\frac{l}{2}} \sum_{r=1}^m |\mathcal{T}_2(g^r(Y_n), h)|^l)] \\ &\leq Ch + C \sum_{l=3}^{\bar{p}} h^l \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{2rl+\bar{p}}] + C \sum_{l=3}^{\bar{p}} h^{\frac{l}{2}} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{rl+\bar{p}}]. \end{aligned} \quad (\text{D.3})$$

Combining (D.2) and (D.3), we obtain

$$\begin{aligned} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_{n+1}|^{\bar{p}}] &\leq Ch + (1 + Ch) \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}] + Ch^2 \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{4r+\bar{p}}] \\ &\quad + Ch^{\tau+1} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{(2r+1)l_1+\bar{p}-1}] \\ &\quad + C \sum_{l=3}^{\bar{p}} h^l \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{2rl+\bar{p}}] + C \sum_{l=3}^{\bar{p}} h^{\frac{l}{2}} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{rl+\bar{p}}]. \end{aligned} \quad (\text{D.4})$$

Choosing $\mathcal{R} = \mathcal{R}(h) = h^{-1/G_1}$ with $G_1 = 6r \vee \frac{(2r+1)l_1-1}{\tau}$, one can immediately show that for all $l = 3, \dots, \bar{p}$,

$$\begin{aligned} \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{(2r+1)l_1+\bar{p}-1} h^\tau &= \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}} (\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\frac{(2r+1)l_1-1}{\tau}} h)^\tau \leq C \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}, \\ \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{4r+\bar{p}} h &= \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}} (\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{4r} h) \leq C \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}, \\ \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{2rl+\bar{p}} h^{l-1} &= \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}} (\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\frac{2rl}{l-1}} h)^{l-1} \leq C \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}, \\ \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{rl+\bar{p}} h^{\frac{l-2}{2}} &= \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}} (\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\frac{2rl}{l-2}} h)^{\frac{l-2}{2}} \leq C \mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}, \end{aligned}$$

where the constants C are independent of the step size h . Thus we have for (D.4)

$$\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n+1}} |Y_{n+1}|^{\bar{p}}] \leq \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_{n+1}|^{\bar{p}}] \leq Ch + (1 + Ch) \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}].$$

The discrete Gronwall inequality [32, Lemma 1.6] shows that

$$\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}] \leq \exp(Cnh)(1 + |Y_0|^{\bar{p}}) \leq C(1 + |Y_0|^{\bar{p}}). \quad (\text{D.5})$$

It remains to estimate $\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}^c} |Y_n|^p]$. It follows from (3.7), (H1) in Assumption 3.1 that

$$\begin{aligned} |Y_{n+1}| &\leq |Y_n| + Ch^{1-\gamma_1} + \sum_{r=1}^m Ch^{-\gamma_2} |W_r(t_{n+1}) - W_r(t_n)| \\ &\leq |Y_0| + (n+1)Ch^{1-\gamma_1} + \sum_{k=0}^n \sum_{r=1}^m Ch^{-\gamma_2} |W_r(t_{k+1}) - W_r(t_k)|. \end{aligned} \quad (\text{D.6})$$

Note that $\mathbf{1}_{\Omega_{\mathcal{R},n}^c} = \mathbf{1}_{\Omega_{\mathcal{R},n-1}^c} + \mathbf{1}_{\Omega_{\mathcal{R},n-1}} \mathbf{1}_{|Y_n| > \mathcal{R}} = \sum_{i=0}^n \mathbf{1}_{\Omega_{\mathcal{R},i-1}} \mathbf{1}_{|Y_i| > \mathcal{R}}$, where we set $\mathbf{1}_{\Omega_{\mathcal{R}(h),-1}} = 1$. This together with the Hölder inequality with $\frac{1}{p'} + \frac{1}{q'} = 1$ for $q' = \frac{\bar{p}}{(p\gamma_1+1)G_1} \vee \frac{\bar{p}}{(1+\frac{1}{2}p+p\gamma_2)G_1} > 1$, due to $p \leq \frac{\bar{p}-G_1}{1+\gamma_1 G_1} \wedge \frac{\bar{p}-G_1}{1+(\frac{1}{2}+\gamma_2)G_1}$, and the Chebyshev inequality give

$$\begin{aligned} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R}(h),n}^c} |Y_n|^p] &= \sum_{i=0}^n \mathbb{E}[|Y_n|^p \mathbf{1}_{\Omega_{\mathcal{R},i-1}} \mathbf{1}_{|Y_i| > \mathcal{R}}] \\ &\leq \sum_{i=0}^n (\mathbb{E}[|Y_n|^{pp'}])^{\frac{1}{p'}} (\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},i-1}} \mathbf{1}_{|Y_i| > \mathcal{R}}])^{\frac{1}{q'}} \\ &= \mathbb{E}[|Y_n|^{pp'}]^{\frac{1}{p'}} \sum_{i=0}^n (\mathbb{P}(\mathbf{1}_{\Omega_{\mathcal{R},i-1}} |Y_i| > \mathcal{R}))^{\frac{1}{q'}} \\ &\leq (\mathbb{E}[|Y_n|^{pp'}])^{\frac{1}{p'}} \sum_{i=0}^n \frac{(\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},i-1}} |Y_i|^{\bar{p}}])^{\frac{1}{q'}}}{\mathcal{R}^{\bar{p}/q'}}. \end{aligned} \quad (\text{D.7})$$

Since $p \leq \frac{\bar{p}-G_1}{1+\gamma_1 G_1} \wedge \frac{\bar{p}-G_1}{1+(\frac{1}{2}+\gamma_2)G_1}$ implies $pp' \leq \bar{p}$, using the Hölder inequality, (D.6) and the elementary inequality implies

$$\begin{aligned} (\mathbb{E}[|Y_n|^{pp'}])^{\frac{1}{p'}} &\leq (\mathbb{E}[|Y_n|^{\bar{p}}])^{\frac{p}{\bar{p}}} \leq C \left(|Y_0|^{\bar{p}} + n^{\bar{p}} h^{\bar{p}(1-\gamma_1)} + h^{-\bar{p}\gamma_2} \mathbb{E} \left[\left| \sum_{k=0}^{n-1} |W_r(t_{k+1}) - W_r(t_k)| \right|^{\bar{p}} \right] \right)^{\frac{p}{\bar{p}}} \\ &\leq C(h^{-p\gamma_1} + h^{-\frac{p}{2}-p\gamma_2}) + C(1 + |X_0|^{\bar{p}})^{\frac{p}{\bar{p}}}. \end{aligned} \quad (\text{D.8})$$

Inserting (D.8) into (D.7) and exploiting the Hölder inequality, $\mathcal{R} = h^{-1/G_1}$ and (D.5), we deduce

$$\begin{aligned} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}^c} |Y_n|^p] &\leq C(n+1)h^{\frac{\bar{p}}{G_1 q'}} (h^{-p\gamma_1} + h^{-\frac{p}{2}-p\gamma_2} + (1 + |X_0|^{\bar{p}})^{\frac{p}{\bar{p}}}) (1 + |X_0|^{\bar{p}})^{\frac{1}{q'}} \\ &\leq C(1 + |X_0|^{\bar{p}})^{\frac{p}{\bar{p}} + \frac{1}{q'}}. \end{aligned} \quad (\text{D.9})$$

This together with (D.5) implies

$$\begin{aligned} \mathbb{E}[|Y_n|^p] &= \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^p] + \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}^c} |Y_n|^p] \\ &\leq (\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}])^{\frac{p}{\bar{p}}} + \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}^c} |Y_n|^p] \\ &\leq C(1 + |X_0|^{\bar{p}})^{\beta} \end{aligned} \quad (\text{D.10})$$

for all integer $p \in [1, \frac{\bar{p}-G_1}{1+\gamma_1 G_1} \wedge \frac{\bar{p}-G_1}{1+(\frac{1}{2}+\gamma_2)G_1}]$. Here $\beta = 1 + \frac{\bar{p}}{pq'} = 1 + \frac{(p\gamma_1+1)G_1}{p} \wedge \frac{(1+\frac{1}{2}p+p\gamma_2)G_1}{p}$. Then, by Jensen's inequality, (3.8) holds for real p as well.

The proof of bounded moments for the scheme (3.7) under (H2') is nearly identical to that under (H2) in Assumption 3.1, except for the second term I_1 on the right-hand side of (D.1). By utilizing (3.3) and assumption (H2'), we can obtain

$$I_1 = \bar{p}h \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} (\langle Y_n, \mathcal{T}_1(f(Y_n), h) \rangle + \frac{(\bar{p}-1)}{2} |\sum_{r=1}^m \mathcal{T}_2(g^r(Y_n), h)|^2)]$$

$$\begin{aligned}
& + \frac{\bar{p}(\bar{p}-1)}{2} \mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}-2} |\mathcal{T}_1(f(Y_n), h)h|^2] \\
\leq & Ch + Ch\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{\bar{p}}] + Ch^2\mathbb{E}[\mathbf{1}_{\Omega_{\mathcal{R},n}} |Y_n|^{4r+\bar{p}}],
\end{aligned}$$

where $\bar{p} \leq p_{\mathcal{T}}$ with $p_{\mathcal{T}}$ from (H2') or (3.6). Similar to the above proof, we choose $\mathcal{R} = h^{-1/\mathbb{G}_1}$, where $\mathbb{G}_1 = 6r$ and we obtain the desired conclusion. \square