

π meson-octet baryon coupling constant obtained by hard-wall AdS/QCD model

Shahin Mamedov ^{a,c,e,f}¹ and Shahnaz Taghiyeva ^{b,d,e}²,

^a *Institute for Physical Problems, Baku State University,
Z. Khalilov street 23, Baku, AZ-1148, Azerbaijan,*

^b *Theoretical Physics Department, Physics Faculty, Baku State University,
Z.Khalilov Street 23, Baku, AZ-1148, Azerbaijan,*

^c *Institute of Physics, Ministry of Science and Education,
H. Javid avenue 33, Baku, AZ-1143, Azerbaijan,*

^d *Shamakhy Astrophysical Observatory, Ministry of Science and Education,
H.Javid 115, Baku, AZ 1000, Azerbaijan,*

^e *Center for Theoretical Physics, Khazar University,
41 Mehseti Street, Baku, AZ-1096, Azerbaijan,*

^f *Lab. for Theor. Cosmology, International Centre of Gravity and Cosmos, Tomsk State
University of Control Systems and Radio Electronics (TUSUR),
634050 Tomsk, Russia*

Abstract

We investigate a strong coupling constant between π meson and octet baryon using the hard-wall AdS/QCD model, where $SU(3)_f$ symmetry-breaking effects are considered. Also, we have mentioned briefly the determining of the profile functions for pseudoscalar meson and baryon fields. The values of couplings obtained here agree with the ones obtained within other theoretical models.

¹ corresponding author: sh.mamedov62@gmail.com

² shahnazilgarzade@gmail.com

I. INTRODUCTION

The anti-de-Sitter/Conformal Field Theory (AdS/CFT) correspondence between two theories is strong-weak, meaning that it relates the strong coupling sector of one theory to the weak coupling sector of the other and vice versa. The AdS/QCD (Anti de-Sitter/Quantum ChromoDynamics) is a simple phenomenological model in five-dimensional curved spacetime, which describes meson and baryon sectors of QCD effectively. In Ref. [1] authors calculated the mass spectra of the baryon octet and their low-lying excited states. They briefly review the holographic model for spin 1/2 baryons. A detailed description of meson-nucleon coupling within the hard-wall AdS/QCD model was given in Ref. [2]. In Ref. [3] a hard-wall model for low-lying spin 1/2 baryons that includes the effect of chiral symmetry breaking was proposed to explain the parity-doublet pattern of excited baryons. The baryon-meson coupling constants were calculated for the spin 1/2 baryon octet and spin 3/2 decuplet in a unified approach relying on symmetry arguments such as the fact that the Yukawa couplings in Ref. [4].

In Ref. [5] the effects of flavor symmetry breaking on three-meson couplings and form factor were investigated and $g_{\rho^n \pi \pi}$, $g_{\rho^n \rho \rho}$, $g_{\rho^n K K}$, $g_{\rho^n K^* K^*}$, $g_{\rho^n D D}$ and $g_{\rho^n D^* D^*}$ strong couplings were predicted using hard-wall holographic model. The two-flavor hard-wall holographic model was extended to four flavors in Ref. [6] and the strong couplings of $(D^{(*)}, D, A)$, $(D^{(*)}, D^{(*)}, V)$, (D_1, D_1, P) , $(\Psi, D^{(*)}, D, P)$, and $(\Psi, D^{(*)}, D, A)$ were calculated after analyzing the wave functions.

We obtain the coupling constants between light pseudoscalar meson π , and octet baryon using the hard-wall AdS/QCD model. The hard-wall model used a sharp cut-off in the extra dimension to realize the confinement and chiral symmetry breaking. In the hard wall model of AdS/QCD the spacetime is given by a five-dimensional anti-de Sitter metric with the additional dimensional labelled as z :

$$ds^2 = g_{MN} dx^M dx^N = \frac{1}{z^2} (\eta_{\mu\nu} dx^\mu dx^\nu - dz^2), \quad (1)$$

where $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$, $\mu, \nu = 0, 1, 2, \dots, d-1$; $g = |\det(g_{MN})| = 1/z^{10}$ the magnitude of the determinant of g_{MN} . Thus, the model is defined in the range: $\epsilon \leq z \leq z_m$, ϵ and z_m are the inverses of the ultraviolet (UV) and infrared (IR) cutoff scales, respectively, in the hard-wall model.

The paper is structured as follows: In the next section, we briefly review our approach-profile functions in the hard-wall model for pseudoscalar and baryon fields. In Sec. III, we define the between π meson- baryon coupling constant in the hard-wall model. Finally, in Sec. IV, we discuss

our results and compare our final results to existing theoretical results and experimental data.

II. PROFILE FUNCTIONS IN THE HARD WALL

A. The wave function of the pseudoscalar meson

Bulk action for the meson sector, which includes the pseudoscalar X field and $A_{L,R}$ gauge fields, is written as below [7]:

$$S = \int d^5x \sqrt{g} \text{Tr} \left[|DX|^2 + 3|X|^2 - \frac{1}{4g_5^2} (F_L^2 + F_R^2) \right], \quad (2)$$

where DX is the covariant derivative of the bulk scalar field: $D^M X = \partial^M X - iA_L^M X + iX A_R^M$ and $F_{L,R}^{MN}$ is the chiral gauge field strength: $F_{L,R}^{MN} = \partial^M A_{L,R}^N - \partial^N A_{L,R}^M - i[A_{L,R}^M, A_{L,R}^N]$. The equation of motion for the pseudoscalar meson is derived using from the least action principle and $A_5 = 0$ gauge condition in this gauge [1]:

$$\partial_z \left(\frac{\partial_z P_\phi}{z} \right) + \frac{4g_5^2 v_u^2}{z^3} (f_\pi - P_\phi) = 0, \quad (3)$$

$$-q^2 \partial_z P_\phi + \frac{4g_5^2 v_u^2}{z^2} \partial_z f_\pi = 0, \quad (4)$$

where $P_\phi(z)$ comes from that of the radial component of the axial-vector field, f_π comes from the KK decomposition of the bulk pion field. As in the calculation of baryon spectrum, we only consider the $m_{u,d} = 0$ case in which the pion as Nambu-Goldstone boson has zero mass ($q^2 = 0$), so we have $f'_\pi(z) = 0$, which also indicates with the normalization condition $f_\pi(z \rightarrow 0) = 0$ that $f_\pi(z) = 0$. Then the equation of motion (4) can be simplified as following:

$$z \partial_z \left(\frac{\partial_z P_\phi}{z} \right) - \frac{g_5^2 v_u^2}{z^2} P_\phi = 0. \quad (5)$$

The solution of the equation (5) is found as below [8]:

$$P_\phi(z) = N \xi z^3 \left[I_{2/3} \left(\frac{\sqrt{2}}{3} \xi z^3 \right) - \frac{I_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right)}{K_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right) K_{2/3} \left(\frac{\sqrt{2}}{3} \xi z^3 \right)} \right]. \quad (6)$$

Here $\xi = 3.77 z_m^{-3}$, $I_{2/3}$, $K_{2/3}$ are Bessel functions. The normalization constant N is determined from the normalization condition [9, 10], and equals:

$$N^{-1/2} = \frac{I_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right)}{2K_{2/3}} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right)^2 \left[\sqrt{3} \pi I_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right) + 3K_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right) \right]. \quad (7)$$

B. Profile function for baryon field

Let us briefly present obtaining the profile function for the baryon field within the hard-wall model. The bulk Lagrangian for the fermion field, which describes the baryons on the boundary theory, is written as below:

$$L_{Baryon} = \sqrt{-g} \left[\frac{i}{2} \bar{N}_1 e_A^M \Gamma^A \nabla_M N_1 - \frac{i}{2} (\nabla_M^\dagger \bar{N}_1) e_A^M \Gamma^A N_1 - m_5 \bar{N}_1 N_1 \right], \quad (8)$$

where $e_A^M = \frac{1}{z} \eta_M^A$ is the vielbein satisfying $g_{MN} = e_A^M e_B^N \eta_{AB}$, $\Gamma^A = (\gamma^\mu, -i\gamma^5)$ are the Dirac matrices and satisfy: $[\Gamma^A, \Gamma^B] = 2\eta^{AB}$ and ∇_M is the gauge and Lorentz covariant derivative:

$$\nabla_M = \partial_M + \frac{i}{4} \omega_M^{AB} \Gamma_{AB} - i(A_L^a)_{Mt^a}. \quad (9)$$

Here $\Gamma^{AB} = \frac{1}{2i}[\Gamma^A, \Gamma^B]$, ω_M^{AB} is spin connection with nonzero components: $\omega_M^{5A} = -\omega_M^{A5} = \frac{1}{z} \delta_M^A$. The Yukawa interaction in the bulk is introduced to include the chiral symmetry breaking:

$$L_{Yukawa} = -g_Y \bar{N}_1 X N_1 - g_Y \bar{N}_2 X^\dagger N_2, \quad (10)$$

where g_Y is the Yukawa constant. The bulk baryon field can be written the chiral components:

$$N_1 = N_{1L} + N_{1R}, \quad (11)$$

where $i\Gamma^5 N_{1L} = N_{1L}$ and $i\Gamma^5 N_{1R} = -N_{1R}$. Then a Kaluza-Klein decomposition for $N_{1L,R}$ is performed to yield the following form:

$$N_{1L,R}(x, z) = \sum_n \int \frac{d^4 p}{(2\pi)^4} f_{1L,R}(z) \psi_{L,R}^{(n)}(p) e^{-ipx}, \quad (12)$$

$$\not{p} \Psi_{(1L,R)}(p) = |p| \Psi_{(1R,L)}(p). \quad (13)$$

We get an equation for $f_{L,R}(z)$:

$$\begin{pmatrix} \partial_z - \frac{\Delta}{z} & -\frac{g_Y v(z)}{z} \\ \frac{g_Y v(z)^\dagger}{z} & \partial_z - \frac{4-\Delta}{z} \end{pmatrix} \begin{pmatrix} f_{1L} \\ f_{2L} \end{pmatrix} = m \begin{pmatrix} f_{1R} \\ f_{2R} \end{pmatrix}, \quad (14)$$

where $v(z)$ has the following form for octet baryon:

$$v_{p,n}(z) = (3c_2 - c_1) v_u(z), \quad (15)$$

$$v_{\Sigma_s}(z) = 2c_2 v_u(z) + (c_2 - c_1) v_s(z), \quad (16)$$

$$v_{\Xi s}(z) = (c_2 - c_1) v_u(z) + 2c_2 v_s(z). \quad (17)$$

In the chiral limit, the f_1 and f_2 sectors are completely decoupled from each other and the equation of motion for f_1 sector is found as follows [2]:

$$\left(\partial_z^2 - \frac{4}{z} + \frac{6 + m_5 - m_5^2}{z^2} \right) f_{1L}^n(z) = -m_n^2 f_{1L}^n(z), \quad (18)$$

$$\left(\partial_z^2 - \frac{4}{z} - \frac{6 + m_5 - m_5^2}{z^2} \right) f_{1R}^n(z) = -m_n^2 f_{1R}^n(z), \quad (19)$$

where m_n represents the KK masses for the baryons. The solutions of the (18), (19) equations can be obtained as:

$$f_{1L,R}^n(z) = z^{\frac{5}{2}} \left[c_{1L} J_{|m_5 \mp \frac{1}{2}|}(m_n z) + c_{2L} Y_{|m_5 \mp \frac{1}{2}|}(m_n z) \right]. \quad (20)$$

The scaling dimension of the baryon operator is $\Delta = \frac{9}{2}$, according to $m_5^2 = (\Delta - 2)^2$ the of m_5 value will be $m_5 = \pm \frac{5}{2}$. Therefore, we can write $f_{L,R}$ as follows:

$$f_{1L}^n = z^{\frac{5}{2}} [c_{1L} J_2(m_n z) + c_{2L} Y_2(m_n z)], \quad (21)$$

$$f_{1R}^n = z^{\frac{5}{2}} [c_{1R} J_3(m_n z) + c_{2R} Y_3(m_n z)]. \quad (22)$$

c_{2L} constant is found from the UV boundary condition $f_{1L}^n(0) = 0$ we get:

$$c_{2L}^n = -\frac{J_2(m_n \varepsilon)}{Y_2(m_n \varepsilon)} c_{1L}^n = \frac{\pi}{2} \left(\frac{m_n \varepsilon}{2} \right)^4 c_{1L}^n \rightarrow 0. \quad (23)$$

Finally, f_{1L} accepts simple form:

$$f_{1L}^n = c_{1L} z^{\frac{5}{2}} J_2(m_n z). \quad (24)$$

f_{1R} is found by taking into account f_{1L} in (12):

$$f_{1R}^n = c_{1R} z^{\frac{5}{2}} J_3(m_n z), \quad (25)$$

$f_{2R,L}$ is found applying $f_{1R}^n(0) = 0$ boundary condition and taking $m_5 = -\frac{5}{2}$:

$$f_{2R}^n = c_{2R} z^{\frac{5}{2}} J_2(m_n z), \quad (26)$$

$$f_{2L}^n = -c_{2L} z^{\frac{5}{2}} J_3(m_n z). \quad (27)$$

$c_{1,2}$ constants are found from the normalization condition [2]:

$$|c_{1,2}| = \frac{\sqrt{2}}{z_m J_2(m_n z_m)}. \quad (28)$$

III. π MESON-OCTET BARYON COUPLING CONSTANT

The action for the interaction between bulk fields in the AdS space can be written as the $5D$ integral:

$$S = \int_0^{z_m} d^4x dz \sqrt{g} L_{int}(x, z). \quad (29)$$

Here L_{int} is the interaction Lagrangian. The interaction Lagrangian for the π meson and octet baryon is given as:

$$L_{\pi bb} = -g_Y \left[\bar{B}_1 X B_2 + \bar{B}_2 X^+ B_1 \right]. \quad (30)$$

Here X is a scalar field and is defined as $X = v(z)e^{iPx}$. $B_{1,2}$ baryon fields can be written as follows by Fourier transformation:

$$B_{1,2L,R} = \frac{1}{(2\pi)^4} \int d^4p f_{1,2L,R} \Psi(p) e^{-ipx}. \quad (31)$$

The map between bulk and boundary quantities is called GKPW (Gubser-Klebanov-Polyakov-Witten) formula which states that:

$$Z_{sugra}|J| \equiv \int_{\Phi \sim J} D\Phi e^{-S[\Phi]} = \langle e^{-\int_{\partial AdS} JO} \rangle \equiv Z_{QFT}, \quad (32)$$

where the left-hand side is the supergravity partition function in anti-de Sitter space, and the right-hand side is the partition function of the Quantum Field Theory (QFT) boundary of the asymptotically this spacetime. As seen from (32) relation, $\Phi \sim J$ means that the field theory side J is a source coupled to an operator O which is dual to Φ . In the limit of small string coupling and large string tension, the equivalence between the partition function of the QFT and the partition function of supergravity will be as follows:

$$e^{-S_{on-shell}[J]} = \langle e^{-\int_{\partial AdS} JO} \rangle_{N \rightarrow \infty, \lambda \rightarrow \infty}. \quad (33)$$

Here, the left-hand side is on-shell supergravity action, and the right-hand side is the QFT partition function in the large N and strong coupling limits. We can define expectation values of CFT observables by taking functional derivatives of this relation

$$\langle O(x) \rangle \sim \frac{\delta S_{on-shell}}{\delta \Phi(x)} \Big|_{\Phi(x)=0}. \quad (34)$$

According to AdS/CFT correspondence the relation (34) can be written for the vacuum expectation value of the baryon current J_μ :

$$\langle J_\mu \rangle = -i \frac{\delta Z_{AdS}}{\delta \tilde{V}_\mu^0} \Big|_{\tilde{V}_\mu^0} = 0, \quad (35)$$

where $\tilde{V}_\mu^0 = \tilde{V}_\mu(q, z=0) = V(q)$ is the UV boundary value of the vector field ($V(z=0) = 1$). On the other side, a current in the QCD theory is defined as:

$$J_5(p', p) = g_{\pi bb} \bar{u}(p') \gamma_5 u(p). \quad (36)$$

The π meson - octet baryon coupling constant $g_{\pi bb}$ defines from a comparison of (35) and (36) equations:

$$g_{\pi bb} = \int_0^{z_m} \frac{dz}{z^5} v(z) g_Y P(z) (f_{1R}(z) f_{2L}(z) - f_{1L}(z) f_{2R}(z)). \quad (37)$$

IV. NUMERICAL RESULTS

Having fixed the parameters σ and m_q the constant $g_{\pi bb}$ (37) can be calculated numerically. We apply the $\sigma = (0.213)^3 \text{ GeV}^3$ and $m_q = 0.0083 \text{ GeV}$ values for these parameters, which were fixed in Ref. [11]. For the constant g_Y we take the value $g_Y = 9.182$ which was established in Ref. [12]. In Table 1 we compare our results to the ones obtained within the chiral soliton model and light cone QCD sum rule.

As seen from the Table 1, $g_{\pi \Sigma^0 \Sigma^0}$ is close to the chiral soliton model result. The numerical value for the between π meson and Ξ^0 baryon coupling constant corresponds to the result obtained within the light cone QCD sum rules framework. For the $g_{\pi \Lambda \Lambda}$ coupling constant hard-wall result is very small ($g_{\pi \Lambda \Lambda} = 4, 59652 \times 10^{-6}$). The result here is reasonable from a physical interpretation point of view. Since the current of the Λ baryon is antisymmetric concerning the u, d quarks, and we use a chiral limit state, this coupling constant is automatically equal to zero.

Table 1. π meson-octet baryon coupling constant

Model	$g_{\pi NN}$	$g_{\pi \Sigma^0 \Sigma^0}$	$g_{\pi \Sigma^- \Sigma^-}$	$g_{\pi \Sigma^+ \Sigma^+}$	$g_{\pi \Xi^0 \Xi^0}$	$g_{\pi \Xi^- \Xi^-}$
Hard-wall model	2.941	3.516	3.554	3.478	2.4799	2.516
Chiral soliton model [13]	3.524 ± 0.012	3.356 ± 0.014	-	-	-1.24 ± 0.009	-
Light cone QCD sum rule [14]	-	2.8 ± 0.3	-	-	2.4 ± 0.2	-

-
- [1] Z. Fang, "Holographic model for the baryon octet," *Physical Review D*, vol. 94, no. 7, p. 074017, 2016.
[2] N. Maru and M. Tachibana, "Meson-nucleon coupling from ads/qcd," *The European Physical Journal C*, vol. 63, pp. 123-132, 2009.

- [3] D. K. Hong, T. Inami, and H.-U. Yee, “Baryons in ads/qcd,” *Physics Letters B*, vol. 646, no. 4, pp. 165–171, 2007.
- [4] L. L. Lopes, K. D. Marquez, and D. P. Menezes, “Baryon coupling scheme in a unified su (3) and su (6) symmetry formalism,” *Physical Review D*, vol. 107, no. 3, p. 036011, 2023.
- [5] A. Ballon-Bayona, G. Krein, and C. Miller, “Strong couplings and form factors of charmed mesons in holographic qcd,” *Physical Review D*, vol. 96, no. 1, p. 014017, 2017.
- [6] S. Momeni and M. Saghebfar, “Erratum to: Charm meson couplings in hard-wall holographic qcd,” *The European Physical Journal C*, vol. 82, no. 8, p. 709, 2022.
- [7] J. Erlich, E. Katz, D. T. Son, and M. A. Stephanov, “Qcd and a holographic model of hadrons,” *Physical Review Letters*, vol. 95, no. 26, p. 261602, 2005.
- [8] E. Katz, A. Lewandowski, and M. D. Schwartz, “Tensor mesons in ads/qcd,” *Physical Review D*, vol. 74, no. 8, p. 086004, 2006.
- [9] —, “Tensor mesons in ads/qcd,” *Physical Review D*, vol. 74, no. 8, p. 086004, 2006.
- [10] S. Mamedov, Z. Hashimli, and S. Jafarzade, “Tensor meson couplings in ads/qcd,” *Physical Review D*, vol. 108, no. 11, p. 114032, 2023.
- [11] A. Cherman, T. D. Cohen, and E. S. Werbos, “Chiral condensate in holographic models of qcd,” *Physical Review C*, vol. 79, no. 4, Apr. 2009. [Online]. Available: <http://dx.doi.org/10.1103/PhysRevC.79.045203>
- [12] D. K. Hong, H.-C. Kim, S. Siwach, and H.-U. Yee, “The electric dipole moment of the nucleons in holographic qcd,” *Journal of High Energy Physics*, vol. 2007, no. 11, p. 036–036, Nov. 2007. [Online]. Available: <http://dx.doi.org/10.1088/1126-6708/2007/11/036>
- [13] G.-S. Yang and H.-C. Kim, “Meson–baryon coupling constants of the su (3) baryons with flavor su (3) symmetry breaking,” *Physics Letters B*, vol. 785, pp. 434–440, 2018.
- [14] T. M. Aliev, K. Azizi, and M. Savcı, “Strong transitions of decuplet to octet baryons and pseudoscalar mesons,” *Nuclear Physics A*, vol. 847, no. 1-2, pp. 101–117, 2010.