

Starving Random Walks

L. Régnier^[0000-0002-3792-1570],
M. Dolgushev^[0000-0003-3306-9840] and
O. Bénichou^[0000-0002-6749-2391]

Abstract In this chapter, we review recent results on the starving random walk (RW) problem, a minimal model for resource-limited exploration. Initially, each lattice site contains a single food unit, which is consumed upon visitation by the RW. The RW starves whenever it has not found any food unit within the previous \mathcal{S} steps.

To address this problem, the key observable corresponds to the inter-visit time τ_k defined as the time elapsed between the finding of the k^{th} and the $(k+1)^{\text{th}}$ food unit. By characterizing the maximum M_n of the inter-visit times $\tau_0, \dots, \tau_{n-1}$, we will see how to obtain the number $N_{\mathcal{S}}$ of food units collected at starvation, as well as the lifetime $T_{\mathcal{S}}$ of the starving RW.

1 Introduction

When it comes to the exploration of a territory by an animal or biological tracers, a fundamental property in quantifying this exploration is the number $\mathcal{N}(t)$ of sites visited by time t [1, 2]. This quantity has been the focus of many mathematical studies, in which the average, variance, and sometimes the full distribution have been explicitly obtained for a wide range of stochastic processes [3, 4, 5, 6, 7, 8, 9]. These processes include nearest-neighbor jumps on a d -dimensional lattice, on fractal media, or RWs with long-range jumps, which model a large variety of biological

L. Régnier
Sorbonne Université, Laboratoire de Physique Théorique de la Matière Condensée (LPTMC), 4
Place Jussieu, 75005 Paris, France e-mail: leo.regnier.pro@outlook.fr

M. Dolgushev
Sorbonne Université, Laboratoire de Physique Théorique de la Matière Condensée (LPTMC), 4
Place Jussieu, 75005 Paris, France e-mail: maxim.dolgushev@sorbonne-universite.fr

O. Bénichou
Sorbonne Université, CNRS, Laboratoire de Physique Théorique de la Matière Condensée
(LPTMC), 4 Place Jussieu, 75005 Paris, France e-mail: benichou@lptmc.jussieu.fr

motions. However, since $\mathcal{N}(t)$ is a cumulative quantity, it does not provide significant information about the past exploration dynamics of the walk: were all the visitations occurring regularly, or were there large time gaps where the RW did not encounter any food units? This differentiation becomes crucial when considering living organisms that rely on finding resources to propel their motion.

Recently, a new quantity has been introduced to enable this differentiation [8, 10, 11, 12, 13]: the time τ_k . This represents the time it takes for a RW to find a site it has never visited before, given that k sites have already been visited. These random variables encapsulate the full dynamics of RW exploration and have been shown to be particularly useful in characterizing the main properties of the starving RW problem [10, 14], a minimal model to describe depletion-controlled dynamics of a RW [15, 16] as encountered in bacteria [17], animals [18], or robots [19]. In the original formulation of the problem, food units are placed on every site of the lattice, and the RW moves through the medium, consuming each food unit upon encounter. The RW starves if \mathcal{S} time steps (the metabolic time) pass without a food encounter (see Fig. 1 for an illustration). This definition establishes a profound connection between τ_k and starvation: starvation occurs the first time we observe a




Fig. 1 The starving RW model. In the lattice, each site initially holds one food unit, which is consumed upon visit by the RW. The RW can make \mathcal{S} steps (here $\mathcal{S} = 50$) without encountering food before starving. Two trajectories (green and red) are depicted. A forager (yellow) has consumed $\mathcal{N}(t) = 1000$ food units. Areas devoid of food are depicted in black. Along the green trajectory, the forager (green) promptly discovers a new food unit, so that $\tau_{1000} < \mathcal{S}$. Conversely, following the red trajectory, it fails to locate food before succumbing to starvation (depicted by the red cross), $\tau_{1000} \geq \mathcal{S}$.

time between visits to new sites, and thus the finding of a new food unit, larger than the metabolic time. Consequently, understanding the long inter-visit times is fundamental for the search for food, necessitating the study of the extreme-value statistics [20, 21, 22, 23, 24]. We will explore this connection further in the following sections.

In this chapter, we examine the visitation properties of general classes of RWs, including anomalous diffusion. We investigate symmetric Markovian RWs that:

- Move in a medium of fractal dimension d_f , such that the number of sites within a ball of radius r grows as r^{d_f} .
- Have a distance to the origin r_t after t jumps that grows as t^{1/d_w} , where d_w is the walk dimension [25]. More precisely, the distribution of $r_t/t^{1/d_w}$ is independent of t for large t .

We note that diffusive RWs in hypercubic lattices of dimension d ($d_w = 2$ and $d_f = d$), as well as Lévy flights (jumps whose length distribution $p(\ell)$ is distributed algebraically with exponent $1 + \alpha$, $p(\ell) \propto 1/|\ell|^{1+\alpha}$, such that $d_w = 1/\alpha$ for $\alpha < 2$) are included in our description.

We begin with a brief summary of results on the visitation process of RWs, focusing particularly on the inter-visit time statistics τ_k of the general RWs under consideration, whose properties depend mainly on the ratio $\mu \equiv d_f/d_w$. Then, we delve into the two original cases in which the starving RW problem was examined [10, 14]: the unidimensional and infinite-dimensional cases, which can be viewed as the two limiting scenarios. Finally, we present the general framework for obtaining the properties of both the number N_S of sites visited at starvation and the lifetime T_S of the starving RW.

2 Visitation properties

In this section, we review the results concerning the exploration dynamics of general Markovian processes which will be useful throughout this chapter.

2.1 Number of sites visited

In the context of symmetric Markovian RWs characterized by a fractal dimension d_f and a walk dimension d_w , it was shown in Refs. [5, 26, 27, 28] that there exist two main different classes of exploration: either $\mu \leq 1$, and the RW visits every site infinitely often, or $\mu > 1$, and the RW will never visit some sites. The first type of RWs is said to be recurrent, while the second type is said to be transient. Of note, the case $\mu = 1$ is differentiated and called marginally recurrent, which we will abbreviate

to marginal in the following.

This distinction is particularly striking when one considers the average number of sites visited by time t (see Refs. [6, 7, 29]),

$$\langle \mathcal{N}(t) \rangle \propto \begin{cases} t^\mu & [\mu < 1] \\ \frac{t}{\ln t} & [\mu = 1] \\ t & [\mu > 1] . \end{cases} \quad (1)$$

The difference between the behavior at $\mu < 1$ and $\mu > 1$ can be understood as follows: while recurrent RWs typically visit the entirety of the ball of volume $r_t^{d_f} \propto t^\mu$ (the number t of steps they perform is much larger than the number of accessible sites $r_t^{d_f} = t^\mu \ll t$), transient RWs rarely return to previous sites and thus visit a new site at almost every time step. Such strong differences also appear in the exploration dynamics for both types of processes, as we proceed to show in the following section.

2.2 Inter-visit times statistics

Below, we summarize the inter-visit time statistics, represented through their probability density $F_k(\tau) \equiv \mathbb{P}(\tau_k = \tau)$, obtained for general Markovian RWs in [11].

Table 1 Summary of the time dependence of $F_k(\tau)$ for the three classes of RWs—recurrent, marginal, and transient. The constants are independent of k and τ . The crossover times t_k and T_k are given up to logarithmic prefactors.

$F_k(\tau)$	t_k	T_k	$1 \ll \tau \ll t_k$	$t_k \ll \tau \ll T_k$	$T_k \ll \tau$
$\mu < 1$ [recurrent]	$k^{1/\mu}$	$k^{1/\mu}$	$\tau^{-(1+\mu)}$	$\exp[-\text{const}(\tau/t_k)^{\mu/(1+\mu)}]$	$\exp[-\text{const}\tau/k^{1/\mu}]$
$\mu = 1$ [marginal]	\sqrt{k}	$k^{3/2}$			
$\mu > 1$ [transient]	1	$k^{(\mu+1)/\mu}$			

The results are as follows:

- For recurrent RWs, $\mu < 1$: there are two distinct time regimes. The first one occurs at times $\tau \ll t_k = k^{1/\mu}$, for which $F_k(\tau)$ decays algebraically with exponent $1+\mu$, independently of k . Then, for $\tau \gg t_k$, $F_k(\tau)$ decays exponentially, at a rate of $1/k^{1/\mu}$. Overall, $F_k(\tau)$ admits a scaling form:

$$F_k(\tau) = \frac{1}{k^{1+\mu}} \psi(\tau/k^{1/\mu}) \quad (2)$$

where ψ is a scaling function that depends only on the process.

- For transient RWs, $\mu > 1$: there are two distinct time regimes once again, but the early time regime $\tau \ll T_k = k^{1+1/\mu}$ is now characterized by an exponential decay in the variable $\tau^{\mu/(1+\mu)}$ (stretched exponential in τ), and the late time regime is an exponential decay in the variable $\tau/k^{1/\mu}$, as for the recurrent case. There is no algebraic decay.
- For marginal RWs, $\mu = 1$: there are now three distinct regimes. Firstly, the distribution $F_k(\tau)$ decays algebraically for $\tau \ll \sqrt{k}$, with an exponent of 2. This is followed by an exponential decay in the variable $\sqrt{\tau/\sqrt{k}}$ for $\sqrt{k} \ll \tau \ll k^{3/2}$, and finally, an exponential decay for $\tau \gg k^{3/2}$ with a rate of $1/k$. More precisely, in the cross-over regime $\tau \sim \sqrt{k}$, which will be particularly useful in the following, it was shown [30] that $F_k(\tau)$ admits a scaling form:

$$F_k(\tau) = \frac{1}{k^{3/2}} \phi(\tau/\sqrt{k}) \quad (3)$$

where ϕ decays algebraically with an exponent of 3 at small arguments and exponentially with $\sqrt{\tau/\sqrt{k}}$ at large arguments. With this we get in the cross-over an algebraic decay τ^{-3} followed by the stretched exponential regime indicated in Table 1. This rich cross-over behavior comes from the analysis of the exit times from a cascade of the largest visited regions, see Ref. [30] for details.

Before applying these fundamental results describing the general τ_k properties to the starving RW problem, let us start with the two simplest cases where the inter-visit time statistics can be derived exactly.

3 Low and high dimensional cases

In this section, we review the two limit cases of unidimensional and infinite dimensional starving RWs, which were solved in [10, 14].

3.1 The 1d case

We start with the case of a symmetric nearest-neighbour RW on a $1d$ line. In this model, the RW jumps with equal probability to one of its two neighbouring lattice sites, on the left or on its right. In this situation, the domain visited by the RW is simple, as it is simply an interval, and the RW starts its new exploration of a new site on the border of the interval (new visits occur only on the left or right border). Thus, the time τ_k to visit a new site when k sites have already been visited is simply given by the time to exit this interval of length k . Consequently, the precise distribution $F_k(\tau)$ of τ_k is known, and given by the first exit time distribution of the interval starting on one of its border [31]:

$$F_k(\tau) = \frac{2\pi}{k^2} \sum_{j=0}^{\infty} (2j+1) \sin\left(\frac{2j+1}{k}\pi\right) \exp\left[-(2j+1)^2\pi^2\tau/2k^2\right]. \quad (4)$$

One can check that indeed Eq. (4) is contained in Table 1 for the value $\mu = 1/2$, as $d_f = 1$ and $d_w = 2$ (the RW is diffusive). Additionally, we note that the τ_k are independent, as the distribution of τ_k only depends on the size k of the visited interval. In the following, we see how the knowledge of the τ_k provides the distribution of the number N_S of sites visited at starvation and then of the lifetime T_S .

3.1.1 Number of sites visited at starvation

First, we connect the inter-visit time statistics to the starving RW problem using the following remark: to have visited at least n sites before starvation, it means that all the n first inter-visit times $\tau_0, \tau_1, \dots, \tau_{n-1}$ are smaller than the metabolic time S . It states that there was no time interval longer than S between two visits. From this remark, we deduce that:

$$\mathbb{P}(N_S \geq n) = \mathbb{P}(\tau_k < S, k = 0, \dots, n-1). \quad (5)$$

From the independence of the τ_k , we set that (in the limit of large n and S):

$$\mathbb{P}(N_S \geq n) = \prod_{k=0}^{n-1} \left(1 - \sum_{\tau=S}^{\infty} F_k(\tau)\right) \sim \exp\left[-\int_0^n dk \int_S^{\infty} d\tau F_k(\tau)\right]. \quad (6)$$

Finally, inserting the exact expression of $F_k(\tau)$ given by Eq. (4) in Eq. (6), we obtain:

$$-\ln \mathbb{P}(N_S \geq n) \sim \sum_{j=0}^{\infty} \int_0^n \frac{4dk}{k} e^{-\frac{(2j+1)^2\pi^2 S}{2k^2}} = \sum_{j=0}^{\infty} \int_1^{\infty} \frac{2dx}{x} e^{-\frac{x(2j+1)^2\pi^2 S}{2n^2}} \quad (7)$$

where we used that $\sin\left(\frac{2j+1}{k}\pi\right) \sim \frac{2j+1}{k}\pi$ at large k . We note that Eq. (7) has a scaling form in the variable S/n^2 ,

$$\mathbb{P}(N_S \geq n) \sim \Xi_{1/2}\left(\frac{S}{n^2}\right) \quad (8)$$

where $\Xi_{1/2}$ is the scaling function, independent of n and S . From Eq. (8), we deduce the scaling with S of the number of sites visited at starvation:

$$\begin{aligned} \langle N_S \rangle &\sim \sum_{n=0}^{\infty} \mathbb{P}(N_S \geq n) \\ &\sim \int_0^{\infty} dn \Xi_{1/2}\left(\frac{S}{n^2}\right) = \sqrt{S} \int_0^{\infty} du \Xi_{1/2}\left(\frac{1}{u^2}\right) \approx 2.9\sqrt{S}. \end{aligned} \quad (9)$$

Additionally, one can derive the asymptotic properties of $\Xi_{1/2}(\mathcal{S}/n^2)$ at large and small arguments. For small number of sites visited at starvation $n/\sqrt{\mathcal{S}} \ll 1$, the sum in Eq. (7) is dominated by the $j = 0$ term, such that:

$$-\ln \mathbb{P}(N_{\mathcal{S}} \geq n) \sim \int_1^\infty \frac{2dx}{x} \exp\left[-x \frac{\pi^2 \mathcal{S}}{2n^2}\right] \sim \frac{4n^2}{\pi^2 \mathcal{S}} \exp\left[-\frac{\pi^2 \mathcal{S}}{2n^2}\right]. \quad (10)$$

In the opposite limit of large number of sites visited at starvation $n/\mathcal{S}^2 \gg 1$, we use that $F_k(\tau)$ converges to the first passage time distribution $F_\infty(\tau)$ to the site at distance one as the RW does not see the other border of the visited interval. The expression of $F_\infty(\tau)$ is well-known and given by [31]:

$$F_\infty(\tau) = \frac{1}{\sqrt{2\pi\tau^3}} e^{-1/2\tau}. \quad (11)$$

Once we insert this expression in (6), we get:

$$-\ln \mathbb{P}(N_{\mathcal{S}} \geq n) = \sum_{k=0}^{n-1} \int_{\mathcal{S}}^\infty F_\infty(\tau) d\tau = n \int_{\mathcal{S}}^\infty \frac{d\tau}{\sqrt{2\pi\tau^3}} e^{-1/2\tau} \sim \frac{n}{\sqrt{8\pi\mathcal{S}}}. \quad (12)$$

3.1.2 Lifetime

We now proceed to determine the lifetime $T_{\mathcal{S}}$. Initially, we calculate the average lifetime of the RW, which is the sum of inter-visit times before starvation plus the metabolic time, representing the final search for a food unit. This can be expressed as:

$$\langle T_{\mathcal{S}} \rangle = \left\langle \sum_{k=1}^{N_{\mathcal{S}}} \tau_k \right\rangle + \mathcal{S} = \sum_{n=1}^{\infty} \left(\sum_{k=1}^n \langle \tilde{\tau}_k \rangle \right) \mathbb{P}(N_{\mathcal{S}} = n) + \mathcal{S} \quad (13)$$

Here, $\tilde{\tau}_k$ denotes the k^{th} inter-visit time conditioned to be less than the metabolic time \mathcal{S} . The average of the conditional inter-visit time is given by:

$$\langle \tilde{\tau}_k \rangle = \frac{\int_0^{\mathcal{S}} \tau F_k(\tau) d\tau}{\int_0^{\mathcal{S}} F_k(\tau) d\tau} \quad (14)$$

Considering the limit $k \rightarrow \infty$ for large visited intervals, τ_k follows an asymptotic distribution noted F_∞ , representing the time to exit a semi-infinite interval starting one unit length from the border as given in Eq. (11). Consequently, the limit as $k \rightarrow \infty$ of Eq. (14) is asymptotically:

$$\langle \tilde{\tau}_\infty \rangle = \frac{\int_0^{\mathcal{S}} \tau F_\infty(\tau) d\tau}{\int_0^{\mathcal{S}} F_\infty(\tau) d\tau} \sim \sqrt{\frac{2}{\pi}} \sqrt{\mathcal{S}}. \quad (15)$$

Utilizing Eq. (13), we obtain:

$$\begin{aligned} \langle T_S \rangle - \mathcal{S} &\sim \sum_{n=1}^{\infty} \left(\sum_{j=1}^n \langle \tilde{\tau}_{\infty} \rangle \right) \mathbb{P}(N_S = n) = \langle \tilde{\tau}_{\infty} \rangle \sum_{n=1}^{\infty} n \mathbb{P}(N_S = n) \\ &= \langle \tilde{\tau}_{\infty} \rangle \langle N_S \rangle \approx 2.3 \mathcal{S} . \end{aligned} \quad (16)$$

We can demonstrate more generally that the n^{th} cumulant of the lifetime T_S scales as \mathcal{S}^n , implying that the random variable T_S/\mathcal{S} is independent of \mathcal{S} for large \mathcal{S} . Denoting by $\kappa_n(\dots)$ the n^{th} cumulant function ($n > 1$), we find that for the n^{th} cumulant of the sum of N_S random variables [32], we have:

$$\kappa_n(T_S) \sim \sum_{k_1, \dots, k_n \in S_n} \frac{n! \kappa_{\sum_{i=1}^n k_i}(N_S)}{k_1! \dots k_n! (1!)^{k_1} \dots (n!)^{k_n}} \kappa_1(\tilde{\tau}_{\infty})^{k_1} \dots \kappa_n(\tilde{\tau}_{\infty})^{k_n} \quad (17)$$

where S_n is the set of all vectors $(k_1, \dots, k_n) \in \mathbb{N}^n$ such that $\sum_{i=1}^n ik_i = n$. Then, from Eq. (11), we find for any n ,

$$\langle \tilde{\tau}_{\infty}^n \rangle = \frac{\int_0^{\mathcal{S}} \tau^n F_{\infty}(\tau) d\tau}{\int_0^{\mathcal{S}} F_{\infty}(\tau) d\tau} \sim \sqrt{\frac{2}{\pi(2n-1)^2}} \mathcal{S}^{n-1/2} \quad (18)$$

which implies that $\kappa_n(\tilde{\tau}_{\infty}) \sim \langle \tilde{\tau}_{\infty}^n \rangle$. Finally, we obtain that for any vector (k_1, \dots, k_n) of S_n , using that N_S is scale-invariant from Eq. (8) such that, for any m , $\kappa_m(N_S) \propto \mathcal{S}^m$,

$$\begin{aligned} \kappa_{\sum_{i=1}^n k_i}(N_S) \kappa_1(\tilde{\tau}_{\infty})^{k_1} \dots \kappa_n(\tilde{\tau}_{\infty})^{k_n} &\propto \left(\sqrt{\mathcal{S}}\right)^{\sum_{i=1}^n k_i} \mathcal{S}^{k_1(1-1/2)} \dots \mathcal{S}^{k_n(n-1/2)} \\ &\propto \mathcal{S}^{\sum_{i=1}^n ik_i} = \mathcal{S}^n \end{aligned} \quad (19)$$

All the terms in the sum of Eq. (17) scale as \mathcal{S}^n . This leads to the following scaling with \mathcal{S} for all the cumulants of the lifetime,

$$\kappa_n(T_S) \propto \mathcal{S}^n . \quad (20)$$

In particular, we deduce that the lifetime is scale-invariant, and scales linearly with the metabolic time.

Several important properties stem from the analysis of the 1d starving RW:

1. The number of sites visited at starvation grows diffusively with the metabolic time \mathcal{S} , $N_S \propto \sqrt{\mathcal{S}}$.
2. The average lifetime grows linearly with the metabolic time \mathcal{S} , $T_S \propto \mathcal{S}$.
3. The distribution of the number of sites visited at starvation and the lifetime are scale-invariant, in the sense that $N_S/\sqrt{\mathcal{S}}$ and T_S/\mathcal{S} are non-

deterministic and independent of \mathcal{S} for large \mathcal{S} . Additionally, the distribution of $N_{\mathcal{S}}$ decays exponentially at large number of distinct sites visited.

These properties are in agreement with what could be expected from the recurrence of the 1D RW (see Sec. 2.1): because the RW is recurrent, the starving process stops rapidly, as the RW often comes back to previously visited sites. Besides, we expect the RW not to find a lot of food units before starving. These two observations are in agreement with the linearity of the lifetime with the metabolic time, which is the minimal dependence with \mathcal{S} (as the RW performs at least \mathcal{S} steps before starving). The number of sites visited at starvation corresponds to the domain visited during this short lifetime (see Eq. (1) for $\mu = 1/2$), $N_{\mathcal{S}} \propto \sqrt{T_{\mathcal{S}}} \propto \sqrt{\mathcal{S}}$. One might wonder whether these results hold when increasing the dimension. As a first example, the other limit case of infinite dimension was considered in [10, 14].

3.2 The infinite dimensional case

In this section, we consider the starving RW in a medium of large dimension d . A first remark is that in the case of a nearest-neighbor RW in dimension d , the constant prefactor in (1) in the transient case can be shown to go to 1 as the dimension increases (see Refs. [33, 5]). This implies that the RW visits a new site at almost every time step, and thus the effect of depletion is small. Consequently, for the RW not visiting a new site, the most likely scenario is to repeat the previous step, returning to its previous position, which occurs with probability $1/(2d)$, a small value. This justifies our assumption that at every time step, the RW visits a new site with probability λ , independent of time, a value that diminishes in larger dimensions (where λ can be interpreted as a generalization of $1/(2d)$ to a RW in a large dimensional space). Additionally, visitation events are assumed to be independent: they occur rarely, and the visited domain contains almost no loops, as it seldom retraces its past trajectory.

These approximations lead to the effective starving RW model represented in Fig. 2 (see Ref. [14]): the effective RW starts at position \mathcal{S} , takes one step closer to 0 with probability λ , and returns to \mathcal{S} with probability $1 - \lambda$ (signifying the finding of a food unit). Similar to the $1d$ case, the derivation begins with determining the number of sites visited at starvation, followed by determining the lifetime.

3.2.1 Number of sites visited at starvation

As a first step, we note that Eq. (5) holds, as one can reexpress the probability of visiting at least n sites before starving via the statistics of the τ_k , specifically the probability that each of the τ_k is smaller than \mathcal{S} for $k = 0, \dots, n - 1$. The probability that $\tau_k \geq \mathcal{S}$ is exactly given by the probability that the effective RW, represented in Fig. 2, arrives at 0 before returning to \mathcal{S} , starting from \mathcal{S} (since the RW just visited




Fig. 2 Effective walk representing the starving RW in infinite dimension. The effective RW starts at position S , takes one step closer to 0 with probability λ .

a new site). This probability, denoted as $1 - P_S \equiv \mathbb{P}(\tau_k \geq S)$, is independent of k and is given by $1 - P_S = \lambda^S$, the probability that none of the S steps lead back to S (this scaling is in agreement with the $\mu \rightarrow \infty$ case of Table 1). Particularly, we note that the τ_k are again independent, as the state at which any visit starts is the same (the effective RW starts at S).

Using Eq. (5), we obtain that:

$$\mathbb{P}(N_S \geq n) = \prod_{k=0}^{n-1} \mathbb{P}(\tau_k < S) = P_S^n = (1 - \lambda^S)^n \sim e^{-n\lambda^S} \quad (21)$$

in the limit of large n and small λ . In particular, we note that the distribution of N_S is exponential with a rate of λ^S , such that:

$$\langle N_S \rangle \sim 1/\lambda^S. \quad (22)$$

Finally, we notice that the distribution of $N_S/\langle N_S \rangle = N_S\lambda^S$ is again independent of S , as in the $1d$ case, see Eq. (8).

3.2.2 Lifetime

Next, we determine the distribution of the lifetime T_S . To do so, we first compute the probability $F(n, t)$ for the effective RW to arrive at 0 for the first time at step t starting from position n . This distribution obeys a recursive relation,

$$F(0, t) = \delta_t, \quad F(n, t) = \lambda F(n-1, t-1) + (1-\lambda)F(S, t-1), \quad (23)$$

where δ_t is the Kronecker symbol (1 if $t = 0$, 0 otherwise). To solve this recursive equation, we Laplace transform Eq. (23), defined for any n and s by $\hat{F}(n, s) = \mathcal{L}(F(n, t)) \equiv \sum_{t=0}^{\infty} F(n, t)e^{-st}$. This leads to the following equation for $\hat{F}(n, s)$:

$$\hat{F}(0, s) = 1, \quad e^{-s} \hat{F}(n, s) = \lambda \hat{F}(n-1, s) + (1-\lambda) \hat{F}(\mathcal{S}, s). \quad (24)$$

The solution is given by:

$$\hat{F}(n, s) = (\lambda e^s)^n \left(1 - \frac{e^s(1-\lambda)}{1-\lambda e^s} \hat{F}(\mathcal{S}, s) \right) + \frac{e^s(1-\lambda)}{1-\lambda e^s} \hat{F}(\mathcal{S}, s). \quad (25)$$

In particular, Eq. (25) gives the average lifetime expression, by noting that $\mathbb{P}(T_{\mathcal{S}} = t) = F(\mathcal{S}, t)$:

$$\langle T_{\mathcal{S}} \rangle = \sum_{t=0}^{\infty} t F(\mathcal{S}, t) = -\partial_s \hat{F}(\mathcal{S}, s)_{s=0} = \lambda^{-\mathcal{S}} \left(\frac{1-\lambda^{\mathcal{S}}}{1-\lambda} \right) \sim 1/\lambda^{\mathcal{S}}, \quad (26)$$

which shows that the average lifetime grows exponentially with the metabolic time \mathcal{S} . Additionally, we note that in the limit $s/\lambda^{\mathcal{S}}$ fixed, small λ and large metabolic times,

$$\mathcal{L}(\mathbb{P}(T_{\mathcal{S}} = t)) \sim \frac{\lambda^{\mathcal{S}}}{s + \lambda^{\mathcal{S}}} \quad (27)$$

so that:

$$\mathbb{P}(T_{\mathcal{S}} = t) \sim \lambda^{\mathcal{S}} e^{-t\lambda^{\mathcal{S}}} = \frac{1}{\langle T_{\mathcal{S}} \rangle} e^{-t/\langle T_{\mathcal{S}} \rangle} \quad (28)$$

which is exponential, and scale-invariant in the sense that $T_{\mathcal{S}}/\langle T_{\mathcal{S}} \rangle$ is a non-deterministic random variable independent of \mathcal{S} .

Several important properties stem from the analysis of the infinite dimensional starving RW:

1. The number of sites visited at starvation grows exponentially with the metabolic time \mathcal{S} .
2. The average lifetime grows exponentially with the metabolic time \mathcal{S} .
3. The distribution of both the number of sites visited at starvation and the lifetime are scale-invariant, in the sense that $N_{\mathcal{S}}\lambda^{\mathcal{S}}$ and $T_{\mathcal{S}}\lambda^{\mathcal{S}}$ are independent of \mathcal{S} for large \mathcal{S} . Additionally, both asymptotic distributions of the rescaled variables decay exponentially at large argument.

These observations are in agreement with what could be expected for a transient RW: it rarely comes back to previously visited sites, such that the starving process lasts for a long time and finds a lot of sites before starving. This is in agreement with the exponential growth with the metabolic time of both the lifetime and the number of sites visited.

Let us now bridge the gap between the one and infinite dimensional cases.

4 General d -dimensional case: the maximum of inter-visit times

As highlighted in the calculations of the unidimensional and infinite dimensional cases, the derivation of the starving RWs' main observables starts with the determination of the probability that the n first intervisit times are smaller than the metabolic time \mathcal{S} , as in Eqs. (5) and (21). The probability of this event can be obtained from the statistics of the maximum $M_n \equiv \max_{k=0, \dots, n-1} \tau_k$:

$$\{\tau_k < \mathcal{S}, k = 0, \dots, n-1\} = \left\{ \max_{k=0, \dots, n-1} \tau_k < \mathcal{S} \right\} = \{M_n < \mathcal{S}\}. \quad (29)$$

However, the τ_k are a priori correlated in the general case: indeed, even though the RW is Markovian, values of τ_k can be representative of different geometries of the visited domain. In the following, we neglect these correlations, because we expect them to be too weak to affect the statistics of the maximum (see [30, 34] for a quantitative argument). This is the reason why we make use of the following approximate expression for the maximum's cumulative distribution function:

$$\mathbb{P}(M_n \leq T) \approx \prod_{k=0}^{n-1} \mathbb{P}(\tau_k \leq T) \sim \exp \left[- \sum_{k=0}^{n-1} \int_T^\infty F_k(\tau) d\tau \right]. \quad (30)$$

4.1 Recurrent walks

For recurrent walks ($\mu < 1$), the probability distribution function of τ_k presents a scaling form (see Eq. (2)), where the scaling function ψ is algebraic at small argument and exponential at large argument (see Table 1). Plugging this information into Eq. (30) implies that the cumulative distribution function of M_n also has a scaling form:

$$\begin{aligned} -\ln \mathbb{P}(M_n \leq T) &\approx \sum_{k=0}^{n-1} \int_T^\infty k^{-1-1/\mu} \psi(\tau/k^{1/\mu}) d\tau \\ &\sim \int_0^{n/T^\mu} \frac{dv}{v} \int_{v^{-1/\mu}}^\infty \psi(u) du = -\ln \Xi_\mu(T/n^{1/\mu}). \end{aligned} \quad (31)$$

Let us now consider the limit behaviours of $\Xi_\mu(m)$ when $m \equiv T/n^{1/\mu} \ll 1$ and $m \gg 1$.

First, in the limit $m \gg 1$, using Table 1,

$$-\ln \Xi_\mu(m) \propto \int_0^{1/m^\mu} \frac{dx'}{x'} e^{-A/x'^{1/\mu}} \propto \int_m^\infty \frac{dx'}{x'} e^{-Ax'} \propto e^{-Am}/m. \quad (32)$$

Then, in the limit $m \ll 1$

$$-\ln \Xi_\mu(m) \propto \text{const.} + \int_1^{1/m^\mu} \frac{dx'}{x'} x' \propto m^{-\mu}. \quad (33)$$

We will see the implications of these scalings on the starving RW problem in Sec. 5.

4.2 Marginal walks

Based on the functional form (3) for $F_k(\tau)$, we expect M_n/\sqrt{n} to converge to a non-trivial distribution at large n . Indeed, for $m > 0$ fixed and $n \rightarrow \infty$, we have that:

$$\begin{aligned} -\ln \mathbb{P}(M_n \leq m\sqrt{n}) &\approx \int_0^n dk \int_{m\sqrt{n}}^\infty d\tau F_k(\tau) \\ &= \int_0^{m^{2/3}n^{1/3}} dk \int_{m\sqrt{n}}^\infty d\tau \exp[-\text{const.} \tau/k] + \int_{m^{2/3}n^{1/3}}^n \frac{dk}{k} \int_{m\sqrt{n}}^\infty \frac{d\tau}{\sqrt{k}} \phi\left(\frac{\tau}{\sqrt{k}}\right) \\ &\sim \int_0^{1/m^2} \frac{du}{u} \int_{1/\sqrt{u}}^\infty dt \phi(t) = -\ln \Xi_1(m). \end{aligned} \quad (34)$$

From this, it is apparent that M_n/\sqrt{n} has an asymptotic limit distribution, denoted by Ξ_1 . The properties of $\Xi_1(m)$ at small and large m are directly derived from the properties of ϕ introduced in Eq. (3) and following similar steps as Eqs. (32) and (33):

$$-\ln \Xi_1(m) \propto \begin{cases} m^{-2} & \text{for } m \ll 1 \\ e^{-A\sqrt{m}} & \text{for } m \gg 1 \end{cases} \quad (35)$$

where A is a process dependent constant and all algebraic prefactors have been neglected in the $m \gg 1$ regime.

4.3 Transient walks

For transient walks, $\mu > 1$, according to Table 1, the decrease of the distribution of τ_k is first exponential in $\tau^{\mu/(1+\mu)}$ (stretched exponential in τ of exponent $\delta = \mu/(1+\mu)$), a regime which lasts up to time $T_k = k^{1+1/\mu}$, and next exponential in $\tau/k^{1/\mu}$. On the other hand, the maximum M_n of n random variables with such stretched exponential tail (but no exponential tail) has an average which grows as $\langle M_n \rangle \sim a (\ln n)^{1/\delta} = a (\ln n)^{1+1/\mu}$ (a a model-dependent constant) and a standard deviation which grows as $\sqrt{\text{Var}(M_n)} = \sqrt{\langle M_n^2 \rangle - \langle M_n \rangle^2} \propto (\ln n)^{1/\delta-1} = (\ln n)^{1/\mu}$, as shown in Ref. [20]. In particular, this means that the maximum is peaked around its mean (standard deviation much smaller than its mean) which is itself small compared

to the time $n^{1+1/\mu}$ (typical value of T_k for k between 0 and $n-1$) at which the stretched exponential regime would stop. We conclude that the asymptotic distribution of the maximum M_n of the τ_k is indeed not influenced by the exponential decay at large times. Thus, the limit law (centered and normalized) of M_n corresponds to the celebrated Gumbel distribution [35] of zero mean and unit variance of cumulative distribution function we note Ξ_∞ . For any value of μ , the cumulative distribution function of M_n centered and normalized is given by:

$$-\ln \Xi_\mu(m) = -\ln \Xi_\infty(m) = e^{-me^{-\gamma_E}} \quad (36)$$

where γ_E is the Euler constant and the rescaled random variable m is defined in Eq. (37).

➤ Important

By considering the rescaled random variable (a a model-dependent constant)

$$m \equiv \begin{cases} M_n/n^{1/\mu} & [\mu < 1] \\ M_n/\sqrt{n} & [\mu = 1] \\ \frac{M_n - a(\ln n)^{1/\mu+1}}{(\ln n)^{1/\mu}} & [\mu > 1] \end{cases} \quad (37)$$

we obtain its asymptotic cumulative distribution Ξ_μ , whose behavior at large and small arguments is summarized in Table 2.

Table 2 Asymptotic properties of the cumulative distribution Ξ_μ of the properly rescaled maximum for different types of processes. All constants have been put to one for simplicity.

	$m \ll 1$	$m \gg 1$
$-\ln \Xi_\mu(m)$		
$\mu < 1$ [recurrent]	$m^{-\mu}$	e^{-m}/m
$\mu = 1$ [marginal]	m^{-2}	$e^{-\sqrt{m}}$
$\mu > 1$ [transient]	e^{-m}	e^{-m}

5 Starving Random Walks

We now show that the knowledge of the cumulative distribution function of M_n allows one to determine the statistics of the number of sites visited at starvation and the lifetime of a starving RW.

5.1 Number of sites visited at starvation

We start from the remark that having M_n smaller than the metabolic time \mathcal{S} is the same as having visited at least n sites before starvation, thus

$$\mathbb{P}(N_{\mathcal{S}} \geq n) = \mathbb{P}(M_n < \mathcal{S}) . \quad (38)$$

This means that one can deduce the distribution of $N_{\mathcal{S}}$ directly from that of M_n .

- For recurrent RWs: The distribution of $N_{\mathcal{S}}$ is scale-invariant (so that $N_{\mathcal{S}}/\mathcal{S}^\mu$ is independent of \mathcal{S}) and its asymptotics is given by:

$$-\ln \mathbb{P}(N_{\mathcal{S}}/\mathcal{S}^\mu \geq x) = -\ln \Xi_\mu(x^{-1/\mu}) \propto \begin{cases} e^{-A/x^{1/\mu}} x^{1/\mu} & \text{for } x \ll 1 \\ x & \text{for } x \gg 1. \end{cases} \quad (39)$$

Of note, this result is consistent with what we found in Sec. 3.1, Eqs. (7), (12) corresponding to $\mu = 1/2$.

- For marginal RWs: As $\mathbb{P}(N_{\mathcal{S}=\sqrt{n}x} \geq n) = \mathbb{P}(M_n \leq \sqrt{nx}) \rightarrow \Xi_1(x)$, with $\Xi_1(x)$ of Eq. (35), we get that $\mathbb{P}(N_{\mathcal{S}}/\mathcal{S}^2 \geq x) \rightarrow \Xi_1(1/\sqrt{x})$. In particular, the distribution has a single-parameter scaling at large starvation index \mathcal{S} and number of visited sites at starvation $N_{\mathcal{S}}$, and its average and standard deviation scale both as \mathcal{S}^2 . Based on Eq. (35), up to logarithmic corrections,

$$-\ln \mathbb{P}(N_{\mathcal{S}}/\mathcal{S}^2 \geq x) \propto \begin{cases} e^{-A/x^{1/4}} & \text{for } x \ll 1 \\ x & \text{for } x \gg 1. \end{cases} \quad (40)$$

- For transient RWs, considering $\langle M_n \rangle \sim a(\ln n)^{\frac{1+\mu}{\mu}}$ and $\sqrt{\text{Var}(M_n)} \sim b(\ln n)^{\frac{1}{\mu}} \equiv \frac{a(1+\mu)\sqrt{6}}{\pi\mu}(\ln n)^{\frac{1}{\mu}}$ [20], the probability $\mathbb{P}\left(\frac{M_n - a(\ln n)^{1/\mu+1}}{b(\ln n)^{1/\mu}} \leq x\right)$ converges to a Gumbel distribution $\Xi_\infty(x)$ with zero mean and unit variance. Consequently, $\mathbb{P}(N_{\mathcal{S}=xb(\ln n)^{1/\mu} + a(\ln n)^{1/\mu+1}} \geq n)$ also converges to $\Xi_\infty(x)$ for large n . By inverting $\mathcal{S} = xb(\ln n)^{1/\mu} + a(\ln n)^{1/\mu+1}$ in the limit of large n , we obtain:

$$\mathbb{P}(N_{\mathcal{S}}/\exp\left[(\mathcal{S}/a)^{\mu/(1+\mu)}\right] \geq x) \sim \Xi_\infty\left(-\frac{\sqrt{6}}{\pi} \ln x\right) = \exp[-xe^{-\gamma E}] . \quad (41)$$

This implies that $N_{\mathcal{S}}$ asymptotically follows a single-parameter scaling: both its average and standard deviation grow exponentially with $\mathcal{S}^{\mu/(1+\mu)}$. Moreover, the asymptotic distribution of $N_{\mathcal{S}}$ is exponential, akin to the infinite-dimensional case (refer to Eq. (21)). Finally, by taking the limit $\mu \rightarrow \infty$ in the expression for the average of $N_{\mathcal{S}}$, we recover the exponential growth of the average number of visited sites at starvation with the metabolic time, as obtained for the infinite-dimensional case in Eq. (22).

The average, standard deviation and asymptotic distribution of N_S in the recurrent, marginal and transient cases are checked numerically in Fig. 3 for representative RWs of each class.




Fig. 3 Number of sites visited at starvation N_S . (a)-(c) Distributions of the rescaled variable $x \equiv N_S / \langle N_S \rangle$ (insets illustrate the small x -values) and (a')-(c') the corresponding averages (red circles) and standard deviations (blue squares) of N_S . The black dashed lines correspond to the best fit of the theory (see Eqs. (39), (40), (41), and (42)). Different universality classes are represented by (a) $1d$ Lévy flights of parameter $\alpha = 1.5$ (recurrent), $S = 14667, 31622$ and 68129 ; (b) nearest neighbour $2d$ RWs, $\mu = 1$ (marginal), $S = 2335, 4832$ and 10^4 ; (c) nearest neighbour $3d$ RWs, $\mu = 3/2$ (transient), $S = 12, 17$ and 22 . Increasing values of S are represented by blue circles, orange stars, and green squares.

➤ Important

By considering the rescaled number of sites visited at starvation $\frac{N_S}{\langle N_S \rangle}$, where the asymptotics of $\langle N_S \rangle$ is given by

$$\langle N_S \rangle \sim \begin{cases} S^\mu & [\mu < 1] \\ S^2 & [\mu = 1] \\ \exp[S^{\mu/(1+\mu)}] & [\mu > 1], \end{cases} \quad (42)$$

we obtain its asymptotic tail distribution $\mathbb{P}\left(\frac{N_S}{\langle N_S \rangle} \geq x\right) = \Phi_\mu(x)$, whose behavior at large and small arguments is summarized in Table 3.

Table 3 Asymptotic properties of the minus log tail distribution Φ_μ of the rescaled number of sites visited at starvation for different type of processes. All constants have been put to one for simplicity.

$-\ln \Phi_\mu(x)$	$x \ll 1$	$x \gg 1$
$\mu < 1$ [recurrent]	$e^{-1/x^{1/\mu}} / x^{1/\mu}$	x
$\mu = 1$ [marginal]	$e^{-1/x^{1/4}}$	x
$\mu > 1$ [transient]	x	x

5.2 Lifetime

We proceed by demonstrating that the distribution of N_S gives access to the distribution of the lifetime T_S .

The lifetime T_S is determined by the accumulation of inter-visit times $\tilde{\tau}_k$ ($k < N_S$), each corresponding to τ_k conditioned on being less than the metabolic time S . Expressing the distribution of T_S yields:

$$\begin{aligned}
\mathbb{P}(T_S = t) &= \int_0^\infty dn \mathbb{P} \left(\sum_{k=0}^{N_S-1} \tau_k + S = t | N_S = n \right) \mathbb{P}(N_S = n) \\
&\approx \int_0^\infty dn \delta(n \langle \tilde{\tau}_\infty \rangle + S - t) \mathbb{P}(N_S = n).
\end{aligned} \tag{43}$$

In this derivation, we exploit the fact that for large k , the distribution of $\tilde{\tau}_k$ becomes independent of k , and the sum follows the law of large numbers, $\sum_{k=0}^{n-1} \langle \tilde{\tau}_k \rangle \sim n \lim_{k \rightarrow \infty} \langle \tilde{\tau}_k \rangle = n \langle \tilde{\tau}_\infty \rangle$. This leads to the tail distribution:

$$\mathbb{P}(T_S \geq t) \approx \mathbb{P} \left(N_S \geq \frac{t - S}{\langle \tilde{\tau}_\infty \rangle} \right). \tag{44}$$

Then, we proceed with the characterization of $\tilde{\tau}_\infty$, which varies for each type of RW.

- For recurrent RWs: At early times, the distribution $F_k(\tau)$ of inter-visit time τ_k behaves as $F_k(\tau) \propto \tau^{-1-\mu}$ (see Table.1). Consequently, the average value of $\tilde{\tau}_\infty$ scales as:

$$\langle \tilde{\tau}_\infty \rangle \propto \frac{\sum_{\tau=1}^S \tau^{-\mu}}{\sum_{\tau=1}^S \tau^{-1-\mu}} \sim c S^{1-\mu} \tag{45}$$

where c is a model-dependent constant. Since $0 < \mu < 1$, the scaling $S^{1-\mu}$ arises from the numerator. By rescaling T_S by S and using Eq. (39), we find:

$$\mathbb{P}(T_S/S \geq t) = \mathbb{P}(N_S \geq c(t-1)S^\mu) = \Xi_\mu((c(t-1))^{-1/\mu}) \tag{46}$$

where Ξ_μ is defined in Eq. (31). This reveals that T_S scales with S irrespective of the value of the exponent $\mu < 1$, generalizing the $1d$ scenario (see Eq. (13)).

- For marginal RWs: Analogously to recurrent RWs, the average value of $\tilde{\tau}_\infty$ is given by:

$$\langle \tilde{\tau}_\infty \rangle \propto \frac{\sum_{\tau=1}^S \tau^{-1}}{\sum_{\tau=1}^S \tau^{-2}} \sim c \ln S . \quad (47)$$

Hence, by scaling T_S by $S^2 \ln S$, we obtain:

$$\mathbb{P}(T_S / (S^2 \ln S) \geq t) = \mathbb{P}(N_S \geq t S^2 / c) = \Xi_1(\sqrt{c/t}) \quad (48)$$

resulting in a scaling form for the distribution of T_S .

- For transient RWs: In this case, the inter-visit times' distribution has finite moments, leading to $\langle \tilde{\tau}_\infty \rangle \rightarrow \langle \tau_\infty \rangle = c$. By rescaling T_S by $\exp[(S/a)^{\mu/(1+\mu)}]$, we derive:

$$\begin{aligned} \mathbb{P}\left(T_S / \exp\left[(S/a)^{\mu/(1+\mu)}\right] \geq t\right) &\sim \mathbb{P}\left(N_S \geq t \exp\left[(S/a)^{\mu/(1+\mu)}\right] / c\right) \\ &\sim \Xi_\infty\left(-\frac{\sqrt{6}}{\pi} \ln(t/c)\right) = \exp\left[-\frac{t}{c} e^{-\gamma E}\right] \end{aligned} \quad (49)$$

where a is the same constant as in Eqs. (37) and (41). The lifetime T_S follows an exponential distribution, akin to the number N_S of sites visited at starvation, as observed in the infinite-dimensional scenario (see Eq. (26)).

The average, standard deviation and asymptotic distribution of T_S in the recurrent, marginal and transient cases are checked numerically in Fig. 4 for representative RWs of each class.

➤ Important

By considering the rescaled lifetime $\frac{T_S}{\langle T_S \rangle}$, where the asymptotic of $\langle T_S \rangle$ is given by

$$\langle T_S \rangle \sim \begin{cases} S & [\mu < 1] \\ S^2 & [\mu = 1] \\ \exp[S^{\mu/(1+\mu)}] & [\mu > 1] , \end{cases} \quad (50)$$

we obtain the asymptotic tail distribution $\mathbb{P}\left(\frac{T_S}{\langle T_S \rangle} \geq t\right) = \Psi_\mu(t)$, which has the properties at large and small arguments summarized in Table 4.




Fig. 4 Lifetime in Starving Random Walks. (a)-(c) Distributions of the rescaled variable $t \equiv T_S / \langle T_S \rangle$ (insets illustrate behavior at small t -values), and (a')-(c') corresponding averages (red circles) and standard deviations (blue squares) of T_S . The black dashed lines denote the best fit from theory (see Eqs. (46), (48), (49), and (50)). Various universality classes are depicted: (a) 1d Lévy flights with parameter $\alpha = 1.5$ (recurrent), $S = 14667, 31622$, and 68129 ; (b) nearest-neighbor 2d random walks, $\mu = 1$ (marginal), $S = 2335, 4832$, and 10^4 ; (c) nearest-neighbor 3d random walks, $\mu = 3/2$ (transient), $S = 12, 17$, and 22 . Increasing values of S are denoted by blue circles, orange stars, and green squares.

Table 4 Asymptotic properties of the minus log tail distribution Ψ_μ of the rescaled lifetime for different type of processes. All constants have been put to one for simplicity.

$-\ln \Psi_\mu(t)$	$x \ll 1$	$x \gg 1$
$\mu < 1$ [recurrent]	$e^{-1/t^{1/\mu}} / t^{1/\mu}$	t
$\mu = 1$ [marginal]	$e^{-1/t^{1/4}}$	t
$\mu > 1$ [transient]	t	t

5.3 Discussion

In this section, we finally bridged the gap between the unidimensional and infinite dimensional starving RWs studied in Sec. 3.

Several important properties stem from the analysis of the general Markovian starving RW:

1. The number of sites visited at starvation grows algebraically with the metabolic time \mathcal{S} for recurrent ($N_S \propto \mathcal{S}^\mu$) and marginal ($\propto \mathcal{S}^2$) RWs, while it grows significantly faster for transient RWs ($\propto \exp \left[\mathcal{S}^{\frac{\mu}{1+\mu}} \right]$).
2. The average lifetime grows linearly with \mathcal{S} for recurrent RWs ($T_S \propto \mathcal{S}$), while it grows quadratically for marginal RWs ($\propto \mathcal{S}^2$) and stretched exponentially for transient RWs ($\propto \exp \left[\mathcal{S}^{\frac{\mu}{1+\mu}} \right]$).

3. The distribution of both the number of sites visited at starvation and the lifetime are scale-invariant, in the sense that $N_S/\langle N_S \rangle$ and $T_S/\langle T_S \rangle$ are independent of S for large S . Additionally, both asymptotic distributions of the rescaled variables decay exponentially at large argument.

6 Conclusion

To conclude, we have shown how to derive key properties of the starving random walk (RW) problem, such as the number of sites visited at starvation (N_S) and the lifetime of the RW (T_S). We have demonstrated that understanding the universal properties of the time between visits (τ_k) leads to universal characteristics for these quantities for general symmetric Markovian RWs. We have highlighted important differences based on a single parameter μ , which characterizes recurrence or transience of RWs. For example, exploration in one or two dimensions (i.e., by recurrent walks) differs significantly from that in three dimensions (i.e., by transient walks), where the lifetime of a tracer with a specific metabolism time changes from algebraic (for recurrent walks) to stretched exponential (for transient walks).

This chapter's methods and findings serve two purposes:

First, they provide a basic framework for studying systems with limited resources, useful in fields like animal ecology [36] or the study of biological species in crowded environments [17]. In these situations, significant memory effects arise because the RW is influenced by its past path, which is affected by resource depletion.

Second, our results furnish a null model against which the exploration dynamics of biological tracers can be assessed. Departures from this model indicate the potential influence of intricate and realistic strategies not encompassed in this study, including the tracer's memory of its trajectory (using biological cues, such as ants or MDCK cells do [2, 1]), renewal of the resources with time [37] or the option for the tracer to abstain from consuming food when unnecessary [38]. Other extensions include the food detection at long distances [39, 40], penalized long moves [41], the intermittent motion [42]. The starving RW problem provides a relatively straightforward model yielding numerous insights and results, with the methods presented in this chapter laying the foundation for a comprehensive understanding of resource-limited motion in a broad range of systems.

References

1. J. d'Alessandro, A. Barbier-Chebbah, V. Cellerin, O. Bénichou, R.M. Mège, R. Voituriez, B. Ladoux, *Nat. Commun.* **12**(1), 4118 (2021)
2. D.M. Gordon, *Anim. Behav.* **49**(3), 649 (1995)
3. A. Dvoretzky, P. Erdős, in *Proc. Second Berkeley Symposium* (1951), pp. 353–367

4. N.C. Jain, W.E. Pruitt, in *Proc. Sixth Berkeley Symposium*, vol. 3 (1972), vol. 3, pp. 31–50
5. B.D. Hughes, *Random Walks and Random Environments* (Oxford University Press, New York, 1995)
6. J.E. Gillis, G.H. Weiss, *J. Math. Phys.* **11**(4), 1307 (1970)
7. J.F. Le Gall, J. Rosen, *Ann. Probab.* pp. 650–705 (1991)
8. L. Régnier, M. Dolgushev, S. Redner, O. Bénichou, *Phys. Rev. E* **105**, 064104 (2022)
9. G.H. Vineyard, *J. Math. Phys.* **4**(9), 1191 (1963)
10. O. Bénichou, S. Redner, *Phys. Rev. Lett.* **113**, 238101 (2014)
11. L. Régnier, M. Dolgushev, S. Redner, O. Bénichou, *Nat. Commun.* **14**(1), 618 (2023)
12. L. Régnier, O. Bénichou, P.L. Krapivsky, *Phys. Rev. Lett.* **130**, 227101 (2023)
13. L. Régnier, M. Dolgushev, O. Bénichou, *Phys. Rev. E* **109**, 064101 (2024)
14. O. Bénichou, M. Chupeau, S. Redner, *J. Phys. A: Math. Theor.* **49**(39), 394003 (2016)
15. D.R. Sowinski, J. Carroll-Nellenback, R.N. Markwick, J. Piñero, M. Gleiser, A. Kolchinsky, G. Ghoshal, A. Frank, *PRX Life* **1**, 023003 (2023)
16. R. Klages, in *Diffusive Spreading in Nature, Technology and Society* (Springer, 2023), pp. 53–74
17. K.M. Passino, in *Innovations and Developments of Swarm Intelligence Applications* (IGI Global, 2012), pp. 219–234
18. C.G. Orlando, A. Tews, P. Banks, C. McArthur, *Biol. Lett.* **16**(7), 20200329 (2020)
19. A.F. Winfield, *Distributed autonomous robotic systems* 8 pp. 185–192 (2009)
20. S.N. Majumdar, A. Pal, G. Schehr, *Phys. Rep.* **840**, 1 (2020)
21. S.N. Majumdar, G. Schehr, *J. Stat. Mech.* **2014**(1), P01012 (2014)
22. S.N. Majumdar, P.L. Krapivsky, *Phys. Rev. E* **62**(6), 7735 (2000)
23. E. Ben-Naim, P.L. Krapivsky, S.N. Majumdar, *Phys. Rev. E* **64**(3), 035101 (2001)
24. T. Kay, L. Giuggioli, *J. Phys. A: Math. Theor.* **56**(34), 345002 (2023)
25. D. Ben-Avraham, S. Havlin, *Diffusion and reactions in fractals and disordered systems* (Cambridge university Press, 2000)
26. G. Pólya, *Mathematische Annalen* **84**(1-2), 149 (1921)
27. L.A. Shepp, *Bull. Amer. Math. Soc.* **70**(6), 540 (1964)
28. J. Bäuml, *Electron. J. Probab.* **28**, 1 (2023)
29. A. Mariz, F. Van Wijland, H. Hilhorst, S. Gomes Júnior, C. Tsallis, *J. Stat. Phys.* **102**, 259 (2001)
30. L. Régnier, M. Dolgushev, O. Bénichou, *Phys. Rev. Lett.* **132**, 127101 (2024)
31. S. Redner, *A Guide to First-Passage Processes* (Cambridge University Press, Cambridge, UK, 2001)
32. N. Niki, N. Shigekazu, H. Inoue, *Commun. Stat. Theory Methods* **19**(5), 1857 (1990)
33. G.N. Watson, *Quart. J. Math. Oxford* **10**, 266 (1939)
34. D. Carpentier, P. Le Doussal, *Phys. Rev. E* **63**(2), 026110 (2001)
35. E.J. Gumbel, *Statistics of extremes* (Columbia university press, 1958)
36. C.G. Orlando, A. Tews, P. Banks, C. McArthur, *Biol. Lett.* **16**(7), 20200329 (2020)
37. M. Chupeau, O. Bénichou, S. Redner, *Phys. Rev. E* **93**(3), 032403 (2016)
38. O. Bénichou, U. Bhat, P.L. Krapivsky, S. Redner, *Phys. Rev. E* **97**, 022110 (2018)
39. H. Sanhedrai, Y. Maayan, L.M. Shekhtman, *Europhys. Lett.* **128**(6), 60003 (2020)
40. H. Sanhedrai, Y. Maayan, *Phys. Rev. E* **103**, 012114 (2021)
41. N. Krishnan, Z.P. Kilpatrick, *Phys. Rev. E* **98**, 052406 (2018)
42. D. Campos, J. Cristín, V. Méndez, *Phys. Rev. E* **103**(5), 052109 (2021)