# (u, v)-ABSORBING (PRIME) HYPERIDEALS IN COMMUTATIVE MULTIPLICATIVE HYPERRINGS

MAHDI ANBARLOEI

ABSTRACT. In this paper, we will introduce the notion of (u, v)-absorbing hyperideals in multiplicative hyperrings and we will show some properties of them. Then we extend this concept to the notion of (u, v)-absorbing prime hyperideals and then we will give some results about them.

### 1. INTRODUCTION

In 1934 [25], the French mathematician F. Marty presented the notion of hyperstructures or multioperations in which an operation yields a set of values rather than a single value. Many authors have contributed to the advancement of this novel area of modern algebra [12, 13, 14, 15, 16, 26, 30, 36]. An importan class of the algebraic hyperstructures called the multiplicative hyperring was defined by Rota in 1982 [32]. In this hyperstructure, the multiplication is a hyperoperation however the addition is an operation. Multiplicative hyperrings are deeply demonstrated and characterized in [3, 4, 5, 22, 28, 29, 31, 35]. Dasgupta studied the prime and primary hyperideals in multiplicative hyperrings in [17]. The notions of 1-absorbing and v-absorbing hyperideals as generalizations of the prime hyperideals were studied in [29] and [5]. The idea of (u, v)-closed hyperideals in a multiplicative hyperring was proposed in [6]. Our purpose of this research work is to introduce and study the notion of (u, v)-absorbing hyperideals as an extension of the *n*-absorbing hyperideals in a commutative multiplicative hyperring. Several specific properties are given to illustrate the structures of the new notion. Furthermore, we extend this concept to (u, v)-absorbing prime hyperideals.

### 2. Multiplicative hyperrings

In this section, we recall some background materials. A hyperoperation " $\circ$ " on non-empty set X is a mapping from  $X \times X$  into  $P^*(X)$  such that  $P^*(X)$  is the family of all non-empty subsets of X. In this case,  $(X, \circ)$  is called hypergroupoid. Let  $X_1, X_2$  be two subsets of X and  $x \in X$ , then  $X_1 \circ X_2 = \bigcup_{x_1 \in X_1, x_2 \in X_2} x_1 \circ x_2$ , and  $X_1 \circ x = X_1 \circ \{x\}$ . This means that the hyperoperation " $\circ$ " on X can be extended to subsets of X. A hypergroupoid  $(X, \circ)$  is called a semihypergroup if  $\bigcup_{a \in y \circ z} x \circ a = \bigcup_{b \in x \circ y} b \circ z$  for all  $x, y, z \in X$  which means  $\circ$  is associative. A semihypergroup X is called a hypergroup if  $x \circ X = X = X \circ x$  for each  $x \in X$ . A non-empty subset Y of a semihypergroup  $(X, \circ)$  refers to a subhypergroup if  $x \circ Y = Y = Y \circ x$  for each  $x \in Y$  [19].

<sup>2010</sup> Mathematics Subject Classification. Primary 20N20; Secondary 16Y99.

Key words and phrases. (u, v)-absorbing hyperideal, AB-(u, v)-absorbing hyperideal, (u, v)-absorbing prime hyperideal.

**Definition 2.1.** [19] An algebraic structure  $(A, +, \circ)$  refers to a commutative multiplicative hyperring if

- (1) (A, +) is a commutative group;
- (2)  $(A, \circ)$  is a semihypergroup;
- (3)  $x \circ (y+z) \subseteq x \circ y + x \circ z$  and  $(y+z) \circ x \subseteq y \circ x + z \circ x$  for every  $x, y, z \in A$ ;
- (4)  $x \circ (-y) = (-x) \circ y = -(x \circ y)$  for every  $x, y \in A$ ;
- (5)  $x \circ y = y \circ x$  for every  $x, y \in A$ .

If in (3), the equality holds then the multiplicative hyperring A is called strongly distributive.

Consider the ring of integers  $(\mathbb{Z}, +, \cdot)$ . For each subset  $T \in P^{\star}(\mathbb{Z})$  with  $|T| \geq 2$ , there exists a multiplicative hyperring  $(\mathbb{Z}_T, +, \circ)$  where  $\mathbb{Z}_T = \mathbb{Z}$  and  $a \circ b = \{a.x.b \mid x \in T\}$  for all  $a, b \in \mathbb{Z}_T$  [17].

**Definition 2.2.** [1] An element  $e \in A$  is considered as an identity element if  $a \in a \circ e$  for all  $a \in A$ .

Throughout this paper, A denotes a commutative multiplicative hyperring with identity 1.

**Definition 2.3.** [1] An element  $x \in A$  is called unit, if there exists  $y \in A$  such that  $1 \in x \circ y$ . Denote the set of all unit elements in A by U(A).

Moreover, a hyperring A refers to a hyperfield if each non-zero element of A is unit.

**Definition 2.4.** [19] A non-empty subset B of A is a hyperideal if

(i)  $x - y \in B$  for all  $x, y \in B$ ;

(ii)  $r \circ x \subseteq B$  for all  $x \in B$  and  $r \in A$ .

**Definition 2.5.** [17] A proper hyperideal B in A refers to a prime hyperideal if  $x \circ y \subseteq B$  for  $x, y \in A$ , then  $x \in B$  or  $y \in B$ .

The intersection of all prime hyperideals of A containing a hyperideal B is said to be the prime radical of B, denoted by rad(B). If the multiplicative hyperring Ahas no prime hyperideal containing B, we define rad(B) = A. Assume that C is the class of all finite products of elements of A that is  $C = \{a_1 \circ a_2 \circ \cdots \circ a_n \mid a_i \in A, n \in$  $\mathbb{N}\} \subseteq P^*(A)$  and B is a hyperideal of A. B refers to a C-hyperideal of A if for each  $C \in C$  and  $B \cap C \neq \emptyset$  imply  $C \subseteq B$ . Notice that  $\{a \in A \mid a^n \subseteq B \text{ for some } n \in$  $\mathbb{N}\} \subseteq rad(A)$ . The equality holds if B is a C-hyperideal of A (see Proposition 3.2 in [17]). Moreover, a hyperideal B of A refers to a strong C-hyperideal if for each  $D \in \mathfrak{U}$  and  $D \cap B \neq \emptyset$  imply  $D \subseteq B$  such that  $\mathfrak{U} = \{\sum_{i=1}^n C_i \mid C_i \in C, n \in \mathbb{N}\}$  and  $C = \{a_1 \circ a_2 \circ \ldots \circ a_n \mid a_i \in A, n \in \mathbb{N}\}$  (for more details see [18]).

**Definition 2.6.** [1] A proper hyperideal B of A is maximal in A if for each hyperideal M of A with  $B \subset M \subseteq A$ , then M = A.

The intersection of all maximal hyperideals of A is denoted by J(A). Also, A refers to a local multiplicative hyperring if it has just one maximal hyperideal.

**Definition 2.7.** [1] Assume that  $B_1$  and  $B_2$  are hyperideals of H. We define  $(B_2:B_1) = \{x \in A \mid x \circ B_1 \subseteq B_2\}.$ 

 $\mathbf{2}$ 

## 3. (u, v)-Absorbing hyperideals

A proper hyperideal Q in A refers to a v-absorbing hyperideal for  $v \in \mathbb{N}$  if whenever  $x_1 \circ \cdots \circ x_{v+1} \subseteq Q$  for  $x_1, \cdots, x_{v+1} \in A$ , then there exist v of the  $x_i$ 's whose product is a subset of Q. In this section, we aim to introduce the class of the (u, v)-absorbing hyperideals as an extension of the v-absorbing hyperideals and present some of their properties.

**Definition 3.1.** Let  $u, v \in \mathbb{N}$  with u > v. A proper hyperideal Q of A is called a (u, v)-absorbing hyperideal if whenever  $x_1 \circ \cdots \circ x_u \subseteq Q$  for  $x_1, \cdots, x_u \in A \setminus U(A)$ , then there are v of the  $x_i$ 's whose product is a subset of Q.

**Example 3.2.** Consider the multiplicative hyperring  $(\mathbb{Z}_T, +, \circ)$  where  $T = \{2, 4\}$ . Then  $Q = \langle 150 \rangle$  is a (4, 3)-absorbing hyperideal of the multiplicative hyperring  $(\mathbb{Z}_T, +, \circ)$ . However, it is not (3, 2)-absorbing as the fact that  $3 \circ 5^2 = \{300, 600, 1200\} \subseteq Q$  but  $3 \circ 5 = \{30, 60\} \notin Q$  and  $5^2 = \{50, 100\} \notin Q$ .

*Remark* 3.3. Assume that Q is a proper hyperideal of A and  $k, r, u, v \in \mathbb{N}$ .

- (i) Let Q be a (u, v)-absorbing hyperideal of A. Then:
  - (1) Q is a (k, r)-absorbing hyperideal for every  $k \ge u$  and  $r \ge v$ . (2) Q is a (u - 1)-absorbing hyperideal of A.
- (ii) Q is a (u, v)-absorbing hyperideal of A if and only if  $x_1 \circ \cdots \circ x_k \subseteq Q$  for  $x_1, \cdots, x_k \in A \setminus U(A)$  and  $k \ge u$  imply that there are v of the  $x_i$ 's whose product is a subset of Q.

**Proposition 3.4.** Let  $Q_i$  be a proper hyperideal of A and  $u_i, v_i \in \mathbb{N}$  for every  $i \in \{1, \dots, k\}$ . If  $Q_i$  is a  $(u_i, v_i)$ -absorbing hyperideal of A for every  $i \in \{1, \dots, k\}$ , then  $Q = \bigcap_{i=1}^k Q_i$  is a (u, v)-absorbing hyperideal of A where  $v = v_1 + \dots + v_k$  and  $u = \max\{u_1, \dots, u_k, v+1\}$ .

Proof. Assume that  $Q_i$  is a  $(u_i, v_i)$ -absorbing hyperideal of A for every  $i \in \{1, \dots, k\}$ . By Remark 3.3 (i), we conclude that  $Q_i$  is a  $(u, v_i)$ -absorbing hyperideal of A where  $v = v_1 + \dots + v_k$  and  $u = \max\{u_1, \dots, u_k, v+1\}$ . Let  $x_1 \circ \dots \circ x_u \subseteq Q$  for  $x_1, \dots, x_u \in A \setminus U(A)$ . Since  $Q_i$  is a  $(u, v_i)$ -absorbing hyperideal of A for every  $i \in \{1, \dots, k\}$  and  $x_1 \circ \dots \circ x_u \subseteq Q_i$ , we get the result that  $\prod_{\alpha \in I_i} x_\alpha \subseteq Q_i$  for each  $i \in \{1, \dots, k\}$  such that  $I_i \subseteq \{1, \dots, u\}$  and  $|I_i| = v_i$ . Put  $I = \bigcup_{i=1}^k I_i$ . Since  $\prod_{\alpha \in I} x_\alpha \subseteq Q$  and  $|I| \leq v$ , we conclude that  $Q = \bigcap_{i=1}^k Q_i$  is a (u, v)-absorbing hyperideal of A.

Let  $u, v \in \mathbb{N}$ . Then the smallest integer not less than  $\frac{u}{v}$  is denoted by  $\left\lfloor \frac{u}{v} \right\rfloor$ .

**Proposition 3.5.** Let Q be a (u, v)-absorbing C-hyperideal of A for  $u, v \in \mathbb{N}$  with u > v. Then:

(i)  $rad(Q) = \{x \in A \mid x^v \subseteq Q\}.$ 

(ii) rad(Q) is a (k, v)-absorbing hyperideal of A where  $k = \max\{\lfloor \frac{u}{v} \rfloor, v+1\}$ .

*Proof.* (i) Assume that Q is a (u, v)-absorbing hyperideal of A. Take any  $x \in rad(Q)$ . Then there exists a smallest element  $n \in \mathbb{N}$  such that  $x^n \subseteq Q$ . Let n > v. Remark 3.3(i)(2) shows that Q is a (u - 1)-absorbing hyperideal of A. Then we conclude that  $n \leq u - 1$  by Theorem 3.6 in [5]. This means  $x^u \subseteq Q$  which means  $x^v \subseteq Q$  as Q is a (u, v)-absorbing hyperideal of A. This is contradicts the fact that n is a smallest positive integer such that  $x^n \subseteq Q$ . Therefore  $n \leq v$  and  $x^v \subseteq Q$ .

(ii) Suppose that  $x_1 \circ \cdots \circ x_k \subseteq rad(Q)$  such that  $k = \max\{\lceil \frac{u}{v} \rceil, v+1\}$  and  $x_1, \cdots, x_k \in A \setminus U(A)$ . Then for any  $a \in x_1 \circ \cdots \circ x_k$ ,  $a^v \subseteq Q$ . Again,  $a^v \subseteq$ 

#### MAHDI ANBARLOEI

 $\begin{array}{l} (x_1 \circ \cdots \circ x_k)^v \subseteq x_1^v \circ \cdots \circ x_k^v \subseteq Q. \text{ So, } (x_1^v \circ \cdots \circ x_k^v) \cap Q \neq \varnothing \text{ and then } x_1^v \circ \cdots \circ x_k^v \subseteq Q \text{ as } Q \text{ is a } \mathcal{C}\text{-hyperideal. The hyperideal } Q \text{ is a } (kv, v)\text{-absorbing hyperideal of } A \text{ because } k \geq \lceil \frac{u}{v} \rceil. \text{ Then we conclude that } x_{i_1}^{r_1} \circ \cdots \circ x_{i_v}^{r_v} \subseteq Q \text{ for some } \{i_1, \cdots, i_v\} \subseteq \{1, \cdots, k\} \text{ such that } 0 \leq i_1, \cdots, i_v \leq v \text{ and } r_1 + \cdots + r_v = v. \text{ This implies that } x_{i_1}^v \circ \cdots \circ x_{i_v}^v \subseteq Q \text{ which means } x_{i_1} \circ \cdots \circ x_{i_v} \subseteq rad(Q). \text{ Consequently, } rad(Q) \text{ is a } (k, v)\text{-absorbing hyperideal of } A. \end{array}$ 

**Theorem 3.6.** Assume that Q is a hyperideal of A and  $u, v \in \mathbb{N}$  with u > v. Then there exists a (u, v)-absorbing hyperideal of A which is minimal in the set of all (u, v)-absorbing hyperideals of A containing Q.

Proof. Suppose that  $\Gamma = \{P \mid P \text{ is a } (u, v)\text{-absorbing hyperideal of } A \text{ and } Q \subseteq P\}$ . Let M be a maximal hyperideal of A containing Q. Then M is a (u, v)-absorbing hyperideal for all  $u, v \in \mathbb{N}$  with u > v and so  $M \in \Gamma$  which means  $\Gamma \neq \emptyset$ . Order  $\Gamma$  by  $\leq$  such that  $P_1 \leq P_2$  if and only if  $P_1 \supseteq P_2$  for every  $P_1, P_2 \in \Gamma$ . Let  $\{P_i\}_{i\in\Delta}$  be a non-empty chain of hyperideals in  $\Gamma$ . Put  $I = \bigcap_{i\in\Delta}P_i$ . Assume that  $x_1 \circ \cdots \circ x_u \subseteq I$  for  $x_1, \cdots, x_u \in A \setminus U(A)$  such that none of the product of the terms v of  $x_i$ 's is a subset of I. Then that there exists  $P_j$  in the chain such that none of the product of the terms v of the  $x_i$ 's whose product is a subset of  $P_j$  which is impossible. Hence there are v of the  $x_i$ 's whose product is a subset of I. Now, by Zorn's lemma, we conclude that  $(\Gamma, \leq)$  has a maximal element which means  $\Gamma$  has a minimal (u, v)-absorbing hyperideal of A containing Q.

As an immediate consequence of the previous theorem, we have the following result.

**Corollary 3.7.** Suppose that Q is a hyperideal of A and  $v \in \mathbb{N}$ . Then there exists a minimal v-absorbing hyperideal P of A such that  $Q \subseteq P$ .

**Theorem 3.8.** Assume that Q is a (u, v)-absorbing hyperideal of A and  $k, r \in \mathbb{N}$  with  $2 \le k \le r \le u - 1$ . If Q has r minimal prime hyperideals, then there are at least  $\frac{r!}{(r-k)!k!}$  k-absorbing hyperideals of A that are minimal over Q.

Proof. Suppose that Q has r minimal prime hyperideals and k = 2. Then there exist the distinct prime hyperideals  $Q_1, \dots, Q_r$  of A that are minimal over Q. Hence  $Q_i \cap Q_j$  is a 2-absorbing hyperideal of A for each  $i, j \in \{1, \dots, r\}$  and  $i \neq j$  by Theorem 3.4 in [5]. Hence we have  $Q \subseteq Q_{ij} \subseteq Q_i \cap Q_j$  for some minmal 2-absorbing hyperideal  $Q_{ij}$  of A by Corollary 3.7. Therefore we conclude that  $Q_i$  and  $Q_j$  are only minimal prime hyperideals over  $Q_{ij}$  by Theorem 3.8 in [5]. Assume that  $Q_{ij}$ 's are not distinct such that  $Q_{ij} = Q_{st}$  for some  $i, j, s, t \in \{1, \dots, r\}$  and  $\{i, j\} \neq \{s, t\}$ . Then we conclude that  $Q_i, Q_j, Q_s$  and  $Q_t$  are distinct minimal prime hyperideals over  $Q_{ij}$  which contradicts the fact that there are are at most two prime hyperideals of A that are minimal over the 2-absorbing hyperideal  $Q_{ij}$ . Hence the  $Q_{ij}$ 's are distinct. Thus we get the result that Q has at least  $\frac{r!}{(r-k)!k!}$  minimal 2-absorbing hyperideals. If k > 2, one can easily complete the proof by using a similar argument.

**Lemma 3.9.** Let Q be a (u, v)-absorbing hyperideals of A,  $Q_1, \dots, Q_v$  be incomparable prime C-hyperideals of A such that  $Q \subseteq \bigcap_{i=1}^{v} Q_i$  and  $k_1, \dots, k_v \in \mathbb{N}$ . If  $x_1^{k_1} \circ \cdots \circ x_v^{k_v} \subseteq Q$  for  $x_i \in Q_i \setminus (\bigcup_{j \neq i} Q_j)$ , then  $x_1 \circ \cdots \circ x_v \subseteq Q$ .

*Proof.* Suppose that  $k_1, \dots, k_v \in \mathbb{N}$ . Put  $k = k_1 + \dots + k_v$ . Let  $k \ge u$ . So Q is a (k, v)-absorbing hyperideal of A by Remark 3.3(i). Then there exist integers

 $r_1, \dots, r_v$  such that  $x_1^{r_1} \circ \dots \circ x_v^{r_v} \subseteq Q, 0 \leq r_i \leq k_i$  and  $r_1 + \dots + r_v = v$ . Assume that one of the  $r_i$ 's is zero. Then we have  $x_1^{r_1} \circ \dots \circ x_{i-1}^{r_{i-1}} \circ x_{i+1}^{r_{i+1}} \circ \dots \circ x_v^{r_v} \subseteq Q \subseteq Q_i$ . Then there exists  $j \in \{1, \dots, i-1, i+1, \dots, v\}$  such that  $x_j \in Q_i$  which is impossible. Hence  $x_1 \circ \dots \circ x_v \subseteq Q$ .

**Theorem 3.10.** Assume that Q is a (u, v)-absorbing strong C-hyperideal of A for  $u, v \in \mathbb{N}$  with  $2 \leq v < u$  and  $Q_1, \dots, Q_v$  are the only minimal prime C-hyperideals over Q. If  $z_i \in Q_i \setminus (\bigcup_{j \neq i} Q_j)$ , then  $Q_j \circ \prod_{i \neq j} z_i \subseteq Q$ .

Proof. Assume that  $x \in P_j$ . If  $x \notin \bigcup_{j \neq i} Q_j$ , then we have  $x \circ \prod_{i \neq j} z_i \subseteq \bigcap_{i=1}^v Q_i = rad(Q)$  as the  $Q_i$ 's are the only minimal prime hyperideals over Q. Therefore we conclude that  $x \circ \prod_{i \neq j} z_i \subseteq Q$  by Proposition 3.5 and Lemma 3.9. Let  $x \in Q_j \cap (\bigcup_{j \neq i} Q_j)$ . Then there exist  $y \in A$  and  $z \in Q_j \setminus (\bigcup_{j \neq i} Q_j)$  such that  $w + x \in Q_j \setminus (\bigcup_{j \neq i} Q_j)$  for every  $w \in y \circ z$  by Corollary 3.12 in [5]. It follows that  $(w + x) \circ \prod_{i \neq j} z_i \subseteq Q$  and so  $w \circ \prod_{i \neq j} z_i + x \circ \prod_{i \neq j} z_i \subseteq Q$  as Q is a strong  $\mathcal{C}$ -hyperideal of A. Since  $w \circ \prod_{i \neq j} z_i \subseteq Q$ , we obtain  $x \circ \prod_{i \neq j} z_i \subseteq Q$  which means  $Q_j \circ \prod_{i \neq j} z_i \subseteq Q$ .  $\Box$ 

Let Q be a u-absorbing hyperideal of A for  $u \in \mathbb{N}$ . We define  $Abs(Q) = \min\{v \mid Q \text{ is a } v\text{-absorbing hyperideal of } A\}$ , otherwise  $Abs(Q) = \infty$  [5]. Moreover, we define  $abs(Q) = \min\{v \mid Q \text{ is an } (Abs(Q) + 1, v)\text{-absorbing hyperideal of } A\}$ . It is clear that  $abs(Q) \leq Abs(Q)$ . Thus, in Proposition 3.4, we get  $abs(\bigcap_{i=1}^{k}) \leq abs(Q_1) + \cdots + abs(Q_k)$  and  $Abs(\bigcap_{i=1}^{k}Q_i) \leq \max\{Abs(Q_1), \cdots, Abs(Q_k), abs(Q_1) + \cdots + abs(Q_k)\}$ . Also, in Proposition 3.5, we have  $abs(rad(Q)) \leq abs(Q)$  and  $Abs(rad(Q)) \leq \max\{\lceil \frac{Abs(Q)+1}{abs(Q)}\rceil - 1, abs(Q)\}$ . Recall from [1] that the hyperideals I and J are coprime if I + J = A.

**Theorem 3.11.** Assume that Q is a (u, v)-absorbing strong C-hyperideal of A for  $u, v \in \mathbb{N}$  with  $2 \leq v < u$  and  $Q_1, \dots, Q_v$  are the only minimal prime C-hyperideals over Q. Then

- (i)  $Q_1 \circ \cdots \circ Q_v \subseteq Q$  and abs(Q) = v.
- (ii) If  $Q_1, \dots, Q_v$  are pairwise coprime, then  $Q_1 \circ \dots \circ Q_v = Q$  and abs(Q) = v.

*Proof.* It can be easily seen that the claim is true in a similar manner to the proofs of Theorem 3.13 and Theorem 3.14 in [5].  $\Box$ 

**Theorem 3.12.** Assume that Q is a radical C-hyperideal of A. Then Q is a (u, v)-absorbing hyperideal of A if and only if Q is a v-absorbing hyperideal of A.

*Proof.* ( $\Longrightarrow$ ) Suppose that Q is a (u, v)-absorbing hyperideal of A and  $x_1 \circ \cdots \circ x_{u-1} \subseteq Q$  for  $x_1, \cdots, x_{u-1} \in A \setminus U(A)$ . Since  $x_1 \circ \cdots \circ x_{k-1} \circ x_k^2 \circ x_{k+1} \circ \cdots \circ x_{u-1} \subseteq Q$ , we conclude that either  $x_{i_1} \circ \cdots \circ x_{i_v} \subseteq Q$  for some  $i_j \in \{1, \cdots, k-1, k, k+1, \cdots, u-1\}$  which implies Q is a (u-1, v)-absorbing hyperideal of A or  $x_k^2 \circ x_{i_1} \circ \cdots \circ x_{i_{v-2}} \subseteq Q$  for some  $i_j \in \{1, \cdots, k-1, k, k+1, \cdots, u-1\}$  which means  $x_k \circ x_{i_1} \circ \cdots \circ x_{i_{v-2}} \subseteq rad(Q)$ . Since Q is a radical hyperideal, we get  $x_k \circ x_{i_1} \circ \cdots \circ x_{i_{v-2}} \subseteq Q$  which implies Q is a (u-1, v)-absorbing hyperideal of A. By continuing this process, we conclude Q is a (v+1, v)-absorbing hyperideal and so it is a v-absorbing hyperideal of A by Remark 3.3(i).

 $(\Leftarrow)$  Let Q is a v-absorbing hyperideal of A and  $x_1 \circ \cdots \circ x_u \subseteq Q$  for  $x_1, \cdots, x_{u-1} \in A \setminus U(A)$ . Take any  $c \in (x_{i_1} \circ \cdots \circ x_{i_{u-v}}) \setminus Q$  such that  $i_j \in \{1, \cdots, u\}$ . Then a product of v of the  $x_i$ 's othere than the  $x_{i_j}$  with c is a subset of Q and so a product of v of the  $x_i$ 's othere than the  $x_{i_j}$  is a subset of Q as Q is a v-absorbing hyperideal of A. Thus Q is a (u, v)-absorbing hyperideal of A.

Let us give the following theorem without proof as a result of Theorem 3.12.

**Corollary 3.13.** Let Q be a (u, v)-absorbing radical C-hyperideal of A. Then abs(Q) = Abs(Q).

Recall from [17] that a proper hyperideal Q of A is primary if for any  $x, y \in A$ ,  $x \circ y \subseteq Q$  implies that  $x \in Q$  or  $y^t \subseteq Q$  for some  $t \in \mathbb{N}$ . Since the radical of a primary C-hyperideal Q is a prime hyperideal of A, say P, Q is referred to as a P-primary C-hyperideal of A.

**Theorem 3.14.** Assume that  $u, v \in \mathbb{N}$  and Q is a P-primary C-hyperideal of A where P is a prime hyperideal of A with  $P^v \subseteq Q$ . Then Q is a (u, v)-absorbing hyperideal of A for any  $u \ge v+1$ . Furthermore, if  $P^v$  is a P-primary C-hyperideal of A, then  $P^v$  is a (u, v)-absorbing hyperideal of A for any  $u \ge v+1$  and  $abs(P^v) = v$ .

*Proof.* The proof follows from Theorem 4.1 in [5].

# 

# 4. AB-(u, v)-Absorbing hyperideals

This section is devoted to the study of a subclass of the (u, v)-absorbing hyperideals called AB-(u, v)-absorbing hyperideals.

**Definition 4.1.** Let  $u, v \in \mathbb{N}$  with u > v. A proper hyperideal Q of A is said to be an AB-(u, v)-absorbing hyperideal if whenever  $x_1 \circ \cdots \circ x_u \subseteq Q$  for  $x_1, \cdots, x_u \in A$ , then there are v of the  $x_i$ 's whose product is a subset of Q.

**Theorem 4.2.** Let Q be a proper strong C-hyperideal of A such that  $a + 1 \notin U(A)$  for some  $a \in Q$  and  $u, v \in \mathbb{N}$  with u > v. Then Q is an AB-(u, v)-absorbing hyperideal if and only if Q is a (u, v)-absorbing hyperideal.

*Proof.*  $(\Longrightarrow)$  It is obvious.

 $\mathbf{6}$ 

 $(\Longleftrightarrow) \text{ Assume that } x_1 \circ \cdots \circ x_u \subseteq Q \text{ for } x_1, \cdots, x_u \in A. \text{ Let } x_1, \cdots, x_{s-1} \text{ be non-unit elements in } A \text{ and } x_s, \cdots, x_u \text{ be unit elements in } A \text{ for some } s \geq v+2. \text{ Since } x_1 \circ \cdots \circ x_u \subseteq Q \text{ and } x_s, \cdots, x_u \text{ are units in } A, \text{ we get } x_1 \circ \cdots, \circ x_s \subseteq Q. \text{ Put } y_i = 1+a \text{ for each } i \in \{1, \cdots, s\}. \text{ Then the } y_i^* \text{s are non-unit. Since } Q \text{ is a } (u, v)\text{-absorbing hyperideal and } x_1 \circ \cdots \circ x_{s-1} \circ y_s \circ \cdots \circ y_u \subseteq Q, \text{ we conclude that there exist } v \text{ elements of the } x_i^* \text{ and the } y_j^* \text{s whose product is a subset of } Q. \text{ Take any } n \in \mathbb{N}. \text{ Assume that } (a+1)^n \subseteq Q. \text{ This implies that } (\sum_{k=0}^n \binom{n}{k} a^{n-k} \circ 1^k) \cap Q \neq \emptyset. \text{ Since } Q \text{ is a strong } C\text{-hyperideal of } A, \text{ we have } (\sum_{k=0}^n \binom{n}{k} a^{n-k} \circ 1^k) \subseteq Q \text{ and so } 1 \in Q, \text{ a contradiction.} \text{ Then } (a+1)^n \notin Q \text{ for all } n \geq 1. \text{ Thus } x_{i_1} \circ \cdots \circ x_{i_h} \circ (a+1)^{v-h} \subseteq Q \text{ for some } h \in \{1, \cdots, v\} \text{ and } i_j \in \{1, \cdots, s-1\}. \text{ Clearly, } \sum_{k=0}^{v-h} \binom{v-h}{k} a^{v-h-k} \circ 1^k = X+1 \text{ where } X \subseteq Q. \text{ Since } x_i_1 \circ \cdots \circ x_{i_h} \circ (X+1) \subseteq Q \cap (x_{i_1} \circ \cdots \circ x_{i_h} \circ X) + (x_{i_1} \circ \cdots \circ x_{i_h} \circ 1), \text{ we obtain } (x_{i_1} \circ \cdots \circ x_{i_h} \circ X) + (x_{i_1} \circ \cdots \circ x_{i_h} \circ 1) \subseteq Q. \text{ Hence we get the result that } x_{i_1} \circ \cdots \circ x_{i_h} \subseteq Q \text{ which implies } Q \text{ is an } AB-(u, v)\text{-absorbing hyperideal of } A. \Box$ 

As a result of the previous theorems, we give the following result.

**Corollary 4.3.** Let Q be a proper strong C-hyperideal of A such that  $Q \nsubseteq J(A)$  and  $u, v \in \mathbb{N}$  with u > v. Then Q is an AB-(u, v)-absorbing hyperideal if and only if Q is a (u, v)-absorbing hyperideal.

(u, v)-ABSORBING (PRIME) HYPERIDEALS IN COMMUTATIVE MULTIPLICATIVE HYPERRINGS7

Let X be a finite sum of finite products of elements of A. Consider the relation  $\gamma$  on a multiplicative hyperring A defined as  $x\gamma y$  if and only if  $\{x, y\} \subseteq X$ , namely,  $x\gamma y$  if and only if  $\{x, y\} \subseteq \sum_{j \in J} \prod_{i \in I_j} z_i$  for some  $z_1, ..., z_n \in A$  and  $I_j, J \subseteq \{1, ..., n\}$ .  $\gamma^*$  denotes the transitive closure of  $\gamma$ . The relation  $\gamma^*$  is the smallest equivalence relation on A such that the set of all equivalence classes, i.e., the quotient  $A/\gamma^*$ , is a fundamental ring. Assume that  $\Sigma$  is the set of all finite sums of products of elements of A. We can rewrite the definition of  $\gamma^*$  on A, namely,  $x\gamma^* y$ if and only if there exist  $z_1, ..., z_n \in A$  such that  $z_1 = x, z_{n+1} = y$  and  $u_1, ..., u_n \in \Sigma$ where  $\{z_i, z_{i+1}\} \subseteq u_i$  for  $1 \leq i \leq n$ . Suppose that  $\gamma^*(x)$  is the equivalence class containing  $x \in A$ . Define  $\gamma^*(x) \oplus \gamma^*(y) = \gamma^*(z)$  for every  $z \in \gamma^*(x) + \gamma^*(y)$  and  $\gamma^*(x) \odot \gamma^*(y) = \gamma^*(w)$  for every  $w \in \gamma^*(x) \circ \gamma^*(y)$ . Then  $(A/\gamma^*, \oplus, \odot)$  is a ring called a fundamental ring of A [37].

**Theorem 4.4.** Let Q be a proper hyperideal of A. Then Q is an AB-(u, v)-absorbing hyperideal of  $(A, +, \circ)$  if and only if  $Q/\gamma^*$  is an AB-(u, v)-absorbing hyperideal of  $(A/\gamma^*, \oplus, \odot)$ .

*Proof.* ( $\Longrightarrow$ ) Let  $x_1 \odot \cdots \odot x_u \in Q/\gamma^*$  for some  $x_1, \cdots, x_u \in A/\gamma^*$ . Then there exist  $a_i \in A$  for every  $i \in \{1, \cdots, u\}$  such that  $x_i = \gamma^*(a_i)$ . This implies that  $x_1 \odot \cdots \odot x_u = \gamma^*(a_1) \odot \cdots \odot \gamma^*(a_u) = \gamma^*(a_1 \circ \cdots \circ a_u)$ . Since  $\gamma^*(a_1 \circ \cdots \circ a_u) \in Q/\gamma^*$ , we get  $a_1 \circ \cdots \circ a_u \subseteq Q$ . Since Q is an AB-(u, v)-absorbing hyperideal of A, we get the result that there is v elements of the  $a_i$ 's, say  $a_{i_1}, \cdots, a_{i_v}$ , whose product is a subset of Q. This implies that  $x_{i_1} \odot \cdots \odot x_{i_v} = \gamma^*(a_{i_1}) \odot \cdots \odot \gamma^*(a_{i_v}) = \gamma^*(a_{i_1} \circ \cdots \circ a_{i_v}) \in Q/\gamma^*$ . Consequently,  $Q/\gamma^*$  is an AB-(u, v)-absorbing hyperideal of  $A/\gamma^*$ .

( $\Leftarrow$ ) Suppose that  $a_1 \circ \cdots \circ a_u \subseteq Q$  for some  $a_1, \cdots, a_u \in A$ . Then we have  $\gamma^*(a_1), \cdots, \gamma^*(a_u) \in A/\gamma^*$  and so  $\gamma^*(a_1) \odot \cdots \odot \gamma^*(a_u) = \gamma^*(a_1 \circ \cdots \circ a_u) \in Q/\gamma^*$ . Since  $Q/\gamma^*$  is an AB-(u, v)-absorbing hyperideal of  $A/\gamma^*$ , there exist v elements of the  $\gamma^*(a_i)$ 's, say  $\gamma^*(a_{i_1}), \cdots, \gamma^*(a_{i_v})$  whose product is in  $Q/\gamma^*$ . This means  $\gamma^*(a_{i_1} \circ \cdots \circ a_{i_v}) = \gamma^*(a_{i_1}) \odot \odot \gamma^*(a_{i_v}) \in Q/\gamma^*$  which implies  $a_{i_1} \circ \cdots \circ a_{i_v} \subseteq Q$ . Thus Q is an AB-(u, v)-absorbing hyperideal of A.

Let  $(A_1, +_1, \circ_1)$  and  $(A_2, +_2, \circ_2)$  be two multiplicative hyperrings with nonzero identity. The set  $A_1 \times A_2$  with the operation + and the hyperoperation  $\circ$  defined as

 $(x_1, x_2) + (y_1, y_2) = (x_1 + y_1, x_2 + y_2)$ 

 $(x_1, x_2) \circ (y_1, y_2) = \{ (x, y) \in A_1 \times A_2 \mid x \in x_1 \circ_1 y_1, y \in x_2 \circ_2 y_2 \}$ 

is a multiplicative hyperring [35]. Now, we present some characterizations of AB-(u, v)-absorbing hyperideals on cartesian product of commutative multiplicative hyperring.

**Theorem 4.5.** Assume that  $A = A_1 \times A_2$  such that  $A_1, A_2$  are commutative multiplicative hyperrings,  $Q = Q_1 \times A_2$  such that  $Q_1$  is a proper hyperideal of  $A_1$  and  $u, v \in \mathbb{N}$  with u > v. Then the following are equivalent.

- (i) Q is a (u, v)-absorbing hyperideal of A.
- (ii)  $Q_1$  is an AB-(u, v)-absorbing hyperideal of  $A_1$ .
- (iii) Q is an AB-(u, v)-absorbing hyperideal of A.

*Proof.* (i)  $\Longrightarrow$  (ii) Suppose on the contrary,  $Q_1$  is not an AB-(u, v)-absorbing hyperideal of  $A_1$ . This means that we have  $x_1 \circ_1 \cdots \circ_1 x_u \subseteq Q_1$  for some  $x_1, \cdots, x_u \in A_1$ and none of product of the terms v of the  $x_i$ 's is not a subset of  $Q_1$ . Since Q is a (u, v)-absorbing hyperideal of A,  $(x_1, 0) \circ \cdots \circ (x_u, 0) \subseteq Q$  and none of the  $(x_i, 0)$ 's

#### MAHDI ANBARLOEI

are unit, there exist v of the  $(x_i, 0)$ 's whose is a subset of Q which implies there exist v of the  $x_i$ 's whose is a subset of  $Q_1$  which is a contradiction. Hence  $Q_1$  is an AB-(u, v)-absorbing hyperideal of  $A_1$ .

(ii)  $\Longrightarrow$  (iii) Let  $(x_1, y_1) \circ \cdots \circ (x_u, y_u) \subseteq Q$  for  $(x_1, y_1), \cdots, (x_u, y_u) \in A$ . This means that  $x_1 \circ_1 \cdots \circ_1 x_u \subseteq Q_1$ . Since  $Q_1$  is an AB-(u, v)-absorbing hyperideal of  $A_1$ , there exist v of the  $x_i$ 's, say  $x_{i_1}, \cdots, x_{i_v}$ , whose product is a subset of  $Q_1$ . It follows that  $(x_{i_1}, y_{i_1}) \circ \cdots \circ (x_{i_v}, y_{i_v}) \subseteq Q$  and so Q is an AB-(u, v)-absorbing hyperideal of A.

(iii)  $\implies$  (i) The claim is true because every AB-(u, v)-absorbing is a (u, v)-absorbing hyperideal.

# 5. (u, v)-Absorbing prime hyperideals

In this section, we aim to extend the class of the (u, v)-absorbing hyperideals to (u, v)-absorbing prime hyperideals.

**Definition 5.1.** Let  $u, v \in \mathbb{N}$  with u > v. A proper hyperideal Q of A refers to a (u, v)-absorbing prime hyperideal if  $x_1 \circ \cdots \circ x_u \subseteq Q$  for  $x_1, \cdots, x_u \in A \setminus U(A)$  implies either  $x_1 \circ \cdots \circ x_v \subseteq Q$  or  $x_{v+1} \circ \cdots \circ x_u \subseteq Q$ .

**Example 5.2.** In Example 3.2, the hyperideal  $\langle 15 \rangle$  is a (3, 1)-absorbing prime hyperideal. However, since  $3^2 \circ 5 = \{180, 360, 720\} \subseteq \langle 15 \rangle$  but  $3, 5 \notin \langle 15 \rangle$ , then  $\langle 15 \rangle$  is not a (3, 1)-absorbing hyperideal

*Remark* 5.3. Let  $u, v \in \mathbb{N}$  with u > v and Q be a C-hyperideal of A.

- (i) Q is a (u, v)-absorbing prime hyperideal of A if and only if Q is a (u, u-v)-absorbing prime hyperideal.
- (ii) If Q is a (u, v)-absorbing prime hyperideal of A, then Q is a (u + 1, v + 1)absorbing prime hyperideal.

**Proposition 5.4.** Assume that Q is a strong C-hyperideal of A such that A is not local and  $u, v \in \mathbb{N}$  with u > v. Then Q is a (u, v)-absorbing prime hyperideal of A if and only if Q is a (u - 1, v - 1)-absorbing prime hyperideal.

*Proof.* ( $\Longrightarrow$ ) Let Q be a (u, v)-absorbing prime hyperideal of A. Suppose on contrary that Q is not a (u-1, v-1)-absorbing prime hyperideal of A. Then there exist  $x_1, \dots, x_{u-1} \in A \setminus U(A)$  such that  $x_1 \circ \dots \circ x_{u-1} \subseteq Q$  but neither  $x_1 \circ \dots \circ x_{v-1} \subseteq Q$  nor  $x_v \circ \dots \circ x_{u-1} \subseteq Q$ . Take any  $y \in A \setminus U(A)$ . Since Q is a (u, v)-absorbing prime hyperideal of A,  $y \circ x_1 \circ \dots \circ x_{u-1} \subseteq Q$  and  $x_v \circ \dots \circ x_{u-1} \not\subseteq Q$ , we have  $y \circ x_1 \circ \dots \circ x_{v-1} \subseteq Q$ . Take any  $z \in U(A)$ . If  $y + z \notin U(A)$ , then we obtain  $(y+z) \circ x_1 \circ \dots \circ x_{v-1} \subseteq Q$  as  $(y+z) \circ x_1 \circ \dots \circ x_{u-1} \subseteq Q$ ,  $x_v \circ \dots \circ x_{u-1} \not\subseteq Q$  and Q is a (u, v)-absorbing prime hyperideal of A. Since Q is a strong C-hyperideal of A and  $(y+z) \circ x_1 \circ \dots \circ x_{v-1} \subseteq y \circ x_1 \circ \dots \circ x_{v-1} + z \circ x_1 \circ \dots \circ x_{v-1}$ , we conclude that  $y \circ x_1 \circ \dots \circ x_{v-1} + z \circ x_1 \circ \dots \circ x_{v-1} \subseteq Q$ . This implies that  $x_1 \circ \dots \circ x_{v-1} \subseteq Q$  because  $y \circ x_1 \circ \dots \circ x_{v-1} \subseteq Q$ . This is a contradiction. Then we have  $y + z \in U(A)$ . By Lemma 2.6 in [28], we get the result that A is local which is a contradiction. Thus Q is a (u-1, v-1)-absorbing prime hyperideal of A.

 $(\Leftarrow)$  It is obvious by Remark 5.3 (ii).

**Proposition 5.5.** Assume that  $u, v \in \mathbb{N}$  with u > v and Q is a strong C-hyperideal of A such that A is not local. Then Q is a (u - v + 1, 1)-absorbing prime hyperideal of A if and only if Q is a prime hyperideal.

*Proof.* ( $\Longrightarrow$ )Let Q be a (u - v + 1, 1)-absorbing prime hyperideal of A. Then we get the result that Q is a (u - v + 1, u - v)-absorbing prime hyperideal of A by Remark 5.3 (i). Hence we conclude that Q is a prime hyperideal by Proposition 5.4. ( $\Leftarrow$ ) Straightforward.

For a C-hyperideal Q of A that the hyperring A is not local, it can be easily proved, by induction, that Q is a (u - 1, v - 1)-absorbing prime hyperideal for  $u, v \in \mathbb{N}$  with u > v if and only if Q is a (u - v + 1, 1)-absorbing prime hyperideal. Thus in view of Proposition 5.4 and Proposition 5.5, we have the following result.

**Corollary 5.6.** In a hyperring A that is not local, every (u, v)-absorbing prime hyperideal is prime and vice versa.

**Theorem 5.7.** Assume that  $Q_1$  and  $Q_2$  are strong *C*-hyperideals of the hyperring  $A_1$  and  $A_2$ , respectively. Let  $u, v \in \mathbb{N}$  with u > v. Then the following are equivalent: (i)  $Q_1 \times Q_2$  is a (u, v)-absorbing prime hyperideal of  $A_1 \times A_2$ .

- (ii)  $Q_1 \times Q_2$  is a prime hyperideal of  $A_1 \times A_2$ .
- (iii)  $Q_1$  is a prime hyperideal of  $A_1$  and  $Q_2 = A_2$  or  $Q_2$  is a prime hyperideal of  $A_2$  and  $Q_1 = A_1$ .

*Proof.* It can be easily seen that the idea is true by Corollary 5.6 as  $A_1 \times A_2$  is not local.

**Proposition 5.8.** Let  $u, v \in \mathbb{N}$  with u > v and Q be a strong C-hyperideal of A. Then Q is a (v + 1, v)-absorbing prime hyperideal of A if and only if Q is a prime hyperideal or A is local such that  $M^v \subseteq Q$  for the only maximal hyperideal M of A.

*Proof.* (⇒) Let *Q* be a (*v*+1, *v*)-absorbing prime hyperideal of *A* and the hyperring *A* is not local. Then *Q* is prime by Corollary 5.6. Suppose that *A* is local and *M* is the only maximal hyperideal of *A* such that  $M^{v} \notin Q$ . If *Q* is not a (*v*, *v* − 1)-absorbing prime hyperideal of *A*, then there exist  $x_{1}, \dots, x_{v} \in A \setminus U(A)$  such that  $x_{1} \circ \dots \circ x_{v} \subseteq Q$  but  $x_{1} \circ \dots \circ x_{v-1} \notin Q$  and  $x_{v} \notin Q$ . Take any  $a \in a_{1} \circ \dots \circ a_{v}$  for every  $a_{1}, \dots, a_{v} \in M$ . Since *Q* is a (*v* + 1, *v*)-absorbing prime hyperideal of *A*,  $a \circ x_{1} \circ \dots \circ x_{v} \subseteq Q$  and  $x_{v} \notin Q$ , we get  $a \circ x_{1} \circ \dots \circ x_{v-1} \subseteq Q$ . Let  $b \in x_{1} \circ \dots \circ x_{v-1}$ . So  $a \circ b \subseteq a \circ x_{1} \circ \dots \circ x_{v-1} \subseteq Q$ . On the other hand,  $a \circ b \subseteq a_{1} \circ \dots \circ a_{v} \circ b$ . Then we get  $a_{1} \circ \dots \circ a_{v} \circ b \subseteq Q$  as *Q* is a strong *C*-hyperideal of *A*. If  $b \in Q$ , then  $x_{1} \circ \dots \circ x_{v-1} \cap Q \neq \emptyset$  and  $s_{1} \circ \dots \circ x_{v-1} \subseteq Q$  which is impossible. Therefore  $b \notin Q$ . Since *Q* is a (*v* + 1, *v*)-absorbing prime hyperideal of *A*. By induction, we conclude that Q is a (*v*, *v* − 1)-absorbing prime hyperideal of *A*. By induction, we conclude that *Q* is prime as  $M^{i} \notin Q$  for any  $i \leq v$ .

 $(\Leftarrow)$  It is obvious.

**Corollary 5.9.** Assume that  $u, v \in \mathbb{N}$  with u > v and Q is a (v + 1, v)-absorbing prime strong C-hyperideal of A that is not prime. If P is a hyperideal of A containing Q, then P is a (u, v)-absorbing prime hyperideal of A.

*Proof.* Suppose that Q is a (v+1, v)-absorbing prime strong C-hyperideal of A that is not prime. By Proposition 5.8, we get the result that A is local and  $M^v \subseteq Q$  where M is the only maximal hyperideal of A. Then  $M^v \subseteq P$  and so we are done by By Proposition 5.8.

**Theorem 5.10.** Let Q be a (u, v)-absorbing prime strong C-hyperideal of A where  $u, v \in \mathbb{N}$  and u > v. Then the following statements hold:

- (i) rad(Q) is a prime hyperideal of A.
- (ii) If Q is not a prime hyperideal of A, then A is local. In particular, if u = v + 1, then rad(Q) = M where M is the only maximal hyperideal of A.

*Proof.* (i) Suppose that Q is a (u, v)-absorbing prime strong  $\mathcal{C}$ -hyperideal of A. Let  $a \circ b \subseteq rad(Q)$  for  $a, b \in A$ . We may assume  $a, b \in A \setminus U(A)$ , without loss of generality. Then we have  $a^n \circ b^n \subseteq Q$  for some  $n \in \mathbb{N}$ . Take any  $x \in a^n$  and  $y \in b^n$ . If x or  $y \in U(A)$ , then we obtain  $b^n \subseteq e \circ b^n \subseteq x^{-1} \circ x \circ b^n \subseteq x^{-1} \circ a^n \circ b^n \subseteq Q$  which means  $b \in rad(Q)$  or  $a^n \subseteq a^n \circ e \subseteq a^n \circ y \circ y^{-1} \subseteq a^n \circ b^n \circ y^{-1} \subseteq Q$  which means  $a \in rad(Q)$ . Now, we assume that  $x, y \notin U(A)$ . Since  $a^{v-1} \circ x \circ b^{u-v-1} \circ y \subseteq Q$  and Q is a (u, v)-absorbing prime hyperideal of A, we get the result that  $a^{v-1} \circ x \subseteq Q$  or  $b^{u-v-1} \circ y \subseteq Q$ . In the first case, since  $(a^{v-1} \circ a^n) \cap Q \neq \emptyset$ , we get  $a^{v+n-1} = a^{v-1} \circ a^n \subseteq Q$  which means  $a \in rad(Q)$ . In the second case, since  $(b^{u-v-1} \circ b^n) \cap Q \neq \emptyset$ , we have  $b^{u-v+n-1} = b^{u-v-1} \circ b^n \subseteq Q$  which implies  $b \in rad(Q)$ . Thus rad(Q) is a prime hyperideal of A.

(ii) Suppose that Q is a (u, v)-absorbing prime strong C-hyperideal of A that is not prime. Then we conclude that A is local by Corollary 5.6. Let u = v + 1 and M be the only maximal hyperideal of A. By Proposition 5.8, we conclude that  $M^v \subseteq Q$  and so rad(Q) = M.

**Lemma 5.11.** Let every hyperideal of A be a prime C-hyperideal. Then A is a hyperfield.

*Proof.* Let  $0 \neq x \in A$ . Since  $\langle x^2 \rangle$  is a prime hyperideal of A and  $x^2 \subseteq \langle x^2 \rangle$ , we get  $x \in \langle x^2 \rangle$  which means  $x \in x^2 \circ a$  for some  $a \in A$ . It follows that  $0 \in x^2 \circ a - x$ . Since  $0 \in x^2 \circ a - x \cap \langle 0 \rangle$ , we get the result that  $(x \circ a - 1) \circ x \subseteq x^2 \circ a - x \subseteq \langle 0 \rangle$ . Let  $t \in x \circ a$ . Since  $\langle 0 \rangle$  is a prime hyperideal,  $(t-1) \circ x \subseteq \langle 0 \rangle$  and  $x \neq 0$ , we obtain t-1=0 and so  $1 \in x \circ a$  which means x is unit. Thus we conclude that A is a hyperfield.

**Theorem 5.12.** Let every hyperideal of A be a strong C-hyperideal, J(A) = M and  $v \in \mathbb{N}$ . Then every proper hyperideal in A is a (v+1, v)-absorbing prime hyperideal if and only if A is local and  $M^v = 0$ .

*Proof.* ( $\Longrightarrow$ ) Suppose on contrary that A is not local. By Corollary 5.6, we conclude that every proper hyperideal in A is prime. Therefore, by Lemma 5.11, we get the result that A is a hyperfield which is impossible. Then A is local. Let M be the only maximal hyperideal of A. Since  $\langle 0 \rangle$  is a (v + 1, v)-absorbing prime hyperideal and A is local, we have  $M^v = 0$  by the hypothesis and Proposition 5.8.

 $(\Leftarrow)$  Let Q be an arbitrary hyperideal and  $x_1 \circ \cdots \circ x_{v+1} \subseteq Q$  for  $x_1, \cdots, x_{v+1} \in A \setminus U(A)$ . Since A is local, M is the only maximal hyperideal of A and  $x_1, \cdots, x_v \in A \setminus U(A)$ , we conclude that  $x_1 \circ \cdots \circ x_v \subseteq M^v = 0$  which means  $x_1 \circ \cdots \circ x_v \subseteq Q$ , as required.  $\Box$ 

**Theorem 5.13.** Let Q be a proper hyperideal of A and  $u, v \in \mathbb{N}$  with u > v. Then Q is a (u, v)-absorbing prime hyperideal of A if and only if for proper hyperideals  $Q_1, \dots, Q_u$  of  $A, Q_1 \circ \dots \circ Q_u \subseteq Q$  implies that  $Q_1 \circ \dots \circ Q_v \subseteq Q$  or  $Q_{v+1} \circ \dots \circ Q_u \subseteq Q$ .

*Proof.* ( $\Longrightarrow$ ) Let Q be a (u, v)-absorbing prime hyperideal of A. Assume that  $Q_1 \circ \cdots \circ Q_u \subseteq Q$  where  $Q_1, \cdots, Q_u$  are proper hyperideals of A such that  $Q_{v+1}, \circ \cdots \circ Q_u \notin Q$ . Then there exists  $x_i \in Q_i$  for each  $i \in \{v+1, \cdots, u\}$  such that  $x_{v+1} \circ \cdots \circ x_u \notin Q$ . Take any  $y_i \in Q_i$  for each  $i \in \{1, \cdots, v\}$ . Therefore we have  $y_1 \circ \cdots y_v \circ x_{v+1} \circ \cdots \circ x_u \subseteq Q$ . This implies that  $y_1 \circ \cdots y_v \subseteq Q$  as Q is a (u, v)-absorbing prime hyperideal of A and  $x_{v+1} \circ \cdots \circ x_u \notin Q$ . Hence  $Q_1 \circ \cdots \circ Q_v \subseteq Q$ , as needed.

 $(\Leftarrow)$  Let  $x_1 \circ \cdots \circ x_u \subseteq Q$  for some  $x_1, \cdots, x_u \in A \setminus U(A)$  such that  $x_1 \circ \cdots \circ x_v \notin Q$ . Q. Put  $Q_i = \langle x_i \rangle$  for each  $i \in \{1, \cdots, u\}$ . Hence we have  $Q_1 \circ \cdots \circ Q_u \subseteq Q$  and  $Q_1 \circ \cdots \circ Q_v \notin Q$ . By the hypothesis, we conclude that  $Q_{v+1} \circ \cdots \circ Q_u \subseteq Q$  which implies  $x_{v+1} \circ \cdots \circ x_u \subseteq Q$ . This means that Q is a (u, v)-absorbing prime hyperideal of A.

**Proposition 5.14.** Let  $u, v \in \mathbb{N}$  with u > v, Q a (u, v)-absorbing prime hyperideal of A and  $x \in A \setminus (Q \cup U(A))$ . Then (Q : x) is a (u - 1, v - 1)-absorbing prime hyperideal of A.

*Proof.* Let  $x_1 \circ \cdots \circ x_{u-1} \subseteq (Q:x)$  for  $x_1, \cdots, x_{u-1} \in A \setminus U(A)$  such that  $x_1 \circ \cdots \circ x_{v-1} \not\subseteq (Q:x)$ . So we have  $x \circ x_1 \circ \cdots \circ x_{u-1} \subseteq Q$ . Since Q is a (u,v)-absorbing prime hyperideal of A and  $x \circ x_1 \circ \cdots \circ x_{v-1} \not\subseteq Q$ , we get the result that  $x_v \circ \cdots x_{u-1} \subseteq Q \subseteq (Q:x)$ . This shows that (Q:x) is a (u-1, v-1)-absorbing prime hyperideal of A.

Assume that A is a multiplicative hyperring. Then the set of all hypermatrices of A is denoted by  $M_m(A)$ . Let  $I = (I_{ij})_{m \times m}$ ,  $J = (J_{ij})_{m \times m} \in P^*(M_m(A))$ . Then  $I \subseteq J$  if and only if  $I_{ij} \subseteq J_{ij}[1]$ .

**Theorem 5.15.** Let  $u, v \in \mathbb{N}$  with u > v and Q be a hyperideal of A. If  $M_m(Q)$  is a (u, v)-absorbing prime hyperideal of  $M_m(A)$ , then Q is a (u, v)-absorbing prime hyperideal of A.

*Proof.* Let  $x_1 \circ \cdots \circ x_u \subseteq Q$  for  $x_1, \cdots, x_u \in A \setminus U(A)$ . Therefore we have

$$\begin{pmatrix} x_1 \circ \cdots \circ x_u & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix} \subseteq M_m(Q).$$

Since  $M_m(Q)$  is a (u, v)-absorbing prime hyperideal of  $M_m(A)$  and

$$\begin{pmatrix} x_1 \circ \cdots \circ x_u & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix} = \begin{pmatrix} x_1 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix} \circ \cdots \circ \begin{pmatrix} x_u & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix}$$

we conclude that

$$\begin{pmatrix} x_1 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix} \circ \cdots \circ \begin{pmatrix} x_v & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix} = \begin{pmatrix} x_1 \circ \cdots \circ x_v & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix} \subseteq M_m(Q)$$

$$\begin{pmatrix} x_{v+1} & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix} \circ \cdots \circ \begin{pmatrix} x_u & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix} = \begin{pmatrix} x_{v+1} \circ \cdots \circ x_u & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix} \subseteq M_m(Q).$$

Thus we get the result that  $x_1 \circ \cdots \circ x_v \subseteq Q$  or  $x_{v+1} \circ \cdots \circ x_u \subseteq Q$ . Hence Q is a (u, v)-absorbing prime hyperideal of A.

Suppose that  $(A, +, \circ)$  is a commutative multiplicative hyperring and x is an indeterminate. Then  $(A[x], +, \diamond)$  is a polynomial multiplicative hyperring where  $tx^n \diamond sx^m = (t \circ s)x^{n+m}$  [11].

**Theorem 5.16.** Let  $u, v \in \mathbb{N}$  with u > v. If Q is a (u, v)-absorbing prime hyperideal of  $(A, +, \circ)$ , then Q[x] is a (u, v)-absorbing prime hyperideal of  $(A[x], +, \diamond)$ .

*Proof.* Assume that  $t_1(x) \diamond \cdots \diamond t_u(x) \subseteq Q[x]$ . Without loss of generality, we may assume that  $t_i(x) = a_i x^{n_i}$  for each  $i \in \{1, \cdots, u\}$  and  $a_i \in A$ . Therefore  $a_1 \circ \cdots \circ a_u x^{n_1 + \cdots + n_u} \subseteq Q[x]$ . This implies that  $a_1 \circ \cdots \circ a_u \subseteq Q$ . Since Q is a (u, v)-absorbing prime hyperideal of  $(A, +, \circ)$ , we obtain  $a_1 \circ \cdots \circ a_v \subseteq Q$  or  $a_{v+1} \circ \cdots \circ a_u \subseteq Q$  which implies  $t_1(x) \diamond \cdots \diamond t_v(x) = a_1 x^{n_1} \diamond \cdots \diamond a_v x^{n_v} = a_1 \circ \cdots \circ a_v x^{n_1 + \cdots + n_v} \subseteq Q[x]$  or  $t_{v+1}(x) \diamond \cdots \diamond t_u(x) = a_{v+1} x^{n_1} \diamond \cdots \diamond a_u x^{n_u} = a_{v+1} \circ \cdots \circ a_u x^{n_{v+1} + \cdots + n_u} \subseteq Q[x]$ . Thus Q[x] is a (u, v)-absorbing prime hyperideal of  $(A[x], +, \diamond)$ .

In view of Theorem 5.16, we get the following result.

**Corollary 5.17.** Suppose that Q is a (u, v)-absorbing prime hyperideal of A. Then Q[x] is a (u, v)-absorbing prime hyperideal of A[x].

Recall from [19] that a mapping  $\theta$  from the multiplicative hyperring  $(A_1, +_1, \circ_1)$ into the multiplicative hyperring  $(A_2, +_2, \circ_2)$  is a hyperring good homomorphism if  $\theta(x +_1 y) = \theta(x) +_2 \theta(y)$  and  $\theta(x \circ_1 y) = \theta(x) \circ_2 \theta(y)$  for all  $x, y \in A_1$ .

**Theorem 5.18.** Assume that  $A_1$  and  $A_2$  are two multiplicative hyperrins such that  $\theta : A_1 \longrightarrow A_2$  is a hyperring good homomorphism and  $\theta(x) \notin U(A_2)$  for every  $x \in A_1 \setminus U(A_1)$ . Let  $u, v \in \mathbb{N}$  with u > v. Then the following hold.

- (i) If  $Q_2$  is a (u, v)-absorbing prime hyperideal of  $A_2$ , then  $\theta^{-1}(Q_2)$  is a (u, v)absorbing prime hyperideal of  $A_1$ .
- (ii) If θ is surjective and Q<sub>1</sub> is a is a (u, v)-absorbing prime C-hyperideal of A<sub>1</sub> with Ker(θ) ⊆ Q<sub>1</sub>, then θ(Q<sub>1</sub>) is a (u, v)-absorbing prime hyperideal of A<sub>2</sub>.

*Proof.* (i) Let  $x_1 \circ_1 \cdots \circ_1 x_u \subseteq \theta^{-1}(Q_2)$  for some  $x_1, \cdots, x_u \in A_1$ . Then we get  $\theta(x_1 \circ_1 \cdots \circ_1 x_u) = \theta(x_1) \circ_2 \cdots \circ_2 \theta(x_u) \subseteq Q_2$  because  $\theta$  is a good homomorphism. Since  $Q_2$  is a (u, v)-absorbing prime hyperideal of  $A_2$ , we obtain  $\theta(x_1 \circ_1 \cdots \circ_1 x_v) = \theta(x_1) \circ_2 \cdots \circ_2 \theta(x_v) \subseteq Q_2$  which means  $x_1 \circ_1 \cdots \circ_1 x_v \subseteq \theta^{-1}(Q_2)$  or  $\theta(x_{v+1} \circ_1 \cdots \circ_1 x_u) = \theta(x_{v+1}) \circ_2 \cdots \circ_2 \theta(x_u) \subseteq Q_2$  which implies  $x_{v+1} \circ_1 \cdots \circ_1 x_u \subseteq \theta^{-1}(Q_2)$ . Thus  $\theta^{-1}(Q_2)$  is a (u, v)-absorbing prime hyperideal of  $A_1$ .

(ii) Let  $y_1 \circ_2 \cdots \circ_2 y_u \subseteq \theta(Q_1)$  for  $y_1, \cdots, y_u \in A_2$ . Then there exists  $x_i \in A_1$  such that  $\theta(x_i) = y_i$  for each  $i \in \{1, \cdots, u\}$  because  $\theta$  is surjective. Therefore  $\theta(x_1 \circ_1 \cdots \circ_1 x_u) = \theta(x_1) \circ_2 \cdots \circ_2 \theta(x_u) \subseteq \theta(Q_1)$ . Now, pick any  $t \in x_1 \circ_1 \cdots \circ_1 x_u$ . Then  $\theta(t) \in \theta(x_1 \circ_1 \cdots \circ_1 x_u) \subseteq \theta(Q_1)$  and so there exists  $s \in Q_1$  such that

or

 $\theta(t) = \theta(s)$ . Then we get  $\theta(t-s) = 0$  which means  $t-s \in Ker(\theta) \subseteq Q_1$  and so  $t \in Q_1$ . Therefore  $x_1 \circ_1 \cdots \circ_1 x_u \subseteq Q_1$  as  $Q_1$  is a  $\mathcal{C}$ -hyperideal. Since  $Q_1$  is a (u, v)-absorbing prime hyperideal of  $A_1$ , we get the result that  $x_1 \circ_1 \cdots \circ_1 x_v \subseteq Q_1$ or  $x_{v+1} \circ_1 \cdots \circ_1 x_u \subseteq Q_1$ . This means  $y_1 \circ_2 \cdots \circ_2 y_v = \theta(x_1 \circ_1 \cdots \circ_1 x_v) \subseteq \theta(Q_1)$ or  $y_{v+1} \circ_2 \cdots \circ_2 y_u = \theta(x_{v+1} \circ_1 \cdots \circ_1 x_u) \subseteq \theta(Q_1)$ . Consequently,  $\theta(Q_1)$  is a (u, v)-absorbing prime hyperideal of  $A_2$ .

Now, we have the following result.

**Corollary 5.19.** Let the hyperideal  $Q_1$  of A be subset the C-hyperideal  $Q_2$  of A,  $x + Q_1 \notin U(A/Q_1)$  for all  $x \in A \setminus U(A)$  and  $u, v \in \mathbb{N}$  with u > v. Then  $Q_2$  is a (u, v)-absorbing prime hyperideal of A if and only if  $Q_2/Q_1$  is a (u, v)-absorbing prime hyperideal of  $A/Q_1$ .

*Proof.* Consider the homomorphism  $\pi : A \longrightarrow A/Q_1$  defined by  $\pi(a) = a + Q_1$ . Since  $\pi$  is a good epimorphism, the claim follows from Theorem 5.18

Let S be a non-empty subset of A. If S is closed under the hypermultiplication and S contains 1, then S refers to a multiplicative closed subset (briefly, MCS)[1]. Consider the set  $(A \times S/ \sim)$  of equivalence classes denoted by  $S^{-1}A$  such that  $(x_1, r_1) \sim (x_2, r_2)$  if and only if there exists  $r \in S$  with  $r \circ r_1 \circ x_2 = r \circ r_2 \circ x_1$ . The equivalence class of  $(x, r) \in A \times S$  is denoted by  $\frac{x}{r}$ . The triple  $(S^{-1}A, \oplus, \odot)$  is a multiplicative hyperring where

The localization map  $\pi: A \longrightarrow S^{-1}A$ , defined by  $a \mapsto \frac{a}{1}$ , is a homomorphism of

The localization map  $\pi : A \longrightarrow S^{-1}A$ , defined by  $a \mapsto \frac{a}{1}$ , is a homomorphism of hyperrings. Moreover, if I is a hyperideal of A, then  $S^{-1}I$  is a hyperideal of  $S^{-1}A$  [27].

**Theorem 5.20.** Assume that Q is a C-hyperideal of A, S a MCS such that  $Q \cap S = \emptyset$  and  $u, v \in \mathbb{N}$  with u > v. If Q is a (u, v)-absorbing prime hyperideal of A, then  $S^{-1}Q$  is a (u - 1, v - 1)-absorbing prime hyperideal of  $S^{-1}A$ .

 $\begin{array}{l} Proof. \ \mathrm{Let} \ \frac{x_1}{r_1} \odot \cdots \odot \frac{x_{u-1}}{r_{u-1}} = \frac{x_1 \circ \cdots \circ x_{u-1}}{r_1 \circ \cdots \circ r_{u-1}} \subseteq S^{-1}Q \ \mathrm{for} \ x_1, \cdots, x_{u-1} \in A \backslash U(A) \ \mathrm{and} \ r_1, \cdots, r_{u-1} \in S \ \mathrm{such} \ \mathrm{that} \ \frac{x_1}{r_1} \odot \cdots \odot \frac{x_{v-1}}{r_{v-1}} \not\subseteq S^{-1}Q. \ \mathrm{Take} \ \mathrm{any} \ a \in x_1 \circ \cdots \circ x_{u-1} \ \mathrm{and} \ r \in r_1 \circ \cdots \circ r_{u-1}. \ \mathrm{Hence} \ \frac{a}{r} \in \frac{x_1 \circ \cdots \circ x_{u-1}}{r_1 \circ \cdots \circ r_{u-1}} \ \mathrm{and} \ \mathrm{so} \ \frac{a}{r} = \frac{a'}{r'} \ \mathrm{for} \ \mathrm{some} \ a' \in Q \ \mathrm{and} \ r' \in S. \\ \mathrm{Then} \ \mathrm{there} \ \mathrm{exists} \ t \in S \ \mathrm{such} \ \mathrm{that} \ t \circ a \circ r' = t \circ a' \circ r. \ \mathrm{This} \ \mathrm{means} \ t \circ a \circ r' \subseteq Q. \\ \mathrm{Since} \ a \in x_1 \circ \cdots \circ x_{u-1}, \ \mathrm{we} \ \mathrm{conclude} \ \mathrm{that} \ t \circ a \circ r' \subseteq t \circ x_1 \circ \cdots \circ x_{u-1} \circ r'. \\ \mathrm{Therefore} \ \mathrm{we} \ \mathrm{have} \ t \circ x_1 \circ \cdots \circ x_{u-1} \circ r' \subseteq Q \ \mathrm{as} \ Q \ \mathrm{is} \ a \ \mathcal{C} \ \mathrm{hyperideal} \ \mathrm{of} \ A \ \mathrm{and} \ t \circ x_1 \circ \cdots \circ x_{u-1} \circ r'. \\ \mathrm{Therefore} \ \mathrm{we} \ \mathrm{have} \ t \circ x_1 \circ \cdots \circ x_{u-1} \circ r' \subseteq Q \ \mathrm{as} \ Q \ \mathrm{is} \ a \ \mathcal{C} \ \mathrm{hyperideal} \ \mathrm{of} \ A \ \mathrm{and} \ t \circ x_1 \circ \cdots \circ x_{u-1} \circ r'. \\ \mathrm{Therefore} \ \mathrm{we} \ \mathrm{have} \ t \circ x_1 \circ \cdots \circ x_{u-1} \circ r' \subseteq Q \ \mathrm{as} \ Q \ \mathrm{is} \ a \ \mathcal{C} \ \mathrm{hyperideal} \ \mathrm{of} \ A \ \mathrm{and} \ t \circ x_1 \circ \cdots \circ x_{u-1} \circ r'. \\ \mathrm{Therefore} \ \mathrm{we} \ \mathrm{have} \ t \circ x_1 \circ \cdots \circ x_{u-1} = \frac{x_1 \circ \cdots \circ x_{u-1}}{r_{v-1}} = \frac{x_1 \circ \cdots \circ x_{u-1}}{r_{v-1} \circ r_{v-1}} \subseteq S^{-1}Q \ \mathrm{which} \ \mathrm{is} \ \mathrm{impossible}. \ \mathrm{Since} \ Q \ \mathrm{is} \ a \ (u, v) \ \mathrm{absorbing} \ \mathrm{prime} \ \mathrm{hyperideal} \ \mathrm{of} \ A, \ \mathrm{sox}_1 \circ \cdots \circ x_{u-1} \subseteq Q \ \mathrm{and} \ \mathrm{sox}_1 \circ \cdots \circ x_{v-1} \not\subseteq Q, \\ \mathrm{we} \ \mathrm{ave} \ x_v \circ \cdots \circ x_{u-1} \subseteq Q \ \mathrm{which} \ \mathrm{implies} \ \frac{x_v}{r_v} \odot \cdots \odot \frac{x_{u-1}}{r_{u-1}} = \frac{x_v \circ \cdots \circ x_{u-1}}{r_v \circ \cdots \circ r_{u-1}} \subseteq S^{-1}Q. \\ \mathrm{This} \ \mathrm{shows} \ \mathrm{that} \ S^{-1}Q \ \mathrm{is} \ \mathrm{a} \ (u-1,v-1) \ \mathrm{absorbing} \ \mathrm{prime} \ \mathrm{hyperideal} \ \mathrm{of} \ S^{-1}A. \ \Box$ 

## References

- R. Ameri, A. Kordi, S. Hoskova-Mayerova, Multiplicative hyperring of fractions and coprime hyperideals, An. St. Univ. ovidious Constanta, 25(2017), No. 1, 5-23.
- [2] R. Ameri, M. Norouzi, On commutative hyperrings, Int. Journal of Algebraic Hyperstructures and its Applications, 1(1) (2014) 45-58.

#### MAHDI ANBARLOEI

- [3] R. Ameri, A. Kordi, On regular multiplicative hyperrings, European Journal of Pure and Applied Mathematics, 9(4)(2016) 402-418.
- [4] R. Ameri, A. Kordi, Clean multiplicative hyperrings, Italian Journal of Pure and Applied Mathematics, (35) (2015) 625-636.
- [5] M. Anbarloei, A generalization of prime hyperideals, Journal of Algebraic Systems, 8 (1) (2020) 113-127.
- M. Anbarloei, (weakly) (s,n)-closed hyperideals in commutative multiplicative hyperrings, Journal of algebraic systems, (2024), https://doi.org/10.22044/JAS.2024.13889.1780.
- [7] D.F. Anderson, A. Badawi. On n-absorbing ideals of commutative rings, Comm. Algebra, 39 (5) (2011) 1646-1672.
- [8] D.F. Anderson, A. Badawi, On (m, n)-closed ideals of commutative rings, Journal of Algebra and Its Applications, 16(1)(2017) 1-21.
- [9] E. Ay, G. Yesilot, D. Sonmez, δ-Primary hyperideals on commutative hyperrings, International Journal Mathematics and Mathematical Sciences (2017): Article ID 5428160, 4 pages.
- [10] A. Badawi, On 2-absorbing ideals of commutative rings, Bull. Austral. Math. Soc., 75 (2007) 417-429.
- [11] R.P. Ciampi, R. Rota, Polynomials over multiplicative hyperrings, J. Discrete Math. Sci. Cryptogr., 6 (2-3) (2003) 217–225.
- [12] P. Corsini, Prolegomena of hypergroup theory, Second ed., Aviani Editore, 1993.
- [13] P. Corsini, V. Leoreanu, Applications of hyperstructures theory, Adv. Math., Kluwer Academic Publishers, 2003.
- [14] J. Chvalina, J., S. Krehlik, M. Novak, Cartesian composition and the problem of generalizing the MAC condition to quasi- multiautomata, An. Stiint. Univ. Ovidius Constanta Ser. Mat., 24(3) (2016) 79-100.
- [15] I. Cristea, S. Jancic-Rasovic, Compositions hyperrings, An. Stiint. Univ. Ovidius Constanta Ser. Mat., 21(2) (2013) 81-94.
- [16] I. Cristea, Regularity of intuitionistic fuzzy relations on hypergroupoids, An. Stiint. Univ. Ovidius Constanta Ser. Mat., 22(1) (2014) 105-119.
- [17] U. Dasgupta, On prime and primary hyperideals of a multiplicative hyperring, Annals of the Alexandru Ioan Cuza University-Mathematics, LVIII (1) (2012) 19-36.
- [18] U. Dasgupta, On certain classes of hypersemirings, PhD Thsis, University of Calcutta, 2012.
- [19] B. Davvaz, V. Leoreanu-Fotea, Hyperring theory and applications, International Academic Press, USA, 2007.
- [20] F. Farzalipour, P. Ghiasvand, Semiprime hyperideals in multiplicative hyperrings, 51th Annual Iranian Mathematics conference, 1 (2021) 801-804.
- [21] P. Ghiasvand, F. Farzalipour, 1-absorbing prime avoidance theorem in a multiplicative hyperring, *Palestine Journal of Mathematics*, 12(1) (2023) 900-908.
- [22] L. Kamali Ardekani, B. Davvaz, Differential multiplicative hyperring, Journal of Algebraic Systems, 2 (1) (2014) 21-35.
- [23] H.A. Khashan, E.Y. Celikel, (m, n)-prime ideals of commutative rings, Preprints 2024, 2024010472. https://doi.org/10.20944/preprints202401.0472.v1
- [24] H.A. Khashan, E.Y. Celikel, On weakly (m, n)-prime ideals of commutative rings, Bull. Korean Math. Soc., In press.
- [25] F. Marty, Sur une generalization de la notion de groupe, In: 8iem Congres des Mathematiciens Scandinaves, Stockholm, (1934) 45-49.
- [26] C. G. Massouros, On the theory of hyperrings and hyperfields, Algebra i Logika, 24 (1985) 728-742.
- [27] A.A. Mena, I. Akray, n-absorbing I-prime hyperideals in multiplicative hyperrings, Journal of Algebraic Systems, 12 (1) (2024) 105-121.
- [28] P. Ghiasvand, F. Farzalipour, On S-prime hyperideals in multiplicative hyperrings, journal of Algebraic Hyperstructures and Logical Algebras, 2 (2) (2021) 25-34.
- [29] P. Ghiasvand, F. Farzalipour, 1-Absorbing prime avoidance theorem in multiplicative hyperrings, *Palestine Journal of Mathematics*, **12**(1) (2023) 900-908.
- [30] M. Novak, n-ary hyperstructures constructed from binary quasi-orderer semigroups, An. Stiint. Univ. Ovidius Constanta Ser. Mat., 22 (3) (2014) 147-168.
- [31] R. Procesi, R. Rota, On some classes of hyperstructures, Combinatorics Discrete Math., 208/209 (1999) 485-497.
- [32] R. Rota, Sugli iperanelli moltiplicativi, Rend. Di Math., Series VII, 4 (1982) 711-724.

14

 $(u,v)\mbox{-}ABSORBING$  (PRIME) HYPERIDEALS IN COMMUTATIVE MULTIPLICATIVE HYPERRING  $\$ 

- [33] E. Sengelen Sevim, B.A. Ersoy, B. Davvaz, Primary hyperideals of multiplicative hyperrings, *Eurasian Bulletin of Mathematics*, 1(1) (2018) 43-49.
- [34] M.K. Sen, U. Dasgupta, Some aspects of G<sub>H</sub>-rings, An. Stiint. Univ. "Al. I. Cuza" Iasi. Mat., 56 (2010) 253-272.
- [35] G. Ulucak, On expansions of prime and 2-absorbing hyperideals in multiplicative hyperrings, Turkish Journal of Mathematics, 43 (2019) 1504-1517.
- [36] T. Vougiouklis, Hyperstructures and their representations, Hadronic Press, Inc., Palm Harber, USA, 1994.
- [37] T. Vougiouklis, Hyperstructures and their representations, *Hadronic Press Inc.*, *Florida*, (1994).

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, IMAM KHOMEINI INTERNATIONAL UNIVERSITY, QAZVIN, IRAN.

Email address: m.anbarloei@sci.ikiu.ac.ir