

The Connes-Chamseddine Hochschild cocycle and the noncommutative integral

Tong Wu^a, Yong Wang^{b,*}

^aDepartment of Mathematics, Northeastern University, Shenyang, 110819, China

^bSchool of Mathematics and Statistics, Northeast Normal University, Changchun, 130024, China

Abstract

In [5], Connes and Chamseddine defined a Hochschild cocycle in the general framework of noncommutative geometry. They computed this Hochschild cocycle for the Dirac operator on 4-dimensional manifolds. We propose a way to study the Connes-Chamseddine Hochschild cocycle from the viewpoint of the noncommutative integral on 6-dimensional manifolds in this paper. We compute several interesting noncommutative integral defined in [8] by the normal coordinated way on n-dimensional manifolds. As a corollary, the Connes-Chamseddine Hochschild cocycle on 6-dimensional manifolds is obtained.

Keywords: The noncommutative geometry; the noncommutative integral; the Connes-Chamseddine Hochschild cocycle; the noncommutative residue.

1. Introduction

Until now, many geometers have studied noncommutative residues. In [9, 18], authors found noncommutative residues are of great importance to the study of noncommutative geometry. Connes showed us that the noncommutative residue on a compact manifold M coincided with the Dixmier's trace on pseudodifferential operators of order $-\dim M$ in [3]. Therefore, the non-commutative residue can be used as integral of noncommutative geometry and become an important tool of noncommutative geometry. In [4], Connes used the noncommutative residue to derive a conformal 4-dimensional Polyakov action analogy. Several years ago, Connes made a challenging observation that the noncommutative residue of the square of the inverse of the Dirac operator was proportional to the Einstein-Hilbert action, which we call the Kastler-Kalau-Walze theorem. In [11], Kastler gave a bruteforce proof of this theorem. In [10], Kalau and Walze proved this theorem in the normal coordinates system simultaneously. Ackermann proved that the Wodzicki residue of the square of the inverse of the Dirac operator $\text{Wres}(D^{-2})$ in turn is essentially the second coefficient of the heat kernel expansion of D^2 in [1].

The many noncommutative residues of Dirac operators have been studied [7, 13, 16, 17]. Wang proved a Kastler-Kalau-Walze type theorem for perturbations of Dirac operators on compact manifolds with or without boundary and gave two kinds of operator-theoretic explanations of the gravitational action on boundary in [13]. In [17], Wang, Wang and Yang gave two kinds of operator-theoretic explanations of the gravitational action about Dirac operators with torsion in the case of 4-dimensional compact manifolds with flat boundary. In [16], Wang, Wang and Wu gave some new spectral functionals which is the extension of spectral functionals to the noncommutative realm with torsion, and related them to the noncommutative residue for manifolds with boundary about Dirac operators with torsion. In [7], Dabrowski, Sitarz and Zalecki recovered the Einstein functional by Dirac operators and the noncommutative residue. In [5], Connes and

*Corresponding author.

Email addresses: wut977@nenu.edu.cn (Tong Wu^a), wangy581@nenu.edu.cn (Yong Wang^{b,*})

Chamseddine defined a Hochschild cocycle which contains the Dirac operator in the general framework of noncommutative geometry. They proved that the inner fluctuations of the spectral action can be computed as residues and give exactly the counterterms for the Feynman graphs with fermionic internal lines. And they computed this Hochschild cocycle for the Dirac operator on 4-dimensioanl manifolds. In [8], Figueroa H, Gracia-Bond“a J M, Lizzi F, et al. introduced a noncommutative integral based on the noncommutative residue [19]. The motivation of this paper is to study the Connes-Chamseddine Hochschild cocycle from the viewpoint of the noncommutative integral on compact oriented 6-dimensional spin manifolds without boundary in the general framework of noncommutative geometry.

The organization of this paper is as follows. In Section 2, we recall some basic facts and formulas about Dirac operator and we introduce the Connes-Chamseddine Hochschild cocycle. In Section 3, using the relationship between the noncommutative integral and the noncommutative residue, that is the formula $\int P ds^n := \text{Wres}(PD^{-n}) := \int_{S^* M} \text{tr}[\sigma_{-n}(PD^{-n})](x, \xi)$, we give the noncommutative integral of the Connes-Chamseddine Hochschild cocycle on n-dimensional manifolds without boundary. To do it, we compute the corresponding symbols for the pseudo-differential operators, and then get trace of the results in the normal coordinate system. And finally, we obtain the Connes-Chamseddine Hochschild cocycle defined by in [5] on 6-dimensional manifolds with boundary.

2. The Dirac operator and the Connes-Chamseddine Hochschild cocycle

Let M be an n-dimensional compact oriented spin manifold with Riemannian metric g , and let ∇^L be the Levi-Civita connection about g . We recall that the Dirac operator D is locally given as follows in terms of an orthonormal section e_i (with dual section θ^k) of the frame bundle of M [11]:

$$\begin{aligned} D &= i\gamma^i \tilde{\nabla}_i = i\gamma^i(e_i + \sigma_i); \\ \sigma_i(x) &= \frac{1}{4}\gamma_{ij,k}(x)\gamma^i\gamma^k = \frac{1}{8}\gamma_{ij,k}(x)[\gamma^j\gamma^k - \gamma^k\gamma^j], \end{aligned} \quad (2.1)$$

where $\gamma_{ij,k}$ represents the Levi-Civita connection ∇ with spin connection $\tilde{\nabla}$, specifically:

$$\begin{aligned} \gamma_{ij,k} &= -\gamma_{ik,j} = \frac{1}{2}[c_{ij,k} + c_{ki,j} + c_{kj,i}], \quad i, j, k = 1, \dots, 4; \\ c_{ij}^k &= \theta^k([e_i.e_j]). \end{aligned} \quad (2.2)$$

Here the γ^i are constant self-adjoint Dirac matrices s.t. $\gamma^i\gamma^j + \gamma^j\gamma^i = -2\delta^{ij}$. In terms of local coordinates x^μ inducing the alternative vierbein $\partial_\mu = S_\mu^i(x)e_i$ (with dual vierbein dx^μ) we have $\gamma^i e_i = \gamma^\mu \partial_\mu$, the γ^μ being now x -dependent Dirac matrices s.t. $\gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu = -2g^{\mu\nu}$ (we use latin sub-(super-) scripts for the basic e_i and greek sub-(super-) scripts for the basis ∂_μ , the type of sub-(super-) scripts specifying the type of Dirac matrices). The specification of the Dirac operator in the greek basis is as follows: one has

$$\begin{aligned} D &= i\gamma^\mu \tilde{\nabla}_\mu = i\gamma^\mu(e_\mu + \sigma_\mu); \\ \sigma_\mu(x) &= S_\mu^i(x)\sigma_i. \end{aligned}$$

Then, using the shorthand: $\Gamma^k = g^{ij}\Gamma_{ij}^k$; $\sigma^j = g^{ij}\sigma_i$, we have respective symbols of D^2 :

Lemma 2.1. [11]

$$\begin{aligned} \sigma_2(D^2) &= |\xi|^2; \\ \sigma_1(D^2) &= i(\Gamma^\mu - 2\sigma^\mu)(x)\xi_\mu; \\ \sigma_0(D^2) &= -g^{\mu\nu}(\partial_\mu\sigma_\nu + \sigma^\mu\sigma_\nu - \Gamma_{\mu\nu}^\alpha\sigma_\alpha)(x) + \frac{1}{4}s(x). \end{aligned} \quad (2.3)$$

We now introduce the Connes-Chamseddine Hochschild cocycle. Firstly, the spectral action is defined as a functional on noncommutative geometries. Such a geometry is specified by a fairly simple data of operator theoretic nature, namely a spectral triple

$$(\mathcal{A}, \mathcal{H}, D)$$

where \mathcal{A} is a noncommutative algebra with involution, acting in the Hilbert space \mathcal{H} while D is a Dirac operator, which is self-adjoint operator with compact resolvent and such that $[D, a]$ is bounded $\forall a \in \mathcal{A}$. Then ρ is a Hochschild cocycle which is given by the following formula defined by Connes and Chamseddine in [5],

$$\begin{aligned} \rho(a^0, a^1, a^2, a^3, a^4) &= \int a^0 [D^2, a^1] [D^2, a^2] [D, a^3] [D, a^4] D^{-6} - \int a^0 [D, a^1] [D^2, a^2] [D^2, a^3] [D, a^4] D^{-6} \\ &\quad + \int a^0 [D, a^1] [D, a^2] [D^2, a^3] [D^2, a^4] D^{-6} - \int a^0 [D^2, a^1] [D, a^2] [D, a^3] [D^2, a^4] D^{-6}, \end{aligned} \quad (2.4)$$

where \int denotes the noncommutative residue.

3. The noncommutative integral

For a pseudo-differential operator P , acting on sections of a spinor bundle over an even n-dimensional compact Riemannian spin manifold M , the analogue of the volume element in noncommutative geometry is the operator $D^{-n} =: ds^n$. And pertinent operators are realized as pseudodifferential operators on the spaces of sections. Extending previous definitions by Connes [6], a noncommutative integral was introduced in [8] based on the noncommutative residue [19], combine (1.4) in [5] and [11], using the definition of the residue:

$$\int P ds^n := \text{Wres} PD^{-n} := \int_{S^* M} \text{tr}[\sigma_{-n}(PD^{-n})](x, \xi), \quad (3.1)$$

where $\sigma_{-n}(PD^{-n})$ denotes the $(-n)$ th order piece of the complete symbols of PD^{-n} , tr as shorthand of trace.

By (3.1), we find the Connes-Chamseddine Hochschild cocycle consists of four noncommutative integrals mentioned above. Nextly, in order to get the result of (2.4), we need to divide the Connes-Chamseddine Hochschild cocycle into four parts and study them separately.

Part i) $P = [D^2, a^1][D^2, a^2][D, a^3][D, a^4]$, and $[D^2, a^1] := A$, $[D^2, a^2][D, a^3][D, a^4] := B$. Let $n = 2m$, by (3.1), we need to compute $\int_{S^* M} \text{tr}[\sigma_{-2m}(ABD^{-2m})](x, \xi)$. Based on the algorithm yielding the principal symbol of a product of pseudo-differential operators in terms of the principal symbols of the factors, we have

$$\begin{aligned} \sigma_{-2m}(ABD^{-2m}) &= \left\{ \sum_{|\alpha|=0}^{\infty} \frac{(-i)^{|\alpha|}}{\alpha!} \partial_\xi^\alpha [\sigma(AB)] \partial_x^\alpha [\sigma(D^{-2m})] \right\}_{-2m} \\ &= \sigma_0(AB)\sigma_{-2m}(D^{-2m}) + \sigma_1(AB)\sigma_{-2m-1}(D^{-2m}) + \sigma_2(AB)\sigma_{-2m-2}(D^{-2m}) \\ &\quad + (-i) \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(AB)] \partial_{x_j} [\sigma_{-2m-1}(D^{-2m})] + (-i) \sum_{j=1}^n \partial_{\xi_j} [\sigma_1(AB)] \partial_{x_j} [\sigma_{-2m}(D^{-2m})] \\ &\quad - \frac{1}{2} \sum_{jl} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(AB)] \partial_{x_j} \partial_{x_l} [\sigma_{-2m}(D^{-2m})]. \end{aligned} \quad (3.2)$$

Lemma 3.1. [12] Let S be pseudo-differential operator of order k and f is a smooth function, $[S, f]$ is a pseudo-differential operator of order $k-1$ with total symbol $\sigma[S, f] \sim \sum_{j \geq 1} \sigma_{k-j}[S, f]$, where

$$\sigma_{k-j}[S, f] = \sum_{|\beta|=1}^j \frac{D_x^\beta(f)}{\beta!} \partial_\xi^\beta (\sigma_{k-(j-|\beta|)}^S). \quad (3.3)$$

By Lemma 2.1 and Lemma 3.1, we get the following lemma.

Lemma 3.2. *The symbols of A and B are given*

$$\begin{aligned}
\sigma_0(A) &= \sum_{j=1}^n \partial_{x_j}(a^1)(\Gamma^j - 2\sigma^j)(x) - \sum_{jl=1}^n \partial_{x_j} \partial_{x_l}(a^1)g^{jl}; \\
\sigma_1(A) &= -2i \sum_{jl=1}^n \partial_{x_j}(a^1)g^{jl}\xi_l; \\
\sigma_0(B) &= \left(\sum_{j=1}^n \partial_{x_j}(a^2)(\Gamma^j - 2\sigma^j)(x) - \sum_{jl=1}^n \partial_{x_j} \partial_{x_l}(a^2)g^{jl} \right) c(da^3)c(da^4) - 2 \sum_{jl=1}^n \partial_{x_l}(a^2)g^{jl}\partial_{x_j}[c(da^3)c(da^4)]; \\
\sigma_1(B) &= -2i \sum_{jl=1}^n \partial_{x_j}(a^2)g^{jl}\xi_l c(da^3)c(da^4).
\end{aligned} \tag{3.4}$$

Further, by the composition formula of pseudodifferential operators, we get the following lemma.

Lemma 3.3. *The symbols of AB are given*

$$\begin{aligned}
\sigma_0(AB) &= \sigma_0(A)\sigma_0(B) + (-i)\partial_{\xi_j}[\sigma_1(A)]\partial_{x_j}[\sigma_0(B)] + (-i)\partial_{\xi_j}[\sigma_0(A)]\partial_{x_j}[\sigma_1(B)] \\
&= \left(\sum_{k=1}^n \partial_{x_k}(a^1)(\Gamma^k - 2\sigma^k)(x) - \sum_{kl=1}^n \partial_{x_k} \partial_{x_l}(a^1)g^{jl} \right) \times \left[\left(\sum_{j=1}^n \partial_{x_j}(a^2)(\Gamma^j - 2\sigma^j)(x) \right. \right. \\
&\quad \left. \left. - \sum_{jl=1}^n \partial_{x_j} \partial_{x_l}(a^2)g^{jl} \right) c(da^3)c(da^4) - 2 \sum_{jl=1}^n \partial_{x_l}(a^2)g^{jl}\partial_{x_j}[c(da^3)c(da^4)] \right] - 2 \sum_{jl=1}^n \left\{ \partial_{x_j}(a^1)g^{jl} \right. \\
&\quad \left[\left(\sum_{r=1}^n \partial_{x_j} \partial_{x_r}(a^2)(\Gamma^r - 2\sigma^r)(x) + \sum_{r=1}^n \partial_{x_r}(a^2)\partial_{x_j}(\Gamma^r - 2\sigma^r)(x) - \sum_{rs=1}^n \partial_{x_j} \partial_{x_r} \partial_{x_s}(a^2)g^{rs} \right. \right. \\
&\quad \left. \left. - \sum_{rs=1}^n \partial_{x_r} \partial_{x_s}(a^2)\partial_{x_j}g^{rs} \right) c(da^3)c(da^4) + \left(\sum_{j=1}^n \partial_{x_j}(a^2)(\Gamma^j - 2\sigma^j)(x) - \sum_{jl=1}^n \partial_{x_j} \partial_{x_l}(a^2)g^{jl} \right) \right. \\
&\quad \left. \partial_{x_j}[c(da^3)c(da^4)] - 2 \sum_{pl=1}^n \partial_{x_j} \partial_{x_l}(a^2)g^{pl}\partial_{x_p}[c(da^3)c(da^4)] - 2 \sum_{pl=1}^n \partial_{x_l}(a^2)g^{pl}\partial_{x_j} \partial_{x_p}[c(da^3)c(da^4)] \right] \\
&\quad \left. - 2 \sum_{jlp} \partial_{x_l}(a^2)\partial_{x_j}(g^{pl})\partial_{x_p}[c(da^3)c(da^4)] \right\};
\end{aligned}$$

$$\begin{aligned}
\sigma_1(AB) &= \sigma_1(A)\sigma_0(B) + \sigma_0(A)\sigma_1(B) + (-i)\partial_{\xi_j}[\sigma_1(A)]\partial_{x_j}[\sigma_1(B)] \\
&= -2i \sum_{pk=1}^n \partial_{x_p}(a^1)g^{pk}\xi_p \left(\sum_{r=1}^n \partial_{x_r}(a^2)(\Gamma^r - 2\sigma^r)(x) - \sum_{rs=1}^n \partial_{x_r} \partial_{x_s}(a^2)g^{rs} \right) c(da^3)c(da^4) \\
&\quad + 4i \sum_{pk=1}^n \partial_{x_p}(a^1)g^{pk}\xi_p \left(\sum_{ql=1}^n \partial_{x_l}(a^2)g^{ql}\partial_{x_q}[c(da^3)c(da^4)] \right) \\
&\quad - 2i \left(\sum_{k=1}^n \partial_{x_k}(a^1)(\Gamma^k - 2\sigma^k)(x) - \sum_{kl=1}^n \partial_{x_k} \partial_{x_l}(a^1)g^{kl} \right) \sum_{ts=1}^n \partial_{x_t}(a^2)g^{ts}\xi_t c(da^3)c(da^4) \\
&\quad + 4i \sum_{j\alpha=1}^n \partial_{x_\alpha}(a^1)g^{\alpha j} \sum_{ts=1}^n \partial_{x_j} \partial_{x_t}(a^2)g^{ts}\xi_t c(da^3)c(da^4) \\
&\quad + 4i \sum_{j\alpha=1}^n \partial_{x_\alpha}(a^1)g^{\alpha j} \sum_{ts=1}^n \partial_{x_t}(a^2)g^{ts}\xi_t \partial_{x_j}[c(da^3)c(da^4)]
\end{aligned}$$

$$\begin{aligned}
& + 4i \sum_{j\alpha=1}^n \partial_{x_\alpha}(a^1) g^{\alpha j} \sum_{ts=1}^n \partial_{x_t}(a^2) \partial_{x_j} g^{ts} \xi_t c(da^3) c(da^4); \\
\sigma_2(AB) & = \sigma_1(A)\sigma_1(B) \\
& = -4 \sum_{jlkst=1}^n \partial_{x_j}(a^1) \partial_{x_l}(a^2) \xi_j \xi_l g^{jk} g^{ls} c(da^3) c(da^4). \tag{3.5}
\end{aligned}$$

By Lemma 2.1, (4.24) in [10] and (3.4) in [14], we get

Lemma 3.4. General dimensional symbols about Dirac operator D are given,

$$\begin{aligned}
\sigma_{-2m}(D^{-2m}) & = |\xi|^{-2m}; \\
\sigma_{-2m-1}(D^{-2m}) & = m|\xi|^{-(2m-2)} \left(-i|\xi|^{-4}\xi_k(\Gamma^k - 2\sigma^k) - 2i|\xi|^{-6}\xi^j\xi_\alpha\xi_\beta\partial_{x_j}g^{\alpha\beta} \right) \\
& + 2i \sum_{k=0}^{m-2} \sum_{\mu=1}^{2m} (k+1-m)|\xi|^{-2m-4}\xi^\mu\xi_\alpha\xi_\beta\partial_\mu^x g^{\alpha\beta}; \\
\sigma_{-2m-2}(D^{-2m})(x_0) & = -\frac{m}{4}|\xi|^{-2m-2}s + \frac{m(m+1)}{3}|\xi|^{-2m-4}\xi_\mu\xi_\nu R_{\mu\alpha\nu\alpha}(x_0), \tag{3.6}
\end{aligned}$$

where s is the scalar curvature.

Proof. See the Appendix. \square

Next, we review here technical tool of the computation, which are the integrals of polynomial functions over the unit spheres. By (32) in [2], we define

$$I_{S_n}^{\gamma_1 \dots \gamma_{2\bar{n}+2}} = \int_{|x|=1} d^n x x^{\gamma_1} \dots x^{\gamma_{2\bar{n}+2}}, \tag{3.7}$$

i.e. the monomial integrals over a unit sphere. Then by Proposition A.2. in [2], polynomial integrals over higher spheres in the n -dimesional case are given

$$I_{S_n}^{\gamma_1 \dots \gamma_{2\bar{n}+2}} = \frac{1}{2\bar{n}+n} [\delta^{\gamma_1 \gamma_2} I_{S_n}^{\gamma_3 \dots \gamma_{2\bar{n}+2}} + \dots + \delta^{\gamma_1 \gamma_{2\bar{n}+1}} I_{S_n}^{\gamma_2 \dots \gamma_{2\bar{n}+1}}], \tag{3.8}$$

where $S_n \equiv S^{n-1}$ in \mathbb{R}^n .

For $\bar{n} = 0$, we have $I^0 = \text{area}(S_n) = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})}$, we immediately get

$$\begin{aligned}
I_{S_n}^{\gamma_1 \gamma_2} & = \frac{1}{n} \text{area}(S_n) \delta^{\gamma_1 \gamma_2}; \\
I_{S_n}^{\gamma_1 \gamma_2 \gamma_3 \gamma_4} & = \frac{1}{n(n+2)} \text{area}(S_n) [\delta^{\gamma_1 \gamma_2} \delta^{\gamma_3 \gamma_4} + \delta^{\gamma_1 \gamma_3} \delta^{\gamma_2 \gamma_4} + \delta^{\gamma_1 \gamma_4} \delta^{\gamma_2 \gamma_3}]; \\
I_{S_n}^{\gamma_1 \gamma_2 \gamma_3 \gamma_4 \gamma_5 \gamma_6} & = \frac{1}{n(n+2)(n+4)} \text{area}(S_n) [\delta^{\gamma_1 \gamma_2} (\delta^{\gamma_3 \gamma_4} \delta^{\gamma_5 \gamma_6} + \delta^{\gamma_3 \gamma_5} \delta^{\gamma_4 \gamma_6} + \delta^{\gamma_3 \gamma_6} \delta^{\gamma_4 \gamma_5}) \\
& + \delta^{\gamma_1 \gamma_3} (\delta^{\gamma_2 \gamma_4} \delta^{\gamma_5 \gamma_6} + \delta^{\gamma_2 \gamma_5} \delta^{\gamma_4 \gamma_6} + \delta^{\gamma_2 \gamma_6} \delta^{\gamma_4 \gamma_5}) + \delta^{\gamma_1 \gamma_4} (\delta^{\gamma_2 \gamma_3} \delta^{\gamma_5 \gamma_6} + \delta^{\gamma_2 \gamma_5} \delta^{\gamma_3 \gamma_6} + \delta^{\gamma_2 \gamma_6} \delta^{\gamma_3 \gamma_5}) \\
& + \delta^{\gamma_1 \gamma_5} (\delta^{\gamma_2 \gamma_3} \delta^{\gamma_4 \gamma_6} + \delta^{\gamma_2 \gamma_4} \delta^{\gamma_3 \gamma_6} + \delta^{\gamma_2 \gamma_6} \delta^{\gamma_3 \gamma_4}) + \delta^{\gamma_1 \gamma_6} (\delta^{\gamma_2 \gamma_3} \delta^{\gamma_4 \gamma_5} + \delta^{\gamma_2 \gamma_4} \delta^{\gamma_3 \gamma_5} + \delta^{\gamma_2 \gamma_5} \delta^{\gamma_3 \gamma_4})]. \tag{3.9}
\end{aligned}$$

Next, we calculate each term of $\int_{S^* M} \text{tr}[\sigma_{-2m}(ABD^{-2m})](x, \xi)$ separately.

(i-1) For $\sigma_0(AB)\sigma_{-2m}(D^{-2m})$:

Using the facts:

$$\Gamma^\mu(x_0) = \sigma^\mu(x_0) = \partial_\mu^x g^{\alpha\beta}(x_0) = 0, \quad \partial_{x_k}(\Gamma^\mu)(x_0) = g^{\alpha\beta} \partial_{x_k}(\Gamma^\mu_{\alpha\beta})(x_0) = \frac{2}{3} R_{k\alpha\mu\alpha}(x_0);$$

$$\partial_{x_k}(\sigma^\mu)(x_0) = \frac{1}{4} \sum_{st=1}^n \partial_{x_k}(<\nabla_{\partial_\mu}^L e_s, e_t>)(x_0) c(e_s)c(e_t) = \frac{1}{8} \sum_{st=1}^n R_{k\mu ts}(x_0) c(e_s)c(e_t). \quad (3.10)$$

In normal coordinates, we get the following result.

$$\begin{aligned} & \sigma_0(AB)\sigma_{-2m}(D^{-2m})(x_0) \\ &= \sum_{kr} \partial_{x_k}^2(a^1) \partial_{x_r}^2(a^2) c(da^3)c(da^4)|\xi|^{-2m} + 2 \sum_{kp} \partial_{x_k}^2(a^1) \partial_{x_p}(a^2) \partial_{x_p}[c(da^3)c(da^4)]|\xi|^{-2m} \\ &- \frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^1) \partial_{x_r}(a^2) R_{j\alpha r\alpha}(x_0) c(da^3)c(da^4)|\xi|^{-2m} + 2 \sum_{jr} \partial_{x_j}(a^1) \partial_{x_r}^2(a^2) \partial_{x_j}[c(da^3)c(da^4)]|\xi|^{-2m} \\ &+ \frac{1}{2} \sum_{jrst} \partial_{x_j}(a^1) \partial_{x_r}(a^2) R_{jrt s}(x_0) c(e_s)c(e_t)c(da^3)c(da^4)|\xi|^{-2m} + 2 \sum_{jr} \partial_{x_j}(a^1) \partial_{x_j} \partial_{x_r}^2(a^2) c(da^3)c(da^4)|\xi|^{-2m} \\ &+ 2 \sum_{jr} \partial_{x_j}(a^1) \partial_{x_j} \partial_{x_r}^2(a^2) c(da^3)c(da^4)|\xi|^{-2m} + 4 \sum_{jp} \partial_{x_j}(a^1) \partial_{x_j} \left(\partial_{x_p}(a^2) \partial_{x_p}[c(da^3)c(da^4)] \right) |\xi|^{-2m}. \quad (3.11) \end{aligned}$$

(i - 1 - a)

$$\begin{aligned} \text{tr} \left(\sum_{kr} \partial_{x_k}^2(a^1) \partial_{x_r}^2(a^2) c(da^3)c(da^4)|\xi|^{-2m} \right) |_{|\xi|=1} &= \sum_{kr} \partial_{x_k}^2(a^1) \partial_{x_r}^2(a^2) \text{tr}[c(da^3)c(da^4)] \\ &= - \sum_{kr} \partial_{x_k}^2(a^1) \partial_{x_r}^2(a^2) g(da^3, da^4) \text{tr}[id], \end{aligned}$$

then

$$\begin{aligned} \int_{|\xi|=1} \text{tr} \left(\sum_{kr} \partial_{x_k}^2(a^1) \partial_{x_r}^2(a^2) c(da^3)c(da^4)|\xi|^{-2m} \right) \sigma(\xi) &= - \sum_{kr} \partial_{x_k}^2(a^1) \partial_{x_r}^2(a^2) g(da^3, da^4) \text{tr}[id] \text{area}(S_n) \\ &= -\Delta(a^1) \Delta(a^2) g(da^3, da^4) \text{tr}[id] \text{area}(S_n), \end{aligned}$$

where $\Delta(a^1)$ denotes a generalized laplacian of a^1 .

(i - 1 - b)

$$\text{tr} \left(2 \sum_{kp} \partial_{x_k}^2(a^1) \partial_{x_p}(a^2) \partial_{x_p}[c(da^3)c(da^4)]|\xi|^{-2m} \right) |_{|\xi|=1} = -2 \sum_{kp} \partial_{x_k}^2(a^1) \partial_{x_p}(a^2) \partial_{x_p}[g(da^3, da^4)] \text{tr}[id],$$

then

$$\begin{aligned} \int_{|\xi|=1} \text{tr} \left(2 \sum_{kp} \partial_{x_k}^2(a^1) \partial_{x_p}(a^2) \partial_{x_p}[c(da^3)c(da^4)]|\xi|^{-2m} \right) \sigma(\xi) &= -2 \sum_{kp} \partial_{x_k}^2(a^1) \partial_{x_p}(a^2) \partial_{x_p}[g(da^3, da^4)] \text{tr}[id] \text{area}(S_n) \\ &= -2\Delta(a^1) g(\nabla(a^2), \nabla g(da^3, da^4)) \text{tr}[id] \text{area}(S_n), \end{aligned}$$

where $\nabla(a^2)$ denotes the gradient of a^2 .

(i - 1 - c)

$$\text{tr} \left(-\frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^1) \partial_{x_r}(a^2) R_{j\alpha r\alpha}(x_0) c(da^3)c(da^4)|\xi|^{-2m} \right) |_{|\xi|=1} = \frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^1) \partial_{x_r}(a^2) R_{j\alpha r\alpha}(x_0) g(da^3, da^4) \text{tr}[id],$$

then

$$\int_{|\xi|=1} \text{tr} \left(-\frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^1) \partial_{x_r}(a^2) R_{j\alpha r\alpha}(x_0) c(da^3)c(da^4)|\xi|^{-2m} \right) \sigma(\xi)$$

$$\begin{aligned}
&= \frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^1) \partial_{x_r}(a^2) R_{j\alpha r\alpha}(x_0) g(da^3, da^4) \text{tr}[id] \text{area}(S_n) \\
&= \frac{4}{3} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^2), e_{\alpha}) g(da^3, da^4) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

($\mathbf{i} - \mathbf{1} - \mathbf{d}$)

$$\text{tr} \left(2 \sum_{jp} \partial_{x_j}(a^1) \partial_{x_p}^2(a^2) \partial_{x_j} [c(da^3)c(da^4)] |\xi|^{-2m} \right) |_{|\xi|=1} = -2 \sum_{jp} \partial_{x_j}(a^1) \partial_{x_p}^2(a^2) \partial_{x_j} [g(da^3, da^4)] \text{tr}[id],$$

then

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(2 \sum_{jp} \partial_{x_j}(a^1) \partial_{x_p}^2(a^2) \partial_{x_j} [c(da^3)c(da^4)] |\xi|^{-2m} \right) \sigma(\xi) \\
&= -2 \sum_{jp} \partial_{x_j}(a^1) \partial_{x_p}^2(a^2) \partial_{x_j} [g(da^3, da^4)] \text{tr}[id] \text{area}(S_n) \\
&= -2 \Delta(a^2) g(\nabla(a^1), \nabla g(da^3, da^4)) \text{tr}[id] \text{area}(S_n),
\end{aligned}$$

($\mathbf{i} - \mathbf{1} - \mathbf{e}$)

$$\begin{aligned}
&\text{tr} \left(\sum_{jr} \partial_{x_j}(a^1) \partial_{x_r}(a^2) \sum_{st} R_{jrts}(x_0) c(e_s) c(e_t) c(da^3) c(da^4) |\xi|^{-2m} \right) |_{|\xi|=1} \\
&= \sum_{jr} \partial_{x_j}(a^1) \partial_{x_r}(a^2) \sum_{st} R_{jrts}(x_0) \text{tr}[c(e_s) c(e_t) c(da^3) c(da^4)],
\end{aligned}$$

where

$$\begin{aligned}
\text{tr}[c(e_s) c(e_t) c(da^3) c(da^4)] &= \sum_{\alpha\beta} e_{\alpha}(a^3) e_{\beta}(a^4) \text{tr}[c(e_s) c(e_t) c(e_{\alpha}) c(e_{\beta})] \\
&= \sum_{\alpha\beta} e_{\alpha}(a^3) e_{\beta}(a^4) \left(-\delta_{st} \text{tr}[c(e_{\alpha}) c(e_{\beta})] + \delta_{s\alpha} \text{tr}[c(e_t) c(e_{\beta})] - \delta_{s\beta} \text{tr}[c(e_t) c(e_{\alpha})] \right),
\end{aligned}$$

then

$$\begin{aligned}
R_{jrts}(x_0) \text{tr}[c(e_s) c(e_t) c(da^3) c(da^4)] &= -R_{jrts}(x_0) e_s(a^3) e_t(a^4) \text{tr}[id] + R_{jrts}(x_0) e_t(a^3) e_s(a^4) \text{tr}[id] \\
&= -2R_{jrts}(x_0) e_s(a^3) e_t(a^4) \text{tr}[id].
\end{aligned}$$

Further,

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(\frac{1}{2} \sum_{jr} \partial_{x_j}(a^1) \partial_{x_r}(a^2) \sum_{st} R_{jrts}(x_0) c(e_s) c(e_t) c(da^3) c(da^4) |\xi|^{-2m} \right) \sigma(\xi) \\
&= - \sum_{jrts} \partial_{x_j}(a^1) \partial_{x_r}(a^2) R_{jrts}(x_0) e_s(a^3) e_t(a^4) \text{tr}[id] \text{area}(S_n) \\
&= -R(\nabla(a^1), \nabla(a^2), \nabla(a^4), \nabla(a^3)) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

($\mathbf{i} - \mathbf{1} - \mathbf{f}$) (See Appendix)

$$\int_{|\xi|=1} \text{tr} \left(2 \sum_{jr} \partial_{x_j}(a^1) \partial_{x_r} \partial_{x_r}^2(a^2) c(da^3) c(da^4) |\xi|^{-2m} \right) \sigma(\xi)$$

$$\begin{aligned}
&= -2 \sum_{jr} \partial_{x_j}(a^1) \partial_{x_j} \partial_{x_r}^2(a^2) g(da^3, da^4) \text{tr}[id] \text{area}(S_n) \\
&= \left(2\nabla(a^1)[\Delta(a^2)] - \frac{4}{3} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^2), e_{\alpha}) \right) g(da^3, da^4) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(**i - 1 - g**) (See Appendix)

$$\int_{|\xi|=1} \text{tr} \left[4 \sum_{jp} \partial_{x_j}(a^1) \partial_{x_j} \left(\partial_{x_p}(a^2) \partial_{x_p} [c(da^3)c(da^4)] \right) |\xi|^{-2m} \right] \sigma(\xi) = -4 \nabla(a^1) \nabla(a^2) [g(da^3, da^4)] \text{tr}[id] \text{area}(S_n).$$

Therefore, we get

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(\sigma_0(AB) \sigma_{-2m}(D^{-2m})(x_0) \right) \sigma(\xi) \\
&= \left(-\Delta(a^1)\Delta(a^2)g(da^3, da^4) - 2\Delta(a^1)g(\nabla(a^2), \nabla g(da^3, da^4)) - R(\nabla(a^1), \nabla(a^2), \nabla(a^4), \nabla(a^3)) \right. \\
&\quad \left. + 2\nabla(a^1)[\Delta(a^2)]g(da^3, da^4) - 4\nabla(a^1)\nabla(a^2)[g(da^3, da^4)] \right) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(**i - 2**) For $\sigma_1(AB)\sigma_{-2m-1}(D^{-2m})$:

By (3.10), in normal coordinates, we get

$$\begin{aligned}
&\sigma_{-2m-1}(D^{-2m})(x_0) \\
&= m|\xi|^{-(2m-2)} \left(-i|\xi|^{-4}\xi_k(\Gamma^k - 2\sigma^k)(x_0) - 2i|\xi|^{-6}\xi^j\xi_{\alpha}\xi_{\beta}\partial_{x_j}g^{\alpha\beta}(x_0) \right) \\
&\quad + 2i \sum_{k=0}^{m-2} \sum_{\mu=1}^{2m} (k+1-m)|\xi|^{-2m-4}\xi^{\mu}\xi_{\alpha}\xi_{\beta}\partial_{\mu}^xg^{\alpha\beta}(x_0) \\
&= 0,
\end{aligned} \tag{3.12}$$

therefore

$$\int_{|\xi|=1} \text{tr} \left(\sigma_1(AB) \sigma_{-2m-1}(D^{-2m})(x_0) \right) \sigma(\xi) = 0.$$

(**i - 3**) For $\sigma_2(AB)\sigma_{-2m-2}(D^{-2m})$:

$$\begin{aligned}
&\sigma_2(AB)\sigma_{-2m-2}(D^{-2m})(x_0) \\
&= m \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \xi_j \xi_l c(da^3)(da^4) |\xi|^{-2m-2} s \\
&\quad - \frac{4m(m+1)}{3} \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) \xi_j \xi_l \xi_{\mu} \xi_{\nu} c(da^3)c(da^4) |\xi|^{-2m-4}.
\end{aligned} \tag{3.13}$$

(**i - 3 - a**)

$$\text{mtr} \left(\sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \xi_j \xi_l c(da^3)c(da^4) |\xi|^{-2m-2} s \right) |_{|\xi|=1} = -m \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \xi_j \xi_l g(da^3, da^4) \text{str}[id],$$

then

$$\int_{|\xi|=1} \text{mtr} \left(\sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \xi_j \xi_l c(da^3)c(da^4) |\xi|^{-2m-2} s \right) \sigma(\xi)$$

$$\begin{aligned}
&= -m \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) g(da^3, da^4) \text{str}[id] \int_{|\xi|=1} \xi_j \xi_l \sigma(\xi) \\
&= -m \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) g(da^3, da^4) \text{str}[id] I_{S_n}^{jl} \\
&= -\frac{1}{2} \sum_j \partial_{x_j}(a^1) \partial_{x_j}(a^2) \text{area}(S_n) g(da^3, da^4) \text{str}[id] \\
&= -\frac{1}{2} g(\nabla(a^1), \nabla(a^2)) \text{area}(S_n) g(da^3, da^4) \text{str}[id].
\end{aligned}$$

(i - 3 - b)

$$\begin{aligned}
&\text{tr} \left(-\frac{4m(m+1)}{3} \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) \xi_j \xi_l \xi_\mu \xi_\nu c(da^3) c(da^4) |\xi|^{-2m-4} \right) |_{|\xi|=1} \\
&= \frac{4m(m+1)}{3} \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) \xi_j \xi_l \xi_\mu \xi_\nu g(da^3, da^4) \text{tr}[id],
\end{aligned}$$

then

$$\begin{aligned}
&\int_{|\xi|=1} -\frac{4m(m+1)}{3} \text{tr} \left(\sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) \xi_j \xi_l \xi_\mu \xi_\nu c(da^3) c(da^4) |\xi|^{-2m-4} \right) \sigma(\xi) \\
&= \frac{4m(m+1)}{3} \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) g(da^3, da^4) \text{tr}[id] \int_{|\xi|=1} \xi_j \xi_l \xi_\mu \xi_\nu \sigma(\xi) \\
&= \frac{4m(m+1)}{3} \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) g(da^3, da^4) \text{tr}[id] I_{S_n}^{jl\mu\nu} \\
&= \frac{1}{3} \left(\sum_{j\mu\alpha} \partial_{x_j}(a^1) \partial_{x_j}(a^2) R_{\mu\alpha\mu\alpha}(x_0) + 2 \sum_{jl\alpha} \partial_{x_j}(a^1) \partial_{x_l}(a^2) R_{j\alpha l\alpha}(x_0) \right) \text{area}(S_n) g(da^3, da^4) \text{tr}[id] \\
&= \frac{1}{3} \left(\sum_{\mu\alpha} g(\nabla(a^1), \nabla(a^2)) R_{\mu\alpha\mu\alpha}(x_0) + 2 \sum_{\alpha} R(\nabla(a^1), e_\alpha, \nabla(a^2), e_\alpha) \right) \text{area}(S_n) g(da^3, da^4) \text{tr}[id] \\
&= \frac{1}{3} \left(g(\nabla(a^1), \nabla(a^2)) s + 2 \sum_{\alpha} R(\nabla(a^1), e_\alpha, \nabla(a^2), e_\alpha) \right) \text{area}(S_n) g(da^3, da^4) \text{tr}[id].
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(\sigma_2(AB) \sigma_{-2m-2}(D^{-2m})(x_0) \right) \sigma(\xi) \\
&= \left(-\frac{1}{6} g(\nabla(a^1), \nabla(a^2)) s + \frac{2}{3} \sum_{\alpha} R(\nabla(a^1), e_\alpha, \nabla(a^2), e_\alpha) \right) \text{area}(S_n) g(da^3, da^4) \text{tr}[id].
\end{aligned}$$

(i - 4) For $-i \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(AB)] \partial_{x_j} [\sigma_{-2m-1}(D^{-2m})]$:

$$\begin{aligned}
&-i \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(AB)] \partial_{x_j} [\sigma_{-2m-1}(D^{-2m})](x_0) \\
&= \frac{8m}{3} \sum_{jplk\alpha} \partial_{x_p}(a^1) \partial_{x_l}(a^2) [\partial_{\xi_j}(\xi_p) \xi_l \xi_k + \partial_{\xi_j}(\xi_l) \xi_p \xi_k] R_{j\alpha k\alpha}(x_0) c(da^3) c(da^4) |\xi|^{-2m-2} \\
&- m \sum_{jplkts} \partial_{x_p}(a^1) \partial_{x_l}(a^2) [\partial_{\xi_j}(\xi_p) \xi_l \xi_k + \partial_{\xi_j}(\xi_l) \xi_p \xi_k] R_{j\alpha k\alpha}(x_0) c(da^3) c(da^4) c(e_s) c(e_t) |\xi|^{-2m-2}
\end{aligned}$$

$$\begin{aligned}
& + \frac{16}{3}m \sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^2) R_{\alpha j\beta r}(x_0) [\partial_{\xi_j}(\xi_p)\xi_l\xi_r\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_r\xi_\alpha\xi_\beta] c(da^3)c(da^4)|\xi|^{-2m-4} \\
& + \frac{16(m-k-1)}{3} \sum_{k=0}^{m-2} \sum_{\mu=1}^{2m} \sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^2) R_{\alpha j\beta\mu}(x_0) [\partial_{\xi_j}(\xi_p)\xi_l\xi_\mu\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_\mu\xi_\alpha\xi_\beta] c(da^3)c(da^4)|\xi|^{-2m-4}.
\end{aligned}$$

(i - 4 - a)

$$\begin{aligned}
& \text{tr} \left(\frac{8}{3}m \sum_{jplk\alpha} \partial_{x_p}(a^1) \partial_{x_l}(a^2) [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] R_{j\alpha k\alpha}(x_0) c(da^3)c(da^4)|\xi|^{-2m-2} \right) |_{|\xi|=1} \\
& = -\frac{8}{3}m \sum_{jplk\alpha} \partial_{x_p}(a^1) \partial_{x_l}(a^2) [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] R_{j\alpha k\alpha}(x_0) g(da^3, da^4) \text{tr}[id],
\end{aligned}$$

then

$$\begin{aligned}
& \int_{|\xi|=1} \text{tr} \left(\frac{8}{3}m \sum_{jplk\alpha} \partial_{x_p}(a^1) \partial_{x_l}(a^2) [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] R_{j\alpha k\alpha}(x_0) c(da^3)c(da^4)|\xi|^{-2m-2} \right) \sigma(\xi) \\
& = -\frac{8}{3}m \sum_{jplk\alpha} \partial_{x_p}(a^1) \partial_{x_l}(a^2) R_{j\alpha k\alpha}(x_0) g(a^3, a^4) \text{tr}[id] \int_{|\xi|=1} [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] \sigma(\xi) \\
& = -\frac{4}{3(m+1)} \sum_{plk\alpha} \partial_{x_p}(a^1) \partial_{x_l}(a^2) R_{p\alpha k\alpha}(x_0) I_{S_n}^{lk} \text{area}(S_n) g(da^3, da^4) \text{tr}[id] \\
& = -\frac{4}{3(m+1)} \sum_{pk\alpha} \partial_{x_p}(a^1) \partial_{x_k}(a^2) R_{p\alpha k\alpha}(x_0) \text{area}(S_n) g(da^3, da^4) \text{tr}[id] \\
& = -\frac{4}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^2), e_{\alpha}) \text{area}(S_n) g(da^3, da^4) \text{tr}[id].
\end{aligned}$$

(i - 4 - b)

$$\begin{aligned}
& \text{tr} \left(-m \sum_{jplkts} \partial_{x_p}(a^1) \partial_{x_l}(a^2) [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] R_{jkts}(x_0) c(da^3)c(da^4)c(e_s)c(e_t)|\xi|^{-2m-2} \right) |_{|\xi|=1} \\
& = 2m \sum_{jplkts} \partial_{x_p}(a^1) \partial_{x_l}(a^2) [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] R_{jkts}(x_0) e_s(da^3)e_t(da^4) \text{tr}[id],
\end{aligned}$$

then

$$\begin{aligned}
& \int_{|\xi|=1} \text{tr} \left(-m \sum_{jplkts} \partial_{x_p}(a^1) \partial_{x_l}(a^2) [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] R_{jkts}(x_0) c(da^3)c(da^4)c(e_s)c(e_t)|\xi|^{-2m-2} \right) \sigma(\xi) \\
& = \frac{1}{2m+2} \text{area}(S^n) \sum_{jlt} \partial_{x_j}(a^1) \partial_{x_l}(a^2) R_{jlt}(x_0) e_s(a^3)e_t(a^4) \text{tr}[id] \\
& + \frac{1}{2m+2} \text{area}(S^n) \sum_{jpts} \partial_{x_p}(a^1) \partial_{x_j}(a^2) R_{jpts}(x_0) e_s(a^3)e_t(a^4) \text{tr}[id] \\
& = \frac{1}{2m+2} \text{area}(S^n) \left(R(\nabla(a^1), \nabla(a^2), \nabla(a^4), \nabla(a^3)) + R(\nabla(a^2), \nabla(a^1), \nabla(a^4), \nabla(a^3)) \right) \text{tr}[id] \\
& = 0.
\end{aligned}$$

(i - 4 - c)

$$\text{tr} \left(\sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^2) R_{\alpha j\beta r}(x_0) [\partial_{\xi_j}(\xi_p)\xi_l\xi_r\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_r\xi_\alpha\xi_\beta] c(da^3)c(da^4)|\xi|^{-2m-4} \right) |_{|\xi|=1}$$

$$= - \sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^2) R_{\alpha j\beta r}(x_0) [\partial_{\xi_j}(\xi_p)\xi_l\xi_r\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_r\xi_\alpha\xi_\beta] c(da^3)c(da^4)|\xi|^{-2m-4} g(da^3,da^4) \text{tr}[id],$$

then

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(\sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^2) R_{\alpha j\beta r}(x_0) [\partial_{\xi_j}(\xi_p)\xi_l\xi_r\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_r\xi_\alpha\xi_\beta] c(da^3)c(da^4)|\xi|^{-2m-4} \right) \sigma(\xi) \\ &= - \sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^2) R_{\alpha j\beta r}(x_0) g(da^3,da^4) \text{tr}[id] \int_{|\xi|=1} [\partial_{\xi_j}(\xi_p)\xi_l\xi_r\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_r\xi_\alpha\xi_\beta] \sigma(\xi) \\ &= - \left(\sum_{jlr\alpha\beta} \partial_{x_j}(a^1) \partial_{x_l}(a^2) R_{\alpha j\beta r}(x_0) I_{S_n}^{l^{r\alpha\beta}} + \sum_{jlr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_j}(a^2) R_{\alpha j\beta r}(x_0) I_{S_n}^{p^{r\alpha\beta}} \right) g(da^3,da^4) \text{tr}[id] \\ &= 0. \end{aligned}$$

(**i – 4 – d**)

Similarly, we get

$$\int_{|\xi|=1} \text{tr} \left[\sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^2) R_{\alpha j\beta\mu}(x_0) [\partial_{\xi_j}(\xi_p)\xi_l\xi_\mu\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_\mu\xi_\alpha\xi_\beta] c(da^3)c(da^4)|\xi|^{-2m-4} \right] \sigma(\xi) = 0.$$

Therefore, we get

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(-i \sum_{j=1}^n \partial_{\xi_j}[\sigma_2(AB)] \partial_{x_j}[\sigma_{-2m-1}(D^{-2m})](x_0) \right) \sigma(\xi) \\ &= -\frac{4}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^2), e_{\alpha}) \text{area}(S_n) g(da^3,da^4) \text{tr}[id]. \end{aligned}$$

(**i – 5**) For $-i \sum_{j=1}^n \partial_{\xi_j}[\sigma_1(AB)] \partial_{x_j}[\sigma_{-2m}(D^{-2m})]$:

$$\partial_{x_j}(|\xi|^{-2m})(x_0) \sigma(\xi) = 0, \quad (3.14)$$

then

$$\int_{|\xi|=1} \text{tr} \left(-i \sum_{j=1}^n \partial_{\xi_j}[\sigma_1(AB)] \partial_{x_j}[\sigma_{-2m}(D^{-2m})] \right) (x_0) \sigma(\xi) = 0.$$

(**i – 6**) For $-\frac{1}{2} \sum_{jl} \partial_{\xi_j} \partial_{\xi_l}[\sigma_2(AB)] \partial_{x_j} \partial_{x_l}[\sigma_{-2m}(D^{-2m})]$:

$$\begin{aligned} & -\frac{1}{2} \sum_{jl} \partial_{\xi_j} \partial_{\xi_l}[\sigma_2(AB)] \partial_{x_j} \partial_{x_l}[\sigma_{-2m}(D^{-2m})](x_0) \\ &= -\frac{4}{3}m \sum_{jlr\alpha\beta} \partial_{x_r}(a^1) \partial_{x_s}(a^2) R_{\alpha j\beta l}(x_0) [\partial_{\xi_l}(\xi_r) \partial_{\xi_j}(\xi_s) + \partial_{\xi_l}(\xi_s) \partial_{\xi_j}(\xi_r)] \xi_{\alpha}\xi_{\beta} c(da^3)c(da^4)|\xi|^{-2m-2}, \end{aligned}$$

so

$$\begin{aligned} & \text{tr} \left(-\frac{4}{3}m \sum_{jlr\alpha\beta} \partial_{x_r}(a^1) \partial_{x_s}(a^2) R_{\alpha j\beta l}(x_0) [\partial_{\xi_l}(\xi_r) \partial_{\xi_j}(\xi_s) + \partial_{\xi_l}(\xi_s) \partial_{\xi_j}(\xi_r)] \xi_{\alpha}\xi_{\beta} c(da^3)c(da^4)|\xi|^{-2m-2} \right) |_{|\xi|=1} \\ &= \frac{4}{3}m \sum_{jlr\alpha\beta} \partial_{x_r}(a^1) \partial_{x_s}(a^2) R_{\alpha j\beta l}(x_0) [\partial_{\xi_l}(\xi_r) \partial_{\xi_j}(\xi_s) + \partial_{\xi_l}(\xi_s) \partial_{\xi_j}(\xi_r)] \xi_{\alpha}\xi_{\beta} g(da^3,da^4) \text{tr}[id], \end{aligned}$$

then

$$\int_{|\xi|=1} -\frac{4}{3}m \text{tr} \left(\sum_{jlr\alpha\beta} \partial_{x_r}(a^1) \partial_{x_s}(a^2) R_{\alpha j\beta l}(x_0) [\partial_{\xi_l}(\xi_r) \partial_{\xi_j}(\xi_s) + \partial_{\xi_l}(\xi_s) \partial_{\xi_j}(\xi_r)] \xi_{\alpha}\xi_{\beta} c(da^3)c(da^4)|\xi|^{-2m-2} \right) \sigma(\xi)$$

$$\begin{aligned}
&= \frac{4}{3}m \sum_{jlr s \alpha \beta} \partial_{x_r}(a^1) \partial_{x_s}(a^2) [\partial_{\xi_l}(\xi_r) \partial_{\xi_j}(\xi_s) + \partial_{\xi_l}(\xi_s) \partial_{\xi_j}(\xi_r)] R_{\alpha j \beta l}(x_0) g(da^3, da^4) \text{tr}[id] \int_{|\xi|=1} \xi_\alpha \xi_\beta \sigma(\xi) \\
&= \frac{4}{3}m \sum_{jlr s \alpha \beta} \partial_{x_r}(a^1) \partial_{x_s}(a^2) [\partial_{\xi_l}(\xi_r) \partial_{\xi_j}(\xi_s) + \partial_{\xi_l}(\xi_s) \partial_{\xi_j}(\xi_r)] R_{\alpha j \beta l}(x_0) g(da^3, da^4) \text{tr}[id] I_{S_n}^{\alpha \beta} \\
&= \frac{2}{3(m+1)} \sum_{jl\alpha} \partial_{x_j}(a^1) \partial_{x_l}(a^2) R_{\alpha j \alpha l}(x_0) \text{area}(S_n) g(da^3, da^4) \text{tr}[id] \\
&= \frac{2}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^2), e_{\alpha}) \text{area}(S_n) g(da^3, da^4) \text{tr}[id].
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(-\frac{1}{2} \sum_{jl} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(AB)] \partial_{x_j} \partial_{x_l} [\sigma_{-2m}(D^{-2m})](x_0) \right) \sigma(\xi) \\
&= \frac{2}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^2), e_{\alpha}) \text{area}(S_n) g(da^3, da^4) \text{tr}[id].
\end{aligned}$$

Finally, we obtain

Proposition 3.5. Let $P = [D^2, a^1][D^2, a^2][D, a^3][D, a^4]$, we have the following noncommutative integral of P based on the Wodzicki residue

$$\begin{aligned}
&\oint a^0 [D^2, a^1][D^2, a^2][D, a^3][D, a^4] D^{-2m} \\
&= 2^m \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \int_M a_0 \left(-\Delta(a^1)\Delta(a^2)g(da^3, da^4) - 2\Delta(a^1)g(\nabla(a^2), \nabla g(da^3, da^4)) - R(\nabla(a^1), \nabla(a^2), \nabla(a^4), \nabla(a^3)) \right. \\
&\quad + 2\nabla(a^1)[\Delta(a^2)]g(da^3, da^4) - 4\nabla(a^1)\nabla(a^2)[g(da^3, da^4)] - \frac{1}{6}g(\nabla(a^1), \nabla(a^2))g(da^3, da^4)s \\
&\quad \left. + \frac{2m}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^2), e_{\alpha})g(da^3, da^4) \right) d\text{Vol}_M.
\end{aligned}$$

Part ii) $P_1 = [D, a^1][D^2, a^2][D^2, a^3][D, a^4]$, $[D, a^1][D^2, a^2] := A_1$, $[D^2, a^3][D, a^4] := B_1$.
Similarly, by Lemma 3.1 in [12] and Lemma 2.1, we get the following lemma.

Lemma 3.6. The symbols of A_1 and B_1 are given

$$\begin{aligned}
\sigma_0(A_1) &= c(da^1) \left(\sum_{j=1}^n \partial_{x_j}(a^2)(\Gamma^j - 2\sigma^j)(x) - \sum_{jl=1}^n \partial_{x_j} \partial_{x_l}(a^2)g^{jl} \right); \\
\sigma_1(A_1) &= -2ic(da^1) \sum_{jl=1}^n \partial_{x_j}(a^2)g^{jl}\xi_l; \\
\sigma_0(B_1) &= \left(\sum_{j=1}^n \partial_{x_j}(a^3)(\Gamma^j - 2\sigma^j)(x) - \sum_{jl=1}^n \partial_{x_j} \partial_{x_l}(a^3)g^{jl} \right) c(da^4) - 2 \sum_{jl=1}^n \partial_{x_l}(a^3)g^{jl}\partial_{x_j}[c(da^4)]; \\
\sigma_1(B_1) &= -2i \sum_{jl=1}^n \partial_{x_j}(a^3)g^{jl}\xi_l c(da^4).
\end{aligned} \tag{3.15}$$

Next, we calculate each term of $\int_{S^* M} \text{tr}[\sigma_{-2m}(A_1 B_1 D^{-2m})](x, \xi)$ separately.
(ii - 1) For $\sigma_0(A_1 B_1) \sigma_{-2m}(D^{-2m})$:

$$\sigma_0(A_1 B_1) \sigma_{-2m}(D^{-2m})(x_0)$$

$$\begin{aligned}
&= \sum_{kr} \partial_{x_k}^2(a^2) \partial_{x_r}^2(a^3) c(da^1) c(da^4) |\xi|^{-2m} + 2 \sum_{kp} \partial_{x_k}^2(a^2) \partial_{x_p}(a^3) c(da^1) \partial_{x_p}[c(da^4)] |\xi|^{-2m} \\
&- \frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^2) \partial_{x_r}(a^3) R_{j\alpha r\alpha}(x_0) c(da^1) c(da^4) |\xi|^{-2m} + 2 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_p}^2(a^3) c(da^1) \partial_{x_j}[c(da^4)] |\xi|^{-2m} \\
&+ \frac{1}{2} \sum_{jrst} \partial_{x_j}(a^2) \partial_{x_r}(a^3) R_{jrst}(x_0) c(e_s) c(e_t) c(da^1) c(da^4) |\xi|^{-2m} + 2 \sum_{jr} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_r}^2(a^3) c(da^1) c(da^4) |\xi|^{-2m} \\
&+ 4 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_p}(a^3) c(da^1) \partial_{x_j} \partial_{x_p}[c(da^4)] |\xi|^{-2m} + 4 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_p}(a^3) c(da^1) \partial_{x_p}[c(da^4)] |\xi|^{-2m}.
\end{aligned} \tag{3.16}$$

(ii - 1 - a)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(\sum_{kr} \partial_{x_k}^2(a^2) \partial_{x_r}^2(a^3) c(da^1) c(da^4) |\xi|^{-2m} \right) \sigma(\xi) &= - \sum_{kr} \partial_{x_k}^2(a^2) \partial_{x_r}^2(a^3) g(da^1, da^4) \text{tr}[id] \\
&= -\Delta(a^2) \Delta(a^3) g(da^1, da^4) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(ii - 1 - b) (See Appendix)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(2 \sum_{kp} \partial_{x_k}^2(a^2) \partial_{x_p}(a^3) c(da^1) \partial_{x_p}[c(da^4)] |\xi|^{-2m} \right) \sigma(\xi) &= -2 \sum_{kp} \partial_{x_k}^2(a^2) \partial_{x_p}(a^3) \sum_{\mu} e_{\mu}(a^1) \partial_{x_p}[e_{\mu}(a^4)] \text{tr}[id] \text{area}(S_n) \\
&= 2\Delta(a^2) g(\nabla(a^1), \nabla_{\nabla(a^3)} \nabla(a^4)) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(ii - 1 - c)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(-\frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^2) \partial_{x_r}(a^3) R_{j\alpha r\alpha}(x_0) c(da^1) c(da^4) |\xi|^{-2m} \right) \sigma(\xi) &= \frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^2) \partial_{x_r}(a^3) R_{j\alpha r\alpha}(x_0) g(da^1, da^4) \text{tr}[id] \text{area}(S_n) \\
&= \frac{4}{3} \sum_{\alpha} R(\nabla(a^2), e_{\alpha}, \nabla(a^3), e_{\alpha}) g(da^1, da^4) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(ii - 1 - d)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(2 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_p}^2(a^3) c(da^1) \partial_{x_j}[c(da^4)] |\xi|^{-2m} \right) \sigma(\xi) &= -2 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_p}^2(a^3) \sum_{\mu} e_{\mu}(a^1) \partial_{x_j}[e_{\mu}(a^4)] \text{tr}[id] \text{area}(S_n) \\
&= 2\Delta(a^3) g(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla(a^4)) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(ii - 1 - e)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(\frac{1}{2} \sum_{jr} \partial_{x_j}(a^2) \partial_{x_r}(a^3) \sum_{st} R_{jrst}(x_0) c(e_s) c(e_t) c(da^1) c(da^4) |\xi|^{-2m} \right) \sigma(\xi) &= - \sum_{jrst} \partial_{x_j}(a^2) \partial_{x_r}(a^3) R_{jrst}(x_0) e_s(a^1) e_t(a^4) \text{tr}[id] \text{area}(S_n)
\end{aligned}$$

$$= -R(\nabla(a^2), \nabla(a^3), \nabla(a^4), \nabla(a^1))\text{tr}[id]\text{area}(S_n).$$

(ii - 1 - f)

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(2 \sum_{jr} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_r}^2(a^3) c(da^1) c(da^4) |\xi|^{-2m} \right) \sigma(\xi) \\ &= -2 \sum_{jr} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_r}^2(a^3) g(da^1, da^4) \text{tr}[id]\text{area}(S_n) \\ &= \left(2\nabla(a^2)[\Delta(a^3)] - \frac{4}{3} \sum_{\alpha} R(\nabla(a^2), e_{\alpha}, \nabla(a^3), e_{\alpha}) \right) g(da^1, da^4) \text{tr}[id]\text{area}(S_n). \end{aligned}$$

(ii - 1 - g)(See Appendix)

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(4 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_p}(a^3) c(da^1) \partial_{x_j} \partial_{x_p} [c(da^4)] |\xi|^{-2m} \right) \sigma(\xi) \\ &= -4 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_p}(a^3) \sum_{\mu} e_{\mu}(a^1) \partial_{x_j} \partial_{x_p} [e_{\mu}(a^4)] \text{tr}[id]\text{area}(S_n) \\ &= -4 \left(g(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla_{\nabla(a^3)} \nabla(a^4)) - g(\nabla_{\nabla(a_1)} \nabla(a_4), \nabla_{\nabla(a_2)} \nabla(a_3)) \right. \\ &\quad \left. - \frac{1}{2} R(\nabla(a^1), \nabla(a^3), \nabla(a^2), \nabla(a^4)) \right) \text{tr}[id]\text{area}(S_n). \end{aligned}$$

(ii - 1 - h)(See Appendix)

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(4 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_p}(a^3) c(da^1) \partial_{x_p} [c(da^4)] |\xi|^{-2m} \right) \sigma(\xi) \\ &= -4 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_p}(a^3) \sum_{\mu} e_{\mu}(a^1) \partial_{x_p} [e_{\mu}(a^4)] \text{tr}[id]\text{area}(S_n) \\ &= 4g(\nabla(a^1), \nabla_{\nabla_{\nabla(a^2)} \nabla(a^3)} \nabla(a^4)) \text{tr}[id]\text{area}(S_n). \end{aligned}$$

Therefore, we get

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(\sigma_0(A_1 B_1) \sigma_{-2m}(D^{-2m})(x_0) \right) \sigma(\xi) \\ &= \left(-\Delta(a^2) \Delta(a^3) g(da^1, da^4) + 2\Delta(a^2) g(\nabla(a^1), \nabla_{\nabla(a^3)} \nabla(a^4)) - R(\nabla(a^2), \nabla(a^3), \nabla(a^4), \nabla(a^1)) \right. \\ &\quad + 2\nabla(a^2)[\Delta(a^3)] g(da^1, da^4) + 2\Delta(a^3) g(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla(a^4)) - 4g(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla_{\nabla(a^3)} \nabla(a^4)) \\ &\quad \left. + 4g(\nabla_{\nabla(a_1)} \nabla(a_4), \nabla_{\nabla(a_2)} \nabla(a_3)) + 2R(\nabla(a^1), \nabla(a^3), \nabla(a^2), \nabla(a^4)) + 4g(\nabla(a^1), \nabla_{\nabla_{\nabla(a^2)} \nabla(a^3)} \nabla(a^4)) \right) \text{tr}[id]\text{area}(S_n). \end{aligned}$$

(ii - 2) For $\sigma_1(A_1 B_1) \sigma_{-2m-1}(D^{-2m})$:

By (3.12), we obtain

$$\int_{|\xi|=1} \text{tr} \left(\sigma_1(A_1 B_1) \sigma_{-2m-1}(D^{-2m})(x_0) \right) \sigma(\xi) = 0.$$

(3) For $\sigma_2(A_1 B_1) \sigma_{-2m-2}(D^{-2m})$:

By (3.10), in normal coordinates, we get the following result.

$$\sigma_2(A_1 B_1) \sigma_{-2m-2}(D^{-2m})(x_0)$$

$$\begin{aligned}
&= m \sum_{jl} \partial_{x_j}(a^2) \partial_{x_l}(a^3) \xi_j \xi_l c(da^1)(da^4) |\xi|^{-2m-2} s \\
&\quad - \frac{4m(m+1)}{3} \sum_{jl} \partial_{x_j}(a^2) \partial_{x_l}(a^3) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) \xi_j \xi_l \xi_\mu \xi_\nu c(da^1) c(da^4) |\xi|^{-2m-4}.
\end{aligned} \tag{3.17}$$

(ii - 3 - a)

$$\begin{aligned}
&\int_{|\xi|=1} m \text{tr} \left(\sum_{jl} \partial_{x_j}(a^2) \partial_{x_l}(a^3) \xi_j \xi_l c(da^1) c(da^4) |\xi|^{-2m-2} s \right) \sigma(\xi) \\
&= -\frac{1}{2} \sum_j \partial_{x_j}(a^2) \partial_{x_j}(a^3) \text{area}(S_n) g(da^1, da^4) \text{str}[id] \\
&= -\frac{1}{2} g(\nabla(a^2), \nabla(a^3)) \text{area}(S_n) g(da^1, da^4) \text{str}[id].
\end{aligned}$$

(ii - 3 - b)

$$\begin{aligned}
&\int_{|\xi|=1} \frac{-4m(m+1)}{3} \text{tr} \left(\sum_{jl} \partial_{x_j}(a^2) \partial_{x_l}(a^3) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) \xi_j \xi_l \xi_\mu \xi_\nu c(da^1) c(da^4) |\xi|^{-2m-4} \right) \sigma(\xi) \\
&= \frac{1}{3} \left(\sum_{j\mu\alpha} \partial_{x_j}(a^2) \partial_{x_j}(a^3) R_{\mu\alpha\mu\alpha}(x_0) + 2 \sum_{jl\alpha} \partial_{x_j}(a^2) \partial_{x_l}(a^3) R_{j\alpha l\alpha}(x_0) \right) \text{area}(S_n) g(da^1, da^4) \text{tr}[id] \\
&= \frac{1}{3} \left(\sum_{\mu\alpha} g(\nabla(a^2), \nabla(a^3)) R_{\mu\alpha\mu\alpha}(x_0) + 2 \sum_{\alpha} R(\nabla(a^2), e_\alpha, \nabla(a^3), e_\alpha) \right) \text{area}(S_n) g(da^1, da^4) \text{tr}[id] \\
&= \frac{1}{3} \left(g(\nabla(a^2), \nabla(a^3)) s + 2 \sum_{\alpha} R(\nabla(a^2), e_\alpha, \nabla(a^3), e_\alpha) \right) \text{area}(S_n) g(da^1, da^4) \text{tr}[id].
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(\sigma_2(A_1 B_1) \sigma_{-2m-2}(D^{-2m})(x_0) \right) \sigma(\xi) \\
&= \left(-\frac{1}{6} g(\nabla(a^2), \nabla(a^3)) s + \frac{2}{3} \sum_{\alpha} R(\nabla(a^2), e_\alpha, \nabla(a^3), e_\alpha) \right) \text{area}(S_n) g(da^1, da^4) \text{tr}[id].
\end{aligned}$$

(ii - 4) For $-i \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(A_1 B_1)] \partial_{x_j} [\sigma_{-2m-1}(D^{-2m})]$:

$$\begin{aligned}
&-i \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(A_1 B_1)] \partial_{x_j} [\sigma_{-2m-1}(D^{-2m})](x_0) \\
&= \frac{8m}{3} \sum_{jplk\alpha} \partial_{x_p}(a^2) \partial_{x_l}(a^3) [\partial_{\xi_j}(\xi_p) \xi_l \xi_k + \partial_{\xi_j}(\xi_l) \xi_p \xi_k] R_{j\alpha k\alpha}(x_0) c(da^1) c(da^4) |\xi|^{-2m-2} \\
&\quad - m \sum_{jplkts} \partial_{x_p}(a^2) \partial_{x_l}(a^3) [\partial_{\xi_j}(\xi_p) \xi_l \xi_k + \partial_{\xi_j}(\xi_l) \xi_p \xi_k] R_{j\alpha k\alpha}(x_0) c(da^1) c(da^4) c(e_s) c(e_t) |\xi|^{-2m-2} \\
&\quad + \frac{16}{3} m \sum_{jplr\alpha\beta} \partial_{x_p}(a^2) \partial_{x_l}(a^3) R_{\alpha j\beta r}(x_0) [\partial_{\xi_j}(\xi_p) \xi_l \xi_r \xi_\alpha \xi_\beta + \partial_{\xi_j}(\xi_l) \xi_p \xi_r \xi_\alpha \xi_\beta] c(da^1) c(da^4) |\xi|^{-2m-4} \\
&\quad + \frac{16(m-k-1)}{3} \sum_{k=0}^{m-2} \sum_{\mu=1}^{2m} \sum_{jplr\alpha\beta} \partial_{x_p}(a^2) \partial_{x_l}(a^3) R_{\alpha j\beta\mu}(x_0) [\partial_{\xi_j}(\xi_p) \xi_l \xi_\mu \xi_\alpha \xi_\beta + \partial_{\xi_j}(\xi_l) \xi_p \xi_\mu \xi_\alpha \xi_\beta] c(da^1) c(da^4) |\xi|^{-2m-4}.
\end{aligned}$$

(ii - 4 - a)

$$\begin{aligned}
& \int_{|\xi|=1} \text{tr} \left(\frac{8m}{3} \sum_{jplka} \partial_{x_p}(a^2) \partial_{x_l}(a^3) [\partial_{\xi_j}(\xi_p) \xi_l \xi_k + \partial_{\xi_j}(\xi_l) \xi_p \xi_k] R_{j\alpha k\alpha}(x_0) c(da^1) c(da^4) |\xi|^{-2m-2} \right) \sigma(\xi) \\
&= -\frac{4}{3(m+1)} \sum_{pk\alpha} \partial_{x_p}(a^2) \partial_{x_k}(a^3) R_{p\alpha k\alpha}(x_0) \text{area}(S_n) g(da^1, da^4) \text{tr}[id] \\
&= -\frac{4}{3(m+1)} \sum_{\alpha} R(\nabla(a^2), e_{\alpha}, \nabla(a^3), e_{\alpha}) \text{area}(S_n) g(da^1, da^4) \text{tr}[id].
\end{aligned}$$

(ii - 4 - b)

$$\begin{aligned}
& \int_{|\xi|=1} \text{tr} \left(-m \sum_{jplkts} \partial_{x_p}(a^2) \partial_{x_l}(a^3) [\partial_{\xi_j}(\xi_p) \xi_l \xi_k + \partial_{\xi_j}(\xi_l) \xi_p \xi_k] R_{jktts}(x_0) c(da^1) c(da^4) c(e_s) c(e_t) |\xi|^{-2m-2} \right) \sigma(\xi) \\
&= \frac{1}{2(2m+2)} \text{area}(S^n) \sum_{jlts} \partial_{x_j}(a^2) \partial_{x_l}(a^3) R_{jlts}(x_0) e_s(a^1) e_t(a^4) \text{tr}[id] \\
&+ \frac{1}{2(2m+2)} \text{area}(S^n) \sum_{jpts} \partial_{x_p}(a^2) \partial_{x_j}(a^3) R_{jpts}(x_0) e_s(a^1) e_t(a^4) \text{tr}[id] \\
&= \frac{1}{2(2m+2)} \text{area}(S^n) \left(R(\nabla(a^2), \nabla(a^3), \nabla(a^4), \nabla(a^1)) + R(\nabla(a^3), \nabla(a^2), \nabla(a^4), \nabla(a^1)) \right) \text{tr}[id] \\
&= 0.
\end{aligned}$$

(ii - 4 - c)

$$\begin{aligned}
& \int_{|\xi|=1} \text{tr} \left(\sum_{jplr\alpha\beta} \partial_{x_p}(a^2) \partial_{x_l}(a^3) R_{\alpha j\beta r}(x_0) [\partial_{\xi_j}(\xi_p) \xi_l \xi_r \xi_{\alpha} \xi_{\beta} + \partial_{\xi_j}(\xi_l) \xi_p \xi_r \xi_{\alpha} \xi_{\beta}] c(da^1) c(da^4) |\xi|^{-2m-4} \right) \sigma(\xi) \\
&= - \left(\sum_{jlr\alpha\beta} \partial_{x_j}(a^2) \partial_{x_l}(a^3) R_{\alpha j\beta r}(x_0) I_{S_n}^{lr\alpha\beta} + \sum_{jr\alpha\beta} \partial_{x_p}(a^2) \partial_{x_j}(a^3) R_{\alpha j\beta r}(x_0) I_{S_n}^{pr\alpha\beta} \right) g(da^1, da^4) \text{tr}[id] \\
&= 0.
\end{aligned}$$

(ii - 4 - d)

Similarly, we get

$$\int_{|\xi|=1} \text{tr} \left(\sum_{jplr\alpha\beta} \partial_{x_p}(a^2) \partial_{x_l}(a^3) R_{\alpha j\beta \mu}(x_0) [\partial_{\xi_j}(\xi_p) \xi_l \xi_{\mu} \xi_{\alpha} \xi_{\beta} + \partial_{\xi_j}(\xi_l) \xi_p \xi_{\mu} \xi_{\alpha} \xi_{\beta}] c(da^1) c(da^4) |\xi|^{-2m-4} \right) \sigma(\xi) = 0.$$

Therefore, we get

$$\begin{aligned}
& \int_{|\xi|=1} \text{tr} \left(-i \sum_{j=1}^n \partial_{\xi_j}[\sigma_2(A_1 B_1)] \partial_{x_j}[\sigma_{-2m-1}(D^{-2m})](x_0) \right) \sigma(\xi) \\
&= -\frac{4}{3(m+1)} \sum_{\alpha} R(\nabla(a^2), e_{\alpha}, \nabla(a^3), e_{\alpha}) \text{area}(S_n) g(da^1, da^4) \text{tr}[id].
\end{aligned}$$

(ii - 5) For $-i \sum_{j=1}^n \partial_{\xi_j}[\sigma_1(A_1 B_1)] \partial_{x_j}[\sigma_{-2m}(D^{-2m})]$:
By (3.14), we get

$$\int_{|\xi|=1} \text{tr} \left(-i \sum_{j=1}^n \partial_{\xi_j}[\sigma_1(A_1 B_1)] \partial_{x_j}[\sigma_{-2m}(D^{-2m})] \right) (x_0) \sigma(\xi) = 0.$$

(ii - 6) For $-\frac{1}{2} \sum_{jl} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(A_1 B_1)] \partial_{x_j} \partial_{x_l} [\sigma_{-2m}(D^{-2m})]$:

$$\begin{aligned} & \int_{|\xi|=1} -\frac{4}{3} mtr \left(\sum_{jlr s \alpha \beta} \partial_{x_r}(a^2) \partial_{x_s}(a^3) R_{\alpha j \beta l}(x_0) [\partial_{\xi_l}(\xi_r) \partial_{\xi_j}(\xi_s) + \partial_{\xi_l}(\xi_s) \partial_{\xi_j}(\xi_r)] \xi_\alpha \xi_\beta c(da^1) c(da^4) |\xi|^{-2m-2} \right) \sigma(\xi) \\ &= \frac{2}{3(m+1)} \sum_{jl\alpha} \partial_{x_j}(a^2) \partial_{x_l}(a^3) R_{\alpha j \alpha l}(x_0) area(S_n) g(da^1, da^4) \text{tr}[id] \\ &= \frac{2}{3(m+1)} \sum_{\alpha} R(\nabla(a^2), e_{\alpha}, \nabla(a^3), e_{\alpha}) area(S_n) g(da^1, da^4) \text{tr}[id]. \end{aligned}$$

Therefore, we get

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(-\frac{1}{2} \sum_{jl} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(A_1 B_1)] \partial_{x_j} \partial_{x_l} [\sigma_{-2m}(D^{-2m})](x_0) \right) \sigma(\xi) \\ &= \frac{2}{3(m+1)} \sum_{\alpha} R(\nabla(a^2), e_{\alpha}, \nabla(a^3), e_{\alpha}) area(S_n) g(da^1, da^4) \text{tr}[id]. \end{aligned}$$

Finally, we obtain

Proposition 3.7. Let $P_1 = [D, a^1][D^2, a^2][D^2, a^3][D, a^4]$, we have the following noncommutative integral of P_1 based on the Wodzicki residue

$$\begin{aligned} & \oint a^0 [D, a^1][D^2, a^2][D^2, a^3][D, a^4] D^{-2m} \\ &= 2^m \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \int_M a_0 \left(-\Delta(a^2) \Delta(a^3) g(da^1, da^4) + 2\Delta(a^2) g(\nabla(a^1), \nabla_{\nabla(a^3)} \nabla(a^4)) - R(\nabla(a^2), \nabla(a^3), \nabla(a^4), \nabla(a^1)) \right. \\ &+ 2\nabla(a^2)[\Delta(a^3)]g(da^1, da^4) + 2\Delta(a^3)g(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla(a^4)) - 4g(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla_{\nabla(a^3)} \nabla(a^4)) \\ &+ 4g(\nabla_{\nabla(a_1)} \nabla(a_4), \nabla_{\nabla(a_2)} \nabla(a_3)) + 2R(\nabla(a^1), \nabla(a^3), \nabla(a^2), \nabla(a^4)) + 4g(\nabla(a^1), \nabla_{\nabla_{\nabla(a^2)}} \nabla_{\nabla(a^3)} \nabla(a^4)) \\ &\left. - \frac{1}{6}g(\nabla(a^2), \nabla(a^3))g(da^1, da^4)s + \frac{2m}{3(m+1)} \sum_{\alpha} R(\nabla(a^2), e_{\alpha}, \nabla(a^3), e_{\alpha}) g(da^1, da^4) \right) d\text{Vol}_M. \end{aligned}$$

Part iii) $P_2 = [D, a^1][D, a^2][D^2, a^3][D^2, a^4]$, $[D, a^1][D, a^2][D^2, a^3] := A_2$, $[D^2, a^4] := B_2$. Similarly, by Lemma 3.1 in [12] and Lemma 2.1, we get the following lemma.

Lemma 3.8. The symbols of A and B are given

$$\begin{aligned} \sigma_0(A_2) &= c(da^1)c(da^2) \left[\sum_{j=1}^n \partial_{x_j}(a^3)(\Gamma^j - 2\sigma^j)(x) - \sum_{jl=1}^n \partial_{x_j} \partial_{x_l}(a^3) g^{jl} \right]; \\ \sigma_1(A_2) &= -2ic(da^1)c(da^2) \sum_{jl=1}^n \partial_{x_j}(a^3) g^{jl} \xi_l; \\ \sigma_0(B_2) &= \sum_{j=1}^n \partial_{x_j}(a^4)(\Gamma^j - 2\sigma^j)(x) - \sum_{jl=1}^n \partial_{x_j} \partial_{x_l}(a^4) g^{jl}; \\ \sigma_1(B_2) &= -2i \sum_{jl=1}^n \partial_{x_j}(a^4) g^{jl} \xi_l. \end{aligned} \tag{3.18}$$

Next, we calculate each term of $\int_{S^* M} \text{tr}[\sigma_{-2m}(A_2 B_2 D^{-2m})](x, \xi)$ separately.
(iii - 1) For $\sigma_0(A_2 B_2) \sigma_{-2m}(D^{-2m})$

$$\sigma_0(A_2 B_2) \sigma_{-2m}(D^{-2m})(x_0)$$

$$\begin{aligned}
&= \sum_{kr} \partial_{x_k}^2(a^3) \partial_{x_r}^2(a^4) c(da^1) c(da^2) |\xi|^{-2m} - \frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^3) \partial_{x_r}(a^4) R_{j\alpha r\alpha}(x_0) c(da^1) c(da^2) |\xi|^{-2m} \\
&+ \frac{1}{2} \sum_{jrst} \partial_{x_j}(a^3) \partial_{x_r}(a^4) R_{jrts}(x_0) c(e_s) c(e_t) c(da^1) c(da^2) |\xi|^{-2m} \\
&+ 2 \sum_{jr} \partial_{x_j}(a^3) \partial_{x_j}^2(a^4) c(da^1) c(da^2) |\xi|^{-2m} \Big) |\xi|^{-2m}. \tag{3.19}
\end{aligned}$$

(iii - 1 - a)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(\sum_{kr} \partial_{x_k}^2(a^3) \partial_{x_r}^2(a^4) c(da^1) c(da^2) |\xi|^{-2m} \right) \sigma(\xi) &= - \sum_{kr} \partial_{x_k}^2(a^3) \partial_{x_r}^2(a^4) g(da^1, da^2) \text{tr}[id] \text{area}(S_n) \\
&= -\Delta(a^3) \Delta(a^4) g(da^1, da^2) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(iii - 1 - b)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(-\frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^3) \partial_{x_r}(a^4) R_{j\alpha r\alpha}(x_0) c(da^1) c(da^2) |\xi|^{-2m} \right) \sigma(\xi) \\
&= \frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^3) \partial_{x_r}(a^4) R_{j\alpha r\alpha}(x_0) g(da^1, da^2) \text{tr}[id] \text{area}(S_n) \\
&= \frac{4}{3} \sum_{\alpha} R(\nabla(a^3), e_{\alpha}, \nabla(a^4), e_{\alpha}) g(da^1, da^2) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(iii - 1 - c)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(\frac{1}{2} \sum_{jr} \partial_{x_j}(a^3) \partial_{x_r}(a^4) \sum_{st} R_{jrts}(x_0) c(e_s) c(e_t) c(da^1) c(da^2) |\xi|^{-2m} \right) \sigma(\xi) \\
&= - \sum_{jrst} \partial_{x_j}(a^3) \partial_{x_r}(a^4) R_{jrts}(x_0) e_s(a^1) e_t(a^2) \text{tr}[id] \text{area}(S_n) \\
&= -R(\nabla(a^3), \nabla(a^4), \nabla(a^2), \nabla(a^1)) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(iii - 1 - d)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(2 \sum_{jr} \partial_{x_j}(a^3) \partial_{x_j} \partial_{x_r}^2(a^4) c(da^1) c(da^2) |\xi|^{-2m} \right) \sigma(\xi) \\
&= -2 \sum_{jr} \partial_{x_j}(a^3) \partial_{x_j} \partial_{x_r}^2(a^4) g(da^1, da^2) \text{tr}[id] \text{area}(S_n) \\
&= \left(2\nabla(a^3)[\Delta(a^4)] - \frac{4}{3} \sum_{\alpha} R(\nabla(a^3), e_{\alpha}, \nabla(a^4), e_{\alpha}) \right) g(da^1, da^2) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(\sigma_0(A_2 B_2) \sigma_{-2m}(D^{-2m})(x_0) \right) \sigma(\xi) \\
&= \left(-\Delta(a^3) \Delta(a^4) - R(\nabla(a^3), \nabla(a^4), \nabla(a^2), \nabla(a^1)) + 2\nabla(a^3)[\Delta(a^4)] \right) g(da^1, da^2) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(iii - 2) For $\sigma_1(A_2 B_2) \sigma_{-2m-1}(D^{-2m})$:

By (3.12), we get

$$\int_{|\xi|=1} \text{tr} \left(\sigma_1(A_2 B_2) \sigma_{-2m-1}(D^{-2m})(x_0) \right) \sigma(\xi) = 0.$$

(iii - 3) For $\sigma_2(A_2B_2)\sigma_{-2m-2}(D^{-2m})$:

$$\begin{aligned} \sigma_2(A_2B_2)\sigma_{-2m-2}(D^{-2m})(x_0) &= m \sum_{jl} \partial_{x_j}(a^3) \partial_{x_l}(a^4) \xi_j \xi_l c(da^1) c(da^2) |\xi|^{-2m-2} s \\ &\quad - \frac{4m(m+1)}{3} \sum_{jl} \partial_{x_j}(a^3) \partial_{x_l}(a^4) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) \xi_j \xi_l \xi_\mu \xi_\nu c(da^1) c(da^2) |\xi|^{-2m-4}. \end{aligned} \quad (3.20)$$

(iii - 3 - a)

$$\begin{aligned} &\int_{|\xi|=1} m \text{tr} \left(\sum_{jl} \partial_{x_j}(a^3) \partial_{x_l}(a^4) \xi_j \xi_l c(da^1) c(da^2) |\xi|^{-2m-2} s \right) \sigma(\xi) \\ &= -\frac{1}{2} \sum_j \partial_{x_j}(a^3) \partial_{x_j}(a^4) \text{area}(S_n) g(da^1, da^2) \text{str}[id] \\ &= -\frac{1}{2} g(\nabla(a^3), \nabla(a^4)) \text{area}(S_n) g(da^1, da^2) \text{str}[id]. \end{aligned}$$

(iii - 3 - b)

$$\begin{aligned} &\int_{|\xi|=1} -\frac{4m(m+1)}{3} \text{tr} \left(\sum_{jl} \partial_{x_j}(a^3) \partial_{x_l}(a^4) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) \xi_j \xi_l \xi_\mu \xi_\nu c(da^1) c(da^2) |\xi|^{-2m-4} \right) \sigma(\xi) \\ &= \frac{1}{3} \left(\sum_{j\mu\alpha} \partial_{x_j}(a^3) \partial_{x_j}(a^4) R_{\mu\alpha\mu\alpha}(x_0) + 2 \sum_{jl\alpha} \partial_{x_j}(a^3) \partial_{x_l}(a^4) R_{j\alpha l\alpha}(x_0) \right) \text{area}(S_n) g(da^1, da^2) \text{tr}[id] \\ &= \frac{1}{3} \left(\sum_{\mu\alpha} g(\nabla(a^3), \nabla(a^4)) R_{\mu\alpha\mu\alpha}(x_0) + 2 \sum_{\alpha} R(\nabla(a^3), e_\alpha, \nabla(a^4), e_\alpha) \right) \text{area}(S_n) g(da^1, da^2) \text{tr}[id] \\ &= \frac{1}{3} \left(g(\nabla(a^3), \nabla(a^4)) s + 2 \sum_{\alpha} R(\nabla(a^3), e_\alpha, \nabla(a^4), e_\alpha) \right) \text{area}(S_n) g(da^1, da^2) \text{tr}[id]. \end{aligned}$$

Therefore, we get

$$\begin{aligned} &\int_{|\xi|=1} \text{tr} \left(\sigma_2(A_2B_2)\sigma_{-2m-2}(D^{-2m})(x_0) \right) \sigma(\xi) \\ &= \left(-\frac{1}{6} g(\nabla(a^3), \nabla(a^4)) s + \frac{2}{3} \sum_{\alpha} R(\nabla(a^3), e_\alpha, \nabla(a^4), e_\alpha) \right) \text{area}(S_n) g(da^1, da^2) \text{tr}[id]. \end{aligned}$$

(iii - 4) For $-i \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(A_2B_2)] \partial_{x_j} [\sigma_{-2m-1}(D^{-2m})]$:

$$\begin{aligned} &-i \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(A_2B_2)] \partial_{x_j} [\sigma_{-2m-1}(D^{-2m})](x_0) \\ &= \frac{8m}{3} \sum_{jplk\alpha} \partial_{x_p}(a^3) \partial_{x_l}(a^4) [\partial_{\xi_j}(\xi_p) \xi_l \xi_k + \partial_{\xi_j}(\xi_l) \xi_p \xi_k] R_{j\alpha k\alpha}(x_0) c(da^1) c(da^2) |\xi|^{-2m-2} \\ &\quad - m \sum_{jplkts} \partial_{x_p}(a^3) \partial_{x_l}(a^4) [\partial_{\xi_j}(\xi_p) \xi_l \xi_k + \partial_{\xi_j}(\xi_l) \xi_p \xi_k] R_{j\alpha k\alpha}(x_0) c(da^1) c(da^2) c(e_s) c(e_t) |\xi|^{-2m-2} \\ &\quad + \frac{16}{3} m \sum_{jplr\alpha\beta} \partial_{x_p}(a^3) \partial_{x_l}(a^4) R_{\alpha j\beta r}(x_0) [\partial_{\xi_j}(\xi_p) \xi_l \xi_r \xi_\alpha \xi_\beta + \partial_{\xi_j}(\xi_l) \xi_p \xi_r \xi_\alpha \xi_\beta] c(da^1) c(da^2) |\xi|^{-2m-4} \end{aligned}$$

$$+ \frac{16(m-k-1)}{3} \sum_{k=0}^{m-2} \sum_{\mu=1}^{2m} \sum_{jplr\alpha\beta} \partial_{x_p}(a^3) \partial_{x_l}(a^4) R_{\alpha j \beta \mu}(x_0) [\partial_{\xi_j}(\xi_p) \xi_l \xi_\mu \xi_\alpha \xi_\beta + \partial_{\xi_j}(\xi_l) \xi_p \xi_\mu \xi_\alpha \xi_\beta] c(da^1) c(da^2) |\xi|^{-2m-4}.$$

(iii - 4 - a)

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(\frac{8m}{3} \sum_{jplk\alpha} \partial_{x_p}(a^3) \partial_{x_l}(a^4) [\partial_{\xi_j}(\xi_p) \xi_l \xi_k + \partial_{\xi_j}(\xi_l) \xi_p \xi_k] R_{j\alpha k\alpha}(x_0) c(da^1) c(da^2) |\xi|^{-2m-2} \right) \sigma(\xi) \\ &= -\frac{4}{3(m+1)} \sum_{pk\alpha} \partial_{x_p}(a^3) \partial_{x_k}(a^4) R_{p\alpha k\alpha}(x_0) \text{area}(S_n) g(da^1, da^2) \text{tr}[id] \\ &= -\frac{4}{3(m+1)} \sum_{\alpha} R(\nabla(a^3), e_{\alpha}, \nabla(a^4), e_{\alpha}) \text{area}(S_n) g(da^1, da^2) \text{tr}[id]. \end{aligned}$$

(iii - 4 - b)

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(-m \sum_{jplkts} \partial_{x_p}(a^3) \partial_{x_l}(a^4) [\partial_{\xi_j}(\xi_p) \xi_l \xi_k + \partial_{\xi_j}(\xi_l) \xi_p \xi_k] R_{jklts}(x_0) c(da^1) c(da^2) c(e_s) c(e_t) |\xi|^{-2m-2} \right) \sigma(\xi) \\ &= \frac{1}{2(2m+2)} \text{area}(S^n) \sum_{jlt}s \partial_{x_j}(a^3) \partial_{x_l}(a^4) R_{jlt}s(x_0) e_s(a^1) e_t(a^2) \text{tr}[id] \\ &+ \frac{1}{2(2m+2)} \text{area}(S^n) \sum_{jpts} \partial_{x_p}(a^3) \partial_{x_j}(a^4) R_{jpts}(x_0) e_s(a^1) e_t(a^2) \text{tr}[id] \\ &= \frac{1}{2(2m+2)} \text{area}(S^n) \left(R(\nabla(a^3), \nabla(a^4), \nabla(a^2), \nabla(a^1)) + R(\nabla(a^4), \nabla(a^3), \nabla(a^2), \nabla(a^1)) \right) \text{tr}[id] \\ &= 0.. \end{aligned}$$

(iii - 4 - c)

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(\sum_{jplr\alpha\beta} \partial_{x_p}(a^3) \partial_{x_l}(a^4) R_{\alpha j \beta r}(x_0) [\partial_{\xi_j}(\xi_p) \xi_l \xi_r \xi_\alpha \xi_\beta + \partial_{\xi_j}(\xi_l) \xi_p \xi_r \xi_\alpha \xi_\beta] c(da^1) c(da^2) |\xi|^{-2m-4} \right) \sigma(\xi) \\ &= - \left(\sum_{jlr\alpha\beta} \partial_{x_j}(a^3) \partial_{x_l}(a^4) R_{\alpha j \beta r}(x_0) I_{S_n}^{l r \alpha \beta} + \sum_{j r \alpha \beta} \partial_{x_p}(a^3) \partial_{x_j}(a^4) R_{\alpha j \beta r}(x_0) I_{S_n}^{p r \alpha \beta} \right) g(da^1, da^2) \text{tr}[id] \\ &= 0. \end{aligned}$$

(iii - 4 - d)

Similarly, we get

$$\int_{|\xi|=1} \text{tr} \left(\sum_{jplr\alpha\beta} \partial_{x_p}(a^3) \partial_{x_l}(a^4) R_{\alpha j \beta \mu}(x_0) [\partial_{\xi_j}(\xi_p) \xi_l \xi_\mu \xi_\alpha \xi_\beta + \partial_{\xi_j}(\xi_l) \xi_p \xi_\mu \xi_\alpha \xi_\beta] c(da^1) c(da^2) |\xi|^{-2m-4} \right) \sigma(\xi) = 0.$$

Therefore, we get

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(-i \sum_{j=1}^n \partial_{\xi_j}[\sigma_2(A_2 B_2)] \partial_{x_j}[\sigma_{-2m-1}(D^{-2m})](x_0) \right) \sigma(\xi) \\ &= -\frac{4}{3(m+1)} \sum_{\alpha} R(\nabla(a^3), e_{\alpha}, \nabla(a^4), e_{\alpha}) \text{area}(S_n) g(da^1, da^2) \text{tr}[id]. \end{aligned}$$

(iii - 5) For $-i \sum_{j=1}^n \partial_{\xi_j}[\sigma_1(A_2 B_2)] \partial_{x_j}[\sigma_{-2m}(D^{-2m})]$:

In normal coordinates, by (3.14), we get

$$\int_{|\xi|=1} \text{tr} \left(-i \sum_{j=1}^n \partial_{\xi_j}[\sigma_1(A_2 B_2)] \partial_{x_j}[\sigma_{-2m}(D^{-2m})] \right) (x_0) \sigma(\xi) = 0.$$

(iii - 6) For $-\frac{1}{2} \sum_{jl} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(A_2 B_2)] \partial_{x_j} \partial_{x_l} [\sigma_{-2m}(D^{-2m})]$:
By (3.10) then in normal coordinates, we get the following result.

$$\begin{aligned} & \int_{|\xi|=1} -\frac{4}{3} m \text{tr} \left(\sum_{jlr s \alpha \beta} \partial_{x_r}(a^3) \partial_{x_s}(a^4) R_{\alpha j \beta l}(x_0) [\partial_{\xi_l}(\xi_r) \partial_{\xi_j}(\xi_s) + \partial_{\xi_l}(\xi_s) \partial_{\xi_j}(\xi_r)] \xi_\alpha \xi_\beta c(da^1) c(da^2) |\xi|^{-2m-2} \right) \sigma(\xi) \\ &= \frac{2}{3(m+1)} \sum_{jl\alpha} \partial_{x_j}(a^3) \partial_{x_l}(a^4) R_{\alpha j \alpha l}(x_0) \text{area}(S_n) g(da^1, da^2) \text{tr}[id] \\ &= \frac{2}{3(m+1)} \sum_{\alpha} R(\nabla(a^3), e_{\alpha}, \nabla(a^4), e_{\alpha}) \text{area}(S_n) g(da^1, da^2) \text{tr}[id]. \end{aligned}$$

Therefore, we get

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(-\frac{1}{2} \sum_{jl} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(A_2 B_2)] \partial_{x_j} \partial_{x_l} [\sigma_{-2m}(D^{-2m})](x_0) \right) \sigma(\xi) \\ &= \frac{2}{3(m+1)} \sum_{\alpha} R(\nabla(a^3), e_{\alpha}, \nabla(a^4), e_{\alpha}) \text{area}(S_n) g(da^1, da^2) \text{tr}[id]. \end{aligned}$$

Finally, we obtain

Proposition 3.9. Let $P_2 = [D, a^1][D, a^2][D^2, a^3][D^2, a^4]$, we have the following noncommutative integral of P_2 based on the Wodzicki residue

$$\begin{aligned} & \oint a^0 [D, a^1][D, a^2][D^2, a^3][D^2, a^4] D^{-2m} \\ &= 2^m \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \int_M a_0 \left[\left(-\Delta(a^3)\Delta(a^4) + 2\nabla(a^3)[\Delta(a^4)] - R(\nabla(a^3), \nabla(a^4), \nabla(a^2), \nabla(a^1)) \right. \right. \\ & \quad \left. \left. - \frac{1}{6}g(\nabla(a^3), \nabla(a^4))s + \frac{2m}{3(m+1)} \sum_{\alpha} R(\nabla(a^3), e_{\alpha}, \nabla(a^4), e_{\alpha})(x_0) \right) g(da^1, da^2) \right] d\text{Vol}_M. \end{aligned}$$

Part iv) $P_3 = [D^2, a^1][D, a^2][D, a^3][D^2, a^4]$, $[D^2, a^1][D, a^2][D, a^3] := A_3$, $[D^2, a^4] := B_3$.
Similarly, by Lemma 3.1 in [12] and Lemma 2.1, we get the following lemma.

Lemma 3.10. The symbols of A and B are given

$$\begin{aligned} \sigma_0(A_3) &= \left[\sum_{j=1}^n \partial_{x_j}(a^1)(\Gamma^j - 2\sigma^j)(x) - \sum_{jl=1}^n \partial_{x_j} \partial_{x_l}(a^1) g^{jl} \right] c(da^2) c(da^3); \\ \sigma_1(A_3) &= -2i \sum_{jl=1}^n \partial_{x_j}(a^1) g^{jl} \xi_l c(da^2) c(da^3); \\ \sigma_0(B_3) &= \sum_{j=1}^n \partial_{x_j}(a^4)(\Gamma^j - 2\sigma^j)(x) - \sum_{jl=1}^n \partial_{x_j} \partial_{x_l}(a^4) g^{jl}; \\ \sigma_1(B_3) &= -2i \sum_{jl=1}^n \partial_{x_j}(a^4) g^{jl} \xi_l. \end{aligned} \tag{3.21}$$

Next, we calculate each term of $\int_{S^* M} \text{tr}[\sigma_{-2m}(A_3 B_3 D^{-2m})](x, \xi)$ separately.
(iv - 1) For $\sigma_0(A_3 B_3) \sigma_{-2m}(D^{-2m})$

$$\begin{aligned} & \sigma_0(A_3 B_3) \sigma_{-2m}(D^{-2m})(x_0) \\ &= \sum_{kr} \partial_{x_k}^2(a^1) \partial_{x_r}^2(a^4) c(da^2) c(da^3) |\xi|^{-2m} + 2 \sum_{kp} \partial_{x_k}^2(a^1) \partial_{x_p}(a^4) \partial_{x_p}[c(da^2) c(da^3)] |\xi|^{-2m} \end{aligned}$$

$$\begin{aligned}
& -\frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^1) \partial_{x_r}(a^4) R_{j\alpha r\alpha}(x_0) c(da^2) c(da^3) |\xi|^{-2m} + \frac{1}{2} \sum_{jrst} \partial_{x_j}(a^1) \partial_{x_r}(a^4) R_{jrts}(x_0) c(e_s) c(e_t) c(da^2) c(da^3) |\xi|^{-2m} \\
& + 2 \sum_{jr} \partial_{x_j}(a^1) \partial_{x_j} \partial_{x_r}^2(a^4) c(da^2) c(da^3) |\xi|^{-2m}.
\end{aligned} \tag{3.22}$$

(iv - 1 - a)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(\sum_{kr} \partial_{x_k}^2(a^1) \partial_{x_r}^2(a^4) c(da^2) c(da^3) |\xi|^{-2m} \right) \sigma(\xi) &= - \sum_{kr} \partial_{x_k}^2(a^1) \partial_{x_r}^2(a^4) g(da^2, da^3) \text{tr}[id] \text{area}(S_n) \\
&= -\Delta(a^1) \Delta(a^4) g(da^2, da^3) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(iv - 1 - b)

$$\begin{aligned}
\int_{|\xi|=1} \text{tr} \left(2 \sum_{kp} \partial_{x_k}^2(a^1) \partial_{x_p}(a^4) \partial_{x_p}[c(da^2) c(da^3)] |\xi|^{-2m} \right) \sigma(\xi) &= -2 \sum_{kp} \partial_{x_k}^2(a^1) \partial_{x_p}(a^4) \partial_{x_p}[g(da^2, da^3)] \text{tr}[id] \text{area}(S_n) \\
&= -2\Delta(a^1) g(\nabla(a^4), \nabla g(da^2, da^3)) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(iv - 1 - c)

$$\begin{aligned}
& \int_{|\xi|=1} \text{tr} \left(-\frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^1) \partial_{x_r}(a^4) R_{j\alpha r\alpha}(x_0) c(da^2) c(da^3) |\xi|^{-2m} \right) \sigma(\xi) \\
&= \frac{4}{3} \sum_{jr\alpha} \partial_{x_j}(a^1) \partial_{x_r}(a^4) R_{j\alpha r\alpha}(x_0) g(da^2, da^3) \text{tr}[id] \text{area}(S_n) \\
&= \frac{4}{3} \sum_{\alpha} R(\nabla a^1, e_{\alpha}, \nabla a^4, e_{\alpha}) g(da^2, da^3) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(iv - 1 - d)

$$\begin{aligned}
& \int_{|\xi|=1} \text{tr} \left(\frac{1}{2} \sum_{jr} \partial_{x_j}(a^1) \partial_{x_r}(a^4) \sum_{st} R_{jrts}(x_0) c(e_s) c(e_t) c(da^2) c(da^3) |\xi|^{-2m} \right) \sigma(\xi) \\
&= - \sum_{jrts} \partial_{x_j}(a^1) \partial_{x_r}(a^4) R_{jrts}(x_0) e_s(a^2) e_t(a^3) \text{tr}[id] \text{area}(S_n) \\
&= -R(\nabla(a^1), \nabla(a^4), \nabla(a^3), \nabla(a^2)) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

(iv - 1 - e)

$$\begin{aligned}
& \int_{|\xi|=1} \text{tr} \left(2 \sum_{jr} \partial_{x_j}(a^1) \partial_{x_j} \partial_{x_r}^2(a^4) c(da^2) c(da^3) |\xi|^{-2m} \right) \sigma(\xi) \\
&= -2 \sum_{jr} \partial_{x_j}(a^1) \partial_{x_j} \partial_{x_r}^2(a^4) g(da^2, da^3) \text{tr}[id] \text{area}(S_n) \\
&= \left(2\nabla(a^1)[\Delta(a^4)] - \frac{4}{3} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^4), e_{\alpha}) \right) g(da^2, da^3) \text{tr}[id] \text{area}(S_n).
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
& \int_{|\xi|=1} \text{tr} \left(\sigma_0(A_3 B_3) \sigma_{-2m}(D^{-2m})(x_0) \right) \sigma(\xi) \\
&= \left(-\Delta(a^1) \Delta(a^4) g(da^2, da^3) - 2\Delta(a^1) g(\nabla(a^4), \nabla g(da^2, da^3)) - R(\nabla(a^1), \nabla(a^4), \nabla(a^3), \nabla(a^2)) \right)
\end{aligned}$$

$$+ 2\nabla(a^1)[\Delta(a^4)]g(da^2, da^3)\Big) \text{tr}[id] \text{area}(S_n).$$

(iv - 2) For $\sigma_1(A_3B_3)\sigma_{-2m-1}(D^{-2m})$:

By (3.12), we get

$$\int_{|\xi|=1} \text{tr}\left(\sigma_1(A_3B_3)\sigma_{-2m-1}(D^{-2m})(x_0)\right) \sigma(\xi) = 0.$$

(iv - 3) For $\sigma_2(A_3B_3)\sigma_{-2m-2}(D^{-2m})$:

$$\begin{aligned} & \sigma_2(A_3B_3)\sigma_{-2m-2}(D^{-2m})(x_0) \\ &= m \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^4) \xi_j \xi_l c(da^2) c(da^3) |\xi|^{-2m-2} s \\ &\quad - \frac{4m(m+1)}{3} \sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^4) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) \xi_j \xi_l \xi_\mu \xi_\nu c(da^2) c(da^3) |\xi|^{-2m-4}. \end{aligned} \quad (3.23)$$

(iv - 3 - a)

$$\begin{aligned} & \int_{|\xi|=1} m \text{tr}\left(\sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^4) \xi_j \xi_l c(da^2) c(da^3) |\xi|^{-2m-2} s\right) \sigma(\xi) \\ &= -\frac{1}{2} \sum_j \partial_{x_j}(a^1) \partial_{x_j}(a^4) \text{area}(S_n) g(da^2, da^3) \text{str}[id] \\ &= -\frac{1}{2} g(\nabla(a^1), \nabla(a^4)) \text{area}(S_n) g(da^2, da^3) \text{str}[id]. \end{aligned}$$

(iv - 3 - b)

$$\begin{aligned} & \int_{|\xi|=1} -\frac{4m(m+1)}{3} \text{tr}\left(\sum_{jl} \partial_{x_j}(a^1) \partial_{x_l}(a^4) \sum_{\mu\nu\alpha} R_{\mu\alpha\nu\alpha}(x_0) \xi_j \xi_l \xi_\mu \xi_\nu c(da^2) c(da^3) |\xi|^{-2m-4}\right) \sigma(\xi) \\ &= \frac{1}{3} \left(\sum_{j\mu\alpha} \partial_{x_j}(a^1) \partial_{x_j}(a^4) R_{\mu\alpha\mu\alpha}(x_0) + 2 \sum_{jl\alpha} \partial_{x_j}(a^1) \partial_{x_l}(a^4) R_{j\alpha l\alpha}(x_0) \right) \text{area}(S_n) g(da^2, da^3) \text{tr}[id] \\ &= \frac{1}{3} \left(\sum_{\mu\alpha} g(\nabla(a^1), \nabla(a^4)) R_{\mu\alpha\mu\alpha}(x_0) + 2 \sum_{\alpha} R(\nabla(a^1), e_\alpha, \nabla(a^4), e_\alpha) \right) \text{area}(S_n) g(da^2, da^3) \text{tr}[id] \\ &= \frac{1}{3} \left(g(\nabla(a^1), \nabla(a^4)) s + 2 \sum_{\alpha} R(\nabla(a^1), e_\alpha, \nabla(a^4), e_\alpha) \right) \text{area}(S_n) g(da^2, da^3) \text{tr}[id]. \end{aligned}$$

Therefore, we get

$$\begin{aligned} & \int_{|\xi|=1} \text{tr}\left(\sigma_2(A_3B_3)\sigma_{-2m-2}(D^{-2m})(x_0)\right) \sigma(\xi) \\ &= \left(-\frac{1}{6} g(\nabla(a^1), \nabla(a^4)) s + \frac{2}{3} \sum_{\alpha} R(\nabla(a^1), e_\alpha, \nabla(a^4), e_\alpha) \right) \text{area}(S_n) g(da^2, da^3) \text{tr}[id]. \end{aligned}$$

(iv - 4) For $-i \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(A_3B_3)] \partial_{x_j} [\sigma_{-2m-1}(D^{-2m})]$:

$$-i \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(A_3B_3)] \partial_{x_j} [\sigma_{-2m-1}(D^{-2m})](x_0)$$

$$\begin{aligned}
&= \frac{8m}{3} \sum_{jplk\alpha} \partial_{x_p}(a^1) \partial_{x_l}(a^4) [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] R_{j\alpha k\alpha}(x_0) c(da^2)c(da^3)|\xi|^{-2m-2} \\
&- m \sum_{jplkts} \partial_{x_p}(a^1) \partial_{x_l}(a^4) [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] R_{jkt}s(x_0) c(da^2)c(da^3)c(e_s)c(e_t)|\xi|^{-2m-2} \\
&+ \frac{16}{3}m \sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^4) R_{\alpha j\beta r}(x_0) [\partial_{\xi_j}(\xi_p)\xi_l\xi_r\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_r\xi_\alpha\xi_\beta] c(da^2)c(da^3)|\xi|^{-2m-4} \\
&+ \frac{16(m-k-1)}{3} \sum_{k=0}^{m-2} \sum_{\mu=1}^{2m} \sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^4) R_{\alpha j\beta\mu}(x_0) [\partial_{\xi_j}(\xi_p)\xi_l\xi_\mu\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_\mu\xi_\alpha\xi_\beta] c(da^2)c(da^3)|\xi|^{-2m-4}.
\end{aligned}$$

(iv - 4 - a)

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(\frac{8m}{3} \sum_{jplk\alpha} \partial_{x_p}(a^1) \partial_{x_l}(a^4) [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] R_{j\alpha k\alpha}(x_0) c(da^2)c(da^3)|\xi|^{-2m-2} \right) \sigma(\xi) \\
&= -\frac{4}{3(m+1)} \sum_{pk\alpha} \partial_{x_p}(a^1) \partial_{x_k}(a^4) R_{p\alpha k\alpha}(x_0) \text{area}(S_n) g(da^2, da^3) \text{tr}[id] \\
&= -\frac{4}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^4), e_{\alpha}) \text{area}(S_n) g(da^2, da^3) \text{tr}[id].
\end{aligned}$$

(iv - 4 - b)

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(-m \sum_{jplkts} \partial_{x_p}(a^1) \partial_{x_l}(a^4) [\partial_{\xi_j}(\xi_p)\xi_l\xi_k + \partial_{\xi_j}(\xi_l)\xi_p\xi_k] R_{jkt}s(x_0) c(da^2)c(da^3)c(e_s)c(e_t)|\xi|^{-2m-2} \right) \sigma(\xi) \\
&= 0.
\end{aligned}$$

(iv - 4 - c)

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(\sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^4) R_{\alpha j\beta r}(x_0) [\partial_{\xi_j}(\xi_p)\xi_l\xi_r\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_r\xi_\alpha\xi_\beta] c(da^2)c(da^3)|\xi|^{-2m-4} \right) \sigma(\xi) \\
&= - \left(\sum_{jlr\alpha\beta} \partial_{x_j}(a^1) \partial_{x_l}(a^4) R_{\alpha j\beta r}(x_0) I_{S_n}^{lra\beta} + \sum_{jr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_j}(a^4) R_{\alpha j\beta r}(x_0) I_{S_n}^{pr\alpha\beta} \right) g(da^2, da^3) \text{tr}[id] \\
&= 0.
\end{aligned}$$

(iv - 4 - d)

Similarly, we get

$$\int_{|\xi|=1} \text{tr} \left(\sum_{jplr\alpha\beta} \partial_{x_p}(a^1) \partial_{x_l}(a^4) R_{\alpha j\beta\mu}(x_0) [\partial_{\xi_j}(\xi_p)\xi_l\xi_\mu\xi_\alpha\xi_\beta + \partial_{\xi_j}(\xi_l)\xi_p\xi_\mu\xi_\alpha\xi_\beta] c(da^2)c(da^3)|\xi|^{-2m-4} \right) \sigma(\xi) = 0.$$

Therefore, we get

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(-i \sum_{j=1}^n \partial_{\xi_j}[\sigma_2(A_3B_3)] \partial_{x_j}[\sigma_{-2m-1}(D^{-2m})](x_0) \right) \sigma(\xi) \\
&= -\frac{4}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^4), e_{\alpha}) \text{area}(S_n) g(da^2, da^3) \text{tr}[id].
\end{aligned}$$

(iv - 5) For $-i\Sigma_{j=1}^n \partial_{\xi_j}[\sigma_1(A_3B_3)] \partial_{x_j}[\sigma_{-2m}(D^{-2m})]$:

By (3.14), we get

$$\int_{|\xi|=1} \text{tr} \left(-i\Sigma_{j=1}^n \partial_{\xi_j}[\sigma_1(A_3B_3)] \partial_{x_j}[\sigma_{-2m}(D^{-2m})] \right) (x_0) \sigma(\xi) = 0.$$

(iv - 6) For $-\frac{1}{2} \sum_{jl} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(A_3 B_3)] \partial_{x_j} \partial_{x_l} [\sigma_{-2m}(D^{-2m})]$:

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(-\frac{4}{3}m \sum_{jlr s \alpha \beta} \partial_{x_r}(a^1) \partial_{x_s}(a^4) R_{\alpha j \beta l}(x_0) [\partial_{\xi_l}(\xi_r) \partial_{\xi_j}(\xi_s) + \partial_{\xi_l}(\xi_s) \partial_{\xi_j}(\xi_r)] \xi_\alpha \xi_\beta c(da^2) c(da^3) |\xi|^{-2m-2} \right) \sigma(\xi) \\ &= \frac{2}{3(m+1)} \sum_{jl\alpha} \partial_{x_j}(a^1) \partial_{x_l}(a^4) R_{\alpha j \alpha l}(x_0) \text{area}(S_n) g(da^2, da^3) \text{tr}[id] \\ &= \frac{2}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^4), e_{\alpha}) \text{area}(S_n) g(da^2, da^3) \text{tr}[id]. \end{aligned}$$

Therefore, we get

$$\begin{aligned} & \int_{|\xi|=1} \text{tr} \left(-\frac{1}{2} \sum_{jl} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(A_3 B_3)] \partial_{x_j} \partial_{x_l} [\sigma_{-2m}(D^{-2m})](x_0) \right) \sigma(\xi) \\ &= \frac{2}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^4), e_{\alpha}) \text{area}(S_n) g(da^2, da^3) \text{tr}[id]. \end{aligned}$$

Finally, we obtain

Proposition 3.11. Let $P_3 = [D^2, a^1][D, a^2][D, a^3][D^2, a^4]$, we have the following noncommutative integral of P_3 based on the Wodzicki residue

$$\begin{aligned} & \oint a^0 [D^2, a^1][D, a^2][D, a^3][D^2, a^4] D^{-2m} \\ &= 2^m \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \int_M a_0 \left(-\Delta(a^1) \Delta(a^4) g(da^2, da^3) - 2\Delta(a^1) g(\nabla(a^4), \nabla g(da^2, da^3)) - R(\nabla(a^1), \nabla(a^4), \nabla(a^3), \nabla(a^2)) \right. \\ & \quad \left. + \left(2\nabla(a^1)[\Delta(a^4)] + \frac{2m}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^4), e_{\alpha}) - \frac{1}{6}g(\nabla(a^1), \nabla(a^4))s \right) g(da^2, da^3) \right) d\text{Vol}_M. \end{aligned}$$

In the same way, we take four different parts for A and B , and then by further calculation, we get the following result.

$$\begin{aligned} & \oint a^0 [D^2, a^1][D^2, a^2][D, a^3][D, a^4] D^{-2m} - \oint a^0 [D, a^1][D^2, a^2][D^2, a^3][D, a^4] D^{-2m} \\ &+ \oint a^0 [D, a^1][D, a^2][D^2, a^3][D^2, a^4] D^{-2m} - \oint a^0 [D^2, a^1][D, a^2][D, a^3][D^2, a^4] D^{-2m} \\ &= 2^m \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \int_M a_0 \left\{ -2\Delta(a^1) g(\nabla(a^2), \nabla g(a^3, a^4)) - 4\nabla(a^1) \nabla(a^2) [g(da^3, da^4)] - 2R(\nabla(a^1), \nabla(a^2), \nabla(a^4), \nabla(a^3)) \right. \\ & \quad \left. + \left(2\nabla(a^1)[\Delta(a^2)] - \Delta(a^1)\Delta(a^2) - \frac{1}{6}g(\nabla(a^1), \nabla(a^2))s + \frac{2m}{3(m+1)} \sum_{\alpha} R(\nabla(a^1), e_{\alpha}, \nabla(a^2), e_{\alpha}) \right) g(da^3, da^4) \right. \\ & \quad \left. + \Delta(a^2)\Delta(a^3)g(da^1, da^4) - 2\Delta(a^2)g(\nabla(a^1), \nabla_{\nabla(a^3)}\nabla(a^4)) + 2R(\nabla(a^2), \nabla(a^3), \nabla(a^4), \nabla(a^1)) \right. \\ & \quad \left. - 2\nabla(a^2)[\Delta(a^3)]g(da^1, da^4) - 2\Delta(a^3)g(\nabla(a^1), \nabla_{\nabla(a^2)}\nabla(a^4)) + 4g(\nabla(a^1), \nabla_{\nabla(a^2)}\nabla_{\nabla(a^3)}\nabla(a^4)) \right. \\ & \quad \left. - 4g(\nabla_{\nabla(a_1)}\nabla(a_4), \nabla_{\nabla(a_2)}\nabla(a_3)) - 2R(\nabla(a^1), \nabla(a^3), \nabla(a^2), \nabla(a^4)) - 4g(\nabla(a^1), \nabla_{\nabla_{\nabla(a^2)}}\nabla(a^3)\nabla(a^4)) \right. \\ & \quad \left. + \frac{1}{6}g(\nabla(a^2), \nabla(a^3))g(da^1, da^4)s - \frac{2m}{3(m+1)} \sum_{\alpha} R(\nabla(a^2), e_{\alpha}, \nabla(a^3), e_{\alpha})g(da^1, da^4) + \left[2\nabla(a^3)[\Delta(a^4)] \right. \right. \\ & \quad \left. \left. - \Delta(a^3)\Delta(a^4) - \frac{1}{6}g(\nabla(a^3), \nabla(a^4))s + \frac{2m}{3(m+1)} \sum_{\alpha} R(\nabla(a^3), e_{\alpha}, \nabla(a^4), e_{\alpha})(x_0) \right) g(da^1, da^2) \right] \\ & \quad + \Delta(a^1)\Delta(a^4)g(da^2, da^3) + 2\Delta(a^1)g(\nabla(a^4), \nabla g(da^2, da^3)) - 2\nabla(a^1)[\Delta(a^4)]g(da^2, da^3) \end{aligned}$$

$$-\left(\frac{2m}{3(m+1)}\sum_{\alpha}R(\nabla(a^1),e_{\alpha},\nabla(a^4),e_{\alpha})-\frac{1}{6}g(\nabla(a^1),\nabla(a^4))s\right)g(da^2,da^3)\Big\}dVol_M. \quad (3.24)$$

When $n = 6$, $m = 3$, we have $area(S_6) = \pi^3$, then we get the following theorem.

Theorem 3.12. *Let $(\mathcal{A}, \mathcal{H}, D)$ be the spectral triple associated to a compact spin Riemannian 6-dimensional manifold M . Then a Connes-Chamseddine Hochschild cocycle is given*

$$\begin{aligned} & \rho(a^0, a^1, a^2, a^3, a^4) \\ &= 8\pi^3 \int_M a_0 \left\{ -2\Delta(a^1)g(\nabla(a^2), \nabla g(a^3, a^4)) - 4\nabla(a^1)\nabla(a^2)[g(da^3, da^4)] - 2R(\nabla(a^2), \nabla(a^3), \nabla(a^4), \nabla(a^1)) \right. \\ &+ \left(2\nabla(a^1)[\Delta(a^2)] - \Delta(a^1)\Delta(a^2) - \frac{1}{6}g(\nabla(a^1), \nabla(a^2))s + \frac{1}{2}\sum_{\alpha}R(\nabla(a^1), e_{\alpha}, \nabla(a^2), e_{\alpha}) \right)g(da^3, da^4) \\ &+ \Delta(a^2)\Delta(a^3)g(da^1, da^4) - 2\Delta(a^2)g(\nabla(a^1), \nabla_{\nabla(a^3)}\nabla(a^4)) + 2R(\nabla(a^2), \nabla(a^3), \nabla(a^4), \nabla(a^1)) \\ &- 2\nabla(a^2)[\Delta(a^3)]g(da^1, da^4) - 2\Delta(a^3)g(\nabla(a^1), \nabla_{\nabla(a^2)}\nabla(a^4)) + 4g(\nabla(a^1), \nabla_{\nabla(a^2)}\nabla_{\nabla(a^3)}\nabla(a^4)) \\ &- 4g(\nabla_{\nabla(a_1)}\nabla(a_4), \nabla_{\nabla(a_2)}\nabla(a_3)) - 2R(\nabla(a^1), \nabla(a^3), \nabla(a^2), \nabla(a^4)) - 4g(\nabla(a^1), \nabla_{\nabla(a^2)}\nabla(a^3)\nabla(a^4)) \\ &+ \frac{1}{6}g(\nabla(a^2), \nabla(a^3))g(da^1, da^4)s - \frac{1}{2}\sum_{\alpha}R(\nabla(a^2), e_{\alpha}, \nabla(a^3), e_{\alpha})g(da^1, da^4) + \left[2\nabla(a^3)[\Delta(a^4)] \right. \\ &- \Delta(a^3)\Delta(a^4) - \frac{1}{6}g(\nabla(a^3), \nabla(a^4))s + \frac{1}{2}\sum_{\alpha}R(\nabla(a^3), e_{\alpha}, \nabla(a^4), e_{\alpha})(x_0) \Big)g(da^1, da^2) \Big] \\ &+ \Delta(a^1)\Delta(a^4)g(da^2, da^3) + 2\Delta(a^1)g(\nabla(a^4), \nabla g(da^2, da^3)) - 2\nabla(a^1)[\Delta(a^4)]g(da^2, da^3) \\ &\left. - \left(\frac{1}{2}\sum_{\alpha}R(\nabla(a^1), e_{\alpha}, \nabla(a^4), e_{\alpha}) - \frac{1}{6}g(\nabla(a^1), \nabla(a^4))s \right)g(da^2, da^3) \right\}dVol_M. \quad (3.25) \end{aligned}$$

4. Appendix

In this appendix, we will prove some facts used in Lemma 3.4 and in the computation of the residue.

Lemma 4.1. *(Also see (4.3) in [7])*

$$\begin{aligned} (1) \sigma_{-2m}(D^{-2m}) &= |\xi|^{-2m}; \\ (2) \sigma_{-2m-1}(D^{-2m}) &= m|\xi|^{-(2m-2)} \left(-i|\xi|^{-4}\xi_k(\Gamma^k - 2\sigma^k) - 2i|\xi|^{-6}\xi^j\xi_{\alpha}\xi_{\beta}\partial_{x_j}g^{\alpha\beta} \right) \\ &+ 2i\sum_{k=0}^{m-2}\sum_{\mu=1}^{2m}(k+1-m)|\xi|^{-2m-4}\xi^{\mu}\xi_{\alpha}\xi_{\beta}\partial_{\mu}^xg^{\alpha\beta}; \\ (3) \sigma_{-2m-2}(D^{-2m})(x_0) &= -\frac{m}{4}|\xi|^{-2m-2}s + \frac{m(m+1)}{3}|\xi|^{-2m-4}\xi_{\mu}\xi_{\nu}R_{\mu\alpha\nu\alpha}(x_0), \end{aligned} \quad (4.1)$$

where s is the scalar curvature.

Proof. (1) holds obviously.

(2) By (3.8) in [15], we have

$$\sigma_{1-\bar{n}}(D^{-\bar{n}+2}) = \frac{\bar{n}-2}{2}\sigma_2^{(-\frac{\bar{n}}{2}+2)}\sigma_{-3}(D^{-2}) - i\sum_{k=0}^{\frac{\bar{n}}{2}-3}\partial_{\xi_{\mu}}\sigma_2^{(-\frac{\bar{n}}{2}+k+2)}\partial_{x_{\mu}}\sigma_2^{-1}(\sigma_2^{-1})^k. \quad (4.2)$$

By plugging $\bar{n} = 2m + 2$ into the above equation, we get

$$\sigma_{-2m-1}(D^{-2m}) = m\sigma_2^{(-m+1)}\sigma_{-3}(D^{-2}) - i \sum_{k=0}^{m-3} \sum_{\mu=1}^{2m} \partial_{\xi_\mu} \sigma_2^{(-m+k+1)} \partial_{x_\mu} \sigma_2^{-1} (\sigma_2^{-1})^k. \quad (4.3)$$

Then, by $\sigma_{-3}(D^{-2}) = -i|\xi|^{-4}\xi_k(\Gamma^k - 2\sigma^k) - 2i|\xi|^{-6}\xi^j\xi_\alpha\xi_\beta\partial_{x_j}g^{\alpha\beta}$, $\partial_{\xi_\mu} \sigma_2^{(-\frac{n}{2}+k+2)} = 2(k+1-m)|\xi|^{-2m+2k}\xi^\mu$ and $\partial_{x_\mu} \sigma_2^{-1} = -|\xi|^{-4}\xi_\alpha\xi_\beta\partial_{x_j}g^{\alpha\beta}$ and (2) holds.

(3) Combined with [10], we make a new arrangement of the proof process of (4.24) in [10]. Firstly, using $\sigma_{-2m}^{\tilde{\Delta}^{-m}} \equiv \sigma_2^{-m}$ we get the recursion relations

$$\begin{aligned} \sigma_{-n}^{\tilde{\Delta}^{-\frac{n}{2}+1}}(x, \xi) &= \sum_{|\alpha|=0}^2 \sum_{k=2}^{4-|\alpha|} (-i)^{|\alpha|} \frac{1}{\alpha!} \partial_\xi^\alpha \sigma_{|\alpha|+k-n}^{\tilde{\Delta}^{-\frac{n}{2}+2}} \partial_x^\alpha \sigma_{-k}^{\tilde{\Delta}^{-1}} \\ &= \sigma_{2-n}^{\tilde{\Delta}^{-\frac{n}{2}+2}} \sigma_2^{-1} + \sigma_{3-n}^{\tilde{\Delta}^{-\frac{n}{2}+2}} \sigma_{-3}^{\tilde{\Delta}^{-1}} + \sigma_{4-n}^{\tilde{\Delta}^{-\frac{n}{2}+2}} \sigma_{-4}^{\tilde{\Delta}^{-1}} \\ &\quad - i \partial_{\xi_\mu} \sigma_{3-n}^{\tilde{\Delta}^{-\frac{n}{2}+2}} \partial_{x_\mu} \sigma_2^{-1} - i \partial_{\xi_\mu} \sigma_2^{-\frac{n}{2}+2} \partial_{x_\mu} \sigma_{-3}^{\tilde{\Delta}^{-1}} \\ &\quad - \frac{1}{2} \partial_{\xi_\mu} \partial_{\xi_\nu} \sigma_2^{-\frac{n}{2}+2} \partial_{x_\mu} \partial_{x_\nu} \sigma_2^{-1}, \end{aligned} \quad (4.4)$$

where $\tilde{\Delta} = D^2$.

After further iterations

$$\sigma_{3-n}^{\tilde{\Delta}^{-\frac{n}{2}+2}}(x_0, \xi) = \left(\frac{1}{2}n - 2\right) \sigma_2^{-\frac{n}{2}+3} \sigma_{-3}^{\tilde{\Delta}^{-1}}(x_0, \xi). \quad (4.5)$$

Write

$$\sigma(\tilde{\Delta}) = \sigma_2 + \sigma_1 + \sigma_0; \quad \sigma(\tilde{\Delta}^{-1}) = \sum_{j=2}^{\infty} b_{-j}. \quad (4.6)$$

By the composition formula of pseudodifferential operators, we have

$$1 = \sigma(\tilde{\Delta} \circ \tilde{\Delta}^{-1}) = \sum_{\alpha} \frac{(-i)^{|\alpha|}}{\alpha!} \partial_\xi^\alpha [\sigma(\tilde{\Delta})] \partial_x^\alpha [\sigma(\tilde{\Delta}^{-1})], \quad (4.7)$$

so

$$\begin{aligned} b_{-2} &= \sigma_2^{-1}; \quad b_{-3} = -\sigma_2^{-1}[\sigma_1\sigma_2^{-1} - i \sum_j \partial_{\xi_j}(\sigma_2)\partial_{x_j}(b_{-2})]; \\ b_{-4} &= -\sigma_2^{-1}[\sigma_1b_{-3} + \sigma_0b_{-2} - i \sum_j \partial_{\xi_j}(\sigma_1)\partial_{x_j}(b_{-2}) - i \sum_j \partial_{\xi_j}(\sigma_2)\partial_{x_j}(b_{-3}) - \frac{1}{2} \sum_j \partial_{\xi_j}\partial_{\xi_k}(\sigma_2)\partial_{x_j}\partial_{x_k}(b_{-2})]. \end{aligned} \quad (4.8)$$

Uingsing $\partial_{x_\mu}(b_{-2})(x_0) = b_{-3}(x_0) = 0$, we have

$$b_{-4}(x_0) = \sigma_{-4}^{\tilde{\Delta}^{-1}}(x_0, \xi) = -\sigma_2^{-1}[\sigma_0b_{-2} - i \sum_j \partial_{\xi_j}(\sigma_2)\partial_{x_j}(b_{-3}) - \frac{1}{2} \sum_j \partial_{\xi_j}\partial_{\xi_k}(\sigma_2)\partial_{x_j}\partial_{x_k}(b_{-2})] \quad (4.9)$$

Then

$$-i \sum_j \partial_{\xi_j}(\sigma_2)\partial_{x_j}(b_{-3}) = -\sigma_2\sigma_{-4}^{\tilde{\Delta}^{-1}} - \sigma_0\sigma_2^{-1} + \frac{1}{2} \sum_j \partial_{\xi_j}\partial_{\xi_k}(\sigma_2)\partial_{x_j}\partial_{x_k}(b_{-2}). \quad (4.10)$$

Inserting Eq. (4.5) and Eq. (4.10) in relation (4.4) yields

$$\begin{aligned}
\sigma_{-n}^{\tilde{\Delta}-\frac{n}{2}+1}(x_0, \xi) &= \sigma_{2-n}^{\tilde{\Delta}-\frac{n}{2}+1} \sigma_2^{-1} + \sigma_2^{-\frac{n}{2}+2} \sigma_{-4}^{\tilde{\Delta}-1} + (2 - \frac{n}{2}) \sigma_2^{-\frac{n}{2}+1} [-\sigma_2^{-\frac{n}{2}+2} \sigma_{-4}^{\tilde{\Delta}-1} - \sigma_0 \sigma_2^{-1} \\
&\quad + \frac{1}{2} \sum_j \partial_{\xi_j} \partial_{\xi_k} (\sigma_2) \partial_{x_j} \partial_{x_k} (b_{-2})] - \frac{1}{2} \sum_j \partial_{\xi_j} \partial_{\xi_k} (\sigma_2^{-\frac{n}{2}+2}) \partial_{x_j} \partial_{x_k} (b_{-2}) \\
&= \sigma_{2-n}^{\tilde{\Delta}-\frac{n}{2}+1} \sigma_2^{-1} + (\frac{n}{2} - 1) \sigma_2^{-\frac{n}{2}+2} \sigma_{-4}^{\tilde{\Delta}-1} + (\frac{n}{2} - 2) \sigma_2^{-\frac{n}{2}} \sigma_0 \\
&\quad + \frac{4}{3} (\frac{n}{2} - 2) (\frac{n}{2} - 1) \sigma_2^{-\frac{n}{2}} \sigma_2^{-2} \sum_{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta}(x_0) \xi_\alpha \xi_\beta \xi_\gamma \xi_\delta.
\end{aligned} \tag{4.11}$$

After the iteration, and by $\sum_{\alpha\beta\mu\nu} \xi_\alpha \xi_\beta \xi_\mu \xi_\nu R_{\alpha\beta\mu\nu}(x_0) = 0$, we obtain

$$\sigma_{-n}^{\tilde{\Delta}-\frac{n}{2}+1}(x_0, \xi) = \frac{n-2}{8} [n \sigma_2^{-\frac{n}{2}+2} \sigma_{-4}^{\tilde{\Delta}-1}(x_0, \xi) + (n-4) \sigma_2^{-\frac{n}{2}} \sigma_0]. \tag{4.12}$$

By plugging $n = 2m + 2$ into the above equation and Eq. (12b) in [11], then in normal coordinates, (3) holds. \square

(i - 1 - f):

$$\begin{aligned}
&\int_{|\xi|=1} \text{tr} \left(2 \sum_{jr} \partial_{x_j} (a^1) \partial_{x_j} \partial_{x_r}^2 (a^2) c(da^3) c(da^4) |\xi|^{-2m} \right) \sigma(\xi) \\
&= \left(2 \nabla(a^1)[\Delta(a^2)](x_0) - \frac{4}{3} \sum_\alpha R(\nabla(a^1), e_\alpha, \nabla(a^2), e_\alpha) \right) g(da^3, da^4) \text{tr}[id] Vol_{S^n}.
\end{aligned} \tag{4.13}$$

Proof. In the normal coordinate system, we can obtain

$$\begin{aligned}
\nabla(a^1)[\Delta(a^2)](x_0) &= \sum_j \partial_{x_j} (a^1) \partial_{x_j} [\Delta(a^2)](x_0) \\
&= \sum_j \partial_{x_j} (a^1) \partial_{x_j} [(- \sum_{\alpha\beta} g^{\alpha\beta} \partial_{x_\alpha} \partial_{x_\beta} + \sum_{\alpha\beta} g^{\alpha\beta} \Gamma_{\alpha\beta}^k \partial_{x_k})(a^2)](x_0) \\
&= \sum_j \partial_{x_j} (a^1) [- \sum_{\alpha\beta} \partial_{x_j} (g^{\alpha\beta}) \partial_{x_\alpha} \partial_{x_\beta} - \sum_{\alpha\beta} g^{\alpha\beta} \partial_{x_j} \partial_{x_\alpha} \partial_{x_\beta} \\
&\quad + g^{\alpha\beta} (x_0) \sum_{jk} \partial_{x_j} (\Gamma_{\alpha\beta}^k) \partial_{x_k} + \sum_{jk} \Gamma_{\alpha\beta}^k \partial_{x_j} \partial_{x_k}] (a^2)(x_0) \\
&= - \sum_{jr} \partial_{x_j} (a^1) \partial_{x_j} \partial_{x_r}^2 (a^2) + \frac{2}{3} \sum_\alpha R(\nabla(a^1), e_\alpha, \nabla(a^2), e_\alpha).
\end{aligned} \tag{4.14}$$

According to the following formula, we have

$$\sum_{jr} \partial_{x_j} (a^1) \partial_{x_j} \partial_{x_r}^2 (a^2) = \frac{2}{3} \sum_\alpha R(\nabla(a^1), e_\alpha, \nabla(a^2), e_\alpha) - \nabla(a^1)[\Delta(a^2)]. \tag{4.15}$$

Then, after the integration, (4.13) holds. \square

(i - 1 - g):

$$\int_{|\xi|=1} \text{tr} \left[4 \sum_{jp} \partial_{x_j} (a^1) \partial_{x_j} \left(\partial_{x_p} (a^2) \partial_{x_p} [c(da^3) c(da^4)] \right) |\xi|^{-2m} \right] \sigma(\xi) = -4 \nabla(a^1) \nabla(a^2) [g(da^3, da^4)] \text{tr}[id] Vol_{S^n}. \tag{4.16}$$

Proof. Take $|\xi| = 1$, we can change the order of trace and derivative. Then

$$\begin{aligned} \text{tr} \left[\sum_{jp} \partial_{x_j}(a^1) \partial_{x_j} \left(\partial_{x_p}(a^2) \partial_{x_p} [c(da^3)c(da^4)] \right) |\xi|^{-2m} \right] &= \nabla(a^1) \sum_p \partial_{x_p}(a^2) \partial_{x_p} [-g(da^3, da^4)] \text{tr}[id] \\ &= -\nabla(a^1) \nabla(a^2) [g(da^3, da^4)] \text{tr}[id]. \end{aligned}$$

Therefore, after the integration, (4.16) holds. \square

(ii - 1 - b):

$$\int_{|\xi|=1} \text{tr} \left(2 \sum_{kp} \partial_{x_k}^2(a^2) \partial_{x_p}(a^3) c(da^1) \partial_{x_p} [c(da^4)] |\xi|^{-2m} \right) \sigma(\xi) = 2\Delta(a^2) g \left(\nabla(a^1), \nabla_{\nabla(a^3)} [\nabla(a^4)] \right) \text{tr}[id] Vol_{S^n}. \quad (4.17)$$

Proof. By $c(da^1) = \sum_\mu e_\mu(a^1)c(e_\mu)$ and $c(da^4) = \sum_\nu e_\nu(a^4)c(e_\nu)$, we have

$$\begin{aligned} \text{tr} \left(\sum_{kp} \partial_{x_k}^2(a^2) \partial_{x_p}(a^3) c(da^1) \partial_{x_p} [c(da^4)] \right) (x_0) &= -\sum_{kp} \partial_{x_k}^2(a^2) \partial_{x_p}(a^3) \sum_\mu e_\mu(a^1) \partial_{x_p} [e_\mu(a^4)] (x_0) \text{tr}[id] \\ &= \Delta(a^2) \sum_p \partial_{x_p}(a^3) \sum_\mu e_\mu(a^1) \partial_{x_p} [e_\mu(a^4)] (x_0) \text{tr}[id] \\ &= \Delta(a^2) \sum_\mu e_\mu(a^1) \sum_p \partial_{x_p}(a^3) \partial_{x_p} [e_\mu(a^4)] (x_0) \text{tr}[id] \\ &= \Delta(a^2) \sum_\mu e_\mu(a^1) \nabla(a^3) [e_\mu(a^4)] (x_0) \text{tr}[id]. \end{aligned}$$

And

$$\begin{aligned} g(\nabla(a^1), \nabla_{\nabla(a^3)} (\nabla(a^4))) (x_0) &= g \left(\sum_\mu e_\mu(a^1) e_\mu, \nabla_{\nabla(a^3)} \left(\sum_l e_l(a^4) e_l \right) \right) (x_0) \\ &= g \left(\sum_\mu e_\mu(a^1) e_\mu, \nabla_{\nabla(a^3)} \left(\sum_l e_l(a^4) e_l \right) \right) (x_0) \\ &= g \left(\sum_\mu e_\mu(a^1) e_\mu, \sum_l \nabla(a^3)(e_l(a^4)) e_l + \sum_l e_l(a^4) \nabla_{\nabla(a^3)} e_l \right) (x_0) \\ &= \sum_\mu e_\mu(a^1) \nabla(a^3) [e_\mu(a^4)]. \end{aligned}$$

Then

$$\Delta(a^2) \sum_\mu e_\mu(a^1) \nabla(a^3) [e_\mu(a^4)] (x_0) \text{tr}[id] = \Delta(a^2) g \left(\nabla(a^1), \nabla_{\nabla(a^3)} [\nabla(a^4)] \right) \text{tr}[id].$$

Therefore, after the integration, (4.17) holds. \square

(ii - 1 - g):

$$\begin{aligned} \int_{|\xi|=1} \text{tr} \left(4 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_p}(a^3) c(da^1) \partial_{x_j} \partial_{x_p} [c(da^4)] |\xi|^{-2m} \right) \sigma(\xi) &= -4 \left(g(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla_{\nabla(a^3)} \nabla(a^4)) - g(\nabla_{\nabla(a_1)} \nabla(a_4), \nabla_{\nabla(a_2)} \nabla(a_3)) \right. \\ &\quad \left. - \frac{1}{2} R(\nabla(a^2), \nabla(a^3), \nabla(a^1), \nabla(a^4)) \right) \text{tr}[id] Vol_{S^n}. \quad (4.18) \end{aligned}$$

Proof. By $c(da^1) = \sum_\mu e_\mu(a^1)c(e_\mu)$ and $c(da^4) = \sum_\nu e_\nu(a^4)c(e_\nu)$, we have

$$\text{tr} \left(\sum_{jp} \partial_{x_j}(a^2) \partial_{x_p}(a^3) c(da^1) \partial_{x_j} \partial_{x_p} [c(da^4)] \right) = - \sum_{jp} \partial_{x_j}(a^2) \partial_{x_p}(a^3) \sum_\mu e_\mu(a^1) \partial_{x_j} \partial_{x_p} [e_\mu(a^4)] \text{tr}[id].$$

Moreover

$$\begin{aligned} & g \left(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla_{\nabla(a^3)} \nabla(a^4) \right) (x_0) \\ &= g \left(\nabla(a^2), \nabla_{\nabla(a^1)} \nabla_{\nabla(a^3)} \left(\sum_\mu e_\mu(a^4) e_\mu \right) \right) (x_0) \\ &= g \left(\nabla(a^2), \sum_\mu \nabla_{\nabla(a^1)} [\nabla(a^3) e_\mu(a^4)] e_\mu + \sum_\mu e_\mu(a^4) \nabla_{\nabla(a^3)} e_\mu \right) (x_0) \\ &= g \left(\nabla(a^2), \sum_\mu \nabla(a^1) [\nabla(a^3) e_\mu(a^4)] e_\mu \right) (x_0) + g \left(\nabla(a^2), \sum_\mu \nabla(a^1) e_\mu(a^4) \nabla_{\nabla(a^3)} e_\mu \right) (x_0) \\ &\quad + g \left(\nabla(a^2), \sum_\mu \nabla(a^3) e_\mu(a^4) \nabla_{\nabla(a^1)} e_\mu \right) (x_0) + g \left(\nabla(a^2), \sum_\mu e_\mu(a^4) \nabla_{\nabla(a^1)} \nabla_{\nabla(a^3)} e_\mu \right) (x_0) \\ &= g \left(\nabla(a^2), \sum_\mu \nabla(a^1) (\nabla(a^3) e_\mu(a^4)) e_\mu \right) (x_0) + g \left(\nabla(a^2), \sum_\mu e_\mu(a^4) \nabla_{\nabla(a^1)} \nabla_{\nabla(a^3)} e_\mu \right) (x_0), \end{aligned}$$

where

$$\begin{aligned} & g \left(\nabla(a^2), \sum_\mu \nabla(a^1) (\nabla(a^3) e_\mu(a^4)) e_\mu \right) (x_0) \\ &= g \left(\partial_{x_j}(a^2) e_j, \sum_{\mu l q} \partial_{x_1}(a^1) \partial_{x_l} [\partial_{x_q}(a^3) \partial_{x_q}(e_\mu(a^4))] e_\mu \right) (x_0) \\ &= \sum_{jlp} \partial_{x_j}(a^2) \partial_{x_l}(a^1) \partial_{x_q}(a^3) \partial_{x_l} [\partial_{x_q}(e_\mu(a^4))] e_\mu + \sum_{jlq} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \partial_{x_q} \partial_{x_l}(a^3) \partial_{x_j} \partial_{x_q}(a^4) \\ &= \sum_{jlp} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \partial_{x_q}(a^3) \partial_{x_l} \partial_{x_q} [e_j(a^4)] + \sum_{jlq} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \partial_{x_q} \partial_{x_l}(a^3) \partial_{x_j} \partial_{x_q}(a^4); \\ & g \left(\nabla_{\nabla(a^1)} \nabla(a^4), \nabla_{\nabla(a^2)} \nabla(a^3) \right) = \sum_{jlq} \partial_{x_j}(a^1) \partial_{x_l}(a^2) \partial_{x_q} \partial_{x_l}(a^3) \partial_{x_j} \partial_{x_q}(a^4); \\ & g \left(\nabla(a^2), \sum_\mu e_\mu(a^4) \nabla_{\nabla(a^1)} \nabla_{\nabla(a^3)} e_\mu \right) (x_0) \\ &= g \left(\nabla(a^2), \sum_\mu e_\mu(a^4) \nabla_{\nabla(a^1)} \sum_{jl} g^{jl} \frac{\partial a^3}{\partial x_j} \nabla_{\partial_{x_l}} e_\mu \right) \\ &= g \left(\nabla(a^2), \sum_\mu e_\mu(a^4) \sum_j \frac{\partial a^3}{\partial x_j}(x_0) \nabla_{\nabla(a^1)} \nabla_{\partial_{x_j}} e_\mu \right) \\ &= g \left(\nabla(a^2), \sum_\mu e_\mu(a^4) \sum_{jl} \frac{\partial a^3}{\partial x_j} \frac{\partial a^1}{\partial x_l} \sum_\alpha (\partial_{x_l} H_{j\alpha\mu} e_\alpha + H_{j\alpha\mu} \nabla_{\partial_{x_l}} e_\alpha)(x_0) \right) \\ &= \frac{1}{2} \sum_{\mu j l \alpha} e_\mu(a^4) e_j(a^3) e_l(a^1) R_{l j \alpha \mu}(x_0) e_\alpha(a^2) \end{aligned}$$

$$= \frac{1}{2} R(\nabla(a^1), \nabla(a^3), \nabla(a^2), \nabla(a^4)),$$

where $H_{j\alpha\mu} = g(\nabla_{\partial_{x_j}} e_\mu, e_\alpha)$. Therefore

$$\begin{aligned} & \sum_{jp} \partial_{x_j}(a^2) \partial_{x_p}(a^3) \sum_\mu e_\mu(a^1) \partial_{x_j} \partial_{x_p}[e_\mu(a^4)] \\ &= g(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla_{\nabla(a^3)} \nabla(a^4)) - g(\nabla_{\nabla(a_1)} \nabla(a_4), \nabla_{\nabla(a_2)} \nabla(a_3)) - \frac{1}{2} R(\nabla(a^1), \nabla(a^3), \nabla(a^2), \nabla(a^4)). \end{aligned}$$

After the integration, (4.18) holds. \square

(ii - 1 - h):

$$\int_{|\xi|=1} \text{tr} \left(4 \sum_{jp} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_p}(a^3) c(da^1) \partial_{x_p}[c(da^4)] |\xi|^{-2m} \right) \sigma(\xi) = -4g(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla_{\nabla(a^3)} \nabla(a^4)) \text{tr}[id] Vol_{S^n}. \quad (4.19)$$

Proof. By $c(da^1) = \sum_\mu e_\mu(a^1)c(e_\mu)$ and $c(da^4) = \sum_\nu e_\nu(a^4)c(e_\nu)$, we have

$$\text{tr} \left(\sum_{jp} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_p}(a^3) c(da^1) \partial_{x_p}[c(da^4)] \right) (x_0) = - \sum_{jp} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_p}(a^3) \sum_\mu e_\mu(a^1) \partial_{x_p}[e_\mu(a^4)] (x_0) \text{tr}[id].$$

Moreover

$$\begin{aligned} \nabla_{\nabla(a^2)} \nabla(a^3)(x_0) &= \sum_j e_j(a^2) \nabla_{e_j} [\sum_l e_l(a^3) e_l](x_0) \\ &= \sum_j e_j(a^2) \sum_l e_j e_l(a^3) e_l(x_0) \\ &= \sum_j \partial_{x_j}(a^2) \sum_l \partial_{x_j} \partial_{x_l}(a^3) \partial_{x_l}(x_0). \end{aligned}$$

Then

$$\begin{aligned} \sum_{jp} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_p}(a^3) \sum_\mu e_\mu(a^1) \partial_{x_p}[e_\mu(a^4)] (x_0) \text{tr}[id] &= \sum_{jp} \partial_{x_j}(a^2) \partial_{x_j} \partial_{x_p}(a^3) g \left(\nabla(a^1), \nabla_{\partial_{x_p}} \nabla(a^4) \right) (x_0) \text{tr}[id] \\ &= g(\nabla(a^1), \nabla_{\nabla(a^2)} \nabla(a^3) \nabla(a^4)) \text{tr}[id]. \end{aligned}$$

After the integration, (4.19) holds. \square

Declarations

Ethics approval and consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and materials: The authors confirm that the data supporting the findings of this study are available within the article.

Competing interests: The authors declare no competing interests.

Funding: This research was funded by National Natural Science Foundation of China: No.11771070, 2023-BSBA-118 and N2405015..

Author Contributions: All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by TW and YW. The first draft of the manuscript was written by TW and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by NSFC No.11771070, 2023-BSBA-118 and N2405015. The authors thank the referee for his (or her) careful reading and helpful comments.

References

References

- [1] Ackermann T. A note on the Wodzicki residue. *J. Geom. Phys.* 1996, 20: 404-406.
- [2] Bochniak A, Sitarz A. Spectral interaction between universes. *J. Cosmol. Astropart. P.* 2022, 4: 055.
- [3] Connes A. The action functional in Noncommutative geometry. *Comm. Math. Phys.* 1998, 117: 673-683.
- [4] Connes A. Quantized calculus and applications. 11th International Congress of Mathematical Physics (Paris,1994). Internat Press, Cambridge, MA. 1995, 15-36.
- [5] Connes A , Chamseddine A H. Inner fluctuations of the spectral action. *J. Geom. Phys.* 2007, 57(1): 1-21.
- [6] Connes A. Noncommutative Geometry. Academic Press, San Diego. 1994.
- [7] Dabrowski L, Sitarz A, Zalecki P. Spectral metric and Einstein functionals. *Adv Math.* 2023, 427: 109128.
- [8] Figueroa H, Gracia-Bond'a J M, Lizzi F, et al. A nonperturbative form of the spectral action principle in noncommutative geometry. *J. Geom. Phys.* 1998, 26(3-4): 329-339.
- [9] Guillemin V W. A new proof of Weyl's formula on the asymptotic distribution of eigenvalues. *Adv. Math.* 1985, 55(2): 131-160.
- [10] Kalau W, Walze M. Gravity, Noncommutative geometry and the Wodzicki residue. *J. Geom. Phys.* 1995, 16: 327-344.
- [11] Kastler D. The Dirac Operator and Gravitation. *Comm. Math. Phys.* 1995, 166: 633-643.
- [12] Ugalde W J. A construction of critical GJMS operators using Wodzicki's residue. *Commun. Math. Phys.* 2006, 261(3): 771-788.
- [13] Wang Y. A Kastler-Kalau-Walze type theorem and the spectral action for perturbations of Dirac operators on manifolds with boundary. *Abstr. Appl. Anal.* Art. 2014, ID 619120, 13 pp.
- [14] Wang Y. General Kastler-Kalau-Walze type theorems for manifolds with boundary II. *Int. J. Geom. Methods M. Phys.* 2019, 16(2): 1950028.
- [15] Wang J, Wang Y. A general Kastler-Kalau-Walze type theorem for manifolds with boundary. *Int. J. Geom. Methods M. Phys.* 2016, 13(1): 1650003.
- [16] Wang J, Wang Y, Wu T. Dirac operators with torsion, spectral Einstein functionals and the noncommutative residue. *J. Math. Phys.* 2023, 64(10). Paper No. 102505, 24 pp.
- [17] Wang J, Wang Y, Yang C L. Dirac operators with torsion and the noncommutative residue for manifolds with boundary. *J. Geom. Phys.* 2014, 81: 92-111.
- [18] Wodzicki M. Local invariants of spectral asymmetry. *Invent. Math.* 1995, 75(1): 143-178.
- [19] Wodzicki M. Noncommutative residue I: Fundamentals, in K-theory, Arithmetic and Geometry, Yu. I. Manin, ed., Lecture Notes in Mathematics Vol. 1289 (Springer, Berlin, 1987).