

MIXED MULTIQUADRATIC SPLITTING FIELDS

FATMA KADER BİNGÖL, ADAM CHAPMAN, AND AHMED LAGHRIBI

ABSTRACT. We study mixed multiquadratic field extensions as splitting fields for central simple algebras of exponent 2 in characteristic 2. As an application, we provide examples of nonexcellent mixed biquadratic field extensions.

CLASSIFICATION (MSC 2020): 11E04, 11E81, 16K20, 13A35

KEYWORDS: quaternion algebra, symbol length, excellent extension, characteristic 2

1. INTRODUCTION

Throughout this short note, F denotes a field of characteristic 2. Let $\mathrm{Br}_2(F)$ denote the 2-torsion subgroup of the Brauer group of F . It is well-known that $\mathrm{Br}_2(F)$ is generated by classes of F -quaternion algebras, see [2, Chap. VII, §9]. Recall that an F -quaternion algebra is a central simple F -algebra of degree 2. For $a \in F$ and $b \in F^\times$, we denote by $[a, b]_F$ the F -quaternion algebra generated by two elements i and j satisfying the relations

$$i^2 + i = a, \quad j^2 = b \quad \text{and} \quad ji = (1 + i)j.$$

The quadratic extension $F(i)$ of F contained in $[a, b]_F$ is a separable extension, while $F(j)$ is inseparable. Any F -quaternion algebra is isomorphic to $[a, b]_F$ for certain $a \in F$ and $b \in F^\times$.

Given a class $[A]$ in $\mathrm{Br}_2(F)$ of a central simple F -algebra A of exponent at most 2, we may ask for the smallest $m \in \mathbb{N}$ such that $[A]$ is equal to the class of a tensor product of m F -quaternion algebras. This numerical invariant m of A is denoted by $\lambda_2(A)$ and called the 2-symbol length of A . In Section 2, we study the behavior of this invariant under multiquadratic extensions of separability degree at most 8 (Proposition 2.6). To this end we start with some descent results, in particular we prove Proposition 2.4 which is based on a transfer of central simple K -algebras induced by the usual Frobenius homomorphism when K is an inseparable quadratic extension of F . The main result of this paper, which is given in Section 3, is the application of our methods to the excellent property for quartic extensions.

Recall that a field extension K/F is called *excellent* if for any quadratic form φ over F , the anisotropic part $(\varphi_K)_{an}$ of the quadratic form φ_K obtained by extending scalars to K is defined over F , i.e. there exists a quadratic form ψ over F such that $(\varphi_K)_{an} \simeq \psi_K$. It is well known that, in arbitrary characteristic, any quadratic extension is excellent [5, Example 29.2]. The same holds for degree 3 extensions, more generally for odd degree extensions, in view of

Springer's Theorem ([5, Corollary 18.5]). The situation changes for quartic extensions. Nonexcellent separable biquadratic extensions exist, see [6, §5], [13] for characteristic not 2 and [9, §6] for characteristic 2. In [9], the excellence property for inseparable quartic extensions is studied. They provide examples of simple inseparable quartic extensions which are nonexcellent. Purely inseparable biquadratic extensions are known to be excellent, see [7]. The remaining case for inseparable quartic extensions is then the case of mixed biquadratic extensions. In Section 3, we provide examples of nonexcellent mixed biquadratic extensions (Example 3.2), hence completing the picture for the study of excellence property for quartic extensions. Our examples are realized as subfields of non-decomposable cyclic algebras of degree 8 and exponent 2 defined over a field of characteristic 2. Our method applies also to simple purely inseparable quartic extensions contained in such an algebra, hence it gives new examples of nonexcellent simple purely inseparable quartic extensions as well.

Our main references are [5] for the theory of quadratic forms, and [2] for the theory of central simple algebras.

2. DESCENT OF QUATERNION ALGEBRAS

For $a_1, \dots, a_n \in F^\times$, let $\langle a_1, \dots, a_n \rangle_b$ denote the non-degenerate symmetric bilinear form

$$F^n \times F^n \rightarrow F, ((x_1, \dots, x_n), (y_1, \dots, y_n)) \mapsto \sum_{i=1}^n a_i x_i y_i.$$

We will be only considering nonsingular quadratic forms, meaning quadratic forms having polar form with trivial radical. Recall from [5, Corollary 7.32] that any such form φ over F decomposes as follows:

$$\varphi \cong [a_1, b_1] \perp \dots \perp [a_n, b_n]$$

for some $a_1, b_1, \dots, a_n, b_n \in F$, where for any scalars $a, b \in F$ we denote by $[a, b]$ the 2-dimensional quadratic form $aX^2 + XY + bY^2$. In particular, a nonsingular quadratic form is of even dimension. The Arf invariant of the form φ is defined to be $\sum_{i=1}^n a_i b_i + \varphi(F)$ in $F/\varphi(F)$, where $\varphi(F) = \{a^2 + a \mid a \in F\}$.

Let $W_q F$ be the Witt group of nonsingular quadratic forms over F , and WF the Witt ring of non-degenerate symmetric bilinear forms over F . The ideal of even-dimensional forms in WF is denoted by IF . The group $W_q F$ is endowed with a WF -module structure induced by tensor product in a natural way. For $n \geq 1$ an integer, let $I_q^n F$ be the product $I^{n-1} F \otimes W_q F$ (we take $I^0 F = WF$). The Hauptsatz of Arason-Pfister [5, Theorem 23.7] asserts that if the Witt class of an anisotropic quadratic form φ lies in $I_q^n F$, then $\dim \varphi \geq 2^n$.

For $n \geq 2$, the quadratic n -fold Pfister form is a nonsingular quadratic form isometric to

$$\langle 1, a_1 \rangle_b \otimes \dots \otimes \langle 1, a_{n-1} \rangle_b \otimes [1, a_n]$$

where $a_1, \dots, a_{n-1} \in F^\times$ and $a_n \in F$. This form is denoted by $\langle\langle a_1, \dots, a_{n-1}, a_n \rangle\rangle$. A quadratic 1-fold Pfister form is isometric to $[1, a]$ for some $a \in F$.

Let us make the following observation relevant to quadratic 2-fold Pfister forms for later use.

Lemma 2.1. *Let $x, y, a, b \in F$ with $x^2 + by^2 \neq 0 \neq b$. There exists some $c \in F$ such that $\langle\langle x^2 + by^2, a \rangle\rangle \simeq \langle\langle b, c \rangle\rangle$.*

Proof. If $x = 0$, then the statement holds with $c = a$. Assume now $x \neq 0$. We have that

$$\begin{aligned} \langle\langle x^2 + by^2, a \rangle\rangle &= [1, a] \perp (x^2 + by^2)[1, a] \\ &\simeq [x^2, x^{-2}a] \perp [x^2 + by^2, (x^2 + by^2)^{-1}a] \\ &\simeq [x^2, x^{-2}a + (x^2 + by^2)^{-1}a] \perp [by^2, (x^2 + by^2)^{-1}a] \\ &\simeq [1, c] \perp b[1, d] \end{aligned}$$

where $c = a(1 + x^2(x^2 + by^2)^{-1})$ and $d = aby^2(x^2 + by^2)^{-1}$. Note that the Arf invariant of $\langle\langle x^2 + by^2, a \rangle\rangle$ is trivial. Hence $c + d \in \wp(F)$. It follows that $\langle\langle x^2 + by^2, a \rangle\rangle \simeq \langle\langle b, c \rangle\rangle$ with $c \in F$. \square

For a finite field extension K/F and an F -linear functional $s : K \rightarrow F$, the transfer of the quadratic form φ (resp. a symmetric bilinear form \mathfrak{b}) over K with respect to s is denoted by $s_*\varphi$ (resp. $s_*\mathfrak{b}$) [5, §20.A]. The functional $s : K \rightarrow F$ induces group homomorphisms $s : I_q^n K \rightarrow I_q^n F$ for any $n \geq 1$; see [5, Corollary 34.17]. We will need the following particular case of Frobenius reciprocity: For a non-degenerate symmetric bilinear form \mathfrak{b} over K and a nonsingular quadratic form φ over F , we have $s_*(\mathfrak{b} \otimes \varphi) \cong s_*\mathfrak{b} \otimes \varphi$; see [5, Proposition 20.2].

Lemma 2.2. *Let K/F be a separable quadratic field extension and $s : K \rightarrow F$ a nonzero F -linear functional such that $s(1) = 0$. Let φ be an anisotropic even-dimensional quadratic form over K . Assume that $s_*\varphi$ is hyperbolic. Then there exists some nonsingular quadratic form ψ over F such that $\varphi \simeq \psi_K$. If furthermore φ has trivial Arf invariant, then ψ can be chosen to have trivial Arf invariant.*

Proof. The first statement follows by [5, Theorem 34.9]. For the second statement, assume further that φ has trivial Arf invariant. By the first part, there exists a nonsingular quadratic form ψ over F such that $\varphi \simeq \psi_K$.

Let $d \in F$ be a representative of the Arf invariant of ψ . Write $\psi \simeq \theta \perp [a, b]$ for a nonsingular quadratic form θ over F and some $a, b \in F$. As φ is anisotropic, we actually have $a, b \in F^\times$. We set $\psi' = \theta \perp [a, b + \frac{d}{a}]$. Then ψ' has trivial Arf invariant and we have that $\psi'_K \simeq \psi_K \simeq \varphi$. \square

Let K/F be a purely inseparable field extension such that $K^2 \subseteq F$. The Frobenius homomorphism $\text{Frob}_{K/F} : K \rightarrow F$, $x \mapsto x^2$ induces a group homomorphism

$$\text{Frob}_{K/F} : \text{Br}_2(K) \rightarrow \text{Br}_2(F), [A] \mapsto [\text{Frob}_{K/F} A].$$

For $x \in K$ and $y \in K^\times$, we have that $\text{Frob}_{K/F}([x, y]_K) = [x^2, y^2]_F$. See [3, §3] for details.

Recall furthermore from [5, Theorem 14.3] the group homomorphism

$$e_2 : I_q^2 F \rightarrow \text{Br}_2(F)$$

which is defined by mapping the Witt class of a quadratic form in $I_q^2 F$ to the Brauer class of its Clifford algebra. The kernel of e_2 is $I_q^3 F$ (see [5, §16]).

Lemma 2.3. *Let K/F be a quadratic field extension with $K^2 \subseteq F$ and let $s : K \rightarrow F$ be a nonzero F -linear functional such that $s(1) = 0$. Then the following diagram commutes:*

$$\begin{array}{ccc} I_q^2 K & \xrightarrow{s_*} & I_q^2 F \\ \downarrow e_2 & & \downarrow e_2 \\ \mathrm{Br}_2(K) & \xrightarrow{\mathrm{Frob}_{K/F}} & \mathrm{Br}_2(F) \end{array}$$

Proof. See [4, Lemma 3.1]. □

For central simple F -algebras A and B , we write $A \sim B$ to indicate that A and B are Brauer equivalent.

Proposition 2.4. *Let K/F be an inseparable quadratic field extension. Let A be a central simple K -algebra of exponent 2 and assume that $m \in \mathbb{N}_+$ is such that A is Brauer equivalent to a tensor product of m K -quaternion algebras. Assume that $\mathrm{Frob}_{K/F} A = 0$. There exist F -quaternion algebras H_1, \dots, H_n where $n \leq 2m - 1$ such that $A \sim \bigotimes_{i=1}^n H_i \otimes_F K$.*

Proof. Let Q_1, \dots, Q_m be K -quaternion algebras with $A \sim \bigotimes_{i=1}^m Q_i$. Let $a_i \in K$, $z_i \in K^\times$ be such that $Q_i \simeq [a_i, z_i]_K$ for $1 \leq i \leq m$. Since $[a_i, z_i]_K \simeq [a_i^2, z_i]_K$, we may assume that $a_i \in F$ for $1 \leq i \leq m$. Fix $b \in F^\times \setminus F^{\times 2}$ with $K = F(\sqrt{b})$. Let $s : K \rightarrow F$ be the F -linear functional with $s(1) = 0$ and $s(\sqrt{b}) = 1$. For $1 \leq i \leq m$, write $z_i = x_i + y_i \sqrt{b}$ with $x_i, y_i \in F$. Let $\lambda_i = y_i^{-1}$ if $y_i \neq 0$, otherwise let $\lambda_i = 1$. Finally we set $\varphi = \perp_{i=1}^m \lambda_i \langle\langle z_i, a_i \rangle\rangle$. Note that we have $e_2(\varphi) = [A]$. Using Frobenius reciprocity, we compute that the Witt class of $s_*\varphi$ is given by $\perp_{i=1}^m \langle\langle x_i^2 + y_i^2 b, a_i \rangle\rangle$. Lemma 2.1 provide $c_1, \dots, c_m \in F$ such that

$$\perp_{i=1}^m \langle\langle x_i^2 + y_i^2 b, a_i \rangle\rangle \simeq \perp_{i=1}^m \langle\langle b, c_i \rangle\rangle$$

which is Witt equivalent to $\langle\langle b, \sum_{i=1}^m c_i \rangle\rangle$. Furthermore by Lemma 2.3, we have

$$e_2(s_*\varphi) = \mathrm{Frob}_{K/F}(e_2(\varphi)) = \mathrm{Frob}_{K/F} A$$

which is trivial by our assumption. Hence the Witt class of $s_*\varphi$ lies in $I_q^3 F$. Since $s_*\varphi$ is Witt equivalent to $\langle\langle b, \sum_{i=1}^m c_i \rangle\rangle$, this implies that $s_*\varphi$ is hyperbolic, by the Hauptsatz of Arason-Pfister. It follows by Lemma 2.2 that $\varphi \simeq \psi_K$ for some nonsingular quadratic form ψ over F with trivial Arf invariant. In particular, we have $\dim \psi = \dim \varphi = 4m$. Then $e_2(\psi) = \sum_{i=1}^{2m-1} [H_i]$ for some F -quaternion algebras H_1, \dots, H_{2m-1} . It follows that

$$[A] = e_2(\varphi) = e_2(\psi_K) = \sum_{i=1}^{2m-1} [H_i \otimes_F K].$$

□

As a corollary, we retrieve the case $p = 2$ from [3, Proposition 4.6].

Corollary 2.5. *Let K/F be an inseparable quadratic field extension. Let A be a central simple F -algebra of exponent 2 and assume that A_K is not split. Then $A \sim B \otimes_F H$ where H is an F -quaternion algebra that splits over K*

and B is a central simple F -algebra with $\lambda_2(B) \leq 2\lambda_2(A_K) - 1$. In particular $\lambda_2(A) \leq 2\lambda_2(A_K)$.

Proof. Set $m = \lambda_2(A_K)$. We have by [3, Proposition 3.2] that $\text{Frob}_{K/F} A_K = 0$. Hence Proposition 2.4 provide F -quaternion algebras H_1, \dots, H_{2m-1} such that $A_K \sim \bigotimes_{i=1}^{2m-1} H_i \otimes_F K$. In particular, $A \otimes_F \bigotimes_{i=1}^{2m-1} H_i$ is split over K , hence is Brauer equivalent to an F -quaternion algebra H_{2m} that splits over K . It follows that $A \sim \bigotimes_{i=1}^{2m} H_i$. \square

Proposition 2.6. *Let A be a central simple F -simple algebra of exponent 2. Assume that A splits over $F(\alpha_1, \dots, \alpha_m, \sqrt{b_1}, \dots, \sqrt{b_n})$ where $\alpha_i^2 - \alpha_i \in F$ for $1 \leq i \leq m$ and $b_1, \dots, b_n \in F^\times \setminus F^{\times 2}$. Then $A \sim B \otimes_F \bigotimes_{i=1}^n [a_i, b_i]_F$ where $a_1, \dots, a_n \in F$ and B is a central simple F -algebra such that the following hold:*

- (1) *If $m = 1$, then $\lambda_2(B) \leq 1$, in particular $\lambda_2(A) \leq n + 1$.*
- (2) *If $m = 2$, then $\lambda_2(B) \leq 2^n + 1$, in particular $\lambda_2(A) \leq 2^n + n + 1$.*
- (3) *If $m = 3$, then $\lambda_2(B) \leq 3 \cdot 2^n + 1$, in particular $\lambda_2(A) \leq 3 \cdot 2^n + n + 1$.*

Proof. We prove the statement by induction on n . In the case where $n = 0$, (1) is clear, and (2) resp. (3) follows by [1] resp. [11]. We explain the proof of the statement only for the case where $m = 3$; the argument is the same for each cases.

Assume now $m = 3$, $n \geq 1$ and set $K = F(\sqrt{b_n})$. We have by induction hypothesis that $A_K \sim B' \otimes_K \bigotimes_{i=1}^{n-1} [a_i, b_i]_K$ for some $a_1, \dots, a_{n-1} \in K$ and a central simple K -algebra B' with $\lambda_2(B') \leq 3 \cdot 2^{n-1} + 1$. Since $[a_i, b_i]_K \simeq [a_i^2, b_i]_K$, we may assume that $a_1, \dots, a_{n-1} \in F$. Hence $(A \otimes_K \bigotimes_{i=1}^{n-1} [a_i, b_i]_F)_K \sim B'$. It follows now by Corollary 2.5 that

$$A \otimes_K \bigotimes_{i=1}^{n-1} [a_i, b_i]_F \sim B \otimes_F [a_n, b_n]_F$$

for $a_n \in F$ and a central simple F -algebra B with $\lambda_2(B) \leq 2(3 \cdot 2^{n-1} + 1) - 1 = 3 \cdot 2^n + 1$. Therefore $A \sim B \otimes_F \bigotimes_{i=1}^n [a_i, b_i]_F$ and we have $\lambda_2(A) \leq 3 \cdot 2^n + 1 + n$. \square

Remark 2.7. Let A be a central simple F -algebra of exponent 2. Assume that A splits over $F(\alpha_1, \dots, \alpha_m, \sqrt{b_1}, \dots, \sqrt{b_n})$ where $\alpha_i^2 - \alpha_i \in F$ for $1 \leq i \leq m$ and $b_1, \dots, b_n \in F^\times \setminus F^{\times 2}$. The bound in Proposition 2.6 for $m = 1$ coincides with the existing one; see [10, Théorème 4] also [3, Proposition 4.2]. For $m = 2$, it is shown in [10, Théorème 5] that $\lambda_2(A) \leq n + 3$, which can also be obtained from methods in [3]. Finally for $m = 3$, the methods in [3] applied to this case would give that $\lambda_2(A) \leq n + 7$. These bounds are better than what we get in (2) and (3) of Proposition 2.6 for $n \geq 2$. They coincide with our bounds from Proposition 2.6 for $n = 1$.

3. EXCELLENCE PROPERTY FOR BIQUADRATIC EXTENSIONS

In this section, we provide examples of nonexcellent mixed biquadratic extensions, completing the study of the excellence property of inseparable quartic extensions done in [9]. The same argument also applies to establish the nonexcellence of simple purely inseparable quartic extensions.

Proposition 3.1. *Let A be a central simple F -algebra of degree 8 and exponent 2. If A contains a mixed biquadratic or simple purely inseparable field extension of F which is excellent, then A decomposes into a tensor product of 3 quaternion algebras over F .*

Proof. Let M/F be an excellent quartic extension of F contained in A as in the statement, i.e. $M = F(\sqrt[4]{b})$ or $M = F(\alpha, \sqrt{b})$ where $\alpha^2 - \alpha = a' \in F$ and $b \in F^\times$. Set $K = F(\sqrt{b})$. Then $A_K \sim [a, x]_K \otimes_K [c, y]_K$ where $c \in F$, $x \in K^\times$, and $y = \sqrt{b}$, $a \in F$ when $M = F(\sqrt[4]{b})$ and $y \in K^\times$, $a = a'$ otherwise. Let $s : K \rightarrow F$ be the F -linear functional with $s(1) = 0$ and $s(\sqrt{b}) = 1$. We choose $\lambda \in F$ such that $s_*(\langle x \rangle_{\mathfrak{b}}) \perp \lambda s_*(\langle y \rangle_{\mathfrak{b}})$ is isotropic. We set $\varphi = \langle\langle x, a \rangle\rangle \perp \lambda \langle\langle y, c \rangle\rangle$ and observe that $e_2(\varphi) = [A_K]$. Using Frobenius reciprocity, we compute that the Witt class of $s_*\varphi$ is given by $s_*(\langle x \rangle_{\mathfrak{b}}) \otimes [1, a] \perp \lambda s_*(\langle y \rangle_{\mathfrak{b}}) \otimes [1, c]$, which is isotropic, since $s_*(\langle x \rangle_b) \perp \lambda s_*(\langle y \rangle_b)$ is isotropic. Therefore $\dim(s_*\varphi)_{an} < 8$. Now Lemma 2.3 implies that

$$e_2(s_*\varphi) = \text{Frob}_{K/F}(e_2(\varphi)) = \text{Frob}_{K/F} A_K$$

which is trivial by [3, Proposition 3.2]. Then the Witt class of $s_*\varphi$ lies in $I_q^3 F$. As $\dim(s_*\varphi)_{an} < 8$, the Hauptsatz of Arason-Pfister implies that $s_*\varphi$ is hyperbolic. Hence by Lemma 2.2, $\varphi \simeq \psi'_K$ for some nonsingular quadratic form ψ' over F with trivial Arf invariant. Note that $\dim(\psi'_M)_{an} = \dim(\varphi_M)_{an} \leq 4$. Since M/F is excellent, we have $(\psi'_M)_{an} \simeq \psi_M$ for some nonsingular quadratic form ψ over F . In particular $\dim \psi \leq 4$. It is easy to see that we can choose ψ with trivial Arf invariant. Then $e_2(\psi) = [H]$ for some F -quaternion algebra H . We have

$$[H_M] = e_2(\psi_M) = e_2(\psi'_M) = e_2(\varphi_M) = [A_M].$$

It follows that $A \otimes_F H$ splits over M . This implies that $A \otimes_F H$ is Brauer equivalent to a tensor product of 2 F -quaternion algebras. We deduce that A decomposes into a tensor product of 3 F -quaternion algebras. \square

Example 3.2. Let C be a cyclic algebra over a field F of characteristic 2 (e.g. a global field of characteristic 2) of degree and exponent equal to 8. Let K be the function field of the Severi-Brauer variety corresponding to $C \otimes_F C$ and set $D(8, 2) = C \otimes_F K$. Then $D(8, 2)$ is a cyclic division algebra of degree 8 and exponent 2 over K , see [12]. In [8], it is shown that $D(8, 2)$ does not decompose into a tensor product of 3 quaternion algebras. Hence by Proposition 3.1, any mixed biquadratic and simple purely inseparable field extension contained in $D(8, 2)$ is not excellent.

Acknowledgement. This work was done during some (online) meetings between the three authors when the first author was visiting the Laboratoire de Mathématiques de Lens in november 2023 in the framework of the project *IEA of CNRS* between the Artois University and the Academic College of Tel-Aviv-Yaffo. The three authors are grateful for the support of both institutions and CNRS. The first author is supported by the 2020 PRIN (project *Derived and underived algebraic stacks and applications*) from MIUR, and by research funds from Scuola Normale Superiore.

REFERENCES

- [1] A.A. Albert. Normal division algebras of degree four over an algebraic field. *Trans. Amer. Math. Soc.* 34 (1932), 363-372.
- [2] A.A. Albert. *Structure of algebras*. AMS Coll. Pub., vol. 24. American Mathematical Society, New York, 1939.
- [3] F.K. Bingöl. Symbol length in positive characteristic. *J. Pure Appl. Algebra* 228 (2024) 107613.
- [4] K.J. Becher, N. Grenier-Boley, J.-P. Tignol. Transfer of quadratic forms and of quaternion algebras over quadratic field extensions. *Arch. Math. (Basel)* 111 (2018), no. 2, 135–143.
- [5] R. Elman, N. Karpenko, A. Merkurjev. *The algebraic and geometric theory of quadratic forms*. Amer. Math. Soc. Colloq. Publ., 56 American Mathematical Society, Providence, RI, 2008. viii+435 pp.
- [6] R. Elman, T.Y. Lam, J.-P. Tignol, A.R. Wadsworth. Witt rings and Brauer groups under multiquadratic extensions. I. *Amer. J. Math.* 105 (1983), no.5, 1119–1170.
- [7] D.W. Hoffmann. Witt kernels of quadratic forms for multiquadratic extensions in characteristic 2. *Proc. Amer. Math. Soc.* 143 (2015), no.12, 5073–5082.
- [8] N.A. Karpenko. Torsion in CH^2 of Severi-Brauer varieties and indecomposability of generic algebras. *Manuscripta Math.* 88 (1995), no.1, 109–117.
- [9] A. Laghribi, D. Mukhija. On the excellence for inseparable quartic extensions. To appear in *Contemporary Mathematics (AMS)*. Proceedings of the Conference NCRA VIII and QFRC II held in Lens, August 2023. *Preprint version*: <https://univ-artois.hal.science/hal-04518918>
- [10] P. Mammone, R. Moresi. Formes quadratiques, algèbres à division et extensions multiquadratiques inséparables. *Bull. Belg. Math. Soc. Simon Stevin* 2 (1995), no.3, 311–319.
- [11] L.H. Rowen. Division algebras of exponent 2 and characteristic 2. *J. Algebra* 90 (1984), no.1, 71–83.
- [12] A. Schofield, M. Van den Bergh. The index of a Brauer class on a Brauer-Severi variety. *Trans. Amer. Math. Soc.* 333 (1992), no. 2, 729–739.
- [13] A.S. Sivatski. Nonexcellence of multiquadratic field extensions. *J. Algebra* 275 (2004), 859-866.

SCUOLA NORMALE SUPERIORE, PIAZZA DEI CAVALIERI 7, 56126 PISA, ITALIA
Email address: fatmakader.bingol@sns.it

SCHOOL OF COMPUTER SCIENCE, ACADEMIC COLLEGE OF TEL-AVIV-YAFFO, RABENU YERUHAM ST., P.O.B 8401, YAFFO, 6818211, ISRAEL
Email address: adam1chapman@yahoo.com

UNIV. ARTOIS, UR 2462, LABORATOIRE DE MATHÉMATIQUES DE LENS (LML), F-62300 LENS, FRANCE
Email address: ahmed.laghribi@univ-artois.fr