REGULARITY PROPERTIES AND DISPERSIVE BLOW-UP FOR THE FIFTH ORDER KORTEWEG-DE VRIES EQUATION ON THE LINE

EDDYE BUSTAMANTE, JOSÉ JIMÉNEZ URREA AND JORGE MEJÍA

ABSTRACT. In this work we prove that the initial value problem (IVP) for the fifth order Korteweg-de Vries equation

 $\left. \begin{array}{ll} u_t + \partial_x^5 u + u \partial_x u = 0, & x \in \mathbb{R}, \ t > 0, \\ u(x,0) = u_0(x), & \end{array} \right\}$

has a unique local solution in time in the Bourgain spaces $X^{s,b}$ for appropriate values of s and b. Besides we prove a regularity property concerning the nonlinear part of that solution. Finally, using the previous property we establish a dispersive blow-up result for global in time solutions of this IVP.

1. INTRODUCTION

In this article we consider the initial value problem (IVP) corresponding to the fifth order Korteweg-de Vries (KdV) equation

$$u_t + \partial_x^5 u + u \partial_x u = 0, \qquad x \in \mathbb{R}, \ t > 0, \\ u(x,0) = u_0(x).$$

$$(1.1)$$

The well-posedness of the Cauchy problem associated with fifth-order KdV-type equations in the functional context of Sobolev spaces has been addressed by various authors in recent years (see, for example, [1], [5], [9], [13], [15], [16], [17], and references therein).

The IVP (1.1) is a particular case of the IVP, studied by Kato in [10] and [11], for Kawahara equation

$$u_t + \alpha \partial_x^5 u + \beta \partial_x^3 u + \gamma \partial_x (u^2) = 0, \qquad x \in \mathbb{R}, \ t > 0, \\ u(x,0) = u_0(x),$$

$$(1.2)$$

where α , β , and γ are real constants with $\alpha \neq 0, \gamma \neq 0$.

The equation in (1.2) models the capillary waves on a shallow layer and the magneto-sound waves in plasma physics (see [12]).

In [10] Kato proves, in the context of Bourgain spaces, that the IVP (1.2) is locally well-posed (LWP) with initial data in the Sobolev space $H^s(\mathbb{R})$ with $s \ge -2$. Moreover, he proves ill-posedness for s < -2.

In [11], using a modified Bourgain space and applying the I-method, Kato manages to show that the IVP (1.2) is globally well-posed in $H^{s}(\mathbb{R})$ with $s \geq -\frac{38}{21}$.

Since in our work we intend to prove a regularity property of the nonlinear part of the solution of the IVP (1.1), using an extra gain of regularity of the bilinear form $\partial_x(uv)$, measured in an appropriate Bourgain space $X^{s,b}$, then our first result is to establish again the local well-posedness of the IVP (1.1) in $X^{s,b}$, when the initial datum u_0 belongs to $H^s(\mathbb{R})$ with $s \ge 0$.

Let us remember that, formally, u is a solution of the IVP (1.1) if and only if

$$u(t) = W(t)u_0 + \int_0^t W(t - t')(-\frac{1}{2}\partial_x(u(t'))^2)dt',$$
(1.3)

²⁰⁰⁰ Mathematics Subject Classification. 35Q53.

Key words and phrases. Aquí van las palabras clave.

where $\{W(t)\}_{t\in\mathbb{R}}$ is the group of unitary operators in the Sobolev spaces $H^s \equiv H^s(\mathbb{R})$ associated to the linear part of the equation in (1.1); i.e.,

$$W(t)u_0]^{\wedge}(\xi) = e^{-it\xi^5} \widehat{u}_0(\xi).$$
(1.4)

(Here, $u(t) \equiv u(\cdot_x, t)$ and \hat{u}_0 is the Fourier transform of the space function u_0).

We are going to work in the context of the Bourgain spaces $X^{s,b} \equiv X^{s,b}(\mathbb{R}^2)$ of tempered distributions u in \mathbb{R}^2_{xt} such that

$$\|\langle \xi \rangle^s \langle \tau + \xi^5 \rangle^b \widehat{u}(\xi, \tau) \|_{L^2_{\xi\tau}} < \infty, \tag{1.5}$$

where $s \in \mathbb{R}$, and $b = \frac{1}{2}^{-}$; i.e., there exists $\epsilon > 0$ such that $\frac{1}{2} - \epsilon < b < \frac{1}{2}$. Here $\langle \cdot \rangle$ is an abbreviation for $1 + |\cdot|$, \hat{u} denotes the Fourier transform of the distribution u, and ξ and τ are the variables in the frequency space corresponding to x and t, respectively.

We prove that, for suitable values of s and b, and for $u_0 \in H^s(\mathbb{R})$, there exists $u \in X^{s,b}(\mathbb{R}^2)$, continuous from \mathbb{R}_t with values in $H^s(\mathbb{R}_x)$, i.e., $u \in X^{s,b}(\mathbb{R}^2) \cap C(\mathbb{R}_t; H^s(\mathbb{R}_x))$, such that, for some T > 0 and $t \in [0,T]$, equation (1.3) holds. In order to achieve this, we introduce a bump function η of the time variable t such that $\eta(\cdot_t) \in C_0^\infty(\mathbb{R}_t)$, supp $\eta \subset [-1,1]$, and $\eta \equiv 1$ in $[-\frac{1}{2}, \frac{1}{2}]$, and we consider the integral equation

$$u(t) = \eta(t)W(t)u_0 + \eta(t)\int_0^t W(t-t')F_T(u(t'))dt',$$
(1.6)

where $F_T(u(t')) = \eta(\frac{t'}{2T})(-\frac{1}{2}\partial_x(u(t'))^2)$. In a precise manner, we prove the following result.

Theorem 1.1. (Local well posedness of the IVP (1.1)). Let $s \ge 0$, and $u_0 \in H^s(\mathbb{R})$. For $\frac{2}{5} < b < \frac{1}{2}$, there exist $T \in (0, \frac{1}{2}]$, and a unique $u \in X^{s,b} \cap C(\mathbb{R}_t; H^s(\mathbb{R}_x))$ solution of (1.6). (The restriction of u to $\mathbb{R} \times [0,T]$ is a local solution in time of the IVP (1.1)).

It can be seen that the result of Theorem 1.1 is also global in time.

Our second result is related to the extra regularity of the nonlinear part of the solution of the IVP (1.1). This type of result was studied in the case of the IVP associated to the generalized Korteweg-de Vries (KdV) equation

$$u_t + \partial_x^3 u + u^k \partial_x u = 0, \qquad k \in \mathbb{Z}^+, \quad k \ge 2, \quad x \in \mathbb{R}, \ t > 0, \\ u(x,0) = u_0(x),$$
 (1.7)

by Linares and Scialom in [20]. Specifically, if we write the solution u(x,t) of the IVP (1.7) in the form

$$u(x,t) = [V(t)u_0](x) + \left[\int_0^t V(t-t')[-u^k(t')\partial_x u(t')]dt'\right](x) \equiv u_1(x,t) + u_2(x,t),$$

where $\{V(t)\}_{t\in\mathbb{R}}$ is the group associated to the linear part of the equation in (1.7), Linares and Scialom prove that the integral part u_2 of the solution u, is smoother than u, when the initial data u_0 is taken in $H^s(\mathbb{R})$ with $s \ge 1$. Precisely they demonstrate that if $u_0 \in H^s(\mathbb{R})$ with $s \ge 1$, then $u_2 \in C(\mathbb{R}_t; H^{s+1}(\mathbb{R}_x))$ and for all T > 0 $\sup_x \|\partial_x^{s+2}u_2\|_{L^2_T} < +\infty$. This result was obtained, using local smoothing effects of Kato type.

Later, in [19], Linares, Ponce and Smith construct an initial data $u_0 \in H^{\frac{3}{2}^-}(\mathbb{R})$ such that $|x|^{\beta}u_0 \in L^2(\mathbb{R})$ for $\beta \in (0, \frac{3}{4})$ in such a way that for the IVP associated to the Korteweg-de Vries equation (k = 1 in the equation in (1.7)) the nonlinear part of the solution of the IVP, given by

$$u_2(x,t) := \left[\int_0^t V(t-t')(-u(t')\partial_x u(t')dt')\right](x)$$

is such that $u_2 \in C([0,T]; H^{\frac{3}{2}^- + \frac{1}{6}}(\mathbb{R}))$. In other words, $u_2(t)$ is smoother than u_0 for every $t \in [0,T]$. In addition to using regularizing effects, in [19] the authors make use of the local well-posedness of the IVP in weighted Sobolev spaces $H^{\frac{3}{2}-\epsilon}(\mathbb{R}) \cap L^2(|x|^{\frac{3}{4}-2\epsilon}dx)$.

In the case of systems of differential equations, recently, Linares and Palacios in [18] prove that for a Schrödinger-Korteweg-de Vries system the nonlinear part of the solution of the IVP associated to this system is smoother REGULARITY PROPERTIES AND DISPERSIVE BLOW-UP FOR THE FIFTH ORDER KORTEWEG-DE VRIES EQUATION ON THE LINB

than the solution. As in [19], they use smoothing properties, and persistence properties for the solutions of the IVP in weighted Sobolev spaces.

Using Bourgain spaces, in [7], Erdoğan and Tzirakis prove that the nonlinear part of the solution of an initial boundary value problem for the cubic nonlinear Schrödinger equation is smoother than the initial data. This result is a consequence of some gain of regularity of the nonlinear term $|u|^2 u$, measured in the norm of certain Bourgain space. With the help of this approach, in the case of the fifth KdV equation, we manage to prove that the nonlinear part of the solution of the IVP (1.1) is smoother than the initial data u_0 . More precisely, we obtain the following result

Theorem 1.2. (Regularity gain of the nonlinear part of the Solution). Let $s \ge 0$, $u_0 \in H^s(\mathbb{R})$, $\frac{2}{5} < b < \frac{1}{2}$, and $0 \le a \le 10b - 4$. On the other hand, let $T \in (0, \frac{1}{2}]$ and $u \in X^{s,b} \cap C(\mathbb{R}_t; H^s(\mathbb{R}_x))$ be as in Theorem 1.1. Then

$$\left\| \int_{0}^{\cdot_{t}} W(\cdot_{t} - t') (-\frac{1}{2} \partial_{x} (u(t'))^{2}) dt' \right\|_{C([0,T];H^{s+a}(\mathbb{R}))} \le C \|u\|_{X^{s,b}}^{2}.$$
(1.8)

Our main objective in this paper is to establish dispersive blow-up for the solutions of the IVP (1.1). The pioneering work in this direction for dispersive equations is due to Benjamin, Bona, and Mahony in [1]. In that paper, the authors show that there exist smooth initial data u_0 , for which the solution of the linearized KdV equation associated with these initial data can develop point singularities in finite time. Other results related to dispersive blow-up for different nonlinear dispersive models can be found in [2], [4], [21], and the references therein.

In [3], Bona and Saut prove the existence of dispersive blow-up for the generalized KdV equation. The idea behind the Bona and Saut proof is to observe that the nonlinear part of the solution is smoother than the solution, which allows to focus attention on the construction of initial data that gives rise to the formation of point singularities for the solution of the linearized IVP. Following this idea, in [19] and [18] it is possible to establish the dispersive blow-up in the case of the generalized KdV equation and in the case of a Schrödinger-KdV system, respectively.

Finally, using the regularity gain in the Theorem 1.2 and by imitating the procedure followed in [19] (see proof of Theorem 1.4 in [19]) we prove a result on dispersive blow-up, which is stated as follows.

Theorem 1.3. (Dispersive blow-up). There exists $u_0 \in H^1(\mathbb{R}) \cap C^{\infty}(\mathbb{R})$ with $||u_0||_{H^1} \ll 1$ so that the solution $u(\cdot, t)$ of the IVP (1.1) satisfies the following properties:

- (i) $u(\cdot, t) \in C^1(\mathbb{R})$ for $t \in (0, +\infty) \setminus \mathbb{Z}^+$;
- (ii) $u(\cdot, t) \in C^1(\mathbb{R} \setminus \{0\}) \setminus C^1(\mathbb{R})$ for $t \in \mathbb{Z}^+$.

2. Linear and non linear estimates

In this section we establish some estimates for the norms in the spaces $X^{s,b}$ of the terms on the right hand side of the integral equation (1.6). Throughout this section the function $\eta(\cdot_t)$ is as before.

Lemma 2.1. Let $s \in \mathbb{R}$, $b \in \mathbb{R}$. Then there exists C > 0 such that, for every function $u_0 \in H^s(\mathbb{R}_x)$,

$$\|\eta(\cdot_t)[W(\cdot_t)u_0](\cdot_x)\|_{X^{s,b}(\mathbb{R}^2)} \le C \|u_0\|_{H^s(\mathbb{R}_x)}.$$
(2.1)

Proof. Let us observe that

$$(\eta(\cdot_t)[W(\cdot_t)u_0](\cdot_x))^{\wedge}(\xi,\tau) = C \int_{\mathbb{R}_t} e^{-i(\tau+\xi^5)t} \eta(t)\widehat{u}_0(\xi)dt = C\widehat{u}_0(\xi)\widehat{\eta}(\tau+\xi^5).$$

Therefore, taking into account that $\widehat{\eta} \in S(\mathbb{R})$,

$$\|\eta(\cdot_{t})[W(\cdot_{t})u_{0}](\cdot_{x})\|_{X^{s,b}(\mathbb{R}^{2})} = C\left(\iint_{\mathbb{R}^{2}}\langle\xi\rangle^{2s}\langle\tau+\xi^{5}\rangle^{2b}|\widehat{u}_{0}(\xi)|^{2}|\widehat{\eta}(\tau+\xi^{5})|^{2}d\tau d\xi\right)^{\frac{1}{2}} \\ = C\left(\int_{\mathbb{R}_{\tau}}\langle\tau\rangle^{2b}|\widehat{\eta}(\tau)|^{2}\right)^{\frac{1}{2}}\|u_{0}\|_{H^{s}(\mathbb{R}_{x})} = C\|u_{0}\|_{H^{s}(\mathbb{R}_{x})}.$$

Remark 1. For $s, b \in \mathbb{R}$, let us denote by $H^{s,b}$ the anisotropic Sobolev space, defined by

$$H^{s,b} := \{ f(\cdot_x, \cdot_t) \in S'(\mathbb{R}^2) : \|f\|_{H^{s,b}} := \|\langle \xi \rangle^s \langle \tau \rangle^b \widehat{f}(\xi, \tau)\|_{L^2_{\xi\tau}} < \infty \}.$$

It can be seen that

$$\|f(\cdot_x, \cdot_t)\|_{X^{s,b}} = \|W(-\cdot_t)f(\cdot_x, \cdot_t)\|_{H^{s,b}}.$$
(2.2)

Remark 2. In [8], Ginibre, Tsutsumi, and Velo proved that if L is the operator defined by

$$(Lf)(t) := \eta\left(\frac{t}{T}\right) \int_0^t f(t')dt',$$

for fixed $T \in (0,1]$, then, for $-\frac{1}{2} < b' \le 0 \le b \le b' + 1$, it is true that

$$\|Lf\|_{H^{b}(\mathbb{R}_{t})} \leq CT^{1-b+b'} \|f\|_{H^{b'}(\mathbb{R}_{t})},$$
(2.3)

where C > 0 is independent of f.

Lemma 2.2. For $s \in \mathbb{R}$, $-\frac{1}{2} < b' \le 0 \le b \le b' + 1$, $0 < T \le 1$, there exists C > 0, such that for every $f \in X^{s,b'}$, $\left\| \eta\left(\frac{\cdot t}{T}\right) \int^{\cdot t} [W(\cdot_t - t')f(t')](\cdot_x)dt' \right\| \le CT^{1-b+b'} \|f\|_{X^{s,b'}}$ (2.4)

$$\left\| \eta\left(\frac{\cdot}{T}\right) \int_{0} \left[W(\cdot_{t} - t')f(t') \right](\cdot_{x})dt' \right\|_{X^{s,b}} \le CT^{1-b+b'} \|f\|_{X^{s,b'}}$$

$$(2.4)$$

Proof. Using (2.2) and (2.3) we obtain

$$\begin{aligned} \left\| \eta\left(\frac{\cdot}{T}\right) \int_{0}^{\cdot t} \left[W(\cdot_{t}-t')f(t')\right](\cdot_{x})dt' \right\|_{X^{s,b}} \\ &= \left\| W(-\cdot_{t}) \left\{ \eta\left(\frac{\cdot}{T}\right) \int_{0}^{\cdot t} \left[W(\cdot_{t}-t')f(t')\right](\cdot_{x})dt' \right\} \right\|_{H^{s,b}} \\ &= \left(\int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \int_{\mathbb{R}_{\tau}} \langle \tau \rangle^{2b} \left| \left\{ \eta\left(\frac{\cdot}{T}\right) \left(\int_{0}^{\cdot t} \left[W(-t')f(t')\right](\cdot_{x})dt' \right)^{\wedge_{x}} \right\}^{\wedge_{t}} \left(\xi, \tau \right) \right|^{2} d\tau d\xi \right)^{\frac{1}{2}} \\ &= \left(\int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \left\| L\left(\left(\left[W(-\cdot_{t})f(\cdot_{t})\right](\cdot_{x})\right)^{\wedge_{x}} \left(\xi \right) \right) \right\|_{H^{b}(\mathbb{R}_{t})}^{2} d\xi \right)^{\frac{1}{2}} \\ &\leq CT^{1-b+b'} \left(\int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \left\| \left(\left[W(-\cdot_{t})f(\cdot_{t})\right](\cdot_{x})\right)^{\wedge_{x}} \left(\xi \right) \right\|_{H^{b'}(\mathbb{R}_{t})}^{2} d\xi \right)^{\frac{1}{2}}. \end{aligned}$$

$$(2.5)$$

Taking into account that

$$\left\| \left(\left[W(-\cdot_t) f(\cdot_t) \right](\cdot_x) \right)^{\wedge_x}(\xi) \right\|_{H^{b'}(\mathbb{R}_t)}^2 = \int_{\mathbb{R}_\tau} \langle \tau \rangle^{2b'} \left| \left(e^{it\xi^5} \left[f(\cdot_x, t) \right]^{\wedge_x}(\xi) \right)^{\wedge_t}(\tau) \right|^2 d\tau,$$

and since

$$\left(e^{it\xi^5}[f(\cdot_x,t)]^{\wedge_x}(\xi)\right)^{\wedge_t}(\tau) = C \int_{\mathbb{R}_t} e^{-it(\tau-\xi^5)} \int_{\mathbb{R}_x} e^{-i\xi x} f(x,t) dx dt$$

we conclude that

$$\left\| \left([W(-\cdot_t)f(\cdot_t)](\cdot_x) \right)^{\wedge_x}(\xi) \right\|_{H^{b'}(\mathbb{R}_t)}^2 = C \int_{\mathbb{R}_\tau} \langle \tau \rangle^{2b'} |\widehat{f}(\xi,\tau-\xi^5)|^2 d\tau = C \int_{\mathbb{R}_\tau} \langle \tau+\xi^5 \rangle^{2b'} |\widehat{f}(\xi,\tau)|^2 d\tau.$$

From (2.5), we obtain

$$\left\| \eta\left(\frac{d}{T}\right) \int_{0}^{t} [W(t_{t} - t')f(t')](t_{x})dt' \right\|_{X^{s,b}} \leq CT^{1-b+b'} \|f\|_{X^{s,b'}}.$$

Lemma 2.3. Let $s \in \mathbb{R}$, b_1, b_2 such that $-\frac{1}{2} < b_1 < b_2 < \frac{1}{2}$, and $T \in (0, 1]$. Then there exists C > 0 such that, for every $F \in X^{s, b_2}$,

$$\left\|\eta\left(\frac{\cdot}{T}\right)F(\cdot_x,\cdot_t)\right\|_{X^{s,b_1}} \le CT^{b_2-b_1}\|F\|_{X^{s,b_2}}.$$
(2.6)

Proof. Let us prove (2.6) for $b_1 = 0$; i.e.,

$$\left\|\eta\left(\frac{\cdot_t}{T}\right)F(\cdot_x,\cdot_t)\right\|_{X^{s,0}} \le CT^{b_2}\|F\|_{X^{s,b_2}}.$$
(2.7)

The general case follows from (2.7), using an interpolation argument. Estimative (2.7) is consequence of the following statements:

(i) $\|\eta\left(\frac{\cdot}{T}\right)F(\cdot_{x},\cdot_{t})\|_{X^{s,0}} \leq CT^{b_{2}} \|\eta\left(\frac{\cdot}{T}\right)F(\cdot_{x},\cdot_{t})\|_{X^{s,b_{2}}},$ (ii) $\|\eta\left(\frac{\cdot}{T}\right)F(\cdot_{x},\cdot_{t})\|_{X^{s,b_{2}}} \leq C\|F\|_{X^{s,b_{2}}};$

where C > 0 is independent of T.

Let us prove (i):

$$\begin{split} \left\| \eta\left(\frac{\cdot}{T}\right)F(\cdot_{x},\cdot_{t})\right\|_{X^{s,0}}^{2} &= \int_{\mathbb{R}}\int_{-\infty}^{+\infty}\langle\xi\rangle^{2s} \left| \left[\eta\left(\frac{\cdot}{T}\right)F\right]^{\wedge}(\xi,\tau)\right|^{2}d\tau d\xi \\ &= \int_{\mathbb{R}}\langle\xi\rangle^{2s}\int_{-\infty}^{+\infty} \left|\eta\left(\frac{t}{T}\right)\left[F(\cdot,t)\right]^{\wedge_{x}}(\xi)\right|^{2}dtd\xi \\ &= \int_{\mathbb{R}_{t}}\left|\eta\left(\frac{t}{T}\right)\right|^{2}\int_{\mathbb{R}_{\xi}}\left|e^{it\xi^{5}}\langle\xi\rangle^{s}\left[F(\cdot,t)\right]^{\wedge_{x}}(\xi)\right|^{2}d\xi dt \\ &= C\int_{-T}^{T}\left|\eta\left(\frac{t}{T}\right)\right|^{2}\int_{\mathbb{R}_{x}}\left|\left[W(-t)J^{s}(F(\cdot,t))\right](x)\right|^{2}dxdt \end{split}$$

where

$$[J^s(F(\cdot,t))]^{\wedge_x}(\xi) := \langle \xi \rangle^s [F(\cdot,t)]^{\wedge_x}(\xi).$$

Hence

$$\left\|\eta\left(\frac{\cdot_{t}}{T}\right)F(\cdot_{x},\cdot_{t})\right\|_{X^{s,0}}^{2} = C\int_{-T}^{T}\int_{\mathbb{R}_{x}}\left|\left[W(-t)J^{s}(\eta\left(\frac{t}{T}\right)F(\cdot,t))\right](x)\right|^{2}dxdt.$$

Since $0 < 2b_2 < 1$, $\frac{1}{2b_2} > 1$. Defining $p := \frac{1}{2b_2}$, and q as the conjugate exponent of p; i.e., $q := \frac{1}{1-2b_2}$, by Hölder's inequality in the t variable, we obtain

$$\begin{split} \left\| \eta\left(\frac{\cdot}{T}\right)F(\cdot_{x},\cdot_{t})\right\|_{X^{s,0}}^{2} &\leq CT^{2b_{2}} \left\{ \int_{-\infty}^{+\infty} \left(\int_{\mathbb{R}} \left| \left[W(-t)J^{s}(\eta\left(\frac{t}{T}\right)F(\cdot,t))\right](x)\right|^{2}dx \right)^{\frac{1}{1-2b_{2}}}dt \right\}^{1-2b_{2}} \\ &\leq CT^{2b_{2}} \int_{\mathbb{R}} \left(\int_{-\infty}^{+\infty} \left| \left[W(-t)J^{s}(\eta\left(\frac{t}{T}\right)F(\cdot,t))\right](x)\right|^{\frac{2}{1-2b_{2}}}dt \right)^{1-2b_{2}}dx \\ &= CT^{2b_{2}} \int_{\mathbb{R}} \left\| \left[W(-\cdot_{t})J^{s}(\eta\left(\frac{\cdot}{T}\right)F(\cdot,\cdot_{t}))\right](x)\right\|_{L^{\frac{2}{1-2b_{2}}}(\mathbb{R}_{t})}^{2}dt. \end{split}$$

Taking into account that $H^{b_2}(\mathbb{R}) \hookrightarrow L^{\frac{2}{1-2b_2}}(\mathbb{R})$ for $0 \leq b_2 < \frac{1}{2}$, it follows that

$$\begin{split} \left\| \eta \left(\frac{\cdot t}{T} \right) F(\cdot_x, \cdot_t) \right\|_{X^{s,0}}^2 &\leq CT^{2b_2} \int_{\mathbb{R}} \int_{-\infty}^{+\infty} \langle \tau \rangle^{2b_2} \left| \left([W(-\cdot_t) J^s(\eta \left(\frac{\cdot t}{T} \right) F(\cdot, \cdot_t))](x) \right)^{\wedge_t}(\tau) \right|^2 d\tau dx \\ &= CT^{2b_2} \int_{-\infty}^{+\infty} \langle \tau \rangle^{2b_2} \int_{\mathbb{R}_{\xi}} \left| \left\{ \left([W(-\cdot_t) J^s(\eta \left(\frac{\cdot t}{T} \right) F(\cdot, \cdot_t))](\cdot_x) \right)^{\wedge_t}(\tau) \right\}^{\wedge_x}(\xi) \right|^2 d\xi d\tau \\ &= CT^{2b_2} \int_{-\infty}^{+\infty} \langle \tau \rangle^{2b_2} \int_{\mathbb{R}_{\xi}} \left| \left(e^{it\xi^5} \langle \xi \rangle^s [\eta \left(\frac{\cdot t}{T} \right) F(\cdot_x, \cdot_t)]^{\wedge_x}(\xi) \right)^{\wedge_t}(\tau) \right|^2 d\xi d\tau \\ &= CT^{2b_2} \int_{-\infty}^{+\infty} \langle \tau \rangle^{2b_2} \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \left| [\eta \left(\frac{\cdot t}{T} \right) F(\cdot_x, \cdot_t)]^{\wedge_{xt}}(\xi, \tau - \xi^5) \right|^2 d\xi d\tau \\ &= CT^{2b_2} \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \int_{-\infty}^{+\infty} \langle \tau + \xi^5 \rangle^{2b_2} \left| [\eta \left(\frac{\cdot t}{T} \right) F(\cdot_x, \cdot_t)]^{\wedge_{xt}}(\xi, \tau) \right|^2 d\tau d\xi \\ &= CT^{2b_2} \left\| \eta \left(\frac{\cdot t}{T} \right) F(\cdot_x, \cdot_t) \right\|_{X^{s,b_2}}^2, \end{split}$$

and estimative (i) is proved.

Now we prove (ii):

$$\left\|\eta\left(\frac{\cdot_{t}}{T}\right)F(\cdot_{x},\cdot_{t})\right\|_{X^{s,b_{2}}}^{2} = \int_{\mathbb{R}}\langle\xi\rangle^{2s}\int_{-\infty}^{+\infty}\langle\tau+\xi^{5}\rangle^{2b_{2}}\left|\left[\eta\left(\frac{\cdot_{t}}{T}\right)F(\cdot_{x},\cdot_{t})\right]^{\wedge}(\xi,\tau)\right|^{2}d\tau d\xi$$

Let us observe that

$$\left[\eta\left(\frac{\cdot_t}{T}\right)F(\cdot_x,\cdot_t)\right]^{\wedge}(\xi,\tau) = C\left[\eta\left(\frac{\cdot_t}{T}\right)(F(\cdot_x,t))^{\wedge_x}(\xi)\right]^{\wedge_t}(\tau) = C\left\{\left[\eta\left(\frac{\cdot_t}{T}\right)\right]^{\wedge_t}*_{\tau}\widehat{F}(\xi,\cdot_{\tau})\right\}(\tau)$$

But,

$$\left[\eta\left(\frac{\cdot t}{T}\right)\right]^{\wedge_t}(\tau) = T\widehat{\eta}(T\tau).$$

Hence

$$\left\|\eta\left(\frac{\cdot t}{T}\right)F(\cdot_{x},\cdot_{t})\right\|_{X^{s,b_{2}}}^{2} = C \int_{\mathbb{R}_{\xi}} \langle\xi\rangle^{2s} \int_{-\infty}^{+\infty} \langle\tau+\xi^{5}\rangle^{2b_{2}} \left|\left[(T\widehat{\eta}(T\cdot)*_{\tau}\widehat{F}(\xi,\cdot)\right](\tau)\right|^{2} d\tau d\xi.$$
(2.8)

Let us estimate the previous integral in the τ variable:

$$\int_{-\infty}^{+\infty} \langle \tau + \xi^5 \rangle^{2b_2} \left| \left[(T\widehat{\eta}(T\cdot) *_{\tau} \widehat{F}(\xi, \cdot)](\tau) \right|^2 d\tau \right]$$

$$\leq C \left\| T\widehat{\eta}(T\cdot) *_{\tau} \widehat{F}(\xi, \cdot) \right\|_{L^2_{\tau}}^2 + C \int_{-\infty}^{+\infty} |\tau|^{2b_2} \left| \left[(T\widehat{\eta}(T\cdot)) *_{\tau} \widehat{F}(\xi, \cdot)](\tau - \xi^5) \right|^2 d\tau \equiv I(\xi) + II(\xi).$$
(2.9)

On the one hand

$$I(\xi) \le C \|T\widehat{\eta}(T \cdot_{\tau})\|_{L^{1}_{\tau}}^{2} \|\widehat{F}(\xi, \cdot_{\tau})\|_{L^{2}_{\tau}}^{2} \le C \|\widehat{F}(\xi, \cdot_{\tau})\|_{L^{2}_{\tau}}^{2},$$
(2.10)

with C independent of T.

||.

On the other hand, to estimate $II(\xi)$, we use Leibniz formula for fractional derivatives (see [14], Theorem A.12):

$$D^{\alpha}(fg) - fD^{\alpha}g\|_{L^{p}(\mathbb{R})} \le C\|g\|_{L^{\infty}(\mathbb{R})}\|D^{\alpha}f\|_{L^{p}(\mathbb{R})} \quad \alpha \in (0,1), \quad 1 (2.11)$$

Let us observe that

$$II(\xi) = C \left\| D_t^{b_2} \left(\mathcal{F}_t^{-1}([T\hat{\eta}(T\cdot) *_{\tau} \hat{F}(\xi, \cdot)](\tau - \xi^5)) \right) \right\|_{L^2_t}^2.$$
(2.12)

But

$$\begin{aligned} \mathcal{F}_t^{-1}[(T\widehat{\eta}(T\cdot)*_\tau\widehat{F}(\xi,\cdot))(\tau-\xi^5)](t) &= C\int_{-\infty}^{+\infty} e^{it\tau} e^{it\xi^5}[T\widehat{\eta}(T\cdot)*_\tau\widehat{F}(\xi,\cdot)](\tau)d\tau\\ &= Ce^{it\xi^5}\mathcal{F}_t^{-1}[T\widehat{\eta}(T\cdot)](t)[F(\cdot_x,t)]^{\wedge_x}(\xi)\\ &= Ce^{it\xi^5}\eta\left(\frac{t}{T}\right)[F(\cdot_x,t)]^{\wedge_x}(\xi). \end{aligned}$$

Then, from (2.12), using (2.11), we have that

$$II(\xi) = C \left\| D_{t}^{b_{2}} \left(e^{i \cdot t \xi^{5}} [F(\cdot_{x}, \cdot_{t})]^{\wedge_{x}}(\xi) \eta\left(\frac{\cdot}{T}\right) \right) \right\|_{L_{t}^{2}}^{2} \\ \leq C \left\| D_{t}^{b_{2}} \left(e^{i \cdot t \xi^{5}} [F(\cdot_{x}, \cdot_{t})]^{\wedge_{x}}(\xi) \eta\left(\frac{\cdot}{T}\right) \right) - e^{i \cdot t \xi^{5}} [F(\cdot_{x}, \cdot_{t})]^{\wedge_{x}}(\xi) D_{t}^{b_{2}} \left(\eta\left(\frac{\cdot}{T}\right) \right) \right\|_{L_{t}^{2}}^{2} \\ + C \left\| e^{i \cdot t \xi^{5}} [F(\cdot_{x}, \cdot_{t})]^{\wedge_{x}}(\xi) D_{t}^{b_{2}} \left(\eta\left(\frac{\cdot}{T}\right) \right) \right\|_{L_{t}^{2}}^{2} \\ \leq C \left\| \eta\left(\frac{\cdot}{T}\right) \right\|_{L_{t}^{\infty}}^{2} \left\| D_{t}^{b_{2}} \left(e^{i \cdot t \xi^{5}} [F(\cdot_{x}, \cdot_{t})]^{\wedge_{x}}(\xi) \right) \right\|_{L_{t}^{2}}^{2} + C \left\| e^{i \cdot t \xi^{5}} [F(\cdot_{x}, \cdot_{t})]^{\wedge_{x}}(\xi) D_{t}^{b_{2}} \left(\eta\left(\frac{\cdot}{T}\right) \right) \right\|_{L_{t}^{2}}^{2} \\ \leq C \left\| D_{t}^{b_{2}} \left(e^{i \cdot t \xi^{5}} [F(\cdot_{x}, \cdot_{t})]^{\wedge_{x}}(\xi) \right) \right\|_{L_{t}^{2}}^{2} + C \left\| e^{i \cdot t \xi^{5}} [F(\cdot_{x}, \cdot_{t})]^{\wedge_{x}}(\xi) D_{t}^{b_{2}} \left(\eta\left(\frac{\cdot}{T}\right) \right) \right\|_{L_{t}^{2}}^{2}. \tag{2.13}$$

From (2.10) it is clear that

$$\int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} I(\xi) d\xi \le C \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \| \widehat{F}(\xi, \cdot_{\tau}) \|_{L^{2}_{\tau}}^{2} d\xi = C \| F \|_{X^{s,0}}^{2}.$$
(2.14)

Besides, from (2.13), using Plancherel's identity, we have that

$$\begin{split} \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} II(\xi) d\xi &\leq C \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \int_{-\infty}^{+\infty} |\tau|^{2b_2} \left| \left(e^{it\xi^5} [F(\cdot_x, t)]^{\wedge_x}(\xi) \right)^{\wedge_t} (\tau) \right|^2 d\tau d\xi \\ &+ C \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \int_{-\infty}^{+\infty} \left| e^{it\xi^5} [F(\cdot_x, t)]^{\wedge_x}(\xi) D_t^{b_2} \left(\eta \left(\frac{t}{T} \right) \right) \right|^2 dt d\xi \\ &\leq C \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \int_{-\infty}^{+\infty} |\tau + \xi^5|^{2b_2} |\widehat{F}(\xi, \tau)|^2 d\tau d\xi \\ &+ C \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \int_{-\infty}^{+\infty} \left| e^{it\xi^5} [F(\cdot_x, t)]^{\wedge_x}(\xi) D_t^{b_2} \left(\eta \left(\frac{t}{T} \right) \right) \right|^2 dt d\xi. \end{split}$$

Using Hölder's inequality in the integral in the t variable with conjugate exponents p and p', we have that

$$\int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} II(\xi) d\xi \le C \|F\|_{X^{s,b_2}}^2 + C \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \left\{ \int_{-\infty}^{+\infty} \left| e^{it\xi^5} [F(\cdot_x, t)]^{\wedge_x}(\xi) \right|^{2p} dt \right\}^{\frac{1}{p}} \left\{ \int_{-\infty}^{+\infty} |D_t^{b_2}\left(\eta\left(\frac{t}{T}\right)\right)|^{2p'} dt \right\}^{\frac{1}{p'}} d\xi.$$

Let us choose p such that $\frac{1}{2} - \frac{1}{2p} = b_2$; i.e., $\frac{1}{2p} = \frac{1-2b_2}{2}$, which is equivalent to $2p = \frac{2}{1-2b_2}$, which implies $H^{b_2}(\mathbb{R}_t) \hookrightarrow L^{2p}(\mathbb{R}_t)$. Then

$$\begin{split} \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} II(\xi) d\xi &\leq C \|F\|_{X^{s,b_{2}}}^{2} + C \int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} \left\| e^{it\xi^{5}} [F(\cdot_{x},t)]^{\wedge_{x}}(\xi) \right\|_{H^{b_{2}}(\mathbb{R}_{t})}^{2} d\xi \left\{ \int_{-\infty}^{+\infty} |D_{t}^{b_{2}}\left(\eta\left(\frac{t}{T}\right)\right)|^{2p'} dt \right\}^{\frac{1}{p'}} \\ &\leq C \|F\|_{X^{s,b_{2}}}^{2} + C \|F\|_{X^{s,b_{2}}}^{2} \left\{ \int_{-\infty}^{+\infty} |D_{t}^{b_{2}}\left(\eta\left(\frac{t}{T}\right)\right)|^{2p'} dt \right\}^{\frac{1}{p'}}. \end{split}$$

Taking into account that the inverse Fourier transform is a bounded operator from $L^{\frac{2p'}{2p'-1}}$ to $L^{2p'}$, it follows that

$$\int_{\mathbb{R}_{\xi}} \langle \xi \rangle^{2s} II(\xi) d\xi \leq C \|F\|_{X^{s,b_{2}}}^{2} + C \|F\|_{X^{s,b_{2}}}^{2} \left\{ \int_{-\infty}^{+\infty} \left| |\tau|^{b_{2}} T \widehat{\eta}(T\tau) \right|^{\frac{2p'}{2p'-1}} d\tau \right\}^{\frac{2p'-1}{p'}} \leq C \|F\|_{X^{s,b_{2}}}^{2}.$$
(2.15)

From (2.8), (2.9), (2.14), and (2.15) we conclude that

$$\left\|\eta\left(\frac{\cdot t}{T}\right)F(\cdot_x,\cdot_t)\right\|_{X^{s,b_2}} \le C\|F\|_{X^{s,b_2}} \quad T \in (0,1],$$

with C independent of T.

In consequence of (i), and (ii), we conclude that

$$\left\|\eta\left(\frac{\cdot_t}{T}\right)F(\cdot_x,\cdot_t)\right\|_{X^{s,0}} \le CT^{b_2}\|F\|_{X^{s,b_2}}.$$

The following calculus inequalities will be used the proof of the estimative of the bilinear form (Lemma 2.5): **Lemma 2.4.** (i) For $\beta \ge \gamma \ge 0$, and $\beta + \gamma > 1$ it follows that

$$\int_{\mathbb{R}} \frac{1}{\langle x - a_1 \rangle^{\beta} \langle x - a_2 \rangle^{\gamma}} dx \le C \frac{\phi_{\beta}(a_1 - a_2)}{\langle a_1 - a_2 \rangle^{\gamma}},\tag{2.16}$$

where

$$\phi_{\beta}(a) \sim \begin{cases} 1 & \text{for } \beta > 1, \\ \log(1 + \langle a \rangle) & \text{for } \beta = 1, \\ \langle a \rangle^{1-\beta} & \text{for } \beta < 1. \end{cases}$$

(For a proof of (2.16), see [6]).

(ii) For $\rho \in (\frac{1}{2}, 1)$,

$$\int_{\mathbb{R}} \frac{1}{\langle x \rangle^{\rho} \sqrt{|x-a|}} \le C \frac{1}{\langle a \rangle^{\rho-\frac{1}{2}}}.$$
(2.17)

(For a proof of (2.17) see [7]).

Lemma 2.5. (Estimative of the bilinear form). Let $s \ge 0$, $\frac{2}{5} \le b < \frac{1}{2}$, and $0 \le a \le 10b-4$. There exists C > 0 such that

$$\|\partial_x(vw)\|_{X^{s+a,-b}} \le C \|v\|_{X^{s,b}} \|w\|_{X^{s,b}}, \qquad (2.18)$$

Proof. We observe that

$$[\partial_x(vw)]^{\wedge}(\xi,\tau) = Ci\xi \int_{\mathbb{R}_2} \widehat{v}(\xi_1,\tau_1)\widehat{w}(\xi-\xi_1,\tau-\tau_1)d\xi_1d\tau_1.$$

Hence

$$\begin{aligned} \|\partial_{x}(vw)\|_{X^{s+a,-b}}^{2} &= C \int_{\mathbb{R}^{2}_{\xi\tau}} \langle\xi\rangle^{2(s+a)} \langle\tau + \xi^{5}\rangle^{-2b} \xi^{2} \left| \int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \widehat{v}(\xi_{1},\tau_{1}) \widehat{w}(\xi - \xi_{1},\tau - \tau_{1}) d\xi_{1} d\tau_{1} \right|^{2} d\xi d\tau \\ &= C \int_{\mathbb{R}^{2}_{\xi\tau}} \left| \int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \langle\xi\rangle^{(s+a)} \langle\tau + \xi^{5}\rangle^{-b} |\xi| \widehat{v}(\xi_{1},\tau_{1}) \widehat{w}(\xi - \xi_{1},\tau - \tau_{1}) d\xi_{1} d\tau_{1} \right|^{2} d\xi d\tau. \end{aligned}$$
(2.19)

Let $h \in L^2(\mathbb{R}^2_{\xi\tau})$ an arbitrary function. If we manage to prove that

$$\left| \int_{\mathbb{R}^{2}_{\xi\tau}} \left[\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \langle \xi \rangle^{(s+a)} \langle \tau + \xi^{5} \rangle^{-b} |\xi| \widehat{v}(\xi_{1},\tau_{1}) \widehat{w}(\xi - \xi_{1},\tau - \tau_{1}) d\xi_{1} d\tau_{1} \right] h(\xi,\tau) d\xi d\tau \right| \leq C \|v\|_{X^{s,b}} \|w\|_{X^{s,b}} \|h\|_{L^{2}(\mathbb{R}^{2})},$$

$$(2.20)$$

then we would have, by a duality argument, that

$$\|\partial_x(vw)\|_{X^{s+a,-b}} \le C \|v\|_{X^{s,b}} \|w\|_{X^{s,b}}.$$
(2.21)

Taking into account that, there exists C > 0, such that for $s \ge 0$

$$\frac{\langle \xi \rangle^s}{\langle \xi_1 \rangle^s \langle \xi - \xi_1 \rangle^s} \le C,$$

then, to establish (2.20), it is enough to prove that

$$\left| \int_{\mathbb{R}^{2}_{\xi\tau}} \left[\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \langle \xi \rangle^{a} \langle \xi_{1} \rangle^{s} \langle \tau + \xi^{5} \rangle^{-b} |\xi| \langle \xi - \xi_{1} \rangle^{s} |\widehat{v}(\xi_{1}, \tau_{1})| |\widehat{w}(\xi - \xi_{1}, \tau - \tau_{1})| d\xi_{1} d\tau_{1} \right] h(\xi, \tau) d\xi d\tau \right| \leq C \|v\|_{X^{s,b}} \|w\|_{X^{s,b}} \|h\|_{L^{2}(\mathbb{R}^{2})}. \tag{2.22}$$

Since

$$\begin{split} & \left| \int_{\mathbb{R}^{2}_{\xi\tau}} \left[\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \langle \xi \rangle^{a} \langle \xi_{1} \rangle^{s} \langle \tau + \xi^{5} \rangle^{-b} |\xi| \langle \xi - \xi_{1} \rangle^{s} |\widehat{v}(\xi_{1},\tau_{1})| |\widehat{w}(\xi - \xi_{1},\tau - \tau_{1})| d\xi_{1} d\tau_{1} \right] h(\xi,\tau) d\xi d\tau \right| \\ & \leq \int_{\mathbb{R}^{2}_{\xi\tau}} \int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle \xi \rangle^{a} |\xi|| h(\xi,\tau)| \langle \xi_{1} \rangle^{s} \langle \tau_{1} + \xi^{5}_{1} \rangle^{b} |\widehat{v}(\xi_{1},\tau_{1})| \langle \xi - \xi_{1} \rangle^{s} \langle \tau - \tau_{1} + (\xi - \xi_{1})^{5} \rangle^{b} |\widehat{w}(\xi - \xi_{1},\tau - \tau_{1})| d\xi_{1} d\tau_{1} d\xi d\tau \\ & \leq \int_{\mathbb{R}^{2}_{\xi\tau}} \left[\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle \xi \rangle^{2a} |\xi|^{2} |h(\xi,\tau)|^{2}}{\langle \tau + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5}_{1} \rangle^{2b} \langle \tau - \tau_{1} + (\xi - \xi_{1})^{5} \rangle^{2b}} d\xi_{1} d\tau_{1} \right]^{\frac{1}{2}} \\ & = \int_{\mathbb{R}^{2}_{\xi\tau}} \int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle \xi \rangle^{2a} |\xi|^{2} |h(\xi,\tau)|^{2}}{\langle \tau + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5}_{1} \rangle^{2b} \langle \tau - \tau_{1} + (\xi - \xi_{1})^{5} \rangle^{2b}} d\xi_{1} d\tau_{1} d\xi d\tau \\ & \leq \left[\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle \xi \rangle^{2a} |\xi|^{2} |h(\xi,\tau)|^{2}}{\langle \tau + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5}_{1} \rangle^{2b} \langle \tau - \tau_{1} + (\xi - \xi_{1})^{5} \rangle^{2b}} d\xi_{1} d\tau_{1} d\xi d\tau \right]^{\frac{1}{2}} \\ & = \left[\int_{\mathbb{R}^{2}_{\xi\tau}} \int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle \xi \rangle^{2a} |\xi|^{2} |h(\xi,\tau)|^{2}}{\langle \tau + \xi^{5} \rangle^{2b} \langle \tau - \tau_{1} + (\xi - \xi_{1})^{5} \rangle^{2b}} d\xi_{1} d\tau_{1} d\xi d\tau \right]^{\frac{1}{2}} \\ & = \left[\int_{\mathbb{R}^{2}_{\xi\tau}} \int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle \xi \rangle^{2a} |\xi|^{2} |h(\xi,\tau)|^{2}}{\langle \tau + \xi^{5} \rangle^{2b} \langle \tau - \tau_{1} + \xi^{5} \rangle^{2b} \langle \tau - \tau_{1} + (\xi - \xi_{1})^{5} \rangle^{2b}} d\xi_{1} d\tau_{1} d\xi d\tau \right]^{\frac{1}{2}} \\ & = \left[\int_{\mathbb{R}^{2}_{\xi\tau}} \int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle \xi \rangle^{2a} |\xi|^{2} |h(\xi,\tau)|^{2} \langle \xi - \xi_{1} \rangle^{2a} \langle \tau - \tau_{1} + (\xi - \xi_{1})^{5} \rangle^{2b}} d\xi_{1} d\tau_{1} \right) d\xi d\tau \right]^{\frac{1}{2}} \\ & = \left[\int_{\mathbb{R}^{2}_{\xi\tau}} |h(\xi,\tau)|^{2} \left(\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle \xi \rangle^{2a} |\xi|^{2} |\xi|^{2} \langle \tau + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5} \rangle^{2b} \langle \tau_{1} - \xi^{5} \rangle^{2a} |\xi|^{2}} d\xi d\tau \\ & = \left[\int_{\mathbb{R}^{2}_{\xi\tau}} |h(\xi,\tau)|^{2} \left(\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle \xi \rangle^{2a} |\xi|^{2} |\xi|^{2} \langle \tau_{1} + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5} \rangle^{2b} \langle \tau_{1} - \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5} \rangle^{2b} \langle \tau_{1} +$$

then, to prove (2.22), it is enough to prove that

$$\sup_{(\xi,\tau)\in\mathbb{R}^2} \int_{\mathbb{R}^2_{\xi_1\tau_1}} \frac{\langle\xi\rangle^{2a}|\xi|^2}{\langle\tau+\xi^5\rangle^{2b}\langle\tau_1+\xi^5_1\rangle^{2b}\langle\tau-\tau_1+(\xi-\xi_1)^5\rangle^{2b}} d\xi_1 d\tau_1 \le C.$$
(2.23)

Let us observe that

$$\int_{\mathbb{R}_{\tau_1}} \frac{\langle \xi \rangle^{2a} |\xi|^2}{\langle \tau + \xi^5 \rangle^{2b} \langle \tau_1 + \xi_1^5 \rangle^{2b} \langle \tau - \tau_1 + (\xi - \xi_1)^5 \rangle^{2b}} d\tau_1 = \frac{\langle \xi \rangle^{2a} |\xi|^2}{\langle \tau + \xi^5 \rangle^{2b}} \int_{\mathbb{R}_{\tau_1}} \frac{1}{\langle \tau_1 - (-\xi_1^5) \rangle^{2b} \langle \tau_1 - (\tau + (\xi - \xi_1)^5) \rangle^{2b}} d\tau_1.$$

Using inequality (2.16), with $\beta = \gamma = 2b < 1$, and $\beta + \gamma = 4b > 1$ $(b > \frac{1}{4})$, we conclude that

$$\int_{\mathbb{R}_{\xi_1}} \int_{\mathbb{R}_{\tau_1}} \frac{\langle \xi \rangle^{2a} |\xi|^2}{\langle \tau + \xi^5 \rangle^{2b} \langle \tau_1 + \xi_1^5 \rangle^{2b} \langle \tau - \tau_1 + (\xi - \xi_1)^5 \rangle^{2b}} d\tau_1 d\xi_1 \le C \int_{\mathbb{R}_{\xi_1}} \frac{\langle \xi \rangle^{2a} |\xi|^2}{\langle \tau + \xi^5 \rangle^{2b} \langle \xi_1^5 + \tau + (\xi - \xi_1)^5 \rangle^{4b-1}} d\xi_1.$$
(2.24)

Let us make the following change of variables in the integral of the right hand side of (2.24):

$$\mu := \xi_1^5 + \tau + (\xi - \xi_1)^5 = \tau + \xi^5 - \frac{5}{2}\xi\xi_1(\xi - \xi_1)[\xi^2 + \xi_1^2 + (\xi - \xi_1)^2]$$

= $\tau + \frac{1}{16}\xi^5 + \frac{5}{2}\xi^3(\xi_1 - \frac{\xi}{2})^2 + 5\xi(\xi_1 - \frac{\xi}{2})^4,$ (2.25)

$$d\mu = -\frac{5}{2}\xi(\xi - 2\xi_1)[\xi^2 + (\xi - 2\xi_1)^2]d\xi_1.$$
(2.26)

Therefore

$$\int_{\mathbb{R}_{\xi_{1}}} \int_{\mathbb{R}_{\tau_{1}}} \frac{\langle \xi \rangle^{2a} |\xi|^{2}}{\langle \tau + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5}_{1} \rangle^{2b} \langle \tau - \tau_{1} + (\xi - \xi_{1})^{5} \rangle^{2b}} d\tau_{1} d\xi_{1} \\
\leq C \frac{\langle \xi \rangle^{2a} |\xi|^{2}}{\langle \tau + \xi^{5} \rangle^{2b}} \int_{A \subset \mathbb{R}_{\mu}} \frac{1}{(1 + |\mu|)^{4b - 1} |\xi| |\xi - 2\xi_{1}| [\xi^{2} + (\xi - 2\xi_{1})^{2}]} d\mu,$$
(2.27)

where $A = [\tau + \frac{1}{16}\xi^5, +\infty)$ when $\xi > 0$.

Using the last equality of (2.25), it can be seen that

$$\left(\xi_1 - \frac{\xi}{2}\right)^2 = -\frac{\xi^2}{4} + \sqrt{\frac{\xi^5 - 4\tau + 4\mu}{20\xi}},$$

then

$$(\xi - 2\xi_1)^2 = -\xi^2 + \sqrt{\frac{4}{5\xi}(\xi^5 - 4\tau + 4\mu)},$$

$$\xi_1 = \frac{\xi}{2} \pm \sqrt{\frac{1}{2}\sqrt{\frac{1}{5\xi}(\xi^5 - 4\tau + 4\mu)} - \frac{1}{4}\xi^2},$$

$$2\xi_1 - \xi = \pm \sqrt{-\xi^2 + \sqrt{\frac{4}{5\xi}(\xi^5 - 4\tau + 4\mu)}}.$$

Therefore, from (2.27), we have that

$$\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle\xi\rangle^{2a}|\xi|^{2}}{\langle\tau+\xi^{5}\rangle^{2b}\langle\tau_{1}+\xi^{5}_{1}\rangle^{2b}\langle\tau-\tau_{1}+(\xi-\xi_{1})^{5}\rangle^{2b}} d\tau_{1}d\xi_{1} \\
\leq C \frac{\langle\xi\rangle^{2a}|\xi|}{\langle\tau+\xi^{5}\rangle^{2b}} \int_{A\subset\mathbb{R}_{\mu}} \frac{1}{(1+|\mu|)^{4b-1}\sqrt{-\xi^{2}+\sqrt{\frac{4}{5\xi}}(\xi^{5}-4\tau+4\mu)}} \sqrt{\frac{4}{5\xi}}(\xi^{5}-4\tau+4\mu)} d\mu \\
\leq C \frac{\langle\xi\rangle^{2a}|\xi|}{\langle\tau+\xi^{5}\rangle^{2b}} \int_{A\subset\mathbb{R}_{\mu}} \frac{\sqrt{\xi^{2}+\sqrt{\frac{4}{5\xi}}(\xi^{5}-4\tau+4\mu)}}{(1+|\mu|)^{4b-1}\sqrt{\frac{-16\tau+16\mu-\xi^{5}}{5\xi}}\sqrt{\frac{4}{5\xi}}(\xi^{5}-4\tau+4\mu)}} d\mu. \tag{2.28}$$

Let us assume, without loss of generality, that $\xi > 0$, and let us observe that the function

$$\mu := \varphi(\xi_1) = \tau + \xi_1^5 + (\xi - \xi_1)^5$$

has its absolute minimum in $\xi_1 = \frac{\xi}{2}$. Then, the integration interval A in (2.28) is

$$A = [\varphi(\frac{\xi}{2}), +\infty) = [\tau + \frac{1}{16}\xi^5, +\infty)$$

This way

$$\mu \ge \tau + \frac{1}{16}\xi^5, \quad \xi^5 - 4\tau + 4\mu \ge \xi^5 - 4\tau + 4\tau + \frac{1}{4}\xi^5 = \frac{5}{4}\xi^5,$$

$$\frac{4}{5\xi}(\xi^5 - 4\tau + 4\mu) \ge \frac{4}{5\xi}\frac{5}{4}\xi^5 = \xi^4,$$

$$\sqrt{\frac{4}{5\xi}}(\xi^5 - 4\tau + 4\mu) \ge \xi^2.$$
(2.29)
(2.30)

Hence, taking into account (2.30), (2.29), from (2.28), we conclude that

$$\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle \xi \rangle^{2a} |\xi|^{2}}{\langle \tau + \xi^{5} \rangle^{2b} \langle \tau_{1} + \xi^{5}_{1} \rangle^{2b} \langle \tau - \tau_{1} + (\xi - \xi_{1})^{5} \rangle^{2b}} d\tau_{1} d\xi_{1} \\
\leq C \frac{\langle \xi \rangle^{2a} |\xi|}{\langle \tau + \xi^{5} \rangle^{2b}} \int_{\mathbb{R}_{\mu}} \frac{1}{\langle \mu \rangle^{4b-1} \sqrt{\frac{-16\tau + 16\mu - \xi^{5}}{5\xi}} |\xi|} d\mu \\
\leq C \frac{\langle \xi \rangle^{2a} |\xi|^{\frac{1}{2}}}{\langle \tau + \xi^{5} \rangle^{2b}} \int_{\mathbb{R}_{\mu}} \frac{1}{\langle \mu \rangle^{4b-1} \sqrt{\frac{-16\tau + 16\mu - \xi^{5}}{5\xi}} |\xi|} d\mu.$$
(2.31)

Using the calculus inequality (2.17) (Lemma 2.4) with $\rho := 4b - 1$ $(4b - 1 \in (\frac{1}{2}, 1) \iff \frac{3}{8} < b < \frac{1}{2}$) in (2.31), it follows that

$$\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle\xi\rangle^{2a}|\xi|^{2}}{\langle\tau+\xi^{5}\rangle^{2b}\langle\tau_{1}+\xi^{5}_{1}\rangle^{2b}\langle\tau-\tau_{1}+(\xi-\xi_{1})^{5}\rangle^{2b}} d\tau_{1}d\xi_{1} \leq C \frac{\langle\xi\rangle^{2a}|\xi|^{\frac{1}{2}}}{\langle\tau+\xi^{5}\rangle^{2b}\langle\tau+\frac{1}{16}\xi^{5}\rangle^{4b-\frac{3}{2}}}.$$
(2.32)

REGULARITY PROPERTIES AND DISPERSIVE BLOW-UP FOR THE FIFTH ORDER KORTEWEG-DE VRIES EQUATION ON THE LINE

It is clear, from (2.32), for $|\xi| \leq 1$, and $\frac{3}{8} < b < \frac{1}{2}$, that

$$\int_{\mathbb{R}^{2}_{\xi_{1}\tau_{1}}} \frac{\langle\xi\rangle^{2a} |\xi|^{2}}{\langle\tau + \xi^{5}\rangle^{2b} \langle\tau_{1} + \xi^{5}_{1}\rangle^{2b} \langle\tau - \tau_{1} + (\xi - \xi_{1})^{5}\rangle^{2b}} d\tau_{1} d\xi_{1} \leq C.$$
(2.33)

Let us assume then that $|\xi| > 1$. We observe that

$$|\xi^5| \le \frac{16}{15} |\tau + \xi^5| + \frac{16}{15} |\tau + \frac{1}{16} \xi^5|.$$

then

$$\frac{1}{2}|\xi^5| \le \frac{16}{15}|\tau + \xi^5| \quad \text{or} \quad \frac{1}{2}|\xi^5| \le \frac{16}{15}|\tau + \frac{1}{16}\xi^5|$$

For $\frac{1}{2}|\xi^5| \le \frac{16}{15}|\tau + \xi^5|$, we have that

$$C\frac{\langle\xi\rangle^{2a}|\xi|^{\frac{1}{2}}}{\langle\tau+\xi^{5}\rangle^{2b}\langle\tau+\frac{1}{16}\xi^{5}\rangle^{4b-\frac{3}{2}}} \le C\frac{|\xi|^{2a+\frac{1}{2}}}{\langle\tau+\xi^{5}\rangle^{2b}} \le C\frac{|\xi|^{2a+\frac{1}{2}}}{\langle\frac{15}{32}|\xi^{5}|\rangle^{2b}} \le C|\xi|^{2a+\frac{1}{2}-10b} \le C,$$
(2.34)

since $2a + \frac{1}{2} - 10b \le 20b - 8 + \frac{1}{2} - 10b = 10b - \frac{15}{2} < 0.$

For $\frac{1}{2}|\xi^5| \le \frac{16}{15}|\tau + \frac{1}{16}\xi^5|$, we have that

$$C \frac{\langle \xi \rangle^{2a} |\xi|^{\frac{1}{2}}}{\langle \tau + \xi^5 \rangle^{2b} \langle \tau + \frac{1}{16} \xi^5 \rangle^{4b - \frac{3}{2}}} \le C \frac{|\xi|^{2a + \frac{1}{2}}}{\langle \tau + \frac{1}{16} \xi^5 \rangle^{4b - \frac{3}{2}}} \le C \frac{|\xi|^{2a + \frac{1}{2}}}{\langle \frac{15}{32} |\xi^5| \rangle^{4b - \frac{3}{2}}} \le C |\xi|^{2a + \frac{1}{2} - 20b + \frac{15}{2}} \le C,$$
(2.35)
since $2a + \frac{1}{2} - 20b + \frac{15}{2} \le 20b - 8 + \frac{1}{2} - 20b + \frac{15}{2} = 0.$

In this way, from (2.32) to (2.35), it follows that, for $\frac{2}{5} < b < \frac{1}{2}$ and $0 \le a \le 10b - 4$, inequality (2.23) holds, which proves (2.21).

3. EXISTENCE OF A LOCAL SOLUTION IN TIME OF IVP (1.1). PROOF OF THEOREM 1.1 Let us demonstrate that there exists $T \in (0, \frac{1}{2}]$ such that the operator Γ_T , defined for $u \in X^{s,b}$ by

$$(\Gamma_T u)(t) := \eta(t)W(t)u_0 + \eta(t) \int_0^T W(t - t')F_T(u(t'))dt', \qquad (3.1)$$

has a fixed point in $X^{s,b}$.

Recall that by inequality (2.1) in Lemma 2.1, we have that

$$\|\eta(\cdot_t)[W(\cdot_t)u_0](\cdot_x)\|_{X^{s,b}(\mathbb{R}^2)} \le C \|u_0\|_{H^s(\mathbb{R}_x)},$$
(3.2)

and by using Lemma 2.2 with $T = 1, b' := -b^*$, where b^* is such that $b < b^* < \frac{1}{2}$,

$$\left\| \eta(\cdot_t) \int_0^{\cdot_t} W(\cdot_t - t') F_T(u(t')) dt' \right\|_{X^{s,b}} \le C \|F_T(u)\|_{X^{s,-b^*}}.$$
(3.3)

Since $-\frac{1}{2} < -b^* < -b < \frac{1}{2}$, by Lemma 2.3, with $2T \leq 1$, and Lemma 2.5, we have that

$$\|F_T(u)\|_{X^{s,-b^*}} \le CT^{-b+b^*}\| - \frac{1}{2}\partial_x(u^2)\|_{X^{s,-b}} \le CT^{-b+b^*}\|u\|_{X^{s,b}}^2.$$
(3.4)

Therefore, combining (3.3) and (3.4), we can affirm that

$$\left\| \eta(\cdot_t) \int_0^{\cdot_t} W(\cdot_t - t') F_T(u(t')) dt' \right\|_{X^{s,b}} \le C T^{-b+b^*} \|u\|_{X^{s,b}}^2.$$
(3.5)

From (3.2) and (3.5) we obtain

$$\|\Gamma_T u\|_{X^{s,b}} \le C(\|u_0\|_{H^s(\mathbb{R}_x)} + T^{-b+b^*} \|u\|_{X^{s,b}}^2).$$
(3.6)

Let us define the closed ball $B_{X^{s,b}}(0,R) \subset X^{s,b}$ centered at 0, with radius $R := 2C \|u_0\|_{H^s(\mathbb{R}_x)} > 0$. Hence, from (3.6), it follows that if $u \in B_{X^{s,b}}(0,R)$,

$$\|\Gamma_T u\|_{X^{s,b}} \le \frac{R}{2} + CT^{-b+b^*} R^2$$

Choosing T > 0 such that

$$CT^{-b+b^*}R^2 \le \frac{R}{2},$$
 (3.7)

we have that Γ_T maps $B_{X^{s,b}}(0,R)$ into itself.

It remains to prove that $\Gamma_T : B_{X^{s,b}}(0,R) \to B_{X^{s,b}}(0,R)$ is a contraction. Let us consider $v, w \in B_{X^{s,b}}(0,R)$. Then, by using Lemmas 2.2, 2.3, and 2.5, we have that

$$\|\Gamma_T v - \Gamma_T w\|_{X^{s,b}} = \left\| \eta(\cdot_t) \int_0^{\cdot_t} W(\cdot_t - t') [F_T(v(t')) - F_T(w(t'))] dt' \right\|_{X^{s,b}} \le CT^{-b+b^*} \|\partial_x (v^2 - w^2)\|_{X^{s,-b}} \le CT^{-b+b^*} \|v + w\|_{X^{s,b}} \|v - w\|_{X^{s,b}} \le 2CRT^{-b+b^*} \|v - w\|_{X^{s,b}}.$$
(3.8)

If we choose T > 0 such that $2CRT^{-b+b^*} < 1$, it is clear that T also satisfies (3.7). In consequence, from (3.8) we conclude that $\Gamma_T: B_{X^{s,b}}(0, R) \to B_{X^{s,b}}(0, R)$ is a contraction.

Let us observe that we can also assume that $T \in (0, \frac{1}{2}]$. Therefore, there exist $T \in (0, \frac{1}{2}]$ and a unique $u \in B_{X^{s,b}}(0, R)$ such that $\Gamma_T u = u$.

Now we prove that $u \in C(\mathbb{R}_t; H^s(\mathbb{R}_x))$. Let us note that the first term on the right hand side of (3.1) is continuous from \mathbb{R}_t , with values in $H^s(\mathbb{R}_x)$. To demonstrate that the second term on the right hand side of (3.1) is continuous form \mathbb{R}_t , with values in $H^s(\mathbb{R}_x)$, it is enough to see that this term belongs to $X^{s,\tilde{b}}$ for some $b > \frac{1}{2}$.

Let us take $\tilde{b} > \frac{1}{2}$ such that $b^* + \tilde{b} \leq 1$. Then

$$-\frac{1}{2} < -b^* < 0 \le \widetilde{b} \le (-b^*) + 1.$$

By applying Lemma 2.2 with T = 1, and Lemmas 2.3 and 2.5, we conclude that

$$\left\| \eta(\cdot_t) \int_0^{\cdot_t} W(\cdot_t - t') F_T(u(t')) dt' \right\|_{X^{s,\tilde{b}}} \leq C \|\eta\left(\frac{\cdot_t}{2T}\right) \partial_x u^2\|_{X^{s,-b^*}} \leq CT^{-b+b^*} \|\partial_x u^2\|_{X^{s,-b}} \leq CT^{-b+b^*} \|u\|_{X^{s,b}}^2 < \infty.$$

In this way, we have proved that $u = \Gamma_T u \in C(\mathbb{R}_t; H^s(\mathbb{R}_x))$. Theorem 1.1 is proved.

In this way, we have proved that $u = \Gamma_T u \in C(\mathbb{R}_t; H^s(\mathbb{R}_x))$. Theorem 1.1 is proved.

4. Regularity gain of the nonlinear part of the solution of IVP (1.1). Proof of Theorem 1.2

Let b^* and \tilde{b} be such that $b < b^* < \frac{1}{2} < \tilde{b}$, and $b^* + \tilde{b} \le 1$. By using the immersion $X^{s+a,\tilde{b}} \hookrightarrow C_b(\mathbb{R}_t; H^{s+a}(\mathbb{R}_x))$ and taking into account that for $0 \le t' \le t \le T \le \frac{1}{2}$, $\eta(t) = 1$ and $-\frac{1}{2}\partial_x(u(t'))^2 = F_T(u(t'))$, we have that

$$\begin{aligned} \left\| \int_{0}^{\cdot t} W(\cdot_{t} - t') (-\frac{1}{2} \partial_{x} (u(t'))^{2}) dt' \right\|_{C([0,T];H^{s+a}(\mathbb{R}_{x}))} \\ &= \left\| \eta(\cdot_{t}) \int_{0}^{\cdot t} W(\cdot_{t} - t') F_{T}(u(t')) dt' \right\|_{C([0,T];H^{s+a}(\mathbb{R}_{x}))} \leq \left\| \eta(\cdot_{t}) \int_{0}^{\cdot t} W(\cdot_{t} - t') F_{T}(u(t')) dt' \right\|_{C_{b}(\mathbb{R}_{t};H^{s+a}(\mathbb{R}_{x}))} \\ &\leq C \left\| \eta(\cdot_{t}) \int_{0}^{\cdot t} W(\cdot_{t} - t') F_{T}(u(t')) dt' \right\|_{X^{s+a,\tilde{b}}}. \end{aligned}$$

$$(4.1)$$

Applying Lemma 2.2 with $-\frac{1}{2} < -b^* \le 0 \le \tilde{b} \le -b^* + 1$, it follows that

$$\left\| \eta(\cdot_t) \int_0^{\cdot_t} W(\cdot_t - t') F_T(u(t')) dt' \right\|_{X^{s+a,\tilde{b}}} \le C \|F_T(u(\cdot))\|_{X^{s+a,-b^*}}.$$
(4.2)

From Lemma 2.3 with $b_1 := -b^*$, $b_2 := -b$, and Lemma 2.5, we can conclude that

$$||F_T(u(\cdot))||_{X^{s+a,-b^*}} \le CT^{-b+b^*} ||\partial_x u^2||_{X^{s+a,-b}} \le C||u||_{X^{s,b}}^2.$$
(4.3)

REGULARITY PROPERTIES AND DISPERSIVE BLOW-UP FOR THE FIFTH ORDER KORTEWEG-DE VRIES EQUATION ON THE LINE

Finally, from (4.1), (4.2) and (4.3), we have that

$$\left\|\int_0^{\cdot t} W(\cdot_t - t')(-\frac{1}{2}\partial_x (u(t'))^2) dt'\right\|_{C([0,T];H^{s+a}(\mathbb{R}))} \le C \|u\|_{X^{s,b}}^2 < +\infty,$$

and Theorem 1.2 is proved.

5. Dispersive blow-up. Proof of Theorem 1.3

Let us define $\phi(x) := e^{-2|x|}$, $x \in \mathbb{R}$. It is clear that $\phi \in C^1(\mathbb{R} \setminus \{0\}) \setminus C^1(\mathbb{R})$, $e^x \phi \in L^2(\mathbb{R})$, and $e^{-x} \phi \in L^2(\mathbb{R})$. Let $\{\alpha_j\}_{j=1}^{\infty}$ be a sequence of positive real numbers such that $\sum_{j=1}^{\infty} \alpha_j e^{4j} < +\infty$, and let us define

$$u_0(x) := \sum_{j=1}^{\infty} \alpha_j [W(-j)\phi](x).$$
(5.1)

Then we have that $u_0 \in H^1(\mathbb{R})$ (in fact, $u_0 \in H^{\frac{3}{2}-}(\mathbb{R})$). Actually,

$$\begin{split} \|W(-j)\phi\|_{H^{\frac{3}{2}-}} &= \left(\int_{-\infty}^{+\infty} (1+\xi^2)^{\frac{3}{2}-} |[W(-j)\phi]^{\wedge}(\xi)|^2 d\xi\right)^{\frac{1}{2}} = \left(\int_{-\infty}^{+\infty} (1+\xi^2)^{\frac{3}{2}-} |e^{ij\xi^5}\widehat{\phi}(\xi)|^2 d\xi\right)^{\frac{1}{2}} \\ &= \left(\int_{-\infty}^{+\infty} (1+\xi^2)^{\frac{3}{2}-} |\widehat{\phi}(\xi)|^2 d\xi\right)^{\frac{1}{2}}. \end{split}$$

Since

$$\widehat{\phi}(\xi) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} e^{-i\xi x} e^{-2|x|} dx = \frac{1}{\sqrt{2\pi}} \left[\frac{1}{2+i\xi} + \frac{1}{2-i\xi} \right] = \frac{4}{\sqrt{2\pi}(\xi^2 + 4)}$$

then

$$\|W(-j)\phi\|_{H^{\frac{3}{2}-}} = \left(\int_{-\infty}^{+\infty} (1+\xi^2)^{\frac{3}{2}-} \frac{16}{2\pi(\xi^2+4)^2} d\xi\right)^{\frac{1}{2}} < +\infty,$$
(5.2)

From (5.1) and (5.2) it can be seen that $u_0 \in H^{\frac{3}{2}-}$, given that

$$\sum_{j=1}^{\infty} \alpha_j \|W(-j)\phi\|_{H^{\frac{3}{2}-}} = \left(\int_{-\infty}^{+\infty} (1+\xi^2)^{\frac{3}{2}-} \frac{16}{2\pi(\xi^2+4)^2} d\xi\right) \sum_{j=1}^{\infty} \alpha_j$$
$$\leq \left(\int_{-\infty}^{+\infty} (1+\xi^2)^{\frac{3}{2}-} \frac{16}{2\pi(\xi^2+4)^2} d\xi\right) \sum_{j=1}^{\infty} \alpha_j e^{4j} < +\infty.$$

It can be seen that the solution u of the IVP (1.1) is global in time and is such that

$$u(t) = W(t)u_0 + \int_0^t W(t - t')(-\frac{1}{2}\partial_x(u(t'))^2)dt' \equiv W(t)u_0 + z(t).$$
(5.3)

By Theorem 1.2, $z(t) \in C^1(\mathbb{R})$ for every $t \ge 0$. (In fact, there exists $\beta > 0$ such that $z(t) \in H^{(\frac{3}{2}-)+\beta}$. Therefore, by taking $\frac{3}{2}$ - close enough to $\frac{3}{2}$ from the left, it follows that $(\frac{3}{2}-)+\beta > \frac{3}{2}$, and as a result $z(t) \in C^1(\mathbb{R})$).

Consequently, the proof of Theorem 1.3 reduces to establish statements (i) and (ii) for $W(t)u_0$ instead of $u(\cdot, t)$. Let us define

$$w(x,t) := e^x [W(t)\phi](x).$$
 (5.4)

Then

$$\frac{\partial_t(e^{-x}w) + \partial_x^5(e^{-x}w) = 0,}{w(x,0) = e^x\phi(x),}$$

$$(5.5)$$

which means that w satisfies the following differential equation:

$$\partial_t w + \partial_x^5 w - 5\partial_x^4 w + 10\partial_x^3 w - 10\partial_x^2 w + 5\partial_x w - w = 0,$$

which indicates that w satisfies the IVP

$$\frac{\partial_t w + (\partial_x - 1)^5 w = 0,}{w(x,0) = e^x \phi(x).}$$
(5.6)

Taking the Fourier transform with respect to the variable x in the differential equation in (5.6), we obtain the IVP

$$\frac{d}{dt}\widehat{w(\cdot,t)} + (i\xi - 1)^5 \widehat{w(\cdot,t)} = 0, \\
\widehat{w(\cdot,0)} = [e^x \phi]^{\wedge}(\xi),$$
(5.7)

whose solution is

$$\widehat{w(\cdot,t)}(\xi) = e^{-i\xi^5 t} e^{5t\xi^4} e^{10it\xi^3} e^{-10t\xi^2} e^{-5it\xi} e^t (e^x \phi)^{\wedge}(\xi).$$
(5.8)

Let us consider

$$\widetilde{w}(x,t) := e^{-x} [W(t)\phi](x).$$
(5.9)

In a similar way as we obtained (5.8), it can be seen that

$$\widehat{\widetilde{w}(\cdot,t)}(\xi) = e^{-i\xi^5 t} e^{-5t\xi^4} e^{10it\xi^3} e^{10t\xi^2} e^{-5it\xi} e^{-t} (e^{-x}\phi)^{\wedge}(\xi).$$
(5.10)

Next, we prove that $u_0 \in C^{\infty}(\mathbb{R})$. For that, it is sufficient to prove that $\sum_{j=1}^{\infty} \alpha_j e^x W(-j) \phi \in C^{\infty}(\mathbb{R})$, which will be true if for every $m \in \mathbb{N} \cup \{0\}$, we have $\sum_{j=1}^{\infty} \alpha_j \|e^x W(-j)\phi\|_{H^m(\mathbb{R})} < +\infty$.

For m = 0, using Plancherel's Theorem and (5.8), it results that

$$\sum_{j=1}^{\infty} \alpha_j \| e^x W(-j)\phi \|_{L^2(\mathbb{R})} = \sum_{j=1}^{\infty} \alpha_j \| w(\cdot_x, -j) \|_{L^2_x} = \sum_{j=1}^{\infty} \| \widehat{w(\cdot_x, -j)}(\cdot_\xi) \|_{L^2_\xi} = \sum_{j=1}^{\infty} \alpha_j \| e^{-5j(\xi^4 - 2\xi^2)} e^{-j} (e^x \phi)^{\wedge}(\xi) \|_{L^2_\xi}$$
$$= \sum_{j=1}^{\infty} \alpha_j \| e^{-5j(\xi^4 - 2\xi^2 + 1)} e^{4j} (e^x \phi)^{\wedge}(\xi) \|_{L^2_\xi} \le \left(\sum_{j=1}^{\infty} \alpha_j e^{4j} \right) \| e^x \phi \|_{L^2_x} < \infty.$$
(5.11)

Now, we prove that, for every $m \in \mathbb{N}$,

$$\sum_{j=1}^{\infty} \alpha_j \|\partial_x^m (e^x W(-j)\phi)\|_{L^2_x} < +\infty.$$
(5.12)

In fact,

$$\begin{split} \sum_{j=1}^{\infty} \alpha_j \|\partial_x^m (e^x W(-j)\phi)\|_{L_x^2} &= \sum_{j=1}^{\infty} \alpha_j \|\xi^m e^{-5j(\xi^2 - 1)^2} e^{4j} (e^x \phi)^{\wedge}(\xi)\|_{L_{\xi}^2} \\ &\leq \sum_{j=1}^{\infty} \alpha_j e^{4j} \|\xi^m e^{-5j(\xi^2 - 1)^2}\|_{L_{\xi}^{\infty}} \|e^x \phi\|_{L_x^2} \leq \sum_{j=1}^{\infty} \alpha_j e^{4j} \frac{\left(1 + \sqrt{1 + \frac{m}{5j}}\right)^{m/2}}{2^{m/2}} \|e^x \phi\|_{L_x^2} \\ &\leq \frac{\left(1 + \sqrt{1 + \frac{m}{5}}\right)^{m/2}}{2^{m/2}} \|e^x \phi\|_{L_x^2} \sum_{j=1}^{\infty} \alpha_j e^{4j} < +\infty. \end{split}$$

From (5.11) and (5.12) we conclude that

$$\sum_{j=1}^{\infty} \alpha_j \| e^x W(-j) \phi \|_{H^m(\mathbb{R})} < +\infty.$$

Now let us prove that, for every $t \in (0, +\infty) \setminus \mathbb{Z}^+$, $W(t)u_0 \in C^1(\mathbb{R})$. Initially, assume that t > 1. Then there exists $j_0 \in \mathbb{Z}^+$ such that $j_0 < t < j_0 + 1$, and

$$W(t)u_0 = \sum_{j=1}^{j_0} \alpha_j W(t-j)\phi + \sum_{j=j_0+1}^{\infty} \alpha_j W(t-j)\phi.$$

For $j \leq j_0$, $W(t-j)\phi = e^x \widetilde{w}(x,t-j)$, and thus, to see that $W(t-j)\phi \in C^1(\mathbb{R})$, it is enough to prove that $\widetilde{w}(x,t-j) \in C^1(\mathbb{R}_x)$. This last statement is true if we prove that $\widetilde{w}(\cdot_x,t-j) \in H^2(\mathbb{R}_x)$. Indeed, for $j \leq j_0 < t$, using (5.11), we have that

$$\begin{split} \|\widetilde{w}(\cdot_{x},t-j)\|_{H^{2}} &\leq C(\|\widetilde{w}(\cdot_{x},t-j)\|_{L^{2}} + \|\partial_{x}^{2}\widetilde{w}(\cdot_{x},t-j)\|_{L^{2}}) \\ &\leq C(\|[\widetilde{w}(\cdot_{x},t-j)]^{\wedge}(\cdot_{\xi})\|_{L^{2}} + \|\xi^{2}[\widetilde{w}(\cdot_{x},t-j)]^{\wedge}(\cdot_{\xi})\|_{L^{2}}) \\ &\leq C\left(\|e^{i(t-j)(-\xi^{5}+10\xi^{3}-5\xi)}e^{-5(t-j)(\xi^{2}-1)^{2}}e^{4(t-j)}(e^{-x}\phi)^{\wedge}(\xi)\|_{L^{2}_{\xi}} + \|\xi^{2}e^{i(t-j)(-\xi^{5}+10\xi^{3}-5\xi)}e^{-5(t-j)(\xi^{2}-1)^{2}}e^{4(t-j)}(e^{-x}\phi)^{\wedge}(\xi)\|_{L^{2}_{\xi}}\right) \\ &\leq C\left(e^{4(t-j)}\|e^{-x}\phi\|_{L^{2}_{x}} + e^{4(t-j)}\|\xi^{2}e^{-5(t-j)(\xi^{2}-1)^{2}}\|_{L^{\infty}_{\xi}}\|e^{-x}\phi\|_{L^{2}_{x}}\right) < \infty. \end{split}$$
(5.13)

From (5.13), we conclude that $\sum_{j=1}^{j_0} \alpha_j W(t-j)\phi \in C^1(\mathbb{R})$. To show that $\sum_{j=j_0+1}^{\infty} \alpha_j W(t-j)\phi \in C^1(\mathbb{R})$, it suffices to demonstrate that $\sum_{j=j_0+1}^{\infty} \alpha_j e^x W(t-j)\phi \in C^1(\mathbb{R})$. For this, it is sufficient to prove that $\sum_{j=j_0+1}^{\infty} \alpha_j e^x W(t-j)\phi \in H^2(\mathbb{R})$. This last statement will be true if we prove that

$$\sum_{j=j_0+1}^{\infty} \alpha_j \| e^x W(t-j)\phi \|_{H^2} < \infty.$$
(5.14)

In fact,

$$\begin{split} \sum_{j=j_0+1} \alpha_j \| e^x W(t-j) \phi \|_{H^2} &= \sum_{j=j_0+1}^{\infty} \alpha_j \| w(\cdot_x, t-j) \|_{H^2} \le C \sum_{j=j_0+1}^{\infty} \alpha_j \left(\| [w(\cdot, t-j)]^{\wedge}(\cdot_{\xi}) \|_{L^2} + \| \xi^2 [w(\cdot, t-j)]^{\wedge}(\cdot_{\xi}) \|_{L^2_{\xi}} \right) \\ &\leq C \sum_{j=j_0+1}^{\infty} \alpha_j \left(e^{-4(t-j)} \| e^x \phi \|_{L^2_x} + \| \xi^2 e^{5(t-j)(\xi^2-1)^2} e^{-4(t-j)} (e^x \phi)^{\wedge}(\xi) \|_{L^2_{\xi}} \right) \\ &\leq C e^{-4t} \sum_{j=j_0+1}^{\infty} \alpha_j \left(e^{4j} \| e^x \phi \|_{L^2_x} + e^{4j} \| \xi^2 e^{5(t-(j_0+1))(\xi^2-1)^2} (e^x \phi)^{\wedge}(\xi) \|_{L^2_{\xi}} \right) \\ &\leq C e^{-4t} \sum_{j=j_0+1}^{\infty} \alpha_j \left(e^{4j} \| e^x \phi \|_{L^2_x} + e^{4j} \| \xi^2 e^{5(t-(j_0+1))(\xi^2-1)^2} \|_{L^\infty_{\xi}} \| e^x \phi \|_{L^2_x} \right) \\ &\leq C e^{-4t} \| e^x \phi \|_{L^2_x} \sum_{j=j_0+1}^{\infty} \alpha_j e^{4j} < +\infty, \end{split}$$

which is precisely (5.14).

Let us observe that, if 0 < t < 1, the analysis is the same as that carried out for the proof of the series $\sum_{j=j_0+1}^{\infty} \alpha_j W(t-j)\phi$.

Thus we conclude that if $t \in (0, +\infty) \setminus \mathbb{Z}^+$, it follows that

$$W(t)u_0 = \sum_{j=1}^{j_0} \alpha_j W(t-j)\phi + \sum_{j=j_0+1}^{\infty} \alpha_j W(t-j)\phi \in C^1(\mathbb{R}).$$

Finally, let us see that, if t = n, for some $n \in \mathbb{Z}^+$, then $W(t)u_0$ belongs to $C^1(\mathbb{R} \setminus \{0\}) \setminus C^1(\mathbb{R})$. That is, let us see that

$$W(n)u_0 = \alpha_n \phi + \sum_{j \in \mathbb{Z}^+ \setminus \{n\}} \alpha_j W(n-j)\phi \equiv \alpha_n \phi + \Phi_n \in C^1(\mathbb{R} \setminus \{0\}) \setminus C^1(\mathbb{R}).$$

Proceeding as in the case $t \notin \mathbb{Z}^+$, it can be shown that

$$\Phi_n \equiv \sum_{j \in \mathbb{Z}^+ \backslash \{n\}} \alpha_n W(n-j) \phi \in C^1(\mathbb{R}),$$

and since $\alpha_n \phi \in C^1(\mathbb{R} \setminus \{0\}) \setminus C^1(\mathbb{R})$, then $W(n)u_0 \notin C^1(\mathbb{R})$. Otherwise, since $\alpha_n > 0$, we would have that $\phi \in C^1(\mathbb{R})$. Moreover, since $\alpha_n \phi \in C^1(\mathbb{R} \setminus \{0\})$, and $\Phi_n \in C^1(\mathbb{R} \setminus \{0\})$, it follows that $W(n)u_0 \in C^1(\mathbb{R} \setminus \{0\})$.

Theorem 1.3 is proved.

Acknowledgments

This work was partially supported by Universidad Nacional de Colombia, Sede-Medellín– Facultad de Ciencias – Departamento de Matemáticas – Grupo de investigación en Matemáticas de la Universidad Nacional de Colombia Sede Medellín, carrera 65 No. 59A -110, post 50034, Medellín Colombia. Proyecto: Análisis no lineal aplicado a problemas mixtos en ecuaciones diferenciales parciales, código Hermes 60827. Fondo de Investigación de la Facultad de Ciencias empresa 3062.

References

- Benjamin, T, Bona, J., Mahony, J., Model equations for long waves in nonlinear dispersive systems, Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences 272 (1972), 47-78.
- [2] Bona, J., Ponce, G., Saut, J.C., Sparber, C., Dispersive blow-up for nonlinear Schrödinger equations revisited, Journal de Mathématiques Pures et Appliquées 102.4 (2014): 782-811.
- Bona, J.L., Saut, J.C., Dispersive blow-up of solutions of generalized Korteweg-de Vries equations, J. Differential equations 103 (1993), 3-57.
- [4] Bona, J.L., Saut, J.C., Dispersive blow-up II. Schrödinger-type equations, optical and oceanic rogue waves, Chinese Annals of Mathematics, Series B 31 (6) (2010), 793-818.
- [5] Bustamante, E., Jiménez, J., Mejía, J., Cauchy problems for fifth-order KdV equations in weighted Sobolev spaces, Electronic Journal of Differential Equations (141) (2015), 1–24.
- [6] Erdoğan, M., Tzirakis, N. Smoothing and global attractors for the Zakharov system on the torus, Analysis & PDE 6 (3) (2013), 723-750.
- [7] Erdoğan, M., Tzirakis, N. Regularity properties of the cubic nonlinear Schrödinger equation on the half line, Journal of Functional Analysis 271 (9) (2016), 2539-2568.
- [8] Ginibre, J., Tsutsumi, Y., Velo, G., On the Cauchy problem for the Zakharov system, J. Funct. Anal. 151 (1997), 384-436.
- [9] Guo, Z., Kwak, C., Kwon, S. Rough solutions of the fifth-order KdV equations, Journal of Functional Analysis 265 (11) (2013), 2791-2829.
- [10] Kato, T., Local well-posedness for Kawahara equation, Adv. Differential Equations 16 (2011), No. 3-4, 257-287.
- [11] Kato, T., Global well-posedness for the Kawahara equation with low regularity data, Communications on Pure and Applied Analysis, 12, 3 (2013), 1321-1339.
- [12] Kawahara, T., Oscillatory solitary waves in dispersive media, J. Phys. Soc. Japan, 33 (1972), 260-264.
- [13] Kenig, C., Pilod, D., Well-posedness for the fifth-order KdV equation in the energy space, Transactions of the American Mathematical Society 367 (4) (2015), 2551-2612.
- [14] Kenig, C., Ponce, G., Vega, L., Well posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle, Communications on pure and applied mathematic 46 (4) (1993), 527-620.
- [15] Kwon, S., On the fifth-order KdV equation: local well-posedness and lack of uniform continuity of the solution map, Journal of Differential Equations 245 (9) (2008), 2627-2659.
- [16] Kwon, S., Well-posedness and ill-posedness of the fifth-order modified KdV equation, Electronic Journal of Differential Equations 2008 (2008) No. 01, 1-15.
- [17] Linares, F., A higher order modified Korteweg-de Vries equation, Comp. Appl. Math 14 No. 3 (1995), 253-267.
- [18] Linares, F., Palacios, J.M., Dispersive blow-up and persistence properties for the Schrödinger-Korteweg-de Vries system, Nonlinearity 32 (2019), 4996-5016.
- [19] Linares, F., Ponce, G., Smith, D., On the regularity of solutions to a class of nonlinear dispersive equations, Math. Ann. 369 (2017), 797–837.
- [20] Linares, F., Scialom, M., On the smoothing properties of solutions to the modified Korteweg-de Vries equation, Journal of Differential Equations, 106 (1993), 141-154.
- [21] Muñoz, A., Pastor, A., Local well-posedness in weighted Sobolev spaces for nonlinear dispersive equations with applications to dispersive blow up, Math. Ann. 386 No. 1 (2023) 207-246.

Eddye Bustamante M., José Jiménez Urrea, Jorge Mejía L.

Departamento de Matemáticas, Universidad Nacional de Colombia

A. A. 3840 Medellín, Colombia

Email address: eabusta0@unal.edu.co, jmjimene@unal.edu.co, jemejia@unal.edu.co