Estimating Treatment Effects under Recommender Interference: A Structured Neural Networks Approach

Ruohan Zhan
rhzhan@ust.hkShichao Han
shichaohan@tencent.comYuchen Hu
yuchenhu@stanford.edu
Stanford UniversityThe Hong Kong University of
Science and TechnologyTencent Inc.Stanford University

Zhenling Jiang* zhenling@wharton.upenn.edu University of Pennsylvania

Recommender systems are essential for content-sharing platforms by curating personalized content. To evaluate updates to recommender systems targeting content creators, platforms frequently rely on creatorside randomized experiments. The treatment effect measures the change in outcomes when a new algorithm is implemented compared to the status quo. We show that the standard difference-in-means estimator can lead to biased estimates due to recommender interference that arises when treated and control creators compete for exposure. We propose a "recommender choice model" that describes which item gets exposed from a pool containing both treated and control items. By combining a structural choice model with neural networks, this framework directly models the interference pathway while accounting for rich viewer-content heterogeneity. We construct a debiased estimator of the treatment effect and prove it is \sqrt{n} -consistent and asymptotically normal with potentially correlated samples. We validate our estimator's empirical performance with a field experiment on Weixin short-video platform. In addition to the standard creator-side experiment, we conduct a costly double-sided randomization design to obtain a benchmark estimate free from interference bias. We show that the proposed estimator yields results comparable to the benchmark, whereas the standard difference-in-means estimator can exhibit significant bias and even produce reversed signs.

Key words: Recommender Interference; Treatment Effect Estimation; Online Content Platforms; Creator-side Randomization; Semiparametric Choice Model; Double/Debiased Estimation and Inference

We are grateful to Susan Athey, Tat Chan, Wanning Chen, Ryan Dew, Yue Fang, Bryan Graham, Ganesh Iyer, Ramesh Johari, Jussi Keppo, Hannah Li, Ilan Lobel, Xiaojie Mao, Tu Ni, Nian Si, Zikun Ye, Dennis Zhang, Jinglong Zhao, conference participants at CODE@MIT 2023, Conference on AI/ML and Business Analytics 2023, and ACIC 2024, and seminar attendees at University of Washington, National University of Singapore, Tsinghua University, The Chinese University of Hong Kong (Shenzhen) for their helpful comments. The short version of this paper has appeared in the ACM Conference on Economics and Computation (EC'24), and we are especially thankful to the anonymous reviewers for their constructive feedback. We are grateful to the WeChat Experimentation Team at Tencent for their generous support, particularly Darwin Yong Wang, who has helped shape and advance this work's applications in the online platform.

^{*}Corresponding author.

1. Introduction

Recommender systems (RecSys) are at the heart of online content platforms by curating viewerspecific content selections based on viewer history and content characteristics. Indeed, recommendation plays a significant role in driving content consumption for viewers (Kiros 2022), and thus determines the exposure and success of content creators. This is particularly the case for shortvideo platforms such as TikTok, where a stream of videos is directly presented to viewers in the For-You feed determined by the recommendation system.

Recognizing the importance of RecSys, platforms regularly develop and assess new algorithms. In this paper, we focus on those aimed at improving the experience of content creators. Platforms rely on creators to generate a diverse and vibrant content pool, which ensures viewer satisfaction and platform success in the long term (Mehrotra et al. 2018, Rosen 1981). By tweaking the recommendation algorithm, platforms can incentivize creators in different ways. For example, recommendation algorithms may boost the visibility of creators who produce higher-quality content or those who invest in paid promotions. Such adjustments can directly influence the engagement and activities of content creators.

To evaluate the performance of these creator-focused algorithm updates, platforms typically rely on experiments with "creator-side randomization". This involves randomly assigning a group of creators in the treatment group, served by the new algorithm, and the others in the control group, served by the status quo algorithm. The treatment effect is the change in outcomes when the new algorithm is implemented compared to the status quo. A standard way to estimate the treatment effect is the difference-in-means (DIM) estimator, which directly calculates the difference in outcomes between treatment vs. control creators.

We show that the DIM estimator can lead to biased treatment effect estimates. This bias arises because treated and control creators compete for exposure through the recommendation algorithm. As a result, outcomes in one group can depend on creators in another group, a phenomenon known as interference or spill-over effects (Bajari et al. 2021, Johari et al. 2022, Bright et al. 2022, Goli et al. 2023). Since a creator's outcomes are influenced not only by their own treatment status but also by the statuses of others, this violates the Stable Unit Treatment Value Assumption (SUTVA) (Imbens 2004) and leads to biased treatment effect estimates. For example, an algorithm that boosts exposure for high-quality creators may produce large effects in a small-scale treatment group, relative to those not receiving the boost in control, but fails to generalize upon full-scale implementation when all high-quality creators receive the boost.

In this paper, we propose an approach to estimate the treatment effect with data from creatorside randomized experiments. A key element in the proposed framework is to model how the algorithm chooses an item to expose to viewers from a set comprised of both treated and control units. This "recommender choice model" captures the interference pathway: outcomes of treated units are affected by the control group because they can be in the same pool competing for exposure. The choice model can rationalize and predict that outcomes can differ between small-scale A/B testing where treated units compete with others in control, and full-scale implementation where all units are treated.

The recommender choice model describes which item gets exposed to a viewer from a set with both treated and control items. The treatment effect estimate is the difference in outcomes between two counterfactual scenarios: when all items are treated vs. all in control. The model leverages an insight that different recommendation algorithms have different ways of calculating the "score" of an item for each viewer, which drives the recommendation process. For example, an algorithm focused on high-quality creators will assign higher scores to these creators. We use separate flexible neural networks to capture the potentially complex mapping from viewer-content pairs to their scores under both treatment and control conditions. One advantage of this method is that it is agnostic to the specific, potentially very complex, recommender system, thus making it broadly applicable to many scenarios.

Methodologically, the proposed semi-parametric approach combines a structural choice model with flexible neural networks. The structural choice model framework is important as it facilitates counterfactual evaluation and inference under alternative treatment assignments (e.g., all treated or all control). This would be difficult to achieve with a fully non-parametric model without the choice model structure. These counterfactual evaluations can be leveraged to calculate the average treatment effect (ATE) across the whole population or for specific subgroups of interest. The neural networks in the choice model are essential for capturing the nuances in how recommendation algorithms score items by accounting for rich viewer-content heterogeneity. This would be very difficult to capture using a parametric model, unlike in settings with only structured attributes such as price and brand.

Building on the double machine learning literature (Farrell et al. 2020, Chernozhukov et al. 2018), we propose a debiased estimator for treatment effect estimation leveraging the semi-parametric recommender choice model. We show that, under regularity conditions, the estimator is \sqrt{n} -consistent and asymptotically normal. Our results build on Chernozhukov et al. (2018), which deals with i.i.d. samples, and extends to scenarios with correlated samples. In our empirical context, sample correlation arises because, for samples with overlapping items, their treatment statuses remain the same instead of being i.i.d. By modeling the data generating process in a sequential manner, we can apply martingale limiting theorems (Hall and Heyde 2014) to establish the asymptotic property of the "oracle estimator" with known nuisance components. Since the debiased estimator satisfies the Neyman orthogonality condition, the difference between the debiased and oracle estimators is shown to be small. Our inference framework can be adapted to other scenarios with correlated samples, such as panel data with repeated observations per customer.

To demonstrate the effectiveness of the proposed approach, we start with Monte Carlo simulations to validate the estimation and inference results of the debiased estimator. Beyond Monte Carlo studies, we apply it to real experimental data from the Weixin short-video platform. In addition to the standard creator-side randomized experiment, we also carry out a double-sided experiment (Ye et al. 2023a, Su et al. 2024). This is done by dividing viewers and creators into three equal-sized *distinct* universes, where viewers can only access the creators from their own universe. One universe runs the creator-side experiment and the other two implement the treatment and control algorithms, respectively. The double-sided design allows us to obtain a benchmark estimate without recommender interference since the treated and control units no longer compete with each other. Using data from the creator-side experiment, we compute the treatment effect with both the standard DIM and the proposed debiased estimators. We show that the proposed estimator yields results comparable to the benchmark, whereas the DIM estimator can exhibit significant bias and even produce reversed signs.

Despite the effectiveness of the double-sided randomization design, implementing such a strategy has several downsides: it is very costly from an engineering perspective; the market size is effectively cut in half for both viewers and creators; and the experimental population cannot be reused for other orthogonal experiments. Viewer-side experiments, where viewers are randomized to treatment vs. control algorithms, do not apply when the focus is on creator outcomes or creator interventions. Viewer-side randomization design can also introduce its own source of interference bias, where outcomes from one condition affect the other condition. For example, if the treatment algorithm favors high-quality creators who receive more exposure and likes, these outcomes may spill over to viewers in the control group.

The proposed estimator uses data from standard creator-side experiments to measure the treatment effect of creator-focused interventions. In particular, it captures the interference coming from units competing for exposure through the recommender system. It does not explicitly consider potential viewer-side temporal interference, where viewer responses might be influenced by previous content. We also abstract away from the potential long-run effect of the treatment, and take the viewer queries and available content as given on the platform.

This paper makes two main contributions. Substantively, it identifies a common source of interference from treated and control units competing for exposure, and proposes a debiased treatment effect estimator. We validate the approach using experimental data on a leading short-video platform by comparing the results to a benchmark from double-sided randomization without interference. Methodologically, the proposed approach combines a structural choice model with neural networks. Our inference results extend the existing results from Chernozhukov et al. (2018), which deals with i.i.d. samples, to correlated samples, which can broaden the applicability of the debiased estimator to other settings with correlated samples, such as panel data.

1.1. Related Literature

There is a growing literature studying interference in randomized experiments, where an individual's outcome is affected by others' treatments, violating the Stable Unit Treatment Value Assumption (SUTVA). Interference can lead to biased treatment effect estimates under standard A/B testing framework (Sävje et al. 2021, Hu et al. 2021, Farias et al. 2022, Johari et al. 2022, Dhaouadi et al. 2023, Zhu et al. 2024). To correctly estimate treatment effect, most of the existing literature leverages innovative *experimental designs* to address the bias from interference. For example, one can use a clustered randomization design if interference primarily occurs within a cluster (e.g., on social networks) and then randomizes treatment at the cluster level (Holtz and Aral 2020, Holtz et al. 2023, Ugander et al. 2013, Hudgens and Halloran 2008). Without a clear cluster structure, a switchback design can be applied that assigns a treatment condition randomly at the market level across different time periods (Bojinov et al. 2023, Hu and Wager 2022, Ni et al. 2023, Xiong et al. 2023). These experimental designs can be very useful but do not apply to all settings, including ours, where the content markets are too interconnected to form clusters and there exist significant time varying factors influencing the outcomes.

Our work aligns more closely with an alternative approach that proposes innovative *estimators* while leveraging standard experimental design. This requires domain-specific analysis of interference types. For example, in experiments for ranking algorithms, Goli et al. (2023) address interference by leveraging items with positions close to those under the counterfactual ranking using historical A/B test data. In marketplaces where the platform matches supply and demand via linear programming, Bright et al. (2022) propose a shadow-price based estimator to mitigate interference bias. For interference bias that arises from consumers impacting others through limited inventory, Farias et al. (2023) model consumer behavior via a Markovian structure to address interference. Our paper proposes an estimator that applies to a common source of interference where treated and control units compete for exposure in online platforms.

This paper draws from recent advances in double machine learning and semiparametric inference (Newey 1994, Chernozhukov et al. 2018, Farrell et al. 2020, Chernozhukov et al. 2019). In particular, the proposed treatment effect estimator directly builds on the doubly robust estimators for semi-parametric models proposed by Farrell et al. (2020), where the parametric outcome model is enriched by non-parametric components. We show that the inference results of the debiased estimator in Farrell et al. (2020) apply to correlated samples, thus extending results from Chernozhukov

et al. (2018), which deals with i.i.d. samples. While in our setting, sample correlation arises from the overlapped items in consideration sets that share treatment status, our inference results can apply to other settings with correlated samples, such as panel settings with repeated observations from the same customer. Since many empirical data can be correlated, this inference result can broaden the applicability of the doubly robust estimator, which has already seen wide adoption in marketing and business research (e.g., Mummalaneni et al. 2022, Ye et al. 2023b, Kim et al. 2023, Cheng et al. 2023, Ellickson et al. 2024).

2. Interference in Recommender System Experiments

In this section, we start with an overview of creator-side recommender experiments and introduce notations to formulate the treatment effect estimation problem. We then discuss how interference arises in the recommendation process. We conclude this section by showing the bias of the commonly applied difference-in-means estimator in our setting.

2.1. Creator-side Recommender Experiments

Recommender systems (RecSys) are designed to serve personalized content for viewers from a large corpus comprised of potentially billions of content items. The overall goal is to score the content items according to their likelihood of engaging the user (e.g., expected viewing time and interactions such as likes and shares). The industry-standard recommendation process is a long pipeline consisting of three main stages: retrieval, ranking, and re-ranking, where the pool of consideration gets decreasingly smaller. In the retrieval stage, the system starts by selecting content items from a vast pool using efficient and scalable algorithms. In the ranking stage, these content items are prioritized with the predicted relevance to the user. Finally, the re-ranking stage fine-tunes the recommendations and often considers other factors such as cold-start or advertising. In our context, the recommender system selects one item in response to each viewer query (e.g., swiping for the next video) instead of presenting an ordered list.

Platforms frequently update their recommender system, which involves anywhere from adjusting algorithm hyper-parameters to implementing a new algorithm. Platforms use experiments to compare against the current benchmark and decide whether to adopt the update. We define several terminologies. *Treatment* denotes an updated recommender system, while *control* is the current status-quo. We use *consideration set* to denote the pool of content for consideration by the recommender system. The treatment vs. control algorithms score content items differently and therefore will lead to different *exposure probabilities* of content items given a consideration set.

The focus of this paper is on creator-side algorithm updates that happen at the re-ranking stage. Because of the focus on creators, platforms frequently rely on creator-side experiments where creators (as opposed to users) are randomized to use the treatment vs. control algorithms for scoring. Because of the focus on the re-ranking stage, the consideration set tends not to be large. However, it can include items from both treatment and control groups that are scored with the corresponding algorithms. One item from the consideration set is exposed to the viewer, and we measure its outcomes from viewer-content interactions, such as view time or likes.

Data from Creator-side Experiments. We now describe the data structure from creatorside experiments and introduce notations. Each observation i is from a viewer query where one item is exposed to the viewer. We observe a tuple $(V_i, \vec{C_i}, \vec{W_i}, k_i^*, Y_i)$, where V_i denotes the viewer and $\vec{C_i} = \{C_{i,1}, \ldots, C_{i,K}\}$ denotes the consideration set of K content items.¹ Content items are randomized into treatment with probability q, or control with probability 1-q, via i.i.d. Bernoulli trials.² $\vec{W_i} = \{W_{i,1}, \ldots, W_{i,K}\}$ indicates the treatment status for each item in the set, with $W_{i,k} = 1$ for treatment and $W_{i,k} = 0$ for control. Content k_i^* is exposed to the viewer, and the outcome variable Y_i measures the viewer's response to the exposed content.

It is useful to clarify what components are affected by the experimental interventions in our framework. The consideration set $\vec{C_i}$ is not affected because we focus on experiments that occur after the retrieval system forms the consideration set. What item gets exposed k_i^* is affected by the experiments. In fact, modeling item exposure conditional on treatment assignment is the focus of the recommender choice model that we introduce in the next section. The experimental interventions impact viewer response Y_i only by affecting which item is exposed. Conditional on exposure, Y_i only depends on the viewer V_i and content C_{i,k^*} pair and not the content's treatment status. We do not consider the temporal interference of viewers and take each viewer query as given.

2.2. Formulate ATE using Policy Value

With creator-side experiments, we introduce notations for treatment effect estimation following the potential outcome framework (Imbens and Rubin 2015). Let $w_c \in \{0, 1\}$ denote the treatment status for content item c.³ Let \mathbf{w}_{-c} collect the treatment statuses of all content items except for c. Let $y(v, c; w_c, \mathbf{w}_{-c})$ denote content c's potential outcome from viewer v with treatment assignment $\mathbf{w} = [w_c, \mathbf{w}_{-c}]$. By convention, we let the potential outcome $y(v, c; w_c, \mathbf{w}_{-c})$ be zero if content c is not recommended to viewer v.

The outcome of interest for content c under treatment assignment (w_c, \mathbf{w}_{-c}) is the expected outcome (or reward) over the distribution of viewer v:

$$r(c; w_c, \mathbf{w}_{-c}) := \mathbb{E}_v \left[y\left(v, c; w_c, \mathbf{w}_{-c}\right) \right].$$

$$\tag{1}$$

¹ It is straightforward to accommodate consideration sets that vary in sizes. We focus on cases with the same size for notation convenience.

 $^{^{2}}$ In practice, randomization typically happens at the creator level with content items inheriting their treatment status. We simplify to content-level randomization for notation convenience, and use creator and content interchangeably in this paper.

 $^{^{3}}$ Our discussion focuses on a binary treatment for notational brevity. The framework can be easily extended to multiple treatments.

Let policy π specify the treatment assignment rule. The policy value $Q(\pi)$ measures the overall outcome across the content set C under treatment assignment policy π (Johari et al. 2022, Goli et al. 2023):

$$Q(\pi) := \mathbb{E}_{\mathbf{w} \sim \pi} \left[\sum_{c \in \mathcal{C}} r(c; w_c, \mathbf{w}_{-c}) \right].$$
⁽²⁾

Let π_1 denote the global treatment policy where all content items use the treatment algorithm $(w_c, \mathbf{w}_{-c}) = \mathbf{1}$ and π_0 denote the global control policy where all items use the control algorithm $(w_c, \mathbf{w}_{-c}) = \mathbf{0}$. The outcome of interest, ATE, is defined as the difference between the policy values of global treatment versus global control.

$$\tau := Q(\pi_1) - Q(\pi_0) = \sum_{c \in \mathcal{C}} \left\{ r(c; \mathbf{1}) - r(c; \mathbf{0}) \right\}.$$
(3)

2.3. Recommender Interference and Bias of DIM Estimator

Interference arises in the recommendation process because content items are "competing for exposure" from the consideration set. Since both treated and control items can appear in the same consideration set, the scoring from the treatment algorithm can influence the exposure probabilities of control content items, and vice versa. As a result, the outcomes of content items depend not only on their own treatment status but also on the treatment status of other items, violating the SUTVA assumption.

Using notations introduced in Section 2.2, interference occurs because the potential outcome for content c, $y(v,c;w_c,\mathbf{w}_{-c})$, depends not only on its treatment status w_c but also on the treatment status of all other content \mathbf{w}_{-c} . Generally, with $\mathbf{w}_{-c}^{(a)} \neq \mathbf{w}_{-c}^{(b)}$, the potential outcomes differ: $y\left(v,c;w_c,\mathbf{w}_{-c}^{(a)}\right) \neq y\left(v,c;w_c,\mathbf{w}_{-c}^{(b)}\right)$. As discussed in Section 2.1, the outcome depends on others' treatment status because they affect the exposure probability of the focal item c.

Because of recommender interference, the standard difference-in-means estimator (DIM) leads to a biased estimate of the treatment effect. Using data from creator-side experiments as described in Section 2.1, the commonly adopted DIM estimator takes the form (Bright et al. 2022, Goli et al. 2023, Johari et al. 2022):

$$\hat{\tau}_{n}^{DIM} := \frac{\sum_{i=1}^{n} \mathbf{1}\left\{W_{i,k_{i}^{*}} = 1\right\} Y_{i}}{nq} - \frac{\sum_{i=1}^{n} \mathbf{1}\left\{W_{i,k_{i}^{*}} = 0\right\} Y_{i}}{n(1-q)},\tag{4}$$

where n is the number of observations. The treatment effect using the DIM estimator, $\hat{\tau}_n^{DIM}$, is the difference in outcomes between exposed items assigned to treatment $(W_{i,k_i^*} = 1)$ relative to those in control $(W_{i,k_i^*} = 0)$. Since the DIM estimator does not consider the "competition" between treated and control items, the estimator does not measure the ground truth ATE of interest, which is the difference in outcomes under global treatment compared to global control (as in Eq. 3).




Figure 1 Illustrative Example of the Bias of the DIM Estimator

We illustrate the bias of the DIM estimator using a simple example. Consider an extreme case where the outcome always equals 1 for the exposed content and 0 otherwise. In such a case, the true ATE is zero because each viewer query always exposes one item and gets an outcome of 1 no matter which algorithm is applied. Suppose the treatment algorithm adds an uplift δ to the item scoring relative to the control. When treated and control units compete for exposure, there will be a boosting effect for treated units when $\delta > 0$, and a suppression effect when $\delta < 0$.

Figure 1 plots the results using this extreme example with the red dotted line denoting the true ATE at zero. We simulate data and compute the DIM estimates with different δ values (detailed simulation settings are described in Section 4.3). The blue line shows the DIM estimates with the 95% confidence intervals. The bias of the DIM estimator can be either positive or negative. When $\delta > 0$, the DIM estimates are positive because treated items are exposed more than control from the boosting effect, despite a true ATE at zero. When $\delta < 0$, the reverse happens with negative DIM estimates when treated items are less likely to be exposed.

This simple example illustrates that the DIM estimator (Eq. 4) does not measure the ATE and the bias can be either positive or negative. We characterize the asymptotic behavior of the DIM estimator as the sample size n grows, and show that it converges to a quantity different than the ATE (Eq. 3). To do so, we introduce an assumption that constrains the number of consideration sets each item appears in, similar to those in prior literature (Viviano 2019, Sävje et al. 2021).

ASSUMPTION 1 (Bounded item appearance). Let n be the sample size. Each item appears in at most a_n consideration sets, with $a_n = o(n^{1/4})$.

This assumption ensures that the number of randomization units (i.e., content items) scales with the sample size. In practical terms, this assumption is often satisfied in real-world applications with creator-side experiments. These experiments typically target items produced by small creators or for advertising purposes, where the number of consideration sets they appear in is relatively small compared to the overall sample size.

PROPOSITION 1 (Convergence of the difference-in-means estimator). Suppose Assumption 1 holds. Under the creator-side randomization with treated probability $q \in (0,1)$, the difference-in-means estimator $\hat{\tau}_n^{DIM}$ converges in probability to a quantity $\tau^{\mathcal{B}(q)}$,

$$\hat{\tau}_{n}^{DIM} \xrightarrow{p} \tau^{\mathcal{B}(q)} \quad where \quad \tau^{\mathcal{B}(q)} := \mathbb{E}\left[\sum_{c \in \mathcal{C}} \left\{ r(c; w_{c} = 1, \mathbf{w}_{-c} \sim \mathcal{B}(q)) - r(c; w_{c} = 0, \mathbf{w}_{-c} \sim \mathcal{B}(q)) \right\} \right], \quad (5)$$

with $\mathbf{w}_{-c} \sim \mathcal{B}(q)$ denoting that the treatment status of all content items except for c is sampled from a Bernoulli distribution with a treatment probability of q.

Detailed proof is given in Appendix A. The quantity $\tau^{\mathcal{B}(q)}$ is often referred to as the direct effect of the treatment under creator-side randomization (Sävje et al. 2021, Hu et al. 2021). The difference between $\tau^{\mathcal{B}(q)}$ and the ATE τ is the interference bias. Prior research has established the bias of the DIM estimator in other contexts, e.g., two-sided matching platforms such as ride-sharing (Bright et al. 2022). Proposition 1 derives the convergence results of the DIM estimator in creator-side randomized experiments.

3. Modeling Interference

In this section, we introduce a recommender choice model to explicitly account for interference during the recommendation phase. This semi-parametric model incorporates neural networks in the choice model in order to flexibly capture personalized recommendations based on diverse viewer preferences and content characteristics. We use counterfactual analysis for treatment effect estimation. We conclude this section by discussing the scope of the proposed approach: what types of interference it does and does not capture.

3.1. Recommender Choice Model

Upon a viewer query, the goal of the recommender system is to choose the most suitable content item for viewer V_i from the consideration set $\vec{C_i}$. Inspired by the concept of *model distillation* (Hinton et al. 2015), we represent the complex recommender system using a choice model, which describes how the recommender system "chooses" an item from the consideration set.

The score or "utility" of content $C_{i,k}$ for viewer V_i takes the form:

$$S_{i,k} = s_0(V_i, C_{i,k}) + W_{i,k} \cdot s_1(V_i, C_{i,k}) + \epsilon_{i,k}.$$
(6)

The baseline score function $s_0(\cdot, \cdot)$ approximates the control algorithm's evaluation of content given viewer V_i and content $C_{i,k}$. For items in the treatment group $(W_{i,k} = 1)$, $s_1(\cdot, \cdot)$ captures the score uplift from the treatment algorithm's evaluation of content $C_{i,k}$ for viewer V_i .⁴ We parameterize the score functions $s_0(\cdot, \cdot)$ and $s_1(\cdot, \cdot)$ as neural networks. Neural networks can flexibly capture the relationship between recommender scoring and the viewer-content pairs, which effectively approximate personalized recommendations for different viewers (Covington et al. 2016).

The error term $\epsilon_{i,k}$ represents the part of content evaluation that score functions s_0 and s_1 do not capture. It accounts for the inherent randomness in recommender systems as well as the approximation errors of these neural networks in representing the actual recommender pipeline.⁵ Randomness in the recommendations can arise from concurrent orthogonal experiments, a common practice among tech companies (Ivaniuk and Duan 2020), as well as system-wide uncertainties, such as viewers being assigned to less sophisticated algorithms in order to reduce latency during peak hours (Taylor and Wexler 2003). We assume that the error term $\epsilon_{i,k}$ follows an i.i.d. Type 1 Extreme Value distribution, and the item with the highest score or "utility" in the consideration set will be exposed to viewer V_i .

With these assumptions, we can write the choice probability using the multinomial logit choice model. The recommender choice model specifies the probability of an item k to be recommended for viewer V_i given consideration set $\vec{C_i}$ and treatment allocation $\vec{W_i}$:

$$P_{i,k}\left(V_i, \overrightarrow{C_i}, \overrightarrow{W_i}; s_0, s_1\right) := \mathbb{P}\left(k_i^* = k\right) = \frac{e^{s_0(V_i, C_{i,k}) + W_{i,k} \cdot s_1(V_i, C_{i,k})}}{\sum_{k'=1}^{K} e^{s_0(V_i, C_{i,k'}) + W_{i,k'} \cdot s_1(V_i, C_{i,k'})}}.$$
(7)

The choice model explicitly models interference in the recommendation process. The exposure probability of an item is affected by not only its own treatment status $W_{i,k}$ but by the treatment status of all others in the consideration set. The recommender choice model serves as the foundation to estimate counterfactual exposure probability under alternative treatment allocation policy, which we describe in detail in Section 3.3.

The recommender choice model is semi-parametric in that it combines the structural choice model with flexible neural networks. Doing so allows us to enjoy the benefits of both. On the one hand, leveraging the choice model enables the treatment effect estimation via counterfactuals under alternative treatment allocations. This would not be feasible with a fully non-parametric black-box model. On the other hand, neural networks can capture flexible relationships so that the recommender choice model better approximates the behavior of the real-world complex recommender system. Employing neural networks also helps mitigate the independence of irrelevant alternatives

⁴ One alternative model specification is to let the score functions s_0 and s_1 depend not only on the focal content item $C_{i,k}$ but the entire consideration set \vec{C}_i . Under the alternative specification, our modeling and inference framework still apply but the computational cost will significantly increase. We explore this alternative in the Appendix G.

⁵ It is also worth noting that it is impractical to simulate outcomes with "offline" recommender systems. In other words, one cannot simulate recommendations and predict viewer response under treatment or control algorithms respectively (Bennett et al. 2007).

(IIA) issue inherent in a standard logit choice model, leading to a proportional substitution pattern. Since neural networks can flexibly account for viewer preferences, adding a new item may disproportionately affect the exposure probability of certain items more than others, depending on content similarity and viewer preferences.

With the score functions parameterized as neural networks, we need an assumption on their boundedness so that the exposure probability of an item is bounded away from zero. We consider universally bounded scores throughout.

ASSUMPTION 2 (Bounded scores). There exists a universal constant C such that the true score functions are bounded: $||s_0||_{\infty} \leq C$ and $||s_1||_{\infty} \leq C$.

This assumption is parallel to the "overlap" condition in propensity score-based debiasing methods like AIPW (Imbens 2004). Under this assumption, we show that each item in the consideration set has some positive probability of being recommended, regardless of the treatment assignment. We formalize this with Lemma 1. The proof is given in Appendix E.1.

LEMMA 1. Under Assumption 2, there exists a universal constant $\delta > 0$ such that each content item in the consideration set has at least δ probability to be recommended:

$$\mathbb{P}\left(k_i^* = k \mid V_i, \overrightarrow{C_i}, \overrightarrow{W_i}; s_0, s_1\right) \ge \delta, \quad \forall \ (V_i, \overrightarrow{C_i}, \overrightarrow{W_i}, k).$$

It is worth mentioning that the recommender choice model can bring standalone practical value for platforms beyond enabling ATE estimation. It represents a recommender distillation process that can potentially replace the complex recommender system to reduce latency when serving viewers online (Tang and Wang 2018). Moreover, the derived score functions approximate the overall value of content items to viewers, which can serve as a useful offline evaluation metric.

3.2. Viewer Response Model

To complete counterfactual evaluations under alternative treatment allocation, we also need a viewer response model that predicts the viewer response when exposed to an item. The viewer's response reflects outcomes that the platform is interested in evaluating, such as view time, likes, or conversion rates. We assume the viewer's response depends on the viewer characteristics V_i and the characteristics of the exposed item C_{i,k_i^*} . Let the viewer's response Y_i be:

$$Y_i = z(V_i, C_{i,k_i^*}) + \zeta_i. \tag{8}$$

 $z(\cdot, \cdot)$ is a viewer response function that we parameterize as a neural net. It predicts a flexible relationship between the input viewer-content pairs and the outcome Y_i . The error term ζ_i is i.i.d. noise. In general, one needs a separate viewer response model for each outcome of interest, but the recommender choice model is commonly shared across different outcomes.

The viewer response model in Eq. (8) assumes that the outcome does not depend on the treatment status of the item conditional on the item being exposed. This is a reasonable assumption since viewers are typically not aware of changes in recommendation algorithms. It also assumes that the outcome does not depend on any previous content the viewers may have seen. The model can be extended to allow previous content interactions as input in addition to viewer characteristics, but doing so will lead to a significantly more complex viewer response model.

In practice, platforms typically have pre-trained viewer response predictions from backend algorithms, which is also the case in our empirical context. Thus, we can directly leverage these preexisting viewer response models and only need to train the recommender choice model. It's worth noting that our framework does not require an unbiased prediction for these viewer responses. As detailed in Section 4, we introduce a debiased estimator with a bias correction term to account for "errors" in the nuisance predictions.

3.3. Counterfactual Analysis for Policy Value and Treatment Effect

With the recommender choice model (Eq. 7) and the viewer response model (Eq. 8), we can represent the policy values (Eq. 2) of any counterfactual treatment assignment policy π as follows (see Appendix B for detailed derivation):

$$Q(\pi) = \mathbb{E}_{(V_i, \vec{C}_i, \vec{W}_i \sim \pi)} \left[\sum_{k=1}^K z(V_i, C_{i,k}) \cdot P_{i,k} \left(V_i, \vec{C}_i, \vec{W}_i; s_0, s_1 \right) \right].$$
(9)

The term in the square bracket captures the expected outcomes of viewer V_i by multiplying the exposure probability and the viewer response conditional on exposure for each item in the consideration set. The policy value $Q(\pi)$ measures the overall outcome by taking the expectation over the distribution of viewer V_i , consideration set \vec{C}_i , and treatment assignment policy \vec{W}_i .

To estimate the treatment effect, we are particularly interested in two specific treatment assignment policies, with all items in the treatment status or all in control. The ATE (Eq. 3) is the difference between the policy values under global treatment $Q(\pi_1)$ vs. global control $Q(\pi_0)$. Using Eq. (9), the ATE can be written as:

$$\tau = Q(\pi_1) - Q(\pi_0) = \mathbb{E}_{(V_i, \vec{C}_i)} \left[\sum_{k=1}^K z(V_i, C_{i,k}) \cdot \left(P_{i,k}(V_i, \vec{C_i}, \vec{W_i} = \mathbf{1}) - P_{i,k}(V_i, \vec{C_i}, \vec{W_i} = \mathbf{0}) \right) \right], \quad (10)$$

where $P_{i,k}(V_i, \overrightarrow{C_i}, \overrightarrow{W_i} = \mathbf{1}) - P_{i,k}(V_i, \overrightarrow{C_i}, \overrightarrow{W_i} = \mathbf{0})$ represents the change in the exposure probability for content item $C_{i,k}$ when going from global control to global treatment scenarios.

Besides looking at the average effect across the whole population, it can be useful for the platform to understand the heterogeneous impact of a new algorithm on different types of creators. Using our framework, it is straightforward to conduct subgroup analysis to estimate heterogeneous treatment effects on different creators. For a particular subgroup C_0 , the corresponding average treatment effect can be calculated as:

$$\tau_{\mathcal{C}_0} = \mathbb{E}_{(V_i,\vec{\mathcal{C}}_i)} \left[\sum_{k=1}^{K} I\left\{ C_{i,k} \in \mathcal{C}_0 \right\} \cdot z(V_i, C_{i,k}) \cdot \left(P_{i,k}(V_i, \vec{C}_i, \vec{W}_i = \mathbf{1}) - P_{i,k}(V_i, \vec{C}_i, \vec{W}_i = \mathbf{0}) \right) \right].$$
(11)

This is close to Eq. (10) except that instead of adding up the expected change in outcomes for all content items in the consideration set, we only do so for the ones that belong in the subgroup of interest, $I \{C_{i,k} \in C_0\}$.

Beyond ATE, one can calculate other outcomes of interest with the estimated model. As a simple example, although Eq. (10) defines the treatment effect as the level difference between treatment and control conditions, $Q(\pi_1) - Q(\pi_0)$, it is just as easy to calculate the treatment effect in relative terms, $\frac{Q(\pi_1)-Q(\pi_0)}{Q(\pi_0)}$. As a more involved example, the model can simulate outcomes as the platform scales up the proportion of traffic allocated to the treatment algorithm. In other words, the framework is flexible to account for any treatment assignment policy π beyond the global treatment and global control.

3.4. Scope of the Proposed Approach

By directly modeling the interference pathway through a recommender choice model, the proposed approach captures interference that arises from items competing for exposure. Since it relies on modeling the exposure probability from a consideration set, the framework is more applicable to experiments that happen in the later stage of the recommender system where the size of the consideration set is relatively small (see Section 2.1). For an experiment that happens in the early stage (e.g., retrieval stage), it will be intrinsically harder to model exposure from a very large pool of items. Since many recommender experiments involve fine-tuning in the later stages, the proposed approach is applicable to many such experiments in practice.

Our study does not explicitly consider potential viewer-side temporal interference, where viewer responses might be influenced by previous content. For example, suppose the treatment algorithm prioritizes high quality content, viewers may be motivated to stay longer with the platform (hence generating more viewer queries). Viewer responses may also be impacted by prior content because the diversity of recommended content can influence viewer behavior (e.g., novelty effect). Addressing these forms of interference requires more nuanced modeling of viewer behavior, such as using reinforcement learning to model viewer behavior as Markov chain processes to deal with viewer-side temporal interference (Farias et al. 2023).

Our study also abstracts away from the potential long-run effect of the treatment. For example, if a treatment algorithm prioritizes certain types of content, it may create a feedback loop that impacts the content production decisions of different creators. In the long term, it may lead to a different pool of available content on the platform. Addressing this involves modeling the content production decisions based on short-term outcomes (e.g., views and likes) that may vary under different algorithms. Our approach focuses on these short-term outcomes by taking the available content and viewer queries as given.

4. Estimation and Inference Procedure

In this section, we describe the estimation and inference procedure of the proposed model. We start with the estimation of the neural network components in the recommender choice and viewer response models. We then construct a debiased estimator for treatment effect estimation and inference. Using Monte Carlo simulations, we show that the proposed approach can effectively recover the true point estimate and uncertainty.

4.1. Estimating the Nuisance Components

We start by describing the estimation of the nuisance components in the recommender choice model, which relies on two score functions: s_0 denotes the baseline score and s_1 is the score uplift from the treatment algorithm. Both score functions are parameterized as neural networks.

Recall that for each viewer query i, we observe that viewer V_i with consideration set $\vec{C_i}$ and treatment status $\vec{W_i}$ gets recommended the k_i^{*th} item, C_{i,k_i^*} . Similar to a fully parametric choice model, estimating the choice model with neural networks relies on maximizing the likelihood function or equivalently, minimizing the loss function. The loss function for any $\tilde{s}_0, \tilde{s}_1 \in \mathcal{F}_s$, where \mathcal{F}_s is the function class for s_0 and s_1 , can be written as follows:

$$\ell_{1}(V_{i}, \vec{C_{i}}, \vec{W_{i}}, k_{i}^{*}; \tilde{s}_{0}, \tilde{s}_{1}) := -\log\left(\mathbb{P}\left(k_{i}^{*} \mid V_{i}, \vec{C_{i}}, \vec{W_{i}}; \tilde{s}_{0}, \tilde{s}_{1}\right)\right)$$

$$= -\left(s_{0}(V_{i}, C_{i, k_{i}^{*}}) + W_{i, k_{i}^{*}} \cdot s_{1}(V_{i}, C_{i, k_{i}^{*}})\right) + \log\left(\sum_{k=1}^{K} e^{s_{0}(V_{i}, C_{i, k}) + W_{i, k} \cdot s_{1}(V_{i}, C_{i, k})}\right).$$

$$(12)$$

This is commonly known as the cross entropy loss. During the training process, the parameters of the neural networks are updated so that we get the estimated (\hat{s}_0, \hat{s}_1) that minimizes the cross entropy loss:

$$(\hat{s}_0, \hat{s}_1) \in \underset{\tilde{s}_0, \tilde{s}_1 \in \mathcal{F}_s}{\operatorname{arg\,min}} \ \frac{1}{n} \sum_{i=1}^n \ell_1\left(V_i, \overrightarrow{C_i}, \overrightarrow{W_i}, k_i^*; \tilde{s}_0, \tilde{s}_1\right).$$
(13)

Intuitively, the score functions \hat{s}_0 and \hat{s}_1 are learned so that the recommender choice model can best approximate the behavior of the actual recommender system.⁶

We then describe the estimation of the viewer response model. Recall that when viewer V_i gets exposed to content item C_{i,k_i^*} , we observe the outcome Y_i . The role of the viewer response model is

 $^{^{6}}$ We discuss the identification of the nuisance components in Appendix D.

to approximate the outcome conditional on the viewer and item pair. This is a standard machine learning task with the loss function determined by the outcome types. For a continuous outcome, the loss function for any $\tilde{z} \in \mathcal{F}_z$, where \mathcal{F}_z is the function class for z, can be written as follows:

$$\ell_2(V_i, C_{i,k_i^*}, Y_i; \tilde{z}) = \left(\tilde{z}(V_i, C_{i,k_i^*}) - Y_i\right)^2.$$
(14)

This is commonly known as the mean square error loss. For a categorical outcome, one can use the cross entropy loss. With either type of loss function, we get the estimated \hat{z} by minimizing the loss:

$$\hat{z} \in \underset{\tilde{z} \in \mathcal{F}_z}{\operatorname{arg\,min}} \frac{1}{n} \sum_{i=1}^n \ell_2(V_i, C_{i,k_i^*}, Y_i; \tilde{z}).$$

$$(15)$$

In practice, platforms often have built models to predict key outcomes of interest. These model predictions are useful for the recommender pipeline since they capture key behaviors of interest for the platform, e.g., watch time and likes. In our empirical application (described in Section 5), we leverage the platform's pre-trained view response models instead of estimating \hat{z} ourselves.

4.2. Debiased Estimator

We now discuss the estimation and inference procedure for the treatment effect. With the estimated recommender choice and viewer response models, one can directly plug in the estimated neural networks $(\hat{s}_0, \hat{s}_1, \hat{z})$ into the ATE formulation in Eq. (10). For each observation with viewer and consideration set pair $(V_i, \overrightarrow{C_i})$, the direct plug-in estimate μ is:

$$\mu(V_i, \overrightarrow{C_i}; \hat{s}_0, \hat{s}_1, \hat{z}) = \sum_{k=1}^{K} \hat{z}(V_i, C_{i,k}) \left\{ \frac{e^{\hat{s}_0(V_i, C_{i,k}) + \hat{s}_1(V_i, C_{i,k})}}{\sum_{k'=1}^{K} e^{\hat{s}_0(V_i, C_{i,k'}) + \hat{s}_1(V_i, C_{i,k'})}} - \frac{e^{\hat{s}_0(V_i, C_{i,k})}}{\sum_{k'=1}^{K} e^{\hat{s}_0(V_i, C_{i,k'})}} \right\}$$

If the components (s_0, s_1, z) are fully parametric, it is straightforward to use μ for treatment effect estimation and inference. With these components being neural networks (or other non-parametric models), the direct plug-in is generally not \sqrt{n} -consistent.

We write down a debiased (DB) estimator for treatment effect following Farrell et al. (2020). The debiased estimator adds a debiasing term to the direct plug-in estimate μ . For each observation, the debiased estimate ψ is:

$$\psi(V_{i}, \vec{C_{i}}, \vec{W_{i}}, k_{i}^{*}, Y_{i}; \hat{s}_{0}, \hat{s}_{1}, \hat{z}, \hat{H}) = \mu(V_{i}, \vec{C_{i}}; \hat{s}_{0}, \hat{s}_{1}, \hat{z})$$

$$-\nabla \mu(V_{i}, \vec{C_{i}}; \hat{s}_{0}, \hat{s}_{1}, \hat{z})^{T} \hat{H}(V_{i}, \vec{C_{i}}; \hat{s}_{0}, \hat{s}_{1}, \hat{z})^{-1} \nabla \ell(V_{i}, \vec{C_{i}}, \vec{W_{i}}, k_{i}^{*}, Y_{i}; \hat{s}_{0}, \hat{s}_{1}, \hat{z}).$$

$$(16)$$

The debiased term has three components. The first component $\nabla \mu$ is the gradient of the direct plug-in estimate μ with respect to the nuisances $(\hat{s}_0, \hat{s}_1, \hat{z})$, which captures the sensitivity of the estimate to the nuisance estimation. Let ℓ be the total loss function which combines the loss ℓ_1 from the recommender choice model and ℓ_2 from the viewer response model. The second component \hat{H} estimates the expected Hessian of the loss function ℓ with respect to the nuisances, with the expectation marginalizing over treatment assignment $\overrightarrow{W_i}$ under the creator-side randomization. The third component $\nabla \ell$ is the gradient of the loss function ℓ with respect to the nuisances, which is scaled by the curvature of the loss function (the second component). We give the explicit expressions of $(\nabla \mu, H, \nabla \ell)$ in Appendix C. The Hessian H is invertible with Assumption 2 (bounded scores), with formal results and proofs in Appendix E.2.

The debiased (DB) estimator $\hat{\tau}_n^{DB}$ is the average of ψ across all observations:

$$\hat{\tau}_{n}^{DB} := \frac{1}{n} \sum_{i=1}^{n} \psi(V_{i}, \overrightarrow{C_{i}}, \overrightarrow{W_{i}}, k_{i}^{*}, Y_{i}; \hat{s}_{0}, \hat{s}_{1}, \hat{z}, \hat{H}).$$
(17)

With ψ and $\hat{\tau}_n^{DB}$, the estimated variance can be calculated as below. The standard error of the DB etimator $\hat{\tau}_n^{DB}$ can be estimated as $n^{-1/2} \hat{V}_n^{1/2}$.

$$\widehat{V}_{n} = \frac{1}{n} \sum_{i=1}^{n} \left(\psi(V_{i}, \overrightarrow{C_{i}}, \overrightarrow{W_{i}}, k_{i}^{*}, Y_{i}; \hat{s}_{0}, \hat{s}_{1}, \hat{z}, \hat{H}) - \hat{\tau}_{n}^{DB} \right)^{2}.$$
(18)

The debiased estimator $\hat{\tau}_n^{DB}$ satisfies the universal Neyman orthogonality with respect to the nuisances. We provide formal results in Appendix E.3. The Neyman orthogonality property ensures that errors in estimating nuisances, e.g., \hat{s}_0 and \hat{s}_1 , impact the treatment effect estimation, $\hat{\tau}_n^{DB}$, only to a second-order degree. This is a key property that effectively mitigates potential bias in the estimator from inaccuracies in nuisance estimation.

Now we present the main estimation and inference guarantees for the debiased estimator $\hat{\tau}_n^{DB}$. Under mild conditions, we show that $\hat{\tau}_n^{DB}$ is \sqrt{n} -consistent and asymptotically normal. The proof builds on Chernozhukov et al. (2018), which deals with i.i.d. sample. In our empirical application, however, the data generating process does not yield an i.i.d. sample, even when each viewer query *i* is treated as independent. This is because for each new observation, if its consideration set overlaps with previous ones, the treatment statuses of the overlapped content items become deterministic. In contrast, for content items without overlap with previous observations, the treatment status for each item follows i.i.d. Bernoulli randomized trials. We model the entire data generating process in a sequential manner and leverage the martingale limiting theorems to characterize the asymptotic behavior for our estimator.

THEOREM 1. Suppose that Assumptions 1 & 2 hold. Assume that the data generating process follows the recommender choice model Equation (7) and the viewer response model Equation (8). Suppose that the nuisance estimates are all bounded by the constant C in Assumption 2 and satisfy the convergence rate: $\|\hat{s}_0 - s_0\|_{L_2} + \|\hat{s}_1 - s_1\|_{L_2} + \|\hat{z} - z\|_{L_2} = o(n^{-1/4})$ and $\|\hat{H} - H\|_{L_2} = o(n^{-1/4})$. Then the debiased estimator $\hat{\tau}_n^{DB}$ in (17) is \sqrt{n} -consistent with $\hat{\tau}_n^{DB} - \tau = O_p(n^{-1/2})$ and asymptotically normal with: $\sqrt{n}(\hat{\tau}_n^{DB} - \tau) / \sqrt{\hat{V}_n} \Rightarrow \mathcal{N}(0, 1)$. The detailed proof for Theorem 1 is provided in Appendix F. At a high level, the proof establishes the asymptotic properties of the debiased estimator $\hat{\tau}_n^{DB}$ by comparing it to an oracle estimator $\tilde{\tau}_n^{DB}$, which assumes the nuisances are known exactly without estimation error. The oracle estimator forms a martingale difference sequence and converges to a normal distribution. Due to Neyman othogonality, the difference between the debiased estimator and the oracle estimator is shown to be negligible, with this difference bounded effectively.

One last detail to note is that, in practice, we use the standard cross-fitting procedure when estimating nuisance components. The idea is to use one subgroup of the data to estimate the nuisance models and use the remaining ones to carry out treatment effect estimation (Chernozhukov et al. 2018, Farrell et al. 2020). Let K be the number of subgroups. We estimate the nuisance models using data excluding group k, $(\hat{s}_0^{(-k)}, \hat{s}_1^{(-k)}, \hat{z}^{(-k)}, \hat{H}^{(-k)})$. Using the estimated nuisance models, we estimate the treatment effect on data from group k, $\hat{\tau}_n^{(k)}$. The cross-fitted ATE estimator can be calculated as the average of the K subgroups: $\hat{\tau}_n^{DB} = \frac{1}{K} \sum_{k=1}^K \hat{\tau}_n^{(k)}$. The variance of the estimator can be calculated in the same way: $\hat{V}_n = \frac{1}{K} \sum_{k=1}^K \hat{V}_n^{(k)}$.

4.3. Monte Carlo Simulations


We use Monte Carlo simulations to validate the estimation and inference results as shown in Theorem 1. We simulate data for n = 1000 viewers with K = 3 items in the consideration sets that are drawn from m = 500 content items. The feature of each content item is characterized in two parts. The first part is a two-dimensional feature vector drawn from uniform distribution, $C_{1,i,k} \sim \mathcal{U}(0,1)^2$. The second part is a binary feature indicating the eligibility of receiving the intervention (e.g., only advertising videos will be boosted under the treatment), $C_{2,i,k} \sim \text{Bernoulli}(0.5)$. Viewers have a two-dimensional feature vector drawn from uniform distribution, $V_i \sim \mathcal{U}(0,1)^2$.

The baseline utility score s_0 is calculated as the inner product between viewer and item feature vectors, $s_0(V_i, C_{i,k}) = V_i \cdot C_{1,i,k}$. The treatment algorithm changes the score of eligible videos when $C_{2,i,k} = 1$ by δ , $s_1(V_i, C_{i,k}) = C_{2,i,k} \cdot s_0 \cdot \delta$. We simulate two scenarios, one with $\delta = 1$ to mimic a boosting treatment, and the other $\delta = -0.5$ to mimic a suppressing treatment. For each viewer *i*, one video is chosen based on the choice model as in Eq. (7). Let the outcome for each view-content pair be the baseline score plus a random Gaussian noise, $Y_i = s_0 + \zeta_i$, where $\zeta_i \sim \mathcal{N}(0, 0.01)$.

We estimate the treatment effects using the proposed debiased (DB) estimator $\hat{\tau}_n^{DB}$ as well as the difference-in-means (DIM) estimator $\hat{\tau}_n^{DIM}$ under the boosting and suppressing scenarios. For each of the 100 Monte Carlo datasets, we calculate the point estimate and standard error. For the DB estimator, the standard error is $n^{-1/2} \hat{V}_n^{1/2}$ as introduced in Theorem 1. We employ 3-fold cross-fitting for nuisance estimation. For the DIM estimator, the standard errors are calculated as the standard deviation of the sample estimates.⁷

⁷ Let $\sigma(a_i)$ denote the sample standard deviation of samples $\{a_i\}$: $\sigma^2(a_i) = n^{-1} \sum_{i=1}^n (a_i - n^{-1} \sum_{i=1}^n a_i)^2$. For the DIM estimator $\hat{\tau}_n^{DIM}$, the standard error estimate is $n^{-1/2} \sigma\left(\left(W_{k_i^*}/q - (1 - W_{k_i^*})/(1 - q)\right)Y_i\right)$.

Figure 2 plots the standardized distribution of the estimates from $\hat{\tau}_n^{DB}$ and $\hat{\tau}_n^{DIM}$. More specifically, we compute the standardized estimates $(\hat{\tau} - \tau)/\hat{\sigma}$, where τ is the true ATE, $\hat{\tau}$ is the point estimate and $\hat{\sigma}$ is the estimated standard error. Essentially, the bias of the estimator is normalized by the estimated standard error. The distribution of these standardized estimates should approximate the standard normal distribution, denoted by the dashed black curve. We see that the distribution from the proposed DB estimator is very close to the standard normal in both cases. The results suggest that the proposed DB estimator is unbiased and asymptotically normal, and thus allows for accurate ATE estimation and inference. The DIM estimator, on the other hand, can either be biased upwards (when $\delta = 1$) or biased downwards (when $\delta = -0.5$).



Note: $(\hat{\tau} - \tau)/\hat{\sigma}$, where $\hat{\tau}$ is the estimate, τ is the true ATE, and $\hat{\sigma}$ is the estimated standard error.

Figure 3 further examines the point estimates and estimated standard errors. The left panel plots the average point estimates, with the horizontal dashed line denoting the truth values, and the 95% confidence intervals computed from the estimated standard errors. As shown in Figure 2, the DB estimator is unbiased while the DIM estimator can have a large bias in either direction. The right panel compares the estimated standard errors with the true uncertainty, measured by Monte Carlo standard errors, alongside the 45-degree dashed line. The DB estimator correctly estimates the uncertainty, whereas the standard error for DIM estimator is underestimated. This occurs because DIM fails to account for the sample correlation caused by overlapping items across consideration sets, which inherit the same treatment status rather than being i.i.d. The DB estimator correctly estimates the standard error since it shares the asymptotic property of the oracle estimator due to its Neyman orthogonality property, as discussed in Section 4.2.




Figure 3 Monte Carlo Results: Point Estimate and Standard Error

5. Empirical Application

In this section, we evaluate the proposed approach using real experimental data. We estimate treatment effects using data from a creator-side field experiment on Weixin's short video platform. As a benchmark, we carry out a blocked double-sided experiment without interference. We find that the ATE estimates from the DB estimator are close to the benchmark, while the DIM estimates are further away from the benchmark, sometimes even having the reversed sign.

5.1. Experimental Design

We apply the proposed debiased estimator to analyze the treatment effect of applying a new recommendation algorithm, relative to the current status quo algorithm, on Weixin's short video platform. The treatment affects the recommended videos by changing their exposure probabilities during the re-ranking stage. Therefore this experiment is a good fit to apply our proposed method because of the relatively small consideration set (see Section 3.4).

Unlike in Monte Carlo simulations, there is no longer a ground truth to compare against in real empirical application. To obtain a benchmark approximating the true treatment effect, we implement a costly double-sided experiment alongside the typical creator-side experiment. The experimental design is summarized in Figure 4. We create three separate experimentation universes. Universe A is used to conduct the creator-side experiment where half of the creators are randomly assigned to use the treatment algorithm and the other half use the control algorithm. We use data from this universe to estimate the treatment effect using the proposed debiased (DB) estimator as well as the difference-in-means (DIM) estimator.

The double-sided randomized experiment is implemented in Universe B (global treatment) and Universe C (global control). Since treated vs. control videos will never appear in the same consideration set, there is no interference coming from the treated vs. control groups competing for exposure. We use data from the double-sided experiment to proxy the ground truth treatment




Figure 4 Experimental Design

effect (Bajari et al. 2021, Johari et al. 2022, Ye et al. 2023a). More specifically, the double-sided (DS) estimator compares the average creator outcomes between the global treatment and global control universes. While effective, the double-sided experimental design is costly to implement and, therefore, not practical for evaluating all creator-focused experiments at scale.


The experiments run for eight days in total. They are carried out simultaneously in the three universes to avoid any time-varying differences in outcomes (e.g., weekends or holidays). The three universes are equal-sized to mitigate any potential impact of market size. They each have about 34,000 videos and a similar number of viewers, with the exact number of viewers omitted for confidentiality reasons.

5.2. Estimate the Recommender Choice Model

To construct the proposed debiased (DB) estimator, we start by estimating a recommender choice model using data from the creator-side randomized experiment. This choice model can then be utilized to estimate the treatment effects for all three outcomes. Below, we provide details on how to estimate the recommender choice model and show that it aligns well with the behavior of the actual recommender system.

We start by defining the consideration set. This is done by empirically identifying the smallest possible consideration set prior to the treatment. Figure 5 plots the eventual exposure probability of items based on their "pre-treatment" ranking by the recommendation algorithm. The exposure probability is the highest for the top-ranked item and quickly goes down to close to 0. We choose a consideration set of K = 15 items, based on the empirical observation that videos outside this range are unlikely to be exposed under any treatment assignment. Thus, these consideration sets capture nearly all relevant videos that have the potential to be exposed.

We then describe the network structure to fit the recommender choice model. The model predicts the choice probability of each item in the consideration set given the data $(V_i, \vec{C}_i, \vec{W}_i)$. Figure 6



provides an illustration. The data inputs to the network are the viewer and content embeddings $(V_i, C_{i,k})$. In practice, we do not directly use the raw embeddings. Instead, we leverage the platform's pre-trained models to evaluate each viewer-content pair. These models are trained pre-experiment and target different engagement metrics, such as view time and likes. The outputs from the pre-trained models go through two fully connected hidden layers. These learned layers get updated through the training process. The score layer has two nodes, one for the baseline score s_0 and the other for the treatment uplift s_1 . The model layer constructs the exposure probability of each item in the consideration set following Eq. (7). Note that treatment assignment \vec{W}_i enter as input in the model layer. We use Python's TensorFlow package to train the model with categorical cross-entropy loss following Eq. (12).

We assess how well the fitted recommender choice model aligns with the actual recommended items observed in the data. Figure 7 plots the ROC curve (receiver operating characteristic curve), with the true positive rate and the false positive rate at different thresholds. The dashed diagonal line represents a classifier with no discriminative power, resulting in an AUC of 0.5 (area under the curve). The fitted model achieves an AUC of 0.97, which indicates that the fitted model can predict the actual recommended items very well. By leveraging the pre-trained models, we can achieve accurate predictions without a very complex neural network.

5.3. Estimate the Treatment Effects

Besides the fitted recommender choice model, we need viewer response models to estimate the treatment effect. We leverage the platform's existing viewer response models, \hat{z} . We track three outcome variables, all measuring binary outcomes.⁸ They share the same recommender choice model but have separate viewer response models, each one to predict its own outcome of interest.

 $^{^{8}}$ The meaning of these outcomes is omitted for confidentiality reasons.





Figure 6 Network structure of the recommender choice model

Figure 7 ROC Curve of the Fitted Recommender Choice Model



To construct the debiased term, we also need to estimate the expected Hessian matrix of the loss function ℓ , with the expectation taken over the treatment assignment. This is done by drawing many realizations of the treatment assignments from the same creator-side treatment assignment rule, and calculating the empirical mean of the Hessian from these data samples (we use 500 samples).

The results from the double-sided estimator $\hat{\tau}^{DS}$, leveraging the double-sided randomized experiment, are taken to approximate the ground truth without interference. The treatment effects measure how the outcomes change under the new recommendation system (treatment) compared




Figure 8 ATE estimation for four metrics.

to the current status quo (control). We compare these benchmark measures with results using our debiased estimator $\hat{\tau}^{DB}$ and the difference-in-means estimator $\hat{\tau}^{DIM}$. We do so for each of the three outcome variables.

Figure 8 presents the results. For outcome 1, the benchmark DS estimator indicates a close-tozero and insignificant treatment effect. The estimate by the proposed DB estimator is also closeto-zero and insignificant. The estimate by the DIM estimator, however, is positive and significant. For outcome 2, the benchmark DS estimator indicates a negative and significant treatment effect. The estimate from the DB estimator is also negative and significant, and close in magnitude to the benchmark. The estimate from the DIM estimator has a reversed sign and is positive and significant. For outcome 3, the benchmark DS estimator indicates a positive and significant estimate. Both the DB estimator and the DIM estimator provide estimates that are close in magnitude.

Across the three outcome variables, the DB estimator produces estimates that are close to the benchmark DS estimator. In contrast, the commonly applied DIM estimator yields significantly different treatment effect estimates. Relying on the DIM estimator can lead to incorrect decisions. For example, to improve outcome 2, the DIM estimator suggests adopting the new recommendation system, even though the current one gives a higher measure. The data from the real experiment, together with a double-sided experimental design, validate the performance of the proposed debiased estimator in treatment effect estimation from creator-side experiments.

6. Conclusion

This paper identifies a common source of interference in randomized experiments that arises from treated and control units competing for exposure. This occurs in online content sharing platforms when running creator-side recommender experiments or other interventions targeting creators. We show that the commonly applied difference-in-means estimator can have either an upward or downward bias. We introduce a novel approach that directly models the interference pathway using data from creator-side experiments. The method combines a structural choice model and nonparametric neural networks to account for rich heterogeneity between viewers and content.

We show that the estimator is \sqrt{n} -consistent and asymptotically normal. Our inference results extend the applicability of prior results to potentially correlated samples. We validate the proposed estimator using experimental data on a leading short-video platform, with a costly blocked doublesided randomization without interference as a benchmark. Results from the proposed debiased estimator are very close to the benchmark, while the DIM estimates can be quite far away from the benchmark, even with the reversed sign. More broadly, future work can leverage similar semiparametric methods by combining structural and machine learning models to solve other challenges ranging from demand estimation to causal inference in diverse marketplaces.

References

- Bajari P, Burdick B, Imbens GW, Masoero L, McQueen J, Richardson T, Rosen IM (2021) Multiple randomization designs. arXiv preprint arXiv:2112.13495.
- Bennett J, Lanning S, et al. (2007) The netflix prize. *Proceedings of KDD cup and workshop*, volume 2007, 35 (New York).
- Bojinov I, Simchi-Levi D, Zhao J (2023) Design and analysis of switchback experiments. *Management Science* 69(7):3759–3777.
- Bright I, Delarue A, Lobel I (2022) Reducing marketplace interference bias via shadow prices. arXiv preprint arXiv:2205.02274 .
- Cervera MR, Dätwyler R, D'Angelo F, Keurti H, Grewe BF, Henning C (2021) Uncertainty estimation under model misspecification in neural network regression. arXiv preprint arXiv:2111.11763.
- Cheng Y, Wang J, Cao X, Shen ZJM, Zhang Y (2023) Selecting creators to sign on a content-sharing platform: A deep-did approach. Available at SSRN 4622422.
- Chernozhukov V, Chetverikov D, Demirer M, Duflo E, Hansen C, Newey W, Robins J (2018) Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal* 21(1):C1–C68.
- Chernozhukov V, Demirer M, Lewis G, Syrgkanis V (2019) Semi-parametric efficient policy learning with continuous actions. Advances in Neural Information Processing Systems 32.
- Covington P, Adams J, Sargin E (2016) Deep neural networks for youtube recommendations. Proceedings of the 10th ACM conference on recommender systems, 191–198.
- Dhaouadi W, Johari R, Weintraub GY (2023) Price experimentation and interference in online platforms. arXiv preprint arXiv:2310.17165.
- Ellickson PB, Kar W, Reeder III JC, Zeng G (2024) Using contextual embeddings to predict the effectiveness of novel heterogeneous treatments. Available at SSRN 4845956 .
- Farias V, Li H, Peng T, Ren X, Zhang H, Zheng A (2023) Correcting for interference in experiments: A case study at douyin. Proceedings of the 17th ACM Conference on Recommender Systems, 455–466.
- Farias VF, Li AA, Peng T, Zheng AT (2022) Markovian interference in experiments. arXiv preprint arXiv:2206.02371 .
- Farrell MH, Liang T, Misra S (2020) Deep learning for individual heterogeneity: An automatic inference framework. arXiv preprint arXiv:2010.14694.
- Foster DJ, Syrgkanis V (2023) Orthogonal statistical learning. The Annals of Statistics 51(3):879–908.
- Goli A, Lambrecht A, Yoganarasimhan H (2023) A bias correction approach for interference in ranking experiments. *Marketing Science*.
- Hall P, Heyde CC (2014) Martingale limit theory and its application (Academic press).

- Hinton G, Vinyals O, Dean J (2015) Distilling the knowledge in a neural network. arXiv preprint arXiv:1503.02531.
- Holtz D, Aral S (2020) Limiting bias from test-control interference in online marketplace experiments. arXivpreprint arXiv:2004.12162.
- Holtz D, Lobel F, Lobel R, Liskovich I, Aral S (2023) Reducing interference bias in online marketplace experiments using cluster randomization: Evidence from a pricing meta-experiment on airbnb. Management Science .
- Hu Y, Li S, Wager S (2021) Average treatment effects in the presence of interference. arXiv preprint arXiv:2104.03802.
- Hu Y, Wager S (2022) Switchback experiments under geometric mixing. arXiv preprint arXiv:2209.00197.
- Hudgens MG, Halloran ME (2008) Toward causal inference with interference. *Journal of the American* Statistical Association 103(482):832–842.
- Imbens GW (2004) Nonparametric estimation of average treatment effects under exogeneity: A review. Review of Economics and statistics 86(1):4–29.
- Imbens GW, Rubin DB (2015) Causal Inference in Statistics, Social, and Biomedical Sciences (Cambridge University Press).
- Ivaniuk A, Duan W (2020) A/b testing at linkedin: Assigning variants at scale. Accessed on Feb 4, 2024. URL https://www.linkedin.com/blog/engineering/ab-testing-experimentation/ a-b-testing-variant-assignment.
- Johari R, Li H, Liskovich I, Weintraub GY (2022) Experimental design in two-sided platforms: An analysis of bias. *Management Science* 68(10):7069–7089.
- Kim D, Jiang Z, Thomadsen R (2023) Tv advertising effectiveness with racial minority representation: Evidence from the mortgage market. Available at SSRN 4521178.
- Kiros H (2022) Hated that video? YouTube's algorithm might push you another just like it. URL https: //www.technologyreview.com/2022/09/20/1059709/youtube-algorithm-recommendations/#: ~:text=YouTube's%20recommendation%20algorithm%20drives%2070,adjust%20what%20it% 20shows%20them.
- Mehrotra R, McInerney J, Bouchard H, Lalmas M, Diaz F (2018) Towards a fair marketplace: Counterfactual evaluation of the trade-off between relevance, fairness & satisfaction in recommendation systems. Proceedings of the 27th acm international conference on information and knowledge management, 2243– 2251.
- Mummalaneni S, Yoganarasimhan H, Pathak V (2022) How do content producers respond to engagement on social media platforms? *Available at SSRN 4173537*.
- Newey WK (1994) The asymptotic variance of semiparametric estimators. *Econometrica: Journal of the Econometric Society* 62(6):1349–1382.

- Newey WK, McFadden D (1994) Chapter 36 large sample estimation and hypothesis testing. volume 4 of handbook of econometrics. *Elsevier* 12:2111–2245.
- Ni T, Bojinov I, Zhao J (2023) Design of panel experiments with spatial and temporal interference. Available at SSRN 4466598 .
- Rosen S (1981) The economics of superstars. The American economic review 71(5):845–858.
- Sävje F, Aronow P, Hudgens M (2021) Average treatment effects in the presence of unknown interference. Annals of statistics 49(2):673.
- Su Y, Wang X, Le EY, Liu L, Li Y, Lu H, Lipshitz B, Badam S, Heldt L, Bi S, et al. (2024) Long-term value of exploration: Measurements, findings and algorithms. *Proceedings of the 17th ACM International Conference on Web Search and Data Mining*, 636–644.
- Tang J, Wang K (2018) Ranking distillation: Learning compact ranking models with high performance for recommender system. Proceedings of the 24th ACM SIGKDD international conference on knowledge discovery & data mining, 2289–2298.
- Taylor S, Wexler J (2003) Where should traffic shaping occur? Accessed on Feb 4, 2024. URL https://web.archive.org/web/20050207081929/http://www.nwfusion.com/newsletters/ frame/2001/00477507.html.
- Ugander J, Karrer B, Backstrom L, Kleinberg J (2013) Graph cluster randomization: Network exposure to multiple universes. Proceedings of the 19th ACM SIGKDD international conference on Knowledge discovery and data mining, 329–337.
- Viviano D (2019) Policy targeting under network interference. arXiv preprint arXiv:1906.10258.
- Xiong R, Chin A, Taylor S (2023) Bias-variance tradeoffs for designing simultaneous temporal experiments. The KDD'23 Workshop on Causal Discovery, Prediction and Decision, 115–131 (PMLR).
- Ye Z, Zhang DJ, Zhang H, Zhang R, Chen X, Xu Z (2023a) Cold start to improve market thickness on online advertising platforms: Data-driven algorithms and field experiments. *Management Science* 69(7):3838– 3860.
- Ye Z, Zhang Z, Zhang D, Zhang H, Zhang RP (2023b) Deep learning based causal inference for large-scale combinatorial experiments: Theory and empirical evidence. Available at SSRN 4375327.
- Zhu Z, Cai Z, Zheng L, Si N (2024) Seller-side experiments under interference induced by feedback loops in two-sided platforms. arXiv preprint arXiv:2401.15811.

Appendix A: Proof of Proposition 1 (Convergence of the DIM Estimator)

Let n be the number of recommendations and $m = |\mathcal{C}|$ be the number of content items. We assume m = O(n) without loss of generality since at most nK contents are considered. Let q be the probability of treatment assignment. For each content c, let w_c denote its treatment status. The DIM estimator is:

$$\hat{\tau}_n^{DIM} = \frac{1}{nq} \sum_{i=1}^n W_{i,k_i^*} Y_i - \frac{1}{n(1-q)} \sum_{i=1}^n (1 - W_{i,k_i^*}) Y_i.$$

We now characterize the asymptotic behavior of $\hat{\tau}^{DIM}$. We have:

$$\begin{aligned} \hat{\tau}_n^{DIM} &= \frac{1}{nq} \sum_{i=1}^n \sum_{c \in \mathcal{C}} w_c \mathbf{1} \left\{ c = C_{i,k_i^*} \right\} Y_i - \frac{1}{n(1-q)} \sum_{i=1}^n \sum_{c \in \mathcal{C}} (1-w_c) \mathbf{1} \left\{ c = C_{i,k_i^*} \right\} Y_i \\ &= \frac{1}{nq} \sum_{c \in \mathcal{C}} w_c \sum_{i=1}^n \mathbf{1} \left\{ c = C_{i,k_i^*} \right\} Y_i - \frac{1}{n(1-q)} \sum_{c \in \mathcal{C}} (1-w_c) \sum_{i=1}^n \mathbf{1} \left\{ c = C_{i,k_i^*} \right\} Y_i \\ &= \frac{1}{q} \sum_{c \in \mathcal{C}} w_c R_c - \frac{1}{1-q} \sum_{c \in \mathcal{C}} (1-w_c) R_c, \end{aligned}$$

where we use $R_c := \frac{1}{n} \sum_{i=1}^n \mathbf{1} \left\{ c = C_{i,k_i^*} \right\} Y_i$ to denote the average viewer-outcome for content c. Let $\tau^{\mathcal{B}(q)}$ be

$$\tau^{\mathcal{B}(q)} := \mathbb{E}\left[\sum_{c \in \mathcal{C}} \left\{ r(c; w_c = 1, \mathbf{w}_{-c} \sim \mathcal{B}(q)) - r(c; w_c = 0, \mathbf{w}_{-c} \sim \mathcal{B}(q)) \right\}\right].$$

We now prove that $\hat{\tau}_n^{DIM}$ converges to $\tau^{\mathcal{B}(q)}$ in probability.

First, we show that $\mathbb{E}\left[\hat{\tau}_{n}^{DIM}\right] = \tau^{\mathcal{B}(q)}$.

$$\begin{split} \mathbb{E}\left[\hat{\tau}_{n}^{DIM}\right] &= \sum_{c \in \mathcal{C}} \frac{1}{q} \mathbb{E}_{\mathbf{w} \sim \mathcal{B}(q)}[w_{c}R_{c}] - \sum_{c \in \mathcal{C}} \frac{1}{1-q} \mathbb{E}_{\mathbf{w} \sim \mathcal{B}(q)}[(1-w_{c})R_{c}] \\ &= \sum_{c \in \mathcal{C}} \frac{1}{q} q \mathbb{E}_{\mathbf{w}_{-c} \sim \mathcal{B}(q)}[R_{c}|w_{c}=1] - \sum_{c \in \mathcal{C}} \frac{1}{1-q}(1-q)\mathbb{E}_{\mathbf{w}_{-c} \sim \mathcal{B}(q)}[R_{c}|w_{c}=0] \\ &= \sum_{c \in \mathcal{C}} \mathbb{E}_{\mathbf{w}_{-c} \sim \mathcal{B}(q)} \left[\frac{1}{n}\sum_{i=1}^{n} \mathbf{1}\left\{c = C_{i,k_{i}^{*}}\right\}Y_{i}|w_{c}=1\right] - \sum_{c \in \mathcal{C}} \mathbb{E}_{\mathbf{w}_{-c} \sim \mathcal{B}(q)} \left[\frac{1}{n}\sum_{i=1}^{n} \mathbf{1}\left\{c = C_{i,k_{i}^{*}}\right\}Y_{i}|w_{c}=0\right] \\ &= \sum_{c \in \mathcal{C}} \mathbb{E}_{\mathbf{w}_{-c} \sim \mathcal{B}(q)} \left[\mathbf{1}\left\{c = C_{i,k_{i}^{*}}\right\}Y_{i}|w_{c}=1\right] - \sum_{c \in \mathcal{C}} \mathbb{E}_{\mathbf{w}_{-c} \sim \mathcal{B}(q)} \left[\mathbf{1}\left\{c = C_{i,k_{i}^{*}}\right\}Y_{i}|w_{c}=0\right] \\ &= \sum_{c \in \mathcal{C}} \mathbb{E}\left[y(v = V_{i},c;w_{c}=1,\mathbf{w}_{-c} \sim \mathcal{B}(q))\right] - \sum_{c \in \mathcal{C}} \mathbb{E}\left[y(v = V_{i},c;w_{c}=0,\mathbf{w}_{-c} \sim \mathcal{B}(q))\right] = \tau^{\mathcal{B}(q)}. \end{split}$$

Next, we characterize the variance of $\hat{\tau}_n^{DIM}.$ By the law of total variance, we have

$$\operatorname{Var}(\hat{\tau}_{n}^{DIM}) = \mathbb{E}\left[\operatorname{Var}\left(\hat{\tau}_{n}^{DIM} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}\right)\right] + \operatorname{Var}\left(\mathbb{E}\left[\hat{\tau}_{n}^{DIM} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}\right]\right)$$

Conditioning on $\{(V_i, \vec{C}_i)\}_{i=1}^n$, for each content *c*, we have:

$$\begin{aligned} \operatorname{Var}\left(\frac{w_{c}R_{c}}{q} - \frac{(1 - w_{c})R_{c}}{1 - q} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}\right) \\ &= \frac{\operatorname{Var}(w_{c}R_{c} \mid \{(V_{i}, \vec{C}_{i}))\}_{i=1}^{n} + \frac{\operatorname{Var}((1 - w_{c})R_{c} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n})}{(1 - q)^{2}} - \frac{2\operatorname{Cov}(w_{c}R_{c}, (1 - w_{c})R_{c} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n})}{q(1 - q)} \\ &= \frac{\mathbb{E}[w_{c}R_{c}^{2} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}] - \mathbb{E}[w_{c}R_{c} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}]^{2}}{q^{2}} + \frac{\mathbb{E}[(1 - w_{c})R_{c}^{2} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}] - \mathbb{E}[(1 - w_{c})R_{c}]^{2}}{(1 - q)^{2} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}} \\ &+ \frac{2\mathbb{E}[w_{c}R_{c} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}]\mathbb{E}[(1 - w_{c})R_{c} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}]}{q(1 - q)} \end{aligned}$$

$$\begin{split} &= \frac{\mathbb{E}[R_c^2 \mid w_c = 1, \{(V_i, \vec{C}_i)\}_{i=1}^n]}{q} - \mathbb{E}[R_c \mid w_c = 1, \{(V_i, \vec{C}_i)\}_{i=1}^n]^2 \\ &+ \frac{\mathbb{E}[R_c^2 \mid w_c = 0, \{(V_i, \vec{C}_i)\}_{i=1}^n]}{1 - q} - \mathbb{E}[R_c \mid w_c = 0, \{(V_i, \vec{C}_i)\}_{i=1}^n]^2 + 2\mathbb{E}[R_c \mid w_c = 1]\mathbb{E}[R_c \mid w_c = 1, \{(V_i, \vec{C}_i)\}_{i=1}^n] \\ &= \frac{\mathbb{E}[R_c^2 \mid w_c = 1, \{(V_i, \vec{C}_i)\}_{i=1}^n]}{q} + \frac{\mathbb{E}[R_c^2 \mid w_c = 0, \{(V_i, \vec{C}_i)\}_{i=1}^n]}{1 - q} \\ &- \left(\mathbb{E}[R_c \mid w_c = 1, \{(V_i, \vec{C}_i)\}_{i=1}^n] - \mathbb{E}[R_c \mid w_c = 1, \{(V_i, \vec{C}_i)\}_{i=1}^n]\right)^2. \end{split}$$

Recall that $R_c := \frac{1}{n} \sum_{i=1}^n \mathbf{1} \{j = k_i^*\} Y_i$. By the boundedness of Y_i and Assumption 1, we have that $R_c = O(\frac{a_n}{n})$, and thus:

$$\operatorname{Var}\left(\frac{w_c R_c}{q} - \frac{(1 - w_c) R_c}{1 - q} \mid \{(V_i, \vec{C}_i)\}_{i=1}^n\right) = O\left(\frac{a_n^2}{n^2}\right).$$

Also conditioning $\{(V_i, \vec{C}_i)\}_{i=1}^n$, define $I_{c_1,c_2} = 1$ if there exists an item c_3 such that item c_1 and item c_3 are present at one consideration set,⁹ and item c_2 and item c_3 are present at one consideration set simultaneously; otherwise set $I_{c_1,c_2} = 0$. Note that if $I_{c_1,c_2} = 0$, there is no interference among the items c_1 & c_2 and thus R_{c_1} and R_{c_2} are independent.

$$\begin{aligned} &\operatorname{Cov}\left(\frac{w_{c_{1}}R_{c_{1}}}{q} - \frac{(1 - w_{c_{1}})R_{c_{1}}}{1 - q}, \frac{w_{c_{2}}R_{c_{2}}}{q} - \frac{(1 - w_{c_{2}})R_{c_{2}}}{1 - q} \middle| \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}\right) \\ &= \begin{cases} 0 & \text{if } I_{c_{1}, c_{2}} = 0, \\ O\left(\sqrt{\operatorname{Var}\left(\frac{w_{c_{1}}R_{c_{1}}}{q} - \frac{(1 - w_{c_{1}})R_{c_{1}}}{1 - q} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}\right)}\sqrt{\operatorname{Var}\left(\frac{w_{c_{2}}R_{c_{2}}}{q} - \frac{(1 - w_{c_{2}})R_{c_{2}}}{1 - q} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}\right)}\right)} & \text{o.w.} \end{aligned}$$

Together, we have

$$\begin{aligned} \operatorname{Var}\left(\hat{\tau}_{n}^{DIM} \mid \{(V_{i},\vec{C}_{i})\}_{i=1}^{n}\right) &\lesssim \sum_{c \in \mathcal{C}}^{m} \operatorname{Var}\left(\frac{w_{c}R_{c}}{q} - \frac{(1-w_{c})R_{c}}{1-q} \mid \{(V_{i},\vec{C}_{i})\}_{i=1}^{n}\right) \\ &+ \sum_{c_{1} \neq c_{2} \in \mathcal{C}} \operatorname{Cov}\left(\frac{w_{c_{1}}R_{c_{1}}}{q} - \frac{(1-w_{c_{1}})R_{c_{1}}}{1-q}, \frac{w_{c_{2}}R_{c_{2}}}{q} - \frac{(1-w_{c_{2}})R_{c_{2}}}{1-q} \mid \{(V_{i},\vec{C}_{i})\}_{i=1}^{n}\right) \\ &\lesssim \frac{ma_{n}^{2}}{n^{2}} + \sum_{c_{1} \neq c_{2} \in \mathcal{C}} I_{c_{1},c_{2}}\frac{a_{n}^{2}}{n^{2}} \lesssim \frac{ma_{n}^{2}}{n^{2}} + \frac{ma_{n}^{4}}{n^{2}} = O\left(\frac{a_{n}^{4}}{n}\right). \end{aligned}$$

Thus,

$$\mathbb{E}\left[\operatorname{Var}\left(\hat{\tau}_{n}^{DIM} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}\right)\right] = O\left(\frac{a_{n}^{4}}{n}\right).$$
(19)

On the other hand, we have

$$\begin{split} \mathbb{E}\left[\hat{\tau}_{n}^{DIM} \mid \{(V_{i},\vec{C}_{i})\}_{i=1}^{n}\right] &= \sum_{c \in \mathcal{C}} \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^{n} \frac{1\left\{c = C_{i,k_{i}^{*}}\right\} w_{c}}{q} Y_{i} - \frac{1}{n} \sum_{i=1}^{n} \frac{1\left\{c = C_{i,k_{i}^{*}}\right\} (1 - w_{c})}{1 - q} Y_{i} \mid \{(V_{i},\vec{C}_{i})\}_{i=1}^{n}\right] \right] \\ &= \frac{1}{n} \sum_{i=1}^{n} \sum_{c \in \mathcal{C}} \left\{ \mathbb{E}\left[\frac{1\left\{c = C_{i,k_{i}^{*}}\right\} w_{c}}{q} Y_{i} \mid \{(V_{i},\vec{C}_{i})\}_{i=1}^{n}\right] - \mathbb{E}\left[\frac{1\left\{c = C_{i,k_{i}^{*}}\right\} (1 - w_{c})}{1 - q} Y_{i} \mid \{(V_{i},\vec{C}_{i})\}_{i=1}^{n}\right] \right] \\ &= \frac{1}{n} \sum_{i=1}^{n} Z_{i}, \\ \text{where} \quad Z_{i} = \mathbb{E}\left[\left(\frac{W_{i,k_{i}^{*}}}{q} - \frac{1 - W_{i,k_{i}^{*}}}{1 - q}\right) Y_{i} \mid \{(V_{i},\vec{C}_{i})\}_{i=1}^{n}\right]. \end{split}$$

⁹ By convention, we say that c_1 and c_3 are in the same consideration set if both refer to the same item.

Note that if sample i_1 and sample i_2 do not share items in the consideration set, i.e., $\vec{C}_{i_1} \cap \vec{C}_{i_2} = \emptyset$, we have $Z_{i_1} \perp Z_{i_2} \mid \{(V_i, \vec{C}_i)\}_{i=1}^n$. Thus,

$$\operatorname{Var}\left(\mathbb{E}\left[\hat{\tau}_{n}^{DIM} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}\right]\right) = \operatorname{Var}\left(\frac{1}{n}\sum_{i=1}^{n}Z_{i}\right) = \frac{\sum_{i=1}^{n}\operatorname{Var}(Z_{i}) + \sum_{i_{1}\neq i_{2}}\operatorname{Cov}(Z_{i_{1}}, Z_{i_{2}})}{n^{2}}$$
$$= \frac{O(n) + na_{n}K \cdot O(1)}{n^{2}} = O\left(\frac{a_{n}}{n}\right). \tag{20}$$

Combining (19) & (20), we have

$$\operatorname{Var}\left(\hat{\tau}_{n}^{DIM}\right) = \mathbb{E}\left[\operatorname{Var}\left(\hat{\tau}_{n}^{DIM} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}\right)\right] + \operatorname{Var}\left(\mathbb{E}\left[\hat{\tau}_{n}^{DIM} \mid \{(V_{i}, \vec{C}_{i})\}_{i=1}^{n}\right]\right) = O\left(\frac{a_{n}^{4}}{n}\right) + O\left(\frac{a_{n}}{n}\right) = O\left(\frac{a_{n}^{4}}{n}\right) = O$$

Recall that $\mathbb{E}[\hat{\tau}_n^{DIM}] = \tau^{\mathcal{B}(q)},$ by Markov inequality we have

$$\hat{\tau}_n^{DIM} \xrightarrow{p} \tau^{\mathcal{B}(q)}.$$
 (21)

Appendix B: Counterfactual Evaluation with the Recommender Choice Model

We write out the explicit expression of counterfactual policy values using the recommender choice model. Given a policy π , the policy value is

$$Q(\pi) = \mathbb{E}_{\mathbf{w} \sim \pi} \left[\sum_{c \in \mathcal{C}} r(c; w_c, \mathbf{w}_{-c}) \right] = \mathbb{E}_{\mathbf{w} \sim \pi} \left[\sum_{c \in \mathcal{C}} \mathbb{E}_v [y(v, c; w_c, \mathbf{w}_{-c})] \right]$$
$$= \mathbb{E}_v \left[\mathbb{E}_{\mathbf{w} \sim \pi} \left[\sum_{c \in \mathcal{C}} y(v, c; w_c, \mathbf{w}_{-c}) \right] \right]$$
$$= \mathbb{E}_v \left[\mathbb{E}_{\overrightarrow{W}_i \sim \pi} \left[Y_i(V_i = v, \overrightarrow{C_i}; \overrightarrow{W_i}) \right] \right] = \mathbb{E}_{V_i, \overrightarrow{W_i} \sim \pi} \left[Y_i(V_i, \overrightarrow{C_i}; \overrightarrow{W_i}) \right]$$
$$= \mathbb{E}_{V_i, \overrightarrow{W_i} \sim \pi} \left[\sum_{k=1}^K z(V_i, C_{i,k}) \cdot \mathbb{P} \left(k_i^* = k \mid V_i, \overrightarrow{C_i}, \overrightarrow{W_i} \right) \right]$$
$$= \mathbb{E}_{V_i, \overrightarrow{W_i} \sim \pi} \left[\sum_{k=1}^K z(V_i, C_{i,k}) \cdot \frac{e^{s_0(V_i, C_{i,k}) + W_{i,k} \cdot s_1(V_i, C_{i,k})}}{\sum_{k'=1}^K e^{s_0(V_i, C_{i,k'}) + W_{i,k'} \cdot s_1(V_i, C_{i,k'})}} \right].$$

Our framework also allows us to instantiate Proposition 1 and explicitly characterize the asymptotic bias of DIM estimator. Let $\delta_{i,k}$ represents the change in exposure probability between the global treatment and global control:

$$\delta_{i,k} := \frac{e^{s_0(V_i, C_{i,k}) + s_1(V_i, C_{i,k})}}{\sum_{k'=1}^{K} e^{s_0(V_i, C_{i,k'}) + s_1(V_i, C_{i,k'})}} - \frac{e^{s_0(V_i, C_{i,k})}}{\sum_{k'=1}^{K} e^{s_0(V_i, C_{i,k'})}}.$$

The bias of the DIM estimator is

$$\Delta^{DIM} := \tau^{\mathcal{B}(q)} - \tau = \mathbb{E}\left[\sum_{k=1}^{K} z(V_i, C_{i,k}) \cdot \left\{\delta_{i,k}^{\mathcal{B}(q)} - \delta_{i,k}\right\}\right],\tag{22}$$

where $\delta_{i,k}^{\mathcal{B}(q)}$ represents the change in exposure probability for content $C_{i,k}$ due to its own treatment when the treatment statuses for others are sampled from the Bernoulli trial $\mathcal{B}(q)$:

$$\delta_{i,k}^{\mathcal{B}(q)} := \mathbb{E}_{W_{i,k'\neq k}\sim\mathcal{B}(q)} \left[\frac{e^{s_0(V_i,C_{i,k})+s_1(V_i,C_{i,k})}}{e^{s_0(V_i,C_{i,k})+s_1(V_i,C_{i,k})} + \sum_{k'\neq k} e^{s_0(V_i,C_{i,k'})+W_{i,k'}\cdot s_1(V_i,C_{i,k'})}} - \frac{e^{s_0(V_i,C_{i,k})}}{e^{s_0(V_i,C_{i,k})} + \sum_{k'\neq k} e^{s_0(V_i,C_{i,k'})+W_{i,k'}\cdot s_1(V_i,C_{i,k'})}} \right].$$
(23)

Appendix C: Explicit Expression of Debiased Estimator

C.1. Debiased Term

We now write out the explicit debiased estimate $\psi(\cdot)$ for each observation, where we drop the subscript i and write the notation as $(V, \vec{C}, \vec{W}, k^*, Y)$ for succinctness. For estimated baseline score function $\hat{s}_0 \in \mathcal{F}_s$. treatment score uplift function $\hat{s}_1 \in \mathcal{F}_s$, and viewer response function $\hat{z} \in \mathcal{F}_z$, we have the estimated exposure probability be

$$\mathbb{P}\left(k^{*}=k \mid V, \overrightarrow{C}, \overrightarrow{W}; \hat{s}_{0}, \hat{s}_{1}\right) = \frac{e^{\hat{s}_{0}(V, C_{k}) + W_{k} \cdot \hat{s}_{1}(V, C_{k})}}{\sum_{k'=1}^{K} e^{\hat{s}_{0}(V, C_{k'}) + W_{k'} \cdot \hat{s}_{1}(V, C_{k'})}} \stackrel{(i)}{=} \frac{e^{\hat{s}_{0}(V, C_{k}) - \hat{s}_{0}(V, C_{1}) + W_{k} \cdot \hat{s}_{1}(V, C_{k})}}{\sum_{k'=1}^{K} e^{\hat{s}_{0}(V, C_{k'}) - \hat{s}_{0}(V, C_{1}) + W_{k'} \cdot \hat{s}_{1}(V, C_{k'})}},$$

where the equation (i) is obtained by normalizing both the numerator and denominator by the exponential of the first content item's baseline score. In other words, for any baseline score vector $(\hat{s}_0(V,C_1),\hat{s}_0(V,C_2),\ldots,\hat{s}_0(V,C_K)),$ if we replace it by $(0,\hat{s}_0(V,C_2)-\hat{s}_0(V,C_1),\ldots,\hat{s}_0(V,C_K)-\hat{s}_0(V,C_1)),$ we will get the same exposure probability result.

This implies that, for any nuisance estimates $(\hat{s}_0, \hat{s}_1, \hat{z})$, the value $\mu(V, \vec{C}; \hat{s}_0, \hat{s}_1, \hat{z})$ can be fully recovered by the vectors

- $\overrightarrow{\hat{S}_0} = (\hat{S}_{0,2}, \dots, \hat{S}_{0,K}) \in \mathbb{R}^{K-1}$ with $\hat{S}_{0,k} = \hat{s}_0(V, C_k) \hat{s}_0(V, C_1);$ $\overrightarrow{\hat{S}_1} = (\hat{S}_{1,1}, \dots, \hat{S}_{1,K}) \in \mathbb{R}^K$ with $\hat{S}_{1,k} = \hat{s}_1(V, C_k);$
- $\vec{\hat{Z}} = (\hat{Z}_{1,1}, \dots, \hat{Z}_{1,K}) \in \mathbb{R}^K$ with $\hat{Z}_{1,k} = \hat{z}(V, C_k)$.

We thus abuse notations and write $\mu(V, \vec{C}; \tilde{s}_0, \tilde{s}_1, \tilde{z})$ as $\mu\left(\vec{\hat{s}_0}, \vec{\hat{s}_1}, \vec{\hat{z}}\right)$. The bias of $\mu\left(\vec{\hat{s}_0}, \vec{\hat{s}_1}, \vec{\hat{z}}\right)$ comes from the deviation of $\left(\vec{\hat{s}_0}, \vec{\hat{s}_1}, \vec{\hat{z}}\right)$ to the true vectors $\left(\vec{\hat{s}_0}, \vec{\hat{s}_1}, \vec{\hat{z}}\right)$ that are defined similarly under the true model (s_0, s_1, z) .

We next follow the double machine learning literature and use the outcome (k^*, Y) to correct the bias of $\mu\left(\vec{\hat{S}}_{0},\vec{\hat{S}}_{1},\vec{\hat{Z}}\right)$ due to the bias of $\left(\vec{\hat{S}}_{0},\vec{\hat{S}}_{1},\vec{\hat{Z}}\right)$ approximating the true $\left(\vec{S}_{0},\vec{S}_{1},\vec{Z}\right)$.

Under the estimates $(\hat{S}_0, \hat{S}_1, \hat{Z})$, we reload the loss function notation and write it as:

$$\ell\left(\overrightarrow{W},k^*,Y;\overrightarrow{\hat{S}_0},\overrightarrow{\hat{S}_1},\overrightarrow{\hat{Z}}\right) = \ell_1\left(\overrightarrow{W},k^*;\overrightarrow{\hat{S}_0},\overrightarrow{\hat{S}_1}\right) + \ell_2\left(k^*,Y;\overrightarrow{\hat{Z}}\right)$$

where

$$\ell_1\left(\vec{W}, k^*; \vec{\hat{S}_0}, \vec{\hat{S}_1}\right) = \begin{cases} -W_1 \hat{S}_{1,1} + \log\left(e^{W_1 \hat{S}_{1,1}} + \sum_{i=2}^K e^{\hat{S}_{0,k} + W_k \hat{S}_{1,k}}\right) & \text{if } k^* = 1\\ -\left(\hat{S}_{0,k} + W_k \hat{S}_{1,k}\right) + \log\left(e^{W_1 \hat{S}_{1,1}} + \sum_{i=2}^K e^{\hat{S}_{0,k} + W_k \hat{S}_{1,k}}\right) & \text{o.w.}; \end{cases}$$

and

$$\ell_2\left(k^*, Y; \overrightarrow{\hat{Z}}\right), = \left(\widehat{Z}_{k^*} - Y\right)^2.$$

We are now ready to introduce the debiased term ψ :

$$\psi\left(\overrightarrow{W},k^*,Y;\overrightarrow{\hat{S}_0},\overrightarrow{\hat{S}_1},\overrightarrow{\hat{Z}},\widehat{H}\right) = \mu\left(\overrightarrow{\hat{S}_0},\overrightarrow{\hat{S}_1},\overrightarrow{\hat{Z}}\right) - \nabla\mu^T \widehat{H}^{-1} \nabla\ell$$

Above, $\nabla \mu$ and $\nabla \ell$ are gradients of μ and ℓ with respect to the nuisance estimates $(\vec{\hat{S}}_0, \vec{\hat{S}}_1, \vec{\hat{Z}})$, and \hat{H} estimates the expected Hessian of ℓ regarding $\left(\vec{\hat{S}}_{0}, \vec{\hat{S}}_{1}, \vec{\hat{Z}}\right)$, where the expectation is taken with respect to the treatments \vec{W} that are assigned following the specified creator-side randomization. The explicit expressions for these derivatives are deferred to Appendix C.2.

C.2. Expressions of Derivatives

C.2.1. Gradient of μ . We have

$$\nabla \mu = \left(\frac{\partial \mu}{\partial \hat{S}_{0}}, \frac{\partial \mu}{\partial \hat{S}_{1}}, \frac{\partial \mu}{\partial \hat{Z}}\right)^{T} = \left(\frac{\partial \mu}{\partial \hat{S}_{0,2}}, \dots, \frac{\partial \mu}{\partial \hat{S}_{0,K}}, \frac{\partial \mu}{\partial \hat{S}_{1,1}}, \dots, \frac{\partial \mu}{\partial \hat{S}_{1,K}}, \frac{\partial \mu}{\partial \hat{Z}_{1}}, \dots, \frac{\partial \mu}{\partial \hat{Z}_{K}}\right)^{T},$$

where

• for each $k = 2, \ldots, K$,

$$\frac{\partial \mu}{\partial \hat{S}_{0,k}} = P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W} \equiv 1) \left\{ \hat{Z}_k - \mathbb{E}[Y \mid \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{\hat{Z}}, \overrightarrow{W} \equiv 1] \right\} - P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W} \equiv 0) \left\{ \hat{Z}_k - \mathbb{E}[Y \mid \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{\hat{Z}}, \overrightarrow{W} \equiv 0] \right\}.$$

• for each $k = 1, \ldots, K$,

$$\begin{split} &\frac{\partial\mu}{\partial\hat{S}_{1,k}} = P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W} \equiv 1) \left\{ \hat{Z}_k - \mathbb{E}[Y \mid \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{\hat{Z}}, \overrightarrow{W} \equiv 1] \right\}, \\ &\frac{\partial\mu}{\partial\hat{Z}_k} = P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W} \equiv 1) - P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W} \equiv 0). \end{split}$$

Above, $\langle\cdot,\cdot\rangle$ denotes inner product between two vectors.

C.2.2. Gradient of ℓ . We have

$$\nabla \ell = \left(\frac{\partial \ell_1}{\partial \hat{S}_0}, \frac{\partial \ell_1}{\partial \hat{S}_1}, \frac{\partial \ell_2}{\partial \hat{Z}}\right)^T = \left(\frac{\partial \ell_1}{\partial \hat{S}_{0,2}}, \dots, \frac{\partial \ell_1}{\partial \hat{S}_{0,K}}, \frac{\partial \ell_1}{\partial \hat{S}_{1,1}}, \dots, \frac{\partial \ell_1}{\partial \hat{S}_{1,K}}, \frac{\partial \ell_2}{\partial \hat{Z}_1}, \dots, \frac{\partial \ell_2}{\partial \hat{Z}_K}\right)^T,$$

where

• for each $k = 2, \ldots, K$:

$$\frac{\partial \ell_1}{\partial \hat{S}_{0,k}} = P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) - \mathbb{I}[k^* = k].$$

• for each $k = 1, \ldots, K$:

$$\begin{split} & \frac{\partial \ell_1}{\partial \hat{S}_{1,k}} = \mathbb{I}\left[W_k = 1\right] \left(P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) - \mathbb{I}[k^* = k]\right), \\ & \frac{\partial \ell_2}{\partial \hat{Z}_k} = \mathbb{I}[k^* = k] \left(\hat{Z}_k - Y\right). \end{split}$$

C.2.3. Hessian of ℓ . We have

$$\nabla^2 \ell = \begin{pmatrix} \frac{\partial^2 \ell_1}{\partial \bar{S}_0^{-2}} & \frac{\partial^2 \ell_1}{\partial \bar{S}_0 \partial \bar{S}_1} & 0\\ \frac{\partial^2 \ell_1}{\partial \bar{S}_1 \partial \bar{S}_0} & \frac{\partial^2 \ell_1}{\partial \bar{S}_1^{-2}} & 0\\ 0 & 0 & \frac{\partial^2 \ell_2}{\partial \bar{Z}^2} \end{pmatrix},$$

where the Hessian of loss function ℓ_1 follows;

$$\frac{\partial^2 \ell_1}{\partial \hat{S}^2_{0,k}} = P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) \left(1 - P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) \right), \qquad k \in \{2: K\}$$

$$\frac{\partial^2 \ell_1}{\partial \hat{S}_{1,k}^2} = W_k P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) \left(1 - P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) \right), \qquad k \in \{1: K\}$$

$$\frac{\partial^2 \ell_1}{\partial \hat{S}_{0,k} \partial \hat{S}_{1,k}} = W_k P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) \left(1 - P(k^* = k; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) \right), \qquad k \in \{2: K\}$$

$$\begin{split} &\frac{\partial^2 \ell_1}{\partial \hat{S}_{0,k_1} \partial \hat{S}_{0,k_2}} = -P(k^* = k_1; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) P(k^* = k_2; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}), \qquad k_1 \neq k_2 \in \{2:K\} \\ &\frac{\partial^2 \ell_1}{\partial \hat{S}_{0,k_1} \partial \hat{S}_{1,k_2}} = -W_{k_2} P(k^* = k_1; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) P(k^* = k_2; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}), \qquad k_1 \neq k_2, k_1 \in \{2:K\}, k_2 \in \{1:K\} \\ &\frac{\partial^2 \ell_1}{\partial \hat{S}_{1,k_1} \partial \hat{S}_{1,k_2}} = -W_{k_1} W_{k_2} P(k^* = k_1; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}) P(k^* = k_2; \overrightarrow{\hat{S}_0}, \overrightarrow{\hat{S}_1}, \overrightarrow{W}), \qquad k_1 \neq k_2 \in \{1:K\}. \end{split}$$

and the Hessian of loss function ℓ_2 follows:

$$\begin{split} &\frac{\partial \ell_2^2}{\partial \hat{Z}_k^2} = \mathbb{I}[k^* = k], \qquad \qquad k \in \{1:K\} \\ &\frac{\partial \ell_2^2}{\partial \hat{Z}_{k_1} \partial \hat{Z}_{k_2}} = 0, \qquad \qquad k_1 \neq k_2 \in \{1:K\}. \end{split}$$

Appendix D: Discussion on Identifying Nuisance Components

We now provide more details on the identifiability of the nuisance components. At a high level, these nuisances can be well estimated if we have a personalized recommender system, as the case for our empirical application. To crystallize this idea, let's consider a concrete example where the score functions are parameterized as follows:

$$s_0(U_i, C_{i,k}) = \beta_0^T \psi(V_i, C_{i,k}) \quad \text{and} \quad s_1(U_i, C_{i,k}) = \beta_1^T \psi(V_i, C_{i,k}).$$
(24)

Above, $\psi(\cdot)$ is some feature mapping function we shall discuss shortly, and β_0, β_1 are the score coefficient vectors we want to identify using the data from a creator-side randomization experiment. With that, the original identification problem (13) is equivalent to the following:

$$\hat{\beta}_{0}, \hat{\beta}_{1} \in \underset{\tilde{\beta}_{0}, \tilde{\beta}_{1}}{\operatorname{arg\,min}} \mathcal{L}_{1}(\tilde{\beta}_{0}, \tilde{\beta}_{1}) := \frac{1}{n} \sum_{i=1}^{n} \ell_{1}\left(V_{i}, \overrightarrow{C_{i}}, \overrightarrow{W_{i}}, C_{i}^{*}, Y_{i}; \tilde{\beta}_{0}, \tilde{\beta}_{1}\right),$$

$$(25)$$

where $\ell_1(\cdot)$ is the cross-entropy loss defined in (12). As a result, $(\hat{\beta}_0, \hat{\beta}_1)$ is the maximum likelihood estimation (MLE). Assuming that the covariance matrix, defined as follows,

$$\Sigma := \mathbb{E}[X_i X_i^{\top}], \quad \text{where} \quad X_i = (\psi(V_i, C_{i,1}); \dots; \psi(V_i, C_{i,K})), \tag{26}$$

has bounded inverse. Then under mild conditions, this MLE is asymptotically normal (Newey and McFadden 1994):

$$\sqrt{n}(\hat{\beta}_0 - \beta_0) \xrightarrow{d} \mathcal{N}(0, \mathcal{I}_0^{-1}) \quad \text{and} \quad \sqrt{n}(\hat{\beta}_1 - \beta_1) \xrightarrow{d} \mathcal{N}(0, \mathcal{I}_1^{-1})$$
(27)

where $\mathcal{I}_0 = \mathbb{E}\left[\frac{\partial^2 \ell_1(\tilde{\beta}_0)}{(\partial \tilde{\beta}_0)^2} \mid \tilde{\beta}_0 = \beta_0\right]$ and $\mathcal{I}_1 = \mathbb{E}\left[\frac{\partial^2 \ell_1(\tilde{\beta}_1)}{(\partial \tilde{\beta}_1)^2} \mid \tilde{\beta}_1 = \beta_1\right]$ are the Fisher information matrices that measure the amount of information data carries to identify the parameter vectors β_0 and β_1 respectively. The regularities of \mathcal{I}_0 and \mathcal{I}_1 are determined by the regularity of Σ , as formalized by the result below.

LEMMA 2. Suppose Assumption 2 holds, and thus there exists a universal constant $\delta > 0$ that lower bounds the item exposure probability by Lemma 1. Suppose the content exposure follows the recommender choice model Eq.(7), with the score form specified in (24). Consider data has been collected from a creator-side randomization experiment with treated probability $q \in (0,1)$. Suppose that the covariate matrix Σ in (26) satisfies that

$$c \cdot I \preceq \Sigma \preceq C \cdot I,$$

for positive constants c, C. Then, the Fisher information matrices $\mathcal{I}_0, \mathcal{I}_1$ in (27) satisfies:

$$\delta^2 c K(K-1) \cdot I \preceq \mathcal{I}_0 \preceq C K(K-1) I, \quad \delta^2 q c K(K-1) \cdot I \preceq \mathcal{I}_1 \preceq q C K(K-1) \cdot I.$$

Therefore, the identifiability of β_0 and β_1 depends on the regularity of the covariate matrix Σ , which is typically well-conditioned in many contexts, including ours, where the recommendation process is highly personalized. This personalization ensures that the covariate X_i varies significantly among viewers V_i , leading to $\Sigma = \mathbb{E}[X_i X_i^{\top}]$ being full-rank with bounded inverse. Furthermore, the identifiability of β_1 also depends on the treatment randomization probability q, with its estimation variance scaling with q^{-1} (also known as inverse propensity), consistent with causal inference literature.

D.1. Proof of Lemma 2

Write $X_{i,k} = \psi(V_i, C_{i,k})$, and $P_{i,k} = \mathbb{P}(k = k^* \mid V_i, \overrightarrow{C_i}, \overrightarrow{W_i}, C_i^*; s_0, s_1)$. We have

$$\frac{\partial \ell_1 \left(V_i, \vec{C_i}, \vec{W_i}, C_i^*, Y_i \right)}{\partial \beta_0} = \sum_{i=1}^K \mathbf{1} \left\{ k = k^* \right\} (-X_{i,k}) + \sum_{k=1}^K P_{i,k} X_{i,k},$$
$$\frac{\partial \ell_1 \left(V_i, \vec{C_i}, \vec{W_i}, C_i^*, Y_i \right)}{\partial \beta_1} = \sum_{i=1}^K \mathbf{1} \left\{ k = k^* \right\} W_{i,k} (-X_{i,k}) + \sum_{k=1}^K W_{i,k} P_{i,k} X_{i,k}$$

Continuing, we have

$$\frac{\partial^2 \ell_1\left(V_i, \overrightarrow{C_i}, \overrightarrow{W_i}, C_i^*, Y_i\right)}{\partial \beta_0^2} = \sum_{i=1}^K P_{i,k} X_{i,k} X_{i,k}^\top - \left(\sum_{k=1}^K P_{i,k} X_{i,k}\right) \left(\sum_{k=1}^K P_{i,k} X_{i,k}\right)^\top,$$
$$\frac{\partial^2 \ell_1\left(V_i, \overrightarrow{C_i}, \overrightarrow{W_i}, C_i^*, Y_i\right)}{\partial \beta_1^2} = \sum_{i=1}^K P_{i,k} W_{i,k} X_{i,k} X_{i,k}^\top - \left(\sum_{k=1}^K W_{i,k} P_{i,k} X_{i,k}\right) \left(\sum_{k=1}^K W_{i,k} P_{i,k} X_{i,k}\right)^\top.$$

Let d be the dimension of $X_{i,k}$. For any unit $x \in \mathbb{R}^d$, we have

$$x^{\top} \frac{\partial^{2} \ell_{1} \left(V_{i}, \overrightarrow{C_{i}}, \overrightarrow{W_{i}}, C_{i}^{*}, Y_{i} \right)}{\partial \beta_{0}^{2}} x = \sum_{k=1}^{K} P_{i,k} (X_{i,k}^{\top} x)^{2} - \left(\sum_{k=1}^{K} P_{i,k} X_{i,k}^{\top} x \right)^{2},$$
$$= \sum_{k_{1}=1}^{K} P_{i,k_{1}} \sum_{k_{2}=1}^{K} P_{i,k_{2}} (X_{i,k_{1}}^{\top} x)^{2} - \left(\sum_{k=1}^{K} P_{i,k} X_{i,k}^{\top} x \right)^{2}$$
$$= \sum_{k_{1} < k_{2}}^{K} P_{i,k_{1}} P_{i,k_{2}} (X_{i,k_{1}}^{\top} x - X_{i,k_{2}}^{\top} x)^{2}.$$

Denote x_{k_1,k_2} as a concatenated vector of K subvectors of length d, with its k_1 -th and k_2 -th sub-vectors being x. Denote $M(k_1, k_2, a_1, a_2)$ as a row-concatenated matrix of K submatrices of size $d \times d$, with its k_1 -th and k_2 -th sub-matrices being a_1I and a_2I . Since $P_{i,k}$ is the exposure probability, and thus we have

$$\begin{aligned} x^{\top} \frac{\partial^{2} \ell_{1} \left(V_{i}, \overrightarrow{C_{i}}, \overrightarrow{W_{i}}, C_{i}^{*}, Y_{i} \right)}{\partial \beta_{0}^{2}} x &\leq \sum_{k_{1} < k_{2}}^{K} (X_{i,k_{1}}^{\top} x - X_{i,k_{2}}^{\top} x)^{2} = x^{\top} \sum_{k_{1} < k_{2}} (X_{i,k_{1}} - X_{i,k_{2}}) (X_{i,k_{1}} - X_{i,k_{2}})^{\top} x \\ &= \sum_{k_{1} < k_{2}} x_{k_{1},k_{2}}^{\top} M(k_{1},k_{2},1,-1)^{\top} \Sigma M(k_{1},k_{2},1,-1) x_{k_{1},k_{2}} \\ &\leq C \sum_{k_{1} < k_{2}} x_{k_{1},k_{2}}^{\top} M(k_{1},k_{2},1,-1)^{\top} M(k_{1},k_{2},1,-1) x_{k_{1},k_{2}} \\ &= C K(K-1) \|x\|_{2}^{2}. \end{aligned}$$

Also under Assumption 2, we have $P_{i,k} \ge \delta$, and thus

$$\begin{aligned} x^{\top} \frac{\partial^{2} \ell_{1} \left(V_{i}, \overrightarrow{C_{i}}, \overrightarrow{W_{i}}, C_{i}^{*}, Y_{i} \right)}{\partial \beta_{0}^{2}} x &\geq \delta^{2} \sum_{k_{1} < k_{2}}^{K} \left(X_{i,k_{1}}^{\top} x - X_{i,k_{2}}^{\top} x \right)^{2} = x^{T} \delta^{2} \sum_{k_{1} < k_{2}} \left(X_{i,k_{1}} - X_{i,k_{2}} \right) \left(X_{i,k_{1}} - X_{i,k_{2}} \right)^{\top} x \\ &= \delta^{2} \sum_{k_{1} < k_{2}} x_{k_{1},k_{2}}^{\top} M(k_{1},k_{2},1,-1)^{\top} \Sigma M(k_{1},k_{2},1,-1) x_{k_{1},k_{2}} \\ &\geq \delta^{2} c \sum_{k_{1} < k_{2}} x_{k_{1},k_{2}}^{\top} M(k_{1},k_{2},1,-1)^{\top} M(k_{1},k_{2},1,-1) x_{k_{1},k_{2}} \\ &= \delta^{2} c K(K-1) \|x\|_{2}^{2}. \end{aligned}$$

Similarly,

$$x^{\top} \frac{\partial^{2} \ell_{1} \left(V_{i}, \overrightarrow{C_{i}}, \overrightarrow{W_{i}}, C_{i}^{*}, Y_{i} \right)}{\partial \beta_{1}^{2}} x = \sum_{k=1}^{K} P_{i,k} (W_{i,k} X_{i,k}^{\top} x)^{2} - \left(\sum_{k=1}^{K} P_{i,k} W_{i,k} X_{i,k}^{\top} x \right)^{2},$$
$$= \sum_{k_{1}=1}^{K} P_{i,k_{1}} \sum_{k_{2}=1}^{K} P_{i,k_{2}} (W_{i,k} X_{i,k_{1}}^{\top} x)^{2} - \left(\sum_{k=1}^{K} P_{i,k} W_{i,k} X_{i,k}^{\top} x \right)^{2}$$
$$= \sum_{k_{1}$$

So we have

$$x^{\top} \frac{\partial^{2} \ell_{1} \left(V_{i}, \overrightarrow{C_{i}}, \overrightarrow{W_{i}}, C_{i}^{*}, Y_{i} \right)}{\partial \beta_{1}^{2}} x \leq x^{T} \sum_{k_{1} < k_{2}} (W_{i,k_{1}} X_{i,k_{1}} - W_{i,k_{2}} X_{i,k_{2}}) (W_{i,k_{1}} X_{i,k_{1}} - W_{i,k_{2}} X_{i,k_{2}})^{\top} x$$
$$= \sum_{k_{1} < k_{2}} x_{k_{1},k_{2}}^{\top} M(k_{1},k_{2}, W_{i,k_{1}}, -W_{i,k_{2}})^{\top} \Sigma M(k_{1},k_{2}, W_{i,k_{1}}, -W_{i,k_{2}}) x_{k_{1},k_{2}} \leq \sum_{k_{1} < k_{2}} C \|x\|_{2}^{2} (W_{i,k_{1}} + W_{i,k_{2}}),$$

so the eigenvalues of \mathcal{I}_1 are upper bounded by $\sum_{k_1 < k_2} 2qC = K(K-1)qC$. Also,

$$\begin{split} x^{\top} \frac{\partial^{2} \ell_{1} \left(V_{i}, \overrightarrow{C_{i}}, \overrightarrow{W_{i}}, C_{i}^{*}, Y_{i} \right)}{\partial \beta_{1}^{2}} x \\ \geq \delta^{2} x^{T} \sum_{k_{1} < k_{2}} (W_{i,k_{1}} X_{i,k_{1}} - W_{i,k_{2}} X_{i,k_{2}}) (W_{i,k_{1}} X_{i,k_{1}} - W_{i,k_{2}} X_{i,k_{2}})^{\top} x \\ = \delta^{2} \sum_{k_{1} < k_{2}} x_{k_{1},k_{2}}^{\top} M(k_{1},k_{2}, W_{i,k_{1}}, -W_{i,k_{2}})^{\top} \Sigma M(k_{1},k_{2}, W_{i,k_{1}}, -W_{i,k_{2}}) x_{k_{1},k_{2}} \\ \geq c \delta^{2} \sum_{k_{1} < k_{2}} \|x\|_{2}^{2} (W_{i,k_{1}} + W_{i,k_{2}}), \end{split}$$

so the eigenvalues of \mathcal{I}_1 are lower bounded by $\delta^2 \sum_{k_1 < k_2} 2qc = K(K-1)qc\delta^2$.

Appendix E: Supporting Results for the Debiased Estimator

E.1. Positive Exposure

We now provide proof for Lemma 1. We have

$$\mathbb{P}\left(k_{i}^{*}=k \mid V_{i}, \overrightarrow{C_{i}}, \overrightarrow{W_{i}}; s_{0}, s_{1}\right) = \frac{e^{s_{0}(V_{i}, C_{i,k}) + W_{i,k} \cdot s_{1}(V_{i}, C_{i,k})}}{\sum_{k'=1}^{K} e^{s_{0}(V_{i}, C_{i,k'}) + W_{i,k'} \cdot s_{1}(V_{i}, C_{i,k'})}} \ge \frac{e^{-2C}}{\sum_{k'=1}^{K} e^{2C}} = e^{-2C-2\log(K)C},$$

where the inequality is by Assumption 2. Set $\delta = e^{-2C - 2\log(K)C}$, concluding the proof.

E.2. Invertible Hessian

LEMMA 3. Suppose Assumptions 2 hold. Suppose that the estimated scores are bounded: $\|\hat{s}_0\|_{\infty} \leq C$ and $\|\hat{s}_1\|_{\infty} \leq C$ for the C in Assumption 2. Under our modeling framework, the expected hessian $H = \mathbb{E}\left[\nabla^2 \ell | V, \vec{C}\right]$ is universally invertible with bounded inverse.

Note that H is a two-block diagonal matrix with the first diagonal block being $H_1 := \mathbb{E}\left[\nabla^2 \ell_1 | V, \vec{C}\right]$ and the second diagonal block being $H_2 := \mathbb{E}\left[\nabla^2 \ell_2 | V, \vec{C}\right]$. It suffices to show that both H_1 and H_2 are universally invertible with bounded inverses.

Regularity of H_1 . Consider the sample (V, \vec{C}, \vec{W}) and estimated nuisance \hat{s}_0, \hat{s}_1 . For notation convenience, we use p_k to represent $P(k^* = k; V, \vec{C}, \vec{W}, \hat{s}_0, \hat{s}_1)$. Define the vector

$$\beta = \begin{pmatrix} p_2 \\ \vdots \\ p_K \\ W_1 p_1 \\ \vdots \\ W_k p_K \end{pmatrix} \in \mathbb{R}^{2K-1},$$

the matrix $A \in \mathbb{R}^{(2K-1) \times (2K-1)}$ with its (K, K)-th entry being $\mathbf{1}[W_1 = 1]p_1$ and others zero and matrix

$$B = \begin{pmatrix} \operatorname{diag}(\sqrt{p_2}, \dots, \sqrt{p_K}) \\ 0, \dots, 0, \\ \operatorname{diag}(\sqrt{p_2}W_2, \dots, \sqrt{p_K}W_k) \end{pmatrix} \in \mathbb{R}^{(2K-1) \times (2K-1)},$$

We thus have

$$H_1 = \mathbb{E}\left[A + BB^T - \beta\beta^T \mid V, \vec{C}\right].$$

We now show that H_1 is positive definite. For any vector $x = (x_1, \ldots, x_{2K-1})$ and any treatment assignment \vec{W} , we have

$$x^{T}(A + BB^{T} - \beta\beta^{T})x$$

$$= p_{1}W_{1}x_{K}^{2} + \sum_{j=1}^{K-1} p_{j+1}(x_{j} + W_{j+1}x_{K+j})^{2} - \left(p_{1}W_{1}x_{K} + \sum_{j=1}^{K-1} p_{j+1}(x_{j} + W_{j+1}x_{K+j})\right)^{2}$$

$$= \left(p_{1}W_{1}x_{K}^{2} + \sum_{j=1}^{K-1} p_{j+1}(x_{j} + W_{j+1}x_{K+j})^{2}\right) \left(\sum_{j=1}^{K} p_{j}\right)$$

$$- \left(p_{1}W_{1}x_{K} + \sum_{j=1}^{K-1} p_{j+1}(x_{j} + W_{j+1}x_{K+j})\right)^{2} \stackrel{(i)}{\geq} 0,$$

$$(28)$$

where (i) is by Cauchy-Swartzh inequality. That is, for any treatment assignment, the Hessian is positive semi-definite, and thus H_1 (the expected Hessian) is positive semi-definite.

We next consider specific assignments of \vec{W} to lower bound the smallest eigenvalue of H_1 . Also note that under the assumption of the boundedness of score functions, there exists a constant $\delta > 0$, such that for any $(U, \vec{V}, \vec{W}, k^*)$, we have $P(k^* | U, \vec{V}, \vec{W}, \hat{s}_0, \hat{s}_1) \ge \delta$.

Consider $\overrightarrow{W}^{(a)} = (0, \dots, 0)$. Then we have (28) instantiate as

$$x^{T}(A + BB^{T} - \beta\beta^{T})x = \sum_{j=1}^{K-1} p_{j+1}^{(a)}x_{j}^{2} - \left(\sum_{j=1}^{K-1} p_{j+1}^{(a)}x_{j}\right)^{2}$$
$$= p_{1}^{(a)}\sum_{j=1}^{K-1} p_{j+1}^{(a)}x_{j}^{2} + \left(\sum_{j=1}^{K-1} p_{j+1}^{(a)}\right)\sum_{j=1}^{K-1} p_{j+1}^{(a)}x_{j}^{2} - \left(\sum_{j=1}^{K-1} p_{j+1}^{(a)}x_{j}\right)^{2} \ge p_{1}\sum_{j=1}^{K-1} p_{j+1}^{(a)}x_{j}^{2} \ge \delta^{2}\sum_{j=1}^{K-1} x_{j}^{2}, \quad (29)$$

where the last inequality is due to Cauchy-Schwartz inequality.

Consider $\overrightarrow{W}^{(b)} = (0, 1, \dots, 1)$. Then we have (28) instantiate as

$$x^{T}(A+BB^{T}-\beta\beta^{T})x = \sum_{j=1}^{K-1} p_{j+1}^{(b)}(x_{j}+x_{K+j})^{2} - \left(\sum_{j=1}^{K-1} p_{j+1}^{(b)}(x_{j}+x_{K+j})\right)^{2}$$
$$= p_{1}^{(b)}\sum_{j=1}^{K-1} p_{j+1}^{(b)}(x_{j}+x_{K+j})^{2} + \left(\sum_{j=1}^{K-1} p_{j+1}^{(b)}\right)\sum_{j=1}^{K-1} p_{j+1}^{(b)}(x_{j}+x_{K+j})^{2} - \left(\sum_{j=1}^{K-1} p_{j+1}^{(b)}(x_{j}+x_{K+j})\right)^{2}$$
$$\stackrel{(i)}{\geq} p_{1}^{(b)}\sum_{j=1}^{K-1} p_{j+1}^{(b)}(x_{j}+x_{K+j})^{2} \ge \delta^{2}\sum_{j=1}^{K-1} (x_{j}+x_{K+j})^{2} = \delta^{2}\sum_{j=1}^{K-1} (x_{j}^{2}+x_{K+j}^{2}+2x_{j}x_{K+j})$$
(30)
$$\stackrel{(ii)}{\geq} \delta^{2}\sum_{j=1}^{K-1} (x_{j}^{2}+x_{K+j}^{2}-\frac{2}{3}x_{K+j}^{2}-\frac{3}{2}x_{j}^{2}) = \delta^{2}\sum_{j=1}^{K-1} \left(-\frac{1}{2}x_{j}^{2}+\frac{1}{3}x_{K+j}^{2}\right),$$

where both (i) and (ii) are due to Cauchy-Schwartz inequality.

Consider $\overrightarrow{W}^{(c)} = (1, 0, \dots, 0)$. We have (28) instantiate as

$$x^{T}(A + BB^{T} - \beta\beta^{T})x = p_{1}^{(c)}x_{K}^{2} + \sum_{j=1}^{K-1} p_{j+1}^{(c)}x_{j}^{2} - \left(p_{1}^{(c)}x_{K} + \sum_{j=1}^{K-1} p_{j+1}^{(c)}x_{j}\right)^{2}$$
(31)

Note that under creator-side randomization, each \vec{W} has assignment probability bounded away from zero, and we denote this lower bound as $\eta > 0$. Also recall that each item exposure probability is lower bounded by $\delta > 0$, with $K\delta \leq 1$. Combining (29), (30), (31), with the fact that the Hessian for any \vec{W} is semi-definite, we have

$$\begin{aligned} x^{T}H_{1}x &= \mathbb{E}[x^{T}(A+BB^{T}-\beta\beta^{T})|V,\vec{C}] \\ &\geq P\left(\vec{W}=\vec{W}^{(a)}\right)\delta^{2}\sum_{j=1}^{K-1}x_{j}^{2}+P\left(\vec{W}=\vec{W}^{(b)}\right)\delta^{2}\sum_{j=1}^{K-1}\left\{-\frac{1}{2}x_{j}^{2}+\frac{1}{3}x_{K+j}^{2}\right\} \\ &+P\left(\vec{W}=\vec{W}^{(c)}\right)\left\{p_{1}^{(c)}x_{K}^{2}+\sum_{j=1}^{K}p_{j+1}^{(c)}x_{j}^{2}-\left(p_{1}^{(c)}x_{K}+\sum_{j=1}^{K-1}p_{j+1}^{(c)}x_{j}\right)^{2}\right\} \\ &\geq \eta\delta^{2}\sum_{j=1}^{K-1}x_{j}^{2}+\frac{\eta\delta^{2}}{2}\sum_{j=1}^{K-1}\left\{-\frac{1}{2}x_{j}^{2}+\frac{1}{3}x_{K+j}^{2}\right\}+\frac{\eta\delta^{2}}{2(1-(K-1)\delta)^{2}}\left\{p_{1}^{(c)}x_{K}^{2}+\sum_{j=1}^{K}p_{j+1}^{(c)}x_{j}^{2}-\left(p_{1}^{(c)}x_{K}+\sum_{j=1}^{K-1}p_{j+1}^{(c)}x_{j}\right)^{2}\right\} \\ &=\frac{\eta\delta^{2}}{2}\sum_{j=1}^{K-1}\left\{\frac{1}{2}x_{j}^{2}+\frac{1}{3}x_{K+j}^{2}\right\}+I, \end{aligned} \tag{32}$$

where $I = \frac{\eta \delta^2}{2(1-(K-1)\delta)^2} \left\{ p_1^{(c)} x_K^2 + \sum_{j=1}^{K-1} (p_{j+1}^{(c)} + (1-(K-1)\delta)^2) x_j^2 - \left(p_1^{(c)} x_K + \sum_{j=1}^{K-1} p_{j+1}^{(c)} x_j \right)^2 \right\}$. We next lower bound term I.

$$I \ge \frac{\eta \delta^2}{2(1 - (K - 1)\delta)^2} \left\{ p_1^{(c)} x_K^2 + \sum_{j=1}^{K-1} (1 + p_1^{(c)}) p_{j+1}^{(c)} x_j^2 - \left(p_1^{(c)} x_K + \sum_{j=1}^{K-1} p_{j+1}^{(c)} x_j \right)^2 \right\}$$

$$= \frac{\eta \delta^2}{2(1 - (K - 1)\delta)^2} \left\{ \left(p_1^{(c)} x_K^2 + \sum_{j=1}^{K-1} (1 + p_1^{(c)}) p_{j+1}^{(c)} x_j^2 \right) \left(p_1^{(c)} + \sum_{j=1}^{K-1} \frac{p_{j+1}^{(c)}}{1 + p_1^{(c)}} + \frac{p_1^{(c)} - (p_1^{(c)})^2}{p_1^{(c)} + 1} \right) - \left(p_1^{(c)} x_K + \sum_{j=1}^{K-1} p_{j+1}^{(c)} x_j \right) \right\}$$

$$\stackrel{(i)}{\ge} \frac{\eta \delta^2}{2(1 - (K - 1)\delta)^2} \left(p_1^{(c)} x_K^2 + \sum_{j=1}^{K-1} (1 + p_1^{(c)}) p_{j+1}^{(c)} x_j^2 \right) \left(\frac{p_1^{(c)} - (p_1^{(c)})^2}{p_1^{(c)} + 1} \right)$$

$$\ge \frac{\eta \delta^2}{2(1 - (K - 1)\delta)^2} \left(p_1^{(c)} x_K^2 + \sum_{j=1}^{K-1} (1 + p_1^{(c)}) p_{j+1}^{(c)} x_j^2 \right) \frac{p_1^{(c)} (1 - p_1^{(c)})}{2}}{2}$$

$$\ge \frac{\eta \delta^2 (p_1^{(c)})^2 (1 - p_1^{(c)})}{4(1 - (K - 1)\delta)^2} x_K^2 \ge \frac{\eta \delta^5}{4(1 - (K - 1)\delta)^2} x_K^2, \qquad (33)$$

where (i) is by Cauchy-Schwarz. Putting (32) and (33) together, for any $x = (x_1, \ldots, x_{2K-1})$, we have

$$x^{T}H_{1}x \ge \frac{\eta\delta^{2}}{2} \sum_{j=1}^{K-1} \left\{ \frac{1}{2}x_{j}^{2} + \frac{1}{3}x_{K+j}^{2} \right\} + \frac{\eta\delta^{5}}{4(1 - (K-1)\delta)^{2}} x_{K}^{2}.$$
(34)

Therefore, H_1 has the smallest eigenvalue of greater than or equal to $\frac{\eta\delta^2}{2}\min\left(\frac{1}{3},\frac{\delta^3}{2(1-(K-1)\delta)^2}\right)$ and thus is invertible with bounded inverse.

Regularity of H_2 . We have H_2 is a diagonal matrix with its k-th diagonal entry being

$$H_2(k,k) = \mathbb{E}\left[\mathbb{E}\left[P(k^* = k \mid U, \vec{V}, \vec{W}, \hat{s}_0, \hat{s}_1)\right] \mid U, \vec{V}\right] \ge \delta.$$

As a result, $H_2 \ge \delta \cdot I$ and thus is invertible with bounded inverse.

E.3. Universal Neyman Orthogonality

We show that the debiased estimate ψ , defined in (16), satisfies the universal Neyman orthogonality (Chernozhukov et al. 2019, Foster and Syrgkanis 2023). This property means that the nuisance estimation error only has a second order effect on the debiased estimate – a key property for achieving the asymptotic normality of the debiased estimator.

PROPOSITION 2 (Universal Orthogonality). The debiased estimator ψ defined in (16) is universally orthogonal with respect to the nuisances in the sense that, for any nuisance components $(\tilde{s}_0, \tilde{s}_1, \tilde{z}, \tilde{H})$,

$$\mathbb{E}[\nabla\psi(V,\vec{C},\vec{W},k^*,Y;\tilde{s}_0=s_0,\tilde{s}_1=s_1,\tilde{z}=z,\tilde{H}=H) \mid V,\vec{C}] = 0,$$

where $(V, \vec{C}, \vec{W}, k^*, Y)$ is sampled from the creator-side randomization experiment, and $\nabla \psi$ is the gradient with respect to the nuisances.

Recall that

$$\begin{split} \psi(V,\vec{C},\vec{W},k^*,Y;\tilde{s}_0,\tilde{s}_1,\tilde{z},\tilde{H}) &= \mu(V,\vec{C};\tilde{s}_0,\tilde{s}_1,\tilde{z}) \\ &- \nabla\mu(V,\vec{C};\tilde{s}_0,\tilde{s}_1,\tilde{z})^T\tilde{H}(V,\vec{C};\tilde{s}_0,\tilde{s}_1,\tilde{z})^{-1}\nabla\ell(V,\vec{C},\vec{W},k^*,Y;\tilde{s}_0,\hat{s}_1,\tilde{z}), \end{split}$$

Let $\tilde{h} = (\tilde{s}_0, \tilde{s}_1, \tilde{z}, \tilde{H}^{-1})$ with the ground truth $h = (s_0, s_1, z, H^{-1})$. It suffices to show that

$$\mathbb{E}[\nabla_{\tilde{h}}\psi(V,\vec{C},\vec{W},k^*,Y;\tilde{h}=h) \mid V,\vec{C}] = 0.$$

We have

_

$$\begin{aligned} \frac{\partial \psi(\tilde{h}=h)}{\partial (\tilde{s}_0, \tilde{s}_1, \tilde{z})} = & \nabla \mu(V, \vec{C}; s_0, s_1, z) - \nabla^2 \mu(V, \vec{C}; s_0, s_1, z) H(V, \vec{C}; s_0, s_1, z)^{-1} \nabla \ell(V, \vec{C}, \vec{W}, k^*, Y; s_0, s_1, z) \\ & - \nabla \mu(V, \vec{C}; s_0, s_1, z) H(V, \vec{C}; s_0, s_1, z)^{-1} \nabla^2 \ell(V, \vec{C}, \vec{W}, k^*, Y; s_0, s_1, z). \end{aligned}$$

Then taking the expectation with respect to $(\overrightarrow{W}, k^*, Y)$, we have

$$\begin{split} \mathbb{E}\left[\frac{\partial\psi(\tilde{h}=h)}{\partial(\tilde{s}_{0},\tilde{s}_{1},\tilde{z})} \mid V,\vec{C}\right] &= \nabla\mu(V,\vec{C};s_{0},s_{1},z) \left(I - H(V,\vec{C};s_{0},s_{1},z)^{-1}\mathbb{E}\left[\nabla^{2}\ell(V,\vec{C},\vec{W},k^{*},Y;s_{0},s_{1},z) \mid V,\vec{C}\right]\right) \\ &- \nabla^{2}\mu(V,\vec{C};s_{0},s_{1},z)H(V,\vec{C};s_{0},s_{1},z)^{-1}\mathbb{E}\left[\nabla\ell(V,\vec{C},\vec{W},k^{*},Y;s_{0},s_{1},z) \mid V,\vec{C}\right] \\ &\stackrel{(i)}{=} \nabla\mu(V,\vec{C};s_{0},s_{1},z) \left(I - H(V,\vec{C};s_{0},s_{1},z)^{-1}H(V,\vec{C};s_{0},s_{1},z)\right) = 0, \end{split}$$

where (i) is because the ground truth (s_0, s_1, z) satisfies the first-order optimality condition of loss function $\ell.$ Similarly, we have

$$\mathbb{E}\left[\frac{\partial\psi(\tilde{h}=h)}{\partial(\tilde{H}^{-1})} \mid V, \vec{C}\right] = -\mathbb{E}\left[\nabla\ell(V, \vec{C}, \vec{W}, k^*, Y; s_0, s_1, z) \mid V, \vec{C}\right] \nabla\mu(V, \vec{C}; s_0, s_1, z)^T = 0.$$

Appendix F: Proof of the Asymptotics of the Debiased Estimator

We prove the below theorem to show the asymptotic regularity of the debiased estimator.

THEOREM 1. Suppose that Assumptions 1 & 2 hold. Assume that the data generating process follows the recommender choice model Equation (7) and the viewer response model Equation (8). Suppose that the nuisance estimates are all bounded by the constant C in Assumption 2 and satisfy the convergence rate: $\|\hat{s}_0 - s_0\|_{L_2} + \|\hat{s}_1 - s_1\|_{L_2} + \|\hat{z} - z\|_{L_2} = o(n^{-1/4})$ and $\|\hat{H} - H\|_{L_2} = o(n^{-1/4})$. Then the debiased estimator $\hat{\tau}_n^{DB}$ in (17) is \sqrt{n} -consistent with $\hat{\tau}_n^{DB} - \tau = O_p(n^{-1/2})$ and asymptotically normal with: $\sqrt{n} (\hat{\tau}_n^{DB} - \tau) / \sqrt{\hat{V}_n} \Rightarrow \mathcal{N}(0, 1)$.

Define the oracle estimator $\tilde{\tau}_n^{DB}$ as

$$\tilde{\tau}_n^{DB} = \frac{1}{n} \sum_{i=1}^n \psi\left(V_i, \overrightarrow{C_i}, \overrightarrow{W_i}, k_i^*, Y_i; s_0, s_1, z, H\right),\tag{35}$$

with that

$$\psi(V_i, \vec{C_i}, \vec{W_i}, k_i^*, Y_i; s_0, s_1, z, H) = \mu(V_i, \vec{C_i}; s_0, s_1, z)$$

$$-\nabla \mu(V_i, \vec{C_i}; s_0, s_1, z)^T H(V_i, \vec{C_i}; s_0, s_1, z)^{-1} \nabla \ell(V_i, \vec{C_i}, \vec{W_i}, k_i^*, Y_i; s_0, s_1, z).$$
(36)

Note that

$$\mathbb{E}[\psi(V_i, \overrightarrow{C_i}, \overrightarrow{W_i}, k_i^*, Y_i; s_0, s_1, z, H) \mid V_i, \overrightarrow{C_i}] = \mu\left(V_i, \overrightarrow{C_i}; s_0, s_1, z\right),\tag{37}$$

by the first order condition of (s_0, s_1, z) in ℓ when conditioning on $(V_i, \vec{C}_i, \vec{W}_i)$.

Our proof follows three main steps:

- (i) Show $\hat{\tau}_n^{DB}$ and $\tilde{\tau}_n^{DB}$ have similar asymptotic behavior.
- (ii) Show asymptotic normality of $\tilde{\tau}_n^{DB}$.
- (iii) Show asymptotically consistent variance estimator \widehat{V}_n .

F.1. Step I: Show $\hat{\tau}_n^{DB}$ and $\tilde{\tau}_n^{DB}$ have similar asymptotic behavior.

This part's proof is largely motivated by the proof pattern for Theorem 3.1 in Chernozhukov et al. (2018), except that we need to additionally deal with the correlation among samples due to the shared items in their consideration sets. For notation convenience, define $\theta_0 = (s_0, s_1, z, H)$ and $\hat{\theta}_0 = (\hat{s}_0, \hat{s}_1, \hat{z}, \hat{H})$. Write $Z_i := (V_i, \overrightarrow{C_i}, \overrightarrow{W_i}, k_i^*, Y_i)$. Note that

$$\mathbb{E}\left[\psi\left(Z_{i};\theta_{0}\right)\mid V_{i},\vec{C}_{i},\vec{W}_{i}\right] \stackrel{(i)}{=} \mathbb{E}\left[\mu\left(V_{i},\vec{C}_{i};s_{0},s_{1},z\right)\mid V_{i},\vec{C}_{i},\vec{W}_{i}\right] = \mu\left(V_{i},\vec{C}_{i};s_{0},s_{1},z\right),\tag{38}$$

where (i) is by the first-order optimality of (s_0, s_1, z) in ℓ . We have

$$\sqrt{n} \left| \hat{\tau}_n^{DB} - \tilde{\tau}_n^{DB} \right| \le I_1 + I_2, \tag{39}$$

where

$$\begin{split} I_1 &:= \left\| \frac{1}{\sqrt{n}} \sum_i \left(\psi(Z_i; \hat{\theta}_0) - \mathbb{E}[\psi(Z_i; \hat{\theta}_0)] \right) - \frac{1}{\sqrt{n}} \sum_i \left(\psi(Z_i; \theta_0) - \mathbb{E}[\psi(Z_i; \theta_0)] \right) \right\|, \\ I_2 &:= \left\| \sqrt{n} \left(\mathbb{E}[\psi(Z_i; \hat{\theta}_0)] - \mathbb{E}[\psi(Z_i; \theta_0)] \right) \right\|. \end{split}$$

We now bound I_1 and I_2 respectively.

Bounding
$$I_1$$
. Write $Q_i := \left(\psi(Z_i; \hat{\theta}_0) - \mathbb{E}[\psi(Z_i; \hat{\theta}_0)]\right) - \left(\psi(Z_i; \theta_0) - \mathbb{E}[\psi(Z_i; \theta_0)]\right)$. We have

$$\mathbb{E}[I_1^2] = \frac{1}{n} \mathbb{E}\left[\left(\sum_i Q_i\right)^2\right] = \frac{1}{n} \sum_i \mathbb{E}[Q_i^2] + \frac{1}{n} \sum_i \sum_{j \neq i} \mathbf{1} \left(\overrightarrow{W_i} \cap \overrightarrow{W_j} \neq \emptyset\right) \mathbb{E}[Q_i Q_j]$$

$$= \frac{O(a_n)}{n} \sum_i \mathbb{E}[Q_i^2] = O\left(\frac{a_n}{n}\right) \sum_i \mathbb{E}\left[\left\{\left(\psi(Z_i; \hat{\theta}_0) - \mathbb{E}[\psi(Z_i; \hat{\theta}_0)]\right) - \left(\psi(Z_i; \theta_0) - \mathbb{E}[\psi(Z_i; \theta_0)]\right)\right\}^2\right] \quad (40)$$

$$\leq O\left(\frac{a_n}{n}\right) \sum_i \mathbb{E}\left[\left(\psi(Z_i; \hat{\theta}_0) - \psi(Z_i; \theta_0)\right)^2\right] = O(a_n \epsilon_n^2). \quad (41)$$

Therefore, by Markov inequality, we have $I_1 = O_p(a_n^{1/2}\epsilon_n)$.

Bounding I_2 . Define the function

$$f(r) := \mathbb{E}[\psi(Z_i; \theta_0 + r(\hat{\theta}_0 - \theta_0))] - \mathbb{E}[\psi(Z_i; \theta_0))], \quad r \in (0, 1).$$
(42)

By Taylor expansion, we have

$$f(1) = f(0) + f'(0) + f''(\tilde{r})/2, \quad \text{for some } \tilde{r} \in (0, 1).$$
(43)

Note that by the Neyman orthogonality (shown in E.3), we have f'(0) = 0. With the bounded inverse of Hessian, we have

$$\mathbb{E}[\|f''(\tilde{r})\|] \le \sup_{r \in (0,1)} \|f''(r)\| = O(\epsilon_n^2).$$
(44)

We have

$$I_{2,k} = O_p\left(\sqrt{n}\epsilon_n^2\right). \tag{45}$$

Combining the bound for I_1 and I_2 , we have

$$\sqrt{n}\left(\hat{\tau}_n^{DB} - \tilde{\tau}_n^{DB}\right) = O_p\left(a_n^{1/2}\epsilon_n + n^{1/2}\epsilon_n^2\right) = o_p(1),\tag{46}$$

with $a_n = O(n^{1/4})$ and $\epsilon_n = o(n^{-1/4})$.

F.2. Step II: Show asymptotic normality of $\tilde{\tau}_n^{DB}$.

Recall the data generating process: when a new viewer V_i arrives, the back-end retrieval system firstly generates the consideration set \vec{C}_i . For content items that have shown in previous samples, the treatment status remains unchanged. For content items that haven't appear, we sample the treatment status from i.i.d. Bernoulli randomized trials. This procedure constructs the treatment collection \vec{W}_i . Then given $(V_i, \vec{C}_i, \vec{W}_i)$, the recommender chooses item k_i^* to expose, yielding the viewer outcome Y_i and the observation tuple $Z_i := (V_i, \vec{C}_i, \vec{W}_i, k_i^*, Y_i)$. We now apply the martingale theorem to analyze the asymptotic behavior of $\tilde{\tau}_n$. Denote the σ -field $\mathcal{F}_i := \sigma(Z_1, \ldots, Z_i)$. We have that

$$\mathbb{E}[\psi(Z_i;\theta_0) \mid \mathcal{F}_{i-1}] = \mathbb{E}\left[\mathbb{E}\left[\psi(Z_i;\theta_0) \mid V_i, \vec{C}_i, \vec{W_i}\right] \mid \mathcal{F}_{i-1}\right] \\ = \mathbb{E}\left[\mathbb{E}\left[\mu(V_i, \vec{C}_i; \theta_0) \mid V_i, \vec{C}_i, \vec{W_i}\right] \mid \mathcal{F}_{i-1}\right] = \mathbb{E}\left[\mu(V_i, \vec{C}_i; \theta_0)\right] = \tau.$$

Therefore $\{\psi(Z_i; \theta_0) - \tau\}$ forms a martingale difference sequence with respect to filtration $\{\mathcal{F}_i\}$. We now apply the following result from Hall and Heyde (2014).

PROPOSITION 3 (Martingale Central Limit Theorem, Theorem 3.2, Hall and Heyde (2014)). Let $\{\xi_i\}$ be a martingale difference sequence with respect to filtration $\{\mathcal{F}_i\}$, and let η^2 be an a.s. finite random variable. Suppose that:

$$\max_{i} |\xi_i| \xrightarrow{p} 0, \tag{47}$$

$$\sum_{i} \xi_{i}^{2} \xrightarrow{p} \eta^{2}, \tag{48}$$

$$\mathbb{E}[\max_{i} \xi_{i}^{2}] \text{ is bounded.}$$

$$\tag{49}$$

Then $\sum_{i=1}^{n} X_i \xrightarrow{d} Z$ (stably), where the random variable Z has characteristic function $\mathbb{E}[\exp(-\frac{1}{2}\eta^2 t^2)]$.

Now let's verify the above conditions for $\sqrt{n}(\tilde{\tau} - \tau)$. Define

$$\xi_i = \frac{1}{\sqrt{n}} \left\{ \psi(Z_i; \theta_0) - \tau \right\}.$$

We have $\mathbb{E}[\xi_i | \mathcal{F}_{i-1}] = 0$. By the bounded inverse of Hessian, we have $|\xi_i| = O(n^{-1/2})$, implying (47) and (49). Section F.3 shows that:

$$\widetilde{V}_n := \sum_i \xi_i^2 = \frac{1}{n} \sum_i (\psi_i(Z_i; \theta_0) - \tau)^2 \xrightarrow{p} \eta^2,$$

where $\eta^2 := \operatorname{Var}(\psi_i(Z_i; \theta_0))$ that is a finite number, verifying (48). We thus have

$$\sqrt{n}(\tilde{\tau}_n^{DB} - \tau) = \sum_i \xi_i \xrightarrow{d} \mathcal{N}(0, \eta^2), \tag{50}$$

where z is a Gaussian random variable $\mathcal{N}(0,\eta^2)$ that has characteristic function $\mathbb{E}[\exp(-\frac{1}{2}\eta^2 t^2)]$. Connecting (50) with (46), by Slutsky's theorem, we have

$$\sqrt{n}\left(\hat{\tau}_{n}^{DB}-\tau\right) = \sqrt{n}\left(\hat{\tau}_{n}^{DB}-\tilde{\tau}_{n}^{DB}\right) + \sqrt{n}\left(\tilde{\tau}_{n}^{DB}-\tau\right) \xrightarrow{d} z,\tag{51}$$

yielding that $\sqrt{n}(\hat{\tau}_n^{DB} - \tau)/\eta$ is asymptotically normal and converges to $\mathcal{N}(0, 1)$.

F.3. Step III: variance convergence and connecting \tilde{V}_n to \hat{V}_n .

We start by showing that \widetilde{V}_n converges in probability to $\eta^2 := \operatorname{Var}(\psi_i(Z_i;\theta_0))$. We first argue that $\mathbb{E}[(\psi_i(Z_i;\theta_0) - \tau)^2]$ has the same value, denoted as η^2 . Note that $\mathbb{E}[\psi_i(Z_i;\theta_0)] = \mathbb{E}[\mathbb{E}[\psi_i(Z_i;\theta_0) | \mathcal{F}_{i-1}]] = \tau$. Therefore $\mathbb{E}[(\psi_i(Z_i;\theta_0) - \tau)^2]$ represents the variance of $\psi_i(Z_i;\theta_0)$. Given n samples, let \vec{w} denote the treatment status of all items involved in the n samples. We have that each entry of \mathbf{w} is i.i.d. sampled from Bernoulli distribution with the treated probability being q, for which we abuse notation and denote as $\mathbf{w} \sim \mathcal{B}(q)$. Therefore,

$$\begin{aligned} \operatorname{Var}(\psi_i(Z_i;\theta_0)) &:= \mathbb{E}\left[(\psi_i(Z_i;\theta_0) - \tau)^2 \right] = \mathbb{E}_{\mathbf{w} \sim \mathcal{B}(q)} \left[\mathbb{E}\left[(\psi_i(Z_i;\theta_0) - \tau)^2 \mid \mathbf{w} \right] \right] \\ &= \mathbb{E}_{\overrightarrow{W}_i \sim \mathcal{B}(q)} \left[\mathbb{E}\left[(\psi_i(Z_i;\theta_0) - \tau)^2 \mid \overrightarrow{W}_i \right] \right] \\ &= \mathbb{E}_{(V_i,\overrightarrow{C}_i),\overrightarrow{W}_i \sim \mathcal{B}(q)} \left[\mathbb{E}\left[(\psi_i(Z_i;\theta_0) - \tau)^2 \mid V_i, \overrightarrow{C}_i, \overrightarrow{W}_i \right] \right]. \end{aligned}$$

Recall that $\{(V_i, \vec{C}_i)\}_{i=1}^n$ are sampled i.i.d. from i.i.d. viewer queries, we thus obtain that $\operatorname{Var}(\psi_i(Z_i; \theta_0))$ across *i* have the same value, denoted as η^2 . Similarly, we have that for any *m*-moment $\mathbb{E}[(\psi_i(Z_i; \theta_0) - \tau)^m]$,

they have the same bounded value across sample i, where the boundedness comes from the boundedness of ψ_i .

We thus have $\mathbb{E}[\widetilde{V}_n] = \eta^2$. We proceed to show $\widetilde{V}_n \xrightarrow{p} \eta^2$. By Markov inequality, given any $\epsilon > 0$, we have

$$P\left(|\widetilde{V}_n - \eta^2| \ge \epsilon\right) \le \frac{\operatorname{Var}(\widetilde{V}_n)}{\epsilon^2}.$$
(52)

Note that

$$\operatorname{Var}\left(\widetilde{V}_{n}\right) = \frac{\sum_{i=1}^{n} \mathbb{E}\left[\left(\psi_{i}(Z_{i};\theta_{0}) - \tau\right)^{4}\right]}{n^{2}} + \frac{\sum_{i \neq j} \mathbf{1}\left(\overrightarrow{W}_{i} \cap \overrightarrow{W}_{j} \neq \emptyset\right) \operatorname{Cov}\left\{\left(\psi_{i}(Z_{i};\theta_{0}) - \tau\right)^{2}, \left(\psi_{j}(Z_{j};\theta_{0}) - \tau\right)^{2}\right\}}{n^{2}}$$
$$\stackrel{(i)}{=} O\left(\frac{n}{n^{2}}\right) + \frac{na_{n}}{n^{2}} = O\left(\frac{a_{n}}{n}\right) = o(n^{-3/4}) = o(1),$$

where (i) uses the boundedness of $\mathbb{E}[(\psi_i(Z_i;\theta_0)-\tau)^4]$ and the Cauchy-Schwartz inequality such that $\operatorname{Cov}\{(\psi_i(Z_i;\theta_0)-\tau)^2,(\psi_j(Z_j;\theta_0)-\tau)^2\} \leq \sqrt{\mathbb{E}[(\psi_i(Z_i;\theta_0)-\tau)^4]\mathbb{E}[(\psi_j(Z_j;\theta_0)-\tau)^4]}.$ Continuing (52), we have

$$P\left(|\widetilde{V}_n - \eta^2| \ge \epsilon\right) \le \frac{\operatorname{Var}(\widetilde{V}_n)}{\epsilon^2} = o(1),$$

implying that $\widetilde{V}_n \xrightarrow{p} \eta^2$. Therefore, $\sqrt{n} \left(\hat{\tau}_n^{DB} - \tau \right) / \eta$ converges in distribution to $\mathcal{N}(0, 1)$.

We now show that \hat{V}_n also converges in probability to η^2 . We have

$$\begin{split} |\widehat{V}_{n} - \widetilde{V}_{n}| &= \left| \frac{1}{n} \sum_{i} \left(\psi(Z_{i}; \hat{\theta}_{0}) - \psi(Z_{i}; \theta_{0}) - \hat{\tau}_{n}^{DB} + \tau \right) \left(\psi(Z_{i}; \hat{\theta}_{0}) + \psi(Z_{i}; \theta_{0}) - \hat{\tau}_{n}^{DB} - \tau \right) \right| \\ &\stackrel{(i)}{\leq} O(1) \cdot \left\{ \frac{1}{n} \sum_{i} \left| \psi(Z_{i}; \hat{\theta}_{0}) - \psi(Z_{i}; \theta_{0}) \right| + \frac{1}{n} \sum_{i} \left| \hat{\tau}_{n}^{DB} - \tau \right| \right\} \\ &= O_{p}(\|\hat{\theta}_{0} - \theta_{0}\|_{L_{2}}) + O_{p}(n^{-1/2}) = O_{p}(\epsilon_{n}) + O_{p}(n^{-1/2}) = o_{p}(1), \end{split}$$

where (i) is by the boundedness of $\psi(Z_i; \hat{\theta}_0)$ and $\psi(Z_i; \theta_0)$.

Therefore,

$$|\widehat{V}_n - \eta^2| \le |\widehat{V}_n - \widetilde{V}_n| + |\eta^2 - \widetilde{V}_n| = o_p(1).$$

Combining the above with $\sqrt{n}(\hat{\tau}^{DB} - \tau)/\eta \xrightarrow{d} \mathcal{N}(0, 1)$, by Slutsky's theorem, we have $\sqrt{n}(\hat{\tau}^{DB} - \tau)/\sqrt{\hat{V}_n} \xrightarrow{d} \mathcal{N}(0, 1)$, concluding the proof.

Appendix G: Discussion on Model Misspecification

Our semi-parametric estimator, $\hat{\tau}_n^{DB}$, differs from the fully non-parametric double/debiased estimators such as the Augmented Inverse Propensity Weighting (AIPW) in the causal inference literature (Imbens 2004), where both the outcome model and the propensity score are non-parametrically defined. In contrast, the recommender choice model (described in Section 3) adopts a semi-parametric multinomial logit (MNL) structure by integrating a non-parametric component that captures the viewer and item heterogeneity, and a parametric component that specifies the role of the treatment. As a result, the estimation and inference guarantees provided by our estimator, $\hat{\tau}_n^{DB}$, depend on the choice model as specified in Eq. (7). If the model is misspecified, the asymptotic properties and results presented in Section 4.2 may no longer hold.

More specifically, the model makes two parametric assumptions. The first assumption is that the choice probability can be written as a logit function. This is not a strong assumption since the softmax function is capable of approximating any categorical distribution (Cervera et al. 2021). Since the recommendation process is stochastic (see discussions in Section 2), we can reasonably approximate the probability of an item being recommended with the logit functional form.

The second assumption is arguably stronger. In our choice model (7), we specify the deterministic part of the score or "utility" using this functional form:

$$\bar{S}_{i,k} = s_0(V_i, C_{i,k}) + W_{i,k} \cdot s_1(V_1, C_{i,k}).$$
(53)

The model specifies that the deterministic part of the score for an item depends only on its characteristics $C_{i,k}$ and its treatment status $W_{i,k}$. Alternatively, one can write down a fully general model, denoted by $s^*(\cdot) = \{s_1^*(\cdot), \ldots, s_K^*(\cdot)\}$, where the deterministic part of the score of each item k depends on the characteristics and treatment status of all items in the consideration set:

$$\bar{S}_{i,k} = s_k^*(V_i, \overrightarrow{C_i}, \overrightarrow{W_i}). \tag{54}$$

The fully general model s^* is more flexible. Consider a consideration set of size K, our choice model in Eq. (53) predicts two sets of scores, K under the control condition (one baseline score s_0 for each item) and K under the treatment condition (s_0 + score uplift s_1 for each item). Under the fully general model, however, the model predicts $2^K K$ scores. This is because the score for each item depends on the treatment assignment of all K items, resulting in 2^K possible combinations.

Suppose the true model is s^* and we impose a more restrictive functional form as in Eq. (53). Under such model misspecification, we essentially approximate the true model within our specified model class:

$$(s_0, s_1) = \underset{\tilde{s}_0, \tilde{s}_1}{\operatorname{arg\,min}} \mathbb{E}\left[\ell_1(V_i, \overrightarrow{C_i}, \overrightarrow{W_i}, k_i^*; \tilde{s}_0, \tilde{s}_1)\right].$$

$$(55)$$

The expectation is taken with respect to $(V_i, \vec{C_i}, \vec{W_i}, k_i^*)$, where the exposed item k_i^* comes from the true model s^* . The proposed debiased estimator consistently estimates the best-in-class choice model parameterized by (s_0, s_1) in Eq. (55), achieving the smallest asymptotic variance (see more discussion on the robustness of doubly robust estimator under model misspecification in Chernozhukov et al. 2019).

Empirically speaking, we are not concerned about model misspecification because of the fit of the recommender choice model (see Section 5.2). Since the current model structure approximates the behavior of the actual recommender very well, there is less of a concern about the flexibility of the score function as defined in Eq. (53). Theoretically speaking, the scoring of a recommender system tends to depend on the fit of the viewer-content pair. The other items of the consideration set typically would not play a role.

In certain applications, if one has prior knowledge of how treatment affects recommendations, the score function can be adjusted based on such knowledge. This can lead to a more flexible function form of the treatment vector \vec{W} beyond the additive linear form as specified in (53), but not fully flexible with 2^{K} combinations of possible treatment assignments. It is worth noting that as long as the score function maintains a parametric relationship with the treatment status, the framework outlined in this paper can be directly applied to construct a debiased estimator and obtain the estimation and inference results.