Local symmetries in partially ordered sets

Christoph Minz*

June 17, 2024

Abstract

Partially ordered sets (posets) have a universal appearance as an abstract structure in many areas of mathematics. Though, even their explicit enumeration remains unknown in general, and only the counts of all partial orders on sets of up to 16 unlabelled elements have been calculated to date, see sequence A000112 in the OEIS.

In this work, we study automorphisms of posets in order to formulate a classification by local symmetries. These symmetries give rise to a division operation on the set of all posets and lead us to the construction of symmetry classes that are easier to characterise and enumerate. Additionally to the enumeration of symmetry classes, I derive polynomial expressions that count certain subsets of posets with a large number of layers (a large height). As an application in physics, I investigate local symmetries (or rather their lack of) in causal sets, which are discrete spacetime models used as a candidate framework for quantum gravity.

1 Introduction

Counting is perhaps the most basic mathematical concept and we may find unexpected connection between different mathematical structures whenever they have an identical enumeration. A fairly ubiquitous structure in various fields of mathematics are partial orders. For example, all subspaces of a von Neumann algebra are partially ordered under inclusion, which implies a partially ordered set of the corresponding subspace projections. Another example are the incidence relations between edges and vertices of a graph, or even the incidences of any (abstract) geometric shape with edges, vertices and higher faces has a representation as a partially ordered set. In physics, the causal structure of any spacetime manifold is a partial order, and as an approach to quantum gravity, causal set theory is based on discrete counterparts to spacetime manifolds where any interval is finite and it is hypothesised that continuum spacetimes are corresponding to or "emerging" from causal sets [Bom+87; BG91; BHS08].

The number of distinct partially ordered sets (posets) on a set of n unlabeled elements is listed as sequence A000112 in the OEIS [Slo+23] (identical to the number of T_0 topological spaces), but only the first sixteen terms of the sequence have been determined to date [BM02], and enumeration results for certain subsets have been studied [Kla69; But72]. Although the asymptotic behaviour of the sequence is known [KR75; Dha80; PST01a; PST01b], finding analytic expression for the number of partial orders remains challenging. This work shall provide a classification of posets by analysing repeating substructures forming local symmetries and making use of the concept of retracts [DR79; DR81]. Although some of the findings might already be known, I use the characterisation of posets with and without symmetries to formulate explicit expressions for partial enumerations, and I illustrate an application in the quantum gravity programme of causal set theory.

^{*}Institut für theoretische Physik, Universität Leipzig, Brüderstraße 16, 04103 Leipzig, Germany; e-mail: christoph.minz@itp.uni-leipzig.de

Any local subset (interval) of a causal set (as a discrete model of a spacetime manifold in physics) is a finite poset. While causal sets that correspond to a given spacetime manifold are usually generated via a random process (called sprinkling), the central idea of the causal set programme is to describe spacetime manifolds as continuous counterparts of the fundamentally discrete structures that are (locally finite) posets. The asymptotic description of generic posets in [KR75], however, showed that these are most likely of a type that is very different from the causal sets obtained by a sprinkling process. Recently, it was shown that the typical Kleitman-Rothschild orders could be sufficiently suppressed in the "sum-over-history" approach leaving those causal sets that are more likely representing discrete spacetimes [CCS23a]. Below, we consider properties to help distinguish between generic posets and those that are more relevant for causal set theory.

The work is structured as follows. First, we recall definitions for the characterization of elements, subsets and entire posets. The definition of local symmetries in posets leads us to the characterisation of and allocation of posets into symmetry classes and their enumeration. Then I derive explicit formulae for the sizes of some subsets of posets with a large number of layers, finally, I comment on the application in causal set theory.

2 Notation and terminology

reflexive:

In this section, we recall some basic notation and terminology that is being used throughout this work. Some readers might be familiar with partial orders, though, I recommend to skim over this section, at least, as I also include some new or less common terms here.

As one may find different notations in the literature, I state the definition of a partially ordered set first.

Definition 2.1. A partially ordered set (*poset*) is a set P with a partial ordering \leq that is (for all $a, b, c \in P$)

transitive:
$$(a \le b \text{ and } b \le c) \Rightarrow a \le c$$
, (1a)

antisymmetric: $(a \le b \text{ and } b \le a) \Rightarrow a = b$, (1b)

$$a \le a$$
. (1c)

For short, I denote a poset by its set P keeping the partial ordering \leq implicit as long as it is not ambiguous. Recall that any isomorphism between (sub)posets A and B, written as $A \xrightarrow{\sim} B$, is order-preserving and order-reflecting. In this work, I consider partial orders on sets of unlabelled elements only, and the set of all posets (up to isomorphisms) is denoted by \mathfrak{P} . The set of all isomorphisms from a poset $P \in \mathfrak{P}$ to itself forms the automorphism group $\operatorname{Aut}(P)$. A subset A of the poset P is fixed under an automorphism $\alpha \in \operatorname{Aut}(P)$ if $\forall a \in A : \alpha(a) = a$.

Definition 2.2 (Element properties). Let P be a poset. The sets of elements preceding, succeeding and related to $a \in P$ are denoted by

$$J^{-}(a) := \{ c \in P \mid c \le a \} ,$$
 (2a)

$$J^{+}(a) := \{ c \in P \mid a \le c \} ,$$
(2b)

respectively, and we have the strict version $J_*^{\pm}(a) = J^{\pm}(a) \setminus \{a\}$. An element *a* is **covered by** *b* and *b* is **covering** *a* written as

$$a \ll b :\Leftrightarrow (a < b \text{ and } \nexists c \in P : a < c < b).$$
 (3)

Let $L^{\pm}(a) \subseteq J^{\pm}(a)$ be the set of all *linked* elements, those covered by and covering a, respectively. Elements a and b are **zic-zac connected** if there is a zic-zac path relating them, that is

$$\exists c_1, c_2, \dots c_n \in P: \qquad a \ge c_1 \le c_2 \dots \le c_n \ge b,$$
(4)

and the elements are **separated** (also *independent* or *incomparable*).

Note that any two related elements are zic-zac connected, while separated elements are only zic-zac connected if they are part of the same poset component.

Remark. Separation between elements describes a symmetric binary relation to be understood as an irreflexive locality relation [GPZ23].

For the application in physics, we consider

Definition 2.3. A causal set (*causet*) is a poset P with an order that is (for all $a, b \in P$)

locally finite:
$$\left| J^+(a) \cap J^-(b) \right| < \infty$$
. (5)

For any element $a \in P$ of a causal set, the sets $J^{\mp}(a)$ are also referred to as *past and future of* a, respectively.

Any subset of a poset is a subposet with the restricted partial order.

Definition 2.4 (Subset properties). A chain is a totally ordered subset of a poset. An **antichain** is a subset of a poset that contains only pairwise separated elements. A subset A of a poset P is **linked** if for all $a, b \in A$:

$$(a < b \Rightarrow a \ll b) \text{ or } (\exists c \in J^+(a) \cap J^-(b) : c \in A).$$
(6)

It is connected if all pairs of elements $a, b \in A$ are zic-zac connected. It is a maximally ordered subset of P if the number of ordered pairs

$$\left| \{ (a,b) \in A \times A \mid a < b \} \right|,\tag{7}$$

is maximal among all subsets of P with cardinality |A|.

Two subsets $A, B \subset P$ are **separated** (components) if all pairs of elements $a \in A$ and $b \in B$ are separated.

In this work, I represent any specific poset by its Hasse diagram, the (directed) graph with vertices for all elements and edges (links) for all covering relations (pointing towards the top of the page). All diagrams are generated with the $\text{LAT}_{\text{E}}X$ -package "causets" [Min20]. For example, all posets with a cardinality n up to n = 4 are

$$\mathfrak{P}_1 = \{\bullet\} , \tag{8a}$$

$$\mathfrak{P}_2 = \{(\bullet \bullet), \mathfrak{l}\}, \tag{8b}$$

$$\mathfrak{P}_{3} = \left\{ \left(\cdot \cdot \cdot \right), \left(\ddagger \cdot \right), \Lambda, \Psi, \ddagger \right\}, \qquad (8c)$$

where posets with separated components are parenthesised. Such parenthesised posets can be constructed from the components by parallel composition. **Definition 2.5.** The series/parallel composition of two posets P and Q, denoted as $P \lor Q$ and $P \sqcup Q$, respectively, is a poset R that is the disjoint union of P and Q such that any two elements both from P or both from Q have the same ordering in R. Any two elements $p \in P$ and $q \in Q$ fulfill p < q in R for the series composition, respective $a \nleq b$ and $b \nleq a$ for the parallel composition.

Definition 2.6. The opposite poset of a poset (P, \leq) is the poset (P, \leq^{op}) such that

$$\forall a, b \in P: \qquad b \leq^{\mathrm{op}} a \Leftrightarrow a \leq b.$$
(9)

A poset is **self-opposite** if it is isomorphic to its opposite.

As a useful tool in the classification, we define a function on the set of posets.

Definition 2.7. The number of layers of a poset P, written as $\ell(P)$, is the length (cardinality) of its longest chain. The layer of an element $a \in P$ is the number of layers of its preceding subset, $\ell_P(a) := \ell(J^-(a))$.

Note the similar term *height* of a poset is one less than the number of layers.

3 Local symmetries in posets

As a pre-cursor to the definition of the local symmetries, note the following fact about subsets of strictly preceding and succeeding elements.

Lemma 3.1 (Common links). Let P be a poset. For all elements $a, b \in P$,

$$\left(J_*^{\pm}(a) = J_*^{\pm}(b)\right) \Leftrightarrow \left(L^{\pm}(a) = L^{\pm}(b)\right).$$
⁽¹⁰⁾

Proof. If the left hand side holds, then we have, for every $c \in J^{\pm}_{*}(a) = J^{\pm}_{*}(b)$,

$$(c \in L^{\pm}(a)) \Leftrightarrow \left(J_{*}^{\pm}(a) \cap J_{*}^{\pm}(c) = \emptyset\right) \Leftrightarrow \left(J_{*}^{\pm}(b) \cap J_{*}^{\pm}(c) = \emptyset\right) \Leftrightarrow \left(c \in L^{\pm}(b)\right), \tag{11}$$

so $L^{\pm}(a) = L^{\pm}(b)$. If the right hand side holds, then the left hand side follows by transitivity of the partial order.

3.1 Symmetric elements in antichains

For the introduction of local symmetries, we start with the simplest case.

Definition 3.1. Let P be a poset. Two elements $a, b \in P$ are singleton-symmetric if

$$L^{\pm}(a) = L^{\pm}(b) \,. \tag{12}$$

For any pair of singleton-symmetric elements $a, b \in P$, also $J_*^{\pm}(a) = J_*^{\pm}(b)$ by Lemma 3.1. Immediate from the definition is that the singleton-symmetry is an equivalence relation (reflexive, symmetric and transitive). Since each pair of elements in an equivalence class has the same order relations to the elements of the other equivalence classes, the quotient inherit the given partial order of the original poset P. The quotient of P by all singleton-symmetries is isomorphic to a *retracted* poset or *retract*, written as $P \oslash \bullet$.

Example 3.1. All antichain posets with more than one element are singleton-symmetric,

$$\bullet = (\bullet \bullet) \oslash \bullet = (\bullet \bullet \bullet) \oslash \bullet = (\bullet \bullet \bullet \bullet) \oslash \bullet = \dots$$
 (13)

Note that the subgroup of automorphisms of an *n*-antichain is isomorphic to the permutation group S_n .

Example 3.2. Similarly, the 2-chain is the retracted poset of

$$\mathbf{I} = \Lambda \oslash \mathbf{\cdot} = \mathbf{v} \oslash \mathbf{\cdot} = \mathbf{\Lambda} \oslash \mathbf{\cdot} = \mathbf{v} \oslash \mathbf{\cdot} = \mathbf{v} \oslash \mathbf{\cdot} = \dots$$
(14)

Example 3.3. The quotient by the singleton-symmetry distributes over parallel compositions of non-singleton posets (and similarly for series compositions) as in

$$\left(\bigwedge \ \sqcup \ \bigwedge \ \sqcup \ \bigvee \ \right) \oslash \cdot = \left(\bigwedge \ \odot \ \cdot \right) \sqcup \left(\bigwedge \ \odot \ \cdot \right) \sqcup \left(\bigwedge \ \odot \ \cdot \right) \sqcup \left(\bigwedge \ \odot \ \cdot \right)$$
(15a)

$$= \bigvee \sqcup \bigvee \sqcup \sqcup \iota . \tag{15b}$$

We will come back to this example after the introduction of more general symmetries.

=

3.2 Cyclic symmetries

The automorphisms that act only on a finite number of elements constitute finite cycles in the automorphism group leading us to the following generalisation.

Definition 3.2. Let P be a poset, Q be a finite poset, and $r \in \mathbb{N}$, $r \geq 2$. For an automorphism $\sigma \in \operatorname{Aut}(P)$, let $\Sigma(\sigma) \subseteq P$ denote the subset of all elements that are not fixed by σ . A (Q, r)-generator is an automorphism $\sigma \in \operatorname{Aut}(P)$ under the following conditions: there exists a sequence of r subsets $S_i \subset \Sigma(\sigma)$, $i \in \{0, \ldots, r-1\}$ with $S_i \cong Q$, and they are the smallest, maximally ordered subsets of $\Sigma(\sigma)$ with

$$\sigma(S_i) = S_{i+1 \mod r}, \qquad \qquad \ell(S_i) = \ell(\Sigma(\sigma)), \qquad \qquad \bigcup_{i=0}^{r-1} S_i = \Sigma(\sigma) \qquad (16)$$

for all $0 \le i < r$. Any distinct pair of these subsets (S_i, S_j) is (Q, r)-symmetric.

Two elements $a, b \in P$ are (Q, r, 0)-symmetric if a = b. They are (Q, r, 1)-symmetric if there exist subsets $A, B \subset P$ that are (Q, r)-symmetric with (Q, r)-generator σ such that $a \in A$ and $b = \sigma^q(a) \in B$ for some $1 \leq q < r$. Recursively setting $n = 2, 3, \ldots$, the elements are (Q, r, n)-symmetric if they are not (Q, r, j)-symmetric for any j < n, but there exists some $c \in P$ and j < n such that a is (Q, r, j)-symmetric to c and b is (Q, r, n - j)-symmetric to b.

For short, a is (Q, r)-symmetric to b if there exists an $n \in \mathbb{N}_0$ such that they are (Q, r, n)-symmetric. For a linked subset $A \cong Q$, the (Q, r)-symmetry group of A is the largest subgroup of (Q, r)-generators such that its action fixes every element $b \in P$ if there exists no element $a \in A$ that is (Q, r)-symmetric to b. The (Q, r)-symmetry set of A is the complement of all elements fixed under the action of the (Q, r)-symmetry group of $A \cong Q$.

Note that the condition of *smallest, maximally ordered* in the definition of a (Q, r)-generator is not strictly necessary, though it reduces the possible representations of an automorphism with a repeated finite poset Q.

Because a (Q, r)-generator is an automorphism $\sigma \in \operatorname{Aut}(P)$ that fixes all elements in the complement of $\Sigma(\sigma)$, any two elements $a, b \in \Sigma(\sigma)$ also share the same (linked) preceding and succeeding elements in $P \setminus \Sigma(\sigma)$,

$$L^{\pm}(a) \smallsetminus \Sigma(\sigma) = L^{\pm}(b) \smallsetminus \Sigma(\sigma), \qquad (17a)$$

$$J^{\pm}(a) \smallsetminus \Sigma(\sigma) = J^{\pm}(b) \smallsetminus \Sigma(\sigma).$$
(17b)

This is a generalisation of the singleton-symmetry, see (12), so the singleton-symmetry can be seen as the special case for $Q = \cdot$ and r = 2.

The (Q, r)-symmetry groups are normal subgroups of automorphisms as (Q, r)-symmetric elements form equivalence classes. For a poset P, the retract $P \otimes_r Q$ is the poset isomorphic to the quotient \tilde{P} that retracts the equivalence classes under all (Q, r)-symmetries such that for all $A, B \in \tilde{P}$:

$$A \leq_{\tilde{P}} B \Leftrightarrow \exists a \in A, b \in B : a \leq_{P} b.$$
⁽¹⁸⁾

If r = 2, a (Q, 2)-symmetry is a reflection of subsets isomorphic to Q, so we say Q-symmetric for short, and drop the index on the division operator, $P \oslash Q$.

Example 3.4. For the parallel composition in Example 3.3, consider further quotients by 2-chainand singleton-symmetries,

$$\left(\bigwedge \sqcup \bigwedge \sqcup \mathfrak{l}\right) \oslash \mathfrak{l} = \mathfrak{l} \sqcup \Lambda \sqcup \mathfrak{l}, \qquad (19a)$$

$$\left(\mathbf{I} \sqcup \mathbf{\Lambda} \sqcup \mathbf{I}\right) \oslash \bullet = \left(\mathbf{I} \quad \mathbf{I} \quad \mathbf{I}\right), \tag{19b}$$

$$(\mathbf{I} \ \mathbf{I} \ \mathbf{I}) \oslash \mathbf{I} = \mathbf{I}.$$
(19c)

Note that the elements of \bigwedge are 1-symmetric and (1,3)-symmetric, both with the 2-chain as retract.

Example 3.5. The smallest poset with a 3-cycle but without a 2-cycle is

$$\bigotimes \bigotimes \otimes_3 \mathbf{i} = \mathbf{i}, \qquad (20)$$

where the diagonal edge across the central element in the Hasse diagram links the corner elements to each other but does *not* link them to the centre.

From the definition, it might be obvious to a groupologist that

Proposition 3.2. The automorphism group of a finite poset P is generated by (Q, r)-symmetries for some set of pairs (Q, r).

Proof. Let $\sigma \in \operatorname{Aut}(P)$ be any non-trivial automorphism. Since any automorphism is part of a cycle that can be represented in a cycle graph of the group, there is some integer $r \geq 2$ such that $\sigma^r = 1_{\operatorname{Aut}(P)}$ (at least as long as P is finite). As in the definition, we denote the set of those elements that are not fixed by σ as $\Sigma(\sigma) \subseteq P$. Consider a sequence of smallest, maximally ordered subsets $S_i \subset \Sigma(\sigma)$ such that (16) holds for all $i \in \{0, 1, \ldots, r-1\}$. Such a sequence can be found by selecting a longest chain S_0 in $\Sigma(\sigma)$ and adding more elements to S_0 , keeping it maximally ordered, until the sequence $(S_0, S_1, \ldots, S_{r-1})$ with $S_i = \sigma^i(S_0)$ covers $\Sigma(\sigma)$. Repeating this process for all choices of longest chains and maximally ordered extensions leads to a choice that minimises $|S_0|$. By this construction, all of the subsets S_i are isomorphic to each other and they are isomorphic to a (finite) poset Q. So the automorphism σ is part of a (Q, r)-symmetry of P. For a finite poset P, we can do this construction for all automorphisms, so the automorphisms corresponding to (Q, r)-symmetries form the entire automorphism group.

3.3 Local symmetry properties

To get towards a classification of posets by these symmetries, we define the following.

Definition 3.3. For a finite poset Q and $r \in \mathbb{N}$, a poset P is **locally** (Q, r)-symmetric and (Q, r)-retractable to the poset \hat{P} if $\hat{P} = P \otimes_r Q \neq P$. The poset P is **locally symmetric** if there exists some finite poset Q and $r \geq 2$ such that P is locally (Q, r)-symmetric, P is retractable to the poset \hat{P} (the retract of P) if there exist some sequence of (Q_i, r_i) -symmetries such that $\hat{P} = P \otimes_{r_1} Q_1 \otimes_{r_2} Q_2 \otimes_{r_3} \ldots \neq P$, and P is **locally unsymmetric** if it is not locally symmetric.

Note that finite, locally unsymmetric posets have a trivial automorphism group.

Example 3.6. All finite chain posets (total orders) are locally unsymmetric, because there is only one element per layer.

Remark (Algebraic structure). Let $\mathfrak{P}_{[m,n]} := \{P \in \mathfrak{P} \mid m \leq |P| \leq n\}$. For any $n \geq 1, r \geq 2$, $(\mathfrak{P}_{[1,n]}, \oslash_r)$ is a magma and the locally unsymmetric posets in $\mathfrak{P}_{[m,n]}$ with $m = \lfloor \frac{n}{r} \rfloor + 1$ are right identities.

As we want to classify (especially finite) posets by their local symmetries, it is of avail to reduce the number of relevant local symmetries, because pairs of elements can be part of different local symmetries simultaneously.

Example 3.7. Any series compositions of n antichains (each with more than one element) is k-chain-retractable to the n-chain for any $k \leq n$, like

So we make a distinction:

Definition 3.4. Let P be a locally (Q, r)-symmetric poset (for some finite poset Q and $r \ge 2$) and let $S \subseteq P$ be a corresponding (Q, r)-symmetry set (for some subset $A \subset P$). The (Q, r)symmetry of S is **composite** if there exists some finite poset $\hat{Q}, \hat{r} \ge 2$, and a corresponding (\hat{Q}, \hat{r}) -symmetry set $\hat{S} \subseteq S$ such that

$$\hat{Q} \subset Q$$
, or (22a)

$$\hat{Q} = Q \text{ and } \hat{r} < r.$$
 (22b)

It is **prime** if it is not composite.

Our main focus shall lie on prime (Q, r)-symmetries, for which all (Q, r)-symmetric subsets are linked. To retract only prime (Q, r)-symmetries, we write a prime as in $P \otimes'_r Q$. In Example 3.7, the only prime retraction is by the singleton-symmetry.

As a consequence of (Q, r)-generators being elements of the automorphism group of a poset, the length of the longest chain subset in the poset and its (Q, r)-retracted version is always the same.

Proposition 3.3 (Common layers). For any poset P, finite poset Q and $r \in \mathbb{N}$, the symmetry quotient P/(Q,r) is layer-preserving,

$$\forall E \in P/(Q, r) : \forall e \in E : \qquad \ell_{P/(Q, r)}(E) = \ell_P(e) . \tag{23}$$

This implies that any locally (Q, r)-symmetric poset P and the retract $P \otimes_r Q$ have the same number of layers, $\ell(P) = \ell(P \otimes_r Q) = \ell(P \otimes Q)$.

For any finite, locally unsymmetric poset Q, the elements of $\bigsqcup_{i=1}^{n} Q$ (with $n \ge 2$) are all pairwise Q-symmetric. More generally, the definition of (Q, r)-symmetric elements simplifies whenever there is some k such that two (Q, r)-symmetric elements are at most (Q, r, k)-symmetric. As an example, consider 2-chain-symmetric elements.

Lemma 3.4. If two elements in a poset P are (1, 2, k)-symmetric, then $k \in \{0, 1\}$.

Proof. Let $a \sim_1 b$ mean that a is (1, 2, 1)-symmetric to b. The only non-trivial part is to show that, for distinct elements $a, b, c \in P$, $a \sim_1 b$ and $b \sim_1 c$ implies $a \sim_1 c$.

By Proposition 3.3, the elements a, b and c have to be on the same layer within P as well as on the same layer within the subsets isomorphic to Q. This means that they are either all on the first or all on the second layer in the 2-chain subsets. Since the 2-chain is self-opposite, let us assume that all elements are on the first level of the 2-chain subsets, otherwise the arguments have to be made with the opposite of P and all signs in the link subsets are reversed.

Let $A = \{a, \hat{a}\}$ and $B_a = \{b, \hat{b}_a\}$ such that $a \ll \hat{a}, b \ll \hat{b}_a, \hat{a} \neq \hat{b}_a$ and the corresponding automorphism α such that $\alpha(A) = B_a$. Similarly, let $B_c = \{b, \hat{b}_c\}$ and $C = \{c, \hat{c}\}$ such that $b \ll \hat{b}_c, c \ll \hat{c}$ and let β be the corresponding 2-chain-reflection such that $\beta(B_c) = C$. If and only if $a < b_a$, then $b < \hat{a}$ and the 2-chain-reflection is not prime but decomposes into two singletonsymmetries. It is obvious from the definition that the singleton-symmetry has a commuting subgroup of automorphisms, so all these automorphisms are reflections. For the remaining part, we restrict to the case that the symmetries $a \sim_1 b$ and $b \sim_1 c$ are prime, so

$$L^{-}(a) = L^{-}(b), \qquad \qquad L^{+}(a) \smallsetminus \{\hat{a}\} = L^{+}(b) \smallsetminus \{\hat{b}_{a}\}, \qquad (24a)$$

$$L^{-}(\hat{a}) \smallsetminus \{a\} = L^{-}(\hat{b}_{a}) \smallsetminus \{b\}, \qquad L^{+}(\hat{a}) = L^{+}(\hat{b}_{a}), \qquad (24b)$$
$$L^{-}(b) = L^{-}(c), \qquad L^{+}(b) \smallsetminus \{\hat{b}_{c}\} = L^{+}(c) \smallsetminus \{\hat{c}\}, \qquad (24c)$$

(c),
$$L^+(b) \smallsetminus \{\hat{b}_c\} = L^+(c) \smallsetminus \{\hat{c}\}, \qquad (24c)$$

$$L^{-}(\hat{b}_{c}) \smallsetminus \{b\} = L^{-}(\hat{c}) \smallsetminus \{c\}, \qquad \qquad L^{+}(\hat{b}_{c}) = L^{+}(\hat{c}). \qquad (24d)$$

so $L^{-}(a) = L^{-}(c)$. For the other subsets, there are two cases to consider, with the simplest examples I I and \bigwedge , respectively.

In the separated case, $\hat{b}_a = \hat{b}_c$, the subsets in (24a) and (24c), as well as (24b) and (24d) are pairwise identical. So $a \sim_1 c$ given the 2-chain-symmetric subsets $A = \{a, \hat{a}\}$ with $a \ll \hat{a}$ and $C = \{c, \hat{c}\}$ with $c \ll \hat{c}$.

In the crossing case, $\hat{b}_a \neq \hat{b}_c$, the first layer relations (24c) and (24a) imply $\hat{b}_a \in L^+(c) \smallsetminus \{\hat{c}\}$ and $\hat{b}_c \in L^+(a) \setminus \{\hat{a}\}$, respectively. Then the second layer (24b) and (24d) yield $c \in L^-(\hat{a})$, $\hat{a} \in L^+(c)$ and $a \in L^-(\hat{c}), \hat{c} \in L^+(a)$. So we find that $\hat{a} = \hat{c}$, because $\hat{a}, \hat{c} \notin L^+(b)$, and $\hat{a} \neq \hat{c}$ would lead to a contradiction at the first layer (24a), (24c) otherwise. With (24c) and (24a), and taking the union with $\{\hat{a}\} = \{\hat{c}\}$, we get

$$\left(L^+(a) \smallsetminus \{\hat{a}, \hat{b}_c\}\right) \cup \{\hat{a}\} = \left(L^+(c) \smallsetminus \{\hat{b}_a, \hat{c}\}\right) \cup \{\hat{c}\}$$

$$(25a)$$

$$L^{+}(a) \smallsetminus \{b_{c}\} = L^{+}(c) \smallsetminus \{b_{a}\}.$$
 (25b)

Similarly, (24b) and (24d) combine to give

$$\left(L^{-}(\hat{b}_{a})\smallsetminus\{b,c\}\right)\cup\{b\}=\left(L^{-}(\hat{b}_{c})\smallsetminus\{a,b\}\right)\cup\{b\}$$
(26a)

$$L^{-}(\hat{b}_{a}) \smallsetminus \{c\} = L^{-}(\hat{b}_{c}) \smallsetminus \{a\}, \qquad (26b)$$

$$L^{+}(\hat{b}_{a}) = L^{+}(\hat{b}_{c}).$$
 (26c)

With $A = \{a, \hat{b}_c\}, C = \{c, \hat{b}_a\}$ as the 2-chains $a \ll \hat{b}_c$ and $c \ll \hat{b}_a$ that are mapped to each other by a 2-chain-reflection, we obtain $a \sim_1 c$.

3.4 Classes of posets with local symmetries

All posets that are (Q, r)-retractable to some poset R form a class of symmetry extensions (or symmetry class)

$$[R \odot_r Q] := \{ P \in \mathfrak{P} \mid P \oslash_r Q = R \neq P \} , \qquad (27)$$

and we write $[R \odot_r Q]'$ to include only prime symmetries. These posets are extensions of R by (prime) (Q, r)-symmetries. In the following, we consider only prime symmetries.

Larger classes are obtained with sequences of symmetry extensions, like

$$[R \odot_{r_1} Q_1 \odot_{r_2} Q_2]' := \{ P \in \mathfrak{P} \mid P \oslash_{r_2}' Q_2 = R_1 \neq P, \\ R_1 \oslash_{r_1}' Q_1 = R \neq R_1 \},$$
(28a)
$$[R \odot_{r_1} Q_1 \odot_{r_2} Q_2 \odot_{r_3} Q_3]' := \{ P \in \mathfrak{P} \mid P \oslash_{r_3}' Q_3 = R_2 \neq P, \\ R_2 \oslash_{r_2}' Q_2 = R_1 \neq R_2, \\ R_1 \oslash_{r_1}' Q_1 = R \neq R_1 \},$$
(28b)

and so on to arbitrary order. Similarly, we write a prime to restrict to prime symmetries. *Example 3.8.* The simplest symmetry class collects all antichains that are singleton-retractable to the singleton, see Example 3.1,

$$[\bullet \odot \bullet]' = \{(\bullet \bullet), (\bullet \bullet \bullet), (\bullet \bullet \bullet \bullet), (\bullet \bullet \bullet \bullet), \dots\}.$$
(29)

Since this symmetry class already includes all 1-layer posets (except for the locally unsymmetric singleton) and the layers are preserved by retractions there are no larger classes of 1-layer posets. In particular,

$$[\bullet \odot \bullet \odot \bullet]' = \emptyset.$$
(30)

More generally, for any poset R that is not singleton-retractable, the class $[R \odot \bullet]'$ includes all extensions by the singleton-symmetry and

$$[R \odot \bullet \odot \bullet]' = \emptyset.$$
⁽³¹⁾

The singleton-symmetry is always prime since it is the smallest symmetry (and the prime is not really necessary, but kept to emphasise the focus on prime symmetries here). Similarly to the antichains symmetry class,

Lemma 3.5 (Repeated parallel composition). For every locally unsymmetric poset R,

$$\left\{ \bigsqcup_{i=1}^{n} R \mid n \ge 2 \right\} \subseteq [R \odot R]'.$$
(32)

The order in the sequence of symmetry retractions in (28) is important, because symmetry retractions do not commute and there are posets that are retractable to more than one locally unsymmetric poset. For example,

$$\bigvee \land \circ \cdot \circ \mathfrak{l} = \bigvee \land \circ \mathfrak{l} = \mathfrak{l}, \qquad (33a)$$

$$\bigvee \bigwedge \oslash \mathbf{i} \oslash \mathbf{i} \oslash \mathbf{i} \oslash \mathbf{i} = \bigvee \bigwedge \oslash \mathbf{i} = \bigwedge .$$
(33b)

This is because, the singleton-symmetry in the poset on the left partially "breaks" the 2-chainsymmetry of \bigwedge . The sequences of symmetry retractions can also be different but still yield the same poset, as in

$$\bigwedge \bigcirc \bigvee \oslash \land = \bigwedge \bigcirc \oslash \land = \bigwedge \bigcirc \land = \bigwedge \bigcirc .$$
(34b)

More so, the retraction sequences do not even have to have the same length, see the example (81) in a later section. This demonstrates that the set of all posets \mathfrak{P} does not partition into locally unsymmetric posets and classes formed from symmetry extensions of these locally unsymmetric posets. Nevertheless, individual poset classes are often easier to enumerate, and as long as we keep track of repeated countings, we can use poset classes to enumerate all posets for a fixed cardinality n.

Define the counting function as

$$c_n: \mathcal{P}(\mathfrak{P}) \to \mathbb{N}_0, \qquad \mathfrak{C} \mapsto \left| \{ P \in \mathfrak{C} \mid |P| = n \} \right|.$$
 (35)

For Example 3.8, we have

$$c_n[\bullet \odot \bullet]' = 1. \tag{36}$$

For any locally unsymmetric poset R with cardinality m := |R|, the counting function of the symmetry class under the singleton-symmetry vanishes for $n \leq m$ and, for n > m,

$$c_n[R \odot \bullet]' = \binom{(n-m)+m-1}{n-m} = \binom{n-1}{m-1},$$
(37)

because there are n - m elements to be distributed into the m equivalence classes formed by the elements of the poset R (while the order in which the elements are distributed does not matter).

4 Posets prime retractable to the 2-chain

In this section, we explore posets (prime) retractable to the 2-chain that form the simplest symmetry classes other than $[\bullet \odot \bullet]'$.

4.1 Complete bipartite graphs and crown posets

The simplest symmetry class of posets prime retractable to the 2-chain are the posets that correspond to complete bipartite graphs, for which all elements of the first and second layer form an equivalence class by the singleton-symmetry,

$$[\mathbf{i} \odot \mathbf{\cdot}]' = \left\{ \Lambda, \mathbf{V}, \Lambda, \mathbf{M}, \mathbf{V}, \mathbf{M}, \mathbf{M}, \mathbf{W}, \mathbf{V}, \ldots \right\}.$$
(38)

The enumeration of this symmetry class is a special case of (37), so for n > 2,

$$c_n[\mathbf{i} \odot \bullet]' = n - 1. \tag{39}$$

For the enumeration of all bipartite graphs, see [Han79].

By Lemma 3.5, the parallel compositions of 2-chains are 2-chain-retractable,

$$[\mathbf{I} \odot \mathbf{I}]' \supseteq \left\{ (\mathbf{I} \ \mathbf{I}), (\mathbf{I} \ \mathbf{I} \ \mathbf{I}), (\mathbf{I} \ \mathbf{I} \ \mathbf{I}), \dots \right\}.$$

$$(40)$$

Furthermore, the prime symmetry class $[\mathbf{1} \odot \mathbf{1}]'$ is complemented by another family of posets.

Definition 4.1 (Crown posets). The *n*-crown poset X_n $(n \ge 2)$ is the poset with two layers of elements a_1, a_2, \ldots, a_n and b_1, b_2, \ldots, b_n and the partial order \le such that

$$\forall i, j \in [1, n]: \qquad i \neq j \Leftrightarrow a_i < b_j. \tag{41}$$

Any crown poset corresponds to a complete bipartite graph where the edges between opposite vertices have been removed. Note that the *n*-crown poset (for n > 2) contains the *k*-crown posets for all k < n.

Lemma 4.1 (Crown retractions). The n-crown poset X_n is 2-chain-retractable to the 2-chain.

Proof. By definition of the *n*-crown, the event a_i is not linked to b_i for all $i \in [1, n]$. Considering the two 2-chains (a_i, b_j) and (a_j, b_i) for any $i \neq j$, the element pairs (a_i, a_j) and (b_i, b_j) are 2-chain symmetric, because a_i and a_j are both linked to all b_k , and similarly b_i and b_j are both linked to all a_k for $k \notin \{i, j\}$. This holds for all $i \neq j$ so that all elements a_i form a single equivalence class, and so do all b_i .

Theorem 4.2.

$$[\mathbf{1} \odot \mathbf{1}]' = \left\{ \bigsqcup_{i=1}^{n} \mathbf{1} \mid n \ge 2 \right\} \cup \{X_n \mid n \ge 3\}$$
(42a)

$$= \left\{ (\mathbf{I} \ \mathbf{I}), (\mathbf{I} \ \mathbf{I} \ \mathbf{I}), (\mathbf{I} \ \mathbf{I} \ \mathbf{I}), \dots, \mathbf{N}, \mathbf{N}, \mathbf{N}, \mathbf{N}, \mathbf{N}, \mathbf{N}, \mathbf{N}, \dots \right\}.$$
(42b)

Proof. For any poset P to be prime 2-chain-retractable to the 2-chain, it has to have two layers by Proposition 3.3 and all elements in the first and second layer each have to form a single equivalence class in P/(1, 2) with all elements in the equivalence classes being (1, 2, 0)-symmetric or (1, 2, 1)-symmetric by Lemma 3.4. This is true for the repeated parallel composition, see Lemma 3.5, and it is true for every crown by Lemma 4.1. It remains to show that there is no other poset in the symmetry class. We will show that any poset P in this symmetry class that is not a parallel composition of 2-chains is a crown poset.

Since P is not a parallel composition of 2-chains, it has to be connected and it must contain two 2-chain subsets such that the second/first layer elements are linked by an element in the first/second layer, so $\bigwedge \bigwedge$ or $\bigvee \bigvee$ is a subset of P, respectively. Under the quotient by the 2-chain, these subsets form two equivalence classes on the first or second layer, respectively, and one equivalence class on the other layer. Only if P has yet another element on the second resp. first layer that is linked to both elements on the other layer, the quotient results in one equivalence class per layer. In either case, this means the 3-crown is a subset of P.

The remaining argument is recursive assuming that we have an *n*-crown as a subset of P for some $n \geq 3$. Note that if P is in the symmetry class, so is its opposite, because the 2-chain is self-opposite. We denote the elements on the first layer a_1, a_2, \ldots, a_n and the elements on the second layer b_1, b_2, \ldots, b_n (as in Definition 4.1). If P contains other elements, there has to be another one, say a_{n+1} linked to the *n*-crown because P is connected, and we assume it is on the first layer (without loss of generality). To keep the first order 2-chain symmetry of all elements, a_{n+1} has to be linked to every element b_i with $i \in [1, n]$. As before, the resulting subset now has two equivalence classes on the first layer when taking the quotient by the 2-chain. Only if there is also another element on the second layer b_{n+1} linked to every element a_i with $i \in [1, n]$ but not linked to a_{n+1} , the quotient is isomorphic to the 2-chain. The subset formed by $a_1, a_2, \ldots, a_{n+1}$, $b_1, b_2, \ldots, b_{n+1}$ and their partial order is the (n + 1)-crown poset. We can repeat this argument until all elements of P are included, so P is a crown poset. Since all crown posets are already included, the symmetry class is complete.

For any cardinality n, there are

$$c_n[\mathbf{1} \odot \mathbf{1}]' = \begin{cases} 2 & \text{if } n \text{ is even and } n \ge 6, \\ 1 & \text{if } n = 4, \\ 0 & \text{otherwise} \end{cases}$$
(43)

posets that are prime 2-chain retractable to the 2-chain.

To get larger symmetry classes (prime) retractable to the 2-chain with further singletonsymmetries and 2-chain-reflections, recall that any repeated extension by singleton-symmetries does not yield any posets, see (31), but we may have alternating symmetry quotients as in Example 3.4, or repeat 2-chain-extensions.

Example 4.1. The following poset retracts to the 2-chain by a sequence of repeated 2-chain quotients,

$$\bigvee \otimes \mathbf{I} \otimes \mathbf{I} \otimes \mathbf{I} = \bigvee \otimes \mathbf{I} = \mathbf{I}.$$
(44)

Next, we consider posets that are (Q, r)-symmetric with Q = 1 and $r \geq 3$.

4.2 Fences and polygons

As any poset that is retractable to the 2-chain has to have two layers (by Proposition 3.3), we analyse another type of connected, 2-layer posets.

Definition 4.2 (Fences). An *n*-fence F_n is a poset of *n* elements a_i that are partially ordered in alternation,

$$a_1 < a_2 > a_3 < \dots a_n$$
 or $a_1 > a_2 < a_3 > \dots a_n$. (45)

It is a **lower** fence (denoted as F_n^{\wedge}) if $\ell_{F_n}(a_1) = 1$ or $\ell_{F_n}(a_n) = 1$, and it is an **upper** fence (denoted as F_n^{\vee}) if $\ell_{F_n}(a_1) = 2$ or $\ell_{F_n}(a_n) = 2$.

Theorem 4.3 (Symmetries of fences). Let

$$f: \mathbb{N} \to \mathbb{N},$$
 $n \mapsto \begin{cases} n & \text{if } n \text{ is even,} \\ \frac{n+1}{2} & \text{if } n \text{ is odd,} \end{cases}$ (46)

and denote the repeated composition as f^{∞} (that terminates when the result is even). Every fence of even cardinality is locally unsymmetric. Every fence of odd cardinality n is retractable to the $f^{\infty}(n)$ -fence.

Proof. The n-fence F_n with even $n \ge 4$ has exactly two elements that have only one link each, a_1 and a_n , that are on different layers so that any automorphism $\alpha \in \operatorname{Aut}(F_n)$ has to leave these fixed. So we could as well remove these elements from the fence and obtain the (n-2)-fence without changing the automorphism group, $\operatorname{Aut}(F_{n-2}) \cong \operatorname{Aut}(F_n)$. As this holds for any even n and the smallest fence, the 2-chain, is locally unsymmetric, the automorphism group of all even fences is trivial.

For odd $n = 2\hat{n} + 1$, the *n*-fences F_n^{\vee} and F_n^{\wedge} have two elements a_1 and a_n on the same layer. The fences F_3^{\vee} and F_3^{\wedge} are singleton-symmetric and retract to the 2-chain, which is F_2 , and f(n) = 2. For $\hat{n} \ge 2$, let A be the \hat{n} -fence subset of $F_n^{\vee}(F_n^{\wedge})$ including a_1 and B be the \hat{n} -fence subset of $F_n^{\vee}(F_n^{\wedge})$ including a_n . The retraction by $F_{\hat{n}}^{\vee}(F_{\hat{n}}^{\wedge})$ folds and matches the subset B onto A, while keeping the central element $a_{\hat{n}+1}$ fixed, and $F_n \oslash Q$ is the upper (lower) f(n)-fence. If f(n) is odd, repeat the retraction to get an $f^2(n)$ -fence, and so on, until the resulting fence has even cardinality $f^{\infty}(n)$ and is locally unsymmetric by the first statement. \Box

Theorem 4.3 implies that, for any $n \in \mathbb{N}$, the upper and the lower $(2^n + 1)$ -fences are retractable to the 2-chain. Except for the 3-fences (\bigwedge and \checkmark), the retractions require a sequence of more than one quotient. Though there are other 2-layer posets that have fence-symmetries and retract to the 2-chain by only *one* quotient. In particular:

Definition 4.3. The *n*-gonal poset G_n is the 2*n*-fence with an additional link that makes it cyclic,

$$a_1 < a_2 > a_3 < \dots a_{2n} > a_1 \,. \tag{47}$$

Note that in the literature you may find these polygonal posets referred to as crowns, different to the generalisation of the 3-gonal poset that I call crown posets, see Definition 4.1. To highlight the abstract representations of regular geometrical objects in terms of posets below, I keep the definition of an polygonal posets. The *n*-gonal poset represents the hull of a regular *n*-gon and a series composition with a singleton on the third layer yields the closure of the polygon that includes the central 2-face. It is obvious from the definition of the *n*-gonal poset that it is retractable to the 2-chain by a (1, n)-symmetry. Though the automorphism groups of polygons also admit subgroups that are generated by reflections only.

Theorem 4.4 (Reflection symmetries of polygonal posets). The n-gonal poset is retractable to the 2-chain by an (n-1)-fence-reflection (followed by a retraction of the singleton-symmetry if n is even).

Proof. The n-gonal poset G_n corresponds to the vertices (first layer elements) and edges (second layer elements) of a regular polygon, see Figure 4.1. Let $A \subset G_n$ such that $A \cong F_{n-1}^{\wedge}$. There are two elements c_1 and c_2 that are linked to some elements in A, which fall on the symmetry axis (dash dotted line) in the third row of Figure 4.1. Note that $B := G_n \setminus A \setminus \{c_1, c_2\}$ is also isomorphic to F_{n-1}^{\wedge} . The subsets A and B are symmetric under reflection about the symmetry axis as shown in Figure 4.1. There are reflected subsets A and B for any symmetry axis that meets an edge element. All such reflections form a subgroup of the dihedral group D_n (symmetry group of the *n*-gon). The subgroup is the entire dihedral group D_n if n is odd and it is half the dihedral group if n is even. The retraction corresponds to reducing the *n*-gon symmetries to the center of the dihedral group $Z(D_n)$, which is only one element (the identity) if n is odd, $G_n \oslash F_{n-1}^{\wedge} = 1$, and two elements if n is even, $G_n \oslash F_{n-1}^{\wedge} = \bigvee \in [1 \odot \bullet]'$.

Taking the quotient by the subgroup of the dihedral group is equivalent to removing the inner automorphisms because $D_n/Z(D_n) \cong Inn(D_n)$ [Mil42]. Under the action of the dihedral (sub)group, any vertex of the *n*-gon is mapped to any other vertex, while any edge is mapped to any other edge only in the odd case. This is graphically represented by connecting the poset elements of the *n*-gon edges to a star *n*-gon that splits into two $\frac{n}{2}$ -gons if *n* is even, see fourth row in Figure 4.1. The resulting equivalence classes in $G_n/(Q, 2)$ are the conjugacy classes of the dihedral group. Similarly, one may take the quotients with upper fences, where the axes of symmetry have to meet at least one vertex element resulting in a 2-chain or the poset $\Lambda \in [\mathbf{1} \odot \mathbf{\cdot}]'$, respectively.



Figure 4.1: The first-layer and second-layer elements of polygonal posets (first row) correspond to the vertices (red) and edges (blue) of regular polygons (second row), respectively. Regular polygons have reflection symmetries (dash dotted axis) that are represented by lower fences (green, third row). Taking the quotient by the fence is equivalent to matching vertex and edge elements for all reflection axes that meet an edge. This results in one equivalence class (fourth row) for the vertices (red cycle) and one or two equivalence classes — the conjugacy classes of the respective dihedral groups — represented by the non-degenerate or degenerate star polygons (blue), respectively. Hence, the quotients are isomorphic to the 2-chain or the Vee-poset (fifth row). The additional element in each poset (first row) above the polygonal poset is the closure including the central 2-face (grey). The element below each poset (light grey) is the dual to the top face, it is the abstract (-1)- or least face (empty shape).

Each polygonal poset is self-opposite, see Figure 4.1. Also when including the element for the 2-face as the closure of the geometric shape (grey, above) as well as its dual, least element (light grey, below), the resulting poset is self-opposite. This is equivalent to the fact that the geometric shape of a regular polygon is self-dual, and each vertex can be uniquely matched with an edge.

The polygon theorem generalises to those polyhedra that are derived from the infinite family of polygons and have dihedral symmetry, too: the self-dual pyramids, bipyramids dual to prisms, and antiprisms dual to trapezohedrons. However, posets corresponding to the vertexedge pairing are not retractable to the 2-chain (but to the N-poset) in general. Exceptions are those polygons that have even larger symmetry groups, like the tetrahedron, the 3-cube or the octahedron.

4.3 Regular polytopes and dimensional retractions

In any dimension $d \ge 3$, there is a finite number of (convex) regular polytopes. In d = 3, we have the five Platonic solids and in d = 4, there are six polytopes including the three special cases of the 24-cell, 120-cell and the 600-cell. There are three families of polytopes that exists in all dimensions $d \ge 3$, the self-dual simplices, and the hypercubes dual to the orthoplexes.

Definition 4.4 (Simplex posets). For any dimension $d \in \mathbb{N}_0$, the *d*-simplex poset S_d is the poset of d + 2 layers with 1 element on the first layer, the least face, preceding d + 1 elements on the second layer for the vertices of the *d*-simplex. The third layer contains an element for each edge of the *d*-simplex such that it succeeds exactly those elements from the first layer that correspond to the boundary (the two vertices) of the edge. This continues to each layer k with an element for each (k - 2)-face succeeding the elements corresponding to its boundary.

The Hasse diagram of the *d*-simplex poset is self-opposite and partitions into two copies of the (d-1)-simplex poset that are offset by 1 layer, giving the lower left and upper right subsets for each of the smallest simplices in the first row of Figure 4.2.

Theorem 4.5 (Reflection symmetries of simplex posets). For any $d \ge 2$, the d-simplex poset is retractable to the (d+2)-chain by the (d-2)-simplex poset.

Proof. For $d \ge 2$, the *d*-simplex has a reflection symmetry about any hyperplane that contains (the closure of) any of the (d-2)-faces. The reflection matches the two adjacent (d-2)-faces and all their boundary faces (except for those meeting the symmetry plane), which are represented by (d-2)-simplex subsets (green in the first row of Figure 4.2) shifted up by one layer. Because the choice of the hyperplane is otherwise arbitrary, there is a matching for any pair of faces of the same dimension. Thus the corresponding quotient by the (d-2)-simplex poset yields a retraction to the (d+2)-chain (for each face type, there is one equivalence class).

Corollary 4.6. For the d-simplex poset, each subset of two layers is retractable to the 2-chain.

Proof. For subsets of all elements from two distinct layers of the *d*-simplex, taking the quotient by the respective subsets of the quotient poset yields posets that retract to the 2-chain. \Box

Example 4.2. The 2-layer subsets of d-simplexes with $d \in \{2, 3, 4\}$ are shown in the lower rows of Figure 4.1. For each example, there is a pair of reflections marked in green. A special case is the 2-layer subset of all vertices and all (d-1)-faces (second row in Figure 4.2), where for each vertex there is only one non-neighbouring (d-1)-face, so that the subset is the (d+1)-crown poset.



Figure 4.2: The simplices from dimension 0 to 4 as posets (first row), where the dimension index of the faces is placed to the left of each layer. The rows below show all (but opposite) distinct subsets formed by the elements of two distinct layers. Green links mark examples of subsets that are reflected.



Figure 4.3: The square, 3-cube and 4-cube represented by their Hasse diagrams (first row), where the dimension of the faces is indicated to the left of each layer. The second row shows the posets that are obtained when taking a quotient by two different types of reflection (hyper)planes. The green subsets correspond to a reflection plane that cut through edges, while the purple subsets correspond to reflections about a diagonal plane that meets vertices.

The retraction of the *d*-simplex to a chain can be understood as a reduction from a higher dimensional order to a 1-dimensional (or total) order. We may take the standard definition of order dimension as in [DM41], though for the application in causal set theory, a dimension of a partial order can be understood equivalently in terms of its embedding Minkowski spacetime.

Definition 4.5. Let \mathbb{M}^{1+d} be (1+d)-dimensional Minkowski space (with the partial ordering determined by its causal structure). The **Minkowski dimension** of a poset P is the smallest number d such that there exists an order-preserving embedding of P in \mathbb{M}^{1+d} .

Cubes and orthoplexes are the other two families of regular polytopes that exist in any dimension $d \ge 3$. Since orthoplexes are the dual polytopes to cubes (they have opposite posets), results about their local symmetries are also dual to the local symmetries of the cubes.

The *d*-cube poset is defined completely analogous to the *d*-simplex. Three examples are shown in Figure 4.3, where two different subsets for symmetry retractions are highlighted in green and purple. Taking a quotient by either of these subsets yields the two retracted posets in the second row. For each *d*-cube, the retract shown on the left is the series composition of the singleton with the (d-1)-simplex, $\cdot \vee S_{d-1}$, while the retract shown on the right is the series composition of the singleton with the 2-antichain and a *d*-chain, see Figure 4.3. Geometrically, the retractions correspond to the symmetry reflections about hyperplanes that pass through edges (and higher faces) but not through vertices, reflecting subsets like the green selections in Figure 4.3. Any vertex of the d-cube is mapped to any other vertex by a sequence of such reflections. While it takes only one reflection to map any vertex to any of its adjacent vertices, opposite pairs of vertices require d reflections.

Any subset of two layers that does not include the vertex layer is retractable to the 2-chain by a single symmetry quotient (magenta selection in Figure 4.3). In terms of local symmetries (Definition 3.2), the *d*-cube is a principal example for elements that are (Q, 2, d)-symmetric (where Q is isomorphic to the magenta selection on the right).

4.4 Local symmetries by complete bipartite graphs

The 2-layer poset composed of the 1- and 2-face-elements of the 3-simplex poset S_3 , see Figure 4.2, is prime wedge-retractable to the 2-chain, in fact it is the only such poset.

Theorem 4.7.

$$\left[\mathbf{I}\odot\Lambda\right]' = \left\{\Lambda\Lambda\right]' = \left\{\Lambda\Lambda\right\}. \tag{48}$$

Proof. Similarly to the proof of Theorem 4.2, we only need to show that there are no other posets in the symmetry class. Any poset P in the symmetry class has to be connected, because the only non-connected posets that are wedge-retractable to the 2-chain are parallel compositions of wedges, but these are also singleton-retractable, so their wedge-symmetries are not prime. Since the connected poset P is wedge-retractable to the 2-chain, it has to contain a subset of two wedges that are linked by another element, for which there are three symmetric options

$$O_1 := \bigwedge \bigwedge, \qquad O_2 := \bigwedge \bigwedge, \qquad O_3 := \bigwedge \bigwedge. \tag{49}$$

More so, all second layer elements have to be first order wedge-symmetric to retract to the 2-chain by the wedge-symmetry, which follows analogously to the proof of Lemma 3.4. For the first option O_1 (lower 7-fence), the two elements from the left on the second layer are first order wedge-symmetric if there are two additional elements on the first layer such that the resulting subset is

$$O_1' := \bigwedge \bigwedge . \tag{50}$$

In the second option O_2 , both second layer elements are already wedge-symmetric. Here we require an additional element on the second layer to obtain the wedge-symmetry between the second and third element from the left on the first layer, leading again to O'_1 .

The wedge-symmetry between the first and second element from the left on the second layer in the third option O_3 requires two additional elements on the first layer that are both linked to the first and third element on the second layer (from the left). This extended subset is the first poset in (51) below, and any further enlargement follows a similar argument as in Theorem 4.2 leading to a sequence of crowns where each element on the first layer is replaced by a 2-antichain:

$$(51)$$

. . .

However, these subsets do not have a prime wedge-symmetry, which renders the third option irrelevant.

The subset O'_1 completes to the edges–2-faces poset of the 3-simplex

$$S_3|_{1,2} = \bigwedge (52)$$

by requiring an element on the second layer linked to the three elements on the first layer with a single link in O'_1 . This is the only choice to make all elements on the first layer wedge-symmetric and there cannot be any other elements in the subset, so that $P = S_3|_{1,2}$.

If there where another element x on the second layer, it would have to be linked to all first-layer elements of $S_3|_{1,2}$ to maintain the wedge-symmetry of these. So x and each other second-layer element y share three of their succeeded elements on the first layer of $S_3|_{1,2}$. Let a_1, a_2, a_3 be the other three elements linked to x but not to y. To have x first-order wedgesymmetric to y, we can at most take two of these elements to form a wedge subset with xat the tip, say a_1 and a_2 . The third element a_3 would also have to be linked to y to make ywedge-symmetric to x, but this is not the case in the subset $S_3|_{1,2}$. So we reached a contradiction.

For an extension by a first-layer element, similarly, it would have to be linked to all secondlayer elements of $S_3|_{1,2}$, so that there must also be another second-layer element as well. With this second layer element, we get the same argumentation as just above. So $S_3|_{1,2}$ is the only connected poset P that is prime wedge-retractable to the 2-chain.

As the vee poset is opposite to the wedge poset, this theorem implies

Corollary 4.8.

$$\begin{bmatrix} \mathbf{I} \odot \mathbf{V} \end{bmatrix}' = \left\{ \bigvee \bigvee \bigvee \right\} . \tag{53}$$

Obviously, their counting functions are

$$c_n \left[\mathbf{i} \odot \mathbf{\Lambda} \right]' = c_n \left[\mathbf{i} \odot \mathbf{\gamma} \right]' = \begin{cases} 1 & \text{if } n = 10, \\ 0 & \text{otherwise.} \end{cases}$$
(54)

Other prime Q-symmetry extensions of the 2-chain where $Q = A_n \vee \cdot$ for some antichain A_n (with $n \geq 2$) or the opposite yield similar symmetry classes (see Figure 4.2). For example, the symmetry class of prime 3-wedge-retractable posets contains two posets of the 4-simplex subsets formed from the 1/3-faces and 2/3-faces,

$$\left[\mathbf{1} \odot \mathbf{\Lambda}\right]' = \left\{ \mathbf{\Lambda}, \mathbf{\Lambda}, \mathbf{\Lambda} \right\}.$$
(55)

With similar arguments as in the proof of Theorem 4.7, it can be shown that there are no *prime* extensions of the 2-chain by Q-symmetries where $Q = A_m \vee A_n$ for some antichains A_m , A_n (with $m, n \ge 2$).

Although the 2-chain is the second simplest locally unsymmetric posets, we can extend other locally unsymmetric posets with symmetries to get locally symmetric posets still smaller than those described above. Using local symmetries for the classification of posets reduces the problem of enumerating all posets to finding and enumerating all locally unsymmetric posets (and then building symmetry classes from these).

5 Enumeration of locally unsymmetric posets by layers

Especially for the application of finite posets as subsets of causal sets, let us compare the numbers of locally unsymmetric posets to all posets with a fixed cardinality, grouped by the number of layers. First, we analyse the distribution of numbers of posets as function of their number of layers and then determine the exact enumeration of posets with a number of layers close to their cardinality. Posets with a small number of layers are dominating for larger cardinalities, so I finish with some remarks on their enumeration.

5.1 Numbers of locally unsymmetric posets vs. all posets

Let \mathfrak{U} denote the set of all locally unsymmetric posets. The smallest of these posets (up to cardinality n = 5) are

$$\mathfrak{U}_1 = \{\bullet\} , \qquad (56a)$$

$$\mathfrak{U}_2 = \{\mathfrak{l}\}, \tag{56b}$$
$$\mathfrak{U}_3 = \{(\mathfrak{l}, \bullet), \mathfrak{l}\}, \tag{56c}$$

$$\mathfrak{U}_{4} = \left\{ \bigwedge, \left(\begin{array}{c} \mathbf{I} & \mathbf{\cdot} \end{array} \right), \bigwedge, \bigvee, \begin{array}{c} \mathbf{I} \\ \mathbf{I} \\ \mathbf{I}_{4} = \left\{ \bigwedge, \left(\begin{array}{c} \mathbf{I} & \mathbf{\cdot} \end{array} \right), \bigwedge, \bigvee, \begin{array}{c} \mathbf{I} \\ \mathbf{I}$$

For the enumeration of locally unsymmetric posets among all posets \mathfrak{P} , we define the following functions and numbers. The number of posets for a given cardinality n and number of layers l is

$$p(n,l) := \left| \left\{ P \in \mathfrak{P} \mid |P| = n, \ell(P) = l \right\} \right|.$$

$$(57)$$

All posets with fixed cardinality but an arbitrary number of layers is the integer sequence in [Slo+23],

$$p_n := \left| \{ P \in \mathfrak{P} \mid |P| = n \} \right| = \sum_{l=1}^n p(n,l) .$$
 (58)

Similarly, the numbers of locally unsymmetric posets $\mathfrak U$ are defined as

$$u(n,l) := \left| \{ P \in \mathfrak{U} \mid |P| = n, \ell(P) = l \} \right|,$$
(59a)

$$u_n := \left| \{ P \in \mathfrak{U} \mid |P| = n \} \right| = \sum_{l=1}^n u(n,l) \,.$$
 (59b)

Explicit values for all numbers up to cardinality n = 7 are listed in Table 5.1 and plotted in Figure 5.1. Note that the peak of the distribution (as functions of the layer number l) for each cardinality $n \leq 7$ has its maximum at or below l = 3, while the peaks of the corresponding distributions for locally unsymmetric posets with the same cardinality are at higher l.

Remark. There is an equivalence between posets and coloured graphs. By assigning a unique colour to each layer in any poset P, from color 1 to color $\ell(P)$, it is easy to see that there is a bijection between the posets (or their Hasse diagrams) and certain colourings of simple graphs. The poset number p(n, l) is equivalent to the number of *l*-coloured, simple graphs where any node with colour k has to be graph connected to at least one node with colour k - 1 if k > 1.

p(n,l)	1	2	3	4	5	6	7	p_n
1	1							1
2	1	1						2
3	1	3	1					5
4	1	8	6	1				16
5	1	20	31	10	1			63
6	1	55	162	84	15	1		318
7	1	163	940	734	185	21	1	2045

(a) Numbers of (unlabelled) posets by layers and in total.

u(n,l)	1	2	3	4	5	6	7	u_n
1	1							1
2	0	1						1
3	0	1	1					2
4	0	1	3	1				5
5	0	1	11	6	1			19
6	0	3	47	41	10	1		102
7	0	9	266	332	106	15	1	729

(b) Numbers of locally unsymmetric (automorphically trivial) posets by layers and in total.

Table 5.1: Number of posets (a) and locally unsymmetric posets (b) by cardinality n (increasing with row number) and layer count l (increasing with column number).



(a) Frequency $p(n,l)/p_n$ vs. the number of layers l for different cardinalities n.



(b) Frequency $u(n,l)/u_n$ vs. the number of layers l for different cardinalities n.

Figure 5.1: Distributions of the number of (locally unsymmetric) posets relative to the total number of (locally unsymmetric) posets vs. the number of layers (length of longest chain) in the posets, all for different cardinalities n of the posets.

Example 5.1. There are three 2-coloured graphs with 3 nodes (n = 3, l = 2); the union of the single node (colour 1) and a graph-connected node pair (colour 1, colour 2); the 3-chain of 3 nodes (colour 1, colour 2, colour 1); as well as the chain of 3 nodes (colour 2, colour 1, colour 2). These correspond to the posets $\mathbf{1} \cdot \mathbf{A}$, and \mathbf{A} , respectively.

For some values of l, we can determine explicit expression for the numbers p(n, l) and u(n, l), starting with all antichain posets, so l = 1,

$$p(n,1) = 1. (60)$$

Here only the singleton is locally unsymmetric and all other antichains are retractable to it,

$$u(1,1) = 1$$
, $\forall n > 1: \quad u(n,1) = 0.$ (61)

The largest possible layer number is l = n that corresponds to the *n*-chain, which is locally unsymmetric,

$$p(n,n) = u(n,n) = 1.$$
 (62)

Next we enumerate the posets with l = n - 1 and l = n - 2.

5.2 Posets with one layer less than their cardinality

The simplest case (apart from the antichains and chains) are the numbers of posets with cardinality n and number of layers n - 1.

Theorem 5.1.

$$p(n, n-1) = \frac{1}{2}n(n-1),$$
 (63a)

$$u(n, n-1) = \frac{1}{2}(n-1)(n-2).$$
(63b)

Proof. All posets with cardinality n and layer number n - 1 are (n - 1)-chains with an extra element that can be separated from the chain, linked once, or linked twice. We denote an arbitrary (possibly empty) chain by a single white vertex, so we have the sequence

$$\underbrace{(\stackrel{\circ}{\mathbf{1}} \cdot)}_{n \ge 2}, \underbrace{\stackrel{\circ}{\underline{\bigwedge}}_{n \ge 3}, \underbrace{\stackrel{\circ}{\underline{\bigvee}}_{n \ge 4}, \underbrace{\stackrel{\circ}{\underline{\bigwedge}}_{n \ge 4}, \underbrace{\stackrel{\circ}{\underline{\bigwedge}}_{n \ge 5}, \underbrace{\stackrel{\circ}{\underline{\bigwedge}}_{n \ge 5}, \underbrace{\stackrel{\circ}{\underline{\bigwedge}}_{n \ge 5}, \underbrace{\stackrel{\circ}{\underline{\bigwedge}}_{n \ge 6}, \underbrace{\stackrel{\circ}{\underline{\frown}}_{n \ge 6}, \underbrace{\stackrel{\bullet}{\underline{\frown}}_{n \ge 6}, \underbrace{\stackrel{\bullet}{\underline{\frown}}_{$$

where the first brace exists for $n \ge 2$ with an (n-2)-chain replacing the white vertex, the second brace exists for $n \ge 3$ with an (n-3)-chain replacing the white vertex, and so on. It follows inductively that the k-th brace exists for $n \ge k+1$ and it encloses k posets. Hence, we have

$$p(n, n-1) = \sum_{k=2}^{n} (k-1) = \frac{1}{2}n(n-1).$$
(65)

Because there is only one element additional to the (n-1)-chain subset, there are n-1 posets that are retractable to the (n-1)-chain whenever the additional element shares an equivalence class with one of the elements in the chain:

$$\underbrace{\mathring{\Lambda}}_{n\geq 2}, \underbrace{\mathring{\Psi}}_{n\geq 3}, \underbrace{\mathring{\Psi}}_{n\geq 4}, \underbrace{\mathring{\Psi}}_{n\geq 5}, \dots$$
(66)

Since the (n-1)-chain is locally unsymmetric, we can at most find a singleton-symmetry with one more element, so there are no other local symmetries possible, and

$$u(n, n-1) = p(n, n-1) - (n-1) = \frac{1}{2}(n-1)(n-2).$$
(67)

This concludes the proof of Theorem 5.1.

The ratio of locally unsymmetric posets to all posets (for $n \ge 2$) goes as

$$\frac{u(n,n-1)}{p(n,n-1)} = 1 - \frac{2}{n} \tag{68}$$

and approaches 1 as $n \to \infty$. Asymptotically, almost all *n*-element posets with n-1 layers are locally unsymmetric.

5.3 Posets with two layers less than their cardinality

The next simplest case concerns the posets with n elements and n-2 layers.

Theorem 5.2. For n < 3, the poset numbers p(n, n-2) and u(n, n-2) vanish, and for $n \ge 3$, they are

$$p(n, n-2) = \frac{1}{6}(n-2)\left(n^3 - 2n^2 - 5n + 12\right)$$
(69a)

$$u(n, n-2) = \frac{1}{6}(n-3)\left(n^3 - 4n^2 + 2n - 2\right)$$
(69b)

Proof. All (n-2)-layer posets with cardinality n contain an (n-2)-chain and two additional elements that are – similarly to the (n-1)-layer posets – each either separated from the chain or linked to the chain with one or two links, while the two elements can be but do not have to be linked to each other as well. Say that an element is linked up/down to the chain at position $i \in [1, n-2]$ if it precedes/succeeds the event at i. For n = 3, there is only the 3-antichain, while for $n \ge 4$, we have the following cases

1. For any poset P from the sequence (64) with cardinality n - 1, we have the parallel composition with the singleton, so there are

$$P \sqcup \bullet$$
: $p(n-1, n-2) = \frac{1}{2}n^2 - \frac{3}{2}n + 1$ (70)

possible combinations.

0

2. One of the additional elements is linked up and the other one is linked down, or both are linked up/down,

$$\bigvee_{i}^{0}, i \bigvee_{i}^{j} \bigvee_{i}^{i}, i \bigvee_{i}^{i} : \qquad (n-3)^{2} + 2\sum_{i=2}^{n-2} \sum_{j=i}^{n-2} 1 = 2n^{2} - 11n + 15. \qquad (71)$$

3. One element is linked to the chain twice and the other element is linked to the chain once, independently,

$$\begin{array}{c}
\stackrel{j}{\underset{i \in \mathcal{N}}{\overset{k}{\underset{i \in \mathcal{N}}{\overset{k}{\underset{i \in \mathcal{N}}{\overset{k}{\underset{i \in \mathcal{N}}{\overset{k}{\underset{i \in \mathcal{N}}{\overset{k}{\underset{i \in \mathcal{N}}{\overset{n-2}{\underset{i = 1}{\overset{n-2}{\underset{i = 1}{\underset{i = 1}{\underset{i = 1}{\overset{n-2}{\underset{i = 1}{\underset{i = 1}$$

4. Both elements are linked to the chain twice, independently,

5. The elements are linked together and at most one of them is linked to the chain once (ignoring the posets already counted with point 2),

$$\begin{pmatrix} \mathring{} & \mathsf{I} \end{pmatrix}, \underbrace{\stackrel{\circ}{\underbrace{i}}}_{i} & \underbrace{i}}_{i} & \underbrace{i}$$

6. The elements are linked together and one of them is linked to the chain twice (ignoring the posets already counted with point 3),

$$\underbrace{2\sum_{i=1}^{n-5}\sum_{j=i+3}^{n-2}1}_{i=i+3} 1 = n^2 - 9n + 20.$$
(75)

7. The elements are linked together and each of them is linked to the chain once (ignoring the posets already counted with point 3),

$$\underbrace{\stackrel{\circ}{j}_{i}}_{i}, \underbrace{\stackrel{\circ}{j}_{i}}_{s}: \qquad 2\sum_{i=1}^{n-5}\sum_{j=i+1}^{n-4} 1 = n^2 - 9n + 20, \text{ and} \qquad (76a)$$

$$\begin{array}{cccc}
\stackrel{j}{\longleftarrow} & , & \stackrel{j}{\bigoplus} & \\
\stackrel{i}{\longleftarrow} & , & \stackrel{j}{\bigoplus} & \\
\stackrel{i}{\longleftarrow} & : & \sum_{i=1}^{n-5} \sum_{j=i+3}^{n-2} 1 + \sum_{i=1}^{n-3} \sum_{j=i+1}^{n-2} 1 = n^2 - 7n + 13. \end{array}$$
(76b)

8. The elements are linked together, one is linked to the chain twice, and the other is linked to the chain once (ignoring the posets already counted with point 4),

9. The elements are linked together and each is linked to the chain twice,

Q

$$: \qquad \sum_{i=1}^{n-5} \sum_{j=i+1}^{n-4} \sum_{k=j+1}^{n-3} \sum_{l=k+1}^{n-2} 1 = \frac{1}{24}n^4 - \frac{7}{12}n^3 + \frac{71}{24}n^2 - \frac{77}{12}n + 5.$$
(78)

The poset number p(n, n-2) as given in (69) is the sum of all the counts from these cases.

Among the p(n, n-2) posets with cardinality n, there are the following locally symmetric posets. First, any locally unsymmetric poset P with cardinality n-1 and n-2 layers form n-1 equivalence classes in P/\bullet , and an additional element can be added to any of these classes so that there are (n-1)u(n-1, n-2) combinations. Second, any singleton-symmetric poset P that is retractable to the (n-2)-chain forms n-2 equivalence classes in P/\bullet , and the remaining two elements are distributed to the classes chosen with repetition, which are $\binom{2+(n-2)-1}{2}$ possibilities. And third, all n-3 posets of the type $C_1 \sqcup (1) \sqcup C_2$ (with chains C_1, C_2 such that $|C_1| + |C_2| = n-4$) are 2-chain-symmetric and retractable to the (n-2)-chain. Together, we have

$$\frac{1}{2}(n-3)(n-2)(n-1) + \frac{1}{2}(n-2)(n-1) + n - 3 = \frac{1}{2}\left(n^3 - 5n^2 + 10n - 10\right)$$
(79)

locally symmetric posets with cardinality n and n-2 layers. The number u(n, n-2) shown in (69) is the difference between the previously computed number p(n, n-2) and this number of locally symmetric posets.

The ratio of locally unsymmetric posets among the number of posets with n-2 layers

$$\frac{u(n,n-2)}{p(n,n-2)} = 1 - \frac{3}{n} + \frac{3}{n^2} + \mathcal{O}(n^{-3})$$
(80)

shows that almost all posets with n-2 layers are locally unsymmetric for large n. Though, this ratio is smaller than the ratio (68) for all n > 3.

5.4 Posets with a small number of layers

. .

Similar calculations could be conducted to get the poset numbers p(n, n - 3), p(n, n - 4), and so on, together with the respective numbers for the locally unsymmetric posets. More relevant applications, however, are the ratios for 2-layer, 3-layer, 4-layer (and higher layer) posets. In the large *n* behaviour, these posets dominate and it was shown that a certain type of 3-layer posets suffice to get the asymptotic enumeration of all posets [KR75]. Such posets referred to as Kleitman-Rothschild orders [LC18] can have local symmetries. The example given in [CCS23a, Fig. 1] has even multiple local symmetries and is retractable to the 3-chain,

$$\langle \mathbf{A} \otimes \mathbf{A}$$

$$\langle \mathbf{\dot{\mathbf{v}}} \rangle \otimes \mathbf{\dot{\mathbf{v}}} = \langle \mathbf{\dot{\mathbf{v}}} \rangle \otimes \mathbf{\dot{\mathbf{v}}} = \mathbf{\dot{\mathbf{t}}}. \tag{81b}$$

In causal set theory, suppression mechanism are considered to remove the dominating low layer posets in favour for posets that may resemble discretized spacetime manifold more likely [LC18; CCS23a; CCS23b]. Spacetime manifolds, as the continuous counterparts to causal sets, do not have local symmetries, because they are distinguishing, meaning any two distinct elements are uniquely identified by their pasts and futures [MS08]. Similarly, elements of (locally) finite, locally unsymmetric posets are uniquely identified by their relations to all other elements in the poset.

Getting an exact quantitative statement about the number of posets with few layers is far more complicated than for the poset numbers computed above. For an estimate of 2-layer posets, we consider a generic 2-layer poset with cardinality $n = n_1 + n_2 \ge 2$ such that $n_1 \ge 1$ elements are on the first layer and $n_2 \ge 1$ elements are on the second layer. All second layer elements have to cover at least one element on the first layer so that the least number of links is n_2 . In any 2-layer poset P with n_2 links, all second layer elements cover k elements where k has to be at least 1 and at most min (n_1, n_2) . For all posets with n_2 elements on the second layer, there are

$$1 + \sum_{k=2}^{\min(n_1, n_2)} \left(\sum_{i_1=1}^{\lfloor n_2/k \rfloor} \sum_{i_2=i_1}^{\lfloor (n_2-i_1)/(k-1) \rfloor} \cdots \sum_{i_{k-1}=i_{k-2}}^{\lfloor (n_2-i_1-\dots-i_{k-2})/2 \rfloor} 1 \right)$$

=
$$1 + \underbrace{\lfloor \frac{n_2}{2} \rfloor}_{n_1, n_2 \ge 2} + \underbrace{\sum_{i_1=1}^{\lfloor n_2/3 \rfloor} \left(\lfloor \frac{n_2-i_1}{2} \rfloor - i_1 + 1 \right)}_{n_1, n_2 \ge 3} + \cdots$$
(82)

possibilities, where $\lfloor \cdot \rfloor$ is the floor function. The first term of this sum (1 possibility) corresponds to the unique poset with n_2 elements all covering the same element on the first layer (k = 1). The second term (k = 2) enumerates the posets such that the second layer elements covering two elements on the first layer — these are the partitions of n_2 into two subsets each with at least one element, so $\lfloor \frac{n_2}{2} \rfloor$ possibilities. This continues recursively up to $k = \min(n_1, n_2)$ partitions. If $n_1 \ge n_2$, the last term in (82) is 1, for the unique poset with n_2 separated 2-chains parallel composed with an $(n_1 - n_2)$ -antichain. All of these posets are at least singleton-symmetric (or 2-chain-symmetric in the last case for $n_1 = n_2$) except for the two special cases \ddagger and $\ddagger \cdot$.

For any of these posets, adding one more link connects two separated components and may break some of the local symmetries, but may also introduce a new local symmetry. For example,

$$\begin{pmatrix} \mathbf{V} & \mathbf{I} \end{pmatrix} \to \mathbf{V} \mathbf{V} \tag{83}$$

breaks an singleton-symmetry but introduces a 2-chain-symmetry. A locally unsymmetric poset is obtained by adding links such that all local symmetries are broken, link by link without introducing new local symmetries. The quickest way to obtain a locally unsymmetric poset in that way is by starting from the poset that is a parallel composition of 2-chains (and a singleton) such that $n_2 = n_1$ (or $n_2 = n_1 - 1$) and adding links until we reach a poset that is a parallel composition of a subset

$$\mathfrak{S} \subset \{\bullet\} \cup \{F_{2j} \mid j \ge 1\} \tag{84}$$

such that \mathfrak{S} contains at least one of the fences F_{2i} . For example,

$$(\mathbf{I} \ \mathbf{I} \$$

Any further link added may or may not reintroduce local symmetries. Since all bipartite posets (38) are singleton-symmetric, reaching a sufficiently large number of links such that

at least two second-layer elements cover all first layer elements means that any such posets are singleton-symmetric again.

While 2-layer posets with a sufficiently small or large number of links are almost always locally symmetric, this cannot be expected for the typical 2-layer posets considered in [KR75]. The expression (82) can be approximated by an *n*-th order polynomial, but there are far more posets with two layers (including those with an arbitrarily large dimension). Consider posets with a cardinality of 4n, 2n elements on each layer and each element linked to n elements. For a singleton-symmetry in the second layer, the chance of an element to cover the same first layer elements as another second layer element is 2^{-n} . Even if such posets may have (Q, r)-symmetries with posets Q larger than the 2-chain, like

$$\bigwedge \bigwedge \oslash' \Lambda = \bigvee, \qquad (86a)$$

it is to be expected that the numbers u(n, 2) and p(n, 2) are almost equal when n becomes large. In general, we may assume that almost all posets with a fixed cardinality and fixed but small number of layers are locally unsymmetric, though, a conclusive statement on this goes beyond this work.

6 Final remarks and applications of local symmetries

Starting from locally unsymmetric posets, we can classify all posets by extensions with local symmetries. I finish with some remarks on local extensions for the construction of symmetry classes, followed by a discussion of local symmetries in causal sets.

6.1 Local retractions and local extensions

A retraction $P \otimes_r Q$ as constructed above, removes all *r*-cyclic, local symmetries described by the finite poset Q from a poset P. For the characterisation of all posets, we may rather want to retract individual (Q, r)-symmetries from a (Q, r)-symmetry set of $A \cong Q$ in P, which we denote as $A^{(Q,r)}$. Let Σ be the subgroup of all automorphisms that leave the elements in $P \smallsetminus A^{(Q,r)}$ fixed. Taking a quotient of P by the subgroup Σ yields a retracted poset \hat{P} that is isomorphic to P up to a finite subset \hat{A} as the retracted version of $A^{(Q,r)}$. This *local retraction* is written as $\hat{P} = (P, A^{(Q,r)}) \triangleright \hat{A}$, and in the Hasse diagram representation, we use a different colour to denote the subset $A^{(Q,r)}$.

Example 6.1. For example,

$$\bigvee \bigvee \triangleright \bullet = \bigvee \bigvee (87)$$

is the local retraction of a two element subset $A^{(\cdot,2)} \cong (\cdot \cdot)$ on the first layer to the singleton, $\hat{A} \cong \cdot$.

Note that such local retractions have an inverse operation by construction. We can *locally* extend a poset \hat{P} at a subset \hat{A} by a (\hat{A}, r) -symmetry to a poset P that is isomorphic to \hat{P} up to a finite subset $A^{(Q,r)}$ with $\hat{A} \cong Q$ if $\hat{P} = (P, A^{(Q,r)}) \triangleright \hat{A}$ where $A^{(Q,r)}$ retracts to \hat{A} . For this inverse operation, we write $P = (\hat{P}, \hat{A}) \triangleleft A^{(Q,r)}$ and use again a different color or a label to indicate the corresponding subsets in the Hasse diagrams.

Example 6.2. For the previous example, we have

$$\bigvee (\cdot \cdot) = \bigvee \land .$$
(88)

With these local retractions and local extensions, it becomes much more clear when and why retractions by local symmetries are commutative.

Any (Q, r)-retraction of a (Q, r)-retractable poset P decomposes into local (Q, r)-retractions (that are prime). If $Q = \bullet$, then the local retractions commute, because any equivalence class in P/\bullet is only affected by a single local retraction.

Example 6.3. Two local singleton-extensions of the crown at different elements can be locally retracted independently,

$$\bigvee \otimes \bullet = \left({}^{2} \bigvee_{1}^{\circ} \triangleright_{1} \bullet \right) \triangleright_{2} \bullet = \left({}^{2} \bigvee_{1}^{\circ} \triangleright_{2} \bullet \right) \triangleright_{1} \bullet = \bigvee \land .$$

$$(89)$$

Commutativity is no longer guaranteed for larger symmetries.

Example 6.4. The order of retractions of the 2-chain-symmetries in the following poset yields different results

$$\bigcirc \mathbf{I} = \left(\bigcirc \mathbf{I} > \mathbf{I} \right) \triangleright \mathbf{I} = \mathbf{I} \bigcirc \mathbf{I}$$
 (90a)

$$\neq \left(\bigcup \bigcup \bigcup \bigcup \bigcup \right) \triangleright \mathbf{I} = \bigwedge , \tag{90b}$$

because the second retraction on the second equation line also involves elements retracted by the first retraction.

By using local extensions to break symmetries of a poset partially, and then retracting the partially broken symmetries followed by retractions of all other symmetries, we can construct larger locally unsymmetric posets.

Example 6.5. Recall that the *n*-gonal poset is an (1, n)-extension of the 2-chain. Any fence of even cardinality n can be constructed recursively from the (n - 1)-gonal poset, breaking the (1, n - 1)-symmetry when locally extending by a 2-antichain and then retracting by the (n - 2)-fence and the singleton-symmetry, for example,

$$\bigvee \bigvee \bigvee (\cdot \cdot) \oslash \land \bigtriangledown \odot \cdot = \land \lor \land .$$
(91)

It remains to prove if all connected, locally unsymmetric posets of a given cardinality n (except for the *n*-chain poset) are constructable from smaller locally unsymmetric posets by a series of local extensions and retractions.

The notation for local extensions can be used also in the construction of symmetry classes for larger posets from symmetry classes of smaller ones.

Example 6.6. All (prime) 2-chain extensions of the following 3-layer poset are

$$\begin{bmatrix} \mathbf{i} \\ \mathbf{i} \\ \mathbf{j} \\ \mathbf{j}$$

Using (43), the enumeration of this class becomes

$$c_n \left[\bigwedge^{\bullet} \odot \mathfrak{l} \right]' = \begin{cases} 4n - 28 & \text{if } n \text{ is odd and } n \ge 11, \\ 10 & \text{if } n = 9, \\ 4 & \text{if } n = 7, \\ 0 & \text{otherwise,} \end{cases}$$
(93)

because the 2-chain extensions always add an even number of elements (so the count for even n vanishes) and the first cases up to n = 11 can be checked directly. For the general induction step $\hat{n} = n + 2$, there are just 4 more posets with a double 2-chain extension (compared to the case n) for each of the two double extension possibilities. So there are 4n - 28 + 8 posets for $\hat{n} = n + 2$, which is $4\hat{n} - 28$.

Similarly, the 2-chain extension class for any (finite) locally unsymmetric poset is constructed by replacing a single pair of linked elements, then two independent pairs of linked elements, then three, and so on. For a 2-chain extension class of a locally symmetric poset, the equivalence classes under the respective local symmetries have to be respected so that elements from one equivalence class are only counted once per local 2-chain extension.

6.2 Sprinkled causal sets and stability of locally unsymmetric posets

To conclude this study of local symmetries in posets with an application, consider causal sets that are generated from a given spacetime manifold via a Poisson process called sprinkling. For any (pre)-compact region U of a spacetime manifold M with volume measure ν and a sprinkling density $\rho > 0$ (a strictly positive real parameter), the probability to obtain a random sprinkle S_U (as a causal set) with cardinality n is given by

$$\Pr\left(|\mathsf{S}_U| = n\right) = e^{-\rho\nu(U)} \frac{\rho^n}{n!} \nu(U)^n , \qquad (94)$$

see [Few+21] for a mathematical review. Especially for Minkowski spacetime, sprinkles are typically not only locally unsymmetric but exhibit further properties stemming from their lack of local symmetries.

Definition 6.1. For any $k \in \mathbb{N}_0$, a poset P is k-stable locally unsymmetric if, for every subset $S \subseteq P$ that has cardinality $0 \le |S| \le k$, the poset $P \smallsetminus S$ is locally unsymmetric. An infinite poset P is total locally unsymmetric if $P \backsim S$ is k-stable locally unsymmetric for every $k \in \mathbb{N}_0$.

Example 6.7. Any chain posets with a minimal or maximal element is total locally unsymmetric.

Theorem 6.1. A sprinkle in d-dimensional Minkowski spacetime is total locally unsymmetric with probability 1.

Proof. In (1 + 0)-dimensional Minkowski spacetime, every sprinkle is a chain poset and thus total locally unsymmetric. If a sprinkle S in Minkowski spacetime \mathbb{M}^{1+d} with d > 0 has a local symmetry, there would have to be a pair of elements $x, y \in S$ that has the same set of succeeding elements (the same future). Since by construction of the sprinkling process, $x \neq y$, we choose a coordinate system such that x is at the origin and y has the same coordinates except for its first spatial coordinate ε , meaning it is displaced by a spatial distance of $\varepsilon > 0$. For any t > 0, let z_t be the point again with identical coordinates except for a time coordinate t and first spatial coordinate $\varepsilon - t$. The spacetime interval I_t between x and z_t is the intersection of the subsets

s(n,l)	1	2	3	4	5	6	7	$ s_n $
1	1							1
2	0	1						1
3	0	0	1					1
4	0	0	1	1				2
5	0	0	0	3	1			4
6	0	0	2	8	6	1		17
7	0	0	4	37	36	10	1	88

Table 6.1: Numbers of 1-stable locally unsymmetric posets by layers and in total.

 $J^+(x)$ and $J^-(z_t)$ under the partial ordering by the causal structure of \mathbb{M}^{1+d} . Since x and y have the same future in the random sprinkle S, the region I_t cannot contain an element of S for any t > 0. The probability for this is

$$\Pr\left(|\mathsf{S} \cap I_t| = 0\right) = e^{-\rho\nu(I_t)}.$$
(95)

The volume $\nu(I_t)$ of the compact region is finite and increases with t linearly for d = 1 and more than linearly for d > 1 by construction. Because, we can choose t arbitrarily large, in the limit $t \to \infty$, the probability (95) vanishes (note that ε can be arbitrarily small but never zero). Hence any sprinkle **S** is locally unsymmetric with probability 1. Moreover, the probability for the region I_t to contain any finite number of elements vanishes in the limit $t \to \infty$ due to an analogous argument. This means that the random sprinkle is total locally unsymmetric with probability 1.

Corollary 6.2. A sprinkle in d-dimensional Minkowski spacetime does not contain singletonsymmetric elements (it does not contain non-Hegelian pairs [WC17]) with probability 1.

Spacetime manifolds that are considered relevant in physics are globally hyperbolic or at least distinguishing, meaning that any two distinct points also have distinct pasts (and futures) [MS08]. Since any pair of elements of a locally unsymmetric (and locally finite) poset is distinct by their relations to all other elements, posets that are locally unsymmetric can be taken as a generalisation of distinguishing spacetimes. The property of total unsymmetry may suffice to rule out infinite posets with a finite number of layers, although this is not shown here. Further investigations are necessary to determine if general (total) locally unsymmetric causets are a candidate for "manifold-like" causets and thus relevant for applications in quantum gravity, which provides an alternative to previous works [MRS09].

It is also an open question, whether local symmetries in finite posets may have direct applications in other parts of causal set theory, in particular, the (covariant) growth dynamics, see [Zal23].

For finite posets, the numbers of 1-stable locally unsymmetric posets s(n, l) by layer are given in Table 6.1. In Example 6.7, we saw that all chains are total locally unsymmetric, so that also s(n, n) = 1. Furthermore, it is easy to see that (similar to the calculation in subsection 2), $s(n, n-1) = \frac{1}{2}(n-2)(n-3)$. Note that up to cardinality n = 7, there is only one 2-layer poset (the 2-chain) that is 1-stable locally unsymmetric.

Local symmetries can be generalised to other types of orders. For example, one may replace the condition of reflexivity of the ordering relation on the set P by irreflexivity making P a strict poset, or by more general conditions like almost reflexivity (defined in [GPZ23]). Because the reflexivity condition is actually not required in the definition, it can be dropped altogether making the order relation \leq merely transitive and antisymmetric.

Acknowledgement

Thanks goes to Steven Carlip and Sumati Surya for discussions about Kleitman-Rothschild orders during the Quantum Gravity conference 2023 at the RRI, India, and to Yasaman Yazdi for suggestions on the presentation.

References

- [Bom+87] L. Bombelli, J. Lee, D. Meyer, and R. Sorkin. "Space-Time as a Causal Set". In: Phys. Rev. Lett. 59 (1987), pp. 521–524.
- [BG91] G. Brightwell and R. Gregory. "Structure of random discrete spacetime". In: Phys. Rev. Lett. 66.3 (1991), pp. 260–263.
- [BHS08] G. Brightwell, J. Henson, and S. Surya. "A 2D model of causal set quantum gravity: the emergence of the continuum". In: *Classical Quantum Gravity* 25 (2008), p. 105025.
- [BM02] G. Brinkmann and B. D. McKay. "Posets on up to 16 points". In: Order 19 (June 2002), pp. 147–179.
- [But72] K. K.-H. Butler. "The Number of Partially Ordered Sets". In: J. Combin. Theory Ser. B 13.3 (1972), pp. 276–289. ISSN: 0095-8956.
- [CCS23a] P. Carlip, S. Carlip, and S. Surya. "Path integral suppression of badly behaved causal sets". In: *Classical Quantum Gravity* 40.9 (Mar. 2023), p. 095004. arXiv: 2209.00327 [gr-qc].
- [CCS23b] P. Carlip, S. Carlip, and S. Surya. The Einstein-Hilbert Action for Entropically Dominant Causal Sets. Nov. 2023. arXiv: 2311.18238 [gr-qc].
- [Dha80] D. Dhar. "Asymptotic enumeration of partially ordered sets". In: Pacific J. Math. 90.2 (1980), pp. 299–305.
- [DR79] D. Duffus and I. Rival. "Retracts of partially ordered sets". In: J. Aust. Math. Soc. 27.4 (1979), pp. 495–506.
- [DR81] D. Duffus and I. Rival. "A structure theory for ordered sets". In: Discrete Math.
 35.1 (1981). Special Volume on Ordered Sets, pp. 53–118. ISSN: 0012-365X.
- [DM41] B. Dushnik and E. W. Miller. "Partially Ordered Sets". In: Amer. J. Math. 63.3 (1941), pp. 600–610.
- [Few+21] C. J. Fewster, E. Hawkins, C. Minz, and K. Rejzner. "Local Structure of Sprinkled Causal Sets". In: Phys. Rev. D 103.8 (2021), p. 086020.
- [GPZ23] L. Guo, S. Paycha, and B. Zhang. "Mathematical Reflections on Locality". In: Jahresber. Dtsch. Math.-Ver. 125.4 (2023), pp. 197–238.
- [Han79] P. Hanlon. "The enumeration of bipartite graphs". In: *Discrete Math.* 28.1 (1979), pp. 49–57.
- [Kla69] D. A. Klarner. "The Number of Graded Partially Ordered Sets". In: J. Combin. Theory 6.1 (1969), pp. 12–19. ISSN: 0021-9800.
- [KR75] D. J. Kleitman and B. L. Rothschild. "Asymptotic enumeration of partial orders on a finite set". In: Trans. Amer. Math. Soc. 205 (1975), pp. 205–220.

- [LC18] S. P. Loomis and S. Carlip. "Suppression of non-manifold-like sets in the causal set path integral". In: *Classical Quantum Gravity* 35.2 (2018), p. 024002. arXiv: 1709.00064 [gr-qc].
- [MRS09] S. Major, D. Rideout, and S. Surya. "Stable Homology as an Indicator of Manifoldlikeness in Causal Set Theory". In: *Classical Quantum Gravity* 26 (2009), p. 175008. arXiv: 0902.0434 [gr-qc].
- [Mil42] G. A. Miller. "Automorphisms of the dihedral groups". In: Proc. Natl. Acad. Sci. USA 28.9 (1942), pp. 368–371.
- [MS08] E. Minguzzi and M. Sánchez. "The causal hierarchy of spacetimes". In: *Recent de*velopments in pseudo-Riemannian geometry. 2008. arXiv: gr-qc/0609119.
- [Min20] C. Minz. causets. LaTeX-package to draw causal set (Hasse) diagrams. Nov. 2020.
- [PST01a] H. J. Prömel, A. Steger, and A. Taraz. "Asymptotic enumeration, global structure, and constrained evolution". In: *Discrete Math.* 229.1–3 (2001), pp. 213–233.
- [PST01b] H. J. Prömel, A. Steger, and A. Taraz. "Phase transitions in the evolution of partial orders". In: J. Combin. Theory 94.2 (2001), pp. 230–275.
- [Slo+23] N. J. A. Sloane et al. A000112: Number of partially ordered sets with n unlabeled elements. The On-Line Encyclopedia of Integer Sequences, OEIS Foundation Inc. 2023.
- [WC17] C. Wüthrich and C. Callender. "What Becomes of a Causal Set?" In: British J. Philos. Sci. 68.3 (2017), pp. 907–925.
- [Zal23] S. Zalel. "Covariant Growth Dynamics". In: Handbook of Quantum Gravity. Ed. by C. Bambi, L. Modesto, and I. Shapiro. Singapore: Springer Nature Singapore, 2023, pp. 1–36.