

NON-ABELIAN EXTENSIONS OF LIE TRIPLE SYSTEMS AND WELLS EXACT SEQUENCES

QINXIU SUN AND SHUANGJIAN GUO*

ABSTRACT. In this paper, we investigate non-abelian extensions and inducibility of pairs of automorphisms of Lie triple systems. First, we introduce non-abelian cohomology groups and classify the non-abelian extensions in terms of non-abelian cohomology groups. Next, we characterize the non-abelian extensions using Maurer-Cartan elements. Furthermore, we explore the inducibility of pairs of automorphisms and derive the analog Wells exact sequences under the circumstance of Lie triple systems. Finally, we state the previous results under the context of abelian extensions of Lie triple systems.

CONTENTS

1. Introduction	1
2. Preliminary on Lie triple systems	2
3. Non-abelian extensions of Lie triple systems	3
4. Non-abelian extensions in terms of Maurer Cartan elements	10
5. Inducibility of pairs of Lie triple system automorphisms	13
6. Wells exact sequences for Lie triple systems	18
7. Particular case: abelian extensions of Lie triple systems	20
References	21

1. INTRODUCTION

Lie triple systems originated from Cartan's studies of Riemannian geometry, in which he employed his classification of the real simple Lie algebras to classify symmetric spaces. The role played by Lie triple systems in the theory of symmetric spaces is like that of Lie algebras in the theory of Lie groups. Lie triple systems were investigated from the algebraic point of view by Jacobson in [15] and studied later by Lister in [19]. Subsequently, Lie triple systems have gained broad attention and a number of works have been gained, see [2, 22, 28] and their references.

Extensions of some mathematical object are useful to understand the underlying structure. The non-abelian extension is a relatively general one among various extensions (e.g. central extensions, abelian extensions, non-abelian extensions etc.). Non-abelian extensions were first developed by Eilenberg and Maclane [5], which led to the low dimensional non-abelian group cohomology. Then numerous works have been devoted to non-abelian extensions of various kinds of algebras, such as Lie algebras, associative algebras, Lie groups, pre-Lie algebras, Leibniz algebras, 3-Lie algebras, Rota-Baxter Lie algebras, conformal algebras and Rota-Baxter Leibniz algebras, see [6, 8, 9, 13, 14, 18, 20, 21, 23, 24, 26] and their references. But little is known

2010 *Mathematics Subject Classification.* 17A30, 17B62, 17B38 .

Key words and phrases. Lie triple system, non-abelian extension, Maurer-Cartan element, inducibility, Wells exact sequence.

about the non-abelian extension of Lie triple systems. This was our first motivation for writing the present paper.

Our second motivation is to explore the inducibility of pairs of Lie triple system automorphisms. The inducibility of a pair of automorphisms has close relationship with extensions of algebras. Such study originated from extensions of abstract groups [27]. This theme was subsequently extended to various algebraic structures, such as Lie algebras, Rota-Baxter Lie algebras, Rota-Baxter groups, associative conformal algebras and Lie-Yamaguti algebras, see [1, 4, 7, 10, 13, 16, 18, 25, 27] and references therein. As byproducts, the Wells short exact sequences were derived for various kinds of algebras [4, 7, 9, 10, 13, 16, 18], which connected the relative automorphism groups and the non-abelian second cohomology groups. Motivated by these results and combined with our study of non-abelian extension of Lie triple systems, naturally we study inducibility of pairs of Lie triple system automorphisms and derive the Wells short exact sequences in the context of non-abelian extensions of Lie triple systems.

The paper is organized as follows. In Section 2, we recall basic knowledge of Lie triple systems. In Section 3, we investigate non-abelian extensions and classified the non-abelian extensions using non-abelian 3-cocycles. In Section 4, we characterize non-abelian extensions in terms of Maurer-Cartan elements. In Section 5, we address the inducibility of pairs of Lie triple system automorphisms and give some equivalent conditions. In Section 6, we consider the analogue of the Wells short exact sequences under the case of Lie triple systems. In Section 7, we explain the previous results in the context of abelian extensions of Lie triple systems.

Throughout the paper, let k be a field. Unless otherwise specified, all vector spaces and algebras are over k .

2. PRELIMINARY ON LIE TRIPLE SYSTEMS

A Lie triple system is a vector space \mathfrak{g} together with a trilinear map $[\cdot, \cdot, \cdot] : \mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g} \longrightarrow \mathfrak{g}$ satisfying

$$(1) \quad [x_1, x_1, x_2] = 0,$$

$$(2) \quad [x_1, x_2, x_3] + [x_2, x_3, x_1] + [x_3, x_1, x_2] = 0,$$

$$(3) \quad [x_1, x_2, [y_1, y_2, y_3]] = [[x_1, x_2, y_1], y_2, y_3] + [y_1, [x_1, x_2, y_2], y_3] + [y_1, y_2, [x_1, x_2, y_3]],$$

for all $x_i, y_i \in \mathfrak{g}$ ($i = 1, 2, 3$).

Eqs. (1) and (2) yield that

$$(4) \quad [x_1, x_2, x_3] + [x_2, x_1, x_3] = 0.$$

A subspace M of \mathfrak{g} is an ideal of \mathfrak{g} if $[M, \mathfrak{g}, \mathfrak{g}] \subseteq M$ and $[\mathfrak{g}, \mathfrak{g}, M] \subseteq M$. An ideal M of \mathfrak{g} is said to be an abelian ideal of \mathfrak{g} if $[M, M, \mathfrak{g}] = [\mathfrak{g}, M, M] = 0$.

A representation of a Lie triple system \mathfrak{g} consists of a vector space V together with a bilinear map $\theta : \mathfrak{g} \wedge \mathfrak{g} \longrightarrow \mathfrak{gl}(V)$ satisfying

$$(5) \quad \theta(x_3, x_4)\theta(x_1, x_2) - \theta(x_2, x_4)\theta(x_1, x_3) - \theta(x_1, [x_2, x_3, x_4]) + D_\theta(x_2, x_3)\theta(x_1, x_4) = 0,$$

$$(6) \quad \theta(x_3, x_4)D_\theta(x_1, x_2) - D_\theta(x_1, x_2)\theta(x_3, x_4) + \theta([x_1, x_2, x_3], x_4) + \theta(x_3, [x_1, x_2, x_4]) = 0,$$

for any $x_i \in \mathfrak{g}$ ($i = 1, 2, 3, 4$), where

$$(7) \quad D_\theta(x_1, x_2) = \theta(x_2, x_1) - \theta(x_1, x_2).$$

One can refer to [11, 12, 19, 29–31] for more information on Lie triple systems and representations.

Let \mathfrak{g} be a Lie triple system and (V, θ) be a representation of it. Denote the $(2n + 1)$ -cochains group of \mathfrak{g} with coefficients in V by $C^{2n+1}(\mathfrak{g}, V)$, where $f \in C^{2n+1}(\mathfrak{g}, V)$ is a multilinear map $f : \mathfrak{g} \times \cdots \times \mathfrak{g} \longrightarrow V$ satisfying

$$(8) \quad f(x_1, \dots, x_{2n-2}, x, x, y) = 0$$

and

$$(9) \quad f(x_1, \dots, x_{2n-2}, x, y, z) + f(x_1, \dots, x_{2n-2}, y, z, x) + f(x_1, \dots, x_{2n-2}, z, x, y) = 0$$

for all $x_i, x, y, z \in \mathfrak{g}$ ($i = 1, \dots, 2n - 2$) and $C^1(\mathfrak{g}, V) = \text{Hom}(\mathfrak{g}, V)$.

The Yamaguti coboundary operator $\delta : C^{2n-1}(\mathfrak{g}, V) \longrightarrow C^{2n+1}(\mathfrak{g}, V)$ is given by

$$\begin{aligned} & (\delta f)(x_1, \dots, x_{2n+1}) \\ &= \theta(x_{2n}, x_{2n+1})f(x_1, \dots, x_{2n-1}) - \theta(x_{2n-1}, x_{2n+1})f(x_1, \dots, x_{2n-2}, x_{2n}) \\ &+ \sum_{i=1}^n (-1)^{i+n} D_\theta(x_{2i-1}, x_{2i})f(x_1, \dots, x_{2i-2}, x_{2i+1}, \dots, x_{2n+1}) \\ &+ \sum_{i=1}^n \sum_{j=2i+1}^{2n+1} (-1)^{i+n+1} f(x_1, \dots, x_{2i-2}, x_{2i+1}, \dots, x_{j-1}, [x_{2i-1}, x_{2i}, x_j], \dots, x_{2n+1}) \end{aligned}$$

for any $x_1, x_2, \dots, x_{2n+1} \in \mathfrak{g}$. Denote the set of all $(2n - 1)$ -cocycles and $(2n - 1)$ -coboundaries respectively by $Z^{2n-1}(\mathfrak{g}, V)$ and $B^{2n-1}(\mathfrak{g}, V)$. Thus, we can define the $(2n - 1)$ -cohomology group by

$$H^{2n-1}(\mathfrak{g}, V) = \begin{cases} Z^1(A, V), & n = 1, \\ Z^{2n-1}(A, V)/B^{2n-1}(A, V), & n > 1. \end{cases}$$

See [17, 31] for more details on Yamaguti cohomology.

3. NON-ABELIAN EXTENSIONS OF LIE TRIPLE SYSTEMS

In this section, we are devoted to considering non-abelian extensions and non-abelian 3-cocycles of Lie triple systems.

Definition 3.1. Let \mathfrak{g} and \mathfrak{h} be two Lie triple systems. A non-abelian extension of \mathfrak{g} by \mathfrak{h} is a Lie triple system $\hat{\mathfrak{g}}$, which fits into a short exact sequence of Lie triple systems

$$\mathcal{E} : 0 \longrightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \longrightarrow 0.$$

When \mathfrak{h} is an abelian ideal of $\hat{\mathfrak{g}}$, the extension \mathcal{E} is called an abelian extension of \mathfrak{g} by \mathfrak{h} . Denote an extension as above simply by $\hat{\mathfrak{g}}$ or \mathcal{E} . A section of p is a linear map $s : \mathfrak{g} \longrightarrow \hat{\mathfrak{g}}$ such that $ps = I_{\mathfrak{g}}$.

Definition 3.2. Let $\hat{\mathfrak{g}}_1$ and $\hat{\mathfrak{g}}_2$ be two non-abelian extensions of \mathfrak{g} by \mathfrak{h} . They are said to be equivalent if there is a homomorphism of Lie triple systems $f : \hat{\mathfrak{g}}_1 \longrightarrow \hat{\mathfrak{g}}_2$ such that the following commutative diagram holds:

$$(10) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathfrak{h} & \xrightarrow{i_1} & \hat{\mathfrak{g}}_1 & \xrightarrow{p_1} & \mathfrak{g} \longrightarrow 0 \\ & & \parallel & & \downarrow f & & \parallel \\ 0 & \longrightarrow & \mathfrak{h} & \xrightarrow{i_2} & \hat{\mathfrak{g}}_2 & \xrightarrow{p_2} & \mathfrak{g} \longrightarrow 0. \end{array}$$

Denote by $\mathcal{E}_{\text{non-abelian}}(\mathfrak{g}, \mathfrak{h})$ the set of all non-abelian extensions of \mathfrak{g} by \mathfrak{h} .

Next, we define a non-abelian cohomology group and show that the non-abelian extensions are classified by the non-abelian cohomology groups.

Definition 3.3. Let \mathfrak{g} and \mathfrak{h} be two Lie triple systems. A non-abelian 3-cocycle on \mathfrak{g} with values in \mathfrak{h} is a triple (ω, θ, ρ) of maps such that $\omega : \mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g} \longrightarrow \mathfrak{h}$ is trilinear, $\theta : \mathfrak{g} \wedge \mathfrak{g} \longrightarrow \mathfrak{gl}(\mathfrak{h})$ is bilinear and $\rho : \mathfrak{g} \longrightarrow \text{Hom}(\mathfrak{h} \wedge \mathfrak{h}, \mathfrak{h})$ is linear, and the following identities are satisfied for all $x_i, y_i, x, y, z, w \in \mathfrak{g}$ ($i = 1, 2, 3$), $a, b, c \in \mathfrak{h}$,

$$(11) \quad \omega(x, y, z) + \omega(y, z, x) + \omega(z, x, y) = 0,$$

$$(12) \quad \begin{aligned} & D_{\theta}(x_1, x_2)\omega(y_1, y_2, y_3) + \omega(x_1, x_2, [y_1, y_2, y_3]_{\mathfrak{g}}) = \omega([x_1, x_2, y_1]_{\mathfrak{g}}, y_2, y_3) \\ & + \theta(y_2, y_3)\omega(x_1, x_2, y_1) + \omega(y_1, [x_1, x_2, y_2]_{\mathfrak{g}}, y_3) - \theta(y_1, y_3)\omega(x_1, x_2, y_2) \\ & + \omega(y_1, y_2, [x_1, x_2, y_3]_{\mathfrak{g}}) + D_{\theta}(y_1, y_2)\omega(x_1, x_2, y_3), \end{aligned}$$

$$(13) \quad \begin{aligned} & D_{\theta}(x, y)\theta(z, w)a - \theta(z, w)D_{\theta}(x, y)a \\ & = \theta([x, y, z]_{\mathfrak{g}}, w)a + \theta(z, [x, y, w]_{\mathfrak{g}})a - D_{\rho}(w)(\omega(x, y, z), a) - \rho(z)(a, \omega(x, y, w)), \end{aligned}$$

$$(14) \quad \theta(x, [y, z, w]_{\mathfrak{g}})a - \rho(x)(a, \omega(y, z, w)) = \theta(z, w)\theta(x, y)a - \theta(y, w)\theta(x, z)a + D_{\theta}(y, z)\theta(x, w)a,$$

$$(15) \quad \begin{aligned} & D_{\theta}(x, y)\rho(z)(a, b) = \rho(z)(D_{\theta}(x, y)a, b) + \rho([x, y, z]_{\mathfrak{g}})(a, b) \\ & + [\omega(x, y, z), a, b]_{\mathfrak{h}} + \rho(z)(a, D_{\theta}(x, y)b), \end{aligned}$$

$$(16) \quad D_{\theta}(y, z)\rho(x)(a, b) = \rho(x)(a, D_{\theta}(y, z)b) - \rho(z)(\theta(x, y)a, b) + \rho(y)(\theta(x, z)a, b),$$

$$(17) \quad \theta(y, z)(\rho(x)(a, b)) = \rho(x)(a, \theta(y, z)b) - D_\rho(z)(\theta(x, y)a, b) - \rho(y)(b, \theta(x, z)a),$$

$$(18) \quad D_\theta(x, y)([a, b, c]_{\mathfrak{h}}) = [D_\theta(x, y)a, b, c]_{\mathfrak{h}} + [a, D_\theta(x, y)b, c]_{\mathfrak{h}} + [a, b, D_\theta(x, y)c]_{\mathfrak{h}},$$

$$(19) \quad \rho(x)(a, \rho(y)(b, c)) = \rho(y)(\rho(x)(a, b), c) + \rho(y)(b, \rho(x)(a, c)) - [\theta(x, y)a, b, c]_{\mathfrak{h}},$$

$$(20) \quad [a, b, \theta(x, y)c]_{\mathfrak{h}} = \theta(x, y)[a, b, c]_{\mathfrak{h}} - D_\rho(y)(D_\rho(x)(a, b), c) - \rho(x)(c, D_\rho(y)(a, b)),$$

$$(21) \quad [a, b, \rho(x)(c, d)]_{\mathfrak{h}} = \rho(x)([a, b, c]_{\mathfrak{h}}, d) - [c, D_\rho(x)(a, b), d]_{\mathfrak{h}} + \rho(x)(c, [a, b, d]_{\mathfrak{h}}),$$

$$(22) \quad \rho(x)(a, [b, c, d]_{\mathfrak{h}}) = [\rho(x)(a, b), c, d]_{\mathfrak{h}} + [b, \rho(x)(a, c), d]_{\mathfrak{h}} + [b, c, \rho(x)(a, d)]_{\mathfrak{h}},$$

where

$$(23) \quad D_\theta(x, y)a - \theta(y, x)a + \theta(x, y)a = 0, \quad \rho(x)(a, b) + D_\rho(x)(a, b) - \rho(x)(b, a) = 0.$$

Definition 3.4. Let $(\omega_1, \theta_1, \rho_1)$ and $(\omega_2, \theta_2, \rho_2)$ be two non-abelian 3-cocycles on \mathfrak{g} with values in \mathfrak{h} . They are said to be equivalent if there exists a linear map $\varphi : \mathfrak{g} \rightarrow \mathfrak{h}$ such that for all $x, y, z \in \mathfrak{g}$ and $a, b \in \mathfrak{h}$, the following equalities hold:

$$(24) \quad \begin{aligned} \omega_1(x, y, z) - \omega_2(x, y, z) &= \theta_2(x, z)\varphi(y) - D_{\theta_2}(x, y)\varphi(z) + \rho_2(x)(\varphi(y), \varphi(z)) - \theta_2(y, z)\varphi(x) \\ &+ D_{\rho_2}(z)(\varphi(x), \varphi(y)) - \rho_2(y)(\varphi(x), \varphi(z)) - [\varphi(x), \varphi(y), \varphi(z)]_{\mathfrak{h}} + \varphi([x, y, z]_{\mathfrak{g}}), \end{aligned}$$

$$(25) \quad \theta_1(x, y)a - \theta_2(x, y)a = \rho_2(x)(a, \varphi(y)) - D_{\rho_2}(y)(a, \varphi(x)) + [a, \varphi(x), \varphi(y)]_{\mathfrak{h}},$$

$$(26) \quad \rho_1(x)(a, b) - \rho_2(x)(a, b) = [a, \varphi(x), b]_{\mathfrak{h}}.$$

Using Eqs. (23), (25) and (26), we get

$$(27) \quad D_{\theta_1}(x, y)a - D_{\theta_2}(x, y)a = \rho_2(y)(\varphi(x), a) - \rho_2(x)(\varphi(y), a) + [\varphi(x), \varphi(y), a]_{\mathfrak{h}},$$

and

$$(28) \quad D_{\rho_1}(x)(a, b) - D_{\rho_2}(x)(a, b) = [b, a, \varphi(x)]_{\mathfrak{h}}.$$

Denote the set of all non-abelian 3-cocycles on \mathfrak{g} with values in \mathfrak{h} by $Z_{nab}^3(\mathfrak{g}, \mathfrak{h})$. The equivalent class of a non-abelian 3-cocycle (ω, θ, ρ) is denoted by $[(\omega, \theta, \rho)]$. The quotient of $Z_{nab}^3(\mathfrak{g}, \mathfrak{h})$ by the above equivalence relation is denoted by $H_{nab}^3(\mathfrak{g}, \mathfrak{h})$, which is called non-abelian cohomology group of \mathfrak{g} with coefficients in \mathfrak{h} .

Using the above notations, we define a trilinear map $[\cdot, \cdot, \cdot]_{(\omega, \theta, \rho)}$ on $\mathfrak{g} \oplus \mathfrak{h}$ by

$$(29) \quad \begin{aligned} [x + a, y + b, z + c]_{(\omega, \theta, \rho)} &= [x, y, z]_{\mathfrak{g}} + \omega(x, y, z) + D_\theta(x, y)c + \theta(y, z)a - \theta(x, z)b \\ &+ D_\rho(z)(a, b) + \rho(x)(b, c) - \rho(y)(a, c) + [a, b, c]_{\mathfrak{h}} \end{aligned}$$

for all $x, y, z \in \mathfrak{g}$ and $a, b, c \in \mathfrak{h}$.

Proposition 3.5. *With the above notions, $(\mathfrak{g} \oplus \mathfrak{h}, [, ,]_{(\omega, \theta, \rho)})$ is a Lie triple system if and only if the triple (ω, θ, ρ) is a non-abelian 3-cocycle. Denote this Lie triple system $(\mathfrak{g} \oplus \mathfrak{h}, [, ,]_{(\omega, \theta, \rho)})$ simply by $\mathfrak{g} \oplus_{(\omega, \theta, \rho)} \mathfrak{h}$.*

Proof. $(\mathfrak{g} \oplus \mathfrak{h}, [, ,]_{(\omega, \theta, \rho)})$ is a Lie triple system if and only if Eqs. (1)-(3) hold for $[, ,]_{(\omega, \theta, \rho)}$. In fact, it is easy to check that Eqs. (1)-(2) hold for $[, ,]_{(\omega, \theta, \rho)}$ if and only if (11), (23) hold. For the Eq. (3), we discuss it for the following cases: for all $x_1, x_2, y_1, y_2, y_3 \in \mathfrak{g} \oplus \mathfrak{h}$,

(I) when all of the five elements x_1, x_2, y_1, y_2, y_3 belongs to \mathfrak{g} , (3) holds if and only if (12) holds.

(II) when x_1, x_2, y_1, y_2, y_3 equal to:

(i) x, y, z, w, a or x, y, z, a, w or x, a, y, z, w respectively, (3) holds if and only if (13) holds.

(ii) x, y, a, z, w or a, x, y, z, w respectively, (3) holds for if and only if (14) holds.

(iii) x, y, z, a, b or x, y, a, b, z or x, y, a, z, b respectively, (3) holds if and only if (15) holds.

(iv) x, a, y, z, b or a, x, y, z, b respectively, (3) holds if and only if (16) holds.

(v) x, a, y, b, z or a, x, y, b, z or x, a, b, y, z or a, x, b, y, z respectively, (3) holds if and only if (17) holds.

(vi) a, b, x, y, z , (3) holds if and only if (15), (17) hold.

(vii) x, y, a, b, c , (3) holds if and only if (18) holds.

(viii) x, a, y, b, c or a, x, y, b, c or x, a, b, c, y or a, x, b, c, y or a, x, b, y, c or x, a, b, y, c , (3) holds if and only if (19) holds.

(ix) a, b, x, c, y or a, b, c, x, y or a, b, x, y, c , (3) holds if and only if (20) holds.

(x) a, b, c, d, x or a, b, c, x, d or a, b, x, c, d , (3) holds if and only if (21) holds.

(xi) a, x, b, c, d or x, a, b, c, d , (3) holds if and only if (22) holds,

where $x, y, z, w \in \mathfrak{g}$ and $a, b, c, d \in \mathfrak{h}$ in all the cases of (i)-(xi).

This completes the proof. □

Let $\mathcal{E} : 0 \longrightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \longrightarrow 0$ be a non-abelian extension of \mathfrak{g} by \mathfrak{h} with a section s of p . Define $\omega_s : \mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{h}$, $\theta_s : \mathfrak{g} \wedge \mathfrak{g} \rightarrow \mathfrak{gl}(\mathfrak{h})$, $\rho_s : \mathfrak{g} \rightarrow \text{Hom}(\mathfrak{h} \wedge \mathfrak{h}, \mathfrak{h})$ respectively by

$$(30) \quad \omega_s(x, y, z) = [s(x), s(y), s(z)]_{\hat{\mathfrak{g}}} - s[x, y, z]_{\mathfrak{g}},$$

$$(31) \quad \theta_s(x, y)a = [a, s(x), s(y)]_{\hat{\mathfrak{g}}}, \quad \rho_s(x)(a, b) = [s(x), a, b]_{\hat{\mathfrak{g}}}.$$

By Eq. (31), we have

$$(32) \quad D_{\theta_s}(x, y)a = [s(x), s(y), a]_{\hat{\mathfrak{g}}}, \quad D_{\rho_s}(x)(a, b) = [a, b, s(x)]_{\hat{\mathfrak{g}}}$$

for any $x, y, z \in \mathfrak{g}$, $a, b \in \mathfrak{h}$.

By direct computations, we have

Proposition 3.6. *With the above notions, $(\omega_s, \theta_s, \rho_s)$ is a non-abelian 3-cocycle on \mathfrak{g} with values in \mathfrak{h} . We call it the non-abelian 3-cocycle corresponding to the extension \mathcal{E} induced by s . Naturally, $(\mathfrak{g} \oplus \mathfrak{h}, [\cdot, \cdot, \cdot]_{(\omega_s, \theta_s, \rho_s)})$ is a Lie triple system. Denote this Lie triple system simply by $\mathfrak{g} \oplus_{(\omega_s, \theta_s, \rho_s)} \mathfrak{h}$.*

In the following, we denote $(\omega_s, \theta_s, \rho_s)$ by (ω, θ, ρ) without ambiguity.

Remark 3.7. When \mathcal{E} is an abelian extension of \mathfrak{g} by \mathfrak{h} , the maps ρ_s given by (31) becomes to zero. Then (\mathfrak{h}, θ_s) is a representation of \mathfrak{g} [31]. Moreover, ω_s is a 3-cocycle of \mathfrak{g} with coefficients in the representation (\mathfrak{h}, θ_s) . We call it the 3-cocycle corresponding to the abelian extension \mathcal{E} induced by s . We always denote ω_s simply by ω if there is no confusion.

Lemma 3.8. *Let $\mathcal{E} : 0 \longrightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \longrightarrow 0$ be a non-abelian extension of \mathfrak{g} by \mathfrak{h} with different sections s_1, s_2 of p . Assume that $(\omega_i, \theta_i, \rho_i)$ is the non-abelian 3-cocycle corresponding to the extension \mathcal{E} induced by s_i ($i=1,2$). Then $(\omega_1, \theta_1, \rho_1)$ and $(\omega_2, \theta_2, \rho_2)$ are equivalent, that is, the equivalent classes of non-abelian 3-cocycles corresponding to a non-abelian extension induced by a section are independent on the choice of sections.*

Proof. Let $\hat{\mathfrak{g}}$ be a non-abelian extension of \mathfrak{g} by \mathfrak{h} . Assume that s_1 and s_2 are two different sections of p , $(\omega_1, \theta_1, \rho_1)$ and $(\omega_2, \theta_2, \rho_2)$ are the corresponding non-abelian 3-cocycles. Define a linear map $\varphi : \mathfrak{g} \longrightarrow \mathfrak{h}$ by $\varphi(x) = s_2(x) - s_1(x)$. Since $p\varphi(x) = ps_2(x) - ps_1(x) = 0$, φ is well defined. Thanks to Eqs. (30)-(32), we get

$$\begin{aligned}
w_1(x, y, z) &= [s_1(x), s_1(y), s_1(z)]_{\hat{\mathfrak{g}}} - s_1[x, y, z]_{\mathfrak{g}} \\
&= [s_2(x) - \varphi(x), s_2(y) - \varphi(y), s_2(z) - \varphi(z)]_{\hat{\mathfrak{g}}} - (s_2[x, y, z]_{\mathfrak{g}} - \varphi([x, y, z]_{\mathfrak{g}})) \\
&= [s_2(x), s_2(y), s_2(z)]_{\hat{\mathfrak{g}}} - [s_2(x), \varphi(y), s_2(z)]_{\hat{\mathfrak{g}}} - [s_2(x), s_2(y), \varphi(z)]_{\hat{\mathfrak{g}}} + [s_2(x), \varphi(y), \varphi(z)]_{\hat{\mathfrak{g}}} \\
&\quad - [\varphi(x), s_2(y), s_2(z)]_{\hat{\mathfrak{g}}} + [\varphi(x), \varphi(y), s_2(z)]_{\hat{\mathfrak{g}}} + [\varphi(x), s_2(y), \varphi(z)]_{\hat{\mathfrak{g}}} - [\varphi(x), \varphi(y), \varphi(z)]_{\hat{\mathfrak{g}}} \\
&\quad - s_2[x, y, z]_{\mathfrak{g}} + \varphi([x, y, z]_{\mathfrak{g}}) \\
&= w_2(x, y, z) + \theta_2(x, z)\varphi(y) - D_{\theta_2}(x, y)\varphi(z) + \rho_2(x)(\varphi(y), \varphi(z)) - \theta_2(y, z)\varphi(x) + D_{\rho_2}(z)(\varphi(x), \varphi(y)) \\
&\quad - \rho_2(y)(\varphi(x), \varphi(z)) - [\varphi(x), \varphi(y), \varphi(z)]_{\hat{\mathfrak{g}}} + \varphi[x, y, z]_{\mathfrak{g}},
\end{aligned}$$

which yields that Eq. (24) holds. Similarly, Eqs. (25) and (26) hold. This finishes the proof. \square

According to Proposition 3.5 and Proposition 3.6, given a non-abelian extension $\mathcal{E} : 0 \longrightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \longrightarrow 0$ of \mathfrak{g} by \mathfrak{h} with a section s of p , we have a non-abelian 3-cocycle $(\omega_s, \theta_s, \rho_s)$ and a Lie triple system $\mathfrak{g} \oplus_{(\omega_s, \theta_s, \rho_s)} \mathfrak{h}$, which yields that

$$\mathcal{E}_{(\omega_s, \theta_s, \rho_s)} : 0 \longrightarrow \mathfrak{h} \xrightarrow{i} \mathfrak{g} \oplus_{(\omega_s, \theta_s, \rho_s)} \mathfrak{h} \xrightarrow{\pi} \mathfrak{g} \longrightarrow 0$$

is a non-abelian extension of \mathfrak{g} by \mathfrak{h} . Since any element $\hat{w} \in \hat{\mathfrak{g}}$ can be written as $\hat{w} = a + s(x)$ with $a \in \mathfrak{h}, x \in \mathfrak{g}$, define a linear map

$$f : \hat{\mathfrak{g}} \longrightarrow \mathfrak{g} \oplus_{(\omega_s, \theta_s, \rho_s)} \mathfrak{h}, \quad f(\hat{w}) = f(a + s(x)) = a + x.$$

It is easy to check that f is a homomorphism of Lie triple systems such that the following commutative diagram holds:

$$\begin{array}{ccccccc} \mathcal{E} : 0 & \longrightarrow & \mathfrak{h} & \xrightarrow{i} & \hat{\mathfrak{g}} & \xrightarrow{p} & \mathfrak{g} \longrightarrow 0 \\ & & \parallel & & \downarrow f & & \parallel \\ \mathcal{E}_{(\omega_s)} : 0 & \longrightarrow & \mathfrak{h} & \xrightarrow{i} & \mathfrak{g} \oplus_{(\omega_s, \theta_s, \rho_s)} \mathfrak{h} & \xrightarrow{\pi} & \mathfrak{g} \longrightarrow 0, \end{array}$$

which indicates that the non-abelian extensions \mathcal{E} and $\mathcal{E}_{(\omega_s, \theta_s, \rho_s)}$ of \mathfrak{g} by \mathfrak{h} are equivalent. On the other hand, if (ω, θ, ρ) is a non-abelian 3-cocycle on \mathfrak{g} with values in \mathfrak{h} , there is a Lie triple system $\mathfrak{g} \oplus_{(\omega, \theta, \rho)} \mathfrak{h}$, which yields the following non-abelian extension of \mathfrak{g} by \mathfrak{h} :

$$\mathcal{E}_{(\omega, \theta, \rho)} : 0 \longrightarrow \mathfrak{h} \xrightarrow{i} \mathfrak{g} \oplus_{(\omega, \theta, \rho)} \mathfrak{h} \xrightarrow{\pi} \mathfrak{g} \longrightarrow 0,$$

where i is the inclusion and π is the projection.

In the sequel, we characterize the relationship between non-abelian cohomology groups and non-abelian extensions in the following.

Proposition 3.9. *Let \mathfrak{g} and \mathfrak{h} be two Lie triple systems. Then the equivalent classes of non-abelian extensions of \mathfrak{g} by \mathfrak{h} are classified by the non-abelian cohomology group, that is, $\mathcal{E}_{nab}(\mathfrak{g}, \mathfrak{h}) \simeq H_{nab}^3(\mathfrak{g}, \mathfrak{h})$.*

Proof. Define a linear map

$$\Theta : \mathcal{E}_{nab}(\mathfrak{g}, \mathfrak{h}) \rightarrow H_{nab}^3(\mathfrak{g}, \mathfrak{h}),$$

where Θ assigns an equivalent class of non-abelian extensions to the class of corresponding non-abelian 3-cocycles. First, we check that Θ is well-defined. Assume that \mathcal{E}_1 and \mathcal{E}_2 are two equivalent non-abelian extensions of \mathfrak{g} by \mathfrak{h} via the map f , that is, the commutative diagram (10) holds. Let $s_1 : \mathfrak{g} \rightarrow \hat{\mathfrak{g}}_1$ be a section of p_1 . Then $p_2 f s_1 = p_1 s_1 = I_{\mathfrak{g}}$, which follows that $s_2 = f s_1$ is a section of p_2 . Let $(\omega_1, \theta_1, \rho_1)$ and $(\omega_2, \theta_2, \rho_2)$ be two non-abelian 3-cocycles induced by the sections s_1, s_2 respectively. Then we have,

$$\begin{aligned} \theta_1(x, y)a &= f(\theta_1(x, y)a) = f([a, s_1(x), s_1(y)]_{\hat{\mathfrak{g}}_1}) \\ &= [f(a), f s_1(x), f s_1(y)]_{\hat{\mathfrak{g}}_2} \\ &= [a, s_2(x), s_2(y)]_{\hat{\mathfrak{g}}_2} \\ &= \theta_2(x, y)a. \end{aligned}$$

By the same token, we have

$$\omega_1(x, y, z) = \omega_2(x, y, z), \quad \rho_1(x)(a, b) = \rho_2(x)(a, b).$$

Thus, $(\omega_1, \theta_1, \rho_1) = (\omega_2, \theta_2, \rho_2)$, which means that Θ is well-defined.

Next, we verify that Θ is injective. Indeed, suppose that $\Theta([\mathcal{E}_1]) = [(\omega_1, \theta_1, \rho_1)]$ and $\Theta([\mathcal{E}_2]) = [(\omega_2, \theta_2, \rho_2)]$. If the equivalent classes $[(\omega_1, \theta_1, \rho_1)] = [(\omega_2, \theta_2, \rho_2)]$, we obtain that the non-abelian 3-cocycles $(\omega_1, \theta_1, \rho_1)$ and $(\omega_2, \theta_2, \rho_2)$ are equivalent via the linear map $\varphi : \mathfrak{g} \rightarrow \mathfrak{h}$, satisfying Eqs. (24)-(26). Define a linear map $f : \mathfrak{g} \oplus_{(\omega_1, \theta_1, \rho_1)} \mathfrak{h} \rightarrow \mathfrak{g} \oplus_{(\omega_2, \theta_2, \rho_2)} \mathfrak{h}$ by

$$f(x + a) = x - \varphi(x) + a, \quad \forall x \in \mathfrak{g}, a \in \mathfrak{h}.$$

According to Eq. (29), for all $x, y, z \in \mathfrak{g}, a, b, c \in \mathfrak{h}$, we get

$$\begin{aligned} & f([x + a, y + b, z + c]_{(\omega_1, \theta_1, \rho_1)}) \\ &= f([x, y, z]_{\mathfrak{g}} + \omega_1(x, y, z) + D_{\theta_1}(x, y)c + \theta_1(y, z)a - \theta_1(x, z)b \\ & \quad + D_{\rho_1}(z)(a, b) + \rho_1(x)(b, c) - \rho_1(y)(a, c) + [a, b, c]_{\mathfrak{h}}) \\ &= [x, y, z]_{\mathfrak{g}} - \varphi([x, y, z]_{\mathfrak{g}}) + \omega_1(x, y, z) + D_{\theta_1}(x, y)c + \theta_1(y, z)a - \theta_1(x, z)b \\ & \quad + D_{\rho_1}(z)(a, b) + \rho_1(x)(b, c) - \rho_1(y)(a, c) + [a, b, c]_{\mathfrak{h}}, \end{aligned}$$

and

$$\begin{aligned} & [f(x + a), f(y + b), f(z + c)]_{(\omega_2, \theta_2, \rho_2)} \\ &= [x - \varphi(x) + a, y - \varphi(y) + b, z - \varphi(z) + c]_{(\omega_2, \theta_2, \rho_2)} \\ &= [x, y, z]_{\mathfrak{g}} + \omega_2(x, y, z) + D_{\theta_2}(x, y)(c - \varphi(z)) + \theta_2(y, z)(a - \varphi(x)) - \theta_2(x, z)(b - \varphi(y)) \\ & \quad + T_2(z)(a - \varphi(x), b - \varphi(y)) + \rho_2(x)(b - \varphi(y), c - \varphi(z)) - \rho_2(y)(a - \varphi(x), c - \varphi(z)) \\ & \quad + [a - \varphi(x), b - \varphi(y), c - \varphi(z)]_{\mathfrak{h}}) \\ &= [x, y, z]_{\mathfrak{g}} + \omega_2(x, y, z) + D_{\theta_2}(x, y)(c - \varphi(z)) + \theta_2(y, z)(a - \varphi(x)) - \theta_2(x, z)(b - \varphi(y)) \\ & \quad + D_{\rho_2}(z)(\varphi(x), \varphi(y)) - D_{\rho_2}(z)(a, \varphi(y)) - D_{\rho_2}(z)(\varphi(x), b) + D_{\rho_2}(z)(a, b) \\ & \quad + \rho_2(x)(\varphi(y), \varphi(z)) - \rho_2(x)(\varphi(y), c) - \rho_2(x)(b, \varphi(z)) + \rho_2(x)(b, c) - \rho_2(y)(\varphi(x), \varphi(z)) \\ & \quad + \rho_2(y)(\varphi(x), c) + \rho_2(y)(a, \varphi(z)) - \rho_2(y)(a, c) - [\varphi(x), \varphi(y), \varphi(z)]_{\mathfrak{h}} + [\varphi(x), \varphi(y), c]_{\mathfrak{h}} \\ & \quad + [\varphi(x), b, \varphi(z)]_{\mathfrak{h}} - [\varphi(x), b, c]_{\mathfrak{h}} + [a, \varphi(y), \varphi(z)]_{\mathfrak{h}} - [a, b, \varphi(z)]_{\mathfrak{h}} - [a, \varphi(y), c]_{\mathfrak{h}} + [a, b, c]_{\mathfrak{h}}. \end{aligned}$$

In view of Eqs. (24)-(28), we have

$$f([x + a, y + b, z + c]_{(\omega_1, \theta_1, \rho_1)}) = [f(x + a), f(y + b), f(z + c)]_{(\omega_2, \theta_2, \rho_2)}.$$

Hence, f is a homomorphism of Lie triple systems. Clearly, the following commutative diagram holds:

$$(33) \quad \begin{array}{ccccccc} \mathcal{E}_{(\omega_1, \theta_1, \rho_1)} : 0 & \longrightarrow & \mathfrak{h} & \xrightarrow{i} & \mathfrak{g} \oplus_{(\omega_1, \theta_1, \rho_1)} \mathfrak{h} & \xrightarrow{\pi} & \mathfrak{g} \longrightarrow 0 \\ & & \parallel & & \downarrow f & & \parallel \\ \mathcal{E}_{(\omega_2, \theta_2, \rho_2)} : 0 & \longrightarrow & \mathfrak{h} & \xrightarrow{i} & \mathfrak{g} \oplus_{(\omega_2, \theta_2, \rho_2)} \mathfrak{h} & \xrightarrow{\pi} & \mathfrak{g} \longrightarrow 0. \end{array}$$

Thus $\mathcal{E}_{(\omega_1, \theta_1, \rho_1)}$ and $\mathcal{E}_{(\omega_2, \theta_2, \rho_2)}$ are equivalent non-abelian extensions of \mathfrak{g} by \mathfrak{h} , which means that $[\mathcal{E}_{(\omega_1, \theta_1, \rho_1)}] = [\mathcal{E}_{(\omega_2, \theta_2, \rho_2)}]$. Thus, Θ is injective.

Finally, we claim that Θ is surjective. For any equivalent class of non-abelian 3-cocycles $[(\omega)]$, by Proposition 3.5, there is a non-abelian extension of \mathfrak{g} by \mathfrak{h} :

$$\mathcal{E}_{(\omega)} : 0 \longrightarrow \mathfrak{h} \xrightarrow{i} \mathfrak{g} \oplus_{(\omega, \theta, \rho)} \mathfrak{h} \xrightarrow{\pi} \mathfrak{g} \longrightarrow 0.$$

Therefore, $\Theta([\mathcal{E}_{(\omega, \theta, \rho)}]) = [(\omega, \theta, \rho)]$, which follows that Θ is surjective. In all, Θ is bijective. This finishes the proof. \square

4. NON-ABELIAN EXTENSIONS IN TERMS OF MAURER CARTAN ELEMENTS

In this section, we classify the non-abelian extensions using Maurer Cartan elements. We start with recalling the Maurer-Cartan elements.

Let $(L = \oplus_i L_i, [,], d)$ be a differential graded Lie algebra. The set $\text{MC}(L)$ of Maurer-Cartan elements of $(L, [,], d)$ is defined by

$$\text{MC}(L) = \{\eta \in L_1 \mid d\eta + \frac{1}{2}[\eta, \eta] = 0\}.$$

Moreover, $\eta_0, \eta_1 \in \text{MC}(L)$ are called gauge equivalent if and only if there exists an element $\varphi \in L_0$ such that

$$\eta_1 = e^{ad_\varphi} \eta_0 - \frac{e^{ad_\varphi} - 1}{ad_\varphi} d\varphi.$$

Let \mathfrak{g} be a vector space. Consider the graded vector space $C^*(\mathfrak{g}, \mathfrak{g}) = \oplus_{n \geq 0} C^n(\mathfrak{g}, \mathfrak{g})$, where $C^n(\mathfrak{g}, \mathfrak{g})$ is the set of linear maps $f \in \text{Hom}(L \otimes \cdots \otimes L \otimes \mathfrak{g}, \mathfrak{g})$, where $L = \mathfrak{g} \otimes \mathfrak{g}$, satisfying Eqs. (8) and (9) for all $X_i \in L$ ($i = 1, 2, \dots, n-1$), $x, y, z \in \mathfrak{g}$. The degree of elements in $C^n(\mathfrak{g}, \mathfrak{g})$ is n . Define

$$(34) \quad [f, g]_{L\mathfrak{g}} = (-1)^{mn} i_f(g) - i_g(f), \quad \forall f \in C^n(\mathfrak{g}, \mathfrak{g}), g \in C^m(\mathfrak{g}, \mathfrak{g}),$$

where $i_f(g) \in C^{n+m}(\mathfrak{g}, \mathfrak{g})$ is given by

$$(35) \quad i_f(g) = \sum_{k=1}^{n+1} (-1)^{m(k-1)} \sum_{\sigma \in \text{sh}(k-1, m)} (-1)^\sigma f \circ_k^\sigma g,$$

where σ is a permutation in $(k-1, m)$ -shuffle and $f \circ_k^\sigma g$ is defined for $k = n+1$ by

$$(36) \quad f \circ_{n+1}^\sigma g(X_1, X_2, \dots, X_{n+m}, z) = f(X_{\sigma(1)}, \dots, X_{\sigma(n)}, g(X_{\sigma(n+1)}, \dots, X_{\sigma(n+m)}, z))$$

and for $1 \leq k \leq n$ by

$$(37) \quad \begin{aligned} & f \circ_k^\sigma g(X_1, \dots, X_{n+m}, z) \\ &= f(X_{\sigma(1)}, \dots, X_{\sigma(k-1)}, g(X_{\sigma(k)}, \dots, X_{\sigma(n+m-1)}, x_{k+m}), y_{k+m}, X_{k+m+1}, \dots, X_{m+n}, z) \\ &+ f(X_{\sigma(1)}, \dots, X_{\sigma(k-1)}, x_{k+m}, g(X_{\sigma(k)}, \dots, X_{\sigma(k+m-1)}, y_{k+m}), X_{k+m+1}, \dots, X_{m+n}, z) \end{aligned}$$

for all $X_i \in L$ ($i = 1, 2, \dots, n+m$) and $z \in \mathfrak{g}$.

Proposition 4.1. [3] *With the above notations, $(C^*(\mathfrak{g}, \mathfrak{g}), [\cdot, \cdot]_{Lts})$ is a graded Lie algebra. $\pi \in C^1(\mathfrak{g}, \mathfrak{g}) = \text{Hom}(\wedge^2 \mathfrak{g} \otimes \mathfrak{g}, \mathfrak{g})$ defines a Lie triple system structure on \mathfrak{g} if and only if $[\cdot, \cdot]_{Lts} = 0$, i.e. π is a Maurer-Cartan element of the graded Lie algebra $(C^*(\mathfrak{g}, \mathfrak{g}), [\cdot, \cdot]_{Lts})$. Furthermore, $(C^*(\mathfrak{g}, \mathfrak{g}), [\cdot, \cdot]_{Lts}, d_\pi)$ is a differential graded Lie algebra, where d_π is given by*

$$(38) \quad d_\pi(f) = (-1)^{n-1} [\pi, f]_{Lts}, \quad \forall f \in C^{n-1}(\mathfrak{g}, \mathfrak{g}).$$

Let $(\mathfrak{g}, \pi_{\mathfrak{g}})$ and $(\mathfrak{h}, \pi_{\mathfrak{h}})$ be two Lie triple systems. Then $(\mathfrak{g} \oplus \mathfrak{h}, \pi_{\mathfrak{g} \oplus \mathfrak{h}})$ is a Lie triple system, where $\pi_{\mathfrak{g} \oplus \mathfrak{h}}$ is defined by

$$\pi_{\mathfrak{g} \oplus \mathfrak{h}}(x+a, y+b, z+c) = \pi_{\mathfrak{g}}(x, y, z) + \pi_{\mathfrak{h}}(a, b, c)$$

for all $x, y, z \in \mathfrak{g}, a, b, c \in \mathfrak{h}$.

In view of Proposition 4.1, $(C^*(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{g} \oplus \mathfrak{h}), [\cdot, \cdot]_{Lts}, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}})$ is a differential graded Lie algebra. Define $C_{>}^n(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}) \subset C^n(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h})$ by

$$C^n(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}) = C_{>}^n(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}) \oplus C^n(\mathfrak{h}, \mathfrak{h}).$$

Denote by $C_{>}(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}) = \bigoplus_n C_{>}^n(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h})$.

Similar to the case of 3-Lie algebras [26], we have

Proposition 4.2. *With the above notations, $(C_{>}(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}), [\cdot, \cdot]_{Lts}, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}})$ is a differential graded Lie subalgebra of $(C^*(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{g} \oplus \mathfrak{h}), [\cdot, \cdot]_{Lts}, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}})$. Moreover, its degree 0 part $C_{>}^0(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}) = \text{Hom}(\mathfrak{h}, \mathfrak{h})$ is abelian.*

Proposition 4.3. *The following conditions are equivalent:*

- (i) $(\mathfrak{g} \oplus \mathfrak{h}, [\cdot, \cdot]_{(\omega, \theta, \rho)})$ is a Lie triple system, which is a non-abelian extension of \mathfrak{g} by \mathfrak{h} ;
- (ii) $\bar{\omega}$ is a Maurer-Cartan element of the differential graded Lie algebra $(C_{>}(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}), [\cdot, \cdot]_{Lts}, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}})$, where

$$\begin{aligned} \bar{\omega}(x+a, y+b, z+c) &= \omega(x, y, z) + D_\theta(x, y)c + \theta(y, z)a - \theta(x, z)b \\ &+ D_\rho(z)(a, b) + \rho(x)(b, c) - \rho(y)(a, c), \end{aligned}$$

for all $x, y, z \in \mathfrak{g}, a, b, c \in \mathfrak{h}$.

Proof. According to the definition of Maurer-Cartan element, $\bar{\omega}$ is a Maurer-Cartan element of the differential graded Lie algebra $(C_{>}(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}), [\cdot, \cdot]_{Lts}, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}})$ if and only if

$$d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \bar{\omega} + \frac{1}{2}[\bar{\omega}, \bar{\omega}]_{Lts} = 0.$$

Thus,

$$d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \bar{\omega} + \frac{1}{2}[\bar{\omega}, \bar{\omega}]_{Lts} = [\pi_{\mathfrak{g} \oplus \mathfrak{h}}, \bar{\omega}]_{Lts} + \frac{1}{2}[\bar{\omega}, \bar{\omega}]_{Lts} = \frac{1}{2}[\pi_{\mathfrak{g} \oplus \mathfrak{h}} + \bar{\omega}, \pi_{\mathfrak{g} \oplus \mathfrak{h}} + \bar{\omega}] = 0,$$

which implies that $\pi_{\mathfrak{g} \oplus \mathfrak{h}} + \bar{\omega}$ is a Maurer-Cartan element of the differential graded Lie algebra $(C^*(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{g} \oplus \mathfrak{h}), [\cdot, \cdot]_{Lts}, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}})$. Combining Proposition 4.1, $(\mathfrak{g} \oplus \mathfrak{h}, [\cdot, \cdot]_{(\omega, \theta, \rho)})$ is a Lie triple system and $\bar{\omega}$ is a Maurer-Cartan element of $(C_{>}(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}), [\cdot, \cdot]_{Lts}, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}})$. \square

Proposition 4.4. *Let \mathfrak{g} and \mathfrak{h} be two Lie triple systems. Then the equivalent classes of non-abelian extensions \mathfrak{g} by \mathfrak{h} are one-to-one correspond to the gauge equivalence classes of Maurer-Cartan elements in the differential graded Lie algebra $(C_{>}(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}), [\cdot, \cdot]_{Lts}, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}})$.*

Proof. Let $\bar{\omega}, \bar{\omega}'$ be two Maurer-Cartan elements of the differential graded Lie algebra $(C_{>}(\mathfrak{g} \oplus \mathfrak{h}, \mathfrak{h}), [\cdot, \cdot]_{Lts}, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}})$, and $\bar{\omega}, \bar{\omega}'$ are gauge equivalent, there is a linear map $\varphi \in \text{Hom}(\mathfrak{g}, \mathfrak{h})$ such that

$$\begin{aligned} \bar{\omega}' &= e^{ad_{\varphi}} \bar{\omega} - \frac{e^{ad_{\varphi}} - 1}{ad_{\varphi}} d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi \\ &= (id + ad_{\varphi} + \frac{1}{2!} ad_{\varphi}^2 + \cdots + \frac{1}{n!} ad_{\varphi}^n + \cdots) \bar{\omega} \\ &\quad - (id + ad_{\varphi} + \frac{1}{2!} ad_{\varphi} + \cdots + \frac{1}{n!} ad_{\varphi}^{n-1} + \cdots) d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi. \end{aligned}$$

For all $w_i = x_i + a_i \in \mathfrak{g} \oplus \mathfrak{h}$ ($i = 1, 2, 3$), thanks to Eqs. (29), (34)-(38), we obtain

$$\begin{aligned} ad_{\varphi} \bar{\omega} = [\varphi, \bar{\omega}]_{Lts}(w_1, w_2, w_3) &= -\bar{\omega}(\varphi(w_1), w_2, w_3) - \bar{\omega}(w_1, \varphi(w_2), w_3) - \bar{\omega}(w_1, w_2, \varphi(w_3)) \\ &= -\theta(x_2, x_3)\varphi(x_1) - D_{\rho}(x_3)(\varphi(x_1), a_2) + \rho(x_2)(\varphi(x_1), a_3) \\ &\quad + \theta(x_1, x_3)\varphi(x_2) - D_{\rho}(x_3)(a_1, \varphi(x_2)) - \rho(x_1)(\varphi(x_2), a_3) \\ &\quad - D_{\theta}(x_1, x_2)\varphi(x_3) - \rho(x_1)(a_2, \varphi(x_3)) + \rho(x_2)(a_1, \varphi(x_3)), \end{aligned}$$

$$\begin{aligned} ad_{\varphi}^2 \bar{\omega} &= [\varphi, [\varphi, \bar{\omega}]_{Lts}]_{Lts}(w_1, w_2, w_3) \\ &= -[\varphi, \bar{\omega}]_{Lts}(\varphi(w_1), w_2, w_3) - [\varphi, \bar{\omega}]_{Lts}(w_1, \varphi(w_2), w_3) - [\varphi, \bar{\omega}]_{Lts}(w_1, w_2, \varphi(w_3)) \\ &= D_{\rho}(x_3)(\varphi(x_1), \varphi(x_2)) - \rho(x_2)(\varphi(x_1), \varphi(x_3)) + D_{\rho}(x_3)(\varphi(x_1), \varphi(x_2)) \\ &\quad + \rho(x_1)(\varphi(x_2), \varphi(x_3)) - \rho(x_2)(\varphi(x_1), \varphi(x_3)) + \rho(x_1)(\varphi(x_2), \varphi(x_3)) \\ &= 2D_{\rho}(x_3)(\varphi(x_1), \varphi(x_2)) - 2\rho(x_2)(\varphi(x_1), \varphi(x_3)) + 2\rho(x_1)(\varphi(x_2), \varphi(x_3)), \end{aligned}$$

$$\begin{aligned}
d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi(w_1, w_2, w_3) &= [\pi_{\mathfrak{g} \oplus \mathfrak{h}}, \varphi]_{Lts}(w_1, w_2, w_3) \\
&= \pi_{\mathfrak{g} \oplus \mathfrak{h}}(\varphi(w_1), w_2, w_3) + \pi_{\mathfrak{g} \oplus \mathfrak{h}}(w_1, \varphi(w_2), w_3) \\
&\quad + \pi_{\mathfrak{g} \oplus \mathfrak{h}}(w_1, w_2, \varphi(w_3)) - \varphi(\pi_{\mathfrak{g} \oplus \mathfrak{h}}(w_1, w_2, w_3)) \\
&= [\varphi(x_1), a_2, a_3]_{\mathfrak{h}} + [a_1, \varphi(x_2), a_3]_{\mathfrak{h}} + [a_1, a_2, \varphi(x_3)]_{\mathfrak{h}} - \varphi([x_1, x_2, x_3]_{\mathfrak{g}}),
\end{aligned}$$

$$\begin{aligned}
ad_{\varphi}(d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi) &= [\varphi, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi]_{Lts}(w_1, w_2, w_3) \\
&= -d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi(\varphi(w_1), w_2, w_3) - d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi(w_1, \varphi(w_2), w_3) - d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi(w_1, w_2, \varphi(w_3)) \\
&= [\varphi(x_1), \varphi(x_2), a_3]_{\mathfrak{h}} + [\varphi(x_1), a_2 \varphi(x_3)]_{\mathfrak{h}} + [\varphi(x_1), \varphi(x_2), a_3]_{\mathfrak{h}} \\
&\quad + [a_1, \varphi(x_2), \varphi(x_3)]_{\mathfrak{h}} - [\varphi(x_1), a_2, \varphi(x_3)]_{\mathfrak{h}} + [a_1, \varphi(x_2), \varphi(x_3)]_{\mathfrak{h}} \\
&= 2[a_1, \varphi(x_2), \varphi(x_3)]_{\mathfrak{h}} + 2[\varphi(x_1), a_2, \varphi(x_3)]_{\mathfrak{h}} + 2[\varphi(x_1), \varphi(x_2), a_3]_{\mathfrak{h}},
\end{aligned}$$

$$\begin{aligned}
ad_{\varphi}^2(d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi) &= [\varphi, [\varphi, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi]_{Lts}]_{Lts}(w_1, w_2, w_3) \\
&= -[\varphi, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi]_{Lts}(\varphi(x_1), w_2, w_3) - [\varphi, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi]_{Lts}(w_1, \varphi(x_2), w_3) \\
&\quad - [\varphi, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi]_{Lts}(\varphi(x_1), w_2, \varphi(x_3)) \\
&= 6[\varphi(x_1), \varphi(x_2), \varphi(x_3)]_{\mathfrak{h}},
\end{aligned}$$

and

$$ad_{\varphi}^n \bar{\omega} = 0, \quad ad_{\varphi}^n(d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \bar{\omega}) = 0, \quad \forall n \geq 3.$$

Therefore, Maurer-Cartan elements $\bar{\omega}'$ and $\bar{\omega}$ are gauge equivalent if and only if

$$\begin{aligned}
\bar{\omega}' &= \bar{\omega} + [\varphi, \pi]_{Lts} + \frac{1}{2!} [\varphi, [\varphi, \pi]_{Lts}]_{Lts} \\
&\quad - (d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi + \frac{1}{2!} [\varphi, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi]_{Lts} + \frac{1}{3!} [\varphi, [\varphi, d_{\pi_{\mathfrak{g} \oplus \mathfrak{h}}} \varphi]_{Lts}]_{Lts}).
\end{aligned}$$

Hence, two Maurer-Cartan elements $\bar{\omega}$ and $\bar{\omega}'$ are equivalent if and only if Eqs. (24)-(26) hold.

The proof is completed. \square

5. INDUCIBILITY OF PAIRS OF LIE TRIPLE SYSTEM AUTOMORPHISMS

In this section, we study inducibility of a pair of Lie triple system automorphisms and characterize them by equivalent conditions.

Let \mathfrak{g} and \mathfrak{h} be two Lie triple systems, and

$$0 \longrightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \longrightarrow 0,$$

be a non-abelian extension of \mathfrak{g} by \mathfrak{h} with a section s of p . For any automorphism $\gamma \in \text{Aut}(\hat{\mathfrak{g}})$, then $\gamma \in \text{Aut}(\mathfrak{h})$ if and only if $\gamma(\mathfrak{h}) \subseteq \mathfrak{h}$. Denote $\text{Aut}_{\mathfrak{h}}(\hat{\mathfrak{g}}) = \{\gamma \in \text{Aut}(\hat{\mathfrak{g}}) \mid \gamma(\mathfrak{h}) = \mathfrak{h}\}$. We define a

linear map $\bar{\gamma} : \mathfrak{g} \longrightarrow \mathfrak{g}$ by

$$\bar{\gamma}(x) = p\gamma s(x), \quad \forall x \in \mathfrak{g}.$$

Assume that s_1 and s_2 are two distinct sections of p , since $ps_1(x) - ps_2(x) = 0$, $s_1(x) - s_2(x) \in \text{Ker } p \cong \mathfrak{h}$, which indicates that $\gamma(s_1(x) - s_2(x)) \in \mathfrak{h}$. Thus, $p\gamma s_1(x) = p\gamma s_2(x)$, which yields that $\bar{\gamma}$ is independent on the choice of sections.

For all $x, y, z \in \mathfrak{g}$, since $p|_{\mathfrak{h}} = 0$,

$$\begin{aligned} \bar{\gamma}([x, y, z]_{\mathfrak{g}}) &= p\gamma(s[x, y, z]_{\mathfrak{g}}) \\ &= p\gamma([s(x), s(y), s(z)]_{\hat{\mathfrak{g}}} - \omega(x, y, z)) \\ &= p\gamma([s(x), s(y), s(z)]_{\hat{\mathfrak{g}}}) \\ &= [p\gamma s(x), p\gamma s(y), p\gamma s(z)]_{\mathfrak{g}} \\ &= [\bar{\gamma}(x), \bar{\gamma}(y), \bar{\gamma}(z)]_{\mathfrak{g}}, \end{aligned}$$

which yields that $\bar{\gamma}$ is a homomorphism of Lie triple systems. It is easy to check that $\bar{\gamma}$ is bijective. Thus, $\bar{\gamma} \in \text{Aut}(\mathfrak{g})$. Then we can define a group homomorphism

$$\lambda : \text{Aut}_{\mathfrak{h}}(\hat{\mathfrak{g}}) \longrightarrow \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h}), \quad \lambda(\gamma) = (\bar{\gamma}, \gamma|_{\mathfrak{h}}).$$

Definition 5.1. A pair $(\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h})$ is said to be inducible if (α, β) is an image of λ .

In the following, we investigate when a pair (α, β) is inducible.

Theorem 5.2. Let $0 \longrightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \longrightarrow 0$ be a non-abelian extension of the Lie triple system \mathfrak{g} by \mathfrak{h} with a section s of p and (ω, θ, ρ) be the corresponding non-abelian 3-cocycle induced by s . A pair $(\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h})$ is inducible if and only if there is a linear map $\varphi : \mathfrak{g} \longrightarrow \mathfrak{h}$ satisfying the following conditions:

(39)

$$\beta(\theta(x, y)a) - \theta(\alpha(x), \alpha(y))\beta(a) = [\beta(a), \varphi(x), \varphi(y)]_{\mathfrak{h}} - D_{\rho}(\alpha(y))(\beta(a), \varphi(x)) + \rho(\alpha(x))(\beta(a), \varphi(y)),$$

(40)

$$\beta(\rho(x)(a, b)) - \rho(\alpha(x))(\beta(a), \beta(b)) = [\beta(a), \varphi(x), \beta(b)]_{\mathfrak{h}},$$

(41)

$$\begin{aligned} \beta\omega(x, y, z) - \omega(\alpha(x), \alpha(y), \alpha(z)) &= D_{\rho}(\alpha(z))(\varphi(x), \varphi(y)) - \rho(\alpha(y))(\varphi(x), \varphi(z)) - \theta(\alpha(y), \alpha(z))\varphi(x) \\ &\quad + \rho(\alpha(x))(\varphi(y), \varphi(z)) + \theta(\alpha(x), \alpha(z))\varphi(y) - D_{\theta}(\alpha(x), \alpha(y))\varphi(z) + \varphi([x, y, z]_{\mathfrak{g}}) - [\varphi(x), \varphi(y), \varphi(z)]_{\mathfrak{h}}, \end{aligned}$$

for all $x, y, z \in \mathfrak{g}$ and $a, b \in \mathfrak{h}$.

Proof. Suppose that $(\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h})$ is inducible, then there is an automorphism $\gamma \in \text{Aut}_{\mathfrak{h}}(\hat{\mathfrak{g}})$ such that $\gamma|_{\mathfrak{h}} = \beta$ and $p\gamma s = \alpha$. For all $x \in \mathfrak{g}$, since s is a section of p , that is, $ps = I_{\mathfrak{g}}$,

$$p(\gamma s - s\alpha)(x) = \alpha(x) - \alpha(x) = 0,$$

which implies that $(\gamma s - s\alpha)(x) \in \ker p \cong \mathfrak{h}$. So we can define a linear map $\varphi : \mathfrak{g} \longrightarrow \mathfrak{h}$ by

$$\varphi(x) = (s\alpha - \gamma s)(x), \quad \forall x \in \mathfrak{g}.$$

For $x, y \in \mathfrak{g}, a \in \mathfrak{h}$, using Eqs. (31)-(32), we get

$$\begin{aligned} & \beta(\theta(x, y)a) - \theta(\alpha(x), \alpha(y))\beta(a) \\ = & \beta[a, s(x), s(y)]_{\hat{\mathfrak{g}}} - [\beta(a), s\alpha(x), s\alpha(y)]_{\hat{\mathfrak{g}}} \\ = & [\beta(a), \beta s(x), \beta s(y)]_{\hat{\mathfrak{g}}} - [\beta(a), s\alpha(x), s\alpha(y)]_{\hat{\mathfrak{g}}} \\ = & [\beta(a), \beta s(x) - s\alpha(x), \beta s(y)]_{\hat{\mathfrak{g}}} + [\beta(a), s\alpha(x), \beta s(y)]_{\hat{\mathfrak{g}}} - [\beta(a), s\alpha(x), s\alpha(y)]_{\hat{\mathfrak{g}}} \\ = & [\beta(a), \varphi(x), \beta s(y)]_{\hat{\mathfrak{g}}} + [\beta(a), s\alpha(x), \varphi(y)]_{\hat{\mathfrak{g}}} \\ = & [\beta(a), \varphi(x), \beta s(y) - s\alpha(y)]_{\hat{\mathfrak{g}}} + [\beta(a), \varphi(x), s\alpha(y)]_{\hat{\mathfrak{g}}} + [\beta(a), s\alpha(x), \varphi(y)]_{\hat{\mathfrak{g}}} \\ = & [\beta(a), \varphi(x), \varphi(y)]_{\hat{\mathfrak{g}}} + [\beta(a), \varphi(x), s\alpha(y)]_{\hat{\mathfrak{g}}} - [s\alpha(x), \beta(a), \varphi(y)]_{\hat{\mathfrak{g}}} \\ = & [\beta(a), \varphi(x), \varphi(y)]_{\mathfrak{h}} + D_{\rho}(\alpha(y))(\beta(a), \varphi(x)) - \rho(\alpha(x))(\beta(a), \varphi(y)), \end{aligned}$$

that is, Eq. (39) holds. Similarly, Eqs. (40) and (41) hold.

Conversely, suppose that $(\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h})$ and there is a linear map $\varphi : \mathfrak{g} \longrightarrow \mathfrak{h}$ satisfying Eqs. (39)-(41). Due to s being a section of p , then all $\hat{w} \in \hat{\mathfrak{g}}$ can be written as $\hat{w} = a + s(x)$ for some $a \in \mathfrak{h}, x \in \mathfrak{g}$. Define a linear map $\gamma : \hat{\mathfrak{g}} \longrightarrow \hat{\mathfrak{g}}$ by

$$\gamma(\hat{w}) = \gamma(a + s(x)) = \beta(a) - \varphi(x) + s\alpha(x).$$

If $\gamma(\hat{w}) = 0$, then $s\alpha(x) = 0$ and $\beta(a) + \varphi(x) = 0$. In view of s and α being injective, we get $x = 0$, it follows that $a = 0$. Thus, $\hat{w} = a + s(x) = 0$, that is γ is injective. For any $\hat{w} = a + s(x) \in \hat{\mathfrak{g}}$,

$$\gamma(\beta^{-1}(a) + \beta^{-1}\varphi\alpha^{-1}(x) + s\alpha^{-1}(x)) = a + s(x) = \hat{w},$$

which yields that γ is surjective. In all, γ is bijective.

Next we show that γ is a homomorphism of Lie triple systems. In fact, for all $\hat{w}_i = a_i + s(x_i) \in \hat{\mathfrak{g}}$ ($i = 1, 2, 3$),

$$\begin{aligned} & [\gamma(\hat{w}_1), \gamma(\hat{w}_2), \gamma(\hat{w}_3)]_{\hat{\mathfrak{g}}} \\ = & [\beta(a_1) - \varphi(x_1) + s\alpha(x_1), \beta(a_2) - \varphi(x_2) + s\alpha(x_2), \beta(a_3) - \varphi(x_3) + s\alpha(x_3)]_{\hat{\mathfrak{g}}} \\ = & [\beta(a_1), \beta(a_2), \beta(a_3)]_{\hat{\mathfrak{g}}} - [\beta(a_1), \beta(a_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} + [\beta(a_1), \beta(a_2), s\alpha(x_3)]_{\hat{\mathfrak{g}}} \\ & - [\beta(a_1), \varphi(x_2), \beta(a_3)]_{\hat{\mathfrak{g}}} + [\beta(a_1), \varphi(x_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} - [\beta(a_1), \varphi(x_2), s\alpha(x_3)]_{\hat{\mathfrak{g}}} \\ & + [\beta(a_1), s\alpha(x_2), \beta(a_3)]_{\hat{\mathfrak{g}}} - [\beta(a_1), s\alpha(x_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} + [\beta(a_1), s\alpha(x_2), s\alpha(x_3)]_{\hat{\mathfrak{g}}} \\ & - [\varphi(x_1), \beta(a_2), \beta(a_3)]_{\hat{\mathfrak{g}}} + [\varphi(x_1), \beta(a_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} - [\varphi(x_1), \beta(a_2), s\alpha(x_3)]_{\hat{\mathfrak{g}}} \\ & + [\varphi(x_1), \varphi(x_2), \beta(a_3)]_{\hat{\mathfrak{g}}} - [\varphi(x_1), \varphi(x_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} + [\varphi(x_1), \varphi(x_2), s\alpha(x_3)]_{\hat{\mathfrak{g}}} \\ & - [\varphi(x_1), s\alpha(x_2), \beta(a_3)]_{\hat{\mathfrak{g}}} + [\varphi(x_1), s\alpha(x_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} - [\varphi(x_1), s\alpha(x_2), s\alpha(x_3)]_{\hat{\mathfrak{g}}} \end{aligned}$$

$$\begin{aligned}
& +[s\alpha(x_1), \beta(a_2), \beta(a_3)]_{\hat{\mathfrak{g}}} - [s\alpha(x_1), \beta(a_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} + [s\alpha(x_1), \beta(a_2), s\alpha(x_3)]_{\hat{\mathfrak{g}}} \\
& - [s\alpha(x_1), \varphi(x_2), \beta(a_3)]_{\hat{\mathfrak{g}}} + [s\alpha(x_1), \varphi(x_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} - [s\alpha(x_1), \varphi(x_2), s\alpha(x_3)]_{\hat{\mathfrak{g}}} \\
& + [s\alpha(x_1), s\alpha(x_2), \beta(a_3)]_{\hat{\mathfrak{g}}} - [s\alpha(x_1), s\alpha(x_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} + [s\alpha(x_1), s\alpha(x_2), s\alpha(x_3)]_{\hat{\mathfrak{g}}} \\
= & \beta([a_1, a_2, a_3]_{\hat{\mathfrak{g}}}) - [\beta(a_1), \beta(a_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} + D_\rho(\alpha(x_3))(\beta(a_1), \beta(a_2)) \\
& - [\beta(a_1), \varphi(x_2), \beta(a_3)]_{\hat{\mathfrak{g}}} + [\beta(a_1), \varphi(x_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} - D_\rho(\alpha(x_3))(\beta(a_1), \varphi(x_2)) \\
& - \rho(\alpha(x_2))(\beta(a_1), \beta(a_3)) + \rho(\alpha(x_2))(\beta(a_1), \varphi(x_3)) + \theta(\alpha(x_2), \alpha(x_3))\beta(a_1) \\
& - [\varphi(x_1), \beta(a_2), \beta(a_3)]_{\hat{\mathfrak{g}}} + [\varphi(x_1), \beta(a_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} - D_\rho(\alpha(x_3))(\varphi(x_1), \beta(a_2)) \\
& + [\varphi(x_1), \varphi(x_2), \beta(a_3)]_{\hat{\mathfrak{g}}} - [\varphi(x_1), \varphi(x_2), \varphi(x_3)]_{\hat{\mathfrak{g}}} + D_\rho(\alpha(x_3))(\varphi(x_1), \varphi(x_2)) \\
& + \rho(\alpha(x_2))(\varphi(x_1), \beta(a_3)) - \rho(\alpha(x_2))(\varphi(x_1), \varphi(x_3)) - \theta(\alpha(x_2), \alpha(x_3))\varphi(x_1) \\
& + \rho(\alpha(x_1))(\beta(a_2), \beta(a_3)) - \rho(\alpha(x_1))(\beta(a_2), \varphi(x_3)) - \theta(\alpha(x_1), \alpha(x_3))\beta(a_2) \\
& - \rho(\alpha(x_1))(\varphi(x_2), \beta(a_3)) + \rho(\alpha(x_1))(\varphi(x_2), \varphi(x_3)) + \theta(\alpha(x_1), \alpha(x_3))\varphi(x_2) \\
& + D_\theta(\alpha(x_1), \alpha(x_2))\beta(a_3) - D_\theta(\alpha(x_1), \alpha(x_2))\varphi(x_3) + \omega(\alpha(x_1), \alpha(x_2), \alpha(x_3)) \\
& + s[\alpha(x_1), \alpha(x_2), \alpha(x_3)]_{\hat{\mathfrak{g}}}
\end{aligned}$$

and

$$\begin{aligned}
& \gamma([\hat{w}_1, \hat{w}_2, \hat{w}_3]_{\hat{\mathfrak{g}}}) \\
= & \gamma([a_1, a_2, a_3]_{\hat{\mathfrak{g}}} + [a_1, a_2, s(x_3)]_{\hat{\mathfrak{g}}} + [a_1, s(x_2), a_3]_{\hat{\mathfrak{g}}} + [a_1, s(x_2), s(x_3)]_{\hat{\mathfrak{g}}} + [s(x_1), a_2, s(x_3)]_{\hat{\mathfrak{g}}} \\
& + [s(x_1), s(x_2), a_3]_{\hat{\mathfrak{g}}} + [s(x_1), a_2, a_3]_{\hat{\mathfrak{g}}} + \omega(x_1, x_2, x_3) + s[x_1, x_2, x_3]_{\hat{\mathfrak{g}}}) \\
= & \beta([a_1, a_2, a_3]_{\hat{\mathfrak{g}}}) + \beta(D_\rho(x_3)(a_1, a_2) - \rho(x_2)(a_1, a_3) + \theta(x_2, x_3)a_1 - \theta(x_1, x_3)a_2 \\
& + D_\theta(x_1, x_2)a_3 + \rho(x_1)(a_2, a_3)) + \beta\omega(x_1, x_2, x_3) - \varphi[x_1, x_2, x_3]_{\hat{\mathfrak{g}}} + s\alpha[x_1, x_2, x_3]_{\hat{\mathfrak{g}}}.
\end{aligned}$$

Thanks to Eqs. (39)-(41), we have

$$\gamma[\hat{w}_1, \hat{w}_2, \hat{w}_3]_{\hat{\mathfrak{g}}} = [\gamma(\hat{w}_1), \gamma(\hat{w}_2), \gamma(\hat{w}_3)]_{\hat{\mathfrak{g}}},$$

which implies that γ is a homomorphism of Lie triple systems. Thus, $\gamma \in \text{Aut}(\hat{\mathfrak{g}})$. Finally, we show that $\gamma|_{\mathfrak{h}} = \beta$ and $p\gamma s = \alpha$. In fact,

$$\gamma(a) = \gamma(a + s(0)) = \beta(a), \quad \forall a \in \mathfrak{h}$$

and

$$(p\gamma s)(x) = p\gamma(0 + s(x)) = p(s\alpha(x) - \varphi(x)) = ps\alpha(x) = \alpha(x), \quad \forall x \in \mathfrak{g}.$$

Therefore, $\gamma|_{\mathfrak{h}} = \beta$ and $p\gamma s = \alpha$. Thus, $(\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h})$ is inducible. \square

Let $0 \longrightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \longrightarrow 0$ be a non-abelian extension of the Lie triple system \mathfrak{g} by \mathfrak{h} with a section s of p and (ω, θ, ρ) be the corresponding non-abelian 3-cocycle induced by s .

For all $(\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h})$. Define maps $\omega_{(\alpha, \beta)} : \mathfrak{g} \times \mathfrak{g} \times \mathfrak{g} \longrightarrow \mathfrak{h}$, $\theta_{(\alpha, \beta)} : \mathfrak{g} \times \mathfrak{g} \longrightarrow \mathfrak{gl}(\mathfrak{h})$, $\rho_{(\alpha, \beta)} : \mathfrak{g} \longrightarrow \text{Hom}(\mathfrak{h} \times \mathfrak{h}, \mathfrak{h})$ respectively by

$$(42) \quad \omega_{(\alpha, \beta)}(x, y, z) = \beta\omega(\alpha^{-1}(x), \alpha^{-1}(y), \alpha^{-1}(z)),$$

$$(43) \quad \theta_{(\alpha, \beta)}(x, y)a = \beta\theta(\alpha^{-1}(x), \alpha^{-1}(y))\beta^{-1}(a),$$

$$(44) \quad \rho_{(\alpha, \beta)}(x)(a, b) = \beta\rho(\alpha^{-1}(x))(\beta^{-1}(a), \beta^{-1}(b)).$$

By direct computations, we get

$$(45) \quad D_{\theta_{(\alpha, \beta)}}(x, y) = \beta D_{\theta}(\alpha^{-1}(x), \alpha^{-1}(y))\beta^{-1}(a), \quad D_{\rho_{(\alpha, \beta)}}(a, b) = \beta D_{\rho}(\alpha^{-1}(x))(\beta^{-1}(a), \beta^{-1}(b)).$$

for all $x, y, z \in \mathfrak{g}$, $a, b \in \mathfrak{h}$.

Proposition 5.3. *With the above notations, $(\omega_{(\alpha, \beta)}, \theta_{(\alpha, \beta)}, \rho_{(\alpha, \beta)})$ is a non-abelian 3-cocycle.*

Proof. For all $x_1, x_2, y_1, y_2, y_3 \in \mathfrak{g}$, using Eqs. (12), (42), (43) and (45), we get

$$\begin{aligned} & D_{\theta_{(\alpha, \beta)}}(x_1, x_2)\omega_{(\alpha, \beta)}(y_1, y_2, y_3) + \omega_{(\alpha, \beta)}(x_1, x_2, [y_1, y_2, y_3]_{\mathfrak{g}}) \\ &= \beta \left(D_{\theta}(\alpha^{-1}(x_1), \alpha^{-1}(x_2))\beta^{-1}(\beta\omega_{(\alpha, \beta)}(\alpha^{-1}(y_1), \alpha^{-1}(y_2), \alpha^{-1}(y_3))) \right. \\ & \quad \left. - \beta(\omega_{(\alpha, \beta)}(\alpha^{-1}(x_1), \alpha^{-1}(x_2), \alpha^{-1}([y_1, y_2, y_3]_{\mathfrak{g}}))) \right) \\ &= \beta\omega([\alpha^{-1}(x_1), \alpha^{-1}(x_2), \alpha^{-1}(y_1)]_{\mathfrak{g}}, \alpha^{-1}(y_2), \alpha^{-1}(y_3)) \\ & \quad + \beta(\theta(\alpha^{-1}(y_2), \alpha^{-1}(y_3))\omega(\alpha^{-1}(x_1), \alpha^{-1}(x_2), \alpha^{-1}(y_1))) \\ & \quad + \beta\omega(\alpha^{-1}(y_1), [\alpha^{-1}(x_1), \alpha^{-1}(x_2), \alpha^{-1}(y_2)]_{\mathfrak{g}}, \alpha^{-1}(y_3)) \\ & \quad - \beta(\theta(\alpha^{-1}(y_1), \alpha^{-1}(y_3))\omega(\alpha^{-1}(x_1), \alpha^{-1}(x_2), \alpha^{-1}(y_2))) \\ & \quad + \beta\omega(\alpha^{-1}(y_1), \alpha^{-1}(y_2), [\alpha^{-1}(x_1), \alpha^{-1}(x_2), \alpha^{-1}(y_3)]_{\mathfrak{g}}) \\ & \quad + \beta(D_{\theta}(\alpha^{-1}(y_1), \alpha^{-1}(y_2))\omega(\alpha^{-1}(x_1), \alpha^{-1}(x_2), \alpha^{-1}(y_3))) \\ &= \omega_{(\alpha, \beta)}([x_1, x_2, y_1]_{\mathfrak{g}}, y_2, y_3) + \theta_{(\alpha, \beta)}(y_2, y_3)\omega_{(\alpha, \beta)}(x_1, x_2, y_1) + \omega_{(\alpha, \beta)}(y_1, [x_1, x_2, y_2]_{\mathfrak{g}}, y_3) \\ & \quad - \theta_{(\alpha, \beta)}(y_1, y_3)\omega_{(\alpha, \beta)}(x_1, x_2, y_2) + \omega_{(\alpha, \beta)}(y_1, y_2, [x_1, x_2, y_3]_{\mathfrak{g}}) + D_{\theta_{(\alpha, \beta)}}(y_1, y_2)\omega_{(\alpha, \beta)}(x_1, x_2, y_3), \end{aligned}$$

which implies that Eq. (12) holds for $\omega_{(\alpha, \beta)}, \theta_{(\alpha, \beta)}$. Take the same procedure, Eqs. (11) and (13)-(22) hold for $\omega_{(\alpha, \beta)}, \theta_{(\alpha, \beta)}, D_{\theta_{(\alpha, \beta)}}, \rho_{(\alpha, \beta)}$ and $D_{\rho_{(\alpha, \beta)}}$. The proof is finished. \square

Theorem 5.4. *Let $0 \longrightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \longrightarrow 0$ be a non-abelian extension of \mathfrak{g} by \mathfrak{h} with a section s of p and (ω, θ, ρ) be the corresponding non-abelian 3-cocycle induced by s . A pair $(\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h})$ is inducible if and only if the non-abelian 3-cocycles (ω, θ, ρ) and $(\omega_{(\alpha, \beta)}, \theta_{(\alpha, \beta)}, \rho_{(\alpha, \beta)})$ are equivalent.*

Proof. Suppose $(\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h})$ is inducible, then by Theorem 5.2, there is a linear map $\varphi : \mathfrak{g} \rightarrow \mathfrak{h}$ satisfying Eqs. (39)-(41). For all $x, y \in \mathfrak{g}, a \in \mathfrak{h}$, there exist $x_0, y_0 \in \mathfrak{g}, a_0 \in \mathfrak{h}$ such that $x = \alpha(x_0), y = \alpha(y_0), a = \beta(a_0)$. Therefore, using Eqs. (39) and (43),

$$\begin{aligned}
& \theta_{\alpha, \beta}(x, y)a - \theta(x, y)a \\
&= \beta(\theta(\alpha^{-1}(x), \alpha^{-1}(y))\beta^{-1}(a)) - \theta(x, y)a \\
&= \beta(\theta(x_0, y_0)a_0) - \theta(\alpha(x_0), \alpha(y_0))\beta(a_0) \\
&= [\beta(a_0), \varphi(x_0), \varphi(y_0)]_{\mathfrak{h}} - D_{\rho}(\alpha(y_0))(\beta(a_0), \varphi(x_0)) + \rho(x_0)(\beta(a_0), \varphi(y_0)) \\
&= [a, \varphi\alpha^{-1}(x), \varphi\alpha^{-1}(y)]_{\mathfrak{h}} - D_{\rho}(y)(a, \varphi\alpha^{-1}(x)) + \rho(x)(a, \varphi\alpha^{-1}(y)),
\end{aligned}$$

which yields that Eq. (25) holds. Analogously, Eqs. (24) and (26) hold. Hence, the non-abelian 3-cocycles $(\omega_{(\alpha, \beta)}, \theta_{(\alpha, \beta)}, \rho_{(\alpha, \beta)})$ and (ω, θ, ρ) are equivalent via the linear map $\varphi\alpha^{-1} : \mathfrak{g} \rightarrow \mathfrak{h}$.

The converse part can be checked similarly. □

6. WELLS EXACT SEQUENCES FOR LIE TRIPLE SYSTEMS

In this section, we are focus on the Wells map associated with non-abelian extensions of Lie triple systems.

Let $\mathcal{E} : 0 \rightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \rightarrow 0$ be a non-abelian extension of \mathfrak{g} by \mathfrak{h} with a section s of p . Suppose that (ω, θ, ρ) is the corresponding non-abelian 3-cocycle induced by s . Define a linear map $W : \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h}) \rightarrow H_{\text{tab}}^3(\mathfrak{g}, \mathfrak{h})$ by

$$(46) \quad W(\alpha, \beta) = [(\omega_{(\alpha, \beta)}, \theta_{(\alpha, \beta)}, \rho_{(\alpha, \beta)}) - (\omega, \theta, \rho)].$$

The map W is called the Wells map. In view of Lemma 3.8, the Wells map W does not depend on the choice of sections.

Theorem 6.1. *Assume that $\mathcal{E} : 0 \rightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \rightarrow 0$ is a non-abelian extension of \mathfrak{g} by \mathfrak{h} with a section s of p . Then there is an exact sequence:*

$$1 \rightarrow \text{Aut}_{\mathfrak{h}}^{\mathfrak{g}}(\hat{\mathfrak{g}}) \xrightarrow{H} \text{Aut}_{\mathfrak{h}}(\hat{\mathfrak{g}}) \xrightarrow{\lambda} \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h}) \xrightarrow{W} H_{\text{tab}}^3(\mathfrak{g}, \mathfrak{h}),$$

where $\text{Aut}_{\mathfrak{h}}^{\mathfrak{g}}(\hat{\mathfrak{g}}) = \{\gamma \in \text{Aut}(\hat{\mathfrak{g}}) | \lambda(\gamma) = (I_{\mathfrak{g}}, I_{\mathfrak{h}})\}$.

Proof. Obviously, $\text{Ker}\lambda = \text{Im}H$ and H is injective. By Theorem 5.4, one can easily check that $\text{Ker}W = \text{Im}\lambda$. This completes the proof. □

Let $\mathcal{E} : 0 \longrightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \longrightarrow 0$ be a non-abelian extension of \mathfrak{g} by \mathfrak{h} with a section s of p . Assume that (ω, θ, ρ) is the non-abelian 3-cocycle corresponding to \mathcal{E} induced by s . Denote

$$(47) \quad Z_{nab}^1(\mathfrak{g}, \mathfrak{h}) = \left\{ \varphi : \mathfrak{g} \rightarrow \mathfrak{h} \left\{ \begin{array}{l} [a, \varphi(x), b]_{\mathfrak{h}} = 0, \\ [a, \varphi(x), \varphi(y)]_{\mathfrak{h}} - D_{\rho}(y)(a, \varphi(x)) + \rho(x)(a, \varphi(y)) = 0, \\ D_{\rho}(z)(\varphi(x), \varphi(y)) - \rho(y)(\varphi(x), \varphi(z)) - \theta(y, z)\varphi(x) \\ + \rho(x)(\varphi(y), \varphi(z)) + \theta(x, z)\varphi(y) - D_{\theta}(x, y)\varphi(z) \\ = [\varphi(x), \varphi(y), \varphi(z)]_{\mathfrak{h}} - \varphi([x, y, z]_{\mathfrak{g}}), \quad \forall x, y, z \in \mathfrak{g}, a, b \in \mathfrak{h} \end{array} \right. \right\}.$$

It is easy to check that $Z_{nab}^1(\mathfrak{g}, \mathfrak{h})$ is an abelian group, which is called a non-abelian 1-cocycle on \mathfrak{g} with values in \mathfrak{h} .

Proposition 6.2. *With the above notations, we have*

(i) *The linear map $K : \text{Ker}\lambda \longrightarrow Z_{nab}^1(\mathfrak{g}, \mathfrak{h})$ given by*

$$(48) \quad K(\gamma)(x) = \varphi_{\gamma}(x) = s(x) - \gamma s(x), \quad \forall \gamma \in \text{Ker}\lambda, x \in \mathfrak{g}$$

is a homomorphism of groups.

(ii) *K is an isomorphism, that is, $\text{Ker}\lambda \simeq Z_{nab}^1(\mathfrak{g}, \mathfrak{h})$.*

Proof. (i) Using Eqs. (30)-(32) and (48), for all $x, y, z \in \mathfrak{g}$, we have,

$$\begin{aligned} & [\varphi_{\gamma}(x), \varphi_{\gamma}(y), \varphi_{\gamma}(z)]_{\mathfrak{h}} - D_{\rho}(z)(\varphi_{\gamma}(x), \varphi_{\gamma}(y)) + \rho(y)(\varphi_{\gamma}(x), \varphi_{\gamma}(z)) + \theta(y, z)\varphi_{\gamma}(x) \\ & - \rho(x)(\varphi_{\gamma}(y), \varphi_{\gamma}(z)) - \theta(x, z)\varphi_{\gamma}(y) + D_{\theta}(x, y)\varphi_{\gamma}(z) - \varphi_{\gamma}(\{x, y, z\}_{\mathfrak{g}}) \\ = & [s(x) - \gamma s(x), s(y) - \gamma s(y), s(z) - \gamma s(z)]_{\hat{\mathfrak{g}}} - [s(x) - \gamma s(x), s(y) - \gamma s(y), s(z)]_{\hat{\mathfrak{g}}} \\ & + [s(y), s(x) - \gamma s(x), s(z) - \gamma s(z)]_{\hat{\mathfrak{g}}} + [s(x) - \gamma s(x), s(y), s(z)]_{\hat{\mathfrak{g}}} - [s(x), s(y) - \gamma s(y), s(z) - \gamma s(z)]_{\hat{\mathfrak{g}}} \\ & - [s(y) - \gamma s(y), s(x), s(z)]_{\hat{\mathfrak{g}}} + [s(x), s(y), s(z) - \gamma s(z)]_{\hat{\mathfrak{g}}} + \gamma s([x, y, z]_{\mathfrak{g}}) - s([x, y, z]_{\mathfrak{g}}) \\ = & \gamma s([x, y, z]_{\mathfrak{g}}) - [\gamma s(x), \gamma s(y), \gamma s(z)]_{\hat{\mathfrak{g}}} + [s(x), s(y), s(z)]_{\hat{\mathfrak{g}}} - s([x, y, z]_{\mathfrak{g}}) \\ = & \omega(x, y, z) - \gamma \omega(x, y, z) \\ = & 0. \end{aligned}$$

Analogously, we can check that φ_{γ} satisfies the other identities in $Z_{nab}^1(\mathfrak{g}, \mathfrak{h})$. Thus, K is well-defined. For any $\gamma_1, \gamma_2 \in \text{Ker}\lambda$ and $x \in \mathfrak{g}$, suppose $K(\gamma_1) = \varphi_{\gamma_1}$ and $K(\gamma_2) = \varphi_{\gamma_2}$. By Eq. (48), we have

$$\begin{aligned} K(\gamma_1\gamma_2)(x) &= s(x) - \gamma_1\gamma_2 s(x) \\ &= s(x) - \gamma_1(s(x) - \varphi_{\gamma_2}(x)) \\ &= s(x) - \gamma_1 s(x) + \gamma_1 \varphi_{\gamma_2}(x) \\ &= \varphi_{\gamma_1}(x) + \varphi_{\gamma_2}(x) \end{aligned}$$

which means that $K(\gamma_1\gamma_2) = K(\gamma_1) + K(\gamma_2)$ is a homomorphism of groups.

(ii) For all $\gamma \in \text{Ker}\lambda$, we obtain that $\lambda(\gamma) = (p\gamma s, \gamma|_{\mathfrak{h}}) = (I_{\mathfrak{g}}, I_{\mathfrak{h}})$. If $K(\gamma) = \varphi_\gamma = 0$, we can get $\varphi_\gamma(x) = s(x) - \gamma s(x) = 0$, that is, $\gamma = I_{\mathfrak{g}}$, which indicates that K is injective. Secondly, we prove that K is surjective. Since all $\hat{x} \in \hat{\mathfrak{g}}$ can be written as $a + s(x)$ for some $a \in \mathfrak{h}, x \in \mathfrak{g}$, for any $\varphi \in Z_{nab}^1(\mathfrak{g}, \mathfrak{h})$, we define a linear map $\gamma : \hat{\mathfrak{g}} \rightarrow \hat{\mathfrak{g}}$ by

$$(49) \quad \gamma(\hat{x}) = \gamma(a + s(x)) = s(x) - \varphi(x) + a, \quad \forall \hat{x} \in \hat{\mathfrak{g}}.$$

It is obviously that $(p\gamma s, \gamma|_{\mathfrak{h}}) = (I_{\mathfrak{g}}, I_{\mathfrak{h}})$. We need to verify that γ is an automorphism of Lie triple system $\hat{\mathfrak{g}}$. One can take the same procedure as the proof of the converse part of Theorem 5.2. It follows that $\gamma \in \text{Ker}\lambda$. Thus, K is surjective. In all, K is bijective. So $\text{Ker}\lambda \simeq Z_{nab}^1(\mathfrak{g}, \mathfrak{h})$. □

Combining Theorem 6.1 and Proposition 6.2, we have

Theorem 6.3. *Let $\mathcal{E} : 0 \rightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \rightarrow 0$ be a non-abelian extension of \mathfrak{g} by \mathfrak{h} . There is an exact sequence:*

$$0 \rightarrow Z_{nab}^1(\mathfrak{g}, \mathfrak{h}) \xrightarrow{i} \text{Aut}_{\mathfrak{h}}(\hat{\mathfrak{g}}) \xrightarrow{\lambda} \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h}) \xrightarrow{W} H_{nab}^3(\mathfrak{g}, \mathfrak{h}).$$

7. PARTICULAR CASE: ABELIAN EXTENSIONS OF LIE TRIPLE SYSTEMS

In this section, we interpret the results of previous section in particular case.

Let $\mathcal{E} : 0 \rightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \rightarrow 0$ be an abelian extension of \mathfrak{g} by \mathfrak{h} with a section s of p . Suppose that ω is a 3-cocycle corresponding to \mathcal{E} induced by s . We have,

Theorem 7.1 ([31]). *(i) The triple $(\mathfrak{g} \oplus \mathfrak{h}, [, ,]_{(\omega, \theta)})$ is a Lie triple system if and only if ω is a 3-cocycle of \mathfrak{g} with coefficients in the representation (\mathfrak{h}, θ) .*

(ii) Abelian extensions of \mathfrak{g} by \mathfrak{h} are classified by the cohomology group $H^3(\mathfrak{g}, \mathfrak{h})$ of \mathfrak{g} with coefficients in (\mathfrak{h}, θ) .

The following statement can be got directly by Theorem 5.2.

Theorem 7.2. *Let $\mathcal{E} : 0 \rightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \rightarrow 0$ be an abelian extension of \mathfrak{g} by \mathfrak{h} with a section s of p . Assume that ω is a 3-cocycle and (\mathfrak{h}, θ) is a representation of \mathfrak{g} associated to \mathcal{E} . A pair $(\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h})$ is inducible with respect to the abelian extension \mathcal{E} if and only if there is a linear map $\varphi : \mathfrak{g} \rightarrow \mathfrak{h}$ satisfying the following conditions:*

$$(50) \quad \begin{aligned} \beta\omega(x, y, z) - \omega(\alpha(x), \alpha(y), \alpha(z)) &= \theta(\alpha(x), \alpha(z))\varphi(y) - \theta(\alpha(y), \alpha(z))\varphi(x) \\ &\quad - D_{\theta}(\alpha(x), \alpha(y))\varphi(z) + \varphi([x, y, z]_{\mathfrak{g}}), \end{aligned}$$

$$(51) \quad \beta(\theta(x, y)a) = \theta(\alpha(x), \alpha(y))\beta(a).$$

According to Eqs. (7) and (51), we have

$$\beta D_{\theta}(x, y)a = D_{\theta}(\alpha(x), \alpha(y))\beta(a).$$

For all $(\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h})$, $\omega_{(\alpha, \beta)}$ may not be a 3-cocycle. Indeed, $\omega_{(\alpha, \beta)}$ is a 3-cocycle if Eq. (51) holds. Thus, it is natural to introduce the space of compatible pairs of automorphisms:

$$C_{\theta} = \left\{ (\alpha, \beta) \in \text{Aut}(\mathfrak{g}) \times \text{Aut}(\mathfrak{h}) \mid \beta(\theta(x, y)a) = \theta(\alpha(x), \alpha(y))\beta(a), \forall x, y \in \mathfrak{g}, a \in \mathfrak{h} \right\}.$$

Analogous to Theorem 5.4, we have

Theorem 7.3. *Let $\mathcal{E} : 0 \rightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \rightarrow 0$ be an abelian extension of \mathfrak{g} by \mathfrak{h} with a section s of p and ω be a 3-cocycle associated to \mathcal{E} induced by s . A pair $(\alpha, \beta) \in C_{\theta}$ is inducible with respect to the abelian extension \mathcal{E} if and only if ω and $\omega_{(\alpha, \beta)}$ are in the same cohomological class.*

In the case of abelian extensions, $Z_{nab}^1(\mathfrak{g}, \mathfrak{h})$ defined by (47) becomes to $H^1(\mathfrak{g}, \mathfrak{h})$ given in Section 2. In the light of Theorem 6.3 and Theorem 7.3, we have the following exact sequence:

Theorem 7.4. *Let $\mathcal{E} : 0 \rightarrow \mathfrak{h} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \rightarrow 0$ be an abelian extension of \mathfrak{g} by \mathfrak{h} . There is an exact sequence:*

$$0 \rightarrow H^1(\mathfrak{g}, \mathfrak{h}) \xrightarrow{i} \text{Aut}_{\mathfrak{h}}(\hat{\mathfrak{g}}) \xrightarrow{\lambda} C_{\theta} \xrightarrow{w} H^3(\mathfrak{g}, \mathfrak{h}).$$

Acknowledgments

The first author is supported by the NSF of China (No. 11871421), the Natural Science Foundation of Zhejiang Province of China (No. LY19A010001) and the Science and Technology Planning Project of Zhejiang Province (No. 2022C01118). The second author is supported by the NSF of China (No. 12161013) and Guizhou Provincial Basic Research Program (Natural Science) (No. ZK[2023]025).

Statements and Declarations

All datasets underlying the conclusions of the paper are available to readers. No conflict of interest exists in the submission of this manuscript.

REFERENCES

- [1] V. G. Bardakov, M. Singh, Extensions and automorphisms of Lie algebras, *J. Algebra Appl.* 16 (2017), 1750162. [2](#)
- [2] S. Chen, Q. Lou, Q. Sun, Cohomologies of Rota-Baxter Lie triple systems and applications, *Commun. Algebra* 51 (10) (2023), 4299-4315. [1](#)
- [3] T. Chtioui, A. Hajjaji, S. Mabrouk, A. Makhoulouf, Cohomologies and deformations of \mathcal{O} -operators on Lie triple systems, *J. Math. Phys.* 64, 081701 (2023). [11](#)
- [4] A. Das, N. Ratheeb, Extensions and automorphisms of Rota-Baxter groups, *J. Algebra* 636 (2023), 626–665. [2](#)

- [5] S. Eilenberg, S. MacLane, Cohomology theory in abstract groups. II. Group extensions with non-abelian kernel, *Ann. Math.* 48 (1947), 326-341. [1](#)
- [6] Y. Frégier, Non-abelian cohomology of extensions of Lie algebras as Deligne groupoid, *J. Algebra* 398 (2014), 243–257. [1](#)
- [7] S. Goswamia, S. K. Mishraa, G. Mukherjee, Automorphisms of extensions of Lie-Yamaguti algebras and inducibility problem, *J. Algebra* 641 (2024), 268–306. [2](#)
- [8] J. Gouray, A differential graded Lie algebra approach to non-abelian extensions of associative algebras, *arXiv:1802.04641*. [1](#)
- [9] Y. Guo, B. Hou, Crossed modules and non-abelian extensions of Rota-Baxter Leibniz algebras, *J. Geom.Phys.* 191 (2023), 104906. [1](#), [2](#)
- [10] S. K. Hazra, A. Habib, Wells exact sequence and automorphisms of extensions of Lie superalgebras, *J. Lie Theory* 30 (2020), 179–199. [2](#)
- [11] B. Harris, Cohomology of Lie triple systems and Lie algebras with involution, *Trans. Amer. Math. Soc.* 98 (1961), 148-162. [3](#)
- [12] L. Hodge, J. Parshall, On the representation theory of Lie triple systems, *Trans. Amer. Math. Soc.* 354 (2002), 4359-4391. [3](#)
- [13] B. Hou, J. Zhao, Crossed modules, non-abelian extensions of associative conformal algebras and Wells exact sequences, *arXiv:2211.10842v1*. [1](#), [2](#)
- [14] N. Inassaridze, E. Khmaladze, M. Ladra, Non-abelian cohomology and extensions of Lie algebras, *J. Lie Theory* 18 (2008), 413–432. [1](#)
- [15] N. Jacobson, Lie and Jordan triple systems, *Amer. J. Math.* 71 (1949), 149-170. [1](#)
- [16] P. Jin, H. Liu, The Wells exact sequence for the automorphism group of a group extension, *J. Algebra* 324 (2010), 1219–1228. [2](#)
- [17] F. Kubo, Y. Taniguchi, A controlling cohomology of the deformation theory of Lie triple systems, *J. Algebra* 278 (2004), 242-250. [3](#)
- [18] S. K. Mishra, A. Das, S. K. Hazra, Non-abelian extensions of Rota-Baxter Lie algebras and inducibility of automorphisms, *Linear. Algebra. Appl.* 669 (2023), 147-174. [1](#), [2](#)
- [19] W. Lister, A structure theory for Lie triple systems, *Trans. Amer. Math. Soc.* 72 (1952), 217-242. [1](#), [3](#)
- [20] J. Liu, Y. Sheng, Q. Wang, On non-abelian extensions of Leibniz algebras, *Commun. Algebra* 46 (2) (2018), 574–587. [1](#)
- [21] Q. Ma, L. Song, Y. Wang, Non-abelian extensions of pre-Lie algebras, *Commun. Algebra* 51 (4) (2023), 1370-1382. [1](#)
- [22] Y. Ma, L. Chen, J. Lin, Systems of quotients of Lie triple systems, *Commun. Algebra* 42 (2014), 3339-3349. [1](#)
- [23] N. Machado, J. Oinert, S. Wagner, Non-abelian extensions of groupoids and their groupoid rings, *arXiv:2207.03369*. [1](#)
- [24] K. H. Neeb, Non-abelian extensions of infinite-dimensional Lie groups, *Ann. Inst. Fourier (Grenoble)* 57 (2007), 209-271. [1](#)
- [25] I. B. S. Passi, M. Singh, M. K. Yadav, Automorphisms of abelian group extensions, *J. Algebra* 324 (2010), 820–830. [2](#)
- [26] L. Song, A. Makhlouf, R. Tang, On non-abelian extensions of 3-Lie algebras, *Commun. Theor. Phys.* 69 (2018), 347-356. [1](#), [11](#)
- [27] C. Wells, Automorphisms of group extensions, *Trans. Amer. Math. Soc.* 155 (1971), 189-194. [2](#)
- [28] R. Yadav, N. Behera, R. Bhutia, Equivariant one-parameter deformations of Lie triple systems, *J. Algebra* 568 (2021), 467-479. [1](#)
- [29] K. Yamaguti, On the cohomology space of Lie triple system, *Kumamoto J. Sci. Ser. A* 5 (1960), 44-52. [3](#)
- [30] K. Yamaguti, On the Lie triple system and its generalization, *J. Sci. Hiroshima Univ. Ser. A* 21, (1957-1958), 155-159. [3](#)
- [31] T. Zhang, Notes on cohomologies of Lie triple systems, *J. Lie Theory* 24(4)(2014), 909-929. [3](#), [7](#), [20](#)

QINXIU SUN, DEPARTMENT OF MATHEMATICS, ZHEJIANG UNIVERSITY OF SCIENCE AND TECHNOLOGY, HANGZHOU, 310023
Email address: qxsun@126.com

SHUANGJIAN GUO, CORRESPONDING AUTHOR, SCHOOL OF MATHEMATICS AND STATISTICS, GUIZHOU UNIVERSITY OF FINANCE AND ECONOMICS, GUIYANG, 550025
Email address: shuangjianguo@126.com