

# ON THE LIE ALGEBRA OF A MALCEV ALGEBRA

OLUFEMI O. OYADARE

Department of Mathematics,  
Obafemi Awolowo University,  
Ile-Ife, 220005, NIGERIA.  
E-mail: *femi\_oya@yahoo.com*

**Abstract.** *This paper develops the structure theory of a Malcev algebra via the consideration of its most important and largest Lie (sub-) algebra. We introduce the notion of a Lie algebra which uniquely corresponds to a Malcev algebra and use this correspondence to derive some basic properties of some types of ideals in the Malcev algebra. We then prove the exact nature of the root-space decomposition of a Malcev algebra.*

**Subject Classification:** 17D10, 17B05, 17B22

**Keywords:** Nonassociative algebra;  $J$ -Nucleus: Malcev algebra: Lie algebra

## §1. Introduction.

One of the most understood and well-developed class of algebras in the whole of mathematics is the class of *Lie algebras*. A Lie algebra is generally defined as an algebra  $\mathfrak{L}$  (over  $\mathbb{R}$  or  $\mathbb{C}$ ) whose product,  $[\cdot, \cdot]$ , is anti-commutative (i.e.,  $[x, x] = 0$ ) and in which the Jacobian,

$$[[x, y], z] + [[y, z], x] + [[z, x], y] =: J(x, y, z),$$

vanishes, for all  $x, y, z \in \mathfrak{L}$ . Numerous articles, monographs and textbooks by notable authors have been published detailing different aspects of a Lie algebra, their inter-relationships and applications to the life sciences. We shall refer to Varadarajan [10.] for orientation. Perhaps, the most singular reason for the extreme utility of a Lie algebra, over and above other algebras, stems out of the fact that every Lie algebra corresponds to a Lie group. This affords results to be exchanged between a Lie algebra (with its *infinitesimal* theory) and its corresponding Lie group (with its *topological* and *manifold* structures),

with one enriching the other in a uniquely determined way. Indeed no other class of algebras has had the privilege of the class of Lie algebras.

Since the *Malcev algebra* (also called the *Moufang-Malcev algebra*) was introduced (See Malcev [5.], Schafer [9.] and Paal [6.]) as a tangent space of an analytic *Moufang loop* and since it was shown that every Lie algebra is a Malcev algebra, mathematicians have been seeking the Lie-algebra type of approach to the understanding of a Malcev algebra. It is believed that a Malcev algebra is a higher version of a Lie algebra in which  $J = J(x, y, z)$  is not necessarily zero. Aspects of the structure theory of a Malcev algebra had been discussed by A. A. Sagle in his University of California thesis [8.]. It had also been found that it is possible to have a form of the Poincare-Birkhoff-Witt Theorem (known for a Lie algebra) for a Malcev algebra, as well as its (nonassociative) universal enveloping algebra (See [2.], [7.] and [9.]). All these partial results ([3.]) were however not enough to give the class of Malcev algebras the esteemed status occupied by one of its sub-classes, the (sub-) class of Lie algebras.

The present paper gives a novel approach to the study of a Malcev algebra via the consideration of a distinguished Lie (sub-) algebra closely attached to it. The paper is organised as follows: §2 considers the structure theory of a Malcev algebra while §3 develops its basic ideal theory. The root-space decomposition of a Malcev algebra is the subject of §4.

## §2. Structure theory of a Malcev algebra

We shall start by giving a precise definition to a Malcev algebra and then give a rapid presentation of aspects of its structure.

**2.1 Definition.** A *Malcev algebra*  $A$  over a field  $k$  is any nonassociative algebra over  $k$  endowed with a distributive multiplication  $[\cdot, \cdot]$  satisfying the relations

$$[a, a] = 0$$

and

$$[J(a, b, c), a] = J(a, b, [a, c]),$$

for all  $a, b, c \in A$ .  $\square$

It should be clear from the above definition that a Lie algebra is in particular a Malcev algebra: Indeed,  $[J(a, b, c), a] = [0, a] = 0 = J(a, b, [a, c])$ . One could then say that a Malcev algebra is a Lie algebra whose Jacobian

may not necessarily vanish but satisfies  $[J(a, b, c), a] = J(a, b, [a, c])$ . This relationship assures us of many examples of a Malcev algebra while examples of a Malcev non-Lie algebra could be found in the literature, Sagle [8.].

The result that got the subject started is the following.

**2.2 Lemma.** ([8.], p. 435) Let  $A$  denote a Malcev algebra and let  $J(A, A, A)$  denote its linear subspace spanned by all elements of the form  $J(a, b, c)$  (for  $a, b, c \in A$ ). Then  $J(A, A, A)$  is an ideal of  $A$   $\square$ .

We shall often simply write  $J$  for the ideal  $J(A, A, A)$ . It should be noted that for a Lie algebra,  $A$ , the ideal  $J(A, A, A)$  is  $\{0\}$ . It then follows that the residue-class algebra  $A/J$  is well defined. In the special case of a Lie algebra  $A$  we have that

$$A/J = A/\{0\} \simeq A.$$

It happens that  $J(A, A, A)$  is the most important ideal for a Malcev non-Lie algebra. Indeed, we have the following.

**2.3 Lemma.** ([8.], p. 435) Let  $A$  denote a Malcev algebra of characteristic not 2 or 3. Then  $J(A, A, A)$  is the smallest ideal of  $A$  for which  $A/J$  is a Lie algebra.  $\square$

This in particular shows that  $J$  is uniquely determined as an ideal of  $A$ . Another distinguished subset of a Malcev algebra is defined in the following.

**2.4 Definition.** Let  $A$  denote a Malcev algebra. The  $J$ -nucleus of  $A$  is given as

$$N = \{x \in A : J(x, A, A) = 0\}. \square$$

This means that  $N$  is the maximal subset of  $A$  for which  $J(N, A, A) = 0$  and may also be shown to an ideal of  $A$ . Hence  $N$  is uniquely determined as an ideal of  $A$ . These two uniquely determined ideals  $J$  and  $N$  of the Malcev algebra  $A$  annihilates each other.

**2.5 Lemma.** ([8.], p. 440) If  $A$  is a Malcev algebra, then  $NJ = 0$ .  $\square$

The third most important ideal of a Malcev algebra is also given below.

**2.6 Definition.** Let  $A$  denote a Malcev algebra and define  $\Delta(x, y)$  as the element of  $A$  for which

$$[z, \Delta(x, y)] = J(z, x, y),$$

for all  $x, y, z \in A$ .

Elementary properties of  $\Delta$  may be found in Sagle[8.] while the crucial result concerning  $\Delta$  is the following.

**2.7 Theorem.** ([8.], p. 432) Let  $A$  denote a Malcev algebra of characteristic not 2 or 3 and let  $\Delta(A, A)$  denote the linear subspace of  $A$  spanned

by all the  $\Delta(x, y)$  (for all  $x, y \in A$ ). Then  $\Delta(A, A)$  is a Lie algebra under commutation.  $\square$

The last result shows that  $\Delta(A, A)$  is not a Lie sub-algebra of  $A$ , since it is only a Lie algebra with respect to the commutation

$$(x, y) = [x, y] - [y, x],$$

$x, y \in A$ . Hence, it is bound to exhibit properties not known to  $A$ . Moreover, by Proposition 5.8 of Sagle [8.], the Lie algebra,  $\Delta(A, A)$  is not uniquely determined from  $A$ .

Our approach in this paper is to employ the well-known fine structure of the Lie (sub-)algebra  $N$  of a Malcev algebra  $A$  to get across to all of  $A$  via the residue-class Lie algebra  $A/J$ . The first major result along this line is the following.

**2.8 Theorem.** ([8.], p. 442) Let  $A$  denote a Malcev algebra such that  $\Delta(A, A)$  is completely reducible in  $A$ , then

$$A = N \oplus J,$$

where  $N$  is completely reduced by  $\Delta(A, A)$  and  $J$  is a semi-simple ideal of  $A$ .  $\square$

The hypotheses of Theorem 1.8 hold if  $A$  is semi-simple. This theorem gives a very useful relationship between the uniquely determined ideals  $N$  and  $J$  (of  $A$ ) in the light of the residue-class Lie algebra  $A/J$ . This is our point of departure in this paper.

### §3. Ideal theory of a Malcev algebra

We use the truth of Theorem 2.8 to define the Lie algebra of a Malcev algebra.

**3.1 Definition.** Let  $A$  denote a Malcev algebra. The Lie algebra corresponding to or of  $A$  is defined as the residue-class (Lie algebra)  $A/J$ .  $\square$

**3.2 Lemma.** Every Malcev algebra has a unique Lie algebra corresponding to it.

**Proof.** Let  $A$  denote any Malcev algebra, then the  $J$ -nucleus  $N$  is non-empty and it is the maximal subset of  $A$  for which  $J(N, A, A) = 0$ . The

direct sum  $A = N \oplus J$  of Theorem 2.8 thus implies that  $A/J \simeq N$ . These respectively imply the existence and uniqueness of the Lie algebra  $A/J$ .  $\square$

In the case where  $A$  is a semi-simple Malcev algebra, its corresponding Lie algebra  $A/J$  is equally semi-simple. We shall denote by  $\phi$  the canonical epimorphism,  $A \twoheadrightarrow A/J$  and shall often make use of the following basic result of *commutative algebra*.

**3.3 Lemma.** ([1.], p. 2) Let  $A$  denote any Malcev algebra. There is a one-to-one order preserving correspondence between the ideals  $\mathfrak{b}$  of  $A$  which contain  $J$  and the ideals  $\bar{\mathfrak{b}}$  of the Lie algebra  $A/J$ , given by  $\mathfrak{b} = \phi^{-1}(\bar{\mathfrak{b}})$ , in which prime ideals correspond to prime ideals.  $\square$

It therefore follows from the last lemma that, even though there are examples to show that it is not always true that the product of ideals of a Malcev algebra  $A$  is an ideal of  $A$  (which is in contrast to the fine property of ideals in a Lie algebra), this failure could now be given a proof and could also be salvaged for ideals of  $A$  contain  $J$  as contained as follows.

**3.4 Proposition.** If  $\mathfrak{a}$  and  $\mathfrak{b}$  are any two ideals of a Malcev algebra,  $A$ , both of which contain  $J$ , then  $\mathfrak{a}\mathfrak{b}$  is an ideal of  $A$ .

**Proof.** The one-to-one correspondence of the last lemma implies that  $\mathfrak{a} = \phi^{-1}(\bar{\mathfrak{a}})$  and  $\mathfrak{b} = \phi^{-1}(\bar{\mathfrak{b}})$ , for some ideals  $\bar{\mathfrak{a}}$  and  $\bar{\mathfrak{b}}$  in the Lie algebra  $A/J$ . Thus

$$\mathfrak{a}\mathfrak{b} = \phi^{-1}(\bar{\mathfrak{a}})\phi^{-1}(\bar{\mathfrak{b}}) = \phi^{-1}(\bar{\mathfrak{a}}\bar{\mathfrak{b}}),$$

where it is known that  $\bar{\mathfrak{a}}\bar{\mathfrak{b}}$  is an ideal in  $A/J \simeq N$ . The crucial Lemma 3.3 thus implies that  $\mathfrak{a}\mathfrak{b}$  is an ideal in  $A$  containing  $J$ .  $\square$

Our approach in this paper will further show that the true correspondence between a Malcev algebra and a Lie algebra is via the process of its residue-class Lie algebra  $A/J$ , which will make the beautiful results of a Lie algebra available to the residue-class (Lie) algebra and then to the Malcev algebra. This could be gleaned through the following commutative diagram:

$$A \xrightarrow{\phi} A/J \xrightarrow{i} N,$$

where  $\phi$  is the canonical map of Lemma 3.3 and  $i$  is the isomorphism of Lemma 3.2. It then means that we have the projection

$$(i \circ \phi) : A \rightarrow N.$$

Clearly,  $i(N + J) = N$ .

The above result, for ideals of  $A$  containing  $J$ , may now be extended to include any subset of the Lie algebra  $A/J \simeq N$ , which would buttress our point of view that it is most profitable to approach the study of a Malcev algebra via its projection map  $(i \circ \phi) : A \rightarrow N$ , into the Lie algebra  $N$ .

**3.5 Lemma.** Let  $\mathfrak{h}$  denote a subalgebra (respectively, an ideal) of  $N$ , then  $H := (i \circ \phi)(\mathfrak{h})$  is a subalgebra (respectively, an ideal) of  $A$ .

**Proof.** The surjectivity of  $\phi$  implies that for any  $n \in N$  there is  $a \in A$  such that  $a = (i \circ \phi)(n)$ .  $\square$

Now that we have a direct access between  $A$  and its Lie algebra,  $N$ , it would be instructive if the extra condition (included in Proposition 3.4) of an ideal containing  $J$  can be removed. This now takes us to the consideration of a special class of ideals of  $N$  meant for this purpose.

**3.6 Definition.** Let  $A$  denote a Malcev algebra. A subset  $X$  of the Lie algebra  $N$  is called an  $i$ -ideal of  $N$  whenever

$$i(n)X \subseteq X,$$

for every  $n \in A/J$ .  $\square$

Lemma 3.5 (together with Lemma 3.3) suggests that the requirement of an  $i$ -ideal  $X$  of  $N$  is equivalent to having  $(i \circ \phi)(a)X \subseteq X$ , for every  $a \in A$ . The following result shows the central importance of  $i$ -ideal of the Lie algebra  $N$  in getting access to and in the understanding of *all* the ideals of a Malcev algebra.

**3.7 Theorem.** Let  $A$  denote any Malcev algebra. Then  $\mathfrak{a}$  is an ideal of  $A$  iff  $(i \circ \phi)(\mathfrak{a})$  is an  $i$ -ideal of  $N$ .

**Proof.** Let  $a \in A/J$  and let  $y \in (i \circ \phi)(\mathfrak{a})$ , then  $y = (i \circ \phi)(x)$ , for some  $x \in \mathfrak{a}$ , an ideal of  $A$ . It is clear (by Lemma 3.5) that  $X := (i \circ \phi)(\mathfrak{a})$  is an additive abelian subgroup of  $N$  and that  $(i \circ \phi)(ax) \subseteq X$ . Note also that

$$(i \circ \phi)(ax) = i(\phi(ax)) = i(a\phi(x)) = i(a)(i \circ \phi)(x) = i(a)y \subseteq i(a)X$$

and that

$$i(a)(i \circ \phi)(x) = (i \circ \phi)(ax) \subseteq X.$$

Hence,  $i(a)X \subseteq X$ , for every  $a \in N$ .

Conversely, let  $\bar{\mathfrak{a}}$  denote an  $i$ -ideal of  $N$ . That is, let  $i(a)\bar{\mathfrak{a}} \subseteq \bar{\mathfrak{a}}$ , for every  $a \in A/J$ . Consider now a subset  $\mathfrak{a}$  of  $A$  as

$$\mathfrak{a} := \{(\phi^{-1} \circ i^{-1})(y) : y \in \bar{\mathfrak{a}}\}.$$

Clearly,  $(i \circ \phi)(\mathfrak{a}) = \bar{\mathfrak{a}}$ . Now let  $r \in A$  and let  $x \in \mathfrak{a}$ , then  $x = (\phi^{-1} \circ i^{-1})(y)$ , for some  $y \in \bar{\mathfrak{a}}$ . Considering the element  $rx$ , we have that  $(i \circ \phi)(rx) = i(\phi(rx)) = i(r\phi(x)) = i(r)(i \circ \phi)(x) \in i(r)(i \circ \phi)(\mathfrak{a}) \in i(r)\bar{\mathfrak{a}} \subseteq \bar{\mathfrak{a}}$ , for all  $r \in A$ . In particular, for all  $r \in N$ . Hence,  $(i \circ \phi)(rx) = i(\phi(rx)) \in \bar{\mathfrak{a}}$ , for all  $r \in N$ . Whence,  $rx \in (\phi^{-1} \circ i^{-1})(\bar{\mathfrak{a}}) = \mathfrak{a}$ .  $\square$

The last Theorem sets up a one-to-one correspondence between (all of) the ideals of a Malcev algebra  $A$  and the  $i$ -ideals of its Lie algebra  $N$ . It equally confirms that the Lie algebra  $N$  is replete with different types of ideals, while the type which is directly concerned with a Malcev algebra is the  $i$ -ideals of  $N$ . We now give a general proof (in contrast to the usual counter-examples) of a major difference on the behaviour of ideals in a Malcev algebra and in a Lie algebra. The next two results show that this negative behaviour of ideals in a Malcev algebra is due to the properties of  $i$ -ideals in  $N$ .

**3.8 Proposition.** The product of two  $i$ -ideals of  $N$  may not always be an  $i$ -ideal of  $N$ .

**Proof.** Every  $i$ -ideal of  $N$  is in particular an ideal in  $A$ . Hence, if  $\bar{\mathfrak{a}}$  and  $\bar{\mathfrak{b}}$  are  $i$ -ideals of  $N$  they also satisfy  $\bar{\mathfrak{a}}\bar{\mathfrak{b}} \subseteq \bar{\mathfrak{a}}$  and  $\bar{\mathfrak{a}}\bar{\mathfrak{b}} \subseteq \bar{\mathfrak{b}}$ . Thus

$$i(a)(\bar{\mathfrak{a}}\bar{\mathfrak{b}}) \subseteq i(a)\bar{\mathfrak{a}} \subseteq \bar{\mathfrak{a}}$$

and

$$i(a)(\bar{\mathfrak{a}}\bar{\mathfrak{b}}) \subseteq i(a)\bar{\mathfrak{b}} \subseteq \bar{\mathfrak{b}},$$

for any  $a \in A/J$ . Therefore,

$$i(a)(\bar{\mathfrak{a}}\bar{\mathfrak{b}}) \subseteq \bar{\mathfrak{a}} \cap \bar{\mathfrak{b}} \not\subseteq \bar{\mathfrak{a}}\bar{\mathfrak{b}}. \quad \square$$

The following may therefore be seen as the first known general result on the product of ideals of a Malcev algebra  $A$ .

**3.9 Theorem.** Product of ideals of a Malcev algebra  $A$  is not always an ideal of  $A$ .

**Proof.** We simply combine Theorem 3.7 with Proposition 3.8.  $\square$

Since there is a one-to-one correspondence between ideals of  $A$  with the  $i$ -ideals of the Lie algebra  $N$ , it follows that among the  $i$ -ideals  $\bar{\mathfrak{a}}$  of  $N$  are those (that form a special class) that corresponds to the ideals  $\phi^{-1}(\bar{\mathfrak{a}})$  of  $A$  that contain  $J$  and therefore satisfy Proposition 3.4. Basic properties of operations on ideals suggest that we seek these (special ones) among the co-prime  $i$ -ideals of  $N$  in order to complete the proof of Proposition 3.8.

**3.10 Lemma.** The product of co-prime  $i$ -ideals of  $N$  is an  $i$ -ideal of  $N$ .

**Proof.** For every  $a \in N$  and for co-prime  $i$ -ideals  $\bar{a}$  and  $\bar{b}$  of  $N$ ,

$$i(a)(\bar{a}\bar{b}) \subseteq \bar{a} \cap \bar{b} \subseteq \bar{a}\bar{b}. \quad \square$$

It is very interesting to confirm if the co-prime  $i$ -ideals of  $N$  correspond to ideals of  $A$  containing  $J$ .

#### §4. Root-space decomposition of a Malcev algebra

We now consider how the root-space decomposition of the Lie algebra  $N$  enriches  $A$ .

**4.1 Definition.** Let  $A$  denote a Malcev algebra and let  $N$  be its corresponding Lie algebra. The *adjoint action* of  $A$  on  $N$  is the map

$$\begin{aligned} ad_{A \times N} : A \times N &\rightarrow N \\ (a, n) &\mapsto ad_{A \times N}(a, n) \\ &=: ad_{A \times N}(a)(n) \end{aligned}$$

defined as

$$ad_{A \times N}(a)(n) = i(ad_{A/J}\phi(a)(i^{-1}(n))).$$

The map  $ad_{A/J} : A/J \rightarrow A/J$  is the adjoint map on the Lie algebra  $A/J$ , given as  $ad_{A/J}(x)(y) = [x, y]$ . The following is immediate.

**4.2 Lemma.** The adjoint action of  $A$  on  $N$  is the adjoint map on  $N$ .

**Proof.**

$$ad_{A \times N}(a)(n) = i(ad_{A/J}\phi(a)(i^{-1}(n))) = i([\phi(a), i^{-1}(n)]) = [(i \circ \phi)(a), n]. \quad \square$$

One of the key results of a Lie algebra is the following.

**4.3 Lemma.** ([4], p. 132) Let  $A$  denote a finite-dimensional Malcev algebra over  $\mathbb{C}$  and let  $\mathfrak{h}$  be a nilpotent Lie subalgebra of the Lie algebra  $N$ , then the generalised weight spaces  $N_\alpha$  of  $N$  relative to  $ad_N \mathfrak{h}$  satisfy

$$(a) \quad N = \bigoplus N_\alpha, \text{ where } N_\alpha \text{ is the set}$$

$$\{X \in N : (ad_H - \alpha(H) \cdot 1)^n X = 0, \forall H \in \mathfrak{h}, \text{ for some integer } n = n(H, X) \geq 0\};$$

$$(b) \quad \mathfrak{h} \subseteq N_0;$$

$$(c) \quad [N_\alpha, N_\beta] \subset N_{\alpha+\beta} \text{ (with the understanding that } N_{\alpha+\beta} \text{ is 0 if } \alpha + \beta \text{ is}$$



not a generalised weight).

We then use these properties of  $N_\alpha$  in  $N$  to develop an equivalent theory for generalised weight spaces of a finite-dimensional Malcev algebra,  $A$ .

**4.4 Theorem.** Let  $A$  denote a finite-dimensional Malcev algebra over  $\mathbb{C}$  and let  $N_\alpha$  be the generalised weight spaces  $N_\alpha$  of  $N$  relative to  $ad_N \mathfrak{h}$  (given in Lemma 4.3 above). Choose a subset  $H$  of  $A$  such that  $\mathfrak{h} := (i \circ \phi)(H)$  is a nilpotent Lie subalgebra of the Lie algebra  $N$ . Then the generalised weight spaces  $A_\alpha$  of  $A$  relative to  $ad_{A \times N} H$  satisfy

(a)  $A = \bigoplus A_\alpha$ , where  $A_\alpha$  is given as the set

$\{a \in A : (ad_{A \times N} h - \alpha(h) \cdot 1)^n a = 0, \forall h \in \mathfrak{h}, \text{ for some integer } n = n(h, a) \geq 0\}$ ;

(b)  $H \subseteq A_0$ ;

(c)  $[A_\alpha, A_\beta] \subset N_{\alpha+\beta}$  (with the understanding that  $N_{\alpha+\beta}$  is 0 if  $\alpha + \beta$  is not a generalised weight).

**Proof.** We set  $A_\alpha = (i \circ \phi)(N_\alpha)$ . Hence,

$$\begin{aligned} [A_\alpha, A_\beta] &= [(i \circ \phi)(N_\alpha), (i \circ \phi)(N_\beta)] = i[\phi(N_\alpha), \phi(N_\beta)] \\ &= i[N_\alpha + J, N_\beta + J] = i([N_\alpha, N_\beta] + J) \subseteq i(N_{\alpha+\beta} + J) = N_{\alpha+\beta}, \end{aligned}$$

since  $N_{\alpha+\beta} \subseteq N$ , in which  $i(N + J) = N$ .  $\square$

There is still a lot of the results of Lie algebras (and Lie groups) that may be proven for a Malcev non-Lie algebra.

## §5. References.

- [1.] Atiyah, M. F. and Macdonald, I. G., *Introduction to commutative algebra*, Addison-Wesley Series in Mathematics, Addison-Wesley, Reading. 1969.
- [2.] Bremner, M. R., Hentzel, I. R., Peresi, L. A., Tvalavadze, M. V., Usefi, H., *Enveloping algebras of Malcev algebras*, *Comment. Math. Univ. Carolin.* **51**, 2 (2010), p. 157 – 174.
- [3.] Harrathi, F., Mabrouk, S., Ncib, O., Silvestrov, S. *Malcev Yang-Baxter equation, weighted  $\mathfrak{D}$ -operators on Malcev algebras and post-Malcev algebras*, arXiv:2206.03629v1 [math.RA] (2022)
- [4.] Knapp, A. W. *Lie Groups Beyond an Introduction*, Progress in Mathematics, **140** (2002)

- [5.] Malcev, A. I., Analytic loops, *Matem. Sb.* **78**, (1955), p. 569 – 578.
- [6.] Paal, E., Note on analytic Moufang loops, *Comment. Math. Univ. Carolin.* **45**, 2 (2004), p. 349 – 354.
- [7.] Perez-Izquierdo, J. M., Shestakov, I. P., An envelope for Malcev algebras, *J. Algebra* **272** (2004), p. 379 – 393.
- [8.] Sagle, A. A., Malcev algebras, *Trans. Amer. Math. Soc.* **101** (1961), p. 426 – 458.
- [9.] Schafer, R. D., *An introduction to nonassociative algebras*, Academic Press, New York, (1966).
- [10.] Varadarajan, V. S., *Lie groups, Lie algebras and their representations*, Springer-Verlag, New York, (1984).